

(19)



(11)

EP 1 836 169 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Claims DE 1
Claims FR 1

(51) Int Cl.:
C07D 213/56 (2006.01)

(86) International application number:
PCT/US2005/047333

(48) Corrigendum issued on:
04.07.2012 Bulletin 2012/27

(87) International publication number:
WO 2006/071960 (06.07.2006 Gazette 2006/27)

(45) Date of publication and mention
of the grant of the patent:
08.02.2012 Bulletin 2012/06

(21) Application number: **05855828.9**

(22) Date of filing: **28.12.2005**

(54) COMPOSITIONS AND METHODS OF TREATING CELL PROLIFERATION DISORDERS

ZUSAMMENSETZUNGEN UND VERFAHREN FÜR DIE BEHANDLUNG VON
ZELLPROLIFERATIONSERKRANKUNGEN

COMPOSITIONS ET PROCÉDÉS DE TRAITEMENT DE TROUBLES IMPLIQUANT UNE
PROLIFÉRATION CELLULAIRE

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR**

(30) Priority: **28.12.2004 US 639834 P**
01.08.2005 US 704551 P
17.10.2005 US 727341 P

(43) Date of publication of application:
26.09.2007 Bulletin 2007/39

(73) Proprietor: **Kinex Pharmaceuticals, LLC**
Buffalo, NY 14203 (US)

(72) Inventor: **HANGAUER, David, G.**
East Amherst, NY 14051 (US)

(74) Representative: **Crump, Julian Richard John et al**
Mintz Levin Cohn Ferris Glovsky & Popeo
Intellectual Property LLP
Alder Castle
10 Noble Street
London EC2V 7JX (GB)

(56) References cited:
EP-A1- 0 463 638 WO-A-96/12473
WO-A-99/01127 WO-A-03/087057
WO-A-2004/056774 WO-A2-03/059903
WO-A2-2005/013914 WO-A2-2005/032493
US-A- 3 868 380 US-A1- 2005 059 713
US-B1- 6 844 367

• **HADJERI, MOHAMED ET AL: "Antimitotic
Activity of 5-Hydroxy-7-methoxy-2- phenyl 4-
quinolones" JOURNAL OF MEDICINAL
CHEMISTRY (2004), 47(20), 4964-4970 CODEN:
JMCMAR; ISSN: 0022-2623, 26 August 2004
(2004-08-26), XP002409072**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 836 169 B9

Description

BACKGROUND OF THE INVENTION

[0001] With more than 563,000 deaths in the United States annually, cancer is the second leading cause of death behind heart disease (UBS Warburg "Disease Dynamics: The Cancer Market," Nov. 8, 2000). Surgery and radiotherapy may be curative if the disease is found early, but current drug therapies for metastatic disease are mostly palliative and seldom offer a long-term cure. Even with the new chemotherapies entering the market, improvement in patient survival is measured in months rather than in years, and the need continues for new drugs effective both in combination with existing agents as first line therapy and as second and third line therapies in treatment of resistant tumors.

[0002] A need remains in the art for improved cell proliferation disorder and cancer treatments.

SUMMARY OF THE INVENTION


[0003] The invention relates to compounds and methods of using the compounds to treat cell proliferation disorders.

[0004] The compounds of the present invention are useful as pharmaceutical agents. For example the compounds may be useful as anti-proliferative agents, for treating mammals, such as for treating humans and animals. The compounds may be used without limitation, for example, as anti-cancer, anti-angiogenesis, anti-metastatic, anti-microbial, anti-bacterial, anti-fungal, anti-parasitic and/or anti-viral agents. The compounds of the invention are useful, for example, in treating lung cancer. The compounds of the invention are also useful, for example, in treating colon cancer. The compounds of the invention are also useful, for example, in treating breast cancer.

[0005] The compounds of the invention are useful in treating diseases and disorders that are modulated by tyrosine kinase inhibition. For example, the compounds of the invention are useful in treating diseases and disorders that are modulated by Src kinase. The compounds of the invention may also be useful in treating diseases and disorders that are modulated by focal adhesion kinase (FAK).

[0006] Compounds of the invention include compounds of Formula I, and salts, solvates or hydrates thereof.

Formula I



where:

[0007] T is absent (*i.e.*, the rings are connected by a bond)

[0008] X_y is CZ, CY, N, or N-O;

[0009] X_z is CZ;

[0010] Y is selected from hydrogen, hydroxyl, halogen, lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl-aryl, and O-benzyl;

[0011] X_a is CR_a , N, or N-O;


[0012] X_b is CR_b , N, or N-O;

[0013] X_c is CN, N, or N-O;

[0014] X_d is CR_d , N, or N-O;

[0015] X_e is CR_e , N, or N-O;


[0016] R_a , R_b , R_c , R_d , R_e , R_4 , R_5 , and R_6 are, independently, hydrogen, hydroxyl, halogen, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl-aryl, O-benzyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-OH, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, COOH, COO-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, SO_2H , SO_2 -lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, or



where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl;

[0017] V is a bond, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- or -OCH₂CH₂CH₂-;

[0018] Z is;




where

[0019] R₁, R₂, and R₃ are independently H or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl;

[0020] n is 0, 1, or 2; and m is 1 or 2;

[0021] R₇, R₈, R₉, R₁₀, and R₁₁ are, independently, hydrogen, hydroxyl, halogen, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, O-lower (C₁, C₄, C₃, C₄, C₅, or C₆) alkyl-aryl, O-benzyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-OH, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-O-C₁, C₂, C₃, C₄, C₅, or C₆ alkyl,




where W is H, or -C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl;

[0022] V is a bond, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂-, or -OCH₂CH₂CH₂-.

[0023] The compounds bi-phenyl-4-carboxylic acid (pyridin-4-ylmethyl)-amide of Synlett; English; 6, 2003; 876-878 and biphenyl-4-carboxylic acid benzylamide of Journal of Heterocyclic Chemistry; English; 18;1981;1305-1308 are excluded from the scope of the invention.

[0024] In certain compounds of the invention, Z is



[0025] Certain compounds of the invention are selected from Compounds 1-136 and 137. For example, the compound of the invention is Compound 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 1, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, or 137 or a compound selected from Table 2.

[0026] Compounds of the invention include Compounds 33, 38, 40, 76, 133, 134, 136 and 137.

[0027] In certain Compounds of Formula I, at least one of X_a , X_b , X_c , X_d and X_e is N


[0028] For example, in the compound of Formula I, X_a is N and each of X_b , X_c , X_d and X_e is CR.

[0029] In certain compounds of Formula I, X_y is CY.

[0030] For example, in certain compounds of Formula I, Y is hydrogen.

[0031] In certain compounds of Formula I, R_b is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. For example, R_b is methoxy or ethoxy. In certain compounds of Formula I, R_b is hydrogen. In other compounds of Formula I, R_b is selected from F, Cl, Br, and I.


[0032] In other compounds of Formula I, R_b is



where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$. For example, V is a bond. In certain compounds of Formula I, W is hydrogen. In other compounds of Formula I, W is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl.

[0033] In certain compounds of Formula I, R_c is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. For example, K is methoxy or ethoxy. In other compounds of Formula I, R_c is hydrogen, F, Cl, Br, or I.

[0034] In other compounds of Formula I, R_c is




where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

[0035] For example, V is a bond. In certain compounds of Formula I, W is hydrogen. In other compounds of Formula I, W is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl.

[0036] In certain compounds of Formula I, R_d is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. For example, R_d is methoxy or ethoxy. In other compounds of Formula I, R_d is hydrogen, F, Cl, Br, or I.

[0037] In other compounds of Formula I, R_d is



where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl; and V is a bond, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- or -OCH₂CH₂CH₂-.

[0038] For example, V is a bond. In certain compounds of Formula I, W is hydrogen. In other compounds of Formula I, W is C₁, C₂, C₃, C₄, C₅, or C₆ alkyl.

[0039] The invention includes a solvate of a compound according to Formula I.







[0040] The invention also includes a hydrate of a compound according to Formula I.


[0041] The invention also includes an acid addition salt of a compound according to Formula I. For example, a hydrochloride salt.

[0042] The invention also includes a pharmaceutically acceptable salt of a compound of Formula I.


[0043] The invention also includes a composition of a compound according to Formula I and at least one pharmaceutically acceptable excipient.

[0044] The invention relates to a compound of Formula I, having a structure according to one of Formulae II-XIII:







; Formula X:




Formula XI:



Formula XII:




Formula XIII:



, or a salt, solvate, hydrate, or prodrug thereof, where:

[0045] R_5 , R_4 , R_5 , R_8 , and R_{10} are, independently, hydrogen, hydroxyl, halogen, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl-aryl, O-benzyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-OH, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, COOH, COO-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, SO₂-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl,




where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl, and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

[0046] For example, in the compound of Formula II-3HB, R_8 is hydrogen, F, Cl, Br, or I. For example, R_8 is F. In certain compounds, R_8 is H.


[0047] In certain compounds of Formula II-XIII, R_b is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. For example, R_b is methoxy or ethoxy.

[0048] In certain compounds of Formula II-XIII, R_b is hydrogen, Cl, Br, or I. In other compounds, in the compound of Formula II-XIII, R_b is




where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl, and V is a bond, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- or -OCH₂CH₂CH₂-.

[0049] In certain compounds of Formula II-308, R₄ is hydrogen, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, F, Cl, Br, or L. In other compounds, in the compound of Formula II-XIII, R₄ is



where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl; and V is a bond, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- or -OCH₂CH₂CH₂-.


[0050] In certain compounds of Formula II-XIII, R₅ is hydrogen, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, F, Cl, Br, or L. In other compounds, in the compound of Formula II-XIII, R₅ is



where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl; and V is a bond, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- or -CH₂CH₂CH₂-.

[0051] In certain compounds of Formula II-XIII, R₁₀ is hydrogen, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, F, Cl, Br, or I. For example, R₁₀ is methoxy, ethoxy or isobutoxy.

[0052] In other compounds of Formula II-XIII, R₁₀ is



where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl; and V is a bond, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- or -OCH₂CH₂CH₂-.

[0053] For example, in the compound of Formula II-XIII, W is hydrogen, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl.

[0054] Certain compounds of the invention include compounds according to Formula II.

[0055] The invention relates to a solvate of a compound according to one of Formulae II-XIII. The invention also relates to a hydrate of a compound according to one of Formulae II-XIII.

[0056] The invention also relates to an acid addition salt of a compound according to one of Formulae II-XIII. For example, a hydrochloride salt.

[0057] The invention also relates to a pharmaceutically acceptable salt of a compound of one of Formulae II-XIII.

[0058] The invention includes compositions comprising a compound according to one of Formulae I-XIII and at least one pharmaceutically acceptable excipient.

[0059] Certain compounds of the invention are non-ATP competitive kinase inhibitors.

[0060] The invention also includes a compound according to one of Formulae I-XIII, or a salt, solvate or hydrate thereof for use in the prevention or treatment of a disease or disorder that is modulated by tyrosine kinase inhibition, wherein the disease or disorder is a cell proliferation disorder or a microbial infection.

[0061] For example, the cell proliferation disorder is selected from cancer, pre-cancer hyperproliferative disorder, psoriasis, diabetic retinopathy, macular degeneration, epidermic and dermoid cysts, lipoma, adenoma, capillary and cutaneous hemangioma, lymphangioma, nevi lesions, teratoma, nephroma, myofibromatosis, osteoplastic tumours and dysplasia.

[0062] The cell proliferation disorder treated or prevented by the compounds of the invention may be a cancer, such as, for example, colon cancer, lung cancer, breast cancer, ovarian cancer, brain cancer, liver cancer, pancreas cancer,

prostate cancer, malignant melanoma, non-melanoma skin cancer, childhood leukemia, lymphoma, multiple myeloma, Hodgkin's disease, acute and chronic leukemia, plasma cell neoplasm, lymphoid neoplasm and cancer associated with AIDS.

[0063] The cell proliferation disorder treated or prevented by the compounds of the invention may be a microbial infection, such as a bacterial, fungal, parasitic or viral infection; ..

[0064] For example, the treatment or prevention of the proliferative disorder may occur through the inhibition of a tyrosine kinase. For example, the tyrosine kinase can be a Src kinase or focal adhesion kinase (FAK).

[0065] The pharmaceutical composition of the invention may modulate a kinase pathway. For example, the kinase pathway is a Src kinase pathway, or a focal adhesion kinase pathway.

[0066] The pharmaceutical composition of the invention may modulate a kinase directly. For example, the kinase is Src kinase, or focal adhesion kinase.

[0067] Certain pharmaceutical compositions of the invention are non-ATP competitive kinase inhibitors.

[0068] Certain pharmaceutical compositions of the invention include a compound elected from Compound 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, and 137. For example, the pharmaceutical composition includes Compound 33, 38, 40, 76, 133, 134, 136 or 137.

[0069] Certain pharmaceutical compositions of the invention include a compound selected from the compounds listed in Table 2.

[0070] A compound of the invention may be used as a pharmaceutical agent. For example, a compound of the invention is used as an anti-proliferative agent, for treating humans and/or animals, such as for treating humans and/or other mammals. The compounds may be used without limitation, for example, as anti-cancer, anti-angiogenesis, anti-microbial, anti-bacterial, anti-fungal, anti-parasitic and/or anti-viral agents. Additionally, the compounds may be used for other cell proliferation-related disorders such as diabetic retinopathy, macular degeneration and psoriasis. Anti-cancer agents include anti-metastatic agents.

[0071] The compound of the invention used as a pharmaceutical agent may be selected from Compounds 1-136 and 137. For example, the compound of the invention used as a pharmaceutical agent is Compound 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, or 137. For example, the compound of the invention used as a pharmaceutical agent is selected from Compounds 33, 38, 40, 76, 133, 134, 136 and 137.

[0072] Certain pharmaceutical agents include a compound selected from the compounds listed in Table 2.

[0073] In one aspect of the invention, a compound of the invention, for example, a compound of Formula I or one of Formulae II-XIII, is used to treat or prevent a cell proliferation disorder in a subject. In one aspect of the embodiment, the cell proliferation disorder is pre-cancer or cancer. In another aspect of the embodiment, the cell proliferation disorder is a hyperproliferative disorder. In another embodiment, prevention or treatment of the cell proliferation disorder, cancer or hyperproliferative disorder occurs through the inhibition of a kinase. In another embodiment, prevention or treatment of the cell proliferation disorder, cancer or hyperproliferative disorder occurs through the inhibition of a tyrosine kinase. In another embodiment, prevention or treatment of the cell proliferation disorder, cancer or hyperproliferative disorder occurs through the inhibition of Src kinase or focal adhesion kinase (FAK). In another embodiment, the subject is a mammal. Preferably, the subject is human.

[0074] The above description sets forth rather broadly the more important features of the present invention in order that the detailed description thereof that follows may be understood, and in order that the present contributions to the art may be better appreciated. Other objects and features of the present invention will become apparent from the following detailed description considered in conjunction with the examples.

DETAILED DESCRIPTION OF THE INVENTION


[0075] The details of one or more embodiments of the invention are set forth in the accompanying description below. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. Other features, objects, and advantages of the invention will be apparent from the description. In the specification, the singular forms also include the plural unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In

the case of conflict, the present specification will control.

[0076] The invention relates to compounds and methods of using compounds to treat cell proliferation disorders.

[0077] The compounds of the present invention are useful as pharmaceutical agents, particularly as anti-proliferative agents, for treating humans and animals, particularly for treating humans and other mammals. The compounds may be used without limitation, for example, as anti-cancer, anti-angiogenesis, anti-metastatic, anti-microbial, anti-bacterial, anti-fungal, anti-parasitic and/or anti-viral agents. The compounds may be used for other cell proliferation-related disorders such as psoriasis.

[0078] Compounds of the invention include compounds of formula I, and salts thereof: Formula I



where:

[0079] T is absent (*i.e.*, the rings are connected by a bond)

[0080] X_y is CZ, CY, N, or N-O;

[0081] X_z is CZ;

[0082] Y is selected from hydrogen, hydroxyl, halogen, lower (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, O-lower (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl-aryl, and O-benzyl;

[0083] X_a is CR_a or N, or N-O;


[0084] X_b is CR_b, N, or N-O;

[0085] X_c is CR_c or N, or N-O;

[0086] X_d is CR_d or N, or N-O;

[0087] X_e is CK_e N, or N-O;

R_a, R_b, R_c, R_d, R_e, R₄, R₅, and R₆ are, independently, hydrogen, hydroxyl, halogen, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, O-lower (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl-aryl, O-benzyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-OH, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-O-lower (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, COOH, COO-lower (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, SO₂H, SO₂-lower (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl,





where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl;

[0088] V is a bond, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- or -OCH₂CH₂CH₂-;

[0089] R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, and R₁₈, are, independently, H or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl;


[0090] Z is:





where R_1, R_2 , and R_3 are independently H or C_1, C_2, C_3, C_4, C_5 , or C_6 alkyl;


[0091] n is 0, 1, or 2 and m is 1 or 2; R_7, R_8, R_9, R_{10} , and R_{11} , are, independently, hydrogen, hydroxyl, halogen, C_1, C_2, C_3, C_4, C_5 , or C_6 alkyl, C_1, C_2, C_3, C_4, C_5 , or C_6 alkoxy, O-lower (C_1, C_2, C_3, C_4, C_5 , or C_6) alkyl-aryl, O-benzyl, C_1, C_2, C_3, C_4, C_5 , or C_6 alkyl-OH, C_1, C_2, C_3, C_4, C_5 , or C_6 alkyl-O- C_1, C_2, C_3, C_4, C_5 , or C_6 alkyl,



where W is H, or C_1, C_2, C_3, C_4, C_5 , or C_6 alkyl, C_1, C_2, C_3, C_4, C_5 , or C_6 alkyl-aryl;

[0092] V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$, or $-OCH_2CH_2CH_2-$.

[0093] In certain compounds of the invention, Z is



[0094] Certain compounds of the invention are selected from Compounds 1-136 and 137. For example, the compound of the invention is Compound 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, or 137 or a compound selected from Table 2.

[0095] Compounds of the invention include Compounds 33, 38, 40, 76, 133, 134, 136 and 137.

[0096] In certain Compounds of Formula I, at least one of X_a, X_b, X_c, X_d and X_e is N.

[0097] For example, in the compound of Formula I, X_a is N and each of X_b, X_c, X_d and X_e is CR.


[0098] In certain compounds of Formula I, X_y is CY.

[0099] For example, in certain compounds of Formula I, Y is hydrogen.

[0100] The compounds of the invention can tolerate a wide variety of functional groups, so various substituted starting materials can be used to synthesize them. The syntheses described herein generally provide the desired final bi-aryl compound at or near the end of the overall process, although it may be desirable in certain instances to further convert the compound to a pharmaceutically acceptable salt or ester thereof.

[0101] In certain compounds of Formula I, R_b is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. For example, R_b is methoxy or ethoxy. In certain compounds of Formula I, R_b is hydrogen. In other compounds of Formula I, R_b is selected from F, Cl, Br, and I. For example, R_b is F.

[0102] In other compounds of Formula I, R_a is



where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$. For example, V is a bond. In certain compounds of Formula I, V is $-CH_2-$, $-CH_2CH_2-$ or $-CH_2CH_2CH_2-$. In other compounds, V is $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.


[0103] In certain compounds of Formula I, W is hydrogen. In other compounds, W is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl. In some compounds, W is methyl.

[0104] In certain compounds of Formula I, R_c is halogen, for example, R_c is F, Cl, Br, or I. In some compounds, n is F. In other compounds, R_c is Cl.

[0105] In some compounds, R_c is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. In some compounds, R_c is methoxy or ethoxy. In some embodiments, R_c is ethoxy.

[0106] In other compounds of Formula I, R is hydrogen.

[0107] In other compounds of Formula I, R_c is




where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl; C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$. In some compounds, V is a bond. In other compounds, V is $-CH_2-$, $-CH_2CH_2-$ or $-CH_2CH_2CH_2-$. In other compounds, V is $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

[0108] In some compounds of Formula I, W is hydrogen. In other compounds, W is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl. In certain compounds, W is methyl.

[0109] In certain compounds of Formula I, R_b is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. For example, R_b is methoxy or ethoxy. In certain compounds of Formula I, R_b is hydrogen. In other compounds of Formula I, R_b is selected from F, Cl, Br, and I. For example, R_b is F.

[0110] In other compounds of Formula I, R_b is



where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$. For example, V is a bond. In certain compounds of Formula I, V is $-CH_2-$, $-CH_2CH_2-$ or $-CH_2CH_2CH_2-$. In other compounds, V is $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

[0111] In certain compounds of Formula I, W is hydrogen. In other compounds, W is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl. In some compounds, W is methyl.


[0112] In certain compounds of Formula I, R_d is halogen, for example, R_d is F, Cl, Br, or I. In some compounds, R_d is F. In other compounds, R_d is Cl.

[0113] In some compounds, R_d is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. In some compounds, R_d is methoxy or ethoxy. In

some embodiments, R_d is ethoxy.

[0114] In other compounds of Formula I, R_d is hydrogen.


[0115] In other compounds of Formula I, R_d is




where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$. In some compounds, V is a bond. In other compounds, V is $-CH_2-$, $-CH_2CH_2-$ or $-CH_2CH_2CH_2-$. In other compounds, V is $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

[0116] In some compounds of Formula I, W is hydrogen. In other compounds, W is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl. In certain compounds, W is methyl.


[0117] The invention relates to a compound of Formula I, having a structure according to one of Formulae II-XIII:





; Formula III:




; Formula IV:




; Formula V:




; Formula VII:




; Formula VIII:




; Formula IX:




; Formula X: ;



Formula XII:




Formula XIII:



, or a salt, solvate, hydrate, or prodrug thereof; where:

[0118] R_b , R_4 , R_5 , R_8 , and R_{10} are, independently, hydrogen, hydroxyl, halogen, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl-aryl, O-benzyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-OH, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, COOH, COO-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, SO_2H , SO_2 -lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl,





where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl, and V is a bond, $-CH_2-$, CH_2CH_2 , $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.


[0119] For example, in the compound of Formula II-XIII, R_8 is hydrogen, F, Cl, Br, or I. For example, R_8 is F. In certain compounds, R_8 is H.

[0120] In certain compounds of Formula II-XIII, R_b is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy. For example, R_b is methoxy or ethoxy. In certain compounds, R_b is ethoxy. In certain compounds, R_b is hydrogen.


[0121] In certain compounds of Formula II-XIII, R_b is Cl, Br, or I. For example, R_b is F or Cl. In other compounds, in the compound of Formula II-XIII, R_b is



[0123] In certain compounds of Formula II-XIII, R₅ is hydrogen, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, F, Cl, Br, or I. For example, R₅ is hydrogen. In some compounds, R₆ is ethoxy. In certain compounds R₅ is F. In other compounds, in the compound of Formula II-XIII, R₅ is



[0125] In other compounds of Formula II-XIII, R₁₀ is











[0126] For example, in the compound of Formula II-XIII W is hydrogen, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl. In some compounds, W is methyl.







[0127] Certain compounds of the invention include compounds according to Formula II.

[0128] Compounds of the invention include those listed in Table 1:







Table 1

Compound #	KX #	Compound
1	1-136	
2	1-305	
3	1-306	
4	1-307	
5	1-308	
6	1-309	
7	1-310	
8	1-311	








(continued)

Compound #	KX #	Compound
9	1-312	
10	1-313	
11	1-314	
12	1-315	
13	1-316	
14	1-317	








(continued)

Compound #	KX #	Compound
15	1-318	
16	1-319	
17	1-320	
18	1-321	
19	1-322	
20	1-323	







(continued)






Compound #	KX #	Compound
21	1-324	
22	1-325	
23	1-326	
24	1-327	
25	1-329	
26	1-357	
27	1-358	

(continued)







Compound #	KX #	Compound
28	2-359	
29	2-368	
30	2-380	
31	2-378	
32		
33	2-381	
34		

(continued)







Compound #	KX #	Compound
35		
36	2-375	
37	2-386	
38	2-377	
39	2-387	
40	2-365	

CC(C)OCc1cc(F)ccc1NC(=O)Cc2ccc(cc2)-c3ccc(OCCN4CCN(C)CC4)cc3







(continued)

Compound #	KX #	Compound
47		
48		
49		
50		
51		
52		







(continued)

Compound #	KX #	Compound
53		
54	2-360	
55	2-369	
56		
57		
58		







(continued)

Compound #	KX #	Compound
59		
60	2-389	
61		
62		
63		
64	2-384	







(continued)

Compound #	KX #	Compound
65		
66	2-388	
67		
68	2-382	
69		
70	2-379	






(continued)

Compound #	KX #	Compound
71		
72	2-373	
73		
74	2-376	
75	2-366	
76	2-361	






(continued)

Compound #	KX #	Compound
77	2-370	
78	2-362	
79	2-363	
80	2-372	
81	2-371	
82	2-364	






(continued)

Compound #	KX #	Compound
83	2-385	
84		
85		
86		
87		

(continued)

Compound #	KX #	Compound
88		
89		
90		
91		
92		






(continued)

Compound #	KX #	Compound
93		
94		
95		
96		
97		







(continued)

Compound #	KX #	Compound
98		 <chem>CCOC1=CC=C(C=C1C2=CC=CC=N2CC(=O)NC3=CC=C(F)C=C3)C4CCN(C)CC4</chem>
99		 <chem>CCOC1=CC=C(C=C1C2=CC=CC=N2CC(=O)NC3=CC(=C(C=C3)F)OC(C)C)C4CCN(C)CC4</chem>
100		 <chem>CCOC1=CC=C(C=C1C2=CC=CC=N2CC(=O)NC3=CC=C(F)C=C3)N4CCOCC4</chem>
101		 <chem>CCOC1=CC=C(C=C1C2=CC=CC=N2CC(=O)NC3=CC(=C(C=C3)F)OC(C)C)N4CCOCC4</chem>
102		 <chem>CCOC1=CC=C(C=C1C2=CC=CC=N2CC(=O)NC3=CC=C(F)C=C3)N4CCOCC4</chem>






(continued)

Compound #	KX #	Compound
103		
104		
105		
106		
107		

(continued)






Compound #	KX #	Compound
108A	1-072 (Chiral Center)	
108B	1-121 (Opposite Enantiomer Of 108A)	
109	1-75	
110	1-62	
111	1-64	
112	1-117	

(continued)







Compound #	KX #	Compound
113		
114	2-390	
115	2-374	
116	2-383	
117		

37





(continued)

Compound #	KX #	Compound
123		
124		
125		
126		
127		

(continued)







Compound #	KX #	Compound
128		
129		
130		
131		
132		
133	2-392	


(continued)


Compound #	KX #	Compound
134	2-391	
135	329-N oxide	
136	2-393	
137	2-394	


[0129] Other Compounds are listed in Table 2.


Table 2


	
	
	














[0130] The invention relates to a solvate of a compound according to one of Formulae I-XIII. The invention also relates to a hydrate of a compound according to one of Formulae .. I-XIII.

[0131] The invention also relates to an acid addition salt of a compound according to one of Formulae I-XIII. For example, a hydrochloride salt.

[0132] The invention also relates to a pharmaceutically acceptable salt of a compound of one of Formulae I-XVIII.

[0133] The invention includes compositions comprising a compound according to one of Formulae I-XIII and at least one pharmaceutically acceptable excipient.

[0134] Certain compounds of the invention are non-ATP competitive kinase inhibitors.

[0135] The invention also includes a method of preventing or treating a cell proliferation disorder by administering a pharmaceutical composition that includes a compound according to one of Formulae I-XIII, or a salt, solvate or hydrate thereof, and at least one pharmaceutically acceptable excipient to a subject in need thereof.

[0136] For example, the cell proliferation disorder is pre-cancer or cancer. The cell proliferation disorder treated or prevented by the compounds of the invention may be a cancer, such as, for example, colon cancer or lung cancer.

[0137] The cell proliferation disorder treated or prevented by the compounds of the invention may be a hyperproliferative disorder

[0138] The cell proliferation disorder treated or prevented by the compounds of the invention may be psoriasis.

[0139] For example, the treatment or prevention of the proliferative disorder may occur through the inhibition of a tyrosine kinase. For example, the tyrosine kinase can be a Src kinase or focal adhesion kinase (FAK).

[0140] The pharmaceutical composition of the invention may modulate a kinase pathway. For example, the kinase pathway is a Src kinase pathway, or focal adhesion kinase pathway.

[0141] The pharmaceutical composition of the invention may modulate a kinase directly. For example, the kinase is Src kinase, or focal adhesion kinase.

[0142] Certain pharmaceutical compositions of the invention are non-ATP competitive kinase inhibitors.

[0143] For example, the compounds of the invention are useful to treat or prevent a microbial infection, such as a bacterial, fungal, parasitic or viral infection.

[0144] Certain pharmaceutical compositions of the invention include a compound selected from Compound 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, and 137. For example, the pharmaceutical composition includes Compound 33, 38, 40, 76, 133, 134, 136 or 137.

[0145] Certain pharmaceutical compositions of the invention include a compound selected from the compounds listed in Table 2.

[0146] A compound of the invention may be used as a pharmaceutical agent. For example, a compound of the invention is used as an anti-proliferative agent, for treating humans and/or animals, such as for treating humans and/or other mammals. The compounds may be used without limitation, for example, as anti-cancer, anti-angiogenesis, anti-microbial, anti-bacterial, anti-fungal, anti-parasitic and/or anti-viral agents. Additionally, the compounds may be used for other cell proliferation-related disorders such as diabetic retinopathy, macular degeneration and psoriasis. Anti-cancer agents include anti-metastatic agents.

[0147] The compound of the invention used as a pharmaceutical agent may be selected from Compounds 1-136 and 137. For example, the compound of the invention used as a pharmaceutical agent is Compound 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102,

103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, or 137. For example, the compound of the invention used as a pharmaceutical agent is selected from Compounds 33, 38, 40, 76, 133, 134, 136 and 137.

[0148] Certain pharmaceutical agents include a compound selected from the compounds listed in Table 2.

[0149] In one aspect of the invention, a compound of the invention, for example, a compound of Formula I or one of Formulae II-XM, is used to treat or prevent a cell proliferation disorder in a subject. In one aspect of the embodiment, the cell proliferation disorder is pre-cancer or cancer. In another aspect of the embodiment, the cell proliferation disorder is a hyperproliferative disorder. In another embodiment, prevention or treatment of the cell proliferation disorder, cancer or hyperproliferative disorder occurs through the inhibition of a kinases. In another embodiment, prevention or treatment of the cell proliferation disorder, cancer or hyperproliferative disorder occurs through the inhibition of a tyrosine kinase. In another embodiment, prevention or treatment of the cell proliferation disorder, cancer or hyperproliferative disorder occurs through the inhibition of Src kinase or focal adhesion kinase (FAK). In another embodiment, the subject is a mammal. Preferably, the subject is human.

[0150] The invention is also drawn to a method of treating or preventing cancer or a proliferation disorder in a subject, comprising administering an effective amount of a compound of the invention, for example, a compound of Formula I or one of Formulae II-XIII. For example, the compound of the invention may be a kinase inhibitor. The compound of the invention may be a non-ATP competitive kinase inhibitor. The compound of the invention may inhibit a kinase directly, or it may affect the kinase pathway.

Definitions

[0151] For convenience, certain terms used in the specification, examples and appended claims are collected here.

[0152] Protein kinases are a large class of enzymes which catalyze the transfer of the γ -phosphate from ATP to the hydroxyl group on the side chain of Ser/Thr or Tyr in proteins and peptides and are intimately involved in the control of various important cell functions, perhaps most notably: signal transduction, differentiation, and proliferation. There are estimated to be about 2,000 distinct protein kinases in the human body, and although each of these phosphorylate particular protein/peptide substrates, they all bind the same second substrate ATP in a highly conserved pocket. About 50% of the known oncogene products are protein tyrosine kinases (PTKs), and their kinase activity has been shown to lead to cell transformation.

[0153] The PTKs can be classified into two categories, the membrane receptor PTKs (e.g. growth factor receptor PTKs) and the non-receptor PTKs (e.g. the Src family of proto-oncogene products and focal adhesion kinase (FAK)). The hyperactivation of Src has been reported in a number of human cancers, including those of the colon, breast, lung, bladder, and skin, as well as in gastric cancer, hairy cell leukemia, and neuroblastoma.

[0154] "Treating", includes any effect, e.g., lessening, reducing, modulating, or eliminating, that results in the improvement of the condition, disease, disorder, etc. "Treating" or "treatment" of a disease state includes: (1) preventing the disease state, i.e. causing the clinical symptoms of the disease state not to develop in a subject that may be exposed to or predisposed to the disease state, but does not yet experience or display symptoms of the disease state; (2) inhibiting the disease state, i.e., arresting the development of the disease state or its clinical symptoms; or (3) relieving the disease state, i.e., causing temporary or permanent regression of the disease state or its clinical symptoms.

[0155] "Disease state" means any disease, disorder, condition, symptom, or indication.

[0156] As used herein, the term "cell proliferative disorder" refers to conditions in which the unregulated and/or abnormal growth of cells can lead to the development of an unwanted condition or disease, which can be cancerous or non-cancerous, for example a psoriatic condition. As used herein, the terms "psoriatic condition" or "psoriasis" refers to disorders involving keratinocyte hyperproliferation, inflammatory cell infiltration, and cytokine alteration.

[0157] In a preferred embodiment, the cell proliferation disorder is cancer. As used herein, the term "cancer" includes solid tumors, such as lung, breast, colon, ovarian, brain, liver, pancreas, prostate, malignant melanoma, non-melanoma skin cancers, as well as hematologic tumors and/or malignancies, such as childhood leukemia and lymphomas, multiple myeloma, Hodgkin's disease, lymphomas of lymphocytic and cutaneous origin, acute and chronic leukemia such as acute lymphoblastic, acute myelocytic or chronic myelocytic leukemia, plasma cell neoplasm, lymphoid neoplasm and cancers associated with AIDS.

[0158] In addition to psoriatic conditions, the types of proliferative diseases which may be treated using the compositions of the present invention are epidermic and dermoid cysts, lipomas, adenomas, capillary and cutaneous hemangiomas, lymphangiomas, nevi lesions, teratomas, nephromas, myofibromatosis, osteoplastic tumors, and other dysplastic masses and the like. The proliferative diseases can include dysplasias and disorders of the like.

[0159] An "effective amount" of a compound of the disclosed invention is the quantity which, when administered to a subject having a disease or disorder, results in regression of the disease or disorder in the subject. Thus, an effective amount of a compound of the disclosed invention is the quantity which, when administered to a subject having a cell proliferation disorder, results in regression of cell growth in the subject. The amount of the disclosed compound to be

administered to a subject will depend on the particular disorder, the mode of administration, co-administered compounds, if any, and the characteristics of the subject, such as general health, other diseases, age, sex, genotype, body weight and tolerances to drugs. The skilled artisan will be able to determine appropriate dosages depending on these and other factors.

[0160] As used herein, the term "effective amount" refers to an amount of a compound, or a combination of compounds, of the present invention effective when administered alone or in combination as an anti-proliferative agent. For example, an effective amount refers to an amount of the compound present in a formulation or on a medical device given to a recipient patient or subject sufficient to elicit biological activity, for example, anti-proliferative activity, such as *e.g.*, anti-cancer activity or anti-neoplastic activity. The combination of compounds optionally is a synergistic combination. Synergy, as described, for example, by Chou and Talalay, *Adv Enzyme Regul.* vol. 22, pp. 27-55 (1984), occurs when the effect of the compounds when administered in combination is greater than the additive effect of the compounds when administered alone as a single agent. In general, a synergistic effect is most clearly demonstrated at sub-optimal concentrations of the compounds. Synergy can be in terms of lower cytotoxicity, or increased anti-proliferative effect, or some other beneficial effect of the combination compared with the individual components.

[0161] "A therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

[0162] A therapeutically effective amount of one or more of the compounds can be formulated with a pharmaceutically acceptable carrier for administration to a human or an animal. Accordingly, the compounds or the formulations can be administered, for example via oral, parenteral, or topical routes, to provide an effective amount of the compound. In alternative embodiments, the compounds prepared in accordance with the present invention can be used to coat or impregnate a medical device, *e.g.*, a stent

[0163] The term "prophylactically effective amount" means an effective amount of a compound or compounds, of the present invention that is administered to prevent or reduce the risk of unwanted cellular proliferation.

[0164] "Pharmacological effect" as used herein encompasses effects produced in the subject that achieve the intended purpose of a therapy. In one preferred embodiment, a pharmacological effect means that primary indications of the subject being treated are prevented, alleviated, or reduced. For example, a pharmacological effect would be one that results in the prevention, alleviation or reduction of primary indications in a treated subject. In another preferred embodiment, a pharmacological effect means that disorders or symptoms of the primary indications of the subject being treated are prevented, alleviated, or reduced. For example, a pharmacological effect would be one that results in the prevention or reduction of primary indications in a treated subject

[0165] With respect to the chemical compounds useful in the present invention, the following terms can be applicable:

[0166] The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (*i.e.*, =O), then 2 hydrogens on the atom are replaced. Keto substituents are not present on aromatic moieties. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (*e.g.*, C=C, C=N, or N=N).

[0167] The present invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium, and isotopes of carbon include C-13 and C-14.

[0168] The compounds described herein may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. *Cis* and *trans* geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic, and geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated. All tautomers of shown or described compounds are also considered to be part of the present invention.

[0169] When any variable (*e.g.*, R₁) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R₁ moieties, then the group may optionally be substituted with up to two R₁ moieties and R¹ at each occurrence is selected independently from the definition of R₁. Also, combinations of substituents and/or variables are permissible, but only if such combinations result in stable compounds.

[0170] When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom in the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible, but only if such combinations result in stable

compounds.

[0171] Compounds of the present invention that contain nitrogens can be converted to N-oxides by treatment with an oxidizing agent (e.g., 3-chloroperoxybenzoic acid (*m*-CPBA) and/or hydrogen peroxides) to afford other compounds of the present invention. Thus, all shown and claimed nitrogen-containing compounds are considered, when allowed by valency and structure, to include both the compound as shown and its N-oxide derivative (which can be designated as N→O or N⁺-O⁻). Furthermore, in other instances, the nitrogens in the compounds of the present invention can be converted to N-hydroxy or N-alkoxy compounds. For example, N-hydroxy compounds can be prepared by oxidation of the parent amine by an oxidizing agent such as *m*-CPBA. All shown and claimed nitrogen-containing compounds are also considered, when allowed by valency and structure, to cover both the compound as shown and its N-hydroxy (i.e., N-OH) and N-alkoxy (i.e., N-OR, wherein R is substituted or unsubstituted C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₃₋₁₄ carbocycle, or 3-14-membered heterocycle) derivatives.

[0172] When an atom or chemical moiety is followed by a subscripted numeric range (e.g., C₁₋₆), the invention is meant to encompass each number within the range as well as all intermediate ranges. For example, "C₁₋₆ alkyl" is meant to include alkyl groups with 1, 2, 3, 4, 5; 6, 1-6, 1-5, 1-4, 1-3, 1-2, 2-6, 2-5, 2-1, 2-3, 3-6, 3-5, 3-4, 4-6, 4-5, and 5-6 carbons.

[0173] As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, C₁₋₆ alkyl is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ alkyl groups. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-pmpyl, n-butyl, s-butyl, t-butyl, n-pentyl, s-pentyl, and n-hexyl. "Alkyl" further includes alkyl groups that have oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more hydrocarbon backbone carbon atoms. In certain embodiments, a straight chain or branched chain alkyl has six or fewer carbon atoms in its backbone (e.g., C₁-C₆ for straight chain, C₃-C₆ for branched chain), and more preferably four or fewer. Likewise, preferred cycloalkyls have from three to eight carbon atoms in their ring structure, and more preferably have five or six carbons in the ring structure.

[0174] Unless the number of carbons is otherwise specified, "lower alkyl" includes an alkyl group, as defined above, but having from one to ten, more preferably from one to six, carbon atoms in its backbone structure. "Lower alkenyl" and "lower alkynyl" have chain lengths of, for example, 2-5 carbon atoms.

[0175] The term "alkyl" also includes both "unsubstituted alkyls" and "substituted alkyls", the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonate, phosphinato, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfanyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Cycloalkyls can be further substituted, e.g., with the substituents described above. An "alkylaryl" or an "aralkyl" moiety is an alkyl substituted with an aryl (e.g., phenylmethyl (benzyl)).

[0176] "Alkenyl" includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double bond. For example, the term "alkenyl" includes straight-chain alkenyl groups (e.g., ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl), branched-chain alkenyl groups, cycloalkenyl (e.g., alicyclic) groups (e.g., cyclopropenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl), alkyl or alkenyl substituted cycloalkenyl groups, and cycloalkyl or cycloalkenyl substituted alkenyl groups. The term "alkenyl" further includes alkenyl groups, which include oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more hydrocarbon backbone carbons. In certain embodiments, a straight chain or branched chain alkenyl group has six or fewer carbon atoms in its backbone (e.g., C₂-C₆ for straight chain, C₃-C₆ for branched chain). Likewise, cycloalkenyl groups may have from three to eight carbon atoms in their ring structure, and more preferably have five or six carbons in the ring structure. The term "C₂-C₆" includes alkenyl groups containing two to six carbon atoms. The term "C₃-C₆" includes alkenyl groups containing three to six carbon atoms.

[0177] The term "alkenyl" also includes both "unsubstituted alkenyls" and "substituted alkenyl", the latter of which refers to alkenyl moieties having substituents replacing a hydrogen on one or more hydrocarbon backbone carbon atoms. Such substituents can include, for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonate, phosphinato, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfanyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0178] "Alkynyl" includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but which contain at least one triple bond. For example, "alkynyl" includes straight-chain alkynyl groups (e.g., ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl), branched-chain alkynyl groups,

and cycloalkyl or cycloalkenyl substituted alkynyl groups. The term "alkynyl" further includes alkynyl groups having oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more hydrocarbon backbone carbons. In certain embodiments, a straight chain or branched chain alkynyl group has six or fewer carbon atoms in its backbone (e.g., C₂-C₆ for straight chain, C₃-C₆ for branched chain). The term "C₂-C₆" includes alkynyl groups containing two to six carbon atoms. The term "C₃-C₆" includes alkynyl groups containing three to six carbon atoms.

[0179] The term "alkynyl" also includes both "unsubstituted alkynyls" and "substituted alkynyl", the latter of which refers to alkynyl moieties having substituents replacing a hydrogen on one or more hydrocarbon backbone carbon atoms. Such substituents can include, for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy, carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonate, phosphinate, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfanyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0180] "Aryl" includes groups with aromaticity, including 5- and 6-membered "unconjugated", or single-ring, aromatic groups that may include from zero to four heteroatoms, as well as "conjugated", or multicyclic, systems with at least one aromatic ring. Examples of aryl groups include benzene, phenyl, pyrrole, furan, thiophene, thiazole, isothiazole, imidazole, triazole, tetrazole, pyrazole, oxazole, isooxazole, pyridine, pyrazine, pyridazine, and pyrimidine, and the like. Furthermore, the term "aryl" includes multicyclic aryl groups, e.g., tricyclic, bicyclic, e.g., naphthalene, benzoxazole, benzodioxazole, benzothiazole, benzoimidazole, benzothiophene, methylenedioxyphenyl, quinoline, isoquinoline, naphthridine, indole, benzofuran, purine, benzofuran, deazapurine, or indolizine. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles", "heterocycles", "heteroaryls" or "heteroaromatics". The aromatic ring can be substituted at one or more ring positions with such substituents as described above, as for example, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkylaminocarbonyl, aralkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aralkylcarbonyl, alkenylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonate, phosphinate, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl groups can also be fused or bridged with alicyclic or heterocyclic rings, which are not aromatic so as to form a multicyclic system (e.g., tetralin, methylenedioxyphenyl).

[0181] As used herein, "halo" or "halogen" refers to fluoro, chloro, bromo, and iodo. The term "perhalogenated" generally refers to a moiety wherein all hydrogens are replaced by halogen atoms.

[O182] "Counterion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, and sulfate.

[0183] The term "non-hydrogen substituent" refers to substituents other than hydrogen. Non-limiting examples include alkyl groups, alkoxy groups, halogen groups, hydroxyl groups, aryl groups, etc.

[0184] As used herein, "carbocycle" or "carbocyclic ring" is intended to mean any stable monocyclic, bicyclic, or tricyclic ring having the specified number of carbons, any of which may be saturated, unsaturated, or aromatic. For example a C₃₋₁₄ carbocycle is intended to mean a mono-, bi-, or tricyclic ring having 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 carbon atoms. Examples of carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantyl, cyclooctyl, cyclooctenyl, cyclooctadienyl, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, and tetrahydronaphthyl. Bridged rings are also included in the definition of carbocycle, including, for example, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane, and [2.2.2]bicyclooMane. A bridged ring occurs when one or more carbon atoms link two non-adjacent carbon atoms. Preferred bridges are one or two carbon atoms. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge. Fused (e.g., naphthyl and tetrahydronaphthyl) and spiro rings are also included.

[0185] As used herein, the term "heterocycle" or "heterocyclic" is intended to mean any stable monocyclic, bicyclic, or tricyclic ring which is saturated, unsaturated, or aromatic and comprises carbon atoms and one or more ring heteroatoms, e.g., 1-2 or 1-3 or 1-4 or 1-5 or 1-6 heteroatoms, independently selected from the group consisting of nitrogen, oxygen, and sulfur. A bicyclic or tricyclic heterocycle may have one or more heteroatoms located in one ring, or the heteroatoms may be located in more than one ring. The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, N→O and S(O)_p where p = 1 or 2). When a nitrogen atom is included in the ring it is either N or NH, depending on whether or not it is attached to a double bond in the ring (*i.e.*, a hydrogen is present if needed to maintain the tri-valency of the nitrogen atom). The nitrogen atom may be substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another substituent, as defined). The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom that results in a stable structures. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen

atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. Bridged rings are also included in the definition of heterocycle. A bridged ring occurs when one or more atoms (*i.e.*, C, O, N, or S) link two non-adjacent carbon or nitrogen atoms. Preferred bridges include, but are not limited to, one carbon atom, two carbon atoms, one nitrogen atom, two nitrogen atoms, and a carbon-nitrogen group. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge. Spiro and fused rings are also included.

[0186] As used herein, the term "aromatic heterocycle" or "heteroaryl" is intended to mean a stable 5, 6, or 7-membered monocyclic or bicyclic aromatic heterocyclic ring or 7, 8, 9, 10, 11, or 12-membered bicyclic aromatic heterocyclic ring which consists of carbon atoms and one or more heteroatoms, *e.g.*, 1 or 1-2 or 1-3 or 1-4 or 1-5 or 1-6 heteroatoms, independently selected from the group consisting of nitrogen, oxygen, and sulfur. In the case of bicyclic heterocyclic aromatic rings, only one of the two rings needs to be aromatic (*e.g.*, 2,3-dihydroindole), though both may be (*e.g.*, quinoline). The second ring can also be fused or bridged as defined above for heterocycles. The nitrogen atom may be substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another substituent, as defined). The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, N→O and S(O)_p, where p = 1 or 2). It is to be noted that total number of S and O atoms in the aromatic heterocycle is not more than 1.

[0187] Examples of heterocycles include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolyl, carbazolyl, 4*H*-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2*H*,6*H*-1,5,2-dithiazinyl, dihydrofuro[2,3-*b*]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, imidazolyl, 1*H*-indazolyl, indolenyl, indolyl, indolizyl, indolyl, 3*H*-indolyl, isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, methylenedioxyphenyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxindolyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2*H*-pyrrolinyl, pyrrolyl, quinazolinyl, quinolinyl, 4*H*-quinolizyl, quinoxalyl, quinuclidinyl, tetrahydrofuran, tetrahydroisoquinolinyl, tetrahydroquinolinyl, tetrazolyl, 6*H*-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl.

[0188] "Acyl" includes compounds and moieties that contain the acyl radical (CH₃CO-) or a carbonyl group. "Substituted acyl" includes acyl groups where one or more of the hydrogen atoms are replaced by for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinate, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0189] "Acylamino" includes moieties wherein an acyl moiety is bonded to an amino group. For example, the term includes alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido groups.

[0190] "Aroyl" includes compounds and moieties with an aryl or heteroaromatic moiety bound to a carbonyl group. Examples of aroyl groups include phenylcarboxy, naphthyl carboxy, etc.

[0191] "Alkoxyalkyl", "alkylaminoalkyl" and "thioalkoxyalkyl" include alkyl groups, as described above, which further include oxygen, nitrogen or sulfur atoms replacing one or more hydrocarbon backbone carbon atoms, *e.g.*, oxygen, nitrogen or sulfur atoms.

[0192] The term "alkoxy" or "alkoxyl" includes substituted and unsubstituted alkyl, alkenyl, and alkynyl groups covalently linked to an oxygen atom. Examples of alkoxy groups (or alkoxy radicals) include methoxy, ethoxy, isopropoxy, propoxy, butoxy, and pentoxy groups. Examples of substituted alkoxy groups include halogenated alkoxy groups. The alkoxy groups can be substituted with groups such as alkenyl, alkenyl halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinate, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moieties. Examples of halogen substituted alkoxy groups include, but are not limited to, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chloromethoxy, dichloromethoxy, and trichloromethoxy.

[0193] The term "thiocarbonyl" or "thiocarboxy" includes compounds and moieties which contain a carbon connected with a double bond to a sulfur atom.

[0194] The term "ether" includes compounds or moieties which contain an oxygen bonded to two different carbon atoms or heteroatoms. For example, the term includes "alkoxyalkyl" which refers to an alkyl, alkenyl, or alkynyl group covalently bonded to an oxygen atom which is covalently bonded to another alkyl group.

[0195] The term "ester" includes compounds and moieties which contain a carbon or a heteroatom bound to an oxygen atom which is bonded to the carbon of a carbonyl group. The term "ester" includes alkoxycarboxy groups such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, pentoxycarbonyl, etc. The alkyl, alkenyl, or alkynyl groups are as defined above.

[0196] The term "thioether" includes compounds and moieties which contain a sulfur atom bonded to two different carbon or heteroatoms. Examples of thioethers include, but are not limited to alkthioalkyls, alkthioalkenyls, and alkthioalkynyls. The term "alkthioalkyls" include compounds with an alkyl, alkenyl, or alkynyl group bonded to a sulfur atom which is bonded to an alkyl group. Similarly, the term "alkthioalkenyls" and alkthioalkynyls" refer to compounds or moieties wherein an alkyl, alkenyl, or alkynyl group is bonded to a sulfur atom which is covalently bonded to an alkynyl group.

[0197] The term "hydroxy" or "hydroxyl," includes groups with an -OH or -O⁻.

[0198] "Polycyclyl" or "polycyclic radical" refers to two or more cyclic rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings. Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle can be substituted with such substituents as described above, as for example, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, alkylaminocarbonyl, aralkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aralkylcarbonyl, alkenylcarbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfanyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0199] An "anionic group," as used herein, refers to a group that is negatively charged at physiological pH. Preferred anionic groups include carboxylate, sulfate, sulfonate, sulfinate, sulfamate, tetrazolyl, phosphate, phosphonate, phosphinate, or phosphorothioate or functional equivalents thereof. "Functional equivalents" of anionic groups are intended to include bioisosteres, *e.g.*, bioisosteres of a carboxylate group. Bioisosteres encompass both classical bioisosteric equivalents and non-classical bioisosteric equivalents. Classical and non-classical bioisosteres are known in the art (see, *e.g.*, Silverman, R. B. *The Organic Chemistry of Drug Design and Drug Action*, Academic Press, Inc.: San Diego, Calif., 1992, pp.19-23). A particularly preferred anionic group is a carboxylate.

[0200] In the present specification, the structural formula of the compound represents a certain isomer for convenience in some cases, but the present invention includes all isomers such as geometrical isomer, optical isomer based on an asymmetrical carbon, stereoisomer, tautomer and the like which occur structurally and an isomer mixture and is not limited to the description of the formula for convenience, and may be any one of isomer or a mixture. Therefore, an asymmetrical carbon atom may be present in the molecule and an optically active compound and a racemic compound may be present in the present compound, but the present invention is not limited to them and includes any one. In addition, a crystal polymorphism may be present but is not limiting, but any crystal form may be single or a crystal form mixture, or an anhydride or hydrate. Further, so-called metabolite which is produced by degradation of the present compound *in vivo* is included in the scope of the present invention.

[0201] "Isomerism" means compounds that have identical molecular formulae but that differ in the nature or the sequence of bonding of their atoms or in the arrangement of their atoms in space. Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereoisomers", and stereoisomers that are non-superimposable mirror images are termed "enantiomers", or sometimes optical isomers. A carbon atom bonded to four nonidentical substituents is termed a "chiral center".

[0202] "Chiral isomer" means a compound with at least one chiral center. It has two enantiomeric forms of opposite chirality and may exist either as an individual enantiomer or as a mixture of enantiomers. A mixture containing equal amounts of individual enantiomeric forms of opposite chirality is termed a "racemic mixture". A compound that has more than one chiral center has 2^{n-1} enantiomeric pairs, where n is the number of chiral centers. Compounds with more than one chiral center may exist as either an individual diastereomer or as a mixture of diastereomers, termed a "diastereomeric mixture". When one chiral center is present, a stereoisomer may be characterized by the absolute configuration (R or S) of that chiral center. Absolute configuration refers to the arrangement in space of the substituents attached to the chiral center. The substituents attached to the chiral center under consideration are ranked in accordance with the Sequence Rule of Cahn, Ingold and Prelog. (Cahn et al, *Angew. Chem Inter. Edit.* 1966, 5, 385; errata 511; Cahn et al., *Angew Chem.* 1966, 78, 413; Cahn and Ingold, *J. Chem Soc.* 1951 (London), 612; Cahn et al., *Experientia* 1956, 12, 81; Cahn, J., *Chem Educ.* 1964, 41, 116).

[0203] "Geometric Isomers" means the diastereomers that owe their existence to hindered rotation about double bonds. These configurations are differentiated in their names by the prefixes *cis* and *trans*, or *Z* and *E*, which indicate that the groups are on the same or opposite side of the double bond in the molecule according to the Cahn-Ingold-

Prelog rules.

[0204] Further, the structures and other compounds discussed in this application include all atropic isomers thereof. "Atropic isomers" are a type of stereoisomer in which the atoms of two isomers are arranged differently in space. Atropic isomers owe their existence to a restricted rotation caused by hindrance of rotation of large groups about a central bond. Such atropic isomers typically exist as a mixture, however as a result of recent advances in chromatography techniques, it has been possible to separate mixtures of two atropic isomers in select cases.

[0205] The terms "crystal polymorphs" or "polymorphs" or "crystal forms" means crystal structures in which a compound (or salt or solvate thereof) can crystallize in different crystal packing arrangements, all of which have the same elemental composition. Different crystal forms usually have different X-ray diffraction patterns, infrared spectral, melting points, density hardness, crystal shape, optical and electrical properties, stability and solubility. Recrystallization solvent, rate of crystallization, storage temperature, and other factors may cause one crystal form to dominate. Crystal polymorphs of the compounds can be prepared by crystallization under different conditions.

[0206] Additionally, the compounds of the present invention, for example, the salts of the compounds, can exist in either hydrated or unhydrated (the anhydrous) form or as solvates with other solvent molecules. Nonlimiting examples of hydrates include monohydrates, dehydrates, etc. Nonlimiting examples of solvates include ethanol solvates, acetone solvates, etc.

[0207] "Solvates" means solvent addition forms that contain either stoichiometric or non stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water the solvate formed is a hydrate, when the solvent is alcohol, the solvate formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water with one of the substances in which the water retains its molecular state as H₂O, such combination being able to form one or more hydrate.

[0208] "Tautomers" refers to compounds whose structures differ markedly in arrangement of atoms, but which exist in easy and rapid equilibrium. It is to be understood that compounds of Formula I may be depicted as different tautomers. It should also be understood that when compounds have tautomeric forms, all tautomeric forms are intended to be within the scope of the invention, and the naming of the compounds does not exclude any tautomer form.

[0209] Some compounds of the present invention can exist in a tautomeric form which are also intended to be encompassed within the scope of the present invention.


[0210] The compounds, salts and prodrugs of the present invention can exist in several tautomeric forms, including the enol and imine form, and the keto and enamine form and geometric isomers and mixtures thereof. All such tautomeric forms are included within the scope of the present invention. Tautomers exist as mixtures of a tautomeric set in solution. In solid form, usually one tautomer predominates. Even though one tautomer may be described, the present invention includes all tautomers of the present compounds

[0211] A tautomer is one of two or more structural isomers that exist in equilibrium and are readily converted from one isomeric form to another. This reaction results in the formal migration of a hydrogen atom accompanied by a switch of adjacent conjugated double bonds. In solutions where tautomerization is possible, a chemical equilibrium of the tautomers will be reached. The exact ratio of the tautomers depends on several factors, including temperature, solvent, and pH. The concept of tautomers that are interconvertable by tautomerizations is called tautomerism.

[0212] Of the various types of tautomerism that are possible, two are commonly observed. In keto-enol tautomerism a simultaneous shift of electrons and a hydrogen atom occurs. Ring-chain tautomerism, is exhibited by glucose. It arises as a result of the aldehyde group (-CHO) in a sugar chain molecule reacting with one of the hydroxy groups (-OH) in the same molecule to give it a cyclic (ring-shaped) form.

[0213] Tautomerizations are catalyzed by: Base: 1. deprotonation; 2. formation of a delocalized anion (e.g. an enolate); 3. protonation at a different position of the anion; Acid: 1. protonation; 2. formation of a delocalized cation; 3. deprotonation at a different position adjacent to the cation.

[0214] Common tautomeric pairs are: ketone - enol, amide - nitrile, lactam - lactim, amide - imidic acid tautomerism in heterocyclic rings (e.g. in the nucleobases guanine, thymine, and cytosine), amine - enamine and enamine - enamine. Examples include:



[0215] It will be noted that the structure of some of the compounds of the invention include asymmetric carbon atoms. It is to be understood accordingly that the isomers arising from such asymmetry (e.g., all enantiomers and diastereomers) are included within the scope of the invention, unless indicated otherwise. Such isomers can be obtained in substantially pure form by classical separation techniques and by stereochemically controlled synthesis. Furthermore, the structures and other compounds and moieties discussed in this application also include all tautomers thereof. Alkenes can include either the E- or Z-geometry, where appropriate. The compounds of this invention may exist in stereoisomeric form, therefore can be produced as individual stereoisomers or as mixtures.

[0216] As used herein, the term "analog" refers to a chemical compound that is structurally similar to another but differs slightly in composition (as in the replacement of one atom by an atom of a different element or in the presence of a particular functional group, or the replacement of one functional group by another functional group). Thus, an analog is a Compound that is similar or comparable in function and appearance, but not in structure or .. origin to the reference compound.

[0217] As defined herein, the term "derivative", refers to compounds that have a common core structure, and are substituted with various groups as described herein. For example, all of the compounds represented by formula I are indole derivatives, and have formula I as a common core.

[0218] The term "bioisostere" refers to a compound resulting from the exchange of an atom or of a group of atoms with another, broadly similar, atom or group of atoms. The objective of a bioisosteric replacement is to create a new compound with similar biological properties to the parent compound. The bioisosteric replacement may be physico-chemically or topologically based. Examples of carboxylic acid bioisosteres include acyl sulfonimides, tetrazoles, sulfonates, and phosphonates. See, e.g., Patani and La Voie, Chem. Rev. 96, 3147-3176 (1996).

[0219] A "pharmaceutical composition" is a formulation containing the disclosed compounds in a form suitable for administration to a subject. In one embodiment, the pharmaceutical composition is in bulk or in unit dosage form. The unit dosage form is any of a variety of forms, including, for example, a capsule, an IV bag, a tablet, a single pump on an aerosol inhaler, or a vial. The quantity of active ingredient (e.g., a formulation of the disclosed compound or salt, hydrate, solvate, or isomer thereof) in a unit dose of composition is an effective amount and is varied according to the particular treatment involved. One skilled in the art will appreciate that it is sometimes necessary to make routine variations to the dosage depending on the age and condition of the patient. The dosage will also depend on the route of administration. A variety of routes are contemplated, including oral, pulmonary, rectal, Parenteral, transdermal, subcutaneous, intravenous, intramuscular, intraperitoneal, inhalational, buccal, sublingual, intrapleural, intrathecal, intranasal, and the like. Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. In a preferred embodiment, the active compound is mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants that are required.

[0220] The term "flash dose" refers to compound formulations that are rapidly dispersing dosage forms.

[0221] The term "immediate release" is defined as a release of compound from a dosage form in a relatively brief period of time, generally up to about 60 minutes. The term "modified release" is defined to include delayed release, extended release, and pulsed release. The term "pulsed release" is defined as a series of releases of drug from a dosage form. The term "sustained release" or "extended release" is defined as continuous release of a compound from a dosage form over a prolonged period.

[0222] A "subject" includes mammals, e.g., humans, companion animals (e.g., dogs, cats, birds, and the like), farm animals (e.g., cows, sheep, pigs, horses, fowl, and the like) and laboratory animals (e.g., rats, mice, guinea pigs, birds, and the like). Most preferably, the subject is human.

[0223] As used herein, the phrase "pharmaceutically acceptable" refers to those compounds, materials, compositions,

carriers, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0224] "Pharmaceutically acceptable excipient" means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes excipient that is acceptable for veterinary use as well as human pharmaceutical use. A "pharmaceutically acceptable excipient" as used in the specification and claims includes both one and more than one such excipient

[0225] The compounds of the invention are capable of further forming salts. All of these forms are also contemplated within the scope of the claimed invention.

[0226] "Pharmaceutically acceptable salt" of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compounds.

[0227] As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines, alkali or organic salts of acidic residues such as carboxylic acids, and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound foiled, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include, but are not limited to, those derived from inorganic and organic acids selected from 2-acetoxybenzoic, 2-hydroxyethane sulfonic, acetic, ascorbic, benzene sulfonic, benzoic, bicarbonic, carbonic, citric, edetic, ethane disulfonic, 1,2-ethane sulfonic, fumaric, glucoheptonic, gluconic, glutamic, glycolic, glycollyarsanilic, hexylresorcinic, hydrabamic, hydrobromic, hydrochloric, hydroiodic, hydroxymaleic, hydroxynaphthoic, isethionic, lactic, lactobionic, lauryl sulfonic, maleic, malic, mandelic, methane sulfonic, napsylic, nitric, oxalic, pamoic; pantothenic, phenylacetic, phosphoric, polygalacturonic, propionic, salicylic, stearic, subacetic, succinic, sulfamic, sulfanilic, sulfuric, tannic, tartaric, toluene sulfonic, and the commonly occurring amine acids, e.g., glycine, alanine, phenylalanine, arginine, etc.

[0228] Other examples include hexanoic acid, cyclopentane propionic acid, pyruvic acid, malonic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo-[2.2.2]-oct-2-ene-1-carboxylic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, muconic acid, and the like. The invention also encompasses salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine, and the like.

[0229] It should be understood that all references to pharmaceutically acceptable salts include solvent addition forms (solvates) or crystal forms (polymorphs) as defined herein, of the same salt

[0230] The pharmaceutically acceptable salts of the present invention can be synthesized from a parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 18th ed. (Mack Publishing Company, 1990). For example, salts can include, but are not limited to, the hydrochloride and acetate salts of the aliphatic amine-containing, hydroxyl amine-containing, and imine-containing compounds of the present invention.

[0231] The compounds of the present invention can also be prepared as esters, for example pharmaceutically acceptable esters. For example a carboxylic acid function group in a compound can be converted to its corresponding ester, e.g., a methyl, ethyl, or other ester. Also, an alcohol group in a compound can be converted to its corresponding ester, e.g., an acetate, propionate, or other ester.

[0232] The compounds of the present invention can also be prepared as prodrugs, for example pharmaceutically acceptable prodrugs. The terms "pro-drug" and "prodrug" are used interchangeably herein and refer to any compound which releases an active parent drug *in vivo*. Since prodrugs are known to enhance numerous desirable qualities of pharmaceuticals (e.g., solubility, bioavailability, manufacturing, etc.) the compounds of the present invention can be delivered in prodrug form. Thus, the present invention is intended to cover prodrugs of the presently claimed compounds, methods of delivering the same and compositions containing the same. "Prodrugs" are intended to include any covalently bonded carriers that release an active parent drug of the present invention *in vivo* when such prodrug is administered to a subject. Prodrugs the present invention are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of the present invention wherein a hydroxy, amino, sulfhydryl, carboxy, or carbonyl group is bonded to any group that, may be cleaved *in vivo* to form a free hydroxyl, free amino, free sulfhydryl, free carboxy or free carbonyl group, respectively.

[0233] Examples of prodrugs include, but are not limited to, esters (e.g., acetate, dialkylaminoacetates, formates,

phosphates, sulfates, and benzoate derivatives) and carbamates (e.g., N,N-dimethylaminocarbonyl) of hydroxy functional groups, esters groups (e.g. ethyl esters, morpholinoethanol esters) of carboxyl functional groups, N-acyl derivatives (e.g. N-acetyl) N-Mannich bases, Schiff bases and enamines of amino functional groups, oximes, acetals, ketals and enol esters of ketone and aldehyde functional groups in compounds of Formula I, and the like, See Bundegaard, H.

"Design of Prodrugs" p1-92, Elsevier, New York-Oxford (1985).

[0234] "Protecting group" refers to a grouping of atoms that when attached to a reactive group in a molecule masks, reduces or prevents that reactivity. Examples of protecting groups can be found in Green and Wuts, Protective Groups in Organic Chemistry, (Wiley, 2nd ed. 1991); Harrison and Harrison et al., Compendium of Synthetic Organic Methods, Vols. 1-8 (John Wiley and Sons, 1971-1996); and Kocienski, Protecting Groups, (Verlag, 3rd ed. 2003).

[0235] The term "amine protecting group" is intended to mean a functional group that converts an amine, amide, or other nitrogen-containing moiety into a different chemical group that is substantially inert to the conditions of a particular chemical reaction. Amine protecting groups are preferably removed easily and selectively in good yield under conditions that do not affect other functional groups of the molecule. Examples of amine protecting groups include, but are not limited to, formyl, acetyl, benzyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, *t*-butyloxycarbonyl (Boc), *p*-methoxybenzyl, methoxymethyl, tosyl, trifluoroacetyl, trimethylsilyl (TMS), fluorenyl-methyloxycarbonyl, 2-trimethylsilyl-ethyloxycarbonyl, 1-methyl-1-(4-biphenyl) ethoxycarbonyl, allyloxycarbonyl, benzyloxycarbonyl (CBZ), 2-trimethylsilyl-ethanesulfonyl (SES), trityl and substituted trityl groups, 9-fluorenylmethyloxycarbonyl (Fmoc), nitro-veratryloxycarbonyl (NVOC), and the like. Other suitable amine protecting groups are straightforwardly identified by those of skill in the art.

[0236] Representative hydroxy protecting groups include those where the hydroxy group is either acylated or alkylated such as benzyl, and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers and allyl ethers.

[0237] "Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

[0238] In the specification, the singular forms also include the plural, unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In the case of conflict, the present specification will control.

[0239] All percentages and ratios used herein, unless otherwise indicated, are by weight.

[0240] "combination therapy" (or "co-therapy") includes the administration of a compound of the invention and at least a second agent as part of a specific treatment regimen intended to provide the beneficial effect from the co-action of these therapeutic agents. The beneficial effect of the combination includes, but is not limited to, pharmacokinetic or pharmacodynamic co-action resulting from the combination of therapeutic agents. Administration of these therapeutic agents in combination typically is carried out over a defined time period (usually minutes, hours, days or weeks depending upon the combination selected). "Combination therapy" may, but generally is not, intended to encompass the administration of two or more of these therapeutic agents as part of separate monotherapy regimens that incidentally and arbitrarily result in the combinations of the present invention.

[0241] "Combination therapy" is intended to embrace administration of these therapeutic agents in a sequential manner, that is, wherein each therapeutic agent is administered at a different time, as well as administration of these therapeutic agents, or at least two of the therapeutic agents, in a substantially simultaneous manner. Substantially simultaneous administration can be accomplished, for example, by administering to the subject a single capsule having a fixed ratio of each therapeutic agent or in multiple, single capsules for each of the therapeutic agents. Sequential or substantially simultaneous administration of each therapeutic agent can be effected by any appropriate route including, but not limited to, oral routes, intravenous routes, intramuscular routes, and direct absorption through mucous membrane tissues. The therapeutic agents can be administered by the same route or by different routes. For example, a first therapeutic agent of the combination selected may be administered by intravenous injection while the other therapeutic agents of the combination may be administered orally. Alternatively, for example, all therapeutic agents may be administered orally or all therapeutic agents may be administered by intravenous injection. The sequence in which the therapeutic agents are administered is not narrowly critical.

[0242] "Combination therapy" also embraces the administration of the therapeutic agents as described above in further combination with other biologically active ingredients and non-drug therapies (e.g., surgery or radiation treatment). Where the combination therapy further comprises a non-drug treatment, the non-drug treatment may be conducted at any suitable time so long as a beneficial effect from the co-action of the combination of the therapeutic agents and non-drug treatment is achieved. For example, in appropriate cases, the beneficial effect is still achieved when the non-drug treatment is temporally removed from the administration of the therapeutic agents, perhaps by days or even weeks.

[0243] Throughout the description, where compositions are described as having, including, or comprising specific components, it is contemplated that compositions also consist essentially of or consist of, the recited components. Similarly, where processes are described as having, including, or comprising specific process steps, the processes also consist essentially of, or consist of, the recited processing steps. Further, it should be understood that the order of steps or order for performing certain actions are immaterial so long as the invention remains operable. Moreover, two or more steps or actions may be conducted simultaneously.

[0244] The compounds, or pharmaceutically acceptable salts thereof is administered orally, nasally, transdermally, pulmonary, inhalationally, buccally, sublingually, intraperitoneally, subcutaneously, intramuscularly, intravenously, rectally, intrapleurally, intrathecally and parenterally. In a preferred embodiment, the compound is administered orally. One skilled in the art will recognize the advantages of certain routes of administration.

[0245] The dosage regimen utilizing the compounds is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration, the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

[0246] Techniques for formulation and administration of the disclosed compounds of the invention can be found in Remington: the Science and Practice of Pharmacy, 19th edition, Mack Publishing Co., Easton, PA (1995). In an embodiment, the compounds described herein, and the pharmaceutically acceptable salts thereof, are used in pharmaceutical preparations in combination with a pharmaceutically acceptable carrier or diluent. Suitable pharmaceutically acceptable carriers include inert solid fillers or diluents and sterile aqueous or organic solutions. The compounds will be present in such pharmaceutical compositions in amounts sufficient to provide the desired dosage amount in the range described herein.

[0247] In one embodiment, the compound is prepared for oral administration, wherein the disclosed compounds or salts thereof are combined with a suitable solid or liquid carrier or diluent to form capsules, tablets, pills, powders, syrups, solutions, suspensions and the like.

[0248] The tablets, pills, capsules, and the like contain from about 1 to about 99 weight percent of the active ingredient and a binder such as gum flu acacias, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch or alginic acid; a lubricant such as magnesium stearate; and/or a sweetening agent such as sucrose, lactose, saccharin, xylitol, and the like. When a dosage unit form is a capsule, it often contains, in addition to materials of the above type, a liquid carrier such as a fatty oil.

[0249] In some embodiments, various other materials are present as coatings or to modify the physical form of the dosage unit. For instance, in some embodiments, tablets are coated with shellac, sugar or both. In some embodiments, a syrup or elixir contains, in addition to the active ingredient, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and a flavoring such as cherry or orange flavor, and the like.

[0250] For some embodiments relating to parental administration, the disclosed compounds, or salts, solvates, tautomers or polymorphs thereof, can be combined with sterile aqueous or organic media to form injectable solutions or suspensions. Injectable compositions are preferably aqueous isotonic solutions or suspensions. The compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. The compositions are prepared according to conventional mixing, granulating or coating methods, respectively, and contain about 0.1 to 75%, preferably about 1 to 50%, of the active ingredient.

[0251] For example, injectable solutions are produced using solvents such as sesame or peanut oil or aqueous propylene glycol, as well as aqueous solutions of water-soluble pharmaceutically-acceptable salts of the compounds. In some embodiments, dispersions are prepared in glycerol, liquid polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. The terms "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal; intraperitoneal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

[0252] For rectal administration, suitable pharmaceutical compositions are, for example, topical preparations, suppositories or enemas. Suppositories are advantageously prepared from fatty emulsions or suspensions. The compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. The compositions are prepared according to conventional mixing, granulating or coating methods, respectively, and contain about 0.1 to 75%, preferably about 1 to 50%, of the active ingredient.

[0253] In some embodiments, the compounds are formulated to deliver the active agent by pulmonary administration, e.g., administration of an aerosol formulation containing the active agent from, for example, a manual pump spray, nebulizer or pressurized metered-dose inhaler. In some embodiments, suitable formulations of this type also include other agents, such as antistatic agents, to maintain the disclosed compounds as effective aerosols.

[0254] A drug delivery device for delivering aerosols comprises a suitable aerosol canister with a metering valve containing a pharmaceutical aerosol formulation as described and an actuator housing adapted to hold the canister and allow for drug delivery. The canister in the drug delivery device has a headspace representing greater than about 15% of the total volume of the canister. Often, the polymer intended for pulmonary administration is dissolved, suspended or emulsified in a mixture of a solvent, surfactant and propellant. The mixture is maintained under pressure in a canister.

that has been sealed with a metering valve.

[0255] For nasal administration, either a solid or a liquid carrier can be used. The solid carrier includes a coarse powder having particle size in the range of, for example, from about 20 to about 500 microns and such formulation is administered by rapid inhalation through the nasal passages; In some embodiments where the liquid carrier is used, the formulation is administered as a nasal spray or drops and includes oil or aqueous solutions of the active ingredients.

[0256] Also contemplated are formulations that are rapidly dispersing dosage forms, also known as "flash dose" forms. In particular, some embodiments of the present invention are formulated as compositions that release their active ingredients within a short period of time, *e.g.*, typically less than about five minutes, preferably less than about ninety seconds, more preferably in less than about thirty seconds and most preferably in less than about ten or fifteen seconds. Such formulations are suitable for administration to a subject via a variety of routes, for example by insertion into a body cavity or application to a moist body surface or open wound.

[0257] Typically, a "flash dosage" is a solid dosage form that is administered orally, which rapidly disperses in the mouth, and hence does not require great effort in swallowing and allows the compound to be rapidly ingested or absorbed through the oral mucosal membranes. In some embodiments, suitable rapidly dispersing dosage forms are also used in other applications, including the treatment of wounds and other bodily insults and diseased states in which release of the medicament by externally-supplied moisture is not possible.

[0258] "Flash dose" forms are known in the art; see for example, effervescent dosage forms and quick release coatings of insoluble microparticles in U.S. Pat. Nos. 5,578,322 and 5,607,697; freeze dried foams and liquids in U.S. Pat. Nos. 4,642,903 and 5,631,023; melt spinning of dosage forms in U.S. Pat. Nos. 4,855,326, 5,380,473 and 5,518,730; solid, free-form fabrication in U.S. Pat. No. 6,471,992; saccaride-based carrier matrix and a liquid binder in U.S. Pat. Nos. 5,587,172, 5,616,344, 6,277,406, and 5,622,719; and other forms known to the art.

[0259] The compounds of the invention are also formulated as "pulsed release" formulations, in which the compound is released from the pharmaceutical compositions in a series of releases (*i.e.*, pulses). The compounds are also formulated as "sustained release" formulations in which the compound is continuously released from the pharmaceutical composition over a prolonged period.


[0260] Also contemplated are formulations, *e.g.*, liquid formulations, including cyclic or acyclic encapsulating or solvating agents, *e.g.*, cyclodextrins, polyethers, or polysaccharides (*e.g.*, methylcellulose), or more preferably, polyanionic β -cyclodextrin derivatives with a sodium sulfonate salt group separate from the lipophilic cavity by an alkyl ether spacer group or polysaccharides. In a preferred embodiment, the agent is methylcellulose. In another preferred embodiment, the agent is a polyanionic β -cyclodextrin derivative with a sodium sulfonate salt separated from the lipophilic cavity by a butyl ether spacer group, *e.g.*, CAPTISOL® (CyDex, Overland, KS). One skilled in the art can evaluate suitable agent/disclosed compound formulation ratios by preparing a solution of the agent in water, *e.g.*, a 40% by weight solution; preparing serial dilutions, *e.g.* to make solutions of 20%, 10, 5%, 2.5%, 0% (control), and the like; adding an excess (compared to the amount that can be solubilized by the agent) of the disclosed compound; mixing under appropriate conditions, *e.g.*, heating, agitation, sonication, and the like; centrifuging or filtering the resulting mixtures to obtain clear solutions; and analyzing the solutions for concentration of the disclosed compound.

[0261] All publications and patent documents cited herein are incorporated herein by reference as if each such publication or document was specifically and individually indicated to be incorporated herein by reference. Citation of publications and patent documents is not intended as an admission that any is pertinent prior art, nor does it constitute any admission as to the contents or date of the same. The invention having now been described by way of written description, those of skill in the art will recognize that the invention can be practiced in a variety of embodiments and that the foregoing description and examples below are for purposes of illustration and not limitation of the claims that follow.


EXAMPLES

Example 1: Syntheses


[0262] Representative syntheses of compounds of the invention are described herein. Synthesis of Compounds 1 and 2 (KX1-136 and KX1-305)

3-benzyloxybenzonitrile

[0263] To a solution of 3-cyanophenol (5.00 g, 42.00 mmol) in acetone (100 ml), potassium carbonate (5.79 g, 42.0 mmol), potassium iodide (335 mg, 21.0 mmol) and benzyl bromide (4.20 ml, 42.00 mmol) were added and the reaction mixture refluxed for 12 hrs (TLC, ethyl acetate:hexane 1:1, $R_f=0.6$), then the solvent removed under vacuum and the residue portioned between water (50 ml) and ethyl acetate (50 ml), the organic layer was washed with water twice and dried over anhydrous sodium sulfate and evaporated under reduced pressure to give the target ether as a yellow oil (8.46g) 96% yield; ^1H NMR (DMSO (dimethylsulfoxide), 400 MHz): δ 7.51-7.33(m, 9H), 5.16(s, 2H).


3-benzyloxybenzylaminehydrochloride

[0264] To a suspension of lithium aluminum hydride, LAH (4.314 g, 113.684 mmol) in dry ether (200 ml) a solution of the 3-benzyloxybenzonitrile in ether (7.92 g, 37.894 mmol) was added drop-wise during 10 min at room temperature, and allowed to stir for 4 hrs (TLC, ethyl acetate: hexane 1:3, $R_f=0.5$), the reaction was quenched with 10 ml ethyl acetate and 10 ml water and filtered. The organic layer was washed with water, dried over Na_2SO_4 and treated with 10 ml conc. HCl to form instant white precipitate (6 g) 68% yield. ^1H NMR (DMSO, 400 MHz): δ 8.33(s, 3H), 7.45-7.37(m, 4H), 7.34-7.30(m, 2H), 7.19(s, 1H), 7.02(t, $J=10\text{Hz}$, 2H), 5.10(s, 2H), 3.97(s, 2H).

N(3-benzyloxy-benzyl)-4-biphenylacetamide

[0265] To a solution of 4-biphenyl acetic acid (2.29 g, 10.45 mmol) in dimethylformamide, DMF, (30 ml) was added diisopropylethylamine, DIPA, (5.47 ml, 31.35 mmol) and stirred at room temperature for 15 min, then benzotriazolyloxy-tris[pyrrolidino]-phosphonium hexafluorophosphate, PyBOP™, (5.43 g, 10.45 mmol) was added and the stirring was continued for further 30 min, then 3-benzyloxybenzylaminehydrochloride (2.6 g, 10.45 mmol) was added and the stirring continued for 24 hrs. The reaction mixture was then poured on to ice cooled water acidified with (10 ml) 1 N HCl and extracted with ethyl acetate (100 ml) and the organic layer washed with saturated solution of NaHCO_3 , water and brine, dried over Na_2SO_4 and the solvent removed under vacuum to give a yellowish-white powder of the desired compound (2.65 g) 62% yield.

[0266] Another procedure involves use amide formation using the acid chloride as shown in the following reaction.




[0267] To 4-biphenylacetic acid (2.5 g) in a flask, thionylchloride (20 ml) was added and heated to reflux for 1 h, cooled, and the excess thionylchloride removed under vacuum to dryness, then the produced crude acid chloride 2.8 g, dissolved in dry DCM (dichloromethane) (30 ml), and added drop wise at 0 °C to equimolar amount of the 3-benzyloxybenzylamine solution in DCM (10 ml) with (1.5 mol) of triethylamine (TEA) and stirred for 5 hrs, then poured onto acidified cold water, the organic layer washed with water, brine and the solvent removed under reduced pressure to give the target amide in 80% yield. ¹H NMR (DMSO, 500 MHz): δ 8.58 (t, J=12Hz 1H), 7.60-7.57 (m, 4H), 7.44-7.29(m, 10H), δ 7.21(t, J=16.5Hz, 2H), 6.85(d, J=6.5Hz, 2H), 6.81(d, J=8.0Hz, 1H), 5.00(s, 2H), 4.24(d, J=6Hz, 2H), 3.51(s, 2H).

15

20

Compound 1: N(3-hydroxy-benzyl)-4-biphenylacetamide




[0268] To remove the benzyl group of this ether (5.00 g, 13.35 mmol) was dissolved in methanol (20 ml), to this solution was added a catalytic amount of 10 % Pd/C (355 mg, 2.21 mmol) in a Parr hydrogenator (55 psi) for 5 hrs, filtered through celite and the solvent removed under vacuum to give the target phenol as yellowish powder (3.20 g) 84% yield, which crystallized from methanol to give (1.5 g) of white crystalline material, mp =169-170 °C. ¹HNMR (DMSO, 400 MHz): 8.9.34(s, 1H), 8.53(s, 1H), 7.63(d, J=8Hz, 2H), 7.58(d, J=8.4Hz, 2H), 7.44(t, J=7.6Hz, 2H), 7.35(d, J=8Hz, 3H), 7.07(t, J=8Hz, 1H), 6.65-6.60(m, 3H), 4.17(d, J=5.6Hz, 2H), 3.5(s, 2H). FAB (fast atom bombardment) HRMS m/e calcd. For (M+H) C₂₁H₂₀NO₂: 318.1449; found: 318.1484.

30

35

40


Compound 2: N(3-fluoro-benzyl)-4-biphenylacetamide



[0269] To a solution of 4-biphenyl acetic acid (2.00 g, 9.42 mmol) in DMF (20 ml) was added DIEA (3.29 ml, 18.84 mmol) and stirred at room temperature for 15 min, then PyBOP (4.90 g, 9.42 mmol) added and the stirring continued for further 30 min, then 3-fluorobenzylamine (1.18 g, 9.42 mmol) added and the stirring continued for 24 hrs, then the reaction mixture poured on to ice cooled water acidified with (10 ml) 1 N HCl and extracted with ethyl acetate (100 ml) and the organic layer washed with saturated solution of NaHCO₃, water and brine, dried over Na₂SO₄ and the solvent removed under vacuum to give a white powder of the desired compound (1.00 g) 33% yield. Another method involves the acid chloride coupling method described below.

50

55



[0270] 4-biphenylacetic acid (2.5 g, 11.78 mmol) charged in a flask then thionylchloride (15 ml) was added and heated to reflux for 1 h, cooled, and the excess thionylchloride removed under vacuum to dryness, then the produced crude acid chloride (2.8 g, 12.13 mmol) dissolved in dry DCM (30 ml), and added drop wise at 0 °C to (1.38 ml, 12.13 mmol) of the 3-fluorobenzylamine solution in DCM (10 ml) along with (1.69 ml, 12.13 mmol) of TEA and stirred for 5 hrs, then poured onto acidified cold water, the organic layer washed with water, brine and the solvent removed under reduced pressure to give the target amide (3.1 g) 80% yield. Recrystallized from methanol, mp = 170-172 °C. ¹H NMR (DMSO, 500 MHz): 8.62(t, *J*=11Hz, 1H), 7.63(d, *J*= 8Hz, 2H), 7.59(d, *J*=8.5Hz, 2H), 7.44(t, *J*=7.5Hz, 2H), 7.37-7.31(m, 4H), 7.08-7.01(m, 3H), 4.28(d, *J*=5.5Hz, 2H), 3.52(s, 2H). FAB HRMS *m/e* calcd. For (M+H) C₂₁H₁₈FNO: 320.1406; found: 320.2, and the base peak found: 342.1262 for (M+Na); calcd. 342.1372.


Synthesis of Compound 3, KX1-306

[0271] The synthesis, outlined in Scheme 1, began with acid chloride formation of biphenylacetic acid followed by amide coupling with 3,5-dibenzoyloxybenzylamine. A large number of impurities were introduced by acid chloride formation. However, other amide coupling procedures such as, for example, PyBOP or carbodiimides, can also be used in this reaction.

[0272] Cleavage of one of the benzyl groups was accomplished under high pressure hydrogen (50-60 psi) for 15 hours. The reaction was monitored by TLC. Silica gel chromatography was used to separate the product from the starting material as well as the dihydroxy side-product

[0273] Biphenyl acetic acid (220 mg, 1.00 mmol) was dissolved in DCM, 5 eq (0.38 mL) of thionyl chloride were added and the reaction was refluxed for 4 hours. Solvents were removed *in vacuo* and the residue was dissolved in DCM. 3,5-Dibenzoyloxybenzylamine (1.1 eq) was added followed by TEA (1 eq). The reaction was then stirred at room temperature overnight. The reaction was diluted to 45 mL (with DCM) and washed with 1 N HCl (3X20 L), saturated sodium bicarbonate (3X20 mL), and brine (3X20 mL). The Reaction was then dried with sodium sulfate and removed *in vacuo* to give 330 mg of crude product Silica gel chromatography (1:1 DCM:EtOAc (ethyl acetate)) gave 220 mg pure product TLC Ref=0.2 (single spot, 7:3 hexanes:EtOAc). LCMS 514.2 (m+H) 536.2(m+Na). ¹H NMR (300 MHz, CDCl₃) δ (ppm) 3.65(s, 2H), 4.50 (d, 5.7Hz, 2H), 4.96 (s, 4H), 5.71 (s, 1H), 6.43 (s, 2H), 6.49 (s, 1H), 7.58-7.26 (m, 19H).

[0274] The dibenzoyloxyamide (1) was dissolved in 15 ml EtOAc (ethyl acetate) with gentle heating in a Part bottle. This was put on the hydrogenator at 50 psi hydrogen for 15 hr. The reaction was filtered through celite and the solvent was removed *in vacuo* to give a crude mixture of starting material and product Silica gel chromatography gave 50 mg 1 and 41 mg desired product KX1-306; LCMS 424.1(m+H), 446.2(m+Na), 847.0(2m+H), 868.9(2m+Na). ¹H NMR (400 MHz, CDCl₃) δ (ppm) 3.66(s, 2H), 4.38 (d, 5.6Hz, 2H), 4.98 (s, 2H), 5.71 (s, 1H), 6.43 (s, 2H), 6.49 (s, 1H), 7.30-7.45 (m, 10H), 7.54-7.57(m, 4H).

Scheme 1


Reagents: i) SOCl_2 , DCM. ii) 3,5-dibenzoyloxybenzylamine (1.0eq), TEA (2.0eq) 20% yield (two steps, with chromatography). iii) 10% Pd/C (10 mol %), H_2 , 55psi, EtOAc 24hr (53%, after chromatography, BORMS).

Synthesis of Compound 4. KX1-307

[0275] The synthesis is outlined in Scheme 2. In one synthesis, the reaction commenced with amide bond formation to give 2, followed by a Suzuki coupling with phenylboronic acid to give the meta-biphenyl product Compound 4, KX1-307. In the Suzuki reaction, the biphenyl product was formed but the reaction did not go to completion (by NMR and LCMS) despite additional, time, heat, and extra catalyst. Using silica gel chromatography, the product could not be separated from the bromo starting material 2. Reversing the Suzuki and amide coupling solved the separation problem and successfully produced the metabiphenyl amide KX1-307 as well as 2'-Fluorobiphenyl-4-acetamide KX1-309 (compound 6, Scheme 3).

[0276] 3-Bromophenylacetic acid (250 mg, 1.163 mmol) and 156 mg (1.1 eq) of phenylboronic acid were dissolved in 6 mL water:isopropanol (6:1). Sodium carbonate (160 mg, 1.3 eq) was dissolved in 0.5 mL distilled water and added to the reaction followed by $\text{Pd}(\text{OH})_2/\text{C}$ (74 mg, 3 mol%). This was rotated in a 65°C water bath for 5 hours. The reaction was filtered through filter paper. Filter paper was washed with 25 mL isopropanol:water:1 N NaOH (35:5:1). Washes were combined and acidified to pH 2 with 1 N sulfuric acid. Isopropanol was removed *in vacuo* and water (10 mL) was added. This aqueous layer was washed with dichloromethane (3X20 mL). Organic washes were combined, dried with sodium sulfate, and removed *in vacuo* to give 215 mg (87% yield) of the biphenyl product 3. TLC Rf=0.7(long streak, 1:1 EtOAc:DCM). ^1H NMR (300 MHz, CDCl_3) δ (ppm) 3.72 (s, 2H), 7.26-7.60 (m, 9H).

[0277] 3-Biphenylacetic acid (3) (100 mg, 0.472 mmol), 3-Fluorobenzylamine (1.1 eq), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, EDCI, (1.1 eq), and HOBT (1-hydroxybenzotriazole, 1.0 eq) were all dissolved in 10 mL anhydrous DCM. After 10 min DIEA (1.1 eq) was added and the reaction was allowed to go overnight. The reaction was diluted to 25 mL and washed with 1N HCl (3X10 mL), saturated sodium bicarbonate (3X10 mL), and brine (2X20 mL). The reaction was dried with sodium sulfate and removed *in vacuo* to give 124 mg pure KX1-307 (83% yield). TLC Rf=0.7 (single spot, 1:1 EtOAc:DCM). ^1H NMR (300 MHz, CDCl_3) δ (ppm) 3.69 (s, 2H) 4.40 (d, 6.0Hz) 5.77 (s, 1H) 6.86-6.96 (m, 3H) 7.10-7.26 (m, 2H) 7.32 (m, 8H).


Scheme 2

Reagents: i) SOCl_2 , DCM. ii) 3-Fluorobenzylamine (1.1 eq), DIEA (2.2 eq) (20% after chromatography). iii) Phenylboronic acid (1.2 eq), 2M sodium carbonate, $\text{Pd}(\text{PPh}_3)_4$ (3 mol %), Toluene (inseparable mixture). iv) Phenylboronic acid (1.1 eq), Na_2CO_3 (1.3 eq), $\text{Pd}(\text{OH})_2/\text{C}$ (3 mol %), 1:6 isopropanol:water (87% yield). v) 3-Fluorobenzylamine (1.1 eq), EDCI (1.1 eq), HOBT (1.0 eq), DIEA (1.1 eq) (83% yield).

Synthesis of Compound 6, KX-309

[0278] The synthesis is outlined in Scheme 3. 4-Bromophenylacetic acid (500 mg, 2.33 mmol) and 358 mg of 2-fluorophenylboronic acid (1.1 eq) were dissolved in 12 mL, 6:1 water:isopropanol. Sodium carbonate (320 mg, 1.3 eq) was dissolved in 1 mL distilled water and added to the reaction followed by $\text{Pd}(\text{OH})_2/\text{C}$ (148 mg, 3 mol %). This was rotated in a 65°C water bath for 5 hours. The reaction was filtered through filter paper. Filter paper was washed with 50 mL isopropanol:water:1 N NaOH (35:5:1). Washes were combined and acidified to pH 2 with 1 N sulfuric acid. Isopropanol was removed *in vacuo*, water (20 mL) was added and washed with dichloromethane (3X30 mL). Organic washes were combined, dried with sodium sulfate, and removed *in vacuo* to give 177 mg (35% yield) of the biphenyl product 4. TLC $R_f=0.7$ (long streak, 1:1 EtOAc:DCM). ^1H NMR (500 MHz, CDCl_3) δ (ppm) 3.73 (s, 2H), 7.16 (t, 10.5Hz, 1H), 7.22 (t, 7.5Hz, 1H), 7.32 (qd, 1.5Hz, 7.5Hz, 1H), 7.38 (d, 8.0Hz, 2H), 7.44 (td, 1.5Hz, 7.5Hz, 1H), 7.54 (d, 8.0Hz, 2H).

[0279] 2'-Fluorobiphenylacetic acid (4) (103 mg, 0.448 mmol), 3-Fluorobenzylamine (1.1 eq), EDCI (1.1 eq), and HOBT (1.0 eq) were all dissolved in 6 mL anhydrous DCM. After 10 min DIEA (1.1 eq) was added and the reaction was allowed to go overnight. Reaction was diluted to 25 mL and washed with 1 N HCl (3X10 L), saturated sodium bicarbonate (3X10 mL), and brine (2X20 mL). The reaction was dried with sodium sulfate and removed *in vacuo* to give 126 mg pure Compound 6, KX1-309 (83% yield). LCMS 360.1 (m+Na) 696.8 (2m+Na). ^1H NMR (300 MHz, CDCl_3) δ (ppm) 3.67 (s, 2H) 4.21 (d, 6.0Hz, 2H) 5.79 (s, 1H) 6.87-6.98 (m, 3H) 7.10-7.44 (m, 7H) 7.53 (dd, 1.5Hz, 7.5Hz, 2H).

Scheme 3

Reagents: i) Phenylboronic acid (1.1 eq), Na_2CO_3 (1.3 eq), $\text{Pd}(\text{OH})_2/\text{C}$ (3 mol %), 1:6 isopropanol: water (35% yield). ii) 3-Fluorobenzylamine (1.1 eq), EDCI (1.1 eq), HOBT (1.0 eq), DIEA (1.1 eq), 83% yield.

Synthesis of Compound 5: N-(3-fluorophenyl)-4-biphenylacetamide, KX1-308.

[0280] Thionyl chloride (0.38 mL, 5.0 mmole) was added to an ice water cooled solution of 4-Biphenylacetic acid (0.2

g, 0.9 mmole) in 5 ml dichloromethane, solution allowed to warm to room temperature then heated under reflux for 1 hr, the solvent and excess thionyl chloride was evaporated under vacuum, the oil formed was redissolved in 5 ml dichloromethane followed by addition of 4-Dimethylaminopyridine (0.12 gm, 1.0 mmole) and 3-Fluoroaniline (0.11 gm, 1.0 mmole), stirred at room temperature over night, then the reaction mixture was diluted with 10 ml dichloromethane and 20 ml water, the organic layer washed with 1 N HCl, saturated NaHCO₃ solution, and saturated NaCl solution, dried using Na₂SO₄ and evaporated dryness (0.2 gm, 72%), ¹H-NMR INOVA-500 (CDCl₃) δ 3.805 (s, 2H), 6.815 (t, J=8.5 Hz, 1H), 7.068(d, J=8.0 Hz, 1H), 7.218-7.284(m, 2H), 7.380-7.499(m, 6H) 7.620-7.664(m, 4H). MS (m/z) 306.2 (M+H)⁺.

Synthesis of Compound 7: N-(3-fluorobenzyl)-4-(3-fluorophenyl)phenylacetamide, KX1-310

[0281] Synthesis of (4'-Fluoro-biphenyl-4-yl)-acetic acid: 4-Bromo-phenylacetic acid (0.5 gm, 2.3 mmole), 3-fluorophenylboronic acid (0.36 gm, 2.4 mmole) and 50% water wet 10% Palladium carbon (0.16 gm, 0.075 mmole Pd) were added to 10 ml of 5:1 water isopropanol mixture, then Na₂CO₃ (0.32 gm, 3 mmole) dissolved in 3 ml of water was added to the above mixture, the reaction was heated at 65-70 °C overnight, the reaction was cooled to room temperature, diluted with 20 ml of 70:15:1 i-PrOH/H₂O/10% NaOH, filtered, the catalyst was washed with 20 mlX3 using the above mixture, the filtrate was acidified using 20% H₂SO₄, filtered and dried (3'-Fluoro-biphenyl-4-yl)-acetic acid: (0.4 gm, 75%) ¹H-NMR INOVA-500 (DMSO d₆) δ 3.623 (s, 2H), 7.192 (m, 1H), 7.358(d, J=8.0 Hz, 2H), 7.474-7.515(m, 3H), 7.652(d, J=8.0 Hz, 2H), 12.316(s, 1H).

[0282] 3-fluorobenzylamine (0.14 ml, 1.1 mmole), PyBOP (0.57 gm, 1.1 mmole), and DIEA (0.36 ml, 2.2 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.22 gm, 76%); ¹H-NMR Nova-500 (DMSO d₆) δ 3.550(s, 2H), 4.303(d, J=6.5Hz, 2H), 7.027-7.097(m, 3H), 7.197(m, 1H), 7.350(m, 1H), 7.389(d, J=8.0Hz, 2H), 7.477-7.518(m, 3H), 7.657(d, J=8.0Hz, 2H), 8.652(t, J=5.5Hz, 1H). MS (m/z) 338.1 (M+H)⁺.

Synthesis of Compound 8, N-(3-fluorobenzyl)-4-(4-fluorophenyl)phenylacetamide, KX1-311

[0283] Synthesis of (4'-Fluoro-biphenyl-4-yl)-acetic acid: 4-Bromo-phenylacetic acid (0.5 gm, 2.3 mmole), 4-fluorophenylboronic acid (0.36 gm, 2.4 mmole) and 50% water wet 10% Palladium carbon (0.16 gm, 0.075 mmole Pd) were added to 10 ml of 5:1 water isopropanol mixture, then Na₂CO₃ (0.32 gm, 3 mmole) dissolved in 3 ml of water was added to the above mixture, the reaction was heated at 65-70 °C overnight, the reaction was cooled to room temperature, diluted with 20 ml of 70:15:1 i-PrOH/H₂O/10% NaOH, filtered, the catalyst was washed with 20 mlX3 using the above mixture, the filtrate was acidified using 20% H₂SO₄, filtered and dried (0.4 gm, 75%) ¹H-NMR INOVA-500 (DMSO d₆) δ 3.621(s, 2H), 7.290(t, J=8.5Hz, 2H), 7.351(d, J=7.5Hz, 2H), 7.593(d, J=7.5Hz, 2H), 7.695(t, J=7Hz, 2H), 12.386(s, 1H).

[0284] (4'-Fluoro-biphenyl-4-yl)-acetic acid (0.2 gm, 0.9 mmole), 3-fluorobenzylamine (0.14 ml, 1.1 mmole), PyBOP (0.57 gm, 1.1 mmole), and DIEA (0.36 ml, 2.2 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.26 gm, 90%); ¹H-NMR INOVA-500 (DMSO d₆) δ 3.541(s, 2H), 4.304(d, J=5.5Hz, 2H), 7.027-7.098(m, 3H), 7.273-7.382(m, 5H), 7.582(d, J=8.0, 2H), 7.694(m, 2H), 8.641 (t, J=5.5Hz, 2H) MS (m/z) 338.1 (M+H)⁺.

Synthesis of Compound 9, N-(3-fluorobenzyl)-N-methyl-4-biphenylacetamide, KX1-312

[0285] 4-biphenylacetic acid (0.25 gm, 1.2 mmole), N-methyl-3-fluorobenzylamine (0.16 gm, 1.2 mmole), EDCI (0.23 gm, 1.2 mmole), and DIEA (0.42 ml, 2.4 mmole) was dissolved in 10 ml DCM and stirred overnight. The reaction mixture was diluted with 10 ml of DCM washed with 10% HCl, saturated NaHCO₃ solution, and saturated NaCl solution, dried using Na₂SO₄ and evaporated to produce viscous clear oil (160 mg, 43%), ¹H-NMR INOVA-500 (DMSO d₆) indicated the presence of a mixture of cis and trans isomers in a ratio of 1:2, running the NMR experiment was run at 50 °C slightly change the value for the chemical shift, but had almost no effect on the ratio. Protons are labeled H_a or H_b to indicate it belongs to one isomer or the other. ¹H-NMR INOVA-500 (DMSO d₆) 2.813(s, 3H_a), 3.000(s, 3 H_b), 3.784(s, 2H_a), 3.841(s, 2H_b), 4.543(s, 2H_b), 4.681(s, 2H_a), 6.931-7.649(m, 13 H_a+13 H_b). MS (m/z) 334.2 (M+H)⁺.

Synthesis of Compound 10, N-(3-fluorobenzyl)-4-phenyl-2-fluorophenylacetamide, KX1-313

[0286] Synthesis of 4-Bromo-2-fluoro-phenylacetamide: 4-Bromo-2-fluorobenzylbromide (5 gm, 18.7 mmole) was dissolved in 30 ml ethanol, to which water solution (10 ml) of KCN (2.43 gm, 37.4 mmole) was added, refluxed overnight, then it was cooled to room temperature, poured into 200 ml of crushed ice, filtered, chromatographed using 1:1 ethyl acetate followed by ethyl acetate (the cyano compound was hydrolyzed on the silica gel to produce the carboxamide), which was evaporated to produce white solid, (1.3 gm, 32%) ¹H NMR INOVA-500 (DMSO d₆) δ 3.436 (s, 2H), 7.005(s, 1H), 7.289(t, J=8.0Hz, 1H), 7.361(d, J=8.0Hz, 1H), 7.478(m, 1H), 7.517(s, 1H).

[0287] Synthesis of 4-Bromo-2-fluoro-phenylacetic acid: 4-Bromo-2-fluoro-phenylacetamide (1.3 gm) was suspended in 100 ml of 30% NaOH, heating at reflux temperature for 24 hrs, cooled to room temperature, washed with DCM and ethyl acetate. The aqueous layer was acidified with conc. HCl, extracted with ethyl acetate, evaporated; the residue was crystallized from isopropanol-water to give needle crystals (0.5 gm, 38%) $^1\text{H-NMR}$ INOVA-500 (DMSO d_6) δ 3.619

(s, 2H), 7.316(t, J=8.0Hz, 1H), 7.379(dd, J=8.0, 1.5Hz, 1H), 7.516(dd, J=8.0, 1.5Hz, 1H), 12.555(s, 1H).
[0288] Synthesis of 4-phenyl-2-fluorophenylacetic acid: 4-Bromo-2-fluoro-phenylacetic acid (0.25 gm, 1.1 mmole), phenylboronic acid (0.15 gm, 1.2 mmole) and 50% water wet 10% Palladium carbon (0.07 gm, 0.033 mmole Pd) were added to 10 ml of 5:1 water isopropanol mixture, then Na_2CO_3 (0.14 gm, 1.3 mmole) dissolved in 3 ml of water was added to the above mixture, the reaction was heated at 65-70 °C overnight, the reaction was cooled to room temperature, diluted with 20 ml of 70:15:1 i-PrOH/ H_2O /10% NaOH, filtered, the catalyst was washed with 20 mlX3 using the above mixture, the filtrate was acidified using 20% H_2SO_4 , filtered and dried (0.2 gm, 83%) $^1\text{H-NMR}$ INOVA-500 (DMSO d_6) δ 3.675(s, 2H), 7.382-7.518(m, 6H), 7.707(d, J=7.5Hz, 2H), 12.498(s, 1H).

[0289] Synthesis of N-(3-fluorobenzyl)-4-phenyl-2-fluorophenylacetamide: 4-phenyl-2-fluorophenylacetic acid (0.2 gm, 0.9 mmole), 3-fluorobenzylamine (0.14 ml, 1.1 mmole), PyBOP (0.57 gm, 1.1 mmole), and DIEA (0.36 ml, 2.2 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.20 gm, 70%); $^1\text{H-NMR}$ INOVA-500 (DMSO d_6) δ 3.612(s, 2H), 4.318(d, J=6Hz, 2H), 7.064-7.117(m, 3H), 7.345-7.503(m, 7H), 7.695(d, J=7.5Hz, 2H), 8.660(t, J=6Hz, 1H). MS (m/z) 338.1 ($\text{M}+\text{H}$)⁺.

Synthesis of Compound 11. N (3-fluorobenzyl)-2-phenylpyridine-5-acetamide, KX1-314

[0290] Synthesis of 2-phenylpyridine-5-acetic acid: 2-chloropyridine-5-acetic acid (0.2 gm, 1.21 mmole), phenylboronic acid (0.16 gm, 1.3 mmole) and 50% water wet 10% Palladium carbon (0.08 gm, 0.036 mmole Pd) were added to 10 ml of 5:1 water isopropanol mixture, then Na_2CO_3 (0.15 gm, 1.4 mmole) dissolved in 3 ml of water was added to the above mixture, the reaction was heated at 65-70 °C overnight, the reaction was cooled to room temperature, diluted with 20 ml or 70:15:1 i-PrOH/ H_2O /10%NaOH, filtered, the catalyst was washed with 20 mlX3 using the above mixture, the filtrate was dried under vacuum and crude mixture was used without any purification in the next step.

[0291] Synthesis of N(3-fluorobenzyl)-2-phenylpyridine-5-acetamide: To the crude from the above reaction, 3-fluorobenzylamine (0.15 gm, 1.2 mmole), PyBOP(0.67 gm, 1.3 mmole), and DIEA (0.32 gm, 2.6 mmole) and was stirred in DMF overnight. The reaction mixture was then poured into water; solid was collected by filtration, re-crystallized using water-methanol (0.06 gm, 18% in two steps). $^1\text{H-NMR}$ INOVA-500 (CDCl_3) δ 3.645(s, 2H), 4.438(d, J=5.5Hz, 2H); 5.867(s, 1H), 6.925-7.009(m, 3H), 7.268(m, 1H), 7.408-7.493(m, 3H), 7.735(m, 2H), 7.965-7.982(m, 2H), 8.582(s, 1H). MS (m/z) 321.2 ($\text{M}+\text{H}$)⁺.

Synthesis of Compound 12, N-(3-Fluoro-benzyl)-2-(4-pyridin-2-yl-phenyl)-acetamide, KX1-315

[0292] Synthesis of 4-(2-Pyridinyl)benzylalcohol: 4-(2-Pyridinyl)benzaldehyde (2 gm, 11 mmole), and NaBH_4 (0.42 gm, 11 mmole) were stirred at room temperature for 2 hr, ethanol was evaporated, residue dissolved in ethyl acetate washed with saturated NaHCO_3 solution, and saturated NaCl solution, dried using Na_2SO_4 and evaporated to produce white solid (1.5 gm, 75%).

[0293] Synthesis of (4-Pyridin-2-yl-phenyl)-acetic acid: The crude of 4-(2-Pyridinyl)benzylalcohol was dissolved in 20 ml DCM, cooled using ice/methanol, triethylamine (1.25 ml, 8.9 mmole) was added followed by methanesulfonylchloride (0.7 ml, 8.9 mmole) added drop wise over 5 minutes. The reaction was allowed to stir at room temperature till the TLC indicated consumption of the starting material (3 hrs), after completion of the reaction, the reaction mixture was washed with water, saturated NaHCO_3 solution, and saturated NaCl solution, dried using Na_2SO_4 and evaporated to produce yellow oil, the oil produced was dissolved in 25 ml of 90% ethanol, KCN (1.05 gm, 16.2 mmole) was then added and it was heated under reflux overnight. Ethanol was evaporated; solid was washed with 50 ml water and filtered. The solid was dissolved in 30 ml of conc. HCl, refluxed for 48 hr; charcoal was added refluxed for 1 hr, filtered. The HCl was evaporated, the solid formed was dissolved in 5 ml of water, NaOH 1 N was added drop wise while extracting with ethyl acetate, the ethyl acetate extract was dried with Na_2SO_4 and evaporated to produce white solid (0.6 gm, 35% in 3 steps) $^1\text{H-NMR}$ INOVA-500 (DMSO d_6) δ 3.641(s, 2H), 7.345(t, J=6.0Hz, 1H), 7.381(d, J=8.5Hz, 2H), 7.879 (t, J=8.0Hz, 1H), 7.951 (d, J=8.0Hz, 1H), 8.034(d, J=8.0Hz, 2H), 8.662(d, J=4.0Hz, 1H), 12.390(s, 1H).

[0294] Synthesis of N-(3-Fluoro-benzyl)-2-(4-pyridin-2-yl-phenyl)-acetamide: (4-Pyridin-2-yl-phenyl)-acetic acid (0.2 gm, 0.9 mmole), 3-fluorobenzylamine (0.14 ml, 1.1 mmole), PyBOP (0.57 gm, 1.1 mmole), and DIEA (0.36 ml, 2.2 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.13 gm, 45%); $^1\text{H-NMR}$ INOVA-500 (DMSO d_6) δ 3.563(s, 2H), 4.305(d, J=6.0Hz, 2H), 7.032-7.095(m, 3H), 7.332-7.360(m, 2H), 7.404(d, J=8.0Hz, 2H), 7.874(t, J=7.0Hz, 1H), 7.948(d, J=8.0Hz, 1H), 8.034(d, J=8.0Hz, 2H), 8.659(d, J=4.0Hz, 2H). MS (m/z) 321.2 ($\text{M}+\text{H}$)⁺.

Synthesis of Compounds 13 and 24


[0295] Syntheses of the pyridyl derivatives, Compound 13, KX1-316, and Compound 24, KX1-327, are shown in Scheme 4. The amide was made first with an EDCI coupling to give amide 5. The Suzuki with 3- or 4-pyridylboronic acids was then performed. The basic nature of the pyridine ring was exploited to purify the product from and remaining starting material. The product was pulled into the aqueous phase away from the starting material using 1 N HCl. After several organic washes the aqueous layer was basified and the product extracted with ethyl acetate. This purification procedure worked well and eliminated the need for chromatography.

KX1-316 (Compound 13)

[0296] A flame dried 50 mL round bottom flask with two condensers was charged with argon. Dimethoxyethane, 15 mL and 1 mL 2 M potassium carbonate was heated to 45 °C while argon was bubbled through the solution. After 1 hour the bromo amide (240 mg, 0.7475 mmol) and 3-pyridylboronic acid (92 mg, 1.1 eq) were added. After one hour, Pd(PPh₃)₄ (43 mg, 5 mol %) was added neat. Reaction was heated at 65-75 °C for 48 hours. The solvent was poured into a round bottom flask, the remaining residue was washed with ethyl acetate. Solvents were combined and removed *in vacuo*. The residue was taken up in 20 mL 1 N HCl and washed with ethyl acetate (3X10 mL). The acid layer was then basified with a combination of 2 N NaOH and saturated sodium bicarbonate to pH 8-9. The aqueous layer was then washed with ethyl acetate (3X20 mL). Solvent extracts were combined, dried with sodium sulfate and removed *in vacuo*. Residue was purified on silica gel column (1:1 DCM: EtOAc) to give 90 mg of the desired product (38% yield). TLC, R_f 0.2 (1:1DCM:EtOAc). LCMS 321.3 (m+H) 640.8 (2m+Na) 662.9 (2M+Na). ¹HNMR (500 MHz, DMSO) 3.54 (s, 2H) 4.29 (d, 6.0Hz, 2H) 7.00-7.08 (m, 3H) 7.34 (q, 8.0 Hz, 1H) 7.40 (d, 10.0Hz, 2H) 7.47 (dd, 6.0Hz, 10.0Hz, 1H) 7.66 (d, 10.0Hz, 2H) 8.05 (dt, 2.5Hz, 10.0Hz, 1H) 8.55 (dd, 2.0Hz, 6.0Hz, 1H) 6.40 (t, 7.0Hz, 1H) 8.78 (d, 2.5Hz, 1H).

KX1-327 (Compound 24)

[0297] A flame dried 50 mL round bottom flask with two condensers was charged with argon. Dimethoxyethane, 15 mL and 1 mL 2 M potassium carbonate was heated to 45 °C while argon was bubbled through the solution. After 1 hour the bromo amide (150 mg, 0.4672 mmol) and 4-pyridylboronic acid (57 mg, 1 eq) were added. After one hour Pd(PPh₃)₄ (27 mg, 5 mol %) was added neat. Reaction was heated at 65-75 °C for 72 hours. The solvent was poured into a round bottom flask, the remaining residue was washed with ethyl acetate. Solvents were combined and removed *in vacuo*. The residue was taken up in 20 mL 1 N HCl and washed with ethyl acetate (3X10 mL). The acid layer was then basified with a combination of 2 N NaOH and saturated sodium bicarbonate to pH 8-9. The aqueous layer was then washed with ethyl acetate (3X20 mL). Solvent extracts were combined, dried with sodium sulfate and removed *in vacuo* to give 71 mg of the desired product (48% yield). TLC, R_f 0.2 (1:1DCM:EtOAc). LCMS 321.3 (m+H). ¹HNMR (500 MHz, DMSO) 3.56 (s, 2H) 4.29 (d, 6.0Hz, 2H) 7.04 (m, 3H) 7.34 (q, 6.5Hz, 1H) 7.42 (d, 8.0Hz, 2H) 7.69 (d, 6.0Hz, 2H) 7.75 (d, 8.5Hz, 2H) 8.61(d, 6.0Hz, 2H) 8.64 (t, 5.5Hz, 1H).

Scheme 4

Reagents: i) 3-Fluorobenzylamine (1.1eq), EDCI (1.1eq), HOBT(1.0eq), DIEA (1.1eq), 88% yield.
 ii) 3(or 4)-Pyridylboronic acid(1.1eq), Na₂CO₃(1.3eq), Pd(PPh₃)₄ (5 mol %), Dimethoxyethane, 2M Na₂CO₃ (2eq).
 KX1-316 (X₁=N, X₂=C) 38%, KX1-327 (X₁=C, X₂=N) 47%.

Synthesis of Compound 14, 2-[6-(3-Chloro-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide, KX1-317

[0298] Synthesis of 2-(3-Chloro-phenyl)-pyridine-5-carboxylic acid: 2-chloropyridine-5-carboxylic acid (0.2 gm, 1.21 mmole), 3-chlorophenylboronic acid (0.2 gm, 1.3 mmole) and 50% water wet 10% Palladium carbon (0.08 gm, 0.036 mmole Pd) were added to 10 ml of 5:1 water isopropanol mixture, then Na₂CO₃ (0.15 gm, 1.4 mmole) dissolved in 3 ml of water was added to the above mixture, the reaction was heated at 65-70 °C overnight, the reaction was cooled to room temperature, diluted with 20 ml of 70:15:1 i-PrOH/H₂O/10%NaOH, filtered, the catalyst was washed with 20 ml X3 using the above mixture, the filtrate was dried under vacuum and crude mixture was used without any purification in the next step.

[0299] Synthesis of 2-[6-(3-Chloro-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide: To the crude from the above reaction, 3-fluorobenzylamine (0.15 gm, 1.2 mmole), PyBOP (0.67 gm, 1.3 mmole), and DIEA (0.32 gm, 2.6 mmole) and was stirred in DMF overnight. The reaction mixture was then poured into water; solid was collected by filtration, re-crystallized using water-methanol (0.02 gm, 6 % in two steps). H¹-NMR INOVA-500 (DMSO *d*₆) δ 3.611(s, 2H), 4.314(d, J=6.0Hz, 2H), 7.048-7.106(m, 3H), 7.364(m, 1H), 7.500-7.545(m, 2H), 7.808(dd, J=8.0, 2.0Hz, 1H), 7.997(d, J=8.0Hz, 1H), 8.046(d, J=8.0Hz, 1H), 8.126(d, J=2.0Hz, 1H), 8.578(s, 1H), 8.699(bs, 1H). MS (*m/z*) 355.2 (M+H)⁺.

Synthesis of Compound 14, 2-[6-(4-Ethyl-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide, KX1-318

[0300] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide: 2-chloropyridine-5-carboxylic acid (0.2 gm, 1.21 mmole), 3-fluorobenzylamine (0.15 ml, 1.2 mmole), PyBOP (0.67 gm, 1.3 mmole), and DIEA (0.43 ml, 2.6 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.3 gm, 85%); H¹-NMR INOVA-500 (CDCl₃) δ 3.562(s, 2H), 4.429(d, J=6.5Hz, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, J=8, 2.5Hz, 1H), 8.280(d, J=2.5Hz, 1H).

[0301] Synthesis of 2-[6-(4-ethyl-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (0.125 gm, 0.5 mmole), 4-ethylbenzeneboronic acid (0.083 gm, 0.55 mmole) was dissolved in dimethoxymethane (DME), Na₂CO₃ (0.11 gm, 1 mmole) in 5 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenylphosphine) (0.029 gm, 0.025 mmole) was added, degassed for additional 15 min, refluxed for 24 hr. The reaction was allowed to cool to room temperature, filtered, solid washed with ethyl acetate; the organic layer was dried, evaporated. The residue was chromatographed using ethyl acetate/ hexane 3:2. The product is white solid (0.08 gm, 47%). H¹-NMR INOVA-500 (DMSO *d*₆) δ 1.228(t, J=7.5 Hz, 3H), 2.669(q, J=7.5Hz, 2H), 3.590(s, 2H), 4.321(d, J=6Hz, 2H), 7.053-7.113(m, 3H), 7.324-7.375(m, 3H), 7.766(dd, J=9.0, 2.0Hz, 1H), 7.887(d, J=8.5Hz, 1H), 7.994(d, J=8.0Hz, 2H), 8.548(s, 1H), 8.696(t, J=5.5Hz, 1H). MS (*m/z*) 349.3 (M+H)⁺.

Synthesis of Compound 16, N-(3-Fluoro-benzyl)-2-(2-fluoro-biphenyl-4-yl)-acetamide, KX1-319

[0302] Synthesis of 2-Fluoro-biphenyl-4-carbaldehyde: 4-Bromo-2-fluoro-biphenyl (2 gm, 8 mmole) was dissolved in 20 ml of anhydrous tetrahydrofuran, THF, cooled to -78 °C under argon (Ar), n-Butyl lithium 2.5 M (3.5 ml, 8.8 mmole) was added drop wise over 10 min, and was stirred for additional 1 hr, DMF anhydrous (0.68 ml, 8.8 mmole) was then added, stirred for additional 1 hr, then warmed to room temperature over 4 hr, It was then quenched with water, extracted with ether, ether was dried, evaporated, the produced compound was purified using 9:1 hexane/ethyl acetate, to produce white solid (1 gm, 62.5%); H¹-NMR INOVA-500 (CDCl₃) δ 7.416-7.495(m, 3H), 7.581-7.661(m, 4H), 7.723(d, J=8.0Hz, 1H), 9.991(s, 1H).

[0303] Synthesis of (2-Fluoro-biphenyl-4-yl)-methanol: 2-Fluoro-biphenyl-4-carbaldehyde (1 gm, 5 mmole), NaBH₄ were dissolved in ethanol stirred for 2 hrs, NaOH 10% was added, ethanol was evaporated, the reaction mixture was extracted with ethyl acetate, the ethyl acetate extract was dried with Na₂SO₄ and evaporated to produce white solid (0.8 gm, 80%). H¹-NMR INOVA-500 (CDCl₃) δ 2.266(s, 1H), 4.683(s, 2H), 7.142-7.168(m, 2H), 7.339-7.442(m, 4H), 7.519.-7.535(m, 2H).

[0304] Synthesis of (2-Fluoro-biphenyl-4-yl)-acetic acid: (2-Fluoro-biphenyl-4-yl)-methanol (0.75 gm, 3.7 mmole) was dissolved in 20 ml DCM, cooled using ice/methanol, triethylamine (0.55 ml, 4.0 mmole) was added followed by methanesulfonylchloride (0.3 ml, 4.0 mmole) added drop wise over 5 minutes. The reaction was allowed to stir at room temperature till the TLC indicated consumption of the starting material (2 hrs), after completion of the reaction, the reaction mixture was washed with water, saturated NaHCO₃ solution, and saturated NaCl solution, dried using Na₂SO₄ and evaporated to produce yellow oil, the oil produced was dissolved in 25 ml of 70% ethanol, KCN (0.4 gm, 6 mmole) was then added and it was heated under reflux overnight. Ethanol was evaporated; solid was washed with 50 ml water and filtered. The solid was dissolved in 20 ml of ethanol, then 20 ml of conc. H₂SO₄ was added, and was refluxed overnight; the solution was allowed to cool to room temperature, poured to 200 ml of crushed ice, the solid was collected by vacuum filtration, suspended in 25 ml of NaOH 30%, heated at reflux temperature for 24 hrs, cooled to room temperature, washed with DCM and ethyl acetate. The aqueous layer was acidified with conc. HCl, extracted with ethyl

acetate, evaporated; the residue was crystallized from isopropanol-water to give white solid (0.15 gm, 18% in 3 steps) $^1\text{H-NMR}$ INOVA-500 ($\text{DMSO } d_6$) δ 3.672(s, 2H), 7.191-7.254(m, 2H), 7.389-7.560(m, 6H), 12.494(s, 1H).

[0305] Synthesis of N-(3-Fluoro-benzyl)-2-(2-fluoro-biphenyl-4-yl)-acetamide: (2-Fluoro-biphenyl-4-yl)-acetic acid (0.12 gm, 0.5 mmole), 3-fluorobenzylamine (0.08 ml, 0.6 mmole), PyBOP (0.34 gm, 0.6 mmole), and DIEA (0.22 ml, 1.3 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.140 gm, 83%); $^1\text{H-NMR}$ INOVA-500 ($\text{DMSO } d_6$) δ 3.580(s, 2H), 4.316(d, $J=5.5\text{Hz}$, 2H), 7.037-7.110(m, 3H), 7.210-7.247(m, 2H), 7.343-7.372(m, 2H), 7.457-7.501(m, 3H), 7.544(d, $J=8.0\text{Hz}$, 2H), 8.660(t, $J=6.0\text{Hz}$, 1H). MS (m/z) 338.1 ($\text{M}+\text{H}$) $^+$.

10 Synthesis of Compound 17, N-(3-Fluoro-benzyl)-2-[6-(4-fluoro-phenyl)-pyridin-3-yl]-acetamide, KX1-320

[0306] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide: 2-chloropyridine-5-acetic acid (0.2 gm, 1.21 mmole), 3-fluorobenzylamine (0.15 ml, 1.2 mole), PyBOP (0.67 gm, 1.3 mole), and DIEA (0.43 ml, 2.6 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.3 gm, 85%); $^1\text{H-NMR}$ INOVA-500 (CDCl_3) δ 3.562(s, 2H), 4.429(d, $J=6.5\text{Hz}$, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, $J=8$, 2.5Hz, 1H), 8.280(d, $J=2.5\text{Hz}$, 1H).

[0307] Synthesis of N-(3-Fluoro-benzyl)-2-[6-(4-fluoro-phenyl)-pyridin-3-yl]-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (0.093 gm, 0.33 mmole), 4-fluorobenzeneboronic acid (0.052 gm, 0.37 mmole) was dissolved in DME, Na_2CO_3 (0.07 gm, 0.66 mmole) in 5 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenyl)phosphine (0.016 gm, 0.015 mmole) was added, degassed for additional 15 min, refluxed for 24 hr. The reaction was allowed to cool to room temperature, filtered, solid washed with ethyl acetate; the organic layer was dried, evaporated. The residue was chromatographed using ethyl acetate/ hexane 3:2. then it crystallized from methanol-water to produce white solid (0.013 gm, 12%). $^1\text{H-NMR}$ INOVA-500 ($\text{DMSO } d_6$) δ 3.587(s, 2H), 4.306(d, $J=5.0\text{Hz}$, 2H), 7.041-7.099(m, 3H), 7.295-7.363(m, 3H), 7.777(d, $J=7.5$, 1H), 7.913(d, $J=8.0\text{Hz}$, 1H), 8.119(s, 2H), 8.546(s, 1H), 8.702(s, 1H). MS (m/z) 339.2 ($\text{M}+\text{H}$) $^+$.

Synthesis of Compound 18, N-(3-Fluoro-benzyl)-2-[6-(3-fluoro-phenyl)-pyridin-3-yl]-acetamide, KX1-321

[0308] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide: 2-chloropyridine-5-acetic acid (0.2 gm, 1.21 mmole), 3-fluorobenzylamine (0.15 ml, 1.2 mmole), PyBOP (0.67 gm, 1.3 mole), and DIEA (0.43 ml, 2.6 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.3 gm, 85%); $^1\text{H-NMR}$ INOVA-500 (CDCl_3) δ 3.562 (s, 2H), 4.429(d, $J=6.5\text{Hz}$, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, $J=8$, 2.5Hz, 1H), 8.280(d, $J=2.5\text{Hz}$, 1H).

[0309] Synthesis of N-(3-Fluoro-benzyl)-2-[6-(3-fluoro-phenyl)-pyridin-3-yl]-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (0.125 gm, 0.5 mmole), 3-fluorobenzeneboronic acid (0.08 gm, 0.55 mmole) was dissolved in DME, Na_2CO_3 (0.11 gm, 1.0 mmole) in 5 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenyl)phosphine (0.029 gm, 0.025 mmole) was added, degassed for additional 15 min, refluxed for 24 hr. The reaction was allowed to cool to room temperature, filtered, solid washed with ethyl acetate; the organic layer was dried, evaporated. The residue was chromatographed using ethyl acetate/ hexane 3:2, then it crystallized from methanol-water to produce white solid (0.075 gm, 45%). $^1\text{H-NMR}$ INOVA-500 ($\text{DMSO } d_6$) δ 3.614(s, 2H), 4.318(d, $J=6.0\text{Hz}$, 2H), 7.053-7.099(m, 3H), 7.273 (t, $J=9.0\text{Hz}$, 1H), 7.367(q, $J=7.0\text{Hz}$, 1H), 7.542(q, $J=7.0\text{Hz}$, 1H), 7.812(d, $J=8.0\text{Hz}$, 1H), 7.891 (d, $J=10.0\text{Hz}$, 1H), 7.942 (d, $J=7.5\text{Hz}$, 1H), 7.992(d, $J=8.0\text{Hz}$, 1H), 8.583(s, 1H), 8.717(s, 1H). MS 339.2 ($\text{M}+\text{H}$) $^+$.

45 Synthesis of Compound 19, 2-[6-(3-Ethoxy-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide, KX1-322

[0310] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide: 2-chloropyridine-5-acetic acid (0.2 gm, 1.21 mmole), 3-fluorobenzylamine (0.15 ml, 1.2 mmole), PyBOP (0.67 gm, 1.3 mmole), and DIEA (0.43 ml, 2.6 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.3 gm, 85%); $^1\text{H-NMR}$ INOVA-500 (CDCl_3) δ 3.562(s, 2H), 4.429(d, $J=6.5\text{Hz}$, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, $J=8$, 2.5Hz, 1H), 8.280(d, $J=2.5\text{Hz}$, 1H).

[0311] Synthesis of N-(3-Fluoro-benzyl)-2-[6-(3-fluoro-phenyl)-pyridin-3-yl]-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (0.15 gm, 0.54 mmole), 3-ethoxybenzeneboronic acid (0.096 gm, 0.6 mmole) was dissolved in DME, Na_2CO_3 (0.11 gm, 1.08 mmole) in 5 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenyl)phosphine (0.031 gm, 0.027 mmole) was added, degassed for additional 15 min, refluxed for 24 hr. The reaction was allowed to cool to room temperature, filtered, solid washed with ethyl acetate; the organic layer was dried, evaporated.

The residue was chromatographed using ethyl acetate/ hexane 3:2. then it crystallized from methanol-water to produce white solid (0.03 gm, 17%). ¹H-NMR INOVA-500 (DMSO *d*₆) δ 1.366(t, J=7.0Hz, 3H), 3.591(s, 2H), 4.110(q, J=7.0Hz, 2H), 4.312(d, J=5.5Hz, 2H), 6.985(d, J=7.5Hz, 1H), 7.048-7.105(m, 3H), 7.342-7.402(m, 2H), 7.621(m, 2H), 7.770(d, J=7.0Hz, 1H), 7.826(d, J=8.0Hz, 1H), 7.942(d, J=7.5Hz, 1H), 8.550(s, 1H), 8.701 (s, 1H). MS (*m/z*) 365.2 (M+H)⁺.

Synthesis of Compound 20, 4-{5-[(3-Fluoro-benzylcarbamoyl)-methyl]-pyridin-2-yl}-benzoic acid, KX1-323

[0312] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide: 2-chloropyridine-5-acetic acid (0.2 gm, 1.21 mmole), 3-fluorobenzylamine (0.15 ml, 1.2 mmole), PyBOP (0.67 gm, 1.3 mmole), and DIEA (0.43 ml, 2.6 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.3 gm, 85%); ¹H-NMR NOVA-500 (CDCl₃) δ 3.562(s, 2H), 4.429(d, J=6.5Hz, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, J=8.25Hz, 1H), 8.280(d, J=2.5Hz, 1H).

[0313] Synthesis of N-(3-Fluoro-benzyl)-2-[6-(3-fluoro-phenyl)-pyridin-3-yl]-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (0.15 gm, 0.54 mole), 4-carboxybenzeneboronic acid (0.096 gm, 0.6 mmole) was dissolved in DME, Na₂CO₃ (0.11 gm, 1.08 mmole) in 5 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenylphosphine) (0.031 gm, 0.027 mmole) was added, degassed for additional 15 min, refluxed for 24 hr. The reaction was allowed to cool to room temperature, filtered, solid washed with ethyl acetate, NaOH 10%, the aqueous layer was washed several times with ethyl acetate, neutralized by drop wise addition of HCl 1% having ethyl acetate in the medium with shaking after each addition of the HCl, ethyl acetate was evaporated and the solid formed was crystallized from methanol-water to produce a white solid (0.07 gm, 40%). ¹H-NMR INOVA-500 (DMSO *d*₆) δ 3.625(s, 2H), 4.318(d, J=5.5Hz, 2H), 7.053-7.111(m, 3H), 7.376(q, J=7.0Hz, 1H), 7.8341(d, J=8.0, 1H), 8.015-8.063(m, 3H), 8.206(d, J=8.0Hz, 1H), 8.613(s, 1H), 8.724(t, J=5.5, 1H). MS (*m/z*) 365.3 (M+H)⁺.

Synthesis of Compound 21, 2-[6-(2-Ethoxy-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide, KX1-324

[0314] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide: 2-chloropyridine-5-acetic acid (0.2 gm, 1.21 mmole), 3-fluorobenzylamine (0.15 ml, 1.2 mmole), PyBOP (0.67 gm, 1.3 mmole), and DIEA (0.43 ml, 2.6 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.3 gm, 85%); ¹H-NMR INOVA-500 (CDCl₃) δ 3.562(s, 2H), 4.429(d, J=6.5Hz, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, J=8.25Hz, 1H), 8.280(d, J=2.5Hz, 1H).

[0315] Synthesis of 2-[6-(2-Ethoxy-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (0.15 gm, 0.54 mmole), 2-ethoxybenzeneboronic acid (0.096 gm, 0.6 mmole) was dissolved in DME, Na₂CO₃ (0.11 gm, 1.08 mmole) in 5 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenylphosphine) (0.031 gm, 0.027 mmole) was added, degassed for additional 15 min, refluxed for 24 hr. The reaction was allowed to cool to room temperature, filtered, solid washed with ethyl acetate; the organic layer was dried, evaporated. The residue was chromatographed using ethyl acetate/ hexane 2:1, then it crystallized from methanol-water to produce a white solid (0.075 gm, 40%). ¹H-NMR INOVA-500 (DMSO *d*₆) δ 1.339(t, J=7.0Hz, 3H), 3.581(s, 2H), 4.112(q, J=7.0Hz, 2H), 4.322(d, J=5.5Hz, 2H), 7.032-7.135(m, 5H), 7.358-7.387(m, 2H), 7.703(d, J=7.0, 1H), 7.748(d, J=7.0Hz, 1H), 7.871(d, J=7.0Hz, 1H), 8.548(s, 1H), 8.725(s, 1H). MS (*m/z*) 365.2 (M+H)⁺.

Synthesis of Compound 22, 2-[6-(4-Ethoxy-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide, KX1-325

[0316] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide: 2-chloropyridine-5-acetic acid (0.2 gm, 1.21 mmole), 3-fluorobenzylamine (0.15 ml, 1.2 mmole), PyBOP (0.67 gm, 1.3 mmole), and DIEA (0.43 ml, 2.6 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol. (0.3 gm, 85%); ¹H-NMR INOVA-500 (CDCl₃) δ 3.562(s, 2H), 4.429(d, J=6.5Hz, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, J=8.25Hz, 1H), 8.280(d, J=2.5Hz, 1H).

[0317] Synthesis of 2-[6-(4-Ethoxy-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (0.15 gm, 0.54 mmole), 4-ethoxybenzeneboronic acid (0.096 gm, 0.6 mmole) was dissolved in DME, Na₂CO₃ (0.11 gm, 1.08 mmole) in 5 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenylphosphine) (0.031 gm, 0.027 mmole) was added, degassed for additional 15 min, refluxed for 24 hr. The reaction was allowed to cool to room temperature, filtered, solid washed with ethyl acetate; the organic layer was dried, evaporated. The residue was chromatographed using ethyl acetate/ hexane 2:1, then it crystallized from methanol-water to produce a white solid (0.08 gm, 42%). ¹H-NMR INOVA-500 (DMSO *d*₆) δ 1.357(t, J=7.0Hz, 3H), 3.564(s, 2H), 4.090(q, J=7.0Hz, 2H), 4.309(d, J=6.0Hz, 2H), 7.012-7.103(m, 5H), 7.361(q, J=7.0Hz, 1H), 7.726(d, J=8.0Hz, 1H), 7.842(d, J=8.0Hz, 1H),

8.012(d, J=8.5Hz, 2H), 8.503(s, 1H), 8.686(s, 1H). MS (m/z) 365.2 (M+H)⁺.

Scale-up synthesis of Compound 22 HCl, 2-[6-(4-Ethoxy-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide HCl, KX1-325 HCl

[0318] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide HCl: 2-chloropyridine-5-acetic acid (6.0 gm, 34 mmole), 3-fluorobenzylamine (4.5 ml, 34 mmole), PyBOP (18 gm, 36 mmole), and DIEA (12.5 ml, 75 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol (6.3 gm, 70%); ¹H-NMR INOVA-500 (CDCl₃) δ 3.562(s, 2H), 4.429(d, J=6.5Hz, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, J=8, 2.5Hz, 1H), 8.280(d, J=2.5Hz, 1H).

[0319] Synthesis of 2-[6-(4-Ethoxy-phenyl)-pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (4.8 gm, 17.2 mmole), 4-ethoxybenzeneboronic acid (3.14 gm, 18.9 mmole) was suspended in DME (100 ml), Na₂CO₃ (3.6 gm, 34.4 mmole) in 15 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenylphosphine) (0.99 gm, 0.86 mmole) was added, degassed for additional 15 min, refluxed overnight. The reaction was allowed to cool to room temperature, filtered, the solid washed with cold ethyl acetate and saturated NaHCO₃ solution, the solid was then recrystallized from methanol to produce white solid (4.8 gm).

[0320] 4.6 gm of the free amine was dissolved in 50 ml ethanol with gentle heating, then 25 ml of 4 N HCl in ethyl acetate was added, the solution was concentrated to 20 ml, then diluted with 100 ml of cold ethyl acetate, the solid formed was filtered washed with more ethyl acetate (50x2) and dried (4.3 gm, 65 %); ¹H-NMR INOVA-500 (DMSO *d*₆) δ 1.386 (t, J=7.0Hz, 3H), 3.822(s, 2H), 4.179(q, J=7.0Hz, 2H), 4.339(d, J=6.0Hz, 2H), 7.074-7.182(m, 5H), 7.374(m, 1H), 8.106(d, J=8.0Hz, 1H), 8.263(d, J=8.0Hz, 1H), 8.312(s, 2H), 8.718(s, 1H), 8.981(s, 1H). MS (m/z) 365.2 (M+H)⁺.

[0321] Melting Point of the free base: 0.1 gm of the HCl salt was stirred in 10 ml of 20% NaOH for 10 min, filtered; the solid was crystallized from ethanol water, dried in the oven at 100 °C for 2 hrs. Melting point was found to be 173-176 °C.

Synthesis of Compound 23, N-(3-Fluoro-benzyl)-2-[6-(4-methanesulfonyl-phenyl)-pyridin-3-yl]-acetamide. KX1-326

[0322] Synthesis of 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide: 2-chloropyridine-5-acetic acid (0.2 gm, 1.21 mmole), 3-fluorobenzylamine (0.15 ml, 1.2 mmole), PyBOP (0.67 gm, 1.3 mmole), and DIEA (0.43 ml, 2.6 mmole) was dissolved in DMF stirred overnight, the reaction mixture was then poured into water, solid was collected by filtration, re-crystallized using water-methanol (0.3 gm, 85%); ¹H-NMR INOVA-500 (CDCl₃) δ 3.562(s, 2H), 4.429(d, J=6.5Hz, 2H), 5.868(s, 1H), 6.929-7.015(m, 3H), 7.300-7.333(m, 2H), 7.668(dd, J=8, 2.5Hz, 1H), 8.280(d, J=2.5Hz, 1H).

[0323] Synthesis of N-(3-Fluoro-benzyl)-2-[6-(4-methanesulfonyl-phenyl)-pyridin-3-yl]-acetamide: 2-(6-Chloro-pyridin-3-yl)-N-(3-fluoro-benzyl)-acetamide and (0.15 gm, 0.54 mmole), 4-methanesulfonyl benzeneboronic acid (0.12 gm, 0.6 mmole) was dissolved in DME, Na₂CO₃ (0.11 gm, 1.08 mmole) in 5 ml of water was added to the DME solution, the solution was then degassed for 30 min (Ar through the solution and vacuum applied for the first 5 min), Palladiumtetrakis(triphenylphosphine) (0.031 gm, 0.027 mmole) was added, degassed for additional 15 min, refluxed for 24 hr. The reaction was allowed to cool to room temperature, filtered, solid washed with ethyl acetate; the organic layer was dried, evaporated. The residue was chromatographed using ethyl acetate/ hexane 2:1, then it crystallized from methanol-water to produce a white solid (0.02 gm, 10%); ¹H-NMR INOVA-500 (DMSO *d*₆) δ 3.341(s, 3H), 3.635(s, 2H), 4.315(d, J=7.0Hz, 2H), 7.047-7.110(m, 3H), 7.366(q, J=9.0Hz, 1H), 7.857(d, J=8.5Hz, 1H), 8.027-8.081(m, 3H), 8.343(d, J=10.5Hz, 2H), 8.631(s, 1H), 8.731(s, 1H). MS (m/z) 399.2 (M+H)⁺.

Synthesis of Compound 24, KX1-327, and Compound 26, KX1-357

[0324] The syntheses are shown in Scheme 5.

Compound 24, KX1-327 HCl


[0325] A solution of 75 mL 1,2-Dimethoxyethane and 16 mL 2 M sodium carbonate was thoroughly degassed by heating at 50 °C with an argon stream through the solvent. 5.00 g of the 4-bromophenyl acetamide (5,15.6 mmol) and 1.95 grams of 4-pyridylboronic acid (1.00 eq) were added to and degassing continued for 1 hour. Tetrakis(triphenylphosphine) palladium (5 mol %) was added neat and the reaction was refluxed for 24 hours. The reaction was cooled and poured into 300 mL distilled water and filtered to give 5.014 g crude product. This crude product was taken up in 1 L of a 1 to 1 mix of 1 N HCl and ethyl acetate. The organic layer was discarded and the aqueous layer was washed two more times with EtOAc. The aqueous layer was the basified with solid sodium bicarbonate to pH 7.5. This was then extracted 3X 300 mL EtOAc to give about 3.25 g of semi-pure product. Pure crystals of the free base were made by dissolving 200 mg in a minimum amount of ethyl acetate with gentle heating and sonication. Hexanes was added to this

solution until it became cloudy. This was heated until clear. Addition of more hexanes followed by heating was repeated two more times. This clear solution was allowed to stand overnight in a sealed vessel. White crystals formed which were washed with hexanes and dried to give about 50 mg (mp 145-146 °C). The rest of the product was dissolved in ethanol and two equivalents of hydrochloric acid (1.1 M in EtOAc) were added. After 1 hour the ethanol was removed and redissolved in the least amount of ethanol at 40 °C. EtOAc was added until the solution became cloudy. The solution was allowed to stand and the desired product crystallized as pure white crystals. The crystals were filtered off, washed with EtOAc and dried to give 2.4 grams (48% overall yield); LCMS 321.3 (m+H). ¹HNMR (500 MHz, DMSO) 3.61 (s, 2H) 4.29 (d, 7.5Hz, 2H) 7.04 (m, 3H) 7.34 (q, 9.5Hz, 1H) 7.50 (d, 10.5Hz, 2H) 7.95 (d, 10.5Hz, 2H) 8.24 (d, 8.0Hz, 2H) 8.70 (s, 1H) 8.87(d, 8.0Hz, 2H).

Compound 26, KX1-357

[0326] 47.0 mg of KX1-327 were dissolved in 5 mL DCM. Meta-chloroperoxybenzoic acid (35.0 mg, 1.4 eq) were added and the reaction was allowed to stir for 13 hours. The reaction was washed 3X 5 mL saturated sodium bicarbonate, dried with sodium sulfate and concentrated to give 45 mg of a yellow solid. NMR revealed the product contained about 15% impurity, which may have been m-chlorobenzoic acid (or the peroxide). The solid was redissolved in 5 mL DCM and washed 3X 5 mL saturated sodium bicarbonate, dried with sodium sulfate and concentrated to give 26 mg of the desired product as a yellow solid; LCMS 337.2 (M+H), 672.9 (2M+H), 694.8 (2M+Na). ¹HNMR (400 MHz, DMSO) 3.54 (s, 2H), 4.28 (d, 6.0Hz, 2H), 7.00-7.08 (m, 3H), 7.34 (q, 8.0Hz, 1H), 7.40 (d, 8.4Hz, 2H), 7.72 (d, 8.4Hz, 2H), 7.75 (d, 7.2Hz, 2H), 8.24 (d, 8.4Hz, 2H), 8.63 (t, 5.6Hz, 1H).

Scheme 5



[0327] 4-Bromophenylacetic acid (6.00 g, 47.9 mmol) was dissolved in 40 mL of anhydrous dichloromethane under an argon atmosphere and cooled in an ice bath. 3-Fluorobenzylamine (1.00 eq) was added and unintended precipitation of the acetic acid/benzylamine salt occurred. More dichloromethane (20 mL) was added followed by DIEA (2.2 eq), HOBT (1.0 eq), and EDCI (1.1 eq). After about 2 hours the solid broke up, 4 hours after that the reaction was finished by TLC. The reaction was diluted with 200 mL of dichloromethane and 200 mL of 1 N hydrochloric acid. Upon shaking in a separatory funnel an emulsion formed. This emulsion was divided in half and dichloromethane was removed. 500 mL ethyl acetate and another 300 mL 1 N HCl was added to each half. The organic layer was washed 2 more times with 1 N HCl, 3X 300 mL saturated sodium bicarbonate, and 3X200 mL with saturated sodium chloride. Organic layers from each extraction were combined and dried with sodium sulfate, and solvent was removed to give 13.12 g (85% yield) desired product; ¹HNMR (500 MHz, CDCl₃) δ (ppm) 3.58 (s, 2H), 4.45 (d, 6.0Hz, 2H), 5.70(bs, 1H) 6.93 (m, 3H), 7.16 (d, 8.1Hz, 2H), 7.26 (m, 1H) 7.48 (d, 8.1Hz, 2H).

Synthesis of Compound 25. KX1-329.

[0328] As shown in Scheme 6, 5-Hydroxy-2-methylpyridine was converted to the triflate, 6, followed by Suzuki reaction to give the 5-phenyl-2-methylpyridine. The methyl pyridine, 7, was deprotonated with n-butyllithium and added to a solution of ethyl carbonate. Saponification followed by amide coupling with PyBOP gave the desired product.

[0329] 5-Hydroxy-2-methylpyridine (3.00 g, 27.5 mmol) was dissolved in 15 mL anhydrous pyridine and cooled to 0 °C. Triflic anhydride (7.76 g, 1.1 eq) was added drop wise over 3 minutes. Following the addition the reaction was removed from the ice bath and allowed to stir for 6 hr. The volume was then reduced to 8 mL *in vacuo*, diluted with 50 mL distilled water, and then extracted with 75 mL EtOAc. The organic layer was then washed with 1 N HCl (3X50 mL), dried with sodium sulfate, and removed *in vacuo* to give 2.78 g (42%) of an amber oil (6); LCMS 242.1 (m+H). ¹HNMR (400 MHz, CDCl₃) 2.58 (s, 3H) 7.26 (d, 8.4Hz, 1H) 7.52 (dd, 2.8Hz, 8.4Hz, 1H) 8.47 (d, 2.8Hz, 1H).


[0330] A flame dried 50 mL round bottom flask with two condensers was charged with argon. Dimethoxyethane, 25 mL and 6 mL 2 M sodium carbonate was heated to 45 °C while argon was bubbled through the solution. After 1 hour, the pyridyl triflate (6) (1.538 g, 6.382 mmol) and phenylboronic acid (856 mg, 1.1 eq) were added. After one hour Pd (PPh₃)₄ (370 mg, 5 mol %) was added, the reaction was heated at 65-75 °C for 48 hours. The solvent was poured into a round bottom flask, the remaining residue was washed with ethyl acetate. Solvents were combined and removed *in vacuo*. The residue was purified by silica gel chromatography (hexanes:EtOAc) to give 702 mg of the desired product 7 (65% yield); LCMS 170.2(m+H). ¹HNMR (400 MHz, CDCl₃) 3.60 (s, 3H) 7.22 (d, 8.0Hz, 1H) 7.38 (t, 7.2Hz, 1H) 7.46 (t, 7.2Hz, 2H) 7.56 (d, 8.0Hz, 2H) 7.77(dd, 2.4Hz, 8.0Hz, 1H) 8.73 (d, 2.4Hz, 1H).

[0331] 5-Phenyl-2-methylpyridine (7,205 mg, 1.223 mmol) was dissolved in freshly distilled THF in flame dried glassware under argon. Cooled to -78 °C in a dry ice/acetone bath for 20 minutes. N-Butyllithium (0.485 mL, 1.0 eq) was added drop wise over 5 minutes. This solution was added to a 1HF solution of ethyl carbonate (1.5 eq) *via* a cannula. The solution was stirred for 2 hours before being quenched with methanol added drop wise. 1 N sodium hydroxide (1 mL) was added before removing the organic solvents *in vacuo*. The remaining aqueous solution was extracted with ether (3X15 mL). Organic layers were combined and dried with sodium sulfate and removed *in vacuo* to give 208 mg 8 (71% yield) ¹HNMR (500 MHz, CDCl₃) 1.30 (m, 3H) 2.61 (s, 2H) 4.20 (m, 3H) 7.22 (d, 8.0Hz, 1H) 7.38 (t, 7.5Hz, 1H) 7.48 (t, 7.5Hz, 2H) 7.58(m, 2H) 7.78 (dd, 2.5H, 8.0Hz, 1H) 8.73 (d, 2.5Hz, 1H)

[0332] Ethyl ester 8 (208 mg, 0.86 mmol) was dissolved in 5 mL THF. 1 N NaOH (about 1 mL) was added and the reaction was put in a 35 °C water bath overnight. The volume of the reaction was reduced to about 1 mL and then acidified with 1 N HCl to precipitate the desired product. The precipitate was isolated by decanting and drying *in vacuo* to give 54 mg (30% yield) of 9; LCMS 214.1 (m+H) 236.0(m+Na). ¹HNMR (400 MHz, CD₃OD) 3.64 (s, 2H) 7.24-7.28 (m, 4H) 7.25 (t, 8.4Hz, 2H) 7.52 (d, 8.4Hz, 2H) 7.87 (dd, 2.0Hz, 8.0Hz, 1H) 8.53 (d, 2.0Hz, 1H).

[0333] Carboxylic acid 9 (54 mg, 0.232 mmol), 3-Fluorobenzylamine (1.1 eq), and PyBOP (1.1 eq) were dissolved in 3 mL anhydrous DMF. After 10 minutes DIEA (1.1 eq) was added and the reaction was allowed to stir overnight. The DMF was removed *in vacuo* and the residue was taken up with methanol and crystallized from methanol/water to give 44 mg Compound 25, KX1-329 (55%) as clear, needle crystals; TLC, R_f 0.2 (1:1DCM:EtOAc). LCMS 321.2 (m+H), 343.1 (m+Na), 662.9 (2m+Na). ¹HNMR (400 MHz, CDCl₃) 3.82 (s, 2H), 4.46 (d, 8.8Hz, 2H), 6.91(t, 9.2Hz, 2H) 6.99 (d, 7.6Hz, 1H), 7.25 (t, 8.4Hz, 2H), 7.34 (d, 8.0Hz, 2H) 7.40 (tt, 1.2Hz, 7.2Hz, 2H) 7.55 (d, 7.6Hz, 2H) 7.80 (b, 1H) 7.86 (dd, 2.0Hz, 7.6Hz, 1H) 8.73 (d, 2.0Hz, 1H).

Scheme 6



Reagents: i) Tf₂O, pyridine (43%). ii) Phenylboronic acid(1.1eq), Na₂CO₃(1.3eq), Pd(PPh₃)₄ (5 mol %), Dimethoxyethane, 2M Na₂CO₃ (2eq) (65% after chromatography). iii) n-Butyl lithium (1.0eq), diethylcarbonate (1.5eq), anhydrous THF. iv) LiOH, THF 30C (18% after crystallization). v) 3-fluorobenzylamine(1.1eq), PyBOP(1.1eq), DIEA(1.1eq), DMF (55% yield).

Synthesis of Compound 27,2-[6-(4-Ethoxy-phenyl)-1-oxo-pyridin-3-yl]-N-(3-fluorobenzyl)-acetamide, KX1-358

[0334] To an ice cooled solution of 0.2 gm of 2-[6-(4-Ethoxy-phenyl) pyridin-3-yl]-N-(3-fluoro-benzyl)-acetamide in 80 mL DCM, 0.13 gm of m-chloroperbenzoic acid was added as solid. After stirring overnight, the reaction was washed with saturated sodium bicarbonate solution, dried with sodium sulfate, evaporated to dryness under vacuum, then chromatographed (silica gel) using ethyl acetate followed by 10% methanol in ethyl acetate to produce 0.16 gm (78%); ¹H¹-NMR INOVA-400 (DMSO d₆) δ 1.357(t, J=7.0Hz, 3H), 3.564(s, 2H), 4.090(q, J=6.8Hz, 2H), 4.309(d, J=5.60Hz, 2H), 7.012-7.103(m, 5H), 7.245(d, J=8.0Hz, 1H), 7.729(m, 1H), 7.529(d, J=8.0Hz, 1H), 7.800(d, J=8.5Hz, 2H), 8.225(s, 1H),

8.663(t, J=5.6 Hz, 1H). MS (m/z) 380 (M+H)⁺.

[0335] For the following syntheses, unless otherwise noted, reagents and solvents were used as received from commercial suppliers. Proton and carbon nuclear magnetic resonance spectra were obtained on a Bruker AC 300 or a Bruker AV 300 spectrometer at 300 MHz for proton and 75 MHz for carbon. Spectra are given in ppm (δ) and coupling constants, J, are reported in Hertz. Tetramethylsilane was used as an internal standard for proton spectra and the solvent peak was used as the reference peak for carbon spectra. Mass spectra and LC-MS mass data were obtained on a Perkin Elmer Sciex 100 atmospheric pressure ionization (APCI) mass spectrometer. LC-MS analyses were obtained using a Luna C8(2) Column (100 × 4.6 mm, Phenomenex) with UV detection at 254 nm using a standard solvent gradient program (Method B). Thin-layer chromatography (TLC) was performed using Analtech silica gel plates and visualized by ultraviolet (UV) light, iodine, or 20 wt % phosphomolybdic acid in ethanol. HPLC analyses were obtained using a Prevail C18 column (53 x 7 mm, Alltech) with UV detection at 254 nm using a standard solvent gradient program (Method A).

Method A:

[0336]

Time (min)	Flow (mL/min)	%A	%B
0.0	3.0	95.0	5.0
10.0	3.0	0.0	100.0
11.0	3.0	0.0	100.0
A = Water with 0.1 v/v Trifluoroacetic Acid B = Acetonitrile with 0.1 v/v Trifluoroacetic Acid			


Method B:

[0337]

Time (min)	Flow (mL/min)	%A	%B
0.0	2.0	95.0	5.0
4.0	2.0	5.0	95.0
A = Water with 0.02 v/v Trifluoroacetic Acid - B = Acetonitrile with 0.02 v/v Trifluoroacetic Acid			

Synthesis of N-benzyl-2-(5-bromopyridin-2-yl)acetamide:


[0338]



[0339] A flask was charged with 5-(5-bromopyridin-2(1H)-ylidene)-2,2-dimethyl-1,3-dioxane-4,6-dione (1.039 g, 3.46 mmol), benzylamine (0.50 mL, 4.58 mmol), and toluene (20 mL). The reaction was brought to reflux under nitrogen for 18 hours, then cooled and placed in a freezer until cold. The product was collected by filtration and washed with hexanes to yield a mass of bright white crystals (1.018 g, 96%).

Synthesis of 4-(2-(4-(4,4,5,5-tetramethyl[1,3,2]dioxaborolan-2-yl)-phenoxy)ethyl)morpholine:


[0340]



[0341] To a stirring solution of 4-(4,4,5,5-tetramethyl[1,3,2]dioxaborolan-2-yl)-phenol (2.55 g, 11.58 mmol), 2-morpholin-4-ylethanol (1.60 mL, 1.73 g, 13.2 mmol) and triphenyl phosphine (3.64 g, 13.9 mmol) in methylene chloride (60 mL) at 0 °C was added dropwise DIAD (2.82 g, 13.9 mmol). The reaction was allowed to warm to room temperature and stir overnight. After 18 hours, additional portions of triphenyl phosphine (1.51 g, 5.8 mmol), 2-morpholin-4-ylethanol (0.70 mL, 5.8 mmol), and DIAD (1.17 g, 5.8 mmol) were added. After stirring an additional 2 hours at room temperature the reaction was concentrated and the residue purified by flash chromatography (5% to 25% EtOAc in CHCl₃) to provide the product as a white solid (2.855 g, 74%).

Synthesis of Compound 134. KX2-391:

[0342]




[0343] A 10 mL reaction tube with a septum closure and stir bar was charged with *N*-benzyl-2-(5-bromopyridin-2-yl)acetamide (123 mg, 0.403 mmol), 4-(2-(4-(4,4,5,5-tetramethyl[1,3,2]dioxaborolan-2-yl)-phenoxy)ethyl)morpholine (171 mg, 0.513 mmol), and FibreCat 1007¹ (30 mg, 0.015 mmol). Ethanol (3 mL) was added, followed by aqueous potassium carbonate solution (0.60 mL, 1.0 M, 0.60 mmol). The tube was sealed and heated under microwave conditions at 150 °C for 10 minutes. The reaction was cooled and concentrated to remove the majority of the ethanol, and then taken up in 10 mL of ethyl acetate and washed successively with water and saturated sodium chloride solution. The organic layer was dried with MgSO₄, filtered and concentrated to a white solid. This white solid was triturated with ethyl ether to give ALB 30349 as a white solid (137 mg, 79%): mp 135-137 °C.; ¹H NMR (300 MHz, CDCl₃) δ 8.70 (d, 1H, *J*=2.0 Hz), 7.81 (dd, 1H, *J*=2.4 Hz, *J*=8.0 Hz), 7.65 (br s, 1H), 7.49 (d, 2H, *J*=8.8 Hz), 7.37-7.20 (m, 6H), 7.01 (d, 2H, *J*=8.8

¹ Polymer bound di(acetato)dicyclohexylphenylphosphinepalladium(II), manufactured by Johnson Matthey, Inc. and available from Aldrich (catalog # 590231). Hz), 4.49 (d, 2H, *J*=5.8 Hz), 4.16 (t, 2H, *J*=5.7 Hz), 3.82 (s, 2H), 3.78-3.72 (m, 4H), 2.84 (t, 2H, *J*=5.7 Hz), 2.62-2.58 (m, 4H; HPLC (Method B) 98.0% (AUC), *t_R* = 1.834 min.; APCI MS *m/z* 432 [M+H]⁺.

(4-bromo-3-fluorophenyl)(morpholino)methanone:


[0344]



[0345] A 500 mL flask was charged with 4-bromo-3-fluorobenzoic acid (5.00 g, 22.83 mmol), 100 mL DMF, morpholine (2.4 mL, 27.5 mmol), and 4-Ethylmorpholine (8.6 mL, 67.9 mmol). HOBt (4.32 g, 32.0 mmol) was added followed by EDC (5.25 g, 27.4 mmol) and the reaction allowed to stir at room temperature for 18 hours. The reaction was concentrated and the resulting orange syrup taken up in 100 mL EtOAc and 100 mL water. The organic layer was washed with 100 mL 2N HCl, 100 mL saturated sodium bicarbonate, and 100 mL saturated sodium chloride. The organic was then dried with MgSO_4 , filtered, and concentrated to give 6.476 g (98%) of a viscous yellow oil. This material was used without further purification.

4-(4-bromo-3-fluorobenzyl)morpholine:

[0346]




[0347] A 250 mL flask was charged with (4-bromo-3-fluorophenyl)(morpholino)methanone (4.569 g, 15.86 mmol) and dissolved in 16 mL of THF. Diphenylsilane (6.2 mL, 33.4 mmol) was added followed by carbonyltris(triphenylphosphine)rhodium(I)hydride (100 mg, 0.109 mmol) and the reaction stirred at room temperature for 20 hours.

[0348] The reaction was diluted with 200 mL of ether and extracted with 1N HCl (2×150 mL). This resulted in the formation of a white precipitate in the separatory funnel. The acid layer and the resulting white precipitate were washed with ether (2×100 mL), and then basified with solid NaOH pellets (23 g). The aqueous layer was then extracted with ether (3×125 mL), dried over MgSO_4 , filtered, and concentrated to give 1.35 g (31%) of a colorless oil. This material was used without further purification.

4-(3-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl)morpholine:


[0349]



[0350] A 10 mL microwave reaction tube with septum closure was charged with 4-(4-bromo-3-fluorobenzyl)morpholine (405 mg, 1.48 mmol), Bis(pinacolato)diboron (516 mg, 2.03 mmol), $\text{Pd(dppf)Cl}_2 \cdot \text{CH}_2\text{Cl}_2$ (62 mg, 0.076 mmol), potassium acetate (659 mg, 6.72 mmol), and DMF (3.6 mL). The vial was placed under nitrogen by evacuation/backfilling (5 cycles) and stirred at 80°C for 8 hours. The reaction was cooled, diluted with ethyl acetate (25 mL) and filtered. The organics were washed with water (25 mL) and saturated sodium chloride (25 mL). The organic layer was then dried over MgSO_4 and concentrated to a dark oil. The product was purified by silica gel chromatography eluting with 2% MeOH in CHCl_3 to give 310 mg (65%) of an off-white solid.

Synthesis of Compound 136. KX2-393:


[0351]



[0352] A 10 mL microwave reaction tube with septum closure was charged with 4-(3-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl)morpholine (307 mg, 0.96 mmol), 2-(5-bromopyridin-2-yl)-N-(3-fluorobenzyl)acetamide (247 mg, 0.77 mmol), and FibreCat 1007 (60 mg, 0.03 mmol). Ethanol (3 mL) was added followed by aqueous potassium carbonate solution (1.2 mL, 1.0 M, 1.2 mmol). The tube was sealed and heated under microwave conditions at 150 °C for 10 minutes. The reaction was cooled and concentrated to remove the majority of the ethanol, and then taken up in 10 mL of ethyl acetate and washed successively with water and saturated sodium chloride solution. The organic layer was dried with MgSO₄, filtered, and concentrated. The material was purified by column chromatography (silica gel, 100:0 CHCl₃/MeOH to 95:5 CHCl₃/MeOH) to provide ALB 30351 as a white solid (240 mg, 74%): mp 91-92 °C.; ¹H NMR (300 MHz, CDCl₃) δ 8.71 (br s, 1H), 7.86-7.84 (m, 1H), 7.78 (br s, 1H), 7.37 (t, 2H, J=7.5Hz), 7.28-7.21 (m, 3H), 7.02 (dd, 1H, J=0.6Hz, J=7.7Hz), 6.98-6.90 (m, 2H), 4.49 (d, 2H, J=5.9Hz), 3.84 (s, 2H), 3.72-3.75 (m, 4H), 3.52 (s, 2H), 2.47-2.50 (m, 4H); HPLC (Method A) 98.7% (AUC), t_R = 3.866 min.; APCI MS m/z 438 [M+H]⁺.

4-(2-(4-bromo-3-fluorophenoxy)ethyl)morpholine:


[0353]



[0354] A flask was charged with 4-bromo-3-fluorophenol (4.999 g, 26.2 mmol) and triphenylphosphine (10298 g, 39.3 mmol). Methylene chloride (120 mL) was added followed by 2-morpholinoethanol (4 mL, 33.0 mmol) and the solution was stirred on an ice water bath to cool. After 5 minutes, diisopropyl azodicarboxylate (7.6 mL, 39.1 mmol) was added over 6 to 8 minutes. The reaction was left stirring on the cold bath to slowly warm to room temperature overnight. The reaction was concentrated and the residue purified by flash chromatography (25% to 100% EtOAc in hexanes) to provide the product as a colorless oil (2.621 g, 33%).

4-(2-(3-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)ethyl)morpholine:


[0355]



[0356] A 40 mL microwave reaction tube with a septum closure and stir bar was charged with 4-(2-(4-bromo-3-fluorophenoxy)ethyl)morpholine (307 mg, 1.0 mmol), Bis(pinacolato)diboron (318 mg, 1.25 mmol), Pd(dppf)Cl₂·CH₂Cl₂ (68 mg, 83 μmol), and Potassium acetate (316 mg, 3.22 mmol). DME (20 mL) was added and the tube sealed. The tube was evacuated / backfilled w. N₂ (5 cycles) and microwaved at 125 °C for 30 minutes. The reaction was cooled to room temperature, concentrated and the residue purified by column chromatography (silica gel, 2% MeOH in CHCl₃) to provide the product as a colorless oil (356 mg, >99%). The ¹H NMR spectrum shows the product to contain a small amount of pinaco1-like impurity. The material was used as-is.

Synthesis of Compound 133. KX2-392:


[0357]



[0358] A 10 mL microwave reaction tube with septum closure was charged with 4-(2-(3-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxabolan-2-yl)phenoxy)ethyl)morpholine (175 mg, 0.50 mmol), 2-(5-bromopyridin-2-yl)-N-(3-fluorobenzyl) acetamide (121 mg, 0.37 mmol), and FibreCat 1007 (30 mg, 0.03 mmol). Ethanol (3 mL) was added followed by aqueous potassium carbonate solution (0.600 mL, 1.0 M, 0.60 mmol). The tube was sealed and heated under microwave conditions at 150 °C for 10 minutes. The reaction was cooled, filtered, and concentrated to remove the majority of the ethanol. The residue was then taken up in 10 mL of ethyl acetate and washed successively with water and saturated sodium chloride solution. The organic layer was dried with MgSO₄, filtered, and concentrated. The material was purified by column chromatography (silica gel, 100:0 CHCl₃/MeOH to 95:5 CHCl₃/MeOH) to provide ALB 30350 as a white solid (70 mg, 40%); mp 126-127 °C.; ¹H NMR (500 MHz, CDCl₃) δ 8.67 (br s, 1H), 7.77-7.85 (m, 2H), 7.21-7.37 (m, 3H), 7.02 (d, 1H, J=7.7Hz), 6.90-6.97 (m, 2H), 6.82 (dd, 1H, J=2.5Hz, J=8.6Hz), 6.76 (dd, 1H, J=2.4Hz, J=12.4Hz), 4.49 (d, 2H, J=5.9Hz), 4.15 (t, 2H, J=5.7Hz), 3.83 (s, 2H), 3.71-3.78 (m, 4H), 2.83 (t, 2H, J=5.7Hz), 2.56-2.63 (m, 4H); HPLC (Method A) >99% (AUC), t_R = 4.026 min.; APCI MS m/z 468 [M+H]⁺.

1-(2-(4-bromo-3-fluorophenoxy)ethyl)-4-methylpiperazine:


[0359]



[0360] A flask was charged with 4-bromo-3-fluorophenol (5.00 g, 26 mmol) and triphenylphosphine (10.30 g, 39 mmol). Methylene chloride (120 mL) was added followed by 2-(4-methylpiperazin-1-yl)ethanol (4.61 g, 32 mmol) and the solution was stirred on an ice water bath to cool. After 5 minutes, diisopropyl azodicarboxylate (7.6 mL, 39.1 mmol) was added over 6 to 8 minutes. The reaction was left stirring on the cold bath to slowly warm to room temperature overnight. The reaction was concentrated and the residue purified by flask chromatography (25% to 100% EtOAc in hexanes) to provide the product as a colorless oil (2.62 g, 33%).

1-(2-(3-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)ethyl)-4-methylpiperazine:

[0361]




[0362] A 40 mL microwave reaction tube with a septum closure and stir bar was charged with 1-(2-(4-bromo-3-fluorophenoxy)ethyl)-4-methylpiperazine (428 mg, 1.35 mmol), Bis(pinacolato)diboron (375 mg, 1.48 mmol), Pd(dppf) Cl₂-CH₂Cl₂ (63 mg, 77 μmol), and Potassium acetate (410 mg, 4.18 mmol). DME (10 mL) was added and the tube sealed

The tube was evacuated / backfilled w. N₂ (5 cycles) and microwaved at 100 °C for 30 minutes. Additional Pd(dppf) Cl₂-CH₂Cl₂ (63 mg, 77 μmol) was added and the reaction microwaved at 100 °C for 60 minutes. The reaction was cooled to room temperature, concentrated and the residue purified by column chromatography (silica gel, 1% to 2% MeOH in CHCl₃) to provide the product as a dark oil (354 mg, 72%).

Synthesis of Compound 137, KX2-394

[0363]



[0364] A 10 mL microwave reaction tube with septum closure was charged with 1-(2-(3-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)ethyl)-4-methylpiperazine (340 mg, 0.93 mmol), 2-(5-bromopyridin-2-yl)-N-(3-fluorobenzyl)acetamide (201 mg, 0.62 mmol), and FibreCat 1007 (125 mg, 0.06 mmol). Ethanol (3 mL) was added followed by aqueous potassium carbonate solution (1.00 mL, 1.0 M, 1.00 mmol). The tube was sealed and heated under microwave conditions at 150 °C for 10 minutes. The reaction was cooled, filtered, and concentrated to remove the majority of the ethanol. The residue was then taken up in 10 mL of ethyl acetate and washed successively with water and saturated sodium chloride solution. The organic layer was dried with MgSO₄, filtered, and concentrated. The material was purified by column chromatography (silica gel, 98:2 CHCl₃/MeOH to 90:10 CHCl₃/MeOH) to provide ALB 30352-2 as a tan gum (28 mg, 9%): is NMR (300 MHz, CDCl₃) δ 8.66 (br s, 1H), 7.78-7.94 (m, 2H), 7.20-7.40 (m, 3H), 6.88-7.06 (m, 3H), 6.70-6.85 (m, 2H), 4.47 (d, 2H, J=5.9Hz), 4.14 (t, 2H, J=5.7Hz), 3.83 (s, 2H), , 2.85 (t, 2H, J=5.1Hz), 2.41-2.77 (m, 8H), 2.34 (s, 3H); HPLC (Method A) >99% (AUC), t_R = 3.778 min.; APCI MS m/z 481 [M+H]⁺.

Example 2: Cell Growth Inhibition

[0365] The drug concentration required to block net cell growth by 50% relative to a control sample is measured as the GI₅₀. The GI₅₀s for several of the compounds of the invention were assayed as described herein.

[0366] The HT29 cell line is a NCI standard human colon carcinoma cell line. HT-29 cells were obtained from ATCC at passage 125 and were used for inhibition studies between passage 126 -151. HT29 cells were routinely cultured in McCoy's 5A medium supplemented with Fetal Bovine Serum (1.5% v/v) and L-glutamine (2 mM).

[0367] The c-Src 3T3 is a mouse fibroblast NIH 3T3 normal cell line that has been transfected with a point-mutant of human c-Src wherein tyrosine 527 has been converted to a phenylalanine. This mutation results in "constitutively active" c-Src because phosphorylation on tyrosine 527 results in auto-inhibition of Src by having it fold back on its own SH2 domain. With a Phe there, this phosphorylation cant occur and therefore auto-inhibition can't occur. Thus, the always fully active mutant Src then converts the normal mouse fibroblasts into rapidly growing tumor cells. Since the hyperactive Src is the main factor driving growth in these cells (particularly when cultured under low growth serum conditions), compounds active in blocking this growth are thought to work by blocking Src signaling (e.g. as a direct Src kinase inhibitor or as an inhibitor acting somewhere else in the Src signaling cascade). The cells were routinely cultured in DMEM supplemented with Fetal Bovine Serum (2.0% v/v), L-glutamine (2 mM) and Sodium Pyruvate (1 mM).

[0368] In the BrdU Assay for cell growth inhibition, quantitation of cell proliferation was based on the measurement of BrdU incorporation during DNA synthesis. The Cell Proliferation ELISA BrdU assay kit (colorimetric) was obtained from Roche Applied Science and performed as per vendor instructions.

[0369] Growth inhibition was expressed as a GI₅₀ where the GI₅₀ is the sample dose that inhibits 50% of cell growth. The growth inhibition (GI) is determined from the formula $GI = (T_0 - T_n \times 100 / T_0 - CON_n)$ where T₀ is the BrdU growth of untreated cells at time "0", T_n is the BrdU growth of treated cells at day "n" and CON. is the control BrdU growth of control cells at day "n". The GI₅₀ was extrapolated and the data plotted using XL-Fit 4.0 software.

[0370] Actively growing cultures were trypsinized and cells were resuspended in 190 μL of appropriate culture medium supplemented with 1.05% FBS in each well of a 96-well culture plate (1000 HT-29 cells; 2500 c-Src 3T3 cells). For 96 well culture plate experiments, c-Src 3T3 medium was supplemented with 10 mM HEPES buffer. HT-29 cells were seeded in standard tissue culture 96-well plates and c-Src 3T3 cells were seeded in 96-well plates coated with Poly-D-lysine (BIOCOAT™). To increase CO₂ diffusion, c-Src 3T3 96-well plates were incubated with their lids raised by 2 mm using sterile rubber caps.

[0371] Seeded 96 well plates were allowed to attach overnight for 18-24 hours, either at 37 °C and 5% CO₂ for HT-29 or at 37 °C and 10% CO₂ for c-Src 3T3. Approx 18-24 hours after seeding, the initial growth of cells (To) was determined for untreated cells using the BrdU assay. Samples were reconstituted in DMSO at 20 mM and intermediate dilutions made using DMEM containing 10% FBS. The final assay concentrations were 1.5% for FBS and 0.05% for DMSO. Samples were added as 10 µL aliquots in triplicate and plates were incubated as above for ~72 hours. Negative (vehicle) and positive controls.(e.g., AZ (KX-328)) were included. Plates were assayed for BrdU and the data analyzed as above for GI₅₀.

[0372] The results are shown in Table 3. In this table, the data is listed as Growth % of Control, such that a lower number-at an indicated concentration indicates a greater potency of the compound in blocking growth of that tumor cell line. All compounds were initially prepared as 20 mM DMSO stock solutions and then diluted into buffer for the *in vitro* tumor growth assays. NG means no cell growth beyond the control and T means the number of cells in the drug treated wells was less than in the control (*i.e.* net cell loss). NT indicates that the test was not performed. Compound AZ (KX-328) is an ATP-competitive tyrosine kinase inhibitor, as described in Plé et al., J. Med. Chem, 47:871-887 (2004).

[0373] As shown in Table 3, GI₅₀s were obtained for a number of the compounds in other cell lines. These GI₅₀'s were determined using the standard tumor growth inhibition assays, similar to that described in detail for the HT29 cell line above, and the following cell lines: colon tumor cell lines KM12, lung cancer cell line H460 and lung cancer cell line A549 (all are NCI standard tumor cell lines).

Table 3:

IOC-#	CMPD	HT-29 Growth, % of Control Mean, n=3			GI ₅₀	cSrc 3T3 Growth. % of Control Mean, n=3		
		5 µM	500 nM	50 nM		10 µM	1.0 µM	100 nM
KX2-328	AZ	T	10.0	73.0	99 nM (c-Src 3T3), 794 nM (HT29)	T	T	13.0
KX1-136		T	T	83.1	53 nM (c-Src 3T3), 484 nM (HT29) 105 nM (KM12) 280 nM (H460) 330 nM (A549)	T	T	48.3
KX1-305	2	T	T	107.7	349 nM (c-Src 373), 877 nM (HT29), 410nM (KM12) 890 nM (H460) 1.03 µM (A549)	T	T	35.0
KX1-307	4	39.4	93.8	85.9		4.2	45.3	65.7
KX1-308	5	32.3	76.1	87.9		67.1	77.7	94.5
KX1-312	9	33.7	67.6	93.7		12.1	94.5	98.5
KX1-306	3	T	T	124.4		T	T	47.0
KX1-313	10	T	T	802		T	T	91.6
KX1-319	16	T	T	101.2		T	T	88.2

EP 1 836 169 B9

(continued)

5	IOC-#	CMPD	HT-29 Growth, % of Control Mean, n=3			GI ₅₀	cSrc 3T3 Growth. % of Control Mean, n=3		
			5 uM	500 nM	50 nM		10 uM	1.0 uM	100 nM
	KX1-309	6	T	T	29.5		T	T	T
	KX1-310	7	T	T	93.3		T	T	101.8
10	I0C1-311	8	T	T	60.4		T	T	81.3
	KX1-327	24	T	T	31.6	>200 nM (c-Src 3T3), 680 nM (HT29)	T	T	81.3
15	KX1-316	13	T	45.1	77.8	>200 nM (c-Src 3T3)	T	T	88.2
20	KX1-315	12	T	50.3	66.0		T	88.1	89.3
	KX1-314	11	14.4	83.7	53.21		39.3	88.4	93.6
	KX1-317	14	T	64.0	83.5		T	85.6	942
	KX1-318	15	T	932	164.7		T	71.0	91.4
25	KX1-320	17	86.2	1320	111.2		73.1	88.5	90.4
	KX1-321	18	23.7	118.1	127.2		55.8	98.2	95.5
	KX1-322	19	T	872	114.1	3,730 nM (Src 3T3)	T	T	94.6
30	KX1-323	20	60.8	108.9	105.8		93.2	97.3	96.8
	KX1-324	21	NG	95.7	91.0		T	90.0	96.0
35	KX1-325	22	T	T	85.0	207 nM (c-Src 3T3), 215 nM (HT29)	T	54.2	97.8
40	KX1-326	23	43.7	73.2	65.4		55.7	87.3	92.2
	KX1-329	25	T	T	101	269 nM (c-Src 3T3), 338 nM (HT29)	T	T	96.0
45	KX1-357	28	NT	NT	NT		9.0	95.4	101.3
	KX1-358	27	NT	NT	NT		82.7	91.4	92.2
50	KX2-359	28	T	T	T	34 nM (c-Src 3T3), 45 nM (HT29)	T	T	T
	KX2-360	54	T	T	91		T	T	106.0
55	KX2-361	76	T	T	T	11 nM (c-Src 3T3), 10 nM (HT29)	T	T	T

EP 1 836 169 B9

(continued)

IOC-#	CMPD	HT-29 Growth, % of Control Mean, n=3			GI ₅₀	cSrc 3T3 Growth. % of Control Mean, n=3		
		5 uM	500 nM	50 nM		10 uM	1.0 uM	100 nM
KX2-362	78	T	T	86	56 nM (c-Src 3T3), 56 nM (HT29)	T	T	101
KX2-363	79	T	67	92		100	70	92
KX2-364	82	T	80	105		T	81	92
KX2-365	40	T	T	88	133 nM (c-Src 313), 93 nM (HT29)	T	T	88
KX2-366	75	T	54	89		T	183	103
KX2-367	41	T	6	64		T	T	102
KX2-368 slightly insoluble	29	T	70	107		27	101	99
KX2-369	55	T	72	87		T	101	100
KX2-370	77	81	93	112		108	105	104
KX2-371	81	18	33	98		18	72	75
KX2-372	80	T	T	T	58 nM (c-Src 3T3); 67nM (HT-29)	T	T	T
KX2-373	72	T	T	64	96 nM (c-Src 3T3); 639 nM (HT-29)	T	T	97
KX2-374	115	T	57	74		T	84	110
KX2-375	38	T	T	99	206 nM (c-Src 3T3); 354 nM (HT-29)	T	T	T
KX2-376	74	T	93	96	>1,600 nM (c-Src 3T3); >400 nM (HT-29)	T	T	T
KX2-377	38	T	T	T	118 nM (c-Src 3T3); 111 nM (HT-29)	T	T	T
KX2-378	31	T	61	88		48	107	122

EP 1 836 169 B9

(continued)

5	IOC-#	CMPD	HT-29 Growth, % of Control Mean, <i>n</i> =3			GI ₅₀	cSrc 3T3 Growth. % of Control Mean, <i>n</i> =3		
			5 uM	500 nM	50 nM		10 uM	1.0 uM	100 nM
	KX2-379	70	T	88	89		T	104	106
	KX2-380	30	T	50	100		T	119	124
10	KX2-381	33	T	T	58	914 nM (c-Src 3T3); 375 nM (HT-29)	T	T	116
15	KX2-382	68	50	97	80		103	114	117
20	KX2-383	116				327 nM (c-Src 3T3); 248 nM (HT-29)			
25	KX2-384	64				1,430 nM (c-Src 3T3); inactive (KT-29)			
30	KX2-385	83				232 nM (c-Src 3T3)			
35	KX2-386	37				897 nM (c-Src 3T3); inactive (HT-29)			
40	KX2-387	38				inactive (c-Src 3T3); 1.860 nM (HT-29)			
45	KX2-388	66				>1,600 nM (c-Src 3T3); 908 nM (HT-29)			
50	KX2-W9	60				Inactive (c-Src 3T3); Inactive (HT-29)			
55	KX1-329 N-oxide	135				inactive (c-Src 3T3); inactive (HT-29)			

(continued)


		HT-29 Growth, % of Control Mean, <i>n</i> =3			cSrc 3T3 Growth. % of Control Mean, <i>n</i> =3			
IOC-#	CMPD	5 uM	500 nM	50 nM	GI ₅₀	10 uM	1.0 uM	100 nM
KX2-380	114				797 nM (c-Src 3T3); 868 nM (Kr-29)			
KX2-391	133				13 nM (c-Src 3T3); 23 nM (HT-29)			
KX2-392	134				13 nM (c-Src 3T3); 21 nM (HT-29)			
KX2-393	138				24 nM (c-Src 3T3); 52 nM (HT-29)			
KX2-394	137				13 nM (c-Src 3T3); 26 nM (HT-29)			
NG =No growth, total growth inhibition; T=Cyt Cytotoxic Effect on Cells, negative growth; NT= Not tested								

Other Embodiments

[0374] While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims. It will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

Claims

1. A compound according to Formula I

Formula I

or a salt, solvate or hydrate thereof,

wherein: T is absent;
X_y is CZ, CY, N, or N-O;

X_z is CZ;

Y is selected from hydrogen, hydroxyl, halogen, lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl-aryl, and O-benzyl;

X_a is CR_a , or N, or N-O;


X_b is CR_b , N, or N-O;

X_c is CR_c , N, or N-O;

X_d is CR_d , N, or N-O;

X_e is CR_e ;


R_a , R_b , R_c , R_d , R_e , R_4 , R_5 , and R_6 are, independently, hydrogen, hydroxyl, halogen, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl-aryl, O-benzyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-OH, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, COOH, COO-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl, SO_2H , SO_2 -lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl,




where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl;

V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$; and

Z is:



R_7 , R_8 , R_9 , R_{10} , and R_{11} are, independently, hydrogen, hydroxyl, halogen, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, O-lower (C_1 , C_2 , C_3 , C_4 , C_5 , or C_6) alkyl-aryl, O-benzyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-OH, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-O- C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl,




R_1 , R_2 , and R_3 are independently H or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl; and n is 1 or 2; and m is 1 or 2; wherein at least one of X_a , X_b , X_c and X_d is N.

2. The compound of claim 1, wherein X_y is CY.

3. The compound of claim 1, wherein Y is hydrogen.


4. The compound of claim 1, wherein Z is



5. The compound of claim 1, wherein R_b is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy.

6. The compound of claim 1, wherein R_b is hydrogen.

7. The compound of claim 1, wherein R_b is





where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

8. The compound of claim 8, wherein V is a bond.

9. The compound of claim 1, wherein X_a is N and each of X_b , X_c , and X_d is CR.

10. A composition comprising a compound according to claim 1 and at least one pharmaceutically acceptable excipient.

11. The compound of claim 1, wherein the compound is





Compound #
12

13

24

25








(continued)

Compound

Compound #








Compound #

47

48

49

50




(continued)


Compound

Compound #


55




56





57





58















Compound #





81


















(continued)


Compound

Compound #


116




117



118






Compound #

119

120

121

(continued)



Compound #

122

123

124


125

(continued)


Compound

Compound #


126




127




128



129






Compound #

130

131

132

133



5

10

15

20

25

30

35

40







45

50

55









(continued)

Compound #



Compound

Compound #



10

15

20

25

30

35

40

45







50

55


(continued)

Compound

Compound #



12. A compound of claim 1, wherein the compound is a compound according to formula II:




or a salt, solvate, or hydrate thereof, wherein:

15 R_b , R_4 , R_5 , R_8 , and R_{10} are as defined in claim 1.

13. The compound of claim 12, wherein R_8 is hydrogen, F, Cl, Br, or I.

14. The compound of claim 12, wherein R_b is C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy.


15. The compound of claim 12, wherein R_b is



30 where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl, and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

16. The compound of claim 12, wherein R_4 is hydrogen, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, F, Cl, Br, or I.


17. The compound of claim 12, wherein R_4 is



40 where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

18. The compound of claim 12, wherein R_5 is hydrogen, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, F, Cl, Br, or I.

19. The compound of claim 12, wherein R_5 is



55 where W is H, or C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkyl-aryl; and V is a bond, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ or $-OCH_2CH_2CH_2-$.

20. The compound of claim 12, wherein R_{10} is hydrogen, C_1 , C_2 , C_3 , C_4 , C_5 , or C_6 alkoxy, F, Cl, Br, or I.

21. The compound of claim 12, wherein R_{10} is



15










35



















40











45

50


55









1




2




3



4




6




(continued)


7




8




9




10




11



14




15




(continued)


16




17




18




19




20



21



22



(continued)

23

5

10

27

15

108A

20

25

108B

30

109

35

40







111

45

112

50

55



5

10

15

20

25

30










35

40











45














50

55



(continued)





127

25. A compound for use according to claim 24, wherein the cell proliferative disorder selected from cancer, pre-cancer, hyperproliferative disorder, psoriasis, diabetic retinopathy, macular degeneration, epidermic and dermoid cysts, lipoma, adenoma, capillary and cutaneous hemangioma, lymphangioma, nevi lesions, teratoma, nephroma, myofibromatosis, osteoplastic tumors, and dysplasia.


26. A compound for use according to claim 25 wherein the cancer is selected from the group consisting of colon cancer, lung cancer, breast cancer, ovarian cancer, brain cancer, liver cancer, pancreas cancer, prostate cancer, malignant melanoma, non-melanoma skin cancer, childhood leukemia, lymphoma, multiple myeloma, Hodgkin's disease, acute and chronic leukemia, plasma cell neoplasm, lymphoid neoplasm and cancer associated with AIDS.

27. A compound for use according to claim 24 wherein the microbial infection is selected from the group consisting of bacterial, fungal, parasitic, or viral infection.

Patentansprüche

1. Verbindung gemäß der Formel I

Formel I



oder ein Salz, Solvat oder Hydrat davon,

worin: T nicht vorliegt;

X_y CZ, CY, N oder N-O ist;

X_z CZ ist;

Y aus Wasserstoff, Hydroxyl, Halogen, Nieder-(C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -)alkyl, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkoxy, O-Nieder-(C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -)alkyl-aryl und O-Benzyl ausgewählt ist;

X_a CR_a oder N oder N-O ist;


X_b CR_b oder N oder N-O ist;

X_c CR_c , N oder N-O ist;

X_d CR_d , N oder N-O ist;

X_e CR_e ist;




R_a , R_b , R_c , R_d , R_e , R_4 , R_5 und R_6 unabhängig Wasserstoff, Hydroxyl, Halogen, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkoxy, O-Nieder-(C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -)alkyl-aryl, O-Benzyl, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl-OH, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl-O-nieder-(C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -)alkyl, COOH, COO-Nieder-(C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -)alkyl, SO_2H , SO_2 -Nieder-(C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -)alkyl,




worin W H oder C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl-aryl ist, sind;

V eine Bindung, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ oder $-OCH_2CH_2CH_2-$ ist und

Z




R₇, R₈, R₉, R₁₀ und R₁₁ unabhängig Wasserstoff, Hydroxyl, Halogen, C₁-, C₂-, C₃-, C₄-, C₅-oder C₆-Alkyl, C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-Alkoxy, O-Nieder-(C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-)alkyl-aryl, O-Benzyl, C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-Alkyl-OH, C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-Alkyl-O-C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-alkyl,



sind, R₁, R₂ und R₃ unabhängig H oder C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-Alkyl sind und n 1 oder 2 ist und m 1 oder 2 ist, wobei mindestens einer von X_a, X_b, X_c und X_d N ist.

2. Verbindung nach Anspruch 1, worin X_y CY ist.
3. Verbindung nach Anspruch 1, worin Y Wasserstoff ist.
4. Verbindung nach Anspruch 1, worin Z




ist.

5. Verbindung nach Anspruch 1, worin R_b C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkoxy ist.

6. Verbindung nach Anspruch 1, worin R_b Wasserstoff ist.

7. Verbindung nach Anspruch 1, worin R_b



ist, wobei W H oder C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl-aryl ist und V eine Bindung, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ oder $-OCH_2CH_2CH_2-$ ist.





8. Verbindung nach Anspruch 8, worin V eine Bindung ist.

9. Verbindung nach Anspruch 1, worin X_a N ist und jeder von X_b , X_c und X_d CR ist.

10. Zusammensetzung, die eine Verbindung gemäß Anspruch 1 und mindestens einen pharmazeutisch verträglichen Träger umfasst.

11. Verbindung nach Anspruch 1, wobei die Verbindung

Verbindung




Verbindung Nr.
12

13


24

25


Verbindung
(fortgesetzt)
Verbindung Nr.




26




28




29



30



31



5

10

15

20

25

30

35

40

45


50

55

(fortgesetzt)

Verbindung

Verbindung Nr.





36

37


38


39







Verbindung Nr.










Verbindung Nr.








Verbindung Nr.














Verbindung Nr.




Verbindung Nr.





Verbindung Nr.





Verbindung Nr.





Verbindung Nr.

(fortgesetzt)

Verbindung





Verbindung Nr.


120


121


122

123









5

10

15

20

25

30

35

40

45

50







55

(fortgesetzt)

Verbindung

Verbindung Nr.



137





(fortgesetzt)

Verbindung


Verbindung Nr.








Verbindung Nr.






Verbindung Nr.

ist.

12. Verbindung nach Anspruch 1, wobei die Verbindung eine Verbindung gemäß der Formel II:




oder ein Salz, Solvat oder Hydrat davon ist, worin:

R_6 , R_4 , R_5 , R_8 , and R_{10} wie im Anspruch 1 definiert sind.

13. Verbindung nach Anspruch 12, worin R_8 Wasserstoff, F, Cl, Br oder I ist.

14. Verbindung nach Anspruch 12, worin R_6 C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkoxy ist.


15. Verbindung nach Anspruch 12, worin R_6



ist, worin W H oder C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl-aryl und V eine Bindung, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ oder $-OCH_2CH_2CH_2-$ ist.

16. Verbindung nach Anspruch 12, worin R_4 Wasserstoff, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkoxy, F, Cl, Br oder I ist.


17. Verbindung nach Anspruch 12, worin R_4



ist, worin W H oder C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl-aryl ist und V eine Bindung, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ oder $-OCH_2CH_2CH_2-$ ist.

18. Verbindung nach Anspruch 12, worin R_5 Wasserstoff, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkoxy, F, Cl, Br oder I ist.


19. Verbindung nach Anspruch 12, worin R_5



ist, worin W H oder C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl, C_1 -, C_2 -, C_3 -, C_4 -, C_5 - oder C_6 -Alkyl-aryl ist und V eine Bindung, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ oder $-OCH_2CH_2CH_2-$ ist.


20. Verbindung nach Anspruch 12, worin R₁₀ Wasserstoff, C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-Alkoxy, F, Cl, Br oder I ist.

21. Verbindung nach Anspruch 12, worin R₁₀























ist, worin W H oder C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-Alkyl, C₁-, C₂-, C₃-, C₄-, C₅- oder C₆-Alkyl-aryl ist und V eine Bindung, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- oder -OCH₂CH₂CH₂- ist.





















22. Verbindung nach Anspruch 1, wobei die Verbindung aus der Gruppe, bestehend aus





ausgewählt ist.


23. Verbindung, ausgewählt aus der Gruppe, bestehend aus









1




2




3




4




6




7



8




9



(fortgesetzt)


10

5



11


10



15

14


20



25

15


30



35

16


40



45

17


50




55

(fortgesetzt)


18




19




20




21




22



23




27



(fortgesetzt)

108A


5



10


108B

15



109


20



25


111

30




112

35




113

40




45

50




55




(fortgesetzt)


5




10




15




20




25




30



35




40



45


50




55

(fortgesetzt)


5




10




15




20




25




30




35



40




45














50

55



(fortgesetzt)



24. Verbindung gemäß Formel oder Formel 11, Anspruch 22 oder 23 oder ein Salz, Solvat oder Hydrat zur Verwendung bei der Behandlung oder Prävention einer Zellproliferationsstörung oder mikrobiellen Infektion.

25. Verbindung zur Verwendung nach Anspruch 24, wobei die Zellproliferationsstörung aus Krebs, Krebsvorstufen, einer hyperproliferativen Störung, Psoriasis, diabetischer Retinopathie, Makuladegeneration, Epidermis- und Dermoidzysten, Lipomen, Adenomen, kapillaren und kutanen Hämangiomen, Lymphangiomen, Nävusläsionen, Teratomen, Nephromen, Myofibromatose, osteoplastischen Tumoren und Dysplasie ausgewählt ist.


26. Verbindung zur Verwendung nach Anspruch 25, wobei der Krebs aus der Gruppe, bestehend aus Dickdarmkrebs, Lungenkrebs, Brustkrebs, Eierstockkrebs, Gehirnkrebs, Leberkrebs, Bauchspeicheldrüsenkrebs, Prostatakrebs, malignem Melanom, nicht-Melanom-Hautkrebs, Leukämie im Kindesalter, Lymphom, multiplem Myelom, Hodgkin-Krankheit, akuter und chronischer Leukämie, Plasmazellenneoplasma, lymphoidem Neoplasma und mit AIDS zusammenhängendem Krebs, ausgewählt ist.

27. Verbindung zur Verwendung nach Anspruch 24, wobei die mikrobielle Infektion aus der Gruppe, bestehend aus bakterieller Infektion, Pilzinfektion, parasitärer Infektion oder viraler Infektion, ausgewählt ist.

Revendications

1. Composé de formule I :

Formule I



ou un sel, solvate ou hydrate de celui-ci,
formule dans laquelle

T est absent ;

X_y est CZ, CY, N, ou N-O ;

X_z est CZ ;

Y est choisi parmi l'hydrogène, les halogènes, hydroxyle, alkyle inférieur (C₁, C₂, C₃, C₄, C₅, ou C₆), alcoxy en C₁, C₂, C₃, C₄, C₅ ou C₆, O-(alkyle inférieur (C₁, C₂, C₃, C₄, C₅ ou C₆))aryle, et O-benzyle ;

X_a est CR_a, ou N, ou N-O ;


X_b est CR_b, N, ou N-O ;

X_c est CR_c, N, ou N-O ;

X_d est CR_d, N, ou N-O ;

X_e est CR_e ;


R_a, R_b, R_c, R_d, R_e, R₄, R₅, et R₆ sont indépendamment l'hydrogène, un halogène, hydroxyle, alkyle en C₁, C₂, C₃, C₄, C₅, ou C₆, alcoxy en C₁, C₂, C₃, C₄, C₅ ou C₆, O-(alkyle inférieur (C₁, C₂, C₃, C₄, C₅ ou C₆))aryle, O-benzyle, (alkyle en C₁, C₂, C₃, C₄, C₅ ou C₆)-OH, (alkyle en C₁, C₂, C₃, C₄, C₅ ou C₆)-O-alkyle inférieur (C₁, C₂, C₃, C₄, C₅ ou C₆), COOH, COO-alkyle inférieur (C₁, C₂, C₃, C₄, C₅ ou C₆), SO₂H, SO₂-alkyle inférieur (C₁, C₂, C₃, C₄, C₅ ou C₆),




où W est H, ou alkyle en C₁, C₂, C₃, C₄, C₅ ou C₆, (alkyle en C₁, C₂, C₃, C₄, C₅ ou C₆)-aryle ;


V est : une liaison, -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -O-CH₂-, -OCH₂CH₂- ou -OCH₂CH₂CH₂- ; et

Z est :





R_7 , R_8 , R_9 , R_{10} , et R_{11} sont indépendamment l'hydrogène, un halogène, hydroxyle, alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , alcoxy en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , 0-(alkyle inférieur (C_1 , C_2 , C_3 , C_4 , C_5 ou C_6))aryle, O-benzyle, (alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6)-OH, (alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6)-O-alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 ,




R_1 , R_2 , et R_3 sont indépendamment H ou alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 ; et n vaut 1 ou 2; et m vaut 1 ou 2 dans lequel au moins l'un de X_a , X_b , X_c et X_d est N.

2. Composé selon la revendication 1, dans lequel X_y est CY.

3. Composé selon la revendication 1, dans lequel Y est l'hydrogène.


4. Composé selon la revendication 1, dans lequel Z est



5. Composé selon la revendication 1, dans lequel R_b est un alcoxy en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 .

6. Composé selon la revendication 1, dans lequel R_b est l'hydrogène.

7. Composé selon la revendication 1, dans lequel R_b est



où W est H, alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , ou (alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6)-aryle; et V est une liaison, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ ou $-OCH_2CH_2CH_2-$.

8. Composé selon la revendication 8, dans lequel V est une liaison.

9. Composé selon la revendication 1, dans lequel X_a est N et chacun de X_b , X_c , et X_d est CR.

10. Composition comprenant un composé selon la revendication 1 et au moins un excipient pharmaceutiquement

acceptable.

11. Composé selon la revendication 1, lequel composé est l'un des suivants :

5

10

15

20

25

30


35

40

45

50







55



(suite)

Composé

Composé N°
26



10

15

20

25

30

35

40

45

50

55

Composé





Composé N°


33


34


35


36















(suite)

Composé





Composé N°


63


64


65


66




































Composé N°
119

120

121

122



10

15

20

25

30

35

40

45





50


55


(suite)

Composé

Composé N°
127







5

10

15

20

25

30

35

40






45

50


55

(suite)

Composé



Composé N°



10

15

20

25

30

35

40

45











50


55

(suite)


Composé

Composé N°





Composé N°



Composé N°

10

15

20

25

30

35

40

45


50

55

(suite)


Composé

Composé N°



12. Composé selon la revendication 1, lequel composé est un composé de formule II :

Formule II




ou un sel, solvate ou hydrate de celui-ci,
formule dans laquelle R_b , R_4 , R_5 , R_8 et R_{10} sont tels que définis dans la revendication 1.

13. Composé selon la revendication 12, dans lequel R_b est l'hydrogène, F, Cl, Br ou I.

14. Composé selon la revendication 12, dans lequel R_b est un alcoxy en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 .


15. Composé selon la revendication 12, dans lequel R_b est



où W est H, ou alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , ou (alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6) -aryle, et V est une liaison, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ ou $-OCH_2CH_2CH_2-$.

16. Composé selon la revendication 12, dans lequel R_4 est l'hydrogène, alcoxy en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , F, Cl, Br ou I.


17. Composé selon la revendication 12, dans lequel R_4 est



où W est H, ou alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , ou (alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6) -aryle ; et V est une liaison, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ ou $-OCH_2CH_2CH_2-$.

18. Composé selon la revendication 12, dans lequel R_5 est l'hydrogène, alcoxy en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , F, Cl, Br ou I.


19. Composé selon la revendication 12, dans lequel R_5 est



où W est H, ou alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , ou (alkyle en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6) -aryle ; et V est une liaison, $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-O-CH_2-$, $-OCH_2CH_2-$ ou $-OCH_2CH_2CH_2-$.



20. Composé selon la revendication 12, dans lequel R_{10} est l'hydrogène, alcoxy en C_1 , C_2 , C_3 , C_4 , C_5 ou C_6 , F, Cl, Br ou I.

21. Composé selon la revendication 12, dans lequel R_{10} est




où W est H, ou alkyle en C₁, C₂, C₃, C₄, C₅ ou C₆, ou (alkyle en C₁, C₂, C₃, C₄, C₅ ou C₆) -aryle ; et V est une liaison, -CH₂-, 1 -CH₂CH₂-, -CH₂CH₂CH₂-, 1 -O-CH₂-, 1 -OCH₂CH₂- ou -OCH₂CH₂CH₂-.











22. Composé selon la revendication 1, lequel composé est choisi dans l'ensemble constitué par les suivants :

























et




23. Composé choisi dans l'ensemble constitué par les suivantes :






1

5




2

10




3

15




4

20




6

25




7

30




8

35




9

40



10


50




55

(suite)


11




14




15




16




17



18



19




(suite)

20


5

10




21

15



22


20



25


23

30



27


35



40


108A

45




108B

50




55

5




10




15


20




25




30




35




40



45




50




55

(suite)


5




10




15




20




25




30




35




40



45




50




55

(suite)


5




10




15




20




25




30




35




40




45









50



55



(suite)



24. Composé de formule I ou de formule II, selon la revendication 22 ou 23, ou un sel ou solvate ou hydrate, pour utilisation dans le traitement ou la prévention d'un trouble de la prolifération cellulaire ou d'une infection microbienne.

25. Composé pour utilisation selon la revendication 24, dans lequel le trouble de la prolifération cellulaire est choisi parmi un cancer, un pré-cancer, un trouble hyperprolifératif, un psoriasis, une rétinopathie diabétique, une dégénérescence maculaire, les kystes épidermiques et dermoïdes, un lipome, un adénome, un hémangiome capillaire ou cutané, un lymphangiome, des lésions de naevius, un tératome, un néphrome, une myofibromatose, les tumeurs ostéoplasiques, et une dysplasie.

26. Composé pour utilisation selon la revendication 25, dans lequel le cancer est choisi dans l'ensemble constitué par le cancer du côlon, le cancer du poumon, le cancer du sein, le cancer ovarien, le cancer du cerveau, le cancer du foie, le cancer du pancréas, le cancer de la prostate, un mélanome malin, le cancer de la peau sans mélanome, la leucémie de l'enfant, un lymphome, un myélome multiple, la maladie de Hodgkin, une leucémie aiguë ou chronique, un néoplasme à plasmocytes, un néoplasme lymphoïde et un cancer associé au SIDA.

27. Composé pour utilisation selon la revendication 24, dans lequel l'infection microbienne est choisie dans l'ensemble constitué par les infections bactériennes, fongiques, parasitaires et virales.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5578322 A [0258]
- US 5607697 A [0258]
- US 4642903 A [0258]
- US 5631023 A [0258]
- US 4855326 A [0258]
- US 5380473 A [0258]
- US 5518730 A [0258]
- US 6471992 B [0258]
- US 5587172 A [0258]
- US 5616344 A [0258]
- US 6277406 B [0258]
- US 5622719 A [0258]

Non-patent literature cited in the description

- **UBS Warburg.** *Disease Dynamics: The Cancer Market*, 08 November 2000 [0001]
- **Synlett**, 2003, vol. 6, 876-878 [0023]
- **Chou ; Talalay.** *Adv Enzyme Regul.*, 1984, vol. 22, 27-55 [0160]
- **Silverman, R. B.** *The Organic Chemistry of Drug Design and Drug Action*. Academic Press, Inc, 1992, 19-23 [0199]
- **Cahn et al.** Sequence Rule of Cahn, Ingold and Prelog. *Angew. Chem Inter. Edit.*, 1966, vol. 5, 385 [0202]
- **Cahn et al.** *Angew Chem.*, 1966, vol. 78, 413 [0202]
- **Cahn ; Ingold.** *J. Chem Soc.*, 1951, 612 [0202]
- **Cahn et al.** *Experientia*, 1956, vol. 12, 81 [0202]
- **Cahn, J.** *Chem Educ.*, 1964, vol. 41, 116 [0202]
- **Patani ; La Voie.** *Chem. Rev.*, 1996, vol. 96, 3147-3176 [0218]
- **Remington's Pharmaceutical Sciences.** Mack Publishing Company, 1990 [0230]
- **Bundegaard, H.** *Design of Prodrugs*. Elsevier, 1985, 1-92 [0233]
- **Green ; Wuts.** *Protective Groups in Organic Chemistry*. 1991 [0234]
- **Harrison ; Harrison et al.** *Compendium of Synthetic Organic Methods*. John Wiley and Sons, 1971, vol. 1-8 [0234]
- **Kocienski.** *Protecting Groups*. Verlag, 2003 [0234]
- **Remington: the Science and Practice of Pharmacy.** Mack Publishing Co, 1995 [0246]
- **Plé et al.** *J. Med. Chem*, 2004, vol. 47, 871-887 [0372]