

(19)



(11)

EP 1 841 458 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:

Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 13, 55, 68

(51) Int Cl.:

A61K 45/00 (2006.01) A61K 45/06 (2006.01)

(48) Corrigendum issued on:

12.12.2012 Bulletin 2012/50

(86) International application number:

PCT/US2006/000308

(45) Date of publication and mention of the grant of the patent:

28.12.2011 Bulletin 2011/52

(87) International publication number:

WO 2006/074296 (13.07.2006 Gazette 2006/28)

(21) Application number: **06717499.5**

(22) Date of filing: **06.01.2006**

(54) **SURFACTANT TREATMENT REGIMEN FOR TREATING OR PREVENTING BRONCHOPULMONARY DYSPLASIA**

DOSIERUNGSSCHEMA UNTER VERWENDUNG EINES TENSIDS ZUR BEHANDLUNG ODER VERHINDERUNG VON LUNGENDYSPLASIEN

REGIME DE TRAITEMENT PAR TENSIOACTIFS POUR LE TRAITEMENT PROPHYLACTIQUE OU THERAPEUTIQUE DE LA DYSPLASIE BRONCHOPULMONAIRE

(84) Designated Contracting States:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR

- **GUARDIA, Carlos**
Philadelphia, Pennsylvania 19103 (US)

(30) Priority: **06.01.2005 US 641805 P**

(43) Date of publication of application:

10.10.2007 Bulletin 2007/41

(74) Representative: **Main, Malcolm Charles Murgitroyd & Company**
Scotland House
165-169 Scotland Street
Glasgow G5 8PL (GB)

(60) Divisional application:

10185423.0 / 2 298 349

(56) References cited:

WO-A-97/19701 WO-A-02/062289
WO-A-2005/115520 WO-A-2006/026237

(73) Proprietor: **Discovery Laboratories, Inc.**
Warrington PA 18976 (US)

- **GORTNER L ET AL: "Early treatment of respiratory distress syndrome with bovine surfactant in very preterm infants: a multicenter controlled clinical trial." PEDIATRIC PULMONOLOGY. SEP 1992, vol. 14, no. 1, September 1992 (1992-09), pages 4-9, XP009065658 ISSN: 8755-6863**

(72) Inventors:

- **KILLIAN, Anthony**
Easton, Pennsylvania 18042 (US)
- **SCHABER, Christopher**
Wrightstown, New Jersey 08562 (US)
- **SEGAL, Robert**
Gwynedd Valley,
Pennsylvania 19437 (US)

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

EP 1 841 458 B9

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**CROSS-REFERENCE TO RELATED APPLICATIONS**

5 [0001] This application claims benefit under 35 U.S.C. § 119(e) to application Serial No. 60/641,805 filed January ,2005,

FIELD

10 [0002] This invention relates to the treatment of pulmonary disorders. In particular, regimens for the treatment or prevention of bronchopulmonary dysplasia and related disorders are provided, featuring the use of pulmonary surfactants.

BACKGROUND

15 [0003] Natural pulmonary surfactants (PS) are protein/lipid compositions that are produced naturally in the lungs and are critical to the lungs' ability to absorb oxygen. They cover the entire alveolar surface of the lungs and the terminal conducting airways leading to the alveoli. Surfactants facilitate respiration by continually modifying the surface tension of the fluid normally present within the alveoli. In the absence of sufficient surfactant, or should the surfactant degrade, the alveoli tend to collapse and the lungs do not absorb sufficient oxygen. By lowering the surface tension of the terminal conducting airways, surfactant maintains patency, *i.e.*, keeps airways open. Loss of patency leads to loss of patency
20 obstruction of the airway and compromised pulmonary function. Human surfactants primarily contain: phospholipids, the major one being dipalmitoyl phosphatidyl-choline (DPPC), and four surfactant polypeptides, A, B, C and D with surfactant protein B (SP-B) being the most essential for respiratory function. Natural and synthetic pulmonary surfactants are commonly used to treat respiratory distress syndrome in premature infants shortly after birth.

25 [0004] Bronchopulmonary dysplasia (BPD or BD), also referred to as chronic lung disease (CLD), is a common, occasionally life threatening, lung disease typically occurring in premature infants who survive respiratory distress syndrome (RDS) and other complications of prematurity. With the near universal adoption of antenatal steroids and post-natal use of exogenous lung surfactants, large numbers of at-risk low birth weight infants now survive the acute lung disease of prematurity only to develop chronic lung disease. Prior to the widespread use of synthetic or animal derived surfactants, BPD was histologically characterized by airway injury and fibrosis. Since the advent of surfactant replacement
30 therapy, infants appear to experience less airway and fibrotic involvement than described in earlier reports but have abnormalities in alveolarization and vascularization. The change in the nature of the condition has led neonatologists to refer to a "new BPD". BPD can also develop in term infants who require respiratory support at birth or soon thereafter.

35 [0005] Surfactant therapy has decreased the frequency with which larger, moderately premature infants develop the disease, but the survival benefits of surfactant in extremely premature infants has resulted in a shift in the incidence of BPD toward smaller, more premature babies. Post-natal steroids in anti-inflammatory doses are known to modestly decrease the physiologic consequences of BPD, but at the expense of neuro-developmental outcomes. Further, steroids may decrease alveolarization. Vitamin A derivatives stimulate alveolarization in animals, but clinical results have been modest. Despite an intriguing rationale, antioxidants have proven to be of little value. Whether inhaled nitric oxide will decrease the likelihood of BPD in very low birth weight infants awaits the outcome of ongoing clinical trials.

40 [0006] Causes of BPD or CLD are probably multi-factorial. However, the pathophysiologic antecedents and consequences of BPD may suggest a new approach to prevention with drugs already in clinical use. Decreased lung compliance as a consequence of initially mild pulmonary edema has been shown to correlate with subsequent development of BPD. Neutrophil migration to the lung increases in the presence of pulmonary edema and hyperoxia. When these cells are activated, they release proteases and inflammatory mediators including oxidants that can further injure the lung. These
45 events degrade pulmonary function necessitating mechanical ventilation for many infants initially weaned from the ventilator and increasing ventilator requirements for those never able to wean. Surfactant is itself a target of proteases, oxidant species and small molecule mediators and can be inactivated by many plasma proteins found in bronchoalveolar fluid in the presence of capillary leak.

50 [0007] There are prior art documents which discuss the use of surfactants such as *Ped. Pulmon.* 14:4-9. (1992) which describes a treatment of Respiratory Distress Syndrome (RDS) in pre-term infants, comprising administering a bolus of surfactant within one hour of birth, and repeating the dosing 12 or 24 hours later.

[0008] WO 97/19701 describes the use of covalent conjugates of glycosaminoglycans administered together with surfactants. In particular, heparin conjugates are used as an adjunctive therapy to the surfactants to prevent Respiratory Distress Syndrome and bronchopulmonary dysplasia. D2 also describes the use of such conjugates to prevent throm-
55 bogenesis, in particular in lung airways, such as those found in infant and adult respiratory distress syndrome and on surfaces in contact with blood.

[0009] WO 02/062289 describes a method for treating infants, children or adults suffering from pulmonary distress caused by low or insufficient production of surfactant. It is particularly suitable for treating premature infants suffering

from Respiratory Distress Syndrome. The method comprises administering a leptin compound to an individual with impaired surfactant production for a time and in an amount sufficient to enhance surfactant production.

[0010] Treatments to prevent or ameliorate BPD and rotated pulmonary disorders are needed. The present invention relates to use of a pulmonary surfactant as claimed hereafter.

[0011] Preferred embodiments of the invention are set forth in the dependent claims.

BRIEF SUMMARY

[0012] There is described a method for treating or preventing BPD in an infant treated with pulmonary surfactant for respiratory distress syndrome ("RDS"). Following the treatment for respiratory distress syndrome with pulmonary surfactant, pulmonary surfactant is administered to the infant in an amount and for a time effective to treat or prevent the BPD. Typically, administration of the pulmonary surfactant for BPD is initiated after the treatment of the respiratory distress syndrome with pulmonary surfactant, for instance, at the next dosage interval following completion of the treatment for RDS, at day 3 of life of the infant, or whenever the treatment for RDS is completed. In certain cases Administration of the pulmonary surfactant may be continued through at least 36 weeks post-menstrual age ("PMA") of the infant or alternatively through at least 28 days post-natal age of the infant. According to the invention, administration of the pulmonary surfactant is continued through at least day 10, 14 or day 18 of life of the infant. In addition treatment is initiated anytime from day 3 of life of the infant to day 10,

[0013] Also described is a method for treating or preventing BPD in an infant requiring respiratory support. The pulmonary surfactant is administered to the infant in an amount and for a time effective to treat or prevent the BPD. The PS can be administered until the infant no longer requires respiratory support. The PS may be administered at or after day 1 of the life of the infant and is continued through at least 28 days post-natal age of the infant. The administering may be initiated at day 1 of life of the infant or at day 3 of life of the infant. according to the present invention, the administering is initiated before the infant has been diagnosed with BPD and the infant is treated with pulmonary surfactant at least once after day 2 or 3 of life of the infant. Administration may be initiated at day 1 of life of the infant or at day 2, of life of the infant and the infant is treated with pulmonary surfactant at least once after day 2, or 3, through to day 10 of life of the infant according to the present invention, treatment is continued through at least day 10, day 14, or day 18 of life of the infant administration of PS can be initiated anytime from day 3 to day 18, or day 3 to day 10 of life of the infant. The infant requiring respiratory support may or may not exhibit respiratory distress syndrome.

[0014] Methods of administering pulmonary surfactant to an infant are known in the art. Administration can be by endotracheal administration, aerosolization and inhalation. The term "inhalation" includes, for example, both inhalation of a dry powder and inhalation of a wet aerosol.

[0015] Use of a pulmonary surfactant according to a preferred embodiment of the invention may be accompanied by another respiratory therapy, such as mechanical ventilation, continuous positive airway pressure (CPAP) including nasal CPAP (nCPAP) or administration of other therapeutic agents.

[0016] Preferably the method comprises administering a synthetic pulmonary surfactant. Particularly preferred is a pulmonary surfactant that contains a peptide having a sequence of SEQ ID NO: 1.

[0017] Other features and advantages of the invention will be understood from the detailed description and examples that follow.

DETAILED DESCRIPTION

[0018] Unless otherwise specified, all medical procedures described or referred to herein are performed in accordance with current standards of care understood by physicians and/or other healthcare practitioners.

[0019] It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting. As used in this specification and the appended claims, the singular forms "a", "an", and "the" include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to "a peptide" includes a combination of two or more peptides, and the like.

[0020] "About" as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of $\pm 20\%$ or $\pm 10\%$, more preferably $\pm 5\%$, even more preferably $\pm 1\%$, and still more preferably $\pm 0.1\%$ from the specified value, as such variations are appropriate to perform the disclosed methods.

[0021] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although any methods and materials similar or equivalent to those described herein can be used in the practice for testing of the present invention, the preferred materials and methods are described herein. In describing and claiming the present invention, the following terminology will be used.

[0022] "Polypeptide," "peptide" "peptoids" and "protein" are used interchangeably to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical

mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid. Non-natural residues are well described in the scientific and patent literature; a few exemplary non-natural compositions useful as mimetics of natural amino acid residues and guidelines are described below. Mimetics of aromatic amino acids can be generated, for example, by replacing by, e.g., D- or L-naphylalanine; D- or L-phenylglycine; D- or L-2 thieneylalanine; D- or L-1, -2,3-, or 4-pyreneylalanine; D- or L-3 thieneylalanine; D- or L-(2-pyridinyl)-alanine; D- or L-(3-pyridinyl)-alanine; D- or L-(2-pyrazinyl)-alanine; D- or L-(4-isopropyl)-phenylglycine; D-(trifluoromethyl)-phenylglycine; D-(trifluoromethyl)-phenylalanine; D-p-fluoro-phenylalanine; D- or L-p-biphenylphenylalanine; K- or L-p-methoxy-biphenylphenylalanine; D- or L-2-indole(alkyl)alanines; and, D- or L-alkylainines, where alkyl can be substituted or unsubstituted methyl, ethyl, propyl, hexyl, butyl, pentyl, isopropyl, iso-butyl, sec-isotyl, iso-pentyl, or a non-acidic amino acids. Aromatic rings of a non-natural amino acid include, e.g., thiazolyl, thiophenyl, pyrazolyl, benzimidazolyl, naphthyl, furanyl, pyrrolyl, and pyridyl aromatic rings.

[0023] "Peptide" includes peptides which are conservative variations of those peptides specifically exemplified herein. "Conservative variation" as used herein denotes the replacement of an amino acid residue by another, biologically similar residue. Examples of conservative variations include, but are not limited to, the substitution of one hydrophobic residue such as isoleucine, valine, leucine, alanine, cysteine, glycine, phenylalanine, proline, tryptophan, tyrosine, norleucine or methionine for another, or the substitution of one polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acids, or glutamine for asparagine, and the like. Neutral hydrophilic amino acids which can be substituted for one another include asparagine, glutamine, serine and threonine. "Conservative variation" also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid provided that antibodies raised to the substituted polypeptide also immunoreact with the unsubstituted polypeptide. Such conservative substitutions are within the definition of the classes of the peptides described herein "Cationic" as used herein refers to any peptide that possesses a net positive charge at pH 7.4. The biological activity of the peptides can be determined by standard methods known to those of skill in the art and described herein.

[0024] Peptides described herein can be synthesized by methods known in the art. For example, in certain embodiments, commonly used methods such as t-BOC or Fmoc protection of alpha-amino groups can be used. Both methods involve stepwise syntheses whereby a single amino acid is added at each step starting from the C terminus of the peptide (See, Coligan et al., Current Protocols in Immunology, Wiley Interscience, 1991, Unit 9). Peptides of the invention can be synthesized, for example, by the well known solid phase peptide synthesis methods described in Merrifield, J. Am. Chem. Soc. 85: 2149, 1962, and Stewart and Young, 1969, Solid Phase Peptides Synthesis, pp. 27-62, using a copoly (styrene-divinylbenzene) containing 0.1-1.0 mMol amines/g polymer. On completion of chemical synthesis, the peptides can be deprotected and cleaved from the polymer by treatment with liquid HF-10% anisole for about 1/4-1 hours at 0°C. After evaporation of the reagents, the peptides are extracted from the polymer with 1% acetic acid solution which is then lyophilized to yield the crude material. This can normally be purified by such techniques as gel filtration on Sephadex G-15 using 5% acetic acid as a solvent. Lyophilization of appropriate fractions of the column will yield the homogeneous peptide or peptide derivatives, which can then be characterized by such standard techniques as amino acid analysis, thin layer chromatography, high performance liquid chromatography, ultraviolet absorption spectroscopy, molar rotation, solubility, and quantitated by the solid phase Edman degradation.

[0025] "Recombinant" when used with reference to a protein indicates that the protein has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein.

[0026] For the purposes of present invention, BPD is diagnosed at 28 days of age or thereabout, 36 weeks PMA or thereabout, or as set forth in Table 1 below. An infant that either remains on mechanical ventilation or requires supplemental oxygen in order to maintain oxygen saturation levels (" SaO_2 ") greater than or equal to 90% (with the exception of infants requiring supplemental oxygen during feedings) at 36 weeks PMA or at 28 days of age is said to have BPD. Generally, chest x-rays will also be performed for infants at 28 days of age in order to confirm the BPD diagnosis. The x-ray of lungs with BPD often have a bubbly, sponge-like appearance. X-rays are diagnostic tests which use invisible electromagnetic energy beams to produce images of internal tissues, bones, and organs onto film.

[0027] Alternatively, BPD can be diagnosed using the criteria set forth in Table 1 below from Jobe and Bancalari, Am J Respir Crit Care Med, 2001, 163:1723-1729

Table 1

Gestational Age	< 32 weeks	≥ 32 weeks
Time point of assessment	• 36 weeks PMA or discharge to home, whichever comes first	• > 28 days but < 56 days post-natal age or discharge to home, whichever comes first

(continued)

Gestational Age	< 32 weeks	≥ 32 weeks
5	<ul style="list-style-type: none"> • Infants have already been treated with oxygen > 21% for at least 28 days plus Breathing room air at 36 weeks PMA or discharge, whichever comes first 	<ul style="list-style-type: none"> • Infants have already been treated with oxygen > 21% for at least 28 days plus Breathing room air by 56 days post-natal age or discharge, whichever comes first
	<p>Mild BPD</p>	
	<p>Moderate BPD</p>	<p>Need for < 30% oxygen at 56 days post-natal age or discharge, whichever comes first</p>
10	<p>Need for < 30% oxygen at 36 weeks PMA or discharge, whichever comes first</p>	<p>Need for < 30% oxygen at 36 weeks PMA or discharge, whichever comes first</p>
	<p>Severe BPD</p>	<p>Need for ≥ 30% oxygen and/or positive pressure (e.g., PPV or NCPAPA) at 56 days post-natal age or discharge, whichever comes first</p>
15	<p>Need for ≥ 30% oxygen and/or positive pressure (e.g., PPV or NCPAPA) at 36 weeks PMA or discharge, whichever comes first</p>	<p>Need for ≥ 30% oxygen and/or positive pressure (e.g., PPV or NCPAPA) at 36 weeks PMA or discharge, whichever comes first</p>

[0028] Infants treated with oxygen > 21% and/or positive pressure for nonrespiratory disease (e.g., central apnea or diaphragmatic paralysis) as provided in Table 1 do not have BPD unless they also develop parenchymal lung disease and exhibit clinical features of respiratory distress. A day of treatment with oxygen > 21% means that the infant received oxygen > 21% for more than 12 hours on that day. Treatment with oxygen > 21% and/or positive pressure at 36 weeks PMA, or at 56 days post-natal age or discharge, should not reflect an "acute" event, but should rather reflect the infant's usual daily therapy for several days preceding and following 36 weeks PMA, 56 days post-natal age, or discharge.

[0029] For the purposes of present invention, "Respiratory Distress Syndrome" is defined as being present in an infant at 24 hours of age with a need for a fraction of inspired oxygen ("FiO₂") ≥ 0.30 combined with the demonstration of a reticulogranular pattern on a chest radiograph obtained between 20 and 28 hours of age.

[0030] "Supplemental oxygen" refers to any requirement for additional oxygen to maintain oxygen saturation levels ≥ 90%. This can include mechanical ventilation and/or CPAP if supplemental oxygen (≥ 21% FiO₂) is being given.

[0031] "Infant" includes neonatal infants. Typically a neonatal infant is an infant born prematurely or otherwise, under 4 weeks old. For the purposes used herein, infants are under 2 years old.

[0032] "Respiratory support" includes any intervention that treats respiratory illness including, for example, the administration of supplemental oxygen, mechanical ventilation, CPAP, albuterol treatment, and the like.

[0033] "Treating" refers to any indicia of success in the treatment or amelioration of the disease or condition, e.g., BPD. Treatment at Day 1 (or DOL 1) is the day on which Time 0 (time of birth) occurs and DOL 2, 3, and so on, begin at 00:00 (midnight) each subsequent day following Time 0.

[0034] "Preventing" refers to the prevention of the disease or condition, e.g., BPD, in the patient. For example, if an infant exhibiting respiratory distress syndrome, or requiring any form of respiratory support at birth, is treated with the methods of the present invention and is not later diagnosed with BPD, e.g., at 28 days of age or at 36 weeks PMA, it will be understood that BPD has been prevented in that infant.

[0035] "Surfactant activity" refers to the ability of any substance, such as an organic molecule, protein or polypeptide, either alone or in combination with other molecules, to lower surface tension at an air/water interface. The measurement can be made with a Wilhelmy Balance or pulsating bubble surfactometer by an *in vitro* assay. See, for example King et al, Am. J. Physiol. 1972, 223:715-726, or Enhorning, J. Appl. Physiol., 1977, 43:198-203 Briefly, the Enhorning Surfactometer (Surfactometer International, Toronto, Ontario) measures the pressure gradient (δP) across a liquid-air interface of a bubble that pulsates at a rate of 20 cycles/min between a maximal (0.55 mm) and minimal (0.4 mm) radius. The bubble, formed in a 37°C, water-enclosed, 20µl sample chamber, is monitored through a microscopic optic while the pressure changes are recorded on a strip chart recorder calibrated for 0 and -2 cm H₂O. In addition, *in vivo* measurements of increases of compliance or airflow at a given pressure of air entering the lung can be readily made, such as in the assay of Robertson, Lung, 1980, 158:57-68, In this assay, the sample to be assessed is administered through an endotracheal tube to fetal rabbits or lambs delivered prematurely by Caesarian section. (These "preemies" lack their own PS, and are supported on a ventilator.) Measurements of lung compliance, blood gases and ventilator pressure provide indices of activity. *In vitro* assays of surfactant activity, which is assessed as the ability to lower the surface tension of a pulsating bubble, and *in vivo* assays utilizing fetal rabbits is described in detail by Revak et al, Am. Rev. Respir. Dis., 1986, 134:1258-1265.

[0036] There is described herein the therapeutic or prophylactic administration of pulmonary surfactant (PS) to infants exhibiting or at risk of developing bronchopulmonary dysplasia (BD or BPD). PS administration can be initiated in an infant exhibiting respiratory distress syndrome, following treatment of such syndrome with pulmonary surfactant or by another means (e.g., ventilation) or a combination thereof. Typically, infant RDS is treated via PS therapy within the first few hours to one or two days after birth. Herein described is the administration of PS for the treatment or prevention of BPD initiated following the RDS treatment, e.g., at day 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 of life or thereafter, or

whenever the treatment for RDS is completed. The infant will may be treated with pulmonary surfactant at the next dosage interval appropriate for such treatment after completion of RDS treatment. For example, where PS is administered as a liquid by intratracheal instillation, treatment with PS for BPD will be about 6 hours following the last treatment of PS for RDS. A clinician will be able to determine, based on administration protocols known in the art, when dosage of PS for the treatment of BPD should be initiated.

[0037] The infant may not receive PS therapy for RDS, but nonetheless can be administered PS to treat or prevent BPD, in accordance with the present invention. The treatment with PS is initiated following the episode of respiratory distress syndrome. In certain preferred embodiments, the administering is initiated before the infant has been diagnosed with BPD and the infant is treated with pulmonary surfactant at least once after day 2, 3, 4, 5, 6, 7, 8, 9, or 10 of life of the infant. The administration of PS can be initiated, for example, at or after day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14 of life or thereabout of the infant. Methods of diagnosing respiratory distress syndrome are known in the art and are thus not described herein in detail.

[0038] Infants to be treated with the methods according to the present invention may require respiratory support but do not necessarily exhibit respiratory distress syndrome. These infants either have not been diagnosed with RDS or have not been treated with PS for RDS. In these infants, treatment can be initiated as soon as it is recognized that these infants require respiratory support. In certain embodiments, this will be at day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 of life or thereabout, or between about day 3 to day 10 of the life of the infant. In certain embodiments, this can be later than day 10 of the life of the infant, *e.g.*, day 14, day 18 or later.

[0039] PS therapy can be provided concomitantly with other forms of respiratory support. Typically, the PS therapy is initiated before the infant has been diagnosed with BPD. In certain embodiments, the present methods will prevent the infant from developing BPD.

[0040] Treatment is continued for a period of time deemed by a physician or other medical practitioner as appropriate to achieve a therapeutic or prophylactic effect. Treatment can be continued, for example, until the infant no longer requires respiratory support or even longer. Typically, treatment is continued for one to two weeks, *e.g.*, until day 14 or day 18 of the infant's life or thereabout. However, the duration of treatment can be shorter, *e.g.*, one week, or longer, *e.g.*, extending three or more weeks, or until the infant is discharged from neonatal intensive care or from the hospital. For example, treatment can be continued through at least 36 weeks PMA of the infant or at least 28 days post-natal age of the infant. In certain embodiments, wherein the infant does not exhibit respiratory distress syndrome at birth but, days later, develops a condition that requires respiratory support, treatment with PS may not be initiated until well after day 3 of life of the infant and thus treatment will typically be continued past day 14 or day 18 of the infant's life. In infants born prematurely, for example, regardless of whether the infant exhibited respiratory distress syndrome at or near birth, treatment can be continued, for example, until 36 weeks PMA or even later.

[0041] In various embodiments, the PS is administered periodically or continuously throughout the treatment period, at dosages and utilizing protocols in accordance with standard and/or manufacturer's instructions (see, *e.g.*, Example 1).

[0042] The PS selected for use in the methods of the invention can be the same as, or different from, the PS utilized for RDS. Any PS currently in use, or hereafter developed for use in RDS and other pulmonary conditions, is suitable for use in the present invention.

[0043] A pulmonary surfactant according to the present invention may comprise a cationic peptide that can be derived from animal sources or synthetically. Exemplary peptides for use herein include naturally and non-naturally occurring pulmonary surfactant polypeptides, such as, for example, one or a combination of animal-derived SP-A, SP-B, SP-C, or SP-D polypeptides; recombinant SP-A, SP-B, SP-C, or SP-D polypeptides; synthetically derived SP-A, SP-B, SP-C, or SP-D polypeptides; SP-A, SP-B, SP-C, and SP-D analogs; SP-A, SP-B, SP-C, and SP-D polypeptide mimics; conservatively modified variants thereof retaining activity; and fragments thereof retaining activity. A pulmonary surfactant polypeptide mimic is generally a polypeptide that is engineered to mimic the essential attributes of human surfactant protein. In The present invention the pulmonary surfactant polypeptide comprises a cationic peptide that consists of at least about 10 amino acid residues, other examples are at least 11 amino acid residues, and no more than about 80, more usually fewer than about 35 and preferably fewer than about 25 amino acid residues.

[0044] Exemplary amino acid sequences of pulmonary surfactant polypeptides for use herein, methods of isolating them, and producing them by genetic engineering techniques are known in the art. See for example, U.S. Patent Nos. 5,874,406; 5,840,527; 4,918,161; 5,827,825; 6,660,833; 5,006,343; 5,455,227; 5,223,481; 5,753,621; 5,891,844; 4,861,756; 5,272,252; 5,024,95; 5,238,920; 5,302,481; 6,022,955; 5,874,406; 5,840,527; 5,827,825; 6,013,619; 6,660,833; and International Publication Nos, WO8603408 and WO08904326. A preferred lung surfactant peptide for use herein is a SP-B or SP-C polypeptide, or polypeptide mimic.


[0045] The synthetic pulmonary surfactant may comprise one or more phospholipids and a polypeptide, in which the polypeptide, when admixed with a phospholipid, forms a synthetic pulmonary surfactant having a surfactant activity greater than the surfactant activity of the phospholipid alone. A particularly preferred pulmonary surfactant polypeptide for use herein is a SP-B polypeptide or polypeptide mimic. SP-B is the protein in natural pulmonary surfactant known to be the most important surfactant protein for surface tension lowering and promoting oxygen exchange. SP-B polypeptide

mimics are small hydrophobic polypeptides, generally less than about 80 amino acids in size. Many SP-B polypeptide mimics possess a repeating hydrophobic cationic motif. Like natural SP-B polypeptide, SP-B mimics, preferably, lower surface tension of the terminal conducting airways and promote oxygen exchange.

[0046] A preferred SP-B mimetic for use in the present invention is KL4 peptide, which is a cationic peptide containing repeating lysine and leucine residues. KL4 is representative of a family of pulmonary surfactant polypeptide mimetics which are described, for example, in U.S. Patent 5,260,273, 5,164,369, 5,407,914 and 6,613,734. Methods of preparing the KL4 peptide can be found in U.S. Patent 5,164,369.


[0047] Pulmonary surfactants polypeptide mimics may refer to polypeptides with an amino acid residue sequence that has a composite hydrophobicity of less than zero, preferably less than or equal to -1, more preferably less than or equal to -2. The composite hydrophobicity value for a peptide is determined by assigning each amino acid residue in a peptide its corresponding hydrophilicity value as described in Hopp et al., Proc. Natl. Acad. Sci. 78: 3824-3829, 1981. For a given peptide, the hydrophobicity values are summed, the sum representing the composite hydrophobicity value. These hydrophobic polypeptides typically perform the function of the hydrophobic region of SP18. Accordingly, the amino acid sequence of the pulmonary surfactant polypeptide mimic may mimic the pattern of hydrophobic and hydrophilic residues of SP18 and perform the function of the hydrophobic region of SF18. SP18 is a known lung surfactant apoprotein, more thoroughly described in Glasser et al., Proc. Natl. Acad. Sci. 84: 4007-4001, 1987. It should be understood, however, that polypeptides and other surfactant molecules described herein are not limited to molecules having sequences like that of native SP18. On the contrary, some preferred surfactant molecules have little resemblance to SP18 with respect to a specific amino acid residue sequence, except that they have similar surfactant activity and alternating charged/uncharged (or hydrophobic/hydrophilic) residue sequences.

[0048] According to the present invention polypeptides for use herein have alternating hydrophobic and hydrophilic amino acid residue regions and are characterized as having at least 10 amino acid residues represented by the formula:




Z and U are amino acid residues such that at each occurrence Z and U are independently selected. Z is a hydrophilic amino acid residue, preferably selected from the group consisting of R, D, E and K. U is a hydrophobic amino acid residue, preferably selected from the group consisting of V, I, L, C, Y, and F. The letters, "a," "b," "c" and "d" are numbers which indicate the number of hydrophilic or hydrophobic residues. The letter "a" has an average value of about 1 to about 5, preferably about 1 to about 3. The letter "b" has an average value of about 3 to about 20, preferably about 3 to about 12, most preferably, about 3 to about 10. The letter "c" is 1 to 10, preferably, 2 to 10, most preferably 3 to 6. The letter "d" has an average value of about 0 to 3, preferably 1 to 2.

[0049] Also described are surfactant polypeptides which include a sequence having alternating groupings of amino acid residues as represented by the formula:




wherein Z is an amino acid residue independently selected from the group consisting of R, D, E, and K; J is an α -aminoaliphatic carboxylic acid; a has an average value of about 1 to about 5; b has an average value of about 3 to about 20; c is 1 to 10; and d is 0 to 3.

[0050] Also described are polypeptides which have alternating groupings of amino acids residue regions as represented by the formula:



wherein B is an amino acid residue independently selected from the group consisting of H, 5-hydroxylysine, 4-hydroxyproline, and 3-hydroxyproline; and U is an amino acid residue independently selected from the group consisting of V, I, L, C, Y, and F. In one preferred variation, B is an amino acid derived from collagen and is preferably selected from the group consisting of 5-hydroxylysine, 4-hydroxyproline, and 3-hydroxyproline; a has an average value of about 1 to about 5; b has an average value of about 3 to about 20; c is 1 to 10; and d is 0 to 3.


[0051] Further described are surfactant polypeptides which include a sequence having alternating groupings of amino acid residues as represented by the formula:



wherein B is an amino acid residue independently selected from the group consisting of H, 5-hydroxylysine, 4-hydroxyproline, and 3-hydroxyproline; and J is an α -aminoaliphatic carboxylic acid; a has an average value of about 1 to about 5; b has an average value of about 3 to about 20; c is 1 to 10; and d is 0 to 3.

[0052] Where "J" is included in the relevant formula, J is an α -aminoaliphatic carboxylic acid having four to six carbons, inclusive. In variations, J is an α -aminoaliphatic carboxylic acid having six or more carbons, inclusive. In other variations, J is selected from the group consisting of α -aminobutanoic acid, α -aminopentanoic acid, α -amino-2-methylpropanoic acid, and α -aminohexanoic acid.

[0053] Surfactant polypeptides of the present invention may comprise a sequence having including a sequence having alternating groupings of amino acid residues as represented by the formula:



wherein Z is an amino acid residue independently selected from the group consisting of R, D, E, and K; and U is an amino acid residue independently selected from the group consisting of V, I, L, C, Y and F; from the group consisting of V, I, L, C and F; or from the group consisting of L and C; a has an average value of about 1 to about 5; b has an average value of about 3 to about 20; c is 1 to 10; and d is 0 to 3.

[0054] In the foregoing formulas, Z and U, Z and J, B and U, and B and J are amino acid residues that, at each occurrence, are independently selected. In addition, in each of the aforementioned formulae, a generally has an average value of about 1 to about 5; b generally has an average value of about 3 to about 20; c is 1 to 10; and d is 0 to 3.

[0055] Z and B may be charged amino acid residues. Z and B may be hydrophilic or positively charged amino acid residues. According to the invention, Z is preferably selected from the group consisting of R, D, E and K. However also described is Z selected from the group consisting of R and K. Further described is B selected from the group consisting of H, 5-hydroxylysine, 4-hydroxyproline, and 3-hydroxyproline. B can be H, or alternatively is a collagen constituent amino acid residue and is selected from the group consisting of 5-hydroxylysine, (δ -hydroxylysine), 4-hydroxyproline, and 3-hydroxyproline.

[0056] U and J may be uncharged amino acid residues. U and J may be hydrophobic amino acid residues. According to the invention U is preferably selected from the group consisting of V, I, L, C, Y, and F. In another preferred embodiment, U is selected from the group consisting of V, I, L, C, and F. However, also described is U selected from the group consisting of L and C. U can be L.

[0057] Similarly, B can be an amino acid preferably selected from the group consisting of H, 5-hydroxylysine, 4-hydroxyproline, and 3-hydroxyproline. Alternatively, B can be selected from the group consisting of collagen-derived amino acids, which includes 5-hydroxylysine, 4-hydroxyproline, and 3-hydroxyproline.

[0058] Charged and uncharged amino acids can be selected from groups of modified amino acids. For example, a charged amino acid can be selected from the group consisting of citrulline, homoarginine, or ornithine, to name a few examples. Similarly, the uncharged amino acid can be selected from the group consisting of α -aminobutanoic acid, α -aminopentanoic acid, α -amino-2-methylpropanoic acid, and α -aminohexanoic acid.

[0059] According to the present invention, items "a", "b", "c" and "d" are numbers which indicate the number of charged or uncharged residues (or hydrophilic or hydrophobic residues). "a" has an average value of about 1 to about 5, preferably about 1 to about 3, more preferably about 1 to about 2, and even more preferably, 1.

[0060] "b" has an average value of about 3 to about 20, preferably about 3 to about 12, more preferably about 3 to about 10, even more preferably in the range of about 4-8. In one example, "b" is about 4.

[0061] "c" is 1 to 10, preferably 2 to 10, more preferably in the range of 3-8 or 4-8, and even more preferably 3 to 6. In one example, "c" is about 4.

[0062] "d" is 0 to 3 or 1 to 3. In one example, "d" is 0 to 2 or 1 to 2; in another example, "d" is 1.

[0063] By stating that an amino acid residue is independently selected, it is meant that at each occurrence, a residue from the specified group is selected. That is, when "a" is 2, for example, each of the hydrophilic residues represented by Z will be independently selected and thus can include RR, RD, RE, RK, DR, DD, DE, DK, and the like. By stating that "a" and "b" have average values, it is meant that although the number of residues within the repeating sequence (e.g., $Z_a U_b$) can vary somewhat within the peptide sequence, the average values of "a" and "b" would be about 1 to about 5 and about 3 to about 20, respectively.

[0064] For example, using the formula $(Z_a U_b)_c Z_d$ for the peptide designated "KL8" in Table 2 below, the formula can be rewritten as $K_1 L_8 K_1 L_8 K_1 L_2$, wherein the average value of "b" is six [i.e., $(8+8+2)/3=6$], c is three and d is zero.

[0065] Polypeptides described herein can also be subject to various changes, such as insertions, deletions and substitutions, either conservative or non-conservative, where such changes provide for certain advantages in their use.

[0066] Additional residues can be added at either terminus of a polypeptide of the present invention, such as for the purpose of providing a "linker" by which such a polypeptide can be conveniently affixed to a label or solid matrix, or carrier. Labels, solid matrices and carriers that can be used with the polypeptides described herein are known in the art.

[0067] Amino acid residue linkers are usually at least one residue and can be 40 or more residues, more often 1 to 10 residues. Typical amino acid residues used for linking are tyrosine, cysteine, lysine, glutamic and aspartic acid, or the like. In addition, a polypeptide sequence described herein can differ from the natural sequence by the sequence being modified by terminal-NH₂ acylation, e.g., acetylation, or thioglycolic acid amidation, terminal-carboxylamidation,

e.g., ammonia, methylamine, and the like.

[0068] Exemplary SP-B polypeptide mimics that can be used in the present invention include, but are not limited to, those shown in the Table 2.

Table 2: Pulmonary Surfactant Mimetic Peptides

Designation ¹	SEQ ID NO	Amino Acid Residue Sequence
KL4	1	KLLLLKLLLLKLLLLKLLLLK
DLA	2	DLLLLDLLLLDLLLLDLLLLD
RLA	3	RLLLLRLLLLRLLLLRLLLLR
RL8	4	RLLLLLLLLRLLLLLLLLRLL
R2L7	5	RRLLLLLLLLRRLLLLLLLLRRL
	6	RLLLLCLLLRLLLLCLLLR
	7	LLLLCLLLRLLLLCLLLRLL
	8	RLLLLCLLLRLLLLCLLLRLLLLCLLLR DLLLLDLLLLDLLLLDLLLLD
RCL1	9	RLLLLCLLLRLLLLCLLLR
RCL2	10	RLLLLCLLLRLLLLCLLLRLL
RCL3	11	RLLLLCLLLRLLLLCLLLRLLLLCLLLR
KL8	12	KLLLLLLLLKLLLLLLLLKLL
KL7	13	KKLLLLLLLLKKLLLLLLLLKKL

¹The designation is an abbreviation for the indicated amino acid residue sequence.

[0069] Also described are a variety of surfactant molecules, including proteins, polypeptides, and molecules including amino acid residues, as well as a variety of surfactant compositions. A wide variety of other molecules, including uncommon but naturally occurring amino acids, metabolites and catabolites of natural amino acids, substituted amino acids, and amino acid analogs, as well as amino acids in the "D" configuration, are useful in molecules and compositions of the present invention. In addition, "designed" amino acid derivatives, analogs and mimics are also useful in various compounds, compositions and methods of the present invention, as well as polymers including backbone structures composed of non-amide linkages.

[0070] For example, in addition to the L-amino acids, amino acid metabolites such as homoarginine, citrulline, ornithine, and α -aminobutanoic acid are also useful in pulmonary surfactants. Thus, in the various formulas described above, "Charged", Z, or B can comprise homoarginine, citrulline, or ornithine, as well as a variety of other molecules as identified herein. Similarly, J can comprise α -aminobutanoic acid (also known as α -aminobutyric acid), α -aminopentanoic acid, α -aminohexanoic acid, and a variety of other molecules identified herein.

[0071] Further, substituted amino acids which are not generally derived from proteins, but which are known in nature, are useful as disclosed herein, include the following examples: L-canavanine; 1-methyl-L-histidine; 3-methyl-L-histidine; 2-methyl L-histidine; a, ϵ -diaminopimelic acid (L form, meso form, or both); sarcosine; L-ornithine betaine; betaine of histidine (herzynine); L-citrulline; L-phosphoarginine; D-octopine; o-carbamyl-D-serine; γ -aminobutanoic acid; and β -lysine. D-amino acids and D-amino acid analogs, including the following, are also useful in proteins, peptides and compositions of the present invention: D-alanine, D-serine, D-valine, D-leucine, D-isoleucine, D-alloisoleucine, D-phenylalanine, D-glutamic acid, D-proline, and D-allohydroxyproline, to name some examples. The foregoing can also be used in surfactant molecules according to the present invention; particularly preferred for use accordingly are those corresponding to the formula $\{(Charged)_a (Uncharged)_b\}_c (Charged)_d$.

[0072] An extensive variety of amino acids, including metabolites and catabolites thereof, can be incorporated into molecules which display a surfactant activity. For example, molecules such as ornithine, homoarginine, citrulline, and α -aminobutanoic acid are useful components of molecules displaying surfactant activity as described herein. Surfactant molecules described herein can also comprise longer straight-chain molecules; α -aminopentanoic acid and α -aminohexanoic acid are two additional examples of such useful molecules.

[0073] Described herein are a wide variety of modified amino acids, including analogs, metabolites, catabolites, and derivatives, irrespective of the time or location at which modification occurs. In essence, one can place modified amino


acids into three categories: (1) catabolites and metabolites of amino acids; (2) modified amino acids generated via posttranslational modification (e.g., modification of side chains); and (3) modifications made to amino acids via non-metabolic or non-catabolic processes (e.g., the synthesis of modified amino acids or derivatives in the laboratory).

[0074] It is also contemplated that one can readily design side chains of the amino acids of residue units that include longer or shortened side chains by adding or subtracting methylene groups in either linear, branched chain, or hydrocarbon or heterocyclic ring arrangements. The linear and branched chain structures can also contain non-carbon atoms such as S, O, or N. Fatty acids can also be useful constituents of surfactant molecules herein. The designed side chains can terminate with (R') or without (R) charged or polar group appendages.

[0075] In addition, analogs, including molecules resulting from the use of different linkers, are also useful as disclosed herein. Molecules with side chains linked together via linkages other than the amide linkage e.g., molecules containing amino acid side chains or other side chains (R- or R'-) wherein the components are linked via carboxy- or phospho-esters, ethylene, methylene, ketone or ether linkages, to name a few examples, are also useful as disclosed herein. In essence, any amino acid side chain, R or R' group-containing molecule can be useful as disclosed herein, as long as the molecule includes alternating hydrophilic and hydrophobic residues (i.e., component molecules) and displays surfactant activity as described herein.

[0076] Further contemplated are molecules comprising peptide dimers joined by an appropriate linker, e.g., peptide dimers linked by cystine molecules. Such linkers or bridges can thus cross-link different polypeptide chains, dimers, trimers, and the like. Other useful linkers which can be used to connect peptide dimers and/or other peptide multimers include those listed above e.g., carboxy- or phospho-ester, ethylene, methylene, ketone or ether linkages, to name a few examples.

[0077] While it is appreciated that many useful polypeptides disclosed herein comprise naturally-occurring amino acids in the "L" form which are joined via peptide linkages, it should also be understood that molecules including amino acid side chain analogs, non-amide linkages (e.g., differing backbones) can also display a significant surfactant activity and can possess other advantages, as well. For example, if it is desirable to construct a molecule (e.g., for use in a surfactant composition) that is not readily degraded, one may wish to synthesize a polypeptide molecule comprising a series of D-amino acids. Molecules comprising a series of amino acids linked via a "retro" backbone, i.e., a molecule that has internal amide bonds constructed in the reverse direction of carboxyl terminus to amino terminus, are also more difficult to degrade and can thus be useful in various applications, as described herein. For example, the following illustrates an exemplary molecule with a "retro" bond in the backbone:



[0078] In another variation, one may wish to construct a molecule that adopts a more "rigid" conformation; one means of accomplishing this would be to add methyl or other groups to the a carbon atom of the amino acids.

[0079] As noted above, other groups besides a CH₃ group can be added to the a carbon atom, that is, surfactant molecules of the present invention are not limited to those incorporating a CH₃ at the a carbon alone. For example, any of the side chains and molecules described above can be substituted for the indicated CH₃ group at the a carbon component.

[0080] As used herein, the terms "analogs" and "derivatives" of polypeptides and amino acid residues are intended to encompass metabolites and catabolites of amino acids, as well as molecules which include linkages, backbones, side-chains or side-groups which differ from those ordinarily found in what are termed "naturally-occurring" L-form amino acids. (The terms "analog" and "derivative" can also conveniently be used interchangeably herein.) Thus, D-amino acids, molecules which mimic amino acids and amino acids with "designed" side chains (i.e., that can substitute for one or more amino acids in a molecule having surfactant activity) are also encompassed by the terms "analogs" and "derivatives" herein.

[0081] A wide assortment of useful surfactant molecules, including amino acids having one or more extended or substituted R or R' groups, is also contemplated by the present invention. Again, one of skill in the art should appreciate from the disclosures that one can make a variety of modifications to individual amino acids, to the linkages, and/or to the chain itself, which modifications will produce molecules possessing the surfactant activity as described herein.

[0082] In certain methods described herein the pulmonary surfactant may comprise one or more lipids. In which case, the surfactant composition can comprise, for example, from as little as about 0.05 to 100 % weight percent lipid, so long as the resulting composition has surfactant activity. By weight percent is meant the percentage of a compound by weight in a composition by weight. Thus, a composition having 50 weight percent lipid contains, for example, 50 grams lipids

per 100 grams total composition. The term "lipid" as used herein refers to a naturally occurring, synthetic or semi-synthetic (*i.e.*, modified natural) compound which is generally amphipathic. The lipids typically comprise a hydrophilic component and a hydrophobic component. Exemplary lipids include, but are not limited to, phospholipids, fatty acids, fatty alcohols, neutral fats, phosphatides, oils, glycolipids, surface-active agents (surfactants), aliphatic alcohols, waxes, terpenes and steroids. The phrase semi-synthetic (or modified natural) denotes a natural compound that has been chemically modified in some fashion. Preferably, the lipids of are fatty acids, alcohols, esters and ethers thereof, fatty amines, or combinations thereof.

[0083] Examples of phospholipids include native and/or synthetic phospholipids. Phospholipids that can be used include, but are not limited to, phosphatidylcholines, phosphatidylglycerols, phosphatidylethanolamines, phosphatidylserines, phosphatidic acids, phosphatidylinositols, sphingolipids, diacylglycerides, cardiolipin, ceramides, cerebroside and the like. Exemplary phospholipids include, but are not limited to, dipalmitoyl phosphatidylcholine (DPPC), dilauryl phosphatidylcholine (DLPC) (C12:0), dimyristoyl phosphatidylcholine (DMPC) (C14:0), distearoyl phosphatidylcholine (DSPC), diphytanoyl phosphatidylcholine, nonadecanoyl phosphatidylcholine, arachidoyl phosphatidylcholine, dioleoyl phosphatidylcholine (DOPC) (C18:1), dipalmitoleoyl phosphatidylcholine (C16:1), linoleoyl phosphatidylcholine (C18:2), myristoyl palmitoyl phosphatidylcholine (MPPC), steroyl myristoyl phosphatidylcholine (SMPC), steroyl palmitoyl phosphatidylcholine (SPPC), palmitoyl oleoyl phosphatidylcholine (POPC), palmitoyl palmitooleoyl phosphatidylcholine (PPoPC), dipalmitoyl phosphatidylethanolamine (DPPE), palmitoyl oleoyl phosphatidylethanolamine (POPE), dioleoyl phosphatidylethanolamine (DOPE), dimyristoyl phosphatidylethanolamine (DMPE), distearoyl phosphatidylethanolamine (DSPE), dioleoyl phosphatidylglycerol (DOPG), palmitoyl oleoyl phosphatidylglycerol (POPG), dipalmitoyl phosphatidylglycerol (DPPG), dimyristoyl phosphatidylglycerol (DMPG), distearoyl phosphatidylglycerol (DSPG), dimyristoyl phosphatidylserine (DMPS), distearoyl phosphatidylserine (DSPS), palmitoyl oleoyl phosphatidylserine (POPS), soybean lecithin, egg yolk lecithin, sphingomyelin, phosphatidylinositols, diphosphatidylglycerol, phosphatidylethanolamine, phosphatidic acids, and egg phosphatidylcholine (EPC).

[0084] Examples of fatty acids and fatty alcohols include, but are not limited to, sterols, palmitic acid, cetyl alcohol, lauric acid, myristic acid, stearic acid, phytanic acid, dipalmitic acid, and the like. Preferably, the fatty acid is palmitic acid and preferably the fatty alcohol is cetyl alcohol.

[0085] Examples of fatty acid esters include, but are not limited to, methyl palmitate, ethyl palmitate, isopropyl palmitate, cholesteryl palmitate, palmityl palmitate sodium palmitate, potassium palmitate, tripalmitin, and the like.

[0086] An example of a semi-synthetic or modified natural lipid is any one of the lipids described above which has been chemically modified. The chemical modification can include a number of modifications; however, a preferred modification is the conjugation of one or more polyethylene glycol (PEG) groups to desired portions of the lipid. Polyethylene glycol (PEG) has been widely used in biomaterials, biotechnology and medicine primarily because PEG is a biocompatible, nontoxic, nonimmunogenic and water-soluble polymer. Zhao and Harris, ACS Symposium Series 680: 458-72, 1997. In the area of drug delivery, PEG derivatives have been widely used in covalent attachment (*i.e.*, "PEGylation") to proteins to reduce immunogenicity, proteolysis and kidney clearance and to enhance solubility. Zalipsky, Adv. Drug Del. Rev. 16: 157-82, 1995.

[0087] Lipids that have been conjugated with PEG are referred to herein as "PEG-lipids." Preferably, when PEG-lipids are used, they are present in alcohols and/or aldehydes.

[0088] The pulmonary surfactant can comprise other excipients, including, but not limited to, various sugars such as dextrose, fructose, lactose, maltose, mannitol, sucrose, sorbitol, trehalose, and the like, surfactants such as, for example, polysorbate-80, polysorbate-20, sorbitan trioleate, tyloxapol and the like, polymers such as PEG, dextran and the like, salts such as NaCl, CaCl₂ and the like, alcohols, such as cetyl alcohol, and buffers.

[0089] Exemplary surfactant compositions can be prepared using methods known in the art. For example, an exemplary surfactant composition comprising lipids and polypeptides can be prepared by admixing a solution of a surfactant polypeptide with a suspension of liposomes, or by admixing the surfactant polypeptide with a suspension of liposomes, or by admixing the surfactant polypeptide and phospholipids directly in the presence of organic solvent.

[0090] Preferably, the pulmonary surfactant comprises phospholipids and free fatty acids or fatty alcohols, *e.g.*, DPPC (dipalmitoyl phosphatidylcholine), POPG (palmitoyl-oleyl phosphatidylglycerol) and palmitic acid (PA). See, for example, U.S. Patent No 5,789,381.

[0091] In certain preferred embodiments, the pulmonary surfactant is lucinactant or another pulmonary surfactant formulation comprising the synthetic surfactant protein KLLLLKLLLLKLLLLKLLLL (KL4; SEQ ID NO:1). Lucinactant, is a combination of DPPC, POPG, palmitic acid (PA) and the KL4 peptide (weight ratio of approximately 7.5 : 2.5 : 1.35 : 0.267). In certain embodiments, the drug product is formulated at concentrations of, for example, 10, 20, and 30 mg/ml of phospholipid content. In certain other embodiments, the drug product is formulated at greater concentrations, *e.g.*, 40, 60, 90, 120 or more mg/ml phospholipid content, with concomitant increases in KL4 concentration.

[0092] Any pulmonary surfactant currently in use, or hereafter developed for use in respiratory distress system and other pulmonary conditions, is suitable for use in the present invention. These include naturally occurring and synthetic pulmonary surfactant. Synthetic PS, as used herein, refers to both protein-free lung surfactants and pulmonary surfactants

comprising synthetic peptides, including peptide mimetics of naturally occurring surfactant protein. Current PS products include, but are not limited to, lucinactant (Surfaxin[®], Discovery Laboratories, Inc., Warrington, PA), bovine lipid surfactant (BLES[®], BLES Biochemicals, Inc. London, Ont), calfactant (Infasurf[®], Forest Pharmaceuticals, St. Louis, MO), natural bovine surfactant (Alveofact[®], Thomae, Germany), bovine surfactant (Surfactant TA[®], Tokyo Tanabe, Japan), poractant alfa (Curosurf[®], Chiesi Farmaceutici SpA, Parma, Italy), pumactant (Alec[®], Britannia Pharmaceuticals, UK), beractant (Survanta[®], Abbott Laboratories, Inc., Abbott Park, IL) and colfosceril palmitate (Exosurf[®], GlaxoSmithKline, plc, Middlesex, U.K.). In a preferred embodiment, the PS is lucinactant or another PS formulation comprising the synthetic surfactant protein KLLLLKLLLLKLLLLKLLLL (KL4; SEQ ID NO:1).

[0093] The treatment regimens described herein can be combined with other respiratory therapies. In certain embodiments, PS administration is performed on infants who are intubated and maintained on ventilation, either conventional ventilation or high frequency ventilation, for a period of time or for the entire duration of the PS treatment.

[0094] Alternative modes of administration can be utilized, as well as alternative PS formulations. For example, PS can be formulated for aerosolization (nebulization) and administered *via* nasal CPAP, nasal or naso-pharyngeal prongs in combination with low-flow oxygen, or *via* face mask or oxygen hood.

[0095] Aerosolized pulmonary surfactant can be administered as provided in copending U.S. application, 11/130,783, filed May 17, 2005. Administration can be in conjunction with another noninvasive pulmonary respiratory therapy involving the administration of positive airway pressure. The term "noninvasive pulmonary respiratory therapy" refers to respiratory therapy which does not use mechanical ventilation and can include CPAP, bilevel positive airway pressure (BiPAP), synchronized intermittent mandatory ventilation (SIMV), and the like. The employment of such therapies involves the use of various respiratory gases, as would be appreciated by the skilled artisan. Respiratory gases used for noninvasive pulmonary respiratory therapy are sometimes referred to herein as "CPAP gas," "CPAP air," "nCPAP", "ventilation gas," "ventilation air," or simply "air." However, those terms are intended to include any type of gas normally used for noninvasive pulmonary respiratory therapy, including but not limited to gases and gaseous combinations listed above for use as the conditioning gas. In certain embodiments, the gas used for noninvasive pulmonary respiratory therapy is the same as the conditioning gas. In other embodiments, the respective gases are different from one another.

[0096] In certain embodiments, the pulmonary delivery methods of this invention are employed in conjunction with CPAP. It has been shown that use of CPAP allows for an increase in functional residual capacity and improved oxygenation. The larynx is dilated and supraglottic airway resistance is normal. There is also an improvement of the synchrony of respiratory thoracoabdominal movements and enhanced Hering-Breuer inflation reflex following airway occlusion. CPAP has been shown to be useful in treating various conditions such as sleep apnea, snoring, ARDS, IRDS, and the like.

[0097] In order to effect administration of CPAP, a pressure source and a delivery device or delivery apparatus are required. CPAP-producing airflow is typically generated in the vicinity of the nasal airways by converting kinetic energy from a jet of fresh humidified gas into a positive airway pressure. A continuous flow rate of breathing gas of about 5 to about 12 liters/minute generates a corresponding CPAP of about 2 to about 10 cm H₂O. Various modifications can be applied to the CPAP system which include sensors that can individualize the amount of pressure based on the patient's need

[0098] Typically, flow rates and pressures suitable for achieving CPAP are based upon the characteristics of the patient being treated. Suitable flow rates and pressures can be readily calculated by the attending clinician. The present invention encompasses the use of a variety of flow rates for the ventilating gas, including low, moderate and high flow rates. Alternatively, the aerosol can be supplied without added positive pressure, *i.e.*, without CPAP as a simultaneous respiratory therapy.

[0099] Preferably, the CPAP-generating air flow being delivered to the patient has a moisture level which will prevent unacceptable levels of drying of the lungs and airways. Thus, the CPAP-generating air is often humidified by bubbling through a hydrator, or the like to achieve a relative humidity of preferably greater than about 70%. More preferably, the humidity is greater than about 85% and still more preferably 98%.

[0100] A suitable source of CPAP-inducing airflow is the underwater tube CPAP (underwater expiratory resistance) unit. This is commonly referred to as a bubble CPAP.

[0101] Another preferred source of pressure is an expiratory flow valve that uses variable resistance valves on the expiratory limb of CPAP circuits. This is typically accomplished via a ventilator.

[0102] Another preferred source is the Infant Flow Driver or "IFD" (Electro Medical Equipment, Ltd., Brighton, Sussex, UK). IFD generates pressure at the nasal level and employs a conventional flow source and a manometer to generate a high pressure supply jet capable of producing a CPAP effect. It is suggested in the literature that the direction of the high pressure supply jet responds to pressures exerted in the nasal cavity by the patient's efforts and this reduces variations in air pressure during the inspiration cycle.

[0103] Other CPAP systems including those that contain similar features to systems just discussed are also contemplated herein.

[0104] The aerosol stream generated in accordance with aerosolized delivery is preferably delivered to the patient via a nasal delivery device which can involve, for example masks, single nasal prongs, binasal prongs, nasopharyngeal

prongs, nasal cannulae and the like. The delivery device is chosen so as to minimize trauma, maintain a seal to avoid waste of aerosol, and minimize the work the patient must perform to breathe.

[0105] When used as an aerosol preparation, the surfactant composition can be supplied in finely divided form, optionally in combination with a suitable propellant. Useful propellants are typically gases at ambient conditions and are condensed under pressure including, for example, lower alkanes and fluorinated alkanes, such as freon. Where the surfactant composition is delivered as an aerosol, the aerosol can be packaged in a suitable container under pressure.

[0106] Suitable dosage of the surfactant, whether aerosolized or delivered as a liquid or dry powder will be dependent on the patient's age and severity of the disorder and will be readily ascertainable by the attending clinician. The actual dosage of pulmonary surfactant will of course vary according to factors such as the extent of exposure and particular status of the subject (e.g., the subject's age, size, fitness, extent of symptoms, susceptibility factors, and the like). By "effective dose" herein is meant a dose that produces effects for which it is administered.

[0107] When the surfactant is supplied to prematurely born infants as a liquid, an aliquot of the surfactant composition is delivered, preferably by intratracheal instillation, to provide an effective dose of surfactant in the lungs of the treated patient. Preferably, a single surfactant dose ranges from, for example, about 20 to about 300 mg total phospholipid (TPL)/kg, more preferably from about 60 to about 175 mg TPL/kg. It being understood, of course, that the exact dose of surfactant will depend upon factors such as the age and condition of the patient, the severity of the condition being treated, and other factors well within the skill of the attending clinician. When the surfactant composition is delivered as an aerosol, such as disclosed in co-pending U.S. Application No. 11/130,783, filed May 17, 2005, the effective dose of lung surfactant can be, for example, from about 1 mg TPL/kg surfactant to about 1000 mg TPL/kg surfactant, preferably from about 2 mg TPL/kg surfactant to about 175 mg TPL/kg surfactant. When the surfactant composition is delivered as a dry powder formulation, the effective dose of lung surfactant can be, for example, from about 1 mg TPL/kg surfactant to about 1000 mg or more TPL/kg surfactant, preferably from about 2 mg TPL/kg surfactant to about 175 mg TPL/kg surfactant. Other methods of delivery include lavage, lung wash, and the like. When so employed dose ranges are well within the skill of one in the art.

[0108] The frequency of dosing can vary, but typically is once every 2-3 days. However, the patient can be dosed more frequently, e.g., every 6-8 hours, twice daily, or once daily, or less frequently, e.g., twice weekly or even once weekly. Alternatively, the patient can be dosed more frequently early in the treatment regimen, and with decreasing frequency later in the treatment regimen, e.g., once every other day for one week, followed by twice weekly until the end of the treatment period. Depending on the dosage form, e.g., aerosol or dry powder as compared with liquid instillate, the patient can be dosed continuously for part or all of the treatment period.

[0109] In use according to the invention, the infant can be treated with other therapeutic, prophylactic or complementary agents, such as steroids, nitric oxide, antioxidants or reactive oxygen scavengers, bronchodilators, diuretics, antimicrobial or anti-infective agents, anti-hypertensive agents, or anti-inflammatory agents (e.g., PLA₂ inhibitors, protease or elastase inhibitors, PDE-4 inhibitors, to name a few), as would be appreciated by one of skill in the art. Such treatment can include concomitant administration of the PS with other therapeutic, prophylactic or complementary agents. Concomitant administration can involve concurrent (i.e. at the same time), prior, or subsequent administration of the complementary agent with respect to the administration of the pulmonary surfactant. When administered in combination, each component can be administered at the same time or sequentially in any order at different points in time. Thus, each component can be administered separately but sufficiently closely in time so as to provide the desired therapeutic effect. Concomitant administration of a PS with other therapeutic, prophylactic or complementary agents means administration of the PS and other agents at such time that both will have a therapeutic effect. A person of ordinary skill in the art, would have no difficulty determining the appropriate timing, sequence and dosages of administration for particular drugs of the present invention.

[0110] The following examples are provided to describe the invention in greater detail. They are intended to illustrate, not to limit, the invention.

EXEMPLARY EMBODIMENTS

EXAMPLE 1: Protocol for Administration of Lucinactant to Premature Infants at Risk for Bronchopulmonary Dysplasia

[0111] This example provides a protocol for administration of lucinactant to very low birthweight (VLBW) premature infants who have been intubated and received surfactants for the prevention or treatment of respiratory distress syndrome (RDS). The protocol has been used in a clinical trial to assess the safety and efficacy of lucinactant in VLBW infants at risk for developing BPD.

[0112] Two dosing regimens of lucinactant (90 mg/kg birthweight or 175 mg/kg birthweight) were utilized. Infants born with a birthweight 600 to 900 grams and who remained intubated at day of life (DOL) 3 in anticipation of worsening lung disease were treated. Treatment was administered on days 3, 5, 7, 10 and 14, if the infant remained intubated. It should

be noted that Day 1 (or DOL 1) is the day on which Time 0 (time of birth) occurs and DOL 2, 3, and so on, begin at 00:00 (midnight) each subsequent day following Time 0.

[0113] Lucinactant was administered in accordance with manufacturer's instructions. Briefly, administration of lucinactant is by syringe typically attached to an end-hole catheter passed through a Bodai valve or equivalent so that that the catheter tip terminates at the distal end of the endotracheal tube. The infant is placed in a head-up left lateral decubitus position and slowly administered half the syringe volume. After allowing a brief recovery period, the infant is placed in the right lateral decubitus position and the remaining contents of the syringe are administered. The dose was administered in quarters (alternating left-right decubitus positioning) if determined to be necessary.

[0114] Efficacy of the treatment protocol was determined by making the following assessments: (1) proportion of infants remaining on mechanical ventilation or oxygen over time; (2) incidence of death or BD at 28 days and 36 weeks post-menstrual age (post-conception); (3) AUC day 3-28 fraction of inspired oxygen (FiO₂) and mean airway pressure; (4) pulmonary compliance as assessed by ventilator derived pressure-volume loops pre, 6 and 24 hours post dose; and (5) surrogate measures of efficacy including assessment of surface tension lowering properties of tracheal aspirates and concentration of growth factors and inflammatory mediators in tracheal aspirates collected as a part of routine neonatal intensive care.

EXAMPLE 2: Protocol for Administration of Lucinactant Delivered as an Aerosol Via nCPAP for the Prevention of RDS in Premature Infants

[0115] This example provides a protocol for administration of lucinactant as an aerosol via nCPA to premature infants. This protocol has been used in a pilot phase 2, open study to evaluate the feasibility, safety, and tolerability of lucinactant delivered as an aerosol via nCPAP for the prevention of RDS in premature neonates.

[0116] Lucinactant was administered using an approved vibrating machine device (Aeroneb-Pro® or equivalent) and delivered via nCPAP within 30 minutes of birth. In a first treatment group, 20 mg/ml of lucinactant was administered within the first 30 minutes of life continually over three hours. Up to three retreatments were permitted over a 48 hour period, with each treatment separated by at least 3 hours (from end of previous treatment). In a second treatment group, 20 mg/ml of lucinactant was administered within the first 30 minutes of life continually over three hours. Up to three retreatments were permitted over a 48 hour period, with each treatment separated by at least one hours (from end of previous treatment).

[0117] Qualified premature neonates were sequentially enrolled and stratified by gestational age (stratum 1: 30-32 completed weeks followed by stratum 2: 28-29 completed weeks) into the first treatment regimen group (3 hours on aerosol treatment / 3 hours off aerosol treatment) and received aerosolized lucinactant via nCPAP. Safety and tolerability data were evaluated after enrollment of every five neonates. Once enrolment in the first treatment group was completed, neonates were sequentially enrolled and stratified into the second treatment group (3 hours on aerosol treatment / 1 hours off aerosol treatment). All neonates enrolled were evaluated through 28 days of age.

[0118] All 17 neonates experienced at least 1 adverse effect regardless of relationship. A total of 83 adverse effects were reported; of these adverse effects, 8 (47.1%) were considered related to study drug, 2 (11.8%) were considered related to the device, and 6 (35.3%) were considered related to both study drug and the device. The most common adverse effects, regardless of relationship to study treatment were apnea (16/17, 94.1%), oxygen saturation decreased (12/17, 70.6%), patent ductus arteriosus (6/17, 35.3%), hypocalcemia (5/17, 29.4%). A total of 5 serious adverse effects were reported during the course of the study. Two of the serious adverse effects were considered related to the device. No deaths were occurred during this study.

[0119] The study showed that it is feasible to deliver lucinactant via nCPAP and the treatment was generally safe and well tolerated. Key observations included that 15 of the 17 neonates had no evidence of HPD at day 28.

[0120] From the foregoing description, various modifications and changes in the compositions and methods will occur to those skilled in the art.

Claims

1. Use of a pulmonary surfactant in the preparation of a medicament for the prevention of bronchopulmonary dysplasia in an infant, wherein the infant has been treated with pulmonary surfactant for respiratory distress syndrome in a dosage regime which comprises administration of the pulmonary surfactant for bronchopulmonary dysplasia after the treatment for respiratory distress syndrome with pulmonary surfactant has been concluded, and wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant for bronchopulmonary dysplasia continued through at least day 10 of the life of the infant.

2. Use as claimed in claim 1 wherein the dosage regime comprises administration of an effective amount of the

pulmonary surfactant initiated at the next dosage interval following completion of the treatment for respiratory distress syndrome with pulmonary surfactant.

- 5 3. Use as claimed in claim 1 or claim 2 wherein the dosing regime comprises administration of an effective amount of the pulmonary surfactant initiated at day 3 of the life of the infant.
4. Use as claimed in claim 1 or claim 2 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant continued through at least day 14 of life of the infant.
- 10 5. Use as claimed in claim 1 or claim 2 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant continued through at least day 18 of life of the infant.
6. Use as claimed in claims 1 or 2 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant initiated anytime from day 3 to day 10 of life of the infant.
- 15 7. Use of a pulmonary surfactant in the preparation of a medicament for the prevention of bronchopulmonary dysplasia in an infant requiring respiratory support and who is at risk of developing bronchopulmonary dysplasia and wherein the pulmonary surfactant is for use in a dosage regime which comprises administration of an effective amount of the pulmonary surfactant through at least day 10 of the life of the infant.
- 20 8. Use as claimed in claim 7 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant initiated at or after day 1 of life of the infant and continued through at least day 28 of life of the infant.
- 25 9. Use as claimed in claim 7 wherein administration of the pulmonary surfactant initiated at or after day 2 of life of the infant.
- 30 10. Use as claimed in claim 7 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant initiated at or after day 3 of life of the infant.
- 35 11. Use as claimed in claim 7 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant continued through at least day 14 of life of the infant.
12. Use as claimed in claim 7 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant continued through at least day 18 of life of the infant.
- 40 13. Use as claimed in claim 7 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant initiated anytime from day 3 to day 18 of life of the infant.
14. Use as claimed in any one of claims 7 to 13 wherein the infant does not exhibit respiratory distress syndrome.
15. Use as claimed in any one of claims 7 to 13 wherein the infant exhibits respiratory distress syndrome.
- 45 16. Use as claimed in any one of the preceding claims wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant by endotracheal administration.
17. Use as claimed in any one of claims 1 to 15 wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant by inhalation.
- 50 18. Use as claimed in any one of the preceding claims wherein the dosage regime comprises administration of an effective amount of the pulmonary surfactant accompanied by another respiratory therapy.
19. Use as claimed in claim 18 wherein the other respiratory therapy is conventional ventilation, high frequency ventilation or continuous positive airway pressure.
- 55 20. Use as claimed in claim 18 wherein the other respiratory therapy is administration of one or more therapeutic agents selected from the group; nitric oxide, steroids, antioxidants, vitamins, vitamin derivatives, reactive oxygen scavengers, bronchodilators, diuretics, antimicrobial agents, anti-infective agents, anti-hypertensive agents or anti-inflam-

matory agents.


21. Use as claimed in any one of the preceding claims wherein the pulmonary surfactant is a synthetic pulmonary surfactant.

22. Use as claimed in claim 21 wherein the synthetic pulmonary surfactant comprises a peptide having the amino acid sequence:

KLLLLKLLLLKLLLLKLLLLK (SEQ ID NO:1)

23. Use as claimed in claim 21 or claim 22 wherein the synthetic pulmonary surfactant comprises a polypeptide mimetic of SP-B.

24. Use as claimed in any one of claims 21 to 23 wherein the synthetic pulmonary surfactant comprises a polypeptide having at least 10 amino acid residues and comprising alternating hydrophobic and hydrophilic amino acid residue regions of the formula:



wherein Z and U are amino acid residues such that at each occurrence are independently selected;

Z is a hydrophilic amino acid residue selected from R, D, E and K;

U is a hydrophobic amino acid residue selected from V, I, L, C, Y and F;

a is between 1 and 5;

b is between 3 and 20;

c is between 1 and 10; and

d is between 0 and 3.

Patentansprüche

1. Eine Verwendung eines Lungensurfactants bei der Herstellung eines Medikaments für die Verhinderung von bronchopulmonaler Dysplasie bei einem Säugling, wobei der Säugling mit Lungensurfactant gegen Atemnotsyndrom behandelt worden ist, in einem Dosierungsschema, das die Verabreichung des Lungensurfactants gegen bronchopulmonale Dysplasie nach Abschluss der Behandlung mit Lungensurfactant gegen Atemnotsyndrom beinhaltet, und wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants gegen bronchopulmonale Dysplasie beinhaltet, die bis mindestens Tag 10 des Lebens des Säuglings fortgesetzt wird.

2. Verwendung gemäß Anspruch 1, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die beim nächsten Dosierungsintervall im Anschluss an die Vollendung der Behandlung mit Lungensurfactant gegen Atemnotsyndrom initiiert wird.

3. Verwendung gemäß Anspruch 1 oder Anspruch 2, wobei das Dosierschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die an Tag 3 des Lebens des Säuglings initiiert wird.

4. Verwendung gemäß Anspruch 1 oder Anspruch 2, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die bis mindestens Tag 14 des Lebens des Säuglings fortgesetzt wird.

5. Verwendung gemäß Anspruch 1 oder Anspruch 2, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die bis mindestens Tag 18 des Lebens des Säuglings fortgesetzt wird.

6. Verwendung gemäß Anspruch 1 oder Anspruch 2, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die zu einem beliebigen Zeitpunkt von Tag 3 bis Tag 10 des Lebens des Säuglings initiiert wird.


7. Eine Verwendung eines Lungensurfactants bei der Herstellung eines Medikaments für die Verhinderung von bronchopulmonaler Dysplasie bei einem Säugling, der Atemhilfe benötigt und bei dem das Risiko besteht, bronchopul-

EP 1 841 458 B9

monale Dysplasie zu entwickeln, und wobei das Lungensurfactant zur Verwendung in einem Dosierungsschema vorgesehen ist, das die Verabreichung einer wirksamen Menge des Lungensurfactants bis mindestens Tag 10 des Lebens des Säuglings beinhaltet.

- 5 **8.** Verwendung gemäß Anspruch 7, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die an oder nach Tag 1 des Lebens des Säuglings initiiert und bis mindestens Tag 28 des Lebens des Säuglings fortgesetzt wird.
- 10 **9.** Verwendung gemäß Anspruch 7, wobei die Verabreichung des Lungensurfactants an oder nach Tag 2 des Lebens des Säuglings initiiert wird.
- 10.** Verwendung gemäß Anspruch 7, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die an oder nach Tag 3 des Lebens des Säuglings initiiert wird.
- 15 **11.** Verwendung gemäß Anspruch 7, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die bis mindestens Tag 14 des Lebens des Säuglings fortgesetzt wird.
- 12.** Verwendung gemäß Anspruch 7, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die bis mindestens Tag 18 des Lebens des Säuglings fortgesetzt wird.
- 20 **13.** Verwendung gemäß Anspruch 7, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die zu einem beliebigen Zeitpunkt von Tag 3 bis Tag 18 des Lebens des Säuglings initiiert wird.
- 25 **14.** Verwendung gemäß einem der Ansprüche 7 bis 13, wobei der Säugling kein Atemnotsyndrom zeigt.
- 15.** Verwendung gemäß einem der Ansprüche 7 bis 13, wobei der Säugling Atemnotsyndrom zeigt.
- 30 **16.** Verwendung gemäß einem der vorhergehenden Ansprüche, wobei das Dosierungsschema die Verabreichung einer wirksamen Menge des Lungensurfactants durch endotracheale Verabreichung beinhaltet.
- 17.** Verwendung gemäß einem der Ansprüche 1 bis 15, wobei das Dosierungsschema die Verabreichung einer wirksamen Menge des Lungensurfactants durch Inhalation beinhaltet.
- 35 **18.** Verwendung gemäß einem der vorhergehenden Ansprüche, wobei das Dosierungsschema eine Verabreichung einer wirksamen Menge des Lungensurfactants beinhaltet, die von einer anderen Atemtherapie begleitet wird.
- 19.** Verwendung gemäß Anspruch 18, wobei die andere Atemtherapie eine herkömmliche Beatmung, eine Hochfrequenzbeatmung oder eine CPAP-Atmung ist.
- 40 **20.** Verwendung gemäß Anspruch 18, wobei die andere Atemtherapie die Verabreichung einer oder mehrerer therapeutischer Mittel, ausgewählt aus der folgenden Gruppe, ist: Stickoxid, Steroide, Antioxidationsmittel, Vitamine, Vitaminderivate, Fänger reaktiven Sauerstoffs, Bronchodilatoren, Diuretika, antimikrobielle Mittel, infektionsverhindernde Mittel, Antihypertonika oder entzündungshemmende Mittel.
- 45 **21.** Verwendung gemäß einem der vorhergehenden Ansprüche, wobei das Lungensurfactant ein synthetisches Lungensurfactant ist.
- 50 **22.** Verwendung gemäß Anspruch 21, wobei das synthetische Lungensurfactant ein Peptid mit der folgenden Aminosäuresequenz beinhaltet:
- KLLLLKLLLLKLLLLKLLLLK (SEQ ID NO: 1)
- 55 **23.** Verwendung gemäß Anspruch 21 oder Anspruch 22, wobei das synthetische Lungensurfactant ein Polypeptid-Mimetikum von SP-B beinhaltet.
- 24.** Verwendung gemäß einem der Ansprüche 21 bis 23, wobei das synthetische Lungensurfactant ein Polypeptid beinhaltet, das mindestens 10 Aminosäurereste aufweist und alternierende hydrophobe und hydrophile Aminosäu-

reestregionen der folgenden Formel beinhaltet:



- 5 wobei Z und U Aminosäurereste sind, so dass jedes Auftreten unabhängig ausgewählt ist;
 Z ein hydrophiler Aminosäurerest ist, ausgewählt aus R, D, E und K;
 U ein hydrophober Aminosäurerest ist, ausgewählt aus V, I, L, C, Y und F; a zwischen 1 und 5 beträgt;
 b zwischen 3 und 20 beträgt;
 c zwischen 1 und 10 beträgt; und
 10 d zwischen 0 und 3 beträgt.

Revendications

- 15 1. Utilisation d'un surfactant pulmonaire dans la préparation d'un médicament pour la prévention d'une dysplasie broncho-pulmonaire chez un nourrisson, dans laquelle le nourrisson a été traité avec un surfactant pulmonaire pour un syndrome de détresse respiratoire dans un régime de dosage qui comprend l'administration du surfactant pulmonaire pour une dysplasie broncho-pulmonaire une fois que le traitement pour le syndrome de détresse respiratoire avec le surfactant pulmonaire a été effectué, et dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire pour la dysplasie broncho-pulmonaire qui est continuée au moins jusqu'au jour 10 de la vie du nourrisson.
- 20 2. Utilisation telle que revendiquée dans la revendication 1 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire initiée à l'intervalle de dosage suivant suite à la fin du traitement pour le syndrome de détresse respiratoire avec le surfactant pulmonaire.
- 25 3. Utilisation telle que revendiquée dans la revendication 1 ou la revendication 2 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire initiée au jour 3 de la vie du nourrisson.
- 30 4. Utilisation telle que revendiquée dans la revendication 1 ou la revendication 2 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire qui est continuée au moins jusqu'au jour 14 de la vie du nourrisson.
- 35 5. Utilisation telle que revendiquée dans la revendication 1 ou la revendication 2 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire qui est continuée au moins jusqu'au jour 18 de la vie du nourrisson.
- 40 6. Utilisation telle que revendiquée dans les revendications 1 ou 2 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire initiée à tout moment du jour 3 au jour 10 de la vie du nourrisson.
- 45 7. Utilisation d'un surfactant pulmonaire dans la préparation d'un médicament pour la prévention d'une dysplasie broncho-pulmonaire chez un nourrisson nécessitant une assistance respiratoire et qui risque de développer une dysplasie broncho-pulmonaire et dans laquelle le surfactant pulmonaire est destiné à être utilisé dans un régime de dosage qui comprend l'administration d'une quantité efficace du surfactant pulmonaire au moins jusqu'au jour 10 de la vie du nourrisson.
- 50 8. Utilisation telle que revendiquée dans la revendication 7 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire initiée au jour 1 ou après le jour 1 de la vie du nourrisson et qui est continuée au moins jusqu'au jour 28 de la vie du nourrisson.
9. Utilisation telle que revendiquée dans la revendication 7 dans laquelle l'administration du surfactant pulmonaire est initiée au jour 2 ou après le jour 2 de la vie du nourrisson.
- 55 10. Utilisation telle que revendiquée dans la revendication 7 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire initiée au jour 3 ou après le jour 3 de la vie du nourrisson.
11. Utilisation telle que revendiquée dans la revendication 7 dans laquelle le régime de dosage comprend l'administration

EP 1 841 458 B9

d'une quantité efficace du surfactant pulmonaire qui est continuée au moins jusqu'au jour 14 de la vie du nourrisson.

5 12. Utilisation telle que revendiquée dans la revendication 7 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire qui est continuée au moins jusqu'au jour 18 de la vie du nourrisson.

13. Utilisation telle que revendiquée dans la revendication 7 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire initiée à tout moment du jour 3 au jour 18 de la vie du nourrisson.

10 14. Utilisation telle que revendiquée dans l'une quelconque des revendications 7 à 13 dans laquelle le nourrisson ne présente pas de syndrome de détresse respiratoire.

15 15. Utilisation telle que revendiquée dans l'une quelconque des revendications 7 à 13 dans laquelle le nourrisson présente un syndrome de détresse respiratoire.

16. Utilisation telle que revendiquée dans l'une quelconque des revendications précédentes dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire par administration endotrachéale.

17. Utilisation telle que revendiquée dans l'une quelconque des revendications 1 à 15 dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire par inhalation.

18. Utilisation telle que revendiquée dans une quelconque des revendications précédentes dans laquelle le régime de dosage comprend l'administration d'une quantité efficace du surfactant pulmonaire accompagnée par une autre thérapie respiratoire.

19. Utilisation telle que revendiquée dans la revendication 18 dans laquelle l'autre thérapie respiratoire est une ventilation traditionnelle, une ventilation à haute fréquence ou une ventilation spontanée en pression positive continue.

20. Utilisation telle que revendiquée dans la revendication 18 dans laquelle l'autre thérapie respiratoire est une administration d'un ou plusieurs agents thérapeutiques sélectionnés dans le groupe comprenant : l'oxyde nitrique, les stéroïdes, les antioxydants, les vitamines, les dérivés de vitamine, les capteurs d'oxygène réactif, les bronchodilatateurs, les diurétiques, les agents antimicrobiens, les agents anti-infectieux, les agents anti-hypertensifs ou les agents anti-inflammatoires.


21. Utilisation telle que revendiquée dans une quelconque des revendications précédentes dans laquelle le surfactant pulmonaire est un surfactant pulmonaire synthétique.

22. Utilisation telle que revendiquée dans la revendication 21 dans laquelle le surfactant pulmonaire synthétique comprend un peptide ayant la séquence d'acides aminés :

40 KLLLLKLLLLKLLLLKLLLLK (SEQ ID NO :1)

23. Utilisation telle que revendiquée dans la revendication 21 ou la revendication 22 dans laquelle le surfactant pulmonaire synthétique comprend un mimétique de polypeptide de SP-B.

45 24. Utilisation telle que revendiquée dans l'une quelconques des revendications 21 à 23 dans laquelle le surfactant pulmonaire synthétique comprend un polypeptide ayant au moins 10 résidus d'acide aminé et comprenant des régions de résidu d'acide aminé hydrophobe et hydrophile en alternance de la formule :



50 dans laquelle Z et U sont des résidus d'acide aminé de telle sorte qu'à chaque occurrence ils sont indépendamment sélectionnés ;

Z est un résidu d'acide aminé hydrophile sélectionné parmi R, D, E et K ;

U est un résidu d'acide aminé hydrophobe sélectionné parmi V, I, L, C, Y et F ; a est compris entre 1 et 5 ;

55 b est compris entre 3 et 20 ;

c est compris entre 1 et 10 ; et

d est compris entre 0 et 3.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 60641805 B [0001]
- WO 9719701 A [0008]
- WO 02062289 A [0009]
- US 5874406 A [0044]
- US 5840527 A [0044]
- US 4918161 A [0044]
- US 5827825 A [0044]
- US 6660833 A [0044]
- US 5006343 A [0044]
- US 5455227 A [0044]
- US 5223481 A [0044]
- US 5753621 A [0044]
- US 5891844 A [0044]
- US 4861756 A [0044]
- US 5272252 A [0044]
- US 502495 A [0044]
- US 5238920 A [0044]
- US 5302481 A [0044]
- US 6022955 A [0044]
- US 6013619 A [0044]
- WO 8603408 A [0044]
- WO 08904326 A [0044]
- US 5260273 A [0046]
- US 5164369 A [0046]
- US 5407914 A [0046]
- US 6613734 B [0046]
- US 5789381 A [0090]
- US 13078305 A [0095] [0107]

Non-patent literature cited in the description

- *Ped. Pulmon.*, 1992, vol. 14, 4-9 [0007]
- **COLIGAN et al.** Current Protocols in Immunology. Wiley Interscience, 1991 [0024]
- **MERRIFIELD.** *J. Am. Chem. Soc.*, 1962, vol. 85, 2149 [0024]
- **STEWART ; YOUNG.** *Solid Phase Peptides Synthesis*, 1969, 27-62 [0024]
- **JOBE ; BANCALARI.** *Am J Respir Crit Care Med*, 2001, vol. 163, 1723-1729 [0027]
- **KING et al.** *Am. J. Physiol.*, 1972, vol. 223, 715-726 [0035]
- **ENHORNING.** *J. Appl. Physiol.*, 1977, vol. 43, 198-203 [0035]
- **ROBERTSON.** *Lung*, 1980, vol. 158, 57-68 [0035]
- **REVAK et al.** *Am. Rev. Respir. Dis.*, 1986, vol. 134, 1258-1265 [0035]
- **HOPP et al.** *Proc. Natl. Acad. Sci.*, 1981, vol. 78, 3824-3829 [0047]
- **GLASSER et al.** *Proc. Natl. Acad. Sci.*, 1987, vol. 84, 4007-4001 [0047]
- **ZHAO ; HARRIS.** *ACS Symposium Series*, 1997, vol. 680, 458-72 [0086]
- **ZALIPSKY.** *Adv. Drug Del. Rev.*, 1995, vol. 16, 157-82 [0086]