



(11) **EP 1 939 296 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
02.07.2008 Bulletin 2008/27

(51) Int Cl.:
C12N 15/55 (2006.01) C12P 13/08 (2006.01)

(21) Application number: **07123936.2**

(22) Date of filing: **17.12.2004**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR**
Designated Extension States:
AL BA HR LV MK RS

(30) Priority: **18.12.2003 PCT/IB03/06456**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
04817630.9 / 1 697 518

(71) Applicant: **BASF SE**
67056 Ludwigshafen (DE)

(72) Inventors:
• **Zelder, Oskar**
67346 Speyer (DE)

- **Klopprogge, Corinna**
68239 Mannheim (DE)
- **Kröger, Burkhard**
67117 Limburgerhof (DE)
- **Schröder, Hartwig**
69226 Nußloch (DE)
- **Haefner, Stefan**
67346 Speyer (DE)
- **Heinzle, Elmar**
66123 Saarbrücken (DE)
- **Wittmann, Christoph**
66127 Saarbrücken (DE)
- **Kiefer, Patrick**
8048 Zürich (CH)

Remarks:

This application was filed on 21-12-2007 as a
divisional application to the application mentioned
under INID code 62.

(54) **Methods for the preparation of a fine chemical by fermentation**

(57) The present invention features methods of in-
creasing the production of a fine chemical, e.g., lysine
from a microorganism, e.g., *Corynebacterium* by way of
deregulating an enzyme encoding gene, i.e., fructose-
1,6-bisphosphatase. In a preferred embodiment, the in-
vention provides methods of increasing the production
of lysine in *Corynebacterium glutamicum* by way of in-
creasing the expression of fructose-1,6-bisphosphatase

activity. The invention also provides a novel process for
the production of lysine by way of regulating carbon flux
towards oxaloacetate (OAA). In a preferred embodiment,
the invention provides methods for the production of
lysine by way of utilizing fructose or sucrose as a carbon
source.

EP 1 939 296 A2

DescriptionBackground of the Invention

[0001] The industrial production of the amino acid lysine has become an economically important industrial process. lysine is used commercially as an animal feed supplement, because of its ability to improve the quality of feed by increasing the absorption of other amino acids, in human medicine, particularly as ingredients of infusion solutions, and in the pharmaceutical industry.

[0002] Commercial production of this lysine is principally done utilizing the gram positive *Corynebacterium glutamicum*, *Brevibacterium flavum* and *Brevibacterium lac-tofermentum* (Kleemann, A., et. al., "Amino Acids," in ULLMANN'S ENCYCLOPEDIA OF INDUSTRIAL CHEMISTRY, vol. A2, pp.57-97, Weinham: VCH-Verlagsgesellschaft (1985)). These organisms presently account for the approximately 250,000 tons of lysine produced annually. A significant amount of research has gone into isolating mutant bacterial strains which produce larger amounts of lysine. Microorganisms employed in microbial process for amino acid production are divided into 4 classes: wild-type strain, auxotrophic mutant, regulatory mutant and auxotrophic regulatory mutant (K. Nakayama et al., in Nutritional Improvement of Food and Feed Proteins, M. Fried-man, ed., (1978), pp. 649-661). Mutants of *Corynebacterium* and related organisms enable inexpensive production of amino acids from cheap carbon sources, e.g., molasses, acetic acid and ethanol, by direct fermentation. In addition, the stereospecificity of the amino acids produced by fermentation (the L isomer) makes the process advantageous compared with synthetic processes.

[0003] Another method in improving the efficiency of the commercial production of lysine is by investigating the correlation between lysine production and metabolic flux through the pentose phosphate pathway. Given the economic importance of lysine production by the fermentive process, the biochemical pathway for lysine synthesis has been intensively investigated, ostensibly for the purpose of increasing the total amount of lysine produced and decreasing production costs (reviewed by Sahm et al., (1996) Ann. N. Y. Acad. Sci. 782:25-39). There has been some success in using metabolic engineering to direct the flux of glucose derived carbons toward aromatic amino acid formation (Flores, N. et al., (1996) Nature Biotechnol. 14:620-623). Upon cellular absorption, glucose is phosphorylated with consumption of phosphoenolpyruvate (phosphotransferase system) (Malin & Bourd, (1991) Journal of Applied Bacteriology 71, 517-523) and is then available to the cell as glucose-6-phosphate. Sucrose is converted into fructose and glucose-6-phosphate by a phosphotransferase system (Shio et al., (1990) Agricultural and Biological Chemistry 54, 1513-1519) and invertase reaction (Yamamoto et al., (1986) Journal of Fermentation Technology 64, 285-291).

[0004] During glucose catabolism, the enzymes glucose-6-phosphate dehydrogenase (EC 1.1.1.49) and glucose-6-phosphate isomerase (EC 5.3.1.9) compete for the substrate glucose-6-phosphate. The enzyme glucose-6-phosphate isomerase catalyses the first reaction step of the Embden-Meyerhof-Parnas pathway, or glycolysis, namely conversion into fructose-6-phosphate. The enzyme glucose-6-phosphate dehydrogenase catalyses the first reaction step of the oxidative portion of the pentose phosphate cycle, namely conversion into 6-phosphogluconolactone.

[0005] In the oxidative portion of the pentose phosphate cycle, glucose-6-phosphate is converted into ribulose-5-phosphate, so producing reduction equivalents in the form of NADPH. As the pentose phosphate cycle proceeds further, pentose phosphates, hexose phosphates and triose phosphates are interconverted. Pentose phosphates, such as for example 5-phosphoribosyl-1-pyrophosphate are required, for example, in nucleotide biosynthesis. 5-Phosphoribosyl-1-pyrophosphate is moreover a precursor for aromatic amino acids and the amino acid L-histidine. NADPH acts as a reduction equivalent in numerous anabolic biosyntheses. Four molecules of NADPH are thus consumed for the biosynthesis of one molecule of lysine from oxalacetic acid. Thus, carbon flux towards oxaloacetate (OAA) remains constant regardless of system perturbations (J. Vallino et al., (1993) Biotechnol. Bioeng., 41, 633-646).

Summary of the Invention

[0006] The present invention is based, at least in part, on the discovery of key enzyme-encoding genes, e.g., fructose-1,6-bisphosphatase, of the pentose phosphate pathway in *Corynebacterium glutamicum*, and the discovery that deregulation, e.g., increasing expression or activity of fructose-1,6-bisphosphatase results in increased lysine production. Furthermore, it has been found that increasing the carbon yield during production of lysine by deregulating, e.g., increasing, fructose-1,6 bisphosphatase expression or activity leads to increased lysine production. In one embodiment, the carbon source is fructose or sucrose. Accordingly, the present invention provides methods for increasing production of lysine by microorganisms, e.g., *C. glutamicum*, where fructose or sucrose is the substrate.

[0007] Accordingly, in one aspect, the invention provides methods for increasing metabolic flux through the pentose phosphate pathway in a microorganism comprising culturing a microorganism comprising a gene which is deregulated under conditions such that metabolic flux through the pentose phosphate pathway is increased. In one embodiment, the microorganism is fermented to produce a fine chemical, e.g., lysine. In another embodiment, fructose or sucrose is used as a carbon source. In still another embodiment, the gene is fructose-1,6-bisphosphatase. In a related embodiment, the

fructose-1,6-bisphosphatase gene is derived from *Corynebacterium*, e.g., *Corynebacterium glutamicum*. In another embodiment, fructose-1,6 bisphosphatase gene is overexpressed. In a further embodiment, the protein encoded by the fructose-1,6-bisphosphatase gene has increased activity.

[0008] In another embodiment, the microorganism further comprises one or more additional deregulated genes. The one or more additional deregulated gene can include, but is not limited to, an ask gene, a dapA gene, an asd gene, a dapB gene, a ddh gene, a lysA gene, a lysE gene, a pycA gene, a zwf gene, a pepCL gene, a gap gene, a zwal gene, a tkt gene, a lad gene, a mqo gene, a tpi gene, a pgk gene, and a sigC gene.

[0009] In a particular embodiment, the gene may be overexpressed or underexpressed,

[0010] Moreover, the deregulated gene can encode a protein selected from the group consisting of a feed-back resistant aspartokinase, a dihydrodipicolinate synthase, an aspartate semialdehyde dehydrogenase, a dihydrodipicolinate reductase, a diaminopimelate dehydrogenase, a diaminopimelate epimerase, a lysine exporter, a pyruvate carboxylase, a phosphate dehydrogenase, a phosphoenolpyruvate carboxylase, a glyceraldehyde-3-phosphate dehydrogenase, an RPF protein precursor, a transketolase, a transaldolase, a menaquinone oxidoreductase, a triosephosphate isomerase, a 3-phosphoglycerate kinase, and an RNA-polymerase sigma factor sigC. In a particular embodiment, the protein may have an increased or a decreased activity.

[0011] In accordance with the methods of the present invention, the one or more additional deregulated genes can also include, but is not limited to, a pepCK gene, a malE gene, a glgA gene, a pgi gene, a dead gene, a menE gene, a citE gene, a mikE17 gene, a poxB gene, a zwa2 gene, and a sucC gene. In a particular embodiment the expression of the at least one gene is upregulated, attenuated, decreased, downregulated or repressed. Moreover, the deregulated gene can encode a protein selected from the group consisting of a phosphoenolpyruvate carboxykinase, a malic enzyme, a glycogen synthase, a glucose-6-phosphate isomerase, an ATP dependent RNA helicase, an *o*-succinylbenzoic acid-CoA ligase, a citrate lyase beta chain, a transcriptional regulator, a pyruvate dehydrogenase, an RPF protein precursor, and a Succinyl-CoA-Synthetase. In a particular embodiment, the protein has a decreased or an increased activity.

[0012] In one embodiment, the microorganisms used in the methods of the invention belong to the genus *Corynebacterium*, e.g., *Corynebacterium glutamicum*.

[0013] In another aspect, the invention provides methods for producing a fine chemical comprising fermenting a microorganism in which fructose-1,6-bisphosphatase is deregulated and accumulating the fine chemical, e.g., lysine, in the medium or in the cells of the microorganisms, thereby producing a fine chemical. In one embodiment, the methods include recovering the fine chemical. In another embodiment, the fructose-1,6-bisphosphatase gene is overexpressed. In yet another embodiment, fructose or sucrose is used as a carbon source.

[0014] In one aspect, fructose-1,6-bisphosphatase is derived from *Corynebacterium glutamicum* and comprises the nucleotide sequence of SEQ ID NO:1 and the amino acid sequence of SEQ ID NO:2.

[0015] Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

[0016]

Figure 1: is a schematic representation of the pentose biosynthetic pathway.

Figure 2: Comparison of relative mass isotopomer fractions of secreted lysine and trehalose measured by GC/MS in tracer experiments of *Corynebacterium glutamicum* ATCC 21526 during lysine production on glucose and fructose.

Figure 3: In vivo carbon flux distribution in the central metabolism of *Corynebacterium glutamicum* ATCC 21526 during lysine production on glucose estimated from the best fit to the experimental results using a comprehensive approach of combined metabolite balancing and isotopomer modeling for ^{13}C tracer experiments with labeling measurement of secreted lysine and trehalose by GC/MS, respectively. Net fluxes are given in square symbols, whereby for reversible reactions the direction of the net flux is indicated by an arrow aside the corresponding black box. Numbers in brackets below the fluxes of transaldolase, transketolase and glucose 6-phosphate isomerase indicate flux reversibilities. All fluxes are expressed as a molar percentage of the mean specific glucose uptake rate ($1.77 \text{ mmol g}^{-1} \text{ h}^{-1}$).

Figure 4: In vivo carbon flux distribution in the central metabolism of *Corynebacterium glutamicum* ATCC 21526 during lysine production on fructose estimated from the best fit to the experimental results using a comprehensive approach of combined metabolite balancing and isotopomer modeling for ^{13}C tracer experiments with labeling measurement of secreted lysine and trehalose by GC/MS, respectively. Net fluxes are given in square symbols, whereby for reversible reactions the direction of the net flux is indicated by an arrow aside the corresponding black box.

Numbers in brackets below the fluxes of transaldolase, transketolase and glucose 6-phosphate isomerase indicate flux reversibilities. All fluxes are expressed as a molar percentage of the mean specific fructose uptake rate ($1.93 \text{ mmol g}^{-1} \text{ h}^{-1}$).

Figure 5: Metabolic network of the central metabolism for glucose-grown (A) and fructose-grown (B) lysine producing *Corynebacterium glutamicum* including transport fluxes, anabolic fluxes and fluxes between intermediary metabolite pools.

Detailed Description of the Invention

[0017] The present invention is based at least in part, on the identification of genes, e.g., *Corynebacterium glutamicum* genes, which encode essential enzymes of the pentose phosphate pathway. The present invention features methods comprising manipulating the pentose phosphate biosynthetic pathway in a microorganism, e.g., *Corynebacterium glutamicum* such that the carbon yield is increased and certain desirable fine chemicals, e.g., lysine, are produced, e.g., produced at increased yields. In particular, the invention includes methods for producing fine chemicals, e.g., lysine, by fermentation of a microorganism, e.g., *Corynebacterium glutamicum*, having deregulated, e.g., increased, fructose-1,6-bisphosphatase expression or activity. In one embodiment, fructose or saccharose is used as a carbon source in the fermentation of the microorganism. Fructose has been established to be a less efficient substrate for the production of fine chemicals, e.g., lysine, from microorganisms. However, the present invention provides methods for optimizing production of lysine by microorganisms, e.g., *C. glutamicum* where fructose or sucrose is the substrate. Deregulation, e.g., amplification, of fructose-1,6-bisphosphatase expression or activity leads to a higher flux through the pentose phosphate pathway, resulting in increased NADPH generation and increased lysine yield.

[0018] The term "pentose phosphate pathway" includes the pathway involving pentose phosphate enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of fine chemicals, e.g., lysine. The pentose phosphate pathway converts glucose molecules into biochemically useful smaller molecule.

[0019] In order that the present invention may be more readily understood, certain terms are first defined herein.

[0020] The term "pentose phosphosphate biosynthetic pathway" includes the biosynthetic pathway involving pentose phosphate biosynthetic genes, enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of fine chemicals, e.g., lysine. The term "pentose phosphosphate biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of fine chemicals, e.g., lysine, in a microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of fine chemicals, e.g., lysine, in vitro. The term "pentose phosphosphate biosynthetic pathway protein" or "pentose phosphosphate biosynthetic pathway enzyme" includes a those peptides, polypeptides, proteins, enzymes, and fragments thereof which are directly or indirectly involved in the pentose phosphosphate biosynthetic pathway, e.g., the fructose-1,6-bisphosphatase enzyme.

[0021] The term "pentose phosphosphate biosynthetic pathway gene" includes a those genes and gene fragments encoding peptides, polypeptides, proteins, and enzymes which are directly or indirectly involved in the pentose phosphosphate biosynthetic pathway, e.g., the fructose-1,6-bisphosphatase gene.

[0022] The term "amino acid biosynthetic pathway gene" is meant to include those genes and gene fragments encoding peptides, polypeptides, proteins, and enzymes, which are directly involved in the synthesis of amino acids, e.g., fructose-1,6-bisphosphatase. These genes may be identical to those which naturally occur within a host cell and are involved in the synthesis of any amino acid, and particularly lysine, within that host cell.

[0023] The term "lysine biosynthetic pathway gene" includes those genes and genes fragments encoding peptides, polypeptides, proteins, and enzymes, which are directly involved in the synthesis of lysine, e.g., fructose-1,6-bisphosphatase. These genes can be identical to those which naturally occur within a host cell and are involved in the synthesis of lysine within that host cell. Alternatively, there can be modifications or mutations of such genes, for example, the genes can contain modifications or mutations which do not significantly affect the biological activity of the encoded protein. For example, the natural gene can be modified by mutagenesis or by introducing or substituting one or more nucleotides or by removing nonessential regions of the gene. Such modifications are readily performed by standard techniques.

[0024] The term "lysine biosynthetic pathway protein" is meant to include those peptides, polypeptides, proteins, enzymes, and fragments thereof which are directly involved in the synthesis of lysine. These proteins can be identical to those which naturally occur within a host cell and are involved in the synthesis of lysine within that host cell. Alternatively, there can be modifications or mutations of such proteins, for example, the proteins can contain modifications or mutations which do not significantly affect the biological activity of the protein. For example, the natural protein can be modified by mutagenesis or by introducing or substituting one or more amino acids, preferably by conservative amino acid substitution, or by removing nonessential regions of the protein. Such modifications are readily performed by standard

techniques. Alternatively, lysine biosynthetic proteins can be heterologous to the particular host cell. Such proteins can be from any organism having genes encoding proteins having the same, or similar, biosynthetic roles.

[0025] The term, "carbon flux" refers to the number of glucose molecules which proceed down a particular metabolic path relative to competing paths. In particular, increased NADPH within a microorganism is achieved by altering the carbon flux distribution between the glycolytic and pentose phosphate pathways of that organism.

[0026] "Fructose-1,6-bisphosphatase activity" includes any activity exerted by a fructose-1,6-bisphosphatase protein, polypeptide or nucleic acid molecule as determined in vivo, or in vitro, according to standard techniques. Fructose-1,6-bisphosphatase is involved in many different metabolic pathways and found in most organisms. Preferably, a fructose-1,6-bisphosphatase activity includes the catalysis of the hydrolysis of fructose 1,6-bisphosphate to fructose 6-phosphate.

[0027] The term 'fine chemical' is art-recognized and includes molecules produced by an organism which have applications in various industries, such as, but not limited to, the pharmaceutical, agriculture, and cosmetics industries. Such compounds include organic acids, such as tartaric acid, itaconic acid, and diaminopimelic acid, both proteinogenic and non-proteinogenic amino acids, purine and pyrimidine bases, nucleosides, and nucleotides (as described e.g. in Kunitake, A. (1996) Nucleotides and related compounds, p. 561-612, in Biotechnology vol. 6, Rehm et al., eds. VCH: Weinheim, and references contained therein), lipids, both saturated and unsaturated fatty acids (e.g., arachidonic acid), diols (e.g., propane diol, and butane diol), carbohydrates (e.g., hyaluronic acid and trehalose), aromatic compounds (e.g., aromatic amines, vanillin, and indigo), vitamins and cofactors (as described in Ullmann's Encyclopedia of Industrial Chemistry, vol. A27, "Vitamins", p. 443-613 (1996) VCH: Weinheim and references therein; and Ong, A.S., Niki, E. & Packer, L. (1995) "Nutrition, Lipids, Health, and Disease" Proceedings of the UNESCO/Confederation of Scientific and Technological Associations in Malaysia, and the Society for Free Radical Research - Asia, held Sept. 1-3, 1994 at Penang, Malaysia, AOCS Press, (1995)), enzymes, polyketides (Cane et al. (1998) Science 282: 63-68), and all other chemicals described in Gutcho (1983) Chemicals by Fermentation, Noyes Data Corporation, ISBN: 0818805086 and references therein. The metabolism and uses of certain of these fine chemicals are further explicated below.

Amino Acid Metabolism and Uses

[0028] Amino acids comprise the basic structural units of all proteins, and as such are essential for normal cellular functioning in all organisms. The term "amino acid" is art-recognized. The proteinogenic amino acids, of which there are 20 species, serve as structural units for proteins, in which they are linked by peptide bonds, while the nonproteinogenic amino acids (hundreds of which are known) are not normally found in proteins (see Ullmann's Encyclopedia of Industrial Chemistry, vol. A2, p. 57-97 VCH: Weinheim (1985)). Amino acids may be in the D- or L- optical configuration, though L-amino acids are generally the only type found in naturally-occurring proteins. Biosynthetic and degradative pathways of each of the 20 proteinogenic amino acids have been well characterized in both prokaryotic and eukaryotic cells (see, for example, Stryer, L. Biochemistry, 3rd edition, pages 578-590 (1988)). The 'essential' amino acids (histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, and valine), so named because they are generally a nutritional requirement due to the complexity of their biosyntheses, are readily converted by simple biosynthetic pathways to the remaining 11 'nonessential' amino acids (alanine, arginine, asparagine, aspartate, cysteine, glutamate, glutamine, glycine, proline, serine, and tyrosine). Higher animals do retain the ability to synthesize some of these amino acids, but the essential amino acids must be supplied from the diet in order for normal protein synthesis to occur.

[0029] Aside from their function in protein biosynthesis, these amino acids are interesting chemicals in their own right, and many have been found to have various applications in the food, feed, chemical, cosmetics, agriculture, and pharmaceutical industries. Lysine is an important amino acid in the nutrition not only of humans, but also of monogastric animals such as poultry and swine. Glutamate is most commonly used as a flavor additive (mono-sodium glutamate, MSG) and is widely used throughout the food industry, as are aspartate, phenylalanine, glycine, and cysteine. Glycine, L-methionine and tryptophan are all utilized in the pharmaceutical industry. Glutamine, valine, leucine, isoleucine, histidine, arginine, proline, serine and alanine are of use in both the pharmaceutical and cosmetics industries. Threonine, tryptophan, and D/L-methionine are common feed additives. (Leuchtenberger, W. (1996) Amino acids - technical production and use, p. 466-502 in Rehm et al. (eds.) Biotechnology vol. 6, chapter 14a, VCH: Weinheim). Additionally, these amino acids have been found to be useful as precursors for the synthesis of synthetic amino acids and proteins, such as N-acetylcysteine, S-carboxymethyl-L-cysteine, (S)-5-hydroxytryptophan, and others described in Ullmann's Encyclopedia of Industrial Chemistry, vol. A2, p. 57-97, VCH: Weinheim, 1985.

[0030] The biosynthesis of these natural amino acids in organisms capable of producing them, such as bacteria, has been well characterized (for review of bacterial amino acid biosynthesis and regulation thereof, see Umbarger, H.E. (1978) Ann. Rev. Biochem. 47: 533-606). Glutamate is synthesized by the reductive amination of α -ketoglutarate, an intermediate in the citric acid cycle. Glutamine, proline, and arginine are each subsequently produced from glutamate. The biosynthesis of serine is a three-step process beginning with 3-phosphoglycerate (an intermediate in glycolysis), and resulting in this amino acid after oxidation, transamination, and hydrolysis steps. Both cysteine and glycine are produced from serine; the former by the condensation of ho-mocysteine with serine, and the latter by the transfer of

the side-chain β -carbon atom to tetrahydrofolate, in a reaction catalyzed by serine transhydroxymethylase. Phenylalanine, and tyrosine are synthesized from the glycolytic and pentose phosphate pathway precursors erythrose 4-phosphate and phosphoenolpyruvate in a 9-step biosynthetic pathway that differ only at the final two steps after synthesis of prephenate. Tryptophan is also produced from these two initial molecules, but its synthesis is an 11-step pathway. Tyrosine may also be synthesized from phenylalanine, in a reaction catalyzed by phenylalanine hydroxylase. Alanine, valine, and leucine are all biosynthetic products of pyruvate, the final product of glycolysis. Aspartate is formed from oxaloacetate, an intermediate of the citric acid cycle. Asparagine, methionine, threonine, and lysine are each produced by the conversion of aspartate. Isoleucine is formed from threonine. A complex 9-step pathway results in the production of histidine from 5-phosphoribosyl-1-pyrophosphate, an activated sugar.

[0031] Amino acids in excess of the protein synthesis needs of the cell cannot be stored, and are instead degraded to provide intermediates for the major metabolic pathways of the cell (for review see Stryer, L. *Biochemistry* 3rd ed. Ch. 21 "Amino Acid Degradation and the Urea Cycle" p. 495-516 (1988)). Although the cell is able to convert unwanted amino acids into useful metabolic intermediates, amino acid production is costly in terms of energy, precursor molecules, and the enzymes necessary to synthesize them. Thus it is not surprising that amino acid biosynthesis is regulated by feedback inhibition, in which the presence of a particular amino acid serves to slow or entirely stop its own production (for overview of feedback mechanisms in amino acid biosynthetic pathways, see Stryer, L. *Biochemistry*, 3rd ed. Ch. 24: "Biosynthesis of Amino Acids and Heme" p. 575-600 (1988)). Thus, the output of any particular amino acid is limited by the amount of that amino acid present in the cell. Vitamin, Cofactor, and Nutraceutical Metabolism and Uses

[0032] Vitamins, cofactors, and nutraceuticals comprise another group of molecules which the higher animals have lost the ability to synthesize and so must ingest, although they are readily synthesized by other organisms such as bacteria. These molecules are either bioactive substances themselves, or are precursors of biologically active substances which may serve as electron carriers or intermediates in a variety of metabolic pathways. Aside from their nutritive value, these compounds also have significant industrial value as coloring agents, antioxidants, and catalysts or other processing aids. (For an overview of the structure, activity, and industrial applications of these compounds, see, for example, Ullman's Encyclopedia of Industrial Chemistry, "Vitamins" vol. A27, p. 443-613, VCH: Weinheim, 1996.) The term "vitamin" is art-recognized, and includes nutrients which are required by an organism for normal functioning, but which that organism cannot synthesize by itself. The group of vitamins may encompass cofactors and nutraceutical compounds. The language "cofactor" includes nonproteinaceous compounds required for a normal enzymatic activity to occur. Such compounds may be organic or inorganic; the cofactor molecules of the invention are preferably organic. The term "nutraceutical" includes dietary supplements having health benefits in plants and animals, particularly humans. Examples of such molecules are vitamins, antioxidants, and also certain lipids (e.g., polyunsaturated fatty acids).

[0033] The biosynthesis of these molecules in organisms capable of producing them, such as bacteria, has been largely characterized (Ullman's Encyclopedia of Industrial Chemistry, "Vitamins" vol. A27, p. 443-613, VCH: Weinheim, 1996; Michal, G. (1999) *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*, John Wiley & Sons; Ong, A.S., Niki, E. & Packer, L. (1995) "Nutrition, Lipids, Health, and Disease" Proceedings of the UNESCO/Confederation of Scientific and Technological Associations in Malaysia, and the Society for Free Radical Research - Asia, held Sept. 1-3, 1994 at Penang, Malaysia, AOCS Press: Champaign, IL X, 374 S).

[0034] Thiamin (vitamin B₁) is produced by the chemical coupling of pyrimidine and thiazole moieties. Riboflavin (vitamin B₂) is synthesized from guanosine-5'-triphosphate (GTP) and ribose-5'-phosphate. Riboflavin, in turn, is utilized for the synthesis of flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD). The family of compounds collectively termed 'vitamin B₆' (e.g., pyridoxine, pyridoxamine, pyridoxa-5'-phosphate, and the commercially used pyridoxin hydrochloride) are all derivatives of the common structural unit, 5-hydroxy-6-methylpyridine. Pantothenate (pantothenic acid, (R)-(+)-N-(2,4-dihydroxy-3,3-dimethyl-1-oxobutyl)- β -alanine) can be produced either by chemical synthesis or by fermentation. The final steps in pantothenate biosynthesis consist of the ATP-driven condensation of β -alanine and pantoic acid. The enzymes responsible for the biosynthesis steps for the conversion to pantoic acid, to β -alanine and for the condensation to pantothenic acid are known. The metabolically active form of pantothenate is Coenzyme A, for which the biosynthesis proceeds in 5 enzymatic steps. Pantothenate, pyridoxal-5'-phosphate, cysteine and ATP are the precursors of Coenzyme A. These enzymes not only catalyze the formation of pantothenate, but also the production of (R)-pantoic acid, (R)-pantolacton, (R)-panthanol (provitamin B₅), pantetheine (and its derivatives) and coenzyme A.

[0035] Biotin biosynthesis from the precursor molecule pimeloyl-CoA in microorganisms has been studied in detail and several of the genes involved have been identified. Many of the corresponding proteins have been found to also be involved in Fecluster synthesis and are members of the nifS class of proteins. Lipoic acid is derived from octanoic acid, and serves as a coenzyme in energy metabolism, where it becomes part of the pyruvate dehydrogenase complex and the α -ketoglutarate dehydrogenase complex. The folates are a group of substances which are all derivatives of folic acid, which in turn is derived from L-glutamic acid, p-amino-benzoic acid and 6-methylpterin. The biosynthesis of folic acid and its derivatives, starting from the metabolism intermediates guanosine-5'-triphosphate (GTP), L-glutamic acid and p-amino-benzoic acid has been studied in detail in certain microorganisms.

[0036] Corrinoids (such as the cobalamines and particularly vitamin B₁₂) and porphyrines belong to a group of chemicals

characterized by a tetrapyrrole ring system. The biosynthesis of vitamin B₁₂ is sufficiently complex that it has not yet been completely characterized, but many of the enzymes and substrates involved are now known.

[0037] Nicotinic acid (nicotinate), and nicotinamide are pyridine derivatives which are also termed 'niacin'. Niacin is the precursor of the important coenzymes NAD (nicotinamide adenine dinucleotide) and NADP (nicotinamide adenine dinucleotide phosphate) and their reduced forms.

[0038] The large-scale production of these compounds has largely relied on cell-free chemical syntheses, though some of these chemicals have also been produced by large-scale culture of microorganisms, such as riboflavin, Vitamin B₆, pantothenate, and biotin. Only Vitamin B₁₂ is produced solely by fermentation, due to the complexity of its synthesis. In vitro methodologies require significant inputs of materials and time, often at great cost.

Purine, Pyrimidine, Nucleoside and Nucleotide Metabolism and Uses

[0039] Purine and pyrimidine metabolism genes and their corresponding proteins are important targets for the therapy of tumor diseases and viral infections. The language "purine" or "pyrimidine" includes the nitrogenous bases which are constituents of nucleic acids, co-enzymes, and nucleotides. The term "nucleotide" includes the basic structural units of nucleic acid molecules, which are comprised of a nitrogenous base, a pentose sugar (in the case of RNA, the sugar is ribose; in the case of DNA, the sugar is D-deoxyribose), and phosphoric acid. The language "nucleoside" includes molecules which serve as precursors to nucleotides, but which are lacking the phosphoric acid moiety that nucleotides possess. By inhibiting the biosynthesis of these molecules, or their mobilization to form nucleic acid molecules, it is possible to inhibit RNA and DNA synthesis; by inhibiting this activity in a fashion targeted to cancerous cells, the ability of tumor cells to divide and replicate may be inhibited. Additionally, there are nucleotides which do not form nucleic acid molecules, but rather serve as energy stores (i.e., AMP) or as coenzymes (i.e., FAD and NAD).

[0040] Several publications have described the use of these chemicals for these medical indications, by influencing purine and/or pyrimidine metabolism (e.g. Christopherson, R.I. and Lyons, S.D. (1990) "Potent inhibitors of de novo pyrimidine and purine biosynthesis as chemotherapeutic agents." *Med. Res. Reviews* 10: 505-548). Studies of enzymes involved in purine and pyrimidine metabolism have been focused on the development of new drugs which can be used, for example, as immunosuppressants or anti-proliferants (Smith, J.L., (1995) "Enzymes in nucleotide synthesis." *Curr. Opin. Struct. Biol.* 5: 752-757; (1995) *Biochem Soc. Transact.* 23: 877-902). However, purine and pyrimidine bases, nucleosides and nucleotides have other utilities: as intermediates in the biosynthesis of several fine chemicals (e.g., thiamine, S-adenosyl-methionine, folates, or riboflavin), as energy carriers for the cell (e.g., ATP or GTP), and for chemicals themselves, commonly used as flavor enhancers (e.g., IMP or GMP) or for several medicinal applications (see, for example, Kuninaka, A. (1996) *Nucleotides and Related Compounds in Biotechnology* vol. 6, Rehms et al., eds. VCH: Weinheim, p. 561-612). Also, enzymes involved in purine, pyrimidine, nucleoside, or nucleotide metabolism are increasingly serving as targets against which chemicals for crop protection, including fungicides, herbicides and insecticides, are developed.

[0041] The metabolism of these compounds in bacteria has been characterized (for reviews see, for example, Zalkin, H. and Dixon, J.E. (1992) "de novo purine nucleotide biosynthesis", in: *Progress in Nucleic Acid Research and Molecular Biology*, vol. 42, Academic Press; p. 259-287; and Michal, G. (1999) "Nucleotides and Nucleosides", Chapter 8 in: *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*, Wiley: New York). Purine metabolism has been the subject of intensive research, and is essential to the normal functioning of the cell. Impaired purine metabolism in higher animals can cause severe disease, such as gout. Purine nucleotides are synthesized from ribose-5-phosphate, in a series of steps through the intermediate compound inosine-5'-phosphate (IMP), resulting in the production of guanosine-5'-monophosphate (GMP) or adenosine-5'-monophosphate (AMP), from which the triphosphate forms utilized as nucleotides are readily formed. These compounds are also utilized as energy stores, so their degradation provides energy for many different biochemical processes in the cell. Pyrimidine biosynthesis proceeds by the formation of uridine-5'-monophosphate (UMP) from ribose-5-phosphate. UMP, in turn, is converted to cytidine-5'-triphosphate (CTP). The deoxy- forms of all of these nucleotides are produced in a one step reduction reaction from the diphosphate ribose form of the nucleotide to the diphosphate deoxyribose form of the nucleotide. Upon phosphorylation, these molecules are able to participate in DNA synthesis. Trehalose Metabolism and Uses

[0042] Trehalose consists of two glucose molecules, bound in α , α -1,1 linkage. It is commonly used in the food industry as a sweetener, an additive for dried or frozen foods, and in beverages. However, it also has applications in the pharmaceutical, cosmetics and biotechnology industries (see, for example, Nishimoto et al., (1998) U.S. Patent No. 5,759,610; Singer, M.A. and Lindquist, S. (1998) *Trends Biotech.* 16: 460-467; Paiva, C.L.A. and Panek, A.D. (1996) *Biotech. Ann. Rev.* 2: 293-314; and Shiosaka, M. (1997) *J. Japan* 172: 97-102). Trehalose is produced by enzymes from many microorganisms and is naturally released into the surrounding medium, from which it can be collected using methods known in the art.

I. Recombinant Microorganisms and Methods for Culturing Microorganisms Such That A Fine Chemical is Produced

[0043] The methodologies of the present invention feature microorganisms, e.g., recombinant microorganisms, preferably including vectors or genes (e.g., wild-type and/or mutated genes) as described herein and/or cultured in a manner which results in the production of a desired fine chemical, e.g. lysine. The term "recombinant" microorganism includes a microorganism (e.g., bacteria, yeast cell, fungal cell, etc.) which has been genetically altered, modified or engineered (e.g., genetically engineered) such that it exhibits an altered, modified or different genotype and/or phenotype (e.g., when the genetic modification affects coding nucleic acid sequences of the microorganism) as compared to the naturally-occurring microorganism from which it was derived. Preferably, a "recombinant" microorganism of the present invention has been genetically engineered such that it overexpresses at least one bacterial gene or gene product as described herein, preferably a biosynthetic enzyme encoding-gene, e.g., fructose-1,6-bisphosphatase, included within a recombinant vector as described herein and/or a biosynthetic enzyme, e.g., fructose-1,6-bisphosphatase expressed from a recombinant vector. The ordinary skilled will appreciate that a microorganism expressing or overexpressing a gene product produces or overproduces the gene product as a result of expression or overexpression of nucleic acid sequences and/or genes encoding the gene product. In one embodiment, the recombinant microorganism has increased biosynthetic enzyme, e.g., fructose-1,6-bisphosphatase, activity.

[0044] In certain embodiments of the present invention, at least one gene or protein may be deregulated, in addition to the fructose-1,6-bisphosphatase gene or enzyme, so as to enhance the production of L-amino acids. For example, a gene or an enzyme of the biosynthesis pathways, for example, of glycolysis, of anaplerosis, of the citric acid cycle, of the pentose phosphate cycle, or of amino acids export may be deregulated. Additionally, a regulatory gene or protein may be deregulated.

[0045] In various embodiments, expression of a gene may be increased so as to increase the intracellular activity or concentration of the protein encoded by the gene, thereby ultimately improving the production of the desired amino acid. One skilled in the art may use various techniques to achieve the desired result. For example, a skilled practitioner may increase the number of copies of the gene or genes, use a potent promoter, and/or use a gene or allele which codes for the corresponding enzyme having high activity. Using the methods of the present invention, for example, overexpressing a particular gene, the activity or concentration of the corresponding protein can be increased by at least about 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400%, 500%, 1000% or 2000%, based on the starting activity or concentration.

[0046] In various embodiments, the deregulated gene may include, but is not limited to, at least one of the following genes or proteins:

- ❖ the ask gene which encodes a feed-back resistant aspartokinase (as disclosed in International Publication No. W02004089996);

- ❖ the dapA gene which encodes dihydrodipicolinate synthase (as disclosed in SEQ ID NOs : 55 and 56, respectively, in International Publication No. W0200100843) ;

- ❖ the asd gene which encodes an aspartate semialdehyde dehydrogenase (as disclosed in SEQ ID NOs : 3435 and 6935, respectively, in European Publication No. 1108790);

- ❖ the dapB gene which encodes a dihydrodipicolinate reductase (as disclosed in SEQ ID NOs : 35 and 36, respectively, in international Publication No. W0200100843);

- ❖ the ddh gene which encodes a diaminopimelate dehydrogenase (as disclosed in SEQ ID NOs : 3444 and 6944, respectively, in European Publication No. 1 108790);

- ❖ the lysA gene which encodes a diaminopimelate epimerase (as disclosed in SEQ ID NOs : 3451 and 6951, respectively, in European Publication No. 1108790);

- ❖ the lysE gene which encodes a lysine exporter (as disclosed in SEQ ID NOs : 3455 and 6955, respectively, in European Publication No. 1108790);

- ❖ the pycA gene which encodes a pyruvate carboxylase (as disclosed in SEQ ID NOs : 765 and 4265, respectively, in European Publication No. 1108790);

- ❖ the zwf gene which encodes a glucose-6-phosphate dehydrogenase (as disclosed in SEQ ID NOs : 243 and 244, respectively, in International Publication No. W0200100844);

- ❖ the pepCL gene which encodes a phosphoenolpyruvate carboxylase (as disclosed in SEQ ID NOs : 3470 and 6970, respectively, in European Publication No. 1108790) ;

- ❖ the gap gene which encodes a glyceraldehyde-3-phosphate dehydrogenase (as disclosed in SEQ ID NOs : 67 and 68, respectively, in International Publication No. W0200100844);

- ❖ the zwf gene which encodes an RPF protein precursor (as disclosed in SEQ ID NOs : 917 and 4417, respectively, in European Publication No. 1108790);

- ❖ the lkt gene which encodes a transketolase (as disclosed in SEQ ID NOs : 247 and 248, respectively, in International Publication No. W0200100844);

- ❖ the tad gene which encodes a transaldolase (as disclosed in SEQ ID NOs: 245 and 246, respectively, in International Publication No. W0200100844) ;
- ❖ the mgo gene which codes for a menaquinone oxidoreductase (as disclosed in SEQ ID NOs: 569 and 570, respectively, in international Publication No. W0200100844) ;
- ❖ the tpi gene which codes for a triosephosphate isomerase (as disclosed in SEQ ID NOs: 61 and 62, respectively, in International Publication No. W0200100844),
- ❖ the pgk gene which codes for a 3-phosphoglycerate kinase (as disclosed in SEQ ID NOs: 69 and 70, respectively, in International Publication No. W0200100844); and
- ❖ the sigC gene which codes for, an : RNA-, polymerase sigma factor sigc (as disclosed in SEQ ID NOs ; 284 and 3784, respectively, in Publication No. 1108790).

[0047] In particular embodiments, the gene may be overexpressed and/or the activity of the protein may be increased.

[0048] Alternatively, in other embodiments, expression of a gene may be attenuated, decreased or repressed so as to decrease, for example, eliminate, the intracellular activity or concentration of the protein encoded by the gene, thereby ultimately improving the production of the desired amino acid. For example, one skilled in the art may use a weak promoter. Alternatively or in combination, a skilled practitioner may use a gene or allele that either codes for the corresponding enzyme having low activity or inactivates the corresponding gene or enzyme. Using the methods of the present invention, the activity or concentration of the corresponding protein can be reduced to about 0 to 50%, 0 to 25%, 0 to 10%, 0 to 9%, 0 to 8%, 0 to 7%, 0 to 6%, 0 to 5%, 0 to 4%, 0 to 3%, 0 to 2% or 0 to 1 % of the activity or concentration of the wild- type protein.

[0049] In certain embodiments, the deregulated gene may include, but is not limited to, at least one of the following genes or proteins:

- ❖ the pepCK gene which codes for the phosphoenolpyruvate carboxykinase (as disclosed in SEQ., B) NOs: 179 and 180, respectively, in International Publication No. W0200100844) ;
- ❖ the mal E gene which codes for the malic enzyme (as disclosed in SEQ ID NOs : 3328 and 6828, respectively, in European Publication No. 1108790);
- ❖ the glgA gene which codes for the glycogen synthase (as disclosed in SEQ ID NOs: 1239 and 4739, respectively, in European Publication No. 1108790); # the pgi gene which codes for the glucose-6-phosphate isomerase (as disclosed in SEQ ID NOs : 41 and 42, respectively, in International Publication No. W0200100844) ;
- ❖ the dead gene which codes for the ATP dependent RNA helicase (as disclosed in SEQ 10 NOs : 1278 and 4778, respectively, in European Publication No. 1108790);
- ❖ the menE gene which codes for the o-succinylbenzoic acid-CoA ligase (as disclosed in SEQ ID NOs : and 4005, respectively, in European Publication No. 1108790),
- ❖ the citE gene which codes for the citrate lyase beta chain (as disclosed in SEQ ID NOs : 547 and 548, respectively, in International Publication No. W0200100844);
- ❖ the mikE17 gene which codes for a transcriptional regulator (as disclosed in SEQ ID NOs : 411 and 3911, respectively, in European Publication No. 1108790);
- ❖ the poxB gene which codes for the pyruvate dehydrogenase (as disclosed in SEQ ID NOs: 85 and 86, respectively, in International Publication No. W0200100844);
- ❖ the zwa2 gene which codes for an RPF protein precursor (as disclosed in European Publication No. ; and
- ❖ the sucC gene which codes for the Succinyl-CoA-Synthetase (as disclosed in European Publication No. 1103611).

[0050] In particular embodiments, the expression of the gene may be attenuated, decreased or repressed and/or the activity of the protein may be decreased.

[0051] The term "manipulated microorganism" includes a microorganism that has been engineered (e.g., genetically engineered) or modified such that results in the disruption or alteration of a metabolic pathway so as to cause a change in the metabolism of carbon. An enzyme is "overexpressed" in a metabolically engineered cell when the enzyme is expressed in the metabolically engineered cell at a level higher than the level at which it is expressed in a comparable wild-type cell. In cells that do not endogenously express a particular enzyme, any level of expression of that enzyme in the cell is deemed an "overexpression" of that enzyme for purposes of the present invention. Overexpression may lead to increased activity of the protein encoded by the gene, e.g., fructose-1,6-bisphosphatase.

[0052] Modification or engineering of such microorganisms can be according to any methodology described herein including, but not limited to, deregulation of a biosynthetic pathway and/or overexpression of at least one biosynthetic enzyme. A "manipulated" enzyme (e.g., a "manipulated" biosynthetic enzyme) includes an enzyme, the expression or production of which has been altered or modified such that at least one upstream or downstream precursor, substrate or product of the enzyme is altered or modified, e.g., has increased activity, for example, as compared to a corresponding

wild-type or naturally occurring enzyme.

[0053] The term "overexpressed" or "overexpression" includes expression of a gene product (e.g., a pentose phosphate biosynthetic enzyme) at a level greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. In one embodiment, the microorganism can be genetically manipulated (e.g., genetically engineered) to overexpress a level of gene product greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. Genetic manipulation can include, but is not limited to, altering or modifying regulatory sequences or sites associated with expression of a particular gene (e.g., by adding strong promoters, inducible promoters or multiple promoters or by removing regulatory sequences such that expression is constitutive), modifying the chromosomal location of a particular gene, altering nucleic acid sequences adjacent to a particular gene such as a ribosome binding site or transcription terminator, increasing the copy number of a particular gene, modifying proteins (e.g., regulatory proteins, suppressors, enhancers, transcriptional activators and the like) involved in transcription of a particular gene and/or translation of a particular gene product, or any other conventional means of deregulating expression of a particular gene routine in the art (including but not limited to use of antisense nucleic acid molecules, for example, to block expression of repressor proteins).

[0054] In another embodiment, the microorganism can be physically or environmentally manipulated to overexpress a level of gene product greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. For example, a microorganism can be treated with or cultured in the presence of an agent known or suspected to increase transcription of a particular gene and/or translation of a particular gene product such that transcription and/or translation are enhanced or increased. Alternatively, a microorganism can be cultured at a temperature selected to increase transcription of a particular gene and/or translation of a particular gene product such that transcription and/or translation are enhanced or increased.

[0055] The term "deregulated" or "deregulation" includes the alteration or modification of at least one gene in a microorganism that encodes an enzyme in a biosynthetic pathway, such that the level or activity of the biosynthetic enzyme in the microorganism is altered or modified. Preferably, at least one gene that encodes an enzyme in a biosynthetic pathway is altered or modified such that the gene product is enhanced or increased, thereby enhancing or increasing the activity of the gene product. The phrase "deregulated pathway" can also include a biosynthetic pathway in which more than one gene that encodes an enzyme in a biosynthetic pathway is altered or modified such that the level or activity of more than one biosynthetic enzyme is altered or modified. The ability to "deregulate" a pathway (e.g., to simultaneously deregulate more than one gene in a given biosynthetic pathway) in a microorganism arises from the particular phenomenon of microorganisms in which more than one enzyme (e.g., two or three biosynthetic enzymes) are encoded by genes occurring adjacent to one another on a contiguous piece of genetic material termed an "operon".

[0056] The term "operon" includes a coordinated unit of gene expression that contains a promoter and possibly a regulatory element associated with one or more, preferably at least two, structural genes (e.g., genes encoding enzymes, for example, biosynthetic enzymes). Expression of the structural genes can be coordinately regulated, for example, by regulatory proteins binding to the regulatory element or by anti-termination of transcription. The structural genes can be transcribed to give a single mRNA that encodes all of the structural proteins. Due to the coordinated regulation of genes included in an operon, alteration or modification of the single promoter and/or regulatory element can result in alteration or modification of each gene product encoded by the operon. Alteration or modification of the regulatory element can include, but is not limited to removing the endogenous promoter and/or regulatory element(s), adding strong promoters, inducible promoters or multiple promoters or removing regulatory sequences such that expression of the gene products is modified, modifying the chromosomal location of the operon, altering nucleic acid sequences adjacent to the operon or within the operon such as a ribosome binding site, increasing the copy number of the operon, modifying proteins (e.g., regulatory proteins, suppressors, enhancers, transcriptional activators and the like) involved in transcription of the operon and/or translation of the gene products of the operon, or any other conventional means of deregulating expression of genes routine in the art (including but not limited to use of antisense nucleic acid molecules, for example, to block expression of repressor proteins). Deregulation can also involve altering the coding region of one or more genes to yield, for example, an enzyme that is feedback resistant or has a higher or lower specific activity.

[0057] A particularly preferred "recombinant" microorganism of the present invention has been genetically engineered to overexpress a bacterially-derived gene or gene product. The term "bacterially-derived" or "derived-from", for example bacteria, includes a gene which is naturally found in bacteria or a gene product which is encoded by a bacterial gene (e.g., encoded by fructose-1,6-bisphosphatase).

[0058] The methodologies of the present invention feature recombinant microorganisms which overexpress one or more genes, e.g., the fructose-1,6-bisphosphatase gene or have increased or enhanced the fructose-1,6-bisphosphatase activity. A particularly preferred recombinant microorganism of the present invention (e.g., *Corynebacterium glutamicum*, *Corynebacterium acetoglutamicum*, *Corynebacterium acetoacidophilum*, and *Corynebacterium thermoamino-* genes, etc.) has been genetically engineered to overexpress a biosynthetic enzyme (e.g., fructose-1, 6-bisphosphatase, the amino acid sequence of SEQ ID NO:2 or encoded by the nucleic acid sequence of SEQ ID NO:1).

[0059] Other preferred "recombinant" microorganisms of the present invention have an enzyme deregulated in the

pentose phosphate pathway. The phrase "microorganism having a deregulated pentose phosphate pathway" includes a microorganism having an alteration or modification in at least one gene encoding an enzyme of the pentose phosphate pathway or having an alteration or modification in an operon including more than one gene encoding an enzyme of the pentose phosphate pathway. A preferred "microorganism having a deregulated pentose phosphate pathway" has been genetically engineered to overexpress a *Corynebacterium* (e.g., *C. glutamicum*) biosynthetic enzyme (e.g., has been engineered to overexpress fructose-1,6-bisphosphatase).

[0060] In another preferred embodiment, a recombinant microorganism is designed or engineered such that one or more pentose phosphate biosynthetic enzyme is overexpressed or deregulated.

[0061] In another preferred embodiment, a microorganism of the present invention overexpresses or is mutated for a gene or biosynthetic enzyme (e.g., a pentose phosphate biosynthetic enzyme) which is bacterially-derived. The term "bacterially-derived" or "derived-from", for example bacteria, includes a gene product (e.g., fructose-1,6-bisphosphatase) which is encoded by a bacterial gene.

[0062] In one embodiment, a recombinant microorganism of the present invention is a Gram positive organism (e.g., a microorganism which retains basic dye, for example, crystal violet, due to the presence of a Gram-positive wall surrounding the microorganism). In a preferred embodiment, the recombinant microorganism is a microorganism belonging to a genus selected from the group consisting of *Bacillus*, *Brevibacterium*, *Corynebacterium*, *Lactobacillus*, *Lactococci* and *Streptomyces*. In a more preferred embodiment, the recombinant microorganism is of the genus *Corynebacterium*. In another preferred embodiment, the recombinant microorganism is selected from the group consisting of *Corynebacterium glutamicum*, *Corynebacterium acetoglutamicum*, *Corynebacterium acetoacidophilum* or *Corynebacterium thermoaminogenes*. In a particularly preferred embodiment, the recombinant microorganism is *Corynebacterium glutamicum*.

[0063] An important aspect of the present invention involves culturing the recombinant microorganisms described herein, such that a desired compound (e.g., a desired fine chemical) is produced. The term "culturing" includes maintaining and/or growing a living microorganism of the present invention (e.g., maintaining and/or growing a culture or strain). In one embodiment, a microorganism of the invention is cultured in liquid media. In another embodiment, a microorganism of the invention is cultured in solid media or semi-solid media. In a preferred embodiment, a microorganism of the invention is cultured in media (e.g., a sterile, liquid media) comprising nutrients essential or beneficial to the maintenance and/or growth of the microorganism. Carbon sources which may be used include sugars and carbohydrates, such as for example glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as for example soy oil, sunflower oil, peanut oil and coconut oil, fatty acids, such as for example palmitic acid, stearic acid and linoleic acid, alcohols, such as for example glycerol and ethanol, and organic acids, such as for example acetic acid. In a preferred embodiment, fructose or saccharose. These substances may be used individually or as a mixture.

[0064] Nitrogen sources which may be used comprise organic compounds containing nitrogen, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya flour and urea or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate. The nitrogen sources may be used individually or as a mixture. Phosphorus sources which may be used are phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding salts containing sodium. The culture medium must furthermore contain metal salts, such as for example magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth-promoting substances such as amino acids and vitamins may also be used in addition to the above-stated substances. Suitable precursors may furthermore be added to the culture medium. The stated feed substances may be added to the culture as a single batch or be fed appropriately during cultivation.

[0065] Preferably, microorganisms of the present invention are cultured under controlled pH. The term "controlled pH" includes any pH which results in production of the desired fine chemical, e.g., lysine. In one embodiment, microorganisms are cultured at a pH of about 7. In another embodiment, microorganisms are cultured at a pH of between 6.0 and 8.5. The desired pH may be maintained by any number of methods known to those skilled in the art. For example, basic compounds such as sodium hydroxide, potassium hydroxide, ammonia, or ammonia water, or acidic compounds, such as phosphoric acid or sulfuric acid, are used to appropriately control the pH of the culture.

[0066] Also preferably, microorganisms of the present invention are cultured under controlled aeration. The term "controlled aeration" includes sufficient aeration (e.g., oxygen) to result in production of the desired fine chemical, e.g., lysine. In one embodiment, aeration is controlled by regulating oxygen levels in the culture, for example, by regulating the amount of oxygen dissolved in culture media. Preferably, aeration of the culture is controlled by agitating the culture. Agitation may be provided by a propeller or similar mechanical agitation equipment, by revolving or shaking the growth vessel (e.g., fermentor) or by various pumping equipment. Aeration may be further controlled by the passage of sterile air or oxygen through the medium (e.g., through the fermentation mixture). Also preferably, microorganisms of the present invention are cultured without excess foaming (e.g., via addition of antifoaming agents such as fatty acid polyglycol esters).

[0067] Moreover, microorganisms of the present invention can be cultured under controlled temperatures. The term "controlled temperature" includes any temperature which results in production of the desired fine chemical, e.g., lysine. In one embodiment, controlled temperatures include temperatures between 15°C and 95°C. In another embodiment,

controlled temperatures include temperatures between 15°C and 70°C. Preferred temperatures are between 20°C and 55°C, more preferably between 30°C and 45°C or between 30°C and 50°C.

[0068] Microorganisms can be cultured (e.g., maintained and/or grown) in liquid media and preferably are cultured, either continuously or intermittently, by conventional culturing methods such as standing culture, test tube culture, shaking culture (e.g., rotary shaking culture, shake flask culture, etc.), aeration spinner culture, or fermentation. In a preferred embodiment, the microorganisms are cultured in shake flasks. In a more preferred embodiment, the microorganisms are cultured in a fermentor (e.g., a fermentation process). Fermentation processes of the present invention include, but are not limited to, batch, fed-batch and continuous methods of fermentation. The phrase "batch process" or "batch fermentation" refers to a closed system in which the composition of media, nutrients, supplemental additives and the like is set at the beginning of the fermentation and not subject to alteration during the fermentation, however, attempts may be made to control such factors as pH and oxygen concentration to prevent excess media acidification and/or microorganism death. The phrase "fed-batch process" or "fed-batch" fermentation refers to a batch fermentation with the exception that one or more substrates or supplements are added (e.g., added in increments or continuously) as the fermentation progresses. The phrase "continuous process" or "continuous fermentation" refers to a system in which a defined fermentation media is added continuously to a fermentor and an equal amount of used or "conditioned" media is simultaneously removed, preferably for recovery of the desired fine chemical, e.g., lysine. A variety of such processes have been developed and are well-known in the art.

[0069] The phrase "culturing under conditions such that a desired fine chemical, e.g., lysine is produced" includes maintaining and/or growing microorganisms under conditions (e.g., temperature, pressure, pH, duration, etc.) appropriate or sufficient to obtain production of the desired fine chemical or to obtain desired yields of the particular fine chemical, e.g., lysine, being produced. For example, culturing is continued for a time sufficient to produce the desired amount of a fine chemical (e.g., lysine). Preferably, culturing is continued for a time sufficient to substantially reach maximal production of the fine chemical. In one embodiment, culturing is continued for about 12 to 24 hours. In another embodiment, culturing is continued for about 24 to 36 hours, 36 to 48 hours, 48 to 72 hours, 72 to 96 hours, 96 to 120 hours, 120 to 144 hours, or greater than 144 hours. In another embodiment, culturing is continued for a time sufficient to reach production yields of a fine chemical, for example, cells are cultured such that at least about 15 to 20 g/L of a fine chemical are produced, at least about 20 to 25 g/L of a fine chemical are produced, at least about 25 to 30 g/L of a fine chemical are produced, at least about 30 to 35 g/L of a fine chemical are produced, at least about 35 to 40 g/L of a fine chemical are produced, at least about 40 to 50 g/L of a fine chemical are produced, at least about 50 to 60 g/L of a fine chemical are produced, at least about 60 to 70 g/L of a fine chemical are produced, at least about 70 to 80 g/L of a fine chemical are produced, at least about 80 to 90 g/L of a fine chemical are produced, at least about 90 to 100 g/L of a fine chemical are produced, at least about 100 to 110 g/L of a fine chemical are produced, at least about 110 to 120 g/L of a fine chemical are produced, at least about 120 to 130 g/L of a fine chemical are produced, at least about 130 to 140 g/L of a fine chemical are produced, or at least about 140 to 160 g/L of a fine chemical are produced. In yet another embodiment, microorganisms are cultured under conditions such that a preferred yield of a fine chemical, for example, a yield within a range set forth above, is produced in about 24 hours, in about 36 hours, in about 40 hours, in about 48 hours, in about 72 hours, in about 96 hours, in about 108 hours, in about 122 hours, or in about 144 hours.

[0070] The methodology of the present invention can further include a step of recovering a desired fine chemical, e.g., lysine. The term "recovering" a desired fine chemical, e.g., lysine includes extracting, harvesting, isolating or purifying the compound from culture media. Recovering the compound can be performed according to any conventional isolation or purification methodology known in the art including, but not limited to, treatment with a conventional resin (e.g., anion or cation exchange resin, non-ionic adsorption resin, etc.), treatment with a conventional adsorbent (e.g., activated charcoal, silicic acid, silica gel, cellulose, alumina, etc.), alteration of pH, solvent extraction (e.g., with a conventional solvent such as an alcohol, ethyl acetate, hexane and the like), dialysis, filtration, concentration, crystallization, recrystallization, pH adjustment, lyophilization and the like. For example, a fine chemical, e.g., lysine, can be recovered from culture media by first removing the microorganisms from the culture. Media is then passed through or over a cation exchange resin to remove unwanted cations and then through or over an anion exchange resin to remove unwanted inorganic anions and organic acids having stronger acidities than the fine chemical of interest (e.g., lysine).

[0071] Preferably, a desired fine chemical of the present invention is "extracted", "isolated" or "purified" such that the resulting preparation is substantially free of other components (e.g., free of media components and/or fermentation byproducts). The language "substantially free of other components" includes preparations of desired compound in which the compound is separated (e.g., purified or partially purified) from media components or fermentation byproducts of the culture from which it is produced.

[0072] In one embodiment, the preparation has greater than about 80% (by dry weight) of the desired compound (e.g., less than about 20% of other media components or fermentation byproducts), more preferably greater than about 90% of the desired compound (e.g., less than about 10% of other media components or fermentation byproducts), still more preferably greater than about 95% of the desired compound (e.g., less than about 5% of other media components or fermentation byproducts), and most preferably greater than about 98-99% desired compound (e.g., less than about

1-2% other media components or fermentation byproducts).

[0073] In an alternative embodiment, the desired fine chemical, e.g., lysine, is not purified from the microorganism, for example, when the microorganism is biologically non-hazardous (e.g., safe). For example, the entire culture (or culture supernatant) can be used as a source of product (e.g., crude product). In one embodiment, the culture (or culture supernatant) supernatant is used without modification. In another embodiment, the culture (or culture supernatant) is concentrated. In yet another embodiment, the culture (or culture supernatant) is dried or lyophilized.

II. Methods of Producing A Fine Chemical Independent of Precursor

Feed Requirements

[0074] Depending on the biosynthetic enzyme or combination of biosynthetic enzymes manipulated, it may be desirable or necessary to provide (e.g., feed) microorganisms of the present invention at least one pentose phosphate pathway biosynthetic precursor such that fine chemicals, e.g., lysine, are produced. The term "pentose phosphate pathway biosynthetic precursor" or "precursor" includes an agent or compound which, when provided to, brought into contact with, or included in the culture medium of a microorganism, serves to enhance or increase pentose phosphate biosynthesis. In one embodiment, the pentose phosphate biosynthetic precursor or precursor is glucose. In another embodiment, the pentose phosphate biosynthetic precursor is fructose. The amount of glucose or fructose added is preferably an amount that results in a concentration in the culture medium sufficient to enhance productivity of the microorganism (e.g., a concentration sufficient to enhance production of a fine chemical e.g., lysine). Pentose phosphate biosynthetic precursors of the present invention can be added in the form of a concentrated solution or suspension (e.g., in a suitable solvent such as water or buffer) or in the form of a solid (e.g., in the form of a powder). Moreover, pentose phosphate biosynthetic precursors of the present invention can be added as a single aliquot, continuously or intermittently over a given period of time.

[0075] Providing pentose phosphate biosynthetic precursors in the pentose phosphate biosynthetic methodologies of the present invention, can be associated with high costs, for example, when the methodologies are used to produce high yields of a fine chemical. Accordingly, preferred methodologies of the present invention feature microorganisms having at least one biosynthetic enzyme or combination of biosynthetic enzymes (e.g., at least one pentose phosphate biosynthetic enzyme) manipulated such that lysine or other desired fine chemicals are produced in a manner independent of precursor feed. The phrase "a manner independent of precursor feed", for example, when referring to a method for producing a desired compound includes an approach to or a mode of producing the desired compound that does not depend or rely on precursors being provided (e.g., fed) to the microorganism being utilized to produce the desired compound. For example, microorganisms featured in the methodologies of the present invention can be used to produce fine chemicals in a manner requiring no feeding of the precursors glucose or fructose.

[0076] Alternative preferred methodologies of the present invention feature microorganisms having at least one biosynthetic enzyme or combination of biosynthetic enzymes manipulated such that L-Lysine or other fine chemicals are produced in a manner substantially independent of precursor feed. The phrase "a manner substantially independent of precursor feed" includes an approach to or a method of producing the desired compound that depends or relies to a lesser extent on precursors being provided (e.g., fed) to the microorganism being utilized. For example, microorganisms featured in the methodologies of the present invention can be used to produce fine chemicals in a manner requiring feeding of substantially reduced amounts of the precursors glucose or fructose.

[0077] Preferred methods of producing desired fine chemicals in a manner independent of precursor feed or alternatively, in a manner substantially independent of precursor feed, involve culturing microorganisms which have been manipulated (e.g., designed or engineered, for example, genetically engineered) such that expression of at least one pentose phosphate biosynthetic enzyme is modified. For example, in one embodiment, a microorganism is manipulated (e.g., designed or engineered) such that the production of at least one pentose phosphate biosynthetic enzyme is deregulated. In a preferred embodiment, a microorganism is manipulated (e.g., designed or engineered) such that it has a deregulated biosynthetic pathway, for example, a deregulated pentose phosphate biosynthesis pathway, as defined herein. In another preferred embodiment, a microorganism is manipulated (e.g., designed or engineered) such that at least one pentose phosphate biosynthetic enzyme, e.g., fructose-1,6-bisphosphatase is overexpressed.

III. High Yield Production Methodologies

[0078] A particularly preferred embodiment of the present invention is a high yield production method for producing a fine chemical, e.g., lysine, comprising culturing a manipulated microorganism under conditions such that lysine is produced at a significantly high yield. The phrase "high yield production method", for example, a high yield production method for producing a desired fine chemical, e.g., lysine, includes a method that results in production of the desired fine chemical at a level which is elevated or above what is usual for comparable production methods. Preferably, a high

yield production method results in production of the desired compound at a significantly high yield. The phrase "significantly high yield" includes a level of production or yield which is sufficiently elevated or above what is usual for comparable production methods, for example, which is elevated to a level sufficient for commercial production of the desired product (e.g., production of the product at a commercially feasible cost). In one embodiment, the invention features a high yield production method of producing lysine that includes culturing a manipulated microorganism under conditions such that lysine is produced at a level greater than 2 g/L, 10 g/L, 15 g/L, 20 g/L, 25 g/L, 30 g/L, 35 g/L, 40 g/L, 45 g/L, 50 g/L, 55 g/L, 60 g/L, 65 g/L, 70 g/L, 75 g/L, 80 g/L, 85 g/L, 90 g/L, 95 g/L, 100 g/L, 110 g/L, 120 g/L, 130 g/L, 140 g/L, 150 g/L, 160 g/L, 170 g/L, 180 g/L, 190 g/L, or 200 g/L.

[0079] The invention further features a high yield production method for producing a desired fine chemical, e.g., lysine, that involves culturing a manipulated microorganism under conditions such that a sufficiently elevated level of compound is produced within a commercially desirable period of time. In an exemplary embodiment, the invention features a high yield production method of producing lysine that includes culturing a manipulated microorganism under conditions such that lysine is produced at a level greater than 15-20 g/L in 5 hours. In another embodiment, the invention features a high yield production method of producing lysine that includes culturing a manipulated microorganism under conditions such that lysine is produced at a level greater than 25-40 g/L in 10 hours. In another embodiment, the invention features a high yield production method of producing lysine that includes culturing a manipulated microorganism under conditions such that lysine is produced at a level greater than 50-100 g/L in 20 hours. In another embodiment, the invention features a high yield production method of producing lysine that includes culturing a manipulated microorganism under conditions such that lysine is produced at a level greater than 140-160 g/L in 40 hours, for example, greater than 150 g/L in 40 hours. In another embodiment, the invention features a high yield production method of producing lysine that includes culturing a manipulated microorganism under conditions such that lysine is produced at a level greater than 130-160 g/L in 40 hours, for example, greater than 135, 145 or 150 g/L in 40 hours. Values and ranges included and/or intermediate within the ranges set forth herein are also intended to be within the scope of the present invention. For example, lysine production at levels of at least 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, and 150 g/L in 40 hours are intended to be included within the range of 140-150 g/L in 40 hours. In another example, ranges of 140-145 g/L or 145-150 g/L are intended to be included within the range of 140-150 g/L in 40 hours. Moreover, the skilled artisan will appreciate that culturing a manipulated microorganism to achieve a production level of, for example, "140-150 g/L in 40 hours" includes culturing the microorganism for additional time periods (e.g., time periods longer than 40 hours), optionally resulting in even higher yields of lysine being produced.

IV. Isolated Nucleic Acid Molecules and Genes

[0080] Another aspect of the present invention features isolated nucleic acid molecules that encode proteins (e.g., *C. glutamicum* proteins), for example, *Corynebacterium* pentose phosphate biosynthetic enzymes (e.g., *C. glutamicum* pentose phosphate enzymes) for use in the methods of the invention. In one embodiment, the isolated nucleic acid molecules used in the methods of the invention are fructose-1,6-bisphosphatase nucleic acid molecules.

[0081] The term "nucleic acid molecule" includes DNA molecules (e.g., linear, circular, cDNA or chromosomal DNA) and RNA molecules (e.g., tRNA, rRNA, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA. The term "isolated" nucleic acid molecule includes a nucleic acid molecule which is free of sequences which naturally flank the nucleic acid molecule (i.e., sequences located at the 5' and 3' ends of the nucleic acid molecule) in the chromosomal DNA of the organism from which the nucleic acid is derived. In various embodiments, an isolated nucleic acid molecule can contain less than about 10 kb, 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb, 0.1 kb, 50 bp, 25 bp or 10 bp of nucleotide sequences which naturally flank the nucleic acid molecule in chromosomal DNA of the microorganism from which the nucleic acid molecule is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular materials when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

[0082] The term "gene," as used herein, includes a nucleic acid molecule (e.g., a DNA molecule or segment thereof), for example, a protein or RNA-encoding nucleic acid molecule, that in an organism, is separated from another gene or other genes, by intergenic DNA (i.e., intervening or spacer DNA which naturally flanks the gene and/or separates genes in the chromosomal DNA of the organism). A gene may direct synthesis of an enzyme or other protein molecule (e.g., may comprise coding sequences, for example, a contiguous open reading frame (ORF) which encodes a protein) or may itself be functional in the organism. A gene in an organism, may be clustered in an operon, as defined herein, said operon being separated from other genes and/or operons by the intergenic DNA. Individual genes contained within an operon may overlap without intergenic DNA between said individual genes. An "isolated gene", as used herein, includes a gene which is essentially free of sequences which naturally flank the gene in the chromosomal DNA of the organism from which the gene is derived (i.e., is free of adjacent coding sequences which encode a second or distinct protein or RNA molecule, adjacent structural sequences or the like) and optionally includes 5' and 3' regulatory sequences, for

example promoter sequences and/or terminator sequences. In one embodiment, an isolated gene includes predominantly coding sequences for a protein (e.g., sequences which encode *Corynebacterium* proteins). In another embodiment, an isolated gene includes coding sequences for a protein (e.g., for a *Corynebacterium* protein) and adjacent 5' and/or 3' regulatory sequences from the chromosomal DNA of the organism from which the gene is derived (e.g., adjacent 5' and/or 3' *Corynebacterium* regulatory sequences). Preferably, an isolated gene contains less than about 10 kb, 5 kb, 2 kb, 1 kb, 0.5 kb, 0.2 kb, 0.1 kb, 50 bp, 25 bp or 10 bp of nucleotide sequences which naturally flank the gene in the chromosomal DNA of the organism from which the gene is derived.

[0083] In one aspect, the methods of the present invention features use of isolated fructose-1,6-bisphosphatase nucleic acid sequences or genes.

[0084] In a preferred embodiment, the nucleic acid or gene is derived from *Bacillus* (e.g., is *Corynebacterium*-derived). The term "derived from *Corynebacterium*" or "*Corynebacterium*-derived" includes a nucleic acid or gene which is naturally found in microorganisms of the genus *Corynebacterium*. Preferably, the nucleic acid or gene is derived from a microorganism selected from the group consisting of *Corynebacterium glutamicum*, *Corynebacterium acetoglutamicum*, *Corynebacterium acetoacidophilum* or *Corynebacterium thermoaminogenes*. In a particularly preferred embodiment, the nucleic acid or gene is derived from *Corynebacterium glutamicum* (e.g., is *Corynebacterium glutamicum*-derived). In yet another preferred embodiment, the nucleic acid or gene is a *Corynebacterium* gene homologue (e.g., is derived from a species distinct from *Corynebacterium* but having significant homology to a *Corynebacterium* gene of the present invention, for example, a *Corynebacterium* fructose-1,6-bisphosphatase gene).

[0085] Included within the scope of the present invention are bacterial-derived nucleic acid molecules or genes and/or *Corynebacterium*-derived nucleic acid molecules or genes (e.g., *Corynebacterium*-derived nucleic acid molecules or genes), for example, the genes identified by the present inventors, for example, *Corynebacterium* or *C. glutamicum* fructose-1,6-bisphosphatase genes. Further included within the scope of the present invention are bacterial-derived nucleic acid molecules or genes and/or *Corynebacterium*-derived nucleic acid molecules or genes (e.g., *C. glutamicum*-derived nucleic acid molecules or genes) (e.g., *C. glutamicum* nucleic acid molecules or genes) which differ from naturally-occurring bacterial and/or *Corynebacterium* nucleic acid molecules or genes (e.g., *C. glutamicum* nucleic acid molecules or genes), for example, nucleic acid molecules or genes which have nucleic acids that are substituted, inserted or deleted, but which encode proteins substantially similar to the naturally-occurring gene products of the present invention. In one embodiment, an isolated nucleic acid molecule comprises the nucleotide sequences set forth as SEQ ID NO:1, or encodes the amino acid sequence set forth in SEQ ID NO:2.

[0086] In another embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 60-65%, preferably at least about 70-75%, more preferably at least about 80-85%, and even more preferably at least about 90-95% or more identical to a nucleotide sequence set forth as SEQ ID NO:1. In another embodiment, an isolated nucleic acid molecule hybridizes under stringent conditions to a nucleic acid molecule having a nucleotide sequence set forth as SEQ ID NO:1. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent (e.g. high stringency) hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature.

[0087] A nucleic acid molecule of the present invention (e.g., a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1) can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, nucleic acid molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) or can be isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed based upon the sequence of SEQ ID NO:1. A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:1.

[0088] In another embodiment, an isolated nucleic acid molecule is or includes a fructose-1,6-bisphosphatase gene, or portion or fragment thereof. In one embodiment, an isolated fructose-1,6-bisphosphatase nucleic acid molecule or gene comprises the nucleotide sequence as set forth in SEQ ID NO:1 (e.g., comprises the *C. glutamicum* fructose-1,6-bisphosphatase nucleotide sequence). In another embodiment, an isolated fructose-1,6-bisphosphatase nucleic acid molecule or gene comprises a nucleotide sequence that encodes the amino acid sequence as set forth in SEQ ID NO:2 (e.g., encodes the *C. glutamicum* fructose-1,6-bisphosphatase amino acid sequence). In yet another embodiment, an isolated fructose-1,6-bisphosphatase nucleic acid molecule or gene encodes a homologue of the fructose-1,6-bisphosphatase protein having the amino acid sequence of SEQ ID NO:2. As used herein, the term "homologue" includes

a protein or polypeptide sharing at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identity with the amino acid sequence of a wild-type protein or polypeptide described herein and having a substantially equivalent functional or biological activity as said wild-type protein or polypeptide. For example, a fructose-1,6-bisphosphatase homologue shares at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identity with the protein having the amino acid sequence set forth as SEQ ID NO: 2 and has a substantially equivalent functional or biological activity (i.e., is a functional equivalent) of the protein having the amino acid sequence set forth as SEQ ID NO:2 (e.g., has a substantially equivalent pantothenate kinase activity). In a preferred embodiment, an isolated fructose-1,6-bisphosphatase nucleic acid molecule or gene comprises a nucleotide sequence that encodes a polypeptide as set forth in SEQ ID NO:2. In another embodiment, an isolated fructose-1,6-bisphosphatase nucleic acid molecule hybridizes to all or a portion of a nucleic acid molecule having the nucleotide sequence set forth in SEQ ID NO:1 or hybridizes to all or a portion of a nucleic acid molecule having a nucleotide sequence that encodes a polypeptide having the amino acid sequence of SEQ ID NOs:2. Such hybridization conditions are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, Ausubel et al., eds., John Wiley & Sons, Inc. (1995), sections 2, 4 and 6. Additional stringent conditions can be found in Molecular Cloning: A Laboratory Manual, Sambrook et al., Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), chapters 7, 9 and 11. A preferred, non-limiting example of stringent hybridization conditions includes hybridization in 4X sodium chloride/sodium citrate (SSC), at about 65-70°C (or hybridization in 4X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 1X SSC, at about 65-70 °C. A preferred, non-limiting example of highly stringent hybridization conditions includes hybridization in 1X SSC, at about 65-70°C (or hybridization in 1X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 0.3X SSC, at about 65-70°C. A preferred, non-limiting example of reduced stringency hybridization conditions includes hybridization in 4X SSC, at about 50-60°C (or alternatively hybridization in 6X SSC plus 50% formamide at about 40-45°C) followed by one or more washes in 2X SSC, at about 50-60°C. Ranges intermediate to the above-recited values, e.g., at 65-70°C or at 42-50°C are also intended to be encompassed by the present invention. SSPE (1X SSPE is 0.15 M NaCl, 10mM NaH₂PO₄, and 1.25 mM EDTA, pH 7.4) can be substituted for SSC (1X SSC is 0.15 M NaCl and 15 mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes each after hybridization is complete. The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, $T_m(^{\circ}\text{C}) = 2(\# \text{ of A} + \text{T bases}) + 4(\# \text{ of G} + \text{C bases})$. For hybrids between 18 and 49 base pairs in length, $T_m(^{\circ}\text{C}) = 81.5 + 16.6(\log_{10}[\text{Na}^+]) + 0.41 (\% \text{G+C}) - (600/\text{N})$, where N is the number of bases in the hybrid, and $[\text{Na}^+]$ is the concentration of sodium ions in the hybridization buffer ($[\text{Na}^+]$ for 1X SSC = 0.165 M). It will also be recognized by the skilled practitioner that additional reagents may be added to hybridization and/or wash buffers to decrease non-specific hybridization of nucleic acid molecules to membranes, for example, nitrocellulose or nylon membranes, including but not limited to blocking agents (e.g., BSA or salmon or herring sperm carrier DNA), detergents (e.g., SDS), chelating agents (e.g., EDTA), Ficoll, PVP and the like. When using nylon membranes, in particular, an additional preferred, non-limiting example of stringent hybridization conditions is hybridization in 0.25-0.5M NaH₂PO₄, 7% SDS at about 65°C, followed by one or more washes at 0.02M NaH₂PO₄, 1 % SDS at 65°C, see e.g., Church and Gilbert (1984) Proc. Natl. Acad. Sci. USA 81:1991-1995, (or, alternatively, 0.2X SSC, 1 % SDS). In another preferred embodiment, an isolated nucleic acid molecule comprises a nucleotide sequence that is complementary to a fructose-1,6-bisphosphatase nucleotide sequence as set forth herein (e.g., is the full complement of the nucleotide sequence set forth as SEQ ID NO:1).

[0089] A nucleic acid molecule of the present invention (e.g., a fructose-1,6-bisphosphatase nucleic acid molecule or gene), can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, nucleic acid molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) or can be isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed based upon the fructose-1,6-bisphosphatase nucleotide sequences set forth herein, or flanking sequences thereof. A nucleic acid of the invention (e.g., a fructose-1,6-bisphosphatase nucleic acid molecule or gene), can be amplified using cDNA, mRNA or alternatively, chromosomal DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques.

[0090] Yet another embodiment of the present invention features mutant fructose-1,6-bisphosphatase nucleic acid molecules or genes. The phrase "mutant nucleic acid molecule" or "mutant gene" as used herein, includes a nucleic acid molecule or gene having a nucleotide sequence which includes at least one alteration (e.g., substitution, insertion, deletion) such that the polypeptide or protein that may be encoded by said mutant exhibits an activity that differs from the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene. Preferably, a mutant nucleic acid molecule or mutant gene (e.g., a mutant fructose-1,6-bisphosphatase gene) encodes a polypeptide or protein having an increased activity (e.g., having an increased fructose-1,6-bisphosphatase activity) as compared to the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene, for example, when assayed under similar conditions

(e.g., assayed in microorganisms cultured at the same temperature). A mutant gene also can have an increased level of production of the wild-type polypeptide.

[0091] As used herein, an "increased or enhanced activity" or "increased or enhanced enzymatic activity" is one that is at least 5% greater than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene, preferably at least 5-10% more, more preferably at least 10-25% more and even more preferably at least 25-50%, 50-75% or 75-100% more than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene. Ranges intermediate to the above-recited values, e.g., 75-85%, 85-90%, 90-95%, are also intended to be encompassed by the present invention. Activity can be determined according to any well accepted assay for measuring activity of a particular protein of interest. Activity can be measured or assayed directly, for example, measuring an activity of a protein isolated or purified from a cell. Alternatively, an activity can be measured or assayed within a cell or in an extracellular medium.

[0092] It will be appreciated by the skilled artisan that even a single substitution in a nucleic acid or gene sequence (e.g., a base substitution that encodes an amino acid change in the corresponding amino acid sequence) can dramatically affect the activity of an encoded polypeptide or protein as compared to the corresponding wild-type polypeptide or protein. A mutant nucleic acid or mutant gene (e.g., encoding a mutant polypeptide or protein), as defined herein, is readily distinguishable from a nucleic acid or gene encoding a protein homologue, as described above, in that a mutant nucleic acid or mutant gene encodes a protein or polypeptide having an altered activity, optionally observable as a different or distinct phenotype in a microorganism expressing said mutant gene or nucleic acid or producing said mutant protein or polypeptide (i.e., a mutant microorganism) as compared to a corresponding microorganism expressing the wild-type gene or nucleic acid or producing said mutant protein or polypeptide. By contrast, a protein homologue has an identical or substantially similar activity, optionally phenotypically indiscernable when produced in a microorganism, as compared to a corresponding microorganism expressing the wild-type gene or nucleic acid. Accordingly it is not, for example, the degree of sequence identity between nucleic acid molecules, genes, protein or polypeptides that serves to distinguish between homologues and mutants, rather it is the activity of the encoded protein or polypeptide that distinguishes between homologues and mutants: homologues having, for example, low (e.g., 30-50% sequence identity) sequence identity yet having substantially equivalent functional activities, and mutants, for example sharing 99% sequence identity yet having dramatically different or altered functional activities.

V. Recombinant Nucleic Acid Molecules and Vectors

[0093] The present invention further features recombinant nucleic acid molecules (e.g., recombinant DNA molecules) that include nucleic acid molecules and/or genes described herein (e.g., isolated nucleic acid molecules and/or genes), preferably *Corynebacterium* genes, more preferably *Corynebacterium glutamicum* genes, even more preferably *Corynebacterium glutamicum* fructose-1,6-bisphosphatase genes.

[0094] The present invention further features vectors (e.g., recombinant vectors) that include nucleic acid molecules (e.g., isolated or recombinant nucleic acid molecules and/or genes) described herein. In particular, recombinant vectors are featured that include nucleic acid sequences that encode bacterial gene products as described herein, preferably *Corynebacterium* gene products, more preferably *Corynebacterium glutamicum* gene products (e.g., pentose phosphate enzymes, for example, fructose-1,6-bisphosphatase).

[0095] The term "recombinant nucleic acid molecule" includes a nucleic acid molecule (e.g., a DNA molecule) that has been altered, modified or engineered such that it differs in nucleotide sequence from the native or natural nucleic acid molecule from which the recombinant nucleic acid molecule was derived (e.g., by addition, deletion or substitution of one or more nucleotides). Preferably, a recombinant nucleic acid molecule (e.g., a recombinant DNA molecule) includes an isolated nucleic acid molecule or gene of the present invention (e.g., an isolated fructose-1,6-bisphosphatase gene) operably linked to regulatory sequences.

[0096] The term "recombinant vector" includes a vector (e.g., plasmid, phage, phasmid, virus, cosmid or other purified nucleic acid vector) that has been altered, modified or engineered such that it contains greater, fewer or different nucleic acid sequences than those included in the native or natural nucleic acid molecule from which the recombinant vector was derived. Preferably, the recombinant vector includes a fructose-1,6-bisphosphatase gene or recombinant nucleic acid molecule including such fructose-1,6-bisphosphatase gene, operably linked to regulatory sequences, for example, promoter sequences, terminator sequences and/or artificial ribosome binding sites (RBSs).

[0097] The phrase "operably linked to regulatory sequence(s)" means that the nucleotide sequence of the nucleic acid molecule or gene of interest is linked to the regulatory sequence(s) in a manner which allows for expression (e.g., enhanced, increased, constitutive, basal, attenuated, decreased or repressed expression) of the nucleotide sequence, preferably expression of a gene product encoded by the nucleotide sequence (e.g., when the recombinant nucleic acid molecule is included in a recombinant vector, as defined herein, and is introduced into a microorganism).

[0098] The term "regulatory sequence" includes nucleic acid sequences which affect (e.g., modulate or regulate) expression of other nucleic acid sequences. In one embodiment, a regulatory sequence is included in a recombinant

nucleic acid molecule or recombinant vector in a similar or identical position and/or orientation relative to a particular gene of interest as is observed for the regulatory sequence and gene of interest as it appears in nature, e.g., in a native position and/or orientation. For example, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence which accompanies or is adjacent to the gene of interest in the natural organism (e.g., operably linked to "native" regulatory sequences, for example, to the "native" promoter). Alternatively, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence which accompanies or is adjacent to another (e.g., a different) gene in the natural organism. Alternatively, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence from another organism. For example, regulatory sequences from other microbes (e.g., other bacterial regulatory sequences, bacteriophage regulatory sequences and the like) can be operably linked to a particular gene of interest.

[0099] In one embodiment, a regulatory sequence is a non-native or non-naturally-occurring sequence (e.g., a sequence which has been modified, mutated, substituted, derivatized, deleted including sequences which are chemically synthesized). Preferred regulatory sequences include promoters, enhancers, termination signals, anti-termination signals and other expression control elements (e.g., sequences to which repressors or inducers bind and/or binding sites for transcriptional and/or translational regulatory proteins, for example, in the transcribed mRNA). Such regulatory sequences are described, for example, in Sambrook, J., Fritsch, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in a microorganism (e.g., constitutive promoters and strong constitutive promoters), those which direct inducible expression of a nucleotide sequence in a microorganism (e.g., inducible promoters, for example, xylose inducible promoters) and those which attenuate or repress expression of a nucleotide sequence in a microorganism (e.g., attenuation signals or repressor sequences). It is also within the scope of the present invention to regulate expression of a gene of interest by removing or deleting regulatory sequences. For example, sequences involved in the negative regulation of transcription can be removed such that expression of a gene of interest is enhanced.

[0100] In one embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes a nucleic acid sequence or gene that encodes at least one bacterial gene product (e.g., a pentose phosphate biosynthetic enzyme, for example fructose-1,6-bisphosphatase) operably linked to a promoter or promoter sequence. Preferred promoters of the present invention include *Corynebacterium* promoters and/or bacteriophage promoters (e.g., bacteriophage which infect *Corynebacterium*). In one embodiment, a promoter is a *Corynebacterium* promoter, preferably a strong *Corynebacterium* promoter (e.g., a promoter associated with a biochemical housekeeping gene in *Corynebacterium* or a promoter associated with a glycolytic pathway gene in *Corynebacterium*). In another embodiment, a promoter is a bacteriophage promoter.

[0101] In another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes a terminator sequence or terminator sequences (e.g., transcription terminator sequences). The term "terminator sequences" includes regulatory sequences which serve to terminate transcription of a gene. Terminator sequences (or tandem transcription terminators) can further serve to stabilize mRNA (e.g., by adding structure to mRNA), for example, against nucleases.

[0102] In yet another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes sequences which allow for detection of the vector containing said sequences (i.e., detectable and/or selectable markers), for example, sequences that overcome auxotrophic mutations, for example, *ura3* or *ilvE*, fluorescent markers, and/or colorimetric markers (e.g., *lacZ*/ β -galactosidase), and/or antibiotic resistance genes (e.g., *amp* or *tet*).

[0103] In yet another embodiment, a recombinant vector of the present invention includes antibiotic resistance genes. The term "antibiotic resistance genes" includes sequences which promote or confer resistance to antibiotics on the host organism (e.g., *Bacillus*). In one embodiment, the antibiotic resistance genes are selected from the group consisting of *cat* (chloramphenicol resistance) genes, *tet* (tetracycline resistance) genes, *erm* (erythromycin resistance) genes, *neo* (neomycin resistance) genes and *spec* (spectinomycin resistance) genes. Recombinant vectors of the present invention can further include homologous recombination sequences (e.g., sequences designed to allow recombination of the gene of interest into the chromosome of the host organism). For example, *amyE* sequences can be used as homology targets for recombination into the host chromosome.

[0104] It will further be appreciated by one of skill in the art that the design of a vector can be tailored depending on such factors as the choice of microorganism to be genetically engineered, the level of expression of gene product desired and the like.

VI. Isolated Proteins

[0105] Another aspect of the present invention features isolated proteins (e.g., isolated pentose phosphate biosynthetic enzymes, for example isolated fructose-1,6-bisphosphatase). In one embodiment, proteins (e.g., isolated pentose phos-

phate enzymes, for example isolated fructose-1,6-bisphosphatase) are produced by recombinant DNA techniques and can be isolated from microorganisms of the present invention by an appropriate purification scheme using standard protein purification techniques. In another embodiment, proteins are synthesized chemically using standard peptide synthesis techniques.

[0106] An "isolated" or "purified" protein (e.g., an isolated or purified biosynthetic enzyme) is substantially free of cellular material or other contaminating proteins from the microorganism from which the protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. In one embodiment, an isolated or purified protein has less than about 30% (by dry weight) of contaminating protein or chemicals, more preferably less than about 20% of contaminating protein or chemicals, still more preferably less than about 10% of contaminating protein or chemicals, and most preferably less than about 5% contaminating protein or chemicals.

[0107] In a preferred embodiment, the protein or gene product is derived from *Corynebacterium* (e.g., is *Corynebacterium*-derived). The term "derived from *Corynebacterium*" or "*Corynebacterium*-derived" includes a protein or gene product which is encoded by a *Corynebacterium* gene. Preferably, the gene product is derived from a microorganism selected from the group consisting of *Corynebacterium glutamicum*, *Corynebacterium acetoglutamicum*, *Corynebacterium acetoacidophilum* or *Corynebacterium thermoaminogenes*. In a particularly preferred embodiment, the protein or gene product is derived from *Corynebacterium glutamicum* (e.g., is *Corynebacterium glutamicum*-derived). The term "derived from *Corynebacterium glutamicum*" or "*Corynebacterium glutamicum*-derived" includes a protein or gene product which is encoded by a *Corynebacterium glutamicum* gene. In yet another preferred embodiment, the protein or gene product is encoded by a *Corynebacterium* gene homologue (e.g., a gene derived from a species distinct from *Corynebacterium* but having significant homology to a *Corynebacterium* gene of the present invention, for example, a *Corynebacterium* fructose-1,6-bisphosphatase gene).

[0108] Included within the scope of the present invention are bacterial-derived proteins or gene products and/or *Corynebacterium*-derived proteins or gene products (e.g., *C. glutamicum*-derived gene products) that are encoded by naturally-occurring bacterial and/or *Corynebacterium* genes (e.g., *C. glutamicum* genes), for example, the genes identified by the present inventors, for example, *Corynebacterium* or *C. glutamicum* fructose-1,6-bisphosphatase genes. Further included within the scope of the present invention are bacterial-derived proteins or gene products and/or *Corynebacterium*-derived proteins or gene products (e.g., *C. glutamicum*-derived gene products) that are encoded bacterial and/or *Corynebacterium* genes (e.g., *C. glutamicum* genes) which differ from naturally-occurring bacterial and/or *Corynebacterium* genes (e.g., *C. glutamicum* genes), for example, genes which have nucleic acids that are mutated, inserted or deleted, but which encode proteins substantially similar to the naturally-occurring gene products of the present invention. For example, it is well understood that one of skill in the art can mutate (e.g., substitute) nucleic acids which, due to the degeneracy of the genetic code, encode for an identical amino acid as that encoded by the naturally-occurring gene. Moreover, it is well understood that one of skill in the art can mutate (e.g., substitute) nucleic acids which encode for conservative amino acid substitutions. It is further well understood that one of skill in the art can substitute, add or delete amino acids to a certain degree without substantially affecting the function of a gene product as compared with a naturally-occurring gene product, each instance of which is intended to be included within the scope of the present invention.

[0109] In a preferred embodiment, an isolated protein of the present invention (e.g., an isolated pentose phosphate biosynthetic enzyme, for example isolated fructose-1,6-bisphosphatase) has an amino acid sequence shown in SEQ ID NO:2. In other embodiments, an isolated protein of the present invention is a homologue of the protein set forth as SEQ ID NO:2, (e.g., comprises an amino acid sequence at least about 30-40% identical, preferably about 40-50% identical, more preferably about 50-60% identical, and even more preferably about 60-70%, 70-80%, 80-90%, 90-95% or more identical to the amino acid sequence of SEQ ID NO:2, and has an activity that is substantially similar to that of the protein encoded by the amino acid sequence of SEQ ID NO:2.

[0110] To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical positions/total # of positions x 100), preferably taking into account the number of gaps and size of said gaps necessary to produce an optimal alignment.

[0111] The comparison of sequences and determination of percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-68, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-77. Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul et al. (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST

program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997) Nucleic Acids Research 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller (1988) Comput Appl Biosci. 4:11-17. Such an algorithm is incorporated into the ALIGN program available, for example, at the GENESTREAM network server, IGH Montpellier, FRANCE (<http://vega.igh.cnrs.fr>) or at the ISREC server (<http://www.ch.embnet.org>). When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

[0112] In another preferred embodiment, the percent homology between two amino acid sequences can be determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. In yet another preferred embodiment, the percent homology between two nucleic acid sequences can be accomplished using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a gap weight of 50 and a length weight of 3.

[0113] This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents, Sequence Listing, Figures, and published patent applications cited throughout this application are incorporated herein by reference.

EXAMPLES

General Methodology:

[0114] Strains. *Corynebacterium glutamicum* ATCC 21526 was obtained from the American Type and Culture Collection (Manassas, USA). This homoserine auxotrophic strain excretes lysine during L-threonine limitation due to the bypass of concerted aspartate kinase inhibition. Precultures were grown in complex medium containing 5 g L⁻¹ of either fructose or glucose. For agar plates the complex medium was additionally amended with 12 g L⁻¹ agar. For the production of cells as inoculum for the tracer experiments and the tracer studies itself a minimal medium amended with 1 mg ml⁻¹ calcium pantothenate-HCl was used (Wittmann, C. and E. Heinzle. 2002. Appl. Environ. Microbiol. 68:5843-5859). In this medium concentrations of carbon source glucose or fructose, of the essential amino acids threonine, methionine and leucine and of citrate were varied as specified below.

[0115] Cultivation. Precultivation consisted of three steps involving (i) a starter cultivation in complex medium with cells from agar plate as inoculum, (ii) a short cultivation for adaption to minimal medium, and (iii) a prolonged cultivation on minimal medium with elevated concentrations of essential amino acids. Pre-cultures inoculated from agar plates were grown overnight in 100 ml baffled shake flasks on 10 ml complex medium. Afterwards cells were harvested by centrifugation (8800 g, 2 min, 30 °C), inoculated into minimal medium, and grown up to an optical density of 2 to obtain exponentially growing cells adapted to minimal medium. Afterwards cells were harvested by centrifugation (8800 g, 30 °C, and 2 min) including a washing step with sterile 0.9 % NaCl. They were then inoculated into 6 ml minimal medium in 50 ml baffled shake flasks with initial concentrations of 0.30 g L⁻¹ threonine, 0.08 g L⁻¹ methionine, 0.20 g L⁻¹ leucine, and 0.57 g L⁻¹ citrate. As carbon source 70 mM glucose or 80 mM fructose were added, respectively. Cells were grown until depletion of the essential amino acids, which was checked by HPLC analysis. At the end of the growth phase cells were harvested, and washed with sterile NaCl (0.9 %). Subsequently they were transferred into 4 ml minimal tracer medium in 25 ml baffled shake flasks for metabolic flux analysis under lysine producing conditions. The tracer medium did not contain any threonine, methionine, leucine and citrate. For each carbon source two parallel flasks were incubated containing (i) 40 mM [1-¹³C] labeled substrate, and (ii) 20 mM [¹³C₆] labeled substrate plus 20 mM of naturally labeled substrate, respectively. All cultivations were carried out on a rotary shaker (Inova 4230, New Brunswick, Edison, NJ, USA) at 30°C and 150 rpm.

[0116] Chemicals. 99% [1-¹³C] glucose, 99% [1-¹³C] fructose, 99% [¹³C₆] glucose and 99% [¹³C₆] fructose were purchased from Campro Scientific (Veenendaal, Netherlands). Yeast extract and tryptone were obtained from Difco Laboratories (Detroit, Michigan USA). All other applied chemicals were from Sigma (St. Louis, MI USA), Merck (Darmstadt, Germany) or Fluka (Buchs, Switzerland), respectively, and of analytical grade.

[0117] Substrate and product analysis. Cell concentration was determined by measurement of cell density at 660 nm (OD_{660nm}) using a photometer (Marsha Pharmacia biotech, Freiburg, Germany) or by gravimetry. The latter was determined by harvesting 10 ml of cells from cultivation broth at room temperature for 10 min at 3700 g, including a washing step with water. Washed cells were dried at 80 °C until weight constancy. The correlation factor (g biomass/OD_{660nm}) between dry cell dry mass and OD_{660nm} was determined as 0.353.

[0118] Concentrations of extracellular substrates and products were determined in cultivation supernatants, obtained via 3 min centrifugation at 16000 g. Fructose, glucose, sucrose, and trehalose were quantified by GC after derivatization into oxime trimethylsilyl derivatives. For this purpose a HP 6890 gas chromatograph (Hewlett Packard, Palo Alto, USA)

with an HP 5MS column (5 % phenyl-methyl-siloxane-diphenyldimethylpolysiloxane, 30 m x 250 μ m, Hewlett Packard, Paolo Alto, CA, USA), and a quadrupole mass selective detector with electron impact ionization at 70 eV (Agilent Technologies, Waldbronn, Germany) was applied. Sample preparation included lyophilization of the culture supernatant, dissolution in pyridine, and subsequent two-step derivatization of the sugars with hydroxylamine and (trimethylsilyl) trifluoroacetamide (BSTFA) (Macherey & Nagel, Düren, Germany) (13, 14). β -D-ribose was used as internal standard for quantification. The injected sample volume was 0.2 μ l. The time program for GC analysis was as follows: 150 °C (0 - 5 min), 8 °C min⁻¹ (5 - 25 min), 310 °C (25 - 35 min). Helium was used as carrier gas with a flow of 1.5 l min⁻¹. The inlet temperature was 310 °C and the detector temperature was 320 °C. Acetate, lactate, pyruvate, 2-oxoglutarate, and dihydroxyacetone were determined by HPLC utilizing an Aminex-HPX-87H Biorad Column (300 x 7.8 mm, Hercules, CA, USA) with 4 mM sulfuric acid as mobile phase at a flow rate of 0.8 ml min⁻¹, and UV-detection at 210 nm. Glycerol was quantified by enzymatic measurement (Boehringer, Mannheim, Germany). Amino acids were analyzed by HPLC (Agilent Technologies, Waldbronn, Germany) utilizing a Zorbax Eclipse-AAA column (150 x 4.6 mm, 5 μ m, Agilent Technologies, Waldbronn, Germany), with automated online derivatization (o-phthalaldehyde + 3-mercaptopropionic acid) at a flow rate of 2 ml min⁻¹, and fluorescence detection. Details are given in the instruction manual. -amino butyrate was used as internal standard for quantification.

[0119] ¹³C labeling analysis. The labeling patterns of lysine and trehalose in cultivation supernatants were quantified by GC-MS. Hereby single mass isotopomer fractions were determined. In the current work they are defined as M_0 (relative amount of non-labelled mass isotopomer fraction), M_1 (relative amount of single labelled mass isotopomer fraction) and corresponding terms for higher labelling. GC-MS analysis of lysine was performed after conversion into the t-butyl-dimethylsilyl (TBDMS) derivate as described previously (Rubino, F. M. 1989. J. Chromatogr. 473:125-133). Quantification of mass isotopomer distributions was performed in selective ion monitoring (SIM) mode for the ion cluster m/z 431-437. This ion cluster corresponds to a fragment ion, which is formed by loss of a t-butyl group from the derivatization residue, and thus includes the complete carbon skeleton of lysine (Wittmann, C., M. Hans and E. Heinzle. 2002. Analytical Biochem. 307:379-382). The labeling pattern of trehalose was determined from its trimethylsilyl (TMS) derivate as described previously (Wittmann, C., H. M. Kim and E. Heinzle. 2003. Metabolic flux analysis at miniaturized scale. submitted). The labeling pattern of trehalose was estimated via the ion cluster at m/z 361-367 corresponding to a fragment ion that contained an entire monomer unit of trehalose and thus a carbon skeleton equal to that of glucose 6-phosphate. All samples were measured first in scan mode therewith excluding isobaric interference between analyzed products and other sample components. All measurements by SIM were performed in duplicate. The experimental errors of single mass isotopomer fractions in the tracer experiments on fructose were 0.85% (M_0), 0.16 % (M_1), 0.27 % (M_2), 0.35 % (M_3), 0.45 % (M_4) for lysine on [1-¹³C] fructose, 0.87 % (M_0), 0.19 % (M_1), 0.44 % (M_2), 0.45 % (M_3), 0.88 % (M_4) for trehalose on [1-¹³C] fructose, and 0.44 % (M_0), 0.54 % (M_1), 0.34 % (M_2), 0.34 % (M_3), 0.19 % (M_4), 0.14 % (M_5) and 0.52 % (M_6) for trehalose on 50 % [¹³C₆] fructose, respectively. The experimental errors of MS measurements in glucose tracer experiments were 0.47 % (M_0), 0.44 % (M_1), 0.21 % (M_2), 0.26 % (M_3), 0.77 % (M_4) for lysine on [1-¹³C] glucose, 0.71 % (M_0), 0.85 % (M_1), 0.17 % (M_2), 0.32 % (M_3), 0.46 % (M_4) for trehalose on [1-¹³C] glucose, and 1.29 % (M_0), 0.50 % (M_1), 0.83 % (M_2), 0.84 % (M_3), 1.71 % (M_4), 1.84 % (M_5) and 0.58 % (M_6) for trehalose on 50 % [¹³C₆] glucose, respectively.

[0120] Metabolic modelling and parameter estimation. All metabolic simulations were carried out on a personal computer. Metabolic network of lysine-producing *C. glutamicum* was implemented in Matlab 6.1 and Simulink 3.0 (Mathworks, Inc., Natick, MA USA). The software implementation included an isotopomer model in Simulink to calculate the ¹³C labeling distribution in the network. For parameter estimation the isotopomer model was coupled with an iterative optimization algorithm in Matlab. Details on the applied computational tools are given by Wittmann and Heinzle (Wittmann, C. and E. Heinzle. 2002. Appl. Environ. Microbiol. 68:5843-5859).

[0121] The metabolic network was based on previous work and comprised glycolysis, pentose phosphate pathway (PPP), tricarboxylic acid (TCA) cycle, anaplerotic carboxylation of pyruvate, biosynthesis of lysine and other secreted products (Tab. 1), and anabolic fluxes from intermediary precursors into biomass. In addition uptake systems for glucose and fructose were alternatively implemented. Uptake of glucose involved phosphorylation to glucose 6-phosphate via a PTS (Ohnishi, J., S. Mitsuhashi, M. Hayashi, S. Ando, H. Yokoi, K. Ochiai and M. A. Ikeda. 2002. Appl. Microbiol. Biotechnol. 58:217-223). For fructose two uptake systems were considered: (i) uptake by PTS_{Fructose} and conversion of fructose into fructose 1,6-bisphosphatase via fructose 1-phosphate and (ii) uptake by PTS_{Mannose} leading to fructose 6-phosphate, respectively (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccain-Bousquet and N. D. Lindley. 1998. Eur. J. Biochem. 254:96-102). In addition fructose-1,6-bisphosphatase was implemented into the model to allow carbon flux in both directions in the upper glycolysis. Reactions regarded reversible were transaldolase and transketolases in the PPP. Additionally glucose 6-phosphate isomerase was considered reversible for the experiments on glucose, whereby the trehalose labeling sensitively reflected the reversibility of this enzyme. In contrast the reversibility of glucose 6-phosphate isomerase could not be determined on fructose. In fructose-grown cells, glucose 6-phosphate is exclusively formed from fructose 6-phosphate leading to identical labeling patterns for the two pools. Therefore interconversion between glucose 6-phosphate and fructose 6-phosphate by a reversible glucose 6-phosphate

isomerase does not result in labeling differences that could be used for the estimation of glucose 6-phosphate isomerase reversibility. The measured labeling of lysine and trehalose was not sensitive towards (i) the reversibility of the flux between the lumped pools of phosphoenolpyruvate/pyruvate and malate/oxaloacetate and (ii) the reversibility of malate dehydrogenase and fumarate hydratase in the TCA cycle. Accordingly these reactions were regarded irreversible. The labeling of alanine from a mixture of naturally labeled and [$^{13}\text{C}_6$] labeled substrate, which is sensitive for these flux parameters, was not available in this study. Based on previous results the glyoxylate pathway was assumed to be inactive (Wittmann, C. and E. Heinzle. 2002. Appl. Environ. Microbiol. 68:5843-5859).

[0122] Stoichiometric data on growth, product formation, and biomass composition of *C. glutamicum* together with mass spectrometric labeling data of secreted lysine and trehalose were used to calculate metabolic flux distributions. The set of fluxes that gave minimum deviation between experimental ($M_{i, \text{exp}}$) and simulated ($M_{i, \text{calc}}$) mass isotopomer fractions of lysine and trehalose of the two parallel experiments was taken as best estimate for the intracellular flux distribution. As described in the appendix the two networks of glucose-grown and fructose-grown cells were over determined. A least square approach was therefore possible. As error criterion a weighted sum of least squares (SLS) was used, where $S_{i, \text{exp}}$ is the standard deviation of the measurements (Eq. 1).

$$SLS = \sum_i \frac{(M_{i, \text{exp}} - M_{i, \text{calc}})^2}{S_{i, \text{exp}}^2} \quad (\text{Equation 1})$$

Multiple parameter initializations were applied to investigate whether an obtained flux distribution represented a global optimum. For all strains the glucose uptake flux during lysine production was set to 100 % and the other fluxes in the network are given as relative molar fluxes normalized to the glucose uptake flux.

[0123] Statistical evaluation. Statistical analysis of the obtained metabolic fluxes was carried out by a Monte-Carlo approach as described previously (Wittmann, C. and E. Heinzle. 2002. Appl. Environ. Microbiol. 68:5843-5859). For each strain, the statistical analysis was carried out by 100 parameter estimation runs, whereby the experimental data, comprising measured mass isotopomer ratios and measured fluxes, were varied statistically. From the obtained data 90 % confidence limits for the single parameters were calculated.

EXAMPLE I: LYSINE PRODUCTION BY *C. GLUTAMICUM* ON FRUCTOSE AND GLUCOSE

[0124] Metabolic fluxes of lysine producing *C. glutamicum* were analyzed in comparative batch cultures on glucose and fructose. For this purpose pre-grown cells were transferred into tracer medium and incubated for about 5 hours. The analysis of substrates and products at the beginning and the end of the tracer experiment revealed drastic differences between the two carbon sources. Overall 11.1 mM lysine was produced on glucose, whereas a lower concentration of only 8.6 mM was reached on fructose. During the incubation over 5 hours, the cell concentration increased from 3.9 g L⁻¹ to 6.0 g L⁻¹ (glucose) and from 3.5 g L⁻¹ to 4.4 g L⁻¹ (fructose). Due to the fact that threonine and methionine were not present in the medium, internal sources were probably utilized by the cells for biomass synthesis. The mean specific sugar uptake rate was higher on fructose (1.93 mmol g⁻¹ h⁻¹) compared to glucose (1.71 mmol g⁻¹ h⁻¹). As depicted in Table 1, the obtained yields of *C. glutamicum* ATCC 21526 differed drastically between fructose and glucose. This involved the main product lysine and various byproducts. Concerning lysine, the yield on fructose was 244 mmol mol⁻¹ and thus was lower compared to the yield on glucose (281 mmol mol⁻¹). Additionally the carbon source had a drastic influence on the biomass yield, which was reduced by almost 50% on fructose in comparison to glucose. The most significant influence of the carbon source on byproduct formation was observed for dihydroxyacetone, glycerol, and lactate. On fructose, accumulation of these byproducts was strongly enhanced. The yield for glycerol was 10 fold higher, whereas dihydroxyacetone and lactate secretion were increased by a factor of six. Dihydroxyacetone was the dominating byproduct on fructose. Due to the lower biomass yield a significantly reduced demand for anabolic precursors resulted for fructose-grown cells (Table 2).

Table 1: Biomass and metabolites in the stage of lysine production by *Corynebacterium glutamicum* ATCC 21526 from glucose (left) and fructose (right). Experimental yields are mean values of two parallel incubations on (i) 40 mM [^{1-13}C] labeled substrate and (ii) 20 mM [$^{13}\text{C}_6$] labeled substrate plus 20 mM naturally labeled substrate with corresponding deviations between the two incubations. All yields are given in (mmol product) (mol)⁻¹ except the yield for biomass, which is given in (mg of dry biomass) (mmol)⁻¹.

Yield	Lysine production on glucose	Lysine production on fructose
Biomass	54.1 ± 0.8	28.5 ± 0.0

(continued)

Yield	Lysine production on glucose	Lysine production on fructose
Lysine	281.0 ± 2.0	244.4 ± 23.3
Valine	0.1 ± 0.0	0.0 ± 0.0
Alanine	0.1 ± 0.0	0.4 ± 0.1
Glycine	6.6 ± 0.0	7.1 ± 0.4
Dihydroxyacetone	26.3 ± 15.3	156.6 ± 25.8
Glycerol	3.8 ± 2.4	38.4 ± 3.9
Trehalose	3.3 ± 0.5	0.9 ± 0.1
-Ketoglutarate	1.6 ± 0.4	6.5 ± 0.3
Acetate	45.1 ± 0.3	36.2 ± 5.7
Pyruvate	1.2 ± 0.4	2.1 ± 0.5
Lactate	7.1 ± 1.7	38.3 ± 3.5

Table 2. Anabolic demand of *Corynebacterium glutamicum* ATCC 21526 for intracellular metabolites in the stage of lysine production from glucose (left) and fructose (right). Experimental data are mean values of two parallel incubations on (i) [1-¹³C] labeled substrate and (ii) a 1:1 mixture of naturally labeled and [¹³C₆] substrate with deviation between the two incubations.

Precursor Demand* mmol (mol glucose) ⁻¹	Lysine production on glucose	Lysine production on fructose
Glucose 6-phosphate	11.09 ± 0.16	5.84 ± 0.05
Fructose 6-phosphate	3.84 ± 0.06	2.02 ± 0.02
Pentose 5-phosphate	47.50 ± 0.70	25.05 ± 0.21
Erythrose 4-phosphate	14.50 ± 0.22	7.64 ± 0.06
Glyceraldehyde 3-phosphate	6.98 ± 0.10	3.68 ± 0.03
3-Phosphoglycerate	59.95 ± 0.89	36.85 ± 0.31
Pyruvate/Phosphoenolpyruvate	107.80 ± 1.60	56.80 ± 0.48
-Ketoglutarate	92.51 ± 1.37	48.73 ± 0.41
Oxaloacetate	48.91 ± 0.72	45.76 ± 0.38
Acetyl CoA	135.30 ± 2.00	71.25 ± 0.60
Diaminopimelate+Lysine**	18.83 ± 0.28	9.92 ± 0.08

*) The estimation of precursor demands was based on the experimental biomass yield obtained for each strain (Tab. 1) and the biomass composition previously measured for *C. glutamicum* (Marx, A., A. A. de Graaf, W. Wiechert, L. Eggeling and H. Sahm. 1996. Biotechnol. Bioeng. 49:111-129).

**) Diaminopimelate and lysine are regarded as separate anabolic precursors. This is due to the fact that anabolic fluxes from pyruvate and oxaloacetate into diaminopimelate (cell wall) and lysine (protein) contribute in addition to the flux of lysine secretion to the overall flux through the lysine biosynthetic pathway.

EXAMPLE II: MANUAL INSPECTION OF ¹³C-LABELING PATTERNS IN TRACER EXPERIMENTS

[0125] Relative mass isotopomer fractions of secreted lysine and trehalose were quantified with GC-MS. These mass isotopomer fractions are sensitive towards intracellular fluxes and therefore display fingerprints for the fluxome of the investigated biological system. As shown in Figure 2, labeling patterns of secreted lysine and trehalose differed significantly between glucose and fructose-grown cells of *C. glutamicum*. The differences were found for both applied tracer labelings and for both measured products. This indicates substantial differences in the carbon flux pattern depending on the applied carbon source. As previously shown, mass isotopomer fractions from two parallel cultivations of *C. glutamicum* on a mixture of [1-¹³C] and [¹³C₆] glucose were almost identical (Wittmann, C., H. M. Kim and E. Heinzle. 2003. Metabolic flux analysis at miniaturized scale. submitted). Therefore, the differences observed can be clearly related to substrate specific differences in metabolic fluxes.

EXAMPLE III: ESTIMATION OF INTRACELLULAR FLUXES

[0126] A central issue of the performed studies was the comparative investigation of intracellular fluxes of *C. glutamicum* during lysine production on glucose and fructose as carbon source, respectively. For this purpose, the experimental data obtained from the tracer experiments were used to calculate metabolic flux distributions for each substrate applying the flux estimation software as described above. The parameter estimation was carried out by minimizing the deviation between experimental and calculated mass isotopomer fractions. The performed approach utilized metabolite balancing during each step of the optimization. This included (i) stoichiometric data on product secretion (Table 2) and (ii) stoichiometric data on anabolic demand for biomass precursors (Table 3). The set of intracellular fluxes that gave the minimum deviation between experimental and simulated labeling patterns was taken as best estimate for the intracellular flux distribution. For both scenarios, identical flux distributions were obtained with multiple initialization values, suggesting that global minima were identified. Obviously, good agreement between experimentally determined and calculated mass isotopomer ratios was achieved (Table 4).

Table 3: Relative mass isotopomer fractions of secreted lysine and trehalose of lysine producing *Corynebacterium glutamicum* ATCC 21526 cultivated on glucose and fructose, respectively. For both carbon sources two parallel tracer experiments on (i) [1-¹³C] labeled and (ii) a 1:1 mixture of naturally ¹³C labeled and [¹³C₆] labeled tracer substrate were carried out. Experimental GC/MS data (exp) and values predicted by the solution of the mathematical model corresponding to the optimized set of fluxes (calc). M₀ denotes the relative amount of non-labelled mass isotopomer fraction, M₁ the relative amount of the single labelled mass isotopomer fraction, and corresponding terms stand for higher labelling

	Lysine (on [1- ¹³ C] labeled substrate)					Trehalose (on [1- ¹³ C] labeled substrate)					Trehalose (on 50 % [¹³ C ₆] labeled substrate)			
	M ₀ M ₄	M ₁ M ₅	M ₂ M ₆	M ₃	M ₄	M ₀	M ₁	M ₂	M ₃	M ₄	M ₀	M ₁	M ₂	M ₃
glucose exp calc	0.234	0.360	0.247	0.110	0.037	0.110	0.551	0.216	0.094	0.023	0.271	0.114	0.087	
	0.115	0.069	0.066	0.279										
fructose exp calc	0.242	0.355	0.245	0.110	0.037	0.114	0.549	0.212	0.094	0.023	0.268	0.113	0.085	
	0.113	0.068	0.064	0.289										
	0.133	0.316	0.304	0.162	0.062	0.212	0.412	0.244	0.092	0.030	0.141	0.103	0.104	
	0.250	0.133	0.110	0.159										
	0.139	0.321	0.298	0.159	0.061	0.195	0.419	0.254	0.094	0.030	0.144	0.103	0.102	
	0.245	0.131	0.111	0.164										

EXAMPLE IV: METABOLIC FLUXES ON FRUCTOSE AND GLUCOSE DURING LYSINE PRODUCTION

[0127] The obtained intracellular flux distributions for lysine-producing *C. glutamicum* on glucose and fructose are shown in Figs. (4, 5). Obviously, the intracellular fluxes differed tremendously depending on the carbon source applied. On glucose, 62 % of the carbon flux was directed towards the PPP, whereas only 36 % were channeled through the glycolytic chain (Fig. 4) Due to this a relatively high amount, 124 % NADPH was generated by the PPP enzymes glucose 6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase. The situation on fructose was completely different (Fig. 5). The performed flux analysis revealed the in vivo activity of two PTS for uptake of fructose, whereby 92.3 % of fructose were taken up by fructose specific PTS_{Fructose}. A comparably small fraction of 7.7 % of fructose was taken up by PTS_{Mannose}. Thus, the majority of fructose entered the glycolysis at the level of fructose 1,6-bisphosphatase, whereas only a small fraction was channeled upstream at fructose 6-phosphate into the glycolytic chain. In comparison to glucose-grown cells, the PPP exhibited a dramatically reduced activity of only 14.4 %. Glucose 6-phosphate isomerase operated in opposite directions on the two carbon sources. In glucose-grown cells 36.2 % net flux were directed from glucose 6-phosphate to fructose 6-phosphate, whereas a backward net flux of 15.2 % was observed on fructose.

[0128] On fructose, the flux through glucose 6-phosphate isomerase and PPP was about twice as high as the flux through the PTS_{Mannose}. However this was not due to a gluconeogenic flux of carbon from fructose-1,6-bisphosphatase to fructose 6-phosphate, which could have supplied extra carbon flux towards the PPP. In fact flux through fructose 1,6-bisphosphatase catalyzing this reaction was zero. The metabolic reactions responsible for the additional flux towards the PPP are the reversible enzymes transaldolase and transketolase in the PPP. About 3.5 % of this additional flux was supplied by transketolase 2, which recycled carbon stemming from the PPP back into this pathway. Moreover 4.2 % of flux was directed towards fructose 6-phosphate and the PPP by the action of transaldolase.

[0129] Depending on the carbon source, completely different flux patterns in lysine producing *C. glutamicum* were also observed around the pyruvate node (Figs. 4, 5). On glucose the flux into the lysine pathway was 30.0 %, whereas a reduced flux of 25.4 % was found on fructose. The elevated lysine yield on glucose compared to fructose is the major reason for this flux difference, but also the higher biomass yield resulting in a higher demand for diaminopimelate for cell wall synthesis and lysine for protein synthesis contributes to it. The anaplerotic flux on glucose was 44.5 % and thus markedly higher compared to the flux on fructose (33.5 %). This is mainly due to the higher demand for oxaloacetate for lysine production, but also to the higher anabolic demands for oxaloacetate and 2-oxoglutarate on glucose. On the other hand, flux through pyruvate dehydrogenase was substantially lower on glucose (70.9 %) compared to fructose (95.2 %). This reduced carbon flux into the TCA cycle resulted in more than 30 % reduced fluxes through TCA cycle enzymes on glucose (Figs. 3, 4).

[0130] Statistical evaluation of the obtained fluxes by a Monte-Carlo approach was used to calculate 90 % confidence intervals for the determined flux parameters. As shown for various key fluxes in Table 5, the confidence intervals were generally narrow. As example the confidence interval for the flux through glucose 6-phosphate dehydrogenase was only 1.2 % for glucose-grown and 3.5 % for fructose-grown cells.

The chosen approach therefore allowed precise flux estimation. It can be concluded that the flux differences observed on glucose and fructose, respectively, are clearly caused by the applied carbon source.

[0131] It has to be noticed that the mean specific substrate uptake of 1.93 mmol g⁻¹ h⁻¹ on fructose was slightly higher than that of 1.77 mmol g⁻¹ h⁻¹ found on glucose. Due to this the absolute intracellular fluxes expressed in mmol g⁻¹ h⁻¹ are slightly increased in relation to glucose compared to the relative fluxes discussed above. The flux distributions of lysine producing *C. glutamicum* on fructose and glucose, respectively, are however so completely different, that all comparisons drawn above also hold for absolute carbon fluxes.

Table 4: Statistical evaluation of metabolic fluxes of lysine producing *Corynebacterium glutamicum* ATCC 21526 grown on fructose (left) and glucose (right) determined by ¹³C tracer studies with mass spectrometry and metabolite balancing: 90 % confidence intervals of key flux parameters were obtained by a Monte-Carlo approach including 100 independent parameter estimation runs for each substrate with statistically varied experimental data.

Flux parameter	Glucose	Fructose
Net Flux		
fructose uptake by PTS _{Frc}	-	[90.0 96.1]
fructose uptake by PTS _{Man}	-	[3.9 10.0]
glucose 6-phosphate isomerase	[35.7 36.8]	[13.4 16.9]
phosphofructokinase	[35.7 36.8]	-
fructose 1,6-bisphosphatase*	-	[-2.1 3.4]
fructose 1,6-bisphosphatase aldolase	[73.7 73.8]	[91.7 92.9]

(continued)

Flux parameter	Glucose	Fructose
glucose 6-phosphate dehydrogenase	[62.5 63.7]	[12.6 16.1]
transaldolase	[19.4 19.8]	[3.6 4.1]
transketolase 1	[19.4 19.8]	[3.6 4.1]
transketolase 2	[17.9 18.3]	[2.9 4.0]
glyceraldehyde 3-phosphate dehydrogenase	[158.1 164.5]	[163.3 174.6]
pyruvate kinase	[156.2 167.4]	[158.9 168.2]
pyruvate dehydrogenase	[69.5 72.5]	[87.1 102.3]
pyruvate carboxylase	[43.7 44.8]	[29.9 37.3]
citrate synthase	[51.2 54.8]	[76.5 91.5]
isocitrate dehydrogenase	[51.2 54.8]	[76.5 91.5]
oxoglutarate dehydrogenase	[41.6 45.6]	[70.9 86.0]
aspartokinase	[29.6 30.3]	[21.8 29.2]
Flux Reversibility**	[4.5 5.1]	-
glucose 6-phosphate isomerase	[4.3 4.9]	[14.5 18.2]
transaldolase	[0.0 0.0]	[0.0 0.1]
transketolase 1	[0.4 0.6]	[0.0 0.1]
transketolase 2		
* The negative flux for the lower confidence boundary is equal to a positive flux in the reverse direction (through phosphofructokinase).		
** Flux reversibility is defined as ratio of back flux to net flux.		

Discussion of Examples I- IV:

A. Substrate specific culture characteristics

[0132] Cultivation of lysine producing *C. glutamicum* on fructose and on glucose, respectively, revealed that growth and product formation strongly depend on the carbon source applied. Significantly reduced yields of lysine and biomass on fructose were previously also reported for another strain of *C. glutamicum*, where lysine and biomass yield were 30 % and 20 % less, respectively, compared to glucose (Kiefer, P., E. Heinzle and C. Wittmann. 2002. *J. Ind. Microbiol. Biotechnol.* 28:338-43). Cultivation of *C. glutamicum* and *C. melassecola* on fructose is linked to higher carbon dioxide production rates in comparison to glucose (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccagn-Bousquet and N. D. Lindley. 1998. *Eur. J. Biochem.* 254:96-102; Kiefer, P., E. Heinzle and C. Wittmann. 2002. *J. Ind. Microbiol. Biotechnol.* 28:338-43). This coincides with the elevated flux through the TCA cycle observed in the present work for this carbon source. Substrate specific differences were also observed for byproducts. The formation of trehalose was lower on fructose compared to glucose. This may be related to different entry points of glucose and fructose into glycolysis (Kiefer, P., E. Heinzle and C. Wittmann. 2002. *J. Ind. Microbiol. Biotechnol.* 28:338-43). Considering the uptake systems in *C. glutamicum*, utilization of glucose leads to the formation of the trehalose precursor glucose 6-phosphate, whereas fructose is converted into fructose 1,6-bisphosphatase and thus enters the central metabolism downstream from glucose 6-phosphate (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccagn-Bousquet and N. D. Lindley. 1998. *Eur. J. Biochem.* 254:96-102). Other byproducts such as dihydroxyacetone, glycerol, and lactate were strongly increased, when fructose was applied as carbon source. From the viewpoint of lysine production, this is not desired, because a substantial fraction of carbon is withdrawn from the central metabolism into the formed byproducts. The specific substrate uptake on fructose ($1.93 \text{ mmol g}^{-1} \text{ h}^{-1}$) was higher than on glucose ($1.77 \text{ mmol g}^{-1} \text{ h}^{-1}$). This result differs from a previous study on exponentially growing *C. melassecola* ATCC 17965 (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccagn-Bousquet and N. D. Lindley. 1998. *Eur. J. Biochem.* 254:96-102), where similar specific uptake rates on fructose and glucose were observed. The higher uptake rate for fructose observed in our study might be due to the fact that the studied strains are different. *C. melassecola* and *C. glutamicum* are related species, but might differ in certain metabolic properties. The strain studied in the present work was previously derived by classical strain optimization. This could have introduced mutations influencing substrate uptake. Another explanation

is the difference in cultivation conditions. Fructose might be more effectively utilized under conditions of limited growth and lysine production.

B. Metabolic flux distributions

[0133] The obtained intracellular flux distributions for lysine-producing *C. glutamicum* on glucose and fructose revealed tremendous differences. Statistical evaluation of the obtained fluxes revealed narrow 90% confidence intervals, so that the observed flux differences can be clearly attributed to the applied carbon sources. One of the most remarkable differences concerns the flux partitioning between glycolysis and PPP. On glucose 62.3 % of carbon was channeled through the PPP. The predominance of the PPP of lysine-producing *C. glutamicum* on this substrate has been previously observed in different studies (Marx, A., A. A. de Graaf, W. Wiechert, L. Eggeling and H. Sahm. 1996. *Biotechnol. Bioeng.* 49:111-129; Wittmann, C. and E. Heinzle. 2001. *Eur. J. Biochem.* 268:2441-2455; Wittmann, C. and E. Heinzle. 2002. *Appl. Environ. Microbiol.* 68:5843-5859). On fructose the flux into the PPP was reduced to 14.4 %. As identified by the performed metabolic flux analysis, this was mainly due to the unfavourable combination of the entry of fructose at the level of fructose 1,6-bisphosphatase and the inactivity of fructose-1,6-bisphosphatase. The observed inactivity of fructose-1,6-bisphosphatase agrees well with enzymatic measurements of *C. melassecola* ATCC 17965 during exponential growth on fructose and on glucose, respectively (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccagn-Bousquet and N. D. Lindley. 1998. *Eur. J. Biochem.* 254:96-102).

[0134] Surprisingly, the flux through glucose 6-phosphate isomerase and PPP was about twice as high as the flux through the PTS_{Mannose} when *C. glutamicum* was cultivated on fructose. Due to the inactivity of fructose-1,6-bisphosphatase this was not caused by a gluconeogenic flux. In fact, *C. glutamicum* possesses an operating metabolic cycle via fructose 6-phosphate, glucose 6-phosphate, and ribose 5-phosphate. Additional flux into the PPP was supplied by transketolase 2, which recycled carbon stemming from the PPP back into this pathway, and by the action of transaldolase, which redirected glyceraldehyde 3-phosphate back into the PPP, thus bypassing gluconeogenesis. This cycling activity may help the cell to overcome NADPH limitation on fructose. The drastically reduced flux arriving at glucose 6-phosphate for fructose-grown *C. glutamicum* might also explain the reduced formation of trehalose on this substrate (Kiefer, P., E. Heinzle and C. Wittmann. 2002. *J. Ind. Microbiol. Biotechnol.* 28:338-43). Glucose 6-phosphate isomerase operated in opposite directions depending on the carbon source. In glucose-grown net flux was directed from glucose 6-phosphate to fructose 6-phosphate, whereas an inverse net flux was observed on fructose. This underlines the importance of the reversibility of this enzyme for metabolic flexibility in *C. glutamicum*.

C. NADPH metabolism

[0135] The following calculations provide a comparison of the NADPH metabolism of lysine producing *C. glutamicum* on fructose and glucose. The overall supply of NADPH was calculated from the estimated flux through glucose 6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase, and isocitrate dehydrogenase. On glucose, the PPP enzymes glucose 6-phosphate dehydrogenase (62.0 %) and glucose 6-phosphate dehydrogenase (62.0 %) supplied the major fraction of NADPH. Isocitrate dehydrogenase (52.9 %) contributed only to a minor extent. A completely different contribution of PPP and TCA cycle to NADPH supply was observed on fructose, where isocitrate dehydrogenase (83.3 %) was the major source for NADPH. Glucose 6-phosphate dehydrogenase (14.4 %) and glucose 6-phosphate dehydrogenase (14.4 %) produced much less NADPH on fructose. NADPH is required for growth and formation of lysine. The NADPH requirement for growth was calculated from a stoichiometric demand of $11.51 \text{ mmol NADPH (g biomass)}^{-1}$, which was assumed to be identical for glucose and fructose (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccagn-Bousquet and N. D. Lindley. 1998. *Eur. J. Biochem.* 254:96-102), and the experimental biomass yield of the present work (Tab. 1). *C. glutamicum* consumed 62.3 % of NADPH for biomass production on glucose, which was much higher as compared to fructose as carbon source (32.8 %). The amount of NADPH required for product synthesis was determined from the estimated flux into lysine (Tab. 1) and the corresponding stoichiometric NADPH demand of $4 \text{ mol (mol lysine)}^{-1}$. It was 112.4 % for lysine production from glucose and 97.6 % for lysine production from fructose. The overall NADPH supply on glucose was significantly higher (176.9 %) compared to fructose (112.1 %), which can be mainly attributed to the increased PPP flux on glucose. The NADPH balance was almost closed on glucose. In contrast a significant apparent deficiency for NADPH of 18.3 % was observed on fructose. This raises the question for enzymes catalyzing metabolic reactions that could supply NADPH in addition to the above mentioned enzymes glucose 6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase and isocitrate dehydrogenase. A likely candidate seems NADPH-dependent malic enzyme. Previously an increased specific activity of this enzyme was detected on fructose-grown *C. melassecola* in comparison to glucose-grown cells (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccagn-Bousquet and N. D. Lindley. 1998. *Eur. J. Biochem.* 254:96-102). However, the flux through this particular enzyme could not be resolved by the experimental setup in the present work. Assuming malic enzyme as missing NADPH generating enzyme, a flux of 18.3 % would be sufficient to supply the apparently missing

NADPH. Detailed flux studies of *C. glutamicum* with glucose as carbon source revealed no significant activity of malic enzyme (Petersen, S., A. A. de Graaf, L. Eggeling, M. M611ney, W. Wiechert and H. Sahm. 2000. J. Biol. Chem. 75: 35932-35941). The situation on fructose might however be coupled to elevated in vivo activity of this enzyme.

5 D. NADH metabolism

[0136] On fructose *C. glutamicum* revealed increased activity of NADH forming enzymes. 421.2 % NADH were formed on fructose by glyceraldehyde 3-phosphate dehydrogenase, pyruvate dehydrogenase, 2-oxoglutarate dehydrogenase, and malate dehydrogenase. On glucose the NADH production was only 322.4 %. Additionally, the anabolic NADH demand was significantly lower on fructose than on glucose. The significantly enhanced NADH production coupled to a reduced metabolic demand could lead to an increased NADH/NAD ratio. For *C. melassecola* it was previously shown that fructose leads to increased NADH/NAD ratio compared to glucose (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccain-Bousquet and N. D. Lindley. 1998. Eur. J. Biochem. 254:96-102). This raises the question for NADH regenerating mechanisms during lysine production on fructose. Fructose-grown cells exhibited an enhanced secretion of dihydroxyacetone, glycerol, and lactate. The increased formation of dihydroxyacetone and glycerol could be due a higher NADH/NAD ratio. NADH was previously shown to inhibit glyceraldehyde dehydrogenase, so that overflow of dihydroxyacetone and glycerol might be related to a reduction of the flux capacity of this enzyme. The reduction of dihydroxyacetone to glycerol could additionally be favored by the high NADH/NAD ratio and thus contribute to regeneration of excess NADH. The NADH demanding lactate formation from pyruvate could have a similar background as the production of glycerol. In comparison to exponential growth, NADH excess under lysine producing conditions, characterized by relatively high TCA cycle activity and reduced biomass yield, might be even higher.

E. Potential targets for optimization of lysine-producing *C. glutamicum* on fructose

[0137] Based on the obtained flux patterns, several potential targets for the optimization of lysine production by *C. glutamicum* on fructose can be formulated. A central point is the supply of NADPH. Fructose-1,6-bisphosphatase is one target for increasing the supply of NADPH. Deregulation, e.g., amplification of its activity leads to a higher flux through the PPP, resulting in increased NADPH generation and increased lysine yield. An increase of the flux through the PPP via amplification of fructose 1,6-bisphosphatase is also be beneficial for aromatic amino acid production (Ikeda, M. 2003. Adv. Biochem. Eng. Biotechnol. 79: 1-36). The inactivity of fructose 1,6-bisphosphatase during growth on fructose is detrimental from the viewpoint of lysine production but not surprising, because this gluconeogenic enzyme is not required during growth on sugars and probably suppressed. In prokaryotes, this enzyme is under efficient metabolic control by e.g. fructose 1,6-bisphosphatase, fructose-2,6 bisphosphatase, metal ions and AMP (Skrypal, I. G. and O. V. lastrebova. 2002. Mikrobiol Z. 64:82-94). It is known that *C. glutamicum* can grow on acetate (Wendisch, V. F., A. A. de Graaf, H. Sahm H. and B. Eikmans. 2000. J. Bacteriol. 182:3088-3096), where this enzyme is essential to maintain gluconeogenesis. Another potential target to increase the flux through the PPP is the PTS for fructose uptake. Modification of flux partitioning between PTS_{Fructose} and PTS_{Mannose} could yield a higher proportion of fructose, which enters at the level of fructose 6-phosphate and thus also lead to an increased PPP flux. Additionally amplification of malic enzyme that probably contributes significantly to NADPH supply on fructose could be an interesting target.

[0138] Another bottleneck comprises the strong secretion of dihydroxyacetone, glycerol, and lactate. The formation of dihydroxyacetone and glycerol could be blocked by deregulation, e.g., deletion of the corresponding enzymes. The conversion of dihydroxyacetone phosphate to dihydroxyacetone could be catalyzed by a corresponding phosphatase. A dihydroxyacetone phosphatase has however yet not been annotated in *C. glutamicum* (see the National Center for Biotechnology Information (NCBI) Taxonomy website: <http://www3.ncbi.nlm.nih.gov/Taxonomy/>). This reaction may be also catalyzed by a kinase, e.g., glycerol kinase. Currently two entries in the genome data base of *C. glutamicum* relate to dihydroxyacetone kinase (see the National Center for Biotechnology Information (NCBI) Taxonomy website: <http://www3.ncbi.nlm.nih.gov/Taxonomy/>).

[0139] Lactate secretion can also be avoided by deregulation, e.g., knockout, of lactate dehydrogenase. Since glycerol and lactate formation could be important for NADH regeneration, negative effects on the overall performance of the organism can however not be excluded. In case carbon flux through the lower glycolytic chain is limited by the capacity of glyceraldehyde 3-phosphate dehydrogenase as previously speculated (Dominguez, H., C. Rollin, A. Guyonvarch, J. L. Guerquin-Kern, M. Coccain-Bousquet and N. D. Lindley. 1998. Eur. J. Biochem. 254:96-102), the suppression of dihydroxyacetone and glycerol production could eventually lead to an activation of fructose-1,6-bisphosphatase and a redirection of carbon flux through the PPP. It should be noticed that dihydroxyacetone is not reutilized during the cultivation of *C. glutamicum* and thus displays wasted carbon with respect to product synthesis, whereas this is not the case for lactate (Coccain-Bousquet, M. and N. D. Lindley. 1995. Enz. Microbiol. Technol. 17:260-267).

[0140] In one embodiment, deregulation of one or more of the above genes in combination is useful in the production of a fine chemical, e.g., lysine.

[0141] In addition, sucrose is also useful as carbon source for lysine production by *C. glutamicum*, e.g., used in conjunction with the methods of the invention. Sucrose is the major carbon source in molasses. As shown previously, the fructose unit of sucrose enters glycolysis at the level of fructose 1,6-bisphosphatase (Dominguez, H. and N. D. Lindley. 1996. Appl. Environ. Microbiol. 62:3878-3880). Therefore this part of the sucrose molecule - assuming an inactive fructose 1,6-bisphosphatase - probably does not enter into the PPP, so that NADPH supply in lysine producing strains could be limited.

EXAMPLE V: CONSTRUCTION OF PLASMID PCIS LYSC

[0142] The first step of strain construction calls for an allelic replacement of the *lysC* wild-type gene in *C. glutamicum* ATCC13032. In it, a nucleotide replacement in the *lysC* gene is carried out, so that, the resulting protein, the amino acid Thr in position 311 is replaced by an Ile. Starting from the chromosomal DNA from ATCC13032 as template for a PCR reaction and using the oligonucleotide primers SEQ ID NO:3 and SEQ ID NO:4, *lysC* is amplified by use of the Pfu Turbo PCR system (Stratagene USA) in accordance with the instructions of the manufacturer. Chromosomal DNA from *C. glutamicum* ATCC 13032 is prepared according to Tauch et al. (1995) Plasmid 33:168-179 or Eikmanns et al. (1994) Microbiology 140:1817-1828. The amplified fragment is flanked at its 5' end by a Sall restriction cut and at its 3' end by a MluI restriction cut. Prior to the cloning, the amplified fragment is digested by these two restriction enzymes and purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg).

SEQ ID NO:3

5'-GAGAGAGAGACGCGTCCCAGTGGCTGAGACGCATC-3'

SEQ ID NO:4

5'-CTCTCTCTGTCGACGAATTCAATCTTACGGCCTG-3'

[0143] The obtained polynucleotide is cloned through the Sall and MluI restriction cuts in pCLIK5 MCS with integrated SacB, referred to in the following as pCIS (SEQ ID NO: 5) and transformed in *E. coli* XL-1 blue. A selection for plasmid-carrying cells is accomplished by plating out on kanamycin (20 µg/mL) - containing LB agar (Lennox, 1955, Virology, 1: 190). The plasmid is isolated and the expected nucleotide sequence is confirmed by sequencing. The preparation of the plasmid DNA is carried out according to methods of and using materials of the company Quiagen. Sequencing reactions are carried out according to Sanger et al. (1977) Proceedings of the National Academy of Sciences USA 74: 5463-5467. The sequencing reactions are separated by means of ABI Prism 377 (PE Applied Biosystems, Weiterstadt) and analyzed. The obtained plasmid pCIS *lysC* is listed as SEQ ID NO:6.

EXAMPLE VI: MUTAGENESIS OF THE LYSC GENE FROM *C. GLUTAMICUM*

[0144] The targeted mutagenesis of the *lysC* gene from *C. glutamicum* is carried out using the QuickChange Kit (Company: Stratagene/USA) in accordance with the instructions of the manufacturer. The mutagenesis is carried out in the plasmid pCIS *lysC*, SEQ ID NO:6. The following oligonucleotide primers are synthesized for the replacement of thr 311 by 311 ile by use of the QuickChange method (Stratagene):

SEQ ID NO:7

5'-CGGCACACCGACATCATCTTCACCTGCCCTCGTTCCG -3'

SEQ ID NO:8

5'-CGGAACGAGGGCAGGTGAAGATGATGTCGGTGGTGCCG -3'

[0145] The use of these oligonucleotide primers in the QuickChange reaction leads, in the *lysC* gene (SEQ ID NO:9), to the replacement of the nucleotide in position 932 (from C to T). The resulting amino acid replacement Thr311Ile in the *lysC* gene is confirmed, after transformation in *E. coli* XL1-blue and plasmid preparation, by [a] sequencing reaction. The plasmid is given the designation pCIS *lysC* thr311ile and is listed as SEQ ID NO:10.

[0146] The plasmid pCIS *lysC* thr311ile is transformed in *C. glutamicum* ATCC13032 by means of electroporation, as described in Liebl, et al. (1989) FEMS Microbiology Letters 53:299-303. Modifications of the protocol are described in DE 10046870. The chromosomal arrangement of the *lysC* locus of individual transformants is checked using standard methods by Southern blot and hybridization, as described in Sambrook et al. (1989), Molecular Cloning. A Laboratory Manual, Cold Spring Harbor. It is thereby established that the transformants involved are those that have integrated the transformed plasmid by homologous recombination at the *lysC* locus. After growth of such colonies overnight in media containing no antibiotic, the cells are plated out on a saccharose CM agar medium (10% saccharose) and incubated at

30°C for 24 hours.

Because the sacB gene contained in the vector pCIS lysC thr311 ile converts saccharose into a toxic product, only those colonies can grow that have deleted the sacB gene by a second homologous recombination step between the wild-type lysC gene and the mutated gene lysC thr311 ile. During the homologous recombination, either the wild-type gene or the mutated gene together with the sacB gene can be deleted. If the sacB gene together with the wild-type gene is removed, a mutated transformant results.

[0147] Growing colonies are picked and examined for a kanamycin-sensitive phenotype. Clones with deleted SacB gene must simultaneously show kanamycin-sensitive growth behavior. Such kanamycin-sensitive clones are investigated in a shaking flask for their lysine productivity (see Example 6). For comparison, the non-treated *C. glutamicum* ATCC13032 is taken. Clones with an elevated lysine production in comparison to the control are selected, chromosomal DNA are recovered, and the corresponding region of the lysC gene is amplified by a PCR reaction and sequenced. One such clone with the property of elevated lysine synthesis and detected mutation in lysC at position 932 is designated as ATCC13032 lysC_{fr}.

EXAMPLE VII: PREPARATION OF THE PLASMID PK19 MOB SACB PEFTU FRUCTOSE-1,6 BISPHTHOSPHATASE

[0148] Chromosomal DNA from *C. glutamicum* ATCC 13032 is prepared according to Tauch et al. (1995) Plasmid 33: 168-179 or Eikmanns et al. (1994) Microbiology 140:1817-1828.

[0149] PCR 1: With the oligonucleotide primers SEQ ID NO 11 and SEQ ID NO 12, the chromosomal DNA as template, and Pfu Turbo polymerase (Company: Stratagene), a region lying upstream of the start codon of the elongation factor TU is amplified by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press.

SEQ ID NO 11

5'- TGGCCGTTACCCTGCGAATG -3'

and

SEQ ID NO 12

5'- TGTATGTCCTCCTGGACTTC -3'

The obtained DNA fragment of approximately 200 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer.

[0151] PCR 2: With the oligonucleotide primers SEQ ID NO 13 and SEQ ID NO 14, the chromosomal DNA as template, and Pfu Turbo polymerase (Company: Stratagene), the 5' region of the gene for fructose-1,6-bisphosphatase is amplified by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press.

SEQ ID NO 13

5'- GAAGTCCAGGAGGACATACAATGAACCTAAAGAACCCCGA -3'

and

SEQ ID NO 14

5'- ATCTACGTCGACCCAGGATGCCCTGGATTTC -3'

The obtained DNA fragment of approximately 740 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer.

[0153] PCR 3: With the oligonucleotide primers SEQ ID NO 15 and SEQ ID NO 16, the chromosomal DNA as template, and Pfu Turbo polymerase (Company: Stratagene), a region lying upstream of the start codon of fructose-1,6-bisphosphatase is amplified by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press.

SEQ ID NO 15

5'- TATCAACGCGTTCTTCATCGGTAGCAGCACC -3'

and

SEQ ID NO 16

5'- CATTGCGAGGGTAACGGCCACTGAAGGGCCTCCTGGG -3'

The obtained DNA fragment of approximately 720 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer.

[0155] PCR 4: With the oligonucleotide primers SEQ ID NO 17 and SEQ ID NO 14, the PCR products from PCR 1

and 2 as template, and Pfu Turbo polymerase (Company: Stratagene), a fusion PCR is carried out by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press.

The obtained DNA fragment of approximately 920 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer.

[0156] PCR 5: With the oligonucleotide primers SEQ ID NO 15 and SEQ ID NO 14, the PCR products from PCR 3 and 4 as template, and Pfu Turbo polymerase (Company: Stratagene), a fusion PCR is carried out by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press.

[0157] The obtained DNA fragment of approximately 1640 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer. Following this, it is cleaved using the restriction enzymes MluI and Sail (Roche Diagnostics, Mannheim) and the DNA fragment is purified using the GFX™ PCR DNA and Gel Band Purification Kit.

[0158] The vector pCIS is cut with the restriction enzymes MluI and Sail and a fragment of 4.3 kb size is isolated, after electrophoretic separation, by use of the GFX™ PCR DNA and Gel Band Purification Kit.

[0159] The vector fragment is ligated together with the PCR fragment from PCR 5 by use of the Rapid DNA Ligation Kit (Roche Diagnostics, Mannheim) in accordance with the instructions of the manufacturer and the ligation batch is transformed in competent *E. coli* XL-1 Blue (Stratagene, La Jolla, USA) according to standard methods, as described in Sambrook et al. (Molecular Cloning. A Laboratory Manual, Cold Spring Harbor, (1989)). A selection for plasmid-carrying cells is accomplished by plating out on kanamycin (20 µg/mL) - containing LB agar (Lennox, 1955, Virology, 1: 190).

[0160] The preparation of the plasmid DNA is carried out according to methods of and using materials of the company Qiagen. Sequencing reactions are carried out according to Sanger et al. (1977) Proceedings of the National Academy of Sciences USA 74:5463-5467. The sequencing reactions are separated by means of ABI Prism 377 (PE Applied Biosystems, Weiterstadt) and analyzed.

[0161] The resulting plasmid pCIS Peftu fructose-1,6-bisphosphatase is listed as SEQ ID NO: 17.

EXAMPLE VIII: PRODUCTION OF LYSINE

[0162] The plasmid pCIS Peftu fructose-1,6-bisphosphatase is transformed in *C. glutamicum* ATCC13032 lysCfbr by means of electroporation, as described in Liebl, et al. (1989) FEMS Microbiology Letters 53:299-303. Modifications of the protocol are described in DE 10046870. The chromosomal arrangement of the fructose-1,6-bisphosphatase gene locus of individual transformants is checked using standard methods by Southern blot and hybridization, as described in Sambrook et al. (1989), Molecular Cloning. A Laboratory Manual, Cold Spring Harbor. It is thereby established that the transformants involve those that have integrated the transformed plasmid by homologous recombination at the fructose-1,6-bisphosphatase gene locus. After growth of such colonies overnight in media containing no antibiotic, the cells are plated out on a saccharose CM agar medium (10% saccharose) and incubated at 30°C for 24 hours.

[0163] Because the sacB gene contained in the vector pCIS Peftu fructose-1,6-bisphosphatase converts saccharose into a toxic product, only those colonies can grow that have deleted the sacB gene by a second homologous recombination step between the wild-type fructose-1,6-bisphosphatase gene and the Peftu fructose-1,6-bisphosphatase fusion. During the homologous recombination, either the wild-type gene or the fusion together with the sacB gene can be deleted. If the sacB gene together with the wild-type gene is removed, a mutated transformant results.

[0164] Growing colonies are picked and examined for a kanamycin-sensitive phenotype. Clones with deleted SacB gene must simultaneously show kanamycin-sensitive growth behavior. Whether the desired replacement of the natural promoter by the Peftu promoter had also taken place is checked by means of the polymerase chain reaction (PCR). For this analysis, chromosomal DNA from the starting strain and the resulting clones is isolated. To this end, the respective clones are removed from the agar plate with a toothpick and suspended in 100 µL of H₂O and boiled up for 10 min at 95°C. In each case, 10 µL of the resulting solution is used as template in the PCR. Used as primers are oligonucleotides that are homologous to the Peftu promoter and to the fructose-1,6-bisphosphatase gene. The PCR conditions are selected as follows: initial denaturation: 5 min at 95°C; denaturation 30 sec at 95°C; hybridization 30 sec at 55°C; amplification 2 min at 72°C; 30 cycles; end extension 5 min at 72°C. In the batch with the DNA of the starting strain, no PCR product could form owing to the selection of the oligonucleotide. Only for clones that had completed the replacement of the natural promoter by Peftu through the 2nd recombination are a band with a size of 340 bp expected. Overall, of the tested clones, 2 clones are positive. The clones are designated as ATCC13032 lysCfbr Peftu fructose-1,6-bisphosphatase 1 and 2.

[0165] In order to investigate the effect of the Peftu fructose-1,6-bisphosphatase construct on the lysine production, the strains ATCC13032, ATCC13032 lysCfbr, and ATCC13032 lysCfbr Peftu fructose-1,6-bisphosphatase 1 are cultivated on CM plates (10.0 g/L D-glucose, 2.5 g/L NaCl, 2.0 g/L urea, 10.0 g/L bacto pepton (Difco), 5.0 g/L yeast extract

(Difco), 5.0 g/L beef extract (Difco), 22.0 g/L agar (Difco), autoclaved (20 min. 121°C)) for 2 days at 30°C. Subsequently, the cells are scraped off the plate and resuspended in saline. For the main culture, 10 mL of medium I and 0.5 g of autoclaved CaCO₃ (Riedel de Haen) are inoculated in a 100 mL Erlenmeyer flask with the cell suspension up to an OD₆₀₀ of 1.5 and incubated for 39 h on a [shaking incubator] of the type Infors AJ118 (Company: Infors, Bottmingen, Switzerland) at 220 rpm. Subsequently, the concentration of the lysine that separated out in the medium is determined.

Medium I:

[0166]

40 g/L saccharose
60 g/L Molasses (calculated with respect to 100% sugar content)
10 g/L (NH₄)₂SO₄
0.4 g/L MgSO₄*7H₂O
0.6 g/L KH₂PO₄
0.3 mg/L thiamine*HCl
1 mg/L biotin (from a 1 mg/mL sterile-filtered stock solution that is adjusted with NH₄OH to pH 8.0)
2 mg/L FeSO₄
2 mg/L MnSO₄

adjusted with NH₄OH to pH 7.8, autoclaved (121°C, 20 min).

In addition, vitamin B12 (hydroxycobalamin Sigma Chemicals) from a stock solution (200 µg/mL, sterile-filtered) is added up to a final concentration of 100 µg/L.

[0167] The determination of the amino acid concentration is conducted by means of high pressure liquid chromatography according to Agilent on an Agilent 1100 Series LC System HPLC. A precolumn derivatization with ortho-phthalaldehyde permits the quantification of the amino acids that are formed; the separation of the amino acid mixture takes place on a Hypersil AA column (Agilent).

EXAMPLE IX : PREPARATION OF THE PLASMID PCIS PSOD FRUCTOSE- 1,6 BISPHOSPHATASE

[0168] Chromosomal DNA from *C. glutamicum* ATGC 13032 is prepared according to Tauch et al. (1995) Plasmid 33 : 168-179 or Eikmanns et al. (1994) Microbiology 140 ; 1817-1828.

[0169] PCR 1 : With the oligonucleotide primers SEQ ID NO 18 and SEQ ID NO 19, the chromosomal DNA as template, and Pfu Turbo polymerase (Company: Stratagene), a region lying upstream of the start codon of the superoxid dismutase is amplified by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press.

SEQ ID NO 18

5'-tagctgccaattattccggg-3'

and

SEQ ID NO 19

5'-GGGTAAAAAATCCTTTTCGTA-3'

[0170] The obtained DNA fragment of approximately 200 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer.

[0171] PCR 2: With the oligonucleotide primers SEQ ID NO 20 and SEQ ID NO 21. the chromosomal DNA as template, and Pfu Turbo polymerase (Company: Stratagene), the 5' region of the gene for fructose-1,6-bisphosphatase is amplified by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press.

SEQ ID NO 20

5'-CCCGGAATAATTGGCAGCTACTGAAGGGCCTCCTGGG-3' and

SEQ ID NO 21

5'-TATCAACGCGTTCTTCATCGGTAGCAGCACC-3'

[0172] The obtained DNA fragment of approximately 720 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer.

[0173] PCR 3: With the oligonucleotide primers SEQ ID NO 22 and SEQ ID NO 23, the chromosomal DNA as template,

and Pfu Turbo polymerase (Company: Stratagene), a region lying upstream of the start codon of fructose-1, 6-bisphosphatase is amplified by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press.

5 SEO ID NO 22
 5'-TACGAAAGGATTTTTTACCCATGAACCTAAAGAACCCCGA-3'
 and
 SEO ID NO 23
 5'-ATCTACGTCGACCCAGGATGCCCTGGATTTC-3'

10 **[0174]** The obtained DNA fragment of approximately 740 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit, (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer.

15 **[0175]** PCR 4: With the oligonucleotide primers SEO ID NO 18 and SEQ ID NO 23, the PCR products from PCR 1 and 3 as template, and Pfu Turbo polymerase (Company: Stratagene), a fusion PCR is carried out by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press.

[0176] The obtained DNA fragment of approximately 930 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer.

20 **[0177]** PCR 5: With the oligonucleotide primers SEQ ID NO 21 and SEO ID NO 23, the PCR products from PCR 2 and 4 as template, and Pfu Turbo polymerase (Company: Stratagene), a fusion PCR is carried out by use of the polymerase chain reaction (PCR) according to standard methods, as described in Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press.

25 **[0178]** The obtained DNA fragment of approximately 1650 bp size is purified using the GFX™ PCR DNA and Gel Band Purification Kit (Amersham Pharmacia, Freiburg) in accordance with the instructions of the manufacturer. Following this, it is cleaved using the restriction enzymes MluI and Sall (Roche Diagnostics, Mannheim) and the DNA fragment is purified using the GFX™ PCR DNA and Gel Band Purification Kit.

[0179] The vector pCIS is cut with the restriction enzymes MluI and Sall and a fragment of 4, 3 kb size is isolated, after electrophoretic separation, by use of the GFX™ PCR DNA and Gel Band Purification Kit.

30 **[0180]** The vector fragment is ligated together with the PCR fragment from PCR 5 by use of the Rapid DNA Ligation Kit (Roche Diagnostics, Mannheim) in accordance with the instructions of the manufacturer and the ligation batch is transformed in competent *E. coli* XL-1 Blue (Stratagene, La Jolla, USA) according to standard methods, as described in Sambrook et al. (Molecular Cloning. A Laboratory Manual, Cold Spring Harbor, (1989)). A selection for plasmid-carrying cells is accomplished by plating out on kanamycin (20 µg/mL)-containing LB agar (Lennox, 1955, Virology, 1: 190).

35 **[0181]** The preparation of the plasmid DNA is carried out according to methods of and using materials of the company Qiagen. Sequencing reactions are carried out according to Sanger et al. (1977) Proceedings of the National Academy of Sciences USA 74: 5463-5467. The sequencing reactions are separated by means of ABI Prism 377 (PE Applied Biosystems, Weiterstadt) and analyzed.

40 **[0182]** The resulting plasmid pCIS Psod fructose-1,6-bisphosphatase is listed as SEQ ID NO : 24.

EXAMPLE X: PRODUCTION OF LYSINE

45 **[0183]** The plasmid pCIS Psod fructose-1,6-bisphosphatase is transformed in *C. glutamicum* ATCC13032 lysC⁺ by means of electroporation, as described in Liebl, et al. (1989) FEMS Microbiology Letters 53: 299-303. Modifications of the protocol are described in DE 10046870. The chromosomal arrangement of the fructose-1,6- bisphosphatase gene locus of individual transformants is checked using standard methods by Southern blot and hybridization, as described in Sambrook et al. (1989), Molecular Cloning. A Laboratory Manual, Cold Spring Harbor, It is thereby established that the transformants involve those that have integrated the transformed plasmid by homologous recombination at the fructose-1, 6-bisphosphatase gene locus. After growth of such colonies overnight in media containing no antibiotic, the

50 cells are plated out on a saccharose CM agar medium (10% saccharose) and incubated at 30°C for 24 hours.
 [0184] Because the sacB gene contained in the vector pCIS Psod fructose-1,6-bisphosphatase converts saccharose into a toxic product, only those colonies can grow that have deleted the sacB gene by a second homologous recombination step between the wild-type fructose-1, 6-bisphosphatase gene and the Psod fructose-1,6- bisphosphatase fusion. During the homologous recombination, either the wild-type gene or the fusion together with the sacB gene can be deleted. If

55 the sacB gene together with the wild-type gene is removed, a mutated transformant results.
 [0185] Growing colonies are picked and examined for a kanamycin-sensitive phenotype, Clones with deleted SacB gene must simultaneously show kanamycin- sensitive growth behavior. Whether the desired replacement of the natural promoter by the Psod promoter had also taken place is checked by means of the polymerase chain reaction (PCR). For

this analysis, chromosomal DNA from the starting strain and the resulting clones is isolated. To this end, the respective clones are removed from the agar plate with a toothpick and suspended in 100 gL of H₂O and boiled up for 10 min at 95°C. In each case, 10 gL of the resulting solution is used as template in the PCR. Used as primers are oligonucleotides that are homologous to the Psod promoter and to the fructose-1, 6-bisphosphatase gene. The PCR conditions are selected as follows : initial denaturation: 5 min at 95°C; denaturation 30 sec at 95°C ; hybridization 30 sec at 55°C ; amplification 2 min at 72°C ; 30 cycles; end extension 5 min at 72°C. In the batch with the DNA of the starting strain, no PCR product could form owing to the selection of the Only three clones that had completed the replacement of the natural promoter by Psod through the 2nd recombination are a band with a size of 350 bp expected. Overall, of the tested clones, 3 clones are positive. The clones are designated as ATCC13032 lysCfbr Psod fructose-1, 6-bisphosphatase 1, 2 and 3.

[0186] In order to investigate the effect of the Psod fructose-, 1, 6-bisphosphatase construct on the lysine production, the strains ATCC13032, ATCC13032 lysCfbr, and ATCC13032 lysCfbr Psod fructose-1, 6-bisphosphatase 1 are cultivated on CM plates (10.0 g/L D-glucose, 2.5 g/L NaCl, 2.0 g/L urea, 10.0 g/L bacto pepton (Difco), 5.0 g/L yeast extract (Difco), 5.0 g/L beef extract (Difco), 22.0 g/L agar (Difco), autoclaved (20 min. 121 °C)) for 2 days at 30°C. Subsequently, the cells are scraped off the plate and resuspended in saline. For the main culture, 10 mL of medium I and 0.5 g of autoclaved CaCO₃ (Riedel de Haen) are inoculated in a 100 mL Erlenmeyer flask with the cell suspension up to an OD₆₀₀ of 1.5 and incubated for 39 h on a shaking incubator of the type Infors AJ118 (Company: Infors, Bottmingen, Switzerland) at 220 rpm.

[0187] Subsequently, the concentration of the lysine that separated out in the medium is determined.

Medium I :

[0188]

40 g/L saccharose
60 g/L Molasses (calculated with respect to 100% sugar content)
10 g/L (NH₄)₂SO₄
0.4 g/L MgSO₄*7H₂O
0.6 g/L KH₂PO₄
0.3 mg/L thiamine*HCl
1 mg/L biotin (from a 1 mg/mL sterile-filtered stock solution that is adjusted with NH₄OH to pH 8. 0)
2mg/L FeSO₄
2 mg/L MnSO₄
adjusted with NH₄OH to pH 7. 8, autoclaved (121°C, 20 min).

[0189] In addition, vitamin B12 (hydroxycobalamin'Sigma Chemicals) from a stock solution (200 µg/mL, sterile-filtered) is added up to a final concentration of 100 µg/L.

[0190] The determination of the amino acid concentration is conducted by means of high pressure liquid chromatography according to Agilent on an Agilent 1100 Series LC System HPLC. A precolumn derivatization with ortho-phthalaldehyde permits the quantification of the amino acids that are formed; the separation of the amino acid mixture takes place on a Hypersil AA column (Agilent).

Equivalents

[0191] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

<110> BASF AKTIENGESELLSCHAFT et al.

<120> METHODS FOR THE PREPARATION OF A
FINE CHEMICAL BY FERMENTATION

<130> BGI-158PC2

<150> PCT/IB2003/006456

<151> 2003-12-18

<160> 24

<170> FastSEQ for windows Version 4.0

<210> 1

<211> 1070

<212> DNA

<213> Corynebacterium glutamicum

<220>

<221> CDS

<222> (22)...(1029)

<400> 1

```

gtgccccagg aggcccttca g atg aac cta aag aac ccc gaa acg cca gac 51
                        Met Asn Leu Lys Asn Pro Glu Thr Pro Asp
                        1                               5          10

cgt aac ctt gct atg gag ctg gtg cga gtt acg gaa gca gct gca ctg 99
Arg Asn Leu Ala Met Glu Leu Val Arg Val Thr Glu Ala Ala Ala Leu
                        15                               20          25

gct tct gga cgt tgg gtt gga cgt ggc atg aag aat gaa ggc gac ggt 147
Ala Ser Gly Arg Trp Val Gly Arg Gly Met Lys Asn Glu Gly Asp Gly
                        30                               35          40

gcc gct gtt gac gcc atg cgc cag ctc atc aac tca gtg acc atg aag 195
Ala Ala Val Asp Ala Met Arg Gln Leu Ile Asn Ser Val Thr Met Lys
                        45                               50          55

ggc gtc gtt gtt atc ggc gag ggc gaa aaa gac gaa gct cca atg ctg 243
Gly Val Val Val Ile Gly Gly Gly Glu Lys Asp Glu Ala Pro Met Leu
                        60                               65          70

tac aac ggc gaa gag gtc gga acc ggc ttt gga cct gag gtt gat atc 291
Tyr Asn Gly Glu Glu Val Gly Thr Gly Phe Gly Pro Glu Val Asp Ile
                        75                               80          85

gca gtt gac cca gtt gac ggc acc acc ctg atg gct gag ggt cgc ccc 339
Ala Val Asp Pro Val Asp Gly Thr Thr Leu Met Ala Glu Gly Arg Pro
                        95                               100          105

aac gca att tcc att ctc gca gct gca gag cgt ggc acc atg tac gat 387
Asn Ala Ile Ser Ile Leu Ala Ala Ala Glu Arg Gly Thr Met Tyr Asp
                        110                              115          120

cca tcc tcc gtc ttc tac atg aag aag atc gcc gtg gga cct gag gcc 435
Pro Ser Ser Val Phe Tyr Met Lys Lys Ile Ala Val Gly Pro Glu Ala
                        125                              130          135

gca ggc aag atc gac atc gaa gct cca gtt gcc cac aac atc aac gcg 483
Ala Gly Lys Ile Asp Ile Glu Ala Pro Val Ala His Asn Ile Asn Ala
                        140                              145          150

gtg gca aag tcc aag gga atc aac cct tcc gac gtc acc gtt gtc gtg 531
Val Ala Lys Ser Lys Gly Ile Asn Pro Ser Asp Val Thr Val Val Val
                        155                              160          165

ctt gac cgt cct cgc cac atc gaa ctg atc gca gac att cgt cgt gca 579
Leu Asp Arg Pro Arg His Ile Glu Leu Ile Ala Asp Ile Arg Arg Ala
                        175                              180          185

ggc gca aag gtt cgt ctc atc tcc gac ggc gac gtt gca ggt gca gtt 627

```

EP 1 939 296 A2

	Gly	Ala	Lys	Val	Arg	Leu	Ile	Ser	Asp	Gly	Asp	Val	Ala	Gly	Ala	Val	
				190					195					200			
5	gca	gca	gct	cag	gat	tcc	aac	tcc	gtg	gac	atc	atg	atg	ggc	acc	ggc	675
	Ala	Ala	Ala	Gln	Asp	Ser	Asn	Ser	Val	Asp	Ile	Met	Met	Gly	Thr	Gly	
			205					210					215				
	gga	acc	cca	gaa	ggc	atc	atc	act	gcg	tgc	gcc	atg	aag	tgc	atg	ggc	723
	Gly	Thr	Pro	Glu	Gly	Ile	Ile	Thr	Ala	Cys	Ala	Met	Lys	Cys	Met	Gly	
			220				225					230					
10	ggc	gaa	atc	cag	ggc	atc	ctg	gcc	cca	atg	aac	gat	ttc	gag	cgc	cag	771
	Gly	Glu	Ile	Gln	Gly	Ile	Leu	Ala	Pro	Met	Asn	Asp	Phe	Glu	Arg	Gln	
						240					245					250	
	aag	gca	cac	gac	gct	ggc	ctg	gtt	ctt	gat	cag	gtt	ctg	cac	acc	aac	819
	Lys	Ala	His	Asp	Ala	Gly	Leu	Val	Leu	Asp	Gln	Val	Leu	His	Thr	Asn	
					255				260						265		
15	gat	ctg	gtg	agc	tcc	gac	aac	tgc	tac	ttc	gtg	gca	acc	ggc	gtg	acc	867
	Asp	Leu	Val	Ser	Ser	Asp	Asn	Cys	Tyr	Phe	Val	Ala	Thr	Gly	Val	Thr	
				270				275						280			
	aac	ggc	gac	atg	ctc	cgt	ggc	gtt	tcc	tac	cgc	gca	aac	ggc	gca	acc	915
	Asn	Gly	Asp	Met	Leu	Arg	Gly	Val	Ser	Tyr	Arg	Ala	Asn	Gly	Ala	Thr	
			285				290						295				
20	acc	cgt	tcc	ctg	gtt	atg	cgc	gca	aag	tca	ggc	acc	atc	cgc	cac	atc	963
	Thr	Arg	Ser	Leu	Val	Met	Arg	Ala	Lys	Ser	Gly	Thr	Ile	Arg	His	Ile	
			300				305					310					
	gag	tct	gtc	cac	cag	ctg	tcc	aag	ctg	cag	gaa	tac	tcc	gtg	gtt	gac	1011
	Glu	Ser	Val	His	Gln	Leu	Ser	Lys	Leu	Gln	Glu	Tyr	Ser	Val	Val	Asp	
						320					325					330	
25	tac	acc	acc	gcg	acc	taa	gagctcttag	ttcgaaaaac	cgccggccat								1059
	Tyr	Thr	Thr	Ala	Thr	*											
					335												
30	tggtggtcggc	g															1070
	<210>	2															
	<211>	335															
	<212>	PRT															
	<213>	Corynebacterium glutamicum															
35	<400>	2															
	Met	Asn	Leu	Lys	Asn	Pro	Glu	Thr	Pro	Asp	Arg	Asn	Leu	Ala	Met	Glu	
	1				5					10					15		
	Leu	Val	Arg	Val	Thr	Glu	Ala	Ala	Ala	Leu	Ala	Ser	Gly	Arg	Trp	Val	
				20					25					30			
	Gly	Arg	Gly	Met	Lys	Asn	Glu	Gly	Asp	Gly	Ala	Ala	Val	Asp	Ala	Met	
			35					40					45				
40	Arg	Gln	Leu	Ile	Asn	Ser	Val	Thr	Met	Lys	Gly	Val	Val	Val	Ile	Gly	
		50					55					60					
	Glu	Gly	Glu	Lys	Asp	Glu	Ala	Pro	Met	Leu	Tyr	Asn	Gly	Glu	Glu	Val	
		65				70					75				80		
	Gly	Thr	Gly	Phe	Gly	Pro	Glu	Val	Asp	Ile	Ala	Val	Asp	Pro	Val	Asp	
				85					90						95		
	Gly	Thr	Thr	Leu	Met	Ala	Glu	Gly	Arg	Pro	Asn	Ala	Ile	Ser	Ile	Leu	
				100					105					110			
45	Ala	Ala	Ala	Glu	Arg	Gly	Thr	Met	Tyr	Asp	Pro	Ser	Ser	Val	Phe	Tyr	
			115					120					125				
	Met	Lys	Lys	Ile	Ala	Val	Gly	Pro	Glu	Ala	Ala	Gly	Lys	Ile	Asp	Ile	
							135					140					
	Glu	Ala	Pro	Val	Ala	His	Asn	Ile	Asn	Ala	Val	Ala	Lys	Ser	Lys	Gly	
						150					155					160	
50	Ile	Asn	Pro	Ser	Asp	Val	Thr	Val	Val	Val	Leu	Asp	Arg	Pro	Arg	His	
					165					170					175		
	Ile	Glu	Leu	Ile	Ala	Asp	Ile	Arg	Arg	Ala	Gly	Ala	Lys	Val	Arg	Leu	
				180					185					190			
	Ile	Ser	Asp	Gly	Asp	Val	Ala	Gly	Ala	Val	Ala	Ala	Ala	Gln	Asp	Ser	
								200					205				
	Asn	Ser	Val	Asp	Ile	Met	Met	Gly	Thr	Gly	Gly	Thr	Pro	Glu	Gly	Ile	
						215						220					
55	Ile	Thr	Ala	Cys	Ala	Met	Lys	Cys	Met	Gly	Gly	Glu	Ile	Gln	Gly	Ile	
						230					235					240	

EP 1 939 296 A2

Leu Ala Pro Met Asn Asp Phe Glu Arg Gln Lys Ala His Asp Ala Gly
 245 250 255
 Leu Val Leu Asp Gln Val Leu His Thr Asn Asp Leu Val Ser Ser Asp
 260 265 270
 Asn Cys Tyr Phe Val Ala Thr Gly Val Thr Asn Gly Asp Met Leu Arg
 275 280 285
 Gly Val Ser Tyr Arg Ala Asn Gly Ala Thr Thr Arg Ser Leu Val Met
 290 295 300
 Arg Ala Lys Ser Gly Thr Ile Arg His Ile Glu Ser Val His Gln Leu
 305 310 315 320
 Ser Lys Leu Gln Glu Tyr Ser Val Val Asp Tyr Thr Thr Ala Thr
 325 330 335

<210> 3
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic construct

<400> 3
 gagagagaga cgcgtccag tggctgagac gcac 35

<210> 4
 <211> 34
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic construct

<400> 4
 ctctctctgt cgacgaattc aatcttacgg cctg 34

<210> 5
 <211> 4323
 <212> DNA
 <213> Corynebacterium glutamicum

<400> 5
 tcgagagggc tgacgtcggg cccggtacca cgcgtcatat gactagtctg gacctagggg 60
 tatcgtcgac atcgatgctc ttctgcgtta attaacaatt gggatcctct agacccggga 120
 tttaaatcgc tagcgggctg cttaaaggaag cggaacacgt agaaagccag tccgcagaaa 180
 cgggtgctgac ccgcatgaa tgcagctac tgggctatct ggacaaggga aaacgcaagc 240
 gcaaagagaa agcaggtagc ttgcagtggg ctacatggc gatagctaga ctggcggtt 300
 ttatggacag caagcgaacc ggaattgccg gctggggcgc cctctggtta ggttgggaag 360
 ccctgcaaaag taaactggat ggcttctctg ccgccaagga tctgatggcg caggggatca 420
 agatctgatc aagagacagg atgaggatcg ttctgcata ttgaacaaga tggattgcac 480
 gcaggttctc cggccgcttg ggtggagagg ctattcggct atgactgggc acaacagaca 540
 atcggctgct ctgatgccgc cgtgttccgg ctgtcagcgc aggggcgcc gggtctttt 600
 gtcaagaccg acctgtccgg tgccctgaat gaactgcagg acgaggcagc gcggctatcg 660
 tggctggcca cgacgggctg tccttgcgca gctgtgctcg acgttgtcac tgaagcggga 720
 agggactggc tgctattggg cgaagtgcg gggcaggatc tcctgtcatc tcaccttgc 780
 cctgccgaga aagtatccat catggctgat gcaatgcggc ggctgcatac gcttgatccg 840
 gctacctgcc cattcgacca ccaagcgaat catcgcatcg agcagacagc tactcgatg 900
 gaagccggct ttgtcgatca ggatgatctg gacgaagagc atcaggggct cgcgccagcc 960
 gaactgttcg ccaggctcaa ggcgcgcatg cccgacggcg aggatctctg cgtgacccat 1020
 ggcgatgcct gcttgccgaa tatcatggtg gaaaatggcc gcttttctg attcatcgac 1080
 tgtggccggc tgggtgtggc ggaccgctat caggacatag cgttggctac ccgtgatatt 1140
 gctgaagagc ttggcgggca atgggctgac cgcttctctg tgctttacg tatcgccgt 1200
 cccgattcgc agcgcacgc cttctatcgc cttcttgacg agttcttctg agcgggactc 1260
 tggggttcga aatgaccgac caagcgacgc ccaacctgcc atcacgagat ttcgattcca 1320
 ccgcccctt ctatgaaagg ttgggcttcg gaatcgttt ccgggacgcc ggtggatga 1380
 tcctccagcg cggggatctc atgctggagt tcttcgccc cgctagcggc gcgccggcg 1440
 gcccggtgtg aaataccgca cagatgcgta aggagaaaat accgcatcag gcgctcttc 1500
 gcttctctgc tcaactgact gctgcgctcg gtcgttcggc tgcggcgagc ggtatcagct 1560
 cactcaaagg cggtaatagc gttatccaca gaatcagggg ataacgcagg aaagaacatg 1620
 tgagcaaaag gccagcaaaa ggccaggaac cgtaaaaagg ccgcttgcct ggcgttttc 1680
 cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 1740
 aaccgacag gactataaag ataccaggcg ttccccctg gaagctccct cgtgcgtct 1800
 cctgttcgga cctgcccgt taccggatac ctgtccgctc ttctcccttc gggaagcgtg 1860
 gcgctttctc atagctcacg ctgtaggtat ctcagttcgg ttaggtcgt tcgctccaag 1920
 ctgggctgtg tgcacgaacc cccggttcag cccgacccgt gcgcttatc cgttaactat 1980
 cgtcttgagt ccaacccggt aagacacgac ttatcgccac tggcagcagc cactggtaac 2040
 aggattagca gagcgaggta tgtaggcggt gctacagagt tcttgaagtg gtggcctaac 2100

	tacggctaca	ctagaaggac	agtatttggg	atctgcgctc	tgctgaagcc	agttaccttc	2160
	ggaaaaagag	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	cggtggtttt	2220
	tttgtttgca	agcagcagat	tacgcgcaga	aaaaaaggat	ctcaagaaga	tcctttgatc	2280
5	ttttctacgg	ggcttgacgc	tcagtggaac	gaaaaactcac	gttaagggat	tttggctcatg	2340
	agattatcaa	aaaggatcct	cacctagatc	cttttaaagg	ccggcccgcg	ccgccatcgg	2400
	cattttcttt	tgcgttttta	tttgttaact	gttaattgtc	cttgttcaag	gatgctgtct	2460
	ttgacaacag	atgttttctt	gcctttgatg	ttcagcagga	agctcggcgc	aaacgttgat	2520
	tgtttgtctg	cgtagaatcc	tctgtttgtc	atatagcttg	taatcacgac	attgtttcct	2580
	ttcgcttgag	gtacagcgaa	gtgtgagtaa	gtaaaggtta	catcgttagg	atcaagatcc	2640
	atttttaaca	caaggccagt	tttgttcagc	ggcttgtatg	ggccagttaa	agaattagaa	2700
10	acataaccaa	gcattgtaaat	atcgttagac	gtaatgccgt	caatcgctcat	ttttgatccg	2760
	cgggagtcag	tgaacaggtta	ccatttgccg	ttcatttttaa	agacgttcgc	gcgttcaatt	2820
	tcattctgtta	ctgtgtttaga	tgcaatcagc	ggtttcatca	cttttttcag	tgtgtaatca	2880
	tcgttttagct	caatcatacc	gagagcgccg	tttgctaact	cagccgtgcg	ttttttatcg	2940
	ctttgcagaa	gtttttgact	ttcttgacgg	aagaatgatg	tgcttttgcc	atagtatgct	3000
	ttgttaataa	aagattcttc	gccttggtag	ccatcttcag	ttccagtgtt	tgcttcaaat	3060
	actaagtatt	tgtggccttt	atcttctacg	tagtgaggat	ctctcagcgt	atgggttgcg	3120
	cctgagctgt	agtgtccctc	atcgatgaac	tgctgtacat	tttgatacgt	ttttccgtca	3180
15	ccgtcaaaga	ttgatttata	atctcttaca	ccgtttagatg	tcaaagagct	gtctgatgct	3240
	gatacgttaa	cttgtgcagt	tgctcagtgtt	tgtttgccgt	aatgtttacc	ggagaaatca	3300
	gtgtagaata	aacggatttt	tccgtcagat	gtaaatgtgg	ctgaacctga	ccattcttgt	3360
	gtttggctct	ttaggataga	atcatttgca	tcgaatttgt	cgctgtcttt	aaagacgcgg	3420
	ccagcgtttt	tcagctgtgc	aatagaagtt	tcgccgactt	tttgatagaa	catgttaaat	3480
	gatgtgtcat	ccgcattttt	aggatctccg	gctaattgcaa	agacgatgtg	gtagccgtga	3540
	tagtttgcga	cagtgccgtc	agcgttttgt	aatggccagc	tgctccaaac	gtccaggcct	3600
20	tttgcagaag	agatatattt	aatgttggag	gaatcaaatt	cagaaacttg	atatatttca	3660
	tttttttgct	gttcagggat	ttgcagcata	tcattggcgtg	taatatggga	aatgccgtat	3720
	gtttctctat	atggctttttg	gttcgttttct	ttcgcaaaccg	cttgagttgc	gcctcttgcc	3780
	agcagtcggt	tagtaaaagt	taatactgtt	gcttgttttg	caaacttttt	gatgttcac	3840
	gttctatgct	ccittttttat	gtactgtgtt	agcggctctg	ttcttccagc	cctctgtttt	3900
	gaagatggca	agtttagttac	gcacaataaa	aaaagaccta	aaatatgtaa	ggggtgacgc	3960
	caaagtatac	actttgacct	ttacacattt	taggtcttgc	ctgctttatc	agtaacaaac	4020
25	ccgcgcgatt	tactttttcga	ctcatttcta	ttagactctc	gtttggattg	caactggctc	4080
	attttctctc	tttgtttgat	agaaaatcat	aaaaggattt	gcagactacg	ggcctaaga	4140
	actaaaaaat	ctatctgttt	cttttcattc	tctgtatttt	ttatagtttc	tgttgcatgg	4200
	gcataaagtt	gccttttttaa	tcacaattca	gaaaatatca	taatatctca	tttcactaaa	4260
	taatagttaa	cggcaggtat	atgtgatggg	ttaaaaagga	tcggcggccg	ctcgatttaa	4320
	atc						4323
30	<210> 6						
	<211> 5860						
	<212> DNA						
	<213> Corynebacterium glutamicum						
	<400> 6						
35	cccggtagca	cgcgtccag	tggtgagac	gcattccgcta	aagccccagg	aaccctgtgc	60
	agaaagaaaa	cactcctctg	gctaggtaga	cacagtttat	aaaggtagag	ttgagcgggt	120
	aactgtcagc	acgtagatcg	aaaggtgcac	aaaggtggcc	ctggctgtac	agaaatatgg	180
	cggttctctg	cttgagagtg	cggaacgcac	tagaaaacgtc	gctgaacgga	tcggttgccac	240
	caagaaggct	ggaaatgatg	tcgtggttgt	ctgctccgca	atgggagaca	ccacggatga	300
	acttctagaa	cttgacagcg	cagtgaaatcc	cggtccgcga	gctcgtgaaa	tggtatgtct	360
	cctgactgct	gtgagcgta	tttctaaccg	tctcgtcgcc	atggctattg	agtcctttgg	420
	cgcagaagcc	caatctttca	cgggctctca	ggctggtgtg	ctcaccaccg	agcggccagg	480
40	aaacgcacgc	attgtttgatg	tcactccagg	tcgtgtgcgt	gaagcactcg	atgagggcaa	540
	gatctgcatt	gttgctgggt	tccagggtgt	taataaaagaa	accgcgcatg	tcaccacggt	600
	gggtcgtggt	ggttctgaca	ccactgcagt	tgctgttgca	gctgctttga	acgctgatgt	660
	gtgtgagatt	tactcggacg	ttgacgggtg	gtataaccgt	gaccgcgcga	tcgttcttaa	720
	tgacagaag	ctggaaaagc	tcagcttcga	agaaatgctg	gaacttgctg	ctgttggctc	780
	caagattttg	gtgctgcgca	gtgttgaaata	cgctcgtgca	ttcaatgtgc	cacttcgctg	840
	acgctcgtct	tatagtaatg	atcccggcac	tttgattgcc	ggctctatgg	aggatattcc	900
45	tgtggaagaa	gcagtcctta	cgggtgtcgc	aaccgacaag	tccgaagcca	aagtaaccgt	960
	tctgggtatt	tccgataaag	caggcgaggc	tgcaaggtt	ttccgtgcgt	tggtgtgatg	1020
	agaaatcaac	attgacatgg	ttctgcagaa	cgctcttctt	gtagaagacg	gcaccaccga	1080
	catcaccttc	acctgccctc	gttccgacgg	ccgcgcgcgc	atggagatct	tgaagaagct	1140
	tcagggttcag	ggcaactgga	ccaatgtgct	ttacgacgac	caggctcgga	aagctctcct	1200
	cgtgggtgct	ggcatgaagt	ctcaccagg	tgttaccgca	gagttcatgg	aagctctgct	1260
	cgatgtcaac	gtgaacatcg	aattgatttc	cacctctgag	attcgtattt	ccgtgctgat	1320
50	ccgtgaagat	gatctggatg	ctgctgcacg	tgcatgtcat	gagcagttcc	agctgggctg	1380
	cgaagacgaa	gccgtcgttt	atgcaggcac	cggacgctaa	agttttaaag	gagtagtttt	1440
	acaatgacca	ccatcgcagt	tggttggtgca	accggccagg	tcggccagggt	tatgcgcacc	1500
	ctttttggaag	agcgcaattt	cccagctgac	actgttctgt	tccttgcctc	cccacgttcc	1560
	gcaggccgta	agattgaatt	cgctgcacatc	gatgctcttc	tgctgttaatt	aacaattggg	1620
	atcctctaga	cccggtgatt	aaatcgctag	cgggctgcta	aagggaagcgg	aacacgtaga	1680
	aagccagtc	gcagaaacgg	tgctgacccc	ggatgaatgt	cagctactgg	gctatctgga	1740
55	caagggaaaa	gcgaagcgca	aagagaaagc	aggtagcttg	cagtgggcct	acatggcgat	1800
	agctagactg	ggcgggtttt	tggaacagaa	gcgaaccgga	attgccagct	ggggcgccct	1860
	ctggttaagg	tgggaagccc	tgcaaaagtaa	actggatggc	tttcttgccg	ccaagatctt	1920

gatggcgag gggatcaaga tctgatcaag agacaggatg aggatcggtt cgcattgattg 1980
 aacaagatgg attgcacgca ggttctccgg ccgcttgggt ggagaggcta ttcggctatg 2040
 actgggcaca acagacaatc ggctgctctg atgcccgggt gttccggctg tcagcgagcg 2100
 ggcgcccggc tctttttgtc aagaccgacc tgtccgggtg cctgaatgaa ctgcaggagc 2160
 aggcagcgcg gctatcgtyg ctggccacga cgggcttcc ttgcgcagct gtgctcgagc 2220
 ttgtcactga agcgggaagg gactggctgc tattggcgca agtgccgggg caggatctcc 2280
 tgtcatctca ccttgctcct gccgagaaag tatccatcat ggctgatgca atgcggcggc 2340
 tgcatacgct tgatccggct acctgcccat tcgaccacca agcgaacat cgcattcgagc 2400
 gagcacgtac tcggatggaa gccggtcttg tcgatcagga tgatctggac gaagagcatc 2460
 aggggctcgc gccagccgaa ctgttcgcca ggctcaaggc gcgcatgccc gacggcgagg 2520
 atctctcgtg gacccatggc gatgctctgt tgccgaatat catggtggaa aatggccgct 2580
 tttctggatt catcgactgt ggccggctgg gtgtggcgga ccgctatcag gacatagcgt 2640
 tggctaccgc tgatattgct gaagagcttg gcggcgaaatg ggctgaccgc ttcctcgtgc 2700
 tttacgggat cgccgctccc gattcgagc gcattgcctt ctatcgctt cttgacgagt 2760
 tcttctgagc gggactctgg ggttcgaaat gaccgaccaa gcgacgccc acctgccatc 2820
 acgagatttc gattccaccg ccgcttctta tgaaagggtt ggcttcggaa tcgttttccg 2880
 ggagcggcgc tggatgatcc tccagcgagg ggaatctcat ctggagttct tcgcccacgc 2940
 tagcggcgcg ccggccggcc cgggtgtaaa taccgcacag atcgtaagg agaaaaatcc 3000
 gcatcaggcg ctcttccgct tcttcgctca ctgactcgct gcgctcggtc gttcggtgc 3060
 ggcgagcggg atcagctcac tcaaaaggcg taatacggtt atccacagaa tcaggggata 3120
 acgcaggaaa gaacatgtga gcaaaaggcc agcaaaaggc caggaaaccgt aaaaaggccg 3180
 cggtgtgagg gtttttccat aggtcccgcc cccctgacga gcatcacaaa aatcgacgtc 3240
 caagtccagag gtggcgaaac ccgacaggac tataaagata ccaggcggtt cccctggaa 3300
 gctccctcgt gcgctctcct gttccgaccc tgccgcttac cggataacctg tccgccttcc 3360
 tcccttcggg aagcgtggcg ctttctcata gctcacgctg taggtatctc agttcgggtg 3420
 aggtcgttcg tcctcaagctg ggctgtgtgc acgaaccccc cgctcagccc gacctcgctg 3480
 ccttatccgg taactatcgt cttgagtcca acccggttaag acacgactta tcgccactgg 3540
 cagcagccac tggtaacagg attagcagag cgaggatgt aggcgggtgct acagagttct 3600
 tgaagtgggt gcctaactac ggctacacta gaaggacagt atttggtatc tgcgtctcgc 3660
 tgaagccagt taccttcgga aaaagagttg gtagctcttg atccggcaaa caaaccacgc 3720
 ctggtagcgg tggttttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc 3780
 aagaagatcc tttgatcttt tctacggggt ctgacgtca gtggaacgaa aactcagctt 3840
 aagggaattt gttcatgaga ttatcaaaaa ggaatctcac ctatgacctt ttaaggccg 3900
 gccgcgcccg ccacggcgat tttcttttgc gtttttatit gtttaactgtt aattgtcctt 3960
 gttcaaggat gctgtctttg acaacagatg ttttcttgcc tttgatgttc agcaggaagc 4020
 tcggcgcaaa cgttgattgt ttgtctcgct agaactcctc tttgtcata tagctgttaa 4080
 tcacgacatt gtttctttc gcttgaggta cagcgaagtg tgagtaagta aaggttacat 4140
 cgttagatcc aagatccatt ttaacacaa ggccagtttt gttcagcggc ttgtatgggc 4200
 cagtttaaga attagaaaca taaccaagca tgtaaatatc gttagacgta atgccgtcaa 4260
 tcgtcatttt tgatccgagg gagtcagtga acaggtacca tttgccgttc attttaaga 4320
 cgttcgcggc ttcaatttca tctgttactg tgtagatgc aatcagcggt ttcatactt 4380
 ttttcagttg ttaactatcg tttagctcaa tcataccgag agcggcggtt gctaactcac 4440
 ccgtgcgttt tttatcgctt tgcagaagtt tttgactttc ttgacggaag aatgatgtgc 4500
 ttttgccata gtatgctttg ttaataaagg attcttcgcc ttggtagcca tcttcagttc 4560
 cagtgtttgc ttcaataact aagtatttgt ggcctttatc ttctacgtag tgaggatctc 4620
 tcagcgtatg gttgtcgctt gagctgtagt tgccttcacg gatgaactgc tgtacatttt 4680
 gatacgtttt tccgtcaccc tcaaagattg atttataatc ctctacaccg ttgatgttca 4740
 aagagctgtc tgatgctgat acgttaactt gtgcagttgt cagtgtttgt ttgccgtaat 4800
 gtttacccgga gaaatcagtg tagaataaac ggatttttcc gtcagatgta aatgtggctg 4860
 aacctgacca ttctgtgtt tggcttttta ggatagaatc atttgcatcg aatttgtcgc 4920
 tgtctttaa gacggcgcca gcgtttttcc agctgtcaat agaagtttcg ccgacttttt 4980
 gatagaacat gtaaatcgat gtgtcatccg catttttagg atctccggct aatgcaaaaga 5040
 cgatgtggta gccgtgatag tttgcgacag tgccgtcagc gttttgtaat ggccagctgt 5100
 cccaaacgtc caggccctttt gcagaagaga tatttttaat tgtggacgaa tcaaatcag 5160
 aaacttgata tttttcattt ttttgcgtgt cagggaattt cagcatatca tggcgtgtaa 5220
 tatgggaaat gccgtatggt tccttatatg gcttttggtt cgtttctttc gcaaaccgctt 5280
 gagttgcgcc tcttgccagc agtgcggtag taaaggttaa tactgttgct tgttttgcaa 5340
 actttttgat gttcatcggt catgtctcct tttttatgta ctgtgttagc ggtctgcttc 5400
 ttccagccct cctgtttgaa gatggcaagt tagttacgca caataaaaaa agacctaaaa 5460
 tatgtaaggg gtgacgcaa agtatacact ttgcccttta cacattttag gtcttgctg 5520
 ctttatcagt acaaaacccg cgcgatttac ttttcgacct cattctatta gactctcggt 5580
 tggattgcaa ctgggtctatt ttctcttttt gtttgataga aaatcataaa aggatttgca 5640
 gactacgggc cttaaagaact aaaaaatcta tctgtttctt ttcattctct gtatttttta 5700
 tagtttctgt tgcattggga taaagttgac tttttaatca caattcagaa aatatcataa 5760
 tatctcattt cactaaataa tagtgaacgg caggtatatg tgatgggtta aaaaggatcg 5820
 gcggccgctc gatttaaatc tcgagaggcc tgacgtcggg 5860

<210> 7
 <211> 38
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic construct

	<400> 7	cggcaccacc gacatcatct tcacctgccc tcgttccg	38
5	<210> 8		
	<211> 38		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Synthetic construct		
10	<400> 8	cggaacgagg gcaggtgaag atgatgtcgg tgggtgccg	38
	<210> 9		
	<211> 1263		
	<212> DNA		
15	<213> Corynebacterium glutamicum		
	<400> 9	gtggccctgg tcgtacagaa atatggcggg tcctcgcttg agagtgcgga acgcattaga 60	
		aacgtcgctg aacggatcgt tgccaccaag aaggctggaa atgatgtcgt gggtgtctgc 120	
		tccgcaatgg gagacaccac ggatgaactt ctagaacttg cagcggcagt gaatcccgtt 180	
		ccgccagctc gtgaaatgga tatgtcctcg actgctggtg agcgtatttc taacgctctc 240	
20		gtcgccatgg ctattgagtc ccttggcgca gaagcccaat ctttcacggg ctctcaggct 300	
		gggtgtgctca ccaccgagcg ccacggaaac gcacgcattg ttgatgtcac tccaggtcgt 360	
		gtgctggaag cactcgatga gggcaagatc tgcattgttg ctggtttcca ggggtgtaat 420	
		aaagaaaccc gcgatgtcac caggttgggt cgtgggtggt ctgacaccac tgcagttgct 480	
		ttggcagctg ctttgaacgc tgatgtgtgt gagatttact cggacgttga cgggtgtgat 540	
		accgtgacc cgcgcacgtt tcctaatgca cagaagctgg aaaagctcag cttcgaagaa 600	
		atgctggaac ttgctgctgt tggctccaag attttgggtc tgcgcagtggt tgaatacgtc 660	
		cgtgcattca atgtgccact tcgcgtacgc tcgtcttata gtaatgatcc cggcactttg 720	
25		attgccggct ctatggagga tattcctgtg gaagaagcag tccttaccgg tgtcgcaacc 780	
		gacaagtccg aagccaaagt aaccgttctg ggtatttccg ataagccagg cgagggtgct 840	
		aagggttttc gtgcgttggc tgatgcagaa atcaacattg acatggttct gcagaacgtc 900	
		tcttctgtag aagacggcac caccgacatc accttcacct gccctcggtc cgacggccgc 960	
		cgcgcgatgg agatcttgaa gaagcttcag gttcagggca actggacca tgtgctttac 1020	
		gacgaccagg tcggcaaaagt ctccctcgtg ggtgctggca tgaagtctca cccaggtggt 1080	
		accgcagagt tcattggaagc tctgcgcgat gtcaacgtga acatcgaatt gatttccacc 1140	
30		tctgagattc gtatttccgt gctgatccgt gaagatgatc tggatgctgc tgcacgtgca 1200	
		ttgcatgagc agttccagct gggcggcgaa gacgaagcgg tcgtttatgc aggcaccgga 1260	
		cgc	1263
	<210> 10		
	<211> 5860		
	<212> DNA		
35	<213> Corynebacterium glutamicum		
	<400> 10	cccggtagca cgcgtccag tggctgagac gcatccgcta aagccccagg aaccctgtgc 60	
		agaaagaaaa cactcctctg gctaggtaga cacagtttat aaaggtagag ttgagcgggt 120	
		aactgtcagc acgtagatcg aaagggtgcac aaagggtggc ctggtcgtac agaaatatgg 180	
		cggttcctcg cttgagagtg cggaaacgat tagaaacgtc gctgaacgga tcgttgccac 240	
40		caagaagggt ggaatatgat tcgtggttgc ctgctccgca atgggagaca ccacggatga 300	
		actttctagaa cttgcagcgg cagtgaatcc cgttccgcca gctcgtgaaa tggatatgct 360	
		cctgactgct ggtgagcgtt tttctaacgc tctcgtcgcc atggctattg agtcccttgg 420	
		cgcagaagcc caatctttca cgggctctca ggctggtgtg ctaccaccg agcggccagg 480	
		aaacgcacgc attgttgatg tcaactccagg tcgtgtgctg gaagcactcg atgagggcaa 540	
		gatctgcatt gttgctggtt tccagggtgt taataaaagaa acccgcgatg tcaccacgtt 600	
		gggtcgtggt ggttctgaca ccaactgcagt tgcgttggca gctgctttga acgctgatgt 660	
45		gtgtgagatt tactcggacg ttgacgggtgt gtataccgct gacccgcgca tcgttcctaa 720	
		tgcacagaag ctggaaaagc tcagcttcga agaaatgctg gaacttgctg ctgttggtct 780	
		caagattttg gtgctgcgca gtgttgaata cgctcgtgca ttcaatgtgc cacttcgctg 840	
		acgctcgtct tatagtaatg atcccggcac ttgtattgcc ggtcttatgg aggatatttc 900	
		tgtggaagaa gcagtcctta ccggtgtcgc aaccgacaag tccgaagcca aagtaaccgt 960	
		tctgggtatt tccgataagc caggcgaggc tgccaaggtt ttccgtgctg tggctgatgc 1020	
		agaaatcaac attgacatgg ttctgcagaa cgtctcttct gtagaagacg gcaccaccga 1080	
50		catcatcttc acctgccctc gttccgacgg ccgccgcgag atggagatct tgaagaagct 1140	
		tcaggttcag ggcaactgga ccaatgtgct ttacgacgac caggctcgga aagtctccct 1200	
		cgtgggtgct ggcatgaagt ctacccagg tgttaccgca gagttcatgg aagctctgct 1260	
		cgatgtcaac gtgaacatcg aattgatttc cactctgag attcgtattt cgtggtgat 1320	
		ccgtgaagat gatctggatg ctgctgcacg tgcattgcat gacgagttcc agctggggcg 1380	
		cgaagacgaa gccgtcggtt atgcaggcac cggacgctaa agttttaaag gagtagtttt 1440	
		acaattgacca ccatcgcagt tgttgggtga accggccagg tcggccagggt tatgcgcacc 1500	
55		cttttggaa gacgcaattt cccagctgac actgttcgtt tctttgcttc cccacgttcc 1560	
		gcaggccgta agattgaatt cgtcgcacatc gatgctcttc tgcgtiaatt aacaattggg 1620	
		atcctctaga cccgggattt aaatcgctag cgggctgcta aaggaagcgg aacacgtaga 1680	

	aagccagttcc	gcagaaacgg	tgctgacccc	ggatgaatgt	cagctactgg	gctatctgga	1740
	caagggaaaa	cgcaagcgca	aagagaaaagc	aggtagcttg	cagtgggctt	acatggcgat	1800
	agctagactg	ggcgggtttta	tggaacagcaa	cggaaccgga	attgccagct	ggggcgccct	1860
5	ctggtaaggt	tggaagcccc	tgcaaaagtaa	actggatggc	tttcttgccg	ccaaggatct	1920
	gatggcgag	gggatcaaga	tctgatcaag	agacaggatg	aggatcggtt	cgcatgattg	1980
	aacaagatgg	attgcacgca	ggttctccgg	ccgcttgggt	ggagaggcta	ttcggctatg	2040
	actggggcaca	acagacaatc	ggctgctctg	atgccgccgt	gttccggctg	tcagcgagg	2100
	ggcgcccggt	ctttttgtc	aagaccgacc	tgiccgggtg	cctgaatgaa	ctgcaggacg	2160
	aggcagcgcg	gctatcggtg	ctggccacga	cgggcggtcc	ttgcgcagct	gtgctcgagc	2220
	ttgtcactga	agcgggaagg	gactggctgc	tattgggcga	agtcccgggg	caggatctcc	2280
10	tgctatctca	ccttgctcct	gccgagaaag	tatccatcat	ggctgatgca	atgcggcggc	2340
	tgcatacgct	tgatccggct	acctgcccc	tcgaccacca	agcgaaacat	cgcatcgagc	2400
	gagcacgtac	tcggatggaa	gccgggtctt	tcgatcagga	tgatctggac	gaagagcatc	2460
	aggggctcgc	gccagccgaa	ctgttcgcca	ggctcaaggc	gcgcagtccc	gacggcgagg	2520
	atctcgtcgt	gacccatggc	gatgcctgct	tgccgaatat	catggtggaa	aatggccgct	2580
	tttctggatt	catcgactgt	ggccggctgg	gtgtggcgga	ccgctatcag	gacatagcgt	2640
	tggtctaccg	tgatattgct	gaagagcttg	gcggcggaatg	ggctgaccgc	ttcctcgctg	2700
	tttacggtat	cgccgctccc	gattcgagc	gcctcgctt	ctatcgctt	cttgacgagt	2760
15	tcttctgagc	gggactctgg	ggttcgaaat	gaccgaccaa	gcgacgcccc	acctgccatc	2820
	acgagatttc	gattccaccg	ccgcttctta	tgaaagggtg	ggcttcggaa	tcgttttccg	2880
	ggacgcccgc	tgatgatcc	tccagcgcg	ggatctcatg	ctggagttct	tcgcccacgc	2940
	tagcgccgcg	ccggccggcc	cggtgtgaaa	taccgcacag	atgcgtaagg	agaaaaatcc	3000
	gcatacggcg	ctcttcgcgt	tcctcgctca	ctgactcgct	gcgctcggtc	gttcgctcgc	3060
	ggcgagcggt	atcagctcac	tcaaaggcgg	taatacgggt	atccacagaa	tcagggggata	3120
	acgcagggaaa	gaacatgtga	gcaaaaggcc	agcaaaaggc	caggaaccgt	aaaaaggccg	3180
20	cgttgtcggc	gtttttccat	aggctccgcc	cccctgacga	gcatacacia	aatcgacgct	3240
	caagtacagag	gtggcgaaac	ccgacaggac	tataaagata	ccaggcggtt	ccccctggaa	3300
	gctccctcgt	gcgctctcct	gttccgaccc	tgccgcttac	cggatacctg	tccgccttcc	3360
	tcccttcggg	aagcgtggcg	ctttctcata	gctcacgctg	taggtatctc	agttcgggtg	3420
	aggctcgttcg	ctccaagctg	ggctgtgtgc	acgaaccccc	cgttcagccc	gaccgctcgc	3480
	ccttatccgg	taactatcgt	cttgagtcca	acccggtaag	acacgactta	tcgccactgg	3540
	cagcagccac	tggtaacagg	attagcagag	cgagggtatg	aggcgggtgct	acagagttct	3600
25	tgaagtggtg	gcctaactac	ggctacacta	gaaggacagt	atttggtatc	tgcgctctgc	3660
	tgaagccagt	taccttcgga	aaaagagttg	gtagctcttg	atccggcaaa	caaaccaccg	3720
	ctggtagcgg	tggttttttt	gtttgcaagc	agcagattac	gcgcagaaaa	aaaggatctc	3780
	aagaagatcc	tttgatcttt	ctacgggggt	ctgacgctca	gtggaaacgaa	aactcacgtt	3840
	aagggatttt	ggtcatgaga	ttatcaaaaa	ggatcttcac	ctagatcctt	ttaaaggccg	3900
	gccgcggccg	ccatcggcgt	tttcttttgc	gtttttattt	gttaactggt	aatgttcctt	3960
30	gttcaaggat	gctgtctttg	acaacagatg	ttttcttgcc	tttgatgttc	agcaggaagc	4020
	tcggcgcaaa	cggtgattgt	ttgtctcggt	agaatcctct	gtttgtcata	tagcttgtaa	4080
	tcacgacatt	gtttcctttc	gcttgaggta	cagcgaagtg	tgagtaagta	aaggttacat	4140
	cgttaggatc	atgatccatt	tttaacacaa	ggccagtttt	gttcagcggc	ttgtatgggc	4200
	cagttaaaga	attagaaaca	taaccaagca	tgtaaataatc	gttagacgta	atgccgtcaa	4260
	tcgtcatctt	tgatccgcgg	gagtcagtga	acaggtacca	tttgccgttc	attttaagaa	4320
	cgtttcgcgg	ttcaatttca	tctgttactg	tgtagatgc	aatcagcggt	ttcatcactt	4380
35	ttttcagtgt	gttaatcatc	tttagctcaa	tcataccgag	agcgccggtt	gctaacctcag	4440
	ccgtgcgttt	tttatcgctt	tgcaagaagt	tttgactttc	ttgacggaag	aatgatgtgc	4500
	ttttgccata	gtagtctttg	ttaaataaag	attcttccgc	ttggtagcca	tcttcagttc	4560
	cagtgtttgc	ttcaataact	aagtatttgt	ggcctttatc	ttctacgtag	tgaggatctc	4620
	tcagcgtatg	gttgcgcct	gagctgtagt	tgcccttcac	gatgaactgc	tgtacatttt	4680
	gatacgtttt	tccgtcaccc	tcaaagattg	atttataatc	ctctacaccg	ttgatgttca	4740
	aagagctgtc	tgatgctgat	acgttaactt	gtgcagttgt	cagtgtttgt	ttgccgtaat	4800
40	gtttaccgga	gaaatcagtg	tagaataaac	ggatttttcc	gtcagatgta	aatgtggctg	4860
	aacctgacca	ttcttgttgt	tggtctttta	ggatagaatc	atttgcatcg	aatttgtcgc	4920
	tgcttttaaa	gacgcggcca	gcgtttttcc	agctgtcaat	agaagtttcg	ccgacttttt	4980
	gatagaacat	gtaaatcgat	gtgtcatccg	catttttagg	atctccggct	aatgcaaaaga	5040
	cgatgtggta	gccgtgatag	tttgcgacag	tgccgtcagc	gttttgtaat	ggccagctgt	5100
	cccaaaccgc	caggcctttt	gcagaagaga	tatttttaat	tgtaggacgaa	tcaaatcag	5160
	aaacttgata	tttttcattt	ttttgctggt	cagggaattg	cagcatatca	tggcgtgtaa	5220
	tatgggaaat	gccgtatggt	tccttatatg	gcttttggtt	cgtttctttc	gcaaaccgctt	5280
45	gagttgcgcc	tcttgccagc	agtgcggtag	taaaggttaa	tactgttgct	tgttttgcaa	5340
	actttttgat	gttcacgtgt	catgtctcct	tttttatgta	ctgtgttagc	ggctcgtctc	5400
	ttccagccct	cctgtttgaa	gatggcaagt	tagttacgca	caataaaaaa	agacctaaaa	5460
	tatgtaaggg	gtgacgccaa	agtatacact	ttgcccttta	cacatttttag	gtcttgccctg	5520
	ctttatcagt	aaacaaaccg	cgcatatttac	ttttcgacct	cattctatta	gactctcggt	5580
	tggattgcaa	ctggtctatt	ttctcttttt	gtttgataga	aaatcataaa	aggatttgca	5640
	gactacgggc	ctaaagaact	aaaaaatcta	tctgtttctt	ttcattctct	gtatttttta	5700
50	tagtttctgt	tgcatgggca	taaagttgcc	tttttaatca	caattcagaa	aatatcataa	5760
	tatctcatct	cactaaataa	tagtgaacgg	caggatatatg	tgatgggtta	aaaaggatcg	5820
	gcggccgctc	gatttaaatc	tcgagaggcc	tgacgtcggg			5860

<210> 11

<211> 20

<212> DNA

<213> Artificial Sequence

<220>
 <223> Synthetic construct

5 <400> 11
 tggccgttac cctgcgaatg 20

<210> 12
 <211> 20
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> Synthetic construct

<400> 12
 tgtatgtcct cctggacttc 20

15 <210> 13
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic construct

20 <400> 13
 gaagtccagg aggacataca atgaacctaa agaaccccga 40

<210> 14
 <211> 31
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> Synthetic construct

<400> 14
 atctacgtcg acccaggatg ccctggattt c 31

30 <210> 15
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Synthetic construct

35 <400> 15
 tatcaacgcg ttcttcatcg gtagcagcac c 31

<210> 16
 <211> 37
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> Synthetic construct

<400> 16
 cattcgcagg gtaacggcca ctgaagggcc tcctggg 37

45 <210> 17
 <211> 5928
 <212> DNA
 <213> Corynebacterium glutamicum

50 <400> 17
 tcgagaggcc tgacgtcggg cccggtacca cgcgttcttc atcggtagca gcacccgaga 60
 ccatgacgcg ggcacgccc agatccatca cacgcagatc acgcacatca gattcctgtg 120
 aggtgtaaat tcccacgtcg tggccatcaa gatcataaga ctgagaaaga tcacgccagc 180
 gagtatcata accagccaca gcatcctcaa cggtttcacc agtttgagtg agctgaatat 240
 agccctcatc tgcggtgaca tatccaacta cagatgcccg ggtgtcatcc accatggtgc 300
 gtcgagctga atttgtggtc cagccttcag gagtttccgg caacctagtt gcatgatcag 360
 tcattgcgcg cgcttcattt gacataaaag tggaagcatc aacttcagg acctgcccac 420
 55 ttccaggggg tcctgtattg aaagaacaca ttcccgtgaa tcccaccgct accaacaatga 480
 tgatcgcgga gactaccaac gagataatca tgtctcgact gccatcaaaa attttcggtc 540

	gtttctcagc	cacccgccta	gtatgtcacg	agtttggtag	gaaacccccct	tttgggtgtc	600
	cagaatccaa	aattccgggc	acaaaagtgc	aacaatagat	gacgtgcggg	ttgatacagc	660
	ccaagcgccg	atacatattat	aatgcgccta	gatacgtgca	acccacgtaa	ccagggtcaga	720
5	tcaagtgtccc	caggaggccc	ttcagtgggc	gttaccctgc	gaatgtccac	agggtagctg	780
	gtagtgtgaa	aatcaacgcc	gttggccctta	ggattcagta	actggcacat	tttgtaatcg	840
	gctagatctg	tgtgtctcagt	cttccaggct	gcttatcaca	gtgaaagcaa	aaccaattcg	900
	tggctgcgaa	agtcgtagcc	accacgaagt	ccaggaggac	atacaatgaa	cctaaagaac	960
	cccgaaacgc	cagaccgtaa	ccttgctatg	gagctggtgc	gagttacgga	agcagctgca	1020
	ctggcttctg	gacgttgggt	tggacgtggc	atgaagaatg	aaggcgacgg	tgccgctggt	1080
	gacgccatgc	gccagctcat	caactcagtg	accatgaagg	gcgtcgttgt	tatcggcgag	1140
	ggcgaaaaag	acgaagctcc	aatgctgtac	aacggcgaag	aggctcgaac	cggctttgga	1200
10	cctgagggtt	atatcgcagt	tgacccagtt	gacggcacca	ccctgatggc	tgagggtcgc	1260
	cccaacgcaa	tttccattct	cgcagctgca	gagcgtggca	ccatgtacga	tccatcctcc	1320
	gtcttctaca	tgaagaagat	cgccgtggga	cctgaggccg	caggcaagat	cgacatcgaa	1380
	gctcagttg	cccacaacat	caacgcggtg	gcaaatgcca	agggaatcaa	cccttcgcac	1440
	gtcaccgttg	tcgtgcttga	ccgtctctgc	cacatcgaac	tgatcgcaga	cattcgtcgt	1500
	gcaggcgcaa	agggttcgtct	catctccgac	ggcgacgttg	caggtgcagt	tgacagcagt	1560
	caggattcca	actccgtgga	catcatgatg	ggcaccggcg	gaaccccgaa	aggcatcatc	1620
15	actgcgtgcg	ccatgaagtg	catgggtggc	gaaatccagg	gcacccctgg	tcgacatcga	1680
	tgctcttctg	cgtaaatata	caattgggat	cctctagacc	cgggatttaa	atcgctagcg	1740
	ggctgctaaa	ggaagcggaa	cacgtagaaa	gccagtcgag	agaaacgggt	ctgaccccg	1800
	atgaatgtca	gctactgggc	tatctggaca	agggaaaaacg	caagcgcaaa	gagaaagcag	1860
	gtagcttgca	gtgggcttac	atggcgatag	ctagactggg	cggttttatg	gacagcaagc	1920
	gaaccggaat	tgccagctgg	ggcgccctct	ggtaaggttg	ggaagccctg	caaagtaaac	1980
	tggtatggctt	tcttgcgccc	aaggatctga	tggcgagggg	gatcaagatc	tgatcaagac	2040
20	acaggatgag	gatcgtttcg	catgattgaa	caagatggat	tgacgcagcg	ttctccggcc	2100
	gcttgggtgg	agaggctatt	cggtctatgac	tgggcacaa	agacaatcgg	ctgctctgat	2160
	gccgcggtgt	tccggctgtc	agcgcagggg	cgcccggttc	ttttgtcaa	gaccgacctg	2220
	tccggtgccc	tgaatgaact	gcaggacgag	gcagcgcggc	tatcgtgggt	ggccagcagc	2280
	ggcgttcctt	gcgcagctgt	gctcgacgtt	gtcactgaag	cgggaaggga	ctggctgcta	2340
	ttgggcgaag	tgccggggca	ggatctcctg	tcattctcacc	ttgctcctgc	cgagaaagta	2400
25	tctgcatggt	ctgatgcaat	cgcgcggtcg	catacgcttg	atccggctac	ctgcccattc	2460
	gaccaccaag	cgaaacatcg	catcgagcga	gcacgtactc	ggatggaagc	cggctctgtc	2520
	gatcaggatg	atctggacga	agagcatcag	gggctcgcgc	cagccgaact	gttcgcccag	2580
	ctcaagcgcg	gcatgccgga	cgcgaggatg	ctcgtctgta	cccatggcga	tgccatcgtg	2640
	ccgaatatca	tggtggaaaa	tggccgcttt	tctggattca	tcgactgtgg	ccggctgggt	2700
	gtggcgggacc	gctatcagga	catagcggtg	gctacccgtg	atatgtgctg	agagcttggc	2760
	ggcgaaatggg	ctgaccgctt	cctcgtgctt	tacgggtatc	ccgctcccca	ttcgcaagcg	2820
30	atcgcttctt	atcgcttctt	tgacgagttc	ttctgagcgg	gactctgggg	ttcgaaatga	2880
	ccgaccaagc	gacgccccac	ctgcctacac	gagatttcga	ttccaccgcc	gccttctatg	2940
	aaagggttggg	cttcggaatc	gttttcgggg	acgcccggctg	gatgatcctc	cagcgcgggg	3000
	atctcatgct	ggagtttctt	gccacgccta	gcggcgccgc	ggccggcccg	gtgtgaaata	3060
	ccgcacagat	gcgtaaggag	aaaataccgc	atcaggcgct	cttcgcgttc	ctcgctcact	3120
	gactcgctgc	gctcggctgt	tcggctgcgg	cgagcggtat	cagctcactc	aaaggcggtg	3180
	atacggttat	ccacagaatc	aggggataac	gcaggaaaga	acatgtgagc	aaaaggccag	3240
35	caaaaggcca	ggaaccgtaa	aaaggccgcg	ttgctggcgt	ttttccatag	gctccgcccc	3300
	cctgacgagc	atcacaaaaa	tcgacgctca	agtcagaggt	ggcgaaaccc	gacaggacta	3360
	taaagatacc	aggcgtttcc	ccctggaagc	tccctcgtgc	gctctcctgt	tccgaccctg	3420
	ccgcttaccg	gatacctgtc	cgcttttctc	ccttcgggaa	gcgtggcgct	ttctcatagc	3480
	tcacgctgta	ggtatctcag	ttcgggtgtag	gtcgttcgct	ccaagctggg	ctgtgtgcac	3540
	gaaccccccg	ttcagcccga	ccgctgcgcc	ttatccggta	actatcgtct	tgagtccaac	3600
	ccggttaagac	acgacttatc	gccactggca	gcagccactg	gtaacaggat	tagcagcagc	3660
40	aggtagtag	gcggtgctac	agagtctctg	aagtgggtgg	ctaactacgg	ctacactaga	3720
	aggacagtat	ttgggtatctg	cgctctgctg	aagccagtta	ccttcggaaa	aagagttggt	3780
	agctcttgat	ccggcaaaaca	aaccaccgct	ggtagcgggt	gtttttttgt	ttgcaagcag	3840
	cagattacgc	gcagaaaaaa	aggatctcaa	gaagatcctt	tgatcttttc	tacggggtct	3900
	gacgctcagt	ggaacgaaaa	ctcaggttaa	gggatttttg	tcatgagatt	atcaaaaagg	3960
	atcttcacct	agatcctttt	aaaggccggc	cgcgcccgcc	atcggcattt	tcttttgcgt	4020
	ttttatttgt	taactgttaa	ttgtccttgt	tcaaggatgc	tgtctttgac	aacagatgtt	4080
45	ttcttgccct	tgatgttcag	caggaagctc	ggcgcaaacg	ttgattgttt	gtctgcgtag	4140
	aatcctctgt	ttgtcatata	gcttgtaatc	acgacattgt	ttcctttcgc	ttgacagata	4200
	gcgaagtgtg	tgtaagtaaa	ggttacatcg	ttaggatcaa	gatccatttt	taacacagg	4260
	ccagttttgt	tcagcggtct	gtatgggcca	gttaaagaat	tagaaacata	accaagcatg	4320
	taaatatcgt	tagacgtaat	gccgtcaatc	gtcatttttg	atccgcgggg	gtcagtgaac	4380
	aggtagcatt	tgccgttcat	tttaaagacg	ttcgcgcggt	caatttcac	tggtactgtg	4440
	ttagatgcaa	tcagcggttt	catcactttt	ttcagtggtg	aatcatcggt	tagctcaatc	4500
	ataccgagag	cgccggtttg	taactcagcc	gtgcgttttt	tatcgctttg	cagaagtttt	4560
50	tgactttctt	gacggaagaa	tgatgtgctt	ttgccatagt	atgctttgtt	aaataaagat	4620
	tcttcgcctt	ggtagccatc	ttcagttcca	gtgtttgctt	caaatactaa	gtattttgtg	4680
	cctttatctt	ctacgtagtg	aggatctctc	agcgtatggt	tgctgcctga	gctgtagtgt	4740
	cctttacatgc	tgaactgctg	tacattttga	tacgtttttc	cgtcacccgtc	aaagattgat	4800
	ttataatcct	ctacaccgtt	gatgttcaaa	gagctgtctg	atgctgatac	gttaacttgt	4860
	gcagttgtca	gtgtttgttt	gccgtaatgt	ttaccggaga	aatcagtgta	gaataaacgg	4920
	atttttccgt	cagatgtaaa	tgtggctgaa	cctgaccatt	cttgtgtttg	gtcttttagg	4980
55	atagaatcat	ttgcatcgaa	tttgcgtgct	tcttttaaaga	cgcgcccgagc	gtttttccag	5040
	ctgtcaatag	aagtttccgc	gactttttga	tagaacatgt	aaatcgatgt	gtcatccgca	5100
	tttttaggat	ctccggctaa	tgcaagagac	atgtggttagc	cgtgatagtt	tgcgacagtg	5160

EP 1 939 296 A2

	ccgtcagcgt	tttghtaatgg	ccagctgtcc	caaacgtcca	ggccttttgc	agaagagata	5220
	tttttaattg	tggaacgaatc	aaattcagaa	acttgatatt	tttcattttt	ttgctgttca	5280
	gggatttgca	gcataatcatg	gcgtgtaata	tggaagaaatgc	cgtatgtttc	cttatatggc	5340
	tttttggttcg	tttcttttcgc	aaacgcttga	gttgcgccctc	ctgccagcag	tgcggtagta	5400
5	aagggttaata	ctggttgcttg	ttttgcaaac	tttttgatgt	tcacgttca	tgtctccttt	5460
	tttatgtact	gtggttagcgg	tctgcttctt	ccagccctcc	tggttgaaga	tggcaagtta	5520
	gttacgcaca	ataaaaaaag	acctaaaaata	tgtaaggggt	gacgccaaaag	tatacacttt	5580
	gccctttaca	catttttaggt	cttgccctgct	ttatcagtaa	caaacccgcg	cgatttactt	5640
	ttcgacctca	ttctattaga	ctctcgtttg	gattgcaact	ggtctatttt	cctcttttgt	5700
	ttgatagaaa	atcataaaaag	gatttgcaga	ctacgggcct	aaagaactaa	aaaatctatc	5760
	tgtttctttt	catttctctgt	attttttata	gttttctgttg	catgggcata	aagttgcctt	5820
10	tttaatcaca	attcagaaaa	tatcataata	tctcatttca	ctaaataata	gtgaacggca	5880
	ggtatatgtg	atgggttaaa	aaggatcggc	ggccgctcga	tttaaatac		5928

	<210> 18	
	<211> 20	
15	<212> DNA	
	<213> Artificial Sequence	

	<220>	
	<223> synthetic construct	

	<400> 18	
20	tagctgcca ttattccggg	20

	<210> 19	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	

	<220>	
25	<223> synthetic construct	

	<400> 19	
	gggtaaaaaa tcctttcgta	20

	<210> 20	
	<211> 37	
30	<212> DNA	
	<213> Artificial Sequence	

	<220>	
	<223> synthetic construct	

	<400> 20	
35	cccgaataa ttggcagcta ctgaagggcc tcctggg	37

	<210> 21	
	<211> 31	
	<212> DNA	
	<213> Artificial Sequence	

	<220>	
40	<223> synthetic construct	

	<400> 21	
	tatcaacgcg ttcttcatcg gtagcagcac c	31

	<210> 22	
	<211> 40	
45	<212> DNA	
	<213> Artificial Sequence	

	<220>	
	<223> synthetic construct	

	<400> 22	
50	tacgaagga ttttttacct atgaacctaa agaaccgccga	40

	<210> 23	
	<211> 31	
	<212> DNA	
55	<213> Artificial Sequence	

	<220>	
--	-------	--

<223> synthetic construct

<400> 23

atctacgtcg acccaggatg ccctggattt c

31

<210> 24

<211> 5920

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 24

cgcggttcttc	atcggtagca	gcacccgaga	ccatgacgcy	ggcatcgccc	agatccatca	60
cacgcagatc	acgcacatca	gattccctgtg	aggtgtaaat	tcccacgtcg	tggccatcaa	120
gatcataaga	ctcagaaaaga	tcacgccagc	gagtatcata	accagccaca	gcatcctcaa	180
cggtttcacc	agtttgagtg	agctgaatat	agccctcatc	tgcggtgaca	tatccaacta	240
cagatgcccgg	ggtgtcatcc	accatgggtgc	gtcagagctga	atttgtggtc	cagccttcag	300
gagtttccgg	caacctagtt	gcatgatcag	tcattgcygc	cgcttccatt	gacataaaag	360
tggaaagcatc	aacttcaggt	acctgccccat	tttcaggggga	tcctgtattg	aaagaacaca	420
ttcccgtgaa	tcccaccgct	accaacatga	tgatcgcgga	gactaccaac	gagataatca	480
tgtctcgact	gccatcaaaa	attttcggtc	gtttctcagc	caccgccta	gtatgtcacg	540
agtttgggtac	gaaacccct	tttgggtgtc	cagaatccaa	aattccgggc	acaaaagtgc	600
aacaatagat	gacgtgcygc	ttgatacagc	ccaagcgccg	atacatttat	aatgcgcta	660
gatacgtgca	acccacgtaa	ccaggtcaga	tcaagtgcgc	caggaggccc	ttcagtagct	720
gccaattatt	ccgggcttgt	gaccgcctac	ccgataaata	ggtcggctga	aaaatttcgt	780
tgcaatatca	acaaaaaggc	ctatcattgg	gaggtgtcgc	accaagtact	tttgcgaagc	840
gccatctgac	ggattttcaa	aagatgtata	tgctcgggtg	ggaaacctac	gaaaggattt	900
tttaccatg	aacctaaaga	accccgaaac	gccagaccgt	aaccttgcta	tggagctggt	960
gcgagttacg	gaagcagctg	cactggcttc	tggacgttgg	gttggacgtg	gcatgaagaa	1020
tgaaggcgac	ggtgccgctg	ttgacgccat	gcgccagctc	atcaactcag	tgaccatgaa	1080
gggctgcgtt	gttatcgcg	agggcgaaaa	agacgaagct	ccaatgctgt	acaacggcga	1140
agaggtcgga	accggctttg	gacctgaggt	tgatatcgca	gttgaccag	ttgacggcac	1200
caccctgatg	gctgagggtc	gcccacacgc	aatttccatt	ctcgcagctg	cagagcggtg	1260
caccatgtac	gatccatcct	ccgtcttcta	catgaagaag	atcgccgtgg	gacctgaggc	1320
cgcaggcaag	atcgacatcg	aagctccagt	tgcccacaac	atcaacgcgg	tggcaaatgc	1380
caagggaatc	aacctttcgc	acgtcacctg	tgctgtgctt	gaccgtcttc	gccacatcga	1440
actgatcgca	gacattcgtc	gtgcaggcgc	aaaggttcgt	ctcatctccg	acggcgacgt	1500
tgacagtgca	gttgacagag	ctcaggattc	caactccgtg	gacatcatga	tgggcaccgg	1560
cggaacccca	gaaggcatca	tactgctgtg	cgccatgaag	tgcatgggtg	gcgaaatcca	1620
gggcatcctg	ggtcgacatc	gatgctcttc	tgctgttaatt	aacaattggg	atcctctaga	1680
cccgggattt	aaatcgctag	cggtctgcta	aaggaagcgg	aacacgtaga	aagccagttc	1740
gcagaaacgg	tgctgacccc	ggatgaatgt	cagctactgg	gctatctgga	caagggaaaa	1800
cgcaagcgca	aagagaaaag	aggtagcttg	cagtgggcct	acatggcgat	agctagactg	1860
ggcggtttta	tggacagcaa	gcgaaccgga	attgccagct	ggggcgccct	ctggtaagggt	1920
tgggaagccc	tgcaaaagta	actggatggc	tttcttgccg	ccaaggatct	gatggcgag	1980
gggcatcaaga	tctgatcaag	agacaggatg	aggatcgctt	cgcatgattg	aacaagattg	2040
attgcacgca	ggttctccgg	ccgcttgggt	ggagaggcta	ttcggctatg	actgggcaca	2100
acagacaatc	ggctgtctcg	atgccgcccgt	gttccggctg	tcagcgagc	ggcgcccggg	2160
tctttttgtc	aagaccgacc	tgtccgggtg	cctgaatgaa	ctgcaggacg	aggcagcgcg	2220
gctatctgtg	ctggccacga	cgggcgcttc	ttgcgcagct	gtgctcgacg	ttgtcatctga	2280
agcgggaagg	gactggctgc	tattgggcga	agtgcggggg	caggatctcc	tgctcatctca	2340
ccttctctct	gccgagaaaag	tatccatcat	ggctgatgca	atgcggcgcc	tgcatatcgt	2400
tgatccggct	acctgcccac	tcgaccacca	agcgaaaacat	cgcatcgagc	gagcagctac	2460
tcggatggaa	gccggtcttg	tcgatcagga	tgatctggac	gaagagcatc	aggggctcgc	2520
gccagccgaa	ctgttcgcca	ggctcaaggc	gcgcatgccc	gacggcgagg	atctcgtcgt	2580
gacccatggc	gatgcctgct	tgccgaatat	catggtggaa	aatggccgct	tttctggatt	2640
catcgactgt	ggccggctgg	gtgtggcgga	ccgctatcag	gacatagcgt	tggctaccgg	2700
tgatattgct	gaagagcttg	gcggcgaaatg	ggctgaccgc	ttcctcgtgc	tttacggtat	2760
cgccgctccc	gattcgacag	gcactgcctt	ctatcgccct	cttgacgagt	tcttctgagc	2820
gggactctgg	ggttcgaaat	gaccgaccaa	gcgacgccc	acctgccatc	acgagatttc	2880
gattccaccg	ccgccttcta	tgaaagggtg	ggcttcggaa	tcgttttccg	ggacgccggc	2940
tggatgatcc	tccagcgcg	ggatctcatg	ctggagtctt	tcgcccacgc	tagcggcgcg	3000
ccggccggcc	cgggtgtgaa	taccgcacag	atgcgtaaag	agaaaatacc	gcatcaggcg	3060
ctcttcgct	tcctcgctca	ctgactcgct	gcgctcggtc	gttcggctgc	ggcgagcggt	3120
atcagctcac	tcaaaaggcg	taatacgggt	atccacagaa	tcaggggata	acgcaggaaa	3180
gaacatgtga	gcaaaaggcc	agcaaaaggc	caggaaccgt	aaaaaggccg	cgttgctggc	3240
gtttttccat	aggctccgcc	cccctgacga	gcatcacaaa	aatcgacgct	caagtcagag	3300
gtggcgaaac	ccgacaggac	tataaagata	ccaggcgctt	ccccctggaa	gctccctcgt	3360
gcgctctcct	gttccgacct	tgccgcttac	cggataacctg	tccgcctttc	tccctctggg	3420
aagcgtggcg	ctttctcata	gctcacgctg	taggtatctc	agttcgggtg	aggtctgttcg	3480
ctccaagctg	ggctgtgtgc	acgaaccccc	cgttcagccc	gaccgctgcy	ccttatccgg	3540
taactatcgt	cttgagtcca	acccggtaag	acacgactta	tcgccactgg	cagcagccac	3600
tggtaacagg	attagcagag	cgaggatgtg	agggcggtgt	acagagttct	tgaagtgggtg	3660
gcctaactac	ggctacacta	gaaggacagt	atttggatgc	tgcgctctgc	tgaagccagt	3720
taccttcgga	aaaagagttg	gtagctcttg	atccggcaaa	caaaccaccg	ctggtagcgg	3780

5 tgggttttttt gtttgcaagc agcagattac ggcgagaaaa aaaggatctc aagaagatcc 3840
 tttgatcttt tctacggggt ctgacgctca gtggaacgaa aactcacgtt aagggatttt 3900
 ggatcatgaga ttatcaaaaa ggatcttcac ctagatcctt ttaaaggccg gccgcggccg 3960
 ccatcggcat tttcttttgc gtttttattt gtttaactgtt aattgtcctt gttcaaggat 4020
 gctgtctttg acaacagatg ttttcttgcc tttgatgttc agcaggaagc tcggcgcaaa 4080
 cgttgattgt ttgtctgctg agaatcctct gtttgtcata tagcttgtaa tcacgacatt 4140
 gtttcctttc gcttgaggta cagcgaagtg tgagtaagta aaggttacat cgtaggagtc 4200
 aagatccatt ttttaacacaa ggccagtttt gttcagcggc ttgtatgggc cagttaaaga 4260
 attagaaaca taaccaagca tgtaaataatc gtttagacgta atgccgtcaa tcgtcatttt 4320
 10 tgatccgcgg gagtcagtga acaggtagca tttgccgttc attttaaaga cgttcgcgcg 4380
 ttcaattttca tctgtttactg tgtagatgac aatcagcggg ttcatacactt ttttcagtgt 4440
 gtaatcatcg ttttagctcaa tcataccgag agcgcggttt gctaactcag ccgtgcgttt 4500
 tttatcgctt tgcagaagtt tttgactttc ttgacggaag aatgatgtgc ttttgccata 4560
 gtatgctttg ttaaataaag attcttcgcc ttggtagcca tcttcagttc cagtgtttgc 4620
 ttcaaatact aagtatttgt ggcctttatc ttctacgtag tgaggatctc tcagcgtagt 4680
 15 gttgtcgctt gagctgtagt tgccttcacg gatgaactgc tgtagctttt gatacgtttt 4740
 tccgtcacccg tcaaagattg atttataatc ctctacaccg ttgatgttca aagagctgtc 4800
 tgatgctgat acgttaactt gtgcagttgt cagtgtttgt ttgccgtaat gtttaccgga 4860
 gaaatcagtg tagaataaac ggatttttcc gtcagatgta aatgtggctg aacctgacca 4920
 ttcttgtgtt tggcttttta ggatagaatc atttgcacg aatttgtcgc tgtctttaaa 4980
 20 gacgcggcca gcgtttttcc agctgtcaat agaagtctcg ccgacttttt gatagaacat 5040
 gtaaatcgat gtgtcatccg catttttagg atctccggct aatgcaaaga cgatgtggta 5100
 gccgtgatag tttgcgacag tgccgtcagc gttttgtaat ggccagctgt cccaaacgtc 5160
 caggcctttt gcagaagaga tatttttaat tgtggacgaa tcaaattcag aaacttgata 5220
 tttttcattt ttttgcgtgt cagggatttg cagcatatca tggcggtgaa tatgggaaat 5280
 gccgtatgtt tccttatatg gcttttggtt cgtttctttc gcaaacgctt gagttgcgcc 5340
 tcctgccagc agtgcggtag taaaggttaa tactgttgct tgttttgcaa actttttgat 5400
 25 gttcatcggt catgtctcct tttttatgta ctgtgttagc ggtctgcttc ttccagccct 5460
 cctgtttgaa gatggcaagt tagttacgca caataaaaaa agacctaaaa tatgtaaggg 5520
 gtgacgcaa agtatacact ttgcccttta cacattttag gtcttgctg ctttatcagt 5580
 aacaacccg cgcgatttac ttttcgacct cattctatta gactctcggt tggattgcaa 5640
 ctggtctatt ttctctttt gtttgataga aaatcataaa aggatttgca gactacgggc 5700
 ctaaagaact aaaaaatcta tctgtttcct ttatttctct gtatttttta tagtttctgt 5760
 30 tgcatgggca taaagttgcc tttttaatca caattcagaa aatatcataa tatctcattt 5820
 cactaaataa tagtgaacgg caggtatatg tgatgggtta aaaaggatcg gcggccgctc 5880
 gatttaaatc tcgagaggcc tgacgtcggg cccggtacca 5920

35

40 **Claims**

1. A method for increasing metabolic flux through the pentose phosphate pathway in a microorganism comprising culturing a microorganism comprising a gene which is deregulated under conditions such that metabolic flux through the pentose phosphate pathway is increased.
2. The method of claim 1, wherein fructose or sucrose is used as a carbon source.
3. The method of claim 1, wherein fructose is used as a carbon source.
4. The method of claim 1, wherein the gene is fructose-1,6-bisphosphatase.
5. The method of claim 4, wherein the fructose-1,6-bisphosphatase gene is derived from *Corynebacterium*.
6. The method of claim 4, wherein the fructose-1,6-bisphosphatase gene is overexpressed.
7. The method of claim 1, wherein the gene encodes fructose-1,6-bisphosphatase.
8. The method of claim 7, wherein fructose-1,6-bisphosphatase has increased activity.

9. The method of claim 1, wherein the microorganism is a Gram positive microorganism.
10. The method of claim 1, wherein the microorganism belongs to the genus *Corynebacterium*.
- 5 11. The method of claim 10, wherein the microorganism is *Corynebacterium glutamicum*.
12. The method of claim 1, wherein the microorganism is fermented to produce a fine chemical.
13. The method of claim 1, wherein the microorganism further comprises one or more additional deregulated genes.
- 10 14. The method of claim 13, wherein the one or more additional deregulated gene is selected from the group consisting of an ask gene, a dapA gene, an asd gene, a dapB gene, a ddh gene, a lysA gene, a lysE gene, a pycA gene, a zwf gene, a pepCL gene, a gap gene, a zwf gene, a tkt gene, a tad gene, a mqo gene, a tpi gene, a pgk gene, and a . sigC gene.
- 15 15. The method of claim 14, wherein the one or more additional deregulated gene is overexpressed.
16. The method of claim 13, wherein the one or more additional deregulated gene encodes a protein selected from the group consisting of a feed-back resistant aspartokinase, a dihydrodipicolinate synthase, an aspartate semialdehyde dehydrogenase, a dihydrodipicolinate reductase, a diaminopimelate dehydrogenase, a diaminopimelate epimerase, a lysine exporter, a pyruvate carboxylase, a glucose-6- phosphate dehydrogenase, a phosphoenolpyruvate carboxylase, a glyceraldehyde-3-, phosphate dehydrogenase, an RPF protein precursor, a transketolase, a transaldolase, a menaquinine oxidoreductase, a triosephosphate isomerase, a 3-phosphoglycerate kinase, and an RNA-polymerase sigma factor sigC.
- 20 17. The method of claim 16, wherein the protein has increased activity.
18. The method of claim 13, wherein the one or more additional deregulated gene is selected from the group consisting of a pepCK gene, a mal E gene, a glgA gene, a pgi gene, la dead gene, a menE gene, a citE gene, a mke17 gene, a poxB gene, a zwa2 gene, and alsuC gene.
- 30 19. The method of claim 18, wherein the one or more additional deregulated gene is attenuated, decreased or repressed.
20. The method of claim 13, wherein the one or more additional deregulated gene encodes a protein selected from the group consisting of a phosphoenolpyruvate carboxykinase, a malic enzyme, a glycogen synthase, a glucose-6-phosphate isomerase, an ATP dependent RNA helicase, an o-succinylbenzoic acid-CoA ligase, a citrate lyase beta chain, a transcriptional regulator, a pyruvate dehydrogenase, an RPF protein precursor, and a Succinyl-CoA-Synthetase.
- 35 21. The method of claim 20, wherein the protein has decreased activity.
22. A method for producing a fine chemical comprising:
 - a) culturing a microorganism in which fructose-1,6-bisphosphatase is deregulated; and
 - b) accumulating the fine chemical in the medium or in the cells of the microorganisms, thereby producing a fine chemical.
- 45 23. A method for producing a fine chemical comprising culturing a microorganism in which at least one pentose phosphate biosynthetic pathway gene or enzyme is deregulated under conditions such that the fine chemical is produced.
- 50 24. The method of claim 23, wherein said biosynthetic gene is fructose-1,6-bisphosphatase.
25. The method of claim 23, wherein said biosynthetic enzyme is fructose-1,6-bisphosphatase.
- 55 26. The method of claim 22 or 24, wherein fructose-1,6-bisphosphatase expression is increased.
27. The method of claim 22 or 25, wherein fructose-1,6-bisphosphatase activity is increased.

28. The method of claim 22 or 23, further comprising recovering the fine chemical.

29. The method of claim 22 or 23, wherein the microorganism is a Gram positive microorganism.

30. The method of claim 29, wherein the one or more additional deregulated gene is selected from the group consisting of an ask gene, a dapA gene, an asd gene, a dapB gene, a ddh gene, a lysA gene, a lysE gene, a pycA gene, a zwf gene, a pepCL gene, a gap gene, a zwal gene, a tkt, gene, a tad gene, a mqo gene, a tpi gene, a pgk gene, and a sigC gene.

31. The method of claim 30, wherein the one or more additional deregulated gene is overexpressed.

32. The method of claim 29, wherein the one or more additional deregulated gene encodes a protein selected from the group consisting of a feed-back resistant aspartokinase, a dihydrodipicolinate synthase, an aspartate semialdehyde dehydrogenase, a dihydrodipicolinate reductase, a diaminopimelate dehydrogenase, a diaminopimelate epimerase, a lysine exporter, a pyruvate carboxylase, a glucose-6- phosphate dehydrogenase, a phosphoenolpyruvate carboxylase, a glyceraldehyde-3- phosphate dehydrogenase, an RPF protein precursor, a transketolase, a transaldolase, a menaquinone oxidoreductase, a triosephosphate isomerase, a 3-phosphoglycerate kinase, and an RNA-polymerase sigma factor sigC.

33. The method of claim 32, wherein the protein has increased activity.

34. The method of claim 29, wherein the one or more additional deregulated gene is selected from the group consisting of a pepCK gene, a mal E gene, a glgA gene, a pgi gene, a dead gene, a menE gene, a citE gene, a mikE17 gene, a pox gene, a zwa2 gene, and a sucC gene. a sucC gene.

35. The method of claim 34, wherein the one or more additional deregulated gene is attenuated, decreased or repressed.

36. The method of claim 29, wherein the one or more additional deregulated gene encodes a protein selected from the group consisting of a phosphoenolpyruvate carboxykinase, a malic enzyme, a glycogen synthase, a glucose-6-phosphate isomerase, an ATP dependent RNA helicase, an o-succinylbenzoic acid-CoA ligase, a citrate lyase beta chain, a transcriptional regulator, a pyruvate dehydrogenase, an RPF. protein precursor, and a Succinyl-CoA-Synthetase.

37. The method of claim 36, wherein the protein has decreased activity.

38. The method of claim 22 or 23, wherein the microorganism is a Gram positive microorganism.

39. The method of claim 22 or 23, wherein the microorganism belongs to the genus Corynebacterium.

40. The method of claim 39, wherein the microorganism is Corynebacterium glutamicum.

41. The method of claim 22 or 23, wherein the fine chemical is lysine.

42. The method of claim 41, wherein lysine is produced at a yield of at least 100 g/L.

43. The method of claim 41, wherein lysine is produced at a yield of at least 150 g/L.

44. The method of claim 22 or 23, wherein fructose or sucrose is used as a carbon source.

45. The method of claim 22 or 23, wherein fructose is used as a carbon source.

46. The method of claim 22 or 24, wherein fructose-1,6-bisphosphatase comprises the nucleotide sequence of SEQ ID NO:1.

47. The method of claim 22 or 24, wherein fructose-1,6-bisphosphatase encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2.

48. A recombinant microorganism which has a deregulated pentose phosphate biosynthesis pathway.

49. A recombinant microorganism comprising a deregulated pentose phosphate biosynthesis gene.

50. The recombinant microorganism of claim 49, wherein said deregulated gene is fructose-1,6-bisphosphatase.

5 51. The recombinant microorganism of claim 50, wherein fructose-1,6-bisphosphatase expression is increased.

52. The recombinant microorganism of claim 50, wherein said fructose-1,6-bisphosphatase gene encodes a fructose-1,6-bisphosphatase protein having increased activity.

10 53. The recombinant microorganism of claim 49, wherein the microorganism belongs to the genus *Corynebacterium*.

54. The recombinant microorganism of claim 53, wherein the microorganism is *Corynebacterium glutamicum*.

15 55. A polypeptide encoded by the nucleotide sequence of SEQ ID NO:1, wherein said polypeptide has fructose-1,6-bisphosphatase activity.

20

25

30

35

40

45

50

55




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5 A




FIGURE 5 B

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5759610 A, Nishimoto [0042]
- WO 2004089996 A [0046]
- WO 200100843 A [0046] [0046]
- EP 1108790 A [0046] [0046] [0046] [0046] [0046] [0046] [0046] [0049] [0049] [0049] [0049]
- WO 200100844 A [0046] [0046] [0046] [0046] [0046] [0046] [0046] [0049] [0049] [0049] [0049]
- EP 1103611 A [0049]
- DE 10046870 [0146] [0162] [0183]

Non-patent literature cited in the description

- Amino Acids. **KLEEMANN, A.** ULLMANN'S ENCYCLOPEDIA OF INDUSTRIAL CHEMISTRY. VCH-Verlagsgesellschaft, 1985, vol. A2, 57-97 [0002]
- **K. NAKAYAMA et al.** *Nutritional Improvement of Food and Feed Proteins*, 1978, 649-661 [0002]
- **SAHM et al.** *Ann. N. Y. Acad. Sci.*, 1996, vol. 782, 25-39 [0003]
- **FLORES, N. et al.** *Nature Biotechnol.*, 1996, vol. 14, 620-623 [0003]
- **MALIN ; BOURD.** *Journal of Applied Bacteriology*, 1991, vol. 71, 517-523 [0003]
- **SHIO et al.** *Agricultural and Biological Chemistry*, 1990, vol. 54, 1513-1519 [0003]
- **YAMAMOTO et al.** *Journal of Fermentation Technology*, 1986, vol. 64, 285-291 [0003]
- **J. VALLINO et al.** *Biotechnol. Bioeng.*, 1993, vol. 41, 633-646 [0005]
- Nucleotides and related compounds. **KUNINAKA, A. et al.** *Biotechnology*. VCH, 1996, vol. 6, 561-612 [0027]
- Vitamins. Ullmann's Encyclopedia of Industrial Chemistry. VCH, 1996, vol. A27, 443-613 [0027]
- Nutrition, Lipids, Health, and Disease. **ONG, A.S. ; NIKI, E. ; PACKER, L.** Proceedings of the UNESCO/Confederation of Scientific and Technological Associations in Malaysia, and the Society for Free Radical Research. AOCS Press, 01 September 1994 [0027]
- **CANE et al.** *Science*, 1998, vol. 282, 63-68 [0027]
- **GUTCHO.** *Chemicals by Fermentation*. Noyes Data Corporation, 1983 [0027]
- Ullmann's Encyclopedia of Industrial Chemistry. VCH, 1985, vol. A2, 57-97 [0028]
- **STRYER, L.** *Biochemistry*. 1988, 578-590 [0028]
- Amino acids - technical production and use. **LEUCHTENBERGER, W. et al.** *Biotechnology*. VCH, 1996, vol. 6, 466-502 [0029]
- Ullmann's Encyclopedia of Industrial Chemistry. 1985, vol. A2, 57-97 [0029]
- **UMBARGER, H.E.** *Ann. Rev. Biochem.*, 1978, vol. 47, 533-606 [0030]
- Amino Acid Degradation and the Urea Cycle. **STRYER, L.** *Biochemistry*. 1988, 495-516 [0031]
- Biosynthesis of Amino Acids and Heme. **STRYER, L.** *Biochemistry*. 1988, 575-600 [0031]
- Vitamins. Ullmann's Encyclopedia of Industrial Chemistry. VCH, 1996, vol. A27, 443-613 [0032] [0033]
- **MICHAL, G.** *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*, 1999 [0033]
- Nutrition, Lipids, Health, and Disease. **ONG, A.S. ; NIKI, E. ; PACKER, L.** Proceedings of the UNESCO/Confederation of Scientific and Technological Associations in Malaysia, and the Society for Free Radical Research. AOCS Press, 01 September 1994, vol. X, 374 [0033]
- **CHRISTOPHERSON, R.I. ; LYONS, S.D.** Potent inhibitors of de novo pyrimidine and purine biosynthesis as chemotherapeutic agents. *Med. Res. Reviews*, 1990, vol. 10, 505-548 [0040]
- **SMITH, J.L.** Enzymes in nucleotide synthesis. *Curr. Opin. Struct. Biol.*, 1995, vol. 5, 752-757 [0040]
- *Biochem Soc. Transact.*, 1995, vol. 23, 877-902 [0040]
- **KUNINAKA, A. et al.** Nucleotides and Related Compounds in Biotechnology. VCH, 1996, vol. 6, 561-612 [0040]
- **ZALKIN, H. ; DIXON, J.E.** de novo purine nucleotide biosynthesis. *Progress in Nucleic Acid Research and Molecular Biology*, 1992, vol. 42, 259-287 [0041]
- Nucleotides and Nucleosides. **MICHAL, G.** *Biochemical Pathways: An Atlas of Biochemistry and Molecular Biology*. Wiley, 1999 [0041]
- **SINGER, M.A. ; LINDQUIST, S.** *Trends Biotech.*, 1998, vol. 16, 460-467 [0042]
- **PAIVA, C.L.A. ; PANEK, A.D.** *Biotech. Ann. Rev.*, 1996, vol. 2, 293-314 [0042]
- **SHIOSAKA, M.** *J. Japan*, 1997, vol. 172, 97-102 [0042]

- Current Protocols in Molecular Biology. John Wiley & Sons, 1989, 6.3.1-6.3.6 [0086]
- **SAMBROOK, J. ; FRITSH, E. F. ; MANIATIS, T.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, 1989 [0087] [0089] [0099]
- Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1995 [0088]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Press, 1989 [0088]
- **CHURCH ; GILBERT.** *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 1991-1995 [0088]
- **KARLIN ; ALTSCHUL.** *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 2264-68 [0111]
- **KARLIN ; ALTSCHUL.** *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 5873-77 [0111]
- **ALTSCHUL et al.** *J. Mol. Biol.*, 1990, vol. 215, 403-10 [0111]
- **ALTSCHUL et al.** *Nucleic Acids Research*, 1997, vol. 25 (17), 3389-3402 [0111]
- **MYERS ; MILLER.** *Comput Appl Biosci.*, 1988, vol. 4, 11-17 [0111]
- **WITTMANN, C. ; E. HEINZLE.** *Appl. Environ. Microbiol.*, 2002, vol. 68, 5843-5859 [0114] [0121] [0123] [0133]
- **RUBINO, F. M.** *J. Chromatogr.*, 1989, vol. 473, 125-133 [0119]
- **WITTMANN, C. ; M. HANS ; E. HEINZLE.** *Analytical Biochem.*, 2002, vol. 307, 379-382 [0119]
- **WITTMANN, C. ; E. HEINZLE.** *Appl. Environ. Microbiol.*, 2002, vol. 68, 5843-5859 [0120]
- **OHNISHI, J. ; S. MITSUHASHI ; M. HAYASHI ; S. ANDO ; H. YOKOI ; K. OCHIAI ; M. A. IKEDA.** *Appl. Microbiol. Biotechnol.*, 2002, vol. 58, 217-223 [0121]
- **DOMINGUEZ, H. ; C. ROLLIN ; A. GUYONVARCH ; J. L. GUERQUIN-KERN ; M. COCAIGN-BOUSQUET ; N. D. LINDLEY.** *Eur. J. Biochem.*, 1998, vol. 254, 96-102 [0121] [0132] [0132] [0133] [0135] [0136] [0139]
- **MARX, A. ; A. A. DE GRAAF ; W. WIECHERT ; L. EGGELING ; H. SAHM.** *Biotechnol. Bioeng.*, 1996, vol. 49, 111-129 [0124] [0133]
- **KIEFER, P., E. HEINZLE ; C. WITTMANN.** *J. Ind. Microbiol. Biotechnol.*, 2002, vol. 28, 338-43 [0132]
- **KIEFER, P. ; E. HEINZLE ; C. WITTMANN.** *J. Ind. Microbiol. Biotechnol.*, 2002, vol. 28, 338-43 [0132] [0134]
- **WITTMANN, C. ; E. HEINZLE.** *Eur. J. Biochem.*, 2001, vol. 268, 2441-2455 [0133]
- **DOMINGUEZ, H. ; C. ROLLIN ; A. GUYONVARCH ; J. L. GUERQUIN-KERN ; M. COCAIGN-BOUSQUET ; N. D. LINDLEY.** *Eur. J. Biochem.*, 1998, vol. 254, 96-102 [0135]
- **PETERSEN, S. ; A. A. DE GRAAF ; L. EGGELING ; M. M611NEY ; W. WIECHERT ; H. SAHM.** *J. Biol. Chem.*, 2000, vol. 75, 35932-35941 [0135]
- **IKEDA, M.** *Adv. Biochem. Eng. Biotechnol.*, 2003, vol. 79, 1-36 [0137]
- **SKRYPAL, I. G. ; O. V. LASTREBOVA.** *Mikrobiol Z.*, 2002, vol. 64, 82-94 [0137]
- **WENDISCH, V. F. ; A. A. DE GRAAF ; H. SAHM H. ; B. EIKMANS.** *J. Bacteriol.*, 2000, vol. 182, 3088-3096 [0137]
- **COCAIGN-BOUSQUET, M. ; N. D. LINDLEY.** *Enz. Microbiol. Technol.*, 1995, vol. 17, 260-267 [0139]
- **DOMINGUEZ, H. ; N. D. LINDLEY.** *Appl. Environ. Microbiol.*, 1996, vol. 62, 3878-3880 [0141]
- **TAUCH et al.** *Plasmid*, 1995, vol. 33, 168-179 [0142] [0148] [0168]
- **EIKMANS et al.** *Microbiology*, 1994, vol. 140, 1817-1828 [0142] [0148] [0168]
- **LENNOX.** *Virology*, 1955, vol. 1, 190 [0143] [0159]
- **SANGER et al.** *Proceedings of the National Academy of Sciences USA*, 1977, vol. 74, 5463-5467 [0143] [0160] [0181]
- **LIEBL et al.** *FEMS Microbiology Letters*, 1989, vol. 53, 299-303 [0146] [0162] [0183]
- **INNIS et al.** PCR Protocols. A Guide to Methods and Applications. Academic Press, 1990 [0149] [0151] [0153] [0155] [0156] [0169] [0177]
- **SAMBROOK et al.** Molecular Cloning. A Laboratory Manual. Cold Spring Harbor, 1989 [0159] [0180] [0183]
- **KNIS et al.** PCR Protocols, A Guide to Methods and Applications. Academic Press, 1990 [0171]
- **LNNIS et al.** PCR Protocols. A Guide to Methods and Applications. Academic Press, 1990 [0173]
- **INNIS et al.** PCR Protocols, A Guide to Methods and Applications. Academic Press, 1990 [0175]
- **LENNOX.** *Virology*, 1955 [0180]