

(19)



(11)

EP 1 963 276 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 4

(48) Corrigendum issued on:
26.09.2012 Bulletin 2012/39

(45) Date of publication and mention
of the grant of the patent:
20.06.2012 Bulletin 2012/25

(21) Application number: **06789243.0**

(22) Date of filing: **03.08.2006**

(51) Int Cl.:
C07D 231/56 ^(2006.01) **C07D 209/48** ^(2006.01)
C07D 209/44 ^(2006.01) **C07D 401/12** ^(2006.01)
C07D 403/12 ^(2006.01) **C07D 471/04** ^(2006.01)
A61K 31/416 ^(2006.01) **A61K 31/4035** ^(2006.01)
A61P 29/00 ^(2006.01)

(86) International application number:
PCT/US2006/030185

(87) International publication number:
WO 2007/073405 (28.06.2007 Gazette 2007/26)

(54) **N-LINKED ARYL HETEROARYL INHIBITORS OF LTA4H FOR TREATING INFLAMMATION**

N-GEBUNDENE ARYLHETEROARYLINHIBITOREN VON LTA4H ZUR BEHANDLUNG VON
ENZÜNDUNGEN

INHIBITEURS ARYLE HETEROARYLE A LIAISON AZOTE DE LTA4H UTILISES DANS LE
TRAITEMENT DE L'INFLAMMATION

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR
Designated Extension States:
AL BA HR MK RS

(30) Priority: **21.12.2005 US 752658 P**

(43) Date of publication of application:
03.09.2008 Bulletin 2008/36

(73) Proprietor: **Decode Genetics EHF**
101 Reykjavik (IS)

(72) Inventors:

- **SANDANAYAKA, Vincent**
Woodridge, Illinois 60517 (US)
- **SINGH, Jasbir**
Woodridge, Illinois 60517 (US)
- **ZHAO, Lei**
Woodridge, Illinois 60517 (US)
- **GURNEY, Mark**
Woodridge, Illinois 60517 (US)

(74) Representative: **Graf von Stosch, Andreas et al**
Graf von Stosch
Patentanwalts-gesellschaft mbH
Prinzregentenstrasse 22
80538 München (DE)

(56) References cited:
EP-A1- 0 418 385 WO-A-2004/065370
WO-A-2005/042496 US-A- 4 448 784
US-A1- 2002 077 329 US-A1- 2004 157 849
US-A1- 2005 043 378 US-A1- 2005 043 379

- **FRAY M JONATHAN ET AL: "Novel Antagonists of Platelet-Activating Factor: 1. Synthesis and Structure-Activity Relationships of Benzodiazepine and Benzazepine Derivatives of 2-Methyl-1-phenylimidazo(4,5-c)pyridine" JOURNAL OF MEDICINAL CHEMISTRY, vol. 38, no. 18, 1995, pages 3514-3523, XP002414336 ISSN: 0022-2623**
- **MASSIMO G ET AL: "BIOLOGICAL STUDIES ON 1 2 BENZISOTHIAZOLE DERIVATIVES III. EVALUATION OF ANTIBACTERIAL ANTIFUNGAL AND DNA-DAMAGING ACTIVITIES OF 1 2 BENZISOTHIAZOLIN-3-ONES" FARMACO (ROME), vol. 45, no. 4, 1990, pages 439-446, XP009077002 ISSN: 0014-827X**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 963 276 B9

- DATABASE BEILSTEIN Beilstein Institut zur Foerderung der chemischen Wissenschaft, Frankfurt am Main, DE; XP002414338 Database accession no. 8581995(BRN) & ARIEF M.M.H; AMINE M.S.; EISSA A. M. F.: EGYPT. J. CHEM., vol. 42, no. 6, 1999, pages 563-572,
- ARTICO M ET AL: "RESEARCH ON COMPOUNDS WITH ANTIBLASTIC ACTIVITY NOTE XXXVI - DERIVATIVES OF 3-P-AMINOPHENYL-2-OXAZOLIDINONE: SYNTHESIS AND ANTINEOPLASTIC ACTIVITIES" FARMACO, EDIZIONE SCIENTIFICA, SOCIETA CHIMICA ITALIANA, PAVIA, IT, vol. 2, no. 24, 1969, pages 179-190, XP001070657 ISSN: 0430-0920
- PENNING T D: "INHIBITORS OF LEUKOTRIENE A4 (LTA4) HYDROLASE AS POTENTIAL ANTI-INFLAMMATORY AGENTS" CURRENT PHARMACEUTICAL DESIGN, BENTHAM SCIENCE PUBLISHERS, SCHIPHOL, NL, vol. 7, no. 3, February 2001 (2001-02), pages 163-179, XP001205198 ISSN: 1381-6128
- PENNING THOMAS D ET AL: "Structure-Activity Relationship Studies on 1-[2-(4- Phenylphenoxy) ethyl]pyrrolidine (SC-22716), a Potent Inhibitor of Leukotriene A4 (LTA4) Hydrolase" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 43, 2000, pages 721-725, XP002197495 ISSN: 0022-2623

Description**Field of the Invention**

[0001] The present invention relates to a chemical genus of biaryl nitrogen-attached heterocycles that are inhibitors of LTA4H (leukotriene A4 hydrolase). They are useful for the treatment and prevention and prophylaxis of inflammatory diseases and disorders.

Background of the Invention

[0002] The end products of the leukotriene pathway are potent inflammatory lipid mediators derived from arachidonic acid. They can potentially contribute to development of atherosclerosis and destabilization of atherosclerotic plaques through lipid oxidation and/or proinflammatory effects. As described elsewhere, a gene on chromosome 13q12 has been identified as playing a major role in myocardial infarction (MI), [Helgadottir et al., Nature Genetics doi:10.1038/ng1311, 8 Feb 2004]. This gene (ALOX5AP), herein after referred to as an MI disease gene, comprises nucleic acid that encodes 5-lipoxygenase activating protein (FLAP), herein after referred to as FLAP. DNA variants in the FLAP gene increase risk for myocardial infarction by 1.8 fold and for stroke by 1.7 fold. The leukotriene pathway, through FLAP, leads to the production of leukotriene B4 by the enzyme leukotriene A4 hydrolase (LTA4H). Leukotriene B4 is one of the most potent chemokine mediators of arterial inflammation. Particular DNA variants in the gene encoding LTA4H also elevate risk for MI and stroke, as described elsewhere [Hakonarsson et al., J.Am.Med.Assoc. 293, 2245-2256 (2005)]. Individuals with a prior history of MI produce more leukotriene B4 when their isolated neutrophils are stimulated with ionomycin. Increased LTB4 production is particularly marked in male patients with a prior history of MI who carry risk variants in the FLAP gene [Helgadottir et al.]. The treatment (prophylactic and/or therapeutic) of certain diseases and conditions (e.g., MI, acute coronary syndrome (ACS), stroke, atherosclerosis) associated with FLAP or with LTA4H can be accomplished by inhibiting LTA4H. Inhibiting LTA4H is advantageous for methods of treatment for MI or susceptibility to MI; for ACS (e.g., unstable angina, non-ST-elevation myocardial infarction (NSTEMI) or ST-elevation myocardial infarction (STEMI)); for decreasing risk of a second MI; for stroke (including transient ischemic attack) or susceptibility to stroke; for atherosclerosis, such as for patients requiring treatment (e.g., angioplasty, stents, coronary artery bypass graft) to restore blood flow in coronary arteries, such as patients requiring treatment for peripheral vascular disease including peripheral occlusive arterial disease, critical limb ischemia (e.g., gangrene, ulceration), and intermittent claudication to restore blood flow in the lower limbs; for atherosclerotic reno-vascular disease; for abdominal aortic aneurysm; and/or for decreasing leukotriene synthesis (e.g., for treatment of MI).


[0003] US Patent Application Publication No. 20050043378 and 20050043379, relate to benzooxazol-2-yl, benzothiazol-2-yl and 1H-benzimidazol-2-yl compounds and derivatives thereof useful as leukotriene A4 hydrolase (LTA4H) inhibitors in treating inflammation and disorders associated with inflammation.

[0004] In other words, US2005/043378 and US2005/043379 disclose leukotriene A4 hydrolase inhibitors (LTA4H inhibitors) exhibiting a phenyl ring attached via a spacer (named Y) to the N-atom of a bicyclic ring. Both documents are exclusively related to compounds exhibiting the spacer Y, which was considered to be indispensable by these documents. The use of any other type of compound is neither suggested nor discussed.


[0005] Penning T. D. (Current Pharmaceutical Design, Bentham Science Publishers, Schiphol, NL, Vol. no. 3, February 2001, pages 163-179) and Penning et al. (Journal of Medicinal Chemistry, American Chemical Society, Washington, US, Vol. 43, 2000, pages 721-725) disclose LTA4H inhibitors exhibiting a phenyl ring directly attached to a C-atom of a monocyclic ring. In addition, Penning et al. discloses a structure-activity relationship study of a linker between the phenyl ring and another phenyl ring (Table 2 of D14) providing evidence that removal of the linker will decrease the affinity. Neither Penning T. D. nor Penning et al. disclose any compound, wherein the phenyl ring is directly attached to an N-atom or any other heteroatom of a ring.

Summary of the Invention

[0006] One aspect of the present invention relates to compounds exhibiting LTA4H enzyme inhibition, having general formula:



10 or



20 wherein

ring (a) is chosen from bicyclic heterocyclyl and bicyclic heterocyclyl substituted with from one to three substituents independently selected from the group consisting of halogen, hydroxyl, loweralkyl, loweracyl, loweralkoxy, fluoroloweralkyl, fluoroloweralkoxy, formyl cyano, phenyl, heteroaryl, nitro and oxo;

25 Q is O;

R⁶ is selected in each occurrence from the group consisting of H and lower alkyl;

30 n is zero or an integer from 1-4;

p is an integer from 1-4;

HET is a 4-7-membered saturated nitrogenous heterocycle; and

35 taken together ZW is H or

Z is (CH₂)₀₋₁₀ in which one or two (CH₂) may optionally be replaced by a C₃-C₆ carbocycle, a C₃-C₆ heterocycle, -O-, -NR¹⁰-, -SO-, -S(O)₂-, -C(=O)- or -C(=O)NH-, provided that said O-, -NR¹⁰-, -SO-, -S(O)₂-, -C(=O)- or -C(=O)NH- are not at the point of attachment to nitrogen and are separated by at least one -(CH₂)₂-;

40 R¹⁰ is selected separately in each occurrence from the group consisting of H and lower alkyl

45 R¹ is chosen from H and lower alkyl; and

W is selected from acyl, hydroxyl, carboxyl, amino, carboxamido, sulfonamide, aminoacyl, -COOalkyl, -CHO, -C(O) fluoroalkyl, -C(O)CH₂C(O)Oalkyl, -C(O)CH₂C(O)Ofuoroalkyl, -SH, -C(O)NH(OH), -C(O)N(OH)R⁴, -N(OH)C(O)OH, -N(OH)C(O)R⁴, heterocyclyl, substituted aryl, and substituted heterocyclyl; and

50 R⁴ is selected from the group consisting of H and lower alkyl.

[0007] A second aspect of the present invention relates to a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound described above.

[0008] A third aspect of the present invention relates to a compound described above for use in inhibiting leukotriene A4 hydrolase where the LTA4H enzyme is contacted with a therapeutically effective amount of a compound described above.


[0009] A fourth aspect of the present invention relates to a compound described above for use in treating a disorder associated with leukotriene A4 hydrolase where a therapeutically effective amount of the compound described above

or a salt, hydrate or ester thereof is administered to a mammal.


Detailed Description of the Invention

[0010] Throughout this specification the substituents are defined when introduced and retain their definitions.

[0011] In one aspect the invention relates to bicyclic and substituted bicyclic heterocycle derivatives useful as LTA4H enzyme inhibitors, having the general formula:






or



Ring (a) is chosen from bicyclic heterocyclyl and bicyclic heterocyclyl substituted with from one to three substituents independently selected from the group consisting of halogen, hydroxyl, loweralkyl, loweracyl, loweralkoxy, fluoroloweralkyl, fluoroloweralkoxy, formyl, cyano, phenyl, heteroaryl, nitro and oxo.


[0012] In some embodiments, ring (a) is an indazole, an indole, or an isoindole ring:






Y is N or CR⁵; R⁷ is chosen from H and loweralkyl; and R⁵ is chosen from H and loweralkyl; n is 1, 2 or 3; and p is 2 or 3.

[0013] An additional embodiment is illustrated by the formula:





or




R² and R³ each are independently selected from the group consisting of H, halogen, hydroxyl, loweralkyl, loweracyl, loweralkoxy, fluoroloweralkyl, fluoroloweralkoxy, formyl, cyano, phenyl, heteroaryl, and nitro; n is 1, 2 or 3; and p is 2 or 3. R² and R³ may be distributed in either ring; for example, both 3,6-dichloroindazole and 5,6-dichloroindazole are encompassed.

[0014] Examples where one or two (CH₂) linkers of Z are optionally replaced by a C₃-C₆ carbocycle or a C₃-C₆ heterocycle include but are not limited to the structures below wherein the wavy lines indicate the point of attachment to HET:







or



[0015] The present invention provides a method for inhibiting leukotriene A4 hydrolase comprising contacting the LTA₄H enzyme with a therapeutically effective amount of a compound according to the general formula



or



[0016] It may be found upon examination that additional species and genera not presently excluded are not patentable to the inventors in this application. In such a case, the exclusion of species and genera in applicants' claims are to be considered artifacts of patent prosecution and not reflective of the inventors' concept or description of their invention. The invention, in a composition aspect, is all compounds of the general formula above, except those that are in the public's possession. The invention, in a method aspect, is a method employing compounds of the general formula above, except those methods that are in the public's possession.

[0017] The present invention provides compounds for use in a method for treating a disorder associated with leukotriene A4 hydrolase comprising administering to a mammal a therapeutically effective amount of a compound or a salt, hydrate or ester thereof according to the general formula given above. In some embodiments the disorder is associated with inflammation. In some embodiments the disorder is selected from allergic inflammation, acute inflammation and chronic inflammation.

[0018] Compounds of the genus represented by the general formula above are inhibitors of LTA₄H enzyme. As such they have utility in treating and preventing inflammatory diseases and disorders, as described above, particularly for such conditions as asthma, chronic obstructed pulmonary disease (COPD), atherosclerosis, rheumatoid arthritis, multiple sclerosis, inflammatory bowel diseases (IBD; including Crohn's disease and ulcerative colitis), or psoriasis, which are each characterized by excessive or prolonged inflammation at some stage of the disease.

[0019] Recent research indicates that the compounds are also useful for treating and preventing atherosclerosis, thrombosis, stroke, acute coronary syndrome, stable angina, peripheral vascular disease, critical leg ischemia, intermittent claudication, abdominal aortic aneurysm and myocardial infarction.

onary syndrome and myocardial infarct.

[0020] The compounds may be presented as salts. The term "pharmaceutically acceptable salt" refers to salts whose counter ion derives from pharmaceutically acceptable non-toxic acids and bases. Suitable pharmaceutically acceptable base addition salts for the compounds of the present invention include, but are not limited to, metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from lysine, N,N-dialkyl amino acid derivatives (e.g. N,N-dimethylglycine, piperidine-1-acetic acid and morpholine-4-acetic acid), N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. When the compounds contain a basic residue, suitable pharmaceutically acceptable base addition salts for the compounds of the present invention include inorganic acids and organic acids. Examples include acetate, benzenesulfonate (besylate), benzoate, bicarbonate, bisulfate, carbonate, camphorsulfonate, citrate, ethanesulfonate, fumarate, gluconate, glutamate, bromide, chloride, isethionate, lactate, maleate, malate, mandelate, methanesulfonate, mucate, nitrate, pamoate, pantothenate, phosphate, succinate, sulfate, tartrate, p-toluenesulfonate, and the like.

[0021] For convenience and clarity certain terms employed in the specification, examples and claims are described herein.

[0022] Alkyl is intended to include linear, branched, or cyclic hydrocarbon structures and combinations thereof. Lower alkyl refers to alkyl groups of from 1 to 6 carbon atoms. Examples of lower alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, s- and t-butyl and the like. Preferred alkyl groups are those of C₂₀ or below. Cycloalkyl is a subset of alkyl and includes cyclic hydrocarbon groups of from 3 to 8 carbon atoms. Examples of cycloalkyl groups include c-propyl, c-butyl, c-pentyl, norbornyl and the like.

[0023] C₁ to C₂₀ hydrocarbon includes alkyl, cycloalkyl, alkenyl, alkynyl, aryl, arylalkyl and combinations thereof. Examples include phenethyl, cyclohexylmethyl, camphoryl, adamantyl and naphthylethyl.

[0024] Alkoxy or alkoxy refers to groups of from 1 to 8 carbon atoms of a straight, branched, cyclic configuration and combinations thereof attached to the parent structure through oxygen. Examples include methoxy, ethoxy, propoxy, isopropoxy, cyclopropyloxy, cyclohexyloxy and the like. Lower-alkoxy refers to groups containing one to four carbons.

[0025] Alkoxyalkyl refers to ether groups of from 3 to 8 atoms of a straight, branched, cyclic configuration and combinations thereof attached to the parent structure through an alkyl. Examples include methoxymethyl, methoxyethyl, ethoxypropyl, and the like.

[0026] Alkoxyaryl refers to alkoxy substituents attached to an aryl, wherein the aryl is attached to the parent structure. Arylalkoxy refers to aryl substituents attached to an oxygen, wherein the oxygen is attached to the parent structure. Substituted arylalkoxy refers to a substituted aryl substituent attached to an oxygen, wherein the oxygen is attached to the parent structure.

[0027] Acyl refers to groups of from 1 to 8 carbon atoms of a straight, branched, cyclic configuration, saturated, unsaturated and aromatic and combinations thereof, attached to the parent structure through a carbonyl functionality. Examples include acetyl, benzoyl, propionyl, isobutyryl, t-butoxycarbonyl, benzyloxycarbonyl and the like. Lower-acyl refers to groups containing one to four carbons.

[0028] Aryl and heteroaryl mean a 5- or 6-membered aromatic or heteroaromatic ring containing 0-3 heteroatoms selected from O, N, or S; a bicyclic 9- or 10-membered aromatic or heteroaromatic ring system containing 0-3 heteroatoms selected from O, N, or S; or a tricyclic 13- or 14-membered aromatic or heteroaromatic ring system containing 0-3 heteroatoms selected from O, N, or S. The aromatic 6- to 14-membered carbocyclic rings include, e.g., benzene and naphthalene, and according to the invention benzoxalane and residues in which one or more rings are aromatic, but not all need be. The 5- to 10-membered aromatic heterocyclic rings include, e.g., imidazole, pyridine, indole, thiophene, benzopyranone, thiazole, furan, benzimidazole, quinoline, isoquinoline, quinoxaline, pyrimidine, pyrazine, tetrazole and pyrazole.

[0029] Arylalkyl refers to a substituent in which an aryl residue is attached to the parent structure through alkyl. Examples are benzyl, phenethyl and the like. Heteroarylalkyl refers to a substituent in which a heteroaryl residue is attached to the parent structure through alkyl. Examples include, e.g., pyridinylmethyl, pyrimidinylethyl and the like. Heterocyclylalkyl refers to a substituent in which a heterocyclyl residue is attached to the parent structure through alkyl. Examples include morpholinoethyl and pyrrolidinylmethyl.

[0030] Heterocycle means a cycloalkyl or aryl residue in which from one to three carbons is replaced by a heteroatom selected from the group consisting of N, O and S. The nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. Examples of heterocycles include pyrrolidine, pyrazole, pyrrole, indole, quinoline, isoquinoline, tetrahydroisoquinoline, benzofuran, benzodioxan, benzodioxole (commonly referred to as methylenedioxyphenyl, when occurring as a substituent), tetrazole, morpholine, thiazole, pyridine, pyridazine, pyrimidine, thiophene, furan, oxazole, oxazoline, isoxazole, dioxane, tetrahydrofuran and the like. It is to be noted that heteroaryl is a subset of heterocycle in which the heterocycle is aromatic. Examples of heterocyclyl residues additionally include piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxo-pyrrolidinyl, 2-oxoazepinyl, azepinyl, 4-piperidinyl, pyrazolidinyl, imidazolyl, imidazolynyl, imidazolidinyl, pyrazinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolyl, quinucridinyl, isothiazolidinyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, tetrahydrofuryl, tetrahydropyranyl,

thienyl, benzothienyl, thiamorpholinyl, thiamorpholinylsulfoxide, thiamorpholinylsulfone, oxadiazolyl, triazolyl and tetrahydroquinolinyl.

[0031] An oxygen heterocycle is a heterocycle containing at least one oxygen in the ring; it may contain additional oxygens, as well as other heteroatoms. A sulphur heterocycle is a heterocycle containing at least one sulphur in the ring; it may contain additional sulphurs, as well as other heteroatoms. A nitrogen heterocycle is a heterocycle containing at least one nitrogen in the ring; it may contain additional nitrogens, as well as other heteroatoms. Oxygen heteroaryl is a subset of oxygen heterocycle; examples include furan and oxazole. Sulphur heteroaryl is a subset of sulphur heterocycle; examples include thiophene and thiazine. Nitrogen heteroaryl is a subset of nitrogen heterocycle; examples include pyrrole, pyridine and pyrazine. A saturated nitrogenous heterocycle is a subset of nitrogen heterocycle. Saturated nitrogenous heterocycle contain at least one nitrogen and may contain additional nitrogens, as well as other heteroatoms. Examples include pyrrolidine, pyrazolidine, piperidine, morpholine, and thiomorpholine.

[0032] Substituted alkyl, aryl, cycloalkyl, heterocyclyl etc. refer to alkyl, aryl, cycloalkyl, or heterocyclyl wherein up to three H atoms in each residue are replaced with halogen, haloalkyl, hydroxy, loweralkoxy, carboxy, carboalkoxy (also referred to as alkoxycarbonyl), carboxamido, cyano, carbonyl, nitro, amino, alkylamino, dialkylamino, mercapto, alkylthio, sulfoxide, sulfone, acylamino, amidino, phenyl, benzyl, heteroaryl, phenoxy, benzyloxy, or heteroaryloxy.

[0033] The terms "halogen" and "halo" refer to fluorine, chlorine, bromine or iodine.

[0034] The term "prodrug" refers to a compound that is made more active *in vivo*.

[0035] Activation *in vivo* may come about by chemical action or through the intermediacy of enzymes. Microflora in the GI tract may also contribute to activation *in vivo*.

[0036] It will be recognized that the compounds of this invention can exist in radiolabeled form, i.e., the compounds may contain one or more atoms containing an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Radioisotopes of hydrogen, carbon, phosphorous, fluorine, and chlorine include ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Compounds that contain those radioisotopes and/or other radioisotopes of other atoms are within the scope of this invention. Tritiated, i.e. ^3H , and carbon-14, i.e., ^{14}C , radioisotopes are particularly preferred for their ease in preparation and detectability. Radiolabeled compounds of the present invention and prodrugs thereof can generally be prepared by methods well known to those skilled in the art. Conveniently, such radiolabeled compounds can be prepared by carrying out the procedures disclosed in the Examples and Schemes by substituting a readily available radiolabeled reagent for a nonradiolabeled reagent.

[0037] As used herein, and as would be understood by the person of skill in the art, the recitation of "a compound" is intended to include salts, solvates, co-crystals and inclusion complexes of that compound.

[0038] The term "solvate" refers to a compound of formula I in the solid state, wherein molecules of a suitable solvent are incorporated in the crystal lattice. A suitable solvent for therapeutic administration is physiologically tolerable at the dosage administered. Examples of suitable solvents for therapeutic administration are ethanol and water. When water is the solvent, the solvate is referred to as a hydrate. In general, solvates are formed by dissolving the compound in the appropriate solvent and isolating the solvate by cooling or using an antisolvent. The solvate is typically dried or azeotroped under ambient conditions. Co-crystals are combinations of two or more distinct molecules arranged to create a unique crystal form whose physical properties are different from those of its pure constituents. Pharmaceutical co-crystals have recently become of considerable interest for improving the solubility, formulation and bioavailability of such drugs as itraconazole [see Remenar et al. J.Am.Chem.Soc. 125, 8456-8457 (2003)] and fluoxetine. Inclusion complexes are described in Remington: The Science and Practice of Pharmacy 19th Ed. (1995) volume 1, page 176-177. The most commonly employed inclusion complexes are those with cyclodextrins, and all cyclodextrin complexes, natural and synthetic, with or without added additives and polymer(s), as described in US Patents 5,324,718 and 5,472,954, are specifically encompassed within the claims.

[0039] The compounds described herein may contain asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms. Each chiral center may be defined, in terms of absolute stereochemistry, as (R)- or (S)-. The present invention is meant to include all such possible isomers, as well as, their racemic and optically pure forms. Optically active (R)- and (S)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. The prefix "*rac*" refers to a racemate. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both *E* and *Z* geometric isomers. The representation of the configuration of any carbon-carbon double bond appearing herein is selected for convenience only, and unless explicitly stated, is not intended to designate a particular configuration. Thus a carbon-carbon double bond depicted arbitrarily as *E* may be *Z*, *E*, or a mixture of the two in any proportion. Likewise, all tautomeric forms are also intended to be included.

[0040] The graphic representations of racemic, ambiscalemic and scalemic or enantiomerically pure compounds used herein are taken from Maehr J. Chem. Ed. 62, 114-120 (1985): solid and broken wedges are used to denote the absolute configuration of a chiral element; wavy lines and single thin lines indicate disavowal of any stereochemical implication which the bond it represents could generate; solid and broken bold lines are geometric descriptors indicating the relative configuration shown but denoting racemic character; and wedge outlines and dotted or broken lines denote enantio-

merically pure compounds of indeterminate absolute configuration.

[0041] Terminology related to "protecting", "deprotecting" and "protected" functionalities occurs throughout this application. Such terminology is well understood by persons of skill in the art and is used in the context of processes that involve sequential treatment with a series of reagents. In that context, a protecting group refers to a group, which is used to mask a functionality during a process step in which it would otherwise react, but in which reaction is undesirable. The protecting group prevents reaction at that step, but may be subsequently removed to expose the original functionality. The removal or "deprotection" occurs after the completion of the reaction or reactions in which the functionality would interfere. Thus, when a sequence of reagents is specified, as it is in the processes of the invention, the person of ordinary skill can readily envision those groups that would be suitable as "protecting groups". Suitable groups for that purpose are discussed in standard textbooks in the field of chemistry, such as Protective Groups in Organic Synthesis by T.W. Greene [John Wiley & Sons, New York, 1991].

[0042] A comprehensive list of abbreviations utilized by organic chemists appears in the first issue of each volume of the Journal of Organic Chemistry. The list is typically presented in a table entitled "Standard List of Abbreviations".

[0043] In general, the compounds of the present invention may be prepared by the methods illustrated in the general reaction schemes as, for example, described below, or by modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants that are in themselves known, but are not mentioned here. The starting materials are either commercially available, synthesized as described in the examples or may be obtained by the methods well known to persons of skill in the art.

[0044] LTA4H inhibitors have been shown to be effective anti-inflammatory agents in pre-clinical studies. For example, oral administration of LTA4H inhibitor SC57461 to rodents resulted in the inhibition of ionophore-induced LTB4 production in mouse blood *ex vivo*, and in rat peritoneum *in vivo* (Kachur et al., 2002, J. Pharm. Exp. Ther. 300(2), 583-587). Furthermore, eight weeks of treatment with the same inhibitor compound significantly improved colitis symptoms in a primate model (Penning, 2001, Curr. Pharm. Des. 7(3): 163-179). The spontaneous colitis that develops in these animals is very similar to human IBD. Therefore persons of skill in the art accept that positive results in LTA4H models are predictive of therapeutic utility in this and other human inflammatory diseases.

[0045] The inflammatory response is characterized by pain, increased temperature, redness, swelling, or reduced function, or by a combination of two or more of these symptoms. The terms inflammation, inflammatory diseases or inflammation-mediated diseases or conditions include, but are not limited to, acute inflammation, allergic inflammation, and chronic inflammation.

[0046] Autoimmune diseases are associated with chronic inflammation. There are about 75 different autoimmune disorders known that may be classified into two types, organ-specific (directed mainly at one organ) and non-organ-specific (affecting multiple organs).

[0047] Examples of organ-specific autoimmune disorders are insulin-dependent diabetes (Type I) which affects the pancreas, Hashimoto's thyroiditis and Graves' disease which affect the thyroid gland, pernicious anemia which affects the stomach, Cushing's disease and Addison's disease which affect the adrenal glands, chronic active hepatitis which affects the liver; polycystic ovary syndrome (PCOS), celiac disease, psoriasis, inflammatory bowel disease (IBD) and ankylosing spondylitis.

[0048] Examples of non-organ-specific autoimmune disorders are rheumatoid arthritis, multiple sclerosis, systemic lupus and myasthenia gravis.

[0049] Furthermore, the compounds, compositions and methods of the present invention are useful in treating cancer. Leukotriene synthesis has been shown to be associated with different types of cancer including esophageal cancer, brain cancer, pancreatic cancer, colon cancer.

[0050] The terms "methods of treating or preventing" mean amelioration, prevention or relief from the symptoms and/or effects associated with lipid disorders. The term "preventing" as used herein refers to administering a medicament beforehand to forestall or obtund an acute episode. The person of ordinary skill in the medical art (to which the present method claims are directed) recognizes that the term "prevent" is not an absolute term. In the medical art it is understood to refer to the prophylactic administration of a drug to substantially diminish the likelihood or seriousness of a condition, and this is the sense intended in applicants' claims. As used herein, reference to "treatment" of a patient is intended to include prophylaxis.

[0051] The term "mammal" is used in its dictionary sense. Humans are included in the group of mammals, and humans would be the preferred subjects of the methods of.

[0052] While it may be possible for the compounds of the present invention to be administered as the raw chemical, it is preferable to present them as a pharmaceutical composition. According to a further aspect, the present invention provides a pharmaceutical composition comprising at least one compound described *supra*, or a pharmaceutically acceptable salt or solvate thereof, together with one or more pharmaceutically carriers thereof and optionally one or more other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

[0053] The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular,

intravenous and intraarticular), rectal and topical (including dermal, buccal, sublingual and intraocular) administration. The most suitable route may depend upon the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing into association at least one compound of the present invention or a pharmaceutically acceptable salt or solvate thereof ("active ingredient") with the carrier, which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

[0054] Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder (including micronized and nanoparticulate powders) or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

[0055] A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, lubricating, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide sustained, delayed or controlled release of the active ingredient therein.

[0056] The pharmaceutical compositions may include a "pharmaceutically acceptable inert carrier", and this expression is intended to include one or more inert excipients, which include starches, polyols, granulating agents, microcrystalline cellulose, diluents, lubricants, binders, disintegrating agents, and the like. If desired, tablet dosages of the disclosed compositions may be coated by standard aqueous or nonaqueous techniques, "Pharmaceutically acceptable carrier" also encompasses controlled release means.

[0057] Compositions of the present invention may also optionally include other therapeutic ingredients, anti-caking agents, preservatives, sweetening agents, colorants, flavors, desiccants, plasticizers, dyes, and the like. Any such optional ingredient must, of course, be compatible with the compound of the invention to insure the stability of the formulation. The dose range for adult humans is generally from 0.1 μ g to 10 g/day orally. Tablets or other forms of presentation provided in discrete units may conveniently contain an amount of compound of the invention which is effective at such dosage or as a multiple of the same, for instance, units containing 0.1 mg to 500 mg, usually around 5 mg to 200 mg. The precise amount of compound administered to a patient will be the responsibility of the attendant physician. However, the dose employed will depend on a number of factors, including the age and sex of the patient, the precise disorder being treated, and its severity. The frequency of administration will depend on the pharmacodynamics of the individual compound and the formulation of the dosage form., which may be optimized by methods well known in the art (e.g. controlled or extended release tablets, enteric coating etc.).


[0058] Combination therapy can be achieved by administering two or more agents, each of which is formulated and administered separately, or by administering two or more agents in a single formulation. Other combinations are also encompassed by combination therapy. For example, two agents can be formulated together and administered in conjunction with a separate formulation containing a third agent. While the two or more agents in the combination therapy can be administered simultaneously, they need not be. For example, administration of a first agent (or combination of agents) can precede administration of a second agent (or combination of agents) by minutes, hours, days, or weeks. Thus, the two or more agents can be administered within minutes of each other or within any number of hours of each other or within any number of days or weeks of each other. In some cases even longer intervals are possible.

[0059] While in many cases it is desirable that the two or more agents used in a combination therapy be present in within the patient's body at the same time, this need not be so. Combination therapy can also include two or more administrations of one or more of the agents used in the combination. For example, if agent X and agent Y are used in a combination, one could administer them sequentially in any combination one or more times, e.g., in the order X-Y-X, X-X-Y, Y-X-Y, Y-Y X, X-X-Y-Y, etc.

[0060] As LTA4H inhibitors, the compounds of the invention have utility in treating and preventing *inter alia* inflammation. The compounds and compositions can be used advantageously in combination with other agents useful in treating and preventing inflammatory conditions and for treating and preventing atherosclerosis, thrombosis, stroke, acute coronary syndrome, stable angina, peripheral vascular disease, critical leg ischemia, intermittent claudication, abdominal aortic aneurysm and myocardial infarction.


[0061] In general, the compounds of the present invention and may be prepared by the methods illustrated in the schemes as, for example, described below, or by modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. The following specific non-limiting examples are illustrative of the invention.

SCHEME I




i) (a) $(\text{COCl})_2$, CH_2Cl_2 , DMF, rt, 3h; (b) ArNH_2 , Et_3N , CH_2Cl_2 , rt, 16h; ii) Zn , NaOH , $\text{MeOH}/\text{H}_2\text{O}$, 90°C , 40h; iii) POCl_3 , CH_3CN , 60°C , 4h; iv) BBr_3 , CH_2Cl_2 , -78°C -rt, 16h; v) K_2CO_3 , RBr , 90°C , 16h

SCHEME II




i) (a) NaH , DMF, 0°C -rt; (b) ROTs, 80°C , 3h; ii) $\text{Br}(\text{CH}_2)_2\text{Cl}$, methyl ethyl ketone, reflux, 36h; iii) CH_3NH_2 (aq), CH_3CN , reflux, 24h; iv) methyl acrylate, CH_2Cl_2 , rt, 24h; v) (a) NaOH , $\text{H}_2\text{O}/\text{MeOH}$; (b) H^+ ; vi) K_2CO_3 , DMF, rt

SCHEME III




i) RR'NH, CH₃CN, 60°C; ii) NaOH, THF/MeOH/H₂O, 18h; iii) (a) (COCl)₂, CH₂Cl₂, (b) RNH₂, CH₂Cl₂;
iv) (a) NaBH₄, THF; (b) HCl

SCHEME IV




i) (a) (COCl)₂, CH₂Cl₂, DMF, rt, 3h; (b) ArNH₂, Et₃N, CH₂Cl₂, rt, 16 h; ii) Zn, NaOH, MeOH/H₂O, 90°C, 40 h;
iii) BBr₃, CH₂Cl₂, -78°C-rt, 16 h; iv) (a) NaH, DMF, 0°C; (b) ROTs, 80°C, 16h

SCHEME V




SCHEME VI



EXAMPLES

Example 1.

[0062]



Step 1

[0063] N-(4-Methoxy-phenyl)-2-nitro-benzamide: To a suspended mixture of 2-nitrobenzoic acid (10.0 g, 60 mmol) and DMF (0.4 mL, 5 mmol) in methylene chloride (200 mL) was slowly added oxalyl chloride (9.11 g, 71 mmol) at room temperature, and then stirred for 4 h. After the solvent was removed, the acid chloride was redissolved in methylene chloride (200 mL). To this solution was subsequently added a solution of p-ansidine (8.1 g, 0.65 mol) in methylene chloride (50 mL) and triethylamine (12.1 g, 0.120 mol). After stirring at room temperature for 18 h, the reaction was quenched with water, and the organic solvent was removed in vacuo. The product was collected on a filter, washed with 1N HCl, water and hexane. After dried under vacuum, the title compound (16 g, 98%) was obtained.

Step 2

[0064] 2-(4-Methoxy-phenyl)-2H-indazol-3-ol: A mixture of the product from step 1 (16 g, 0.059 mol), NaOH (9.4 g, 0.235 mol) and zinc (23 g, 0.352 mol) in methylene chloride (250 mL) was heated to 80°C for 24 h. After the methanol was removed in vacuo, the aqueous residue was acidified with 10% HCl (aq.) to pH to ~2. The product was collected on a filter, washed with 1N HCl, water and hexane. After dried under vacuum, the title compound (12 g, 86%) was obtained; MS (ESI+) m/z 241 (M+1, 100).

Step 3

[0065] 3-Chloro-2-(4-methoxy-phenyl)-2H-indazole: To a mixture of the product from step 2 (12.0 g, 0.05 mol) in acetonitrile (250 mL) was added POCl₃ (11.5 g, 0.74 mol), and heated to 80 °C for 4 h. After the reaction mixture was cooled to room temperature, water (500 mL) was added. The product was collected on a filter, and washed with water. After dried under vacuum, the title compound (8 g, 67%) was obtained; MS (ESI+) m/z 259 (M+1, 100).


Step 4

[0066] 4-(3-Chloro-indazol-2-yl)-phenol: To a solution of product from step 3 (5.0 g, 19.3 mmol) in CH₂Cl₂ (100 mL) was added boron tribromide (14.53 g, 58 mmol) at -78 °C over 5 min. The reaction mixture was stirred at -78 °C for 1 h, and then at room temperature for 18 h. The reaction was quenched with saturated aqueous NaHCO₃ at 0 °C, the product was collected on a filter, and washed with water. After dried under vacuum, the title compound (4.7 g, 99%) was obtained; MS (ESI+) m/z 245 (M+1, 100).

Step 5

[0067] 3-Chloro-2-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-2H-indazole: A mixture of product from step 4 (0.12 g, 0.47 mmol), 1-(2-chloroethyl)piperidine hydrochloride (0.17 g, 0.95 mmol), K₂CO₃ (0.20 g, 1.42 mmol), in DMF (1 mL) was stirred at 90 °C for 18 h. After the reaction mixture was quenched with water, the reaction mixture was extracted with CH₂Cl₂, washed 10% NaOH, and brine, dried over Na₂SO₄. After concentrated in vacuo, the residue was purified by a column chromatography on silica gel eluting with 25% ethyl acetate/hexane to yield the title compound (0.13 g 77%); ¹H NMR (400 MHz, CDCl₃) δ 7.70 (d, J = 8 Hz, 1H), 7.62 (d, J = 8 Hz, 1H), 7.59 (d, J = 8 Hz, 2H), 7.35 (m, 1H), 7.15 (m, 1H), 7.06 (d, J = 8 Hz, 2H), 4.18 (t, J = 6 Hz, 2H), 2.82 (t, J = 6 Hz, 2H), 2.53 (br, 4H), 1.63 (m, 4H), 1.46 (m, 2H); LC/MS (ESI+) m/z: 93%, 356 (M+1).

Example 2.

[0068]

Step 1

[0069] 3-Chloro-2-[4-(2-chloro-ethoxy)-phenyl]-2H-indazole: A mixture of the product from step 4 of the example 1 (0.5g, 2.0mmol), 1-bromo-2-chloroethane (1.2 g, 8.2 mmol), K₂CO₃ (1.4 g, 10.2 mmol), in MEK (10 mL) was refluxed for 36 h. After the reaction mixture was diluted with water, the resulting aqueous mixture was extracted with ethyl acetate and sequentially washed with water, saturated aq. NaHCO₃, water and brine. The combined organic portions were dried over Na₂SO₄, filtered and concentrated in vacuo to afford the title compound (0.5 g, 88%).

Step 2

[0070] {2-[4-(3-Chloro-indazol-2-yl)-phenoxy]-ethyl}-methyl-amine hydrochloride: A mixture of the product from Step 1 (0.1 g, 0.33 mmol), methylamine (40% in water, 3 mL, 81 mmol) in acetonitrile (2 mL) was heated to 60 °C for 7 h, and then room temperature for 16 h. The mixture was extracted with ethyl acetate. The combined organic portions were acidified with 10% HCl to pH to 1, and stirred at 0 °C for 1 h. The product was collected on a filter, washed with 1N HCl, water and hexane. After dried under vacuum, the title compound (100 mg, 91%) was obtained as hydrochloride salt; MS (APCI) m/z 302 (M+1, 100).

Step 3

[0071] 3-({2-[4-(3-Chloro-indazol-2-yl)-phenoxy]-ethyl}-methyl-amino)-propionic acid methyl ester: A mixture of the product from step 2 (0.1 g, 0.27 mmol), methyl acrylate (0.15 g, 1.7 mmol) in methylene chloride (2 mL) was stirred at room temperature for 24 h. After the inorganic salts were filtered off, the solvent was removed in vacuo. The crude


product was purified by a column chromatography on silica gel eluting with 5% methanol in methylene chloride to afford the ester (100 mg, 60%) MS (APCI) m/z 388 ($M+1$, 100).

Step 4

[0072] 3-({2-[4-(3-Chloro-indazol-2-yl)-phenoxy]-ethyl}-methyl-amino)-propionic acid hydrochloride: A solution of {3-[4-(3-chloro-indazol-2-yl)-phenoxy]-propyl}-methyl-amine hydrochloride (0.1 g, 0.26 mmol), NaOH (50% in water, 0.041 g, 0.52 mmol) in THF/MeOH/H₂O (1:1:1, 6 mL) was stirred at room temperature for 16 h. After the organic solvents were removed, the residue was diluted with water, acidified with 10% HCl to pH to 2, extracted with ethyl acetate and sequentially washed with water, and brine. The combined organic portions were dried over Na₂SO₄, filtered and concentrated in vacuo to afford the title compound (0.07 g, 88%) after it triturated with ether; ¹H NMR (400 MHz, DMSO) δ 7.68-7.59 (m, d, J = 8.8 Hz, 4H), 7.42-7.38 (m, 1H), 7.24-7.17 (m, d, J = 8.8 Hz, 3H), 4.17 (t, J = 6 Hz, 2H), 3.27 (m, 4H), 2.77 (m, s, 5H), 2.18 (m, 2H); LC/MS (ESI+) m/z : 93%.

Example 3.

[0073]



Step 1


[0074] 4-({2-[4-(3-Chloro-indazol-2-yl)-phenoxy]-ethyl}-methyl-amino)butyric acid ethyl ester: A mixture of prepared from the product from step 3 of the Example 4 (0.3 g, 0.89 mmol), ethyl 4-bromobutyrate (0.19 g, 0.98 mmol) and K₂CO₃ (powder) (0.49 g, 3.55 mmol) in DMF was stirred at room temperature for 18 h. After the DMF was removed, the residue was partitioned with methylene chloride and water, washed with brine. The combined organic phases were dried over Na₂SO₄ and concentrated. The crude material was purified by a chromatography on silica gel eluting with 10% ethyl acetate in hexane to provide the ester compound (150 mg, 45%).

Step 2

[0075] 4-({2-[4-(3-Chloro-indazol-2-yl)-phenoxy]-ethyl}-methyl-amino)butyric acid hydrochloride salt: A solution of this ester from step 1 (0.1 g, 0.23 mmol) and NaOH (20 mg, 0.47 mmol) in EtOH/water (1:1.5 mL) was stirred at room temperature for 16 h. After the ethanol was removed, the aqueous solution was acidified with 10% HCl to pH = 2, extracted with ethyl acetate. The combined organic phases were dried over Na₂SO₄ and concentrated. The crude material was triturated with ether to yield the title compound (48 mg, 50%); ¹H NMR (400 MHz, DMSO) δ 9.01 (s, 1H), 8.01 (d, J = 9.2 Hz, 2H), 7.76 (d, J = 8 Hz, 1H), 7.68 (d, J = 8 Hz, 1H), 7.30 (m, 1H), 7.15 (d, J = 8 Hz, 2H), 7.08 (m, 1H), 4.15 (t, J = 12, 6 Hz, 2H), 3.27 (m, 4H), 2.84 (m, 2H), 2.76 (s, 3H), 2.18 (m, 2H); LC/MS (ESI+) m/z : 98%.

Example 4.

[0076]




Step 1

[0077] 3-Chloro-2-[4-(3-chloro-propoxy)-phenyl]-2H-indazole: The title compound was prepared from 3-chloro-2-(4-methoxy-phenyl)-2H-indazole (1.2 g, 4.9 mmol) and 1-bromo-3-chloropropane (3.1 g, 19.6 mmol), K_2CO_3 (2.8 g, 20.4 mmol), in methyl ethyl ketone (50 mL) was refluxed for 36 h. After the reaction mixture was diluted with water, the resulting aqueous mixture was extracted with ethyl acetate and sequentially washed with water, saturated aq. $NaHCO_3$, water and brine. The combined organic portions were dried over Na_2SO_4 , filtered and concentrated in *vacuo* to afford the title compound (1.2g, 80%). MS (APCI) m/z: 321 (M+1, 100).

Step 2

[0078] {3-[4-(3-Chloro-indazol-2-yl)-phenoxy]-propyl}-methyl-amine hydrochloride: The title compound was prepared from product from step 1 (0.6 g, 1.9 mmol) and methyl amine (40% in water, 16.2 mL, 0.47 mol) in acetonitrile (6 mL) was heated to 60 °C for 7 h, and then room temperature for 16 h. The mixture was extracted with ethyl acetate. The combined organic portions were acidified with 10% HCl to pH to 1, and stirred at 0 °C for 1 h. The product was collected on a filter, washed with 1N HCl, water and hexane. After drying under vacuum, the title compound (0.5 g, 79%) as a hydrochloride salt. 1H NMR (400 MHz, CD_3OD) δ 7.66-7.60 (m, d, J = 9.2 Hz, 4H), 7.43-7.39 (m, 1H), 7.23-7.18 (m, d, J = 9.2 Hz, 3H), 4.24 (t, J = 6 Hz, 2H), 3.27 (t, J = 6 Hz, 2H), 2.77 (s, 3H), 2.24 (m, 2H); MS (APCI) m/z 316 (M+1, 100); LC/MS (APCI) m/z: 99%.

Example 5.

[0079]


Step 1

[0080] 3-((3-(4-(3-Chloro-indazol-2-yl)-phenoxy)-propyl)-methyl-amino)-propionic acid methyl ester: The title compound was prepared from product from step 2 of the Example 6 (0.2 g, 0.5 mmol) and methyl acrylate (0.29 g, 3.4 mmol) using the procedure of Step 3 of the Example 6 to provide the title compound (0.2 g, 60%) MS (ESI+) m/z 302 (M+1, 100).

Step 2

[0081] 3-((3-(4-(3-Chloro-indazol-2-yl)-phenoxy)-propyl)-methyl-amino)-propionic acid hydrochloride salt The title compound was prepared from product from step 1 (0.2 g, 0.5 mmol) and NaOH (50% in water, 0.080 g, 1.0 mmol) using the procedure of Step 6 of the Example 4 to provide the title compound (0.2 g, 60%). 1H NMR (400 MHz, DMSO) 7.68-7.59 (m, d, J = 8.8 Hz, 4H), 7.42-7.38 (m, 1H), 7.24-7.17 (m, d, J = 8.8 Hz, 3H), 4.17 (t, J = 6 Hz, 2H), 3.27 (m, 4H), 2.77 (m, s, 5H), 2.18 (m, 2H); LC/MS (ESI+) m/z: 99%.

Example 6.

[0082]


Step 1

[0083] 4-({3-[4-(3-Chloro-indazol-2-yl)-phenoxy]-propyl}-methyl-amino)butyric acid ethyl ester The title compound was prepared from product from step 1 of the Example 4 (0.1 g, 0.5 mmol) and ethyl 4-bromobutyrate (0.061 g, 0.31 mmol) using the procedure of step 1 of the Example 3 to provide the title compound (0.2 g, 60%).

Step 2


[0084] 4-({3-[4-(3-Chloro-indazol-2-yl)-phenoxy]-propyl}-methyl-amino)butyric acid Hydrochloride: The title compound was prepared from the product of step 1 (0.1 g, 0.28 mmol) and NaOH (50% in water, 0.080 g, 1.0 mmol) using the procedure of step 2 of the Example 3 to provide the title compound (30 mg, 30%) as a hydrochloride salt. ¹H NMR (400 MHz, DMSO) δ 7.68-7.62 (m, 4H), 7.43-7.38 (m, 1H), 7.28 (d, *J* = 8.8 Hz, 2H), 7.17 (m, 1H), 7.05 (d, *J* = 8.8 Hz, 2H), 4.23 (t, *J* = 5 Hz, 2H), 3.33 (br t, *J* = 8.8 Hz, 2H), 3.17 (br t, *J* = 8 Hz, 2H), 2.86 (s, 3H), 2.53 (t, *J* = 6.4 Hz, 2H), 2.45 (m, 2H), 2.15 (m, 2H) LC/MS (APCI) *m/z*: 98%.

Example 7.

[0085]

3-({3-[4-(3-chloro-1H-indazol-2-yl)-phenoxy]-propyl}-methyl-amino)-propionic acid: To the product from step 2 of the Example 6 (0.1 g, 0.26 mmol) in THF (2 mL) was added *n*-butyllithium (2.5 M in hexane, 0.33 mL, 0.82 mmol) at -78 °C, and continued to at -78 to -20 °C for an additional one hour. After the reaction was quenched with water, it was acidified with 10% HCl to pH to 1, extracted with 2% MeOH in methylene chloride, dried over Na₂SO₄, and concentrated in vacuo to yield the title compound (85 mg, 85%). ¹H NMR (400 MHz, DMSO) δ 9.01 (s, 1H), 8.01 (d, *J* = 9.2 Hz, 2H), 7.76 (d, *J* = 8 Hz, 1H), 7.68 (d, *J* = 8 Hz, 1H), 7.30 (m, 1H), 7.15 (d, *J* = 8 Hz, 2H), 7.08 (m, 1H), 4.15 (t, *J* = 6 Hz, 2H), 3.27 (m, 4H), 2.84 (m, 2H), 2.76 (s, 3H), 2.18 (m, 2H); LC/MS (APCI) *m/z*: 91%.

Example 8.

[0086]

Step 1

[0087] Methyl 1-({3-[4-(3-chloro-indazol-2-yl)-phenoxy]-propyl}-piperidine-4-carboxylate: The title compound was prepared from 3-chloro-2-[4-(3-chloro-propoxy)-phenyl]-2H-indazole (0.5 g, 1.6 mmol) and ethyl isonipecotatate (0.37 g, 2.3 mmol) using the procedure of step 4 of the Example 4 to provide the title compound (0.56 g, 75%) as a hydrochloride salt.


Step 2

[0088] 1-({3-[4-(3-Chloro-indazol-2-yl)-phenoxy]-propyl}-piperidine-4-carboxylic acid hydrochloride: A solution of the ester (0.45 g, 0.26 mmol), NaOH (50% in water, 0.041 g, 0.52 mmol) in THF/EtOH/H₂O (1:1:1, 15 mL) was stirred at room temperature for 16 h. After the organic solvents were removed, the residue was diluted with water, acidified with 10% HCl to pH to 2, and stirred at 0 °C for 1 h. The product was collected on a filter, and dried in vacuo to afford the

title compound (0.33 g, 78%) as a hydrochloride salt. ^1H NMR (400 MHz, CD_3OD) δ 7.65(m, 1H), 7.59 (d, J = 8.8 Hz, 4H), 7.42-7.38 (m, 1H), 7.24-7.17 (m, d, J = 8.8 Hz, 3H), 4.23 (t, J = 6 Hz, 2H), 3.71 (br, 1H), 3.38 (m, 2H), 3.08 (br, 1H), 2.33 (m, 3H), 1.91(br, 2H), LC/MS (ESI+) m/z : 98%.

Example 9.


[0089]



1-{3-[4-(3-Chloro-indazol-2-yl)-phenoxy]-propyl}-piperidine-4-carboxylic acid methylamide: To a solution of 1-{3-[4-(3-chloro-indazol-2-yl)-phenoxy]-propyl}-piperidine-4-carboxylic acid hydrochloride (0.11 g, 0.25 mmol) in methylene chloride (11 mL) was added oxalyl chloride (0.12 g, 0.98 mmol) at room temperature, and stirred at room temperature for 3h. After the solvent was removed, the acid chloride was dissolved in THF and methylamine (40% in H_2O , 3.8 g, 49 mmol) was added in one portion. The resulting reaction was stirred at room temperature for 3 h, and the THF was removed. The product was collected on a filter and air-dry for 8 h to provide the title compound (0.1 g, 99%). ^1H NMR (400 MHz, CDCl_3) δ 7.71 (d, J = 10.8 Hz, 1H), 7.62-7.56 (m, 3H), 7.37-7.33 (m, 1H), 7.17-7.14 (m, 2H), 7.04 (d, J = 12Hz, 2H), 5.48 (br, 1H), 4.10 (t, J = 6 Hz, 2H), 3.00 (br d, J = 12Hz, 1H), 2.82 (d, J = 4.8Hz, 3H), 2.52 (t, J = 6 Hz, 2H), 2.10-1.73 (m, 9H), LC/MS (APCI) m/z : 99%.

Example 10.


[0090]



1-{3-[4-(3-Chloro-indazol-2-yl)-phenoxy]-propyl}-piperidine-4-carboxylic acid amide: 1-{3-[4-(3-Chloro-indazol-2-yl)-phenoxy]-propyl}-piperidine-4-carboxylic acid hydrochloride (0.1g, 0.25mmol) and ammonia (28% in H_2O , 3.0g, 49mmol) using the procedure of step 3 of the Example 11 to provide the title compound (0.07g, 70%). ^1H NMR (400 MHz, CDCl_3) δ 7.71(d, J = 8.8 Hz, 1H), 7.59-7.56 (m, 3H), 7.37-7.33 (m, 1H), 7.17-7.14 (m, 2H), 7.04 (d, J = 8.8Hz, 2H), 5.46 (br, 1H), 5.33 (br, 1H), 4.10 (t, J = 6 Hz, 2H), 3.01 (br d, J = 12Hz, 1H), 2.54 (t, J = 6 Hz, 2H), 2.10-1.73 (m, 9H), LC/MS (APCI) m/z : 97%.

Example 11.

[0091]




(1-{3-[4-(3-Chloro-indazol-2-yl)-phenoxy]-propyl}piperidin-4-yl)-methanol: To a solution of methyl 1- {3-[4-(3-chloro-in-

dazol-2-yl)-phenoxy]-propyl)-piperidine-4-carboxylate (0.1 g, 0.21 mmol) in THF (10 mL) was added NaBH₄ (23 mg, 0.63 mmol) at 0 °C in portions. The reaction mixture was stirred at room temperature for 4 h. After it was quenched with water, the mixture was extracted with ethyl acetate, washed with brine and water, dried over Na₂SO₄ and concentrated in vacuo. The crude product was purified by a column chromatography on silica gel eluting with 20% ethyl acetate in hexane to the title compound (0.43 g 50%). ¹H NMR (400 MHz, CDC13) δ 7.71 (d, *J* = 8.8 Hz, 1H), 7.59-7.56 (m, 3H), 7.37-7.33 (m, 1H), 7.17-7.14 (m, 2H), 7.04 (d, *J* = 8.8Hz, 2H), 4.10 (t, *J* = 6 Hz, 2H), 3.22 (m, 2H), 3.01 (br d, *J* = 12Hz, 1H), 2.54 (t, *J* = 6 Hz, 2H), 2.10-1.73 (m, 9H).

Example 12.


[0092]



3-Chloro-2-[4-((R)-1-pyrrolidin-2-ylmethoxy)-phenyl]-2H-indazole: To a mixture of NaH (60% in mineral oil, 13 mg, 0.32 mmol) in DMF (1 mL) was added a solution of 4-(3-Chloro-indazol-2-yl)-phenol of the Example 1 (0.06 g, 0.25 mmol) in DMF (1 mL) at 0 °C. The resulting slurry was stirred at 0 °C for 30 minutes and at room temperature for 30 minutes before a solution of (R)-2-(toluene-4-sulfonyloxymethyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (0.174 g, 0.49 mmol) in DMF (1 mL) was added. The mixture was stirred at 80 °C for 4 h. The reaction mixture was poured over ice and then concentrated under reduced pressure. The crude residue was extracted into ethyl acetate and sequentially washed with water, saturated aq. NaHCO₃, water and brine. The combined organic portions were dried over Na₂SO₄, filtered and concentrated in vacuo. The crude product was purified by a column chromatography on silica gel eluting with 10% ethyl acetate in hexane to afford the Boc-protected compound, which was dissolved in dioxane. To this solution was added 4M HCl in dioxane (0.5 mL, 2.2 mmol), and stirred at room temperature for 4 h. After the solvent was removed, the crude material was triturated with ether to afford the title compound (20 mg, 20%); ¹H NMR (400 MHz, CDC13) δ 7.67 (d, *J* = 8.8 Hz, 1H), 7.57 (d, *J* = 9.2 Hz, 2H), 7.60-7.58 (m, 1H), 7.35-7.31 (m, 1H), 7.15 (d, *J* = 9.2 Hz, 1H), 7.15-7.12 (m, 1H), 4.45 (m, 1H), 4.34 (m, 1H), 4.01 (m, 1H), 3.41 (m, 2H), 2.25-1.98 (m, 4H); LC/MS (ESI+) *m/z*: 95%; 328 (M+1, 100).

Example 13.


[0093]



3-Chloro-2-[4-((S)-1-pyrrolidin-2-ylmethoxy)-phenyl]-2H-indazole hydrochloride: The title compound was prepared from 4-(3-Chloro-indazol-2-yl)-phenol (2.7 g, 11.0 mmol) using the procedure of the Example 12 to provide the title compound (1.2g, 80%); ¹H NMR (400 MHz, CDCl₃) δ 10.23 (br, 1H), 9.57 (br, 1H), 7.71 (d, *J* = 8.8 Hz, 1H), 7.62 (m, 3H), 7.53 (m, 1H), 7.25 (m, 3H), 4.47 (m, 2H), 4.09 (m, 1H), 3.45 (m, 2H), 2.25-1.98 (m, 4H); LC/MS (ESI+) *m/z*: 95%; 328(M+1).

Example 14.

[0094]



Step 1

[0095] 4-Chloro-N-(4-methoxy-phenyl)-2-nitro-benzamide: The title compound was prepared from 4-chloro-2-nitro-benzoic acid (6.0 g, 30 mmol) and p-anisidine (4.0 g, 33 mmol) using the procedure of step 1 of the Example 1 to provide the title compound (8.5g, 90%).

Step 2

[0096] 6-Chloro-2-(4-methoxy-phenyl)-2H-indazol-3-ol: The title compound was prepared from the product from step 1 (5.0 g, 16.3 mmol) using the procedure of step 2 of the Example 1 to provide the title compound (3.7g, 82%), MS (ESI-) m/z 273 (M-1, 100).


Step 3

[0097] 6-Chloro-2-(4-hydroxy-phenyl)-2H-indazol-3-ol: The title compound was prepared from the product of step 2 (0.5 g, 1.8 mmol) using The procedure of step 4 of the Example 1 to provide the title compound (0.3 g, 64%). MS (ESI+) m/z 261 (M+1, 100).

Step 4

[0098] 2-[4-((R)-1-Pyrrolidin-2-ylmethoxy)-phenyl]-2H-indazol-3-ol hydrochloride: The title compound was prepared from the product of step 3 (0.1 g, 0.38 mmol) using The procedure of the Example 2 to provide the title compound (0.07 g, 58%). ¹H NMR (400 MHz, CDCl₃) δ 7.76 (br d, J = 8 Hz, 1H), 7.47 (d, J = 8.8 Hz, 1H), 7.45 (br, 2H), 7.04 (d, J = 2 Hz, 1H), 6.94 (d, J = 9.2 Hz, 2H), 6.88 (dd, J = 8.4, 2 Hz, 1H), 4.45 (m, 1H), 4.34 (m, 1H), 4.01 (m, 1H), 3.41 (m, 2H), 2.25-1.98 (m, 4H), LC/MS (ESI+) m/z 95%.

Example 15.

[0099]

Step 1

[0100] {(R)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]pyrrolidin-1-yl}-acetic acid t-butyl ester: The title compound was prepared from 3-chloro-2-[4-((R)-1-pyrrolidin-2-ylmethoxy)-phenyl]-2H-indazole (0.4 g, 1.1 mmol) and t-butyl bromoacetate (0.27 g, 1.4 mmol) using the procedure of step 7 of the Example 5 to provide the title compound (0.4 g, 80%).


Step 2

[0101] {(R)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]pyrrolidin-1-yl}-acetic acid Hydrochloride: A mixture of the product of step 1 (0.4 g, 0.91 mmol) and HCl (4 M in dioxane, 2.3 mL, 9.1 mmol) in dioxane (5 mL) was stirred at room temperature for 16 h. After the solvent was removed, the residue was triturated with ether to provide the title compound (280 mg, 62%) as a hydrochloride salt. ¹H NMR (400 MHz, DMSO) δ 7.71-7.66 (m, 4H), 7.40 (m, 1H), 7.22 (m, 3H),

4.4.50-4.27 (m, 4H), 4.04 (br, 1H), 3.70 (br, 2H), 2.28 (m, 1H), 2.04 (m, 2H), 1.98 (m, 1H), LC/MS (APCI) m/z: 94%.

Example 16.

[0102]



Step 1


[0103] 3-((R)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]methyl)-pyrrolidin-1-yl-propionic acid methyl ester: The title compound was prepared from product from 3-chloro-2-[4-((R)-1-pyrrolidin-2-ylmethoxy)-phenyl]-2H-indazole (0.4 g, 1.1 mmol) and methyl acrylate (0.25 g, 3.4 mmol) using the procedure of step 5 of the Example 4 to provide the title compound (0.37 g, 79%).

Step 2

[0104] 3-((R)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]methyl)-pyrrolidin-1-yl-propionic acid Hydrochloride: A mixture of the product of step 1 (0.37 g, 0.89 mmol), water (1 mL) and HCl (4 M in dioxane, 2.3 mL, 9.1 mmol) in dioxane (5 mL) was stirred at room temperature for 16 h. After the solvent was removed, the residue was triturated with ether to provide the title compound (200 mg, 44%) as a hydrochloride salt. ¹H NMR (400 MHz, CDCl₃) δ 7.72-7.66 (m, 4H), 7.41 (m, 1H), 7.23 (m, 3H), 4.44 (br, 2H), 4.03 (br, 1H), 3.71 (br, 2H), 3.35 (br, 1H), 3.20 (br, 1H), 2.87 (t, J = 8 Hz, 2H), 2.27 (m, 1H), 2.09-1.98 (br, 3H), LC/MS (APCI) m/z: 85%.

Example 17.

[0105]



Step 1

[0106] 4-((R)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]methyl)-pyrrolidin-1-yl-butyric acid methyl ester: The title compound was prepared from product from 3-chloro-2-[4-((R)-1-pyrrolidin-2-ylmethoxy)-phenyl]-2H-indazole (0.4 g, 1.1 mmol) and methyl 4-bromobutyrate (0.25 g, 1.4 mmol) using the procedure of step 7 of the Example 5 to provide the title compound (0.38 g, 79%).


Step 2

[0107] 4-((R)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]methyl)-pyrrolidin-1-yl-butyric acid Hydrochloride: The title compound was prepared from compound the product of step 1 (0.38 g, 0.89 mmol) using the procedure step 2 of Example 16 to provide the title compound (150 mg, 40%). ¹H NMR (400 MHz, CDCl₃) δ 7.71 (m, 4H), 7.41 (m, 1H), 7.24 (m, 3H), 4.45 (m, 2H), 3.98 (br, 1H), 3.66 (m, 1H), 3.52 (m, 1H), 3.17 (m, 2H), 2.40 (t, J = 7.2 Hz, 2H), 2.23 (m, 2H), 2.08-1.84

(br, 4H), LC/MS (APCI) m/z: 85%.

Example 18.

[0108]



Step 1


[0109] ((S)-2-[4-(3-Chloro-indazol-2-yl)phenoxy]methyl-pyrrolidin-1-yl)-acetic acid methyl ester: The title compound was prepared from product from 3-chloro-2-[4-((S)-1-pyrrolidin-2-ylmethoxy)-phenyl]-2H-indazole (0.33 g, 0.94 mmol) and t-butyl bromoacetate (0.22 g, 1.1 mmol) using the procedure of step 1 of the Example 15 to provide the title compound (0.258 g, 63%).

Step 2

[0110] ((S)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]methyl-pyrrolidin-1-yl) -acetic acid Hydrochloride: The title compound was prepared from compound the product of step 1 (0.25 g, 0.6 mmol) using the procedure Step 2 of Example 15 to provide the title compound (110 mg, 30%). ¹H NMR (400 MHz, CDCl₃) δ 7.71 (d, J= 8.8 Hz, 1H), 7.58 (d, J= 8.8 Hz, 1H), 7.53 (d, J= 8 Hz, 1H), 7.35 (m, 1H), 7.15 (m, 3H), 4.87 (br, 1H), 4.37 (br, 3H), 4.19 (br, 1H), 4.01 (br, 1H), 3.45 (br, 1H), 2.25-1.98 (br, 4H), LC/MS (APCI) m/z: 99%.

Example 19.

[0111]



Step 1


[0112] 4-((S)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]methyl-pyrrolidin-1-yl)-butyric acid methyl ester: The title compound was prepared from product from 3-chloro-2-[4-((S)-1-pyrrolidin-2-ylmethoxy)-phenyl]-2H-indazole (0.33 g, 0.94 mmol) and methyl 4-bromobutyrate (0.25 g, 1.4 mmol) using the procedure of Step 1 of the Example 17 to provide the title compound (0.30 g, 75%).

Step 2

[0113] 4-((S)-2-[4-(3-Chloro-indazol-2-yl)-phenoxy]methyl-pyrrolidin-1-yl)-butyric acid Hydrochloride: The title compound was prepared from compound the product of step 1 (0.30 g, 0.68 mmol) using the procedure step 2 of Example 17 to provide the title compound (200 mg, 52%). ¹H NMR (400 MHz, CDCl₃) δ 7.71 (m, 4H), 7.41 (m, 1H), 7.24 (m, 3H), 4.45 (m, 2H), 3.95 (br, 3H), 3.64 (m, 1H), 3.42 (m, 1H), 3.16 (m, 1H), 2.94 (m, 2H), 2.27 (m, 1H), 2.08-1.84 (br, 3H), LC/MS (APCI) m/z: 99%.

Example 20

[0114]



Step 1

[0115] 1-(4-Methoxy-phenyl)-1H-indazole: A mixture of 1H-Indazole (0.50 g, 4.23 mmol), 4-methoxyphenylboronic acid (1.30 g, 8.5 mmol), triethylamine (0.86 g, 8.5 mmol) and cupric acetate ((0.77 g, 4.23 mmol) in CH_2Cl_2 (50 mL) was stirred at room temperature for 48 h. The reaction mixture was diluted with water and passed through a pad of Celite. The filtrate was extracted with EtOAc, washed with brine, and dried over Na_2SO_4 . After being concentrated, the crude material was purified by a chromatography on silica gel to yield the title compound (0.32 g, 30%).

Step 2


[0116] 4-Indazol-1-yl-phenol: The title compound was prepared from the product of the step 1 (0.15 g, 0.67 mmol) and BBr_3 (0.50 g, 2.0 mmol) in CH_2Cl_2 (3 mL) by using the step 4 procedure of Example 1 with 90% yield (0.13 g).

Step 3

[0117] 1-[4-((R)-1-Pyrrolidin-2-ylmethoxy)-phenyl]-1H-indazole: The title compound was prepared from the product of the step 2 (0.12 g, 0.62 mmol) and (R)-2-(toluene-4-sulfonyloxymethyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (0.25 g, 0.7 mmol) in DMF (3 mL) by using the step 1 procedure of Example 2 with 30% yield (0.06 g). ^1H NMR (400 MHz, CDCl_3) δ 8.15 (s, 1H), 7.78 (d, $J = 8.4$ Hz, 1H), 7.60 (m, 2H), 7.40 (t, $J = 8$ Hz, 1H), 7.20 (m, 4H), 4.30 (br, 2H), 4.11 (br, 1H), 3.45 (br, 2H), 2.30-1.94 (br, 4H), LC/MS (APCI) m/z : 99%.

Example 21

[0118]




Step 1

[0119] (R)-2-[4-(1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-phenoxy]methylpyrrolidine-1-carboxylic acid tert-butyl ester 1: To a 25 mL vial which contained a suspension of NaH (60% in mineral oil, 30 mg, 0.75 mmol) in DMF (3 mL) was added 2-(4-Hydroxy-phenyl)-isoindole-1,3-dione (105 mg, 0.5 mmol) at 0°C . The mixture was allowed to warm to rt and stir at rt for 30 min then cooled to 0°C . To this reaction mixture was added (R)-2-(toluene-4-sulfonyloxymethyl)-pyrrolidine-1-carboxylic acid tert-butyl ester (173 mg, 0.5 mmol) at 0°C . The resulting mixture was allowed to warm to rt and stir at rt for 30 min and then was heated to 90°C and stirred at 90°C for 16 h. After cooled to rt, the mixture was poured into 150 mL ice-water solution and then was extracted with EtOAc (3x50 mL). The combined organic layers were washed with water (3x50 mL), brine (30 mL) and dried over Na_2SO_4 . After concentrated in vacuo, the residue was purified by a column chromatography on silica gel to yield the title product, (120 mg, 55%)

Step 2


[0120] 2-[4-((R)-1-Pyrrolidin-2-ylmethoxy)-phenyl]-isoindole-1,3-dione hydrochloride 2: To a 20 mL vial which contained a solution of the product from step 1 (40 mg, 0.1 mmol) in dioxane (2 mL) was added HCl (4 N in dioxane, 2 mL) at 0 °C. The mixture was allowed to warm to rt and stir at rt for 16 h. The solvent was removed and the crude was purified by recrystallization from MeOH-ether to yield the title product (30 mg, 90 %); LCMS; 98%, ESI⁺ Calcd: 322.3, found m/z: 323.6 (M+1); ¹HNMR (400 MHz, DMSO-*d*₆): δ 1.73-1.84 (m, 1H), 1.86-2.04 (m, 2H), 2.10-2.20 (m, 1H), 3.18-3.28 (m, 2H), 3.88-3.98 (m, 1H), 4.16-4.21 (m, 1H), 4.33 (dd, *J* = 10.8 Hz, 3.6 Hz, 1H), 7.14 (dd, *J* = 6.8 Hz, 2.0 Hz, 2H), 7.40 (dd, *J* = 6.8 Hz, 2.4 Hz, 2H), 7.89-7.97 (m, 4H).

Example 22

[0121]

2-[4-(2-Piperidin-1-yl-ethoxy)-phenyl]-isoindole-1,3-dione: To a 50 mL vial which contained a solution of 2-(4-Hydroxy-phenyl)-isoindole-1,3-dione (239 mg, 1 mmol) and 1-(2-chloroethyl) piperidine hydrochloride (390 mg, 2 mmol) in DMF (15 mL) was added K₂CO₃ (600 mg, 4.5 mmol) at rt. The reaction mixture was heated to 85 °C and stirred at 85 °C for 72 h. After cooled to rt, the reaction mixture was poured into 50 mL ice-water mixture and then was extracted with EtOAc (3 x 30 mL). The combined organic layers were washed with water (3 x 30 mL), brine (30 mL) and dried over Na₂SO₄. After concentrated in vacuo, the residue was purified by a column chromatography on silica gel to yield the title product, (90 mg, 28%); LCMS; 97%, ESI⁺; Calcd: 350.4; Found m/z: 351.6 (M+1); ¹HNMR (400 MHz, CDCl₃) δ 1.40-1.68 (m, 6H), 2.52 (br, 4H), 2.80 (t, *J* = 6.0 Hz, 2H), 4.15 (t, *J* = 6.0 Hz, 2H), 7.02 (d, *J* = 9.2 Hz, 2H), 7.32 (d, *J* = 9.2 Hz, 2H), 7.77-7.79 (m, 2H), 7.84-7.96 (m, 2H).

Example 23

[0122]

Step 1

[0123] (R)-2-[4-((1,3-Dioxo-1,3-dihydro-isoindol-2-yl)-phenoxy)methyl]-pyrrolidine-1-carboxylic acid tert-butyl ester 3: To a 25 mL vial which contained a suspension of NaH (60% in mineral oil, 30 mg, 0.75 mmol) in DMF (3 mL) was added 2-(4-Hydroxy-phenyl)-isoindole-1,3-dione 1 (105 mg, 0.5 mmol), at 0 °C. The mixture was allowed to warm to rt and stir at rt for 30 min and then cooled to 0 °C. To this reaction mixture was added (R)-2-(toluene-4-sulfonyloxymethyl)-pyrrolidine-1-carboxylic acid tert-butyl ester 2 (173 mg, 0.5 mmol) at 0 °C. The resulting mixture was allowed warm to rt and stir at rt for 30 min and then was heated to 95 °C and stirred at 95 °C for 16 h. After cooled to rt, the mixture was poured onto 20 mL ice-water solution and this solution was allowed to stir at 0 °C for 30 min. The solid which formed was filtered out, dried through air to afford the title product, (120 mg, 55 %).

Step 2

[0124] 2-[4-((R)-1-Methyl-pyrrolidin-2-ylmethoxy)-phenyl]-2,3-dihydro-1H-isoindole 4: To a 25 mL press resistant vial which contained a suspension of LiAlH₄ (100 mg, 3 mmol) in anhydrous THF (15 mL) was added the product from step

1 (120 mg, 0.3 mmol) at -78 °C. The reaction mixture was allowed to warm to rt and stir at rt for 2 h then was heated to 78 °C and stirred at 78 °C for 24 h. After cooled to 5 °C, 1 eq of water was added to the mixture which was followed by addition of 1 eq of 15% NaOH and then 3 eq of water. The solid which formed was filtered out and washed with THF water (2 x 20 mL). The combined organic solvent was removed under vacuo to obtain the crude product which was purified by recrystallization with ether-EtOAc-hexane to afford the title product, (50 mg, 57 %); LCMS; 99%, ESI⁺ Calcd: 308.4 m/z, found: 309.7 m/z (M+1);

¹H NMR (400 MHz, DMSO-d₆); δ 1.65-2.08 (m, 4H), 2.17-2.32 (m, 1H), 2.49 (s, 3H), 2.56-2.61 (m, 1H), 3.09-3.13 (m, 1H), 3.85 (dd, *J* = 8.8 Hz, 6.0 Hz, 1H), 3.98 (dd, *J* = 9.2 Hz, 5.2 Hz, 1H), 4.41 (s, 4H), 6.62 (d, *J* = 9.2 Hz, 2H), 6.93 (d, *J* = 8.8 Hz, 2H), 7.27-7.34 (m, 4H).

[0125] (1) *In vitro* assay testing inhibitory activity against purified recombinant human LTA₄ hydroase:

[0126] A human LTA₄ hydrolase full-length cDNA clone (NM_000895) was purchased from OriGene Technologies (Rockville, MD). The gene was amplified by polymerase chain reaction and transferred via pDONR201 into the bacterial expression vector pDEST17 by recombination (both plasmids from Invitrogen, Carlsbad, CA). The resulting construct was transformed into *Escherichia coli* BL21-AI (Invitrogen), and expression was induced by chemical induction with arabinose. The recombinant enzyme was purified by chromatography on a FPLC system (Amersham Biosciences, Uppsala, Sweden) using immobilized metal affinity chromatography (Ni-NTA Superflow, Qiagen, Hilden, Germany) and anion exchange chromatography (MonoQ HR 10/10, Amersham Biosciences).

[0127] The compounds of the invention were incubated in a series of dilutions with 200 nM of recombinant enzyme in assay buffer (100 mM Tris-HCl, 100 mM NaCl, 5 mg/ml fatty-acid free BSA, 10% DMSO, pH 8.0) for 10 min at room temperature to allow binding between LTA₄ hydrolase and the inhibitors. LTA₄ was prepared by alkaline hydrolysis of LTA₄ methyl ester (Biomol, Plymouth Meeting, PA, or Cayman Chemicals, Ann Arbor, MI). A solution of 10 μg of the ester was dried under a nitrogen stream and redissolved in 60 μl of a solution of 80% acetone and 20% 0.25 M NaOH.

[0128] After incubation for 40 min at room temperature the resulting approximately 500 μM stock of LTA₄ was kept at -80 °C for no more than a few days prior to use.

[0129] Immediately before the assay, LTA₄ was diluted to a concentration of 10 μM in assay buffer (without DMSO) and added to the reaction mixture to a final concentration of 2 μM to initiate the enzyme reaction. After incubation for 120 sec at room temperature, the reaction was stopped by adding 2 volumes of chilled quenching buffer, containing acetonitrile with 1% acetic acid and 225 nM LTB₄-d₄ (Biomol). The samples were then kept at 4 °C overnight to complete protein precipitation and centrifuged for 15 min at 1800 g. LTB₄ formed was measured by LC-MS/MS using LTB₄-d₄ as an internal standard and an external LTB₄ standard (Biomol) as reference. Briefly, the analyte was separated from LTB₄ isomers formed by spontaneous hydrolysis of LTA₄ using isocratic elution (modified protocol from Mueller et al. (1996), J. Biol. Chem. 271, 24345-24348) on a HPLC system (Waters, Milford, MA) and analyzed on a tandem quadrupole mass spectrometer (Waters). MRM transitions followed on 2 channels were 335.2 > 195.3 (LTB₄) and 339.2 > 197.3 (LTB₄-d₄). Based on the amounts of LTB₄ found at each inhibitor concentration, a dose-response curve was fitted to the data and an IC₅₀ value was calculated.

[0130] (1) *In vitro* assay testing inhibitory activity against purified recombinant human LTA₄ hydroase:

[0131] A human LTA₄ hydrolase full-length cDNA clone (NM_000895) was purchased from OriGene Technologies (Rockville, MD). The gene was amplified by polymerase chain reaction and transferred via pDONR201 into the bacterial expression vector pDEST17 by recombination (both plasmids from Invitrogen, Carlsbad, CA). The resulting construct was transformed into *Escherichia coli* BL21-AI (Invitrogen), and expression was induced by chemical induction with arabinose. The recombinant enzyme was purified by chromatography on a FPLC system (Amersham Biosciences, Uppsala, Sweden) using immobilized metal affinity chromatography (Ni-NTA Superflow, Qiagen, Hilden, Germany) and anion exchange chromatography (MonoQ HR 10/10, Amersham Biosciences).

[0132] The compounds of the invention were incubated in a series of dilutions with 200 nM of recombinant enzyme in assay buffer (100 mM Tris-HCl, 100 mM NaCl, 5 mg/ml fatty-acid free BSA, 10% DMSO, pH 8.0) for 10 min at room temperature to allow binding between LTA₄ hydrolase and the inhibitors. LTA₄ was prepared by alkaline hydrolysis of LTA₄ methyl ester (Biomol, Plymouth Meeting, PA, or Cayman Chemicals, Ann Arbor, MI). A solution of 10 μg of the ester was dried under a nitrogen stream and redissolved in 60 μl of a solution of 80% acetone and 20% 0.25 M NaOH. After incubation for 40 min at room temperature the resulting approximately 500 μM stock of LTA₄ was kept at -80 °C for no more than a few days prior to use.

[0133] Immediately before the assay, LTA₄ was diluted to a concentration of 10 μM in assay buffer (without DMSO) and added to the reaction mixture to a final concentration of 2 μM to initiate the enzyme reaction. After incubation for 120 sec at room temperature, the reaction was stopped by adding 2 volumes of chilled quenching buffer, containing acetonitrile with 1% acetic acid and 225 nM LTB₄-d₄ (Biomol). The samples were then kept at 4 °C overnight to complete protein precipitation and centrifuged for 15 min at 1800 g. LTB₄ formed was measured by LC-MS/MS using LTB₄-d₄ as an internal standard and an external LTB₄ standard (Biomol) as reference. Briefly, the analyte was separated from LTB₄ isomers formed by spontaneous hydrolysis of LTA₄ using isocratic elution (modified protocol from Mueller et al. (1996), J. Biol. Chem. 271, 24345-24348) on a HPLC system (Waters, Milford, MA) and analyzed on a tandem quadrupole mass

spectrometer (Waters). MRM transitions followed on 2 channels were 335.2 > 195.3 (LTB₄) and 339.2 > 197.3 (LTB₄-d₄). Based on the amounts of LTB₄ found at each inhibitor concentration, a dose-response curve was fitted to the data and an IC₅₀ value was calculated.


[0134] (2) *Ex vivo* assay testing inhibitory activity in human whole blood after stimulation with calcium ionophor:

[0135] Human blood was collected in heparin-containing Vacutainer tubes. For each sample, 200 μ l of blood were dispensed into a pre-warmed plate and 188 μ l of RPMI-1640 medium (Invitrogen) containing 20 μ g/ml Indomethacin (Sigma, St. Louis, MO) were added. Then 4 μ l of a series of compound dilutions (in DMSO) were added, followed by a 15 min incubation at 37 °C with gentle shaking. After that, blood samples were stimulated by adding Ionomycin (Calbiochem) to a final concentration of 20 μ M.

[0136] After another incubation at 37 °C for 30 min, samples were centrifuged for 5 min at 1800 g and 4 °C. Supernatants were taken and LTB₄ concentrations were determined using a commercially available enzyme-linked immunoassay (R&D Systems, Minneapolis, MN) according to the manufacturer's instructions. Results obtained for different concentrations of hydrolase inhibitor were then used to fit a dose-response curve and calculate an IC₅₀ value.


[0137]

Table I

					
Example	m	n	X	R2	IC50 (μ M)(hLTA4H)
2	2	2	Cl	CO ₂ H	A
3	3	2	Cl	CO ₂ H	A
4	0	3	Cl	H	A
5	2	3	Cl	CO ₂ H	A
6	3	3	Cl	CO ₂ H	A
7	2	3	H	CO ₂ H	ND
A = < 5 μ M; ND = Not Determined					

[0138]

Table II


			
Example	n	R	IC50 (μ M)(hLTA4H)
1	2	H	A
8	3	CO ₂ H	A
9	3	CONHCH ₃	A
10	3	CONH ₂	A

(continued)

Example	n	R	IC50 (μ M)(hLTA4H)
11	3	CH ₂ OH	A
A = < 5 μ M			

[0139]

Table III

					
Example	X	Y	*	R	IC50 (μ M)(hLTA4H)
12	H	Cl	R	H	A
13	H	Cl	S	(CH ₂) ₃ CO ₂ H	A
14	Cl	OH	R	H	ND
15	H	Cl	R	CH ₂ CO ₂ H	A
16	H	Cl	R	(CH ₂) ₂ CO ₂ H	A
17	H	Cl	R	(CH ₂) ₃ CO ₂ H	A
18	H	Cl	S	H	A
19	H	Cl	S	(CH ₂) ₃ CO ₂ H	A
A = < 5 μ M; ND = Not Determined					






[0140]

Table IV

Example	hLTA4H Enzyme IC 50	Example	hLTA4H Enzyme IC 50
1	A	10	A
2	A	11	A
3	A	12	A
4	A	13	A
5	A	14	—
6	A	15	A
7	—	16	A
8	A	17	A
9	A	18	A
		19	A
A = < 5 μ M			


[0141]

Table V


Compound	hLTA4H Enzyme IC ₅₀ (μM)
	A
	A
	B
	B
	C
	C
A < 5 μM B = 5-20 μM C > 20 μM.	

Claims

1. A compound of formula:



or



wherein

ring (a) is chosen from bicyclic heterocyclyl and bicyclic heterocyclyl substituted with from one to three substituents independently selected from the group consisting of halogen, hydroxyl, alkyl groups of from 1 to 6 carbon atoms, acyl groups containing one to four carbon atoms, alkoxy groups containing one to four carbon atoms, fluoroalkyl groups of from 1 to 6 carbon atoms, fluoroalkoxy groups containing one to four carbon atoms, formyl, cyano, phenyl, heteroaryl, nitro and oxo;

Q is O;

R¹ is chosen from H and alkyl groups of from 1 to 6 carbon atoms;

n is zero or an integer from 1-4;

p is an integer from 1-4;

HET is a 4-7-membered saturated nitrogenous heterocycle; and

taken together ZW is H or

Z is $(CH_2)_{0-10}$ in which one or two (CH_2) may optionally be replaced by a C_3-C_6 carbocycle, a C_3-C_6 heterocycle, $-O-$, $-NR^{10}-$, $-SO-$, $-S(O)_2-$, $-C(=O)-$ or $-C(=O)(NH)-$, provided that said $O-$, $-NR^{10}-$, $-SO-$, $-S(O)_2-$, $-C(=O)-$ or $-C(=O)(NH)-$ are not at the point of attachment to nitrogen and are separated by at least one $-(CH_2)_2-$;


R¹⁰ is selected separately in each occurrence from the group consisting of H and alkyl groups of from 1 to 6 carbon atoms;

W is selected from acyl, hydroxyl, carboxyl, amino, carboxamido, sulfonamide, aminoacyl, $-COOalkyl$, $-CHO$, $-C(O)fluoroalkyl$, $-C(O)CH_2C(O)Oalkyl$, $-C(O)CH_2C(O)Ofuoroalkyl$, $-SH$, $-C(O)NH(OH)$, $-C(O)N(OH)R^4$, $-N(OH)C(O)OH$, $-N(OH)C(O)R^4$, heterocyclyl, substituted aryl, and substituted heterocyclyl; and


R⁴ is selected from the group consisting of H and alkyl groups of from 1 to 6 carbon atoms.

2. A compound according to claim 1, wherein ring (a) is chosen from an indazole, an indole and an isoindole ring.

3. A compound according to claim 2 of formula:



or



wherein

R² and R³ each are independently selected from the group consisting of H, halogen, hydroxyl, alkyl groups of from 1 to 6 carbon atoms, acyl groups containing one to four carbon atoms, alkoxy groups containing one to four carbon atoms, fluoroalkyl groups of from 1 to 6 carbon atoms, fluoroalkoxy groups containing one to four carbon atoms, formyl, cyano, phenyl, heteroaryl, and nitro;

n is 1, 2 or 3; and

p is 2 or 3.


4. A compound according to claim 1, wherein

HET is chosen from piperidine and pyrrolidine;

Z is (CH₂)₀₋₁₀; in which one or two (CH₂) may optionally be replaced by -O-, -NR¹⁰-, -SO-, -S(O)₂-, -C(=O)- or -C=O (NH), and

W is selected from the group consisting of hydrogen, acyl, hydroxy, carboxy, amino, carboxamido and aminoacyl.

5. A compound according to claim 4 of formula



wherein


R² is chosen from H, hydroxyl, and halogen;

R³ is chosen from H and halogen;

Z is chosen from (CH₂)₀₋₃ and

W is selected from the group consisting of hydrogen, acyl, hydroxyl, carboxyl, amino, carboxamido, aminoacyl and COOR⁴.

6. A compound according to claim 4 of formula



wherein


R² is chosen from H, hydroxyl, and halogen;

R³ is chosen from H and halogen;


Z is (CH₂)₀₋₃; and

W is selected from the group consisting of hydrogen, hydroxyl, carboxyl and carboxamido.

7. A compound according to claim 2 of formula:



or



wherein

R² is selected from the group consisting of H, alkyl groups of from 1 to 6 carbon atoms, fluoroalkyl groups of from 1 to 6 carbon atoms, phenyl and oxo;

R³ is selected from the group consisting of H, halogen, alkyl groups of from 1 to 6 carbon atoms, acyl groups containing one to four carbon atoms, alkoxy groups containing one to four carbon atoms, fluoroalkyl groups of from 1 to 6 carbon atoms, fluoroalkoxy groups containing one to four carbon atoms, formyl, cyano, phenyl, heteroaryl and nitro;

n is from 1-3; and

p is 2 or 3.


8. A compound according to claim 7, wherein

HET is chosen from piperidine and pyrrolidine;

Z is (CH₂)₀₋₁₀; in which one or two (CH₂) may optionally be replaced by -O-, -NR¹⁰-, -SO-, -S(O)₂-, -C(=O)- or -C=O (NH), and

W is selected from the group consisting of hydrogen, acyl, hydroxyl, carboxyl, amino, carboxamido and aminoacyl.

9. A compound according to claim 2 of formula:



wherein

Y is N or CR⁵;

R⁷ is chosen from H and alkyl groups of from 1 to 6 carbon atoms; and

R⁵ is chosen from H and alkyl groups of from 1 to 6 carbon atoms.

10. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one compound according to any of claims 1-9.

11. A compound according to any of claims 1-9 for use in inhibiting leukotriene A4 hydrolase where the the LTA4H enzyme is contacted with a therapeutically effective amount of a compound according to any of claims 1-9.


12. A compound according to any of claims 1-9 for use in treating a disorder associated with leukotriene A4 hydrolase where a therapeutically effective amount of a compound or a salt, hydrate or ester thereof according to any of claims 1-9 is administered to a mammal.

13. The compound according to claim 12 wherein said disorder is associated with inflammation, in particular a disorder selected from allergic inflammation, acute inflammation and chronic inflammation.


14. The compound according to claim 12 wherein said disorder is chosen from asthma, chronic obstructive pulmonary disease (COPD), rheumatoid arthritis, multiple sclerosis, inflammatory bowel diseases (IBD), psoriasis, atherosclerosis, thrombosis, stroke, acute coronary syndrome, stable angina, peripheral vascular disease, critical leg ischemia, intermittent claudication, abdominal aortic aneurysm and myocardial infarction.

Patentansprüche

1. Verbindung entsprechend Formel



oder



wobei

der Ring (a) ausgewählt ist aus einem bicyklischen Heterozyklyl-Rest und einem bicyklischen Heterozyklyl-Rest, der mit von einem bis drei Substituenten substituiert ist, die unabhängig ausgewählt sind aus der Gruppe bestehend aus Halogen, Hydroxyl, Alkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen, Acyl-Gruppen enthaltend ein bis vier Kohlenstoffatome, Alkoxy-Gruppen enthaltend ein bis vier Kohlenstoffatome, Fluoroalkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen, Fluoroalkoxy-Gruppen enthaltend ein bis vier Kohlenstoffatome, Formyl, Cyano, Phenyl, Heteroaryl, Nitro und Oxo;

Q O ist;

R¹ ausgewählt ist aus H und Alkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen;

n null oder eine ganze Zahl von 1-4 ist;

p eine ganze Zahl von 1-4 ist;

HET ein 4-7-gliedriger gesättigter stickstoffhaltiger Heterozyklus ist; und

ZW zusammengefasst H ist oder

Z ein $(\text{CH}_2)_{0-10}$ ist, bei welchem optional ein oder zwei (CH_2) durch einen $\text{C}_3\text{-C}_6$ Carbonzyklus, einen $\text{C}_3\text{-C}_6$ Heterozyklus, $-\text{O}-$, $-\text{NR}^{10}-$, $-\text{SO}-$, $-\text{S}(\text{O})_2-$, $-\text{C}(=\text{O})-$ oder $-\text{C}(=\text{O})(\text{NH})-$, ersetzt sein können, vorausgesetzt, dass jene $\text{O}-$, $-\text{NR}^{10}-$, $-\text{SO}-$, $-\text{S}(\text{O})_2-$, $-\text{C}(=\text{O})-$ oder $-\text{C}(=\text{O})(\text{NH})-$ -Reste nicht an der Verbindungsstelle zum Stickstoff sind und wenigstens durch ein $-(\text{CH}_2)_2$ -getrennt sind;


jeder auftretende R^{10} separate ausgewählt ist aus der Gruppe bestehend aus H und Alkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen;

W ausgewählt ist aus Acyl, Hydroxyl, Carboxyl, Amino, Carboxamido, Sulfonamide, Aminoacyl, $-\text{COOalkyl}$, $-\text{CHO}$, $-\text{C}(\text{O})\text{fluororalkyl}$, $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{Oalkyl}$, $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{Ofluoroalkyl}$, $-\text{SH}$, $-\text{C}(\text{O})\text{NH}(\text{OH})$, $-\text{C}(\text{O})\text{N}(\text{OH})\text{R}^4$, $-\text{N}(\text{OH})\text{C}(\text{O})\text{OH}$, $-\text{N}(\text{OH})\text{C}(\text{O})\text{R}^4$, Heterocyclyl, substituiertem Aryl, und substituiertem Heterocyclyl; und


R^4 ausgewählt ist aus der Gruppe bestehend aus H und Alkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen.

2. Verbindung entsprechend Anspruch 1, wobei der Ring (a) ausgewählt ist aus einem Indazol-, einem Indol- und einem Isoindolring.

3. Verbindung entsprechend Anspruch 2 gemäß der Formel



oder



wobei R^2 und R^3 jeweils unabhängig ausgewählt sind aus der Gruppe bestehend aus H, Halogen, Hydroxyl, Alkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen, Acyl-Gruppen enthaltend ein bis vier Kohlenstoffatome, Alkoxy-Gruppen enthaltend ein bis vier Kohlenstoffatome, Fluoralkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen, Fluoralkoxy-Gruppen enthaltend ein bis vier Kohlenstoffatome, Formyl, Cyano, Phenyl, Heteroaryl, und Nitro;

n 1, 2 oder 3 ist; und

p 2 oder 3 ist.


4. Verbindung entsprechend Anspruch 1, wobei

HET ausgewählt ist aus Piperidin und Pyrrolidin;

Z $(\text{CH}_2)_{0-10}$ ist, bei welchem optional ein oder zwei (CH_2) durch $-\text{O}-$, $-\text{NR}^{10}-$, $-\text{SO}-$, $-\text{S}(\text{O})_2-$, $-\text{C}(=\text{O})-$ oder $-\text{C}(=\text{O})(\text{NH})$ ersetzt sein können; und

W ausgewählt ist aus der Gruppe bestehend aus Wasserstoff, Acyl, Hydroxy, Carboxy, Amino, Carboxamido und Aminoacyl.

5. Verbindung entsprechend Anspruch 4 gemäß der Formel



wobei


R^2 ausgewählt ist aus H, Hydroxyl, und Halogen;

R^3 ausgewählt ist aus H und Halogen;

Z ausgewählt ist aus $(CH_2)_{0-3}$; und

W ausgewählt ist aus der Gruppe bestehend aus Wasserstoff, Acyl, Hydroxyl, Carboxyl, Amino, Carboxamido, Aminoacyl und $COOR^4$.

6. Verbindung entsprechend Anspruch 4 gemäß der Formel



wobei


R^2 ausgewählt ist aus H, Hydroxyl, und Halogen;

R^3 ausgewählt ist aus H und Halogen;


Z $(CH_2)_{0-3}$ ist; und

W ausgewählt ist aus der Gruppe bestehend aus Wasserstoff, Hydroxyl, Carboxyl und Carboxamido.

7. Verbindung entsprechend Anspruch 2 gemäß Formel



oder



wobei

R^2 ausgewählt ist aus der Gruppe bestehend aus H, Alkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen, Fluoralkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen, Phenyl und Oxo;


R^3 ausgewählt ist aus der Gruppe bestehend aus H, Halogen, Alkylgruppen aus von 1 bis 6 Kohlenstoffatomen, Acyl-Gruppen enthaltend ein bis vier Kohlenstoffatome, Alkoxy-Gruppen enthaltend ein bis vier Kohlenstoffatome, Fluoralkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen, Fluoralkoxy-Gruppen enthaltend ein bis vier Kohlenstoffatome, Formyl, Cyano, Phenyl, Heteroaryl und Nitro;

n von 1-3 ist; und

p 2 oder 3 ist.

8. Verbindung entsprechend Anspruch 7, wobei HET ausgewählt ist aus Piperidin und Pyrrolidin;
Z $(CH_2)_{0-10}$ ist; bei welchem optional ein oder zwei (CH_2) durch -O-, $-NR^{10}$ -, $-SO$ -, $-S(O)_2$ -, $-C(=O)$ - oder $-C(=O)NH$ - ersetzt sein können; und
W ausgewählt ist aus der Gruppe bestehend aus Wasserstoff, Acyl, Hydroxyl, Carboxyl, Amino, Carboxamido und Aminoacyl.

9. Verbindung entsprechend Anspruch 2 gemäß Formel



wobei

Y N oder CR^5 ist;


R^7 ausgewählt ist aus H und Alkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen; und

R^5 ausgewählt ist aus H und Alkyl-Gruppen aus von 1 bis 6 Kohlenstoffatomen.


10. Pharmazeutische Zusammensetzung umfassend einen pharmazeutisch akzeptablen Träger und eine therapeutisch effektive Menge von wenigstens einer Verbindung entsprechend einem der Ansprüche 1 bis 9.
11. Verbindung entsprechend einem der Ansprüche 1 bis 9 zur Verwendung bei der Inhibierung der Leukotrien A4 Hydrolase, wobei das LTA4H Enzym mit einer therapeutisch effektiven Menge einer Verbindung entsprechend einem der Ansprüche 1 bis 9 in Kontakt gebracht wird.
12. Verbindung entsprechend einem der Ansprüche 1 bis 9 zur Verwendung bei der Behandlung einer Erkrankung im Zusammenhang mit der Leukotrien A4 Hydrolase, wobei einem Säuger eine therapeutisch effektive Menge einer Verbindung entsprechend einem der Ansprüche 1 bis 9 oder eines Salzes, Hydrats oder Esters davon verabreicht wird.
13. Verbindung entsprechend Anspruch 12, wobei die Erkrankung in Zusammenhang mit einer Entzündung steht, insbesondere eine Erkrankung ausgewählt aus allergischen Entzündungen, akuten Entzündungen und chronischen Entzündungen.
14. Verbindung entsprechend Anspruch 12, wobei die Erkrankung ausgewählt ist aus Asthma, chronisch obstruktiver Lungenerkrankung (COPD), rheumatischer Arthritis, Multipler Sklerose, entzündliche Darmerkrankung (IBD), Psoriasis, Arteriosklerose, Thrombose, Schlaganfall, akute Erkrankung der Herzkranzgefäße, stabile Angina, periphere Verschlusskrankheit, kritische Beinischämie; intermittierende Klaudikation, abdominales Aortenaneurysma und Herzinfarkt.

Revendications

1. Composé de formule :



ou



où

le cycle (a) est choisi parmi hétérocyclyle bicyclique et hétérocyclyle bicyclique substitué avec de un à trois substituants choisis indépendamment dans le groupe consistant en halogène, hydroxyle, les groupes alkyle de 1 à 6 atomes de carbone, les groupes acyle contenant un à quatre atomes de carbone, les groupes alcoxy contenant un à quatre atomes de carbone, les groupes fluoroalkyle de 1 à 6 atomes de carbone, les groupes fluoroalcoxy contenant un à quatre atomes de carbone, formyle, cyano, phényle, hétéroaryle, nitro et oxo;

Q est O;

R¹ est choisi parmi H et les groupes alkyle de 1 à 6 atomes de carbone;

n est zéro ou un entier de 1-4;

p est un entier de 1-4;

HET est un hétérocycle azoté saturé à 4-7 chaînons ; et

ZW pris dans son ensemble est H ou

Z est (CH₂)₀-₁₀ où un ou deux (CH₂) peuvent éventuellement être remplacés par un C₃-C₆ carbocycle, un C₃-C₆ hétérocycle, -O-, -NR¹⁰-, -SO-, -S(O)₂-, -C(=O)- ou -C(=O)(NH)-, à condition que lesdits -O-, -NR¹⁰-, -SO-, -S(O)₂-, -C(=O)- ou -C(=O)(NH)- ne soient pas le point de liaison à l'azote et soient séparés par au moins un -(CH₂)₂-;


R¹⁰ est choisi séparément dans chaque cas dans le groupe consistant en H et les groupes alkyle de 1 à 6 atomes de carbone;

W est choisi parmi acyle, hydroxyle, carboxyle, amino, carboxamido, sulfonamide, aminoacyle, -COOalkyle, -CHO, -C(O)fluoroalkyle, -C(O)CH₂C(O)Oalkyle, -C(O)CH₂C(O)Ofuoroalkyle, -SH, -C(O)NH(OH), -C(O)N(OH)R⁴, -N(OH)C(O)OH, -N(OH)C(O)R⁴, hétérocyclyle, aryle substitué et hétérocyclyle substitué ; et


R⁴ est choisi dans le groupe consistant en H et les groupes alkyle de 1 à 6 atomes de carbone.

2. Composé selon la revendication 1, où le cycle (a) est choisi parmi un cycle indazole, un cycle indole et un cycle isoindole.

3. Composé selon la revendication 2 de formule:



ou



où

R² et R³ sont choisis chacun indépendamment dans le groupe consistant en H, halogène, hydroxyle, les groupes alkyle de 1 à 6 atomes de carbone, les groupes acyle contenant un à quatre atomes de carbone, les groupes alcoxy contenant un à quatre atomes de carbone, les groupes fluoroalkyle de 1 à 6 atomes de carbone, les groupes fluoroalcoxy contenant un à quatre atomes de carbone, formyle, cyano, phényle, hétéroaryle et nitro;
n est 1, 2 ou 3; et
p est 2 ou 3.


4. Composé selon la revendication 1, où

HET est choisi parmi pipéridine et pyrrolidine;

Z est (CH₂)₀₋₁₀ où un ou deux (CH₂) peuvent éventuellement être remplacés par -O-, -NR¹⁰-, -SO-, -S(O)₂-, -C(=O)- ou -C(=O)NH-, et

W est choisi dans le groupe consistant en hydrogène, acyle, hydroxy, carboxy, amino, carboxamido et aminoacyle.

5. Composé selon la revendication 4 de formule



où


R² est choisi parmi H, hydroxyle et halogène;

R³ est choisi parmi H et halogène;

Z est choisi parmi (CH₂)₀₋₃ et

W est choisi dans le groupe consistant en hydrogène, acyle, hydroxyle, carboxyle, amino, carboxamido, aminoacyle et COOR⁴.

6. Composé selon la revendication 4 de formule



où


R² est choisi parmi H, hydroxyle et halogène;

R³ est choisi parmi H et halogène;


Z est $(\text{CH}_2)_{0-3}$; et

W est choisi dans le groupe consistant en hydrogène, hydroxyle, carboxyle et carboxamido.

7. Composé selon la revendication 2 de formule:



ou



où

R^2 est choisi dans le groupe consistant en H, les groupes alkyle de 1 à 6 atomes de carbone, les groupes fluoroalkyle de 1 à 6 atomes de carbone, phényle et oxo;

R^3 est choisi dans le groupe consistant en H, halogène, les groupes alkyle de 1 à 6 atomes de carbone, les groupes acyle contenant un à quatre atomes de carbone, les groupes alcoxy contenant un à quatre atomes de carbone, les groupes fluoroalkyle de 1 à 6 atomes de carbone, les groupes fluoroalcoxy contenant un à quatre atomes de carbone, formyle, cyano, phényle, hétéroaryle et nitro;

n est de 1-3; et

p est 2 ou 3.


8. Composé selon la revendication 7, où

HET est choisi parmi pipéridine et pyrrolidine;

Z est $(\text{CH}_2)_{0-10}$ où un ou deux (CH_2) peuvent éventuellement être remplacés par $-\text{O}-$, $-\text{NR}^{10}-$, $-\text{SO}-$, $-\text{S}(\text{O})_2-$, $-\text{C}(=\text{O})-$ ou $-\text{C}=\text{O}(\text{NH})-$, et

W est choisi dans le groupe consistant en hydrogène, acyle, hydroxyle, carboxyle, amino, carboxamido et aminoacyle.

9. Composé selon la revendication 2 de formule:



où

Y est N ou CR⁵;

R⁷ est choisi parmi H et les groupes alkyle de 1 à 6 atomes de carbone; et

R⁵ est choisi parmi H et les groupes alkyle de 1 à 6 atomes de carbone.

10. Composition pharmaceutique comprenant un vecteur pharmaceutiquement acceptable et une quantité thérapeutiquement efficace d'au moins un composé selon l'une quelconque des revendications 1-9.

11. Composé selon l'une quelconque des revendications 1-9 destiné à être utilisé dans l'inhibition de la leucotriène A₄ hydrolase où l'enzyme LTA₄H est mise en contact avec une quantité thérapeutiquement efficace d'un composé selon l'une quelconque des revendications 1-9.

12. Composé selon l'une quelconque des revendications 1-9 destiné à être utilisé dans le traitement d'un trouble associé avec la leucotriène A₄ hydrolase où une quantité thérapeutiquement efficace d'un composé ou d'un sel, hydrate ou ester de celui-ci selon l'une quelconque des revendications 1-9 est administrée à un mammifère.

13. Composé selon la revendication 12 où ledit trouble est associé avec une inflammation, en particulier un trouble choisi parmi une inflammation allergique, une inflammation aiguë et une inflammation chronique.

14. Composé selon la revendication 12 où ledit trouble est choisi parmi l'asthme, la bronchopneumopathie chronique obstructive (COPD), la polyarthrite rhumatoïde, la sclérose en plaques, les maladies inflammatoires chroniques de l'intestin (MICI), le psoriasis, l'athérosclérose, la thrombose, l'attaque, le syndrome coronarien aigu, l'angor stable, les maladies vasculaires périphériques, l'ischémie critique des jambes, la claudication intermittente, l'anévrisme de l'aorte abdominale et l'infarctus du myocarde.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20050043378 A [0003]
- US 20050043379 A [0003]
- US 2005043378 A [0004]
- US 2005043379 A [0004]
- US 5324718 A [0038]
- US 5472954 A [0038]

Non-patent literature cited in the description

- **HELGADOTTIR et al.** *Nature Genetics*, 08 February 2004 [0002]
- **HAKONARSSON et al.** *J.Am.Med.Assoc.*, 2005, vol. 293, 2245-2256 [0002]
- **PENNING T. D.** *Current Pharmaceutical Design*. Bentham Science Publishers, February 2001, vol. 3, 163-179 [0005]
- **PENNING et al.** *Journal of Medicinal Chemistry*. American Chemical Society, 2000, vol. 43, 721-725 [0005]
- **REMENAR et al.** *J.Am.Chem.Soc.*, 2003, vol. 125, 8456-8457 [0038]
- Remington: *The Science and Practice of Pharmacy*. 1995, vol. 1, 176-177 [0038]
- *Maehr J. Chem. Ed.*, 1985, vol. 62, 114-120 [0040]
- **T.W. GREENE.** *Protective Groups in Organic Synthesis*. John Wiley & Sons, 1991 [0041]
- **KACHUR et al.** *J. Pharm. Exp. Ther.*, 2002, vol. 300 (2), 583-587 [0044]
- **PENNING.** *Curr. Pharm. Des.*, 2001, vol. 7 (3), 163-179 [0044]
- **MUELLER et al.** *J. Biol. Chem.*, 1996, vol. 271, 24345-24348 [0129] [0133]