



(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

- (15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 16, 17, 18, 29, 46,
55, 81, 102, 113, 126, 142
- (51) Int Cl.:
C07F 15/00^(2006.01) A61K 31/28^(2006.01)
- (86) International application number:
PCT/CH2007/000234
- (48) Corrigendum issued on:
30.12.2015 Bulletin 2015/53
- (87) International publication number:
WO 2007/128158 (15.11.2007 Gazette 2007/46)
- (45) Date of publication and mention
of the grant of the patent:
05.08.2015 Bulletin 2015/32
- (21) Application number: **07720131.7**
- (22) Date of filing: **09.05.2007**

(54) **TRANSITION METAL COMPLEXES FOR INHIBITING RESISTANCE IN THE TREATMENT OF
CANCER AND METASTASIS**

ÜBERGANGSMETALLKOMPLEXE ZUM INHIBIEREN VON RESISTENZ BEI DER BEHANDLUNG
VON KREBS UND METASTASEN

COMPLEXES DE MÉTAUX DE TRANSITION PERMETTANT d'INHIBER LA RÉSISTANCE DANS LE
TRAITEMENT DU CANCER ET DES MÉTASTASES

- | | |
|---|---|
| <p>(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE
SI SK TR</p> <p>(30) Priority: 09.05.2006 US 799433 P</p> <p>(43) Date of publication of application:
21.01.2009 Bulletin 2009/04</p> <p>(73) Proprietor: Ecole Polytechnique Fédérale de
Lausanne (EPFL)
1015 Lausanne (CH)</p> <p>(72) Inventors:
• DYSON, Paul, Joseph
CH-1024 Ecublens (CH)
• ANGE, Wee, Han
CH-1024 Ecublens (CH)</p> | <p>(74) Representative: Schneiter, Sorin
Abrema Agence Brevets & Marques
Ganguillet
Avenue du Théâtre 16
P.O. Box 5027
1002 Lausanne (CH)</p> <p>(56) References cited:
• ANG, WEE HAN ET AL: "Rational Design of
Platinum(IV) Compounds to Overcome
Glutathione-S- Transferase Mediated Drug
Resistance" JOURNAL OF THE AMERICAN
CHEMICAL SOCIETY , 127(5), 1382-1383 CODEN:
JACSAT; ISSN: 0002-7863, 2005, XP002422722
• SERLI, B. ET AL.: "Is the aromatic fragment of
piano-stool ruthenium compounds an essential
feature for anticancer activity? The development
of new Ru(II)-[9]aneS3 analogues" EUR. J.
INORG. CHEM., vol. 2005, 2005, pages 4323-3434,
XP002422723</p> |
|---|---|

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] This invention relates to organometallic compounds of the metals Ru, Os, Rh, and Ir, to their use in medicine, particularly for the treatment and/or prevention of cancer, and more particularly in the treatment of metastasis. This new class of compounds contains, covalently bound to any of the ligands of the metal, or directly to the metal, a bioactive organic compound which is an inhibitor of resistance pathway.

Background and Objectives of the Invention

[0002] Resistance against bioactive principles that directly target the origin of a disease are widespread, and one or more of a number of known resistance pathways are at their origin. For example, the Glutathione-S-Transferase (GST) is a superfamily class of isozymes that constitute the main cellular defense against xenobiotics, including bioactive principle designed for the treatment of a disease. They catalyse the conjugation of endogenous glutathione (GSH) with the electrophilic groups of substrates, the first step in the mercapturic acid pathway that leads to elimination of toxic compounds. The overexpression of several subclasses of GST, namely GST- π and GST- α , has been linked to the multidrug resistance phenomenon of certain anticancer drugs, such as *Cisplatin* and *adriamycin*. More recently, GSTP1-1 (GST- π subclass) was found to mediate the c-Jun N-Terminal Kinase (JNK) signal transduction pathway, an important control of cell survival. It was found to have a significant affinity for the C terminus of JNK and therefore could potentially interfere with and suppress downstream induction of cellular apoptosis. Clearly, GST is a potential target for chemotherapeutic drug design, in order to inhibit resistance against anti-cancer drugs.

[0003] Ruthenium-based compounds have shown some potential as anticancer drugs. For example, US 4980473 discloses 1,10-phenanthroline complexes of ruthenium(II) and cobalt(II) which are said to be useful for the treatment of tumour cells in a subject.

[0004] Some other ruthenium(II) and ruthenium(III) complexes which have been shown to exhibit antitumour activity are mentioned in Guo et al, *Inorganica Chimica Acta*, 273 (1998), 1-7, specifically *trans*-[RuCl₂(DMSO)₄], *trans*-[RuCl₄(imidazole)₂] and *trans*-[RuCl₄(indazole)₂]. Clarke *et al* have reviewed the anticancer, and in particular the antimetastatic, activity of ruthenium complexes: *Chem. Rev.*, 1999, 99, 251-253. Also, Sava has reviewed the antimetastatic activity in "Metal Compounds in Cancer Therapy" Ed by S P Fricker, Chapman and Hall, London 1994, p. 65-91.

[0005] Dale et al, *Anti-Cancer Drug Design*, (1992), 7, 3-14, describes a metronidazole complex of ruthenium(II) ie, [(C₆H₅)RuCl₂(metronidazole)] and its effect on DNA and on *E. coli* growth rates. Metronidazole sensitises hypoxic tumour cells to radiation and appears to be an essential element of the complexes of Dale et al. There is no indication that the complexes would be at all effective in the absence of the metronidazole ligand.

[0006] Kramer et al, *Chem Eur J.*, 1996, 2, No. 12, p. 1518-1526 discloses half sandwich complexes of ruthenium with amino esters, Bennett et al, *Canadian Journal of Chemistry*, (2001), 79, 655-669 discloses certain ruthenium(II) complexes with acetylacetonate ligands. Oro et al, *J Chem Soc, Dalton Trans*, (1990), 1463 describes ruthenium(II) complexes containing -p-cymene and acetylacetonate ligands WO 01/130790 discloses ruthenium(II) compounds and their use as anticancer agents. The compounds have neutral N-donor ligands and the resulting ruthenium complex is generally positively charged.

[0007] WO 02/102572 also discloses ruthenium(II) compounds that have activity against cancer cell lines. Again, the complexes are generally positively charged. Complexes are disclosed containing a bidentate ligand which is a neutral diamine ligand.

[0008] Chen et al, *J. Am. Chem. Soc.*, volume 124, no 12, 3064, (2002), describes the mechanism of binding of ruthenium complexes to guanine bases. The binding model requires NH bonds from a diamino ligand to be present in the complex for hydrogen bonding to the guanine base. Similarly, Morris et al, *J. Med. Chem.*, volume 44, 3616-3621, (2001), describes the selectivity of ruthenium(II) complexes for binding to guanine bases.

[0009] Further references concerned with Ruthenium, complexes for treatment of cancer are WO 06/018649, US 2006/0058270, US 2005/0239765.

[0010] Serli et al, *Eur. J. Inorg. Chem.*, no 17, 3423, (2005), demonstrates that the Ru^{II}-arene in Ruthenium organometallic complexes ("piano-stool" complexes) is not an essential feature for the *in vitro* anticancer activity and may be replaced by another ligand with a low steric demand, such as [9]aneS₃.

[0011] Ang et al, *J. Am. Chem. Soc.*, volume 127, no 5, 1382, (2005), provides a compound (ethacraplatin) having two functional properties: anticancer activity and inhibiting activity of GST, wherein the cytotoxicity of the cisplatin anticancer drug is improved by linking a GST inhibitor to said drug. The anticancer activity of ethacraplatin is linked to the presence of the cisplatin. The linked GST inhibitor is able to reverse the intrinsic drug-associated resistance of cisplatin-resistant cancer cell lines in the presence of cisplatin in the ethacraplatin compound.

[0012] Very few, if any, of the compounds and complexes of the prior art cited above have resulted in clinical phase studies, not to mention actual therapies. The reason for the poor performance of these principles are manifold and may be linked to toxicity problems or un-sufficient efficiency in treatment.

[0013] It is thus an objective of the present invention to explore new ways for treating cancer, for example based on work done in the area of complexes of transition metals.

[0014] It is a further aspect of the present invention to increase the activity and/or efficiency of therapies against diseases and in particular cancer. More particularly, it is an objective to reduce the resistance intrinsic to or developed against therapies, and in particular against resistance in cancer chemotherapies. By reducing resistance to bioactive principles against diseases and in particular cancer, it is hoped to increase the overall efficiency of the therapy. With increased efficiency, lower levels of the bioactive principle needs to be administered, which may further reduce side effects linked to the treatment.


[0015] Tumors of various kinds can be removed surgically, the most relevant problem of these cancers being the development of metastasis. It is thus an objective of the present invention to prevent and/or treat metastasis. In particular, it is an objective of the invention to assist the treatment for prevention and/or treatment of metastasis.

Summary of the Invention

[0016] The present invention relates to complexes of transition metals comprising ligands in any form and of any nature, with the proviso that at least one of the ligands comprises at least one bioactive organic compound selected from inhibitors of resistance pathways and/or pharmaceutically acceptable derivatives thereof. Inhibitors may be directly attached to the transition metal as a ligand, or it may be covalently bound to a ligand of the transition metal.


[0017] Remarkably, the present inventors observed low toxicity and high inhibition of resistance pathways when administering the organometallic compounds of the present invention. Surprisingly, high proliferation inhibition of the organometallic compounds of the present invention on carcinoma cells was observed. Moreover, toxicity against healthy, normal cells remained very low.

[0018] Accordingly, in a first aspect, the present invention provides, as claimed in claim 1, an organometallic compound of formula (XIV),




in which A is an arene, optionally substituted and optionally comprising one or more heteroatoms; R_{18} , R_{19} , R_{20} , are ligands of the central ruthenium atom which are, independently of each other, selected from halogens and/or N-, O-, S-, or P- donor ligands; R_{17} is an optional residue selected from an alkyl, alkenyl, alkynyl, aryl, optionally substituted and optionally comprising one or more heteroatoms; whereby at least one bioactive organic compound constitutes at least one selected from R_{17} - R_{20} , or is covalently linked to any of the R_{17} - R_{20} , with the proviso that residues selected from R_{17} - R_{20} , which constitute the bioactive organic compound, or which are covalently bound the bioactive organic compound, are not halogens; and wherein said bioactive organic compound is selected from ethacrynic acid, peptidomimetics based on glutathione, p-chlorophenoxyisobutyrate, Gossypol, indomethacin, non-steroidal anti-inflammatory compounds based on ibuprofen and ketoprofen, misonidazole, Piroprost, Sulfasalazine, sulfoxime-based compounds such as buthionone sulfoxime and methionone sulfoxime, S-sulfocysteine, S-sulfohomocysteine, cystamine, quinidine, vinblastine, terfernadine, tamoxifen, verapamil, cyclosporin, amitriptyline, progesterone, pleurotin, azelaic acid, bischloroethylnitrosourea, or palmarumycin, and their pharmaceutical acceptable derivatives and salts.

[0019] In an embodiment, the arene A of this organometallic compound is an arene of the formula (VI)




(VI),

in which R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} are, independently of each other, hydrogen, alkyl, alkenyl, alkynyl, aryl, optionally substituted and optionally comprising one or more heteroatoms, and in which two or more residues selected from R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} may be covalently linked with each other thus forming a bi- or tri-, or polycyclic system, and in which any of the residues R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} optionally constitutes or is optionally bound to the bioactive compound. In a preferred embodiment, the organometallic compound of the invention comprises the structure of formula (X):




(X),

in which, the arene A is an optionally substituted, benzyl or cyclopentadienyl, B is an optional linking group, which may be selected from alkyl, alkenyl, alkynyl and aryl, which is optionally substituted, which optionally comprises one or more heteroatoms and has 0-15, preferably 1-8 carbon atoms; and X is a functional group of B or the cyclic system, through which the bioactive compound is bound to B or to the cyclic system. Preferably, the organometallic compound of the invention comprises the structure of formula (XI) or (XII):




(XI)




(XII), in which m is 1-10.

[0020] In a further embodiment, the N-donor ligand of the organometallic compound of the invention comprises the structure of formula (II).




(II),

in which M is the transition metal Ru; the dashed line represents bonding, as a monodentate ligand, to the transition metal Ru; the circle represents a mono- or polycyclic system, comprising at least one N-heteroatom, indicated as N, which provides an electron pair enabling attachment to M, the cyclic system being optionally further substituted; B is an optional linking group, which is selected from alkyl, alkenyl, alkynyl and aryl, is optionally substituted, and optionally comprises one or more heteroatoms and has 0-15 carbon atoms; and X is a functional group of B or the cyclic system, through which the bioactive compound is bound to B or to the cyclic system. Preferably, the N-donor ligand of the above organometallic compound comprises the structure of formula (III) below,




to which the bioactive compounds is linked by means of an amide bond and in which, M is the transition metal Ru; the dashed line represents bonding, as a monodentate ligand, to the transition metal Ru; the circle represents a mono- or polycyclic system comprising at least two N-heteroatoms; n is 1-10, preferably 2-6, most preferably 3-4; and X is as defined above.

[0021] According to a further embodiment, the P-donor ligand of the organometallic compound of the invention comprises the structure of formula (XIII):



in which R_{14} R_{15} R_{16} may be the same or different, are each C1-C6 alkyl, alkenyl, alkynyl, aryl, optionally substituted and optionally comprising one or more heteroatoms, or R_{14} R_{15} R_{16} may together with the phosphorous atom form a cycloalkyl group, such group being optionally heterocyclic. Preferably, the P-donor ligand of formula (XIII) in the organometallic compound of the invention is a residue suitable of increasing the solubility of the organometallic compound of claim 1 or 2 in water.

[0022] In a preferred embodiment, the organometallic compound of the invention comprises the structure of formula (XV):



in which X is a halogen; Y is a halogen or N-, P-, O- or S-donor ligand as defined above; and, in which, if the bioactive compound is bound to R_{17} , R_{17} corresponds to "B-X-" as defined in formula (X) above, and R_{18} is a N-, P-, O- or S-donor ligand as defined above; or, if the bioactive compound is bound to R_{18} , or constitutes R_{18} , R_{18} is, respectively, a N-, P-, O- or S-donor ligand as defined above to which the bioactive compound is covalently bound; or R_{18} constitutes the bioactive compound, the bioactive compound comprising itself N-, P-, O- or S-atoms functioning as donors suitable to attach it to the central Ru.

[0023] In a further embodiment of the invention, these organometallic compounds of the invention are provided for use as a medicament.

[0024] In a further embodiment, these organometallic compounds are provided for use in the treatment and/or prevention of cancer and/or metastasis.

[0025] In a preferred embodiment, these organometallic compounds are provided for use in reducing resistance of cancers and/or metastasis against anti-cancer drugs.

[0026] In a further aspect, the invention provides a composition comprising an anti-cancer drug and the organometallic compound of the invention for use in treating and/or preventing metastasis.

[0027] According to another aspect, the invention provides the use of these organometallic compounds for the manufacture of a medicament for treating and/or preventing metastasis. In a further aspect, the invention provides the use of the composition of the invention for the manufacture of a medicament for treating and/or preventing metastasis.

[0028] Further embodiments are provided in the appended claims and will be apparent from the detailed description of preferred embodiments below.

[0029] Aside from the present invention as claimed in the aspects and embodiments above, an organometallic compound of the general formula (I) below is also described,



which may be charged or neutral, and which may be present in the form of a salt and/or an optimally resolved enantiomer, in which,

- M is a transition metal selected from the group of Ru, Os, Rh, and Ir;
- $R_1, R_2, R_3, R_4, R_5, R_6$ are neutral or charged ligands of the transition metal, whereby two, three or more of the ligands $R_1, R_2, R_3, R_4, R_5, R_6$ may be present in the form of one or more single compounds, the single compound being
 - a bi-, tri- or polydentate compound, and/or,
 - an alkene, alkyne, cyclopentadienyl and/or an arene, the alkene, alkyne, cyclopentadienyl and/or an arene being optionally substituted and optionally I comprising one or more heteroatoms;

in which at least one bioactive organic compound selected from inhibitors of resistance pathways, related compounds, and/or derivatives of any of the fore-mentioned, is present in the organometallic compound, whereby the bioactive organic compound is directly attached to the metal, thus constituting at least one of the ligands selected from R_1 - R_6 , and/or is covalently bound to any of the ligands selected from R_1 - R_6 . It is also shown that said organometallic compounds may be for use as a medicament. They even may be provided in the preparation of a medicament for the inhibition of resistance pathways and/or for treating cancer, and in particular metastasis. Further a method for treating and/or preventing metastasis comprising the step of administering to an individual an effective amount of said organometallic compound is also described. A further method for treating and/or preventing metastasis is also shown and comprises the step of administering to an individual an effective amount of an anti-cancer drug and, in parallel, an effective amount of said organometallic compound.

[0030] In the figures,

Figure 1 shows a single crystal X-ray diffraction structure of compound (3), which is an example of an organometallic compound of the invention.

Figure 2 shows a single crystal X-ray diffraction structure of compound (6), which is an example of an organometallic compound of the invention.

Figure 3 shows a single crystal X-ray diffraction structure of compound (8), which is an organometallic compound having, amongst others, oxalate as a bivalent ligand.

Figure 4. illustrates the process for preparing (3), an example of an organometallic compound of the invention.

Figure 6 illustrates the process for preparing (6), an example of an organometallic compound of the invention.

Figure 7 illustrates the process for preparing (7), an example of an charged organometallic compound of the invention, present as its tetrafluoroborate salt.

Figure 8 illustrates the process for preparing (8) and (9), organometallic compounds that are used as controls for evaluating the activity of organometallic compounds of the invention.

Detailed Description of the Preferred Embodiments of the Invention

[0031] The present invention relates to organometallic compounds and their use as medicaments. The organometallic compound comprises a transition metal M, and, linked to the transition metal, a number of ligands. Ligands include donors of electron pairs and/or donors of π -orbitals from unsaturated bonds in organic molecules, for example.

[0032] The term "comprise" or "comprising", for the purpose of the present invention is intended to mean "including amongst other". It is not intended to mean, "consisting only of".

[0033] The transition metal M is selected from the group of Ru, Os, Rh, and Ir. Preferably, M is Ruthenium. The transition metal may be present in any oxidation state known with respect to the specific transition metal. For example, the oxidation state may be II or III. Preferably, M is Ruthenium (II).

[0034] $R_1, R_2, R_3, R_4, R_5, R_6$ are neutral or charged ligands attached to the transition metal. Suitable ligands include halogen ions, such as F, Cl⁻, Br⁻ and I⁻, preferably chloride. One, two or more ligands selected from R_1 - R_6 may be provided by halogens.

[0035] The term "selected from" a group indicated by " R_1 - R_6 ", for example, or " R^{N1} - R^{N3} " as in the above

paragraph or indicated below, respectively, refers to the fact that one or more selected from all the individuals of the group may be selected independently of each other. Accordingly, if one specimen of R_1 - R_6 is to be selected, any one of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 can be selected.

[0036] One, two, three or more of the ligands R_1 , R_2 , R_3 , R_4 , R_5 , R_6 may be selected from ligands providing electron pairs. For example, the ligands may be selected from N-, P-, O- or S-donor ligands. Preferably, at least one ligand of the organometallic compound of the invention is selected from N-, P-, O- or S-donor ligands. For example, at least one N-donor and/or at least one P-donor ligand is present.

[0037] Examples of N-donor ligands are nitrile ligands ($N\equiv C-R$); azo ligands ($N=N-R$); aromatic N-donor ligands; amine ligands ($NR^{N1}R^{N2}R^{N3}$); azide (N_3^-); cyanide ($N\equiv C^-$); isothiocyanate (NCS^-).


[0038] In both nitrile and azo ligands, R may be selected from alkyl, alkenyl, alkynyl, and aryl, optionally substituted and optionally comprising one or more heteroatoms.

[0039] Aromatic N-donor ligands include optionally substituted pyridine, pyridazine, pyrimidine, purine and pyrazine, for example. Substituents may be selected from alkyl, alkenyl, alkynyl, and aryl, optionally substituted and optionally comprising one or more heteroatoms.

[0040] R^{N1} , R^{N2} , and R^{N3} may, independently of each other, be selected from H, alkyl, alkenyl, alkynyl, aryl, optionally substituted and optionally comprising one or more heteroatoms.

[0041] Preferably, two or all three of R^{N1} - R^{N3} may be fused to form a cyclic N-donor compound. Furthermore, any of R^{N1} - R^{N3} may be linked covalently to any other ligand of M, to provide bi-, tridentate or other polyvalent ligands. Preferably, R^{N1} , R^{N2} , and R^{N3} are, independently of each other, selected from H and C_{1-8} alkyl, optionally substituted.

[0042] According to a preferred embodiment, at least one of the ligands selected from R_1 - R_6 is an N-donor ligand comprising the structure of formula (II) below,



in which,

M is the transition metal;

the circle represents a mono- or polycyclic system, comprising at least one N-heteroatom, indicated as N, which provides an electron pair enabling attachment to M;

B is an optional linking group, which may be selected from alkyl, alkenyl, alkynyl and aryl, which is optionally substituted, which optionally comprises one or more heteroatoms and has 0-15, preferably 1-8 carbon atoms;

X is a functional group of B or the cyclic system, through which the bioactive compound is bound to B or to the cyclic system.


[0043] The cyclic system may further be substituted.

[0044] Preferably, in formula (II), the circle is a heterocyclic ring, for example a heterocyclic arene.

[0045] Preferably, it is a 5- or 6-membered heterocyclic ring, for example arene. Preferably, A is a C_1 - C_{16} , more preferably a C_2 - C_4 alkylene.

[0046] Preferably, X is selected from -O- and from NH-. Accordingly, X preferably represents an ether, ester or peptide group, of which the carbonyl part, if applicable, may be part of the "bioactive compound" or of B.


[0047] According to a particularly preferred embodiment, any one of the ligands selected from R_1 - R_6 is a N-donor ligand comprising the structure of formula (III) below, to which the bioactive compound is linked by means of an amide bond:



in which,

the dashed line represents bonding, as a monodentate ligand, to the transition metal;
 the circle represents a mono- or polycyclic system comprising at least two N-heteroatoms;
 n is 1-10, preferably 2-6, most preferably 3-4;
 X is as defined above for (III).

[0048] The cyclic system may, for example, be selected from imidazole, purine, pyrazine, pyrimidine, 1,8-naphthydrin, chinoxaline, chinazoline, pteridine. Preferably, the cyclic system is imidazole, resulting in N-donor ligands comprising structures as illustrated in formula (IV) and (V) below.



in which n is as indicated for (III) above.

[0049] The bioactive compound comprises, in its released and/or original, active form, a carboxy-group, which is linked to the N-atom or O-atom indicated in (IV) or (V), upon formation of an amide bond.

[0050] Examples of P-donor ligands are $PR^{P1}R^{P2}R^{P3}$, in which R^{P1} , R^{P2} , and R^{P3} are defined as R^{N1} , R^{N2} , and R^{N3} above, wherein a fused P-donor ligand may arise if two or all three of R^{N1} - R^{N3} are fused.

[0051] A preferred example of a P-donor ligand is PTA (1,3,5-triaza-7-phospha-adamantane).

[0052] S-donor ligands are ligands which bind to M via a sulphur atom. Examples include thiosulfate ($S_2O_3^{2-}$); isothiocyanate (NCS^-); sulfoxide ligands ($R^{S1}R^{S2}SO$); thioether ligands ($R^{S1}R^{S2}S$); thiolate ligands ($R^{S1}S^-$); sulfinat ligands ($R^{S1}SO_2^-$); and sulfenat ligands ($R^{S1}SO^-$), wherein R^{S1} and R^{S2} are independently selected from alkyls, alkenyls, alkynyls, aryls, optionally substituted and optionally comprising one or more heteroatoms.

[0053] O-donor ligands are ligands which bind to M via an oxygen atom. Examples include carbonate (CO_3^{2-}); carboxylate ligands ($R^C CO_2^-$); nitrate (NO_3^-); sulfate (SO_4^{2-}) and sulphonate ($R^{S1}O_3^-$), wherein R^C is selected from alkyls, alkenyls, alkynyl, aryls, optionally substituted and optionally comprising one or more heteroatoms.

[0054] According to the organometallic complex of the invention, two, three or more of the ligands R_1 , R_2 , R_3 , R_4 , R_5 , R_6 may be present in the form of one or more single compounds, the single compound being

- a bi-, tri- or polydentate compound, and/or,
- an alkene, alkyne, cyclopentadienyl and/or an arene, the alkene, alkyne, cyclopentadienyl and/or an arene being optionally substituted and optionally comprising one or more heteroatoms.

[0055] Bi-, tri- or polydentate ligands generally comprise at least two donor ligands, such as N-, P-, O- or S-donor ligands as defined above, for example. A bi-, tri- or polydentate ligand may, furthermore, comprise different donor ligands within the same compound.

[0056] An example of a bidentate N-donor ligand is 2,2'-bipyridine, optionally substituted. An example of a bidentate O-donor ligand is oxalate. A well known example of a polydentate ligand is EDTA (ethylene diamine tetraacetic acid), which comprise 6 donor locations. In this case, all residues R_1 - R_6 would be provided by one single polydentate compound.

[0057] Two or more substituents of R_1 - R_6 may be present in the form of one or more single compounds being a alkene, alkyne, cyclopentadienyl, and/or arene.

[0058] In these cases, double bonds of the ligands may play a role in the formation of the bond with the central M, thus giving rise, if the ligands are formed by a cyclic compound, to sandwich or half-sandwich ("piano-stool") configurations, for example.

[0059] Accordingly, in a preferred embodiment, three ligands of the organometallic compound of the present invention selected from R_1 , R_2 , R_3 , R_4 , R_5 , and R_6 , are present in the form of an alkene, alkyne, cyclopentadienyl and/or arene, optionally substituted and optionally comprising one or more heteroatoms, optionally bound covalently to the bioactive compound.

[0060] Examples of linear alkenes that function as suitable ligands to M include alkene, propene, 1,3-butadiene.

[0061] Examples of cyclic alkenes that function as suitable ligands to M include cyclohexa-1,4-diene and cycloocta-1,5-diene.

[0062] Preferably, three ligands selected from R_1 - R_6 are formed by an organic molecule such as an arene. Preferably,

the selected ligands result in a pseudo-octahedral arrangement around a central M, although other geometries are also possible, e.g. pentagonal bipyramid, square pyramid, tetrahedral and square planar or intermediate structures thereof.


[0063] The compound of the invention may thus be a half-sandwich compound. Preferably, three ligands selected from R_1 - R_6 are formed by a cyclic alkene, cyclopentadienyl and/or by an arene, optionally substituted, and optionally comprising one or more heteroatoms. The at least one cyclic "tridentate" ligand may be a mono-, bi-, tri- or polycyclic compound. Preferably, it is monocyclic. Preferably, it is an arene. Arenes are aromatic hydrocarbons. They may be substituted and comprise one or more heteroatoms.

[0064] Examples of mono-cyclic arenes are benzene and cyclopentadienyl ($C_5H_5^-$). The later is considered by the present inventors being an arene, but is often mentioned explicitly for the sake of avoiding doubts. Examples of monocyclic arenes comprising at least one N-heteroatom are pyridine, pyrazine, pyrimidine, pyridazine, for example. Of course, other heteroatoms may be present in the arene besides or instead of N, such as those mentioned above.

[0065] Accordingly, the arene may be polycyclic. Examples of polycyclic arenes are pentalene, indene, naphthalene, azulene, and so forth. Examples of polycyclic arenes comprising N-heteroatoms are indolizine, +H-indole, 2H-isindole, 3H-indole, 1H-indazole, /H-purine, indoline, isoindoline, 4H-quinolizine, quinoline, isoquinoline, pteridine, phtalazine, naphthydrine, quinazoline, cinnoline. Of course, other heteroatoms may be present in the polycyclic arene besides or instead of N, such as those mentioned above.

[0066] Most preferably, the organometallic compound of the present invention comprises, as at least one ligand, at least one are selected, independently, from benzene and cyclopentadienyl, which are optionally substituted by alkyl, alkenyl, alkynyl or aryl residues, optionally further substituted and comprising one or more heteroatoms.

[0067] According to a preferred embodiment of the present invention, three adjacent ligands of the organometallic compound of the invention, selected from R_1 , R_2 , R_3 , R_4 , R_5 , R_6 are formed by an arene of formula (VI)



(VI),


in which R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} may, independently of each other, be the same or different, are each hydrogen, alkyl, alkenyl, alkynyl, or aryl, which are, if applicable, optionally substituted and which optionally comprise one or more heteroatoms, and in which two or more residues selected from R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} may be covalently linked with each other thus forming a bi- or tri-, or polycyclic system, and in which any of the residues R_8 - R_{13} is optionally bound to the bioactive compound.

[0068] Monocyclic examples of arenes according to the compound of formula (VI) without heteroatoms are benzene, methylbenzene, cymene, for example.

[0069] Monocyclic examples of arenes according to the compound of formula (VI) comprising any one or more residues selected from R_8 - R_{13} being an alkyl, alkenyl or alkynyl that is substituted and/or comprising at least one heteroatom, the remaining residues being hydrogens are benzyl alcohol, 2-phenylethanol, 3-phenylpropanol, 4-phenylbutanol, benzylamine, 2-phenylethanamine, 3-phenylpropanamine, 4-phenylbutanamine, 2-phenylethanaminium, N,N,-dimethyl-2-phenylethanamine, all of which may be substituted to be bound to the bioactive organic compound.

[0070] The residues R_8 - R_{13} may thus comprise charges, but are preferably non-charged.


[0071] One or more residues selected from R_8 - R_{13} may comprise heteroatoms with electron pairs or double bonds that are linked to the central M, thus forming one of the ligands R_1 - R_6 , which is not yet occupied. An example of a compound falling in this category is reproduced by formula (VII) below:



(VII),

in which the bioactive organic compound may be linked to any of residues R_5 or R_6 or to an optional further residue of the benzene ring.


[0072] Examples of bicyclic variants of compound (VI), in which two residues, R_{12} and R_{13} are covalently linked forming a bicyclic systems are indicated with formulae (VIII) and (IX) below



in which R_8 - R_{11} have the same meaning as above in formulae (VI).

[0073] Further substituents may be present on the cycle formed by R_{12} and R_{13} , which are not shown here.

[0074] According to a particularly preferred embodiment of the present invention, the compound of formula (VI) comprises the structure of formula (X):



in which,

A is a optionally substituted, benzyl or cyclopentadienyl,

B is an optional linking group, which may be selected from alkyl, alkenyl, alkynyl and aryl, which is optionally substituted, which optionally comprises one or more heteroatoms and has 0-15, preferably 1-8 carbon atoms;

X is a functional group of B or the cyclic system, through which the bioactive compound is bound to B or to the cyclic system.

[0075] Preferably, in (X), A is benzene.


[0076] Preferably, in (X), B is a C_1 - C_{16} , more preferably a C_2 - C_4 alkyl. Preferably the alkyl only comprises X as a substituent and is otherwise unsubstituted.

[0077] Preferably, X is selected from -O- and from NH-. Accordingly, X preferably represents ether or part of an ester or amide group, of which the carbonyl part, if applicable, may be part of the "bioactive compound" or of the alkylene. Preferably, the bioactive compound, in its released and/or original, active form, comprises a carboxy-group and is linked to the N-or O-heteroatom indicated as X in formula (X) by means of an amide bond or ester bond, respectively.

[0078] Accordingly, in case of an ester bond, a hydroxy group of B may be esterified with a carboxy group of the bioactive compound, or, vice versa, a hydroxy group of the bioactive compound may be esterified with a carboxy-group optionally present on B. Both alternatives allow for hydrolytic cleavage and release of the bioactive compound.

[0079] In the case of an amide bond, an amino group of B may be linked to a carboxy group of the bioactive compound, or, vice versa, an amine group of the bioactive compound may be esterified with a carboxy-group optionally present on B. Both alternatives allow for hydrolytic cleavage and release of the bioactive compound.

[0080] According to further, particularly preferred embodiments, the compound of formula (VI) comprises the structures of formula (XI) and (XII):



in which,

m is 1-10, preferably 2-8.

[0081] Preferably, -O- and -NH- represent ester and peptide bonds, respectively, as defined for X in formula (X) above. More preferably, the bioactive compound, in its released and/or original, active form, comprises a carboxy-group and is linked to the O- or the N-atom indicated in (XI) and (XII), respectively, by means of an ester and peptide, bond, respectively.

[0082] Most preferably, m in formula (XI) is 1 and in (XII) is 2.

[0083] According to a preferred embodiment, at least one of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 of the organometallic compound of the present invention is a residue suitable of increasing the solubility of the complex of formula (I) in water (H_2O). This residue is preferably selected specifically for this purpose. The bioactive organic compound may be linked to this residue. Generally, hydrocarbons having a high heteroatom:carbon ratio are suitable to increase the solubility.

[0084] Accordingly, at least one of the residues R_1 - R_6 has the purpose of increasing the solubility of the organometallic compound of the invention in water. Preferably, the at least one ligand has a hydrophilic group. Hydrophilic groups include (-OH), (=O), (-COOH), (-NH₂), (-NHR-), (-O-), (-SH₂), (-S-), (-SO₃-), for example, with R being optionally substituted alkyl, alkenyl or aryl.

[0085] For example, the at least one residue selected from R_1 - R_6 for increasing solubility in water may be a hydrocarbon comprising one or several groups capable of engaging in hydrogen bonding with water, such as, for example, hydroxy groups.

[0086] In a preferred embodiment of the organometallic compound of the present invention, at least one organometallic compound of claim 1, in which any of R_1 , R_2 , R_3 , R_4 , R_5 , R_6 has the formula (XIII).




in which,

R_{14} , R_{15} , R_{16} may be the same or different, are each C_1 - C_6 alkyl, aryl or substituted aryl, or R_{14} , R_{15} , R_{16} may together with the phosphorous atom form a cycloalkyl group, such group being optionally heterocyclic. Preferably, R_{14} , R_{15} , R_{16} together form a fused ring system comprising 1-5 heteroatoms, preferably N. This residue, if present, preferably increases the solubility of the organometallic compound of the present invention in water.

[0087] An example of a residue capable of increasing solubility is 1,3,5-triaza-7-phospha-adamantane (PTA), 3-methyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane (MePTA), 3,7-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane (DAPTA), which may be linked via the P-atom to the central M in the compound of the present invention. An example of a bivalent ligand capable of increasing solubility of the compound of the present invention in water is oxalate.

[0088] In a preferred embodiment the organometallic compound of the present invention comprises the structure (XIV) below,



in which,

A is an arene, optionally substituted and optionally comprising one or more heteroatoms;

R_{18} , R_{19} , R_{20} , are ligands of the central Ruthenium atom which are, independently of each other, selected from halogens and/or N-, O-, S-, or P- donor ligands;


R_{17} is an optional residue selected from an alkyl, alkenyl, alkynyl, aryl, optionally substituted and optionally comprising one or more heteroatoms;

whereby the at least one bioactive organic compound constitutes at least one selected from R_{17} - R_{20} , or is covalently linked to any of the R_{17} - R_{20} , with the proviso that residues selected from R_{17} - R_{20} , which constitute the bioactive organic compound, or which are covalently bound the bioactive organic compound, are not halogens.

[0089] Preferred A and residues R_{17} - R_{20} are as indicated above. Accordingly, the grouping of A- R_{17} may be selected from formulae (X) - (XII), and the residues R_{18} - R_{20} , may be selected from halogens mentioned above and from N-, O-,

S-, or P- donor ligands mentioned above, for example. If the bioactive compound is bound to one or more of R_{18} - R_{20} , this/these residue(s) may be selected from formulae (III)-(V), for example.

[0090] According to a still preferred embodiment, the organometallic compounds of the present invention comprise the structures (XV) below:



in which,

X is a halogen;

Y is a halogen or N-, P-, O- or S-donor ligand as defined above; and,

in which,

if the bioactive compound is bound to R_{17} , R_{17} corresponds to "B-X-" as defined in formula (X) above, and R_{18} is a N-, P-, O- or S-donor ligand as defined above; and/or,

if the bioactive compound is bound to R_{18} , or constitutes R_{18} , R_{18} is, respectively, a N-, P-, O- or S-donor ligand as defined above to which the bioactive compound is covalently bound; or R_{18} constitutes the bioactive compound, the bioactive compound comprising itself N-, P-, O- or S-atoms functioning as donors suitable to attach it to the central Ru.

[0091] Preferably, Y in (XV) is a halogen.

[0092] Preferably, R_{18} is P, R_{14} , R_{15} , R_{16} as defined above. Preferably, it is PTA.

[0093] Preferably, in (XV), the bioactive compound is linked to R_{17} by an amide or ester bond. For example, the benzene to which R_{17} is bound, R_{17} and the bioactive compound together may correspond to the structure illustrated in formulae (XI) and (XII), for example.

[0094] If the bioactive compound is linked to R_{18} , R_{18} and the bioactive compound preferably correspond to any of formula (II) - (V) above.

[0095] If the bioactive compound is linked to R_{18} , R_{18} is preferably a N-, P-, O-, or S-donor ligand, to which the bioactive compound is linked. Accordingly, R_{18} preferably carries a functional group to which the bioactive compound can be linked, such as amide and ester bonds, as mentioned above.

[0096] In this alternative case, with R_{18} being covalently bound to the bioactive compound, R_{17} may represent one or more residues such as R_8 - R_{13} in the compound of formula (VI) above.

[0097] The present invention also envisages the possibility that the organometallic compound comprises more than one bioactive compound covalently bound to a ligand selected from R_1 - R_6 . Accordingly, in the above example of formula (XV), both, R_{17} and R_{18} may be covalently bound to a bioactive organic compound, whereby R_{18} may, instead constitute the bioactive compound. The bioactive compounds may then be the same or different. If they are different, the second bioactive compound may have a biological activity different from the bioactive compound generally used in the context of the present invention. Preferably, however, if more than one bioactive compounds are present, all are useful in the treatment of cancer in general and/or metastasis in particular.

[0098] Many residues, and in particular ligands attached to M detailed above are indicated to be substituted. For the purpose of the present invention, substituents are preferably selected, independently from each other if there are more than one substituents, from alkyls, alkenyls, alkynyls, aryls, the alkyls, alkenyls, alkynyls, aryls, optionally comprising one or more heteroatoms, and functional groups such as, for example, imine-groups ($=NH$), amino groups ($-NH_2$), hydroxy groups ($-OH$), thiol groups ($-SH$), carbonyl groups ($=O$), thio groups ($=S$), carboxyl groups ($-COOH$), nitrile groups ($-C\equiv N$), nitro groups ($-NO_2$), any other functional group and, if applicable, charged derivatives (for example, if pH dependent) and salts of the functional groups, for example. The substituent comprising a functional group may be substituted at the functional group.

[0099] Substituents may be branched and/or be further substituted.

[0100] The present invention, for example in claim 1, refers to an alkene, alkyne, cyclopentadienyl, and/or arene. The alkene or alkyne is considered to be an unsaturated C_2 - C_{30} , more preferably a C_3 - C_{15} and most preferably C_4 - C_8 hydrocarbon. An alkene comprises at least one ($C=C$)-double bond, whereas the alkyne comprises at least one ($C\equiv C$)-triple bond. The arene is preferably an aromatic C_5 - C_{35} , more preferably C_6 - C_{20} , most preferably C_6 - C_{15} hydrocarbon.

The alkene or alkyne may also be cyclic, if there are ≥ 3 carbons. The alkene or alkyne may be branched, if there are ≥ 4 or ≥ 5 carbons, respectively. The alkene, alkyne, cyclopentadienyl and/or arene (or aryl radical) may further be substituted, as indicated above, and optionally comprising one or more heteroatoms.

[0101] Alkenyl, alkynyl and/or aryl residues or substituents, as mentioned, for example, with respect to nitrile and azo ligands, aromatic N-donor ligands, amine ligands, P-donor ligands, S-donor ligands, O-donor ligands, substituents of benzene or cyclopentadienyl ligands, substituents selected from R_8 - R_{13} , R_{17} in formula (III), substituents of the alkene, alkyne, cyclopentadienyl and/or arene mentioned in the above paragraph, substituents of an alkyl or with respect to substituents in general are radicals of the alkenes, alkynes, cyclopentadienyls and/or arenes as defined in the paragraph above.

[0102] Alkyl substituents, as mentioned, for example, with respect to substituents in general as defined above, N-donor ligands, in particular nitrile and azo ligands, aromatic N-donor ligands, amine ligands, substituents of B in formula (II), P-donor ligands, S-donor ligands, O-donor ligands, substituents of the alkene, alkyne and/or arene of claim 1, substituents of benzene and/or cyclopentadienyl, substituents R_8 - R_{13} of formulae (VI), substituents of B in formula (IX), substituents selected from R_{14} - R_{15} , R_{17} in formula (III), are preferably C_1 - C_{30} alkyls, more preferably C_2 - C_{25} alkyls, even more preferably C_4 - C_{10} alkyls. If the alkyl comprises more than 3 carbons, it may be cyclic or branched. Alkyls that are cyclic and branched are also encompassed, if the comprise more than 6 carbons. Alkyls may generally be further substituted.

[0103] A heteroatom, for the purpose of the present invention may be any heteroatom, but is preferably selected from B, Si, N, P, As, O, S, Se, T, and halogens. If several heteroms are present, they may be the same or different. More preferably, heteroatoms are selected from N, O, P, S, and halogens. If the heteroatom is present in a substituent, alkene, alkyne, cyclopentadienyl or arene, it may change the chemical class of the compound. For example, an O present in a linear alkyl results in an ether. For the purpose of the present invention, this example would be considered to be an alkyl comprising one O heteroatom.

[0104] The present invention provides an organometallic compound having a bioactive agent linked to at least one of the ligands of the central M.

[0105] Preferably, the bioactive organic compound is selected from inhibitors of resistance pathways and/or a pharmaceutically acceptable derivatives thereof. Inhibiting activity may, if it is not yet described, be assessed by specific, commercially obtainable or described assays. The skilled person is capable of using meaningful concentrations of an presumed inhibitor in such an assay. An example of an assay for testing Glutathione S-transferase inhibiting activity is mentioned in Example 6.

[0106] According to a preferred embodiment, the bioactive organic compound is an inhibitor of resistance selected from Glutathione S-transferase (GST) inhibitors, γ -Glutamyl Cysteine Synthetase (γ -GCS) inhibitors, multidrug resistance protein (MRP)/P-glycoproteins (PgP) inhibitors, inhibitors of cell signalling pathways.

[0107] Examples of inhibitors of Glutathione S-transferase inhibitors comprise, but not limited to, ethacrynic acid, peptidomimetics based on glutathione, p-chlorophenoxyisobutyrate, Gossypol, indomethacin, non-steroidal anti-inflammatory compounds based on ibuprofen and ketoprofen, misonidazole, Piroprost, Sulfasalazine, and their derivatives.

[0108] Examples of inhibitors of γ -Glutamyl Cysteine Synthetase comprise, but not limited to, sulfoxime-based compounds such as buthinone sulfoxime and methinone sulfoxime, S-sulfocysteine, S-sulfohomocysteine, cystamine, and their derivatives.

[0109] Examples of inhibitors of the multidrug resistance protein comprise, but not limited to, quinidine, vinblastine, terfemadine, tamoxifen, verapamil, cyclosporin, amitriptyline, progesterone, and their derivatives.

[0110] Examples of inhibitors of cell signaling pathways comprise, but not limited to, pleurotin, azelaic acid, bischloroethylnitrosourea, palmarumycin, and their derivatives.

[0111] Optically pure enantiomers, mixtures of enantiomers such as racemates, diastereomers, mixtures of diastereomers, diastereomeric racemates, mixtures of diastereomeric racemates, and the meso-form, as well as pharmaceutically acceptable salts, solvent complexes and morphological forms of the bioactive organic compounds are also encompassed by the present invention.

[0112] The expression pharmaceutical acceptable derivatives and derivatives also encompasses, but is not limited to, salts.

[0113] Salts comprise, for example, salts with inorganic acids or organic acids like hydrochloric or hydrobromic acid, sulphuric acid, phosphoric acid, citric acid, formic acid, acetic acid, maleic acid, tartaric acid, benzoic acid, methanesulfonic acid, p-toluenesulfonic acid, and the like that are non toxic to living organisms or in case the compound (I) is acidic in nature with an inorganic base like an alkali or earth alkali base, e.g. sodium hydroxide, potassium hydroxide, calcium hydroxide and the like.

[0114] According to the present invention, the bioactive compound constitutes a ligand of the central M selected from R_1 - R_6 , or is covalently bound to a ligand to the central M of the organometallic compound of the invention. If the bioactive compound is covalently bound to the ligands, the covalent bond can be a carbon-carbon alkyl, alkenyl or alkynyl bonds, or carbon-heteronuclear bonds such as amide ($-\text{CONH}-$), ester ($-\text{CO}_2-$), ether ($-\text{CH}_2\text{O}-$), thioether ($-\text{CH}_2\text{S}-$), amine

(-CH₂N-), imine (-CH=N-), phosphorous (-CH₂-P-). Preferably, the covalent bond is a carbon heteronuclear bond.

[0115] According to a preferred embodiment of the present invention, the organometallic compounds of the present invention are used as medicaments.

[0116] More particularly, according to a preferred embodiment, the organometallic compound according to any of the preceding claims in the treatment and/or prevention of cancer and/or metastasis. The effectiveness of the present compounds for the treatment of metastasis is remarkable. With many cancers, tumors may be removed surgically, with the occurrence of metastasis remaining the principle problem to which so far no convincing remedy has been found. Surprisingly, the compounds of the present invention are effective in overcoming the resistance of many cancer cells in metastasis to anti-cancer drugs. Thanks to the compounds of the present invention, anti-cancer drugs that are inefficient due to the onset of resistance against it, these very drugs may again be effectively employed. In addition, due to the increase of efficiency against cancer drugs, lower doses of the later may be applied, thus reducing the occurrence and severity of side effects.

[0117] Accordingly, according to a preferred embodiment, the organometallic compound according to the invention are useful for reducing resistance of cancers and/or metastasis against anti-cancer drugs.

[0118] According to a further embodiment, the present invention comprises a composition comprising an anti-cancer drug and the organometallic compound of any of the preceding claims for treating and/or preventing metastasis. The organometallic compound of the invention is most effective if administered together with an anti-cancer drug which may be conventional, and to which cancer cells have developed resistance, or a capable of developing resistance.

[0119] As becomes clear from the above, the principle of the present invention does not only encompass a single, specific compound but is more general. In particular, with respect to the different ligands bound to the central M, and to one of which the bioactive compound is covalently bound, a large variability is provided. The bioactive compound may also be directly attached to the central atom, increasing the variability. If the bioactive compound is bound to a ligand, it may be linked to ligands of different general structure and to ligands that are linked to the central M in different ways, for example as free-electron-pair-donors or as donors of π -bonds that are capable of complexing or associating to a transition metal. Preferably, however, the central M is Ruthenium (Ru), and the overall compound has a sandwich or half-sandwich structure.

[0120] Therefore, a high variability exists with respect to the ligands. For examples, the arene-part shown in exemplary ligands of formula (VII) and (VIII) may be selected from a large number of possible arenes that can be used in sandwich- or half-sandwich compounds. These arenes may, of course, be substituted, for example for improving the physico-chemical properties of the overall compound, such as solubility, or for improving effectiveness and reducing toxicity to normal cells. Such substituents are not shown here in all the detail, they may be selected at the discretion of the skilled person and are not essential for the general principle described herein.

[0121] The following examples illustrate the principle of the present invention on the basis of ethacrynic acid, which functions as the bioactive compound and which is shown to be effective in the treatment of metastasis. The examples also illustrate the structural variability of connecting the bioactive compound to ligands of different structures and being linked to the central M in different ways.

A. Examples 1-5: Synthesis of the Organometallic Compounds of the Invention

Example 1

Ethacrynic-(cyclohexa-1,4-dienyl)methylamide (1)

[0122] Ethacrynic acid (500 mg, 1.65 mmol) was refluxed in oxalyl chloride (5 ml) for 30 mins. Unreacted oxalyl chloride was removed *in vacuo* and dichloromethane (10 ml) was added to redissolve the residual colourless oil. (cyclohexa-1,4-dienyl)methylamine (109 mg, 1.00 mmol) and triethylamine (1 ml) was added sequentially and the reaction mixture was refluxed for a further 6 h. On completion, a dark brown solution was obtained. The reaction mixture was washed with 5% NaHCO₃ (50 ml), brine (2 x 50 ml) and dried *in vacuo*. The residual oil was separated on silica using 5:95 MeOH:dichloromethane as the eluent to yield a colourless oil which crystallizes on standing (yield : 250 mg, 63.3 %). ¹H NMR (CDCl₃, 400.13 MHz) 7.19 (d, 1H, Ar_{EA}-H, ³J_{HH} = 8.4 Hz), 6.87 (d, 1H, Ar_{EA}-H, ³J_{HH} = 8.4 Hz), 6.81 (b, 1H, -CH₂NH-), 5.95 (s, 1H, =CH₂), 5.64-5.71 (m, 3H, C=CH-C), 5.58 (s, 1H, =CH₂), 4.60 (s, 2H, -OCH₂CO₂-). 3.91 (d, 2H, -CH₂NH-, ³J_{HH} = 6.0 Hz), 2.83 (t, 2H, -CH₂NH-, ³J_{HH} = 6.0 Hz), 2.61-2.72 (m, 4H, =C-CH₂-C=), 2.47 (q, 2H, -CH₂CH), ³J_{HH} = 7.6 Hz), 1.15 (t, 3H, -CH₂CH₃, ³J_{HH} = 7.6 Hz).

[(Ethacrynic- η^6 :benzylamide)RuCl(μ -Cl)]₂ (2)

[0123] (1) (0.43 g, 3.32 mmol) was refluxed with RuCl₃·3H₂O (50 mg, 0.192 mmol) in degassed EtOH (25 ml) for 6 hours and left to stand at -4°C for 12 h, during which a reddish-orange precipitate separates from the dark green solution.

The precipitate was filtered, dissolved in dichloromethane and precipitated using diethyl ether to yield a light orange precipitate. The product was dried in vacuo (yield : 76.0 mg, 70.4 %) . ^1H NMR (CDCl_3 , 400.13 MHz) 8.11 (t, 1H, $-\text{CH}_2\text{NH}-$), 7.16 (d, 1H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4$ Hz), 6.90 (d, 1H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4$ Hz), 5.97 (s, 1H, $=\text{CH}_2$), 5.77 (t, 1H, $p\text{-Ar}_{\text{Ph}}\text{-H}$, $^3J_{\text{HH}} = 6.0$ Hz), 5.63 (s, 1H, $=\text{CH}_2$), 5.56 (dd, 2H, $m\text{-Ar}_{\text{Ph}}\text{-H}$, $^3J_{\text{HH}} = 6.0$, 6.0 Hz), 5.43 (d, 2H, $o\text{-Ar}_{\text{Ph}}\text{-H}$, $^3J_{\text{HH}} = 6.0$ Hz), 4.80 (s, 2H, $-\text{OCH}_2\text{CO}_2-$), 4.58 (d, 2H, $-\text{CH}_2\text{NH}-$, $^3J_{\text{HH}} = 6.0$ Hz), 2.47 (q, 2H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz), 1.15 (t, 3H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz). Anal. ($\text{C}_{40}\text{H}_{38}\text{Cl}_8\text{N}_2\text{O}_6\text{Ru}_2$) C 42.57, H 3.39, N 2.48, Found C 42.48, H 3.52, N 2.70.

(Ethacrynic- η^6 :benzylamide)Ru(PTA) Cl_2 (3)

[0124] (2) (28.4 mg, 0.025 mmol) was refluxed with PTA (9.3 mg, 0.059 mmol) in 1:1 MeOH/dichloromethane (15 ml) for 2 hours. The solvent was removed and the residual was recrystallised from dichloromethane/diethyl ether to yield an orange precipitate (yield : 31 mg, 85.3 %). ^1H NMR (CDCl_3 , 400.13 MHz) 7.87 (t, 1H, $-\text{CH}_2\text{NH}-$), 7.18 (d, 1H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4$ Hz), 6.90 (d, 1H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4$ Hz), 5.96 (s, 1H, $=\text{CH}_2$), 5.07 (t, 1H, $p\text{-Ar}_{\text{Ph}}\text{-H}$, $^3J_{\text{HH}} = 5.6$ Hz), 5.60 (s, 1H, $=\text{CH}_2$), 5.70 (m, 2H, $m\text{-Ar}_{\text{Ph}}\text{-H}$), 5.52 (d, 2H, $o\text{-Ar}_{\text{Ph}}\text{-H}$, $^3J_{\text{HH}} = 6.0$ Hz), 4.69 (s, 2H, $-\text{OCH}_2\text{CO}_2-$), 4.56 (d, 2H, $-\text{CH}_2\text{NH}-$, $^3J_{\text{HH}} = 6.0$ Hz), 4.51 (s, 6H, PTA- $\text{N-CH}_2\text{-N}$), 4.32 (s, 6H, PTA- $\text{P-CH}_2\text{-N}$), 2.47 (q, 2H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz), 1.15 (t, 3H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz). Anal. ($\text{C}_{26}\text{H}_{31}\text{Cl}_4\text{N}_4\text{O}_3\text{PRu}$) C 42.23, H 4.50, N 7.58, Found C 42.54, H 4.40, N 7.74.

Example 2

(Ethacrynic- η^6 :phenylethanoate)Ru(PTA) Cl_2 (4)

[0125] Ethacrynic acid (200 mg, 0.66 mmol), N,N-dicyclohexylcarbodiimide (200 mg, 0.98 mmol), N,N-dimethylaminopyridine (120 mg, 1.10 mmol) and (η^6 :phenylethanol)Ru(PTA) Cl_2 (90 mg, 0.20 mmol) was dissolved in dichloromethane (50 ml) and stirred for 96 h. The reaction mixture was filtered through celite to remove the urea precipitate and separated on silica gel using acetone. The product was triturated in diethyl ether and recrystallised from dichloromethane/diethyl ether to yield a brown precipitate (yield: 50 mg, 34.0%). ^1H NMR (CDCl_3 , 400.13 MHz) 7.07 (d, 1H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4$ Hz), 6.72 (d, 1H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4$ Hz), 6.00 (s, 1H, $=\text{CH}_2$), 5.62 (s, 1H, $=\text{CH}_2$), 5.48 (m, 2H, $m\text{-Ar}_{\text{Ph}}\text{-H}$), 5.22 (d, 2H, $o\text{-Ar}_{\text{Ph}}\text{-H}$, $^3J_{\text{HH}} = 5.6$ Hz), 5.13 (t, 1H, $p\text{-Ar}_{\text{Ph}}\text{-H}$, $^3J_{\text{HH}} = 5.2$ Hz), 4.81 (s, 2H, $-\text{OCH}_2\text{CO}_2-$), 4.55 (s, 6H, PTA- $\text{N-CH}_2\text{-N}$), 4.50 (t, 2H, $-\text{CO}_2\text{CH}_2\text{CH}_2-$, $^3J_{\text{HH}} = 6.0$ Hz), 4.33 (s, 6H, PTA- $\text{P-CH}_2\text{-N}$), 2.83 (t, 2H, $-\text{CO}_2\text{CH}_2\text{CH}_2-$, $^3J_{\text{HH}} = 6.0$ Hz), 2.47 (q, 2H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz), 1.16 (t, 3H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz). Anal. ($\text{C}_{27}\text{H}_{32}\text{Cl}_4\text{N}_3\text{O}_4\text{PRu}$) C 44.03, H 4.38, N 5.71, Found C 43.94, H 4.40, N 5.72.

Example 3

N-[2-(1H-imidazol-1-yl)propyl]ethacrynic-amide (5)

[0126] Ethacrynic acid (349 mg, 1.16 mmol) was refluxed in dichloromethane (20 ml) with an excess of oxalyl chloride (2 ml) for 1 h. Unreacted oxalyl chloride was removed under vacuum and the reaction mixture concentrated to yield ethacrynic acid chloride as a colourless oil. The acid chloride was taken up in dichloromethane (20 ml) and N-(amino-propyl)imidazole (500 mg, 4.00 mmol) was added. The reaction mixture was then refluxed for 2 h. 5% NaHCO_3 solution (25 ml) was added to quench the reaction and the aqueous phase was extracted with dichloromethane (3 x 25 ml). The organic phases were combined and washed with brine (2 x 25 ml) and dried over Na_2SO_4 . The solvent was removed and the product was separated on silica using 20% EtOH/80% CHCl_3 . The solvent was removed to yield a colourless oil (yield: 378 mg, 79.5%). ^1H NMR (CDCl_3 , 400.13 MHz) 7.52 7.07 6.96 (s, 3H, $\text{Ar}_{\text{imidazole}}\text{-H}$), 7.20 6.86 (d, 2H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4$ Hz), 6.82 (m, 1H, NH), 5.96 5.59 (s, 2H, $=\text{CH}_2$), 4.57 (s, 2H, $-\text{OCH}_2\text{CO}_2-$), 4.04 (t, 2H, $\text{Im-CH}_2\text{-CH}_2$), 3.42 (dt, 2H, $\text{CH}_2\text{-CH}_2\text{-NH}$), 2.47 (q, 2H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz), 2.10 (tt, 2H, $\text{CH}_2\text{-CH}_2\text{-CH}_2$), 1.16 (t, 3H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz).

(η^6 -cymene)Ru Cl_2 (ethacrynic-propylamide-imidazole) (6)

[0127] [(η^6 -cymene)RuCl($\mu\text{-Cl}$)] $_2$ (68 mg, 0.11 mmol) and (5) (94 mg, 0.23 mmol) was dissolved in dichloromethane (25 ml) and stirred for 12 h. The reaction mixture was concentrated and pentane was added to precipitate the product. The product was recrystallised from dichloromethane/pentane to yield an orange precipitate (yield: 144 mg, 90.3%). ^1H NMR (CDCl_3 , 400.13 MHz) 7.95 7.30 6.91 (s, 3H, imidazole-H), 7.19 6.90 (d, 2H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4$ Hz), 6.94 (t, 1H, N-H), 5.96 5.61 (s, 2H, $=\text{CH}_2$), 5.45 5.26 (dd, 4H, $\text{Ar}_{\text{Ph}}\text{-H}$, $^3J_{\text{HH}} = 6.0$ Hz), 4.60 (s, 2H, $-\text{OCH}_2\text{CO}_2-$), 3.89 (t, 2H, $-\text{CH}_2\text{CH}_2\text{-imidazole}$, $^3J_{\text{HH}} = 7.2$ Hz), 3.36 (q, 2H, $\text{NHCH}_2\text{CH}_2-$, $^3J_{\text{HH}} = 7.2$ Hz), 2.96 (septet, 1H, $-\text{CH}_3\text{CHCH}_3$, $^3J_{\text{HH}} = 8.0$ Hz), 2.47 (q, 2H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6$ Hz), 2.16 (s, 3H, $-\text{ArCH}_3$), 1.93 (m, 2H, $\text{NHCH}_2\text{CH}_2-$), 1.26 (d, 6H, $-\text{CH}(\text{CH}_3)_2$,

$^3J_{\text{HH}} = 8.0 \text{ Hz}$, 1.15 (t, 3H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6 \text{ Hz}$). Anal. ($\text{C}_{29}\text{H}_{35}\text{Cl}_4\text{N}_3\text{O}_3\text{Ru}$) C 48.67, H 4.93, N 5.88, Found C 48.75, H 4.95, N 5.79.

Example 4

$[(\eta^6\text{-cymene})\text{RuCl}(\text{PTA})(\text{imidazolium-ethacrynic amide})]\text{BF}_4^-$ (7)

[0128] $(\eta^6\text{-cymene})\text{Ru}(\text{PTA})\text{Cl}_2$ (96 mg, 0.21 mmol), NaBF_4 (92 mg, 0.84 mmol) and (5) (124 mg, 0.30 mmol) was suspended in methanol (15 ml) and refluxed for 2 h, during which a yellow solution was obtained. The reaction mixture was cooled to room temperature and the solvent removed. The residue was extracted with dichloromethane (2 x 25 ml) and filtered through celite. The dichloromethane extracts was concentrated and diethyl ether was added to precipitate the product. The product was recrystallised from dichloromethane/diethyl ether to yield a yellow precipitate (yield: 150 mg, 77.2%). ^1H NMR (CDCl_3 , 400.13 MHz) 8.28 7.42 7.04 (s, 3H, imidazole-H), 7.22 (t, 1H, N-H), 7.17 6.99 (d, 2H, $\text{Ar}_{\text{EA}}\text{-H}$, $^3J_{\text{HH}} = 8.4 \text{ Hz}$), 5.96 5.58 (s, 2H, $=\text{CH}_2$), 5.81 5.61 (dd, 4H, $\text{Ar}_{\text{Ph}}\text{-H}$), 4.72 (m, 2H, $-\text{OCH}_2\text{CO}_2^-$), 4.41 (m, 6H, $\text{PTA-N-CH}_2\text{-N}$), 4.10 (m, 6H, $\text{PTA-P-CH}_2\text{-N}$), 3.89 (t, 2H, $-\text{CH}_2\text{CH}_2\text{-imidazole}$, $^3J_{\text{HH}} = 7.2 \text{ Hz}$) 3.36 (q, 2H, $\text{NHCH}_2\text{CH}_2-$, $^3J_{\text{HH}} = 7.2 \text{ Hz}$), 2.96 (septet, 1H, $-\text{CH}_3\text{CHCH}_3$, $^3J_{\text{HH}} = 8.0 \text{ Hz}$), 2.47 (q, 2H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6 \text{ Hz}$), 2.16 (s, 3H, $-\text{ArCH}_3$), 1.93 (m, 2H, $\text{NHCH}_2\text{CH}_2-$), 1.26 (d, 6H, $-\text{CH}(\text{CH}_3)_2$, $^3J_{\text{HH}} = 8.0 \text{ Hz}$), 1.15 (t, 3H, $-\text{CH}_2\text{CH}_3$, $^3J_{\text{HH}} = 7.6 \text{ Hz}$). ESI-MS (CH_2Cl_2 , +ve mode) m/z 840 $[\text{M}]^+$. Anal. ($\text{C}_{35}\text{H}_{47}\text{BCl}_3\text{F}_4\text{N}_6\text{O}_3\text{PRu}$) C 45.45, H 5.12, N 9.09, Found C 45.60, H 5.15, N 9.07.

Example 5

$(\eta^6\text{-cymene})\text{Ru}(\text{PTA})(\text{C}_2\text{O}_4)$ (8)

[0129] $[(\eta^6\text{-cymene})\text{RuCl}(\mu\text{-Cl})_2]$ (196.8 mg, 0.322 mmol) and silver oxalate (240 mg, 0.797 mmol) were stirred in water for 12 h. The mixture was then filtered through celite to remove the AgCl precipitate. The solvent was removed under vacuum and the residue was redissolved in methanol (25 ml). PTA (120 mg, 0.764 mmol) was added and the reaction was stirred for 2 h. The solvent was reduced to ca. 5% of its original volume and diethyl ether (25 ml) was added. The slurry was cooled to 4°C for 12 h to complete precipitation of the product. The precipitate was filtered and recrystallised from methanol-diethyl ether to yield a light orange precipitate (yield : 285 mg, 89 %). ^1H NMR (D_2O , 400.13 MHz) 5.98 5.89 (dd, 4H, Ar-H), 4.57 (s, 6H, $\text{PTA-N-CH}_2\text{-N}$), 4.15 (s, 6H, $\text{PTA-P-CH}_2\text{-N}$), 2.61 (septet, 1H, $-\text{CH}(\text{CH}_3)_2$), 2.05 (s, 3H, $-\text{CH}_3$). 1.22 (d, $-\text{CH}(\text{CH}_3)_2$). $^{13}\text{C}\{^1\text{H}\}$ NMR (D_2O , 100.63 MHz) 166.2 ($-\text{CO}_2$), 105.0 97.7 87.3 86.8 (Ar-C), 70.7 ($\text{N-CH}_2\text{-N}$), 48.7 ($\text{P-CH}_2\text{-N}$). 30.8 ($-\text{ArCH}_3$), 21.5 ($-\text{CH}(\text{CH}_3)_2$), 17.3 ($-\text{CH}(\text{CH}_3)_2$). $^{31}\text{P}\{^1\text{H}\}$ NMR (D_2O , 400.13 MHz) -33.39. Anal. ($\text{C}_{18}\text{H}_{26}\text{N}_3\text{O}_4\text{PRu}\cdot 0.5\text{H}_2\text{O}$) C 44.08, H 5.55, N 8.57, Found C 44.24, H 5.58, N 8.69.

$(\eta^6\text{-cymene})\text{Ru}(\text{PTA})(\text{C}_6\text{H}_6\text{O}_4)$ (9)

[0130] $[(\eta^6\text{-cymene})\text{RuCl}(\mu\text{-Cl})_2]$ (228 mg, 0.373 mmol) and silver 1,1-cyclobutanedicarboxylate (300 mg, 0.838 mmol) were stirred in acetonitrile (50 ml) for 12 h, during which a yellow precipitate was formed. The solvent was removed and the residue was redissolved in methanol (25 ml). The mixture was filtered through celite to remove the AgCl precipitate. PTA (130 mg, 0.828 mmol) was added to the filtrate and the solution was stirred for 2 hours. The solvent was reduced to ca. 5% of its original volume and diethyl ether (25 ml) was added. The slurry was cooled to 4°C for 4 hours to complete precipitation of the product. The precipitate was filtered and recrystallised from dichloromethane-diethyl ether to yield an orange precipitate (yield : 288 mg, 72.2 %). ^1H NMR (CDCl_3 , 400.13 MHz) 5.54 5.43 (dd, 4H, Ar-H), 4.49 (s, 6H, $\text{PTA-N-CH}_2\text{-N}$), 4.15 (s, 6H, $\text{PTA-P-CH}_2\text{-N}$), 2.76 2.66 (t, 4H, $-\text{CH}_2(\text{CH}_2)_2$), 2.58 (septet, 1H, $-\text{CH}(\text{CH}_3)_2$), 2.02 (s, 3H, $-\text{CH}_3$), 1.94 (quintet, 2H, $-\text{CH}_2(\text{CH}_2)_2$), 1.24 (d, $-\text{CH}(\text{CH}_3)_2$). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 100.63 MHz) 178.7 ($-\text{CO}_2$), 102.5 96.1 87.9 85.3 (Ar-C), 72.9 ($\text{N-CH}_2\text{-N}$), 50.8 ($\text{P-CH}_2\text{-N}$), 30.9 ($-(\text{O}_2\text{C})\text{C}(\text{CH}_2)_2$), 30.9 ($\text{C}(\text{CH}_2)_{\text{top}}\text{CH}_2$), 30.9 ($\text{C}(\text{CH}_2)_{\text{bottom}}\text{CH}_2$), 30.9 ($-(\text{CH}_2)_2\text{CH}_2$), 30.9 ($-\text{ArCH}_3$), 22.5 ($-\text{CH}(\text{CH}_3)_2$), 17.3 ($-\text{CH}(\text{CH}_3)_2$). $^{31}\text{P}\{^1\text{H}\}$ NMR (CDCl_3 , 400.13 MHz) -30.16. Anal. ($\text{C}_{22}\text{H}_{32}\text{N}_3\text{O}_4\text{PRu}\cdot \text{H}_2\text{O}$) C 47.73, H 6.19, N 7.59, Found C 47.99, H 6.45, N 7.72.

B. Examples 6-9: In vitro and in vivo Biological Data

Example 6: Determination of Inhibition of Glutathione-S-Transferase Activity

[0131] Human A549 lung carcinoma cells, known to express high levels of Glutathione-S-Transferase, were routinely grown in flasks with DMEM medium containing 4.5 g/l glucose, 10% foetal calf serum (FCS) and antibiotics at 37°C and 6% CO_2 . When the cells are confluent, they are trypsinised and concentrated in a centrifuge at 4°C . The cells were then diluted in phosphate-buffer saline (PBS) containing protease inhibitor cocktail (final concentration of 1 $\mu\text{g/ml}$) and ho-

mogenised by repeatedly freezing in liquid nitrogen and thawing (4 cycles). The homogenised cell samples were centrifuged at 4°C and the supernatant, which is the cell lysates, was separated. The cell lysates are stored at -56 °C.

[0132] The Ru compounds are weighed, and dissolved in DMSO to 100 mM. They are diluted in water to 100 µM, such that the DMSO concentration did not exceed 0.1%. The cell lysates were exposed to the drug solutions for 90 mins. Control, containing the cell lysates with water/0.1% DMSO was also prepared.

[0133] The GST activity in the treated cell lysates was determined using the glutathione-CDNB (1-chloro-2,4-dinitrobenzene) assay. Glutathione and CDNB were dissolved in deionised water and ethanol respectively to make up a 100 mM solutions. A developing solution, containing 50 mM phosphate buffer solution (pH 6.5), was prepared and was added 1 % v/v, the 100 mM glutathione and 100 mM CDNB solutions such that their final concentrations are both 1 mM. The developing solution was added to a 96-well plate at 200 µl per well, followed by the treated cell lysates at 2 µl per well. The absorbance at 340 nm was monitored continuously for 5 mins at 15 s intervals. The average slope of the change in absorbance was determined as a fraction of the control as the percentage of residual GST activity.

Results:

[0134] Cells lysates treated with complexes with good GST-inhibition ability should exhibit low residual GST activity. The summary of the results are shown in Table 1 below:

Table 1: GST activity in lung carcinoma cells

Treatment	Residual GST activity (% control)
Ethacrynic acid	86.32 ± 13.5
(Ethacrynic-η ⁶ :phenylmethanamide)Ru(PTA)Cl ₂ (3)	63.57 ± 2.0
(Ethacrynic-η ⁶ :phenylethanol ester)Ru(PTA)Cl ₂ (4)	42.77 ± 13.9
(η ⁶ -cymene)RuCl ₂ (ethacrynic-propylamide-imidazole) (6)	23.70 ± 10.7
[(η ⁶ -cymene)RuCl(PTA)(ethacrynic-propylamide-imidazole)]BF ₄ ⁻ (7)	62.58 ± 8.8

Example 7: Determination of Cell Growth Proliferation Inhibition

[0135] The cells were routinely grown in flasks with DMEM medium containing 4.5 g/l glucose, 10% foetal calf serum (FCS) and antibiotics at 37°C and 6% CO₂. When the cell are confluent, they are trypsinised and seeded in 48-well plates as monolayers for 24 h.

[0136] For (3), (4), (5) and (7), the Ru compounds are weighed, and dissolved in DMSO to 100 mM. They are diluted in excess medium and diluted to 100 µM, such that the DMSO concentration did not exceed 0.12%. The 100 µM Ru complex solution was then serially diluted to make up 50 µM, 25 µM, 12.5 µM, 6.3 µM, 3.1 µM, 1.6 µM solutions. The media is removed from the cell plates and the drug solutions applied in triplicate. A set of control cells, with media containing 1.0% DMSO, was left on each plate. The cells were exposed to the drugs for 72 hours.

[0137] For RAPTA-C, (8) and (9), the Ru compounds are weighed, and dissolved directly in medium to 1600 µM, then serially diluted to make up 800 µM, 400 µM, 200 µM, 100 µM, 50 µM, 25 µM solutions. The media is removed from the cell plates and the drug solutions applied in triplicate. A set of control cells, with media containing 1.0% DMSO, was left on each plate. The cells were exposed to the drugs for 72 hours.

[0138] Cell viability was determined using the standard MTT assay protocol, which allows the quantification of the mitochondrial activity in metabolically active cells. MTT (final concentration 0.2 mg/ml) was added to the cells for 2 h, then the culture medium was aspirated and the violet formazan precipitate dissolved in 0.1 N HCl in 2-propanol. The optical density, which is directly proportional to number of surviving cells, was quantified at 540 nm using a multiwell plate reader and the fraction of surviving cells was calculated from the absorbance of untreated control cells.

Results:

[0139] Using the above protocol, the compounds were tested against human T47D and MCF-7 breast carcinoma, A549 lung carcinoma and HT-29 colon carcinoma cell lines (see Figure 7).

Table 2: Comparison of activity between RAPTA-C and Ru-ethacrynic acid derivatives

Complexes	IC ₅₀ (μM)			
	HT29	A549	T47D	MCF7
(η ⁶ -cymene)Ru(PTA)Cl ₂	436	1029	1063	>1600
Ethacrynic acid	73.5	50.7	7.73	66.0
(Ethacrynic-η ⁶ :phenylmethanamide)Ru(PTA)Cl ₂ (3)	50.7	32.3	2.91	19.9
(Ethacrynic-η ⁶ :phenylethanoate)Ru(PTA)Cl ₂ (4)	105.5	66.7	6.28	104.7
(η ⁶ -cymene)RuCl ₂ (ethacrynic-propylamide-imidazole)(6)	39.1	34.0	4.80	10.7
[(η ⁶ -cymene)RuCl(PTA)(ethacrynic-propylamide-imidazole)]BF ₄ ⁻ (7)	64.2	65.1	5.97	19.9

Table 3: Comparison of activity between RAPTA-C and RAPTA-C carboxylate derivatives

Complex	IC ₅₀ (μM)			
	HT29	A549	T47D	MCF7
(η ⁶ -cymene)Ru(PTA)Cl ₂	436	1029	1063	>1600
(η ⁶ -cymene)Ru(PTA)(C ₂ O ₄) (8)	267	1130	1174	>1600
(η ⁶ -cymene)Ru(PTA)(C ₆ H ₆ O ₄) (9)	265	1567	1088	>1600

Example 8: IC₅₀ values of compounds tested on Tumor cell lines

[0140] Tumor cell lines derived from human and mouse tumours listed below were stabilised for in vitro propagation.

Table 4: Tumor cell lines for in vitro study.

Identification	Tumor histotype
HT-29 ¹	Human colorectal carcinoma
MIA-PaCa-2 ²	Human pancreatic cancer
H460M ³	Human lung carcinoma/metastatic
A2780	Human ovarian carcinoma
A2780/CDDP ⁴	Human ovarian carcinoma resistant to Cisplatin
TLX5 ⁵	Mouse lymphoma

¹) Obtained from CRO, Aviano, Italy (Dr. P. Spessotto); ²) Obtained from Istituto Nazionale per lo Studio e la Cura dei Tumori, Milan (kindly supplied by Dr. D. Coradini); ³), as ²), but kindly supplied by Dr. G. Pratesi; ⁴) A2780cis is the variant of the parental A2780 resistant to Cisplatin (ECACC No. 93112517); ⁵) Originally obtained from Chester Beatty Institute, London, UK.

[0141] IC₅₀ values are determined as follows: Tumour cells are incubated in the appropriate complete medium at 37°C and under controlled atmosphere (5% CO₂). Test compounds are tested at doses in the range of 100 nM and 100 μM. Cytotoxicity is determined by the MTT test, by measuring cell viability as the cell metabolic capacity to transform the tetrazolium salt of MTT in the blue formazan, by mitochondrial dehydrogenases; the blue colour is read at 570 nm with a spectrophotometer (Alley MC, Scudiero DA, Monks A, Hursey ML, Czerwinski MJ, Fine DL, Abbott BJ, Mayo JG, Shoemaker RH, Boyd MR. Feasibility of drug screening with panels of human tumor cell lines using a microculture tetrazolium assay. Cancer Res 48: 598-601, 1988. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. J Immunol Methods 65: 55-63, 1983). Briefly, on day 0, in 96-well plates, 2-4000 cells/well/100 μl complete medium are plated. On day 1, test compounds are added at the test concentrations of 10⁻⁷ M; 3*10⁻⁷ M; 10⁻⁶ M; 3*10⁻⁶ M; 10⁻⁵ M; 3*10⁻⁵ M; 10⁻⁴ M. Each concentration is studied in triplicate. Cell viability, as determined by the MTT test is done on Day 4, after 3 days treatment. Each well is added with 10 μl/100 μl MTT (prepared previously by dissolving 5 mg/ml MTT in PBS, sterilized with filters at a cut-off of 0.22 μm and stored at

4°C). Plates are put in an incubator at 37°C for 4hrs, then medium is eliminated and each well is added with 200µl DMSO (Sigma Chemical Co., USA). Plates are read with a spectrophotometer (Spectra count Packard Bell, Meriden, CT, USA) at 570 nm wave length. IC₅₀ is calculated with the GraphPad Prism4.

[0142] The table below shows the IC₅₀ values for the tested compounds. The reported values are the most representative of two separate experiments. Test concentrations have been done by serial dilutions of each compound starting from a mother solution of 10⁻² M prepared by dissolving Compound 3, Compound 4, Ethacrynic acid and cisplatin in dimehtylsulfoxide, and RAPTA-T (the moiety of the test compounds without the Ethacrynic acid moiety) in sterile apirogen water.

Table 5: IC₅₀ values (µM)

	A2780	A2780 /CDDP	H460M	HT-29	MIA PaCa-2	TLX5
Comp. 3	7.1 ÷ 9.5	4.6 ÷ 53	2.5 ÷ 3.8	2.3 ÷ 3.8	14.7 ÷ 20.1	0.8 ÷ 3.0
Comp. 4	~ 30	36.7 ÷ 38.7	29.1 ÷ 31.5	30.0 ÷ 32.6	81.5 ÷ 92.7	3.6 ÷ 4.3
Ethacrynic acid	91.1 ÷ 94.8	84.3 ÷ 97.6	58.1 + 63.3	35.2 ÷ 44.0	> 100	1.3 ÷ 1.8
Cisplatin	10.5 ÷ 11.8	> 100	1.3 ÷ 3.0	4.9 ÷ 32.6	10.6 ÷ 12.9	0.02 ÷ 0.2
RAPTA-T	> 100	> 100	> 100	> 100	> 100	64.1 ÷ 92.9

[0143] It can be seen from the table above that both compounds 3 and 4 of the present invention are active, in particular also against the Cisaplatin-resistant cell line A2780/CDDP.

Example 9: In-vivo study

[0144] Tumour cells (TLX5, see above in Example 8) were derived from mouse tumours and stabilised for in vivo propagation. For the animal study, TLX5 lymphoma cells were injected in the peritoneal cavity of CBA brown-grey inbred mice (Harlan Nossan, San Giovanni al Natisone, Udine, Italy) at day 0. Test compounds were applied at day 3 at various doses, also by the intraperitoneal route, with each drug (test compound or reference drug) diluted in appropriate physiological solution and injected in a volume of 0.1 ml/10g body weight.

[0145] Results from the in vivo study are shown in the table below.

Table 6: Survival time in an in vivo TXL5 lymphoma model


	Mean survival time (days ± SD)	% increase vs Controls
Controls	10.0 ± 0.82	-
Compound 3 at 200 mg/kg	12.2 ± 1.79**	22
Cisplatin 8 mg/kg	11.8 ± 1.79*	18

[0146] As can be seen from the table above, control mice die 10 after tumor implant. The treatment with compound 3 raises the mean survival time to 12.2 days, corresponding to +22% versus control, a result statistically significant (** p=0.0089, Logrank test). Cisplatin, known to have severe side effects, was also effective at the dose indicated, although to a lesser extent (* p=0.0182).

[0147] While these results indicate that Compound 3 of the present invention is active *in vivo*, it is to be noted that the compound was administered as a suspension of the compound in carboxymethylcellulose. This fact has most probably reduced the bioavailability of the compound 3.

Claims

1. An organometallic compound of formula (XIV),



in which,


A is an arene, optionally substituted and optionally comprising one or more heteroatoms; R_{18} , R_{19} , R_{20} , are ligands of the central ruthenium atom which are, independently of each other, selected from halogens and/or N-, O-, S-, or P- donor ligands;

R_{17} is an optional residue selected from an alkyl, alkenyl, alkynyl, aryl, optionally substituted and optionally comprising one or more heteroatoms;

whereby at least one bioactive organic compound constitutes at least one selected from R_{17} - R_{20} , or is covalently linked to any of the R_{17} - R_{20} , with the proviso that residues selected from R_{17} - R_{20} , which constitute the bioactive organic compound, or which are covalently bound the bioactive organic compound, are not halogens; and

wherein said bioactive organic compound is selected from ethacrynic acid, peptidomimetics based on glutathione, p-chlorophenoxyisobutyrate, Gossypol, indomethacin, non-steroidal anti-inflammatory compounds based on ibuprofen and ketoprofen, misonidazole, Piriprost, Sulfasalazine, sulfoxime-based compounds such as buthionine sulfoxime and methionine sulfoxime, S-sulfocysteine, S-sulfohomocysteine, cystamine, quinidine, vinblastine, terfernadine, tamoxifen, verapamil, cyclosporin, amitriptyline, progesterone, pleurotin, azelaic acid, bis-chloroethylnitrosourea, or palmarumycin, and their pharmaceutical acceptable derivatives and salts.

2. The organometallic compound of claim 1, in which A is an arene of the formula (VI)




(VI),

in which R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} are, independently of each other, hydrogen, alkyl, alkenyl, alkynyl, aryl, optionally substituted and optionally comprising one or more heteroatoms, and

in which two or more residues selected from R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} may be covalently linked with each other thus forming a bi- or tri-, or polycyclic system, and in which any of the residues R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} optionally constitutes or is optionally bound to the bioactive compound.

3. The organometallic compound according to any one of the preceding claims, wherein the N-donor ligand comprises the structure of formula (II):



(II)

in which,


M is the transition metal Ru;

the dashed line represents bonding, as a monodentate ligand, to the transition metal Ru the circle represents a mono- or polycyclic system, comprising at least one N-heteroatom, indicated as N, which provides an electron pair enabling attachment to M, the cyclic system being optionally further substituted;

B is an optional linking group, which is selected from alkyl, alkenyl, alkynyl and aryl, is optionally substituted, and optionally comprises one or more heteroatoms and has 0-15 carbon atoms; and

X is a functional group of B or the cyclic system, through which the bioactive compound is bound to B or to the cyclic system.

4. The organometallic compound according to any one of the preceding claims, wherein the N-donor ligand comprises the structure of formula (III) below, to which the bioactive compound is linked by means of an amide bond:



(III)

in which,

M is the transition metal Ru;


the dashed line represents bonding, as a monodentate ligand, to the transition metal Ru;

the circle represents a mono- or polycyclic system comprising at least two N-heteroatoms;

n is 1-10, preferably 2-6, most preferably 3-4;

X is as defined above.

5. The organometallic compound according to any one of claims 1-2 wherein the P-donor ligand comprises the structure of formula (XIII),




in which,

$R_{14} R_{15} R_{16}$ may be the same or different, are each C1-C6 alkyl, alkenyl, alkynyl, aryl, optionally substituted and optionally comprising one or more heteroatoms, or $R_{14} R_{15} R_{16}$ may together with the phosphorous atom form a cycloalkyl group, such group being optionally heterocyclic.

6. The organometallic compound according to claim 5, wherein the P-donor ligand of formula (XIII) is a residue suitable of increasing the solubility of the organometallic compound of claim 1 or 2 in water.

7. The organometallic compound according to claim 2 comprises the structure of formula (X):




in which,

A is an optionally substituted, benzyl or cyclopentadienyl,

B is an optional linking group, which may be selected from alkyl, alkenyl, alkynyl and aryl, which is optionally substituted, which optionally comprises one or more heteroatoms and has 0-15, preferably 1-8 carbon atoms; and


X is a functional group of B or the cyclic system, through which the bioactive compound is bound to B or to the cyclic system.

8. The organometallic compound according to any one of claims 2 and 7 comprises the structure of formula (XI) or (XII):



in which,
m is 1-10.

9. The organometallic compounds according to claim 1 comprises the structure of formula (XV):



in which,

X is a halogen;

Y is a halogen or N-, P-, O- or S-donor ligand as defined above; and,

in which,

if the bioactive compound is bound to R_{17} , R_{17} corresponds to "B-X-" as defined in formula (X) above, and R_{18} is a N-, P-, O- or S-donor ligand as defined above; or,

if the bioactive compound is bound to R_{18} , or constitutes R_{18} , R_{18} is, respectively, a N-, P-, O- or S-donor ligand as defined above to which the bioactive compound is covalently bound; or R_{18} constitutes the bioactive compound, the bioactive compound comprising itself N-, P-, O- or S-atoms functioning as donors suitable to attach it to the central Ru.

10. The organometallic compound according to any of the preceding claims for use as a medicament.


11. The organometallic compound according to any of the preceding claims for use in the treatment and/or prevention

of cancer and/or metastasis.

12. The organometallic compound according to any of the preceding claims for use in reducing resistance of cancers and/or metastasis against anti-cancer drugs.
13. A composition comprising an anti-cancer drug and the organometallic compound of any of the preceding claims for use in treating and/or preventing metastasis.
14. Use of the organometallic compound according to any one of claims 1-12 for the manufacture of a medicament for treating and/or preventing metastasis.
15. Use of the composition according to claim 13 for the manufacture of a medicament for treating and/or preventing metastasis.

Patentansprüche

1. Metallorganische Verbindung der Formel (XIV),



in der

A für ein Aren steht, das gegebenenfalls substituiert ist und gegebenenfalls ein oder mehrere Heteroatome umfasst;


R₁₈, R₁₉, R₂₀ für Liganden des zentralen Rutheniumatoms stehen, die unabhängig voneinander aus Halogenen und/oder N-, O-, S- oder P-Donorliganden ausgewählt sind;

R₁₇ für einen fakultativen Rest steht, der aus einem Alkyl, Alkenyl, Alkynyl, Aryl, das gegebenenfalls substituiert ist und gegebenenfalls ein oder mehrere Heteroatome umfasst, ausgewählt ist;

wobei mindestens eine bioaktive organische Verbindung mindestens einen der Reste R₁₇-R₂₀ darstellt oder kovalent an einen beliebigen der Reste R₁₇-R₂₀ gebunden ist, mit der Maßgabe, dass aus R₁₇-R₂₀ ausgewählte Reste, die die bioaktive organische Verbindung darstellen oder die kovalent an die bioaktive organische Verbindung gebunden sind, nicht für Halogen stehen; und

wobei die bioaktive organische Verbindung aus Ethacrynsäure, Peptidomimetika auf Basis von Glutathion, p-Chlorphenoxyisobutyrat, Gossypol, Indomethacin, nichtsteroidalen Antiphlogistika auf Basis von Ibuprofen und Ketoprofen, Misonidazol, Piroprost, Sulfasalazin, S-Sulfocystein, S-Sulfohomocystein, Cystamin, Quinidin, Vinblastin, Terferadin, Tamoxifen, Verapamil, Cyclosporin, Amitriptylin, Progesteron, Pleurotin, Azelainsäure, Bischlormethylnitrosoharnstoff oder Palmarumycin und deren pharmazeutisch unbedenklichen Derivaten und Salzen ausgewählt ist.


2. Metallorganische Verbindung nach Anspruch 1, in der A für ein Aren der Formel (VI)



(VI)

steht, in der R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} unabhängig voneinander für Wasserstoff, Alkyl, Alkenyl, Alkynyl, Aryl, das gegebenenfalls substituiert ist und gegebenenfalls ein oder mehrere Heteroatome umfasst, stehen und in der zwei oder mehr aus R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} ausgewählte Reste unter Bildung eines bi- oder tri- oder polycyclischen Systems kovalent miteinander verknüpft sein können und in der ein beliebiger der Reste R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} gegebenenfalls die bioaktive Verbindung darstellt oder gegebenenfalls an die bioaktive Verbindung gebunden ist.

3. Metallorganische Verbindung nach einem der vorhergehenden Ansprüche, wobei der N-Donorligand die Struktur der Formel (II) umfasst:



(II)

in der

M für das Übergangsmetall Ru steht;


die gestrichelte Linie eine Bindung, als einzähniger Ligand, an das Übergangsmetall Ru darstellt;

der Kreis ein mono- oder polycyclisches System darstellt, das mindestens ein als N angegebenes N-Heteroatom umfasst, welches ein die Anbindung an M ermöglichendes Elektronenpaar bereitstellt, wobei das cyclische System gegebenenfalls weiter substituiert ist;

B für eine fakultative Brückengruppe steht, die aus Alkyl, Alkenyl, Alkynyl und Aryl ausgewählt ist, gegebenenfalls substituiert ist und gegebenenfalls ein oder mehrere Heteroatome umfasst und 0-15 Kohlenstoffatome aufweist; und

X für eine funktionelle Gruppe von B oder dem cyclischen System steht, über die die bioaktive Verbindung an B bzw. an das cyclische System gebunden ist.

4. Metallorganische Verbindung nach einem der vorhergehenden Ansprüche, wobei der N-Donorligand die Struktur der nachstehenden Formel (III) umfasst, mit der die bioaktive Verbindung über eine Amidbindung verknüpft ist:



(III)

in der

M für das Übergangsmetall Ru steht;

die gestrichelte Linie eine Bindung, als einzähniger Ligand, an das Übergangsmetall Ru darstellt;
der Kreis ein mono- oder polycyclisches System darstellt, das mindestens 2 N-Heteroatome umfasst;
n für 1-10, vorzugsweise 2-6, ganz besonders bevorzugt 3-4, steht; X wie oben definiert ist.

5. Metallorganische Verbindung nach einem der Ansprüche 1-2, wobei der P-Donorligand die Struktur der Formel (XIII) umfasst:




in der

R_{14} , R_{15} , R_{16} gleich oder verschieden sein können und jeweils für C1-C6-Alkyl, Alkenyl, Alkynyl, Aryl, das gegebenenfalls substituiert ist und gegebenenfalls ein oder mehrere Heteroatome umfasst, stehen oder R_{14} , R_{15} , R_{16} zusammen mit dem Phosphoratom eine Cycloalkylgruppe bilden können, wobei eine derartige Gruppe gegebenenfalls heterocyclisch ist.

6. Metallorganische Verbindung nach Anspruch 5, wobei es sich bei dem P-Donorliganden der Formel (XIII) um einen zur Erhöhung der Löslichkeit der metallorganischen Verbindung nach Anspruch 1 oder 2 in Wasser geeigneten Rest handelt.

7. Metallorganische Verbindung nach Anspruch 2, die die Struktur der Formel (X) umfasst:




in der

A für ein gegebenenfalls substituiertes Benzyl oder Cyclopentadienyl steht,

B für eine fakultative Brückengruppe steht, die aus Alkyl, Alkenyl, Alkynyl und Aryl ausgewählt ist, gegebenenfalls substituiert ist und gegebenenfalls ein oder mehrere Heteroatome umfasst und 0-15 und vorzugsweise 1-8 Kohlenstoffatome aufweist; und

X für eine funktionelle Gruppe von B oder dem cyclischen System steht, über die die bioaktive Verbindung an B bzw. an das cyclische System gebunden ist.


8. Metallorganische Verbindung nach einem der Ansprüche 2 und 7, die die Struktur der Formel (XI) oder (XII) umfasst:



in der

m für 1-10 steht.

9. Metallorganische Verbindung nach Anspruch 1, die die Struktur der Formel (XV) umfasst:



(XV)

in der

X für ein Halogen steht;

Y für ein Halogen oder einen N-, P-, O- oder S-Donorliganden gemäß obiger Definition steht; und
in der

dann, wenn die bioaktive Verbindung an R₁₇ gebunden ist, R₁₇ "B-X-" gemäß der Definition in obiger Formel (X) entspricht und R₁₈ für einen N-, P-, O- oder S-Donorliganden gemäß obiger Definition steht; oder
dann, wenn die bioaktive Verbindung an R₁₈ gebunden ist oder R₁₈ darstellt, R₁₈ für einen N-, P-, O- oder S-Donorliganden gemäß obiger Definition steht, an den die bioaktive Verbindung kovalent gebunden ist; bzw. R₁₈ die bioaktive Verbindung darstellt, wobei die bioaktive Verbindung selbst N-, P-, O- oder S-Atome umfasst, die als zur Verknüpfung der bioaktiven Verbindung an das zentrale Ru geeignete Donoren fungieren.

10. Metallorganische Verbindung nach einem der vorhergehenden Ansprüche zur Verwendung als Medikament.

11. Metallorganische Verbindung nach einem der vorhergehenden Ansprüche zur Verwendung bei der Behandlung und/oder Prävention von Krebs und/oder Metastasen.

12. Metallorganische Verbindung nach einem der vorhergehenden Ansprüche zur Verwendung bei der Verringerung der Resistenz von Krebserkrankungen und/oder Metastasen gegen Antikrebsmittel.


13. Zusammensetzung, umfassend ein Antikrebsmittel und die metallorganische Verbindung nach einem der vorhergehenden Ansprüche, zur Verwendung bei der Behandlung und/oder Prävention von Metastasen.

14. Verwendung der metallorganischen Verbindung nach einem der Ansprüche 1-12 zur Herstellung eines Medikaments zur Behandlung und/oder Prävention von Metastasen.

15. Verwendung der Zusammensetzung nach Anspruch 13 zur Herstellung eines Medikaments zur Behandlung und/oder Prävention von Metastasen.

Revendications

1. Composé organométallique de formule (XIV) :




(XIV)

dans laquelle

A est un arène, éventuellement substitué et comprenant éventuellement un ou plusieurs hétéroatomes ;
 R₁₈, R₁₉ et R₂₀ sont des ligands de l'atome de ruthénium central qui sont, indépendamment les uns des autres, choisis parmi les halogènes et/ou les ligands donneurs de N, O, S ou P ;
 R₁₇ est un résidu facultatif choisi parmi les résidus alkyle, alcényle, alcynyle et aryle, éventuellement substitués et comprenant éventuellement un ou plusieurs hétéroatomes ;
 au moins un composé organique bioactif constituant au moins un résidu choisi parmi R₁₇- R₂₀ ou étant lié de façon covalente à l'un quelconque de R₁₇ - R₂₀, à condition que les résidus choisis parmi R₁₇ - R₂₀, qui constituent le composé organique bioactif, ou qui sont liés de façon covalente au composé organique bioactif, ne soient pas des halogènes ; et ledit composé organique bioactif étant choisi parmi l'acide étacrynique, les peptidomimétiques basés sur la glutathione, le p-chlorophénoxyisobutyrate, le gossypol, l'indométhacine, les composés anti-inflammatoires non stéroïdiens basés sur l'ibuprofène et le kétoprofène, le misonidazole, le piroprost, la sulfasalazine, les composés à base de sulfoxime tels que le sulfoxime de buthinone et le sulfoxime de méthinone, la S-sulfocystéine, la S-sulfohomocystéine, la cystamine, la quinidine, la vinblastine, la terfénadine, le tamoxifène, le vérapamil, la cyclosporine, l'amitriptyline, la progestérone, la pleurotine, l'acide azélaïque, la bischloroéthylnitrosourée et la palmarumycine, et leurs dérivés et sels pharmaceutiquement acceptables.

2. Composé organométallique selon la revendication 1, dans lequel A est un arène de la formule (VI) :




(VI)

dans laquelle R₈, R₉, R₁₀, R₁₁, R₁₂ et R₁₃ sont, indépendamment les uns des autres, un hydrogène, un alkyle, un alcényle, un alcynyle ou un aryle, éventuellement substitué et comprenant éventuellement un ou plusieurs hétéroatomes, et

dans laquelle au moins deux résidus choisis parmi R₈, R₉, R₁₀, R₁₁, R₁₂ et R₁₃ peuvent être liés de façon covalente les uns avec les autres, formant ainsi un système bi-, tri- ou polycyclique, et dans laquelle l'un quelconque des résidus R₈, R₉, R₁₀, R₁₁, R₁₂ et R₁₃ constitue éventuellement ou est éventuellement lié au composé bioactif.

3. Composé organométallique selon l'une quelconque des revendications précédentes, dans lequel le ligand donneur de N comprend la structure de formule (II) :




(II)

dans laquelle

M est le métal de transition Ru ;
 la ligne en tirets représente une liaison, comme un ligand monodenté, avec le métal de transition Ru ;
 le cercle représente un système mono- ou polycyclique, comprenant au moins un hétéroatome N, indiqué par N, qui fournit une paire d'électrons permettant la fixation à M, le système cyclique étant éventuellement aussi substitué ;
 B est un groupe de liaison facultatif, qui est choisi parmi les groupes alkyle, alcényle, alcynyle et aryle, est éventuellement substitué, et comprend éventuellement un ou plusieurs hétéroatomes et a 0-15 atomes de carbone ; et
 X est un groupe fonctionnel de B ou du système cyclique, par lequel le composé bioactif est lié à B ou au

système cyclique.

4. Composé organométallique selon l'une quelconque des revendications précédentes, dans lequel le ligand donneur de N comprend la structure de formule (III) ci-dessous, à laquelle le composé bioactif est lié au moyen d'une liaison amide :



dans laquelle

M est le métal de transition Ru ;
la ligne en tirets représente une liaison, comme un ligand monodenté, avec le métal de transition Ru ;
le cercle représente un système mono- ou polycyclique comprenant au moins deux hétéroatomes N ;
n vaut 1-10, de préférence 2-6, idéalement 3-4 ;
X est tel que défini ci-dessus.


5. Composé organométallique selon l'une quelconque des revendications 1 et 2, dans lequel le ligand donneur de P comprend la structure de formule (XIII) :



dans laquelle

R_{14} , R_{15} et R_{16} peuvent être identiques ou différents, sont chacun un alkyle, alcényle, alcynyle ou aryle en C_1 - C_6 , éventuellement substitué et comprenant éventuellement un ou plusieurs hétéroatomes, ou bien R_{14} , R_{15} et R_{16} peuvent, conjointement avec l'atome de phosphore, former un groupe cycloalkyle, un tel groupe étant éventuellement hétérocyclique.


6. Composé organométallique selon la revendication 5, dans lequel le ligand donneur de P de formule (XIII) est un résidu approprié pour augmenter la solubilité du composé organométallique de la revendication 1 ou 2 dans l'eau.
7. Composé organométallique selon la revendication 2, qui comprend la structure de formule (X) :



dans laquelle


A est un benzyle ou cyclopentadiényle éventuellement substitué,
B est un groupe de liaison facultatif, qui peut être choisi parmi les groupes alkyle, alcényle, alcynyle et aryle, qui est éventuellement substitué, qui comprend éventuellement un ou plusieurs hétéroatomes et qui a 0-15, de préférence 1-8 atomes de carbone ; et
X est un groupe fonctionnel de B ou du système cyclique, par lequel le composé bioactif est lié à B ou au système cyclique.

8. Composé organométallique selon l'une quelconque des revendications 2 et 7, qui comprend la structure de formule (XI) ou (XII) :



Composé bioactif

(XI)




Composé bioactif

(XII)

dans lesquelles
m vaut 1-10.

9. Composé organométallique selon la revendication 1, qui comprend la structure de formule (XV) :



(XV)

dans laquelle

X est un halogène ;

Y est un halogène ou un ligand donneur de N, P, O ou S tel que défini ci-dessus ; et

dans lesquels

si le composé bioactif est lié à R₁₇, R₁₇ correspond à « B-X- » tel que défini dans la formule (X) ci-dessus, et R₁₈ est un ligand donneur de N, P, O ou S tel que défini ci-dessus ; ou

si le composé bioactif est lié à R₁₈, ou constitue R₁₈, R₁₈ est respectivement un ligand donneur de N, P, O ou S tel que défini ci-dessus auquel le composé bioactif est lié de façon covalente, ou R₁₈ constitue le composé bioactif, le composé bioactif comprenant lui-même des atomes N, P, O ou S fonctionnant comme donneurs appropriés pour le fixer au Ru central.

10. Composé organométallique selon l'une quelconque des revendications précédentes à utiliser comme un médicament.

11. Composé organométallique selon l'une quelconque des revendications précédentes à utiliser dans le traitement et/ou la prévention du cancer et/ou de la métastase.

12. Composé organométallique selon l'une quelconque des revendications précédentes à utiliser dans la réduction de la résistance de cancers et/ou de métastases contre des médicaments anticancéreux.

13. Composition comprenant un médicament anticancéreux et le composé organométallique de l'une quelconque des revendications précédentes à utiliser dans le traitement et/ou la prévention de la métastase.

14. Utilisation du composé organométallique selon l'une quelconque des revendications 1 à 12 pour la préparation d'un médicament destiné à traiter et/ou prévenir la métastase.

15. Utilisation de la composition selon la revendication 13 pour la préparation d'un médicament destiné à traiter et/ou prévenir la métastase.



Figure 1**Figure 2**







Figure 3**Figure 4****Figure 5**

Figure 6**Figure 7****Figure 8**

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4980473 A [0003]
- WO 01130790 A [0006]
- WO 02102572 A [0007]
- WO 06018649 A [0009]
- US 20060058270 A [0009]
- US 20050239765 A [0009]

Non-patent literature cited in the description

- **GUO et al.** *Inorganica Chimica Acta*, 1998, vol. 273, 1-7 [0004]
- *Chem. Rev.*, 1999, vol. 99, 251-253 [0004]
- Metal Compounds in Cancer Therapy. Chapman and Hall, 1994, 65-91 [0004]
- **DALE et al.** *Anti-Cancer Drug Design*, 1992, vol. 7, 3-14 [0005]
- **KRAMER et al.** *Chem Eur J*, 1996, vol. 2 (12), 1518-1526 [0006]
- **BENNETT et al.** *Canadian Journal of Chemistry*, 2001, vol. 79, 655-669 [0006]
- **ORO et al.** *J Chem Soc, Dalton Trans*, 1990, 1463 [0006]
- **CHEN et al.** *J. Am. Chem. Soc.*, 2002, vol. 124 (12), 3064 [0008]
- **MORRIS et al.** *J. Med. Chem.*, 2001, vol. 44, 3616-3621 [0008]
- **SERLI et al.** *Eur. J. Inorg. Chem.*, 2005, 3423 [0010]
- **ANG et al.** *J. Am. Chem. Soc.*, 2005, vol. 127 (5), 1382 [0011]
- **ALLEY MC ; SCUDIERO DA ; MONKS A ; HURSEY ML ; CZERWINSKI MJ ; FINE DL ; ABBOTT BJ ; MAYO JG ; SHOEMAKER RH ; BOYD MR.** Feasibility of drug screening with panels of human tumor cell lines using a microculture tetrazolium assay. *Cancer Res*, 1988, vol. 48, 598-601 [0141]
- **MOSMANN T.** Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*, 1983, vol. 65, 55-63 [0141]