

(19)



(11)

EP 2 062 897 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 22

(51) Int Cl.:
C07D 489/08 (2006.01)

(48) Corrigendum issued on:
05.12.2012 Bulletin 2012/49

(45) Date of publication and mention
of the grant of the patent:
19.09.2012 Bulletin 2012/38

(21) Application number: **08169873.0**

(22) Date of filing: **25.11.2008**

(54) **Demethylation of 14-hydroxy substituted alkaloid derivatives**

Demethylierung von 14-Hydroxysubstituierten Alkaloidderivaten

Déméthylation de dérivés d'alkaloïde 14-hydroxy substitués

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT
RO SE SI SK TR**

(30) Priority: **26.11.2007 EP 07121537**

(43) Date of publication of application:
27.05.2009 Bulletin 2009/22

(73) Proprietor: **Siegfried AG
4800 Zofingen (CH)**

(72) Inventors:

- **Lauterbach, Erik Heinz, Dr.
5034 Suhr (CH)**
- **Dinkel, Thomas
4802 Strengelbach (CH)**
- **Heller, Sabrina
4665 Oftringen (CH)**

- **Bertogg, Andreas, Dr.
8005, Zürich (CH)**

(74) Representative: **Isarpatent
Patent- und Rechtsanwälte
Postfach 44 01 51
80750 München (DE)**

(56) References cited:
WO-A-2006/084389 GB-A- 1 124 441

- **IJIMA I. ET AL.: "Studies in the (+) morphinan series. V. Synthesis and biological properties of (+) naloxone" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON; US, vol. 21, no. 4, 1 January 1978 (1978-01-01), pages 398-400, XP002428747 ISSN: 0022-2623**

EP 2 062 897 B9

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention is directed to a method for demethylating 14-hydroxy substituted alkaloid derivatives, in particular of 14-hydroxy-17-methyl-4,5-epoxymorphinan-6-on-derivatives. This is achieved by reacting a starting compound with a compound of general formula $R^1OOC-N=N-COOR^2$ in a suitable solvent.

Background of the Invention

[0002] Naltrexone is an opioid receptor antagonist used primarily in the management of alcohol dependence and opioid dependence. It is available on the market in form of its hydrochloric salt, i.e. naltrexone hydrochloride. Naltrexone and its active metabolite 6- β -naltrexol are competitive antagonists at μ - and κ -opioid receptors, and to a lesser extent at δ -opioid receptors. The plasma half-life of naltrexone is about 4 hours, for 6- β -naltrexol 13 hours.

[0003] One important step in the synthesis of naltrexone is a demethylation step for removing the methyl group from the nitrogen atom of the alkaloid molecule, as it is present for example in oxycodone and oxymorphone.

[0004] One way to achieve this demethylation step is disclosed in GB-1,124,441. Therein, a process of removing a methyl group from a tertiary nitrogen atom of an alkaloid molecule is disclosed, comprising reacting the methylated compound with a lower alkyl azodicarboxylate, wherein the lower alkyl group has from 1-8 carbon atoms, for a time sufficient to bring about a reaction and to provide a demethylated alkaloid derivative. However, the patent specification is only related to alkaloid compounds, which do not carry an OH-group at C14 of the alkaloid backbone. For example, GB-1,124,441 is silent on how to perform a demethylation on oxycodone and oxymorphone, respectively.

[0005] The paper titled "Reactions of Azodicarboxylic Esters with Amines" of S. HOSZTAFI, Österreichische Apotheker-Verlags-ges.m.b.H., Wien, 1987, describes the reactions of aliphatic and aromatic amines and alkaloids with azodicarboxylic esters.

[0006] Further documents are available which are related to the demethylation of alkaloids, for example, EP 0 295 783 discloses diethyl azodicarboxylate for demethylating alkaloids. On page 8 of the description, it is disclosed to use diethyl azodicarboxylate in acetonitrile for the N-dealkylation of a compound of the indicated formula. However, in the course of this reaction, a bridging group is introduced between 2N-atoms as it can be seen from formula (VII). Furthermore, EP 0 295 783 is not related to alkaloids having a 14-OH group.

[0007] The like, GB-1,179,479 discloses removing a methyl group by treatment with a di-lower alkyl azodicarboxylate, wherein the lower alkyl group has from 1-4 carbon atoms, followed by treatment of dilute mineral acid. This reaction, however, is disclosed in the context of general formula (I) as indicated in GB-1,179,479, which does not possess a 14-OH group.

[0008] In the prior art, there are also chemical processes disclosed, which allow the demethylation of alkaloid compounds which are carrying an OH-group at C14. In these methods, however, the OH group is being protected in order to avoid a reaction with the usual demethylation agents.

[0009] For example, oxycodone may be reacted with Ac_2O in order to achieve acetyloxycodone (having a protected OH-group) which is further converted by well-known demethylating agents ($BrCN/H_2SO_4$) to *noroxycodone* (which is demethylated).

[0010] The same step can be performed in the synthesis of *noroxymorphone*, starting from oxymorphone. There, oxymorphone is reacted with Ac_2O resulting in a protected OH-group at C14 (oxymorphone diacetate). After reacting with $BrCN$ and H_2SO_4 , *noroxymorphone* is yielded. The conversion by means of $BrCN$ is described in IJIMA et al., "Studies in the (+)-Morphinan Series. 5. Synthesis and Biological Properties of (+)-Naloxone", Journal of Medicinal Chemistry, 1978, Vol. 21, No. 4.

[0011] However, there is not any disclosure in the art for a process for demethylating alkaloid derivatives which are carrying the 14-OH group, which 14-OH group is not being protected during the reaction. This might be due to the fact that conventional demethylating agents, such as $BrCN$, do not function under these circumstances as proper demethylating agents: The free OH group at C14 is too close to the cyanide and can perform a 5-exo-dig-cyclization which would not allow a proper performance of the reaction. See in this connection, CURRIE, A.C.; NEWBOLD, G.T. et al., Roy. Coll. Sci., Technol., Glasgow, UK, Journal of the Chemical Society, abstracts (1961), 4693-4700.

[0012] Further, see also GB-975,601, published in 1964. Here, it is disclosed that 14-acetoxy-N-cyanonorcocodeine acetate may be produced by reacting 14-acetoxycodeine acetate with cyanogen bromide and may be converted to 14-hydroxynorcocodeine by means of lithium aluminium hydride.

[0013] Iijima et al "Studies in the (+) morphinan series. V. Synthesis and biological properties of (+) naloxone" JOURNAL OF MEDICINAL CHEMISTRY; AMERICAN CHEMICAL SOCIETY; WASHINGTON; US, vol. 21, no. 4, 1 January 1978 (1978-01-01) and WO 2006/084389 are directed to the conversion of 3, 14-dihydroxydihydromorphinone into 3, 14-dihydroxyhydronormorphinone. 3, 14-diacetoxydihydromorphinone is reacted with either $BrCN$ or ethylchloroformate.

[0014] There is no indication that azodicarboxylic acid dialkyl ester should be used as a reagent in this conversion reaction.

Summary of the invention

[0015] Therefore, there is a need remaining to provide a method of demethylating 14-OH substituted alkaloid derivatives, and in particular, oxycodone and oxymorphone without the need of protecting/deprotecting the 14-OH group. Furthermore, it is an object of the present invention to provide a method of demethylating alkaloid derivatives which is environment-friendly and allows a conversion of 14-OH-alkaloid derivatives to the respective nor-alkaloid-derivatives in a high yield.

[0016] These problems are solved by the subject-matter of the independent claim. Embodiments are set forth in the dependent claims.

[0017] By the present invention, for the first time, a way of demethylating alkaloid derivatives having a 14-OH group is provided without the need of protecting the OH group during the reaction. This, surprisingly, could be achieved by using azodicarboxylic acid dialkyl esters of general formula $R^1OOC-N=N-COOR^2$ in a suitable solvent. This is insofar surprising as the prior art teaches that the 14-OH group is too reactive for using the conventional demethylating agents as discussed above without having protected the 14-OH group before.

[0018] By the new process of demethylation, two additional steps can be avoided, i.e. protecting and deprotecting the 14-OH-group of the alkaloid derivative. Furthermore, by using azodicarboxylic acid dialkyl ester compounds as, for example, DIAD or DEAD, substances, which are potentially harmful to the environment (like BrCN) may be avoided.

[0019] In order to give an overview over the improvements achieved by the present invention, it is referred to the enclosed Figures 1 and 2 showing two conventional ways for the synthesis of naltrexone base. Here, oxycodone is converted to acetyloxycodone in order to provide a protective group for 14 OH, and, then, demethylation is achieved by means of BrCN in order to achieve *noroxycodone* or *noroxymorphone*, respectively.


[0020] Figure 3 describes the reaction of the present invention. Here, oxycodone is converted directly to *noroxycodone*, as it is also the case for oxymorphone to *noroxymorphone* (see Figure 4).

[0021] It could be shown by the inventors that the yield of this reaction step is quite high, for example, for the conversion of oxymorphone to *noroxymorphone* the yield is about 80-90%.


Detailed description of the invention

[0022] In particular, the present invention is directed to the following:

According to a first aspect, the invention is directed to a method of producing a compound of formula (I)



or a salt thereof, comprising
reacting a compound of formula (II)



15 with an azodicarboxylic acid dialkyl ester of general formula $R^1OOC-N=N-COOR^2$ in a suitable solvent wherein

X is selected from H, alkyl, silyl or acetyl;

20 R^1 and R^2 are independently selected from linear or branched substituted or unsubstituted alkyl, preferably C_1 - C_6 -alkyl, more preferably methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, pentyl, hexyl, benzyl and

wherein the bond between atoms 7 and 8 is a single or a double bond, in order to obtain the compound of formula (I).

25 **[0023]** Salts of the above compound of formula (I) include carboxylate salts and others that are within a reasonable benefit/risk ratio, pharmacologically effective and suitable for contact with the tissues of patients without undue toxicity, irritation, or allergic response. Representative salts include hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylale, mesylate, glucoheptonate, lactibionate, and laurylsulfonate. See

30 **[0024]** The hydrochloride salt is preferred.

[0025] The azodicarboxylic acid dialkyl ester of formula $R^1OOC-N=N-COOR^2$ preferably is selected from compounds, wherein R^1 and R^2 are each ethyl or isopropyl, respectively, termed diethyl azodicarboxylate (DEAD) and diisopropyl azodicarboxylate (DIAD), or mixtures thereof.

35 **[0026]** In the reaction mixture, these compounds preferably are used in an amount of 1.5 to 3.0 eq.

[0027] In a further embodiment, the solvent is an aprotic polar or dipolar solvent and is preferably selected from methanol, ethanol, acetone, toluene, dimethylformamide, N,N-dimethylacetamide, acetonitrile, acetic acid ethylester and methyl-tert-butylether. Other solvents which fall into this category are methylene chloride, chloroform, tetrahydrofuran, dioxane, 1,3-dimethyl-2-imidazolidinone, dimethylsulfoxide, nitromethane or hexamethylphosphoric triamide. The

40 starting material may be completely or partly dissolved within the aprotic polar or dipolar solvent.

[0028] The most preferred solvent is dimethylformamide.

[0029] The above demethylation reaction from a compound of formula (II) to a compound of formula (I) preferably is performed at a temperature in the range of room temperature (20°C) to 100°C , preferably in the range of from 30 - 90°C , more preferably in the range of from 40 - 80°C and most preferably, in the range of 50 - 70°C .

45 **[0030]** This temperature is maintained for at least one hour, preferably at least two hours, more preferably at least three hours and most preferably at least four hours.

[0031] In a further preferred embodiment, after reacting the compound of formula (II) with an azodicarboxylic acid dialkyl ester, the reaction solution is supplemented with 5,5-dimethylcyclohexane-1,3-dione (dimedone) or hydrazines and methanol.

50 **[0032]** The reason for using dimedone or hydrazines is that during the demethylation reaction an aminal is formed, which is hydrolyzed with methanol to yield the secondary amine (N-H) and formaldehyde-dimethylacetal (or formaldehyde following hydrolysis). In order to remove formaldehyde from the reaction mixture, a compound suitable for capturing the same will enhance the purity of the reaction. Any other compound suitable of capturing formaldehyde may be used in addition or in place of dimedone or a hydrazines.

55 **[0033]** Dimedone preferably is used in an amount of at least 2 Eq, for example 2-3.5 Eq regarding the presumed amount of formaldehyde (-dimethylacetal). An example of a preferred amount is about 3.0. Eq.

[0034] Preferably, the reaction solution is maintained at a temperature in the range of room temperature (20°C) to 100°C , preferably in the range of 30 - 80°C , more preferably in the range of 40 - 70°C over a time of 1-10 hours, preferably

2-5 hours after adding dimedone/hydrazines and methanol. A most preferred temperature range is 50-65°C.

[0035] In a further preferred embodiment, following reacting the compound of formula (II) and optionally reacting with dimedone and methanol, an acid is added to the reaction solution. The reason for adding an acid is to protonate the reaction product in order to bring it in the polar (water) phase. The remaining substances (for example DEAD or DIAD) are separated into the organic phase and, thus, removed.

[0036] In more detail, when the reaction of the compound of formula II with $R^1OOC-N=N-COOR^2$ is complete, an acid as mentioned above (for example hydrochloric acid), water and an organic solvent is added. The organic solvent is not restricted in its kind and is preferably methylenchloride. Thus, two phases arise, wherein the organic phase (for example methylenchloride) will incorporate the remaining amounts of $R^1OOC-N=N-COOR^2$ (DIAD or DEAD, for example) and formaldehyde/dimedone.

[0037] The aqueous phase in turn receives the reaction product as a hydrochloride, and, thus, is separating it from the reaction mixture.

[0038] Basically, any conceivable type of acid may be used for this purpose, however, hydrochloric acid turned out to be most promising. The hydrochloric acid preferably has a concentration of about 5% V/V.

[0039] In a preferred embodiment of the invention, the reaction solution further comprises at least one antioxidant. It was found that the addition an antioxydant can prevent the formation of oxidative by-products. In the inventive method, oxidative by-products can be formed, depending on the reaction conditions. The removal of these by-products might require additional purification steps. Especially difficult to remove are 2,2'-bis(14-hydroxy-14-methyl-4.5-epoxymorphine-6-one)-derivatives which can only be removed by tedious purification procedures. In preferred embodiments, the amount of antioxidants in the reaction mixture is 1 to 50 mol-%, 5 to 40 mol-% or 10 to 30 mol-%. The addition of antioxidants can thus significantly improve the purity of the reaction product and reduce the process steps and costs of the reaction.

[0040] Suitable antioxidants are ascorbic acid, citric acid, tartaric acid, polyhydroxy butyric acid (PHB) esters, butylated hydroxyanisol (BHA) and butylated hydroxytoluene (BHT), preferably 2,6-di-tert-butyl-4-methylphenol. Further suitable antioxidants are selected from the group consisting of organic acids and carboxylic acids, acid salts of amino acids, sodium metabisulphite, malic acid, isoascorbic acid, sodium sulphite, sodium bisulphate, tocopherol, water- and fat-soluble derivatives of tocopherol, sulphites, bisulphites and hydrogen sulphites, 2,6-di-t-butyl-alpha-dimethylamino-p-cresol, t-butylhydroquinone, di-t-amylhydroquinone, di-t-butylhydroquinone, butylhydroxytoluene, pyrocatechol, pyrogallol, propyl/gallate, and nordihydroguaiaretic acid, phosphoric acids, sorbic and benzoic acids, esters and derivatives thereof.

[0041] Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

[0042] The present invention now is described in more detail by means of Figures and Examples.

[0043] In the Figures, the following is shown:

Fig. 1 shows a reaction scheme showing one conventional way of the synthesis of naltrexone base. The demethylation step from acetyloxycodone to *noroxycodone* is done by means of conventional means ($BrCN/H_2SO_4$). The 14 OH-group is protected before the demethylation step is performed.

Fig. 2 shows a further reaction scheme showing one conventional way of the synthesis of naltrexone base. The demethylation step from acetyloxymorphone to *noroxymorphone* is done by means of conventional means ($BrCN/H_2SO_4$). The 14 OH-group is protected before the demethylation step is performed.

Fig. 3 shows a reaction scheme showing an example of the synthesis of the present invention. The demethylation step from acetyloxycodone to *noroxycodone* is done by means of DEAD or DIAD. The 14 OH-group is not protected before the demethylation step is performed.


Fig. 4 shows a reaction scheme showing a further example of the synthesis of the present invention. The demethylation step from acetyloxymorphone to *noroxymorphone* is done by means of DEAD or DIAD. The 14 OH-group the like is not protected before the demethylation step is performed.

Fig. 5 shows a comparison of a way of synthesis according to a conventional approach (1st synthesis) and according to one embodiment of the present invention (2nd synthesis).

Examples:

[0044] The following is an illustration of one way to carry out the invention. The Example is related to the demethylation of oxymorphone, however can also be performed for any conceivable compound reflected by formula (II).

[0045] N-demethylation with an azodicarboxylate:



INPUT:

[0046]

15

Amount in g	Name of material	mmol	Content in %
50.0	Oxycodone	154.8	93,3 ^{w/w}
233.0	Dimethyl formamide pure		
62.6	Diisopropylazodicarboxylat (DIAD)	309.6	
65.1	5,5-Dimethylcyclohexane-1,3-dione (Dimedone)	464.3	
19.8	Methanol pure	618.8	
610.0	Dichlormethane		
27.4	Hydrochloride acid approximately 32% technical	240.5	32
450.0	Deionised water		
26.2	Ammonia solution approximately 25%, pure	384.6	25
120.0	Acetone		
1664.1	Sum	---	---

20

25

30

OUTPUT:

[0047]

35

Amount in g	Name of material	mol	Content in %	1
42.6	Noroxycodone	130.3	87.9	287.31

40

PROCESS:

[0048]

1

50.0 g Oxycodone are solved at room temperature in
233.0 g dimethylformamide at room temperature and supplemented with
62.6 g diisopropylazodicarboxylate.

The solution is heated to 55°C and a yellow to red mixture is formed. The solution is stirred for 4 hours at this temperature.

2

Progress of the reaction is controlled by HPLC for example.

EP 2 062 897 B9

(continued)

3			To the reaction mixture
5	65.1	g	dimedone and
	19.8	g	methanol are added starting at 55°C.
4			The mixture is kept at a temperature of 60°C whereas viscosity drops. The reaction mixture is stirred for 4 hours at a temperature of 60°C.
5			The reaction mixture is kept at a temperature of 20°C and is supplemented with
15	460.0	g	dichlormethane,
	200.0	g	deionised water and
	27.4	g	hydrochloric acid 32% and is stirred for at least 5min.
			Two clear phases are formed: a redish organic and a yellow aqueous phase.
6			The aqueous phase is separated.
7			The aqueous phase is washed with
20	150.0	g	dichlormethane. The phases are separated.
8			To the aqueous phase
25	26.2	g	aqueuos ammonia solution (25% w/w) are added at 20°C under stirring. A suspension is formed.
30	[0049] The further steps in the procedure comprise purification as used in chemistry. The steps described illustrate a possible way.		
	PURIFICATION:		
35	[0050]		
9			The suspension is brought to a temperature of 15°C and stirred for at least two hours.
10			The suspension is vacuum-filtered, dried and the residue is slurried with
40	250.0	g	water at a temperature of 20°C.
11			The suspension is vacuum-filtered and dried well by aspiration.
45			The filter residue is slurried with acetone at a temperature of 20°C, vacuum-filtered and is dried well by aspiration.
12	100.0	g	
50			The filter residue is again washed with acetone and well dried by aspiration.
13	20.0	g	

55

EP 2 062 897 B9

(continued)

14

The product is dried in a vacuum drying oven at 60°C.

5

15

42.6 g product is yielded as a fawn solid.

[0051] As it can be derived from table 1, showing the influence of the solvents and the amount of DIAD on the yield of demethylated products, dimethylacetamide, dimethylformamide and mixtures of dimethylformamide/toluene brought about the highest reaction yield. Further, the amount of DIAD used preferably was in the range of 1.5 to 3.0 eq.

10

Table 1

15

Noroxycodone : Examining the influence of the solvent and eq DIAD at 50 °C. Data indicated as analysed by HPLC.

20

25

30

35

40

45

50

55

Solvent Solvent System	conc. of educt	eq DIAD	HPLC after	Educt	Product
Toluene	10%	1.6	2.5 h	82.4%	11.1%
			20 h	19.7%	61.4%
Toluene	10%	2.5	2.5 h	72.2%	26.2%
			20 h	1.7%	66.7%
Toluene	10%	4.1	2.5 h	57.3%	38.3%
			20 h	0.00%	79.8%
Acetone	10%	1.6	2.5 h	72.1%	25.0%
			20 h	11.0%	76.2%
Acetone	10%	2.5	2.5 h	52.0%	42.3%
			20 h	0.00%	86.0%
Acetone	10%	4.1	2.5 h	36.0%	56.1%
			20 h	1.2%	86.3%
Methanol	10%	1.6	2.5 h	47.1%	20.3%
			20 h	37.3%	8.5%
Methanol	10%	2.5	2.5 h	3.0%	41.6%
			20 h	1.5%	18.8%
Methanol	10%	4.1	2.5 h	2.5%	50.4%
			20 h	1.4%	20.0%
Ethanol	10%	1.6	2.5 h	36.3%	38.5%

EP 2 062 897 B9

(continued)

Noroxycodone : Examining the influence of the solvent and eq DIAD at 50 °C. Data indicated as analysed by HPLC.					
Solvent Solvent System	conc. of educt	eq DIAD	HPLC after	Educt	Product
			20 h	2.1%	31.8%
Ethanol	10%	2.5	2.5 h	13.4%	56.0%
			20 h	0.00%	26.0%
Ethanol	10%	4.1	2.5 h	8.3%	63.2%
MTBE	10%	3.0	16 h	26.5%	69.3%
DMF	17%	3.0	16 h	1.0%	86.3%
			42 h	1.6%	84.5%
DMAc	17%	3.0	16 h	0.7%	91.0%
			42 h	0.5%	92.5%
ACN	17%	3.0	16 h	1.0%	80.0%
EE	17%	3.0	16 h	3.5%	81.0%
Toluene	33%	2.0	16 h	4.0%	69.0%
Toluene	7%	2.0	16 h	28.0%	62.0%
Toluene	33%	4.0	16 h	1.0%	71.0%
			42 h	1.0%	58.0%
Toluene	7%	4.0	16 h	7.0%	79.0%
Toluene	20%	3.0	5d 20°C	2.0%	83.0%
DMF	17%	3.0	17 h	1.0%	79.0%
DMF	10%	3.0	17 h	1.0%	91.0%
DMF	28%	3.0	17 h	1.0%	85.0%
DMF	17%	2.0	17 h	1.0%	89.0%
1DMF	17%	1.5	17 h	1.0%	89.0%

EP 2 062 897 B9

(continued)

Noroxycodone : Examining the influence of the solvent and eq DIAD at 50 °C. Data indicated as analysed by HPLC.					
Solvent Solvent System	conc. of educt	eq DIAD	HPLC after	Educt	Product
DMF/Toluene 6/1 (w/w)	15%	3.0	17 h	1.0%	88.0%
DMF/Toluene 1/1 (w/w)	15%	3.0	17 h	1.0%	86.0%
DMF/Toluene 1/6 (w/w)	15%	3.0	17 h	2.0%	85.0%
DMF/Toluene 1/1 (w/w)	15%	1.5	16 h	0.0%	89.0%
DMF/Toluene 1/1 (w/w)	15%	2.0	16 h	1.0%	89.0%
DMF/Toluene 1/6 (w/w)	15%	1.5	16 h	11.0%	76.0%
DMF/Toluene 1/6 (w/w)	15%	2.0	16 h	3.0%	83.0%
DMF/Toluene 1/1 (w/w)	7%	1.5	16 h	4.0%	90.0%
DMF/Toluene 1/1 (w/w)	7%	2.0	16 h	0.0%	92.0%
DMF/Toluene 1/6 (w/w)	7%	1.5	16 h	23.0%	70.0%
DMF/Toluene 1/6 (w/w)	7%	2.0	16 h	12.0%	81.0%
DMF	15%	1.0	16 h	9.0%	83.0%
Toluene dry	7%	2.0	20 h	5.0%	69.0%
Toluene/water 95/5 (w/w)	7%	2.0	20 h	11.0%	62.0%
DMF/water 98/2 (w/w)	14%	2.0	16 h	0.7%	79.7%

Table 2

Noroxycodone : Examining the influence of the solvent and eq DEAD at 50 °C. Data indicated as analysed by HPLC.					
Solvent	conc. of educt	eq DEAD	HPLC after	Educt	Product
Toluene	10%	2.0	16 h	10.0%	64.0%

Table 3

Noroxymorphone : Examining the influence of the solvent and eq DIAD at 50 °C. Data indicated as analysed by HPLC.					
Solvent	conc. of educt	eq DIAD	HPLC after	Educt	Product
DMF	13%	2.0	17 h	4.6%	89.1%
DMF	13%	2.0	16 h	1.5%	73.2%


[0052] As shown in the table 4 below 2,2'-bisoroxymorphone is formed as an unwanted side-product in the transformation of oxymorphone to noroxymorphone following the method described above. The amount of the by-products depends on the reaction conditions. In the presence of 25 mol-% of butylated hydroxytoluene (BHT), the formation of by-products, especially of 2,2'-bisoroxymorphone, which is hard to remove otherwise, can be suppressed.

Addition of BHT	Reaction time	Noroxymorphone (reaction product)	2,2'-Bisoroxymorphone (impurity)
Yes	4h	92.6 % ^{*)}	0.0 % ^{*)}
No	4h	90.9 %	0.7 %
Yes	16h	92.7 %	0.0 %
No	16h	89.7 %	1.5 %

^{*)} =area-% with HPLC


Claims

1. A method of producing a compound of formula (I)



or a salt thereof, comprising:

reacting a compound of formula (II)



15 with an azodicarboxylic acid dialkyl ester of general formula $R^1OOC-N=N-COOR^2$ in a suitable solvent wherein X is selected from H, alkyl, silyl or acetyl;

R^1 and R^2 are independently selected from linear or branched substituted or unsubstituted alkyl; and wherein the bond between atoms 7 and 8 is single or a double bond.

- 20
2. The method according to claim 1, wherein R^1 and/or R^2 are independently selected from C_1 - C_6 -alkyl, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, pentyl, hexyl, benzyl or are independently selected from isopropyl and ethyl.
- 25
3. The method of claim 2, wherein R^1 and R^2 are identical.
4. The method of one or more of the preceding claims, wherein the solvent is an aprotic dipolar solvent.
5. The method of claim 4, wherein the solvent is selected from methanol, ethanol, acetone, toluene, dimethylformamide, N,N-dimethylacetamide, acetonitrile, acetic acid ethylester and methyl-tert-butylether.
- 30
6. The method of one or more of the preceding claims, wherein the reaction is performed at a temperature in the range of room temperature (20°C) to 100°C .
- 35
7. The method of claim 6, wherein the reaction is performed at a temperature in the range of from 30 - 90°C .
8. The method of claim 6 or 7, wherein the temperature is maintained for at least one hour.
9. The method of one or more of the preceding claims, wherein after reacting the compound of formula (II) with an azodicarboxylic acid dialkyl ester, the reaction solution is supplemented with 5,5-dimethylcyclohexane-1,3-dione (dimedone) and methanol or hydrazines and methanol.
- 40
10. The method of claim 9, wherein the reaction solution is maintained at a temperature in the range of room temperature (20°C) to 100°C .
- 45
11. The method of one or more of the preceding claims, wherein following reacting the compound of formula (II) and optionally reacting with dimedone/hydrazines and methanol, an acid is added to the reaction solution.
12. The method of claim 11, wherein the acid is a hydrochloric acid.
- 50
13. The method of one or more of the preceding claims, wherein the hydrochloric acid has a concentration of about 5% V/V.
14. The method of one or more of the preceding claims, wherein the reaction solution further comprises at least one antioxidant.
- 55
15. The method of one or more of the preceding claims, wherein the antioxidant is selected from the group consisting of ascorbic acid, citric acid, tartaric acid, polyhydroxy butyric acid (PHB) esters, butylated hydroxyanisol (BHA) and


butylated hydroxytoluene (BHT), or 2,6-di-tert-butyl-4-methylphenol.

Patentansprüche

5

1. Verfahren zum Herstellen einer Verbindung der Formel (I)

10




15

oder eines Salzes davon, wobei das Verfahren Folgendes umfasst:

20

Umsetzen einer Verbindung der Formel (II)

25



30

35

mit einem Azodicarbonsäure-dialkylester der allgemeinen Formel $R^1OOC-N=N-COOR^2$ in einem geeigneten Lösungsmittel, wobei X aus H, Alkyl, Silyl oder Acetyl ausgewählt wird; R^1 und R^2 unabhängig voneinander aus linearem oder verzweigtem substituierten oder unsubstituierten Alkyl ausgewählt werden; und wobei die Bindung zwischen Atomen 7 und 8 eine Einfach- oder eine Doppelbindung ist.

40

2. Verfahren nach Anspruch 1, wobei R^1 und/oder R^2 unabhängig voneinander ausgewählt werden aus C_1 - C_6 -Alkyl, Methyl, Ethyl, n-Propyl, Isopropyl, n-Butyl, sek.-Butyl, tert.-Butyl, Pentyl, Hexyl, Benzyl oder unabhängig voneinander aus Isopropyl und Ethyl ausgewählt werden.

45

3. Verfahren nach Anspruch 2, wobei R^1 und R^2 identisch sind.

4. Verfahren nach einem oder mehreren der vorangehenden Ansprüche, wobei das Lösungsmittel ein aprotisches dipolares Lösungsmittel ist.

50

5. Verfahren nach Anspruch 4, wobei das Lösungsmittel aus Methanol, Ethanol, Aceton, Toluol, Dimethylformamid, N,N-Dimethylacetamid, Acetonitril, Essigsäureethylester und Methyl-tert-butylether ausgewählt wird.

55


6. Verfahren nach einem oder mehreren der vorangehenden Ansprüche, wobei die Reaktion bei einer Temperatur im Bereich von Raumtemperatur (20 °C) bis 100 °C ausgeführt wird.

7. Verfahren nach Anspruch 6, wobei die Reaktion bei einer Temperatur im Bereich von 30 bis 90 °C ausgeführt wird.

8. Verfahren nach Anspruch 6 oder 7, wobei die Temperatur mindestens eine Stunde lang gehalten wird.
9. Verfahren nach einem oder mehreren der vorangehenden Ansprüche, wobei nach dem Umsetzen der Verbindung der Formel (II) mit einem Azodicarbonsäure-dialkylester die Reaktionslösung mit 5,5-Dimethylcyclohexan-1,3-dion (Dimedon) und Methanol oder Hydrazinen und Methanol ergänzt wird.
10. Verfahren nach Anspruch 9, wobei die Reaktionslösung bei einer Temperatur im Bereich von Raumtemperatur (20 °C) bis 100 °C gehalten wird.
11. Verfahren nach einem oder mehreren der vorangehenden Ansprüche, wobei im Anschluss an das Umsetzen der Verbindung der Formel (II) und das optionale Umsetzen mit Dimedon/Hydrazinen und Methanol eine Säure zu der Reaktionslösung gegeben wird.
12. Verfahren nach Anspruch 11, wobei die Säure eine Chlorwasserstoffsäure ist.
13. Verfahren nach einem oder mehreren der vorangehenden Ansprüche, wobei die Chlorwasserstoffsäure eine Konzentration von etwa 5 % V/V hat.
14. Verfahren nach einem oder mehreren der vorangehenden Ansprüche, wobei die Reaktionslösung des Weiteren mindestens ein Antioxidationsmittel umfasst.
15. Verfahren nach einem oder mehreren der vorangehenden Ansprüche, wobei das Antioxidationsmittel aus folgender Gruppe ausgewählt wird: Ascorbinsäuren, Citronensäure, Tartrinsäure, Polyhydroxybuttersäure (PHB)-Estern, butyliertem Hydroxyanisol (BHA) und butyliertem Hydroxytoluen (BHT) oder 2,6-Di-tert-butyl-4-methylphenol.


Revendications

1. Procédé de production d'un composé de formule (I)



ou d'un sel associé, comprenant :

la mise à réagir d'un composé de formule (II)



EP 2 062 897 B9

avec un acide azodicarboxylique ester de dialkyle de formule générale $R^1OOC-N=N-COOR^2$ dans un solvant adapté dans laquelle X est choisi parmi H, l'alkyle, le silyle ou l'acétyle ;
R¹ et R² sont choisis indépendamment parmi un alkyle substitué ou non substitué linéaire ou ramifié ; et dans laquelle la liaison entre les atomes 7 et 8 est une simple ou une double liaison.

- 5
2. Procédé selon la revendication 1, dans lequel R¹ et/ou R² sont choisis indépendamment parmi un alkyle en C₁ à C₆, le méthyle, l'éthyle, le n-propyle, l'iso-propyle, le n-butyle, le secbutyle, le tert-butyle, le pentyle, l'hexyle, le benzyle ou sont choisis indépendamment parmi l'isopropyle et l'éthyle.
- 10
3. Procédé selon la revendication 2, dans lequel R¹ et R² sont identiques.
4. Procédé selon l'une ou plusieurs des revendications précédentes, dans lequel le solvant est un solvant dipolaire aprotique.
- 15
5. Procédé selon la revendication 4, dans lequel le solvant est choisi parmi le méthanol, l'éthanol, l'acétone, le toluène, le diméthylformamide, le N,N-diméthylacétamide, l'acétonitrile, l'acide acétique éthylester et le méthyl-tert-butyléther.
- 20
6. Procédé selon l'une ou plusieurs des revendications précédentes, dans lequel la réaction est effectuée à une température située dans la plage de la température ambiante (20 °C) à 100 °C.
7. Procédé selon la revendication 6, dans lequel la réaction est effectuée à une température située dans la plage de 30 à 90 °C.
- 25
8. Procédé selon la revendication 6 ou 7, dans lequel la température est maintenue durant au moins une heure.
9. Procédé selon l'une ou plusieurs des revendications précédentes, dans lequel après la mise à réagir du composé de formule (II) avec un acide azodicarboxylique ester de dialkyle, la solution réactionnelle est complétée avec de la 5,5-diméthylcyclohexane-1,3-dione (dimédone) et du méthanol ou des hydrazines et du méthanol.
- 30
10. Procédé selon la revendication 9, dans lequel la solution réactionnelle est maintenue à une température située dans la plage de la température ambiante (20 °C) à 100 °C.
- 35
11. Procédé selon l'une ou plusieurs des revendications précédentes, dans lequel après la mise à réagir du composé de formule (II) et éventuellement la mise à réagir avec de la dimédone/hydrazines et du méthanol, un acide est ajouté à la solution réactionnelle.
12. Procédé selon la revendication 11, dans lequel l'acide est un acide chlorhydrique.
- 40
13. Procédé selon l'une ou plusieurs des revendications précédentes, dans lequel l'acide chlorhydrique a une concentration d'environ 5 % V/V.
14. Procédé selon l'une ou plusieurs des revendications précédentes, dans lequel la solution réactionnelle comprend au moins un antioxydant.
- 45
15. Procédé selon l'une ou plusieurs des revendications précédentes, dans lequel l'antioxydant est choisi dans le groupe constitué par l'acide ascorbique, l'acide citrique, l'acide tartrique, les esters d'acide polyhydroxybutyrique (PHB), l'hydroxyanisole butylé (HAB) et l'hydroxytoluène butylé (HTB), ou le 2,6-di-tert-butyl-4-méthylphénol.
- 50

55









Fig. 1



MeSO_3H ,
 methionine
 Andre, Dormoy &
 Heymes
 76%

Ac_2O
 Iijima, et al
 90%



standard
procedure

1. BrCN
 2. H_2SO_4

Iijima (J. Med. Chem.
 1978, 398)
 91% after recryst
 + 95%

Fig. 2




Fig. 3




Fig. 4







Fig. 5

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- GB 1124441 A [0004]
- EP 0295783 A [0006]
- GB 1179479 A [0007]
- GB 975601 A [0012]
- WO 2006084389 A [0013]

Non-patent literature cited in the description

- **S. HOSZTAFI.** Reactions of Azodicarboxylic Esters with Amines. Österreichische Apotheker-Verlagsges., 1987 [0005]
- **IJIMA et al.** Studies in the (+)-Morphinan Series. 5. Synthesis and Biological Properties of (+)-Naloxone. *Journal of Medicinal Chemistry*, 1978, vol. 21 (4) [0010]
- **CURRIE, A.C. ; NEWBOLD, G.T. et al.** Journal of the Chemical Society. Roy. Coll. Sci., Technol, 1961, 4693-4700 [0011]
- Studies in the (+) morphinan series. V. Synthesis and biological properties of (+) naloxone. **IJIMA et al.** JOURNAL OF MEDICINAL CHEMISTRY. AMERICAN CHEMICAL SOCIETY, 01 January 1978, vol. 21 [0013]
- **S. M. BERGE et al.** Pharmaceutical Salts. *J. Pharm. Sci., J.*, vol. 977 (66), 1-19 [0023]