



(11) **EP 2 125 007 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Claims EN 1-2, 29, 33-34

(51) Int Cl.:
A61K 39/085 ^(2006.01) **A61K 39/39** ^(2006.01)
A61K 31/4745 ^(2006.01) **A61P 11/06** ^(2006.01)

(48) Corrigendum issued on:
16.04.2014 Bulletin 2014/16

(86) International application number:
PCT/US2008/001631

(45) Date of publication and mention of the grant of the patent:
08.01.2014 Bulletin 2014/02

(87) International publication number:
WO 2008/115319 (25.09.2008 Gazette 2008/39)

(21) Application number: **08799591.6**

(22) Date of filing: **07.02.2008**

(54) **CONJUGATES OF SYNTHETIC TLR AGONISTS AND USES THEREFOR**

KONJUGATE SYNTHETISCHER TLR-AGONISTEN UND VERWENDUNG DAFÜR

CONJUGUÉS D'AGONISTES SYNTHÉTIQUES DE TLR ET UTILISATIONS DE CEUX-CI

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR

• **DANIELS, Gregory A.**
La Jolla, California 92093 (US)

(30) Priority: **07.02.2007 US 888699 P**

(74) Representative: **Graf von Stosch, Andreas et al**
Graf von Stosch
Patentanwalts-gesellschaft mbH
Prinzregentenstrasse 22
80538 München (DE)

(43) Date of publication of application:
02.12.2009 Bulletin 2009/49

(60) Divisional application:
12004181.9 / 2 510 946

(56) References cited:
WO-A-2005/060966 US-A1- 2005 004 144

(73) Proprietor: **The Regents of The University of California**
Oakland, CA 94607-5200 (US)

• **WILLE-REECE ULRIKE ET AL: "HIV Gag protein conjugated to a toll-like receptor 7/8 agonist improves the magnitude and quality of Th1 and CD8(+) T cell responses in nonhuman primates"**
PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol. 102, no. 42, October 2005 (2005-10), pages 15190-15194, XP002500309
ISSN: 0027-8424

(72) Inventors:
• **COTTAM, Howard B.**
Escondido, California 92027 (US)
• **CARSON, Dennis A.**
La Jolla, California 92037 (US)
• **WU, Christina C. N.**
Escondido, California 92026 (US)
• **WRASIDLO, Wolfgang**
La Jolla, California 92037 (US)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 125 007 B9

- JIN ET AL: "Synthesis and immunostimulatory activity of 8-substituted amino 9-benzyladenines as potent Toll-like receptor 7 agonists" **BIOORGANIC & MEDICINAL CHEMISTRY LETTERS**, PERGAMON, ELSEVIER SCIENCE, GB, vol. 16, no. 17, 1 September 2006 (2006-09-01), pages 4559-4563, XP005593685 ISSN: 0960-894X
- ZAKS KAREN ET AL: "Efficient immunization and cross-priming by vaccine adjuvants containing TLR3 or TLR9 agonists complexed to cationic liposomes" **JOURNAL OF IMMUNOLOGY, AMERICAN ASSOCIATION OF IMMUNOLOGISTS**, US, vol. 176, no. 12, 15 June 2006 (2006-06-15), pages 7335-7345, XP002485614 ISSN: 0022-1767
- WU CHRISTINA C N ET AL: "Immunotherapeutic activity of a conjugate of a Toll-like receptor 7 ligand" **PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA**, vol. 104, no. 10, March 2007 (2007-03), pages 3990-3995, XP002500310 ISSN: 0027-8424
- "Figure 12-8" In: David E. Metzler: "Biochemistry. The Chemical reactions of living cells", 1977, Academic Press, Inc., London

Description**Cross-Reference to Related Applications**

5 **[0001]** The present application claims benefit of the filing date of U.S. application Serial No. 60/888,699, filed on February 7, 2007.

Statement of Government Rights

10 **[0002]** The invention described herein was made with government support under Grant Number AI056453 awarded by the National Institutes of Health. The United States Government has certain rights in the invention.

Background

15 **[0003]** A great deal has been learned about the molecular basis of innate recognition of microbial pathogens in the last decade. It is generally accepted that many somatic cells express a range of pattern recognition receptors that detect potential pathogens independently of the adaptive immune system (see Janeway et al., *Annu. Rev. Immunol.*, 20:197 (2002)). These receptors are believed to interact with microbial components termed pathogen associated molecular patterns (PAMPs). Examples of PAMPs include peptidoglycans, lipoteichoic acids from gram-positive cell walls, the sugar
20 mannose (which is common in microbial carbohydrates but rare in humans), bacterial DNA, double-stranded RNA from viruses, and glucans from fungal cell walls. PAMPs generally meet certain criteria that include, (a) their expression by microbes but not their mammalian hosts, (b) conservation of structure across the wide range of pathogens, and (c) the capacity to stimulate innate immunity. Toll-like Receptors (TLRs) have been found to play a central role in the detection of PAMPs and in the early response to microbial infections (see Underhill et al., *Curr. Opin. Immunol.*, 14:103 (2002)).
25 Ten mammalian TLRs and a number of their agonists have been identified. For example, TLR7 and TLR9 recognize and respond to imiquimod and immunostimulatory CpG oligonucleotides (ISS-ODN), respectively. The synthetic immunomodulator R-848 (resiquimod) activates both TLR7 and TLR8. While TLR stimulation initiates a common signaling cascade (involving the adaptor protein MyD88, the transcription factor NF- κ B, and pro-inflammatory and effector cytokines), certain cell types tend to produce certain TLRs. For example, TLR7 and TLR9 are found predominantly on the
30 internal faces of endosomes in dendritic cells (DCs) and B lymphocytes (in humans; mouse macrophages express TLR7 and TLR9). TLR8, on the other hand, is found in human blood monocytes (see Homung et al., *J. Immunol.*, 168:4531 (2002)).

[0004] In this respect, the synthesis and immunostimulatory activity of 8-substituted amino 9-benzyladenines as potent Toll-like receptor 7 agonist has been disclosed (see Jin et al., *Bioorg Med Chem Lett.* 16: 4559-63 (2006)).

35 **[0005]** Beyond that, it has been also reported that conjugation of an HIV Gag protein to a TLR7/8 agonist improves the magnitude and quality of Th1 and CD8+ T cell responses in nonhuman primates (see Wille-Reece et al., *Proc Natl Acad Sci U S A*, 102 (42): 15190-4, (2005)).

[0006] Interferons (INFs) are also involved in the efficient induction of an immune response, especially after viral infection (Brassard et al., *J. Leukoc. Biol.*, 71 :568 (2002)). However, many viruses produce a variety of proteins that
40 block interferon production or action at various levels. Antagonism of interferon is believed to be part of a general strategy to evade innate, as well as adaptive immunity (see Levy et al., *Cytokine Growth Factor Rev.*, 12:143 (2001)). While TLR agonists may be sufficiently active for some methods of treatment, in some instances the microbial interferon antagonists could mitigate the adjuvant effects of synthetic TLR agonist.

[0007] A more specific response to microbial infections is based on active or passive immunization. If universal immunization is not considered cost-effective (or pharmaco-economically viable), identification of a population at-risk that
45 would benefit from immunoprophylaxis may be cost-effective, although identifying that population may be not straightforward. Nevertheless, there are some clearly defined at-risk populations for certain bacterial infections, such as staphylococcal infections, including dialysis patients, patients with ventriculoperitoneal shunts, patients at-risk of infective endocarditis, and residents of nursing homes, all of which suffer from chronic conditions that place them at a prolonged
50 increased risk from staphylococcal infections. Many of these patients are also at increased risk for acquiring healthcare-associated methicillin-resistant *Staphylococcus aureus* (HA-MRSA). Blocking colonization of *Staphylococcus*, however, may be more achievable than protecting against infection.

[0008] Passive immunoprophylaxis using either polyclonal antibodies (Capparelli et al., *Antimicrob. Agents Chemo.*, 49:4121 (2005)) or monoclonal antibodies (mAbs) (www.biosynexus.com/productcandidates.html) may provide immediate (although short-term) protection for patients who either cannot wait for a vaccine effect to occur or whose
55 immune systems are too compromised to mount a response to a vaccine. One potential indication for passive immunoprophylaxis is a hospital outbreak of MRSA-related infections. In such cases, exposed individuals may benefit from immediate prophylaxis, whereas individuals residing on the same ward or chronic care facility may benefit from active

immunization. Moreover, intensive care unit patients are potential beneficiaries of passive immunoprophylaxis, as each of them would likely acquire one or more risk factors for staphylococcal infections.

Summary of the Invention

5

[0009] The present application discloses conjugates of a synthetic TLR agonist linked via a stable covalent bond to a macromolecule and compositions having those conjugates, as well as methods of using the conjugates. The conjugates may include macromolecules directly linked to a synthetic TLR agonist, e.g., TLR7 or TLR9 agonists, or linked via a linker to the TLR agonist, for instance, linked via an amino group, a carboxy group or a succinamide group. For instance, the disclosed conjugates include a synthetic TLR agonist (pharmacophore) covalently bound to a macromolecule such as, for instance, a peptide, polypeptide, e.g., an antibody or an antigen binding fragment thereof, lipid, a polymer such as polyethylene glycol, a bead, such as a polystyrene bead, or dendrimer. The disclosed conjugates are broad-spectrum, long-lasting, and non-toxic synthetic immunostimulatory agents, which are useful for activating the immune system of a mammal, e.g., a human. In particular, the disclosed conjugates optimize the immune response while limiting undesirable systemic side effects associated with unconjugated TLR agonists.

10

15

[0010] The synthetic TLR agonist may help direct the conjugate to TLRs within the endosomes of target cells and enhance delivery of the macromolecule. In one aspect, the synthetic TLR agonist is specific for endosomal TLRs. In one aspect, the TLR agonist may be a TLR7, TLR8, TLR3, or TLR9 agonist. Moreover, the synthetic TLR agonist may enhance the response to the macromolecule (e.g., immune response). Likewise, the macromolecule may be useful for activating the immune system and/or may direct the conjugate to particular cells. Thus, the macromolecule, e.g., one with a primary amino group that is linked to a synthetic TLR agonist, may enhance the activity of the synthetic TLR agonist or have a separate desirable activity. For example, the macromolecule may enhance the activity of the TLR agonist by helping to direct the agonist to the TLR within the endosomes of target cells, by enhancing signal transduction induced by the TLR agonist, or by cross-linking the receptor, or any combination thereof. In one example, the macromolecule is a lipid which is spontaneously incorporated into liposomes. In one example, the macromolecule is a nanoparticle which has amine groups on its surface. Once coupled to a TLR agonist, the TLR agonist-nanoparticle conjugate may be of a size, for instance, about 100 nm, that may reside (be present) in endosomes.

20

25

[0011] Hospital acquired *Staphylococcus aureus* (SA) infections are a major cause of morbidity and mortality. However, vaccines are not used in acute settings because they take too long to act and they are not effective in immunocompromised patients. The invention provides a vaccine for the rapid vaccination of patients at-risk for gram-positive bacterial infections, e.g., SA infections, which employs Toll-like receptor-7 (TLR7) agonists and one or more antigens (immunogens) of a gram-positive bacteria. The use of the vaccines of the invention induces immunity in about 6 days, which provides for applications not amenable to standard vaccination protocol (e.g., acute care settings).

30

35

[0012] As disclosed herein, a composition comprising a gram-positive bacteria, *Bacillus anthracis* (BA), and a synthetic TLR7 agonist was prepared. The composition induced IL12 and IL6 secretion (indicative of activation of bone marrow derived macrophages (BMDM)) *in vitro* and protected mice versus subsequent, otherwise lethal intra-pulmonary BA challenge *in vivo*. In particular, the administration of a composition containing a TLR7 agonist conjugate and an immunogen (UC-IV199-albumin/irradiated BA spores) induced protective immunity to BA within 6 days. In contrast, injection of the animals with BA spores alone, or with BA plus a conventional adjuvant, i.e., cholera toxin (CT), did not protect the animals from a lethal challenger. The rapidity of a protective immune response in a naïve animal was unexpected.

40

[0013] The application thus discloses immunogenic compositions. In one aspect, an immunogenic composition includes a synthetic TLR agonist such as a TLR7 agonist, e.g., UC-IV199, coupled to a gram-positive bacterial cell, for instance, coupled to free amino groups on killed *Staphylococcus aureus*; a synthetic TLR agonist such as a TLR7 agonist coupled to a bacterial extract of isolated gram-positive bacterial antigens; a synthetic TLR agonist such as a TLR7 agonist coupled to isolated gram-positive bacterial protein, e.g., recombinant protein; or a synthetic TLR agonist such as a TLR7 agonist coupled to isolated gram-positive bacterial carbohydrates. For example, a synthetic TLR7 agonist may be coupled to bacterial carbohydrates using methods to attach *Staphylococcus aureus* polysaccharides to protein carriers (such as those used for tetanus toxoid). The killed bacterial preparation may be prepared using gamma irradiation, heat or chemical treatment. In another aspect, an immunogenic composition includes a synthetic TLR agonist such as a TLR7 agonist coupled to an adjuvant and a preparation comprising killed gram-positive bacterial cells; a synthetic TLR agonist such as a TLR7 agonist coupled to an adjuvant and a preparation comprising a gram-positive bacterial extract; or a synthetic TLR agonist such as a TLR7 agonist coupled to an adjuvant and a preparation comprising an isolated gram-positive bacterial antigen, e.g., recombinant protein. For example, the immunogenic composition may include UC-IV 199 coupled to albumin and a preparation with killed gram-positive bacteria, e.g., killed *Staphylococcus aureus*. In one aspect, an immunogenic composition includes a synthetic TLR7 agonist coupled to an adjuvant and a preparation comprising a recombinant gram-positive bacterial antigen, such as an isolated gram-positive bacterial protein or a peptide thereof, or isolated gram-positive bacterial carbohydrate. In one aspect, a single dose of the immunogenic composition may show very potent activity, e.g., provide protective immunity, in a short period of time, e.g., less than about 10 days.

45

50


55

[0014] In one aspect, a sterilized vaccine is administered to a subject from 0 to 7 days prior to hospitalization. In one aspect, the vaccine is administered intramuscularly. In one aspect, the vaccine is administered at a dosage between 10 μ g and 10 mg.

[0015] The use of conjugates of synthetic TLR agonists such as TLR7 agonists is advantageous as accessible and versatile chemistry permits conjugation to any antigen, and modifiable conjugates have defined stoichiometry. The conjugates are inexpensive to prepare and are potent, and so provide rapid protection, enabling use in acute settings such as trauma, burn, pre-surgery, or bio-terrorism.

[0016] The application includes the use of conjugates of a synthetic TLR agonist and a macromolecule, as well as TLR agonist conjugates and another molecule. The conjugate may be useful to prevent, inhibit or treat disorders including, but not limited to, allergic asthma, infectious diseases such as respiratory viral infections, e.g., those associated with influenza virus or respiratory syncytial virus (RSV) infection, lupus and other autoimmune diseases, and as a vaccine, e.g., for cancer or infectious diseases. In one aspect, a single dose of the conjugate may show very potent activity in stimulating the immune response. Moreover, because of the low toxicity of the conjugates, in some circumstances higher doses may be administered, e.g., systemically, while under other circumstances lower doses may be administered, e.g., due to localization of the conjugate. In one aspect, when administered at high doses, the synthetic TLR agonist conjugates may elicit an a composition for use in antagonistic response, and so may be useful to inhibit or treat asthma or autoimmune diseases. A first dose may elicit a hyperresponse which, in turn, suppresses the immune response, thereby avoiding inflammation. Thus, the use of higher doses and readministration may result in inhibition of an immune response.

[0017] In one embodiment, the invention provides a composition for use in a method to prevent or inhibit a gram-positive bacterial infection in a mammal, comprising administering to the mammal an effective amount of the composition comprising a bacterial antigen of a gram-positive bacteria and an amount of a compound having formula (IA):



wherein X^1 is -O-, -S-, or NR^c ;

wherein R^c is hydrogen, C_{1-10} alkyl, or substituted C_{1-10} alkyl, or R^c and R^1 taken together with the nitrogen atom can form a heterocyclic ring or a substituted heterocyclic ring, wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C_{1-6} alkyl, hydroxy C_{1-6} alkylene, C_{1-6} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkoxy C_{1-6} alkylene, amino, cyano, halogen, or aryl;

R^1 is hydrogen, (C_1-C_{10}) alkyl, substituted (C_1-C_{10}) alkyl, C_{6-10} aryl, or substituted C_{6-10} aryl, C_{5-9} heterocyclic, substituted C_{5-9} heterocyclic; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C_{1-6} alkyl, hydroxy C_{1-6} alkylene, C_{1-6} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkoxy C_{1-6} alkylene, amino, cyano, halogen, or aryl;

each R^2 is independently hydrogen, -OH, (C_1-C_6) alkyl, substituted (C_1-C_6) alkyl, (C_1-C_6) alkoxy, substituted (C_1-C_6) alkoxy, -C(O)- (C_1-C_6) alkyl (alkanoyl), substituted -C(O)- (C_1-C_6) alkyl, -C(O)- (C_6-C_{10}) aryl (aroyl), substituted -C(O)- (C_6-C_{10}) aryl, -C(O)OH (carboxyl), -C(O)O (C_1-C_6) alkyl (alkoxycarbonyl), substituted -C(O)O (C_1-C_6) alkyl, - NR^aR^b , -C(O)NR $^aR^b$ (carbamoyl), substituted -C(O)NR $^aR^b$, halo, nitro, or cyano; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C_{1-6} alkyl, hydroxy C_{1-6} alkylene, C_{1-6} alkoxy, C_{3-6} cycloalkyl, C_{1-6} alkoxy C_{1-6} alkylene, amino, cyano, halogen, or aryl;

each R^a and R^b is independently hydrogen, (C_1-C_6) alkyl, (C_3-C_8) cycloalkyl, (C_1-C_6) alkoxy, halo (C_1-C_6) alkyl, (C_3-C_8) cycloalkyl (C_1-C_6) alkyl, (C_1-C_6) alkanoyl, hydroxy (C_1-C_6) alkyl, aryl, aryl (C_1-C_6) alkyl, Het, Het (C_1-C_6) alkyl, or (C_1-C_6) alkoxycarbonyl;

X^2 is a bond or a linking group; n is 1, 2, 3, or 4; and R^3 is a macromolecule comprising a heterologous peptide, heterologous protein, heterologous lipid, bead, such as a polystyrene bead, heterologous nucleic acid molecule or dendrimer;


or a pharmaceutically acceptable salt thereof, including hydrates thereof.

[0018] In certain non-claimed embodiments, the definition of the group R^1 can be used interchangeably with the definition of the group R^1 for any other formula described herein.

[0019] Non-limiting examples of macromolecules or linkers therefore include not only an oxygen atom, a sulfur atom,

a nitrogen atom or a carbon atom (and appropriately appended hydrogen atoms when necessary to fill valences) but also macromolecules or linkers with side chains that increase solubility, such as, for example, groups containing morpholino, piperidino, pyrrolidino, or piperazino rings and the like; amino acids, polymers of amino acids (proteins or peptides), e.g., dipeptides or tripeptides, and the like; carbohydrates (polysaccharides), nucleotides such as, for example, PNA, RNA and DNA, and the like; polymers of organic materials, such as, for example, polyethylene glycol, polylactide and the like; monomeric and polymeric lipids; insoluble organic nanoparticles; non-toxic body substances such as, for example, cells, lipids, antigens such as, for example microbes, such as, for example, viruses, bacteria, fungi, and the like. The antigens can include inactivated whole organisms, or sub-components thereof and the like.

[0020] Also provided is a compound for use in a method to prevent or inhibit a gram-positive bacterial infection in a mammal, comprising administering to the mammal an effective amount of the compound having formula (IB):



wherein X¹ is -O-, -S-, or -NR^c;

wherein R^c is hydrogen, C₁₋₁₀alkyl, or substituted C₁₋₁₀alkyl, or R^c and R¹ taken together with the nitrogen atom can form a heterocyclic ring or a substituted heterocyclic ring, wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆-cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

R¹ is hydrogen, (C₁₋₁₀)alkyl, substituted (C₁₋₁₀)alkyl, C₆₋₁₀aryl, or substituted C₆₋₁₀aryl, C₅₋₉heterocyclic, substituted C₅₋₉heterocyclic; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆-cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

each R² is independently hydrogen, -OH, (C₁₋₆)alkyl, substituted (C₁₋₆)alkyl, (C₁₋₆)alkoxy, substituted (C₁₋₆)alkoxy, -C(O)-(C₁₋₆)alkyl (alkanoyl), substituted -C(O)-(C₁₋₆)alkyl, -C(O)-(C₆₋₁₀)aryl (aroyl), substituted -C(O)-(C₆₋₁₀)aryl, -C(O)OH (carboxyl), -C(O)O(C₁₋₆)alkyl (alkoxycarbonyl), substituted -C(O)O(C₁₋₆)alkyl, -NR^aR^b, -C(O)NR^aR^b (carbamoyl), substituted -C(O)NR^aR^b, halo, nitro, or cyano; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

each R^a and R^b is independently hydrogen, (C₁₋₆)alkyl, (C₃₋₈)cycloalkyl, (C₁₋₆)alkoxy, halo(C₁₋₆)alkyl, (C₃₋₈)cycloalkyl(C₁₋₆)alkyl, (C₁₋₆)alkanoyl, hydroxy(C₁₋₆)alkyl, aryl, aryl(C₁₋₆)alkyl, Het, Het (C₁₋₆)alkyl, or (C₁₋₆)alkoxycarbonyl;

X² is a bond or a linking group; n is 1, 2, 3, or 4; and R³ is the gram-positive bacteria, an isolated antigenic protein or peptide of the gram-positive bacteria, or an isolated polysaccharide of the gram-positive bacteria; or a pharmaceutically acceptable salt thereof, including hydrates thereof, wherein the gram-positive bacterium is not *Bacillus* or *Listeria*. In one embodiment, the gram-positive bacteria is a *Staphylococcus*.

In one embodiment, R³ is an isolated antigenic protein or peptide of *Staphylococcus* and that compound is administered with a preparation of killed *Staphylococcus*.

[0021] The application discloses a compound of the application for use in medical therapy (e.g., for prophylaxis of bacterial diseases such as in a vaccine). These compounds can also be used for biodefense, e., against *B. anthrax*.

[0022] Further provided are non-claimed compositions and a compound of the application for use in medical therapy, e.g., to treat asthma or viral infections or prevent viral infection, as well as the use of the conjugates for the manufacture of a medicament for the treatment of a TLR associated condition or symptom or one in which an augmented immune response or a suppressed immune response is indicated

[0023] In addition, the application also discloses a non-claimed pharmaceutical composition comprising at least one disclosed compound herein, or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable diluent or carrier, optionally in combination with a preparation of a selected gram-positive bacteria, e.g., a killed preparation or an extract, isolated protein of a selected gram-positive bacteria, or isolated carbohydrate (polysac-

charide) of a selected gram-positive bacteria.

[0024] The application includes the use of conjugates of TLR7 agonists and a macromolecule, e.g., one with a primary amino group, that enhances the activity of the agonist, e.g., albumin, or has a separate desirable activity, e.g., is an antigen of a gram-positive bacterium. The conjugates may include macromolecules directly linked to a TLR7 agonist, or linked via a linker to the TLR7 agonist. The conjugates may optimize the immune response while limiting undesirable systemic side effects of TLR7 agonists.

[0025] In one embodiment, the invention provides compound for use in a method for preventing or treating a gram-positive bacterial infection in a mammal, such as a human. The use includes administering to a mammal in need of such therapy, an effective amount of a compound of the application, or a pharmaceutically acceptable salt thereof, conjugated to an adjuvant or to at least one antigen of the gram-positive bacterium.

[0026] Also disclosed is a non-claimed method to identify conjugates useful to prevent, inhibit or treat a particular condition or symptom, e.g., by identifying the cytokine profile induced by the conjugate *in vitro* or *in vivo* or by identifying the endosome, e.g., early, middle or late, having the TLR for the TLR agonist. As different cells have different endosomes, the identification of endosome patterns in cells may allow for targeting of conjugates to specific cell types or improve access of conjugates to endosomes.

Brief Description of Figures

[0027]

Figure 1 shows a TLR agonist/alphagalactosyl-ceramide conjugate.

Figure 2 illustrates a UC-1V150 conjugate of G1 PAMAM with an ethylene diamine core.

Figure 3A illustrates a linker (SANH) for conjugating a macromolecule and a synthetic TLR agonist.

Figure 3B illustrates synthesis of UC-1V150.

Figure 3C illustrates conjugation of UC-1V-150 to MSA. 200 μ L of MSA (25 mg/mL) is mixed with 100 μ L of conjugation buffer (1M phosphate, pH=7.2) and 690 μ L of PBS. 844 μ g of SANH in 10 μ L of DMF (40-fold molar excess to MSA) is added to protein solution (Final concentration of NP in reaction mixture is 5 mg/mL). After gentle mixing, reaction is allowed to proceed at room temperature for 2 hours. To remove excess of SANH the reaction mixture is loaded on a microcon spin filter device (YM-3, Millipore) and concentrated to about 70 μ L. 460 μ g of UC-1V150 dissolved in 10 μ L of DMF was added to MSA modified with SANH and the reaction mixture was incubated at RT overnight. To remove excess UC-1V150 the reaction mixture was first concentrated to 50 μ L using a microcon spin filter device (Millipore: YM-3) and loaded on a G25 micro-spin column (GE Healthcare).

Figure 4 is a graphic illustration of the absorption profile (at about 350 nm) of a compound of formula I (an OVA/SANH/UC-1V150 conjugate).

Figure 5 illustrates the absorbance profile of a conjugation reaction of a synthetic TLR7 agonist, UC-1V150, to mouse serum albumin (MSA). The UC-1V150 to MSA ratio is approximately 5:1.

Figure 6 illustrates a TLR agonist/phospholipid conjugate.

Figure 7 shows the effect of UC-1V199/lipid administration at different doses and timing.

Figures 8A-B show that UC-1V199/lipid inhibits TLR7 and TLR2 signaling.

Figure 9 illustrates biphasic dose-response to an ultrapotent TLR7 agonist (a pico/nanomolar agonist and micro-molecular antagonist).

Figures 10A and B illustrate that the UC-1V150/MSA conjugates activate both murine bone marrow-derived macrophages (panel A) and human peripheral blood mononuclear cells (panel B). Cells were incubated with various concentrations of the conjugates from 0.5 nM to 10 μ M with BMDM or from 0.1 to 10 μ M with PBMC. Culture supernatants were harvested after 24 hours and cytokine levels were analyzed by Luminex.

Figures 11A, B, and C illustrate the *in vivo* efficacy of a TLR7 agonist conjugate. C57BL/6 mice were injected (i.v. via the tail vein) with various amounts of UC-1V150 (aldehyde-modified SM-360320) or UC-1V150/MSA per mouse. Serum samples were collected and cytokine levels were analyzed by Luminex. The effect from the unconjugated synthetic TLR7 agonist, SM-360320, lasted for only 2 hours whereas UC-1V150/MSA extended the effect to at least 6 hours.

Figure 12 shows sustained *in vivo* local activity of a UC-1V150/MSA conjugate without a systemic effect. C57BL/6 mice were anesthetized and administered (i.t.) with 3 nmol of UC-1V150/MSA. At the indicated time points, mice were sacrificed, and BALF and sera collected. The data were combined from two separate experiments with at least six mice per group. The results show the mean values \pm SEM.

Figure 13 shows cytokine induction in BMDM by irradiated anthrax spore-TLR7 agonist conjugate.

Figure 14 provides survival graphs after immunization of mice with UC-1V150/MSA and challenge with spores. A) Age matched female A/J mice were administered i.n. saline only or saline containing MSA (an amount equivalent to UC-1V150/MSA), UC-1 V 150 or UC-1 V 150/MSA at 0.75 nmole/mouse 1 day before *B. anthracis* infection, and

survival was assessed up to 13 days. B) Balb/c mice were administered i.n. saline or UC-1V150/MSA at 5 nmole/mouse 1 day before influenza virus infection, and survival followed for 21 days. In each model, Kaplan-Meier survival curves and log-rank tests were performed to determine significance. At least 8 mice were tested in each group. Figure 15 provides a graph of percent survival after a single dose of vaccine with TLR7 agonists and conjugates.

Figure 16 illustrates that protection against anthrax spore exposure depends on CD4+ cells.

Figure 17 shows a local cytokine profile in mice. C57BL/6 mice were administered i.t. with a UC-1V150/MSA conjugate or unconjugated UC-1V136 at 3 nmole or 500 nmole per mouse, respectively. BALF and sera were collected at the indicated time points and cytokine levels determined by multiplex immunoassay. Mean values from at least 3-5 mice per group are shown \pm SEM.

Figure 18 illustrates the absorbance spectrum for direct conjugation of SIV particles to UC-1V150.

Figure 19 illustrates the cytokine induction in BMDC by a conjugate of a synthetic TLR7 agonist and virus particles. Figures 20A-B illustrate the effects of a UC-1V150/inactivated SIV conjugate (panel A) or UC-1V150/OVA/ODN (panel B) on IL-12 production. Myeloid BMDC were incubated for 24 hours under various conditions with 0.1 μ g/mL as indicated. IL-12 levels in the cell supernatant were measured by ELISA.

Figure 21 is a graphic illustration of the stimulation of bone marrow derived dendritic cells (BMDC) with OVA/UC-1V150 or OVA/ODN (ODN= oligodeoxynucleotide).

Figure 22 is an illustration of the UV spectrum of a double-conjugate, (OVA/UC-1V150/ODN 1043).

Figure 23 is an illustration of the induction of IL-12 in BMDC using OVA/ODN/UC-1V150 conjugates. OVA/1043 and OVA/1018 are ODN conjugates.

Figure 24 illustrates conjugation of a synthetic TLR agonist to a lipid component of a liposome. Self-assembly of the TLR conjugate coupled via spacer-linker to a C-15 lipid, resulted in the formation of 100 nM nanoparticles. TLR agonist and an NHS-ester of the lipid were reacted in equimolar amounts in DMF and 1 equivalent of triethylamine for 6 hours. The reaction mixture was purified by preparing HPLC under isocratic conditions in 50:50 acetonitrile/water.

Figure 25 shows a schematic of a TLR agonist/liposome conjugate. Liposomes are formed with cholesterol:DOPE:DSPC:mPEG2000-DSPE:TLR-DSPE:BODIPY-DOPE 30:30:30:5:5:1.5; DSPE = distearoylphosphatidylethanolamine; DOPE = dioleoylphosphatidylethanolamine; BODIPY = 6-(((4-(4-difluoro-5-(2-thienyl)-4-bora-3a, 4a-diaza-s-indacene-3-yl)styryloxy)acetyl)amino)hexanamido-DOPE. Cholesterol:DOPE:DSPC:DSPE-TLRagonist:DSPE-mPEG (in 1:1:1:0.16:0.16 molar ratio) in chloroform were taken in 30 mL glass culture tubes, dried under a stream of nitrogen gas and vacuum-dessicated for a minimum of 6 h to remove any residual organic solvent. The dried lipid film was hydrated in sterile deionized water in a total volume of 1 mL for a minimum of 12 hours. Liposomes were vortexed for 2-3 minutes to remove any adhering lipid film and sonicated in a bath sonicator (ULTRASONIK 28X) for 2-3 minutes at room temperature to produce multilamellar vesicles (MLV). MLVs were then sonicated with a Ti-probe (using a Branson 450 sonifier at 100% duty cycle and 25 W output power) in an ice bath for 1-2 minutes to produce small unilamellar vesicles (SUVs) as indicated by the formation of a clear translucent solution. The solution was pressure filtered in sequence through 200 and then 100 nm nucleopore polycarbonate membranes to obtain liposome nanoparticles of 100 nm with a polydispersity factor of less than 0.1.

Figure 26 shows synthesis of lipid conjugate WW-109. 0.45 mg (1 μ mole) of IV-199 was added to 100 μ L of a 10 mM solution of DOPE in chloroform. To this solution was added 0.1 mg of triethylamine from a chloroform stock. The mixture was reacted at room temperature for 24 hours and the chloroform was rotavaped. The white solid residue was washed three times in 60%/methanol/hexane and centrifuged to obtain a white solid. The m/z by Mass spec was 1086 and the compound had a uv max absorption at 268 nm. Fatty acid moieties of various chain lengths can be used to prepare the analogous compounds, including C₁₄-C₂₂ carboxylic acids with one, two, three, or four sites of unsaturation, epoxidation, hydroxylation, or a combination thereof, at any feasible locations of the carboxylic acid carbon chain. In one specific embodiment, the fatty acid moieties are C₁₇ carboxylic acids with a site of unsaturation at C₈-C₉. In another specific embodiment, the fatty acid moieties are C₁₈ carboxylic acids with a site of unsaturation at C₉-C₁₀. The carboxylic acid moieties of each fatty acid moiety can be the same, or they can be different (see, e.g., Figure 6).

Figure 27 illustrates a schematic of a silica particle with TLR agonists covalently bonded thereto.

Figure 28 provides exemplary compounds for preparing conjugates of the invention or for use in the methods of the invention. Other conjugates include TLR agonists coupled to human serum albumin, e.g., HSA/UC-1V150, or DOPE/UC-1V199. UC-1X-51 increases TNF-alpha levels three fold (110 ng/mL)

Detailed Description of Invention

Definitions

[0028] As used herein, the term "antibody" refers to a protein having one or more polypeptides substantially encoded by immunoglobulin genes or fragments of immunoglobulin genes. The recognized immunoglobulin genes include the

kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as the myriad of immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

[0029] The basic immunoglobulin (antibody) structural unit is known to comprise a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kD) and one "heavy" chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (V_L) and variable heavy chain (V_H) refer to these light and heavy chains respectively.

[0030] Antibodies may exist as intact immunoglobulins, or as modifications in a variety of forms including, for example, FabFc₂, Fab, Fv, Fd, (Fab')₂, an Fv fragment containing only the light and heavy chain variable regions, a Fab or (Fab')₂ fragment containing the variable regions and parts of the constant regions, a single-chain antibody, e.g., scFv, CDR-grafted antibodies and the like. The heavy and light chain of a Fv may be derived from the same antibody or different antibodies thereby producing a chimeric Fv region. The antibody may be of animal (especially mouse or rat) or human origin or may be chimeric or humanized. As used herein the term "antibody" includes these various forms.

[0031] A composition is comprised of "substantially all" of a particular compound, or a particular form a compound (e.g., an isomer) when a composition comprises at least about 90%, and preferably at least about 95%, 99%, and 99.9%, of the particular composition on a weight basis. A composition comprises a "mixture" of compounds, or forms of the same compound, when each compound (e.g., isomer) represents at least about 10% of the composition on a weight basis. A purine analog of the invention, or a conjugate thereof, can be prepared as an acid salt or as a base salt, as well as in free acid or free base forms. In solution, certain of the compounds of the invention may exist as zwitterions, wherein counter ions are provided by the solvent molecules themselves, or from other ions dissolved or suspended in the solvent.

[0032] As used herein, the term "isolated" refers to *in vitro* preparation, isolation and/or purification of a nucleic acid molecule, a peptide or protein, or other molecule so that it is not associated with *in vivo* substances or is present in a form that is different than is found in nature. Thus, the term "isolated" when used in relation to a nucleic acid, as in "isolated oligonucleotide" or "isolated polynucleotide" refers to a nucleic acid sequence that is identified and separated from at least one contaminant with which it is ordinarily associated in its source. An isolated nucleic acid is present in a form or setting that is different from that in which it is found in nature. In contrast, non-isolated nucleic acids (e.g., DNA and RNA) are found in the state they exist in nature. For example, a given DNA sequence (e.g., a gene) is found on the host cell chromosome in proximity to neighboring genes; RNA sequences (e.g., a specific mRNA sequence encoding a specific protein), are found in the cell as a mixture with numerous other mRNAs that encode a multitude of proteins. Hence, with respect to an "isolated nucleic acid molecule", which includes a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, the "isolated nucleic acid molecule" (1) is not associated with all or a portion of a polynucleotide in which the "isolated nucleic acid molecule" is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence. The isolated nucleic acid molecule may be present in single-stranded or double-stranded form. When a nucleic acid molecule is to be utilized to express a protein, the nucleic acid contains at a minimum, the sense or coding strand (i.e., the nucleic acid may be single-stranded), but may contain both the sense and anti-sense strands (i.e., the nucleic acid may be double-stranded).

[0033] The term "amino acid" as used herein, comprises the residues of the natural amino acids (e.g., Ala, Arg, Asn, Asp, Cys, Glu, Gln, Gly, His, Hyl, Hyp, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val) in D or L form, as well as unnatural amino acids (e.g., phosphoserine, phosphothreonine, phosphotyrosine, hydroxyproline, gamma-carboxyglutamate; hippuric acid, octahydroindole-2-carboxylic acid, statine, 1,2,3,4,-tetrahydroisoquinoline-3-carboxylic acid, penicillamine, ornithine, citrulline, -methyl-alanine, para-benzoylphenylalanine, phenylglycine, propargylglycine, sarcosine, and tert-butylglycine). The term also comprises natural and unnatural amino acids bearing a conventional amino protecting group (e.g., acetyl or benzyloxycarbonyl), as well as natural and unnatural amino acids protected at the carboxy terminus (e.g., as a (C₁-C₆)alkyl, phenyl or benzyl ester or amide; or as an -methylbenzyl amide). Other suitable amino and carboxy protecting groups are known to those skilled in the art (see for example, T.W. Greene, *Protecting Groups In Organic Synthesis*; Wiley: New York, 1981, and references cited therein). For instance, an amino acid can be linked to the remainder of a compound of formula I through the carboxy terminus, the amino terminus, or through any other convenient point of attachment, such as, for example, through the sulfur of cysteine.

[0034] The term "toll-like receptor" (TLR) refers to a member of a family of receptors that bind to pathogen associated molecular patterns (PAMPs) and facilitate an immune response in a mammal. Ten mammalian TLRs are known, e.g., TLR1-10.

[0035] The term "toll-like receptor agonist" (TLR agonist) refers to a molecule that binds to a TLR. Synthetic TLR agonists are chemical compounds that are designed to bind to a TLR and activate the receptor. Exemplary synthetic TLR agonists provided herein include "TLR-7 agonist", "TLR" agonist", "TLR-3 agonist" and "TLR-9 agonist." TLR agonists include imiquimod, resiquimod, broprimine and loxoribine.

[0036] The term "nucleic acid" as used herein, refers to DNA, RNA, single-stranded, double-stranded, or more highly aggregated hybridization motifs, and any chemical modifications thereof. Modifications include, but are not limited to,

those providing chemical groups that incorporate additional charge, polarizability, hydrogen bonding, electrostatic interaction, and fluxionality to the nucleic acid ligand bases or to the nucleic acid ligand as a whole. Such modifications include, but are not limited to, peptide nucleic acids (PNAs), phosphodiester group modifications (e.g., phosphorothioates, methylphosphonates), 2'-position sugar modifications, 5-position pyrimidine modifications, 7-position purine modifications, 8-position purine modifications, 9-position purine modifications, modifications at exocyclic amines, substitution of 4-thiouridine, substitution of 5-bromo or 5-iodo-uracil; backbone modifications, methylations, unusual base-pairing combinations such as the isobases, isocytidine and isoguanidine and the like. Nucleic acids can also include non-natural bases, such as, for example, nitroindole. Modifications can also include 3' and 5' modifications such as capping with a BHQ, a fluorophore or another moiety.

[0037] As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

[0038] The pharmaceutically acceptable salts of the compounds useful in the present invention can be synthesized from the parent compound, which contains a basic or acidic moiety, by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, PA, p. 1418 (1985).

[0039] The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication commensurate with a reasonable benefit/risk ratio.

[0040] The following definitions are used, unless otherwise described: halo or halogen is fluoro, chloro, bromo, or iodo. Alkyl, alkoxy, alkenyl, alkynyl, etc. denote both straight and branched groups; but reference to an individual radical such as "propyl" embraces only the straight chain radical, a branched chain isomer such as "isopropyl" being specifically referred to. Aryl denotes a phenyl radical or an ortho-fused bicyclic carbocyclic radical having about nine to ten ring atoms in which at least one ring is aromatic. Het can be heteroaryl, which encompasses a radical attached via a ring carbon of a monocyclic aromatic ring containing five or six ring atoms consisting of carbon and one to four heteroatoms each selected from the group consisting of non-peroxide oxygen, sulfur, and N(X) wherein X is absent or is H, O, (C₁-C₄)alkyl, phenyl or benzyl, as well as a radical of an ortho-fused bicyclic heterocycle of about eight to ten ring atoms derived therefrom, particularly a benz-derivative or one derived by fusing a propylene, trimethylene, or tetramethylene diradical thereto.

[0041] It will be appreciated by those skilled in the art that compounds of the invention having a chiral center may exist in and be isolated in optically active and racemic forms. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties described herein, it being well known in the art how to prepare optically active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase) and how to determine agonist activity using the standard tests described herein; or using other similar tests which are well known in the art. It is also understood by those of skill in the art that the compounds described herein include their various tautomers, which can exist in various states of equilibrium with each other.

[0042] "Therapeutically effective amount" is intended to include an amount of a compound useful in the present invention or an amount of the combination of compounds claimed, e.g., to treat or prevent the disease or disorder, or to treat the symptoms of the disease or disorder, in a host. As used herein, "treating" or "treat" includes (i) preventing a pathologic condition from occurring (e.g. prophylaxis); (ii) inhibiting the pathologic condition or arresting its development; (iii) relieving the pathologic condition; and/or diminishing symptoms associated with the pathologic condition.

[0043] As used herein, the term "patient" refers to organisms to be treated by the methods of the present invention. Such organisms include, but are not limited to, mammals such as humans. In the context of the invention, the term "subject" generally refers to an individual who will receive or who has received treatment (e.g., administration of a compound of the invention).

[0044] "Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive

isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent. Only stable compounds are contemplated by the present invention.

The TLR Agonists and Conjugates of the application and Uses Thereof

5

[0045] In one embodiment, the application discloses a therapeutic method for preventing or treating a pathological condition or symptom in a mammal, such as a human, wherein the activity of a TLR agonist is implicated and its action is desired. The method includes administering to a mammal in need of such therapy, an effective amount of a compound of the application, or a pharmaceutically acceptable salt thereof. Non-limiting examples of pathological conditions or symptoms that are suitable for treatment include cancers, inflammatory diseases of the gastrointestinal tract, brain, skin, joints, and other tissues, bacterial or viral diseases, autoimmune diseases, and treating Crohn's Disease. The compounds of the application can also be used to prepare vaccines against bacteria, viruses, cancer cells, or cancer-specific peptides, or to enhance anti-cancer monoclonal antibodies, as a CNS stimulant, or for biodefense. The application thus discloses a compound of the invention for use in medical therapy (e.g., for use as an anti-cancer agent, to treat bacterial diseases, to treat for viral diseases, such as hepatitis C and hepatitis B, to treat Crohn's Disease, and generally as therapeutic agents for treating immunologic disease). Furthermore, compounds of the application may prevent carcinogenesis, e.g., by hepatitis C and hepatitis B viruses, and may be used for the manufacture of a medicament useful for the treatment of cancer, viral or bacterial infection, Crohn's Disease, and immunologic disorders in a mammal, such as a human.

10

15

[0046] In one aspect, the present application discloses a use for treating a viral infection in a mammal by administering a TLR agonist conjugate of the invention. The viral infection may be caused by a RNA virus, a product of the RNA virus that acts as a TLR agonist, and/or a DNA virus. An exemplary DNA virus is hepatitis B virus. In one example, the viral infection is caused by a coronavirus that causes Severe Acute Respiratory Syndrome (SARS), a Hepatitis B virus, or a Hepatitis C Virus.

20

[0047] In one aspect, the present application discloses a use for treating cancer by administering an effective amount of a TLR agonist conjugate of the invention. The cancer may be an interferon sensitive cancer, such as, for example, a leukemia, a lymphoma, a myeloma, a melanoma, or a renal cancer. Specific cancers that can be treated include melanoma, superficial bladder cancer, actinic keratoses, intraepithelial neoplasia, and basal cell skin carcinoma, squamous, and the like. In addition, the method of the invention includes treatment for a precancerous condition such as, for example, actinic keratoses or intraepithelial neoplasia, familial polyposis (polyps), cervical dysplasia, cervical cancers, superficial bladder cancer, and any other cancers associated with infection (e.g., lymphoma Karposi's sarcoma, or leukemia); and the like.

25

30

[0048] In another aspect, the present application discloses a use for treating an autoimmune disease by administering a therapeutically effective amount of a TLR agonist conjugate of the invention or a pharmaceutically acceptable salt of such a compound. Exemplary autoimmune diseases are Multiple Sclerosis, lupus, rheumatoid arthritis and the like.

35

[0049] In another aspect, the present application disclose a use of treating Crohn's Disease by administering a TLR agonist conjugate of the invention.

[0050] The TLR agonist conjugates may include a homofunctional TLR agonist polymer, e.g., formed of a TLR7 agonist or a TLR3 agonist. The TLR7 agonist can be a 7-thia-8-oxoguanosinyl (TOG) moiety, a 7-deazaguanosinyl (7DG) moiety, a resiquimod moiety, or an imiquimod moiety. In another aspect, the TLR agonist conjugate may include a heterofunctional TLR agonist polymer. The heterofunctional TLR agonist polymer may include a TLR7 agonist and a TLR3 agonist or a TLR9 agonist, or all three agonists. The heterofunctional TLR agonist polymer can include a TLR8 agonist and a TLR9 agonist.

40

[0051] The application includes covalently conjugating a synthetic TLR agonist with selected macromolecules to achieve, for example, a desired molecular shape, size and valence in order to optimize the immunologic properties of the resulting conjugate, and/or to target or deliver the conjugate to desired cells and tissues. As described herein, the conjugates are designed to be useful in a variety of medical applications including, but not limited to, allergic asthma, respiratory viral infections (influenza and RSV), lupus and other autoimmune diseases, and as antigen-adjuvant combinations for vaccines against cancer and infectious diseases. The conjugates provide an optimum immune response while limiting undesirable systemic side effects by tethering the immune activator (the synthetic TLR agonist) to a macromolecule by a strong covalent bond. The macromolecule may serve as a targeting entity and/or an integral part of the immune response, such as the antigen in an adjuvant-antigen conjugate. A major advantage when administering a stable conjugate in a localized environment is that only very small amounts of TLR agonist are released over time into the systemic environment.

45

50

[0052] In one aspect, the macromolecule is selected from products, such as proteins, lipids or dendrimers, or polymers having amino groups on their surfaces, such as polystyrene "amino beads," each having primary amino groups available for conjugation to a linker such as SANH, or for direct conjugation to the synthetic TLR7 agonist. For example, following conjugation of the linker and macromolecule, the TLR7 agonist, such as UC-1V150, is reacted with the NHS ester of the SANH-macromolecule conjugate to provide a TLR7 agonist-SANH-macromolecule conjugate.

55

[0053] Vaccines are not generally used in acute settings because (1) they take too long to act and (2) they are not effective in immune compromised patients. For example, *Staphylococcus aureus* (SA) infections are a major cause of morbidity and mortality in hospitalized patients. Groups particularly at risk are those with immune suppression due to burns, trauma, catheter placement, dialysis, or advanced age in nursing homes. Moreover, many strains of hospital

acquired SA are resistant to conventional antibiotics.

[0054] The present invention overcomes these two barriers to treatment. The use of compositions, including synthetic TLR7 agonists and conjugates with synthetic TLR7 agonists, in combination with gram-positive bacterial antigens is provided herein. TLR7 ligands generally have poor pharmacokinetics, and rapid systemic absorption and excretion. Due to systemic dispersal, they result in cytokine syndrome. Effective adjuvants must create an "immune gradient" of cytokines and chemokines. In one embodiment, conjugation of potent synthetic TLR7 agonists to macromolecules enhances delivery properties, improves pharmacokinetics, and avoids systemic toxicity by localized exposure.

[0055] In one embodiment, the invention provides a composition for use in method to prevent or inhibit a gram-positive bacterial infection in a mammal, comprising administering to the mammal an effective amount of a composition comprising a bacterial antigen of a gram-positive bacteria and an amount of a synthetic TLR7 agonist. In another embodiment, the invention provides a compound for use in method to prevent or inhibit a gram-positive bacterial infection in a mammal, comprising administering to the mammal an effective amount of a synthetic TLR7 agonist conjugated to a gram-positive bacterial antigen. For example, a 1V150-MSA conjugate retains its TLR7 agonist activity, has enhanced potency and reduced toxicity, causes local activation of innate immunity, and induces T cell dependent immune protection within 6 days after a single vaccination with a bacterial antigen.

[0056] In one embodiment, a synthetic TLR7 agonist is administered with or conjugated to one or more antigens of *S. aureus*. Table 1 provides exemplary antigens for *S. aureus* for use with synthetic TLR7 agonists, particularly in acute care settings. The vaccines of the invention may unexpectedly provide a rapid and effective immune response.

Table 1

| <i>Staphylococcus aureus</i> immunogens | |
|--|-----------------------------------|
| Weapon | |
| | Exfoliative toxin B |
| | Exfoliative toxin A |
| | Toxic shock-syndrome toxin |
| | Enterotoxin A-E, H-U |
| | Bone sialoprotein-binding protein |
| | Collagen-binding protein |
| | Clumping factor A |
| | Clumping factor B |
| | α -hemolysin |
| | γ -hemolysin |
| | Protein A |
| | Clumping factor A |
| | Fibronectin-binding protein A |
| | Fibronectin-binding protein B |
| | Collagen-binding protein |
| | Lipoteichoic acid |
| | Peptidoglycan |
| | Protein A |
| | Fibronectin-binding protein B |
| | α -hemolysin |
| | Panton valentine leukocidin |
| | Collagen-binding protein |
| | Lipoteichoic acid |
| | Peptidoglycan |
| | Capsular polysaccharide |
| | Clumping factor A |
| | Protein A |


(continued)

Staphylococcus aureus immunogens


Weapon

Fibronectin-binding proteins

[0057] In one aspect, the application discloses the following conjugates




a compound of formula (II)

Thiazolopyrimidines

a compound of formula (III)

PurinesX¹ = -O-, -S-, or NR^c,wherein R^c hydrogen, C₁₋₁₀alkyl, or C₁₋₁₀alkyl substituted by C₃₋₆ cycloalkyl, or R^c and R¹ taken together with the nitrogen atom can form a heterocyclic ring or a substituted heterocyclic ring, wherein the substituents are hydroxy, C₁₋₆ alkyl, hydroxy C₁₋₆ alkylene, C₁₋₆ alkoxy, C₁₋₆ alkoxy C₁₋₆ alkylene, or cyano;wherein R¹ is (C₁₋₁₀)alkyl, substituted (C₁₋₁₀)alkyl, C₆₋₁₀ aryl, or substituted C₆₋₁₀ aryl, C₅₋₉ heterocyclic, substituted C₅₋₉ heterocyclic; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆ alkyl, hydroxy C₁₋₆ alkylene, C₁₋₆ alkoxy, C₁₋₆ alkoxy C₁₋₆ alkylene, amino, cyano, halogen, or aryl;each R² is independently hydrogen, -OH, (C₁₋₆)alkyl, substituted (C₁₋₆)alkyl, (C₁₋₆)alkoxy, substituted (C₁₋₆)alkoxy, -C(O)-(C₁₋₆)alkyl (alkanoyl), substituted -C(O)-(C₁₋₆)alkyl, -C(O)-(C₆₋₁₀)aryl (aroyl), substituted -C(O)-(C₆₋₁₀)aryl, -C(O)OH (carboxyl), -C(O)O(C₁₋₆)alkyl (alkoxycarbonyl), substituted -C(O)O(C₁₋₆)alkyl, -NR^aR^b, -C(O)NR^aR^b (carbamoyl), -O-C(O)NR^aR^b, -(C₁₋₆)alkylene-NR^aR^b, -(C₁₋₆)alkylene-C(O)NR^aR^b, halo, nitro, or cyano;wherein each R^a and R^b is independently hydrogen, (C₁₋₆)alkyl, (C₃₋₈)cycloalkyl, (C₁₋₆)alkoxy, halo(C₁₋₆)alkyl, (C₃₋₈)cycloalkyl(C₁₋₆)alkyl, (C₁₋₆)alkanoyl, hydroxy(C₁₋₆)alkyl, aryl, aryl(C₁₋₆)alkyl, aryl, aryl(C₁₋₆)alkyl, Het, Het (C₁₋₆)alkyl, or (C₁₋₆)alkoxycarbonyl; wherein X² is a bond or a linking group; wherein R³ is a macromolecule; wherein n is 1, 2, 3, or 4; wherein m is 1 or 2; wherein q is 1 to 1,000, 10⁴, 10⁵, 10⁶ or more; or a pharmaceutically acceptable salt thereof. The macromolecule groups can include organic molecules, composed of carbon, oxygen, hydrogen, nitrogen, sulfur, phosphorous, or combinations thereof, which are not harmful to body tissues (e.g., they are non-toxic, and/or do not cause inflammation) and may include, but not be limited to, dendrimers, proteins, peptides, lipids and their formulations (e.g., liposome nanoparticles), with or without linkers (X² groups), and amino-modified polymers, such as polystyrene beads, as well as α-galactosylceramides (see Figure 1).


[0058] The compounds of the application can be prepared using compounds having formula (IA-1):



(IA-1)

where X² is a group that can react to a specific group of compounds, e.g., those disclosed in U.S. Patent No. 6,329,381 (Kurimoto et al.), or form a bond to a linking group or react to form a bond to a macromolecule, and the remaining variables are as defined above for formula (IA). Non-limiting examples of macromolecules include those with side chains that increase solubility, such as, for example, groups containing morpholino, piperidino, pyrrolidino, or piperazino rings and the like; amino acids, polymers of amino acids (proteins or peptides), e.g., dipeptides or tripeptides, and the like; carbohydrates (polysaccharides), nucleotides such as, for example, PNA, RNA and DNA, and the like; polymers of organic materials, such as, for example, polyethylene glycol, poly-lactide and the like; monomeric and polymeric lipids; insoluble organic nanoparticles; non-toxic body substances such as, for example, cells, lipids, vitamins, co-factors, antigens such as, for example microbes, such as, for example, viruses, bacteria, fungi, and the like. The antigens can include inactivated whole organisms, or sub-components thereof, e.g., cells and the like.

[0059] In one aspect, a compound of the application has formula (IC):



wherein

X is N or CR^x wherein R^x is hydrogen, halogen, substituted alkyl, unsubstituted alkyl, substituted heteroalkyl, or unsubstituted heteroalkyl;

Y is S or N;

the dashes (----) indicate optional bonds; wherein:

when the bond between Y and the carbon marked by an asterisk is a double bond, Q² is not present;

when the bond between Q¹ and the carbon marked by an asterisk is a double bond, Q¹ is O, S, NY¹, or NNY²Y³; and

when the bond between Q¹ and the carbon marked by an asterisk is a single bond, Q¹ is hydrogen, cyano, nitro, O-Y², S-Y², NY¹Y², or NY²NY³Y⁴; wherein

Y¹ is hydrogen, substituted alkyl, unsubstituted alkyl, substituted cycloalkyl, unsubstituted cycloalkyl, substituted heteroalkyl, unsubstituted heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl, -C(=O)-substituted alkyl, -C(=O)-unsubstituted alkyl, -C(=O)O-substituted alkyl, -C(=O)O-unsubstituted alkyl, cyano, nitro, hydroxyl, or O-Y²;

Y², Y³, and Y⁴, are each independently hydrogen, substituted alkyl, unsubstituted alkyl, substituted heteroalkyl, unsubstituted heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl;

Z is O, S, or NY⁵ wherein Y⁵ is hydrogen, substituted alkyl, unsubstituted alkyl, substituted heteroalkyl, unsubstituted heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl;

Q² and Q³ are each independently hydrogen, substituted alkyl, unsubstituted alkyl, substituted heteroalkyl, unsubstituted heteroalkyl, substituted aryl, unsubstituted aryl, substituted heteroaryl, unsubstituted heteroaryl;

X¹ is -O-, -S-, or -NR^c-;

R^c is hydrogen, C₁₋₁₀alkyl, or substituted C₁₋₁₀alkyl, or R^c and R¹ taken together with the nitrogen atom can form a heterocyclic ring or a substituted heterocyclic ring;

R¹ is hydrogen, (C₁-C₁₀)alkyl, substituted (C₁-C₁₀)alkyl, C₆₋₁₀aryl, or substituted C₆₋₁₀aryl, C₅₋₉heterocyclic, or substituted C₅₋₉heterocyclic ring;

each R² is independently hydrogen, -OH, (C₁-C₆)alkyl, substituted (C₁-C₆)alkyl, (C₁-C₆)alkoxy, substituted (C₁-C₆)alkoxy, -C(O)-(C₁-C₆)alkyl (alkanoyl), substituted -C(O)-(C₁-C₆)alkyl, -C(O)-(C₆-C₁₀)aryl (aroyl), substituted -C(O)-(C₆-C₁₀)aryl, -C(O)OH (carboxyl), -C(O)O(C₁-C₆)alkyl (alkoxycarbonyl), substituted -C(O)O(C₁-C₆)alkyl, -NR^aR^b, -C(O)NR^aR^b (carbamoyl), -O-C(O)NR^aR^b, -(C₁-C₆)alkylene-NR^aR^b, -(C₁-C₆)alkylene-C(O)NR^aR^b, halo, nitro, or cyano;

each R^a and R^b is independently hydrogen, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₁-C₆)heteroalkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl, (C₃-C₈)cycloalkyl(C₁-C₆)alkyl, (C₁-C₆)alkanoyl, hydroxy(C₁-C₆)alkyl, aryl, aryl(C₁-C₆)alkyl, Het, Het

(C₁-C₆)alkyl, or (C₁-C₆)alkoxycarbonyl;

wherein the substituents on any alkyl, cycloalkyl, heteroalkyl, amino, alkoxy, alkanoyl, aryl, heteroaryl, or heterocyclic groups are one or more (e.g., 1, 2, 3, 4, 5, or 6) hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, heterocycle (such as piperidinyl or morpholinyl), or aryl;

X² is a bond or a linking group;

k is 0, 1, 2, 3, or 4;

n is 0, 1, 2, 3, or 4; and

R³ is a macromolecule comprising a cell, virus, vitamin, cofactor, peptide, protein, nucleic acid molecule, lipid, bead or particle, such as a polystyrene bead or nanoparticles, or a dendrimer;

or a pharmaceutically acceptable salt thereof, including hydrates thereof.

[0060] In certain examples, the groups X²-R³ can form a linker to a second formula (IC) moiety so as to form a dimer. For example, the linker can be any linker as described herein, such as a divalent aryl or heteroaryl, bis-amide aryl, bis-amide heteroaryl, bis-hydrazide aryl, bis-hydrazide heteroaryl, or the like. Alternatively, Q¹ can form a linker to a second formula (IC) moiety so as to form a dimer through a disulfide linkage. See for example, Figure 28.

[0061] In cases where compounds are sufficiently basic or acidic to form acid or base salts, use of the compounds as salts may be appropriate. Examples of acceptable salts are organic acid addition salts formed with acids which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, α -ketoglutarate, and α -glycerophosphate. Suitable inorganic salts may also be formed, including hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

[0062] Acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

[0063] Alkyl includes straight or branched C₁₋₁₀ alkyl groups, e.g., methyl, ethyl, propyl, butyl, pentyl, isopropyl, isobutyl, 1-methylpropyl, 3-methylbutyl, hexyl, and the like.

[0064] Lower alkyl includes straight or branched C₁₋₆ alkyl groups, e.g., methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl, and the like.

[0065] The term "alkylene" refers to a divalent straight or branched hydrocarbon chain (e.g., ethylene: -CH₂-CH₂).

[0066] C₃₋₇ Cycloalkyl includes groups such as, cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like, and alkyl-substituted C₃₋₇ cycloalkyl group, preferably straight or branched C₁₋₆ alkyl group such as methyl, ethyl, propyl, butyl or pentyl, and C₅₋₇ cycloalkyl group such as, cyclopentyl or cyclohexyl, and the like.

[0067] Lower alkoxy includes C₁₋₆ alkoxy groups, such as methoxy, ethoxy or propoxy, and the like.

[0068] Lower alkanoyl includes C₁₋₆ alkanoyl groups, such as formyl, acetyl, propanoyl, butanoyl, pentanoyl or hexanoyl, and the like.

C₇₋₁₁ aryl, includes groups such as benzoyl or naphthoyl;

Lower alkoxycarbonyl includes C₂₋₇ alkoxycarbonyl groups, such as methoxycarbonyl, ethoxycarbonyl or propoxycarbonyl, and the like.

[0069] Lower alkylamino group means amino group substituted by C₁₋₆ alkyl group, such as, methylamino, ethylamino, propylamino, butylamino, and the like.

[0070] Di(lower alkyl)amino group means amino group substituted by the same or different and C₁₋₆ alkyl group (e.g., dimethylamino, diethylamino, ethylmethylamino).

[0071] Lower alkylcarbamoyl group means carbamoyl group substituted by C₁₋₆ alkyl group (e.g., methylcarbamoyl, ethylcarbamoyl, propylcarbamoyl, butylcarbamoyl).

[0072] Di(lower alkyl)carbamoyl group means carbamoyl group substituted by the same or different and C₁₋₆ alkyl group (e.g., dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl).

[0073] Halogen atom means halogen atom such as fluorine atom, chlorine atom, bromine atom or iodine atom.

[0074] Aryl refers to a C₆₋₁₀ monocyclic or fused cyclic aryl group, such as phenyl, indenyl, or naphthyl, and the like.

[0075] Heterocyclic or heterocycle refers to monocyclic saturated heterocyclic groups, or unsaturated monocyclic or fused heterocyclic group containing at least one heteroatom, e.g., 0-3 nitrogen atoms (-NR^d where R^d is H, alkyl, or Y² as defined herein), 0-1 oxygen atom (-O-), and 0-1 sulfur atom (-S-). Non-limiting examples of saturated monocyclic heterocyclic group includes 5 or 6 membered saturated heterocyclic group, such as tetrahydrofuranyl, pyrrolidinyl, morpholinyl, piperidyl, piperazinyl or pyrazolidinyl. Non-limiting examples of unsaturated monocyclic heterocyclic group includes 5 or 6 membered unsaturated heterocyclic group, such as furyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, thienyl, pyridyl or pyrimidinyl. Non-limiting examples of unsaturated fused heterocyclic groups includes unsaturated bicyclic

heterocyclic group, such as indolyl, isoindolyl, quinolyl, benzothiazolyl, chromanyl, benzofuranyl, and the like. A Het group can be a saturated heterocyclic group or an unsaturated heterocyclic group, such as a heteroaryl group.

[0076] R^c and R¹ taken together with the nitrogen atom to which they are attached can form a heterocyclic ring. Non-limiting examples of heterocyclic rings include 5 or 6 membered saturated heterocyclic rings, such as 1-pyrrolidinyl, 4-morpholinyl, 1-piperidyl, 1-piperazinyl or 1-pyrazolidinyl, 5 or 6 membered unsaturated heterocyclic rings such as 1-imidazolyl, and the like.

[0077] The alkyl, aryl, heterocyclic groups of R¹ can be optionally substituted with one or more substituents, wherein the substituents are the same or different, and include lower alkyl; cycloalkyl, hydroxyl; hydroxy C₁₋₆ alkylene, such as hydroxymethyl, 2-hydroxyethyl or 3-hydroxypropyl; lower alkoxy; C₁₋₆ alkoxy C₁₋₆ alkyl, such as 2-methoxyethyl, 2-ethoxyethyl or 3-methoxypropyl; amino; alkylamino; dialkyl amino; cyano; nitro; acyl; carboxyl; lower alkoxy carbonyl; halogen; mercapto; C₁₋₆ alkylthio, such as, methylthio, ethylthio, propylthio or butylthio; substituted C₁₋₆ alkylthio, such as methoxyethylthio, methylthioethylthio, hydroxyethylthio or chloroethylthio; aryl; substituted C₆₋₁₀ monocyclic or fused-cyclic aryl, such as 4-hydroxyphenyl, 4-methoxyphenyl, 4-fluorophenyl, 4-chlorophenyl or 3,4-dichlorophenyl; 5-6 membered unsaturated heterocyclic, such as furyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, thienyl, pyridyl or pyrimidinyl; and bicyclic unsaturated heterocyclic, such as indolyl, isoindolyl, quinolyl, benzothiazolyl, chromanyl, benzofuranyl or phthalimino. In certain examples, one or more of the above groups can be expressly excluded as a substituent of various other groups of the formulas.

[0078] In some examples, the five-membered ring of the formula is a thiazole ring, e.g., where Y of formula IA above is S and Q² is absent.

[0079] The alkyl, aryl, heterocyclic groups of R² can be optionally substituted with one or more substituents, wherein the substituents are the same or different, and include hydroxyl; C₁₋₆ alkoxy, such as methoxy, ethoxy or propoxy; carboxyl; C₂₋₇ alkoxy carbonyl, such as methoxycarbonyl, ethoxycarbonyl or propoxycarbonyl) and halogen.

[0080] The alkyl, aryl, heterocyclic groups of R^c can be optionally substituted with one or more substituents, wherein the substituents are the same or different, and include C₁₋₆ cycloalkyl; hydroxyl; C₁₋₆ alkoxy; amino; cyano; aryl; substituted aryl, such as 4-hydroxyphenyl, 4-methoxyphenyl, 4-chlorophenyl or 3,4-dichlorophenyl; nitro and halogen.

[0081] The heterocyclic ring formed together with R^c and R¹ and the nitrogen atom to which they are attached can be optionally substituted with one or more substituents, wherein the substituents are the same or different, and include C₁₋₆ alkyl; hydroxy C₁₋₆ alkylene; C₁₋₆ alkoxy C₁₋₆ alkylene; hydroxyl; C₁₋₆ alkoxy; and cyano.

[0082] In some examples, when Q¹ is O-Y², Y² is not hydrogen.

[0083] A specific value for X is N.

[0084] Another specific value for X is CH.

[0085] Of course, only one of the two bonds indicated by dashed lines may be present in one molecule of a compound of the indicated formula. In one example, the bond between Y and the carbon marked by an asterisk is a double bond. In another example, the bond between Q¹ and the carbon marked by an asterisk is a double bond.

[0086] A specific value for Q¹ is O.

[0087] Another specific value for Q¹ is S.

[0088] Another specific value for Q¹ is NY¹, for example, =NH.

[0089] Another specific value for Q¹ is NNY²Y³.

[0090] In one example, the bond between Q¹ and the carbon marked by an asterisk is a single bond.

[0091] A specific value for Q¹ is hydrogen.

[0092] Another specific value for Q¹ is NH₂.

[0093] Another specific value for Q¹ is O-Y².

[0094] A specific value for Y¹ is hydrogen.

[0095] Another specific value for Y¹ is alkyl, for example, (C₁-C₆)alkyl, such as methyl.

[0096] Another specific value for Y¹ is aryl, such as phenyl.

[0097] A specific value for each of Y², Y³, and Y⁴ is hydrogen.

[0098] Another specific value for each of Y², Y³, and Y⁴ (independently) is alkyl, for example, (C₁-C₆)alkyl, such as methyl.

[0099] Another specific value for each of Y², Y³, and Y⁴ (independently) is aryl, such as phenyl.

[0100] A specific value for Z is O.

[0101] Another specific value for Z is S.

[0102] Another specific value for Z is NY⁵ wherein Y⁵ is hydrogen, methyl, or phenyl.

[0103] A specific value for Q² is hydrogen.

[0104] Another specific value for Q² is methyl, or phenyl.




[0105] A specific value for Q³ is hydrogen.

[0106] Another specific value for Q³ is methyl, or phenyl.


[0107] A specific value for X¹ is a sulfur atom, an oxygen atom or -NR^c-.


[0108] Another specific X¹ is a sulfur atom.

- [0109] Another specific X¹ is an oxygen atom.
 [0110] Another specific X¹ is -NR^c-.
 [0111] Another specific X¹ is -NH-.
 [0112] A specific value for Y is N.
 5 [0113] Another specific value for Y is S.
 [0114] A specific value for R^c is hydrogen, C₁₋₄ alkyl or substituted C₁₋₄ alkyl.
 [0115] A specific value for R¹ and R^c taken together is when they form a heterocyclic ring or a substituted heterocyclic ring.
 10 [0116] Another specific value for R¹ and R^c taken together is substituted or unsubstituted morpholino, piperidino, pyrrolidino, or piperazino ring
 [0117] A specific value for R¹ is hydrogen, C₁₋₄alkyl, or substituted C₁₋₄alkyl.
 [0118] Another specific R¹ is 2-hydroxyethyl, 3-hydroxypropyl, 4-hydroxybutyl, 2-aminoethyl, 3-aminopropyl, 4-aminobutyl, methoxymethyl, 2-methoxyethyl, 3-methoxypropyl, ethoxymethyl, 2-ethoxyethyl, methylthiomethyl, 2-methylthioethyl, 3-methylthiopropyl, 2-fluoroethyl, 3-fluoropropyl, 2,2,2-trifluoroethyl, cyanomethyl, 2-cyanoethyl, 3-cyanopropyl, 15 methoxycarbonylmethyl, 2-methoxycarbonylethyl, 3-methoxycarbonylpropyl, benzyl, phenethyl, 4-pyridylmethyl, cyclohexylmethyl, 2-thienylmethyl, 4-methoxyphenylmethyl, 4-hydroxyphenylmethyl, 4-fluorophenylmethyl, or 4-chlorophenylmethyl.
 [0119] Another, specific R¹ is hydrogen, CH₃-, CH₃-CH₂-, CH₃CH₂CH₂-, hydroxyC₁₋₄alkylene, or C₁₋₄alkoxyC₁₋₄alkylene.
 20 [0120] Another specific value for R¹ is hydrogen, CH₃-, CH₃-CH₂-, CH₃-O-CH₂CH₂- or CH₃-CH₂-O-CH₂CH₂-.
 [0121] A specific value for R² is hydrogen, halogen, or C₁₋₄alkyl.
 [0122] Another specific value for R² is hydrogen, chloro, bromo, CH₃-, or CH₃-CH₂-.
 [0123] Specific substituents for substitution on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkylene, C₃₋₆cycloalkyl, amino, cyano, halogen, or aryl.
 25 [0124] A specific value for X² is a bond or a chain having up to about 24 atoms; wherein the atoms are selected from the group consisting of carbon, nitrogen, sulfur, non-peroxide oxygen, and phosphorous.
 [0125] Another specific value for X² is a bond or a chain having from about 4 to about 12 atoms.
 [0126] Another specific value for X² is a bond or a chain having from about 6 to about 9 atoms.
 [0127] Another specific value for X² is




50 or






or



15

[0128] Another specific value for X² is



25

[0129] In certain examples, the linker or the group X² is not a linker disclosed in PCT Application Publication No. WO 2007/024707. Additionally, in some examples, R³ is not an auxiliary disclosed in PCT Application Publication No. WO 2007/024707.

30

[0130] A specific macromolecule is an amino acid, a carbohydrate, a peptide, a protein, an antigen, a nucleic acid, a lipid, a dendrimer, a body substance, or a cell such as a microbe.

[0131] A specific peptide, has from 2 to about 20 amino acid residues.

[0132] Another specific peptide, has from 10 to about 20 amino acid residues.

[0133] A specific macromolecule includes a carbohydrate.

[0134] A specific nucleic acid is DNA, RNA or PNA.

35

[0135] A specific macromolecule is a cell, lipid, vitamin, lipid, or co-factor.

[0136] A specific antigen is a microbe.

[0137] A specific microbe is a virus, bacteria, or fungi.

[0138] Another specific microbe is a virus or a bacteria.

[0139] Specific bacteria are *Bacillus anthracis*, *Listeria monocytogenes*, *Francisella tularensis*, *Salmonella*, or *Staphylococcus*.

40

[0140] Specific *Salmonella* are *S. typhimurium* or *S. enteritidis*.

[0141] Specific *Staphylococcus* include *S. aureus*.

[0142] Specific viruses are RNA viruses, including RSV and influenza virus, a product of the RNA virus, or a DNA virus, including herpes virus.

45

[0143] A specific DNA virus is hepatitis B virus.

[0144] In other examples, the macromolecule is not an amino acid, a carbohydrate, a peptide, an antigen such as a microbe, for example, a virus (for example, RNA viruses, e.g., SIV, hepatitis C virus or a coronavirus, a product of the RNA virus, or a DNA virus, such as Hepatitis B virus, fungi, or bacteria such as *Bacillus anthracis* (anthrax), *Listeria monocytogenes*, *Francisella tularensis*, or *Salmonella* (e.g., typhimurium or enteritidis), a nucleic acid such as DNA, RNA, PNA, or a body substance such as a cell or lipid.

50

[0145] A specific value for k is 0. Another specific value for k is 1. Another specific value for k is 2. In some embodiments, k is not 1.

[0146] Specific compounds of the application have the general formula

55

IA-L-A¹;

IA-L-(A¹)₂;

IA-L-A¹-A¹;IA-L-A¹-L-A¹;

5

(IA)₂-L-A¹-A¹;(IA)₂-L-A¹-L-A¹;

10

(IA)₂-L-A¹; or(IA)₂-L-(A¹)₂;


wherein IA is as disclosed herein; L is absent or is a linking group; and each A¹ group independently represents a macromolecule.

[0147] The application discloses compositions of a compound of the invention optionally in combination with other active agents, e.g., ribavirin, mizoribine, and mycophenolate mofetil. Other non-limiting examples are known and are disclosed in U.S. published patent application No. 20050004144.


[0148] Processes for preparing compounds of the application for preparing intermediates useful for preparing compounds of the application are provided as further examples of the application. Intermediates useful for preparing compounds of the application are also provided as further examples of the application.

[0149] For example, the compounds (conjugates) of the application can be prepared using standard synthetic methods known in the art. A general ester and aldehyde synthesis is illustrated below. UC-1V150 was synthesized in seven steps from 2,6-dichloropurine. The free aldehyde group on the benzyl moiety of UC-1V150 enabled us to couple the agonist to many different auxiliary chemical entities, including proteins, oligonucleotides, aromatic molecules, lipids, viruses, and cells, through a linker molecule that contained a hydrazine or amino group.


General Synthesis



NHS Ester Synthesis



Aldehyde Synthesis



[0150] Chemistry of UC-1V150. The synthesis of UC-1V150 and the preparation of the indicated compounds 2-8 was as follows. **Compound 2:** 4-(2,6-dichloropurin-9-ylmethyl)benzoxonitrile. 2,6-dichloro-9H-purine (1, 16 mmol) was dissolved in DMF (50 mL) with potassium carbonate (50 mmol) added, and the mixture was stirred at ambient temperature for 16 hours after adding α -Bromo-*p*-tolunitrile (22 mmol). After filtration to remove insoluble inorganic salts, the filtrate was poured into water (1500 mL) and extracted with ethyl acetate (2 x 400 mL), dried over magnesium sulfate and evaporated to yield a residue which was subjected to flash silica gel chromatography using 1:2:10 ethyl acetate/acetone/hexanes. Yield 3.33 g (69%). UV, NMR and MS were consistent with structure assignment. **Compound 3:** 4-(6-amino-2-chloropurin-9-ylmethyl)benzoxonitrile. **Compound 2** (1.9 g) was placed in a steel reaction vessel and methanolic ammonia (80 mL, 7 N) was added. The sealed vessel was heated at 60°C for 12 hours, cooled in ice and the solid product filtered off. Yield 1.09 g. UV, NMR and MS were consistent with assigned structure. **Compound 4:** 4-[6-amino-2-(2-methoxyethoxy)purin-9-ylmethyl]benzoxonitrile. The sodium salt of 2-methoxyethanol was first generated by dissolving sodium metal (81 mg) in 2-methoxyethanol (30 mL) with heat, and then **compound 3** (1.0 g) dissolved in methoxyethanol was added (300 mL, with heat). The reaction mixture was heated for 8 hours at 115°C bath temperature, concentrated *in vacuo* to near dryness and the residue partitioned between ethyl acetate and water. Flash silica gel chromatography of the organic layer using 5% methanol in dichloromethane gave 763 mg product. NMR was consistent with structure assignment. **Compound 5:** 4-[6-amino-8-bromo-2-(2-methoxyethoxy)purin-9-ylmethyl]benzoxonitrile. **Compound 4** (700

mg) was dissolved in dichloromethane (400 mL) and bromine (7 mL) was added dropwise. The mixture was stirred overnight at room temperature and extracted first with aqueous sodium thiosulfate (2 L of 0.1 M) solution, then with aqueous sodium bicarbonate (500 mL, saturated). The residue from the organic layer was chromatographed on silica gel using 3% methanol in dichloromethane to yield 460 mg of bromo product. NMR, W and MS were consistent with structure assignment. **Compound 6:** 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzotriazole. Sodium methoxide was generated by reaction of sodium metal (81 mg) in dry methanol (30 mL) and combined with a solution of **compound 5** (700 mg) dissolved in dry dimethoxyethane and the temperature raised to 100°C. After overnight reaction, the mixture was concentrated *in vacuo* and the residue was chromatographed on silica using 5% methanol in dichloromethane. Yield 120 mg. NMR was consistent with structure assignment. **Compound 7:** 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzaldehyde. **Compound 6** (100 mg) was dissolved in dry THF (3 mL) and cooled to 0°C under argon. The reducing agent, lithium N,N'-(dimethylethylenediamino)aluminum hydride, used to convert the nitrile to the aldehyde function. A 0.5 M solution in dry THF was prepared and 0.72 mL of such was added to the reaction flask. The mixture was stirred at 0-5°C for 1 hour, quenched by addition of 3 M HCl, extracted with ethyl acetate followed by dichloromethane, and then concentrated *in vacuo* to yield 85 mg. NMR was consistent with structure assignment. **Compound 8:** 4-[6-amino-8-hydroxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzaldehyde (UC-1V150). **Compound 7** (800 mg) was combined with sodium iodide (504 mg) and acetonitrile (40 mL), and chlorotrimethylsilane (0.5 mL) was slowly added. The mixture was heated at 70°C for 3.5 hours, cooled and filtered. The solid product was washed with water, then ether, to yield 406 mg. NMR, UV, and MS were consistent with structure assignment.

[0151] Additional examples for preparing specific compounds are included herein.

[0152] As described in the examples herein, a soluble TLR7 agonist capable of covalent coupling to primary amines under physiologic conditions was prepared. The *in vitro* activity of several compounds and antigen-adjutant complexes was then tested utilizing bone marrow-derived murine or peripheral blood mononuclear cell-derived dendritic cells (DC) to characterize DC maturation and cytokine secretion (e.g., IL-12, IL-6, TGF-beta, and IFN-gamma). Immunocompetent syngeneic C57/B1 mice were prophylactically vaccinated with intradermal antigen-TLR7 agonist complexes and challenged with B 16 melanoma tumor cells expressing the cOVA transgene.

[0153] The effective concentration (EC₅₀) for each compound generally followed a bell shaped distribution with higher doses being inhibitory. Maximal stimulation occurred between 10 and 1000 nM. Covalently coupled adjuvant molecules to TLR agonist retained activity but with generally lower EC₅₀ values. Coupling UC-1V199 to chicken ovalbumin nearly doubled median survival from 22 to 35 days following subcutaneous tumor challenge compared with chicken ovalbumin alone.

[0154] Thus, covalent linkage of a TLR7 agonist to a tumor antigen stimulated DC cytokine production and protected mice from tumor challenge. The use of a suitable TLR7 agonist which retains its immune stimulating properties under physiologic conditions following coupling to a macromolecule, such as an antigen, may be useful in the development of an *in situ* vaccine in solid tumor therapy.

[0155] Various purines, pyridines, and imidazoquinolines, with molecular weights of 200-400 kD, have been shown to activate TLR7 and compounds that were specific TLR7 ligands were 100-1000 fold more powerful than imiquimod on a molar basis (Lee et al., *infra*). Because these TLR agonists are structurally very similar to normal component of nucleotides, they are very unlikely to induce a haptenic immune reaction after repeated administration.

[0156] An adenine based TLR7 pharmacore may need to be covalently tied to an "auxiliary group" (macromolecule) to promote uptake into the endosomes of dendritic cells, where TLR7 is expressed, and to retain the TLR agonist. Accordingly, the TLR7 agonist UC-1V150 was prepared and coupled via its aldehyde function and a linker to free amino groups on various proteins, including mouse albumin (MSA) (Figure 3). The conjugates were 100-fold more potent *in vitro* and *in vivo* than the uncoupled adenine analog. Moreover, intrapulmonary administration of the albumin conjugate (UC-1V150/MSA) to mice induced local cytokine production in the bronchial alveolar lavage fluid (BALF) without systemic cytokine release. In marked contrast, the delivery of the untethered drug to the airways quickly triggered cytokine release in the bloodstream.

[0157] In one embodiment, a TLR7 agonist maximizes the production of Th1 stimulating cytokines (interferons and IL-12) compared to TNF α and IL-1. TLR7 is localized on the inner surfaces of the endosomal vesicles that are constantly synthesized and undergo maturation in DC. For example, to prevent asthma, a stable and potent TLR agonist that traffics to the early endosomes of dendritic cells and induces primarily Type I interferons is preferred. A TLR agonist was covalently attached to a phospholipid auxiliary group with the expectation that the conjugate, UC-1V199/L (Figure 6), would quickly and stably insert into lipid membranes of cells, including endosomal vesicles. Remarkably, as little as 30 picomolar UC-1V199/L induced cytokine synthesis in bone marrow derived mouse mononuclear cells. Figures 7-8 show data for IL-12 synthesis.

[0158] The TLR7 ligands that are purines or imidazoquinolines have a peculiar property, i.e., a biphasic dose response curve. At high concentrations, the drug fails to induce cytokine synthesis. The biphasic effect is observed in highly purified dendritic cells, and appears to be cell autonomous. However, the remarkable potency of UC-1V199/L enabled re-examination of the phenomenon, using pharmacologically acceptable drug concentrations (Figure 9). Maximal cytokine

production was observed with 10 nM UC-1V199/L while higher concentrations induced progressively less IL-12 (and TNF) release.

5 [0159] High and sustained concentrations of TLR7 agonists are known to induce refractoriness to TLR re-stimulation that can last 24 hours or more. Such a complex system of regulation is apparently part of a fail-safe mechanism that prevents cells and tissues from self-destruction during inflammatory responses. Thus, it was of interest to determine if concentrations of UC-1V199/L that failed to induce significant cytokine synthesis could nonetheless induce "TLR tolerance." Indeed, when bone marrow derived mononuclear cells were exposed to a non-activating concentration UC-1V199/L (1 μ M), and then re-stimulated 24 hours later with the same compound, with UC-1V150 or with pam3Cys (P3C, a TLR2 activator), they displayed a markedly diminished cytokine response. In contrast, the UC-1V199/L treated cells retained responsiveness to ligands of TLR3 and TLR4, which go through the TRIF pathway (results not shown). Preliminary experiments indicated that non-responsiveness was also induced *in vivo*. Thus, daily administration of UC-1V199/L, and related drugs, may suppress inflammation induced by MyD88-dependent stimuli, without the systemic side effects associated with TLR activation.

15 [0160] In one aspect, the conjugates of the application may be useful for preventing, inhibiting or treating asthma. Asthma is characterized by episodes of intermittent reversible airway constriction, bronchial smooth muscle hyperplasia and chronic inflammation. Atopic disease predisposes to asthma but up to half of the affected patients are not atopic. Other environmental risk factors for asthma include tobacco smoke and air pollutants. Moreover, disease flares in affected asthmatic patients may be triggered not only by allergens but also by airway irritants, temperature changes and infections.

20 [0161] The initial development of an allergic response is partly regulated by a balance between Th1 and Th2 lymphocytes, and their respective cytokines, especially the interferons and IL-4. Vaccination of animals with an allergen in conjunction with a TLR7 or TLR9 agonist preferentially expands allergen-specific Th1 memory cells. Consequently, subsequent immunization with antigen in conjunction with a Th2 biased adjuvant does not readily elicit an IgE response. Mice that were vaccinated with antigen and TLR7 or TLR9 agonists were resistant to experimental asthma.

25 [0162] A different approach is needed for the treatment of asthma with TLR agonists versus the prevention of asthma. In affected patients, the airways and pulmonary tissues are already infiltrated with a diverse population of inflammatory cells, including many subsets of lymphocytes, macrophages, dendritic cells, mast cells, eosinophils, and neutrophils. In this situation, TLR agonists can potentially exacerbate disease, by augmenting the release of inflammatory mediators such as TNF-alpha and IL-1. Indeed, the ability of various microbial agents to activate TLRs may explain why they trigger asthmatic attacks.

30 [0163] A TLR agonist for the prevention of asthma preferably is confined to the lungs but also maximizes the production of Th1 stimulating cytokines (interferons and IL-12), compared to TNF-alpha and IL-1. Both TLR7 and TLR9 are localized on the inner surfaces of the endosomal vesicles that are constantly synthesized and undergo maturation in dendritic cells. TLR9 activating oligonucleotides that are aggregated phosphodiester oligonucleotides stay longer in early endosomal vesicles and therefore induce more type I interferons than nonaggregated phosphorothioate oligonucleotides, which go to mature vesicles. The results imply that the spatial organization of the TLR agonist governs its trafficking and its pattern of induced cytokine synthesis. To prevent asthma, a stable, potent and molecularly characterized TLR agonist that trafficks to the early endosomes of dendritic cells and induces primarily Type I interferons is preferred.

35 [0164] To study the effect of conjugates of the application on allergic asthma, airway inflammation is induced by sensitizing mice via subcutaneous injection of 20 μ g of ovalbumin absorbed with 500 μ g alum per mouse in saline on day 0 and day 7. On days 16 and 21, mice are challenged *i.n.* with 5 μ g ovalbumin per mouse. Conjugates are administered *i.n.*, *p.o.* or *i.v.* at different time points prior to the first ovalbumin challenge on day 16. Twenty-four hours after the last challenge (day 22), airway responsiveness is measured, mice are sacrificed and BALF cells, lung and spleen samples collected. Naive mice and ovalbumin/alum-sensitized mice serve as controls. The total number of cells in BALF are counted and stained with Wright-Giemsa to determine numbers of eosinophils, lymphocytes, neutrophils and mast cells. Cytokine levels in the BALF are determined by Luminex assays. Airway responsiveness to methacholine is assessed 24 hours after the last challenge using a single chamber, whole body plethysmograph. The Penh, a dimensionless value that correlates well with pulmonary resistance measured by conventional two chamber plethysmography in ventilated mice, is used to monitor airway responsiveness.

45 [0165] The compounds of this application are administered in a therapeutically effective amount to a subject in need of treatment. Administration of the compositions of the invention can be via any of suitable route of administration, particularly parenterally, for example, intravenously, intra-arterially, intraperitoneally, intrathecally, intraventricularly, intraurethrally, intrastemally, intracranially, intramuscularly, or subcutaneously. Such administration may be as a single bolus injection, multiple injections, or as a short- or long-duration infusion. Implantable devices (e.g., implantable infusion pumps) may also be employed for the periodic parenteral delivery over time of equivalent or varying dosages of the particular formulation. For such parenteral administration, the compounds are preferably formulated as a sterile solution in water or another suitable solvent or mixture of solvents. The solution may contain other substances such as salts, sugars (particularly glucose or mannitol), to make the solution isotonic with blood, buffering agents such as acetic, citric, and/or phosphoric acids and their sodium salts, and preservatives.

[0166] The compounds of the invention can be formulated as pharmaceutical compositions and administered to a mammalian host, such as a human patient in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

[0167] Thus, the present compounds may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the active compound may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1 % of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

[0168] The tablets, troches, pills, capsules, and the like may also contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring may be added. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the active compound, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and devices.

[0169] The active compound may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active compound or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[0170] The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0171] Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

[0172] For topical administration, the present compounds may be applied in pure form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

[0173] Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Useful liquid carriers include water, alcohols or glycols or water-alcohol/glycol blends, in which the present compounds can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent pads, used to impregnate bandages and other dressings, or sprayed onto the affected area using pump-type or aerosol sprayers.

[0174] Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for application directly to the skin of the user.

[0175] In addition, in one embodiment, the invention provides various dosage formulations of the conjugates for inhalation delivery. For example, formulations may be designed for aerosol use in devices such as metered-dose inhalers, dry powder inhalers and nebulizers.

[0176] Examples of useful dermatological compositions which can be used to deliver the compounds of the invention to the skin are known to the art; for example, see Jacquet et al. (U.S. Pat. No. 4,608,392), Geria (U.S. Pat. No. 4,992,478), Smith et al. (U.S. Pat. No. 4,559,157) and Wortzman (U.S. Pat. No. 4,820,508).

[0177] Useful dosages of the compounds of the invention can be determined by comparing their *in vitro* activity, and *in vivo* activity in animal models. Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art; for example, see U.S. Pat. No. 4,938,949. The ability of a compound of the invention to act as a TLR agonist may be determined using pharmacological models which are well known to the art, including the procedures disclosed by Lee et al., Proc. Natl. Acad. Sci. USA, 100: 6646 (2003).

[0178] Generally, the concentration of the compound(s) of the invention in a liquid composition, such as a lotion, will be from about 0.1-25 wt-%, preferably from about 0.5-10 wt-%. The concentration in a semi-solid or solid composition such as a gel or a powder will be about 0.1-5 wt-%, preferably about 0.5-2.5 wt-%.

[0179] The active ingredient may be administered to achieve peak plasma concentrations of the active compound of from about 0.5 to about 75 μ M, preferably, about 1 to 50 μ M, most preferably, about 2 to about 30 μ M. This may be achieved, for example, by the intravenous injection of a 0.05 to 5% solution of the active ingredient, optionally in saline, or orally administered as a bolus containing about 1-100 mg of the active ingredient. Desirable blood levels may be maintained by continuous infusion to provide about 0.01-5.0 mg/kg/hr or by intermittent infusions containing about 0.4-15 mg/kg of the active ingredient(s).

[0180] The amount of the compound, or an active salt or derivative thereof, required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being treated and the age and condition of the patient and will be ultimately at the discretion of the attendant physician or clinician. In general, however, a suitable dose will be in the range of from about 0.5 to about 100 mg/kg, e.g., from about 10 to about 75 mg/kg of body weight per day, such as 3 to about 50 mg per kilogram body weight of the recipient per day, preferably in the range of 6 to 90 mg/kg/day, most preferably in the range of 15 to 60 mg/kg/day.

[0181] The compound is conveniently administered in unit dosage form; for example, containing 5 to 1000 mg, conveniently 10 to 750 mg, most conveniently, 50 to 500 mg of active ingredient per unit dosage form.

[0182] The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or more sub-doses per day. The sub-dose itself may be further divided, e.g., into a number of discrete loosely spaced administrations; such as multiple inhalations from an insufflator or by application of a plurality of drops into the eye. The dose, and perhaps the dose frequency, will also vary according to the age, body weight, condition, and response of the individual patient. In general, the total daily dose range for a compound or compounds of formula (I), for the conditions described herein, may be from about 50 mg to about 5000 mg, in single or divided doses. Preferably, a daily dose range should be about 100 mg to about 4000 mg, most preferably about 1000-3000 mg, in single or divided doses, e.g., 750 mg every 6 hr of orally administered compound. This can achieve plasma levels of about 500-750 μ M, which can be effective to kill cancer cells. In managing the patient, the therapy should be initiated at a lower dose and increased depending on the patient's global response.

[0183] As described above, compositions that contain a compound of the invention, are useful in the treatment or prevention of a disease or disorder in, for example, humans or other mammals (e.g., bovine, canine, equine, feline, ovine, and porcine animals), and perhaps other animals as well. Depending on the particular compound, the composition will, for example, be useful for treating cancer, an infection, enhancing adaptive immunity (e.g., antibody production, T cell activation, etc.), as vaccines, and/or stimulating the central nervous system.

[0184] The invention will be further described by the following non-limiting examples.

Example I

[0185] Processes for preparing compounds of formula (I) are provided as further embodiments of the invention and are illustrated by the following procedures in which the meanings of the generic radicals are as given above unless otherwise qualified.

[0186] General Chemistry. Reagents and solvents were acquired from Aldrich, Milwaukee, WI. Uncorrected melting points were determined on a Laboratory Device Mel-Temp II capillary melting point apparatus. Proton nuclear magnetic resonance spectra were recorded on a Varian Unity 500 NMR spectrophotometer at 499.8 MHz or on a Varian Mercury NMR spectrophotometer at 400.06 MHz. The chemical shifts were reported in ppm on the scale from the indicated reference. Positive and negative ion loop mass spectra were performed by Department of Chemistry UCSD, San Diego, CA. Elemental analyses were performed by NuMega Resonance Labs, San Diego, CA. Column chromatography was conducted on E Merck silica gel (230-400 mesh) with the indicated solvent system. Analytical thin layer chromatography (TLC) was conducted on silica gel 60 F-254 plates (EM Reagents).

[0187] Preparation of 4-(2,6-dichloropurin-9-ylmethyl)benzonitrile. 2,6-dichloro-9H-purine (16 mmol) is dissolved in DMF (50 mL) and potassium carbonate (50 mmol) is added. α -Bromo-*p*-tolunitrile (22 mmol) is then added and the mixture is stirred at ambient temperature for 16 hours. After filtration to remove insoluble inorganic salts, the filtrate is poured into water (1500 mL) and extracted with ethyl acetate (2 x 400 mL), dried over magnesium sulfate and evaporated to yield a residue which is subjected to flash silica gel chromatography using 1:2:10 ethyl acetate/acetone/hexanes. Yield 3.33 g (69%). UV, NMR and MS were consistent with structure assignment.

[0188] Preparation of 4-(6-amino-2-chloropurin-9-ylmethyl)benzonitrile. The product above (1.9 g) is placed in a steel reaction vessel and methanolic ammonia (80 mL, 7 N) is added. The sealed vessel is heated at 60°C for 12 hours, cooled in ice and the solid product filtered off. Yield 1.09 g. UV, NMR and MS were consistent with assigned structure.

[0189] Preparation of 4-[6-amino-2-(2-methoxyethoxy)purin-9-ylmethyl]benzonitrile. Sodium salt of 2-methoxyethanol is generated by dissolving sodium metal (81 mg) in 2-methoxyethanol (30 mL) with heat. To this solution is added the product of example 2 (1.0 g) dissolved in methoxyethanol (300 mL, with heat). The reaction mixture is heated for 8 hours at 115°C bath temperature, concentrated *in vacuo* to near dryness and the residue partitioned between ethyl acetate and water. Flash silica gel chromatography of the organic layer using 5% methanol in dichloromethane gave 763 mg product. NMR is consistent with structure assignment.

[0190] Preparation of 4-[6-amino-8-bromo-2-(2-methoxyethoxy)purin-9-ylmethyl]benzonitrile. The product immediately above (700 mg) is dissolved in dichloromethane (400 mL) and bromine (7 mL) is added dropwise. The mixture is stirred overnight at room temperature and extracted with aqueous sodium thiosulfate (2 L of 0.1 M) solution and then with aqueous sodium bicarbonate (500 mL, saturated). The residue from the organic layer is chromatographed on silica gel using 3% methanol in dichloromethane) to yield 460 mg of bromo product. NMR, UV and MS are consistent with structure assignment.

[0191] Preparation of 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzonitrile. Sodium methoxide is generated by reaction of sodium metal (81 mg) in dry methanol (30 mL). The product immediately above (700 mg) is dissolved in dry dimethoxyethane and the temperature raised to 100°C. After overnight reaction, the mixture is concentrated *in vacuo* and the residue is chromatographed on silica using 5% methanol in dichloromethane. Yield 120 mg. NMR is consistent with structure assignment.

[0192] Preparation of Lithium N,N'-(dimethylethylenediamino)aluminum hydride. This reducing agent used to convert the nitrile to the aldehyde function is prepared essentially as described in Bull. Korean Chem. Soc., 23:1697 (2002). A 0.5 M solution in dry THF is prepared.

[0193] Preparation of 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzaldehyde. 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzonitrile (100 mg) is dissolved in dry THF (3 mL) and cooled to 0°C under argon. The aluminum hydride reagent generated above (0.72 mL) is added to the reaction flask and the mixture is stirred at 0-5°C for 1 hour and then quenched by addition of 3 M HCl. The mixture is then extracted with ethyl acetate and then dichloromethane and concentrated *in vacuo* to yield 85 mg. NMR is consistent with structure assignment.

[0194] Preparation of 4-[6-amino-8-hydroxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzaldehyde (UC-1V150). The product immediately above (800 mg) is combined with sodium iodide (504 mg) and acetonitrile (40 mL), and then chlorotrimethylsilane (0.5 mL) is slowly added. The mixture is heated at 70°C for 3.5 hours, cooled and filtered. The solid product is washed with water, then ether to yield 406 mg. NMR, UV, MS are consistent with structure assignment. This material is suitable for conjugation reactions between linkers and macromolecules.

[0195] Preparation of methyl 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzoate. The procedure is as described by Jayachitra, et al., Synth. Comm., 33:3461 (2003). 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzonitrile (1 mmol) is dissolved in dry methanol (5 mL) and freshly distilled BF₃ etherate (4 mmol) is added to the solution. The resulting mixture is refluxed under argon for 20 hours. The solvent is removed *in vacuo* and the residue is taken up in dichloromethane (10 mL) and extracted with dilute aqueous sodium bicarbonate (2 x 10 mL) and the organic layer is dried over magnesium sulfate. After evaporation the product is purified by silica gel column chromatography using 5% methanol in dichloromethane to yield 0.8 mmol.

[0196] Preparation of 4-[6-amino-8-hydroxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzoic acid. 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzoate (100 mg) is combined with sodium iodide (63 mg) and acetonitrile (10 mL), and then chlorotrimethylsilane (120 mL) is slowly added. The mixture is heated at 70°C for 6 hours, cooled and filtered. The solid product is washed with water, then ether to yield 51 mg.

[0197] Preparation of 2,5-dioxopyrrolidin-1-yl 4-[6-amino-8-hydroxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzoate. 4-[6-amino-8-methoxy-2-(2-methoxyethoxy)purin-9-ylmethyl]benzoate (2 mmol) is dissolved in dichloromethane or dioxane (10 mL) and EDC (2 mmol) is added. To this solution is added N-hydroxysuccinimide (2 mmol) and resulting mixture is stirred at room temperature for 1 hour. The mixture is taken to dryness *in vacuo* and the crude product is purified by silica gel chromatography to yield 2 mmol of product that is suitable for conjugation reactions involving primary amines.

Example II

[0198] UC-1V 150 was covalently coupled to MSA first modified with a succinimidyl 6-hydrazino-nicotinamide acetone hydrazone (SANH) linker to yield a stable molecule with a characteristically altered UV spectrum. The UC-1V150/MSA conjugate was identified by a UV absorption peak at 342 nm due to hydrazone formation, whereas SANH alone absorbed at 322 nm. Quantification of UC-1V150 molecules conjugated per MSA was extrapolated from a standard curve of UC-1V150-SANH (Figure 1). Consistently the UC-1V150/MSA conjugates were obtained at a ratio of about 5:1. The biological studies reported here were done by using 5:1 UC-1V150/MSA.

[0199] Modification of MSA with SANH. 200 μ L of MSA (25 mg/mL) was mixed with 100 μ L of conjugation buffer (1M NaPi, pH=7.2) and 690 μ L of PBS. 844 μ g of SANH in 10 μ L of DFM (40-fold molar excess to MSA) was added to protein solution (final concentration of MSA in reaction mixture is 5 mg/mL). After gentle mixing reaction was proceeded at room temperature for 2 hours. To remove excess of SANH the reaction mixture was loaded on NAP-10 column equilibrated with PBS and modified MSA was eluted with 1.5 mL of PBS.

[0200] Attachment of IV150 to MSA modified with SANH. 460 μ g of IV150 dissolved in 10 μ L of DMF was added to MSA modified with SANH and the reaction mixture was incubated at RT overnight. To remove excess of IV150 the reaction mixture was firstly concentrated to 1 mL using micro-spin column (Millipore: BIOMAX 5K) and loaded on NAP-10 column as mentioned above.

[0201] TLR7 agonists were also conjugated to oligodeoxynucleotides (ODNs) (Figures 20-21), a virus (Figures 18-19), and to a lipid component which can then be incorporated into a liposome (Figures 24-25).

[0202] Synthesis of spatially regulated TLR7 agonist. Each conjugate is prepared by standard techniques well known in bioconjugation chemistry. Characterization of each by quantitative UV, LC/MS, and PAGE methods determines the "valence" or ratio of TLR agonist to its auxiliary group (macromolecule). From this information, the size and shape of the conjugates are easily estimated by modeling techniques. The diversity in size, shape, and valence of the conjugates is introduced through the selection of the macromolecule, represented in the structure scheme as R3. For example, when R3 is a dendrimer, such as of the common poly(amidoamine) variety, the number of surface functional groups for attachment of the TLR agonist is precisely defined based on the number of branching points or generations of that particular dendrimer. A first generation (G1) has 8 surface amino groups, a G2 has 16, and so on, thus resulting in a high level of control over valence and size of the conjugates (see Figure 2). Additionally, some dendrimer nanoparticles may contain both a targeting ligand and the TLR7 agonist. The TLR7 agonist-lipid conjugates may also have a variety of "valences" depending on the selection of the lipids. For example, the potent conjugate UV-1V199/L (Figure 6) was prepared by coupling a carboxy derivative of the TLR7 agonist (UC-1V199) to the ethanolamino group of the commercially available dioleanylphosphatidylethanolamine (DOPE).

[0203] These lipid conjugates are formulated into various liposome nanoparticles by combining with cholesterol, DOPE, and other lipids to produce particles having a hydrodynamic diameter of about 100 nm (Figures 24-25). The hexagons in the figure represent UC-1V199/L and related TLR7 agonist with phospholipid tails.

[0204] TLR7 agonists and dimers, as well as TLR conjugates have been shown to have cytokine releasing and/or cytokine activity *in vivo* as determined by assays such as those disclosed herein. For instance, imiquimod, broprimine, UC-1V138, UC-1V136, UC-1V150, UC-1X105, UC-1V199, UC-1W236, UC-1X51, UC-1W247, UC-1X113, UC-1V199/L, UC-1V150/BSA, conjugates of UC-1V150 with or without a linker and MSA, OVA, virions, and/or ODN, conjugates of UC-1V199 and DOPE, silica, lipid, or irradiated spores, and conjugates of UC-1V1043 and UC-1V1018 with OVA all have shown activity.

Example III**Materials and Methods**

[0205] Compound evaluation *in vitro*. The ability of TLR7 conjugates to stimulate and/or to inhibit cytokine production is assessed in murine bone marrow derived mononuclear cells (BMDM) that are highly enriched in dendritic cells, as well as in human peripheral blood mononuclear cells (PBMC) cells. BMDM are plated in 96 well plates and treated in triplicate with vehicle or various doses starting from 10 μ M diluted in 3-fold increments down to picomolar concentrations. After 24 hours the supernatants are harvested and assayed for up to 30 different cytokines, chemokines and other mediators, using a Luminex bead assay system, and commercially available reagents. The cytokine/chemokine ELISA results are supplemented with quantitative mRNA expression measurements and with two-dimensional phosphoprotein analyses, to gain insight into the scope and mechanism of tolerance induction. At the time of supernatant harvest, media are replaced in the wells with MTT, as a colorimetric assessment of cell survival. Human PBMC are isolated from commercial blood packs and treated similarly.

[0206] To assess the trafficking of the TLR agonist-conjugated nanoliposomes and dendrimers, the respective nanoparticles are loaded or modified with a fluorochrome. Subcellular localization is determined microscopically, in some

cases in cells that have been treated with inhibitors of endosomal maturation.

[0207] To compare the anti-inflammatory activities of the TLR7 conjugates with different auxiliary groups, BMDM are treated first with the most potent compounds, at previously determined concentrations that had minimal effects on pro-inflammatory cytokine stimulation (TNF α , IL-1). After 24 hours, the medium is replaced, and the cells are challenged with activating ligands of different TLR family members (Pam3Cys for TLR2, poly(I:C) for TLR3, LPS for TLR4, flagellin for TLR5, Malp-2 for TLR6, UC-1V150 for TLR7, R848 for TLR7/8, CpG oligonucleotides for TLR9, and the like) at concentrations that effectively induce cytokine production in mock treated cells. The cells are assessed by multiplex immunoassay, quantitative PCR and phosphoprotein blotting. To better understand the kinetics of induction and maintenance of tolerance, TLR7 conjugate-primed cells are also challenged at different time intervals and analyzed for the pattern of cytokine production.

[0208] Compound evaluation *in vivo*. Production of bronchoalveolar lavage fluid (BALF) versus systemic cytokines after administration to the airways of mice is assessed. Anesthetized female age-matched C57BL/6 mice are administered nasally (i.n.) orally (p.o.), or intravenously (i.v.), with various amounts of the TLR7 conjugates as previously described, or with the liposomes or dendrimers in appropriate vehicles. After recovery and at different time points, sera and BALF are collected and analyzed for cytokines and chemokines by Luminex assay. The weights, temperatures, and fluid intake patterns of the treated animals are recorded, as a clinical surrogate for a systemic "cytokine syndrome."

[0209] Subsequent experiments assess the ability of the different agents to produce local and systemic refractoriness (TLR tolerance) to TLR activation after high dose administration i.n., p.o. or i.v., as determined by sera and BALF cytokines. High doses of the various TLR7 conjugates, which do not induce significant cytokines *in vivo*, nor clinical signs of a cytokine syndrome, are selected. Mice are treated with the selected high doses given by the different routes of administration, and then challenged with activators of different TLRs at various time points. Serum and BALF are collected and analyzed and clinical symptoms are recorded. The anti-inflammatory activities of the conjugates are confirmed with a lethal shock model previously used to study LPS and CpG. In this model, Balb/c mice that have been previously injected i.p. with D-galactosamine succumb after systemic challenge with different TLR activators, due to cytokine stimulation and liver damage. Active anti-inflammatory drugs fail to induce clinical symptoms in sensitized animals, and will also prevent shock caused by other TLR ligands. With a defined endpoint, this model is especially useful for determining the kinetics and duration of TLR tolerance.

Example IV

Materials and Methods

[0210] Mice. Female C57BL/6 mice (5-6 weeks of age) were obtained from Harlan West Coast (Germantown, CA), and female A/J mice (6-8 weeks of age) were purchased from The Jackson Laboratories (Bar Harbor, ME). A/J mice were used for infection with the Sterne strain of *B. anthracis* (Kenney et al., J. Infect. Dis., 190:774 (2004)). The mice were bred and maintained under standard conditions in the University of California at San Diego Animal Facility, which is accredited by the American Association for Accreditation of Laboratory Animal Care. All animal protocols received prior approval by the Institutional Review Board. For the H1N1 influenza study, female BALB/c mice (16-18 g) were obtained from Charles River Laboratories (Wilmington, MA) and maintained in the American Association for Accreditation of Laboratory Animal Care-accredited Laboratory Animal Research Center of Utah State University.

[0211] *In Vitro* Stimulation of BMDM. BMDM were isolated from various strains of mice were seeded in 96-well plates at a density of 5×10^4 cells per well. Compounds were added to 10-day-old cultures at a final concentration ranging from 0.01 to 10 μ M or as otherwise indicated. After 24 hours of incubation, culture supernatants were collected and assayed for cytokine inductions by either sandwich ELISA (BD Pharmingen, San Diego, CA) or multiplex Luminex (Austin, TX) assay using the Beadlyte Mouse MultiCytokine customized kit (Upstate, Charlottesville, VA, and eBiosciences, San Diego, CA), according to the manufacturer's instructions.

[0212] Administration of Compounds to Mice. Female age-matched C57BL/6 mice were injected with 100 μ L of saline solution containing UC-1V150 or UC-1V150/MSA, each containing the equivalent of 0.38-38 nmol of the pharmacore via the tail vein. For intrapulmonary administration, mice were anesthetized with i.p. Avertin solution and shaved around the neck area. The trachea were exposed with a small incision and injected with 50 μ L of saline solution containing various amounts of UC-1V 150/MSA or the unconjugated drug. After recovery and at different time points, serum and BALF were collected and analyzed for IL-6, IL-12p40, IFN- γ , RANTES, and MCP-1 by Luminex assay. In other experiments, mice were anesthetized with an intramuscular ketamine/xylene solution and administered the same amount of UC-1V150/MSA in i.t. doses of 50 μ l or i.n. doses of 20 μ L. Because similar cytokine levels were observed in the BALF 24 hours after administration by either method, the more convenient i.n. route was used in infectious model studies.

[0213] Infection of A/J Mice with *B. anthracis* Spores. Spores were prepared from the Sterne strain of *B. anthracis* (pXO1⁺pXO2⁻) as previously described (Sabet et al., FEMS Immunol. Med. Microbiol., 47:369 (2006); Guidi-Rontani et al., Mol. Microbiol., 42:931 (2001)). Purified spores were stored in PBS at 1×10^8 to 4×10^8 cfu/mL at 4°C. Before

infection, the spores were heated to 65°C for 30 minutes to initiate germination. A/J mice were anesthetized intramuscularly with ketamine/xylene solution and administered i.n. with 0.75 nmol of UC-1V150 or UC-1V150/MSA per mouse 1 day before anthrax infection. Control mice received saline only or saline containing MSA at equivalent amounts as in UC-1V 150/MSA. Infection was carried out i.n. with 2×10^5 to 8×10^5 spores of *B. anthracis* in a 20 μ L volume. Survival was observed for 13 days, because the majority of the saline-treated mice died within 3-6 days. Results were obtained from eight mice per group.

[0214] Infection of Balb/c mice with influenza virus. Influenza A/New Caledonia/20/99 (H1N1) virus was obtained from the Centers for Disease Control and Prevention (Atlanta, GA). The virus was propagated twice in Madin Darby canine kidney (MDCK) cells, further passaged 7 times in mice to make it virulent followed by another passage in cell culture to amplify it. Mice were anesthetized i.p. with ketamine (100 mg/kg) and infected i.n. with virus at approximately $10^{5.0}$ cell culture infectious doses per mouse in a 50 μ L inoculum volume. A single intranasal dose of 75 μ L in either saline alone or containing UC-1V10/MSA to 5 nmole per mouse was given 24 hours prior to virus exposure. Ten infected mice per treated group and 20 placebo control animals were followed for survival for 21 days.

[0215] Statistics. Cytokine levels were compared by the Mann Whitney U-test with $p \leq 0.05$ to determine statistical significance. Kaplan-Meier survival curves and log rank tests were performed using GraphPad Prism software version 4.0c (San Diego, CA) to compare differences in survival.

Results

[0216] Potent *in vitro* and *in vivo* cytokine release in response to UC-1V 150/MSA conjugates. Incubation of bone-marrow-derived macrophages (BMDM) with UC-1V150 alone stimulated cytokine release (Figure 10). When conjugated to MSA, similar or higher levels of cytokines were detected with a 10-fold lower equivalent concentration of the TLR7 agonist. Experiments with TLR transformants, performed as described previously, confirmed that UC-1V150, similar to the compound lacking the aldehyde modification (UC-1V136), was a specific TLR7 agonist (Lee et al., Proc. Natl. Acad. Sci. USA, 103:1828 (2006)). After i.v. injection into mice, UC-1V150 induced serum cytokine levels that peaked at about 2 hours after injection and then quickly declined to near background levels (data not shown). Comparison of the cytokine production profiles of UC-1V150 versus the UC-1V150/MSA 2 hours after i.v. injection at various dosages demonstrated that the MSA conjugate enhanced the potency by 10- to 100-fold (Figure 11). Sera from saline or MSA control groups revealed little or no detectable cytokine levels (data not shown).

[0217] UC-1V150/MSA Conjugates Provide Prolonged and Localized Pulmonary Activity. To ensure adequate delivery of the TLR7 agonists to the respiratory system, the drugs were initially directly into the trachea. Substantial cytokine induction was found in bronchial alveolar lavage fluid (BALF) extracted from mice treated intratracheally (i.t.) with UC-1V 150/MSA, whereas serum cytokines were very low and near background levels in the same animals (Figure 12). In marked contrast, similar levels of cytokine were observed in both BALF and sera of mice injected i.t. with small-molecule TLR7 agonists, which sometimes induced behavioral changes, such as hair standing on end and shivering, suggestive of a cytokine syndrome (Table 2). Subsequent studies with UC-1V150 showed that intranasal (i.n.) delivery also induced selective cytokine production in the BALF, probably due to drug aspiration. Accordingly, i.n. administration was used to evaluate the UC-1V150 conjugates in two infectious animal models of pneumonitis. Mice pretreated i.n. with UC-1V 150/MSA one day before infection with *B. anthracis* spores had an extended mean survival of 7.5 days compared with 5 days in control mice ($P < 0.025$) (Figure 14A). In contrast, no significant difference was observed in mice treated with either saline; the equivalent amount of MSA, or with UC-1V150 alone. These data confirmed that the UC-1V150 conjugate, but not the free drug, had intrapulmonary immunotherapeutic activity. Thus, conjugation of the TLR7 agonist to MSA enhanced its potency and reduced its toxicity after local delivery to the respiratory tract.

[0218] In another study, BALB/c mice were pretreated i.n. with the UC-1V150/MSA conjugate 1 day before influenza virus infection (H1N1 strain). The mean survival of the treated mice was extended to 11.5 days compared with 7 days in untreated controls ($P < 0.0001$) (Figure 14B). Together these results suggest that conjugation of the TLR7 agonist to MSA enhanced its potency and reduced its toxicity after local delivery to the respiratory tract.

[0219] UC-1V 150/MSA was administered i.n. prior to anthrax infection followed by treatment with ciprofloxacin (25 mg/kg) on day 4. Placebo treatment followed by ciprofloxacin treatment resulted in about 15-25% survival, while treatment with a conjugate and ciprofloxacin resulted in about 90% survival. Thus, the conjugate is particularly useful as a coadjuvant with an anthrax vaccine.

Discussion

[0220] The compound UC-1V150 is one of the most potent and versatile synthetic small-molecule TLR7 ligands yet discovered because (i) it is active at nanomolar concentrations; (ii) it can be coupled to a variety of macromolecules with enhancement of activity in some cases; and (iii) its pharmacokinetic properties can be changed by modification of the auxiliary groups. The TLR7-protein conjugate UC-1V150/MSA was characterized as having approximately five small

molecules covalently linked to each MSA protein molecule. The conjugate retained TLR7 agonist activity and indeed was both more potent and had a longer duration of action, compared with the free monomeric drug. Moreover, this conjugate could be delivered effectively to the respiratory system by i.n. or i.t. administration. Drug delivery by i.n. proved to be effective in a mouse model of a bacterial infection. When considering delivery to the respiratory system, a potentially important advantage of preparing the TLR7 agonists as conjugates of macromolecules is that systemic side effects may be avoided by confining the immunostimulatory activity to the local mucosal environment.

[0221] The macromolecular conjugate would be expected to be absorbed into the systemic circulation more slowly than the free drug and, indeed, may be avidly scavenged by resident macrophages and dendritic cells expressing TLR7. Accordingly, the conjugate should mitigate the type of severe side effects that have been associated with systemic delivery of TLR7/8 agonists. The UC-1V150/MSA conjugate may also provide beneficial immunotherapeutic activity when administered to mucosal sites, such as the genitourinary and gastrointestinal tracts, for the control of infectious, allergic, or malignant diseases. The macromolecular carrier of the TLR7 agonist may also provide an improved approach for selective delivery of the immunotherapeutic to a specific organ or tissue. For example, the lipid conjugates of UC-1V150 can be incorporated into liposomes of different size and composition, whereas protein conjugates of the TLR7 agonist may target different dendritic cell subsets. Differences in the intracellular trafficking of the UC-1V150 conjugate may induce distinct patterns of cytokine production, analogous to the effects observed with TLR9-activating oligonucleotides (Rothenfusser et al., Hum. Immunol., 63:111 (2002)).

[0222] One potential problem that has been observed with drugs conjugated to proteins is the development of antibodies against the low-molecular-weight hapten-like portion of the molecule. However, UC-1V150, unlike the TLR7/8 vaccine conjugates studied earlier, has a simple adenine-like structure that is unlikely to induce hypersensitivity reactions. Indeed, anti-UC-1V150 antibodies were not observed after administration of the protein conjugates, except after repeated administration of a keyhole limpet hemocyanin carrier in complete Freud's adjuvant (unpublished data).

[0223] New agents for the prevention and treatment of influenza virus infections are being sought, particularly with the spread of highly pathogenic strains from Asia. Morbidity and mortality from commonly circulating strains is high each year. Treatment of the infection can be accomplished by approved antiviral drugs, which are moderately effective if started early. Enhancement of the immune system is also being investigated as a strategy that could accelerate protective antiviral responses, especially in immune compromised hosts. It is possible that systemic immune activation via TLR signaling does not create a local cytokine and chemokine gradient required to mobilize immune cells to the site of infection. In support of this hypothesis, the unconjugated UC-1V150, which is rapidly absorbed through the mucosa, failed to protect mice from *B. anthracis* infection, whereas the UC-1V150 conjugate was effective.

[0224] *B. anthracis* has become an agent of bioterrorism. A rapid response against microbial pathogens is critical for effective biodefense. In general an antibody or cellular immune response may protect against these pathogens; however, generating these protective responses quickly requires prior exposure to specific antigens for each organism. Although it is known that influenza virus engages TLR7 (Barchet et al., Eur. J. Immunol., 35:360 (2005)), bacterial anthrax most likely can engage TLR2, TLR4, and TLR9. In addition to being a common signaling intermediary for the TLRs, MyD88 has also been shown to be necessary for resistance to infection in a mouse model of anthrax (Hughes et al., Infect. Immun., 73:7535 (2005)). Because the UC-1V150 conjugate works effectively as an adjuvant against infections that use different pathways, it can be applied as a biodefense strategy that would not need be specific to the antigens of a particular microbe and that would be useful in mixed as well as single agent attacks.

Example V


[0225] There is no known SA vaccine that is potent enough or that can act quickly enough to prevent SA infections in "at-risk" patients prior to hospitalization. A single injection of a potent TLR7 agonist and killed gram-positive bacteria, e.g., SA, or a subunit thereof, may boost protective immunity to the bacteria within one week of administration. The injection may include, for example, 1) a TLR7 agonist such UC-IV199 conjugated directly to free amino groups on killed gram-positive bacteria, 2) a TLR7 agonist such as UC-IV199 conjugated to albumin in combination with killed gram-positive bacteria, 3) a TLR7 agonist such as UC-IV 199 conjugated to a recombinant gram-positive bacterial protein, or 4) a TLR7 agonist such as UC-IV199 conjugated to gram-positive bacterial polysaccharides (e.g., via a linker known to the art, such as that used the StaphVax®).

[0226] As described hereinabove, a TLR7 agonist was conjugated to lethally irradiated spores of the Sterne vaccine strain of *Bacillus anthracis* (BA). Like SA, BA is a gram-positive bacteria. Compared to spores alone, the conjugated bacterium was a potent activator of mouse bone marrow derived macrophages (BMDM) as measured by cytokine (IL-12 and IL-6) secretion. In another experiment, a single injection into mice of lethally irradiated spores of the Sterne strain of BA, mixed with a TLR7 agonist conjugated to mouse albumin (MSA), protected the animals against lethal intrapulmonary BA challenge given only six days later. In contrast, injection of the animals with BA spores alone, or with BA plus a conventional adjuvant, cholera toxin (CT), did not protect the animals. Thus, a TLR7-agonist albumin/irradiated spore vaccine induced protective immunity to *Bacillus anthracis* within 6 days. This rapidity of response in a naive animal

was totally unexpected. The same vaccine technology will likely protect humans from hospital-acquired SA infection.

Claims

- 5
1. A composition for use in a method to prevent or inhibit a gram-positive bacterial infection in a mammal, comprising administering to the mammal an effective amount of the composition comprising a bacterial antigen of a gram-positive bacteria and an amount of a compound having formula (IA):



wherein X¹ is -O-, -S-, or -NR^c-;

wherein R^c is hydrogen, C₁₋₁₀alkyl, or substituted C₁₋₁₀alkyl, or R^c and R¹ taken together with the nitrogen atom can form a heterocyclic ring or a substituted heterocyclic ring, wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

R¹ is hydrogen, (C₁₋₁₀)alkyl, substituted (C₁₋₁₀)alkyl, C₆₋₁₀aryl, or substituted C₆₋₁₀aryl, C₅₋₉heterocyclic, substituted C₅₋₉heterocyclic; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

each R² is independently hydrogen, -OH, (C₁₋₆)alkyl, substituted (C₁₋₆)alkyl, (C₁₋₆)alkoxy, substituted (C₁₋₆)alkoxy, -C(O)-(C₁₋₆)alkyl (alkanoyl), substituted -C(O)-(C₁₋₆)alkyl, -C(O)-(C₆₋₁₀)aryl (aroyl), substituted -C(O)-(C₆₋₁₀)aryl, -C(O)OH (carboxyl), -C(O)O(C₁₋₆)alkyl (alkoxycarbonyl), substituted -C(O)O(C₁₋₆)alkyl, -NR^aR^b, -C(O)NR^aR^b (carbamoyl), substituted -C(O)NR^aR^b, halo, nitro, or cyano; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;


each R^a and R^b is independently hydrogen, (C₁₋₆)alkyl, (C₃₋₈)cycloalkyl, (C₁₋₆)alkoxy, halo(C₁₋₆)alkyl, (C₃₋₈)cycloalkyl(C₁₋₆)alkyl, (C₁₋₆)alkanoyl, hydroxy(C₁₋₆)alkyl, aryl, aryl(C₁₋₆)alkyl, Het, Het (C₁₋₆)alkyl, or (C₁₋₆)alkoxycarbonyl;

X² is a bond or a linking group; n = 1, 2, 3 or 4; and

R³ is a macromolecule comprising a heterologous peptide, heterologous protein, heterologous lipid, bead, such as a polystyrene bead, heterologous nucleic acid molecule or dendrimer;

or a pharmaceutically acceptable salt thereof, including hydrates thereof.

- 45
50
55
2. A compound for use in a method to prevent or inhibit a gram-positive bacterial infection in a mammal, comprising administering to the mammal an effective amount of the compound having formula (IB):



(IB)

wherein X^1 is -O-, -S-, or -NR^c-;

wherein R^c is hydrogen, C₁₋₁₀alkyl, or substituted C₁₋₁₀alkyl, or R^c and R¹ taken together with the nitrogen atom can form a heterocyclic ring or a substituted heterocyclic ring, wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆-cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

R¹ is hydrogen, (C_{1-C10})alkyl, substituted (C_{1-C10})alkyl, C₆₋₁₀aryl, or substituted C₆₋₁₀aryl, C₅₋₉heterocyclic, substituted C₅₋₉heterocyclic; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

each R² is independently hydrogen, -OH, (C_{1-C6})alkyl, substituted (C_{1-C6})alkyl, (C_{1-C6})alkoxy, substituted (C_{1-C6})alkoxy, -C(O)-(C_{1-C6})alkyl (alkanoyl), substituted -C(O)-(C_{1-C6})alkyl, -C(O)-(C_{6-C10})aryl (aroyl), substituted -C(O)-(C_{6-C10})aryl, -C(O)OH (carboxyl), -C(O)O(C_{1-C6})alkyl (alkoxycarbonyl), substituted -C(O)O(C_{1-C6})alkyl, -NR^aR^b, -C(O)NR^aR^b (carbamoyl), substituted -C(O)NR^aR^b, halo, nitro, or cyano; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

each R^a and R^b is independently hydrogen, (C_{1-C6})alkyl, (C₃₋₈)cycloalkyl, (C_{1-C6})alkoxy, halo(C_{1-C6})alkyl, (C₃₋₈)cycloalkyl(C_{1-C6})alkyl, (C_{1-C6})alkanoyl, hydroxy(C_{1-C6})alkyl, aryl, aryl(C_{1-C6})alkyl, Het, Het (C_{1-C6})alkyl, or (C_{1-C6})alkoxycarbonyl;

X² is a bond or a linking group; n = 1, 2, 3 or 4; and


R³ is the gram-positive bacteria, an isolated antigenic protein or peptide of the gram-positive bacteria, or an isolated polysaccharide of the gram-positive bacteria;

or a pharmaceutically acceptable salt thereof, including hydrates thereof, wherein the gram-positive bacterium is not Bacillus or Listeria.

3. The composition or compound for use of claim 1 or 2, wherein X¹ is sulfur atom.
4. The composition or compound for use of claim 1 or 2, wherein X¹ is oxygen atom.
5. The composition or compound for use of claim 1 or 2, wherein X¹ is -NR^c-, wherein R^c is hydrogen, C₁₋₆ alkyl or substituted C₁₋₆ alkyl; wherein the alkyl substituents are C₃₋₆cycloalkyl, hydroxy, C₁₋₆alkoxy, amino, cyano, or aryl.
6. The composition or compound for use of claim 5, wherein X¹ is -NH-.
7. The composition or compound for use of any one of claims 1 to 6, wherein R¹ and R^c taken together form a heterocyclic ring or a substituted heterocyclic ring.
8. The composition or compound for use of claim 7, wherein R¹ and R^c taken together form a substituted or unsubstituted morpholino, piperidino, pyrrolidino, or piperazino ring.
9. The composition or compound for use of any one of claims 1 to 6, wherein R¹ is hydrogen, C₁₋₄alkyl, or substituted C₁₋₄alkyl.
10. The composition or compound for use of claim 9, wherein R¹ is hydrogen, CH₃-, CH₃-CH₂-, CH₃CH₂CH₂-, hydroxyC₁₋₄alkylene, or C₁₋₄alkoxyC₁₋₄alkylene.

EP 2 125 007 B9


11. The composition or compound for use of claim 10, wherein R¹ is hydrogen, CH₃-, CH₃-CH₂-, CH₃-O-CH₂CH₂- or CH₃-CH₂-O-CH₂CH₂-.
12. The composition or compound for use of any one of claims 1 to 11, wherein R² is hydrogen, halogen, or C₁₋₄alkyl.
13. The composition or compound for use of claim 12, wherein R² is hydrogen, chloro, bromo, CH₃-, or CH₃-CH₂-.
14. The composition or compound for use of any one of claims 1 to 13, wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkylene, C₃₋₆cycloalkyl, amino, cyano, halogen, or aryl.
15. The composition or compound for use of any one of claims 1 to 14, wherein X² is a bond or a chain having up to about 24 atoms; wherein the atoms are selected from the group consisting of carbon, nitrogen, sulfur, non-peroxide oxygen, and phosphorous.
16. The composition or compound for use of claim 15, wherein X² is a bond or a chain having from about 4 to about 12 atoms.
17. The composition or compound for use of claim 15, wherein X² is a bond or a chain having from about 6 to about 9 atoms.
18. The composition or compound for use of claim 15, wherein X² is



19. The composition for use of any one of claims 1 or 3 to 18, wherein the macromolecule is a phospholipid, a liposome, or a dendrimer.
20. The composition for use of any one of claims 1 or 3 to 18, wherein the heterologous peptide has from 2 to about 200 amino acid residues.
21. The compound for use of any one of claims 2 to 18, wherein the peptide has from 2 to about 200 amino acid residues.
22. The composition for use of any one of claims 1 or 3 to 18, wherein the heterologous peptide has from 10 to about 200 amino acid residues.
23. The compound for use of any one of claims 2 to 18, wherein the peptide has from 10 to about 200 amino acid residues.
24. The composition for use of any one of claims 1 or 3 to 18, wherein the macromolecule comprises carbohydrate.
25. The composition for use of any one of claims 1 or 3 to 18, wherein the nucleic acid molecule comprises DNA, RNA or PNA.
26. The composition or compound for use of any one of claims 1 to 25, wherein the gram-positive bacteria is *Staphylococcus aureus*.
27. The composition or compound for use of any one of claims 1 to 26, wherein the mammal is a human.
28. The composition or compound for use of any one of claims 1 to 27, wherein the mammal is at-risk of a gram-positive bacterial infection.
29. The composition or compound for use of any one of claims 1 to 28, wherein the mammal is immunocompromised.
30. The composition or compound for use of any one of claims 1 to 29, wherein the mammal has an effective immune

response to the administration by about 6 days after administration.

31. The composition for use of any one of claims 1 or 3 to 30, wherein the composition comprises a preparation of killed gram-positive bacteria.
32. The composition or compound for use of any one of claims 1 to 31, wherein the amount is effective to prevent infection by about 6 days after administration.
33. A vaccine comprising a composition comprising a bacterial antigen of a *S. aureus* bacteria and an amount of a compound having formula (IA):



wherein X¹ is -O-, -S-, or -NR^c-;

wherein R^c is hydrogen, C₁₋₁₀alkyl, or substituted C₁₋₁₀alkyl, or R^c and R¹ taken together with the nitrogen atom can form a heterocyclic ring or a substituted heterocyclic ring, wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

R¹ is hydrogen, (C₁₋₁₀)alkyl, substituted (C₁₋₁₀)alkyl, C₆₋₁₀aryl, or substituted C₆₋₁₀aryl, C₅₋₉heterocyclic, substituted C₅₋₉heterocyclic; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

each R² is independently hydrogen, -OH, (C₁₋₆)alkyl, substituted (C₁₋₆)alkyl, (C₁₋₆)alkoxy, substituted (C₁₋₆)alkoxy, -C(O)-(C₁₋₆)alkyl (alkanoyl), substituted -C(O)-(C₁₋₆)alkyl, -C(O)-(C₆₋₁₀)aryl (aroyl), substituted -C(O)-(C₆₋₁₀)aryl, -C(O)OH (carboxyl), -C(O)O(C₁₋₆)alkyl (alkoxycarbonyl), substituted -C(O)O(C₁₋₆)alkyl, -NR^aR^b, -C(O)NR^aR^b (carbamoyl), substituted -C(O)NR^aR^b, halo, nitro, or cyano; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;


each R^a and R^b is independently hydrogen, (C₁₋₆)alkyl, (C₃₋₈)cycloalkyl, (C₁₋₆)alkoxy, halo(C₁₋₆)alkyl, (C₃₋₈)cycloalkyl(C₁₋₆)alkyl, (C₁₋₆)alkanoyl, hydroxy(C₁₋₆)alkyl, aryl, aryl(C₁₋₆)alkyl, Het, Het(C₁₋₆)alkyl, or (C₁₋₆)alkoxycarbonyl;

X² is a bond or a linking group; n = 1, 2, 3 or 4; and

R³ is a macromolecule comprising a heterologous peptide, heterologous protein, heterologous lipid, bead, such as a polystyrene bead, heterologous nucleic acid molecule or dendrimer;

or a pharmaceutically acceptable salt thereof, including hydrates thereof.

34. A vaccine comprising a compound having formula (IB):



wherein X¹ is -O-, -S-, or -NR^c-;

wherein R^c is hydrogen, C₁₋₁₀alkyl, or substituted C₁₋₁₀alkyl, or R^c and R¹ taken together with the nitrogen atom can form a heterocyclic ring or a substituted heterocyclic ring, wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

R¹ is hydrogen, (C_{1-C10})alkyl, substituted (C_{1-C10})alkyl, C₆₋₁₀aryl, or substituted C₆₋₁₀aryl, C₅₋₉heterocyclic, substituted C₅₋₉heterocyclic; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

each R² is independently hydrogen, -OH, (C_{1-C6})alkyl, substituted (C_{1-C6})alkyl, (C_{1-C6})alkoxy, substituted (C_{1-C6})alkoxy, -C(O)-(C_{1-C6})alkyl (alkanoyl), substituted -C(O)-(C_{1-C6})alkyl, -C(O)-(C_{6-C10})aryl (aroyl), substituted -C(O)-(C_{6-C10})aryl, -C(O)OH (carboxyl), -C(O)O(C_{1-C6})alkyl (alkoxycarbonyl), substituted -C(O)O(C_{1-C6})alkyl, -NR^aR^b, -C(O)NR^aR^b (carbamoyl), substituted -C(O)NR^aR^b, halo, nitro, or cyano; wherein the substituents on the alkyl, aryl or heterocyclic groups are hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkylene, C₁₋₆alkoxy, C₃₋₆cycloalkyl, C₁₋₆alkoxyC₁₋₆alkylene, amino, cyano, halogen, or aryl;

each R^a and R^b is independently hydrogen, (C_{1-C6})alkyl, (C₃₋₈)cycloalkyl, (C_{1-C6})alkoxy, halo(C_{1-C6})alkyl, (C₃₋₈)cycloalkyl(C_{1-C6})alkyl, (C_{1-C6})alkanoyl, hydroxy(C_{1-C6})alkyl, aryl, aryl(C_{1-C6})alkyl, Het, Het (C_{1-C6})alkyl, or (C_{1-C6})alkoxycarbonyl;


X² is a bond or a linking group; n = 1, 2, 3 or 4; and

R³ is the *S. aureus* bacteria, an isolated antigenic protein or peptide of the *S. aureus* bacteria, or an isolated polysaccharide of the *S. aureus* bacteria;

or a pharmaceutically acceptable salt thereof, including hydrates thereof.

Patentansprüche

1. Eine Zusammensetzung zur Verwendung in einem Verfahren zur Verhinderung oder Inhibierung einer Infektion durch gram-positive Bakterien in einem Säugetier, umfassend die Verabreichung an das Säugetier einer effektiven Menge der Zusammensetzung, umfassend ein bakterielles Antigen eines gram-positiven Bakteriums und eine Menge einer Verbindung, welche die Formel (IA) aufweist:



wobei X¹ -O-, -S- oder -NR^c- ist;

wobei R^c Wasserstoff, C₁₋₁₀Alkyl oder substituiertes C₁₋₁₀Alkyl ist, oder R^c und R¹ zusammen mit dem Stickstoffatom einen heterozyklischen Ring oder einen substituierten heterozyklischen Ring bilden können, wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;

R¹ Wasserstoff, (C₁-C₁₀)Alkyl, substituiertes (C₁-C₁₀)Alkyl, C₆₋₁₀Aryl oder substituiertes C₆₋₁₀Aryl, ein C₅₋₉Heterozyklus, ein substituiertes C₅₋₉Heterozyklus ist; wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;


jedes R² unabhängig Wasserstoff, -OH, (C₁-C₆)Alkyl, substituiertes (C₁-C₆)Alkyl, (C₁-C₆)Alkoxy, substituiertes (C₁-C₆)Alkoxy, -C(O)-(C₁-C₆)Alkyl (Alkanoyl), substituiertes -C(O)-(C₁-C₆)Alkyl, -C(O)-(C₆-C₁₀)Aryl (Aroyl), substituiertes -C(O)-(C₆-C₁₀)Aryl, -C(O)OH (Carboxyl), -C(O)O(C₁-C₆)Alkyl (Alkoxy-carbonyl), substituiertes -C(O)O(C₁-C₆)Alkyl, -NR^aR^b, -C(O)NR^aR^b (Carbamoyl), substituiertes -C(O)NR^aR^b, Halo, Nitro oder Cyano ist; wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind; jedes R^a und R^b unabhängig Wasserstoff, (C₁-C₆)Alkyl, (C₃-C₈)Cycloalkyl, (C₁-C₆)Alkoxy, Halo(C₁-C₆)alkyl, (C₃-C₈)Cycloalkyl(C₁-C₆)alkyl, (C₁-C₆)Alkanoyl, Hydroxy(C₁-C₆)alkyl, Aryl, Aryl(C₁-C₆)alkyl, Het, Het(C₁-C₆)alkyl oder (C₁-C₆)Alkoxy-carbonyl ist;

X² eine Bindung oder eine verbindende Gruppe ist; n = 1, 2, 3 oder 4 ist und

R³ ein Makromolekül ist, umfassend ein heterologes Peptid, heterologes Protein, heterologes Lipid, Kügelchen, wie etwa Polystyrolkügelchen, heterologes Nukleinsäuremolekül oder Dendrimer;

oder ein pharmazeutisch akzeptables Salz davon, einschließlich Hydrate davon.

2. Eine Verbindung zur Verwendung in einem Verfahren zur Vorbeugung oder Inhibierung einer Infektion durch gram-positive Bakterien in einem Säugetier, umfassend die Verabreichung an das Säugetier einer effektiven Menge der Verbindung, welche die Formel (IB) aufweist:



wobei X¹ -O-, -S- oder -NR^c- ist;

wobei R^c Wasserstoff, C₁₋₁₀Alkyl oder substituiertes C₁₋₁₀Alkyl ist, oder R^c und R¹ zusammen mit dem Stickstoffatom einen heterozyklischen Ring oder einen substituierten heterozyklischen Ring bilden können, wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;

R¹ Wasserstoff, (C₁-C₁₀)Alkyl, substituiertes (C₁-C₁₀)Alkyl, C₆₋₁₀Aryl oder substituiertes C₆₋₁₀Aryl, ein C₅₋₉Heterozyklus, ein substituiertes C₅₋₉Heterozyklus ist; wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₁₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;

jedes R² unabhängig Wasserstoff, -OH, (C₁-C₆)Alkyl, substituiertes (C₁-C₆)Alkyl, (C₁-C₆)Alkoxy, substituiertes (C₁-C₆)Alkoxy, -C(O)-(C₁-C₆)Alkyl (Alkanoyl), substituiertes -C(O)-(C₁-C₆)Alkyl, -C(O)-(C₆-C₁₀)Aryl (Aroyl), substituiertes -C(O)-(C₆-C₁₀)Aryl, -C(O)OH (Carboxyl), -C(O)O(C₁-C₆)Alkyl (Alkoxy-carbonyl), substituiertes -C(O)O(C₁-C₆)Alkyl, -NR^aR^b, -C(O)NR^aR^b (Carbamoyl), substituiertes -C(O)NR^aR^b, Halo, Nitro oder Cyano ist; wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind; jedes R^a und R^b unabhängig Wasserstoff, (C₁-C₆)Alkyl, (C₃-C₈)Cycloalkyl, (C₁-C₆)Alkoxy, Halo(C₁-C₆)alkyl, (C₃-C₈)Cycloalkyl(C₁-C₆)alkyl, (C₁-C₆)Alkanoyl, Hydroxy(C₁-C₆)alkyl, Aryl, Aryl(C₁-C₆)alkyl, Het, Het(C₁-C₆)alkyl oder (C₁-C₆)Alkoxy-carbonyl ist;


X² eine Bindung oder eine verbindende Gruppe ist; n = 1, 2, 3 oder 4 ist und

R³ ein gram-positives Bakterium, ein isoliertes antigenes Protein oder Peptid des gram-positiven Bakteriums oder ein isoliertes Polysaccharid des gram-positiven Bakteriums ist;

oder ein pharmazeutisch akzeptables Salz davon, einschließlich Hydrate davon, wobei das gram-positive Bakterium nicht Bacillus oder Listeria ist.

EP 2 125 007 B9

3. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 1 oder 2, wobei X¹ ein Schwefelatom ist.
4. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 1 oder 2, wobei X¹ ein Sauerstoffatom ist.
5. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 1 oder 2, wobei X¹ -NR^c- ist, wobei R^c Wasserstoff, C₁₋₆Alkyl oder substituiertes C₁₋₆Alkyl ist; wobei die Alkylsubstituenten C₃₋₆Cycloalkyl, Hydroxy, C₁₋₆Alkoxy, Amino, Cyano oder Aryl sind.
6. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 5, wobei X¹ -NH- ist.
7. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 6, wobei R¹ und R^c zusammen einen heterozyklischen Ring oder einen substituierten heterozyklischen Ring bilden.
8. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 7, wobei R¹ und R^c zusammen einen substituierten oder unsubstituierten Morpholino-, Piperidino-, Pyrrolidino oder Piperazino-Ring bilden.
9. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 6, wobei R¹ Wasserstoff, C₁₋₄Alkyl oder substituiertes C₁₋₄Alkyl ist.
10. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 9, wobei R¹ Wasserstoff, CH₃-, CH₃-CH₂-, CH₃CH₂CH₂-, HydroxyC₁₋₄alkylen oder C₁₋₄AlkoxyC₁₋₄alkylen ist.
11. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 10, wobei R¹ Wasserstoff, CH₃-, CH₃-CH₂-, CH₃-O-CH₂CH₂- oder CH₃-CH₂-O-CH₂CH₂- ist.
12. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 11, wobei R² Wasserstoff, Halogen oder C₁₋₄Alkyl ist.
13. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 12, wobei R² Wasserstoff, Chloro, Bromo, CH₃- oder CH₃-CH₂- ist.
14. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 13, wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₁₋₆AlkoxyC₁₋₆alkylen, C₃₋₆Cycloalkyl, Amino, Cyano, Halogen oder Aryl sind;
15. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 14, wobei X² eine Bindung oder eine Kette ist, die bis zu etwa 24 Atome hat; wobei die Atome ausgewählt sind aus der Gruppe, bestehend aus Kohlenstoff, Stickstoff, Schwefel, nicht-peroxidem Sauerstoff und Phosphor.
16. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 15, wobei X² eine Bindung oder eine Kette ist, die von etwa 4 bis etwa 12 Atome hat.
17. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 15, wobei X² eine Bindung oder eine Kette ist, die von etwa 6 bis etwa 9 Atome hat.
18. Die Zusammensetzung oder Verbindung zur Verwendung nach Anspruch 15, wobei X²




ist.

19. Die Zusammensetzung zur Verwendung nach einem der Ansprüche 1 oder 3 bis 18, wobei das Makromolekül ein Phospholipid, ein Liposom oder ein Dendrimer ist.

EP 2 125 007 B9

20. Die Zusammensetzung zur Verwendung nach einem der Ansprüche 1 oder 3 bis 18, wobei das heterologe Peptid von 2 bis etwa 200 Aminosäurereste hat.
- 5 21. Die Verbindung zur Verwendung nach einem der Ansprüche 2 bis 18, wobei das Peptid von 2 bis etwa 200 Aminosäurereste hat.
22. Die Zusammensetzung zur Verwendung nach einem der Ansprüche 1 oder 3 bis 18, wobei das heterologe Peptid von 10 bis etwa 200 Aminosäurereste hat.
- 10 23. Die Verbindung zur Verwendung nach einem der Ansprüche 2 bis 18, wobei das Peptid von 10 bis etwa 200 Aminosäurereste hat.
24. Die Zusammensetzung zur Verwendung nach einem der Ansprüche 1 oder 3 bis 18, wobei das Makromolekül ein Kohlenhydrat umfasst.
- 15 25. Die Zusammensetzung zur Verwendung nach einem der Ansprüche 1 oder 3 bis 18, wobei das Nukleinsäuremolekül DNA, RNA oder PNA umfasst.
- 20 26. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 25, wobei das gram-positive Bakterium *Staphylococcus aureus* ist.
27. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 26, wobei das Säugetier ein Mensch ist.
- 25 28. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 27, wobei das Säugetier ein Risiko für eine Infektion mit gram-positiven Bakterien aufweist.
29. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 28, wobei das Säugetier immunsupprimiert ist.
- 30 30. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 29, wobei das Säugetier eine effektive Immunantwort auf die Verabreichung nach etwa 6 Tagen nach Verabreichung hat.
31. Die Zusammensetzung zur Verwendung nach einem der Ansprüche 1 oder 3 bis 30, wobei die Zusammensetzung eine Zubereitung abgetöteter gram-positiver Bakterien umfasst.
- 35 32. Die Zusammensetzung oder Verbindung zur Verwendung nach einem der Ansprüche 1 bis 31, wobei die Menge effektiv ist, einer Infektion nach etwa 6 Tagen nach Verabreichung vorzubeugen.
- 40 33. Eine Vakzine, umfassend eine Zusammensetzung, umfassend ein bakterielles Antigen eines *S. aureus* Bakteriums und eine Menge einer Verbindung, welche die Formel (IA) aufweist:



wobei X¹ -O-, -S- oder -NR^c- ist;


wobei R^c Wasserstoff, C₁₋₁₀Alkyl oder substituiertes C₁₋₁₀Alkyl ist, oder R^c und R¹ zusammen mit dem Stickstoffatom einen heterozyklischen Ring oder einen substituierten heterozyklischen Ring bilden können, wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;

R¹ Wasserstoff, (C₁-C₁₀)Alkyl, substituiertes (C₁-C₁₀)Alkyl, C₆₋₁₀Aryl oder substituiertes C₆₋₁₀Aryl, ein C₅₋₉Heterozyklus, ein substituiertes C₅₋₉Heterozyklus ist; wobei die Substituenten auf den Alkyl-, Aryl- oder

heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;
 jedes R² unabhängig Wasserstoff, -OH, (C₁₋₆)Alkyl, substituiertes (C₁₋₆)Alkyl, (C₁₋₆)Alkoxy, substituiertes (C₁₋₆)Alkoxy, -C(O)-(C₁₋₆)Alkyl (Alkanoyl), substituiertes -C(O)-(C₁₋₆)Alkyl, -C(O)-(C₆₋₁₀)Aryl (Aroyl),
 5 substituiertes -C(O)-(C₆₋₁₀)Aryl, -C(O)OH (Carboxyl), -C(O)O(C₁₋₆)Alkyl (Alkoxy-carbonyl), substituiertes -C(O)O(C₁₋₆)Alkyl, -NR^aR^b, -C(O)NR^aR^b (Carbamoyl), substituiertes -C(O)NR^aR^b, Halo, Nitro oder Cyano ist;
 wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;
 jedes R^a und R^b unabhängig Wasserstoff, (C₁₋₆)Alkyl, (C₃₋₈)Cycloalkyl, (C₁₋₆)Alkoxy, Halo(C₁₋₆)alkyl, (C₃₋₈)Cycloalkyl(C₁₋₆)alkyl, (C₁₋₆)Alkanoyl, Hydroxy(C₁₋₆)alkyl, Aryl, Aryl(C₁₋₆)alkyl, Het, Het(C₁₋₆)alkyl oder (C₁₋₆)Alkoxy-carbonyl ist;
 X² eine Bindung oder eine verbindende Gruppe ist; n = 1, 2, 3 oder 4 ist und
 R³ ein Makromolekül ist, umfassend ein heterologes Peptid, heterologes Protein, heterologes Lipid, Kügelchen, wie z.B. Polystyrolkügelchen, heterologes Nukleinsäuremolekül oder Dendrimer;

oder ein pharmazeutisch akzeptables Salz davon, einschließlich Hydrate davon.

34. Eine Vakzine, umfassend eine Verbindung der Formel (IB):



wobei X¹ -O-, -S- oder -NR^c- ist;

wobei R^c Wasserstoff, C₁₋₁₀Alkyl oder substituiertes C₁₋₁₀Alkyl ist, oder R^c und R¹ zusammen mit dem Stickstoffatom einen heterozyklischen Ring oder einen substituierten heterozyklischen Ring bilden können, wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;

R¹ Wasserstoff, (C₁₋₁₀)Alkyl, substituiertes (C₁₋₁₀)Alkyl, C₆₋₁₀Aryl oder substituiertes C₆₋₁₀Aryl, ein C₅₋₉Heterozyklus, ein substituiertes C₅₋₉Heterozyklus ist; wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;

jedes R² unabhängig Wasserstoff, -OH, (C₁₋₆)Alkyl, substituiertes (C₁₋₆)Alkyl, (C₁₋₆)Alkoxy, substituiertes (C₁₋₆)Alkoxy, -C(O)-(C₁₋₆)Alkyl (Alkanoyl), substituiertes -C(O)-(C₁₋₆)Alkyl, -C(O)-(C₆₋₁₀)Aryl (Aroyl), substituiertes -C(O)-(C₆₋₁₀)Aryl, -C(O)OH (Carboxyl), -C(O)O(C₁₋₆)Alkyl (Alkoxy-carbonyl), substituiertes -C(O)O(C₁₋₆)Alkyl, -NR^aR^b, -C(O)NR^aR^b (Carbamoyl), substituiertes -C(O)NR^aR^b, Halo, Nitro oder Cyano ist;

wobei die Substituenten auf den Alkyl-, Aryl- oder heterozyklischen Gruppen Hydroxy, C₁₋₆Alkyl, HydroxyC₁₋₆alkylen, C₁₋₆Alkoxy, C₃₋₆Cycloalkyl, C₁₋₆AlkoxyC₁₋₆alkylen, Amino, Cyano, Halogen oder Aryl sind;
 jedes R^a und R^b unabhängig Wasserstoff, (C₁₋₆)Alkyl, (C₃₋₈)Cycloalkyl, (C₁₋₆)Alkoxy, Halo(C₁₋₆)alkyl, (C₃₋₈)Cycloalkyl(C₁₋₆)alkyl, (C₁₋₆)Alkanoyl, Hydroxy(C₁₋₆)alkyl, Aryl, Aryl(C₁₋₆)alkyl, Het, Het(C₁₋₆)alkyl oder (C₁₋₆)Alkoxy-carbonyl ist;

X² eine Bindung oder eine verbindende Gruppe ist; n = 1, 2, 3 oder 4 ist und


R³ das *S. aureus* Bakterium, ein isoliertes antigenes Protein oder Peptid des *S. aureus* Bakteriums, oder ein isoliertes Polysaccharid des *S. aureus* Bakteriums ist;

oder ein pharmazeutisch akzeptables Salz davon, einschließlich Hydrate davon.

Revendications

1. Composition pour son utilisation dans une méthode de prévention ou d'inhibition d'une infection bactérienne à Gram

positif chez un mammifère, comprenant l'administration au mammifère d'une quantité efficace de la composition comprenant un antigène bactérien d'une bactérie à Gram positif et d'une quantité de composé répondant à la formule (IA) :



dans laquelle X¹ est -O-, -S-, ou -NR^c- ;

dans laquelle R^c est hydrogène, alkyle en C₁ à C₁₀ ou alkyle en C₁ à C₁₀ substitué, ou R^c et R¹, pris conjointement avec l'atome d'azote, peuvent former un cycle hétérocyclique ou un cycle hétérocyclique substitué, dans lequel les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à C₆, hydroxyalkylène en C₁ à C₆, alcoxy en C₁ à C₆, cycloalkyle en C₃ à C₆, alcoxy en C₁ à C₆alkylène en C₁ à C₆, amino, cyano, halogène ou aryle ;

R¹ est hydrogène, alkyle (en C₁ à C₁₀), alkyle (en C₁ à C₁₀) substitué, aryle en C₆ à C₁₀, ou aryle en C₆ à C₁₀ substitué, hétérocyclique en C₅ à C₉, hétérocyclique en C₅ à C₉ substitué ; dans laquelle les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à C₆, hydroxyalkylène en C₁ à C₆, alcoxy en C₁ à C₆, cycloalkyle en C₃ à C₆, alcoxy en C₁ à C₆alkylène en C₁ à C₆, amino, cyano, halogène, ou aryle ;

chaque R² est indépendamment hydrogène, -OH, alkyle (en C₁ à C₆), alkyle (en C₁ à C₆) substitué, alcoxy (en C₁ à C₆), alcoxy (en C₁ à C₆) substitué, -C(O)-alkyle (en C₁ à C₆) (alcanoyle), -C(O)-alkyle (en C₁ à C₆) substitué, -C(O)-aryle (en C₆ à C₁₀) (aroyle), -C(O)-aryle (en C₆ à C₁₀) substitué, -C(O)OH (carboxyle), -C(O)Oalkyle (en C₁ à C₆) (alcoycarbonyle), -C(O)Oalkyle (en C₁ à C₆) substitué, -NR^aR^b, -C(O)NR^aR^b (carbamoyle), -C(O)NR^aR^b substitué, halogéno, nitro ou cyano ; dans laquelle les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à C₆, hydroxyalkylène en C₁ à C₆, alcoxy en C₁ à C₆, cycloalkyle en C₃ à C₆, alcoxy en C₁ à C₆alkylène en C₁ à C₆, amino, cyano, halogène ou aryle ;


chaque R^a et R^b est indépendamment hydrogène, alkyle (en C₁ à C₆), cycloalkyle (en C₃ à C₈), alcoxy (en C₁ à C₆), halogénoalkyle (en C₁ à C₆), cycloalkyl (en C₃ à C₈)alkyle (en C₁ à C₆), alcanoyle (en C₁ à C₆), hydroxyalkyle (en C₁ à C₆), aryle, arylalkyle (en C₁ à C₆), Het, Hetalkyle (en C₁ à C₆) ou alcoycarbonyle (en C₁ à C₆) ;

X² est une liaison ou un groupe de liaison ; n = 1, 2, 3 ou 4 ; et

R³ est une macromolécule comprenant un peptide hétérologue, une protéine hétérologue, un lipide hétérologue, une bille, telle qu'une bille de polystyrène, une molécule d'acide nucléique hétérologue ou un dendrimère ;

ou un sel pharmaceutiquement acceptable de celui-ci, y compris ses hydrates.

2. Composé pour son utilisation dans une méthode de prévention ou d'inhibition d'une infection bactérienne à Gram positif chez un mammifère, comprenant l'administration au mammifère d'une quantité efficace du composé répondant à la formule (IB) :



dans laquelle X¹ est -O-, -S-, ou -NR^c- ;

dans laquelle R^c est hydrogène, alkyle en C₁ à 10 ou alkyle en C₁ à 10 substitué, ou R^c et R¹, pris conjointement avec l'atome d'azote, peuvent former un cycle hétérocyclique ou un cycle hétérocyclique substitué, dans lequel les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à 6, hydroxyalkylène en C₁ à 6, alcoxy en C₁ à 6, cycloalkyle en C₃ à 6, alcoxy en C₁ à 6 alkylène en C₁ à 6, amino, cyano, halogène ou aryle ;

R¹ est hydrogène, alkyle (en C₁ à C₁₀), alkyle (en C₁ à C₁₀) substitué, aryle en C₆ à 10, ou aryle en C₆ à 10 substitué, hétérocyclique en C₅ à 9, hétérocyclique en C₅ à 9 substitué ; dans laquelle les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à 6, hydroxyalkylène en C₁ à 6, alcoxy en C₁ à 6, cycloalkyle en C₃ à 6, alcoxy en C₁ à 6 alkylène en C₁ à 6, amino, cyano, halogène, ou aryle ;

chaque R² est indépendamment hydrogène, -OH, alkyle (en C₁ à C₆), alkyle (en C₁ à C₆) substitué, alcoxy (en C₁ à C₆), alcoxy (en C₁ à C₆) substitué, -C(O)-alkyle (en C₁ à C₆) (alcanoyle), -C(O)-alkyle (en C₁ à C₆) substitué, -C(O)-aryle (en C₆ à C₁₀) (aroyle), -C(O)-aryle (en C₆ à C₁₀) substitué, -C(O)OH (carboxyle), -C(O)Oalkyle (en C₁ à C₆) (alcoxycarbonyle), -C(O)Oalkyle (en C₁ à C₆) substitué, -NR^aR^b, -C(O)NR^aR^b (carbamoyle), -C(O)NR^aR^b substitué, halogéno, nitro ou cyano ; dans laquelle les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à 6, hydroxyalkylène en C₁ à 6, alcoxy en C₁ à 6, cycloalkyle en C₃ à 6, alcoxy en C₁ à 6 alkylène en C₁ à 6, amino, cyano, halogène ou aryle ;

chaque R^a et R^b est indépendamment hydrogène, alkyle (en C₁ à C₆), cycloalkyle (en C₃ à C₆), alcoxy (en C₁ à C₆), halogénoalkyle (en C₁ à C₆), cycloalkyl (en C₃ à C₆) alkyle (en C₁ à C₆), alcanoyle (en C₁ à C₆), hydroxyalkyle (en C₁ à C₆), aryle, arylalkyle (en C₁ à C₆), Het, Hetalkyle (en C₁ à C₆) ou alcoxycarbonyle (en C₁ à C₆) ;

X² est une liaison ou un groupe de liaison ; n = 1, 2, 3 ou 4 ; et

R³ est la bactérie Gram positive, une protéine antigénique isolée ou un peptide de la bactérie à Gram positif, ou un polysaccharide isolé de la bactérie à Gram positif ;

ou un sel pharmaceutiquement acceptable de celui-ci, y compris ses hydrates, dans lequel la bactérie à Gram positif n'est pas *Bacillus* ou *Listeria*.

3. Composition ou composé pour son utilisation selon la revendication 1 ou 2, dans laquelle ou lequel X¹ est un atome de soufre.
4. Composition ou composé pour son utilisation selon la revendication 1 ou 2, dans laquelle ou lequel X¹ est un atome d'oxygène.
5. Composition ou composé pour son utilisation selon la revendication 1 ou 2, dans laquelle ou lequel X¹ est -NR^c-, dans laquelle ou lequel R^c est hydrogène, alkyle en C₁ à 6 ou alkyle en C₁ à 6 substitué ; dans laquelle ou lequel les substituants alkyle sont cycloalkyle en C₃ à 6, hydroxy, alcoxy en C₁ à 6, amino, cyano ou aryle.
6. Composition ou composé pour son utilisation selon la revendication 5, dans laquelle ou lequel X¹ est -NH-.
7. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 6, dans laquelle ou lequel R¹ et R^c pris ensemble forment un cycle hétérocyclique ou un cycle hétérocyclique substitué.
8. Composition ou composé pour son utilisation selon la revendication 7, dans laquelle ou lequel R¹ et R^c pris ensemble forment un cycle morpholino, pipéridino, pyrrolidino ou pipérazino substitué ou non substitué.
9. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 6, dans laquelle ou lequel R¹ est hydrogène, alkyle en C₁ à 4, ou alkyle en C₁ à 4 substitué.
10. Composition ou composé pour son utilisation selon la revendication 9, dans laquelle ou lequel R¹ est hydrogène, CH₃-, CH₃-CH₂-, -CH₃CH₂CH₂-, hydroxyalkylène en C₁ à 4 ou alcoxy en C₁ à 4 alkylène en C₁ à 4.
11. Composition ou composé pour son utilisation selon la revendication 10, dans laquelle ou lequel R¹ est hydrogène, CH₃-, CH₃-CH₂-, CH₃-O-CH₂CH₂- ou CH₃-CH₂-O-CH₂CH₂-.
12. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 11, dans laquelle ou lequel R² est hydrogène, halogène ou alkyle en C₁ à 4.

EP 2 125 007 B9

13. Composition ou composé pour son utilisation selon la revendication 12, dans laquelle ou lequel R² est hydrogène, chloro, bromo, CH₃- ou CH₃-CH₂-.


14. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 ou 13, dans laquelle ou lequel les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à 6, hydroxyalkylène en C₁ à 6, alcoxy en C₁ à 6, alcoxy en C₁ à 6, alkylène en C₁ à 6, cycloalkyle en C₃ à 6, amino, cyano, halogène ou aryle.

15. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 14, dans laquelle ou lequel X² est une liaison ou une chaîne ayant jusqu'à environ 24 atomes ; dans laquelle ou lequel les atomes sont choisis dans le groupe consistant en le carbone, l'azote, le soufre, l'oxygène non-péroxyde et le phosphore.

16. Composition ou composé pour son utilisation selon la revendication 15, dans laquelle ou lequel X² est une liaison ou une chaîne ayant d'environ 4 à environ 12 atomes.

17. Composition ou composé pour son utilisation selon la revendication 15, dans laquelle ou lequel X² est une liaison ou une chaîne ayant d'environ 6 à environ 9 atomes.

18. Composition ou composé pour son utilisation selon la revendication 15, dans laquelle ou lequel X² est



19. Composition pour son utilisation selon l'une quelconque des revendications 1 ou 3 à 18, dans laquelle la macromolécule est un phospholipide, un liposome ou un dendrimère.

20. Composition pour son utilisation selon l'une quelconque des revendications 1 ou 3 à 18, dans laquelle le peptide hétérologue a de 2 à environ 200 résidus d'acides aminés.

21. Composé pour son utilisation selon l'une quelconque des revendications 2 à 18, dans lequel le peptide a de 2 à environ 200 résidus d'acides aminés.

22. Composition pour son utilisation selon l'une quelconque des revendications 1 ou 3 à 18, dans laquelle le peptide hétérologue a de 10 à environ 200 résidus d'acides aminés.

23. Composé pour son utilisation selon l'une quelconque des revendications 2 à 18, dans lequel le peptide a de 10 à environ 200 résidus d'acides aminés.

24. Composition pour son utilisation selon l'une quelconque des revendications 1 ou 3 à 18, dans laquelle la macromolécule comprend un glucide.

25. Composition pour son utilisation selon l'une quelconque des revendications 1 ou 3 à 18, dans laquelle la molécule d'acide nucléique comprend l'ADN, l'ARN ou l'APN.


26. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 25, dans laquelle ou lequel la bactérie à Gram positif est *Staphylococcus aureus*.

27. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 26, dans laquelle ou lequel le mammifère est un humain.

28. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 27, dans laquelle ou lequel le mammifère présente un risque d'infection bactérienne à Gram positif.

29. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 28, dans laquelle ou lequel le mammifère est immunocompromis.

30. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 29, dans laquelle ou lequel le mammifère a une réponse immunitaire efficace à l'administration environ 6 jours après administration.
31. Composition pour son utilisation selon l'une quelconque des revendications 1 ou 3 à 30, dans laquelle la composition comprend une préparation de bactéries à Gram positif tuées.
32. Composition ou composé pour son utilisation selon l'une quelconque des revendications 1 à 31, dans laquelle ou lequel la quantité est efficace pour prévenir une infection environ 6 jours après administration.
33. Vaccin comprenant une composition comprenant un antigène bactérien d'une bactérie *S. aureus* et une quantité d'un composé répondant à la formule (IA) :



dans laquelle X¹ est -O-, -S-, ou -NR^c- ;

dans laquelle R^c est hydrogène, alkyle en C₁ à 10 ou alkyle en C₁ à 10 substitué, ou R^c et R¹, pris conjointement avec l'atome d'azote, peuvent former un cycle hétérocyclique ou un cycle hétérocyclique substitué, dans lequel les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à 6, hydroxyalkylène en C₁ à 6, alcoxy en C₁ à 6, cycloalkyle en C₃ à 6, alcoxy en C₁ à 6 alkylène en C₁ à 6, amino, cyano, halogène ou aryle ;

R¹ est hydrogène, alkyle (en C₁ à C₁₀), alkyle (en C₁ à C₁₀) substitué, aryle en C₆ à 10, ou aryle en C₆ à 10 substitué, hétérocyclique en C₅ à 9, hétérocyclique en C₅ à 9 substitué ; dans laquelle les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à 6, hydroxyalkylène en C₁ à 6, alcoxy en C₁ à 6, cycloalkyle en C₃ à 6, alcoxy en C₁ à 6 alkylène en C₁ à 6, amino, cyano, halogène, ou aryle ;

chaque R² est indépendamment hydrogène, -OH, alkyle (en C₁ à C₆), alkyle (en C₁ à C₆) substitué, alcoxy (en C₁ à C₆), alcoxy (en C₁ à C₆) substitué, -C(O)-alkyle (en C₁ à C₆) (alcanoyle), -C(O)-alkyle (en C₁ à C₆) substitué, -C(O)-aryle (en C₆ à C₁₀) (aroyle), -C(O)-aryle (en C₆ à C₁₀) substitué, -C(O)OH (carboxyle), -C(O)Oalkyle (en C₁ à C₆) (alcoxycarbonyle), -C(O)Oalkyle (en C₁ à C₆) substitué, -NR^aR^b, -C(O)NR^aR^b (carbamoyle), -C(O)NR^aR^b substitué, halogéno, nitro ou cyano ; dans laquelle les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à 6, hydroxyalkylène en C₁ à 6, alcoxy en C₁ à 6, cycloalkyle en C₃ à 6, alcoxy en C₁ à 6 alkylène en C₁ à 6, amino, cyano, halogène ou aryle ;


chaque R^a et R^b est indépendamment hydrogène, alkyle (en C₁ à C₆), cycloalkyle (en C₃ à C₈), alcoxy (en C₁ à C₆), halogénoalkyle (en C₁ à C₆), cycloalkyl (en C₃ à C₈) alkyle (en C₁ à C₆), alcanoyle (en C₁ à C₆), hydroxyalkyle (en C₁ à C₆), aryle, arylalkyle (en C₁ à C₆), Het, Hetalkyle (en C₁ à C₆) ou alcoxycarbonyle (en C₁ à C₆) ;

X² est une liaison ou un groupe de liaison ; n = 1, 2, 3 ou 4 ; et

R³ est une macromolécule comprenant un peptide hétérologue, une protéine hétérologue, un lipide hétérologue, une bille, telle qu'une bille de polystyrène une molécule d'acide nucléique hétérologue ou un dendrimère ;

ou un sel pharmaceutiquement acceptable de celui-ci, y compris ses hydrates.

34. Vaccin comprenant un composé répondant à la formule (IB) :



dans laquelle X¹ est -O-, -S-, ou -NR^c ;

15 dans laquelle R^c est hydrogène, alkyle en C₁ à C₁₀ ou alkyle en C₁ à C₁₀ substitué, ou R^c et R¹, pris conjointement avec l'atome d'azote, peuvent former un cycle hétérocyclique ou un cycle hétérocyclique substitué, dans lequel les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à C₆, hydroxyalkylène en C₁ à C₆, alcoxy en C₁ à C₆, cycloalkyle en C₃ à C₆, alcoxy en C₁ à C₆alkylène en C₁ à C₆, amino, cyano, halogène ou aryle ;

20 R¹ est hydrogène, alkyle (en C₁ à C₁₀), alkyle (en C₁ à C₁₀) substitué, aryle en C₆ à C₁₀, ou aryle en C₆ à C₁₀ substitué, hétérocyclique en C₅ à C₉, hétérocyclique en C₅ à C₉ substitué ; dans laquelle les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à C₆, hydroxyalkylène en C₁ à C₆, alcoxy en C₁ à C₆, cycloalkyle en C₃ à C₆, alcoxy en C₁ à C₆alkylène en C₁ à C₆, amino, cyano, halogène, ou aryle ;

25 chaque R² est indépendamment hydrogène, -OH, alkyle (en C₁ à C₆), alkyle (en C₁ à C₆) substitué, alcoxy (en C₁ à C₆), alcoxy (en C₁ à C₆) substitué, -C(O)-alkyle (en C₁ à C₆) (alcanoyle), -C(O)-alkyle (en C₁ à C₆) substitué, -C(O)-aryle (en C₆ à C₁₀) (aroyle), -C(O)-aryle (en C₆ à C₁₀) substitué, -C(O)OH (carboxyle), -C(O)Oalkyle (en C₁ à C₆) (alcoxycarboyle), -C(O)Oalkyle (en C₁ à C₆) substitué, -NR^aR^b, -C(O)NR^aR^b (carbamoyle), -C(O)NR^aR^b substitué, halogéno, nitro ou cyano ; dans laquelle les substituants sur les groupes alkyle, aryle ou hétérocyclique sont hydroxy, alkyle en C₁ à C₆, hydroxyalkylène en C₁ à C₆, alcoxy en C₁ à C₆, cycloalkyle en C₃ à C₆, alcoxy en C₁ à C₆alkylène en C₁ à C₆, amino, cyano, halogène ou aryle ;

30 chaque R^a et R^b est indépendamment hydrogène, alkyle (en C₁ à C₆), cycloalkyle (en C₃ à C₈), alcoxy (en C₁ à C₆), halogénoalkyle (en C₁ à C₆), cycloalkyl (en C₃ à C₈)alkyle (en C₁ à C₆), alcanoyle (en C₁ à C₆), hydroxyalkyle (en C₁ à C₆), aryle, arylalkyle (en C₁ à C₆), Het, Hetalkyle (en C₁ à C₆) ou alcoxycarboyle (en C₁ à C₆) ;

X² est une liaison ou un groupe de liaison ; n = 1, 2, 3 ou 4 ; et

35 R³ est la bactérie *S. aureus*, une protéine antigénique isolée ou un peptide de la bactérie *S. aureus*, ou un polysaccharide isolé de la bactérie *S. aureus* ;

ou un sel pharmaceutiquement acceptable de celui-ci, y compris ses hydrates.




FIG. 1





FIG. 2



UC-1V150

FIG. 3B




FIG. 3C




FIG. 4




FIG. 5




FIG. 6

UC-1V199/L can Act as an Inhibitor of TLR7 Signaling




FIG. 7

UC-1V199/L Inhibits TLR7 Signaling




FIG. 8A

UC-1V199/L Inhibits TLR2 Signaling




FIG. 8B




FIG. 9




FIG. 10B

FIG. 10A




FIG. 11A

FIG. 11B

FIG. 11C




FIG. 12

Cytokine induction in BMDM by irradiated anthrax spore conjugate




FIG. 13





FIG. 14

UC-1V150/MSA as anthrax spore vaccine adjuvant: (I) One immunization 6 days prior to infection



FIG. 15

Protection elicited by gamma-irradiated anthrax spores depends on CD4+ cells




FIG. 16

| Cytokine | Time (h) | Conjugated | | | | Unconjugated | | | | Ratio | |
|----------|----------|------------|------|-------|------|--------------|------|-------|------|-------|-----|
| | | BALF | | Serum | | BALF | | Serum | | | |
| | | ng/ml | SEM | ng/ml | SEM | ng/ml | SEM | ng/ml | SEM | | |
| IL-6 | 2 | 0.25 | 0.09 | 0.28 | 0.00 | 0.9 | 0.73 | 0.40 | 5.00 | 1.14 | 0.1 |
| | 4 | 1.34 | 0.38 | 0.06 | 0.02 | 22.1 | 1.09 | 0.64 | 2.93 | 1.53 | 0.4 |
| | 6 | 3.50 | 1.07 | 0.14 | 0.05 | 24.3 | 3.00 | 0.76 | 3.95 | 1.24 | 0.8 |
| | 24 | 4.06 | 0.41 | 0.03 | 0.00 | 117.2 | 4.39 | 0.84 | 1.08 | 0.31 | 4.1 |
| IL-12p40 | 2 | 0.06 | 0.00 | 0.23 | 0.03 | 0.2 | 0.11 | 0.09 | 0.53 | 0.53 | 0.2 |
| | 4 | 0.11 | 0.00 | 0.50 | 0.39 | 0.2 | 0.06 | 0.03 | 0.23 | 0.23 | 0.3 |
| | 6 | 0.29 | 0.04 | 0.25 | 0.06 | 1.1 | 0.00 | 0.00 | 2.82 | 0.46 | 0.0 |
| | 24 | 5.05 | 3.70 | 0.11 | 0.00 | 45.1 | 0.01 | 0.01 | 0.49 | 0.47 | 0.0 |
| TNF-a | 2 | 1.25 | 1.02 | 0.08 | 0.01 | 16.4 | 2.62 | 0.83 | 0.52 | 0.21 | 5.0 |
| | 4 | 9.67 | 1.01 | 0.02 | 0.01 | 580.1 | 1.53 | 0.73 | 0.32 | 0.16 | 4.8 |
| | 6 | 8.28 | 1.72 | 0.02 | 0.00 | 427.5 | 2.03 | 0.67 | 0.39 | 0.15 | 5.2 |
| | 24 | 4.22 | 2.80 | 0.01 | 0.00 | 551.8 | 0.93 | 0.47 | 0.15 | 0.04 | 6.2 |

FIG. 17




FIG. 18




FIG. 19




FIG. 20A




FIG. 20B




FIG. 21




FIG. 22




FIG. 24




FIG. 25





FIG. 26

FIG. 27












FIG. 28



Imiquimod




Bropirimine




UC-1V199


FIG. 28 (Cont'd.)



UC-1W236




UC-1X51




Loxoribine


Fig. 28 (Cont'd.)



UC-1W247



UC-1X113



UC-1V186

FIG. 28 (Cont'd)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 88869907 P [0001]
- US 6329381 B, Kurimoto [0058]
- WO 2007024707 A [0129]
- US 20050004144 A [0147]
- US 4608392 A, Jacquet [0176]
- US 4992478 A, Geria [0176]
- US 4559157 A, Smith [0176]
- US 4820508 A, Wortzman [0176]
- US 4938949 A [0177]

Non-patent literature cited in the description

- **JANEWAY et al.** *Annu. Rev. Immunol.*, 2002, vol. 20, 197 [0003]
- **UNDERHILL et al.** *Curr. Opin. Immunol.*, 2002, vol. 14, 103 [0003]
- **HOMUNG et al.** *J. Immunol.*, 2002, vol. 168, 4531 [0003]
- **JIN et al.** *Bioorg Med Chem Lett.*, 2006, vol. 16, 4559-63 [0004]
- **WILLE-REECE et al.** *Proc Natl Acad Sci USA*, 2005, vol. 102 (42), 15190-4 [0005]
- **BRASSARD et al.** *J. Leukoc. Biol.*, 2002, vol. 71, 568 [0006]
- **LEVY et al.** *Cytokine Growth Factor Rev.*, 2001, vol. 12, 143 [0006]
- **CAPPARELLI et al.** *Antimicrob. Agents Chemo.*, 2005, vol. 49, 4121 [0008]
- Protecting Groups. **T.W. GREENE.** *Organic Synthesis.* Wiley, 1981 [0033]
- Remington's Pharmaceutical Sciences. Mack Publishing Company, 1985, 1418 [0038]
- **LEE et al.** *Proc. Natl. Acad. Sci. USA*, 2003, vol. 100, 6646 [0177]
- *Bull. Korean Chem. Soc.*, 2002, vol. 23, 1697 [0192]
- **JAYACHITRA et al.** *Synth. Comm.*, 2003, vol. 33, 3461 [0195]
- **KENNEY et al.** *J. Infect. Dis.*, 2004, vol. 190, 774 [0210]
- **SABET et al.** *FEMS Immunol. Med. Microbiol.*, 2006, vol. 47, 369 [0213]
- **GUIDI-RONTANI et al.** *Mol. Microbiol.*, 2001, vol. 42, 931 [0213]
- **LEE et al.** *Proc. Natl. Acad. Sci. USA*, 2006, vol. 103, 1828 [0216]
- **ROTHENFUSSER et al.** *Hum. Immunol.*, 2002, vol. 63, 111 [0221]
- **BARCHET et al.** *Eur. J. Immunol.*, 2005, vol. 35, 360 [0224]
- **HUGHES et al.** *Infect. Immun.*, 2005, vol. 73, 7535 [0224]