



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
06.01.2010 Bulletin 2010/01

(21) Application number: **09164393.2**

(22) Date of filing: **15.11.2004**

(51) Int Cl.:
A61K 31/42 ^(2006.01) **A61K 31/422** ^(2006.01)
A61K 31/426 ^(2006.01) **A61K 31/427** ^(2006.01)
C07D 277/56 ^(2006.01) **C07D 417/12** ^(2006.01)
C07D 417/04 ^(2006.01) **C07D 417/06** ^(2006.01)
C07D 263/34 ^(2006.01)

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LU MC NL PL PT RO SE SI SK TR

(30) Priority: **14.11.2003 US 520355 P**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
04811321.1 / 1 682 127

(71) Applicant: **Vertex Pharmaceuticals Incorporated**
Cambridge, MA 02139 (US)

(72) Inventors:
• **Hadida Ruah, Sarah. S.**
La Jolla, CA 92109 (US)

• **Miller, Mark T.**
San Diego, CA 92109 (US)
• **Grootenhuys, Peter D.J.**
San Diego, CA 92130 (US)
• **Hamilton, Matthew**
Hackettstown, NJ 07840 (US)


(74) Representative: **Cohausz & Florack**
Patent- und Rechtsanwälte
Bleichstraße 14
40211 Düsseldorf (DE)

Remarks:

This application was filed on 02-07-2009 as a divisional application to the application mentioned under INID code 62.

(54) **Thiazoles and oxazoles useful as modulators of atp-binding cassette transporters**

(57) The present invention relates to modulators of ATP-Binding Cassette ("ABC") transporters or fragments thereof, including Cystic Fibrosis Transmembrane Regulator ("CFTR"), compositions thereof, and methods therewith. The present invention also relates to methods of treating ABC transporter mediated diseases using such modulators.



DescriptionTECHNICAL FIELD OF THE INTENTION

[0001] The present invention relates to modulators of ATP-Binding Cassette ("ABC") transporters or fragments thereof, including cystic fibrosis transmembrane conductance regulator ("CFTR"), compositions thereof, and methods therewith. The present invention also relates to methods of treating ABC transporter mediated diseases using such modulators.

BACKGROUND OF THE INVENTION

[0002] ABC transporters are a group of membrane transporter proteins that play a major role in the transport and protection of cells against a wide variety of pharmacological agents, potentially toxic drugs, and xenobiotics. ABC transporters are homologous membrane proteins that bind and use cellular adenosine triphosphate (ATP) for their specific activities. Some of these transporters were discovered as multidrug resistance proteins (like the MDR1-P glycoprotein, or the multidrug resistance protein, N4RP1), defending malignant cancer cells against chemotherapeutic agents. Up until the present time, 48 human ABC transporters have been identified, and these have been arranged into 7 families based on their sequence identity and function.

[0003] ABC transporters play a variety of important physiological roles within the body, as well as providing a defense against harmful compounds from the environment. Moreover they represent important potential drug targets both in their own right, as well as, because in many cases therapeutic drugs are also transported out of the target cell by these molecules.

[0004] One of the members of the ABC transporter family, namely, CFTR, is believed be the chloride channel responsible for cAMP-mediated chloride secretion in epithelial cells, and to play a key role in the secretion of chloride and maintenance of normal electrolyte transport throughout the body. CFTR is a protein of approximately 1480 amino acids made up of two repeated elements, each comprising six transmembrane segments and a nucleotide-binding domain. The two repeats are separated by a large, polar, regulatory (R)-domain containing multiple potential phosphorylation sites.

[0005] The gene associated with CFTR has been identified and sequenced (See Gregory, R. J. et al. (1990) Nature 347:382-386; Rich, D. P. et al. (1990) Nature 347:358-362), (Riordan, J. R. et al. (1989) Science 245:1066-1073). A defect in this gene leads to cystic fibrosis (hereinafter "CF"), the most common fatal genetic disease in humans affecting approximately one in every 2,500 infants born in the United States. Within the general United States population, up to 10 million people carry a single copy of the defective gene without apparent ill effects. In contrast, individuals with two copies of the CF associated gene suffer from the chronic effects of CF, including chronic lung destruction and death.

[0006] In patients with CF, expression of the CF associated gene in airway cells, leads to reduced cellular apical chloride conductance causing an imbalance in ion and fluid transport. It is widely believed that this leads to the abnormal mucus secretion in pancreatic ductules and in the airways that ultimately results in the pulmonary infections and epithelial cell damage typically associated with disease progression in CF. In addition to respiratory problems, CF patients typically suffer from gastrointestinal problems, and pancreatic insufficiency. Males are almost uniformly infertile and fertility is decreased in females. In contrast to the severe effects of two copies of the CF associated gene, individuals with a single copy of the CF associated gene exhibit increased resistance to cholera and to dehydration resulting from diarrhea - perhaps explaining the relatively high frequency of the CF gene within the population.

[0007] Sequence analysis of the CFTR gene of CF chromosomes has revealed a variety of disease causing mutations (Cutting, G. R. et al. (1990) Nature 346:366-369; Dean, M. et al. (1990) Cell 61:863:870; and Kerem, B-S. et al. (1989) Science 245:1073-1080; Kerem, B-S et al. (1990) Proc. Natl. Acad. Sci. USA 87:8447-8451). At present, more than 1000 mutations in the CF gene have been identified (<http://www.genet.sickkids.on.ca/cftr/>), but population studies have indicated that the most common CF mutation, a deletion of the 3 nucleotides that encode phenylalanine at position 508 of the CFTR amino acid sequence, is associated with approximately 70% of the cases of cystic fibrosis. The mutated CFTR protein is referred to as $\Delta F508$.

[0008] It is believed that the deletion of residue 508 in $\Delta F508$ -CFTR prevents the nascent protein from folding correctly, resulting in the inability of this mutant protein to exit the endoplasmic reticulum (hereinafter "ER"), and traffic to the plasma membrane. As a result, insufficient amounts of the mature protein are present at the plasma membrane and chloride transport within epithelial tissues is significantly reduced (Quinton, P. M. (1990), FASEB J. 4: 2709-2727). Hence, the cellular phenomenon of defective ER processing of other proteins like CFTR, by the ER machinery, has been shown to be the underlying basis for a wide range of isolated and inherited diseases. The two ways that the ER machinery can malfunction is either by loss of coupling to ER export of the proteins leading to degradation, or by the ER accumulation of these defective/misfolded proteins [Aridor M, et al., Nature Med., 5 (7), pp 745-751 (1999); Shastri, B.S., et al., Neurochem. International, 43, pp 1-7 (2003); Rutishauser, J., et al., Swiss Med Wkly, 132, pp 211-222 (2002); Morello, JP et al., TIPS, 21, pp. 466- 469 (2000); Bross P., et al., Human Mut., 14, pp. 186-198 (1999)]. Studies have shown, however, that $\Delta F508$ -CFTR, when presented at the plasma membrane is functional as a cAMP-responsive

Cl-Channel;(Dalemans et al. (1991), Nature Lond. 354: 526-528 Denning et al., supra.; Pasyk and Foskett (1995), J. Cell. Biochem. 270: 12347-50).

[0009] Although CFTR transports a variety of molecules in addition to anions, this role of transporting anions represents an important element in the overall cellular machinery for transporting ions and water across the epithelium. The other elements include the epithelial Na⁺ channel, ENaC, Na⁺/2Cl⁻/K⁺ co-transporter, Na⁺-K⁺-ATPase pump and the basolateral membrane K⁺ channels, that are responsible for the uptake of chloride into the cell.

[0010] These elements work together to achieve directional transport across the epithelium via their selective expression and localization within the cell. Chloride absorption takes place by the coordinated activity of: (i) ENaC and CFTR present on the apical membrane; and (ii) the Na⁺-K⁺-ATPase pump and Cl⁻ channels expressed on the basolateral surface of the cell. Secondary active transport of chloride from the luminal side leads to the accumulation of intracellular chloride, which can then passively leave the cell via Cl⁻ channels, resulting in a vectorial transport. Arrangement of Na⁺/2Cl⁻/K⁺ co-transporter, Na⁺-K⁺-ATPase pump and the basolateral membrane K⁺ channels on the basolateral surface and CFTR on the luminal side coordinate the secretion of chloride via CFTR on the luminal side. Because water is probably never actively transported itself, its flow across epithelia depends on tiny transepithelial osmotic gradients generated by the bulk flow of sodium and chloride.

[0011] In addition to CF, modulation of CFTR activity may be beneficial for other diseases not directly caused by mutations in CFTR, such as secretory diseases and other protein folding diseases mediated by CFTR. These include, but are not limited to, chronic obstructive pulmonary disease (hereinafter "COPD"), dry eye disease, and Sjögren's Syndrome.

[0012] COPD is characterized by airflow limitation that is progressive and not fully reversible. The airflow limitation is due to mucus hypersecretion, emphysema, and bronchiolitis. Activators of mutant or wild-type CFTR offer a potential treatment of mucus hypersecretion and impaired mucociliary clearance that is common in COPD. Specifically, increasing anion secretion across CFTR may facilitate fluid transport into the airway surface liquid to hydrate the mucus and optimized periciliary fluid viscosity. This would lead to enhanced mucociliary clearance and to a reduction in the symptoms associated with COPD. Dry eye disease is characterized by a decrease in tear aqueous production and abnormal tear film lipid, protein and mucin profiles. There are many causes of dry eye, some of which include age, Lasik eye surgery, arthritis, medications, chemical/thermal burns, allergies, and diseases, such as CF and Sjögren's syndrome. Increasing anion secretion via CFTR would enhance fluid transport from the corneal endothelial cells and secretory glands surrounding the eye to increase corneal hydration. This would help to alleviate the symptoms associated with dry eye disease. Sjögren's syndrome is an autoimmune disease in which the immune system attacks moisture-producing glands throughout the body, including the eye, mouth, skin, respiratory tissue, liver, vagina, and gut. Symptoms include dry eye, mouth, and vagina, as well as lung disease. The disease is also associated with rheumatoid arthritis, systemic lupus, systemic sclerosis, and polymyositis/dermatomyositis. Defective protein trafficking is believed to cause the disease for which treatment options are limited. Modulators of CFTR activity may hydrate the various organs afflicted by the disease and help to elevate the associated symptoms.

[0013] As discussed above, it is believed that the deletion of residue 508 in $\Delta F508$ -CFTR prevents the nascent protein from folding correctly, resulting in the inability of this mutant protein to exit the ER, and traffic to the plasma membrane. As a result, insufficient amounts of the mature protein are present at the plasma membrane and chloride transport within epithelial tissues is significantly reduced. In fact, this cellular phenomenon of defective ER processing of ABC transporters by the ER machinery, has been shown to be the underlying basis not only for CF disease, but for a wide range of other isolated and inherited diseases. The two ways that the ER machinery can malfunction is either by loss of coupling to ER export of the proteins leading to degradation, or by the ER accumulation of these defective/misfolded proteins [Aridor M, et al., Nature Med., 5(7), pp 745- 751 (1999); Shastry, B.S., et al., Neurochem. International, 43, pp 1-7 (2003); Rutishauser, J., et al., Swiss Med Wkly, 132, pp 211-222 (2002); Morello, JP et al., TIPS, 21, pp. 466- 469 (2000); Bross P., et al., Human Mut., 14, pp. 186-198 (1999)].

[0014] The diseases associated with the first class of ER malfunction are CF (due to misfolded $\Delta F508$ -CFTR), hereditary emphysema (due to $\alpha 1$ -antitrypsin; non Piz variants), hereditary hemochromatosis, coagulation-fibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, mucopolysaccharidoses (due to lysosomal processing enzymes), Sandhof/Tay-Sachs (due to β -hexosaminidase), Crigler-Najjar type II (due to UDP-glucuronyl-sialyl-transferase), polyendocrinopathy/ hyperinsulemia, Diabetes mellitus (due to insulin receptor), Laron dwarfism (due to growth hormone receptor), myeloperoxidase deficiency, primary hypoparathyroidism (due to preproparathyroid hormone), melanoma (due to tyrosinase). The diseases associated with the latter class of ER malfunction are glycanosis CDG type 1, hereditary emphysema (due to $\alpha 1$ -antitrypsin (PiZ variant), congenital hyperthyroidism, osteogenesis imperfecta (due to Type I, II, IV procollagen), hereditary hypofibrinogenemia (due to fibrinogen), ACT deficiency (due to $\alpha 1$ -antichymotrypsin), Diabetes insipidus (DI), neurophyseal DI (due to Vasopressin hormone/V2-receptor), neprogenic DI (due to aquaporin II), Charcot-Marie Tooth syndrome (due to Peripheral myelin protein 22), Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease

(due to β APP and presenilins), Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders such as Huntington, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluyian, and myotonic dystrophy, as well as Spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to Prion protein processing defect), Fabry disease (due to lysosomal α -galactosidase A) and Straussler-Scheinker syndrome (due to Prp processing defect).

[0015] In CF, chloride transport mediated by the CFTR is reduced resulting in the abnormal mucus secretion that characterizes the disease. By contrast in secretory diarrheas epithelial water transport is dramatically increased as a result of secretagogue activated chloride transport. The mechanism involves elevation of CAMP and stimulation of CFTR.

[0016] Although there are numerous causes of diarrhea, the major consequences of diarrheal diseases, resulting from excessive chloride transport are common to all, and include dehydration, acidosis, death and impaired growth.

[0017] Acute and chronic diarrheas represent a major medical problem in many areas of the world. Diarrhea is both a significant factor in malnutrition and the leading cause of death (5,000,000 deaths/year) in children less than five years old.

[0018] Secretory diarrheas are also a dangerous condition in patients of acquired immunodeficiency syndrome (AIDS) and chronic inflammatory bowel disease (IBD). Sixteen million travelers to developing countries from industrialized nations every year develop diarrhea, with the severity and number of cases of diarrhea varying depending on the country and area of travel.

[0019] Diarrhea in barn animals and pets such as cows, pigs and horses, sheep, goats, cats and dogs, also known as scours, is a major cause of death in these animals. Diarrhea can result from any major transition, such as weaning or physical movement, as well as in response to a variety of bacterial or viral infections and generally occurs within the first few hours of the animal's life.

[0020] The most common diarrheal causing bacteria is enterotoxigenic E-coli (ETEC) having the K99 pilus antigen. Common viral causes of diarrhea include rotavirus and coronavirus. Other infectious agents include cryptosporidium, giardia lamblia, and salmonella, among others.

[0021] Symptoms of rotaviral infection include excretion of watery feces, dehydration and weakness. Coronavirus causes a more severe illness in the newborn animals, and has a higher mortality rate than rotaviral infection. Often, however, a young animal may be infected with more than one virus or with a combination of viral and bacterial micro-organisms at one time. This dramatically increases the severity of the disease.

[0022] Accordingly, there is a need for modulators of an ABC transporter activity, and compositions thereof, that can be used to modulate the activity of the ABC transporter in the cell membrane of a mammal.

[0023] There is a need for methods of treating ABC transporter mediated diseases using such modulators of ABC transporter activity.

[0024] There is a need for methods of modulating an ABC transporter activity in an *ex vivo* cell membrane of a mammal.


[0025] There is a need for modulators of CFTR activity that can be used to modulate the activity of CFTR in the cell membrane of a mammal.

[0026] There is a need for methods of treating CFTR-mediated diseases using such modulators of CFTR activity.

[0027] There is a need for methods of modulating CFTR activity in an *ex vivo* cell membrane of a mammal.

SUMMARY OF THE INVENTION

[0028] It has now been found that compounds of this invention, and pharmaceutically acceptable compositions thereof, are useful as modulators of ABC transporter activity. These compounds have the general formula I:



or a pharmaceutically acceptable salt thereof, wherein X, Z, R¹, R², R³, and R⁴ are described generally and in classes and subclasses below.


[0029] These compounds and pharmaceutically acceptable compositions are useful for treating or lessening the severity of a variety of diseases, disorders, or conditions, including, but not limited to, cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-cibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, and others.

teinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, secretory diarrhea or polycystic kidney disease, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, hereditary emphysema, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidolulsian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Strausler-Scheinker syndrome, COPD, dry eye disease, or Sjogren's disease.

DETAILED DESCRIPTION OF THE INVENTION

I. General Description of Compounds of the Invention:

[0030] The present invention relates to compounds of formula I useful as modulators of ABC transporter activity:




or a pharmaceutically acceptable salt thereof,

wherein X is O or S; Z is a bond or is an optionally substituted C_{1-4} alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by $-NR-$, $-S-$, $-O-$, $-SO_2NR-$, $-NRSO_2-$, $-SO_2-$, or $-CO-$; R^1 is hydrogen, or is a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R^1 is optionally substituted at one or more carbon or nitrogen atoms with x independent occurrences of $-QR^x$; wherein x is 0-5; wherein Q is a bond or is a C_1-C_6 alkylidene chain wherein up to two methylene units of Q are optionally and independently replaced by $-CO-$, $-CO_2-$, $-COCO-$, $-CONR-$, $-OCONR-$, $-NRNR-$, $-NRNRCO-$, $-NRCO-$, $-NRCO_2-$, $-NRCONR-$, $-SO-$, $-SO_2-$, $-SO_2-$, $-SO_2NR-$, $-NRSO_2NR-$, $-O-$, $-S-$; or $-NR-$; and each occurrence of R^x is independently R' , halogen, NO_2 , or CN , or $-QR^x$ is $=O$, $=S$, or $=NR'$; each occurrence of R is independently hydrogen or an optionally substituted C_{1-6} aliphatic group; and each occurrence of R' is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or R and R' , two occurrences of R, or two occurrences of R' , are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; R^2 is halogen, $-CN$, $-CF_3$, $-NO_2$ or $-TqR$; R^3 is U_mR' and R^4 is V_pCy^1 ,

wherein m, p, and q are each independently 0 or 1, and U, V, and T are each independently an optionally substituted C_1-C_4 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by $-CO-$, $-CO_2-$, $-COCO-$, $-CONR-$, $-OCONR-$, $-NRNR-$, $-NRNRCO-$, $-NRCO-$, $-NRCO_2-$, $-NRCONR-$, $-SO-$, $-SO_2-$, $-NRSO_2-$, $-SO_2NR-$, $-NRSO_2NR-$, $-O-$, $-S-$; or $-NR-$; Cy^1 is a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein Cy^1 is optionally substituted at one or more carbon or nitrogen atoms with y independent occurrences of $-WR^w$;

wherein y is 0-5; wherein W is a bond or is a C_1-C_6 alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by $-CO-$, $-CO_2-$, $-COCO-$, $-CONR-$, $-OCONR-$, $-NRNR-$, $-NRNRCO-$, $-NRCO-$, $-NRCO_2-$, $-NRCONR-$, $-SO-$, $-SO_2-$, $-NRSO_2-$, $-SO_2NR-$, $-NRSO_2NR-$, $-O-$, $-S-$; or $-NR-$; and each occurrence of R^w is independently R' , halogen, NO_2 , or CN , or $-WR^w$ is $=O$, $=S$, or $=NR'$; or R^3 and R^4 , taken together with the nitrogen atom to which they are bound form an optionally substituted 5-, 6- or 7-


membered saturated, partially unsaturated or fully unsaturated ring having the structure:



wherein r is 0, 1, or 2; one of X^3 , X^4 , or X^5 is $\text{CH-V}_p\text{-Cy}^1$ or $\text{N-V}_p\text{-Cy}^1$, and the others of X^3 , X^4 , or X^5 are CHR' or NR' ; and each occurrence of X^1 , when present, and X^2 are each independently $\text{C(R}')_2$, $-\text{CO}-$, or $-\text{CS}-$.

[0031] In some embodiments, for compounds of formula I as described generally above:


a) when X is S , Z is CH_2 , and R^2 is hydrogen, then R^1 is not:



b) when X is S or O ; and R^2 is formyl, 4,5-dihydro-2-oxazolyl, $-\text{COOR}'$, or $-\text{COSR}'$; then:

- i) when Z is $-\text{CH}_2-$, R^1 is not phenyl optionally substituted with $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_1\text{-C}_4$ alkylthio, $\text{C}_1\text{-C}_4$ haloalkylthio, halogen, cyano, or nitro; or
- ii) when Z is C_{1-6} alkyl, then R^1 is not $\text{C}_3\text{-C}_6$ cycloalkyl optionally substituted with $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_1\text{-C}_4$ alkylthio, $\text{C}_1\text{-C}_4$ haloalkylthio, halogen, cyano, or nitro;


c) when X is S , R^2 is H , Z is $-\text{CH}_2-$, and R^1 is unsubstituted phenyl, then when R^3 is hydrogen, R^4 is not:



d) when X is S , R^2 is SMe or Me , Z is CH_2 and R^1 is unsubstituted phenyl,


- i) when R^3 is hydrogen, then R^4 is not unsubstituted benzyl; and
- ii) when R^3 is ethyl, then R^4 is not unsubstituted phenyl;

e) when X is O , Z is CH_2 , R^1 is unsubstituted phenyl, R^2 is Me , and R^3 is hydrogen, then R^4 is not




f) when X is O , Z is CH_2 , R^1 is unsubstituted phenyl, R^2 is hydrogen, and R^3 is hydrogen, then R^4 is not unsubstituted phenyl, 4-Me-phenyl, or unsubstituted benzyl; and

g) when X is O , Z is $-\text{CH}(\text{iPr})-$, R^1 is




and R^3 is hydrogen, then R^4 is not $-(CH_2)_2-S-Phenyl$;

h) when $Z-R^1$ is methyl, X is S, R^3 is *i*Bu, then R^4 is not:



and

i) when $Z-R^1$ is phenyl, X is S, and R_3 is $-(CH_2)_2$ -pyrid-2-yl, then R^4 is not



[0032] 2. Compounds and Definitions:

[0033] Compounds of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry", 5th Ed., Ed.: Smith, M.B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.

[0034] As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the invention. It will be appreciated that the phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." In general, the term "substituted", whether preceded by the term "optionally" or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term "stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and preferably their recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture, light, or other chemically reactive conditions, for at least a week.

[0035] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle" "cycloaliphatic" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups contain 1-20 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-10 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-8 aliphatic carbon atoms. In still other embodiments, aliphatic groups

contain 1-6 aliphatic carbon atoms, and in yet other embodiments aliphatic groups contain 1-4 aliphatic carbon atoms. In some embodiments, "cycloaliphatic" (or "carbocycle" or "cycloalkyl") refers to a monocyclic C₃-C₈ hydrocarbon or bicyclic C₈-C₁₂ hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic ring system has 3-7 members. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

[0036] The term "heteroaliphatic", as used herein, means aliphatic groups wherein one or two carbon atoms are independently replaced by one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon. Heteroaliphatic groups may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and include "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" groups.

[0037] The term "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" as used herein means non-aromatic, monocyclic, bicyclic, or tricyclic ring systems in which one or more ring members are an independently selected heteroatom. In some embodiments, the "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" group has three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the system contains 3 to 7 ring members.

[0038] The term "heteroatom" means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or NR⁺ (as in N-substituted pyrrolidinyl)).

[0039] The term "unsaturated", as used herein, means that a moiety has one or more units of unsaturation.

[0040] The term "alkoxy", or "thioalkyl", as used herein, refers to an alkyl group, as previously defined, attached to the principal carbon chain through an oxygen ("alkoxy") or sulfur ("thioalkyl") atom.

[0041] The terms "haloalkyl", "haloalkenyl" and "haloalkoxy" means alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms. The term "halogen" means F, Cl, Br, or I.

[0042] The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term "aryl" may be used interchangeably with the term "aryl ring". The term "aryl" also refers to heteroaryl ring systems as defined herein-below.

[0043] The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

[0044] An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl and heteroarylalkoxy and the like) group may contain one or more substituents and thus may be "optionally substituted". Unless otherwise defined above and herein, suitable substituents on the unsaturated carbon atom of an aryl or heteroaryl group are generally selected from halogen; -R^o; -OR^o; -SR^o; phenyl (Ph) optionally substituted with R^o; -O(Ph) optionally substituted with R^o; -(CH₂)₁₋₂(Ph), optionally substituted with R^o; -CH=CH(Ph), optionally substituted with R^o; -NO₂; -CN; -N(R^o)₂; -NR^oC(O)R^o; -NR^oC(S)R^o; -NR^oC(O)N(R^o)₂; -NR^oC(S)N(R^o)₂; -NR^oCO₂R^o; -NR^oNR^oC(O)R^o; -NR^oNR^oC(O)N(R^o)₂; -NR^oNR^oCO₂R^o; -C(O)C(O)R^o; -C(O)CH₂C(O)R^o; -CO₂R^o; -C(O)R^o; -C(S)R^o; -C(O)N(R^o)₂; -C(S)N(R^o)₂; -OC(O)N(R^o)₂; -OC(O)R^o; -C(O)N(OR^o)R^o; -C(NOR^o)R^o; -S(O)₂R^o; -S(O)R^o; -SO₂N(R^o)₂; -S(O)R^o; -NR^oSO₂N(R^o)₂; -NR^oSO₂R^o; -N(OR^o)R^o; -C(=NH)-N(R^o)₂; -P(O)₂R^o; -PO(R^o)₂; -OPO(R^o)₂; -(CH₂)₀₋₂NHC(O)R^o; phenyl (Ph) optionally substituted with R^o; -O(Ph) optionally substituted with R^o; -(CH₂)₁₋₂(Ph), optionally substituted with R^o; or -CH=CH(Ph), optionally substituted with R^o; wherein each independent occurrence of R^o is selected from hydrogen, optionally substituted C₁₋₆ aliphatic, an unsubstituted 5-6 membered heteroaryl or heterocyclic ring, phenyl, -O(Ph), or -CH₂(Ph), or, notwithstanding the definition above, two independent occurrences of R^o, on the same substituent or different substituents, taken together with the atom(s) to which each R^o group is bound, to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0045] Optional substituents on the aliphatic group of R^o are selected from NH₂, NH(C₁₋₄aliphatic), N(C₁₋₄aliphatic)₂, halogen, C₁₋₄aliphatic, OH, O(C₁₋₄aliphatic), NO₂, CN, CO₂H, CO₂(C₁₋₄aliphatic), O(haloC₁₋₄ aliphatic), or haloC₁₋₄aliphatic, wherein each of the foregoing C₁₋₄aliphatic groups of R^o is unsubstituted.

[0046] An aliphatic or heteroaliphatic group, or a non-aromatic heterocyclic ring may contain one or more substituents and thus may be "optionally substituted". Unless otherwise defined above and herein, suitable substituents on the saturated carbon of an aliphatic or heteroaliphatic group, or of a non-aromatic heterocyclic ring are selected from those listed above for the unsaturated carbon of an aryl or heteroaryl group and additionally include the following: =O, =S,

=NNHR*, =NN(R*)₂, =NNHC(O)R*, =NNHCO₂(alkyl), =NNHSO₂(alkyl), or =NR*, where each R* is independently selected from hydrogen or an optionally substituted C₁₋₆ aliphatic group.


[0047] Unless otherwise defined above and herein, optional substituents on the nitrogen of a non-aromatic heterocyclic ring are generally selected from -R⁺, -N(R⁺)₂, -C(O)R⁺, -CO₂R⁺, -C(O)C(O)R⁺, -C(O)CH₂C(O)R⁺, -SO₂R⁺, -SO₂N(R⁺)₂, -C(=S)N(R⁺)₂, -C(=NH)-N(R⁺)₂, or -NR⁺SO₂R⁺; wherein R⁺ is hydrogen, an optionally substituted C₁₋₆ aliphatic, optionally substituted phenyl, optionally substituted -O(Ph), optionally substituted -CH₂(Ph), optionally substituted -(CH₂)₁₋₂(Ph); optionally substituted -CH=CH(Ph); or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring having one to four heteroatoms independently selected from oxygen, nitrogen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R⁺, on the same substituent or different substituents, taken together with the atom(s) to which each R⁺ group is bound, form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0048] Optional substituents on the aliphatic group or the phenyl ring of R⁺ are selected from -NH₂, -NH(C₁₋₄ aliphatic), -N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, -OH, -O(C₁₋₄ aliphatic), -NO₂, -CN, -CO₂H, -CO₂(C₁₋₄ aliphatic), -O(halo C₁₋₄ aliphatic), or halo(C₁₋₄ aliphatic), wherein each of the foregoing C₁₋₄ aliphatic groups of R⁺ is unsubstituted.


[0049] The term "alkylidene chain" refers to a straight or branched carbon chain that may be fully saturated or have one or more units of unsaturation and has two points of attachment to the rest of the molecule.

[0050] As detailed above, in some embodiments, two independent occurrences of R[°] (or R⁺, R, R' or any other variable similarly defined herein), are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0051] Exemplary rings that are formed when two independent occurrences of R[°] (or R⁺, R, R' or any other variable similarly defined herein), are taken together with the atom(s) to which each variable is bound include, but are not limited to the following: a) two independent occurrences of R[°] (or R⁺, R, R' or any other variable similarly defined herein) that are bound to the same atom and are taken together with that atom to form a ring, for example, N(R[°])₂, where both occurrences of R[°] are taken together with the nitrogen atom to form a piperidin-1-yl, piperazin-1-yl, or morpholin-4-yl group; and b) two independent occurrences of R[°] (or R⁺, R, R' or any other variable similarly defined herein) that are bound to different atoms and are taken together with both of those atoms to form a ring, for example where a phenyl group is substituted with two occurrences of OR[°]



these two occurrences of R[°] are taken together with the oxygen atoms to which they are bound to form a fused 6-membered oxygen containing ring:



It will be appreciated that a variety of other rings can be formed when two independent occurrences of R[°] (or R⁺, R, R' or any other variable similarly defined herein) are taken together with the atom(s) to which each variable is bound and that the examples detailed above are not intended to be limiting.

[0052] Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ¹³C- or ¹⁴C-enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

[0053] The term "ABC-transporter" as used herein means an ABC-transporter protein or a fragment thereof comprising



at least one binding domain, wherein said protein or fragment thereof is present *in vivo* or *in vitro*. The term "binding domain" as used herein means a domain on the ABC-transporter that can bind to a modulator. See, e.g., Hwang, T. C. et al., J. Gen. Physiol. (1998): 111(3), 477-90.

[0054] The term "CFTR" as used herein means cystic fibrosis transmembrane conductance regulator or a mutation thereof capable of regulator activity, including, but not limited to, $\Delta F508$ CFTR and G551D CFTR (see, e.g., <http://www.genet.sickkids.on.ca/cftr/>, for CFTR mutations).

[0055] The term "modulating" as used herein means increasing or decreasing by a measurable amount.

[0056] 3. Description of Exemplary Compounds:

[0057] As described generally above, for compounds of the invention, X is O or S and compounds of formula **I-A** or **I-B** are provided:









**I-A****I-B**


[0058] As described generally above, Z is a bond or an optionally substituted C_{1-4} alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by -NR-, -S-, -O-, or -CO-. In certain exemplary embodiments, Z is $-[C(R^5)_2]_n-$, wherein n is 0, 1, 2, or 3, and each occurrence of R^5 is independently halogen, CN, NO_2 , or -YR, wherein Y is a bond or is an optionally substituted C_1-C_4 alkylidene chain wherein up to two methylene units of Y are optionally and independently replaced by -O-, -NR-, -CO-, -S-, -SO-, or -SO₂-. In other exemplary embodiments Z is $-[C(R^5)_2]_nO-$, wherein n is 0, 1, 2, or 3, and each occurrence of R^5 is independently halogen, CN, NO_2 , or -YR, wherein Y is a bond or is an optionally substituted C_1-C_4 alkylidene chain wherein up to two methylene units of Y are optionally and independently replaced by -O-, -NR-, -CO-, -S-, -SO-, or -SO₂-. In still other exemplary embodiments Z is $-[C(R^5)_2]_nS-$, wherein n is 1, 2, or 3, and each occurrence of R^5 is independently halogen, CN, NO_2 , or -YR, wherein Y is a bond or is an optionally substituted C_1-C_4 alkylidene chain wherein up to two methylene units of Y are optionally and independently replaced by -O-, -NR-, -CO-, -S-, -SO-, or -SO₂-.

[0059] In certain other embodiments, Z is a bond. In yet other embodiments, Z is $-C(R^5)_2-$.


[0060] In certain embodiments, Z is $-CH_2-$ or $-CH_2-CH_2-$. In some embodiments, Z is $-CH_2-$.

[0061] As described generally above, R^1 is hydrogen, or is a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R^1 is optionally substituted at one or more carbon atoms with x independent occurrences of -QR^x; wherein x is 0-5. In some embodiments R^1 is hydrogen. In other embodiments R^1 is selected from one of the following groups:


**a****b****c****d****e****f****g****h**




i




j




k




l




m




n




o




p




q




r




s




t




u




v




w




x




y




z




aa




bb




cc



dd



ee




[0062] In some embodiments R^1 is one of rings a, b, c, d, m, n, o, ee, gg, or pp. In other embodiments R^1 is ring a. In certain other embodiments, R^1 is ring gg.

[0063] It will be appreciated that, as described generally above, R^1 is optionally substituted at one or more carbon or nitrogen atoms with x independent occurrences of $-QR^x$; wherein x is 0-5; wherein Q is a bond or is a C_1 - C_6 alkylidene chain wherein up to two methylene units of Q are optionally and independently replaced by $-CO-$, $-CO_2-$, $-COCO-$, $-CONR-$, $-OCONR-$, $-NRNR-$, $-NRNRCO-$, $-NRCO-$, $-NRCO_2-$, $-NRCONR-$, $-SO-$, $-SO_2-$, $-NRO_2-$, $-SO_2NR-$, $-NRSO_2NR-$, $-O-$, $-S-$; or $-NR-$; and each occurrence of R^x is independently R' , halogen, NO_2 , or CN , or $-QR^x$ is $=O$, $=S$, or $=NR'$. In certain embodiments, x is 0, 1, 2 or 3, and Q is a bond or is an optionally substituted C_1 - C_6 alkylidene chain wherein one or two methylene units are optionally and independently replaced by O , NR , S , SO_2 , or CO_2 , CO , and R^x is R' or halogen. In other embodiments, x is 0, 1, 2, or 3 and each occurrence of QR^x , when present, is independently $-C_{1-3}alkyl$, $-O(C_{1-3}alkyl)$, $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)_2N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, or SO_2NRR' .

[0064] As described generally above, R^2 is halogen, $-CF_3$, $-CN$, $-NO_2$, or $-TqR$. In certain embodiments, R^2 is TqR , wherein T is a bond or is a C_1 - C_4 alkylidene chain, wherein one or two methylene units of T are optionally replaced by $-CO-$, $-CONR-$, $-SO_2-$, $-NRSO_2-$, $-SO_2NR-$, $-O-$, $-S-$, or $-NR$. In other preferred embodiments, R^2 is hydrogen or is optionally substituted C_1 - C_4 alkyl. In still other preferred embodiments, R^2 is hydrogen, $-CF_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH_2OR-(CH_2)_2OR$, $-(CH_2)_3OR$, $-CH_2N(R)_2$, $-(CH_2)_2N(R)_2$, $-(CH_2)_3N(R)_2$, $-CH_2NRCOR-(CH_2)_2NRCOR$, or $-(CH_2)_3NRCOR$.


[0065] As described generally above, R^3 is U_mR' and R^4 is V_pCy^1 ; wherein m , p , U , V , and Cy^1 are defined generally above and herein; or R^3 and R^4 , taken together with the nitrogen atom to which they are bound form an optionally substituted 5-, 6- or 7-membered saturated, partially unsaturated or fully unsaturated ring having the structure:



wherein r is 0, 1, or 2; one of X^3 , X^4 , or X^5 is $\text{CH-V}_p\text{-Cy}^1$ or $\text{N-V}_p\text{-Cy}^1$, and the others of X^3 , X^4 , or X^5 is CHR' or NR' ; each occurrence of X^1 , when present, and X^2 are each independently $\text{C(R}')_2$, $-\text{CO}-$, or $-\text{CS}-$.

[0066] In certain embodiments for compounds of general formulas **I**, **I-A**, or **I-B**, R^3 is $\text{U}_m\text{R}'$ and R^4 is V_pCy^1 . In some embodiments, R^3 is hydrogen. In yet other embodiments, R^3 is optionally substituted C_{1-4} alkyl. In yet other embodiments R^3 is $\text{U}_m\text{R}'$ where m is 1 and U is $-\text{CH}_2-$ and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain exemplary embodiments, R' is an optionally substituted phenyl or pyridyl group. In some embodiments, p is 1 and V is $-\text{CH}_2\text{C(R}^6)_2-$ or $-\text{C(R}^6)_2-$, wherein each occurrence of R^6 is independently halogen, CN , NO_2 , or $-\text{YR}$, wherein Y is a bond or is an optionally substituted $\text{C}_1\text{-C}_4$ alkylidene chain wherein up to two methylene units of Y are optionally and independently replaced by $-\text{O}-$, $-\text{NR}-$, $-\text{CO}-$, $-\text{S}-$, $-\text{SO}-$, or $-\text{SO}_2-$, or wherein the two occurrences of R^6 on the same carbon atom are taken together to form an optionally substituted 3-6-membered spiro ring having 0-3 heteroatoms. In certain embodiments, R^6 is hydrogen, methyl, or two occurrences of R^6 on the same carbon atom are taken together to form an optionally substituted 3-6-membered spiro ring having 0, 1 or 2 heteroatoms selected from nitrogen, oxygen, or sulfur. In still other embodiments, p is 0 and R^4 is Cy^1 .


[0067] In yet other embodiments, R^3 and R^4 , taken together with the nitrogen atom to which they are bound form an optionally substituted 5-, 6- or 7-membered saturated, partially unsaturated or fully unsaturated ring having the structure:




wherein r is 0, 1, or 2; one of X^3 , X^4 , or X^5 is $\text{CH-V}_p\text{-Cy}^1$ or $\text{N-V}_p\text{-Cy}^1$, and the others of X^3 , X^4 , or X^5 is CHR' or NR' ; each occurrence of X^1 , when present, and X^2 are each independently $\text{C(R}')_2$, $-\text{CO}-$, or $-\text{CS}-$. In certain embodiments, r is 1, X^1 , X^2 , X^3 and X^5 are each CH_2 , and X^4 is $\text{CH-V}_p\text{-Cy}^1$, or $\text{N-V}_p\text{-Cy}^1$. In certain other embodiments, p is 1 and V is SO_2 , $-\text{NRSO}_2$, CO , or NRCO . In still other embodiments, X^4 is $\text{N-V}_p\text{-Cy}^1$, p is 1 and V is SO_2 or CO .


[0068] As described generally above, Cy^1 is a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein Cy^1 is optionally substituted at one or more carbon or nitrogen atoms with y independent occurrences of WR^w ; wherein y is 0-5; wherein W is a bond or is a $\text{C}_1\text{-C}_6$ alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by $-\text{CO}-$, $-\text{CO}_2-$, $-\text{COCO}-$, $-\text{CONR}-$, $-\text{OCONR}-$, $-\text{NRNR}-$, $-\text{NRNRCO}-$, $-\text{NRCO}-$, $-\text{NRCO}_2-$, $-\text{NRCONR}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NRSO}_2-$, $-\text{SO}_2\text{NR}-$, $-\text{NRSO}_2\text{NR}-$, $-\text{O}-$, $-\text{S}-$; or $-\text{NR}-$; and each occurrence of R^w is independently R' , halogen, NO_2 or CN , or $-\text{WR}^w$ is $=\text{O}$, $=\text{S}$, or $=\text{NR}'$.

[0069] In certain embodiments, Cy^1 is selected from one of the following rings:







ix




x




xi




xii




xiii




xiv




xv




xvi




xvii




xviii



xix




xx




xxi




xxii




xxiii




xxiv



xxv




xxvi




xxvii




xxviii




xxix



xxx



xxxi




[0070] It will be appreciated that, as described generally above, Cy¹ is optionally substituted at one or more carbon or nitrogen atoms with y independent occurrences of -WR^W; y is 0-5;

wherein W is a bond or is a C₁-C₆ alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by -CO-, -CO₂-, -COCO-, -CONR-, -OCONR-, -NRNR-, -NRNRCO-, -NRCO-, -NRCO₂-, -NRCONR-, -SO-, -SO₂-, -NRSO₂-, -SO₂NR-, -NRSO₂NR-, -O-, -S-, or -NR-; and each occurrence of R^W is independently R', halogen, NO₂, or CN, or -WR^W is =O, =S, or =NR'. In certain embodiments, y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C₁-C₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO₂, or CO₂, CO, and R^W is R' or halogen. In other embodiments, y is 0, 1, 2, or 3 and each occurrence of WR^W, when present, is independently -C₁₋₃alkyl, -O(C₁₋₃alkyl), -CF₃, -OCF₃, -SCF₃, -F, -Cl, -Br, -COOR', -COR', -O(CH₂)₂N(R)(R'), -O(CH₂)N(R)(R'), -CON(R)(R'), -(CH₂)₂OR', -(CH₂)OR', optionally substituted phenyl, optionally substituted benzyl, -N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), or SO₂NRR'.


[0071] In addition to the compounds and subsets described above, certain additional compounds are of interest.

[0072] For example, in certain embodiments, Z is CH₂, R² and R³ are each hydrogen, and R⁴ is V_pCy¹, where p is 1 and V is -CH₂C(R⁶)₂ and compounds have the general formula II:




where R¹, X, R⁶ and Cy¹ are as defined generally above and in subsets herein.

[0073] As described generally above, X is S or O and compounds of formula II-A or II-B are provided:




II-A




II-B


[0074] In some embodiments, for compounds of general formulae II, II-A, or II-B, R¹ is selected from one of the following groups:




a




b




c




d




e




f




g




h




i




j




k




l




m




n




o




p




q




r



s




t




u




v




w




x




y




z




aa




bb




cc




dd




ee




ff




gg




hh




ii




jj




kk




ll




mm




nn




oo



pp



qq



















rr


[0075] In other embodiments R^1 is one of rings **a**, **b**, **c**, **d**, **m**, **n**, **o**, **ee**, **gg**, or **pp**. In other embodiments R^1 is ring **a**. In certain other embodiments, R^1 is ring **gg**.

[0076] In certain embodiments, for compounds of formulae **II**, **II-A**, or **II-B**, x is 0, 1, 2 or 3, and Q is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^x is R' or halogen. In other embodiments, x is 0, 1, 2, or 3 and each occurrence of QR^x , when present, is independently $-C_{1-3}$ alkyl, $-O(C_{1-3}$ alkyl), $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, - or SO_2NRR' .


[0077] In yet other embodiments, for compounds of general formulae **II**, **II-A**, or **II-B**, R^6 is hydrogen, methyl, or two occurrences of R^6 on the same carbon atom are taken together to form an optionally substituted 3-6-membered spiro ring having 0, 1 or 2 heteroatoms selected from nitrogen, oxygen, or sulfur.

[0078] In certain embodiments, for compounds of general formulae **II**, **II-A**, or **II-B**, Cy^1 is selected from one of the following rings:


**i****ii****iii****iv****v****vi****vii****viii****ix****x****xi****xii****xiii****xiv****xv****xvi**




xvii




xviii



xix




xx




xxi




xxii




xxiii




xxiv



xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv






xxxvi



xxxvii






xxxviii



xxxix

xL

xLi




xLii

xLiii

xLiv



[0079] In certain embodiments, for compounds of formulae **II**, **II-A**, or **II-B**, y is 0-5; wherein W is a bond or is a C_1 - C_6 alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by $-CO-$, $-CO_2-$, $-COCO-$, $-CONR-$, $-OCONR-$, $-NRNR-$, $-NRNRCO-$, $-NRCO-$, $-NRCO_2-$, $-NRCONR-$, $-SO-$, $-SO_2-$, $-NRSO_2-$, $-SO_2NR-$, $-NRSO_2NR-$, $-O-$, $-S-$; or $-NR-$; and each occurrence of R^W is independently R' , halogen, NO_2 , or CN , or $-WR^W$ is $=O$, $=S$, or $=NR'$. In certain embodiments, y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two methylene units are optionally and independently replaced by O , NR , S , SO_2 or CO_2 , and R^W is R' or halogen. In other embodiments, y is 0, 1, 2, or 3 and each occurrence of WR^W , when present, is independently $-C_{1-3}$ alkyl, $-O(C_{1-3}$ alkyl), $-OPh$, $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)_2N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, or SO_2NRR' .

[0080] In yet other embodiments, Z is a bond and R^1 is 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; R^2 is hydrogen; R^3 is U_mR' where m is 0 or 1, and U , when present, is $-CH_2-$ and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and R^4 is $-CH_2Cy^1$ and compounds of formula **III** are provided:

**III**


where R^1 , X , U , m , R' , and Cy^1 are as defined generally above and in subsets herein.

[0081] As described generally above, X is S or O and compounds of formula **III-A** or **III-B** are provided:


**III-A****III-B**

[0082] In some embodiments, for compounds of general formulae **III**, **III-A**, or **III-B**, R^1 is selected from one of the following groups:


5




a



b




c




d


10




e



f




g




h


15




i



j




k




l

20


25




m



n




o




p

30


35




q



r




s




t

40


45




u



v























w



x

50

55



























[0083] In other embodiments R^1 is one of rings **a**, **b**, **c**, **d**, **m**, **n**, **o**, **ee**, **gg**, or **pp**. In other embodiments R^1 is ring **a**. In certain other embodiments, R^1 is ring **gg**.


[0084] In certain embodiments, for compounds of formulae **III**, **III-A**, or **III-B**, x is 0, 1, 2 or 3, and Q is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^X is R' or halogen. In other embodiments, x is 0, 1, 2, or 3 and each occurrence of QR^X , when present, is independently $-C_{1-3}$ alkyl, $-O(C_{1-3}$ alkyl), $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, - or SO_2NRR' .

[0085] As described generally above for compounds of formula **III**, R^3 is U_mR' where m is 0 or 1, and U , when present, is $-CH_2-$ and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring


having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, R' is an optionally substituted phenyl or pyridyl group.

[0086] In certain embodiments, for compounds of general formulae **III**, **III-A**, or **III-B**, Cy¹ is selected from one of the following rings:


**i****ii****iii****iv****v****vi****vii****viii****ix****x****xi****xii****xiii****xiv****xv****xvi****xvii****xviii****xix****xx****xxi****xxii****xxiii****xxiv**



xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi



xLii



xLiii




xLiv

[0087] In certain embodiments, for compounds of formulae III, III-A, or III-B, y is 0-5;



wherein W is a bond or is a C₁-C₆ alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by -CO-, -CO₂-, -COCO-, -CONR-, -OCONR-, -NRNR-, -NRNRCO-, -NRCO-, -NRCO₂-, -NRCONR-, -SO-, -SO₂-, -NRSO₂-, -SO₂NR-, -NRSO₂NR-, -O-, -S-, or -NR-; and each occurrence of R^W is independently R', halogen, NO₂, or CN, or -WR^W is =O, =S, or =NR'. In certain embodiments, y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C₁₋₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO₂ or CO₂ CO, and R^W is R' or halogen. In other embodiments, y is 0, 1, 2, or 3 and each occurrence of WR^W, when present, is independently -C₁₋₃alkyl, -O(C₁₋₃alkyl), -CF₃, -OCF₃, -SCF₃, -F, -Cl, -Br, -COOR', -COR', -O(CH₂)₂N(R)(R'), -O(CH₂)N(R)(R'), -CON(R)(R'), -(CH₂)₂OR', -(CH₂)OR', optionally substituted phenyl, optionally substituted benzyl, -N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), or SO₂NRR'.

[0088] In still other embodiments, Z-R¹ is -CH₃; R² is hydrogen; R³ is U_mR' where m is 0 or 1, and U, when present, is -CH₂- and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and R⁴ is -CH₂Cy¹ and compounds of formula **IV** are provided:

**IV**













where X, U, m, R', and Cy¹ are as defined generally above and in subsets herein.


[0089] As described generally above, X is S or O and compounds of formula **IV-A** or **IV-B** are provided:

**IV-A****IV-B**


[0090] As described generally above for compounds of formula **IV**, R³ is U_mR' where m is 0 or 1, and U, when present, is -CH₂- and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, R' is an optionally substituted phenyl or pyridyl group.

[0091] In certain embodiments, for compounds of general formulae **IV**, **IV-A**, or **IV-B**, Cy¹ is selected from one of the following rings:


**i****ii****iii****iv****v****vi****vii****viii****ix****x****xi****xii**




xiii




xiv




xv




xvi




xvii




xviii



xix




xx




xxi




xxii




xxiii




xxiv



xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi



xLii



xLiii




xLiv


[0092] In certain embodiments, for compounds of formulae **IV**, **IV-A**, or **IV-B**, y is 0-5;

wherein W is a bond or is a C_1 - C_6 alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by $-CO-$, $-CO_2-$, $-COCO-$, $-CONR-$, $-OCONR-$, $-NRNR-$, $-NRNRCO-$, $-NRCO-$, $-NRCO_2-$, $-NRCONR-$, $-SO-$, $-SO_2-$, $-NRSO_2-$, $-SO_2NR-$, $-NRSO_2NR-$, $-O-$, $-S-$; or $-NR-$; and each occurrence of R^W is independently R' , halogen, NO_2 , or CN , or $-WR^W$ is $=O$, $=S$, or $=NR'$. In certain embodiments, y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two methylene units are optionally and independently replaced by O , NR , S , SO_2 , or CO_2 , CO , and R^W is R' or halogen. In other embodiments, y is 0, 1, 2, or 3 and each occurrence of WR^W , when present, is independently $-C_{1-3}$ alkyl, $-O(C_{1-3}$ alkyl), $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, or SO_2NRR' .

[0093] In yet other embodiments, Z is $-CH_2-$; and R^3 and R^4 , taken together with the nitrogen atom to which they are bound form an optionally substituted 5-, 6- or 7-membered saturated, partially unsaturated or fully unsaturated ring having the structure:




wherein r is 0, 1, or 2; one of X^3 , X^4 , or X^5 is $CH-V_p-Cy^1$ or $N-V_p-Cy^1$, and the others of X^3 , X^4 , or X^5 is CHR' or NR' ; each occurrence of X^1 , when present, and X^2 are each independently $C(R')_2$, $-CO-$, or $-CS-$, and compounds of formula **V** are provided:




V

where R^1 , X , R^2 , X^1 , X^2 , X^3 , X^4 , X^5 and r are as described generally above and in subsets herein.

[0094] As described generally above, X is S or O and compounds of formula **V-A** or **V-B** are provided:




V-A




V-B


[0095] In some embodiments, for compounds of general formulae **V**, **V-A**, or **V-B**, R^1 is selected from one of the following groups:




a




b




c




d




e




f




g




h




i




j




k




l




m




n




o




p




q




r



s




t




u




v




w




x




y




z




aa




bb




cc




dd




ee




ff




gg




hh




ii




jj




kk




ll




mm




nn




oo



pp



qq



















rr


[0096] In other embodiments R^1 is one of rings **a**, **b**, **c**, **d**, **m**, **n**, **o**, **ee**, **gg**, or **pp**. In other embodiments R^1 is ring **a**. In certain other embodiments, R^1 is ring **gg**.

[0097] In certain embodiments, for compounds of formulae **V**, **V-A**, or **V-B**, x is 0, 1, 2 or 3, and Q is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^X is R' or halogen. In other embodiments, x is 0, 1, 2, or 3 and each occurrence of QR^X , when present, is independently $-C_{1-3}$ alkyl, $-O(C_{1-3}$ alkyl), $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, $-$ or SO_2NRR' .


[0098] In certain embodiments, for compounds of general formulae **V**, **V-A**, or **V-B**, r is 1, X^1 , X^2 , X^3 and X^5 are each CH_2 , and X^4 is $CH-V_p-Cy^1$, or $N-V_p-Cy^1$. In certain other embodiments, p is 1 and V is SO_2 , $-NRSO_2$, CO, or $NRCO$. In still other embodiments, X^4 is $N-V_p-Cy^1$, p is 1 and V is SO_2 or CO. In yet other embodiments, for compounds of general formula **V-A** or **V-B**, X^4 is $N-V_p-Cy^1$, p is 1 and V is SO_2 .

[0099] In certain embodiments, for compounds of general formulae **V**, **V-A**, or **V-B**, Cy^1 is selected from one of the following rings:


**i****ii****iii****iv****v****vi****vii****viii****ix****x****xi****xii****xiii****xiv****xv****xvi**




xvii




xviii



xix




xx




xxi




xxii




xxiii




xxiv



xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi



xLii




xLiii



xLiv



[0100] In certain embodiments, for compounds of general formulae **V**, **V-A**, or **V-B**, y is 0-5; wherein W is a bond or is a C_1 - C_6 alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by $-CO-$, $-CO_2-$, $-COCO-$, $-CONR-$, $-OCONR-$, $-NRNR-$, $-NRNRCO-$, $-NRCO-$, $-NRCO_2-$, $-NRCONR-$, $-SO-$, $-SO_2-$, $-NRSO_2-$, $-SO_2NR-$, $-NRSO_2NR-$, $-O-$, $-S-$; or $-NR-$; and each occurrence of R^W is independently R' , halogen, NO_2 , or CN , or $-WR^W$ is $=O$, $=S$, or $=NR'$. In certain embodiments, y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two methylene units are optionally and independently replaced by O , NR , S , SO_2 , or CO_2 , CO , and R^W is R' or halogen. In other embodiments, y is 0, 1, 2, or 3 and each occurrence of WR^W , when present, is independently $-C_{1-3}$ alkyl, $-O(C_{1-3}$ alkyl), $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, or SO_2NRR' .

[0101] In another embodiment, the present invention provides compounds of formula **VI**:

**VI;**

wherein: ring A is a 3-7 membered ring having up to 3 heteroatoms selected from O, S, or NR; R , R^1 , Z , X , and Cy_1 are as defined above.


[0102] As described generally above, X is S or O, and compounds of formula **VI-A** or **VI-B** are provided:

**VI-A****VI-B.**


[0103] In one embodiment, ring A is a 3-7 membered cycloalkyl ring. In another embodiment, ring A is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In certain embodiments, ring A is cyclopentyl or cyclohexyl.

[0104] In another embodiment, ring A has one oxygen ring heteroatom. Or, ring A has one sulfur ring heteroatom. Or, ring A has one nitrogen ring heteroatom. Preferably, R (in NR) is hydrogen.


[0105] In some embodiments, for compounds of general formulae **VI**, **VI-A**, or **VI-B**, R^1 is selected from one of the following groups:




a




b




c




d




e




f




g




h




i




j




k




l




m




n




o




p




q




r




s




t




u























v



w



x





[0106] In other embodiments R^1 is one of rings **a**, **b**, **c**, **d**, **m**, **n**, **o**, **ee**, or **pp**. In yet other embodiments R^1 is phenyl (ring **a**).

[0107] In certain embodiments, for compounds of formulae **VI**, **VI-A**, or **VI-B**, x is 0, 1, 2 or 3, and Q is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^X is R' or halogen. In other embodiments, x is 0, 1, 2, or 3 and each occurrence of QR^X , when present, is independently $-C_{1-3}alkyl$, $-O(C_{1-3}alkyl)$, $-OPh$, $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, or SO_2NRR' .


[0108] In certain embodiments, for compounds of general formulae **VI**, **VI-A**, or **VI-B**, Cy^1 is selected from one of the

following rings:







XXV




XXVI




XXVII




XXVIII




XXIX




XXX




XXXI




XXXII




XXXIII




XXXIV




XXXV




XXXVI




XXXVII




XXXVIII




XXXIX




XL




xLi



xLii



xLiii





xLiv

[0109] In one embodiment, Cy_1 is ring i or ring xxxiii above.

[0110] In certain embodiments, for compounds of general formulae VI, VI-A, or VI-B, y is 0-5; wherein W is a bond or is a C_1 - C_6 alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by -CO-, -CO₂-, -COCO-, -CONR-, -OCONR-, -

Table 1. Examples of Compounds of Formula I:

Cmpd #	Compound	Cmpd #	Compound
1		2	

NRNR-, -NRNRCO-, -NRCO-, -NRCO₂-, -NRCONR-, -SO-, -SO₂-, -NRSO₂-, -SO₂NR-, -NRSO₂NR-, -O-, -S-; or -NR-; and each occurrence of R^W is independently R', halogen, NO₂, or CN, or -WR^W is =O, =S, or =NR'. In certain embodiments, y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C₁₋₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO₂, or CO₂, CO, and R^W is R' or halogen. In other embodiments, y is 0, 1, 2, or 3 and each occurrence of WR^W, when present, is independently -C₁₋₃alkyl, -O(C₁₋₃alkyl), -CF₃, -OCF₃, -SCF₃, -F, -Cl, -Br, -COOR', -COR', -O(CH₂)₂N(R)(R'), -O(CH₂)N(R)(R'), -CON(R)(R'), -(CH₂)₂OR', -(CH₂)OR', optionally substituted phenyl, optionally substituted benzyl, -N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), - or SO₂NRR'.



[0111] In one embodiment, Z is -CH₂- or -CH₂-CH₂-. Or, Z is -CH₂-.





[0112] In another embodiment, R¹ is phenyl optionally substituted with up to three substituents selected from -C₁₋₃alkyl, -O(C₁₋₃alkyl), -CF₃, -OCF₃, -SCF₃, -F, -Cl, -Br, -COOR', -COR', -O(CH₂)N(R)(R'), -O(CH₂)N(R)(R'), -CON(R)(R'), -(CH₂)₂OR', -(CH₂)OR', optionally substituted phenyl, optionally substituted benzyl, -N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), - or SO₂NRR'.





[0113] Representative examples of compounds of formula I are set forth below in Table 1.





[0114]





Table 1. Examples of Compounds of Formula I:





Cmpd #	Compound	Cmpd #	Compound
1		2	





Cmpd #	Compound	Cmpd #	Compound
3		5	
4		6	





Cmpd #	Compound	Cmpd #	Compound
7		9	
8		10	





Cmpd #	Compound	Cmpd #	Compound
11		13	
12		14	





Cmpd #	Compound	Cmpd #	Compound
15		17	
16		18	





Cmpd #	Compound	Cmpd #	Compound
19		21	
20		22	





Cmpd #	Compound	Cmpd #	Compound
23		25	
24		26	





Cmpd #	Compound	Cmpd #	Compound
27		29	
28		30	





Cmpd #	Compound	Cmpd #	Compound
31		33	
32		34	





Cmpd #	Compound	Cmpd #	Compound
35		37	
36		38	





Cmpd #	Compound	Cmpd #	Compound
39		41	
40		42	





Cmpd #	Compound	Cmpd #	Compound
43		45	
44		46	





Cmpd #	Compound	Cmpd #	Compound
47		49	
48		50	





Cmpd #	Compound	Cmpd #	Compound
51		53	
52		54	





Cmpd #	Compound	Cmpd #	Compound
55		57	
56		58	





Cmpd #	Compound	Cmpd #	Compound
59		61	
60		62	





Cmpd #	Compound	Cmpd #	Compound
63		65	
64		66	





Cmpd #	Compound	Cmpd #	Compound
67		69	
68		70	





Cmpd #	Compound	Cmpd #	Compound
71		73	
72		74	

Cmpd #	Compound	Cmpd #	Compound
75		77	
76		78	

Cmpd #	Compound	Cmpd #	Compound
79		81	
80		82	

Cmpd #	Compound	Cmpd #	Compound
83		85	
84		86	

Cmpd #	Compound	Cmpd #	Compound
87		89	
88		90	

Cmpd #	Compound	Cmpd #	Compound
91		93	
92		94	

Cmpd #	Compound	Cmpd #	Compound
95	 <chem>CC1=CC=C(C=C1)CCNC(=O)c2nc3c(s3)CCc4cc(Cl)c(Cl)cc4</chem>	97	 <chem>ClC1=CC=C(C=C1)CCNC(=O)c2nc3c(s3)CCc4cc(Cl)c(Cl)cc4</chem>
96	 <chem>COC1=CC=C(C=C1)CCNC(=O)c2nc3c(s3)CCc4cc(Cl)c(Cl)cc4</chem>	98	 <chem>COC1=CC=C(C=C1)CCNC(=O)c2nc3c(s3)CCc4cc(Cl)c(Cl)cc4</chem>

Cmpd #	Compound	Cmpd #	Compound
99		100	


[0115] 4. General Synthetic Methodology:


[0116] The compounds of this invention may be prepared in general by methods known to those skilled in the art for analogous compounds, as illustrated by the general scheme below, and the preparative examples that follow.

[0117] Schemes I and II below depict the synthesis of thiazole and oxazole starting materials where Z is $-\text{CH}_2-$ and R^1 is optionally substituted phenyl.

[0118]

[00119] Scheme I:




Scheme II:**References:**

[0119] Org. Lett., 2000, 2(8), 1165-1168; A. J. Phillips et al. Chem Pharm Bull, 1986, 34(7), 2851; K. Meguro et al.

[0120] **Scheme III:**

[0121] Scheme III depicts general reaction conditions for the coupling of the thiazole or oxazole starting materials (as depicted generally above) and an appropriate amine to generate thiazole or oxazole amides as represented generally by compounds of formula I.



[0122] Although certain exemplary embodiments are depicted and described above and herein, it will be appreciated that a compounds of the invention can be prepared according to the methods described generally above using appropriate starting materials by methods generally available to one of ordinary skill in the art.

[0123] **5. Uses, Formulation and Administration**

[0124] *Pharmaceutically acceptable compositions*

[0125] As discussed above, the present invention provides compounds that are useful as modulators of ABC transporters and thus are useful in the treatment of disease, disorders or conditions such as cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-cibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, secretory diarrhea or polycystic kidney disease, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, palyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, hereditary emphysema, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perliazeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear plasy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebular ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Strausler-Scheinker syndrome, COPD, dry eye disease, or Sjogren's disease.

[0126] Accordingly, in another aspect of the present invention, pharmaceutically acceptable compositions are provided, wherein these compositions comprise any of the compounds as described herein, and optionally comprise a pharmaceutically acceptable carrier, adjuvant or vehicle. In certain embodiments, these compositions optionally further comprise

one or more additional therapeutic agents.

[0127] It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, a pharmaceutically acceptable derivative includes, but is not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or any other adduct or derivative which upon administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

[0128] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. A "pharmaceutically acceptable salt" means any non-toxic salt or salt of an ester of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an active metabolite or residue thereof.

[0129] Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, *et al.* describe pharmaceutically acceptable salts in detail in J. Pharmaceutical Sciences, 1977, 66, 1-19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxyethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[0130] As described above, the pharmaceutically acceptable compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, adjuvant, or vehicle, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutically acceptable compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutically acceptable composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, or potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

[0131] *Uses of Compounds and Pharmaceutically Acceptable Compositions*

[0132] In yet another aspect, the present invention provides a method of treating a condition, disease, or disorder

implicated by ABC transporter activity. In certain embodiments, the present invention provides a method of treating a condition, disease, or disorder implicated by a deficiency of ABC transporter activity, the method comprising administering a composition comprising a compound of formula (I) to a subject, preferably a mammal, in need thereof.

[0133] In certain preferred embodiments, the present invention provides a method of treating cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-cibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, secretory diarrhea or polycystic kidney disease, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, hereditary emphysema, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebular ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Straussler-Scheinker syndrome, COPD, dry eye disease, or Sjogren's disease, comprising the step of administering to said mammal an effective amount of a composition comprising a compound of formula (I), or a preferred embodiment thereof as set forth above.

[0134] According to an alternative preferred embodiment, the present invention provides a method of treating cystic fibrosis comprising the step of administering to said mammal a composition comprising the step of administering to said mammal an effective amount of a composition comprising a compound of formula (I), or a preferred embodiment thereof as set forth above.

[0135] According to the invention an "effective amount" of the compound or pharmaceutically acceptable composition is that amount effective for treating or lessening the severity of one or more of cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-cibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, secretory diarrhea or polycystic kidney disease, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, hereditary emphysema, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebular ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Straussler-Scheinker syndrome, COPD, dry eye disease, or Sjogren's disease.

[0136] The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating or lessening the severity of one or more of cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-cibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, secretory diarrhea or polycystic kidney disease, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, hereditary emphysema, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebular ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Straussler-Scheinker syndrome, COPD, dry eye disease, or Sjogren's disease. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the

specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts. The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

[0137] The pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[0138] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0139] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[0140] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0141] In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[0142] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[0143] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[0144] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in

a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0145] The active compounds can also be in microencapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[0146] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms are prepared by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[0147] As described generally above, the compounds of the invention are useful as modulators of ABC transporters. Thus, without wishing to be bound by any particular theory, the compounds and compositions are particularly useful for treating or lessening the severity of a disease, condition, or disorder where hyperactivity or inactivity of ABC transporters is implicated in the disease, condition, or disorder. When hyperactivity or inactivity of an ABC transporter is implicated in a particular disease, condition, or disorder, the disease, condition, or disorder may also be referred to as a "ABC transporter-mediated disease, condition or disorder". Accordingly, in another aspect, the present invention provides a method for treating or lessening the severity of a disease, condition, or disorder where hyperactivity or inactivity of an ABC transporter is implicated in the disease state.

[0148] The activity of a compound utilized in this invention as a modulator of an ABC transporter may be assayed according to methods described generally in the art and in the Examples herein.

[0149] It will also be appreciated that the compounds and pharmaceutically acceptable compositions of the present invention can be employed in combination therapies, that is, the compounds and pharmaceutically acceptable compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another agent used to treat the same disorder), or they may achieve different effects (e.g., control of any adverse effects). As used herein, additional therapeutic agents that are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

[0150] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

[0151] The compounds of this invention or pharmaceutically acceptable compositions thereof may also be incorporated into compositions for coating an implantable medical device, such as prostheses, artificial valves, vascular grafts, stents and catheters. Accordingly, the present invention, in another aspect, includes a composition for coating an implantable device comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. In still another aspect, the present invention includes an implantable device coated with a composition comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. Suitable coatings and the general preparation of coated implantable devices are described in US Patents 6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethylsiloxane, polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings

may optionally be further covered by a suitable topcoat of fluorosilicone, polysaccharides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition.

[0152] Another aspect of the invention relates to modulating ABC transporter activity in a biological sample or a patient (e.g., *in vitro* or *in vivo*), which method comprises administering to the patient, or contacting said biological sample with a compound of formula I or a composition comprising said compound. The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

[0153] Modulation of ABC transporter activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, the study of ABC transporters in biological and pathological phenomena; and the comparative evaluation of new modulators of ABC transporters.

[0154] In yet another embodiment, a method of modulating activity of an anion channel *in vitro* or *in vivo*, is provided comprising the step of contacting said channel with a compound of formula (I). In preferred embodiments, the anion channel is a chloride channel or a bicarbonate channel. In other preferred embodiments, the anion channel is a chloride channel.

[0155] According to an alternative embodiment, the present invention provides a method of increasing the number of functional ABC transporters in a membrane of a cell, comprising the step of contacting said cell with a compound of formula (I). The term "functional ABC transporter" as used herein means an ABC transporter that is capable of transport activity. In preferred embodiments, said functional ABC transporter is CFTR.

[0156] According to another preferred embodiment, the activity of the ABC transporter is measured by measuring the transmembrane voltage potential. Means for measuring the voltage potential across a membrane in the biological sample may employ any of the known methods in the art, such as optical membrane potential assay or other electrophysiological methods.

[0157] The optical membrane potential assay utilizes voltage-sensitive FRET sensors described by Gonzalez and Tsien (See, Gonzalez, J. E. and R. Y. Tsien (1995) "Voltage sensing by fluorescence resonance energy transfer in single cells" *Biophys J* 69(4): 1272-80, and Gonzalez, J. E. and R. Y. Tsien (1997) "Improved indicators of cell membrane potential that use fluorescence resonance energy transfer" *Chem Biol* 4(4): 269-77) in combination with instrumentation for measuring fluorescence changes such as the Voltage/Ion Probe Reader (VIPR) (See, Gonzalez, J. E., K. Oades, et al. (1999) "Cell-based assays and instrumentation for screening ion-channel targets" *Drug Discov Today* 4(9): 431-439).

[0158] These voltage sensitive assays are based on the change in fluorescence resonant energy transfer (FRET) between the membrane-soluble, voltage-sensitive dye, DiSBAC₂(3), and a fluorescent phospholipid, CC2-DMPE, which is attached to the outer leaflet of the plasma membrane and acts as a FRET donor. Changes in membrane potential (V_m) cause the negatively charged DiSBAC₂(3) to redistribute across the plasma membrane and the amount of energy transfer from CC2-DMPE changes accordingly. The changes in fluorescence emission can be monitored using VIPR™ II, which is an integrated liquid handler and fluorescent detector designed to conduct cell-based screens in 96- or 384-well microtiter plates.

[0159] In another aspect the present invention provides a kit for use in measuring the activity of a ABC transporter or a fragment thereof in a biological sample *in vitro* or *in vivo* comprising

- (i) a composition comprising a compound of formula (I); and (ii) instructions for a) contacting the composition with the biological sample and b) measuring activity of said ABC transporter or a fragment thereof. In one embodiment, the kit further comprises instructions for a) contacting an additional composition with the biological sample; b) measuring the activity of said ABC transporter or a fragment thereof in the presence of said additional compound, and c) comparing the activity of the ABC transporter in the presence of the additional compound with the density of the ABC transporter in the presence of a composition of formula (I). In preferred embodiments, the kit is used to measure the density of CFTR.

[0160] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

EXAMPLES

[0161] A): Preparation of certain exemplary compounds of the invention:

[0162] General Experimental Procedures:

[0163] Amides Preparation: If the appropriate acid chloride was commercially available, it was added to one equivalent of the appropriate secondary amine in minimum of 1,4-dioxane containing two equivalents triethylamine. The reaction mixture was allowed to stir overnight at room temperature. The reaction mixture was then filtered and evaporated to dryness. The crude product was purified by reverse phase preparative mass-directed liquid chromatography / mass

spectrometry.

[0164] If the appropriate acid chloride was not commercially available the appropriate carboxylic acid was added to a solution containing one equivalent of the appropriate amine in a minimum of acetonitrile containing two equivalents of triethylamine. *O*-(7-Azabenzotriazol-1-yl)-*N,N,N,N*-tetramethyluronium hexafluorophosphate (HATU, 1.2 eq.) is added, and the reaction is stirred overnight. The crude product was then purified by reverse phase preparative mass-directed liquid chromatography / mass spectrometry.

[0165] Specific Examples:

[0166] General. All reagents and solvents were used as received without further purification. Thin layer chromatography was performed on glass-backed silica gel 60 plates pre-coated with a fluorescent dye from EM Science. Mass spectrometry was performed in the positive mode on a PE SCIEX EX150 mass spectrometer. Purity was determined by the observed total ion current, and the ultraviolet absorption at 220 nm and 254 nm

[0167] Preparation of Amines:

[0168] C-[1-(3,4-Dimethoxy-phenyl)-cyclopentyl]-methylamine: (3,4-Dimethoxyphenyl)-acetonitrile (5.00 g, 28.2 mmol) was dissolved in 60 mL of anhydrous tetrahydrofuran in a 250 mL round bottom flask. Sodium hydride (2.03 g, 84.6 mmol) was slowly added and the reaction mixture was warmed to 50-60°C. 1,4-Dichlorobutane (4.30 g, 33.9 mmol) was then added and the reaction mixture was heated to reflux for 16 hours. An additional aliquot of 1,4-dichlorobutane (4.30 g, 33.9 mmol) was added and the reaction mixture was refluxed for an additional 24 hours. The reaction mixture was cooled to room temperature and quenched with the slow addition of methanol. The reaction mixture was evaporated to dryness and purified by column chromatography on silica gel to yield a pale yellow oil (1.61 g, 6.98 mmol, 24.8 %). The resulting 1-(3,4-dimethoxy-phenyl)-cyclopentanecarbonitrile (363 mg, 1.57 mmol) was dissolved in dry ether (4 mL) and cooled to 0°C under an atmosphere of nitrogen. Lithium aluminum hydride (1.57 mL, 1M in ether) was slowly added and the reaction mixture was allowed to warm to room temperature and stirred for 16 hours. The reaction mixture was quenched with the slow addition of methanol. The reaction mixture was washed with a saturated aqueous sodium chloride solution, separated, and evaporated to dryness to give a colorless oil (356 mg, 1.38 mmol, 87.9 %). ESI-MS *m/z* calc. 235.3, found 236.2 (M+1)⁺. Retention time of 1.64 minutes.

[0169] Preparation of Amides:

[0170] 2-(4-Methoxy-benzyl)-thiazole-4-carboxylic acid [1-(3,4-dimethoxy-phenyl)-cyclopentylmethyl]-amide: 2-(4-Methoxy-benzyl)-thiazole-4-carboxylic acid (101 mg, 0.405 mmol) and C-[1-(3,4-Dimethoxy-phenyl)-cyclopentyl]-methylamine (96.5 mg, 0.410 mmol) were dissolved in acetonitrile (2 mL) containing triethylamine (84.1 μL, 0.600 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N,N*-tetramethyluronium hexafluorophosphate (171 mg, 0.450 mmol) was added and the solution was allowed to stir for 16 hours. The reaction mixture was evaporated to dryness and purified by column chromatography on silica gel using a gradient of 5-30 % ethyl acetate in hexanes. The pure fractions were combined and evaporated to dryness to yield a yellow solid (84.0 mg, 0.180 mmol, 43.9 %). ESI-MS *m/z* calc. 466.6, found 467.2 (M+1)⁺. Retention time of 8.32 minutes. ¹H NMR (400 MHz, CD₃CN) δ 1.68 - 2.07 (m, 8H), 3.50 (d, *J* = 6.3 Hz, 2H), 3.82 (s, 9H), 4.20 (s, 2H), 6.83 - 7.29 (m, 8H), 7.87 (s, 1H).

[0171] 2-Benzyl-thiazole-4-carboxylic acid [2-(3,4-dimethoxy-phenyl)-ethyl]-amide. Rink amide resin (0.627 g, 0.752 mmol, 1.20 mmol/g) was suspended and allowed to swell for 10 minutes in 4 mL of *N,N*-dimethylformamide (DMF). Phenylacetic acid (0.15 g, 1.1 mmol) and triethylamine (0.21 mL, 1.5 mmol) were then added to the reaction mixture. *O*-(7-Azabenzotriazol-1-yl)-*N,N,N,N*-tetramethyluronium hexafluorophosphate (HATU, 0.46 g, 1.2 mmol) was added and the reaction mixture was swirled for two hours, filtered, and washed with DMF and dichloromethane. The resin was then suspended in 5 mL of toluene and 2,4-bis-(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (Lawesson's Reagent, 0.93 g, 2.3 mmol) was added to the suspension. The reaction mixture was then swirled for 2 hours at 65°C, filtered, and washed with DMF and dichloromethane. The resin (0.50 g, 0.60 mmol) was then allowed to swell in 5 mL of tetrahydrofuran for 10 minutes and bromopyruvic acid (0.060 g, 0.36 mmol) was added to the reaction mixture. The mixture was then subjected to microwave irradiation for 20 minutes at 135°C, and followed by filtration to yield crude 2-benzyl-thiazole-4-carboxylic acid. MS *m/z* calc. 219.0, found (ESI); 220.2 (M+H⁺). Retention time 2.38 minutes. The crude acid was dissolved in 2 mL of acetonitrile containing triethylamine (0.0836 mL, 0.600 mmol) and 2-(3,4-dimethoxy-phenyl)-ethylamine (0.0332 mL, 0.200 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N,N*-tetramethyluronium hexafluorophosphate (0.0836 g, 0.220 mmol) was added and the solution was allowed to stir for 16 hours. The reaction mixture was then purified by reverse-phase preparative liquid chromatography to yield (0.011 g, 0.029 mmol, 4.8 %) a colorless oil. MS *m/z* calc. 382.1, found (ESI); 383.2 (M+H⁺) Retention time 2.97 minutes. ¹H NMR (400 MHz, CD₃CN) δ 2.85 (t, *J* = 7.0 Hz, 2H), 3.53-3.65 (m, 2H), 3.78 (s, 3H), 3.80 (s, 3H), 4.33 (s, 2H), 6.78-6.91 (m, 3H), 7.29-7.42 (m, 5H), 7.52 (s, 1H), 7.92 (s, 1H)

[0172] 2-(4-Methoxy-benzyl)-oxazole-4-carboxylic acid [1-(3,4-dimethoxy-phenyl)-cyclopentylmethyl]-amide:

[0173] 3-Hydroxy-2-[2-(4-methoxy-phenyl)-acetylaminol-propionic acid methyl ester: to a suspension of DL-serine HCl (2.0 g, 12.8 mmol) in dichloromethane (10 mL) at 0°C were added dropwise triethylamine (3.58 mL, 25.7 mmol) and (4-methoxyphenyl)-acetyl chloride (1.96 mL, 12.8 mmol). The mixture was stirred at 25°C for 12 hours. The reaction mixture was washed with a saturated aqueous sodium chloride solution, separated, and evaporated to dryness and purified by

column chromatography (hexanes: ethyl acetate from 25 to 100%). 3-Hydroxy-2-[2-(4-methoxy-phenyl)-acetylami-
no]-propionic acid methyl ester was isolated as a white solid (2g, 58 %). ESI-MS m/z calc. 267.3, found 268.2 (M+1)⁺.
Retention time of 1.83 minutes. ¹H NMR (400 MHz, CDCl₃) δ 3.64 (s, 2H), 3.78 (s, 3H), 3.83 (s, 3H), 3.93 (qd, J = 14.6
and 4.0 Hz, 2H), 4.66 (m, 1H), 6.40 (m, 1H), 7.01 (d, J = 8.7 Hz, 2H), 7.23 (d, J = 8.7 Hz, 2H).

[0174] 2-(4-Methoxy-benzyl)-oxazole-4-carboxylic acid methyl ester: DAST (diethylaminosulfur trifluoride) was
added to a cold (-78°C) solution of 3-hydroxy-2-[2-(4-methoxy-phenyl)-acetylaminol-propanoic acid methyl ester (400
mg, 1.5 mmol) in dichloromethane (8 mL). After stirring for 1h at -78°C, CCl₃Br (149 μ L, 1.5 mmol) and DBU (226 μ L,
1.5 mmol) were added and the reaction was allowed to warm to 25°C and stirred at this temperature for 12h. The reaction
was poured into saturated aqueous NaHCO₃ and the biphasic mixture was extracted with DCM. The combined organic
extracts were dried with MgSO₄ and purified by column chromatography (hexanes: ethyl acetate from 25 to 100%). 2-(4-
Methoxybenzyl)-oxazole-4-carboxylic acid methyl ester was isolated as a white solid (80 mg, 21%). ESI-MS m/z calc.
247.5, found 248.2 (M+1)⁺. Retention time of 2.64 minutes. ¹H NMR (400 MHz, CDCl₃) δ 3.80 (s, 3H), 3.92 (s, 3H), 4.12
(s, 2H), 6.87 (d, J = 8.0 Hz, 2H), 7.24 (d, J = 8.0 Hz, 2H).

[0175] 2-(4-Methoxy-benzyl)-oxazole-4-carboxylic acid: A mixture of 2-(4-Methoxybenzyl)-oxazole-4-carboxylic
acid methyl ester (40 mg, 0.16 mmol), methanol (2 mL) and 1N NaOH (1 mL) was heated at 60°C for 2 hours and allowed
to stand at 25°C for 30 minutes. After dilution with water, the mixture was adjusted to pH 2 with 2N HCl and extracted
with ether. The organic layers were combined, dried with MgSO₄ and evaporated to dryness to yield 25 mg (66%) of
2-(4-Methoxy-benzyl)-oxazole-4-carboxylic acid as a white solid. ESI-MS m/z calc. 233.2, found 234.2 (M+1)⁺. Retention
time of 2.27 minutes.

[0176] 2-(4-Methoxy-benzyl)-oxazole-4-carboxylic acid [1-(3,4-dimethoxy-phenyl)-cyclopentylmethyl]-amide:
Starting from 2-(4-Methoxy-benzyl)-oxazole-4-carboxylic acid (25g, 0.1 mmol) and [2-(3,4-Dimethoxy-phenyl)-2-
methyl]-propylamine (47 mg, 0.2 mmol) following a similar procedure reported for the preparation of 2-(4-Methoxy-
benzyl)-thiazole-4-carboxylic acid[1-(3,4-dimethoxy-phenyl)-cyclopentylmethyl]-amide, the amide (30 mg, 66%)
was obtained as a white solid. ESI-MS m/z calc. 450.5, found 451.2(M+1)⁺. Retention time of 3.51 minutes. ¹H NMR
(400 MHz, CDCl₃) δ 1.68 - 2.19 (m, 8H), 3.50 (d, J = 6.3 Hz, 2H), 3.81 (s, 3H), 3.88 (s, 3H), 3.92 (s, 3H), 4.00 (s, 2H),
6.73 - 6.88 (m, 6H), 7.20 (d, J = 8.7 Hz, 2H), 8.07 (s, 1H).

[0177] Other compounds of formula I have been prepared by methods substantially similar to those described above.
Table 2 below recites analytical data for selected compounds of formula I.

Table 2

Cmpd #	LC-MS (M+1) ⁺	LC-RT (min)
6	451.20	3.60
10	468.00	3.17
12	502.00	3.25
13	473.00	4.84
14	457.00	4.66
15	454.00	4.19
16	499.00	4.25
17	468.00	3.17
18	468.00	3.23
19	513.20	3.16
20	422.00	4.32

Cmpd #	LC-MS (M+1) ⁺	LC-RT (min)
21	377.00	4.46
22	512.00	4.57
23	501.00	4.96
24	560.20	4.33
25	323.00	4.14
26	368.00	4.12
27	354.00	3.18
28	357.00	3.90
29	343.00	3.40
30	479.00	4.28
31	465.40	3.52

Cmpd #	LC-MS (M+1) ⁺	LC-RT (min)
32	505.40	4.24
33	489.20	4.46
34	413.20	3.62
35	427.00	3.45
36	467.20	4.23
37	451.20	4.42
38	417.20	3.87
39	431.40	3.73
40	471.20	4.46
41	455.20	4.68
42	437.20	3.24

Cmpd #	LC-MS (M+1)	LC-RT (min)
43	423.20	3.40
44	361.00	2.91
45	403.40	3.25
46	417.20	3.17
47	343.00	3.53
48	457.40	3.84
49	424.20	2.53
50	423.20	3.55
51	407.60	3.71
52	437.40	3.78
53	451.40	3.64
54	481.20	3.65
55	471.20	3.67
56	501.20	3.67
57	497.20	3.49
58	437.20	3.51
59	467.40	3.63
60	467.40	3.54
61	497.40	3.33
62	497.40	3.55

Cmpd #	LC-MS (M+1) ⁺	LC-RT (min)
63	497.20	3.58
64	451.20	3.77
65	451.40	3.76
66	471.20	3.72
67	505.40	3.78
68	505.40	3.80
69	505.20	3.80
70	527.20	3.35
71	481.40	3.50
72	477.20	3.48
73	526.00	3.37
74	530.20	3.62
75	439.40	3.17
76	483.40	2.90
77	441.20	3.25
78	488.20	3.03
79	492.20	3.27
80	538.20	3.32
81	451.20	3.88
82	443.20	3.51

Cmpd #	LC-MS (M+1) ⁺	LC-RT (min)
83	443.40	3.51
84	481.40	3.74
85	475.20	4.09
86	392.00	2.28
87	397.00	3.60
88	391.00	3.70
89	405.20	3.84
90	421.00	3.75
91	427.20	3.85
92	405.20	3.87
93	421.00	3.73
94	425.00	3.89
95	405.40	3.87
96	421.20	3.63
97	425.20	3.89
98	451.00	3.81
99	483.40	4.12
100	481.20	4.19

[0178] B) Assays for Detecting and Measuring Δ F508-CFTR Correction Properties of Compounds

[0179] I) Membrane potential optical methods for assaying Δ F508-CFTR modulation properties of compounds

[0180] The optical membrane potential assay utilized voltage-sensitive FRET sensors described by Gonzalez and Tsien (See, Gonzalez, J. E. and R. Y. Tsien (1995) "Voltage sensing by fluorescence resonance energy transfer in single cells" Biophys J 69(4): 1272-80, and Gonzalez, J. E. and R. Y. Tsien (1997) "Improved indicators of cell membrane potential that use fluorescence resonance energy transfer" Chem Biol 4(4): 269-77) in combination with instrumentation for measuring fluorescence changes such as the Voltage/Ion Probe Reader (VIPR) (See, Gonzalez, J. E., K. Oades, et al. (1999) "Cell-based assays and instrumentation for screening ion-channel targets" Drug Discov Today 4(9): 431-439).

[0181] These voltage sensitive assays are based on the change in fluorescence resonant energy transfer (FRET) between the membrane-soluble, voltage-sensitive dye, DiSBAC₂(3), and a fluorescent phospholipid, CC2-DMPE, which is attached to the outer leaflet of the plasma membrane and acts as a FRET donor. Changes in membrane potential (V_m) cause the negatively charged DiSBAC₂(3) to redistribute across the plasma membrane and the amount of energy transfer from CC2-DMPE changes accordingly. The changes in fluorescence emission were monitored using VIPR™ II, which is an integrated liquid handler and fluorescent detector designed to conduct cell-based screens in 96- or 384-well microtiter plates.

[0182] Identification of Correction Compounds

[0183] To identify small molecules that correct the trafficking defect associated with Δ F508-CFTR; a single-addition HTS assay format was developed. The cells were incubated in serum-free medium for 16 hrs at 37 °C in the presence or absence (negative control) of test compound. As a positive control, cells plated in 384-well plates were incubated for 16 hrs at 27 °C to "temperature-correct" Δ F508-CFTR. The cells were subsequently rinsed 3X with Krebs Ringers solution and loaded with the voltage-sensitive dyes. To activate Δ F508-CFTR, 10 μ M forskolin and the CFTR potentiator, genistein (20 μ M), were added along with Cl⁻-free medium to each well. The addition of Cl⁻-free medium promoted Cl⁻ efflux in response to Δ F508-CFTR activation and the resulting membrane depolarization was optically monitored using the FRET-based voltage-sensor dyes.

[0184] Identification of Potentiator Compounds

[0185] To identify potentiators of Δ F508-CFTR, a double-addition HTS assay format was developed. During the first addition, a Cl⁻-free medium with or without test compound was added to each well. After 22 sec, a second addition of Cl⁻-free medium containing 2 - 10 μ M forskolin was added to activate Δ F508-CFTR. The extracellular Cl⁻ concentration following both additions was 28 mM, which promoted Cl⁻ efflux in response to Δ F508-CFTR activation and the resulting membrane depolarization was optically monitored using the FRET-based voltage-sensor dyes.

Solutions

Bath Solution #1: (in mM)

NaCl 160, KCl 4.5, CaCl₂ 2, MgCl₂ 1, HEPES 10, pH 7.4 with NaOH.

EP 2 140 865 A1

(continued)

Chloride-free bath solution: salts. Chloride salts in Bath Solution #1 are substituted with gluconate
CC2-DMPE: Prepared as a 10 mM stock solution in DMSO and stored at -20°C.
DiSBAC₂(3): Prepared as a 10 mM stock in DMSO and stored at -20°C.

[0186] Cell Culture

[0187] NIH3T3 mouse fibroblasts stably expressing $\Delta F508$ -CFTR are used for optical measurements of membrane potential. The cells are maintained at 37 °C in 5% CO₂ and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β -ME, 1 X pen/strep, and 25 mM HEPES in 175 cm² culture flasks. For all optical assays, the cells were seeded at 30,000/well in 384-well matrigel-coated plates and cultured for 2 hrs at 37 °C before culturing at 27 °C for 24 hrs. for the potentiator assay. For the correction assays, the cells are cultured at 27 °C or 37 °C with and without compounds for 16 - 24 hoursB) Electrophysiological Assays for assaying $\Delta F508$ -CFTR modulation properties of compounds

[0188] 1.Ussing Chamber Assay

[0189] Ussing chamber experiments were performed on polarized epithelial cells expressing $\Delta F508$ -CFTR to further characterize the $\Delta F508$ -CFTR modulators identified in the optical assays. FRT $\Delta F508$ -CFTR epithelial cells grown on Costar Snapwell cell culture inserts were mounted in an Ussing chamber (Physiologic Instruments, Inc., San Diego, CA), and the monolayers were continuously short-circuited using a Voltage-clamp System (Department of Bioengineering, University of Iowa, IA, and, Physiologic Instruments, Inc., San Diego, CA). Transepithelial resistance was measured by applying a 2-mV pulse. Under these conditions, the FRT epithelia demonstrated resistances of 4 K Ω /cm² or more. The solutions were maintained at 27 °C and bubbled with air. The electrode offset potential and fluid resistance were corrected using a cell-free insert. Under these conditions, the current reflects the flow of Cl⁻ through $\Delta F508$ -CFTR expressed in the apical membrane. The I_{SC} was digitally acquired using an MP100A-CE interface and AcqKnowledge software (v3.2.6; BIOPAC Systems, Santa Barbara, CA).

[0190] Identification of Correction Compounds

[0191] Typical protocol utilized a basolateral to apical membrane Cl⁻ concentration gradient. To set up this gradient, normal ringer was used on the basolateral membrane, whereas apical NaCl was replaced by equimolar sodium gluconate (titrated to pH 7.4 with NaOH) to give a large Cl⁻ concentration gradient across the epithelium. All experiments were performed with intact monolayers. To fully activate $\Delta F508$ -CFTR, forskolin (10 μ M) and the PDE inhibitor, IBMX (100 μ M), were applied followed by the addition of the CFTR potentiator, genistein (50 μ M).

[0192] As observed in other cell types, incubation at low temperatures of FRT cells stably expressing $\Delta F508$ -CFTR increases the functional density of CFTR in the plasma membrane. To determine the activity of correction compounds, the cells were incubated with 10 μ M of the test compound for 24 hours at 37°C and were subsequently washed 3X prior to recording. The cAMP - and genistein-mediated I_{SC} in compound-treated cells was normalized to the 27°C and 37°C controls and expressed as percentage activity. Preincubation of the cells with the correction compound significantly increased the cAMP- and genistein-mediated I_{SC} compared to the 37°C controls.

[0193] Identification of Potentiator Compounds

[0194] Typical protocol utilized a basolateral to apical membrane Cl⁻ concentration gradient. To set up this gradient, normal ringers was used on the basolateral membrane and was permeabilized with nystatin (360 μ g/ml), whereas apical NaCl was replaced by equimolar sodium gluconate (titrated to pH 7.4 with NaOH) to give a large Cl⁻ concentration gradient across the epithelium. All experiments were performed 30 min after nystatin permeabilization. Forskolin (10 μ M) and all test compounds were added to both sides of the cell culture inserts. The efficacy of the putative $\Delta F508$ -CFTR potentiators was compared to that of the known potentiator, genistein.

[0195]

Solutions

Basolateral solution (in mM): NaCl (135), CaCl₂ (1.2), MgCl₂ (1.2), K₂HPO₄ (2.4), KHPO₄ (0.6), N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) (10), and dextrose (10).
The solution was titrated to pH 7.4 with NaOH.

Apical solution (in mM): Same as basolateral solution with NaCl replaced with Na Gluconate (135).

[0196] Cell Culture

[0197] Fisher rat epithelial (FRT) cells expressing $\Delta F508$ -CFTR (FRT $\Delta F508$ -CFTR) were used for Ussing chamber experiments for the putative $\Delta F508$ -CFTR modulators identified from our optical assays. The cells were cultured on Costar Snapwell cell culture inserts and cultured for five days at 37 °C and 5% CO₂ in Coon's modified Ham's F-12 medium supplemented with 5% fetal calf serum, 100 U/ml penicillin, and 100 μ g/ml streptomycin. Prior to use for characterizing

the potentiator activity of compounds, the cells were incubated at 27 °C for 16 - 48 hrs to correct for the $\Delta F508$ -CFTR. To determine the activity of correction compounds, the cells were incubated at 27 °C or 37 °C with and without the compounds for 24 hours.

[0198] 2. Whole-cell recordings

[0199] The macroscopic $\Delta F508$ -CFTR current ($I_{\Delta F508}$) in temperature- and test compound-corrected NIH3T3 cells stably expressing $\Delta F508$ -CFTR were monitored using the perforated-patch, whole-cell recording. Briefly, voltage-clamp recordings of $I_{\Delta F508}$ were performed at room temperature using an Axopatch 200B patch-clamp amplifier (Axon Instruments Inc., Foster City, CA). All recordings were acquired at a sampling frequency of 10 kHz and low-pass filtered at 1 kHz. Pipettes had a resistance of 5 - 6 M Ω when filled with the intracellular solution. Under these recording conditions, the calculated reversal potential for Cl⁻ (E_{Cl}) at room temperature was -28 mV. All recordings had a seal resistance > 20 G Ω and a series resistance < 15 M Ω . Pulse generation, data acquisition, and analysis were performed using a PC equipped with a Digidata 1320 A/D interface in conjunction with Clampex 8 (Axon Instruments Inc.). The bath contained < 250 μ l of saline and was continuously perfused at a rate of 2 ml/min using a gravity-driven perfusion system.

[0200] Identification of Correction Compounds

[0201] To determine the activity of correction compounds for increasing the density of functional $\Delta F508$ -CFTR in the plasma membrane, we used the above-described perforated-patch-recording techniques to measure the current density following 24-hr treatment with the correction compounds. To fully activate $\Delta F508$ -CFTR, 10 μ M forskolin and 20 μ M genistein were added to the cells. Under our recording conditions, the current density following 24-hr incubation at 27 °C was higher than that observed following 24-hr incubation at 37 °C. These results are consistent with the known effects of low-temperature incubation on the density of $\Delta F508$ -CFTR in the plasma membrane. To determine the effects of correction compounds on CFTR current density, the cells were incubated with 10 μ M of the test compound for 24 hours at 37 °C and the current density was compared to the 27 °C and 37 °C controls (% activity). Prior to recording, the cells were washed 3X with extracellular recording medium to remove any remaining test compound. Preincubation with 10 μ M of correction compounds significantly increased the cAMP- and genistein-dependent current compared to the 37 °C controls.

[0202] Identification of Potentiator Compounds

[0203] The ability of $\Delta F508$ -CFTR potentiators to increase the macroscopic $\Delta F508$ -CFTR Cl⁻ current ($I_{\Delta F508}$) in NIH3T3 cells stably expressing $\Delta F508$ -CFTR was also investigated using perforated-patch-recording techniques. The potentiators identified from the optical assays evoked a dose-dependent increase in $I_{\Delta F508}$ with similar potency and efficacy observed in the optical assays. In all cells examined, the reversal potential before and during potentiator application was around -30 mV, which is the calculated E_{Cl} (-28 mV).

[0204]

Solutions

Intracellular solution (in mM): Cs-aspartate (90), CsCl (50), MgCl₂ (1), HEPES (10), and 240 μ g/ml amphotericin-B (pH adjusted to 7.35 with CsOH).
Extracellular solution (in mM): N-methyl-D-glucamine (NMDG)-Cl (150), MgCl₂ (2), CaCl₂ (2), HEPES (10) (pH adjusted to 7.35 with HCl).

[0205] Cell Culture

[0206] NIH3T3 mouse fibroblasts stably expressing $\Delta F508$ -CFTR are used for whole-cell recordings. The cells are maintained at 37 °C in 5% CO₂ and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β -ME, 1 X pen/strep, and 25 mM HEPES in 175 cm² culture flasks. For whole-cell recordings, 2,500 - 5,000 cells were seeded on poly-L-lysine-coated glass coverslips and cultured for 24 - 48 hrs at 27 °C before use to test the activity of potentiators; and incubated with or without the correction compound at 37 °C for measuring the activity of correctors.

[0207] 3. Single-channel recordings

[0208] The single-channel activities of temperature-corrected $\Delta F508$ -CFTR stably expressed in NIH3T3 cells and activities of potentiator compounds were observed using excised inside-out membrane patch. Briefly, voltage-clamp recordings of single-channel activity were performed at room temperature with an Axopatch 200B patch-clamp amplifier (Axon Instruments Inc.). All recordings were acquired at a sampling frequency of 10 kHz and low-pass filtered at 400 Hz. Patch pipettes were fabricated from Coming Kovar Sealing #7052 glass (World Precision Instruments, Inc., Sarasota, FL) and had a resistance of 5 - 8 M Ω when filled with the extracellular solution. The $\Delta F508$ -CFTR was activated after excision, by adding 1 mM Mg-ATP, and 75 nM of the cAMP-dependent protein kinase, catalytic subunit (PKA; Promega Corp. Madison, WI). After channel activity stabilized, the patch was perfused using a gravity-driven microperfusion system. The inflow was placed adjacent to the patch, resulting in complete solution exchange within 1 - 2 sec. To maintain $\Delta F508$ -CFTR activity during the rapid perfusion, the nonspecific phosphatase inhibitor F⁻ (10 mM NaF) was added to

the bath solution. Under these recording conditions, channel activity remained constant throughout the duration of the patch recording (up to 60 min). Currents produced by positive charge moving from the intra-to extracellular solutions (anions moving in the opposite direction) are shown as positive currents. The pipette potential (V_p) was maintained at 80 mV.

[0209] Channel activity was analyzed from membrane patches containing ≤ 2 active channels. The maximum number of simultaneous openings determined the number of active channels during the course of an experiment. To determine the single-channel current amplitude, the data recorded from 120 sec of $\Delta F508$ -CFTR activity was filtered "off-line" at 100 Hz and then used to construct all-point amplitude histograms that were fitted with multigaussian functions using Bio-Patch Analysis software (Bio-Logic Comp. France). The total microscopic current and open probability (P_o) were determined from 120 sec of channel activity. The P_o was determined using the Bio-Patch software or from the relationship $P_o = I/i(N)$, where I = mean current, i = single-channel current amplitude, and N = number of active channels in patch.

[0210]

Solutions

Extracellular solution (in mM): NMDG (150), aspartic acid (150), CaCl_2 (5), MgCl_2 (2), and HEPES (10) (pH adjusted to 7.35 with Tris base).

Intracellular solution (in mM): NMDG-Cl (150), MgCl_2 (2), EGTA (5), TES (10), and Tris base (24) (pH adjusted to 7.35 with HCl).


[0211] Cell Culture

[0212] NIH3T3 mouse fibroblasts stably expressing $\Delta F508$ -CFTR are used for excised-membrane patch-clamp recordings. The cells are maintained at 37 °C in 5% CO_2 and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β -ME, 1 X pen/strep, and 25 mM HEPES in 175 cm^2 culture flasks. For single channel recordings, 2,500 - 5,000 cells were seeded on poly-L-lysine-coated glass coverslips and cultured for 24 - 48 hrs at 27 °C before use.

[0213] Compounds of the invention are useful as modulators of ATP binding cassette transporters, particularly CFTR. In some embodiments, certain compounds depicted in Table 1 exhibit a relative modulating efficacy of greater than 30%:

Claims

1. A compound of formula I:



or a pharmaceutically acceptable salt thereof,
wherein X is O or S;

R^1 is hydrogen, or is a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R^1 is optionally substituted at one or more carbon or nitrogen atoms with x independent occurrences of

- QR^x ; wherein x is 0-5; wherein Q is a bond or is a C_1 - C_6 alkylidene chain wherein up to two methylene units of Q are optionally and independently replaced by $-\text{CO}-$, $-\text{CO}_2-$, $-\text{COCO}-$, $-\text{CONR}-$, $-\text{OCONR}-$, $-\text{NRNR}-$, $-\text{NRN}-$, $-\text{RCO}-$, $-\text{NRCO}-$, $-\text{NRCO}_2-$, $-\text{NRCONR}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NRSO}_2-$, $-\text{SO}_2\text{NR}-$, $-\text{NRSO}_2\text{NR}-$, $-\text{O}-$, $-\text{S}-$; or $-\text{NR}-$; and each occurrence of R^x is independently R' , halogen, NO_2 , or CN , or $-\text{QR}^x$ is $=\text{O}$, $=\text{S}$, or $=\text{NR}'$;
 Z is a bond or is $-\text{CH}_2-$, provided that when Z is a bond, R^1 is not hydrogen;
each occurrence of R is independently hydrogen or an optionally substituted C_1 - C_6 aliphatic group; and each

occurrence of R' is independently hydrogen or an optionally substituted C₁-C₆ aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or R and R', two occurrences of R, or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; R² is halogen, -CN, -CF₃, -NO₂, or -TqR; R³ is U_mR' and R⁴ is V_pCy¹,


wherein m, p, and q are each independently 0 or 1, and U, V, and T are each independently an optionally substituted C₁-C₄alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by -CO-, -CO₂-, -COCO-, -CONR-,

- OCONR-, -NRNR-, -NRNRCO-, -NRCO-, -NRCO₂-, -NRCONR-, -SO-, -SO₂-, -NRSO₂-, -SO₂NR-, -NRSO₂NR-, -O-, -S-, or -NR-;

Cy¹ is a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein Cy¹ is optionally substituted at one or more carbon or nitrogen atoms with y independent occurrences of WR^W;


wherein y is 0-5; wherein W is a bond or is a C₁-C₆ alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by -CO-, -CO₂-, -COCO-, -CONR-, -OCONR-, -NRNR-, -NRNRCO-, -NRCO-, -NRCO₂-, -NRCONR-, -SO-, -SO₂-, -NRSO₂-, -SO₂NR-, -NRSO₂NR-, -O-, -S-, or -NR-; and each occurrence of R^W is independently R', halogen, NO₂, or CN, or -WR^W is =O, =S, or =NR'; or

R³ and R⁴, taken together with the nitrogen atom to which they are bound form an optionally substituted 5-, 6- or 7-membered saturated, partially unsaturated or fully unsaturated ring having the structure:



wherein r is 0, 1, or 2; one of X³, X⁴, or X⁵ is CH-V_p-Cy¹ or N-V_p-Cy¹, and the others of X³, X⁴, or X⁵ are CHR' or NR'; and each occurrence of X¹, when present, and X² are each independently C(R')₂, -CO-, or -CS-; provided that:


a) when X is S, Z is CH₂, and R² is hydrogen, then R¹ is not:



b) when X is S or O; and R² is formyl, 4,5-dihydro-2-oxazolyl, -COOR', or -COSR'; then:


i) when Z is -CH₂-, R¹ is not phenyl optionally substituted with C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄haloalkylthio, halogen, cyano, or nitro; or ii) when Z is C₁₋₆alkyl, then R¹ is not C₃-C₆cycloalkyl optionally substituted with C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄haloalkylthio halogen, cyano, or nitro; c) when X is S, R² is H, Z is -CH₂-, and R¹ is unsubstituted

phenyl, then when R³ is hydrogen, R⁴ is not:




d) when X is S, R² is SMe or Me, Z is CH₂ and R¹ is unsubstituted phenyl,

- i) when R³ is hydrogen, then R⁴ is not unsubstituted benzyl; and
 ii) when R³ is ethyl, then R⁴ is not unsubstituted phenyl; e) when X is O, Z is CH₂, R¹ is unsubstituted phenyl, R² is Me, and R³ is hydrogen, then R⁴ is not




f) when X is O, Z is CH₂, R¹ is unsubstituted phenyl, R² is hydrogen, and R³ is hydrogen, then R⁴ is not unsubstituted phenyl, 4-Me-phenyl, or unsubstituted benzyl; and g) when X is O, Z is -CH(iPr)-, R¹ is




and R³ is hydrogen, then R⁴ is not

-(CH₂)₂-S-Phenyl; h) when Z-R¹ is methyl, X is S, R³ is iBu, then R⁴ is not:




and



i) when Z-R¹ is phenyl, X is S, and R₃ is -(CH₂)₂-pyrid-2-yl, then R⁴ is not



2. The compound of claim 1, wherein R² and R³ are each hydrogen, and R⁴ is V_pCy¹, where p is 1 and V is -CH₂C(R⁶)₂ and compounds have the general formula II:

















**II**


3. The compound of claim 2, wherein X is S or O and compounds of formula II-A or II-B are provided:

**II-A****II-B**


4. The compound of claim 3, wherein one or more of the compound variables are selected from one of more of:

a) R¹ is:


**a****b****c****d****e****f****g****h****i****j****k****l****m****n****o****p**




q




r




s




t




u




v




w




x




y




z




aa




bb




cc




dd




ee




ff




gg




hh




ii




jj




kk




ll






mm



nn



oo

**pp****qq****rr;**

b) R¹ is one of rings **a**, **b**, **c**, **d**, **m**, **n**, **o**, **ee**, or **pp**;

















c) R¹ is phenyl (ring **a**);

d) x is 0, 1, 2 or 3, and Q is a bond or is an optionally substituted C₁-C₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO₂, or CO₂, CO, and R^x is R' or halogen;


e) x is 0, 1, 2, or 3 and each occurrence of QR^x, when present, is independently -C₁₋₃alkyl, -O(C₁₋₃alkyl), -CF₃, -OCF₃, -SCF₃, -F, -Cl, -Br, -COOR', -COR', -O(CH₂)₂N(R)(R'), -O(CH₂)N(R)(R'), -CON(R)(R'), -(CH₂)₂OR', -(CH₂)OR', optionally substituted phenyl, optionally substituted benzyl, -N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), -or SO₂NRR';

f) R⁶ is hydrogen, methyl, or two occurrences of R⁶ on the same carbon atom are taken together to form an optionally substituted 3-6-membered spiro ring having 0, 1 or 2 heteroatoms selected from nitrogen, oxygen, or sulfur;


g) Cy¹ is:

**i****ii****iii****iv****v****vi****vii****viii****ix****x****xi****xii****xiii****xiv****xv****xvi**


5




xvii



xviii




xix



xx


10




xxi



xxii




xxiii




xxiv

15


20



xxv



xxvi




xxvii




xxviii

25


30



xxix




xxx




xxxi

35


40




xxxii



xxxiii




xxxiv




xxxv

45


50



xxxvi




xxxvii




xxxviii


55




xxxix




xL




xLi



xLii



xLiii




xLiv;

h) y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C_1 - C_6 alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^W is R' or halogen; or


i) y is 0, 1, 2, or 3 and each occurrence of WR^W , when present, is independently $-C_1$ - C_3 alkyl, $-O(C_1$ - C_3 alkyl), $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CR_2)_2N(R)(R')$, $-O(CH_2)_2N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, or SO_2NRR' .

5. The compound of claim 1, wherein R^2 is hydrogen; R^3 is U_mR' where m is 0 or 1, and U , when present, is $-CH_2-$ and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and R^4 is $-CH_2Cy^1$ and compounds of formula III are provided:




III.

6. The compound of claim 5, wherein X is S or O and compounds of formula III-A or III-B are provided:




III-A




III-B.

7. The compound of claim 5, herein one or more of the compound variables are selected from one of more of:


a) R^1 is:




a




b




c




d




e




f




g




h




i




j




k




l




m




n




o




p




q




r




s




t




u























v



w



x

**y****z****aa****bb****cc****dd****ee****ff****gg****hh****ii****jj****kk****ll****mm****nn****oo****pp****qq****rr;**

b) R^1 is one of rings **a**, **b**, **c**, **d**, **m**, **n**, **o**, **ee**, or **pp**;

c) R^1 is phenyl (ring **a**);


d) x is 0, 1, 2 or 3, and Q is a bond or is an optionally substituted C_1 - C_6 alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^X is R' or halogen;

e) m is 0 or 1, and U , when present, is $-CH_2-$ and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen,


oxygen, or sulfur;

f) R' is an optionally substituted phenyl or pyridyl group;


g) Cy¹ is:




i




ii




iii




iv




v




vi




vii




viii




ix




x




xi




xii




xiii




xiv




xv




xvi




xvii




xviii



xix




xx




xxi




xxii




xxiii




xxiv



xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi



xLii



xLiii




xLiv;

h) y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C_1 - C_6 alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^W is R' or halogen; or

i) y is 0, 1, 2, or 3 and each occurrence of WR^W , when present, is independently $-C_{1-3}$ alkyl, $-O(C_{1-3}$ alkyl), $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, or SO_2NRR' .

8. The compound of claim 1, wherein, $Z-R^1$ is $-CH_3$; R^2 is hydrogen; and R^4 is $-CH_2Cy^1$ and compounds of formula IV


are provided:




IV

wherein m is 0 or 1, and U, when present, is -CH₂- and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

9. The compound of claim 8, wherein X is S or O and compounds of formula III-A or III-B are provided:




IV-A




IV-B

10. The compound of claim 8, wherein one or more of the compound variables are selected from one of more of:


- m is 0 or 1, and U, when present, is -CH₂- and R' is a 5- or 6-membered optionally substituted saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;
- R' is an optionally substituted phenyl or pyridyl group;
- Cy¹ is:




i




ii




iii




iv




v




vi




vii




viii




ix




x




xi




xii




xiii




xiv




xv




xvi




xvii




xviii




xix




xx




xxi




xxii




xxiii




xxiv




xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi



xLii



xLiii




xLiv;

d) y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C₁-C₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO₂, or CO₂, CO, and R^W is R' or halogen; or


e) y is 0, 1, 2, or 3 and each occurrence of WR^W, when present, is independently -C₁₋₃alkyl, -O(C₁₋₃alkyl), -CF₃, -OCF₃, -SCF₃, -F, -Cl, -Br, -COOR', -COR', -O(CH₂)₂N(R)(R'), -O(CH₂)N(R)(R'), -CON(R)(R'), -(CH₂)₂OR', -(CH₂)OR', optionally substituted phenyl, optionally substituted benzyl, -N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), or -SO₂NRR'.

11. The compound of claim 1, wherein compounds of formula V are provided:




V.

12. The compound of claim 11, wherein X is S or O and compounds of formula V-A or V-B are provided:




V-A




V-B

13. The compound of claim 11, wherein one or more of the compound variables are selected from one of more of:


a) R¹ is:




a




b




c




d




e




f




g




h




i




j




k




l




m




n




o




p




q




r




s




t




u























v



w



x

**y****z****aa****bb****cc****dd****ee****ff****gg****hh****ii****jj****kk****ll****mm****nn****oo****pp****qq****rr;**

b) R^1 is one of rings **a**, **b**, **c**, **d**, **m**, **n**, **o**, **ee**, or **pp**;

c) R^1 is phenyl (ring **a**);


d) x is 0, 1, 2 or 3, and Q is a bond or is an optionally substituted C_1 - C_6 alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^X is R' or halogen;

e) r is 1, X^1 , X^2 , X^3 and X^5 are each CH_2 , and X^4 is $CH-V_p-Cy^1$, or $N-V_p-Cy^1$, p is 1 and V is SO_2 , $-NRSO_2$, CO, or $NRCO$;


f) X^4 is $N-V_p-Cy^1$, p is 1 and V is SO_2 or CO;

g) X^4 is $N-V_p-Cy^1$, p is 1 and V is SO_2 ;


h) Cy^1 is:




i




ii




iii




iv




v




vi




vii




viii




ix




x




xi




xii




xiii




xiv




xv




xvi




xvii




xviii



xix




xx




xxi




xxii




xxiii




xxiv




xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi



xLii



xLiii




xLiv;

i) y is 0, 1, 2 or 3, and W is a bond or is an optionally substituted C_1 - C_6 alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, SO_2 , or CO_2 , CO, and R^W is R' or halogen; or

j) y is 0, 1, 2, or 3 and each occurrence of WR^W , when present, is independently $-C_1$ - C_3 alkyl, $-O(C_1$ - C_3 alkyl), $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-CON(R)(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, optionally substituted benzyl, $-N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, or SO_2NRR' .

14. The compound according to claim 1, having formula VI:




VI;


wherein:

ring A is a 3-7 membered ring having 0-3 heteroatoms independently selected from O, S, or NR.

15. The compound according to claim 14, wherein said compound has formula VI-A or formula VI-B:




VI-A




VI-B.

16. The compound according to claim 15, wherein ring A is a 3-7 membered cycloalkyl ring, in particular ring A is cyclopentyl or cyclohexyl.


17. The compound according to claim 14, wherein R¹ is selected from one of the following groups:




a




b




c




d




e




f




g




h




i




j




k




l




m




n




o




p




q




r




s




t




u




v




w




x




y




z




aa




bb




cc




dd




ee




ff




gg




hh




ii




jj




kk




ll




mm




nn




oo




pp




qq




rr




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi



xLii









xLiii











xLiv.





18. The compound according to claim 1, wherein said compound is selected from Table 1.











Cmpd #	Compound	Cmpd #	Compound
3		5	
4		6	





Cmpd #	Compound	Cmpd #	Compound
7		9	
8		10	





Cmpd #	Compound	Cmpd #	Compound
11		13	
12		14	





Cmpd #	Compound	Cmpd #	Compound
15		17	
16		18	





Cmpd #	Compound	Cmpd #	Compound
19		21	
20		22	





Cmpd #	Compound	Cmpd #	Compound
23		25	
24		26	





Cmpd #	Compound	Cmpd #	Compound
27		29	
28		30	





Cmpd #	Compound	Cmpd #	Compound
31		33	
32		34	





Cmpd #	Compound	Cmpd #	Compound
35		37	
36		38	





Cmpd #	Compound	Cmpd #	Compound
39		41	
40		42	





Cmpd #	Compound	Cmpd #	Compound
43		45	
44		46	





Cmpd #	Compound	Cmpd #	Compound
47		49	
48		50	





Cmpd #	Compound	Cmpd #	Compound
51		53	
52		54	





Cmpd #	Compound	Cmpd #	Compound
55		57	
56		58	





Cmpd #	Compound	Cmpd #	Compound
59		61	
60		62	





Cmpd #	Compound	Cmpd #	Compound
63		65	
64		66	





Cmpd #	Compound	Cmpd #	Compound
67		69	
68		70	





Cmpd #	Compound	Cmpd #	Compound
71		73	
72		74	

Cmpd #	Compound	Cmpd #	Compound
75		77	
76		78	



Cmpd #	Compound	Cmpd #	Compound
79		81	
80		82	

Cmpd #	Compound	Cmpd #	Compound
83		85	
84		86	

Cmpd #	Compound	Cmpd #	Compound
87		89	
88		90	

Cmpd #	Compound	Cmpd #	Compound
91		93	
92		94	

Cmpd #	Compound	Cmpd #	Compound
95		97	
96		98	

Cmpd #	Compound	Cmpd #	Compound
99		100	

19. A pharmaceutical composition comprising:

- (i) a compound according to claim 1; and
- (ii) a pharmaceutically acceptable carrier.



EUROPEAN SEARCH REPORT

Application Number
EP 09 16 4393

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X,P	WO 2004/084842 A2 (IRM LLC, BERMUDA) 7 October 2004 (2004-10-07) * example 190 *	1-19	INV. A61K31/42 A61K31/422 A61K31/426 A61K31/427 C07D277/56 C07D417/12 C07D417/04 C07D417/06 C07D263/34
X,P	DE 102 50 110 A1 (BAYER CROPS SCIENCE A.-G., GERMANY) 13 May 2004 (2004-05-13) * example 7 *	1-19	
X	HAHN, HOH-GYU ET AL: "Synthesis of new trifluoromethylated 1,3-oxazolecarboxanilides and their fungicidal activities" AGRICULTURAL CHEMISTRY AND BIOTECHNOLOGY (ENGLISH EDITION) , 45(1), 37-42 CODEN: ACBTFF; ISSN: 1229-2737, 2002, XP009046397 examples	1-19	
X	EP 0 652 229 A2 (ISAGRO S.P.A., ITALY) 10 May 1995 (1995-05-10) * page 19; examples 86,87 *	1-19	
X	EP 0 419 944 A2 (BASF A.-G., GERMANY) 3 April 1991 (1991-04-03) pages 19-20, compounds for which R4 = isopropyl	1-19	TECHNICAL FIELDS SEARCHED (IPC) A61K C07D
X	GB 1 318 291 A (SHELL INTERNATIONALE RESEARCH MAATSCHAPPIJ N. V.) 23 May 1973 (1973-05-23) page 12, last compound of table 2	1-19	
A	US 6 630 482 B1 (BECQ FREDERIC ET AL) 7 October 2003 (2003-10-07) * page 1, column 1 * * claim 1 *	1-19	
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 6 November 2009	Examiner Fanni, Stefano
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

 2
EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 09 16 4393

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-11-2009

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2004084842	A2	07-10-2004	US 2004198780 A1	07-10-2004
DE 10250110	A1	13-05-2004	AU 2003271728 A1	25-05-2004
			BR 0315771 A	20-09-2005
			CA 2503643 A1	13-05-2004
			CN 1708487 A	14-12-2005
			WO 2004039789 A1	13-05-2004
			EP 1558592 A1	03-08-2005
			JP 2006508088 T	09-03-2006
			KR 20050056210 A	14-06-2005
			MX PA05004466 A	26-07-2005
			US 2006155122 A1	13-07-2006
			ZA 200503278 A	28-06-2006
EP 0652229	A2	10-05-1995	AT 193303 T	15-06-2000
			AT 229971 T	15-01-2003
			AU 676602 B2	13-03-1997
			AU 7441194 A	27-04-1995
			DE 69424623 D1	29-06-2000
			DE 69424623 T2	08-02-2001
			DE 69431932 D1	30-01-2003
			DE 69431932 T2	24-12-2003
			DK 652229 T3	21-08-2000
			DK 0976760 T3	10-03-2003
			ES 2145094 T3	01-07-2000
			ES 2186300 T3	01-05-2003
			GR 3033702 T3	31-10-2000
			IT M1932113 A1	05-04-1995
			JP 3710150 B2	26-10-2005
			JP 7215993 A	15-08-1995
			NZ 264612 A	24-06-1997
			PT 652229 E	29-09-2000
			PT 976760 E	28-02-2003
			US 6084065 A	04-07-2000
EP 0419944	A2	03-04-1991	BR 9004803 A	10-09-1991
			CA 2026131 A1	27-03-1991
			DE 3932052 A1	04-04-1991
			HU 56084 A2	29-07-1991
			JP 3145478 A	20-06-1991
GB 1318291	A	23-05-1973	CH 552339 A	15-08-1974
			DE 2117807 A1	28-10-1971
			ES 390121 A1	01-05-1974
			FR 2089546 A5	07-01-1972
			JP 54007857 B	10-04-1979

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 09 16 4393

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-11-2009

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1318291	A	NL 7104858 A	19-10-1971
US 6630482	B1	07-10-2003	
		CA 2258924 A1	12-02-1998
		DE 69710208 D1	14-03-2002
		DE 69710208 T2	18-07-2002
		EP 0937044 A1	25-08-1999
		FR 2751969 A1	06-02-1998
		WO 9805642 A1	12-02-1998
		JP 2000515863 T	28-11-2000

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 6099562 A [0151]
- US 5886026 A [0151]
- US 5304121 A [0151]

Non-patent literature cited in the description

- **Gregory, R. J. et al.** *Nature*, 1990, vol. 347, 382-386 [0005]
- **Rich, D. P. et al.** *Nature*, 1990, vol. 347, 358-362 [0005]
- **Riordan, J. R. et al.** *Science*, 1989, vol. 245, 1066-1073 [0005]
- **Cutting, G. R. et al.** *Nature*, 1990, vol. 346, 366-369 [0007]
- **Dean, M. et al.** *Cell*, 1990, vol. 61, 863-870 [0007]
- **Kerem, B-S. et al.** *Science*, 1989, vol. 245, 1073-1080 [0007]
- **Kerem, B-S et al.** *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 8447-8451 [0007]
- **Quinton, P. M.** *FASEB J.*, 1990, vol. 4, 2709-2727 [0008]
- **Aridor M et al.** *Nature Med.*, 1999, vol. 5 (7), 745-751 [0008] [0013]
- **Shastri, B.S. et al.** *Neurochem. International*, 2003, vol. 43, 1-7 [0008] [0013]
- **Rutishauser, J. et al.** *Swiss Med Wkly*, 2002, vol. 132, 211-222 [0008] [0013]
- **Morello, JP et al.** *TIPS*, 2000, vol. 21, 466-469 [0008] [0013]
- **Bross P. et al.** *Human Mut.*, 1999, vol. 14, 186-198 [0008] [0013]
- **Dalemans et al.** *Nature Lond.*, 1991, vol. 354, 526-528 [0008]
- **Pasyk ; Foskett.** *J. Cell. Biochem.*, 1995, vol. 270, 12347-50 [0008]
- **Thomas Sorrell.** *Organic Chemistry.* University Science Books, 1999 [0033]
- *March's Advanced Organic Chemistry.* John Wiley & Sons, 2001 [0033]
- **Hwang, T. C. et al.** *J. Gen. Physiol.*, 1998, vol. 111 (3), 477-90 [0053]
- *Org. Lett.*, 2000, vol. 2 (8), 1165-1168 [0119]
- **A. J. Phillips et al.** *Chem Pharm Bull*, 1986, vol. 34 (7), 2851 [0119]
- *J. Pharmaceutical Sciences*, 1977, vol. 66, 1-19 [0129]
- **E. W. Martin.** *Remington's Pharmaceutical Sciences.* Mack Publishing Co, 1980 [0130]
- **Gonzalez, J. E. ; R. Y. Tsien.** Voltage sensing by fluorescence resonance energy transfer in single cells. *Biophys J*, 1995, vol. 69 (4), 1272-80 [0157] [0180]
- **Gonzalez, J. E. ; R. Y. Tsien.** Improved indicators of cell membrane potential that use fluorescence resonance energy transfer. *Chem Biol*, 1997, vol. 4 (4), 269-77 [0157] [0180]
- **Gonzalez, J. E. ; K. Oades et al.** Cell-based assays and instrumentation for screening ion-channel targets. *Drug Discov Today*, 1999, vol. 4 (9), 431-439 [0157] [0180]