



(11) **EP 2 149 303 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
03.02.2010 Bulletin 2010/05

(51) Int Cl.:
A01N 43/80 (2006.01) A01N 43/90 (2006.01)

(21) Application number: **09170115.1**

(22) Date of filing: **29.09.2006**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR**

(30) Priority: **04.10.2005 EP 05292073**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
**07119656.2 / 1 925 202
06255041.3 / 1 772 056**

(71) Applicant: **Rohm and Haas Company
Philadelphia,
Pennsylvania 19106-2399 (US)**

(72) Inventors:
• **Warwick, Eileen Fleck
Lansdale, PA 19446 (US)**

- **Diehl, Megan Anne
Line Lexington, PA 18932 (US)**
- **Shaw, Dalores Ann
Collegeville, PA 19426 (US)**
- **Levy, Richard
06560, France (FR)**

(74) Representative: **Buckley, Guy Julian
Patent Outsourcing Limited
1 King Street
Bakewell
Derbyshire DE45 1DZ (GB)**

Remarks:

This application was filed on 11-09-2009 as a
divisional application to the application mentioned
under INID code 62.

(54) **Microbicidal composition**

(57) Synergistic microbicidal compositions containing N-(n-butyl)-1,2-benzisothiazolin-3-one or N-methyl-1,2-benzisothiazolin-3-one.

EP 2 149 303 A1

Description

[0001] This invention relates to a synergistic combination of selected microbicides having greater activity than would be observed for the individual microbicides.

[0002] In some cases, commercial microbicides cannot provide effective control of microorganisms, even at high use concentrations, due to weak activity against certain types of microorganisms, e.g., those resistant to some microbicides, or due to aggressive environmental conditions. Combinations of different microbicides are sometimes used to provide overall control of microorganisms in a particular end use environment. For example, U.S. Pat. App. Pub. No. 2004/0014799 discloses a synergistic combination of N-(n-butyl)-1,2-benzisothiazolin-3-one (BBIT) and 2-methyl-4-isothiazolin-3-one (MI) over the limited range of ratios of BBIT to MI of 10:1 to 1.67:1. However, there is a need for additional combinations of microbicides having enhanced activity against various strains of microorganisms to provide effective control of the microorganisms. Moreover, there is a need for combinations containing lower levels of individual microbicides for environmental and economic benefit. The problem addressed by this invention is to provide such additional combinations of microbicides.

STATEMENT OF THE INVENTION

[0003] The present invention is directed to a microbicidal composition comprising: (a) N-(n-butyl)-1,2-benzisothiazolin-3-one; and (b) 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride.

[0004] The present invention is further directed to a microbicidal composition comprising: (a) N-methyl-1,2-benzisothiazolin-3-one; and (b) 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride.

DETAILED DESCRIPTION OF THE INVENTION

[0005] "BBIT" is N-(n-butyl)-1,2-benzisothiazolin-3-one. "MBIT" is N-methyl-1,2-benzisothiazolin-3-one.

[0006] As used herein, the following terms have the designated definitions, unless the context clearly indicates otherwise. The term "microbicide" refers to a compound capable of killing, inhibiting the growth of or controlling the growth of microorganisms at a locus; microbicides include bactericides, fungicides and algacides. The term "microorganism" includes, for example, fungi (such as yeast and mold), bacteria and algae. The term "locus" refers to an industrial system or product subject to contamination by microorganisms. The following abbreviations are used throughout the specification: ppm = parts per million by weight (weight/weight), mL = milliliter, ATCC = American Type Culture Collection, MBC = minimum biocidal concentration, and MIC = minimum inhibitory concentration. Unless otherwise specified, temperatures are in degrees centigrade (°C), and references to percentages (%) are by weight. Amounts of organic microbicides are given on an active ingredient basis in ppm (w/w).

[0007] The compositions of the present invention unexpectedly have been found to provide enhanced microbicidal efficacy at a combined active ingredient level lower than that of the individual microbicides. In one embodiment of the invention, those antimicrobial compositions which contain halogenated 3-isothiazolones contain relatively low levels thereof, preferably no more than 1000 ppm, more preferably no more than 500 ppm, more preferably no more than 100 ppm, and most preferably no more than 50 ppm. Concentrations of halogenated 3-isothiazolones in the composition of this invention are based on the total weight of active ingredients in the composition, i.e., the microbicides exclusive of any amounts of solvents, carriers, dispersants, stabilizers or other materials which may be present. In one embodiment of the invention, the antimicrobial composition contains less than 1000 ppm of 5-chloro-2-methyl-4-isothiazolin-3-one, more preferably no more than 500 ppm, more preferably no more than 100 ppm, and most preferably no more than 50 ppm.

[0008] In one embodiment of the invention, the antimicrobial composition comprises N-(n-butyl)-1,2-benzisothiazolin-3-one and 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride. Preferably, a weight ratio of N-(n-butyl)-1,2-benzisothiazolin-3-one to 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride is from 1:0.5 to 1:200, more preferably from 1:1 to 1:200.

[0009] In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and cis-1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride. Preferably, a weight ratio of N-methyl-1,2-benzisothiazolin-3-one to cis-1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride is from 1:0.2 to 1:400, more preferably from 1:0.3 to 1:200.

[0010] The microbicides in the composition of this invention may be used "as is" or may first be formulated with a solvent or a solid carrier. Suitable solvents include, for example, water; glycols, such as ethylene glycol, propylene glycol, diethylene glycol, dipropylene glycol, polyethylene glycol, and polypropylene glycol; glycol ethers; alcohols, such as methanol, ethanol, propanol, phenethyl alcohol and phenoxypropanol; ketones, such as acetone and methyl ethyl ketone; esters, such as ethyl acetate, butyl acetate, triacetyl citrate, and glycerol triacetate; carbonates, such as propylene carbonate and dimethyl carbonate; and mixtures thereof. It is preferred that the solvent is selected from water, glycols, glycol ethers, esters and mixtures thereof. Suitable solid carriers include, for example, cyclodextrin, silicas, diatomaceous

earth, waxes, cellulosic materials, alkali and alkaline earth (e.g., sodium, magnesium, potassium) metal salts (e.g., chloride, nitrate, bromide, sulfate) and charcoal.

[0011] When a microbicide component is formulated in a solvent, the formulation may optionally contain surfactants. When such formulations contain surfactants, they are generally in the form of emulsive concentrates, emulsions, micro-emulsive concentrates, or microemulsions. Emulsive concentrates form emulsions upon the addition of a sufficient amount of water. Microemulsive concentrates form microemulsions upon the addition of a sufficient amount of water. Such emulsive and microemulsive concentrates are generally well known in the art; it is preferred that such formulations are free of surfactants. U.S. Patent No. 5,444,078 may be consulted for further general and specific details on the preparation of various microemulsions and microemulsive concentrates.

[0012] A microbicide component also can be formulated in the form of a dispersion. The solvent component of the dispersion can be an organic solvent or water, preferably water. Such dispersions can contain adjuvants, for example, co-solvents, thickeners, anti-freeze agents, dispersants, fillers, pigments, surfactants, biodispersants, sulfosuccinates, terpenes, furanones, polycations, stabilizers, scale inhibitors and anti-corrosion additives.

[0013] When both microbicides are each first formulated with a solvent, the solvent used for the first microbicide may be the same as or different from the solvent used to formulate the other commercial microbicide, although water is preferred for most industrial biocide applications. It is preferred that the two solvents are miscible.

[0014] Those skilled in the art will recognize that the microbicide components of the present invention may be added to a locus sequentially, simultaneously, or may be combined before being added to the locus. It is preferred that the first microbicide and the second microbicide component be added to a locus simultaneously or sequentially. When the microbicides are added simultaneously or sequentially, each may individual components may contain adjuvants, such as, for example, solvent, thickeners, anti-freeze agents, colorants, sequestrants (such as ethylenediamine-tetraacetic acid, ethylenediaminedisuccinic acid, iminodisuccinic acid and salts thereof), dispersants, surfactants, biodispersants, sulfosuccinates, terpenes, furanones, polycations, stabilizers, scale inhibitors and anti-corrosion additives.

[0015] The microbicidal compositions of the present invention can be used to inhibit the growth of microorganisms or higher forms of aquatic life (such as protozoans, invertebrates, bryozoans, dinoflagellates, crustaceans, mollusks, etc) by introducing a microbicidally effective amount of the compositions onto, into, or at a locus subject to microbial attack. Suitable loci include, for example: industrial process water; electrocoat deposition systems; cooling towers; air washers; gas scrubbers; mineral slurries; wastewater treatment; ornamental fountains; reverse osmosis filtration; ultrafiltration; ballast water; evaporative condensers; heat exchangers; pulp and paper processing fluids and additives; starch; plastics; emulsions; dispersions; paints; latices; coatings, such as varnishes; construction products, such as mastics, caulks, and sealants; construction adhesives, such as ceramic adhesives, carpet backing adhesives, and laminating adhesives; industrial or consumer adhesives; photographic chemicals; printing fluids; household products, such as bathroom and kitchen cleaners; cosmetics; toiletries; shampoos; soaps; detergents; industrial cleaners; floor polishes; laundry rinse water; metalworking fluids; conveyor lubricants; hydraulic fluids; leather and leather products; textiles; textile products; wood and wood products, such as plywood, chipboard, wallboard, flakeboard, laminated beams, oriented strandboard, hardboard, and particleboard; petroleum processing fluids; fuel; oilfield fluids, such as injection water, fracture fluids, and drilling muds; agriculture adjuvant preservation; surfactant preservation; medical devices; diagnostic reagent preservation; food preservation, such as plastic or paper food wrap; food, beverage, and industrial process pasteurizers; toilet bowls; recreational water; pools; and spas.

[0016] Preferably, the microbicidal compositions of the present invention are used to inhibit the growth of microorganisms at a locus selected from one or more of mineral slurries, pulp and paper processing fluids and additives, starch, emulsions, dispersions, paints, latices, coatings, construction adhesives, such as ceramic adhesives, carpet backing adhesives, photographic chemicals, printing fluids, household products such as bathroom and kitchen cleaners, cosmetics, toiletries, shampoos, soaps, detergents, industrial cleaners, floor polishes, laundry rinse water, metal working fluids, textile products, agriculture adjuvant preservation, surfactant preservation, diagnostic reagent preservation, food preservation, and food, beverage, and industrial process pasteurizers.

[0017] The specific amount of the composition of this invention necessary to inhibit or control the growth of microorganisms and higher aquatic life forms in a locus depends upon the particular locus to be protected. Typically, the amount of the composition of the present invention to control the growth of microorganisms in a locus is sufficient if it provides from 0.1 to 1,000 ppm of the isothiazoline ingredient of the composition in the locus. It is preferred that the isothiazolone ingredients of the composition be present in the locus in an amount of at least 0.5 ppm, more preferably at least 4 ppm and most preferably at least 10 ppm. It is preferred that the isothiazolone ingredients of the composition be present in the locus in an amount of no more than 1000 ppm, more preferably no more than 500 ppm, and most preferably no more than 200 ppm.

[0018] In one embodiment of the invention, the composition is substantially free of enzymatic biocides. Preferably, when BBIT or MBIT and either methylparaben or ethylparaben are combined, the composition is substantially free of enzymatic biocides. Enzymatic biocides are enzymes having activity against microbes, as defined, e.g., in U.S. Pat. App. Pub. No. 2002/0028754.

Materials and Methods

[0019] The synergism of the combination of the present invention was demonstrated by testing a wide range of concentrations and ratios of the compounds.

[0020] One measure of synergism is the industrially accepted method described by Kull, F.C.; Eisman, P.C.; Sylwestrowicz, H.D. and Mayer, R.L., in Applied Microbiology 9:538-541 (1961), using the ratio determined by the formula:

$$Q_a/Q_A + Q_b/Q_B = \text{Synergy Index ("SI")}$$

wherein:

Q_A = concentration of compound A (first component) in ppm, acting alone, which produced an end point (MIC of Compound A).

Q_a = concentration of compound A in ppm, in the mixture, which produced an end point.

Q_B = concentration of compound B (second component) in ppm, acting alone, which produced an end point (MIC of Compound B).

Q_b = concentration of compound B in ppm, in the mixture, which produced an end point.

[0021] When the sum of Q_a/Q_A and Q_b/Q_B is greater than one, antagonism is indicated. When the sum is equal to one, additivity is indicated, and when less than one, synergism is demonstrated. The lower the SI, the greater the synergy shown by that particular mixture. The minimum inhibitory concentration (MIC) of a microbicide is the lowest concentration tested under a specific set of conditions that prevents the growth of added microorganisms.

[0022] Synergy tests were conducted using standard microtiter plate assays with media designed for optimal growth of the test microorganism. Minimal salt medium supplemented with 0.2% glucose and 0.1% yeast extract (M9GY medium) was used for testing bacteria; Potato Dextrose Broth (PDB medium) was used for testing yeast and mold. In this method, a wide range of combinations of microbicides and other personal care raw materials was tested by conducting high resolution MIC assays in the presence of various concentrations of BBIT or MBIT. High resolution MICs were determined by adding varying amounts of microbicide to one column of a microtitre plate and doing subsequent ten-fold dilutions using an automated liquid handling system to obtain a series of endpoints ranging from 2 ppm to 10,000 ppm active ingredient.

[0023] The synergy of the combinations of the present invention was determined against a bacterium, *Escherichia coli* (E. coli - ATCC #8739), a yeast, *Candida albicans* (C. albicans -- ATCC 10231), and a mold, *Aspergillus niger* (A. niger-ATCC 16404). The bacteria were used at a concentration of about 5×10^6 bacteria per mL and the yeast and mold at 5×10^5 fungi per mL. These microorganisms are representative of natural contaminants in many consumer and industrial applications. The plates were visually evaluated for microbial growth (turbidity) to determine the MIC after various incubation times at 25° C (yeast and mold) or 30° C (bacteria).

[0024] The test results for demonstration of synergy of the BBIT combinations of the present invention are shown below in Table 1. In each test, First Component (A) was BBIT and the Second Component (B) was the other commercial microbicide. Each table shows the specific combinations of BBIT and the second component; results against the microorganisms tested with incubation times; the end-point activity in ppm measured by the MIC for BBIT alone (Q_A), for the second component alone (Q_B), for BBIT in the mixture (Q_a) and for second component in the mixture (Q_b); the calculated SI value; and the range of synergistic ratios for each combination tested (BBIT/second component or A/B).

[0025] The test results for demonstration of synergy of the MBIT combinations of the present invention are shown below in Table 2. In each test, First Component (A) was MBIT and the Second Component (B) was the other commercial microbicide. Each table shows the specific combinations of MBIT and the second component; results against the microorganisms tested with incubation times; the end-point activity in ppm measured by the MIC for MBIT alone (Q_A), for the second component alone (Q_B), for MBIT in the mixture (Q_a) and for second component in the mixture (Q_b); the calculated SI value; and the range of synergistic ratios for each combination tested (MBIT/second component or A/B).

EP 2 149 303 A1

Table 1

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)

Second Component (B) = Quaternium-15 (active - 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride)

Microorganism	Q _a	Q _b	SI	A/B
E. coli 8739 - M9GY (72 hours)	0	200	1.00	----
	40	50	0.43	1/1
	40	60	0.43	1/1
	40	80	0.44	1/2
	40	100	0.45	1/3
	40	200	0.50	1/5
	40	300	0.55	1/8
	40	400	0.60	1/10
	40	500	0.65	1/13
	40	600	0.70	1/15
	40	800	0.80	1/20
	200	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
C. albicans 10231 - PDB (24 hours)	0	600	1.00	-----
	5	50	0.75	1/10
	5	60	0.77	1/12
	5	80	0.80	1/16
	5	100	0.83	1/20
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
A. niger 16404 - PDB (4 days)	0	3000	1.00	-----
	10	1000	0.58	1/100
	10	2000	0.92	1/200
	20	1000	0.83	1/50
	40	0	1.00	-----

[0026] The ratios of BBIT/1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride tested ranged from 1/0.02 to 1/5000.

Table 2

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Quaternium-15 (active - 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride)

Microorganism	Q _a	Q _b	SI	A/B
E. coli 8739 - M9GY (24 hours)	0	50	1.00	-----
	5	20	0.90	1/400
	7.5	2	0.79	1/0.3
	7.5	3	0.81	1/0.4
	7.5	4	0.83	1/0.5
	7.5	5	0.85	1/0.7
	7.5	6	0.87	1/0.8
	7.5	8	0.91	1/1
	7.5	10	0.95	1/1
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
C. albicans 10231 - PDB	0	500	1.00	-----

EP 2 149 303 A1

(continued)

Microorganism (24 hours)	Q _a	Q _b	SI	A/B
	2.5	500	1.33	1/200
	5	300	1.27	1/60
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
A. niger 16404 - PDB	0	3000	1.00	-----
(3 days)	10	1000	0.58	1/100
	10	2000	0.92	1/200
	20	1000	0.83	1/50
	40	0	1.00	-----

[0027] The ratios of MBIT/1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride tested ranged from 1/0.01 to 1/4000.

Claims

1. A microbicidal composition comprising:

- (a) N-(n-butyl)-1,2-benzisothiazolin-3-one; and
- (b) 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride.

2. The microbicidal composition of claim 1, wherein the weight ratio of N-(n-butyl)-1,2-benzisothiazolin-3-one to 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride is 1:0.5 to 1:200.

3. The microbicidal composition of claim 2, wherein the weight ratio of N-(n-butyl)-1,2-benzisothiazolin-3-one to 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride is 1:1 to 1:200.

4. A microbicidal composition comprising:

- (a) N-methyl-1,2-benzisothiazolin-3-one; and
- (b) 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride.

5. The microbicidal composition of claim 4, wherein the weight ratio of N-methyl-1,2-benzisothiazolin-3-one to cis-1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride is 1:0.2 to 1:400.

6. The microbicidal composition of claim 5, wherein the weight ratio of N-methyl-1,2-benzisothiazolin-3-one to cis-1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride is 1:0.3 to 1:200.



EUROPEAN SEARCH REPORT

Application Number
EP 09 17 0115

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y	EP 0 375 367 A2 (ROHM & HAAS [US]) 27 June 1990 (1990-06-27) * abstract * * tables 15, 18, 19 * -----	1-6	INV. A01N43/80 A01N43/90
Y	EP 1 245 153 A2 (BODE CHEMIE GMBH & CO [DE]) 2 October 2002 (2002-10-02) * page 2, paragraph 8 * -----	1-6	
Y	EP 0 787 430 A1 (RIEDEL DE HAEN AG [DE]) 6 August 1997 (1997-08-06) * page 2, line 26 - line 29 * -----	1-6	
			TECHNICAL FIELDS SEARCHED (IPC)
			A01N
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 18 December 2009	Examiner Davies, Maxwell
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

2
EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 09 17 0115

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18-12-2009

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0375367	A2	27-06-1990	AT 134477 T	15-03-1996
			AT 173880 T	15-12-1998
			AU 624884 B2	25-06-1992
			AU 4686689 A	28-06-1990
			BR 8906719 A	11-09-1990
			DE 68925801 D1	04-04-1996
			DE 68925801 T2	06-02-1997
			DE 68928874 D1	14-01-1999
			DE 68928874 T2	29-07-1999
			ES 2083977 T3	01-05-1996
			ES 2124479 T3	01-02-1999
			FI 94207 B	28-04-1995
			HU 58469 A2	30-03-1992
			IL 92728 A	25-01-1994
			JP 2221205 A	04-09-1990
			JP 2886226 B2	26-04-1999
			PT 92653 A	29-06-1990
			US 4964892 A	23-10-1990
EP 1245153	A2	02-10-2002	AT 282315 T	15-12-2004
			AT 329488 T	15-07-2006
			DE 10112755 A1	02-10-2002
			DK 1249166 T3	14-03-2005
			DK 1245153 T3	16-10-2006
			EP 1249166 A1	16-10-2002
			ES 2233728 T3	16-06-2005
			ES 2265461 T3	16-02-2007
EP 0787430	A1	06-08-1997	DE 19548710 A1	26-06-1997
			JP 9278611 A	28-10-1997

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20040014799 A [0002]
- US 5444078 A [0011]
- US 20020028754 A [0018]

Non-patent literature cited in the description

- Kull, F.C. ; Eisman, P.C. ; Sylwestrowicz, H.D. ; Mayer, R.L. *Applied Microbiology*, 1961, vol. 9, 538-541 [0020]