



(11) **EP 2 197 900 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

- (15) Correction information:  
**Corrected version no 1 (W1 B1)**  
**Corrections, see**  
**Description Paragraph(s) 11, 13, 80, 90, 268**  
**Claims DE 5, 6, 11**  
**Claims EN 10, 11**  
**Claims FR 10, 11**  
**Numerous spelling errors of minor importance**
- (51) Int Cl.:  
**C07K 7/64** <sup>(2006.01)</sup> **C07K 14/705** <sup>(2006.01)</sup>  
**A61K 38/12** <sup>(2006.01)</sup> **A61K 38/16** <sup>(2006.01)</sup>
- (86) International application number:  
**PCT/EP2008/006932**
- (87) International publication number:  
**WO 2009/027063 (05.03.2009 Gazette 2009/10)**

(48) Corrigendum issued on:  
**09.01.2013 Bulletin 2013/02**

(45) Date of publication and mention  
of the grant of the patent:  
**01.08.2012 Bulletin 2012/31**

(21) Application number: **08801676.1**

(22) Date of filing: **22.08.2008**

(54) **MUTANT DOUBLE CYCLIZED RECEPTOR PEPTIDES INHIBITING BETA 1-ADRENOCEPTOR ANTIBODIES**

BETA 1-ADRENOCEPTOR-ANTIKÖRPER INHIBIERENDE, MUTIERTE, DOPPELT CYCLISIERTE REZEPTORPEPTIDE

RÉCÉPTEURS PEPTIDIQUES CYCLISÉS DOUBLE MUTANTS INHIBITEURS D'ANTICORPS ANTI-RÉCÉPTEURS BÊTA 1-ADRÉNERGIQUES

- (84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT  
RO SE SI SK TR**
- (30) Priority: **24.08.2007 EP 07016637**
- (43) Date of publication of application:  
**23.06.2010 Bulletin 2010/25**
- (60) Divisional application:  
**12170992.7 / 2 497 777  
12178671.9**
- (73) Proprietor: **Julius-Maximilians-Universität  
Würzburg  
97070 Würzburg (DE)**
- (72) Inventors:  
• **JAHNS, Roland  
97080 Würzburg (DE)**
- **JAHNS, Valérie  
97080 Würzburg (DE)**
- **LOHSE, Martin  
97074 Würzburg (DE)**
- **NIKOLAEV, Viacheslav  
97084 Würzburg (DE)**
- (74) Representative: **Vossius & Partner  
Siebertstrasse 4  
81675 München (DE)**
- (56) References cited:  
**EP-A- 1 270 030 WO-A-01/21660  
WO-A-01/83693 WO-A-2006/103101**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 2 197 900 B9**

- JAHNS R ET AL: "A NEW CYCLIC RECEPTOR-PEPTIDE PREVENTS DEVELOPMENT OF HEART DILATATION AND FAILURE INDUCED BY ANTIBODIES ACTIVATING CARDIAC .BETA.1-ADRENERGIC RECEPTORS" CIRCULATION, AMERICAN HEART ASSOCIATION, DALLAS, TX, vol. 112, no. 17, SUPPL. S, 1 October 2005 (2005-10-01), page U53, ABSTRACT 120, XP009072196 ISSN: 0009-7322
- FRIDELL E ET AL: "A cyclized peptide for studies of human parvovirus B19 infection" JOURNAL OF IMMUNOLOGICAL METHODS, ELSEVIER SCIENCE PUBLISHERS B.V., AMSTERDAM, NL, vol. 138, no. 1, 8 April 1991 (1991-04-08), pages 125-128, XP023986994 ISSN: 0022-1759 [retrieved on 1991-04-08]
- BUCK S H ET AL: "The dogfish peptides scyliorhinin I and scyliorhinin II bind with differential selectivity to mammalian tachykinin receptors" EUROPEAN JOURNAL OF PHARMACOLOGY, ELSEVIER BV, NL, vol. 144, no. 1, 24 November 1987 (1987-11-24), pages 109-111, XP023742211 ISSN: 0014-2999 [retrieved on 1987-11-24]

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

## Description

**[0001]** The present invention relates to novel  $\beta$ -AR homologous cyclopeptide-mutants comprising only two cysteine residues able to form an intramolecular linkage. Disclosed herein are also linear peptides that can form these cyclopeptide-mutants and nucleic acid molecules encoding these cyclopeptide-mutants and linear peptides. Moreover, vectors and recombinant host cells comprising said nucleic acid molecule are disclosed. A method for producing the disclosed cyclopeptide-mutants is provided. Further provided is a composition comprising the peptides of the invention. The present invention also relates to therapeutic and diagnostic means, methods and uses taking advantage of the peptides of the invention and to means, methods and uses for detecting anti- $\beta$ -adrenergic receptor antibodies like anti- $\beta_1$ -adrenergic receptor antibodies.

**[0002]** Progressive cardiac dilatation and pump failure of unknown etiology has been termed "idiopathic" dilated cardiomyopathy (DCM) (Richardson 1996 Circulation, 93, 841-842). DCM represents one of the main causes of severe heart failure with an annual incidence of up to 100 patients and a prevalence of 300-400 patients per million (AHA report 2007). Mutations in genes encoding myocyte structural proteins (Morita 2005) and several cardiotoxins, including alcohol, anthracyclines, and, more recently, therapeutically used monoclonal antibodies (e.g., trastuzumab) account for about one third of DCM cases (Chien 2000, Fabrizio and Regan 1994). The etiology of the remaining two thirds is still poorly understood, however. At present the large majority of DCM is thought to arise from an initial (mostly viral) infection leading to acute myocarditis which upon activation of the immune system may progress to (chronic) autoimmune myocarditis resulting in cardiac dilatation and severe congestive heart failure; the latter progression occurs particularly, when associated (a) with the development of autoantibodies against distinct myocyte sarcolemmal or membrane proteins which are essential for cardiac function (Freedman 2004, Jahns 2006), or (b) with chronic inflammation of the myocardium and viral persistence (Kühl 2005). These recent findings are further strengthened by the fact, that patients with DCM often have alterations in both cellular and humoral immunity (Jahns 2006, Limas 1997, Luppi 1998, Mahrholdt 2006). Under such conditions an initial acute inflammatory reaction may proceed into a kind of low-grade inflammation (MacLellan 2003) facilitating the development of abnormal or misled immune responses to the primary infectious trigger (Freedman 2004, Kühl 2005, MacLellan and Lysis 2003, Maekawa 2007, Smulski 2006).

**[0003]** In the context of their humoral response a substantial number of DCM patients have been found to develop cross-reacting antibodies and/or autoantibodies to various cardiac antigens, including mitochondrial proteins (e.g., adenine nucleotide translocator, lipoamide and pyruvate dehydrogenase (Pohlner 1997, Schultheiss 1985, Schultheiss 1988, Schulze 1999)), sarcolemmal proteins (e.g., actin, laminin, myosin, troponin (Caforio 2002, Göser 2006, Li 2006, Neumann 1990, Okazaki 2003)), and membrane proteins (e.g., cell surface adrenergic or muscarinic receptors (Christ. 2006, Fu 1993, Jahns 1999b, Magnusson 1994). From these, only a few selected antibodies appear to be able to cause myocardial tissue injury and to induce severe congestive heart failure by itself, however. In addition, the individual genetic predisposition (including the respective human leucocyte antigen (HLA)- and the major histocompatibility complex (MHC)-phenotype (Limas 1996)) may also significantly contribute to the susceptibility to self-directed immune reactions and the phenotypic expression of the disease (Limas 2004, MacLellan 2003).

**[0004]** Homologies between myocyte surface molecules such as membrane receptors and viral or bacterial proteins have been proposed as a mechanism for the elaboration of endogenous cardiac autoantibodies by antigen mimicry (Hoebeke 1996, Mobini 2004). Chagas' heart disease, a slowly evolving inflammatory cardiomyopathy, is one of the most prominent examples for this mechanism (Elies 1996, Smulski 2006). The disease originates from an infection with the protozoon *Trypanosoma cruzi*; molecular mimicry between the ribosomal P2 $\beta$ -protein of *T. cruzi* and the N-terminal half of the second extracellular loop of the  $\beta_1$ -adrenergic receptor ( $\beta_1$ -AR) results in generation of cross-reacting antibodies in about 30% of the Chagas' patients (Ferrari 1995). Because receptor-autoantibodies from patients with DCM preferentially recognize the C-terminal half of the same loop (Wallukat 1995), it was speculated that these antibodies might originate from molecular mimicry between the  $\beta_1$ -AR and a hitherto unidentified viral pathogen (Magnusson 1996). Another -probably more relevant- mechanism leading to the production of endogenous cardiac autoantibodies would be primary cardiac injury followed by (sudden or chronic) liberation of a "critical amount" of antigenic determinants from the myocyte membrane or cytoplasm, previously hidden to the immune system. Such injury most likely occurs upon acute infectious (myocarditis), toxic, or ischemic heart disease (myocardial infarction) resulting in myocyte apoptosis or necrosis (Caforio 2002, Rose 2001). Presentation of myocardial self-antigens to the immune system may then induce an autoimmune response, which in the worst case results in perpetuation of immune-mediated myocyte damage involving either cellular (e.g., T-cell), or humoral (e.g., B-cell) immune responses, or co-activation of both the innate and the adaptive immune system (Eriksson 2003, Rose 2001).

**[0005]** From a pathophysiological point of view, it seems reasonable to link the harmful (e.g., cardiomyopathy-inducing) potential of a heart-specific autoantibody to the *accessibility* and to the *functional relevance* of the corresponding target. Myocyte surface receptors are easily accessible to autoantibodies (Okazaki 2005). The two most promising candidates are the cardiac  $\beta_1$ -AR (representing the predominant adrenoceptor subtype in the heart) and the M2-muscarinic acetylcholine receptor; against both receptors autoantibodies have been detected in DCM patients (Fu. 1993, Jahns 1999b,

Matsui 1995). Whereas anti-muscarinic antibodies (exhibiting an agonist-like action on the cardiac M2 acetylcholine-receptor) have been mainly associated with negative chronotropic effects at the sinuatrial level (e.g., sinus node dysfunction, atrial fibrillation (Baba 2004, Wang. 1996)), agonistic anti- $\beta_1$ -AR antibodies have been associated with both the occurrence of severe arrhythmia at the ventricular level (Christ 2001, Iwata 2001a), and the development of (maladaptive) left ventricular hypertrophy, finally switching to left ventricular enlargement and progressive heart failure (Iwata 2001b, Jahns 1999b, Khoyneshad 2007). Both autoantibodies appear to be directed against the second extracellular loop of the respective receptors. To generate an autoimmune response, myocyte membrane proteins (e.g., receptors) must be degraded to small oligopeptides able to form a complex with a MHC or HLA class II molecule of the host (Hoebeke 1996). In case of the human  $\beta_1$ -AR computer-based analysis for potential immunogenic amino-acid stretches has shown, that the only portion of the receptor molecule containing B- and T-cell epitopes and being accessible to antibodies was in fact the predicted second extracellular receptor loop ( $\beta_1$ -EC<sub>II</sub>) (Hoebeke 1996). This might explain the successful use of second loop-peptides for the generation of  $\beta_1$ -specific receptor antibodies in different animal-models (Iwata 2001 b, Jahns. 2000, Jahns 1996). Moreover, in the last decade several groups have independently demonstrated that second loop antibodies preferentially recognize intact native  $\beta_1$ -AR in various immunological assays (whole cell-ELISA, immunoprecipitation, immunofluorescence), indicating that they are "conformational" (Hoebeke 1996, Jahns 2006). Functional testing revealed that the same antibodies also affected receptor function, such as intracellular cAMP-production and/or cAMP-dependent protein kinase (PKA) activity, suggesting that they may act as allosteric regulators of  $\beta_1$ -AR activity (Jahns 2000, Jahns 2006). The structure of the  $\beta_1$ -AR was also analyzed by Warne (2008 Nature. DOI: 10. 1038).

**[0006]** Following Witebsky's postulates (Witebsky 1957) *indirect* evidence for the autoimmune etiology of a disease requires identification of the trigger (e.g., the responsible self-antigen), and induction of a self antigen-directed immune response in an experimental animal, which then must develop a similar disease. *Direct* evidence, however, requires reproduction of the disease by transfer of homologous pathogenic antibodies or autoreactive T-cells from one to another animal of the same species (Rose 1993).

**[0007]** To analyze the pathogenetic potential of anti- $\beta_1$ -AR antibodies, Jahns et al. has chosen an experimental *in vivo* approach, which met the Witebsky criteria for direct evidence of autoimmune diseases. DCM was induced by immunizing inbred rats against  $\beta_1$ -EC<sub>II</sub> (100% sequence homology between human and rat; *indirect evidence*); then the disease was reproduced in healthy animals by isogenic transfer of rat anti- $\beta_1$ -AR "autoantibodies" (*direct evidence*) (Jahns 2004). The animals developed progressive left ventricular (LV)-dilatation and dysfunction, a relative decrease in LV wall-thickness, and selective downregulation of  $\beta_1$ -AR, a feature that is also seen in human DCM (Lohse 2003).

**[0008]** These results, together with an agonist-like short-term effect of the antibodies *in vivo* (Jahns 2004), suggest that both the induced and the transferred cardiomyopathic phenotypes can be attributed mainly to the mild but sustained receptor activation achieved by stimulatory anti- $\beta_1$ -AR antibodies. This hypothesis is supported by the large body of data available on the cardiotoxic effects of excessive and/or long term  $\beta_1$ -AR activation seen after genetic or pharmacological manipulation (Engelhardt 1999, Woodiwiss 2001). Therefore, anti- $\beta_1$ -AR induced dilated immune-cardiomyopathy (DiCM) can now be regarded as a pathogenetic disease entity of its own, together with other established receptor-directed autoimmune diseases such as myasthenia gravis or Graves' disease (Freedman 2004, Hershko 2005, Jahns 2004, Jahns 2006).

**[0009]** The clinical importance of cardiac autoantibodies is difficult to assess, since low titers of such antibodies can also be detected in the healthy population as a part of the natural immunologic repertoire (Rose 2001). However, regarding functionally active anti- $\beta_1$ -AR antibodies previous data from Jahns et al. has demonstrated that their prevalence is almost negligible in healthy individuals (<1%) provided that a screening procedure based on cell-systems presenting the target (e.g., the  $\beta_1$ -AR) in its natural conformation is used (Jahns 1999b). By employing the latter screening method, occurrence of anti- $\beta_1$ -AR autoantibodies could also be excluded in patients with chronic valvular or hypertensive heart disease (Jahns 1999a). In contrast, the prevalence of stimulating anti- $\beta_1$ -AR was ~10% in ischemic (ICM) and ~30% in dilated cardiomyopathy (DCM) (Jahns 1999b), which was significantly higher than in healthy controls, but in the lower range of previous reports on DCM collectives (33% to 95% prevalence) (Limas 1992, Magnusson 1994, Wallukat 1995). It seems conceivable that differences in screening methods aiming to detect functionally active anti- $\beta_1$ -AR autoantibodies most likely account for the wide range of prevalences reported in the past (Limas 1992). In fact, only a minor fraction of ELISA-defined human anti- $\beta_1$ -AR autoantibodies was able to bind to cell surface located native  $\beta$ -AR. Only this fraction recognized (as determined by immunofluorescence) and activated (as determined by increases in cellular cAMP and/or PKA activity) human  $\beta_1$ -AR expressed in the membrane of intact eukaryotic cells (Jahns 2000, Jahns 1999b). Therefore, cell systems presenting the target in its natural conformation represent an essential tool in the screening for functionally relevant anti- $\beta$ -AR autoantibodies (Nikolaev 2007).

**[0010]** Clinically, the presence of anti- $\beta_1$ -AR autoantibodies in DCM has been shown to be associated with a more severely depressed cardiac function (Jahns 1999b), the occurrence of more severe ventricular arrhythmia (Chiale 2001), and a higher incidence of sudden cardiac death (Iwata 2001a). Recent data comparing antibody-positive with antibody-negative DCM patients over a follow-up period of more than 10 years not only confirmed a higher prevalence of ventricular

arrhythmia in the presence of activating anti- $\beta_1$ -AR, but also revealed that antibody-positivity predicted an almost three-fold increased cardiovascular mortality-risk (Störk 2006). Taken together, the available clinical data underscore the pathophysiological relevance of functionally active anti- $\beta_1$ -AR antibodies in DCM.

**[0011]** One today generally accepted pharmacological strategy would be the use of beta-blocking agents in order to attenuate or even abolish the autoantibody-mediated stimulatory effects, at least if  $\beta$ -blockers can indeed prevent the antibody-induced activation of  $\beta_1$ -AR (Freedman 2004, Jahns 2000, Matsui 2001, Jahns 2006). New therapeutic approaches actually include elimination of stimulatory anti- $\beta_1$ -AR by non-selective or selective immunoadsorption (Hershko 2005, Wallukat 2002), or direct targeting of the anti- $\beta_1$ -EC<sub>II</sub> antibodies and/or the anti- $\beta_1$ -EC<sub>II</sub> producing B-cells themselves (that is, induction of immune tolerance) (Anderton 2001). Non-selective immunoadsorption, however, because of an increased risk of infection after immunoglobulin depletion, requires the substitution of human IgG on the ground of safety (Felix 2000) with all possible side effects of substituted human proteins known in the art including severe anaphylactic reactions and death.

**[0012]** WO 01/21660 discloses certain peptides homologous to epitopes of the 1<sup>st</sup> and the 2<sup>nd</sup> loop of  $\beta_1$ -AR, and proposes to apply these peptides for medical intervention of dilatative cardiomyopathy (DCM). Even if WO 01/21660 mentions marginally that peptides may be modified in order to protect them against serum proteases, for example by cyclization, corresponding examples and embodiments are not given and any in vitro or in vivo effect of the proposed peptides on the course of DCM or on the course of receptor-antibody titers is not shown. Moreover, in WO 01/21660 intends to rely on the above mentioned non-selective immunoadsorption approaches bearing the correspondingly mentioned risks.


**[0013]** In contrast thereto, the newly developed  $\beta_1$ -EC<sub>II</sub>-homologous cyclopeptides (e.g.  $\beta_1$ -EC<sub>II</sub>-CPs) were employed six weeks after the active induction of stimulatory anti- $\beta_1$ -EC<sub>II</sub> antibodies.  $\beta_1$ -EC<sub>II</sub>-CPs are cyclopeptides containing 3 cysteine residues and hence, can form intramolecular bonds, whereby there is a potential option to form two intramolecular bonds (besides the cyclization between the N- and C-terminus), individually.  $\beta_1$ -EC<sub>II</sub>-CP significantly reduced the amount of circulating anti- $\beta_1$ -EC<sub>II</sub> antibodies and effectively prevented development of cardiac dilatation and dysfunction (Boivin 2005). The above-mentioned  $\beta_1$ -EC<sub>II</sub>-CPs were also disclosed in WO 2006/103101.

**[0014]** In view of the present art, the technical problem underlying the present invention is the provision of improved and easily obtainable means and methods for the medical intervention of diseases related to anti- $\beta$ -AR antibodies, particular to anti- $\beta_1$ -EC<sub>II</sub> antibodies.

**[0015]** The technical problem is solved by provision of the embodiments characterized in the claims.

**[0016]** Accordingly, in a first aspect, the present invention relates to  $\beta$ -AR homologous cyclopeptide-mutants (also termed herein as "cyclic peptides" or "cyclopeptides" and the like), particularly to  $\beta_1$ -AR homologous cyclopeptide-mutants, namely  $\beta_1$ -EC<sub>II</sub> homologous cyclopeptide-mutants. The structure of these cyclopeptide-mutants/cyclic peptides is characterized by being able to form only one individual intramolecular disulphide bond.

**[0017]** Particularly, in the first aspect, the present invention relates to a cyclic peptide of formula I:



wherein

- a) x is an amino acid other than Cys;
- b) h is any integer from 1 to 15;
- c) i is any integer from 0 to 14;
- d) x<sup>c</sup> is Pro; and
- e) y is an amino acid other than Cys, and

wherein said cyclic peptide consists of at least 16 and of at most 25 amino acids and comprises the amino acid stretch

**Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 45); or**  
**Glu-Ser-Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 46),**

wherein Xxx<sub>1</sub>, Xxx<sub>3</sub> and Xxx<sub>4</sub> is an amino acid other than Cys, and  
 wherein said cyclic peptide

- i) is capable of reducing an antibody-mediated activation of  $\beta_1$ -adrenergic receptor ( $\beta_1$ -AR);

- ii) is capable of binding (auto)antibodies against the second extracellular loop (EC<sub>II</sub>) of  $\beta_1$ -AR; and/or  
 iii) is capable of inhibiting the interaction between  $\beta_1$ -AR and (auto)antibodies against EC<sub>II</sub> of  $\beta_1$ -AR.

**[0018]** Particular preferred embodiments, as discussed below, are specific cyclic peptides as depicted in formulas VII, IX, IX', VI or VIII.

**[0019]** The present invention solves the above identified technical problem since, as documented herein below and in the appended examples, it was surprisingly found that mutant cyclic peptides containing only two cysteines, which can form one single defined, individual intramolecular disulfide bond, are also able to inhibit anti- $\beta_1$ -AR antibodies, and are useful in inhibiting stimulatory anti- $\beta_1$ -AR antibodies.

**[0020]** It was furthermore surprisingly found in context of the present invention that the peptide-mutants with a cyclic structure (i.e. the cyclic peptides) as described and provided herein are superior to their linear counterparts in terms of both, the recognition or scavenging of conformational anti- $\beta_1$ -AR antibodies and their antibody-neutralizing (i.e. pharmaceutical) potential. These findings were obtained by the exemplarily employment of ELISA competition assays and functional (cAMP) FRET-assays, respectively.

**[0021]** In addition, the inventive cyclic peptides comprising only two cysteines, which can form one single defined, individual intramolecular disulfide bond, can easily be obtained/manufactured, biochemically characterized and purified. This is particularly true when pure fractions of the same cyclopeptide isomers are required. In context of this invention, a mixture of cyclopeptide isomers, i.e. stereo-isomers, comprising cyclopeptide isomers with different intramolecular disulphide bonds is avoided. As documented herein below, because of this avoidance a specific and clean medical product (fulfilling GMP standards) comprising isomers all with the same intramolecular disulfide bond can be obtained.

**[0022]** It was a further, surprising finding in context of the present invention, and as illustrated in the appended examples, that the exact nature of the exchange of one of the cysteine residues with a serine residue markedly determined the antibody neutralizing potency of cyclic peptides derived from  $\beta_1$ -EC<sub>II</sub>.

**[0023]** Particularly, a Cys→Ser exchange like that at position 18 of the herein exemplarily and preferably disclosed 25-meric cyclopeptide (formulas VII/IX), at position 17 of the herein exemplarily and preferably disclosed 22-meric cyclopeptide (formula IX') or at position 14 of the herein exemplarily and preferably disclosed 18-meric cyclopeptide (formulas VI/VIII), respectively, yields cyclic peptides (Cys-Ser cyclic peptides) with excellent antibody-neutralizing and pharmacological effects in vitro (Figs. 4-11 and 27), whereas the Cys→Ser exchange at position 17, 16 or 13 of the herein exemplarily disclosed 25-meric, 22-meric or 18-meric cyclic peptide, respectively (Ser-Cys cyclic peptides), had, surprisingly, almost no inhibitory effect. This inhibitory effect could neither be detected regarding their properties as antibody-scavengers nor in terms of their capability of inhibiting functional antibody-effects; as neutralization of receptor-stimulation in vitro as shown in, for example, Figs. 4-10).

**[0024]** It was a further finding in context of the present invention that an almost perfect steric imitation of the EC<sub>II</sub>- $\beta_1$ -AR domain can be obtained by a second loop-homologous cyclized peptide comprising 22 amino acids, for example 21 amino-acids of the published original primary sequence of the human  $\beta_1$ -AR, i.e. amino-acids 200 (R) to 221 (T) (numbering according to Frielle et al. 1987, PNAS 84, pages 7920-7924), with an additional amino acid residue (for example glycine (G)) to close the synthetic cycle at position 222 to form a 22 AA cyclopeptide.

**[0025]** Without being bound by theory, the cardio-protective and immunomodulating activity of the cyclic peptides largely depends on their conformation. It was additionally found out in context of this invention that an introduction of the smallest naturally occurring amino-acid glycine at the (predicted) ring closure site (or at the position corresponding thereto) leads to an enhanced binding of anti- $\beta_1$ -AR autoantibodies, i.e. apparently further enhances the similarity of the 22 AA cyclopeptide with the EC<sub>II</sub>- $\beta_1$ -AR domain. Particularly, the appended examples, inter alia, indicate that the cyc22AA cyclopeptides have a significantly higher antibody-blocking efficiency *in vivo* than other EC<sub>II</sub>-imitating cyclopeptides larger (i.e., cyc25AA peptides) or smaller (i.e., cyc18AA peptides). Computer-aided modelling studies with said 22 AA cyclopeptide confirmed an excellent imitation of the predicted second extracellular loop structure with a calculated difference in size of only 4.5 Angström (4.5 Å) at the base of the cyclopeptide (opposed to the assumed antibody-binding site), when compared with the predicted native second extracellular loop backward helix (see also appended Fig. 24). Moreover, it was demonstrated herein and in the appended examples that particularly said 22 AA cyclopeptide reduces the titer of anti- $\beta_1$ -AR autoantibodies with an extraordinary high efficiency.

**[0026]** Since replacement of one of the three cysteines present in the cyclic 22 AA peptide allows for the introduction of a reinforced disulfide bridge (as a second "internal" cycle, generated by double cyclization) between the two remaining cysteines, the resultant cyclic 22 AA cyclopeptide also represents a biochemically unambiguously defined product (see also Figs. 25 and 26).

**[0027]** It was also surprisingly found that a Gln↔D-Glu exchange at position 25 (25-meric cyclopeptide-mutants) or 18 (18-meric cyclopeptide-mutants) did not significantly influence the blocking capacity of the cyclopeptides, regardless of their length; i.e., 25 versus 18 amino-acids as shown in Figs. 6, 7 and 9).

**[0028]** The examples below also document that the cyclic peptides as disclosed herein show improved features, for example as compared to peptides comprising three Cys residues (for example the Cys/Cys cyclic peptides disclosed in

WO 2006/103101). Examples of improved features of the cyclic peptides of this invention are an extremely good capacity for blocking anti- $\beta_1$ -AR antibodies and their advanced producibility according to GMP standards.

[0029] In context of the present invention, the *in vitro* findings were generally confirmed in *in vivo* tests (Figs. 12-16 and 28/29). Interestingly, the difference in the blocking efficiency of the Cys<sub>18,17 or 14</sub>→Ser<sub>18,17 or 14</sub> mutated cyclopeptides compared with that of the linear peptides was even more pronounced *in vivo* (Figs. 5, 7, and 14-16).

[0030] The established rat model of anti-betal-adrenergic antibody-induced autoimmune-cardiomyopathy (Jahns, 2004) served to assess the efficacy of the generated beta1-ECII homologous cyclopeptide mutants *in vivo*. The *in vivo* data indicate, that the efficiency of the disclosed mutated cyclopeptides (e.g. 18AA Cys/Ser cyclopeptide) might equally depend on the administered dose (Figs. 14-16).

[0031] In addition, the *in vivo* experiments demonstrated that the antibody-blocking capacity of mutant cyclopeptides is seemingly not affected by a reduction in the number of amino acids from a 25-meric to a 18-meric cyclopeptide; both *in vitro* and *in vivo* data demonstrate an excellent comparability of these two 2 cysteine-containing single disulfide bond 25AA Cys/Ser or 18AA Cys/Ser cyclopeptide mutants. It should be noted, however, that both 1.0 mg/kg 25AA-meric Cys/Ser as well as high dose (i.e., 4.0 mg/kg Bw) 18AA-meric Cys/Ser mutants led to an initial transient increase in antibody-titers, and thus postponed a significant reduction in receptor antibody titers to the third or fourth cyclopeptide-application (third or fourth month of therapy). This phenomenon did not occur with either 1.0 or 2.0 mg/kg Bw doses of 18AA Cys/Ser cyclopeptide mutants (Fig. 16B,C).

[0032] Animals to which particularly the 18meric or 25meric cyclic peptide as disclosed herein was administered showed no signs of abnormalities, and only the desired effect of the administered peptide, namely the blockage of anti- $\beta_1$ -AR antibodies, was detected. Accordingly, the peptides as provided herein display no undesired side effects or toxicity at the applied dosage regimen. This was further demonstrated herein by showing that no toxicity on the kidney was exerted by the cyclic peptides of the invention (no mechanical obstruction of glomerular membranes was detected; Fig. 23). In addition, the routine laboratory parameters indicative of kidney function remained normal under 12 months of CP-application and did not differ from untreated control animals. (Fig. 22A, B).

[0033] The antibody-blocking capacity of mutated cyclopeptides of this invention is not affected by the length of the peptide, as long as the peptide is not shorter than 18 AA and not longer than 25 AA. This was exemplarily demonstrated by the reduction of the number of amino acids of the peptide from 25 to 18. Within the range of 18 to 25 amino acids, cyclic peptides having 22 amino acids are most effective in accordance with this invention and, accordingly, are a particular preferred embodiment. An example of such a particular preferred 22mer cyclic peptide is shown in formula IX'.

[0034] One advantage of the cyclopeptide mutants of the present invention is -by mutating one particular cysteine (preferably the Cys corresponding to Cys 216 of the amino acid sequence of  $\beta_1$ -AR) to a serine-residue and by reinforcing formation of the unique possible intramolecular S-S bridge through a second S-S specific cyclization procedure- that their conformational restraint is increased. In comparison to peptides known in the art, this increased restraint of the inventive peptides leads to a molecule that better mimics the epitope presented in the native conformation of the second  $\beta_1$ -EC<sub>II</sub> loop on the cell surface.

[0035] Beta blockers, such as bisoprolol, which are used in the art for the treatment of DCM and other diseases which are caused by stimulatory anti- $\beta_1$ -AR antibodies, significantly reduce both heart rate and blood pressure. In contrast thereto, an *in vivo*-application of the mutant cyclopeptides of the present invention has no negative impact on lung function, heart rate or blood pressure (Figs. 20 and 21). In addition, a number of important laboratory parameters to assess liver and kidney function were not influenced by the repeated cyclopeptide injections (Figs. 22a/b and 23). Therefore, the cyclic peptides of the present invention are, *inter alia*, particularly suitable for the treatment of distinct patient groups which otherwise could not be treated by using a beta blocker, i.e. patients who, for example, already suffer from bradycardia or for whom the use of beta blockers is not possible because of contraindications (like those suffering from obstructive lung disease or hypotension).

[0036] As mentioned, a further advantage of the means and methods of the present invention, particularly over means and methods taking advantage of (cyclic)peptides derived from  $\beta_1$ -EC<sub>II</sub> still having 3 cysteines (as, for example disclosed in WO 2006/103101), is that the formation of mixtures of cyclopeptide isomers can be avoided.

[0037] The biochemical characterization of a mixture of different cyclopeptide isomers, formed during cyclization of peptides comprising three or more Cys residues, is laborious. Accordingly, the production of pure cyclic peptide fractions containing only one sort of a cyclopeptide isomer is time and cost intensive, when taking advantage of peptides comprising three or more Cys residues. This is particularly true, when the cyclic peptides are produced under GMP standards.

[0038] In contrast thereto, the cyclic peptides of the present invention can easily be characterized and produced as pure fractions of the same isomer. This leads to a high reproducibility. The particular advantage of the peptides of the present invention is that mixtures of isomers, which have to be separated and must be characterized in laborious testings, are avoided, and that at least one further production step (separation and/or biochemical characterization) is finally omitted (see also Sewald 2002).

[0039] The present invention is, *inter alia*, based on the experiments described in the appended examples.

[0040] In context of these examples, one of the cysteines either at position 17 or at position 18 of the  $\beta_1$ -EC<sub>II</sub> 25AA-

cyclopeptide was replaced by a serine residue (Cys<sub>17</sub> or 18-Ser<sub>17</sub> or 18 mutation), so that only one individual, single intramolecular disulfide bond (S-S) can be formed (Fig. 1). Measures like this provides the potential to reduce side effects and to maintain or to increase the biological efficacy of the constructs of the present invention. The cyclic peptides of this invention can be obtained, in contrast to the peptides of the prior art which form mixtures of isomers, by simple, robust and highly reproducible manufacturing processes. These can be scaled up efficiently. Furthermore these processes avoids separation of isomers mixtures and are suitable for GMP standards. The appended examples provide for corresponding manufacturing/production methods.

**[0041]** In the appended examples, the cyclization of the inventive peptides was, inter alia, obtained by the introduction of a "DGLu" mutation, e.g. at the (ring) closure site of the cyclic peptide; Gln↔DGLu mutation as shown in Fig. 2).

**[0042]** Furthermore, the number of amino acids (AA) was reduced from 25AA to 22AA and further to 18AA in further sets of cyclopeptide-mutants of the present invention. This measure provides the potential to minimize the potential immunologic side effects of the constructs. The 18AA cyclopeptide-mutants contained a cysteine→serine exchange either at position 14 or at position 13 (18AA containing Cys<sub>13</sub>-Ser<sub>14</sub> or Ser<sub>13</sub>-Cys<sub>14</sub> mutant cyclopeptides, respectively), either combined with a (further) glutamine-exchange/D-glutamic acid, e.g. at the ring closure site of the cyclic peptide (Gln→D-Glu mutation), or not (Fig. 2). The 22AA cyclopeptide-mutants contained a cysteine→serine exchange at position 17 (22AA containing Cys<sub>16</sub>-Ser<sub>17</sub>), optionally combined with the introduction of a Gly residue at position 22 (a possible ring closure site of the cyclic peptide; Fig. 24).

**[0043]** Taken together, the herein provided experimental *in vitro* data as well as the *in vivo* data clearly demonstrate that the antibody-blocking capacity of the disclosed mutant cyclopeptides is not affected by the reduction of the number of amino acids from a 25-meric to a 18-meric cyclopeptide when using a dose ranging from 0,25 to 5,0 mg/kg body weight (Bw) and in particular from 1.0 to 2.0 mg/kg Bw. *In vitro* and *in vivo* data demonstrate an excellent comparability of the two 2 cysteine-containing single disulfide bond 25AA Cys/Ser (formulas VII/IX) or 18AA Cys/Ser (formulas VI/VIII) cyclopeptide mutants at a dose of 1.0 mg/kg Bw; Figs. 16 to 21. However, "intermediate" cyclic peptides of 19 to 24 AA exhibit an increased activity in accordance with this invention. Particularly, cyclic peptides of 22 AA exhibit an increased activity in accordance with this invention. A preferred example of such an "intermediate" cyclic peptide is a cyclic peptide comprising or consisting of the amino acid residues as shown in formula IX<sup>-</sup>.

**[0044]** Moreover, the exact nature of the exchange of one of the cysteine residues with a serine residue (i.e., Cys/Ser or Ser/Cys-mutation) markedly determined the potency of the disclosed cyclic peptides *in vitro* and also *in vivo* (Figs. 6-10 and 14-16).

**[0045]**  $\beta$ -adrenergic receptors ( $\beta$ -AR), particularly ( $\beta_1$ -adrenergic receptors ( $\beta_1$ -AR), are well known in the art. For example, the nucleotide and amino acid sequence (SEQ ID NO. 40) of the human  $\beta_1$ -AR (also known as adrenergic  $\beta_1$ -receptor (ADRB1)) can be obtained from databank entry NM\_000684 or NP\_000675.  $\beta$ -ARs are known to form two extracellular domains termed herein as EC<sub>I</sub> and EC<sub>II</sub> or  $\beta_{(1)}$ -EC<sub>I</sub> and  $\beta_{(1)}$ -EC<sub>II</sub>. As mentioned above, the cyclic peptides of the present invention share sequence similarity with  $\beta_1$ -EC<sub>II</sub>, particularly with the amino acid stretch DEARRCYNDP-KCCDFV (SEQ ID NO. 33) or RAESDEARRCYNDPKCCDFVTNR (SEQ ID NO. 34) of the human  $\beta_1$ -AR (amino acid positions 204 to 219 or 200 to 222, respectively) or, particularly, with the amino acid stretch DEARRCYNDPK (SEQ ID NO. 45) or ESDEARRCYNDPK (SEQ ID NO. 46) of the human  $\beta_1$ -AR. The term " $\beta$ -AR" as used herein preferably refers to a  $\beta_1$ -adrenergic receptor ( $\beta_1$ -AR), more preferably to the human  $\beta_1$ -AR as described above.

**[0046]** A cyclic peptide provided herein has at least one of the features selected from the group consisting of:

- a) being capable of binding (auto-)antibodies against the EC<sub>II</sub> loop of  $\beta_1$ -adrenergic receptor ( $\beta_1$ -AR);
- b) being capable of inhibiting the interaction between  $\beta_1$ -AR and (auto-)antibodies against the EC<sub>II</sub> loop of  $\beta_1$ -AR; and
- c) being capable of reducing an antibody-mediated activation of  $\beta_1$ -AR.

**[0047]** The structure of  $\beta_1$ -AR was, inter alia, analyzed by Wame (2008 Nature. DOI: 10. 1038).

**[0048]** As mentioned above, the cyclic peptide of the present invention is defined by the general formula cyclo(x-x<sub>n</sub>-Cys-x-x-x-x<sup>c</sup>-x-cys-y-x<sub>i</sub>-x) (formula I). In this formula, "y" may be any amino acid residue but Cys, preferably "y" may be any amino acid residue except Pro and except Cys. Generally, "y" may be any amino acid, as long as this amino acid does not form an intramolecular linkage (e.g. a disulphide bond) with another amino acid of the cyclic peptide provided herein (e.g. with another Cys of the cyclic peptide provided herein). Preferably, "y" may be any amino acid similar to Cys (i.e. having a similar chemical structure and/or a similar behavior within a 3 dimensional peptide structure), with the exception that it does not form an intramolecular linkage (e.g. a disulphide bond) with another amino acid of the cyclic peptide provided herein (e.g. with another Cys of the cyclic peptide provided herein). More preferably, "y" may be any polar amino acid but Cys, like Thr or Ser. Most preferably, in the cyclic peptide provided herein, "y" is Ser or a Ser analogue. "Ser analogue" in this context means a residue, particularly an amino acid residue, having a structural character similar to that of Ser. "Ser analogue" refers to, for example, a(n) (amino acid) residue having a similar chemical structure like that of Ser and/or a similar behavior within a 3 dimensional peptide structure like that of Ser. As a further example, "y" may also be selenocysteine or an analogue thereof.



**[0049]** In general, the meaning of terms like "any amino acid (residue) but Cys" or "amino acid (residue) other than Cys" is clear to the skilled person. Particularly, as used throughout this invention, such terms refer to any amino acid, as long as this amino acid does not form an intramolecular linkage (e.g. a disulphide bond) with another amino acid of the cyclic peptide provided herein (e.g. with another Cys of the cyclic peptide provided herein).

**[0050]** As mentioned, one main feature of the cyclic peptides of this invention is that they comprise only two Cys able to form an intramolecular linkage. Such cyclic peptides can, for example, be obtained by substituting a third Cys of a peptide homologous to the  $\beta_1$ -EC<sub>II</sub> by a different amino acid. Thereby, the Cys to be substituted is the one corresponding to the 2<sup>nd</sup> or, which is preferred, 3<sup>rd</sup> Cys of the  $\beta_1$ -EC<sub>II</sub> which lie in direct proximity to each other (amino acid position 215 and 216 of human  $\beta_1$ -AR (see also NP\_000675 and SEQ ID NO. 40). These two Cys residues, are referred to herein as "Cys-Cys", "Cys/Cys", "Cys<sub>215</sub>-Cys<sub>216</sub>" or "Cys<sub>215</sub>/Cys<sub>216</sub>" and the like).

**[0051]** The resulting mutant peptides or mutations as disclosed herein are accordingly termed as "Cys-Ser", "Cys/Ser", "Cys<sub>13, 16</sub> or 17-Ser<sub>14, 17</sub> or 18" or "Cys<sub>13, 16</sub> or 17/Ser<sub>14, 17</sub> or 18" mutant peptides or mutations or "Ser-Cys", "Ser/Cys", "Ser<sub>13</sub> or 17-Cys<sub>14</sub> or 18" or "Ser<sub>13</sub> or 17/Cys<sub>14</sub> or 18" mutant peptides or mutations, depending which of the Cys is replaced and how many amino acids the mutant peptide comprises.

**[0052]** Alternatively, the mutant peptides as disclosed herein are defined by referring to the particular amino acid exchanges at a certain position. Then, the mutant peptides/mutations are, for example, termed "Cys<sub>14, 17</sub> or 18→Ser<sub>14, 17</sub> or 18" mutant peptides/mutations or "Cys<sub>13</sub> or 17→Ser<sub>13</sub> or 17" mutant peptides/mutations, depending on whether the Cys corresponding to Cys<sub>216</sub> or the Cys corresponding to the Cys<sub>215</sub>, respectively, of  $\beta_1$ -AR is replaced by a different amino acid. The indices "14, 17 or 18" or "13 or 17" relate to the position in the exemplified cyclic peptide of the invention, whereby position 1 corresponds to the first "x" as defined in formula I, i.e. cyclo(x-x<sub>h</sub>-Cys-x-x-x<sup>c</sup>-x-Cys-y-x<sub>i</sub>-x).

**[0053]** Accordingly, terms like "Cys<sub>13</sub>-Ser<sub>14</sub>" or "Cys<sub>13</sub>/Ser<sub>14</sub>" mutant peptides/mutations are used in the same sense as "Cys<sub>14</sub>→Ser<sub>14</sub>" mutant peptides/mutations and, in this particular example, refer to 18mer peptides disclosed herein. Terms like "Cys<sub>16</sub>-Ser<sub>17</sub>" or "Cys<sub>16</sub>/Ser<sub>17</sub>" mutant peptides/mutations are used in the same sense as "Cys<sub>17</sub>→Ser<sub>17</sub>" mutant peptides/mutations and, in this particular example, refer to 22mer peptides disclosed herein. Terms like "Cys<sub>17</sub>-Ser<sub>18</sub>" or "Cys<sub>17</sub>/Ser<sub>18</sub>" mutant peptides/mutations are used in the same sense as "Cys<sub>18</sub>→Ser<sub>18</sub>" mutant peptides/mutations and, in this particular example, refer to 25mer peptides disclosed herein.

**[0054]** Analogously, terms like "Ser<sub>13</sub>-Cys<sub>14</sub>" or "Ser<sub>13</sub>/Cys<sub>14</sub>" mutant peptides/mutations are used in the same sense as "Cys<sub>13</sub>→Ser<sub>13</sub>" mutant peptides/mutations and, in this particular example, refer to 18mer peptides disclosed herein, and terms like "Ser<sub>17</sub>-Cys<sub>18</sub>" or "Ser<sub>17</sub>/Cys<sub>18</sub>" mutant peptides/mutations are used in the same sense as "Cys<sub>17</sub>→Ser<sub>17</sub>" mutant peptides/mutations and, in this particular example, refer to 25mer peptides disclosed herein.

**[0055]** The exemplarily indices given above refer to the position of the indicated amino acid within the herein disclosed particular 18mer, 22mer or 25mer peptide, respectively. In context of this invention, the starting point with respect to an indicated amino acid position given for a cyclic peptide disclosed herein is the N-terminal amino acid of the linearized backbone the cyclic peptide (like the first "x" in formula I, see above). The starting point with respect to an indicated amino acid position given for a linear peptide disclosed herein is its N-terminal amino acid.

**[0056]** The findings as provided herein and in the appended examples demonstrate a comparability of 25AA, 22AA and 18AA cyclopeptides without any Cys mutation with the cyclic 25AA, 22AA or 18AA Cys<sub>18, 17</sub> or 14→Ser<sub>18, 17</sub> or 14 (Cys-Ser) mutants, but not with the cyclic 25AA or 18AA Cys<sub>17</sub> or 13→Ser<sub>17</sub> or 13 (Ser-Cys) mutants.

**[0057]** As also mentioned above, in the formulas of the cyclic peptide of the present invention, h can be any integer from 1 to 15, preferably from 5 to 9, and/or i can be any integer from 0 to 14, preferably from 1 to 14, more preferably from 0 to 6 and even more preferably from 1 to 6. Accordingly, h can be 1, 2, 3, 4, 5, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 and/or i can be 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14. Preferably, h is 5, 8 or 9 and/or i is 3, 4 or 6. More preferably, h is 8 and/or i is 4. In particularly preferred embodiments of this invention, x<sub>h</sub> stands for the particular amino acid stretches DEARR (SEQ ID NO. 35), AESDEARR (SEQ ID NO. 47) or RAESDEARR (SEQ ID NO. 36) and/or x<sub>i</sub> stands for the particular amino acid stretches DFV (SEQ ID NO. 37), DFVT (SEQ ID NO. 48) or DFVTNR (SEQ ID NO. 38). In more preferred particular embodiments of this invention, x<sub>h</sub> stands for the particular amino acid stretch AESDEARR (SEQ ID NO. 47) and/or x<sub>i</sub> stands for the particular amino acid stretch DFVT (SEQ ID NO. 48).

**[0058]** The cyclic peptide of the present invention (or the cyclic part thereof) may consist of at least 18 amino acids and of at most 25 amino acids. Accordingly, the cyclic peptide of the present invention may consist of 18, 19, 20, 21, 22, 23, 24 or 25 amino acids, whereby particularly 18 or 25 amino acids are preferred and particularly 22 amino acids are most preferred. In a less preferred embodiment, also smaller peptides, i.e. peptides comprising 16 or 17 amino acids are envisaged.

**[0059]** A particularly preferred cyclic peptide in context of this invention is one of (21+1=)22AA length, having a defined maximum and minimum size of the cyclic molecule dependent on the respective amino-acid composition, constituted by 21 amino-acids from the original primary sequence of the human betal-adrenergic receptor (i.e., amino-acids 200 to 221; Frielle 1987, PNAS 84, 7920-7924) with an additional glycine as 22<sup>nd</sup> amino-acid at the assumed ring closure site (position 222).

**[0060]** Without being bound by theory, the number of amino acids and thus the length of the primary structure (i.e. the amino acid backbone) of cyclic peptides binding anti- $\beta_1$ -AR antibodies is crucial for their biological effects and/or successful/effective manufacture.

**[0061]** A peptide-length equal or above 26 amino acids (primary structure) may be stimulating directly (that is, without the use of carrier proteins) immunocompetent T-cells and thus may provoke an undesired paradoxal increase of anti- $\beta_1$ -receptor antibody production through T-cell mediated B-cell stimulation.



**[0062]** A peptide-length below 16 amino acids (primary structure) leads to undesired crystallization during the production process and problems in dissolving the synthesized products in an aqueous solution, e.g. for purposes of i.v. or s.c. injections

**[0063]** In a less preferred embodiment of this invention also cyclic peptides falling under the above given definitions a) to e) of formula I and consisting of only 16 amino acids or, even less preferred, consisting of only 17 amino acids are provided. A non-limiting example of such a less preferred cyclic peptide is the peptide cyclo(Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Tyr-Gln/DGlu) (formable by an amino acid backbone as depicted in SEQ ID NO. 39).



**[0064]** It is particularly preferred herein that the disclosed cyclic peptide contains only one Pro. Accordingly, it is particularly preferred that neither the y nor an x of the formulas depicted herein, except of  $x^c$ , is Pro. Within the amino acid stretch,  $x^c$  as depicted in formula I (or other formulas) is Pro.

**[0065]** It is particularly envisaged herein that the acidic amino acid Asp precedes the Pro contained in the cyclic peptide of the invention. Accordingly it is preferred that x as depicted in formula I (or other formulas) which precedes  $x^c$  is the acidic amino acid Asp.

**[0066]** More specifically, the cyclic peptide of the present invention may be defined by formula I' or I'':



**[0067]** Even more specifically, the cyclic peptide of the present invention may be defined by formula I''' or I''':



**[0068]** In general,  $x_I$  and  $x_{II}$  as depicted in formula I' and I'' (and in the other formulas depicted herein) may, as mentioned, any amino acid but Cys. However, particularly when the ring closure of the cyclic peptides of the invention occurs between  $x_I$  and  $x_{II}$ , it is particularly envisaged that  $x_I$  and  $x_{II}$  are such amino acids able to form a peptide bond, or the like, with each other under conditions of a "head to tail" cyclization. "Head to tail" cyclizations are known in the art (e.g. Kates and Albericio: Solid phase synthesis, CRC-Press, 2000; Williams, Chemical Approaches to the Synthesis of Peptides, CRC-Press 1997; Benoiton: Chemistry of Peptide Synthesis. CRC-Press, 2005) and examples thereof are given in the experimental part. Possible examples of amino acids that may be  $x_I$  are Gly, Val, Thr, Ser and, preferably, Ala. Possible examples of amino acids that may be  $x_{II}$  are Glu and, preferably, Gln. Less preferred,  $x_{II}$  may also be Asp or Asn. Most preferred,  $x_I$  is Ala and  $x_{II}$  is Gln or Glu (preferably DGlu).

**[0069]** Accordingly, in the cyclic peptides of this invention  $x_{II}$  as referred to in formula I' and I''' can be Gln or Glu, wherein Glu may also be DGlu (D-Glu; D-Glutamic acid). However, naturally amino acids are preferred herein. Therefore, it is more preferred that  $x_{II}$  is Gln.

**[0070]** The skilled person is able chose amino acid residues appropriate to be  $x_I$  and/or  $x_{II}$  of formula I' and I''' in accordance with this invention on the basis of the teaching provided herein and his knowledge of the art (e.g. Williams, Chemical Approaches to the Synthesis of Peptides, CRC-Press 1997; Benoiton: Chemistry of Peptide Synthesis. CRC-Press, 2005).

**[0071]**  $x_{III}$  and  $x_{IV}$  as depicted in formula I'' and I''' (and in the other formulas depicted herein) may, as mentioned, any amino acid but Cys. However; particularly when the ring closure of the cyclic peptides of the invention occurs between  $x_{III}$  and  $x_{IV}$ , it is particularly envisaged that  $x_{III}$  and  $x_{IV}$  are such amino acids able to form a peptide bond, or the like, with each other under conditions of a "head to tail" cyclization. A possible example of an amino acid that may be  $x_{III}$  is Arg. One possible, and most preferred, example of an amino acid that may be  $x_{IV}$  is Gly or a Gly analogue. "Gly analogue" in this context means a residue, particularly an amino acid residue, having a structural character similar than that of Gly. Particularly, "Gly analogue" refers to, for example, a(n) (amino acid) residue having the same (or even a smaller) size than a Gly residue. It was surprisingly found in context of this invention that particularly a small (amino acid) residue like Gly at the " $x_{IV}$ " position leads to an improved mimicking of the ECII of  $\beta_1$ -AR by the corresponding cyclic peptides of the invention. The skilled person is able chose amino acid residues appropriate to be  $x_{III}$  and/or  $x_{IV}$  of formula I'' and I''' in accordance with this invention on the basis of the teaching provided herein and his knowledge of the art (e.g. Williams,

Chemical Approaches to the Synthesis of Peptides, CRC-Press 1997; Benoiton: Chemistry of Peptide Synthesis. CRC-Press, 2005).

**[0072]** It is further particularly preferred herein that the cyclic peptides of this invention lack Trp and/or His. Accordingly, it is particularly envisaged in context of the invention, that neither an x nor y as depicted in any of Formula I to I'" is Trp or His.

**[0073]** Furthermore, it is preferred that the provided cyclic peptides lack sites susceptible for hydrolysis or cleaving proteases, like, for example, serum proteases. The meanings - of the terms "hydrolysis" and "(serum) proteases" are well known in the art.

**[0074]** A peptide as provided herein can also be described as a peptide consisting of or comprising an amino acid sequence homologous to SEQ ID NO. 33 (representing a wild type amino acid stretch comprising epitopes of  $\beta_1$ -EC<sub>II</sub>), wherein (a) the amino acid corresponding to position 13 (or, less preferred, corresponding to position 12) of SEQ ID NO. 33 is not Cys and the amino acid corresponding to positions 6 and 12 (or, less preferred, corresponding to position 6 and 13) of SEQ ID NO. 33 is Cys, (b) wherein said amino acid sequence contains no further Cys able to form an intramolecular linkage within the peptide, i.e. within that part of the peptide being homologous to SEQ ID NO. 33, and wherein the peptide can function as a cyclic peptide in accordance with this invention, e.g. is able to block anti- $\beta$ -AR antibodies, or wherein the peptide can form such a cyclic peptide. Optionally, the further provisions given herein with respect to the structure of the disclosed linear and/or cyclic peptides apply here, mutatis mutandis. The so defined peptide consists of a stretch of 16 amino acids being homologous to SEQ ID NO. 33 flanked at the N- and C-terminus by one or more amino acids, preferably naturally occurring amino acids, like the "x<sub>I</sub>"/"x<sub>III</sub>" at position 1 and the "x<sub>II</sub>"/"x<sub>IV</sub>" at the last position of formulas I' to I'" given herein.

**[0075]** In context of the invention, and in particular in context of the (wild type) SEQ ID NO. 33, "homologous" means identical on amino acid level for at least 18,75%, at least 37,5%, at least 50%, at least 56,25%, at least 62,5%, at least 68,75%, at least 75%, at least 81,25%, at least 87,5% or 93,75%, wherein the higher values are preferred.




**[0076]** In general, the meaning of the term "amino acid" or "amino acid residue" is known in the art and is used herein accordingly. Thereby, it is of note that when an "amino acid" is a component of a peptide/protein the term "amino acid" is used herein in the same sense than "amino acid residue".

**[0077]** Particularly, an "amino acid" or "amino acid residue" as referred to herein is preferably envisaged to be a naturally occurring amino acid, more preferably a naturally occurring L-amino acid (except the above mentioned D<sub>Glu</sub>). However, albeit less preferred, an "amino acid" or "amino acid residue" in context of this invention may also be a non-naturally occurring (i.e. a synthetic) amino acid, like, for example, norleucine or  $\beta$ -alanine, or, particularly in case of "y" of the formulas depicted herein, selenocysteine or an analog thereof.

**[0078]** Also known in the art is the meaning of the terms "acidic amino acid(s)", "basic amino acid(s)", "aliphatic amino acid(s)" and "polar amino acid(s)" (for example, Stryer, Biochemie, Spectrum Akad. Verlag, 1991, Item I. 2.). These terms are correspondingly used throughout this invention. Thereby, the particular provisos given herein with respect to the cyclic peptides of the invention also apply.

**[0079]** Particularly, the term "acidic amino acid(s)" as used herein is intended to mean an amino acid selected from the group comprising Asp, Asn, Glu, and Gln, preferably Asp and Glu; the term "basic amino acid(s)" as used herein is intended to mean an amino acid selected from the group comprising Arg, Lys and His, preferably Arg and Lys; the term "aliphatic amino acid(s)" as used herein is intended to mean any amino acid selected from the group comprising Gly, Ala, Ser, Thr, Val, Leu, Ile, Asp, Asn, Glu, Gln, Arg, Lys, Cys and Met; and the term "polar amino acid(s)" as used herein is intended to mean any amino acid selected from the group comprising Cys, Met, Ser, Tyr, Gln, Asn and, less preferred, Trp.


**[0080]** In a more general aspect, the cyclic peptide as disclosed herein may be a cyclic peptide of formula II, III or III':



wherein

- a) x<sub>1</sub> is individually and independently selected from the group consisting of acidic amino acids; and/or
- b) x<sub>2</sub> is individually and independently selected from the group consisting of basic amino acids.

**[0081]** In a more specific aspect, the cyclic peptide as disclosed herein may be a cyclic peptide of formula IV, V or V':



cyclo(x<sub>1</sub>-x<sub>2</sub>-x<sub>4</sub>-x<sub>1</sub>-x<sub>4</sub>-x<sub>1</sub>-x<sub>4</sub>-x<sub>2</sub>-x<sub>2</sub>-Cys-x<sub>3</sub>-x<sub>5</sub>-x-x<sup>c</sup>-x<sub>2</sub>-Cys-y-x<sub>1</sub>-x<sub>3</sub>-x<sub>3</sub>-x<sub>4</sub>-x<sub>5</sub>-x<sub>2</sub>-x<sub>II</sub>) (V);

cyclo(x<sub>III</sub>-x<sub>4</sub>-x<sub>1</sub>-x<sub>4</sub>-x<sub>1</sub>-x<sub>4</sub>-x<sub>2</sub>-x<sub>2</sub>-Cys-x<sub>3</sub>-x<sub>5</sub>-x-x<sup>c</sup>-x<sub>2</sub>-Cys-y-x<sub>1</sub>-x<sub>3</sub>-x<sub>3</sub>-x<sub>4</sub>-x<sub>IV</sub>) (V'),

5 wherein

- a) x<sub>1</sub> is individually and independently selected from the group consisting of acidic amino acids;
- b) x<sub>2</sub> is individually and independently selected from the group consisting of basic amino acids;
- c) x<sub>3</sub> is individually and independently selected from the group consisting of Leu, lie, Val, Met, Trp, Tyr and Phe;
- d) x<sub>4</sub> is individually and independently selected from the group consisting of Ser, Thr, Ala and Gly; and/or
- e) x<sub>5</sub> is individually and independently selected from the group consisting of Gln and Asn.

15 **[0082]** In a further aspect, the cyclic peptides comprise an amino acid stretch as defined by amino acid position 2-12 or 2-14 of formula II or IV, an amino acid stretch as defined by amino acid position 4-16 or 4-18 of formula III or V or an amino acid stretch as defined by amino acid position 3-15 or 3-17 of formula or III' or V'. In a more general aspect, the cyclic peptide as disclosed herein may be a cyclic peptide of formula II, III or III'

**[0083]** In a further particular aspect, the cyclic peptide as disclosed herein may comprise the amino acid stretch  
 aci-Glu-Ala-bas-bas-Cys-Tyr-neu-aci-neu-bas;  
 aci-neu-aci-Glu-Ala-bas-bas-Cys-Tyr-neu-aci-neu-bas;  
 20 aci-Glu-Ala-bas-bas-Cys-Tyr-neu-aci-neu-bas-Cys-Ser; or  
 aci-neu-aci-Glu-Ala-bas-bas-Cys-Tyr-neu-aci-neu-bas-Cys-Ser,  
 wherein "aci" stands for acidic amino acid, "neu" stands for neutral amino acid and "bas" stands for basic amino acid. Each amino acid residue of the above two amino acid stretches may also be defined independently as the corresponding amino acid residue of any one of formulas I, II, III, III', IV, V, and V' as provided herein.

25 **[0084]** The cyclic peptide as provided herein comprises the amino acid stretch  
 Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 45) or  
 Gtu-Ser-Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 46),  
 wherein Xxx<sub>1</sub> is defined as "x" or "x<sub>1</sub>", Xxx<sub>3</sub> is defined as "x" or "x<sub>3</sub>" and/or Xxx<sub>4</sub> is defined as "x" or "x<sub>4</sub>" as mentioned in the above depicted formulas. For example, the above-mentioned amino acid stretch may be  
 30 Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys (SEQ ID NO. 45) or  
 Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys (SEQ ID NO. 46).

**[0085]** It is particularly envisaged herein that the cyclic peptides of this invention comprise one or more epitopes beared by β<sub>1</sub>-EC<sub>II</sub>, like, for example, epitopes beared by any of the above mentioned amino acid stretches (or by parts of the disclosed cyclic peptides comprising these amino acid stretches). In this context, the term "epitope" particularly refers to an amino acid stretch to which an (auto)anti-β<sub>1</sub>-AR antibody binds. Particularly, an epitope in context of this invention consists of at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13 or at least 14 amino acids. The person skilled in the art is in the position to deduce which particular amino acid residue(s) of β<sub>1</sub>-EC<sub>II</sub> contribute to (an) epitope(s) to which (auto)anti-β<sub>1</sub>-AR antibodies bind. Hence, he can deduce which particular amino acid residues have at least to be comprised in the cyclic peptides of this invention in order to ensure that these peptide bind (auto)anti-β<sub>1</sub>-AR antibodies. For this purpose, several means and methods known in the art can be employed (for example, means and methods for epitope mapping (like PepSpots™, Biacore, Amino acid scans (like alanine scans)).

**[0086]** Non limiting examples of a cyclic peptide according to this invention are cyclic peptides selected from the group consisting of:

- a) cyclic peptides formable or formed by the amino acid sequence as depicted in any one of SEQ ID NO. 41, 43, 1 to 4 and 17 to 20;
- b) cyclic peptides formable or formed by an amino acid sequence as encoded by a nucleotide sequence as depicted in any one of SEQ ID NO. 42, 44, 9 to 12, 25 to 28, 49, 50, 53 and 54; and
- 50 c) cyclic peptides formable or formed by an amino acid sequence as encoded by a nucleotide sequence which differs from the nucleotide sequence as depicted in any one of SEQ ID NO. 42, 44, 9 to 12, 25 to 28, 49, 50, 53 and 54 due to the degeneracy of the genetic code.

**[0087]** According to this invention, those cyclic peptides being Cys-Ser mutant peptides, i.e. having the Cys corresponding to the third Cys of the β<sub>1</sub>-EC<sub>II</sub> (the Cys at position 216 of β<sub>1</sub>-AR) exanged by Ser, are particularly preferred. The above given examples refer to such particularly preferred cyclic peptides. As demonstrated in the appended examples, such cyclic peptides are particularly useful in inhibiting or diagnosing anti-β<sub>1</sub>-AR antibodies.

**[0088]** The particular structure of the above exemplified particularly preferred cyclic peptides is given by any one of

the following formulas VI to IX':

cyclo(Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Gly) (IX');

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Gln) (VI);

cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-Gln) (VII);

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-DGlu) (VIII); and

cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-DGlu) (IX).

**[0089]** Non limiting examples of less preferred cyclic peptides are cyclic peptides selected from the group consisting of:

- a) cyclic peptides formable or formed by the amino acid sequence as depicted in any one of SEQ ID NO. 5 to 8 and 21 to 24;
- b) cyclic peptides formable or formed by an amino acid sequence as encoded by a nucleotide sequence as depicted in any one of SEQ ID NO. 13 to 16 and 29 to 32; and
- c) cyclic peptides formable or formed by an amino acid sequence as encoded by a nucleotide sequence which differs from the nucleotide sequence as depicted in any one of SEQ ID NO. 29 to 32 due to the degeneracy of the genetic code.

**[0090]** The particular structure of the above exemplified less preferred cyclic peptides is given by any one of the following formulas X to XIII:

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Ser-Cys-Asp-Phe-Val-Gln) (X);

cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Ser-Cys-Asp-Phe-Val-Thr-Asn-Arg-Gln) (XI);

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Ser-Cys-Asp-Phe-Val-DGlu) (XII);

cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Ser-Cys-Asp-Phe-Val-Thr-Asn-Arg-DGlu) (XIII).

**[0091]** The above peptides are less preferred since peptides as the "Cys-Ser" mutant peptides ("Cys-Ser" cyclic peptides) are, inter alia, in vivo more functional than the herein defined less preferred "Ser-Cys" mutant peptides ("Ser-Cys" cyclic peptides).

**[0092]** In this context it is of note that most preferred examples of the cyclic peptides according to this invention are particularly those cyclic peptides, the pharmacological and/or diagnostic function of which has been demonstrated in the appended examples (e.g. those characterized by any one of formula VI to IX').

**[0093]** It will be understood that for the various peptides of the present invention a certain flexibility and variability in the primary sequence, i. e. the amino acid sequence backbone, is possible as long as the overall secondary and tertiary structure of the respective peptides, which is defined by at least some fixed amino acid residues and by their spatial arrangement, is ensured (see, e.g., formula I, supra).

**[0094]** Based on the teaching provided herein, the skilled person is, on the one hand, readily in the position to find out corresponding variants of the peptides of the invention. On the other hand, the skilled person is able to test whether a given variant of peptides of the present invention still has the desired function, for example the ability to specifically bind to  $\beta_1$ -AR antibodies, and therefore has the potential for a corresponding medical intervention, like the therapeutic and diagnostic applications described and provided herein. Corresponding experimental guidance for such tests, i.e. respective assays, are exemplarily provided and described herein, particularly in the appended examples.

**[0095]** Accordingly, also disclosed herein are variants of the herein disclosed and described peptides, given that, first, these variants are still functionally active in accordance with this invention, i. e. functionally active as binding partners for anti- $\beta_1$ -AR antibodies, particularly for anti- $\beta_1$ -AR antibodies against the  $\beta_1$ -EC<sub>II</sub>, more particularly functionally active as inhibitors of  $\beta_1$ -AR and even more preferably active in inhibiting the interaction between  $\beta_1$ -AR and anti- $\beta_1$ -AR antibodies against the  $\beta_1$ -EC<sub>II</sub>, more preferably auto-anti- $\beta_1$ -AR antibodies against the  $\beta_1$ -EC<sub>II</sub>; and, second, that these variants are not present in form of isomers mixtures or do not form isomers mixtures when cyclised in accordance with a production method as disclosed herein. These variants are envisaged to have only two certain Cys residues forming or being able to form only one individual intramolecular linkage (e.g. disulphide bond).

**[0096]** Within the variants of the peptides of the present invention it is, for example, envisaged that one or more amino acids of said peptides are replaced by other one or more naturally occurring or synthetic amino acids. In this context, it is preferred that this/these amino acid exchange(s) is/are (a)conservative amino acid exchange(s); i.e. that the replacement amino acid belongs to the same category of amino acids than the amino acid to be replaced. For example, an acidic amino acid may be replaced by another acidic amino acid, a basic amino acid may be replaced by another basic amino acid, an aliphatic amino acid may be replaced by another aliphatic amino acid, and/or a polar amino acid may be replaced by another polar amino acid.

**[0097]** Accordingly, particularly preferred and provided variants of the (cyclo) peptides of the present invention are variants wherein at least one of an acidic amino acid is replaced by a different amino acid selected from the group consisting of acidic amino acids, at least one of the basic amino acids is replaced by a different amino acid selected from the group consisting of basic amino acids, at least one of a polar amino acid is replaced by a different amino acid selected from the group consisting of polar amino acids and/or at least one of an aliphatic amino acid is replaced by a different amino acid selected from the group consisting of aliphatic amino acids (given that the above mentioned-requirements are fulfilled).

**[0098]** It is particularly envisaged that the amino acid exchanges which lead to variants of the disclosed (cyclic) peptides are such, that the pattern of polarity and charge within the tertiary structure of the resulting variant still substantially mimics the three-dimensional structure of the corresponding EC<sub>II</sub> epitope(s) of  $\beta_1$ -AR.

**[0099]** With respect to the "Variants" of the (cyclo) peptides of the present invention the herein defined Cys may also be replaced by other amino acids, as long as the replacement still leads to an individual intramolecular linkage, like that of a disulphide bond, within the cyclopeptide, i.e. the avoidance of isomers mixtures formation during cyclization and/or a correct mimicry of the EC<sub>II</sub> of  $\beta_1$ -AR. Such amino acid may, inter alia, be a non-naturally occurring amino acid, like a non-naturally occurring amino acid having an -SH group able to form a disulphide bond. However, it is preferred herein that the Cys given in formula I, above, is indeed a naturally occurring amino acid, preferably Cys itself.

**[0100]** It will also be acknowledged by the one skilled in the art that one or several of the amino acids forming the (cyclic) peptide of the present invention may be modified. In accordance therewith any amino acid as used herein may also represent its modified form. For example, an alanine residue as used herein may comprise a modified alanine residue. Such modifications may, among others, be a methylation or acylation or the like, whereby such modification or modified amino acid is preferably comprised by the present invention as long as the thus modified amino acid and more particularly the peptide containing said thus modified amino acid is still functionally active as defined herein, like functionally active as an inhibitor of anti- $\beta_1$ -AR antibodies, preferably active in inhibiting the interaction between  $\beta_1$ -AR and antibodies, and more preferably auto-antibodies directed against  $\beta_1$ -AR. Respective assays for determining whether such a peptide, i. e. a peptide comprising one or several modified amino acids, fulfils this requirement, are known to the one skilled in the art and, among others, also described herein, particularly in the example part hereof.

**[0101]** The invention also provides derivatives of the disclosed (cyclic) peptides such as salts with physiological organic

and anorganic acids like HCl, H<sub>2</sub>SO<sub>4</sub>, H<sub>3</sub>PO<sub>4</sub>, malic acid, fumaric acid, citronic acid, tatratic acid, acetic acid.

**[0102]** As used herein, the sequences of the peptides disclosed are indicated from the N-terminus to the C-terminus, whereby the N-terminus is at the left side and the C-terminus is at the right side of the respective depicted amino acid sequence. When referring to cyclic peptides, the corresponding sequences are indicated from the side corresponding to the left side of formula I to the side corresponding to the right side of formula I.

**[0103]** A "cyclic peptide" or "cyclopeptide" and the like in accordance with the present invention is a peptide intramolecularly forming a molecular ring structure within its amino acid backbone/primary amino acid sequence by at least one, preferably by at least two, more preferably by exactly two intramolecular linkages having covalent character. The forming of this molecular ring structure is, in context of this invention, also termed "cyclization".

**[0104]** In one particularly preferred embodiment, the cyclic peptide of this invention has two intramolecular linkages having covalent character, wherein one of these linkages is an intramolecular linkage between the N- and C-terminal ends of a peptide being the amino acid backbone/primary amino acid sequence of the cyclic peptide disclosed and the other one is an intramolecular linkage between two non-terminal amino acids of this peptide. As mentioned, these two non terminal amino acids may be two Cys.

**[0105]** Generally, "cyclization" in accordance with this invention may occur by at least one linkage which is a covalent binding selected from the group consisting of S-S linkages, peptide bonds, carbon bonds such as C-C or C=C, ester bonds, ether bonds, azo bonds, C-S-C linkages, C-N-C linkages and C=N-C linkages.

**[0106]** Particularly, the peptide bond as mentioned throughout this invention can be formed by the NH<sub>2</sub> group of an N-terminal amino acid and the COOH group of an C-terminal amino acid of a peptide forming the amino acid backbone/primary amino acid sequence of the cyclic peptide disclosed.

**[0107]** Preferably, an intramolecular linkage between the N- and C-terminal ends of a peptide forming the amino acid backbone/primary amino acid sequence of the cyclic peptide disclosed is a peptide bond and an intramolecular linkage between two non-terminal amino acids of this peptide is an S-S linkage (i. e. disulphide bond).

**[0108]** In context of this invention, an intramolecular S-S linkage within the cyclic peptide provided can be formed between two Cys residues within the amino acid backbone/primary amino acid sequence of said cyclic peptide.

**[0109]** Within the cyclic peptides of this invention, not only the above mentioned two particular intramolecular covalent linkage may be formed but also further intramolecular linkages may occur, with the proviso that the herein described functionality of the cyclic peptides is maintained and that the cyclic peptides can still easily be characterized biochemically, which, e.g., means that no isomers mixtures are formed during cyclization of the corresponding amino acid backbone/primary amino acid sequence.

**[0110]** For example, such further intramolecular linkages are additional bonds formed by a side chain of NH<sub>2</sub> groups and COOH groups of the constituent amino acids.

**[0111]** Terms like "amino acid backbone" or "primary amino acid sequence" as used throughout the present invention refer, on the one hand, to that structural component or part of a cyclic peptide which is formable or formed by its corresponding amino acid sequence. On the other hand, these terms refer to the linear peptides able to form the cyclic peptides of this invention by cyclization.

**[0112]** In one particular aspect, a cyclic peptide is disclosed, which is obtainable by the method for producing a cyclic peptide as provided herein. The definitions given herein-above also apply with respect to this particularly disclosed cyclic peptide.

**[0113]** In one aspect, also such peptides are disclosed, the disclosed cyclic peptides are formable by or are formed by. Particularly these peptides are the linear peptides forming or being able to form the herein disclosed cyclic peptides, i.e. the amino acid backbone/primary amino acid sequence thereof.

**[0114]** In general, such a linear peptide can be any peptide, the covalent linkage of the N- and C-terminus of which results in a cyclic peptide as disclosed in accordance with the present invention. For example, such a linear peptide may be some kind of an intermediate compound in an procedure of producing the cyclic peptides of this invention, like the method for producing a cyclic peptide as disclosed herein.

**[0115]** In general, the N- and C-terminal end of a linear peptide disclosed herein may be any amino acid pair lying in direct proximity to each other within the amino acid backbone of a cyclic peptide disclosed in context of this invention. In other words, cyclization (ring closure) of the cyclic peptide of this invention may generally occur between any of said amino acid pairs. The skilled person is readily in the position to find out such particular amino acid pairs which are effective/suitable to act as N- and C-terminal ends of a herein disclosed linear peptide, i.e. which are effective/suitable to act as an amino acid pair being involved in the ring closure/cyclization as defined in context of this invention.

**[0116]** In one preferred but non-limiting example, the cyclization (ring closure) of a linear peptide of this invention may occur between Ala and Gln or Glu, i.e. the N-terminal amino acid of this linear peptide would be Ala and the C-terminal amino acid would be Gln or Glu. Examples of such linear peptides able to form the cyclic peptide of the present invention are SEQ ID NO. 1 to 4 and, less preferred SEQ ID NO. 5 to 8.

**[0117]** In another preferred but non-limiting example, the cyclization (ring closure) of a linear peptide of this invention may occur between Lys and Pro, i.e. the N-terminal amino acid of this linear peptide would be Lys and the C-terminal

amino acid would be Pro. Examples of such linear peptides able to form the cyclic peptide of the present invention are SEQ ID NO. 17 to 20 and, less preferred SEQ ID NO. 21 to 24.

**[0118]** In a more preferred but non-limiting example, particularly when a 22mer cyclic peptide is provided, the cyclization (ring closure) of a linear peptide as disclosed herein may occur between Arg and Gly, i.e. the N-terminal amino acid of this linear peptide would be Arg and the C-terminal amino acid would be Gly. An examples of such a linear peptide able to form the cyclic peptide of the present invention is SEQ ID NO. 41.

**[0119]** In another more preferred but non-limiting example, particularly when a 22mer cyclic peptide is provided, the cyclization (ring closure) of a linear peptide as disclosed herein may occur between Lys and Pro, i.e. the N-terminal amino acid of this linear peptide would be Lys and the C-terminal amino acid would be Pro. An examples of such a linear peptide able to form the cyclic peptide of the present invention is SEQ ID NO. 43.

**[0120]** Besides their amino acid backbone, the cyclic peptides of the invention may further comprise (e.g. have covalently bound) (a) further substituent(s), like labels, anchors (like proteinaceous membrane anchors), tags (like HIS tags) and the like. Appropriate substituents and methods for adding them to the cyclic peptide of this invention are known to those of ordinary skill in the art.

**[0121]** Examples of labels in this context include, inter alia, fluorochromes (like fluorescein, rhodamine, Texas Red, etc.), enzymes (like horse radish peroxidase,  $\beta$ -galactosidase, alkaline phosphatase), radioactive isotopes (like  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{35}\text{S}$ ,  $^{125}\text{I}$  or  $^{123}\text{I}$ ,  $^{135}\text{I}$ ,  $^{124}\text{I}$ ,  $^{11}\text{C}$ ,  $^{15}\text{O}$ ), biotin, digoxigenin, colloidal metals, chemi- or bioluminescent compounds (like dioxetanes, luminol or acridiniums). One particularly envisaged label that may be bound to the peptide of this invention is a fluorochrome belonging to a FRET pair of fluorochromes, for example a GFP variant (e.g. GFP, eGFP, EYFP or ECFP).

**[0122]** A variety of techniques are available for labeling biomolecules, are well known to the person skilled in the art and are considered to be within the scope of the present invention and comprise, inter alia, covalent coupling of enzymes or biotinyl groups, phosphorylations, biotinylations, random priming, nick-translations, tailing (using terminal transferases). Such techniques are, e.g., described in Tijssen, "Practice and theory of enzyme immunoassays", Burden and von Knippenburg (Eds), Volume 15 (1985); "Basic methods in molecular biology", Davis LG, Diber MD, Battey Elsevier (1990); Mayer, (Eds) "Immunochemical methods in cell and molecular biology" Academic Press, London (1987); or in the series "Methods in Enzymology", Academic Press, Inc. Detection methods comprise, but are not limited to, autoradiography, fluorescence microscopy, direct and indirect enzymatic reactions, etc.

**[0123]** The substituent(s) can be bound (e.g. covalently) to the cyclic peptides of the invention directly or via linkers. The skilled person is readily in the position to find out appropriate linkers to be employed in this context.

**[0124]** In a further aspect, disclosed herein is a nucleic acid molecule comprising a nucleotide sequence encoding the amino acid backbone/primary amino acid sequence of a cyclic peptide as disclosed in context of this invention.

**[0125]** Further disclosed herein is a nucleic acid molecule comprising a nucleotides sequence encoding the linear peptides as described herein.

**[0126]** For example, such nucleic acid molecule may comprise a nucleotide sequence as depicted in any one of SEQ ID NO. 42, 44, 9 to 12, 25 to 28, 49, 50, 53 and 54 or, less preferred, SEQ ID NO. 13 to 16 and 29 to 32 or a nucleotide sequence which differs therefrom due to the degeneracy of the genetic code.

**[0127]** The meanings of the terms "nucleic acid molecule(s)", "nucleic acid sequence(s)" and "nucleotide sequence(s)" and the like are well known in the art and are used accordingly in context of the present invention.

**[0128]** For example; when used throughout this invention, these terms refer to all forms of naturally occurring or recombinantly generated types of nucleotide sequences and/or nucleic acid sequences/molecules as well as to chemically synthesized nucleotide sequences and/or nucleic acid sequences/molecules. These terms also encompass nucleic acid analogues and nucleic acid derivatives such as e. g. locked DNA, PNA, oligonucleotide thiophosphates and substituted ribo-oligonucleotides. Furthermore, these terms also refer to any molecule that comprises nucleotides or nucleotide analogues.

**[0129]** Preferably, the terms "nucleic acid molecule(s)", "nucleic acid sequence(s)" and "nucleotide sequence(s)" and the like refer to deoxyribonucleic acid (DNA) or ribonucleic acid (RNA). The "nucleic acid molecule(s)", "nucleic acid sequence(s)" and "nucleotide sequence(s)" may be made by synthetic chemical methodology known to one of ordinary skill in the art, or by the use of recombinant technology, or may be isolated from natural sources, or by a combination thereof. The DNA and RNA may optionally comprise unnatural nucleotides and may be single or double stranded. "Nucleic acid molecule(s)", "nucleic acid sequence(s)" and "nucleotides sequence(s)" also refer to sense and anti-sense DNA and RNA, that is, a nucleotide sequence which is complementary to a specific sequence of nucleotides in DNA and/or RNA.

**[0130]** Furthermore, the terms "nucleic acid molecule(s)", "nucleic acid sequence(s)" and "nucleotide sequence(s)" and the like may refer to DNA or RNA or hybrids thereof or any modification thereof that is known in the state of the art (see, e.g., US 5525711, US 4711955, US 5792608 or EP 302175 for examples of modifications). These molecules may be single- or double-stranded, linear or circular, natural or synthetic, and without any size limitation. For instance, the "nucleic acid molecule(s)", "nucleic acid sequence(s)" and/or "nucleotide sequence(s)" may be genomic DNA, cDNA, mRNA, antisense RNA, ribozymal or a DNA encoding such RNAs or chimeroplasts (Cole-Strauss Science 1996 273



(5280) 1386-9). They may be in the form of a plasmid or of viral DNA or RNA. "Nucleic acid molecule(s)", "nucleic acid sequence(s)" and "nucleotide sequence(s)" and the like may also refer to (an) oligonucleotide(s), wherein any of the state of the art modifications such as phosphothioates or peptide nucleic acids (PNA) are included.

**[0131]** The nucleic acid-molecules as disclosed herein are particularly useful for producing a cyclic peptide of the invention, for example by the corresponding method disclosed herein.

**[0132]** In another aspect, disclosed herein also is a vector comprising the nucleic acid molecule as disclosed herein and described above.

**[0133]** Said vector may be a cloning vector or an expression vector, for example, a phage, plasmid, viral or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host/cells. The herein disclosed nucleic acid molecule may be joined to a particular vector containing selectable markers for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate or rubidium, chloride precipitate, or in a complex with a charged lipid or in carbon-based clusters, such as fullerenes. Should the vector be a virus, it may be packaged in vitro using an appropriate packaging cell line prior to application to host cells.

**[0134]** Preferably, the nucleic acid molecule as disclosed herein is operatively linked to expression control sequences (e.g. within the herein disclosed vector) allowing expression in prokaryotic or eukaryotic cells or isolated fractions thereof. Expression of said polynucleotide comprises transcription of the nucleic acid molecule, preferably into a translatable mRNA. Regulatory elements ensuring expression in eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art. They usually comprise regulatory sequences ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the *lac*, *trp* or *tac* promoter in *E. coli*, and examples for regulatory elements permitting expression in eukaryotic host cells are the *AOX1* or *GAL1* promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (Invitrogen), pSPORT1 (GIBCO BRL). Preferably, said vector is an expression vector and/or a gene transfer vector. Expression vectors derived from viruses such as retroviruses, adenoviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the polynucleotides or vector as disclosed herein into a targeted cell population. Methods which are well known to those skilled in the art can be used to construct a vector as disclosed herein; see, for example, the techniques described in Sambrook, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1994). Alternatively, the polynucleotides and vectors as disclosed herein can be reconstituted into liposomes for delivery to target cells.

**[0135]** The term "isolated fractions thereof" refers to fractions of eukaryotic or prokaryotic cells or tissues which are capable of transcribing or transcribing and translating RNA from the vector disclosed herein. Said fractions comprise proteins which are required for transcription of RNA or transcription of RNA and translation of said RNA into a polypeptide. Said isolated fractions may be, e.g., nuclear and cytoplasmic fractions of eukaryotic cells such as of reticulocytes. Kits for transcribing and translating RNA which encompass the said isolated fractions of cells or tissues are commercially available, e.g., as TNT reticulolysate (Promega).

**[0136]** Again, like the disclosed nucleic acid molecules, also the vectors as described herein are particularly useful for producing a cyclic peptide of the invention, for example by the corresponding method disclosed herein.

**[0137]** In a further aspect, disclosed herein is a recombinant host cell comprising the nucleic acid molecule and/or the vector as disclosed herein. In context of this aspect, the nucleic acid molecule and/or the vector as disclosed herein can, inter alia, be used for genetically engineering host cells, e.g., in order to express and isolate the amino acid backbone/primary amino acid sequence of the cyclic peptide disclosed herein, and hence, the linear peptide as disclosed herein.

**[0138]** Said host cell may be a prokaryotic or eukaryotic cell; see supra. The nucleic acid molecule or vector which is present in the host cell may either be integrated into the genome of the host cell or it may be maintained extrachromosomally.

**[0139]** The host cell can be any prokaryotic or eukaryotic cell, such as a bacterial, insect, fungal, plant, animal, mammalian or, preferably, human cell. Preferred fungal cells are, for example, those of the genus *Saccharomyces*, in particular those of the species *S. cerevisiae*, or those belonging to the group of hyphal fungi, for example several *penicillia* or *aspergilla* strains. The term "prokaryotic" is meant to include all bacteria which can be transformed or transfected with a nucleic acid molecule for the expression of an amino acid backbone/primary amino acid sequence of the cyclic peptide disclosed herein, and hence, the linear peptide disclosed herein. Prokaryotic hosts may include gram negative as well as gram positive bacteria such as, for example, *E. coli*, *S. typhimurium*, *Serratia marcescens* and *Bacillus subtilis*. A nucleic acid molecule coding for an amino acid backbone/primary amino acid sequence of the cyclic peptide disclosed

herein, and hence, the herein disclosed linear peptide can be used to transform or transfect a host using any of the techniques commonly known to those of ordinary skill in the art. Methods for preparing fused, operably linked genes and expressing them in bacteria or animal cells are well-known in the art (Sambrook, supra). The genetic constructs and methods described therein can be utilized for expression of the above mentioned amino acid backbone/primary amino acid sequence and linear peptide in, e.g., prokaryotic hosts.

**[0140]** In general, expression vectors containing promoter sequences which facilitate the efficient transcription of the inserted polynucleotide are used in connection with the host. The expression vector typically contains an origin of replication, a promoter, and a terminator, as well as specific genes which are capable of providing phenotypic selection of the transformed cells. The transformed prokaryotic hosts can be grown in fermentors and cultured according to techniques known in the art to achieve optimal cell growth. The expressed peptides can then be isolated from the grown medium, cellular lysates, or cellular membrane fractions. The isolation and purification of the microbially or otherwise expressed peptides may be by any conventional means such as, for example, preparative chromatographic separations and immunological separations such as those involving the use of monoclonal or polyclonal antibodies (Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1994)).

**[0141]** Again, like the nucleic acid molecules and the vectors as described herein, also the corresponding host cells are particularly useful for producing a cyclic peptide of the invention, for example by the corresponding method disclosed herein.

**[0142]** In yet another aspect, the present invention relates to a method for producing a cyclic peptide of the present invention, comprising the steps of

a)

- (i) culturing a recombinant host cell as disclosed herein under conditions such that the amino acid backbone of the herein disclosed cyclic peptide (or the herein disclosed linear peptide) is expressed, and recovering said amino acid backbone (or said linear peptide); or
- (ii) chemically synthesizing the amino acid backbone of the herein disclosed cyclic Peptide (or the herein disclosed linear peptide); and

b) cyclization of said amino acid backbone (or said linear peptide) to form the herein disclosed cyclic peptide.

**[0143]** The definitions given herein-above with respect to the term "cyclization" apply here, mutatis mutandis. In the particular context of the above method, the meaning of the term "cyclization" encompasses both, forming of the intramolecular bridge (disulphide bond) and the ring closure by covalently connecting the N- and C-termini of the backbones of the cyclic peptides to be produced.

**[0144]** The definitions given herein with respect to the cyclic peptide, its amino acid backbone or the corresponding linear peptide according to this invention, as well as with respect to the host cell disclosed herein, apply here, mutatis mutandis.

**[0145]** As already mentioned above with respect to the disclosed cyclic and linear peptides, in a preferred embodiment of this further aspect of the invention, the N-terminal amino acid of the amino acid backbone/linear peptide to be cyclized in order to produce a cyclic peptide of this invention is Ala, Arg or Lys and the corresponding C-terminal amino acid is Gln, Gly or Glu (also DGLu is possible) or Pro. However, as mentioned-above, also other N- and C-terminal amino acids are envisaged, i.e. also other cyclization (ring closure) sites can be employed in context of the disclosed method.

**[0146]** The person skilled in the art is readily able to put the herein disclosed method for producing a cyclic peptide into practice, based on his common general knowledge and the prior art like WO 2006/103101, which discloses a general methodology how to synthesize peptides and, particularly cyclic peptides. Also, the teachings of the invention, for example in the appended experimental part (example 1), provides for enabling technical guidance.

**[0147]** In the non-limiting example 1 of the invention, the cyclopeptide mutants were first synthesized in form of their linear peptides/amino acid backbones (for example by applying a chemical synthesis approach, like the Fmoc / tert butyl strategy (as described in WO 2006/103101; Chen W.C. and White P.D.: Fmoc Solid Phase Peptide Synthesis, Oxford University Press 2003)), and were then cyclized covalently on the backbone by condensation of the C-terminal carboxyl group with the amino group of the N-terminal amino acid ("head to tail" cyclization; Kates S. and Albericio F.: Solid phase synthesis, CRC-Press, 2000).

**[0148]** Subsequently, a disulphide bond is established between those two cysteine residues of the linear peptides which are able to form a disulphide bond (e.g. between Cys<sub>7</sub> and Cys<sub>13</sub> of the 18mer peptide, between Cys<sub>10</sub> and Cys<sub>16</sub> of the 22mer peptide or between Cys<sub>11</sub> and Cys<sub>17</sub> of the 25mer peptide) by chemical interaction known in the art (e.g. Benoiton N.L.: Chemistry of Peptide Synthesis. CRC-Press, 2005).

**[0149]** In general, in context of the "cyclization" step of the above described method; the ring closure of the linear backbone of the cyclic peptides to be produced may be performed before or after the formation of the S-S bridge. In

other words, the S-S bridge between the two Cys residues of the AA chain of the peptides may be the first step in the "cyclization" procedure of the described production process and the ring closure may be the second step, or vice versa. The skilled person is able to find out which of these particular approaches is appropriate for a given setup of the production preconditions.

**[0150]** As mentioned above, the linear peptides/amino acid backbones of the cyclic peptides to be produced can also be produced by recombinant engineering techniques. Such techniques are well known in the art (e. g. Sambrook, supra). As also mentioned above, by this kind of production of said linear peptides/amino acid backbones particular advantage can be taken of the herein disclosed and described nucleic acid molecules, vectors and/or host cells. The definitions correspondingly given above apply here, *mutatis mutandis*.

**[0151]** Several approaches of peptide synthesis particular synthesis approaches of cyclic peptides are known in the art. (e.g. Williams, Chemical Approaches to the Synthesis of Peptides, CRC-Press 1997; Benoiton: Chemistry of Peptide Synthesis. CRC-Press, 2005). The skilled person is readily in the position to apply the prior art knowledge to the particular requirements of the disclosed method for producing cyclic peptides, based of the herein provided teaching.

**[0152]** As already mentioned above, also disclosed herein is a cyclic peptide obtainable or obtained by the above described method, but also a corresponding linear peptide (amino acid backbone/ primary sequence of the corresponding cyclic peptide) obtainable or obtained by the above described method as some kind of an intermediate product (particularly a product obtainable or obtained by step a) of the above described method).

**[0153]** In general, the cyclic peptide according to the present invention may, inter alia, be for use in medical intervention approaches. Such approaches comprise the use as or in a diagnostic agent and for the manufacture of a medicament for the treatment of diseases or the use in or as a composition, preferably a pharmaceutical composition, a diagnostic composition or a diagnostic kit, preferably for the detection of anti- $\beta$ -AR antibodies, more preferably for the detection of anti- $\beta_1$ -AR antibodies.

**[0154]** As mentioned, the antibodies as defined or described herein are preferably autoantibodies.

**[0155]** Non-limiting uses and applications of the disclosed compounds, particularly the cyclic peptide according to the present invention, are described herein, for example in the following.

**[0156]** The present invention also relates to a composition comprising a cyclic peptide as provided in the context of the present invention, and optionally a carrier.

**[0157]** In one particular embodiment of this aspect, said composition is a pharmaceutical composition and said carrier is a pharmaceutically acceptable carrier.

**[0158]** The composition of this invention, particularly pharmaceutical composition of this invention, is particularly for the use in the treatment, amelioration or prevention as described and defined herein. Accordingly, the pharmaceutical composition of this invention may be for use in the treatment, amelioration or prevention of a disease where the activity of a  $\beta$ -AR is enhanced or for the treatment of a patient having antibodies against a  $\beta$ -AR. Moreover, the pharmaceutical composition of this invention may be for use in inducing immune tolerance of a patient, particularly immune tolerance of a patient with respect to immunogenic stretches of the endogenous  $\beta_1$ -AR.

**[0159]** Apart from containing at least one cyclic peptide of the present invention, the (pharmaceutical) composition provided may either comprise two or a plurality (like at least 3 or at least 5) of cyclic peptides of the present invention.

**[0160]** Likewise, not only one, but also two or a plurality (like at least 3 or at least 5) of said cyclic peptides may be for use in administering to a patient in need of medical intervention in accordance with the present invention. Thereby, the administration of said more than one of cyclic peptides may be simultaneously or successively.

**[0161]** Moreover, in one particular embodiment, the present invention relates to the pharmaceutical composition, the method or uses for medical intervention or the cyclic peptide or the pharmaceutical composition as disclosed herein, wherein said cyclic peptide is to be administered with or said pharmaceutical composition comprises at least one further pharmaceutically active agent.

**[0162]** Said at least one further pharmaceutically active agent may be a  $\beta$  receptor blocker, preferably a selective  $\beta$ -AR blocker, like, for example, a  $\beta_1$ -AR blocker selected from the group consisting of atenolol, metoprolol, nebivolol, bisoprolol and the like. Without being bound by theory, this kind of particular combination provides for protection from antibody-induced, selective  $\beta_1$ -AR downregulation by the herein provided cyclic peptides, since  $\beta_1$ -AR is at the same time upregulated by betablockers, like bisoprolol or metoprolol, and ultimately results in a synergistic effect of the cyclic peptides and the additional  $\beta$ -blocker(s).

**[0163]** The carrier optionally comprised in the (pharmaceutical) composition of the invention or to be administered together with the (pharmaceutical) composition or the cyclic peptide of the invention may particularly be a pharmaceutically acceptable carrier, excipient or diluent.

**[0164]** Such carriers are well known in the art. The skilled person is readily in the position to find out such carriers which are particularly suitable to be employed in accordance with the present invention.

**[0165]** In the following, several non-limiting administration schemes and the use of correspondingly suitable pharmaceutically acceptable carrier are described.

**[0166]** For use in an administration of the pharmaceutical composition and/or the cyclic peptides in accordance with

this invention via subcutaneous (s.c.) or intravenous (i.v.) injection, compounds of the invention may be formulated in aqueous solution, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiologically saline buffer. For transmucosal and transpulmonary administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

**[0167]** The use of pharmaceutical acceptable carriers to formulate the compounds according to the present invention into dosages or pharmaceutical compositions suitable for systemic, i.e. intravenous/intraarterial, or subcutaneous administration is within the scope of the present invention. With proper choice of carrier and suitable manufacturing practice, the compositions of the present invention, in particular those formulated as solutions, may be administered parenterally, such as by intravenous injection. The compounds can be readily formulated using pharmaceutically acceptable carriers well known in the art into dosages suitable for subcutaneous or oral administration. Such carriers enable the compounds according to the present invention to be formulated as tablets, pills, capsules, dragees, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a subject to be treated.

**[0168]** Compounds according to the present invention, or medicaments comprising them, intended to be administered intracorporally/intracellularly may be administered using techniques well known to those of ordinary skill in the art. For example, such agents may be encapsulated into liposomes, then administered as described above. Liposomes are spherical lipid bilayers with aqueous interiors. All molecules present in an aqueous solution at the time of liposome formation are incorporated into the aqueous interior. The liposomal contents are both protected from the external microenvironment and, because liposomes fuse with cell membranes, are efficiently delivered near the cell surface. Delivery systems involving liposomes are disclosed in U.S. Patent No. 4,880,635 to Janoff et al. The publications and patents provide useful descriptions of techniques for liposome drug delivery.

**[0169]** Pharmaceutical compositions comprising a compound according to the present invention for use in parenteral and/or subcutaneous administration include aqueous solutions of the active compound(s) in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil or castor oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain compounds which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, dextran, or the like. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions and to allow for a constantly slow release of the substance in the organism.

**[0170]** It is clear to the skilled person that, in accordance with the present invention, the disclosed pharmaceutical composition or cyclic peptide may be administered in a pharmaceutically/therapeutically effective dose, which means that a pharmaceutically/therapeutically effective amount of the compound administered is reached. Preferably, a pharmaceutically/therapeutically effective dose refers to that amount of the compound administered (active ingredient) that produces amelioration of symptoms or a prolongation of survival of a subject which can be determined by the one skilled in the art doing routine testing.

**[0171]** It is of note that the dosage regimen of the compounds to be administered in accordance with the present invention will be determined by the attending physician and clinical factors. As is well known in the medical arts, that dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. A person skilled in the art is aware of and is able to test the relevant doses, the compounds to be medically applied in accordance with the present invention are to be administered in.

**[0172]** As shown herein, the effect of the cyclic peptides provided herein, namely the blockage of anti- $\beta_1$ -AR antibodies, can be obtained in a dose dependent manner. Thereby, the efficiency of the Cys→Ser mutated cyclopeptides as disclosed herein depends on a threshold concentration (Figs. 14, 15, and 16 B,C).

**[0173]** Accordingly, the disclosed pharmaceutical composition or cyclic peptide may particularly be administered in a manner that it is present in a concentration, i.e. reaches a threshold concentration, of at least 0,05 mg per kg body weight, preferably in a concentration of at least 0,1 mg per kg body weight, more preferably in a range of 0,1 mg per kg body weight (100  $\mu$ g/kg) to 100 mg per kg body weight, more preferably in a range of 1 mg per kg body weight to 100 mg per kg body weight and most preferably in a range of 1 mg per kg body weight to 10 mg per kg body weight. Particularly, the effective dose of the disclosed pharmaceutical composition or cyclic peptide may be at about 1 mg per kg body weight. Also higher concentrations of the disclosed pharmaceutical composition or cyclic peptide are generally envisaged to be reached by correspondingly applied administration schemes. For example, such higher concentrations may be at least 2, 3, 4 or 5 mg per kg body weight. Concentrations of at least 1 mg per kg body weight or at least 2 mg per kg body weight are preferred.

**[0174]** One particularly preferred, non-limiting administration scheme to be applied in context of this invention is an s.c. or i.v. application every two or four weeks.

**[0175]** In this context, it is of note that in the rat model employed herein a dose of 1 to 4 mg/kg i.v. every other month were sufficient to obtain therapeutic levels of the compounds according to the present invention, with the respective dosage for humans preferably being about 0,3-10 mg/kg i.v. or s.c., more preferably being about 1-10 mg/kg i.v. or s.c.,

even more preferably being about 1-5 mg/kg i.v. or s.c.

**[0176]** As demonstrated herein, the administration of the disclosed cyclic peptides may initially trigger a transient opposite immune response, in particular when applied in higher doses. Such transient immune responses in the long run are compensated by the antibody-inactivating activity of the administered cyclic peptides. This may lead to an decelerated effect of the administered cyclic peptides, i.e. a decelerated elimination of anti- $\beta_1$ -AR antibodies and hence a decelerated reduction of (aberrant)  $\beta_1$ -AR activity.

**[0177]** The present invention also relates to a cyclic peptide or a pharmaceutical composition as disclosed herein, and optionally a pharmaceutically acceptable carrier, for use in

- a) the treatment, amelioration or prevention of a disease where the activity of a  $\beta$ -AR, preferably  $\beta_1$ -AR, is enhanced;
- b) the treatment of a patient having antibodies against a  $\beta$ -AR, preferably against  $\beta_1$ -AR, or suffering from or being at risk to develop a disease as disclosed herein; or
- c) inducing immune tolerance.

**[0178]** The diseases to be medically intervened (treated, ameliorated, prevented or diagnosed) in accordance with this invention or the diseases the patient as defined and described herein suffers from are preferably those, where the  $\beta_1$ -AR is activated in a non-physiological manner, more preferably is activated by antibodies, more preferably by auto-antibodies which are directed against the  $\beta_1$ -AR.

**[0179]** Exemplarily and preferably, the diseases to be medically intervened in accordance with this invention or the diseases the patient as defined and described herein suffers from comprise, however, are not limited thereto, the group of heart diseases.

**[0180]** Particularly, the heart diseases to be medically intervened in accordance with this invention or the heart diseases the patient as defined and described herein suffers from may comprise but are not limited to infectious and non-infectious heart disease, ischemic and non-ischemic heart disease, inflammatory heart disease and myocarditis, cardiac dilatation, idiopathic cardiomyopathy, (idiopathic) dilated cardiomyopathy (DCM), immune-cardiomyopathy, heart failure, and any cardiac arrhythmia including ventricular and/or supraventricular premature capture beats as well as any atrial arrhythmia including atrial fibrillation and/or atrial flutter.

**[0181]** In other words, the heart disease as referred to in the descriptions and definitions given herein with respect to the methods or the cyclic peptide or the pharmaceutical composition of the invention may be heart diseases selected from the group comprising infectious and non-infectious heart disease, ischemic and non-ischemic heart disease, inflammatory heart disease and myocarditis, cardiac dilatation, idiopathic cardio-myopathy, (idiopathic) dilated cardiomyopathy (DCM), immune-cardiomyopathy, heart failure, and any cardiac arrhythmia including ventricular and/or supraventricular premature capture beats as well as any atrial arrhythmia including atrial fibrillation and/or atrial flutter.

**[0182]** It is of note that the most preferred disease to be medically intervened (treated, ameliorated, prevented or diagnosed) in accordance with this invention or the most preferred disease the patient as defined and described herein suffers from is DCM, preferably idiopathic DCM.

**[0183]** A particular subgroup of the "patients" for the purpose of the present invention are those patients suffering from any of the diseases described herein, more particularly the group of heart diseases described herein and having at the same time antibodies directed against  $\beta$ -ARs, more preferably antibodies against the  $\beta_1$ -AR, whereby the antibodies are preferably auto-antibodies.

**[0184]** A disease to be medically intervened (treated, ameliorated, prevented or diagnosed) in accordance with this invention or a disease the patient as defined and described herein suffers from is intended to be induced by antibodies against a  $\beta$ -AR, preferably by antibodies against  $\beta_1$ -AR. Preferably, these antibodies are auto-antibodies.

**[0185]** The means and methods provided herein are particularly useful when provided in the prophylaxis/prevention of a disease as defined herein. This means that a patient may be treated with the cyclic peptide and/or pharmaceutical composition of the invention prior to the onset (of symptoms) of a disease as defined herein. For example, this preventive treatment may follow a preceding diagnostic application that, e.g., takes advantage of the diagnostic means and methods provided herein. Thereby, it is preferred that a preventive treatment taking advantage of the therapeutic means and methods of this invention is applied, when the risk to develop a disease as defined herein is diagnosed, e.g. when anti- $\beta$ -AR (auto-)antibodies are detected.

**[0186]** In this context, a preferred "patient" is one bearing at risk to develop a disease as defined herein. Particularly, such a patient is one having anti- $\beta$ -AR (auto-)antibodies, preferably anti- $\beta_1$ -AR (auto-)antibodies, but not (yet) suffering from a disease as defined herein, or symptoms thereof.

**[0187]** The immune tolerance to be induced in context of this invention is envisaged to be particularly obtained by suppression of the production of antibodies against immunogenic stretches of the  $\beta$ -AR molecule, which, without being bound by theory, may be due to a blockade of the antigen-recognition sites of the antibody-producing early (mature) B-cells and memory B-cells.

**[0188]** It is within the present invention that the provided pharmaceutical composition or cyclic peptide is particularly

for use in the treatment, prevention and/or amelioration of any of the diseases and patient groups or patients as defined herein including the detection of anti- $\beta$ -AR antibodies in these patients by using the aforementioned compounds.

**[0189]** A "patient" for the purposes of the present invention, i. e. to whom a compound according to the present invention is to be administered or who suffers from the disease as defined and described herein or who is intended to be diagnosed in accordance with this invention, includes both humans and other animals and organisms. Thus the compounds and methods of this invention are applicable to or in connection with both human therapy and veterinary applications including diagnostic(s), diagnostic procedures and methods as well as staging procedures and methods. In the preferred embodiment the patient is a mammal, and in the most preferred embodiment the patient is human.

**[0190]** The mutant cyclic peptides according to the present invention may also be used for the preparation of a medicament for the treatment, prevention and/or amelioration of any of the diseases and patient groups/patients as defined herein. What is said herein for the pharmaceutical composition applies also to the medicament for the manufacture of which the peptides of the present invention may be used.

**[0191]** In a still further aspect, disclosed herein is a diagnostic agent comprising or being a cyclic peptide or a composition according to this invention, and optionally at least one further biologically active compound.

**[0192]** Preferably the herein disclosed diagnostic agent consists of or comprises a mutant peptide of the present invention, whereby the mutant peptide comprises a label. Such label may be selected from the group comprising radioactive labels and fluorescent labels. Respective labels are known to the ones skilled in the art. The definitions and descriptions of labels as given herein-above apply here, mutatis mutandis. Typically, the peptide is the part of the diagnostic agent conferring specific binding characteristics to the diagnostic agent, preferably binding to anti- $\beta_1$ -AR antibodies, whereas the label confers the signalling characteristics to the diagnostic agent.

**[0193]** The diagnostic agent as disclosed herein may comprise, apart from (a) labelled or unlabelled mutant peptide(s) of the present invention, a further biologically active compound. Such further biologically active compound may be a means to confer signalling characteristics to the diagnostic agent, particularly in case the mutant peptides of the present invention are unlabelled. For example, the further biologically active compound can be an antibody, preferably a monoclonal antibody, and more preferably a labelled antibody specifically binding to a mutant peptide of the present invention or to a complex consisting of a mutant peptide of the present invention and an anti- $\beta$ -AR antibody, preferably an anti- $\beta_1$ -AR antibody.

**[0194]** In a further aspect, disclosed herein is a method for diagnosing a disease as defined and described herein comprising the steps of

- a) detecting antibodies against a  $\beta$ -AR (for example in a sample) using the cyclic peptide or the composition or the diagnostic agent of the present invention; and
- b) diagnosing for said disease, when the titer of said antibodies is increased.

**[0195]** In a further aspect, disclosed herein are the mutant peptides, pharmaceutical compositions and medicaments according to the present invention for use in a method for diagnosing a patient which can be treated using said compounds according to the present invention. In context of this particular

**[0196]** method also a step of detecting antibodies against a  $\beta$ -AR (for example in a sample) using the compounds of the present invention and/or a step of considering whether the outcome of said detection step indicates a disease as defined herein, may be employed. As mentioned, a disease as defined herein or the risk to develop a disease as defined herein is indicated, when the titer of said anti- $\beta$ -AR antibodies is increased.

**[0197]** In another aspect, disclosed herein is a cyclic peptide, a composition or a diagnostic agent as provided and described herein for use in diagnosing (for example in a sample) a disease as defined herein. Again, a disease as defined herein or the risk to develop a disease as defined herein is indicated by an increased titer of anti- $\beta$ -AR antibodies.

**[0198]** In context of the present invention the term "increased titer of anti- $\beta$ -AR antibodies" means that the titer of anti- $\beta$ -AR antibodies (for example in a sample derived from a patient to be diagnosed in accordance with this invention) is higher than that of a healthy control patient, i.e. a patient not suffering from a disease as defined herein and/or a patient lacking anti- $\beta$ -AR antibodies.

**[0199]** As mentioned, in healthy patients, anti- $\beta$ -AR antibodies are usually hardly or not at all present or detectable. Accordingly, an "increased titer of anti- $\beta$ -AR antibodies" in accordance with the present invention preferably refers to any occurrence of anti- $\beta$ -AR antibodies, i.e. any occurrence of a detectable amount of anti- $\beta$ -AR antibodies.

**[0200]** A suitable "sample" in accordance with the present invention includes, but is not limited to, (a) biological or medical sample(s), like, e.g. (a) sample(s) comprising cell(s) or tissue(s). For example, Such (a) sample(s) may comprise (s) biological material of biopsies. The meaning of "biopsies" is known in the art. For instance, biopsies comprise cell(s) or tissue(s) taken, e. g. by the attending physician, from a patient/subject as described herein. Exemplarily, but not limiting, the biological or medical sample to be analysed in context of the present invention is or is derived from blood, plasma, white blood cells, urine, semen, sputum, cerebrospinal fluid, lymph or lymphatic tissues or cells, muscle cells, heart cells, cells from veins or arteries, nerve cells, cells from spinal cord, brain cells, liver cells, kidney cells, cells from

the intestinal tract, cells from the testis, cells from the urogenital tract, colon cells, skin, bone, bone marrow, placenta, amniotic fluid, hair, hair and/or follicles, stem cells (embryonic, neuronal, and/or others) or primary or immortalized cell lines (lymphocytes, macrophages, or cell lines). Preferred "samples" in accordance with the present invention are those derived from blood or plasma. The biological or medical sample as defined herein may also be or be derived from

biopsies, for example biopsies derived from heart tissue, veins or arteries.

**[0201]** In a further aspect, disclosed herein is a diagnostic kit, for example a diagnostic kit for the detection of antibodies against a  $\beta$ -AR, comprising the herein disclosed cyclic peptide, composition or diagnostic agent.

**[0202]** The disclosed kit comprises at least one of the compounds as disclosed according to the invention, like, for example a cyclic or linear peptide a nucleic acid molecule, vector or host cell or a composition or diagnostic agent.

**[0203]** In one particular aspect, the kit further comprises an instruction leaflet, and/or a buffer for use in the application of the kit, and/or at least one reaction vessel for carrying out the detection reaction for which the kit is or is to be used. In a further aspect, at least one, some or all of the reagents used in connection with the application of said kit are present as portions useful in carrying out the reaction(s) for which the kit is to be used.

**[0204]** The cyclic peptides, diagnostic agents or kits as disclosed herein may also be applied for the detection of  $\beta_1$ -AR. Accordingly, these cyclic peptides, diagnostic agents or kits may particularly be useful for identifying/detecting bound anti- $\beta_1$ -AR antibodies on cell- and/or tissue surfaces. For example, the cyclic peptides, diagnostic agents or kits as disclosed herein may be used in imaging purposes, like single photon emission computed tomography (SPECT), MiBi, PET, magnetic resonance tomography (MRT) or other diagnostic imaging techniques employed in medicine. Due to the  $^{131}\text{I}$ -labelled CP-distribution pattern in vivo (Fig. 30), the disclosed cyclic peptides, diagnostic agents may, for example, be particularly useful as organ-specific tracers.

**[0205]** One particular approach for using the compounds according to the present invention as a diagnostic and in a diagnostic method, respectively, is a three-step screening procedure. For example, this method comprises performing an ELISA with the cyclic peptides according to the present invention as well as determining immunofluorescence and determining cAMP responses in cells expressing native human  $\beta$ -AR. It is to be acknowledged that each and any of the aforementioned steps can as such be preformed for the detection of said antibodies using the cyclic peptides according to the present invention. A large number of patients, for example heart failure patients, may thus be screened for functionally active anti- $\beta_1$ -AR antibodies. In connection with such (but also with other herein disclosed) diagnostic methods, the definition of functionally active anti- $\beta_1$ -AR antibodies is preferably based on their effects on receptor-mediated signalling, that is, their effects on cellular cAMP levels and on the activity of the cAMP-dependent protein kinase (PKA). Cyclic AMP is an universal second messenger of many G protein-coupled receptors including the  $\beta$ -AR family. It exerts its effects via PKA, cAMP-gated ion channels, phosphodiesterases, and exchange proteins directly activated by cAMP, known as Epac1 and 2. The prior art describes several fluorescence methods for measuring cAMP in intact cells which can all be used in connection with the herein disclosed diagnostic method. Fluorescence resonance energy transfer (FRET) between green fluorescent protein (GFP) variants fused to the regulatory and catalytic subunits of PKA has been described to study the spatio-temporal dynamics of cAMP in neurons (Hernpel CM, Vincent P, Adams SR, Tsien RY, Sclerston AI. *Nature*. 1996;384:113-114) or cardiac myocytes. (Zaccolo M, Pozzan T., *Science*. 2002;295: 1711-1715).

**[0206]** More recently, single chain fluorescence indicators have been described in the art which are characterized by having an enhanced cyan (CFP) or yellow fluorescent protein (YFP) directly fused to the cAMP-binding domain of Epac-proteins, which allowed to achieve a higher sensitivity and better temporal resolution of the cAMP measurements. Such system is, among others described in WO2005/052186. Such system can be used in connection with any diagnostic procedure using the cyclic peptides or other corresponding compounds according to the present invention. Also such system can be used for, however is not limited thereto, analyzing the prevalence of functionally active anti- $\beta_1$ -AR antibodies. Preferably such diagnostic method is applied to a cohort of previously antibody-typed DCM patients or any individual to be assessed insofar or any individual suspected of suffering from any of the diseases described herein or being at risk to suffer therefrom. In a further step of the diagnostic method and screening method, the ability of  $\beta$ -blockers to inhibit anti- $\beta_1$ -AR antibodies-induced receptor activation may be assessed and determined, respectively.

**[0207]** The afore described assay which is a FRET-based methods as described in WO 2005/052186 making use of the peptides according to the present invention is advantageous insofar as it is simpler, less time consuming, and at the same time discloses or identifies all DCM patients previously considered anti- $\beta_1$ -EC<sub>II</sub> antibody-positive. This embodiment of a FRET based method of diagnosing making use of one or several of the peptides according to the present invention is based on detecting antibody-induced increases in cAMP.

**[0208]** Taken together, screening by Epac-FRET appears to represent a very sensitive single step approach, allowing to detect activating antibodies directed against the human  $\beta_1$ -AR. Therefore, also disclosed herein is the use of one or several of the peptides according to the present invention for use in an Epac-FRET assay. More preferably such Epac-FRET assay is for use in diagnosis, even more preferably in diagnosis of patients suffering from or suspected of suffering from any of the disease described herein.

**[0209]** In view of the above, it is particularly preferred to use or apply the FRET technology, particularly a FRET-based

detection system, in accordance with this invention.

[0210] In a further aspect, the present invention relates to a method for detecting of antibodies against a  $\beta$ -AR (for example in a sample as defined herein) comprising the step of contacting the cyclic peptide of the invention with said antibodies to be detected.

[0211] In a further aspect, the present invention relates to the cyclic peptide, composition or diagnostic agent as disclosed herein for detecting (for example in a sample as defined herein) antibodies against a  $\beta$ -AR.

[0212] The above method for detecting of antibodies or cyclic peptide, composition or diagnostic agent is particularly useful to be employed in context of the diagnostic applications as described herein

[0213] Throughout the present application, the following abbreviations shall have the following meanings: Ab or ab: antibody, Abs or abs: antibodies, AR: adrenergic receptor, EC extra cellular domain of a  $\beta$ -AR, EC<sub>II</sub> extra cellular domain II of a  $\beta$ -AR and AA amino acid.

[0214] The present invention is further described by reference to the following non-limiting figures and examples.

[0215] The Figures show:

**Figure 1** is a diagram depicting the scheme of the  $\beta_1$ -EC<sub>II</sub>-25 amino-acid (AA) cyclopeptide and the mutated  $\beta_1$ -EC<sub>II</sub>-25AA cyclopeptides (black rings with the original Cys-residues (white balls) or the Ser mutated Cysteines (black balls; Cys/Ser or Ser/Cys, respectively), and the corresponding high liquid pressure liquid chromatographic elution profiles detected at a wave length of 210 nm or 220nm, respectively.

For the 3 Cys-containing  $\beta_1$ -EC<sub>II</sub>-25AA Cys/Cys cyclopeptide, HPLC was carried out in a Waters Separation Modul 2690 together with a Waters Dual Lambda absorbance detector; absorbance was read at 220 nm. After peptide synthesis and cyclization, the samples were dissolved in H<sub>2</sub>O/5% acetonitril (ACN) and loaded on a Nucleosil 100-5/C18 column (Macherey-Nagel Inc., Germany; column length 250 mm, lumen 4 mm) with a flow of 1 ml/min; then a separation-gradient from 5% to 60% ACN in the presence of 0.2% TFA was run. The remaining faint amount of linear  $\beta_1$ -EC<sub>II</sub>-25AA peptide yielded a small peak, typically between 14 and 16 min, whereas the fractions containing the  $\beta_1$ -EC<sub>II</sub>-25AA Cys/Cys cyclopeptide appeared in a range from 18 to 22 min.

HPLC of the mutant 25AA-cyclopeptides was performed with a Silica C18 column (15 $\mu$ m, 120A, length 250 mm, lumen 4 mm) with a flow of 1 ml/min followed by a separation-gradient from 5% to 60% ACN in the presence of 0.1 % TFA. The fractions containing the cyclic  $\beta_1$ -EC<sub>II</sub>-25AA mutants were monitored by UV-absorption (210 nm) and showed a sharp single elution peak appearing at -10.5 min (25AA Cys/Ser-mutant, left panel) or -16.5 min (25AA Ser/Cys-mutant, right panel).

**Figure 2** is a diagram depicting the scheme of the mutated  $\beta_1$ -EC<sub>II</sub>-25AA or 18AA-cyclo-peptides (black rings with the original Cys-residues (white balls) or the Ser mutated Cysteines (black balls; Cys/Ser or Ser/Cys, respectively), together with the amino-acids involved in forming the primary ring structure after head-to-tail closure (closure site either Ala-DGlu, or Pro-Lys).

For the synthesis of cyclic  $\beta_1$ -EC<sub>II</sub>-18AA (or  $\beta_1$ -EC<sub>II</sub>-25AA) peptides on the solid phase, Fmoc-Glu-ODmab or another Fmoc amino acid having a side chain protecting group which can be selectively cleaved off in an orthogonal manner, is incorporated at the C-terminal end of the linear peptide. The cleaving off of the cyclic peptide from the synthesis resin generates a peptide amide (in the case of D-Glu $\rightarrow$ Gln) and the removal of the protective groups of the side chain is done by treating the resin with trifluoro acetate acid/triisopropylsilane/ethandithiole/water for several hours.

**Figure 3** shows six panels demonstrating the HPLC elution profiles of two linear (left panels, 18AA Cys/Ser and 25AA Cys/Cys, respectively) and four of the mutant cyclopeptides of the present invention, all of them Gln-containing cyclopeptides with a Pro-Lys closure site. HPLC of the mutant 25AA- or 18AA-cyclopeptides was carried out in a Waters Separation Modul together with a UV absorbance detector; absorbance was read at 210 nm. After peptide-synthesis and cyclization, the samples were dissolved in H<sub>2</sub>O/5% acetonitril (ACN) and loaded on a Silica C18 column (15 $\mu$ m, 120A, column length 250 mm, lumen 4 mm) with a flow of 1 ml/min; then a separation-gradient from 5% to 60% ACN in the presence of 0.1 % TFA was run. As shown for their linear counterparts, the fractions containing the cyclic  $\beta_1$ -EC<sub>II</sub>-18AA mutants (upper row, middle panel (Cys/Ser, -14 min), right panel (Ser/Cys, -16,5 min); or  $\beta_1$ -EC<sub>II</sub>-25AA mutants (lower row, middle panel (Cys/Ser, -10,6), right panel (Ser/Cys, ~16,5 )) gave sharp single elution peaks appearing at the indicated time points.

**Figure 4** is a diagram depicting the blocking capacity of  $\beta_1$ -EC<sub>II</sub>-18AA cyclopeptide mutants having a D-Glu ring closure (Cys/Ser mutation, white columns; Ser/Cys mutation, diagonally right hatched columns) compared with the 3 Cys-containing 18AA cyclopeptide (black columns) in an ELISA-competition assay using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen. Representative results obtained with IgG-fractions isolated from the sera of 12 different immunized antibody-positive rats are depicted. The y-axis represents the blocking efficiency of the various peptides used given in % of blocked versus non-blocked ELISA-reactivity of the sera after preincubation



(12h over-night, 4°C, rotating incubator) with the indicated cyclopeptides.

**Figure 5** is a diagram depicting the blocking capacity of  $\beta_1$ -EC<sub>II</sub>-18AA cyclopeptide mutants having a D-Glu ring closure (Cys/Ser mutation, white diamonds; Ser/Cys mutation, black diamonds) compared with the 3 Cys-containing linear 25AA Cys/Cys-peptide (black squares) in an ELISA-competition assay using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen. A single representative serum from an antibody-positive cardiomyopathic rat was used (Fig.4, rat number 4). The y-axis represents the concentration of specific anti- $\beta_1$ -EC<sub>II</sub> IgG antibodies as determined by ELISA, the x-axis corresponds to the molar excess of linear or cyclic peptides used to preincubate the IgG-fractions (12h, 4° C, rotating incubator) assuming a 1:1 stoichiometry (one cyclic (2.1 kDa molecular mass (MM)) or linear peptide (3.0 kDa MM) was assumed to block one IgG-antibody (150 kDa MM)).

**Figure 6** is a diagram depicting the blocking capacity of  $\beta_1$ -EC<sub>II</sub>-18AA cyclopeptide mutants having a D-Glu ring closure on  $\beta_1$ -receptor-mediated signalling (functional cAMP-assay) using an approach by fluorescence resonance energy transfer (FRET). The effect of the preincubation (12h, 4° C, rotating incubator) of anti- $\beta_1$ -EC<sub>II</sub> IgG antibodies of a representative rat (serum) (Fig. 4, rat number 4) with  $\beta_1$ -EC<sub>II</sub>-18AA-cyclopeptide mutants (Cys/Ser mutation, dark blue (3); Ser/Cys mutation, light blue (4)) was compared with the effect of a 3 Cys-containing 25AA Cys/Cys cyclopeptide (red (2)) or the result obtained with anti- $\beta_1$ -EC<sub>II</sub> IgG antibodies in the absence of blocking peptides (black (1)). The y-axis represents the normalized YFP/CFP-ratio of the registered FRET emission signals, the x-axis corresponds to the registration time given in seconds (s).

**Figure 7** is a diagram resuming the blocking effect of cyclopeptide mutants having a D-Glu ring closure after preincubation (12h, 4° C, rotating incubator) with IgG isolated from 78 sera from immunized antibody-positive rats in an ELISA-competition assay using the linear 3 Cys-containing 25AA Cys/Cys-peptide as an antigen. Columns represent the results obtained with mutant  $\beta_1$ -EC<sub>II</sub>-18AA cyclopeptides (Cys/Ser mutation black column; Ser/Cys mutation, white column) compared with the 3 Cys-containing 18AA cyclopeptide (vertically hatched column), the 3 Cys-containing 25AA cyclopeptide (horizontally hatched column), or the 3 Cys-containing linear 25AA peptide (diagonally right hatched column). Error bars indicate the standard error of the mean ( $\pm$  SEM). The y-axis represents the blocking efficiency of the various peptides used given in % of blocked versus non-blocked ELISA- reactivity of the sera.

**Figure 8** is a diagram resuming the blocking capacity of  $\beta_1$ -EC<sub>II</sub>-18AA cyclopeptides mutants (having a D-Glu ring closure) in an ELISA competition assay performed with sera from n=82 immunized antibody-positive rats. About 95% of the sera were efficiently blocked by the  $\beta_1$ -EC<sub>II</sub>-18AA Cys/Ser mutated cyclopeptide alone (schematically depicted on the top of the left white column), whereas about 5% of the sera were blocked by both, the 18AA Cys/Ser- and the 18AA Ser/Cys-mutants (the latter schematically depicted on the top of the right black column).

**Figure 9** is a diagram composed of two major (upper and lower) panels resuming the blocking effect of both 25AA- and 18AA-cyclopeptide mutants having a Gln closure site, as well as 18AA cyclopeptide mutants having a D-Glu closure site after preincubation (12h, 4° C, rotating incubation) with sera isolated from 69 different immunized antibody-positive rats in an ELISA-competition assay using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen. The upper panel depicts rat sera preferentially reacting with the Cys/Ser mutated cyclopeptides (type1 reaction, n=64), separated in cyc25AA(Gln)-peptides (left) and cyc18AA(Gln and D-Glu)-peptides (right). The lower panel depicts the rat sera reacting with both the Cys/Ser and the Ser/Cys mutated cyclopeptides (type2 reaction, n=5), again separated in the results obtained with cyc25AA(Gln)-peptides (left) and cyc18AA(Gln and D-Glu)-peptides (right).

The first three columns on the left side within the two panels represent the results obtained with the (non mutated) 3 Cys-containing 25AA (Gln-)cyclopeptide (black columns) and the mutant  $\beta_1$ -EC<sub>II</sub>-25AA (Gln-)cyclopeptides (Cys/Ser mutation, white columns; Ser/Cys mutation, horizontally hatched columns).

The five columns on the right side within the two panels represent the results obtained with the (non-mutated) 3 Cys-containing 18AA (Gln-)cyclopeptide (black columns) compared with the different 2 Cys-containing mutant 18AA cyclopeptides (18AA Cys/Ser mutant with a Gln closure site, white columns; 18AA Cys/Ser mutant with a D-Glu closure site, diagonally left hatched columns; 18AA Ser/Cys mutant with a Gln closure site, diagonally right hatched columns; 18AA Ser/Cys mutant with a D-Glu closure site, vertically hatched columns).

The error bars represent the standard error of the mean ( $\pm$  SEM). The y-axis represents the blocking efficiency of the various peptides used given in % of blocked versus non-blocked ELISA reactivity of the sera.

**Figure 10** is a diagram resuming the blocking capacity of  $\beta_1$ -EC<sub>II</sub>-18AA (Gln-) cyclopeptide mutants in an ELISA competition assay performed with sera from n=69 immunized anti-body-positive rats. About 93% (n=64) of the sera

were efficiently blocked by the  $\beta_1$ -ECII-18AA Cys/Ser mutated cyclopeptide alone (schematically depicted on the top of the left white column), whereas about 7% (n=5) of the sera were blocked by both, the 18AA Cys/Ser- and the 18AA Ser/Cys-mutant (the latter schematically depicted on the top of the right black column).

**Figure 11** is a diagram resuming the dose-dependent (x-axis, abscissa: -fold molar excess of specific peptides) blocking capacity of various linear and cyclic beta1-ECII-peptides given in % of the unblocked antibody-titer (y-axis, ordinate), including 25AA Cys/Cys linear peptides (black squares), 25AA Cys/Ser cyclopeptide mutants (white squares), 18AA Cys/Cys cyclo-peptides (black diamonds), 18AA Cys/Ser cyclopeptide mutants (white diamonds), and 18AA Cys/Ser linear peptide mutants (vertically hatched diamonds) in an ELISA competition assay performed with sera from n=6 randomly chosen immunized antibody-positive rats. All sera were most efficiently blocked by the beta1-ECII-18AA Cys/Ser mutant cyclopeptide followed by the non-mutant 18AA Cys/Cys cyclopeptide and the 25AA Cys/Ser mutant cyclopeptide. All cyclopeptides were largely superior to their linear counterparts (with or without mutation) in terms of their antibody blocking capacities ( $P < 0.005$ ).

**Figure 12** is a diagram resuming the in vivo blocking capacity of in total five (prophylactic) applications of various linear and cyclic beta1-ECII-peptides, started 3 months after the first immunization (and two subsequent beta1-ECII/GST-antigen-boosts, corresponding to a *prevention protocol*). Serum-titers of the beta1-receptor antibodies were determined before and 18-20h after each peptide injection (abscissa, time in months) and are given in % of the corresponding antibody-titers of immunized untreated rats (y-axis, ordinate). The injected peptides were: 25AA Cys/Cys linear peptide (black squares), 25AA Cys/Cys cyclopeptide (white squares), 18AA Cys/Cys cyclopeptide (black diamond), 18AA Cys/Ser cyclopeptide mutant (white diamonds), and the 18AA Cys/Ser linear peptide mutant (vertically hatched diamonds). Also *in vivo*, the efficiency of the cyclic peptides was largely superior to their linear counterparts. The highest efficiency in terms of antibody-neutralization was achieved with 1.0 mg/kg body weight (Bw) of non mutant 25AA Cys/Cys or 18AA Cys/Cys-cyclo-peptides ( $87.7 \pm 2\%$  or  $89.9 \pm 3\%$  decrease after 5 cyclopeptide injections, compared with untreated immunized animals; both  $P < 0.005$ ), followed by the 18AA Cys/Ser mutant cyclopeptide ( $54.5 \pm 2\%$  decrease after 5 cyclopeptide injections;  $P < 0.05$ ), whereas linear 25AA Cys/Cys peptides or linear 18AA Cys/Ser mutants at a same concentration had no significant blocking effects ( $25.8 \pm 3\%$  or  $4.5 \pm 11\%$  decrease after 5 injections,  $P = 0.16$  or  $P = 0.8$ , respectively). Black circles indicate antibody-titers of untreated regularly (every 4 weeks) immunized animals serving as reference, set at 100%.

**Figure 13** shows the results of ELISPOT-assays carried out with B-cells prepared from either the bone marrow (left columns) or the spleen (right columns) of immunized anti-beta1-positive cardiomyopathic untreated animals (Beta1 untreated, black columns) compared with those isolated from immunized anti-beta1 antibody-positive cardiomyopathic animals prophylactically treated with the 25AA-ECII Cys/Cys cyclopeptides (25cyc. Cys/Cys, vertically hatched columns), the 18AA-ECII Cys/Ser cyclopeptide mutant (18cyc. Cys/Ser, diagonally right hatched columns), or the linear 18AA-ECII Cys/Ser peptide mutant (18lin. Cys/Ser, horizontally hatched columns). For the assays, ELISPOT plates were coated overnight with the specific antigen (GST/beta1-ECII-FP) in 0.05 mol/l Tris buffer, pH 9.4.; then the plates were washed 3 times and blocked with BSA for 3 hours at 37°C. Subsequently, the plates were incubated overnight at 37°C with B-cells from either spleen or bone marrow (cultured in RPMI 1640/X-VIVO-15 medium supplemented with 10% fetal calf serum (FCS)) with  $1 \times 10^6$  to  $1 \times 10^3$  cells per well. After 12 hours the B-cells were discarded and the plates with the B-cell secreted IgG bound were washed several times (PBS/0.5% Tween) before the addition of alkaline phosphatase conjugated secondary anti-rat IgG (0.3 µg/ml) to detect bound rat IgG. Then the plates were incubated for another 3 hours at 37°C, washed several times with PBS/0.5% Tween, and developed using LMP/BICP 5:1 (1 ml per well; "LMP" means low melting agarose, and "BICP" means 5-bromo-4-chloro-3-indolyl phosphate *p*-toluidine salt, a blue-colored dye) allowing for a quantification of the blue spots obtained, with each spot representing an antigen-specific IgG secreting spleen or bone-marrow cell.

**Figure 14** is a diagram resuming the in-vivo blocking effect of both 25AA and 18AA cyclo-peptide mutants with a Gln closure site, determined after the first intravenous (i.v.) injection of 1.0 mg/kg body weight (Bw) into immunized antibody-positive rats. Sera were drawn 18-20 hours after i.v. injection of 1.0 or 0.25 mg/kg Bw of the indicated peptides and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen. The first row within the panel represents the group of rats (n=5) treated with the mutant 2 Cys-containing 25AA Cys/Ser (Gln)-cyclopeptide (schematically depicted on the top of the first row, hatched circles), the second and third row represent groups of rats treated with the mutant 18AA Cys/Ser (Gln)-cyclopeptide at two different concentrations (n=40, 1 mg/kg Bw; n=9; 0.25 mg/kg Bw, scheme of the cyclic peptide depicted on the top of the second row, white circles), the fourth row represents n=4 animals treated with the mutant 18AA Ser/Cys (Gln)-cyclopeptide (1 mg/kg Bw, scheme of the cyclic peptide depicted on the top of the fourth row, black diamonds), and the fifth row represents the results obtained with i.v. injected (mutant) 2 Cys-containing linear 18AA Cys/Ser (Gln)-peptides (scheme of the

linear peptide depicted on the top of the fifth row, black circles).

The bars and numbers (in boxes) of each row represent the mean values of the blocking capacity of the respectively indicated peptide given in % of the ELISA-immunoreactivity of the sera before and 18-20 hours after i.v. peptide injection (y-axis).

**Figure 15** is a diagram resuming the *in-vivo* blocking effect of both 25AA and 18AA cyclo-peptide mutants with a Gln closure site, determined after the first intravenous (i.v.) injection of 1 mg/kg Bw or 0.25 mg/kg/Bw into immunized antibody-positive rats. Sera were drawn 18-20 hours after i.v. injection of the various peptides and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys peptide as an antigen.

The graph depicts the relative decrease (or increase) in specific anti- $\beta_1$ -receptor antibody-titers in sera from antibody-positive immunized rats after injection of the various peptides and shows the respective mean value of the blocking capacity of the indicated peptide given in % of the ELISA-immunoreactivity of the sera before and 18-20 hours after i.v. injection (y-axis). The symbols on the right side of the panel represent: white square, 18AA Cys/Ser mutant (Gln-)cyclopeptide (1 mg/kg Bw); white diamond, 18AA Cys/Ser mutant (Gln-)cyclopeptide (0.25 mg/kg Bw); black diamond, 18AA Ser/Cys mutant (Gln-)cyclopeptide (1 mg/kg Bw); horizontally hatched square, 18AA Cys/Ser mutant linear (Gln-)peptide (1 mg/kg Bw); black square, 25AA Cys/Ser mutant (Gln-)cyclopeptide (1 mg/kg Bw). For treasons of charity error bars are not shown in the graph.

**Figure 16A** is a diagramm resuming the *in vivo* blocking effect of both 25AA and 18AA cyclopeptide mutants with a Gln closure site, determined after a total of nine intravenous (i.v.) in-jections of 1.0 mg/kg body weight (Bw) of the indicated peptides into immunized antibody-positive rats. Sera were drawn before and 18-20 hours after i.v. injection of the various peptides every 4 weeks (abscissa: time in months of treatment) and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen.

The graph depicts the relative decrease (or increase) in specific anti-beta1-receptor antibody-titers in sera from antibody-positive immunized rats after injection of the indicated peptides and shows the respective mean value of the blocking capacity of the peptide given in % of the initial ELISA-immunoreactivity before starting treatment (y-axis, ordinate).

The symbols indicate:

black circles, untreated regularly (every 4 weeks) immunized animals (n=9); white diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=20); black squares, 25AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=5).

**Figure 16B** is a diagramm resuming the *in vivo* blocking effect of various concentrations of 18AA cyclopeptide mutants with a Gln closure site, determined after a total of nine intra-venous (i.v.) injections of 0.25, 1.0, 2.0, and 4.0 mg/kg body weight (Bw) into immunized antibody-positive rats, irrespective of the cyclopeptide "responder-state" of individual animals. Sera were drawn before and 18-20 hours after i.v. injection of the various peptides every 4 weeks (abscissa: time in months), and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen.

The graph depicts the relative decrease (or increase) in specific anti-beta1-receptor antibody-titers in sera from antibody-positive immunized rats after injection of the indicated peptides and shows the respective mean values of the blocking capacity of the peptides given in % of the initial ELISA-immunoreactivity before starting treatment (y-axis, ordinate).

The symbols indicate black circles, untreated regularly (every 4 weeks) immunized animals (n=9); white circles, 18AA Cys/Ser mutant (Gln-)cyclopeptide (0.25 mg/kg Bw, n=4);

white diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=20); vertically hatched diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (2.0 mg/kg Bw, n=5); black diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (4.0 mg/kg Bw, n=9).

**Figure 16C** is a diagramm resuming the *in vivo* blocking effect of various concentrations of 18AA cyclopeptide mutants with a Gln closure site, determined after a total of nine intra-venous (i.v.) injections of 0.25, 1.0, 2.0, and 4.0 mg/kg body weight (Bw) into immunized antibody-positive rats, respecting only cyclopeptide-sensitive "responders", defined as animals having , after 7 cyclopeptide-injections, a maximum remaining receptor anti-body level equal or inferior to 80% of the respective titer at start of therapy (compare the curves between. 16c. and Fig. 16b., the latter representing the naturally occuring inhomogenous response of unselected animals). Sera were drawn as described above and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen.

The graph depicts the relative decrease in specific anti-beta1-receptor antibody-titers in sera from antibody-positive immunized responders after injection of the indicated peptides giving the blocking capacity in % of the initial ELISA-

immunoreactivity (y-axis, ordinate).

The symbols indicate (number of responders in bold):

black circles, untreated regularly (every 4 weeks) immunized animals (n=9); white circles, 18AA Cys/Ser mutant (Gln-)cyclopeptide (0.25 mg/kg Bw, n=3/4); white diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=16/20); vertically hatched diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (2.0 mg/kg Bw, n=2/5); black diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (4.0 mg/kg Bw, n=6/9).

**Figure 17.A** is a diagram showing the time course (month 0 to 20) of the internal end-systolic and end-diastolic left ventricular diameters (LVES, LVED) of GST/ $\beta_1$ -EC<sub>II</sub>-immunized un-treated (black circles) versus GST/ $\beta_1$ -EC<sub>II</sub>-immunized animals treated with the indicated various cyclopeptides (see legend)- as determined by echocardiography (echocardiographic system: Visual Sonics, Vevo 770 (version V2.2.3), equipped with a 15-17.5 MHz transducer), whereby LVES/LVED is left ventricular end-systolic diameter/left ventricular end-diastolic diameter.

The symbols indicate:

white circles, untreated 0.9% NaCl-injected non immunized control animals (n=10);

black circles, untreated regularly (every 4 weeks) immunized animals (n=9); white diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=20); black squares, 25AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=5); white squares, 18AA Cys/Ser mutant linear (Gln-)peptide (1.0 mg/kg Bw, n=5).

**Figure 17.B** is a similar diagram showing the time course (month 0 to 20) of the internal end-systolic and end-diastolic left ventricular diameters (LVES, LVED) of GST/ $\beta_1$ -EC<sub>II</sub>-immunized untreated (black circles) versus GST/ $\beta_1$ -EC<sub>II</sub>-immunized animals, treated with different concentrations of the 18AA Cys/Ser cyclopeptide mutant (see legend).

The symbols indicate:

white circles, untreated 0.9% NaCl-injected non immunized control animals (n=10);

black circles, untreated regularly (every 4 weeks) immunized animals (n=9); white squares, 18AA Cys/Ser mutant (Gln-)cyclopeptide (0.25 mg/kg Bw, n=4); white diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=20); vertically hatched diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (2.0 mg/kg Bw, n=5); black diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (4.0 mg/kg Bw, n=9).

**Figure 18** is a diagram indicating the titer course (month 0 to 9) of specific anti- $\beta_1$ -EC<sub>II</sub> antibodies in GST/ $\beta_1$ -EC<sub>II</sub>-immunized versus 0.9% NaCl-injected rats, whereby "Beta1" means immunized animals (before starting treatment with peptides according to the present invention), and "NaCl controls" means corresponding NaCl-injected control animals.

**Figure 19A** is a diagram depicting the time course (month 0 to 20) of the "Cardiac index" (CI) in ml/min/g (body weight) as determined by echocardiography (echocardiographic system see legend to figure 17a.).

The symbols indicate:

white circles, untreated 0.9% NaCl-injected non immunized control animals (n=10);

black circles, untreated regularly (every 4 weeks) immunized animals (n=9); white diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=20); black squares, 25AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=5); white squares, 18AA Cys/Ser mutant linear (Gln-)peptide (1.0 mg/kg Bw, n=5).

**Figure 19B** is a similar diagram showing the time course (month 0 to 20) of the "Cardiac index" (CI) in ml/min/g (body weight) as determined by echocardiography (echocardiographic system see legend to figure 17a.).

The symbols indicate:

white circles, untreated 0.9% NaCl-injected non immunized control animals (n=10);

black circles, untreated regularly (every 4 weeks) immunized animals (n=9); white squares, 18AA Cys/Ser mutant (Gln-)cyclopeptide (0.25 mg/kg Bw, n=4); white diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (1.0 mg/kg Bw, n=20); vertically hatched diamonds, 18AA Cys/Ser mutant (Gln-)cyclopeptide (2.0 mg/kg Bw, n=5).

**Figure 20** shows three rows of panels (a, b, c) with hemodynamic parameters obtained in the therapy study after 10 months of treatment, in detail in the first row shows in each panel of the first row (a) on the left side the heart frequency (HF) given in beats per minute (=bpm), and on the right side the LV systolic blood pressure (LV press.) given in mmHg; in each panel of the second row (b) on the left side the contractility (+ dP/dt) in mmHg/s, and on the right side the relaxation (-dP/dt) in -mmHg/s; the third row (c) shows the left ventricular end-diastolic pressure (LVEDP) as determined by cardiac catheterization in mmHg.

Left and right panels within each row separate data obtained with (left panels, constantly 1.0 mg/kg Bw of the different peptides) cyclic 18AA Cys/Ser (diagonally right hatched columns, n=20 animals), linear Cys/Ser mutants (horizontally

hatched columns, n= 5 animals), and cyclic 25AA Cys/Ser mutants (vertically hatched columns, n= 5 animals). Columns in the right panels represent data obtained with various concentrations of the cyc18AA Cys/Ser mutant; with white filled, black dotted columns corresponding to 0.25 mg/kg Bw (n= 4 animals), diagonally right hatched columns to 1.0 mg/kg Bw (n= 10 animals), diagonally left hatched columns to 2.0 mg/kg Bw (n= 5 animals), and black structured columns to 4.0 mg/kg/Bw (n= 9 animals). Black and white columns in each panel serve as a reference and correspond to either untreated regularly (every 4 weeks) immunized animals (positive control, black, n= 9), or to 0.9% NaCl-injected non immunized control animals (negative control, white, n=10).

In the legends "Beta1 untreated" means immunized anti-beta1 antibody positive cardiomyopathic not treated animals (n=9, black columns), "Controls" means the 0.9% NaCl-injected control group (n=10, white columns), "18cyc Cys/Ser." means immunized anti-beta1-positive cardiomyopathic animals therapeutically treated with the indicated linear 18lin Cys/Ser (n=5 [1.0 mg/kg Bw] or cyclic 18cyc Cys/Ser mutants (n=10 [1.0 mg/kg Bw], or cyclic 25AA Cys/Ser peptide mutants (n=4 [1.0 mg/kg Bw] after 9 months of immunization. Panels on the right side depict the effects of different doses of intravenously injected beta1-ECII 18AA Cys/Ser cyclopeptide mutants (n=4 [0.25 mg/kg Bw]; n=20 [1.0 mg/kg Bw], n=5 [2.0 mg/kg Bw], and n=9 [4.0mg/kg Bw], respectively). Differences between the groups were assessed by one way ANOVA; n.s. = not significant, \* $P < 0.05$ , \*\* $P < 0.005$ .

**Figure 21** shows two panels (a and b) with macro anatomic parameters of the animals from the therapy study as columns:

Upper panels (a.) show the relative wet weights of the indicated organs (from the left to the right: heart, spleen, right kidney, left kidney, lung, and liver) given in g/kg body weight, whereby "Beta1 untreated" means immunized anti-beta1 antibody positive cardiomyopathic not treated animals (n= 9, black columns), "Controls" means the 0.9% NaCl-injected control group (n=10, white columns), "18cyc Cys/Ser" means immunized anti-beta1-positive cardiomyopathic animals therapeutically treated with the indicated linear 18lin Cys/Ser (n=5 [1.0 mg/kg Bw], horizontally hatched columns), or cyclic 18cyc Cys/Ser mutants (n=20 [1.0 mg/kg Bw], diagonally right hatched columns), or cyclic 25AA Cys/Ser peptide mutants (n=4 [1.0 mg/kg Bw], vertically hatched columns) after 9 months of immunization.

Lower panels (b.) show the relative wet weights of the indicated organs (from the left to the right: heart, spleen, right kidney, left kidney, lung, and liver) given in g/kg body weight of immunized anti-beta1-positive cardiomyopathic animals therapeutically treated with the indicated doses of beta1-ECII 18AA Cys/Ser cyclopeptide mutants (n=4 [0.25 mg/kg Bw], white filled black dotted columns; n=20 [1.0 mg/kg Bw] diagonally right hatched columns, n=5 [2.0 mg/kg Bw], diagonally left hatched columns; n=9 [4.0mg/kg Bw], black structured columns) after 9 months of immunization, whereby "Beta1 untreated" means immunized anti-beta1 antibody positive cardiomyopathic not treated animals (n= 9, black columns), and "Controls" means the 0.9% NaCl-injected control group (n=10, white columns).

Kidney R means right and Kidney L means left. Differences between the groups were assessed by one way ANOVA; n.s. = not significant, \* $P < 0.05$ , \*\* $P < 0.005$ .

**Figure 22** shows two panels (a and b) with different laboratory parameters determined in the serum of animals after 10 months of treatment. "Beta1 untreated" and "Controls" in both panels means immunized anti-beta1 antibody positive cardiomyopathic not treated animals (n=5, black columns, positive control), and 0.9% NaCl-injected controls (n=6, white columns, negative control), respectively.

Upper panels (a) show the parameters of immunized anti-beta1-positive cardiomyopathic animals therapeutically treated with the indicated linear 18lin Cys/Ser (n=5 [1.0 mg/kg Bw], horizontally hatched columns), or cyclic 18cyc Cys/Ser mutants (n=20 [1.0 mg/kg Bw], diagonally right hatched columns), or cyclic 25AA Cys/Ser peptide mutants (n=4 [1.0 mg/kg Bw], vertically hatched columns) after 9 months of immunization.

Lower panels (b) show the parameters of immunized anti-beta1-positive cardiomyopathic animals therapeutically treated with the indicated doses of beta1-ECII 18AA Cys/Ser cyclo-peptide mutants (n=4 [0.25 mg/kg Bw], white filled black dotted columns; n=20 [1.0 mg/kg Bw] diagonally right hatched columns, n=5 [2.0 mg/kg Bw], diagonally left hatched columns; n=9 [4.0mg/kg Bw], black structured columns) after 9 months of immunization; Crea means creatinine; GOT means glutamic oxaloacetic transaminase; GPT means glutamic pyruvate transaminase; LDH means lactate dehydrogenase.

**Figure 23** shows the distribution pattern of texas red (fluorochrom-)labeled 18AA-ECII Cys/Ser cyclopeptide mutants ("CP-1") after i.v.-injection of 1.0 mg/kg body weight (Bw) of the labeled cyclopeptide into either non-immunized 0.9% NaCl treated control animals (left panel) or immunized antibody-positive cardiomyopathic Lewis-rats (550g Bw). The photographs depict the subcellular distribution of texas red-labeled 18AA Cys/Ser cyclopeptide mutants in the kidney (2 $\mu$ m sections of the cortical kidney region). The images show, that no toxicity on the kidney was exerted by the cyclic peptides of the invention and no mechanical obstruction of glomerular membranes was observed.

**Figure 24** is a diagram depicting the scheme of mutated cysteine-containing beta1-ECII-homologous cyclo-peptides (amino-acids (AA) are represented as white balls with the corresponding AA letter code written in each ball). Cysteine molecules and their substitutes are depicted as black balls. The assumed localization of the disulfide bridge is represented by a bold black line.

Left side: scheme depicting the original sequence of the ECII-loop of the human beta1 adrenergic receptor; middle: cyclic 22AA ECII-homologous peptide with the glycine mutation at the assumed ring closure site (Position 222).

The right panel depicts an example of a cyclic 22AA peptide-mutant containing only two cysteines (i.e., position 209 and 215). The Cys/Ser mutant of the cysteine at position 216 is shown (Cyclic 22AA beta1-ECII peptide Cys<sub>216</sub>→Ser<sub>216</sub>).

Numbers given indicate the numbering of the amino-acids in the original primary sequence according to Frielle et al. 1987, PNAS 84, pages 7920-7924.

**Figure 25** shows two panels demonstrating the high pressure liquid chromatography (HPLC) elution profiles of two cyclic (22+1)=22 AA peptides; the first panel corresponds to the 3 cysteine-containing construct cyc22AA Cys/Cys (25A), and the second to the 2 cysteine-containing mutant cyc22AA Cys/Ser (25B) of the present invention, all of them cyclopeptides with a Gly closure site. HPLC was carried out in a Hewlett Packard Series 1050 analytical HPLC-system (Agilent Technologies Germany GmbH, Böblingen) equipped with a dual wavelength UV absorbance detector; absorbance was read at 216 nm. After peptide-synthesis and cyclization, the samples were dissolved in H<sub>2</sub>O/0.1% tri-fluoro-acid (TFA) and loaded on a analytical HPLC-column (Waters GmbH, Eschborn) XBridge BEH130, C18, 3,5 µm (column length 50 mm, lumen 4.6 mm) with a flow of 2 ml/min; then a separation-gradient from 0% to 75% acetonitril (ACN) in the pre-sence of 0.1% TFA was run over 5 minutes.

The fractions containing the cyclic beta1-ECII-22AA Cys/Cys peptides with three freely accessible cysteine molecules exhibit the typical mountain-like elution pattern indicating the presence of cystein-racemates (25A, elution between 2.6 and 3.8 minutes). In contrast, the mutant cyclic 22AA peptide containing only two cysteines (connected by a second reinforced disulfide bridge) gave a sharp single elution peak appearing at 2.63 (cyc22AA Cys/Ser, 25B).

**Figure 26** shows two representative panels depicting the characterization of the 22AA-ECII cyclic peptides by mass spectroscopy (MALDI). The first panel corresponds to the 3 cysteine-containing construct cyc22AA Cys/Cys (26A), and the second to the 2 cysteine-containing mutant cyc22AA Cys/Ser (26B) of the present invention, all of them cyclopeptides with a Gly closure site. The panels show representative-MALDI-tracings of the indicated cyclic beta1-ECII 22AA peptides.

The ordinate of each graph shows measured signal intensities ("a.u." means arbitrary units), the abscissa indicates the molecular mass (given in m/z). 26A corresponds to the cyc22AA Cys/Cys-peptide (2518.28 m/z) and 26B to the cyc22AA Cys/Ser-mutant (2502.31 m/z). The MALDI-analysis was carried out using a reflex II-mass spectroscope (Bruker Daltonic GmbH, Bremen), equipped with a Scout-26 sample carrier. In each case the simply protonated molecule was analyzed at 2200 m/z.

**Figure 27A** and **B** depict the *in vitro* blocking (=neutralization) capacity of various cysteine-containing cyclopeptide variants of the second extracellular loop (ECII) of the human beta1-adrenergic receptor, determined by testing n=6 individual sera (27A) of immunized beta1-ECII-antibody-positive rats after over-night incubation with the indicated cyclopeptides (12-14h, 4°C) by ELISA. Columns in 27A represent the receptor-antibody blocking efficiency of the indicated cyclopeptides in % of the antibody-(ELISA)-signals obtained with unblocked antibody-positive rat sera. Columns in 27B represent the mean blocking efficiency for each cyclopeptide, error bars indicate ± SEM.

white columns: cyc18AA Cys/Ser mutant (blocking-efficiency 60.0±8.3%, P=0.0014 when tested for significance against unblocked sera by two-sided t-test); vertically hatched columns: cyc 18AA Cys/Cys (blocking-efficiency 66.1 ±7.0%, P=0.00025); black columns: cyc 22AA Cys/Cys (blocking-efficiency 82.0±5.0%, P=0.00046); diagonally (right) hatched columns: cyc 22AA Cys/Ser mutant (blocking-efficiency 74.9±5.0%, P=0.00026); horizontally hatched columns: cyc25AA Cys/Cys (blocking-efficiency 73.4±5.0%, P=0.00011).

**Figure 28A** and **B** depict the *in vivo* blocking (=neutralization) capacity of two cysteine-containing 18AA or 22AA cyclopeptide-mutants of the second extracellular loop (ECII) of the human beta1-adrenergic receptor upon therapeutic injection of the different constructs into rats regularly immunized since 8 months (first = basic immunization followed by 7 antigen-boosts every 4 weeks). The effects of four to five subsequent cyclopeptide-injections every 4 weeks are shown. Fig. 28A depicts the mean values ± SEM of each of the treated groups of immunized beta1-ECII-antibody positive cardiomyopathic rats (animal number per group is given in the legend).

28A shows the mean effect of 4 subsequent cyclopeptide-injections, determined 20-22 hours after application of the indicated constructs. The remaining receptor antibody-titers after each injection in % of the antibody-titers at start therapy (month 8) are depicted (columns). Error bars indicate ± SEM. Numbers in columns indicate the number

of (subsequent) monthly injection.

Black columns: untreated antibody-positive animals (reference-titer after in total 8+4 (=12) antigen-boosts (compared to the titer at start of therapy)  $110.7 \pm 5.6\%$ ; n= 5, positive control).

White columns: cyc18AA Cys/Ser mutant, n= 5 animals (antibody-titer remaining after 4 injections in per cent of the titer at start of therapy:  $76.0 \pm 23.0\%$ ,  $P= 0.44$  when tested for significance against the antibody-titer of untreated antibody-positive animals by two-sided *t*-test).

Diagonally (right) hatched columns: cyc 22AA Cys/Ser mutant, n=8 animals (antibody-titer remaining after 4 injections in per cent of the titer at start of therapy:  $9.0 \pm 2.2\%$ ,  $P= 3.0 \times 10^{-7}$ ).

**Figure 28B** depicts the time course of antibody-titers after 4 subsequent cyclopeptide-injections, determined *before* and 20-22 hours *after* application of the indicated constructs. Values are given in per cent of increase or decrease in the respective antibody-titers after each cyclopeptide-injection compared with the antibody-titer determined at start of therapy (month 8).

black circles: untreated antibody-positive animals (n=5, positive control); white squares: cyc18AA Cys/Ser mutant, 4 injections, n= 5 animals; black diamonds: cyc 22AA Cys/Ser mutant, 4 injections, n=8 animals.

**Figure 29A** is a diagram showing the time course (month 0 to 12) of the internal end-systolic and end-diastolic left ventricular diameters (LVES, LVED) of GST/beta1-ECII-immunized untreated (black circles) versus GST/beta1-ECII-immunized animals treated with the indicated various cyclopeptides (see legend) as determined by 2D- and M-mode echocardiography (echocardiographic system: Visual Sonics, Vevo 770 (version V2.2.3), equipped with a 15-17.5 MHz transducer), whereby LVES/LVED is left ventricular end-systolic diameter/left ventricular end-diastolic diameter.

white circles, untreated 0.9% NaCl-injected non immunized control animals (n=5); black circles, untreated regularly (every 4 weeks) immunized antibody-positive animals (n=6); white squares, cyc18AA Cys/Ser mutant (1.0 mg/kg Bw, n=5); black diamonds, cyc 22AA Cys/Ser mutant (Gly-)peptide (1.0 mg/kg Bw, n=8).

**Figure 29B** is a diagram depicting the time course (month 0 to 12) of the "Cardiac index" (CI) in ml/min/g (body weight) as determined by 2D- and Doppler-echocardiography (echocardiographic system see above).

white circles, untreated 0.9% NaCl-injected non immunized control animals (n=5) black circles, untreated regularly (every 4 weeks) immunized antibody-positive animals (n=6)

white squares, cyc18AA Cys/Ser mutant (1.0 mg/kg Bw, n=5);

black diamonds, cyc22AA Cys/Ser mutant (Gly-)peptide (1.0 mg/kg Bw, n=8)

**Figure 30** shows the pattern of accumulated radioactivity in the indicated organs 20 min after i.v.-injection of either non-immunized 0.9% NaCl treated control animals (left panel) or immunized antibody-positive Lewis rats (350-400g Bw) with 0.5-1.0 MBq of iodine131-labeled 18AA-ECII Cys/Ser cyclopeptide mutants. Values are given in % activity of the initially injected radioactivity (ID) per g of organ wet weight.

[0216] The following, non-limiting examples illustrate the invention.

#### Example 1: Synthesis of mutant cyclopeptides

[0217] Three particular examples of the herein disclosed cyclopeptides which can form only one single individual disulfide bond are composed of 18, 22 or 25 amino acids (AA): EC<sub>II</sub>-18AA Cys/Ser mutant (Gln-)cyclopeptide, EC<sub>II</sub>-22AA Cys/Ser mutant (Gly-)cyclopeptide and EC<sub>II</sub>-25AA Cys/Ser mutant (Gln-)cyclopeptide, respectively. The primary sequence is partially homologous to the human sequence of the  $\beta_1$ -AR (amino acid positions 204 through 219, 200 through 220 and 200 through 222, respectively). By restricting conformational flexibility through head-to-tail cyclization of the linear peptide followed by a second (single) disulfide-bond stabilizing cyclization procedure, the 18AA, 22 or 25AA cyclopeptide mutant adopts a conformation which more closely mimics that of the epitope as presented on the surface of the native  $\beta_1$ -EC<sub>II</sub> protein loop. Furthermore, cyclization has often been employed as a tool to prolong the duration of action of peptide, since in general cyclic peptides are more stable to proteolysis than their linear counterparts.

[0218] In detail, the peptide sequence of the Cyclo(K-18-P) Cyclic S-S, Cys/Ser mutant is: Cyclo-Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Gln; Cyclization can occur between Cys<sub>7</sub> and Cys<sub>13</sub> (disulphide bond) and Ala<sub>1</sub> and Gln<sub>18</sub> (ring closure).

[0219] In detail, the peptide sequence of the Cyclo(K-22-P) Cyclic S-S, Cys/Ser mutant is: Cyclo-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Gly; Cyclization can occur between Cys<sub>10</sub> and Cys<sub>16</sub> (disulphide bond) and Arg<sub>1</sub> and Gly<sub>22</sub> (ring closure).

[0220] In detail, the peptide sequence of the Cyclo(K-25-P) Cyclic S-S, Cys/Ser mutant is: Cyclo-Ala-Arg-Ala-Glu-Ser-

Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-Gln; Cyclization can occur between Cys<sub>11</sub> and Cys<sub>17</sub> (disulphide bond) and Ala<sub>4</sub> and Gln<sub>25</sub> (ring closure).

**[0221]** The cyclopeptide mutants of the present invention are first synthesized as linear peptides, and are then cyclized covalently on the backbone by condensation of the C-terminal carboxyl group with the amino group of the N-terminal amino acid. Subsequently, a disulphide bond between cysteine residues 7 and 13 (18mer cyclopeptide), cysteine residues 10 and 16 (22mer cyclopeptide) and cysteine residues 11 and 17 (25mer cyclopeptide) is established.

**[0222]** The linear peptide is assembled by stepwise solid phase peptide synthesis using an Fmoc/tert butyl strategy. Chlorotrityl is used as a starting resin. The first amino acid (Fmoc-Pro-OH) is coupling with DIEA in DMF, the second with PYBOP/HOBT/DIEA in DMF and the following amino acids with diisopropylcarboimide, HOBT in DMF. The peptide quality is monitored online by UV detection. Deprotection / coupling (two-fold excess) procedure is described below:

Table 1: Deprotection / coupling (two-fold excess) procedure

Step	Solvents		Cycle
1	Coupling / DMF	(*) min	Coupling
2	DMF	3 x 1 min	Wash
3	Piperidine 25% / DMF	1 min	Deprotection
4	Piperidine 25% / DMF	2 x 15 min	Deprotections
5	DMF	7x1min	Wash
* coupling time is determined by Kaiser test			

**[0223]** For assembly, the following amino acids were used (exemplarily provided for the 18mer cyclic Cys/Ser peptide mutant):

Table 2: amino acids used for F-moc synthesis of the 18mer cyclic Cys/Ser peptide mutant

Amino acid		
Fmoc-ASP (OtBu)-OH	Fmoc-Asn(Trt)-OH	Fmoc-Tyr(tBu)-OH
Fmoc-Cyc(Trt)-OH	Fmoc-Arg(Pbf)-OH	Fmoc-Ala-OH
Fmoc-Glu(OtBu)-OH	Fmoc-D-Glu(OtBu)-OH	Fmoc-Val-OH
Fmoc-Phe-OH	Fmoc-Ser(tBu)-OH	Fmoc-Lys(Boc)-OH

**[0224]** The fully protected peptide with reactive N-terminal amino- and C-terminal carboxylgroups is cleaved from the resin by treatment with hexafluoroisopropanol/dichloromethane.

**[0225]** Cyclization is carried out thereafter in solution according to the following protocol:

**[0226]** The "head to tail"-cyclization of the protected- peptide- is- performed with PyBOP/NaHCO<sub>3</sub> in high DMF dilution (10 mmol of linear peptide / 1L of DMF). The cyclization is completed after 3 days. After DMF evaporation, the peptide is washed with 5% NaHCO<sub>3</sub>, H<sub>2</sub>O and pure H<sub>2</sub>O. The reaction mixture is cooled down, and the peptide is deprotected excepting the cysteine groups. Afterwards, the partially protected peptide is isolated by precipitation with methyl t-butyl ether.

**[0227]** The crude peptide is pre-purified by liquid chromatography:

Stationary phase: silica C 18, 15 µm, 120 Å  
 Eluant: H<sub>2</sub>O acetonitrile + 0.1 % TFA  
 Detection: UV (210 nm)

**[0228]** The disulfide cyclization is performed in H<sub>2</sub>O (2 mg/mL) with the presence of dimethyl sulfoxide (3%). The cyclization reaction is completed after 3 days.

**[0229]** The peptide is purified by HPLC, using the conditions described above.

**[0230]** The fractions with a purity greater than 95% are pooled. The peptide is exchanged on an ion exchange resin (Dowex 1X2) and the final solution lyophilized. The peptide content is determined by amino acid analysis (Edman sequencing).

**[0231]** The HPLC elution profiles in Figs. 1, 3, 25 and 26 clearly demonstrate the sharp and well defined elution peaks



obtained with mutant cyclopeptides all containing the same single disulfide bond in comparison to the relatively large (mountain-like) elution profile obtained with a 3 Cys-containing 18AA Cys<sub>13</sub>-Cys<sub>14</sub>, 22AA Cys<sub>16</sub>-Cys<sub>17</sub> and 25AA Cys<sub>17</sub>-Cys<sub>18</sub> cyclopeptide.

**[0232]** All following *in vitro* and *in vivo* studies were carried out with these cyclopeptide mutants.

## Example 2: *In vitro* ELISA competition assay

**[0233]** The blocking capacity of  $\beta_1$ -EC<sub>II</sub>-18AA cyclopeptide mutants (Cys<sub>13</sub>-Ser<sub>14</sub> or Ser<sub>13</sub>-Cys<sub>14</sub> mutation having an additional D-Glu→Gln exchange, e.g. at the ring closure site) was compared with the 3 Cys-containing 25AA or 18AA Cys/Cys cyclopeptides after preincubation (12h, 4° C, rotating incubation over-night) of different numbers of sera from immunized antibody-positive rats in an ELISA-competition assay using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen. Figs. 4, 5, and 7-10 show the results from measurements performed with sera of n=12 up to n=82 immunized antibody-positive rats using different cyclopeptides of the present invention.

**[0234]** The results obtained with IgG-fractions isolated from the sera of 12 different immunized antibody-positive rats revealed that, surprisingly, only the Cys/Ser but not the Ser/Cys cyclopeptide mutants were able to significantly block antibody-binding to the  $\beta_1$ -EC<sub>II</sub>-antigen. The blocking effect of the 18AS Cys/Ser mutant cyclopeptide was comparable or (in more than half of the sera analyzed n=8/12) was even more effective than the blocking capacity of the 3 Cys-containing  $\beta_1$ -EC<sub>II</sub>-18AA Cys/Cys cyclopeptide (Fig. 4). This finding was confirmed by dose-titration studies demonstrating a clear dose dependent blocking effect of the  $\beta_1$ -EC<sub>II</sub>-18AA Cys/Ser cyclopeptide mutant, which at two to eight-fold molar excess (assuming a 1:1 stoichiometry for both cyclic (2.1 kDa molecular mass (MM)) or linear peptides (3.0 kDa MM), considering a molecular weight of 150 kDa for an IgG-molecule) was consistently largely superior in blocking receptor antibodies than either its Ser/Cys mutated counterpart or a 3 Cys-containing linear 25AA Cys/Cys peptide (Fig. 5).

**[0235]** Fig. 7 resumes the blocking effect of different cyclopeptide mutants after preincubation with IgG isolated from n=78 immunized antibody-positive rats in an ELISA-competition assay with the linear 3 Cys-containing 25AA Cys/Cys-peptide as an antigen. The blocking capacity of 18AA Cys/Ser mutants was highest (69±2%), followed by 25AA Cys/Cys (67±2%), 18AA Cys/Cys cyclopeptides (60±4%), and 3 Cys-containing linear 25AA peptides (55±4%) compared with the totally inefficient cyclic 25AA or 18AA Ser/Cys mutants (6±2% or 1±2%, respectively;  $P < 5 \times 10^{-30}$  or  $P < 3.7 \times 10^{-33}$ ). Only less than 5% of the tested rat sera (n=4) were also partially blocked by the cyclic EC<sub>II</sub>-18AA Ser/Cys mutants, whereas the large majority (>95%; n=78) was efficiently blocked by the EC<sub>II</sub>-18AA Cys/Ser mutant (Fig. 8).

**[0236]** The *in vitro* blocking effect of cyclopeptide mutants having either a Gln-Ala closure site or an D-Glu-Ala closure site was analyzed by ELISA-competition assay (with the 3 Cys-containing linear 25AA Cys/Cys peptide as an antigen) after preincubation with sera isolated from 69 different immunized antibody-positive rats. The results are shown in Fig. 9.

**[0237]** Again, there were two patterns of reaction of the rat sera tested (type1/type2). A major fraction of the sera (93%, Fig. 9, upper panel, type1 and Fig. 10) was very efficiently blocked by the 25AA or 18AA Cys/Ser mutant cyclopeptides (69±2% or 68±3% (Gln-closure)/ 69±2% (D-Glu-closure), respectively) which were even superior to the corresponding 3-Cys-containing 25AA or 18AA Cys/Cys cyclopeptide (65±2% or 59±2%, respectively), whereas the 25AA or 18AA Ser/Cys mutants had almost no inhibitory effect, irrespective of the amino-acid at the closure site (6±2%, 25AA Ser/Cys,  $P < 5 \times 10^{-30}$ ; 1±2%, 18AA Ser/Cys with Gln- ( $P < 3.7 \times 10^{-33}$ ) or 1±2% with D-Glu-closure, ( $P < 6 \times 10^{-55}$ ).

**[0238]** A minor fraction of the sera (7%, Fig. 9, lower panel, type 2 and Fig.10) was blocked similarly by Cys/Ser or Ser/Cys mutated peptides - although to a lesser extent in terms of inhibitory capacity; in addition, for both 25AA and 18AA cyclopeptides the mutants were less effective than their 3 Cys-containing 25AA or 18AA Cys/Cys counterparts. Blocking capacity of the 25AA or 18AA Cys/Ser mutant cyclopeptides was 48±6% or 50±8% (Gln-closure)/47±5% (D-Glu-closure), respectively, which was constantly inferior to that of the corresponding 3-Cys-containing 25AA or 18AA Cys/Cys cyclopeptides (72±5% or 67±6%, respectively); however, in these animals the 25AA or 18AA Ser/Cys mutants revealed blocking capacities which were almost comparable to those of Cys/Ser-mutants (37±7% 25AA Ser/Cys,  $P=0.22$  n.s.; 47±3% 18AA Ser/Cys with Gln- or 33±5% with D-Glu-closure,  $P=0.7$  n.s. or  $P=0.08$  n.s., respectively).

**[0239]** Subsequently, the dose-dependent blocking capacity of various linear and cyclic beta1-EC<sub>II</sub>-peptides *in vitro* was analyzed by using the same ELISA competition assay (Fig. 11):

**[0240]** Experiments including linear 25AA Cys/Cys peptides, cyclic 25AA Cys/Ser peptide mutants, cyclic 18AA Cys/Cys peptides, cyclic 18AA Cys/Ser peptide mutants and a linear 18AA Cys/Ser peptide mutant revealed, that all sera from n=6 randomly chosen immunized antibody-positive rats were best blocked in a dose-dependent manner by beta1-EC<sub>II</sub>-18AA Cys/Ser mutant cyclopeptides, followed by non-mutant 18AA Cys/Cys cyclopeptides, and the 25AA Cys/Ser cyclopeptide mutant. All cyclopeptides were largely superior to their linear- counterparts (with or without mutation) in terms of antibody neutralizing capacity ( $P < 0.005$ ; Fig. 11), yielding a dose-dependent decrease in circulating free anti-beta1-EC<sub>II</sub> antibodies of 53% with an 8-fold excess, 66% with an 20-fold excess and about 85% with an 80-fold excess of 18AA Cys/Ser cyclopeptide mutants. The corresponding results for cyclic 18AA Cys/Cys- or cyclic 25AA Cys/Ser-peptides were: 46/30% [8-fold excess], 56/49% [20-fold excess], and 71/83% at a 80-fold excess. Linear peptides were clearly less efficient, yielding a dose-dependent decrease in receptor antibody-titers of only 24% [8-fold excess], 35%

[20-fold excess], and about 50% at a 80-fold excess for both, linear 25AA Cys/Cys and linear 18AA Cys/Ser peptides.

**[0241]** Additionally, the *in vitro* blocking (=neutralization) capacity of various cyclopeptide variants of the second extracellular loop (ECII) of the human beta1-adrenergic receptor was tested with sera of immunized beta1-ECII antibody-positive rats after incubation for 12-14h at 4°C. The *in vitro* blocking efficiency of the 22AA cyclopeptide cyc22AA Cys/Cys (blocking-efficiency  $82.0 \pm 5.0\%$  versus unblocked sera,  $P=0.000046$ ) and of 22AA cyclopeptide mutants cyc22AA Cys/Ser (blocking-efficiency  $74.9 \pm 5.0\%$ ,  $P=0.00026$ ) was even higher than the blocking capacity of previously described 3 Cys-containing cyclopeptides, i.e., cyc25AA Cys/Cys (blocking-efficiency  $73.4 \pm 5.0\%$ ,  $P=0.00011$ ) or cyc18AA Cys/Cys (blocking-efficiency  $66.1 \pm 7.0\%$  versus unblocked sera,  $P=0.00025$ ; see Figs. 27A/B)

### Example 3: In vitro functional FRET-assay

**[0242]** The blocking capacity of  $\beta_1$ -EC<sub>II</sub> 25AA or 18AA cyclopeptide mutants (having a D-Glu/Gln at the ring closure site or not) on  $\beta_1$ -receptor-mediated signalling (functional cAMP-assay) was assayed using an approach by fluorescence resonance energy transfer (FRET) (Fig. 6).

**[0243]** The effect of the pre-incubation (12h, 4° C, rotating incubator) of anti- $\beta_1$ -EC<sub>II</sub> IgG antibodies of a representative rat with  $\beta_1$ -EC<sub>II</sub>-18AA cyclopeptide mutants (Cys/Ser or Ser/Cys mutations, respectively) was compared with the inhibitory effect of a 3 Cys-containing 25AA Cys/Cys cyclopeptide or with the effect of anti- $\beta_1$ -EC<sub>II</sub> IgG antibodies not incubated with blocking peptides. The normalized YFP/CFP-ratio of the registered FRET emission signals served to quantify the effect of the cyclopeptide mutants of the present invention in terms of blockade (in per cent) of antibody-induced cellular cAMP-production of transiently Epac1-transfected stably  $\beta_1$ -AR expressing human embryonic kidney cells (HEK 293- $\beta_1$  cells). The x-axis in Fig. 6 corresponds to the registration time given in seconds (s).

**[0244]** The inhibitory effect of  $\beta_1$ -EC<sub>II</sub>-18AA cyclopeptide mutants on the antibody-induced stimulation of  $\beta_1$ -adrenergic transmembrane signalling was analyzed using an approach by fluorescence resonance energy transfer (FRET). Again, also in terms of inhibiting measurable functional antibody-effects (blocking intracellular cAMP-increases) the cyclic  $\beta_1$ -EC<sub>II</sub>-18AA Cys/Ser mutant was largely superior to its Ser/Cys counterpart, and even slightly more effective than a 3 Cys-containing 25AA Cys/Cys cyclopeptide (Fig. 6).

**[0245]** Taken together, the results of the tests performed herein demonstrate that the antibody-blocking capacity of mutated cyclopeptides was not affected by the reduction of the number of amino-acids from a 25-meric to a 18-meric peptide. The results also demonstrate an excellent comparability of 25AA Cys/Cys and 18AA Cys/Cys cyclopeptides with the cyclic 25AA or 18AA Cys/Ser mutants, but not with the cyclic 25AA or 18AA Ser/Cys mutants. Surprisingly, the exact nature of the exchange of one single cysteine residue with a serine residue markedly determines the neutralizing potency of the mutated peptides: the Cys→Ser exchange at position 18 (25-AA cyclopeptide) or at position 14 (18-AA cyclopeptide), respectively, yielded cyclic peptides with excellent antibody-neutralizing and pharmacological effects *in vitro* (Figs. 6-10), whereas the Cys<sub>17</sub>→Ser<sub>17</sub> or Cys<sub>13</sub>→Ser<sub>13</sub> mutants (25-AA or 18-AA peptide, respectively) had almost no inhibitory effect, neither regarding their properties as antibody-scavengers nor in terms of their capability of inhibiting functional antibody-effects (neutralization of receptor-stimulation *in vitro*; Figs. 6-10 and Example 3). The D-Glu/Gln exchange at position 25 (25AA cyclopeptide-mutants) or 18 (18AA cyclopeptide-mutants) did not significantly influence the blocking capacity of the cyclopeptides, regardless of their length (i.e., 25 versus 18 amino-acids; Figs. 7-10).

### Example 4: Animal model, "in vivo" blockade of receptor antibodies

**[0246]** The animal model used in this example and any other example described herein, if not indicated to the contrary, is the human analogue rat model. Prior to evaluating and testing, respectively, this human analogue rat model was treated as described herein-below using the various compounds of the present invention, more particularly compounds of formula VI, VII, VIII and IX, and, as controls, a linear EC<sub>II</sub>-18AA Cys/Ser mutated (Gln<sub>18</sub>-)peptide (with the following amino-acid sequence: Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Gln) and a linear non-mutated 3 Cys-containing EC<sub>II</sub>-25AA Cys/Cys (Gln<sub>25</sub>) peptide (with the following amino-acid sequence: Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Cys-Asp-Phe-Val-Thr-Asn-Arg-Gln).

**[0247]** The *in-vivo* blocking effects of both 25AA and 18AA Cys/Ser mutant cyclopeptides (with a Gln closure site), the 18AA Ser/Cys mutant cyclopeptide, and a mutated linear 18AA Cys/Ser peptide were analyzed after intravenous (i.v.) injection of 1.0 mg/kg body weight (Bw) of each construct into freshly immunized antibody-positive rats (i.e., use of cyclo-peptides in a kind of "prevention" study), with a first cyclopeptide-application 3 months after the initial immunization (and two subsequent boost at months 2 and 3). In total, five prophylactic applications of the various constructs were given at 4-weekly intervals, always two weeks after the monthly continued antigen boost. Sera were drawn 18-20 hours after i.v. injection and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen (Fig. 12).

**[0248]** This first (prophylactic) *in vivo* cyclopeptide-applications demonstrated, that the highest efficiency in terms of antibody-neutralization was achieved with 1.0 mg/kg body weight (Bw) of non mutant 25AA Cys/Cys or 18AA Cys/Cys-

cyclopeptides ( $87.7 \pm 2\%$  or  $89.9 \pm 3\%$  decrease after 5 cyclo-peptide injections, compared with untreated immunized animals; both  $P < 0.005$ ), followed by the 18AA Cys/Ser mutant cyclopeptide ( $54.5 \pm 2\%$  decrease after 5 cyclopeptide injections;  $P < 0.05$ ), whereas linear 25AA Cys/Cys peptides or linear 18AA Cys/Ser mutants at a same concentration had no significant blocking effects ( $25.8 \pm 3\%$  or  $4.5 \pm 11\%$  antibody-titer decrease after 5 injections,  $P = 0.16$  or  $P = 0.8$ ; Fig. 12).

**[0249]** This finding was confirmed by ELISpot analysis of bone marrow and spleen cell preparations of selected cyclopeptide-treated *versus* untreated immunized rats ( $n=3$ ). Fig. 13 shows a significant decrease in the number of specific anti-beta1-ECII antibody-secreting cells (ASC), both in the spleen and - to a lesser extent - in the bone marrow only in rats treated with 25AA Cys/Cys cyclopeptide or with 18AA Cys/Ser cyclopeptide mutants ( $n=3$  or 4, respectively), whereas the linear 18AA Cys/Ser peptide mutant ( $n=3$ ) had no effect on ASC (neither spleen nor bone marrow).

**[0250]** Moreover, the *in vivo* blocking effects of *therapeutically* used 25AA and 18AA Cys/Ser cyclo-peptide mutants, the 18AA Ser/Cys mutant (Gln)-cyclopeptide, and a mutated linear 18AA Cys/Ser peptide were assessed after a first intravenous (i.v.) dose (i.e., 1.0 mg/kg body weight (Bw)) of each construct injected into long term immunized anti-beta1-ECII antibody-positive rats, yet presenting a cardiomyopathic phenotype (after nine months of 1x monthly immunization with the beta1-ECII/GST antigen; Figs. 14-16 and 17). Sera were drawn 18-20 hours after the first i.v. injection of the various constructions and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen. "Therapeutic" application of various cyclopeptides in cardiomyopathic antibody-positive rats revealed a higher *in vivo* blocking capacity of 18AA Cys/Ser cyclopeptide mutants (1mg/kg/Bw) compared with either 25AA Cys/Ser cyclopeptide mutants or the clearly less efficient 18AA Ser/Cys cyclopeptide mutants at a same concentration (Figs. 14 and 15). Again, the *in vivo* efficiency of the 18AA Cys/Ser cyclopeptide mutant was largely superior to that of the linear 18AA Cys/Ser peptide mutant. However, when decreasing the applied dose of cyclic 18AA Cys/Ser mutants to 0.25 mg/kg body weight (Bw), no relevant decrease in receptor-antibodies was achieved, suggesting a dose-and-effect relation for cyclopeptide mutants.

**[0251]** Repeated therapeutic injections of mutant single S-S cyclopeptides every 4 weeks into long-term immunized rats with antibody-induced immune-cardiomyopathy confirmed a kind of "critical minimal dose"-and-effect relation for single S-S cyclopeptide mutants: a dose of 0.25 mg/kg Bw of the 18AA Cys/Ser cyclopeptide-albeit capable of scavenging receptor-anti-bodies to some extent - was clearly less efficient in terms of both, (1) the achieved decrease in circulating receptor-antibodies (even when respecting only cyclopeptide-sensitive-"responders", defined as animals having, after 7 cyclopeptide-injections, a maximum remaining antibody-level equal or inferior to 80% of the titer at start of therapy (Figs. 16b and c), and (2) in the achieved cardioprotective effect (Figs. 17b, 19b, 20, and 21b) compared with either a dose of 1.0 or 2.0 mg/kg body weight 18AA Cys/Ser cyclopeptide. The latter doses were almost equally efficient in terms of both, neutralization of circulating receptor antibodies (Figs. 16b and c), and reversal of antibody-induced cardiomyopathic features (Figs. 17b, 19b, 20, and 21 b). A further increase in the applied dose to 4.0 mg/kg body weight, however, did not result in higher efficiency - neither regarding antibody scavenging capacity (Figs. 16b and c), nor regarding cardioprotective effects (Figs. 17b, 19b, 20, and 21b).

**[0252]** Upon injection of the peptides no serious local or systemic side effects were observed. In addition, after injection of the various mutant cyclopeptides, both the heart rate and the blood pressure of the animals were not affected (Fig. 20a). In addition, no obvious changes in routine laboratory parameters occurred associated with the application of the cyclopeptide-mutants (Fig. 22a and b).

**[0253]** In order to generate anti- $\beta_1$ -receptor antibodies the animals were immunized with a fusion protein containing bacterial glutathione-S-transferase and the sequence of the second extra-cellular loop of the human  $\beta_1$ -adrenergic receptor (GST/ $\beta_1$ -EC<sub>II</sub>). Before treatment of the animals with mutated cyclopeptides according to the present invention, progressive dilated immune cardiomyopathy is observed after 6 to 8 months of regular immunization every four weeks (Fig. 17). All of the immunized animals developed high titers of stimulatory anti- $\beta_1$ -EC<sub>II</sub> antibodies. The specific anti- $\beta_1$ -EC<sub>II</sub> titer reached a maximum between 6 and 8 months of continuously boosting the animals every 4 weeks, whereas NaCl-injected control animals developed no specific receptor antibodies (Fig. 18).

**[0254]** Such immunized animals were used for the application of mutant cyclopeptides according to the present invention. The *in-vivo* blocking effect of both 25AA and 18AA Cys/Ser mutant cyclopeptides (with a Gln closure site) were determined after a first intravenous (i.v.) injection of 1.0 mg/kg body weight (Bw) (for 18AA Cys/Ser cyclopeptides also 0.25 mg/kg/Bw) into immunized antibody-positive rats. Sera were drawn 18-20 hours after i.v.- injection of the different peptides and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen.

**[0255]** As mentioned, the *in-vivo* blocking effects of both 25AA and 18AA Cys/Ser mutated cyclopeptides (with a Gln closure site), the 18AA Ser/Cys mutant (Gln)-cyclopeptide, and a mutated linear 18AA Cys/Ser peptide were analyzed after a first intravenous (i.v.) injection of 1 mg/kg body weight (Bw) of each construct into immunized antibody-positive rats. Sera were drawn 18-20 hours after i.v. injection and assayed for reactivity by ELISA using the 3 Cys-containing linear 25AA Cys/Cys-peptide as an antigen.

**[0256]** The *in vivo* data confirmed a higher blocking capacity of the 18AA Cys/Ser mutated cyclo-peptides (1mg/kg/Bw) compared with either 25AA Cys/Ser mutants or the clearly less effective 18AA Ser/Cys mutated cyclopeptides at a same

concentration (Figs. 14 and 15). The *in vivo* efficiency of the 18AA Cys/Ser cyclopeptide was also largely superior to that of the linear 18AA Cys/Ser peptide.

**[0257]** Upon injection of the peptides no serious local or systemic side effects were observed. In addition, after injection of the various mutant cyclopeptides, both the heart rate and the blood pressure of the animals were not affected.

**[0258]** However, the *in vivo* data also indicate, that the efficiency of the 18AA Cys/Ser mutated (Gln)-cyclopeptide seems to depend also on the applied dose; an injection of 0.25mg/kg/Bw was less efficient in terms of antibody-neutralization than the same construct given at a concentration of 1 mg/kg/Bw (Figs. 14 and 15).

**[0259]** The *in vitro* findings described in Examples 2 and 3 were generally confirmed *in vivo* (e.g. Figs. 14-16). Interestingly, the difference in the blocking efficiency of the Cys/Ser mutated cyclopeptides compared with that of the linear peptides was even more pronounced *in vivo* (Figs. 5, 7, and 14-16).

**[0260]** However, the *in vivo* data also indicate, that the efficiency of the 18AA Cys/Ser mutated cyclopeptide might equally depend on the applied dose (Figs. 14-16). The obtained results are compatible with a (minimal) dose-and-effect relation for single S-S cyclopeptide mutants: a dose of 0.25 mg/kg of the 18AACys/Ser cyclopeptide mutant was largely less efficient in terms of both, the achieved decrease in circulating receptor-antibodies and in the achieved cardioprotective effect compared with either a dose of 1.0 or 2.0 mg/kg body weight (Bw) (Figs. 14-16). These doses were almost equally efficient in terms of both, neutralization of circulating receptor antibodies and reversal of antibody-induced cardiomyopathic features (Figs. 16-20). A further increase in the applicated dose to 4.0 mg/kg Bw, however, did not result in higher efficiency - neither regarding antibody scavenging (Fig. 16) capacity nor regarding cardioprotective effects *in vivo* (Figs. 17-21).

**[0261]** A high dose (=4.0 mg/kg Bw) of cyc18AA Cys/Ser mutants did not increase the efficiency; instead, it led to an transient increase in antibody-titers, allowing for significant reductions in receptor-antibody titers only after the third or fourth cyclopeptide-injection. Most notably, the effect on the antibody-neutralizing capacity of the different injected concentrations of cyc18AA Cys/Ser mutant cyclopeptides was also confirmed in terms of reversal of antibody-induced cardiomyopathic features in the course of the study with the best cardioprotection achieved by 1.0 or 2.0 mg/kg Bw 18AA Cys/Ser cyclopeptide mutants (Figs. 17B, 19B, 20, 21B).

**[0262]** As mentioned, both 1.0 mg/kg Bw of 25AA-meric Cys/Ser as well as high dose (=4.0 mg/kg Bw) of 18AA-meric Cys/Ser mutants led to an transient increase in antibody-titers, compatible with an initial immune reaction, allowing for significant reductions in receptor-antibody titers only after the third or fourth cyclopeptide-injection (third or fourth month of therapy; see Figs. 16a and b). This phenomenon did not occur with either 1.0 or 2.0 mg/kg doses of 18AA Cys/Ser cyclopeptide mutants, resulting in higher absolute decreases in antibody-titers after 9 months of treatment (1.0 mg/kg:  $-59 \pm 14\%$  or 2.0 mg/kg:  $-59 \pm 12\%$ , respectively,  $P < 0.0005$  versus immunized untreated animals) compared to only  $-37 \pm 13\%$  (1.0 mg/kg 25AA Cys/Ser-CP;  $P = 0.36$  versus immunized untreated animals) or  $-39 \pm 14\%$  (4.0 mg/kg 18AA Cys/Ser-CP,  $P = 0.24$  versus immunized untreated animals) of the respective antibody-titers at start of therapy. A dose of 1.0 or 2.0mg/kg Bw cyc18AA Cys/Ser peptides were thus almost equally efficient in terms of neutralizing circulating receptor antibodies (Figs. 16b and c), and in the course of the study also in terms of reversal of antibody-induced cardiomyopathic features (Figs. 17b, 19b, 20, and 21 b).

**[0263]** In addition, the *in vivo* experiments demonstrated that the antibody-blocking capacity of mutant cyclopeptides is seemingly not affected by a reduction in the number of amino acids from a 25-meric to a 18-meric cyclopeptide; both *in vitro* and *in vivo* data demonstrate an excellent comparability of the two 2 cysteine-containing single disulfide bond 25AA Cys/Ser or 18AA Cys/Ser cyclopeptide mutants (Fig. 17a). It should be noted, however, that both 1.0 mg/kg 25AA-meric Cys/Ser as well as high dose (i.e., 4.0 mg/kg Bw) 18AA-meric Cys/Ser mutants led to an initial transient increase in antibody-titers (Fig. 16a and b), and thus postponed a significant reduction in receptor antibody titers to the third or fourth cyclopeptide-application (third or fourth month of therapy). This phenomenon did not occur with either 1.0 or 2.0 mg/kg Bw doses of 18AA Cys/Ser cyclopeptide mutants.

**[0264]** The *in vivo* blocking (=neutralization) capacity of the cyc22AA Cys/Ser mutants of the second extracellular loop (ECII) of the human beta1-adrenergic receptor was also tested by "therapeutic" injection into rats which had been regularly immunized over 8 months (basic immunization and seven subsequent antigen-boosts every 4 weeks, see Figs. 28A/B), and compared with the effects of the described cyc18AA Cys/Ser-mutant.

**[0265]** After four to five regular cyclopeptide injections every 4 weeks, the titers in untreated antibody-positive animals increased to  $110.7 \pm 5.6\%$  of the values at start of therapy ( $n=5$ , positive controls; reference-titer after in total 8 + 4 (=12) antigen-applications compared with the antibody-titers at month 8). In contrast, 4 injections of cyc22AA Cys/Ser mutants ( $n=8$  animals) decreased the antibody titers to  $9.0 \pm 2.2\%$  of the antibody-titers at start of therapy ( $P = 3.0 \times 10^{-7}$ , when tested for significance against the antibody-titers of untreated antibody-positive animals by two-sided *t*-test). The *in vivo*-efficiency of the cyc22AA-mutants is thus further enhanced compared to the described Cys/Ser cyclopeptide-mutant having a length of 18 amino-acids (cyc18AA Cys/Ser;  $n=5$  animals), which after 4 injections decreased the antibody-titers to  $76.0 \pm 23.0\%$  of the titers at start of therapy ( $P = 0.44$  versus untreated antibody-positive animals, n.s.; see Fig. 28A).

**[0266]** In addition, echocardiographic follow-up data after 4 months of treatment also show a superiority of cyc22AA

mutants compared with the cyc18AA Cys/Ser mutant regarding their cardioprotective effects *in vivo*, as assessed by the decrease in both left ventricular end-diastolic (LVED) and end-systolic (LVES) diameters (Fig. 29A), and an increase in "Cardiac Index" (CI, given in ml/min/g body weight; see Fig. 29B), as determined by 2 dimensional- and Doppler-echocardiography using a Visual Sonics echocardiographic system (Vevo 770, version V2.2.3), equipped with a 17.5 MHz transducer).

**[0267]** Taken together, because the cardioprotective and immunomodulating activity of the ECII-homologous cyclic peptides appears to depend largely on their conformation, an intramolecularly localized disulfide bridge is essential to stabilize and maintain the three-dimensional structure of the construction. In the cyclic 21+1 (=22) AA peptide, the remaining cysteines (i.e. in position 209 and 215, in case 216 has been mutated to Ser) maintain a defined intramolecular distance, further strengthened by introduction of the smallest naturally occurring amino-acid glycine at the (predicted) ring closure site, in order to allow for the formation of a structure-defining intramolecular disulfide bridge.

**[0268]** The present invention refers to the following nucleotide and amino acid sequences:

SEQ ID No. 1:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>14</sub>→Ser<sub>14</sub>); Cyclization may occur between Ala<sub>1</sub> and Gln<sub>18</sub> Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Gln

SEQ ID No. 2:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>18</sub>→Ser<sub>18</sub>); Cyclization may occur between Ala<sub>1</sub> and Gln<sub>25</sub> Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-Gln

SEQ ID No. 3:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>14</sub>→Ser<sub>14</sub>; Gln<sub>18</sub>→DGlu<sub>18</sub>); Cyclization may occur between Ala<sub>1</sub> and DGlu<sub>18</sub> Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-DGlu

SEQ ID No. 4:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>18</sub>→Ser<sub>18</sub>; Gln<sub>25</sub>→DGlu<sub>25</sub>); Cyclization may occur between Ala<sub>1</sub> and DGlu<sub>25</sub> Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-DGlu

SEQ ID No. 5:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>13</sub>→Ser<sub>13</sub>); Cyclization may occur between Ala<sub>1</sub> and Gln<sub>18</sub> Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Ser-Cys-Asp-Phe-Val-Gln

SEQ ID No. 6:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>17</sub>→Ser<sub>17</sub>); Cyclization may occur between Ala<sub>1</sub> and Gln<sub>25</sub> Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Ser-Cys-Asp-Phe-Val-Thr-Asn-Arg-Gln

SEQ ID No. 7:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>13</sub>→Ser<sub>31</sub>; Gln<sub>18</sub>→DGlu<sub>18</sub>); Cyclization may occur between Ala<sub>1</sub> and DGlu<sub>18</sub> Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Ser-Cys-Asp-Phe-Val-DGlu

SEQ ID No. 8:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>17</sub>→Ser<sub>17</sub>; Gln<sub>25</sub>→DGlu<sub>25</sub>); Cyclization may occur between Ala<sub>1</sub> and DGlu<sub>25</sub> Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Ser-Cys-Asp-Phe-Val-Thr-Asn-Arg-DGlu

SEQ ID No. 9:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>14</sub>→Ser<sub>14</sub>) gcngacgaggcgccgctgctacaacgaccccaagtgcSERgacttcgtccar

SEQ ID No. 10:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>18</sub>→Ser<sub>18</sub>) gcncgggaggagagcgacgaggcgccgctgctacaacgaccccaagtgcSERgacttcgtcaccaaccggcar

SEQ ID No. 11:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>14</sub>→Ser<sub>14</sub>; Gln<sub>18</sub>→D<sub>G</sub>L<sub>u</sub><sub>18</sub>) gcngacgaggcgccgctgctacaacgaccccaagtgcSERgacttcgtcgar

SEQ ID No. 12:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>18</sub>-Ser<sub>18</sub>; Gln<sub>25</sub>→D<sub>G</sub>L<sub>u</sub><sub>25</sub>) gcncgggaggagagcgacgaggcgccgctgctacaacgaccccaagtgcSERgacttcgtcaccaaccgggar

SEQ ID No. 13:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>13</sub>→Ser<sub>13</sub>) gcngacgaggcgccgctgctacaacgaccccaagSERTgctgacttcgtccar

SEQ ID No. 14:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>17</sub>→Ser<sub>17</sub>) gcncgggaggagagcgacgaggcgccgctgctacaacgaccccaagSERTgctgacttcgtcaccaaccggcar

SEQ ID No. 15:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>13</sub>→Ser<sub>13</sub>; Gln<sub>18</sub>→D<sub>G</sub>L<sub>u</sub><sub>18</sub>) gcngacgaggcgccgctgctacaacgaccccaagSERTgctgacttcgtcgar

SEQ ID No. 16:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>17</sub>→Ser<sub>17</sub>; Gln<sub>25</sub>→D<sub>G</sub>L<sub>u</sub><sub>25</sub>) gcncgggaggagagcgacgaggcgccgctgctacaacgaccccaagSERTgctgacttcgtcaccaaccgggar

SEQ ID No. 17:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>13</sub>-Ser<sub>3</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>18</sub> Lys-Cys-Ser-Asp-Phe-Val-Gln-Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 18:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>3</sub>→Ser<sub>3</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>25</sub> Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-Gln-Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 19:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>3</sub>→Ser<sub>3</sub>; Gln<sub>7</sub>→D<sub>G</sub>L<sub>u</sub><sub>7</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>18</sub> Lys-Cys-Ser-Asp-Phe-Val-D<sub>G</sub>L<sub>u</sub>-Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 20:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>3</sub>→Ser<sub>3</sub>; Gln<sub>10</sub>→DGlu<sub>10</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>25</sub> Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-DGlu-Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 21:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>2</sub>→Ser<sub>2</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>18</sub> Lys-Ser-Cys-Asp-Phe-Val-Gln-Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 22:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>2</sub>→Ser<sub>2</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>25</sub> Lys-Ser-Cys-Asp-Phe-Val-Thr-Asn-Arg-Gln-Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 23:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>2</sub>→Ser<sub>2</sub>; Gln<sub>7</sub>→DGlu<sub>7</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>18</sub> Lys-Ser-Cys-Asp-Phe-Val-DGlu-Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 24:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>2</sub>→Ser<sub>2</sub>; Gln<sub>10</sub>→DGlu<sub>10</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>25</sub> Lys-Ser-Cys-Asp-Phe-Val-Thr-Asn-Arg-DGlu-Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 25:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>3</sub>→Ser<sub>3</sub>) aagtgcSERgacttcgtccargcngacgaggcgccgctgctacaacgacccc

SEQ ID No. 26:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>3</sub>→Ser<sub>3</sub>) aagtgcSERgacttcgtccaccaaccggcargcncggcgagagcgacgaggcgccgctgctacaacgacccc

SEQ ID No. 27:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>3</sub>→Ser<sub>3</sub>; Gln<sub>7</sub>→DGlu<sub>7</sub>) aagtgcSERgacttcgtcgargcngacgaggcgccgctgctacaacgacccc

SEQ ID No. 28:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA; Cys<sub>3</sub>→Ser<sub>3</sub>; Gln<sub>10</sub>→DGlu<sub>10</sub>) aagtgcSERgacttcgtccaccaaccgggargcncggcgagagcgacgaggcgccgctgctacaacgacccc

SEQ ID No. 29:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (18AA; Cys<sub>2</sub>→Ser<sub>2</sub>) aagSERTgcgacttcgtccargcngacgaggcgccgctgctacaacgacccc

SEQ ID No. 30:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (25AA;

## EP 2 197 900 B9

Cys<sub>2</sub>→Ser<sub>2</sub>) aagSERTgcgacttcgtcaccaaccggcargcncgggaggagcgacgagggcgccgctgctacaacgacccc

SEQ ID No. 31:

5 Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (18AA; Cys<sub>2</sub>→Ser<sub>2</sub>; Gln<sub>7</sub>→DGlu<sub>7</sub>) aagSERTgcgacttcgtcgargcngacgaggcgccgctgctacaacgacccc

SEQ ID No. 32:

10 Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (25AA; Cys<sub>2</sub>→Ser<sub>2</sub>; Gln<sub>10</sub>-DGlu<sub>10</sub>) aagSERTgcgacttcgtcaccaaccgggargcncgggaggagcgacgagggcgccgctgctacaacgacccc

SEQ ID No. 33:

15 Amino acid sequence of an EC<sub>II</sub> epitope bearing portion of human β<sub>1</sub>-AR (16AA; AA positions 204 to 219) DEARRCYNDPKCCDFV

SEQ ID No. 34:

20 Amino acid sequence of an EC<sub>II</sub> epitope bearing portion of human β<sub>1</sub>-AR (23AA; AA positions 200 to 222) RAESDEARRCYNDPKCCDFVTNR

SEQ ID No. 35:

25 Amino acid sequence of an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR DEARR

SEQ ID No. 36:

30 Amino acid sequence of an EC<sub>II</sub> epitope (bearing portion) of human β<sub>1</sub>-AR RAESDEARR

SEQ ID No. 37:

35 Amino acid sequence of an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR DFV

SEQ ID No. 38:

Amino acid sequence of an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR DFVTNR

40 SEQ ID No. 39:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (16AA; Cys<sub>11</sub>→Ser<sub>11</sub>; N-terminal AA: Gln<sub>16</sub> or DGlu<sub>16</sub>); Cyclization may occur between Ala, and Gln<sub>16</sub>/DGlu<sub>16</sub> Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Tyr-Gln/DGlu

45 SEQ ID No. 40:

Amino acid sequence of the human β<sub>1</sub>-AR

50

55



1 mgagvlvlga sepgnlssaa plpdgaataa rllvpasppa slppasesp eplsqgwttag  
 61 mgllmalivl livagnvlvi vaiaktprlq tltnlfimsl asadlvmgll vvpfgativv  
 121 wgrweygsff celwtsvdvl cvtasietlc vialdrylai tspfryqsl1 trararglvc  
 181 tvwaisalvs flpilmhwwr aesdearrcy ndpkccdfvt nrayaiassv vsfyvplcim  
 241 afvylrvfre aqkqvkkids cerrflggpa rppspspspv papapppgpp rpaaaaatap  
 301 langragkrr psrlvalreq kalktlgiim gvftlcwlpf flanvvkafh relvpdrfv  
 361 ffnwlgyans afnpiiycrs pdfrkafggl lccarraarr rhathgdrpr asgclarpgp  
 421 pspgaasdd ddddvvgatp parllepwag cnggaaadsd ssldepcrpg faseskv

SEQ ID No. 41:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (22AA; Cys<sub>17</sub>→Ser<sub>17</sub>); Cyclization may occur between Arg<sub>1</sub> and Gly<sub>22</sub> Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Gly

SEQ ID No. 42:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (22AA; Cys<sub>17</sub>→Ser<sub>17</sub>) cgggcgagagcgacgagggcgccgctgctacaacgacccaagtgcSERgacttcgtcaccGLY

SEQ ID No. 43:

Amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (22AA; Cys<sub>3</sub>→Ser<sub>3</sub>); Cyclization may occur between Lys<sub>1</sub> and Pro<sub>22</sub> Lys-Cys-Ser-Asp-Phe-Val-Thr-Gly-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro

SEQ ID No. 44:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (22AA; Cys<sub>3</sub>→Ser<sub>3</sub>) aagtgcSERgacttcgtcaccGLYcgggcgagagcgacgagggcgccgctgctacaacgacccc

SEQ ID No. 45:

Amino acid sequence of an EC<sub>II</sub> epitope (bearing portion) of human  $\beta_1$ -AR DEARRCYNDPK

SEQ ID No. 46:

Amino acid sequence of an EC<sub>II</sub> epitope (bearing portion) of human  $\beta_1$ -AR ESDEARRCYNDPK

SEQ ID No. 47:

Amino acid sequence of an EC<sub>II</sub> epitope of human  $\beta_1$ -AR AESDEARR

SEQ ID No. 48:

Amino acid sequence of an EC<sub>II</sub> epitope of human  $\beta_1$ -AR DFVT

SEQ ID No. 49:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (19AA; Cys<sub>15</sub>→Ser<sub>15</sub> (18AA; Cys<sub>14</sub>→Ser<sub>14</sub>)) gcnagcgacgagggcgccgctgctacaacgacccaagtgcSERgacttcgtccar

SEQ ID No. 50:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human  $\beta_1$ -AR (19AA;

Cys<sub>15</sub>→Ser<sub>15</sub> (18AA; Cys<sub>14</sub>→Ser<sub>14</sub>); Gln<sub>18</sub>→DGLu<sub>18</sub>) gcnagcgacgaggcgccgctgctacaacgaccccaagtgcSER-gacttcgtcgar

SEQ ID No. 51:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (19AA; Cys<sub>14</sub>→Ser<sub>14</sub> (18AA; Cys<sub>13</sub>→Ser<sub>13</sub>)) gcnagcgacgaggcgccgctgctacaacgaccccaagSERTgcgacttcgtccar

SEQ ID No. 52:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (19AA; Cys<sub>14</sub>→Ser<sub>14</sub> (18AA; Cys<sub>13</sub>→Ser<sub>13</sub>); Gln<sub>18</sub>→DGLu<sub>18</sub>) gcnagcgacgaggcgccgctgctacaacgaccccaagSERTgcgacttcgtcgar

SEQ ID No. 53:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (19AA; Cys<sub>3</sub>→Ser<sub>3</sub> (18AA; Cys<sub>3</sub>→Ser<sub>3</sub>)) aagtgcSERTgacttcgtccargcnagcgacgaggcgccgctgctacaacgacccc

SEQ ID No. 54:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (19AA; Cys<sub>3</sub>→Ser<sub>3</sub> (18AA; Cys<sub>3</sub>→Ser<sub>3</sub>); Gln<sub>7</sub>→DGLu<sub>7</sub>) aagtgcSERTgacttcgtcgargcnagcgacgaggcgccgctgctacaacgacccc

SEQ ID No. 55:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (19AA; Cys<sub>2</sub>→Ser<sub>2</sub>; (18AA; Cys<sub>2</sub>→Ser<sub>2</sub>)) aagSERTgcgacttcgtccargcnagcgacgaggcgccgctgctacaacgacccc

SEQ ID No. 56:

Nucleotide sequence encoding an amino acid sequence homologous to an EC<sub>II</sub> epitope of human β<sub>1</sub>-AR (19AA; Cys<sub>2</sub>→Ser<sub>2</sub>; (18AA; Cys<sub>2</sub>→Ser<sub>2</sub>); Gln<sub>7</sub>→DGLu<sub>7</sub>) aagSERTgcgacttcgtcgargcnagcgacgaggcgccgctgctacaacgacccc

**[0269]** In the nucleotide sequences, "SER" stands for any nucleotide triplet coding for Ser (serine), i.e. for tcn or agy; and "GLY" stands for any nucleotide triplet coding for Gly (Glycine), i.e. for ggn.

**[0270]** n stands for any nucleotide, particularly a, c, g or t, y stands for t or c and r stands for a or g.

**[0271]** As used herein, the sequences of the various peptides are indicated from the N-terminus to the C-terminus, whereby the N-terminus is at the left side and the C-terminus is at the right side of the respective depicted amino acid sequence.

**[0272]** The following additional abbreviations are used herein:

amino acid :	3-letter code:	1-letter code:
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartic acid	Asp	D
Cysteine	Cys	C
Glutamic acid	Glu	E
Glutamine	Gln	Q
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L

(continued)

	amino acid :	3-letter code:	1-letter code:
	Lysine	Lys	K
5	Methionine	Met	M
	Phenylalanine	Phe	F
	Proline	Pro	P
	Serine	Ser	S
10	Threonine	Thr	T
	Tryptophan	Trp	W
	Tyrosine	Tyr	Y
	Valine	Val	V
	Asparagine or aspartic acid	Asx	B
15	Glutamine or glutamic acid	Glx	Z
	Leucine or Isoleucine	Xle	J
	Unspecified or unknown amino acid	Xaa	X

**Additional references cited****[0273]**

American, Heart, Association: 2007. Dallas; Heart disease and stroke statistics - 2007 update. Circulation 115: e69-e171

Anderton 2001 Immunology 104: 367-376

Baba 2004 Eur. Heart J. 25: 1108-1115

Boivin 2005 Eur. J. Heart Fail. 4, suppl.1: 24 (104)

Caforio 2002 Eur. J. Heart Fail. 4: 411-417

Chiale 2001 Circulation 103: 1765-1771

Chien 2000 Oncol. 27: 9-17

Christ 2001 J. Mol. Cell. Cardiol. 33: 1515-1525

Christ 2006 J. Mol. Cell Cardiol. 41: 716-723

Elies 1996 J. Immunol. 157: 4203-4211

Engelhardt 1999 Proc. Natl. Acad. Sci. USA 96: 7059-7064

Eriksson 2003 Nat. Med. 9: 1484-1490

Fabrizio 1994 Drugs Ther. 8: 89-94

Felix 2000 J. Am. Coll. Cardiol. 35: 1590-1598

Ferrari 1995 J. Exp. Med. 182: 59-65

Freedman 2004 J. Clin. Invest. 113: 1379-1382

Fu 1993 J. Clin. Invest. 91: 1964-1968

Göser 2006 Circulation 114: 1693-1702

Hershko 2005 Ann. N.Y. Acad. Sci. 1051: 635-646

Hoerbeke 1996 Int. J. Cardiol. 54: 103-111

Iwata 2001 a J. Am. Coll. Cardiol. 37: 418-424

Iwata 2001 b Circ. Res. 88: 578-586

Jahns 1996 Eur. J. Pharmacol. 316: 111-121

Jahns 1999a J. Am. Coll. Cardiol. 34: 1545-1551

Jahns 1999b Circulation 99: 649-654

Jahns 2000 J. Am. Coll. Cardiol. 36: 1280-1287

Jahns 2004 J. Clin. Invest. 113: 1419-1429

Jahns 2006 Trends Cardiovasc Med 16: 20-24

Khoynezhad 2007 Eur. J. Heart Fail. 9: 120-123

Kühl 2005 Circulation 112: 1965-1970

Li 2006 177: 8234-8240

Limas 1992 Am. Heart J. 123: 967-970

Limas 1996 Int. J. Cardiol. 54: 113-116

Limas 1997 Circulation 95: 1979-1980

- 5 Limas 2004 Am. J. Cardiol. 93: 1189-1191  
 Lohse 2003 Circ. Res. 93: 896-906  
 Luppi 1998 Circulation 98: 777-785  
 MacLellan 2003 Nat. Med. 9: 1455-1456  
 Maekawa 2007 Circulation 115: 5-8  
 Magnusson 1994 Circulation 89: 2760-2767  
 Magnusson 1996 Int. J. Cardiol. 54: 137-141  
 Mahrholdt 2006 Circulation 114: 1581-1590  
 Matsui 1995 Autoimmunity 21: 85-88  
 10 Matsui 2001 Autoimmunity 43: 217-220  
 Mobini 2004 Autoimmunity Rev. 3: 277-284  
 Morita 2005 J. Clin. Invest. 115: 518-526  
 Neumann 1990 J. Am. Coll. Cardiol. 16: 839-846  
 Nikolaev 2007 J. Am. Coll. Cardiol. 50: 423-431  
 15 Okazaki 2003 Nat. Med. 9: 1477-1483  
 Okazaki 2005 Trends Mol Med 11: 322-326  
 Pohlner 1997 Am. J. Cardiol. 80: 1040-1045  
 Rose 1993 Immunol. Today 14: 426-430  
 Rose 2001 J. Clin. Invest. 107: 943-944  
 20 Schultheiss 1985 J. Mol. Cell Cardiol. 17: 603-617  
 Schultheiss 1988 J. Exp. Med. 168: 2102-2109  
 Schulze 1999 Cardiovasc. Res. 44: 91-100  
 Sewald 2002 Peptides: Chemistry and Biology, Wiley-VCH.  
 Smulski 2006 FASEB J. 20: 1396-1406  
 25 Störk 2006 Am. Heart J. 152: 697-704  
 Wallukat 1995 J. Mol. Cell Cardiol. 27: 397-406  
 Wallukat 2002 N. Engl. J. Med. 347: 1806  
 Wang 1996 Blood Pressure 3: 25-27  
 Witebsky 1957 J. Am. Med. Assoc. 164: 1439-1447  
 30 Woodiwiss 2001 Circulation 103: 155-160

## SEQUENCE LISTING

**[0274]**

- 35 <110> Julius-Maximilians-Universität Würzburg  
 <120> Mutant double cyclized receptor peptides inhibiting betal-adrenoceptor antibodies  
 40 <130> M3119 PCT S3  
 <160> 56  
 <170> PatentIn version 3.4  
 45 <210> 1  
 <211> 18  
 <212> PRT  
 <213> Artificial  
 50 <220>  
 <223> Amino acid sequence homologous to an ECII epitope of human betal-AR  
 <400> 1  
 55

# EP 2 197 900 B9

Ala Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys Cys Ser Asp Phe  
1 5 10 15

5 Val Gln

<210> 2

<211> 25

<212> PRT

10 <213> Artificial

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

15 <400> 2

Ala Arg Ala Glu Ser Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys  
1 5 10 15

20

Cys Ser Asp Phe Val Thr Asn Arg Gln  
20 25

25 <210> 3

<211> 18

<212> PRT

<213> Artificial

30 <220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

<221> VARIANT

35 <222> (18)..(18)

<223> /replace = "DGlu"

<400> 3

40

Ala Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys Cys Ser Asp Phe  
1 5 10 15

45 Val Glu

<210> 4

<211> 25

<212> PRT

50 <213> Artificial

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

55 <220>

<221> VARIANT

<222> (25)..(25)

<223> /replace = "DGlu"

# EP 2 197 900 B9

<400> 4

Ala Arg Ala Glu Ser Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys  
1 5 10 15

Cys Ser Asp Phe Val Thr Asn Arg Glu  
20 25

<210> 5  
<211> 18  
<212> PRT  
<213> Artificial

<220>  
<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<400> 5

Ala Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys Ser Cys Asp Phe  
1 5 10 15

Val Gln

<210> 6  
<211> 25  
<212> PRT  
<213> Artificial

<220>  
<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<400> 6

Ala Arg Ala Glu Ser Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys  
1 5 10 15

Ser Cys Asp Phe Val Thr Asn Arg Gln  
20 25

<210> 7  
<211> 18  
<212> PRT  
<213> Artificial

<220>  
<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<220>  
<221> VARIANT  
<222> (18)..(18)  
<223> /replace ="DGlu"

<400> 7

# EP 2 197 900 B9

Ala Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys Ser Cys Asp Phe  
1 5 10 15

5 Val Glu

<210> 8

<211> 25

<212> PRT

10 <213> Artificial

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

15 <220>

<221> VARIANT

<222> (25)..(25)

<223> /replace ="DGlu"

20 <400> 8

Ala Arg Ala Glu Ser Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys  
1 5 10 15

25

Ser Cys Asp Phe Val Thr Asn Arg Glu  
20 25

30 <210> 9

<211> 54

<212> DNA

<213> Artificial

35 <220>

<223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

<221> misc\_feature

40 <222> (3)..(3)

<223> n is a, c, g, t or u

<220>

<221> misc\_feature

45 <222> (40)..(42)

<223> /replace ="tcc"

/replace ="agc"

/replace ="tca"

/replace ="tcg"

50 /replace ="agt"

<220>

<221> misc\_feature

<222> (54)..(54)

55 <223> r ia a or g

<400> 9

gcngacgagg cgcgccgctg ctacaacgac cccaagtgtct ctgacttgt ccar 54

<210> 10  
 <211> 75  
 <212> DNA  
 <213> Artificial

5

<220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

10

<220>  
 <221> misc\_feature  
 <222> (3)..(3)  
 <223> n is a, c, g, t or u

15

<220>  
 <221> misc\_feature  
 <222> (52)..(54)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 20 /replace ="tcg"  
 /replace ="agt"

25

<220>  
 <221> misc\_feature  
 <222> (75)..(75)  
 <223> r is a or g

<400> 10

30

gcncggggcgg agagcgacga ggcgcgcgcg tgctacaacg accccaagtg ctctgacttc 60  
 gtcaccaacc ggcar . 75

35

<210> 11  
 <211> 54  
 <212> DNA  
 <213> Artificial

40

<220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

45

<220>  
 <221> misc\_feature  
 <222> (3)..(3)  
 <223> n is a, c, g, t or u

50

<220>  
 <221> misc\_feature  
 <222> (40)..(42)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 /replace ="tcg"  
 55 /replace ="agt"

<220>  
 <221> misc\_feature



# EP 2 197 900 B9

<222> (54)..(54)  
 <223> r is a or g

<400> 11  
 5 gcngacgagg cgcgccgctg ctacaacgac cccaagtgtct ctgacttcgt cgar 54

<210> 12  
 <211> 75  
 <212> DNA  
 10 <213> Artificial

<220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

15 <220>  
 <221> misc\_feature  
 <222> (3)..(3)  
 <223> n is a, c, g, t or u

20 <220>  
 <221> misc\_feature  
 <222> (52)..(54)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 25 /replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"

30 <220>  
 <221> misc\_feature  
 <222> (75)..(75)  
 <223> r is a or g

35 <400> 12

gcncggggcgg agagcgacga ggcgcgccgc tgctacaacg accccaagtg ctctgacttc 60  
 gtcaccaacc gggar 75

40 <210> 13  
 <211> 54  
 <212> DNA  
 <213> Artificial

45 <220>  
 .<223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

50 <220>  
 <221> misc\_feature  
 <222> (3)..(3)  
 <223> n is a, c, g, t or u

55 <220>  
 <221> misc\_feature  
 <222> (37)..(39)  
 <223> /replace ="tcc"  
 /replace ="agc"

/replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"

5 <220>  
 <221> misc\_feature  
 <222> (54)..(54)  
 <223> r is a or g

10 <400> 13  
 gcngacgagg cgcgccgctg ctacaacgac cccaagtctt gcgacttcgt ccar 54

<210> 14  
 <211> 75

15 <212> DNA  
 <213> Artificial

<220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

20 <220>  
 <221> misc\_feature  
 <222> (3)..(3)  
 <223> n is a, c, g, t or u

25 <220>  
 <221> misc\_feature  
 <222> (49)..(51)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"

30 <220>  
 <221> misc\_feature  
 <222> (75)..(75)  
 <223> r is a or g

35 <400> 14

gcncggggcgg agagcgacga ggcgcgccgc tgctacaacg accccaagtc ttgcgacttc 60

45 gtcaccaacc ggcar 75

<210> 15  
 <211> 54

50 <212> DNA  
 <213> Artificial

<220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

55 <220>  
 <221> misc\_feature  
 <222> (3)..(3)

<223> n is a, c, g, t or u  
 <220>  
 <221> misc\_feature  
 5 <222> (37)..(39)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 /replace ="tcg"  
 10 /replace ="agt"  
 <220>  
 <221> misc\_feature  
 <222> (54)..(54)  
 15 <223> r is a or g  
 <400> 15  
 gcngacgagg cgcgccgctg ctacaacgac cccaagtctt gcgacttcgt cgar 54  
 20 <210> 16  
 <211> 75  
 <212> DNA  
 <213> Artificial  
 25 <220>  
 <223> SEQ ID No. 16:  
 Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR  
 <220>  
 30 <221> misc\_feature  
 <222> (3)..(3)  
 <223> n is a, c, g, t or u  
 <220>  
 35 <221> misc\_feature  
 <222> (49)..(51)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 40 /replace ="tcg"  
 /replace ="agt"  
 <220>  
 <221> misc\_feature  
 45 <222> (75)..(75)  
 <223> r is a or g  
 <400> 16  
 50 gcncggggcgg agagcgacga ggcgcgccgc tgctacaacg accccaagtc ttgcgacttc 60  
 gtcaccaacc gggar 75  
 55 <210> 17  
 <211> 18  
 <212> PRT  
 <213> Artificial

# EP 2 197 900 B9

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<400> 17

5

Lys Cys Ser Asp Phe Val Gln Ala Asp Glu Ala Arg Arg Cys Tyr Asn  
1 5 10 15

10

Asp Pro

<210> 18

<211> 25

15

<212> PRT

<213> Artificial

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

20

<400> 18

25

Lys Cys Ser Asp Phe Val Thr Asn Arg Gln Ala Arg Ala Glu Ser Asp  
1 5 10 15

30

Glu Ala Arg Arg Cys Tyr Asn Asp Pro  
20 25

<210> 19

<211> 18

<212> PRT

<213> Artificial

35

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

40

<221> VARIANT

<222> (7)..(7)

<223> /replace ="DGlu"

<400> 19

45

Lys Cys Ser Asp Phe Val Glu Ala Asp Glu Ala Arg Arg Cys Tyr Asn  
1 5 10 15

50

Asp Pro

<210> 20

<211> 25

55

<212> PRT

<213> Artificial

<220>

# EP 2 197 900 B9

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

<221> VARIANT

<222> (10)..(10)

<223> /replace ="DGlu"

<400> 20

Lys Cys Ser Asp Phe Val Thr Asn Arg Glu Ala Arg Ala Glu Ser Asp  
1 5 10 15

Glu Ala Arg Arg Cys Tyr Asn Asp Pro  
20 25

<210> 21

<211> 18

<212> PRT

<213> Artificial

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<400> 21

Lys Ser Cys Asp Phe Val Gln Ala Asp Glu Ala Arg Arg Cys Tyr Asn  
1 5 10 15

Asp Pro

<210> 22

<211> 25

<212> PRT

<213> Artificial

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<400> 22

Lys Ser Cys Asp Phe Val Thr Asn Arg Gln Ala Arg Ala Glu Ser Asp  
1 5 10 15

Glu Ala Arg Arg Cys Tyr Asn Asp Pro  
20 25

<210> 23

<211> 18

<212> PRT

<213> Artificial

<220>

# EP 2 197 900 B9

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

<221> VARIANT

<222> (7)..(7)

<223> /replace ="DGlu"

<400> 23

Lys Ser Cys Asp Phe Val Glu Ala Asp Glu Ala Arg Arg Cys Tyr Asn  
1 5 10 15

Asp Pro

<210> 24

<211> 25

<212> PRT

<213> Artificial

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

<221> VARIANT

<222> (10)..(10)

<223> /replace ="DGlu"

<400> 24

Lys Ser Cys Asp Phe Val Thr Asn Arg Glu Ala Arg Ala Glu Ser Asp  
1 5 10 15

Glu Ala Arg Arg Cys Tyr Asn Asp Pro  
20 25

<210> 25

<211> 54

<212> DNA

<213> Artificial

<220>

<223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

<221> misc\_feature

<222> (7)..(9)

<223> /replace =" .tcc"

/replace ="agc"

/replace ="tca"

/replace ="tcg"

/replace ="agt"

<220>

<221> misc\_feature

<222> (21)..(21)  
 <223> r is a or g

<220>  
 5 <221> misc\_feature  
 <222> (24)..(24)  
 <223> n is a, c, g, t or u

<400> 25  
 10 aagtgctctg acttcgtcca rgcngacgag gcgcgccgct gctacaacga cccc 54

<210> 26  
 <211> 75  
 <212> DNA  
 15 <213> Artificial

<220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

20 <220>  
 <221> misc\_feature  
 <222> (7)..(9)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 25 /replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"

30 <220>  
 <221> misc\_feature  
 <222> (30)..(30)  
 <223> r is a or g

35 <220>  
 <221> misc\_feature  
 <222> (33)..(33)  
 <223> n is a, c, g, t or u

<400> 26  
 40

aagtgctctg acttcgtcac caaccggcar gcncggggcg agagcgacga ggcgcgccgc 60  
 tgctacaacg acccc 75

45 <210> 27  
 <211> 54  
 <212> DNA  
 <213> Artificial

50 <220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

55 <220>  
 <221> misc\_feature  
 <222> (7)..(9)  
 <223> /replace ="tcc"  
 /replace ="agc"

/replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"

5       <220>  
       <221> misc\_feature  
       <222> (21)..(21)  
       <223> r is a or g

10       <220>  
       <221> misc\_feature  
       <222> (24)..(24)  
       <223> n is a, c, g, t or u

15       <400> 27  
       aagtgctctg acttcgtcga rgcngacgag gcgcgccgct gctacaacga cccc       54

20       <210> 28  
       <211> 75  
       <212> DNA  
       <213> Artificial

25       <220>  
       <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

30       <220>  
       <221> misc\_feature  
       <222> (7)..(9)  
       <223> /replace ="tcc"  
       /replace ="agc"  
       /replace ="tca"  
       /replace ="tcg"  
       /replace ="agt"

35       <220>  
       <221> misc\_feature  
       <222> (30)..(30)  
       <223> r is a or g

40       <220>  
       <221> misc\_feature  
       <222> (33)..(33)  
       <223> n is a, c, g, t or u

45       <400> 28

50       aagtgctctg acttcgtcac caaccgggar gcncgggcgg agagcgacga ggcgcgccgc       60  
       tgctacaacg accccc       75

55       <210> 29  
       <211> 54  
       <212> DNA  
       <213> Artificial

      <220>  
       <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR



<220>  
 <221> misc\_feature  
 <222> (4)..(6)  
 <223> /replace ="tcc"  
 5 /replace ="agc"  
 /replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"

10 <220>  
 <221> misc\_feature  
 <222> (21)..(21)  
 <223> r is a or g

15 <220>  
 <221> misc\_feature  
 <222> (24)..(24)  
 <223> n is a, c, g, t or u

20 <400> 29  
 aagttcttgcg acttcgtcca rgcngacgag gcgcgccgct gctacaacga cccc 54

<210> 30  
 <211> 75  
 25 <212> DNA  
 <213> Artificial.

30 <220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

35 <220>  
 <221> misc\_feature  
 <222> (4)..(6)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"

40 <220>  
 <221> misc\_feature  
 <222> (30)..(30)  
 <223> r is a or g

45 <220>  
 <221> misc\_feature  
 <222> (33)..(33)  
 <223> n is a, c, g, t or u

50 <400> 30  
 aagttcttgcg acttcgtcac caaccgggcar gcncggggcgg agagcgacga ggcgcgccgc 60

55 .tgctacaacg accccc 75

<210> 31  
 <211> 54  
 <212> DNA  
 <213> Artificial  
 5  
 <220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR  
 <220>  
 10 <221> misc\_feature  
 <222> (4)..(6)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 15 /replace ="tcg"  
 /replace ="agt"  
 <220>  
 <221> misc\_feature  
 20 <222> (21)..(21)  
 <223> r is a or g  
 <220>  
 <221> misc\_feature  
 25 <222> (24).. (24)  
 <223> n is a, c, g, t or u  
 <400> 31  
 aagttctgcg actctgctga rgcngacgag gcgcgccgct gctacaacga cccc 54  
 30  
 <210> 32  
 <211> 75  
 <212> DNA  
 <213> Artificial  
 35  
 <220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR  
 <220>  
 40 <221> misc\_feature  
 <222> (4)..(6)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 45 /replace ="tcg"  
 /replace ="agt"  
 <220>  
 <221> misc\_feature  
 50 <222> (30)..(30)  
 <223> r is a or g  
 <220> -  
 <221> misc\_feature  
 55 <222> (33)..(33)  
 <223> n is a, c, g, t or u  
 <400> 32

aagtcttgcg acttcgtcac caaccgggar gcncggggcg agagcgacga ggcgcgccgc 60

tgctacaacg acccc 75

5

<210> 33  
 <211> 16  
 <212> PRT  
 <213> Artificial

10

<220>  
 <223> Amino acid sequence of an ECII epitope of human betal-AR

15

<400> 33

Asp	Glu	Ala	Arg	Arg	Cys	Tyr	Asn	Asp	Pro	Lys	Cys	Cys	Asp	Phe	Val
1				5					10					15	.

20

<210> 34  
 <211> 23  
 <212> PRT  
 <213> Artificial

25

<220>  
 <223> Amino acid sequence of an ECII epitope of human betal-AR

<400> 34

30

Arg	Ala	Glu	Ser	Asp	Glu	Ala	Arg	Arg	Cys	Tyr	Asn	Asp	Pro	Lys	Cys
1				5					10					15	

35

Cys	Asp	Phe	Val	Thr	Asn	Arg
			20			

<210> 35  
 <211> 5  
 <212> PRT  
 <213> Artificial

40

<220>  
 <223> Amino acid sequence of an ECII epitope of human betal-AR

45

<400> 35

Asp	Glu	Ala	Arg	Arg
1			5	

50

<210> 36

<211> 9  
 <212> PRT  
 <213> Artificial

55

<220>  
 <223> Amino acid sequence of an ECII epitope of human betal-AR

<400> 36

5 Arg Ala Glu Ser Asp Glu Ala Arg Arg  
1 5

<210> 37

<211> 3

<212> PRT

10 <213> Artificial

<220>

<223> Amino acid sequence of an ECII epitope of human betal-AR

15 <400> 37

Asp Phe Val

1

20

<210> 38

<211> 6

<212> PRT

<213> Artificial

25

<220>

<223> Amino acid sequence of an ECII epitope of human betal-AR

<400> 38

30

Asp Phe Val Thr Asn Arg

1

5

35

<210> 39

<211> 16

<212> PRT

<213> Artificial

40

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

<221> VARIANT

45

<222> (16)..(16)

<223> /replace ="DGlu"

/replace ="Glu"

<400> 39

50

Ala Arg Arg Cys Tyr Asn Asp Pro Lys Cys Ser Asp Phe Val Tyr Glx

1

5

10

15

55

<210> 40

<211> 477

<212> PRT

<213> Homo sapiens

## EP 2 197 900 B9

&lt;400&gt; 40

5 Met Gly Ala Gly Val Leu Val Leu Gly Ala Ser Glu Pro Gly Asn Leu  
 1 5 10 15  
 Ser Ser Ala Ala Pro Leu Pro Asp Gly Ala Ala Thr Ala Ala Arg Leu  
 20 25 30  
 10 Leu Val Pro Ala Ser Pro Pro Ala Ser Leu Leu Pro Pro Ala Ser Glu  
 35 40 45  
 15 Ser Pro Glu Pro Leu Ser Gln Gln Trp Thr Ala Gly Met Gly Leu Leu  
 50 55 60  
 20 Met Ala Leu Ile Val Leu Leu Ile Val Ala Gly Asn Val Leu Val Ile  
 65 70 75 80  
 Val Ala Ile Ala Lys Thr Pro Arg Leu Gln Thr Leu Thr Asn Leu Phe  
 85 90 95  
 25 Ile Met Ser Leu Ala Ser Ala Asp Leu Val Met Gly Leu Leu Val Val  
 100 105 110  
 30 Pro Phe Gly Ala Thr Ile Val Val Trp Gly Arg Trp Glu Tyr Gly Ser  
 115 120 125  
 Phe Phe Cys Glu Leu Trp Thr Ser Val Asp Val Leu Cys Val Thr Ala  
 130 135 140  
 35 Ser Ile Glu Thr Leu Cys Val Ile Ala Leu Asp Arg Tyr Leu Ala Ile  
 145 150 155 160  
 40 Thr Ser Pro Phe Arg Tyr Gln Ser Leu Leu Thr Arg Ala Arg Ala Arg  
 165 170 175  
 45 Gly Leu Val Cys Thr Val Trp Ala Ile Ser Ala Leu Val Ser Phe Leu  
 180 185 190  
 50 Pro Ile Leu Met His Trp Trp Arg Ala Glu Ser Asp Glu Ala Arg Arg  
 195 200 205

55

EP 2 197 900 B9

	Cys	Tyr	Asn	Asp	Pro	Lys	Cys	Cys	Asp	Phe	Val	Thr	Asn	Arg	Ala	Tyr	
	210						215					220					
5	Ala	Ile	Ala	Ser	Ser	Val	Val	Ser	Phe	Tyr	Val	Pro	Leu	Cys	Ile	Met	
	225					230					235					240	
10	Ala	Phe	Val	Tyr	Leu	Arg	Val	Phe	Arg	Glu	Ala	Gln	Lys	Gln	Val	Lys	
					245					250					255		
15	Lys	Ile	Asp	Ser	Cys	Glu	Arg	Arg	Phe	Leu	Gly	Gly	Pro	Ala	Arg	Pro	
				260					265					270			
20	Pro	Ser	Pro	Ser	Pro	Ser	Pro	Val	Pro	Ala	Pro	Ala	Pro	Pro	Pro	Gly	
				275				280					285				
25	Pro	Pro	Arg	Pro	Ala	Ala	Ala	Ala	Ala	Thr	Ala	Pro	Leu	Ala	Asn	Gly	
	290					295						300					
30	Arg	Ala	Gly	Lys	Arg	Arg	Pro	Ser	Arg	Leu	Val	Ala	Leu	Arg	Glu	Gln	
	305				310						315				320		
35	Lys	Ala	Leu	Lys	Thr	Leu	Gly	Ile	Ile	Met	Gly	Val	Phe	Thr	Leu	Cys	
				325						330					335		
40	Trp	Leu	Pro	Phe	Phe	Leu	Ala	Asn	Val	Val	Lys	Ala	Phe	His	Arg	Glu	
				340					345					350			
45	Leu	Val	Pro	Asp	Arg	Leu	Phe	Val	Phe	Phe	Asn	Trp	Leu	Gly	Tyr	Ala	
			355					360					365				
50	Asn	Ser	Ala	Phe	Asn	Pro	Ile	Ile	Tyr	Cys	Arg	Ser	Pro	Asp	Phe	Arg	
	370					375						380					
55	Lys	Ala	Phe	Gln	Gly	Leu	Leu	Cys	Cys	Ala	Arg	Arg	Ala	Ala	Arg	Arg	
	385					390					395				400		
60	Arg	His	Ala	Thr	His	Gly	Asp	Arg	Pro	Arg	Ala	Ser	Gly	Cys	Leu	Ala	
					405					410					415		
65	Arg	Pro	Gly	Pro	Pro	Pro	Ser	Pro	Gly	Ala	Ala	Ser	Asp	Asp	Asp	Asp	
				420					425					430			
70	Asp	Asp	Val	Val	Gly	Ala	Thr	Pro	Pro	Ala	Arg	Leu	Leu	Glu	Pro	Trp	
			435					440					445				

# EP 2 197 900 B9

Ala Gly Cys Asn Gly Gly Ala Ala Ala Asp Ser Asp Ser Ser Leu Asp  
450 455 460

5

Glu Pro Cys Arg Pro Gly Phe Ala Ser Glu Ser Lys Val  
465 470 475

10

<210> 41  
<211> 22  
<212> PRT  
<213> Artificial

15

<220>  
<223> Amino acid sequence homologous to an ECII epitope of human betal-AR  
  
<400> 41

20

Arg Ala Glu Ser Asp Glu Ala Arg Arg Cys Tyr Asn Asp Pro Lys Cys  
1 5 10 15

Ser Asp Phe Val Thr Gly  
20

25

<210> 42  
<211> 66  
<212> DNA  
<213> Artificial

30

<220>  
<223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

35

<220>  
<221> misc\_feature  
<222> (49)..(51)  
<223> /replace ="tcc"  
/replace ="agc"  
/replace ="tca"  
/replace ="tcg"  
/replace ="agt"

40

45

<220>  
<221> misc\_feature  
<222> (66)..(66)  
<223> n is a, c, g, t or u

<400> 42

50

cgggcggaga gcgacgaggc gcgccgctgc tacaacgacc ccaagtgtc tgacttcgtc 60  
accggn 66

55

<210> 43  
<211> 22  
<212> PRT  
<213> Artificial

# EP 2 197 900 B9

<220>

<223> Amino acid sequence homologous to an ECII epitope of human betal-AR

<400> 43

5

Lys Cys Ser Asp Phe Val Thr Gly Arg Ala Glu Ser Asp Glu Ala Arg  
1 5 10 15

10

Arg Cys Tyr Asn Asp Pro  
20

<210> 44

15

<211> 66

<212> DNA

<213> Artificial

<220>

20

<223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

<220>

<221> misc\_feature

<222> (7)..(9)

25

<223> /replace ="tcc"

/replace ="agc"

/replace ="tca"

/replace ="tcg"

/replace ="agt"

30

<220>

<221> misc\_feature

<222> (24)..(24)

<223> n is a, c, g, t or u

35

<400> 44

aagtgtctctg acttcgtcac cggncggggcg gagagcgacg aggcgcgcgcg ctgctacaac 60

40

gacccc 66

<210> 45

<211> 11

45

<212> PRT

<213> Artificial

<220>

<223> Amino acid sequence of an ECII epitope (bearing portion) of human betal-AR

50

<220>

<221> misc\_feature

<222> (2)..(3)

<223> Xaa can be any naturally occurring amino acid

55

<220>

<221> misc\_feature

<222> (7)..(7)



# EP 2 197 900 B9

<223> Xaa can be any naturally occurring amino acid

<400> 45

5

Asp Xaa Xaa Arg Arg Cys Xaa Asn Asp Pro Lys  
1 5 10

<210> 46

10

<211> 13

<212> PRT

<213> Artificial

<220>

15

<223> Amino acid sequence of an ECII epitope (bearing portion) of human betal-AR

<220>

<221> misc\_feature

<222> (4)..(5)

20

<223> Xaa can be any naturally occurring amino acid

<220>

<221> misc\_feature

<222> (9)..(9)

25

<223> Xaa can be any naturally occurring amino acid

<400> 46

30

Glu Ser Asp Xaa Xaa Arg Arg Cys Xaa Asn Asp Pro Lys  
1 5 10

<210> 47

<211> 8

35

<212> PRT

<213> Artificial

<220>

<223> Amino acid sequence of an ECII epitope of human betal-AR

40

<400> 47

45

Ala Glu Ser Asp Glu Ala Arg Arg  
1 5

<210> 48

<211> 4

<212> PRT

50

<213> Artificial

<220>

<223> Amino acid sequence of an ECII epitope of human betal-AR

55

<400> 48

EP 2 197 900 B9

Asp Phe Val Thr  
1

5       <210> 49  
          <211> 57  
          <212> DNA  
          <213> Artificial

10       <220>  
          <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

          <220>  
          <221> misc\_feature

15       <222> (3)..(3)  
          <223> n is a, c, g, t or u

          <220>  
          <221> misc\_feature

20       <222> (43)..(45)  
          <223> /replace ="tcc"  
          /replace ="agc"  
          /replace ="tca"  
          /replace ="tcg"

25       /replace ="agt"

          <220>  
          <221> misc\_feature

30       <222> (57)..(57)  
          <223> r is a or g

          <400> 49  
          gcnagcgacg aggcgcgcgcg ctgctacaac gaccccaagt gctctgactt cgtccar       57

35       <210> 50  
          <211> 57  
          <212> DNA  
          <213> Artificial

40       <220>  
          <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

          <220>  
          <221> misc\_feature

45       <222> (3)..(3)  
          <223> n is a, c, g, t or u

          <220>  
          <221> misc\_feature

50       <222> (43)..(45)  
          <223> /replace ="tcc"  
          /replace ="agc"  
          /replace ="tca"  
          /replace ="tcg"

55       /replace ="agt"

          <220>  
          <221> misc\_feature

# EP 2 197 900 B9

<222> (57)..(57)

<223> r is a or g

<400> 50

5 gcnagcgacg aggcgcgccg ctgctacaac gaccccaagt gctctgactt cgtcgar 57

<210> 51

<211> 57

<212> DNA

10 <213> Artificial

<220>

<223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

15 <220>

<221> misc\_feature

<222> (3)..(3)

<223> n is a, c, g, t or u

20 <220>

<221> misc\_feature

<222> (40)..(42)

<223> /replace ="tcc"

/replace ="agc"

25 /replace ="tca"

/replace ="tcg"

/replace ="agt"

<220>

30 <221> misc\_feature

<222> (57)..(57)

<223> r is a or g

<400> 51

35 gcnagcgacg aggcgcgccg ctgctacaac gaccccaagt cttgcgactt cgtccar 57

<210> 52

<211> 57

<212> DNA

40 <213> Artificial

<220>

<223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

45 <220>

<221> misc\_feature

<222> (3)..(3)

<223> n is a, c, g, t or\_u

50 <220>

<221> misc\_feature

<222> (40)..(42)

<223> /replace ="tcc"

/replace ="agc"

55 /replace ="tca"

/replace ="tcg"

/replace ="agt"

## EP 2 197 900 B9

<220>  
 <221> misc\_feature  
 <222> (57)..(57)  
 <223> r is a or g  
 5  
 <400> 52  
 gcnagcgacg aggcgcgcgcg ctgctacaac gaccccaagt ctgcgactt cgtcgar 57  
 <210> 53  
 10 <211> 57  
 <212> DNA  
 <213> Artificial  
 <220>  
 15 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betatI-AR  
 <220>  
 <221> misc\_feature  
 <222> (7)..(9)  
 20 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"  
 25  
 <220>  
 <221> misc\_feature  
 <222> (21)..(21)  
 <223> r is a or g  
 30  
 <220>  
 <221> misc\_feature  
 <222> (24)..(24)  
 <223> n is a, c, g, t or u  
 35  
 <400> 53  
 aagtgtcttg acttcgtcca rgcnagcgac gaggcgcgcc gctgtacaa cgacccc 57  
 <210> 54  
 40 <211> 57  
 <212> DNA  
 <213> Artificial  
 <220>  
 45 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betatI-AR  
 <220>  
 <221> misc\_feature  
 <222> (7)..(9)  
 50 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 /replace ="tcg"  
 /replace ="agt"  
 55  
 <220>  
 <221> misc\_feature  
 <222> (21)..(21)

<223> r is a or g

<220>  
 <221> misc\_feature  
 5 <222> (24)..(24)  
 <223> n is a, c, g, t or u

<400> 54  
 10 aagtgtctg acttcgtcga rgcnagcgac gaggcgcgcc gctgtacaa cgacccc 57

<210> 55  
 <211> 57  
 <212> DNA  
 <213> Artificial  
 15

<220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

<220>  
 20 <221> misc\_feature  
 <222> (4)..(6)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 25 /replace ="tcg"  
 /replace ="agt"

<220>  
 30 <221> misc\_feature  
 <222> (21)..(21)  
 <223> r is a or g

<220>  
 35 <221> misc\_feature  
 <222> (24)..(24)  
 <223> n is a, c, g, t or u

<400> 55  
 40 aagtcttgcg acttcgtcca rgcnagcgac gaggcgcgcc gctgtacaa cgacccc 57

<210> 56  
 <211> 57  
 <212> DNA  
 <213> Artificial  
 45

<220>  
 <223> Nucleotide sequence encoding an amino acid sequence homologous to an ECII epitope of human betal-AR

<220>  
 50 <221> misc\_feature  
 <222> (4)..(6)  
 <223> /replace ="tcc"  
 /replace ="agc"  
 /replace ="tca"  
 55 /replace ="tcg"  
 /replace ="agt"

<220>

<221> misc\_feature

<222> (21)..(21)

<223> r is a or g

5

<220>

<221> misc\_feature

<222> (24)..(24)

<223> n is a, c, g, t or u

10


<400> 56

aagtcttgcg acttcgtcga rgcnagcgac gaggcgcgcc gctgctacaa cgacccc 57

## Claims

15

1. A cyclic peptide of formula I:



20

wherein

a) x is an amino acid other than Cys;

b) h is any integer from 1 to 15;

c) i is any integer from 0 to 14;

25

d)  $X^c$  is Pro;

e) y is an amino acid other than Cys, and

wherein said cyclic peptide consists of at least 16 and of at most 25 amino acids and comprises the amino acid stretch Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 45); or

30

Glu-Ser-Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 46),

wherein Xxx<sub>1</sub>, Xxx<sub>3</sub> and Xxx<sub>4</sub> is an amino acid other than Cys, and

wherein said cyclic peptide

i) is capable of reducing an antibody-mediated activation of  $\beta_1$ -adrenergic receptor ( $\beta_1$ -AR);

35

ii) is capable of binding (auto)antibodies against the second extracellular loop (EC<sub>II</sub>) of  $\beta_1$ -AR; and/or

iii) is capable of inhibiting the interaction between  $\beta_1$ -AR and (auto)antibodies against EC<sub>II</sub> of  $\beta_1$ -AR.

2. The cyclic peptide of claim 1, wherein y is individually and independently selected from the group consisting of polar amino acids.



40

3. The cyclic peptide of any one of claims 1 or 2, wherein y is Ser or a Ser analogue.

4. The cyclic peptide of any one of claims 1 to 3, wherein h is 5, 8 or 9 and/or wherein i is 3, 4 or 6.

45

5. The cyclic peptide of any one of claims 1 to 4, being a cyclic peptide of formula I' or I'':





50

wherein  $x_i$  is Ala, Gly, Val, Thr or Ser and  $x_{III}$  is Arg.

6. The cyclic peptide of any one of claims 1 to 5, being a cyclic peptide of formula I''' or I''':

55






wherein  $x_{II}$  is Gln, Glu, Asp or Asn and  $x_{IV}$  is Gly or a Gly analogue.

7. The cyclic peptide of claim 5 or 6, wherein  $x_I$  is Ala.

5 8. The cyclic peptide of any one of claims 5 to 7, wherein  $x_{II}$  is Gln or Glu.

9. The cyclic peptide of any one of claims 1 to 8, wherein  $y$  is not Pro and/or  $x$  except  $x^c$  is not Pro.




10 10. The cyclic peptide of any one of claims 1 to 9 being a cyclic peptide of formula II, III or III':



wherein

- 20 a)  $x_1$  or  $Xxx_1$  is individually and independently selected from the group consisting of acidic amino acids; and/or  
b)  $x_2$  is individually and independently selected from the group consisting of basic amino acids.

11. The cyclic peptide of any one of claims 1 to 10 being a cyclic peptide of formula IV, V or V':



30 wherein



- a)  $x_1$  or  $Xxx_1$  is individually and independently selected from the group consisting of acidic amino acids;  
b)  $x_2$  is individually and independently selected from the group consisting of basic amino acids;  
c)  $x_3$  or  $Xxx_3$  is individually and independently selected from the group consisting of Leu, Ile, Val, Met, Trp, Tyr and Phe; and/or  
35 d)  $x_4$  or  $Xxx_4$  is individually and independently selected from the group consisting of Ser, Thr, Ala and Gly.

12. The cyclic peptide of any one of claims 1 to 11, comprising the amino acid stretch

40 Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys; or  
Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys.

13. The cyclic peptide of any one of claims 1 to 12 being a cyclic peptide selected from the group consisting of:

- a) a cyclic peptide formable or formed by the amino acid sequence as depicted in any one of SEQ ID NO. 41, 43, 1 to 4 and 17 to 20;  
b) a cyclic peptide formable or formed by an amino acid sequence as encoded, by a nucleotide sequence as depicted in any one of SEQ ID NO. 42, 44, 9 to 12, 25 to 28, 49, 50, 53 and 54;  
c) a cyclic peptide formable or formed by an amino acid sequence as encoded by a nucleotide sequence which differs from the nucleotide sequence as depicted in any one of SEQ ID NO. 42, 44, 9 to 12, 25 to 28, 49, 50, 53 and 54 due to the degeneracy of the genetic code; and  
50 d) a cyclic peptide of any one of formula VI to IX':



**cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-Gln) (VII);**

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-DGlu) (VIII);

**cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-DGlu) (IX).**

14. The cyclic peptide of claim 13, wherein at least one of acidic amino acids is replaced by a different amino acid selected from the group consisting of acidic amino acids.
15. The cyclic peptide of claim 13 or 14, wherein at least one of the basic amino acids is replaced by a different amino acid selected from the group consisting of basic amino acids.
16. The cyclic peptide of any one of claims 13 to 15, wherein at least one of the aliphatic amino acid residues is replaced by a different amino acid selected from the group consisting of aliphatic amino acids.
17. A method for producing a cyclic peptide of any one of claims 1 to 16, comprising the steps of
  - a)
    - (i) culturing a recombinant host cell comprising a nucleic acid molecule encoding the amino acid backbone of the cyclic peptide of any one of claims 1 to 16 or a vector comprising said nucleic acid molecule under conditions such that the amino acid backbone of the polypeptide of any one of claims 1 to 16 is expressed, and recovering said amino acid backbone; or
    - (ii) chemically synthesizing the amino acid backbone of the polypeptide of any one of claims 1 to 16; and
  - b) cyclization of said amino acid backbone to form the cyclic peptide of any one of claims 1 to 16.
18. The method of claim 17, wherein said nucleotide sequence is a nucleotide sequence as depicted in any one of SEQ ID NO. 9 to 12, 25 to 28, 42 and 44 or a nucleotide sequence which differs therefrom due to the degeneracy of the genetic code.
19. A composition comprising a cyclic peptide of any one of claims 1 to 16 and optionally a carrier.
20. The composition of claim 19, wherein said composition is a pharmaceutical composition and said carrier is a pharmaceutically acceptable carrier.
21. A cyclic peptide of any one of claims 1 to 16 or a pharmaceutical composition of claim 20, and optionally a pharmaceutically acceptable carrier, for use in
  - a) the treatment, amelioration or prevention of a disease where the activity of a  $\beta$ -AR is enhanced;
  - b) the treatment of a patient having antibodies against a  $\beta$ -AR; or
  - c) inducing immune tolerance.
22. The cyclic peptide or pharmaceutical composition of claim 21, wherein said disease where the activity of a  $\beta$ -adrenergic receptor is enhanced is a heart disease or wherein said patient suffers from a heart disease.
23. The cyclic peptide or pharmaceutical composition of claim 22, wherein said heart disease is selected from the group consisting of infectious and non-infectious heart disease, ischemic and non-ischemic heart disease, inflammatory




heart disease and myocarditis, cardiac dilatation, idiopathic cardio-myopathy, (idiopathic) dilated cardiomyopathy (DCM), immune-cardiomyopathy, heart failure, and any cardiac arrhythmia including ventricular and/or supraventricular premature capture beats as well as any atrial arrhythmia including atrial fibrillation and/or atrial flutter.

24. The cyclic peptide or the pharmaceutical composition of claim 22 or 23, wherein said heart disease is (idiopathic) DCM.
25. The cyclic peptide or the pharmaceutical composition of any one of claims 21 to 24, wherein said disease is induced by antibodies against a  $\beta$ -AR.
26. The cyclic peptide or the pharmaceutical composition of any one of claims 21 to 25, wherein said cyclic peptide or said pharmaceutical composition is administered so that at least 0,05 mg of said cyclic peptide per kg body weight is reached.
27. Method for detecting antibodies against a  $\beta$ -AR (in a sample) comprising the step of contacting the cyclic peptide of any one of claims 1 to 16 with said antibodies to be detected.
28. The cyclic peptide, the pharmaceutical composition or the method of any one of claims 21 to 27, wherein said  $\beta$ -AR is  $\beta_1$ -AR.

## Patentansprüche

1. Cyclisches Peptid mit der Formel I:



wobei

- a) x eine andere Aminosäure als Cys ist;
- b) h eine ganze Zahl von 1 bis 15 ist;
- c) i eine ganze Zahl von 0 bis 14 ist;
- d)  $x^c$  Pro ist;
- e) y eine andere Aminosäure als Cys ist; und



wobei das cyclische Peptid aus mindestens 16 und höchstens 25 Aminosäuren besteht umfassend den Aminosäurebereich

Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 45); oder  
Glu-Ser-Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 46),  
wobei Xxx<sub>1</sub>, Xxx<sub>3</sub> und Xxx<sub>4</sub> eine andere Aminosäure als Cys ist, und

wobei das cyclische Peptid


- i) in der Lage ist, eine Antikörper-vermittelte Aktivierung von  $\beta_1$ -adrenergem Rezeptor ( $\beta_1$ -AR) zu reduzieren;
- ii) in der Lage ist, (Auto)-Antikörper gegen die zweite extrazelluläre Schleife (EC<sub>II</sub>) von  $\beta_1$ -AR zu binden; und/oder
- iii) in der Lage ist, die Interaktion zwischen  $\beta_1$ -AR und (Auto)-Antikörpern gegen EC<sub>II</sub> von  $\beta_1$ -AR zu inhibieren.

2. Cyclisches Peptid nach Anspruch 1, wobei y individuell und unabhängig aus der Gruppe bestehend aus polaren Aminosäuren ausgewählt ist.
3. Cyclisches Peptid nach einem der Ansprüche 1 oder 2, wobei y Ser oder ein Ser-Analog ist.
4. Cyclisches Peptid nach einem der Ansprüche 1 bis 3, wobei h 5, 8 oder 9 ist und/oder wobei i 3, 4 oder 6 ist.
5. Cyclisches Peptid nach einem der Ansprüche 1 bis 4 mit der Formel I' oder I'':



wobei  $x_I$  Ala, Gly, Val, Thr oder Ser ist und  $x_{III}$  Arg ist.

6. Cyclisches Peptid nach einem der Ansprüche 1 bis 5 mit der Formel I''' oder I''':




wobei  $x_{II}$  Gln, Glu, Asp, oder Asn ist und  $x_{IV}$  Gly oder ein Gly-Analog ist.

7. Cyclisches Peptid nach Anspruch 5 oder 6, wobei  $x_I$  Ala ist.

8. Cyclisches Peptid nach einem der Ansprüche 5 bis 7, wobei  $x_{II}$  Gln oder Glu ist.

9. Cyclisches Peptid nach einem der Ansprüche 1 bis 8, wobei  $y$  nicht Pro ist und/oder  $x$  ausser  $x^c$  nicht Pro ist.


10. Cyclisches Peptid nach einem der Ansprüche 1 bis 9 mit der Formel II, III oder III':



wobei

- a)  $x_1$  oder  $Xxx_1$  individuell und unabhängig aus der Gruppe bestehend aus sauren Aminosäuren ausgewählt ist; und/oder
- b)  $x_2$  individuell und unabhängig aus der Gruppe bestehend aus basischen Aminosäuren ausgewählt ist.

11. Cyclisches Peptid nach einem der Ansprüche 1 bis 10 mit der Formel IV, V oder V':



wobei

- a)  $x_1$  oder  $Xxx_1$  individuell und unabhängig aus der Gruppe bestehend aus sauren Aminosäuren ausgewählt ist; und/oder
- b)  $x_2$  individuell und unabhängig aus der Gruppe bestehend aus basischen Aminosäuren ausgewählt ist;
- c)  $x_3$  oder  $Xxx_3$  individuell und unabhängig aus der Gruppe bestehend aus Leu, Ile, Val, Met, Trp, Tyr und Phe ausgewählt ist; und/oder
- d)  $x_4$  oder  $Xxx_4$  individuell und unabhängig aus der Gruppe bestehend aus Ser, Thr, Ala und Gly ausgewählt ist.

12. Cyclisches Peptid nach einem der Ansprüche 1 bis 11, umfassend den Aminosäurebereich

Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys; oder  
Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys.

13. Cyclisches Peptid nach einem der Ansprüche 1 bis 12 ausgewählt aus der Gruppe bestehend aus:

- a) einem cyclischen Peptid formbar oder gebildet aus der Aminosäuresequenz wie in einer der SEQ ID NO. 41, 43, 1 bis 4 und 17 bis 20 dargestellt;
- b) einem cyclischen Peptid formbar oder gebildet aus der Aminosäuresequenz, die durch die Nucleotidsequenz wie in einer der SEQ ID NO. 42, 44, 9 bis 12, 25 bis 28, 49, 50, 53 und 54 dargestellt codiert wird;
- c) einem cyclischen Peptid formbar oder gebildet aus einer Aminosäuresequenz, die durch eine Nucleotidse-

quenz codiert wird, die sich von der Nucleotidsequenz wie in einer der SEQ ID NO. 42, 44, 9 bis 12, 25 bis 28, 49, 50, 53 und 54 dargestellt aufgrund der Degeneration des genetischen Codes unterscheidet; und  
d) einem cyclischen Peptid nach einer der Formeln VI bis IX':

cyclo(Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Gly) (IX')

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Gln) (VI);

cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-Gln) (VII);

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-DGlu) (VIII);  
cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-DGlu) (IX).

14. Cyclisches Peptid nach Anspruch 13, wobei mindestens eine der sauren Aminosäuren durch eine andere Aminosäure ausgewählt aus der Gruppe bestehend aus sauren Aminosäuren ersetzt ist.

15. Cyclisches Peptid nach Anspruch 13 oder 14, wobei mindestens eine der basischen Aminosäuren durch eine andere Aminosäure ausgewählt aus der Gruppe bestehend aus basischen Aminosäuren ersetzt ist.

16. Cyclisches Peptid nach einem der Ansprüche 13 bis 15, wobei mindestens eine der aliphatischen Aminosäuren durch eine andere Aminosäure ausgewählt aus der Gruppe bestehend aus aliphatischen Aminosäuren ersetzt ist.

17. Verfahren zur Produktion eines cyclischen Peptids nach einem der Ansprüche 1 bis 16, umfassend die Schritte:

a)

- (i) Züchten einer rekombinanten Wirtszelle umfassend ein Nucleinsäuremolekül, das das Aminosäurerückgrat des cyclischen Peptids nach einem der Ansprüche 1 bis 16 codiert, oder einen Vektor, der das Nucleinsäuremolekül umfasst, unter solchen Bedingungen, dass das Aminosäurerückgrat des Polypeptids nach einem der Ansprüche 1 bis 16 exprimiert wird, und Gewinnung des Aminosäurerückgrats; oder
- (ii) chemische Synthese des Aminosäurerückgrats des Polypeptids nach einem der Ansprüche 1 bis 16; und

b) Cyclisieren des Aminosäurerückgrats, um das cyclische Peptid nach einem der Ansprüche 1 bis 16 zu bilden.

18. Verfahren nach Anspruch 17, wobei die Nucleotidsequenz eine Nucleotidsequenz wie in einer der SEQ ID NO. 9 bis 12, 25 bis 28, 42 und 44 dargestellt ist oder eine Nucleotidsequenz, die sich davon aufgrund der Degeneration des genetischen Codes unterscheidet.

19. Zusammensetzung umfassend ein cyclisches Peptid nach einem der Ansprüche 1 bis 16 und gegebenenfalls einen Träger.

20. Zusammensetzung nach Anspruch 19, wobei die Zusammensetzung ein Arzneimittel ist und der Träger ein pharmazeutisch verträglicher Träger ist.

21. Cyclisches Peptid nach einem der Ansprüche 1 bis 16 oder Arzneimittel nach Anspruch 20 und gegebenenfalls ein pharmazeutisch verträglicher Träger zur Verwendung in

a) der Behandlung, Verbesserung oder Vorbeugung einer Krankheit, bei der die Aktivität eines  $\beta$ -AR gesteigert

ist;

b) der Behandlung eines Patienten, der Antikörper gegen einen  $\beta$ -AR hat; oder

c) der Induzierung von Immuntoleranz.

22. Cyclisches Peptid oder Arzneimittel nach Anspruch 21, wobei die Krankheit, in der die Aktivität eines  $\beta$ -adrenergen Rezeptors gesteigert ist, eine Herzerkrankung ist oder wobei der Patient an einer Herzerkrankung leidet.

23. Cyclisches Peptid oder Arzneimittel nach Anspruch 22, wobei die Herzerkrankung ausgewählt ist aus der Gruppe bestehend aus infektiöser und nicht-infektiöser Herzerkrankung, ischämischer und nicht-ischämischer Herzerkrankung, entzündlicher Herzerkrankung und Myokarditis, Herzdilatation, idiopathischer Myokardiopathie, (idiopathischer) dilatativer Myokardiopathie (DCM), Immun-Myokardiopathie, Herzversagen und jeder Herzarrhythmie einschließlich ventrikulärer und/oder supraventrikulärer verfrühter Capture Beats sowie jeder atrialer Arrhythmie einschließlich Vorhofflimmern und/oder Vorhofflattern.

24. Cyclisches Peptid oder Arzneimittel nach Anspruch 22 oder 23, wobei die Herzerkrankung (idiopathische) DCM ist.

25. Cyclisches Peptid oder Arzneimittel nach einem der Ansprüche 21 bis 24, wobei die Krankheit durch Antikörper gegen einen  $\beta$ -AR ausgelöst wird.


26. Cyclisches Peptid oder Arzneimittel nach einem der Ansprüche 21 bis 25, wobei das cyclische Peptid oder das Arzneimittel so verabreicht wird, dass mindestens 0,05 mg des cyclischen Peptids pro kg Körpergewicht erreicht wird.

27. Verfahren zum Nachweis von Antikörpern gegen einen  $\beta$ -AR (in einer Probe), umfassend den Schritt des Inkontaktbringens des cyclischen Peptids nach einem der Ansprüche 1 bis 16 mit den Antikörpern, die nachgewiesen werden sollen.

28. Cyclisches Peptid, Arzneimittel oder Verfahren nach einem der Ansprüche 1 bis 27, wobei der  $\beta$ -AR  $\beta_1$ -AR ist.

## Revendications

1. Peptide cyclique selon la formule I :



dans lequel :

a) x est un acide aminé différent de Cys ;

b) h est un quelconque nombre entier compris entre 1 et 15 ;

c) i est un quelconque nombre entier compris entre 0 et 14 ;

d)  $x^c$  est Pro ;

e) y est un acide aminé différent de Cys ; et

dans lequel ledit peptide cyclique consiste en au moins 16 et au plus 25 acides aminés et comprend l'étirement d'acide aminé

Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 45) ; ou Glu-Ser-Asp-Xxx<sub>1</sub>-Xxx<sub>4</sub>-Arg-Arg-Cys-Xxx<sub>3</sub>-Asn-Asp-Pro-Lys (SEQ ID NO. 46),

où Xxx<sub>1</sub>, Xxx<sub>3</sub> et Xxx<sub>4</sub> sont un acide aminé différent de Cys ; et

dans lequel ledit peptide cyclique :

i) est capable de réduire une activation à médiation anticorps du récepteur  $\beta_1$ -adrénergique ( $\beta_1$ -AR) ;

ii) est capable de lier des (auto)anticorps contre la deuxième boucle extracellulaire (EC<sub>II</sub>) de  $\beta_1$ -AR ; et/ou



iii) est capable d'inhiber l'interaction entre  $\beta_1$ -AR et les (auto)anticorps contre (EC<sub>II</sub>) de  $\beta_1$ -AR.

2. Peptide cyclique selon la revendication 1, dans lequel y est sélectionné individuellement et indépendamment dans le groupe consistant en des acides aminés polaires.

3. Peptide cyclique selon l'une quelconque des revendications 1 ou 2, dans lequel y est Ser ou un analogue de Ser.



4. Peptide cyclique selon l'une quelconque des revendications 1 à 3, dans lequel h est égal à 5, 8 ou 9 et/ou dans lequel i est égal à 3, 4 ou 6.

5. Peptide cyclique selon l'une quelconque des revendications 1 à 4, qui est un peptide cyclique selon la formule I' ou I'' :



où  $x_I$  est Ala, Gly, Val, Thr ou Ser, et  $x_{III}$  est Arg.

6. Peptide cyclique selon l'une quelconque des revendications 1 à 5, qui est un peptide cyclique selon la formule I''' ou I'''' :






où  $x_{II}$  est Gln, Glu, Asp ou Asn, et  $x_{IV}$  est Gly ou un analogue de Gly.

7. Peptide cyclique selon la revendication 5 ou 6, dans lequel  $x_I$  est Ala.

8. Peptide cyclique selon l'une quelconque des revendications 5 à 7, dans lequel  $x_{II}$  est Gln ou Glu.

9. Peptide cyclique selon l'une quelconque des revendications 1 à 8, dans lequel y n'est pas Pro et/ou x, à l'exception de  $x^c$ , n'est pas Pro.

10. Peptide cyclique selon l'une quelconque des revendications 1 à 9, qui est un peptide cyclique selon la formule II, III ou III' :






où :

a)  $x_1$  ou  $Xxx_1$  est sélectionné individuellement et indépendamment dans le groupe consistant en des acides aminés acides ; et/ou

b)  $x_2$  est sélectionné individuellement et indépendamment dans le groupe consistant en des acides aminés basiques.

11. Peptide cyclique selon l'une quelconque des revendications 1 à 10, qui est un peptide cyclique selon la formule IV, V ou V' :



où :

- a)  $x_1$  ou  $Xxx_1$  est sélectionné individuellement et indépendamment dans le groupe consistant en des acides aminés acides ;  
 b)  $x_2$  est sélectionné individuellement et indépendamment dans le groupe consistant en des acides aminés basiques ;  
 5 c)  $x_3$  ou  $Xxx_3$  est sélectionné individuellement et indépendamment dans le groupe consistant en Leu, Ile, Val, Met, Trp, Tyr et Phe ; et/ou  
 d)  $x_4$  ou  $Xxx_4$  est sélectionné individuellement et indépendamment dans le groupe consistant en Ser, Thr, Ala et Gly.

10 **12.** Peptide cyclique selon l'une quelconque des revendications 1 à 11, comprenant l'étirement d'acide aminé Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys ; ou  
 Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys.

15 **13.** Peptide cyclique selon l'une quelconque des revendications 1 à 12, qui est un peptide cyclique sélectionné dans le groupe consistant en :

- a) un peptide cyclique formé ou pouvant être formé par la séquence d'acide aminé telle que décrite dans l'une quelconque des SEQ ID NO. 41, 43, 1 à 4 et 17 à 20 ;  
 b) un peptide cyclique formé ou pouvant être formé par une séquence d'acide aminé telle que codée par une  
 20 séquence de nucléotides telle que décrite dans l'une quelconque des SEQ ID NO. 42, 44, 9 à 12, 25 à 28, 49, 50, 53 et 54 ;  
 c) un peptide cyclique formé ou pouvant être formé par une séquence d'acide aminé telle que codée par une séquence de nucléotides qui diffère de la séquence de nucléotides telle que décrite dans l'une quelconque des  
 25 SEQ ID NO. 42, 44, 9 à 12, 25 à 28, 49, 50, 53 et 54 en raison de la dégénérescence du code génétique ; et  
 d) un peptide cyclique selon l'une quelconque des formules VI à IX' :

cyclo(Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-  
 30 Cys-Ser-Asp-Phe-Val-Thr-Gly) (IX') ;

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-  
 35 Phe-Val-Gln) (VI) ;

cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-  
 40 Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-Gln) (VII) ;

cyclo(Ala-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-Lys-Cys-Ser-Asp-  
 45 Phe-Val-DGlu) (VIII) ;

cyclo(Ala-Arg-Ala-Glu-Ser-Asp-Glu-Ala-Arg-Arg-Cys-Tyr-Asn-Asp-Pro-  
 50 Lys-Cys-Ser-Asp-Phe-Val-Thr-Asn-Arg-DGlu) (IX).

**14.** Peptide cyclique selon la revendication 13, dans lequel au moins l'un des acides aminés acides est remplacé par un acide aminé différent sélectionné dans le groupe consistant en des acides aminés acides.

55 **15.** Peptide cyclique selon la revendication 13 ou 14, dans lequel au moins l'un des acides aminés basiques est remplacé par un acide aminé différent sélectionné dans le groupe consistant en des acides aminés basiques.

**16.** Peptide cyclique selon l'une quelconque des revendications 13 à 15, dans lequel au moins l'un des résidus d'acides

aminés aliphatiques est remplacé par un acide aminé différent sélectionné dans le groupe constitué par des acides aminés aliphatiques.

- 5 17. Procédé destiné à produire un peptide cyclique selon l'une quelconque des revendications 1 à 16, comprenant les étapes consistant à :

a)

10 (i) cultiver une cellule hôte recombinante comprenant une molécule d'acide nucléique codant la chaîne principale d'acide aminé du peptide cyclique selon l'une quelconque des revendications 1 à 16, ou un vecteur comprenant ladite molécule d'acide nucléique dans des conditions telles que la chaîne principale d'acide aminé du polypeptide selon l'une quelconque des revendications 1 à 16 est exprimée, et récupérer ladite chaîne principale d'acide aminé ; ou

15 (ii) synthétiser chimiquement la chaîne principale d'acide aminé du polypeptide selon l'une quelconque des revendications 1 à 16 ; et

b) effectuer une cyclisation de ladite chaîne principale d'acide aminé afin de former le peptide cyclique selon l'une quelconque des revendications 1 à 16.

- 20 18. Procédé selon la revendication 17, dans lequel ladite séquence de nucléotides est une séquence de nucléotides telle que décrite dans l'une quelconque des SEQ ID NO. 9 à 12, 25 à 28, 42 et 44, ou une séquence de nucléotides qui diffère de celle-ci en raison de la dégénérescence du code génétique.

- 25 19. Composition comprenant un peptide cyclique selon l'une quelconque des revendications 1 à 16 et éventuellement un support.

- 20 20. Composition selon la revendication 19, dans laquelle ladite composition est une composition pharmaceutique et ledit support est un support pharmaceutiquement acceptable.

- 30 21. Peptide cyclique selon l'une quelconque des revendications 1 à 16 ou composition pharmaceutique selon la revendication 20, et éventuellement support pharmaceutiquement acceptable, destiné à une utilisation dans :

a) le traitement, l'amélioration ou la prévention d'une maladie où l'activité d'un  $\beta$ -AR est accrue ;

b) le traitement d'un patient présentant des anticorps contre un  $\beta$ -AR ; ou

35 c) l'induction d'une tolérance immunitaire.

22. Peptide cyclique ou composition pharmaceutique selon la revendication 21, dans lequel ladite maladie où l'activité d'un récepteur  $\beta$ -adrénergique est accrue est une cardiopathie ou dans lequel ledit patient souffre d'une cardiopathie.

- 40 23. Peptide cyclique ou composition pharmaceutique selon la revendication 22, dans lequel ladite cardiopathie est sélectionnée dans le groupe consistant en une cardiopathie infectieuse et non infectieuse, une cardiopathie ischémique et non ischémique, une cardiopathie inflammatoire et une myocardite, une dilatation cardiaque, une cardiomyopathie idiopathique, une cardiomyopathie dilatée (CMD) (idiopathique), une cardiomyopathie immune, une insuffisance cardiaque, et toute arythmie cardiaque y compris les battements de capture ventriculaire et/ou supra-ventriculaire prématurée, ainsi que toute arythmie auriculaire y compris la fibrillation auriculaire et/ou le flutter auriculaire.

- 50 24. Peptide cyclique ou composition pharmaceutique selon la revendication 22 ou 23, dans lequel ladite cardiopathie est une CMD (idiopathique).

25. Peptide cyclique ou composition pharmaceutique selon l'une quelconque des revendications 21 à 24, dans lequel ladite maladie est induite par des anticorps contre un  $\beta$ -AR.

- 55 26. Peptide cyclique ou composition pharmaceutique selon l'une quelconque des revendications 21 à 25, dans lequel ledit peptide cyclique ou ladite composition pharmaceutique est administré(e) de telle sorte que ledit peptide cyclique atteigne au moins 0,05 mg par kilogramme de poids corporel.

27. Procédé destiné à détecter des anticorps contre un  $\beta$ -AR (dans un échantillon) comprenant l'étape où le peptide

cyclique selon l'une quelconque des revendications 1 à 16 est mis en contact avec lesdits anticorps à détecter.

- 28.** Peptide cyclique, composition pharmaceutique ou procédé selon l'une quelconque des revendications 21 à 27, dans lequel ledit  $\beta$ -AR est un  $\beta_1$ -AR.

5

10

15

20

25

30

35

40


45

50

55



Figure 1.






Figure 2.

Figure 3.






Figure 4.








Figure 6.

Figure 7.




Figure 8.






Figure 9.





Figure 10.







Figure 12.

Course of ELISA immunoreactivity, % of untreated immunized rats






Figure 13.

Figure 14.




Figure 15.




Figure 16.A






Figure 16.B




Figure 16.C




Figure 17. A





Figure 17. B






Figure 18.




Figure 19

B




Figure 20 A






Figure 20B




Figure 20C




Figure 21A.





Figure 21B.







Figure 22A.

Figure 22B.







Figure 23.

Figure 24.

$\beta_1$ -EC<sub>II</sub> sequence



Cyclopeptide Cys/Cys 22 AA



Figure 25A.




Figure 25B.




Figure 26A.




Figure 26B.





Figure 27.

A



B





Figure 28.

A



B




Figure 29A




Figure 29B







Figure 30



## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- WO 0121660 A [0012]
- WO 2006103101 A [0013] [0028] [0036] [0146] [0147]
- US 5525711 A [0130]
- US 4711955 A [0130]
- US 5792608 A [0130]
- EP 302175 A [0130]
- US 4880635 A, Janoff [0168]
- WO 2005052186 A [0206] [0207]

## Non-patent literature cited in the description

- **RICHARDSON.** *Circulation*, 1996, vol. 93, 841-842 [0002]
- **WARNE.** *Nature*, 2008, 1038 [0005]
- **FRIELLE et al.** *PNAS*, 1987, vol. 84, 7920-7924 [0024] [0215]
- **FRIELLE.** *PNAS*, 1987, vol. 84, 7920-7924 [0059]
- **KATES ; ALBERICIO.** Solid phase synthesis. CRC-Press, 2000 [0068]
- **WILLIAMS.** Chemical Approaches to the Synthesis of Peptides. CRC-Press, 1997 [0068] [0070] [0071] [0151]
- **BENOITON.** Chemistry of Peptide Synthesis. CRC-Press, 2005 [0068] [0070] [0071] [0151]
- **STRYER.** Biochemie. Spectrum Akad. Verlag, 1991 [0078]
- **TIJSSEN.** Practice and theory of enzyme immunoassays. 1985, vol. 15 [0122]
- **DAVIS LG ; DIBMER MD.** Basic methods in molecular biology. Battey Elsevier, 1990 [0122]
- Immunochemical methods in cell and molecular biology. Academic Press, 1987 [0122]
- Methods in Enzymology. Academic Press, Inc, [0122]
- **COLE-STRAUSS.** *Science*, 1996, vol. 273 (5280), 1386-9 [0130]
- **SAMBROOK.** Molecular Cloning A Laboratory Manual. Cold Spring Harbor Laboratory, 1989 [0134]
- **AUSUBEL.** Current Protocols in Molecular Biology. Green Publishing Associates and Wiley Interscience, 1994 [0134] [0140]
- **CHEN W.C. ; WHITE P.D.** *Fmoc Solid Phase Peptide Synthesis*, 2003 [0147]
- **KATES S. ; ALBERICIO F.** *Solid phase synthesis*, 2000 [0147]
- **BENOITON N.L.** Chemistry of Petide Synthesis. CRC-Press, 2005 [0148]
- **HERNPOL CM ; VINCENT P ; ADAMS SR ; TSIEN RY ; SELVERSTON AL.** *Nature*, 1996, vol. 384, 113-114 [0205]
- **ZACCOLO M ; POZZAN T.** *Science*, 2002, vol. 295, 1711-1715 [0205]
- Heart disease and stroke statistics - 2007 update. *Circulation*, 2007, vol. 115, e69-e171 [0273]
- **ANDERTON.** *Immunology*, 2001, vol. 104, 367-376 [0273]
- **BABA.** *Eur. Heart J.*, 2004, vol. 25, 1108-1115 [0273]
- **BOIVIN.** *Eur. J. Heart Fail.*, 2005, vol. 4 (1), 24 [0273]
- **CAFORIO.** *Eur. J. Heart Fail.*, 2002, vol. 4, 411-417 [0273]
- **CHIALE.** *Circulation*, 2001, vol. 103, 1765-1771 [0273]
- **CHIEN.** *Oncol.*, 2000, vol. 27, 9-17 [0273]
- **CHRIST.** *J. Mol. Cell. Cardiol.*, 2001, vol. 33, 1515-1525 [0273]
- **CHRIST.** *J. Mol. Cell Cardiol.*, 2006, vol. 41, 716-723 [0273]
- **ELIES.** *J. Immunol.*, 1996, vol. 157, 4203-4211 [0273]
- **ENGELHARDT.** *Proc. Natl. Acad. Sci. USA*, 1999, vol. 96, 7059-7064 [0273]
- **ERIKSSON.** *Nat. Med.*, 2003, vol. 9, 1484-1490 [0273]
- **FABRIZIO.** *Drugs Ther.*, 1994, vol. 8, 89-94 [0273]
- **FELIX.** *J. Am. Coll. Cardiol.*, 2000, vol. 35, 1590-1598 [0273]
- **FERRARI.** *J. Exp. Med.*, 1995, vol. 182, 59-65 [0273]
- **FREEDMAN.** *J. Clin. Invest.*, 2004, vol. 113, 1379-1382 [0273]
- **FU.** *J. Clin. Invest.*, 1993, vol. 91, 1964-1968 [0273]
- **GÖSER.** *Circulation*, 2006, vol. 114, 1693-1702 [0273]
- **HERSHKO.** *Ann. N.Y. Acad. Sci.*, 2005, vol. 1051, 635-646 [0273]
- **HOEBEKE.** *Int. J. Cardiol.*, 1996, vol. 54, 103-111 [0273]
- **IWATA.** *J. Am. Coll. Cardiol.*, 2001, vol. 37, 418-424 [0273]
- **IWATA.** *Circ. Res.*, 2001, vol. 88, 578-586 [0273]

- **JAHNS.** *Eur. J. Pharmacol.*, 1996, vol. 316, 111-121 [0273]
- **JAHNS.** *J. Am. Coll. Cardiol.*, 1999, vol. 34, 1545-1551 [0273]
- **JAHNS.** *Circulation*, 1999, vol. 99, 649-654 [0273]
- **JAHNS.** *J. Am. Coll. Cardiol.*, 2000, vol. 36, 1280-1287 [0273]
- **JAHNS.** *J. Clin. Invest.*, 2004, vol. 113, 1419-1429 [0273]
- **JAHNS.** *Trends Cardiovasc Med*, 2006, vol. 16, 20-24 [0273]
- **KHOYNEZHAD.** *Eur. J. Heart Fail.*, 2007, vol. 9, 120-123 [0273]
- **KÜHL.** *Circulation*, 2005, vol. 112, 1965-1970 [0273]
- **LIMAS.** *Am. Heart J.*, 1992, vol. 123, 967-970 [0273]
- **LIMAS.** *Int. J. Cardiol.*, 1996, vol. 54, 113-116 [0273]
- **LIMAS.** *Circulation*, 1997, vol. 95, 1979-1980 [0273]
- **LIMAS.** *Am. J. Cardiol.*, 2004, vol. 93, 1189-1191 [0273]
- **LOHSE.** *Circ. Res.*, 2003, vol. 93, 896-906 [0273]
- **LUPPI.** *Circulation*, 1998, vol. 98, 777-785 [0273]
- **MACLELLAN.** *Nat. Med.*, 2003, vol. 9, 1455-1456 [0273]
- **MAEKAWA.** *Circulation*, 2007, vol. 115, 5-8 [0273]
- **MAGNUSSON.** *Circulation*, 1994, vol. 89, 2760-2767 [0273]
- **MAGNUSSON.** *Int. J. Cardiol.*, 1996, vol. 54, 137-141 [0273]
- **MAHRHOLDT.** *Circulation*, 2006, vol. 114, 1581-1590 [0273]
- **MATSUI.** *Autoimmunity*, 1995, vol. 21, 85-88 [0273]
- **MATSUI.** *Autoimmunity*, 2001, vol. 43, 217-220 [0273]
- **MOBINI.** *Autoimmunity Rev.*, 2004, vol. 3, 277-284 [0273]
- **MORITA.** *J. Clin. Invest.*, 2005, vol. 115, 518-526 [0273]
- **NEUMANN.** *J. Am. Coll. Cardiol.*, 1990, vol. 16, 839-846 [0273]
- **NIKOLAEV.** *J. Am. Coll. Cardiol.*, 2007, vol. 50, 423-431 [0273]
- **OKAZAKI.** *Nat. Med.*, 2003, vol. 9, 1477-1483 [0273]
- **OKAZAKI.** *Trends Mol Med*, 2005, vol. 11, 322-326 [0273]
- **POHLNER.** *Am. J. Cardiol.*, 1997, vol. 80, 1040-1045 [0273]
- **ROSE.** *Immunol. Today*, 1993, vol. 14, 426-430 [0273]
- **ROSE.** *J. Clin. Invest.*, 2001, vol. 107, 943-944 [0273]
- **SCHULTHEISS.** *J. Mol. Cell Cardiol.*, 1985, vol. 17, 603-617 [0273]
- **SCHULTHEISS.** *J. Exp. Med.*, 1988, vol. 168, 2102-2109 [0273]
- **SCHULZE.** *Cardiovasc. Res.*, 1999, vol. 44, 91-100 [0273]
- **SEWALD.** *Peptides: Chemistry and Biology.* Wiley-VCH, 2002 [0273]
- **SMULSKI.** *FASEB J.*, 2006, vol. 20, 1396-1406 [0273]
- **STÖRK.** *Am. Heart J.*, 2006, vol. 152, 697-704 [0273]
- **WALLUKAT.** *J. Mol. Cell Cardiol.*, 1995, vol. 27, 397-406 [0273]
- **WALLUKAT.** *N. Engl. J. Med.*, 2002, vol. 347, 1806 [0273]
- **WANG.** *Blood Pressure*, 1996, vol. 3, 25-27 [0273]
- **WITEBSKY.** *J. Am. Med. Assoc.*, 1957, vol. 164, 1439-1447 [0273]
- **WOODIWISS.** *Circulation*, 2001, vol. 103, 155-160 [0273]