

(19)



(11)

**EP 2 258 397 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:

**11.10.2017 Bulletin 2017/41**

(21) Application number: **09722008.1**

(22) Date of filing: **17.03.2009**

(51) Int Cl.:

<b>A61K 47/60</b> <sup>(2017.01)</sup>	<b>A61K 47/64</b> <sup>(2017.01)</sup>
<b>A61K 31/352</b> <sup>(2006.01)</sup>	<b>A61K 31/405</b> <sup>(2006.01)</sup>
<b>A61K 31/407</b> <sup>(2006.01)</sup>	<b>A61K 31/704</b> <sup>(2006.01)</sup>
<b>A61K 38/00</b> <sup>(2006.01)</sup>	<b>A61P 29/00</b> <sup>(2006.01)</sup>
<b>A61P 35/00</b> <sup>(2006.01)</sup>	<b>C08G 69/44</b> <sup>(2006.01)</sup>
<b>C08G 69/48</b> <sup>(2006.01)</sup>	<b>C08G 69/10</b> <sup>(2006.01)</sup>
<b>C08G 69/40</b> <sup>(2006.01)</sup>	

(86) International application number:

**PCT/JP2009/055115**

(87) International publication number:

**WO 2009/116509 (24.09.2009 Gazette 2009/39)**

**(54) POLYMER CONJUGATE OF PHYSIOLOGICALLY ACTIVE SUBSTANCE**

POLYMERKONJUGAT AUS EINEM PHYSIOLOGISCH AKTIVEN STOFF

CONJUGUÉ DE POLYMÈRE ET DE SUBSTANCE PHYSIOLOGIQUEMENT ACTIVE

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR**

(30) Priority: **18.03.2008 JP 2008069067**

(43) Date of publication of application:

**08.12.2010 Bulletin 2010/49**

(73) Proprietor: **Nippon Kayaku Kabushiki Kaisha Tokyo 100-0005 (JP)**

(72) Inventors:

- **KITAGAWA, Masayuki Tokyo 115-8588 (JP)**
- **SENO, Chieko Tokyo 115-8588 (JP)**

(74) Representative: **Wablat Lange Karthaus**

**Anwaltssozietät  
Potsdamer Chaussee 48  
14129 Berlin (DE)**

(56) References cited:

<b>EP-A1- 1 580 216</b>	<b>EP-A1- 2 042 195</b>
<b>EP-B1- 1 401 473</b>	<b>WO-A1-99/30727</b>
<b>WO-A1-2005/066214</b>	<b>WO-A1-2007/080898</b>
<b>WO-A1-2008/010463</b>	<b>WO-A2-02/065988</b>
<b>JP-A- 61 243 026</b>	<b>JP-T- 2000 516 948</b>
<b>JP-T- 2005 508 832</b>	

- **GREENWALD R B ET AL: "Effective drug delivery by PEGylated drug conjugates", ADVANCED DRUG DELIVERY REVIEWS, ELSEVIER BV, AMSTERDAM, NL, vol. 55, no. 2, 10 February 2003 (2003-02-10), pages 217-250, XP002275066, ISSN: 0169-409X, DOI: 10.1016/S0169-409X(02)00180-1**
- **NAKANISHI, T. ET AL.: 'Development of the polymer micelle carrier system for doxorubicin' J. CONTROL. RELEASE vol. 74, no. 1-3, 2001, pages 295 - 302, XP004297538**

**EP 2 258 397 B1**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**Description**

## Technical Field

5 **[0001]** The present invention relates to a high-molecular weight conjugate of physiologically active substances in which a residue of the physiologically active substance having a carboxy group is bonded, through a specific linker, to a carboxy group of a block copolymer having a polyethylene glycol moiety and a polymer moiety having two or more carboxylic acid groups and the use thereof. The description also discloses a method for manufacturing the same, as well as conjugates wherein the residue of the physiologically active substance has an amino group.

## Background Art

15 **[0002]** High-molecular weight conjugates obtained by bonding physiologically active substances such as an anticancer agent or an anti-inflammatory agent, especially a physiologically active substances with low solubility in water, to a high-molecular weight carrier have been studied extensively since the high-molecular weight conjugates improve the in vivo pharmacokinetics of the physiologically active substances themselves, the water solubility thereof and the like, and are expected to thus improve the efficacy of the substances as drugs. Particularly block copolymers in which a hydrophilic polymer and a hydrophobic polymer are bonded are characterized by being capable of maintaining the water solubility as the whole polymer even if the amount of a physiologically active substances carried thereon is increased, by forming a micelle having the hydrophobic polymer carrying the physiologically active substances as the inner shell and the hydrophilic polymer covering the surface.

20 **[0003]** Patent Document 1 discloses a compound in which a drug is bonded to a block copolymer of a polyethylene glycol and a polyaspartic acid and which forms micelles and has water solubility. Patent Document 2 discloses a high-molecular weight derivative of camptothecins in which a side chain carboxy group of a block copolymer of a polyethylene glycol and a polyglutamic acid is bonded to a phenolic hydroxyl group of the camptothecins.

25 **[0004]** Patent Document 3 discloses a compound in which a drug is bonded to a polyethylene glycol through a specific linker to cause a benzyl elimination reaction. Patent Document 4 discloses a compound in which a drug is bonded to a branched amino acid such as aspartic acid bonded to a polyethylene glycol, through a specific linker to cause a benzyl elimination reaction.

30 Patent Document 1: Japanese Patent No. 2694923

Patent Document 2: WO 2004/39869

Patent Document 3: Japanese Patent Application Laid-Open

Publication (Kohyo) No. 2002-508400

35 Patent Document 4: Japanese Patent Application Laid-Open

Publication (Kohyo) No. 2004-532289

40 **[0005]** EP 1 580 216 A1 teaches a high-molecular weight derivative of Camptothecin having a structure wherein a carboxylic acid group of a copolymer of polyethylene glycol and a polymer having a carboxylic acid group at the side chain, is combined with a phenolic hydroxyl group of Camptothecin via an ester bond. The phenolic hydroxyl group of Camptothecin is directly ester-bonded to a carboxylic acid group of a block copolymer without a linker in EP 1 580 216 A1. EP 1 580 216 A1 does not disclose the feature that the carboxy group of the physiologically active substance is bound to the carboxylic acid group of the block copolymer via a specific linker.

45 **[0006]** WO 02/065988 A2 discloses terminally-branched polymeric prodrug platform and more specifically a polymeric conjugate in which an amino group of a physiologically active substance is bonded to a polymer having a polyethylene glycol moiety and a single aspartic acid moiety through a specific linker.

50 **[0007]** Greenwald R. B. et al describe in "Effective drug delivery by PEGylated drug conjugates" (Advanced Drug Delivery Reviews, Elsevier BV, Amsterdam, NL, Vol. 55, No. 2, 2003, pages 217 - 250) a polymeric derivative in which a physiologically active substance having a hydroxy group or an amino group is bound to a polymer via a linker. There is no description that a physiologically active substance having a carboxy group is bound to a conjugate via a linker in this document.

55 **[0008]** Nakanishi, T. et al describe in "Development of the polymer micelle carrier system for doxorubicin" (J. Control. Release, Vol. 74, No. 1 - 3, 2001, pages 295 - 302) a polymeric conjugate in which an amino group of doxorubicin is directly amide-bonded to a carboxylic acid group of a block copolymer without a linker. There is no description that a physiologically active substance having a carboxy group is bound to a conjugate via a linker in this document. Since the amide bond is a chemically stable, the conjugate of the above cited document cannot release the drug by the hydrolysis and the efficacy cannot be obtained.

**[0009]** EP 2 042 195 A1 teaches a high-molecular weight derivative of Combretastatin having a structure wherein a

carboxylic acid group of a copolymer of polyethylene glycol and a polymer having a carboxylic acid group at the side chain, is combined with a phenolic hydroxyl group of Combretastatin via an ester bond. The phenolic hydroxyl group of Combretastatin is directly ester-bonded to a carboxylic acid group of a block copolymer without a linker in EP 2 042 195 A1. EP 2 042 195 A1 does not disclose the feature that the carboxy group of the physiologically active substance is bound to the carboxylic acid group of the block copolymer via a specific linker.

**[0010]** EP 1 401 473 B1 discloses the reaction between the chloroacetate ester of 4-hydroxymethyl-2,6-dimethylphenol (compound 14i) and retinoid acid (compound 14ii) and the reaction occurs between the hydroxymethyl group of compound 14i and the carboxy group of compound 14ii. However, EP 1 401 473 B1 does not disclose the feature that the carboxy group of the physiologically active substance is bound to the carboxylic acid group of the block copolymer via a specific linker.

#### Disclosure of the Invention

#### Problems to be Solved by the Invention

**[0011]** In the case of the doxorubicin conjugate disclosed in Patent Document 1, a block copolymer and doxorubicin are directly bonded through an amide bond. However, the amide bond is a chemically stable bonding mode, and therefore the efficacy of such a conjugate is questionable since the in vivo release of the drug by the hydrolysis is very slow.

In fact, as a Comparative Example in Examples of the specification of the present application, the antitumor effect was measured when doxorubicin was bonded directly to a block copolymer to find rarely discernible antitumor activity.

**[0012]** In a high-molecular weight derivative of camptothecins disclosed in Patent Document 2, the bonded drugs are sustainably released by selecting drugs having a phenolic hydroxyl group exhibiting a higher hydrolysis reactivity than ordinary phenolic hydroxyl groups and forming an ester of the phenolic hydroxyl group with a carboxylic acid group of polyglutamic acid. However, this method cannot be applied to drugs having an amino group or a carboxy group.

**[0013]** In order to release a drug from a high-molecular weight conjugate disclosed in Patent Document 3, a bond between a polyethylene glycol and a linker needs to be degraded by hydrolyzing enzymes in the body. However, such hydrolyzing enzymes in the body are known to greatly vary not only among species but also among individuals even in the same species, and therefore there is also a concern that the effect of the drug release from the conjugate would be greatly different among individuals when the cleavage of the bond to the drug is dependent on the hydrolyzing enzymes.


**[0014]** A high-molecular weight conjugate disclosed in Patent Document 4 includes aspartic acid as a part of a block copolymer, and a linker is bonded to the aspartic acid. However, the bond between the aspartic acid and the linker of the high-molecular weight conjugate needs to be hydrolyzed by hydrolyzing enzymes in the body as in Patent Document 3, and therefore there is also a concern that release of the drug may vary among species as well as among individuals. The high-molecular weight conjugate disclosed in Patent Document 4 does not form a micelle, and then if the amount of the drug carried on the conjugate is increased, the water solubility may not be maintained.

#### Means for Solving the Problems

**[0015]** As a result of intensive and extensive studies on a solution for the aforementioned problems, the present inventors have found that a compound in which a residue of a physiologically active substance having a carboxy group or a residue of a physiologically active substance having an amino group is bonded to a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety through a specific linker to cause a benzyl elimination reaction can release the physiologically active substance under physiological conditions without depending on hydrolyzing enzymes. This finding leads to the completion of the present invention. In the compound, the polyaspartic acid or the polyglutamic acid may presumably form a 5- or 6-membered cyclic imide structure under physiological conditions, whereby an ester bond may be cleaved, and then, the benzyl elimination reaction proceeds to release the physiologically active substance.


**[0016]** Specifically, the present invention relates to the following (1) to (15):

(1) A high-molecular weight conjugate of physiologically active substances, comprising a structure in which a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety is bonded to a substituent represented by the general formula (I):




wherein Ar is represented by one of the following formulae:


EP 2 258 397 B1



5 R<sup>15</sup> and R<sup>16</sup> represent a hydrogen atom; and O-C(=O)-A represents a residue of the physiologically active substance having a carboxy group, wherein the conjugate is represented by the general formula (II):



25 wherein R<sup>1</sup> represents a hydrogen atom or a (C1 to C6) alkyl group; R<sup>2</sup> represents a linkage group; R<sup>3</sup> represents a hydrogen atom or a (C1 to C6) acyl group; R<sup>4</sup> represents a substituent represented by the general formula (III):




wherein Ar<sup>1</sup> is represented by one of the following formulae:




40 R<sup>17</sup> and R<sup>18</sup> represent a hydrogen atom; and O-C(=O)-A represents a residue of the physiologically active substance having a carboxy group;  
R<sup>5</sup> represents a substituent selected from the group consisting of a hydroxymethylphenoxy group optionally having a substituent, a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group, an amino acid with a protected carboxy group, and -NR<sup>6</sup>CONHR<sup>7</sup> wherein R<sup>6</sup> and R<sup>7</sup>, which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group; a represents an integer of 5 to 11,500; d, e, f, g, h, i and j each represent an integer of 0 to 200; d+e represents an integer of 1 to 200; and d+e+f+g+h+i+j represents an integer of 2 to 200; and respective units of the polyaspartic acid are bonded in any order,

50 or wherein the conjugate is represented by the general formula (IV):


55



wherein  $R^8$  represents a hydrogen atom or a (C1 to C6) alkyl group;  $R^9$  represents a linkage group;  $R^{10}$  represents a hydrogen atom or a (C1 to C6) acyl group;  $R^{11}$  represents a substituent represented by the general formula (V):



wherein  $Ar^2$  is represented by one of the following formulae:



$R^{19}$  and  $R^{20}$  represent a hydrogen atom; and  $O-C(=O)-A$  represents a residue of the physiologically active substance having a carboxy group;

$R^{12}$  represents a substituent selected from the group consisting of a hydroxymethylphenoxy group optionally having a substituent, a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group, an amino acid with a protected carboxy group, and  $-NR^{13}CONHR^{14}$  wherein  $R^{13}$  and  $R^{14}$ , which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group;  $b$  represents an integer of 5 to 11,500;  $k$  represents an integer of 1 to 200;  $m$  and  $n$  each represent an integer of 0 to 200; and  $k+m+n$  represents an integer of 2 to 200; and respective units of the polyglutamic acid are bonded in any order. (2) The high-molecular weight conjugate of physiologically active substances according to (1) above, wherein  $R^1$  is a (C1 to C3) alkyl group;  $R^2$  is a (C2 to C6) alkylene group;  $R^3$  is a (C1 to C3) acyl group;  $Ar^1$  in the general formula (III) as  $R^4$  is a phenyl group in which the bond of  $Ar^1$  with a polymer is present in the ortho or para position to the bond with  $CR^{17}R^{18}$ ; and  $a$  is an integer of 100 to 300;  $d$ ,  $e$ ,  $f$ ,  $g$ ,  $h$ ,  $i$  and  $j$  are each an integer of 0 to 100;  $d + e$  is an integer of 1 to 100; and  $d + e + f + g + h + i + j$  is an integer of 6 to 100.

(3) The high-molecular weight conjugate of physiologically active substances according to (2) above, wherein  $R^1$  is a methyl group;  $R^2$  is a trimethylene group;  $R^3$  is an acetyl group; and  $R^5$  is  $-NR^6CONHR^7$  wherein  $R^6$  and  $R^7$  are both a cyclohexyl group or an isopropyl group.

(4) The high-molecular weight conjugate of physiologically active substances according to (1) above, wherein  $R^8$  is a (C1 to C3) alkyl group;  $R^9$  is a (C2 to C6) alkylene group;  $R^{10}$  is a (C1 to C3) acyl group;  $Ar^2$  in the general formula (V) as  $R^{11}$  is a phenyl group in which the bond of  $Ar^2$  with a polymer is present in the ortho or para position to the bond with  $CR^{19}R^{20}$ ; and  $b$  is an integer of 100 to 300;  $k$  represents an integer of 1 to 90;  $m$  and  $n$  each represent an integer of 0 to 90; and  $k + m + n$  is an integer of 6 to 90.

(5) The high-molecular weight conjugate of physiologically active substances according to (4) above, wherein  $R^8$  is a methyl group;  $R^9$  is a trimethylene group;  $R^{10}$  is an acetyl group; and  $R^{12}$  is  $-NR^{13}CONHR^{14}$  wherein  $R^{13}$  and  $R^{14}$  are both a cyclohexyl group or an isopropyl group.

(6) A high-molecular weight conjugate of physiologically active substances, obtained by ester-bonding a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety to a phenolic hydroxyl group of a hydroxybenzyl alcohol compound, and bonding an alcoholic hydroxyl group of the resultant ester compound to a carboxy group of the physiologically active substance having the carboxy group.

(7) The high-molecular weight conjugate of physiologically active substances according to (1) above, wherein the physiologically active substance is an anticancer agent.

(8) The high-molecular conjugate of physiologically active substances according to (1) above, wherein the physiologically active substance is a physiologically active peptide.

(9) The high-molecular conjugate of physiologically active substances according to (8) above, wherein the physio-

logically active peptide is bestatin.

(10) The high-molecular conjugate of physiologically active substances according to (1) above, wherein the physiologically active substance is an anti-inflammatory agent.

(11) The high-molecular conjugate of physiologically active substances according to (1) above, wherein the physiologically active substance having a carboxy group is indomethacin, etodolac, mefenamic acid, diclofenac, flurbiprofen, ibuprofen, methotrexate and DMXAA.

(12) The high-molecular conjugate of physiologically active substances according to (1) above, wherein the high-molecular weight forms a micelle in water.

(13) A pharmaceutical product comprising a high-molecular conjugate of physiologically active substances according to (1) above as an active ingredient.

(14) An anticancer agent comprising a high-molecular conjugate of physiologically active substances according to (7) above as an active ingredient.


(15) An anti-inflammatory agent comprising a high-molecular conjugate of physiologically active substances according to (10) above as an active ingredient.

#### Effects of the Invention


**[0017]** The high-molecular weight conjugate of physiologically active substances according to the present invention is a compound in which a residue of the physiologically active substance having a carboxy group is bonded to a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety through a specific linker to cause a benzyl elimination reaction. As being capable of releasing the physiologically active substance under physiological conditions without depending on hydrolyzing enzymes in the body, the high-molecular weight conjugate of the present invention is expected to achieve efficacious therapeutic effects of the physiologically active substance without being affected by differences among individuals. Furthermore, the release rate can be regulated suitably by using polyaspartic acid or polyglutamic acid which can be selected in accordance with the physiologically active substances.

#### Best Mode for Carrying Out the Invention

**[0018]** In the high-molecular weight conjugate of physiologically active substances according to the present disclosure, a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety is bonded to a substituent represented by the general formula (I):



wherein Ar is represented by one of the following formulae:




; R<sup>15</sup> and R<sup>16</sup> independently represent a hydrogen atom or a (C1 to C6) alkyl group optionally having a substituent; and A represents a residue of the physiologically active substance having a carboxy group or a residue of the physiologically active substance having an amino group.

**[0019]** The block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety in the high-molecular weight conjugate of physiologically active substances according to the present disclosure is preferably formed by bonding a polyethylene glycol moiety to a polymer in which aspartic acid or glutamic acid or a derivative thereof forms a polymer backbone through amido bonds and which has carboxy groups in the side chain.

**[0020]** In the general formula (I), Ar includes aromatic hydrocarbon groups optionally having a substituent, or aromatic heterocyclic groups optionally having a substituent, and the substituents include a (C1 to C6) alkyl group, a (C1 to C6) alkoxy group, a nitro group and a halogen atom, and examples thereof include a methyl group, an ethyl group, a tert-butyl group, a methoxy group and a bromine atom. The number of substituents and the substitution positions of the substituents are not particularly limited.

**[0021]** Above all, examples of -Ar- include the following groups.



wherein N represents a nitrogen atom; and J represents an oxygen atom or a sulfur atom. According to the invention, -Ar- is represented by the following groups which are unsubstituted phenyl groups.




[0022] In the general formula (I), R<sup>15</sup> and R<sup>16</sup> independently include a hydrogen atom or a (C1-C6) alkyl group optionally having a substituent, and examples of "the (C1-C6) alkyl group optionally having a substituent" include a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an s-butyl group, an isobutyl group, an n-pentyl group and an n-hexyl group. Examples of substituents in "the (C1-C6) alkyl group optionally having a substituent" include an amino group, a methylamino group, a dimethylamino group, an ethylamino group and a diethylamino group. Preferably, R<sup>15</sup> and R<sup>16</sup> are both a hydrogen atom.

[0023] In one aspect of the disclosure, in the general formula (I), AH (H is a hydrogen atom) refers to a physiologically active substance having an amino group and is not particularly limited. However, in a preferable high-molecular weight conjugate, an amino group of the physiologically active substance having the amino group is bonded through a carbonyl group to an alcoholic hydroxyl group of a hydroxybenzyl alcohol compound which is bonded to a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety. Therefore the physiologically active substance is preferably a physiologically active substance having a primary or secondary amino group.


[0024] Examples of the physiologically active substance having an amino group (reference) include anthracycline-based anticancer agents and cytidine-based anticancer agents. Particularly preferred aspects of the disclosure are doxorubicin, daunorubicin, epirubicin, pirarubicin, amrubicin, ethynyl cytidine, CNDAC (2'-cyano-2'-deoxy-1-β-D-arabino-furanosyl cytosine), gemcitabine and the like.

Doxorubicin



## Daunorubicin


5



10

15


## Epirubicin



20

25

## Pirarubicin




30

35

40

## Amrubicin




45

50 **[0025]** The physiologically active substance in the general formula (I) may be physiologically active peptides. Using an amino group (reference) or a carboxy group at the terminals of the peptides or at other positions in the peptides, high-molecular weight conjugates carrying the peptides can be prepared. Despite their useful physiological activity in vitro, many of the peptides may not exhibit the effects in vivo because they may rapidly be degraded by hydrolyzing enzymes or other enzymes in vivo. Such peptides are expected to exhibit their effects even in vivo by being incorporated

55 **[0026]** Examples of the peptides include bestatin and bestatin methyl ester shown below, and Glufanide, Ghrelin, Tertomotide, PR1, Octreotide, Lanreotide and Pasireotide.

Bestatin


5



10

Bestatin methyl ester

15




20

**[0027]** According to the invention, in the general formula (I), the residue of a physiologically active substance having a carboxy group as A means that the formula "HO-C(=O)-A" represents the physiologically active substance. The physiologically active substance includes, but is not particularly limited to, nonsteroidal anti-inflammatory agents and anti-cancer agents, such as, for example, indomethacin, etodolac, mefenamic acid, diclofenac, flurbiprofen, ibuprofen, methotrexate and DMXAA.

25

Indomethacin


30



35

Etodolac


40



45

DXMAA


50



**[0028]** The high-molecular weight conjugate of physiologically active substances according to the present disclosure is preferably a compound represented by the aforementioned general formula (II), wherein R<sup>1</sup> represents a hydrogen atom or a (C1 to C6) alkyl group; R<sup>2</sup> represents a linkage group; R<sup>3</sup> represents a hydrogen atom or a (C1 to C6) acyl group;


55

R<sup>4</sup> represents a substituent represented by the general formula (III):



wherein

is represented by one of the following formulae:



$\text{R}^{17}$  and  $\text{R}^{18}$  independently represent a hydrogen atom or a (C1 to C6) alkyl group optionally having a substituent; and A represents a residue of the physiologically active substance having a carboxy group or a residue of the physiologically active substance having an amino group;

$\text{R}^5$  represents a substituent selected from the group consisting of a hydroxymethylphenoxy group optionally having a substituent, a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group, an amino acid with a protected carboxy group, and  $-\text{NR}^6\text{CONHR}^7$  wherein  $\text{R}^6$  and  $\text{R}^7$ , which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group; and a represents an integer of 5 to 11,500; d, e, f, g, h, i and j each represent an integer of 0 to 200; d + e represents an integer of 1 to 200; and d + e + f + g + h + i + j represents an integer of 2 to 200.

**[0029]** The (C1 to C6) alkyl group in  $\text{R}^1$  in the general formula (II) includes (C1 to C6) straight chain or branched alkyl groups, and preferred are (C1 to C4) alkyl groups, including, for example, a methyl group, an ethyl group, an n-propyl group and an n-butyl group. Particularly preferred is a methyl group.

**[0030]** The linkage group as  $\text{R}^2$  in the general formula (II) is preferably a (C2 to C6) alkylene group, including, for example, an ethylene group, a trimethylene group and a tetramethylene group. A trimethylene group is particularly preferred.

**[0031]** The (C1 to C6) acyl group as  $\text{R}^3$  in the general formula (II), not particularly limited, preferably includes (C1 to C3) acyl groups, including a formyl group, an acetyl group and an n-propionyl group. An acetyl group is particularly preferred.

**[0032]** The  $\text{Ar}^1$  in the general formula (III) as  $\text{R}^4$  in the general formula (II) includes the same groups as Ar in the aforementioned general formula (I), and preferable groups thereof are also the same.

**[0033]** The  $\text{R}^{17}$  and  $\text{R}^{18}$  in the general formula (III) as  $\text{R}^4$  in the general formula (II) include the same groups as the  $\text{R}^{15}$  and  $\text{R}^{16}$  in the aforementioned general formula (I), and preferable groups thereof are similarly a hydrogen atom.

**[0034]** The group A in the general formula (III) as  $\text{R}^4$  in the general formula (II) includes the same compounds as A in the aforementioned general formula (I), and preferable compounds thereof are also the same.

**[0035]** The  $\text{R}^5$  in the general formula (II) represents a group selected from the group consisting of a hydroxymethylphenoxy group optionally having a substituent, a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group, an amino acid with a protected carboxy group, and  $-\text{NR}^6\text{CONHR}^7$  wherein  $\text{R}^6$  and  $\text{R}^7$ , which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group. The groups selected as  $\text{R}^5$  in the general formula (II) may be the same or different in one molecule, and a block copolymer used in the high-molecular weight conjugate according to the present invention may include a single-type  $\text{R}^5$  or may exist as a mixture of compounds having different substituents as  $\text{R}^5$  in each molecule.

**[0036]** Examples of the hydroxymethylphenoxy group which optionally having a substituent include a 4-hydroxymethylphenoxy group, a 2-hydroxymethylphenoxy group, a 4-hydroxymethyl-2-methoxyphenoxy group, a 4-hydroxymethyl-2,6-dimethoxyphenoxy group, a 4-hydroxymethyl-2,6-di-tert-butylphenoxy group, a 2-hydroxymethyl-4-nitrophenoxy group, a 2-hydroxymethyl-6-methoxyphenoxy group and a 2-hydroxymethyl-4-bromophenoxy group. A 4-hydroxymethylphenoxy group, and a 2-hydroxymethylphenoxy group are preferred.

**[0037]** The (C1 to C30) alkoxy group includes straight-chain or branched (C1 to C30) alkoxy groups, and preferred is a straight-chain or branched (C1 to C10) alkoxy group, including a methoxy group, an ethoxy group, an n-propoxy group, an i-propoxy group, an n-butoxy group and a t-butoxy group. Examples of the (C1 to C30) aralkyloxy group include straight-chain or branched (C1 to C30) aralkyloxy groups, and preferred is a straight-chain or branched (C1 to C12) aralkyloxy groups, including a 4-phenylbutoxy group.

**[0038]** Examples of the (C1 to C30) alkylamino group or the di(C1 to C30) alkylamino group include straight-chain or branched (C1 to C30) alkylamino groups or a straight-chain or branched di(C1 to C30) alkylamino groups, and preferred is a straight-chain or branched (C1 to C20) alkylamino group or straight-chain or branched di(C1 to C20) alkylamino groups, including a methylamino group, an ethylamino group, an n-propylamino group, an i-propylamino group, an n-

butylamino group, a t-butylamino group, a dimethylamino group, a diethylamino group and a di(n-butyl)amino group.

**[0039]** Examples of the amino acid with a protected carboxy group include an amino acid generally used in peptide synthesis, in which a carboxyl group is protected, including, for example, a phenylalanine benzyl ester.

**[0040]** Examples of the group  $-NR^6CONHR^7$  as  $R^5$  in the general formula (II) wherein  $R^6$  and  $R^7$ , which may be the same or different, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group, include, but are not particularly limited to, for example a cyclohexylaminocarbonyl-cyclohexylamino group and an isopropylaminocarbonylisopropylamino group.

**[0041]** The polyaspartic acid moiety in the high-molecular weight conjugate of physiologically active substances represented by the aforementioned general formula (II) according to the present disclosure includes constituting units of  $\alpha$ -amino acid type,  $\beta$ -amino acid type and cyclized type. These constituting units may be bonded in any order without any particular limitation, and each type of the amino acids may be bonded to form a block-type form or a random-type form.

**[0042]** The total number of aspartic acid in the polyaspartic acid moiety in the high-molecular weight conjugate of physiologically active substances represented by the aforementioned general formula (II) is represented by "d + e + f + g + h + i + j". The total number of aspartic acids may be determined, for example, based on the amount of aspartic acid derivatives used in the preparation of the block copolymer and is about 2 to 200, preferably about 6 to 100, more preferably about 15 to 90.

**[0043]** The proportion of the number of aspartic acid residues bonded to the physiologically active substance (d + e) based on the total number of aspartic acid (d + e + f + g + h + i + j) is 1 to 100%, preferably 1 to 90%, more preferably 2 to 60%. In addition, the number of aspartic acid residues (d + e) is about 1 to 200, preferably about 1 to 100, more preferably about 1 to 90.


**[0044]** The proportion of the  $\alpha$ -amino acid type (d + f + h) based on the total number of aspartic acid residues (d + e + f + g + h + i + j) is 10 to 100%, preferably 20 to 100%. The proportion of the  $\beta$ -amino acid type (e + g + i) is 0 to 90%, preferably 0 to 80%. The proportions can appropriately be changed, for example, by suitably selecting a deprotection condition for a protecting group in the polyaspartic acid.

**[0045]** The molecular weight of the polyethylene glycol moiety in the aforementioned general formula (II) is about 300 to 500,000, preferably about 500 to 100,000, more preferably about 1,000 to 50,000. "a" in the aforementioned general formula (II) is an integer of about 5 to 11,500, preferably an integer of about 8 to 2,300, more preferably an integer of about 100 to 300.

**[0046]** Also preferred as the high-molecular weight conjugate of physiologically active substances according to the present disclosure is a compound represented by the aforementioned general formula (IV) wherein  $R^8$  represents a hydrogen atom or a (C1 to C6) alkyl group;


$R^9$  represents a linkage group;  $R^{10}$  represents a hydrogen atom or a (C1 to C6) acyl group;

$R^{11}$  represents a substituent represented by the general formula (V):



wherein  $Ar^2$

is represented by one of the following formulae:



$R^{19}$  and  $R^{20}$  independently represent a hydrogen atom or a (C1 to C6) alkyl group optionally a substituent; and A represents a residue of the physiologically active substance having a carboxy group or a residue of the physiologically active substance having an amino group;

$R^{12}$  represents a substituent selected from the group consisting of a hydroxymethylphenoxy group optionally having a substituent, a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group, an amino acid with a protected carboxy group, and  $-NR^{13}CONHR^{14}$  wherein  $R^{13}$  and  $R^{14}$ , which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group; and b represents an integer of 5 to 11,500; k represents an integer of 1 to 200; m and n each represent an integer of 0 to 200, and k + m + n represents an integer of 2 to 200.

**[0047]** The (C1 to C6) alkyl group as  $R^8$  in the general formula (IV) includes the same groups as in  $R^1$  in the aforementioned general formula (II), and preferable groups thereof are the same, and a methyl group is particularly preferred.

**[0048]** The linkage group as  $R^9$  in the general formula (IV) is preferably a (C2 to C6) alkylene group, and examples

thereof include an ethylene group, a trimethylene group and a tetramethylene group. Above all, a trimethylene group is particularly preferred.

[0049] The (C1 to C6) acyl group as R<sup>10</sup> in the general formula (IV) includes the same groups as in R<sup>3</sup> in the aforementioned general formula (II), and preferable groups thereof are the same, and an acetyl group is particularly preferred.

[0050] Ar<sup>2</sup> in the general formula (V) as R<sup>11</sup> in the general formula (IV) includes the same groups as Ar in the aforementioned general formula (I), and preferable groups thereof are the same.

[0051] The R<sup>19</sup> and R<sup>20</sup> in the general formula (V) as R<sup>11</sup> in the general formula (IV) include the same groups as R<sup>15</sup> and R<sup>16</sup> in the aforementioned general formula (I), and a preferable group thereof is similarly a hydrogen atom.

[0052] "A" in the general formula (V) as R<sup>11</sup> in the general formula (IV) includes the same compounds as A in the aforementioned general formula (I), and preferable compounds thereof are also the same.

[0053] The R<sup>12</sup> in the general formula (IV) represents a group selected from the group consisting of a hydroxymethylphenoxy group optionally having a substituent, a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group, an amino acid with a protected carboxy group, and -NR<sup>13</sup>CONHR<sup>14</sup> wherein R<sup>13</sup> and R<sup>14</sup>, which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group. The groups selected as R<sup>12</sup> in the general formula (IV) may be the same or different in one molecule, and a block copolymer used in the high-molecular weight conjugate according to the present invention may include a single-type R<sup>12</sup> or may exist as a mixture of compounds having different substituents as R<sup>12</sup> in each molecule.

[0054] The hydroxymethylphenoxy group optionally having a substituent includes the same groups as the hydroxymethylphenoxy group which may have a substituent in R<sup>5</sup> in the aforementioned general formula (II), and preferable groups thereof are the same.

[0055] The (C1 to C30) alkoxy group, the (C1 to C30) aralkyloxy group, the (C1 to C30) alkylamino group and the di(C1 to C30) alkylamino group as R<sup>12</sup> in the general formula (IV) include the same groups as the (C1 to C30) alkoxy group, the (C1 to C30) aralkyloxy group, the (C1 to C30) alkylamino group and the di(C1 to C30) alkylamino group as R<sup>5</sup> in the aforementioned general formula (II).

[0056] The amino acid with a protected carboxy group for R<sup>12</sup> in the general formula (IV) includes amino acids having a carboxy group protected, used in the common peptide synthesis, and examples thereof include phenylalanine benzyl ester.

[0057] The group -NR<sup>13</sup>CONHR<sup>14</sup> wherein R<sup>13</sup> and R<sup>14</sup>, which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group, for R<sup>12</sup> in the general formula (IV) includes the same groups as the -NR<sup>6</sup>CONHR<sup>7</sup> as R<sup>5</sup> in the general formula (II), and preferable groups thereof are also the same.

[0058] The total number of glutamic acid residues of the polyglutamic acid moiety in the high-molecular weight conjugate of physiologically active substances represented by the aforementioned general formula (IV) according to the present invention is represented by k + m + n and is about 2 to 200, preferably about 6 to 90, more preferably 6 to 60.

[0059] The proportion of the number of glutamic acid residues (k) to which the physiologically active substance is bonded on the basis of the total number of glutamic acid residues (k + m + n) is 1 to 100%, preferably 3 to 90%, more preferably 4 to 60%. The number of glutamic acid residues (k) is 1 to 200, preferably about 1 to 90, more preferably about 2 to 30.

[0060] Each constituting units of the glutamic acid moiety in the high-molecular weight conjugate of physiologically active substances represented by the aforementioned general formula (IV) may be bonded in any order, and amino acids of each type may be bonded to form a block-type form or a random-type form.

[0061] "b" in the aforementioned general formula (IV) is an integer of about 5 to 11,500, preferably an integer of about 8 to 2,300, more preferably an integer of about 100 to 300.

[0062] The molecular weight of the polyethylene glycol moiety in the aforementioned general formula (IV) is about 300 to 500,000, preferably about 500 to 100,000, more preferably about 1,000 to 50,000.

[0063] The molecular weight of the high-molecular weight conjugate of physiologically active substances according to the present invention is about 500 to 600,000, preferably about 600 to 110,000, more preferably about 1,100 to 80,000. According to the present invention, the term "molecular weight" refers to a weight average molecular weight determined by the GPC method.

[0064] The high-molecular weight conjugate of physiologically active substances according to the present invention may form a micelle having a polyethylene glycol moiety as the outer shell in water.

[0065] The high-molecular weight conjugate of physiologically active substances according to the present disclosure is obtained by ester-bonding a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety to a phenolic hydroxyl group of a hydroxybenzyl alcohol compound having an alcoholic hydroxyl group protected by using a dehydrating condensation agent in an organic solvent to form an ester compound, and deprotecting the protecting group, and then, bonding the alcoholic hydroxyl group of the ester compound to a carboxy group of the physiologically active substance having the carboxy group or bonding the alcoholic

hydroxyl group of the ester compound to an amino group of the physiologically active substance having the amino group through a carbonyl group. The manufacturing method of the high-molecular weight conjugate according to the present invention is also included in the present disclosure. An intermediate to which the hydroxybenzyl alcohol compound is bonded is also included in the present disclosure. That is, the high-molecular weight conjugate of the present invention is obtained by a method described below. For example, a block copolymer of a polyethylene glycol moiety and a polyaspartic acid prepared according to the method disclosed in Japanese Patent No. 3268913 and a hydroxybenzyl alcohol compound in which an alcoholic hydroxyl group and functional groups other than phenolic hydroxyl groups are protected as necessary are subjected to a reaction using a dehydrating condensation agent such as dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIPC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride salt (WSC) and 1-ethoxycarbonyl-2-ethoxy-1,2-dihydroxyquinolinone (EEDQ) at a temperature of 0 to 180°C, preferably 5 to 50°C in an organic solvent in which both of the block copolymer and the hydroxybenzyl alcohol compound are dissolved, preferably in an aprotic polar solvent such as N,N-dimethylformamide (DMF), 1,3-dimethyl-2-imidazolidinone (DMI) and N-methylpyrrolidone (NMP). In the case where a physiologically active substance having a carboxy group (according to the invention) is bonded thereafter, the protecting group of the alcoholic hydroxyl group at the benzyl-position is deprotected, and then, the alcoholic hydroxyl group at the benzyl-position is ester-bonded to the carboxy group of the physiologically active substance in which functional groups other than the carboxy group have been protected as necessary by using the aforementioned dehydrating condensation agent in the aforementioned solvent, and the protecting groups are deprotected as necessary. In the case where a physiologically active substance having an amino group (reference) is bonded, the protecting group of the alcoholic hydroxyl group at the benzyl-position is deprotected, and thereafter, the alcoholic hydroxyl group at the benzyl-position is condensed through a carbonyl group with the amino group of the physiologically active substance in which functional groups other than the amino group have been protected as necessary by using disuccinyl carbonate, carbonyldiimidazole or the like in the aforementioned solvent at 0 to 30°C, and the protecting groups are deprotected as necessary.

**[0066]** In the manufacturing method of the present disclosure, for example, a block copolymer of a polyethylene glycol moiety and a polyglutamic acid prepared by the method according to Japanese Patent Laid-Open (Kokai) No. 5-955 may be used in place of the aforementioned block copolymer of a polyethylene glycol moiety and a polyaspartic acid.

**[0067]** In the condensation reaction, a reaction aid such as N,N-dimethylaminopyridine (DMAP) may be used.

**[0068]** In addition, a high-molecular weight conjugate of physiologically active substances in which R<sup>5</sup> in the general formula (II) is the group -NR<sup>6</sup>CONHR<sup>7</sup> wherein R<sup>6</sup> and R<sup>7</sup>, which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group, or a high-molecular weight conjugate of a physiologically active substance in which R<sup>12</sup> in the general formula (IV) is the group -NR<sup>13</sup>CONHR<sup>14</sup> wherein R<sup>13</sup> and R<sup>14</sup>, which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group may also be obtained by a reaction using the aforementioned carbodiimides as a condensation agent.

**[0069]** As a method for manufacturing a compound represented by the general formula (II) or (IV) in which R<sup>5</sup> or R<sup>12</sup> in the compound is a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group or an amino acid with a protected carboxy group, there can be mentioned a method in which a carboxy group of the block copolymer is first activated by a method used in an ordinary condensation reaction, and then reacted with a corresponding alcohol, a corresponding amine, a corresponding amino acid with a protected carboxy group in an amount to be introduced under basic conditions; and a method in which a corresponding alcohol, a corresponding amine, a corresponding amino acid with a protected carboxy group is first activated by a method used in an ordinary condensation reaction, and is then reacted with a polymer.

**[0070]** It is also possible to re-activate any unreacted carboxy groups remained after the bonding of the hydroxybenzyl alcohol compound in a similar reaction, and then to introduce into the re-activated carboxy groups a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group, an amino acid with a protected carboxy group. Alternatively, different alcohols, amines and the like may be repeatedly reacted to synthesize a mixture of compounds having various substituents as the R<sup>5</sup> or the R<sup>12</sup>, with which the hydroxybenzyl alcohol compound may subsequently be condensed.

**[0071]** However, the method for manufacturing the high-molecular weight conjugate of physiologically active substances according to the present invention is not limited to the aforementioned methods.

**[0072]** The high-molecular weight conjugate of physiologically active substances according to the present invention can be used as a pharmaceutical product which is indicated for a disease for which the physiologically active substance carried on the conjugate have an efficacy. Examples of the pharmaceutical products include anticancer agents and anti-inflammatory agents. The high-molecular weight conjugate according to the present invention can be used in a dosage form which is conventionally used, including injections, tablets, and powders. For formulation process, pharmaceutically acceptable carriers which are conventionally used, for example, binding agents, lubricating agents, disintegrating agents, solvents, vehicles, solubilizing agents, dispersing agents, stabilizing agents, suspending agents, preservatives, soothing agents, colorants and flavors can also be used.

**[0073]** The high-molecular weight conjugate of physiologically active substances according to the present invention is used preferably as an injection, and usually water, a physiological saline, a 5% glucose or mannitol liquid, a water-soluble organic solvent (for example, glycerol, ethanol, dimethylsulfoxide, N-methylpyrrolidone, polyethylene glycol, cremophor, and a mixture thereof) or a mixture of water and the water-soluble organic solvents can be used.

**[0074]** The dosage of the high-molecular weight conjugate of physiologically active substances according to the present invention can vary as a matter of course, depending on the physiologically active substance as well as the sexuality, ages, physiological conditions, pathological conditions and the like of patients. The conjugate is parenterally administered, typically at a dose of 0.01 to 500 mg/m<sup>2</sup>, preferably 0.1 to 250 mg/m<sup>2</sup>, as an active ingredient per day in an adult. The administration by injection is performed intravenously, intraarterially, or into an affected site (a tumor site, an inflammation site, and the like), for example.

#### Examples

**[0075]** Hereinafter, the present invention will be illustrated more specifically with reference to Examples.

**[0076]** The Gaussian distribution analysis was conducted using a ZetaPotential/Particlerizer NICOMP 380ZLS (manufactured by Particle Sizing Systems Co.).

#### Example 1

Synthesis of Compound 1 (a phenyl ester conjugate of a block copolymer consisting of a methoxypolyethylene glycol moiety having a molecular weight of 5,000 and a polyaspartic acid moiety having a polymerization number of 30 and 4-(hydroxymethyl)phenol: in the general formula (II), R<sup>1</sup> = Me (a methyl group), R<sup>2</sup> = a trimethylene group, R<sup>3</sup> = Ac (an acetyl group), R<sup>4</sup> = 4-(hydroxymethyl)phenoxy group, R<sup>5</sup> = an isopropylaminocarbonylisopropylamino group, and d + e + f + g + h + i + j = 30, and a = 113)

**[0077]** A methoxypolyethylene glycol-polyaspartic acid block copolymer (a polymerization number of aspartic acid: 30, 1.0 g) prepared by the method according to Japanese Patent No. 3268913, and 4-(tert-butyltrimethylsilyloxyethyl)-phenol (847 mg) prepared by the method according to Patent Document 4 were dissolved in DMF (8 ml), and DMAP (43 mg) and DIPC (1.11 ml) were added at 15°C. The mixture was stirred for 20 hours. To the reaction solution, ethanol (12 ml), ethyl acetate (12 ml) and diisopropyl ether (96 ml) were added, the mixture was stirred at room temperature for 2 hours, and thereafter, a precipitate was collected by filtration and washed with ethanol/diisopropyl ether (1/4 (v/v), 10 ml). The resultant precipitate was dissolved in acetonitrile (15 ml), and then 4N hydrochloric acid/dioxane (0.38 ml) was added at 0°C, and the mixture was stirred at 0°C for 15 min. To the reaction solution, ethanol (45 ml) and diisopropyl ether (180 ml) were added, and the mixture was stirred at room temperature for 30 min. The supernatant liquid was separated, and ethanol/diisopropyl ether (1/4 (v/v), 100 ml) was further added to the residue, and the mixture was stirred for 15 min. Then the supernatant liquid was again separated, and the residue was dried under reduced pressure to obtain Compound 1 (880 mg).

**[0078]** The content of 4-(hydroxymethyl) phenol bonded in Compound 1 was determined by analyzing, by HPLC (high performance liquid chromatography), 4-(hydroxymethyl)phenol released by adding 1N sodium hydroxide aqueous solution to Compound 1, agitating the solution at 40°C for 1 hour and then adding acetic acid to neutralize the solution. As a result, the content of 4-(hydroxymethyl)phenol was determined as 9.1% (w/w), and the proportion of d + e based on d + e + f + g + h + i + j was 22.5%.

**[0079]** According to this method, an isopropylaminocarbonylisopropylamino group may be added as R<sup>5</sup>, and an existing ratio of the group could be determined by <sup>1</sup>H-NMR (hydrogen nuclear magnetic resonance spectrum) using Compound 1 dissolved in sodium deuterioxide/deuterium oxide/deuterated acetonitrile. The proportion of the isopropylaminocarbonylisopropylamino group in the polyaspartic acid, that is, the proportion of f + g based on d + e + f + g + h + i + j was 7.6%. The remaining aspartic acid residues were in the form of a free carboxylic acid (h + i) or a cyclic structure (j).

#### Example 2

(reference)

Synthesis of Compound 2 (a conjugate of Compound 1 and doxorubicin)

**[0080]** Compound 1 (500 mg) obtained in Example 1 and disuccinyl carbonate (933 mg) were dissolved in DMF (30 ml), and triethylamine (0.254 ml) was added thereto at 15°C. The mixture was stirred for 20 hours. To the reaction solution, ethanol (90 ml) and diisopropyl ether (360 ml) were added, and the mixture was stirred at room temperature for 30 min. The supernatant liquid was separated, and ethanol/diisopropyl ether (1/4 (v/v), 100 ml) was further added to

the residue, and the mixture was stirred for 15 min. Thereafter, the supernatant liquid was again separated, and then the residue was dried under reduced pressure. The thus obtained solid was dissolved in DMF (25 ml), and doxorubicin hydrochloride (116 mg) and triethylamine (0.028 ml) dissolved in DMF (10 ml) were added thereto at 15°C, and the mixture was stirred for 2 hours. To the reaction solution, ethanol (100 ml) and diisopropyl ether (400 ml) were added, and the mixture was stirred at room temperature for 1 hour. The supernatant liquid was separated, and ethanol/diisopropyl ether (1/4 (v/v), 150 ml) was further added to the residue. The mixture was stirred for 15 min. Thereafter, the supernatant liquid was again separated, and the residue was dried under reduced pressure. The resulting solid was dissolved in acetonitrile (30 ml) and water (3 ml), and the solution was applied to a column of an ion exchange resin (5 ml of Dowex 50(H<sup>+</sup>), manufactured by The Dow Chemical Co.) and eluted with acetonitrile/water (10/1 (v/v), 15 ml). To the resulting elution fraction, water (60 ml) was added, and acetonitrile was then distilled off under reduced pressure, followed by lyophilization to obtain Compound 2 (564 mg).

**[0081]** The content of doxorubicin bonded in Compound 2 was determined by analyzing by HPLC (high performance liquid chromatography) an aglycon (adriamycinone) of doxorubicin released by adding 1N hydrochloric acid (1 ml) to Compound 2 (3.38 mg) and agitating the mixture at 40°C for 1 hour, and comparing the analysis result with a calibration curve of the aglycon obtained by subjecting doxorubicin to the same treatment. As a result, the content of doxorubicin bonded was determined as 11.5% (w/w). In Compound 2, no free doxorubicin was detected.

**[0082]** Gaussian distribution analysis using an aqueous solution of Compound 2 (1 mg/ml) revealed that the average particle diameter is 23 nm (volume weighting). Therefore, Compound 2 was considered to form micelles in water.

### Example 3

Synthesis of Compound 3 (a phenyl ester conjugate of a block copolymer consisting of a methoxypolyethylene glycol moiety having a molecular weight of 12,000 and a polyglutamic acid moiety having a polymerization number of 23 and 4-(hydroxymethyl)phenol: in the general formula (IV), R<sup>8</sup> = Me (a methyl group), R<sup>9</sup> = a trimethylene group, R<sup>10</sup> = Ac (an acetyl group), R<sup>11</sup> = a 4-(hydroxymethyl)phenoxy group, R<sup>12</sup> = an isopropylaminocarbonylisopropylamino group, k + m + n = 23, and b = 273)

**[0083]** A methoxypolyethylene glycol-polyglutamic acid block copolymer (a polymerization number of glutamic acid: 23, 1.0 g) prepared by the method according to Japanese Patent Laid-Open No. 5-955 and 4-(tert-butyltrimethylsilyloxy-methyl)phenol (295 mg) prepared by the method according to Patent Document 4 were dissolved in DMF (15 ml), and DMAP (19 mg) and DIPC (0.485 ml) were added at 27°C. The mixture was stirred at the same temperature for 20 hours. To the reaction solution, ethanol (23 ml), ethyl acetate (23 ml) and diisopropyl ether (180 ml) were added, and the mixture was stirred at room temperature for 2 hours. Thereafter, the precipitate was collected by filtration and washed with ethanol/diisopropyl ether (1/4 (v/v), 20 ml). The resultant precipitate was dissolved in acetonitrile (15 ml), and 4N hydrochloric acid/dioxane (0.38 ml) was then added at 0°C. The mixture was stirred for 15 min. To the reaction solution, ethanol (45 ml) and diisopropyl ether (180 ml) were added, and the mixture was stirred at room temperature for 30 min. The supernatant liquid was separated, ethanol/diisopropyl ether (1/4 (v/v), 100 ml) was further added to the residue, and the mixture was stirred for 15 min. Thereafter, the supernatant liquid was again separated, and the residue was dried under reduced pressure to obtain Compound 3 (1.1 g).

**[0084]** The content of 4-(hydroxymethyl)phenol bonded in Compound 3 was determined by analyzing by HPLC (high performance liquid chromatography) 4-(hydroxymethyl)phenol released by adding a 1N sodium hydroxide aqueous solution to Compound 3, agitating the mixture at 40°C for 1 hour and then adding acetic acid to neutralize the solution. As a result, the content of 4-(hydroxymethyl)phenol was determined as 9.4% (w/w), and the proportion of k based on k + m + n was determined as 53.8%.

**[0085]** According to this method, an isopropylaminocarbonylisopropylamino group can be added as R<sup>12</sup>, and an existing ratio of the group could be determined by <sup>1</sup>H-NMR (hydrogen nuclear magnetic resonance spectrum) using Compound 3 dissolved in sodium deuterioxide/deuterium oxide/deuterated acetonitrile. The proportion of the isopropylaminocarbonylisopropylamino group to polyglutamic acid, that is, the proportion of m based on k + m + n was 32.0%. The remaining glutamic acid residues were in the form of a free carboxylic acid (n).

### Example 4 (reference)

Synthesis of Compound 4 (a conjugate of Compound 3 and doxorubicin)

**[0086]** Compound 3 (500 mg) obtained in Example 3 and disuccinyl carbonate (966 mg) were dissolved in DMF (30 ml), and triethylamine (0.263 ml) was added at 15°C. The mixture was stirred for 20 hours. To the reaction solution, ethanol (90 ml) and diisopropyl ether (360 ml) were added, and the mixture was stirred at room temperature for 30 min. The supernatant liquid was separated, ethanol/diisopropyl ether (1/4 (v/v), 100 ml) was further added to the residue,

## EP 2 258 397 B1

and the mixture was stirred for 15 min. Thereafter, the supernatant liquid was again separated, and the residue was dried under reduced pressure. The resulting solid was dissolved in DMF (25 ml), and doxorubicin hydrochloride (120 mg) and triethylamine (0.029 ml) dissolved in DMF (10 ml) were added at 15°C. The mixture was stirred for 2 hours. To the reaction solution, ethanol (100 ml) and diisopropyl ether (400 ml) were added, and the mixture was stirred at room temperature for 1 hour. The supernatant liquid was separated, and ethanol/diisopropyl ether (1/4 (v/v), 150 ml) was further added to the residue, and the mixture was stirred for 15 min. Thereafter, the supernatant liquid was again separated, and the residue was dried under reduced pressure. The resulting solid was dissolved in acetonitrile (30 ml) and water (3 ml), and the resulting solution was applied to a column of an ion exchange resin (5 ml of Dowex 50(H<sup>+</sup>), made by The Dow Chemical Co.), and eluted with acetonitrile/water (10/1 (v/v), 15 ml). To the resulting elution fraction, water (60 ml) was added, and then, acetonitrile was distilled off under reduced pressure. Thereafter, the resultant fraction was lyophilized to obtain Compound 4 (611 mg).

**[0087]** The content of doxorubicin bonded in Compound 4 was determined by analyzing by HPLC (high performance liquid chromatography) an aglycon (adriamycinone) of doxorubicin released by adding 1N hydrochloric acid (1 ml) to Compound 4 (3.13 mg) and agitating the mixture at 40°C for 1 hour, and comparing the analysis result with a calibration curve of the aglycon obtained by subjecting doxorubicin to the same treatment. As a result, the content of doxorubicin bonded was determined as 17.1% (w/w). In Compound 4, no free doxorubicin was detected.

**[0088]** Gaussian distribution analysis using an aqueous solution of Compound 4 (1 mg/ml) revealed that the average particle diameter is 43 nm (volume weighting). Therefore, Compound 4 was considered to form micelles in water.

### Example 5 (reference)

Synthesis of Compound 5 (a conjugate of Compound 3 and bestatin methyl ester)

**[0089]** Compound 3 (50 mg) obtained in Example 3 and disuccinyl carbonate (97 mg) were dissolved in DMF (3 ml), and triethylamine (0.026 ml) was added at 15°C. The mixture was stirred for 20 hours. To the reaction solution, ethanol (9 ml) and diisopropyl ether (36 ml) were added, and the mixture was stirred at room temperature for 30 min. The supernatant liquid was separated, ethanol/diisopropyl ether (1/4 (v/v), 10 ml) was further added to the residue, and the mixture was stirred for 15 min. Thereafter, the supernatant liquid was again separated, and the residue was dried under reduced pressure. The resulting solid was dissolved in DMF (2 ml), and bestatin methyl ester hydrochloride (8.8 mg) disclosed in Japanese Patent Laid-Open (Kokai) No. 2001-131066 and triethylamine (0.003 ml) dissolved in DMF (0.5 ml) were added at 15°C. The mixture was stirred for 2 hours. To the reaction solution, ethanol (7.5 ml) and diisopropyl ether (30 ml) were added, and the mixture was stirred at room temperature for 1 hour. The supernatant liquid was separated, and ethanol/diisopropyl ether (1/4 (v/v), 10 ml) was further added to the residue. The mixture was stirred for 15 min. Thereafter, the supernatant liquid was again separated, and the residue was dried under reduced pressure. The resulting solid was dissolved in acetonitrile (5 ml) and water (0.5 ml), and the solution was applied to a column of an ion exchange resin (1 ml of Dowex 50(H<sup>+</sup>), manufactured by The Dow Chemical Co.), and eluted with acetonitrile/water (10/1 (v/v), 3 ml). To the resulting elution fraction, water (6 ml) was added, and then, acetonitrile was distilled off under reduced pressure. Thereafter, the resulting fraction was lyophilized to obtain Compound 5 (55 mg).

**[0090]** The content of bestatin methyl ester bonded in Compound 5 was determined by analyzing by HPLC (high performance liquid chromatography) bestatin methyl ester released by dissolving Compound 5 (4.89 mg) in a 1 mol/l sodium methoxide methanol solution (1 ml) and allowing a reaction at room temperature for 1 hour, and then comparing the analysis result with a calibration curve of bestatin methyl ester. As a result, the content of bestatin methyl ester bonded was determined as 14.6% (w/w). In Compound 5, no free bestatin methyl ester was detected.

**[0091]** Gaussian distribution analysis using an aqueous solution of Compound 5 (1 mg/ml) revealed that the average particle diameter is 41 nm (volume weighting). Therefore, Compound 5 was considered to form micelles in water.

### Example 6

Synthesis of Compound 6 (a conjugate of Compound 3 and etodolac)

**[0092]** Compound 3 (50 mg) obtained in Example 3 and etodolac (manufactured by Wako Pure Chemical Industries, Ltd., 10.8 mg) were dissolved in DMF (0.5 ml), and DMAP (0.5 mg) and DIPC (0.012 ml) were added at 15°C. The mixture was stirred for 20 hours. To the reaction solution, ethanol (1.5 ml) and diisopropyl ether (6 ml) were added, and the mixture was stirred at room temperature for 1 hour. Thereafter, the precipitate was collected by filtration and washed with ethanol/diisopropyl ether (1/4 (v/v), 3 ml). The resultant precipitate was dissolved in acetonitrile (3 ml) and water (0.3 ml), and then applied to a column of an ion exchange resin (0.5 ml of Dowex 50(H<sup>+</sup>), manufactured by The Dow Chemical Co.) and eluted with acetonitrile/water (10/1 (v/v), 3 ml). To the resultant elution fraction, water (6 ml) was added, and acetonitrile was distilled off under reduced pressure. Thereafter, the resultant fraction was lyophilized to

## EP 2 258 397 B1

obtain Compound 6 (56 mg).

[0093] The content of etodolac bonded in Compound 6 was determined by analyzing by HPLC (high performance liquid chromatography) etodolac released by dissolving Compound 6 (10.57 mg) in a 1 mol/l sodium hydroxide aqueous solution (10 ml) and allowing the solution to react at room temperature for 1 hour to neutralize the solution with acetic acid, and comparing the analysis result with a calibration curve of etodolac. As a result, the content of etodolac bonded was determined as 11.6% (w/w). In Compound 6, no free etodolac was detected.

[0094] Gaussian distribution analysis using an aqueous solution of Compound 6 (1 mg/ml) revealed that the average particle diameter is 49 nm (volume weighting). Therefore, Compound 6 was considered to form micelles in water.

### Example 7

Synthesis of Compound 7 (a conjugate of Compound 3 and indomethacin)

[0095] Compound 3 (50 mg) obtained in Example 3 and indomethacin (manufactured by Wako Pure Chemical Industries, Ltd., 13.5 mg) were dissolved in DMF (0.5 ml), and DMAP (0.5 mg) and DIPC (0.012 ml) were added at 15°C. The mixture was stirred for 20 hours. To the reaction solution, ethanol (1.5 ml) and diisopropyl ether (6 ml) were added, and the mixture was stirred at room temperature for 1 hour. Thereafter the precipitate was collected by filtration and washed with ethanol/diisopropyl ether (1/4 (v/v), 3 ml). The resultant precipitate was dissolved in acetonitrile (3 ml) and water (0.3 ml), and then applied to a column of an ion exchange resin (0.5 ml of Dowex 50(H<sup>+</sup>), manufactured by The Dow Chemical Co.) and eluted with acetonitrile/water (10/1 (v/v), 3 ml). To the resultant elution fraction, water (6 ml) was added, and then, acetonitrile was distilled off under reduced pressure. Thereafter, the resultant fraction was lyophilized to obtain Compound 7 (58 mg).

[0096] The content of indomethacin bonded to Compound 7 was determined by analyzing by HPLC (high performance liquid chromatography) a decomposed substance (a 4-chlorobenzoic acid eliminated substance) of indomethacin released by dissolving Compound 7 (8.59 mg) in a 1 mol/l sodium hydroxide aqueous solution (10 ml), allowing the solution to react at room temperature for 1 hour and then neutralizing the solution with acetic acid, and comparing the analysis result with a calibration curve of the same decomposed substance obtained from indomethacin by the same treatment. As a result, the content of indomethacin bonded was determined as 13.3% (w/w). In Compound 7, no free indomethacin was detected.

[0097] Gaussian distribution analysis using an aqueous solution of Compound 7 (1 mg/ml) revealed that the average particle diameter is 61 nm (volume weighting). Therefore, Compound 7 was considered to form a micelle in water. Comparative Example 1

Synthesis of Comparative Compound 1 (an amide conjugate of a block copolymer consisting of a methoxypolyethylene glycol moiety having a molecular weight of 5,000 and a polyaspartic acid moiety having a polymerization number of 30 and doxorubicin)

[0098] A methoxypolyethylene glycol-polyaspartic acid block copolymer (a polymerization number of aspartic acid: 30, 240 mg) prepared by the method according to Japanese Patent No. 3268913 was dissolved in DMF (10 ml), and N-hydroxysuccinic acid imide (197 mg) and DIPC (0.267 ml) were then added at 35°C and stirred for 30 min. To the reaction solution, ethyl acetate (20 ml) and diisopropyl ether (100 ml) were added, and the mixture was stirred at room temperature for 1 hour. Thereafter, the precipitate was filtered, washed with ethyl acetate/diisopropyl ether (1/4 (v/v), 10 ml), and dried under reduced pressure. The resultant precipitate was dissolved in DMF (6 ml), and doxorubicin hydrochloride (130 mg) and triethylamine (0.038 ml) dissolved in DMF (6 ml) were added at 25°C. The mixture was stirred for 2 hours. To the reaction solution, ethyl acetate (24 ml) and diisopropyl ether (96 ml) were added, and the mixture was stirred at room temperature for 1 hour. The precipitate was filtered, washed with ethyl acetate/diisopropyl ether (1/4 (v/v), 10 ml), and dried under vacuum. The resultant precipitate was dissolved in acetonitrile (30 ml) and water (3 ml) and then applied to a column of an ion exchange resin (5 ml of Dowex 50(H<sup>+</sup>), manufactured by The Dow Chemical Co.) and eluted with acetonitrile/water (10/1 (v/v), 15 ml). To the resultant elution fraction, water (36 ml) was added, and then, acetonitrile was distilled off under reduced pressure. Thereafter, the resultant fraction was lyophilized to obtain Comparative Compound 1 (338 mg).

[0099] The content of doxorubicin bonded in Comparative Compound 1 was determined by analyzing by HPLC (high performance liquid chromatography) an aglycon (adriamycinone) of doxorubicin released by adding 1N hydrochloric acid (1 ml) to Comparative Compound 1 (5.16 mg) and stirring the mixture at 40°C for 1 hour, and comparing the analysis result with a calibration curve of the aglycon obtained by subjecting doxorubicin to the same treatment. As a result, the content of doxorubicin bonded was determined as 31.6% (w/w). In Comparative Compound 1, no free doxorubicin was detected.

Test Example 1 (reference)

Drug Release in the Absence of Enzymes (Doxorubicin)

5 **[0100]** Compound 2 and Compound 4 which are the high-molecular weight conjugates according to the present invention, and Comparative Compound 1 were each dissolved in a concentration of 1 mg/ml in PBS (phosphate-buffered physiological saline, pH: 7.1), and incubated at 37°C. Doxorubicin released from the high-molecular weight conjugate was analyzed and quantified by HPLC in comparison with a calibration curve. The proportion of the quantified value to the total amount of the drug determined from the content of the drug in the high-molecular weight conjugate is shown in Figure 1.

10 **[0101]** As is clear from Figure 1, the high-molecular weight conjugates (Compound 2 and Compound 4) according to the present invention significantly released doxorubicin even in the absence of hydrolyzing enzymes. Particularly Compound 2 having a polyaspartic acid moiety released doxorubicin more rapidly than Compound 4 having a glutamic acid moiety. Incidentally, in Compound 2, the amount of doxorubicin released after 24 hours appeared to be decreased probably because doxorubicin released into the solution was partially degraded in the solution. By contrast, release of doxorubicin from Comparative Compound 1 having the amido bond was not confirmed even after 24 hours. These results indicate an excellent drug release performance of the high-molecular weight conjugate according to the present invention in the absence of hydrolyzing enzymes.

20 Test Example 2

(reference)

Drug Release in the Absence of Enzymes (Bestatin methyl ester)

25 **[0102]** Compound 5 which is a high-molecular weight conjugate according to the present invention was dissolved in PBS (phosphate-buffered physiological saline, pH: 7.1) at a concentration of 1 mg/ml and incubated at 37°C. Bestatin methyl ester released from the high-molecular weight conjugate was analyzed and quantified by HPLC in comparison with a standard curve. The proportion of the quantified value based on the total amount of the drug determined from the drug content in the high-molecular weight conjugate is shown in Figure 2.

30 **[0103]** As is clear from Figure 2, the high-molecular weight conjugate according to the present invention significantly released bestatin methyl ester even without hydrolyzing enzymes. This result indicates the excellent drug release performance of the high-molecular weight conjugate according to the present invention in the absence of hydrolyzing enzymes.

35 Test Example 3

Drug Release in the Absence of Hydrolyzing Enzymes

40 (Etodolac and Indomethacin)

**[0104]** Compound 6 and Compound 7 were each dissolved in PBS (phosphate-buffered physiological saline, pH: 7.1) at a concentration of 1 mg/ml and incubated at 37°C. Etodolac or indomethacin released from the high-molecular weight conjugate was analyzed and quantified by HPLC in comparison with a calibration curve. The proportion of the quantified value based on the total amount of the drug determined from the drug content of the high-molecular weight conjugate is shown in Figure 3.

45 **[0105]** As is clear from Figure 3, the high-molecular weight conjugate according to the present invention significantly released etodolac or indomethacin even without hydrolyzing enzymes. These results demonstrate an excellent drug release performance of the high-molecular weight conjugate according to the present invention in the absence of hydrolyzing enzymes.

50 Test Example 4 (reference) Antitumor activity

**[0106]** Mouse colon cancer, Colon 26, maintained by serial subcutaneous subculture in mice, was minced into about 2-mm cubic fragments, and the fragments were subcutaneously transplanted on the dorsal part of female CDF1 mice with a trocar. Eight days after the tumor transplantation, the high-molecular weight conjugate according to the present invention (Compound 2 and Compound 4) or a control drug (Comparative Compound 1 and doxorubicin hydrochloride) was each administered once into the mouse tail vein in a way such that each mouse received the same dose per body

## EP 2 258 397 B1

weight in terms of doxorubicin. Each compound was dissolved in a 5% liquid glucose for injection and used. After the administration, the major axis (L mm) and the minor axis (W mm) of the tumor were measured using a caliper, and the tumor volume was calculated by the formula  $(L \times W^2)/2$ . Table 1 shows the relative tumor based on the tumor volume on the day of administration.

[Table 1]

	Dose	Days after administration				
		0	3	7	10	14
Compound 2	60 mg/kg	1.00	0.73	0.22	0.22	0.72
	30 mg/kg	1.00	1.58	2.94	3.59	6.67
Compound 4	120 mg/kg	1.00	0.89	0.33	0.17	0.03
	60 mg/kg	1.00	1.65	1.51	0.78	0.71
	30 mg/kg	1.00	1.96	2.76	2.17	3.44
Comparative Compound 1	240 mg/kg	1.00	3.19	9.89	22.68	42.49
Doxorubicin	15 mg/kg	1.00	1.79	3.89	5.27	12.17
Control		1.00	4.37	12.55	17.94	30.69

**[0107]** The lethal dose of doxorubicin used as the control drug was 30 mg/kg, and therefore doxorubicin was administered at a dose of 15 mg/kg to perform the antitumor test. As is clear from Table 1, the high-molecular weight conjugate according to the present invention had antitumor activity at a broadened range of the doses and exhibited a higher antitumor effect compared with doxorubicin. By contrast, Comparative Compound 1 exhibited no antitumor effect at all even in a dose of 240 mg/kg.

### Brief Description of the Drawings


#### **[0108]**

Figure 1 (reference) shows the proportion of the amount of doxorubicin released from Compound 2, Compound 4 and Comparative Compound 1, based on the total amounts of bound doxorubicin in PBS solutions (pH: 7.1, 37°C). Figure 2 (reference) shows the proportion of the amount of bestatin methyl ester from Compound 5, based on the total amount of bound bestatin methyl ester in a PBS solution (pH: 7.1, 37°C).


Figure 3 shows the proportion of the amount of etodolac or indomethacin released from Compound 6 or Compound 7, based on the total amount of the bound etodolac or indomethacin in a PBS solution (pH: 7.1, 37°C).

### Claims

1. A high-molecular weight conjugate of physiologically active substances, comprising a structure in which a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety is bonded to a substituent represented by the general formula (I):



wherein Ar is represented by one of the following formulae:




R<sup>15</sup> and R<sup>16</sup> represent a hydrogen atom; and O-C(=O)-A represents a residue of the physiologically active substance having a carboxy group, wherein the conjugate is represented by the general formula (II):



wherein Ar<sup>2</sup>

is represented by one of the following formulae:

5



10

; R<sup>19</sup> and R<sup>20</sup> represent a hydrogen atom; and O-C(=O)-A represents a residue of the physiologically active substance having a carboxy group;

15

R<sup>12</sup> represents a substituent selected from the group consisting of a hydroxymethylphenoxy group optionally having a substituent, a (C1 to C30) alkoxy group, a (C1 to C30) aralkyloxy group, a (C1 to C30) alkylamino group, a di(C1 to C30) alkylamino group, an amino acid with a protected carboxy group, and -NR<sup>13</sup>CONHR<sup>14</sup> wherein R<sup>13</sup> and R<sup>14</sup>, which may be the same or different from each other, represent a (C3 to C6) cyclic alkyl group or a (C1 to C5) alkyl group which may be substituted with a tertiary amino group; b represents an integer of 5 to 11,500; k represents an integer of 1 to 200; m and n each represent an integer of 0 to 200; and k+m+n represents an integer of 2 to 200; and respective units of the polyglutamic acid are bonded in any order.

20

2. The high-molecular weight conjugate of physiologically active substances according to claim 1, wherein R<sup>1</sup> is a (C1 to C3) alkyl group; R<sup>2</sup> is a (C2 to C6) alkylene group; R<sup>3</sup> is a (C1 to C3) acyl group; Ar<sup>1</sup> in the general formula (III) of R<sup>4</sup> is a phenyl group in which the bond of Ar<sup>1</sup> with a polymer is present in the ortho or para position to the bond with CR<sup>17</sup>R<sup>18</sup>; and a is an integer of 100 to 300; d, e, f, g, h, i and j are each an integer of 0 to 100; d+e is an integer of 1 to 100; and d+e+f+g+h+i+j is an integer of 6 to 100.

25

3. The high-molecular weight conjugate of physiologically active substances according to claim 2, wherein R<sup>1</sup> is a methyl group; R<sup>2</sup> is a trimethylene group; R<sup>3</sup> is an acetyl group; and R<sup>5</sup> is -N R<sup>6</sup>CONHR<sup>7</sup> wherein R<sup>6</sup> and R<sup>7</sup> are both a cyclohexyl group or an isopropyl group.

30

4. The high-molecular weight conjugate of physiologically active substances according to claim 1, wherein R<sup>8</sup> is a (C1 to C3) alkyl group; R<sup>9</sup> is a (C2 to C6) alkylene group; R<sup>10</sup> is a (C1 to C3) acyl group; Ar<sup>2</sup> in the general formula (V) as R<sup>11</sup> is a phenyl group in which the bond of Ar<sup>2</sup> with a polymer is present in the ortho or para position to the bond with CR<sup>19</sup>R<sup>20</sup>; and b is an integer of 100 to 300, k represents an integer of 1 to 90, m and n each represent an integer of 0 to 90; and k+m+n is an integer of 6 to 90.

35

5. The high-molecular weight conjugate of physiologically active substance according to claim 4, wherein R<sup>8</sup> is a methyl group; R<sup>9</sup> is a trimethylene group; R<sup>10</sup> is an acetyl group; and R<sup>12</sup> is -NR<sup>13</sup>CONHR<sup>14</sup> wherein R<sup>13</sup> and R<sup>14</sup> are both a cyclohexyl group or an isopropyl group.

40

6. The high-molecular weight conjugate of physiologically active substances according to claim 1, obtained by:

ester-bonding a side chain carboxy group of a block copolymer having a polyethylene glycol moiety and a polyaspartic acid moiety or a polyglutamic acid moiety to a phenolic hydroxyl group of a hydroxybenzyl alcohol compound; and

45

bonding an alcoholic hydroxyl group of the resultant ester compound with a carboxy group of the physiologically active substance having the carboxy group.

50

7. The high-molecular weight conjugate of physiologically active substances according to claim 1, wherein the physiologically active substance is an anticancer agent.

8. The high-molecular weight conjugate of physiologically active substances according to claim 1, wherein the physiologically active substance is a physiologically active peptide.

9. The high-molecular weight conjugate of physiologically active substances according to claim 8, wherein the physiologically active peptide is bestatin.


55

10. The high-molecular weight conjugate of physiologically active substances according to claim 1, wherein the physiologically active substance is an anti-inflammatory agent.


11. The high-molecular weight conjugate of physiologically active substances according to claim 1, wherein the physiologically active substance having a carboxy group is indomethacin, etodolac, mefenamic acid, diclofenac, flurbiprofen, ibuprofen, methotrexate and DMXAA.
12. The high-molecular weight conjugate of physiologically active substances according to claim 1, wherein the high-molecular weight conjugate forms a micelle in water.
13. A pharmaceutical product comprising a high-molecular weight conjugate of physiologically active substances according to claim 1 as an active ingredient.
14. An anticancer agent comprising a high-molecular weight conjugate of physiologically active substances according to claim 7 as an active ingredient.
15. An anti-inflammatory agent comprising a high-molecular weight conjugate of physiologically active substances according to claim 10 as an active ingredient.

Patentansprüche


1. Hochmolekulares Konjugat von physiologisch aktiven Substanzen, umfassend eine Struktur, in der eine Seitenketten-Carboxygruppe eines Blockcopolymers, das eine Polyethylenglykoleinheit und eine Polyasparaginsäureeinheit oder eine Polyglutaminsäureeinheit aufweist, an einen durch die allgemeine Formel (I) dargestellten Substituenten gebunden ist:




wobei Ar durch eine der folgenden Formeln dargestellt wird:



R<sup>15</sup> und R<sup>16</sup> ein Wasserstoffatom darstellen; und O-C(=O)-A einen Rest der physiologisch aktiven Substanz mit einer Carboxygruppe darstellt, wobei das Konjugat dargestellt ist durch die allgemeine Formel (II):




wobei R<sup>1</sup> ein Wasserstoffatom oder eine (C1- bis C6-) Alkylgruppe darstellt; R<sup>2</sup> eine Bindungsgruppe darstellt; R<sup>3</sup> ein Wasserstoffatom oder eine (C1- bis C6-) Acylgruppe darstellt; R<sup>4</sup> einen durch die allgemeine Formel (III) dargestellten Substituenten darstellt:



wobei Ar<sup>1</sup>  
 durch eine der folgenden Formeln dargestellt wird:

5



10

R<sup>17</sup> und R<sup>18</sup> ein Wasserstoffatom darstellen; und O-C(=O)-A einen Rest der physiologisch aktiven Substanz mit einer Carboxygruppe darstellt;


R<sup>5</sup> einen Substituenten darstellt, der ausgewählt ist aus der Gruppe, bestehend aus einer Hydroxymethylphenoxygruppe, die gegebenenfalls einen Substituenten aufweist, einer (C1- bis C30-) Alkoxygruppe, einer (C1- bis C30-) Aralkyloxygruppe, einer (C1- bis C30-) Alkylaminogruppe, einer Di-(C1-bis C30-) Alkylaminogruppe, einer Aminosäure mit einer geschützten Carboxylgruppe und -NR<sup>6</sup>CONHR<sup>7</sup>, wobei R<sup>6</sup> und R<sup>7</sup>, die gleich oder verschieden voneinander sein können, eine zyklische (C3- bis C6-) Alkylgruppe oder eine (C1- bis C5-) Alkylgruppe darstellen, die mit einer tertiären Aminogruppe substituiert sein kann; a eine ganze Zahl von 5 bis 11.500 darstellt; d, e, f, g, h, i und j jeweils eine ganze Zahl von 0 bis 200 darstellen; d + e eine ganze Zahl von 1 bis 200 darstellt; und d + e + f + g + h + i + j eine ganze Zahl von 2 bis 200 darstellt; und jeweilige Einheiten der Polyasparaginsäure in beliebiger Reihenfolge gebunden sind,

15

20

oder wobei das Konjugat dargestellt ist durch die allgemeine Formel (IV):


25



30


wobei R<sup>8</sup> ein Wasserstoffatom oder eine (C1- bis C6-) Alkylgruppe darstellt; R<sup>9</sup> eine Bindungsgruppe darstellt; R<sup>10</sup> ein Wasserstoffatom oder eine (C1- bis C6-) Acylgruppe darstellt; R<sup>11</sup> einen durch die allgemeine Formel (V) dargestellten Substituenten darstellt:

35



wobei Ar<sup>2</sup>  
 durch eine der folgenden Formeln dargestellt wird:

40



45

R<sup>19</sup> und R<sup>20</sup> ein Wasserstoffatom darstellen; und O-C(=O)-A einen Rest der physiologisch aktiven Substanz mit einer Carboxygruppe darstellt;

R<sup>12</sup> einen Substituenten darstellt, der ausgewählt ist aus der Gruppe, bestehend aus einer Hydroxymethylphenoxygruppe, die gegebenenfalls einen Substituenten aufweist, einer (C1- bis C30-) Alkoxygruppe, einer (C1- bis C30-) Aralkyloxygruppe, einer (C1- bis C30-) Alkylaminogruppe, einer Di-(C1-bis C30-) Alkylaminogruppe, einer Aminosäure mit einer geschützten Carboxygruppe und -NR<sup>13</sup>CONHR<sup>14</sup>, wobei R<sup>13</sup> und R<sup>14</sup>, die gleich oder verschieden voneinander sein können, eine zyklische (C3- bis C6-) Alkylgruppe oder eine (C1- bis C5-) Alkylgruppe darstellen, die mit einer tertiären Aminogruppe substituiert sein kann; b eine ganze Zahl von 5 bis 11.500 darstellt; k eine ganze Zahl von 1 bis 200 darstellt; m und n jeweils eine ganze Zahl von 0 bis 200 darstellen; und k + m + n eine ganze Zahl von 2 bis 200 darstellt; und jeweilige Einheiten der Polyglutaminsäure in beliebiger Reihenfolge gebunden sind.

50


55

## EP 2 258 397 B1


2. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 1, wobei  $R^1$  eine (C1- bis C3-) Alkylgruppe darstellt;  $R^2$  eine (C2- bis C6-) Alkylgruppe darstellt;  $R^3$  eine (C1- bis C3-) Acylgruppe darstellt;  $Ar^1$  in der allgemeinen Formel (III) von  $R^4$  eine Phenylgruppe darstellt, in der die Bindung von  $Ar^1$  mit einem Polymer in ortho- oder paraStellung zu der Bindung mit  $CR^{17}R^{18}$  vorliegt; und a eine ganze Zahl von 100 bis 300 darstellt; d, e, f, g, h, i und j jeweils eine ganze Zahl von 0 bis 100 darstellen; d + e eine ganze Zahl von 1 bis 100 darstellt; und d + e + f + g + h + i + j eine ganze Zahl von 6 bis 100 darstellt.
3. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 2, wobei  $R^1$  eine Methylgruppe darstellt;  $R^2$  eine Trimethylengruppe darstellt;  $R^3$  eine Acetylgruppe darstellt; und  $R^5$  - $NR^6CONHR^7$  darstellt, wobei  $R^6$  und  $R^7$  beide eine Cyclohexylgruppe oder eine Isopropylgruppe sind.
4. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 6, wobei  $R^8$  eine (C1- bis C3-) Alkylgruppe darstellt;  $R^9$  eine (C2- bis C6-) Alkylgruppe darstellt;  $R^{10}$  eine (C1- bis C3-) Acylgruppe darstellt;  $Ar^2$  in der allgemeinen Formel (V) als  $R^{11}$  eine Phenylgruppe darstellt, in der die Bindung von  $Ar^2$  mit einem Polymer in ortho- oder paraStellung zu der Bindung mit  $CR^{19}R^{20}$  vorliegt; und b eine ganze Zahl von 100 bis 300 darstellt; k eine ganze Zahl von 1 bis 90 darstellt, m und n jeweils eine ganze Zahl von 0 bis 90 darstellen; und k + m + n eine ganze Zahl von 6 bis 90 darstellt.
5. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 4, wobei  $R^8$  eine Methylgruppe darstellt;  $R^9$  eine Trimethylengruppe darstellt;  $R^{10}$  eine Acetylgruppe darstellt; und  $R^{12}$  - $NR^{13}CONHR^{14}$  darstellt, wobei  $R^{13}$  und  $R^{14}$  beide eine Cyclohexylgruppe oder eine Isopropylgruppe sind.
6. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 1, erhalten durch:
  - 25 Ester-Bindung einer Seitenketten-Carboxygruppe eines Blockcopolymers mit einer Polyethylenglykoleinheit und einer Polyasparaginsäureeinheit oder einer Polyglutaminsäureeinheit an eine phenolische Hydroxylgruppe einer Hydroxybenzylalkoholverbindung; und
  - Bindung einer alkoholischen Hydroxylgruppe der resultierenden Esterverbindung mit einer Carboxygruppe der physiologisch aktiven Substanz, welche die Carboxygruppe aufweist.
7. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 1, wobei die physiologisch aktive Substanz ein Antikrebsmittel ist.
8. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 1, wobei die physiologisch aktive Substanz ein physiologisch aktives Peptid ist.
9. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 8, wobei das physiologisch aktive Peptid Bestatin ist.
10. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 1, wobei die physiologisch aktive Substanz ein entzündungshemmendes Mittel ist.
11. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 1, wobei die physiologisch aktive Substanz, die eine Carboxygruppe aufweist, Indomethacin, Etodolac, Mefenaminsäure, Diclofenac, Flurbiprofen, Ibuprofen, Methotrexat und DMXAA ist.
12. Hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 1, wobei das hochmolekulare Konjugat in Wasser eine Mizelle bildet.
13. Pharmazeutisches Produkt, umfassend ein hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 1 als Wirkstoff.
14. Antikrebsmittel, umfassend ein hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 7 als Wirkstoff.
15. Entzündungshemmendes Mittel, umfassend ein hochmolekulares Konjugat von physiologisch aktiven Substanzen nach Anspruch 10 als Wirkstoff.

Revendications

1. Conjugué de haut poids moléculaire de substances physiologiquement actives, comprenant une structure dans laquelle un groupe carboxy de chaîne latérale d'un copolymère séquencé ayant une fraction polyéthylène glycol et une fraction acide polyaspartique ou une fraction acide polyglutamique est lié à un substituant représenté par la formule générale (I) :




dans laquelle Ar est représenté par l'une des formules suivantes :




ou



R<sup>15</sup> et R<sup>16</sup> représentent un atome d'hydrogène ; et O-(C=O)-A représente un résidu de la substance physiologiquement active ayant un groupe carboxy, où le conjugué est représenté par la formule générale (II) :



dans laquelle R<sup>1</sup> représente un atome d'hydrogène ou un groupe alkyle en C1 à C6 ; R<sup>2</sup> représente un groupe de liaison ; R<sup>3</sup> représente un atome d'hydrogène ou un groupe acyle en C1 à C6 ; R<sup>4</sup> représente un substituant représenté par la formule générale (III) :



dans laquelle Ar<sup>1</sup> est représenté par une des formules suivantes :




ou

EP 2 258 397 B1




5 ; R<sup>17</sup> et R<sup>18</sup> représentent un atome d'hydrogène ; et O-C(=O)A représente un résidu de la substance physiologiquement active ayant un groupe carboxy ;  
 R<sup>5</sup> représente un substituant choisi dans le groupe constitué par un groupe hydroxyméthylphénoxy ayant éventuellement un substituant, un groupe alcoxy en C1 à C30, un groupe aralkyloxy en C1 à C30, un groupe alkylamino en C1 à C30, un groupe alkylamino en di (C1 à C30), un acide aminé avec un groupe carboxy protégé, et -NR<sup>6</sup>CONHR<sup>7</sup> dans laquelle R<sup>6</sup> et R<sup>7</sup>, qui peuvent être identiques ou différents l'un de l'autre, représentent un groupe alkyle cyclique en C3 à C6 ou un groupe alkyle en C1 à C5 qui peut être substitué avec un groupe amino tertiaire ; a représente un nombre entier de 5 à 11500 ; d, e, f, g, h, i et j représentent chacun un nombre entier de 0 à 200 ; d + e représente un nombre entier de 1 à 200 ; et d + e + f + g + h + i + j représente un nombre entier de 2 à 200 ; et les unités respectives de l'acide polyaspartique sont liées dans n'importe quel ordre,

ou bien où le conjugué est représenté par la formule générale (IV) :



dans laquelle R<sup>8</sup> représente un atome d'hydrogène ou un groupe alkyle en C1 à C6 ; R<sup>9</sup> représente un groupe de liaison ; R<sup>10</sup> représente un atome d'hydrogène ou un groupe acyle en C1 à C6 ; R<sup>11</sup> représente un substituant représenté par la formule générale (V) :



dans laquelle Ar<sup>2</sup> est représenté par une des formules suivantes :



40 ou



; R<sup>19</sup> et R<sup>20</sup> représentent un atome d'hydrogène ; et OC(=O)-A représente un résidu de la substance physiologiquement active ayant un groupe carboxy ;

R<sup>12</sup> représente un substituant choisi dans le groupe constitué par un groupe hydroxyméthylphénoxy ayant éventuellement un substituant, un groupe alcoxy en C1 à C30, un groupe aralkyloxy en C1 à C30, un groupe alkylamino en C1 à C30, un groupe alkylamino di(C1 à C30), un acide aminé avec un groupe carboxy protégé, et -NR<sup>13</sup>CONHR<sup>14</sup> dans laquelle R<sup>13</sup> et R<sup>14</sup>, qui peuvent être identiques ou différents l'un de l'autre, représentent un groupe alkyle cyclique en C3 à C6, ou un groupe alkyle en C1 à C5 qui peut être substitué avec un groupe amino tertiaire ; b représente un nombre entier de 5 à 11500 ; k représente un nombre entier de 1 à 200 ; m et n représentent chacun un nombre entier de 0 à 200 ; et k + m + n représente un nombre entier de 2 à 200 ; et les unités respectives de l'acide polyglutamique sont liées dans n'importe quel ordre.

2. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, dans lequel R<sup>1</sup> est un groupe alkyle en C1 à C3 ; R<sup>2</sup> est un groupe alkylène en C2 à C6 ; R<sup>3</sup> est un groupe acyle en C1 à C3 ;

## EP 2 258 397 B1

Ar<sup>1</sup> dans la formule générale (III) de R<sup>4</sup> est un groupe phényle dans lequel la liaison de Ar<sup>1</sup> avec un polymère est présente en position ortho ou para à la liaison avec CR<sup>17</sup>R<sup>18</sup>; et a est un nombre entier de 100 à 300 ; d, e, f, g, h, i et j sont chacun un nombre entier de 0 à 100 ; d + e est un nombre entier de 1 à 100 ; et d + e + f + g + h + i + j est un nombre entier de 6 à 100.

5

3. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 2, dans lequel R<sup>1</sup> est un groupe méthyle ; R<sup>2</sup> est un groupe triméthylène ; R<sup>3</sup> est un groupe acétyle ; et R<sup>5</sup> est -NR<sup>6</sup>CONHR<sup>7</sup> dans laquelle R<sup>6</sup> et R<sup>7</sup> sont tous les deux un groupe cyclohexyle ou un groupe isopropyle.

10

4. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, dans lequel R<sup>8</sup> est un groupe alkyle en C1 à C3 ; R<sup>9</sup> est un groupe alkylène en C2 à C6 ; R<sup>10</sup> est un groupe acyle en C1 à C3 ; Ar<sup>2</sup> dans la formule générale (V) comme R<sup>11</sup> est un groupe phényle dans lequel la liaison de Ar<sup>2</sup> avec un polymère est présente en position ortho ou para à la liaison avec CR<sup>19</sup>R<sup>20</sup>; et b est un nombre entier de 100 à 300, k représente un nombre entier de 1 à 90, met n représentent chacun un nombre entier de 0 à 90 ; et k + m + n est un nombre entier de 6 à 90.

15

5. Conjugué de haut poids moléculaire de substance physiologiquement active selon la revendication 4, dans lequel R<sup>8</sup> est un groupe méthyle ; R<sup>9</sup> est un groupe triméthylène ; R<sup>10</sup> est un groupe acétyle ; et R<sup>12</sup> est -NR<sup>13</sup>CONHR<sup>14</sup> dans laquelle R<sup>13</sup> et R<sup>14</sup> sont tous les deux un groupe cyclohexyle ou un groupe isopropyle.

20

6. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, obtenu par :

liaison ester d'un groupe carboxy de chaîne latérale d'un copolymère séquencé ayant une fraction polyéthylène glycol et une fraction acide polyaspartique ou une fraction acide polyglutamique à un groupe hydroxyle phénolique d'un composé alcool hydroxybenzylique ; et liaison d'un groupe hydroxyle alcoolique du composé ester résultant avec un groupe carboxy de la substance physiologiquement active ayant le groupe carboxy.

25

7. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, dans lequel la substance physiologiquement active est un agent anti-cancéreux.

30

8. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, dans lequel la substance physiologiquement active est un peptide physiologiquement actif.

35

9. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, dans lequel le peptide physiologiquement actif est la bestatine.

10. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, dans lequel la substance physiologiquement active est un agent anti-inflammatoire

40

11. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, dans lequel la substance physiologiquement active ayant un groupe carboxy est l'indométhacine, l'étodolac, l'acide méfénamique, le diclofénac, le flurbiprofène, l'ibuprofène, le méthotraxate et le DMXAA.

45

12. Conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1, dans lequel le conjugué de haut poids moléculaire forme une micelle dans l'eau.

13. Produit pharmaceutique comprenant un conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 1 comme principe actif.

50

14. Agent anti-cancéreux comprenant un conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 7 comme principe actif.

15. Agent anti-inflammatoire comprenant un conjugué de haut poids moléculaire de substances physiologiquement actives selon la revendication 10 comme principe actif.

55

Figure 1




Figure 2





Figure 3



**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Patent documents cited in the description**

- JP 2694923 B [0004]
- WO 200439869 A [0004]
- JP 2002508400 A [0004]
- JP 2004532289 A [0004]
- EP 1580216 A1 [0005]
- WO 02065988 A2 [0006]
- EP 2042195 A1 [0009]
- EP 1401473 B1 [0010]
- JP 3268913 B [0065] [0077] [0098]
- JP 5000955 A [0066] [0083]
- JP 2001131066 A [0089]

**Non-patent literature cited in the description**

- Effective drug delivery by PEGylated drug conjugates. **GREENWALD R. B. et al.** Advanced Drug Delivery Reviews. Elsevier BV, 2003, vol. 55, 217-250 [0007]
- **NAKANISHI, T. et al.** Development of the polymer micelle carrier system for doxorubicin. *J. Control. Release*, 2001, vol. 74 (1 - 3), 295-302 [0008]