



(11) **EP 2 321 640 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
30.11.2016 Bulletin 2016/48

(51) Int Cl.:
A61K 38/48 ^(2006.01) **A61K 9/20** ^(2006.01)
A61P 1/00 ^(2006.01) **A61P 3/00** ^(2006.01)

(21) Application number: **09808534.3**

(86) International application number:
PCT/US2009/004791

(22) Date of filing: **21.08.2009**

(87) International publication number:
WO 2010/021752 (25.02.2010 Gazette 2010/08)

(54) **FORMULATION FOR ORAL ADMINISTRATION OF PROTEINS**

FORMULIERUNG ZUR ORALEN VERABREICHUNG VON PROTEINEN

FORMULATION POUR UNE ADMINISTRATION ORALE DE PROTÉINES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL
PT RO SE SI SK SM TR**

(30) Priority: **21.08.2008 US 90831 P**

(43) Date of publication of application:
18.05.2011 Bulletin 2011/20

(73) Proprietor: **Immunogenics LLC
Corona del Mar, CA 92526 (US)**

(72) Inventors:
• **SIEGEL, Matthew, John
San Mateo
CA 94403 (US)**
• **BERNER, Bret
Half Moon Bay
CA 94019 (US)**

(74) Representative: **Brasnett, Adrian Hugh et al
Mewburn Ellis LLP
City Tower
40 Basinghall Street
London EC2V 5DE (GB)**

(56) References cited:
WO-A1-99/41361 WO-A2-2008/115428

US-A- 4 477 434 US-A- 5 466 547
US-A1- 2003 007 959 US-A1- 2003 215 438
US-A1- 2004 058 870 US-A1- 2004 248 821
US-A1- 2005 249 719 US-A1- 2006 019 941
US-A1- 2006 083 727 US-A1- 2007 140 978
US-A1- 2007 197 439 US-A1- 2007 243 254
US-A1- 2008 125 437 US-A1- 2008 193 436
US-B1- 6 291 533

- **GASS ET AL: "Combination Enzyme Therapy for Gastric Digestion of Dietary Gluten in Patients With Celiac Sprue", GASTROENTEROLOGY, ELSEVIER, PHILADELPHIA, PA, vol. 133, no. 2, 3 August 2007 (2007-08-03), pages 472-480, XP022198873, ISSN: 0016-5085, DOI: 10.1053/J.GASTRO.2007.05.028**
- **BETHUNE M T ET AL: "Heterologous Expression, Purification, Refolding, and Structural-Functional Characterization of EP-B2, a Self-Activating Barley Cysteine Endoprotease", CHEMISTRY AND BIOLOGY, CURRENT BIOLOGY, LONDON, GB, vol. 13, no. 6, June 2006 (2006-06), pages 637-647, XP027991171, ISSN: 1074-5521 [retrieved on 2006-06-01]**
- **SIEGEL M ET AL: "Rational Design of Combination Enzyme Therapy for Celiac Sprue", CHEMISTRY AND BIOLOGY, CURRENT BIOLOGY, LONDON, GB, vol. 13, no. 6, June 2006 (2006-06), pages 649-658, XP027991172, ISSN: 1074-5521 [retrieved on 2006-06-01]**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 321 640 B1

Description

FIELD OF THE INVENTION

[0001] This invention relates to methods and compositions for stabilizing proteases intended for oral administration against degradation in the stomach and in gastric fluid and in other physiological fluids that may oxidize and degrade these proteins. The present invention therefore relates to the fields of biology, chemistry, molecular biology, medicinal chemistry, medicine, and pharmacology.

DESCRIPTION OF RELATED DISCLOSURES

[0002] Oral delivery of proteins to the gastrointestinal tract is known to be problematic, as a result of poor protein stability due to both gastric pH and proteolytic digestion by enzymes, as well as poor absorption. Many attempts have been made to deliver proteins by the oral route or locally to the stomach. US Patents 7,291,598; 7,265,097; and 7,244,709, for example, all use nanoparticles containing chitosan to improve the absorption and stability of proteins in the gastrointestinal (GI) tract. US Patent 6,541,606 utilizes crystalline proteins to avoid degradation in the stomach and for treatment of diseases of the GI tract and discusses potential bioadhesion to Peyer's patches to improve protein absorption from the GI tract. US Patent 7,351,741 describes additives to improve absorption of proteins from the GI tract. These patents describe the problems of stability of proteins in the GI tract due to acidic pH in the stomach and due to enzymatic digestion as well as poor absorption from the GI tract. The instability of proteins in the GI tract or stomach due to oxidation or non-proteolytic digestion is not described by any of the aforementioned references.

[0003] Lactase, a β -galactosidase that metabolizes lactose, and papain, a cysteine protease, are both enzymes administered orally as dietary supplements. Lactase is administered as a digestive aid for individuals that are deficient in the ability to digest lactose. For example, lactase from *Aspergillus oryzae* is administered as Lactaid® (Pleasantville, NJ) as a dietary supplement, and US Patents 6,660,313 and 6,562,339 describe formulations of lactase for oral administration in dietary supplements. Papain, a cysteine protease, is administered as a digestive aid. However, its optimal temperature is 65°C, and it is not clear that papain is an active enzyme in the stomach.

[0004] US Patents 7,320,788 and 7,303,871 describe the use of glutenases, alone and in combination, as orally administered enzymes to digest gluten for treatment (and prevention of symptoms) of Celiac sprue. One of the enzymes described as a glutenase is a recombinant version of a cysteine endoprotease originally isolated from barley. In 1953, it was first recognized that ingestion of gluten, a common dietary protein present in wheat, barley and rye causes disease in sensitive individuals. Gluten

is a complex mixture of glutamine- and proline-rich glutenin and prolamine molecules, which is thought to be responsible for disease induction. Ingestion of such proteins by sensitive individuals produces flattening of the normally luxurious, rug-like, epithelial lining of the small intestine known to be responsible for efficient and extensive terminal digestion of peptides and other nutrients. Clinical symptoms of celiac sprue include fatigue, chronic diarrhea, malabsorption of nutrients, weight loss, abdominal distension, anemia, as well as a substantially enhanced risk for the development of osteoporosis and intestinal malignancies (lymphoma and carcinoma). The disease has an incidence of approximately 1 in 200 in European populations.

[0005] A related disease is dermatitis herpetiformis, which is a chronic eruption characterized by clusters of intensely pruritic vesicles, papules, and urticaria-like lesions. IgA deposits occur in almost all normal-appearing and perilesional skin. Asymptomatic gluten-sensitive enteropathy is found in 75 to 90% of dermatitis herpetiformis patients and in some of their relatives. Onset is usually gradual. Itching and burning are severe, and scratching often obscures the primary lesions with eczematization of nearby skin, leading to an erroneous diagnosis of eczema. Strict adherence to a gluten-free diet for prolonged periods may control the disease in some patients. Dapsone, sulfapyridine and colchicines are sometimes prescribed for relief of itching due to dermatitis herpetiformis.

[0006] Celiac sprue and dermatitis herpetiformis are generally considered to be an autoimmune disease, and the antibodies found in the serum of the patients indicate an immunological basis of the disease. Antibodies to tissue transglutaminase (tTG) and gliadin appear in almost 100% of the patients with active Celiac sprue, and the presence of such antibodies, particularly of the IgA class, has been used in diagnosis of the disease.

[0007] The large majority of Celiac sprue and dermatitis herpetiformis patients express the HLA-DQ2 [DQ(a1*0501, b1*02)] and/or DQ8 [DQ(a1*0301, b1*0302)] molecules. It is believed that intestinal damage is caused by interactions between specific gliadin oligopeptides and HLA-DQ2 or DQ8, which in turn induce proliferation of T lymphocytes in the sub-epithelial layers. T helper 1 cells and cytokines apparently play a major role in a local inflammatory process leading to villus atrophy of the small intestine.

[0008] At the present time, the only treatment for Celiac sprue is the strict avoidance of all foods containing gluten, although clinical trials of glutenases are ongoing. While gluten withdrawal has transformed the prognosis for children diagnosed with Celiac sprue and substantially improved it for adult patients, some people still die of the disease, mainly adults who had severe disease at the outset. An important cause of death is lymphoreticular disease (especially intestinal lymphoma). It is not known whether a gluten-free diet diminishes this risk. Apparent clinical remission is often associated with histologic relapse that is detected only by review biopsies or by in-

creased EMA titers.

[0009] Gluten is so widely used, for example in commercial soups, sauces, ice creams, hot dogs, and other foods, that patients need detailed lists of foodstuffs to avoid and expert advice from a dietitian familiar with Celiac sprue. Ingesting even small amounts of gluten may prevent remission or induce relapse. Supplementary vitamins, minerals, and hematinics may also be required, depending on the degree of deficiency experienced by a particular patient. A few patients respond poorly or not at all to gluten withdrawal, either because the diagnosis is incorrect or because the disease is refractory. In the latter case, oral corticosteroids (e.g., prednisone 10 to 20 mg bid) may induce response.

[0010] Perhaps the most promising new therapy in clinical development is the use of glutenases as described in US Patents 7,320,788 and 7,303,871 (see also PCT Pub. Nos. 2008/115428; 2008/115411; 2007/044906; and 2007/047303; and US 20080193436) to prevent and/or treat the symptoms of Celiac sprue and/or dermatitis herpetiformis by decreasing the levels of toxic gluten oligopeptides in foodstuffs, either prior to or after ingestion by a patient. These patent publications disclose that certain gluten oligopeptides are resistant to cleavage by gastric and pancreatic enzymes, that the presence of such peptides in the small intestine results in toxic effects in Celiac sprue (and dermatitis herpetiformis) patients, and that enzymatic treatment can remove such peptides and their toxic effects. By digestion with glutenases, these toxic oligopeptides are cleaved into fragments, thereby preventing or relieving their toxic effects in Celiac sprue or dermatitis herpetiformis patients.

[0011] Many glutenases comprise cysteine and/or methionine residues, and proteins containing cysteine or methionine can be subject to oxidation during storage either as bulk enzyme or as dosage forms, and enzymes with cysteine in the active site, specifically cysteine proteases, can be inactivated by oxidation of cysteine. The stability of these proteins during storage or after lyophilization may be improved by adding acetylcysteine or methionine to scavenge free radical attack.

[0012] A variety of antioxidants are known to be useful as preservatives during storage of pharmaceuticals and foods, including sulfur-containing antioxidants, free thiols, such as cysteine, homocysteine, thioglycerol, acetylcysteine, as well as sodium sulfite, sodium metabisulfite, sodium thiosulfate, and the like, disulfides (such as dithiothreitol and α -lipoic acid where free thiols can be generated), and other classes of antioxidants, such as chelating agents, such as EDTA, ascorbic acid, gallic acid and its derivatives, tocopherol and its derivatives, and others known in the art to prevent oxidation. Selection of an antioxidant that will work for a particular protein is often not predictable and involves trial and error.

[0013] Because prevention of oxidation in dosage forms involves limited volumes of drug, small amounts of antioxidant can be effective, because the concentration of the antioxidant in the dosage form is relatively

large. However, the amount of antioxidant in the dosage form would have negligible activity in the larger volume of the stomach, which can be 1 liter in volume or more.

[0014] Certain cysteine proteases, in particular, papain and certain cathepsins, have been reported to be deactivated by S-nitrosylation of the active site cysteine (see Wang et al. (2000) JBC 2002 277(21):18568-73; Ascenzi et al. (2001) Curr. Protein Pept. Sci. 2: 137-153.; and Venturini et al. (2000) Biochem. Biophys. Res. Commun. 270: 437-441.). Certain sulfhydryl containing NO donors have been demonstrated to reversibly inhibit this enzyme by S-NO bond formation at the active site, Cys25. This inhibition was reversed by dithiothreitol, but not by ascorbic acid. The researchers presumed that a disulfide bond between the papain and the S-NO donor was responsible for the inhibition.

[0015] Nitrate in saliva has been shown, in rats, to be converted to nitrite, which can then be further converted to NO at fasted gastric pH of 2 (Bjorne et al. (2004) J. Clin. Invest. 113: 106-114). Under such conditions, increased NO formation from nitrites and increased blood flow, presumably mediated by NO, were reportedly observed. It is known that, from the serosal side, nitric oxide acts on the stomach to influence gastrointestinal motility, possibly by forming S-nitrosyl thiol protein intermediates.

[0016] Thus, there remains a need for pharmaceutical formulations and unit dosage forms that can improve stability of a protein in the fed stomach. The present invention meets this and other needs.

SUMMARY OF THE INVENTION

[0017] The present invention provides pharmaceutical formulations of a protein suitable for oral administration that stabilizes the protein in the fed stomach. The present invention provides a solid pharmaceutical formulation of a barley endoproteinase B2 (EPB2) or a recombinant form thereof suitable for oral administration in unit dose form that comprises a sulfur-containing antioxidant selected from a sulfate, a free thiol, and the group consisting of sodium sulfite, sodium metabisulfite, N-acetylcysteine, homocysteine, cysteine, monothioglycerol, sodium thiosulfate, α -lipoic acid, and dithiothreitol. In various embodiments, the pharmaceutical formulation optionally comprises a chelating agent. In one embodiment, the chelating agent is EDTA. In other embodiments, the chelating agent is citric acid.

[0018] In one embodiment, the barley endoprotease EPB2 is admixed with a prolyl endopeptidase (PEP). In one embodiment, the PEP is *Sphingomonas capsulata* PEP or a modified form of that protease, or a recombinant version of either. In one embodiment, the PEP is *Aspergillus niger* PEP or a modified form of that protease, or a recombinant version of either.

[0019] The present invention is useful in a method for preventing and/or treating Celiac sprue and/or dermatitis herpetiformis in a patient, said method comprising the step of orally administering a stabilized cysteine protease

containing pharmaceutical formulation of the invention to said patient, contemporaneously with the patient's ingestion of a gluten-containing foodstuff.

[0020] In a further aspect, the present invention provides unit dose forms of the novel compositions of the invention as defined in the claims. In various embodiments, the unit dose form can conveniently be used in the method of the invention. The dosage forms and pharmaceutical formulations of the invention can be used for the treatment of mammals as well as humans.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021]

Figure 1 shows the activity of ALV001* (the active form of the proenzyme ALV001) during incubation with postprandial gastric fluid.

Figure 2 shows the activity of ALV001 during incubation with postprandial gastric fluid.

Figure 3 is a western blot of ALV001 showing stabilization of ALV001 in postprandial gastric fluid with 1-thioglycerol.

Figure 4 shows the activity of ALV003 during incubation with postprandial gastric fluid in the absence or presence of 1-thioglycerol.

Figure 5 shows the reversal of loss of enzyme activity with antioxidant.

Figure 6 shows the stabilization of ALV001 in postprandial gastric fluid with reducing agents.

Figure 7 shows stabilization as in Figure 6, with sodium thiosulfate at 0.5 mM, sodium metabisulfite at 0.5 mM and lipoic acid at 5 mM.

Figure 8 shows a time course for stabilization with sodium sulfite.

Figure 9 shows a titration of sodium metabisulfite.

Figure 10 shows a restoration of activity after the addition of reducing agents.

Figure 11 shows a dose dependence of sodium metabisulfite in stabilization of ALV001.

Figure 12 shows a metabisulfite titration at 2X ALV001 enzyme concentration.

Figure 13 shows that sodium metabisulfite stabilizes ALV001 activity in orange juice. 3 mg/ml ALV001 was reconstituted in orange juice with or without 840 μ M sodium metabisulfite. Chromogenic enzyme activity was measured at the indicated timepoints to determine the relative amount of functional enzyme. Figure 14 shows that cysteine stabilizes ALV001 activity under simulated postprandial gastric conditions. A gluten-free meal was chewed and spit into resting gastric fluid. ALV003 and varying levels of cysteine were added at time 0 and the stability of ALV001 activity was tested by monitoring chromogenic enzyme activity over time. Gastric acid secretion was simulated throughout the experiment.

Figure 15 shows that ALV001 stability study. 150 mg of enzyme was formulated with 300 mg. citric acid

monohydrate, and 8 mg. sodium bisulfite, placed in polypropylene bottle, and sealed in foil pouches. A chromogenic assay was measured at specified time points to assess stability.

DETAILED DESCRIPTION OF THE INVENTION

[0022] In the current art of oral delivery (the administration of a substance by mouth, i.e., oral administration) of proteins, stabilization in the GI tract to prevent protein oxidation has not been discussed; instead, the focus of past efforts has been on the potential instability of the orally ingested protein to acid or to proteolysis by digestive enzymes. Thus, the need to provide additional stabilization of orally administered proteins to oxidation has not been appreciated prior to the present invention. Moreover, the benefits of such stabilization can be particularly beneficial, particularly for the oral delivery of enzymes to the stomach for the purpose of enhancing proteolysis. The present invention arose at least in part from the discovery that oxidation can play a key role in degradation of such orally administered enzymes.

[0023] The enzyme ALV001, a recombinant, modified form of the cysteine protease known as barley EPB2, was rapidly inactivated in *ex vivo* extracts of fed human gastric fluid and at rates that were significantly faster than observed at similar pH (3 to 5) and pepsin content. One aspect of the present invention arises from the discovery that the addition of antioxidants, particularly sulfur-containing antioxidants, such as sulfites or free thiols or agents that generate free thiols in gastric fluid, stabilize ALV001 (and thus, other cysteine proteases and cysteine-containing proteins) in gastric fluid. Surprisingly, these sulfur-containing antioxidants can also regenerate active enzyme from inactivated ALV001 in gastric fluid. Other antioxidants, such as ascorbic acid and methionine (in which the sulfhydryl group is methylated) did not stabilize (preserve the activity of) ALV001. EDTA did help stabilize ALV001 in *ex vivo* gastric fluid, albeit to a lesser extent than the antioxidants, and chelating agents generally therefore provide some stabilization, alone or in combination with an antioxidant, in accordance with the invention. Among the antioxidants, sodium sulfite and sodium metabisulfite are particularly effective in stabilizing ALV001.

[0024] As used herein, an "antioxidant" is a molecule capable of slowing or preventing the oxidation of other molecules. Oxidation is a chemical reaction that transfers electrons from a substance to an oxidizing agent. Antioxidants include, without limitation, reducing agents, ascorbic acid (vitamin C), lipoic acid, melatonin, uric acid, carotenes, retinols, tocopherols and tocotrienols, e.g. α -tocopherol (vitamin E), ubiquinone (coenzyme Q), and the like.

[0025] As used herein a "reducing agent" is any compound in the subclass of physiologically acceptable antioxidant compounds that reduces another species in a redox (reduction-oxidation) reaction and, in so doing, be-

come oxidized and so serves as an electron donor in the redox reaction. Reducing agents include, without limitation, methionine, glutathiol, dithiothreitol, dithioerythritol, β -mercaptoethanol, sodium metabisulfite, thioglycolate, cysteine, N-acetylcysteine, homocysteine, monothioglycerol, sodium sulfite, potassium bisulfite, sodium thiosulfite, and the like. Some such agents comprise a free thiol (*i.e.* mercaptan) group, *e.g.* monothioglycerol, sodium thiosulfite, sodium metabisulfite, and the like. Agents that are in an oxidized form but that comprise a free thiol in acidic include, without limitation, α -lipoic acid, dithiothreitol, glutathione, and the like.

[0026] For certain antioxidants, the levels required to act in a one liter volume of the stomach would require an amount of antioxidant in the dosage form (or to be taken accompanying the dosage form) that are larger than currently accepted for toxicological reasons in the FDA's Inactive Ingredients Guide. Accordingly, their use in the compositions of the invention would likely require additional clinical testing of the antioxidant itself. In contrast, the sulfites and metabisulfites, especially their sodium and potassium salts and combinations, with their long history as preservatives in the food industry and their unusual potency as reducing agents for cysteine proteases as described herein are effective and accepted pharmaceutical excipients.

[0027] In some embodiments of the invention, the antioxidant is selected from sodium sulfite, sodium bisulfite, potassium bisulfate, potassium metabisulfite, sodium thiosulfate, glutathione, cysteine, homocysteine, sodium dithionite, thioglycerol, and acetylcysteine, alone or in combination. In other embodiments, a sulfur containing antioxidant is selected from the group consisting of sodium sulfite, sodium metabisulfite, N-acetylcysteine, homocysteine, cysteine, monothioglycerol, sodium thiosulfate, α -lipoic acid, and dithiothreitol. In some embodiments, the antioxidant is sodium metabisulfite.

[0028] The dosage of an antioxidant for purposes of the present invention may be empirically determined by assays known in the art, *e.g.* by testing a range of doses for stabilizing effect on a known concentration of protein. It will be understood by one of skill in the art that the dosage range will depend on the specific antioxidant. In some embodiments, the dosage of antioxidant is at least about 2 mg per dosage form; at least about 5 mg per dosage form; at least about 10 mg per dosage form, at least about 50 mg per dosage form; and not more than about 500 mg per dosage form, usually not more than about 300 mg per dosage form.

[0029] In some embodiments, the antioxidant is sodium metabisulfite, which may be present at a concentration of from about 2 - 50 mg per dosage form, preferably at from about 4 - 16 mg per dosage form, and most preferably at from about 6 - 10 mg per dosage form. In other embodiments, the antioxidant is cysteine, which may be present at a concentration of from about 5 - 400 mg per dosage form, preferably at from about 10 - 350 per dosage form, and most preferably at from about 50 - 300 mg

per dosage form.

[0030] In various embodiments, the pharmaceutical formulation optionally comprises a chelating agent. The chelating agent is a physiologically acceptable component. Examples of the chelating agent include, without limitation, ethylenediaminetetraacetic acid (EDTA), sodium ethylenediaminetetraacetate, EGTA, phytic acid, citric acid, *e.g.* anhydrous citric acid, and the like. These can be used alone or in admixture. In one embodiment, the chelating agent is EDTA.

[0031] The dosage of a chelating agent may be empirically determined by assays known in the art, *e.g.* by testing a range of doses for stabilizing effect on a known concentration of protein, usually in combination with a selected antioxidant. It will be understood by one of skill in the art that the dosage range will depend on the specific chelating agent and protein to be administered. In some embodiments, the dosage is at least about 5 mg per dosage form; at least about 10 mg per dosage form; at least about 50 mg per dosage form, at least about 100 mg per dosage form; and not more than about 500 mg per dosage form, usually not more than about 300 mg per dosage form.

[0032] In some embodiments the chelating agent is citric acid, *e.g.* anhydrous citric acid, citric acid monohydrate, which may be present at a dose of from about 10 to about 500 mg per dosage form, usually at a dose of from about 25 to 350 mg per dosage form, and may be provided at a dose of from about 45 - 275 mg per dosage form.

[0033] In the formulations of the present invention, an antioxidant, *e.g.* a reducing agent as described above, is provided at a concentration effective to stabilize the protease active agent when exposed to gastric fluid, particularly when exposed to postprandial gastric fluid. The formulations are intended for *in vivo* use. However, for convenience, the stability may be tested *in vitro*, *e.g.* by combining the protease with postprandial gastric fluid and assessing proteolytic activity over a suitable period of time, *e.g.* for about 10 minutes, about 20 minutes, about 30 minutes, about 1 hour, or more. An effective concentration is sufficient to maintain at least about 25% of the enzyme starting activity, at least about 50% of the enzyme starting activity, at least about 75% of the enzyme starting activity, or at least about 90% of the enzyme starting activity, or is sufficient substantially to maintain the original activity for at least about 10 minutes, at least about 30 minutes, or longer.

[0034] Thus, the present invention provides novel formulations of proteins suitable for oral administration, which proteins are more resistant to oxidative degradation than currently available compositions. Those of skill in the art will appreciate that this discovery may enhance the therapeutic potential of a wide variety of proteins for oral administration, as prior testing or properties of a protein may have indicated that the protein would not provide therapeutic benefit on oral administration. For example, while papain is marketed as a nutraceutical, it has not

found use as a therapeutic. Despite its high optimal temperature, papain may provide therapeutic benefit if administered in a formulation of the invention.

[0035] Those of skill in the art will therefore appreciate that the present invention provides pharmaceutical formulations of a protein as defined in the claims suitable for oral administration that stabilize the protein in the fed stomach. In one embodiment, the formulation comprises, in addition to the protein and sulfur-containing antioxidant, a chelating agent.

[0036] Cysteine proteases have been reported to be useful for a variety of therapeutic purposes. For example, U.S. Patent No. 6,241,973, describes cysteine proteases useful in teeth whitening. PCT Publication No. WO 2004/058816, describes cysteine proteases useful in vaccines against liver fluke. U.S. Patent App. Pub. No. 20080039400 describes cysteine proteases useful in treating inflammatory bowel disease and other conditions. U.S. Patent App. Pub. No. 20070264311, describes cysteine proteases useful in treating food intolerance. U.S. Patent App. Pub. No. 20070148267 describes cysteine proteases useful in regulating food intake. U.S. Patent App. Pub. Nos. 20060134017 and 20060134018 describe cysteine proteases useful in the prevention of bacterial adhesion to biological surfaces, i.e., for prevention of plaque formation. Cysteine proteases have also been described for systemic treatments, including, for example, induction of apoptosis in cancers. The compositions of the invention are accordingly useful in such methods.

[0037] The barley endoprotease EPB2 or a recombinant form thereof is a protease capable of digesting a gluten peptide that is toxic to a Celiac sprue patient into non-toxic fragments, either alone or in combination with another protease.

[0038] As used herein, a "pharmaceutical formulation", or "dosage form" includes any formulation intended for administration to humans or mammals and so includes not only formulations that undergo clinical testing and approval by regulatory authorities, such as the FDA, but also formulations that do not require regulatory approval, such as nutraceuticals.

[0039] Thus, for example, the compositions of the invention, including pharmaceutical formulations and dosage forms, can also include, depending on the formulation desired, pharmaceutically-acceptable, non-toxic carriers or diluents, which are defined as vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. The diluent is selected so as not to adversely affect the biological activity of the protease. Examples of such diluents are distilled water, buffered water, physiological saline, PBS, Ringer's solution, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation can include other carriers, adjuvants, or non-toxic, nontherapeutic, nonimmunogenic stabilizers, excipients and the like. The compositions can also include additional substances to approximate physiological conditions, such

as pH adjusting and buffering agents, toxicity adjusting agents, wetting agents and detergents.

[0040] The compositions of the invention can also include any of a variety of stabilizing agents, such as an antioxidant for example. The proteins or polypeptides in a composition of the invention can also be complexed with molecules that enhance their *in vivo* attributes. Such molecules include, for example, carbohydrates, polyamines, amino acids, other peptides, ions (e.g., sodium, potassium, calcium, magnesium, manganese), and lipids.

[0041] Further guidance regarding formulations that are suitable for various types of administration can be found in Remington's Pharmaceutical Sciences, Mace Publishing Company, Philadelphia, PA, 17th ed. (1985). For a brief review of methods for drug delivery, see, Langer, Science 249:1527-1533 (1990).

[0042] For oral administration, the active ingredient can be administered in solid dosage forms, such as capsules, tablets, and powders, such as in sachets or sprinkles, or in liquid dosage forms, such as elixirs, syrups, and suspensions. The active component(s) can be encapsulated in gelatin or hypromellose capsules together with inactive ingredients and powdered carriers, such as glucose, lactose, sucrose, mannitol, starch, cellulose or cellulose derivatives, magnesium stearate, stearic acid, sodium saccharin, talcum, magnesium carbonate. Examples of additional inactive ingredients that may be added to provide desirable color, taste, stability, buffering capacity, dispersion or other known desirable features are red iron oxide, silica gel, sodium lauryl sulfate, titanium dioxide, and edible white ink. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric-coated for selective disintegration in the gastrointestinal tract. Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

[0043] The components used to formulate the pharmaceutical compositions of the invention are preferably of high purity and are substantially free of potentially harmful contaminants (e.g., typically at least National Food (NF) grade, generally at least analytical grade, and more typically at least pharmaceutical grade). To the extent that a given compound must be synthesized prior to use, the resulting product is typically substantially free of any potentially toxic agents, particularly any endotoxins, which may be present during the synthesis or purification process.

[0044] Nutraceutical formulations are defined as "a food or part of a food that offers medical and/or health benefits including prevention or treatment of disease." (Dr. Stephen DeFelice, director of Foundation for Innovation In Medicine). Products range from isolated nutrients, dietary supplements and diets, to genetically engi-

neered designer foods, functional foods, herbal products and processed foods such as cereal, soup and beverages. Functional foods, the most popular term among consumers but not a clearly delineated product category, are defined by Clare Hasler, Ph.D., of the University of Illinois, as foods that include "any modified food or food ingredients that may provide a health benefit beyond the traditional nutrients it contains." Nutraceutical formulations of interest include foods for veterinary or human use, such as health food bars, drinks and drink supplements, and the like. These foods are enhanced by the inclusion of a biologically active protease in a composition as provided herein.

[0045] When the dosage form or dosage forms are for treating celiac sprue and contain the enzyme ALV001, at least one dosage form may contain ALV001 alone or in combination with ALV002. The total content of ALV001 administered with a meal when ALV001 is administered without ALV002 is from 50 to 2000 mg of ALV001 protein, preferably between 100 and 1000 mg, and still more preferably between 100 and 600 mg of ALV001. When ALV001 and ALV002 are administered in a combination either in the same or separate dosage forms, they may be given in a ratio of 10:1 to 1:10, more preferably 3:1 to 1:3, and most preferably at a ratio of 1.5:1 to 1: 1.5. If administered at approximately a 1:1 ratio, the total enzyme content given with a meal may be 50 to 2000 mg, more preferably 50 to 900 mg, and most preferably 100 to 400 mg.

[0046] The proteolytic activity of ALV001 at 25°C against the chromogenic substrate Z-Phe-Arg-pNA is assessed by measuring the rate of change in the absorbance of light at 410 nm. The method is based on the release of para-nitroaniline (pNA, 4-nitroaniline) into solution, which absorbs light at 410 nm. The specification is ≥ 5000 u/mg for ALV001.

[0047] The proteolytic activity of ALV002 at 25°C against the chromogenic substrate Z-Gly-Pro-pNA is assessed by measuring the rate of change in the absorbance of light at 410 nm. The method is based on the release of para-nitroaniline (pNA, 4-nitroaniline) into solution, which absorbs light at 410 nm. The specification is ≥ 3000 u/mg for ALV002.

[0048] The present invention is useful in a method for preventing and/or treating Celiac sprue and/or dermatitis herpetiformis in a patient, said method comprising the step of orally administering a stabilized cysteine protease containing pharmaceutical formulation of the invention, wherein the formulation is stabilized with an antioxidant, and wherein the protease is capable of degrading gluten into fragments non-toxic to Celiac sprue patients upon oral administration, to said patient contemporaneously with the patient's ingestion of a gluten-containing foodstuff. U.S. Patent Nos. 7,320,788 and 7,303,871 describe methods for identifying proteases capable of degrading gluten into fragments non-toxic to Celiac sprue patients.

[0049] In a further aspect, the present invention provides pharmaceutical formulations and dosage forms,

i.e., unit dose forms, of the novel compositions described herein. In various embodiments, the unit dose form can conveniently be used in the method of the present disclosure.

EXAMPLES

1. Demonstration of Instability of ALV001 in Postprandial Human Gastric Fluid

[0050] Frozen human postprandial gastric fluids were thawed and centrifuged at 21,000g for 5 minutes to pellet insoluble material, and the supernatant was used for this example. Between 50 and 200 μ l of gastric fluid were added to the wells of a 96-well plate and incubated at 37 °C. When antioxidants were added, they were added at this step. ALV001 or activated ALV001* (see PCT patent application No. 2008/003425

regarding how ALV001* (the active form of ALV001) is produced upon proteolytic cleavage of ALV001) was dissolved at a ratio of 20 μ l water/mg powder, the concentration was measured by the absorbance at 280 nm, and the protein was added to the gastric fluid at a concentration of 0.1 to 0.3 mg/ml. At the specified sampling time, a 1.6 μ l sample of gastric fluid was diluted in assay buffer to give a final volume of 280 μ l, and the concentration was measured by a chromogenic activity assay using Z-Phe-Arg-p-nitroaniline (pNA) as the substrate. The method is based on the ability of pNA to absorb at 410 nm when released from the substrate. ALV001 is the proenzyme or zymogen for Endoprotease B (Isoform 2) that cleaves the C-terminal side of the arginine in the sequence Phe-Arg. Enzymatic cleavage and subsequent release of pNA may be monitored by the change in absorbance at 410 nm. To determine the reaction rate, the absorbance at 410 nm was measured every 10 seconds for 1 minute. The slope of the regression line divided by the enzyme concentration was the reported enzyme activity.

[0051] Stability of ALV001* in human gastric fluid was analyzed for 4 healthy volunteers fasting and at 15, 30, 45, 60, 75, and 90 minutes after eating a meal. The gastric fluid was withdrawn from these subjects through a nasogastric tube. The enzymatic activity of ALV001 was measured after 1.5, 5, 10, 20, and 60 minutes after addition to the gastric fluid. Western blots were also performed to investigate proteolytic degradation of ALV001* in the gastric fluid.

[0052] Figure 1 shows that ALV001* activity was only detectable in the 15 and 30 minute postprandial gastric fluid of Subject 4 and even then disappeared after 20 minutes of incubation. Similar results were observed for the other 3 subjects and with the addition of the zymogen, ALV001. In the fasting gastric fluid, rapid enzymatic degradation of ALV001* was expected due to low pH, for example, pH 1.3 for subject 4, and pepsin content. Postprandial gastric fluid has a transient increase in pH to the range of 3.5 or 4 or even greater, and at this greater pH,

pepsin activity is reduced, and ALV001* is expected to be more resistant to proteolytic digestion and to degradation by acidity.

[0053] As shown in Figure 2, for subject 1 in the 15 minute postprandial sample at pH 4.0, the enzyme ALV001, was shown by the chromogenic assay to be rapidly inactivated, but the enzyme was still intact as measured by the Western blot, shown in Figure 3. In the same subject, in the 30 and 60 minute postprandial gastric samples, which were pH 2.5 and 1.2, respectively, the enzyme was both inactive and proteolyzed. Similar data were obtained with the other patient samples. The conclusion was that an unexpected loss of enzymatic activity of this cysteine protease ALV001 in early postprandial gastric fluid samples resulted from a mechanism that was not related to proteolysis.

2. Demonstration of Stabilization of ALV001 in Postprandial Gastric Fluid by 1-Thioglycerol (monothioglycerol)

[0054] Figure 4 shows the results of adding a range of concentrations of monothioglycerol (MTG), from 0.01 to 50 mM, to the wells of 15 minute postprandial gastric fluid as described above. A control without MTG was also included. The rate of loss of enzymatic activity of ALV001 decreased with increasing MTG concentrations with no loss at 50 mM MTG. Thus, the antioxidant MTG prevented the loss of enzyme activity of this cysteine protease.

[0055] Moreover, as shown in Figure 5, pre-incubation of ALV003, a combination of ALV001 and ALV002 (a recombinant form of *Sphingomonas capsulata* prolyl endopeptidase) at approximately 1:1 enzyme ratio, in 15 minute postprandial gastric fluid followed by addition of 50 mM MTG versus no addition showed that the antioxidant MTG could reverse the loss of enzymatic activity of ALV001*. After approximately 20 minutes addition of MTG to nonfunctional pre-incubated ALV001*, nearly full enzymatic activity was restored.

3. Demonstration of Stabilization of ALV001 in Postprandial Gastric Fluid by Selected Reducing Agents

[0056] Numerous reducing agents, in particular, N-acetylcysteine, L-cysteine, homocysteine, 2-mercaptoethanol, dithiothreitol, tris(2-Carboxyethyl) phosphine hydrochloride (TCEP), lipoic acid, sodium thiosulfate, sodium sulfite, and sodium metabisulfite, were all demonstrated to stabilize ALV001* in early postprandial gastric fluid. In contrast, L-methionine and ascorbic acid did not stabilize the enzyme. EDTA enhanced the stability of ALV001*, suggesting a dependency on metals for the inactivation.

[0057] Shown in Figure 6, selected reducing agents were added at 5 mM to ex vivo postprandial gastric fluid that was obtained from a healthy volunteer 15 minutes after eating and was incubated at 37 °C. The proenzyme ALV001 was added to the gastric fluid. Acetate buffer at pH 4.5 was used as a control, because ALV001 is fully

activated under these conditions. Samples were taken to determine ALV001 activity after 1.5, 5, 10, 20, 45 minutes.

[0058] As can be seen from the figure, the cysteine protease ALV001 was fully active in acetate buffer. In contrast, the enzyme is rapidly inactivated in water, gluten, and even with ascorbic acid, an antioxidant, added. There is evidence of stabilization with the chelating agent, EDTA. Substantial improvements in stability were observed with the reducing agents, cysteine and N-acetylcysteine, and the most effective stabilization was demonstrated with the reducing agent sodium sulfite.

[0059] Shown in Figure 7, in a second set of tests performed in the same manner, sodium thiosulfate at 0.5 mM, sodium metabisulfite at 0.5 mM, and lipoic acid at 5 mM were demonstrated to improve the stability of ALV001 in postprandial gastric fluid.

[0060] Shown in Figure 8, sodium sulfite was a particularly effective reducing agent to stabilize the ALV001, with the effective persisting down to 0.05 mM. Sodium metabisulfite improved stability of ALV001 even at concentrations of 50 μM.

[0061] Shown in Figure 9, in another set of tests, the ability of these reducing agents and a chelating agent to reverse the degradation of ALV001 in gastric fluid was examined. ALV001 was added to postprandial gastric fluid and incubated for 15 minutes at 37 °C. Reducing agents were added at 5 mM to the gastric fluid and then samples were taken to examine ALV001 activity at 1.5, 5, 10, 20, and 40 minutes after the addition of the reducing agents. The buffer control and EDTA did not restore the enzyme activity, and sodium sulfite and acetylcysteine partially restored the enzyme activity.

[0062] Shown in Figure 11, the dose dependence of sodium metabisulfite and improved stability of ALV001 at 0.12 mg/ml was measured in postprandial gastric fluid. Samples were taken at 1.5, 5, 10, 20, and 40 minutes. While sodium metabisulfite provided considerable stabilization even at 25 μM, the best stabilization was observed at 50 μM. This test was repeated at twice the enzyme concentration to ensure that the effect was not dependent on enzyme concentration, Figure 12.

4. Dosage Form Stability Testing

[0063] 65 mg of ALV003, a mixture consisting of 38 mg ALV001 (Lot A031207) and 27 mg of ALV002 (Lot B061707) were weighed into each hypromellose capsules size 0 (Qualicaps). For one set of capsules 0.8 mg per capsule sodium metabisulfite (Spectrum, Lot WG 0855) was first added to the ALV003 powder and it was mixed by hand with a spatula before placing in the capsules. The hypromellose capsules were placed in 30 ml polypropylene vials with 1g desiccant canisters (Sorb-it). The vials were transferred into 25°C/60%RH and 40 °C/75% RH stability chambers. Samples were withdrawn at Initial (T=0), 2, 4, and 8 week intervals for enzyme activity analysis.

5. Sodium Metabisulfite in Grape Juice as Antioxidant

[0064] The compositions of the invention include food-stuffs modified to contain a therapeutic protease and an anti-oxidant, reducing agent, and/or chelating agent. This example illustrates the use of the reducing agent sodium metabisulfite in combination with the cysteine protease ALV001 taken or admixed with grape and other juices. To demonstrate the ability to stabilize ALV001 in gastric fluid by grape juice, as an example, the following experiment was performed. ALV001 was prepared in water at 20 μ l/mg powder or approximately 9.5 mg/ml. Into a 96 well plate was added 54 μ l 15 minute postprandial gastric fluid pre-incubated to 37 °C, 5 μ l of juice and 2.66 μ l of ALV001 diluted 1:1 with water. The juices tested were sparkling white grape juice (Welch's), white grape juice (Welch's), white grape juice (Safeway), lime juice (Real-ime), and lemon juice (Safeway). These fruit juices contain metabisulfite as a preservative. Water was used as a control. The water control showed the usual rapid degradation of ALV001 in postprandial gastric fluid. The lime and lemon juice showed slower degradation of ALV001, and all 3 grape juices were stable over the 20 minute time course of the test.

[0065] Thus, in one embodiment of the invention, ALV003 or another glutenase is administered (for example, 300 mg of drug) in an HPMC capsule and is taken with a glass, approximately 100 to 250 ml or more, of grape juice.

6. Sodium Metabisulfite in Orange Juice as Antioxidant

[0066] 3 mg/ml ALV001 was reconstituted in orange juice with or without 840 μ M sodium metabisulfite. Chromogenic enzyme activity was measured as described above, at the indicated timepoints, to determine the relative amount of functional enzyme, as shown in Figure 13. The data show increased stability in the presence of sodium metabisulfite.

7. Cysteine as Antioxidant

[0067] A gluten-free meal was chewed and spit into resting gastric fluid. ALV003 and varying levels of cysteine were added at time 0, and the stability of ALV001 activity was measured by monitoring chromogenic enzyme activity over time. Gastric acid secretion was simulated throughout the experiment. As shown in Figure 14, cysteine stabilizes ALV001 activity under these simulated postprandial gastric conditions.

8. Citric Acid Monohydrate and Sodium Bisulfite as Antioxidant

[0068] 150 mg of enzyme (ALV001) was formulated with 300 mg. citric acid monohydrate, and 8 mg. sodium metabisulfite, placed in polypropylene bottle, and sealed in foil pouches. A chromogenic assay was measured at

specified time points to assess stability, as shown in Figure 15.

9. Dosage Forms

[0069] The present invention provides a dosage form that contains 300 mg or 900 mg ALV003, optionally in a tablet form, with added mannitol and microcrystalline cellulose and 8 mg sodium metabisulfite.

[0070] The present invention provides a dosage form that contains a lyophilized powder of ALV001 in which sodium metabisulfite had been added to the solution prior to lyophilization instead of monothioglycerol at a ratio from 100:8 (w/w) to a ratio of 900:4 (w/w).

[0071] The present invention provides a dosage form that contains a spray dried powder of ALV001 in which sodium metabisulfite had been added to the solution prior to spray drying instead of monothioglycerol at a ratio from 100:8 (w/w) to a ratio of 900:4 (w/w).

[0072] The present invention provides a dosage form that contains immediate release ALV001 in a range of 100 to 900 mg and sodium metabisulfite in a range of 1 to 10 mg.

[0073] The present invention provides a dosage form that contains controlled release ALV001 in a range of 100 to 900 mg and sodium metabisulfite in a range of 1 to 10 mg released from 30 minutes to 6 hours.

[0074] The present invention provides a dosage form that contains controlled release ALV001 in a range of 100 to 900 mg and sodium metabisulfite in a range of 1 to 10 mg released from 30 minutes to 6 hours.

[0075] The present invention provides a dosage form that contains pulsed release ALV001 in a range of 100 to 900 mg and sodium metabisulfite in a range of 1 to 10 mg in an immediate release pulse and a second pulse released from 20 minutes to 6 hours later.

[0076] The present disclosure provides a dosage form that contains a protease and a quantity of an antioxidant that achieves an antioxidant concentration of at least 30, 50, 100, or 200 μ M in the stomach.

[0077] The present disclosure provides a dosage form that contains a protease and a quantity of a compound that generates a concentration of free thiol of at least 100, 200, or 500 μ M in the stomach.

[0078] The present invention provides a dosage form that contains sodium metabisulfite and may contain both ALV001 and ALV002 where one enzyme component is formulated to provide an immediate release and the other enzyme component is either a pulsed release or to provide a controlled release.

[0079] All of the ALV001 dosage forms above can also be modified to include a second protease. In one embodiment, the second protease is a prolyl endopeptidase (PEP). In one embodiment the PEP is *Sphingomonas capsulata* PEP, for example and without limitation, as described in PCT Pub. No. 2008/115411. In various embodiments, between 1-2000 mg of the PEP and ALV001 are dosed at a PEP:ALV001 weight ratio of between

1:100 to 100:1, more preferably between 1:20 to 20:1, more preferably between 1:5 and 5:1, and most preferably at a 1:1 ratio. In one embodiment, the dosage form is constructed so that the ALV001 and antioxidant, e.g., sodium metabisulfite, are immediately released, and the PEP is released either immediately; or is released in one or more short, delayed pulses (from 10 minutes to 3 hours); or is released in sustained release over 10 minutes to 3 hours.

[0080] In one embodiment, the present disclosure provides a dosage form in which the antioxidant is an antioxidant other than sodium metabisulfite.

[0081] The present disclosure provides oral dosage forms comprising a protein and an antioxidant selected from the group consisting of sodium sulfite, sodium bisulfite, potassium bisulfate, potassium metabisulfite, alone or combination; sodium thiosulfate; glutathione, cysteine, homocysteine, sodium dithionite, thioglycerol; and acetylcysteine.

[0082] The present disclosure may provide roller compacted granules or pellets of the enzyme. The antioxidant may be either contained in the granules or pellets or blended with the granules or pellets and either filled or compressed into a dosage form.

[0083] The pharmaceutical formulations of the invention can be in the form of, for example and without limitation, particles, particles in capsule or sachet, or tablet. Tablets may be single layer, bilayer, or multilayer and may be coated or uncoated. The formulation can, for example and without limitation, be added to a food or drink and then administered, for example, as a sprinkled powder or granule formulation or as a spread in the form of a jam or powder. A capsule of low water content may be desired for stability, and hypromellose capsules, HPMC, of size 1, 0, or 00, can be used. Capsules can be packaged in a dry environment either with desiccant or desiccant packs or if in blisters under dry nitrogen or other dry environment.

[0084] The pharmaceutical formulations of the invention can comprise a lubricant such as magnesium stearate, stearic acid, sodium stearyl fumarate, or sodium stearyl lactylate, hydrogenated vegetable oil (such as hydrogenated and refined triglycerides of stearic and palmitic acids). These may be at 0.3 to 5% of weight of the dosage form. If mannitol is contained at a high concentration in the lyophilized powder, then higher concentrations of lubricant may be used.

[0085] The cysteine protease (or other protein active pharmaceutical ingredient) powder may be blended with lubricant or other excipients such as a filler or binder and granulated. If the cysteine protease is unstable with water and temperature, then these can be roller compacted into granules, if necessary using chilled rollers for stability. One may optionally include an agent that modifies or controls pH, at least for the first few minutes after the dosage form is in the GI tract, to facilitate activation of zymogen proteins such as ALV001.

[0086] Fillers such as dicalcium phosphate, microcrystalline cellulose, maltodextrins, mannitol, lactose, sucrose, or trehalose may be included and blended with the powders or included in the lyophilized powder or spray-dried powder. More hydrophilic fillers such as microcrystalline cellulose may be avoided for certain enzymes, such as ALV001.

talline cellulose, maltodextrins, mannitol, lactose, sucrose, or trehalose may be included and blended with the powders or included in the lyophilized powder or spray-dried powder. More hydrophilic fillers such as microcrystalline cellulose may be avoided for certain enzymes, such as ALV001.

[0087] Controlled-release excipients may be blended in to form polymeric drug-containing matrices. These matrices may be from about 1 mm in diameter to the size of a full tablet 10 to 12 mm in width and even 1.8 cm or more in length. These matrices can provide extended-release into the stomach being retained with food for ½ to 8 hours depending on the size. These matrices may or may not be swellable. If swellable, extended-release hydrophilic polymers that are appropriate include cellulose polymers and their derivatives (such as for example, hydroxyethylcellulose, hydroxypropylcellulose, carboxymethylcellulose, and microcrystalline cellulose, polysaccharides and their derivatives, polyalkylene oxides, polyethylene glycols, chitosan, poly(vinyl alcohol), xanthan gum, maleic anhydride copolymers, poly(vinyl pyrrolidone), starch and starch-based polymers, poly(2-ethyl-2-oxazoline), poly(ethyleneimine), polyurethane hydrogels, and crosslinked polyacrylic acids and their derivatives. Further examples are copolymers of the polymers listed in the preceding sentence, including block copolymers and grafted polymers. Extended-release coatings could also be prepared on these particles using some of the above polymers.

Claims

1. A solid pharmaceutical formulation of a barley endoproteinase B2 (EPB2) or a recombinant form thereof suitable for oral administration in unit dose form that comprises a sulfur-containing antioxidant selected from a sulfate, a free thiol, and the group consisting of sodium sulfite, sodium metabisulfite, N-acetylcysteine, homocysteine, cysteine, monothioglycerol, sodium thiosulfate, α -lipoic acid, and dithiothreitol.
2. The pharmaceutical formulation of claim 1, wherein said anti-oxidant is present in a dosage of at least about 2 mg per dosage form to not more than about 500 mg per dosage form.
3. The pharmaceutical formulation of either of claims 1 or 2, wherein said sulfur-containing antioxidant is sodium metabisulfite, which is present at about 2 - 50 mg per dosage form.
4. The pharmaceutical formulation of either of claims 1 or 2, wherein said sulfur-containing antioxidant is cysteine, which is present at about 5 - 400 mg per dosage form.

5. The pharmaceutical formulation of any one of the preceding claims that comprises a chelator.
6. The pharmaceutical formulation of claim 5, wherein the chelator is EDTA or citric acid.
7. The pharmaceutical formulation of claim 6, wherein said chelator is present in a dosage of at least about 5 mg per dosage form to not more than about 500 mg/dosage form.
8. The pharmaceutical formulation of claim 7, wherein said chelator is citric acid, which is present at 10 to 500 mg per dosage form.
9. The pharmaceutical formulation of any one of the preceding claims wherein the unit dose further comprises a prolyl endopeptidase.
10. The pharmaceutical formulation of claim 9 wherein said prolyl endopeptidase is a *Sphingomonas capsulata* PEP or a recombinant form thereof.
11. The pharmaceutical formulation of any one of the preceding claims, wherein the formulation is a capsule, tablet, powder, sachet or sprinkles.
12. The formulation of any one of claims 1-11 for use in a method for preventing and/or treating Celiac sprue and/or dermatitis herpetiformis in a patient, said method comprising the step of orally administering said formulation to said patient contemporaneously with the patient's ingestion of a gluten-containing foodstuff.

Patentansprüche

1. Feste pharmazeutische Formulierung eines Gersten-Endoproteinase-B2 (EPB2) oder einer rekombinanten Form davon, geeignet für eine orale Verabreichung in Einheitsdosierform, die ein schwefelhaltiges Antioxidans umfasst, das aus einem Sulfat, einem ungebundenen Thiol und der Gruppe bestehend aus Natriumsulfit, Natriummetabisulfit, N-Acetylcystein, Homocystein, Cystein, Monothioglycerin, Natriumthiosulfat, α -Liponsäure und Dithiothreitol ausgewählt ist.
2. Pharmazeutische Formulierung nach Anspruch 1, worin das Antioxidans in einer Dosierung von zumindest etwa 2 mg pro Dosierform bis nicht mehr als etwa 500 mg pro Dosierform enthalten ist.
3. Pharmazeutische Formulierung nach Anspruch 1 oder 2, worin das schwefelhaltige Antioxidans Natriummetabisulfit ist, das in etwa 2-50 mg pro Dosierform enthalten ist.

4. Pharmazeutische Formulierung nach Anspruch 1 oder 2, worin das schwefelhaltige Antioxidans Cystein ist, das in etwa 5-400 mg pro Dosierform enthalten ist.
5. Pharmazeutische Formulierung nach einem der vorangegangenen Ansprüche, die einen Chelatbildner umfasst.
6. Pharmazeutische Formulierung nach Anspruch 5, worin der Chelatbildner EDTA oder Citronensäure ist.
7. Pharmazeutische Formulierung nach Anspruch 6, worin der Chelatbildner in einer Dosierung von zumindest etwa 5 mg pro Dosierform bis nicht mehr als etwa 500 mg/Dosierform enthalten ist.
8. Pharmazeutische Formulierung nach Anspruch 7, worin der Chelatbildner Citronensäure ist, die bei 10 bis 500 mg pro Dosierform vorhanden ist.
9. Pharmazeutische Formulierung nach einem der vorangegangenen Ansprüche, worin die Einheitsdosis ferner eine Prolylendopeptidase umfasst.
10. Pharmazeutische Formulierung nach Anspruch 9, worin die Prolylendopeptidase eine *Sphingomonas capsulata*-PEP oder eine rekombinante Form davon ist.
11. Pharmazeutische Formulierung nach einem der vorangegangenen Ansprüche, worin die Formulierung eine Kapsel, eine Tablette, ein Pulver, ein Päckchen oder Sprühnebel ist.
12. Formulierung nach einem der Ansprüche 1-11 zur Verwendung in einem Verfahren zur Prävention und/oder Behandlung von Zöliakie und/oder Dermatitis herpetiformis in einem Patienten, wobei das Verfahren den Schritt des oralen Verabreichens der Formulierung an den Patienten zur selben Zeit mit der Aufnahme eines glutenhaltigen Nahrungsmittels durch den Patienten umfasst.

Revendications

1. Formulation pharmaceutique solide d'une endoprotéinase d'orge B2 (EPB2) ou d'une forme recombinée de celle-ci adaptée à une administration orale sous une forme galénique unitaire qui comprend un antioxydant contenant du soufre choisi parmi un sulfate, un thiol libre, et le groupe constitué du sulfite de sodium, du métabisulfite de sodium, de la N-acétylcystéine, de l'homocystéine, de la cystéine, du monothioglycérol, du thiosulfate de sodium, de l'acide α -lipoïque et du dithiothréitol.

2. Formulation pharmaceutique selon la revendication 1, dans laquelle ledit antioxydant est présent en une dose d'au moins environ 2 mg par forme galénique à pas plus d'environ 500 mg par forme galénique. 5
3. Formulation pharmaceutique selon l'une ou l'autre des revendications 1 ou 2, dans laquelle ledit antioxydant contenant du soufre est le métabisulfite de sodium, lequel est présent à raison d'environ 2 à 50 mg par forme galénique. 10
4. Formulation pharmaceutique selon l'une ou l'autre des revendications 1 ou 2, dans laquelle ledit antioxydant contenant du soufre est la cystéine, laquelle est présente à raison d'environ 5 à 400 mg par forme galénique. 15
5. Formulation pharmaceutique selon l'une quelconque des revendications précédentes qui comprend un chélateur. 20
6. Formulation pharmaceutique selon la revendication 5, dans laquelle le chélateur est l'EDTA ou l'acide citrique. 25
7. Formulation pharmaceutique selon la revendication 6, dans laquelle ledit chélateur est présent est présent en une dose d'au moins environ 5 mg par forme galénique à pas plus d'environ 500 mg/forme galénique. 30
8. Formulation pharmaceutique selon la revendication 7, dans laquelle ledit chélateur est l'acide citrique, lequel est présent à raison de 10 à 500 mg par forme galénique. 35
9. Formulation pharmaceutique selon l'une quelconque des revendications précédentes dans laquelle la dose unitaire comprend en outre une prolyl endopeptidase. 40
10. Formulation pharmaceutique selon la revendication 9 dans laquelle ladite prolyl endopeptidase est une *Sphingomonas capsulata* PEP ou une forme recombinée de celle-ci. 45
11. Formulation pharmaceutique selon l'une quelconque des revendications précédentes, laquelle formulation est une capsule, un comprimé, une poudre, un sachet ou des comprimés à saupoudrer. 50
12. Formulation selon l'une quelconque des revendications 1 à 11 pour utilisation dans un procédé de prévention et/ou de traitement d'une maladie coeliaque et/ou d'une dermatite herpétiforme chez un patient, ledit procédé comprenant l'étape d'administration orale de ladite formulation audit patient simultanément avec l'ingestion par le patient d'un aliment con-

tenant du gluten.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14






FIGURE 15


A




B



C



D



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 7291598 B [0002]
- US 7265097 B [0002]
- US 7244709 B [0002]
- US 6541606 B [0002]
- US 7351741 B [0002]
- US 6660313 B [0003]
- US 6562339 B [0003]
- US 7320788 B [0004] [0010] [0048]
- US 7303871 B [0004] [0010] [0048]
- US 2008115428 PCT [0010]
- US 2008115411 A [0010]
- US 2007044906 A [0010]
- US 2007047303 A [0010]
- US 20080193436 A [0010]
- US 6241973 B [0036]
- WO 2004058816 PCT [0036]
- US 20080039400 A [0036]
- US 20070264311 A [0036]
- US 20070148267 A [0036]
- US 20060134017 A [0036]
- US 20060134018 A [0036]
- WO 2008003425 PCT [0050]
- WO 2008115411 PCT [0079]

Non-patent literature cited in the description

- **WANG et al.** *JBC* 2002, 2000, vol. 277 (21), 18568-73 [0014]
- **ASCENZI et al.** *Curr. Protein Pept. Sci.*, 2001, vol. 2, 137-153 [0014]
- **VENTURINI et al.** *Biochem. Biophys. Res. Commun.*, 2000, vol. 270, 437-441 [0014]
- **BJORNE et al.** *J. Clin. Invest.*, 2004, vol. 113, 106-114 [0015]
- Remington's Pharmaceutical Sciences. Mace Publishing Company, 1985 [0041]
- **LANGER.** *Science*, 1990, vol. 249, 1527-1533 [0041]
- **CLARE HASLER.** Ph.D. University of Illinois [0044]