



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
07.03.2012 Bulletin 2012/10

(51) Int Cl.:
A61K 31/58 (2006.01) A61P 7/10 (2006.01)

(21) Application number: **11009239.2**

(22) Date of filing: **22.06.2010**

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR

(72) Inventor: **Bar-Or, David**
Englewood
Colorado 80110 (US)

(30) Priority: **22.06.2009 US 219185**
18.03.2010 US 315350 P

(74) Representative: **Schaeberle, Steffen**
Hoefer & Partner
Patentanwälte
Pilgersheimer Strasse 20
81543 München (DE)

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
10792567.9 / 2 326 332

Remarks:

This application was filed on 22-11-2011 as a divisional application to the application mentioned under INID code 62.

(71) Applicant: **DMI Acquisition Corp.**
Greenwood Village, CO 80111 (US)

(54) **Method for treatment of diseases**

(57) The present invention relates to a vascular-hyperpermeability-inhibiting amount of a danazol com-

pound, for use for inhibiting vascular hyperpermeability in an animal in need thereof.




FIGURE 1

Description

[0001] This application claims benefit of provisional application 61/219,185, filed June 22, 2009, and provisional application 61/315,350, filed March 18, 2010, the complete disclosures of both of which are incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The invention relates to a method of inhibiting vascular hyperpermeability and the edema and other adverse effects that result from it. The invention also relates to a method of modulating the cytoskeleton of endothelial cells. Both methods comprise administering a danazol compound to an animal.

BACKGROUND

[0003] The vascular endothelium lines the inside of all blood vessels. It acts as the interface between the blood and the tissues and organs. The endothelium forms a semipermeable barrier that maintains the integrity of the blood fluid compartment, but permits passage of water, ions, small molecules, macromolecules and cells in a regulated manner. Dysregulation of this process produces vascular leakage into underlying tissues. Leakage of fluid into tissues causing edema can have serious and life threatening consequences in a variety of diseases. Accordingly, it would be highly desirable to have a method for reducing edema, preferably at its earliest stage, and restoring the endothelial barrier to physiological.

SUMMARY OF THE INVENTION

[0004] The invention provides such a method. In particular, the invention provides a method of inhibiting vascular hyperpermeability and the edema and other adverse effects that result from it. The method comprises administering a vascular-hyperpermeability-inhibiting amount of a danazol compound to an animal in need thereof. Inhibition of vascular hyperpermeability according to the invention includes inhibition of paracellular-caused hyperpermeability and transcytosis-caused hyperpermeability. Recent evidence indicates that transcytosis-caused hyperpermeability is the first step of a process that ultimately leads to tissue and organ damage in many diseases and conditions. Accordingly, the present invention provides a means of early intervention in these diseases and conditions which can reduce, delay or even potentially prevent the tissue and organ damage seen in them.

[0005] The invention also provides a method of modulating the cytoskeleton of endothelial cells in an animal. The method comprises administering an effective amount of a danazol compound to the animal.

[0006] "Vascular hyperpermeability" is used herein to mean permeability of a vascular endothelium that is increased as compared to basal levels. "Vascular hyperpermeability," as used herein, includes paracellular-caused hyperpermeability and transcytosis-caused hyperpermeability.

[0007] "Paracellular-caused hyperpermeability" is used herein to mean vascular hyperpermeability caused by paracellular transport that is increased as compared to basal levels. Other features of "paracellular-caused hyperpermeability" are described below.

[0008] "Paracellular transport" is used herein to mean the movement of ions, molecules and fluids through the inter-endothelial junctions (IEJs) between the endothelial cells of an endothelium.

[0009] "Transcytosis-caused hyperpermeability" is used herein to mean vascular hyperpermeability caused by transcytosis that is increased as compared to basal levels.

[0010] "Transcytosis" is used herein to mean the active transport of macromolecules and accompanying fluid-phase plasma constituents across the endothelial cells of the endothelium. Other features of "transcytosis" are described below.

[0011] "Basal level" is used herein to refer to the level found in a normal tissue or organ.

[0012] "inhibiting, "inhibit" and similar terms are used herein to mean to reduce, delay or prevent.

[0013] An animal is "in need of" treatment according to the invention if the animal, presently has a disease or condition mediated by vascular hyperpermeability, exhibits early signs of such a disease or condition, or has a predisposition to develop such a disease or condition.

[0014] "Mediated" and similar terms are used here to mean caused by, causing, involving or exacerbated by, vascular hyperpermeability.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015]

Figure 1 shows the OD levels measured after incubation of HUVEC cells with danazol as a measure of its ability to

prevent initial proliferation of endothelial cells.

Figure 2 shows photographs of HUVEC cells taken after incubation with danazol as a measure of its ability to prevent tube formation of endothelial cells. A = control; B = 1 μ M danazol, C = 10 μ M danazol, D = 50 μ M danazol and E = 50 μ M LY294002.

Figure 3 shows the fluorescence measured after treatment of HUVEC cells with danazol as a measure of their ability to prevent endothelial cell invasion.

DETAILED DESCRIPTION OF THE PRESENTLY-PREFERRED EMBODIMENTS OF THE INVENTION

[0016] The endothelium is a key gatekeeper controlling the exchange of molecules from the blood to the tissue parenchyma. It largely controls the permeability of a particular vascular bed to blood-borne molecules. The permeability and selectivity of the endothelial cell barrier is strongly dependent on the structure and type of endothelium lining the microvasculature in different vascular beds. Endothelial cells lining the microvascular beds of different organs exhibit structural differentiation that can be grouped into three primary morphologic categories: sinusoidal, fenestrated and continuous.

[0017] Sinusoidal endothelium (also referred to as "discontinuous endothelium") has large intercellular and intracellular gaps and no basement membrane, allowing for minimally restricted transport of molecules from the capillary lumen into the tissue and vice versa. Sinusoidal endothelium is found in liver, spleen and bone marrow.

[0018] Fenestrated endothelia are characterized by the presence of a large number of circular transcellular openings called fenestrae with a diameter of 60 to 80 nm. Fenestrated endothelia are found in tissues and organs that require rapid exchange of small molecules, including kidney (glomeruli, peritubular capillaries and ascending vasa recta), pancreas, adrenal glands, endocrine glands and intestine. The fenestrae are covered by thin diaphragms, except for those in mature, healthy glomeruli. See Ichimura et al., J. Am. Soc. Nephrol., 19:1463-1471 (2008).

[0019] Continuous endothelia do not contain fenestrae or large gaps. Instead, continuous endothelia are characterized by an uninterrupted endothelial cell monolayer. Most endothelia in the body are continuous endothelia, and continuous endothelium is found in, or around, the brain (blood brain barrier), diaphragm, duodenal musculature, fat, heart, some areas of the kidneys (papillary microvasculature, descending vasa recta), large blood vessels, lungs, mesentery, nerves, retina (blood retinal barrier), skeletal muscle, testis and other tissues and organs of the body.

[0020] Endothelial transport in continuous endothelium can be thought of in a general sense as occurring by paracellular and transcellular pathways. The paracellular pathway is the pathway between endothelial cells, through the interendothelial junctions (IEJs). In unperturbed continuous endothelium, water, ions and small molecules are transported paracellularly by diffusion and convection. A significant amount of water (up to 40%) also crosses the endothelial cell barrier transcellularly through water-transporting membrane channels called aquaporins. A variety of stimuli can disrupt the organization of the IEJs, thereby opening gaps in the endothelial barrier. The formation of these intercellular gaps allows passage of fluid, ions, macromolecules (e.g., proteins) and other plasma constituents between the endothelial cells in an unrestricted manner. This paracellular-caused hyperpermeability produces edema and other adverse effects that can eventually result in damage to tissues and organs.

[0021] The transcellular pathway is responsible for the active transport of macromolecules, such as albumin and other plasma proteins, across the endothelial cells, a process referred to as "transcytosis." The transport of macromolecules occurs in vesicles called caveolae. Almost all continuous endothelia have abundant caveolae, except for continuous endothelia located in brain and testes which have few caveolae. Transcytosis is a multi-step process that involves successive caveolae budding and fission from the plasmalemma and translocation across the cell, followed by docking and fusion with the opposite plasmalemma, where the caveolae release their contents by exocytosis into the interstitium. Transcytosis is selective and tightly regulated under normal physiological conditions.

[0022] There is a growing realization of the fundamental importance of the transcellular pathway. Transcytosis of plasma proteins, especially albumin which represents 65% of plasma protein, is of particular interest because of its ability to regulate the transvascular oncotic pressure gradient. As can be appreciated, then, increased transcytosis of albumin and other plasma proteins above basal levels will increase the tissue protein concentration of them which, in turn, will cause water to move across the endothelial barrier, thereby producing edema.

[0023] Low density lipoproteins (LDL) are also transported across endothelial cells by transcytosis. In hyperlipidemia, a significant increase in transcytosis of LDL has been detected as the initial event in atherogenesis. The LDL accumulates in the subendothelial space, trapped within the expanded basal lamina and extracellular matrix. The subendothelial lipoprotein accumulation in hyperlipidemia is followed by a cascade of events resulting in atheromatous plaque formation. Advanced atherosclerotic lesions are reported to be occasionally accompanied by the opening of IEJs and massive uncontrolled passage of LDL and albumin.

[0024] Vascular complications are a hallmark of diabetes. At the level of large vessels, the disease appears to be expressed as an acceleration of an atherosclerotic process. With respect to microangiopathy, alterations in the microvasculature of the retina, renal glomerulus and nerves cause the greatest number of clinical complications, but a con-

tinuously increasing number of investigations show that diabetes also affects the microvasculature of other organs, such as the mesentery, skin, skeletal muscle, heart, brain and lung, causing additional clinical complications. In all of these vascular beds, changes in vascular permeability appear to represent a hallmark of the diabetic endothelial dysfunction.

[0025] In continuous endothelium, capillary hyperpermeability to plasma macromolecules in the early phase of diabetes is explained by an intensification of transendothelial vesicular transport (*i.e.*, by increased transcytosis) and not by the destabilization of the IEs. In addition, the endothelial cells of diabetics, including those of the brain, have been reported to contain an increased number of caveolae as compared to normals, and glycated proteins, particularly glycated albumin, are taken up by endothelial cells and transcytosed at substantially greater rates than their native forms. Further, increased transcytosis of macromolecules is a process that continues beyond the early phase of diabetes and appears to be a cause of edema in diabetic tissues and organs throughout the disease if left untreated. This edema, in turn, leads to tissue and organ damage. Similar increases in transcellular transport of macromolecules have been reported in hypertension.

[0026] Paracellular-caused hyperpermeability is also a factor in diabetes and the vascular complications of diabetes. The IEs of the paracellular pathway include the adherens junctions (AJs) and tight junctions (TJs). Diabetes alters the content, phosphorylation and localization of certain proteins in both the AJs and TJs, thereby contributing to increased endothelial barrier permeability.

[0027] In support of the foregoing discussion and for further information, see Frank et al., *Cell Tissue Res.*, 335:41-47 (2009), Simionescu et al., *Cell Tissue Res.*, 335:27-40 (2009); van den Berg et al., *J. Cyst. Fibros.*, 7(6): 515-519 (2008); Viazzi et al., *Hypertens. Res.*, 31:873-879 (2008); Antonetti et al., Chapter 14, pages 340-342, in *Diabetic Retinopathy* (edited by Elia J. Duh, Humana Press, 2008), Felinski et al., *Current Eye Research*, 30:949-957 (2005), Pascariu et al., *Journal of Histochemistry & Cytochemistry*, 52(1):65-76 (2004); Bouchard et al., *Diabetologia*, 45:1017-1025 (2002); Arshi et al., *Laboratory Investigation*, 80(8):1171-1184 (2000); Viores et al., *Documenta Ophthalmologica*, 97:217-228 (1999); Oomen et al., *European Journal of Clinical Investigation*, 29:1035-1040 (1999); Viores et al., *Pathol. Res. Pract.*, 194:497-505 (1998); Antonetti et al., *Diabetes*, 47:1953-1959 (1998), Popov et al., *Acta Diabetol.*, 34:285-293 (1997); Yamaji et al., *Circulation Research*, 72:947-957 (1993); Viores et al., *Histochemical Journal*, 25:648-663 (1993); Beals et al., *Microvascular Research*, 45:11-19 (1993); Caldwell et al., *Investigative Ophthalmol Visual Sci.*, 33(5):1610-1619 (1992).

[0028] Endothelial transport in fenestrated endothelium also occurs by transcytosis and the paracellular pathway. In addition, endothelial transport occurs by means of the fenestrae. Fenestrated endothelia show a remarkably high permeability to water and small hydrophilic solutes due to the presence of the fenestrae.

[0029] The fenestrae may or may not be covered by a diaphragm. The locations of endothelium with diaphragmed fenestrae include endocrine tissue (*e.g.*, pancreatic islets and adrenal vortex), gastrointestinal mucosa and renal peritubular capillaries. The permeability to plasma proteins of fenestrated endothelium with diaphragmed fenestrae does not exceed that of continuous endothelium.

[0030] The locations of endothelium with nondiaphragmed fenestrae include the glomeruli of the kidneys. The glomerular fenestrated endothelium is covered by a glycocalyx that extends into the fenestrae (forming so-called "seive plugs") and by a more loosely associated endothelial cell surface layer of glycoproteins. Mathematical analyses of functional permselectivity studies have concluded that the glomerular endothelial cell glycocalyx, including that present in the fenestrae, and its associated surface layer account for the retention of up to 95% of plasma proteins within the circulation.

[0031] Loss of fenestrae in the glomerular endothelium has been found to be associated with proteinuria in several diseases, including diabetic nephropathy, transplant glomerulopathy, pre-eclampsia, diabetes, renal failure, cyclosporine nephropathy, serum sickness nephritis and Thy-1 nephritis. Actin rearrangement and, in particular, depolymerization of stress fibers have been found to be important for the formation and maintenance of fenestrae.

[0032] In support of the foregoing discussion of fenestrated endothelia and for additional information, see Satchell et al., *Am. J. Physiol. Renal Physiol.*, 296:F947-F956 (2009); Haraldsson et al., *Curr. Opin. Nephrol. Hypertens.*, 18:331-335 (2009); Ichimura et al., *J. Am. Soc. Nephrol.*, 19:1463-1471 (2008); Ballermann, *Nephron Physiol.*, 106:19-25 (2007); Toyoda et al., *Diabetes*, 56:2155-2160 (2007); Stan, "Endothelial Structures Involved In Vascular Permeability," pages 679-688, *Endothelial Biomedicine* (ed. Aird, Cambridge University Press, Cambridge, 2007); Simionescu and Antohe, "Functional Ultrastructure of the Vascular Endothelium: Changes in Various Pathologies," pages 42-69, *The Vascular Endothelium I* (eds. Moncada and Higgs, Springer-Verlag, Berlin, 2006).

[0033] Endothelial transport in sinusoidal endothelium occurs by transcytosis and through the intercellular gaps (interendothelial slits) and intracellular gaps (fenestrae). Treatment of sinusoidal endothelium with actin filament-disrupting drugs can induce a substantial and rapid increase in the number of gaps, indicating regulation of the porosity of the endothelial lining by the actin cytoskeleton. Other cytoskeleton altering drugs have been reported to change the diameters of fenestrae. Therefore, the fenestrae-associated cytoskeleton probably controls the important function of endothelial filtration in sinusoidal endothelium. In liver, defenestration (loss of fenestrae), which causes a reduction in permeability of the endothelium, has been associated with the pathogenesis of several diseases and conditions, including aging, atherogenesis, atherosclerosis, cirrhosis, fibrosis, liver failure and primary and metastatic liver cancers. In support of

the foregoing and for additional information, see Yokomori, *Med. Mol. Morphol.*, 41:1-4 (2008); Stan, "Endothelial Structures Involved In Vascular Permeability," pages 679-688, *Endothelial Biomedicine* (ed. Aird, Cambridge University Press, Cambridge, 2007); DeLeve, "The Hepatic Sinusoidal Endothelial Cell," pages 1226-1238, *Endothelial Biomedicine* (ed. Aird, Cambridge University Press, Cambridge, 2007); Pries and Kuebler, "Normal Endothelium," pages 1-40, *The Vascular Endothelium I* (eds. Moncada and Higgs, Springer-Verlag, Berlin, 2006); Simionescu and Antohe, "Functional Ultrastructure of the Vascular Endothelium: Changes in Various Pathologies," pages 42-69, *The Vascular-Endothelium I* (eds. Moncada and Higgs, Springer-Verlag, Berlin, 2006); Braet and Wisse, *Comparative Hepatology*, 1:1-17 (2002); Kanai et al., *Anat. Rec.*, 244:175-181 (1996); Kempka et al., *Exp. Cell Res.*, 176:38-48 (1988); Kishimoto et al., *Am. J. Anat.*, 178:241-249 (1987).

[0034] The invention provides a method of inhibiting vascular hyperpermeability present in any tissue or organ containing or surrounded by continuous endothelium. As noted above, continuous endothelium is present in, or around, the brain (blood brain barrier), diaphragm, duodenal musculature, fat, heart, some areas of the kidneys (papillary microvasculature, descending vasa recta), large blood vessels, lungs, mesentery, nerves, retina (blood retinal barrier), skeletal muscle, skin, testis, umbilical vein and other tissues and organs of the body. Preferably, the continuous endothelium is that found in or around the brain, heart, lungs, nerves or retina.

[0035] The invention also provides a method of inhibiting vascular hyperpermeability present in any tissue or organ containing or surrounded by fenestrated endothelium. As noted above, fenestrated endothelium is present in, or around, the kidney (glomeruli, peritubular capillaries and ascending vasa recta), pancreas, adrenal glands, endocrine glands and intestine. Preferably, the fenestrated endothelium is that found in the kidneys, especially that found in the glomeruli of the kidneys.

[0036] Further, any disease or condition mediated by vascular hyperpermeability can be treated by the method of the invention. Such diseases and conditions include diabetes, hypertension and atherosclerosis.

[0037] In particular, the vascular complications of diabetes, including those of the brain, heart, kidneys, lung, mesentery, nerves, retina, skeletal muscle, skin and other tissues and organs containing continuous or fenestrated endothelium, can be treated by the present invention. These vascular complications include edema, accumulation of LDL in the subendothelial space, accelerated atherosclerosis, and the following: brain (accelerated aging of vessel walls), heart (myocardial edema, myocardial fibrosis, diastolic dysfunction, diabetic cardiomyopathy), kidneys (diabetic nephropathy), lung (retardation of lung development in the fetuses of diabetic mothers, alterations of several pulmonary physiological parameters and increased susceptibility to infections), mesentery (vascular hyperplasia), nerves (diabetic neuropathy), retina (macular edema and diabetic retinopathy) and skin (redness, discoloration, dryness and ulcerations).

[0038] Diabetic retinopathy is a leading cause of blindness that affects approximately 25% of the estimated 21 million Americans with diabetes. Although its incidence and progression can be reduced by intensive glycemic and blood pressure control, nearly all patients with type 1 diabetes mellitus and over 60% of those with type 2 diabetes mellitus eventually develop diabetic retinopathy. There are two stages of diabetic retinopathy. The first, non-proliferative retinopathy, is the earlier stage of the disease and is characterized by increased vascular permeability, microaneurysms, edema and eventually vessel closures. Neovascularization is not a component of the nonproliferative phase. Most visual loss during this stage is due to the fluid accumulating in the macula, the central area of the retina. This accumulation of fluid is called macular edema and can cause temporary or permanent decreased vision. The second stage of diabetic retinopathy is called proliferative retinopathy and is characterized by abnormal new vessel formation. Unfortunately, this abnormal neovascularization can be very damaging because it can cause bleeding in the eye, retinal scar tissue, diabetic retinal detachments or glaucoma, any of which can cause decreased vision or blindness. Macular edema can also occur in the proliferative phase.

[0039] Diabetic neuropathy is a common serious complication of diabetes. There are four main types of diabetic neuropathy: peripheral neuropathy, autonomic neuropathy, radiculoplexus neuropathy and mononeuropathy. The signs and symptoms of peripheral neuropathy, the most common type of diabetic neuropathy, include numbness or reduced ability to feel pain or changes in temperature (especially in the feet and toes), a tingling or burning feeling, sharp pain, pain when walking, extreme sensitivity to the lightest touch, muscle weakness, difficulty walking, and serious foot problems (such as ulcers, infections, deformities and bone and joint pain). Autonomic neuropathy affects the autonomic nervous system that controls the heart, bladder, lungs, stomach, intestines, sex organs and eyes, and problems in any of these areas can occur. Radiculoplexus neuropathy (also called diabetic amyotrophy, femoral neuropathy or proximal neuropathy) usually affects nerves in the hips, shoulders or abdomen, usually on one side of the body. Mononeuropathy means damage to just one nerve, typically in an arm, leg or the face. Common complications of diabetic neuropathy include loss of limbs (e.g., toes, feet or legs), charcot joints, urinary tract infections, urinary incontinence, hypoglycemia unawareness (may even be fatal), low blood pressure, digestive problems (e.g., constipation, diarrhea, nausea and vomiting), sexual dysfunction (e.g., erectile dysfunction), and increased or decreased sweating. As can be seen, symptoms can range from mild to painful, disabling and even fatal.

[0040] Diabetic nephropathy is the most common cause of end-stage renal disease in the United States. It is a vascular complication of diabetes that affects the glomerular capillaries of the kidney and reduces the kidney's filtration ability.

Nephropathy is first indicated by the appearance of hyperfiltration and then microalbuminuria. Heavy proteinuria and a progressive decline in renal function precede end-stage renal disease. Typically, before any signs of nephropathy appear, retinopathy has usually been diagnosed. Renal transplant is usually recommended to patients with end-stage renal disease due to diabetes. Survival rate at 5 years for patients receiving a transplant is about 60% compared with only 2% for those on dialysis.

[0041] Hypertension typically develops over many years, and it affects nearly everyone eventually. Uncontrolled hypertension increases the risk of serious health problems, including heart attack, congestive heart failure, stroke, peripheral artery disease, kidney failure, aneurysms, eye damage, and problems with memory or understanding.

[0042] Atherosclerosis also develops gradually. Atherosclerosis can affect the coronary arteries, the carotid artery, the peripheral arteries or the microvasculature, and complications of atherosclerosis include coronary artery disease (which can cause angina or a heart attack), coronary microvascular disease, carotid artery disease (which can cause a transient ischemic attack or stroke), peripheral artery disease (which can cause loss of sensitivity to heat and cold or even tissue death), and aneurysms.

[0043] Additional diseases and conditions that can be treated according to the invention include acute lung injury, acute respiratory distress syndrome (ARDS), age-related macular degeneration, cerebral edema, choroidal edema, choroiditis, coronary microvascular disease, cerebral microvascular disease, Eals disease, edema caused by injury (e.g., trauma or burns), edema associated with hypertension, glomerular vascular leakage, hemorrhagic shock, Irvine Gass Syndrome, ischemia, macular edema (e.g., caused by vascular occlusions, post-intraocular surgery (e.g., cataract surgery), uveitis or retinitis pigmentosa, in addition to that caused by diabetes), nephritis (e.g., glomerulonephritis, serum sickness nephritis and Thy-1 nephritis), nephropathies, nephrotic edema, nephrotic syndrome, neuropathies, organ failure due to tissue edema (e.g., in sepsis or due to trauma), pre-eclampsia, pulmonary edema, pulmonary hypertension, renal failure, retinal edema, retinal hemorrhage, retinal vein occlusions (e.g., branch or central vein occlusions), retinitis, retinopathies (e.g., arteriosclerotic retinopathy, hypertensive retinopathy, radiation retinopathy, sickle cell retinopathy and retinopathy of prematurity, in addition to diabetic retinopathy), silent cerebral infarction, systemic inflammatory response syndromes (SIRS), transplant glomerulopathy, uveitis, vascular leakage syndrome, vitreous hemorrhage and Von Hippel Lindau disease. In addition, certain drugs, including those used to treat multiple sclerosis, are known to cause vascular hyperpermeability, and danazol can be used to reduce this unwanted side effect when using these drugs. Hereditary and acquired angioedema are expressly excluded from those diseases and conditions that can be treated according to the invention.

[0044] "Treat," "treating" or "treatment" is used herein to mean to reduce (wholly or partially) the symptoms, duration or severity of a disease or condition, including curing the disease, or to prevent the disease or condition.

[0045] Recent evidence indicates that transcytosis-caused hyperpermeability is the first step of a process that ultimately leads to tissue and organ damage in many diseases and conditions. Accordingly, the present invention provides a means of early intervention in these diseases and conditions which can reduce, delay or even potentially prevent the tissue and organ damage seen in them. For instance, an animal can be treated immediately upon diagnosis of one of the disease or conditions treatable according to the invention (those diseases and conditions described above). Alternatively, preferred is the treatment of animals who have early signs of, or a predisposition to develop, such a disease or condition prior to the existence of symptoms. Early signs of, and risk factors for, diabetes, hypertension and atherosclerosis are well known, and treatment of an animal exhibiting these early signs or risk factors can be started prior to the presence of symptoms of the disease or condition (i.e., prophylactically).

[0046] For instance, treatment of a patient who is diagnosed with diabetes can be started immediately upon diagnosis. In particular, diabetics should preferably be treated with a danazol compound prior to any symptoms of a vascular complication being present, although this is not usually possible, since most diabetics show such symptoms when they are diagnosed (see below). Alternatively, diabetics should be treated while nonproliferative diabetic retinopathy is mild (i.e., mild levels of microaneurysms and intraretinal hemorrhage). See Diabetic Retinopathy, page 9 (Ed. Elia Duh, M.D., Human Press, 2008). Such early treatment will provide the best chance of preventing macular edema and progression of the retinopathy to proliferative diabetic retinopathy. Also, the presence of diabetic retinopathy is considered a sign that other microvascular complications of diabetes exist or will develop (see *Id.*, pages 474-477), and early treatment may also prevent or reduce these additional complications. Of course, more advanced diseases and conditions that are vascular complications of diabetes can also be treated with beneficial results.

[0047] However, as noted above, vascular complications are often already present by the time diabetes is diagnosed. Accordingly, it is preferable to prophylactically treat a patient who has early signs of, or a predisposition to develop, diabetes. These early signs and risk factors include fasting glucose that is high, but not high enough to be classified as diabetes ("prediabetes"), hyperinsulinemia, hypertension, dyslipidemia (high cholesterol, high triglycerides, high low-density lipoprotein, and/or low level of high-density lipoprotein), obesity (body mass index above 25), inactivity, over 45 years of age, inadequate sleep, family history of diabetes, minority race, history of gestational diabetes and history of polycystic ovary syndrome.


[0048] Similarly, treatment of a patient who is diagnosed with hypertension can be started immediately upon diagnosis.

Hypertension typically does not cause any symptoms, but prophylactic treatment can be started in a patient who has a predisposition to develop hypertension. Risk factors for hypertension include age, race (hypertension is more common blacks), family history (hypertension runs in families), overweight or obesity, lack of activity, smoking tobacco, too much salt in the diet, too little potassium in the diet, too little vitamin D in the diet, drinking too much alcohol, high levels of stress, certain chronic conditions (e.g., high cholesterol, diabetes, kidney disease and sleep apnea) and use of certain drugs (e.g., oral contraceptives, amphetamines, diet pills, and some cold and allergy medications).

[0049] Treatment of a patient who is diagnosed with atherosclerosis can be started immediately upon diagnosis. However, it is preferable to prophylactically treat a patient who has early signs of, or a predisposition to develop, atherosclerosis. Early signs and risk factors for atherosclerosis include age, a family history of aneurysm or early heart disease, hypertension, high cholesterol, high triglycerides, insulin resistance, diabetes, obesity, smoking, lack of physical activity, unhealthy diet, and high level of C-reactive protein.

[0050] The method of the invention for inhibiting vascular hyperpermeability comprises administering an effective amount of a danazol compound to an animal in need thereof to inhibit the vascular hyperpermeability. As used here, "a danazol compound" means danazol, prodrugs of danazol and pharmaceutically acceptable salts of danazol and its prodrugs.

[0051] Danazol (17 α -pregna-2,4-dien-20-yno[2,3-d]-isoxazol-17 β -ol) is a known synthetic steroid hormone. Its structure is:



Danazol

[0052] Methods of making danazol are known in the art. See e.g., U.S. Patents No. 3,135,743, and GB Patent No. 905,844. Also, danazol is available commercially from many sources, including Barr Pharmaceuticals, Inc., Lannett Co., Inc., sanofi-aventis Canada, Sigma-Aldrich, and Parchem Trading Ltd.

[0053] "Prodrug" means any compound which releases an active parent drug (danazol in this case) *in vivo* when such prodrug is administered to an animal. Prodrugs of danazol include danazol wherein the hydroxyl group is bonded to any group that may be cleaved *in vivo* to generate the free hydroxyl. Examples of danazol prodrugs include esters (e.g., acetate, formate, and benzoate derivatives) of danazol.

[0054] The pharmaceutically-acceptable salts of danazol and its prodrugs include conventional non-toxic salts, such as salts derived from inorganic acids (such as hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, and the like), organic acids (such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, glutamic, aspartic, benzoic, salicylic, oxalic, ascorbic acid, and the like) or bases (such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation or organic cations derived from N,N-dibenzylethylenediamine, D-glucosamine, or ethylenediamine). The salts are prepared in a conventional manner, e.g., by neutralizing the free base form of the compound with an acid. In particular, isoxazoles, such as danazol, are weakly basic substances and will form acid-addition salts upon addition of strong acids and quaternary ammonium salts upon addition of esters of strong acids (e.g., an ester of a strong inorganic or organic sulfonic acid, preferably a lower-alkyl, lower alkenyl or lower aralkyl ester, such as methyl iodide, ethyl iodide, ethyl bromide, propyl bromide, butyl bromide, allyl bromide, methyl sulfate, methyl benzenesulfonate, methyl-p-toluene-sulfonate, benzyl chloride and the like). See U.S. Patent No. 3,135,743.

[0055] As noted above, a danazol compound can be used to inhibit vascular hyperpermeability and to treat a disease or condition mediated by vascular hyperpermeability. To do so, the danazol compound is administered to an animal in need of treatment. Preferably, the animal is a mammal, such as a rabbit, goat, dog, cat, horse or human. Most preferably, the animal is a human.

[0056] Effective dosage forms, modes of administration and dosage amounts for the compounds of the invention (*i.e.*, danazol, a prodrug of danazol or a pharmaceutically-acceptable salt of either one of them) may be determined empirically

using the guidance provided herein. It is understood by those skilled in the art that the dosage amount will vary with the particular disease or condition to be treated, the severity of the disease or condition, the route(s) of administration, the duration of the treatment, the identity of any other drugs being administered to the animal, the age, size and species of the animal, and like factors known in the medical and veterinary arts. In general, a suitable daily dose of a compound of the present invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. However, the daily dosage will be determined by an attending physician or veterinarian within the scope of sound medical judgment. If desired, the effective daily dose may be administered as two, three, four, five, six or more sub-doses, administered separately at appropriate intervals throughout the day. Administration of the compound should be continued until an acceptable response is achieved.

[0057] Danazol compounds have previously been reported to inhibit angiogenesis. See PCT application WO 2007/009087. Surprisingly and quite unexpectedly, it has been found that danazol compounds can be used in the practice of the present invention at optimum doses that are about 100-1000 times lower than those previously reported for inhibiting angiogenesis and substantially less than those amounts currently administered to patients for the treatment of other diseases and conditions (typically 200-800 mg/day for an adult human). Uses of these lower doses of danazol compounds should avoid any significant side effects, perhaps all side effects, which will be especially advantageous for early or prophylactic treatment of diseases and conditions according to the present invention.

[0058] In particular, an effective dosage amount of a danazol compound for inhibiting vascular hyperpermeability will be from 0.1 ng/kg/day to 35 mg/kg/day, preferably from 40 ng/kg/day to 5.0 mg/kg/day, most preferably from 100 ng/kg/day to 1.5 mg/kg/day. An effective dosage amount will also be that amount that will result in a concentration in a relevant fluid (*e.g.*, blood) from 0.0001 μ M to 5 μ M, preferably from 0.1 μ M to 1.0 μ M, more preferably from 0.1 μ M to 0.5 μ M, most preferably about 0.1 μ M. An effective dosage amount will also be that amount that will result in a concentration in the tissue or organ to be treated of about 0.17% (weight/weight) or less, preferably from 0.00034% to 0.17%, most preferably 0.0034% to 0.017%. When given topically or locally, the danazol compound will preferably be administered at a concentration from 0.0001 μ M to 5 μ M, preferably from 0.1 μ M to 1.0 μ M, more preferably from 0.1 μ M to 0.5 μ M, most preferably about 0.1 μ M, or at a concentration of about 0.17% (weight/weight) or less, preferably from 0.00034% to 0.17%, most preferably 0.0034% to 0.017%. When given orally to an adult human, the dose will preferably be from about 1 ng/day to about 100 mg/day, more preferably the dose will be from about 1 mg/day to about 100 mg/day, most preferably the dose will be from about 10 mg/day to about 90 mg/day, preferably given in two equal doses per day. Further, danazol is expected to accumulate in cells and tissues, so that an initial (loading) dose (*e.g.* 100 mg per day) may be reduced after a period of time (*e.g.*, 2-4 weeks) to a lower maintenance dose (*e.g.* 1 mg per day) which can be given indefinitely without significant side effects, perhaps without any side effects. As used herein, a "vascular-hyperpermeability-inhibiting amount" of a danazol compound is defined to mean those amounts set forth above in this paragraph.

[0059] The invention also provides a method of modulating the cytoskeleton of endothelial cells in an animal. This embodiment of the invention is based on the discoveries that danazol inhibits F-actin stress fiber formation, causes the formation of cortical actin rings, enhances and prolongs the formation of cortical actin rings by sphingosine-1 phosphate (SIP), inhibits RhoA, increases phosphorylation of VE-cadherin, appears to activate barrier-stabilizing GTPases and appears to stabilize microtubules. Modulation of the cytoskeleton can reduce vascular hyperpermeability and increase vascular hypopermeability (*i.e.*, permeability below basal levels), thereby returning the endothelium to homeostasis. Accordingly, those diseases and conditions mediated by vascular hyperpermeability can be treated (see above) and those diseases and conditions mediated by vascular hypopermeability can also be treated. The latter type of diseases and conditions include aging liver, atherogenesis, atherosclerosis, cirrhosis, fibrosis of the liver, liver failure and primary and metastatic liver cancers.

[0060] The method of modulating the cytoskeleton of endothelial cells comprises administering an effective amount of a danazol compound to the animal. "Danazol compound" and "animal" have the same meanings as set forth above.

[0061] Effective dosage forms, modes of administration and dosage amounts for the compounds of the invention (*i.e.*, danazol, a prodrug of danazol or a pharmaceutically-acceptable salt of either one of them) for modulating the cytoskeleton may be determined empirically using the guidance provided herein. It is understood by those skilled in the art that the dosage amount will vary with the particular disease or condition to be treated, the severity of the disease or condition, the route(s) of administration, the duration of the treatment, the identity of any other drugs being administered to the animal, the age, size and species of the animal, and like factors known in the medical and veterinary arts. In general, a suitable daily dose of a compound of the present invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. However, the daily dosage will be determined by an attending physician or veterinarian within the scope of sound medical judgment. If desired, the effective daily dose may be administered as two, three, four, five, six or more sub-doses, administered separately at appropriate intervals throughout the day. Administration of the compound should be continued until an acceptable response is achieved.

[0062] In particular, an effective dosage amount of a danazol compound for modulating the cytoskeleton of endothelial cells will be from 0.1 ng/kg/day to 35 mg/kg/day, preferably from 40 ng/kg/day to 5.0 mg/kg/day, most preferably from

100 ng/kg/day to 1.5 mg/kg/day. An effective dosage amount will also be that amount that will result in a concentration in a relevant fluid (e.g., blood) from 0.0001 μ M to 5 μ M, preferably from 0.1 μ M to 1.0 μ M, more preferably from 0.1 μ M to 0.5 μ M, most preferably about 0.1 μ M. An effective dosage amount will also be that amount that will result in a concentration in the tissue or organ to be treated of about 0.17% (weight/weight) or less, preferably from 0.00034% to 0.17%, most preferably 0.0034% to 0.017%. When given topically or locally, the danazol compound will preferably be administered at a concentration from 0.0001 μ M to 5 μ M, preferably from 0.1 μ M to 1.0 μ M, more preferably from 0.1 μ M to 0.5 μ M, most preferably about 0.1 μ M, or at a concentration of about 0.17% (weight/weight) or less, preferably from 0.00034% to 0.17%, most preferably 0.0034% to 0.017%. When given orally to an adult human, the dose will preferably be from about 1 ng/day to about 100 mg/day, more preferably the dose will be from about 1 mg/day to about 100 mg/day, most preferably the dose will be from about 10 mg/day to about 90 mg/day, preferably given in two equal doses per day. Further, danazol is expected to accumulate in cells and tissues, so that an initial (loading) dose (e.g. 100 mg per day) may be reduced after a period of time (e.g., 2-4 weeks) to a lower maintenance dose (e.g. 1 mg per day) which can be given indefinitely without significant side effects, perhaps without any side effects.

[0063] The compounds of the present invention (i.e., danazol, prodrugs thereof and pharmaceutically-acceptable salts of either of them) may be administered to an animal patient for therapy by any suitable route of administration, including orally, nasally, parenterally (e.g., intravenously, intraperitoneally, subcutaneously or intramuscularly), transdermally, intraocularly and topically (including buccally and sublingually). Generally preferred is oral administration for any disease or condition treatable according to the invention. The preferred routes of administration for treatment of diseases and conditions of the eye are orally, intraocularly and topically. Most preferred is orally. It is quite unexpected and surprising that diseases of the eye can be treated by oral administration of a danazol compound, since successful treatment of such diseases and conditions by oral administration of a drug has not been previously reported. The preferred routes of administration for treatment of diseases and conditions of the brain are orally and parenterally. Most preferred is orally.

[0064] While it is possible for a compound of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical formulation (composition). The pharmaceutical compositions of the invention comprise a compound or compounds of the invention as an active ingredient in admixture with one or more pharmaceutically-acceptable carriers and, optionally, with one or more other compounds, drugs or other materials. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the animal. Pharmaceutically-acceptable carriers are well known in the art. Regardless of the route of administration selected, the compounds of the present invention are formulated into pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art. See, e.g., *Remington's Pharmaceutical Sciences*.

[0065] Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, powders, granules or as a solution or a suspension in an aqueous or non-aqueous liquid, or an oil-in-water or water-in-oil liquid emulsions, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia), and the like, each containing a predetermined amount of a compound or compounds of the present invention as an active ingredient. A compound or compounds of the present invention may also be administered as bolus, electuary or paste.

[0066] In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the active ingredient (i.e., danazol, a prodrug of danazol, a pharmaceutically-acceptable salt of either one of them, or combinations of the foregoing) is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, cetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

[0067] A tablet may be made by compression or molding optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

[0068] The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to

provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active ingredient can also be in microencapsulated form.

[0069] Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically-acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

[0070] Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

[0071] Suspensions, in addition to the active ingredient, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

[0072] The invention also provides pharmaceutical products suitable for treatment of the eye. Such pharmaceutical products include pharmaceutical compositions, devices and implants (which may be compositions or devices).

[0073] Pharmaceutical formulations (compositions) for intraocular injection of a compound or compounds of the invention into the eyeball include solutions, emulsions, suspensions, particles, capsules, microspheres, liposomes, matrices, etc. See, *e.g.*, U.S. Patent No. 6,060,463, U.S. Patent Application Publication No. 2005/0101582, and PCT application WO 2004/043480, the complete disclosures of which are incorporated herein by reference. For instance, a pharmaceutical formulation for intraocular injection may comprise one or more compounds of the invention in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or non-aqueous solutions, suspensions or emulsions, which may contain antioxidants, buffers, suspending agents, thickening agents or viscosity-enhancing agents (such as a hyaluronic acid polymer). Examples of suitable aqueous and nonaqueous carriers include water, saline (preferably 0.9%), dextrose in water (preferably 5%), buffers, dimethylsulfoxide, alcohols and polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like). These compositions may also contain adjuvants such as wetting agents and emulsifying agents and dispersing agents. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as polymers and gelatin. Injectable depot forms can be made by incorporating the drug into microcapsules or microspheres made of biodegradable polymers such as polylactide-polyglycolide. Examples of other biodegradable polymers include poly(orthoesters), poly(glycolic) acid, poly(lactic) acid, polycaprolactone and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes (composed of the usual ingredients, such as dipalmitoyl phosphatidylcholine) or microemulsions which are compatible with eye tissue. Depending on the ratio of drug to polymer or lipid, the nature of the particular polymer or lipid components, the type of liposome employed, and whether the microcapsules or microspheres are coated or uncoated, the rate of drug release from microcapsules, microspheres and liposomes can be controlled.

[0074] The compounds of the invention can also be administered surgically as an ocular implant. For instance, a reservoir container having a diffusible wall of polyvinyl alcohol or polyvinyl acetate and containing a compound or compounds of the invention can be implanted in or on the sclera. As another example, a compound or compounds of the invention can be incorporated into a polymeric matrix made of a polymer, such as polycaprolactone, poly(glycolic) acid, poly(lactic) acid, poly(anhydride), or a lipid, such as sebacic acid, and may be implanted on the sclera or in the eye. This is usually accomplished with the animal receiving a topical or local anaesthetic and using a small incision made behind the cornea. The matrix is then inserted through the incision and sutured to the sclera.

[0075] The compounds of the invention can also be administered topically to the eye, and a preferred embodiment of the invention is a topical pharmaceutical composition suitable for application to the eye. Topical pharmaceutical compositions suitable for application to the eye include solutions, suspensions, dispersions, drops, gels, hydrogels and ointments. See, *e.g.*, U.S. Patent No. 5,407,926 and PCT applications WO 2004/058289, WO 01/30337 and WO 01/68053, the complete disclosures of all of which are incorporated herein by reference.

[0076] Topical formulations suitable for application to the eye comprise one or more compounds of the invention in an aqueous or nonaqueous base. The topical formulations can also include absorption enhancers, permeation enhancers, thickening agents, viscosity enhancers, agents for adjusting and/or maintaining the pH, agents to adjust the osmotic pressure, preservatives, surfactants, buffers, salts (preferably sodium chloride), suspending agents, dispersing agents, solubilizing agents, stabilizers and/or tonicity agents. Topical formulations suitable for application to the eye will preferably comprise an absorption or permeation enhancer to promote absorption or permeation of the compound or compounds of the invention into the eye and/or a thickening agent or viscosity enhancer that is capable of increasing the residence

time of a compound or compounds of the invention in the eye. See PCT applications WO 2004/058289, WO 01/30337 and WO 01/68053. Exemplary absorption/permeation enhancers include methysulfonylmethane, alone or in combination with dimethylsulfoxide, carboxylic acids and surfactants. Exemplary thickening agents and viscosity enhancers include dextran, polyethylene glycols, polyvinylpyrrolidone, polysaccharide gels, Gelrite®, cellulosic polymers (such as hydroxypropyl methylcellulose), carboxyl-containing polymers (such as polymers or copolymers of acrylic acid), polyvinyl alcohol and hyaluronic acid or a salt thereof.

[0077] Liquid dosage forms (*e.g.*, solutions, suspensions, dispersions and drops) suitable for treatment of the eye can be prepared, for example, by dissolving, dispersing, suspending, etc. a compound or compounds of the invention in a vehicle, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol and the like, to form a solution, dispersion or suspension. If desired, the pharmaceutical formulation may also contain minor amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, pH buffering agents and the like, for example sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, etc.

[0078] Aqueous solutions and suspensions suitable for treatment of the eye can include, in addition to a compound or compounds of the invention, preservatives, surfactants, buffers, salts (preferably sodium chloride), tonicity agents and water. If suspensions are used, the particle sizes should be less than 10 μm to minimize eye irritation. If solutions or suspensions are used, the amount delivered to the eye should not exceed 50 μl to avoid excessive spillage from the eye.

[0079] Colloidal suspensions suitable for treatment of the eye are generally formed from microparticles (*i.e.*, microspheres, nanospheres, microcapsules or nanocapsules, where microspheres and nanospheres are generally monolithic particles of a polymer matrix in which the formulation is trapped, adsorbed, or otherwise contained, while with microcapsules and nanocapsules the formulation is actually encapsulated). The upper limit for the size of these microparticles is about 5 μm to about 10 μm .

[0080] Ophthalmic ointments suitable for treatment of the eye include a compound or compounds of the invention in an appropriate base, such as mineral oil, liquid lanolin, white petrolatum, a combination of two or all three of the foregoing, or polyethylene-mineral oil gel. A preservative may optionally be included.

[0081] Ophthalmic gels suitable for treatment of the eye include a compound or compounds of the invention suspended in a hydrophilic base, such as Carbopol-940 or a combination of ethanol, water and propylene glycol (*e.g.*, in a ratio of 40:40:20). A gelling agent, such as hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose or ammoniated glycyrrhizinate, is used. A preservative and/or a tonicity agent may optionally be included.

[0082] Hydrogels suitable for treatment of the eye are formed by incorporation of a swellable, gel-forming polymer, such as those listed above as thickening agents or viscosity enhancers, except that a formulation referred to in the art as a "hydrogel" typically has a higher viscosity than a formulation referred to as a "thickened" solution or suspension. In contrast to such preformed hydrogels, a formulation may also be prepared so to form a hydrogel *in situ* following application to the eye. Such gels are liquid at room temperature but gel at higher temperatures (and thus are termed "thermoreversible" hydrogels), such as when placed in contact with body fluids. Biocompatible polymers that impart this property include acrylic acid polymers and copolymers, N-isopropylacrylamide derivatives and ABA block copolymers of ethylene oxide and propylene oxide (conventionally referred to as "poloxamers" and available under the Pluronic® tradename from BASF-Wayndotte).

[0083] Preferred dispersions are liposomal, in which case the formulation is enclosed within liposomes (microscopic vesicles composed of alternating aqueous compartments and lipid bilayers).

[0084] Eye drops can be formulated with an aqueous or nonaqueous base also comprising one or more dispersing agents, solubilizing agents or suspending agents. Drops can be delivered by means of a simple eye dropper-capped bottle or by means of a plastic bottle adapted to deliver liquid contents dropwise by means of a specially shaped closure.

[0085] The compounds of the invention can also be applied topically by means of drug-impregnated solid carrier that is inserted into the eye. Drug release is generally effected by dissolution or bioerosion of the polymer, osmosis, or combinations thereof. Several matrix-type delivery systems can be used. Such systems include hydrophilic soft contact lenses impregnated or soaked with the desired compound of the invention, as well as biodegradable or soluble devices that need not be removed after placement in the eye. These soluble ocular inserts can be composed of any degradable substance that can be tolerated by the eye and that is compatible with the compound of the invention that is to be administered. Such substances include, but are not limited to, poly(vinyl alcohol), polymers and copolymers of polyacrylamide, ethylacrylate and vinylpyrrolidone, as well as cross-linked polypeptides or polysaccharides, such as chitin.

[0086] Dosage forms for the other types of topical administration (*i.e.*, not to the eye) or for transdermal administration of compounds of the invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches, drops and inhalants. The active ingredient may be mixed under sterile conditions with a pharmaceutically-acceptable carrier, and with any buffers, or propellants which may be required. The ointments, pastes, creams and gels may contain, in addition to the active ingredient, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof. Powders and sprays can contain, in addition to the active ingredient, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder or mixtures of these substances. Sprays can additionally contain

customary propellants such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane. Transdermal patches have the added advantage of providing controlled delivery of compounds of the invention to the body. Such dosage forms can be made by dissolving, dispersing or otherwise incorporating one or more compounds of the invention in a proper medium, such as an elastomeric matrix material. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can be controlled by either providing a rate-controlling membrane or dispersing the compound in a polymer matrix or gel. A drug-impregnated solid carrier (*e.g.*, a dressing) can also be used for topical administration.

[0087] Pharmaceutical formulations include those suitable for administration by inhalation or insufflation or for nasal administration. For administration to the upper (nasal) or lower respiratory tract by inhalation, the compounds of the invention are conveniently delivered from an insufflator, nebulizer or a pressurized pack or other convenient means of delivering an aerosol spray. Pressurized packs may comprise a suitable propellant such as dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide, or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount.

[0088] Alternatively, for administration by inhalation or insufflation, the composition may take the form of a dry powder, for example, a powder mix of one or more compounds of the invention and a suitable powder base, such as lactose or starch. The powder composition may be presented in unit dosage form in, for example, capsules or cartridges, or, *e.g.*, gelatin or blister packs from which the powder may be administered with the aid of an inhalator, insufflator or a metered-dose inhaler.

[0089] For intranasal administration, compounds of the invention may be administered by means of nose drops or a liquid spray, such as by means of a plastic bottle atomizer or metered-dose inhaler. Liquid sprays are conveniently delivered from pressurized packs. Typical of atomizers are the Mistometer (Wintrop) and Medihaler (Riker).

[0090] Nose drops may be formulated with an aqueous or nonaqueous base also comprising one or more dispersing agents, solubilizing agents or suspending agents. Drops can be delivered by means of a simple eye dropper-capped bottle or by means of a plastic bottle adapted to deliver liquid contents dropwise by means of a specially shaped closure.

[0091] Pharmaceutical compositions of this invention suitable for parenteral administrations comprise one or more compounds of the invention in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or non-aqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents. Also, drug-coated stents may be used.

[0092] Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[0093] These compositions may also contain adjuvants such as wetting agents, emulsifying agents and dispersing agents. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like in the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

[0094] In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally-administered drug is accomplished by dissolving or suspending the drug in an oil vehicle.

[0095] Injectable depot forms are made by forming microcapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissue. The injectable materials can be sterilized for example, by filtration through a bacterial-retaining filter.

[0096] The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampules and vials, and may be stored in a lyophilized condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the type described above.

[0097] A danazol compound may be given alone to treat a disease or condition involving vascular hyperpermeability or dysfunction of the cytoskeleton. Alternatively, the danazol compound may be given in combination with one or more other treatments or drugs suitable for treating the disease or condition. For instance, the danazol compound can be administered prior to, in conjunction with (including simultaneously with), or after the other treatment or drug. In the case

of another drug, the drug and the danazol compound may be administered in separate pharmaceutical compositions or as part of the same pharmaceutical composition. Suitable drugs are described in U.S. Application Number 12/820,325, the complete disclosure of which is incorporated herein by reference.

[0098] As used herein, "a" or "an" means one or more.

[0099] Additional objects, advantages and novel features of the present invention will become apparent to those skilled in the art by consideration of the following non-limiting examples.

EXAMPLES

Example 1: Danazol's Effects on Angiogenesis (Comparative)

A. HUVEC Cell Proliferation

Protocol:

[0100] Primary human umbilical vein endothelial cells (HUVECs) and EGM-2 growth medium were obtained from Cambrex (Walkersville, MD). The cells were passaged in medium supplemented with 2% fetal calf serum (FCS) in tissue culture flasks at 37°C and 5% CO₂. Subculturing was performed using trypsin when 60-80% confluence was obtained as specified by the supplier.

[0101] Cryopreserved ampoules of passage 2 HUVECs were thawed and plated in 96 well tissue culture plates at 5,000 cells/cm². A 50 mM stock solution of danazol was prepared in ethanol and the FCS in the medium was increased to 5% to keep danazol in solution. The cells were treated with medium containing final concentrations of danazol ranging from 0.1 to 100 μM in triplicates. 24, 48, and 72 hour incubations were performed and cell proliferation was determined utilizing Celltiter 96 AQueous One Solution Cell Proliferation assay from Promega (Madison, WI). In short, medium was aspirated from each well and the cells were washed with 200 μl Hepes buffered saline (HBSS) from Cambrex warmed to 37°C. 100 μl diluted celltiter solution (15 μl stock + 85 μl EGM-2 containing 0.1% FCS) were added to each well and incubated for an additional 4 hours. Optical density was determined by microplate reader using a 530 nm filter after blank subtraction and data presented as OD ± standard deviation. The final concentration of ethanol in the wells was less than 0.2% and had no effect on cell proliferation or viability.

[0102] All data are presented as representative experiments done in triplicate. Differences between subsets were analyzed using student t-test in Microsoft Excel. P < 0.05 was considered statistically significant.

Results, Observations and Discussion:

[0103] Culturing primary HUVECs in the presence of danazol decreased the OD obtained from the Promega celltiter proliferation assay in a time and dose dependent fashion (Figure 1). The celltiter assay is based on the reduction of the assay solution by dehydrogenase enzymes to a formazan dye that directly correlates to cell number.

[0104] Danazol treatment at 24 hours seemed to be effective only at very high doses. Significant decreases (p value < 0.05) in assay OD were seen at 10 μM or greater concentrations of danazol. The OD detected in the nil wells was 0.414 ± 0.06 and treatment with 10 μM danazol decreased the OD to 0.288 ± 0.037 while 100 μM to 0.162 ± 0.017, equating to percent inhibitions of 30% and 65% respectively.

[0105] At 48 hours, the inhibition observed was significant even at levels of 1 μM. The nil reading obtained after 48 hours in culture increased to 0.629 ± 0.095 and was reduced to 0.378 ± 0.037 by 1 μM, 0.241 ± 0.012 by 10 μM, and 0.19 ± 0.033 by 100 μM (or percent inhibitions of 40%, 61 %, and 70% respectively).

[0106] After 72 hours, all danazol treatments tested exhibited significant reduction in HUVEC proliferation. The OD obtained in nil wells was 1.113 ± 0.054 and after 0.1 μM treatment fell to 0.798 ± 0.037, 1 μM to 0.484 ± 0.022, 10 μM to 0.229 ± 0.016, and 100 μM to 0.156 ± 0.018 (inhibitions of 28%, 57%, 80%, and 86% respectively).

[0107] Examination of the OD obtained from all 100 μM danazol doses was consistent at all time points indicating a complete arrest of cell proliferation at this concentration.

[0108] In summary, danazol exhibited strong inhibition of endothelial cell proliferation.

B. HUVEC Tube Formation

Protocol:

[0109] To investigate the formation of capillary-like structures by HUVECs, the Angiogenesis System: Endothelial Cell Tube Formation Assay was purchased from BD Biosciences (San Jose, CA) and used according to the manufacturer's protocol. In brief, 100,000 HUVECs were seeded onto rehydrated matrigel plugs in 96 well tissue culture plates in the

presence of 5% FCS to induce tube formation. Danazol was added to final concentrations of 1 μ M, 10 μ M, or 50 μ M and LY294002 (positive control) was added at 50 μ M. After 18 hours the wells were photographed using a Kodak DCS Pro SLR/N digital camera (Rochester, NY) mounted on an inverted microscope. Ethanol treated wells were included to determine if the vehicle had any effects on cell differentiation.

Results, Observations and Discussion:

[0110] To elucidate if danazol can prevent the formation of tube-like structures by HUVEC, 96 well plates containing matrigel plugs were used. Endothelial cells when cultured in the presence of angiogenic substances and supplied with an extracellular matrix scaffold will differentiate into structures loosely resembling capillary vessels. HUVECs grown with danazol exhibited fewer organized structures with thin and less defined interconnections than controls (see Figure 2, in which A = control, B = 1 μ M danazol, C = 10 μ M danazol, D = 50 μ M danazol, and E = 50 μ M LY294002). Treatment with 50 μ M danazol led to isolated colonies of HUVEC located in the plug with very few, thin connections or vessel lumen spaces. The effect of danazol was very similar to the positive control compound LY294002. To ensure that the vehicle used had no effect, wells were treated with ethanol at concentrations corresponding to the highest dose of danazol used and no effect on tube formation was observed (data not shown). These data indicate that danazol is an effective inhibitor of tube formation at 50 μ M. Danazol had no effect on tube formation at 1 μ M or 10 μ M.

C. HUVEC Invasion

Protocol:

[0111] BioCoat Matrigel Invasion Chambers were purchased from BD Biosciences (San Jose, CA). Inserts were rehydrated at 37°C with 500 μ l HBSS for 2 hours prior to use in humidified incubator. Trypsinized HUVECs were washed twice with warm EGM-2 containing 0.1% FCS and added to the upper chamber of the invasion insert at 100,000 cells in a total volume of 250 μ l. Danazol and control compounds were added to the upper reservoir at final concentrations of 10 μ M and 100 μ M. 750 μ l EGM-2 supplemented with 5% FCS was added to the bottom chamber to initiate invasion and the plates were incubated for 24 hours. Non-invasive cells were removed from the upper chamber with moistened cotton swabs and then the inserts were washed twice with HBSS. The inserts were then submerged in 10 μ M calcein AM prepared in HBSS and incubated for 4 hours. Fluorescence was determined in a microplate reader at 485 nm excitation and 595 nm emission. LY294002 and the structurally similar but inactive compound LY303511 served as positive and negative controls respectively for this experiment.

Results:

[0112] The results are presented in Figure 3. All data is presented as representative experiment done in triplicate. Differences between subsets were analyzed using student t-test in Microsoft Excel. $P < 0.05$ was considered statistically significant.

[0113] Porous, matrigel coated inserts were used to determine if danazol can interfere with the invasion or migration of endothelial cells (Figure 3). In the system used for the study, a significant increase in cells was detected by fluorescent dye after the addition of FCS to the chamber opposite the endothelial cells ($5674 \text{ FU} \pm 77$ to 7143 ± 516). Danazol at concentrations of 10 μ M and 100 μ M had no effect, while LY294002 showed almost complete attenuation of cell invasion (5814 ± 153). These data indicate that factors present in the FCS induce the production by HUVECs of proteases that digest extracellular matrix, followed by migration along a chemotactic gradient. Danazol has no apparent inhibitory effect on invasion and migration of HUVECs in this model.

D. HUVEC Migration

Protocol:

[0114] Assays were performed to determine the effect of danazol on the migration of HUVECs in a scratch migration assay. Passage 8 HUVECs, lot number 8750 (obtained from Lonza), were plated in 6-well plates (ICS BioExpress) in endothelial growth medium-2 (EGM-2) complete medium (obtained from Lonza). The plates were cultured in a 37°C incubator with 5% CO₂ for 48-72 hours to achieve confluent monolayers. The monolayers were then "scratched" with a 1000 μ l pipet tip and washed two times with warm EGM-2 medium. The final wash medium was aspirated and replaced with fresh EGM-2 medium or fresh EGM-2 medium containing a range of concentrations of danazol concentrations (Sigma, # D8399). Photographs of the damaged monolayers were taken and the plates were incubated in a 37°C incubator with 5% CO₂ for another 24 hours. The wells were photographed again. The gaps were measured in each

photograph using Adobe Photoshop software, and gap measurements are presented as the number of pixels in the gap.

Results:

[0115] The results of three separate experiments are presented in Table 1 below. As can be seen from Table 1, danazol, at 50 μM , 75 μM and 100 μM , was found to significantly inhibit HUVEC migration in this assay. The EGM-2 culture medium used in this assay contains a cocktail of growth factors as compared to the FCS used in the Matrigel model described in section C above. This difference in the growth factors may account for the difference in the results obtained using the two models.

TABLE 1

Compound(s)	Danazol Concentration	Mean pixels	Mean % Inhibition	STD	SEM
Diluent control (ethanol)		1264.00			
Danazol	10 μM	1004.00	21.14	14.87	8.59
Danazol	25 μM	1184.00	5.50	8.80	5.08
Danazol	50 μM	895.33	27.64	17.63	10.18
Danazol	75 μM	317.33	74.62	6.80	3.93
Danazol	100 μM	178.67	85.90	0.92	0.53

Example 2: Danazol Effect on Vascular Permeability of HUVEC Monolayers

Protocol:

[0116] Assays were performed to determine the effect of danazol on permeability of HUVEC monolayers. Passage 5-10 HUVECs, lot number 7016 (obtained from Lonza), were seeded onto 1-micron-pore-size inserts located in the wells of a 24-well plate (Grcincr BioOnc 24-well Thincert cell culture inserts, #662610, or ISC BioExprcss, # T-3300-15) using endothelial growth medium-2 (EGM-2) (obtained from Lonza). The plates were cultured in a 37°C incubator with 5% CO₂ for 48-72 hours to achieve confluence and develop tight monolayers. The medium was then removed and replaced with fresh medium or fresh medium containing a range of danazol concentrations (Sigma, # D8399). Tumor necrosis factor α (TNF α Pierce Biotechnology, # RTNFA1) and interleukin-1 β (IL-1 β ;Sigma, # I-9401) were added to appropriate wells at final concentrations of 10 ng/ml each. TNF α and IL-1 β induce permeability; they can cause up to a ten-fold increase in permeability. Finally, streptavidin conjugated to horseradish peroxidase (HRP) (Pierce Biotechnology, # N100, 1.25 mg/ml) was added to each well at a final dilution of 1:250. HRP is a large molecule having a molecular weight of about 44,000. Final volumes were 300 μl in the upper chambers and 700 μl in the bottom chambers of each well. The plates were incubated for an additional 24 hours in the 37°C incubator with 5% CO₂. After this incubation, the inserts were removed and discarded. Visual examination of the cells on the inserts indicated that all of the monolayers were still intact.

[0117] To evaluate HRP flow-through, 15 μl of the resulting solutions in the bottom chambers were transferred to 96-well ELISA plates (each reaction performed in triplicate). Next, 100 μl of tetramethylbenzidine (TMB) solution (Pierce) were added to each well, and the color developed for 5 minutes at room temperature. Color development was stopped by adding 100 μl of 0.18 N acidic solution. OD was determined for each well using a microplate reader set at 450 nm minus 530 nm. The percent inhibition of permeability was calculated versus controls, and the means for three separate experiments are presented in Table 2.

[0118] As can be seen from Table 2, danazol at concentrations of 25.0 μM or higher actually increased vascular permeability. A concentration of 10.0 μM had little or no effect on vascular permeability. Danazol at concentrations from 0.1 μM to 5.0 μM , with 0.1 μM to 0.5 μM being optimal, decreased vascular permeability. The dose-response curve is very interesting as there is a second peak of inhibition at concentrations from 0.001 μM (or perhaps even lower) to 0.005 μM . Thus, danazol exhibits a very surprising and unexpected dose response curve for vascular permeability.

[0119] As shown in Example 1, a concentration of 50 μM to 100 μM would be required to obtain inhibition of HUVEC proliferation, migration and tube formation after 18-24 hours of incubation with danazol. As shown in this Example 2, these optimal concentrations for inhibiting angiogenesis would dramatically increase vascular permeability after 24 hours (see Table 2). Conversely, optimal concentrations for use to inhibit vascular permeability (0.1 μM to 0.5 μM) have insignificant effects on angiogenesis at 24 hours.

TABLE 2

Compound(s)	Danazol Concentration	Mean % Inhibition	STD	SEM
Danazol	0.001 μ M	19.35	5.39	3.11
Danazol	0.005 μ M	16.37	8.04	4.64
Danazol	0.01 μ M	-2.74	14.56	8.40
Danazol	0.05 μ M	7.67	8.83	5.10
Danazol	0.1 μ M	35.59	23.08	11.54
Danazol	0.5 μ M	30.95	12.01	6.01
Danazol	1.0 μ M	21.20	31.13	13.92
Danazol	5.0 μ M	14.63	15.30	7.65
Danazol	10.0 μ M	14.29	36.85	13.03
Danazol	25.0 μ M	-1.06	22.60	11.30
Danazol	50.0 μ M	-377.36	384.50	171.95
TNF α + IL-1 β + Danazol	0.1 μ M	31.30	25.26	12.63
TNF α + IL-1 β + Danazol	1.0 μ M	29.22	16.17	7.23
TNF α + IL-1 β + Danazol	10.0 μ M	8.47	20.45	9.14
TNF α + IL-1 β + Danazol	25.0 μ M	-39.93	15.53	7.76
TNF α + IL-1 β + Danazol	50.0 μ M	-117.16	29.20	14.60

Example 3: Danazol Effect on Vascular Permeability

[0120] Passage 9 human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA) were passaged in EGM-2 medium (Lonza, Walkersville, MD) until 80% confluence was obtained. The cells were then released from the passage flask using Trypsin-EDTA, and the cells in the resulting suspension were counted to determine both viability and cell numbers. Viability of the cell suspension was greater than 90% in this experiment.

[0121] The cells were then seeded onto inserts (1 micron pore size) located in the wells of a 24-well plate (Grcinc BioOnc 24-well Thincrt cell culture inserts, #662610) in 300 μ l EGM-2 complete medium (obtained from Lonza). Then, 700 μ l EGM-2 was placed in the bottom chamber, and the plates were cultured in a 37°C incubator with 5% CO₂ for 48 hours to achieve confluent monolayers. Transendothelial electrical resistance (TER) measurements were taken using an STX 100 electrode attached to EVOM² voltohmmeter (both from World Precision Instruments) for all inserts to confirm establishment of a semipermeable barrier. To perform the measurements, one probe was placed in each well with one electrode in the upper chamber and one in the lower chamber.

[0122] Then, the cells were treated in duplicate as follows. EGM-2 medium was carefully decanted from the inserts and replaced with IMDM medium containing 0.5% fetal bovine serum and EGM-2 supplements, except for VEGF and hydrocortisone (all from Lonza). In some wells, the IMDM medium contained danazol (Sigma, # D8399) in a ten-fold serial dilution. The plates were incubated in a 37°C incubator at 5% CO₂ for four hours before 30 μ l of a solution containing 4% fluorescent-labelled human serum albumin was added to the upper chamber of each well. The plates were incubated in a 37°C incubator with 5% CO₂ for an additional 18 hours.

[0123] After this incubation, the inserts were removed and discarded, and 200 μ l of the medium from the bottom chamber was transferred to 96-well black fluoro-plates (Falcon) in triplicate. The fluorescence of each well was then measured at an excitation wavelength of 340 nm and an emission wavelength of 470 nm. Mean fluorescence units (FU) for each insert were then calculated, and duplicate readings were averaged. The results are presented in Table 3.

TABLE 3

Danazol Concentration	Mean FU	STD
None	767.13	8.38
0.01 μ M	688.50	14.94

(continued)

Danazol Concentration	Mean FU	STD
0.1 μ M	743.90	8.95
1.0 μ M	783.39	14.59
10.0 μ M	768.99	18.85

[0124] As can be seen, the lowest concentration of danazol (0.01 μ M) gave the greatest inhibition (about 10%). Control wells run with no cells gave over 4000 FU in the lower chamber, showing that the retinal endothelial monolayers were selectively permeable.

Example 4: Effect of Danazol on TER of Three Different Endothelial Cell Monolayers

[0125] Assays were performed to determine the effect of danazol on transendothelial electrical resistance (TER) of human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA). To do so, 150,000 passage 14 human retinal endothelial cells were seeded onto inserts (1 micron pore size) located in the wells of a 24-well plate (Greiner BioOne 24-well Thincert cell culture inserts, #662610) in 300 μ l EGM-2 complete medium (obtained from Lonza). Then, the plates were cultured in a 37°C incubator with 5% CO₂ for 24 hours. After the incubation, the culture medium was carefully decanted and replaced with either fresh EGM-2 or fresh EGM-2 containing danazol at a final concentration of 1 μ M. The plates were placed back in the incubator and cultured for an additional 144 hours. Assays were also performed in the same manner using passage 8 human brain endothelial cells and passage 8 human umbilical vein endothelial cells.

[0126] An initial TER measurement was taken for each insert using EVOM² voltohmmeter connected to an STX100 electrode (both from World Precision Instruments). Measurements were also taken at 24, 48, 72 and 144 hours. The results are presented in Tables 4, 5 and 6 below. All data are presented as TER measurements/cm² of insert with TER of blank inserts subtracted.

TABLE 4

Human Retinal Endothelial Cells					
Danazol Concentration	0 Hours	24 Hours	48 Hours	72 Hours	144 Hours
None	32.3	96.0	144.4	148.0	219.7
1.0 μ M	21.7	132.3	182.3	217.7	234.8

TABLE 5

Human Brain Endothelial Cells					
Danazol Concentration	0 Hours	24 Hours	48 Hours	72 Hours	144 Hours
None	41.4	115.7	176.3	154.0	151.5
1.0 μ M	32.3	139.9	188.4	149.5	125.8

TABLE 6

Human Umbilical Vein Endothelial Cells					
Danazol Concentration	0 Hours	24 Hours	48 Hours	72 Hours	144 Hours
None	82.3	217.2	276.3	226.8	227.3
1.0 μ M	70.2	246.0	364.1	270.7	286.4

[0127] As can be seen, danazol enhanced TER measurements (reduced ion permeability) in the retinal and umbilical vein endothelial cell monolayers. Danazol did not appear to have much effect on the TER of the brain endothelial cell monolayers, except at the earliest time point. TER is a measurement of the electrical resistance across cellular monolayers.

ayers. It is an indication of barrier integrity and correlates with ion permeability.

Example 5: Danazol Effect on Akt Phosphorylation

[0128] Assays were performed to determine the effect of danazol on phosphorylation of Akt in human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA). The cells were grown in a 25 cm² flask to near confluence in EGM-2 medium (Lonza, Walkersville, MD) containing 2% fetal calf serum (Lonza). The cells were then released from the passage flask using Trypsin/EDTA. The cells in the resulting suspension were counted and seeded on a 96-well plate at 1×10^4 cells/well in EGM-2 medium. The plate was incubated at 37°C with 5% CO₂ for 24 hours. Then, 200 µl of either EGM-2 medium (control) or various concentrations of danazol were added, and the plates were incubated for an additional 2 hours. After this incubation, the cells were fixed immediately with 4% formaldehyde, refrigerated, and the extent of phosphorylation of Akt determined using the Akt Cellular Activation of Signaling ELISA Kit (CASE™ Kit for AKT S473; SABiosciences, Frederick, MD) following the manufacturer's protocols. The CASE™ Kit for AKT S473 quantifies the amount of activated (phosphorylated) Akt protein relative to total Akt protein in parallel assays using a conventional ELISA format with colorimetric detection. The Akt phosphorylation site is serine 473 and is recognized by one of the antibodies used in one of the two parallel assays to provide a measure of activated Akt protein. The other antibody used in the other parallel assay recognizes Akt to provide a measure of total Akt protein. Both primary antibodies are detected using a horseradish peroxidase-labeled secondary antibody. Addition of the manufacturer's Developing Solution for 10 minutes, followed by addition of the manufacturer's Stop Solution, produces the result which can be measured colorimetrically.

[0129] The results are presented in Table 7 below. As can be seen there, all of the concentrations of danazol caused an increase in Akt phosphorylation (activation).

TABLE 7

TREATMENT	PERCENT INCREASE IN AKT PHOSPHORYLATION VERSUS CONTROL	STANDARD DEVIATION
0.5 µM danazol	73.8%	92.9%
1.0 µM danazol	66.7%	11.7%
2.0 µM danazol	101.6%	9.1%
5.0 µM danazol	40.5%	17.7%
10.0 µM danazol	115.3%	112.9%
20.0 µM danazol	161.3%	128.7%
50.0 µM danazol	98.6%	61.2%

[0130] It is believed that these results provide a possible explanation for the vascular permeability dose response curve obtained in Example 2. As shown in Example 2, low doses of danazol reduced permeability, while high doses increased permeability. It is believed that a certain level of phosphorylation of Akt at S473 reduces permeability (the 0.5-5.0 µM concentrations in this experiment), while hyperphosphorylation of Akt at S473 causes increased permeability (the 10-50 µM concentrations in this experiment).

Example 6: Effect of Danazol and Steroid Receptor Antagonists on TER of Retinal Endothelial Cell Monolayers

[0131] Assays were performed to determine the effect of danazol and steroid receptor antagonists on transendothelial electrical resistance (TER) of human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA). To do so, Greiner tissue culture well inserts (Greiner BioOne 24-well Thincert cell culture inserts, #662610) were coated with 5 µg/cm² fibronectin (Sigma). Then, passage 12 human retinal endothelial cells were seeded into the upper chamber of the wells at 120,000 cells per insert in a volume of 300 µl of EGM-2 medium (Lonza). The volume for the lower chamber was 700 µl of EGM-2 medium (Lonza). The plates were cultured in a 37°C incubator with 5% CO₂ for 48 hours to establish intact monolayers. At the end of the incubation, TER measurements were taken using an STX 2 probe attached to EVOM² voltohmmeter (both from World Precision Instruments) for all inserts to confirm integrity of the endothelial barrier. All inserts exhibited elevated resistance as compared to inserts without cells.

[0132] Then, the culture medium was carefully decanted and replaced with fresh EGM-2, with and without several additives. The additives were danazol, hydroxyflutamide (androgen receptor antagonist), fluestrant (estrogen antagonist) and PI3 kinase inhibitor LY 294002 (control). Stock solutions of all additives, except danazol, were made at 10 mM

in DMSO. The danazol stock solution was 10 mM in ethanol. Working 200 μ M dilutions of all additives were made in same solvents. Then, 200 nM dilutions of each additive, and of equivalent dilutions of ethanol and DMSO (controls), were made in EGM-2 medium, and danazol and each of the other additives or medium (control) were added to the wells in the combinations and final concentrations shown in the table below. The plates were then placed back into the incubator, and TER measurements were taken as described above for each insert at 30 minutes, 60 minutes, 120 minutes and 24 hours. TER was calculated by subtracting the background measurement (empty insert) from the reading of an insert and dividing by the surface area of the insert (0.33 cm²). The results are presented in Table 8 below.

TABLE 8

Treatment	TER at 30 minutes	TER at 60 minutes	TER at 120 minutes	TER at 24 Hours
None	216.22	249.25	234.23	312.31
0.1 μ M Danazol	255.26	267.27	249.35	366.37
0.1 μ M Hydroxyflutamide	177.18	186.19	201.20	297.30
0.1 μ M Fluevestrant	228.23	270.27	258.26	336.34
0.1 μ M Hydroxyflutamide followed by 0.1 μ M Danazol	237.24	276.28	240.24	363.36
0.1 μ M Fluevestrant followed by 0.1 μ M Danazol	195.20	309.31	255.26	393.39
10.0 μ M LY294002	297.30	354.35	276.28	345.35
10.0 μ M LY294002 followed by 0.1 μ M Danazol	243.24	342.34	270.27	336.34

[0133] As can be seen from Table 8, danazol and fluevestrant increased the TER measurements (reduced permeability), while hydroxyflutamide reduced the readings (increased permeability), compared to the control (no treatment). Danazol prevented the reduction caused by hydroxyflutamide. This could be evidence that danazol is occupying the androgen receptor in these cells. Danazol and fluevestrant showed additive results at some time points.

Example 7: Effect of Danazol on Actin Stress Fiber Formation

[0134] The IEJs of the paracellular pathway include AJs and TJs. The actin cytoskeleton is bound to each junction and control the integrity of the junctions through actin remodeling. Reorganization of the actin filaments into stress fibers results in application of mechanical forces to the junctions that pull them apart, cause cellular contraction and changes in morphology. The process of actin polymerization is very dynamic, which allows for the rapid reorganization of actin structures and the transition from the quiescent phenotype, characterized by thick cortical actin ring and the absence of stress fibers, to the activated cell phenotype with thin or no cortical actin and abundant stress fibers. The actin cytoskeleton appears also to be involved in transcytosis, perhaps by regulating the movement of caveolae.

[0135] Human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA) were seeded into Falcon Optilux assay plates (BD Biosciences) at 1000 cells per well in a total volume of 200 μ l of EGM-2 medium (Lonza). The plates were cultured in a 37°C incubator with 5% CO₂ for 48 hours. Then, the medium was removed and replaced with 200 μ l of IMDM medium supplemented with 0.1% fetal bovine serum (all from Lonza), and the cells were cultured under these growth factor and serum starved conditions for one hour to suppress actin polymerization. Then danazol (0.1 μ M or 10 μ M final concentrations) or the PI3 kinase inhibitor LY294002 (10 μ M final concentration) (positive control) were added. Immediately following addition of these compounds, TNF α (final concentration of 50 ng/ml) was added. After incubation for 30 minutes in a 37°C incubator with 5% CO₂, the medium was aspirated, and the cells were fixed with 3.6% formaldehyde in phosphate buffered saline (PBS) for ten minutes at room temperature. All wells were then washed two times with 100 μ l PBS. The cells were permeabilized using a 0.1 % Triton X-100 in PBS for 5 minutes. All wells were then washed two times with 100 μ l PBS, and 50 μ l of a 1:40 dilution of rhodamine-phalloidin (Invitrogen) in PBS was added to the cells to stain for F-actin and left on the cells for 20 minutes at room temperature. All wells were then washed two times with 100 μ l PBS. Then 100 μ l PBS was added to each well and the cells were observed and photographed using an inverted microscope with rhodamine filters (ex530/em590).

[0136] The results showed that danazol affected the ability of stress fibers to develop. When treated with danazol, the cells exhibited different staining patterns, dependent on the dosage. At the lower danazol dose (0.1 μ M), diffuse staining throughout the cytoplasm was observed, possibly indicative of a stabilizing event or of a resting phenotype. At the higher

danazol dose (10.0 μ M), stress fibers with multiple focal points were detected. These findings correlate with previous results (see previous examples) that lower danazol doses inhibit permeability and higher danazol doses increase permeability. TNF α stimulated the cells and led to strong stress fiber development with intensely staining focal points. Danazol and LY294002 decreased the number of cells exhibiting stress fiber development with TNF α .

Example 8: Effect of Danazol on Actin Stress Fiber Formation

[0137] Human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA) were seeded into Falcon Optilux assay plates (BD Biosciences) coated with 1 μ g/cm² fibronectin at 3000 cells per well in a total volume of 200 μ l of EGM-2 medium (Lonza). The plates were cultured in a 37°C incubator with 5% CO₂ for 48 hours. Then, the medium was removed and replaced with 200 μ l of Ultraculture medium supplemented with 2.0% fetal bovine serum (all from Lonza), and the cells were cultured under these growth factor and serum starved conditions overnight to suppress actin polymerization. Then, the medium was removed and replaced with fresh Ultraculture medium supplemented with 2.0% fetal bovine serum containing danazol (0.1 μ M, 1 μ M or 10 μ M) or the PI3 kinase inhibitor LY294002 (10 μ M) (positive control). After incubation with these compounds for 30 minutes in a 37°C incubator with 5% CO₂, vascular endothelial growth factor (VEGF) (final concentration of 25 ng/ml) was added. After incubation for an additional 30 minutes in a 37°C incubator with 5% CO₂, the medium was aspirated, and the cells were fixed using 3.6% formaldehyde in phosphate buffered saline (PBS) for ten minutes at room temperature. All wells were then washed two times with 100 μ l PBS. The cells were permeabilized using a 0.1% Triton X-100 in PBS for 5 minutes. All wells were then washed two times with 100 μ l PBS, and 50 μ l of a 1:40 dilution of rhodamine-phalloidin (Invitrogen) in PBS was added to the cells to stain for F-actin and left on the cells for 20 minutes at room temperature. All wells were then washed two times with 100 μ l PBS. To counter-stain for nuclei, 100 μ l of a 3 μ M DAPI (4,6-diamidino-2-phenylindole, dilactate (Invitrogen)) solution was added to each well. After 5 minutes, the cells were washed two times with 100 μ l PBS. Then 100 μ l PBS was added to each well and the cells were observed and photographed using an inverted microscope using rhodamine (ex530/em590) and DAPI (ex350/em460) filters.

[0138] The results showed that danazol affected the ability of stress fibers to develop. When treated with danazol, the cells exhibited different stress fiber formation patterns, dependent on the dosage applied. At the lowest danazol dose (0.1 μ M), diffuse F-actin staining throughout the cytoplasm was observed. At 1.0 μ M danazol, the diffuse staining persisted, but stress fibers and focal points around the perimeter of most cells were visible. At the highest danazol dose (10.0 μ M), there was no longer any diffuse staining, and stress fiber development and focal points were seen. The staining seen with the lower doses of danazol exhibited a perinuclear staining pattern, indicating microtubule stabilization similar to that observed with paclitaxel (a Taxol compound known to stabilize and polymerize microtubules). With VEGF, there was strong stress fiber development. Danazol changed the VEGF pattern in a dose-dependent manner: (i) the lowest 0.1 μ M danazol dose made the stress fibers less pronounced and some diffuse staining appeared; (ii) the 1.0 μ M dose showed fewer thick stress fibers, but focal points were seen on contact surfaces; and (iii) the highest 10.0 μ M danazol dose showed strong stress fiber development with focal points. LY294002 prevented the strong stress fiber development seen with VEGF and exhibited diffuse staining.

Example 9: Effect of Danazol on Vascular Endothelial Cadherin (VE-Cadherin) Phosphorylation

[0139] Passage 12 human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA) were grown to confluence on fibronectin-coated (1 μ g/cm²) 10 cm² tissue culture plates using EGM-2 culture medium (Lonza) in a 37°C incubator with 5% CO₂. When complete confluence was achieved, the medium was replaced with Ultraculture medium supplemented with 0.5% fetal bovine serum and L-glutamine (all from Lonza), and the cells were cultured under these growth factor and serum starved conditions for 24 hours. Then, the medium was removed and replaced with fresh Ultraculture medium supplemented with 0.5% fetal bovine serum and L-glutamine containing danazol (0.1 μ M, 1 μ M or 10 μ M) or ethanol (vehicle control). After incubation for 15 minutes in a 37°C incubator with 5% CO₂, vascular endothelial growth factor (VEGF) (final concentration of 50 ng/ml) was added, and the plates incubated for an additional 15 minutes in a 37°C incubator with 5% CO₂.

[0140] The plates were immediately treated to lyse the cells as follows. PBS and the lysis buffer (PBS containing 1% Triton X-100 supplemented with phosphatase inhibitor solutions 1 and 2 (Sigma), protease inhibitor (Sigma) and sodium orthovanadate at a final concentration of 2 mM) were cooled to 4°C. The cells were washed two times with 5 ml of the ice cold PBS and then lysed in 500 μ l of the ice cold lysis buffer. The resulting protein extracts were transferred to 1.7 ml microcentrifuge tubes, and cell debris was removed by spinning at 4°C at 10,000 rpm for 10 minutes. Then, 450 μ l of the cleared solution was transferred to tubes containing 25 μ l of Protein Dynabeads (Invitrogen) coated with 10 μ l C19 anti-VE cadherin polyclonal antibody (Santa Cruz Biotechnology) (coating performed following manufacturer's protocol). The extracts and beads were then incubated overnight at 4°C on an orbital shaker to capture VE cadherin from the extracts. The beads were then washed four times with ice cold lysis buffer. To release the protein from the

beads, they were heated for 10 minutes at 75°C in SDS loading dye containing 20% reducing dye (Invitrogen).

[0141] The released proteins were separated in 4-20% polyacrylamide gels (Invitrogen) at 120 volts for 1 hour. To determine phosphorylation and total protein in the gels, Pro-Q diamond (Invitrogen) and SYPRO ruby (Invitrogen) protein staining were sequentially performed following the manufacturer's protocol. The gels were photographed and densitometry performed using a Kodak imaging station. The results are presented in Table 9 below.

TABLE 9

VE-Cadherin				
	Nil (ethanol control)	0.1 μ M danazol	VEGF	0.1 μ M danazol followed by VEGF
Relative intensity - ProQ results (phosphorylated protein)	1.00	1.51	1.89	1.38
Relative intensity - SYPRO results (total protein)	1.00	0.89	0.84	0.83
Ratio phosphorylated: total protein	0.215	0.365 (1.70 fold increase)	0.481 (2.24 fold increase)	0.358 (1.66 fold increase)

[0142] As can be seen, danazol caused an increase in VE-cadherin phosphorylation. VEGF caused an even greater increase in VE-cadherin phosphorylation (hyperphosphorylation), which was reversed by danazol. VE-cadherin is a component of AJs, and phosphorylation of VE-cadherin can have a variety of effects depending on the residue. In general, tyrosine phosphorylation of VE-cadherin leads to AJ disassembly and increased permeability. Serine 665 phosphorylation, however, causes a rapid but reversible internalization of VE-cadherin associated with reduced barrier function. A feedback loop appears to exist in which internalized VE-cadherin drives an increase in cytoplasmic p120, a scaffolding protein that complexes to AJs. This up-regulation induces a decrease in active RhoA in association with an increase in the barrier-stabilizing GTPases like Rac1, Rap-1, and Cdc42. It is believed that the increase in VE-cadherin phosphorylation observed in this experiment following low dose danazol treatment leads to the activation of barrier stabilizing GTPases. In addition, danazol may prevent the destabilizing phosphorylation events induced by VEGF.

Example 10: Effect of Danazol and Steroid Receptor Antagonists on TER of Retinal Endothelial Cell Monolayers

[0143] Assays were performed to determine the effect of danazol and steroid receptor antagonists on transendothelial electrical resistance (TER) of human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA). To do so, Greiner tissue culture well inserts (Greiner BioOne 24-well Thincert cell culture inserts, #662610) were coated with 5 μ g/cm² fibronectin. Then, passage 13 human retinal endothelial cells were seeded into the upper chamber of the wells at 120,000 cells per insert in a volume of 300 μ l of EGM-2 medium (Lonza). The volume for the lower chamber was 700 μ l of EGM-2 medium (Lonza). The plates were cultured in a 37°C incubator with 5% CO₂ for 48 hours to establish intact monolayers. At the end of the incubation, TER measurements were taken using an STX 2 probe attached to EVOM² voltohmmeter (both from World Precision Instruments) for all inserts to confirm integrity of the endothelial barrier. All inserts exhibited elevated resistance as compared to inserts without cells.

[0144] Then, the culture medium was carefully decanted and replaced with fresh EGM-2, with and without several additives. The additives were danazol, hydroxyflutamide (androgen receptor antagonist), fluvastatin (estrogen antagonist), testosterone, estradiol and PI3 kinase inhibitor LY 294002 (control). Stock solutions of all additives, except danazol, were made at 10 mM in DMSO. The danazol stock solution was 10 mM in ethanol. Working 200 μ M dilutions of all additives were made in same solvents. Then, 200 nM dilutions of each additive, and of equivalent dilutions of ethanol and DMSO (controls), were made in EGM-2 medium, and danazol and each of the other additives or medium (control) were added to the wells in the combinations and final concentrations shown in the table below. The plates were then placed back into the incubator, and TER measurements were taken as described above for each insert at 5 minutes, 30 minutes, 60 minutes and 24 hours. TER was calculated by subtracting the background measurement (empty insert) from the reading of an insert and dividing by the surface area of the insert (0.33 cm²). The results are presented in Table 10 below.

[0145] As can be seen from Table 10, danazol increased the TER measurements, hydroxyflutamide reduced the

readings, testosterone reduced the readings very slightly, and fluvestrant had essentially no effect, compared to the control (no treatment). Danazol prevented the reduction caused by hydroxyflutamide and the very slight reduction seen with testosterone. As with the results in Example 6, this could be evidence that danazol is occupying the androgen receptor in these cells.

TABLE 10

Treatment	TER at 5 minutes	TER at 30 minutes	TER at 60 minutes	TER at 24 Hours
None	250.30	262.31	251.00	287.09
0.1 μ M Danazol	280.03	311.56	313.06	348.35
0.1 μ M Hydroxyflutamide	190.44	207.46	215.97	267.27
0.1 μ M Hydroxyflutamide followed by 0.1 μ M Danazol	230.48	275.53	262.01	312.31
0.1 μ M Fluvestrant	223.47	251.50	243.99	279.28
0.1 μ M Fluvestrant followed by 0.1 μ M Danazol	219.47	279.53	273.02	343.34
10 nM Testosterone	257.51	240.49	225.98	267.27
100 nM Testosterone followed by 0.1 μ M Danazol	273.52	287.54	259.01	283.28
10 nM Estradiol	246.50	245.50	250.00	328.33
10 nM Estradiol followed by 0.1 μ M Danazol	276.53	307.56	282.03	363.36

Example 11: Effect of Danazol on Actin Stress Fiber Formation

[0146] Passage 6 human renal glomerular microvascular endothelial cells (ACBRI 128, Cell Systems Corporation (exclusive distributor for Applied Cell Biology Research Institute), Kirkland, WA) and passage 12 human retinal endothelial cells (ACBRI 181, Cell Systems Corporation (exclusive distributor for Applied Cell Biology Research Institute), Kirkland, WA) were seeded into 16-chamber glass slides coated with 5 μ g/cm² fibronectin at 2000 cells per well in a total volume of 200 μ l of EGM-2 medium (Lonza). The plates were cultured in a 37°C incubator with 5% CO₂ for 48 hours with daily medium changes. Then, the test compounds (danazol, TNF α and S1P), diluted in Hanks Balanced Salt Solution (HBSS; Lonza), were added to give the following final concentrations: danazol (1 μ M) (Sigma), TNF α (1 ng/ml) (Sigma), and S1P (1 μ M) (Sigma). The slides were incubated with the test compounds for 15 minutes, 30 minutes or 24 hours in a 37°C incubator with 5% CO₂. After this incubation, the medium was aspirated, and the cells were fixed using 3.6% formaldehyde in phosphate buffered saline (PBS) for ten minutes at room temperature. All wells were then washed two times with 100 μ l PBS. The cells were permeabilized using a 0.1% Triton X-100 in PBS for 5 minutes. All wells were then washed two times with 100 μ l PBS, and 50 μ l of a 1:40 dilution of rhodamine-phalloidin (Invitrogen) in PBS was added to the cells to stain for F-actin and left on the cells for 20 minutes at room temperature. All wells were then washed two times with 100 μ l PBS. Then 100 μ l PBS was added to each well and the cells were observed and photographed using an inverted microscope using a rhodamine (ex530/em590) filter.

[0147] The results showed that danazol affected the ability of stress fibers to develop in the renal glomerular microvascular endothelial cells. When treated with danazol alone, the cells exhibited perinuclear staining at 15 minutes, diffuse staining throughout the cells with ruffled edges on many of the cells at 3 hours, and staining similar to untreated controls at 24 hours. With TNF α alone, stress fibers were seen at all times, with the number of cells exhibiting stress fibers and the thickness of the fibers increasing with time. Danazol decreased the stress fiber formation and the thickness of the fibers at all times, and cortical actin rings and ruffled edges were visible beginning at 3 hours. Cells treated with S1P alone showed actin cortical rings, with development beginning at 15 minutes and strongest at 3 hours. The cells were returning to morphology similar to untreated controls at 24 hours. Danazol seemed to enhance the cortical rings. Also, diffuse staining was observed, especially at 15 minutes and 24 hours.

[0148] For the retinal endothelial cells treated with danazol alone, the cells exhibited perinuclear staining at 15 minutes, diffuse staining throughout the cells with ruffled edges on many of the cells at 3 hours, and staining similar to untreated controls at 24 hours. With TNF α alone, stress fibers were seen at all times, with the number of cells exhibiting stress fibers and the thickness of the fibers increasing from 15 minutes to 3 hours and being reduced after 24 hours of incubation.

Danazol decreased the stress fiber formation and/or the thickness of the fibers at all times. Diffuse staining was observed at 15 minutes and 24 hours, and cortical actin rings were visible at 3 hours. Cells treated with S1P alone showed actin cortical rings, with development beginning at 15 minutes and strongest at 3 hours. The cells were returning to morphology similar to untreated controls at 24 hours. Danazol seemed to enhance the cortical rings at 3 hours. Also, diffuse staining was observed, especially at 15 minutes and 24 hours.

[0149] S1P (sphingosine-1 phosphate) plays a very important function in the formation and maintenance of vascular endothelium. S1P is a constitutive signaling input that facilitates the organization and barrier function of the vascular endothelium through its effects on the actin cytoskeleton. In particular, S1P is involved in the formation of cortical actin fibers and organization of the adherens junctions. Depletion of S1P leads to vascular leak and edema, and S1P can reverse endothelial dysfunction and restore barrier function.

[0150] In this experiment, danazol exhibited an ability to strengthen the protective effects of S1P in both retinal and glomerular endothelial cells. Danazol also reversed the formation of stress fibers induced by TNF α in both of these types of endothelial cells. Diffuse perinuclear staining is seen in cells treated with danazol alone.

Example 12: Effect of Danazol on ECIS

[0151] Assays were performed to determine the effect of danazol on transendothelial electrical resistance (TER) of human renal glomerular microvascular endothelial cells (ACBRI 128, Cell Systems Corporation (exclusive distributor for Applied Cell Biology Research Institute), Kirkland, WA) or human retinal endothelial cells (ACBRI 181, Cell Systems Corporation (exclusive distributor for Applied Cell Biology Research Institute), Kirkland, WA). Electrical resistance was measured using the electric cell-substrate impedance sensing (ECIS) system (ECISZ θ , obtained from Applied Biophysics) with 8-well multiple electrode plates (8W10E). Each well of the plates was coated with 5 μ g/cm² fibronectin in HBSS by adding the fibronectin in a volume of 100 μ l per well and incubating the plates for 30 minutes in a 37°C incubator with 5% CO₂. The fibronectin solution was removed, and 400 μ l of EGM-2 culture medium (Lonza) was added to each well. The plates were connected to the ECISZ θ system and were electrically stabilized. The EGM-2 medium was aspirated and replaced with 200 μ l of EGM-2 culture medium containing 100,000 cells per well. The plates were reconnected to the ECISZ θ system and incubated for 24 hours in a 37°C incubator with 5% CO₂. The EGM-2 medium was aspirated and replaced with 400 μ l of fresh EGM-2 culture medium per well. The plates were reconnected to the ECISZ θ system and incubated for 6 hours in a 37°C incubator with 5% CO₂. Concentrated solutions of the test compounds in HBSS were prepared and placed in the incubator to equilibrate. The test compounds were then added to appropriate wells at the following final concentrations: danazol (1 μ M) (Sigma) and S1P (1 μ M) (Sigma). ECIS (resistance) was monitored for 90 hours.

[0152] In the retinal endothelial cells, 1.0 μ M danazol alone showed an increase of ECIS as compared to untreated cells starting about 1.5-2.0 hours after treatment and persisting for 5 hours. S1P alone showed an increase of ECIS as compared to untreated cells which started within the first 15 minutes after treatment and persisted for about 3 hours. Danazol and S1P in combination increased the ECIS as compared to S1P alone and untreated cells, and this increased ECIS persisted for about 90 hours. Thus, danazol exhibited an ability to enhance the early effects of S1P and to maintain a higher resistance throughout the experiment when S1P was present.

[0153] Glomerular endothelial cells exhibited a different pattern. Danazol alone had no effect on ECIS until about 30 hours after treatment. Danazol alone increased ECIS compared to untreated cells from about 30 to about 90 hours, with the greatest increase occurring between about 60-90 hours. S1P alone also had no effect on ECIS until about 30 hours after treatment. S1P alone increased ECIS compared to untreated cells from about 30 to about 60 hours. The combination of danazol and S1P had no effect on ECIS until about 30 hours after treatment. This combination increased ECIS compared to untreated cells, S1P alone and danazol alone. In particular, the combination increased ECIS compared to untreated cells from about 30 to about 70 hours, increased ECIS compared to S1P alone from about 30 to 75 hours, and increased ECIS compared to danazol alone from about 30 to about 50 hours.

Example 13: Effect of Danazol on ECIS

[0154] Assays were performed to determine the effect of danazol on transendothelial electrical resistance (TER) of human renal glomerular microvascular endothelial cells (ACBRI 128, Cell Systems Corporation (exclusive distributor for Applied Cell Biology Research Institute), Kirkland, WA). Electrical resistance was measured using the electric cell-substrate impedance sensing (ECIS) system (ECISZ θ , obtained from Applied Biophysics) with 8-well multiple electrode plates (8W10E). Each well of the plates was coated with 5 μ g/cm² fibronectin in HBSS by adding the fibronectin in a volume of 50 μ l per well and incubating the plates for 30 minutes in a 37°C incubator with 5% CO₂. The fibronectin solution was removed, and 200 μ l of EGM-2 culture medium (Lonza) was added to each well. The plates were connected to the ECISZ θ system and were electrically stabilized. The EGM-2 medium was aspirated and replaced with 200 μ l of EGM-2 culture medium containing 40,000 passage 6 cells per well. The plates were reconnected to the ECISZ θ system

and incubated for 24 hours in a 37°C incubator with 5% CO₂. The EGM-2 medium was aspirated and replaced with 200 µl of fresh EGM-2 culture medium per well. The plates were reconnected to the ECISZ0 system and incubated for an additional 24 hours in a 37°C incubator with 5% CO₂. The EGM-2 medium was aspirated and replaced with 200 µl of fresh EGM-2 culture medium without dexamethasone per well. The plates were reconnected to the ECISZ0 system and incubated overnight in a 37°C incubator with 5% CO₂. Finally, the EGM-2 medium was aspirated and replaced with 200 µl of fresh EGM-2 culture medium without dexamethasone per well. The plates were reconnected to the ECISZ0 system and incubated 2 hours in a 37°C incubator with 5% CO₂. Concentrated solutions of the test compounds in HBSS were prepared and placed in the incubator to equilibrate. The test compounds were then added to appropriate wells at the following final concentrations: danazol (1 µM) (Sigma) and dexamethasone (1 µM) (Sigma). ECIS (resistance) was monitored for 90 hours.

[0155] Danazol alone increased ECIS compared to untreated cells beginning at about 3 hours and persisting for about 90 hours. The increase was greatest from about 12 to about 15 hours. When compared to dexamethasone, danazol exhibited a similar pattern, but the enhancement of ECIS (TER) was not as great.

Example 14: Effect of Danazol on RhoA

[0156] Remodeling of the endothelial cell cytoskeleton is central to many functions of the endothelium. The Rho family of small GTP-binding proteins have been identified as key regulators of F-actin cytoskeletal dynamics. The Rho family consists of three isoforms, RhoA, RhoB and RhoC. The activation of RhoA activity leads to prominent stress fiber formation in endothelial cells. Stimulation of endothelial cells with thrombin increases Rho GTP and myosin phosphorylation, consistent with increased cell contractility. Inhibition of RhoA blocks this response and the loss of barrier function, demonstrating a critical role for Rho in vascular permeability.

[0157] This experiment was performed using a commercially-available Rho activation assay (GLISA) purchased from Cytoskeleton, Denver, Colorado, following the manufacturer's protocol. Briefly, passage 8 or 12 human retinal endothelial cells (ACBRI 181, Applied Cell Biology Research Institute, Kirkland, WA) were cultured on fibronectin-coated (1 µg/cm²) 6-well tissue culture plates using EGM-2 culture medium (Lonza) for 24 hours in a 37°C incubator with 5% CO₂ (30,000 cells /well in total volume of 3 ml). Then, the medium was aspirated and replaced with Ultraculture medium supplemented with 0.1% fetal bovine serum, L-glutamine, sodium pyruvate, penicillin/streptomycin and ITSS (insulin, transferrin sodium selenium) (all from Lonza) to serum starve the cells and reduce the background level of RhoA. The cells were cultured for 24 hours in a 37°C incubator with 5% CO₂. Test compounds diluted in HBSS were placed in the incubator to equilibrate before addition to the cells. Then, 150 µl of each test compound was added to the appropriate culture wells, and the plates were incubated in the incubator for an additional 15 minutes. Then, thrombin was added to appropriate wells. After 1 minute, the cells were washed one time with 1.5 ml phosphate buffered saline and were then lysed with 100 µl GLISA lysis buffer supplemented with protease inhibitors. The extracts were scraped, transferred to microcentrifuge tubes and transferred to ice to preserve the active form of RhoA. All extracts were then cleared of debris by spinning at 10,000 rpms for 2 minutes at 4°C. The supernatants were transferred to new tubes and placed back on ice. Aliquots of each extract were removed for the GLISA assay and for protein determinations. All protein concentrations were within 10%, and the extracts were used at the achieved concentrations (equates to 15 µg total protein per well). The GLISA assay was performed using the reagents supplied in the kit.

[0158] The results for the passage 12 retinal endothelial cells are presented in Table 11 below. As expected, the active Rho A levels induced by thrombin were very high. All of the test compounds inhibited the thrombin-induced activation of Rho A.

[0159] The results for the passage 8 retinal endothelial cells are presented in Table 12 below. As expected, the active Rho A levels induced by thrombin were very high. All of the test compounds inhibited the thrombin-induced activation of Rho A.

TABLE 11

Treatment	Mean OD	Percent Inhibition vs. Untreated Control	Percent Inhibition vs. Thrombin
Untreated	0.455	---	---
1.0 µM Danazol	0.424	6.82	---
1.0 µM Dexamethasone	0.428	5.83	---
10.0 µM PI3 kinase inhibitor LY 294002	0.370	18.70	---
1.0 µM Src-1 Inhibitor*	0.349	23.21	---

(continued)

Treatment	Mean OD	Percent Inhibition vs. Untreated Control	Percent Inhibition vs. Thrombin
0.1 U/ml Thrombin	1.013	---	---
0.1 U/ml Thrombin + 1.0 μ M Danazol	0.859	---	27.57
0.1 U/ml Thrombin + 1.0 μ M Dexamethasone	0.826	---	33.48
0.1 U/ml Thrombin + 10.0 μ M PI3 kinase inhibitor LY294002	0.685	---	58.73
0.1 U/ml Thrombin + 1.0 μ M Src-I Inhibitor	0.534	---	85.85
* Obtained from Sigma.			

TABLE 12

Treatment	Mean OD	Percent Inhibition vs. Untreated Control	Percent Inhibition vs. Thrombin
Untreated	0.102	---	---
1.0 μ M Danazol	0.027	73.89	---
10.0 μ M PI3 kinase inhibitor LY 294002	0.056	45.32	---
0.1 U/ml Thrombin	0.561	---	---
0.1 U/ml Thrombin + 1.0 μ M Danazol	0.373	---	41.02
0.1 U/ml Thrombin + 10.0 μ M PI3 kinase inhibitor LY294002	0.433	---	27.86

Example 15: Animal Model Of Vascular Hyperpermeability

[0160] New Zealand white rabbits received 0.215 mg/kg of danazol orally twice per day for 7 days. The rabbits were then injected once intravitreally with vascular endothelial growth factor A (VEGF-A) to produce vascular leakage in the retina. Then, 24 hours later, fluorescein sodium was injected, and the fluorescence of the eyes was measured using a Fluorotron (Ocumetrics) (five measurements averaged). A single control (placebo) rabbit had 250 fluorescence units in the retina, indicating vascular leakage there. A single danazol-treated rabbit gave 16 fluorescence units, which represents a 94% reduction in vascular leakage caused by the danazol.

Claims

1. A vascular-hyperpermeability-inhibiting amount of a danazol compound, for use for inhibiting vascular hyperpermeability in an animal in need thereof.
2. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to claim 1 for treating a disease or condition mediated by vascular hyperpermeability.
3. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 2 wherein the disease or condition is a vascular complication of diabetes.
4. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 3 wherein the

vascular complication of diabetes is diabetic macular edema, diabetic retinopathy, diabetic nephropathy, diabetic neuropathy, edema, diabetic cardiomyopathy, accumulation of low density lipoproteins in subendothelial space, accelerated atherosclerosis, accelerated aging of vessel walls in the brain, myocardial edema, myocardial fibrosis, diastolic dysfunction, retardation of lung development in the fetuses of diabetic mothers, alterations of one or more pulmonary physiological parameters, increased susceptibility to infections, vascular hyperplasia in the mesentery, or redness, discoloration, dryness and ulcerations of the skin.

5. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 4 wherein the vascular complication is diabetic macular edema.
6. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 4 wherein the vascular complication is diabetic retinopathy.
7. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 6 wherein the diabetic retinopathy is nonproliferative diabetic retinopathy.
8. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 4 wherein the vascular complication is diabetic nephropathy.
9. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 4 wherein the vascular complication is edema or diabetic cardiomyopathy.
10. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 2 wherein the disease or condition is diabetes, atherosclerosis, hypertension, an acute lung injury, acute respiratory distress syndrome, age-related macular degeneration, cerebral edema, choroidal edema, choroiditis, coronary microvascular disease, cerebral microvascular disease, Eals disease, edema caused by injury, edema associated with hypertension, glomerular vascular leakage, hemorrhagic shock, Irvine Gass Syndrome, ischemia, macular edema, nephritis, nephropathies, nephrotic edema, nephrotic syndrome, neuropathy, organ failure due to edema, pre-eclampsia, pulmonary edema, pulmonary hypertension, renal failure, retinal edema, retinal hemorrhage, retinal vein occlusion, retinitis, retinopathy, silent cerebral infarction, systemic inflammatory response syndrome, transplant glomerulopathy, uveitis, vascular leakage syndrome, vitreous hemorrhage, or Von Hippel Lindau disease.
11. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 10 wherein the disease or condition is diabetes.
12. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 10 wherein the disease or condition is a macular edema, a retinopathy, or a neuropathy.
13. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 1 for use for treating a disease or condition mediated by vascular hyperpermeability at the appearance of one or more early signs of, or a predisposition to develop, the disease or condition.
14. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 1 for use for treating a disease or condition mediated by vascular hyperpermeability immediately upon diagnosis of the disease or condition.
15. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 13 or 14 wherein the disease or condition is diabetes, hypertension or atherosclerosis.
16. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 1 wherein the vascular hyperpermeability is vascular hyperpermeability of a continuous endothelium found in, or around, a brain, diaphragm, duodenal musculature, fat, heart, kidney, large blood vessel, lung, mesentery, nerve, retina, skeletal muscle, skin or testis or vascular hyperpermeability of a fenestrated endothelium found in, or around, a kidney, a pancreas, an adrenal, an endocrine gland or an intestine.
17. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 16 wherein the continuous endothelium is found in, or around, a brain, heart, lung, nerve or retina or wherein the fenestrated endothelium is found in a kidney.

18. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to any one of Claims 1-17 wherein the danazol compound is danazol.
19. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to any one of Claims 1-18 wherein the danazol compound is administered orally.
20. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to any one of Claims 1-19 wherein the animal is a human.
21. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 20 wherein the vascular-hyperpermeability-inhibiting amount is from 1 ng to 100 mg of the danazol compound per day, preferably from 1 mg to 100 mg of the danazol compound per day, and more preferably from 10 mg to 90 mg of the danazol compound per day.
22. A vascular-hyperpermeability-inhibiting amount of a danazol compound for use according to Claim 21 wherein the vascular-hyperpermeability-inhibiting amount is from 10 mg to 90 mg of the danazol compound per day.
23. An effective amount of a danazol compound for use for modulating a cytoskeleton of an endothelial cell in an animal.
24. An effective amount of a danazol compound for use according to Claim 23 wherein the modulation of the cytoskeleton includes inhibition of actin stress fiber formation, increasing or prolonging the formation of cortical actin rings, or inhibition of RhoA.
25. Pharmaceutical composition comprising an effective amount of a danazol compound for use according to any of claims 1 to 24.




FIGURE 1




FIGURE 2




FIGURE 3



EUROPEAN SEARCH REPORT

Application Number
EP 11 00 9239

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	WO 2007/009087 A2 (DMI BIOSCIENCES INC [US]; BAR-OR DAVID [US]) 18 January 2007 (2007-01-18) * page 1, line 10 - line 29 * * page 3, line 16 - line 20 * * page 21, line 25 - page 22, line 6 * * examples 4-6 *	1-25	INV. A61K31/58 A61P7/10
X	----- TOMINO Y ET AL: "CLINICAL EFFECT OF DANAZOL IN PATIENTS WITH IGA NEPHROPATHY", JAPANESE JOURNAL OF MEDICINE, vol. 26, no. 2, 1987, pages 162-166, XP009152793, ISSN: 0021-5120 * page 165 *	1,2,10, 13,14, 16-20, 23-25	
X	----- HOEPFL F ET AL: "LONG-TERM DANAZOL THERAPY FOR HEREDITARY ANGIO-EDEMA", DMW (DEUTSCHE MEDIZINISCHE WOCHENSCHRIFT), vol. 115, no. 4, 1990, pages 133-138, XP001526011, ISSN: 0012-0472 * abstract * * "Fall 1" on page 135 * * last sentence on page 137 *	1,2,10, 13,14, 16,18-25	
			TECHNICAL FIELDS SEARCHED (IPC)
			A61K A61P
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 25 January 2012	Examiner Loher, Florian
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

1
EPO FORM 1503 03/82 (P04C01)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61219185 B [0001]
- US 61315350 B [0001]
- US 3135743 A [0052] [0054]
- GB 905844 A [0052]
- WO 2007009087 A [0057]
- US 6060463 A [0073]
- US 20050101582 A [0073]
- WO 2004043480 A [0073]
- US 5407926 A [0075]
- WO 2004058289 A [0075] [0076]
- WO 0130337 A [0075] [0076]
- WO 0168053 A [0075] [0076]
- US 820325 A [0097]

Non-patent literature cited in the description

- **ICHIMURA et al.** *J. Am. Soc. Nephrol.*, 2008, vol. 19, 1463-1471 [0018] [0032]
- **FRANK et al.** *Cell Tissue Res.*, 2009, vol. 335, 41-47 [0027]
- **SIMIONESCU.** *Cell Tissue Res.*, 2009, vol. 335, 27-40 [0027]
- **VAN DEN BERG et al.** *J. Cyst. Fibros.*, 2008, vol. 7 (6), 515-519 [0027]
- **VIAZZI et al.** *Hypertens. Res.*, 2008, vol. 31, 873-879 [0027]
- **ANTONETTI et al.** *Diabetic Retinopathy.* Humana Press, 2008, 340-342 [0027]
- **FELINSKI et al.** *Current Eye Research*, 2005, vol. 30, 949-957 [0027]
- **PASCARIU et al.** *Journal of Histochemistry & Cytochemistry*, 2004, vol. 52 (1), 65-76 [0027]
- **BOUCHARD et al.** *Diabetologia*, 2002, vol. 45, 1017-1025 [0027]
- **ARSHI et al.** *Laboratory Investigation*, 2000, vol. 80 (8), 1171-1184 [0027]
- **VINOES et al.** *Documenta Ophthalmologica*, 1999, vol. 97, 217-228 [0027]
- **OOMEN et al.** *European Journal of Clinical Investigation*, 1999, vol. 29, 1035-1040 [0027]
- **VINOES et al.** *Pathol. Res. Pract.*, 1998, vol. 194, 497-505 [0027]
- **ANTONETTI et al.** *Diabetes*, 1998, vol. 47, 1953-1959 [0027]
- **POPOV et al.** *Acta Diabetol.*, 1997, vol. 34, 285-293 [0027]
- **YAMAJI.** *Circulation Research*, 1993, vol. 72, 947-957 [0027]
- **VINOES et al.** *Histochemical Journal*, 1993, vol. 25, 648-663 [0027]
- **BEALS et al.** *Microvascular Research*, 1993, vol. 45, 11-19 [0027]
- **CALDWELL et al.** *Investigative Ophthalmol Visual Sci.*, 1992, vol. 33 (5), 16101619 [0027]
- **SATCHELL et al.** *Am. J. Physiol. Renal Physiol.*, 2009, vol. 296, F947-F956 [0032]
- **HARALDSSON et al.** *Curr. Opin. Nephrol. Hypertens.*, 2009, vol. 18, 331-335 [0032]
- **BALLERMANN.** *Nephron Physiol.*, 2007, vol. 106, 19-25 [0032]
- **TOYODA et al.** *Diabetes*, 2007, vol. 56, 2155-2160 [0032]
- Endothelial Structures Involved In Vascular Permeability. **STAN.** *Endothelial Biomedicine.* Cambridge University Press, 2007, 679-688 [0032] [0033]
- Functional Ultrastructure of the Vascular Endothelium: Changes in Various Pathologies. **SIMIONESCU ; ANTOHE.** *The Vascular Endothelium I.* Springer-Verlag, 2006, 42-69 [0032]
- **YOKOMORI.** *Med. Mol. Morphol.*, 2008, vol. 41, 1-4 [0033]
- The Hepatic Sinusoidal Endothelial Cell. **DELEVE.** *Endothelial Biomedicine.* Cambridge University Press, 2007, 1226-1238 [0033]
- Normal Endothelium. **PRIES ; KUEBLER.** *The Vascular Endothelium I.* Springer-Verlag, 2006, 1-40 [0033]
- Functional Ultrastructure of the Vascular Endothelium: Changes in Various Pathologies. **SIMIONESCU ; ANTOHE.** *The Vascular-Endothelium I.* Springer-Verlag, 2006, 42-69 [0033]
- **BRAET ; WISSE.** *Comparative Hepatology*, 2002, vol. 1, 1-17 [0033]
- **KANAI et al.** *Anat. Rec.*, 1996, vol. 244, 175-181 [0033]
- **KEMPKA et al.** *Exp. Cell Res.*, 1988, vol. 176, 38-48 [0033]
- **KISHIMOTO et al.** *Am. J. Anat.*, 1987, vol. 178, 241-249 [0033]
- *Diabetic Retinopathy.* Human Press, 2008 [0046]