

(19)



(11)

EP 2 433 637 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 54
Claims EN 1, 12, 13

(51) Int Cl.:
A61K 31/517^(2006.01)

(86) International application number:
PCT/ES2010/070338

(48) Corrigendum issued on:
10.06.2015 Bulletin 2015/24

(87) International publication number:
WO 2010/133742 (25.11.2010 Gazette 2010/47)

(45) Date of publication and mention
of the grant of the patent:
11.06.2014 Bulletin 2014/24

(21) Application number: **10732398.2**

(22) Date of filing: **20.05.2010**

(54) **USE OF QUINAZOLINE DERIVATIVES FOR NEURODEGENERATIVE DISEASES**

Verwendung von Quinazolinderivaten zur neurodegenerative Erkrankungen.

UTILISATION DE DÉRIVÉS DE QUINAZOLINES DANS DES MALADIES NEURODÉGÉNÉRATIVES

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO SE SI SK SM TR**

(74) Representative: **Durán-Corretjer, S.L.P.**
Còrsega, 329
(Paseo de Gracia/Diagonal)
08037 Barcelona (ES)

(30) Priority: **20.05.2009 ES 200930189**

(56) References cited:
WO-A1-2008/113881

(43) Date of publication of application:
28.03.2012 Bulletin 2012/13

(73) Proprietor: **Consejo Superior de Investigaciones
Científicas
(CSIC)**
28006 Madrid (ES)

- **MENNITI F S ET AL: "Phosphodiesterases in the CNS: Targets for drug development" NATURE REVIEWS DRUG DISCOVERY, vol. 5, no. 8, August 2006 (2006-08), pages 660-670, XP002601485 ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL DOI: 10.1038/NRD2058**
- **CASTANO TANIA ET AL: "Synthesis, structural analysis, and biological evaluation of thioxoquinazoline derivatives as phosphodiesterase 7 inhibitors" CHEMMEDCHEM, vol. 4, no. 5, 11 May 2009 (2009-05-11), pages 866-876, XP009138976 WILEY - VCH VERLAG., WEINHEIM, DE ISSN: 1860-7179 DOI: 10.1002/CMDC.200900043 [retrieved on 2009-04-06]**

- (72) Inventors:
- **MARTÍNEZ GIL, Ana**
E-28006 Madrid (ES)
 - **GIL AYUSO-GONTAN, Carmen**
E-28006 Madrid (ES)
 - **PÉREZ MARTÍN, Concepción**
E-28006 Madrid (ES)
 - **PÉREZ CASTILLO, Ana**
E-28029 Madrid (ES)
 - **MORALES GARCÍA, José**
E-28029 Madrid (ES)
 - **REDONDO SANCHO, Miriam**
E-28006 Madrid (ES)
 - **SANZ SAN CRISTÓBAL, Marina**
E-28029 Madrid (ES)

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

EP 2 433 637 B9

Description**TECHNICAL FIELD**

5 **[0001]** The invention is directed to the field of medical chemistry and more specifically to heterocyclic derivatives of quinazolines and their potential for treating neurodegenerative and/or neurological diseases, among others Parkinson's disease, and therefore relates to the pharmaceutical sector.

STATE OF THE ART

10 **[0002]** Neurodegenerative diseases are one of the main causes of death in the Western world. Parkinson's disease (PD) is the second most common neurodegenerative disease after Alzheimer's disease (Dauer, W.; Przedborski, S., Parkinson's disease: mechanisms and models. *Neuron* 2003, 39,889-909), affecting approximately 15% of people aged over 65. Currently only symptomatic therapies are available which, although effective in the first stages of the disease have considerable long-term secondary effects. Therefore it is necessary to find new effective and safe therapies for treating this pathology.


15 **[0003]** PD is characterised by progressive death of the dopaminergic neurons of the *substantia nigra pars compacta* (SNpc) (Lang, AE.; Lozano, AM., Parkinson's disease. First of two parts. *The New England Journal of Medicine* 1998, 339,1044-1053 and Lang, AE.; Lozano, AM., Parkinson's disease. Second of two parts. *The New England Journal of Medicine* 1998, 339,1130-1143) and by inclusion bodies (Lewy's bodies) containing α -synuclein. The main consequence of this neuronal loss is a marked reduction in the brain's availability of dopamine in the *caudate nucleus* and *putamen*, areas where the dopaminergic neurons of SNpc are projected. This provokes a considerable dysfunction in the regulation of the main brain structures involved in controlling movement, the basal ganglia. Classic Parkinson's symptoms include dyskinesias (tremors of the hands, arms, legs and face), rigidity of the trunk and extremities, bradykinesia (slowness of movement) and postural instability with balancing problems.

20 **[0004]** The finding that PD was characterised by a loss of dopamine led to the discovery of therapies aimed at correcting this deficiency. These are palliative therapies, aimed at treating the symptoms of the disease but none of them manages to halt its progression (Savitt, JM.; Dawson, VL.; Dawson, TM., Diagnosis and treatment of Parkinson disease: molecules to medicine. *J Clin Invest* 2006, 116,1744-1754). Currently, the dopamine precursor, levodopa, is the most effective treatment in PD (LeWitt, P.A., Levodopa therapeutics for Parkinson's disease: new developments. *Parkinsonism Relat Disord.* 2009, Suppl 1,S31-4). This treatment is sometimes combined with other drugs such as peripheral decarboxylase inhibitors (carbidopa) (Abdel-Salam, O.M., Drugs used to treat Parkinson's disease, present status and future directions. *CNS Neurol Disord Drug Targets.* 2008, 7, 321-442), catechol-O-methyl-transferase (COMT) inhibitors, which prolong the half life of levodopa (Gallagher, D. A.; Schrag, A., Impact of newer pharmacological treatments on quality of life in patients with Parkinson's disease. *CNS Drugs.* 2008, 22, 563-586) and dopamine agonists that directly stimulate the postsynaptic dopamine receptors (Kiebertz, K., Therapeutic strategies to prevent motor complications in Parkinson's disease. *J Neurol.* 2008, 255, Suppl 4, 42-45). Despite all these advances in the treatment of PD symptoms, their efficacy declines over time fundamentally due to the development of motor complications such as dyskinesias and dystonias.

25 **[0005]** In view of the major limitations presented by current therapies, both pharmacological and surgical, there is a need to develop other alternatives to slow down or stop the progress of the disease (Poewe, W., Treatments for Parkinson disease - past achievements and current clinical needs. *Neurology* 2009, 72, 7 Suppl, S65-73). Current research is focused on preventing neuronal dopaminergic degeneration and on the discovery of new drugs, levodopa alternatives, to manage to stop the progress of the disease and even to generate new dopaminergic type neurons. Recently different chemical families have emerged with potential in this pathology, phosphodiesterase inhibitors being one of them (Use of pde7 inhibitors for the treatment of movement disorders, USPTO Application: 20080260643, WO2008/113881). Given our prior experience in this area (Castaño, T.; Wang, H.; Campillo, N.E.; Ballester, S.; González-García, C.; Hernández, J.; Pérez, C.; Cuenca, J.; Pérez-Castillo, A.; Martínez, A.; Huertas, O.; Gelpí, J.L.; Luque, F.J.; Ke, H.; Gil, C., Synthesis, structural analysis, and biological evaluation of thioxoquinazoline derivatives as phosphodiesterase 7 inhibitors. *ChemMedChem.* 2009, 4, 866-876), we demonstrate the use of heterocyclic quinazoline derivatives as effective agents in Parkinson models *in vivo* and propose their use not only in this neurodegenerative pathology but also in other neurological diseases due to their neuroprotective effect in all pathologies of the nervous system occurring with neuronal death.

BRIEF DESCRIPTION OF THE INVENTION

30 **[0006]** An object of the present invention is the use of a compound of formula (I):



wherein:

A, is fused carbocycle or heterocycle optionally substituted with 5, 6 or 7 members saturated or unsaturated, --- may be a double bond;

X and Y, are selected independently from the group consisting of hydrogen, alkyl, =O, =S, aryl, O-alkyl, O-aryl, S-alkyl or -S-aryl; and


R, R¹, R² and R³ selected independently from the group consisting of hydrogen, halogen, alkyl, haloalkyl, aryl, cycloalkyl, (CH₂)_n-aryl, heteroaryl, -OR³, -C(O)OR³, (CH₂)_n-C(O)OR³ or -S(O)_t, wherein n is greater than or equal to 0 and wherein t is 1 or 2,

or a pharmaceutically acceptable salt, prodrug, solvate or stereoisomer thereof, to produce a medicament for the treatment and/or prevention of neurodegenerative diseases, including, Parkinson's disease Alzheimer's disease, amyotrophic lateral sclerosis, brain ischemia, and/or of pathologies or neurological diseases wherein the dopaminergic system is involved, including, but not limited to, post-encephalitic parkinsonism, dystonias, Tourette's syndrome, periodic limb movement disorders, restless legs syndrome, attention deficit disorders with hyperactivity.

DETAILED DESCRIPTION OF THE INVENTION

[0007] The present invention is based on the fact that the inventors have demonstrated that the compounds of formula (I) are neuroprotective in primary cultures of astrocytes and glia and/or dopaminergic cell lines (Examples 1-2), as well as in an *in vivo* model of lipopolysaccharide (LPS)-induced neurotoxicity (Example 3). In addition, the inventors have demonstrated that the compounds are capable of crossing the blood-brain barrier *in vitro* and have antioxidant properties, meaning that they can be used in the production of pharmaceutical compositions for the treatment and/or prevention of neurodegenerative diseases and/or neurological diseases.

[0008] Therefore, one object of the present invention is the use of a compound of formula (I):



wherein:

A, is a fused carbocycle or heterocycle optionally substituted with 5, 6 or 7 members saturated or unsaturated, --- may be a double bond;

X and Y, are independently selected from the group consisting of hydrogen, alkyl, =O, =S, aryl, O-alkyl, O-aryl, S-alkyl or -S-aryl; and


R, R¹, R² and R³ are independently selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, aryl, cycloalkyl, (CH₂)_n-aryl, heteroaryl, -OR³, -C(O)OR³, (CH₂)_n-C(O)OR³ or -S(O)_t, wherein n is greater than or equal to 0 and wherein t is 1 or 2,

or a pharmaceutically acceptable salt, prodrug, solvate or stereoisomer thereof,

to produce a medicament for the treatment and/or prevention of neurodegenerative and/or neurological diseases.

[0009] Preferably, A is a carbocycle with 6 members. More preferably, A is phenyl.

[0010] In a preferred embodiment, the invention relates to the use of a compound of formula (II):



(II)

wherein

X and Y are independently selected from O or S; and

R and R¹ are independently selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, aryl, cycloalkyl, (CH₂)_n-aryl, heteroaryl, -OR³;-C(O)OR³, (CH₂)_n-C(O)OR³ or -S(O)_t-, wherein R³ is selected between hydrogen, alkyl, aryl or cycloalkyl, n is greater than or equal to 0 and t is 1 or 2,

or a pharmaceutically acceptable salt, prodrug, solvate or stereoisomer thereof,


to produce a medicament for the treatment and/or prevention of neurodegenerative and/or neurological diseases.

[0011] Preferably, X is O and Y is S.

[0012] Preferably, X is S and Y is S.

[0013] Preferably, R is selected between H or C₁-C₆ alkyl. More preferably, R is methyl.

[0014] In another preferred embodiment, the invention relates to the use of a compound of formula (III):



(III)

wherein:

X is selected between O or S and Y is selected between O-aryl or S-alkyl; and

R and R¹ are independently selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, aryl, cycloalkyl, (CH₂)_n-aryl, heteroaryl, -OR³;-C(O)OR³, (CH₂)_n-C(O)OR³ or -S(O)_t-, wherein R³ is selected between hydrogen, alkyl, aryl or cycloalkyl, n is greater than or equal to 0 and t is 1 or 2

or a pharmaceutically acceptable salt, prodrug, solvate or stereoisomer thereof,

to produce a medicament for the treatment and/or prevention of neurodegenerative and/or neurological diseases.

[0015] Preferably, X is O.

[0016] Preferably, X is S.

[0017] Preferably, Y is S-alkyl C₁-C₆. More preferably, Y is S-CH₃.

[0018] Preferably, R is selected between H or C₁-C₆ alkyl. More preferably, R is methyl.

[0019] Preferably, R₁ is phenyl substituted or non-substituted. More preferably, R₁ is phenyl substituted by at least one halogen selected between Br or F.

[0020] A particular object of the invention consists of the use of a compound of the invention of formula (I) selected from the following group:

- 3-Phenyl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline,
- 3-(2,6-Difluorophenyl)-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline,
- 3-(2-Bromophenyl)-8-methyl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline
- 3-(2,6-Difluorophenyl)-8-methyl-2-methylthio-4-oxo-3,4-dihydroquinazoline,
- 3-(2-Bromophenyl)-8-methyl-2-methylthio-4-oxo-3,4-dihydroquinazoline,

- 3-Phenyl-2,4-dithioxo-1,2,3,4-tetrahydroquinazoline,
- 3-Phenyl-2-methylthio-4-thioxo-3,4-dihydroquinazoline,
- 3-(2,6-Difluorophenyl)-2,4-dithioxo-1,2,3,4-tetrahydroquinazoline.

5 or a pharmaceutically acceptable salt, prodrug, solvate or stereoisomer thereof, to produce a medicament for the treatment and/or prevention of neurodegenerative and/or neurological diseases.

[0021] The neurodegenerative and/or neurological disease is selected between Parkinson's disease, Alzheimer's disease, amyotrophic lateral sclerosis, brain ischemia, post-encephalitic parkinsonism, dystonias, Tourette's syndrome, periodic limb movement disorders, restless legs syndrome, attention deficit disorders with hyperactivity.

10 [0022] "Carbocycle" refers to a cyclic system consisting of just carbon and hydrogen atoms. For the purpose of this invention, the carbocycle may be a monocyclic, bicyclic or tricyclic system and may include fused systems and the cycle may be partially saturated or aromatic. Examples of these carbocycles include, but are not limited to, cycloalkyls, as defined herein, phenyl, naphthyl, anthracenyl, indanyl and similar.

15 [0023] "Heterocycle" refers in the present invention to a stable radical having a ring of 3 to 15 members comprising carbon atoms and one to five heteroatoms selected from the group consisting of nitrogen, oxygen and sulphur, preferably a 4 to 8-member ring with one or more heteroatoms, more preferably a 5 or 6-member ring with one or more heteroatoms. For the purpose of this invention, the heterocycle may be a monocyclic, bicyclic or tricyclic ring system, which can include condensed ring systems, and the nitrogen, carbon or sulphur atom in the heterocyclic radical can be optionally oxidised, the nitrogen atom can be optionally quaternised, and the heterocyclic radical may be partially or totally saturated or be

20 aromatic. Examples of such heterocycles include, but are not limited to azepines, benzimidazole, benzothiazole, furane, isothiazole, imidazole, indole, piperidine, piperazine, purine, quinoline, thiadiazole, tetrahydrofurane, cumane, morpholine, pyrrole, pyrazole, oxazole, isoxazole, triazole, imidazole, etc.

[0024] The term "alkyl" refers in the present invention to radicals of hydrocarbonated chains, linear or branched, having 1 to 10 carbon atoms, preferably 1 to 6, and that bind to the rest of the molecule through a simple bond, for example,

25 methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *tert*-butyl, *sec*-butyl, *n*-pentyl, *n*-hexyl etc. The alkyl groups can be optionally substituted by one or more substituents such as halogen (known as haloalkyl), hydroxyl, alkoxy, carboxyl, carbonyl, cyan, acyl, alkoxy carbonyl, amino, nitro, mercapto and alkylthio.

[0025] "Cycloalkyl" refers in the present invention to a stable monocyclic or bicyclic radical of 3 to 10 members, preferably of 3 to 8 members and more preferably of 6 members, which is saturated or partially saturated, and which

30 only consists of carbon and hydrogen atoms, such as cyclopentyl, cyclohexyl or adamantyl which can be optionally substituted with one or more groups such as alkyl, halogen, hydroxyl, alkoxy, carboxyl, cyan, carbonyl, acyl, alkoxy carbonyl, amino, nitro, mercapto and alkylthio.

[0026] The term "aryl" refers in the present invention to a radical phenyl, naphthyl, indenyl, phenantryl, or anthracyl. The radical aryl may be optionally substituted by one or more substituents such as alkyl, haloalkyl, aminoalkyl, dialkylamino, hydroxyl, alkoxy, phenyl, mercapto, halogen, nitro, cyan and alkoxy carbonyl.

35 [0027] "Heteroaryl" refers to an aryl having at least one heterocycle.

[0028] "Halogen" refers to fluorine, chlorine, bromine or iodine.

[0029] The compounds of the present invention represented by the formula (I), and more specifically, the specific compounds belonging to this previously described general formula may include isomers, depending on the presence of

40 multiple bonds (for example, *Z*, *E*), including optic isomers or enantiomers, depending on the presence of chiral centres. The individual isomers, enantiomers or diastereoisomers and the mixtures thereof fall within the scope of the present invention. The individual enantiomers or diastereoisomers, as well as their mixtures, can be separated by conventional techniques.

[0030] Also, the scope of this invention includes the prodrugs of the compounds of formula (I). The term "prodrug" as used herein includes any compound derived from a compound of formula (I), for example, esters, including esters of

45 carboxylic acids, amino acid esters, phosphate esters, metal salt sulfonate esters, etc., carbamates, amides, etc., which, when administered to an individual is capable of providing, directly or indirectly, said compound of formula (I) in said individual. Advantageously, said derivative is a compound that increases the bioavailability of the compound of formula (I) when administered to an individual or that potentiates the release of the compound of formula (I) in a biological

50 compartment. The nature of said derivative is not critical on condition that it can be administered to an individual and provides the compound of formula (I) in a biological compartment of an individual. The aforesaid prodrug can be produced following conventional methods known to experts in the art.

[0031] The compounds of the invention may be in crystalline form as free compounds or in solvate form, intending both forms to be within the scope of the present invention. In this sense, the term "solvate", as used herein, includes

55 both pharmaceutically acceptable solvates, in other words, solvates of the compound of formula (I) which can be used to produce a medicament, as well as pharmaceutically unacceptable solvates, which can be useful to produce pharmaceutically acceptable solvates or salts. The nature of the pharmaceutically acceptable solvate is not critical on condition that it is pharmaceutically acceptable. In a particular embodiment, the solvate is a hydrate. The solvates can be obtained

by conventional solvation methods known to technicians in the art.


[0032] For therapeutic application, the compounds of formula (I), their isomers, salts, prodrugs or solvates, will come preferably in a pharmaceutically acceptable or substantially pure form, in other words, having a pharmaceutically acceptable level of purity excluding standard drugs such as diluents and carriers, and not including material considered toxic at standard dose levels. The purity levels for the active principle are preferably higher than 50%, more preferably higher than 70%, more preferably higher than 90%. In a preferred embodiment, they are higher than 95% of the compound of formula (I), or the salts, solvates or prodrugs thereof.

[0033] Unless otherwise specified, the compounds of the invention also include compounds that differ only in the presence of one or more isotopically-enriched atoms. For example, compounds having said structure, excluding the substitution of hydrogen with a deuterium or a tritium, or the substitution of a carbon by a carbon enriched with ^{13}C or ^{14}C or nitrogen enriched with ^{15}N , are within the scope of the invention.

[0034] The compounds of formula (I) for therapeutic use are prepared in solid form or aqueous suspension, in a pharmaceutically acceptable diluent. These preparations may be administered via any appropriate route of administration, wherefore said preparation will be formulated in the suitable pharmaceutical form for the selected route of administration. In a particular embodiment, the compound of formula (I) provided by this invention is administered orally, topically, rectally or parenterally (including subcutaneously, intraperitoneally, intradermally, intramuscularly, intravenously, etc.). A review of the different pharmaceutical forms for administering medicaments and the excipients required to obtain them can be found, for example, in the "Treaty of Galenic Pharmacy", C. Faulí i Trillo, 1993, Luzán 5, S.A. Ediciones, Madrid, or in other standard or similar Pharmacopoeias of Spain and the US.

[0035] The compounds described in the present invention, their pharmaceutically acceptable salts, prodrugs and/or solvates as well as the pharmaceutical compositions containing them can be used in conjunction with other additional drugs in order to provide a combination therapy. Said additional drugs may form part of the same pharmaceutical composition or, alternatively, may be provided in the form of a separate composition for administration simultaneously or not with the administration of the pharmaceutical composition comprising a compound of formula (I), or pharmaceutically acceptable prodrug, solvate, derivative or salt thereof.

[0036] Another additional object of the present invention consists of a method for treating a neurodegenerative and/or neurological disease comprising the administration of a therapeutically effective amount of a compound of formula (I):



wherein:

A, is a fused carbocycle or heterocycle optionally substituted with 5, 6 or 7 members saturated or unsaturated, --- may be a double bond;

X and Y, are independently selected between the group consisting of hydrogen, alkyl, =O, =S, aryl, O-alkyl, O-aryl, S-alkyl or -S-aryl; and

R, R¹, R² and R³ are independently selected between the group consisting of hydrogen, halogen, alkyl, haloalkyl, aryl, cycloalkyl, (CH₂)_n-aryl, heteroaryl, -OR³; -C(O)OR³, (CH₂)_n-C(O)OR³ or -S(O)_t, wherein n is greater than or equal to 0 and wherein t is 1 or 2,

or a pharmaceutically acceptable salt, prodrug, solvate or stereoisomer thereof.

[0037] In the sense used in this description, the expression "therapeutically effective amount" refers to the quantity of agent or compound capable of having a neuroprotective effect on primary cultures of astrocytes and glia and/or on dopaminergic cell lines, calculated to produce the required effect *in vivo* and, in general, will be determined, among other aspects, by the inherent properties of the compounds, including the age, state of the patient, severity of the alteration or disorder, and route and frequency of administration.

[0038] The neurodegenerative and/or neurological diseases to be treated in said method are selected between Parkinson's disease, Alzheimer's disease, amyotrophic lateral sclerosis, brain ischemia, post-encephalitic parkinsonisms, dystonias, Tourette's syndrome, periodic limb movement disorders, restless legs syndrome, attention deficit disorders with hyperactivity.

[0039] The compound of the invention is compatible for use in protocols wherein the compounds of formula (I) or their

mixtures are used alone or in combination with other treatments or medical procedures.

DESCRIPTION OF THE DRAWINGS

5 [0040]

Figure 1.- Neuroprotective effect of the compounds on lipopolysaccharide (LPS)-stimulated primary astrocyte cultures.

Figure 2.- Neuroprotective effect of the compounds on LPS-stimulated primary microglia cultures.

10 **Figure 3.-** Neuroprotection of the SH-SY5Y dopaminergic cell line

Figure 4.- (A) Viability of the dopaminergic cells of "*substantia nigra*" neurons following LPS-induced damage. (B) Microglial activation in response to LPS.

EXAMPLES OF THE INVENTION

15 [0041] In the following examples, the following compounds have been tested having a formula comprised within formula (I):

- 3-phenyl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline (**Comp. 1**),
- 20 - 3-(2,6-difluorophenyl)-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline (**Comp. 2**),
- 3-(2-bromophenyl)-8-methyl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline (**Comp. 3**),
- 3-(2,6-difluorophenyl)-8-methyl-2-methylthio-4-oxo-3,4-dihydroquinazoline (**Comp. 4**),
- 3-(2-bromophenyl)-8-methyl-2-methylthio-4-oxo-3,4-dihydroquinazoline (**Comp. 5**),
- 3-phenyl-2,4-dithioxo-1,2,3,4-tetrahydroquinazoline (**Comp. 6**),
- 25 - 3-phenyl-2-methylthio-4-thioxo-3,4-dihydroquinazoline (**Comp. 7**),
- 3-(2,6-difluorophenyl)-2,4-dithioxo-1,2,3,4-tetrahydroquinazoline (**Comp 8**).

Example 1.- Measurement of the neuroprotective effect of the compounds of formula (I) according to nitrite production by the different cell lines

30 [0042] Essentially, primary cultures of microglia and astrocytes were used (Luna-Medina, R.; Cortes-Canteli, M.; Alonso, M.; Santos, A.; Martinez, A.; Perez-Castillo, A., Regulation of inflammatory response in neural cells in vitro by thiazolidinones derivatives through peroxisome proliferator-activated receptor gamma activation. J. Biol. Chem. 2005, 280, 21453-21462). The primary cultures of astrocytes and microglia were obtained from the cortex and hippocampus of postnatal 2-day-old mice. After dissecting the cortex and hippocampus and clearing them of the meninges, cells are disintegrated by mechanical grinding and incubated with 0.25% trypsin/EDTA at 37°C during 45 minutes. DMEM with 10% foetal serum is added to stop the digestion with trypsin and mechanical grinding of the tissue is completed. Centrifugation is applied at 800xg/5 min and the precipitate washed 3 times in EBSS; finally, cells are resuspended in DMEM plus 10% foetal serum and seeded at a density of 0.5x10⁵ cells/cm². Cells are incubated for 10-12 days after which a monolayer of astrocytes is observed to which microglia cells lightly adhere. To isolate microglia cells culture bottles are incubated in a rotary agitator at 37°C during 4 hours at 250 rpm and the medium containing microglia centrifuged at 1500xg/5 min. Microglia cells are resuspended in DMEM/10% FBS and seeded at a density of 2-4x10⁵ cells/cm². After 1 hour of incubation, to allow them to adhere to the plate, they are washed with TD and incubated in DMEM/10% FBS during 24 hours, after which cells are used in the various experiments. The level of purity of these cultures is determined by immunocytochemistry assays with neuron-specific antibodies β-tubulin and MAP2), astrocytes (GFAP), oligodendrocytes (CNPase) and microglia (OX42).

45 [0043] Cell cultures of astrocytes and microglia are treated with LPS (10 μg/ml) in the absence and presence of the different compounds. The compounds are added 1 hour before the inflammatory stimulus. At different times after incubation, cells are washed and gathered for corresponding measurement of the effect of the compounds on NO (nitric oxide) production by iNOS (inducible nitric oxide synthase) as an indicator of neural damage due to inflammatory processes (Kroncke K. D.; Fehsel K.; Kolb-Bachofen V., Nitric oxide: cytotoxicity versus cytoprotection--how, why, when, and where? Nitric Oxide 1997, 1, 107-120). To this effect, after 24 hours of incubation the quantity of nitrites, one of the oxidation products of NO, is determined. To achieve this, the method based on the Griess reaction is used (Griess, P. Bemerkungen zu der abhandlung der H.H. Weselsky und Benedikt Ueber einige azoverbindungen. Chem. Ber. 1879, 12, 426-428): 100 μl of supernatant from the cultures is mixed with 100 μl of Griess reagent in a 96-well plate and incubated during 15 min at room temperature. Next, absorbance is measured at 540 nm in a microplate reader. The amount of nitrites produced is determined using a standard sodium nitrite curve.

55 Reference compounds in the assay: Rolipram and BRL50481.

See Figures 1 and 2

Example 2.- Neuroprotection assay of SH-SY5Y neuroblastoma cells

5 [0044] The human neuroblastoma cell line SH-SY5Y is exposed to a 50 μ M concentration of neurotoxin 6-hydroxy-dopamine (6-OHDA) in the presence or absence of S14 or TC3.6. 6-OHDA is a highly toxic substance for these cells and incubation with it provokes significant cell death, it being commonly used as an *in vitro* cell model of parkinsonism (Mendez, J.S.; Finn, B.W., Use of 6-hydroxydopamine to create lesions in catecholamine neurons in rats. J Neurosurg. 1975, 42, 166-173.). Cellular viability is determined using the bromide compound of 3(4,5-dimethylthiazol-2)-2,5-diphenyltetrazolium (MTT) which measures the integrity of the mitochondrial function. Each data represents the average \pm SD of five replicates in three different experiments. * $P < 0.05$ and ** $P < 0.01$, statistically significant differences between the different experiments. See figure 3.

Example 3.- Neuronal survival in vivo

15 [0045] The lipopolysaccharide LPS, 5 μ g) or vehicle is injected unilaterally into the *substantia nigra* (SN) of adult rats. One of the animal groups additionally receives 5 nmol of S14 in the injection. Three days after the operation, the animals are given an intracardial perfusion of 4% paraformaldehyde in PBS, during approximately 45-60 minutes. Following perfusion the brain is extracted and left in the paraformaldehyde solution during 24 hours after which it is changed to a solution of 4% paraformaldehyde/30% sucrose for approximately 48 hours. Brains fixed in this way are frozen in carbonic snow taking cuts of 25 μ m in a cryostat. For single-antibody staining, the ABC system is used (ABC Elite kit, Vector). Selected cuts are washed in PBS and treated with a methanol/PBS/H₂O/H₂O₂ solution in order to block endogenous peroxidases. After washing with PBS cuts are blocked with 0.1 M lysine/ 5% serum/ 0.1 % triton X-100/ 4% BSA in PBS. Incubation with the corresponding primary antibody is carried out at 4°C during 18 hours in PBS with 4% BSA, 1% serum and 0.1% triton X-100. After washing with PBS, cuts are incubated during 1 hour at room temperature with the corresponding biotinylated secondary antibody in PBS containing 4% BSA, 1% serum and 0.1% triton X-100, at a 1:200 dilution. After incubation the ABC kit is added and cuts are incubated during 1 hour. After washing with PBS they are revealed with DAB and mounted onto gelatine-coated slides and dehydrated for subsequent microscopic observation. Neuronal integrity is measured through conventional Nissl staining and the survival of dopaminergic cells by staining the cuts with an anti-tyrosine hydroxylase (TH) specific antibody. Glial activation is determined using anti-GFAP antibodies (stains reactive astrocytes) and anti-CD11 b (OX-42, stains activated microglia). See figure 4

Example 4: Measurement of the antioxidant effect of the compounds of formula (I)

35 [0046] Given that in many neurodegenerative diseases neuronal death occurs due to oxidative stress, the capacity of these compounds to capture free radicals has been evaluated using the ORAC methodology.
 [0047] Determination is following the ORAC-FL method of Ou et al. partially adapted by BMG LABTECH (BMG LABTECH Application note 148 (2006): ORAC Assay on the FLUOstar OPTIMA to Determine Antioxidant Capacity. http://www.bmglabtech.com/application-notes/fluorescence-intensity/orac_148.cfm). The FLUOstar Optima (BMG Labtechnologies GmbH, Offenburg, Germany) plate reader was used with an excitation filter at 485 nm excitation and 520 nm emission. 2,2'-Azobis-(amidinopropane) dihydrochloride (AAPH), (\pm)-6-hydroxy-2,5,7,8- tetramethylchromane-2-carboxylic acid (trolox) and fluorescein (FL) are purchased from Sigma- Aldrich. The reaction is carried out in 75 mM phosphate buffer (pH 7.4) and a final reaction volume of 200 μ L. The antioxidant (25 μ L) and fluorescein (150 μ L; 10 nM) are placed on black 96-well plates (96F untreated, NuncTM). The mixture is pre-incubated during 30 minutes at 37 °C afterwards rapidly adding a solution of AAPH (25 μ L, 240 mM), using a multi-channel pipette. The microplate is rapidly placed on the reader and fluorescence is measured every 90 sec during 90 min. The 96-well plate is agitated automatically before each reading. The compounds are tested at 4 different concentrations (10-1 μ M). A blank is also placed (FL + AAPH in phosphate buffer) and as reference compound four concentrations of trolox (10-1 μ M). All reactions are carried out in duplicate, and at least three different tests per compound. Data are exported to an Excel spreadsheet (Microsoft), representing absorbance versus time and subsequently the area under curve is calculated following the formula (1)

$$(1) \text{ AUC} = f_1 / f_0 + f_j / f_0 + \dots + f_{34} / f_0 + (f_{35} / f_0)$$

55

wherein: AUC = area under curve
 f_0 = fluorescence measured at time 0

f_i = fluorescence measured at time t

[0048] The net AUC values of each compound are determined by subtracting from the AUC value of the compound the AUC value of the blank (FL + AAPH in phosphate buffer).

[0049] The measurement of antioxidant is expressed in trolox equivalents. Relative trolox equivalents are calculated according to formula (2):

$$(2) \left(\frac{AUC_{sample} - AUC_{blank}}{AUC_{Trolox} - AUC_{blank}} * \frac{[Trolox]}{[sample]} \right)$$

[0050] The results are shown in table 1.

Table 1.- Antioxidant capacity of the compounds in trolox equivalents

Compound	Trolox equivalents
1	1.19
2	Not to 10 μ M
3	1.31
4	0.13
5	Not to 10 μ M
6	0.46
7	0.37
8	n.d.
n.d.= not determined	

Example 5: Measurement of the blood brain permeability of the compounds of formula (I)

[0051] The passing of the blood brain barrier is determined *in vitro* by means of a permeability assay using an artificial membrane, PAMPA (*Parallel Artificial Membrane Permeation Assay*), following the procedure previously optimised in the authors' laboratories. Two 96-well plates (Millipore) are used on a *sandwich* type assembly. The donating microplate contains a PVDF filter at the bottom of each well where a lipid extract of pig is placed, acting as the biological barrier. Each assay is validated with 8 commercial drugs whose CNS penetration is known.

[0052] Initially, wavelengths are determined of the absorption spectrum of a solution of each compound (1 mg in 5 ml of PBS/ EtOH 70:30), whose absorbance levels must be between 1-0.8 so as to, subsequently, measure permeability.

[0053] Once the required wavelengths have been selected, 180 μ L of each solution (1 mg of compound in 5 mL of PBS/ EtOH 70:30) is added to each well of the donor plate. The artificial membrane is coated with 4 μ L of PBL solution in 20 mg/ml dodecane. 180 μ l of the buffer solution (pH=7.4) is added to the acceptor plate, next forming a "sandwich"-type system in such a way that the donor plate is above the acceptor plate through the artificial membrane. This system is incubated for 4 h without agitation at room temperature. The absorbance levels of the acceptor plate are measured using a UV spectrophotometer on a 96-well plate with 130 μ l in each one. The results are expressed as the average value of three independent tests each one containing three repetitions of each derivative to be analysed. The differences between final and initial absorbance levels are correlated with permeability.

[0054] The method employed was validated in the same conditions using 9 commercial products whose capacity to penetrate the CNS is known. In this way, a good linear relationship was obtained between the experimental and described permeability. Based on this correlation it was possible to establish limits to predict the passing of the blood brain barrier. Therefore it is considered that a molecule passes the blood brain barrier (CNS +) when its permeability (Pe) is above 4 10^{-6} .cm.s $^{-1}$ and therefore, it would be capable of reaching its therapeutic target situated in the CNS. At the same time, a molecule is considered not to pass the blood brain barrier (CNS -) when Pe is below 2 10^{-6} .cm.s $^{-1}$. Intermediate values present uncertainty (CNS+/-).


[0055] The data is shown in table 2. Most of the compounds evaluated are capable of passing the blood brain barrier according to this methodology.

Table 2.- Passage of the blood brain barrier of the compounds

Compound	CNS Penetration	Bibliography
Chlorpromazine	+	6.5
Desipramine	+	12
Enoxacin	-	1.8
Hydrocortisone	-	1.9
Imipramine	+	13
Ofloxacin	-	0.8
Promazine	+	8.1
Verapamil	+	16
Atenolol	-	0.8
Comp. 1	+	
Comp. 2	+	
Comp. 3	+	
Comp. 4	n.d.	
Comp. 5	-	
Comp. 6	+	
Comp. 7	n.d.	
Comp. 8	+	
n.d.= not determined due to lack of solubility in the experimental medium		

Claims

1. Use of a compound of formula (I):



(I)

wherein:

A, is a fused carbocycle or heterocycle optionally substituted with 5, 6 or 7 members saturated or unsaturated, --- may be a double bond;

X and Y, are independently selected between the group consisting of hydrogen, alkyl, =O, =S, aryl, O-alkyl, O-aryl, S-alkyl or -S-aryl; and

R, R¹, R² and R³ are independently selected between the group consisting of hydrogen, halogen, alkyl, haloalkyl, aryl, cycloalkyl, (CH₂)_n-aryl, -, -OR³; -C(O)OR³, (CH₂)_n-C(O)OR³ or -S(O)_t, wherein n is greater than or equal to 0 and wherein t is 1 or 2,


or a pharmaceutically acceptable salt, prodrug, solvate or stereoisomer thereof, **wherein the prodrug is an ester of carboxylic acid, an amino acid ester, a phosphate ester, a metal salt sulfonate ester, a carbamate or an**

amide derived of the compound of formula (I) which when administered to an individual is capable of providing, directly or indirectly, said compound of formula (I),

to produce a medicament for the treatment of neurodegenerative and/or neurological diseases, characterized in that the neurodegenerative and/or neurological disease is selected from the following group: Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, brain ischemia, post-encephalitic parkinsonisms, dystonias, Tourette's syndrome, periodic limb movement disorders, restless legs syndrome, attention deficit disorders with hyperactivity.

2. Use according to claim 1 wherein A is a carbocycle with 6 members, preferably phenyl.

3. Use according to claim 1, characterized in that the compound of formula (I) is a compound of formula (II):



(II)

wherein

X and Y are independently selected between O or S; and


R and R¹ are independently selected between the group consisting of hydrogen, halogen, alkyl, haloalkyl, aryl, cycloalkyl, (CH₂)_n-aryl, heteroaryl, -OR³; -C(O)OR³, (CH₂)_n-C(O)OR³ or -S(O)_t, wherein R³ is selected between hydrogen, alkyl, aryl or cycloalkyl, n is greater than or equal to 0 and t is 1 or 2;

4. Use according to claim 3 wherein X is O and Y is S.

5. Use according to claim 3 wherein X is S and Y is S.

6. Use according to any of claims 3 to 5 wherein R is selected between H or alkyl C₁-C₆, preferably methyl.

7. Use according to claim 1, characterized in that the compound of formula (I) is a compound of formula (III):



(III)

wherein:

X is selected between O or S and Y is selected between O-aryl or S-alkyl; and

R and R¹ are independently selected between the group consisting of hydrogen, halogen, alkyl, haloalkyl, aryl, cycloalkyl, (CH₂)_n-aryl, -OR³; -C(O)OR³, (CH₂)_n-C(O)OR³ or -S(O)_t, wherein R³ is selected between hydrogen, alkyl, aryl or cycloalkyl, n is greater than or equal to 0 and t is 1 or 2

8. Use according to claim 7, wherein X is O.

9. Use according to claim 7, wherein X is S.


10. Use according to any of claims 7 to 9, wherein Y is S-alkyl C₁-C₆, preferably S-CH₃.

11. Use according to any of the preceding claims wherein R is selected between H or alkyl C1-C6, preferably methyl.
12. Use according to any of the previous claims wherein R¹ is phenyl substituted or non-substituted.
- 5 13. Use according to the preceding claim wherein R¹ is phenyl substituted with at least one halogen selected between Br or F.
14. Use according to claim 1, **characterized in that** the compound of formula (I) is a compound selected from the following group:

- 10
- 3-phenyl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline,
 - 3-(2,6-difluorophenyl)-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline,
 - 3-(2-Bromophenyl)-8-methyl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazoline,
 - 15 - 3-(2,6-Difluorophenyl)-8-methyl-2-methylthio-4-oxo-3,4-dihydroquinazoline,
 - 3-(2-Bromophenyl)-8-methyl-2-methylthio-4-oxo-3,4-dihydroquinazoline,
 - 3-Phenyl-2,4-dithioxo-1,2,3,4-tetrahydroquinazoline, and
 - 3-Phenyl-2-methylthio-4-thioxo-3,4-dihydroquinazoline
 - 3-(2,6-Difluorophenyl)-2,4-dithioxo-1,2,3,4-tetrahydroquinazoline

20 **Patentansprüche**

1. Verwendung einer Verbindung mit der Formel (I):



35 wobei

A ein fusionierter Carbocyclus oder Heterocyclus ist, der optional mit 5, 6 oder 7 gesättigten oder ungesättigten Glieder substituiert ist,

40 --- eine Doppelbindung sein kann;

X und Y unabhängig ausgewählt sind aus der Gruppe bestehend aus Wasserstoff, Alkyl =O, =S, Aryl, O-Alkyl, O-Aryl, S-Alkyl oder -S-Aryl; und

R, R¹, R² und R³ unabhängig ausgewählt sind aus der Gruppe bestehend aus Wasserstoff, Halogen, Alkyl, Haloalkyl, Aryl, Cycloalkyl, (CH₂)_n-Aryl, -OR³; -C(O)OR³, (CH₂)_n-C(O)OR³ oder -S(O)_t-, wobei n größer oder

45 gleich 0 ist und t 1 oder 2 ist,

oder ein pharmazeutisch akzeptables Salz, Prodrug, Solvat oder Stereoisomer davon ist, wobei das Prodrug ein Carbonsäureester, ein Aminosäureester, ein Phosphatester, ein Metallsalzsulfonatester, ein Carbamat oder ein Amid ist, abgeleitet von der Verbindung mit der Formel (I), die, wenn sie an ein Individuum verabreicht wird, imstande


50 ist direkt oder indirekt die besagte Verbindung mit der Formel (I) bereitzustellen,

um ein Medikament zur Behandlung von neurodegenerativen und/oder neurologischen Erkrankungen herzustellen, **dadurch gekennzeichnet, dass** die neurodegenerativen und/oder neurologischen Erkrankungen ausgewählt sind aus der folgenden Gruppe: Alzheimer-Krankheit, Parkinson-Krankheit, amyotrophe Lateralsklerose, Hirnischämie, post-enzephalitisches Parkinsonsyndrom, Dystonie, Tourette-Syndrom, Syndrome der Bewegung der Gliedmaßen, Syndrom der ruhelosen Beine, Aufmerksamkeitsdefizit-Syndrome mit Hyperaktivität.

55

2. Verwendung nach Anspruch 1, wobei A ein Carbocyclus mit 6 Gliedern, vorzugsweise Phenyl, ist.

3. Verwendung nach Anspruch 1, **dadurch gekennzeichnet, dass** die Verbindung mit der Formel (I) ein Verbindung mit der Formel (II) ist:




(II)

wobei

15 X und Y unabhängig ausgewählt sind aus O oder S; und
 R und R¹ unabhängig ausgewählt sind aus der Gruppe bestehend aus Wasserstoff, Halogen, Alkyl, Haloalkyl, Aryl, Cycloalkyl, (CH₂)_n-Aryl, Heteroaryl, -OR³; -C(O)OR³, (CH₂)_n-C(O)OR³ oder -S(O)_t, wobei R³ ausgewählt ist aus Wasserstoff, Alkyl, Aryl oder Cycloalkyl, n größer oder gleich 0 ist und t 1 oder 2 ist.

- 20
4. Verwendung nach Anspruch 3, wobei X O und Y S ist.
5. Verwendung nach Anspruch 3, wobei X S und Y S ist.
- 25 6. Verwendung nach einem der Ansprüche 3 bis 5, wobei R ausgewählt ist aus H oder Alkyl C₁-C₆, vorzugsweise Methyl.
7. Verwendung nach Anspruch 1, **dadurch gekennzeichnet, dass** die Verbindung mit der Formel (I) ein Verbindung mit der Formel (III) ist:



(III)

40 wobei

45 X ausgewählt ist aus O oder S und Y ausgewählt ist aus O-Aryl oder S-Alkyl; und
 R und R¹ unabhängig ausgewählt sind aus der Gruppe bestehend aus Wasserstoff, Halogen, Alkyl, Haloalkyl, Aryl, Cycloalkyl, (CH₂)_n-Aryl, -OR³; -C(O)OR³, (CH₂)_n-C(O)OR³ oder -S(O)_t, wobei R³ ausgewählt ist aus Wasserstoff, Alkyl, Aryl oder Cycloalkyl, n größer oder gleich 0 ist und t 1 oder 2 ist.

8. Verwendung nach Anspruch 7, wobei X O ist.
9. Verwendung nach Anspruch 7, wobei X S ist.
- 50 10. Verwendung nach einem der Ansprüche 7 bis 9, wobei Y S-Alkyl C₁-C₆, vorzugsweise S-CH₃, ist.
11. Verwendung nach einem der vorhergehenden Ansprüche, wobei R ausgewählt ist aus H oder Alkyl C₁-C₆, vorzugsweise Methyl.
- 55 12. Verwendung nach einem der vorhergehenden Ansprüche, wobei R¹ substituiertes oder unsubstituiertes Phenyl ist.
13. Verwendung nach einem der vorhergehenden Ansprüche, wobei R¹ Phenyl ist, das mit wenigstens einem Halogen


ausgewählt aus Br oder F substituier ist.

14. Verwendung nach Anspruch 1, **dadurch gekennzeichnet, dass** die Verbindung mit der Formel (I) eine Verbindung ist, die ausgewählt ist aus der folgenden Gruppe:

- 3-Phenyl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazolin,
- 3-(2,6-Difluorphenyl)-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazolin,
- 3-(2-Bromphenyl)-8-methyl-4-oxo-2-thioxo-1,2,3,4-tetrahydroquinazolin,
- 3-(2,6-Difluorphenyl)-8-methyl-2-methylthio-4-oxo-3,4-dihydroquinazolin,
- 3-(2-Bromphenyl)-8-methyl-2-methylthio-4-oxo-3,4-dihydroquinazolin,
- 3-Phenyl-2,4-dithioxo-1,2,3,4-tetrahydroquinazolin, und
- 3-Phenyl-2-methylthio-4-thioxo-3,4-dihydroquinazolin
- 3-(2,6-Difluorphenyl)-2,4-dithioxo-1,2,3,4-tetrahydroquinazolin.

Revendications

1. Utilisation d'un composé de formule (I) :



dans laquelle :

A représente un carbocycle ou un hétérocycle condensé éventuellement substitué par 5, 6 ou 7 chaînons saturé ou insaturé ;

--- peut représenter une double liaison ;

X et Y sont choisis indépendamment dans le groupe constitué d'un atome d'hydrogène, des groupes alkyle, =O, =S, aryle, 0-alkyle, 0-aryle, S-alkyle ou -S-aryle ; et


R, R¹, R² et R³ sont choisis indépendamment dans le groupe constitué d'un atome d'hydrogène, d'halogène, des groupes alkyle, halogénoalkyle, aryle, cycloalkyle, (CH₂)_n-aryle, -OR³, -C(O)OR³, (CH₂)_n-C(O)OR³ ou -S(O)_t-, où n est supérieur ou égal à 0 et où t vaut 1 ou 2,

ou un sel pharmaceutiquement acceptable, un précurseur, un solvate ou un stéréoisomère de celui-ci, où le précurseur est un ester d'acide carboxylique, un ester d'acide aminé, un ester phosphate, un ester sulfonate de sel métallique, un carbamate ou un amide dérivé du composé de formule (I) qui, lorsqu'il est administré à un individu, est capable de fournir, directement ou indirectement, ledit composé de formule (I),

pour produire un médicament destiné au traitement de maladies neurodégénératives et/ou neurologiques, **caractérisée en ce que** la maladie neurodégénérative et/ou neurologique est choisie dans le groupe suivant : la maladie d'Alzheimer, la maladie de Parkinson, la sclérose latérale amyotrophique, l'ischémie cérébrale, les parkinsonismes post-encéphaliques, les dystonies, le syndrome de Tourette, les troubles des mouvements périodiques des membres, le syndrome des jambes sans repos, les troubles de déficit de l'attention avec hyperactivité.

2. Utilisation selon la revendication 1, où A représente un carbocycle avec 6 chaînons, de préférence un cycle phényle.

3. Utilisation selon la revendication 1, **caractérisé en ce que** le composé de formule (I) est un composé de formule (II) :




dans laquelle

15 X et Y sont choisis indépendamment entre O et S ; et

R et R¹ sont choisis indépendamment dans le groupe constitué d'un atome d'hydrogène, d'halogène, des groupes alkyle, halogénoalkyle, aryle, cycloalkyle, (CH₂)_n-aryle, hétéroaryle, -OR³, -C(O)OR³, (CH₂)_n-C(O)OR³ ou -S(O)_t-, où R³ est choisi entre un atome d'hydrogène, un groupe alkyle, aryle ou cycloalkyle, n est supérieur ou égal à 0 et où t vaut 1 ou 2.

- 20
4. Utilisation selon la revendication 3, où X représente O et Y représente S.
5. Utilisation selon la revendication 3, où X représente S et Y représente S.
- 25
6. Utilisation selon l'une quelconque des revendications 3 à 5, où R est choisi entre H ou un groupe alkyle en C₁ à C₆, de préférence un groupe méthyle.
7. Utilisation selon la revendication 1, **caractérisée en ce que** le composé de formule (I) est un composé de formule (III) :



40 dans laquelle :

X est choisi entre O et S et Y est choisi entre un groupe O-aryle ou S-alkyle ; et

45 R et R¹ sont choisis indépendamment dans le groupe constitué d'un atome d'hydrogène, d'halogène, des groupes alkyle, halogénoalkyle, aryle, cycloalkyle, (CH₂)_n-aryle, -OR³, -C(O)OR³, (CH₂)_n-C(O)OR³ ou -S(O)_t-, où R³ est choisi entre un atome d'hydrogène, un groupe alkyle, aryle ou cycloalkyle, n est supérieur ou égal à 0 et où t vaut 1 ou 2.

8. Utilisation selon la revendication 7, où X représente O.
- 50
9. Utilisation selon la revendication 7, où X représente S.
10. Utilisation selon l'une quelconque des revendications 7 à 9, où Y représente un groupe S-alkyle en C₁ à C₆, de préférence un groupe S-CH₃.
- 55
11. Utilisation selon l'une quelconque des revendications précédentes, où R est choisi entre H ou un groupe alkyle en C₁ à C₆, de préférence un groupe méthyle.
12. Utilisation selon l'une quelconque des revendications précédentes, où R¹ représente un cycle phényle substitué ou

non substitué.

13. Utilisation selon la revendication précédente, où R¹ représente un groupe phényle substitué par au moins un atome d'halogène choisi entre Br ou F.

5

14. Utilisation selon la revendication 1, **caractérisée en ce que** le composé de formule (I) est un composé choisi dans le groupe suivant :

10

la 3-phényl-4-oxo-2-thio-1,2,3,4-tétrahydroquinazoline,
la 3-(2,6-difluorophényl)-4-oxo-2-thio-1,2,3,4-tétrahydroquinazoline,
la 3-(2-bromophényl)-8-méthyl-4-oxo-2-thio-1,2,3,4-tétrahydroquinazoline,
la 3-(2,6-difluorophényl)-8-méthyl-2-méthylthio-4-oxo-3,4-dihydroquinazoline,
la 3-(2-bromophényl)-8-méthyl-2-méthylthio-4-oxo-3,4-dihydroquinazoline,
la 3-phényl-2,4-dithio-1,2,3,4-tétrahydroquinazoline,
la 3-phényl-2-méthylthio-4-thio-3,4-dihydroquinazoline, et
la 3-(2,6-difluorophényl)-2,4-dithio-1,2,3,4-tétrahydroquinazoline.

15

20

25

30

35



40

45

50

55

FIG. 1








FIG. 2







FIG. 3






FIG. 4

A)



B)



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20080260643 A [0005]
- WO 2008113881 A [0005]

Non-patent literature cited in the description

- **DAUER, W. ; PRZEDBORSKI, S.** Parkinson's disease: mechanisms and models. *Neuron*, 2003, vol. 39, 889-909 [0002]
- **LANG, AE. ; LOZANO, AM.** Parkinson's disease. First of two parts. *The New England Journal of Medicine*, 1998, vol. 339, 1044-1053 [0003]
- **LANG, AE. ; LOZANO, AM.** Parkinson's disease. Second of two parts. *The New England Journal of Medicine*, 1998, vol. 339, 1130-1143 [0003]
- **SAVITT, JM. ; DAWSON, VL. ; DAWSON, TM.** Diagnosis and treatment of Parkinson disease: molecules to medicine. *J Clin Invest*, 2006, vol. 116, 1744-1754 [0004]
- **LEWITT, P.A.** Levodopa therapeutics for Parkinson's disease: new developments. *Parkinsonism Relat Disord.*, 2009, vol. 1, 31-4 [0004]
- **ABDEL-SALAM, O.M.** Drugs used to treat Parkinson's disease, present status and future directions. *CNS Neurol Disord Drug Targets.*, 2008, vol. 7, 321-442 [0004]
- **GALLAGHER, D. A. ; SCHRAG, A.** Impact of newer pharmacological treatments on quality of life in patients with Parkinson's disease. *CNS Drugs*, 2008, vol. 22, 563-586 [0004]
- **KIEBURTZ, K.** Therapeutic strategies to prevent motor complications in Parkinson's disease. *J Neurol.*, 2008, vol. 255 (4), 42-45 [0004]
- **POEWE, W.** Treatments for Parkinson disease - past achievements and current clinical needs. *Neurology*, vol. 72 (7), 65-73 [0005]
- **CASTAÑO, T. ; WANG, H. ; CAMPILLO, N.E ; BALLESTER, S. ; GONZÁLEZ-GARCÍA, C. ; HERNÁNDEZ, J. ; PÉREZ, C. ; CUENCA, J. ; ; PÉREZ-CASTILLO, A. ; MARTÍNEZ, A.** Synthesis, structural analysis, and biological evaluation of thioxoquinazoline derivatives as phosphodiesterase 7 inhibitors. *ChemMedChem.*, 2009, vol. 4, 866-876 [0005]
- Treaty of Galenic Pharmacy. 1993 [0034]
- **LUNA-MEDINA, R. ; CORTES-CANTELI, M. ; ALONSO, M ; SANTOS, A. ; MARTINEZ, A. ; PE-REZ-CASTILLO, A.** Regulation of inflammatory response in neural cells in vitro by thiadiazolidinones derivatives through peroxisome proliferator-activated receptor gamma activation. *J. Biol. Chem.*, 2005, vol. 280, 21453-21462 [0042]
- **KRONCKE K. D. ; FEHSEL K. ; KOLB-BACHOFEN V.** Nitric oxide: cytotoxicity versus cytoprotection--how, why, when, and where?. *Nitric Oxide*, 1997, vol. 1, 107-120 [0043]
- **GRIESS, P.** Bemerkungen zu der abhandlung der H.H. Weselsky und Benedikt Ueber einige azo-verbindingen. *Chem. Ber.*, vol. 1879 (12), 426-428 [0043]
- **MENDEZ, J.S. ; FINN, B.W.** Use of 6-hydroxy-dopamine to create lesions in catecholamine neurons in rats. *J Neurosurg.*, 1975, vol. 42, 166-173 [0044]