



(11) **EP 2 433 940 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Claims EN 1, 2, 3, 10, 11
Numerous spelling errors of minor importance

(48) Corrigendum issued on:
28.10.2015 Bulletin 2015/44

(45) Date of publication and mention
of the grant of the patent:
24.09.2014 Bulletin 2014/39

(21) Application number: **10769728.6**

(22) Date of filing: **27.04.2010**

(51) Int Cl.:
C07D 471/10 ^(2006.01) **A61K 31/438** ^(2006.01)
A61K 31/444 ^(2006.01) **A61K 31/496** ^(2006.01)
A61K 31/5355 ^(2006.01) **A61P 5/20** ^(2006.01)
A61P 7/08 ^(2006.01) **A61P 19/02** ^(2006.01)
A61P 19/08 ^(2006.01) **A61P 43/00** ^(2006.01)
C07D 487/10 ^(2006.01)

(86) International application number:
PCT/JP2010/057432

(87) International publication number:
WO 2010/126030 (04.11.2010 Gazette 2010/44)

(54) **SPIROIMIDAZOLONE DERIVATIVE**

SPIROIMIDAZOLONDERIVAT

DÉRIVÉ DE SPIROIMIDAZOLONE

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL
PT RO SE SI SK SM TR

(30) Priority: **28.04.2009 JP 2009109256**

(43) Date of publication of application:
28.03.2012 Bulletin 2012/13

(73) Proprietor: **Chugai Seiyaku Kabushiki Kaisha**
Kita-ku
Tokyo 115-8543 (JP)

(72) Inventors:
• **ESAKI, Toru**
Gotenba-shi
Shizuoka 412-8513 (JP)
• **NISHIMURA, Yoshikazu**
Gotenba-shi
Shizuoka 412-8513 (JP)
• **ISSHIKI, Yoshiaki**
Gotenba-shi
Shizuoka 412-8513 (JP)
• **OKAMOTO, Naoki**
Gotenba-shi
Shizuoka 412-8513 (JP)

- **FURUTA, Yoshiyuki**
Gotenba-shi
Shizuoka 412-8513 (JP)
- **MIZUTANI, Akemi**
Gotenba-shi
Shizuoka 412-8513 (JP)
- **OHTA, Masateru**
Gotenba-shi
Shizuoka 412-8513 (JP)
- **LAI, Wayne Wen**
Wuxi City
Jiangsu Province 214135 (CN)
- **KOTAKE, Tomoya**
Gotenba-shi
Shizuoka 412-8513 (JP)

(74) Representative: **Vossius & Partner**
Patentanwälte Rechtsanwälte mbB
Siebertstrasse 3
81675 München (DE)

(56) References cited:
WO-A1-2007/135417 WO-A1-2008/148689
WO-A2-2007/149873 WO-A2-2009/074575
JP-T- 2004 523 583 JP-T- 2008 515 895

EP 2 433 940 B9

DescriptionTechnical Field

5 **[0001]** The present invention relates to spiroimidazolone derivatives and use thereof.

Background Art

10 **[0002]** Parathyroid hormone (PTH) is a major regulator of calcium homeostasis and its main target organs are considered to be the bones and kidneys. Native human parathyroid hormone is a polypeptide consisting of 84 amino acids. This hormone is secreted from the parathyroid gland in response to low blood calcium levels, and acts on osteoblasts (bone-building cells) in the bones and tubular epithelial cells in the kidneys. This hormone interacts with a cell surface receptor molecule called PTH-1 receptor or PTH/PTHrP receptor, which is expressed by both osteoblasts and renal tubular cells.

15 **[0003]** PTHrP (PTH-related protein), the major cause of humoral hypercalcemia of malignancy (HHM), also has normal functions including developmental roles. PTHrP has 141 amino acids, although mutants also occur that result from alternative gene splicing mechanisms. PTHrP plays a key role in the formation of the skeleton through a process that also involves binding to the PTH-1 receptor (Non Patent Literature 1, Non Patent Literature 2).

20 **[0004]** Regulation of calcium concentrations is necessary for normal functions of the gastrointestinal system, skeletal system, nervous system, neuromuscular system and cardiovascular system. Synthesis and release of PTH are primarily controlled by the serum calcium level. Synthesis and release of PTH are stimulated at low serum calcium levels, and synthesis and release of PTH are suppressed at high serum calcium levels. PTH, in turn, maintains the serum calcium level by directly or indirectly promoting calcium entry into the blood at three calcium exchange sites: intestine, bone and kidney. PTH contributes to net gastrointestinal absorption of calcium by assisting in the renal synthesis of active vitamin D. PTH promotes calcium mobilization from the bone to serum by stimulating differentiation of osteoclasts that are bone-resorbing cells. This also mediates at least three main effects in the kidney (stimulation of tubular calcium resorption; enhancement of phosphate clearance; and promotion of an increase in the enzyme that completes the synthesis of active vitamin D). PTH is considered to exert these effects primarily through receptor-mediated activation of adenylate cyclase and/or phospholipase C.

25 **[0005]** Disruption of calcium homeostasis may produce many clinical disorders (e.g., serious bone disease, anemia, renal dysfunction, ulcers, myopathy and neuropathy), and this usually results from conditions that produce an alteration in the level of parathyroid hormone. Hypercalcemia is a condition characterized by an elevated serum calcium level. This is often associated with primary hyperparathyroidism in which excessive PTH production occurs as a result of parathyroid gland lesions (e.g., adenoma, hyperplasia or carcinoma). Humoral hypercalcemia of malignancy (HHM), another type of hypercalcemia, is the most common paraneoplastic syndrome. This appears to result in most instances from the production of a certain protein hormone that shares amino acid homology with PTH by tumors (e.g., squamous cell carcinoma, renal carcinoma, ovarian carcinoma or bladder carcinoma). These PTHrPs appear to mimic the effects of PTH on the kidney and skeleton in some degree, and are considered to interact with the PTH receptor in these tissues. PTHrP is usually found at low levels in many tissues including keratinocytes, brain, pituitary gland, parathyroid gland, adrenal cortex, medulla, fetal liver, osteoblast-like cells and lactating mammary tissues. For many HHM malignant tumors, high levels of PTHrP are observed in the circulatory system, and this leads to elevated calcium levels associated with HHM.

35 **[0006]** The pharmacological profiles of PTH and PTHrP are nearly identical in most in vitro assay systems, and elevated blood levels of PTH (i.e., primary hyperparathyroidism) or PTHrP (i.e., HHM) have comparable effects on inorganic ion homeostasis (Non Patent Literature 3, Non Patent Literature 4). The similarities in the biological activities of the two ligands can be explained by their interaction with the PTH/PTHrP receptor, a common receptor expressed abundantly in the bones and kidneys (Non Patent Literature 5).

40 **[0007]** The PTH-1 receptor is homologous in primary structure to some other receptors binding to peptide hormones, such as secretin (Non Patent Literature 6), calcitonin (Non Patent Literature 7) and glucagon (Non Patent Literature 8); these receptors together form a distinct family called receptor family B (Non Patent Literature 9). Within this family, the PTH-1 receptor is unique in that it binds to two peptide ligands and thereby regulates two separate biological processes. A recently identified PTH receptor subtype called PTH-2 receptor binds to PTH but not to PTHrP (Non Patent Literature 10). This finding has implied that the structural differences in the PTH and PTHrP ligands determine the selectivity for interaction with the PTH-2 receptor. The PTH-2 receptor has been detected by RNA methods in the brain, pancreas and vasculature; however, its biological functions have not been determined (Non Patent Literature 10). The family B receptors are assumed to use a common molecular mechanism for engagement with their own cognate peptide hormone (Non Patent Literature 11).

55 **[0008]** The PTH-1 receptor binds to both PTH and PTHrP and causes not only intracellular cAMP accumulation and adenylyl cyclase (AC) activation but also signal transduction to phospholipase C (PLC), thereby leading to the production

of inositol trisphosphate (IP₃), diacylglycerol (DAG) and intracellular calcium (iCa²⁺) (Non Patent Literature 12, Non Patent Literature 13).

[0009] Osteoporosis is a potentially crippling bone disease and is observed in a substantial portion of the elderly population, in pregnant women and even in juveniles. The term "osteoporosis" refers to a group of disorders consisting of different constituents. Osteoporosis is clinically classified into type I and type II. Type I osteoporosis occurs primarily in middle-aged women and is associated with menopausal estrogen loss, while type II osteoporosis is associated with the elderly. Patients with osteoporosis are considered to benefit from novel therapies designed to promote fracture repair, or therapies designed to prevent or reduce fractures associated with the disease.

[0010] This disease is characterized by reduced bone mass, decreased bone mineral density (BMD), decreased bone strength and an increased risk of fracture. Currently, there is no effective cure for osteoporosis, although estrogen, calcitonin, and etidronate and alendronate that are bisphosphonates are used to treat the disease with various levels of success. These agents act to decrease bone resorption.

[0011] PTH(1-34) (teriparatide) has a strong bone anabolic effect and induces significant increases in bone mineral density and bone strength. Subcutaneous administration of human PTH(1-34) increased the spine bone mineral density (BMD) by 8% in one year and decreased the risks of vertebral fracture and nonvertebral fracture by 65% and 55% in two years, respectively (Non Patent Literature 14). Subcutaneous administration of human PTH(1-84) also increased the spine bone mineral density (BMD) by 6.9% in 18 months and decreased the risk of vertebral fracture by 58% (Non Patent Literature 15). Parathyroid hormone is currently believed to be one of the most effective treatments for osteoporosis (Non Patent Literature 16). Importantly, hPTH(1-34) must be administered in a pulsed manner (e.g., subcutaneous injection once daily) to achieve its bone-forming effect. Longer administration of PTH(1-34) such as by use of a continuous infusion pump mechanism activates bone-resorptive responses mediated by osteoclasts much stronger than bone-forming responses mediated by osteoblasts, and thus PTH(1-34) exerts a net degradation effect on the bone.

[0012] Although parathyroid hormone is believed to be one of the most effective treatments for osteoporosis, only less than 1% of patients with osteoporosis use teriparatide and the average duration of teriparatide is 12 months (Non Patent Literature 16). Teriparatide must be administered by self-injection. The fact that it is difficult to use a pen-type device for self-administration is the principal cause of the low compliance of teriparatide-administered patients. It is obvious that noninvasively, preferably orally, available compounds having a PTH-like effect with clinical efficacy in osteoporosis similar to that of parathyroid hormone can considerably improve the compliance of patients with regard to the administration, and that the compounds can be the most useful therapeutic option for patients with osteoporosis.

[0013] There are many low molecular weight agonists for the GPCR family A; however, only a limited number of low molecular weight ligands for the GPCR family B have been reported (Non Patent Literature 17). Low molecular weight agonists have been reported for the GLP-1 receptor, calcitonin receptor and PTH1 receptor belonging to the GPCR family B; however, there is no compound used in clinical applications for the treatment of diseases.

[Citation List]

[Non Patent Literature]

[0014]

[Non Patent Literature 1] Karaplis, A. C. et al., Genes and Dev. 8:277-289 (1994)

[Non Patent Literature 2] Lanske, B. et al., Science 273:663-666 (1996)

[Non Patent Literature 3] Broadus, A. E. & Stewart, A. F., "Parathyroid hormone-related protein: Structure, processing and physiological actions," in Basic and Clinical Concepts, Bilzikian, J. P. et al., eds., Raven Press, New York (1994), pp. 259-294

[Non Patent Literature 4] Kronenberg, H. M. et al., "Parathyroid hormone: Biosynthesis, secretion, chemistry and action," in Handbook of Experimental Pharmacology, Mundy, G. R. & Martin, T. J., eds., Springer-Verlag, Heidelberg (1993), pp. 185-201

[Non Patent Literature 5] Urena, P. et al., Endocrinology 134:451-456 (1994)

[Non Patent Literature 6] Ishihara, T. et al., EMBO J. 10:1635-1641 (1991)

[Non Patent Literature 7] Lin, H. Y. et al., Science 254:1022-1024 (1991)

[Non Patent Literature 8] Jelinek, L. J. et al., Science 259:1614-1616 (1993)

[Non Patent Literature 9] Kolakowski, L. F., Receptors and Channels 2:1-7 (1994)

[Non Patent Literature 10] Usdin, T. et al., J. Biol. Chem. 270:15455-15458 (1995)

[Non Patent Literature 11] Bergwitz, C. et al., J. Biol. Chem. 271:26469-26472 (1996)

[Non Patent Literature 12] Abou-Samra, A. B et al., Pro. Natl. Acad. Sci. USA, 89:2732-2736, 1992

[Non Patent Literature 13] Bringham F. R. et al., Endocrinology 132:2090-2098, 1993


[Non Patent Literature 14] Neer RM et al., N. Eng. J. Med. 344:1434-1441, 2003

- i) a hydrogen atom and
- ii) C1-C10 alkyl optionally substituted with a halogen atom(s);

Y is selected from:

- 1) an oxygen atom,
- 2) a sulfur atom and
- 3) =NR₃₇,

or 4) Y is -NR₃₈R₃₉ shown in the following formula (A):



which can be tautomeric;

R₃₇ is selected from:

- 1) hydrogen,
- 2) hydroxy and
- 3) C1-C10 alkoxy;

R₃₈ and R₃₉ are independently selected from hydrogen or C1-C10 alkyl optionally substituted with cycloalkyl, or R₃₈ and R₃₉ may be bonded to each other to form a ring selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with C1-C10 alkyl; m represents an integer of 0 to 2;

n represents an integer of 0 to 2;

R₁ is selected from:

- 1) hydrogen,
- 2) cycloalkyl optionally substituted with a group(s) selected from R₄, R₅ and R₆,
- 3) a heterocycle optionally substituted with a group(s) selected from R₂₅, R₄, R₅ and R₆,
- 4) aryl optionally substituted with a group (s) selected from R₃, R₄, R₅ and R₆ and
- 5) heteroaryl optionally substituted with a group(s) selected from R₂₅, R₄, R₅ and R₆;

R₃ is selected from:

- 1) -CONR₇R₈,
- 2) -OR₉,
- 3) -NR₉R₁₀,
- 4) -N(R₉)COR₁₁,
- 5) -N(R₉) SO₂R₁₂,
- 6) -SO₂R₁₅,
- 7) C1-10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, -COR₁₆ and -NR₁₃R₁₄,
- 8) heteroaryl optionally having C1-10 alkyl and/or C1-10 alkoxy as a substituent and
- 9) -N(R₉)CSR₁₁;

R₄ is selected from:

- 1) a halogen atom,
- 2) cyano,
- 3) nitro,

- 4) amino,
- 5) -NHCOR₂₆,
- 6) C1-C10 alkyl optionally substituted with a group(s) independently selected from hydroxycarbonyl, C1-C10 alkoxy, carbonyl and aminocarbonyl,
- 7) C1-C10 haloalkyl,
- 8) C1-C10 alkoxy,
- 9) C1-C10 haloalkylcarbonyl,
- 10) -COR₁₆,
- 11) C1-C10 hydroxyalkyl and
- 12) C1-C10 heteroalkyl;

R₅ is selected from a halogen atom, C1-C10 alkyl, C1-C10 haloalkyl and C1-C10 alkoxy;

R₆ is selected from a halogen atom, C1-C10 alkyl and C1-C10 haloalkyl;

R₇ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from amino and C1-C10 alkylamino,
- 3) C1-C10 hydroxyalkyl,
- 4) C1-C10 haloalkyl,
- 5) C1-C10 heteroalkyl,
- 6) C1-C10 heteroalkyl optionally substituted with a group(s) selected from a hydroxyl group, C1-C10 alkylamino and C2-C10 alkenyl,
- 7) aryl,
- 8) heteroaryl,
- 9) aryl C1-C10 alkyl,
- 10) a heterocycle optionally substituted with C1-C10 alkyl,
- 11) - (CH₂)_L COR₁₆ (wherein L represents an integer of 1 to 4),
- 12) C1-C10 alkoxy,
- 13) C2-C10 alkenyl and
- 14) -NR₄₀R₄₁;

R₄₀ and R₄₁ are independently selected from hydrogen, C1-C10 alkyl and C1-C10 alkylcarbonyl, or R₄₀ and R₄₁ may be bonded to each other to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholinyl, and the heterocycle is optionally substituted with C1-C10 alkyl;

R₈ is selected from hydrogen and C1-C10 alkyl optionally substituted with a halogen atom(s) and/or a hydroxyl group(s);

R₇ and R₈ may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from O, N, S, SO and SO₂, and the heterocycle optionally contains carbonyl, and the heterocycle is optionally substituted with a substituent(s) independently selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally having C1-C10 alkylamino as a substituent,
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 8) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 9) a heterocycle optionally substituted with C1-C10 alkyl,
- 10) heteroaryl optionally substituted with C1-C10 alkyl,
- 11) heterocyclyl C1-C10 alkyl,
- 12) -COR₁₆,
- 13) -NR₁₉R₂₀,
- 14) -SO₂R₂₁,

- 15) C1-C10 alkoxy-C1-C10 alkyl optionally having a hydroxyl group(s) as a substituent(s) and
 16) C1-C10 hydroxyalkoxy, wherein the hydrogen atom of the hydroxyl group is optionally replaced by C1-C10 hydroxyalkyl, and

the heterocycle may further form a spiro ring together with a 4- to 6-membered heterocycle, and the bonded 4- to 6-membered heterocycle optionally contains O and N as ring-forming elements in addition to carbon atoms, and the carbon atom(s) may be oxidized to form carbonyl, and the 4- to 6-membered heterocycle is optionally further substituted with C1-C10 alkyl;

R₁₆ is selected from:

- 1) a hydroxyl group,
- 2) C1-C10 alkoxy,
- 3) NR₁₇R₁₈ and
- 4) C1-C10 alkyl optionally substituted with a substituent(s) selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, carbonyl or C1-C10 alkylamino;

R₁₇ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) selected from aryl, amino, C1-C10 alkylamino, C1-C10 alkylcarbonylamino and a hydroxyl group,
- 3) heteroaryl and
- 4) C1-C10 alkoxy;

R₁₈ is selected from hydrogen, C1-C10 alkyl and C1-C10 hydroxyalkyl;

R₁₇ and R₁₈ may be bonded to each other to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl, a halogen atom and C1-C10 alkoxy, carbonyl;

R₁₉ is selected from hydrogen, C1-C10 alkyl, C1-C10 haloalkyl, C1-C10 alkylcarbonyl, C1-C10 hydroxyalkyl, C1-C10 aminoalkyl, C1-C10 alkoxy, carbonyl and C1-C10 heteroalkyl;

R₂₀ is selected from hydrogen and C1-C10 alkyl;

R₁₉ and R₂₀ may be bonded to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom;

R₂₁ is selected from:

- 1) C1-C10 alkyl optionally substituted with aryl,
- 2) amino,
- 3) C1-C10 alkylamino and
- 4) aryl optionally substituted with C1-C10 alkyl;

R₉ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from R₂₃,
- 3) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 4) cycloalkyl optionally substituted with a halogen atom(s) or a hydroxyl group(s),
- 5) a heterocycle optionally substituted with a group(s) independently selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy, C1-C10 alkoxy, carbonyl, amino and a halogen atom,
- 6) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom and a hydroxyl group,
- 7) heteroaryl optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy, carbonyl and a halogen atom and
- 8) cycloalkenyl optionally substituted with a group(s) selected from C1-C10 alkoxy, C1-C10 alkylamino, amino, a hydroxyl group and a halogen atom, wherein the cycloalkenyl optionally contains a carbonyl group;

R₂₃ is independently selected from:

- 1) a halogen atom,
- 2) a hydroxyl group,
- 3) a C1-C10 alkylcarbonyloxy group,
- 4) -COR₁₆,
- 5) amino,
- 6) C1-C10 alkylamino,
- 7) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy carbonyl and a halogen atom and
- 8) cyano;

R₁₀ is selected from:

- 1) hydrogen and
- 2) C1-C10 alkyl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group and aryl;

R₉ and R₁₀ may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from N, O, S, SO, SO₂, carbonyl and thiocarbonyl, and the heterocycle is optionally substituted with a substituent(s) independently selected from R₂₄;

R₂₄ is independently selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from C1-C10 alkylamino and C1-C10 alkylcarbonylamino,
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 8) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 9) a heterocycle optionally substituted with C1-C10 alkyl,
- 10) heteroaryl,
- 11) heterocyclyl C1-C10 alkyl,
- 12) -COR₁₆,
- 13) -NR₁₉R₂₀ and
- 14) -SO₂R₂₁;

R₁₁ is selected from:

- 1) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a hydroxyl group,
- ii) -NR₁₇R₁₈,
- iii) a C1-C10 alkoxy group,
- iv) a halogen atom,
- v) C1-C10 alkoxy carbonyl,
- vi) aminocarbonyl and
- vii) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, amino, C1-C10 alkylamino and -COR₂₂,

- 2) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, amino, C1-C10 alkylamino and -COR₂₂,
- 3) cycloalkyl optionally substituted with a halogen atom(s),
- 4) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy carbonyl and a halogen atom,

- 5) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino and a hydroxyl group,
 6) amino,
 7) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino, hydroxycarbonyl and a hydroxyl group and
 8) C2-C10 alkenyl;

R₂₂ is selected from C1-C10 alkoxy, a hydroxyl group, amino and C1-C10 alkylamino;

R₁₂ is selected from:

- 1) C1-C10 alkyl,
 2) amino and
 3) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from amino, C1-C10 alkylamino and a hydroxyl group;

R₁₃ is selected from:

- 1) hydrogen,
 2) C1-C10 alkyl,
 3) C1-C10 alkylcarbonyl, wherein the alkyl is optionally substituted with a hydroxyl group(s),
 4) C1-C10 alkoxycarbonyl,
 5) aminocarbonyl,
 6) C1-C10 alkylaminocarbonyl and
 7) heterocyclic carbonyl optionally substituted with C1-C10 alkyl;

R₁₄ is selected from:

- 1) hydrogen and
 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino;

R₁₃ and R₁₄ may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from O, N, S, SO and SO₂, and the heterocycle optionally contains carbonyl, and the heterocycle is optionally substituted with C1-C10 alkyl;

R₁₅ is selected from:

- 1) C1-C10 alkyl and
 2) -NR₃₅R₃₆;

R₃₅ is selected from:

- 1) hydrogen,
 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
 ii) a hydroxyl group,
 iii) C1-C10 alkylcarbonylamino,
 iv) -COR₁₆,
 v) amino,
 vi) C1-C10 alkylamino,
 vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),
 viii) heteroaryl optionally substituted with a C1-C10 alkyl group(s) and
 ix) a heterocycle,

3) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,

4) cycloalkyl optionally substituted with a group(s) independently selected from a halogen atom and a

hydroxyl group,

5) a heterocycle optionally substituted with a group(s) independently selected from C1-C10 alkyl, a halogen atom and aryl C1-C10 alkyl,

6) heteroaryl optionally substituted with C1-C10 alkyl and

7) C1-C10 alkylcarbonyl;

R₃₆ is selected from:

1) hydrogen and

2) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group and aryl;

R₃₅ and R₃₆ may be bonded to each other to form a ring selected from azetidiny, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom;

R₂₅ is selected from:

1) a halogen atom,

2) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

i) a halogen atom,

ii) aryl,

iii) heteroaryl,

iv) a heterocycle optionally substituted with a C1-C10 alkyl group(s),

v) -COR₁₆,

vi) -NR₁₃R₁₄ and

vii) -SO₂R₂₁,

3) C1-C10 heteroalkyl optionally substituted with a hydroxyl group(s),

4) C1-C10 hydroxyalkyl, wherein each hydroxyl group may be independently substituted with a group(s) selected from C1-C10 alkyl, aryl C1-C10 alkyl and C1-C10 alkylcarbonyl,

5) -COR₁₆,

6) -SO₂R₂₁,

7) aryl and

8) cyano;

R₂ is selected from:

1) C1-C10 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a substituent(s) independently selected from R₄₂,

2) C2-C10 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a substituent(s) independently selected from R₄₂,

3) C2-C10 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a substituent(s) independently selected from R₄₂,

4) cycloalkyl optionally substituted with a group(s) independently selected from:

i) a halogen atom,

ii) C2-C10 alkenyl or C1-C10 alkyl,

iii) aryl optionally substituted with 1 to 3 substituents independently selected from C1-C10 alkyl, a halogen atom, C1-C10 alkoxy, C1-C10 alkylamino and C1-C10 alkylcarbonyl,

iv) cycloalkyl,

v) C2-C10 alkenyl optionally substituted with halogen,

vi) C1-C10 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with a halogen atom(s),

vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),

viii) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom or C1-C10 alkoxy optionally substituted with a halogen atom(s),

ix) C2-C10 alkynyl and

x) $-\text{Si}(\text{R}_{43})_3$,

5) a heterocycle, wherein the heterocycle is optionally substituted with a group(s) independently selected from:

- i) a C1-C10 alkyl group,
- ii) C1-C10 alkylcarbonyl, wherein the alkyl group is optionally substituted with R_{27} ,
- iii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- iv) heteroarylcarbonyl,
- v) C1-C10 alkoxycarbonyl, wherein the alkyl group is optionally substituted with a group(s) independently selected from a halogen atom, aryl and C1-C10 alkoxy,
- vi) aryloxy carbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) and/or C1-C10 alkyl,
- vii) $-\text{CONR}_{28}\text{R}_{29}$,
- viii) $-\text{SO}_2\text{R}_{21}$,
- ix) a halogen atom,
- x) cycloalkylcarbonyl optionally fused with an aryl group and
- xi) C2-C10 alkenylcarbonyl, wherein the alkenyl group is optionally substituted with aryl, wherein the aryl is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl or C1-C10 alkoxy,

6) aryl optionally substituted with a group(s) independently selected from R_{44} ,

7) heteroaryl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) C1-C10 alkyl and
- iii) C1-C10 alkoxy;

8) C1-C10 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a substituent(s) independently selected from R_{42} ,

9) $-\text{S}(\text{O})_q\text{R}_{43}$ (wherein q is an integer of 0 to 2) and

10) cycloalkenyl optionally substituted with C1-C10 alkyl;

R_{44} is selected from:

- 1) a halogen atom,
- 2) cyano,
- 3) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a hydroxyl group,
- ii) $-\text{OR}_{26}$,
- iii) cyano,
- iv) aryloxy optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl optionally substituted with a halogen atom(s) or C1-C10 alkoxy optionally substituted with a halogen atom(s) and
- v) a halogen atom,

4) cycloalkyl optionally substituted with a group(s) independently selected from a halogen atom or C1-C10 alkyl optionally substituted with a halogen atom(s),

5) C1-C10 alkoxy optionally substituted with a halogen atom(s) or a C2-C6 alkenyl group,

6) $-\text{COR}_{30}$,

7) C1-C10 alkylcarbonylamino,

8) C1-C10 alkoxycarbonylamino, wherein the alkoxy group is optionally substituted with aryl,

9) C1-C10 heteroalkyl optionally substituted with a halogen atom(s),

10) aryl optionally substituted with a substituent(s) independently selected from:

- i) a halogen atom,

- ii) C1-C10 alkyl,
- iii) C1-C10 alkoxy and
- iv) aryl optionally substituted with aryl optionally substituted with C1-C10 alkyl,

- 5 11) heteroaryl optionally substituted with a C1-C10 alkyl group(s),
- 12) $-\text{SO}_2\text{R}_{43}$,
- 13) $-\text{SOR}_{43}$,
- 14) C1-C10 alkylthio optionally substituted with a halogen atom(s),
- 15) $-\text{Si}(\text{R}_{43})_3$ and
- 10 16) $-\text{SF}_5$;

R_{42} is selected from:

- 1) hydrogen,
- 15 2) aryl optionally substituted with a' group(s) independently selected from C1-C10 alkyl optionally substituted with halogen, a halogen atom and C1-C10 alkoxy,
- 3) hydroxycarbonyl,
- 4) C1-C10 alkoxycarbonyl,
- 5) aminocarbonyl,
- 20 6) C1-C10 alkylaminocarbonyl,
- 7) C1-C10 alkoxycarbonylamino,
- 8) amino,
- 9) a hydroxyl group and
- 10) oxetane, tetrahydrofuran or tetrahydropyran optionally substituted with C1-C10 alkyl;

R_{43} represents a C1-C10 alkyl group;

R_{26} is aryl, or C1-C10 alkyl optionally substituted with a halogen atom(s);

R_{27} is selected from:

- 30 1) aryl optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- 2) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with aryl,
- 3) a hydroxyl group,
- 4) amino,
- 35 5) C1-C10 alkylamino,
- 6) hydroxycarbonyl,
- 7) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl and/or aryl, and
- 8) heteroaryloxy;

R_{28} is selected from hydrogen or C1-C10 alkyl optionally substituted with aryl;

R_{29} is selected from hydrogen or C1-C10 alkyl optionally substituted with aryl;

R_{28} and R_{29} may be bonded to form a ring selected from azetidiny, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom;

R_{30} is selected from a hydroxyl group, C1-C10 alkoxy and $-\text{NR}_{31}\text{R}_{32}$;

R_{31} and R_{32} are independently selected from:


- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with aryl and
- 50 3) aryl;

R_{31} and R_{32} may be bonded to form a ring selected from azetidiny, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl, a halogen atom and C1-C10 alkoxycarbonyl; and R_{33} and R_{34} are independently selected from:


- 55 1) hydrogen and
- 2) C1-C10 alkyl], or a pharmacologically acceptable salt thereof;

or


a compound represented by the following general formula (2):



[wherein W, Y, m, n, R₄, R₅, R₆, R₇, R₁₁, R₁₆, R₁₉, R₂₀, R₂₁, R₃₃, R₃₄ and R₄₄ are as defined for the formula (1) ;
U represents a bond, C1-C10 alkylene or any group selected from groups represented by the following formula:



A is selected from O, NH and CH₂;
R₄₆ is selected from hydrogen or R₄₄;
T is selected from aryl and heteroaryl;
V is selected from:



E is a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) selected from O, N, S, SO and SO₂, and the heterocycle is optionally substituted with a group(s) selected from:

- 1) hydrogen,
- 2) a halogen atom,
- 3) C1-C10 alkyl optionally having a group(s) independently selected from C1-C10 alkylamino, a halogen atom and a hydroxyl group,
- 4) a hydroxyl group,
- 5) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 6) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group,

amino and C1-C10 alkylamino,

7) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,

8) a heterocycle optionally substituted with C1-C10 alkyl,

9) heteroaryl optionally substituted with C1-C10 alkyl,

10) heterocyclyl C1-C10 alkyl,

11) $-\text{COR}_{16}$,

12) $-\text{NR}_{19}\text{R}_{20}$ and


13) $-\text{SO}_2\text{R}_{21}$;

Z is a divalent group selected from:

1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s) and/or a hydroxyl group(s), wherein the carbon atom(s) may be oxidized to form carbonyl;

2) C2-C10 alkenylene or C2-C10 heteroalkenylene optionally substituted with a halogen atom(s) and/or a hydroxyl group(s), wherein the carbon atom(s) may be oxidized to form carbonyl; and

3) a group selected from:



G is a divalent group selected from:

1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s); and

2) C2-C10 alkenylene or C2-C10 heteroalkenylene optionally substituted with a halogen atom(s);

J is a divalent group selected from:

1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s); and

2) C2-C10 alkenylene or C2-C10 heteroalkenylene optionally substituted with a halogen atom(s);

B is selected from a heterocycle or heteroaryl; and R_{45} is selected from hydrogen or C1-C10 alkyl], or a pharmacologically acceptable salt thereof.

[2] The compound or a pharmacologically acceptable salt thereof according to [1], wherein

W is selected from:

1) a single bond,

2) C1-C10 alkylene optionally containing a carbonyl group, wherein the alkylene is optionally substituted with a halogen atom(s) or hydroxy,

3) C2-C10 alkenylene optionally substituted with a halogen atom(s),

4) C2-C10 alkynylene,

5) arylene,

6) heteroarylene,

7) $-\text{NH}-$, $-\text{NHCH}_2-$ or $-\text{NHCH}_2\text{CH}_2-$,

8) cycloalkylene and

9) $-(\text{cycloalkylene})-\text{CH}_2-$;

X is selected from the following bond or groups:

1) a single bond,

- 2) C1-C10 alkylene optionally substituted with cycloalkyl,
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) C1-C10 oxyalkylene;

R₁ is selected from:

- 1) hydrogen,
- 2) cycloalkyl optionally substituted with a group selected from R₄,
- 3) a heterocycle optionally substituted with a group(s) selected from R₂₅ and R₄,
- 4) aryl optionally substituted with a group(s) selected from R₃, R₄, R₅ and R₆ and
- 5) heteroaryl optionally substituted with a group(s) selected from R₂₅, R₄ and R₅;

R₉ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from R₂₃,
- 3) cycloalkyl optionally substituted with a halogen atom(s) or a hydroxyl group(s),
- 4) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy, C1-C10 alkoxycarbonyl, amino and a halogen atom,
- 5) C1-C10 heteroalkyl optionally substituted with a group(s) selected from a halogen atom and a hydroxyl group,
- 6) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxycarbonyl and a halogen atom and
- 7) cycloalkenyl optionally substituted with a group(s) selected from C1-C10 alkoxy, C1-C10 alkylamino, amino, 1 to 3 hydroxyl groups and 1 to 4 halogen atoms, wherein the cycloalkenyl optionally contains a carbonyl group;

R₁₀ is selected from:

- 1) hydrogen and
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group and aryl;

R₉ and R₁₀ may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from N, O, S, SO, SO₂, carbonyl and thiocarbonyl, and the heterocycle is optionally substituted with a substituent(s) independently selected from R₂₄;

R₂₄ is selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from C1-C10 alkylamino and C1-C10 alkylcarbonylamino,
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 8) C1-C10 heteroalkyl optionally substituted with 1 to 2 groups selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 9) -COR₁₆ and
- 10) -NR₁₉R₂₀;

R₁₁ is selected from:

- 1) C1-C10 alkyl optionally substituted with 1 to 3 substituents independently selected from:

- i) a hydroxyl group,
- ii) $-NR_{17}R_{18}$,
- iii) a C1-C10 alkoxy group,
- iv) a halogen atom,
- v) C1-C10 alkoxycarbonyl and
- vi) aminocarbonyl,

- 2) aryl,
- 3) aryl C1-C10 alkyl,
- 4) cycloalkyl optionally substituted with a halogen atom(s),
- 5) a heterocycle optionally substituted with C1-C10 alkyl,
- 6) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino and a hydroxyl group,
- 7) amino,
- 8) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino, hydroxycarbonyl and a hydroxyl group and
- 9) C2-C10 alkenyl; and

R_{33} and R_{34} are hydrogen,
wherein R_3 , R_4 , R_5 , R_6 , R_{16} , R_{17} , R_{18} , R_{19} , R_{20} , R_{23} and R_{25} are as defined in [1], respectively.

[3] The compound or a pharmacologically acceptable salt thereof according to [1] or [2], wherein

W is selected from:

- 1) a single bond,
- 2) C1-C10 alkylene optionally substituted with a halogen atom(s),
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) heteroarylene,

X is selected from the following bond or groups:

- 1) a single bond,
- 2) C1-C10 alkylene,
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) C1-C10 oxyalkylene, wherein the oxyalkylene is bonded to a 1,3,8-triaza-spiro[4.5]dec-1-en-4-one ring or a 1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione ring through a carbon atom in the oxyalkylene;

R_1 is selected from:

- 1) aryl optionally substituted with a group(s) selected from R_3 , R_4 and R_5 and
- 2) heteroaryl optionally substituted with a group(s) selected from R_{25} and R_4 ;

R_3 is selected from:

- 1) $-CONR_7R_8$,
- 2) $-OR_9$,
- 3) $-NR_9R_{10}$,
- 4) $-N(R_9)COR_{11}$,
- 5) $-N(R_9)SO_2R_{12}$,
- 6) $-SO_2R_{15}$,
- 7) C1-C10 alkyl optionally substituted with a group(s) selected from $-COR_{16}$ and $-NR_{13}R_{14}$ and
- 8) $-N(R_9)CSNH_2$;

R_4 is selected from:

- 1) halogen,
- 2) cyano,
- 3) amino,
- 4) C1-C10 alkyl,
- 5) C1-C10 haloalkyl,
- 6) C1-C10 alkoxy,
- 7) C1-C10 haloalkylcarbonyl,
- 8) -COR₁₆ and
- 9) C1-C10 heteroalkyl;

R₂ is selected from:

- 1) C1-C10 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a group selected from R₄₂,
- 2) C2-C10 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a group selected from R₄₂,
- 3) C2-C10 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a group selected from R₄₂,
- 4) cycloalkyl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) C2-C10 alkenyl or C1-C10 alkyl,
- iii) aryl optionally substituted with a group(s) independently selected from C1-C10 alkyl, a halogen atom and C1-C10 alkoxy,
- iv) cycloalkyl,
- v) C2-C10 haloalkenyl or C1-C10 haloalkyl,
- vi) C1-C10 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with a halogen atom(s),
- vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),
- viii) C1-C10 alkyl substituted with C1-C10 alkoxy, wherein the alkyl and/or the alkyl in the alkoxy is optionally substituted with a halogen atom(s),
- ix) C2-C10 alkynyl and
- x) -Si(R₄₃)₃,

- 5) a heterocycle, wherein the heterocycle is optionally substituted with a group(s) selected from:

- i) a C1-C10 alkyl group,
- ii) C1-C10 alkylcarbonyl, wherein the alkyl group is optionally substituted with R₂₇,
- iii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- iv) heteroarylcarbonyl,
- v) C1-C10 alkoxycarbonyl, wherein the alkyl group is optionally substituted with a group(s) independently selected from a halogen atom, aryl and C1-C10 alkoxy,
- vi) aryloxycarbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) and/or C1-C10 alkyl,
- vii) -CONR₂₈R₂₉ and
- viii) -SO₂R₂₁,

- 6) aryl optionally substituted with a group(s) independently selected from R₄₄,

- 7) heteroaryl optionally substituted with any of the following groups:

- i) C1-C10 alkyl,

- 8) C1-C10 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a group selected from R₄₂,

- 9) -S(O)_qR₄₃ (wherein q is an integer of 0 to 2) and

- 10) cycloalkenyl optionally substituted with C1-C10 alkyl, R₄₄ is selected from:

- 1) a halogen atom,
- 2) cyano,
- 3) C1-C10 alkyl optionally substituted with any of the following groups:

- i) a hydroxyl group,
- ii) -OR₂₆,
- iii) cyano and
- iv) aryloxy optionally substituted with a group(s) selected from a halogen atom, C1-C10 alkyl, C1-C10 haloalkyl or C1-C10 haloalkoxy,

- 4) C1-C10 haloalkyl,
- 5) cycloalkyl optionally substituted with a group(s) selected from a halogen atom and C1-C10 haloalkyl,
- 6) C1-C10 alkoxy optionally substituted with a halogen atom(s) or a C2-C6 alkenyl group,
- 7) -COR₃₀,
- 8) C1-C10 heteroalkyl optionally substituted with a halogen atom(s),
- 9) aryl optionally substituted with a group(s) independently selected from:

- i) C1-C10 alkyl and
- ii) aryl,

- 10) heteroaryl optionally substituted with a C1-C10 alkyl group(s),
- 11) -SO₂R₄₃,
- 12) C1-C10 alkylthio optionally substituted with a halogen atom(s),
- 13) -Si(R₄₃)₃ and
- 14) -SF₅; and

R₂₇ is selected from:

- 1) aryl optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- 2) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with aryl,
- 3) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl and aryl and
- 4) heteroaryloxy,

wherein R₅, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₂₁, R₂₅, R₂₆, R₂₈, R₂₉, R₃₀, R₄₂ and R₄₃ are as defined in [1] or [2] from which [3] depends, respectively.

[4] The compound or a pharmacologically acceptable salt thereof according to any of [1] to [3], wherein

W is selected from:

- 1) C1-C6 alkylene optionally substituted with a fluorine atom(s),
- 2) C1-C6 alkenylene and
- 3) thiophene,

X is selected from the following bond or groups:

- 1) a single bond,
- 2) C1-C6 alkylene and
- 3) C1-C6 oxyalkylene optionally substituted with a halogen atom(s), wherein the oxyalkylene is bonded to a 1,3,8-triaza-spiro[4.5]dec-1-en-4-one ring or a 1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione ring through a carbon atom in the oxyalkylene;

Y represents an oxygen atom;
m represents 1; and
n represents 1.

[5] The compound or a pharmacologically acceptable salt thereof according to any of [1] to [4], wherein

W is selected from:

- 1) ethylene,
- 2) vinylene and
- 3) thiophene,

X represents a single bond;


R₃ is selected from:

- 1) -CONR₇R₈,
- 2) -OR₉,
- 3) -NR₉R₁₀,
- 4) -N(R₉)-COR₁₁,
- 5) -N(R₉)SO₂R₁₂,
- 6) -SO₂R₁₅ and
- 7) C1-C6 alkyl optionally substituted with a group(s) selected from -COR₁₆ and -NR₁₃R₁₄;

R₂ is selected from:

- 1) C1-C10 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a group selected from R₄₂,
- 2) C2-C10 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a group selected from R₄₂,
- 3) C2-C10 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a group selected from R₄₂,
- 4) cycloalkyl optionally substituted with a group(s) independently selected from:
 - i) a halogen atom,
 - ii) C2-C6 alkenyl or C1-C6 alkyl,
 - iii) aryl optionally substituted with a group(s) independently selected from C1-C6 alkyl, a halogen atom, C1-C6 alkoxy, C1-C6 alkylamino and C1-C6 alkylcarbonyl,
 - iv) cycloalkyl,
 - v) C2-C6 haloalkenyl or C1-C6 haloalkyl,
 - vi) C1-C6 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with a halogen atom(s),
 - vii) C1-C6 alkoxy optionally substituted with a halogen atom(s),
 - viii) C1-C6 alkyl substituted with C1-C6 alkoxy, wherein the alkyl and/or the alkyl in the alkoxy is optionally substituted with halogen,
 - ix) C2-C6 alkynyl and
 - x) -Si(R₄₃)₃,

5) a group represented by the following general formula (B):



(wherein Ra represents a group selected from:

- i) C1-C6 alkylcarbonyl, wherein the alkyl group is optionally substituted with R₂₇,
- ii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C6 alkyl and C1-C6 alkoxy,
- iii) C1-C6 alkoxycarbonyl, wherein the alkyl group is optionally substituted with a group(s) selected from a halogen atom, aryl and C1-C6 alkoxy,
- iv) aryloxy carbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) or C1-C6

alkyl,
v) $-\text{CONR}_{28}\text{R}_{29}$ and
vi) $-\text{SO}_2\text{R}_{21}$,

5 6) aryl optionally substituted with a group(s) independently selected from R_{44} ,
7) heteroaryl optionally substituted with any of the following groups:

10 i) a halogen atom,
ii) C1-C6 alkyl and
iii) C1-C6 alkoxy;

8) C1-C6 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a group selected from R_{42} ,

9) $-\text{S}(\text{O})_q\text{R}_{43}$ (wherein q is an integer of 0 to 2) and

15 10) cycloalkenyl optionally substituted with C1-C6 alkyl; and

R_{44} is selected from:

20 1) a halogen atom,
2) cyano,
3) C1-C6 alkyl optionally substituted with any of the following groups:

25 i) a hydroxyl group,
ii) $-\text{OR}_{26}$,
iii) cyano and
iv) aryloxy optionally substituted with a group(s) selected from a halogen atom, C1-C6 alkyl, C1-C6 haloalkyl or C1-C6 haloalkoxy,

30 4) C1-C6 haloalkyl,
5) cycloalkyl optionally substituted with a group(s) selected from a halogen atom and C1-C6 haloalkyl,
6) C1-C6 alkoxy optionally substituted with a halogen atom(s),
7) $-\text{COR}_{30}$,
8) C1-C6 heteroalkyl optionally substituted with a halogen atom(s),
9) aryl optionally substituted with a group(s) independently selected from:

35 i) C1-C6 alkyl and
ii) aryl,


40 10) heteroaryl optionally substituted with a C1-C6 alkyl group(s),
11) $-\text{SO}_2\text{R}_{43}$,
12) C1-C6 alkylthio optionally substituted with a halogen atom(s),
13) $-\text{Si}(\text{R}_{43})_3$ and
14) $-\text{SF}_5$,

45 wherein R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{21} , R_{26} , R_{27} , R_{28} , R_{29} , R_{30} , R_{42} and R_{43} are as defined in [1] to [4] from which [5] depends, respectively.

[6] The compound or a pharmacologically acceptable salt thereof according to [3], wherein R_1 is a group represented by the following general formula (3):


50

55




wherein R_3 , R_4 and R_5 are as defined for R_3 , R_4 and R_5 in [3].

[7] The compound or a pharmacologically acceptable salt thereof according to [3], wherein R_1 is a group represented by the following general formula (4):




wherein R_4 and R_{25} are as defined for R_4 and R_{25} in [3].

[8] The compound or a pharmacologically acceptable salt thereof according to [5], wherein R_1 is a group represented by the following general formula (3):



wherein R_3 , R_4 and R_5 are as defined for R_3 , R_4 and R_5 in [5].

[9] The compound or a pharmacologically acceptable salt thereof according to [5], wherein R_1 is a group represented by the following general formula (4):




(4)

wherein R_4 and R_{25} are as defined for R_4 and R_{25} in [5].

[10] Compounds of Compound Nos. (1) to (1446) described herein or pharmacologically acceptable salts thereof.

[11] The compound or a pharmacologically acceptable salt thereof according to any of [1] to [5], wherein U represents C1-C6 alkylene or any group selected from groups represented by the following formula:




;

A is O;

R_{46} is selected from hydrogen, C1-C10 alkyl, C1-C10 haloalkyl and C1-C10 hydroxyalkyl;

V is selected from:



;

E is pyrrolidine or piperidine optionally substituted with a hydroxyl group(s); and

R_7 is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl and
- 3) C1-C10 hydroxyalkyl.

[12] A pharmaceutical composition comprising the compound or a pharmacologically acceptable salt thereof according to any of [1] to [11] as an active ingredient.

[13] A prophylactic or therapeutic agent for use in the treatment or prevention of osteoporosis, fracture, osteomalacia, arthritis, thrombocytopenia, hypoparathyroidism, hyperphosphatemia or tumoral calcinosis, comprising the compound or a pharmacologically acceptable salt thereof according to any of [1] to [11] as an active ingredient.

[14] Use of the compound or a pharmacologically acceptable salt thereof according to any of [1] to [11] for the manufacture of a prophylactic or therapeutic agent for use in the treatment or prevention of osteoporosis, fracture, osteomalacia, arthritis, thrombocytopenia, hypoparathyroidism, hyperphosphatemia or tumoral calcinosis.

[15] The compound or a pharmacologically acceptable salt thereof according to any of [1] to [11] for use in the treatment or prevention of osteoporosis, fracture, osteomalacia, arthritis, thrombocytopenia, hypoparathyroidism, hyperphosphatemia or tumoral calcinosis.

[0017] In the description of each claim, a substituent not particularly defined is as defined for the same substituent in another claim from which the claim depends.

[0018] In the present specification and claims translated into languages such as English, description with indefinite articles (e.g., "a", "an" in English), definite articles (e.g., "the" in English) and the like includes singular and plural aspects unless otherwise defined. For example, "a group" in English includes one or more groups.

[Advantageous Effects of Invention]

[0019] The compounds or pharmacologically acceptable salts thereof according to the present invention have a parathyroid hormone-like effect involving bone anabolism which can considerably improve the compliance of patients as compared with parathyroid hormone peptide agonists. [Description of Embodiments]

[0020] The present invention relates to spiroimidazolone derivatives and use thereof. The present inventors have synthesized a compound represented by the above formula (1) or (2) or a pharmacologically acceptable salt thereof for the first time and have found that the compound or a salt thereof is a compound having a parathyroid hormone (PTH)-like effect.

[0021] The "alkyl" herein refers to a monovalent group derived by removing any one hydrogen atom from an aliphatic hydrocarbon, and covers a subset of hydrocarbyl or hydrocarbon group structures not containing a heteroatom or an unsaturated carbon-carbon bond and containing hydrogen and carbon atoms in the backbone. Examples of the alkyl group include those of linear or branched structures. The alkyl group is preferably an alkyl group having 1 to 10 carbon atoms (C₁-C₁₀; "C_p-C_q" hereinafter means that the group has p to q carbon atoms), more preferably a C₁-C₆ alkyl group. In particular, it is preferably a C₁-C₃ alkyl group in R₃₈ and R₃₉, a C₁-C₃ alkyl group in R₃, a C₁-C₃ alkyl group in R₄, a C₁-C₃ alkyl group in R₅, a C₁-C₃ alkyl group in R₆, a C₁-C₃ alkyl group in R₇, a C₁-C₃ alkyl group in R₄₀ and R₄₁, a C₁-C₃ alkyl group in R₈, a C₁-C₃ alkyl group in a substituent on a heterocycle where R₇ and R₈ are bonded to each other to form the heterocycle or a substituent on a spiro ring where the spiro ring is formed with the heterocycle, a C₁-C₃ alkyl group in R₁₆, a C₁-C₅ alkyl group in R₁₇, a C₁-C₃ alkyl group in R₁₈, a C₁-C₃ alkyl group in a substituent on a heterocycle where R₁₇ and R₁₈ are bonded to each other to form the heterocycle, a C₁-C₃ alkyl group in R₁₉, a C₁-C₃ alkyl group in R₂₀, a C₁-C₃ alkyl group in R₂₁, a C₁-C₄ alkyl group in R₉, a C₁-C₃ alkyl group in R₂₃, a C₁-C₃ alkyl group in R₁₀, a C₁-C₃ alkyl group in R₂₄, a C₁-C₄ alkyl group in R₁₁, a C₁-C₃ alkyl group in R₁₂, a C₁-C₃ alkyl group in R₁₃, a C₁-C₄ alkyl group in R₁₄, a C₁-C₃ alkyl group in a substituent on a heterocycle where R₁₃ and R₁₄ are bonded to each other to form the heterocycle, a C₁-C₃ alkyl group in R₁₅, a C₁-C₄ alkyl group in R₃₅, a C₁-C₃ alkyl group in R₃₆, a C₁-C₄ alkyl group in R₂₅, a C₁-C₁₃ alkyl group in R₂, a C₁-C₅ alkyl group in R₄₄, a C₁-C₃ alkyl group in R₄₂, a C₁-C₃ alkyl group in R₄₃, a C₁-C₃ alkyl group in R₂₆, a C₁-C₃ alkyl group in R₂₇ and a C₁-C₆ alkyl group in R₂₈.

[0022] Specific examples of the alkyl include a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, an s-butyl group, a t-butyl group, a pentyl group, an isopentyl group, a 2,3-dimethylpropyl group, a 3,3-dimethylbutyl group, a hexyl group, a 2,3-dimethylhexyl group, a 1,1-dimethylpentyl group, a heptyl group and an octyl group.

[0023] The "alkenyl" herein refers to a monovalent group having at least one double bond (two adjacent SP² carbon atoms). Depending on the configuration of the double bond and the substituent (if present), the geometry of the double bond can be an entgegen (E) or zusammen (Z) configuration or a cis or trans configuration. Examples of the alkenyl group include linear or branched groups, including straight chains that include internal olefins. Preferred examples include C₂-C₁₀ alkenyl groups, and more preferred examples include C₂-C₆ alkenyl groups. In particular, it is preferably a C₂-C₅ alkenyl group in R₇ and a C₁-C₉ alkenyl group in R₂.

[0024] Specific examples of such alkenyl include a vinyl group, an allyl group, a 1-propenyl group, a 2-propenyl group, a 1-butenyl group, a 2-butenyl group (including cis and trans), a 3-butenyl group, a pentenyl group and a hexenyl group.

[0025] The "alkynyl" herein refers to a monovalent group having at least one triple bond (two adjacent SP carbon atoms). Examples include linear or branched alkynyl groups, including internal alkynes. Preferred examples include C₂-C₁₀ alkynyl groups, and more preferred examples include C₂-C₆ alkynyl groups. In particular, it is preferably a C₂-C₉ alkynyl group in R₂.

[0026] Specific examples of the alkynyl include an ethynyl group, a 1-propynyl group, a propargyl group, a 3-butylnyl group, a pentynyl group, a hexynyl group, a 3-phenyl-2-propynyl group, a 3-(2'-fluorophenyl)-2-propynyl group, a 2-hydroxy-2-propynyl group, a 3-(3-fluorophenyl)-2-propynyl group and a 3-methyl-(5-phenyl)-4-pentynyl group.

[0027] The alkenyl or alkynyl can have one or more double bonds or triple bonds, respectively.

[0028] The "cycloalkyl" herein refers to a saturated cyclic monovalent aliphatic hydrocarbon group and includes single rings, fused rings, bicyclo rings and spiro rings. Preferred examples include C₃-C₁₀ cycloalkyl groups. Specific examples

of the cycloalkyl group include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group and a bicyclo[2.2.1]heptyl group.

[0029] The "cycloalkenyl" herein refers to a cyclic aliphatic hydrocarbon group having at least one double bond and includes single rings, fused rings, bicyclo rings and spiro rings. Preferred examples include C3-C10 cycloalkynyl groups, and more preferred examples include C3-C6 alkenyl groups. It is preferably C3-C5 cycloalkenyl in R_g. Specific examples of the cycloalkenyl group include a cyclopropenyl group, a cyclobutenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a cyclooctenyl group and a tetralinyl group.

[0030] The "heteroatom" herein refers to a nitrogen atom (N), an oxygen atom (O) or a sulfur atom (S).

[0031] The "halogen atom" herein refers to a fluorine atom, a chlorine atom, a bromine atom or an iodine atom.

[0032] The "haloalkyl" herein represents a group in which preferably 1 to 9, more preferably 1 to 5, of the same or different above "halogen atoms" are bonded to the above "alkyl". The haloalkyl is preferably C1-C10 haloalkyl, more preferably C1-C6 haloalkyl. In particular, it is preferably C1-C3 haloalkyl in R₄ and C1-C3 haloalkyl in R₄₄.

[0033] Specific examples include a fluoromethyl group, a difluoromethyl group and a trifluoromethyl group.

[0034] The "haloalkenyl" herein represents a group in which preferably 1 to 9, more preferably 1 to 5, of the same or different above "halogen atoms" are bonded to the above "alkenyl".

[0035] The "alkylcarbonyl" herein refers to a carbonyl group to which the above-defined "alkyl" is bonded, and is preferably C1-C10 alkylcarbonyl, more preferably C1-C6 alkylcarbonyl. In particular, it is preferably C1-C3 alkylcarbonyl in R₄₀ and R₄₁, C1-C3 alkylcarbonyl in R₁₉, C1-C3 alkylcarbonyl in R₉, C1-C3 alkylcarbonyl in R₂₃, C1-C3 alkylcarbonyl in R₁₂, C1-C3 alkylcarbonyl in R₃₅ and C1-C5 alkylcarbonyl in R₂.

[0036] The "Cn-Cm alkylcarbonyl" herein means that the alkyl therein is a "Cn-Cm" alkyl in terms of the number of carbon atoms. Hereinafter, the same applies to a group containing "alkylcarbonyl".

[0037] Specific examples include an acetyl group, an ethylcarbonyl group, a 1-propylcarbonyl group, a 2-propylcarbonyl group and a 2,2-dimethylpropylcarbonyl group.

[0038] The "haloalkylcarbonyl" herein refers to a carbonyl group to which the above-defined "haloalkyl" is bonded. The haloalkylcarbonyl is preferably C1-C10 haloalkylcarbonyl, more preferably C1-C6 haloalkylcarbonyl. In particular, it is preferably C1-C3 haloalkylcarbonyl in R₄.

[0039] The "cycloalkylcarbonyl" herein refers to a carbonyl group to which the above-defined "cycloalkyl" is bonded.

[0040] The "alkenylcarbonyl" herein refers to a carbonyl group to which the above-defined "alkenyl" is bonded, and is preferably C2-C10 alkenyl, more preferably C2-C6 alkenylcarbonyl. In particular, it is preferably C2-C3 alkenylcarbonyl in R₂.

[0041] The "Cn-Cm alkenylcarbonyl" herein means that it includes an alkenyl having "Cn-Cm" carbon atoms. Hereinafter, the same applies to a group containing "alkenylcarbonyl".

[0042] The "alkoxy" herein refers to an oxy group to which the above-defined "alkyl" is bonded, and is preferably a C1-C10 alkoxy group, more preferably a C1-C6 alkoxy group. In particular, it is preferably a C1-C3 alkoxy group in R₃₇, a C1-C3 alkoxy group in R₃, a C1-C3 alkoxy group in R₄, a C1-C3 alkoxy group in R₅, a C1-C3 alkoxy group in R₇, a C1-C3 alkoxy group in a substituent on a heterocycle where R₇ and R₈ are bonded to each other to form the heterocycle, a C1-C4 alkoxy group in R₁₆, a C1-C3 alkoxy group in R₁₇, a C1-C3 alkoxy group in R₁₁, a C1-C4 alkoxy group in R₂ and a C1-C4 alkoxy group in R₂₇. Specific examples include a methoxy group, an ethoxy group, a 1-propoxy group, a 2-propoxy group, an n-butoxy group, an i-butoxy group, a sec-butoxy group, a t-butoxy group, a 1-pentyloxy group, a 2-pentyloxy group, a 3-pentyloxy group, a 2-methyl-1-butyloxy group, a 3-methyl-1-butyloxy group, a 2-methyl-2-butyloxy group, a 3-methyl-2-butyloxy group, a 2,2-dimethyl-1-propyloxy group, a 1-hexyloxy group, a 2-hexyloxy group, a 3-hexyloxy group, a 2-methyl-1-pentyloxy group, a 3-methyl-1-pentyloxy group, a 4-methyl-1-pentyloxy group, a 2-methyl-2-pentyloxy group, a 3-methyl-2-pentyloxy group, a 4-methyl-2-pentyloxy group, a 2-methyl-3-pentyloxy group, a 3-methyl-3-pentyloxy group, a 2,3-dimethyl-1-butyloxy group, a 3,3-dimethyl-1-butyloxy group, a 2,2-dimethyl-1-butyloxy group, a 2-ethyl-1-butyloxy group, a 3,3-dimethyl-2-butyloxy group, a 2,3-dimethyl-2-butyloxy group and a 1-methyl-cyclopropylmethoxy group.

[0043] The "alkylcarbonyloxy" herein refers to an oxy group to which the above-defined "alkylcarbonyl" is bonded, and is preferably a C1-C10 alkylcarbonyloxy group, more preferably a C1-C6 alkylcarbonyloxy group. In particular, it is preferably a C1-C3 alkylcarbonyloxy group in R₂₃.

[0044] The "alkoxycarbonyl" herein refers to a carbonyl group to which the above-defined "alkoxy" is bonded. The alkoxycarbonyl is preferably C1-C10 alkoxycarbonyl, more preferably C1-C6 alkoxycarbonyl. It is preferably C1-C3 alkoxycarbonyl in R₄, C1-C3 alkoxycarbonyl in R₁₆, C1-C4 alkoxycarbonyl in R₁₉, C1-C4 alkoxycarbonyl in R₉, C1-C3 alkoxycarbonyl in R₂₃, C1-C3 alkoxycarbonyl in R₁₁, C1-C4 alkoxycarbonyl in R₁₃, C1-C5 alkoxycarbonyl in R₂ and C1-C3 alkoxycarbonyl in R₄₂. Examples include -CO₂tBu (t-butoxycarbonyl) and -CO₂Me (methoxycarbonyl).

[0045] The "Cn-Cm alkoxycarbonyl" herein means that the alkyl in the alkoxy is a "Cn-Cm" alkyl in terms of the number of carbon atoms. Hereinafter, the same applies to a group containing "alkoxycarbonyl".

[0046] The "heteroalkyl" herein refers to a group containing 1 to 5 heteroatoms in the above-defined "alkyl" backbone and is preferably C1-C10 heteroalkyl, more preferably C1-C6 heteroalkyl. In particular, it is preferably C1-C5 heteroalkyl

in R₉, C1-C5 heteroalkyl in R₁₁ and C1-C6 heteroalkyl in R₂₅. Examples include -CH₂OCH₃, -CH₂OCH₂CH₃, -CH(Me)OCH₃ and -CH₂CH₂NMe₂.

[0047] The "heteroalkenyl" herein refers to a group containing preferably 1 to 5 heteroatoms in the above-defined "alkenyl" backbone.

[0048] The "alkylene" herein refers to a divalent group having a basic skeleton represented by -(CH₂)_n- (preferably n = 1 to 10), and may contain a branched chain. Specific examples include C1-C5 alkylene (n = 1 to 5). More specific examples include a methylene group, a dimethylmethylene group, an ethylene group, a propylene group, a butylene group and a pentamethylene group. In particular, it is preferably C2-C5 alkylene in W, C1-C9 alkylene in X, C1-C10 alkylene in U, C1-C10 alkylene in Z and C1-C5 alkylene in G.

[0049] The "alkylidene" herein refers to a divalent group produced by removing two hydrogen atoms from the same carbon atom of a ring, the free valencies of which are part of a double bond. The geometry of the double bond can be an entgegen (E) or zusammen (Z) configuration or a cis or trans configuration. Examples of the alkylidene include linear or branched groups. Preferred examples include C1-C10 alkylidene, and more preferred examples include C1-C6 alkylidene. In particular, it is preferably C1-C4 alkylidene in R₂. Specific examples include methylene (=CH₂), ethylidene (=CHCH₃), isopropylidene (=C(CH₃)₂) and propylidene (=CHCH₂CH₃)

[0050] The "alkenylene" herein refers to a divalent group having at least one double bond (two adjacent SP² carbon atoms). Depending on the configuration of the double bond and the substituent (if present), the geometry of the double bond can be an entgegen (E) or zusammen (Z) configuration or a cis or trans configuration. Examples of the alkenylene include linear or branched groups. Preferred examples include C2-C10 alkenylene, and more preferred examples include C2-C6 alkenylene. Specific examples include a vinylene group, a 1-propenylene group, a 1-butenylene group and a 1-pentenylene group. In particular, it is preferably C2-C5 alkenylene in W, C2-C9 alkenylene in X, C2-C10 alkenylene in Z and C2-C5 alkenylene in G.

[0051] The "alkynylene" herein refers to a divalent group having at least one triple bond (two adjacent SP carbon atoms). Examples include linear or branched alkynylenes. Preferred examples include C2-C10 alkynylene, and more preferred examples include C2-C6 alkynylene. In particular, it is preferably C2-C5 alkynylene in W and C2-C9 alkynylene in X.

[0052] The "cycloalkylene" herein refers to a saturated cyclic divalent aliphatic hydrocarbon group and includes single rings, bicyclo rings and spiro rings. Preferred examples include C3-C10 cycloalkylene. Specific examples of the cycloalkyl group include a cyclopropylene group, a cyclobutylene group, a cyclopentylene group, a cyclohexylene group, a cycloheptylene group, a cyclooctylene group and a bicyclo[2.2.1]heptylene group.

[0053] The "oxyalkylene" herein refers to a divalent C1-C10 group in which one end of the above-defined alkylene is an oxygen atom. Examples include -CH₂O-, -C(Me)₂O-, -CH₂CH₂O-, -CH₂CH(Me)O- and -CH₂C(Me)₂O-. In X, the oxyalkylene is preferably bonded to a 1,3,8-triaza-spiro[4.5]dec-1-en-4-one ring or a 1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione ring through a carbon atom in the oxyalkylene. In particular, it is preferably C1-C5 oxyalkylene in X.

[0054] The "heteroalkylene" herein refers to a divalent, preferably C1-C10, group containing 1 to 5 heteroatoms in the above-defined "alkylene" backbone, and may contain a branched chain. Examples include -CH₂OCH₂-, -CH₂OCH₂CH₂-, -CH(Me)OCH₂-, -CH₂CH₂NHCH₂- and -CH₂CH₂N(Me)CH₂-. In particular, it is preferably C2-C5 heteroalkylene in W, C1-C8 heteroalkylene in Z and C1-C4 heteroalkylene in G.

[0055] The "heteroalkenylene" herein refers to a divalent, preferably C1-C10, group containing 1 to 5 heteroatoms in the above-defined "alkenylene" backbone, and may contain a branched chain. In particular, it is preferably C2-C8 heteroalkenylene in Z.

[0056] The "aryl" herein refers to a monovalent aromatic hydrocarbon ring, and may be partially saturated insofar as it is aromatic. Preferred examples include C6-C10 aryl. Specific examples of the aryl include a phenyl group, a naphthyl group (e.g., a 1-naphthyl group, a 2-naphthyl group) and a tetrahydronaphthyl group.

[0057] The "heteroaryl" herein refers to a monovalent group of an aromatic ring containing preferably 1 to 5 heteroatoms in the ring-forming atoms, and may be partially saturated. The saturated carbon atom(s) may be oxidized to form carbonyl. The ring may be a single ring or two fused rings (e.g., a bicyclic heteroaryl obtained by fusion with a benzene ring or monocyclic heteroaryl ring). The number of the ring-forming carbon atoms is preferably 1 to 10 (C1-C10 heteroaryl).

[0058] Specific examples of the heteroaryl include a furyl group, a thienyl group, a pyrrolyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, an oxadiazolyl group, a thiadiazolyl group, a triazolyl group, a tetrazolyl group, a pyridyl group, a pyrimidyl group, a pyridazinyl group, a pyrazinyl group, a triazinyl group, a benzofuranyl group, a benzothienyl group, a benzothiadiazolyl group, a benzothiazolyl group, a benzoxazolyl group, a benzoxadiazolyl group, a benzimidazolyl group, an indolyl group, an isoindolyl group, an indazolyl group, a quinolyl group, an isoquinolyl group, a cinnolyl group, a quinazolinyl group, a quinoxalinyl group, a benzodioxolyl group, an indolizinyl group and an imidazopyridyl group.

[0059] The "arylene" herein refers to a divalent group derived by further removing any one hydrogen atom from the above-defined "aryl". Preferred examples include C6-C10 arylene. Specific examples include a 1,3-phenylene group and a 1,4-phenylene group.

[0060] The "heteroarylene" herein refers to a divalent group derived by further removing any one hydrogen atom from the above-defined "heteroaryl". Specific examples include a 2,5-thiophenediyl group and a 2,6-pyridinediyl group.

[0061] The "heterocycle" herein refers to a C1-10 nonaromatic cycloalkyl, wherein the cycloalkyl is a monovalent group containing preferably 1 to 5 heteroatoms in the ring-forming atoms, the cycloalkyl may have a double bond in the ring, the carbon atom(s) may be oxidized to form carbonyl, the heteroatoms may form an oxo group, and the cycloalkyl may contain two fused rings. Specific examples of the heterocycle include azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, oxazolidone, a 1,4-benzodioxanyl group, a tetrahydropyranyl group, a 1,3-dioxolanyl group, a 1,3-thiazolidinyl group, a hydantoyl group, a benzoxazolinonyl group, a benzothiazolinonyl group, a 2,4-(1H,3H)quinazolinonyl group, an indolinyl group, an oxindolyl group, a 1,3-benzoxolyl group, an imidazolidinyl group, a pyrazolidinyl group, an oxazolidinyl group, an isoxazolidinyl group, a thiomorpholinyl group, a dihydrothiazolyl group, an oxetanyl group, a 2-oxa-6-aza-spiro[3.3]heptanyl group, a 1,2,3,4-tetrahydroquinolyl group, an imidazolidonyl group, a pyrazolidonyl group, an oxazolidonyl group, a succinimidyl group, a 2-azetidiny group, a 2-oxopiperazinyl group, a 3,5-dioxomorpholinyl group, a 2-oxomorpholinyl group, a 2,5-dehydrouraciny group, a 2-pyrrolidinonyl group, a 2-piperidinonyl group, a 4-piperidinonyl group, a 3-isoxazolidone group, a 1,1,3-trioxo[1,2,5]thiadiazolidinone group, a 1,1-dioxo-1 λ ⁶-thiomorphonyl group and an imidazolidine-2,4-dione group. In these groups, the carbon atom(s) may be oxidized to form carbonyl, and the heteroatoms may have an oxo group.

[0062] The "heterocyclic carbonyl" herein refers to a carbonyl group to which the above-defined "heterocycle" is bonded.

[0063] The "alkylamino" herein refers to an amino group to which one or two of the above-defined "alkyl" groups are bonded. Preferred examples include C1-C10 monoalkylamino and C1-C10 dialkylamino, and more preferred examples include C1-C6 monoalkylamino and C1-C6 dialkylamino. Two alkyl groups in the dialkylamino may be the same or different. In particular, it is preferably C1-C3 monoalkylamino or C1-C3 dialkylamino in R₇, C1-C3 monoalkylamino or dialkylamino in a substituent on a heterocycle where R₇ and R₈ are bonded to each other to form the heterocycle, C1-C3 monoalkylamino or C1-C3 dialkylamino in R₁₆, C1-C3 monoalkylamino or C1-C3 dialkylamino in R₁₇, C1-C3 monoalkylamino or C1-C3 dialkylamino in R₂₁, C1-C3 monoalkylamino or C1-C3 dialkylamino in R₂₃, C1-C3 monoalkylamino or C1-C3 dialkylamino in R₂₄, C1-C3 monoalkylamino or C1-C3 dialkylamino in R₁₁, C1-C3 monoalkylamino or C1-C3 dialkylamino in R₁₂, C1-C3 monoalkylamino or C1-C3 dialkylamino in R₁₄ and C1-C3 monoalkylamino or C1-C3 dialkylamino in R₃₅. The "alkyl" in the alkylamino may have the above-defined "aryl" as a substituent(s). Examples of the alkylamino include -NHCH₃, -N(CH₃)₂, -N(CH₃)CH₂CH₃ and -NHCH₂Ph.

[0064] The "amino" herein refers to a monovalent group having two hydrogen atoms on a nitrogen atom (a group represented by -NH₂).

[0065] The "arylalkyl" herein refers to a group in which any hydrogen atom in the above-defined "alkyl" is replaced by the above-defined "aryl". Preferred examples of the arylalkyl include C6-C10 aryl C1-C10 alkyl. In particular, it is preferably C6-C10 aryl C1-C3 alkyl in R₂₁, C6-C10 aryl C1-C3 alkyl in R₃₅ and C6-C10 aryl C1-C3 alkyl in R₂₅. Specific examples include a benzyl group, a phenethyl group and a 3-phenyl-1-propyl group.

[0066] The "heterocyclic alkyl" herein refers to a group in which any hydrogen atom in the above-defined "alkyl" is replaced by the above-defined "heterocycle". Specific examples include a morpholin-4-yl-methyl group, a 2-(morpholin-4-yl)ethyl group, a 4-hydroxy-piperidin-1-yl-methyl group, a 2-(4-hydroxy-piperidin-1-yl)ethyl group, a 4-methyl-piperazin-1-yl-methyl group and a 2-(4-methyl-piperazin-1-yl)ethyl group.

[0067] The "hydroxyalkyl" herein refers to a group in which any hydrogen atom(s) in the above-defined "alkyl" is replaced by preferably 1 to 4 hydroxyl groups, and it is preferably a C1-C10 hydroxyalkyl group, more preferably a C1-C6 hydroxyalkyl group. In particular, it is preferably a C1-C4 hydroxyalkyl group in R₂₅. Specific examples include hydroxymethyl, 2-hydroxyethyl, 3-hydroxypropyl and 2,3-dihydroxypropyl.

[0068] The "alkylcarbonylamino" herein refers to an amino group to which one or two of the above-defined "alkylcarbonyl" groups are bonded. Preferred examples include C1-C10 monoalkylcarbonylamino and C1-C10 dialkylcarbonylamino, and more preferred examples include C1-C6 monoalkylcarbonylamino and C1-C6 dialkylcarbonylamino. Two alkyl groups in the dialkylcarbonylamino may be the same or different. In particular, it is preferably C1-C3 alkylcarbonylamino in R₁₇, C1-C3 alkylcarbonylamino in R₂₄ and C1-C3 alkylcarbonylamino in R₃₅. Examples include CH₃CONH-.

[0069] The "alkoxycarbonylamino" herein refers to an amino group to which one or two of the above-defined "alkoxycarbonyl" groups are bonded. Preferred examples include C1-C10 monoalkoxycarbonylamino and C1-C10 dialkoxycarbonylamino. Two alkoxy groups in the dialkoxycarbonylamino may be the same or different. In particular, it is preferably C1-C4 monoalkoxycarbonyl or C1-C4 dialkoxycarbonylamino in R₄₂.

[0070] The "alkylaminocarbonyl" herein refers to a carbonyl group to which the above-defined "alkylamino" is bonded, and is preferably C1-C10 alkylaminocarbonyl, more preferably C1-C6 alkylaminocarbonyl. In particular, it is preferably C1-C3 alkylaminocarbonyl in R₁₃. Examples include CH₃NHCO-.

[0071] The "C_n-C_m alkylaminocarbonyl" herein means that the alkyl therein is a "C_n-C_m" alkyl in terms of the number of carbon atoms. Hereinafter, the same applies to a group containing "alkylaminocarbonyl".

[0072] The "arylcarbonyl" herein refers to a carbonyl group to which the above-defined "aryl" is bonded.

[0073] The "aryloxy" herein refers to an oxy group to which the above-defined "aryl" is bonded.

[0074] The "aryloxy" herein refers to a carbonyl group to which the above-defined "aryloxy" is bonded.

[0075] The "heteroaryloxy" herein refers to an oxy group to which the above-defined "heteroaryloxy" is bonded.

[0076] The "heteroaryloxy" herein refers to a carbonyl group to which the above-defined "heteroaryloxy" is bonded.

[0077] The "hydroxycarbonyl" herein refers to $-\text{CO}_2\text{H}$ (carboxyl).

[0078] The "aminocarbonyl" herein refers to a carbonyl group to which the above-defined "amino" is bonded.

[0079] The "hydroxyalkylamino" herein refers to an amino group to which one or two of the above-defined "hydroxyalkyl" groups are bonded. Examples include mono(hydroxyalkyl)amino and di(hydroxyalkyl)amino. Two hydroxyalkyl groups in the di(hydroxyalkyl)amino may be the same or different. $-\text{NHCH}_2-$ or $-\text{NHCH}_2\text{CH}_2-$ in W herein is preferably bonded through the nitrogen atom to the sulfonyl group in the formula (1).

[0080] The "hydroxyalkylaminoalkyl" herein refers to a group in which any hydrogen atom in the above-defined "alkyl" is replaced by the above-defined "hydroxyalkylamino".

[0081] The "alkoxyalkyl" herein refers to a group in which any hydrogen atom in the above-defined "alkyl" is replaced by the above-defined "alkoxy", and is preferably C1-C10 alkoxy-C1-C10 alkyl, more preferably C1-C6 alkoxy-C1-C6 alkyl. In particular, it is preferably C1-C3 alkoxy-C1-C3 alkyl in a substituent on a heterocycle where R_7 and R_8 are bonded to each other to form the heterocycle.

[0082] The "hydroxyalkyloxy" herein refers to a group in which any hydrogen atom in the above-defined "alkoxy" is replaced by a hydroxyl group, and is preferably C1-C10 hydroxyalkyloxy, more preferably C1-C6 hydroxyalkyloxy.

[0083] The "thiocarbonyl" herein refers to a group represented by $\text{C}=\text{S}$.

[0084] The "alkylthio" herein refers to a thio group to which the above-defined "alkyl" is bonded, and is preferably a C1-C10 alkylthio group, more preferably a C1-C6 alkylthio group.

[0085] The "B optionally substituted with A" herein denotes that any hydrogen atom(s) in B may be replaced with any number of As.

[0086] In the present invention, the number of substituents is not limited unless otherwise indicated. For example, the number of substituents may be 1 to 7, 1 to 4, 1 to 3, 1 to 2, or 1.

[0087] The "PTH-like effect" herein refers to activity of increasing intracellular CAMP (CAMP: cyclic adenosine monophosphate) by action on the PTH receptor or action on the signal transduction pathway through the PTH receptor.

[0088] Herein, "*" in a chemical formula denotes a bonding position.

[0089] The compounds according to the present invention, whether free forms or pharmacologically acceptable salts, are included in the present invention. Examples of such "salts" include inorganic acid salts, organic acid salts, inorganic base salts, organic base salts and acidic or basic amino acid salts.

[0090] Preferred examples of the inorganic acid salts include hydrochlorides, hydrobromides, sulfates, nitrates and phosphates. Preferred examples of the organic acid salts include acetates, succinates, fumarates, maleates, tartrates, citrates, lactates, stearates, benzoates, methanesulfonates and p-toluenesulfonates.

[0091] Preferred examples of the inorganic base salts include alkali metal salts such as sodium salts and potassium salts, alkaline earth metal salts such as calcium salts and magnesium salts, aluminum salts and ammonium salts. Preferred examples of the organic base salts include diethylamine salts, diethanolamine salts, meglumine salts and N,N-dibenzylethylenediamine salts.

[0092] Preferred examples of the acidic amino acid salts include aspartates and glutamates. Preferred examples of the basic amino acid salts include arginine salts, lysine salts and ornithine salts.

[0093] The compounds of the present invention may absorb moisture, have adsorbed water or form hydrates when left in the air. Such hydrates are also included in the salts of the present invention.

[0094] Further, the compounds I of the present invention may absorb certain other solvents to form solvates. Such salts are also encompassed in the present invention as salts of the compounds of the formula (1) or (2).

[0095] Herein, a structural formula of a compound may represent a certain isomer for the sake of convenience. However, the compounds of the present invention include all isomers such as geometric isomers, optical isomers based on asymmetric carbons, stereoisomers and tautomers as well as mixtures of these isomers which occur due to the structures of the compounds, without being limited to the formulas described for the sake of convenience, and may be either one of isomers or a mixture thereof. Thus, the compounds of the present invention may have an asymmetric carbon atom in the molecule and may be present as optically active forms and racemates, but the present invention is not limited to either of them and includes both of them.

[0096] The present invention includes all isotopes of the compounds represented by the formula (1) or (2). In the isotopes of the compounds of the present invention, at least one atom is replaced by an atom having the same atomic number (proton number) but having a different mass number (sum of the number of protons and the number of neutrons). Examples of the isotopes contained in the compounds of the present invention include a hydrogen atom, a carbon atom, a nitrogen atom, an oxygen atom, a phosphorus atom, a sulfur atom, a fluorine atom and a chlorine atom, including ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl , respectively. In particular, radioisotopes that decay by emitting radioactivity such as ^3H and ^{14}C are useful in body tissue distribution tests for pharmaceuticals or compounds. Stable isotopes do not decay, are almost equal in abundance and do not emit radioactivity, and thus they can be used

safely. The isotopes of the compounds of the present invention can be converted according to conventional methods by substituting a reagent containing a corresponding isotope for a reagent used for synthesis.

[0097] The compounds according to the present invention may exhibit crystalline polymorphism, but are not particularly limited to any one of these, but may be in any one of these crystal forms or exist as a mixture of two or more crystal forms.

[0098] The compounds according to the present invention include prodrugs thereof. The prodrugs are derivatives of the compounds of the present invention which have chemically or metabolically decomposable groups and are converted back to the original compounds after administration in vivo to exhibit their original efficacy, including complexes not formed with covalent bonds, and salts.

[0099] The compounds represented by the above formula (1) or (2) according to the present invention are preferably as follows.

[0100] W is preferably selected from:

- 1) a single bond,
- 2) C1-C10 alkylene optionally containing a carbonyl group, wherein the alkylene is optionally substituted with a halogen atom(s) and/or a hydroxyl group(s),
- 3) C2-C10 alkenylene optionally substituted with a halogen atom(s),
- 4) C2-C10 alkynylene,
- 5) arylene optionally substituted with a halogen atom(s),
- 6) heteroarylene optionally substituted with a halogen atom(s),
- 7) C1-C10 heteroalkylene optionally substituted with a halogen atom(s),
- 8) -NH-, -NHCH₂- or -NHCH₂CH₂-,
- 9) cycloalkylene and
- 10) -(cycloalkylene)-CH₂-.

[0101] More preferably, the above W is selected from:

- 1) a single bond,
- 2) C1-C10 alkylene optionally containing a carbonyl group, wherein the alkylene is optionally substituted with a halogen atom(s) or hydroxy,
- 3) C2-C10 alkenylene optionally substituted with a halogen atom(s),
- 4) C2-C10 alkynylene,
- 5) arylene,
- 6) heteroarylene,
- 7) -NH-, -NHCH₂- or -NHCH₂CH₂-,
- 8) cycloalkylene and
- 9) -(cycloalkylene)-CH₂-.

[0102] Still more preferably, the above W is selected from:

- 1) a single bond,
- 2) C1-C10 alkylene optionally substituted with a halogen atom(s),
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) heteroarylene.

[0103] Particularly preferably, the above W is selected from:

- 1) C1-C6 alkylene optionally substituted with a fluorine atom(s),
- 2) C1-C6 alkenylene and
- 3) thiophene.

[0104] More particularly preferably, the above W is selected from:

- 1) ethylene,
- 2) vinylene and
- 3) thiophene.

[0105] The above X is preferably selected from the following bond or groups:

- 1) a single bond,
 - 2) C1-C10 alkylene optionally substituted with a halogen atom(s) or cycloalkyl,
 - 3) C2-C10 alkenylene optionally substituted with a halogen atom(s),
 - 4) C2-C10 alkynylene optionally substituted with a halogen atom(s),
 - 5) C1-10 oxyalkylene optionally substituted with a halogen atom(s) and
 - 6) -NR₄₇-
- wherein R₄₇ is selected from:

- i) a hydrogen atom and
- ii) C1-C10 alkyl optionally substituted with a halogen atom(s).

[0106] More preferably, the above X is selected from the following bond or groups:

- 1) a single bond,
- 2) C1-C10 alkylene optionally substituted with cycloalkyl,
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) C1-C10 oxyalkylene.

[0107] Still more preferably, the above X is selected from the following bond or groups:

- 1) a single bond,
- 2) C1-C10 alkylene,
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) C1-C10 oxyalkylene, wherein the oxyalkylene is bonded to a 1,3,8-triaza-spiro[4.5]dec-1-en-4-one ring or a 1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione ring through a carbon atom in the oxyalkylene.


[0108] Particularly preferably, the above X is selected from the following bond or groups:

- 1) a single bond,
- 2) C1-C6 alkylene and
- 3) C1-C6 oxyalkylene optionally substituted with a halogen atom(s), wherein the oxyalkylene is bonded to a 1,3,8-triaza-spiro[4.5]dec-1-en-4-one ring or a 1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione ring through a carbon atom in the oxyalkylene.

[0109] More particularly preferably, the above X is a single bond.

[0110] The above Y is preferably selected from:

- 1) an oxygen atom,
- 2) a sulfur atom and
- 3) =NR₃₇,
- or 4) Y is -NR₃₈R₃₉ represented by the following formula (A) :



and can form tautomers;

R₃₇ is selected from:

- 1) hydrogen,
- 2) hydroxy and

3) C1-C10 alkoxy; and

R_{38} and R_{39} are independently selected from hydrogen or C1-C10 alkyl optionally substituted with cycloalkyl, or R_{38} and R_{39} may be bonded to each other to form a ring selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with C1-C10 alkyl.

[0111] More preferably, the above Y is an oxygen atom.

[0112] The above m is preferably an integer of 0 to 2, more preferably 1.

[0113] The above n is preferably an integer of 0 to 2, more preferably 1.

[0114] The above R_1 is preferably selected from:

- 1) hydrogen,
- 2) cycloalkyl optionally substituted with a group(s) selected from R_4 , R_5 and R_6 ,
- 3) a heterocycle optionally substituted with a group(s) selected from R_{25} , R_4 , R_5 and R_6 ,
- 4) aryl optionally substituted with a group(s) selected from R_3 , R_4 , R_5 and R_6 and
- 5) heteroaryl optionally substituted with a group(s) selected from R_{25} , R_4 , R_5 and R_6 .


[0115] More preferably, the above R_1 is selected from:

- 1) hydrogen,
- 2) cycloalkyl optionally substituted with a group selected from R_4 ,
- 3) a heterocycle optionally substituted with a group(s) selected from R_{25} and R_4 ,
- 4) aryl optionally substituted with a group(s) selected from R_3 , R_4 , R_5 and R_6 and
- 5) heteroaryl optionally substituted with a group(s) selected from R_{25} , R_4 and R_5 .


[0116] Still more preferably, the above R_1 is selected from:

- 1) aryl optionally substituted with a group(s) selected from R_3 , R_4 and R_5 and
- 2) heteroaryl optionally substituted with a group(s) selected from R_{25} and R_4 .

[0117] Particularly preferably, the above R_1 is the following general formula (3) or (4).



(3),



(4)

[0118] The above R_3 is preferably selected from:

- 1) $-\text{CONR}_7\text{R}_8$,
- 2) $-\text{OR}_9$,
- 3) $-\text{NR}_9\text{R}_{10}$,
- 4) $-\text{N}(\text{R}_9)\text{COR}_{11}$,
- 5) $-\text{N}(\text{R}_9)\text{SO}_2\text{R}_{12}$,
- 6) $-\text{SO}_2\text{R}_{15}$,
- 7) C1-10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, $-\text{COR}_{16}$ and $-\text{NR}_{13}\text{R}_{14}$,
- 8) heteroaryl optionally having C1-10 alkyl and/or C1-10 alkoxy as a substituent(s) and
- 9) $-\text{N}(\text{R}_9)\text{CSR}_{11}$.

[0119] More preferably, the above R_3 is selected from:

- 1) -CONR₇R₈,
- 2) -OR₉,
- 3) -NR₉R₁₀,
- 4) -N(R₉)COR₁₁,
- 5) -N(R₉)SO₂R₁₂,
- 6) -SO₂R₁₅,
- 7) C1-C10 alkyl optionally substituted with a group(s) selected from -COR₁₆ and -NR₁₃R₁₄ and
- 8) -N(R₉)CSNH₂.

[0120] Still more preferably, the above R₃ is selected from:

- 1) -CONR₇R₈,
- 2) -OR₉,
- 3) -NR₉R₁₀,
- 4) -N(R₉)COR₁₁,
- 5) -N(R₉)SO₂R₁₂,
- 6) -SO₂R₁₅ and
- 7) C1-C6 alkyl optionally substituted with a group(s) selected from -COR₁₆ and -NR₁₃R₁₄.

[0121] The above R₄ is preferably selected from:

- 1) halogen,
- 2) cyano,
- 3) nitro,
- 4) amino,
- 5) -NHCOR₂₆,
- 6) C1-C10 alkyl optionally substituted with a group(s) independently selected from hydroxycarbonyl, C1-C10 alkoxy, carbonyl and aminocarbonyl,
- 7) C1-C10 haloalkyl,
- 8) C1-C10 alkoxy,
- 9) C1-C10 haloalkylcarbonyl,
- 10) -COR₁₆,
- 11) C1-C10 hydroxyalkyl and
- 12) C1-C10 heteroalkyl.

[0122] More preferably, the above R₄ is selected from:

- 1) halogen,
- 2) cyano,
- 3) amino,
- 4) C1-C10 alkyl,
- 5) C1-C10 haloalkyl,
- 6) C1-C10 alkoxy,
- 7) C1-C10 haloalkylcarbonyl,
- 8) -COR₁₆ and
- 9) C1-C10 heteroalkyl.

[0123] The above R₅ is preferably selected from a halogen atom, C1-C10 alkyl, C1-C10 haloalkyl and C1-C10 alkoxy.

[0124] The above R₆ is preferably selected from a halogen atom, C1-C10 alkyl and C1-C10 haloalkyl.

[0125] The above R₇ is preferably selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from amino and C1-C10 alkylamino,
- 3) C1-C10 hydroxyalkyl,
- 4) C1-C10 haloalkyl,
- 5) C1-C10 heteroalkyl,
- 6) C1-C10 heteroalkyl optionally substituted with a group(s) selected from a hydroxyl group, C1-C10 alkylamino and C2-C10 alkenyl,

- 7) aryl,
- 8) heteroaryl,
- 9) aryl C1-C10 alkyl,
- 10) a heterocycle optionally substituted with C1-C10 alkyl,
- 11) $-(CH_2)_L COR_{16}$ (wherein L represents an integer of 1 to 4),
- 12) C1-C10 alkoxy,
- 13) C2-C10 alkenyl and
- 14) $-NR_{40}R_{41}$; and

R_{40} and R_{41} are independently selected from hydrogen, C1-C10 alkyl and C1-C10 alkylcarbonyl, or R_{40} and R_{41} may be bonded to each other to form a ring selected from azetidiny, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl, and the heterocycle is optionally substituted with C1-C10 alkyl.

[0126] The above R_8 is preferably selected from hydrogen and C1-C10 alkyl optionally substituted with a halogen atom(s) and/or a hydroxyl group(s).

[0127] The above R_7 and R_8 may be bonded, to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from O, N, S, SO and SO_2 , and the heterocycle optionally contains carbonyl, and the heterocycle is optionally substituted with a substituent(s) independently selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally having C1-C10 alkylamino as a substituent(s),
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 8) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 9) a heterocycle optionally substituted with C1-C10 alkyl,
- 10) heteroaryl optionally substituted with C1-C10 alkyl,
- 11) heterocyclyl C1-C10 alkyl,
- 12) $-COR_{16}$,
- 13) $-NR_{19}R_{20}$,
- 14) $-SO_2R_{21}$,
- 15) C1-C10 alkoxy-C1-C10 alkyl optionally having a hydroxyl group(s) as a substituent(s) and
- 16) C1-C10 hydroxyalkyloxy, wherein the hydrogen atom of the hydroxyl group may be replaced by C1-C10 hydroxyalkyl, and

the heterocycle may further form a spiro ring together with a 4- to 6-membered heterocycle, and the bonded 4- to 6-membered heterocycle optionally contains O and N as ring-forming elements in addition to carbon atoms, and the carbon atom(s) may be oxidized to form carbonyl, and the 4- to 6-membered heterocycle is optionally further substituted with C1-C10 alkyl.

[0128] The above R_{16} is preferably selected from:

- 1) a hydroxyl group,
- 2) C1-C10 alkoxy,
- 3) $NR_{17}R_{18}$ and
- 4) C1-C10 alkyl optionally substituted with a substituent(s) selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, carbonyl or C1-C10 alkylamino.

[0129] The above R_{17} is preferably selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) selected from aryl, amino, C1-C10 alkylamino, C1-C10 alkylcarbonylamino and a hydroxyl group,
- 3) heteroaryl and
- 4) C1-C10 alkoxy.

[0130] The above R_{18} is preferably selected from hydrogen, C1-C10 alkyl and C1-C10 hydroxyalkyl.

[0131] The above R_{17} and R_{18} may be bonded to each other to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl, a halogen atom and C1-C10 alkoxy carbonyl.

[0132] The above R_{19} is preferably selected from hydrogen, C1-C10 alkyl, C1-C10 haloalkyl, C1-C10 alkyl carbonyl, C1-C10 hydroxyalkyl, C1-C10 aminoalkyl, C1-C10 alkoxy carbonyl and C1-C10 heteroalkyl.

[0133] The above R_{20} is preferably selected from hydrogen and C1-C10 alkyl.

[0134] The above R_{19} and R_{20} may be bonded to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom.

[0135] The above R_{21} is preferably selected from:

- 1) C1-C10 alkyl optionally substituted with aryl,
- 2) amino,
- 3) C1-C10 alkylamino and
- 4) aryl optionally substituted with C1-C10 alkyl.

[0136] The above R_9 is preferably selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from R_{23} ,
- 3) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 4) cycloalkyl optionally substituted with a halogen atom(s) or a hydroxyl group(s),
- 5) a heterocycle optionally substituted with a group(s) independently selected from C1-C10 alkyl, C1-C10 alkyl carbonyl, C1-C10 alkoxy, C1-C10 alkoxy carbonyl, amino and a halogen atom,
- 6) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom and a hydroxyl group,
- 7) heteroaryl optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkyl carbonyl, C1-C10 alkoxy carbonyl and a halogen atom and
- 8) cycloalkenyl optionally substituted with a group(s) selected from C1-C10 alkoxy, C1-C10 alkylamino, amino, a hydroxyl group and a halogen atom, wherein the cycloalkenyl may contain a carbonyl group.

[0137] More preferably, the above R_9 is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from R_{23} ,
- 3) cycloalkyl optionally substituted with a halogen atom(s) or a hydroxyl group(s),
- 4) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkyl carbonyl, C1-C10 alkoxy, C1-C10 alkoxy carbonyl, amino and a halogen atom,
- 5) C1-C10 heteroalkyl optionally substituted with a group(s) selected from a halogen atom and a hydroxyl group,
- 6) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl, C1-C10 alkyl carbonyl, C1-C10 alkoxy carbonyl and a halogen atom and
- 7) cycloalkenyl optionally substituted with a group(s) selected from C1-C10 alkoxy, C1-C10 alkylamino, amino, 1 to 3 hydroxyl groups and 1 to 4 halogen atoms, wherein the cycloalkenyl may contain a carbonyl group.

[0138] The above R_{23} is preferably selected from:

- 1) a halogen atom,
- 2) a hydroxyl group,
- 3) a C1-C10 alkyl carbonyloxy group,
- 4) $-COR_{16}$,
- 5) amino,
- 6) C1-C10 alkylamino,
- 7) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkyl carbonyl, C1-C10 alkoxy carbonyl and a halogen atom and
- 8) cyano.

[0139] The above R_{10} is preferably selected from:

- 1) hydrogen and
- 2) C1-C10 alkyl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group and aryl.

[0140] R_9 and R_{10} may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from N, O, S, SO, SO_2 , carbonyl and thiocarbonyl, and the heterocycle is optionally substituted with a substituent(s) independently selected from R_{24} .

[0141] The above R_{24} is preferably selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from C1-C10 alkylamino and C1-C10 alkylcarbonylamino,
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 8) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 9) a heterocycle optionally substituted with C1-C10 alkyl,
- 10) heteroaryl,
- 11) heterocyclyl C1-C10 alkyl,
- 12) $-COR_{16}$,
- 13) $-NR_{19}R_{20}$ and
- 14) $-SO_2R_{21}$.

[0142] More preferably, the above R_{24} is selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from C1-C10 alkylamino and C1-C10 alkylcarbonylamino,
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 8) C1-C10 heteroalkyl optionally substituted with one to two types of groups selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 9) $-COR_{16}$ and
- 10) $-NR_{19}R_{20}$.

[0143] The above R_{11} is preferably selected from:

- 1) C1-C10 alkyl optionally substituted with a group(s) independently selected from:
 - i) a hydroxyl group,
 - ii) $-NR_{17}R_{18}$,
 - iii) a C1-C10 alkoxy group,
 - iv) a halogen atom,
 - v) C1-C10 alkoxy carbonyl,
 - vi) aminocarbonyl and
 - vii) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, amino, C1-C10 alkylamino and $-COR_{22}$,

- 2) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, amino, C1-C10 alkylamino and -COR₂₂,
 3) cycloalkyl optionally substituted with a halogen atom(s),
 4) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy, amino, C1-C10 alkylamino and a hydroxyl group,
 5) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino and a hydroxyl group,
 6) amino,
 7) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino, hydroxycarbonyl and a hydroxyl group and
 8) C2-C10 alkenyl.

[0144] More preferably, the above R₁₁ is selected from:

- 1) C1-C10 alkyl optionally substituted with 1 to 3 substituents independently selected from:
- i) a hydroxyl group,
 - ii) -NR₁₇R₁₈,
 - iii) a C1-C10 alkoxy group,
 - iv) a halogen atom,
 - v) C1-C10 alkoxy, amino, C1-C10 alkylamino and a hydroxyl group,
 - vi) aminocarbonyl,
- 2) aryl,
 3) aryl C1-C10 alkyl,
 4) cycloalkyl optionally substituted with a halogen atom(s),
 5) a heterocycle optionally substituted with C1-C10 alkyl,
 6) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino and a hydroxyl group,
 7) amino,
 8) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino, hydroxycarbonyl and a hydroxyl group and
 9) C2-C10 alkenyl.

[0145] The above R₂₂ is preferably selected from C1-C10 alkoxy, a hydroxyl group, amino and C1-C10 alkylamino.

[0146] The above R₁₂ is preferably selected from:

- 1) C1-C10 alkyl,
- 2) amino and
- 3) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from amino, C1-C10 alkylamino and a hydroxyl group.

[0147] The above R₁₃ is preferably selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl,
- 3) C1-C10 alkylcarbonyl, wherein the alkyl is optionally substituted with a hydroxyl group(s),
- 4) C1-C10 alkoxy, amino, C1-C10 alkylamino and a hydroxyl group,
- 5) aminocarbonyl,
- 6) C1-C10 alkylaminocarbonyl and
- 7) heterocyclic carbonyl optionally substituted with C1-C10 alkyl.

[0148] The above R₁₄ is preferably selected from:

- 1) hydrogen and
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino.

[0149] Further, R_{13} and R_{14} may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from O, N, S, SO and SO_2 , and the heterocycle optionally contains carbonyl, and the heterocycle is optionally substituted with C1-C10 alkyl.

[0150] The above R_{15} is preferably selected from:

- 1) C1-C10 alkyl and
- 2) $-NR_{35}R_{36}$.

[0151] The above R_{35} is preferably selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) a hydroxyl group,
- iii) C1-C10 alkylcarbonylamino,
- iv) $-COR_{16}$,
- v) amino,
- vi) C1-C10 alkylamino,
- vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),
- viii) heteroaryl optionally substituted with a C1-C10 alkyl group(s) and
- ix) a heterocycle,

3) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,

4) cycloalkyl optionally substituted with a group(s) independently selected from a halogen atom and a hydroxyl group,

5) a heterocycle optionally substituted with a group(s) independently selected from C1-C10 alkyl, a halogen atom and aryl C1-C10 alkyl,

6) heteroaryl optionally substituted with C1-C10 alkyl and

7) C1-C10 alkylcarbonyl.

[0152] The above R_{36} is preferably selected from:

- 1) hydrogen and
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group and aryl.

[0153] The above R_{35} and R_{36} may be bonded to each other to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom.

[0154] The above R_{25} is preferably selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) aryl,
- iii) heteroaryl,
- iv) a heterocycle optionally substituted with a C1-C10 alkyl group(s),
- v) $-COR_{16}$,
- vi) $-NR_{13}R_{14}$ and
- vii) $-SO_2R_{21}$,

3) C1-C10 heteroalkyl optionally substituted with a hydroxyl group(s),

4) C1-C10 hydroxyalkyl, wherein each hydroxyl group may be independently substituted with a group(s) selected from C1-C10 alkyl, aryl C1-C10 alkyl and C1-C10 alkylcarbonyl,

5) $-COR_{16}$,

6) $-SO_2R_{21}$,

- 7) aryl and
8) cyano.

[0155] The above R_2 is preferably selected from:

- 1) C1-C10 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a substituent(s) independently selected from R_{42} ,
- 2) C2-C10 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a substituent(s) independently selected from R_{42} ,
- 3) C2-C10 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a substituent(s) independently selected from R_{42} ,
- 4) cycloalkyl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) C2-C10 alkenyl or C1-C10 alkyl,
- iii) aryl optionally substituted with 1 to 3 substituents independently selected from C1-C10 alkyl, a halogen atom, C1-C10 alkoxy, C1-C10 alkylamino and C1-C10 alkylcarbonyl,
- iv) cycloalkyl,
- v) C2-C10 alkenyl optionally substituted with halogen,
- vi) C1-C10 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the , alkylidene is optionally substituted with a halogen atom(s),
- vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),
- viii) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom or C1-C10 alkoxy optionally substituted with a halogen atom(s),
- ix) C2-C10 alkynyl and
- x) $-\text{Si}(\text{R}_{43})_3$,

5) a heterocycle, wherein the heterocycle is optionally substituted with a group(s) independently selected from:

- i) a C1-C10 alkyl group,
- ii) C1-C10 alkylcarbonyl, wherein the alkyl group is optionally substituted with R_{27} ,
- iii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- iv) heteroarylcarbonyl,
- v) C1-C10 alkoxycarbonyl, wherein the alkyl group is optionally substituted with a group(s) independently selected from a halogen atom, aryl and C1-C10 alkoxy,
- vi) aryloxycarbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) and/or C1-C10 alkyl,
- vii) $-\text{CONR}_{28}\text{R}_{29}$,
- viii) $-\text{SO}_2\text{R}_{21}$,
- ix) a halogen atom,
- x) cycloalkylcarbonyl optionally fused with an aryl group and
- xi) C2-C10 alkenylcarbonyl, wherein the alkenyl group is optionally substituted with aryl, wherein the aryl is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl or C1-C10 alkoxy,

6) aryl optionally substituted with a group(s) independently selected from R_{44} ,

7) heteroaryl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) C1-C10 alkyl and
- iii) C1-C10 alkoxy;

8) C1-C10 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a substituent(s) independently selected from R_{42} ,

9) $-\text{S}(\text{O})_q\text{R}_{43}$ (wherein q is an integer of 0 to 2) and

10) cycloalkenyl optionally substituted with C1-C10 alkyl.

[0156] More preferably, the above R_2 is selected from:

1) C1-C10 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a group selected from R_{42} ,

2) C2-C10 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a group selected from R_{42} ,

3) C2-C10 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a group selected from R_{42} ,

4) cycloalkyl optionally substituted with a group(s) independently selected from:

i) a halogen atom,

ii) C2-C10 alkenyl or C1-C10- alkyl,

iii) aryl optionally substituted with a group(s) independently selected from C1-C10 alkyl, a halogen atom and C1-C10 alkoxy,

iv) cycloalkyl,

v) C2-C10 haloalkenyl or C1-C10 haloalkyl,

vi) C1-C10 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with a halogen atom(s),

vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),

viii) C1-C10 alkyl substituted with C1-C10 alkoxy, wherein the alkyl and/or the alkyl in the alkoxy is optionally substituted with a halogen atom(s),

ix) C2-C10 alkynyl and

x) $-\text{Si}(\text{R}_{43})_3$,

5) a heterocycle, wherein the heterocycle is optionally substituted with a group(s) selected from:

i) a C1-C10 alkyl group,

ii) C1-C10 alkylcarbonyl, wherein the alkyl group is optionally substituted with R_{27} ,

iii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,

iv) heteroarylcarbonyl,

v) C1-C10 alkoxy carbonyl, wherein the alkyl group is optionally substituted with a group(s) independently selected from a halogen atom, aryl and C1-C10 alkoxy,

vi) aryloxy carbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) and/or C1-C10 alkyl,

vii) $-\text{CONR}_{28}\text{R}_{29}$ and

viii) $-\text{SO}_2\text{R}_{21}$,

6) aryl optionally substituted with a group(s) independently selected from R_{44} ,

7) heteroaryl optionally substituted with any of the following groups:

i) C1-C10 alkyl,

8) C1-C10 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a group selected from R_{42} ,

9) $-\text{S}(\text{O})_q\text{R}_{43}$ (wherein q is an integer of 0 to 2) and

10) cycloalkenyl optionally substituted with C1-C10 alkyl.

[0157] Still more preferably, the above R_2 is selected from:

1) C1-C13 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a group selected from R_{42} ,

2) C2-C13 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a group selected from R_{42} ,

3) C2-C13 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a group selected from R_{42} ,

4) cycloalkyl optionally substituted with a group(s) independently selected from:

i) a halogen atom,

ii) C2-C6 alkenyl or C1-C6 alkyl,

iii) aryl optionally substituted with a group(s) independently selected from C1-C6 alkyl, a halogen atom, C1-C6

alkoxy, C1-C6 alkylamino and C1-C6 alkylcarbonyl,

iv) cycloalkyl,

v) C2-C6 haloalkenyl or C1-C6 haloalkyl,

vi) C1-C6 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with a halogen atom(s),


vii) C1-C6 alkoxy optionally substituted with a halogen atom(s),

viii) C1-C6 alkyl substituted with C1-C6 alkoxy, wherein the alkyl and/or the alkoxy is optionally substituted with halogen,

ix) C2-C6 alkynyl and

x) $-\text{Si}(\text{R}_{43})_3$,

5) a group represented by the following general formula (B) :



(wherein Ra represents a group selected from:

i) C1-C6 alkylcarbonyl, wherein the alkyl group is optionally substituted with R_{27} ,

ii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C6 alkyl and C1-C6 alkoxy,

iii) C1-C6 alkoxycarbonyl, wherein the alkyl group is optionally substituted with a group(s) selected from a halogen atom, aryl and C1-C6 alkoxy,

iv) aryloxycarbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) or C1-C6 alkyl,

v) $-\text{CONR}_{28}\text{R}_{29}$ and

vi) $-\text{SO}_2\text{R}_{21}$),

6) aryl optionally substituted with a group(s) independently selected from R_{44} ,

7) heteroaryl optionally substituted with any of the following groups:

i) a halogen atom,

ii) C1-C6 alkyl and


iii) C1-C6 alkoxy;

8) C1-C6 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a group selected from R_{42} ,

9) $-\text{S}(\text{O})_q\text{R}_{43}$ (wherein q is an integer of 0 to 2) and

10) cycloalkenyl optionally substituted with C1-C6 alkyl.

[0158] When the above R_2 is a "cycloalkyl optionally substituted with 1 to 3 substituents" and the substituent is "alkylidene (wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with 1 to 5 halogen atoms)", examples of the R_2 include the following groups.



[0159] R_{44} is preferably selected from:

1) a halogen atom,

2) cyano,

3) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a hydroxyl group,
- ii) $-OR_{26}$,
- iii) cyano,
- iv) aryloxy optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl optionally substituted with a halogen atom(s) or C1-C10 alkoxy optionally substituted with a halogen atom(s) and
- v) a halogen atom,

4) cycloalkyl optionally substituted with a group(s) independently selected from a halogen atom or C1-C10 alkyl optionally substituted with a halogen atom(s),

5) C1-C10 alkoxy optionally substituted with a halogen atom(s) or a C2-C6 alkenyl group(s),

6) $-COR_{30}$,

7) C1-C10 alkylcarbonylamino,

8) C1-C10 alkoxycarbonylamino, wherein the alkoxy group is optionally substituted with aryl,

9) C1-C10 heteroalkyl optionally substituted with a halogen atom(s),

10) aryl optionally substituted with a substituent(s) independently selected from:

- i) a halogen atom,

- ii) C1-C10 alkyl,

- iii) C1-C10 alkoxy and

- iv) aryl optionally substituted with aryl optionally substituted with C1-C10 alkyl,

11) heteroaryl optionally substituted with a C1-C10 alkyl group(s),

12) $-SO_2R_{43}$,

13) $-SOR_{43}$,

14) C1-C10 alkylthio optionally substituted with a halogen atom(s),

15) $-Si(R_{43})_3$ and

16) $-SF_5$.

[0160] More preferably, R_{44} is selected from:

- 1) a halogen atom,

- 2) cyano,

- 3) C1-C10 alkyl optionally substituted with any of the following groups:

- i) a hydroxyl group,

- ii) $-OR_{26}$,

- iii) cyano and

- iv) aryloxy optionally substituted with a group(s) selected from a halogen atom, C1-C10 alkyl, C1-C10 haloalkyl or C1-C10 haloalkoxy,

4) C1-C10 haloalkyl,

5) cycloalkyl optionally substituted with a group(s) selected from a halogen atom and C1-C10 haloalkyl,

6) C1-C10 alkoxy optionally substituted with a halogen atom(s) or a C2-C6 alkenyl group(s),

7) $-COR_{30}$,

8) C1-C10 heteroalkyl optionally substituted with a halogen atom(s),

9) aryl optionally substituted with a group(s) independently selected from:

- i) C1-C10 alkyl and

- ii) aryl,

10) heteroaryl optionally substituted with a C1-C10 alkyl group(s),

11) $-SO_2R_{43}$,

12) C1-C10 alkylthio optionally substituted with a halogen atom(s),

13) $-Si(R_{43})_3$ and

14) $-SF_5$.

[0161] Still more preferably, R_{44} is selected from:

- 1) a halogen atom,
- 2) cyano,
- 3) C1-C6 alkyl optionally substituted with any of the following groups:

- i) a hydroxyl group,
- ii) $-OR_{26}$,
- iii) cyano and
- iv) aryloxy optionally substituted with a group(s) selected from a halogen atom, C1-C6 alkyl, C1-C6 haloalkyl or C1-C6 haloalkoxy,

- 4) C1-C6 haloalkyl,
- 5) cycloalkyl optionally substituted with a group(s) selected from a halogen atom and C1-C6 haloalkyl,
- 6) C1-C6 alkoxy optionally substituted with a halogen atom(s),
- 7) $-COR_{30}$,
- 8) C1-C6 heteroalkyl optionally substituted with a halogen atom(s),
- 9) aryl optionally substituted with a group(s) independently selected from:

- i) C1-C6 alkyl and
- ii) aryl,

- 10) heteroaryl optionally substituted with a C1-C6 alkyl group(s),
- 11) $-SO_2R_{43}$,
- 12) C1-C6 alkylthio optionally substituted with a halogen atom(s),
- 13) $-Si(R_{43})_3$ and
- 14) $-SF_5$.

[0162] R_{42} is preferably selected from:

- 1) hydrogen,
- 2) aryl optionally substituted with a group(s) independently selected from C1-C10 alkyl optionally substituted with halogen, a halogen atom and C1-C10 alkoxy,
- 3) hydroxycarbonyl,
- 4) C1-C10 alkoxycarbonyl,
- 5) aminocarbonyl,
- 6) C1-C10 alkylaminocarbonyl,
- 7) C1-C10 alkoxycarbonylamino,
- 8) amino,
- 9) a hydroxyl group and
- 10) oxetane, tetrahydrofuran or tetrahydropyran optionally substituted with C1-C10 alkyl.

[0163] R_{43} preferably represents a C1-C10 alkyl group.

[0164] R_{26} is preferably aryl, or C1-C10 alkyl optionally substituted with a halogen atom(s).

[0165] R_{27} is preferably selected from:

- 1) aryl optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- 2) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with aryl,
- 3) a hydroxyl group,
- 4) amino,
- 5) C1-C10 alkylamino,
- 6) hydroxycarbonyl,
- 7) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl and/or aryl, and
- 8) heteroaryloxy.

[0166] More preferably, R_{27} is selected from:

- 1) aryl optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,

- 2) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with aryl,
- 3) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl and aryl and
- 4) heteroaryloxy.

[0167] The above R_{28} is preferably selected from hydrogen or C1-C10 alkyl optionally substituted with aryl.

[0168] The above R_{29} is preferably selected from hydrogen or C1-C10 alkyl optionally substituted with aryl.

[0169] The above R_{28} and R_{29} may be bonded to form a ring selected from azetidiny, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom.

[0170] The above R_{30} is preferably selected from a hydroxyl group, C1-C10 alkoxy and $-NR_{31}R_{32}$.

[0171] Preferably, the above R_{31} and R_{32} are independently selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with aryl and
- 3) aryl.


[0172] The above R_{31} and R_{32} may be bonded to form a ring selected from azetidiny, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl, a halogen atom and C1-C10 alkoxy carbonyl.

[0173] Preferably, the above R_{33} and R_{34} are independently selected from:


- 1) hydrogen and
- 2) C1-C10 alkyl.

[0174] More preferably, the above R_{33} and R_{34} are hydrogen.

[0175] In the above formula (2), U preferably represents a bond, C1-C10 alkylene or any group selected from groups represented by the following formula.



[0176] More preferably, U is C1-C6 alkylene or any group selected from groups represented by the following formula.




[0177] A is preferably selected from O, NH and CH_2 and is more preferably O.


[0178] R_{46} is preferably selected from hydrogen or R_{44} , more preferably selected from hydrogen, C1-C10 alkyl, C1-C10 haloalkyl and C1-C10 hydroxyalkyl, and still more preferably selected from C1-C10 alkyl, C1-C10 haloalkyl and C1-C10 hydroxyalkyl.

[0179] T is preferably selected from aryl and heteroaryl.

[0180] V is preferably selected from:



[0181] More preferably, V is selected from:




[0182] E is preferably a 4- to 7-membered heterocycle optionally containing 1 to 2 additional elements or groups selected from O, N, S, SO and SO₂, and the heterocycle is optionally substituted with one substituent selected from:

- 25
- 1) hydrogen,
 - 2) a halogen atom,
 - 3) C1-C10 alkyl optionally having a group(s) independently selected from C1-C10 alkylamino, a halogen atom and a hydroxyl group,
 - 4) a hydroxyl group,
 - 30
 - 5) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
 - 6) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
 - 35
 - 7) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
 - 8) a heterocycle optionally substituted with C1-C10 alkyl,
 - 9) heteroaryl optionally substituted with C1-C10 alkyl,
 - 40
 - 10) heterocyclyl C1-C10 alkyl,
 - 11) -COR₁₆,
 - 12) -NR₁₉R₂₀ and
 - 13) -SO₂R₂₁.

[0183] More preferably, E is pyrrolidine or piperidine optionally substituted with a hydroxyl group.

[0184] Z is preferably a divalent group selected from:

- 45
- 1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s) and/or a hydroxyl group(s), wherein the carbon atom(s) may be oxidized to form carbonyl;
 - 2) C1-C10 alkenylene or C1-C10 heteroalkenylene optionally substituted with a halogen atom(s) and/or a hydroxyl group(s), wherein the carbon atom(s) may be oxidized to form carbonyl; and
 - 50
 - 3) a group selected from:



[0185] G is preferably a divalent group selected from:

- 1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s); and
- 2) C1-C10 alkenylene or a C1-C10 heteroalkenylene optionally substituted with a halogen atom(s).

[0186] J is preferably a divalent group selected from:

- 1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s); and
- 2) C1-C10 alkenylene or a C1-C10 heteroalkenylene optionally substituted with a halogen atom(s).

[0187] B is preferably selected from a heterocycle or heteroaryl.

[0188] R₄₅ is preferably selected from hydrogen or C1-C10 alkyl.

[0189] R₇ is preferably selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from amino and C1-C10 alkylamino,
- 3) C1-C10 hydroxyalkyl,
- 4) C1-C10 heteroalkyl,
- 5) C1-C10 heteroalkyl optionally substituted with 1 to 3 groups selected from a hydroxyl group, C1-C10 alkylamino and C2-C10 alkenyl,
- 6) aryl,
- 7) heteroaryl,
- 8) aryl C1-C10 alkyl,
- 9) a heterocycle optionally substituted with C1-C10 alkyl,
- 10) -(CH₂)_LCOR₁₆ (wherein L represents an integer of 1 to 4),
- 11) C1-C10 alkoxy,
- 12) C2-C10 alkenyl and
- 13) -NR₄₀R₄₁.

[0190] More preferably, R₇ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl and
- 3) C1-C10 hydroxyalkyl

[0191] Specific examples of the compound represented by the formula (1) according to the present invention include the following compounds:

- (1) 8-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (2) 8-(3-chloro-benzenesulfonyl)-2-pyridin-4-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (3) 8-(3-chloro-benzenesulfonyl)-2-propyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (4) 8-(3-chloro-benzenesulfonyl)-2-isopropyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (5) 8-(3-chloro-benzenesulfonyl)-2-(3-methoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (6) 8-(3-chloro-benzenesulfonyl)-2-(3-chloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (7) 2-benzyl-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (8) 8-(3-chloro-benzenesulfonyl)-2-methyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (10) 2-biphenyl-2-yl-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (11) 8-(3-chloro-benzenesulfonyl)-2-o-tolyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (12) 8-(3-chloro-benzenesulfonyl)-2-(4-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (13) 8-(3-chloro-benzenesulfonyl)-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (14) 8-(3-chloro-benzenesulfonyl)-2-(1-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (15) 2-(1-acetyl-piperidin-4-yl)-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (16) 2-tert-butyl-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (17) 8-(3-chloro-benzenesulfonyl)-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (18) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butyl ester;
- (19) 8-(3-chloro-benzenesulfonyl)-2-(3-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (20) 8-(3-chloro-benzenesulfonyl)-2-(2-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (21) 8-(3-chloro-benzenesulfonyl)-2-(4-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (22) 2-(1R,2S,4S)-bicyclo[2.2.1]hept-2-yl-8-(3-chlorobenzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (23) 2-(1R,2R,4S)-bicyclo[2.2.1]hept-2-yl-8-(3-chlorobenzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (24) 8-(3-chloro-benzenesulfonyl)-2-(4-methoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (25) 8-(3-chloro-benzenesulfonyl)-2-(4-methoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (26) 2-(4-tert-butyl-cyclohexyl)-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (27) 8-(3-chloro-benzenesulfonyl)-2-(4-fluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (28) 8-(3-chloro-benzenesulfonyl)-2-cyclohexylmethyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (29) 8-(3-chloro-benzenesulfonyl)-2-phenethyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (30) 8-(3-chloro-benzenesulfonyl)-2-(2-cyclohexyl-ethyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (31) 8-(3-chloro-benzenesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (32) 8-(3-chloro-benzenesulfonyl)-2-(3-methanesulfonyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (33) 8-(3-chloro-benzenesulfonyl)-2-(1-phenyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (34) 8-(3-chloro-benzenesulfonyl)-2-(2-naphthalen-1-yl-ethyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (35) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (36) 2-tert-butyl-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (37) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (38) 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butyl ester;
- (39) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(2-naphthalen-1-yl-ethanesulfonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (40) 8-(3-chloro-benzenesulfonyl)-2-[1-(propane-1-sulfonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (41) 2-(1-benzenesulfonyl-piperidin-3-yl)-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (42) 8-(3-chloro-benzenesulfonyl)-2-(1-phenylmethanesulfonyl-piperidin-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (43) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-(1-phenylmethanesulfonyl-piperidin-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (44) 8-(4-chloro-benzenesulfonyl)-2-(2,4-dichlorophenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (45) 2-(2,4-dichloro-phenyl)-8-(2-trifluoromethyl-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (46) 8-(butane-1-sulfonyl)-2-(2,4-dichloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (47) 2-(2,4-dichloro-phenyl)-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (48) 2-(2,4-dichloro-phenyl)-8-(quinoline-8-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (49) 8-(3-chloro-4-fluoro-benzenesulfonyl)-2-(2,4-dichloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (50) 8-(3-chloro-benzenesulfonyl)-2-(2,4-dichlorophenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (51) 2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-benzoic acid methyl ester;
- (52) 2-cyclohexyl-8-(5-methyl-3-phenyl-isoxazole-4-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (53) 8-(benzo[b]thiophene-3-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (54) 8-(benzo[b]thiophene-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (55) 8-(5-chloro-thiophene-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (56) 2-cyclohexyl-8-(thiophene-2-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (57) 2-cyclohexyl-8-(naphthalene-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (58) 2-cyclohexyl-8-(2,4-dimethyl-thiazole-5-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (59) 2-cyclohexyl-8-(3,5-dimethyl-isoxazole-4-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (60) 2-cyclohexyl-8-(2,3-dihydro-benzo[1,4]dioxine-6-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (61) 2-cyclohexyl-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (62) 3-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-thiophene-2-carboxylic acid methyl ester;
- (63) 5-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-4-methyl-thiophene-2-carboxylic acid methyl ester;
- (64) 2-cyclohexyl-8-(2,5-dimethyl-thiophene-3-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (65) 8-(5-bromo-thiophene-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (66) 8-(5-chloro-thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (67) 8-(naphthalene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (68) 8-(benzo[b]thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (69) 8-(5-bromo-thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (70) 8-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.6]undec-1-en-4-one;
- (71) 7-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,7-triaza-spiro[4.5]dec-1-en-4-one;
- (72) 7-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,7-triaza-spiro[4.4]non-1-en-4-one;
- (73) 4-[2-[2-(2,4-dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,N,N-trimethyl-benza-

mide;

(74) 4-[2-(2-tert-butyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;

(75) 3,N,N-trimethyl-4-(2-[4-oxo-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzamide;

(76) 3,N,N-trimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(77) 4-[2-(2-cyclopentyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;

(78) 4-{2-[2-(2,6-difluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(79) 4-{2-[2-(2,6-dimethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(80) 4-{2-[2-(3-methoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(81) 4-{2-[2-(3-chloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(82) 4-{2-[2-(3,5-bis-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(83) 4-[2-(2-benzyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;

(84) 3,N,N-trimethyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzamide;

(85) 4-{2-[2-(2-chloro-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(86) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(87) 4-(2-[2-(4-chloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3,N,N-trimethyl-benzamide;

(88) 4-{2-[2-(2,3-dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(89) 4-{2-[2-(3-chloro-4-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(90) 4-{2-[2-(2-chloro-4-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(91) 4-{2-[2-(3-bromo-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(92) 4-{2-[2-(2,2-difluoro-benzo[1,3]dioxol-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(93) 4-{2-[2-(3-chloro-2-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(94) 3,N,N-trimethyl-4-(2-{4-oxo-2-[4-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;

(95) 4-{2-[2-(4-chloro-3-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(96) 4-{2-[2-(3-fluoro-4-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(97) 3,N,N-trimethyl-4-{2-[2-(6-methyl-pyridin-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(98) 4-{2-[2-(3,4-dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(99) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(100) 4-{2-[2-(2,4-bis-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(101) 3,N,N-trimethyl-4-[2-(4-oxo-2-phenethyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzamide;

(102) 4-{2-[2-(2,4-dimethyl-thiazol-5-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(103) 4-[2-(2-cyclohexylmethyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;

(104) 3,N,N-trimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(105) 3,N,N-trimethyl-4-{2-[4-oxo-2-(2-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

- (106) 3,N,N-trimethyl-4-{2-[4-oxo-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (107) 4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- 5 (108) 4-{2-[2-(2-cyclohexyl-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (109) 4-{2-[2-(3-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (110) 4-{2-[2-(3-methanesulfonyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- 10 (111) 3,N,N-trimethyl-4-{2-[2-(2-methyl-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (112) 3,N,N-trimethyl-4-{2-[2-(2-methyl-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- 15 (113) 4-{2-[2-(2,3-dimethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (114) 4-{2-[2-(3-fluoro-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (115) 4-{2-[2-(3-fluoro-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- 20 (116) 4-{2-[2-(2-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (117) 4-{2-[2-(4-fluoro-3-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- 25 (118) 4-{2-[2-(4-difluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (119) 4-{2-[2-(2-methoxy-pyridin-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (120) 3,N,N-trimethyl-4-{2-[2-(5-methyl-pyrazin-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- 30 (121) 3,N,N-trimethyl-4-{2-[2-(4-oxo-2-thiazol-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (122) 3,N,N-trimethyl-4-{2-[2-(1-methyl-1H-imidazol-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (123) 3,N,N-trimethyl-4-{2-[4-oxo-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- 35 (124) 4-{2-[2-(4-chloro-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (125) 4-{2-[2-(4-fluoro-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- 40 (126) 4-{2-[2-(4-chloro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (127) 4-{2-[2-(3-chloro-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (128) 4-{2-[2-(4-chloro-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- 45 (129) 4-{2-[2-(4-chloro-2-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (130) 4-{2-[2-(3-isopropoxymethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- 50 (131) 4-{2-[2-(2,4-dichloro-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (133) 4-{2-[2-(3-chloro-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (134) 4-{2-[2-(2,4-dichloro-benzyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- 55 (135) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-benzyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (136) 3,N,N-trimethyl-4-{2-[4-oxo-2-(1-trifluoromethyl-cyclopropyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

nyl]-ethyl)-benzamide;

(137) 3,N,N-trimethyl-4-{2-[4-oxo-2-(1-trifluoromethyl-cyclobutyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(138) 3,N,N-trimethyl-4-{2-[4-oxo-2-(1-trifluoromethyl-cyclopentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(139) 4-{2-[2-(2-fluoro-4-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(140) 3,N,N-trimethyl-4-{2-[4-oxo-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(141) 3,N,N-trimethyl-4-{2-[4-oxo-2-((E)-propenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(142) 3,N,N-trimethyl-4-{2-[4-oxo-2-((E)-styryl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(143) 4-{2-[2-(2-benzothiazol-6-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide};

(144) 4-{2-[2-(4-methoxy-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(145) N-{3-methyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-acetamide;

(146) N-(4-{2-[2-(2,3-dimethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(147) N-(4-{2-[2-(2,3-dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(148) N-(3-methyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(149) N-(2-hydroxy-ethyl)-N-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(150) acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester;

(151) acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-[2-[4-oxo-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester;

(152) acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-[2-[4-oxo-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester;

(153) acetic acid (S)-1-acetoxymethyl-2-[acetyl-(4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-amino]-ethyl ester;

(154) 8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(155) 8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(156) 8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(157) 2-(3-chloro-phenyl)-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(158) 2-(4-chloro-phenyl)-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(159) 8-{2-[4-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(160) 8-{2-[4-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(161) 8-{2-[4-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(162) {4-[2-(2-tert-butyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-carbamic acid tert-butyl ester;

(163) (4-{2-[2-(4,4-difluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzyl)-carbamic acid tert-butyl ester;

(164) (4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzyl)-carbamic acid tert-butyl ester;

(165) (3-methyl-4-[2-[4-oxo-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzyl)-carbamic acid tert-butyl ester;

(166) 8-[2-(3-amino-phenyl)-ethanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(167) N-14-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl)-acetamide;

- (168) 3,N,N-trimethyl-4-(2-{4-oxo-2-[3-(2,2,2-trifluoro-ethoxymethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;
- (169) 4-(2-{2-[3-(2,2-difluoro-ethoxymethyl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide;
- 5 (170) 3,N,N-trimethyl-4-(2-{4-oxo-2-[3-(2,2,3,3-tetrafluoro-propoxymethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;
- (171) 4-(2-{2-[3-(3,5-dimethyl-isoxazol-4-yl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide;
- (172) 4-[2-(2-biphenyl-3-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;
- 10 (173) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-pyridin-3-yl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (174) 3-{8-[2-(4-dimethylcarbamoyl-2-methyl-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl}-benzoic acid methyl ester;
- (175) 4-(2-{2-[1-(2,4-dichloro-phenoxy)-1-methyl-ethyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide;
- 15 (176) 3,N,N-trimethyl-4-{2-[4-oxo-2-(2,3,4-trifluoro-phenoxy-methyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (177) 2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic acid (3-ethyl-phenyl)-amide;
- (178) 2-cyclohexyl-8-[(E)-2-(1H-indol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 20 (179) 2-cyclohexyl-8-[(E)-2-(2-trifluoromethyl-phenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (180) 2-cyclohexyl-8-[(E)-2-(3-methoxy-phenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (181) 2-cyclohexyl-8-[(E)-2-(2-methoxy-phenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (182) 2-cyclohexyl-8-[(E)-2-(1H-indol-4-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (183) 2-cyclohexyl-8-[(E)-2-(2-fluoro-6-trifluoromethylphenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 25 (184) 2-cyclohexyl-8-[(E)-2-(2,3-difluoro-phenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (185) 2-cyclohexyl-8-[(E)-2-(3-fluoro-2-trifluoromethylphenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (186) 2-cyclohexyl-8-[(E)-2-o-tolyl-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 30 (187) 2-cyclohexyl-8-[(E)-2-[4-(2-hydroxy-ethyl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (188) 2-cyclohexyl-8-[(E)-2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (189) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indole-1-carboxylic acid dimethylamide;
- 35 (190) N-[2-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-phenyl]-acetamide;
- (191) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (192) 2-cyclohexyl-8-[(E)-2-[1-(2-morpholin-4-yl-ethyl)-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 40 (193) 2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-N,N-dimethyl-acetamide;
- (194) 3-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethylphenyl}-1,1-dimethyl-urea;
- 45 (195) cyclopropanecarboxylic acid {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl}-amide;
- (196) N-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl]-2-hydroxy-acetamide;
- (197) {3-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-4-methyl-phenyl}-carbamic acid methyl ester;
- 50 (198) 1-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl}-3-(2-hydroxy-ethyl)-urea;
- (199) 2-cyclohexyl-8-[(E)-2-[5-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 55 (200) 2-cyclohexyl-8-[(E)-2-[2-(2-hydroxy-ethylamino)-6-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (201) 2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenylamino}-N,N-dimethyl-acetamide;

- (202) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-3-fluoro-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (203) 2-cyclohexyl-8-[(E)-2-[2-methyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (204) N-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethylphenyl]-acetamide;
- (205) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-benzyl}-carbamic acid tert-butyl ester;
- (206) 2-cyclohexyl-8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (207) 2-cyclohexyl-8-[(E)-2-[2-methyl-4-[(2-oxo-oxazolidin-5-ylmethyl)-amino]-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (208) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-N-(2-dimethylamino-ethyl)-acetamide;
- (209) 3-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzonitrile;
- (210) 8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (211) 8-[(E)-2-[4-(2-hydroxy-ethylamino)-2-trifluoromethyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (212) 8-[(E)-2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (213) 3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (214) N-(2-hydroxy-ethyl)-3,N-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (215) N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (216) 8-[(E)-2-[2-methyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethenesulfonyl]-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (217) 3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-N-(2,2,2-trifluoro-ethyl)-benzenesulfonamide;
- (218) 3-methyl-N-(2-morpholin-4-yl-ethyl)-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (219) 3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-N-pyridin-3-yl-benzenesulfonamide;
- (220) 8-[(E)-2-[4-((R)-4-hydroxy-2-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (221) 8-[(E)-2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (222) 8-[(E)-2-[4-((R)-5-hydroxymethyl-2-oxo-oxazolidin-3-yl)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (223) 8-[(E)-2-[4-[3-(3-dimethylamino-propoxy)-azetidine-1-carbonyl]-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (224) N-(4-hydroxy-butyl)-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (225) N-methyl-N-(4-methyl-3-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (226) 8-[(E)-2-(3-hydroxy-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (227) 8-[(E)-2-(5-hydroxy-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (228) N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-(1-methyl-piperidin-4-yl)-acetamide;
- (229) N-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-methanesulfonamide;
- (230) 8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (231) 3,N,N-trimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vi-

nyl)-benzamide;

(232) N-(2-hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

(233) 3-fluoro-N,N-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(234) 8-((E)-2-[2,5-dichloro-4-(3,4-dihydroxy-butoxy)-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(235) 8-((E)-2-[4-(3-dimethylamino-propoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(236) N-methyl-N-(2-methyl-3-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

(237) N-methyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

(238) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

(239) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide;

(240) 2-cycloheptyl-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(241) 2-(4,4-difluoro-cyclohexyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(242) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(243) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(2-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(244) 2-(2-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(245) 8-((E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(246) 8-((E)-2-[1-(2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(247) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(2-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(248) 2-(2-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(249) 2-(2,4-dichloro-phenyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(250) 2-(4-chloro-phenyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(251) 2-cyclohexyl-8-((E)-4-(1H-indol-4-yl)-but-3-ene-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(252) 2-cyclohexyl-8-((E)-5-(1H-indol-4-yl)-pent-4-ene-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(253) 2-cyclohexyl-8-((E)-3-[1-(2,3-dihydroxy-propyl)-1H-indol-4-yl]-prop-2-ene-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(254) 8-((E)-2-[3-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(255) 8-((E)-2-[5-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(256) 8-((E)-2-[3-(2-hydroxy-ethoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(257) N-(2-hydroxy-ethyl)-N-(4-methyl-3-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

(258) 8-((E)-2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(259) N-(4-((E)-2-[2-(2-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(260) N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-piperidin-4-yl-acetamide;

(261) 2-cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-

one;

(262) 2-cyclohexyl-8-[5-(1H-indol-4-yl)-pentane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(263) 2-cyclohexyl-8-[4-(1H-indol-4-yl)-butane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(264) N-methyl-N-(2-methyl-3-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(265) N-methyl-N-(4-methyl-3-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(266) 8-[2-[3-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(267) 8-[2-[5-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(268) 8-[2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(269) 2-cyclohexyl-8-[2-(1H-indol-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(270) 2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenylamino]-N,N-dimethyl-acetamide;

(271) cyclopropanecarboxylic acid {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl}-amide;

(272) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-benzyl}-carbamic acid tert-butyl ester;

(273) 3-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl]-1,1-dimethyl-urea;

(274) {3-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-4-methyl-phenyl}-carbamic acid methyl ester;

(275) 2-cyclohexyl-8-[2-[5-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(276) N-[2-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamide;

(277) 2-cyclohexyl-8-(2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(278) 1-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethylphenyl]-3-(2-hydroxy-ethyl)-urea;

(279) 2-cyclohexyl-8-[2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(280) 2-cyclohexyl-8-[2-[4-(2-hydroxy-ethyl)-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(281) 2-cyclohexyl-8-[2-[2-methyl-4-[(2-oxo-oxazolidin-5-ylmethyl)-amino]-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(282) 2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-N-(2-dimethylamino-ethyl)-acetamide;

(283) 2-cyclohexyl-8-[2-[2-methyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(284) 8-[2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(285) 8-[2-[4-(4-hydroxy-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(286) 8-[2-[4-(2-hydroxy-ethylamino)-2-trifluoromethylphenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(287) N-(4-hydroxy-butyl)-3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzenesulfonamide;

(288) 8-[2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(289) 8-[2-[2-methyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(290) 3-methyl-4-[2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-N-(2,2,2-trifluoro-ethyl)-benzenesulfonamide;

(291) 3-methyl-4-[2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzenesulfonamide;

(292) 3-methyl-N-(2-morpholin-4-yl-ethyl)-4-[2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzenesulfonamide;

- (293) 3-fluoro-N,N-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (294) 3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-pyridin-3-yl-benzenesulfonamide;
- (295) 8-{2-[4-(3-dimethylamino-propoxy)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (296) 8-{2-[4-((R)-5-hydroxymethyl-2-oxo-oxazolidin-3-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (297) 8-{2-[4-((R)-4-hydroxy-2-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (298) N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (299) N-methyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (300) 2-hydroxy-N-methyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (301) 1-methyl-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-urea;
- (302) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-acetamide;
- (303) 2-cyclohexyl-8-[2-(1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (304) 2-(3-trifluoromethoxy-phenyl)-8-[2-(2-trifluoromethyl-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (305) 8-[2-(1H-indol-4-yl)-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (306) 8-[2-(1H-indol-4-yl)-ethanesulfonyl]-2-(4-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (307) 8-[2-(2,2-difluoro-benzo[1,3]dioxol-4-yl)-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (308) 8-[2-(1-methyl-1H-indol-7-yl)-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (309) 2-cyclohexyl-8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (310) 4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-indole-1-carboxylic acid dimethylamide;
- (311) 8-[3-(1H-indol-4-yl)-propane-1-sulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (312) 2-cyclohexyl-8-[2-(1-methanesulfonyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (313) 2-cyclohexyl-8-[2-[1-(2-dimethylamino-ethyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (314) 8-[2-(4-amino-2-trifluoromethyl-phenyl)-ethanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (315) 2-cyclohexyl-8-[3-(3-trifluoromethyl-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (316) 2-cyclohexyl-8-(3-m-tolyl-propane-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (317) 2-cyclohexyl-8-[3-(3-hydroxy-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (318) 2-cyclohexyl-8-[3-(2-hydroxy-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (319) 8-(2-benzo[b]thiophen-3-yl-ethanesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (320) 2-cyclohexyl-8-(2-isoquinolin-5-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (321) 2-cyclohexyl-8-[2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (322) 8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl]-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (323) 2-cyclohexyl-8-[2-(2-trifluoromethyl-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (324) 4-[3-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-propyl]-N,N-dimethyl-benzamide;
- (325) 8-[2-[4-((R)-2,3-dihydroxy-propoxy)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (326) N-(2-hydroxy-ethyl)-N-(4-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (327) N-(2-hydroxy-ethyl)-N-(2-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (328) 8-[3-(3-amino-phenyl)-propane-1-sulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (329) N,N-dimethyl-4-{5-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-

2-yl)-benzamide;

(330) N,N-dimethyl-3-{5-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-benzamide;

(331) 3,N,N-trimethyl-4-{5-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-benzamide;

(332) N-{4-[5-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-thiophen-2-yl]-3-methyl-phenyl}-acetamide;

(333) 4-{5-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-3,N,N-trimethyl-benzamide;

(334) 3-{5-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-4,N,N-trimethyl-benzamide;

(335) 2-methyl-3'-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-biphenyl-4-carboxylic acid dimethylamide;

(336) 2-methyl-4'-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-biphenyl-4-carboxylic acid dimethylamide;

(337) 3,N,N-trimethyl-4-{6-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-pyridin-2-yl}-benzamide;

(338) 2-cyclohexyl-8-{2-[2-(3,5-dimethyl-isoxazol-4-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(339) 2-cyclohexyl-8-{2-[2-(6-methoxy-pyridin-3-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(340) 2-cyclohexyl-8-{2-[2-(1-methyl-1H-pyrazol-4-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(341) 8-{5-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-thiophene-2-sulfonyl}-2-(4-fluoro-3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(342) N-(4-{5-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(343) 2-cyclohexyl-8-((E)-2-thiazol-2-yl-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(344) 2-cyclohexyl-8-((E)-2-cyclopentyl-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(345) 2-cyclohexyl-8-(2-thiazol-2-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(346) 2-cyclohexyl-8-[3-(4-methoxy-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(347) N-benzyl-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-benzamide;

(348) 8-(3-chloro-benzenesulfonyl)-2-[3-(morpholine-4-carbonyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(349) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-N-methyl-N-phenyl-benzamide;

(350) N-benzyl-3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-benzamide;

(351) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-N,N-dimethyl-benzamide;

(352) 8-{2-[4-(4-methanesulfonyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(353) N-[(R)-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoyl)-pyrrolidin-3-yl]-acetamide;

(354) 8-{2-[2-methyl-4-((S)-2-trifluoromethyl-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(355) 8-{2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one

(356) 8-{2-[4-((S)-2-hydroxymethyl-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(357) 8-(2-{4-[4-(2-hydroxy-ethyl)-piperidine-1-carbonyl]-2-methyl-phenyl}-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(358) 8-(2-{2-methyl-4-[4-(3-methyl-[1,2,4]oxadiazol-5-yl)-piperidine-1-carbonyl]-phenyl}-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(359) 8-{2-[2-methyl-4-(4-pyrimidin-2-yl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(360) 4-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoyl)-piperazine-1-sulfonic acid dimethylamide;

(361) 8-{2-[2-methyl-4-(4-pyridin-2-yl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(362) 8-[2-(4-{4-[2-(2-hydroxy-ethoxy)-ethyl]-piperazine-1-carbonyl}-2-methyl-phenyl)-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(363) 8-(2-{2-methyl-4-[4-(2-morpholin-4-yl-ethyl)-piperazine-1-carbonyl]-phenyl}-ethanesulfonyl)-2-(3-trifluorome-

thyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(364) 8-{2-[2-methyl-4-(4-thiazol-2-yl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(365) 8-{2-[4-(4,4-difluoro-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(366) 8-{2-[4-(3-hydroxy-propyl)-piperazine-1-carbonyl]-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(367) (S)-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoyl)-pyrrolidine-2-carboxylic acid amide;

(368) 8-{2-[2-methyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(369) (R)-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoyl)-pyrrolidine-2-carboxylic acid amide;

(370) 8-{2-[4-((S)-3-hydroxy-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(371) 8-{2-[4-((R)-2-hydroxymethyl-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(372) 8-{2-[4-((S)-3-dimethylamino-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(373) 8-{2-[4-(4-tert-butyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(374) 8-(2-[4-(4-(3-dimethylamino-propyl)-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(375) 8-(2-[4-(4-(4-fluoro-phenyl)-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(376) 8-{2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(377) 8-{2-[4-(3-hydroxy-azetidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(378) 8-{2-[4-(3-fluoro-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(379) 8-{2-[4-(3-fluoro-azetidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(380) 8-{2-[2-methyl-4-(piperidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(381) 8-{2-[4-(azetidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(382) 4-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoyl)-piperazine-1-carboxylic acid dimethylamide;

(383) 3,5,N,N-tetramethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(384) 8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(385) 8-((E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(386) 8-((E)-2-[4-(4-hydroxy-4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(387) 8-((E)-2-[4-((3R,4R)-3-dimethylamino-4-hydroxy-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(388) 8-((E)-2-[2,6-dimethyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(389) N-(3-hydroxy-propyl)-3,5,N-trimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(390) N-(2-dimethylamino-ethyl)-3,5,N-trimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(391) N-(3-dimethylamino-propyl)-3,5,N-trimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(392) N-carbamoylmethyl-3,5,N-trimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-

ene-8-sulfonyl]-vinyl)-benzamide;

(393) 3,5,N-trimethyl-N-(1-methyl-piperidin-4-yl)-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(394) 8-((E)-2-[4-(4-acetyl-piperazine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(395) 4-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoyl)-piperazine-1-carboxylic acid dimethylamide;

(396) 4-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoyl)-piperazine-1-carboxylic acid amide;

(397) 8-((E)-2-[4-(4-(3-dimethylamino-propyl)-piperazine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(398) 8-((E)-2-[2,6-dimethyl-4-((R)-3-methylamino-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(399) 8-((E)-2-[4-((R)-3-dimethylamino-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(400) 8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(401) 8-((E)-2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(402) 8-((E)-2-[4-((S)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(403) 8-((E)-2-[4-(4-hydroxy-4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(404) 8-((E)-2-[4-((R)-3-amino-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(405) 8-((E)-2-[4-(4-dimethylamino-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(406) 8-((E)-2-[4-((3S,4S)-3-hydroxy-4-isopropylamino-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(407) 8-((E)-2-[4-(4-(2-dimethylamino-ethoxy)-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(408) 8-((E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(409) 2,N,N-trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(410) 2-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(411) 8-{2-[2-methyl-3-(morpholine-4-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(412) 4,N,N-trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(413) N-(2-hydroxy-ethyl)-4,N-dimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(414) 8-{2-[2-methyl-5-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(415) 4,N-dimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(416) 8-(2-[4-(4-(2-hydroxy-ethyl)-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(417) 8-{2-[2-methyl-4-(thiazolidine-3-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(418) 2-fluoro-5,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(419) 8-{2-[5-fluoro-4-(4-hydroxy-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(420) N-benzyl-2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-1H-indol-3-yl}-acetamide;

(421) 2-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-N-methyl-benzamide;

- (422) 3-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-N-pyridin-4-yl-benzamide;
- (423) 8-(3-chloro-benzenesulfonyl)-2-[1-(3,3-dimethyl-butyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (424) 8-(3-chloro-benzenesulfonyl)-2-[1-(2-hydroxy-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (425) 8-(3-chloro-benzenesulfonyl)-2-[1-(4-chloro-benzoyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (426) 8-(3-chloro-benzenesulfonyl)-2-[1-(3-methoxybenzoyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (428) 2-[1-(1H-indole-5-carbonyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (429) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(3-phenyl-propionyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (430) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(1,2,3,4-tetrahydro-naphthalene-2-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (431) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(quinoline-6-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (432) 2-[1-(2-3H-imidazol-4-yl-acetyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (433) 2-[1-(2,5-dimethyl-thiazol-4-yl)-acetyl]-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (434) 2-[1-(2-(5-methyl-2-phenyl-oxazol-4-yl)-acetyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (435) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(2-pyridin-2-yl-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (436) 2-[1-(4-benzyloxy-butyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (437) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(4-phenyl-butyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (438) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(2-pyridin-4-yl-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (439) 2-[1-(2-tert-butoxy-acetyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (441) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-((E)-(3-phenyl-acryloyl))-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (442) 2-[1-(2-amino-acetyl)-piperidin-3-yl]-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (444) 8-[2-[2-methyl-4-(3-methylamino-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (445) 8-[2-[4-(3-amino-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (446) 8-((E)-2-[2,6-dimethyl-4-(piperazine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (447) 8-((E)-2-[4-[4-(2-hydroxy-acetyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (448) 4-[4-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoyl)-piperazin-1-yl]-4-oxo-butyric acid methyl ester;
- (449) 8-((E)-2-[4-[4-(4-dimethylamino-butyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (450) 2-methoxy-N-methyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (451) 2-hydroxy-N-methyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (452) [(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoyl)-methyl-amino]-acetic acid;
- (453) 8-(3-chloro-benzenesulfonyl)-2-[1-(3,3-dimethylbutyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (454) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butylamide;
- (455) 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid phenethyl-amide;
- (456) 8-(3-chloro-benzenesulfonyl)-2-[1-(piperidine-1-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (457) 3-(2-dimethylamino-ethyl)-1-methyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea;
- (458) 1-methyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea;
- 5 (459) 1-{3-[2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-2-methyl-phenyl}-3-methyl-urea;
- (460) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid 2-methoxy-ethyl ester;
- (461) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid benzyl ester;
- 10 (462) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid isobutyl ester;
- (463) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid methyl ester;
- 15 (464) 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid benzyl ester;
- (465) 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid 2,2-dimethyl-propyl ester; .
- (466) methyl-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-carbamic acid 2-dimethylamino-ethyl ester;
- 20 (467) methyl-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-carbamic acid methyl ester;
- (468) methyl-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-carbamic acid methyl ester;
- 25 (469) 2-cyclohexyl-8-(2-o-tolyl-ethanesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-ene-4-thione;
- (470) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (471) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- 30 (472) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (474) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-thiourea;
- (475) 8-{2-[2,6-dimethyl-4-(methyl-thiazol-2-yl-amino)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- 35 (476) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-acetic acid methyl ester;
- (477) 2-cyclohexyl-8-((E)-2-[1-(morpholine-4-carbonyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- 40 (478) 2-cyclohexyl-8-((E)-2-[1-(4-hydroxy-butyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- (479) 2-cyclohexyl-8-[(E)-2-(1H-indol-7-yl)-ethenesulfonyl]-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- (480) 2-cyclohexyl-8-((E)-2-[1-(3,4-dihydroxy-butyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- 45 (481) N-(2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-ethyl)-acetamide;
- (482) N-(1-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-ethyl)-acetamide;
- (483) 2-cyclohexyl-8-[(E)-2-(1-thiazol-2-ylmethyl-1H-indol-4-yl)-ethenesulfonyl]-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- 50 (484) 3-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-propane-1-sulfonic acid amide;
- (485) 4-methyl-piperazine-1-carboxylic acid {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl}-amide;
- 55 (486) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-benzyl}-carbamic acid isobutyl ester;
- (487) 2-cyclohexyl-8-((E)-2-{1-[2-(2-hydroxy-ethoxy)-ethyl]-1H-indol-5-yl}-ethenesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one;

- (488) 2-cyclohexyl-8-[(E)-2-(6-methyl-1H-indol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (489) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-benzyl}-2-hydroxy-acetamide;
- (490) 3-(2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-ethyl)-1,1-dimethyl-urea;
- (491) 8-[(E)-2-(4-amino-2,6-bis-trifluoromethyl-phenyl)-ethenesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (492) 2-cyclohexyl-8-[(E)-2-(6-trifluoromethyl-1H-indol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (493) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenylamino}-acetic acid methyl ester;
- (494) 2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenylamino}-N-(2-hydroxy-ethyl)-acetamide;
- (495) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-N-(2-hydroxy-ethyl)-N-methyl-3-trifluoromethyl-benzenesulfonamide;
- (496) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-N,N-bis-(2-hydroxy-ethyl)-3-trifluoromethyl-benzenesulfonamide;
- (497) 2-cyclohexyl-8-[(E)-2-[4-(2-dimethylamino-ethylamino)-2-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (498) N-(2-acetyl-amino-ethyl)-2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenylamino}-acetamide;
- (499) 2-cyclohexyl-8-[(E)-2-(6-trifluoromethyl-1H-benzimidazol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (500) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-5-fluoro-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (501) 2-cyclohexyl-8-[(E)-2-{4-[(2-hydroxy-ethyl)-methyl-amino]-2-trifluoromethyl-phenyl}-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (502) 2-cyclohexyl-8-[(E)-2-[4-(1,1-dioxo-1 λ ⁶-thiomorpholine-4-carbonyl)-2-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (503) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-N-(2-dimethylamino-ethyl)-3-methyl-benzenesulfonamide;
- (504) 3-(3-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (505) 1-methyl-3-(3-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (506) N-(3-isopropoxy-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (507) 3-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (508) 3-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1,5,5-trimethyl-imidazolidine-2,4-dione;
- (509) N-(3-ethoxy-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (510) N-(3-ethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (511) 5,5-dimethyl-3-(3-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (512) N-(3-methoxymethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (513) 2-(4-methyl-cyclohexyl)-8-[(E)-2-[2-methyl-4-(2-oxo-oxazolidin-3-yl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (514) N-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-(2-methoxy-ethyl)-acetamide;
- (515) 8-[(E)-2-[2,6-dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (516) 8-[(E)-2-[2,6-dimethyl-4-(2-oxo-piperidin-1-yl)-phenyl]-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (517) N-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzyl)-N-methyl-acetamide;

- (518) 3-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl)-imidazolidine-2,4-dione;
- (519) 3-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- 5 (520) 8-((E)-2-[2,6-dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethenesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (521) 1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl)-pyrrolidine-2,5-dione;
- (522) N-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl)-acetamide;
- 10 (523) N-(3,5-difluoro-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide;
- (524) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 15 (525) [(S)-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-2,5-dioxo-pyrrolidin-3-yl]-carbamic acid tert-butyl ester;
- (526) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (527) 8-((E)-2-[2-methyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 20 (528) 8-((E)-2-[2-methyl-4-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethylamino]-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (529) 2-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenylamino)-acetamide;
- 25 (530) 3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-N-pyridin-4-yl-benzenesulfonamide;
- (531) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (532) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3-bis-(2-hydroxy-ethyl)-urea;
- 30 (533) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3-bis-(2-hydroxy-ethyl)-3-methyl-urea;
- (534) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-(tetrahydro-pyran-4-yl)-methanesulfonamide;
- 35 (535) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (536) 3-{3-methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-imidazolidine-2,4-dione;
- (537) 3-{4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethylphenyl}-imidazolidine-2,4-dione;
- 40 (538) 4-{3-methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-morpholine-3,5-dione;
- (539) 3-{3,5-dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-imidazolidine-2,4-dione;
- 45 (540) 5,5-dimethyl-3-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-imidazolidine-2,4-dione;
- (541) 8-((E)-2-[1-((R)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (542) N-(2-methoxy-ethyl)-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-acetamide;
- 50 (543) N-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-N-(2-methoxy-ethyl)-acetamide;
- (544) N-[2-(2-hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-acetamide;
- 55 (545) N-[2-(2-fluoro-ethoxy)-ethyl]-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-acetamide;
- (546) N-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-N-methyl-acetamide;

- (547) 3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzoic acid;
- (548) 3-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-imidazolidine-2,4-dione;
- (549) 8-((E)-2-{4-[4-(2-fluoro-ethyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl}-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (550) 8-((E)-2-[2,6-dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (551) N-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-N-[2-(2-fluoro-ethoxy)-ethyl]-acetamide;
- (552) 3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzoic acid N'-acetyl-hydrazide;
- (553) 8-[(E)-2-(4-hydroxymethyl-2,6-dimethyl-phenyl)-ethenesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (554) 3-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-imidazolidine-2,4-dione;
- (555) 3-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-5,5-dimethyl-imidazolidine-2,4-dione;
- (556) 8-[(E)-2-[2,6-dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethenesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (557) 1-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-pyrrolidine-2,5-dione;
- (558) N-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-acetamide;
- (559) N-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-N-methyl-acetamide;
- (560) 3-(4-[(E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3-methyl-phenyl)-imidazolidine-2,4-dione;
- (561) N-(4-[(E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-N-[2-(2-methoxy-ethoxy)-ethyl]-acetamide;
- (562) N-(4-[(E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-N-[2-(2-fluoro-ethoxy)-ethyl]-acetamide;
- (563) 3-(4-[(E)-2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3-methyl-phenyl)-imidazolidine-2,4-dione;
- (564) 3-(4-[(E)-2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;
- (565) 3-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-benzyl]-1,1-dimethyl-urea;
- (566) (2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-ethyl)-carbamic acid methyl ester;
- (567) 4-methyl-piperazine-1-carboxylic acid 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-benzylamide;
- (568) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-phenyl}-N',N'-dimethyl-sulfamide;
- (569) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-benzyl}-(3-dimethylamino-propyl)-carbamic acid tert-butyl ester;
- (570) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-N-(2,2-dimethyl-propyl)-acetamide;
- (571) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-N-(2-diisopropylamino-ethyl)-acetamide;
- (572) 2-cyclohexyl-8-[(E)-2-[2-methyl-4-(2-oxo-azetidin-1-yl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (573) 2-cyclohexyl-8-[(E)-2-[2,6-dimethyl-4-(thiazol-2-ylamino)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (574) N-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethylphenyl]-N-(3-methyl-oxetan-3-ylmethyl)-acetamide;
- (575) 3-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;

- (576) N-(3-ethoxymethyl-4-((E)-2-[2-(4-methyl cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (577) 2-(4-methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(4-oxo-oxazolidin-3-yl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 5 (578) 4-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-3-oxo-piperazine-1-carboxylic acid tert-butyl ester;
- (579) 2-(4-methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(4-methyl-2-oxo-oxazolidin-3-yl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 10 (580) 2-(4-methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(3-oxo-isoxazolidin-2-yl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (581) 3-(2,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione;
- (582) 3-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-5-methyl-imidazolidine-2,4-dione;
- 15 (583) 3-(2,6-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione;
- (584) N-(2,6-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide;
- (585) 8-((E)-2-(3-methoxy-2-methyl-phenyl)-ethenesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 20 (586) 8-((E)-2-{4-[(1H-imidazol-2-yl)-methyl-amino]-2,6-dimethyl-phenyl}-ethenesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (587) 1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-dihydro-pyrimidine-2,4-dione;
- 25 (588) 2-(4-methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(5-oxo-pyrazolidin-1-yl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (589) 5-tert-butoxymethyl-3-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione;
- 30 (590) N,N-dimethyl-2-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonylamino)-acetamide;
- (591) 3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-N-pyridin-3-ylmethyl-benzenesulfonamide;
- (592) N-(4-hydroxy-cyclohexyl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonamide;
- 35 (593) 8-((E)-2-(4-methanesulfonyl-2-methyl-phenyl)-ethenesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (594) N-acetyl-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonamide;
- (595) N-(2-hydroxy-1,1-bis-hydroxymethyl-ethyl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonamide;
- 40 (596) N-(1-benzyl-piperidin-4-yl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonamide;
- (597) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 45 (598) 8-((E)-2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (599) N-((R)-2,3-dihydroxy-propyl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonamide;
- 50 (600) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-3,5-difluoro-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (601) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-3,5-dimethylphenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (602) 4-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester;
- 55 (603) N-isopropyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (604) 4-[acetyl-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester;

- (605) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-(2-hydroxy-ethyl)-urea;
- (606) 8-((E)-2-[4-((R)-2-(isopropylamino-methyl)-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (607) N-(2-dimethylamino-ethyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (608) 2-dimethylamino-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide;
- (609) N-(2-dimethylamino-ethyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methanesulfonamide;
- (610) N-[(R)-1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-5-oxo-pyrrolidin-2-ylmethyl]-acetamide;
- (611) 8-((E)-2-[4-((R)-2-dimethylaminomethyl-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (612) N-cyanomethyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (613) N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-(2,2,2-trifluoro-ethyl)-acetamide;
- (614) 3-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,5,5-trimethyl-imidazolidine-2,4-dione;
- (615) 3-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (616) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-(tetrahydro-pyran-4-yl)-acetamide;
- (617) 8-((E)-2-[2-methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (618) 3-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione
- (619) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (620) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methylphenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (621) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methylphenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (622) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (623) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (624) 1-(4-((E)-2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methylphenyl)-imidazolidine-2,4-dione;
- (625) 1-(4-((E)-2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methylphenyl)-5-methyl-imidazolidine-2,4-dione;
- (626) 1-(4-((E)-2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methylphenyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (627) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethylphenyl]-ethenesulfonyl)-2-nonyl-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- (628) 3,5-dimethyl-4-((E)-2-(2-nonyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-benzoic acid;
- (629) 3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid trimethylhydrazide;
- (630) 8-((E)-2-[1-((R)-2,3-dihydroxy-propyl)-4,6-dimethyl-1H-indol-5-yl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (631) 3-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethylphenyl)-imidazolidine-2,4-dione;
- (632) N-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethylphenyl)-N-(3-methyl-oxetan-3-ylmethyl)-acetamide;
- (633) 2-cyclohexyl-8-((E)-2-quinolin-8-yl-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (634) 2-cyclohexyl-8-((E)-2-(2-oxo-2,3-dihydro-1H-indol-4-yl)-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (635) 3-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione;
- (636) 3-(4-((E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;
- (637) 2-cyclohexyl-8-[(E)-2-(2-methyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (638) 2-cyclohexyl-8-[(E)-2-(1-methyl-1,2,3,4-tetrahydro-quinolin-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (639) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethylphenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (640) 4-((E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-[1,1,1-²H₃]methyl-carbamic acid tert-butyl ester;
- (641) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (642) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (643) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (644) 8-((E)-2-[2,6-dimethyl-4-(2-oxa-7-aza-spiro[3.5]nonane-7-carbonyl)-phenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (645) 2-cyclohexyl-8-[2-(1,2,3,4-tetrahydro-quinolin-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (646) 4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-1H-indole-3-carbonitrile;
- (647) 7-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-indole-1-carboxylic acid dimethylamide;
- (648) 2-cyclohexyl-8-(2-quinolin-5-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (649) 2-cyclohexyl-8-[2-[3-(2,2,2-trifluoro-acetyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (650) 2-cyclohexyl-8-[2-(1-isopropyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (651) 2-cyclohexyl-8-[2-[1-(4-hydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (652) N-{3-cyano-4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-acetamide;
- (653) 8-(2-isoquinolin-5-yl-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (654) 8-(2-quinolin-5-yl-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (655) N-(3-methoxy-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;
- (656) 8-[2-[4-(3,3-dimethyl-1,1,4-trioxo-1λ⁶-[1,2,5]thiadiazolidin-2-yl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (657) 8-[2-[2-methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (658) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzyl}-carbamic acid isobutyl ester;
- (659) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzyl}-2-hydroxy-acetamide;
- (660) 2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenylamino}-N-(4-hydroxy-butyl)-acetamide;
- (661) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethylphenyl}-carbamic acid isobutyl ester;
- (662) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethylphenyl}-benzamide;
- (663) 2-cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzothiazol-6-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (664) N-(2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzenesulfonylamino}-ethyl)-acetamide;
- (665) N-(2-dimethylamino-ethyl)-2-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenylamino)-acetamide;
- (666) 8-[2-[4-((2R,6S)-2,6-dimethyl-morpholine-4-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (667) N-(2,2,3,3,4,4,4-heptafluoro-butyl)-3-methyl-4-[2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzenesulfonamide;
- (668) 8-[2-(3-methoxy-2-methyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (669) (2-methoxy-3,5-dimethyl-4-{2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-carbamic acid tert-butyl ester;
- (670) 8-{2-[4-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (671) 8-{2-(5,7-dimethyl-2-oxo-2,3-dihydro-benzoxazol-6-yl)-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (672) (2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-ethyl)-carbamic acid methyl ester;
- (673) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-methyl-carbamic acid tert-butyl ester;
- (674) 8-(2-{4-[3-(2-hydroxy-ethyl)-2-oxo-imidazolidin-1-yl]-2-methyl-phenyl}-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (675) 2-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonylamino)-acetamide;
- (676) 8-{2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (677) 3-(4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethylphenyl)-imidazolidine-2,4-dione;
- (678) N-cyclopentyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (679) 1-(2,3-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (680) 1-(2,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (681) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester;
- (682) 1-(3-methoxy-5-methyl-4-{2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (683) 1-(3-chloro-5-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (684) 1-(2-methoxy-3,5-dimethyl-4-{2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (685) 5,7-dimethyl-6-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-1H-quinazoline-2,4-dione;
- (686) 8-[(E)-2-[1-(2-amino-ethyl)-1H-indol-4-yl]-ethenesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (687) 8-[(E)-2-(4-aminomethyl-phenyl)-ethenesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (688) (S)-2-amino-3-hydroxy-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-propionamide;
- (689) N-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-piperidin-4-yl-methanesulfonamide;
- (690) 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (691) 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (692) 1-{3,5-dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-1-methyl-urea;
- (693) 1-(2-methoxy-3,5-dimethyl-4-[(E)-2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (694) 1-[3,5-dimethyl-4-[(E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl]-phenyl]-1-(2-fluoro-ethyl)-urea;
- (695) 1-(3-chloro-5-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (696) 1-[3,5-dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (697) 8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (698) 8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-ethoxy]-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-

propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(699) N-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide;

(700) N-[2-(2-hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-acetamide;

(701) N-[3,5-dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-N-(2-hydroxy-ethyl)-acetamide;

(702) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(704) 8-[(E)-2-[1-[2-((S)-2,3-dihydroxy-propoxy)-ethyl]-1H-indol-4-yl]-ethenesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(705) 8-[(E)-2-[2,6-dimethyl-4-((2S,3S)-2,3,4-trihydroxy-butoxy)-phenyl]-ethenesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(706) 2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-8-[(E)-2-[1-[2-((2S,3S)-2,3,4-trihydroxy-butoxy)-ethyl]-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(707) N-((S)-2,3-dihydroxy-propyl)-N-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;

(708) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester;

(709) N-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl]-N-(4-hydroxy-cyclohexyl)-acetamide;

(710) 2-cyclohexyl-8-[(E)-2-[4-[(R)-2-(2-hydroxy-ethoxymethyl)-5-oxo-pyrrolidin-1-yl]-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(711) N-(2-hydroxy-ethyl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-isobutylamide;

(712) 8-[(E)-2-[4-[(R)-5-hydroxymethyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl]-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(713) 8-[(E)-2-[4-[(R)-5-hydroxymethyl-3-methyl-2-oxo-pyrrolidin-1-yl]-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(714) 3-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-(2-hydroxy-ethyl)-5,5-dimethyl-imidazolidine-2,4-dione;

(715) N-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-(4-hydroxy-cyclohexyl)-methanesulfonamide;

(716) N-(4-[(E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide;

(717) N-(2-fluoro-5-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(718) N-(4-[(E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(719) N-(2-hydroxy-ethyl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-methanesulfonamide;

(720) N-(2-hydroxy-ethyl)-N-[3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-methanesulfonamide;

(721) N-(4-[2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(722) 2-hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;

(723) 1-[3,5-dimethyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl]-1-methyl-urea;

(724) 1-[3,5-dimethyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl]-1-[2-(2-hydroxy-ethoxy)-ethyl]-urea;

(725) 1-[3,5-dimethyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl]-1-(2-methoxy-ethyl)-urea;

(726) 1-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-isopropyl-urea;

(727) 1-(3-ethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;

(728) 1-(3-methoxy-5-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfo-

nyl]-vinyl)-phenyl)-1-methyl-urea;

(729) 1-cyanomethyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea;

(730) 1-cyclopentyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea;

(731) 1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-(2,2,2-trifluoro-ethyl)-urea;

(732) 1-(4-[2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(733) 1-((S)-2,3-dihydroxy-propyl)-1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea;

(734) N-[4-((E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-phenyl]-N-cyclopentyl-acetamide;

(735) (S)-2-amino-N-[4-((E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl]-3-methyl-butylamide;

(736) 2-[4-((E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl]-N-pyridin-4-yl-acetamide;

(737) 2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

(738) 2-cyclohexyl-8-((E)-2-[4-(4,5-dihydro-thiazol-2-ylamino)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(739) 2-cyclohexyl-8-((E)-2-[2-methyl-4-(3-methyl-oxetan-3-ylmethoxy)-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(740) 1-(3,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropyl]-phenyl)-1-methyl-urea;

(741) 1-(3,5-dimethyl-4-((E)-1-methyl-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;

(742) N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-sulfamide;

(743) N-(3-hydroxy-propyl)-N'-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-sulfamide;

(744) N-methyl-N-(3-methyl-4-((E)-3-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-propenyl)-phenyl)-acetamide;

(745) 2-cyclohexyl-8-[2-(2-methyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(746) 8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(747) 2-cyclohexyl-8-[2-(1-methyl-1,2,3,4-tetrahydro-quinolin-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(748) 2-cyclohexyl-8-[2-[1-(3,4-dihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(749) 3-(2,6-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-imidazolidine-2,4-dione;

(750) 2-(3-methyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(751) N-(3-ethoxy-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(752) N-(3-ethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(753) N-(3-ethoxymethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(754) N-(3-methoxymethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(755) N-(2-methoxy-ethyl)-N-[3-methyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-acetamide;

(756) N-[3,5-dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-N-(2-methoxy-ethyl)-acetamide;

(757) N-[2-(2-fluoro-ethoxy)-ethyl]-N-[3-methyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-acetamide;

- (758) N-[2-(2-hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;
- (759) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-N-methyl-acetamide;
- 5 (760) 3-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-imidazolidine-2,4-dione;
- (761) 8-(2-{4-[4-(2-fluoro-ethyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl}-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (764) 8-{2-[2,6-dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 10 (765) 8-{2-[2,6-dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (766) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-N-[2-(2-fluoro-ethoxy)-ethyl]-acetamide;
- 15 (767) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-5-methyl-imidazolidine-2,4-dione;
- (768) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzoic acid;
- (769) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzoic acid N'-acetyl-hydrazide;
- 20 (770) 8-[2-(4-hydroxymethyl-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (771) 8-{2-[2,6-dimethyl-4-(2-oxo-piperidin-1-yl)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 25 (772) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-acetamide;
- (773) N-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzyl)-acetamide;
- (774) 3-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-imidazolidine-2,4-dione;
- 30 (775) 3-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-5,5-dimethyl-imidazolidine-2,4-dione;
- (776) 8-{2-[2,6-dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 35 (777) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-pyrrolidine-2,5-dione;
- (778) 3-(2,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;
- (779) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzyl)-imidazolidine-2,4-dione;
- 40 (780) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (781) 8-{2-[2,6-dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 45 (782) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzyl)-pyrrolidine-2,5-dione;
- (783) N-(2,6-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-N-methyl-acetamide;
- (784) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-N-methyl-acetamide;
- 50 (785) N-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzyl)-N-methyl-acetamide;
- (786) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-dihydro-pyrimidine-2,4-dione;
- 55 (787) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzoic acid trimethylhydrazide;
- (788) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-(8,8,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (789) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (790) 3-{3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-imidazolidine-2,4-dione;
- (791) 8-{2-[1-((R)-2,3-dihydroxy-propyl)-4,6-dimethyl-1H-indol-5-yl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (792) 3-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;
- (793) 8-{2-[2-methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (794) N-(4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;
- (795) 3-{4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl}-imidazolidine-2,4-dione;
- (796) 3,5-dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzoic acid;
- (797) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (798) 8-{2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethylphenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (799) 3-(3,5-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;
- (800) 3-(4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;
- (801) 8-{2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethylphenyl]-ethanesulfonyl}-2-(4-fluoro-3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (802) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (803) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (804) 8-{2-[2,6-dimethyl-4-(2-oxa-7-aza-spiro[3.5]nonane-7-carbonyl)-phenyl]-ethanesulfonyl}-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (812) N-(3-isopropoxy-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (817) 1-{3,5-dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- (818) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-(2-fluoro-ethyl)-urea;
- (820) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide;
- (821) 8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethylphenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (822) 8-(2-{1-[2-((S)-2,3-dihydroxy-propoxy)-ethyl]-1H-indol-4-yl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (823) 2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-8-(2-{1-[2-((2S,3S)-2,3,4-trihydroxy-butoxy)-ethyl]-1H-indol-4-yl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (824) N-((S)-2,3-dihydroxy-propyl)-N-(3,5-dimethyl-4-{2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (825) N-[2-(2-hydroxy-ethoxy)-ethyl]-N-(3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl)-acetamide;
- (826) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-[2-(2-hydroxy-ethoxy)-ethyl]-urea;
- (827) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-(2-methoxy-ethyl)-urea;
- (828) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-isopropyl-urea;
- (829) 1-(3-ethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

- (830) 1-((S)-2,3-dihydroxy-propyl)-1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-urea;
- (831) 8-{2-[4-(4-hydroxy-4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 5 (832) 3-{3,5-dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-imidazolidine-2,4-dione;
- (833) 5-tert-butoxymethyl-3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;
- 10 (834) 2-(4-methyl-cyclohexyl)-8-{2-[2-methyl-4-(5-oxo-pyrazolidin-1-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (835) 2-cyclohexyl-8-[2-(1H-indol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (836) 2-cyclohexyl-8-[2-(6-trifluoromethyl-1H-indol-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (837) 2-cyclohexyl-8-[2-(6-methyl-1H-indol-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 15 (838) 2-cyclohexyl-8-[2-{1-[2-(2-hydroxy-ethoxy)-ethyl]-1H-indol-5-yl}-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (839) 2-cyclohexyl-8-[2-(6-trifluoromethyl-1H-benzimidazol-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (840) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 20 (841) 2-cyclohexyl-8-[2-[4-(1,1-dioxo-1 λ ⁶-thiomorpholine-4-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (842) 2-cyclohexyl-8-[2-[4-(2-dimethylamino-ethylamino)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (843) 8-{2-[2-methyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 25 (844) 4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-N-(2-dimethylamino-ethyl)-3-methyl-benzenesulfonamide;
- (845) 8-(2-{2-methyl-4-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethylamino]-phenyl}-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 30 (846) 3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-pyridin-4-yl-benzenesulfonamide;
- (847) 2-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenylamino)-acetamide;
- (848) N-(1-benzyl-piperidin-4-yl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;
- 35 (849) 8-{2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (850) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 40 (851) N-((R)-2,3-dihydroxy-propyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;
- (852) 3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-pyridin-3-ylmethyl-benzenesulfonamide;
- (853) N-(4-hydroxy-cyclohexyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;
- 45 (854) N-(2-hydroxy-1,1-bis-hydroxymethyl-ethyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;
- (855) 8-{2-[4-(3,4-dihydroxy-butoxy)-3,5-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 50 (856) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (857) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (858) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[4-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 55 (859) 4-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester;
- (860) 8-{2-[4-(3,4-dihydroxy-butoxy)-3,5-difluoro-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

za-spiro[4.5]dec-1-en-4-one;

(861) 2-cyclohexyl-8-{2-[2-methyl-4-(2-oxo-azetidin-1-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(862) 1-methyl-3-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(863) 5,5-dimethyl-3-(3-methyl-4-{2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(864) 5,5-dimethyl-3-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(865) 2-(4-methyl-cyclohexyl)-8-{2-[2-methyl-4-(2-oxo-oxazolidin-3-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(866) 2-(4-methyl-cyclohexyl)-8-{2-[2-methyl-4-(4-oxo-oxazolidin-3-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(867) 8-{2-[1-((R)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(868) 8-{2-[4-[(1H-imidazol-2-yl)-methyl-amino]-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(869) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;

(870) 2-(4-methyl-cyclohexyl)-8-{2-[2-methyl-4-(2-oxo-piperazin-1-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(872) 2-cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(873) N-(2-fluoro-5-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(874) N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-sulfamide;

(875) N-(3-hydroxy-propyl)-N'-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-sulfamide;

(876) 1-cyanomethyl-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-urea;

(877) 1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-(2,2,2-trifluoro-ethyl)-urea;

(878) 1-(4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione;

(879) 1-(4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-5-methyl-imidazolidine-2,4-dione;

(880) 3-(4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione;

(881) 3-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(882) 3-(4-{2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione;

(883) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(884) 3-(4-{2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;

(885) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1,5,5-trimethyl-imidazolidine-2,4-dione;

(886) (S)-2-amino-3-hydroxy-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-propionamide;

(887) 1-(4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;

(888) 3,5-dimethyl-4-{2-[4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoic acid N,N'-dimethyl-hydrazide;

(889) 8-{2-[2-methyl-4-(piperidin-4-yloxy)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;

(890) 4-{2-[2-(4,4-difluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-

benzamide;

(891) 4-{2-[2-(4-methanesulfonyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(892) 3,N,N-trimethyl-4-(2-{4-oxo-2-[(E)-2-(3-trifluoromethyl-phenyl)-vinyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;

(893) 3,N,N-trimethyl-4-[2-(4-oxo-2-phenylethynyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzamide;

(894) 4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(895) 4-{2-[2-(4-tert-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(896) 4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(897) 3,N,N-trimethyl-4-(2-{4-oxo-2-[4-(4,4-trifluorobutyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;

(898) 4-{2-[4-(cyclohexylmethyl-amino)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-diene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(899) 4-{2-[4-dimethylamino-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-diene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(900) 4-{2-[4-[(Z)-hydroxyimino]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(901) 4-{2-[4-[(Z)-methoxyimino]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(902) 4-(2-{2-[4-(2-methoxy-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(903) N-{4-[2-(2-cyclopentyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamide;

(904) N-(3-methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(905) N-(4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(906) N-(4-{2-[2-(4-cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(907) N-(4-{2-[2-(3-cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(908) N-(4-{2-[2-(2-cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(909) N-(4-{2-[2-(4-tert-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(910) N-(4-{2-[2-(4-ethoxymethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(911) N-(3-methyl-4-{2-[4-oxo-2-(4-propoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(912) N-(4-{2-[2-(4-butoxy-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(913) N-(4-{2-[2-(4-isopropoxymethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(914) N-[4-(2-{2-[4-(3-fluoro-propoxy)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(915) N-[4-(2-{2-[4-(3-fluoro-propoxy)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(916) N-[4-(2-{2-[4-((E)-3,3-difluoro-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(917) N-{4-[2-(2-cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamide;

(918) N-14-[2-(2-adamantan-1-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl)-acetamide;

(919) N-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethoxymethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-acetamide;

(920) N-[4-(2-{2-[4-(4-chloro-phenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

- (921) N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propoxy)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;
- (922) N-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;
- 5 (923) N-(3-methyl-4-{2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-acetamide;
- (924) N-(3-methyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-acetamide;
- (925) N-(4-{2-[2-(4,4-dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;
- 10 (926) N-(3-methyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-acetamide;
- (927) N-[4-(2-{2-[4-(2-methoxy-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;
- 15 (928) [3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl]-carbamic acid tert-butyl ester;
- (929) N-(2-hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-isobutylamide;
- (930) 2-hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-acetamide;
- 20 (931) N-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-2,2,2-trifluoro-N-methyl-acetamide;
- (932) N-(3,5-dimethyl-4-{2-[2-((1S,3R)-3-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-2,2,2-trifluoro-N-methyl-acetamide;
- 25 (933) N-(3,5-dimethyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-2,2,2-trifluoro-N-methyl-acetamide;
- (934) 1-{4-[2-(2-cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;
- (935) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- 30 (936) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoroethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (937) 1-[2-chloro-3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- 35 (938) 1-(4-{2-[2-(4,4-dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (939) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propylidene)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (940) 1-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3,5-dimethyl-phenyl]-1-methyl-urea;
- 40 (941) 1-(3,5-dimethyl-4-{2-[8-oxo-6-(3-trifluoromethoxyphenyl)-2,5,7-triaza-spiro[3.4]oct-5-ene-2-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- (942) N-(4-((E)-1-fluoro-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-methyl-acetamide;
- 45 (943) 3-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-imidazolidine-2,4-dione;
- (944) 3-[3-methyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-imidazolidine-2,4-dione;
- (945) 3-[4-(2-{2-[4-((E)-3,3-difluoro-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidine-2,4-dione;
- 50 (946) 3-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidine-2,4-dione;
- (947) 3-(3-methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-imidazolidine-2,4-dione;
- 55 (948) 3-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoroethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-imidazolidine-2,4-dione;
- (949) 3-(3-methyl-4-{2-[4-oxo-2-((1S,3R)-3-propylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-imidazolidine-2,4-dione;

- (950) 3-(3-methyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;
- (951) 3-(4-{2-[2-(4,4-dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione;
- (952) 3-(3-methyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;
- (953) N-(2-chloro-4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-5-methyl-phenyl)-acetamide;
- (954) N-(2-chloro-4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;
- (955) N-[4-(2-[2-[4-(3,3-difluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;
- (956) 3-[4-(2-[2-[4-(3,3-difluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidine-2,4-dione;
- (957) 2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-N-phenyl-benzamide;
- (958) 8-(2-{2-methyl-4-[4-(1-methyl-piperidin-4-yl)-piperazine-1-carbonyl]-phenyl}-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (959) 8-((E)-2-[4-((R)-3-fluoro-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (960) 8-((E)-2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (961) 3,5,N,N-tetramethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;
- (962) 3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;
- (963) 8-((E)-2-[4-(3-fluoro-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (964) 8-[2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (965) 8-[2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (966) 8-[2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (967) 8-((E)-2-[2,6-dimethyl-4-(2-oxa-6-aza-spiro[3.3]heptane-6-carbonyl)-phenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (968) 8-((E)-2-[2,6-dimethyl-4-(3-oxo-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (969) 8-((E)-2-[4-((3R,5S)-3,5-dimethyl-piperazine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (970) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(3-hydroxy-3-methyl-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (971) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (972) 8-((E)-2-[2,6-dimethyl-4-(4-oxo-piperidine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (973) 2-[4-(3,3-difluoro-allyl)-cyclohexyl]-8-[2-[4-(4-fluoro-4-hydroxymethyl-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (974) 2-[4-(3,3-difluoro-allyl)-cyclohexyl]-8-[2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (975) 2-(4-ethyl-cyclohexyl)-8-[2-[4-[4-(2-fluoro-ethyl)-piperazine-1-carbonyl]-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (976) 2-(4-ethyl-cyclohexyl)-8-[2-[2-methyl-4-(4-oxetan-3-yl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (977) 8-[2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (978) 8-[2-[4-(azetidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (979) 8-[2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-nonyl-1,3,8-triaza-

spiro[4.5]dec-1-en-4-one;

(980) 8-{2-[2,6-dimethyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(981) 8-{2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(982) 8-{2-[4-(azetidine-1-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(983) 8-{2-[2,6-dimethyl-4-(piperidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(984) 8-{2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(985) 8-{2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(986) 8-{2-[2,6-dimethyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(987) N,N-dimethyl-2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

(988) N-methoxy-2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

(989) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(990) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(991) 8-((E)-2-[2,6-dimethyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(992) 8-((E)-2-[4-(isoxazolidine-2-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(993) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl)-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(994) 3,5,N,N-tetramethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(995) 8-{2-[2,6-dimethyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(996) 3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid hydrazide;

(997) 8-((E)-2-[2,6-dimethyl-4-(pyrazolidine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(998) N-methoxy-3,5,N-trimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(999) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1000) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1001) 8-((E)-2-[4-(isoxazolidine-2-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1002) 8-((E)-2-[4-(1,1-dioxo-1 λ ⁶-thiomorpholine-4-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1003) 3,5,N,N-tetramethyl-4-((E)-2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;

(1006) 2-cyclohexyl-8-((E)-2-[4-(4-hydroxy-4-trifluoromethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1007) 2-cyclohexyl-8-((E)-2-[2,6-dimethyl-4-(2-oxo-oxazolidine-3-carbonyl)-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1008) 2-(4-butyl-cyclohexyl)-8-((E)-2-[4-(2-hydroxy-ethoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl)-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1009) 2-(4-butyl-cyclohexyl)-8-[(E)-2-(4-[2-(2-hydroxy-ethoxy)-ethoxy]-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1010) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl)-ethanesulfonyl)-

2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
 (1011) 2-amino-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide;
 (1012) (3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methyl-carbamic acid 2-hydroxy-ethyl ester;
 (1013) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3,3-trimethyl-urea;
 (1014) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3-dimethyl-urea;
 (1015) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
 (1016) 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
 (1017) 1-(4-((E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;
 (1018) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
 (1019) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[3-(4,4,4-trifluoro-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
 (1020) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(4,4,4-trifluoro-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
 (1021) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(4-pentylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
 (1022) 1-(3,5-dimethyl-4-((E)-2-[2-(7-methylsulfanyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
 (1023) 1-[3,5-dimethyl-4-((E)-2-[2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
 (1024) 1-[3,5-dimethyl-4-[(E)-2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-1-methyl-urea;
 (1025) 1-[3,5-dimethyl-4-[(E)-2-(2-non-3-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-1-methyl-urea;
 (1026) 1-[3,5-dimethyl-4-((E)-2-[2-[10-(3-methyl-oxetan-3-yl)-decyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
 (1027) 1-[3,5-dimethyl-4-[(E)-2-(2-non-1-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-1-methyl-urea;
 (1028) 1-(4-{2-[2-(4-fluoro-3-trifluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
 (1029) 1-(4-{2-[2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
 (1030) 1-(4-{2-[2-(4-tert-butyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
 (1031) 1-[3,5-dimethyl-4-(2-{2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
 (1032) 1-[3,5-dimethyl-4-(2-{2-[10-(3-methyl-oxetan-3-yl)-decyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
 (1033) 2-(8-{2-[4-(tert-butoxycarbonyl-methyl-amino)-2,6-dimethyl-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-pyrrolidine-1-carboxylic acid tert-butyl ester;
 (1034) (4-{2-[2-(1,1-difluoro-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-methyl-carbamic acid tert-butyl ester;
 (1035) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
 (1036) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
 (1037) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(4-isopropylidene-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
 (1038) 1-[3,5-dimethyl-4-(2-[4-oxo-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
 (1039) 1-(3,5-dimethyl-4-[2-[4-oxo-2-(5,6,7,8-tetrahydro-naphthalen-2-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfo-

nyl]-ethyl)-phenyl)-1-methyl-urea;

(1040) 1-(4-{2-[2-((1S,3S,5R)-3,5-dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1041) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(1-propyl-butyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1042) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

(1043) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(6,6,7,7,7-pentafluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1044) N-(3,5-dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-N-methyl-acetamide;

(1045) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1046) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1047) N-(3,5-dimethyl-4-{2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-N-methyl-acetamide;

(1048) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[1-(4,4,4-trifluorobutyl)-cyclopropyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

(1050) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1051) 1-(4-{2-[2-(cis-3,4-dimethyl-cyclopentyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1052) 1-{4-[2-(2-dicyclopropylmethyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;

(1053) 1-{4-[2-(2-bicyclo[3.3.1]non-9-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;

(1054) 1-{4-[2-((1R,5S)-2-bicyclo[3.2.1]oct-3-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;

(1055) 1-(3,5-dimethyl-4-{2-[2-(3-methyl-cyclopentyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1056) 1-(4-{2-[2-(bicyclo[2.2.1]hept-7-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;

(1057) 1-(4-{2-[2-((1S,2S)-2-hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1058) 1-{3,5-dimethyl-4-[2-(4-oxo-2-spiro[2.5]oct-6-yl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;

(1059) 1-(4-{2-[2-(4-difluoromethylene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1060) N-[4-(2-{2-[4-(2,2-difluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;

(1061) N-[4-(2-{2-[4-(2-fluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;

(1062) N-(4-{2-[2-(4-butyl-4-fluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(1063) N-[4-(2-{2-[4-((E)-3-fluoro-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide;

(1064) N-[4-(2-{2-[4-(3-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide;

(1065) N-[4-(2-{2-[4-(3-chloro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide;

(1066) N-[4-(2-{2-[4-(3-fluoro-propylidene)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide;

(1067) N-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide;

(1068) N-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide;

- (1069) N-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;
- (1070) N-[4-(2-{2-[4-((E)-3-fluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1071) N-[4-(2-{2-[4-(2,2-difluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1072) N-[4-(2-{2-[4-(2-fluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1073) N-[4-(2-{2-[4-fluoro-4-(3-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1074) N-[4-(2-{2-[4-ethynyl-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1075) N-[4-(2-{2-[4-(2-difluoromethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1076) N-[4-(2-{2-[4-(3,3-difluoro-propylidene)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1077) N-[4-(2-{2-[4-(2-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1078) N-[4-(2-{2-[4-(1-fluoro-1-methyl-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1079) N-[4-(2-{2-[4-(4-butylidene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;
- (1080) N-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;
- (1081) 1-{3,5-dimethyl-4-[2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- (1082) 1-{3,5-dimethyl-4-[2-(2-non-3-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- (1083) 1-{3,5-dimethyl-4-[2-(2-non-1-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- (1084) 1-(3,5-dimethyl-4-{1-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropylmethyl}-phenyl)-1-methyl-urea;
- (1085) 1-(4-{2,2-difluoro-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1086) 1-[4-{2,2-difluoro-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl}-3,5-dimethylphenyl]-1-methyl-urea;
- (1087) 1-{3,5-dimethyl-4-[2-(4-oxo-2-[1,1';2',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl}-1-methyl-urea;
- (1088) 1-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea;
- (1089) 1-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[3-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea;
- (1090) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-propyl}-phenyl)-1-methyl-urea;
- (1091) 2-[4-fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1092) 2-[4-fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1093) 2-(4-chloro-3-trifluoromethoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1094) 2-(3-fluoro-4-trifluoromethoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1095) 2-(3,4-bis-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1096) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1097) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,2-trifluoro-ethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1098) 2-(4-fluoro-3-methyl-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1099) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 5 (1100) 8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1101) 8-(1,1-difluoro-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1102) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-[(E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 10 (1103) 8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(4,4,5,5,5-pentafluoropentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1104) 6-(4-methyl-cyclohexyl)-2-(2-naphthalen-1-yl-ethanesulfonyl)-2,5,7-triaza-spiro[3.4]oct-5-en-8-one;
- (1106) (11-{8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl}-undecyl)-carbamic acid tert-butyl ester;
- 15 (1107) 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(9-hydroxy-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1108) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- 20 (1109) 14-(8-[(E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecanoic acid;
- (1110) 1-(3,5-dimethyl-4-[(E)-2-[2-(4-[1,1,1-²H₃]methyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- (1111) 1-(4-[(E)-2-[2-(4-fluoro-3-trifluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- 25 (1112) 1-(3,5-dimethyl-4-[(E)-2-[2-(3-nonafluorobutyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1113) 1-(4-[(E)-2-[2-(4-chloro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- 30 (1114) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1115) 1-(4-[(E)-2-[2-(6-ethoxy-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1116) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-[3-(6,6,6-trifluoro-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- 35 (1117) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-[4-(6,6,6-trifluoro-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- (1118) 1-(4-[(E)-2-[2-(11-fluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- 40 (1119) 1-{4-[(E)-2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-1-methyl-urea;
- (1120) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-((E)-6-phenyl-hex-5-enyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1121) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-((E)-9-phenyl-non-8-enyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- 45 (1122) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(5-propylsulfanyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1123) 1-(4-[(E)-2-[2-(7-methoxy-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- 50 (1124) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(5-propoxy-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1125) 1-(4-[(E)-2-[2-(11,11-difluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1126) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-propyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- 55 (1127) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(2-propyl-benzofuran-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1128) 1-(4-[(E)-2-[2-(3-methoxy-4-pentyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-

dimethyl-phenyl)-1-methyl-urea;

(1129) 1-(4-((E)-2-[2-(4-[1,1,2,2,2-²H₅]ethyl-cyclohex-3-enyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1130) 1-(4-((E)-2-[2-(4-[1,1,2,2,2-²H₅]ethyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1131) 12-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid;

(1132) 12-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid ethyl ester;

(1133) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;

(1134) 1-(3,5-dimethyl-4-((Z)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;

(1135) 1-[3,5-dimethyl-4-(2-[4-oxo-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

(1136) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1137) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(4-trimethylsilanyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1138) 1-(4-{2-[2-((1R,3R,5S)-3,5-bis-trifluoromethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1139) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(4-trimethylsilanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1140) 1-(3,5-dimethyl-4-[2-(4-oxo-2-tridecyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl)-1-methyl-urea;

(1141) 1-(3,5-dimethyl-4-[2-(4-oxo-2-undecyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl)-1-methyl-urea;

(1142) 1-(3,5-dimethyl-4-[2-(2-octyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl)-1-methyl-urea;

(1143) 1-(4-{2-[2-((1S,3R)-3-hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1144) 1-(3,5-dimethyl-4-(2-[2-[3-(3-methyl-butyl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1145) 1-(4-{2-[2-((1S,3R)-3-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1146) 1-[3,5-dimethyl-4-(2-{2-[(1S,3S)-3-(3-methyl-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

(1147) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3,3,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1148) 1-(4-{2-[2-(4-[1,1,2,2,2-²H₅]ethyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1149) 1-(4-{2-[2-(4-[1,1,2,2,2-²H₅]ethyl-cyclohex-3-enyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1150) 1-(4-{2-[2-(4-chloro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1151) 1-(3,5-dimethyl-4-(2-[4-oxo-2-(1,9,9,9-tetrafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1152) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1153) 1-(4-{2-[2-(11-fluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1154) 1-(4-{2-[2-(6-ethylsulfanyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1155) 1-(3,5-dimethyl-4-[2-(4-oxo-2-[1,1';3',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1156) 1-(4-{2-[2-(11,11-difluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1157) 1-(3,5-dimethyl-4-[(E)-2-(4-oxo-2-[1,1';3',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vi-

nyl]-phenyl)-1-methyl-urea;

(1158) 8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1159) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1160) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1161) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-((E)-2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1162) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1163) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1164) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1165) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1166) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl)-2-(4-fluoro-3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1167) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1168) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-acetic acid;

(1169) [3-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-ureido]-acetic acid;

(1170) 12-(8-[2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid;

(1171) 10-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid;

(1172) 10-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid amide;

(1173) 12-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid amide;

(1174) 12-(8-[2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid amide;

(1175) 14-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecanoic acid amide;

(1176) 14-(8-[2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecanoic acid amide;

(1178) 8-[2-(2-amino-5,7-dimethyl-benzoxazol-6-yl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1179) 2-cyclohexyl-8-[2-[2,6-dimethyl-4-(2-oxo-oxazolidine-3-carbonyl)-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1180) N-methoxy-3,5,N-trimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzamide;

(1181) 8-[2-[4-(isoxazolidine-2-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1182) 8-[2-[4-(isoxazolidine-2-carbonyl)-2,6-dimethylphenyl]-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1183) 8-[2-[4-(1,1-dioxo-1 λ ⁶-thiomorpholine-4-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(4-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1184) 8-(2-[4-[4-((R)-2,3-dihydroxy-propoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1185) 2-(3,4-dichloro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1186) 2-(3-chloro-4-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1187) [3-(8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-phenyl]-acetonitrile;

- (1188) 2-(3-chloro-4-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1189) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-pentafluorosulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1190) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-pentafluorosulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1191) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(3-trifluoromethyl-phenoxy-methyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1192) 2-(4-fluoro-2,3-dimethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1193) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1194) 2-benzo[1,3]dioxol-5-yl-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1195) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-pentafluoroethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1196) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-pentafluoroethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1197) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-3-trifluoromethylphenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1198) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1199) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,3,3-tetrafluoro-propoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1200) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,2-trifluoro-ethoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1201) 2-[3-chloro-4-(2,2,3,3-tetrafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1202) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(4,4,5,5,5-pentafluoro-pentyloxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1203) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,3,3,3-pentafluoro-propoxy)-3-trifluoromethylphenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1204) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(4,4,4-trifluoro-butoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1205) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(3-trifluoromethyl-phenoxy-methyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1206) 2-[3-(1,1-difluoro-ethyl)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1207) 2-[3-fluoro-4-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1208) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(1H-indol-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1209) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-4-trifluoromethylphenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1210) 2-[4-chloro-3-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1211) 2-(3-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-4-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1212) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(2,2,3,3-tetrafluoro-2,3-dihydro-benzo[1,4]dioxin-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1213) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1214) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-methyl-4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1215) 2-[4-fluoro-3-(3-fluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1216) 2-[3-chloro-4-(2,2,2-trifluoro-ethoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1217) 2-(3-difluoromethyl-4-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1218) 2-[4-fluoro-3-(4,4,4-trifluoro-butoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1219) 2-(4-difluoromethyl-3-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1220) 2-(4-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1221) 2-(4-[1,1,2,2,2-²H₅]ethoxy-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1222) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-[1,2,2,2,2,2-²H₇]isopropoxy-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1223) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1224) 2-(3-[1,1,2,2,2-²H₅]ethoxy-4-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1225) 2-(4-fluoro-3-[1,2,2,2,2,2-²H₇]isopropoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1226) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,3,3-tetrafluoro-propoxy)-5-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1227) 2-[4-chloro-3-(4,4,4-trifluoro-butoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1228) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,3,3-pentafluoro-propoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1229) 2-[4-chloro-3-(2,2,3,3-tetrafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1230) 2-[4-chloro-3-(2,2,3,3-pentafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1231) 2-(3-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-4-chlorophenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1232) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(4,4,4-trifluoro-butoxy)-4-trifluoromethoxy-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1233) 2-(3-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-5-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1234) 8-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1235) 8-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1236) 2-(3-chloro-4-trifluoromethyl-phenyl)-8-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1237) 2-(3,4-dimethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1238) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-methoxy-3-methyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1239) 2-(4-fluoro-2,5-dimethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1240) 2-(4-fluoro-2,3-dimethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1241) 2-(3-chloro-4-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1242) 2-(3-chloro-4-fluoro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1243) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-pentafluoroethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1244) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-pentafluoroethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1245) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-pentafluorosulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1246) 2-[3-chloro-4-(2,2,2-trifluoro-ethoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1247) 2-[3-chloro-4-(2,2,3,3-tetrafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 5 (1248) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,2-trifluoro-ethoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1249) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,3,3,3-pentafluoro-propoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1250) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(4,4,4-trifluoro-butoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 10 (1251) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,3,3-tetrafluoro-propoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1252) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(4,4,5,5,5-pentafluoro-pentyloxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 15 (1253) 2-(4-[1,1,2,2,2-²H₅]ethoxy-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1254) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1255) 2-(4-difluoromethyl-3-fluoro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 20 (1256) 2-(3-difluoromethyl-4-fluoro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1257) 2-(3,4-dichloro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 25 (1258) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-methyl-4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1259) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(2-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1260) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-fluoro-4-(2,2,2-trifluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 30 (1261) 2-(4-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1262) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-[1,2,2,2,2,2-²H₇]isopropoxy-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 35 (1263) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1264) 2-[4-fluoro-3-(3-fluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1265) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 40 (1266) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1267) 2-[3-(1,1-difluoro-ethyl)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 45 (1268) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(1H-indol-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1269) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1270) 2-[4-chloro-3-(2,2,3,3,3-pentafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 50 (1271) 2-[4-chloro-3-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1272) 2-(3-[1,1,2,2,2-²H₅]ethoxy-4-fluoro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 55 (1273) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(4,4,4-trifluoro-butoxy)-4-trifluoromethoxy-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1274) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1275) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1276) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 5 (1277) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1278) 2-(3-chloro-4-trifluoromethyl-phenyl)-8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1279) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 10 (1280) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1281) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(2,2,3,3-tetrafluoro-2,3-dihydro-benzo[1,4]dioxin-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 15 (1282) [3-(8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-phenyl]-acetonitrile;
- (1283) 8-((E)-2-[2-methyl-4-(piperidin-4-yloxy)-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1284) 8-{2-(4-aminomethyl-2-methyl-phenyl)-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 20 (1285) 2-(4-methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(2-oxo-piperazin-1-yl)-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1286) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-pyrrolidin-2-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- 25 (1287) 2-(1,1-difluoro-ethyl)-8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (1288) 8-[(E)-2-(2,6-dimethyl-4-[1,1,1-²H₃]-methylamino-phenyl)-ethanesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (1289) 8-[(E)-2-(2,6-dimethyl-4-methylaminomethyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 30 (1290) 8-[(E)-2-(4-aminomethyl-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1291) 8-[(E)-2-(4-aminomethyl-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 35 (1292) 8-[(E)-2-(2,6-dimethyl-4-methylaminomethyl-phenyl)-ethanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1293) 8-[2-(4-amino-3-chloro-2-methyl-phenyl)-ethanesulfonyl]-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 40 (1294) N-(1-acetyl-piperidin-4-yl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (1295) 8-{2-[4-(4,5-dihydro-thiazol-2-ylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1296) N-[4-(2-[2-[4-(4-fluoro-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;
- 45 (1297) 1-(3,5-dimethyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1298) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- 50 (1299) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-5-hydroxymethyl-imidazolidine-2,4-dione;
- (1300) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl]-1-methyl-urea;
- (1301) 1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl)-1-methyl-urea;
- 55 (1302) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1303) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfo-

nyl]-vinyl)-phenyl)-1-methyl-urea;

(1304) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;

(1305) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;

(1306) 1-(3,5-dimethyl-4-[2-[4-oxo-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(1307) 1-(4-((E)-2-[2-(11-hydroxy-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1308) [11-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-undecyl]-carbamic acid tert-butyl ester;

(1309) 1-(4-((E)-2-[2-(9-hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1310) 1-(4-[2-[2-(4-isopropylidene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1311) 1-(4-[2-[2-(1,1-difluoro-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1312) 1-(4-((E)-2-[2-(4-fluoro-3-trifluoromethylphenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-[1,1,1-²H₃] methyl-urea;

(1313) [3,5-dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl]-urea;

(1314) (3,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzyl)-urea;

(1315) 1-(4-((E)-2-[2-(11-amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1316) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[5-(propane-1-sulfinyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;

(1317) 1-(4-[2-[2-(6-ethanesulfinyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1318) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[5-(propane-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;

(1319) 1-(4-((E)-2-[2-(7-methanesulfonyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1320) 1-(4-[2-[2-(6-ethanesulfonyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1321) 1-(4-((E)-2-[2-(9,9-difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1322) 1-(4-[2-[2-(9-hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1323) 1-(4-[2-[2-(9,9-difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1324) 1-(4-[2-[2-(9-amino-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1325) 1-(4-((E)-2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1326) 1-(4-((Z)-2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1327) 1-(4-[2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1328) 8-[2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(4,4,5,5,5-pentafluoro-pentyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1329) 8-[2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(4-fluoro-3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1330) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1331) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1332) 8-[2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl]-2-(4-fluoro-3-trifluoromethoxy-phenyl)-

1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1333) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1334) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-{2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1335) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1336) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1337) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-{2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1338) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[3-(4,4,4-trifluoro-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea;

(1339) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea;

(1340) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1341) 1-(3,5-dimethyl-4-{2-[2-(3-nonafluorobutyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1342) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(4-pentyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1343) 1-(3,5-dimethyl-4-{2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1344) 1-(4-{2-[2-(6-ethoxy-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1345) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[3-(6,6,6-trifluoro-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea;

(1346) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(6,6,6-trifluoro-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

(1347) 12-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid ethyl ester;

(1348) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(9-phenyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1349) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(6-phenyl-hexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1350) 14-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecanoic acid;

(1351) 1-(4-{2-[2-(7-methoxy-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1352) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(5-propoxy-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1353) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-propyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1354) 1-(4-{2-[2-(3-methoxy-4-pentyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1355) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(2-propyl-benzofuran-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1356) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[5-(propane-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1357) 1-(4-{2-[2-(7-methanesulfonyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1358) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1359) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[3-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1360) [11-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-undecyl]-carbamic acid tert-butyl ester;

(1361) 1-(4-{2-[2-(11-hydroxy-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

nyl)-1-methyl-urea;

(1362) 10-(8-[2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid;

(1363) 10-(8-[2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid amide;

(1364) 1-(3,5-dimethyl-4-{2-[2-(4-[1,1,1-²H₃]-methyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;

(1365) 1-(3,5-dimethyl-4-{(E)-2-[4-oxo-2-(9-phenyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea;

(1366) 1-(4-{2-[2-(11-amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1367) 3-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-4-ethoxy-cyclobut-3-ene-1,2-dione;

(1368) 3-amino-4-[(3,5-dimethyl-4-{2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dione;

(1369) 3-dimethylamino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dione;

(1370) N-(4-{2-[2-(4-ethynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methylphenyl)-acetamide;

(1371) 1-(4-{2-[4-{(Z)-hydroxyimino}-2-(4-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1372) 2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic acid 2-methyl-benzylamide;

(1373) 2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic acid (2-o-tolyl-ethyl)-amide;

(1374) 2-cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-2-hydroxy-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1375) 2-cyclohexyl-8-(2-oxo-2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1376) 2-cyclohexyl-8-(2-o-tolyl-ethynesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1377) 2-cyclohexyl-8-{2-(1H-indol-4-yl)-ethynesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1378) 3,5,N,N-tetramethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzamide;

(1379) 3,5,N,N-tetramethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(1380) 8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1381) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1382) 8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1383) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1384) 8-(2-[4-{(R)-2,3-dihydroxy-propoxy}-piperidine-1-carbonyl]-2,6-dimethyl-phenyl)-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1385) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzoic acid hydrazide;

(1386) 8-{2-[2,6-dimethyl-4-(pyrazolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1387) N,N-dimethyl-2-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(1388) N-methoxy-2-(3-methyl-4-{2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(1389) 1-[3,5-dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl]-1-methyl-urea;

(1390) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzyl)-1-methyl-urea;

(1391) 1-(4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-[1,1,1-²H₃]methyl-urea;

(1392) 2-cyclohexyl-8-{2-[1-((2S,3S)-2,3,4-trihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1393) 8-{2-[1-((2S,3S)-4-benzyloxy-2,3-dihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl}-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
 (1394) N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-acetamide;
 5 (1395) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 (1396) 3,N,N-trimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 (1397) 3,N,N-trimethyl-4-{2-[2-(methyl-phenyl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 10 (1398) 4-{2-[2-(benzyl-methyl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (1399) 4-{2-[2-(cyclohexylmethyl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 15 (1400) 3,N,N-trimethyl-4-{2-[2-[methyl-(4-trifluoromethyl-phenyl)-amino]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 (1401) N-(3-methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
 (1402) N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
 20 (1403) 4-{2-[2-(4-butyl-piperidin-1-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (1404) 4-{2-[2-(4-butyl-cyclohexylamino)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 25 (1405) 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethylphenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-N-pent-4-enyl-benzamide;
 (1419) N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide;
 (1443) dec-9-enoic acid {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-amide; or
 30 (1446) hept-6-enoic acid {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-amide,

or a pharmacologically acceptable salt thereof.

35 **[0192]** Such compounds represented by the above formula (1) or pharmacologically acceptable salts thereof according to the present invention are useful as compounds having a PTH-like effect, preferably PTH1 receptor agonists, and are useful for the prevention and/or treatment of osteoporosis, fracture, osteomalacia, arthritis, thrombocytopenia, hypoparathyroidism, hyperphosphatemia, tumoral calcinosis or the like, or stem cell mobilization.

40 **[0193]** The compounds or salts thereof according to the present invention can be formulated by conventional methods into tablets, powders, fine granules, granules, coated tablets, capsules, syrups, troches, inhalations, suppositories, injections, ointments, ophthalmic ointments, ophthalmic preparations, nasal preparations, ear preparations, cataplasms, lotions and the like. Commonly used excipients, binders, lubricants, colorants, correctives, and as necessary, stabilizers, emulsifiers, absorption promoters, surfactants, pH adjusters, preservatives, antioxidants and the like can be used for formulation, and they are blended with ingredients commonly used as raw materials of pharmaceutical preparations and formulated by conventional methods.

45 **[0194]** For example, oral preparations are manufactured by adding, to the compound or a pharmacologically acceptable salt thereof according to the present invention, an excipient, and as necessary, a binder, a disintegrant, a lubricant, a colorant, a corrective and the like and then formulating them into powder, fine granules, granules, tablets, coated tablets, capsules and the like by a conventional method.

50 **[0195]** Examples of these ingredients include animal and vegetable oils such as soybean oil, beef tallow and synthetic glyceride; hydrocarbons such as liquid paraffin, squalane and solid paraffin; ester oils such as octyldodecyl myristate and isopropyl myristate; higher alcohols such as cetostearyl alcohol and behenyl alcohol; silicone resin; silicone oil; surfactants such as polyoxyethylene fatty acid ester, sorbitan fatty acid ester, glycerol fatty acid ester, polyoxyethylene sorbitan fatty acid ester, polyoxyethylene hydrogenated castor oil and a polyoxyethylene-polyoxypropylene block copolymer; water-soluble polymers such as hydroxyethylcellulose, polyacrylic acid, a carboxyvinyl polymer, polyethylene glycol, polyvinylpyrrolidone and methylcellulose; lower alcohols such as ethanol and isopropanol; polyhydric alcohols such as glycerol, propylene glycol, dipropylene glycol and sorbitol; sugars such as glucose and sucrose; inorganic powders such as silicic anhydride, magnesium aluminum silicate and aluminum silicate; and purified water.

[0196] Examples of the excipients include lactose, corn starch, white soft sugar, glucose, mannitol, sorbitol, microcrystalline cellulose and silicon dioxide.

[0197] Examples of the binders include polyvinyl alcohol, polyvinyl ether, methylcellulose, ethylcellulose, acacia, tragacanth, gelatin, shellac, hydroxypropylmethylcellulose, hydroxypropylcellulose, polyvinylpyrrolidone, a polypropylene glycol-polyoxyethylene block polymer and meglumine

[0198] Examples of the disintegrants include starch, agar, gelatin powder, microcrystalline cellulose, calcium carbonate, sodium bicarbonate, calcium citrate, dextrin, pectin and carboxymethylcellulose calcium.

[0199] Examples of the lubricants include magnesium stearate, talc, polyethylene glycol, silica and hydrogenated vegetable oil.

[0200] Colorants used are those approved as additives to pharmaceuticals. Correctives used are cocoa powder, peppermint camphor, empassm, mentha oil, borneol, powdered cinnamon bark and the like.

[0201] Obviously, these tablets and granules may be sugar-coated or otherwise coated appropriately as necessary. Liquid preparations such as syrups and injectable preparations are manufactured by adding a pH adjuster, a solubilizer, a tonicity adjusting agent and the like, and as necessary, a solubilizing agent, a stabilizer and the like to the compound or a pharmacologically acceptable salt thereof according to the present invention and formulating them by a conventional method.

[0202] The method of manufacturing external preparations is not limited and they can be manufactured by conventional methods. Specifically, various raw materials commonly used for pharmaceuticals, quasi drugs, cosmetics and the like can be used as base materials for formulation. Specific examples of the base materials used include raw materials such as animal and vegetable oils, mineral oils, ester oils, waxes, higher alcohols, fatty acids, silicone oil, surfactants, phospholipids, alcohols, polyhydric alcohols, water-soluble polymers, clay minerals and purified water. Further, pH adjusters, antioxidants, chelators, preservatives and fungicides, colorants, flavors and the like may be added as necessary. The base materials for external preparations according to the present invention are not limited to these materials.

[0203] Ingredients such as ingredients having a differentiation-inducing effect, blood flow promoters, bactericides, anti-inflammatory agents, cell activators, vitamins, amino acids, humectants and keratolytic agents may also be blended as necessary. The aforementioned base materials are added in an amount corresponding to the concentration usually chosen for the manufacture of external preparations.

[0204] The mode of administration of the compounds or salts thereof, or hydrates of the compounds or salts according to the present invention is not particularly limited, and they may be orally or parenterally administered by methods commonly used. For example, they can be formulated into preparations such as tablets, powders, granules, capsules, syrups, troches, inhalations, suppositories, injections, ointments, ophthalmic ointments, ophthalmic preparations, nasal preparations, ear preparations, cataplasms and lotions and administered.

[0205] The dosage of the medicine according to the present invention can be appropriately selected depending on the severity of the symptom, the age, the sex, the body weight, the mode of administration, the type of the salt, the specific type of the disease, and the like.

[0206] Although the dosage significantly varies according to the type of the disease and the severity of the symptom of the patient, the age of the patient, the sex difference and the difference in sensitivity to drugs between the patients, and the like, the dosage is usually about 0.03 to 1000 mg, preferably 0.1 to 500 mg and more preferably 0.1 to 100 mg per day for adults and is administered divided into one to several doses a day. For injections, the dosage is usually about 1 μ g/kg to 3000 μ g/kg, preferably about 3 μ g/kg to 1000 μ g/kg.

[0207] In the manufacture of the compounds of the present invention represented by the above formula (1), raw material compounds and various reagents may form salts, hydrates or solvates, all vary according to the starting material, the solvent used, and the like, and are not particularly limited insofar as they do not inhibit the reaction.

[0208] The solvent used also varies according to the starting material, the reagent and the like, and is not particularly limited insofar as it does not inhibit the reaction and dissolves the starting material to a certain extent, obviously.

[0209] Various isomers (e.g., geometric isomers, optical isomers based on asymmetric carbons, rotamers, stereoisomers and tautomers) can be purified and isolated using common separation means, e.g., recrystallization, diastereomeric salt methods, enzymatic resolution methods and various chromatography methods (e.g., thin-layer chromatography, column chromatography, high performance liquid chromatography and gas chromatography).

[0210] The compounds according to the present invention obtained as free forms can be converted to salts that may be formed by the compounds or to hydrates of the compounds according to conventional methods. The compounds according to the present invention obtained as salts or hydrates of the compounds can also be converted to free forms of the compounds according to conventional methods.

[0211] The compounds according to the present invention can be isolated and purified by applying common chemical operations such as extraction, concentration, evaporation, crystallization, filtration, recrystallization and various chromatography methods.

[0212] All prior art documents cited herein are hereby incorporated by reference.


[0213] General manufacturing methods for the compounds of the present invention and examples will be shown below.

General synthesis methods

[0214] The compounds of the present invention can be synthesized by various methods, some of which will be described with reference to the following schemes. The schemes are illustrative and the present invention is not limited only by the chemical reactions and conditions explicitly indicated. Although some substituents are excluded in the following schemes for the sake of clarity, such exclusion is not intended to limit the disclosure of the schemes. Representative compounds of the present invention can be synthesized using appropriate intermediates, known compounds, and reagents. R_1 , R_2 , R_{33} , R_{34} , W , X , Y , m and n in the formulas in the following general synthesis methods are as defined for R_1 , R_2 , R_{33} , R_{34} , W , X , Y , m and n in the compounds represented by the above general formula (1) (compounds represented by the formula I in the following general synthesis methods).

[0215] The compounds of the general formula (1) according to the present invention can be synthesized by the manufacturing methods shown below.


Scheme 1 (Method A)



[0216] Scheme 1 (Method A) is a method of reacting a spiro-amine derivative (1) with various sulfonyl chlorides (2) in an appropriate solvent such as dichloromethane or tetrahydrofuran in the presence of an appropriate base such as triethylamine or pyridine. The reaction temperature is 0°C to room temperature, for example, and the reaction time is 0.5 to 24 hours. The resulting sulfonamide derivative (formula I) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0217] The spiro-amine derivative (1) shown in Scheme 1 can be synthesized from an acylamino-nitrile derivative (3) or acylamino-amide derivative (4). Scheme 2 shows a method of synthesizing the spiro-amine derivative (1).

Scheme 2



(In the scheme, $Y = \text{O}$.)


[0218] Step 2 is a method of cyclizing an acylamino-nitrile derivative (3) in an appropriate solvent such as an aqueous ethanol solution or an aqueous dimethyl sulfoxide solution in the presence of an aqueous sodium hydroxide solution and aqueous hydrogen peroxide solution. The reaction temperature is reflux temperature, for example, and the reaction time is 1 to 24 hours. The resulting cyclized derivative (5) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0219] The cyclized derivative (5) can also be synthesized by two-step reaction (Step 3, Step 4). Step 3 is a method of converting the nitrile group to an amido group under basic hydrolysis conditions in the presence of hydrogen peroxide. (This reaction can be performed with reference to Chemistry-A European Journal (2002), 8(2), 439-450, for example.) Step 4 is a method of cyclizing an acylamino-amide derivative (4) in an appropriate solvent such as ethanol, tert-butanol or dimethyl sulfoxide in the presence of an appropriate base such as an aqueous sodium hydroxide solution or potassium t-butoxide. The reaction temperature is room temperature to reflux temperature, for example, and the reaction time is 1 to 24 hours. The resulting cyclized derivative (5) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0220] Step 5 is a reaction of deprotecting the t-butoxycarbonyl group with an appropriate acid such as trifluoroacetic acid or hydrochloric acid in an appropriate solvent such as dichloromethane, dioxane or methanol. (This reaction can be performed with reference to Protective Groups in Organic Synthesis, Wiley-Interscience, for example.)

[0221] The acylamino-nitrile derivative (3) or acylamino-amide derivative (4) shown in Scheme 2 can be synthesized from an amino-nitrile derivative (8a) or amino-amide derivative (8b). Scheme 3 shows a method of synthesizing the acylamino-nitrile derivative (3) or acylamino-amide derivative (4).

Scheme 3




(In the scheme, Y = O.)

[0222] Step 6 is a method of reacting an acid chloride derivative (6) with an amino-nitrile derivative (8a) or amino-amide derivative (8b), respectively, in an appropriate solvent such as dichloromethane or tetrahydrofuran in the presence of an appropriate base such as triethylamine or pyridine. The reaction temperature is 0°C to room temperature, for example, and the reaction time is 0.5 to 24 hours. The resulting acylamino-nitrile derivative (3) or acylamino-amide derivative (4) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography. The acid chloride derivative (6) used for the reaction can be purchased or can be synthesized from a carboxylic acid derivative (7) by the method described in March, Advanced Organic Chemistry, 5th Edition, John Wiley and Sons, New York, P523-P524, for example.

[0223] Step 7 is a method of reacting a carboxylic acid derivative (7) with amino-nitrile (8a) or amino-amide (8b). Examples of the coupling reagent include N,N'-dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC), O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU) and 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride n-hydrate (DMT-MM). Examples of the base include triethylamine or N,N-diisopropylethylamine. If necessary, 4-(dimethylamino)pyridine (DMAP) may be used as a catalyst. Examples of the appropriate solvent include dichloromethane or N,N-dimethylformamide. Examples of the appropriate solvent used in the case of DMT-MM include methanol, ethanol and acetonitrile. The reaction temperature is 0°C to room temperature, for example, and the reaction time is 0.5 to 24 hours. The resulting acylamino-nitrile derivative (3) or acylamino-amide derivative (4) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0224] The sulfonyl chloride derivative (2) shown in Scheme 1 can be purchased or can be synthesized as shown in Scheme 4a and Scheme 4b.

Scheme 6




(In the scheme, Y = O.)

[0230] Step 1 is a method of reacting a keto-amine derivative (17) with a sulfonyl chloride derivative (2). Step 13 is a Strecker synthesis of converting a ketone derivative (18) to an amino-nitrile derivative (19). Specifically, this is a method of reacting a ketone derivative (18) with sodium cyanide or potassium cyanide and ammonium chloride or ammonium acetate in an appropriate solvent such as methanol, ethanol or tetrahydrofuran in the presence/absence of water. The reaction temperature is room temperature to 80°C, for example, and the reaction time is 2 to 72 hours. The resulting amino-nitrile derivative (19) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0231] The cyano-amide derivative (15) can be synthesized by the same method as in Step 6 or Step 7 in Scheme 3. Step 3 is a method of synthesizing an amido-amide compound (16) by hydrolysis of the cyano-amide derivative (15).

[0232] The spiro-amine derivative of the formula I, in particular, the aryl-ethenylsulfonamide derivative of the formula II (o = 0) and the aryl-propenylsulfonamide derivative of the formula II (o = 1), can be synthesized by a Heck reaction of an olefinated sulfonamide derivative (20) with an aryl halide (21) in Scheme 7.

Scheme 7 (Method C)




(In the scheme, Y = O.)

[0233] Step 14 is a method of synthesizing an arylethenylsulfonamide derivative (formula II) by coupling an olefinated sulfonamide derivative (20) with an aryl halide derivative (21) in an appropriate solvent such as N,N-dimethylacetamide (DMA), N,N-dimethylformamide (DMF) or 1,4-dioxane in the presence of a palladium catalyst such as palladium(II) acetate (Pd(OAc)₂) or tetrakis(triphenylphosphine)palladium(0) (Pd(PPh₃)₄), in the presence or absence of a phosphine ligand such as triphenylphosphine (PPh₃) or tri-o-tolylphosphine ((o-tol)₃P) and in the presence of an appropriate base such as triethylamine, respectively, in an N₂ atmosphere. The reaction temperature is 90°C to reflux temperature. This reaction can be performed under microwave irradiation. The resulting arylethenylsulfonamide derivative (formula II) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0234] The spiro-amine derivative (formula I), in particular, the ethenylsulfonamide derivative (formula III), can also be synthesized by coupling a Horner-Wadsworth-Emmons reagent with an aldehyde derivative (24) as shown in Scheme 8.

Scheme 8 (Method D)




(In the scheme, Y = O.)

[0235] Step 15 is a method of synthesizing a Horner-Wadsworth-Emmons reagent (23) by coupling a methanesulfonamide derivative (22) with diethyl chlorophosphate in an appropriate solvent such as tetrahydrofuran or diethyl ether in the presence of a base such as lithium hexamethyldisilazide (LHMDS) or lithium diisopropylamide (LDA). The reaction is performed at -78°C to room temperature for 1 to 24 hours in an N_2 atmosphere. The resulting Horner-Wadsworth-Emmons reagent (23) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography. This reaction can be performed with reference to Tetrahedron 2001, 57(37), 7899-7907, for example.

[0236] Step 16 is a method of synthesizing an ethenylsulfonamide derivative (formula III) by reacting the Horner-Wadsworth-Emmons reagent (23) with an aldehyde derivative (24) under Horner-Wadsworth-Emmons reaction conditions. This reaction can be performed with reference to Synlett 2005, 5, 834-838; Tetrahedron 2001, 57(37), 7899-7907, for example.

[0237] The spiro-amine derivative (formula I), in particular, the aryl-alkylsulfonamide derivative (formula IV), can be synthesized by reduction of an olefin of the formula II in Scheme 9.

Scheme 9 (Method E)




(In the scheme, Y = O.)

[0238] Step 17 is a method of hydrogenating an olefin of the formula II in an inert solvent such as methanol, ethanol, dimethylformamide or dimethylacetamide in the presence of a catalyst such as palladium carbon or palladium hydroxide carbon, respectively, under an H_2 atmosphere. The reaction temperature is room temperature to 80°C, and the reaction may be performed under pressure. The resulting aryl-alkylsulfonamide derivative (formula IV) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0239] The cyclized derivative (5) amide used in the above reaction can be converted to a thioamide (Step 18) and used for the reaction of Step 5 or Step 1. This reaction can be performed with reference to March, Advanced Organic Chemistry, 5th Edition, for example.


Scheme 10



(In the scheme, Y = O.)

[0240] The ketone derivative (18) shown in Scheme 6 can be derived from a keto-amine derivative (17) through an ethenesulfonamide derivative (26). It can also be derived from a ketal-amine derivative (27) through a ketal-ethenesulfonamide derivative (28).

Scheme 11




[0241] Step 19 is a method of reacting a keto-amine derivative (17) or ketal-amine derivative (27) with chloroethanesulfonyl chloride in an appropriate solvent such as dichloromethane in the presence of an appropriate base such as triethylamine. The reaction temperature is 0°C to 40°C, for example, and the reaction time is 0.1 to 1 hour. The resulting ethenesulfonamide derivative (26) or ketal-ethenesulfonamide derivative (28) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0242] The ketone derivative (18) can be synthesized from the ethenesulfonamide derivative (26) by the same method as in Step 14 in Scheme 7.

[0243] The ketone derivative (18) can also be synthesized by converting the ketal-ethenesulfonamide derivative (28) to a ketal-sulfonamide derivative (29) by the same method as in Step 14 in Scheme 7 and then deprotecting the ketal by the method of Step 20. Step 20 is a method of reacting the ketal-sulfonamide derivative (29) with an acid such as trifluoroacetic acid or hydrochloric acid in an appropriate solvent such as aqueous acetone or aqueous ethanol. The reaction temperature is 55°C to 80°C (boiling point of the solvent), for example, and the reaction time is 1 to 24 hours. The resulting ketone derivative (18) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0244] The olefinated sulfonamide derivative (20) shown in Scheme 7 can be derived from a spiro-amine derivative (1).

Scheme 12




[0245] Step 21 is a method of reacting a spiro-amine derivative (1) with a sulfonyl chloride reagent (e.g., chlo-

roethanesulfonyl chloride or 2-propene-1-sulfonyl chloride) in an appropriate solvent such as dichloromethane in the presence of an appropriate base such as triethylamine. The reaction temperature is 0°C to 40°C, for example, and the reaction time is 0.1 to 1 hour. The resulting olefinated sulfonamide derivative (20) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0246] The methanesulfonamide derivative (22) shown in Scheme 8 can be derived from a spiro-amine derivative (1).


Scheme 13



[0247] Step 22 is a method of reacting a spiro-amine derivative (1) with methanesulfonyl chloride in an appropriate solvent such as dichloromethane in the presence of an appropriate base such as triethylamine. The reaction temperature is 0°C to room temperature, for example, and the reaction time is 0.1 to 1 hour. The methanesulfonamide derivative (22) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0248] The compound of the general formula (1) wherein Y is a sulfur atom (formula VI) can be synthesized from a thioamide intermediate (25) in Scheme 10 by the reactions of Step 5-Step 1 as in the case of the compound wherein Y is an oxygen atom, for example. It can also be synthesized from an amide derivative wherein Y is an oxygen atom (formula V) by Step 18.


Scheme 14



[0249] Step 18 is a method of reacting an amide derivative (formula V) with a Lawesson's reagent in an appropriate solvent such as toluene. The reaction temperature is room temperature to the boiling point of the solvent, for example, and the reaction time is several hours to 24 hours. The thioamide derivative (formula VI) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography. This conversion reaction of carbonyl to thiocarbonyl can be performed with reference to March, Advanced Organic Chemistry, 5th Edition, for example.

[0250] The compound of the general formula (1) wherein Y is a nitrogen atom (formula VII, formula VIII) can be synthesized by converting the thioamido group of a thioamide intermediate (25) to an amidino group (step 23) to provide an amidino intermediate (31, 32) and then subjecting the intermediate to the reactions of Step 5 and subsequent Step 1 as in the case of the compound of the general formula (1) wherein Y is an oxygen atom, for example.


Scheme 15



[0251] Step 23 is a method of reacting a thioamide intermediate (25) with a primary amine or secondary amine in an appropriate solvent such as methanol. The reaction temperature is room temperature to the boiling point of the solvent, for example, and the reaction time is several hours to 24 hours. The amidino intermediate (31, 32) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0252] The derivative of the general formula (1) wherein X is a single bond and R₂ is optionally substituted aryl or heteroaryl (formula IX) can also be synthesized from a thiohydantoin derivative (33).


Scheme 16



[0253] Step 24 is a method of reacting a thiohydantoin derivative (33) with an optionally substituted arylboronic acid (34) in an appropriate solvent such as N-methylpyrrolidone in the presence of a copper catalyst such as CuTC or a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0). The reaction temperature is room temperature to the boiling point of the solvent, for example, and the reaction time is 0.5 to 24 hours. The substituted phenyl derivative (formula IX) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0254] The thiohydantoin derivative (33) can be synthesized from an amino-nitrile derivative (19) through Step 3 and Step 25.


Scheme 17



[0255] Step 3 is a method of synthesizing an amino-amide derivative (35) by hydrolyzing an amino-nitrile derivative (19). Step 25 is a reaction of converting the amino-amide derivative (35) to a thiohydantoin derivative (33). Step 25 is a method of reacting the amino-amide derivative (35) with a thiocarbonylating reagent such as di(2-pyridyl) thionocarbonate in an appropriate solvent such as tetrahydrofuran. The reaction temperature is 0°C to room temperature, for example, and the reaction time is 0.5 hour to several hours. The thiohydantoin derivative (33) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0256] The alkynyl derivative of the general formula (1) wherein W is acetylene and R₁ is optionally substituted aryl or heteroaryl (formula XI) can be synthesized through an acetophenone derivative (formula X).

Scheme 18



[0257] Step 26 is a method of condensation with an aryl ester (36) in an appropriate solvent such as tetrahydrofuran or diethyl ether in the presence of a base such as lithium hexamethyldisilazide (LHMDS) or lithium diisopropylamide (LDA) preferably with the addition of DMPU. The reaction is performed at -78°C to room temperature for 1 to 24 hours in an N₂ atmosphere. The resulting acetophenone derivative (formula X) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.


[0258] Step 27 is a method of synthesizing an alkynyl derivative (formula XI) by subjecting the acetophenone derivative (formula X) to dehydration reaction. Specifically, 1 to 10 equivalents of a dehydrating agent, preferably 2-chloro-1-methylpyridinium iodide, and an appropriate base, preferably triethylamine, are added and reacted in an appropriate solvent such as dichloromethane at 0°C to a temperature under heating. The resulting alkynyl derivative (formula XI) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0259] The substituted alkylene derivative of the general formula (1) wherein W is branched alkylene or haloalkylene (formula XII) can be synthesized as follows, for example.

[0260] The substituted alkylene derivative (formula XII) can be obtained by nucleophilic substitution reaction with a

ketal-sulfonamide derivative (37) as a raw material to introduce an alkyl group or a halogen atom onto the carbon adjacent to the sulfonyl group (Step 28), ketal deprotection reaction (Step 20) and the steps shown in Schemes 5 and 6.

Scheme 19




[0261] Step 28 is a reaction of a ketal-alkylenesulfonamide derivative (37) with an electrophilic reagent such as an alkyl halide or NFSI (N-fluorodibenzenesulfonimide) in an appropriate solvent such as tetrahydrofuran in the presence of a base such as n-butyllithium or lithium diisopropylamide. The reaction temperature is -78°C to room temperature, for example, and the reaction time is 0.5 hour to several hours. The ketal-sulfonamide derivative (38) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography. In this reaction, one or two substituents are introduced onto the carbon atom adjacent to the sulfonyl group, and the equivalents of the base are controlled according to need. In the introduction of two substituents, two substituents can be introduced all at once using an excess of a base; however, it is desirable to once obtain a compound having one substituent introduced therein by purification and then introduce the other substituent.

[0262] The guanidine derivative of the general formula (1) wherein X is a nitrogen atom (formula XIII) can be synthesized through a guanidine intermediate (41).

[0263] The guanidine intermediate (41) can be synthesized from a thiohydantoin derivative (39) through an isothiurea derivative (40). The guanidine derivative (formula XIII) can be synthesized from the guanidine intermediate (41) by the reactions of Step 5 and subsequent Step 1, for example.

Scheme 20




[0264] Step 29 is S-alkylation reaction of a thiohydantoin derivative (39). Specifically, this is a method of reacting a thiohydantoin derivative (39) with an alkyl halide reagent such as methyl iodide in an appropriate solvent such as methanol in the presence of a base such as sodium hydroxide. The reaction temperature is room temperature to the boiling point of the solvent, for example, and the reaction time is several hours to 24 hours. The isothiurea derivative (40) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0265] Step 30 is a reaction of converting isothiurea to guanidine. Specifically, this is a method of reacting the isothiurea derivative (40) with a substituted primary amine or substituted secondary amine in an appropriate solvent such as dimethylacetamide in the presence of an acid catalyst such as acetic acid. The reaction temperature is room temperature to the boiling point of the solvent, for example, and the reaction time is 0.5 hour to several hours. The

guanidine derivative (41) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0266] The derivative compound represented by the general formula (2), wherein Z includes alkenylene, can be synthesized from a diolefin derivative (formula XIV) by olefin metathesis reaction (Step 31).


Scheme 21



[0267] Step 31 is a method of cyclizing a diolefin derivative (formula XIV) using a Grubbs reagent in an appropriate solvent such as dichloroethane. The reaction temperature is room temperature to the boiling point of the solvent, for example, and the reaction time is several hours to 24 hours. The macrocyclic derivative (formula (2)) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0268] The derivative compound represented by the general formula (2), wherein Z includes an amido group, can also be synthesized from a derivative having a primary amine or secondary amine at one end of the compound and a carboxylic acid at the other end (formula XV) by amidation reaction (Step 7).


Scheme 22



[0269] The compound of the formula (2) is synthesized by cyclizing a derivative having an amine and a carboxylic acid at the ends (formula XV) using the above-described method of Step 7.

[0270] The derivative compound represented by the general formula (2), wherein Z includes triazole, can also be synthesized from a derivative (42) having an alkyne at one end of the compound and an azide at the other end using click chemistry (Step 32).

Scheme 23



[0271] Step 32 is a method of reacting an azide and an alkyne in a derivative (42) having the alkyne and the azide at the ends in an appropriate solvent such as acetonitrile or tetrahydrofuran in the presence of a copper catalyst such as CuI with the addition of a base such as diisopropylethylamine or 2,6-lutidine and ascorbic acid if necessary. The reaction temperature is room temperature to the boiling point of the solvent, for example, and the reaction time is 1 to 24 hours. The derivative represented by the general formula (2) is isolated by a common technique and, if necessary, may be purified by crystallization or chromatography.

[0272] Derivatives can be synthesized from compounds obtained in Schemes 21, 22 and 23 by olefin hydrogenation reaction, olefin oxidation reaction, sulfur atom oxidation reaction, deprotection reaction of various protecting groups and the like as necessary.

[0273] Compounds containing functional groups such as alkenyl, amine, carboxylic acid, alkynyl and azido functional groups at the ends, the raw materials used in Schemes 21, 22 and 23 and the like for synthesizing the general formula (2), may be synthesized by previously introducing these functional groups into a carboxylic acid derivative (7), a bromide derivative (9), a styrene derivative (11), an aryl halide derivative (21), an aldehyde derivative (24), a substituted phenylboronic acid (34) and the like and subjecting them to the same method as the method of manufacturing the general formula (1), or may be synthesized by synthesizing intermediates of the general formula (1), in which functional groups such as carboxylic acid and phenol functional groups are introduced into R₁ or R₂, and introducing the functional groups from these intermediates into these derivatives by an appropriate reaction.

[0274] If the functional groups introduced at the ends are previously introduced into a carboxylic acid derivative (7), a bromide derivative (9), a styrene derivative (11), an aryl halide derivative (21), an aldehyde derivative (24), a substituted phenylboronic acid (34) and the like, then these functional groups are protected and deprotected by an appropriate method as necessary in a process of synthesis by the same method as the method of manufacturing the general formula (1).

[0275] If the functional groups introduced at the ends are introduced into these derivatives by synthesizing intermediates of the general formula (1), in which a functional group such as carboxylic acid and phenol functional groups is introduced into R₁ or R₂, and introducing the functional groups from these intermediates by an appropriate reaction, then amidation, Mitsunobu reaction or the like is preferred as such an appropriate reaction. The functional groups are protected and deprotected by an appropriate method as necessary. For example, to introduce alkenyl or azido into R₁, amidation reaction is performed between a compound of the general formula (1), in which carboxylic acid is introduced into R₁, and an alkylamine with alkenyl or azido bonded thereto. To introduce alkenyl or alkynyl into R₂, Mitsunobu reaction is performed between a compound of the general formula (1), in which phenol is introduced into R₂, and an alkylamine with alkenyl, alkynyl or a protected amine bonded thereto. Raw materials for synthesizing the general formula (2) can be synthesized by combining these reactions of introducing functional groups into R₁ or R₂.

[0276] The compound (7), compound (9), compound (11), compound (21), compound (24) or compound (34) used in the above reactions can be synthesized from known compounds using appropriate reagents and reactions. For example, an amino group, if present in R₁ or R₂, may be alkylated, acylated, carbamated, converted to ureas or sulfonamidated from known compounds. Carboxylic acids or esters, if present, may be amidated under general conditions. Sulfonyl chlorides, if present, may be condensed with amines and sulfonamidated. Alcohols, if present, may be etherified or carbamated. Aryl halides, if present, may be coupled with arylboric acids or aryl boronates under general Suzuki conditions. Olefins, if present, may be reduced or converted to diols. Thioether groups, if present, may be oxidized to sulfoxides or sulfones. If ketones or carbonyl groups are present, the carbon chain may be extended by Wittig reaction, Horner-Wadsworth-Emmons reaction, aldol reaction or the like. In the introduction of fluorine atoms, reagents containing fluorine atoms may be introduced by these reactions, or aldehydes, ketones or carboxylic acids may be reacted with diethylaminosulfur trifluoride, for example.

[0277] The techniques for introducing these groups can be performed with reference to March, Advanced Organic Chemistry, 5th Edition, John Wiley and Sons, New York; J. Med. Chem., 2005, 48, 6066-6083; Organic Syntheses (1951), 31, 8-11; Bioorg. Med. Chem. Lett., 2003, 13, 837-840; Chem. Rev. 2002, 102, 1359; J. Organomet. Chem. 1999, 576, 147; Chem. Rev. 1995, 95, 2457, for example. These groups may be protected with protecting groups under general conditions, if necessary. This can be performed with reference to Protective Groups in Organic Synthesis, Wiley-Interscience, for example.

[0278] Some of the compounds of the present invention are useful not only as compounds having a PTH-like effect but also as intermediates for the synthesis of additional compounds of the present invention. For example, amines may be alkylated, acylated, carbamated, converted to ureas, sulfonamidated or sulfamidated under general conditions. Carboxylic acid and ester moieties may be converted to amides under general conditions. Amido groups may be converted to thioamido groups. Olefins may be reduced or converted to diols. Thioether groups, if present, may be oxidized to sulfoxides or sulfones. The techniques for introducing these groups can be performed with reference to March, Advanced Organic Chemistry, 5th Edition, John Wiley and Sons, New York; J. Med. Chem., 2005, 48, 6066-6083; Organic Syntheses (1951), 31, 8-11; Bioorg. Med. Chem. Lett., 2003, 13, 837-840, for example. Protecting groups may be deprotected under general conditions. This can be performed with reference to Protective Groups in Organic Synthesis,

Wiley-Interscience, for example. Aryl halides may be coupled with arylboric acids or aryl boronates under general Suzuki conditions. This can be performed with reference to Chem. Rev. 2002, 102, 1359; J. Organomet. Chem. 1999, 576, 147; Chem. Rev. 1995, 95, 2457, for example.

5 Examples

[0279] The content of the present invention will be described in more detail by the following examples and test example; however, the present invention is not limited to the content of the examples and test example. All starting materials and reagents were obtained from commercial suppliers or synthesized using known methods. ¹H-NMR spectra were measured using EX270 (manufactured by JEOL), Mercury300 (manufactured by Varian), ARX-3000 (manufactured by Bruker), ECP-400 (manufactured by JEOL) or 400-MR (manufactured by Varian) with or without Me₄Si as the internal standard (s = singlet, d = doublet, t = triplet, brs = broad singlet, m = multiplet). Mass spectrometry measurement was performed using a mass spectrometer, LCQ Classic (manufactured by Thermo Electron), ZQ2000 (manufactured by Waters), 3100 (manufactured by Waters), ZMD4000 (manufactured by Waters), SQD (manufactured by Waters) or 2020 (manufactured by Shimadzu). Microwave irradiation was performed using InitiatorTM (manufactured by Biotage). In LCMS and HPLC, measurement of the retention time and mass spectrometry were performed by the following apparatuses and analysis conditions.

20

25

30

35

40

45

50

55

[Table 1]

LCMS, HPLC condition No.	Apparatus	Column (ID. × length) (mm)	Mobile phase	Gradient (A/B)	Flow rate	Column temperature (°C)	Wavelength
LCMS A-1	Agilent 1100/ LCQ Classic	Cadenza CD-C18 3µm (3.0 × 30)	A) 0.05% TFA, H ₂ O B) 0.05% TFA, MeCN	95/5 => 0/100 (3.5 min) 0/100 (1 min)	1.5 mL/min	35	210-400 nm PDA total
LCMS A-2	Agilent 1100/ LCQ Classic	Cadenza CD-C18 3µm (3.0×30)	A) 0.05% TFA, H ₂ O B) 0.05% TFA, MeCN	95/5 => 0/100 (9.5 min) 0/100 (2.5 min)	1.0 mL/min	35	210-400 nm PDA total
LCMS-B-1	Alliance 2795 HT/ 996 PDA/ ZMD4000	Cadenza CD-C18 3µm (3.0 × 30)	A) 0.05% TFA, H ₂ O B) 0.05% TFA, MeCN	95/5 => 0/100 (3.5 min) 0/100 (1 min)	1.5 mL/min	35	210-400 nm PDA total
LCMS-C-1	2525 BGM/ 2996 PDA/ ZQ2000	Chromolith Flash RP-18e (46 × 25)	A) 10mM AcONH ₄ H ₂ O B) MeOH	95/5 => 0/100 (3 min) 0/100 (2 min)	2.0 mL/min	Room temperature	210-400 nm PDA total
LCMS-C-2	2525 BGM/ 2996 PDA/ ZQ2000	Chromolith Flash RP-18e (46 × 25)	10mM AcONH ₄ , H ₂ O B) MeOH	95/5=> 0/100 (3min) 0/100 (2 min)	2.0 mL/min	Room temperature	210-400 nm PDA total
LCMS-C-3	2525 BGM/ 2996 PDA/ ZQ2000	Chromolith Flash RP-18e (46 × 25)	A) 10mM AcONH ₄ H ₂ O B) MeOH	50/50 => 0/100 (3 min) 0/100 (2 min)	2.0 mL/min	Room temperature	210-400 nm PDA total

(continued)

LCMS, HPLC condition No.	Apparatus	Column (ID. × length) (mm)	Mobile phase	Gradient (A/B)	Flow rate	Column temperature (°C)	Wavelength
LCMS-D-1	2545 BGM/ 2996 PDA/ 3100	Sunfire™ C18 5µm (46 × 50)	A) 0.50% TFA, H ₂ O B) 0.05% TFA, MeCN	90/10 => 10/90 (5 min)	4.0 mL/min	25	210-400 nm PDA total
LCMS-E-1	Agilent 1100	Waters X-Bridge C18 Sum (2.1 × 50)	A) 0.01% NH ₃ , H ₂ O B) 0.01% NH ₃ MeCN	95/5 => 5/95 (5 min)	1.2 mL/min	40	190-400 nm PDA total
LCMS-E-2	Alliance 2795/ ZQ2000	Waters X- Bridge C18 Sum (2.1 × 50)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	95/5=> 35/65 (5 min)	1.2 mL/min	40	190-400nm PDA total
LCMS-E-3	Alliance 2795/ ZQ2000	Waters X- Bridge C18 5µm (2.1 × 50)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	95/5 (0 min) => 95/5 (0.5 min) => 5/95 (5 min)	1.2 mL/min	5.5	190-400 nm PDA total
LCMS-E-4	Alliance 2795/ ZQ2000	Waters X- Bridge C18 5µm (2.1 × 50)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	95/5 (0 min) => 95/5 (0.5 min) => 5/95 (5 min)	1.2 mL/min	45	190-400 nm PDA total
LCMS-E-5	Alliance 2795/ ZQ2000	Waters X-Bridge C18 5µm (2.1 × 50)	A) 0.01% NH ₃ H ₂ O B) 0 0.1%NaH ₃ MeCN	95/5(0 min) => 95/5 (0.5 min) 35/65 (5 min) => 5/95 (5.5 min)	1.2 mL/min	45	190-400nm PDA total

(continued)

LCMS, HPLC condition No.	Apparatus	Column (ID. × length) (mm)	Mobile phase	Gradient (A/B)	Flow rate	Column temperature (°C)	Wavelength
LCMS-E-6	Alliance 2795/ ZQ2000	Waters X- Bridge C18 5µm (2.1 × 50)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	95/5=> 5/95 (5 min)	1.2 mL/min	40	190-400 nm PDA total
LCMS-E-7	Alliance 2795/ ZQ2000	Waters X- Bridge C18 3.5µm (2.1 × 50)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	95/5(0 min) => 95/5 (3 min)	1.2 mL/min	45	190-400 nm PDA total
LCMS E-8	Alliance 2795/ ZQ2000	Waters X- badge C18 5µm (2.1 × 50)	A) 0.1% 7FA, H ₂ O B) 0.1% TFA, MeCN	95/5 (0 min) => 95/5 (0.5 min) => 35/65 (5 min) => 5/95 (5.5 min)	1.2 mL/min	45	190-400 nm PDA total
LCMS-F-1	Acquity/ SQD	Ascentis Express C18 (2.1 × 50)	A) 10mM AcONH ₄ H ₂ O B) MeOH	95/5 => 0/100 (1 min) 0/100 (0.4 min)	1.0 mL/min	35	210-400 nm PDA total
LCMS-F-2	Acquity/ SQD	Ascentis Express C18 (2.1 × 50)	A) 0.1% HCO ₂ H, H ₂ O B) 0.1% HCO ₂ H, MeCN	95/5 => 0/100 (1 min) 0/100 (0.4 min)	1.0 mL/min	35	210-400 nm PDA total
LCMS-G-1	UFLC XR/2020	Acquity (2.1 × 50)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	95/5 => 0/100 (1.5 min) 0/100 (0.5 min)	1.0 mL/min	35	305nm, bandwidth 95nm


(continued)

LCMS, HPLC condition No.	Apparatus	Column (ID. × length) (mm)	Mobile phase	Gradient (A/B)	Flow rate	Column temperature (°C)	Wavelength
HPLC-A-1	LC-2010A (SHIMAZU)	YMC-ODSA (6.0 × 150)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	90/20 => 10/80 (40 min)	1.0 mL/min	25	UV 254, 225 nm
HPLC-A-2	LC-2010A (SHIMAZU)	YMC-ODSA (6.0 × 150)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	90/30 => 10/70 (40 min)	1.0 mL/min	25	UV 254, 225 nm
HPLC-A-3	LC-2010A (SHIMAZU)	YMC-ODSA (6.0 × 150)	A) 0.1% TFA, H ₂ O B) 0.1% TFA, MeCN	90/10=> 10/90 (25 min)	1.0 mL/min	2.5	UV 254, 225 nm

Example 1


8-(3-Chloro-benzenesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1)

[0280]

(Reaction 1-1)


[0281] Potassium carbonate (13.96 g, 101.1 mmol) and 3-chlorobenzenesulfonyl chloride were continuously added to a two-phase solution of 4-piperidone hydrochloride hydrate (6.06 g, 39.48 mmol) in chloroform (47.4 mL) and - water (47.4 mL), and the mixture was stirred at room temperature. A saturated aqueous sodium bicarbonate solution was added, and the organic layer and the aqueous layer were separated. The aqueous layer was then further extracted with dichloromethane. The organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting solid was washed with n-hexane and then collected by filtration and dried under reduced pressure to give 1-(3-chloro-benzenesulfonyl)-piperidin-4-one as a white solid (10.5 g, 97%).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 2.55 (4H, t, $J = 6.3$ Hz), 3.42 (4H, t, $J = 6.0$ Hz), 7.48 (1H, t, $J = 8.0$ Hz), 7.58 (1H, dt, $J = 8.0, 1.7$ Hz), 7.67 (1H, dt, $J = 7.7, 1.7$ Hz), 7.77 (1H, t, $J = 1.9$ Hz).

(Reaction 1-2)

[0282] Ammonium chloride (790 mg, 14.77 mmol) and a 28% aqueous ammonia solution (2.2 mL) were added to a solution of 1-(3-chloro-benzenesulfonyl)-piperidin-4-one (3.11 g, 11.36 mmol) in dimethylformamide (15 mL), and the mixture was stirred at room temperature for one hour. Thereafter, sodium cyanide (724 mg, 14.77 mmol) was added, and the mixture was further stirred for 17 hours and then quenched with a saturated aqueous sodium carbonate solution. The organic layer and the aqueous layer were separated, and the aqueous layer was then further extracted with ethyl acetate:n-hexane (4:1). The organic layers were combined, washed with water ($\times 4$), and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (n-hexane:ethyl acetate = 1:3) to give 4-amino-1-(3-chloro-benzenesulfonyl)-piperidine-4-carbonitrile as a white solid (2.41 g, 71%).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 1.75 (2H, s), 1.80-1.90 (2H, m), 2.11-2.14 (2H, m), 2.87-2.96 (2H, m), 3.54-3.62 (2H, m), 7.47 (1H, t, $J = 8.1$ Hz), 7.58-7.66 (2H, m), 7.75 (1H, t, $J = 1.8$ Hz).


(Reaction 1-3)

[0283] A solution of cyclohexanecarbonyl chloride (118 μL , 0.880 mmol) in chloroform (0.25 mL) was added to a mixed

solution of 4-amino-1-(3-chloro-benzenesulfonyl)-piperidine-4-carbonitrile (120 mg, 0.400 mmol) in chloroform (1.25 mL) and a saturated aqueous sodium carbonate solution (1.25 mL), and the mixture was vigorously stirred at room temperature for 16 hours. Cyclohexanecarbonyl chloride (51 μ L) was further added and the mixture was stirred for 2.5 hours. The organic layer and the aqueous layer were then separated, and the aqueous layer was further extracted with chloroform. The organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The residue was washed with n-hexane to give cyclohexanecarboxylic [1-(3-chloro-benzenesulfonyl)-4-cyano-piperidin-4-yl]-amide as a white solid. This was used in the next step without further purification.

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 1.20-1.96 (14H, m), 2.04-2.14 (1H, m), 2.55 (2H, brd, $J = 13$ Hz), 2.76-2.87 (2H, m), 5.58 (1H, s), 7.51 (1H, t, $J = 7.9$ Hz), 7.60-7.66 (2H, m), 7.75 (1H, t, $J = 1.8$ Hz). MS (ESI) $m/z = 410$ ($\text{M}+\text{H}$) $^+$.

(Reaction 1-4)



[0284] A 6 M aqueous sodium hydroxide solution (0.74 mL) and a 30% aqueous hydrogen peroxide solution (0.25 mL) were added to a solution of cyclohexanecarboxylic [1-(3-chloro-benzenesulfonyl)-4-cyano-piperidin-4-yl]-amide (100 mg, 0.244 mmol) in ethanol (1.60 mL), and the mixture was heated under reflux for 4.5 hours. The reaction mixture was cooled to room temperature and water was then added, followed by concentration under reduced pressure. The residue was neutralized with a saturated aqueous ammonium chloride solution, followed by extraction with dichloromethane. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane:methanol = 90:10) to give 8-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a white solid (52 mg, 52%).











$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 1.13-1.40 (7H, m), 1.54-1.75 (7H, m), 2.22-2.30 (1H, m), 2.72-2.80 (2H, m), 3.53-3.59 (2H, m), 7.67-7.85 (4H, m), 10.80 (1H, s). MS (ESI) $m/z = 410$ ($\text{M}+\text{H}$) $^+$.

[0285] The example compounds shown below were synthesized by operations similar to those in Example 1 using appropriate reagents and starting materials.

[Table 2]











Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
2		LCMS-A-1	1.92	405 ($\text{M}+\text{H}$) $^+$
3		LCMS-A-1	1.9	370 ($\text{M}+\text{H}$) $^+$
4		LCMS-A-1	1.87	370 ($\text{M}+\text{H}$) $^+$
5		LCMS-A-1	2.32	434 ($\text{M}+\text{H}$) $^+$
6		LCMS-A-1	2.65	438 ($\text{M}+\text{H}$) $^+$

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
7		LCMS-A-1	2.22	418 (M+H)+
8		LCMS-A-1	1.79	342 (M+H)+
10		LCMS-C-1	2.82	480 (M+H)+
11		LCMS-C-1	2.66	418 (M+M)+
12		LCMS-C-1	2.89	472 (M+H)+
13		LCMS-A-1	2.82	488 (M+H)+
14		LCMS-C-1	2.86	424 (M+H)+
15		LCMS-A-1	1.84	453 (M+H)+
16		LCMS-A-1	1.97	384 (M+H)+
17		LCMS-C-1	2.81	478 (M+H)+

EP 2 433 940 B9

(continued)


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
18		LCMS-C-1	2.8	511 (M+H)+
19		LCMS-A-1	2.22	424 (M+H)+
20		LCMS-C-1	2.78	424 (M+H)+
21		LCMS-C-1	3.1	452 (M+H)+
22		LCMS-C-1	2.76	422 (M+H)+
23		LCMS-C-1	2.74	422 (M+H)+
24		LCMS-C-1	2.47	440 (M+H)+
25		LCMS-C-1	2.45	440 (M+H)+
26		LCMS-C-1	3.14	466 (M+H)+
27		LCMS-C-1	2.61	442 (M+H)+

(continued)


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
28		LCMS-C-1	2.81	424 (M+H)+
29		LCMS-C-1	2.65	432 (M+H)+
30		LCMS-A-1	2.4	438 (M+H)+
31		LCMS-A-1	2.79	472 (M+H)+
32		LCMS-C-1	2.3	482 (M+H)+
33		LCMS-C-1	3.08	486 (M+H)+
34		LCMS-C-1	2.87	482 (M+H)+

Example 2


8-(2-Naphthalen-1-yl-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 35)

[0286](Reaction 2-1)

[0287] 1-(2-Naphthalen-1-yl-ethanesulfonyl)-piperidin-4-one was synthesized by the procedure described in Reaction 1-1 of Example 1 using 2-naphthalen-1-yl-ethanesulfonyl chloride as a reagent.
MS (ESI) m/z = 318 (M+H)+.


(Reaction 2-2)

[0288] 4-Amino-1-(2-naphthalen-1-yl-ethanesulfonyl)-piperidine-4-carbonitrile was synthesized by operations similar to those in Reaction 1-2 of Example 1 using THF-CH₃CN as a solvent and using appropriate reagents and starting material. MS (ESI) *m/z* = 344 (M+H)⁺.

(Reaction 2-3)

[0289] 3-Trifluoromethyl-benzoyl chloride (57 μ L, 0.378 mmol) was added to a solution of 4-amino-1-(2-naphthalen-1-yl-ethanesulfonyl)-piperidine-4-carbonitrile (100 mg, 0.291 mmol) and Et₃N (61 μ L) in CH₂Cl₂ (3 mL). The reaction mixture was stirred at room temperature for four hours and then diluted with CH₂Cl₂, and the organic layer was washed with water. The organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The residue was used in the next step without further purification.

MS (ESI) *m/z* = 516 (M+H)⁺.

(Reaction 2-4)

[0290] 8-(2-Naphthalen-1-yl-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 1-4 of Example 1 using appropriate reagents and starting material. ¹H-NMR (400 MHz, CDCl₃) δ 1.65-1.75 (2H, m), 2.07-2.16 (2H, m), 3.32-3.38 (2H, m), 3.42-3.55 (2H, m), 3.63-3.70 (2H, m), 3.82-3.90 (2H, m), 7.40-7.47 (2H, m), 7.50-7.55 (1H, m), 7.56-7.65 (2H, m), 7.77-7.782 (2H, m), 7.90 (1H, d, *J* = 4.0 Hz), 8.02-8.10 (2H, m), 8.18 (1H, s); MS (ESI) *m/z* = 516 (M+H)⁺.

[0291] The example compounds shown below were synthesized by operations similar to those in Example 2 using appropriate reagents and starting materials.

[Table 3]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (<i>m/z</i>)
36		LCMS-A-1	2.24	428 (M+H) ⁺

(continued)


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
37		LCMS-A-1	2.55	522 (M+H)+

Example 3

3-[8-(2-Naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butyl ester (Compound 38)

[0292]**(Reaction 3-1)**

[0293] DCC (1.3 eq) and DMAP (5 mol%) were added to a mixture of 4-amino-1-(2-naphthalen-1-yl-ethanesulfonyl)-piperidine-4-carbonitrile and piperidine-1,3-dicarboxylic acid 1-tert-butyl ester (1.3 eq) in DMF, and the mixture was stirred at room temperature overnight. Ethyl acetate was added to the reaction solution, and the organic layer was then sequentially washed with 1 N NaOH, water and saturated brine. The organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The residue was used in the next step without further purification.

(Reaction 3-2)

[0294] 3-[8-(2-Naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 1-4 of Example 1 using appropriate reagents and starting material.


MS (ESI) m/z = 555 (M+H)+.

Example 4

8-(2-Naphthalen-1-yl-ethanesulfonyl)-2-[1-(2-naphthalen-1-yl-ethanesulfonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 39)

[0295]


(Reaction 4-1)



[0296] Trifluoroacetic acid (10 eq) was added dropwise to a solution of 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butyl ester in CH_2Cl_2 . The reaction mixture was stirred at room temperature overnight and then concentrated under reduced pressure to give 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-piperidin-3-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate as a pale yellow form (70%).

$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ 0.98 (3H, t), 1.15-1.22 (1H, m), 1.44-1.52 (2H, m), 1.56-1.67 (2H, m), 1.69-1.82 (2H, m), 1.86-2.03 (3H, m), 2.63-2.71 (1H, m), 2.79-2.85 (2H, m), 2.86-3.05 (3H, m), 3.49-3.65 (3H, m), 3.65-3.76 (2H, m), 7.44 (1H, t, $J = 7.83$ Hz), 7.50-7.55 (1H, m), 7.61 (1H, d, $J = 8.08$ Hz), 7.71 (1H, t, $J = 1.77$ Hz). MS (ESI) $m/z = 517$ ($\text{M}+\text{H}$) $^+$.

(Reaction 4-2)



[0297] 8-(2-Naphthalen-1-yl-ethanesulfonyl)-2-[1-(2-naphthalen-1-yl-ethanesulfonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 2-3 of Example 2 using appropriate reagents and starting material.


MS (ESI) $m/z = 673$ ($\text{M}+\text{H}$) $^+$.

[0298] The example compounds shown below were synthesized by operations similar to those in Example 3 and Example 4 using appropriate reagents and starting materials.

[Table 4]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
40		LCMS-E-2	3.76	517 ($\text{M}+\text{H}$) $^+$
41		LCMS-E-2	4.13	551 ($\text{M}+\text{H}$) $^+$
42		LCMS-E-2	4.11	565 ($\text{M}+\text{H}$) $^+$

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
43		LCMS-M-2	3.58	609 (M+H)+


Example 5

8-(4-Chloro-benzenesulfonyl)-2-(2,4-dichloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 44)

[0299](Reaction 5-1)


[0300] 4-Cyano-4-(2,4-dichloro-benzoylamino)-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 2-3 of Example 2 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.47 (9H, s), 1.87-1.97 (2H, m), 2.45-2.55 (2H, m), 3.32-3.43 (2H, m), 3.90-4.05 (2H, m), 6.48 (1H, brs), 7.38 (1H, dd, J = 8.4, 2.0 Hz), 7.45 (1H, d, J = 2.0 Hz), 7.78 (1H, d, J = 8.4 Hz).

(Reaction 5-2)


[0301] 2-(2,4-Dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 1-4 of Example 1 using appropriate reagents and starting material.

MS (ESI) m/z = 490 (M+H)+.

(Reaction 5-3)

[0302] 4 N HCl-dioxane (20 ml, 80 mmol) was added to a solution of 2-(2,4-dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (3.11 g, 7.81 mmol) in CH_2Cl_2 (60 mL), and the mixture was stirred at room temperature for four hours. The reaction mixture was diluted with CH_2Cl_2 -hexane, and the precipitated solid was then filtered. The resulting solid was washed with ethyl acetate and then dried under reduced pressure to give 2-(2,4-dichloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride (3.18 g) as a colorless solid. MS (ESI) m/z = 298 (M+H)⁺.

(Reaction 5-4)



[0303] Triethylamine (88 μl , 0.632 mmol) and 4-chlorobenzenesulfonyl chloride (70 mg, 0.332 mmol) were added to a mixed solution of 2-(2,4-dichloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride (100 mg, 0.299 mmol) in dichloromethane (3 ml). The reaction solution was stirred at room temperature for 16 hours and then diluted with dichloromethane, and the organic layer was washed with water. The organic layer was dried over sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate:hexane) to give 8-(4-chloro-benzenesulfonyl)-2-(2,4-dichloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (70.8 mg, 96%).

¹H-NMR (400 MHz, DMSO- d_6) δ 1.57-1.67 (2H, m), 1.85-1.95 (2H, m), 2.74-2.83 (2H, m), 3.64-3.72 (2H, m), 7.57-7.60 (1H, m), 7.61-7.65 (1H, m), 7.75-7.79 (2H, m), 7.81-7.86 (3H, m), 11.5 (1H, brs). MS (ESI) m/z = 472 (M+H)⁺.

[0304] The example compounds shown below were synthesized by operations similar to those in Example 5 using appropriate reagents and starting materials.


[Table 5]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
45		LCMS-C-3	4.59	468, 470 (M+H) ⁺
46		LCMS-A-1	2.45	418 (M+H) ⁺
47		LCMS-A-1	2.94	516 (M+H) ⁺

Example 6

2-(2,4-Dichloro-phenyl)-8-(quinoline-8-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 48)

[0305]

(Reaction 6-1)

[0306] 2-(2,4-Dichloro-phenyl)-8-(quinoline-8-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 4-2 of Example 4 using appropriate reagents and starting material and using pyridine as a base and solvent.

MS (ESI) m/z = 490 (M+H)+.

[0307] The example compounds shown below were synthesized by operations similar to those in Example 6 using appropriate reagents and starting materials.


[Table 6]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
49		LCMS-C-1	9.77	490 (M+H)+
50		LCMS-C-1	9.57	472 (M+H)+

Example 7


2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-benzoic acid methyl ester (Compound 51)

[0308]

(Reaction 7-1)

[0309] 2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 5-1 and Reaction 5-2 of Example 5 using appropriate reagents and starting material. MS (ESI) m/z = 358 (M+Na)+.


(Reaction 7-2)



10 [0310] Trifluoroacetic acid (20 ml) was added to a solution of 2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (4.43 g, 13.2 mmol) in dichloromethane, and the mixture was stirred at room temperature for five hours. The reaction mixture was concentrated under reduced pressure, and the residue was then triturated with CH₂Cl₂. The resulting solid was collected by filtration and dried under reduced pressure to give 2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate (5.66 g, 92%).

MS (ESI) m/z = 236 (M+H)+.

15 (Reaction 7-3)



25 [0311] 2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-benzoic acid methyl ester was synthesized by operations similar to those in Reaction 5-4 of Example 5 using appropriate reagents and starting material.











MS (ESI) m/z = 434 (M+H)+.

30 [0312] The example compounds shown below were synthesized by operations similar to those in Example 7 using appropriate reagents and starting materials.

[Table 7]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
52		LCMS-E-1	3.584	458 (M+H)+
53		LCMS-E-1	3.587	433 (M+H)+
54		LCMS-E-1	3.727	433 (M+H)+
55		LCMS-E-1	3.585	416 (M+H)+

(continued)


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
56		LCMS-E-1	3.143	382 (M+H)+
57		LCMS-E-1	3.604	427 (M+H)+
58		LCMS-E-2	1.22	411 (M+H)+
59		LCMS-E-1	3.065	396 (M+H)+
60		LCMS-E-1	3.141	434 (M+H)+
61		LCMS-E-1	3.782	454 (M+H)+
62		LCMS-C-1	2.35	440 (M+H)+
63		LCMS-C-1	2.64	454 (M+H)+
64		LCMS-A-1	2.00	470 (M+H)+
65		LCMS-C-2	2.13	461 (M+H)+

Example 8

8-(5-Chloro-thiophene-2-sulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 66)

[0313]

(Reaction 8-1)



[0314] 8-(5-Chloro-thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Example 5 using appropriate reagents and starting material.

MS (ESI) m/z = 478 (M+H)+.

[0315] The example compounds shown below were synthesized by operations similar to those in Example 8 using appropriate reagents and starting materials.


[Table 8]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
67		LCMS-C-1	2.90	488 (M+H)+
68		LCMS-C-1	2.92	494 (M+H)+
69		LCMS-C-2	2.38	523 (M+H)+

Example 9

8-(3-Chloro-benzenesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.6]undec-1-en-4-one (Compound 70)

[0316]

(Reaction 9-1)

[0317] 8-(3-Chloro-benzenesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.6]undec-1-en-4-one was synthesized by operations similar to those in Reaction 1-2 of Example 1 and Example 7 using appropriate reagents and starting material. MS (ESI) $m/z = 424$ (M+H)+.

[0318] The example compounds shown below were synthesized by operations similar to those in Example 9 using appropriate reagents and starting materials.


[Table 9]

Compound	Structure	LCMS or H PLC condition	Retention time (min)	MS (m/z)
71		LCMS-E-2	2.78	410 (M+H)+
72		LCMS-E-2	2.81	396 (M+H)+

Example 10


4-{2-[2-(2,4-Dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide (Compound 73)

[0319]

(Reaction 10-1)


[0320] 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride n-hydrate (n = about 2.7) (2.42 g, 7.53 mmol) was added to a mixture of 4-bromo-3-methylbenzoic acid (1.58 g, 7.37 mmol), EtOH (26 ml) and a 40% aqueous dimethylamine solution (0.75 ml, 7.4 mmol), and the mixture was stirred at room temperature for 24 hours. The reaction mixture was concentrated under reduced pressure, and the resulting residue was then dissolved in ethyl acetate. The organic layer was washed with water, and then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to give 4-bromo-3-methyl-N,N-trimethylbenzamide as a colorless solid (1.16 g, 65%).

¹H-NMR (300 MHz) (CDCl₃) δ 2.42 (3H, s), 2.98 (3H, br s), 3.10 (3H, br s), 7.08 (1H, dd, J = 8.4 and 2.1 Hz), 7.30 (1H, d, J = 2.1 Hz), 7.55 (1H, d, J = 8.4 Hz). MS (ESI) m/z = 243 (M+H)⁺.

(Reaction 10-2)

[0321] A mixture of 4-bromo-3-methyl-N,N-trimethylbenzamide (798 mg, 3.30 mmol), potassium vinyltrifluoroborate (579 mg, 4.32 mmol), PdCl₂ (59.0 mg, 0.333 mmol), PPh₃ (265 mg, 1.01 mmol) and Cs₂CO₃ (3.22 g, 9.90 mmol) in THF (6.5 ml)-H₂O (0.65 ml) was heated with stirring at 85°C for 21 hours in a sealed test tube in an N₂ atmosphere. The reaction mixture was cooled to room temperature and then extracted with ether. The organic layer was washed with water, and then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to give 3-methyl-4-vinyl-N,N-trimethylbenzamide (565 mg, 90%).

¹H-NMR (400 MHz) (CDCl₃) δ 2.36 (3H, s), 3.00 (3H, br s), 3.10 (3H, br s), 5.35 (1H, dd, J = 11.0 and 1.0 Hz), 5.68 (1H, dd, J = 17.5 and 1.0 Hz), 6.93 (1H, dd, J = 17.5 and 11.0 Hz), 7.21 (1H, d, J = 8.0 Hz), 7.22 (1H, s), 7.48 (1H, d, J = 8.0 Hz). MS (ESI) m/z = 190 (M+H)⁺.


(Reaction 10-3)

[0322] A mixture of 3-methyl-4-vinyl-N,N-trimethylbenzamide (706 mg, 3.73 mmol), triphenylsilanethiol (1.76 g, 6.00 mmol)

and AIBN (185 mg, 1.13 mmol) in toluene (16 ml) was heated with stirring at 88°C for two hours in a sealed test tube in an N₂ atmosphere. The reaction mixture was cooled to room temperature and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to give 3,N,N-trimethyl-4-(2-triphenylsilylsulfanyl-ethyl)-benzamide (1.24 g, 69%).

¹H-NMR (400 MHz) (CDCl₃) δ 2.08 (3H, s), 2.61 (2H, m), 2.75 (2H, m), 2.95 (3H, br s), 3.07 (3H, br s), 6.89 (1H, d, J = 7.8 Hz), 7.08 (1H, d, J = 7.8 Hz), 7.12 (1H, s), 7.37-7.47 (9H, m), 7.66-7.69 (6H, m). MS (ESI) m/z = 482 (M+H)⁺.


(Reaction 10-4)



[0323] Citric acid monohydrate (110 mg, 0.523 mmol) and potassium carbonate (52.8 mg, 0.382 mmol) were added to a solution of 3,N,N-trimethyl-4-(2-triphenylsilylsulfanyl-ethyl)-benzamide (767 mg, 1.59 mmol) in MeOH (27 ml) at room temperature, and the mixture was stirred for one hour. The reaction mixture was concentrated under reduced pressure, and the residue was then dissolved in dichloromethane. The organic layer was washed with water, and then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to give 4-(2-mercapto-ethyl)-3,N,N-trimethyl-benzamide (351 mg, 99%).


¹H-NMR (400 MHz) (CDCl₃) δ 2.33 (3H, s), 2.74 (2H, dt, J = 7.5 and 7.5 Hz), 2.94 (2H, t, J = 7.5 Hz), 2.99 (3H, br s), 3.10 (3H, br s), 7.16 (1H, d, J = 7.8 Hz), 7.18 (1H, d, J = 7.8 Hz), 7.23 (1H, s). MS (ESI) m/z = 224 (M+H)⁺.

(Reaction 10-5)



[0324] Potassium nitrate (583 mg, 5.77 mmol) was added to a solution of 4-(2-mercapto-ethyl)-3,N,N-trimethyl-benzamide (514 mg, 2.30 mmol) in MeCN (23 ml) at room temperature. The mixture was cooled to -40°C, and sulfonyl chloride (1.68 M solution in dichloromethane, 3.46 ml, 5.81 mmol) was then added dropwise over 15 minutes. After stirring at -40°C to -20°C for 2.5 hours, the reaction mixture was diluted with dichloromethane (80 ml) and quenched with a saturated aqueous sodium bicarbonate solution (20 ml). The organic layer and the aqueous layer were separated, and the organic layer was then washed with a saturated aqueous sodium chloride solution (30 ml), dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/1 → 1/2) to give 2-(4-(dimethylcarbamoyl)-2-methyl-phenyl)-ethanesulfonyl chloride as a colorless solid (491 mg, 74%).

¹H-NMR (300 MHz) (CDCl₃) δ 2.39 (3H, s), 2.99 (3H, br s), 3.11 (3H, br s), 3.36 (2H, m), 3.83 (2H, m), 7.19 (1H, d, J = 7.5 Hz), 7.23 (1H, dd, J = 7.5 and 1.5 Hz), 7.28 (1H, d, J = 1.5 Hz). MS (ESI) m/z = 290 (M+H)⁺.

(Reaction 10-6)

[0325] 4-{2-[2-(2,4-Dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide was synthesized by operations similar to those in Reaction 5-4 of Example 5 using appropriate reagents and starting material.

MS (ESI) $m/z = 551$ (M+H)+.

[0326] The example compounds shown below were synthesized by operations similar to those in Example 10 using appropriate reagents and starting materials.

(Compounds 74 to 144)

[0327]











[Table 10]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
74		LCMS-C-2	1.57	463 (M+H)+
75		LCMS-C-1	2.58	551 (M+H)+
76		LCMS-C-1	2.50	503 (M+H)+
n		LCMS-C-1	2.22	475 (M+h)+
78		LCMS-C-1	2.13	519 (M+H)+
79		LCMS-A-1	1.88	543 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
80		LCMS-C-1	2.32	513 (M+H)+
81		LCMS-C-1	2.50	5170 (M+H)+
82		LCMS-A-1	2.72	619 (M+H)+
83		LCMS-C-1	2.23	497 (M+H)+
84		LCQ-01	2.01	497 (M+H)+
85		LCMS-C-1	2.47	585 (M+H)+
86		LCMS-C-1	2.57	551 (M+H)+
87		LCMS-C-1	2.48	517 (M+H)+
88		LCMS-C-1	2.43	551 (M+H)+
89		LCMS-C-1	2.53	535 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
90		LCMS-C-1	2.30	535 (M+H)+
91		LCMS-B-1	2.01	561 (M+H)+
92		LCMS-C-2	1.87	563 (M+H)+
93		LCMS-B-1	1.97	535 (M+H)+
94		LCMS-C-1	2.52	599 (M+H)+
95		LCMS-C-1	2.60	531 (M+H)+
96		LCMS-C-1	2.38	569 (M+H)+
97		LCMS-C-1	2.22	498 (M+H)+
98		LCMS-C-1	2.67	551 (M+H)+
99		LCMS-C-1	2.60	567 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
100		LCMS-C-1	2.63	619 (M+H)+
101		LCMS-C-1	2.33	511 (M+H)+
102		LCMS-C-1	2.07	518 (m+H)+
103		LCMS-A-1	2.02	503 (M+H)+
104		LCMS-A-1	2.12	557 (M+H)+
105		LCMS-C-1	2.40	567 (M+H)+
106		LCMS-C-1	2.62	567 (M+H)+
107		LCMS-C-2	1.97	569 (M+H)+
108		LCMS-C-1	2.65	517 (M+H)+
109		LCMS-A-1	2.10	501 (M+H)+

(continued)



Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
110		LCMS-A-1	1.96	561 (M+H)+
111		LCMS-C-1	2.52	565 (M+H)+
112		LCMS-C-1	2.52	565 (M+H)+
113		LCMS-B-1	1.78	511 (M+H)+
114		LCMS-B-1	1.87	515 (M+H)+
115		LCMS-C-1	2.67	569 (M+H)+
116		LCMS-C-1	2.50	569 (M+H)+
117		LCMS-A-1	2.12	515 (M+H)+
118		LCMS-C-1	2.38	549 (M+H)+
119		LCMS-C-1	2.27	514 (M+H)+
120		LCMS-C-1	2.10	499 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
121		LCMS-C-1	1.98	490 (M+H)+
122		LCMS-C-1	2.05	487 (M+H)+
123		LCMS-C-1	1.95	491 (M+H)+
124		LCMS-C-1	2.48	547 (M+H)+
125		LCMS-C-1	2.32	531 (M+H)+
126		LCMS-C-1	2.72	585 (M+H)+
127		LCMS-C-1	2.78	585 (M+H)+
128		LCMS-C-1	2.42	531 (M+H)+
129		LCMS-C-1	2.42	535 (M+H)+
130		LCMS-A-1	2.19	555 (M+H)+







131

(continued)







Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
143		LCMS-C-1	2.18	540 (M+H)+
144		LCMS-A-1	2.26	581 (M+H)+

[0328] The spiroamine reagents used in the synthesis of Compounds 74 to 85 and shown below were synthesized by operations similar to those in Reaction 7-1 and Reaction 7-2 using appropriate reagents and starting materials.

[Table 11]



Target Compound	Spiroamine reagent	Spiroamine reagent MS(m/z)
74		This compound was directly used in the next step (Reaction 10-6).
75		298 (M+H)+
76		250 (M+H)+
77		222 (M+H)+
78		266 (M+H)+
79		290 (M+H)+

(continued)





Target Compound	Spiroamine reagent	Spiroamine reagent MS(m/z)
80	 2TFA	260 (M+H)+
81	 2TFA	264 (M+H)+
82	 2TFA	366 (M+H)+
83	 2TFA	244 (M+H)+
84	 2TFA	244 (M+H)+
85	 2TFA	332 (M+H)+

[0329] The spiroamine reagents used in the synthesis of Compounds 86 to 91 and shown below were synthesized by operations similar to those in Example 8 using appropriate reagents and starting materials.

[Table 12]


Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
86	 2HCl	298 (M+H)+
87	 2HCl	264 (M+H)+

(continued)

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
88	 2HCl	298 (M+H)+
89	 2HCl	282 (M+H)+
90	 2HCl	282 (M+H)+
91	 2HCl	308 (M+H)+

[0330] The spiroamine reagent used in the synthesis of Compound 92 (2-(2,2-difluoro-benzo[1,3]dioxol-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized as follows.


(Reaction 10-7)



[0331] 4-Cyano-4-[(2,2-difluoro-benzo[1,3]dioxole-4-carbonyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 2-3 of Example 2 using appropriate reagents and starting material.

MS (ESI) m/z = 410 (M+H)+.


(Reaction 10-8)



[0332] A 1 N aqueous NaOH solution (0.274 ml, 0.274 mmol) and a 35% aqueous H_2O_2 solution (0.051 ml, 0.52 mmol) were added to a solution of 4-cyano-4-[(2,2-difluoro-benzo[1,3]dioxole-4-carbonyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester (56.0 mg, 0.137 mmol) in THF (0.23 ml) at room temperature. The reaction mixture was stirred at room temperature for 43 hours, and a 35% aqueous H_2O_2 solution (0.030 ml, 0.31 mmol) was then further added at room temperature. After stirring at room temperature for 48 hours, the mixture was quenched with a 1 N aqueous HCl solution (0.2 ml) and concentrated under reduced pressure. A 6 N aqueous NaOH solution (0.33 ml, 2.0 mmol) was added to a suspension of the resulting residue in EtOH (1.2 ml) at room temperature, and the mixture was stirred at room temperature for 24 hours. The reaction mixture was quenched with a saturated aqueous NH_4Cl solution (0.4 ml) and then concentrated under reduced pressure. The resulting residue was dissolved in ethyl acetate and washed with water, and then dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate = 3/2) to give 2-(2,2-difluoro-benzo[1,3]dioxol-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (51.2 mg, 91%).

$^1\text{H-NMR}$ (300 MHz) (CDCl_3) δ 1.50 (9H, s), 1.55 (2H, m), 1.95 (2H, m), 3.46 (2H, m), 4.02 (2H, br), 7.23-7.25 (2H, m), 7.88-7.94 (1H, m), 8.41 (1H, br s).


(Reaction 10-9)



[0333] 2-(2,2-Difluoro-benzo[1,3]dioxol-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 7-2 of Example 7 using appropriate reagents and starting material. (This compound was directly used in the next reaction.)

[0334] The spiroamine reagent used in the synthesis of Compound 93 (2-(3-chloro-2-fluoro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride) was synthesized as follows.


(Reaction 10-10)



[0335] 4-(3-Chloro-2-fluoro-benzoylamino)-4-cyano-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 2-3 of Example 2 using appropriate reagents and starting material.

MS (ESI) m/z = 404 ($\text{M}+\text{Na}$) $^+$.


(Reaction 10-11)



[0336] 1 N NaOH (8.60 ml, 8.60 mmol) and a 30% H₂O₂ solution (4.30 ml) were added to a solution of 4-(3-chloro-2-fluoro-benzoylamino)-4-cyano-piperidine-1-carboxylic acid tert-butyl ester (1.63 g, 4.28 mmol) in THF (8.60 ml) at room temperature, and the mixture was stirred at room temperature for two hours. The reaction mixture was adjusted to pH 6 by adding 2 N HCl and then extracted with ethyl acetate three times. The organic layers were sequentially washed with H₂O (x 2) and saturated brine, and then dried over Na₂SO₄ and concentrated under reduced pressure. The resulting residue was triturated with H₂O, and the solid was collected by filtration. The resulting solid was washed with Et₂O and then dried under reduced pressure to give 4-carbamoyl-4-(3-chloro-2-fluoro-benzoylamino)-piperidine-1-carboxylic acid tert-butyl ester as a white powder (1.30 g, 76%).

MS (ESI) m/z = 400 (M+H)+.


(Reaction 10-12)



[0337] Potassium t-butoxide (1.01 g, 8.97 mmol) was added to a solution of 4-carbamoyl-4-(3-chloro-2-fluoro-benzoylamino)-piperidine-1-carboxylic acid tert-butyl ester (1.20 g, 2.99 mmol) in tBuOH (30.0 ml) at room temperature, and the mixture was stirred at 40°C for six hours. The reaction mixture was adjusted to pH 6 by adding 2 N HCl and then extracted with AcOEt three times. The organic layers were sequentially washed with H₂O (x 2) and saturated brine, and then dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (hexane/AcOEt = 90:10 -> 50:50) to give 2-(3-chloro-2-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester as a colorless form (1.13 g, 99%).

MS (ESI) m/z = 382 (M+H)+.

(Reaction 10-13)




[0338] 2-(3-Chloro-2-fluoro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride was synthesized by operations similar to those in Reaction 5-3 of Example 5 using appropriate reagents and starting material.

MS (ESI) m/z = 282 (M+H)+.

[0339] The spiroamine reagent used in the synthesis of Compound 94 (2-[4-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride) was synthesized as follows.


(Reaction 10-14)



[0340] N,N,N',N'-Tetramethyl-O-(7-azabenzotriazol-1-yl)uronium hexafluorophosphate (1.60 g, 4.20 mmol) was added to a solution of 4-(1,1,2,2-tetrafluoro-ethoxy)-benzoic acid (1.00 g, 4.20 mmol), 4-amino-4-cyano-piperidine-1-carboxylic acid tert-butyl ester (995 mg, 4.42 mmol) and N,N-diisopropylethylamine (1.46 ml, 8.39 mmol) in DMF (8.8 ml) at 0°C. The mixture was gradually warmed to room temperature and stirred for 28.5 hours. An aqueous ammonium chloride solution was added to the reaction mixture, followed by extraction with ethyl acetate three times. The organic layers were sequentially washed with H₂O (× 2) and saturated brine, and then dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (hexane:AcOEt = 90:10 → 30:70) to give 4-cyano-4-[4-(1,1,2,2-tetrafluoro-ethoxy)-benzoylamino]-piperidine-1-carboxylic acid tert-butyl ester as a light brown powder (1.60 g, 86%).

MS (ESI) m/z = 446 (M+H)+.

(Reaction 10-15)



[0341] 2-[4-(1,1,2,2-Tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride was synthesized by operations similar to those in Reaction 5-2 and Reaction 5-3 of Example 5 using appropriate reagents and starting material.





MS (ESI) m/z = 346 (M+H)+.

[0342] The following spiroamine reagents used in the synthesis of Compounds 95 to 99 were synthesized by operations similar to those in Reaction 10-14 and Reaction 10-15 using appropriate reagents and starting materials.

[Table 13]





Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
95		278 (M+H)+

(continued)










Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
96		316 (M+H)+
97		245 (M+H)+
98		298 (M+H)+
99		314 (M+H)+

[0343] The following spiroamine reagents used in the synthesis of Compounds 100 to 114 were synthesized by operations similar to those in Reaction 10-14, Reaction 5-2 and Reaction 7-2 using appropriate reagents and starting materials.



[Table 14]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
100		366 (M+H)+
101		258 (M+H)+
102		265 (M+H)+
103		250 (M+H)+

(continued)




Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
104	 2TFA	304 (M+H)+
105	 2TFA	314 (M+H)+
106	 2TFA	314 (M+H)+
107	 2TFA	315 (M+H)+
108	 2TFA	264 (M+H)+
109	 2TFA	248 (M+H)+
110	 2TFA	308 (M+H)+
111	 2TFA	312 (M+H)+
112	 2TFA	312 (M+H)+

(continued)

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
113	 2TFA	258 (M+H)+
114	 2TFA	262 (M+H)+


[0344] The following spiroamine reagents used in the synthesis of Compounds 115 to 117 were synthesized by operations similar to those in Reaction 10-14, Reaction 10-8 and Reaction 7-2 using appropriate reagents and starting materials.

[Table 15]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
115	 2TFA	315 (M+H)+
116	 2TFA	315 (M+H)+
117	 2TFA	262 (M+H)+

[0345] The following spiroamine reagent used in the synthesis of Compound 118 (2-(4-difluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one) was synthesized by operations similar to those in Reaction 10-14, Reaction 10-8 and Reaction 5-3 using appropriate reagents and starting material.


[Table 16]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
118	 2HCl	296 (M+H)+

[0346] The spiroamine reagent used in the synthesis of Compound 119 (2-(2-methoxy-pyridin-4-yl)-1,3,8-triaza-

spiro[4.5]dec-1-en-4-one dinitrfluoroacetate) was synthesized as follows.


(Reaction 10-16)



[0347] 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride 2.7-hydrate (1.06 g, 3.26 mmol) was added to a solution of 2-methoxy-isonicotinic acid (500 mg, 3.26 mmol) and 4-amino-4-cyano-piperidine-1-carboxylic acid tert-butyl ester (669 mg, 2.97 mmol) in EtOH (8.0 ml) at room temperature, and the mixture was stirred for 46.5 hours. An aqueous NaHCO₃ solution was added to the reaction mixture, followed by extraction with AcOEt three times. The organic layers were washed with saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by column chromatography (CH₂Cl₂/MeOH = 99:1 to 95:5) to give 4-cyano-4-[(2-methoxy-pyridine-4-carbonyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester as a colorless form (901 mg, 84%).

MS (ESI) m/z = 361 (M+H)+.

(Reaction 10-17)












[0348] 2-(2-Methoxy-pyridin-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dinitrfluoroacetate was synthesized by operations similar to those in Reaction 5-2 and Reaction 7-2 using appropriate reagents and starting material. (This compound was directly used in the next reaction.)

[0349] The following spiroamine reagents used in the synthesis of Compounds 120 to 131 were synthesized by the procedure described in Reaction 10-16 and Reaction 10-17 using appropriate reagents and starting materials.



[Table 17]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
120		246 (M+H)+

(continued)


Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
121	 2TFA	This compound was directly used in the next step (Reaction 10-6).
122	 3TFA	234 (M+H)+
123	 2TFA	This compound was directly used in the next step (Reaction 10-6).
124	 2TFA	294 (M+H)+
125	 2TFA	278 (M+H)+
126	 2TFA	332 (M+H)+
127	 2TFA	331 (M+H)+
128	 2TFA	278 (M+H)+
129	 2TFA	282 (M+H)+

(continued)

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
130	 2TFA	302 (M+H)+
131	 2TFA	328 (M+H)+


[0350] The following spiroamine reagent used in the synthesis of Compound 133 was synthesized by operations similar to those in Reaction 10-16, Reaction 10-8 and Reaction 5-3 using appropriate reagents and starting material.

[Table 18]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
133	 2HCl	378 (M+H)+

[0351] The spiroamine reagent used in the synthesis of Compound 134 (2-(2,4-dichloro-benzyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized as follows.


(Reaction 10-18)



[0352] (2,4-Dichloro-phenyl)-acetic acid (218 mg, 1.07 mmol), 1-ethyl-3-(3'-dimethylamino-propyl)carbodiimide hydrochloride (255 mg, 1.33 mmol), 1-hydroxybenzotriazole hydrate (136 mg, 0.88 mmol) and N,N-diisopropylethylamine (0.378 ml, 2.22 mmol) were sequentially added to a solution of 4-amino-4-cyano-piperidine-1-carboxylic acid tert-butyl ester (200 mg, 0.888 mmol) in DMF (4 ml) at room temperature, and the mixture was stirred at room temperature for 16 hours. H₂O (20 ml) was added to the reaction mixture, followed by extraction with AcOEt (40 ml and 20 ml). The organic layers were sequentially washed with H₂O (20 ml), 1 N HCl (20 ml), H₂O (20 ml) and saturated brine (20 ml), and then dried over Na₂SO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane/AcOEt) to give 4-cyano-4-[2-(2,4-dichlorophenyl)-acetylamino]-piperidine-1-carboxylic acid tert-butyl ester as a white powder (285 mg, 78%).

¹H-NMR (270 MHz, CDCl₃) δ 1.45 (9H, s), 1.72 (2H, ddd, J = 13.2, 10.7, 3.9 Hz), 2.34-2.37 (2H, m), 3.20-3.27 (2H, m), 3.68 (2H, s), 3.81-3.97 (2H, m), 5.55 (1H, s), 7.28 (1H, dd, J = 7.8, 2.0 Hz), 7.30 (1H, d, 7.8 Hz), 7.45 (1H, d, J = 2.0 Hz). MS (ESI) m/z = 412 (M+H)+.

(Reaction 10-19)



[0353] 2-(2,4-Dichloro-benzyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 5-2 and Reaction 7-2 using appropriate reagents and starting material.

MS (ESI) *m/z* = 312 (M+H)+.


[0354] The following spiroamine reagent used in the synthesis of Compound 135 was synthesized by operations similar to those in Reaction 10-18 and Reaction 10-19 using appropriate reagents and starting material.

[Table 19]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (<i>m/z</i>)
135	2TFA	312 (M+H)+

[0355] The spiroamine reagent used in the synthesis of Compound 136 (2-(1-trifluoromethyl-cyclopropyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized as follows.


(Reaction 10-20)



[0356] Oxalyl chloride (0.20 ml, 2.3 mmol) and dimethylformamide (8 μ l) were added to a solution of 1-trifluoromethyl-cyclopropanecarboxylic acid (308 mg, 2.00 mmol) in dichloromethane (2.1 ml) at 0°C. The mixture was stirred at 0°C for 30 minutes and then stirred at room temperature for two hours. The reaction mixture was concentrated under reduced pressure. A solution of the resulting residue in dichloromethane (1.5 ml) was added dropwise to a solution of 4-amino-4-cyanopiperidine-1-carboxylic acid tert-butyl ester (377 mg, 1.67 mmol) and diisopropylethylamine (0.42 ml, 2.4 mmol) in dichloromethane (2.0 ml) over three minutes at 0°C, and the mixture was stirred at room temperature for 13 hours. The reaction mixture was diluted with dichloromethane, and the organic layer was then washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 3/1 $\rightarrow$ 2/1) to give 4-cyano-4-[(1-trifluoromethyl-cyclopropanecarbonyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester as a colorless solid (519 mg, 86%).

¹H-NMR (400 MHz) (CDCl₃) δ 1.46 (9H, s), 1.29 (2H, dd, *J* = 7.5 and 4.5 Hz), 1.56 (2H, m), 1.80 (2H, m), 2.40 (2H, m), 3.30 (2H, m), 3.93 (2H, br), 6.07 (1H, br s). *R_f* = 0.62 in TLC (developer; hexane:AcOEt = 1:1).

(Reaction 10-21)



[0357] 2-(1-Trifluoromethyl-cyclopropyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 5-2 and Reaction 7-2 using appropriate reagents and starting material. (This compound was directly used in Reaction 10-6.)


[0358] The following spiro-amine reagents used in the synthesis of Compounds 137 to 138 were synthesized by operations similar to those in Reaction 10-20 and Reaction 10-21 using appropriate reagents and starting materials. (These compounds were directly used in Reaction 10-6.)

[Table 20]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
137		This compound was directly used in the next step (Reaction 10-6).
138		This compound was directly used in the next step (Reaction 10-6).

[0359] The spiroamine reagent used in the synthesis of Compound 139 (2-(2-fluoro-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized as follows.


(Reaction 10-22)



[0360] HATU (939 mg, 2.47 mmol) and DIPEA (525 μ L, 3.09 mmol) were added to a solution of 4-amino-4-carbamoyl-piperidine-1-carboxylic acid tert-butyl ester (500 mg, 2.06 mmol) and 2-fluoro-4-trifluoromethyl-benzoic acid (514 mg, 2.47 mmol) in DMF (10 mL). The mixture was stirred at room temperature for 19 hours and then quenched with a saturated aqueous ammonium chloride solution. The reaction mixture was diluted with EtOAc, and the organic layer was then washed with H₂O and saturated brine, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was triturated with n-hexane and EtOAc and then collected by filtration to give 4-carbamoyl-4-(2-fluoro-

4-trifluoromethyl-benzoylamino)-piperidine-1-carboxylic acid tert-butyl ester as a white solid. This was used in the next step without further purification.


(Reaction 10-23)



[0361] A 6 N aqueous NaOH solution (54.8 μ L, 323 μ mol) was added to a solution of 4-carbamoyl-4-(2-fluoro-4-trifluoromethyl-benzoylamino)-piperidine-1-carboxylic acid tert-butyl ester (100 mg, 231 μ mol) in DMSO (0.3 mL), and the mixture was stirred at room temperature for 27 hours. The reaction mixture was quenched with a saturated aqueous ammonium chloride solution and then diluted with EtOAc, and the organic layer was sequentially washed with H₂O and saturated brine. The organic layer was dried over MgSO₄ and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (CH₂Cl₂/MeOH = 95:5) to give 2-(2-fluoro-4-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester as a white solid (54.5 mg, 57%).

MS (ESI) m/z = 438 (M+Na)+.

(Reaction 10-24)



[0362] 2-(2-Fluoro-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 7-2 using appropriate reagents and starting material.



MS (ESI) m/z = 315 (M+H)+.

[0363] The following spiro-amine reagents used in the synthesis of Compounds 140 to 143 were synthesized by operations similar to those in Reaction 10-22, Reaction 10-23 and Reaction 10-24 using appropriate reagents and starting materials.

[Table 21]


Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
140	 2TFA	346 (M+H)+
141	 2TFA	194 (M+H)+

(continued)

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
142	 2TFA	256 (M+H)+
143	 2TFA	287 (M+H)+

[0364] The spiroamine reagent used in the synthesis of Compound 144 (2-(4-methoxy-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized as follows.


(Reaction 10-25)



[0365] 4-Carbamoyl-4-(4-methoxy-3-trifluoromethyl-benzoylamino)-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 2-3 using 4-amino-4-carbamoyl-piperidine-1-carboxylic acid tert-butyl ester as a starting material amine.

MS (ESI) m/z = 446 (M+H)+.

(Reaction 10-26)



[0366] 2-(4-Methoxy-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 10-23 and Reaction 7-2 using appropriate reagents and starting material.


MS (ESI) m/z = 328 (M+H)+.

Example 11

N-{3-Methyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-acetamide (Compound 145)


[0367]

(Reaction 11-1)



[0368] 2-(4-Acetylamino-2-methyl-phenyl)-ethanesulfonyl chloride was synthesized by operations similar to those in Reaction 10-2, Reaction 10-3, Reaction 10-4 and Reaction 10-5 using appropriate reagents and starting material. MS (ESI) $m/z = 276$ ($M+H$) $^+$.

(Reaction 11-2)





[0369] N-{3-Methyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-acetamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material. MS (ESI) $m/z = 483$ ($M+H$) $^+$.

[0370] The example compounds shown below were synthesized by operations similar to those in Example 11 using appropriate reagents and starting materials.

[Table 22]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
146		LCMS-C-1	2.33	497 ($M+H$) $^+$

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
147		LCMS-C-1	2.40	537 (M+H)+
148		LCMS-C-1	2.45	489 (M+H)+


[0371] The spiroamine reagent used in the synthesis of Compound 148 (2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride) was synthesized as follows.

(Reaction 11-3)



[0372] 2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 5-1 and Reaction 5-2 using appropriate reagents and starting material. MS (ESI) m/z = 372 (M+Na)+.

(Reaction 11-4)




[0373] 2-(4-Methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride was synthesized by operations similar to those in Reaction 7-2 and Reaction 5-3 using appropriate reagents and starting material. MS (ESI) m/z = 250 (M+H)+.


Example 12

N-(2-Hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide (Compound 149)


[0374]

(Reaction 12-1)

[0375] A mixture of 2-bromo-5-iodotoluene (1.60 g, 5.40 mmol), 2-aminoethanol (0.49 mL, 8.14 mmol), CuI (53.3 mg, 0.28 mmol), L-proline (63.4 mg, 0.55 mmol) and K₂CO₃ (1.49 g, 10.8 mmol) in DMSO (3.24 mL) was stirred at 60°C for 12 hours. The reaction mixture was cooled and then diluted with AcOEt, and the organic layer was sequentially washed with H₂O and saturated brine. The organic layer was dried over Na₂SO₄ and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/AcOEt = 1/1) to give 2-(4-bromo-3-methyl-phenylamino)-ethanol as a brown solid (1.00 g, 81%). MS (ESI) m/z = 230, 232 (M+H)⁺.

(Reaction 12-2)

[0376] Pyridine (158.4 mL, 1.958 mol) was added to a solution of 2-(4-bromo-3-methyl-phenylamino)-ethanol (19.28 g, 83.788 mmol) in Ac₂O (158.4 mL, 1.676 mol). The mixture was stirred at room temperature for 18 hours and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (CH₂Cl₂/MeOH = 100/0 to 95/5) to give acetic acid 2-[acetyl-(4-bromo-3-methyl-phenyl)-amino]-ethyl ester as a brown viscous oil (21.44 g, 81%). MS (ESI) m/z = 314, 316 (M+H)⁺.


(Reaction 12-3)

[0377] Acetic acid 2-[acetyl-[4-(2-chlorosulfonyl-ethyl)-3-methyl-phenyl]-amino]-ethyl ester was synthesized by oper-

ations similar to those in Reaction 10-2, Reaction 10-3, Reaction 10-4 and Reaction 10-5 using appropriate reagents and starting material.

MS (ESI) $m/z = 362$ (M+H)+.


(Reaction 12-4)



[0378] Acetic acid 2-[acetyl-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester was synthesized by operations similar to those in Reaction 5-4 of Example 5 using appropriate reagents and starting material.

MS (ESI) $m/z = 639$ (M+H)+.

(Reaction 12-5)



[0379] K_2CO_3 (9.1 mg, 66.0 μ mol) was added to a solution of acetic acid 2-[acetyl-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester (28.1 mg, 44.0 μ mol) in MeOH (0.5 mL). The mixture was stirred at room temperature for two hours. H_2O was then added and the mixture was diluted with CH_2Cl_2 . The organic layer was washed with saturated brine, and then dried over $MgSO_4$ and concentrated under reduced pressure. The resulting residue was purified by column chromatography ($CH_2Cl_2/MeOH = 15/1$) to give N-(2-hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide as a white amorphous (24.0 mg, 92%).


^1H -NMR (400 MHz, $CDCl_3$) δ 1.71-1.75 (2H, m), 1.90 (3H, s), 2.09-2.04 (2H, m), 2.40 (3H, s), 3.16-3.24 (5H, m), 3.46-3.53 (4H, m), 3.77-3.87 (4H, m), 7.04-7.06 (2H, m), 7.24-7.26 (1H, m), 7.42-7.44 (1H, m), 7.58 (1H, t, $J = 8.3$ Hz), 7.79-7.81 (1H, m), 7.86 (1H, m). MS (ESI) $m/z = 597$ (M+H)+.

Example 13

Acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester (Compound 150)


[0380]

(Reaction 13-1)



[0381] Acetic acid (S)-1-acetoxymethyl-2-[acetyl-4-(2-chlorosulfonyl-ethyl)-3-methyl-phenyl]-amino-ethyl ester was synthesized by operations similar to those in Reaction 12-1, Reaction 12-2, Reaction 10-2, Reaction 10-3, Reaction 10-4 and Reaction 10-5 using appropriate reagents and starting material.
MS (ESI) $m/z = 434$ (M+H)+.

(Reaction 13-2)



[0382] Acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-amino]-ethyl ester was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.
MS (ESI) $m/z = 695$ (M+H)+.


[0383] The example compounds shown below were synthesized by operations similar to those in Example 13 using appropriate reagents and starting materials.

[Table 23]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	ms (m/z)
151		LCMS-A-1	2.53	695 (M+H)+
152		LCMS-A-1	2.50	711 (M+H)+
153		LCMS-A-1	2.57	713 (M+H)+

Example 14

8-{2-[4-((S)-2,3-Dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 154)

[0384]**(Reaction 14-1)**

[0385] A 1.2 M aqueous KOH solution (0.5 mL) was added to a solution of acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-amino]-ethyl ester (40.8 mg, 0.0587 mmol) in MeOH (3 mL). The reaction mixture was stirred at 50°C for 1.5 hours and then cooled to room temperature. Dowex 50W × 4 (237.6 mg) was added. The mixture was further stirred at room temperature for two hours and then filtered, and the organic layer was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (CH₂Cl₂/MeOH = 10/1) to give 8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a white powder (30.3 mg, 91%).

¹H-NMR (400 MHz, CD₃OD) δ 1.72-1.75 (2H, m), 1.99-2.04 (2H, m), 2.28 (3H, s), 2.98-3.06 (3H, m), 3.19-3.28 (2H, m), 3.46-3.61 (5H, m), 3.77-3.80 (3H, m), 6.49-6.53 (2H, m), 6.98 (1H, d, J = 8.3 Hz), 7.85 (2H, d, J = 8.3 Hz), 8.12 (2H, d, J = 8.3 Hz). MS (ESI) m/z = 569 (M+H)+.

[0386] The example compound shown below was synthesized by operations similar to those in Example 14 using appropriate reagents and starting material.

[Table 24]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
155		LCMS-A-1	1.93	569 (M+H)+

Example 15

8-{2-[4-((S)-2,3-Dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 156)

[0387]

(Reaction 15-1)



[0388] 8-{2-[4-((S)-2,3-Dihydroxy-propylamino)-2-methylphenyl]-ethanesulfonyl}-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 5-4 and Reaction 14-1 using appropriate reagents and starting material.

MS (ESI) m/z = 585 (M+H)+.

[0389] The example compounds shown below were synthesized by operations similar to those in Example 15 using appropriate reagents and starting materials.

[Table 25]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
157		LCMS-B-1	1.54	535 (M+H)+
158		LCMS-C-1	2.35	535 (M+H)+

Example 16

8-{2-[4-(2-Hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 159)

[0390]

(Reaction 16-1)



[0391] 8-{2-[4-(2-Hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 5-4 and Reaction 14-1 using appropriate reagents and starting material.

MS (ESI) m/z = 555 (M+H)+.

[0392] The example compounds shown below were synthesized by operations similar to those in Example 16 using appropriate reagents and starting materials.


[Table 26]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
160		LCMS-B-1	1.73	539 (M+H)+
161		LCMS-B-1	1.47	539 (M+H)+

Example 17


{4-[2-(2-tert-Butyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-carbamic acid tert-butyl ester (Compound 162)

[0393]

(Reaction 17-1)


[0394] NaBH_4 (1.45 g, 38.25 mmol) was added in small portions to a mixture of 4-bromo-3-methyl-benzonitrile (2.50 g, 12.8 mmol), NiCl_2 (1.65 g, 12.8 mmol) and Boc_2O (5.57 g, 25.5 mmol) in anhydrous MeOH (130 ml) at 0°C . The mixture was stirred at room temperature for two hours and then concentrated under reduced pressure. Ethyl acetate and water were added to the resulting residue, and the mixture was filtered through celite. The two-layer solution was separated, and the aqueous layer was then further extracted with ethyl acetate. The organic layers were combined, dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/0 $\rightarrow$ 4/1) to give (4-bromo-3-methyl-benzyl)-carbamic acid tert-butyl ester as a white solid (2.42 g, 63%).

MS (ESI) m/z = 322 ($\text{M}+\text{Na}$) $^+$.

(Reaction 17-2)

[0395] [4-(2-Chlorosulfonyl-ethyl)-3-methyl-benzyl]-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-2, Reaction 10-3, Reaction 10-4 and Reaction 10-5 using appropriate reagents and starting material.

MS (ESI) m/z = 292 ($\text{M-tBu}+\text{Hx}_2$) $^+$.

(Reaction 17-3)

[0396] {4-[2-(2-tert-Butyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-carbamic acid

tert-butyl ester was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 521 (M+H)^+$.

[0397] The example compounds shown below were synthesized by operations similar to those in Example 17 using appropriate reagents and starting materials.

[Table 27]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
163		LCMS-C-1	2.67	583 (M+H)+
164		LCMS-C-2	2.35	625 (M-H)-
165		LCMS-B-1	1.92	549 (M+H)+


[0398] The spiroamine reagent used in the synthesis of Compound 163 (2-(4,4-difluoro-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized by operations similar to those in Reaction 10-14, Reaction 5-2 and Reaction 7-2 using appropriate reagents and starting material.

Example 18

8-[2-(3-Amino-phenyl)-ethanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 166)

[0399]


(Reaction 18-1)



[0400] A mixture of 1-(2-bromo-ethyl)-3-nitro-benzene (4 g, 17.4 mmol) and thiourea (1.5 g, 19.1 mmol) in ethanol (20 mL) was heated under reflux for one hour. The reaction mixture was concentrated under reduced pressure to give Compound 18b as a pale yellow solid. Further, NCS (7.66 g, 57.4 mmol) was added to a mixed solution of this solid in acetic acid (43.5 ml) and H₂O (14.5 ml) on an ice bath, and the mixture was stirred at 5 to 10°C for 50 minutes. The

reaction mixture was diluted with CH_2Cl_2 , and the organic layer was then washed with water, dried over Na_2SO_4 and concentrated under reduced pressure. Compound 7c (5.2 g, 11.2 mmol) and Et_3N (6.3 mL, 44.9 mmol) were added to a solution of the resulting Compound 18c in CH_2Cl_2 , and the mixture was stirred at room temperature for four hours. The reaction mixture was concentrated under reduced pressure, and the residue was then purified by silica gel flash chromatography to give 2-cyclohexyl-8-[2-(3-nitro-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (18d) as a white solid (1 g, yield 20% (three steps)). This compound was directly used in the next step.

(Reaction 18-2)




[0401] 10% Pd-C (1 g) was added to a solution of Compound 18d (1 g, 2.23 mmol) in ethanol (10 ml), and the mixture was stirred at room temperature for two days in an H_2 atmosphere. The reaction mixture was filtered, and the filtrate was then concentrated under reduced pressure to give 8-[2-(3-amino-phenyl)-ethanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (920 mg, 98%). MS (ESI) $m/z = 419$ ($\text{M}+\text{H}$) $^+$.

Example 19


N-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-acetamide (Compound 167)

[0402]

(Reaction 19-1)




[0403] 8-[2-(4-Amino-phenyl)-ethanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 5-4 and Reaction 18-2 using appropriate reagents and starting material. This compound was directly used in the next step.

(Reaction 19-2)


[0404] Acetic anhydride (45 mg, 0.44 mmol) was added to a solution of 8-[2-(4-amino-phenyl)-ethanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (92 mg, 0.22 mmol) in CH_2Cl_2 (5 mL). Triethylamine (55 mg, 0.5 mmol) was then added on an ice bath, and the mixture was stirred at room temperature for one hour. The reaction mixture was diluted with dichloromethane, and the organic layer was then sequentially washed with water and saturated brine and dried over Na_2SO_4 . The organic layer was concentrated under reduced pressure, and the resulting residue was then purified by P-TLC to give N-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-acetamide (55 mg, 54.3%).
MS (ESI) $m/z = 461$ (M+H)+.

Example 20


3,N,N-Trimethyl-4-(2-{4-oxo-2-[3-(2,2,2-trifluoro-ethoxymethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide (Compound 168)

[0405](Reaction 20-1)

[0406] 4-(3-Chloromethyl-benzoylamino)-4-cyano-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.
MS (ESI) $m/z = 342$ (M+H)+.

(Reaction 20-2)

[0407] Sodium hydride (60% oil suspension, 191 mg, 4.77 mmol) was added to a solution of 2,2,2-trifluoro-ethanol (347 μL , 4.77 mmol) and 4-(3-chloromethyl-benzoylamino)-4-cyano-piperidine-1-carboxylic acid tert-butyl ester (600 mg, 1.59 mmol) in DMF (8 mL) at 0°C . The mixture was stirred at room temperature overnight, and then quenched with water and diluted with ethyl acetate. The organic layer was sequentially washed with a saturated aqueous NaHCO_3 solution, water (x 2) and saturated brine, and then dried over Na_2SO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1 $\rightarrow$ 1/2) to give 4-cyano-4-[3-(2,2,2-trifluoro-ethoxymethyl)-benzoylamino]-piperidine-1-carboxylic acid tert-butyl ester as a white solid (366 mg, 52%).

MS (ESI) $m/z = 442$ (M+H)+.(Reaction 20-3)

[0408] 3,N,N-Trimethyl-4-(2-{4-oxo-2-[3-(2,2,2-trifluoro-ethoxymethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide was synthesized by operations similar to those in Reaction 5-2, Reaction 7-2 and Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 595$ (M+H)+.

[0409] The example compounds shown below were synthesized by operations similar to those in Example 20 using appropriate reagents and starting materials.


[Table 28]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
169		LCMS-C-1	2.23	577 (M+H)+
170		LCMS-C-1	2.53	627 (M+H)+

Example 21


4-(2-{2-[3-(3,5-Dimethyl-isoxazol-4-yl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide (Compound 171)

[0410]

(Reaction 21-1)

[0411] A mixture of 2-(3-bromo-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (100 mg, 0.245 mmol), 3,5-dimethyl-isoxazole-4-boronic acid (51.8 mg, 0.367 mmol), tetrakis-(triphenylphosphine)palladium(0) (28 mg, 0.0245 mmol) and K_3PO_4 (104 mg, 0.490 mmol) in dioxane (1.2 mL) was heated with stirring at 100°C for one hour in a nitrogen atmosphere. The reaction mixture was cooled, and then quenched with water and extracted with ethyl acetate ($\times 3$). The organic layers were combined and sequentially washed with water ($\times 2$) and saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane:ethyl acetate = 2:1) to give 2-[3-(3,5-dimethyl-isoxazol-4-yl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester as a pale yellow solid (88.3 mg, 85%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.51 (9H, s), 1.52-1.63 (2H, m), 1.89-2.27 (2H, m), 2.31 (3H, s), 2.45 (3H, s), 3.38-3.55 (2H, m), 3.94-4.12 (2H, m), 7.45 (1H, d, $J = 7.8$ Hz), 7.61 (1H, dd, $J = 7.8, 7.8$ Hz), 7.85 (1H, s), 7.92 (1H, d, $J = 7.8$ Hz), 10.20 (1H, brs).



(Reaction 21-2)

[0412] 4-(2-(2-[3-(3,5-Dimethyl-isoxazol-4-yl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3,N,N-trimethyl-benzamide was synthesized by operations similar to those in Reaction 5-3 and Reaction 5-4 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ 1.73-1.82 (2H, m), 1.98-2.07 (2H, m), 2.27 (3H, s), 2.42 (3H, s), 2.43 (3H, s), 3.00 (3H, s), 3.09 (3H, s), 3.14-3.22 (2H, m), 3.32-3.38 (2H, m), 3.45-3.55 (2H, m), 3.75-3.84 (2H, m), 7.23 (1H, d, $J = 7.8$ Hz), 7.26 (1H, s), 7.34 (1H, d, $J = 7.8$ Hz), 7.60 (1H, d, $J = 7.8$ Hz), 7.66 (1H, d, $J = 7.8$ Hz), 7.92 (1H, s), 7.99 (1H, d, $J = 7.8$ Hz). MS (ESI) $m/z = 578$ ($\text{M}+\text{H}$) $^+$.

[0413] The example compounds shown below were synthesized by operations similar to those in Example 21 using appropriate reagents and starting materials.


[Table 29]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
172		LCMS-B-1	2.11	559 (M+H)+
173		LCMS-B-1	1.50	560 (M+H)+


Example 22

3-{8-[2-(4-Dimethylcarbamoyl-2-methyl-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl}-benzoic acid methyl ester (Compound 174)

[0414]

(Reaction 22-1)

[0415] 2-(3-Carboxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-14 and Reaction 2-4 using appropriate reagents and starting material. MS (ESI) m/z = 374 (M+H)+.


(Reaction 22-2)

[0416] (Trimethylsilyl)diazomethane (2.0 M in hexane, 4.0 ml, 8.0 mmol) was added dropwise to a solution of 2-(3-carboxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (200 mg, 0.54 mmol) in meth-

anol (10 ml). The mixture was stirred at room temperature for one hour, and (trimethylsilyl)diazomethane (1.0 ml, 2.0 mmol) was then further added, followed by stirring for one hour. The reaction mixture was concentrated under reduced pressure, and the resulting solid was then washed with a solution of hexane/ethyl acetate = 5/1 to give 2-(3-methoxycarbonyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (91.5 mg, 44%).

MS (ESI) m/z = 388 (M+H)⁺

(Reaction 22-3)



[0417] 3-{8-[2-(4-Dimethylcarbamoyl-2-methyl-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl}-benzoic acid methyl ester was synthesized by operations similar to those in Reaction 4-1 and Reaction 5-4 using appropriate reagents and starting material.


MS (ESI) m/z = 541 (M+H)⁺.

Example 23

4-(2-{2-[1-(2,4-Dichloro-phenoxy)-1-methyl-ethyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide (Compound 175)


[0418]

(Reaction 23-1)




[0419] 2,4-Dichlorophenol (448 mg, 2.75 mmol) and K_2CO_3 (775 mg, 5.61 mmol) were continuously added to 2-bromo-2-methyl-propionic acid ethyl ester (800 mg, 4.10 mmol) in N,N-dimethylacetamide (4 ml) at room temperature. The mixture was stirred at 110°C for 14 hours, and saturated NH_4Cl and H_2O were then added, followed by extraction with AcOEt ($\times 2$). The organic layers were combined and sequentially washed with H_2O and saturated brine, and then dried over Na_2SO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane/AcOEt) to give 2-(2,4-dichloro-phenoxy)-2-methyl-propionic acid ethyl ester (389 mg, 50%).

^1H -NMR (270 MHz, $CDCl_3$): δ : 1.28 (3H, t, J = 7.3 Hz), 1.60 (6H, s), 4.25 (2H, q, J = 7.3 Hz), 6.86 (1H, d, J = 8.8 Hz), 7.10 (1H, dd, J = 8.8, 2.4 Hz), 7.38 (1H, d, J = 2.4 Hz).




(Reaction 23-3)




[0422] The example compound shown below was synthesized by operations similar to those in Example 23 using appropriate reagents and starting material.

[Table 30]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
176		LCMS-C-1	2.42	567 (M+H)+


Example 24

2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic (3-ethylphenyl)amide (Compound 177)

[0423](Reaction 24-1)

[0424] 2-Bromoethanol (0.28 mL, 4.00 mmol) was added to a solution of chlorosulfonyl isocyanate (0.35 mL, 4.00 mmol) in dichloromethane (1.8 mL) at 0°C. After stirring for 90 minutes, a solution of 3-ethylaniline (0.55 mL, 4.40 mmol) and triethylamine (1.23 mL, 8.80 mmol) in dichloromethane (3.6 mL) was added. The mixture was stirred for 90 minutes and then quenched with a 2 N aqueous hydrochloric acid solution. The mixed solution was separated, and the aqueous layer was then extracted with ether. The organic layers were combined and washed with water and saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was washed with ether (5 mL) to give 2-oxo-oxazolidine-3-sulfonic acid (3-ethylphenyl)amide (LCMS yield 80%).

MS (ESI) m/z = 271 (M+H)+.

(Reaction 24-2)

[0425] A solution of 2-oxo-oxazolidine-3-sulfonic acid (3-ethylphenyl)amide (92 mg, 0.340 mmol) and 2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate (93 mg, 0.395 mmol) in acetonitrile (0.80 mL) was irradiated with microwaves (150°C, 15 min). The reaction mixture was filtered, and the resulting filtrate was then concentrated under reduced pressure. Further, the resulting residue was purified by silica gel chromatography to give 2-cyclohexyl-4-oxo-

1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonic acid (3-ethylphenyl)amide as a white amorphous (33 mg, 23%).


$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 1.24 (3H, t, $J = 7.8$ Hz), 1.31-1.93 (14H, m), 2.36-2.40 (1H, m), 2.64 (2H, q, $J = 7.8$ Hz), 3.34-3.44 (2H, m), 3.69-3.76 (2H, m), 6.54 (1H, s), 6.96-7.02 (3H, m), 7.20-7.24 (1H, m), 8.27 (1H, s). MS (ESI) $m/z = 419$ ($M+H$) $^+$.

Example 25

2-Cyclohexyl-8-[(E)-2-(1H-indol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 178)

[0426]


(Reaction 25-1)



[0427] 2-Chloro-ethanesulfonyl chloride (440 μl , 4.21 mmol) was added to a solution of 2-cyclohexyl-1,3,8-triazaspiro[4.5]dec-1-en-4-one dihydrochloride (1.50 g, 3.24 mmol) and triethylamine (2.7 ml, 19.4 mmol) in CH_2Cl_2 (30 ml) at room temperature in an N_2 atmosphere. The mixture was stirred at room temperature for 30 minutes, and then washed with water, dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was triturated with AcOEt-hexane, and the solid was then collected by filtration and dried to give 2-cyclohexyl-8-ethenesulfonyl-1,3,8-triazaspiro[4.5]dec-1-en-4-one as a colorless solid (692 mg, 66%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.25-1.45 (6H, m), 1.70-2.05 (8H, m), 2.40-2.47 (1H, m), 3.21-3.30 (2H, m), 3.61-3.69 (2H, m), 6.03 (1H, d, $J = 8.0$ Hz), 6.26 (1H, d, $J = 16.0$ Hz), 6.49 (1H, dd, $J = 16.0, 8.0$ Hz), 8.17 (1H, brs). MS (ESI) $m/z = 326$ ($M+H$) $^+$.

(Reaction 25-2)



[0428] 2-Cyclohexyl-8-ethenesulfonyl-1,3,8-triazaspiro[4.5]dec-1-en-4-one (60.0 mg, 0.184 mmol), 5-bromo-indole (72.0 mg, 0.367 mmol), palladium(II) acetate (4.1 mg, 0.0183 mmol), tris(o-tolyl)phosphine (11.2 mg, 0.0368 mmol), triethylamine (0.077 ml, 0.552 mmol) and DMA (0.6 ml) were mixed in a sealed test tube in an N_2 atmosphere. This mixture was irradiated with microwaves (190°C, 20 min). The reaction mixture was cooled, and then quenched with saturated brine and extracted with ethyl acetate three times. The organic layers were combined, sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (CH_2Cl_2 -MeOH) to give 2-cyclohexyl-8-[(E)-2-(1H-indol-5-yl)-ethenesulfonyl]-1,3,8-triazaspiro[4.5]dec-1-en-4-one as a yellow form (40.6 mg, 50%).










$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 1.20-1.45 (5H, m), 1.52-1.95 (7H, m), 1.98-2.11 (2H, m), 2.35-2.48 (1H, m), 3.20-3.31 (2H, m), 3.68-3.79 (2H, m), 6.59-6.63 (1H, m), 6.65 (1H, d, $J = 16$ Hz), 7.25-7.28 (1H, m), 7.33-7.44 (2H, m), 7.60 (1H, d, $J = 16$ Hz), 7.77-7.79 (1H, m), 8.33 (1H, brs), 8.37 (1H, brs). MS (ESI) $m/z = 441$ ($M+H$) $^+$.

[0429] The example compounds shown below were synthesized by operations similar to those in Example 25 using appropriate reagents and starting materials.









(Compounds 179 to 203)

[0430]

[Table 31]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
179		LCMS-E-8	3.68	470 (M+H)+
180		LCMS-E-5	3.14	432 (M+H)+
181		LCMS-C-4	2.91	432 (M+H)+
182		LCMS-C-1	2.48	441 (M+H)+
183		LCMS-E-6	1.66	488 (M+H)+
184		LCMS-E-6	1.53	438 (M+H)+
185		LCMS-E-6	1.7	488 (M+H)+
186		LCMS-C-1	2.73	416 (M+H)+
187		LCMS-C-1	2.37	446 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
188		LCMS-A-1	1.84	459 (M+H)+
189		LCMS-C-1	2.53	512 (M+H)+
190		LCMS-A-1	1.87	473 (M+H)+
191		LCMS-C-1	2.35	515 (M+H)+
192		LCMS-C-1	2.65	554 (M+H)+
193		LCMS-C-1	2.37	526 (M+H)+
194		LCMS-C-1	2.60	556 (M+H)+
195		LCMS-C-1	2.75	553 (M+H)+


(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
196		LCMS-C-1	2.48	543 (M+H)+
197		LCMS-A-1	2.19	489 (M+H)+
198		LCMS-C-1	2.48	572 (M+H)+
199		LCMS-A-1	1.89	475 (M+H)+
200		LCMS-A-1	2.09	475 (M+H)+
201		LCMS-A-1	2.19	570 (M+H)+
202		LCMS-C-1	2.44	573 (M+H)+
203		LCMS-A-1	2.33	585 (M+H)+

[0431] The aryl bromide reagent used in the synthesis of Compound 191 ((S)-3-(4-bromo-indol-1-yl)-propane-1,2-diol)


was synthesized as follows.

(Reaction 25-3)



[0432] NaH (382 mg, 9.55 mmol, 60% oily suspension) was added to a solution of 4-bromo-indole (1.0 ml, 7.97 mmol) and (R)-(-)-2,2-dimethyl-1,3-dioxolan-4-ylmethyl p-toluenesulfonate (2.74 g, 9.57 mmol) in dimethylformamide (20 ml) at 0°C. The mixture was stirred at 0°C for two hours and at room temperature for 18 hours. NaH (190 mg, 4.75 mmol, 60% oily suspension) was further added, and the mixture was stirred at room temperature for six hours. The reaction mixture was diluted with AcOEt, and the organic layer was then washed with water ($\times 2$), dried over sodium sulfate and concentrated under reduced pressure. The resulting 4-bromo-1-((S)-2,2-dimethyl-1,3-dioxolan-4-ylmethyl)-1H-indole was used in the next step without further purification.


(Reaction 25-4)



[0433] A 2 N aqueous HCl solution (15 ml) was added to a solution of the above mixture (4-bromo-1-((S)-2,2-dimethyl-1,3-dioxolan-4-ylmethyl)-1H-indole) in tetrahydrofuran (30 ml), and the mixture was stirred at room temperature for eight hours. The reaction mixture was concentrated, and the residue was then diluted with AcOEt. This organic layer was sequentially washed with water ($\times 2$) and a saturated aqueous NaCl solution, and then dried and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (CH_2Cl_2 -MeOH) to give (S)-3-(4-bromo-indol-1-yl)-propane-1,2-diol as a colorless solid (2.05 g, 95%).
MS (ESI) m/z = 270, 272 ($M+H$) $^+$.

[0434] The aryl bromide reagent used in the synthesis of Compound 192 (4-bromo-1-(2-morpholin-4-yl-ethyl)-1H-indole) was synthesized as follows.

(Reaction 25-5)




[0435] 4-Bromo-1-(2-morpholin-4-yl-ethyl)-1H-indole was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 309, 311 ($M+H$) $^+$.

[0436] The aryl bromide reagent used in the synthesis of Compound 193 (2-(4-bromo-indol-1-yl)-N,N-dimethylaceta-

mide) was synthesized as follows.

(Reaction 25-6)




[0437] 2-(4-Bromo-indol-1-yl)-N,N-dimethyl-acetamide was synthesized by operations similar to those in Reaction 25-3, Reaction 23-2 and Reaction 10-18 using appropriate reagents, solvent and starting material.

MS (ESI) $m/z = 281, 283 (M+H)^+$.

[0438] The aryl bromide reagent used in the synthesis of Compound 194 (3-(4-bromo-3-trifluoromethyl-phenyl)-1,1-dimethyl-urea) was synthesized as follows.

(Reaction 25-7)




[0439] 3-(4-Bromo-3-trifluoromethyl-phenyl)-1,1-dimethyl-urea was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

MS (ESI) $m/z = 311, 313 (M+H)^+$.

[0440] The aryl bromide reagent used in the synthesis of Compound 195 (cyclopropanecarboxylic (4-bromo-3-trifluoromethyl-phenyl)-amide) was synthesized as follows.

(Reaction 25-8)




[0441] Cyclopropanecarboxylic (4-bromo-3-trifluoromethylphenyl)-amide was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

MS (ESI) $m/z = 308, 310 (M+H)^+$.

[0442] The aryl bromide reagent used in the synthesis of Compound 196 (N-(4-bromo-3-trifluoromethyl-phenyl)-2-

hydroxy-acetamide) was synthesized as follows.

(Reaction 25-9)




[0443] N-(4-Bromo-3-trifluoromethyl-phenyl)-2-hydroxy-acetamide was synthesized by operations similar to those in Reaction 2-3 and Reaction 23-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 298, 300 (M+H)^+$.

[0444] The aryl bromide reagent used in the synthesis of Compound 197 ((3-bromo-4-methyl-phenyl)-carbamic acid methyl ester) was synthesized as follows.


(Reaction 25-10)



[0445] Methyl chloroformate (0.202 ml, 2.62 mmol) was added to a solution of 3-bromo-4-methyl-phenylamine (243 mg, 1.31 mmol) in pyridine (2 ml), and the mixture was stirred at room temperature overnight. H_2O was added to the reaction mixture, followed by extraction with AcOEt (x 2). The organic layers were combined and sequentially washed with H_2O and saturated brine, and then dried over Na_2SO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel chromatography (n-hexane/AcOEt) to give (3-bromo-4-methyl-phenyl)-carbamic acid methyl ester (288 mg, 90%).

$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 2.34 (3H, s), 3.77 (3H, s), 6.51 (1H, br. s), 7.14 (1H, d, $J = 7.2$ Hz), 7.20 (1H, dd, $J = 7.4, 2.0$ Hz), 7.63 (1H, d, $J = 2.0$ Hz).


[0446] The aryl bromide reagent used in the synthesis of Compound 198 (1-(4-bromo-3-trifluoromethyl-phenyl)-3-(2-hydroxy-ethyl)-urea) was synthesized as follows.

(Reaction 25-11)

[0447] p-Nitrophenyl chloroformate (437 mg, 2.17 mmol) was added to a solution of 4-bromo-3-trifluoromethyl-aniline (400 μ l, 1.67 mmol) and pyridine (202 μ l, 2.50 mmol) in CH₂Cl₂ (6.2 ml) at 0°C. The mixture was stirred at 0°C for one hour, and 2-amino-ethanol (150 μ l, 2.50 mmol) was then added, followed by further stirring at 0°C for two hours. Triethylamine (210 μ l, 1.51 mmol) was added to the mixture, and the mixture was stirred at 0°C for one hour. 1 N HCl was added to the reaction mixture, followed by extraction with CH₂Cl₂/AcOEt. The organic layer was washed with water ($\times$ 2), and then dried over sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (CH₂Cl₂-MeOH) to give 1-(4-bromo-3-trifluoromethyl-phenyl)-3-(2-hydroxy-ethyl)-urea as a white powder (520 mg, 73%).

MS (ESI) m/z = 327, 329 (M+H)⁺.


[0448] The aryl bromide reagent used in the synthesis of Compound 199 (2-(3-bromo-4-methyl-phenylamino)-ethanol) was synthesized as follows.

(Reaction 25-12)

[0449] Triethylamine (0.28 mL, 2.00 mmol) and bromoethanol (0.14 mL, 1.98 mmol) were added to a solution of 3-bromo-4-methyl-phenylamine (240 mg, 1.29 mmol) in toluene (2 ml). The mixture was stirred at 100°C overnight and H₂O was then added, followed by extraction with AcOEt ($\times$ 2). The organic layers were combined and sequentially washed with H₂O and saturated brine, and then dried over Na₂SO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane/AcOEt) to give 2-(3-bromo-4-methyl-phenylamino)-ethanol (185 mg, 62%).

¹H-NMR (270 MHz, CDCl₃) δ 1.68 (1H, br, OH), 2.27 (3H, s, Me), 3.26 (2H, dd, J = 5.3, 5.1 Hz), 3.82 (2H, dd, J = 5.3, 5.1 Hz), 3.90 (1H, br, NH), 6.52 (1H, dd, J = 8.2, 2.5 Hz), 6.85 (1H, d, J = 2.5 Hz), 7.01 (1H, d, J = 8.2 Hz). MS (ESI) m/z = 230, 232 (M+H)⁺.


[0450] The aryl bromide reagent used in the synthesis of Compound 200 (2-(2-bromo-3-methyl-phenylamino)-ethanol) was synthesized as follows.

(Reaction 25-13)

[0451] 2-(2-Bromo-3-methyl-phenylamino)-ethanol was synthesized by operations similar to those in Reaction 25-12 using appropriate reagents and starting material.

MS (ESI) m/z = 230, 232 (M+H)+.


[0452] The aryl bromide reagent used in the synthesis of Compound 201 (2-(4-bromo-3-trifluoromethyl-phenylamino)-N,N-dimethyl-acetamide) was synthesized as follows.

(Reaction 25-14)

[0453] 2-(4-Bromo-3-trifluoromethyl-phenylamino)-N,N-dimethyl-acetamide was synthesized by operations similar to those in Reaction 25-12, Reaction 14-1 and Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 325, 327 (M+H)+.


[0454] The aryl bromide reagent used in the synthesis of Compound 202 (4-bromo-1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-3-fluoro-1H-indole) was synthesized as follows.

(Reaction 25-15)

[0455] Diethylaminotrifluorosulfur (1.5 mL, 11.06 mmol) was added to a solution of 4-bromo-1H-indole-2,3-dione (1.0 g, 4.4 mmol) in dichloromethane (44 mL) at 0°C. The mixture was stirred at room temperature for 54 hours and then quenched with methanol-water. The organic layer and the aqueous layer were separated, and the aqueous layer was then extracted with dichloromethane. The organic layers were combined, dried over sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-bromo-3,3-difluoro-1,3-dihydro-indol-2-one as a yellow solid (559 mg, 51%).

MS (ESI) m/z = 246 (M-H)-.

(Reaction 25-16)




[0456] Synthesis of a 1.3 M solution of BH_3F in tetrahydrofuran (Reagent A): Boron trifluoride etherate (2 mL) was added dropwise to a suspension of sodium borohydride (340 mg, 4.5 mmol) in tetrahydrofuran (12 mL) at 0°C . The mixture was stirred for 90 minutes to give Reagent A.

[0457] Reagent A (2.85 mL, 3.709 mmol) was added dropwise to a solution of 4-bromo-3,3-difluoro-1,3-dihydro-indol-2-one (400 mg, 1.61 mmol) in tetrahydrofuran (8.1 mL) at 0°C . The mixture was stirred at 0°C for 3.5 hours and at room temperature for 16 hours. Further, Reagent A (3.0 mL) was added to the reaction mixture, followed by stirring at room temperature for three hours. The reaction mixture was quenched with 3 M HCl (4.8 mL) and then extracted with ethyl acetate ($\times 2$). The organic layers were combined and sequentially washed with water and saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-bromo-3-fluoro-1H-indole as a yellow oil (132 mg, 38%).

MS (ESI) m/z = 212 (M-H)-.

(Reaction 25-17)




[0458] 4-Bromo-1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-3-fluoro-1H-indole was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.33 (3H, s), 1.40 (3H, s), 3.65 (1H, dd, J = 5.9, 8.8 Hz), 4.04 (1H, dd, J = 6.1, 8.8 Hz), 4.14 (2H, t, J = 4.9 Hz), 4.37-4.42 (1H, m), 7.02-7.07 (2H, m), 7.23-7.27 (2H, m).

[0459] The aryl bromide reagent used in the synthesis of Compound 203 ((4-bromo-3-methyl-phenyl)-(3,3,4,4-tetrafluoro-pyrrolidin-1-yl)-methanone) was synthesized as follows.

(Reaction 25-18)



[0460] (4-Bromo-3-methyl-phenyl)-(3,3,4,4-tetrafluoro-pyrrolidin-1-yl)-methanone was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.


MS (ESI) m/z = 340, 342 (M+H)+.

Example 26

N-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-acetamide
(Compound 204)

[0461]

(Reaction 26-1)



[0462] A mixture of 2-cyclohexyl-8-ethenesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (100.0 mg, 0.307 mmol), N-(4-bromo-3,5-dimethyl-phenyl)-acetamide (112. mg, 0.461 mmol), palladium(II) acetate (10 mg, 0.0461 mmol), tris(o-tolyl)phosphine (28 mg, 0.0922 mmol), triethylamine (0.128 ml, 0.922 mmol) and DMA (1.5 ml) was added to a sealed test tube in an N₂ atmosphere. This mixture was heated with stirring at 130°C for 13.5 hours. Palladium(II) acetate (10 mg, 0.0461 mmol), tris(o-tolyl)phosphine (28 mg, 0.0922 mmol) and triethylamine (0.128 ml, 0.922 mmol) were further added to the reaction mixture at room temperature in an N₂ atmosphere, and the mixture was heated with stirring at 130°C for 14 hours. The reaction mixture was cooled and water was then added. The aqueous layer was extracted with ethyl acetate (× 3). The organic layers were combined and sequentially washed with water (× 2) and saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (CH₂Cl₂-MeOH) to give N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-acetamide (61.4 mg, 41%).

¹H-NMR (400 MHz, CD₃OD) δ 1.20-1.57 (6H, m), 1.59-1.79 (3H, m), 1.80-2.00 (6H, m), 2.12 (3H, s), 2.39 (6H, s), 3.20-3.40 (2H, m), 3.58-3.75 (2H, m), 6.58 (1H, d, J = 16 Hz), 7.35 (2H, s), 7.57 (1H, d, J = 16 Hz). MS (ESI) m/z = 487 (M+H)⁺.

[0463] The example compounds shown below were synthesized by operations similar to those in Example 26 using appropriate reagents and starting materials.




(Compounds 205 to 208)

[0464]

[Table 32]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
205		LCMS-C-1	2.87	599 (M+H) ⁺

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
206		LCMS-C-1	2.38	520 (M+H)+
207		LCMS-C-1	2.23	530 (M+H)+
208		LCMS-C-1	2.12	530 (M+H)+

[0465] The aryl bromide reagent used in the synthesis of Compound 204 (N-(4-bromo-3,5-dimethyl-phenyl)-acetamide) was synthesized as follows.

(Reaction 26-2)




[0466] Benzyltrimethylammonium tribromide (BTMA-Br₃) (7.8 g, 20.21 mmol) was added to a solution of N-(3,5-dimethylphenyl)-acetamide (3.0 g, 18.38 mmol) in CH₂Cl₂/MeOH (90 ml/90 ml) at room temperature in an Ar atmosphere. The reaction mixture was stirred at room temperature for 10 minutes. The reaction mixture was concentrated under reduced pressure, and CH₂Cl₂ was then added to the resulting residue. The organic layer was washed with H₂O, and then dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane:EtOAc= 1:1) to give N-(4-bromo-3,5-dimethyl-phenyl)-acetamide (4.0 g, yield 90%).

MS (ESI+) m/z = 242, 244 (M+H)+.

[0467] The aryl bromide reagent used in the synthesis of Compound 205 ((4-bromo-3-methyl-benzyl)-carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 26-3)




[0468] NaBH₄ (1.45 g, 38.25 mmol) was added in small portions to a mixture of 4-bromo-3-methyl-benzonitrile (2.50 g, 12.8 mmol), NiCl₂ (1.65 g, 12.8 mmol) and BoC₂O (5.57 g, 25.5 mmol) in anhydrous MeOH (130 ml) at 0°C. The reaction mixture was stirred at room temperature for two hours and then concentrated under reduced pressure. Ethyl

acetate and water were added to the resulting residue, and the mixture was filtered through celite. The organic layer and the aqueous layer were separated, and the aqueous layer was then extracted with ethyl acetate. The organic layers were combined, dried over Na_2SO_4 and then concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate = 1/0 $\rightarrow$ 4/1) to give (4-bromo-3-methyl-benzyl)-carbamic acid tert-butyl ester as a white solid (2.42 g, 63%).

MS (ESI) m/z = 322 ($M+\text{Na}$) $^+$.

[0469] The aryl bromide reagent used in the synthesis of Compound 206 ((R)-3-(4-bromo-3,5-dimethyl-phenoxy)-propane-1,2-diol) was synthesized as follows.


(Reaction 26-4)



[0470] A mixture of 4-bromo-3,5-dimethyl-phenol (500 mg, 2.49 mmol), (R)-(-)-2,2-dimethyl-1,3-dioxolan-4-ylmethyl p-toluenesulfonate (856 mg, 2.99 mmol) and K_2CO_3 (1.03 g, 7.45 mmol) in dimethylformamide (5 ml) was stirred at 100°C for two hours. The reaction mixture was diluted with AcOEt, and the organic layer was then washed with water ($\times 2$), dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (AcOEt-hexane) to give (S)-4-(4-bromo-3,5-dimethyl-phenoxy)-2,2-dimethyl-1,3-dioxolane as a colorless solid (749 mg, 95%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.40 (3H, s), 1.46 (3H, s), 2.37 (6H, s), 3.85-3.92 (2H, m), 3.98-4.03 (1H, m), 4.13-4.18 (1H, m), 4.42-4.48 (1H, m), 6.66 (2H, s).

(Reaction 26-5)




[0471] (R)-3-(4-Bromo-3,5-dimethyl-phenoxy)-propane-1,2-diol was synthesized by operations similar to those in Reaction 25-4 using appropriate reagents and starting material.

MS (ESI) m/z = 275, 277 ($M+\text{H}$) $^+$.

[0472] The aryl bromide reagent used in the synthesis of Compound 207 (5-[(4-bromo-3-methyl-phenylamino)-methyl]-oxazolidin-2-one) was synthesized as follows.

(Reaction 26-6)




[0473] 5-[(4-Bromo-3-methyl-phenylamino)-methyl]-oxazolidin-2-one was synthesized by operations similar to those in Reaction 25-12 using appropriate reagents and starting material.

MS (ESI) m/z = 285, 287 ($M+\text{H}$) $^+$.

[0474] The aryl bromide reagent used in the synthesis of Compound 208 (2-(4-bromo-phenyl)-N-(2-dimethylaminoethyl)acetamide) was synthesized as follows.

(Reaction 26-7)




[0475] 2-(4-Bromo-phenyl)-N-(2-dimethylaminoethyl)acetamide was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.
MS (ESI) m/z = 285, 287 ($M+H$)⁺.

Example 27

3-[(E)-2-[2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzonitrile (Compound 209)

[0476]

(Reaction 27-1)




[0477] 3-[(E)-2-[2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzonitrile was synthesized by operations similar to those in Reaction 25-1 and Reaction 25-2 using appropriate reagents and starting material.
MS (ESI) m/z = 441 ($M+H$)⁺.

Example 28

8-[(E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 210)

[0478]

(Reaction 28-1)



[0479] 8-[(E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-1 and Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) m/z = 441 (M+H)+.

[0480] The example compounds shown below were synthesized by operations similar to those in Example 28 using appropriate reagents and starting materials.



(Compounds 211 to 214)

[0481]

[Table 33]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
211		LCMS-C-2	2.02	591 (M+H)+
212		LCMS-B-1	2.25	575 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
213		LCMS-C-1	2.37	557 (M+H)+
214		LCMS-C-1	2.42	579 (M+H)+

[0482] The aryl bromide reagent used in the synthesis of Compound 210 (4-(4-bromo-3-methyl-phenoxy)-butane-1,2-diol) was synthesized as follows.

(Reaction 28-2)




[0483] 4-(4-Bromo-3-methyl-phenoxy)-butane-1,2-diol was synthesized by operations similar to those in Reaction 26-4 and Reaction 25-4 using appropriate reagents and starting material.

MS (ESI) m/z = 275, 277 (M+H)+.

[0484] The aryl bromide reagent used in the synthesis of Compound 211 (2-(4-bromo-3-trifluoromethyl-phenylamino)-ethanol) was synthesized as follows.

(Reaction 28-3)




[0485] 2-(4-Bromo-3-trifluoromethyl-phenylamino)-ethanol was synthesized by operations similar to those in Reaction 12-1 using appropriate reagents and starting material.

MS (ESI) m/z = 284, 286 ($M+H$)⁺.

[0486] The aryl bromide reagent used in the synthesis of Compound 212 ((4-bromo-3-methyl-phenyl)-pyrrolidin-1-yl-methanone) was synthesized as follows.

(Reaction 28-4)




[0487] (4-Bromo-3-methyl-phenyl)-pyrrolidin-1-yl-methanone was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 268, 270 ($M+H$)⁺.

[0488] The aryl bromide reagent used in the synthesis of Compound 213 (4-bromo-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 28-5)




[0489] A 28% aqueous NH₃ solution (2.0 ml) was added to a solution of 4-bromo-3-methyl-benzenesulfonyl chloride (250 mg, 0.927 mmol) in THF (2.0 ml) at 0°C. The mixture was stirred at 0°C for 6.5 hours. The reaction mixture was quenched with 1 N HCl and extracted with CH₂Cl₂. The organic layer was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (CH₂Cl₂-AcOEt) to give 4-bromo-3-methyl-benzenesulfonamide as a white powder (126 mg, 54%).

MS (ESI) m/z = 272, 274 ($M+Na$)⁺.

[0490] The aryl bromide reagent used in the synthesis of Compound 214 (4-bromo-N-(2-hydroxy-ethyl)-3,N-dimethyl-benzamide) was synthesized as follows.

(Reaction 28-6)



[0491] 4-Bromo-N-(2-hydroxy-ethyl)-3,N-dimethyl-benzamide was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 272, 274 ($M+H$)⁺.


Example 29

N-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phe-

nyl)-acetamide (Compound 215)

[0492]

(Reaction 29-1)



[0493] N-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) m/z = 535 (M+H)+.

[0494] The example compounds shown below were synthesized by operations similar to those in Example 29 using appropriate reagents and starting materials.






(Compounds 216 to 223)

[0495]

[Table 34]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
216		LCMS-C-1	2.55	604 (M+H)+
217		LCMS-C-1	2.70	639 (M+H)+
218		LCMS-C-1	2.57	670 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
219		LCMS-C-1	1.95	634 (M+H)+
220		LCMS-C-1	2.42	577 (M+H)+
221		LCMS-C-1	2.42	591 (M+H)+
222		LCMS-C-1	2.45	593 (M+H)+
223		LCMS-C-1	2.50	676 (M+H)+

[0496] The aryl bromide reagent used in the synthesis of Compound 216 ((4-bromo-3-methyl-phenyl)-(4-methyl-piperazin-1-yl)-methanone) was synthesized as follows.

(Reaction 29-2)






[0497] (4-Bromo-3-methyl-phenyl)-(4-methyl-piperazin-1-yl)-methanone was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 297, 299 (M+H)+.


[0498] The aryl bromide reagents used in the synthesis of Compounds 217 to 219 were synthesized by operations similar to those in Reaction 28-5 using appropriate reagents and starting materials.

[Table 35]

Target Compound	Aryl bromide reagent	Aryl bromide reagent MS (m/z)
217		354, 355 (M+Na)+
218		363, 365 (M+H)+
219		327, 329 (M+H)+

[0499] The aryl bromide reagent used in the synthesis of Compound 220 ((R)-1-(4-bromo-3-methyl-phenyl)-4-hydroxy-pyrrolidin-2-one) was synthesized as follows.


(Reaction 29-3)



[0500] A mixture of 2-bromo-5-iodotoluene (500 mg, 1.68 mmol), (R)-4-hydroxy-pyrrolidinone (204 mg, 2.02 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (58.5 mg, 0.101 mmol), tris(dibenzylideneacetone)-dipalladium(0)-chloroform adduct (35.0 mg, 0.034 mmol) and cesium carbonate (769 mg, 2.36 mmol) in 1,4-dioxane (degassed, 5 ml) was stirred at 110°C overnight in a nitrogen stream. The reaction mixture was treated with H_2O and extracted with AcOEt (x 2). The organic layers were combined and sequentially washed with H_2O and saturated brine, and then dried over Na_2SO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel chromatography (n-hexane/AcOEt) to give (R)-1-(4-bromo-3-methyl-phenyl)-4-hydroxy-pyrrolidin-2-one as a pale brown solid (173 mg, 38%).

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 2.29 (1H, d, $J = 17.1$ Hz), 2.34 (3H, s), 2.82 (1H, dd, $J = 17.1, 6.4$ Hz), 3.57 (1H, d, $J = 10.3$ Hz), 4.01 (1H, dd, $J = 10.3, 4.9$ Hz), 4.36-4.40 (1H, m), 5.35 (1H, d, $J = 3.4$ Hz, OH), 7.51 (1H, dd, $J = 8.8, 2.4$ Hz), 7.55 (1H, d, $J = 8.8$ Hz), 7.62 (1H, br. s). MS (ESI) $m/z = 270, 272$ (M+H)+.


[0501] The aryl bromide reagent used in the synthesis of Compound 221 ((R)-1-(4-bromo-3-methyl-phenyl)-5-hydroxymethyl-pyrrolidin-2-one) was synthesized as follows.

(Reaction 29-4)

[0502] (R)-1-(4-Bromo-3-methyl-phenyl)-5-hydroxymethyl-pyrrolidin-2-one was synthesized by operations similar to those in Reaction 29-3 using appropriate reagents and starting material.

MS (ESI) m/z = 284, 286 (M+H)⁺.


[0503] The aryl bromide reagent used in the synthesis of Compound 222 ((R)-3-(4-bromo-3-methyl-phenyl)-5-hydroxymethyl-oxazolidin-2-one) was synthesized as follows.

(Reaction 29-5)

[0504] A mixture of (S)-3-(4-bromo-3-methyl-phenylamino)-propane-1,2-diol (202 mg, 0.777 mmol), diethyl carbonate (3 ml), sodium methoxide (28% in MeOH, 0.160 ml) and MeOH (4 ml) was stirred at 130°C overnight. The reaction mixture was treated with saturated NH₄Cl and H₂O and extracted with AcOEt (× 2). The organic layers were combined and sequentially washed with H₂O and saturated brine, and then dried over Na₂SO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane/AcOEt) to give (R)-3-(4-bromo-3-methyl-phenyl)-5-hydroxymethyl-oxazolidin-2-one (170 mg, 77%).


¹H-NMR (400 MHz, DMSO-d₆) δ 2.35 (3H, s), 3.53-3.57 (1H, m), 3.65-3.68 (1H, m), 3.82 (1H, dd, J = 8.8, 6.4 Hz), 4.06 (1H, dd, J = 9.3, 8.8 Hz), 4.67-4.72 (1H, m), 5.22 (1H, br. s), 7.42 (1H, dd, J = 8.8, 2.9 Hz), 7.54 (1H, d, J = 2.5 Hz), 7.56 (1H, d, J = 8.8 Hz). MS (ESI) m/z = 286, 288 (M+H)⁺.

[0505] The aryl bromide reagent used in the synthesis of Compound 223 ((4-bromo-3,5-dimethyl-phenyl)-[3-(3-dimethylamino-propoxy)-azetidin-1-yl]-methanone) was synthesized as follows.

(Reaction 29-6)

[0506] (4-Bromo-3,5-dimethyl-phenyl)-[3-(3-hydroxy-azetidin-1-yl)-methanone was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 284, 286 (M+H)⁺.


(Reaction 29-7)

[0507] NaH (110 mg, 2.75 mmol, 60% oily suspension) and NaI (274 mg, 1.83 mmol) were added to a solution of (4-bromo-3,5-dimethyl-phenyl)-(3-hydroxy-azetidin-1-yl)-methanone (130 mg, 0.458 mmol) and (3-chloro-propyl)-dimethyl-amine (289 mg, 1.83 mmol) in toluene (1.8 ml). The mixture was stirred at 110°C for 15 hours. The reaction mixture was diluted with AcOEt, and the organic layer was then sequentially washed with a saturated aqueous NaHCO₃ solution, water and a saturated aqueous NaCl solution. Further, the organic layer was dried over sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane/AcOEt) to give (4-bromo-3,5-dimethyl-phenyl)-[3-(3-dimethylamino-propoxy)-azetidin-1-yl]-methanone (46 mg, 27%).

MS (ESI) m/z = 369, 371 (M+H)⁺.

Example 30

N-(3-Hydroxy-propyl)-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide (Compound 224)

[0508](Reaction 30-1)

[0509] N-(3-Hydroxy-propyl)-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide was synthesized by operations similar to those in Reaction 25-1 and Reaction 25-2 using appropriate reagents and starting material.


MS (ESI) m/z = 645 (M+H)⁺.

[0510] The example compound shown below was synthesized by operations similar to those in Example 30 using appropriate reagents and starting material.

(Compound 225)


[0511]

[Table 36]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
225		LCMS-D-1	3.3	567 (M+H)+

[0512] The aryl bromide reagent used in the synthesis of Compound 224 (4-bromo-N-(3-hydroxy-propyl)-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 30-2)




[0513] 4-Bromo-N-(3-hydroxy-propyl)-3-methyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 322, 324 (M+H)+.

[0514] The aryl bromide reagent used in the synthesis of Compound 225 (N-(3-bromo-4-methyl-phenyl)-N-methyl-acetamide) was synthesized as follows.

(Reaction 30-3)



[0515] N-(3-Bromo-4-methyl-phenyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.


MS (ESI) m/z = 242, 244 (M+H)+.

Example 31

8-[(E)-2-(3-Hydroxy-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 226)

[0516]

(Reaction 31-1)



[0517] 8-[(E)-2-(3-Hydroxy-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 510$ (M+H) $^+$.

[0518] The example compounds shown below were synthesized by operations similar to those in Example 31 using appropriate reagents and starting materials.

(Compounds 227 to 239)

[0519]


[Table 37]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
227		HPLC-A-2	11.5	510 (M+H) $^+$
228		LCMS-A-1	2.12	648 (M+H) $^+$
229		LCMS-D-1	3.3	601 (M+H) $^+$
230		LCMS-A-1	2.25	621 (M+H) $^+$
231		LCMS-C-1	2.60	565 (M+H) $^+$

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
232		LCMS-C-1	2.52	595 (M+H)+
233		LCMS-C-1	2.57	567 (M-1)-
234		LCMS-C-1	2.73	652 (M+1)+
235		LCMS-C-1	2.67	595 (M+1)+
236		LCMS-D-1	3.1	565 (M+H)+
237		LCMS-C-1	2.63	565 (M+H)+
238		HPLC-A-2	12.8	565 (M+H)+
239		HPLC-A-2	14.0	579 (M+H)+


[0520] The aryl bromide reagent used in the synthesis of Compound 230 ((4-bromo-3-methyl-phenyl)-(4-hydroxy-piperidin-1-yl)-methanone) was synthesized as follows.

(Reaction 31-2)

[0521] (4-Bromo-3-methyl-phenyl)-(4-hydroxy-piperidin-1-yl)-methanone was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 298, 300 ($M+H$) $^+$.


[0522] The aryl bromide reagent used in the synthesis of Compound 231 (4-bromo-3,N,N-trimethyl-benzamide) was synthesized as follows.

(Reaction 31-3)

[0523] 4-Bromo-3,N,N-trimethyl-benzamide was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 264, 266 ($M+H$) $^+$.


[0524] The aryl bromide reagent used in the synthesis of Compound 232 (N-(4-bromo-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide) was synthesized as follows.

(Reaction 31-4)

[0525] N-(4-Bromo-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 12-5 using appropriate reagents and starting material.

MS (ESI) m/z = 272, 274 ($M+H$) $^+$.

[0526] The aryl bromide reagent used in the synthesis of Compound 233 (4-bromo-3-fluoro-N,N-dimethyl-benzamide) was synthesized as follows.


(Reaction 31-5)

[0527] 4-Bromo-3-fluoro-N,N-dimethyl-benzamide was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 246, 248 ($M+H$) $^+$.

[0528] The aryl bromide reagent used in the synthesis of Compound 234 (4-(4-bromo-2,5-dichloro-phenoxy)-butane-1,2-diol) was synthesized as follows.

(Reaction 31-6)




[0529] 4-(4-Bromo-2,5-dichloro-phenoxy)-butane-1,2-diol was synthesized by operations similar to those in Reaction 26-4 and Reaction 31-6 using appropriate reagents and starting material.

MS (ESI) m/z = 351, 353 ($M+H$) $^+$.

[0530] The aryl bromide reagent used in the synthesis of Compound 235 ([3-(4-bromo-3-methyl-phenoxy)-propyl]-dimethyl-amine) was synthesized as follows.

(Reaction 31-7)




[0531] 3-Dimethylamino-propan-1-ol (251 μ L, 2.14 mmol) and DEAD (973 μ L, 2.14 mmol) were added to a solution of 4-bromo-3-methyl-phenol (200 mg, 1.07 mmol) and PPh_3 (561 mg, 2.14 mmol) in THF (10 mL) at 0°C. The mixture was stirred for two hours and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give [3-(4-bromo-3-methyl-phenoxy)-propyl]-dimethyl-amine (176 mg, 61%).

MS (ESI) m/z = 273, 275 ($M+H$) $^+$.

[0532] The aryl bromide reagent used in the synthesis of Compound 236 (N-(3-bromo-2-methyl-phenyl)-N-methyl-acetamide) was synthesized as follows.

(Reaction 31-8)




[0533] N-(3-Bromo-2-methyl-phenyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 242, 244 ($M+H$)⁺.

[0534] The aryl bromide reagent used in the synthesis of Compound 237 (N-(4-bromo-3-methyl-phenyl)-N-methyl-acetamide) was synthesized as follows.

(Reaction 31-9)




[0535] N-(4-Bromo-3-methyl-phenyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 264, 266 ($M+H$)⁺.

[0536] The aryl bromide reagent used in the synthesis of Compound 239 (N-(4-bromo-3,5-dimethyl-phenyl)-N-methyl-acetamide) was synthesized as follows.

(Reaction 31-10)




[0537] N-(4-Bromo-3,5-dimethyl-phenyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 256, 258 ($M+H$)⁺.

Example 32

2-Cycloheptyl-8-[(E)-2-[1-[(S)-2,3-dihydroxy-propyl]-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 240)

[0538]

(Reaction 32-1)

[0539] 2-Cycloheptyl-8-[(E)-2-[1-[(S)-2,3-dihydroxy-propyl]-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14, Reaction 1-4, Reaction 11-4, Reaction 25-1 and Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) m/z = 529 (M+H)+.

[0540] The example compound shown below was synthesized by operations similar to those in Example 32 using appropriate reagents and starting material.

(Compound 241)

[0541]

[Table 38]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS(m/z)
241		LCMS-A-1	1.90	551 (M+H)+

Example 33

8-[(E)-2-[4-[(R)-2,3-Dihydroxy-propoxy]-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 242)

[0542]

(Reaction 33-1)



[0543] 8-((E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-1 and Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 600$ (M+H)+.

[0544] The example compounds shown below were synthesized by operations similar to those in Example 33 using appropriate reagents and starting materials.

(Compounds 243 to 246)

[0545]

[Table 39]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
243		LCMS-A-1	2.36	600 (M+H)+
244		LCMS-C-1	2.52	637 (M+H)+
245		LCMS-A-1	2.92	562 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
246		LCMS-E-4	2.82	557 (M+H)+

[0546] The aryl bromide reagent used in the synthesis of Compound 244 ((4-bromo-3,5-dimethyl-phenyl)-(4-hydroxy-piperidin-1-yl)-methanone) was synthesized as follows.

(Reaction 33-2)



[0547] (4-Bromo-3,5-dimethyl-phenyl)-(4-hydroxy-piperidin-1-yl)-methanone was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.


MS (ESI) m/z = 312, 314 (M+H)+.

Example 34

8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(2-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 247)

[0548]

(Reaction 34-1)



[0549] 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(2-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-1 and Reaction

26-1 using appropriate reagents and starting material.




MS (ESI) $m/z = 600$ (M+H)+.

[0550] The example compounds shown below were synthesized by operations similar to those in Example 34 using appropriate reagents and starting materials.

(Compounds 248 to 250)

[0551]

[Table 40]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
248		LCMS-C-1	2.62	593 (M+H)+
249		LCMS-B-1	1.96	577 (M+H)+
250		LCMS-B-1	1.85	543 (M+H)+

Example 35

2-Cyclohexyl-8-[(E)-4-(1H-indol-4-yl)-but-3-ene-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 251)

[0552]

(Reaction 35-1)




20 **[0553]** 2-Cyclohexyl-8-[(E)-4-(1H-indol-4-yl)-but-3-ene-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-1 and Reaction 25-2 using appropriate reagents and starting material. MS (ESI) m/z = 469 (M+H)+.

Example 36

25 2-Cyclohexyl-8-[(E)-5-(1H-indol-4-yl)-pent-4-ene-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 252)

[0554]

(Reaction 36-1)




55 **[0555]** 2-Cyclohexyl-8-[(E)-5-(1H-indol-4-yl)-pent-4-ene-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-1 and Reaction 25-2 using appropriate reagents and starting material. MS (ESI) m/z = 483 (M+H)+.

Example 37

2-Cyclohexyl-8-[(E)-3-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-prop-2-ene-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 253)

[0556]

(Reaction 37-1)

[0557] 2-Cyclohexyl-8-[(E)-3-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-prop-2-ene-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-1, Reaction 25-2 and Reaction 25-4 using appropriate reagents and starting material.


MS (ESI) m/z = 529 ($\text{M}+\text{H}$) $^+$.

Example 38

8-[(E)-2-[3-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 254)

[0558]

(Reaction 38-1)



[0559] 8-((E)-2-[3-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 31-7, Reaction 26-1 and Reaction 25-4 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 1.7 (2H, d, $J = 13.73$ Hz), 1.97 (1H, brs), 2.00 (2H, m), 2.18 (2H, dt, $J = 3.05, 13.73$ Hz), 2.30 (3H, s), 2.51 (1H, brs), 3.35 (2H, dt, $J = 3.05, 11.83$ Hz), 3.59 (1H, m), 3.75 (1H, m), 3.80 (2H, d, $J = 11.83$ Hz), 4.06 (1H, brs), 4.18 (2H, m), 6.66 (1H, d, $J = 15.64$ Hz), 6.95 (1H, d, $J = 7.63$ Hz), 7.17 (1H, t, $J = 7.63$ Hz), 7.21 (1H, d, $J = 7.63$ Hz), 7.42 (1H, d, $J = 8.01$ Hz), 7.54 (1H, dd, $J = 7.63, 8.01$ Hz), 7.75 (1H, d, $J = 7.63$ Hz), 7.81 (1H, s), 7.81 (1H, d, $J = 15.64$ Hz), 9.75 (1H, s). MS (ESI) $m/z = 598$ (M+H)+.

[0560] The example compound shown below was synthesized by operations similar to those in Example 38 using appropriate reagents and starting material.

(Compound 255)

[0561]


[Table 41]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
255		HPLC-A-3	11.56	598 (M+H)+


Example 39

8-((E)-2-[3-(2-Hydroxy-ethoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 256)

[0562]

(Reaction 39-1)

[0563] 8-((E)-2-((3-(2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy)-2-methyl-phenyl)-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material. This compound was used as such in the next step without purification.

(Reaction 39-2)

[0564] Tetrabutylammonium fluoride (0.11 ml, 0.11 mmol, 1 M in THF) was added to a solution of 8-((E)-2-((3-(2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy)-2-methyl-phenyl)-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one obtained above in anhydrous THF (1 ml) at room temperature in an Ar atmosphere. The mixture was stirred at room temperature for two hours and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane:EtOAc = 1:1) to give 8-((E)-2-((3-(2-hydroxy-ethoxy)-2-methyl-phenyl)-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (3.2 mg, yield in two steps: 48%).

¹H-NMR (300 MHz, DMSO-d₆) δ 11.75 (1H, s), 7.99 (1H, d, J = 7.5 Hz), 7.90 (1H, s), 7.63 (3H, m), 7.37 (1H, d, J = 7.5 Hz), 7.23 (2H, m), 7.06 (1H, d, J = 7.9 Hz), 4.85 (1H, t, J = 5.6 Hz), 4.00 (2H, m), 3.74 (2H, m), 3.60 (2H, m), 3.20 (2H, m), 2.27 (3H, s), 1.88 (2H, m), 1.63 (2H, m). MS (ESI+) m/z = 554 (M+H)⁺.

[0565] The example compounds shown below were synthesized by operations similar to those in Example 39 using appropriate reagents and starting materials.

(Compounds 257 to 258)


[0566]

[Table 42]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
257		LCMS-D-1	3.1	595 (M+H)+
258		LCMS-D-1	3.10	621 (M+H)+

[0567] The aryl bromide reagent used in the synthesis of Compound 257 (N-(3-bromo-4-methyl-phenyl)-N-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-acetamide) was synthesized as follows.


(Reaction 39-3)



[0568] N-(3-Bromo-4-methyl-phenyl)-N-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.
MS (ESI) m/z = 386, 388 (M+H)+.

[0569] The aryl bromide reagent used in the synthesis of Compound 258 ((R)-1-(4-bromo-3,5-dimethyl-phenyl)-5-(tert-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one) was synthesized as follows.

(Reaction 39-4)




[0570] (R)-1-(4-Bromo-3,5-dimethyl-phenyl)-5-(tert-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one was synthesized by operations similar to those in Reaction 29-3 using appropriate reagents and starting material.

¹H-NMR (300 MHz, CDCl₃) δ -0.06 (3H, s), -0.03 (3H, s), 0.86 (9H, s), 2.10 (1H, m), 2.26 (1H, m), 2.40 (6H, s), 2.48 (1H, ddd, J = 4.6, 10.3, 16.8 Hz), 2.68 (1H, ddd, J = 8.0, 9.9, 17.9 Hz), 3.56 (2H, dq, J = 3.8, 10.7 Hz), 4.15 (1H, m), 7.10 (2H, s).

Example 40

N-(4-((E)-2-[2-(2-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide (Compound 259)

[0571]

(Reaction 40-1)


[0572] N-(4-((E)-2-[2-(2-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 12-5 using appropriate reagents and starting material.

MS (ESI) m/z = 597 (M+H)⁺.

Example 41


N-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-piperidin-4-yl-acetamide (Compound 260)

[0573]


(Reaction 41-1)

[0574] Acetic acid (4.9 eq) and sodium triacetoxyborohydride (2.0 eq) were sequentially added to a solution of 4-bromo-3-methylaniline (246 mg, 1.32 mmol) and 1-(tert-butoxycarbonyl)-4-piperidone (350 mg, 1.76 mmol) in 1,2-dichloroethane (10 ml). The mixture was stirred at room temperature for 3.5 hours and then quenched with a saturated aqueous sodium carbonate solution. The reaction mixture was extracted with dichloromethane, and the organic layer was then concentrated under reduced pressure to 4-(4-bromo-3-methyl-phenylamino)-piperidine-1-carboxylic acid tert-butyl ester as a white solid (586 mg, 100%). This compound was used in the next step without further purification.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.32 (m, 2H), 1.42 (s, 9H), 2.01 (d, $J = 13.2$ Hz, 2H), 2.31 (s, 3H), 2.92 (t, $J = 11.6$ Hz, 2H), 3.45 (br, 2H), 4.04 (br, 1H), 6.32 (dd, $J = 2.4$ Hz, 8.4 Hz, 1H), 6.48 (d, $J = 2.8$ Hz, 1H), 7.28 (m, 1H). MS (ESI) $m/z = 369$ ($\text{M}+\text{H}$) $^+$.

(Reaction 41-2)


[0575] 4-[Acetyl-(4-bromo-3-methyl-phenyl)-aminol-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 12-2 using the compound obtained above as a starting material. MS (ESI) $m/z = 411, 413$ ($\text{M}+\text{H}$) $^+$.

(Reaction 41-3)

[0576] N-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-piperidin-4-yl-acetamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 4-1 using appropriate reagents and starting material. MS (ESI) $m/z = 634$ ($\text{M}+\text{H}$) $^+$.

Example 42

2-Cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 261)

[0577]**(Reaction 42-1)**

[0578] 10% Pd-C (28 mg) was added to a solution of 2-cyclohexyl-8-[(E)-2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (56.6 mg, 0.123 mmol) in MeOH-DMF (4 ml, 1:1). The mixture was stirred at room temperature overnight in a hydrogen atmosphere. The reaction mixture was filtered through celite, and the filtrate was then concentrated under reduced pressure. The resulting residue was diluted with ethyl acetate, and the organic layer was then washed with water (x 2), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, AcOEt-hexane) to give 2-cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a colorless foam (46.9 mg, 83%).
¹H-NMR (400 MHz, CDCl₃) δ 1.21-1.45 (6H, m), 1.50-1.60 (2H, m), 1.65-1.85 (4H, m), 1.90-1.96 (2H, m), 2.38-2.48 (1H, m), 3.25-3.40 (6H, m), 3.65-3.73 (2H, m), 7.01 (2H, d, J = 8.0 Hz), 7.13 (1H, t, J = 8 Hz), 8.59 (1H, brs), 9.03 (1H, brs).
 MS (ESI) m/z = 461 (M+H)+.

[0579] The example compounds shown below were synthesized by operations similar to those in Example 42 using appropriate reagents and starting materials.

(Compounds 262 to 267)

[0580]

[Table 43]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
262		LCMS-E-5	3.3	485 (M+H)+
263		LCMS-E-4	2.89	471 (M+H)+
264		LCMS-D-1	3.1	567 (M+H)+
265		LCMS-D-1	3.3	567 (M+H)+
266		HPLC-A-3	11.35	600 (M+H)+
267		HPLC-A-3	11.03	600 (M+H)+

Example 43

8-{2-[4-((R)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 268)

[0581]

(Reaction 42-2)



[0582] The following reaction was performed by utilizing a continuous-flow hydrogenation reactor H-Cube® Type HC-2 (ThalesNano Nanotechnology Inc.).

[0583] 8-{(E)-2-[4-((R)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethenesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (28.5 mg, 48.3 μmol) was dissolved in EtOH/DMF 4:1 (concentration 10 mg/ml). The mixture was allowed to pass through 10% Pd/C (CatCart™) at a flow rate of 2 ml/min under the conditions of 30 bar and 40°C in a hydrogen atmosphere, and was subjected to hydrogenation reaction. The resulting reaction solution was concentrated under reduced pressure. The residue was purified by preparative TLC ($\text{CH}_2\text{Cl}_2/\text{MeOH} = 20:1$) to give 8-{2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a white powder (12.9 mg, 45%).

$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 1.60-1.63 (2H, m), 1.83-1.89 (2H, m), 1.97-2.04 (1H, m), 2.12-2.22 (1H, m), 2.32 (3H, s), 2.28-2.36 (1H, m), 2.52-2.59 (1H, m), 2.97-3.01 (2H, m), 3.29-3.40 (6H, m), 3.67-3.70 (2H, m), 4.24-4.29 (1H, m), 4.80 (1H, t, $J = 5.4$ Hz), 7.22-7.29 (2H, m), 7.79 (1H, br t, $J = 7.8$ Hz), 7.98 (1H, br. d, $J = 7.8$ Hz), 8.29 (1H, br. d, $J = 7.3$ Hz), 8.33 (1H, br. s), 11.81 (1H, br. s). MS (ESI) $m/z = 593$ (M+H)+.

[0584] The example compounds shown below were synthesized by operations similar to those in Example 43 using appropriate reagents and starting materials.








(Compounds 269 to 302)

[0585]








[Table 44]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
269		LCMS-A-1	2.07	443 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
270		LCMS-C-1	2.55	572 (M+H)+
271		LCMS-C-1	2.67	555 (M+H)+
272		LCMS-C-1	2.87	601 (M+H)+
273		LCMS-C-1	2.53	558 (M+H)+
274		LCMS-C-3	1.23	491 (M+H)+
275		LCMS-C-3	1.06	477 (M+H)+
276		LCMS-C-3	0.86	475 (M+H)+








(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
277		LCMS-C-1	2.73	418 (M+H)+
278		LCMS-C-1	2.42	574 (M+H)+
279		LCMS-C-1	2.37	522 (M+H)+
280		LCMS-C-1	2.33	448 (M+H)+
281		LCMS-B-1	1.45	532 (M+H)+
282		LCMS-C-1	2.12	532 (M+H)+
283		LCMS-C-1	2.70	587 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
284		LCMS-C-2	1.85	584 (M+H)+
285		LCMS-A-1	2.23	623 (M+H)+
286		LCMS-C-2	2.03	593 (M+H)+
287		LCMS-B-1	2.10	647 (M+H)+
288		LCMS-C-1	2.67	577 (M+H)+
289		LCMS-C-1	2.55	606 (M+H)+
290		LCMS-C-1	2.65	641 (M+H)+
291		LCMS-C-1	2.35	559 (M+H)+

(continued)


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
292		LCMS-C-1	2.52	672 (M+H)+
293		LCMS-C-1	2.55	571 (M+H)+
294		LCMS-A-1	2.19	636 (M+H)+
295		LCMS-C-1	2.67	597 (M+H)+
296		LCMS-C-1	2.45	595 (M+H)+
297		LCMS-B-1	1.91	579 (M+H)+
298		LCMS-C-1	2.52	537 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
299		LCMS-C-1	2.47	567 (M+H)+
300		LCMS-A-1	2.32	583 (M+H)+
301		LCMS-A-1	2.28	568 (M+H)+
302		LCMS-B-1	1.61	489 (M+H)+

Example 44

2-Cyclohexyl-8-[2-(1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 303)

[0586](Reaction 44-1)

[0587] 2-Cyclohexyl-8-[2-(1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 42-1 using appropriate reagents and starting material. MS (ESI) $m/z = 443$ (M+H)+.

[0588] The example compounds shown below were synthesized by operations similar to those in Example 44 using appropriate reagents and starting materials.










(Compounds 304 to 320)

[0589]


[Table 45]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
304		LCMS-A-1	2.92	550 (M+H)+
305		LCMS-A-1	2.60	521 (M+H)+
306		LCMS-A-1	2.15	457 (M+H)+
307		LCMS-A-1	2.92	562 (M+H)+
308		LCMS-A-1	2.89	535 (M+H)+
309		LCMS-C-1	2.38	517 (M+H)+
310		LCMS-C-1	2.52	514 (M+H)+
311		LCMS-E-7	1.53	471 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
312		LCMS-C-1	2.57	521 (M+H)+
313		LCMS-A-1	1.65	514 (M+H)+
314		LCMS-C-1	2.48	487 (M+H)+
315		HPLC-A-1	13.4	486 (M+H)+
316		HPLC-A-2	11.6	432 (M+H)+
317		LCMS-D-1	1.8	434 (M+H)+
318		LCMS-D-1	3.0	434 (M+H)+
319		LCMS-C-1	2.87	460 (M+H)+
320		LCMS-A-1	1.49	455 (M+H)+

[0590] The aryl bromide reagent used in the synthesis of Compound 313 ([2-(4-bromo-indol-1-yl)-ethyl]-dimethylamine) was synthesized as follows.

(Reaction 44-2)


[0591] [2-(4-Bromo-indol-1-yl)-ethyl]-dimethyl-amine was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 267, 269 (M+H)+.

Example 45

2-Cyclohexyl-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 321)

[0592]

(Reaction 45-1)

[0593] 2-Cyclohexyl-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 42-2 using appropriate reagents and starting material.


MS (ESI) m/z = 527 (M+H)+.

[0594] The example compound shown below was synthesized by operations similar to those in Example 45 using appropriate reagents and starting material.

(Compound 322)


[0595]

[Table 46]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
322		LCMS-C-1	2.48	585 (M+H)+

[0596] The aryl bromide reagent used in the synthesis of Compound 321 ((S)-3-(4-bromo-3-methyl-phenylamino)-propane-1,2-diol) was synthesized as follows.

(Reaction 45-2)




[0597] (S)-3-(4-Bromo-3-methyl-phenylamino)-propane-1,2-diol was synthesized by operations similar to those in Reaction 26-4 and Reaction 25-4 using appropriate reagents and starting material.
MS (ESI) m/z = 260, 262 (M+H)+.

Example 46

2-Cyclohexyl-8-[2-(2-trifluoromethyl-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 323)

[0598]

(Reaction 46-1)



[0599] 2-Cyclohexyl-8-[2-(2-trifluoromethyl-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 42-1 using appropriate reagents and starting material.

MS (ESI) m/z = 472 (M+H)+.

[0600] The example compound shown below was synthesized by operations similar to those in Example 46 using appropriate reagents and starting material.

(Compound 324)

[0601]


[Table 47]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
324		LCMS-D-1	2.7	489 (M+H)+

Example 47

8-{2-[4-((R)-2,3-Dihydroxy-propoxy)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 325)


[0602]

(Reaction 47-1)

[0603] 8-{2-[4-((R)-2,3-Dihydroxy-propoxy)-2-methylphenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 42-2 using appropriate reagents and starting material.

MS (ESI) m/z = 586 ($\text{M}+\text{H}$)⁺.

[0604] The aryl bromide reagent used in the synthesis of Compound 325 ((R)-3-(4-bromo-3-methyl-phenoxy)-propane-1,2-diol) was synthesized as follows.

(Reaction 47-2)

[0605] (R)-3-(4-Bromo-3-methyl-phenoxy)-propane-1,2-diol was synthesized by operations similar to those in Reaction 26-4 and Reaction 25-4 using appropriate reagents and starting material.


MS (ESI) m/z = 283, 285 ($\text{M}+\text{Na}$)⁺.

Example 48

N-(2-Hydroxy-ethyl)-N-(4-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide (Compound 326)

[0606]

(Reaction 48-1)




15 **[0607]** N-(2-Hydroxy-ethyl)-N-(4-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide was synthesized by operations similar to those in Reaction 42-1 and Reaction 39-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 597$ (M+H)+.

20 Example 49

N-(2-Hydroxy-ethyl)-N-(2-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide (Compound 327)


25 **[0608]**

(Reaction 49-1)



55 **[0609]** N-(2-Hydroxy-ethyl)-N-(2-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide was synthesized by operations similar to those in Reaction 26-1, Reaction 42-1 and Reaction 39-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 597$ (M+H)+.


[0610] The aryl bromide reagent used in the synthesis of Compound 327 (N-(3-bromo-2-methyl-phenyl)-N-[2-(tert-butyl-dimethyl-silyloxy)-ethyl]-acetamide) was synthesized as follows.

(Reaction 49-2)

[0611] N-(3-Bromo-2-methyl-phenyl)-N-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.
MS (ESI) m/z = 386, 388 (M+H)⁺


Example 50

8-[3-(3-Amino-phenyl)-propane-1-sulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 328)

[0612](Reaction 50-1)

[0613] N-(3-(3-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-propyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 42-1 using appropriate reagents and starting material.

MS (ESI) m/z = 475 (M+H)⁺.

(Reaction 50-2)


[0614] Conc. HCl (0.5 ml) was added to a solution of N-{3-[3-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-propyl]-phenyl}-acetamide (5.0 mg, 0.0105 mmol) in MeOH (1 ml) at room temperature. The mixture was stirred at 30 to 40°C for four hours. The reaction mixture was concentrated under reduced pressure, and the resulting residue was then purified by preparative TLC (CH₂Cl₂:MeOH = 10:1) to give 8-[3-(3-amino-phenyl)-propane-1-sulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (2.5 mg, yield 55%).

Example 51

N,N-Dimethyl-4-{5-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-benzamide (Compound 329)


[0615]

(Reaction 51-1)



[0616] 8-(5-Bromo-thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material. MS (ESI) *m/z* = 523 (M+H)+.

(Reaction 51-2)



[0617] A mixture of 8-(5-bromo-thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (11.1 mg, 0.0212 mmol), 4-(N,N-dimethylaminocarbonyl)phenylboronic acid (8.0 mg, 0.041 mmol), Pd (PPh₃)₄ (3.8 mg, 0.0033 mmol) and Na₂CO₃ (22.0 mg, 0.208 mmol) in toluene (0.12 ml)-EtOH (0.12 ml)-H₂O (0.12 ml) was stirred at 85°C for 20 hours in a sealed test tube in an N₂ atmosphere. The reaction mixture was cooled to room temperature and extracted with AcOEt. The organic layer was washed with a saturated aqueous NH₄Cl solution, and then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (AcOEt) to give N,N-dimethyl-4-{5-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-benzamide (9.7 mg, 77%).








¹H-NMR (300 MHz) (CDCl₃) δ 1.74 (2H, br d, *J* = 13.5 Hz), 2.21 (2H, ddd, *J* = 13.5, 11.0, and 4.0 Hz), 3.04 (3H, br s), 3.15 (3H, br s), 3.24 (2H, ddd, *J* = 11.5, 11.0, and 3.0 Hz), 3.82 (2H, ddd, *J* = 11.5, 4.0 and 4.0 Hz), 7.38 (1H, d, *J* = 4.0 Hz), 7.50 (2H, d, *J* = 8.4 Hz), 7.57 (1H, d, *J* = 4.0 Hz), 7.60 (1H, t, *J* = 8.1 Hz), 7.66 (2H, d, *J* = 8.4 Hz), 7.78 (1H, d, *J* = 8.1 Hz), 7.99 (1H, d, *J* = 8.1 Hz), 8.12 (1H, s), 9.61 (1H, br s). MS (ESI) *m/z* = 591 (M+H)+.

[0618] The example compounds shown below were synthesized by operations similar to those in Example 51 using appropriate reagents and starting materials.


(Compounds 330 to 337)

[0619]

[Table 48]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
330		LCMS-C-2	2.08	591 (M+H)+
331		LCMS-C-2	2.15	605 (M+H)+
332		LCMS-C-2	1.92	529 (M+H)+
333		LCMS-C-2	2.20	623 (M+H)+
334		LCMS-C-2	2.20	623 (M+H)+
335		LCMS-D-1	3.37	615 (M+H)+
336		LCMS-D-1	3.35	615 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
337		LCMS-D-1	3.27	616 (M+H)+

[0620] The aryl boronate reagent used in the synthesis of Compounds 331, 333, 335, 336 and 337 (3,N,N-trimethyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide) was synthesized as follows.

(Reaction 51-3)




[0621] A mixture of 4-bromo-3,N,N-trimethylbenzamide (203 mg, 0.838 mmol), 1,1'-bis(diphenylphosphino)-ferrocene (dppf) (27.9 mg, 0.0503 mmol), PdCl₂(dppf)-CH₂Cl₂ (41.6 mg, 0.0509 mmol), AcOK (245 mg, 2.50 mmol) and bis(pinacolato)diboron (286 mg, 1.13 mmol) in dioxane (5.5 ml) was stirred at 85°C for six hours in a sealed test tube in an N₂ atmosphere. The reaction mixture was cooled to room temperature and extracted with AcOEt. The organic layer was washed with water, and then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to give 3,N,N-trimethyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide (135 mg, 56%).

MS (ESI) m/z = 290 (M+H)+.

[0622] The aryl boronate reagent used in the synthesis of Compound 334 (4,N,N-trimethyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide) was synthesized as follows.

(Reaction 51-4)




[0623] 4,N,N-Trimethyl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzamide was synthesized by operations similar to those in Reaction 51-3 using appropriate reagents and starting material.

MS (ESI) m/z = 242 (M+H)+.

[0624] The aryl boronate reagent used in the synthesis of Compound 332 (N-[3-methyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetamide) was synthesized as follows.

(Reaction 51-5)



[0625] N-[3-Methyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-acetamide was synthesized by operations similar to those in Reaction 51-3 using appropriate reagents and starting material.

MS (ESI) m/z = 276 (M+H)+.

[0626] The following aryl bromide reagents used in the synthesis of Compounds 332, 333, 334, 335, 336 and 337 were synthesized by operations similar to those in Reaction 51-1 using appropriate reagents and starting materials.

[Table 49]


Target Compound	Aryl bromide	MS
332		460, 462 (M+H)+
333 334		540, 542 (M+H)+
335		532, 534 (M+H)+
336		532, 534 (M+H)+
337		533, 535 (M+H)+

Example 52

2-Cyclohexyl-8-{2-[2-(3,5-dimethyl-isoxazol-4-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 338)

[0627]


(Reaction 52-1)



[0628] Triphenylphosphine (7.83 g, 29.8 mmol) and carbon tetrabromide (12.4 g, 37.3 mmol) were added to a solution of 2-(2-bromophenyl)ethanol (5.00 g, 24.9 mmol) in dichloromethane (123 mL). The mixture was stirred at room temperature for 15 hours, and a saturated aqueous sodium carbonate solution was then added. The organic layer and the aqueous layer were separated, and the organic layer was then concentrated under reduced pressure. The resulting residue was triturated with ethyl acetate:n-hexane (1:4, 200 mL) and then filtered, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give 1-bromo-2-(2-bromoethyl)benzene as a colorless oil (6.23 g, 95%).

$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 3.30 (2H, t, $J = 7.6$ Hz), 3.60 (2H, t, $J = 7.3$ Hz), 7.10-7.17 (1H, m), 7.26-7.28 (2H, m), 7.55 (1H, d, $J = 8.1$ Hz).


(Reaction 52-2)



[0629] A solution of 1-bromo-2-(2-bromoethyl)benzene (6.23 g, 23.6 mmol) in ethanol (20.5 mL) was added to a solution of sodium sulfite (3.12 g, 24.7 mmol) in water (25.0 mL). The mixture was heated at 100°C for 24 hours. The reaction mixture was filtered, and the filtrate was then left to stand at 3°C overnight. The resulting white crystals were collected by filtration and dried to give sodium 2-(2-bromo-phenyl)ethanesulfonate (4.00 g, 59%).


$^1\text{H-NMR}$ (270 MHz, $\text{d}_6\text{-DMSO}$) δ 2.60-2.67 (2H, m), 2.94-3.00 (2H, m), 7.09-7.15 (1H, m), 7.25-7.33 (2H, m), 7.55 (1H, d, $J = 8.6$ Hz).

(Reaction 52-3)



[0630] N,N -Dimethylformamide (4.2 mL) and thionyl chloride (5.1 mL, 69.7 mmol) were sequentially added to a suspension of sodium 2-(2-bromo-phenyl)ethanesulfonate (4.00 g, 13.9 mmol) in toluene. The mixture was stirred at 100°C for 66 hours and then poured into ice water. The organic layer and the aqueous layer were separated, and the aqueous layer was extracted with ether. The organic layers were combined and sequentially washed with water and saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure to give 2-(2-bromophenyl)ethanesulfonyl chloride (4.10 g). This was used in the next step without further purification.

(Reaction 52-4)



[0631] 2-Cyclohexyl-8-{2-[2-(3,5-dimethyl-isoxazol-4-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 5-4 and Reaction 21-1 using appropriate reagents and starting material.

MS (ESI) m/z = 499 (M+H)+.

[0632] The example compounds shown below were synthesized by operations similar to those in Example 52 using appropriate reagents and starting materials.

(Compounds 339 to 340)

[0633]


[Table 50]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
339		LCMS-E-5	4.26	511 (M+H)+
340		LCMS-E-5	3.73	484 (M+H)+


Example 53

8-{5-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-thiophene-2-sulfonyl}-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 341)

[0634]

(Reaction 53-1)

[0635] 2-{4-[2-(2,2-Dimethyl-[1,3]dioxolan-4-yl)-ethoxy]-2-methyl-phenyl}-4,4,5,5-tetramethyl-[1,3,2]dioxaborolane was synthesized by operations similar to those in Reaction 51-3 using appropriate reagents and starting material. MS (ESI) $m/z = 257$ ($\text{M}-\text{C}_3\text{H}_6\text{O}+\text{H}$)⁺.


(Reaction 53-2)

[0636] 8-{5-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-thiophene-2-sulfonyl}-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 51-2 and Reaction 25-4 using appropriate reagents and starting material. MS (ESI) $m/z = 656$ ($\text{M}+\text{H}$)⁺.


Example 54

N-(4-{5-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide (Compound 342)

[0637]

(Reaction 54-1)

[0638] Acetic acid 2-{acetyl-[3-methyl-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-amino}-ethyl ester was synthesized by operations similar to those in Reaction 51-3 using appropriate reagents and starting material.
MS (ESI) $m/z = 315 (M+H)^+$.

(Reaction 54-2)


[0639] N-(4-{5-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 51-2 and Reaction 12-5 using appropriate reagents and starting material.
MS (ESI) $m/z = 653 (M+H)^+$.

Example 55

2-Cyclohexyl-8-((E)-2-thiazol-2-yl-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 343)

[0640]


(Reaction 55-1)



[0641] 2-Cyclohexyl-8-methanesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 314$ ($M+H$)⁺.

(Reaction 55-2)



[0642] LHMDS (1.1 ml, 1.11 mmol) was added to a solution of 2-cyclohexyl-8-methanesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (100 mg, 0.32 mmol) in THF (3 ml) at -20°C in an N_2 atmosphere. The mixture was stirred at -20°C for 30 minutes, and diethyl chlorophosphate (48 μl , 0.34 mmol) was then added. Further, the mixture was stirred at -20°C for 60 minutes, and 2-thiazolecarboxaldehyde (31 μl , 0.35 mmol) was then added. The reaction mixture was stirred at room temperature for one hour, and ethyl acetate (10 ml) and an aqueous NH_4Cl solution (5 ml) were then added. The organic layer and the aqueous layer were separated, and the aqueous layer was then extracted with ethyl acetate (10 ml). The organic layers were combined and washed with saturated brine, and then dried over Na_2SO_4 and concentrated under reduced pressure. The resulting residue was recrystallized from ethyl acetate to give 2-cyclohexyl-8-((E)-2-thiazol-2-yl-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (87 mg, yield 67%).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.38 (1H, s), 7.94 (1H, d, $J = 3.4$ Hz), 7.57 (1H, d, $J = 15.3$ Hz), 7.51 (1H, d, $J = 3.3$ Hz), 7.11 (1H, d, $J = 15.3$ Hz), 3.75 (2H, m), 3.31 (2H, m), 2.46-2.36 (1H, m), 2.07-1.97 (2H, m), 1.92-1.88 (2H, m), 1.83-1.80 (2H, m), 1.75-1.50 (4H, m), 1.50-1.20 (4H, m).

[0643] The example compound shown below was synthesized by operations similar to those in Example 55 using appropriate reagents and starting material.

(Compound 344)

[0644]


[Table 51]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
344		HPLC-A-1	13.5	394 ($M+H$) ⁺

Example 56

2-Cyclohexyl-8-(2-thiazol-2-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 345)

[0645]




[0646] 2-Cyclohexyl-8-(2-thiazol-2-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 42-1 using appropriate reagents and starting material.

¹H-NMR (300 MHz, CDCl₃) δ 9.04 (1H, s), 7.70 (1H, d, J = 3.5 Hz), 7.25 (1H, d, J = 3.0 Hz), 3.75 (2H, m), 3.57-3.44 (4H, m), 3.37 (2H, m), 2.46-2.38 (1H, m), 2.01-1.90 (4H, m), 1.84-1.70 (3H, m), 1.56-1.50 (2H, m), 1.48-1.26 (5H, m). MS (ESI) m/z = 411 (M+H)⁺.

Example 57

2-Cyclohexyl-8-[3-(4-methoxy-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 346)

[0647]




[0648] 2-Cyclohexyl-8-[3-(4-methoxy-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 42-1 using appropriate reagents and starting material. MS (ESI) m/z = 448 (M+H)⁺.

Example 58

N-Benzyl-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-benzamide (Compound 347)

[0649]

(Reaction 58-1)



[0650] N-Benzyl-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-benzamide was synthesized by operations similar to those in Reaction 5-4, Reaction 23-2 and Reaction 10-18 using appropriate reagents and starting material.


MS (ESI) $m/z = 509$ (M+H)+.

Example 59

8-(3-Chloro-benzenesulfonyl)-2-[3-(morpholine-4-carbonyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 348)

[0651]

(Reaction 59-1)



[0652] 8-(3-Chloro-benzenesulfonyl)-2-[3-(morpholine-4-carbonyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-18 using appropriate reagents and starting material. MS (ESI) $m/z = 517$ (M+H)+.

[0653] The example compounds shown below were synthesized by operations similar to those in Example 59 using appropriate reagents and starting materials.

(Compounds 349 to 351)

[0654]


[Table 52]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
349		LCMS-E-2	4.11	537 (M+H)+
350		LCMS-E-6	1.76	537 (M+H)+
351		LCMS-E-6	1.42	475 (M+H)+

Example 60

8-{2-[4-(4-Methanesulfonyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 352)

[0655]

(Reaction 60-1)

[0656] 8-{2-[4-(4-Methanesulfonyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1, Reaction

10

[0657] The example compounds shown below were synthesized by operations similar to those in Example 60 using appropriate reagents and starting materials.

[0658]

[Table 53]

15

20

25

30

35

40

45








50

55

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (<i>m/z</i>)
359		LCMS-A-1	2.45	670 (M+H)+
360		LCMS-C-1	2.58	699 (M+H)+
361		LCMS-C-1	2.78	669 (M+H)+
362		LCMS-C-1	2.47	680 (M+H)+
363		LCMS-C-1	2.53	705 (M+H)+
364		LCMS-C-1	2.72	675 (M+H)+
365		LCMS-A-1	2.67	627 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (<i>m/z</i>)
366		LCMS-A-1	1.95	650 (M+H)+
367		LCMS-C-1	2.42	620 (M+H)+
368		LCMS-A-1	2.81	649 (M+H)+
369		LCMS-C-1	2.42	620 (M+H)+
370		LCMS-C-1	2.42	593 (M+H)+
371		LCMS-C-1	2.55	607 (M+H)+
372		LCMS-C-1	2.57	620 (M+H)+

(continued)


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (<i>m/z</i>)
373		LCMS-A-1	2.08	648 (M+H)+
374		LCMS-C-1	2.43	677 (M+H)+
375		LCMS-A-1	2.79	686 (M+H)+
376		LCMS-A-1	2.03	634 (M+H)+
377		LCMS-C-1	2.40	579 (M+H)+
378		LCMS-B-1	2.16	595 (M+H)+
379		LCMS-B-1	2.15	581 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (<i>m/z</i>)
380		LCMS-B-1	2.38	591 (M+H)+
381		LCMS-B-1	2.12	563 (M+H)+
382		LCMS-C-1	2.52	663 (M+H)+

Example 61

3,5,N,N-Tetramethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide (Compound 383)

[0659]**(Reaction 61-1)**

[0660] 3,5,N,N-Tetramethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) *m/z* = 563 (M+H)+.

[0661] The example compounds shown below were synthesized by operations similar to those in Example 61 using appropriate reagents and starting materials.

(Compounds 384 to 393)

[0662]

[Table 54]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
384		LCMS-C-1	2.53	619 (M+H)+
385		LCMS-C-1	2.52	591 (M+H)+
386		LCMS-C-1	2.48	649 (M+H)+
387		LCMS-C-1	2.50	648 (M+H)+
388		LCMS-C-1	2.77	589 (M+H)+
389		LCMS-C-1	2.53	607 (M+H)+
390		LCMS-C-1	2.60	620 (M+H)+
391		LCMS-C-1	2.48	634 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
392		LCMS-C-1	2.28	606 (M+H)+
393		LCMS-C-1	2.62	646 (M+H)+

Example 62

8-[(E)-2-[4-(4-Acetyl-piperazine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 394)

[0663]**(Reaction 62-1)**

[0664] 8-[(E)-2-[4-(4-Acetyl-piperazine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 10-14 using appropriate reagents and starting material.
MS (ESI) m/z = 662 (M+H)+.

[0665] The example compounds shown below were synthesized by operations similar to those in Example 62 using appropriate reagents and starting materials.






(Compounds 395 to 408)

[0666]

[Table 55]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
395		LCMS-C-1	2.67	691 (M+H)+
396		LCMS-C-1	2.52	663 (M+H)+
397		LCMS-C-1	2.50	705 (M+H)+
398		LCMS-D-1	2.7	634 (M+H)+
399		LCMS-D-1	2.7	648 (M+H)+
400		LCMS-D-1	3.0	635 (M+H)+
401		LCMS-D-1	3.0	621 (M+H)+
402		LCMS-D-1	3.0	621 (M+H)+
403		LCMS-D-1	2.9	665 (M+H)+

(continued)


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
404		LCMS-D-1	2.7	620 (M+H)+
405		LCMS-D-1	2.7	662 (M+H)+
406		LCMS-D-1	2.7	679 (M+H)+
407		LCMS-D-1	2.7	707 (M+H)+
408		LCMS-D-1	3.0	607 (M+H)+

Example 63

2,N,N-Trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide (Compound 409)

[0667]

(Reaction 63-1)



[0668] 2,N,N-Trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide was synthesized by operations similar to those in Reaction 26-1, Reaction 42-1 and Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) $m/z = 567$ (M+H)+.

[0669] The example compounds shown below were synthesized by operations similar to those in Example 63 using appropriate reagents and starting materials.

(Compounds 410 to 411)

[0670]


[Table 56]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
410		LCMS-D-1	2.9	539 (M+H)+
411		LCMS-D-1	3.1	609 (M+H)+

Example 64

4,N,N-Trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide (Compound 412)

[0671]

(Reaction 64-1)

[0672] 4,N,N-Trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide was synthesized by operations similar to those in Reaction 26-1, Reaction 42-1 and Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) $m/z = 567$ (M+H)+.

[0673] The example compounds shown below were synthesized by operations similar to those in Example 64 using appropriate reagents and starting materials.

(Compounds 413 to 415)

[0674]

[Table 57]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
413		HPLC-A-1	11.5	597 (M+H)+
414		HPLC-A-1	9.4	311 (M+H)+
415		HPLC-A-2	9.5	553 (M+H)+

Example 65

8-(2-{4-[4-(2-Hydroxy-ethyl)-piperazine-1-carbonyl]-2-methyl-phenyl}-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 416)

[0675]

(Reaction 65-1)



[0676] 8-(2-{4-[4-(2-Hydroxy-ethyl)-piperazine-1-carbonyl]-2-methyl-phenyl}-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1, Reaction 42-1 and Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 652 (M+H)+.

[0677] The example compound shown below was synthesized by operations similar to those in Example 64 using appropriate reagents and starting material.

(Compound 417)

[0678]


[Table 58]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
417		LCMS-C-2	2.31	611 (M+H)+

Example 66


2-Fluoro-5,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide (Compound 418)

[0679]

(Reaction 66-1)


[0680] $n\text{-BuLi}$ (5.0 ml, 8.0 mmol, 1.6 M in hexane) was added dropwise to a solution of $n\text{-Bu}_2\text{Mg}$ (8.0 ml, 8.0 mmol, 1.0 M in heptane) at room temperature for 10 minutes. The mixture was stirred at room temperature for 15 minutes and then cooled to $-10 \pm 2^\circ\text{C}$. A solution of 2,5-dibromo-4-fluorotoluene (5.466 g, 19.59 mmol) in toluene (30 ml)-THF (6 ml) was added dropwise to this mixed reaction solution over 30 minutes, and the mixture was then stirred at 0°C for one hour. The reaction mixture was added dropwise to a solution cooled to -10°C of DMF (2.1 ml, 27 mmol) in toluene (7.6 ml) over 15 minutes. Further, this mixture was stirred at -10 to -5°C for 30 minutes, and then quenched with an aqueous citric acid solution (2.3 M, 16 ml, 37 mmol) and extracted with Et_2O . The organic layer was washed with water, and then dried over MgSO_4 and concentrated under reduced pressure to give 4-bromo-2-fluoro-5-methyl-benzaldehyde (3.74 g, 88%).

$^1\text{H-NMR}$ (300 MHz) (CDCl_3) δ 2.42 (3H, s), 7.42 (1H, d, $J = 9.6$ Hz), 7.72 (1H, d, $J = 7.2$ Hz), 10.29 (1H, s).

(Reaction 66-2)

[0681] NaH_2PO_4 (418 mg, 3.48 mmol) in H_2O (17.6 ml), a 35% aqueous H_2O_2 solution (2.5 ml, 25.7 mmol) and NaClO_2 (2.23 g, 24.7 mmol) in H_2O (34.9 ml) were sequentially added to a mixture of 4-bromo-2-fluoro-5-methyl-benzaldehyde (3.74 g, 17.2 mmol) in MeCN (52 ml) at 0°C . The mixture was stirred at room temperature for 14 hours, and then made acidic (pH 3) with a 10% aqueous HCl solution and extracted with ethyl acetate (3 x 100 ml). The organic layers were washed with H_2O (70 ml), and then dried over MgSO_4 and concentrated under reduced pressure to give 4-bromo-2-fluoro-5-methyl-benzoic acid as a pale orange solid (4.02 g, 100%).

$^1\text{H-NMR}$ (300 MHz) (CDCl_3) δ 2.42 (3H, s), 7.41 (1H, d, $J = 9.9$ Hz), 7.88 (1H, d, $J = 7.5$ Hz). MS (ESI) $m/z = 231$ (M-H) $^-$.


(Reaction 66-3)

[0682] (Trimethylsilyl)diazomethane (4.0 ml, 8.0 mmol, 2 M in Et_2O) was added dropwise to a solution of 4-bromo-2-fluoro-5-methyl-benzoic acid (1.88 g, 8.05 mmol) in benzene (7.5 ml)- MeOH (5.6 ml) at $10 \pm 2^\circ\text{C}$ over 10 minutes. The

mixture was stirred at room temperature for 30 minutes and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 40/1) to give methyl 4-bromo-2-fluoro-5-methyl-benzoate (1.62 g, 82%).

¹H-NMR (300 MHz) (CDCl₃) δ 2.40 (3H, s), 3.93 (3H, s), 7.37 (1H, d, J = 9.9 Hz), 7.80 (1H, d, J = 7.8 Hz).

(Reaction 66-4)



[0683] 2-Fluoro-5,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide was synthesized by operations similar to those in Reaction 10-2, Reaction 10-3, Reaction 10-4, Reaction 10-5, Reaction 5-4, Reaction 23-2 and Reaction 10-1 using appropriate reagents and starting material. MS (ESI) m/z = 569 (M+H)+.

[0684] The example compound shown below was synthesized by operations similar to those in Example 66 using appropriate reagents and starting material.

(Compound 419)

[0685]

[Table 59]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
419		LCMS-C-2	1.78	625 (M+H)+

Example 67

N-Benzyl-2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-1H-indol-3-yl]-acetamide (Compound 420)

[0686]

(Reaction 67-1)



[0687] N-Benzyl-2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-1H-indol-3-yl]-acetamide was synthesized by operations similar to those in Reaction 25-2, Reaction 42-1, Reaction 23-2 and Reaction 10-1 using appropriate reagents and starting material.


MS (ESI) $m/z = 590$ ($\text{M}+\text{H}$)⁺.

Example 68

2-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-N-methyl-benzamide (Compound 421)

[0688]

(Reaction 68-1)



[0689] 2-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec 1-ene-8-sulfonyl)-ethyl]-N-methyl-benzamide was synthesized by operations similar to those in Reaction 25-2, Reaction 42-1, Reaction 23-2 and Reaction 10-18 using appropriate reagents and starting material.

MS (ESI) $m/z = 461$ ($\text{M}+\text{H}$) $^+$.

[0690] The example compound shown below was synthesized by operations similar to those in Example 68 using appropriate reagents and starting material.

(Compound 422)

[0691]


[Table 60]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
422		LCMS-E-6	3.33	525 ($\text{M}+\text{H}$) $^+$

Example 69

8-(3-Chloro-benzenesulfonyl)-2-[1-(3,3-dimethyl-butyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 423)

[0692]

(Reaction 69-1)

[0693] 8-(3-Chloro-benzenesulfonyl)-2-[1-(3,3-dimethyl-butyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-18 using appropriate reagents and starting material. MS (ESI) $m/z = 509 (M+H)^+$.

[0694] The example compound shown below was synthesized by operations similar to those in Example 69 using appropriate reagents and starting material.

(Compound 424)

[0695]


[Table 61]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
424		LCMS-E-2	2.9	469 (M+H)+

Example 70

8-(3-Chloro-benzenesulfonyl)-2-[1-(4-chloro-benzoyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 425)

[0696]

(Reaction 70-1)

[0697] 8-(3-Chloro-benzenesulfonyl)-2-[1-(4-chlorobenzoyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.36-1.67 (m, 5 H), 1.84-2.06 (m, 4 H), 2.66-2.84 (m, 1 H), 2.90-3.10 (m, 2 H), 3.16-3.34 (m, 1 H), 3.41-3.55 (m, 1 H), 3.55-3.67 (m, 2 H), 4.04-4.27 (m, 1 H), 7.23-7.29 (m, 2 H), 7.30-7.37 (m, 2 H), 7.43 (t, $J = 7.83$ Hz, 1 H), 7.50-7.55 (m, 1 H), 7.59-7.65 (m, 1 H), 7.72 (t, $J = 1.77$ Hz, 1 H). MS (ESI) $m/z = 549 (M+H)^+$.

[0698] The example compound shown below was synthesized by operations similar to those in Example 70 using appropriate reagents and starting material.

(Compound 426)

[0699]


[Table 62]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
426		LCMS-E-2	3.82	545 (M+H)+

Example 72

2-[1-(1H-Indol-5-carbonyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 428)

[0700]

(Reaction 72-1)

[0701] 2-[1-(1H-Indol-5-carbonyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-18 using appropriate reagents and starting material.

MS (ESI) m/z = 598 (M+H)+.

[0702] The example compounds shown below were synthesized by operations similar to those in Example 72 using appropriate reagents and starting materials.

(Compounds 429 to 439)

[0703]

[Table 63]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
429		LCMS-E-3	3.5	587 (M+H)+
430		LCMS-E-3	3.67	613 (M+H)+


(continued)

Compound	Structure	LCMS or HPLC condition	Retentiontime (min)	MS (m/z)
431		LCMS-E-3	1.57	610 (M+H)+
432		LCMS-E-2	2.47	563 (M+H)+
433		LCMS-E-2	3.79	608 (M+H)+
434		LCMS-E-2	4.49	654 (M+H)+
435		LCMS-E-2	2.97	574 (M+H)+
436		LCMS-E-2	4.48	631 (M+H)+
437		LCMS-E-2	4.57	601 (M+H)+
438		LCMS-E-2	2.68	574 (M+H)+
439		LCMS-E-2	4.02	569 (M+H)+

Example 73

8-(2-Naphthalen-1-yl-ethanesulfonyl)-2-{1-[(E)-(3-phenyl-acryloyl)]-piperidin-3-yl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 441)

[0704](Reaction 73-1).



10 **[0705]** 8-(2-Naphthalen-1-yl-ethanesulfonyl)-2-{1-[(E)-(3-phenyl-acryloyl)]-piperidin-3-yl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.


MS (ESI) m/z = 585 (M+H) +.

15 Example 74

2-[1-(2-Amino-acetyl)-piperidin-3-yl]-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 442)

20 **[0706]**

(Reaction 74-1)




35 **[0707]** 2-[1-(2-Amino-acetyl)-piperidin-3-yl]-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-18 and Reaction 4-1 using appropriate reagents and starting material.

MS (ESI) m/z = 468 (M+H)+.

45 Example 75

8-{2-[2-Methyl-4-(3-methylamino-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 444)

50 **[0708]**

(Reaction 75-1)

[0709] 8-((E)-2-((4-(2-(3-(trifluoromethyl)phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one)-2-yl)ethanesulfonyl)-2-(3-(trifluoromethyl)phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 4-1 using appropriate reagents and starting material.

MS (ESI) m/z = 606 (M+H)+.

[0710] The example compounds shown below were synthesized by operations similar to those in Example 75 using appropriate reagents and starting materials.

(Compounds 445 to 446)

[0711]

[Table 64]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
445		LCMS-B-1	1.70	592 (M+H)+
446		LCMS-C-1	2.50	620 (M+H)+

Example 76

8-((E)-2-((4-(2-(4-(2-Hydroxy-acetyl)-piperazine-1-carbonyl)-2,6-dimethyl-phenyl)-ethanesulfonyl)-2-(3-(trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 447)

[0712]

(Reaction 76-1)



15 **[0713]** 8-((E)-2-{4-[4-(2-Hydroxy-acetyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl}-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 678 (M+H)⁺.

20 **[0714]** The example compounds shown below were synthesized by operations similar to those in Example 76 using appropriate reagents and starting materials.

(Compounds 448 to 449)

25 **[0715]**


[Table 65]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
30 448		LCMS-C-1	2.62	734 (M+H) ⁺
35 449		LCMS-C-1	2.43	733 (M+H) ⁺

45 Example 77

2-Methoxy-N-methyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide (Compound 450)


50 **[0716]**

(Reaction 77-1)

[0717] 2-Methoxy-N-methyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 10-18 using appropriate reagents and starting material.
MS (ESI) $m/z = 595$ (M+H)+.

Example 78

2-Hydroxy-N-methyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide (Compound 451)


[0718](Reaction 78-1)

[0719] 2-Hydroxy-N-methyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 2-3 and Reaction 12-5 using appropriate reagents and starting material.
MS (ESI) $m/z = 581$ (M+H)+.

Example 79

[(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoyl)-methyl-amino]-acetic acid (Compound 452)


[0720]

(Reaction 79-1)

[0721] [(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoyl)-methyl-amino]-acetic acid was synthesized by operations similar to those in Reaction 10-14 and Reaction 23-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 607 (M+H)^+$

Example 80

8-(3-Chloro-benzenesulfonyl)-2-[1-(3,3-dimethyl-butyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 453)

[0722](Reaction 80-1)

[0723] Triethylamine (2 eq), 3,3-dimethyl-butylaldehyde (1 eq) and sodium triacetoxyborohydride (1.5 eq) were added to a solution of 8-(3-chloro-benzenesulfonyl)-2-piperidin-3-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one trifluoroacetate (126 mg, 0.24 mmol) in dichloromethane (5 ml). The mixture was stirred at room temperature overnight and then concentrated under reduced pressure. The resulting residue was purified by HPLC to give 8-(3-chloro-benzenesulfonyl)-2-[1-(3,3-dimethyl-butyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (35 mg, yield 30%).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.85 (s, 9 H), 0.87-0.93 (m, 2 H), 1.43-1.56 (m, 4 H), 1.58-1.70 (m, 3 H), 1.75-1.99 (m, 6 H), 2.93-3.03 (m, 3 H), 3.54-3.63 (m, 3 H), 7.43 (t, $J = 7.83$ Hz, 1 H), 7.49-7.54 (m, 1 H), 7.62 (d, $J = 7.58$ Hz, 1 H), 7.72 (t, $J = 1.77$ Hz, 1 H). MS (ESI) $m/z = 495 (M+H)^+$.

Example 81

3-[8-(3-Chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butylamide (Compound 454)

[0724]

(Reaction 81-1)



[0725] A solution of triphosgene (345 mg, 1.16 mmol) in CH_2Cl_2 (7 ml) was added to a solution of tert-butylamine (331 μl , 3.14 mmol) and triethylamine (876 μl , 6.29 mmol) in CH_2Cl_2 (10 ml) at -78°C . The mixture was stirred at room temperature for 10 minutes, followed by addition of a solution of 8-(3-chloro-benzenesulfonyl)-2-piperidin-3-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one trifluoroacetate (165 mg, 0.314 mmol) and triethylamine (876 μl , 6.29 mmol) in CH_2Cl_2 (2 ml). Further, the reaction mixture was stirred at room temperature for 10 minutes and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 3-[8-(3-Chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butylamide as a colorless oil (40 mg, yield 25%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.25 (s, 9 H), 1.47-1.63 (m, 2 H), 1.71-1.84 (m, 2 H), 1.83-2.07 (m, 6 H), 3.02-3.13 (m, 2 H), 3.14-3.25 (m, 2 H), 3.53-3.65 (m, 2 H), 3.84-3.95 (m, 1 H), 7.43 (t, $J = 7.83$ Hz, 1 H), 7.49-7.55 (m, 1 H), 7.61 (d, $J = 7.83$ Hz, 1 H), 7.72 (t, $J = 1.77$ Hz, 1 H). MS (ESI) $m/z = 510$ ($\text{M}+\text{H}$) $^+$.

[0726] The example compound shown below was synthesized by operations similar to those in Example 81 using appropriate reagents and starting material.

(Compound 455)

[0727]


[Table 66]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
455		LCMS-E-2	3.34	602 ($\text{M}+\text{H}$) $^+$

Example 82

8-(3-Chloro-benzenesulfonyl)-2-[1-(piperidine-1-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 456)

[0728]


(Reaction 82-1)

[0729] TEA (1.143 mmol, 3 eq) and piperidine-1-carbonyl chloride (0.457 mmol, 1.2 eq) were sequentially added to a mixed solution of 8-(3-chloro-benzenesulfonyl)-2-piperidin-3-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one trifluoroacetate (200 mg, 0.38 mmol) in dichloromethane (3 ml). The resulting mixture was stirred at room temperature overnight and then concentrated under reduced pressure. The resulting residue was purified by HPLC to give 8-(3-chloro-benzenesulfonyl)-2-[1-(piperidine-1-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (30 mg, yield 15%).

¹H-NMR (400 MHz, CDCl₃) δ 1.34-1.48 (m, 2 H), 1.49-1.66 (m, 6 H), 1.66-1.81 (m, 2 H), 1.84-1.96 (m, 1 H), 1.97-2.12 (m, 2 H), 2.79-2.96 (m, 1 H), 2.98-3.25 (m, 8 H), 3.25-3.42 (m, 2 H), 3.62-3.89 (m, 3 H), 7.50 (t, J = 7.83 Hz, 1 H), 7.57-7.62 (m, 1 H); 7.69 (d, J = 7.58 Hz, 1 H), 7.80 (t, J = 1.77 Hz, 1 H). MS (ESI) m/z = 522 (M+H)⁺.

Example 83

3-(2-Dimethylamino-ethyl)-1-methyl-1-(3-methyl-4-{(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-urea (Compound 457)

[0730](Reaction 83-1)

[0731] 4-Nitrophenyl chloroformate (35 mg, 0.17 mmol) was added to a solution of 8-[(E)-2-(2-methyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (80 mg, 0.15 mmol) in THF (1 ml) at room temperature, and the mixture was then stirred at 70°C for 30 minutes. The reaction mixture was extracted with AcOEt, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting intermediate (83a) (20 mg, 0.029 mmol) was dissolved in DMA (0.1 ml), and N,N-dimethylethylenediamine (0.1 ml, 0.91 mmol) was

added. The mixture was then stirred at 140°C for one hour and at 100°C for one hour. The resulting reaction mixture was purified by silica gel column chromatography to give 3-(2-dimethylamino-ethyl)-1-methyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea as an amorphous (12 mg, yield 66%).

¹H-NMR (400 MHz, DMSO-d₆) δ 1.63-1.66 (m, 2H), 1.85-1.90 (m, 2H), 2.09 (s, 6H), 2.28 (t, J = 6.8 Hz, 2H), 2.40 (s, 3H), 3.06-3.17 (m, 7H), 3.58-3.61 (m, 2H), 6.18 (t, J = 5.8 Hz, 1H), 7.20 (d, J = 8.3 Hz, 1H), 7.22 (s, 1H), 7.26 (d, J = 15.6 Hz, 1H), 7.55 (d, J = 15.6 Hz, 1H), 7.57-7.67 (m, 2H), 7.82 (d, J = 8.3 Hz, 1H), 7.91 (br, 1H), 8.01 (br, 1H), 11.8 (br, 1H). MS (ESI) m/z = 637 (M+H)⁺.

[0732] The example compound shown below was synthesized by operations similar to those in Example 83 using appropriate reagents and starting material.

(Compound 458)

[0733]

[Table 67]


Compound	structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
458		LCMS-B-1	2.04	566 (M+H) ⁺

Example 84

1-{3-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-2-methyl-phenyl}-3-methyl-urea (Compound 459)

[0734]

(Reaction 84-1)




[0735] 1-[3-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-2-methyl-phenyl]-3-methyl-urea was synthesized by operations similar to those in Reaction 25-2, Reaction 42-1 and Reaction 84-1 using appropriate reagents and starting material.
MS (ESI) $m/z = 490$ (M+H)+.

Example 85

3-[8-(3-Chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid 2-methoxy-ethyl ester (Compound 460)

[0736]

(Reaction 85-1)



[0737] 3-[8-(3-Chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid 2-methoxy-ethyl ester was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.21-1.33 (m, 2 H), 1.34-1.51 (m, 3 H), 1.52-1.71 (m, 2 H), 1.73-2.02 (m, 3 H), 2.45-2.62 (m, 1 H), 2.81-2.95 (m, 3 H), 3.26 (s, 3 H), 3.43-3.51 (m, 2 H), 3.54-3.65 (m, 2 H), 3.95-4.04 (m, 1 H), 4.05-4.15 (m, 2 H), 7.35-7.44 (m, 1 H), 7.45-7.51 (m, 1 H), 7.56 (d, $J = 7.83$ Hz, 1 H), 7.66 (t, $J = 1.77$ Hz, 1 H). MS (ESI) $m/z = 513$ (M+H)+.

[0738] The example compounds shown below were synthesized by operations similar to those in Example 85 using appropriate reagents and starting materials.

(Compounds 461 to 465)

[0739]

[Table 68]


Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
461		LCMS-E-6	1.86	545 (M+H)+
462		LCMS-E-6	1.84	511 (M+H)+
463		LCMS-E-6	1.46	469 (M+H)+

(continued)

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
464		LCMS-E-3	3.65	589 (M+H)+
465		LCMS-E-3	3.75	569 (M+H)+

Example 86

Methyl-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-carbamic acid 2-dimethylamino-ethyl ester (Compound 466)

[0740](Reaction 86-1)

[0741] Methyl-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-carbamic acid 2-dimethylamino-ethyl ester was synthesized by operations similar to those in Reaction 83-1 using appropriate reagents and starting material.


MS (ESI) m/z = 638 (M+H)+.

[0742] The example compound shown below was synthesized by operations similar to those in Example 86 using appropriate reagents and starting material.

(Compound 467)


[0743]

[Table 69]

Compound	Structure	LCMS or HPLC condition	Retention time (min)	MS (m/z)
467		LCMS-A-1	2.74	581 (M+H)+

Example 87

Methyl-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-carbamate (Compound 468)


[0744]**(Reaction 87-1)**

[0745] Methyl-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-carbamate was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

MS (ESI) m/z = 583 (M+H)+.

Example 88

2-Cyclohexyl-8-(2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione (Compound 469)

[0746]


[0747] 2-Cyclohexyl-8-(2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (24.2 mg, 0.058 mmol), Lawesson's reagent (48.3 mg, 0.116 mmol) and toluene (1.16 ml) were added to a sealed test tube and stirred at 110°C overnight. The reaction mixture was cooled to ambient temperature, and the solvent was then distilled off under reduced pressure. The residue was purified by column chromatography (silica gel, CH₂Cl₂-MeOH) to give 2-cyclohexyl-8-(2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione (11.2 mg, 45%).

¹H-NMR (400 MHz, CD₃OD) δ 1.29-1.58 (7H, m), 1.73-2.15 (7H, m), 2.36 (3H, s), 2.55 (1H, tt, J = 4, 12 Hz), 3.09-3.13 (2H, m), 3.25-3.27 (2H, m), 3.30-3.31 (2H, m), 3.80-3.83 (2H, m), 7.13-7.23 (4H, m). MS (ESI) m/z = 434 (M+H)+.

Example 89


1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 470)

[0748]

(Reaction 89-1)


[0749] 8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) m/z = 473 (M+H)⁺.

(Reaction 89-2)

[0750] Sodium cyanate (15 mg, 0.243 mmol) was added to a solution of 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (23 mg, 0.0487 mmol) and acetic acid (1.3 ml) in dichloromethane (0.5 ml) at room temperature, and the mixture was then stirred for two hours. The reaction mixture was diluted with dichloromethane, and the organic layer was then washed with water and a saturated aqueous sodium bicarbonate solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by preparative TLC (silica gel, MeOH/AcOEt/CH₂Cl₂) to give 1-(3,5-dimethyl-4-{(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea (22.5 mg, 90%).

¹H-NMR (400 MHz, CDCl₃) δ 0.92 (3H, d, J = 4.0 Hz), 0.95-1.1 (2H, m), 1.35-1.50 (3H, m), 1.65-1.75 (2H, m), 1.80-1.85 (2H, m), 1.90-2.00 (4H, m), 2.30-2.40 (1H, m), 2.38 (6H, s), 3.26 (3H, s), 3.35-3.45 (2H, m), 3.60-3.75 (2H, m), 4.54 (2H, brs), 6.39 (1H, d, J = 16.0 Hz), 7.03 (2H, s), 7.54 (1H, d, J = 16.0 Hz), 8.10 (1H, brs). MS (ESI) m/z = 516 (M+H)⁺.

(Reaction 89-3)

[0751] 1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 42-2 using appropriate reagents and starting material.


¹H-NMR (400 MHz, CD₃OD) δ 0.92 (3H, d, J = 8.0 Hz), 0.95-1.06 (2H, m), 1.35-1.50 (3H, m), 1.65-1.75 (2H, m), 1.80-1.86 (2H, m), 1.88-2.00 (4H, m), 2.30-2.40 (1H, m), 2.36 (6H, s), 2.95-3.02 (2H, m), 3.15-3.22 (2H, m), 3.23 (3H, s), 3.45-3.52 (2H, m), 3.68-3.77 (2H, m), 4.47 (2H, brs), 6.95 (2H, s), 8.06 (1H, brs). MS (ESI) m/z = 518 (M+H)⁺.

Example 90

1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (Compound 471)

[0752]

(Reaction 90-1)



[0753] 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 5-3, Reaction 25-1, Reaction 26-1, Reaction 7-2 and Reaction 89-2 using appropriate reagents and starting material.


MS (ESI) $m/z = 570$ (M+H)+.

Example 91

1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 472)

[0754]

(Reaction 91-1)




[0755] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 42-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 572$ (M+H)+.


Example 92

(3-Methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-sulfamide (Compound 473)

[0756]

(Reaction 92-1)

[0757] 8-[(E)-2-(4-amino-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was obtained by operations similar to those in Reaction 26-1 using 8-ethenesulfonyl-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a starting material.
MS (ESI) m/z = 509 (M+H)+.

(Reaction 92-2)

[0758] A solution of tert-butanol (71.9 mg, 0.97 mmol) in dichloromethane (1.5 ml) was added to a solution of chloro-sulfonyl isocyanate (137 mg, 0.97 mmol) in dichloromethane (3 ml) with stirring under ice-cooling. The mixture was stirred at 0°C for 10 minutes. A solution of 8-[(E)-2-(4-amino-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (400 mg, 0.81 mmol) and triethylamine (164 mg, 1.62 mmol) in dichloromethane (3 ml) was then added, and the mixture was further stirred for one hour. The mixed reaction solution was quenched with water and then extracted with dichloromethane. The organic layer was washed with saturated brine, and then dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate-hexane) to give N-(tert-butoxycarbonyl)-N'-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)sulfamide (334 mg, 60.0%).

¹H-NMR (400 MHz, CDCl₃) δ 1.42 (9H, s), 1.78 (2H, dt, J = 14.2, 3.9 Hz), 2.04-2.14 (2H, m), 2.40 (3H, s), 3.43 (2H, ddd, J = 12.7, 9.8, 2.9 Hz), 3.74 (2H, dt, J = 12.2, 4.4 Hz), 6.64 (1H, d, J = 15.6 Hz), 7.08-7.11 (2H, m), 7.38 (1H, d, J = 8.3 Hz), 7.48-7.54 (2H, m), 7.68 (1H, d, J = 15.1 Hz), 7.73 (1H, d, J = 7.8 Hz), 7.76 (1H, s), 9.62 (1H, s);

MS (ESI) m/z = 688 (M+H)+.


Example 93

1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-thiourea (Compound 474)

[0761] Benzoyl isothiocyanate (37.8 mg, 0.23 mmol) was added to a solution of 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (100 mg, 0.21 mmol) in acetone (3 ml) in a nitrogen stream. The mixture was heated under reflux for one hour and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 3-benzoyl-1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-thiourea (140 mg).

¹H-NMR (400 MHz, CDCl₃) δ 0.92 (3H, d, J = 6.8 Hz), 0.94-1.07 (2H, m), 1.32-1.46 (3H, m), 1.52-1.64 (2H, m), 1.82 (2H, dd, J = 13.7, 2.4 Hz), 1.90-2.00 (4H, m), 2.28-2.33 (1H, m), 2.35 (6H, s), 2.90-3.00 (2H, m), 3.08-3.16 (2H, m), 3.36-3.45 (2H, m), 3.64-3.78 (5H, m), 7.01 (2H, s), 7.37-7.62 (5H, m);

MS (ESI) m/z = 638 (M+H)+.

(Reaction 93-2)


[0762] Hydrazine monohydrate (55 mg, 1.1 mmol) was added to a solution of 3-benzoyl-1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-thiourea (140 mg, 0.22 mmol) in ethanol (7 ml). The mixture was stirred at room temperature for 15 hours and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 1-(3,5-dimethyl-4-{2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-thiourea (119 mg).

¹H-NMR (400 MHz, CDCl₃) δ 0.92 (3H, d, J = 6.3 Hz), 1.00 (2H, ddd, J = 25.4, 13.7, 2.9 Hz), 1.32-1.47 (3H, m), 1.59-1.68 (2H, m), 1.82 (2H, dd, J = 10.7, 3.4 Hz), 1.90-1.98 (4H, m), 2.28-2.35 (1H, m), 2.35 (6H, s), 2.95-3.02 (2H, m), 3.15-3.22 (2H, m), 3.45 (2H, ddd, J = 12.2, 9.3, 3.4 Hz), 3.55 (3H, s), 3.74 (2H, dt, J = 13.1, 4.4 Hz), 5.63 (1H, brs), 6.94 (2H, s), 8.14 (1H, s);

MS (ESI) m/z = 534 (M+H)⁺.

Example 94

8-{2-[2,6-Dimethyl-4-(methyl-thiazol-2-yl-amino)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 475)

[0763](Reaction 94-1)

[0764] Bromoacetaldehyde diethylacetal (44.2 mg, 0.224 mmol) was added to a solution of 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-thiourea (100 mg, 0.187 mmol) in acetic acid (2 ml) in a nitrogen stream. The mixture was heated under reflux for two hours and then concentrated under reduced pressure. The resulting residue was diluted with dichloromethane, and the organic layer was then sequentially washed with a saturated aqueous sodium bicarbonate solution and saturated brine and dried over anhydrous magnesium sulfate. The organic layer was concentrated, and the resulting residue was then purified by silica gel column chromatography (dichloromethane-methanol) to give 8-{2-[2,6-Dimethyl-4-(methyl-thiazol-2-yl-amino)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (7.1 mg).

¹H-NMR (400 MHz, CDCl₃) δ 0.92 (3H, d, J = 6.8 Hz), 0.95-1.07 (2H, m), 1.33-1.48 (3H, m), 1.56-1.67 (2H, m), 1.82 (2H, d, J = 11.2 Hz), 1.91-2.03 (4H, m), 2.29-2.35 (1H, m), 2.37 (6H, s), 2.99-3.06 (2H, m), 3.14-3.21 (2H, m), 3.43 (2H, ddd, J = 12.7, 9.3, 2.9 Hz), 3.49 (3H, s), 3.77 (2H, dt, J = 12.2, 4.4 Hz), 6.48 (1H, d, J = 3.4 Hz), 7.06 (2H, s), 7.22 (1H,

d, J = 3.4 Hz), 8.33 (1H, s);
MS (ESI) m/z = 558 (M+H)+.

Example 95

[0765] The example compounds shown below were obtained by operations similar to those in Reaction 25-2 using appropriate reagents and starting materials.

(Compounds 476 to 503)

[0766]

[Table 70]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
476		LCMS-C-1	2.53	513 (M+H)+
477		LCMS-C-1	2.55	554 (M+H)+
478		LCMS-C-1	2.62	513 (M+H)+
479		LCMS-A-1	2.25	441 (M+H)+
480		LCMS-C-1	2.42	529 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
481		LCMS-C-1	2.30	485 (M-H)-
482		LCMS-C-1	2.30	485 (M-H)-
483		LCMS-C-1	2.62	536 (M-H)-
484		LCMS-C-1	2.38	562 (M+H)+
485		LCMS-C-1	2.68	611 (M+H)+
486		LCMS-C-1	2.75	529 (M-H)-
487		LCMS-A-1	2.00	529 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
488		LCMS-A-1	2.17	455 (M+H)+
489		LCMS-C-1	2.30	489 (M+H)+
490		LCMS-C-1	2.38	516 (M+H)+
491		LCMS-C-1	2.68	551 (M-H)-
492		LCMS-A-1	2.30	509 (M+H)+
493		LCMS-C-1	2.65	557 (M+H)+
494		LCMS-A-1	1.90	586 (M+H)+

(continued)


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
495		LCMS-C-1	2.60	607 (M+H)+
496		LCMS-C-1	2.48	637 (M+H)+
497		LCMS-C-1	2.32	502 (M+H)+
498		LCMS-C-1	2.37	627 (M+H)+
499		LCMS-C-1	2.55	510 (M+H)+
500		LCMS-A-1	2.35	573 (M+H)+
501		LCMS-A-1	2.14	543 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
502		LCMS-C-1	2.18	577 (M+H)+
503		LCMS-C-1	2.28	566 (M+H)+

[0767] The aryl bromide reagent used in the synthesis of Compound 477 ((4-bromo-indol-1-yl)-morpholin-4-yl-methanone) was synthesized as follows.

(Reaction 95-1)




[0768] (4-Bromo-indol-1-yl)-morpholin-4-yl-methanone was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 3.61 (4H, t, J = 4.8 Hz), 3.78 (4H, t, J = 4.8 Hz), 6.69 (1H, d, J = 3.6 Hz), 7.17 (1H, t, J = 7.9 Hz), 7.35 (2H, d, J = 3.6 Hz), 7.38 (2H, d, J = 7.9 Hz), 7.65 (1H, d, J = 7.9 Hz).

[0769] The aryl bromide reagent used in the synthesis of Compound 478 (4-(4-bromo-indol-1-yl)-butan-1-ol) was synthesized as follows.

(Reaction 95-2)




[0770] 4-(4-Bromo-indol-1-yl)-butan-1-ol was synthesized by operations similar to those in Reaction 29-7 and Reaction

25-4 using appropriate reagents and starting material.

MS (ESI) m/z = 268, 270 ($M+H$)⁺.

[0771] The aryl bromide reagent used in the synthesis of Compound 480 (4-(4-bromo-indol-1-yl)-butane-1,2-diol) was synthesized as follows.


(Reaction 95-3)



[0772] 4-(4-Bromo-indol-1-yl)-butane-1,2-diol was synthesized by operations similar to those in Reaction 25-3 and Reaction 25-4 using appropriate reagents and starting material. MS (ESI) m/z = 284, 286 ($M+H$)⁺.

[0773] The aryl bromide reagent used in the synthesis of Compound 483 (4-bromo-1-thiazol-2-ylmethyl-1H-indole) was synthesized as follows.


(Reaction 95-4)



[0774] 4-Bromo-1-thiazol-2-ylmethyl-1H-indole was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material. MS (ESI) m/z = 293, 295 ($M+H$)⁺.

[0775] The aryl bromide reagent used in the synthesis of Compound 484 (3-(4-bromo-indol-1-yl)-propane-1-sulfonic amide) was synthesized as follows.


(Reaction 95-5)



[0776] 3-(4-Bromo-indol-1-yl)-propane-1-sulfonyl chloride was synthesized as a crude product by operations similar

to those in Reaction 25-3 and Reaction 52-3 using 4-bromoindole (0.20 ml, 1.59 mmol) as a starting material and using THF as a solvent.

(Reaction 95-6)




[0777] 3-(4-Bromo-indol-1-yl)-propane-1-sulfonyl chloride obtained as a crude product was all dissolved in diethyl ether (3.0 ml). A 28% aqueous ammonia solution (3.0 ml) was then added dropwise and the mixture was stirred at room temperature for 2.5 hours. Water was added to the reaction system, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give 3-(4-bromo-indol-1-yl)-propane-1-sulfonamide (55.6 mg, 11% in three steps).

MS (ESI) m/z = 317, 319 (M+H)⁺.

[0778] The aryl bromide reagent used in the synthesis of Compound 485 (4-methyl-piperazine-1-carboxylic (4-bromo-3-trifluoromethyl-phenyl)-amide) was synthesized as follows.

(Reaction 95-7)




[0779] 4-Methyl-piperazine-1-carboxylic (4-bromo-3-trifluoromethyl-phenyl)-amide was synthesized by operations similar to those in Reaction 25-11 using appropriate reagents and starting material.

MS (ESI) m/z = 366, 368 (M+H)⁺.

[0780] The aryl bromide reagent used in the synthesis of Compound 486 ((4-bromo-benzyl)-carbamic acid isobutyl ester) was synthesized as follows.

(Reaction 95-8)




[0781] (4-Bromo-benzyl)-carbamic acid isobutyl ester was synthesized by operations similar to those in Reaction 25-10

using appropriate reagents and starting material.

MS (ESI) m/z = 286, 288 (M+H)+.

[0782] The aryl bromide reagent used in the synthesis of Compound 487 (2-[2-(5-bromo-indol-1-yl)-ethoxy]-ethanol) was synthesized as follows.

(Reaction 95-9)




[0783] 2-[2-(5-Bromo-indol-1-yl)-ethoxy]-ethanol was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 7.74 (1H, d, J = 1.6 Hz), 7.28 (1H, dd, J = 8.7, 1.8 Hz), 7.23 (1H, d, J = 8.6 Hz), 7.15 (1H, d, J = 3.1 Hz), 6.44 (1H, d, J = 3.1 Hz), 4.29 (2H, t, J = 5.4 Hz), 3.79 (2H, t, J = 5.4 Hz), 3.65-3.59 (2H, m), 3.48-3.44 (2H, m), 1.67 (1H, t, J = 6.1 Hz).

[0784] The aryl bromide reagent used in the synthesis of Compound 488 (5-bromo-6-methyl-1H-indole) was synthesized as follows.


(Reaction 95-10)



[0785] (4-Bromo-2-iodo-5-methyl-phenyl)-carbamic acid methyl ester (605 mg, 1.64 mmol) was dissolved in THF (6 ml). (Trimethylsilyl)acetylene (0.70 ml, 4.95 mmol), copper iodide (33.5 mg, 0.175 mmol), $\text{Pd(PPh}_3)_2\text{Cl}_2$ (56.5 mg, 0.081 mmol) and triethylamine (0.690 ml, 4.95 mmol) were added and the mixture was stirred at room temperature for four hours. Water was added to the reaction system, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane-ethyl acetate) to give (4-bromo-5-methyl-2-trimethylsilyl-ethynyl-phenyl)-carbamic acid methyl ester (548 mg, 98%).

MS (ESI) m/z = 340 (M+H)+.

(Reaction 95-11)




[0786] (4-Bromo-5-methyl-2-trimethylsilyl-ethynyl-phenyl)-carbamic acid methyl ester (506 mg, 1.49 mmol) was dissolved in ethanol (6 ml). Sodium ethoxide (20% solution in ethanol, 1.17 ml, 2.97 mmol) was added and the mixture was stirred at 70°C overnight. The reaction solution was poured into ice water, and 1 N hydrochloric acid and saturated brine were added to this mixture, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane-ethyl acetate) to give 5-bromo-6-methyl-1H-indole (207 mg, 66%).

¹H-NMR (270 MHz, CDCl₃) δ 8.04 (1H, br s), 7.81 (1H, s), 7.27 (1H, s), 7.16-7.14 (1H, m), 6.46-6.43 (1H, m), 2.49 (3H, s).


[0787] The aryl bromide reagent used in the synthesis of Compound 489 (N-(4-bromo-benzyl)-2-hydroxy-acetamide) was synthesized as follows.

(Reaction 95-12)



[0788] Acetic acid (4-bromo-benzylcarbamoyl)-methyl ester was synthesized as a crude product by operations similar to those in Reaction 2-3 using 4-bromobenzylamine hydrochloride (200 mg, 0.899 mmol) as a starting material and using pyridine as a base.

(Reaction 95-13)




[0789] Acetic acid (4-bromo-benzylcarbamoyl)-methyl ester obtained as a crude product was all dissolved in THF (2.0 ml). A 2 N aqueous sodium hydroxide solution (2.0 ml) was then added and the mixture was stirred at room temperature for five hours. Water was added to the reaction system, followed by extraction with ethyl acetate. The organic layer was washed with saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica

gel column chromatography (dichloromethane-methanol) to give N-(4-bromo-benzyl)-2-hydroxy-acetamide (56.0 mg, 25% for two steps).

MS (ESI) m/z = 244, 246 (M+H)+.

[0790] The aryl bromide reagent used in the synthesis of Compound 490 (3-[2-(4-bromo-phenyl)-ethyl]-1,1-dimethyl-urea) was synthesized as follows.

(Reaction 95-14)




[0791] 3-[2-(4-Bromo-phenyl)-ethyl]-1,1-dimethyl-urea was synthesized by operations similar to those in Reaction 82-1 using appropriate reagents and starting material.

MS (ESI) m/z = 271, 273 (M+H)+.

[0792] The aryl bromide reagent used in the synthesis of Compound 492 (5-bromo-6-trifluoromethyl-1H-indole) was synthesized as follows.


(Reaction 95-15)



[0793] 4-Bromo-3-trifluoromethyl-phenylamine (1.00 g, 4.17 mmol) was dissolved in acetic acid (5 ml). Iodine monochloride (1 M solution in dichloromethane, 5 ml) was added and the mixture was stirred at 60°C overnight. The reaction solution was poured into a mixture of ice and a saturated aqueous sodium bicarbonate solution and then extracted with ethyl acetate. The organic layer was washed with an aqueous sodium bicarbonate solution, an aqueous sodium thiosulfate solution, water and saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane-ethyl acetate) to give 4-bromo-2-iodo-5-trifluoromethyl-phenylamine (889 mg, 58%).

$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 7.92 (1H, s), 7.00 (1H, s), 4.31 (2H, br s).

(Reaction 95-16)




[0794] 5-Bromo-6-trifluoromethyl-1H-indole was synthesized by operations similar to those in Reaction 25-10, Reaction 95-10 and Reaction 95-11 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 8.41 (1H, s), 7.96 (1H, s), 7.78 (1H, s), 7.38-7.36 (1H, m), 6.57-6.54 (1H, m).

[0795] The aryl bromide reagent used in the synthesis of Compound 493 ((4-bromo-3-trifluoromethyl-phenylamino)-acetic acid methyl ester) was synthesized as follows.

(Reaction 95-17)




[0796] N,N -Diisopropylethylamine (1.22 ml, 7.00 mmol) and methyl bromoacetate (1.00 g, 6.54 mmol) were sequentially added to a solution of 4-bromo-3-(trifluoromethyl)aniline (1.40 g, 5.83 mmol) in DMF (10 ml), and the mixture was heated with stirring at 80°C for 25 hours. The reaction mixture was cooled and water was then added, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was triturated with hexane:dichloromethane = 9:1 to give (4-bromo-3-trifluoromethyl-phenylamino)-acetic acid methyl ester (1.22 g, 67%).

MS (ESI) m/z = 312, 314 ($\text{M}+\text{H}$) $^+$


[0797] The aryl bromide reagent used in the synthesis of Compound 494 (2-(4-bromo-3-trifluoromethyl-phenylamino)- N -(2-hydroxy-ethyl)-acetamide) was synthesized as follows.

(Reaction 95-18)



[0798] A 2 N aqueous NaOH solution (15.0 ml, 30.0 mmol) was added to a solution of (4-bromo-3-trifluoromethyl-phenylamino)-acetic acid methyl ester (4.30 g, 13.8 mmol) in methanol-THF (6:1, 35.0 ml), and the mixture was stirred at room temperature for 18 hours. The reaction mixture was made acidic with hydrochloric acid and extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over MgSO_4 and concentrated under reduced pressure to give (4-bromo-3-trifluoromethyl-phenylamino)-acetic acid (4.03 g, 98%).
MS (ESI) m/z = 298, 300 ($\text{M}+\text{H}$)⁺

(Reaction 95-19)




[0799] 2-(4-Bromo-3-trifluoromethyl-phenylamino)-N-(2-hydroxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

¹H-NMR (CDCl_3) δ 2.11 (1H, t, J = 5.1 Hz), 3.48 (2H, q, J = 5.2 Hz), 3.73 (2H, q, J = 5.2 Hz), 3.84 (2H, d, J = 5.4 Hz), 4.51-4.57 (1H, m), 6.61 (1H, dd, J = 8.7, 2.9 Hz), 6.72 (1H, s), 6.93 (1H, d, J = 2.9 Hz), 7.49 (1H, d, J = 8.7 Hz).

[0800] The aryl bromide reagent used in the synthesis of Compound 495 (4-bromo-N-(2-hydroxy-ethyl)-N-methyl-3-trifluoromethyl-benzenesulfonamide) was synthesized as follows.

(Reaction 95-20)




[0801] 4-Bromo-N-(2-hydroxy-ethyl)-N-methyl-3-trifluoromethyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 362, 364 ($\text{M}+\text{H}$)⁺.

[0802] The aryl bromide reagent used in the synthesis of Compound 496 (4-bromo-N,N-bis-(2-hydroxy-ethyl)-3-trifluoromethyl-benzenesulfonamide) was synthesized as follows.

(Reaction 95-21)




[0803] 4-Bromo-N,N-bis-(2-hydroxy-ethyl)-3-trifluoromethyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 392, 394 ($M+H$)⁺.

[0804] The aryl bromide reagent used in the synthesis of Compound 497 (N'-(4-bromo-3-methyl-phenyl)-N,N-dimethylethane-1,2-diamine) was synthesized as follows.

(Reaction 95-22)




[0805] 2-Chloro-N,N-dimethylethylamine hydrochloride (372 mg, 2.58 mmol), potassium iodide (428 mg, 2.58 mmol) and triethylamine (0.719 ml, 5.16 mmol) were added to a solution of 4-bromo-3-methylaniline (400 mg, 2.15 mmol) in toluene (5.0 ml), and the mixture was heated with stirring at 110°C for 17 hours. The reaction mixture was cooled and water was then added, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give N'-(4-bromo-3-methyl-phenyl)-N,N-dimethylethane-1,2-diamine (100 mg, 18%).

MS (ESI) m/z = 257, 259 ($M+H$)⁺.

[0806] The aryl bromide reagent used in the synthesis of Compound 498 (N-(2-acetylamino-ethyl)-2-(4-bromo-3-trifluoromethyl-phenylamino)-acetamide) was synthesized as follows.

(Reaction 95-23)




[0807] N-(2-Acetylamino-ethyl)-2-(4-bromo-3-trifluoromethyl-phenylamino)-acetamide was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 382, 384 ($M+H$)+.

[0808] The aryl bromide reagent used in the synthesis of Compound 499 (5-bromo-6-trifluoromethyl-1H-benzimidazole) was synthesized as follows.

(Reaction 95-24)




[0809] 4-Bromo-5-trifluoromethyl-benzene-1,2-diamine (200 mg, 0.785 mmol) was dissolved in formic acid (3 ml), and the mixture was stirred at 120°C for six hours. The reaction solution was concentrated, and water was added to the resulting residue, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 5-bromo-6-trifluoromethyl-1H-benzimidazole (201 mg) as a crude compound.

MS (ESI) m/z = 265 ($M+H$)+.

[0810] The aryl bromide reagent used in the synthesis of Compound 500 (4-bromo-1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-5-fluoro-1H-indole) was synthesized as follows.


(Reaction 95-25)



[0811] A solution of n-butyllithium (2.1 mL, 3.39 mmol) in tetrahydrofuran (6.8 mL) was cooled to -78°C, and 2,2,6,6-tetramethylpiperidine (0.57 mL, 3.39 mmol) and a 1.0 M solution of potassium t-butoxide in tetrahydrofuran (3.4 mL, 3.39 mmol) were added. After stirring for 15 minutes, a solution of 5-fluoro-1-triisopropylsilanyl-1H-indole (494 mg, 1.70 mmol) in tetrahydrofuran (5 mL) was added dropwise, and the mixture was stirred at -78°C for 2.5 hours. 1,2-Dibromo-1,1,2,2-tetrafluoroethane (38 mL, 0.319 mmol) was added, and the mixture was warmed to -40°C over 35 minutes and further warmed to 22°C over 12 hours. Silica gel (17 g) was added and the solvent was then distilled off. The residue was subjected to silica gel column chromatography to give a pale yellow oily substance (380 mg) as a mixture of 4-bromo-5-fluoro-1-triisopropylsilanyl-1H-indole:5-fluoro-1-triisopropylsilanyl-1H-indole = 1:1.1.

¹H-NMR (CDCl₃) δ 7.37-7.33 (2H, m), 6.94 (1H, dd, J = 9.0, 4.5 Hz), 6.68 (1H, d, J = 3.9 Hz), 1.71-1.63 (3H, m), 1.14 (18H, d, J = 7.3 Hz).

(Reaction 95-26)




[0812] 4-Bromo-1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-5-fluoro-1H-indole was synthesized by operations similar to those in Reaction 39-2 and Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 328, 330 (M+H)+.

[0813] The aryl bromide reagent used in the synthesis of Compound 501 (2-[(4-bromo-3-trifluoromethyl-phenyl)-methyl-amino]-ethanol) was synthesized as follows.


(Reaction 95-27)



[0814] [(4-Bromo-3-trifluoromethyl-phenyl)-methyl-amino]-acetic acid methyl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 326, 328 (M+H)+.

(Reaction 95-28)




[0815] A solution of [(4-bromo-3-trifluoromethyl-phenyl)-methyl-amino]-acetic acid methyl ester (77.6 mg, 0.238 mmol) in THF (1.0 ml) was added dropwise to a suspension of lithium aluminum hydride (372 mg, 2.58 mmol) in THF (1.5 ml) at 0°C. The mixture was stirred for 14 hours while gradually warming from 0°C to room temperature. A 2 N aqueous HCl solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give 2-[(4-bromo-3-trifluoromethyl-phenyl)-methyl-amino]-ethanol (58.0 mg, 82%).

MS (ESI) m/z = 298, 300 (M+H)+.

[0816] The aryl bromide reagent used in the synthesis of Compound 502 ((4-bromo-3-methyl-phenyl)-(1,1-dioxo-1 λ^6 -thiomorpholin-4-yl)-methanone) was synthesized as follows.

(Reaction 95-29)




[0817] (4-Bromo-3-methyl-phenyl)-(1,1-dioxo-1 λ^6 -thiomorpholin-4-yl)-methanone was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 332, 334 (M+H)+.

[0818] The aryl bromide reagent used in the synthesis of Compound 503 (4-bromo-N-(2-dimethylamino-ethyl)-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 95-30)



[0819] 4-Bromo-N-(2-dimethylamino-ethyl)-3-methyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 321, 323 (M+H)+.







Example 96

[0820] The example compounds shown below were obtained by operations similar to those in Reaction 25-2 using appropriate reagents and starting materials.

(Compounds 504 to 523)

[0821]

[Table 71]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
504		LCMS-C-1	2.4	528 (M+H) ⁺
505		LCMS-C-1	2.47	542 (M+H) ⁺
506		LCMS-D-1	1.91	531 (M+H) ⁺
507		LCMS-D-1	1.9	570 (M+H) ⁺
508		LCMS-D-1	1.98	584 (M+H) ⁺
509		LCMS-D-1	1.9	517 (M+H) ⁺

(continued)


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
510		LCMS-D-1	1.82	501 (M+H)+
511		LCMS-F-1	0.92	556 (M+H)+
512		LCMS-D-1	1.84	517 (M+H)+
513		LCMS-C-1	2.47	515 (M+H)+
514		LCMS-D-1	2.09	559 (M+H)+
515		LCMS-D-1	1.83	543 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
516		LCMS-D-1	2.22	541 (M+H)+
517		LCMS-D-1	2.23	529 (M+H)+
518		LCMS-D-1	2.13	556 (M+H)+
519		LCMS-D-1	2.4	584 (M+H)+
520		LCMS-D-1	2.28	541 (M+H)+
521		LCMS-D-1	2.27	555 (M+H)+
522		LCMS-D-1	2.45	515 (M+H)+
523		LCMS-D-1	2.33	523 (M+H)+

[0822] The aryl bromide reagent used in the synthesis of Compound 504 (3-(4-bromo-3-methyl-phenyl)-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 96-1)




[0823] Triethylamine (20 ml, 145 mmol) and 4-bromo-3-methyl-phenylamine (4.49 g, 24.14 mmol) were added to a solution of triphosgene (1.0 ml, 8.05 mmol) in THF (70 ml) at 0°C, and the mixture was stirred at the same temperature for 40 minutes. Further, the reaction mixture was warmed to room temperature and stirred at the same temperature for one hour. Water was added to the reaction solution, and the mixture was then concentrated under reduced pressure, followed by extraction with ethyl acetate. The organic phase was sequentially washed with water and saturated brine and then concentrated under reduced pressure to give [3-(4-bromo-3-methyl-phenyl)-ureido]-acetic acid ethyl ester (5.07 g) as a crude product. This crude product was used in the next reaction without purification.

[0824] 4 N HCl-dioxane (7.5 ml, 30 mmol) was added to a solution of the crude product [3-(4-bromo-3-methyl-phenyl)-ureido]-acetic acid ethyl ester (5.07 g) in dioxane (60 ml), and the mixture was heated with stirring at 80°C for 17 hours. The reaction mixture was cooled to room temperature and then concentrated under reduced pressure, and water was added, followed by extraction with ethyl acetate. The organic phase was washed with saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 3-(4-bromo-3-methyl-phenyl)-imidazolidine-2,4-dione (2.40 g, 37%).

MS (ESI) m/z = 269, 271 ($M+H$)⁺.

[0825] The aryl bromide reagent used in the synthesis of Compound 505 (3-(4-bromo-3-methyl-phenyl)-1-methyl-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 96-2)




[0826] (3-(4-Bromo-3-methyl-phenyl)-1-methyl-imidazolidine-2,4-dione) was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 283, 285 ($M+H$)⁺.

[0827] The aryl bromide reagent used in the synthesis of Compound 506 (N-(4-bromo-3-isopropoxyphenyl)acetamide) was synthesized as follows.

(Reaction 96-3)




[0828] N-(4-Bromo-3-isopropoxyphenyl)acetamide was synthesized by operations similar to those in Reaction 26-2 and Reaction 26-4 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 7.64 (d, 1H, J = 2.29 Hz), 7.42 (d, 1H, J = 8.39 Hz), 7.15 (brs, 1H), 6.72 (dd, 1H, J = 8.39 Hz, J = 2.29 Hz), 4.57 (m, 1H, J = 6.1 Hz), 2.17 (s, 3H), 1.38 (d, 6H, J = 6.1 Hz).

[0829] The aryl bromide reagents used in the synthesis of Compound 507 and Compound 508 (3-(4-bromo-3,5-dimethylphenyl)-5,5-dimethylimidazolidine-2,4-dione and 3-(4-bromo-3,5-dimethylphenyl)-1,5,5-trimethylimidazolidine-2,4-dione) were synthesized as follows.


(Reaction 96-4)



[0830] Triphosgene (2.23 g, 7.52 mmol) was added to a solution of 4-bromo-3,5-dimethyl-phenylamine (4.3 g, 21.49 mmol) in dioxane (71 ml). After stirring at 100°C for 15 hours, water was added to the reaction solution. After extraction with ethyl acetate, the organic phase was sequentially washed with water and saturated brine and concentrated under reduced pressure to give 2-bromo-5-isocyanato-1,3-dimethylbenzene (2.0 g, 41%).

MS (ESI) m/z = 226, 228 ($M+H$)⁺.


(Reaction 96-5)



[0831] Triethylamine (1.8 ml, 13.27 mmol) was added dropwise to a solution of 2-bromo-5-isocyanato-1,3-dimethylbenzene (2.0 g, 8.84 mmol) in anhydrous methanol (30 ml) at 0°C. The reaction solution was gradually warmed to room temperature and stirred for one hour. The reaction solution was concentrated under reduced pressure, and the residue was diluted with ethyl acetate. The organic layer was then sequentially washed with water and saturated brine and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane-ethyl

acetate) to give methyl (4-bromo-3,5-dimethylphenyl)carbamate (1.8 g, 79%).
MS (ESI) m/z = 258, 260 ($M+H$)⁺.

(Reaction 96-6)



[0832] α -Aminoisobutyric acid methyl ester (3.2 g, 20.92 mmol) and N,N-diisopropylethylamine (6.1 ml, 34.86 mmol) were added to a solution of methyl (4-bromo-3,5-dimethylphenyl)carbamate (900 mg, 3.48 mmol) in anhydrous DMA (17.5 ml, 0.2 M). After irradiation with microwaves at 170°C for 30 minutes, water was added to the reaction solution, followed by extraction with ethyl acetate. The organic phase was sequentially washed with water and saturated brine and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 3-(4-bromo-3,5-dimethylphenyl)-5,5-dimethylimidazolidine-2,4-dione (304 mg, 28%)

MS (ESI) m/z = 311, 313 ($M+H$)⁺


and

3-(4-bromo-3,5-dimethylphenyl)-1,5,5-trimethylimidazolidine-2,4-dione (245 mg, 23%).

MS (ESI) m/z = 325, 327 ($M+H$)⁺

[0833] The aryl bromide reagent used in the synthesis of Compound 509 (N-(4-bromo-3-ethoxyphenyl)acetamide) was synthesized as follows.

(Reaction 96-7)




[0834] N-(4-Bromo-3-ethoxyphenyl)acetamide was synthesized by operations similar to those in Reaction 26-4 using appropriate reagents and starting material.

^1H -NMR ($CDCl_3$) δ 7.48 (d, 1H, J = 2.29 Hz), 7.42 (d, 1H, J = 8.39 Hz), 7.22 (brs, 1H), 6.72 (dd, 1H, J = 8.39 Hz, J = 2.29 Hz), 4.10 (q, 2H, J = 6.87 Hz), 2.17 (s, 3H), 1.46 (t, 3H, J = 6.87 Hz).

[0835] The aryl bromide reagent used in the synthesis of Compound 510 (N-(4-bromo-3-ethylphenyl)acetamide) was synthesized as follows.

(Reaction 96-8)




[0836] N-(4-Bromo-3-ethylphenyl)acetamide was synthesized by operations similar to those in Reaction 26-2 using

appropriate reagents and starting material.

$^1\text{H-NMR}$ (CDCl_3) δ 7.44 (d, 1H, $J = 8.39$ Hz), 7.39 (d, 1H, $J = 2.29$ Hz), 7.23 (dd, 1H, $J = 8.39$ Hz, $J = 2.29$ Hz), 7.17 (brs, 1H), 2.72 (q, 2H, $J = 7.25$ Hz), 2.17 (s, 3H), 1.21 (t, 3H, $J = 7.25$ Hz).

[0837] The aryl bromide reagent used in the synthesis of Compound 511 (3-(4-bromo-3-methyl-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione) was synthesized as follows.


(Reaction 96-9)



[0838] 2-Amino-N-(4-bromo-3-methyl-phenyl)-2-methyl-propionamide was synthesized by operations similar to those in Reaction 10-1 and Reaction 39-2 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (CDCl_3) δ 1.46 (6H, s), 2.37 (3H, s), 7.29 (1H, dd, $J = 8.8$, 2.4 Hz), 7.44 (1H, d, $J = 8.8$ Hz), 7.59 (1H, d, $J = 2.4$ Hz), 9.84 (1H, br s).

(Reaction 96-10)




[0839] Bis(trichloromethyl) carbonate (50.6 mg, 0.171 mmol) was added to a solution of 2-amino-N-(4-bromo-3-methyl-phenyl)-2-methyl-propionamide (132 mg, 0.487 mmol) and triethylamine (0.204 ml, 1.46 mmol) in dichloromethane (5.0 ml) at 0°C, and the mixture was stirred for 21 hours while gradually warming to room temperature. An aqueous ammonium chloride solution was added to the reaction mixture, followed by extraction with dichloromethane. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give 3-(4-bromo-3-methyl-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione (146 mg, 88%).

MS (ESI) $m/z = 297$, 299 ($\text{M}+\text{H}$) $^+$.

[0840] The aryl bromide reagent used in the synthesis of Compound 512 (N-(4-bromo-3-methoxymethyl-phenyl)-acetamide) was synthesized as follows.

(Reaction 96-11)




[0841] N-(4-Bromo-3-methoxymethyl-phenyl)-acetamide was synthesized by operations similar to those in Reaction 20-2, Reaction 19-2 and Reaction 26-2 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (CDCl_3) δ 7.12 (t, 1H, $J = 7.8$ Hz), 6.70 (m, 2H), 6.60 (dd, 1H, $J = 7.5$ Hz, 2.1 Hz), 4.42 (s, 2H), 3.53 (s, 3H), 2.08 (s, 3H).

[0842] The aryl bromide reagent used in the synthesis of Compound 513 (3-(4-bromo-3-methyl-phenyl)-oxazolidin-2-one) was synthesized as follows.


(Reaction 96-12)



[0843] 2-(4-Bromo-3-methyl-phenylamino)-ethanol was synthesized by operations similar to those in Reaction 12-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 230, 232$ ($\text{M}+\text{H}$) $^+$.

(Reaction 96-13)




[0844] Diethyl carbonate (18 ml), and a 28% solution of sodium methoxide in methanol (1.1 ml, 5.70 mmol) were added to 2-(4-bromo-3-methyl-phenylamino)-ethanol (1.21 g, 5.27 mmol), and the mixture was heated with stirring at 110°C for 15 hours. Further, methanol (16 ml) was added to the reaction solution, and the mixture was heated with stirring at 110°C for one hour. An aqueous ammonium chloride solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic phase was sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was triturated with hexane to give 3-(4-bromo-3-methyl-phenyl)-oxazolidin-2-one (989 mg, 73%).

MS (ESI) m/z = 256, 258 (M+H)+.

[0845] The aryl bromide reagent used in the synthesis of Compound 514 (N-(4-bromo-3,5-dimethyl-phenyl)-N-(2-methoxy-ethyl)-acetamide) was synthesized as follows.

(Reaction 96-14)




[0846] N-(4-Bromo-3,5-dimethyl-phenyl)-N-(2-methoxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 300, 302 (M+H)+.

[0847] The aryl bromide reagent used in the synthesis of Compound 515 (4-(4-bromo-3,5-dimethyl-phenyl)-morpholin-3-one) was synthesized as follows.


(Reaction 96-15)



[0848] Bromo-acetic acid 2-[acetyl-(4-bromo-3,5-dimethylphenyl)-amino]-ethyl ester was synthesized by operations


similar to those in Reaction 25-3, Reaction 39-2 and Reaction 10-1 using appropriate reagents and starting material.
 $^1\text{H-NMR}$ (CDCl_3) δ 6.95 (s, 2H), 4.04 (s, 2H), 3.86-3.74 (m, 4H), 2.43 (s, 6H), 1.89 (s, 3H).

(Reaction 96-16)




[0849] 6 N NaOH (1.5 ml) was added to a solution of bromo-acetic acid 2-[acetyl-(4-bromo-3,5-dimethyl-phenyl)-amino]-ethyl ester (302 mg, 0.74 mmol) in EtOH (6 ml). The reaction solution was heated under reflux overnight, cooled to room temperature and then diluted with ethyl acetate. The organic phase was sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 2-(4-bromo-3,5-dimethyl-phenylamino)-ethanol (181 mg, 68%).
 $^1\text{H-NMR}$ (CDCl_3) δ 6.41 (s, 2H), 3.82 (t, 2H, $J = 5.1$ Hz), 3.27 (t, 2H, $J = 5.1$ Hz), 2.34 (s, 6H).

(Reaction 96-17)



[0850] 2-Bromo-N-(4-bromo-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)acetamide was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.
 $^1\text{H-NMR}$ (CDCl_3) δ 7.01 (s, 2H), 3.89-3.78 (m, 6H), 2.44 (s, 6H).

(Reaction 96-18)




[0851] Potassium t -butoxide (53 mg, 0.44 mmol) was added to a solution of 2-bromo-N-(4-bromo-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide (147 mg, 0.40 mmol) in t -BuOH (2 ml), and the mixture was heated under reflux overnight. The reaction solution was cooled to room temperature and water was then added, followed by extraction with

ethyl acetate. The organic phase was washed with saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-(4-bromo-3,5-dimethyl-phenyl)-morpholin-3-one (100%).

¹H-NMR (CDCl₃) δ 7.05 (s, 2H), 4.33 (s, 2H), 4.02 (t, 2H, J = 5.0 Hz), 3.72 (t, 2H, J = 5.0 Hz), 2.42 (s, 6H).

[0852] The aryl bromide reagent used in the synthesis of Compound 516 (1-(4-bromo-3,5-dimethylphenyl)piperidin-2-one) was synthesized as follows.


(Reaction 96-19)



[0853] 5-Bromo-pentanoic (4-bromo-3,5-dimethyl-phenyl)-amide was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 7.25 (s, 2H), 7.19 (brs, 1H), 3.43 (t, 2H, J = 6.49 Hz), 2.37 (t, 2H, J = 6.87 Hz), 2.37 (s, 6H), 1.9 (m, 4H).

(Reaction 96-20)




[0854] Sodium hydride (37 mg, 0.925 mmol) was added to a solution of 5-bromo-pentanoic (4-bromo-3,5-dimethyl-phenyl)-amide (320 mg, 0.881 mmol) in DMF (8 ml) at 0°C, and the mixture was stirred at room temperature for two days. The reaction solution was diluted with ethyl acetate, and the organic layer was sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give 1-(4-bromo-3,5-dimethylphenyl)piperidin-2-one (240 mg, 97%).

¹H-NMR (CDCl₃) δ 6.97 (s, 2H), 3.58 (t, 2H, J = 6.87 Hz), 2.55 (t, 2H, J = 6.87 Hz), 2.40 (s, 6H), 1.93 (m, 4H).

[0855] The aryl bromide reagent used in the synthesis of Compound 517 (N-(4-bromo-3,5-dimethyl-benzyl)-N-methyl-acetamide) was synthesized as follows.


(Reaction 96-21)



[0856] Methylamine (12.3 ml, 24.60 mmol, 2.0 M solution in methanol) was added dropwise to a solution of 2-bromo-5-iodomethyl-1,3-dimethylbenzene (400 mg, 1.23 mmol) in anhydrous DMF (12 ml), and the mixture was stirred at room temperature for two days. The reaction solution was concentrated under reduced pressure, and the residue was

diluted with ethyl acetate. The organic layer was then sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 1-(4-bromo-3,5-dimethylphenyl)-N-methylmethanamine (280 mg, 100%).
MS (ESI) m/z = 228, 230 ($M+H$)⁺.

(Reaction 96-22)




[0857] N-(4-Bromo-3,5-dimethyl-benzyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

MS (ESI) m/z = 270, 272 ($M+H$)⁺.

[0858] The aryl bromide reagent used in the synthesis of Compound 518 (3-(4-bromo-3,5-dimethyl-benzyl)-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 96-23)




[0859] 3-(4-Bromo-3,5-dimethyl-benzyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 52-1 (using iodine as a reagent) and Reaction 25-3 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 7.10 (s, 2H), 5.63 (s, 1H), 4.55 (s, 2H), 3.97 (d, 2H, J = 1.14 Hz), 2.38 (s, 6H).

[0860] The aryl bromide reagent used in the synthesis of Compound 519 (3-(4-bromo-3,5-dimethyl-benzyl)-5,5-dimethyl-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 96-24)




[0861] 3-(4-Bromo-3,5-dimethyl-benzyl)-5,5-dimethyl-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 325, 327 ($M+H$)⁺.

[0862] The aryl bromide reagent used in the synthesis of Compound 520 (1-(4-bromo-3,5-dimethyl-benzyl)-pyrrolidin-2-one) was synthesized as follows.

(Reaction 96-25)




[0863] 1-(4-Bromo-3,5-dimethyl-benzyl)-pyrrolidin-2-one was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

¹H-NMR ($CDCl_3$) δ 6.94 (s, 2H), 4.33 (s, 2H), 3.25 (t, 2H, J = 7.24 Hz), 2.44 (t, 2H, J = 8.01 Hz), 2.38 (s, 6H), 2.04-1.93 (m, 2H).

[0864] The aryl bromide reagent used in the synthesis of Compound 521 (1-(4-bromo-3,5-dimethyl-benzyl)-pyrrolidine-2,5-dione) was synthesized as follows.

(Reaction 96-26)




[0865] 1-(4-Bromo-3,5-dimethyl-benzyl)-pyrrolidine-2,5-dione was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 7.09 (s, 2H), 4.54 (s, 2H), 2.70 (s, 4H), 2.37 (s, 6H).

[0866] The aryl bromide reagent used in the synthesis of Compound 522 (N-(4-bromo-3,5-dimethyl-benzyl)-acetamide) was synthesized as follows.

(Reaction 96-27)




[0867] N-(4-Bromo-3,5-dimethyl-benzyl)-acetamide was synthesized by operations similar to those in Reaction 96-21 and Reaction 2-3 using appropriate reagents and starting material.

MS (ESI) m/z = 256, 258 (M+H)⁺.

[0868] The aryl bromide reagent used in the synthesis of Compound 523 (N-(4-bromo-3,5-difluoro-phenyl)-N-methyl-acetamide) was synthesized as follows.

(Reaction 96-28)



[0869] N-(4-Bromo-3,5-difluoro-phenyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 19-2 and Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 264, 266 (M+H)⁺.



Example 97

[0870] The example compounds shown below were obtained by operations similar to those in Reaction 25-2 using appropriate reagents and starting materials.

(Compounds 524 to 525)


[0871]

[Table 72]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
524		LCMS-B-1	2.03	577 (M+H) ⁺
525		LCMS-F-1	0.98	690 (M+H) ⁺

[0872] The aryl bromide reagent used in the synthesis of Compound 525 ([*(S)*]-1-(4-bromo-3-methyl-phenylcarbamoyl)-2-carbamoyl-ethyl]-carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 97-1)



[0873] [*(S)*]-1-(4-Bromo-3-methyl-phenylcarbamoyl)-2-carbamoyl-ethyl]-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.
MS (ESI) m/z = 400 (M+H)⁺.

Example 98

[0874] The example compounds shown below were obtained by operations similar to those in Reaction 25-2 using appropriate reagents and starting materials.

(Compounds 526 to 534)

[0875]

[Table 73]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
526		LCMS-B-1	2.5	593 (M+H)+
527		LCMS-C-1	2.6	620 (M+H)+
528		LCMS-C-1	2.65	649 (M+H)+
529		LCMS-C-1	2.45	566 (M+H)+
530		LCMS-C-1	2.38	650 (M+H)+
531		LCMS-C-1	2.63	598 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
532		LCMS-D-1	2.16	654 (M+H)+
533		LCMS-D-1	2.35	668 (M+H)+
534		LCMS-D-1	3.1	685 (M+H)+

[0876] The aryl bromide reagent used in the synthesis of Compound 528 (2-(4-bromo-3-methyl-phenylamino)-1-(4-methyl-piperazin-1-yl)-ethanone) was synthesized as follows.

(Reaction 98-1)




[0877] 2-(4-Bromo-3-methyl-phenylamino)-1-(4-methyl-piperazin-1-yl)-ethanone was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 326 (M+H)+.

[0878] The aryl bromide reagent used in the synthesis of Compound 529 (2-(4-bromo-3-methyl-phenylamino)-acetamide) was synthesized as follows.

(Reaction 98-2)




[0879] 2-(4-Bromo-3-methyl-phenylamino)-acetamide was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) $m/z = 243$ ($M+H$) $^+$.

[0880] The aryl bromide reagent used in the synthesis of Compound 530 (4-bromo-3-methyl-N-pyridin-4-yl-benzenesulfonamide) was synthesized as follows.

(Reaction 98-3)




[0881] 4-Bromo-3-methyl-N-pyridin-4-yl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 327$ ($M+H$) $^+$.

[0882] The aryl bromide reagent used in the synthesis of Compound 532 (1-(4-bromo-3,5-dimethylphenyl)-1,3-bis(2-hydroxyethyl)urea) was synthesized as follows.


(Reaction 98-4)



[0883] N-(2-(Benzyloxy)ethyl)-4-bromo-3,5-dimethylaniline was synthesized by operations similar to those in Reaction 25-3 and Reaction 96-16 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 2.32 (s, 6H), 3.27 (t, $J = 5.39$ Hz, 2H), 3.68 (t, $J = 5.37$ Hz, 2H), 3.95 (brs, 1H), 4.54 (s, 2H), 6.37 (s, 2H), 7.25-7.38 (m, 5H).


(Reaction 98-5)



[0884] Phosgene (3.5 ml, 6.64 mmol, 20% solution in toluene) and N,N-diisopropylethylamine (1.2 ml, 6.64 mmol) were added to a solution of N-(2-(benzyloxy)ethyl)-4-bromo-3,5-dimethylaniline (740 mg, 2.21 mmol) in anhydrous toluene (11 ml) at 0°C. The reaction solution was gradually warmed to room temperature and stirred at the same temperature for three hours. Water was added to the reaction mixture, followed by extraction with ethyl acetate. The organic phase was sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane) to give N-(2-(benzyloxy)ethyl)-N-(4-bromo-3,5-dimethylphenyl)carbamic acid chloride (860 mg, 98%).

¹H-NMR (300 MHz, CDCl₃), δ 2.36 (s, 6H), 3.67 (t, J = 5.42 Hz, 2H), 3.89 (t, J = 4.95 Hz, 2H), 4.50 (s, 2H), 6.95 (s, 2H), 7.27-7.38 (m, 5H).


(Reaction 98-6)



[0885] 1-(2-(Benzyloxy)ethyl)-1-(4-bromo-3,5-dimethylphenyl)-3-(2-hydroxyethyl)urea was synthesized by operations similar to those in Reaction 82-1 using appropriate reagents and starting material.

MS (ESI) m/z = 421, 423 (M+H)⁺.

(Reaction 98-7)




[0886] Boron trichloride (0.93 ml, 0.93 mmol, 1.0 M solution in dichloromethane) was added to a solution of 1-(2-(benzyloxy)ethyl)-1-(4-bromo-3,5-dimethylphenyl)-3-(2-hydroxyethyl)urea (98 mg, 0.23 mmol) in anhydrous dichloromethane (4.6 ml) at -78°C, and the mixture was stirred at the same temperature for two hours. A saturated aqueous sodium

bicarbonate solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic phase was sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 1-(4-bromo-3,5-dimethylphenyl)-1,3-bis(2-hydroxyethyl)urea (64 mg, 83%).

¹H-NMR (300 MHz, CDCl₃) δ 2.43 (s, 6H), 2.94 (t, J = 5.11 Hz, 1H), 3.33 (q, J = 5.23 Hz, 2H), 3.60 (t, J = 4.78 Hz, 1H), 3.66 (q, J = 4.91 Hz, 2H), 3.71-3.81 (m, 4H), 4.73 (brs, 1H), 7.01 (s, 2H).

[0887] The aryl bromide reagent used in the synthesis of Compound 533 (1-(4-bromo-3,5-dimethyl-phenyl)-1,3-bis-(2-hydroxy-ethyl)-3-methyl-urea) was synthesized as follows.

(Reaction 98-8)




[0888] 1-(4-Bromo-3,5-dimethyl-phenyl)-1,3-bis-(2-hydroxy-ethyl)-3-methyl-urea was synthesized by operations similar to those in Reaction 82-1 and Reaction 98-7 using appropriate reagents and starting material.

MS (ESI) m/z = 345, 347 (M+H)⁺.

[0889] The aryl bromide reagent used in the synthesis of Compound 534 (N-(4-bromo-3,5-dimethyl-phenyl)-N-(tetrahydro-pyran-4-yl)-methanesulfonamide) was synthesized as follows.

(Reaction 98-9)



[0890] N-(4-Bromo-3,5-dimethyl-phenyl)-N-(tetrahydro-pyran-4-yl)-methanesulfonamide was synthesized by operations similar to those in Reaction 41-1 and Reaction 6-1 using appropriate reagents and starting material.

MS (ESI) m/z = 362, 364 (M+H)⁺.


Example 99

[0891] The example compound shown below was obtained by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

(Compound 535)

[0892]


[Table 74]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
535		LCMS-C-1	2.52	637 (M+H)+

Example 100

3-{3-Methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-imidazolidine-2,4-dione (Compound 536)


[0893]

(Reaction 100-1)

[0894] 8-Ethenesulfonyl-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-20, Reaction 1-4, Reaction 4-1, Reaction 5-3 and reaction 25-1 using appropriate reagents and starting material.

MS (ESI) m/z = 370 (M+H)+.

(Reaction 100-2)



15 **[0895]** 3-{3-Methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 558$ (M+H)+.

20 **[0896]** The example compounds shown below were synthesized by operations similar to those in Reaction 100-2 using appropriate reagents and starting materials.

(Compounds 537 to Compound 539)


25 **[0897]**

[Table 75]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
537		LCMS-A-1	2.63	612 (M+H)+
538		LCMS-A-1	2.6	573 (M+H)+
539		LCMS-A-1	2.43	572 (M+H)+

55 **[0898]** The aryl bromide reagent used in the synthesis of Compound 537 (3-(4-bromo-3-trifluoromethyl-phenyl)-imidazolidine-2,4-dione) was synthesized as follows.


(Reaction 100-3)



[0899] 3-(4-Bromo-3-trifluoromethyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 84-1 and Reaction 96-1 using appropriate reagents and starting material.
MS (ESI) $m/z = 321$ (M-H)-.

[0900] The aryl bromide reagent used in the synthesis of Compound 538 (4-(4-bromo-3-methyl-phenyl)-morpholine-3,5-dione) was synthesized as follows.


(Reaction 100-4)



[0901] [1,4]Dioxane-2,6-dione (312 mg, 2.69 mmol) was added to a solution of 4-bromo-3-methyl-phenylamine (500 mg, 2.69 mmol) in $t\text{BuOMe}$ (7.0 ml), and the mixture was stirred at room temperature overnight. The reaction solution was concentrated under reduced pressure. Acetic anhydride (4.0 ml, 42.3 mmol) and sodium acetate (35 mg, 0.427 mmol) were added to the resulting residue, and the mixture was heated with stirring at 60°C for three hours. Water was added to the reaction solution, and collection by filtration and trituration with water gave 4-(4-bromo-3-methyl-phenyl)-morpholine-3,5-dione (577 mg, 73%).
MS (ESI) $m/z = 284, 286$ (M+H)+.

[0902] The aryl bromide reagent used in the synthesis of Compound 539 (3-(4-bromo-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 100-5)



[0903] 3-(4-Bromo-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in

Reaction 84-1 and Reaction 96-1 using appropriate reagents and starting material.


MS (ESI) m/z = 283, 285 ($M+H$)⁺.

Example 101

5,5-Dimethyl-3-[3-methyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-imidazolidine-2,4-dione (Compound 540)

[0904]


(Reaction 101-1)



[0905] $n\text{-Butyllithium}$ (1.6 M solution in hexane, 26 ml, 41.6 mmol) was added to a solution of triphenyl-(3,3,3-trifluoro-propyl)-phosphonium iodide (20.25 g, 41.64 mmol) in THF (141 ml) at -78°C over 13 minutes, and the mixture was stirred at the same temperature for 20 minutes. A solution of 4-oxo-cyclohexanecarboxylic acid ethyl ester (6.56 g, 38.54 mmol) in THF (22 ml) was added to the reaction solution at -78°C over 17 minutes, and the mixture was stirred at the same temperature for one hour. A 50% saturated aqueous ammonium chloride solution was added, followed by extraction with dichloromethane. The organic layer was then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give ethyl 4-(3,3,3-trifluoropropylidene)-cyclohexanecarboxylate (9.25 g, 96%).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.25 (3H, t, $J = 7.1$ Hz), 1.57 (2H, m), 1.90 (1H, m), 2.01 (2H, m), 2.11 (1H, m), 2.31 (1H, m), 2.49 (2H, m), 2.80 (2H, m), 4.13 (2H, t, $J = 7.1$ Hz), 5.15 (1H, t, $J = 7.6$ Hz).

(Reaction 101-2)




[0906] 4-Carbamoyl-4-[[4-(3,3,3-trifluoro-propyl)-cyclohexanecarbonyl]-amino]-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 18-2 (using PtO_2 as a catalyst), Reaction 95-18 and Reaction 10-14 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 0.98 (2H, m), 1.25-1.70 (9H, m), 1.45 (9H, s), 1.88 (4H, m), 2.10 (3H, m), 3.08 (2H, m), 3.81 (2H, m), 5.30 (1H, br), 5.40 (1H, s), 7.15 (1H, br).

(Reaction 101-3)


[0907] LiOH·H₂O (1.55 g, 36.9 mmol) was added to a solution of 4-carbamoyl-4-[(4-(3,3,3-trifluoro-propyl)-cyclohexanecarbonyl]-amino)-piperidine-1-carboxylic acid tert-butyl ester (5.53 g, 12.3 mmol) in ethanol (123 mL), and the mixture was stirred at 85°C for two hours. A 50% saturated aqueous ammonium chloride solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by reprecipitation with hexane-ethyl acetate = 3:1 to give 4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (4.88 g, 92%). ¹H-NMR (400 MHz, CDCl₃) δ 1.05 (2H, m), 1.25-1.60 (7H, m), 1.45 (9H, s), 1.81 (2H, m), 1.90 (2H, m), 2.02 (2H, m), 2.11 (2H, m), 2.36 (1H, m), 3.40 (2H, m), 3.90 (2H, m), 8.10 (1H, br).

(Reaction 101-4)

[0908] 8-Ethenesulfonyl-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 4-1, Reaction 5-3 and Reaction 25-1 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.05 (2H, m), 1.25-1.70 (7H, m), 1.89 (2H, m), 2.00 (4H, m), 2.11 (2H, m), 2.39 (1H, m), 3.25 (2H, m), 3.67 (2H, m), 6.03 (1H, d, J = 10.0 Hz), 6.26 (1H, d, J = 16.0 Hz), 6.03 (1H, dd, J = 16.0 and 10.0 Hz), 8.50 (1H, br).

(Reaction 101-5)



[0909] 5,5-Dimethyl-3-[3-methyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) m/z = 638 ($M+H$) $^+$.

[0910] The example compounds shown below were synthesized by operations similar to those in Example 101 using appropriate reagents and starting materials.

(Compounds 541 to Compound 559)

[0911]

[Table 76]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
541	Chiral	LCMS-F-1	0.94	611 ($M+H$) $^+$
542		LCMS-D-1	2.30	627 ($M+H$) $^+$
543		LCMS-D-1	2.42	641 ($M+H$) $^+$

10

15

20

25

30








35

40




45

50

(continued)


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
550		LCMS-D-1	2.23	625 (M+H) ⁺
551		LCMS-D-1	2.55	673 (M+H) ⁺
552		LCMS-D-1	2.02	626 (M+H) ⁺
553		LCMS-D-1	2.40	556 (M+H) ⁺
554		LCMS-D-1	2.45	638 (M+H) ⁺
555		LCMS-D-1	2.67	666 (M+H) ⁺
556		LCMS-D-1	2.58	623 (M+H) ⁺

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
557		LCMS-D-1	2.42	637 (M+H)+
558		LCMS-D-1	2.78	597 (M+H)+
559		LCMS-D-1	2.45	611 (M+H)+

[0912] The aryl bromide reagent used in the synthesis of Compound 542 (N-(4-bromo-3-methyl-phenyl)-N-(2-methoxy-ethyl)-acetamide) was synthesized as follows.

(Reaction 101-6)




[0913] (N-(4-Bromo-3-methyl-phenyl)-N-(2-methoxy-ethyl)-acetamide) was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 286, 288 (M+H)+.

[0914] The aryl bromide reagent used in the synthesis of Compound 543 (N-(4-bromo-3,5-dimethyl-phenyl)-N-(2-methoxy-ethyl)-acetamide) was synthesized as follows.

(Reaction 101-7)




[0915] N-(4-Bromo-3,5-dimethyl-phenyl)-N-(2-methoxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 6.93 (s, 2H), 3.82 (t, 2H, J = 5.7 Hz), 3.49 (t, 2H, J = 5.7 Hz), 3.30 (s, 2H), 2.42 (s, 6H), 1.85 (s, 3H).

[0916] The aryl bromide reagent used in the synthesis of Compound 544 (N-(4-bromo-3-methyl-phenyl)-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide) was synthesized as follows.


(Reaction 101-8)



[0917] Sodium hydride (60% oil suspension, 100 mg, 2.63 mmol) was added to a solution of 4-bromo-3-methylphenylacetamide (500 mg, 2.19 mmol), tert-butyl[2-(2-chloroethoxy)ethoxy]dimethylsilane (excess) and sodium iodide (324 mg, 2.19 mmol) in dimethylformamide (20 ml). The mixture was heated with stirring at 100°C for 17 hours. The reaction solution was cooled and then concentrated. The resulting residue was purified by silica gel column chromatography (ethyl acetate-hexane) to give N-(4-bromo-3-methyl-phenyl)-N-[2-(2-(tert-butyldimethyl-silanyloxy)-ethoxy)-ethyl]-acetamide (555 mg, 65%).

[0918] ¹H-NMR (CDCl₃) δ 0.04 (s, 6H), 0.87 (s, 9H), 1.83 (s, 3H), 2.39 (s, 3H), 3.40-3.50 (m, 2H), 3.49-3.66 (m, 2H), 3.63-3.77 (m, 2H), 3.73-3.89 (m, 2H), 6.76-7.00 (m, 1H), 6.98-7.16 (m, 1H), 7.38-7.63 (m, 1H).

(Reaction 101-9)




[0919] N-(4-Bromo-3-methyl-phenyl)-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide was synthesized by operations similar to those in Reaction 39-2 using appropriate reagents and starting material.

MS (ESI) m/z = 316, 318 (M+H)⁺.

[0920] The aryl bromide reagent used in the synthesis of Compound 545 (N-(4-bromo-3-methyl-phenyl)-N-[2-(2-fluoroethoxy)-ethyl]-acetamide) was synthesized as follows.

(Reaction 101-10)




[0921] N-(4-Bromo-3-methyl-phenyl)-N-[2-(2-fluoro-ethoxy)-ethyl]-acetamide was synthesized by operations similar to those in Reaction 25-15 using appropriate reagents and starting material.

MS (ESI) m/z = 318, 320 ($M+H$)⁺.

[0922] The aryl bromide reagent used in the synthesis of Compound 548 (3-(4-bromo-3,5-dimethyl-phenyl)-1-methyl-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 101-11)




[0923] 3-(4-Bromo-3,5-dimethyl-phenyl)-1-methyl-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 96-6 using appropriate reagents and starting material.

MS (ESI) m/z = 297, 299 ($M+H$)⁺.

[0924] The aryl bromide reagent used in the synthesis of Compound 549 ((4-bromo-3,5-dimethyl-phenyl)-[4-(2-fluoro-ethyl)-piperazin-1-yl]-methanone) was synthesized as follows.

(Reaction 101-12)




[0925] (4-Bromo-3,5-dimethyl-phenyl)-[4-(2-fluoro-ethyl)-piperazin-1-yl]-methanone was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

¹H-NMR (CDCl_3) δ 7.10 (s, 2H), 4.66 (t, 1H, J = 4.96 Hz), 4.51 (t, 1H, J = 4.96 Hz), 3.79 (s, 2H), 3.47 (s, 2H), 2.79 (t, 1H, J = 4.96 Hz), 2.70 (t, 1H, J = 4.96 Hz), 2.56 (brs, 4H), 2.43 (s, 6H).


[0926] The aryl bromide reagent used in the synthesis of Compound 551 (N-(4-bromo-3,5-dimethyl-phenyl)-N-[2-(2-fluoroethoxy)ethyl]acetamide) was synthesized as follows.

(Reaction 101-13)



[0927] 2-[2-[Acetyl-(4-bromo-3,5-dimethyl-phenyl)amino]ethoxy]ethyl methanesulfonate was synthesized by operations similar to those in Reaction 39-2 and Reaction 5-4 using appropriate reagents and starting material. MS (ESI) m/z = 408, 410 ($\text{M}+\text{H}$) $^+$.

(Reaction 101-14)




[0928] Potassium fluoride (180 mg, 3.11 mmol) was added to a solution of 2-[2-[acetyl-(4-bromo-3,5-dimethyl-phenyl)amino]ethoxy]ethyl methanesulfonate (254 mg, 0.622 mmol) in PEG200 (2 ml), and the mixture was irradiated with microwaves at 100°C for 10 minutes. The reaction solution was diluted with ethyl acetate, and the organic layer was sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give N-(4-bromo-3,5-dimethyl-phenyl)-N-[2-(2-fluoroethoxy)ethyl]acetamide (144 mg, 70%).

MS (ESI) m/z = 332, 334 ($\text{M}+\text{H}$) $^+$.

[0929] The aryl bromide reagent used in the synthesis of Compound 552 (N'-(4-bromo-3,5-dimethyl-benzoyl)-hydrazinecarboxylic acid tert-butyl ester) was synthesized as follows.

(Reaction 101-15)




[0930] N'-[4-Bromo-3,5-dimethyl-benzoyl]-hydrazinecarboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.
MS (ESI) m/z = 343, 345 ($M+H$)⁺.

Example 102


3-(4-[(E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3-methyl-phenyl)-imidazolidine-2,4-dione (Compound 560)

[0931]

(Reaction 102-1)



[0932] 2-(4-Butyl-cyclohexyl)-8-ethenesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14, Reaction 1-4, Reaction 4-1, Reaction 5-3 and Reaction 25-1 using appropriate reagents and starting material.
MS (ESI) m/z = 382 ($M+H$)⁺.

(Reaction 102-2)

[0933] 3-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) m/z = 570 (M+H)+.

[0934] The example compounds shown below were synthesized by operations similar to those in Reaction 102-2 using appropriate reagents and starting materials.

(Compounds 561 to Compound 562)


[0935]

[Table 77]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
561		LCMS-D-1	2.17	645 (M+H)+
562		LCMS-D-1	3.25	633 (M+H)+

[0936] The aryl bromide reagent used in the synthesis of Compound 561 (N-(4-bromo-3,5-dimethyl-phenyl)-N-[2-(2-methoxy-ethoxy)-ethyl]-acetamide) was synthesized as follows.

(Reaction 102-3)



[0937] N-(4-Bromo-3,5-dimethyl-phenyl)-N-[2-(2-methoxy-ethoxy)-ethyl]-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.


MS (ESI) m/z = 334, 336 ($M+H$)⁺.

Example 103

3-(4-((E)-2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidine-2,4-dione (Compound 563)


[0938]

(Reaction 103-1)



[0939] 8-Ethenesulfonyl-2-(4-isopropyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14, Reaction 1-4, Reaction 4-1, Reaction 5-3 and Reaction 25-1 using appropriate reagents and starting material.

MS (ESI) m/z = 368 ($M+H$)⁺.

(Reaction 103-2)

[0940] 3-(4-((E)-2-[2-(4-Isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) m/z = 556 (M+H)+.

[0941] The example compound shown below was synthesized by operations similar to those in Reaction 103-2 using appropriate reagents and starting material.

(Compound 564)

[0942]

[Table 78]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
564		LCMS-C-1	2.72	570 (M+H)+







Example 104

[0943] The example compounds shown below were obtained by operations similar to those in Reaction 26-1 using appropriate reagents and starting materials.





(Compounds 565 to 574)

[0944]

[Table 79]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
565		LCMS-C-1	2.25	502 (M+H)+
566		LCMS-C-1	2.47	503 (M+H)+
567		LCMS-C-1	2.28	555 (M+H)-
568		LCMS-C-1	2.43	538 (M+H)+
569		LCMS-C-1	2.8	628 (M-H)-
570		LCMS-C-1	2.6	527 (M-H)-

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
571		LCMS-C-1	2.17	584 (M-H)-
572		LCMS-C-1	2.47	485 (M+H)+
573		LCMS-D-1	2.82	528 (M+H)+
574		LCMS-D-1	2.82	571 (M+H)+

[0945] The aryl bromide reagent used in the synthesis of Compound 567 (4-methyl-piperazine-1-carboxylic acid 4-bromo-benzylamide) was synthesized as follows.

(Reaction 104-1)




[0946] 4-Methyl-piperazine-1-carboxylic acid 4-bromo-benzylamide was synthesized by operations similar to those in

Reaction 82-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 312$ (M+H)+.

[0947] The aryl bromide reagent used in the synthesis of Compound 568 (N'-(4-bromo-3-methyl-phenyl)-N,N-dimethyl-sulfamide) was synthesized as follows.

(Reaction 104-2)




[0948] N'-(4-Bromo-3-methyl-phenyl)-N,N-dimethyl-sulfamide was synthesized by operations similar to those in Reaction 82-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 293$ (M+H)+.

[0949] The aryl bromide reagent used in the synthesis of Compound 569 ((4-bromo-3-methyl-benzyl)-(3-dimethylamino-propyl)-carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 104-3)




[0950] (4-Bromo-3-methyl-benzyl)-(3-dimethylamino-propyl)-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) $m/z = 385, 387$ (M+H)+.

[0951] The aryl bromide reagent used in the synthesis of Compound 570 (2-(4-bromo-phenyl)-N-(2,2-dimethyl-propyl)-acetamide) was synthesized as follows.

(Reaction 104-4)




[0952] 2-(4-Bromo-phenyl)-N-(2,2-dimethyl-propyl)-acetamide was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 285, 287 ($M+H$)⁺.

[0953] The aryl bromide reagent used in the synthesis of Compound 571 (2-(4-bromo-phenyl)-N-(2-diisopropylamino-ethyl)acetamide) was synthesized as follows.

(Reaction 104-5)




[0954] 2-(4-Bromo-phenyl)-N-(2-diisopropylamino-ethyl)acetamide was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 342, 344 ($M+H$)⁺.

[0955] The aryl bromide reagent used in the synthesis of Compound 572 (1-(4-bromo-3-methyl-phenyl)-azetidin-2-one) was synthesized as follows.

(Reaction 104-6)




[0956] 1-(4-Bromo-3-methyl-phenyl)-azetidin-2-one was synthesized by operations similar to those in Reaction 29-3 using appropriate reagents and starting material.

MS (ESI) m/z = 240, 242 ($M+H$)⁺.

[0957] The aryl bromide reagent used in the synthesis of Compound 573 ((4-bromo-3,5-dimethyl-phenyl)-thiazol-2-yl-amine) was synthesized as follows.


(Reaction 104-7)



20 **[0958]** A solution of 4-bromo-3,5-dimethyl-phenylamine (200 mg) and benzoyl isothiocyanate (0.14 ml) in acetone (2 ml) was heated under reflux for 30 minutes. After cooling the reaction solution, a 1 N aqueous sodium hydroxide solution (2.19 ml) was added and the mixture was stirred at 50°C for 12 hours. The mixture was extracted with dichloromethane, and the organic layer was then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was triturated with hexane to give (4-bromo-3,5-dimethyl-phenyl)-thiourea (146 mg, 57%).

25 ¹H-NMR (300 MHz, DMSO-d₆) δ 9.62 (1H, s), 7.8-7.2 (2H, br), 7.19 (2H, s), 2.32 (6H, s).

(Reaction 104-8)




40 **[0959]** (4-Bromo-3,5-dimethyl-phenyl)-thiazol-2-yl-amine was synthesized by operations similar to those in Reaction 94-1 using appropriate reagents and starting material.

MS (ESI) m/z = 285 (M+H)⁺.

[0960] The aryl bromide reagent used in the synthesis of Compound 574 (N-(4-bromo-3,5-dimethyl-phenyl)-N-(3-methyl-oxetan-3-ylmethyl)-acetamide) was synthesized as follows.

(Reaction 104-9)




[0961] N-(4-Bromo-3,5-dimethyl-phenyl)-N-(3-methyl-oxetan-3-ylmethyl)-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.
MS (ESI) m/z = 326, 328 (M+H)+.

Example 105

3-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione (Compound 575)

[0962]

(Reaction 105-1)



[0963] 3-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.
MS (ESI) m/z = 542 (M+H)+.

[0964] The example compounds shown below were synthesized by operations similar to those in Example 105 using appropriate reagents and starting materials.

(Compounds 576 to Compound 589)

[0965]







[Table 80]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
576		LCMS-D-1	2.82	531 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
577		LCMS-C-1	2.62	515 (M+H)+
578		LCMS-A-1	2.44	628 (M+H)+
579		LCMS-A-1	2.23	529 (M+H)+
580		LCMS-C-1	2.72	515 (M+H)+
581		LCMS-C-1	2.42	542 (M+H)+

(continued)


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
582		LCMS-A-1	2.05	556 (M+H)+
583		LCMS-A-1	1.95	542 (M+H)+
584		LCMS-D-1	1.93	515 (M+H)+
585		LCMS-F-1	1.05	460 (M+H)+
586		LCMS-A-1	1.83	539 (M+H)+
587		LCMS-D-1	2.02	556 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
588		LCMS-C-1	2.53	515 (M+H)+
589		LCMS-C-1	2.83	628 (M+H)+

[0966] The aryl bromide reagent used in the synthesis of Compound 577 (3-(4-bromo-3-methyl-phenyl)-oxazolidin-4-one) was synthesized as follows.


(Reaction 105-2)



[0967] Acetic acid chlorocarbonylmethyl ester (1.73 ml) was added to a solution of 4-bromo-3-methyl-phenylamine (2.0 g, 10.7 mmol) and pyridine (5.21 ml) in dichloromethane (20 ml), and the mixture was stirred at 40°C for 2.5 hours. The mixture was cooled, and then quenched with water and extracted with dichloromethane. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure to give acetic acid (4-bromo-3-methyl-phenylcarbamoyl)-methyl ester (3.19 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.24 (3H, s), 2.39 (3H, s), 4.68 (2H, s), 7.22-7.25 (1H, m), 7.46-7.50 (2H, m), 7.70 (1H, brs).


(Reaction 105-3)



[0968] N-(4-Bromo-3-methyl-phenyl)-2-hydroxy-acetamide was synthesized by operations similar to those in Reaction 12-5 using appropriate reagents and starting material.

MS (ESI) m/z = 244, 246 ($M+H$)⁺.

(Reaction 105-4)




[0969] Sodium hydride (81 mg, 1.80 mmol) was added to a solution of N-(4-bromo-3-methyl-phenyl)-2-hydroxy-acetamide (200 mg, 0.819 mmol) in DMF (4.0 ml), and the mixture was stirred at room temperature for 50 minutes. Further, dibromomethane (0.114 ml, 1.64 mmol) was added to the reaction solution, and the mixture was heated with stirring at 110°C for two hours. Cooling to room temperature and subsequent purification by silica gel column chromatography (hexane-ethyl acetate) gave 3-(4-bromo-3-methyl-phenyl)-oxazolidin-4-one (45 mg, 21%).

MS (ESI) m/z = 256, 258 ($M+H$)⁺.

[0970] The aryl bromide reagent used in the synthesis of Compound 578 (4-(4-bromo-3-methyl-phenyl)-3-oxo-piperazine-1-carboxylic acid tert-butyl ester) was synthesized as follows.

(Reaction 105-5)




[0971] 4-(4-Bromo-3-methyl-phenyl)-3-oxo-piperazine-1-carboxylic acid tert-butyl ester was synthesized by operations

similar to those in Reaction 29-3 using appropriate reagents and starting material.

MS (ESI) m/z = 369, 371 (M+H)+.

[0972] The aryl bromide reagent used in the synthesis of Compound 579 (3-(4-bromo-3-methyl-phenyl)-4-methyl-oxazolidin-2-one) was synthesized as follows.

(Reaction 105-6)




[0973] 3-(4-Bromo-3-methyl-phenyl)-4-methyl-oxazolidin-2-one was synthesized by operations similar to those in Reaction 12-1 and Reaction 96-13 using appropriate reagents and starting material.

MS (ESI) m/z = 270, 272 (M+H)+.

[0974] The aryl bromide reagent used in the synthesis of Compound 580 (2-(4-bromo-3-methyl-phenyl)-isoxazolidin-3-one) was synthesized as follows.


(Reaction 105-7)



[0975] 3-Chloro-propionyl (157 μL , 1.65 mmol) was added to a mixture of N-m-tolyl-hydroxylamine (235 mg, 1.69 mmol) and potassium carbonate (223 mg, 1.69 mmol) in N,N-dimethylformamide (1.7 mL) at -10°C . The mixture was stirred at room temperature for 2.5 hours, and water and ethyl acetate were then added. The organic layer and the aqueous layer were separated, and the aqueous layer was repeatedly extracted with ethyl acetate three times. The organic layers were combined, washed with water twice and saturated brine, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 2-m-tolyl-isoxazolidin-3-one as a pale yellow solid (213 mg, 73%).

MS (ESI) m/z = 178 (M+H)+.

(Reaction 105-8)




[0976] 2-(4-Bromo-3-methyl-phenyl)-isoxazolidin-3-one was synthesized by operations similar to those in Reaction 26-2 using appropriate reagents and starting material.

MS (ESI) m/z = 297, 299 (M+H)+.

[0977] The aryl iodide reagent used in the synthesis of Compound 581 (3-(4-iodo-2,5-dimethyl-phenyl)-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 105-9)




[0978] 3-(4-Iodo-2,5-dimethyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 84-1 and Reaction 96-1 using appropriate reagents and starting material.

MS (ESI) m/z = 331 (M+H)+.

[0979] The aryl bromide reagent used in the synthesis of Compound 582 (3-(4-bromo-3,5-dimethyl-phenyl)-5-methyl-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 105-10)




[0980] 3-(4-Bromo-3,5-dimethyl-phenyl)-5-methyl-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 10-1, Reaction 39-2 and Reaction 96-10 using appropriate reagents and starting material.

MS (ESI) m/z = 297, 299 (M+H)+.

[0981] The aryl bromide reagent used in the synthesis of Compound 583 (3-(4-bromo-2,6-dimethyl-phenyl)-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 105-11)




[0982] Ethyl isocyanatoacetate (581 mg, 4.50 mmol) and N,N-diisopropylethylamine (426 mg, 1.65 mmol) were added to a solution of 4-bromo-2,6-dimethylaniline (600 mg, 3.00 mmol) in toluene (6 ml) with stirring in a nitrogen stream, and the mixture was heated with stirring at 120°C. After 30 minutes, the reaction solution was brought to room temperature, and the precipitate was collected by filtration, washed with toluene and then dried under reduced pressure. The resulting solid was suspended in toluene (6 ml). DBU (68.4 mg, 2.25 mmol) was added and the mixture was heated with stirring at 120°C. After 30 minutes, the reaction solution was brought to room temperature and diluted with ethyl acetate, and the organic layer was washed with a 1 N aqueous hydrochloric acid solution and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the magnesium sulfate was then removed by filtration. The filtrate was concentrated under reduced pressure. The residue was subjected to column chromatography (hexane-ethyl acetate) to give 3-(4-bromo-2,6-dimethyl-phenyl)-imidazolidine-2,4-dione (570 mg, 67%).

MS (ESI) $m/z = 283, 285 (M+H)^+$.

[0983] The aryl bromide reagent used in the synthesis of Compound 584 (N-(4-bromo-2,6-dimethyl-phenyl)-N-methyl-acetamide) was synthesized as follows.

(Reaction 105-12)




[0984] N-(4-Bromo-2,6-dimethyl-phenyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 19-2 (using DMAP as a base) and Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) $m/z = 256, 258 (M+H)^+$.

[0985] The aryl bromide reagent used in the synthesis of Compound 586 ((4-bromo-3,5-dimethyl-phenyl)-(1H-imidazol-2-yl)-methyl-amine) was synthesized as follows.

(Reaction 105-13)




[0986] Iodomethane (260 mg, 9.15 mmol) was added to a solution of 1-(4-bromo-3,5-dimethyl-phenyl)-1-methyl-thiourea (500 mg, 1.83 mmol) in acetone (10 ml) with stirring in a nitrogen stream, and the mixture was heated with stirring at 50°C for two hours. The reaction solution was concentrated under reduced pressure, and a mixture of the resulting residue and aminoacetaldehyde dimethylacetal (250 mg, 2.38 mmol) in iso-BuOH (8.3 ml) was then heated under reflux for four hours. The reaction mixture was concentrated under reduced pressure. Concentrated hydrochloric acid (3 ml) was then added to the resulting residue, and the mixture was heated with stirring at 90°C for 30 minutes. The reaction mixture was cooled and then adjusted to pH 10 with a 2 N aqueous sodium hydroxide solution, followed by extraction with dichloromethane. The organic layer was dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 1-(4-bromo-3,5-dimethyl-phenyl)-(1H-imidazol-2-yl)-methyl-amine (145 mg, 28%).

¹H-NMR (400 MHz, DMSO-d₆) δ 2.30 (6H, s), 3.25 (3H, s), 6.71 (1H, s), 6.83 (2H, s), 6.87 (1H, s).

MS (ESI) m/z = 280 (M+H)⁺.

[0987] The aryl bromide reagent used in the synthesis of Compound 587 (1-(4-bromo-3,5-dimethyl-phenyl)-dihydropyrimidine-2,4-dione) was synthesized as follows.

(Reaction 105-14)




[0988] 3-[1-(4-Bromo-3,5-dimethyl-phenyl)-ureido]-propionic acid ethyl ester was synthesized by operations similar to those in Reaction 25-12 (using 1,4-dioxane as a solvent and 2,6-lutidine as a base) and Reaction 89-2 using appropriate

reagents and starting material.

$^1\text{H-NMR}$ (CDCl_3) δ 6.99 (s, 2H), 4.43 (brs, 2H), 4.05 (q, 2H, $J = 7.25$ Hz), 3.94 (t, 2H, $J = 7.25$ Hz), 2.57 (t, 2H, $J = 7.25$ Hz), 2.42 (s, 6H), 1.20 (t, 3H, $J = 7.25$ Hz).

(Reaction 105-15)




[0989] A solution of sodium ethoxide (15.7 mg, 0.033 mmol) in ethanol (0.1 ml) was added to a solution of 3-[1-(4-bromo-3,5-dimethyl-phenyl)-ureido]-propionic acid ethyl ester (11.4 mg, 0.033 mmol) in ethanol (0.9 ml), and the mixture was stirred at room temperature for 24 hours. The mixture was adjusted to pH 4 with 1 N hydrochloric acid and then extracted with dichloromethane. The organic layer was dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give 1-(4-bromo-3,5-dimethyl-phenyl)-dihydro-pyrimidine-2,4-dione (9.8 mg, 99%).

$^1\text{H-NMR}$ (CDCl_3) δ 7.59 (s, 1H), 7.03 (s, 2H), 3.83 (t, 2H, $J = 6.49$ Hz), 2.83 (t, 2H, $J = 6.49$ Hz), 2.42 (s, 6H).

[0990] The aryl bromide reagent used in the synthesis of Compound 588 (2-(4-bromo-3-methyl-phenyl)-pyrazolidin-3-one) was synthesized as follows.

(Reaction 105-16)




[0991] Triethylamine (625 μL , 13.1 mmol) and acrylic acid ethyl ester (1.82 mL, 5.74 mmol) were added to a solution of (4-bromo-3-methyl-phenyl)-hydrazine (1.05 g, 5.22 mmol) in EtOH (26.1 mL) at room temperature in a nitrogen atmosphere, and the mixture was stirred at 80°C for 18 hours. The reaction solution was cooled, and 50% NaH (501 mg, 10.4 mmol) was then added to the reaction solution at 0°C. The mixture was stirred at 0°C for 30 minutes, and 50% NaH (251 mg, 5.20 mmol) was then further added, followed by further stirring for 30 minutes. The reaction mixture was quenched with a saturated aqueous ammonium chloride solution and extracted with ethyl acetate three times. The organic layers were combined and washed with a mixed solution of water:saturated brine (1:1). After separation, the organic layer was dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 2-(4-bromo-3-methyl-phenyl)-pyrazolidin-3-one as a brown form (684 mg, 60%).

MS (ESI) $m/z = 255, 257$ ($\text{M}+\text{H}$) $^+$.

[0992] The aryl bromide reagent used in the synthesis of Compound 589 (3-(4-bromo-3,5-dimethyl-phenyl)-5-tert-butoxymethyl-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 105-17)




[0993] 3-(4-Bromo-3,5-dimethyl-phenyl)-5-tert-butoxymethyl-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 105-11 using appropriate reagents and starting material.
MS (ESI) m/z = 367, 369 (M-H)-.

Example 106

N,N-Dimethyl-2-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonylamino)-acetamide (Compound 590)

[0994]

(Reaction 106-1)








[0995] N,N-Dimethyl-2-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonylamino)-acetamide was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.
MS (ESI) m/z = 642 (M+H)+.

[0996] The example compounds shown below were synthesized by operations similar to those in Example 106 using appropriate reagents and starting materials.

(Compounds 591 to Compound 595)


[0997]

[Table 81]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
591		LCMS-C-1	2.57	648 (M+H)+
592		LCMS-C-1	2.48	655 (M+H)+
593		LCMS-C-1	2.48	556 (M+H)+
594		LCMS-A-1	2.43	599 (M+H)+
595		LCMS-C-1	2.35	661 (M-H)+

[0998] The aryl bromide reagent used in the synthesis of Compound 590 (2-(4-bromo-3-methyl-benzenesulfonylamino)-N,N-dimethyl-acetamide) was synthesized as follows.


(Reaction 106-2)



[0999] 2-(4-Bromo-3-methyl-benzenesulfonylamino)-N,N-dimethyl-acetamide was synthesized by operations similar to those in Reaction 5-4, Reaction 95-18 and Reaction 10-14 using appropriate reagents and starting material. MS (ESI) $m/z = 335, 337 (M+H)^+$.

[1000] The aryl bromide reagent used in the synthesis of Compound 591 (4-bromo-3-methyl-N-pyridin-3-ylmethyl-benzenesulfonamide) was synthesized as follows.


(Reaction 106-3)



[1001] 4-Bromo-3-methyl-N-pyridin-3-ylmethyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material. MS (ESI) $m/z = 341, 343 (M+H)^+$.

[1002] The aryl bromide reagent used in the synthesis of Compound 592 (4-bromo-N-(4-hydroxy-cyclohexyl)-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 106-4)




[1003] 4-Bromo-N-(4-hydroxy-cyclohexyl)-3-methyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 348, 350 (M+H)⁺.

[1004] The aryl bromide reagent used in the synthesis of Compound 594 (N-acetyl-4-bromo-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 106-5)




[1005] N-Acetyl-4-bromo-3-methyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 95-6 and Reaction 12-2 using appropriate reagents and starting material.

MS (ESI) m/z = 292, 294 (M+H)⁺.

[1006] The aryl bromide reagent used in the synthesis of Compound 595 (4-bromo-N-(2-hydroxy-1,1-bis-hydroxymethyl-ethyl)-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 106-6)



[1007] 4-Bromo-N-(2-hydroxy-1,1-bis-hydroxymethyl-ethyl)-3-methyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.


MS (ESI) m/z = 354, 356 (M+H)⁺.

Example 107

N-(1-Benzyl-piperidin-4-yl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonamide (Compound 596)

[1008]

(Reaction 107-1)



[1009] N-(1-Benzyl-piperidin-4-yl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzenesulfonamide was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) m/z = 746 (M+H)+.

[1010] The example compounds shown below were synthesized by operations similar to those in Example 107 using appropriate reagents and starting materials.

(Compounds 597 to Compound 618)

[1011]








[Table 82]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
597		LCMS-C-1	2.58	593 (M+H)+







(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
598		LCMS-C-1	2.77	648 (M+H)+
599		LCMS-C-1	2.45	647 (M+H)+
600		LCMS-C-1	2.6	620 (M+H)+
601		LCMS-C-1	2.67	612 (M+H)+
602		LCMS-C-1	3.17	693 (M+H)+
603		LCMS-B-1	2.4	593 (M+H)+
604		LCMS-A-1	2.85	734 (M+H)+


(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
605		LCMS-D-1	2.97	610 (M+H)+
606		LCMS-D-1	1.8	662 (M+H)+
607		LCMS-D-1	2.32	636 (M+H)+
608		LCMS-D-1	2.78	622 (M+H)+
609		LCMS-D-1	1.82	672 ((H+H)+)
610		LCMS-D-1	3.03	662 (M+H)+
611		LCMS-D-1	2.7	648 (M+H)+

(continued)


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
612		LCMS-C-1	2.55	590 (M+H)+
613		LCMS-C-1	2.8	633 (M+H)+
614		LCMS-D-1	3.43	648 (M+H)+
615		LCMS-D-1	3.27	634 (M+H)+
616		LCMS-D-1	2.15	649 (M+H)+
617		LCMS-F-1	0.93	593 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
618		LCMS-F-1	0.9	592 (M+H)+

[1012] The aryl bromide reagent used in the synthesis of Compound 596 (N-(1-benzyl-piperidin-4-yl)-4-bromo-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 107-2)




[1013] N-(1-Benzyl-piperidin-4-yl)-4-bromo-3-methyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 423, 425 (M+H)+.

[1014] The aryl bromide reagent used in the synthesis of Compound 597 ((S)-3-(5-bromo-indol-1-yl)-propane-1,2-diol) was synthesized as follows.

(Reaction 107-3)




[1015] (S)-3-(5-Bromo-indol-1-yl)-propane-1,2-diol was synthesized by operations similar to those in Reaction 25-3 and Reaction 25-4 using appropriate reagents and starting material.

MS (ESI) m/z = 270, 272 (M+H)+.

[1016] The aryl bromide reagent used in the synthesis of Compound 598 ((4-bromo-3-methyl-phenyl)-(4-isopropyl-piperazin-1-yl)-methanone) was synthesized as follows.

(Reaction 107-4)




[1017] 4-Bromo-3-methyl-phenyl-(4-isopropyl-piperazin-1-yl)-methanone was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) m/z = 325, 327 ($M+H$)⁺.

[1018] The aryl bromide reagent used in the synthesis of Compound 599 (4-bromo-N-((R)-2,3-dihydroxy-propyl)-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 107-5)




[1019] 4-Bromo-N-((R)-2,3-dihydroxy-propyl)-3-methyl-benzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 324, 326 ($M+H$)⁺.

[1020] The aryl bromide reagent used in the synthesis of Compound 600 (4-(4-bromo-2,6-difluoro-phenoxy)-butane-1,2-diol) was synthesized as follows.

(Reaction 107-6)




[1021] 4-(4-Bromo-2,6-difluoro-phenoxy)-butane-1,2-diol was synthesized by operations similar to those in Reaction 23-1 and Reaction 25-4 using appropriate reagents and starting material.

^1H -NMR (400 MHz, $CDCl_3$) δ 1.90-1.94 (2H, m), 3.57 (1H, dd, J = 12.0, 8.0 Hz), 3.74 (1H, dd, J = 12.0, 4.0 Hz), 4.06-4.26 (1H, m), 4.26-4.33 (2H, m), 7.07-7.12 (2H, m).

[1022] The aryl bromide reagent used in the synthesis of Compound 601 (4-(4-bromo-2,6-dimethyl-phenoxy)-butane-1,2-diol) was synthesized as follows.

(Reaction 107-7)




[1023] 4-(4-Bromo-2,6-dimethyl-phenoxy)-butane-1,2-diol was synthesized by operations similar to those in Reaction 23-1 and Reaction 31-6 using appropriate reagents and starting material.

^1H -NMR (400 MHz, CD_3OD) δ 1.77-1.82 (1H, m), 2.00-2.09 (1H, m), 2.45 (6H, s), 3.50-3.54 (2H, m), 3.84-3.95 (3H, m).

[1024] The aryl bromide reagent used in the synthesis of Compound 602 (4-(4-bromo-3-methyl-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester) was synthesized as follows.

(Reaction 107-8)




[1025] 4-(4-Bromo-3-methyl-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 31-7 using appropriate reagents and starting material.

^1H -NMR (400 MHz, $CDCl_3$) δ 1.50 (9H, s), 1.68-1.76 (2H, m), 1.87-1.92 (2H, m), 2.35 (1H, s), 3.30-3.36 (2H, m), 3.64-3.71 (2H, m), 4.38-4.43 (1H, m), 6.61 (1H, dd, J = 8.0, 4.0 Hz), 6.81 (1H, d, J = 4.0 Hz), 7.39 (1H, d, J = 8.0 Hz).

[1026] The aryl bromide reagent used in the synthesis of Compound 603 (N-(4-bromo-3-methyl-phenyl)-N-isopropyl-acetamide) was synthesized as follows.

(Reaction 107-9)




[1027] N-(4-Bromo-3-methyl-phenyl)-N-isopropyl-acetamide was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

MS (ESI) m/z = 270, 272 ($M+H$)⁺.

[1028] The aryl bromide reagent used in the synthesis of Compound 604 (4-[acetyl-(4-bromo-3-methyl-phenyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester) was synthesized as follows.

(Reaction 107-10)




[1029] 4-[Acetyl-(4-bromo-3-methyl-phenyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 41-1 and Reaction 12-2 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.13-1.30 (2H, m), 1.40 (9H, s), 1.70-1.80 (2H, m), 1.76 (3H, s), 2.42 (3H, s), 2.72-2.84 (2H, m), 4.07-4.20 (2H, m), 4.70-4.80 (1H, m), 6.77 (1H, dd, J = 2.8, 8.4 Hz), 6.94 (1H, d, J = 2.8 Hz), 7.57 (1H, d, J = 8.4 Hz).

[1030] The aryl bromide reagent used in the synthesis of Compound 605 (1-(4-bromo-3,5-dimethyl-phenyl)-1-(2-hydroxy-ethyl)-urea) was synthesized as follows.

(Reaction 107-11)




[1031] 1-(4-Bromo-3,5-dimethyl-phenyl)-1-(2-hydroxy-ethyl)-urea was synthesized by operations similar to those in Reaction 82-1 and Reaction 98-7 using appropriate reagents and starting material.

MS (ESI) m/z = 287, 289 ($M+H$)⁺.


[1032] The aryl bromide reagent used in the synthesis of Compound 606 ((R)-1-(4-bromo-3,5-dimethyl-phenyl)-5-(isopropylamino-methyl)-pyrrolidin-2-one) was synthesized as follows.

(Reaction 107-12)



[1033] Methanesulfonic acid (R)-1-(4-bromo-3,5-dimethylphenyl)-5-oxo-pyrrolidin-2-ylmethyl ester was synthesized by operations similar to those in Reaction 39-2 and Reaction 5-4 using appropriate reagents and starting material. ¹H-NMR (CDCl₃) δ 7.12 (s, 2H), 4.43 (m, 1H), 4.19 (m, 2H), 2.93 (s, 3H), 2.70 (m, 1H), 2.60 (m, 1H), 2.41 (s, 6H), 2.40 (m, 1H), 2.16 (m, 1H).

(Reaction 107-13)




[1034] Sodium iodide (catalytic amount) and isopropylamine (1.37 g, 23 mmol) were added to a solution of methanesulfonic acid (R)-1-(4-bromo-3,5-dimethylphenyl)-5-oxo-pyrrolidin-2-ylmethyl ester (150 mg, 0.399 mmol) in THF (3 ml) at room temperature, and the mixture was heated with stirring at 70°C for 1.5 days. After cooling to room temperature, the reaction solution was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give (R)-1-(4-bromo-3,5-dimethylphenyl)-5-(isopropylamino-methyl)-pyrrolidin-2-one (83 mg, 61%).

¹H-NMR (CDCl₃) δ 7.12 (s, 2H), 4.23 (m, 1H), 2.67 (m, 4H), 2.53 (m, 1H), 2.41 (s, 6H), 2.30 (m, 1H), 2.03 (m, 1H), 0.97 (dd, 6H, J = 2.67 Hz, J = 6.1 Hz).

[1035] The aryl bromide reagent used in the synthesis of Compound 607 (N-(4-bromo-3,5-dimethylphenyl)-N-(2-dimethylamino-ethyl)-acetamide) was synthesized as follows.

(Reaction 107-14)




[1036] N-(4-Bromo-3,5-dimethylphenyl)-N-(2-dimethylaminoethyl)-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 313, 315 ($M+H$)⁺.

[1037] The aryl bromide reagent used in the synthesis of Compound 608 (N-(4-bromo-3,5-dimethyl-phenyl)-2-dimethylamino-N-methyl-acetamide) was synthesized as follows.

(Reaction 107-15)




[1038] N-(4-Bromo-3,5-dimethyl-phenyl)-2-dimethylamino-N-methyl-acetamide was synthesized by operations similar to those in Reaction 2-3 and Reaction 95-17 using appropriate reagents and starting material.

MS (ESI) m/z = 299, 301 ($M+H$)⁺.

[1039] The aryl bromide reagent used in the synthesis of Compound 609 (N-(4-bromo-3,5-dimethyl-phenyl)-N-(2-dimethylamino-ethyl)-methanesulfonamide) was synthesized as follows.

(Reaction 107-16)




[1040] N-(4-Bromo-3,5-dimethyl-phenyl)-N-(2-dimethylaminoethyl)-methanesulfonamide was synthesized by operations similar to those in Reaction 6-1 and Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 349, 351 ($M+H$)⁺.

[1041] The aryl bromide reagent used in the synthesis of Compound 610 ((R)-5-(aminomethyl)-1-(4-bromo-3,5-dimethylphenyl)pyrrolidin-2-one) was synthesized as follows.

(Reaction 107-17)




[1042] (R)-5-(Aminomethyl)-1-(4-bromo-3,5-dimethylphenyl)pyrrolidin-2-one was synthesized by operations similar to those in Reaction 107-13 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 7.10 (s, 2H), 4.20 (m, 1H), 2.80 (m, 2H), 2.60 (m, 2H), 2.41 (s, 6H), 2.30 (m, 1H), 2.03 (m, 1H).

[1043] The aryl bromide reagent used in the synthesis of Compound 611 ((R)-1-(4-bromo-3,5-dimethyl-phenyl)-5-dimethylaminomethyl-pyrrolidin-2-one) was synthesized as follows.

(Reaction 107-18)




[1044] (R)-1-(4-Bromo-3,5-dimethyl-phenyl)-5-dimethylaminomethyl-pyrrolidin-2-one was synthesized by operations similar to those in Reaction 80-1 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 7.14 (s, 2H), 4.22 (m, 1H), 2.58 (m, 2H), 2.4 (s, 6H), 2.37 (m, 2H), 2.28 (m, 1H), 2.23 (s, 6H), 2.10 (m, 1H).

[1045] The aryl bromide reagent used in the synthesis of Compound 612 (N-(4-bromo-3-methyl-phenyl)-N-cyanomethyl-acetamide) was synthesized as follows.


(Reaction 107-19)



[1046] (4-Bromo-3-methyl-phenylamino)-acetonitrile was synthesized by operations similar to those in Reaction 95-17 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 2.36 (3H, s), 3.86-3.93 (1H, m), 4.09 (2H, d, J = 7.8 Hz), 6.43 (1H, dd, J = 8.8, 3.8 Hz), 6.60 (1H, d, J = 3.8 Hz), 7.38 (1H, d, J = 8.8 Hz).

(Reaction 107-20)




[1047] Acetic anhydride (4.99 ml) was added to (4-bromo-3-methyl-phenylamino)-acetonitrile (594 mg, 2.64 mmol),

and the mixture was heated with stirring at 115°C for one hour. The reaction solution was cooled and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give N-(4-bromo-3-methyl-phenyl)-N-cyanomethyl-acetamide (697 mg, 99%).
MS (ESI) m/z = 267, 269 (M+H)+.


[1048] The aryl bromide reagent used in the synthesis of Compound 613 (N-(4-bromo-3-methyl-phenyl)-N-(2,2,2-trifluoro-ethyl)-acetamide) was synthesized as follows.

(Reaction 107-21)



[1049] Potassium carbonate (2.50 g, 18.1 mmol) and trifluoro-methanesulfonic acid 2,2,2-trifluoro-ethyl ester (2.36 ml, 16.4 mmol) were added to a solution of 4-bromo-3-methylaniline (1.68 g, 9.03 mmol) in MeCN (39.0 ml) at room temperature, and the mixture was stirred at 80°C overnight. The reaction mixture was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography (hexane-ethyl acetate) to give (4-bromo-3-methyl-phenyl)-(2,2,2-trifluoro-ethyl)-amine (2.20 g, 91%).
MS (ESI) m/z = 268, 270 (M+H)+.


(Reaction 107-22)



[1050] N-(4-Bromo-3-methyl-phenyl)-N-(2,2,2-trifluoro-ethyl)-acetamide was synthesized by operations similar to those in Reaction 107-20 using appropriate reagents and starting material.
MS (ESI) m/z = 310, 312 (M+H)+.

[1051] The aryl bromide reagent used in the synthesis of Compound 616 (N-(4-bromo-3,5-dimethyl-phenyl)-N-(tetrahydro-pyran-4-yl)-acetamide) was synthesized as follows.

(Reaction 107-23)




[1052] N-(4-Bromo-3,5-dimethyl-phenyl)-N-(tetrahydro-pyran-4-yl)-acetamide was synthesized by operations similar to those in Reaction 41-1, Reaction 19-2 (using DMAP as a base) and Reaction 26-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 326, 328 (M+H)^+$.

[1053] The aryl bromide reagent used in the synthesis of Compound 617 (4-(4-bromo-3-methyl-phenyl)-morpholin-3-one) was synthesized as follows.

(Reaction 107-24)



[1054] 4-(4-Bromo-3-methyl-phenyl)-morpholin-3-one was synthesized by operations similar to those in Reaction 10-1 and Reaction 96-18 using appropriate reagents and starting material.


MS (ESI) $m/z = 270, 272 (M+H)^+$.

Example 108

2-(4-Fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 619)

[1055]

(Reaction 108-1)



[1056] 2-(4-Fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 593 (M+H)^+$.

[1057] The example compounds shown below were synthesized by operations similar to those in Example 108 using appropriate reagents and starting materials.


(Compounds 620 to Compound 621)

[1058]

[Table 83]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
620		LCMS-C-1	2.48	609 ($M+H$) ⁺
621		LCMS-C-1	2.43	609 ($M+H$) ⁺


[1059] The aryl bromide reagent used in the synthesis of Compound 621 ((S)-1-(4-bromo-3-methyl-phenyl)-5-hydroxymethyl-pyrrolidin-2-one) was synthesized as follows.

(Reaction 108-2)

[1060] (S)-1-(4-Bromo-3-methyl-phenyl)-5-hydroxymethyl-pyrrolidin-2-one was synthesized by operations similar to those in Reaction 29-3 using appropriate reagents and starting material.
MS (ESI) m/z = 284 (M+H)+.

Example 109


8-{{(E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl}-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 622)

[1061](Reaction 109-1)

[1062] 2,2,6,6-Tetramethylpiperidine 1-oxyl (202 mg, 1.29 mmol) and (diacetoxyiodo)benzene (3.33 g, 10.4 mmol) were added to a solution of 8,8,8-trifluoro-octanol (- 8.63 mmol) in dichloromethane (34.5 mL) at 0°C in an N₂ atmosphere, and the mixture was stirred at 0°C for five minutes and at room temperature for 1.5 hours. The reaction solution was diluted with dichloromethane (200 mL), and the organic layer was sequentially washed with a saturated aqueous sodium sulfite solution (100 mL), a saturated aqueous sodium bicarbonate solution (100 mL) and saturated brine (100 mL). The organic layer was dried over sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 8,8,8-trifluoro-octanal as a colorless oily substance (310 mg, two steps, 20%).

¹H-NMR (400 MHz, CDCl₃) δ 1.35-1.50 (4H, br-m), 1.58-1.62 (2H, br-m), 1.68-1.73 (2H, br-m), 2.05-2.17 (2H, br-m), 2.49 (2H, t, J = 7.1 Hz), 9.82 (1H, s).

¹⁹F-NMR (376 MHz, CDCl₃) δ -66.3 (3F, s).


(Reaction 109-2)

[1063] 1,8-Diazabicyclo[5.4.0]undec-7-ene (436 μ L, 2.92 mmol) was added to a solution of (diethoxy-phosphoryl)-acetic acid ethyl ester (633 μ L, 2.43 mmol) and lithium chloride (144 mg, 3.41 mmol) in acetonitrile (20.0 mL) at 0°C, and the mixture was stirred at 0°C for 10 minutes. A solution of 8,8,8-trifluoro-octanal (2.43 mmol) in acetonitrile (4.3 mL) was added dropwise to the reaction solution at 0°C, and the mixture was stirred for 10 minutes. Thereafter, the reaction mixture was stirred at room temperature for one hour and diluted with methyl tert-butyl ether (200 mL). The organic layer was sequentially washed with a saturated aqueous ammonium chloride solution (30 mL), H₂O (30 mL) and saturated brine (30 mL), dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give (E)-10,10,10-trifluoro-dec-2-enoic acid ethyl ester as a colorless oily substance (426.6 mg, 70%).

¹H-NMR (400 MHz, CDCl₃) δ 1.29 (3H, t, J = 7.2 Hz), 1.31-1.41 (4H, m), 1.41-1.49 (2H, m), 1.51-1.59 (2H, m), 1.99-2.12 (2H, m), 2.20 (2H, ddd, J = 14.5, 7.2, 1.5 Hz), 4.18 (2H, q, J = 7.1 Hz), 5.81 (1H, dt, J = 15.7, 1.6 Hz), 6.95 (1H, dt, J = 15.6, 7.0 Hz).


¹⁹F-NMR (376 MHz, CDCl₃) δ -66.4 (3F, s).

(Reaction 109-3)



[1064] 8-Ethenesulfonyl-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 18-2, Reaction 23-2, Reaction 10-14, Reaction 1-4, Reaction 4-1, Reaction 5-3 and Reaction 25-1 using appropriate reagents and starting material.


MS (ESI) m/z = 424 (M+H)⁺.

(Reaction 109-4)


[1065] 8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.
MS (ESI) $m/z = 613$ (M+H)+.

Example 110


8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one (Compound 623)

[1066](Reaction 110-1)

[1067] Triethylborane (43.8 mL, 438 mmol) and 1,1,1,2,2-pentafluoro-2-iodo-ethane (8.52 mL, 657 mmol) were added to a solution of hex-5-enoic acid (5.21 mL, 438 mmol) in hexane (219 mL) at room temperature, and the mixture was stirred at room temperature for five days. The reaction solution was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 7,7,8,8,8-pentafluoro-5-iodooctanoic acid (purity 80%) as a colorless oily substance (2.63 g, 17%).
 $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.72-1.82 (1H, m), 1.83-1.92 (2H, m), 1.86-1.98 (1H, m), 2.36-2.46 (2H, m), 2.67-2.96 (2H, m), 4.65-4.34 (1H, m).


(Reaction 110-2)

[1068] 7,7,8,8,8-Pentafluoro-octanal was synthesized by operations similar to those in Reaction 95-28 and Reaction 109-1 using appropriate reagents and starting material. This was used in the next reaction without complete purification.

(Reaction 110-3)


[1069] 1,8-Diazabicyclo[5.4.0]undec-7-ene (1.08 mL, 7.26 mmol) was added to a solution of (diethoxy-phosphor-yl)-acetic acid ethyl ester (1.57 mL, 7.87 mmol) and lithium chloride (359 mg, 8.47 mmol) in acetonitrile (60.5 mL) at 0°C , and the mixture was stirred at 0°C for 10 minutes. A solution of 7,7,8,8,8-pentafluoro-octanal (1.32 g, 6.05 mmol) in acetonitrile (20.5 mL) was added dropwise to the reaction solution at 0°C , and the mixture was stirred for 10 minutes. Thereafter, the reaction mixture was stirred at room temperature for one hour and diluted with methyl tert-butyl ether (300 mL). The organic layer was sequentially washed with 2 N hydrochloric acid (50 mL), H_2O (50 mL) and saturated brine (50 mL), dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give (E)-9,9,10,10,10-pentafluoro-dec-2-enoic acid ethyl ester (purity 80%) as a colorless oily substance (49.7 mg, 47%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.29 (3H, t, $J = 7.2$ Hz), 1.37-1.44 (1H, m), 1.46-1.52 (1H, m), 1.56-1.68 (2H, m), 1.93-2.08 (2H, m), 2.18-2.26 (2H, m), 4.19 (2H, q, $J = 7.2$ Hz), 5.79-5.86 (1H, m), 6.88-6.98 (1H, m).

(Reaction 110-4)


[1070] 9,9,10,10,10-Pentafluoro-decanoic acid was synthesized by operations similar to those in Reaction 18-2 and Reaction 95-18 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.31-1.44 (6H, br-m), 1.53-1.68 (4H, m), 1.92-2.08 (2H, br-m), 2.36 (2H, t, $J = 7.6$ Hz).

(Reaction 110-5)

[1071] 8-Ethenesulfonyl-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14, Reaction 1-4, Reaction 4-1, Reaction 5-3 and Reaction 25-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 460$ ($\text{M}+\text{H}$) $^+$.


(Reaction 110-6)

[1072] 8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) m/z = 649 (M+H)+.

Example 111

1-(4-((E)-2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidine-2,4-dione (Compound 624)


[1073](Reaction 111-1)

[1074] 8-Ethenesulfonyl-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14, Reaction 10-11, Reaction 10-12, Reaction 4-1, Reaction 5-3 and Reaction 25-1 using

appropriate reagents and starting material.

MS (ESI) m/z = 352 (M-H) $^-$.

(Reaction 111-2)



[1075] 1-(4-((E)-2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) m/z = 542 (M+H) $^+$.

[1076] The example compounds shown below were synthesized by operations similar to those in Reaction 111-2 using appropriate reagents and starting materials.

(Compounds 625 to Compound 626)


[1077]

[Table 84]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
625		LCMS-C-1	2.7	556 (M+H) $^+$
626		LCMS-C-1	2.73	571 (M+H) $^+$

[1078] The aryl bromide reagent used in the synthesis of Compound 624 (1-(4-bromo-3-methyl-phenyl)-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 111-3)




[1079] (2-Chloro-acetyl)-carbamic acid ethyl ester (356 mg, 2.15 mmol) was added to 4-bromo-3-methyl-phenylamine (400 mg, 2.15 mmol) and dimethylphenylamine (273 μ L, 2.15 mmol) at room temperature. The mixture was stirred at 130°C for five hours, and the reaction solution was then cooled. The precipitate was collected by filtration and washed with CH₃CN to give 1-(4-bromo-3-methyl-phenyl)-imidazolidine-2,4-dione as a colorless solid (314 mg, 54%).

MS (ESI) m/z = 267, 269 (M+H)⁺.

[1080] The aryl bromide reagent used in the synthesis of Compound 625 (1-(4-bromo-3-methyl-phenyl)-5-methyl-imidazolidine-2,4-dione) was synthesized as follows.


(Reaction 111-4)



[1081] 2-Bromo-propionic acid ethyl ester (973 mg, 5.22 mmol) was added to 4-bromo-3-methyl-phenylamine (1.00 g, 5.37 mmol) and dimethylphenylamine (682 μ L, 5.37 mmol) at room temperature, and the mixture was stirred at 60°C for 15 hours. The reaction solution was then cooled and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 2-(4-bromo-3-methyl-phenylamino)-propionic acid ethyl ester as a yellow form (1.38 g, 90%).

MS (ESI) m/z = 286, 288 (M+H)⁺.

(Reaction 111-5)




[1082] KOCN (326 mg, 4.02 mmol) was added to a mixed solution of 2-(4-bromo-3-methyl-phenylamino)-propionic acid ethyl ester (383 mg, 1.34 mmol) in EtOH (5.30 mL) and H₂O (2.68 mL). The mixture was stirred at room temperature for three hours and at 60°C for 11 hours, and AcOH (1 mL) was then added, followed by further stirring for two hours. KOCN (163 mg, 2.01 mmol) was added, followed by further stirring for three hours. The reaction solution was cooled. H₂O (50 mL) was added to the reaction solution at room temperature, and this aqueous layer was then extracted with ethyl acetate (20 mL × 3). The organic layers were concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 1-(4-bromo-3-methyl-phenyl)-5-methylimidazolidine-2,4-dione as a yellow form (134 mg, 35%).

MS (ESI) m/z = 283, 285 (M+H)⁺.

[1083] The aryl bromide reagent used in the synthesis of Compound 626 (1-(4-bromo-3-methyl-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 111-6)




[1084] 1-(4-Bromo-3-methyl-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 111-4 and Reaction 111-5 using appropriate reagents and starting material.

MS (ESI) m/z = 297, 299 (M+H)⁺.

Example 112

8-[(E)-2-[4-(3,4-Dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 627)

[1085]

(Reaction 112-1)

[1086] 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material. MS (ESI) $m/z = 578$ (M+H)+.

[1087] The example compound shown below was synthesized by operations similar to those in Example 112 using appropriate reagents and starting material.

(Compound 628)

[1088]

[Table 85]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
628		LCMS-F-1	0.96	518 (M+H)+

Example 113

3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzoic acid trimethylhydrazide (Compound 629)

[1089]

(Reaction 113-1)



[1090] 3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzoic acid trimethylhydrazide was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) m/z = 626 (M+H)+.

[1091] The example compound shown below was synthesized by operations similar to those in Example 113 using appropriate reagents and starting material.

(Compound 630)


[1092]

[Table 86]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
630		LCMS-D-1	2.37	639 (M+H)+

[1093] The aryl bromide reagent used in the synthesis of Compound 629 (4-bromo-3,5-dimethyl-benzoic acid trimethylhydrazide) was synthesized as follows.

(Reaction 113-2)




[1094] 4-Bromo-3,5-dimethyl-benzoic acid trimethylhydrazide was synthesized by operations similar to those in Reaction 10-14 and Reaction 41-1 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (CDCl_3) δ 7.20 (s, 2H), 3.02 (s, 2H), 2.48 (s, 6H), 2.42 (s, 6H).

[1095] The aryl bromide reagent used in the synthesis of Compound 630 ((R)-3-(5-bromo-4,6-dimethyl-indol-1-yl)-propane-1,2-diol) was synthesized as follows.


(Reaction 113-3)



[1096] N-(4-Bromo-3,5-dimethyl-phenyl)-N-(2,2-diethoxy-ethyl)-methanesulfonamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (CDCl_3) δ 7.06 (s, 2H), 4.57 (t, 1H, $J = 5.72$ Hz), 3.71 (d, 2H, $J = 5.72$ Hz), 3.64 (m, 2H), 3.49 (m, 2H), 2.95 (s, 3H), 2.40 (s, 6H), 1.15 (t, 6H, $J = 7.24$ Hz).


(Reaction 113-4)



[1097] A 1 M solution of titanium(IV) chloride in dichloroethane (5.3 ml, 5.3 mmol) was added to a solution of N-(4-bromo-3,5-dimethyl-phenyl)-N-(2,2-diethoxy-ethyl)-methanesulfonamide (2.09 g, 5.3 mmol) in toluene (17 ml) at room temperature, and the mixture was heated with stirring at 100°C for two hours. An aqueous sodium bicarbonate solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic phase was washed with

saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 5-bromo-4,6-dimethyl-1-methanesulfonyl-indole (1.28 g, 80%). ¹H-NMR (CDCl₃) δ 7.66 (s, 1H), 7.37 (d, 1H, J = 3.81 Hz), 6.69 (dd, 1H, J = 3.81, 0.76 Hz), 3.07 (s, 3H), 2.57 (d, 6H, J = 13.73 Hz).

(Reaction 113-5)




[1098] (R)-3-(5-Bromo-4,6-dimethyl-indol-1-yl)-propane-1,2-diol was synthesized by operations similar to those in Reaction 14-1 and Reaction 25-3 using appropriate reagents and starting material.
MS (ESI) m/z = 298, 300 (M+H)+.

Example 114

3-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imida-
zolidine-2,4-dione (Compound 631)

[1099]

(Reaction 114-1)



[1100] 3-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.
MS (ESI) m/z = 584 (M+H)+.

[1101] The example compound shown below was synthesized by operations similar to those in Example 114 using appropriate reagents and starting material.

(Compound 632)


[1102]

[Table 87]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
632		LCMS-D-1	3.22	627 (M+H)+

Example 115

2-Cyclohexyl-8-((E)-2-quinolin-8-yl-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 633)

[1103](Reaction 115-1)

[1104] 2-Cyclohexyl-8-ethenesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (20 mg, 61.5 μ mol), 8-bromoquinoline (19 mg, 91.3 μ mol), POPd1 (Combiphos, 2.9 mg, 3.1 μ mol) and sodium acetate (7.6 mg, 92.6 μ mol) in dimethylacetamide (0.6 ml) were mixed in a sealed vessel in a nitrogen atmosphere. This mixture was irradiated in a microwave apparatus (190°C, 40 min). The reaction mixture was cooled, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 2-cyclohexyl-8-((E)-2-quinolin-8-yl-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (15.3 mg, 16%).

MS (ESI) m/z = 453 (M+H)+.

[1105] The example compound shown below was synthesized by operations similar to those in Example 115 using appropriate reagents and starting material.

(Compound 634)

[1106]

[Table 88]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
634		LCMS-1	2.3	457 (M+H)+

Example 116

3-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione (Compound 635)

[1107]

(Reaction 116-1)



[1108] 3-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 26-1 (using NMP as a solvent) using appropriate reagents and starting material.


MS (ESI) m/z = 606 (M+H)+.

Example 117

3-(4-((E)-2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione (Compound 636)

[1109]

(Reaction 117-1)



[1110] 3-(4-((E)-2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 26-1 (using NMP as a solvent) using appropriate reagents and starting material.


MS (ESI) m/z = 608 (M+H)+.

Example 118

2-Cyclohexyl-8-[(E)-2-(2-methyl-1H-indol-4-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 637)

[1111]

(Reaction 118-1)



[1112] A mixture of 2-cyclohexyl-8-ethenesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (100 mg, 0.307 mmol), trifluoro-methanesulfonic acid 2-methyl-1H-indol-4-yl ester (129 mg, 0.462 mmol), tetrakis(triphenylphosphine) palladium(0) (35 mg, 30.2 μ mol) and triethylamine (130 μ L, 0.933 mmol) in 1,4-dioxane (1.5 ml) was heated with stirring at 100°C for 18 hours. The reaction mixture was cooled, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 2-cyclohexyl-8-[(E)-2-(2-methyl-1H-indol-4-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (19.3 mg, 14%).

MS (ESI) m/z = 455 (M+H)+.

[1113] The example compound shown below was synthesized by operations similar to those in Example 118 using appropriate reagents and starting material.

(Compound 638)


[1114]

[Table 89]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
638		LCMS-A-1	1.99	471 (M+H)+

[1115] Toluene-4-sulfonic acid 1-methyl-1,2,3,4-tetrahydro-quinolin-5-yl ester used in the synthesis of Compound 638 was synthesized as follows.

(Reaction 118-2)



[1116] Toluene-4-sulfonic acid 1-methyl-1,2,3,4-tetrahydro-quinolin-5-yl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (270 MHz, CDCl_3) δ 1.92-2.02 (2H, m), 2.75-2.82 (2H, m), 2.92 (3H, s), 3.22-3.28 (2H, m), 6.50-6.55 (2H, m), 7.03-7.10 (1H, dd, $J = 8.1, 8.1$ Hz).

Example 119

8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 639)

[1117]

(Reaction 119-1)



[1118] 8-Ethenesulfonyl-2-(4-fluoro-3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (57.6 mg, 0.142 mmol), 4-(4-bromo-3,5-dimethyl-phenoxy)-butane-1,2-diol (49.3 mg, 0.170 mmol), bis(dibenzylideneacetone)palladium(0) (8 mg, 0.014 mmol) and tri-tert-butylphosphine tetrafluoroborate (4 mg, 0.014 mmol) were placed in a vial. NMP (0.5 ml) and N-methyldicyclohexylamine (36.1 μl , 0.170 mmol) were sequentially added in a nitrogen atmosphere, and the mixture was heated with stirring at 100°C for 2.5 hours. A saturated aqueous NH_4Cl solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by thin layer chromatography (ethyl acetate:dichloromethane:methanol = 10:10:1) to give 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (42.8 mg, 49%).






MS (ESI) m/z = 614 (M+H)+.

[1119] The example compounds shown below were synthesized by operations similar to those in Example 119 using appropriate reagents and starting materials.

5 (Compounds 640 to Compound 644)


[1120]

[Table 90]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
640		LCMS-F-1	1.13	642 (M+H)+
641		LCMS-F-1	0.99	669 (M+H)+
642		LCMS-C-1	2.7	651 (M+H)+
643		LCMS-C-1	2.6	595 (M+H)+
644		LCMS-F-1	1	663 (M+H)+

[1121] The aryl bromide reagent used in the synthesis of Compound 640 ((4-bromo-3,5-dimethyl-phenyl)-[1,1,1-²H₃]methyl-carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 119-2)




[1122] (4-Bromo-3,5-dimethyl-phenyl)-[1,1,1-²H₃]methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) *m/z* = 317 (M+H)⁺.

[1123] The aryl bromide reagent used in the synthesis of Compound 641 ((4-bromo-3,5-dimethyl-phenyl)-(4-fluoromethyl-4-hydroxy-piperidin-1-yl)-methanone) was synthesized as follows.


(Reaction 119-3)



[1124] Tetrabutylammonium fluoride (1.0 M in THF, 5.6 ml, 5.6 mmol) was added to a solution of 1-oxa-6-aza-spiro[2.5]octane-6-carboxylic acid tert-butyl ester (400 mg, 1.87 mmol) in tetrahydrofuran (5 ml), and the mixture was heated under reflux for 36 hours. The reaction solution was cooled and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate-hexane) to give 4-fluoromethyl-4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester (87 mg, 19%).

¹H-NMR (CDCl₃) δ 1.46 (s, 9H), 2.93-3.31 (m, 2H), 3.69-3.94 (m, 2H), 4.14 (s, 1H), 4.30 (s, 1H).

(Reaction 119-4)




[1125] (4-Bromo-3,5-dimethyl-phenyl)-(4-fluoromethyl-4-hydroxy-piperidin-1-yl)-methanone was synthesized by operations similar to those in Reaction 5-3 and Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) *m/z* = 344, 346 (M+H)⁺.

[1126] The aryl bromide reagent used in the synthesis of Compound 642 ((4-bromo-3,5-dimethyl-phenyl)-(4-hydroxy-4-methyl-piperidin-1-yl)-methanone) was synthesized as follows.

(Reaction 119-5)




[1127] (4-Bromo-3,5-dimethyl-phenyl)-(4-hydroxy-4-methyl-piperidin-1-yl)-methanone was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 326, 328 (M+H)⁺.

[1128] The aryl bromide reagent used in the synthesis of Compound 644 ((4-bromo-3,5-dimethyl-phenyl)-(2-oxa-7-aza-spiro[3.5]non-7-yl)-methanone) was synthesized as follows.

(Reaction 119-6)




[1129] (4-Bromo-3,5-dimethyl-phenyl)-(2-oxa-7-aza-spiro[3.5]non-7-yl)-methanone was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 338, 340 (M+H)⁺.

Example 120

2-Cyclohexyl-8-[2-(1,2,3,4-tetrahydro-quinolin-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 645)

[1130]

(Reaction 120-1)

[1131] 2-Cyclohexyl-8-[2-(1,2,3,4-tetrahydro-quinolin-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 42-1 using appropriate reagents and starting material.



MS (ESI) m/z = 459 (M+H)+.

[1132] The example compounds shown below were synthesized by operations similar to those in Example 120 using appropriate reagents and starting materials.

(Compounds 646 to Compound 652)

[1133]

[Table 91]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
646		LCMS-C-1	2.42	468 (M+H)+
647		LCMS-C-1	2.60	514 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
648		LCMS-C-1	2.47	455 (M+H) +
649		LCMS-C-1	2.70	539 (M+H)+
650		LCMS-C-1	2.88	485 (M+H)+
651		LCMS-A-1	2.05	515 (M+H)+
652		LCMS-D-1	2.07	486 (M+H)+

[1134] The aryl bromide reagent used in the synthesis of Compound 647 ((7-bromo-indole-1-carboxylic acid dimethylamide) was synthesized as follows.

(Reaction 120-2)




[1135] 7-Bromo-indole-1-carboxylic acid dimethylamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 267, 269 (M+H)+.

[1136] The aryl bromide reagent used in the synthesis of Compound 649 (1-(4-bromo-1H-indol-3-yl)-2,2,2-trifluoroeth-

anone) was synthesized as follows.

(Reaction 120-3)




[1137] Trifluoroacetic anhydride (850 μ L, 6.12 mmol) was added to a solution of 4-bromoindole (1.00 g, 5.10 mmol) in N,N-dimethylformamide (2.0 mL), and the mixture was stirred at room temperature for 1.5 hours. Water was added, followed by extraction with a mixed solvent of ethyl acetate:hexane = 4:1. The organic layers were combined, washed with a saturated aqueous sodium bicarbonate solution, water and saturated brine and dried over sodium sulfate, and the solvent was then distilled off. The residue was purified by silica gel column chromatography to give 1-(4-bromo-1H-indol-3-yl)-2,2,2-trifluoro-ethanone (353 mg, 24%).

MS (ESI) m/z = 292 (M+H) $^{+}$.

[1138] The aryl bromide reagent used in the synthesis of Compound 650 (4-bromo-1-isopropyl-1H-indole) was synthesized as follows.

(Reaction 120-4)




[1139] 4-Bromo-1-isopropyl-1H-indole was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) m/z = 238, 240 (M+H) $^{+}$.

[1140] The aryl bromide reagent used in the synthesis of Compound 651 (4-(4-bromo-indol-1-yl)-butan-1-ol) was synthesized as follows.

(Reaction 120-5)




[1141] 4-(4-Bromo-indol-1-yl)-butan-1-ol was synthesized by operations similar to those in Reaction 25-3 and Reaction 25-4 using appropriate reagents and starting material.

MS (ESI) m/z = 268, 270 ($M+H$).

[1142] The aryl bromide reagent used in the synthesis of Compound 652 (N-(4-bromo-3-cyano-phenyl)-acetamide) was synthesized as follows.

(Reaction 120-6)



[1143] N-(4-Bromo-3-cyano-phenyl)-acetamide was synthesized by operations similar to those in Reaction 19-2 (using DMAP as a base) using appropriate reagents and starting material.


^1H -NMR (DMSO- d_6) δ 10.39 (s, 1H), 8.18 (d, 1H, J = 2.28 Hz), 7.79 (d, 1H, J = 8.74 Hz), 7.70 (dd, 1H, J = 9.15, 2.67 Hz), 2.07 (s, 3H).

Example 121

8-(2-Isoquinolin-5-yl-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 653)

[1144]

(Reaction 121-1)



[1145] 8-(2-Isoquinolin-5-yl-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 42-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 533$ (M+H)+.

[1146] The example compound shown below was synthesized by operations similar to those in Example 121 using appropriate reagents and starting material.

(Compound 654)

[1147]


[Table 92]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
654		LCMS-A-1	2.02	533 (M+H)+

Example 122


N-(3-Methoxy-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide (Compound 655)

[1148]

(Reaction 122-1)

[1149] N-(3-Methoxy-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 503$ ($\text{M}+\text{H}$) $^+$.

(Reaction 122-2)

[1150] 20% $\text{Pd}(\text{OH})_2\text{-C}$ (30 mg) was placed into a solution of N-(3-methoxy-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide (31 mg, 0.0617 mmol) in methanol (5 ml), and the atmosphere was replaced with hydrogen. The mixture was stirred at room temperature for 16 hours. The reaction mixture was filtered, and the filtrate was then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give N-(3-methoxy-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-acetamide (20 mg).


MS (ESI) $m/z = 505$ ($\text{M}+\text{H}$) $^+$.

[1151] The example compound shown below was synthesized by operations similar to those in Example 122 using appropriate reagents and starting material.

(Compound 656)

[1152]


[Table 93]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
656		LCMS-D-1	1.7	608 (M+H)+

The aryl bromide reagent used in the synthesis of Compound 656 (5-(4-bromo-3,5-dimethyl-phenyl)-4,4-dimethyl-1,1-dioxo-1λ⁶-[1,2,5]thiadiazolidin-3-one) was synthesized as follows.

[1153]


(Reaction 122-3)



[1154] Ethyl-2-bromoisobutyric acid (3.7 ml, 24.99 mmol) and sodium bicarbonate (630 mg, 7.49 mmol) were added to 4-bromo-3,5-dimethyl-aniline (500 mg, 2.49 mmol), and the mixture was irradiated with microwaves at 130°C for 15 minutes. The reaction mixture was diluted with ethyl acetate, and the organic layer was sequentially washed with water and saturated brine and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give ethyl 2-[(4-bromo-3,5-dimethyl-phenyl)amino]-2-methyl-propanoate (230 mg, 29%).

MS (ESI) m/z = 314, 316 (M+H)+.


(Reaction 122-4)



[1155] Ethyl 2-[N-(4-bromo-3,5-dimethyl-phenyl)-N-sulfamoyl-amino]-2-methyl-propanoate was synthesized by operations similar to those in Reaction 92-2 and Reaction 7-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 505 (M+H)^+$.

(Reaction 122-5)




[1156] A 2 M solution of sodium methoxide in methanol (4 ml, 8 mmol) was added to a solution of ethyl 2-[N-(4-bromo-3,5-dimethyl-phenyl)-N-sulfamoyl-amino]-2-methyl-propanoate (100 mg, 0.254 mmol) in methanol (12 ml), and the mixture was irradiated with microwaves at 65°C for 10 minutes. The reaction mixture was diluted with ethyl acetate, and the organic layer was sequentially washed with water and saturated brine and concentrated under reduced pressure to give 5-(4-bromo-3,5-dimethyl-phenyl)-4,4-dimethyl-1,1-dioxo-1 λ^6 -[1,2,5]thiadiazolidin-3-one (80 mg, 90%).

$^1\text{H-NMR}$ (CDCl_3) δ 7.23 (s, 2H), 2.48 (s, 6H), 1.31 (s, 6H).

Example 123

8-{2-[2-Methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 657)

[1157](Reaction 123-1)

[1158] 8-{2-[2-Methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 122-2 using appropriate reagents and starting material.


MS (ESI) $m/z = 561$ (M+H)+.

Example 124

{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzyl}-carbamic acid isobutyl ester (Compound 658)

[1159]

(Reaction 124-1)



[1160] {4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzyl}-carbamic acid isobutyl ester was synthesized by operations similar to those in Reaction 25-2 and Reaction 42-2 using appropriate reagents and starting material.







MS (ESI) $m/z = 533$ (M+H)+.

[1161] The example compounds shown below were synthesized by operations similar to those in Example 124 using appropriate reagents and starting materials.

(Compounds 659 to Compound 664)


[1162]

[Table 94]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
659		LCMS-A-1	1.62	491 (N+H)+
660		LCMS-C-1	2.40	616 (M+H)+
661		LCMS-C-1	2.98	587 (M+H)+
662		LCMS-C-1	2.82	591 (M+H)+
663		LCMS-C-1	2.37	477 (M+H)+
664		LCMS-C-1	2.20	582 (M+H)+

[1163] The aryl bromide reagent used in the synthesis of Compound 658 ((4-bromo-benzyl)-carbamic acid isobutyl ester) was synthesized as follows.

(Reaction 124-2)




[1164] Isobutyl chloroformate (0.152 ml, 1.17 mmol) was added to a solution of 4-bromo-benzylamine hydrochloride (200 mg, 0.899 mmol) and pyridine (0.182 ml, 2.25 mmol) in DMF (2.0 ml), and the mixture was stirred at room temperature for 1.5 hours. A 1 N aqueous HCl solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give (4-bromo-benzyl)-carbamic acid isobutyl ester (180 mg, 70%).

MS (ESI) m/z = 286, 288 ($M+H$) $^+$.

[1165] The aryl bromide reagent used in the synthesis of Compound 660 (2-(4-bromo-3-trifluoromethyl-phenylamino)-N-(4-hydroxy-butyl)-acetamide) was synthesized as follows.

(Reaction 124-3)




[1166] 2-(4-Bromo-3-trifluoromethyl-phenylamino)-N-(4-hydroxy-butyl)-acetamide was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 369 ($M+H$) $^+$.

[1167] The aryl bromide reagent used in the synthesis of Compound 661 ((4-bromo-3-trifluoromethyl-phenyl)-carbamic acid isobutyl ester) was synthesized as follows.

(Reaction 124-4)




[1168] (4-Bromo-3-trifluoromethyl-phenyl)-carbamic acid isobutyl ester was synthesized by operations similar to those in Reaction 124-2 using appropriate reagents and starting material.

MS (ESI) m/z = 338 ($M-H$) $^-$.

[1169] The aryl bromide reagent used in the synthesis of Compound 662 (N-(4-bromo-3-trifluoromethyl-phenyl)-ben-

zamide) was synthesized as follows.

(Reaction 124-5)




[1170] N-(4-Bromo-3-trifluoromethyl-phenyl)-benzamide was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

MS (ESI) m/z = 344 ($\text{M}+\text{H}$) $^+$.

[1171] The aryl bromide reagent used in the synthesis of Compound 664 (N-[2-(4-bromo-3-methyl-benzenesulfonylamino)-ethyl]-acetamide) was synthesized as follows.

(Reaction 124-6)



[1172] N-[2-(4-Bromo-3-methyl-benzenesulfonylamino)-ethyl]-acetamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.


MS (ESI) m/z = 335 ($\text{M}+\text{H}$) $^+$.

Example 125

N-(2-Dimethylamino-ethyl)-2-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenylamino)-acetamide (Compound 665)

[1173]

(Reaction 125-1)



[1174] N-(2-Dimethylamino-ethyl)-2-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenylamino)-acetamide was synthesized by operations similar to those in Reaction 25-2 and Reaction 42-2 using appropriate reagents and starting material.


MS (ESI) $m/z = 639 \text{ (M+H)}^+$.

[1175] The example compound shown below was synthesized by operations similar to those in Example 125 using appropriate reagents and starting material.

(Compound 666)


[1176]

[Table 95]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
666		LCMS-A-1	2.60	637 (M+H) ⁺

[1177] The aryl bromide reagent used in the synthesis of Compound 666 ((4-bromo-3-methyl-phenyl)-(cis-2,6-dimethyl-morpholin-4-yl)-methanone) was synthesized as follows.

(Reaction 125-2)



[1178] (4-Bromo-3-methyl-phenyl)-(cis-2,6-dimethyl-morpholin-4-yl)-methanone was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.


MS (ESI) m/z = 312, 314 ($M+H$)⁺.

Example 126

N-(2,2,3,3,4,4,4-Heptafluoro-butyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide (Compound 667)

[1179]

(Reaction 126-1)




[1180] N-(2,2,3,3,4,4,4-Heptafluoro-butyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 42-1 using appropriate reagents and starting material.

MS (ESI) m/z = 741 ($M+H$)⁺.

[1181] The aryl bromide reagent used in the synthesis of Compound 667 (4-bromo-N-(2,2,3,3,4,4,4-heptafluoro-butyl)-3-methyl-benzenesulfonamide) was synthesized as follows.

(Reaction 126-2)




[1182] 4-Bromo-N-(2,2,3,3,4,4,4-heptafluorobutyl)-3-methylbenzenesulfonamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.
MS (ESI) $m/z = 454, 456 (M+\text{Na})^+$.

Example 127

8-[2-(3-Methoxy-2-methyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 668)

[1183]

(Reaction 127-1)



[1184] 8-[2-(3-Methoxy-2-methyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 122-2 using appropriate reagents and starting material.


MS (ESI) $m/z = 462 (M+\text{H})^+$.

[1185] The example compound shown below was synthesized by operations similar to those in Example 127 using appropriate reagents and starting material.

(Compound 669)


[1186]

[Table 96]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
669		LCMS-F-1	1.1	605 (M+H)+

[1187] The aryl bromide reagent used in the synthesis of Compound 669 ((4-bromo-2-methoxy-3,5-dimethyl-phenyl)-methyl-carbamic acid tert-butyl ester) was synthesized as follows.


(Reaction 127-2)



[1188] Di-tert-butyl dicarbonate (1.32 ml, 5.75 mmol) was added to a solution of 2-amino-4,6-dimethyl-phenol (731.8 mg, 5.335 mmol) in THF (3.7 ml), and the mixture was stirred at room temperature for five hours. The reaction mixture was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography (hexane-ethyl acetate) to give (2-hydroxy-3,5-dimethyl-phenyl)-carbamic acid tert-butyl ester as a red purple solid (1.234 g, 97%).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 1.52 (9H, s), 2.21 (3H, s), 2.24 (3H, s), 6.54 (1H, br), 6.71 (1H, s), 6.76 (1H, s), 7.74 (1H, br).


(Reaction 127-3)



[1189] (2-Methoxy-3,5-dimethyl-phenyl)-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 26-4 (using cesium carbonate as a base) using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.30-1.56 (9H, br), 2.24 (3H, s), 2.25 (3H, s), 3.16 (3H, s), 3.69 (3H, s), 6.76 (1H, br), 6.87 (1H, s).

(Reaction 127-4)



[1190] N-Bromosuccinimide (326 mg, 1.83 mmol) was added to a solution of (2-methoxy-3,5-dimethyl-phenyl)-methyl-carbamic acid tert-butyl ester (456 mg, 1.72 mmol) in dichloromethane (1.8 ml) at 0°C , and the mixture was stirred at room temperature for 50 minutes. The reaction mixture was diluted with dichloromethane and adjusted to pH 9 and washed with water and a 1 N aqueous NaOH solution (1 ml), and the organic layer was dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give (4-bromo-2-methoxy-3,5-dimethyl-phenyl)-methyl-carbamic acid tert-butyl ester as a colorless solid (545 mg, 92%).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.30-1.56 (9H, br), 2.35 (3H, s), 2.37 (3H, s), 3.14 (3H, s), 3.69 (3H, s), 6.82-7.07 (1H, br).

Example 128

8-{2-[4-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 670)

[1191]

(Reaction 128-1)



[1192] 8-{2-[4-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 122-2 using appropriate reagents and starting material.


MS (ESI) m/z = 575 ($\text{M}+\text{H}$) $^+$.

Example 129

8-[2-(5,7-Dimethyl-2-oxo-2,3-dihydro-benzoxazol-6-yl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one (Compound 671)

[1193]

(Reaction 129-1)




[1194] 8-[2-(5,7-Dimethyl-2-oxo-2,3-dihydro-benzoxazol-6-yl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 122-2 (using MeCN-DMF as a solvent) using appropriate reagents and starting material.

MS (ESI) m/z = 503 (M+H)+.

[1195] The aryl bromide reagent used in the synthesis of Compound 129 (6-bromo-5,7-dimethyl-3H-benzoxazol-2-one) was synthesized as follows.

(Reaction 129-2)




[1196] 6-Bromo-5,7-dimethyl-3H-benzoxazol-2-one was synthesized by operations similar to those in Reaction 127-4 (using acetic acid as a solvent) using appropriate reagents and starting material.

MS (ESI) m/z = 242, 244 (M+H)+.

Example 130

(2-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-ethyl)-carbamic acid methyl ester (Compound 672)

[1197]

(Reaction 130-1)

[1198] (2-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-ethyl)-carbamic acid methyl ester was synthesized by operations similar to those in Reaction 26-1 and Reaction 42-2 using appropriate reagents and starting material.

MS (ESI) m/z = 505 (M+H)+.

[1199] The example compound shown below was synthesized by operations similar to those in Example 130 using appropriate reagents and starting material.

(Compound 673)


[1200]

[Table 97]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
673		LCMS-C-1	2.93	559 (M-H)-

[1201] The aryl bromide reagent used in the synthesis of Compound 673 ((4-bromo-3-methyl-benzyl)-methyl-carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 130-2)




[1202] (4-Bromo-3-methyl-benzyl)-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.
MS (ESI) m/z = 336, 338 ($M+Na$)⁺.

Example 131

8-(2-{4-[3-(2-Hydroxy-ethyl)-2-oxo-imidazolidin-1-yl]-2-methyl-phenyl}-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 674)

[1203]


(Reaction 131-1)



[1204] 8-(2-{4-[3-(2-Hydroxy-ethyl)-2-oxo-imidazolidin-1-yl]-2-methyl-phenyl}-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 42-2 using appropriate reagents and starting material.
MS (ESI) m/z = 608 ($M+H$)⁺.

[1205] The aryl bromide reagent used in the synthesis of Compound 674 (1-(4-bromo-3-methyl-phenyl)-3-(2-hydroxy-ethyl)-imidazolidin-2-one) was synthesized as follows.

(Reaction 131-2)




[1206] 1-(4-Bromo-3-methyl-phenyl)-3-(2-hydroxy-ethyl)-imidazolidin-2-one was synthesized by operations similar to those in Reaction 29-3 using appropriate reagents and starting material.
MS (ESI) m/z = 299, 301 ($M+H$)⁺.

Example 132

2-(3-Methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonylamino)-acetamide (Compound 675)

[1207]

(Reaction 132-1)



[1208] 2-(3-Methyl-4-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonylamino)-acetamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 42-2 using appropriate reagents and starting material.
MS (ESI) m/z = 632 ($M+H$)⁺.


[1209] The example compound shown below was synthesized by operations similar to those in Example 132 using

appropriate reagents and starting material.

(Compound 676)


[1210]

[Table 98]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
676		LCMS-C-1	2.70	614 (M+H)+

[1211] The aryl bromide reagent used in the synthesis of Compound 675 (2-(4-bromo-3-methyl-benzenesulfonylamino)-acetamide) was synthesized as follows.

(Reaction 132-2)




[1212] 2-(4-Bromo-3-methyl-benzenesulfonylamino)-acetamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.
MS (ESI) m/z = 307, 309 (M+H)+.

Example 133

3-(4-{2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione (Compound 677)

[1213]

(Reaction 133-1)




[1214] 3-(4-{2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 26-1 and Reaction 42-2 using appropriate reagents and starting material.
MS (ESI) m/z = 586 ($\text{M}+\text{H}$) $^+$.

Example 134

N-Cyclopentyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide (Compound 678)

[1215]

(Reaction 134-1)




[1216] N-Cyclopentyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide was synthesized by operations similar to those in Reaction 26-1, Reaction 42-2 and Reaction 12-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 621$ (M+H)+.

[1217] The aryl bromide reagent used in the synthesis of Compound 678 ((4-bromo-3-methyl-phenyl)-cyclopentyl-amine) was synthesized as follows.

(Reaction 134-2)



[1218] (4-Bromo-3-methyl-phenyl)-cyclopentyl-amine was synthesized by operations similar to those in Reaction 41-1 using appropriate reagents and starting material.


MS (ESI) $m/z = 254, 256$ (M+H)+.

Example 135

1-(2,3-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 679)

[1219]

(Reaction 135-1)



[1220] 1-(2,3-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 26-1, Reaction 42-2 and Reaction 89-2 using appropriate reagents and starting material.


MS (ESI) $m/z = 518$ (M+H)+.

[1221] The example compound shown below was synthesized by operations similar to those in Example 135 using appropriate reagents and starting material.

(Compound 680)


[1222]

[Table 99]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
680		LCMS-C-1	2.58	518 (M+H)+

[1223] The aryl iodide reagent used in the synthesis of Compound 679 (4-iodo-N,2,3-trimethyl-aniline) was synthesized as follows.

(Reaction 135-2)




[1224] A 28% solution of sodium methoxide in methanol (0.694 ml, 6.07 mmol) was added to a solution of 4-iodo-2,3-dimethyl-aniline (500 mg, 2.02 mmol) and paraformaldehyde (121 mg, 4.05 mmol) in methanol (8.0 ml), and the mixture was stirred at room temperature for 17 hours. Sodium borohydride (153 mg, 4.05 mmol) was further added, and the mixture was stirred at room temperature for four hours. A 1 N aqueous NaOH solution was added to the reaction mixture, followed by extraction with dichloromethane. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-iodo-N,2,3-trimethyl-aniline (146 mg, 88%).

MS (ESI) $m/z = 262$ (M+H)+.

[1225] The aryl iodide reagent used in the synthesis of Compound 680 ((4-iodo-2,5-dimethyl-phenyl)-methyl-amine) was synthesized as follows.

(Reaction 135-3)



[1226] (4-Iodo-2,5-dimethyl-phenyl)-methyl-amine was synthesized by operations similar to those in Reaction 135-2 using appropriate reagents and starting material.


MS (ESI) $m/z = 262$ (M+H)+.

Example 136

{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester (Compound 681)

[1227]

(Reaction 136-1)




[1228] {4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 26-1, Reaction 42-2 and Reaction 39-2 using appropriate reagents and starting material.

MS (ESI) m/z = 589 (M-H)-.

[1229] The aryl bromide reagent used in the synthesis of Compound 681 ((4-bromo-3-methyl-benzyl)-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 136-2)



[1230] (4-Bromo-3-methyl-benzyl)-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.


MS (ESI) m/z = 472, 474 (M+H)+.

Example 137

1-(3-Methoxy-5-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 682)

[1231]

(Reaction 137-1)




[1232] 1-(3-Methoxy-5-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 26-1 (using DMI as a solvent), Reaction 122-2 (using acetonitrile as a solvent), Reaction 5-3 and Reaction 89-2 (using KOCN as a reagent) using appropriate reagents and starting material.

MS (ESI) $m/z = 534$ ($\text{M}+\text{H}$) $^+$.

[1233] The aryl bromide reagent used in the synthesis of Compound 682 ((4-bromo-3-methoxy-5-methyl-phenyl)-methyl-carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 137-2)



[1234] (4-Bromo-3-methoxy-5-methyl-phenyl)-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 127-2 (using toluene as a solvent), Reaction 26-2 and Reaction 26-4 (using cesium carbonate as a base) using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.50 (9H, s), 2.42 (3H, s), 3.26 (3H, s), 3.90 (3H, s), 6.71 (1H, d, $J = 4.0$ Hz), 6.77 (1H, $J = 4.0$ Hz).

Example 138

1-(3-Chloro-5-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 683)

[1235]


(Reaction 138-1)



[1236] 1-(3-Chloro-5-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 26-1, Reaction 42-1, Reaction 5-3 and Reaction 89-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 538$ (M+H)+.

[1237] The aryl bromide reagent used in the synthesis of Compound 683 ((4-bromo-3-chloro-5-methyl-phenyl)-methyl-carbamic acid tert-butyl ester) was synthesized as follows.


(Reaction 138-2)



[1238] Iron (2.34 g, 41.8 mmol) and acetic acid (0.80 mL, 14.0 mmol) were added to a mixed solution of 2-bromo-1-chloro-3-methyl-5-nitrobenzene (3.50 g, 14.0 mmol) in ethanol (15 mL)-water (31 mL) at room temperature. The mixture was stirred at 100°C for one hour, and a saturated aqueous sodium bicarbonate solution was then added at 0°C. The mixture was filtered through celite, and the filtrate was washed with ethyl acetate and water. The filtrate was concentrated under reduced pressure. Ethyl acetate was then added, and the organic layer and the aqueous layer were separated. The aqueous layer was repeatedly extracted with ethyl acetate three times, and the organic layers were then dried over sodium sulfate. The resulting residue was concentrated under reduced pressure to give 4-bromo-3-chloro-5-methylphenylamine as a pale brown solid (3.01 g, 98%).

MS (ESI) $m/z = 220, 222$ (M+H)+.

(Reaction 138-3)



[1239] (4-Bromo-3-chloro-5-methyl-phenyl)-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 127-2 (using toluene as a solvent) and Reaction 26-4 (using cesium carbonate as a base) using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.46 (9H, s), 2.44 (3H, s), 3.22 (3H, s), 7.06 (1H, d, J = 2.4 Hz), 7.23 (1H, d, J = 2.4 Hz).

[1240] The example compound shown below was synthesized by operations similar to those in Example 138 using appropriate reagents and starting material.

(Compound 684)


[1241]

[Table 100]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
684		LCMS-F-1	0.94	548 (M+H)+


Example 139

5,7-Dimethyl-6-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-1H-quinazoline-2,4-dione (Compound 685)

[1242](Reaction 139-1)

[1243] Benzyltrimethylammonium dichloroiodate (324 mg, 0.930 mmol) was added to a mixture of 8-[2-(4-amino-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride (530 mg, 1.03 mmol) and calcium carbonate (517 mg, 5.17 mmol) in methanol (6 mL)-dichloromethane (15 mL) at room temperature. The mixture was stirred at room temperature for 22 hours, and an aqueous sodium bicarbonate solution and ethyl acetate were then added. The organic layer and the aqueous layer were separated, and the aqueous layer was repeatedly extracted with ethyl acetate three times. The organic layers were combined and washed with saturated brine, and the insoluble matter was then filtered off through celite. The filtrate was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography to give 8-[2-(4-amino-3-iodo-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a pale yellow solid (337 mg, 62%).


MS (ESI) m/z = 587 (M+H)+.

(Reaction 139-2)

[1244] A mixture of 8-[2-(4-amino-3-iodo-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (15 mg, 0.0256 mmol), palladium acetate (1.1 mg, 5.11 μmol), 1,1'-bis(diphenylphosphino)fer-

rocene (2.2 mg, 5.11 μmol), tert-butyl isocyanate (12 μL , 0.0767 mmol) and N,N-diisopropylethylamine (13 μL , 0.0767 mmol) in tetrahydrofuran (1 mL) was heated with stirring for 12 hours in a pressure bottle sealed under the conditions of 4 atm and 80°C in a carbon monoxide atmosphere. After returning to room temperature, palladium acetate (2.2 mg, 10.22 μmol), 1,1'-bis(diphenylphosphino)ferrocene (4.4 mg, 10.22 μmol), tert-butyl isocyanate (24 μL , 0.153 mmol) and N,N-diisopropylethylamine (26 μL , 0.153 mmol) were added to the reaction solution, and the mixture was heated with stirring for 14 hours in a pressure bottle sealed under the conditions of 4 atm and 80°C in a carbon monoxide atmosphere. After returning to room temperature, the reaction solution was concentrated under reduced pressure, and the resulting residue was purified by preparative TLC to give 3-tert-butyl-5,7-dimethyl-6-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-1H-quinazoline-2,4-dione as a pale yellow solid (3.3 mg, 22%). MS (ESI) m/z = 586 (M+H)+.

(Reaction 139-3)




[1245] An aqueous hydrogen bromide solution (80 μL) was added to a mixed solution of 3-tert-butyl-5,7-dimethyl-6-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-1H-quinazoline-2,4-dione (3.3 mg, 3.41 μmol) in acetic acid (80 μL), and the mixture was heated with stirring at 100°C for one hour. The reaction solution was returned to room temperature and then concentrated under reduced pressure, and the resulting residue was purified by preparative TLC to give 5,7-dimethyl-6-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-1H-quinazoline-2,4-dione as a pale yellow solid (1.2 mg, 44%). MS (ESI) m/z = 530 (M+H)+.

Example 140

8-[(E)-2-[1-(2-Amino-ethyl)-1H-indol-4-yl]-ethenesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 686)

[1246]

(Reaction 140-1)




[1247] 8-[(E)-2-[1-(2-Amino-ethyl)-1H-indol-4-yl]-ethenesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 7-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 484$ (M+H)+.

[1248] The aryl bromide reagent used in the synthesis of Compound 686 ([2-(4-bromo-indol-1-yl)-ethyl]-carbamic acid tert-butyl ester) was synthesized as follows.


(Reaction 140-2)



[1249] Tetrabutylammonium hydrogen sulfate (41.0 mg, 0.121 mmol) and sodium hydroxide (210 mg, 5.25 mmol) were added to a solution of 4-bromo-1H-indole (0.30 ml, 2.39 mmol) in acetonitrile (0.80 mL), and the mixture was stirred at room temperature for 20 minutes. Subsequently, 2-chloroethylamine hydrochloride (334 mg, 2.88 mmol) was added and the mixture was heated with stirring at 100°C for seven hours. After cooling, water was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 2-(4-bromo-indol-1-yl)-ethylamine (222 mg, 39%).

MS (ESI) $m/z = 239, 241$ (M+H)+.

(Reaction 140-3)



[1250] A 2 N aqueous NaOH solution (0.47 ml, 0.94 mmol) and di-tert-butyl dicarbonate (223 mg, 1.02 mmol) were sequentially added to a solution of 2-(4-bromo-indol-1-yl)-ethylamine (222 mg, 0.928 mmol) in dioxane (0.47 ml), and the mixture was stirred at room temperature for 19 hours. Water was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was washed with saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane only) to give [2-(4-bromo-indol-1-yl)-ethyl]-carbamic acid tert-butyl ester (292 mg, 93%).


MS (ESI) $m/z = 361, 363$ (M+Na)+.

[1251] The example compound shown below was synthesized by operations similar to those in Example 140 using appropriate reagents and starting material.

(Compound 687)

[1252]


[Table 101]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
687	 2TFA	LCMS-C-1	1.98	431 (M+H)+

Example 141

(S)-2-Amino-3-hydroxy-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-propionamide (Compound 688)

[1253]


(Reaction 141-1)

[1254] (S)-2-Amino-3-hydroxy-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-propionamide was synthesized by operations similar to those in Reaction 25-2 and Reaction 7-2 using appropriate reagents and starting material.

MS (ESI) m/z = 580 (M+H)+.

[1255] The aryl bromide reagent used in the synthesis of Compound 688 ((S)-1-(4-bromo-3-methyl-phenylcarbamoyl)-2-hydroxy-ethyl]-carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 141-2)




[1256] [(S)-1-(4-Bromo-3-methyl-phenylcarbamoyl)-2-hydroxy-ethyl]-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-1 using appropriate reagents and starting material.
MS (ESI) m/z = 317, 219 ($M-tBu+H+H$)⁺.

Example 142

N-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-piperidin-4-yl-methanesulfonamide (Compound 689)

[1257]


(Reaction 142-1)



[1258] N-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-piperidin-4-yl-methanesulfonamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 7-2 using appropriate reagents and starting material.
MS (ESI) m/z = 684 ($M+H$)⁺.

[1259] The aryl bromide reagent used in the synthesis of Compound 142 (4-[(4-bromo-3,5-dimethyl-phenyl)-methanesulfonyl-amino]-piperidine-1-carboxylic acid tert-butyl ester) was synthesized as follows.

(Reaction 142-2)



[1260] 4-[(4-Bromo-3,5-dimethyl-phenyl)-methanesulfonyl-amino]-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 41-1 and Reaction 6-1 using appropriate reagents and starting material.


MS (ESI) $m/z = 461, 463 (M+H)^+$.

Example 143

8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 690)

[1261]

(Reaction 143-1)




[1262] 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 7-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 557 (M+H)^+$.

[1263] The aryl bromide reagent used in the synthesis of Compound 690 ((4-bromo-3,5-dimethyl-phenyl)-methyl-

carbamic acid tert-butyl ester) was synthesized as follows.

(Reaction 143-2)




[1264] Iodomethane (20.5 ml, 329 mmol) was added to a solution of (4-bromo-3,5-dimethyl-phenyl)-carbamic acid tert-butyl ester (47.5 g, 158 mmol) and cesium carbonate (80.6 g, 247 mmol) in DMF (165 ml) at room temperature, and the mixture was stirred for 27 hours. Further, cesium carbonate (26.9 g, 82.6 mmol) and iodomethane (20.5 ml, 329 mmol) were added at room temperature, and the mixture was further stirred for three days. An aqueous ammonium chloride solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give (4-bromo-3,5-dimethyl-phenyl)-methyl-carbamic acid tert-butyl ester (9.34 g, 92%).

MS (ESI) m/z = 258, 260 ($M-tBu+H+H$)⁺.

Example 144

8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 691)

[1265]

(Reaction 144-1)


[1266] 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 7-2 using appropriate reagents and starting material.

MS (ESI) m/z = 593 ($\text{M}+\text{H}$)⁺.

Example 145

1-{3,5-Dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-1-methyl-urea (Compound 692)

[1267]


(Reaction 145-1)

[1268] 1-{3,5-Dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-1-methyl-urea was synthesized by operations similar to those in Reaction 25-2, Reaction 7-2 and Reaction 89-2 using appropriate reagents and starting material.
MS (ESI) m/z = 546 ($\text{M}+\text{H}$)⁺.

Example 146

1-(2-Methoxy-3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea (Compound 693)

[1269]

(Reaction 146-1)

[1270] 1-(2-Methoxy-3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 26-1, Reaction 7-2 and Reaction 89-2 using appropriate reagents and starting material.


MS (ESI) m/z = 546 ($\text{M}+\text{H}$)⁺.

Example 147

1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-(2-fluoro-ethyl)-urea (Compound 694)

[1271]

(Reaction 147-1)




[1272] 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-phenyl]-1-(2-fluoro-ethyl)-urea was synthesized by operations similar to those in Reaction 26-1, Reaction 7-2 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) m/z = 630 (M+H)+.

[1273] The aryl bromide reagent used in the synthesis of Compound 694 ((4-bromo-3,5-dimethyl-phenyl)-(2-fluoro-ethyl)-carbamic acid tert-butyl ester) was synthesized as follows.


(Reaction 147-2)



[1274] A solution of 4-bromo-3,5-dimethyl-phenylamine (400 mg, 2 mmol), toluene-4-sulfonic acid 2-fluoro-ethyl ester (567 mg, 2.6 mmol) and 2,6-lutidine (429 mg, 4.0 mmol) in DMA (5 ml) was heated with stirring at 120°C for four hours. The mixture was quenched with water and then extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate-hexane) to give (4-bromo-3,5-dimethyl-phenyl)-(2-fluoro-ethyl)-amine (262 mg, 53%).

$^1\text{H-NMR}$ (CDCl_3) δ 6.39 (s, 2H), 4.68 (t, 1H, J = 4.96 Hz), 4.52 (t, 1H, J = 4.96 Hz), 3.92 (brd, 1H), 3.45 (t, 1H, J = 4.96 Hz), 3.36 (t, 1H, J = 4.96 Hz), 2.34 (s, 3H).

(Reaction 147-3)




[1275] (4-Bromo-3,5-dimethyl-phenyl)-(2-fluoro-ethyl)-carbamate tert-butyl ester was synthesized by operations similar to those in Reaction 19-2 (using DMAP as a base) using appropriate reagents and starting material. $^1\text{H-NMR}$ (CDCl_3) δ 6.97 (s, 2H), 4.64 (t, 1H, $J = 4.96$ Hz), 4.48 (t, 1H, $J = 4.96$ Hz), 3.9 (t, 1H, $J = 4.96$ Hz), 3.82 (t, 1H, $J = 4.96$ Hz), 2.39 (s, 6H), 1.45 (s, 9H).

Example 148

1-(3-Chloro-5-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (Compound 695)

[1276]

(Reaction 148-1)




[1277] 1-(3-Chloro-5-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 26-1, Reaction 50-2 and Reaction 89-2 using appropriate reagents and starting material. MS (ESI) $m/z = 630$ ($\text{M}+\text{H}^+$).

Example 149

1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea (Compound 696)

[1278]

(Reaction 149-1)


[1279] 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 25-2, Reaction 7-2, Reaction 89-2 and Reaction 42-2 using appropriate reagents and starting material.

MS (ESI) m/z = 600 (M+H)+.

Example 150

8-[(E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 697)

[1280]

(Reaction 150-1)

[1281] 8-[(E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 25-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 616 (M+H)^+$.

[1282] The example compound shown below was synthesized by operations similar to those in Example 150 using appropriate reagents and starting material.

(Compound 698)

[1283]

[Table 102]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
698		LCMS-D-1	2.56	660 ($M+H$) ⁺

Example 151

N-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide (Compound 699)

[1284]

(Reaction 151-1)




[1285] N-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide was synthesized by operations similar to those in Reaction 25-2 and Reaction 39-2 using appropriate reagents and starting material.

MS (ESI) m/z = 671 (M+H)+.

[1286] The aryl bromide reagent used in the synthesis of Compound 699 (N-(4-bromo-3,5-dimethyl-phenyl)-N-[2-[2-(tert-butyl-dimethyl-silanyloxy)-ethoxy]-ethyl]-acetamide) was synthesized as follows.

(Reaction 151-2)



[1287] N-(4-Bromo-3,5-dimethyl-phenyl)-N-[2-[2-(tert-butyl-dimethyl-silanyloxy)-ethoxy]-ethyl]-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (CDCl_3) δ 6.95 (s, 2H), 3.81 (dd, 2H, J = 6.10, 5.72 Hz), 3.70 (dd, 2H, J = 4.95, 5.34 Hz), 3.58 (dd, 2H, J = 5.72, 6.10 Hz), 3.46 (dd, 2H, J = 5.72, 4.95 Hz), 2.41 (s, 6H), 1.83 (s, 3H), 0.86 (s, 9H).

Example 152

N-[2-(2-Hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-acetamide (Compound 700)

[1288]

(Reaction 152-1)



[1289] N-[2-(2-Hydroxy-ethoxy)-ethyl]-N-{3-methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-acetamide was synthesized by operations similar to those in Reaction 25-2 and Reaction 12-5 using appropriate reagents and starting material.

MS (ESI) $m/z = 605$ (M+H) $^+$.

[1290] The example compound shown below was synthesized by operations similar to those in Example 152 using appropriate reagents and starting material.

(Compound 701)

[1291]


[Table 103]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
701		LCMS-A-1	2.44	575 (M+H) $^+$

[1292] The aryl bromide reagent used in the synthesis of Compound 701 (acetic acid 2-[acetyl-(4-bromo-3,5-dimethyl-

phenyl)-amino]-ethyl ester) was synthesized as follows.

(Reaction 152-2)




[1293] Acetic acid 2-[acetyl-(4-bromo-3,5-dimethyl-phenyl)-amino]-ethyl ester was synthesized by operations similar to those in Reaction 25-12 and Reaction 12-2 using appropriate reagents and starting material.
MS (ESI) m/z = 328, 330 ($M+H$)⁺.

Example 153

2-Cyclohexyl-8-[(E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 702)


[1294]

(Reaction 153-1)



[1295] 2-Cyclohexyl-8-[(E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-5-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.


MS (ESI) m/z = 555 ($M+H$)⁺.

(Reaction 153-2)

[1296] 2-Cyclohexyl-8-((E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-5-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (41.9 mg, 0.0755 mmol) was dissolved in a methylene chloride-methanol mixed solution (1:1, 1.5 ml). Trifluoroacetic acid (0.35 ml) was added and the mixture was stirred for two days. A saturated aqueous sodium bicarbonate solution and water were added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by P-TLC (ethyl acetate-methanol) to give 2-cyclohexyl-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (33.4 mg, 86%).

MS (ESI) m/z = 515 (M+H)+.

[1297] The aryl bromide reagent used in the synthesis of Compound 702 (5-bromo-1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indole) was synthesized as follows.

(Reaction 153-3)

[1298] 5-Bromo-1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indole was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.


MS (ESI) m/z = 310 (M+H)+.

Example 154

8-((E)-2-[1-[2-((S)-2,3-Dihydroxy-propoxy)-ethyl]-1H-indol-4-yl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 704)

[1299]

(Reaction 154-1)



[1300] 8-((E)-2-{1-[2-((S)-2,3-Dihydroxy-propoxy)-ethyl]-1H-indol-4-yl}-ethenesulfonyl)-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 and Reaction 25-4 using appropriate reagents and starting material.

MS (ESI) m/z = 655 (M+H)+.

[1301] The example compounds shown below were synthesized by operations similar to those in Example 154 using appropriate reagents and starting materials.

(Compounds 705 to Compound 706)

[1302]

[Table 104]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
705		LCMS-D-1	2.14	646 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
706		LCMS-D-1	2.15	685 (M+H)+

[1303] The aryl bromide reagent used in the synthesis of Compound 704 (4-bromo-1-[2-((R)-2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-ethyl]-1H-indole) was synthesized as follows.


(Reaction 154-2)



[1304] 4-Bromo-1-[2-((R)-2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy)-ethyl]-1H-indole was synthesized by operations similar to those in Reaction 25-3, Reaction 39-2 and Reaction 20-2 using appropriate reagents and starting material. MS (ESI) m/z = 354,356 (M+H)+.

[1305] The aryl bromide reagent used in the synthesis of Compound 705 ([[(4S,5S)-5-(4-bromo-3,5-dimethyl-phenoxy)methyl]-2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy]-tert-butyl-dimethyl-silane) was synthesized as follows.

(Reaction 154-3)




[1306] [[(4S,5S)-5-(4-Bromo-3,5-dimethyl-phenoxy)methyl]-2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy]-tert-butyl-dimethyl-silane was synthesized by operations similar to those in Reaction 26-4 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 6.67 (s, 2H), 4.27 (m, 1H), 4.13-3.73 (m, 5H), 2.36 (s, 6H), 1.44 (s, 3H), 1.43 (s, 3H), 0.88 (s, 9H),

0.06 (s, 6H).

[1307] The aryl bromide reagent used in the synthesis of Compound 706 ((4S,5S)-5-[2-(4-bromo-indol-1-yl)-ethoxymethyl]-2,2-dimethyl-[1,3]dioxolan-4-yl)-methanol) was synthesized as follows.

(Reaction 154-4)




[1308] ((4S,5S)-5-[2-(4-Bromo-indol-1-yl)-ethoxymethyl]-2,2-dimethyl-[1,3]dioxolan-4-yl)-methanol was synthesized by operations similar to those in Reaction 6-1 and Reaction 20-2 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 7.28 (m, 2H), 7.20 (d, 1H, J = 3.3 Hz), 7.06 (t, 1H, J = 7.8 Hz), 6.55 (d, 1H, J = 3.0 Hz), 4.30 (t, 2H, J = 5.4 Hz), 3.94 (m, 1H), 3.84-3.41 (m, 7H), 1.91 (br s, 1H), 1.39 (s, 3H), 1.36 (s, 3H).

Example 155

N-((S)-2,3-Dihydroxy-propyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide (Compound 707)

[1309]

(Reaction 155-1)

[1310] N-((S)-2,3-Dihydroxy-propyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 26-1 and reaction 25-4 using appropriate reagents and starting material.


MS (ESI) m/z = 659 ($M+H$)⁺.

Example 156

{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester (Compound 708)

[1311]

(Reaction 156-1)



[1312] {4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 26-1 and Reaction 39-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 589 (M+H)^+$.

[1313] The example compounds shown below were synthesized by operations similar to those in Example 156 using appropriate reagents and starting materials.

(Compounds 709 to Compound 710)

[1314]


[Table 105]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
709		LCMS-D-1	2.34	585 ($M+H$) ⁺
710		LCMS-D-1	2.53	587 ($M+H$) ⁺

[1315] The aryl bromide reagent used in the synthesis of Compound 708 ((4-bromo-3-methyl-benzyl)-[2-(tert-butyl-

dimethyl-silanyloxy)-ethyl]-carbamic acid tert-butyl ester) was synthesized as follows.


(Reaction 156-2)



[1316] (4-Bromo-3-methyl-benzyl)-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.
MS (ESI) m/z = 472, 474 ($M+H$)⁺.

[1317] The aryl bromide reagent used in the synthesis of Compound 709 (N-(4-bromo-3,5-dimethyl-phenyl)-N-[4-(tert-butyl-dimethyl-silanyloxy)-cyclohexyl]-acetamide) was synthesized as follows.


(Reaction 156-3)



[1318] N-(4-Bromo-3,5-dimethyl-phenyl)-N-[4-(tert-butyl-dimethyl-silanyloxy)-cyclohexyl]-acetamide was synthesized by operations similar to those in Reaction 41-1, Reaction 19-2 (using DMAP as a base) and Reaction 26-2 using appropriate reagents and starting material.
MS (ESI) m/z = 454, 456 ($M+H$)⁺.

[1319] The aryl bromide reagent used in the synthesis of Compound 710 ((R)-1-(4-bromo-3,5-dimethyl-phenyl)-5-[2-(tert-butyl-dimethyl-silanyloxy)-ethoxymethyl]-pyrrolidin-2-one) was synthesized as follows.

(Reaction 156-4)



[1320] (R)-1-(4-Bromo-3,5-dimethyl-phenyl)-5-[2-(tert-butyl-dimethyl-silanyloxy)-ethoxymethyl]-pyrrolidin-2-one was synthesized by operations similar to those in Reaction 20-2 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (CDCl_3) δ 7.07 (s, 2H), 4.19 (m, 1H), 3.64 (t, 2H, $J = 4.96$ Hz), 3.44 (m, 2H), 3.40 (t, 2H, $J = 4.96$ Hz), 2.52 (m, 2H), 2.36 (s, 6H), 2.17 (m, 2H), 0.87 (s, 9H), 0.04 (s, 6H).

Example 157

N-(2-Hydroxy-ethyl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-isobutylamide (Compound 711)

[1321]

(Reaction 157-1)



[1322] N-(2-Hydroxy-ethyl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-isobutylamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 39-2 using appropriate reagents and starting material.





MS (ESI) $m/z = 623$ ($\text{M}+\text{H}$) $^+$.

[1323] The example compounds shown below were synthesized by operations similar to those in Example 157 using appropriate reagents and starting materials.

(Compounds 712 to Compound 715)


[1324]

[Table 106]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
712		LCMS-D-1	2.90	649 (M+H) ⁺
713		LCMS-D-1	2.90	635 (M+H) ⁺
714		LCMS-D-1	3.20	678 (M+H) ⁺
715		LCMS-D-1	3.30	699 (M+H) ⁺


[1325] The aryl bromide reagent used in the synthesis of Compound 711 (N-(4-bromo-3-methyl-phenyl)-N-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-isobutylamide) was synthesized as follows.

(Reaction 157-2)



[1326] Triethylamine (0.39 ml, 2.80 mmol), dimethylaminopyridine (13 mg, 0.11 mmol) and tert-butyl-dimethyl-chlorosilane (319 mg, 2.11 mmol) were added to a solution of 2-(4-bromo-3-methyl-phenylamino)-ethanol (430 mg, 1.87 mmol) in dichloromethane (3.8 ml) at room temperature, and the mixture was stirred at room temperature for 24 hours. The reaction mixture was diluted with dichloromethane, and the organic layer was washed with water, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give (4-bromo-3-methyl-phenyl)-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-amine (629 mg, 98%). $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 0.08 (6H, s), 0.91 (9H, s), 2.32 (3H, s), 3.18 (2H, dd, $J = 5.7$ and 5.1 Hz), 3.80 (2H, t, $J = 5.1$ Hz), 4.01 (1H, dull t, $J = 5.7$ Hz), 6.34 (1H, dd, $J = 8.7$, 2.5 Hz), 6.51 (1H, d, $J = 2.5$ Hz), 7.27 (1H, d, $J = 8.7$ Hz).

(Reaction 157-3)




[1327] N-(4-Bromo-3-methyl-phenyl)-N-[2-(tert-butyl-dimethyl-silyloxy)-ethyl]-isobutylamide was synthesized by operations similar to those in Reaction 105-2 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 0.04 (6H, s), 0.87 (9H, s), 1.02 (6H, d, $J = 6.6$ Hz), 2.41 (3H, s), 2.47 (1H, sept, $J = 6.6$ Hz), 3.74 (4H, s), 6.92 (1H, dd, $J = 8.4, 2.3$ Hz), 7.13 (1H, d, $J = 2.3$ Hz), 7.53 (1H, d, $J = 8.4$ Hz).

[1328] The aryl bromide reagents used in the synthesis of Compound 712 and Compound 713 ((R)-1-(4-bromo-3,5-dimethylphenyl)-5-(tert-butyl-dimethyl-silyloxymethyl)-3,3-dimethylpyrrolidin-2-one and (R)-1-(4-bromo-3,5-dimethylphenyl)-5-(tert-butyl-dimethyl-silyloxymethyl)-3-methyl-pyrrolidin-2-one) were synthesized as follows.


(Reaction 157-4)



[1329] 1 M LHMDS (0.61 ml, 0.61 mmol) was added dropwise to a solution of (R)-1-(4-bromo-3,5-dimethylphenyl)-5-(((tert-butyl)dimethylsilyloxy)methyl)pyrrolidin-2-one (119 mg, 0.29 mmol) in THF (2.4 ml) at -78°C in a nitrogen atmosphere, and the mixture was stirred at -78°C for 15 minutes. A solution of iodomethane (38 μL , 0.62 mmol) in THF (0.5 ml) was added at -78°C , and the mixture was stirred at -78°C for 15 minutes, warmed to room temperature and further stirred for three hours. A saturated aqueous ammonium chloride solution was added to the reaction mixture, and the mixture was diluted with dichloromethane. The organic layer was washed with saturated brine, and then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give a mixture of (R)-1-(4-bromo-3,5-dimethylphenyl)-5-(tert-butyl-dimethyl-silyloxymethyl)-3,3-dimethylpyrrolidin-2-one (minor) and (R)-1-(4-bromo-3,5-dimethylphenyl)-5-(tert-butyl-dimethyl-silyloxymethyl)-3-methylpyrrolidin-2-one (major) (629 mg, 98%). This was used in Heck reaction without complete separation and purification.

[1330] The aryl bromide reagent used in the synthesis of Compound 714 ((3-(4-bromo-3,5-dimethylphenyl)-1-[2-(tert-butyl-dimethyl-silyloxy)-ethyl]-5,5-dimethyl-imidazolidine-2,4-dione) was synthesized as follows.

(Reaction 157-5)




[1331] (3-(4-Bromo-3,5-dimethyl-phenyl)-1-[2-(tert-butyl-dimethyl-silyloxy)-ethyl]-5,5-dimethyl-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 7.11 (s, 2H), 3.85 (t, 2H, d = 6.0 Hz), 3.43 (t, 2H, d = 6.0 Hz), 2.42 (s, 6H), 1.49 (s, 6H), 0.90 (s, 9H), 0.07 (s, 6H).

[1332] The aryl bromide reagent used in the synthesis of Compound 715 (N-(4-bromo-3,5-dimethyl-phenyl)-N-[4-(tert-butyl-dimethyl-silanyloxy)-cyclohexyl]-methanesulfonamide) was synthesized as follows.

(Reaction 157-6)



[1333] N-(4-Bromo-3,5-dimethyl-phenyl)-N-[4-(tert-butyl-dimethyl-silanyloxy)-cyclohexyl]-methanesulfonamide was synthesized by operations similar to those in Reaction 41-1 and Reaction 6-1 using appropriate reagents and starting material.


MS (ESI) m/z = 490, 492 (M+H)⁺.

Example 158

N-4-[(E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide (Compound 716)

[1334]

(Reaction 158-1)




[1335] N-4-[(E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide was synthesized by operations similar to those in Reaction 26-1 and

Reaction 39-2 using appropriate reagents and starting material.

MS (ESI) m/z = 631 (M+H)+.

[1336] The aryl bromide reagent used in the synthesis of Compound 716 (N-(4-bromo-3,5-dimethyl-phenyl)-N-{2-[2-(tert-butyl-dimethyl-silanyloxy)-ethoxy]-ethyl}-acetamide) was synthesized as follows.

(Reaction 158-2)



[1337] N-(4-Bromo-3,5-dimethyl-phenyl)-N-{2-[2-(tert-butyl-dimethyl-silanyloxy)-ethoxy]-ethyl}-acetamide was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (CDCl_3) δ 6.95 (s, 2H), 3.81 (dd, 2H, J = 6.10, 5.72 Hz), 3.70 (dd, 2H, J = 4.95, 5.34 Hz), 3.58 (dd, 2H, J = 5.72, 6.10 Hz), 3.46 (dd, 2H, J = 5.72, 4.95 Hz), 2.41 (s, 6H), 1.83 (s, 3H), 0.86 (s, 9H).

Example 159

N-(2-Fluoro-5-methyl-4-{(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-N-(2-hydroxy-ethyl)-acetamide (Compound 717)

[1338]

(Reaction 159-1)




[1339] N-(2-Fluoro-5-methyl-4-{(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfo-

nyl]-vinyl)-phenyl)-N-(2-hydroxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 12-5 using appropriate reagents and starting material.

MS (ESI) m/z = 613 (M+H)+.

[1340] The aryl iodide reagent used in the synthesis of Compound 717 (acetic acid 2-[acetyl-(2-fluoro-4-iodo-5-methyl-phenyl)-amino]-ethyl ester) was synthesized as follows.


(Reaction 159-2)



[1341] Pyridine (0.87 mL, 10.78 mmol) was added to a solution of 2-fluoro-4-iodo-5-methyl-phenylamine (1082.9 mg, 4.314 mmol) in THF (10.8 mL). 2-Chloroethyl chloroformate (0.47 mL, 4.53 mmol) was then added dropwise and the mixture was stirred overnight. Potassium hydroxide (968.2 mg, 17.25 mmol) and ethanol (10.8 mL) were subsequently added, and the mixture was heated under reflux overnight. The reaction mixture was then quenched by adding a saturated aqueous ammonium chloride solution and extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 2-(2-fluoro-4-iodo-5-methyl-phenylamino)-ethanol as a pale brown solid (1217.0 mg, 96%).

MS (ESI) m/z = 296 (M+H)+.

(Reaction 159-3)



[1342] Acetic acid 2-[acetyl-(2-fluoro-4-iodo-5-methyl-phenyl)-amino]-ethyl ester was synthesized by operations similar to those in Reaction 12-2 using appropriate reagents and starting material.


MS (ESI) m/z = 402 (M+Na)+.

Example 160

N-(4-((E)-2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide (Compound 718)

[1343]

(Reaction 160-1)




[1344] N-(4-((E)-2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 26-1 and Reaction 12-5 using appropriate reagents and starting material.
MS (ESI) m/z = 611 (M+H)+.

Example 161

N-(2-Hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methanesulfonamide (Compound 719)


[1345]

(Reaction 161-1)



[1346] N-[2-(tert-Butyl-dimethyl-silanyloxy)-ethyl]-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methanesulfonamide was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.


¹H-NMR (300 MHz, CDCl₃) δ 0.04 (6H, s), 0.87 (9H, s), 1.70 (2H, m), 2.21 (2H, m), 2.44 (3H, s), 3.02 (3H, s), 3.32 (2H, m), 3.71 (2H, t, J = 5.7 Hz), 3.81 (4H, m), 6.67 (1H, J = 15.3 Hz), 7.27 (2H, m), 7.42 (1H, m), 7.56 (2H, m), 7.71 (1H, d, J = 15.3 Hz), 7.80 (1H, d, J = 8.0 Hz), 7.84 (1H, s), 10.24 (1H, brs).

(Reaction 161-2)

[1347] A 10% aqueous citric acid solution (0.14 ml, 0.067 mmol) was added to a solution of N-[2-(tert-butyl-dimethyl-silyloxy)-ethyl]-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methanesulfonamide (14.3 mg, 0.0192 mmol) in acetonitrile (0.2 ml), and the mixture was stirred at 60°C for 2.5 hours. The reaction mixture was diluted with ethyl acetate, and the organic layer was washed with an aqueous sodium bicarbonate solution, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate) to give N-(2-hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methanesulfonamide (12.5 mg, 100%).

MS (ESI) m/z = 631 (M+H)+.

[1348] The aryl bromide reagent used in the synthesis of Compound 719 (N-(4-bromo-3-methyl-phenyl)-N-[2-(tert-butyl-dimethyl-silyloxy)-ethyl]-methanesulfonamide) was synthesized as follows.

(Reaction 161-3)

[1349] N-(4-Bromo-3-methyl-phenyl)-N-[2-(tert-butyl-dimethyl-silyloxy)-ethyl]-methanesulfonamide was synthesized by operations similar to those in Reaction 6-1 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 0.04 (6H, s), 0.87 (9H, s), 2.40 (3H, s), 2.96 (3H, s), 3.68 (2H, m), 3.75 (2H, m), 7.05 (1H, ddd, J = 8.5, 2.5, 0.6 Hz), 7.25 (1H, d, J = 2.5 Hz), 7.54 (1H, d, J = 8.5 Hz).

Example 162

N-(2-Hydroxy-ethyl)-N-{3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-methanesulfonamide (Compound 720)

[1350]

(Reaction 162-1)




[1351] N-(2-Hydroxy-ethyl)-N-{3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-methanesulfonamide was synthesized by operations similar to those in Reaction 26-1, Reaction 39-2 and Reaction 122-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 599$ (M+H)+.

Example 163

N-(4-{2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide (Compound 721)

[1352]

(Reaction 163-1)


[1353] N-(4-{2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide was synthesized by operations similar to those in Reaction 26-1, Reaction 12-5 and Reaction 42-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 599$ ($\text{M}+\text{H}$) $^+$.

Example 164

2-Hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide (Compound 722)

[1354]


(Reaction 164-1)



[1355] 2-Hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 26-1, Reaction 161-2 and Reaction 12-5 using appropriate reagents and starting material.
MS (ESI) m/z = 611 (M+H)+.

[1356] The aryl bromide reagent used in the synthesis of Compound 722 (acetic acid ((4-bromo-3-methyl-phenyl)-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-carbamoyl)-methyl ester) was synthesized as follows.

(Reaction 164-2)



[1357] Acetic acid ((4-bromo-3-methyl-phenyl)-[2-(tert-butyl-dimethyl-silanyloxy)-ethyl]-carbamoyl)-methyl ester was synthesized by operations similar to those in Reaction 105-2 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 0.03 (6H, s), 0.86 (9H, s), 2.13 (3H, s), 2.41 (3H, s), 3.76 (4H, s), 4.36 (2H, s), 6.99 (1H, dd, J = 8.4, 2.4 Hz), 7.19 (1H, d, J = 2.4 Hz), 7.56 (1H, d, J = 8.4 Hz).

Example 165

1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea (Compound 723)

[1358]

(Reaction 165-1)




[1359] 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 26-1 and Reaction 81-1 using appropriate reagents and starting material.
MS (ESI) m/z = 598 ($\text{M}+\text{H}$) $^+$.

Example 166

1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-[2-(2-hydroxy-ethoxy)-ethyl]-urea (Compound 724)

[1360]

(Reaction 166-1)



[1361] 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-[2-(2-hydroxy-ethoxy)-ethyl]-urea was synthesized by operations similar to those in Reaction 25-2 and Reaction 81-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 672$ (M+H)⁺

[1362] The example compound shown below was synthesized by operations similar to those in Example 166 using appropriate reagents and starting material.

(Compound 725)


[1363]

[Table 107]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
725		LCMS-D-1	2.32	642 (M+H) ⁺

[1364] The aryl bromide reagent used in the synthesis of Compound 724 (2-[2-(4-bromo-3,5-dimethyl-phenylamino)-ethoxy]-ethanol) was synthesized as follows.

(Reaction 166-2)




[1365] 2-[2-(4-Bromo-3,5-dimethyl-phenylamino)-ethoxy]-ethanol was synthesized by operations similar to those in Reaction 39-2 and Reaction 96-16 using appropriate reagents and starting material.

MS (ESI) m/z = 288, 290 ($M+H$)⁺.

[1366] The aryl bromide reagent used in the synthesis of Compound 725 ((4-bromo-3,5-dimethyl-phenyl)-(2-methoxy-ethyl)-amine) was synthesized as follows.

(Reaction 166-3)



[1367] (4-Bromo-3,5-dimethyl-phenyl)-(2-methoxy-ethyl)-amine was synthesized by operations similar to those in Reaction 25-3 and Reaction 96-16 using appropriate reagents and starting material.


MS (ESI) m/z = 258, 260 ($M+H$)⁺.

Example 167

1-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-isopropyl-urea (Compound 726)

[1368]

(Reaction 167-1)



[1369] 1-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-isopropyl-urea was synthesized by operations similar to those in Reaction 25-2 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 544$ ($M+H$) $^+$.

[1370] The example compound shown below was synthesized by operations similar to those in Example 167 using appropriate reagents and starting material.

(Compound 727)


[1371]

[Table 108]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
727		LCMS-D-1	2.68	516 ($M+H$) $^+$

[1372] The aryl bromide reagent used in the synthesis of Compound 726 ((4-bromo-3,5-dimethyl-phenyl)-isopropyl-amine) was synthesized as follows.

(Reaction 167-2)




[1373] (4-Bromo-3,5-dimethyl-phenyl)-isopropyl-amine was synthesized by operations similar to those in Reaction 41-1 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (CDCl_3) δ 6.33 (s, 2H), 3.57 (q, 1H, $J = 6.6$ Hz), 2.32 (s, 6H), 1.18 (d, 6H, $J = 6.6$ Hz).

[1374] The aryl bromide reagent used in the synthesis of Compound 727 ((4-bromo-3-ethyl-phenyl)-methyl-amine) was synthesized as follows.

(Reaction 167-3)



[1375] (4-Bromo-3-ethyl-phenyl)-methyl-amine was synthesized by operations similar to those in Reaction 25-3 and Reaction 96-16 using appropriate reagents and starting material.


MS (ESI) $m/z = 214, 216$ ($M+H$) $^+$.

Example 168

1-(3-Methoxy-5-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (Compound 728)

[1376]

(Reaction 168-1)



[1377] 1-(3-Methoxy-5-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 26-1 (using DMI as a solvent) and Reaction 89-2 (using KOCN as a reagent) using appropriate reagents and starting material.


MS (ESI) $m/z = 532$ ($M+H$) $^+$.

Example 169

1-Cyanomethyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea (Compound 729)

[1378]

(Reaction 169-1)



[1379] 1-Cyanomethyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea was synthesized by operations similar to those in Reaction 26-1 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) m/z = 591 (M+H)+.

[1380] The example compound shown below was synthesized by operations similar to those in Example 169 using appropriate reagents and starting material.

(Compound 730)

[1381]

[Table 109]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
730		LCMS-B-1	2.36	620 (M+H)+

Example 170

1-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-(2,2,2-trifluoro-ethyl)-urea (Compound 731)

[1382]

(Reaction 170-1)



[1383] 1-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-(2,2,2-trifluoro-ethyl)-urea was synthesized by operations similar to those in Reaction 26-1 and Reaction 81-1 using appropriate reagents and starting material.


MS (ESI) m/z = 634 ($\text{M}+\text{H}$) $^+$.

Example 171

1-(4-{2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 732)

[1384]

(Reaction 171-1)




[1385] 1-(4-{2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 25-2, Reaction 89-2 and Reaction 42-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 532$ (M+H)+.

Example 172

1-((S)-2,3-Dihydroxy-propyl)-1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea (Compound 733)

[1386]

(Reaction 172-1)




[1387] 1-((S)-2,3-Dihydroxy-propyl)-1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea was synthesized by operations similar to those in Reaction 25-2, Reaction 89-2 and Reaction 25-4 using appropriate reagents and starting material.

MS (ESI) m/z = 576 (M+H)+.

[1388] The aryl bromide reagent used in the synthesis of Compound 733 ((4-bromo-3,5-dimethyl-phenyl)-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-amine) was synthesized as follows.

(Reaction 172-2)



[1389] (4-Bromo-3,5-dimethyl-phenyl)-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-amine was synthesized by operations similar to those in Reaction 96-16 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (CDCl_3) δ 6.39 (s, 2H), 4.34 (m, 1H), 4.09 (dd, 1H, J = 8.2, 6.3 Hz), 3.75 (dd, 1H, J = 8.2, 6.3 Hz), 3.29-3.11 (m, 2H), 2.33 (s, 6H), 1.45 (s, 3H), 1.37 (s, 3H).

Example 173

N-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-phenyl}-N-cyclopentyl-acetamide trifluoroacetate (Compound 734)

[1390]

(Reaction 173-1)



[1391] N-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-phenyl}-N-cyclopentyl-acetamide trifluoroacetate was synthesized by operations similar to those in Reaction 26-1 and Reaction 12-2 (using HPLC for purification) using appropriate reagents and starting material.


MS (ESI) $m/z = 541$ (M+H)+.

Example 174

(S)-2-Amino-N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-3-methyl-butylamide (Compound 735)

[1392]


(Reaction 174-1)



[1393] ((S)-1-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl-carbamoyl}-2-methyl-propyl)-carbamic acid 9H-fluoren-9-ylmethyl ester was synthesized by operations similar to those in Reaction 26-1 and Reaction 10-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 766$ (M+H)+.

(Reaction 174-2)



[1394] Piperidine (1 ml) was added to a solution of ((S)-1-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenylcarbamoyl}-2-methyl-propyl)-carbamic acid 9H-fluoren-9-ylmethyl ester (84 mg, 0.11 mmol) in dichloromethane (4 ml), and the mixture was stirred at room temperature for three hours. The reaction mixture was quenched with water and then extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give (S)-2-amino-N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-3-methyl-butylamide (20 mg, 33%).


MS (ESI) $m/z = 544$ (M+H)+.

Example 175

2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-N-pyridin-4-yl-acetamide (Compound 736)

[1395]

(Reaction 175-1)




[1396] 2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-N-pyridin-4-yl-acetamide was synthesized by operations similar to those in Reaction 25-2, Reaction 23-2 and Reaction 10-22 using appropriate reagents and starting material.

MS (ESI) $m/z = 575$ ($M+H$) $^+$.

[1397] The aryl bromide reagent used in the synthesis of Compound 736 ((4-bromo-indol-1-yl)-acetic acid ethyl ester) was synthesized as follows.

(Reaction 175-2)




[1398] (4-Bromo-indol-1-yl)-acetic acid ethyl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

MS (ESI) $m/z = 282$ ($M+H$) $^+$.

Example 176


2-(3-Methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide (Compound 737)

[1399]

(Reaction 176-1)

[1400] (3-Methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetic acid was synthesized by operations similar to those in Reaction 25-2 and Reaction 23-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 488$ ($\text{M}+\text{H}$) $^+$.


(Reaction 176-2)

[1401] N,N-Diisopropylethylamine (31.4 μL , 0.185 mmol) and 2-chloro-1-methylpyridinium iodide (18.9 mg, 0.074 mmol) were added to a solution of (3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetic acid (30.0 mg, 0.062 mmol) in dichloromethane (0.5 mL) and DMF (0.1 mL), and the mixture was stirred for 10 minutes. A 2.0 M ammonia-methanol solution (0.15 mL, 0.308 mmol) and DMAP (0.8 mg, 0.006 mmol) were then added, and the mixture was stirred overnight. The reaction mixture was then quenched by adding a saturated aqueous ammonium chloride solution and extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide as a white powder (18.3 mg, 61%).

MS (ESI) $m/z = 487$ ($\text{M}+\text{H}$) $^+$.

[1402] The aryl bromide reagent used in the synthesis of Compound 737 (2-(4-bromo-3-methyl-phenyl)-malonic acid dimethyl ester) was synthesized as follows.


(Reaction 176-3)




[1403] 2-(4-Bromo-3-methyl-phenyl)-malonic acid dimethyl ester was synthesized by operations similar to those in Reaction 12-1 using appropriate reagents and starting material.
MS (ESI) m/z = 302 (M+H)⁺.

Example 177

2-Cyclohexyl-8-[(E)-2-[4-(4,5-dihydro-thiazol-2-ylamino)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triazaspiro[4.5]dec-1-en-4-one (Compound 738)

[1404](Reaction 177-1)

[1405] 8-[(E)-2-(4-Amino-2,6-dimethyl-phenyl)vinyl]sulfonyl-3-cyclohexyl-2,4,8-triazaspiro[4.5]dec-3-en-1-one was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.
MS (ESI) m/z = 445 (M+H)⁺.

(Reaction 177-2)

[1406] 2-(Methylthio)-2-thiazoline (19 μ L, 0.17 mmol) and acetic acid (1.2 ml) were added to a solution of 8-[(E)-2-(4-amino-2,6-dimethyl-phenyl)vinyl]sulfonyl-3-cyclohexyl-2,4,8-triazaspiro[4.5]dec-3-en-1-one (74 mg, 0.16 mmol) in EtOH (2.5 ml) at room temperature, and the mixture was heated with stirring at 80°C for 16 hours. The reaction solution was diluted with ethyl acetate, and the organic layer was then sequentially washed with water and saturated brine and


concentrated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate) to give 2-cyclohexyl-8-((E)-2-[4-(4,5-dihydro-thiazol-2-ylamino)-2,6-dimethylphenyl]-ethenesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one (46.1 mg, 52%).
MS (ESI) m/z = 530 (M+H)+.

Example 178

2-Cyclohexyl-8-((E)-2-[2-methyl-4-(3-methyl-oxetan-3-ylmethoxy)-phenyl]-ethenesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one (Compound 739)

[1407]

(Reaction 178-1)




[1408] 2-Cyclohexyl-8-((E)-2-[2-methyl-4-(3-methyl-oxetan-3-ylmethoxy)-phenyl]-ethenesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 25-2 and Reaction 26-4 using appropriate reagents and starting material.
MS (ESI) m/z = 516 (M+H)+.


Example 179

1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropyl}-phenyl)-1-methyl-urea (Compound 740) and 1-(3,5-dimethyl-4-{1-methyl-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea (Compound 741)

[1409]


(Reaction 179-1)

[1410] (3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.
MS (ESI) $m/z = 573$ ($\text{M}+\text{H}$)⁺.

(Reaction 179-2)

[1411] A 1 M solution of sodium bis(trimethylsilyl)amide in tetrahydrofuran (0.31 mL, 0.306 mmol) was added to a mixture of trimethylsulfoxonium iodide (29 mg, 0.131 mmol) in 1,3-dimethyl-2-imidazolidinone (2 mL) at room temperature. The reaction solution was stirred at room temperature for 0.5 hour. A mixed solution of (3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methyl-carbamic acid tert-butyl ester (50 mg, 0.0873 mmol) in 1,3-dimethyl-2-imidazolidinone (2 mL) was then added at room temperature, and the mixture was heated with stirring at 50°C for 15 hours. After returning to room temperature, an aqueous ammonium chloride solution and ethyl acetate were added to the reaction solution. The organic layer and the aqueous layer were separated, and the aqueous layer was repeatedly extracted with ethyl acetate three times. The organic layers were combined, washed with water twice and saturated brine, and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give a mixture of (3,5-dimethyl-4-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropyl)-phenyl)-methyl-carbamic acid tert-butyl ester. This mixture was used in the next reaction as such without further purification.
MS (ESI) $m/z = 587$ ($\text{M}+\text{H}$)⁺.

(Reaction 179-3)



[1412] 1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropyl-phenyl}-1-methyl-urea

MS (ESI) $m/z = 530$ (M+H)+

and

1-(3,5-dimethyl-4-{1-methyl-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea

MS (ESI) $m/z = 530$ (M+H)+


were synthesized by operations similar to those in Reaction 7-2 and Reaction 89-2 using appropriate reagents and starting material.

Example 180

N-(3-Methyl-4-{(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-sulfamide (Compound 742)


[1413]

(Reaction 180-1)



[1414] 8-[(E)-2-(4-Amino-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.


MS (ESI) $m/z = 509$ (M+H)+

(Reaction 180-2)

[1415] A solution of tBuOH (71.9 mg, 0.97 mmol) in dichloromethane (1.5 ml) was added to a solution of chlorosulfonyl isocyanate (137 mg, 0.97 mmol) in dichloromethane (3 ml) with stirring under ice-cooling. The mixture was stirred at 0°C for 10 minutes and then added to a solution of 8-[(E)-2-(4-amino-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (400 mg, 0.81 mmol) and triethylamine (164 mg, 1.62 mmol) in dichloromethane (3 ml). The mixture was stirred for one hour, and then quenched with water and extracted with dichloromethane. The organic layer was washed with saturated brine, and then dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (methyl acetate-hexane) to give N-tert-butoxycarbonyl-N'-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-sulfamide (334 mg, 60%).

¹H-NMR (400 MHz, CDCl₃) δ 1.42 (9H, s), 1.78 (2H, dt, J = 14.2, 3.9 Hz), 2.04-2.14 (2H, m), 2.40 (3H, s), 3.43 (2H, ddd, J = 12.7, 9.8, 2.9 Hz), 3.74 (2H, dt, J = 12.2, 4.4 Hz), 6.64 (1H, d, J = 15.6 Hz), 7.08-7.11 (2H, m), 7.38 (1H, d, J = 8.3 Hz), 7.48-7.54 (2H, m), 7.68 (1H, d, J = 15.1 Hz), 7.73 (1H, d, J = 7.8 Hz), 7.76 (1H, s), 9.62 (1H, s);

MS (ESI) m/z = 668 (M+H)⁺.

(Reaction 180-3)

[1416] N-(3-Methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-sulfamide was synthesized by operations similar to those in Reaction 4-1 using appropriate reagents and starting material.


MS (ESI) m/z = 588 (M+H)⁺.

Example 181

N-(3-Hydroxy-propyl)-N'-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-sulfamide (Compound 743)

[1417]

(Reaction 181-1)




[1418] N-(3-Hydroxy-propyl)-N'-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-sulfamide was synthesized by operations similar to those in Reaction 31-7, Reaction 4-1 and Reaction 39-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 646$ ($\text{M}+\text{H}$) $^+$.

Example 182

N-Methyl-N-(3-methyl-4-((E)-3-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-propenyl)-phenyl)-acetamide (Compound 744)

[1419]

(Reaction 182-1)


[1420] N-Methyl-N-(3-methyl-4-((E)-3-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-propenyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 5-4, Reaction 55-2 and Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) m/z = 515 (M+H)⁺.

Example 183

2-Cyclohexyl-8-[2-(2-methyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-2-en-4-one (Compound 745)

[1421]

(Reaction 183-1)

[1422] 2-Cyclohexyl-8-[2-(2-methyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 745) was obtained by operations similar to those in Reaction 18-2 using Compound 637 as a starting material.

MS (ESI) m/z = 457 (M+H)⁺.

[1423] The example compounds shown below were obtained by operations similar to those in Example 183 using appropriate solvents (methanol or dimethylformamide or a methanol-dimethylformamide mixed solution) and starting compounds.

(Compounds 746 to Compound 749)

[1424]

Example 184

[1425]

[1427] The example compounds shown below were obtained by operations similar to those in Example 184 using appropriate solvents (acetonitrile or methanol or an acetonitrile-methanol mixed solution) and starting compounds.







455

[Table 111]







Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
509	751		LCMS-D-1	2.57	519 (M+H) ⁺
510	752		LCMS-D-1	2.53	503 (M+H) ⁺
576	753		LCMS-D-1	2.48	533 (M+H) ⁺
512	754		LCMS-D-1	2.33	519 (M+H) ⁺
542	755		LCMS-D-1	2.30	629 (M+H) ⁺

[1428]

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
543	756		LCMS-D-1	2.43	643 (M+H)+
545	757		LCMS-D-1	2.30	661 (M+H)+
544	758		LCMS-D-1	2.11	659 (M+H)+
546	759		LCMS-D-1	1.56	599 (M+H)+
548	760		LCMS-D-1	2.08	640 (M+H)+
549	761		LCMS-D-1	1.62	686 (M+H)+

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
550	764		LCMS-D-1	2.22	627 (M+H) ⁺
515	765		LCMS-D-1	1.80	545 (M+H) ⁺
551	766		LCMS-D-1	2.42	675 (M+H) ⁺
582	767		LCMS-A-1	2.03	558 (M+H) ⁺
547	768		LCMS-D-1	2.57	572 (M+H) ⁺
552	769		LCMS-D-1	2.03	628 (M+H) ⁺

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
553	770		LCMS-D-1	2.38	558 (M+H)+
516	771		LCMS-D-1	2.58	543 (M+H)+
554	772		LCMS-D-1	2.47	599 (M+H)+
552	773		LCMS-D-1	2.08	517 (M+H)+
554	774		LCMS-D-1	2.32	640 (M+H)+
555	775		LCMS-D-1	2.57	668 (M+H)+






(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
556	776		LCMS-D-1	2.50	625 (M+H)+
557	777		LCMS-D-1	2.45	639 (M+H)+
581	778		LCMS-C-1	2.42	544 (M+H)+
518	779		LCMS-D-1	1.93	558 (M+H)+
519	780		LCMS-D-1	2.20	586 (M+H)+
520	781		LCMS-D-1	2.10	543 (M+H)+






(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
557	782		LCMS-D-1	1.98	557 (M+H) ⁺
584	783		LCMS-D-1	1.96	517 (M+H) ⁺
559	784		LCMS-D-1	2.47	613 (M+H) ⁺
517	785		LCMS-D-1	2.10	531 (M+H) ⁺
587	786		LCMS-D-1	1.72	558 (M+H) ⁺
629	787		LCMS-D-1	2.40	628 (M+H) ⁺
623	788		LCMS-C-1	2.85	651 (M+H) ⁺

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
622	789		LCMS-C-1	2.80	615 (M+H) ⁺
536	790		LCMS-C-1	2.82	560 (M+H) ⁺
630	791		LCMS-D-1	2.45	641 (M+H) ⁺
618	792		LCMS-F-1	0.89	594 (M+H) ⁺
617	793		LCMS-F-1	0.93	595 (M+H) ⁺

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
718	794		LCMS-F-1	0.94	613 (M+H) ⁺
537	795		LCMS-A-1	2.53	614 (M+H) ⁺
628	796		LCMS-F-1	0.96	520 (M+H) ⁺
535	797		LCMS-F-1	0.94	639 (M+H) ⁺
627	798		LCMS-A-1	2.43	580 (M+H) ⁺

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
635	799		LCMS-F-1	0.91	608 (M+H) ⁺
636	800		LCMS-F-1	0.91	610 (M+H) ⁺
639	801		LCMS-F-1	0.99	616 (M+H) ⁺
641	802		LCMS-F-1	0.99	671 (M+H) ⁺
642	803		LCMS-F-1	1.00	653 (M+H) ⁺







(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
644	804		LCMS-F-1	1.00	665 (M+H)+
506	812		LCMS-D-1	2.20	533 (M+H)+
692	817		LCMS-A-1	2.47	548 (M+H)+
694	818		LCMS-D-1	2.37	632 (M+H)+
699	820		LCMS-D-1	2.25	673 (M+H)+



(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
697	821		LCMS-D-1	2.21	618 (M+H)+
704	822		LCMS-D-1	2.28	657 (M+H)+
706	823		LCMS-D-1	2.25	687 (M+H)+
707	824		LCMS-D-1	2.40	661 (M+H)+
700	825		LCMS-A-1	2.35	607 (M+H)+
724	826		LCMS-D-1	2.12	674 (M+H)+

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
725	827		LCMS-D-1	2.37	644 (M+H) ⁺
726	828		LCMS-D-1	2.32	546 (M+H) ⁺
727	829		LCMS-D-1	2.68	518 (M+H) ⁺
733	830		LCMS-D-1	1.52	578 (M+H) ⁺
403	831		LCMS-F-1	0.91	667 (M+H) ⁺
539	832		LCMS-A-1	2.43	574 (M+H) ⁺

(continued)


Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
589	833		LCMS-B-2	4.38	631 (M+H)+
588	834		LCMS-C-1	2.47	517 (M+H)+

Example 185

2-Cyclohexyl-8-[2-(1H-indol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 835)

[1429]

(Reaction 185-1)








[1430] 2-Cyclohexyl-8-[2-(1H-indol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 835) was obtained by operations similar to those in Reaction 42-2 using Compound 479 as a starting material.

MS (ESI) m/z = 443 (M+H)⁺.

[1431] The example compounds shown below were obtained by operations similar to those in Example 185 using appropriate solvents (an ethanol-dimethylformamide mixed solution or ethanol) and starting compounds.





(Compounds 836 to Compound 879)

[Table 112]

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
492	836		LCMS-A-1	2.25	511 (M+H)+
488	837		LCMS-A-1	2.12	457 (M+H)+
487	838		LCMS-A-1	1.95	531 (M+H)+
499	839		LCMS-A-1	1.66	512 (M+H)+
524	840		LCMS-A-1	2.25	579 (M+H)+

[1432]



(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
502	841		LCMS-C-1	2.15	579 (M+H)+
497	842		LCMS-C-1	2.22	504 (M+H)+
527	843		LCMS-C-1	2.63	622 (M+H)+
844	844		LCMS-C-1	2.20	568 (M+H)+





(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
528	845		LCMS-C-1	2.58	651 (M+H)+
530	846		LCMS-C-1	2.37	652 (M+H)+
529	847		LCMS-C-1	2.48	568 (M+H)+
596	848		LCMS-C-1	2.90	748 (M+H)+
598	849		LCMS-C-1	2.77	650 (M+H)+

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
597	850		LCMS-C-1	2.58	595 (M+H)+
599	851		LCMS-C-1	2.42	649 (M+H)+
591	852		LCMS-C-1	2.53	650 (M+H)+
592	853		LCMS-C-1	2.43	657 (M+H)+






(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
595	854		LCMS-C-1	2.35	663 (M+H)+
601	855		LCMS-C-1	2.68	614 (M+H)+
619	856		LCMS-C-1	2.72	595 (M+H)+
620	857		LCMS-B-1	1.95	611 (M+H)+

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
621	858		LCMS-C-1	2.47	611 (M+H)+
602	859		LCMS-C-1	3.15	695 (M+H)+
600	860		LCMS-C-1	2.58	622 (M+H)+
572	861		LCMS-C-1	2.47	487 (M+H)+
505	862		LCMS-C-1	2.45	544 (M+H)+

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
511	863		LCMS-F-1	0.93	558 (M+H) ⁺
540	864		LCMS-F-1	0.96	640 (M+H) ⁺
513	865		LCMS-C-1	2.55	517 (M+H) ⁺
577	866		LCMS-C-1	2.60	517 (M+H) ⁺
541	867		LCMS-F-1	0.96	613 (M+H) ⁺

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
586	868		LCMS-A-1	1.84	541 (M+H)+
507	869		LCMS-F-1	0.96	572 (M+H)+
1185	870		LCMS-A-1	1.65	530 (M+H)+
702	872		LCMS-C-1	2.30	517 (M+H)+

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
717	873		LCMS-B-1	2.04	615 (M+H)+
473	874		LCMS-C-1	2.42	590 (M+H)+
743	875		LCMS-C-1	2.43	648 (M+H)+
729	876		LCMS-C-1	2.20	580 (M+H)+

(continued)


Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
731	877		LCMS-C-1	2.70	636 (M+H)+
624	878		LCMS-C-1	2.65	544 (M+H)+
625	879		LCMS-C-1	2.68	558 (M+H)+

Example 186

3-(4-{2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione (Compound 880)

[1433]

(Reaction 186-1)



[1434] 3-(4-{2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione (Compound 880) was obtained by operations similar to those in Reaction 91-1 using Compound 560 as a starting material.

MS (ESI) m/z = 572 (M+H)+.

[1435] The example compounds shown below were obtained by operations similar to those in Example 186 using appropriate solvents (an ethanol-dimethylformamide mixed solution or ethanol) and starting compounds.





(Compounds 881 to Compound 887)

[1436]

[Table 113]


Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
504	881		LCMS-C-1	2.33	530 (M+H)+
563	882		LCMS-C-1	2.67	558 (M+H)+
575	883		LCMS-C-1	2.43	544 (M+H)+

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
564	884		LCMS-C-1	2.70	572 (M+H)+
508	885		LCMS-F-1	0.97	586 (M+H)+
688	886		LCMS-F-1	0.86	582 (M+H)+
626	887		LCMS-C-1	2.70	572 (M+H)+

Example 187

3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzoic acid N,N'-dimethyl-hydrazide (Compound 888)

[1437]**(Reaction 187-1)**

[1438] 3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzoic acid N,N'-dimethyl-hydrazide (Compound 888) was obtained by operations similar to those in Reaction 10-14


using Compound 768 as a starting material and using dichloromethane as a solvent.
MS (ESI) m/z = 614 (M+H)⁺.

Example 188

8-{2-[2-Methyl-4-(piperidin-4-yloxy)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride (Compound 889)

[1439]

(Reaction 188-1)




[1440] 8-[2-[2-Methyl-4-(piperidin-4-yloxy)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one hydrochloride (Compound 889) was obtained by operations similar to those in Reaction 42-2 and Reaction 5-3 using Compound 602 as a starting material.
MS (ESI) m/z = 595 (M+H)+.

Example 189

4-{2-[2-(4,4-Difluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide
(Compound 890)

[1441]

(Reaction 189-1)



[1442] 4-{2-[2-(4,4-Difluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-

benzamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 523 (M-H)-.

[1443] The example compounds shown below were synthesized by operations similar to those in Reaction 189-1 using appropriate reagents and starting materials.






(Compounds 891 to 901)

[1444]

[Table 114]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
891		LCMS-C-1	2.07	561 (M+H)+
892		LCMS-A-1	2.28	577 (M+H)+
893		LCMS-C-1	2.42	507 (M+H)+
894		LCMS-C-1	2.93	545 (M+H)+
895		LCMS-C-1	2.85	545 (M+H)+
896		LCMS-C-1	2.63	517 (M+H)+

(continued)




Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
897		LCMS-B-2	4.91	599 (M+H)+
898		LCMS-C-2	2.88	660 (M-H)-
899		LCMS-A-1	2.24	594 (M+H)+
900		LCMS-C-2	1.93	582 (M+H)+
901		LCMS-C-2	2.33	594 (M-H)-

[1445] The spiroamine reagents used in the synthesis of Compounds 890, 891, 895 and 897 and shown below were synthesized by operations similar to those in Reaction 10-14, Reaction 1-4 and Reaction 4-1 using appropriate reagents and Compound 5a as a starting material.


[Table 115]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
890	 2TFA	272 (M+H)+

(continued)


Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
891		308 (M+H)+
895		292 (M+H)+
897		346 (M+H)+

[1446] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 897 (4-(4,4,4-trifluoro-but-1-enyl)-cyclohexanecarboxylic acid methyl ester) was synthesized by the method shown below.

(Reaction 189-2)


[1447] A 1.6 M solution of *n*-butyllithium in hexane (2.5 mL) was added dropwise to a suspension of triphenyl-(3,3,3-trifluoro-propyl)-phosphonium iodide (1.90 g, 3.91 mmol) in tetrahydrofuran (14 mL) at 0°C. The mixture was stirred at 0°C for 35 minutes, and a solution of 4-formyl-cyclohexanecarboxylic acid methyl ester (605 mg, 3.55 mmol) in tetrahydrofuran (8.0 mL) was then added dropwise to the reaction solution at -78°C. The mixture was stirred for 45 minutes, and a saturated aqueous ammonium chloride solution was then added, followed by extraction with *tert*-butyl methyl ether. The organic layer was washed with water and a saturated aqueous sodium chloride solution and washed with sodium sulfate. After concentration, the residue was purified by silica gel column chromatography to give 4-(4,4,4-trifluoro-but-1-enyl)-cyclohexanecarboxylic acid methyl ester (657 mg, 67%) as a colorless oily substance and geometric isomer mixture.

$^1\text{H-NMR}$ (CDCl_3) δ 5.69 (1.0H, t, J = 10.4 Hz), 5.51 (0.2H, dt, J = 13.7, 2.9 Hz), 5.32 (1.2H, tt, J = 9.2, 3.3 Hz), 3.69 (2.8H, dd, J = 3.0, 2.6 Hz), 3.67 (0.6H, d, J = 0.6 Hz), 2.90-2.78 (2.5H, m), 2.59-2.54 (1.0H, m), 2.40-2.31 (1.0H, m), 2.25-2.20 (0.5H, m), 2.06-1.98 (2.6H, m), 1.75-1.13 (8.0H, m).


(Reaction 189-3)

[1448] 4-(4,4,4-Trifluorobutyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 18-2 and Reaction 95-18 (using potassium hydroxide as a base) using appropriate reagents and starting material. This was used as such in the next reaction.

[1449] The spiroamine reagent used in the synthesis of Compound 892 (2-[(E)-2-(3-trifluoromethyl-phenyl)-vinyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride) was synthesized by the method shown below.


(Reaction 189-4)

[1450] 4-Carbamoyl-4-[(E)-3-(3-trifluoromethyl-phenyl)-acryloylamino]-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material. MS (ESI) $m/z = 440$ (M-H)-.

(Reaction 189-5)

[1451] A 6 N aqueous sodium hydroxide solution was added to a solution of 4-carbamoyl-4-[(E)-3-(3-trifluoromethyl-phenyl)-acryloylamino]-piperidine-1-carboxylic acid tert-butyl ester (961 mg, 2.27.mmol) in ethanol (20 ml), and the mixture was stirred at room temperature for 22 hours. The reaction solution was quenched with saturated ammonium chloride and then extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-oxo-2-[(E)-2-(3-trifluoromethyl-phenyl)-vinyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (706 mg, 73%).
MS (ESI) m/z = 422 (M-H)-.


(Reaction 189-6)



[1452] 2-[(E)-2-(3-Trifluoromethyl-phenyl)-vinyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride was synthesized by operations similar to those in Reaction 5-3 using appropriate reagents and starting material.
MS (ESI) m/z = 324 (M+H)+.


[1453] The spiroamine reagent used in the synthesis of Compound 893 (2-phenylethynyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized by the method shown below.

(Reaction 189-7)




[1454] 2-Phenylethynyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 10-14, Reaction 189-5 and Reaction 4-1 using appropriate reagents and starting material.
MS (ESI) m/z = 254 (M+H)+.

[1455] The spiroamine reagent used in the synthesis of Compound 898 (cyclohexylmethyl-[2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-dien-4-yl]-amine ditrifluoroacetate) was synthesized as follows.

(Reaction 189-8)


[1456] 4-Thioxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 88-1 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.42 (2H, dull d, J = 16.0 Hz), 1.50 (9H, s), 2.14 (2H, td, J = 16.0, 4.0 Hz), 3.33 (2H, br), 4.18 (2H, br), 7.44 (1H, m), 7.58 (1H, t, J = 8.0 Hz), 7.79 (1H, d, J = 8.0 Hz), 7.82 (1H, s), 10.30 (1H, br).

(Reaction 189-9)

[1457] Cyclohexyl-methylamine (0.044 ml, 0.34 mmol) was added to a solution of 4-thioxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (14.6 mg, 0.0340 mmol) in methanol (0.1 ml), and the mixture was stirred at 60°C for 24 hours and at 70°C for 11 hours. The reaction mixture was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-(cyclohexylmethyl-amino)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-diene-8-carboxylic acid tert-butyl ester (16.3 mg, 94%).

¹H-NMR (400 MHz, CDCl₃) δ 1.02 (2H, m), 1.24 (3H, m), 1.44 (2H, d, J = 13.2 Hz), 1.50 (9H, s), 1.65 (4H, m), 1.76 (4H, m), 3.40 (4H, m), 4.17 (2H, br), 5.12 (1H, br), 7.28 (1H, m), 7.44 (1H, t, J = 8.0 Hz), 7.08 (1H, dull s), 8.16 (1H, m).




(Reaction 189-10)

[1458] Cyclohexylmethyl-[2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-dien-4-yl]-amine ditrifluoroac-

etate was synthesized by operations similar to those in Reaction 4-1 using appropriate reagents and starting material. MS (ESI) m/z = 255 (M+H)+.

[1459] The spiroamine reagents used in the synthesis of Compounds 899 to 901 and shown below were synthesized by operations similar to those in Reaction 189-9 and Reaction 189-10 using appropriate reagents and starting materials.

[Table 116]


Target Compound	Spiroamine reagent	Spiroamine reagent $^1\text{H-NMR}$
899	 2TFA	$^1\text{H NMR}$ (400 MHz, CDCl_3) δ : 2.06 (4H, m), 3.24 (2H, m), 3.60 (3H, s), 3.70 (3H, s), 4.10 (2H, br), 7.60 (1H; d, J = 8 Hz), 7.65 (1H, t, J = 8 Hz), 8.32 (1H, s), 8.49 (1H, d, J = 8 Hz).
900	 2TFA	This was used in the next reaction without purification.
901	 2TFA	This was used in the next reaction without purification.

Example 190

4-(2-{2-[4-(2-Methoxy-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide (Compound 902)

[1460]

(Reaction 190-1)




[1461] 2-(4-Dimethylcarbamoyl-2-methyl-phenyl)-ethanesulfonyl chloride (22.2 mg) was added to a solution of 2-[4-(2-methoxy-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate (63.9 μmol) and magnesium oxide (20 mg) in tetrahydrofuran-water (4:1 (v/v), 640 μL), and the mixture was stirred at room temperature for 30 minutes. 2-(4-Dimethylcarbamoyl-2-methyl-phenyl)-ethanesulfonyl chloride (22.2 mg) was further added and the mixture was stirred for one hour. The reaction mixture was quenched with water and extracted with dichloromethane. The organic layer was concentrated, and the resulting residue was purified by silica gel column chromatography (ethyl acetate-hexane) to give 4-(2-{2-[4-(2-methoxy-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide (32.7 mg, 94%).


MS (ESI) $m/z = 547 (M+H)^+$.

[1462] The spiroamine reagent used in the synthesis of Compound 902 and shown below (2-[4-(2-methoxy-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized by operations similar to those in Reaction 10-14, Reaction 1-4 and Reaction 4-1 using appropriate reagents and Compound 5a as a starting material.

[Table 117]


Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
902	 2TFA	294 ($M+H$) ⁺

[1463] The carboxylic acid (4-(2-methoxy-ethyl)-cyclohexanecarboxylic acid) necessary for the synthesis of the spiroamine reagent used for Compound 902 (2-[4-(2-methoxy-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized by the method shown below.

(Reaction 190-2)

[1464] Potassium carbonate (2.61 g, 18.9 mmol) and benzyl bromide (2.24 mL, 18.9 mmol) were added to a solution of 4-hydroxymethyl-cyclohexanecarboxylic acid (cis-trans = 2.9:1 mixture) (2.49 g, 15.7 mmol) in DMF (31.5 mL) at room temperature, and the mixture was stirred at 60°C for one hour. The reaction solution was cooled, and H₂O (60 mL) was then added to the reaction solution, followed by extraction with hexane:ethyl acetate (2:1) (300 mL) twice. The organic layers were dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-hydroxymethyl-cyclohexanecarboxylic acid benzyl ester (cis-trans = 3.5:1 mixture) as a colorless oily substance (3.79 g, 97%).


MS (ESI) $m/z = 249 (M+H)^+$.

(Reaction 190-3)

[1465] 2,2,6,6-Tetramethylpiperidine 1-oxyl (309 mg, 1.98 mmol) and (diacetoxyiodo)benzene (7.01 g, 21.8 mmol) were added to a solution of 4-hydroxymethyl-cyclohexanecarboxylic acid benzyl ester (cis-trans = 4:1 mixture) (4.91 g,


19.8 mmol) at 0°C in an N₂ atmosphere, and the mixture was stirred at 0°C for one hour and at room temperature for seven hours. The reaction solution was diluted with dichloromethane (200 mL), and the organic layer was sequentially washed with a saturated aqueous sodium sulfite solution (100 mL), a saturated aqueous sodium bicarbonate solution (100 mL) and saturated brine (100 mL). The organic layer was dried over sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-formyl-cyclohexanecarboxylic acid benzyl ester (cis-trans = 4:1 mixture) as a colorless oily substance (4.44 g, 91%). MS (ESI) m/z = 247 (M+H)⁺.

(Reaction 190-4)



[1466] NaHMDS (1.0 M in THF) (466 μ L, 466 μ mol) was added to a solution of methoxymethyltriphenylphosphonium chloride (160 mg, 466 μ mol) in tetrahydrofuran (3.88 mL) at 0°C in an N₂ atmosphere, and the mixture was stirred at 0°C for one hour. A solution of 4-formyl-cyclohexanecarboxylic acid benzyl ester (cis-trans = 4:1 mixture) (95.6 mg, 388 μ mol) in tetrahydrofuran (2.00 mL) was added dropwise to the reaction solution at 0°C, and the mixture was stirred for 30 minutes. Thereafter, the reaction mixture was stirred at room temperature for 20 hours and quenched with a saturated aqueous ammonium chloride solution (5 mL). H₂O (20 mL) was then added, followed by extraction with dichloromethane (50 mL). The organic layer was dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-(2-methoxy-vinyl)-cyclohexanecarboxylic acid benzyl ester (trans-cis = 4:1 and E-Z = 2:1 mixture) as a yellow oily substance (49.7 mg, 47%). MS (ESI) m/z = 275 (M+H)⁺.

(Reaction 190-5)



[1467] 4-(2-Methoxy-ethyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 18-2 using appropriate reagents and starting material.





¹H-NMR (400 MHz, CDCl₃) δ 0.89-1.02 (2H, ddd, J = 3.8, 13.2, 24.9 Hz), 1.30-1.62 (5H, m), 1.79-1.85 (2H, br-m), 1.92-2.04 (2H, br-m), 2.25 (0.8H, tt, J = 3.4, 12.2 Hz), 2.58 (0.2H, quintet, J = 4.9 Hz), 3.32 (0.6H, s), 3.33 (2.4H, s), 3.41 (2H, t, J = 6.8 Hz).

Example 191








N-{4-[2-(2-Cyclopentyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamide (Compound 903)

[1468]



[Table 118]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
904		LCMS-C-1	2.45	543 (M+H)+
905		LCMS-C-1	2.85	531 (M+H)+
906		LCMS-A-1	2.05	494 (M+H)+
907		LCMS-A-1	2.05	494 (M+H)+

(continued)


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
908		LCMS-A-1	1.99	494 (M+H) ⁺
909		LCMS-C-1	2.85	531 (M+H) ⁺
910		LCMS-C-1	2.38	533 (M+H) ⁺
911		LCMS-C-1	2.48	533 (M+H) ⁺
912		LCMS-C-1	2.6	547 (M+H) ⁺
913		LCMS-C-1	2.52	547 (M+H) ⁺
914		LCMS-C-2	1.68	551 (M+H) ⁺

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
915		LCMS-C-2	1.65	551 (M+H)+
916		LCMS-C-2	1.85	551 (M+H)+

[1472] The spiroamine reagent used in the synthesis of Compound 906 (4-(4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-benzonitrile dihydrochloride) was synthesized by the following method.


(Reaction 191-2)



[1473] 2-(4-Carbamoyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-14 and Reaction 10-12 using appropriate reagents and starting material.

MS (ESI) m/z = 373 (M+H)+.


(Reaction 191-3)



10 **[1474]** Trifluoroacetic anhydride (0.287 ml, 2.07 mmol) was added to a solution of 2-(4-carbamoyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (350 mg, 0.94 mmol) and pyridine (0.303 ml) in dioxane (1.1 ml) at 0°C. The mixture was stirred for 30 minutes and then stirred at room temperature for one hour. An aqueous NaHCO₃ solution was added to the reaction mixture, followed by extraction with dichloromethane. The organic layer was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give 2-(4-cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester as a white powder (235 mg, 71%).

15 MS (ESI) m/z = 353 (M-H)-.

20 (Reaction 191-4)



30 **[1475]** 4-(4-Oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-benzonitrile dihydrochloride was synthesized by operations similar to those in Reaction 5-3 using appropriate reagents and starting material.

MS (ESI) m/z = 255 (M+H)+.






35 **[1476]** The spiroamine reagents used in the synthesis of Compounds 907 to 908 and shown below were synthesized by operations similar to those in Reaction 10-14, Reaction 10-12 and Reaction 5-3 using appropriate reagents and Compound 10ag as a starting material.

40 [Table 119]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
907	 2HCl	255 (M+H)+
908	 2HCl	255 (M+H)+


55 **[1477]** The spiroamine reagents used in the synthesis of Compounds 910, 911, 912 and 913 and shown below were synthesized by operations similar to those in Reaction 10-14, Reaction 1-4 and Reaction 4-1 using appropriate reagents and Compound 5a as a starting material.

[Table 120]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z) or ¹ H-NMR
910	 2TFA	292 (M-H)-
911	 2TFA	292 (M-H)-
912	 2TFA	306 (M-H)-
913	 2TFA	306 (M-H)-
916	 2TFA	¹ H-NMR (400MHz, CD ₃ OD) δ 1.24-1.33 (2H, m), 1.56-1.64 (2H, m), 1.93-1.96 (4H, m), 2.06-2.13 (5H, m), 2.50-2.70 (1H, m), 3.34-3.42 (2H, m), 3.51-3.57 (2H, m), 5.61-5.70 (1H, m), 5.96-6.25 (2H, m)


[1478] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 910 (4-ethoxymethyl-cyclohexanecarboxylic acid) was synthesized by the method shown below.

(Reaction 191-5)




[1479] 4-Ethoxymethyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 20-2 and Reaction 95-18 using appropriate reagents and starting material. This was used as such in the next reaction.

[1480] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 911 (4-propoxycyclohexanecarboxylic acid) was synthesized by the method shown below.

(Reaction 191-6)


[1481] 4-Propoxy-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 20-2 and Reaction 95-18 using appropriate reagents and starting material. This was used as such in the next reaction.

[1482] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 912 (4-butoxycyclohexanecarboxylic acid) was synthesized by the method shown below.

(Reaction 191-7)


[1483] 4-Butoxy-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 20-2 and Reaction 95-18 using appropriate reagents and starting material. This was used as such in the next reaction.

[1484] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 913 (4-isopropoxymethyl-cyclohexanecarboxylic acid) was synthesized by the method shown below.

(Reaction 191-8)


[1485] 4-Isopropoxymethyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 20-2 and Reaction 95-18 using appropriate reagents and starting material. This was used as such in the next reaction.

[1486] The spiroamine reagent used in the synthesis of Compound 914 (2-[4-(3-fluoro-propoxy)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized by the following method.

(Reaction 191-9)


[1487] N,N-Dimethylformamide di-tert-butyl acetal (7.4 ml, 31 mmol) was added to a solution of trans-4-hydroxy-cyclohexanecarboxylic acid (1.484 g, 10.29 mmol) in toluene (8.5 ml), and the mixture was stirred at 80°C for 25 hours. The reaction mixture was diluted with ether, and the organic layer was sequentially washed with water, an aqueous sodium bicarbonate solution and saturated brine, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give trans-4-hydroxy-cyclohexanecarboxylic acid tert-butyl ester as a colorless solid (838 m, 41%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.28 (2H, m), 1.43 (9H, s), 1.45 (2H, m), 1.99 (4H, m), 2.14 (1H, m), 3.60 (1H, m).

(Reaction 191-10)

[1488] cis-4-(3-Hydroxy-propoxy)-cyclohexanecarboxylic acid tert-butyl ester was obtained by operations similar to those in Reaction 20-2 and Reaction 39-2 using the compound obtained above and appropriate reagents.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.43 (9H, s), 1.53 (2H, m), 1.62 (2H, m), 1.83 (6H, m), 2.25 (1H, m), 2.60 (1H, t, $J = 5.4$ Hz), 3.46 (1H, m), 3.61 (2H, t, $J = 5.9$ Hz), 3.79 (2H, q, $J = 5.4$ Hz).


(Reaction 191-11)

[1489] Deoxo-Fluor (5 mg, 0.02 mmol) was added to a solution of cis-4-(3-hydroxy-propoxy)-cyclohexanecarboxylic

acid tert-butyl ester (3.9 mg, 0.015 mmol) in dichloromethane (0.1 ml), and the mixture was stirred at room temperature for two hours. An aqueous sodium bicarbonate solution was added to the reaction mixture, followed by extraction with dichloromethane. The organic layer was then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give cis-4-(3-fluoro-propoxy)-cyclohexanecarboxylic acid tert-butyl ester (3.1 mg, 79%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.44 (9H, s), 1.51 (2H, m), 1.60 (2H, m), 1.79 (2H, m), 1.94 (2H, m), 2.25 (1H, m), 2.60 (1H, t, $J = 5.4$ Hz), 3.43 (1H, m), 3.51 (2H, t, $J = 6.1$ Hz), 4.56 (2H, dt, $J = 47.4, 5.9$ Hz).

(Reaction 191-12)




[1490] 2-[4-(3-Fluoro-propoxy)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 4-1 (further adding water), Reaction 10-14, Reaction 10-12 and Reaction 4-1 using appropriate reagents and starting material. This was used as such in the next reaction.

[1491] The spiroamine reagent used in the synthesis of Compound 915 and shown below was synthesized by operations similar to those in Reaction 10-14, Reaction 10-12 and Reaction 4-1 using appropriate reagents and Compound 10ag as a starting material.

[Table 121]

Target Compound	Spiroamine reagent	Spiroamine reagent $^1\text{H-NMR}$
915	2TFA	This was used as such in the next reaction.

[1492] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 915 (4-(3-fluoro-propoxy)-cyclohexanecarboxylic acid) was synthesized by the method shown below.

(Reaction 191-13)


[1493] 4-(3-Fluoro-propoxy)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 20-2, Reaction 39-2, Reaction 191-11 and Reaction 4-1 (further adding water) using appropriate reagents and starting material. This was used as such in the next reaction.

[1494] The spiroamine reagent used in the synthesis of Compound 916 and shown below was synthesized by operations similar to those in Reaction 10-14, Reaction 1-4 and Reaction 4-1 using appropriate reagents and Compound 5a as a starting material.

[Table 122]


Target Compound	Spiroamine reagent	Spiroamine reagent $^1\text{H-NMR}$
916	2TFA	$^1\text{H-NMR}$ (400MHz, CD_3OD) δ 1.24-1.33 (2H, m), 1.56-1.64 (2H, m), 1.93-1.96 (4H, m), 2.06-2.13 (5H, m), 2.50-2.70 (1H, m), 3.34-3.42 (2H, m), 3.51-3.57 (2H, m), 5.61-5.70 (1H, m), 5.96-6.25 (2H, m)

[1495] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 916 (4-((E)-3,3-difluoro-propenyl)-cyclohexanecarboxylic acid) was synthesized by the method shown below.


(Reaction 191-14)

[1496] Potassium t-butoxide (68.3 mg, 609 μmol) was added to a solution of (1,3-dioxolan-2-ylmethyl)-triphenylphos-

phonium bromide (267 mg, 609 μmol) in THF (2.0 ml) at 0°C, and the mixture was stirred at 0°C for 1.5 hours in an N_2 atmosphere. A solution of 4-formyl-cyclohexanecarboxylic acid benzyl ester (50.0 mg, 203 μmol) in THF (1.5 ml) was added to the reaction mixture at 0°C, and the mixture was stirred at room temperature for 1.5 hours. Thereafter, the reaction mixture was quenched by adding a saturated aqueous ammonium chloride solution at 0°C and then extracted with ethyl acetate three times. The organic layers were sequentially washed with H_2O (x 2) and saturated brine, and then dried over MgSO_4 and concentrated under reduced pressure. The residue was used in the next step without further purification.


(Reaction 191-15)

[1497] 1 N hydrochloric acid (406 μl , 406 μl) was added to a solution of the residue obtained in Reaction 191-14 in THF (2.0 ml) at 0°C, and the mixture was stirred at room temperature for 4.5 hours. The reaction solution was quenched by adding a saturated aqueous sodium bicarbonate solution at 0°C and then extracted with ethyl acetate three times. The organic layers were sequentially washed with H_2O (x 2) and saturated brine, and then dried over MgSO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (hexane-ethyl acetate) to give 4-((E)-3-oxo-propenyl)-cyclohexanecarboxylic acid benzyl ester as a colorless oil (38.2 mg, 69%). MS (ESI) m/z = 273 (M+H)+.

(Reaction 191-16)

[1498] 4-((E)-3,3-Difluoro-propenyl)-cyclohexanecarboxylic acid benzyl ester was synthesized by operations similar to those in Reaction 191-11 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.46-1.52 (2H, m), 1.61-1.67 (3H, m), 1.88-2.30 (4H, m), 2.60-2.70 (1H, m), 5.58-5.64 (1H, m), 5.88-6.17 (2H, m).

(Reaction 191-17)

[1499] 4-((E)-3,3-Difluoro-propenyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material.


¹H-NMR (400 MHz, CDCl₃) δ 1.46-1.52 (2H, m), 1.61-1.67 (3H, m), 1.88-2.30 (4H, m), 2.60-2.70 (1H, m), 5.58-5.64 (1H, m), 5.88-6.17 (2H, m).

Example 192

N-{4-[2-(2-Cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamide (Compound 917)

[1500]

(Reaction 192-1)



[1501] N-{4-[2-(2-Cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamide was synthesized by operations similar to those in Reaction 190-1 using appropriate reagents and starting material. MS (ESI) m/z = 489 (M+H)+.

[1502] The example compounds shown below were synthesized by operations similar to those in Reaction 192-1 using appropriate reagents and starting materials.

(Compounds 918 to 919)

[1503]


[Table 123]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
918		LCMS-C-1	2.65	527 (M+H)+
919		LCMS-B-1	2.03	587 (M+H)+

[1504] The spiroamine reagent used in the synthesis of Compound 918 (2-adamantan-1-yl-1,3,8-triaza-spiro[4.5]dec-

1-en-4-one ditrifluoroacetate) was synthesized by the following method.

(Reaction 192-2)



[1505] 2-Adamantan-1-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 10-14, Reaction 10-11, Reaction 10-12 and Reaction 4-1 using appropriate reagents and starting material.


MS (ESI) m/z = 288 (M+H)⁺.

[1506] The spiroamine reagent used in the synthesis of Compound 919 was synthesized by operations similar to those in Reaction 10-14, Reaction 1-4 and Reaction 4-1 using appropriate reagents and Compound 5a as a starting material.

[Table 124]


Target Compound	Spiroamine reagent	Spiroamine reagent MS(m/z)
919	2TFA	348 (M+H) ⁺

[1507] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 919 (4-(2,2,2-trifluoro-ethoxymethyl)-cyclohexanecarboxylic acid) was synthesized by the method shown below.

(Reaction 192-3)

[1508] 2,2,2-Trifluoro-ethanol (288 μL , 4.03 mmol) was added to a mixed solution of 4-hydroxymethyl-cyclohexanecarboxylic acid benzyl ester (100 mg, 0.403 mmol), 1,1'-azobis(N,N-dimethylformamide) (139 mg, 0.805 mmol) and tributyl-phosphine (199 μL , 0.805 mmol) in toluene (1.2 mL) at 0°C . The mixture was stirred at 65°C for 1.5 hours and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-(2,2,2-trifluoro-ethoxymethyl)-cyclohexanecarboxylic acid benzyl ester as a colorless liquid (126 mg, 95%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.94-1.06 (0.4H, m), 1.23-1.47 (1.8H, m), 1.40-1.52 (0.4H, m), 1.55-1.68 (3.2H, m), 1.69-1.80 (0.8H, m), 1.82-1.91 (0.4H, m), 1.97-2.08 (2H, m), 2.25-2.34 (0.2H, m), 2.58-2.65 (0.8H, m), 3.41 (0.4H, d, $J = 6.8$ Hz), 3.43 (1.6H, d, $J = 6.8$ Hz), 3.78 (2H, q, $J = 8.8$ Hz), 5.11 (0.4H, s), 5.13 (1.6H, s), 7.29-7.40 (5H, m).

(Reaction 192-4)


[1509] 4-(2,2,2-Trifluoro-ethoxymethyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 18-2 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.96-2.23 (9H, m), 2.30-2.90 (1H, m), 3.37-3.49 (2H, m), 3.79 (2H, q, $J = 8.8$ Hz), 9.56 (1H, brs).

Example 193

N-[4-(2-{2-[4-(4-Chloro-phenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide (Compound 920)

[1510]

(Reaction 193-1)

[1511] N-[4-(2-(2-[4-(4-Chloro-phenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 586 (M+H)^+$.

[1512] The example compounds shown below were synthesized by operations similar to those in Reaction 193-1 using appropriate reagents and starting materials.




(Compounds 921 to 926)

[1513]

[Table 125]



Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
921		LCMS-B-1	1.79	587 ($M+H$) ⁺
922		LCMS-C-1	2.53	557 ($M+H$) ⁺
923		LCMS-C-1	2.78	517 ($M+H$) ⁺

(continued)


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
924		LCMS-C-1	2.48	489 (M+H)+
925		LCMS-C-1	2.63	503 (M+H)+
926		LCMS-C-1	2.85	531 (M+H)+

[1514] The spiroamine reagents used in the synthesis of Compounds 920 and 921 and shown below were synthesized by operations similar to those in Reaction 10-14, Reaction 1-4 and Reaction 4-1 using appropriate reagents and Compound 5a as a starting material.

[Table 126]


Target Compound	Spiroamine reagent	Spiroamine reagent MS(m/z)
920	 2TFA	346 (M+H)+
921	 2TFA	348 (M+H)+

[1515] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 921 (4-(3,3,3-trifluoro-propoxy)-cyclohexanecarboxylic acid) was synthesized by the method shown below.

(Reaction 193-2)

[1516] 4-(3,3,3-Trifluoro-propoxy)-benzoic acid benzyl ester was synthesized by operations similar to those in Reaction 31-7 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.63 (2H, qt, $J = 10.4$, 6.8 Hz), 4.23 (2H, t, $J = 6.0$ Hz), 5.32 (2H, s), 6.90 (1H, d, $J = 8.8$ Hz), 7.30-7.43 (5H, m), 8.02 (1H, d, $J = 8.8$ Hz).

(Reaction 193-3)


[1517] 10% Rh-C (14.7 mg) was added to a solution of 4-(3,3,3-trifluoro-propoxy)-benzoic acid benzyl ester (147.3 mg, 0.454 mmol) in iPrOH (1.5 mL). The hydrogen pressure was adjusted to 5 atm, and the mixture was then heated with stirring at 80°C overnight. The reaction mixture was filtered through celite, and the filtrate was then diluted with ethyl acetate. A saturated aqueous sodium bicarbonate solution was added, and the organic layer and the aqueous layer were separated. The aqueous layer was adjusted to pH 1 with 1 N hydrochloric acid and then extracted with ethyl acetate. The organic layers were sequentially washed with water and saturated brine, and then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give 4-(3,3,3-trifluoro-propoxy)-cyclohexanecarboxylic acid as a colorless transparent oily substance (70.2 mg, 64%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.19-2.08 (8H, m), 2.27-2.43 (3H, m), 3.23 (0.2H, tt, $J = 11.2$, 4.0 Hz), 3.45-3.49 (0.8H, m), 3.59 (1.6H, t, $J = 6.8$ Hz), 3.66 (0.4H, t, $J = 6.8$ Hz).


[1518] The spiroamine reagent used in the synthesis of Compound 922 (2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized by the method shown below.

(Reaction 193-4)


[1519] DMF (10 mL) was added to a reaction vessel containing 4-formyl-benzoic acid methyl ester (501.1 mg, 3.053 mmol) and potassium acetate (15.0 mg, 0.153 mmol), and the mixture was cooled to 0°C. Trimethyl(trifluoromethyl)silane (0.96 mL, 6.105 mmol) was added dropwise and the mixture was stirred for 50 minutes. 2 N hydrochloric acid (10 mL) was then added to the reaction mixture, and the mixture was stirred at room temperature overnight and then diluted with ethyl acetate. A saturated aqueous sodium bicarbonate solution was added, and the organic layer and the aqueous layer were separated. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-(2,2,2-trifluoro-1-hydroxy-ethyl)-benzoic acid methyl ester (680.8 mg, 95%).
¹H-NMR (400 MHz, CDCl₃) δ 2.63 (1H, d, J = 5.2 Hz), 3.92 (3H, s), 5.06-5.12 (1H, m), 7.55 (2H, d, J = 8.4 Hz), 8.07 (2H, d, J = 8.4 Hz).

(Reaction 193-5)

[1520] Toluene (26 mL) was added to a reaction vessel containing 4-(2,2,2-trifluoro-1-hydroxy-ethyl)-benzoic acid methyl ester (607.2 mg, 2.593 mmol), DMAP (633.6 mg, 5.186 mmol) and Molecular Sieve 4A (916.1 mg). Phenyl chlorothioformate (0.54 mL, 3.889 mmol) was added dropwise and the mixture was stirred overnight. The reaction mixture was filtered through celite, and the filtrate was then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-(2,2,2-trifluoro-1-phenoxythiocarbonyloxy-ethyl)-benzoic acid methyl ester as a colorless oily substance (900.5 mg, 94%).
¹H-NMR (400 MHz, CDCl₃) δ 3.93 (3H, s), 6.62 (1H, q, J = 6.4 Hz), 7.06-7.09 (2H, m), 7.27-7.31 (1H, m), 7.38-7.42 (2H, m), 7.60 (2H, d, J = 8.4 Hz), 8.11 (2H, d, J = 8.4 Hz).


(Reaction 193-6)

[1521] 4-(2,2,2-Trifluoro-1-phenoxythiocarbonyloxy-ethyl)-benzoic acid methyl ester (462.8 mg, 1.25 mmol) and AIBN (41.0 mg, 0.25 mmol) were dissolved in ultrasonically degassed toluene (12.5 mL). Tri-n-butyltin hydride (0.50 mL, 1.874 mmol) was added and the mixture was heated with stirring at 80°C for two hours. The reaction solution was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography to give 4-(2,2,2-trifluoro-ethyl)-benzoic acid methyl ester as white crystals (254.6 mg, 93%).
¹H-NMR (400 MHz, CDCl₃) δ 3.41 (2H, q, J = 10.8 Hz), 3.91 (3H, s), 7.36 (2H, d, J = 8.4 Hz), 8.02 (2H, d, J = 8.4 Hz).

(Reaction 193-7)

[1522] 4-(2,2,2-Trifluoro-ethyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 95-18 and Reaction 193-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.03-2.08 (11H, m), 2.28 (0.33H, tt, $J = 12.0, 3.2$ Hz), 2.61-2.64 (0.66H, m).

(Reaction 193-8)

[1523] 2-[4-(2,2,2-Trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 10-14, Reaction 10-8 and Reaction 4-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 318$ ($\text{M}+\text{H}$) $^+$.


[1524] The spiroamine reagent used in the synthesis of Compound 923 and shown below was synthesized by operations similar to those in Reaction 10-14, Reaction 10-8 and Reaction 4-1 using appropriate reagents and Compound 5a as a starting material.

[Table 127]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
923	 2TFA	278 ($\text{M}+\text{H}$) $^+$

[1525] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 923 (3-propyl-cyclohexanecarboxylic acid) was synthesized by the method shown below.


(Reaction 193-9)



[1526] A suspension solution of ethyltriphenylphosphonium bromide (1079.3 mg, 2.907 mmol) in THF (10 mL) was cooled to 0°C. LHMDS (2.781 mL, 2.781 mmol, 1.0 M in THF) was added dropwise, and the mixture was stirred for 30 minutes. A solution of 3-formyl-benzoic acid methyl ester (415.0 mg, 2.528 mmol) in THF (2.5 mL) was then added dropwise, and the mixture was stirred for 10 minutes and then stirred at room temperature overnight. The reaction mixture was quenched by adding a saturated aqueous ammonium chloride solution and then extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 3-propenyl-benzoic acid methyl ester as a yellow transparent oily substance (218.2 mg, 49%).

¹H-NMR (400 MHz, CDCl₃) δ 1.89-1.92 (3H, m), 3.92 (1H, s), 3.93 (2H, s), 5.86 (0.66H, dq, J = 11.6, 7.2 Hz), 6.32 (0.33H, dq, J = 15.6, 6.4 Hz), 6.41-6.47 (1H, m), 7.34-7.43 (1H, m), 7.47-7.51 (1H, m), 7.84-7.90 (1H, m), 7.97-8.01 (1H, m).


(Reaction 193-10)



[1527] 3-Propyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 95-18 and Reaction 193-3 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 0.80-2.05 (16H, m), 2.33 (0.6H, tt, J = 12.4, 3.2 Hz), 2.67-2.70 (0.4H, m).



[1528] The spiroamine reagent used in the synthesis of Compound 924 (2-(3-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate) was synthesized by the method shown below.

(Reaction 193-11)


[1529] 2-(3-Methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 176-2, Reaction 10-8 and Reaction 4-1 using appropriate reagents and starting material. MS (ESI) $m/z = 250$ (M+H)+.

[1530] The spiroamine reagents used in the synthesis of Compounds 925 and 926 and shown below were synthesized by operations similar to those in Reaction 10-14, Reaction 10-8 and Reaction 4-1 using appropriate reagents and starting materials.

[Table 128]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
925	 2TFA	264 (M+H)+
926	 2TFA	292 (M+H)+

[1531] The carboxylic acid necessary for the synthesis of the spiroamine reagent used for Compound 926 (3,3,5,5-tetramethyl-cyclohexanecarboxylic acid) was synthesized by the method shown below.

(Reaction 193-12)


[1532] A solution of [1,3]dithian-2-yl-trimethyl-silane (566.4 mg, 2.944 mmol) in THF (6 mL) was cooled to 0°C. nBuLi (1.78 mL, 2.845 mmol, 1.6 M in n-hexane) was added dropwise and then the mixture was stirred for 10 minutes. The reaction solution was cooled to -78°C. A solution of 3,3,5,5-tetramethyl-cyclohexanone (302.7 mg, 1.962 mmol) in THF (2 mL) was then added dropwise, and the mixture was stirred for two hours. The reaction mixture was quenched by adding a saturated aqueous ammonium chloride solution and extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure.

[1533] The resulting residue was dissolved in acetonitrile (2.1 mL). Water (0.52 mL) and trifluoroacetic acid (0.51 mL) were added and the mixture was heated with stirring at 65°C for three hours. The reaction solution was cooled to room temperature. A 30% aqueous hydrogen peroxide solution (3.2 mL) was then added and the mixture was heated with stirring at 80°C for one hour. The reaction solution was cooled to room temperature, and a 5 M aqueous sodium hydroxide solution (15.7 mL) was then added, followed by extraction with ether. A saturated aqueous sodium bicarbonate solution was added, and the organic layer and the aqueous layer were separated. The aqueous layer was adjusted to pH 1 with 2 N hydrochloric acid and then extracted with ethyl acetate. The organic layers were sequentially washed with water and saturated brine, and then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give 3,3,5,5-tetramethyl-cyclohexanecarboxylic acid as a white powder (342.3 mg, 95% in two steps).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.93 (6H, s), 1.01 (6H, s), 1.06-1.28 (4H, m), 1.68-1.71 (2H, m), 2.65 (1H, tt, $J = 12.8, 3.2$ Hz).

Example 194

N-[4-(2-{4-(2-Methoxy-ethyl)-cyclohexyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide (Compound 927)

[1534](Reaction 194-1)


[1535] N-[4-(2-{4-(2-Methoxy-ethyl)-cyclohexyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3-methyl-phenyl]-acetamide was synthesized by operations similar to those in Reaction 190-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 533$ ($\text{M}+\text{H}$) $^+$.

Example 195

[3-Methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-carbamic acid tert-butyl ester (Compound 928)

[1536]

(Reaction 195-1)


[1537] [3-Methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.
MS (ESI) m/z = 643 ($\text{M}+\text{H}$) $^+$.

Example 196

N-(2-Hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-isobutylamide (Compound 929)

[1538]

(Reaction 196-1)




[1539] N-(2-Hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-isobutylamide was synthesized by operations similar to those in Reaction 5-4, Reaction 96-16, Reaction 157-2, Reaction 105-2 and Reaction 39-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 625$ ($M+H$)⁺.

Example 197

2-Hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide (Compound 930)

[1540]

(Reaction 197-1)


[1541] 2-Hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide was synthesized by operations similar to those in Reaction 105-2, Reaction 39-2 and Reaction 12-5 using appropriate reagents and starting material.
MS (ESI) $m/z = 613$ (M+H)⁺.

Example 198

N-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-2,2,2-trifluoro-N-methyl-acetamide (Compound 931)

[1542]


(Reaction 198-1)



[1543] 2-(2,6-Dimethyl-4-[methyl-(2,2,2-trifluoro-acetyl)-amino]-phenyl)-ethanesulfonyl chloride was synthesized by operations similar to those in Reaction 10-2, Reaction 4-1, Reaction 19-2, Reaction 10-3, Reaction 10-4 and Reaction 10-5 using appropriate reagents and starting material.

MS (ESI) $m/z = 358$ ($M+H$) $^+$.

(Reaction 198-2)




[1544] N-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-2,2,2-trifluoro-N-methyl-acetamide was synthesized by operations similar to those in Reaction 190-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 571$ ($M+H$) $^+$.

Example 199

N-(3,5-Dimethyl-4-{2-[2-(3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-2,2,2-trifluoro-N-methyl-acetamide (Compound 932)

[1545]

(Reaction 199-1)

[1546] N-(3,5-Dimethyl-4-{2-[2-(3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-2,2,2-trifluoro-N-methyl-acetamide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 571 (M+H)+.

[1547] The example compound shown below was synthesized by operations similar to those in Reaction 199-1 using appropriate reagents and starting material.

(Compound 933)

[1548]

[Table 129]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
933		LCMS-C-1	3.13	613 (M+H)+

Example 200

1-{4-[2-(2-Cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea (Compound 934)

[1549]

(Reaction 200-1)



[1550] 1-{4-[2-(2-Cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-1-methyl-urea was synthesized by operations similar to those in Reaction 5-4, Reaction 12-5 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.

MS (ESI) $m/z = 518 \text{ (M+H)}^+$.

[1551] The example compounds shown below were synthesized by operations similar to those in Reaction 200-1 using appropriate reagents and starting materials.

(Compounds 935, 938 and 941)

[1552]


[Table 130]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
935		LCMS-A-1	2.35	582 (M+H)+
938		LCMS-C-1	2.70	532 (M+H)+

(continued)


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
941		LCMS-F-1	0.93	554 (M+H)+

[1553] The spiroamine reagent used in the synthesis of Compound 941 (6-(3-trifluoromethoxy-phenyl)-2,5,7-triaza-spiro[3.4]oct-5-en-8-one ditrifluoroacetate) was synthesized as follows.

(Reaction 200-2)


[1554] Diallylamine (0.31 ml, 2.5 mmol) and trimethylsilylnitrile (0.155 ml, 1.25 mmol) were added to a solution of 3-oxo-azetidine-1-carboxylic acid tert-butyl ester (171 mg, 1.00 mmol) in acetic acid (1.7 ml, 30 mmol), and the mixture was stirred at 60°C for four hours. A Saturated aqueous sodium bicarbonate solution (11.5 ml) was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 3-cyano-3-diallylamino-azetidine-1-carboxylic acid tert-butyl ester (212 mg, 76%).

¹H-NMR (400 MHz, CDCl₃) δ 1.45 (9H, s), 3.10 (4H, d, J = 7.0 Hz), 4.01 (2H, d, J = 8.6 Hz), 4.09 (2H, d, J = 8.6 Hz), 5.19 (1H, d, J = 10.2 Hz), 5.30 (1H, d, J = 17.0 Hz), 5.82 (1H, m).

(Reaction 200-3)

[1555] A solution of 3-cyano-3-diallylamino-azetidine 1-carboxylic acid tert-butyl ester (143.5 mg, 0.5174 mmol), 1,3-dimethylbarbituric acid (242.5 mg, 1.553 mmol) and tetrakis(triphenylphosphine)palladium(0) (30.3 mg, 0.0262 mmol) in dichloromethane (1.3 ml) was stirred at 40°C for five hours. A saturated aqueous sodium bicarbonate solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography

(hexane-ethyl acetate) to give 3-amino-3-cyano-azetidine-1-carboxylic acid tert-butyl ester (98 mg, 96%).
¹H-NMR (400 MHz, CDCl₃) δ 1.44 (9H, s), 2.03 (2H, br), 3.88 (2H, d, J = 8.8 Hz), 4.34 (2H, d, J = 8.8 Hz).

(Reaction 200-4)

[1556] 6-(3-Trifluoromethoxy-phenyl)-2,5,7-triaza-spiro[3.4]oct-5-en-8-one ditrifluoroacetate was synthesized by operations similar to those in Reaction 10-11, Reaction 10-14, Reaction 10-12 and Reaction 4-1 using appropriate reagents and starting material.


¹H-NMR (400 MHz, CDCl₃) δ 4.31 (2H, d, J = 12.0 Hz), 4.40 (2H, d, J = 12.0 Hz), 7.56 (1H, d, J = 8.2 Hz), 7.66 (1H, t, J = 8.2 Hz), 7.95 (1H, d, J = 8.2 Hz), 7.96 (1H, s).

Example 201

1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea (Compound 936) and 1-[2-chloro-3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoroethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea (Compound 937)

[1557]


(Reaction 201-1)



[1558] A mixture of [4-(2-chlorosulfonyl-ethyl)-3,5-dimethyl-phenyl]-methyl-carbamic acid tert-butyl ester and [2-chloro-4-(2-chlorosulfonyl-ethyl)-3,5-dimethyl-phenyl]-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-2 (using RuPhos as a ligand), Reaction 10-3, Reaction 10-4 and Reaction 10-5 using appropriate reagents and starting material.

(R=H : R=Cl = 0.6 : 0.4)

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.32-1.54 (9H, m), 2.32-2.47 (6H, m), 3.08-3.25 (3H, m), 3.29-3.44 (2H, m), 3.60-3.74 (2H, m), 6.96 (1.6H, m).

(Reaction 201-2)

[1559] 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea

MS (ESI) m/z = 586 (M+H)⁺

and

1-[2-chloro-3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea

MS (ESI) m/z = 620 (M+H)⁺


were obtained by operations similar to those in Reaction 5-4, Reaction 4-1 and Reaction 89-2 (using KOCN) using the starting material obtained above and appropriate reagents.

Example 202

1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propylidene)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea (Compound 939)

[1560]

(Reaction 202-1)



[1561] 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propylidene)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 5-4, Reaction 5-3 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.

MS (ESI) $m/z = 598 (M+H)^+$.

[1562] The example compound shown below was synthesized by operations similar to those in Reaction 202-1 using appropriate reagents and starting material.


(Compound 940)

[1563]

[Table 131]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
940		LCMS-B-1	1.91	580 ($M+H$) ⁺

[1564] The spiroamine reagent used in the synthesis of Compound 939 (2-[4-(3,3,3-trifluoro-propylidene)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride) was synthesized as follows.

(Reaction 202-2)

[1565] 2-[4-(3,3,3-Trifluoro-propylidene)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride was synthesized by operations similar to those in Reaction 10-14, Reaction 10-12 and Reaction 5-3 using appropriate reagents and starting material.

MS (ESI) $m/z = 330 (M+H)^+$.


[1566] The spiroamine reagent used in the synthesis of Compound 940 and shown below was synthesized by operations similar to those in Reaction 10-14, Reaction 10-12 and Reaction 5-3 using appropriate reagents and Compound 10ag as a starting material.

[Table 132]

Target Compound	Spiroamine reagent	Spiroamine reagent MS (m/z)
940	2HCl	312 ($M+H$) ⁺

(Reaction 202-3)


[1567] The carboxylic acid derivative necessary for the synthesis of the spiroamine reagent used in the synthesis of Compound 940 (4-(3,3-difluoro-allyl)-cyclohexanecarboxylic acid) was synthesized in the following manner.



[1568] 4-(2-Oxo-ethyl)-cyclohexanecarboxylic acid benzyl ester (trans:cis = 4:1) was synthesized by operations similar to those in Reaction 25-4 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.04 (1.6H, m), 1.30 (0.4H, m), 1.51 (1.6H, m), 1.62 (0.8H, m), 1.84 (1.6H, m), 1.89 (0.8H, m), 2.10 (2.2H, m), 2.32 (2.8H, m), 2.60 (0.2H, m), 5.11 (1.6H, s), 5.13 (0.4H, s), 7.35 (5H, m), 9.75 (0.2H, t, J = 2.0 Hz), 9.76 (0.8H, t, J = 2.0 Hz).


(Reaction 202-4)



[1569] A solution of 4-(2-oxo-ethyl)-cyclohexanecarboxylic acid benzyl ester (trans:cis = 4:1) (21.4 mg, 0.082 mmol) in dimethylformamide (0.3 ml) was added to a solution of sodium chlorodifluoroacetate (34.1 mg, 0.224 mmol) and triphenylphosphine (59.9 mg, 0.228 mmol) in dimethylformamide (0.41 ml) at 90 to 95°C over five minutes, and the mixture was stirred at 130°C for four hours. Sodium chlorodifluoroacetate (34.0 mg, 0.22 mmol) was then added to the reaction mixture at the same temperature, and the mixture was further stirred for two hours. The reaction mixture was diluted with ether, and the organic layer was washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-(3,3-difluoro-allyl)-cyclohexanecarboxylic acid benzyl ester (trans:cis = 4:1) (14.1 m, 58%).

¹H-NMR (400 MHz, CDCl₃) δ 0.95 (1.6H, m), 1.26 (0.4H, m), 1.44 (1.6H, m), 1.55 (0.8H, m), 1.81 (1.6H, m), 1.87 (2.2H, m), 2.20 (2 H, m), 2.28 (0.8H, m), 2.60 (0.2H, m), 4.09 (0.2H, dtd, J = 25.4, 8.3, 2.9 Hz), 4.12 (0.8H, dtd, J = 25.4, 7.8, 2.9 Hz), 5.11 (1.6H, s), 5.13 (0.4H, s), 7.34 (5H, m).

(Reaction 202-5)




[1570] A 1 N aqueous NaOH solution (0.084 ml, 0.084 mmol) was added to a solution of 4-(3,3-difluoro-allyl)-cyclohexanecarboxylic acid benzyl ester (trans:cis = 4:1) (14.1 mg, 0.0478 mmol) in methanol (1.0 mL). The mixture was stirred at room temperature for 1.5 hours, and then adjusted to pH 6 with a 1 N aqueous HCl solution and concentrated under reduced pressure. The resulting residue was adjusted to pH 3 with dilute hydrochloric acid and extracted with dichloromethane, and the organic layer was dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was dissolved in THF (0.2 ml)-H₂O (0.2 ml), and LiOH·H₂O (7.7 mg, 0.18 mmol) was added. The mixture was stirred at room temperature for three hours, and then adjusted to pH 3 with a 1 N aqueous HCl solution and extracted with dichloromethane. The organic layer was then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by column chromatography (hexane-ethyl acetate) to give 4-(3,3-difluoro-allyl)-cyclohexanecarboxylic acid (8.3 g, 86%).

¹H-NMR (400 MHz, CDCl₃) δ 0.97 (1.8H, m), 1.27 (0.4H, m), 1.43 (1.8H, m), 1.58 (1H, m), 1.82 (2H, m), 1.88 (2H, m), 2.03 (2H, m), 2.25 (0.9H, m), 2.61 (0.1H, m), 4.13 (1H, dtd, J = 25.4, 7.8, 2.9 Hz).

Example 203


N-(4-((E)-1-Fluoro-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-methyl-acetamide (Compound 942)

[1571]

(Reaction 203-1)


[1572] Tributyl(1-ethoxyvinyl)tin (1.07 mmol, 3.18 mmol) and dichlorobis(triphenylphosphine)palladium(II) (101 mg, 0.145 mmol) were added to a solution of N-(4-bromo-3-methylphenyl)-N-methylacetamide (700 mg, 2.89 mmol) in 1,4-dioxane (7 mL), and the mixture was heated with stirring at 90°C for 12 hours in a nitrogen stream. The reaction mixture was cooled and then filtered through celite. The solution was concentrated under reduced pressure, and the residue was then silica gel column chromatography (hexane-ethyl acetate) to give N-[4-(1-ethoxy-vinyl)-3-methyl-phenyl]-N-methyl-acetamide (440 mg, 65%).

MS (ESI) m/z = 234 (M+H)+.


(Reaction 203-2)

[1573] N-[4-(2-Bromo-1,1-difluoroethyl)-3-methylphenyl]-N-methylacetamide was synthesized by operations similar to those in Reaction 127-4 and Reaction 191-11 using appropriate reagents and starting material.


MS (ESI) m/z = 306, 308 (M+H)+.

(Reaction 203-3)

[1574] t-Dodecanethiol (0.227 mL, 0.96 mmol) was added to a solution of potassium t-butoxide (108 mg, 0.96 mmol) in DMF (2 mL), and the mixture was stirred at room temperature. A solution of N-[4-(2-bromo-1,1-difluoroethyl)-3-methylphenyl]-N-methylacetamide (245 mg, 0.800 mmol) in DMF (2 mL) was added to the mixture which was then stirred at room temperature for one hour. Saturated NH_4Cl was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was washed with saturated brine, and then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give N-[4-(2-(1,1-dimethyldecylsulfanyl)-1,1-difluoroethyl)-3-methylphenyl]-N-methylacetamide (287 mg, 84%). MS (ESI) $m/z = 428$ ($\text{M}+\text{H}^+$).

(Reaction 203-4)

[1575] 2 N HCl (0.4 mL) was added to a solution of N-[4-(2-(1,1-dimethyldecylsulfanyl)-1,1-difluoroethyl)-3-methylphenyl]-N-methylacetamide (102 mg, 0.239 mmol) in MeCN (1 mL) at 0°C . After stirring for five minutes, t-butyl hypochlorite (0.135 mL, 1.20 mmol) was added in small portions at -10°C . The mixture was stirred for 15 minutes, and saturated NH_4Cl was then added, followed by extraction with ethyl acetate. The organic layer was washed with saturated brine. The organic layer was dried over MgSO_4 and then concentrated under reduced pressure to give a mixture containing 2-[4-(acetylmethylamino)-2-methylphenyl]-2,2-difluoroethanesulfonyl chloride (121 mg). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.72 (1H, s), 7.62 (1H, m), 7.30 (1H, m), 4.47-4.57 (2H, m), 3.18 (3H, s), 2.49-2.53 (3H, m), 1.80 and 1.82 (3H, s).


(Reaction 203-5)

[1576] N-(4-((E)-1-Fluoro-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 6-1 using appropriate reagents and the starting material obtained above.

MS (ESI) m/z = 519 (M+H)+.


Example 204

3-[3-Methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-imidazolidine-2,4-dione (Compound 943)

[1577](Reaction 204-1)


[1578] Potassium vinyltrifluoroborate (356 mg, 242 μ mol), ethyldiisopropylamine (48 μ L, 279 μ mol) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium dichloromethane adduct (15.1 mg, 18.6 μ mol) were added to a solution of 3-(4-bromo-3-methyl-phenyl)-imidazolidine-2,4-dione (50 mg, 186 μ mol) in n-PrOH (372 μ L) at room temperature in an N_2 atmosphere. The mixture was stirred at 100°C for 1.5 hours, and the reaction solution was then cooled. The reaction solution was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 3-(3-methyl-4-vinyl-phenyl)-imidazolidine-2,4-dione as a yellow brown form (33 mg, 82%).

MS (ESI) m/z = 217 (M+H)+.

(Reaction 204-2)

[1579] 2-[4-(2,5-Dioxo-imidazolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl chloride was synthesized by operations similar to those in Reaction 10-3, Reaction 10-4 and Reaction 10-5 using appropriate reagents and starting material.

MS (ESI) m/z = 317, 319 ($M+H$)⁺.

(Reaction 204-3)

[1580] 3-[3-Methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and the starting material obtained above.





MS (ESI) m/z = 612 ($M+H$)⁺.

[1581] The example compounds shown below were synthesized by operations similar to those in Reaction 204-3 using appropriate reagents and starting materials.

(Compounds 944 to 947)

[1582]


[Table 133]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
944		LCMS-F-1	0.95	626 (M+H)+
945		LCMS-F-1	0.85	592 (M+H)+
946		LCMS-B-1	1.85	592 (M+H)+
947		LCMS-F-1	0.84	584 (M+H)+

Example 205

3-[3-Methyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-imidazolidine-2,4-dione (Compound 948)

[1583](Reaction 205-1)



[1584] 3-[3-Methyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfo-
nyl)-ethyl)-phenyl]-imidazolidine-2,4-dione was synthesized by operations similar to those in Reaction 5-4 using appro-
priate reagents and starting material.

MS (ESI) m/z = 598 (M+H)⁺

[1585] The example compounds shown below were synthesized by operations similar to those in Reaction 205-1 using appropriate reagents and starting materials.

(Compounds 949 to 952)

[1586]


[Table 134]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
949		LCMS-A-1	2.16	558 (M+H) ⁺
950		LCMS-C-1	2.35	530 (M+H) ⁺
951		LCMS-C-1	2.52	544 (M+H) ⁺
952		LCMS-C-1	2.73	572 (M+H) ⁺

Example 206


N-(2-Chloro-4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-5-methyl-phenyl)-acetamide (Compound 953) and N-(2-chloro-4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide (Compound 954)

[1587]

(Reaction 206-1)

[1588] A sulfonyl chloride reagent (a mixture of 2-(4-acetylamino-5-chloro-2-methyl-phenyl)-ethanesulfonyl chloride and 2-(4-acetylamino-3-chloro-2-methyl-phenyl)-ethanesulfonyl chloride) was synthesized by operations similar to those in Reaction 10-5 using appropriate reagents and starting material.

MS (ESI) m/z = 310, 312, 314 (M+H)+.

(Reaction 206-2)

[1589] N-(2-Chloro-4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-5-methyl-phenyl)-acetamide

MS (ESI) m/z = 537 (M+H)+

and

N-(2-chloro-4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide


MS (ESI) m/z = 537 (M+H)+

were obtained by operations similar to those in Reaction 5-4 using appropriate reagents and the starting material obtained above.

Example 207

N-[4-(2-{2-[4-(3,3-Difluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide (Compound 955)

[1590]

(Reaction 207-1)

[1591] N-[4-(2-{2-[4-(3,3-Difluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide (Compound 955) was obtained by operations similar to those in Reaction 18-2 using Compound as a starting material.

MS (ESI) m/z = 553 (M+H)+.

[1592] The example compound shown below was obtained by operations similar to those in Reaction 207-1 using an appropriate starting compound.

(Compound 956)

[1593]

[Table 135]


Raw material Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
945	956		LCMS-F-1	0.87	594 (M+H)+

Example 208

2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-N-phenyl-benzamide (Compound 957)

[1594]

(Reaction 208-1)




[1595] 2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-N-phenyl-benzamide was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.
MS (ESI) m/z = 495 (M+H)+.

Example 209

8-(2-{2-Methyl-4-[4-(1-methyl-piperidin-4-yl)-piperazine-1-carbonyl]-phenyl}-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 958)

[1596]

(Reaction 209-1)




[1597] 8-(2-{2-Methyl-4-[4-(1-methyl-piperidin-4-yl)-piperazine-1-carbonyl]-phenyl}-ethanesulfonyl)-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.
MS (ESI) m/z = 689 (M+H)+.

Example 210

8-((E)-2-[4-((R)-3-Fluoro-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 959)

[1598]

(Reaction 210-1)



[1599] 8-((E)-2-[4-((R)-3-Fluoro-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using

appropriate reagents and starting material.




MS (ESI) $m/z = 623$ (M+H)+.

[1600] The example compounds shown below were synthesized by operations similar to those in Reaction 210-1 using appropriate reagents and starting materials.

(Compounds 960 to 962)

[1601]

[Table 136]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
960		LCMS-B-1	2.15	649 (M+H)+
961		LCMS-C-1	2.72	579 (M+H)+
962		LCMS-C-1	2.62	551 (M+H)+

Example 211

8-[(E)-2-[4-(3-Fluoro-azetidine-1-carbonyl)-2,6-dimethylphenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-tri-
aza-spiro[4.5]dec-1-en-4-one (Compound 963)

[1602]

(Reaction 211-1)



[1603] 8-((E)-2-[4-(3-Fluoro-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 25-15 using appropriate reagents and starting material.


MS (ESI) $m/z = 609$ (M+H)+.

Example 212

8-{2-[4-(4-Hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 964)

[1604]

(Reaction 212-1)



[1605] 8-{2-[4-(4-Hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 122-2 using appropriate reagents and starting material.



MS (ESI) $m/z = 637$ (M+H)+.

[1606] The example compounds shown below were synthesized by operations similar to those in Reaction 212-1 using appropriate reagents and starting materials.

(Compounds 965 to 966)

[1607]


[Table 137]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
965		LCMS-F-1	0.91	609 (M+H)+
966		LCMS-F-1	0.94	651 (M+H)+

Example 213


8-((E)-2-[2,6-Dimethyl-4-(2-oxa-6-aza-spiro[3.3]heptane-6-carbonyl)-phenyl]-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 967)

[1608]

(Reaction 213-1)

[1609] 4-((E)-2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-benzoic acid was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.


MS (ESI) m/z = 554 (M+H)+.

(Reaction 213-2)

15 **[1610]** 8-((E)-2-((2,6-Dimethyl-4-(2-oxa-6-aza-spiro[3.3]heptane-6-carbonyl)-phenyl)-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.
MS (ESI) m/z = 635 (M+H)+.

Example 214

20 8-((E)-2-((2,6-Dimethyl-4-(3-oxo-piperazine-1-carbonyl)-phenyl)-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 968)

[1611](Reaction 214-1)

40 **[1612]** 8-((E)-2-((2,6-Dimethyl-4-(3-oxo-piperazine-1-carbonyl)-phenyl)-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-18 using appropriate reagents and starting material.
MS (ESI) m/z = 636 (M+H)+.

45 **[1613]** The example compounds shown below were synthesized by operations similar to those in Reaction 214-1 using appropriate reagents and starting materials.

(Compounds 969 to 972)

[1614]

Example 215

[1615]


Chemical reaction scheme showing the synthesis of compound **215c** from compound **215a** and compound **215b**.

Compound **215a** (a piperidine ring substituted with a 2,2,2-trifluoroethyl group and a 2,2,2-trifluoroethyl group) reacts with compound **215b** (a 4-methoxyphenyl sulfonate derivative) in the presence of TEA and CH₂Cl₂ to form compound **215c** (a piperidine ring substituted with a 2,2,2-trifluoroethyl group and a 4-methoxyphenyl sulfonate derivative).

[1616] 4-(2-{2-[4-(3,3-Difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-benzoic acid methyl ester was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 552 (M+H)^+$.


(Reaction 215-2)



[1617] Potassium t-butoxide (15.6 mg) was added to a solution of 4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-benzoic acid methyl ester (25.6 mg, 46.4 μmol) in t-butanol (464 μL) and tetrahydrofuran (464 μL), and the mixture was stirred at room temperature for two days. The reaction mixture was diluted with tert-butyl methyl ether and then adjusted to pH 1 with 2 N hydrochloric acid, followed by extraction with ethyl acetate. The organic layer was concentrated under reduced pressure, and the resulting residue was then dried to give 4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-benzoic acid (27.5 mg, 91%).

MS (ESI) $m/z = 538 (M+H)^+$.

(Reaction 215-3)



[1618] 2-[4-(3,3-Difluoro-allyl)-cyclohexyl]-8-{2-[4-(4-fluoro-4-hydroxymethyl-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.


MS (ESI) $m/z = 653 (M+H)^+$.

[1619] The example compound shown below was synthesized by operations similar to those in Reaction 215-3 using appropriate reagents and starting material.

(Compound 974)


[1620]

[Table 139]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
974		LCMS-B-1	1.93	653 (M+H)+

[1621] The amine reagent used for Compound 973 ((4-fluoro-piperidin-4-yl)-methanol hydrochloride) was synthesized by the following method.


(Reaction 215-4)



[1622] 1-Oxa-6-aza-spiro[2.5]octane-6-carboxylic acid tert-butyl ester (144 mg, 679 μmol), triethylamine (1.10 mL, 6.79 mmol) and triethylamine trihydrofluoride (2.85 mL, 20.4 mmol) were mixed in a sealed test tube. This mixture was stirred at 120°C for 6.5 hours. The reaction mixture was cooled, and then quenched with a 2 N aqueous NaOH solution and extracted with ethyl acetate three, times. The organic layers were combined, washed with saturated brine, dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-AcOEt) to give 4-fluoro-4-hydroxymethyl-piperidine-1-carboxylic acid tert-butyl ester (24.9 mg, 16%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.44-1.64 (2H, m), 1.82-1.96 (2H, m), 3.04-3.17 (2H, m), 3.61 (2H, d, $J = 20.0$ Hz), 3.84-3.98 (2H, br-m).

(Reaction 215-5)




[1623] A 4 N solution of hydrochloric acid in 1,4-dioxane (213 μL) was added to a solution of 4-fluoro-4-hydroxymethyl-piperidine-1-carboxylic acid tert-butyl ester (24.9 mg, 0.107 μmol) in MeOH (213 μL) at room temperature, and the mixture was stirred at room temperature for two hours. The reaction solution was concentrated under reduced pressure to give (4-fluoro-piperidin-4-yl)-methanol hydrochloride as a brown form (19.6 mg).

$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ 1.80-2.08 (2H, m), 2.10-2.20 (2H, m), 3.17-3.30 (2H, m), 3.30-3.45 (2H, m), 3.63 (2H, d, $J = 19.6$ Hz).

[1624] The sulfonyl chloride reagent used for Compound 973 (4-(2-chlorosulfonyl-ethyl)-3-methyl-benzoic acid methyl ester) was synthesized by the following method.

(Reaction 215-6)




[1625] 4-(2-Chlorosulfonyl-ethyl)-3-methyl-benzoic acid methyl ester was synthesized by operations similar to those in Reaction 10-2, Reaction 10-3, Reaction 10-4 and Reaction 10-5 using appropriate reagents and starting material. MS (ESI) $m/z = 299$ ($\text{M}+\text{Na}$) $^+$.

Example 216


2-(4-Ethyl-cyclohexyl)-8-(2-{4-[4-(2-fluoro-ethyl)-piperazine-1-carbonyl]-2-methyl-phenyl}-ethanesulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one (Compound 975)

[1626]

(Reaction 216-1)

[1627] 4-{2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid was synthesized by operations similar to those in Reaction 5-4 and Reaction 95-18 using appropriate reagents and starting material.

MS (ESI) m/z = 490 (M+H)+.

(Reaction 216-2)

[1628] 2-(4-Ethyl-cyclohexyl)-8-(2-(4-(2-fluoro-ethyl)-piperazine-1-carbonyl)-2-methyl-phenyl)-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.


MS (ESI) m/z = 604 (M+H)+.

[1629] The example compound shown below was synthesized by operations similar to those in Reaction 216-2 using appropriate reagents and starting material.

(Compound 976)


[1630]

[Table 140]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
976		LCMS-A-1	1.81	614 (M+H)+

[1631] The amine reagent used for Compound 976 (1-oxetan-3-yl-piperazine) was synthesized by the following method.

(Reaction 216-3)



[1632] 1-Oxetan-3-yl-piperazine was synthesized by operations similar to those in Reaction 41-1 and Reaction 18-2 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.27 (4H, br s), 2.89-2.91 (4H, m), 3.42-3.48 (1H, m), 4.58-4.65 (4H, m).

Example 217

8-{2-[2-Methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 977)


[1633]

(Reaction 217-1)



[1634] 3-Methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzoic acid was synthesized by operations similar to those in Reaction 5-4 and Reaction 95-18 using appropriate reagents and , starting material. MS (ESI) m/z = 506 (M+H)+.

(Reaction 217-2)



[1635] 8-{2-[2-Methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material. MS (ESI) m/z = 559 (M+H)+.

[1636] The example compounds shown below were synthesized by operations similar to those in Reaction 217-2 using appropriate reagents and starting materials.

(Compounds 978 to 979)

[1637]

[Table 141]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
978		LCMS-A-1	2.47	545 (M+H)+
979		LCMS-A-1	2.29	575 (M+H)+

Example 218

8-{2-[2,6-Dimethyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 980)


[1638]

(Reaction 218-1)



[1639] 3,5-Dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzoic acid was synthesized by operations similar to those in Reaction 26-1 and Reaction 122-2 using appropriate reagents and starting material. MS (ESI) $m/z = 520$ (M+H)+.

(Reaction 218-2)



[1640] 8-{2-[2,6-Dimethyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material. MS (ESI) $m/z = 573$ (M+H)+.

[1641] The example compounds shown below were synthesized by operations similar to those in Reaction 218-2 using appropriate reagents and starting materials.




(Compounds 981 to 986)

[1642]

[Table 142]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
981		LCMS-F-1	1.03	589 (M+H)+
982		LCMS-F-1	1.09	559 (M+H)+
983		LCMS-F-1	1.13	587 (M+H)+

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
984		LCMS-F-1	1.04	603 (M+H)+
985		LCMS-F-1	1.03	575 (M+H)+
986		LCMS-F-1	1.08	602 (M+H)+

Example 219

N,N-Dimethyl-2-(3-methyl-4-((E)-2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide (Compound 987)

[1643]**(Reaction 219-1)**

[1644] N,N-Dimethyl-2-(3-methyl-4-((E)-2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide was synthesized by operations similar to those in Reaction 10-18 using appropriate reagents and starting material.

MS (ESI) m/z = 515 (M+H)+.

[1645] The example compound shown below was synthesized by operations similar to those in Reaction 219-1 using appropriate reagents and starting material.

(Compound 988)

[1646]


[Table 143]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
988		LCMS-C-1	2.47	517 (M+H)+

Example 220


8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 989)

[1647]

(Reaction 220-1)

[1648] 3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.

MS (ESI) m/z = 570 (M+H)+.

(Reaction 220-2)

[1649] 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 667 (M+H)+.


[1650] The example compounds shown below were synthesized by operations similar to those in Reaction 220-2 using appropriate reagents and starting materials.

[Table 144]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
990		LCMS-D-1	2.28	685 (M+H)+
991		LCMS-D-1	1.60	652 (M+H)+
992		LCMS-D-1	2.57	625 (M+H)+
993		LCMS-D-1	1.88	727 (M+H)+
994		LCMS-D-1	1.87	597 (M+H)+

[1652] The amine reagent used for Compound 993 ((R)-3-(piperidin-4-yloxy)-propane-1,2-diol hydrochloride) was synthesized by the following method.


(Reaction 220-3)



[1653] 4-((S)-2,2-Dimethyl-[1,3]dioxolan-4-ylmethoxy)-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 25-3 using appropriate reagents and starting material.

¹H-NMR (CDCl₃) δ 1.35 (s, 3H), 1.41 (s, 3H), 1.44 (s, 9H), 1.48-1.53 (m, 2H), 1.75-1.87 (m, 2H), 2.95-3.15 (m, 2H), 3.39-3.50 (m, 2H), 3.51-3.58 (m, 2H), 3.67-3.80 (m, 2H), 4.00-4.09 (m, 1H), 4.17-4.32 (m, 1H).

(Reaction 220-4)




[1654] (R)-3-(Piperidin-4-yloxy)-propane-1,2-diol hydrochloride was synthesized by operations similar to those in Reaction 5-3 using appropriate reagents and starting material. This was used in the next reaction without purification.

Example 221

8-{2-[2,6-Dimethyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 995)

[1655]

(Reaction 221-1)



[1656] 8-{2-[2,6-Dimethyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoropropyl)-cy-

clohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 122-2 using appropriate reagents and starting material.


MS (ESI) m/z = 654 (M+H)+.

5 Example 222

3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid hydrazide (Compound 996)

10 **[1657]**

(Reaction 222-1)



[1658] 3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid hydrazide was synthesized by operations similar to those in Reaction 10-14 and Reaction 4-1 using appropriate reagents and starting material.

MS (ESI) m/z = 584 (M+H)+.

[1659] The example compound shown below was synthesized by operations similar to those in Reaction 222-1 using appropriate reagents and starting material.

40 (Compound 997)

[1660]

[Table 145]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
997		LCMS-D-1	2.12	624 (M+H)+

Example 223

N-Methoxy-3,5,N-trimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide (Compound 998)


[1661]

(Reaction 223-1)



[1662] 3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material. MS (ESI) m/z = 488 (M+H)⁺.

(Reaction 223-2)



[1663] N-Methoxy-3,5,N-trimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.

MS (ESI) m/z = 531 (M+H)⁺.

[1664] The example compounds shown below were synthesized by operations similar to those in Reaction 223-2 using appropriate reagents and starting materials.

(Compounds 999 to 1003)

[1665]


[Table 146]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
999		LCMS-D-1	1.99	585 (M+H)+
1000		LCMS-D-1	1.96	603 (M+H)+
1001		LCMS-D-1	2.48	543 (M+H)+
1002		LCMS-D-1	1.67	605 (M+H)+
1003		LCMS-D-1	1.91	515 (M+H)+

Example 224

8-{2-[2,6-Dimethyl-4-(pyrazolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one

[1666]


(Reaction 224-1)

[1667] 8-{2-[2,6-Dimethyl-4-(pyrazolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14, Reaction 122-2 and Reaction 4-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 544$ (M+H)+.

Example 225

8-{(E)-2-[2,6-Dimethyl-4-(pyrazolidine-1-carbonyl)-phenyl]-ethenesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one

[1668](Reaction 225-1)


[1669] 8-[(E)-2-[2,6-Dimethyl-4-(pyrazolidine-1-carbonyl)-phenyl]-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 4-1 using appropriate reagents and starting material.
MS (ESI) $m/z = 542$ (M+H)+.

Example 226

2-Cyclohexyl-8-[(E)-2-[4-(4-hydroxy-4-trifluoromethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1006)


[1670]

(Reaction 226-1)



[1671] 4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-benzoic acid was synthesized by operations similar to those in Reaction 25-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 474$ (M+H)+.

(Reaction 226-2)




[1672] 2-Cyclohexyl-8-[(E)-2-[4-(4-hydroxy-4-trifluoromethyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.
MS (ESI) $m/z = 625$ (M+H)+.

Example 227

2-Cyclohexyl-8-[(E)-2-[2,6-dimethyl-4-(2-oxo-oxazolidine-3-carbonyl)-phenyl]-ethenesulfonyl]-1,3,8-triazaspiro[4.5]dec-1-en-4-one (Compound 1007)

[1673]


(Reaction 227-1)



[1674] DMT-MM (181 mg, 0.50 mmol) was added to a solution of 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethylbenzoic acid (160 mg, 0.33 mmol) in anhydrous ethanol (3.3 ml), and the mixture was stirred at room temperature for 15 hours. The mixture was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography (dichloromethane-methanol) to give 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethylbenzoic acid 4,6-dimethoxy-[1,3,5]triazin-2-yl ester (109 mg, 53%).

MS (ESI) m/z = 613 (M+H)+.

(Reaction 227-2)




[1675] Oxazolidin-2-one (47 mg, 0.53 mmol) was added to a solution of 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethylbenzoic acid 4,6-dimethoxy-[1,3,5]triazin-2-yl ester (109 mg, 0.17 mmol) and triethylamine (0.12 ml, 0.88 mmol) in anhydrous acetonitrile (1 ml), and the mixture was heated with stirring at 80°C for 15 hours. The mixture was cooled and water was then added, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, and then dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 2-cyclohexyl-8-[(E)-2-[2,6-dimethyl-4-(2-oxo-oxazolidine-3-carbonyl)-phenyl]-ethenesulfonyl]-1,3,8-triazaspiro[4.5]dec-1-en-4-one (31 mg, 32%).

MS (ESI) m/z = 543 (M+H)+.


Example 228

2-(4-Butyl-cyclohexyl)-8-((E)-2-{4-[4-(2-hydroxy-ethoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl}-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1008)

[1676]


(Reaction 228-1)

[1677] 4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethylbenzoic acid was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material. MS (ESI) $m/z = 530$ ($\text{M}+\text{H}$)⁺.

(Reaction 228-2)

[1678] 2-(4-Butyl-cyclohexyl)-8-((E)-2-{4-[4-(2-hydroxy-ethoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl}-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material. MS (ESI) $m/z = 657$ ($\text{M}+\text{H}$)⁺.

[1679] The amine reagent used for Compound 1008 (2-(piperidin-4-yloxy)-ethanol hydrochloride) was synthesized by the following method.

(Reaction 228-3)

[1680] 2-(Piperidin-4-yloxy)-ethanol hydrochloride was synthesized by operations similar to those in Reaction 20-2 and Reaction 5-3 using appropriate reagents and starting material.

¹H-NMR (300 MHz, DMSO-d₆) δ 1.50-1.62 (m, 2H), 1.80-1.89 (m, 2H), 2.87-2.93 (m, 2H), 3.10-3.16 (m, 2H), 3.69-3.76 (m, 1H), 5.00 (s, 1H), 8.74-8.90 (m, 2H).

[1681] The example compound shown below was synthesized by operations similar to those in Reaction 228-2 using appropriate reagents and starting material.


(Compound 1009)

[1682]

[Table 147]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1009		LCMS-D-1	2.42	701 (M+H) ⁺


[1683] The amine reagent used for Compound 1009 (2-[2-(piperidin-4-yloxy)-ethoxy]-ethanol hydrochloride) was synthesized by the following method.

(Reaction 228-4)

[1684] 4-{2-[2-(tert-Butyl-dimethyl-silyloxy)-ethoxy]-ethoxy}-piperidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 20-2 using appropriate reagents and starting material.

¹H-NMR (300 MHz, CDCl₃) δ 0.07 (s, 6H), 0.89 (s, 9H), 1.43-1.59 (m, 11H), 1.79-1.88 (m, 2H), 2.99-3.13 (m, 2H), 3.44-3.52 (m, 1H), 3.54-3.59 (m, 2H), 3.60-3.68 (s, 4H), 3.72-3.84 (m, 4H).

(Reaction 228-5)




[1685] 2-[2-(Piperidin-4-yloxy)-ethoxy]-ethanol hydrochloride was synthesized by operations similar to those in Reaction 5-3 using appropriate reagents and starting material. This was used in the next reaction without purification.

Example 229

8-((E)-2-{4-[4-((R)-2,3-Dihydroxy-propoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl}-ethenesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1010)

[1686]


(Reaction 229-1)



[1687] 3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(9,9,9-trifluoro-nonyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzoic acid was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) m/z = 572 (M+H)+.

(Reaction 229-2)




[1688] 8-((E)-2-{4-[4-((R)-2,3-Dihydroxy-propoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl}-ethenesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material.
MS (ESI) m/z = 729 (M+H)+.

Example 230


2-Amino-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide (Compound 1011)

[1689]

(Reaction 230-1)




[1690] 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 26-1 using appropriate reagents and starting material.
MS (ESI) m/z = 537 (M+H)+.

(Reaction 230-2)

[1691] 2-Chloro-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide was synthesized by operations similar to those in Reaction 2-3 using appropriate reagents and starting material.

MS (ESI) $m/z = 613 (M+H)^+$.


(Reaction 230-3)

[1692] Ammonia (6 N solution in ethanol, 0.3 ml) was added to a solution of 2-chloro-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide (38 mg, 0.06 mmol) in anhydrous ethanol (0.5 ml), and the mixture was stirred at 50 to 60°C for five hours. The mixed solution was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography (dichloromethane) to give 2-amino-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide (11 mg, 31%).


MS (ESI) $m/z = 594 (M+H)^+$.

Example 231


(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methyl-carbamic acid 2-hydroxy-ethyl ester (Compound 1012)

[1693](Reaction 231-1)

[1694] Phosgene (20% solution in toluene, 35 μL , 67 μmol) was added to a mixed solution of 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (30 mg, 56 μmol) and triethylamine (15 μL , 84 μmol) in dichloromethane (1.5 ml) and dimethylformamide (0.5 ml) at 0°C. The mixture was stirred at room temperature for three hours, and then quenched with water and extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give N-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-methyl-chloroformamide (28 mg, 85%). MS (ESI) m/z = 599 (M+H)+.

(Reaction 231-2)

[1695] Sodium hydride (60% oil suspension, 5.6 mg, 0.14 mmol) was added to a solution of N-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-methyl-chloroformamide (30 mg, 50.1 μmol) in tetrahydrofuran (1.0 ml) at 0°C, and the mixture was stirred for 15 minutes. 2-(tert-Butyl-dimethyl-silyloxy)-ethanol (30 μL , 0.14 mmol) was then added and the mixture was stirred at 40°C for two hours. The mixture was cooled, and then quenched with water and extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give (3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-methyl-carbamic acid 2-(tert-butyl-dimethyl-silyloxy)-ethyl ester (29 mg, 78%). MS (ESI) m/z = 739 (M+H)+.


(Reaction 231-3)

[1696] (3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-methyl-carbamic acid 2-hydroxy-ethyl ester was synthesized by operations similar to those in Reaction 39-2 using appropriate reagents and starting material. MS (ESI) m/z = 625 (M+H)+.

Example 232

1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1,3,3-trimethyl-urea (Compound 1013)

[1697]

(Reaction 232-1)

[1698] 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3,3-trimethyl-urea was synthesized by operations similar to those in Reaction 231-2 using appropriate reagents and starting material.

MS (ESI) m/z = 608 (M+H)+.

[1699] The example compounds shown below were synthesized by operations similar to those in Reaction 232-1 using appropriate reagents and starting materials.

(Compounds 1014 to 1015)

[1700]

[Table 148]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1014		LCMS-D-1	3.02	594 (M+H)+
1015		LCMS-D-1	2.92	580 (M+H)+

Example 233

8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(3-trifluoromethylsulfonyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1016)

[1701]


(Reaction 233-1)



[1702] {4-[(E)-2-(1,4-Dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 25-1 and Reaction 26-1 using appropriate reagents and starting material.

MS (ESI) $m/z = 467$ (M+H)+.


(Reaction 233-2)



[1703] Trifluoroacetic acid (5.3 ml, 71.79 mmol) was added to a solution of {4-[(E)-2-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester (670 mg, 1.43 mmol) in acetone-water (8.0 ml-8.0 ml) at room temperature, and the mixture was heated with stirring at 50°C for 18 hours. The mixed reaction solution was cooled and then concentrated under reduced pressure. The residue was neutralized by adding a saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over Na_2SO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate-hexane) to give 1-[(E)-2-(N,2,6-trimethylaniline)-ethanesulfonyl]-piperidin-4-one (364 mg, 78%).


MS (ESI) $m/z = 323$ (M+H)+.

(Reaction 233-3)




[1704] Ammonium acetate (686 mg, 8.91 mmol) and potassium cyanide (541 mg, 8.31 mmol) were added to a solution of 1-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-piperidin-4-one (1.91 g, 5.94 mmol) in MeOH (20 ml), and the mixture was heated with stirring at 60°C for three hours. The mixed reaction solution was cooled and a saturated aqueous NaHCO₃ solution was then added, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over Na₂SO₄ and concentrated under reduced pressure. The resulting residue was triturated with hexane:CH₂Cl₂ = 7:3 to give 4-amino-1-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-piperidine-4-carbonitrile (1.86 g, 89%).
MS (ESI) m/z = 349 (M+H)+.

(Reaction 233-4)




[1705] DMSO (0.9 ml, 12.8 mmol), a 1 N aqueous NaOH solution (1.06 ml, 1.06 mmol) and 30% aqueous hydrogen peroxide (0.72 ml, 6.40 mmol) were sequentially added to a solution of 4-amino-1-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-piperidine-4-carbonitrile (1.85 g, 5.33 mmol) in MeOH (30 ml) at 0°C, and the mixture was stirred at room temperature for 2.5 hours. A saturated Na₂S₂O₃ solution was added to the reaction mixture, and the precipitated solid was obtained by suction filtration. The resulting solid was washed with water, dissolved in a CH₂Cl₂-MeOH (3:2) solution, dried over Na₂SO₄ and concentrated under reduced pressure. The resulting residue was triturated with hexane:CH₂Cl₂ = 4:1 to give 4-amino-1-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-piperidine-4-carboxylic amide (1.36 g, 70%).
MS (ESI) m/z = 367 (M+H)+.

(Reaction 233-5)

[1706] 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 189-5 using appropriate reagents and starting material.
MS (ESI) $m/z = 553 (M+H)^+$.

Example 234

1-(4-[(E)-2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1017)

[1707](Reaction 234-1)

[1708] 1-(4-[(E)-2-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 189-5

and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 582$ (M+H)+.

[1709] The example compounds shown below were synthesized by operations similar to those in Reaction 234-1 using appropriate reagents and starting materials.

(Compounds 1018 to 1021)


[1710]

[Table 149]

Compound	Structure	LCMS condition	Retention time(min)	MS (m/z)
1018		LCMS-F-1	0.95	612 (M+H)+
1019		LCMS-F-1	0.99	622 (M+H)+
1020		LCMS-F-1	0.97	622 (M+H)+
1021		LCMS-F-1	1.04	566 (M+H)+

[1711] The carboxylic acid reagent used in the synthesis of Compound 1019 (3-(4,4,4-trifluorobutoxy)-benzoic acid) was synthesized by the following method.

(Reaction 234-2)




[1712] 3-(4,4,4-Trifluorobutoxy)-benzoic acid was synthesized by operations similar to those in Reaction 26-4 (using Cs_2CO_3 as a base) and Reaction 189-5 using appropriate reagents and starting material.

MS (ESI) m/z = 247 (M-H)-.

[1713] The carboxylic acid reagent used in the synthesis of Compound 1020 (4-(4,4,4-trifluorobutoxy)-benzoic acid) was synthesized by the following method.

(Reaction 234-3)




[1714] 4-(4,4,4-Trifluorobutoxy)-benzoic acid was synthesized by operations similar to those in Reaction 26-4 (using Cs_2CO_3 as a base) and Reaction 189-5 using appropriate reagents and starting material.

MS (ESI) m/z = 249 (M+H)+.

Example 235

1-(3,5-Dimethyl-4-((E)-2-[2-(7-methylsulfanyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (Compound 1022)


[1715]

(Reaction 235-1)

[1716] 1-(3,5-Dimethyl-4-((E)-2-[2-(7-methylsulfanyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 10-12 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) *m/z* = 564 (M+H)+.

[1717] The carboxylic acid reagent used in the synthesis of Compound 1022 (8-(methylthio)octanoic acid) was synthesized by the following method.

(Reaction 235-2)


[1718] Sodium thiomethoxide (942 mg, 13.44 mmol) was added to a solution of 8-bromooctanoic acid (500 mg, 2.24 mmol) in methanol (5.6 mL), and the mixture was heated under reflux overnight. The reaction mixture was concentrated under reduced pressure, adjusted to pH 1 by adding 1 N hydrochloric acid and then extracted with ethyl acetate. The organic layer was washed with saturated brine, and then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane:methanol = 20:1) to give 8-(methylthio)octanoic acid as a colorless oily substance (426.5 mg, 100%).

¹H-NMR (300 MHz, CDCl₃) δ 1.31-1.41 (m, 6H), 1.54-1.66 (m, 4H), 2.09 (s, 3H), 2.35 (t, 2H, *J* = 7.2 Hz), 2.48 (t, 2H, *J* = 7.2 Hz).


Example 236

1-[3,5-Dimethyl-4-((E)-2-[2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea (Compound 1023)


[1719]

(Reaction 236-1)

[1720] 1-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-4-[9-(3-methyl-oxetan-3-yl)-nonanoylamino]-piperidine-4-carboxylic amide was synthesized by operations similar to those in Reaction 10-14 using appropriate reagents and starting material. This was used in the next reaction without purification.

(Reaction 236-2)


[1721] LiOH.H₂O (16.6 mg, 0.396 mmol) was added to a solution of 1-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-4-[9-(3-methyl-oxetan-3-yl)-nonanoylamino]-piperidine-4-carboxylic amide (96 mg, 0.098 mmol) in ethanol (1.0 mL), and the mixture was stirred at 50°C for two hours. A 50% saturated aqueous ammonium chloride solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate) to give 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-[8-(3-methyl-oxetan-3-yl)-octyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (54.8 mg, 100%). ¹H-NMR (400 MHz, CD₃OD) δ 1.26 (3H, s), 1.34 (10H, m), 1.63 (6H, m), 1.95 (2H, m), 2.37 (6H, s), 2.44 (2H, m), 2.77 (3H, s), 3.15 (2H, m), 3.63 (2H, m), 4.31 (2H, d, J = 5.6 Hz), 4.40 (2H, d, J = 5.6 Hz), 6.34 (1H, d, J = 15.6 Hz), 6.34 (2H, s), 7.63 (1H, d, J = 15.6 Hz).

(Reaction 236-3)

[1722] 1-[3,5-Dimethyl-4-((E)-2-{2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 89-2 (using KOCN) using appropriate reagents and starting material.


MS (ESI) m/z = 602 (M+H)+.

[1723] The carboxylic acid reagent used in the synthesis of Compound 1023 (9-(3-methyl-oxetan-3-yl)-nonanoic acid) was synthesized by the following method.

(Reaction 236-4)

[1724] (9-Bromo-nonyloxymethyl)-benzene was synthesized by operations similar to those in Reaction 20-2 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.25-1.45 (10H, m), 1.61 (2H, m), 1.85 (2H, m), 3.40 (2H, t, J = 6.8 Hz), 3.46 (2H, t, J = 6.8 Hz), 4.50 (2H, s), 7.27-7.35 (5H, m).


(Reaction 236-5)

[1725] Methyl-malonic acid diethyl ester (0.850 ml, 4.99 mmol) was added to a suspension of sodium hydride (55% oily suspension, 139.5 mg, 3.197 mmol) in THF (0.8 ml) over seven minutes under ice-cooling, and the mixture was stirred until foaming was terminated at room temperature (for about 25 minutes). A solution of (9-bromo-nonyloxymethyl)-benzene (593 mg, 1.89 mmol) in THF (0.12 ml) was added to the reaction solution at room temperature over 15 minutes, and the mixture was then stirred at 90°C for five hours. The reaction mixture was diluted with ether and water

was then added, followed by extraction with ether. The organic layer was dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 30/1 $\rightarrow$ 20/1) to give 2-(9-benzyloxy-nonyl)-2-methyl-malonic acid diethyl ester (735 mg, 96%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.24 (6H, t, $J = 6.8$ Hz), 1.27 (12H, m), 1.39 (3H, s), 1.60 (2H, m), 1.83 (2H, m), 3.46 (2H, t, $J = 6.8$ Hz), 4.20 (4H, m), 4.50 (2H, s), 7.27-7.34 (5H, m).


(Reaction 236-6)



[1726] 2-(9-Benzyloxy-nonyl)-2-methyl-propane-1,3-diol was synthesized by operations similar to those in Reaction 95-28 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.82 (3H, s), 1.28 (14H, m), 1.61 (2H, m), 2.14 (2H, t, $J = 4.2$ Hz), 3.46 (2H, t, $J = 6.6$ Hz), 3.54 (4H, m), 4.50 (2H, s), 7.27-7.35 (5H, m).


(Reaction 236-7)



[1727] n -Butyllithium (2.6 M solution in hexane, 0.295 ml, 0.767 mmol) was added to a solution of 2-(9-benzyloxy-nonyl)-2-methyl-propane-1,3-diol (221 mg, 0.686 mmol) in THF (5.1 ml) at 0°C over three minutes, and the mixture was then stirred at the same temperature for 30 minutes. A solution of TsCl (138 mg, 0.723 mmol) in THF (0.91 ml) was added to the reaction solution at 0°C over eight minutes, and the mixture was then stirred at the same temperature for one hour. n -Butyllithium (2.6 M solution in hexane, 0.295 ml, 0.767 mmol) was added dropwise to the reaction mixture at 0°C , and the mixture was then stirred at 60°C for six hours. The reaction mixture was diluted with ether and water was then added, followed by extraction with ether. The organic layer was dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 15/1) to give 3-(9-benzyloxy-nonyl)-3-methyl-oxetane (188 mg, 90%).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.27 (3H, s), 1.29 (12H, m), 1.61 (4H, m), 3.47 (2H, t, $J = 6.6$ Hz), 4.32 (2H, d, $J = 5.4$ Hz), 4.41 (2H, d, $J = 5.4$ Hz), 4.50 (2H, s), 7.27-7.35 (5H, m).

(Reaction 236-8)



[1728] 9-(3-Methyl-oxetan-3-yl)-nonan-1-ol was synthesized by operations similar to those in Reaction 122-2 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.27 (3H, s), 1.29 (12H, m), 1.59 (4H, m), 3.64 (2H, t, J = 6.6 Hz), 4.33 (2H, d, J = 5.8 Hz), 4.41 (2H, d, J = 5.8 Hz).

(Reaction 236-9)

[1729] TEMPO (3.4 mg, 0.022 mmol) and iodobenzene diacetate (69.7 mg, 0.216 mmol) were added to a solution of 9-(3-methyl-oxetan-3-yl)-nonan-1-ol (21 mg, 0.098 mmol) in acetonitrile (0.2 ml)-water (0.1 ml) at room temperature, and the mixture was stirred at the same temperature for two hours. Water (0.1 ml) was then added to the reaction mixture at room temperature, and the mixture was stirred at the same temperature for one hour. A 10% aqueous citric acid solution (0.45 ml) was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water, a 10% aqueous sodium thiosulfate solution and saturated brine, dried over MgSO₄ and concentrated under reduced pressure to give 9-(3-methyl-oxetan-3-yl)-nonanoic acid (22 mg, 100%).

¹H-NMR (400 MHz, CDCl₃) δ 1.27 (3H, s), 1.31 (10H, m), 1.63 (4H, m), 2.35 (2H, t, J = 7.6 Hz), 4.33 (2H, d, J = 5.4 Hz), 4.42 (2H, d, J = 5.4 Hz).

[1730] The example compounds shown below were synthesized by operations similar to those in Reaction 236-1, Reaction 236-2 and Reaction 236-3 using appropriate reagents and starting materials.

(Compounds 1024 to 1027)


[1731]

[Table 150]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1024		LCMS-C-2	1.97	542 (M+H) ⁺
1025		LCMS-F-1	1.01	542 (M+H) ⁺
1026		LCMS-C-2	2.13	630 (M+H) ⁺
1027		LCMS-C-2	2.22	540 (M-H) ⁻

[1732] The carboxylic acid reagent used in the synthesis of Compound 1024 (dec-5-ynoic acid) was synthesized by the following method.


(Reaction 236-10)



[1733] 1-Hexyne (43 μl , 0.38 mmol) and ethyl 4-bromobutyrate (40 μl , 0.28 mmol) were added to a suspension of 1,3-bis(1-adamantyl)imidazolium chloride (5.2 mg, 0.014 mmol), copper iodide (4.1 mg, 0.022 mmol), allylpalladium(II) chloride dimer (2.6 mg, 0.071 mmol) and cesium carbonate (127 mg, 0.390 mmol) in anhydrous ether (0.37 ml)-anhydrous DMF (0.185 ml) at room temperature in a nitrogen stream, and the mixture was stirred at 45°C for 17 hours. The reaction mixture was extracted with pentane. The insoluble matter was removed by filtration, and the filtrate was then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ether = 50/1) to give dec-5-ynoic acid ethyl ester (40 mg, 73%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.90 (3H, t, $J = 7.2$ Hz), 1.26 (3H, t, $J = 7.2$ Hz), 1.42 (4H, m), 1.80 (2H, m), 2.14 (2H, m), 2.22 (2H, m), 2.42 (2H, t, $J = 7.6$ Hz), 4.13 (2H, q, $J = 7.2$ Hz).


(Reaction 236-11)



[1734] Dec-5-ynoic acid was synthesized by operations similar to those in Reaction 189-5 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.91 (3H, t, $J = 7.2$ Hz), 1.43 (4H, m), 1.82 (2H, m), 2.15 (2H, m), 2.25 (2H, m), 2.50 (2H, t, $J = 7.2$ Hz).

[1735] The carboxylic acid reagent used in the synthesis of Compound 1025 (dec-4-ynoic acid) was synthesized by the following method.


(Reaction 236-12)

[1736] Dec-4-ynoic acid was synthesized by operations similar to those in Example 236-10, Reaction 39-2 and Reaction 236-9 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 0.89 (3H, t, J = 6.8 Hz), 1.32 (4H, m), 1.47 (2H, m), 2.13 (2H, m), 2.49 (2H, m), 2.57 (2H, m).

[1737] The carboxylic acid reagent used in the synthesis of Compound 1026 (11-(3-methyl-oxetan-3-yl)-undecanoic acid) was synthesized by the following method.

(Reaction 236-13)




[1738] 11-(3-Methyl-oxetan-3-yl)-undecanoic acid was synthesized by operations similar to those in Reaction 20-2, Reaction 236-5, Reaction 95-28, Reaction 236-7, Reaction 122-2 and Reaction 236-9 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.27 (3H, s), 1.28 (14H, m), 1.62 (4H, m), 2.35 (2H, t, $J = 7.6$ Hz), 4.34 (2H, d, $J = 5.6$ Hz), 4.43 (2H, d, $J = 5.6$ Hz).

[1739] The carboxylic acid reagent used in the synthesis of Compound 1027 (dec-2-ynoic acid) was synthesized by the following method.

(Reaction 236-14)



[1740] Dec-2-ynoic acid was synthesized by operations similar to those in Reaction 236-9 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.88 (3H, t, $J = 7.2$ Hz), 1.28 (6H, m), 1.40 (2H, m), 1.59 (2H, m), 2.35 (2H, t, $J = 7.2$ Hz).

Example 237

1-(4-{2-[2-(4-Fluoro-3-trifluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1028)

[1741]

(Reaction 237-1)



[1742] 1-(4-{2-[2-(4-Fluoro-3-trifluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 189-5, Reaction 89-2 and Reaction 184-1 using appropriate reagents and starting material.



MS (ESI) m/z = 600 (M+H)⁺.

[1743] The example compounds shown below were synthesized by operations similar to those in Reaction 237-1 using appropriate reagents and starting materials.

(Compounds 1029 to 1030)


[1744]

[Table 151]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1029		LCMS-F-1	0.95	578 (M+H)+
1030		LCMS-F-1	0.99	554 (M+H)+

Example 238

1-[3,5-Dimethyl-4-(2-[2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea (Compound 1031)

[1745]**(Reaction 238-1)**

[1746] 1-[3,5-Dimethyl-4-(2-[2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 184-1 using appropriate reagents and starting material.


MS (ESI) m/z = 604 (M+H)+.

[1747] The example compound shown below was synthesized by operations similar to those in Reaction 238-1 using appropriate reagents and starting material.

(Compound 1032)

[1748]

[Table 152]


Compound	Structure	LCMS condition	Retention time (min)	MS(m/z)
1032		LCMS-C-2	2.13	630 (M-H)-

Example 239

2-(8-{2-[4-(tert-Butoxycarbonyl-methyl-amino)-2,6-dimethyl-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-pyrrolidine-1-carboxylic acid tert-butyl ester (Compound 1033)

[1749]

(Reaction 239-1)




[1750] {4-[2-(4-Amino-4-carbamoyl-piperidine-1-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl

ester was synthesized by operations similar to those in Reaction 184-1, Reaction 233-2, Reaction 19-2, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.

MS (ESI) m/z = 469 (M+H)+.

(Reaction 239-2)



[1751] 2-(8-{2-[4-(tert-Butoxycarbonyl-methyl-amino)-2,6-dimethyl-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-pyrrolidine-1-carboxylic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-14 and Reaction 189-5 using appropriate reagents and starting material.

MS (ESI) m/z = 648 (M+H)+.

[1752] The example compound shown below was synthesized by operations similar to those in Reaction 239-2 using appropriate reagents and starting material.

(Compound 1034)

[1753]


[Table 153]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1034		LCMS-F-1	1.01	541 (M-H)-

Example 240

8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1035)

[1754]

(Reaction 240-1)

[1755] 8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14, Reaction 189-5 and Reaction 4-1 using appropriate reagents and starting material.

MS (ESI) m/z = 595 (M+H)+.

[1756] The example compound shown below was synthesized by operations similar to those in Reaction 240-1 using appropriate reagents and starting material.

(Compound 1036)

[1757]


[Table 154]

Compound	Structure	LCMS condition	Retention time (min)	MS(m/z)
1036		LCMS-C-1	2.93	555 (M+H)+

Example 241


8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(4-isopropylidene-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1037)

[1758]

(Reaction 241-1)

[1759] (4-{2-[2-(4-isopropylidene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethylphenyl)-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 10-14 and Reaction 189-5 using appropriate reagents and starting material.


MS (ESI) m/z = 601 (M+H)+.

(Reaction 241-2)

[1760] About 40 4AMS beads were added to a solution of (4-{2-[2-(4-isopropylidene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethylphenyl)-methyl-carbamic acid tert-butyl ester (85.7 mg, 143 μmol) in dichloromethane (1.4 ml), and the mixture was stirred at room temperature for 10 minutes. Thereafter, $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (90.2 μl , 715 μmol) was added to the reaction mixture at 0°C, and the mixture was stirred at room temperature for three hours. The reaction mixture was quenched by adding triethylamine and diluted with ethyl acetate. The organic layer was then washed with a saturated aqueous sodium bicarbonate solution and water, and then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane/methanol = 100/0 $\rightarrow$ 92/8) to give 8-[2-(2,6-dimethyl-4-methylaminophenyl)-ethanesulfonyl]-2-(4-isopropylidene-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (66.0 mg, 92%).

MS (ESI) m/z = 501 (M+H)+.


[1761] The carboxylic acid reagent used in the synthesis of Compound 1037 (4-isopropylidene-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 241-3)

[1762] 4-Isopropylidene-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 191-14 and Reaction 189-5 using appropriate reagents and starting material.
MS (ESI) $m/z = 169$ (M+H)+.

Example 242

1-[3,5-Dimethyl-4-(2-{4-oxo-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-1-methyl-urea (Compound 1038)

[1763](Reaction 242-1)


[1764] 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 189-5, Reaction 4-1 and Reaction 89-2 using appropriate reagents and starting material.
MS (ESI) $m/z = 614$ (M+H)+.

Example 243

1-(3,5-Dimethyl-4-{2-[4-oxo-2-(5,6,7,8-tetrahydro-naphthalen-2-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 1039)

[1765]

(Reaction 243-1)



[1766] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(5,6,7,8-tetrahydro-naphthalen-2-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 236-2, Reaction 4-1 and Reaction 89-2 using appropriate reagents and starting material.




MS (ESI) m/z = 552 (M+H)⁺.

[1767] The example compounds shown below were synthesized by operations similar to those in Reaction 243-1 using appropriate reagents and starting materials.

(Compounds 1040 to 1042)


[1768]

[Table 155]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1040		LCMS-A-1	2.06	532 (M+H)+
1041		LCMS-A-1	2.00	520 (M+H)+
1042		LCMS-A-1	2.26	636 (M+H)+

[1769] The carboxylic acid reagent used in the synthesis of Compound 1042 (4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 243-2)




[1770] 4-(2,2,3,3,3-Pentafluoro-propyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 193-4, Reaction 193-5, Reaction 193-6 and Reaction 95-18 using appropriate reagents and starting material. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.64 (0.6H, qui, $J = 4.9$ Hz), 2.29 (0.4H, tt, $J = 12.2, 3.4$ Hz), 2.09-1.06 (11H, m). (cis/trans = ca 6 : 4)

Example 244

1-(3,5-Dimethyl-4-{2-[4-oxo-2-(6,6,7,7,7-pentafluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 1043)


[1771]

(Reaction 244-1)

[1772] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(6,6,7,7,7-pentafluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-1, Reaction 10-12, Reaction 4-1 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.

MS (ESI) $m/z = 610$ (M+H)+.

[1773] The carboxylic acid reagent used in the synthesis of Compound 1043 (7,7,8,8,8-pentafluoro-octanoic acid) was synthesized by the following method.



[1774] 7,7,8,8,8-Pentafluoro-octanoic acid was synthesized by operations similar to those in Reaction 18-2 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.42-1.48 (2H, m), 1.57-1.73 (4H, m), 2.03 (2H, tt, $J = 6.8, 18.2$ Hz), 2.39 (2H, t, $J = 7.4$ Hz).

Example 245

N-(3,5-Dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-N-methyl-acetamide (Compound 1044) and 1-(3,5-dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 1045)

[1775]

(Reaction 245-1)



[1776] N-(3,5-Dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-N-methyl-acetamide

MS (ESI) $m/z = 573$ (M+H)⁺

and


1-(3,5-dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea

MS (ESI) $m/z = 574$ (M+H)⁺

were synthesized by operations similar to those in Reaction 10-1, Reaction 10-12, Reaction 4-1 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.

[1777] The carboxylic acid reagent used in the synthesis of Compound 1044 and Compound 1045 (8,8,8-trifluoro-octanoic acid) was synthesized by the following method.

(Reaction 245-2)



[1778] 8,8,8-Trifluoro-octanoic acid was synthesized by operations similar to those in Reaction 109-1, Reaction 101-1, Reaction 18-2 and Reaction 109-1 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.32-1.46 (4H, br-m), 1.52-1.68 (2H, m), 1.66-1.72 (2H, m), 2.00-2.14 (2H, m), 2.38 (2H, t, $J = 7.2$ Hz).

[1779] The example compounds shown below were synthesized by operations similar to those in Reaction 245-1 using appropriate reagents and starting materials.

(Compounds 1046 to Compound 1047)

[1780]

[Table 156]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1046		LCMS-B-1	2.03	602 (M+H)+
1047		LCMS-B-1	2.16	601 (M+H)+

Example 246

1-[3,5-Dimethyl-4-(2-{4-oxo-2-[1-(4,4,4-trifluoro-butyl)-cyclopropyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-1-methyl-urea (Compound 1048)

[1781]

(Reaction 246-1)




[1782] 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[1-(4,4,4-trifluorobutyl)-cyclopropyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 10-12, Reaction 4-1 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material. MS (ESI) $m/z = 572$ ($M+H$) $^+$.

Example 247

1-[3,5-Dimethyl-4-(2-{4-oxo-2-(3-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-1-methyl-urea (Compound 1050)


[1783]

(Reaction 247-1)

[1784] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 189-5, Reaction 5-3 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) m/z = 572 (M+H)+.

[1785] The carboxylic acid reagent used in the synthesis of Compound 1050 (3-trifluoromethyl-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 247-2)

[1786] 3-Trifluoromethyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 193-3 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.22-2.40 (9.57H, m), 2.88 (0.43H, m) (cis : trans = 1.3 : 1)

[1787] The example compounds shown below were synthesized by operations similar to those in Reaction 247-1 using appropriate reagents and starting materials.


(Compounds 1051 to 1058)

[1788]


[Table 157]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1051		LCMS-F-1	0.93	518 (M+H)+
1052		LCMS-F-1	0.87	516 (M+H)+
1053		LCMS-F-1	0.99	544 (M+H)+
1054		LCMS-F-1	0.93	530 (M+H)+
1055		LCMS-F-1	0.89	504 (M+H)+
1056		LCMS-F-1	0.91	516 (M+H)+
1057		LCMS-F-1	1.08	588 (M+H)+

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1058		LCMS-F-1	0.94	530 (M+H)+


[1789] The carboxylic acid reagent used in the synthesis of Compound 1051 (cis-3,4-dimethyl-cyclopentanecarboxylic acid) was synthesized by the following method.

(Reaction 247-3)

[1790] cis-3,4-Dimethyl-cyclopentanecarboxylic acid was synthesized by operations similar to those in Reaction 193-12 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 0.85 (3H, d, J = 7.0 Hz), 0.89 (3H, d, J = 7.0 Hz), 1.58-1.68 (2H, m), 2.00-2.15 (4H, m), 2.78 (0.4H, dd, J = 17.2, 8.8 Hz), 2.92-3.00 (0.6H, m) (cis : trans = 6 : 4).


[1791] The carboxylic acid reagent used in the synthesis of Compound 1052 (dicyclopropyl-acetic acid) was synthesized by the following method.

(Reaction 247-4)

[1792] Dicyclopropyl-acetic acid was synthesized by operations similar to those in Reaction 193-12 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.24-0.33 (4H, m), 0.48-0.52 (2H, m), 0.58-0.62 (2H, m), 1.05-1.10 (3H, m).

[1793] The carboxylic acid reagent used in the synthesis of Compound 1053 (bicyclo[3.3.1]nonane-9-carboxylic acid) was synthesized by the following method.


(Reaction 247-5)

[1794] Bicyclo[3.3.1]nonane-9-carboxylic acid was synthesized by operations similar to those in Reaction 193-12 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.48-1.63 (4H, m), 1.70-1.80 (2H, m), 1.83-1.96 (6H, m), 2.33 (2H, br), 2.46 (1H, br).

[1795] The carboxylic acid reagent used in the synthesis of Compound 1057 (2-hexyl-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 247-6)




25 **[1796]** 2-Hexyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 101-1, Reaction 95-18 and Reaction 193-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.88 (3H, t, $J = 7.2$ Hz), 1.20-1.55 (13H, m), 1.60-1.90 (6H, m), 2.56-2.59 (1H, m).

[1797] The carboxylic acid reagent used in the synthesis of Compound 1058 (spiro[2.5]octane-6-carboxylic acid) was synthesized by the following method.


30 (Reaction 247-7)



40 **[1798]** 4-Methylene-cyclohexanecarboxylic acid benzyl ester was synthesized by operations similar to those in Reaction 193-9 using appropriate reagents and starting material.

MS (ESI) $m/z = 231$ ($\text{M}+\text{H}$) $^+$.

45 (Reaction 247-8)




55 **[1799]** Et_2Zn (1.08 M solution in hexane, 5.23 ml, 5.65 mmol) was added to a solution of 4-methylene-cyclohexanecarboxylic acid benzyl ester (86.3 mg, 375 μmol) in toluene (1.4 ml), and the mixture was stirred at room temperature for 30 minutes. CH_2I_2 (500 μl , 6.22 mmol) was added to the reaction mixture at 0°C , and the mixture was stirred at 60°C for 28 hours. Thereafter, Et_2Zn (1.08 M solution in hexane, 2.60 ml, 2.22 mmol) and CH_2I_2 (260 μl , 3.23 mmol) were added to the reaction mixture, and the mixture was stirred at 60°C for four days. Further, Et_2Zn (1.08 M solution in hexane, 2.60 ml, 2.22 mmol) and CH_2I_2 (500 μl , 6.22 mmol) were added to the reaction mixture, and the mixture was

stirred at 60°C for one day. The reaction mixture was quenched by adding a 1% aqueous HCl solution and diluted with ethyl acetate and Et₂O. The organic layer was then washed with a 1% aqueous HCl solution, a saturated aqueous sodium bicarbonate solution and saturated brine, and then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 100/0 → 65/35) to give

spiro[2.5]octane-6-carboxylic acid benzyl ester (70.6 mg, 77%).
¹H-NMR (400 MHz, CDCl₃) δ 0.19-0.22 (2H, m), 0.26-0.30 (2H, m), 0.94-1.00 (2H, m), 1.60-1.72 (4H, m), 1.90-1.95 (2H, m), 2.37-2.42 (1H, m), 5.12 (2H, s), 7.30-7.38 (5H, m).

(Reaction 247-9)



[1800] Spiro[2.5]octane-6-carboxylic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material.


¹H-NMR (400 MHz, CDCl₃) δ 0.19-0.24 (2H, m), 0.28-0.30 (2H, m), 0.97-1.03 (2H, m), 1.60-1.72 (4H, m), 1.92-1.95 (2H, m), 2.35-2.42 (1H, m).

Example 248

1-(4-{2-[2-(4-Difluoromethylene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1059)

[1801]

(Reaction 248-1)




[1802] 1-(4-{2-[2-(4-Difluoromethylene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 10-12, Reaction 241-2 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) m/z = 552 (M+H)+.

[1803] The carboxylic acid reagent used in the synthesis of Compound 1059 (4-difluoromethylene-cyclohexanecarboxylic acid) was synthesized by the following method.


(Reaction 248-2)



[1804] HMPT (7.40 ml, 39.5 mmol) was added to a solution of 4-oxo-cyclohexanecarboxylic acid benzyl ester (1.50 g, 6.59 mmol) and CF_2Br_2 (1.8 ml, 19.8 mmol) in THF (30 ml) at 0°C, and the reaction mixture was stirred at room temperature for 22 hours. Water was added, followed by extraction with ethyl acetate. The organic layer was then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 100/0 - > 75/25) to give 4-difluoromethylene-cyclohexanecarboxylic acid benzyl ester (166 mg, 9%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.57 (2H, ddd, J = 12, 12, 4 Hz), 1.80-1.90 (2H, m), 1.97-2.05 (2H, m), 2.43-2.49 (3H, m), 5.12 (2H, s), 7.32-7.40 (5H, m).

(Reaction 248-3)



[1805] 4-Difluoromethylene-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 215-2 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.52-1.63 (2H, m), 1.85-1.94 (2H, m), 1.97-2.05 (2H, m), 2.43-2.49 (3H, m).

Example 249

N-[4-(2-{2-[4-(2,2-Difluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamide (Compound 1060)

[1806]


(Reaction 249-1)



[1807] Trifluoroacetic acid (6.17 mL, 83.0 mmol) was added to a solution of 4-amino-4-carbamoyl-piperidine-1-carboxylic acid tert-butyl ester (2.02 g, 8.30 mmol) in dichloromethane (16.6 mL), and the mixture was stirred at room temperature for 1.5 hours. The reaction solution was concentrated under reduced pressure. The residue was dissolved in methanol (2.00 mL), repeatedly concentrated under reduced pressure twice and dried under reduced pressure to give 4-amino-piperidine-4-carboxylic amide 2TFA salt as a colorless substance (3.25 g).

$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ 2.10-2.19 (2H, br-m), 2.56-2.65 (2H, m), 3.34-3.45 (4H, m).


(Reaction 249-2)



[1808] 1-[2-(4-Acetylamino-2-methyl-phenyl)-ethanesulfonyl]-4-amino-piperidine-4-carboxylic amide was synthesized by operations similar to those in Reaction 5-4 using appropriate reagents and starting material.

MS (ESI) m/z = 383 (M+H)+.

(Reaction 249-3)




[1809] N-[4-(2-{2-[4-(2,2-Difluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide was synthesized by operations similar to those in Reaction 10-14 and Reaction 189-5 using appropriate reagents and starting material.

MS (ESI) m/z = 539 (M+H)+.

[1810] The carboxylic acid reagent used in the synthesis of Compound 1060 (4-(2,2-difluoro-ethyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 249-4)



[1811] 4-(2,2-Difluoro-ethyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 191-11 and Reaction 95-18 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.00-2.04 (11H, m), 2.27 (0.25H, tt, J = 12.4, 3.2 Hz), 2.60-2.64 (0.75H, m), 5.68-6.01 (1H, m) (cis : trans = 3 : 1).

[1812] The example compounds shown below were synthesized by operations similar to those in Reaction 249-3 using appropriate reagents and starting materials.

(Compounds 1061 to 1063)


[1813]

[Table 158]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1061		LCMS-C-1	2.40	521 (M+H)+
1062		LCMS-B-1	1.98	549 (M+H)+
1063		LCMS-B-1	1.76	533 (M+H)+


[1814] The carboxylic acid reagent used in the synthesis of Compound 1061 (4-(2-fluoro-ethyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 249-5)



[1815] Sodium borohydride (89 mg, 2.35 mmol) was added to a solution of 4-(2-oxo-ethyl)-cyclohexanecarboxylic acid benzyl ester (306 mg, 1.18 mmol) in methanol (6 ml) at 0°C. The mixture was stirred at 0°C for one hour, and then quenched with a saturated aqueous ammonium chloride solution and extracted with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-(2-hydroxy-ethyl)-cyclohexanecarboxylic acid benzyl ester (299 mg).


¹H-NMR (400 MHz, CDCl₃) δ 1.20-1.32 (2H, m), 1.40-1.63 (9H, m), 1.95-2.05 (2H, m), 2.25-2.33 (0.2H, m), 2.55-2.62 (0.8H, m), 3.67 (1.6H, t, J = 6.8 Hz), 3.69 (0.4H, t, J = 6.4 Hz), 7.30-7.40 (5H, m) (cis : trans = 4 : 1).

(Reaction 249-6)


[1816] 4-(2-Fluoro-ethyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 191-11 and Reaction 95-18 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.93-2.03 (11H, m), 2.26 (0.2H, tt, $J = 12.0, 3.2$ Hz), 2.57-2.62 (0.8H, m), 4.39-4.56 (2H, m) (cis : trans = 4 : 1).

[1817] The carboxylic acid reagent used in the synthesis of Compound 1062 (4-butyl-4-fluoro-cyclohexanecarboxylic acid) was synthesized by the following method.


(Reaction 249-7)

[1818] A 2.6 M solution of $n\text{-BuLi}$ in THF (681 μl , 1.77 mmol) was added to a solution of 4-oxo-cyclohexanecarboxylic acid ethyl ester (186 μl , 1.18 mmol) in Et_2O (4.0 ml) at -60°C in an N_2 atmosphere, and the mixture was stirred at -60°C for four hours. The reaction mixture was quenched by adding water and then diluted with ethyl acetate. The organic layer was washed with water, and then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was used in the next step without purification.


(Reaction 249-8)

[1819] 4-Butyl-cyclohex-3-enecarboxylic acid ethyl ester was synthesized by operations similar to those in Reaction 25-15 using appropriate reagents and starting material.

MS (ESI) $m/z = 211$ ($\text{M}+\text{H}$) $^+$.

(Reaction 249-9)


[1820] HF-Py (0.5 ml) was added to 4-butyl-cyclohex-3-enecarboxylic acid ethyl ester (43.0 mg, 205 μ mol) at room temperature, and the reaction mixture was stirred at room temperature for two hours. The reaction mixture was diluted by adding dichloromethane and then quenched by adding a saturated aqueous sodium bicarbonate solution and solid sodium bicarbonate at 0°C. The organic layer was washed with 2 N HCl, and then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 100/0 $\rightarrow$ 97/3) to give 4-butyl-4-fluoro-cyclohexanecarboxylic acid ethyl ester (22.5 mg, 48%, cis:trans = 1:3). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.91 (3H, t, J = 8.0 Hz), 1.25 (3H, t, J = 8.0 Hz), 1.28-1.38 (6H, m), 1.51-1.84 (6H, m), 1.84-2.00 (2H, m), 2.20-2.26 (0.75H, m), 2.48-2.54 (0.25H, m), 4.10-4.16 (2H, m).

(Reaction 249-10)

[1821] 4-Butyl-4-fluoro-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.91 (3H, t, J = 7.0 Hz), 1.25-1.46 (6H, m), 1.50-1.62 (2H, m), 1.69-2.05 (6H, m), 2.25-2.35 (0.75H, m), 2.58-2.70 (0.25H, m).

[1822] The carboxylic acid reagent used in the synthesis of Compound 1063 (4-((E)-3-fluoro-propenyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 249-11)

[1823] 4-((E)-3-Fluoro-propenyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material.


tion 249-5, Reaction 191-11 and Reaction 95-18 using appropriate reagents and starting material. This was used in the next step without complete purification.

Example 250

N-[4-(2-{2-[4-(3-Fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide (Compound 1064) and N-[4-(2-{2-[4-(3-chloro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide (Compound 1065)

[1824]

(Reaction 250-1)



[1825] N-[4-(2-{2-[4-(3-Fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide


MS (ESI) m/z = 535 (M+H)⁺

and N-[4-(2-{2-[4-(3-chloro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide

MS (ESI) m/z = 551 (M+H)⁺

were synthesized by operations similar to those in Reaction 10-14 and Reaction 189-5 using appropriate reagents and starting material.

[1826] The carboxylic acid reagent used in the synthesis of Compound 1064 and Compound 1065 (a mixture of 4-(3-fluoro-propyl)-cyclohexanecarboxylic acid and 4-(3-chloro-propyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 250-2)


[1827] A mixture of 4-(3-fluoro-propyl)-cyclohexanecarboxylic acid and 4-(3-chloro-propyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 101-1, Reaction 39-2, Reaction 5-4, Reaction 119-3, Reaction 18-2 (using platinum oxide) and Reaction 95-18 using appropriate reagents and starting material. This was used in the next step without complete purification.

Example 251

N-[4-(2-[2-[4-(3-Fluoro-propylidene)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamide (Compound 1066)

[1828]

(Reaction 251-1)



[1829] N-[4-(2-{2-[4-(3-Fluoro-propylidene)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamide was synthesized by operations similar to those in Reaction 10-14 and Reaction 10-12 using appropriate reagents and starting material.

MS (ESI) $m/z = 533 (M+H)^+$.

[1830] The example compounds shown below were synthesized by operations similar to those in Reaction 251-1 using appropriate reagents and starting materials.







(Compounds 1067 to 1077)

[1831]




[Table 159]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1067		LCMS-C-2	2.00	551 (M+H)+
1068		LCMS-C-2	2.03	551 (M+H)+

(continued)


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1069		LCMS-C-2	2.08	607 (M+H)+
1070		LCMS-B-1	1.84	533 (M+H)+
1071		LCMS-C-1	2.43	539 (M+H)+
1072		LCMS-C-1	2.40	521 (M+H)+
1073		LCMS-F-1	0.87	553 (M+H)+
1074		LCMS-B-1	1.63	499 (M+H)+

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1075		LCMS-F-1	0.85	525 (M+H)+
1076		LCMS-B-1	1.79	551 (M+H)+
1077		LCMS-F-1	0.91	535 (M+H)+


[1832] The carboxylic acid reagent used in the synthesis of Compound 1066 (a mixture of 4-(3-fluoro-propylidene)-cyclohexanecarboxylic acid and 4-(3-chloro-propylidene)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 251-2)




[1833] A mixture of 4-(3-fluoro-propylidene)-cyclohexanecarboxylic acid and 4-(3-chloro-propylidene)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material. This was used in the next step without complete purification.

[1834] The carboxylic acid reagent used in the synthesis of Compound 1069 (4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 251-3)


[1835] 4-Trifluoromethanesulfonyloxymethyl-cyclohexanecarboxylic acid benzyl ester (cis:trans = 4:1) was synthesized by operations similar to those in Reaction 12-2 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.10 (0.4H, m), 1.34 (1.6H, m), 1.44-1.72 (4H, m), 1.83 (0.2H, m), 1.92 (0.8H, m), 2.11 (2H, m), 2.32 (0.2H, m), 2.68 (0.8H, m), 4.33 (1.6H, d, $J = 6.8$ Hz), 4.39 (0.4H, d, $J = 5.8$ Hz), 5.12 (0.4H, s), 5.14 (1.6H, s), 7.35 (5H, m).

(Reaction 251-4)

[1836] (Pentafluoroethyl)trimethylsilane (105 mg, 0.548 mmol) was added to a solution of 4-trifluoromethanesulfonyloxymethyl-cyclohexanecarboxylic acid benzyl ester (cis:trans = 4:1) (67.0 mg, 0.176 mmol) in DME (0.88 ml) at -30°C , and tetramethylammonium fluoride (21 mg, 0.22 mmol) was then added at -30 to -27°C over one hour. The mixture was stirred for four hours while warming from -30°C to 0°C and further stirred at 0°C for one hour. Water was added to the reaction mixture, followed by extraction with dichloromethane. The organic layer was then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 150/1) to give 4-(2,2,3,3,3-pentafluoropropyl)-cyclohexanecarboxylic acid benzyl ester (cis:trans = 11:1) (11.4 mg, 18%).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.32-1.75 (6H, m), 1.80-2.10 (5H, m), 2.26 (0.08H, m), 2.61 (0.92H, m), 5.13 (2H, s), 7.35 (5H, m).

(Reaction 251-5)

[1837] $\text{LiOH}\cdot\text{H}_2\text{O}$ (3.8 mg, 0.091 mmol) was added to a solution of 4-(2,2,3,3,3-pentafluoropropyl)-cyclohexanecarboxylic acid benzyl ester (cis:trans = 11:1) (11.4 mg, 0.0325 mmol) in THF (0.15 ml)- H_2O (0.15 mL). The mixture was stirred at room temperature for 16 hours, and then adjusted to pH 2 with a 1 N aqueous HCl solution and extracted with dichloromethane. The organic layer was dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was dissolved in tert-butanol (0.4 ml), and potassium tert-butoxide (10 mg, 0.089 mmol) was added. The mixture was stirred at room temperature for two hours, and then adjusted to pH 3 with a 1 N aqueous HCl solution and extracted with dichloromethane. The organic layer was then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by column chromatography (hexane/ethyl acetate = 5/1) to give 4-(2,2,3,3,3-pentafluoropropyl)-cyclohexanecarboxylic acid (cis:trans = 9:1) (8.5 mg, 100%).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.11 (0.2H, m), 1.30-1.78 (5.8H, m), 1.83-2.10 (5H, m), 2.26 (0.1H, m), 2.64 (0.9H, m).

[1838] The carboxylic acid reagent used in the synthesis of Compound 1070 (4-((E)-3-fluoro-allyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 251-6)

[1839] A 3.3 M Red-Al solution in toluene (0.25 ml, 0.82 mmol) was added to a solution of 4-(3,3-difluoro-allyl)-cyclohexanecarboxylic acid benzyl ester (cis:trans = 5.4:1) (42.8 mg, 0.145 mmol) in toluene (0.12 mL), and the mixture was stirred at 85°C for 17 hours. The reaction mixture was poured into ice water, adjusted to pH 3 with a 4 N aqueous H₂SO₄ solution and extracted with ether. The organic layer was dried over MgSO₄ and concentrated under reduced pressure to give [4-((E)-3-fluoro-allyl)-cyclohexyl]-methanol (25.0 mg, 100%).


¹H-NMR (400 MHz, CDCl₃) δ 0.92 (0.84H, m), 1.20-1.84 (9.16H, m), 1.87 (1.68H, dd, J = 7.8, 6.8 Hz), 2.10 (0.32H, t, J = 7.3 Hz), 3.45 (0.32H, d, J = 5.8 Hz), 3.54 (1.68H, d, J = 6.8 Hz), 5.31 (1H, m), 6.47 (1H, dd, J = 86.0, 11.0 Hz).

(Reaction 251-7)

[1840] 4-((E)-3-Fluoro-allyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 109-1 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 0.80-2.10 (11H, m), 2.26 (0.15H, m), 2.61 (0.85H, m), 4.71 (0.16H, dm, J = 40.6 Hz), 5.30 (0.84H, m), 6.47 (1H, dd, J = 86.0, 11.0 Hz) (cis : trans = 85 : 15).

[1841] The carboxylic acid reagent used in the synthesis of Compound 1073 (4-fluoro-4-(3-fluoro-propyl)-cyclohexanecarboxylic acid) was synthesized by the following method.


(Reaction 251-8)

[1842] 4-Fluoro-4-(3-fluoro-propyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 249-9 and Reaction 95-18 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 2.36-2.17 (3H, m), 2.05-1.66 (8H, m), 1.47-1.25 (2H, m).

[1843] The carboxylic acid reagent used in the synthesis of Compound 1074 (4-ethynyl-cyclohexanecarboxylic acid) was synthesized by the following method.


(Reaction 251-9)



[1844] Dimethyl (1-diazo-2-oxopropyl)phosphonate (Bestmann reagent) (450 mg, 2.34 mmol) was added to a mixture of 4-formyl-cyclohexanecarboxylic acid benzyl ester (390 mg, 1.58 mmol) and potassium carbonate (323 mg, 2.34 mmol) in methanol (10 mL) at 0°C and the mixture was stirred for five hours. Further, the reaction solution was stirred at room temperature for two hours. A saturated aqueous ammonium chloride solution and ethyl acetate were then added at 0°C, and the organic layer and the aqueous layer were separated. The aqueous layer was repeatedly extracted with ethyl acetate three times, and the organic layers were then combined, washed with saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give a mixture of 4-ethynyl-cyclohexanecarboxylic acid methyl ester and 4-ethynyl-cyclohexanecarboxylic acid benzyl ester as a colorless liquid (220 mg, 78%).

^1H -NMR (400 MHz, $CDCl_3$) δ 1.34-1.48 (1.6H, m), 1.48-1.62 (1H, m), 1.65-1.87 (2.6H, m), 1.87-2.10 (3.8H, m), 2.20-2.42 (1.4H, m), 2.65-2.75 (0.6H, m), 3.66 (1.1H, s), 3.68 (1.4H, s), 5.10 (0.05H, s), 5.13 (0.3H, s), 7.29-7.40 (0.9H, m) (Me : Bn = 0.85 : 0.15, cis : trans = 0.6 : 0.4).

(Reaction 251-10)




[1845] 4-Ethynyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material.

^1H -NMR (400 MHz, CD_3OD) δ 1.37-1.50 (1.3H, m), 1.53-1.67 (1.6H, m), 1.67-1.82 (2.8H, m), 1.82-2.05 (2.5H, m), 2.18-2.38 (2.1H, m), 2.62-2.73 (0.7H, m).

[1846] The carboxylic acid reagent used in the synthesis of Compound 1077 (4-(2-fluoro-propyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 251-11)




[1847] Methylmagnesium bromide (1 M solution in THF, 0.526 mL, 0.526 mmol) was added dropwise to 4-(2-oxoethyl)-cyclohexanecarboxylic acid benzyl ester (114 mg, 0.439 mmol) in THF (2.2 mL) at -78°C, and the mixture was stirred at the same temperature for 30 minutes. Water was added to the reaction mixture at the same temperature and extracted with ethyl acetate. The organic layer was sequentially washed with a saturated aqueous ammonium chloride

solution, water and saturated brine and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-(2-hydroxy-propyl)-cyclohexanecarboxylic acid benzyl ester (80.7 mg, 67%).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 1.18 (3H, d, J = 6.1 Hz), 1.22-1.49 (5H, m), 1.50-1.67 (6H, m), 1.95-2.05 (2H, m), 2.58 (1H, dt, J = 9.1, 4.9 Hz), 3.82-3.94 (1H, m), 5.13 (2H, s), 7.29-7.39 (5H, m);

MS (ESI) m/z = 259 ($\text{M-H}_2\text{O}+\text{H}$) $^+$.

(Reaction 251-12)



[1848] 4-(2-Fluoro-propyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 191-11 and Reaction 95-18 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.23-2.07 (11H, m), 1.31 (3H, dd, J = 23.9, 7.0 Hz), 2.55-2.64 (1H, m), 4.63-4.87 (1H, m).

Example 252

N-[4-(2-{2-[4-(1-Fluoro-1-methyl-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide (Compound 1078)

[1849]


(Reaction 252-1)



[1850] N-[4-(2-{2-[4-(1-Fluoro-1-methyl-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide was synthesized by operations similar to those in Reaction 10-14 and Reaction 101-3 using appropriate reagents and starting material.


MS (ESI) m/z = 535 ($\text{M}+\text{H}$) $^+$.

[1851] The carboxylic acid reagent used in the synthesis of Compound 1078 (4-(1-fluoro-1-methyl-ethyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 252-2)


[1852] 4-(1-Hydroxy-ethyl)-cyclohexanecarboxylic acid benzyl ester was synthesized by operations similar to those in Reaction 251-11 using appropriate reagents and starting material.

MS (ESI) $m/z = 263$ ($M+H$) $^+$.

(Reaction 252-3)

[1853] Dess-Martin reagent (151 mg, 0.355 mmol) was added to a solution of 4-(1-hydroxy-ethyl)-cyclohexanecarboxylic acid benzyl ester (71.6 mg, 0.273 mmol) in anhydrous dichloromethane (0.91 ml) at 0°C . The mixture was stirred at the same temperature for 10 minutes, and further warmed to room temperature and stirred for 4.5 hours. An aqueous sodium thiosulfate solution was added to the reaction solution, followed by extraction with dichloromethane. The organic phase was concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 4-acetyl-cyclohexanecarboxylic acid benzyl ester (60.2 mg, 85%).

MS (ESI) $m/z = 261$ ($M+H$) $^+$.

(Reaction 252-4)

[1854] 4-(1-Fluoro-1-methyl-ethyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 251-11, Reaction 191-11 and Reaction 95-18 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (300 MHz, CD_3OD) δ 1.20-1.32 (2H, m), 1.24 (6H, d, $J = 21.9$ Hz), 1.45-1.58 (3H, m), 1.62-1.70 (2H, m),


2.17-2.26 (2H, m), 2.60-2.65 (1H, m).

Example 253

5 N-(4-{2-[2-(4-Butylidene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide (Compound 1079)

[1855]

(Reaction 253-1)




[1856] N-(4-{2-[2-(4-Butylidene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methylphenyl)-acetamide was synthesized by operations similar to those in Reaction 10-1 and Reaction 10-12 using appropriate reagents and starting material.

MS (ESI) m/z = 529 (M+H)+.

[1857] The carboxylic acid reagent used in the synthesis of Compound 1079 (4-butylidene-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 253-2)



[1858] 4-Butylidene-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 191-14 and Reaction 95-18 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.88 (3H, t, $J = 7.3$ Hz), 1.28-1.40 (2H, m), 1.42-1.62 (2H, m), 1.77-1.88 (1H, m), 1.89-2.12 (5H, m), 2.20-2.29 (1H, m), 2.46-2.55 (1H, m), 2.55-2.63 (1H, m), 5.14 (1H, t, $J = 7.3$ Hz).

Example 254

N-[3-Methyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide (Compound 1080)

[1859]

(Reaction 254-1)



[1860] N-[3-Methyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sul-

fonyl)-ethyl)-phenyl]-acetamide was synthesized by operations similar to those in Reaction 10-14 and Reaction 101-3 using appropriate reagents and starting material.


MS (ESI) m/z = 607 (M+H)+.

Example 255

1-{3,5-Dimethyl-4-[2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea (Compound 1081)

[1861]


(Reaction 255-1)



[1862] 4-Amino-1-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-piperidine-4-carboxylic amide was synthesized by operations similar to those in Reaction 184-1 using appropriate reagents and starting material.

MS (ESI) m/z = 369 (M+H)+.

(Reaction 255-2)



[1863] 1-{3,5-Dimethyl-4-[2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 101-3 and Reaction 89-2 using appropriate reagents and starting material.



MS (ESI) m/z = 544 (M+H)+.

[1864] The example compounds shown below were synthesized by operations similar to those in Reaction 255-2 using appropriate reagents and starting materials.

(Compounds 1082 to 1083)

[1865]


[Table 160]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1082		LCMS-F-1	1.01	544 (M+H)+
1083		LCMS-C-2	2.27	544 (M+H)+

Example 256

1-(3,5-Dimethyl-4-{1-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropylmethyl}-phenyl)-1-methyl-urea (Compound 1084)


[1866]

(Reaction 256-1)

[1867] n-Butyllithium (1.6 M solution in hexane, 0.58 ml, 0.93 mmol) was added to a solution of {4-[2-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-methyl-carbamic acid tert-butyl ester (195.7 mg, 0.4176 mmol) in THF (4.2 ml) at -78°C over six minutes, and the mixture was stirred at the same temperature for 20 minutes. 1-Bromo-2-chloro-ethane (105 µl, 1.26 mmol) was added to the reaction solution at -78°C within 10 minutes. The mixture was then stirred while warming from -78°C to 0°C over one hour, and further stirred at room temperature for one hour. A 50% saturated aqueous ammonium chloride solution was added, followed by extraction with ethyl acetate. The organic layer was then dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 5/2) to give {4-[4-chloro-2-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-butyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester (133 mg, 60%).


¹H-NMR (400 MHz, CDCl₃) δ 1.44 (9H, s), 1.80 (4H, t, J = 6.0 Hz), 1.85 (1H, m), 2.28 (1H, m), 2.33 (6H, s), 3.00 (1H, dd, J = 14.0, 11.6 Hz), 3.18 (1H, dd, J = 14.0, 4.4 Hz), 3.22 (3H, s), 3.33 (1H, m), 3.52 (4H, t, J = 6.0 Hz), 3.60 (2H, m), 3.98 (4H, s), 6.92 (2H, s).

(Reaction 256-2)



[1868] n -Butyllithium (1.6 M solution in hexane, 345 μl , 0.552 mmol) was added to a solution of {4-[4-chloro-2-(1,4-dioxaspiro[4.5]decane-8-sulfonyl)-butyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester (115.0 mg, 0.2165 mmol) in THF (3.0 ml) at -78°C over three minutes. The mixture was stirred at the same temperature for five minutes, and then warmed and further stirred at room temperature for one hour. A 50% saturated aqueous ammonium chloride solution was added, followed by extraction with ethyl acetate. The organic layer was then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 2/1) to give {4-[1-(1,4-dioxaspiro[4.5]decane-8-sulfonyl)-cyclopropylmethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester and {4-[1-(1,4-dioxaspiro[4.5]decane-8-sulfonyl)-cyclopropylmethyl]-3,5-dimethyl-phenyl}-methyl-amine as a mixture (46.0 mg, 43% and 19.6 mg, 23%). This was used in the next reaction without complete purification.


(Reaction 256-3)




[1869] A 5 N aqueous HCl solution (2.0 ml, 10 mmol) was added to a solution of a mixture of 4-[1-(1,4-dioxaspiro[4.5]decane-8-sulfonyl)-cyclopropylmethyl]-3,5-dimethyl-phenyl-methyl-carbamic acid tert-butyl ester (54.7 mg, 0.111 mmol) and {4-[1-(1,4-dioxaspiro[4.5]decane-8-sulfonyl)-cyclopropylmethyl]-3,5-dimethyl-phenyl}-methyl-amine (23.1 mg, 0.0585 mmol) in EtOH (2.0 mL), and the mixture was stirred at 80°C for two hours. The reaction mixture was neutralized by adding a 5 N aqueous NaOH solution (1.9 ml) and then extracted with ethyl acetate. The organic layer was dried over MgSO_4 and concentrated under reduced pressure to give 1-[1-(2,6-dimethyl-4-methylamino-benzyl)-cyclopropanesulfonyl]-4-piperidinone (58.8 mg, 99%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.44 (2H, m), 1.21 (2H, m), 2.22 (6H, s), 2.60 (4H, t, $J = 6.0$ Hz), 2.79 (3H, s), 3.27 (2H, s), 3.78 (4H, t, $J = 6.0$ Hz), 3.99 (4H, s), 6.28 (2H, s).

(Reaction 256-4)



[1870] {4-[1-(4-Amino-4-carbamoyl-piperidine-1-sulfonyl)-cyclopropylmethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 127-2 (using toluene as a solvent), Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material. $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.37 (2H, m), 1.19 (2H, m), 1.45 (9H, s), 1.59 (2H, m), 2.24 (2H, m), 2.30 (6H, s), 3.20 (3H, s), 3.36 (2H, s), 3.43 (2H, ddd, $J = 13.0, 9.6$ and 3.2 Hz), 3.77 (2H, ddd, $J = 13.0, 5.2$ and 4.8 Hz), 5.39 (1H, br), 6.89 (2H, s), 7.27 (1H, br).


(Reaction 256-5)

[1871] 1-(3,5-Dimethyl-4-{1-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropylmethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 101-3, Reaction 4-1 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) m/z = 608 (M+H)+.

Example 257

1-(4-{2,2-Difluoro-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1085)


[1872](Reaction 257-1)

[1873] n-Butyllithium (1.54 M solution in hexane, 5.6 ml, 8.67 mmol) was added to a solution of diisopropyl-amine (1.45 mL, 10.4 mmol) in tetrahydrofuran (30 mL) at 0°C, and the mixture was stirred for 20 minutes. The reaction solution

was brought to -78°C , and a solution of {4-[2-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester (2.71 g, 5.78 mmol) in tetrahydrofuran (10 mL) was then added dropwise slowly. The reaction solution was stirred at -78°C for 0.5 hour, and N-fluorobenzenesulfonimide (2.73 g, 8.67 mmol) was then added at -78°C , followed by stirring for one hour. A saturated aqueous ammonium chloride solution was added to the reaction solution at -78°C , and the mixture was brought to room temperature. Ethyl acetate was then added, and the organic layer and the aqueous layer were separated. The aqueous layer was repeatedly extracted with ethyl acetate three times. The organic layers were then combined and washed with saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give {4-[2-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-2,2-difluoro-ethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester as a white foamy solid (205 mg, 7%).

MS (ESI) $m/z = 505$ (M+H) $^{+}$.


(Reaction 257-2)



[1874] {4-[2-(4-Amino-4-carbamoyl-piperidine-1-sulfonyl)-2,2-difluoro-ethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester was synthesized by operations similar to those in Reaction 233-2, Reaction 19-2, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 505$ (M+H) $^{+}$.

(Reaction 257-3)




[1875] 1-(4-{2,2-Difluoro-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 10-12, Reaction 4-1 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 618 (M+H)^+$.

Example 258

1-{4-[2,2-Difluoro-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea (Compound 1086)

[1876]


(Reaction 258-1)

[1877] 1-{4-[2,2-Difluoro-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 101-3, Reaction 4-1 and Reaction 89-2 using appropriate reagents and starting material.
MS (ESI) m/z = 584 (M+H)+.

Example 259

1-{3,5-Dimethyl-4-[2-(4-oxo-2-[1,1';2',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea (Compound 1087)


[1878](Reaction 259-1)



[1879] 1-{3,5-Dimethyl-4-[2-(4-oxo-2-[1,1';2',1'']terphenyl-3-yl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 10-11, Reaction 10-12 (using ethanol as a solvent), Reaction 89-2 and Reaction 122-2 using appropriate reagents and starting material. MS (ESI) $m/z = 650$ (M+H)+.

[1880] The carboxylic acid reagent used in the synthesis of Compound 1087 ([1,1';2',1'']terphenyl-3"-carboxylic acid) was synthesized by the following method.

(Reaction 259-2)




[1881] 3-Boronobenzoic acid (300 mg, 1.81 mmol), palladium acetate (40.4 mg, 0.18 mmol), triphenylphosphine (94.6 mg, 0.36 mmol) and potassium carbonate (374.6 mg, 2.71 mmol) were added to a solution of 2-bromo-1,1'-biphenyl (0.3 ml, 1.81 mmol) in acetonitrile (10 mL)-water (2.5 ml), and the mixture was heated with stirring at 100°C overnight. The reaction mixture was cooled and then filtered through celite, and the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane:ethyl acetate = 1:1) to give [1,1';2',1'']terphenyl-3"-carboxylic acid (191 mg, 39%).

MS (ESI) m/z = 275 (M+H)+.

Example 260

5 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-phenyl]-1-methyl-urea (Compound 1088)

[1882]




[1883] 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 1-4 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) m/z = 606 (M+H)+.

[1884] The carboxylic acid reagent used in the synthesis of Compound 1088 (4-(4,4,4-trifluoro-butyl)-benzoic acid) was synthesized by the following method.

(Reaction 260-2)



[1885] 4-(4,4,4-Trifluoro-butyl)-benzoic acid was synthesized by operations similar to those in Reaction 191-14, Reaction 18-2 and Reaction 95-18 using appropriate reagents and starting material.

MS (ESI) m/z = 233 (M+H)+.

[1886] The example compound shown below was synthesized by operations similar to those in Reaction 260-1 using appropriate reagents and starting material

(Compound 1089)


[1887]

[Table 161]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1089		LCMS-D-1	2.63	606 (M+H)+

[1888] The carboxylic acid reagent used in the synthesis of Compound 1089 (3-(4,4,4-trifluoro-butyl)-benzoic acid) was synthesized by the following method.

(Reaction 260-3)



[1889] 3-(4,4,4-Trifluoro-butyl)-benzoic acid was synthesized by operations similar to those in Reaction 191-14, Reaction 18-2 and Reaction 95-18 using appropriate reagents and starting material.


MS (ESI) $m/z = 233$ (M+H)+.

Example 261

1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-propyl}-phenyl)-1-methyl-urea (Compound 1090)

[1890]

(Reaction 261-1)




[1891] {4-[2-(4-Amino-4-carbamoyl-piperidine-1-sulfonyl)-propyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-

butyl ester was synthesized by operations similar to those in Reaction 256-1 (using LDA as a base), Reaction 233-2 (using hydrochloric acid as an acid), Reaction 127-2, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.21 (3H, d, J = 6.4 Hz), 1.46 (9H, s), 1.57 (2H, m), 2.19 (2H, m), 2.32 (6H, s), 2.90 (1H, dd, J = 14.2 and 12.2 Hz), 3.21 (3H, s), 3.26 (2H, m), 3.40 (2H, m), 3.68 (2H, m), 5.33 (1H, br), 6.91 (2H, s), 7.23 (1H, br).

(Reaction 261-2)




[1892] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-propyl)-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 101-3, Reaction 4-1 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) m/z = 596 (M+H)⁺.


Example 262

2-[4-Fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1091)

[1893]

(Reaction 262-1)


[1894] 4-Amino-1-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-piperidine-4-carboxylic acid was synthesized by operations similar to those in Reaction 119-1, Reaction 233-2, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.
MS (ESI) m/z = 465 (M+H)+.

(Reaction 262-2)

[1895] 2-[4-Fluoro-3-(4,4,5,5,5-pentafluoropentyloxy)-phenyl]-8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 10-12 using appropriate reagents and starting material.
MS (ESI) m/z = 745 (M+H)+.

[1896] The carboxylic acid reagent used in the synthesis of Compound 1091 (4-fluoro-3-(4,4,5,5,5-pentafluoropentyloxy)-benzoic acid) was synthesized by the following method.

(Reaction 262-3)



[1897] 4-Fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-benzoic acid was synthesized by operations similar to those in Reaction 31-7 and Reaction 95-18 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ 0.21-0.14 (2H, m), 2.30-2.42 (2H, m), 4.19 (2H, t, $J = 6.0$ Hz), 7.20 (1H, dd, $J = 10.8, 8.4$ Hz), 7.64-7.68 (1H, m), 7.72 (1H, dd, $J = 8.4, 2.0$ Hz).

Example 263

2-[4-Fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1092)


[1898]

(Reaction 263-1)



[1899] 4-Amino-1-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-piperidine-4-carboxylic amide was synthesized by operations similar to those in Reaction 10-18, Reaction 119-1, Reaction

233-3 and Reaction 233-4 using appropriate reagents and starting material.
MS (ESI) $m/z = 479$ (M+H)+.

(Reaction 263-2)

[1900] 2-[4-Fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 10-12 using appropriate reagents and starting material.

MS (ESI) $m/z = 759$ (M+H)+.

[1901] The example compounds shown below were synthesized by operations similar to those in Reaction 263-2 using appropriate reagents and starting materials.






(Compounds 1093 to 1099)

[1902]


[Table 162]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1093		LCMS-F-1	1.03	683 (M+H)+
1094		LCMS-F-1	1.01	667 (M+H)+

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1095		LCMS-F-1	1.03	701 (M+H)+
1096		LCMS-F-1	1.05	741 (M+H)+
1097		LCMS-C-1	2.60	647 (M+H)+
1098		LCMS-C-1	2.60	597 (M+H)+
1099		LCMS-F-1	1.01	667 (M+H)+


[1903] The carboxylic acid reagent used in the synthesis of Compound 1096 (3-(4,4,5,5,5-pentafluoro-pentyloxy)-benzoic acid) was synthesized by the following method.

(Reaction 263-3)

[1904] 3-(4,4,5,5,5-Pentafluoro-pentyloxy)-benzoic acid was synthesized by operations similar to those in Reaction 31-7 and Reaction 95-18 using appropriate reagents and starting material.

MS (ESI) $m/z = 297$ (M-H)-.

[1905] The carboxylic acid reagent used in the synthesis of Compound 1097 (3-(2,2,2-trifluoro-ethyl)-benzoic acid) was synthesized by the following method.

(Reaction 263-4)


[1906] 3-(2,2,2-Trifluoro-ethyl)-benzoic acid was synthesized by operations similar to those in Reaction 193-4, Reaction 193-5, Reaction 193-6 and Reaction 95-18 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.43 (2H, q, $J = 10.8$ Hz), 3.93 (3H, s), 7.43-7.51 (2H, m), 7.99-8.04 (2H, m).

Example 264

8-[(E)-2-[4-(4-Hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1100)


[1907]

(Reaction 264-1)

[1908] 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 189-5 using appropriate reagents and starting material.
MS (ESI) m/z = 681 (M+H)+.


Example 265

8-{1,1-Difluoro-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1101)

[1909](Reaction 265-1)


[1910] Dehydrated tert-butanol (0.19 ml, 2.0 mmol), 4-dimethylaminopyridine (81 mg, 0.67 mmol) and EDCI (255 mg, 1.33 mmol) were added to a solution of 4-bromo-3,5-dimethyl-benzoic acid (123 mg, 0.535 mmol) in dichloromethane (1.0 ml) at 0°C, and the mixture was stirred at room temperature for 25 hours. The reaction mixture was diluted with dichloromethane, and the organic layer was washed with water, dried over MgSO₄ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 100/1 → 50/1) to give 4-bromo-3,5-dimethyl-benzoic acid tert-butyl ester (106 mg, 70%).
¹H-NMR (400 MHz, CDCl₃) δ 1.58 (9H, s), 2.45 (6H, s), 7.66 (2H, s).

(Reaction 265-2)



[1911] 4-Amino-1-[1,1-difluoro-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-piperidine-4-carboxylic amide was synthesized by operations similar to those in Reaction 26-1, Reaction 184-1, Reaction 257-1, Reaction 257-1, Reaction 233-2, Reaction 10-14, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.

¹H-NMR (400 MHz, CDCl₃) δ 1.07 (2H, br), 1.53 (2H, m), 1.84 and 1.95 (each 1H, br), 2.21 (2H, m), 2.36 (6H, s), 3.21 and 3.33 (each 1H, br), 3.48 (2H, m), 3.68 (2H, dd, J = 20.4 and 18.4 Hz), 3.70 (1H, br), 3.86 (2H, m), 3.97 (1H, m), 4.20 (1H, br), 5.33 (1H, br), 7.07 (2H, s), 7.19 (1H, br).


(Reaction 265-3)

[1912] 8-{1,1-Difluoro-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 101-3 using appropriate reagents and starting material.

MS (ESI) *m/z* = 673 (M+H)+.

Example 266

2-(4-Fluoro-3-trifluoromethoxy-phenyl)-8-[(E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1102)


[1913]**(Reaction 266-1)**

[1914] 4-Amino-1-[(E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-piperidine-4-car-

boxylic acid was synthesized by operations similar to those in Reaction 119-1, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 437$ (M+H)⁺.

(Reaction 266-2)




[1915] 2-(4-Fluoro-3-trifluoromethoxy-phenyl)-8-((E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 10-12 using appropriate reagents and starting material.

MS (ESI) $m/z = 625$ (M+H)⁺.

Example 267

8-[(E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethylphenyl]-ethenesulfonyl]-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1103)


[1916]

(Reaction 267-1)

[1917] 4-Amino-1-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-piperidine-4-carboxylic amide was synthesized by operations similar to those in Reaction 26-1, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 428$ ($\text{M}+\text{H}$) $^+$.

(Reaction 267-2)




[1918] 8-[(E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-[3-(4,4,5,5,5-pentafluoropentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 10-1 and Reaction 189-5 using appropriate reagents and starting material.
MS (ESI) m/z = 690 (M+H)+.

Example 268

6-(4-Methyl-cyclohexyl)-2-(2-naphthalen-1-yl-ethanesulfonyl)-2,5,7-triaza-spiro[3.4]oct-5-en-8-one (Compound 1104)

[1919]


(Reaction 268-1)



[1920] 3-Amino-1-(2-naphthalen-1-yl-ethanesulfonyl)-, azetidine-3-carboxylic amide was synthesized by operations similar to those in Reaction 190-1, Reaction 109-1, Reaction 200-2, Reaction 200-3 and Reaction 233-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 334$ (M+H)+.

(Reaction 268-2)



[1921] 6-(4-Methyl-cyclohexyl)-2-(2-naphthalen-1-yl-ethanesulfonyl)-2,5,7-triaza-spiro[3.4]oct-5-en-8-one was synthesized by operations similar to those in Reaction 10-14 and Reaction 10-12 using appropriate reagents and starting material.


MS (ESI) $m/z = 440$ (M+H)+.

Example 269

8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(11-hydroxy-undecyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one

[1922]

(Reaction 269-1)



[1923] HATU (57 mg, 0.149 mmol) was added to a solution of 12-hydroxy-dodecanoic acid (33 mg, 0.149 mol), 4-amino-1-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-piperidine-4-carboxylic amide (50 mg, 0.136 mmol) and diisopropylethylamine (71 μ L, 0.47 mmol) in DMF (1.3 ml) at 0°C, and the mixture was stirred at room temperature for 1.5 hours. Ethanol (2.6 ml) and potassium t-butoxide (76 mg, 0.678 mmol) were added to the reaction mixture, and the mixture was heated with stirring at 70°C for three hours. The reaction mixture was quenched with a saturated aqueous ammonium chloride solution, and water was then added, followed by extraction with ethyl acetate. The organic layer was sequentially washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-methanol) to give 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(11-hydroxy-undecyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (58.7 mg, 79%).

MS (ESI) $m/z = 547$ (M+H)+.

[1924] The example compounds shown below were synthesized by operations similar to those in Reaction 269-1 using appropriate reagents and starting materials.


(Compounds 1106 to 1107)

[1925]

[Table 163]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1106		LCMS-C-1	3.12	646 (M+H)+

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1107		LCMS-G-1	0.93	519 (M+H) ⁺

Example 270

1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (Compound 1108)





[1926]**(Reaction 270-1)**

[1927] 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-((1S,3R)-3-propylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.
MS (ESI) m/z = 544 (M+H)⁺.

[1928] The example compounds shown below were synthesized by operations similar to those in Reaction 270-1 using appropriate reagents and starting materials.

(Compounds 1109 to 1113)

[Table 164]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1109		LCMS-C-1	2.72	646 (M+H)+
1110		LCMS-F-1	0.93	520 (M+H)+
1111		LCMS-F-1	0.93	598 (M+H)+
1112		LCMS-F-1	1.03	714 (M+H)+

5

10

15

20

25

30

35


40

45

50


55

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1113		LCMS-F-1	0.99	598 (M+H)+

[1930] The carboxylic acid reagent used in the synthesis of Compound 1110 (4-[1,1,1-²H₃]methyl-[4-²H₁]cyclohexanecarboxylic acid) was synthesized by the following method.


(Reaction 270-2)



[1931] 4-[1,1,1-²H₃]Methylene-cyclohexanecarboxylic acid ethyl ester was synthesized by operations similar to those in Reaction 101-1 using appropriate reagents and starting material.


¹H-NMR (400 MHz, CDCl_3) δ 1.25 (3H, t, $J = 7.2$ Hz), 1.52-1.64 (2H, m), 1.95-2.10 (4H, m), 2.34 (2H, ddd, $J = 13.6, 4.4, 4.4$ Hz), 2.44 (1H, dddd, $J = 10.8, 10.8, 3.6, 3.6$ Hz), 4.13 (2H, q, $J = 7.6$ Hz).

(Reaction 270-3)



[1932] 20% w/w Pd/C (2.6 mg) was added to a solution of 4-[1,1,1-²H₃]methylene-cyclohexanecarboxylic acid ethyl ester (26.0 mg, 153 μmol) in MeOH (1 ml) in an N_2 atmosphere. After deuterium substitution, the reaction mixture was stirred at room temperature for one hour. The reaction mixture was filtered through celite, and the filtrate was concentrated under reduced pressure to give 4-[1,1,1-²H₃]methyl-[4-²H₁]cyclohexanecarboxylic acid ethyl ester. This was used in the next step without further purification.

(Reaction 270-4)




[1933] 4-[1,1,1-²H₃]Methyl-[4-²H₁]cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material. This was used in the next step without further purification.

Example 271

1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (Compound 1114)


[1934]

(Reaction 271-1)

[1935] 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 614$ ($M+H$) $^+$.

[1936] The carboxylic acid reagent used in the synthesis of Compound 1114 (4'-propyl-biphenyl-3-carboxylic acid) was synthesized by the following method.

(Reaction 271-2)

[1937] 4'-Propyl-biphenyl-3-carboxylic acid was synthesized by operations similar to those in Reaction 259-2 using appropriate reagents and starting material.









MS (ESI) $m/z = 241$ ($M+H$) $^+$.

[1938] The example compounds shown below were synthesized by operations similar to those in Reaction 259-1 using appropriate reagents and starting materials.

(Compounds 1115 to 1130)

[1939]


[Table 165]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1115		LCMS-D-1	1.84	548 (M+H)+
1116		LVMS-D-1	2.82	634 (M+H)+
1117		LCMS-D-1	2.75	634 (M+H)+
1118		LCMS-F-1	1.07	592 (M+H)+
1119		LCMS-F-1	0.92	502 (M+H)+
1120		LCMS-F-1	1.02	578 (M+H)+
1121		LCMS-F-1	1.11	620 (M+H)+
1122		LCMS-D-1	1.75	564 (M+H)+


(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1123		LCMS-D-1	1.81	548 (M+H)+
1124		LCMS-D-1	1.98	548 (M+H)+
1125		LCMS-F-1	1.05	610 (M+H)+
1126		LCMS-D-1	2.48	588 (M+H)+
1127		LCMS-D-1	2.12	578 (M+H)+
1128		LCMS-D-1	2.98	596 (M+H)+
1129		LCMS-F-1	0.95	533 (M+H)+

(continued)


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1130		LCMS-F-1	0.97	536 (M+H) ⁺

[1940] The carboxylic acid reagent used in the synthesis of Compound 1115 (7-ethoxy-heptanoic acid) was synthesized by the following method.



[1941] 0.5 M sodium ethoxide (1.6 ml, 4.31 mmol) was added to a solution of ethyl 7-bromohexanoate (300 mg, 1.44 mmol) in ethanol (7.0 ml), and the mixture was heated under reflux for two hours. The reaction mixture was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography (hexane:ethyl acetate = 10:1) to give ethyl 7-ethoxyhexanoate (102.4 mg, 40%).


¹H-NMR (300 MHz, DMSO-d₆) δ 1.18 (t, 3H, J = 7.3 Hz), 1.24 (t, 3H, J = 7.3 Hz), 1.29-1.38 (m, 4H), 1.52-1.68 (m, 4H), 2.28 (t, 2H, J = 7.6 Hz), 3.45-3.51 (m, 4H), 4.05-4.17 (m, 2H).



[1942] 7-Ethoxy-heptanoic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material.


¹H-NMR (300 MHz, DMSO-d₆) δ 1.13 (m, 3H), 1.35 (m, 4H), 1.58 (m, 4H), 2.23 (m, 2H), 3.49 (m, 4H), 12.36 (s, 1H).

[1943] The carboxylic acid reagent used in the synthesis of Compound 1116 (3-(6,6,6-trifluoro-hexyl)-benzoic acid) was synthesized by the following method.

(Reaction 271-5)

[1944] Acrolein (180 μl , 2.71 mmol), tetrabutylammonium bromide (385 mg, 1.19 mmol), palladium acetate (5 mmol%) and potassium carbonate (450 mg, 3.26 mmol) were added to a solution of methyl 3-iodobenzoate (300 mg, 1.08 mmol) in DMF (6.0 ml), and the mixture was heated with stirring at 80°C for two hours. The reaction mixture was cooled to room temperature and then diluted with ethyl acetate, and the organic layer was washed with water and saturated brine. The organic layer was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography (hexane:ethyl acetate = 8:1) to give ethyl 3-(3-oxoprop-1-en-1-yl)benzoate as a white solid (280 mg, 96%).


MS (ESI) m/z = 205 (M+H)+.

(Reaction 271-6)

[1945] 3-(6,6,6-Trifluoro-hexyl)-benzoic acid was synthesized by operations similar to those in Reaction 191-14, Reaction 18-2 and Reaction 95-18 using appropriate reagents and starting material.

MS (ESI) m/z = 261 (M+H)+.

[1946] The carboxylic acid reagent used in the synthesis of Compound 1117 (4-(6,6,6-trifluoro-hexyl)-benzoic acid) was synthesized by the following method.


(Reaction 271-7)

[1947] 4-(6,6,6-Trifluoro-hexyl)-benzoic acid was synthesized by operations similar to those in Reaction 191-14, Reaction 18-2 and Reaction 95-18 using appropriate reagents and starting material.

MS (ESI) m/z = 261 (M+H)+.

[1948] The carboxylic acid reagent used in the synthesis of Compound 1120 ((E)-7-phenyl-hept-6-enoic acid) was synthesized by the following method.

(Reaction 271-8)




[1949] (E)-7-Phenyl-hept-6-enoic acid was synthesized by operations similar to those in Reaction 26-1 (using NMP as a solvent) using appropriate reagents and starting material.

MS (ESI) m/z = 203 (M-H)-.

[1950] The carboxylic acid reagent used in the synthesis of Compound 1121 ((E)-10-phenyl-dec-9-enoic acid) was synthesized by the following method.

(Reaction 271-9)




[1951] (E)-10-Phenyl-dec-9-enoic acid was synthesized by operations similar to those in Reaction 26-1 (using NMP as a solvent) using appropriate reagents and starting material.

MS (ESI) m/z = 245 (M-H)-.

[1952] The carboxylic acid reagent used in the synthesis of Compound 1122 (6-propylsulfanyl-hexanoic acid) was synthesized by the following method.

(Reaction 271-10)




[1953] A solution of propanethiol (0.609 ml, 6.72 mmol) in anhydrous THF (10 ml) was cooled to -10°C in a nitrogen

atmosphere. 2 M nBuLi (4.03 ml, 8.07 mmol) was added dropwise and the mixture was then stirred for 10 minutes. A solution of ethyl 6-bromohexanoate in anhydrous THF (5 ml) was then added and the mixture was stirred for 40 minutes. The reaction mixture was quenched by adding water and then extracted with ethyl acetate. The organic layer was washed with water and saturated brine and dried over sodium sulfate. The organic layer was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography (hexane:ethyl acetate = 15:1) to give ethyl 6-(propylthio)hexanoate as a colorless oily substance (1.46 g, 100%).

¹H-NMR (300 MHz, CDCl₃) δ 0.98 (t, 3H, J = 7.6 Hz), 1.25 (t, 3H, J = 7.2 Hz), 1.46-1.36 (m, 2H), 1.69-1.54 (m, 6H), 2.30 (t, 2H, J = 7.2 Hz), 2.49 (dd, 4H, J = 7.2, 14.3 Hz), 4.12 (dd, 2H, J = 7.2, 14.1 Hz).

(Reaction 271-11)




[1954] 6-Propylsulfanylhexanoic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material.

¹H-NMR (300 MHz, DMSO-d₆) δ 0.92 (m, 3H), 1.34 (m, 2H), 1.51 (m, 6H), 2.19 (m, 2H), 2.45 (m, 4H).

[1955] The carboxylic acid reagent used in the synthesis of Compound 1123 (8-methoxy-octanoic acid) was synthesized by the following method.

(Reaction 271-12)




[1956] 8-Methoxy-octanoic acid was synthesized by operations similar to those in Reaction 271-3 using appropriate reagents and starting material.

¹H-NMR (300 MHz, CDCl₃) δ 1.39 (m, 6H), 1.69 (m, 4H), 2.13 (s, 1H), 2.42 (m, 2H), 3.33 (s, 3H), 3.39 (m, 2H).

[1957] The carboxylic acid reagent used in the synthesis of Compound 1124 (6-propoxy-hexanoic acid) was synthesized by the following method.


(Reaction 271-13)




[1958] Sodium (354 mg, 15.38 mmol) was added to a solution of 6-bromohexanoic acid (300 mg, 1.54 mmol) in propyl alcohol (15 ml), and the mixture was heated under reflux for two hours. The reaction mixture was cooled to room temperature and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane:methanol = 30:1) to give 6-propoxy-hexanoic acid (209 mg, 78%).

¹H-NMR (300 MHz, CDCl₃) δ 0.88 (m, 3H), 1.37 (m, 2H), 1.62 (m, 6H), 2.44 (m, 2H), 3.52 (m, 4H).

[1959] The carboxylic acid reagent used in the synthesis of Compound 1126 (4-propyl-decanoic acid) was synthesized by the following method.

(Reaction 271-14)

10 **[1960]** 4-Hydroxy-decanoic acid was synthesized by operations similar to those in Reaction 95-18 using appropriate reagents and starting material. This was used in the next step without complete purification.


(Reaction 271-15)

20 **[1961]** PCC (1.2 g, 2.71 mmol) was added to a solution of 4-hydroxydecanoic acid (830 mg, 4.4 mmol) in dichloromethane (30 ml), and the mixture was stirred at room temperature for five hours. The reaction mixture was adjusted to pH 1 by adding 1 N hydrochloric acid and then extracted with ethyl acetate. The organic layer was washed with water and saturated brine and then concentrated under reduced pressure. The resulting residue was then dissolved in ethanol (15 ml). Five drops of sulfuric acid were added and the mixture was stirred at 80°C for 18 hours. The reaction mixture was

25 concentrated under reduced pressure, and the resulting residue was purified by silica gel column chromatography (hexane:ethyl acetate = 2:1) to give ethyl 4-oxodecanoate (440 mg, 46% in two steps).

¹H-NMR (300 MHz, CDCl₃) δ 0.86 (t, 3H, J = 7.2 Hz), 1.26 (m, 9H), 1.57 (m, 2H), 2.43 (t, 2H, J = 7.2 Hz),


[1962] 2.56 (m, 2H), 2.70 (m, 2H), 4.11 (dt, 2H, J = 7.2, 7.2 Hz).

(Reaction 271-16)

50 **[1963]** 4-Propyl-decanoic acid was synthesized by operations similar to those in Reaction 191-14, Reaction 95-18 and Reaction 18-2 using appropriate reagents and starting material.

¹H-NMR (300 MHz, CDCl₃) δ 0.88 (m, 6H), 1.27 (m, 15H), 2.38 (m, 2H), 2.61 (m, 2H), 8.91 (br, 1H).


[1964] The carboxylic acid reagent used in the synthesis of Compound 1127 (2-propyl-benzofuran-6-carboxylic acid) was synthesized by the following method.

(Reaction 271-17)

[1965] 2-Propyl-benzofuran-6-carboxylic acid was synthesized by operations similar to those in Reaction 95-10 (using PdCl₂(PPh₃)₂ as a catalyst) and Reaction 95-18 using appropriate reagents and starting material.

¹H-NMR (300 MHz, CDCl₃) δ 1.02 (m, 3H), 1.80 (m, 2H), 2.79 (m, 2H), 6.46 (s, 1H), 7.53 (m, 1H), 7.96 (m, 1H), 8.16 (s, 1H).

[1966] The carboxylic acid reagent used in the synthesis of Compound 1128 (3-methoxy-4-pentyl-benzoic acid) was synthesized by the following method.


(Reaction 271-18)

[1967] 3-Methoxy-4-pentyl-benzoic acid was synthesized by operations similar to those in Reaction 26-4, Reaction 95-10 (using PdCl₂(PPh₃)₂ as a catalyst), Reaction 122-2 and Reaction 95-18 using appropriate reagents and starting material.

MS (ESI) m/z = 223 (M+H)⁺.


[1968] A mixture of the carboxylic acid reagent used in the synthesis of the compound 1129 (4-([1,1,2,2,2-²H₅]ethyl)-cyclohex-3-enecarboxylic acid) and the carboxylic acid reagent used in the synthesis of the compound 1130 (4-([1,1,2,2,2-²H₅]ethyl)-[4-²H]-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 271-19)



[1969] A solution of iodo-ethane-d₅ (3.00 g, 18.6 mmol) and triphenylphosphine (14.6 mg, 55.8 mmol) in toluene (15 ml) was stirred at 110°C for 21 hours. The reaction mixture was filtered, and the solid was washed with toluene and dried to give [1,1,2,2,2-²H₅]ethyl-triphenylphosphonium iodide as a white solid (7.85 g, 100%). This was used in the next reaction without complete purification.

(Reaction 271-20)




[1970] 4-([1,1,2,2,2-²H₅]Ethyl)-cyclohex-3-enecarboxylic acid and 4-([1,1,2,2,2-²H₅] ethyl)-[4-²H]-cyclohexanecarboxylic acid were synthesized as a mixture by operations similar to those in Reaction 101-1, Reaction 18-2 and Reaction 95-18 using appropriate reagents and starting material. This was used in the next step without complete purification.

Example 272

12-(8-((E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid (Compound 1131) and 12-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid ethyl ester (Compound 1132)

[1971]

(Reaction 272-1)



[1972] 12-(8-((E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid

MS (ESI) m/z = 618 (M+H)+

and

12-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid ethyl ester

MS (ESI) m/z = 646 (M+H)+


were synthesized by operations similar to those in Reaction 269-1 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.

Example 273

1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea trifluoroacetate (Compound 1133) and 1-(3,5-dimethyl-4-((Z)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea trifluoroacetate (Compound 1134)

[1973]

(Reaction 273-1)



[1974] 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea trifluoroacetate
MS (ESI) m/z = 566 (M+H)⁺
and

1-(3,5-dimethyl-4-((Z)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea trifluoroacetate

MS (ESI) m/z = 566 (M+H)⁺


were synthesized by operations similar to those in Reaction 269-1 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.

Example 274

1-[3,5-Dimethyl-4-(2-{4-oxo-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea (Compound 1135)

[1975]

(Reaction 274-1)



[1976] 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[3-(4,4,5,5,5-pentafluoropentyloxy)-phenyl]-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1, Reaction 89-2 (using KOCN) and Reaction 184-1 using appropriate reagents and starting material.


MS (ESI) m/z = 674 (M+H)⁺.

Example 275

1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 1136)

[1977]

(Reaction 275-1)




[1978] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1, Reaction 89-2 (using KOCN) and Reaction 184-1 using appropriate reagents and starting material.

MS (ESI) m/z = 616 (M+H)+.

[1979] The carboxylic acid reagent used in the synthesis of Compound 1136 (3'-propyl-biphenyl-3-carboxylic acid) was synthesized by the following method.

(Reaction 275-2)



[1980] 3'-Propyl-biphenyl-3-carboxylic acid was synthesized by operations similar to those in Reaction 259-2, Reaction 191-14, Reaction 18-2 and Reaction 95-18 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.36 (t, 1H, $J = 1.5$ Hz), 8.10 (dt, 1H, $J = 7.6, 1.5$ Hz), 7.85 (dt, 1H, $J = 7.6, 1.5$ Hz), 7.56 (t, 1H, $J = 7.6$ Hz), 7.46 (dt, 1H, $J = 7.3, 1.5$ Hz), 7.45 (d, 1H, $J = 7.3$ Hz), 7.39 (t, 1H, $J = 7.3$ Hz), 7.23 (dt, 1H, $J = 7.3, 1.5$ Hz), 2.68 (t, 2H, $J = 7.6$ Hz), 1.71 (m, 2H), 0.99 (t, 3H, $J = 7.6$ Hz).

Example 276

1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4-trimethylsilanyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 1137)

[1981]


(Reaction 276-1)



[1982] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4-trimethylsilylanyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1 (using LiOH), Reaction 4-1 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.
MS (ESI) $m/z = 576$ (M+H)+.

[1983] The carboxylic acid reagent used in the synthesis of Compound 1137 (4-trimethylsilylanyl-cyclohexanecarboxylic acid) was synthesized by the following method.


(Reaction 276-2)



[1984] 1-Bromo-4-trimethylsilyl-benzene (0.426 ml, 2.18 mmol) was dissolved in THF (20 ml), and n -butyllithium (1.59 M solution in n -hexane, 1.51 ml, 1.40 mmol) was added dropwise at -78°C . After stirring for 20 minutes, crushed dry ice (excess) was added. The reaction solution was stirred at room temperature for one hour, and 1 M hydrochloric acid and water were then added, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n -hexane/ethyl acetate) to give 4-trimethylsilylanyl-benzoic acid (363 mg, 86%).

$^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 12.93 (1H, br s), 7.91 (2H, d, $J = 8.3$ Hz), 7.65 (2H, d, $J = 7.8$ Hz), 0.27 (9H, t, $J = 3.4$ Hz).

(Reaction 276-3)



[1985] 4-Trimethylsilylanyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 193-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.73-2.66 (0.6H, m), 2.33 (0.4H, tt, $J = 11.0, 3.7$ Hz), 2.15-0.58 (8H, m), -0.06 (9H, s) (cis : trans = ca 6 : 4).

[1986] The example compounds shown below were synthesized by operations similar to those in Reaction 276-1 using appropriate reagents and starting materials.

(Compounds 1138 to 1142)


[1987]

[Table 166]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1138		LCMS-A-1	2.44	640 (M+H)+
1139		LCMS-A-1	2.47	570 (M+H)+
1140		LCMS-A-1	2.94	604 (M+H)+
1141		LCMS-A-1	2.66	576 (M+H)+
1142		LCMS-A-1	2.31	534 (M+H)+

[1988] The carboxylic acid reagent used in the synthesis of Compound 1138 (3,5-bis-trifluoromethyl-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 276-4)



[1989] 3,5-Bis-trifluoromethyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 193-3 using appropriate reagents and starting material.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.51-2.41 (1H, m), 2.35-2.27 (2H, m), 2.24-2.12 (3H, m), 1.52-1.42 (2H, m), 1.41-1.30 (1H, m).

Example 277

1-(4-{2-[2-((1S,3R)-3-Hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1143)

[1990]

(Reaction 277-1)




[1991] 1-(4-{2-[2-((1S,3R)-3-Hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1, Reaction 5-3 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.

MS (ESI) m/z = 588 ($\text{M}+\text{H}$) $^+$.

[1992] The carboxylic acid reagent used in the synthesis of Compound 1143 (3-hexyl-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 277-2)



[1993] 3-Hexyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 101-1, Reaction 95-18 and Reaction 193-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.84-2.03 (22H, m), 2.32 (0.6H, tt, $J = 11.6, 2.8$ Hz), 2.67-2.68 (0.4H, m) (cis : trans = 3 : 2).

[1994] The example compounds shown below were synthesized by operations similar to those in Reaction 277-1 using appropriate reagents and starting materials.

(Compounds 1144 to 1149)

[1995]

[Table 167]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1144		LCMS-C-1	2.93	568 (M+H)+
1145		LCMS-C-1	2.93	560 (M+H)+
1146		LCMS-C-1	3.03	574 (M+H)+

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1147		LCMS-F-1	0.98	638 (M+H)+
1148		LCMS-F-1	0.97	538 (M+H)+
1149		LCMS-F-1	0.95	535 (M+H)+


[1996] The carboxylic acid reagent used in the synthesis of Compound 1144 (3-(3-methyl-butyl)-benzoic acid) was synthesized by the following method.

(Reaction 277-3)



[1997] 3-(3-Methyl-butyl)-benzoic acid was synthesized by operations similar to those in Reaction 101-1, Reaction 95-18 and Reaction 18-2 using appropriate reagents and starting material.
MS (ESI) m/z = 191 (M-H)-.


[1998] The carboxylic acid reagent used in the synthesis of Compound 1145 (3-butyl-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 277-4)

[1999] 3-Butyl-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 193-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.84-2.03 (18H, m), 2.33 (0.6H, m), 2.68 (0.4H, m) (cis : trans = 3 : 2).


[2000] The carboxylic acid reagent used in the synthesis of Compound 1146 (3-(3-methyl-butyl)-cyclohexanecarboxylic acid) was synthesized by the following method.

(Reaction 277-5)

[2001] 3-(3-Methyl-butyl)-cyclohexanecarboxylic acid was synthesized by operations similar to those in Reaction 193-3 using appropriate reagents and starting material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.81-2.04 (20H, m), 2.33 (0.66H, tt, $J = 12.0, 3.2$ Hz), 2.67-2.68 (0.33H, m) (cis : trans = 2 : 1).


[2002] The carboxylic acid reagent used in the synthesis of Compound 1147 (4,4,10,10,10-pentafluoro-decanoic acid) was synthesized by the following method.

(Reaction 277-6)

[2003] One piece of I_2 was added to a solution of magnesium (204 mg, 8.40 mmol) in THF (5 ml) in a nitrogen atmosphere, and the reaction mixture was stirred at 45°C for 20 minutes. A solution of 6-bromo-1,1,1-trifluorohexane (1.53 g, 7.00 mmol) in THF (2 ml) was added and the reaction mixture was stirred at 45°C for one hour to give Compound **277m** (0.875 M solution in THF).

[2004] This Compound **277m** (0.875 M solution in THF, 5.71 ml, 5.00 mmol) was added dropwise to succinylethyl chloride (1.00 g, 5.00 mmol) and CuI (57.9 mg, 304 μmol) in THF (17 ml) at 0°C . The reaction mixture was stirred at 0°C for 30 minutes. The reaction mixture was quenched with a saturated aqueous ammonium chloride solution and diluted with ethyl acetate. The organic layer was then washed with a saturated aqueous sodium bicarbonate solution, water and saturated brine, and then dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ethyl acetate = 100/0 $\rightarrow$ 80/20) to give 10,10,10-trifluoro-4-oxo-decanoic acid ethyl ester (926 mg, 69%).

MS (ESI) $m/z = 269$ ($\text{M}+\text{H}$) $^+$.

(Reaction 277-7)

[2005] 4,4,10,10,10-Pentafluoro-decanoic acid was synthesized by operations similar to those in Reaction 191-11 and Reaction 95-18 using appropriate reagents and starting material.


¹H-NMR (400 MHz, CDCl₃) δ 1.41-1.45 (2H, m), 1.50-1.64 (4H, m), 1.79-1.92 (2H, m), 2.02-2.25 (4H, m), 2.58-2.62 (2H, m).

Example 278

1-(4-{2-[2-(4-Chloro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1150)

[2006]

(Reaction 278-1)




[2007] 1-(4-{2-[2-(4-Chloro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1, Reaction 4-1 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.
MS (ESI) m/z = 600 (M+H)+.

Example 279

1-(3,5-Dimethyl-4-(2-[4-oxo-2-(1,9,9,9-tetrafluorononyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea (Compound 1151)

[2008]

(Reaction 279-1)




[2009] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(1,9,9,9-tetrafluorononyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1, Reaction 5-3 and Reaction 89-2 (using KOCN) using appropriate reagents and starting material.

MS (ESI) m/z = 620 (M+H)+.

[2010] The carboxylic acid reagent used in the synthesis of Compound 1151 (2,10,10,10-tetrafluoro-decanoic acid) was synthesized by the following method.

(Reaction 279-2)



[2011] 2,10,10,10-Tetrafluoro-decanoic acid was synthesized by operations similar to those in Reaction 26-4, Reaction 257-1 (using KHMDS as a base) and Reaction 215-2 using appropriate reagents and starting material.


¹H-NMR (400 MHz, CDCl₃) δ 1.30-1.40 (6H, m), 1.49-1.57 (4H, m), 1.90-2.13 (4H, m), 4.97 (1H, ddd, J = 48.8, 5.2, 5.2 Hz).

Example 280

1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 1152)

[2012]

(Reaction 280-1)



[2013] 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea was synthesized by operations similar to those in Reaction 269-1 and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) $m/z = 616 (M+H)^+$.

[2014] The example compounds shown below were synthesized by operations similar to those in Reaction 280-1 using appropriate reagents and starting materials.



(Compounds 1153 to 1156)

[2015]

[Table 168]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1153		LCMS-F-1	1.08	594 ($M+H$) ⁺
1154		LCMS-D-1	1.72	566 ($M+H$) ⁺

(continued)

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1155		LCMS-D-1	3.12	650 (M+H)+
1156		LCMS-F-1	1.05	612 (M+H)+

[2016] The carboxylic acid reagent used in the synthesis of Compound 1154 (7-ethylsulfanyl-heptanoic acid) was synthesized by the following method.

(Reaction 280-2)



[2017] 7-Ethylsulfanyl-heptanoic acid was synthesized by operations similar to those in Reaction 271-10 (using NaOEt as a base) and Reaction 95-18 using appropriate reagents and starting material.


¹H-NMR (300 MHz, DMSO-d₆) δ 1.24 (m, 3H), 1.39 (m, 4H), 1.56 (m, 4H), 2.24 (m, 2H), 2.49 (m, 4H), 12.05 (s, 1H).

Example 281

1-{3,5-Dimethyl-4-[(E)-2-(4-oxo-2-[1,1';3',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-1-methyl-urea (Compound 1157)

[2018]

(Reaction 281-1)




[2019] 1-{3,5-Dimethyl-4-[(E)-2-(4-oxo-2-[1,1';3',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-1-methyl-urea was synthesized by operations similar to those in Reaction 10-14, Reaction 10-11, Reaction 10-12 (using ethanol as a solvent) and Reaction 89-2 using appropriate reagents and starting material.

MS (ESI) m/z = 648 (M+H)+.

[2020] The carboxylic acid reagent used in the synthesis of Compound 1157 ([1,1';3',1'']terphenyl-3-carboxylic acid) was synthesized by the following method.

(Reaction 281-2)




[2021] [1,1';3',1'']Terphenyl-3-carboxylic acid was synthesized by operations similar to those in Reaction 259-2 using appropriate reagents and starting material.

MS (ESI) m/z = 275 (M+H)+.

Example 282

8-[(E)-2-[4-(4-Hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1158)

[2022]

(Reaction 282-1)

25 **[2023]** 8-((E)-2-[4-(4-Hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 269-1 using appropriate reagents and starting material.

MS (ESI) m/z = 727 (M+H)+.

[2024] The example compounds shown below were synthesized by operations similar to those in Reaction 282-1 using appropriate reagents and starting materials.

(Compounds 1159 to 1160)

[2025]

[Table 169]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1159		LCMS-B-1	2.16	653 (M+H)+
1160		LCMS-B-1	2.02	631 (M+H)+

Example 283

2-(4-Fluoro-3-trifluoromethoxy-phenyl)-8-((E)-2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1161)

[2026]


(Reaction 283-1)



[2027] 4-Amino-1-((E)-2-[4-(4-hydroxymethyl)piperidine-1-carbonyl]-2,6-dimethylphenyl)-ethanesulfonyl-piperidine-4-carboxylic amide was synthesized by operations similar to those in Reaction 119-1, Reaction 10-14, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.

MS (ESI) $m/z = 479$ (M+H)+.

(Reaction 283-2)



[2028] 2-(4-Fluoro-3-trifluoromethoxy-phenyl)-8-((E)-2-[4-(4-hydroxymethyl)piperidine-1-carbonyl]-2,6-dimethylphenyl)-ethanesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 269-1 using appropriate reagents and starting material.


MS (ESI) $m/z = 667$ (M+H)+.

[2029] The example compound shown below was synthesized by operations similar to those in Reaction 283-2 using appropriate reagents and starting material.

(Compound 1162)

[2030]


[Table 170]

Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1162		LCQ-A-1	2.51	651 (M+H)+

Example 284

2-(2,2-Difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1163)

[2031]

(Reaction 284-1)

[2032] 2-(2,2-Difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethylphenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 269-1 using appropriate reagents and starting material.


MS (ESI) m/z = 645 (M+H)+.

Example 285

8-((E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethylphenyl]-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1164)

[2033]

(Reaction 285-1)



[2034] 8-[(E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(4-fluoro-3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 269-1 using appropriate reagents and starting material.
MS (ESI) m/z = 616 (M+H)+.

[2035] The example compound shown below was synthesized by operations similar to those in Reaction 285-1 using appropriate reagents and starting material.

(Compound 1165)

[2036]

[Table 171]


Compound	Structure	LCMS condition	Retention time(min)	MS (m/z)
1165		LCMS-F-1	0.94	594 (M+H)+

Example 286

8-[(E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]ethenesulfonyl]-2-(4-fluoro-3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1166)


[2037]

(Reaction 286-1)



[2038] 4-Amino-1-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-methylphenyl]-ethenesulfonyl)-piperidine-4-carboxylic amide was synthesized by operations similar to those in Reaction 26-1, Reaction 233-2, Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.
MS (ESI) $m/z = 428$ (M+H) $^+$.

(Reaction 286-2)



[2039] 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethoxy-phenyl)-

1,3,8-triaza-spiro[4.5]dec-1-en-4-one was synthesized by operations similar to those in Reaction 269-1 using appropriate reagents and starting material.

MS (ESI) m/z = 616 (M+H)+.

[2040] The example compounds shown below were synthesized by operations similar to those in Reaction 286-2 using appropriate reagents and starting materials.

(Compound 1167)

[2041]

[Table 172]


Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1167		LCMS-C-1	2.60	594 (M+H)+

Example 287

4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl]-acetic acid (Compound 1168)

[2042]



(Reaction 287-1)



[20433] {4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-acetic acid (Compound 1168) was obtained by operations similar to those in Reaction 95-18 using Compound 476 as a starting material. MS (ESI) m/z = 499 (M+H)⁺.

[2044] The example compounds shown below were obtained by operations similar to those in Reaction 287-1 using appropriate starting compounds.

[Table 173]


Raw material Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
504	1169		LCMS-A-1	1.94	546 (M+H) ⁺
1347	1170		LCMS-C-1	2.52	620 (M+H) ⁺

Example 288

10-(8-((E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid (Compound 1171)

[2045]

(Reaction 288-1)




[2046] 10-(8-((E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid (Compound 1171) was obtained by operations similar to those in Reaction 95-17 (using DMI as a solvent), Reaction 269-1 and Reaction 89-2 (using KOCN as a reagent) using appropriate reagents and starting material. MS (ESI) $m/z = 590$ ($M+H$)⁺.

Example 289

10-(8-((E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic amide (Compound 1172)

[2047]





(Reaction 289-1)

[2048] 10-(8-((E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic amide was obtained by operations similar to those in Reaction 10-14 using Compound 1171 as a starting material.

MS (ESI) m/z = 589 (M+H)⁺.

[2049] The example compounds shown below were obtained by operations similar to those in Reaction 289-1 using appropriate starting compounds.

[Table 174]


Raw material Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1131	1173		LCMS-A-1	2.12	617 (M+H)+
1170	1174		LCMS-A-1	2.10	619 (M+H)+
1109	1175		LCMS-C-1	2.82	645 (M+H)+
1350	1176		LCMS-C-1	2.80	647 (M+H)+

Example 290

8-[2-(2-Amino-5,7-dimethyl-benzoxazol-6-yl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one (Compound 1178)

[2050]


(Reaction 290-1)



[2051] Triethylamine (34.7 μ L, 249 μ mol) and di-tert-butyl dicarbonate (32.6 mg, 149 μ mol) were added to a solution of 8-[2-(5,7-dimethyl-2-oxo-2,3-dihydro-benzoxazol-6-yl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one (25 mg, 49.7 μ mol) in dichloromethane/acetonitrile/DMF (1:1:1) (1.0 mL) at room temperature, and the mixture was stirred at room temperature for three hours. The reaction solution was concentrated under reduced pressure, and the resulting residue was dissolved in methanol (1.0 mL) without purification. Potassium carbonate (34.3 mg, 249 μ mol) was added to the solution, and the mixture was stirred at room temperature for two hours. H₂O (3 mL) was added, followed by extraction with dichloromethane (10 mL) twice. The organic layers were dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by PTLC (CH₂Cl₂-MeOH) to give (2-hydroxy-3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-carbamic acid tert-butyl ester as a yellow substance (7.0 mg, 24%).

MS (ESI) m/z = 577, 579 (M+H)⁺.

(Reaction 290-2)



[2052] A mixed solution of (2-hydroxy-3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-carbamic acid tert-butyl ester (7.0 mg, 12.1 μ mol) and dichloromethane/TFA (2:1) (750 μ L) was prepared and stirred at room temperature for one hour. The reaction solution was concentrated under reduced pressure, and the resulting residue was dissolved in ethanol (1.00 mL) without purification. Bromocyanide (3.9 mg, 36.3 μ mol) and sodium bicarbonate (6.1 mg, 72.6 μ mol) were added to the solution, and the mixture was stirred at room temperature for five hours. H₂O (2 mL) were added, followed by extraction with ethyl acetate (10 mL) twice. The organic layers were dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by PTLC (CH₂Cl₂-MeOH-DMF) to give 8-[2-(2-amino-5,7-dimethyl-benzoxazol-6-yl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one as a yellow substance (0.7 mg, 9%).

¹H-NMR (400 MHz, CDCl₃) δ 0.87 (3H, d, J = 8.0 Hz), 0.85-1.01 (2H, m), 1.20-1.52 (5H, m), 1.68-1.78 (4H, br-m),


1.82-1.90 (2H, br-m), 2.24 (1H, tt, J = 3.6, 12.0 Hz), 2.31 (3H, s), 2.33 (3H, s), 2.90-3.06 (2H, br-m), 3.11-3.19 (2H, br-m), 3.20-3.38 (2H, br-m), 3.60-3.68 (2H, br-m), 6.87 (1H, s), 7.24 (2H, s), 10.83 (1H, s).

Example 291

2-Cyclohexyl-8-{2-[2,6-dimethyl-4-(2-oxo-oxazolidine-3-carbonyl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1179)

[2053]

(Reaction 291-1)



[2054] 2-Cyclohexyl-8-{2-[2,6-dimethyl-4-(2-oxo-oxazolidine-3-carbonyl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1179) was obtained by operations similar to those in Reaction 122-2 using Compound as a starting material.

MS (ESI) m/z = 545 (M+H)+.

[2055] The example compounds shown below were obtained by operations similar to those in Reaction 291-1 using appropriate solvents (acetonitrile or methanol or an acetonitrile-methanol mixed solution) and starting compounds.

[Table 175]


Raw material Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
998	1180		LCMS-D-1	2.02	533 (M+H)+
992	1181		LCMS-D-1	2.77	627 (M+H)+
1001	1182		LCMS-D-1	1.93	545 (M+H)+
1002	1183		LCMS-D-1	1.72	607 (M+H)+
1010	1184		LCMS-D-1	2.31	731 (M+H)+

Example 292

2-(3,4-Dichloro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methylpiperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1185)

[2056]


(Reaction 292-1)



[2057] 4-Amino-1-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-piperidine-4-carboxylic amide was obtained by operations similar to those in Reaction 233-3 and Reaction 233-4 using appropriate reagents and starting material.

MS (ESI) m/z = 479 (M+H)+.


(Reaction 292-2)



[2058] Di(2-pyridyl) thionocarbonate (0.97 g, 4.2 mmol) was added to a solution of 4-amino-1-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-piperidine-4-carboxylic amide (1.82 g, 3.8 mmol) in THF (7.6 ml), and the mixture was stirred at 50°C for one hour. The reaction mixture was purified by column chromatography (amine-loaded silica gel, dichloromethane/methanol = 99:1 -> 88:12) to give 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-thioxo-1,3,8-triaza-spiro[4.5]decan-4-one as a colorless solid (1.55 g, 78%).

MS (ESI) m/z = 521 (M+H)+.

(Reaction 292-3)



[2059] A solution of 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-thioxo-1,3,8-triaza-spiro[4.5]decan-4-one (25 mg, 0.048 mmol), 3,4-dichlorophenylboronic acid (27.5 mg, 0.144 mmol), palladium tetrakis(triphenyl)phosphine (11.1 mg, 0.0096 mmol) and CuTC (36.8 mg, 0.192 mmol) in NMP (0.1 mL) was

heated with stirring at 80°C for 30 minutes in a nitrogen atmosphere. After cooling to room temperature, N-acetylcysteine (33 mg, 0.2 mmol) was added to the reaction mixture. The reaction mixture was purified by silica gel column chromatography (NH silica gel, methylene chloride:methanol = 100:0 → 90:10) to give 2-(3,4-dichloro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a white solid (13.4 mg, 44%).

MS (ESI) m/z = 633 (M+H)+.

[2060] The example compounds shown below were obtained by operations similar to those in Reaction 292-3 using appropriate starting compounds.







(Compounds 1186 to Compound 1238)

[2061]







[Table 176]

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1186		LCMS-F-2	0.72	617 (M+H)+
1187		LCMS-F-2	0.63	604 (M+H)+
1188		LCMS-F-2	0.79	667 (M+H)+
1189		LCMS-F-2	0.77	691 (M+H)+
1190		LCMS-F-2	0.76	691 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1191		LCMS-F-2	0.84	739 (M+H)+
1192		LCMS-F-2	0.69	611 (M+H)+
1193		LCMS-F-2	0.78	647 (M+H)+
1194		LCMS-F-2	0.61	609 (M+H)+
1195		LCMS-F-2	0.79	683 (M+H)+
1196		LCMS-F-2	0.78	683 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1197		LCMS-F-2	0.84	759 (M+H)+
1198		LCMS-G-1	1.11	649 (M+H)+
1199		LCMS-G-1	1.10	763 (M+H)+
1200		LCMS-G-1	1.10	731 (M+H)+
1201		LCMS-G-1	1.13	729 (M+H)+
1202		LCMS-G-1	1.15	809 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1203		LCMS-G-1	1.17	781 (M+H)+
1204		LCMS-G-1	1.15	759 (M+H)+
1205		LCMS-G-1	1.13	739 (M+H)+
1206		LCMS-G-1	1.09	629 (M+H)+
1207		LCMS-G-1	1.17	759 (M+H)+
1208		LCMS-F-2	0.57	604 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1209		LCMS-F-2	0.86	759 (M+H)+
1210		LCMS-F-2	0.83	725 (M+H)+
1211		LCMS-F-2	0.80	664 (M+H)+
1212		LCMS-F-2	0.81	695 (M+H)+
1213		LCMS-F-2	0.74	681 (M+H)+
1214		LCMS-F-2	0.79	663 (M+H)+
1215		LCMS-F-2	0.72	659 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1216		LCMS-G-1	1.09	697 (M+H)+
1217		LCMS-F-2	0.72	633 (M+H)+
1218		LCMS-F-2	0.79	709 (M+H)+
1219		LCMS-F-2	0.72	633 (N+H)+
1220		LCMS-F-2	0.86	714 (M+H)+
1221		LCMS-F-2	0.76	682 (M+H)+
1222		LCMS-F-2	0.80	698 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1223		LCMS-F-2	0.83	763 (M+H)+
1224		LCMS-F-2	0.71	632 (M+H)+
1225		LCMS-F-2	0.74	648 (M+H)+
1226		LCMS-F-2	0.83	763 (M+H)+
1227		LCMS-F-2	0.84	725 (M+H)+
1228		LCMS-F-2	0.87	781 (M+H)+
1229		LCMS-F-2	0.79	729 (M+H)+

(continued)


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1230		LCMS-F-2	0.84	747 (M+H)+
1231		LCMS-F-2	0.85	681 (M+H)+
1232		LCMS-F-2	0.87	775 (M+H)+
1233		LCMS-F-2	0.90	714 (M+H)+
1234		LCMS-G-1	1.10	651 (M+H)+
1235		LCMS-G-1	1.13	649 (M+H)+
1236		LCMS-G-1	1.15	669 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1237		LCMS-C-1	2.63	593 (M+H)+
1238		LCMS-C-1	2.58	609 (M+H)+

[2062] The arylboronic acid reagent used in the synthesis of Compound 1195 (4-pentafluoroethylphenylboronic acid) was synthesized by the following method.


(Reaction 292-4)



[2063] A solution of 4-bromoiodobenzene (500 mg, 1.77 mmol), trimethylsilylpentafluoroethane (679 mg, 3.53 mmol), copper iodide (672 mg, 3.53 mmol) and potassium fluoride (205 mg, 3.53 mmol) in N-methylpyrrolidone (1.0 mL) was heated with stirring at 100°C for three hours in a sealed reaction vessel. After cooling to room temperature, the reaction mixture was purified by silica gel column chromatography (hexane:ethyl acetate = 20:1) to give 1-bromo-4-pentafluoroethylbenzene as a colorless liquid (253 mg, 52%).

¹H-NMR (270 MHz, CDCl₃) δ 7.46 (2H, d, J = 8.6 Hz), 7.65 (2H, d, J = 8.6 Hz).

(Reaction 292-5)




[2064] A 1.5 M solution of n-butyllithium in tetrahydrofuran (0.57 mL) was added to a solution of 1-bromo-4-pentafluoroethylbenzene (180 mg, 0.65 mmol) in diethyl ether (1.0 mL) at -78°C, and the mixture was stirred for 20 minutes. Thereafter, trimethyl borate (101 mg, 3.28 mmol) was added and the mixture was stirred at -78°C for 10-minutes and at room temperature for 30 minutes. 6 N aqueous hydrochloric acid (200 μL) was added to the reaction mixture, and the reaction was terminated. The mixture was then purified by silica gel column chromatography to give 4-pentafluoroethylphenylboronic acid as a white solid (111 mg, 71%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.87 (2H, d, $J = 8.0$ Hz), 7.64 (2H, d, $J = 8.0$ Hz), 4.61 (s, 2H).

[2065] The arylboronic acid reagent used in the synthesis of Compound 1196 (3-pentafluoroethylphenylboronic acid) was synthesized by the following method.

(Reaction 292-6)




[2066] 3-Pentafluoroethylphenylboronic acid was obtained by operations similar to those in Reaction 292-4 and Reaction 292-5 using 1-bromo-3-iodobenzene as a starting material.

MS (ESI) $m/z = 239$ (M-H)-.

[2067] The arylboronic acid reagent used in the synthesis of Compound 1197 (4-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-3-trifluoromethylphenylboronic acid) was synthesized by the following method.


(Reaction 292-7)



[2068] Potassium tert-butoxide (236 mg, 2.1 mmol) was added to a solution of 5-bromo-2-fluorobenzotrifluoride (485 mg, 2.0 mmol) and 2-trifluoromethyl-2-propanol (0.24 mL, 2.2 mmol) in DMI (0.5 mL) at room temperature, and the mixture was stirred at 100°C for two hours. The reaction solution was purified by silica gel column chromatography to give 4-bromo-1-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-2-trifluoromethylbenzene (349 mg, 50%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.73 (1H, d, $J = 2.4$ Hz), 7.59 (1H, dd, $J = 2.4, 8.9$ Hz), 7.14 (1H, d, $J = 8.9$ Hz), 1.53 (s, 6H).

(Reaction 292-8)



















[2069] 4-(2,2,2-Trifluoro-1,1-dimethyl-ethoxy)-3-trifluoromethylphenylboronic acid was obtained by operations similar to those in Reaction 292-5 using 4-bromo-1-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-2-trifluoromethylbenzene as a starting material.







MS (ESI) $m/z = 315$ (M-H)-.

[2070] The arylboronic acid reagents shown below were synthesized by operations similar to those in Reaction 292-7 and Reaction 292-5 using appropriate starting compounds and used in the synthesis of the compounds in Table 176.

[Table 177]







Target Compound	Raw material	Arylboronic acid structure	MS (m/z)
1199			319 (M-H)-
1200			287 (M-H)-
1201			285 (M-H)-
1202			365 (M-H)-
1203			337 (M-H)-
1204			315 (M-H)-
1216			253 (M-H)-
1220			270 (M-H)-

(continued)


Target Compound	Raw material	Arylboronic acid structure	MS (m/z)
1221			238 (M-H)-
1222			254 (M-H)-
1232			331 (M-H)-

[2071] The arylboronic acid reagents shown below were synthesized by operations similar to those in Reaction 292-5 using appropriate starting compounds and used in the synthesis of the compounds in Table 176.

[Table 178]

Target Compound	Raw material	Arylboronic acid structure	MS (m/z)
1214			219 (M-H)-
1217			189 (M-H)-
1219			189 (M-H)-


[2072] The arylboronic acid reagent used in the synthesis of Compound 1229 (4-chloro-3-(2,2,3,3-tetrafluoropropoxy)-phenylboronic acid) was synthesized by the following method.

(Reaction 292-9)

[2073] 4-Bromo-1-chloro-2-(2,2,3,3-tetrafluoro-propoxy)-benzene was obtained by operations similar to those in Reaction 292-7 using appropriate starting compound and reagents.

¹H-NMR (400 MHz, CDCl₃) δ 7.27 (1H, d, J = 8.3 Hz), 7.15 (1H, dd, J = 2.0, 8.3 Hz), 7.07 (1H, d, J = 2.0 Hz), 6.15 (1H, dt, J = 5.4, 53.2 Hz), 4.39 (2H, t, J = 11.2 Hz).

(Reaction 292-10)



[2074] A 1.5 M solution of n-butyllithium in tetrahydrofuran (0.99 mL) was added to a solution of 4-bromo-1-chloro-2-(2,2,3,3-tetrafluoro-propoxy)-benzene (434 mg, 1.35 mmol) and triisopropyl borate (382 mg, 2.03 mmol) in anhydrous tetrahydrofuran (2.0 mL) at -78°C, and the mixture was stirred for 10 minutes. The reaction mixture was warmed to room temperature and stirred for 30 minutes. 6 N aqueous hydrochloric acid was then added to the reaction mixture, and the reaction was terminated, followed by extraction with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was treated with a mixed solution of dichloromethane and hexane to give 4-chloro-3-(2,2,3,3-tetrafluoro-propoxy)-phenylboronic acid as a white solid (246 mg, 64%).







MS (ESI) m/z = 285 (M-H)-.

[2075] The arylboronic acid reagents shown below were synthesized by operations similar to those in Reaction 292-7 and Reaction 292-10 using appropriate starting compounds and used in the synthesis of the compounds in Table 176.

[Table 179]


Target Compound	Raw material	Arylboronic acid structure	MS (m/z)
1226			319 (M-H)-
1227			281 (M-H)-
1228			337 (M-H)-

(continued)

Target Compound	Raw material	Arylboronic acid structure	MS (m/z)
1230			303 (M-H)-
1231			236 (M-H)-
1233			270 (M-H)-

[2076] The arylboronic acid reagent used in the synthesis of Compound 1213 (4-(1,1,2,2-tetrafluoro-ethoxy)phenylboronic acid) was synthesized by the following method.

(Reaction 292-11)




[2077] 4-(1,1,2,2-Tetrafluoro-ethoxy)phenylboronic acid was obtained by operations similar to those in Reaction 292-10 using appropriate starting compound and reagents.

MS (ESI) m/z = 237 (M-H)-.

[2078] The arylboronic acid reagent used in the synthesis of Compound 1223 (3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluoromethylphenylboronic acid) was synthesized by the following method.


(Reaction 292-12)



[2079] 4-Bromo-2-(2,2,3,3-tetrafluoro-propoxy)-1-trifluoromethyl-benzene was obtained by operations similar to those in Reaction 292-7 using appropriate starting compound and reagents.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.74 (1H, d, J = 2.4 Hz), 7.65 (1H, dd, J = 2.4, 8.8 Hz), 6.88 (1H, d, J = 8.8 Hz), 6.06 (1H, dt, J = 5.4, 53.2 Hz), 4.40 (2H, t, J = 11.2 Hz).


(Reaction 292-13)



[2083] Toluene was added to a mixture of 5-bromo-2-fluoro-phenol (382 mg, 2.0 mmol), ethanol- d^5 (0.104 mL, 2.0 mmol) and N,N,N',N'-tetramethylazodicarboxamide (465 mg, 2.7 mmol). Tributylphosphine (0.622 mL, 2.5 mmol) was added at 0°C and the mixture was stirred for 14 hours. The reaction solution was purified by silica gel column chromatography to give 4-bromo-2-[1,1,2,2,2- $^2H_5$]ethoxy-1-fluoro-benzene (416 mg, 93%).

^1H -NMR (400 MHz, $CDCl_3$) δ 7.07 (1H, dd, J = 2.1, 7.4 Hz), 7.00 (1H, ddd, J = 2.1, 4.1, 8.4 Hz), 6.94 (1H, dd, J = 8.4, 10.7 Hz).

(Reaction 292-15)



[2084] 4-Bromo-3-[1,1,2,2,2- $^2H_5$]ethoxy-1-fluorophenylboronic acid was obtained by operations similar to those in Reaction 292-5 using 4-bromo-2-[1,1,2,2,2- $^2H_5$]ethoxy-1-fluoro-benzene as a starting material.

MS (ESI) m/z = 188 (M-H)-.


[2085] The arylboronic acid reagents shown below were synthesized by operations similar to those in Reaction 292-14 and Reaction 292-5 using appropriate starting compounds and used in the synthesis of the compounds in Table 176.

[Table 181]

Target Compound	Raw material	Arylboronic acid structure	MS (m/z)
1207			315 (M-H)-
1211			220 (M-H)-
1218			265 (M-H)-
1225			204 (M-H)-

[2086] The arylboronic acid reagent used in the synthesis of Compound 1215 (3-fluoro-4-(3-fluoro-propoxy)phenylboronic acid) was synthesized by the following method.


(Reaction 292-16)



[2087] 4-Bromo-2-fluorophenol (382 mg, 2.0 mmol) was dissolved in DMI (0.5 mL), and potassium tert-butoxide (224 mg, 2.0 mmol) was added at room temperature. 1-Iodo-3-fluoropropane (376 mg, 2.0 mmol) was added to the reaction solution, and the mixture was heated to 60°C and stirred for six hours. The reaction solution was purified by silica gel column chromatography to give 4-bromo-2-fluoro-1-(3-fluoro-propoxy)-benzene (410 mg, 82%).

¹H-NMR (400 MHz, CDCl₃) δ 7.10 (1H, dd, J = 2.1, 7.4 Hz), 7.03 (1H, ddd, J = 2.4, 3.9, 8.6 Hz), 6.95 (1H, dd, J = 8.7, 10.7 Hz), 4.66 (2H, dt, J = 5.6, 46.9 Hz), 4.15 (2H, t, J = 6.1 Hz), 2.20 (2H, ddt, J = 5.8, 5.8, 26.1 Hz).

(Reaction 292-17)




[2088] 3-Fluoro-4-(3-fluoro-propoxy)phenylboronic acid was obtained by operations similar to those in Reaction 292-5 using 4-bromo-2-fluoro-1-(3-fluoro-propoxy)-benzene as a starting material.

MS (ESI) m/z = 215 (M-H)-.

[2089] The arylboronic acid reagent used in the synthesis of Compound 1212 (2,2,3,3-tetrafluoro-2,3-dihydro-benzo[1,4]dioxin-6-yl-boronic acid) was synthesized by the following method.

(Reaction 292-18)



[2090] 2,2,3,3-Tetrafluoro-1,4-benzodioxane (484 mg), bis (pinacolato) diboron (295 mg), [Ir(COD)(OMe)₂]₂ (15.4 mg) and 4,4'-di-tert-butyl-2,2'-dipyridyl (12.5 mg) were mixed. 1,4-Dioxane (0.5 mL) was added in a nitrogen atmosphere and stirred at 100°C for two hours. MeOH (0.5 mL) was added to the reaction solution, and metaperiodic acid (1.06 g) was added in four portions under icecooling. Water was added to the reaction solution, followed by extraction with ethyl acetate and concentration. The resulting mixture was purified by silica gel column chromatography to give 2,2,3,3-tetrafluoro-2,3-dihydro-benzo[1,4]dioxin-6-yl-boronic acid (230 mg, 39%).

MS (ESI) m/z = 251 (M-H)-.


Example 293

2-(4-Fluoro-2,5-dimethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phe-

nyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1239)


[2091]

(Reaction 293-1)



[2092] 8-((E)-2-(4-(4-Fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl)-ethenesulfonyl)-2-thioxo-1,3,8-triaza-spiro[4.5]decan-4-one was obtained by operations similar to those in Reaction 119-1, Reaction 233-3, Reaction 233-4 and Reaction 292-2 using 1-ethenesulfonyl-piperidin-4-one as a starting material. MS (ESI) m/z = 539 (M+H)+.

(Reaction 293-2)



[2093] 2-(4-Fluoro-2,5-dimethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one was obtained by operations similar to those in Reaction using 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-thioxo-1,3,8-triaza-spiro[4.5]decan-4-one as a starting material.

MS (ESI) m/z = 629 (M+H)+.







[2094] The example compounds shown below were obtained by operations similar to those in Reaction 293-2 using appropriate starting compounds.

(Compounds 1240 to 1281)








[2095]

704

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1247		LCMS-F-1	1.01	747 (M+H) ⁺
1248		LCMS-F-1	1.01	749 (M+H) ⁺
1249		LCMS-B-1	2.48	799 (M+H) ⁺
1250		LCMS-B-1	2.41	777 (M+H) ⁺
1251		LCMS-F-1	1.00	781 (M+H) ⁺
1252		LCMS-F-1	1.06	827 (M+H) ⁺

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1253		LCMS-F-1	0.99	700 (M+H)+
1254		LCMS-F-1	1.00	665 (M+H)+
1255		LCMS-F-1	0.93	651 (M+H)+
1256		LCMS-F-1	0.92	651 (M+H)+
1257		LCMS-F-1	1.00	651 (M+H)+
1258		LCMS-F-1	1.01	681 (M+H)+
1259		LCMS-F-1	0.96	665 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1260		LCMS-F-1	0.95	699 (M+H)+
1261		LCMS-F-1	1.04	732 (M+H)+
1262		LCMS-F-1	1.01	716 (M+H)+
1263		LCMS-F-1	1.05	777 (M+H)+
1264		LCMS-F-1	0.95	677 (M+H)+
1265		LCMS-F-1	1.04	777 (M+H)+

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1266		LCMS-F-1	1.04	777 (M+H)+
1267		LCMS-F-1	0.93	647 (M+H)+
1268		LCMS-F-1	0.88	622 (M+H)+
1269		LCMS-F-1	0.98	667 (M+H)+
1270		LCMS-F-1	1.04	765 (M+H)+
1271		LCMS-F-1	1.04	743 (M+H)+
1272		LCMS-F-1	0.95	650 (M+H)+

(continued)


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1273		LCMS-F-1	1.06	793 (M+H)+
1274		LCMS-F-1	1.02	781 (M+H)+
1275		LCMS-G-1	1.11	665 (M+H)+
1276		LCMS-G-1	1.11	669 (M+H)+
1277		LCMS-G-1	1.13	667 (M+H)+
1278		LCMS-G-1	1.16	687 (M+H)+
1279		LCMS-G-1	1.17	783 (M+H)+

(continued)


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1280		LCMS-G-1	1.11	663 (M+H)+
1281		LCMS-F-1	1.03	715 (M+H)+

Example 294

[3-(8-((E)-2-[4-(4-Hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-phenyl]-acetonitrile (Compound 1282)

[2096](Reaction 294-1)

[2097] 8-((E)-2-[4-(4-Hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-thioxo-1,3,8-triaza-spiro[4.5]decan-4-one was obtained by operations similar to those in Reaction 292-2 (using 1,1'-thiocarbonyldiimidazole) using appropriate reagents and starting material.
MS (ESI) m/z = 507 (M+H)+.

(Reaction 294-2)


[2098] [3-(8-((E)-2-[4-(4-Hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-phenyl]-acetonitrile was obtained by operations similar to those in Reaction 292-3 using 8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-thioxo-1,3,8-triaza-spiro[4.5]decan-4-one as a starting material.
MS (ESI) m/z = 590 (M+H)+.

Example 295

8-[(E)-2-[2-Methyl-4-(piperidin-4-yloxy)-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one trifluoroacetate (Compound 1283)

[2099]

(Reaction 295-1)



[2100] 8-[(E)-2-[2-Methyl-4-(piperidin-4-yloxy)-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one trifluoroacetate (Compound 1283) was obtained by operations similar to those in Reaction 4-1 using Compound 602 as a starting material.

MS (ESI) m/z = 593 (M+H)+.

[2101] The example compounds shown below were obtained by operations similar to those in Reaction 295-1 using appropriate starting compounds. Compound 1285 was obtained as a free form by desalination post-treatment

(Compounds 1284 to Compound 1285)

[2102]

[Table 183]


Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
928	1284		LCMS-B-1	1.67	543 (M+H)+
578	1285		LCMS-A-1	1.69	528 (M+H)+

Example 296

8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-pyrrolidin-2-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride (Compound 1286)

[2103]

(Reaction 296-1)



[2104] 8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-pyrrolidin-2-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one dihydrochloride (Compound 1286) was obtained by operations similar to those in Reaction 5-3 using Compound 1033 as a starting material.

MS (ESI) $m/z = 448$ (M+H)+.

[2105] The example compounds shown below were obtained by operations similar to those in Reaction 296-1 using appropriate starting compounds.

(Compounds 1287 to Compound 1288)

[2106]


[Table 184]

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1034	1287		LCMS-B-1	1.44	443 (M+H)+
640	1288		LCMS-F-1	1.05	542 (M+H)+

Example 297

8-[(E)-2-(2,6-Dimethyl-4-methylaminomethyl-phenyl)-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1289)

[2107]

(Reaction 297-1)

10 **[2108]** 8-[(E)-2-(2,6-Dimethyl-4-methylaminomethyl-phenyl)-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1289) was obtained by operations similar to those in Reaction 50-2 (conversion to a free form by post-treatment) using Compound 517 as a starting material.

MS (ESI) $m/z = 487$ (M+H)+.

15 **[2109]** The example compounds shown below were obtained by operations similar to those in Reaction 297-1 using appropriate starting compounds.

(Compounds 1290 to Compound 1292)


[2110]

[Table 185]

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
558	1290		LCMS-D-1	2.27	555 (M+H)+
522	1291		LCMS-D-1	1.93	473 (M+H)+
559	1292		LCMS-D-1	1.77	569 (M+H)+

Example 298

8-[(E)-2-(4-Amino-3-chloro-2-methyl-phenyl)-ethanesulfonyl]-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1293)

[2111](Reaction 298-1)

[2112] 8-[2-(4-Amino-3-chloro-2-methyl-phenyl)-ethanesulfonyl]-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1293) was obtained by operations similar to those in Reaction 12-5 using Compound 954 as a starting material.


MS (ESI) m/z = 495 (M+H)+.

Example 299

N-(1-Acetyl-piperidin-4-yl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide (Compound 1294)

[2113]

(Reaction 299-1)



[2114] N-(1-Acetyl-piperidin-4-yl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide (Compound 1294) was obtained by operations similar to those in Reaction 4-1 and Reaction 12-2 using Compound 604 as a starting material.


MS (ESI) m/z = 676 (M+H)+.

Example 300

8-[2-[4-(4,5-Dihydro-thiazol-2-ylamino)-2-methyl-phenyl]-ethanesulfonyl]-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1295)

[2115]

(Reaction 300-1)




[2116] 8-{2-[4-(4,5-Dihydro-thiazol-2-ylamino)-2-methylphenyl]-ethanesulfonyl}-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1295) was obtained by operations similar to those in Reaction 12-5, Reaction 18-2 and Reaction 177-2 using Compound 953 as a starting material.
 MS (ESI) $m/z = 546 (M+H)^+$.

Example 301

N-[4-(2-{2-[4-(4-Fluoro-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl}-3-methyl-phenyl]-acetamide (Compound 1296)

[2117]


(Reaction 301-1)



[2118] 2-[4-(4-Fluoro-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester was obtained by operations similar to those in Reaction 101-1, Reaction 23-2, Reaction 18-2, Reaction 10-1 and Reaction 189-5 using 4-oxo-cyclohexanecarboxylic acid ethyl ester as a starting material.

¹H-NMR (400 MHz, CDCl₃) δ 0.97-1.05 (1H, m), 1.23-1.35 (4H, m), 1.35-1.50 (3H, m), 1.47 (9H, s), 1.60-1.75 (4H, m), 1.75-1.85 (2H, m), 1.85-1.95 (2H, m), 1.95-2.05 (2H, m), 2.35-2.45 (1H, m), 3.35-3.45 (2H, m), 3.90-4.05 (2H, m), 4.35-4.42 (1H, m), 4.45-4.52 (1H, m), 8.85 (1H, s).

(Reaction 301-2)



[2119] N-[4-(2-[2-[4-(4-Fluoro-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methylphenyl]-acetamide (Compound 1296) was obtained by operations similar to those in Reaction 4-1 and Reaction 5-4 using 2-[4-(4-fluorobutyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester as a starting material.


MS (ESI) m/z = 549 (M+H)+.

Example 302

1-(3,5-Dimethyl-4-{2-[2-(3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 1297)

[2120]

(Reaction 302-1)



[2121] 1-(3,5-Dimethyl-4-{2-[2-(3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea (Compound 1297) was obtained by operations similar to those in Reaction 12-5 and Reaction 89-2 (using KOCN) using Compound 932 as a starting material.

MS (ESI) m/z = 518 (M+H)+.

5

5

5

10



20

25

30

30

35



50


55

Example 305

1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-1-methyl-urea (Compound 1300)

[2126]

(Reaction 305-1)









[2127] 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-1-methyl-urea (Compound 1300) was obtained by operations similar to those in Reaction 89-2 (using KOCN) using Compound 1193 as a starting material.

MS (ESI) m/z = 612 (M+H)+.

[2128] The example compounds shown below were obtained by operations similar to those in Reaction 305-1 using appropriate starting compounds.

(Compounds 1301 to Compound 1312)

[Table 186]


Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1289	1301		LCMS-D-1	1.97	530 (M+H)+
1035	1302		LCMS-F-1	1.01	638 (M+H)+
691	1303		LCMS-F-1	1.01	636 (M+H)+
690	1304		LCMS-F-1	1.00	600 (M+H)+
1016	1305		LCMS-C-1	2.75	596 (M+H)+
1036	1306		LCMS-F-1	0.98	598 (M+H)+

[2129]

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1105	1307		LCMS-C-1	2.70	590 (M+H)+
1106	1308		LCMS-C-1	2.98	689 (M+H)+
1107	1309		LCMS-F-1	0.92	562 (M+H)+
1037	1310		LCMS-A-1	2.10	544 (M+H)+
1287	1311		LCMS-B-1	1.71	486 (M+H)+


(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1288	1312		LCMS-F-1	0.99	585 (M+H)+

Example 306

[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-urea (Compound 1313)

[2130]

(Reaction 306-1)

[2131] [3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-urea (Compound 1313) was obtained by operations similar to those in Reaction 89-2 (using KOCN) and Reaction 122-2 using Compound 1290 as a starting material.

MS (ESI) m/z = 600 (M+H)+.

[2132] The example compound shown below was obtained by operations similar to those in Reaction 306-1 using an appropriate starting compound.

(Compound 1314)

[2133]

[Table 187]


Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1291	1314		LCMS-D-1	1.78	518 (M+H)+

Example 307

1-(4-{(E)-2-[2-(11-Amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1315)

[2134]

(Reaction 307-1)



[2135] 1-(4-((E)-2-[2-(11-Amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1315) was obtained by operations similar to those in Reaction 4-1 using Compound 1209 as a starting material.


MS (ESI) m/z = 589 (M+H)+.

Example 308

1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[5-(propane-1-sulfinyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea (Compound 1316)

[2136]

(Reaction 308-1)



[2137] 30% aqueous hydrogen peroxide (0.013 ml) was added to a mixed solution of (E)-1-(3,5-dimethyl-4-(2-((4-oxo-2-(5-(propylthio)pentyl)-1,3,8-triazaspiro[4.5]dec-1-en-8-yl)sulfonyl)vinyl)phenyl)-1-methyl-urea (55 mg) and molybdenum(IV) dichloride dioxide (3 mg) in acetone (1.5 ml)-water (0.5 ml), and the mixture was stirred at room temperature for five minutes in a nitrogen stream. The reaction mixture was quenched with a saturated aqueous sodium bicarbonate solution and then extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (silica gel, CH_2Cl_2 -MeOH) to give 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[5-(propane-1-sulfinyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea (64 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.77 (s, 1H), 7.55 (d, 1H, J = 15.6 Hz), 7.02 (s, 2H), 6.38 (d, 1H, J = 15.6 Hz), 4.50 (s, 2H), 3.71-3.64 (m, 2H), 3.41-3.32 (m, 2H), 3.26 (s, 3H), 2.76-2.54 (m, 4H), 2.52-2.43 (m, 2H), 2.38 (s, 6H), 2.00-1.61 (m, 12H), 1.09 (t, 3H, J = 7.4 Hz).

MS (ESI) m/z = 580 (M+H)+.

[2138] The example compound shown below was obtained by operations similar to those in Reaction 308-1 using an appropriate starting compound.

(Compound 1317)

5

10

15

20

25

30

35

40


45

50

55

[2139]

[Table 188]


Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1154	1317		LCMS-D-1	1.42	582 (M+H)+

Example 309

1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[5-(propane-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea (Compound 1318)

[2140]

(Reaction 309-1)



[2141] 30% aqueous hydrogen peroxide (0.047 ml) was added to a mixed solution of (E)-1-(3,5-dimethyl-4-(2-((4-oxo-2-(5-(propylthio)pentyl)-1,3,8-triazaspiro[4.5]dec-1-en-8-yl)sulfonyl)vinyl)phenyl)-1-methylurea (61 mg) and molybdenum(IV) dichloride dioxide (6.5 mg) in acetonitrile (1 ml), and the mixture was stirred at room temperature for two hours in a nitrogen stream. The reaction mixture was quenched with a saturated aqueous sodium bicarbonate solution and then extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (silica gel, CH_2Cl_2 -MeOH) to give 1-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[5-(propane-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea (64 mg).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.96 (s, 1H), 7.54 (d, 1H, $J = 15.6$ Hz), 7.02 (s, 2H), 6.39 (d, 1H, $J = 15.6$ Hz), 4.68 (s, 2H), 3.68-3.61 (m, 2H), 3.45-3.37 (m, 2H), 3.25 (s, 3H), 2.98-2.91 (m, 4H), 2.48 (t, 2H, $J = 7.4$ Hz), 2.37 (s, 6H), 1.98-1.84 (m, 6H), 1.81-1.70 (m, 4H), 1.66-1.57 (m, 2H), 1.09 (t, 3H, $J = 7.4$ Hz).



MS (ESI) $m/z = 596$ ($\text{M}+\text{H}$) $^+$.

[2142] The example compounds shown below were obtained by operations similar to those in Reaction 309-1 using appropriate starting compounds.

(Compounds 1319 to Compound 1320)

[2143]


[Table 189]

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1022	1319		LCMS-D-1	1.52	596 (M+H)+
1154	1320		LCMS-D-1	1.49	598 (M+H)+

Example 310


1-(4-((E)-2-[2-(9,9-Difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1321)

[2144]

(Reaction 310-1)

[2145] NMO (22.0 mg, 0.192 mmol), Molecular Sieves 4A (25.0 mg) and TPAP (0.700 mg, 0.00213 mmol) were added to a solution of 1-(4-((E)-2-[2-(9-hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3,5-dimethyl-phenyl)-1-methyl-urea (24.0 mg, 0.0426 mmol) in CH_2Cl_2 (850 μl) at room temperature. The mixture was stirred at room temperature for one hour and then filtered through celite, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography to give 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(9-oxo-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (17.0 mg, 71%).

MS (ESI) m/z = 562 ($M+H$) $^+$.

(Reaction 310-2)


[2146] 1-(4-((E)-2-[2-(9,9-Difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1321) was obtained by operations similar to those in Reaction 191-11 using 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(9-oxo-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea as a starting material.

MS (ESI) m/z = 582 ($M+H$) $^+$.

Example 311

1-(4-{2-[2-(9-Hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1322)

[2147]

(Reaction 311-1)


[2148] 1-(4-{2-[2-(9-Hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethylphenyl)-1-methyl-urea (Compound 1322) was obtained by operations similar to those in Reaction 184-1 using Compound 1210 as a starting material and acetonitrile as a solvent.

MS (ESI) m/z = 564 (M+H)⁺.

Example 312

1-(4-{2-[2-(9,9-Difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1323)

[2149]

(Reaction 312-1)


[2150] 1-(4-{2-[2-(9,9-Difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1323) was obtained by operations similar to those in Reaction 310-1 and Reaction 191-11 using Compound 1260 as a starting material.
MS (ESI) m/z = 584 (M+H)+.

Example 313

1-(4-{2-[2-(9-Amino-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1324)

[2151]

(Reaction 313-1)




[2152] 1-(4-{2-[2-(9-Amino-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethylphenyl)-1-methyl-urea (Compound 1324) was obtained by operations similar to those in Reaction 310-1 and Reaction 80-1 (using NaBH_3CN as a reducing agent and methanol as a solvent) using Compound 1260 as a starting material.
MS (ESI) m/z = 563 (M+H)+.

Example 314

1-(4-{(E)-2-[2-(9-Fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethylphenyl)-1-methyl-urea (Compound 1325) and 1-(4-{(Z)-2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1326)

[2153]

(Reaction 314-1)

[2154] 1-(4-((E)-2-[2-(9-Fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethylphenyl)-1-methyl-urea (Compound 1325)

MS (ESI) m/z = 564 (M+H)⁺

and 1-(4-((Z)-2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethylphenyl)-1-methyl-urea (Compound 1326)


MS (ESI) m/z = 564 (M+H)⁺

were obtained by operations similar to those in Reaction 191-11 using Compound 1260 as a starting material.

Example 315

1-(4-{2-[2-(9-Fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1327)

[2155]

(Reaction 315-1)


[2156] 1-(4-{2-[2-(9-Fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1327) was obtained by operations similar to those in Reaction 191-11 using Compound 1260 as a starting material.

MS (ESI) m/z = 566 (M+H)⁺.

Example 316

8-{2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1328)

[2157]

(Reaction 316-1)

[2158] 8-{2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1328) was obtained by operations similar to those in Reaction 122-2 using Compound 1003 as a starting material.

MS (ESI) m/z = 692 (M+H)⁺.

[2159] The example compounds shown below were obtained by operations similar to those in Reaction 316-1 using appropriate solvents (acetonitrile or methanol or an acetonitrile-methanol mixed solution) and starting compounds.

(Compounds 1329 to Compound 1364)

[Table 190]

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1164	1329		LCMS-F-1	0.95	618 (M+H)+
1165	1330		LCMS-F-1	0.93	596 (M+H)+
1167	1331		LCMS-F-1	0.94	596 (M+H)+
1166	1332		LCMS-F-1	0.96	618 (M+H)+
1160	1333		LCMS-F-1	0.93	633 (M+H)+
1159	1334		LCMS-F-1	0.95	655 (M+H)+

[2160]

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1099	1335		LCMS-F-1	1.01	669 (M+H) ⁺
1163	1336		LCMS-F-1	1.00	647 (M+H) ⁺
1102	1337		LCMS-F-1	0.96	627 (M+H) ⁺
1019	1338		LCMS-F-1	0.98	624 (M+H) ⁺
1020	1339		LCMS-F-1	0.97	624 (M+H) ⁺

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1133	1340		LCMS-F-1	1.01	568 (M+H) ⁺
1112	1341		LCMS-F-1	1.07	716 (M+H) ⁺
1021	1342		LCMS-F-1	1.06	568 (M+H) ⁺
1108	1343		LCMS-F-1	1.01	546 (M+H) ⁺
1115	1344		LCMS-D-1	1.76	550 (M+H) ⁺
1116	1345		LCMS-D-1	2.82	636 (M+H) ⁺

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1117	1346		LCMS-D-1	2.73	636 (M+H) ⁺
1132	1347		LCMS-C-1	3.02	648 (M+H) ⁺
1121	1348		LCMS-C-1	3.17	624 (M+H) ⁺
1120	1349		LCMS-C-1	2.87	582 (M+H) ⁺
1109	1350		LCMS-C-1	2.73	648 (M+H) ⁺
1123	1351		LCMS-D-1	1.77	550 (M+H) ⁺


(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1124	1352		LCMS-D-1	1.58	550 (M+H) ⁺
1126	1353		LCMS-D-1	2.45	590 (M+H) ⁺
1128	1354		LCMS-D-1	2.98	598 (M+H) ⁺
1127	1355		LCMS-D-1	2.10	580 (M+H) ⁺
1318	1356		LCMS-D-1	1.55	598 (M+H) ⁺
1319	1357		LCMS-D-1	1.52	598 (M+H) ⁺

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1088	1358		LCMS-D-1	2.40	608 (M+H) ⁺
1089	1359		LCMS-D-1	2.48	608 (M+H) ⁺
1308	1360		LCMS-C-1	2.98	691 (M+H) ⁺
1307	1361		LCMS-C-1	2.68	592 (M+H) ⁺
1171	1362		LCMS-F-1	0.85	592 (M+H) ⁺
1172	1363		LCMS-F-1	0.88	591 (M+H) ⁺


(continued)

Starting Compound	1110	Target Compound	1364	Structure	LCMS condition	Retention time (min)	MS (m/z)
					LCMS-F-1	0.93	522 (M+H) ⁺

Example 317

1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(9-phenyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (Compound 1365)

[2161]

(Reaction 317-1)


[2162] 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(9-phenyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea (Compound 1365) was obtained by operations similar to those in Reaction 18-2 using Compound 1121 as a starting material.

MS (ESI) m/z = 622 (M+H)+.

Example 318

1-(4-{2-[2-(11-Amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1366)

[2163]

(Reaction 318-1)

[2164] 1-(4-{2-[2-(11-Amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-me-

thyl-urea (Compound 1366) was obtained by operations similar to those in Reaction 4-1 using Compound 1360 as a starting material.


MS (ESI) m/z = 591 (M+H)+.

Example 319

3-[(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-4-ethoxy-cyclobut-3-ene-1,2-dione (Compound 1367)

[2165]

(Reaction 319-1)




[2166] 3-[(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-4-ethoxy-cyclobut-3-ene-1,2-dione (Compound 1367) was obtained by operations similar to those in Reaction 12-5 and Reaction 95-17 (using ethanol as a solvent) using N-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-2,2,2-trifluoro-N-methyl-acetamide as a starting material.

MS (ESI) m/z = 599 (M+H)+.

Example 320

3-Amino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dione (Compound 1368)


[2167]

(Reaction 320-1)

[2168] 3-Amino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dione (Compound 1368) was obtained by operations similar to those in Reaction 230-3 using Compound 1367 as a starting material.
MS (ESI) m/z = 570 ($\text{M}+\text{H}$)⁺.

Example 321

3-Dimethylamino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dione (Compound 1369)


[2169](Reaction 321-1)

[2170] 3-Dimethylamino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dione (Compound 1369) was obtained by operations similar to those in Reaction 230-3 using Compound 1367 as a starting material.
MS (ESI) m/z = 598 ($\text{M}+\text{H}$)⁺.

Example 322

N-(4-{2-[2-(4-Ethynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide (Compound 1370)

[2171]

(Reaction 322-1)


[2172] Lithium hydroxide monohydrate (4.3 mg, 0.102 mmol) was added to a mixed solution of N-(4-{2-[2-(4-ethynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide (17 mg, 0.0341 mmol) in ethanol (1.25 mL) at room temperature. The mixture was stirred at 60°C for 14 hours and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give N-(4-{2-[2-(4-ethynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1371) as a white solid (17 mg, 99%). MS (ESI) m/z = 499 (M+H)⁺.

Example 323

1-(4-{2-[4-[(E)-Hydroxyimino]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1371)

[2173]

(Reaction 323-1)



[2174] 1-(4-{2-[4-[(E)-Hydroxyimino]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea (Compound 1371) was obtained by operations similar to those in Reaction 184-1, Reaction 88-1, Reaction 89-2 (using KOCN) and Reaction 189-9 using 3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-methyl-carbamic acid tert-butyl ester as a starting material.


MS (ESI) $m/z = 533 (M+H)^+$.

Example 324

2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic acid 2-methyl-benzylamide (Compound 1372)

[2175]

(Reaction 324-1)



[2176] 2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic acid 2-methyl-benzylamide (Compound 1372) was obtained by operations similar to those in Reaction 24-2 using 2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a starting material.


MS (ESI) $m/z = 433$ (M+H)+.

Example 325

2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic (2-o-tolyl-ethyl)-amide (Compound 1373)

[2177]

(Reaction 325-1)



[2178] 2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic (2-o-tolyl-ethyl)-amide (Compound 1373) was obtained by operations similar to those in Reaction 24-2 using 2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a starting material.


MS (ESI) $m/z = 448$ (M+H)+.

Example 326

2-Cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-2-hydroxy-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1374)


[2179]

(Reaction 326-1)




[2180] 1H-Indole-4-carbaldehyde (1.81 g, 12.5 mmol) and cesium carbonate (8.15 g, 25.0 mmol) were added to a solution of methanesulfonic acid (R)-2,2-dimethyl-[1,3]dioxolan-4-yl methyl ester (3.40 g, 16.1 mmol) in 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (30.8 mL), and the mixture was stirred at 90°C for 40 hours. Water was added, followed by extraction with hexane:ethyl acetate (1:4). The organic layer was washed with water four times and then dried over sodium sulfate. After concentration, the residue was purified by silica gel column chromatography to give 1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indole-4-carbaldehyde (2.88 g, 88%) as a yellow oily substance.

MS (ESI) $m/z = 260$ (M+H)+.

(Reaction 326-2)


[2181] A suspension of 2-cyclohexyl-8-methanesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (100 mg, 0.319 mmol) in 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (0.66 mL) was cooled to 0°C. A 1 M solution of lithium hexamethyldisilazide in tetrahydrofuran (0.989 mL) was then added and the mixture was stirred at room temperature for 30 minutes. After cooling again to 0°C, a solution of 1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indole-4-carbaldehyde (87 mg, 0.335 mmol) in tetrahydrofuran (0.4 mL) was added and the mixture was stirred at 0°C for five hours. Water was added and the aqueous layer was extracted with ethyl acetate. The organic layer was washed with water and then dried over sodium sulfate. After concentration, the residue was purified by silica gel column chromatography to give 2-cyclohexyl-8-{2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-4-yl]-2-hydroxy-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (113 mg, 62%) as a pale yellow solid. MS (ESI) m/z = 573 (M+H)+.

(Reaction 326-3)

[2182] 2-Cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-2-hydroxy-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1374) was synthesized by operations similar to those in Reaction 4-1 using appropriate reagents and starting material. MS (ESI) m/z = 573, 533 (M+H)+.

Example 327

2-Cyclohexyl-8-(2-oxo-2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1375)

[2183](Reaction 327-1)

[2184] 2-Cyclohexyl-8-(2-oxo-2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1375) was obtained by operations similar to those in Reaction 326-2 using 2-cyclohexyl-8-methanesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a starting material.


MS (ESI) $m/z = 432$ (M+H)+.

Example 328

2-Cyclohexyl-8-(2-o-tolyl-ethynesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1376)

[2185]

(Reaction 328-1)




[2186] 2-Chloro-1-methyl-pyridinium iodide (18 mg, 0.070 mmol) and triethylamine (0.28 mL, 1.98 mmol) were added to a solution of 2-cyclohexyl-8-(2-oxo-2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (20 mg, 0.046 mmol) in methylene chloride (1.0 mL), and the mixture was stirred at room temperature for 19 hours. 2-Chloro-1-methylpyridinium iodide (18 mg, 0.070 mmol) and triethylamine (0.28 mL, 1.98 mmol) were further added, and the mixture was stirred at room temperature for five hours. A 1 M aqueous sodium hydroxide solution was added to the reaction mixture, and the mixture was stirred at room temperature for 20 minutes. The aqueous layer was extracted with methylene chloride, and the organic layer was washed with a 1 M aqueous sodium hydroxide solution, water and saturated brine and dried over sodium sulfate. The organic layer was concentrated, and the residue was then silica gel column chromatography to give 2-cyclohexyl-8-(2-o-tolyl-ethynesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (15 mg, 79%) as a white solid.

$^1\text{H-NMR}$ (CDCl_3) δ 8.39 (1H, s), 7.60 (1H, dd, $J = 7.6, 1.2$ Hz), 7.38 (1H, td, $J = 7.6, 1.4$ Hz), 7.28-7.16 (2H, m), 3.80-3.75 (2H, m), 3.42-3.36 (2H, m), 2.52 (3H, s), 2.46-2.38 (1H, m), 2.14-2.07 (2H, m), 1.94-1.56 (8H, m), 1.47-1.22 (6H, m); MS (ESI) $m/z = 414$ (M+H)+.

Example 329

2-Cyclohexyl-8-[2-(1H-indol-4-yl)-ethynesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1377)


[2187]

(Reaction 329-1)

[2188] 2-Cyclohexyl-8-[2-(1H-indol-4-yl)-ethynesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1377) was obtained by operations similar to those in Reaction 326-2 and Reaction 328-1 using 2-cyclohexyl-8-methanesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a starting material.
MS (ESI) $m/z = 439$ (M+H)+.

Example 330

3,5,N,N-Tetramethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide (Compound 1378)

[2189](Reaction 330-1)

[2190] 3,5,N,N-Tetramethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide (Compound 1378) was obtained by operations similar to those in Reaction 122-2 using Compound 994 as a starting material.
MS (ESI) $m/z = 599$ (M+H)+.

[2191] The example compounds shown below were obtained by operations similar to those in Reaction 330-1 using appropriate solvents (acetonitrile or methanol or an acetonitrile-methanol mixed solution) and starting compounds.

(Compounds 1379 to Compound 1391)

[Table 191]




Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1003	1379		LCMS-D-1	1.91	517 (M+H)+
989	1380		LCMS-D-1	2.31	669 (M+H)+
990	1381		LCMS-D-1	2.26	687 (M+H)+
999	1382		LCMS-C-1	2.62	585 (M-H)-
1000	1383		LCMS-D-1	2.88	605 (M+H)+

[2192]

(continued)

Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
993	1384		LCMS-D-1	2.17	729 (M+H)+
996	1385		LCMS-D-1	1.87	586 (M+H)+
997	1386		LCMS-D-1	2.00	626 (M+H)+
987	1387		LCMS-F-1	0.94	517 (M+H)+
988	1388		LCMS-F-1	0.90	519 (M+H)+

(continued)


Starting Compound	Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1300	1389		LCMS-D-1	2.37	614 (M+H)+
1301	1390		LCMS-D-1	1.95	532 (M+H)+
1312	1391		LCMS-F-1	0.98	587 (M+H)+

Example 331

2-Cyclohexyl-8-{2-[1-((2S,3S)-2,3,4-trihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1392) and 8-{2-[1-((2S,3S)-4-benzyloxy-2,3-dihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl}-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1393)

[2193]

(Reaction 331-1)



[2194] 2-Cyclohexyl-8-{2-[1-((2S,3S)-2,3,4-trihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1392)

MS (ESI) m/z = 547 (M+H)+

and 8-{2-[1-((2S,3S)-4-benzyloxy-2,3-dihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl}-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (Compound 1393)


MS (ESI) m/z = 637 (M+H)+

were obtained by operations similar to those in Reaction 26-1, Reaction 4-1 and Reaction 42-2 using 2-cyclohexyl-8-ethanesulfonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as a starting material.

Example 332

N-[3-Methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-acetamide (Compound 1394)


[2195]

(Reaction 332-1)

[2196] N-[3-Methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-acetamide (Compound 1394) was obtained by operations similar to those in Reaction 12-2 and Reaction 14-1 (using NaOMe as a base) using Compound 1284 as a starting material.
MS (ESI) m/z = 585 (M+H)⁺.


Example 333

3,N,N-Trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide (Compound 1395)

[2197](Reaction 333-1)


[2198] N,N-Diisopropylethylamine (1.67 ml, 9.84 mmol) was added to a solution of 4-amino-4-carbamoyl-piperidine-1-carboxylic acid tert-butyl ester (1.0 g, 4.1 mmol) in THF (10 ml) at 0°C, and thiophosgene (0.376 ml, 4.9 mmol) was further added dropwise slowly. The reaction solution was warmed to room temperature and stirred overnight. A 10% aqueous citric acid solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layers were combined and dried over magnesium sulfate, and the solvent was then distilled off. The residue was purified by silica gel column chromatography to give 4-oxo-2-thioxo-1,3,8-triaza-spiro[4.5]decane-8-carboxylic acid tert-butyl ester (944 mg, 81%).
MS (ESI) m/z = 284 (M-H)⁻.

(Reaction 333-2)



10 [2199] Iodomethane (0.329 ml, 5.28 mmol) and a 1 N aqueous NaOH solution (3.3 ml, 3.3 mmol) were sequentially added to a solution of 4-oxo-2-thioxo-1,3,8-triaza-spiro[4.5]decane-8-carboxylic acid tert-butyl ester (944 mg, 3.3 mmol) in methanol (33 ml) at room temperature, and the mixture was stirred at the same temperature overnight. Water was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layers were combined and dried over magnesium sulfate, and the solvent was then distilled off. The residue was purified by silica gel column chromatography to give 2-methylsulfanyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (762 mg, 77%).


15 MS (ESI) m/z = 322 (M+Na)+.



30 [2200] Acetic acid (0.275 ml, 4.8 mmol) was added to a solution of 2-methylsulfanyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (72 mg, 0.24 mmol) and *m*-trifluoromethylaniline (0.150 ml, 1.2 mmol) in DMA (1.0 ml), and the mixture was irradiated with microwaves at 150°C for 20 minutes. A saturated aqueous sodium bicarbonate solution was added to the reaction mixture, followed by extraction with ethyl acetate. The organic layers were combined, washed with saturated brine and dried over magnesium sulfate, and the solvent was then distilled off. The residue was purified by silica gel column chromatography to give 4-oxo-2-(3-(trifluoromethyl)phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (50 mg, 51%).

35 MS (ESI) m/z = 313 (M-(Boc+H)+H)+.

40 (Reaction 333-4)




25 **[2201]** 3,N,N-Trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide was synthesized by operations similar to those in Reaction 4-1 and Reaction 190-1 using appropriate reagents and starting material.
MS (ESI) $m/z = 566$ (M+H)⁺.

30 Example 334

3,N,N-Trimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide (Compound 1396)

35 **[2202]**

(Reaction 334-1)



[2203] 3,N,N-Trimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide was synthesized by operations similar to those in Reaction 333-3, Reaction 5-3 and Reaction 190-1 using appropriate reagents and starting material.



MS (ESI) $m/z = 566$ (M+H)+.

[2204] The example compounds shown below were synthesized by operations similar to those in Reaction 334-1 using appropriate reagents and starting materials.



(Compounds 1397 to Compound 1400)

[2205]

[Table 192]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1397		LCMS-C-1	2.08	510 (M-H)-
1398		LCMS-C-1	2.20	524 (M-H)-

(continued)

Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1399		LCMS-C-1	2.17	518(M+H)+
1400		LCMS-B-1	1.83	580(M+H)+

Example 335

N-(3-Methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide (Compound 1401)

[2206](Reaction 335-1)

[2207] N-(3-Methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide was synthesized by operations similar to those in Reaction 190-1 using appropriate reagents and starting material.

MS (ESI) m/z = 552 (M+H)+.

[2208] The example compounds shown below were synthesized by operations similar to those in Reaction 335-1 using appropriate reagents and starting materials.

(Compound 1402)

[2209]

[Table 193]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1402		LCMS-B-1	1.85	552(M+H)+

Example 336

4-{2-[2-(4-Butyl-piperidin-1-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide (Compound 1403)

[2210]

(Reaction 336-1)



[2211] Acetic acid (0.115 ml, 1.336 mmol) and 2-methylsulfanyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (20 mg, 0.0668 mmol) were added to a solution of 4-butyl-piperidine hydrochloride (41 mg, 0.200 mmol) and a 10N aqueous sodium hydroxide solution (0.036 ml, 0.360 mmol) in DMI (0.3 ml), and the mixture was stirred at 110°C overnight. The reaction mixture was purified by silica gel column chromatography to give a mixture of 2-(4-butyl-piperidin-1-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (22.3 mg).

4-{2-[2-(4-Butyl-piperidin-1-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide (23.0 mg, 63% in three steps) was synthesized by operations similar to those in Reaction 4-1 and Reaction 190-1 using this mixture as a starting material.


MS (ESI) m/z = 546 (M+H)+.

[2212] The example compounds shown below were synthesized by operations similar to those in Reaction 336-1 using appropriate reagents and starting materials.

(Compound 1404)


[2213]

[Table 194]


Target Compound	Structure	LCMS condition	Retention time (min)	MS (m/z)
1404		LCMS-B-1	2.24	560(M+H)+

Example 337

4-{2-[2-(3-But-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-N-pent-4-enyl-benzamide (Compound 1405)

[2214]**(Reaction 337-1)**

[2215] 3-Hydroxy-5-trifluoromethylbenzoic acid (2.17 g, 10.6 mmol), potassium carbonate (8.73 g, 63.2 mmol) and 4-bromo-1-butene (4.34 ml, 43.7 mmol) were dissolved in DMF (21 ml), and this mixture was irradiated in a microwave apparatus (100°C, 60 min). The reaction solution was poured into a cooled aqueous dilute hydrochloric acid solution, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting crude product was purified by silica gel column chromatography to give 3-but-3-enyloxy-5-trifluoromethylbenzoic acid but-3-enyl ester (2.64 g, 80%). ^1H -NMR (400 MHz, $CDCl_3$) δ 7.86 (1H, s), 7.71 (1H, s), 7.31 (1H, s), 5.95-5.81 (2H, m), 5.22-5.11 (4H, m), 4.40 (2H, t, $J = 6.6$ Hz), 4.10 (2H, t, $J = 6.6$ Hz), 2.60-2.51 (4H, m).


(Reaction 337-2)

[2216] 3-But-3-enyloxy-5-trifluoromethylbenzoic acid but-3-enyl ester (2.64 g, 8.39 mmol) was dissolved in methanol.

A 5 N aqueous sodium hydroxide solution (5.1 ml, 25.2 mmol) was added and the mixture was stirred at room temperature for two hours. The reaction solution was cooled, quenched with 2 N hydrochloric acid (20 ml, 40 mmol) and then extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 3-but-3-enyloxy-5-trifluoromethyl-benzoic acid (2.13 g, 98%).

¹H-NMR (400 MHz, CDCl₃) δ 7.95 (1H, s), 7.78 (1H, s), 7.38 (1H, s), 5.96-5.86 (1H, m), 5.23-5.14 (2H, m), 4.12 (2H, t, J = 6.6 Hz), 2.59 (2H, q, J = 6.5 Hz).

(Reaction 337-3)




[2217] DMF (one drop) was added to a solution of 3-but-3-enyloxy-5-trifluoromethyl-benzoic acid (1.73 g, 6.65 mmol) in methylene chloride (6.8 ml). Oxalyl dichloride (0.566 ml, 6.60 mmol) was then added dropwise under ice-cooling, and the mixture was stirred at room temperature for three hours.

[2218] The reaction solution obtained above was added dropwise to a solution of 4-amino-4-cyano-piperidine-1-carboxylic acid tert-butyl ester (1.49 g, 6.65 mmol) and triethylamine (1.85 ml, 13.3 mmol) in methylene chloride (10 ml) under ice-cooling, and the mixture was stirred at room temperature for two hours. The reaction solution was cooled and water and 2 N hydrochloric acid were then sequentially added, followed by extraction with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give 4-(3-but-3-enyloxy-5-trifluoromethyl-benzoylamino)-4-cyano-piperidine-1-carboxylic acid tert-butyl ester as a crude product (3.0 g). This compound was used in the next reaction without further purification.

MS (ESI) m/z = 368 (M-Boc+H)+;

HPLC retention time: 3.32 min (analysis condition LCMS-A-1).


(Reaction 337-4)



[2219] 4-(3-But-3-enyloxy-5-trifluoromethyl-benzoylamino)-4-cyano-piperidine-1-carboxylic acid tert-butyl ester (3.0 g) was dissolved in ethanol, and a 5 N aqueous sodium hydroxide solution (6.9 ml, 34.5 mmol) and a 30% aqueous hydrogen peroxide solution (3 ml) were added. After stirring at room temperature for two hours, DMSO (19 ml) was added to the reaction solution, and the mixture was stirred at 50°C for four hours. The reaction solution was cooled, and then quenched with a saturated aqueous ammonium chloride solution and extracted with ethyl acetate. The organic layer was sequentially washed with a saturated aqueous ammonium chloride solution, water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (2.39 g, 67% in two steps).


¹H-NMR (400 MHz, CDCl₃) δ 10.10 (1H, s), 7.76 (1H, s), 7.63 (1H, s), 7.32 (1H, s), 5.96-5.86 (1H, m), 5.24-5.15 (2H, m), 4.15 (2H, t, J = 6.6 Hz), 4.01 (2H, s), 3.52 (2H, t, J = 11.2 Hz), 2.60 (2H, q, J = 6.7 Hz), 1.96-1.89 (2H, m), 1.65-1.55

(2H, m), 1.50 (9H, s);

MS (ESI) m/z = 368 (M-Boc+H)⁺, 412 (M-tBu+H)⁺.(Reaction 337-5)

[2220] Trifluoroacetic acid (27 ml) was added to a solution of 2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-carboxylic acid tert-butyl ester (2.39 g, 5.12 mmol) in methylene chloride (54 ml), and the mixture was stirred at room temperature for 1.5 hours. The reaction solution was concentrated under reduced pressure with azeotropic distillation with toluene, and the resulting residue (trifluoroacetate) was then dissolved in methanol (50 ml). A 4 N solution of hydrochloric acid in dioxane (16 ml) was added and the mixture was concentrated under reduced pressure. The resulting residue was dissolved in a mixed solution of ethyl acetate (100 ml)-ethanol (5 ml), followed by washing with a 1 N aqueous K₃PO₄ solution. The organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give 2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (1.92 g). This compound was used in the next reaction without further purification.


¹H-NMR (CDCl₃) δ 7.73 (1H, s), 7.63 (1H, s), 7.29 (1H, s), 5.97-5.87 (1H, m), 5.21 (2H, d, J = 17.1 Hz); 5.15 (2H, d, J = 10.3 Hz), 4.15 (2H, t, J = 6.8 Hz), 3.25-3.10 (4H, m), 2.60 (2H, q, J = 6.7 Hz), 1.95-1.85 (2H, m), 1.60-1.57 (2H, m); MS(ESI) m/z = 368 (M+H)⁺.

(Reaction 337-6)

[2221] Triethylamine (1.27 ml, 9.11 mmol) and 4-(2-chlorosulfonyl-ethyl)-3-methyl-benzoic acid methyl ester (1.01 g, 3.65 mmol) were added to a solution of 2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (1.40 g, 3.83 mmol) in methylene chloride (35 ml) at 0°C. The mixture was stirred at room temperature for two hours, and then quenched with a saturated aqueous ammonium chloride solution and extracted with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give 4-12-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl-3-methyl-benzoic acid methyl ester (2.10 g). This compound was used in the next reaction without further purification.

¹H-NMR (400 MHz, CDCl₃) δ 7.87 (1H, s), 7.85-7.84 (1H, m), 7.68 (1H, s), 7.57 (1H, s), 7.31 (1H, s), 7.26-7.25 (1H, m), 5.96-5.85 (1H, m), 5.22-5.17 (2H, m), 4.14 (2H, t, J = 6.6 Hz), 3.91 (3H, s), 3.83 (2H, td, J = 8.2, 3.9 Hz), 3.54-3.49 (2H, m), 3.25-3.15 (4H, m), 2.60 (1H, q, J = 6.7 Hz), 2.42 (3H, s), 2.13-2.06 (2H, m), 1.77-1.73 (2H, m);


MS (ESI) m/z = 608 (M+H)⁺.

(Reaction 337-7)

[2222] A 5 N aqueous sodium hydroxide solution (6.6 ml, 33 mmol) was added to a solution of 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid methyl ester (2.10 g) in methanol (22 ml), and the mixture was stirred at room temperature for two hours. The reaction solution was cooled and then quenched with 2 N hydrochloric acid (25 ml), followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid (1.78 g).

¹H-NMR (400 MHz, CD₃OD) δ 7.85-7.74 (3H, m), 7.41-7.32 (3H, m), 5.99-5.89 (1H, m), 5.19 (1H, dd, J = 17.3, 1.7 Hz), 5.11 (1H, dd, J = 10.3, 2.0 Hz), 4.17 (2H, t, J = 6.6 Hz), 3.84-3.76 (2H, m), 3.56-3.46 (2H, m), 3.41-3.17 (4H, m), 2.58 (2H, q, J = 6.7 Hz), 2.44 (3H, s), 2.07-1.97 (2H, m), 1.78-1.69 (2H, m);

MS (ESI) m/z = 594 (M+H)+.

(Reaction 337-8)


[2223] HATU (194 mg, 0.510 mmol), N,N-diisopropylethylamine (143 μL) and methyl-pent-4-enyl-amine (80 mg) were added to a solution of 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid (200 mg, 0.337 mmol) in DMF (3 ml), and the mixture was stirred at room temperature overnight. A saturated aqueous ammonium chloride solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-N,N-dimethyl-N-pent-4-enyl-benzamide (192 mg, 84%).

MS (ESI) m/z = 675 (M+H)+;

HPLC retention time: 3.13 min (analysis condition LCMS-A-1).

[2224] Methyl-pent-4-enyl-amine used in the above Reaction 337-8 was synthesized by the following method (Angewandte Chemie, International Edition (2004), 43(41), 5542-5546).

(Reaction 337-9)



[2225] A 40% solution of methylamine in methanol (2.74 ml, 26.8 mmol) and NaI (20 mg, 0.134 mmol) were added to a solution of 5-bromo-1-butene (318 μ L, 2.68 mmol) in ethanol (2 ml), and the mixture was stirred at 60°C overnight in a sealed tube. The reaction solution was cooled and concentrated hydrochloric acid (2.4 ml) was then added. The mixture was concentrated under reduced pressure. The resulting residue was washed with tert-butyl methyl ether and then made basic with a 5 N aqueous sodium hydroxide solution under ice-cooling, followed by extraction with tert-butyl methyl ether (x 3). The organic layers were dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give methyl-pent-4-enyl-amine (80 mg, 30% as an object).


¹H-NMR (400 MHz, CDCl₃) δ 5.88-5.77 (1H, m), 5.05-4.99 (1H, m), 4.98-4.93 (1H, m), 2.58 (2H, t, J = 7.1 Hz), 2.43 (3H, s), 2.12-2.03 (2H, m), 1.62-1.49 (2H, m).

Example 338

Compound 1406

[2226]

(Reaction 338-1)




Compound 1405

Compound 1406

[2227] Grubbs catalyst 2nd generation (44 mg, 0.0519 mmol) was added to a solution of 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-N-pent-4-enyl-benzamide (175 mg, 0.259 mmol) in 1,2-dichloroethane (260 ml), and the mixture was stirred at 40°C overnight in an argon stream. The reaction solution was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography to give a macrocyclic olefin compound (Compound 1406) (157 mg, 94%).

¹H-NMR (400 MHz, CDCl₃) δ 9.76 (0.2H, s), 9.59 (0.8H, s), 8.19 (1H, s), 8.12 (1H, s), 7.35-7.10 (4H, m), 5.61-5.48 (2H, m), 4.20 (0.8H, t, J = 5.4 Hz), 4.09 (0.2H, t, J = 5.1 Hz), 3.67-3.05 (10H, m), 3.03 (0.6H, s), 2.98 (2.4H, s), 2.65-2.48 (2H, m), 2.47 (2.4H, s), 2.41 (0.6H, s), 2.33-2.18 (2H, m), 1.80-1.22 (6H, m);

MS (ESI) m/z = 647 (M+H)⁺.

(Reaction 340-2)

[2232] A macrocyclic olefin compound (Compound 1408) was obtained by the same method as in Reaction 338-1 using N-allyl-4-{2-[2-(3-butoxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide (151 mg, 0.233 mmol) as a starting material.


MS (ESI) m/z = 619 (M+H)⁺;

HPLC retention time: 2.69 min (analysis condition LCMS-A-1).

Example 341

Compound 1409

[2233]

(Reaction 341-1)

[2234] A saturated macrocyclic compound (Compound 1409) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1408) as a starting material.


MS (ESI) m/z = 621 (M+H)⁺;

HPLC retention time: 2.72 min (analysis condition LCMS-A-1).

Example 342

Compounds 1410 and Compound 1411

[2235]


(Reaction 342-1)

[2236] 4-{2-[2-(3-But-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-hex-5-enyl-3,N-dimethyl-benzamide was obtained by the same method as in Reaction 337-8 using 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methylbenzoic acid and hex-5-enyl-methyl-amine as starting materials.

MS (ESI) m/z = 689 (M+H)+;


HPLC retention time: 3.32 min (analysis condition LCMS-A-1).

[2237] Hex-5-enyl-methyl-amine used in the above Reaction 342-1 was synthesized in the following manner.

(Reaction 342-2)

[2238] Hex-5-enyl-methyl-amine was obtained by the same method as in Reaction 337-9 using 6-bromo-1-hexene (437 mg, 2.68 mmol) as a raw material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 5.86-5.76 (1H, m), 5.03-4.93 (2H, m), 2.57 (2H, t, J = 7.0 Hz), 2.43 (3H, s), 2.10-2.04 (2H, m), 1.54-1.38 (4H, m).

(Reaction 342-3)

[2239] A macrocyclic olefin compound (Compound 1410, E/Z = 98:2) and a macrocyclic olefin compound (Compound 1411, E/Z = 59:41) were obtained by the same method as in Reaction 338-1 using 4-{2-[2-(3-but-3-enyloxy-5-trifluor-

omethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-N-hex-5-enyl-3,N-dimethyl-benzamide as a starting material.

Compound 1410

MS (ESI) m/z = 661 (M+H)+;

HPLC retention time: 3.09 min (analysis condition LCMS-A-1).

Compound 1411

MS (ESI) m/z = 661 (M+H)+;


HPLC retention time: 3.08 min (analysis condition LCMS-A-1).

Example 343

Compound 1412

[2240]

(Reaction 343-1)



[2241] A saturated macrocyclic compound (Compound 1412) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1411) as a starting material.

MS (ESI) m/z = 663 (M+H)+;


HPLC retention time: 3.22 min (analysis condition LCMS-A-1).

Example 344

Compounds 1413 and Compound 1414

[2242]

(Reaction 344-1)




[2243] 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-hept-6-enyl-3,N-dimethyl-benzamide was obtained by the same method as in Reaction 337-8 using 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid and hept-6-enyl-methyl-amine as starting materials.

MS (ESI) m/z = 703 (M+H)+;

HPLC retention time: 3.45 min (analysis condition LCMS-A-1).

[2244] Hept-6-enyl-methyl-amine used in the above Reaction 344-1 was synthesized as follows.


(Reaction 344-2)



[2245] Hept-6-enyl-methyl-amine was obtained by the same method as in Reaction 337-9 using 7-bromo-1-heptene as a raw material.

¹H-NMR (CDCl₃) δ 5.86-5.76 (1H, m), 5.02-4.97 (1H, m), 4.95-4.92 (1H, m), 2.56 (2H, t, J = 7.1 Hz), 2.43 (3H, s), 2.05 (2H, q, J = 7.0 Hz), 1.52-1.29 (6H, m).

(Reaction 344-3)



[2246] A macrocyclic olefin compound (E/Z mixture) was obtained by the same method as in Reaction 338-1 using 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-hept-6-enyl-3,N-dimethyl-benzamide as a starting material. This mixture was purified by HPLC to give Compound 1413 (E/Z = 97:3) and Compound 1414 (E/Z = 10:90).

Compound 1413

MS (ESI) m/z = 675 (M+H)+;

HPLC retention time: 3.20 min (analysis condition LCMS-A-1).

Compound 1414

MS (ESI) m/z = 675 (M+H)+;


HPLC retention time: 3.18 min (analysis condition LCMS-A-1).

Example 345

Compound 1415

[2247]

(Reaction 345-1)



[2248] A saturated macrocyclic compound (Compound 1415) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (a mixture of Compound 1413 and Compound 1414) as a starting material.

MS (ESI) $m/z = 677$ (M+H)+;


HPLC retention time: 3.34 min (analysis condition LCMS-A-1).

Example 346

Compounds 1416 and Compound 1417

[2249]

(Reaction 346-1)




[2250] 4-{2-[2-(3-But-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-N-oct-7-enyl-benzamide was obtained by the same method as in Reaction 337-8 using 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methylbenzoic acid and methyl-oct-7-enyl-amine as starting materials.

MS (ESI) $m/z = 717$ (M+H)+;

HPLC retention time: 3.55 min (analysis condition LCMS-A-1)

[2251] Methyl-oct-7-enyl-amine used in the above Reaction 346-1 was synthesized as follows.


(Reaction 346-2)



[2252] Methyl-oct-7-enyl-amine was obtained by the same method as in Reaction 337-9 using 8-bromo-1-octene as a raw material.

¹H-NMR (400 MHz, CDCl₃) δ 5.86-5.76 (1H, m), 5.02-4.96 (1H, m), 4.95-4.91 (1H, m), 2.56 (2H, t, J = 7.1 Hz), 2.43 (3H, s), 2.07-2.01 (2H, m), 1.50-1.30 (8H, m).

(Reaction 346-3)



[2253] A macrocyclic olefin compound (E/Z mixture) was obtained by the same method as in Reaction 338-1 using 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-N-oct-7-enyl-benzamide as a starting material. The resulting mixture was purified by HPLC (MeOH/MeCN/H₂O) to give Compound 1416 (E/Z = 96:4) and Compound 1417 (E/Z = 19:81).

Compound 1416

MS (ESI) m/z = 689 (M+H)⁺;

HPLC retention time: 3.38 min (analysis condition LCMS-A-1).

Compound 1417


MS (ESI) m/z = 689 (M+H)⁺;

HPLC retention time: 3.26 min (analysis condition LCMS-A-1).

Example 347

Compound 1418

[2254]

(Reaction 347-1)

[2255] A saturated macrocyclic compound (Compound 1418) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1416) as a starting material.


MS (ESI) $m/z = 691$ (M+H)⁺;

HPLC retention time: 3.56 min (analysis condition LCMS-A-1).

Example 348

Compound 1419

[2256]


(Reaction 348-1)

[2257] N-(2-Allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide was obtained by the same method as in Reaction 337-8 using 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid and (2-allyloxy-ethyl)-methyl-amine as starting materials.

MS (ESI) $m/z = 691$ (M+H)⁺;

HPLC retention time: 3.08 min (analysis condition LCMS-A-1).

[2258] (2-Allyloxy-ethyl)-methyl-amine used in the above Reaction 348-1 was synthesized in the following manner.

(Reaction 348-2)

[2259] (2-Allyloxy-ethyl)-methyl-amine was obtained by the same method as in Reaction 337-9 using, as a starting material, methanesulfonic acid 2-allyloxy-ethyl ester synthesized from 2-allyloxy-ethanol by the method described in Journal of Organic Chemistry (2006), 71(21), 8183-8189.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 5.97-5.87 (1H, m), 5.30-5.24 (1H, m), 5.20-5.17 (1H, m), 4.00 (2H, br d, $J = 5.9$ Hz), 3.55 (2H, t, $J = 5.4$ Hz), 2.76 (2H, t, $J = 5.1$ Hz), 2.45 (3H, s).

Example 349

Compound 1420

[2260]

(Reaction 349-1)



[2261] A macrocyclic olefin compound (Compound 1420) and its isomer A (349a) and isomer B (349b) were obtained by the same method as in Reaction 338-1 using N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluoromethylphenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide as a starting material.

Compound 1420

MS (ESI) $m/z = 663$ ($M+H$) $^+$;

HPLC retention time: 2.84 min (analysis condition LCMS-A-1)

Isomer A (349a)

MS (ESI) $m/z = 663$ ($M+H$) $^+$

HPLC retention time: 2.77 min (analysis condition LCMS-A-1).

Isomer B (349b)


MS (ESI) $m/z = 663$ ($M+H$) $^+$

HPLC retention time: 2.96 min (analysis condition LCMS-A-1).

Example 350

Compound 1421

[2262]

(Reaction 350-1)


[2263] A saturated macrocyclic compound (Compound 1421) was obtained by the same method as in Reaction 339-1 using macrocyclic olefin mixture (Compounds 1420, 349a and 349b) as a starting material.

MS (ESI) $m/z = 665$ ($M+H$)⁺;

HPLC retention time: 2.95 min (analysis condition LCMS-A-1).

Example 351

Compounds 1422 and Compound 1423

[2264](Reaction 351-1)


[2265] N-Allylcarbamoylmethyl-4-{2-[2-(3-butenyloxy)-5-(trifluoromethyl)-phenyl]-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide was obtained by the same method as in Reaction 337-8 using 4-{2-[2-(3-butenyloxy)-5-(trifluoromethyl)-phenyl]-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid and N-allyl-2-methylamino-acetamide hydrochloride as starting materials.

MS (ESI) $m/z = 704$ ($M+H$)⁺;

HPLC retention time: 2.80 min (analysis condition LCMS-A-1).

[2266] N-Allyl-2-methylamino-acetamide hydrochloride used in the above reaction was synthesized by the following

method.


(Reaction 351-2)

[2267] Allylamine (0.377 ml, 5.03 mmol) and 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride n-hydrate (DMT-MM) (1.89 g, 6.04 mmol) were added to a solution of Boc-sarcosine (1.0 g, 5.29 mmol) in ethanol, and the mixture was stirred at room temperature for 18 hours. A saturated aqueous sodium bicarbonate solution and water were added to the reaction solution, followed by extraction with ether. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give allylcarbamoylmethyl-methyl-carbamic acid tert-butyl ester (712 mg).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 6.27 (0.5H, br s), 6.02 (0.5H, br s), 5.88-5.79 (1H, m), 5.18 (1H, br d, $J = 17.6$ Hz), 5.15 (1H, br d, $J = 11.2$ Hz), 3.91 (2H, br t, $J = 5.6$ Hz), 3.88 (2H, s), 2.95 (3H, s), 1.47 (9H, s).

[2268] Trifluoroacetic acid (7 ml) was added to a solution of the resulting allylcarbamoylmethyl-methyl-carbamic acid tert-butyl ester in methylene chloride (14 ml), and the mixture was stirred at room temperature for three hours. The reaction solution was concentrated under reduced pressure, and 4 N hydrochloric acid-dioxane was then added to the resulting residue. The mixture was concentrated under reduced pressure again to give N-allyl-2-methylamino-acetamide hydrochloride (577 mg). This was used in the next reaction without complete purification.

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 8.96 (2H, br s), 8.68 (1H, br t, $J = 5.6$ Hz), 5.86-5.76 (1H, m), 5.19 (1H, dq, $J = 17.1$, 1.6 Hz), 5.10 (1H, dq, $J = 10.4$, 1.5 Hz), 3.79-3.75 (2H, m), 3.71 (2H, br s), 2.55 (2H, br s).

(Reaction 351-3)

[2269] A macrocyclic olefin compound (Compound 1422) and its isomer (Compound 1423) were obtained by the same method as in Reaction 338-1 using N-allylcarbamoylmethyl-4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide as a starting material.

Compound 1422

MS (ESI) $m/z = 676$ ($\text{M}+\text{H}$) $^+$;

HPLC retention time: 2.47 min (analysis condition LCMS-A-1).

Compound 1423

MS (ESI) $m/z = 676$ ($\text{M}+\text{H}$) $^+$;


HPLC retention time: 2.61 min (analysis condition LCMS-A-1).

Example 352

Compound 1424

[2270]

(Reaction 352-1)




[2271] 8-{2-[4-((3S,4S)-3-Allyloxy-4-hydroxy-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one were obtained by the same method as in Reaction 337-8 using 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid and (3S,4S)-4-allyloxy-pyrrolidin-3-ol hydrochloride as starting materials.

MS (ESI) m/z = 719 (M+H)+;

HPLC retention time: 2.81 min (analysis condition LCMS-A-1).


[2272] (3S,4S)-4-Allyloxy-pyrrolidin-3-ol hydrochloride used in the above reaction was synthesized by the following method.

(Reaction 352-2)



[2273] (3S,4S)-3-Allyloxy-4-hydroxy-pyrrolidine-1-carboxylic acid tert-butyl ester synthesized by the method described in the patent literature (DE4234330) (139 mg, 0.57 mmol) was dissolved in methylene chloride (2.4 ml). A 4 N solution of hydrochloric acid in dioxane (0.628 ml, 2.45 mmol) was added and the mixture was stirred at room temperature for two hours. The reaction solution was concentrated under reduced pressure to give (3S,4S)-4-allyloxy-pyrrolidin-3-ol hydrochloride (105 mg). This was used in the next reaction without further purification.

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 9.43 (2H, br s), 5.93-5.83 (1H, m), 5.72 (1H, br d, J = 2.4 Hz), 5.28 (1H, dq, J = 17.3, 1.8 Hz), 5.17 (1H, dq, J = 10.5, 1.5 Hz), 4.26 (1H, br s), 4.04-4.02 (2H, m), 3.95 (1H, d, J = 4.4 Hz), 3.32-3.06 (4H, m).

(Reaction 352-3)

[2274] A macrocyclic olefin compound (Compound 1424) was obtained by the same method as in Reaction 338-1 (using Hoveyda-Grubbs 2nd generation as a catalyst) using 8-{2-[4-((3S,4S)-3-allyloxy-4-hydroxy-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (106 mg, 0.148 mmol) as a starting material.


MS (ESI) m/z = 691 (M+H)⁺;

HPLC retention time: 2.48 min (analysis condition LCMS-A-1).

Example 353

Compound 1425

[2275]

(Reaction 353-1)

[2276] A saturated macrocyclic compound (Compound 1425) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1424) as a starting material.

MS (ESI) m/z = 693 (M+H)⁺;


HPLC retention time: 2.54 min (analysis condition LCMS-A-1).

Example 354

Compound 1426

[2277]

(Reaction 354-1)




[2278] N-(2-Allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzamide was obtained by the same method as in Reaction 337-8 using 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methylbenzoic acid and 2-allyloxy-ethylamine as starting materials.

MS (ESI) $m/z = 677$ (M+H)+;

HPLC retention time: 1.08 min (analysis condition LCMS-F-1).

(Reaction 354-2)



[2279] A macrocyclic olefin compound (Compound 1426) was obtained by the same method as in Reaction 338-1 using N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzamide as a starting material.

MS (ESI) $m/z = 649$ (M+H)+;


HPLC retention time: 2.83 min (analysis condition LCMS-C-1).

Example 355

Compound 1427

[2280]

(Reaction 355-1)



[2281] A saturated macrocyclic compound (Compound 1427) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1426) as a starting material.

MS (ESI) m/z = 651 (M+H)⁺;


HPLC retention time: 1.07 min (analysis condition LCMS-F-1).

Example 356

Compound 1428

[2282]

(Reaction 356-1)




[2283] N-(2-Allyloxy-ethyl)-4-{2-[2-(3-butenyloxy)-5-(trifluoromethyl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-(2-hydroxy-ethyl)-3-methyl-benzamide was obtained by the same method as in Reaction 337-8 using 4-{2-[2-(3-butenyloxy)-5-(trifluoromethyl)phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid and 2-(2-allyloxy-ethylamino)-ethanol as starting materials.


MS (ESI) m/z = 721 (M+H)⁺;

HPLC retention time: 1.05 min (analysis condition LCMS-F-1).

[2284] 2-(2-Allyloxy-ethylamino)-ethanol used in the above Reaction 356-1 was synthesized by the following method.

(Reaction 356-2)

[2285] 2-(2-Allyloxy-ethylamino)-ethanol was obtained by the same method as in Reaction 337-9 using methanesulfonic acid 2-allyloxy-ethyl ester as a raw material.

(Reaction 356-3)


[2286] A macrocyclic olefin compound (Compound 1428) was obtained by the same method as in Reaction 338-1 (using Hoveyda-Grubbs 2nd generation as a catalyst) using N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-(2-hydroxy-ethyl)-3-methyl-benzamide as a starting material.

MS (ESI) *m/z* = 693 (M+H)⁺;

HPLC retention time: 1.06 min (analysis condition LCMS-F-1).

Example 357

Compound 1429

[2287](Reaction 357-1)


[2288] A saturated macrocyclic compound (Compound 1429) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1428) as a starting material.

MS (ESI) *m/z* = 695 (M+H)⁺;

HPLC retention time: 1.09 min (analysis condition LCMS-F-1).

HPLC retention time: 0.61 min (analysis condition LCMS-F-1).

(Reaction 359-2)




[2293] $Pd(OH)_2 \cdot C$ (20.0 g) was added to a solution of 4-[(E)-2-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-vinyl]-3-methylbenzoic acid (20.0 g, 54.43 mmol) in THF (600 ml)-methanol (200 ml), and the mixture was stirred at room temperature overnight in a hydrogen atmosphere. The reaction mixture was filtered through celite, and the filtrate was then concentrated under reduced pressure to give 4-[2-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-ethyl]-3-methylbenzoic acid as a white solid (18.23 g, 90.7%).

MS (ESI) m/z = 370 ($M+H$)⁺;

HPLC retention time: 1.85 min (analysis condition LCMS-B-1).

(Reaction 359-3)




[2294] A 6 M aqueous hydrochloric acid solution (217.9 ml, 1307.4 mmol) was slowly added to a suspension of 4-[2-(1,4-dioxo-8-aza-spiro[4.5]decane-8-sulfonyl)-ethyl]-3-methylbenzoic acid (16.1 g, 43.58 mmol) in acetone (485 ml) at 0°C, and the mixture was warmed to room temperature and stirred overnight. The reaction mixture was filtered off, and the filtrate was then concentrated under reduced pressure. The precipitated solid was filtered off again. The solids filtered off were combined and dried to give 3-methyl-4-[2-(4-oxo-piperidine-1-sulfonyl)-ethyl]-benzoic acid as a white solid (13.62 g, 91.8%).

MS (ESI) m/z = 326 ($M+H$)⁺;

HPLC retention time: 1.57 min (analysis condition LCMS-B-1).

(Reaction 359-4)




[2295] HATU (91 mg, 0.239 mmol) was added to a solution of 3-methyl-4-[2-(4-oxo-piperidine-1-sulfonyl)-ethyl]-benzoic acid (50 mg, 0.154 mmol), (2-allyloxy-ethyl)-methylamine (36 mg, 0.312 mmol) and diisopropylethylamine (0.065 ml, 0.384 mmol) in DMF (0.5 ml), and the mixture was stirred at room temperature overnight. Water (12 ml) and 1 N hydrochloric acid (1.5 ml) were added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was sequentially washed with 0.1 N hydrochloric acid, water and saturated brine, and then dried over anhydrous

sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give N-(2-allyloxy-ethyl)-3,N-dimethyl-4-[2-(4-oxo-piperidine-1-sulfonyl)-ethyl]-benzamide (76 mg, 100%).

MS (ESI) $m/z = 423$ (M+H)+;

HPLC retention time: 2.15 min (analysis condition LCMS-C-1).

(Reaction 359-5)




[2296] Potassium cyanide (382 mg, 5.87 mmol) and ammonium acetate (513 mg, 6.65 mmol) were added to a solution of N-(2-allyloxy-ethyl)-3,N-dimethyl-4-[2-(4-oxo-piperidine-1-sulfonyl)-ethyl]-benzamide (1.65 g, 3.91 mmol) in methanol (20 ml), and the mixture was stirred at 65°C for three hours. Sodium bicarbonate (290 mg) was added to the reaction solution, and the mixture was then concentrated under reduced pressure. Water was added to the resulting residue, followed by extraction with methylene chloride. The organic layer was washed with saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give N-(2-allyloxy-ethyl)-4-[2-(4-amino-4-cyano-piperidine-1-sulfonyl)-ethyl]-3,N-dimethyl-benzamide (1.57 g) as a crude compound.

[2297] A 1 N aqueous sodium hydroxide solution (1.41 ml) and 30% aqueous hydrogen peroxide (0.95 ml) were added to a solution of the resulting N-(2-allyloxy-ethyl)-4-[2-(4-amino-4-cyano-piperidine-1-sulfonyl)-ethyl]-3,N-dimethyl-benzamide in methanol (24 ml)-DMSO (1.3 ml), and the mixture was stirred at room temperature for one hour. A 10% (w/w) aqueous sodium sulfite solution (2.64 ml) was added to the reaction solution, and the mixture was stirred at room temperature for 40 minutes. The precipitated insoluble matter was then removed by filtration. The resulting filtrate was concentrated under reduced pressure to give 1-(2-{4-[(2-allyloxy-ethyl)-methyl-carbamoyl]-2-methyl-phenyl}-ethanesulfonyl)-4-amino-piperidine-4-carboxylic amide (2.13 g). This was used in the next reaction without further purification.

MS (ESI) $m/z = 467$ (M+H)+;

HPLC retention time: 1.59 min (analysis condition LCMS-A-1).

(Reaction 359-6)



[2298] Potassium tert-butoxide (816 mg, 7.28 mmol) was added to a solution of 3-fluoro-4-(trifluoromethyl)benzoic acid (682 mg, 3.28 mmol) and benzyl alcohol (471 mg, 4.36 mmol) in DMSO (7.3 ml), and the mixture was stirred at room

temperature for 16 hours. The reaction solution was made acidic by adding concentrated-hydrochloric acid, and the precipitated insoluble matter was then filtered off. The resulting solid was washed with water and then dried to give 3-benzyloxy-4-trifluoromethyl-benzoic acid as a crude compound.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.77-7.70 (3H, m), 7.48-7.32 (5H, m), 5.27 (2H, s);

MS (ESI) m/z = 295 (M-H) $^-$;

HPLC retention time: 2.37 min (analysis condition LCMS-C-1).


[2299] 10% Pd/C (590 mg) was added to 3-benzyloxy-4-trifluoromethyl-benzoic acid in a methanol-ethyl acetate mixed solvent (1:1), and the mixture was stirred at room temperature for two days in a hydrogen atmosphere. The reaction solution was filtered through celite, and the filtrate was then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane-ethyl acetate) to give 3-hydroxy-4-trifluoromethyl-benzoic acid (582 mg, 86%).

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 13.26 (1H, br s), 10.94 (1H, br s), 7.63 (1H, d, J = 8.3 Hz), 7.59 (1H, s), 7.46 (1H, d, J = 8.8 Hz);

MS (ESI) m/z = 205 (M-H) $^-$;

HPLC retention time: 1.15 min (analysis condition LCMS-C-1).


(Reaction 359-7)



[2300] 3-But-3-enyloxy-4-trifluoromethyl-benzoic acid but-3-enyl ester was obtained by the same method as in Reaction 337-1 using 3-hydroxy-4-trifluoromethyl-benzoic acid as a raw material.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.66-7.61 (3H, m), 5.98-5.81 (2H, m), 5.22-5.15 (2H, m), 5.14-5.10 (2H, m), 4.40 (2H, t, J = 6.8 Hz), 4.16 (2H, t, J = 6.6 Hz), 2.62-2.51 (4H, m).


(Reaction 359-8)



[2301] 3-But-3-enyloxy-4-trifluoromethyl-benzoic acid was obtained by the same method as in Reaction 337-2 using 3-but-3-enyloxy-4-trifluoromethyl-benzoic acid but-3-enyl ester as a raw material.

$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ 7.73 (1H, br s), 7.68 (2H, br s), 6.00-5.90 (1H, m), 5.21-5.15 (1H, m), 5.11-5.08 (1H, m), 4.19 (2H, t, J = 6.6 Hz), 2.60-2.55 (2H, m).

(Reaction 359-9)




[2302] HATU (245 mg, 0.644 mmol) was added to a solution of 1-(2-{4-[(2-allyloxy-ethyl)-methyl-carbamoyl]-2-methyl-phenyl}-ethanesulfonyl)-4-amino-piperidine-4-carboxylic amide (250 mg), 3-but-3-enyloxy-4-trifluoromethyl-benzoic acid (155 mg, 0.596 mmol) and diisopropylethylamine (0.140 ml, 0.812 mmol) in DMF (5 ml), and the mixture was stirred at room temperature for 1.5 hours. Water was added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 1-(2-{4-[(2-allyloxy-ethyl)-methyl-carbamoyl]-2-methyl-phenyl}-ethanesulfonyl)-4-(3-but-3-enyloxy-4-trifluoromethyl-benzoylamino)-piperidine-4-carboxylic amide as a crude compound (381 mg).

[2303] Potassium tert-butoxide (302 mg, 2.69 mmol) was added to a solution of the resulting 1-(2-{4-[(2-allyloxy-ethyl)-methyl-carbamoyl]-2-methyl-phenyl}-ethanesulfonyl)-4-(3-but-3-enyloxy-4-trifluoromethyl-benzoylamino)-piperidine-4-carboxylic amide in ethanol (6 ml), and the mixture was stirred at 80°C for 30 minutes. A saturated aqueous ammonium chloride solution, water and saturated brine were sequentially added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-4-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide (176 mg).

MS (ESI) m/z = 691 ($M+H$)⁺;

HPLC retention time: 3.03 min (analysis condition LCMS-C-1).

(Reaction 359-10)



[2304] A macrocyclic olefin compound (Compound 1431) (E/Z = 1:2) was obtained by the same method as in Reaction 338-1 (using Hoveyda-Grubbs 2nd generation as a catalyst) using N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-4-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide as a starting material.

MS (ESI) m/z = 663 ($M+H$)⁺;


HPLC retention time: 2.90 min (analysis condition LCMS-C-1).

Example 360

Compound 1432

[2305]

(Reaction 360-1)



[2306] A saturated macrocyclic compound (Compound 1432) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1431) as a starting material.

MS (ESI) m/z = 665 (M+H)⁺;


HPLC retention time: 1.06 min (analysis condition LCMS-C-1).

Example 361

Compound 1433


[2307]

(Reaction 361-1)




[2308] Benzyl bromide (2.17 ml, 18.3 mmol) was added to a solution of 4-formyl-3-hydroxybenzoic acid (1.01 g, 6.10 mmol) and potassium carbonate (3.37 g, 24.4 mmol) in DMF (10 ml), and the mixture was stirred at 50°C for five hours. The reaction solution was poured into ice water and extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 3-benzyloxy-4-formylbenzoic acid benzyl ester (2.01 g, 95%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 10.58 (1H, s), 7.89 (1H, d, J = 8.3 Hz), 7.78 (1H, d, J = 1.5 Hz), 7.73 (1H, br d, J = 8.3 Hz), 7.46-7.35 (10H, m), 5.38 (2H, s), 5.24 (2H, s).

(Reaction 361-2)


[2309] n-Butyllithium (1.65 M solution in n-hexane, 4.22 ml, 6.92 mmol) was added dropwise to a solution of triphenyl-n-propyl-phosphonium bromide (2.91 g, 7.54 mmol) in THF (49 ml) at 0°C, and the mixture was stirred for 30 minutes. Further, a solution of 3-benzyloxy-4-formyl-benzoic acid benzyl ester (2.01 g, 5.80 mmol) in THF (4.9 ml) was added dropwise and then the mixture was stirred at 0°C for 30 minutes and at room temperature for 16 hours. A saturated aqueous ammonium chloride solution, water and saturated brine were added to the reaction solution, and this mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (n-hexane-ethyl acetate) to give 3-benzyloxy-4-((E/Z)-but-1-enyl)-benzoic acid benzyl ester (E/Z = 2:3, 1.74 g, 80%).

¹H-NMR (400 MHz, CDCl₃) δ 7.68-7.60 (2H, m), 7.50-7.30 (11H, m), 6.79 (0.4H, d, J = 16.1 Hz), 6.57 (0.6H, d, J = 11.7 Hz), 6.40 (0.4H, dt, J = 16.0, 6.6 Hz), 5.78 (0.6H, dt, J = 13.7, 5.9 Hz), 5.36 (1.2H, s), 5.35 (0.8H, s), 5.14 (0.8H, s), 5.14 (1.2H, s), 2.32-2.22 (2H, m), 1.09 (1.2H, t, J = 7.3 Hz), 1.04 (1.8H, t, J = 7.6 Hz).

(Reaction 361-3)

[2310] 10% Pd/C (174 mg) was added to 3-benzyloxy-4-((E/Z)-but-1-enyl)-benzoic acid benzyl ester (1.74 g, 4.67 mmol) in a methanol-ethyl acetate mixed solvent (1:1), and the mixture was stirred at room temperature for 21 hours in a hydrogen atmosphere. The reaction solution was filtered through celite, and the filtrate was then concentrated under reduced pressure to give 4-butyl-3-hydroxy-benzoic acid (922 mg) as a crude compound.


¹H-NMR (400 MHz, CD₃OD) δ 7.42-7.39 (2H, m), 7.14-7.11 (1H, m), 2.64 (2H, t, J = 7.6 Hz), 1.62-1.54 (2H, m), 1.42-1.33 (2H, m), 0.94 (3H, t, J = 7.3 Hz).

(Reaction 361-4)

[2311] 3-But-3-enyloxy-4-butyl-benzoic acid but-3-enyl ester was obtained by the same method as in Reaction 337-1 using 4-butyl-3-hydroxy-benzoic acid as a raw material.

¹H-NMR (400 MHz, CDCl₃) δ 7.56 (1H, dd, J = 7.8, 1.5 Hz), 7.47 (1H, d, J = 1.5 Hz), 7.17 (1H, d, J = 7.3 Hz), 5.97-5.82 (2H, m), 5.21-5.14 (2H, m), 5.13-5.08 (2H, m), 4.35 (2H, t, J = 6.8 Hz), 4.07 (2H, t, J = 6.3 Hz), 2.64 (2H, t, J = 7.8 Hz), 2.60-2.49 (4H, m), 1.60-1.52 (2H, m), 1.40-1.30 (2H, m), 0.92 (3H, t, J = 7.3 Hz).


(Reaction 361-5)



[2312] 3-But-3-enyloxy-4-butyl-benzoic acid (243 mg, 93%) was obtained by the same method as in Reaction 337-2 using 3-but-3-enyloxy-4-butyl-benzoic acid but-3-enyl ester (318 mg, 1.05 mmol) as a raw material.

¹H-NMR (400 MHz, CD₃OD) δ 7.53 (1H, dd, J = 7.6, 1.7 Hz), 7.50 (1H, d, J = 1.5 Hz), 7.19 (1H, d, J = 7.8 Hz), 6.01-5.91 (1H, m), 5.21-5.15 (1H, m), 5.11-5.07 (1H, m), 4.08 (2H, t, J = 6.1 Hz), 2.65 (2H, t, J = 7.6 Hz), 2.59-2.54 (2H, m), 1.61-1.53 (2H, m), 1.40-1.31 (2H, m), 0.94 (3H, t, J = 7.3 Hz).

(Reaction 361-6)




[2313] N-(2-Allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-4-butyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl-ethyl]-3,N-dimethyl-benzamide was obtained by the same method as in Reaction 359-9 using 3-but-3-enyloxy-4-butyl-benzoic acid (117 mg, 0.471 mmol) and 1-(2-{4-[(2-allyloxy-ethyl)-methyl-carbamoyl]-2-methyl-phenyl}-ethanesulfonyl)-4-amino-piperidine-4-carboxylic amide as starting materials.

MS (ESI) m/z = 677 (M-H)⁻;

HPLC retention time: 3.23 min (analysis condition LCMS-C-1).

(Reaction 361-7)



[2314] A macrocyclic olefin compound (Compound 1433) (E/Z = 1:2) was obtained by the same method as in Reaction 338-1 (using Hoveyda-Grubbs 2nd generation as a catalyst) using N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-4-butyl-

phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3,N-dimethyl-benzamide as a starting material.

[2315] MS (ESI) m/z = 651 (M+H)⁺;


HPLC retention time: 3.13 min (analysis condition LCMS-C-1).

Example 362

Compound 1434

[2316]

(Reaction 362-1)



[2317] A saturated macrocyclic compound (Compound 1434) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1433) as a starting material.

MS (ESI) m/z = 653 (M+H)⁺;


HPLC retention time: 1.12 min (analysis condition LCMS-F-1).

Example 363

Compound 1435


[2318]

(Reaction 363-1)




[2319] 3-But-3-enyloxy-benzoic acid but-3-enyl ester was obtained by the same method as in Reaction 337-1 using 3-hydroxy-benzoic acid as a raw material.

¹H-NMR (400 MHz, CDCl₃) δ 7.62 (1H, br d, J = 7.8 Hz), 7.55 (1H, br s), 7.33 (1H, t, J = 8.1 Hz), 7.09 (1H, br d, J = 8.3 Hz), 5.96-5.82 (2H, m), 5.20-5.10 (4H, m), 4.37 (2H, t, J = 6.8 Hz), 4.06 (2H, t, J = 6.8 Hz), 2.60-2.50 (4H, m).

(Reaction 363-2)

[2320] 3-But-3-enyloxy-benzoic acid was obtained by the same method as in Reaction 337-2 using 3-but-3-enyloxy-benzoic acid but-3-enyl ester as a raw material.

¹H-NMR (400 MHz, CD₃OD) δ 7.61-7.58 (1H, m), 7.53-7.52 (1H, m), 7.36 (1H, t, J = 8.1 Hz), 7.15-7.12 (1H, m), 5.99-5.88 (1H, m), 5.20-5.14 (1H, m), 5.11-5.07 (1H, m), 4.06 (2H, t, J = 6.6 Hz), 2.57-2.51 (2H, m).


(Reaction 363-3)

[2321] N-(2-Allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide was obtained by the same method as in Reaction 359-9 using 3-but-3-enyloxy-benzoic acid and 1-(2-{4-[(2-allyloxy-ethyl)-methyl-carbamoyl]-2-methyl-phenyl}-ethanesulfonyl)-4-amino-piperidine-4-carboxylic amide as starting materials.

MS (ESI) m/z = 621 (M-H)-;

HPLC retention time: 2.85 min (analysis condition LCMS-C-1).

(Reaction 363-4)



[2322] A macrocyclic olefin compound (Compound 1435) E/Z = 1:2) was obtained by the same method as in Reaction 338-1 (using Hoveyda-Grubbs 2nd generation as a catalyst) using N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide as a starting material.

MS (ESI) m/z = 595 (M+H)+;


HPLC retention time: 2.70 min (analysis condition LCMS-C-1).

Example 364

Compound 1436

[2323]

(Reaction 364-1)




[2324] A solution of 3-fluoro-5-trifluoromethyl-benzoic acid (400 mg, 1.92 mmol) in DMF (2 ml) was added dropwise to a suspension of sodium hydride (60% oily, 235 mg, 5.88 mmol) and pent-4-en-1-ol (506 mg, 5.88 mmol) in DMF (12 ml), and the mixture was stirred at 60°C. Further, this mixture was irradiated in a microwave apparatus (150°C, 20 min). The reaction solution was poured into 0.2 N aqueous hydrochloric acid and then extracted with ethyl acetate. The organic layer was washed with saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give 3-pent-4-enyloxy-5-trifluoromethyl-benzoic acid (290 mg, 55%).

¹H-NMR (CDCl₃) δ 7.94 (1H, s), 7.77 (1H, s), 7.37 (1H, s), 5.91-5.81 (1H, m), 5.11-5.06 (1H, m), 5.05-5.02 (1H, m), 4.07 (2H, t, J = 6.3 Hz), 2.30-2.24 (2H, m), 1.97-1.90 (2H, m);

MS (ESI) m/z = 273 (M-H)-;

HPLC retention time: 2.50 min (analysis condition LCMS-C-1).

(Reaction 364-2)




[2325] N-(2-Allyloxy-ethyl)-3,N-dimethyl-4-{2-[4-oxo-2-(3-pent-4-enyloxy-5-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide was obtained by the same method as in Reaction 359-9 using 3-pent-4-enyloxy-5-trifluoromethyl-benzoic acid and 1-(2-{4-[(2-allyloxy-ethyl)-methyl-carbamoyl]-2-methylphenyl}-ethanesulfonyl)-4-amino-piperidine-4-carboxylic amide as starting materials.

MS (ESI) $m/z = 705 (M+H)^+$

HPLC retention time: 1.12 min (analysis condition LCMS-F-1).

(Reaction 364-3)



[2326] A macrocyclic olefin compound (Compound 1436) was obtained by the same method as in Reaction 338-1 (using Hoveyda-Grubbs 2nd generation as a catalyst) using N-(2-allyloxy-ethyl)-3,N-dimethyl-4-{2-[4-oxo-2-(3-pent-4-enyloxy-5-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide as a starting material.

MS (ESI) $m/z = 677 (M+H)^+$;


HPLC retention time: 1.07 min (analysis condition LCMS-F-1).

Example 365

Compound 1437

[2327]

(Reaction 365-1)



[2328] A saturated macrocyclic compound (Compound 1437) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1436) as a starting material.

MS (ESI) m/z = 679 (M+H)⁺;


HPLC retention time: 1.11 min (analysis condition LCMS-F-1).

Example 366

Compound 1438

[2329]

(Reaction 366-1)




[2330] 3-Methyl-4-[2-(4-oxo-piperidine-1-sulfonyl)-ethyl]-benzoic acid (1.23 g, 3.78 mmol) was suspended in toluene (20.0 ml). Di-tert-butoxymethyl-dimethyl-amine (3.63 ml, 15.12 mmol) was added and the mixture was stirred at 80°C for 30 minutes. Thereafter, di-tert-butoxymethyl-dimethyl-amine (2.70 ml, 11.34 mmol) was added again and the mixture was stirred at 80°C for 30 minutes. After completion of the reaction, the reaction solution was left to cool and diluted with ethyl acetate. The organic layer was then washed with an aqueous sodium bicarbonate solution and saline. The organic layer was dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (hexane:ethyl acetate) to give 3-methyl-4-[2-(4-oxo-piperidine-1-sulfonyl)-ethyl]-benzoic acid tert-butyl ester as a white solid (1.01 g, 70.0%).

MS (ESI) m/z = 382 (M+H)⁺;


HPLC retention time: 2.54 min (analysis condition LCMS-B-1)

(Reaction 366-2)




[2331] 3-Methyl-4-[2-(4-oxo-piperidine-1-sulfonyl)-ethyl]-benzoic acid tert-butyl ester (969.2 mg, 2.54 mmol), ammonium acetate (469.9 mg, 6.10 mmol) and potassium cyanide (330.9 mg, 5.08 mmol) were dissolved in methanol (12.0 ml), and the mixture was stirred at 65°C for one hour. After completion of the reaction, the mixture was left to cool and an aqueous sodium bicarbonate solution was added, followed by extraction with ethyl acetate. The organic layer was


washed with saline, and then dried over anhydrous magnesium sulfate and concentrated under reduced pressure to give 4-[2-(4-amino-4-cyano-piperidine-1-sulfonyl)-ethyl]-3-methyl-benzoic acid tert-butyl ester as an amorphous (1.07 g).
MS (ESI) m/z = 408 (M+H)⁺;
HPLC retention time: 2.54 min (analysis condition LCMS-F-1).

(Reaction 366-3)

[2332] 4-[2-(4-Amino-4-cyano-piperidine-1-sulfonyl)-ethyl]-3-methyl-benzoic acid tert-butyl ester (500.0 mg, 1.23 mmol), 3-benzyloxy-5-trifluoromethyl-benzoic acid (436.1 mg, 1.47 mmol) and DIPEA (0.321 ml, 1.85 mmol) were dissolved in DMF (5.50 ml) HATU (561.2 mg, 1.47 mmol) was added and the mixture was stirred at room temperature for one hour. The reaction solution was diluted with diethyl ether, and the organic layer was washed with a 1 M aqueous hydrochloric acid solution, an aqueous sodium bicarbonate solution and saline. The organic layer was dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (hexane-ethyl acetate) to give 4-[2-[4-(3-benzyloxy-5-trifluoromethyl-benzoylamino)-4-cyano-piperidine-1-sulfonyl]-ethyl]-3-methyl-benzoic acid tert-butyl ester as a yellow amorphous (796.2 mg, 94.4% in two steps).
MS (ESI) m/z = 685 (M+H)⁺;
HPLC retention time: 1.16 min (analysis condition LCMS-F-1).

(Reaction 366-4)


[2333] 4-[2-[4-(3-Benzyloxy-5-trifluoromethyl-benzoylamino)-4-cyano-piperidine-1-sulfonyl]-ethyl]-3-methyl-benzoic acid tert-butyl ester (1.40 g, 2.04 mmol) was dissolved in DMSO (0.188 ml, 2.65 mmol) and methanol (7.00 ml). A 1 M aqueous sodium hydroxide solution (0.204 ml, 0.204 mmol) and aqueous hydrogen peroxide (30%, 0.265 ml, 2.65 mmol) were added under ice-cooling. The mixture was warmed to room temperature and stirred as such for two hours. After completion of the reaction, an aqueous sodium thiosulfate solution and an aqueous ammonium chloride solution were added, followed by extraction with ethyl acetate. The organic layer was washed with saline, dried over anhydrous magnesium sulfate and then concentrated under reduced pressure to give 4-[2-[4-(3-benzyloxy-5-trifluoromethyl-benzoylamino)-4-carbamoyl-piperidine-1-sulfonyl]-ethyl]-3-methyl-benzoic acid tert-butyl ester as a pale yellow amorphous (1.48 g).
MS (ESI) m/z = 704 (M+H)⁺;
HPLC retention time: 1.15 min (analysis condition LCMS-F-1).

(Reaction 366-5)

[2334] 4-{2-[4-(3-Benzyloxy-5-trifluoromethyl-benzoylamino)-4-carbamoyl-piperidine-1-sulfonyl]-ethyl}-3-methyl-benzoic acid tert-butyl ester (1.23 g, 1.75 mmol) was dissolved in methanol (17.5 ml). A 1 M aqueous sodium hydroxide solution (1.75 ml, 1.75 mmol) was added and the mixture was stirred at 60°C for six hours. After completion of the reaction, the reaction solution was left to cool and an aqueous ammonium chloride solution was added, followed by extraction with ethyl acetate. The organic layer was washed with saline, dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The residue obtained by concentration was purified by column chromatography (hexane:ethyl acetate) to give 4-{2-[2-(3-benzyloxy-5-trifluoromethylphenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid tert-butyl ester as a pale yellow amorphous (952.1 mg, yield in two steps: 79.3%).

MS (ESI) m/z = 686 (M+H)+;


HPLC retention time: 1.19 min (analysis condition LCMS-F-1).

(Reaction 366-6)

[2335] 4-{2-[2-(3-Benzyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid tert-butyl ester (1.14 g, 1.66 mmol) was dissolved in THF (8.0 ml). DMAP (60.8 mg, 0.498 mmol) and di-tert-butyl dicarbonate (725.6 mg, 3.32 mmol) were added and the mixture was stirred at room temperature for one hour. Thereafter, di-tert-butyl dicarbonate (181.0 mg, 0.830 mmol) was further added and the mixture was stirred at room temperature for 30 minutes. After completion of the reaction, the reaction solution was diluted with ethyl acetate, and the organic layer was washed with saline. The organic layer was dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (hexane:ethyl acetate) to give 2-(3-benzyloxy-5-trifluoromethyl-phenyl)-8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester as a white solid (1.11 g, 85.1%).

MS (ESI) m/z = 786 (M+H)+;

HPLC retention time: 1.20 min (analysis condition LCMS-F-1).


(Reaction 366-7)

[2336] 2-(3-Benzyloxy-5-trifluoromethyl-phenyl)-8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester (1.11 g, 1.41 mmol) was dissolved in ethyl acetate (45.0 ml)-THF (15.0 ml). Pd-C (222 mg) was added and the mixture was stirred at room temperature and for one hour in a hydrogen atmosphere. After completion of the reaction, the black solid was filtered off through celite, and the filtrate was concentrated under reduced pressure to give 8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-(3-hydroxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester as an amorphous (1.01 g, 100%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.80-7.82 (2H, m), 7.22-7.24 (4H, m), 6.83 (1H, br), 3.68-3.74 (2H, m), 3.11-3.27 (6H, m), 2.50 (3H, s), 2.05-2.14 (2H, m), 1.63-1.71 (2H, m), 1.38 (9H, s);

MS (ESI) m/z = 696 ($\text{M}+\text{H}$) $^+$;

HPLC retention time: 1.13 min (analysis condition LCMS-F-1).


(Reaction 366-8)

[2337] 2-[2-(2-Chloro-ethoxy)-ethoxy]-ethanol (2.00 mL, 13.8 mmol) was dissolved in ethanol (14.0 mL). A solution of methylamine in methanol (40%, 14.0 mL, 138 mmol) and sodium iodide (103 mg, 0.67 mmol) were added, and the mixture was stirred at 60°C for 18 hours and then stirred at 75°C for seven hours in a nitrogen atmosphere. The reaction solution was concentrated under reduced pressure to give 2-[2-(2-methylamino-ethoxy)-ethoxy]-ethanol as a crude product.

[2338] 2-[2-(2-Methylamino-ethoxy)-ethoxy]-ethanol was dissolved in THF (6.88 mL)-water (6.88 mL). Di-tert-butyl dicarbonate (9.01 g, 41.3 mmol) and potassium carbonate (5.71 g, 41.3 mmol) were added at 0°C, and the mixture was stirred at room temperature for 15 hours. A 1 M aqueous hydrochloric acid solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate and saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give {2-[2-(2-hydroxy-ethoxy)-ethoxy]-ethyl}-methyl-carbamate (1.53 g, 42%).

$^1\text{H-NMR}$ (CDCl_3) δ 3.74-3.72 (2H, m), 3.67-3.60 (8H, m), 3.42-3.39 (2H, br m), 2.91 (3H, s), 1.45 (9H, s).

(Reaction 366-9)




20 **[2339]** 8-[2-(4-tert-Butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-(3-hydroxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester (30.0 mg, 43.1 μ mol), {2-[2-(2-hydroxy-ethoxy)-ethoxy]-ethyl}-methyl-carbamic acid tert-butyl ester (22.7 mg, 86.2 μ mol) and triphenylphosphine (22.6 mg, 86.2 μ mol) were dissolved in THF (0.22 mL). TMAD (14.8 mg, 86.2 μ mol) was added and the mixture was stirred at 60°C for 30 minutes in a nitrogen atmosphere. The reaction solution was concentrated under reduced pressure, and the resulting residue was then purified by silica gel column chromatography (hexane-ethyl acetate) to give 2-[3-(2-[2-(2-tert-butoxycarbonyl-methyl-amino)-ethoxy]-ethoxy)-ethoxy]-5-trifluoromethyl-phenyl]-8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester (37.4 mg, 86%).

25 MS (ESI) m/z = 941 (M+H)⁺;
HPLC retention time: 1.18 min (analysis condition LCMS-F-1).


30

(Reaction 366-10)



50 **[2340]** 2-[3-(2-[2-(2-tert-Butoxycarbonyl-methyl-amino)-ethoxy]-ethoxy)-ethoxy]-5-trifluoromethyl-phenyl]-8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester (37.4 mg, 39.7 μ mol) was dissolved in 4 N hydrochloric acid-dioxane (0.79 mL). Water (14.3 μ L, 795 μ mol) was added and the mixture was stirred at room temperature for three hours. The reaction solution was concentrated under reduced pressure to give 3-methyl-4-[2-[2-(3-[2-[2-(2-methylamino-ethoxy)-ethoxy]-ethoxy)-5-trifluoromethyl-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzoic acid hydrochloride (28.0 mg, 98%).

55 MS (ESI) m/z = 685 (M+H)⁺;
HPLC retention time: 1.88 min (analysis condition LCMS-B-1).

(Reaction 366-11)

[2341] 3-Methyl-4-{2-[2-(3-{2-[2-(2-methylamino-ethoxy)-ethoxy]-ethoxy}-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoic acid hydrochloride (26.0 mg, 36.1 μ mol) was dissolved in DMF (7.21 mL). DIPEA (62.8 μ L, 361 μ mol) and HATU (68.6 mg, 180 μ mol) were added and the mixture was stirred at 70°C for two hours. A 3 M aqueous hydrochloric acid solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with water, saturated aqueous sodium bicarbonate and saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (dichloromethane-ethyl acetate) to give a saturated macrocyclic compound (Compound 1438) (20.8 mg, 87%).


$^1\text{H-NMR}$ (CD_3OD) δ 7.93 (1H, s), 7.88 (1H, s), 7.61 (1H, s), 7.32 (1H, s), 7.25-7.22 (2H, m), 7.14 (1H, d, J = 7.4 Hz), 4.15 (2H, t, J = 5.4 Hz), 3.78-3.35 (16H, m), 3.11-3.06 (2H, m), 2.97 (3H, s), 2.37 (3H, s), 1.98 (2H, td, J = 13.0, 4.0 Hz), 1.56 (2H, d, J = 12.9 Hz);

MS (ESI) m/z = 667 ($M+H$) $^+$;

HPLC retention time: 2.25 min (analysis condition LCMS-B-1).

Example 367

Compound 1439

[2342](Reaction 367-1)

[2343] {(4R,5S)-5-[2-(tert-Butyl-diphenyl-silanyloxy)-ethyl]-2,2-dimethyl-[1,3]dioxolan-4-yl}-methanol (159 mg, 0.38 mmol) was dissolved in dichloromethane (1.15 mL). Diisopropylethylamine (200 μ L, 1.15 mmol) and Tf_2O (77.4 μ L, 0.46 mmol) were added at 0°C, and the mixture was stirred at room temperature for one hour in a nitrogen atmosphere. Saturated aqueous ammonium chloride was added to the reaction solution, followed by extraction with ethyl acetate.


The organic layer was washed with saturated aqueous sodium bicarbonate and saturated brine, dried over sodium sulfate and filtered. The filtrate was concentrated under reduced pressure to give trifluoro-methanesulfonic acid (4R,5S)-5-[2-(tert-butyl-diphenyl-silanyloxy)-ethyl]-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl ester as a crude product.

[2344] (2-Hydroxy-ethyl)-methyl-carbamic acid tert-butyl ester (73.9 mg, 0.42 mmol) was dissolved in THF (1.15 mL). Sodium hydride (50%, 22.1 mg, 0.46 mmol) was added at 0°C and the mixture was stirred at 0°C for 10 minutes in a nitrogen atmosphere. Trifluoro-methanesulfonic acid (4R,5S)-5-[2-(tert-butyl-diphenyl-silanyloxy)-ethyl]-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl ester was added to the reaction solution, and the mixture was stirred at room temperature for two hours in a nitrogen atmosphere. Saturated aqueous ammonium chloride was added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate and saturated brine, dried over sodium sulfate and filtered, and the filtrate was concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give (2-[(4R,5S)-5-[2-(tert-butyl-diphenyl-silanyloxy)-ethyl]-2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy]-ethyl)-methyl-carbamic acid tert-butyl ester (82.2 mg, 38%).

MS (ESI) m/z = 572 (M+H)⁺;


HPLC retention time: 1.27 min (analysis condition LCMS-F-1).

(Reaction 367-2)



[2345] (2-[(4R,5S)-5-[2-(tert-Butyl-diphenyl-silanyloxy)-ethyl]-2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy]-ethyl)-methyl-carbamic acid tert-butyl ester (82.2 mg, 0.14 mmol) was dissolved in THF (0.19 mL). TBAF (1.0 M solution in THF, 0.19 mL, 0.19 mmol) was added at 0°C and the mixture was stirred at room temperature for one hour in a nitrogen atmosphere. Saturated aqueous ammonium chloride was added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate and saturated brine, and then dried over sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane-ethyl acetate) to give {2-[(4R,5S)-5-(2-hydroxyethyl)-2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy]-ethyl}-methyl-carbamic acid tert-butyl ester (41.7 mg, 88%).


¹H-NMR (CDCl₃) δ 4.34-4.31 (1H, m), 4.25 (1H, q, J = 6.1 Hz), 3.84-3.74 (2H, m), 3.54-3.47 (6H, m), 2.90 (3H, s), 2.46 (1H, dd, J = 7.8, 3.3 Hz), 1.82-1.78 (2H, br m), 1.45-1.44 (12H, m), 1.35 (3H, s).

(Reaction 367-3)

[2346] 2-(3-(2-((4S,5R)-5-((2-(tert-Butoxycarbonyl(methyl)amino)ethoxy)methyl)-2,2-dimethyl-1,3-dioxolan-4-yl)ethoxy)-5-(trifluoromethyl)phenyl)-8-(4-(tert-butoxycarbonyl)-2-methylphenethylsulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester was obtained by the same method as in Reaction 366-9 using 8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-(3-hydroxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester and {2-[(4R,5S)-5-(2-hydroxy-ethyl)-2,2-dimethyl-[1,3]dioxolan-4-ylmethoxy]-ethyl}-methyl-carbamic acid tert-butyl ester as starting materials.

MS (ESI) m/z = 1012 (M+H)⁺;


HPLC retention time: 1.21 min (analysis condition LCMS-F-1).

(Reaction 367-4)

[2347] 4-[2-(2-{3-[(3S,4R)-3,4-Dihydroxy-5-(2-methylamino-ethoxy)-pentyloxy]-5-trifluoromethyl-phenyl}-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzoic acid hydrochloride was obtained by the same method as in Reaction 366-10 using 2-(3-(2-((4S,5R)-5-((2-(tert-butoxycarbonyl(methyl)amino)ethoxy)methyl)-2,2-dimethyl-1,3-dioxolan-4-yl)ethoxy)-5-(trifluoromethyl)phenyl)-8-(4-(tert-butoxycarbonyl)-2-methylphenethylsulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester as a starting material.

MS (ESI) m/z = 729 (M+H)⁺;

HPLC retention time: 0.80 min (analysis condition LCMS-F-1).

(Reaction 367-5)

[2348] A saturated macrocyclic compound (Compound 1439) was obtained by the same method as in Reaction 366-11 using 4-[2-(2-{3-[(3S,4R)-3,4-dihydroxy-5-(2-methylamino-ethoxy)-pentyloxy]-5-trifluoromethyl-phenyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzoic acid hydrochloride as a starting material.


MS (ESI) m/z = 697 (M+H)⁺;

HPLC retention time: 0.92 min (analysis condition LCMS-F-1).

Example 368


Compound 1440

[2349]

(Reaction 368-1)


[2350] BOC-sarcosine (900 mg, 4.76 mmol) was dissolved in ethanol (10 ml). 2-(2-Aminoethoxy)ethanol (0.477 ml, 4.76 mmol) and 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride n-hydrate (DMT-MM) (1.79 g, 5.71 mmol) were added and the mixture was stirred at room temperature overnight. The reaction solution was concentrated under reduced pressure, and water was added to the resulting residue, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was subjected to silica gel column chromatography (dichloromethane-methanol) to give [2-(2-hydroxy-ethoxy)-ethylcarbamoyl]-methyl]-methyl-carbamic acid tert-butyl ester. This was used in the next reaction without complete purification.

(Reaction 368-2)




[2351] 8-[2-(4-tert-Butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-[3-(2-{2-[2-(tert-butoxycarbonyl-methyl-amino)-acetyl-amino]-ethoxy}-ethoxy)-5-trifluoromethyl-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester was obtained by the same method as in Reaction 366-9 using 8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-(3-hydroxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester and {[2-(2-hydroxy-ethoxy)-ethyl-carbamoyl]-methyl}-methyl-carbamic acid tert-butyl ester as starting materials. This was used in the next reaction without complete purification.

(Reaction 368-3)



[2352] 3-Methyl-4-{2-[2-(3-{2-[2-(2-methylamino-acetyl-amino)-ethoxy]-ethoxy}-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoic acid hydrochloride was obtained by the same method as in Reaction 366-10 using 8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-[3-(2-{2-[2-(tert-butoxycarbonyl-methyl-amino)-acetyl-amino]-ethoxy}-ethoxy)-5-trifluoromethyl-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester (13 mg, 0.0136 mmol) as a starting material. This was used in the next reaction without complete purification.

(Reaction 368-4)

[2353] A saturated macrocyclic compound (Compound 1440) was obtained by the same method as in Reaction 366-11 using 3-methyl-4-{2-[2-(3-{2-[2-(2-methylamino-acetylamino)-ethoxy]-ethoxy}-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-tri-


MS (ESI) $m/z = 680$ ($M+H$) $^{+}$;

HPLC retention time: 0.89 min (analysis condition LCMS-F-1).

Example 369


Compound 1441

[2354]

(Reaction 369-1)

[2355] A mixture of 2-(methylamino)ethanol (5.2 g, 69.2 mmol) and tert-butyl dicarbonate (15.8 g, 72.7 mmol) in methylene chloride (200 mL) was stirred at room temperature for 16 hours. The reaction mixture was, diluted with methylene chloride, and the organic layer was then washed with water, dried over $MgSO_4$ and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give (2-hydroxyethyl)methylcarbamic acid 1,1-dimethylethyl ester (12.1 g, yield 100%).

^1H -NMR (400 MHz, $CDCl_3$) δ 1.47 (9H, m), 2.91 (3H, s), 3.39-3.41 (2H, m), 3.72-3.76 (2H, m).


(Reaction 369-2)

[2356] 4-Dimethylaminopyridine (634 mg, 5.2 mmol) was added to a solution of 2,2,3,3,4,4-hexafluoro-pentane-1,5-diol (1 g, 4.7 mmol) and tert-butyldimethylsilyl chloride (708 mg, 4.7 mmol) in methylene chloride (10 ml), and the mixture was stirred at room temperature for 16 hours. The reaction mixture was concentrated under reduced pressure, and the residue was then purified by silica gel column chromatography to give 5-(tert-butyl-dimethyl-silanyloxy)-2,2,3,3,4,4-

hexafluoro-pentan-1-ol (770 mg, 50%).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.10 (6H, s), 0.90 (9H, m), 4.07-4.11 (4H, m).

(Reaction 369-3)




[2357] (2-Hydroxyethyl)methylcarbamic acid 1,1-dimethylethyl ester (350 mg, 2.0 mmol) was dissolved in pyridine (2 ml). Mesyl chloride (0.229 ml, 2.1 mmol) was added at room temperature, and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was diluted with methylene chloride, and the organic layer was washed with a saturated aqueous sodium bicarbonate solution, dried over MgSO_4 and concentrated under reduced pressure to give methanesulfonic acid 2-(tert-butoxycarbonyl-methyl-amino)-ethyl ester. This was used in the next reaction without complete purification.

[2358] 5-(tert-Butyl-dimethyl-silanyloxy)-2,2,3,3,4,4-hexafluoro-pentan-1-ol (400 mg, 1.23 mmol) was dissolved in dimethylformamide (2 ml). Sodium hydride (51.5 mg, 1.29 mmol) was added at room temperature, and the mixture was stirred at room temperature for 30 minutes. A solution of methanesulfonic acid 2-(tert-butoxycarbonyl-methyl-amino)-ethyl ester obtained above in dimethylformamide (0.5 ml) was then added at room temperature, and the mixture was stirred at room temperature overnight. The reaction mixture was extracted with ethyl acetate, and the organic layer was then washed with water, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give {2-[5-(tert-butyl-dimethyl-silanyloxy)-2,2,3,3,4,4-hexafluoro-pentyloxy]-ethyl}-methyl-carbamic acid tert-butyl ester. This was used in the next reaction without complete purification.

[2359] {2-[5-(tert-Butyl-dimethyl-silanyloxy)-2,2,3,3,4,4-hexafluoro-pentyloxy]-ethyl}-methyl-carbamic acid tert-butyl ester obtained above was dissolved in a 1 M solution of tetrabutylammonium fluoride in tetrahydrofuran (0.2 mL), and the mixture was reacted at room temperature for one hour. The reaction mixture was concentrated under reduced pressure, and the residue was then purified by silica gel column chromatography to give [2-(2,2,3,3,4,4-hexafluoro-5-hydroxy-pentyloxy)-ethyl]-methyl-carbamic acid tert-butyl ester (70 mg). This was used in the next reaction without complete purification.

[2360] [2-(2,2,3,3,4,4-Hexafluoro-5-hydroxy-pentyloxy)-ethyl]-methyl-carbamic acid tert-butyl ester obtained above was dissolved in methylene chloride (0.5 ml). Triethylamine (0.0528 ml, 0.38 mmol) and trifluoromethanesulfonyl chloride (0.0212 ml, 0.20 mmol) were added at room temperature, and the mixture was stirred at room temperature for 64 hours. The reaction mixture was extracted with ethyl acetate, and the organic layer was washed with saturated brine, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give Trifluoro-methanesulfonic acid 5-[2-(tert-butoxycarbonyl-methyl-amino)-ethoxy]-2,2,3,3,4,4-hexafluoro-pentyl ester (66 mg).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.44 (9H, s), 2.89 (3H, s), 3.35-3.42 (m, 2H), 3.65-3.72 (m, 2H), 3.90-3.94 (m, 2H), 4.73-4.83 (m, 2H).


(Reaction 369-4)

[2361] 4-{2-[2-(3-Benzyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid tert-butyl ester (952.0 mg, 1.39 mmol) was dissolved in ethyl acetate (15.0 ml)-THF (5.0 ml). Pd-C (190.4 mg) was added and the mixture was stirred at room temperature for one hour in a hydrogen atmosphere. After completion of the reaction, the precipitated solid was dissolved by adding dichloromethane and methanol, and the remaining black solid was then filtered off through celite. The filtrate was concentrated under reduced pressure to give 4-{2-[2-(3-hydroxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid tert-butyl ester as a white solid (776.2 mg, 93.8%).

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ 11.7 (1H, br), 10.6 (1H, br), 7.66-7.80 (4H, m), 7.39 (1H, d, J 8.0 Hz), 7.25 (1H, s), 3.59-3.70 (2H, m), 3.27-3.43 (4H, m), 3.04-3.08 (2H, m), 2.39 (3H, s), 1.79-1.88 (2H, m), 1.59 (2H, m), 1.55 (9H, s);

MS (ESI) m/z = 596 ($M+H$) $^+$;


HPLC retention time: 1.12 min (analysis condition LCMS-F-1).

(Reaction 369-5)

[2362] 4-{2-[2-(3-Hydroxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid tert-butyl ester (77.0 mg, 0.13 mmol) and trifluoro-methanesulfonic acid 5-[2-(tert-butoxycarbonylmethyl-amino)-ethoxy]-2,2,3,3,4,4-hexafluoropentyl ester (58.0 mg, 0.13 mmol) were dissolved in DMF (1 ml). Potassium carbonate (53.8 mg, 0.39 mmol) was added and the mixture was stirred at 60°C overnight. The reaction mixture was extracted with ethyl acetate, and the organic layer was then washed with saturated brine, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give 4-{2-[2-(3-[5-[2-(tert-butoxycarbonylmethyl-amino)-ethoxy]-2,2,3,3,4,4-hexafluoropentyloxy]-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid tert-butyl ester as a white solid.

nyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3-methyl-benzoic acid tert-butyl ester (57 mg, 46%).
MS (ESI) m/z = 947 (M+H)+.

(Reaction 369-6)



[2363] 4-[2-[2-(3-[5-[2-(tert-Butoxycarbonyl-methyl-amino)-ethoxy]-2,2,3,3,4,4-hexafluoro-pentyloxy]-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3-methyl-benzoic acid tert-butyl ester (56 mg, 0.059 mmol) was dissolved in water (0.1 ml) and 4 N hydrochloric acid-dioxane (1 ml), and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated under reduced pressure, and the residue was then dissolved in DMF (5 ml). Triethylamine (0.053 ml, 0.384 mmol) and O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (144 mg, 0.384 mmol) were added at room temperature, and the mixture was heated with stirring at 70°C for two hours. The reaction mixture was extracted with ethyl acetate, and the organic layer was washed with water, dried over MgSO_4 and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give a saturated macrocyclic compound (Compound 1441) (12 mg, 21%).

MS (ESI) m/z = 773 (M+H)+;


HPLC retention time: 6.52 min (analysis condition LCMS-A-2).

Example 370

Compound 1442

[2364]


(Reaction 370-1)



[2365] 8-[2-(4-tert-Butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-(3-but-3-ynyloxy-5-trifluoromethylphenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester (37.9 mg, 71%) was obtained by the same method as in Reaction 366-9 using 8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-(3-hydroxy-5-trifluoromethylphenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester and 3-butyne-1-ol as starting materials. MS (ESI) m/z = 592 (M-Boc-tBu)+;

HPLC retention time: 3.79 min (analysis condition LCMS-A-1).

(Reaction 370-2)




[2366] 4 M hydrochloric acid-dioxane (1.80 ml) and water (0.0173 ml, 0.960 mmol) were added to 8-[2-(4-tert-butoxycarbonyl-2-methyl-phenyl)-ethanesulfonyl]-2-(3-but-3-ynyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-3-carboxylic acid tert-butyl ester (35.9 mg, 0.048 mmol), and the mixture was stirred at room temperature for five hours. The reaction solution was concentrated under reduced pressure to give 4-{2-[2-(3-but-3-ynyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid as a white solid.

MS (ESI) m/z = 592 (M+H)+;

HPLC retention time: 0.91 min (analysis condition LCMS-F-1).


(Reaction 370-3)



[2367] 4-{2-[2-(3-But-3-ynyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzoic acid, (3-azido-propyl)-methylamine hydrochloride (15.3 mg, 0.101 mmol) and DIPEA (0.044 ml, 0.255 mmol) were dissolved in DMF (0.400 ml). HATU (0.038 mg, 0.101 mmol) was added and the mixture was stirred at room temperature for one hour. The reaction solution was diluted with ethyl acetate, and the organic layer was then washed with a 1 M aqueous hydrochloric acid solution, an aqueous sodium bicarbonate solution and saline. The organic layer was dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane-ethyl acetate) to give N-(3-azido-propyl)-4-{2-[2-(3-but-3-ynyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzamide as a colorless oily substance.

MS (ESI) m/z = 688 (M+H)+;

HPLC retention time: 1.05 min (analysis condition LCMS-F-1).

(Reaction 370-4)

[2368] N-(3-Azido-propyl)-4-{2-[2-(3-but-3-ynoxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide was dissolved in acetonitrile (17.0 ml)-THF (4.0 ml). DMSO (0.022 ml), DIPEA (0.012 ml, 0.069 mmol), 2,6-lutidine (0.0053 ml, 0.046 mmol) and copper(I) iodide (13.2 mg, 0.069 mmol) were added, and the mixture was stirred at room temperature overnight. Thereafter, copper(I) iodide (13.2 mg, 0.069 mmol) was further added and the mixture was stirred at room temperature for one hour. The reaction solution was concentrated under reduced pressure. Ethyl acetate was added and the precipitated solid was filtered off. The filtrate was washed with a 1 M aqueous hydrochloric acid solution, an aqueous sodium bicarbonate solution and saline. The organic layer was dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The resulting residue was purified by column chromatography (dichloromethane-ethyl acetate and dichloromethane-methanol) to give a macrocyclic compound (Compound 1442) as a white solid (3.2 mg, 20.2% in three steps).

MS (ESI) m/z = 688 (M+H)⁺;


HPLC retention time: 0.94 min (analysis condition LCMS-F-1).

Example 371

Compound 1443

[2369]

(Reaction 371-1)



[2370] {4-[2-(4-Carbamoyl-4-hept-6-enoylamino-piperidine-1-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester was obtained as a crude compound by the same method as in Reaction 359-9 using hept-6-enoic acid and {4-[2-(4-amino-4-carbamoyl-piperidine-1-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester as starting materials.

MS (ESI) m/z = 579 (M+H)⁺;

HPLC retention time: 2.49 min (analysis condition LCMS-B-1).


[2371] {4-[2-(4-Carbamoyl-4-hept-6-enoylamino-piperidine-1-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester obtained above (Crude compound, 1.00 g) was dissolved in ethanol. Lithium hydroxide monohydrate (188 mg, 4.48 mmol) was added and the mixture was stirred at 40°C for 19 hours. A saturated aqueous ammonium chloride solution and water were added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with water and saturated brine, and then dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-carbamic acid tert-butyl ester (821 mg, 1.47 mmol).

¹H-NMR (400 MHz, CDCl₃) δ 7.85 (1H, br s), 6.92 (2H, s), 5.83-5.73 (1H, m), 5.04-4.96 (2H, m), 3.80-3.74 (2H, m), 3.45-3.38 (2H, m), 3.22 (3H, s), 3.16-3.12 (2H, m), 3.01-2.97 (2H, m), 2.44 (2H, t, J = 7.6 Hz), 2.34 (6H, s), 2.10 (2H, q, J = 7.2 Hz), 2.02-1.95 (2H, m), 1.71-1.59 (4H, m), 1.46 (9H, s);

MS (ESI) m/z = 561 (M+H)⁺;

HPLC retention time: 2.61 min (analysis condition LCMS-A-1).

(Reaction 371-2)



[2372] {4-[2-(2-Hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-methyl-carbamic acid tert-butyl ester (820 mg, 1.46 mmol) was dissolved in methylene chloride (16 ml). Trifluoroacetic acid (10 ml) was added and the mixture was stirred at room temperature for two hours. The reaction solution was concentrated, and a saturated aqueous sodium bicarbonate solution was added to the residue. This mixture was extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-hex-5-enyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (686 mg) as a crude compound.

[2373] The resulting crude product 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-hex-5-enyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one (160 mg, 0.35 mmol) was dissolved in methylene chloride. 9-Decenoyl chloride (0.35 mmol) (prepared by allowing oxalyl chloride and a catalytic amount of DMF to act on 9-decenoic acid in methylene chloride) and triethylamine (0.195 ml, 1.4 mmol) were added, and the mixture was stirred at room temperature for 17 hours. A saturated aqueous ammonium chloride solution and water were added to the reaction solution, and the resulting mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, dried over anhydrous sodium sulfate and then concentrated under reduced pressure. The resulting residue was purified by silica gel column chromatography to give dec-9-enoic acid {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-amide (Compound 1443) (100 mg, 48% in two steps).

MS (ESI) m/z = 613 (M+H)⁺;


HPLC retention time: 5.80 min (analysis condition LCMS-C-1).

Example 372

Compound 1444

[2374]

(Reaction 372-1)




15 **[2375]** A macrocyclic olefin compound (Compound 1444) was obtained by the same method as in Reaction 338-1 using dec-9-enoic acid {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-methyl-amide as a starting material.
 MS (ESI) m/z = 585 ($M+H$)⁺;
 HPLC retention time: 5.15 min (analysis condition LCMS-B-2).

20 Example 373

Compound 1445

25 **[2376]**

(Reaction 373-1)




40 **[2377]** A saturated macrocyclic compound (Compound 1445) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1444) as a starting material.
 MS (ESI) m/z = 587 ($M+H$)⁺;
 HPLC retention time: 2.70 min (analysis condition LCMS-A-1).

45 Example 374

Compound 1446

50 **[2378]**

(Reaction 374-1)


[2379] Hept-6-enoic acid {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-methyl-amide (Compound 1446) was obtained by the same method as in Reaction 371-2 using 6-heptenoic acid and 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-hex-5-enyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one as starting materials.

MS (ESI) $m/z = 571$ (M+H)+;

HPLC retention time: 2.49 min (analysis condition LCMS-A-1).

Example 375

Compound 1447

[2380](Reaction 375-1)

[2381] A macrocyclic olefin compound (Compound 1447) was obtained by the same method as in Reaction 338-1 using hept-6-enoic acid {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-methyl-amide as a starting material.


MS (ESI) $m/z = 543$ (M+H)+;

HPLC retention time: 2.25 min (analysis condition LCMS-A-1).

Example 376

Compound 1448

[2382]

(Reaction 376-1)

[2383] A saturated macrocyclic compound (Compound 1448) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1447) as a starting material.


MS (ESI) $m/z = 545 (M+H)^+$;

HPLC retention time: 2.34 min (analysis condition LCMS-A-1).

Example 377

Compound 1449

[2384]

(Reaction 377-1)

[2385] A saturated macrocyclic compound (Compound 1449) was obtained by the same method as in Reaction 339-1 using a macrocyclic olefin compound (Compound 1422) as a starting material.

MS (ESI) $m/z = 678 (M+H)^+$;

HPLC retention time: 2.50 min (analysis condition LCMS-A-1).

Biological Experimental Example

Experimental Example A: In vitro cAMP signal activity of compounds in human PTH1 receptor

Materials and method

(Peptides)

[2386] Human PTH(1-34) and calcitonin were purchased from Peptide Institute, Inc. (Osaka, Japan), dissolved in 10 mM acetic acid to 1 mM and stored in a -80°C freezer.

(Cell culture)

[2387] Cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum

(Hyclone), 100 units/ml penicillin G and 100 µg/ml streptomycin sulfate (Invitrogen Corp) at 37°C in a humidified atmosphere containing 5% CO₂.

[2388] cAMP signal transduction analysis utilized LLC-PK1 cells not expressing the PTH1 receptor, and HKRK-B7 cells, that is, LLC-PK1 cells overexpressing the human PTH1 receptor at 9.5×10^5 receptors/cell (Takasu et al., J. Bone. Miner. Res. 14:11-20, 1999).

(cAMP stimulation)

[2389] HKRK-B7 or LLC-PK1 cells were seeded in a 96-well plate at 1×10^5 cells/well and incubated overnight. On the following day, 50 µl of cAMP assay buffer (DMEM, 2 mM IBMX, 0.2 mg/ml bovine serum albumin, 35 mM Hepes-NaOH, pH 7.4) containing human PTH(1-34) or Compound was added and the plate was placed in a 37°C incubator. The cells were incubated for 20 minutes. After removing the medium, the cells were washed with 100 µl of cAMP assay buffer once. The plate was placed on dry ice powder to freeze the cells and then removed from the dry ice. The cells were lysed with 40 µl of 50 mM HCl and frozen again on dry ice. The amount of intracellular cAMP produced was measured using a commercially available cAMP EIA kit (Biotrack cAMP EIA system, GE health care).

[2390] The compounds of the present invention demonstrated a significant cAMP response in HKRK-B7 cells. Table 195 shows percentage values obtained by dividing the amount of cAMP produced by the compound of the present invention in HKRK-B7 cells at 1×10^{-3} M (*at 3×10^{-4} M for Compound 15) by the amount of cAMP produced by hPTH(1-34) as a positive control at 100 nM. The degree of cAMP response in LLC-PK1 cells was lower than the degree in HKRK-B7 cells.

[Table 195]

Compound	cAMP production activity (%)
1	42
2	1.0
3	20
4	11
5	2.5
6	4.2
7	13
8	2.6
10	2.4
11	16
12	12
13	31
14	38
15	1.8*
16	41
17	39
18	44
19	35
20	38
21	42
22	43
23	41
24	18
25	18
26	36
27	42
28	31
29	28
30	26
31	41
32	26
33	4.4
34	16
35	56
36	3.5
37	54
38	52
39	25
40	19
41	21
42	27
43	39
44	25
45	22
46	17
47	45
48	25
49	29
50	26

Compound	cAMP production activity (%)
51	38
52	23
53	33
54	35
55	55
56	55
57	35
58	39
59	56
60	34
61	43
62	56
63	51
64	45
65	65
66	57
67	53
68	51
69	61
70	31
71	3.7
72	1.5
73	66
74	48
75	75
76	73
77	69
78	31
79	8.2
80	71
81	57
82	65
83	41
84	51
85	76
86	66
87	6.7
88	97
89	96
90	88
91	73
92	97
93	111
94	62
95	76
96	70
97	4.3
98	75
99	80

Compound	cAMP production activity (%)
100	61
101	49
102	23
103	80
104	79
105	78
106	94
107	110
108	62
109	62
110	25
111	93
112	77
113	111
114	105
115	81
116	94
117	61
118	73
119	60
120	3.1
121	5.8
122	3.0
123	23
124	64
125	62
126	79
127	72
128	57
129	3.7
130	65
131	72
133	82
134	77
135	67
136	72
137	50
138	49
139	64
140	77
141	32
142	63
143	4.5
144	59
145	129
146	122
147	105
148	79
149	56

Compound	cAMP production activity (%)
150	62
151	53
152	47
153	59
154	82
155	45
156	64
157	70
158	62
159	96
160	65
161	69
162	43
163	41
164	45
165	37
166	56
167	44
168	69
169	71
170	77
171	36
172	102
173	71
174	68
175	73
176	74
177	28
178	29
179	49
180	60
181	19
182	38
183	68
184	37
185	33
186	51
187	12
188	70
189	54
190	61
191	57
192	52
193	65
194	56
195	36
196	66
197	41
198	31

Compound	cAMP production activity (%)
199	46
200	37
201	56
202	27
203	25
204	110
205	47
206	70
207	36
208	22
209	24
210	79
211	60
212	59
213	74
214	84
215	81
216	84
217	41
218	72
219	60
220	80
221	103
222	43
223	85
224	54
225	47
226	83
227	87
228	8.3
229	68
230	66
231	96
232	69
233	13
234	78
235	49
236	40
237	74
238	90
239	80
240	49
241	44
242	75
243	80
244	83
245	34
246	39
247	81

Compound	cAMP production activity (%)
248	66
249	71
250	62
251	28
252	28
253	54
254	97
255	64
256	67
257	42
258	87
259	67
260	24
261	70
262	26
263	41
264	69
265	55
266	81
267	42
268	99
269	43
270	55
271	57
272	67
273	55
274	74
275	72
276	63
277	38
278	59
279	67
280	57
281	92
282	29
283	63
284	82
285	65
286	54
287	58
288	82
289	99
290	76
291	66
292	58
293	38
294	106
295	95
296	65

Compound	cAMP production activity (%)
297	91
298	63
299	83
300	73
301	72
302	95
303	76
304	47
305	73
306	45
307	58
308	72
309	72
310	76
311	67
312	49
313	63
314	68
315	26
316	20
317	62
318	52
319	31
320	33
321	55
322	75
323	53
324	30
325	61
326	76
327	84
328	41
329	33
330	23
331	55
332	90
333	87
334	34
335	28
336	28
337	17
338	60
339	66
340	67
341	62
342	93
343	13
344	35
345	21

Compound	cAMP production activity (%)
346	28
347	23
348	5.6
349	11
350	5.5
351	7.4
352	19
353	120
354	27
355	84
356	78
357	78
358	71
359	65
360	62
361	82
362	97
363	67
364	78
365	81
366	85
367	50
368	38
369	43
370	66
371	72
372	51
373	70
374	79
375	57
376	74
377	64
378	60
379	59
380	75
381	72
382	36
383	72
384	61
385	94
386	86
387	97
388	84
389	75
390	22
391	26
392	83
393	44
394	59

Compound	cAMP production activity (%)
395	88
396	85
397	126
398	52
399	64
400	76
401	83
402	85
403	51
404	88
405	7.7
406	100
407	49
408	117
409	55
410	96
411	54
412	54
413	56
414	69
415	56
416	66
417	97
418	84
419	96
420	31
421	68
422	14
423	42
424	2.5
425	17
426	16
428	8.8
429	38
430	23
431	14
432	5.1
433	18
434	25
435	17
436	28
437	45
438	14
439	33
441	24
442	2.5
444	46
445	83
446	49

Compound	cAMP production activity (%)
447	89
448	64
449	94
450	56
451	96
452	58
453	16
454	19
455	31
456	28
457	25
458	77
459	55
460	18
461	51
462	47
463	28
464	54
465	66
466	23
467	60
468	90
469	47
470	90
471	111
472	104
474	89
475	84
476	61
477	31
478	33
479	15
480	44
481	59
482	38
483	41
484	47
485	7.7
486	59
487	49
488	48
489	37
490	26
491	58
492	37
493	50
494	66
495	21
496	24

Compound	cAMP production activity (%)
497	38
498	53
499	40
500	61
501	59
502	14
503	67
504	60
505	61
506	83
507	43
508	24
509	70
510	57
511	29
512	49
513	33
514	75
515	113
516	73
517	58
518	68
519	72
520	28
521	55
522	82
523	89
524	90
525	14
526	83
527	93
528	55
529	68
530	18
531	69
532	80
533	69
534	39
535	83
536	72
537	3.9
538	133
539	80
540	22
541	79
542	66
543	76
544	80
545	86

Compound	cAMP production activity (%)
546	75
547	52
548	88
549	86
550	124
551	92
552	64
553	80
554	82
555	40
556	19
557	50
558	74
559	72
560	66
561	52
562	74
563	69
564	68
565	45
566	19
567	24
568	39
569	4.2
570	66
571	39
572	36
573	35
574	42
575	57
576	95
577	74
578	13
579	55
580	25
581	75
582	104
583	85
584	24
585	39
586	82
587	53
588	77
589	22
590	70
591	34
592	87
593	28
594	69

Compound	cAMP production activity (%)
595	63
596	40
597	51
598	74
599	59
600	67
601	64
602	3.0
603	69
604	21
605	54
606	28
607	6.5
608	20
609	46
610	85
611	82
612	62
613	44
614	25
615	46
616	94
617	96
618	121
619	61
620	112
621	80
622	134
623	123
624	36
625	47
626	53
627	5.3
628	48
629	87
630	4.1
631	65
632	51
633	37
634	29
635	93
636	88
637	38
638	46
639	101
640	26
641	85
642	87
643	94

Compound	cAMP production activity (%)
644	75
645	55
646	99
647	104
648	61
649	40
650	55
651	54
652	63
653	67
654	50
655	74
656	14
657	124
658	84
659	46
660	60
661	45
662	23
663	27
664	77
665	54
666	51
667	40
668	40
669	58
670	123
671	81
672	47
673	27
674	68
675	68
676	69
677	69
678	86
679	65
680	101
681	55
682	81
683	74
684	101
685	46
686	22
687	25
688	55
689	27
690	86
691	69
692	101

Compound	cAMP production activity (%)
693	103
694	77
695	78
696	132
697	60
698	62
699	101
700	121
701	140
702	84
704	68
705	76
706	90
707	124
708	38
709	58
710	76
711	64
712	16
713	55
714	36
715	20
716	62
717	111
718	74
719	77
720	82
721	92
722	60
723	95
724	74
725	58
726	75
727	52
728	87
729	45
730	74
731	54
732	45
733	104
734	47
735	32
736	16
737	96
738	79
739	47
740	123
741	91
742	50

Compound	cAMP production activity (%)
743	54
744	19
745	67
746	120
747	55
748	61
749	77
750	87
751	83
752	79
753	104
754	89
755	74
756	79
757	79
758	98
759	79
760	93
761	104
764	124
765	101
766	88
767	83
768	79
769	55
770	105
771	80
772	69
773	86
774	80
775	70
776	79
777	71
778	57
779	53
780	48
781	30
782	14
783	50
784	84
785	92
786	57
787	81
788	142
789	157
790	88
791	6.1
792	110
793	124

Compound	cAMP production activity (%)
794	76
795	97
796	64
797	88
798	101
799	7.1
800	77
801	103
802	100
803	103
804	78
812	105
817	111
818	79
820	98
821	82
822	99
823	103
824	140
825	114
826	90
827	78
828	92
829	79
830	73
831	4.7
832	84
833	34
834	76
835	50
836	56
837	66
838	75
839	57
840	98
841	45
842	81
843	77
844	86
845	68
846	47
847	71
848	77
849	124
850	82
851	83
852	58
853	63
854	80

Compound	cAMP production activity (%)
855	82
856	81
857	89
858	100
859	26
860	50
861	36
862	55
863	67
864	100
865	8.9
866	47
867	71
868	77
869	65
870	63
872	109
873	77
874	61
875	65
876	22
877	35
878	25
879	70
880	68
881	48
882	70
883	56
884	59
885	58
886	68
887	58
888	86
889	26
890	61
891	8.6
892	51
893	14
894	85
895	90
896	83
897	85
898	4.3
899	12
900	90
901	1.9
902	67
903	56
904	69

Compound	cAMP production activity (%)
905	75
906	78
907	85
908	74
909	78
910	64
911	71
912	98
913	81
914	68
915	61
916	59
917	69
918	63
919	68
920	70
921	59
922	84
923	84
924	76
925	69
926	102
927	80
928	51
929	76
930	92
931	72
932	66
933	60
934	87
935	112
936	98
937	120
938	97
939	111
940	86
941	21
942	31
943	74
944	71
945	77
946	102
947	89
948	68
949	92
950	59
951	93
952	95
953	77

Compound	cAMP production activity (%)
954	81
955	79
956	87
957	17
958	49
959	77
960	84
961	92
962	86
963	16
964	119
965	115
966	82
967	44
968	69
969	45
970	112
971	83
972	89
973	112
974	111
975	74
976	73
977	80
978	91
979	145
980	85
981	106
982	96
983	91
984	133
985	120
986	96
987	54
988	50
989	86
990	87
991	64
992	65
993	64
994	87
995	98
996	85
997	74
998	92
999	61
1000	86
1001	64
1002	50

Compound	cAMP production activity (%)
1003	67
1006	52
1007	20
1008	73
1009	70
1010	96
1011	17
1012	87
1013	48
1014	84
1015	83
1016	92
1017	101
1018	109
1019	72
1020	81
1021	137
1022	105
1023	92
1024	66
1025	114
1026	68
1027	82
1028	75
1029	104
1030	115
1031	111
1032	88
1033	22
1034	54
1035	77
1036	82
1037	87
1038	111
1039	103
1040	111
1041	38
1042	102
1043	99
1044	86
1045	106
1046	101
1047	82
1048	96
1050	92
1051	85
1052	62
1053	70
1054	80

Compound	cAMP production activity (%)
1055	84
1056	94
1057	100
1058	133
1059	116
1060	58
1061	55
1062	65
1063	72
1064	73
1065	83
1066	83
1067	69
1068	68
1069	79
1070	69
1071	60
1072	54
1073	66
1074	66
1075	69
1076	88
1077	74
1078	74
1079	91
1080	81
1081	53
1082	22
1083	113
1084	13
1085	100
1086	151
1087	97
1088	95
1089	99
1090	118
1091	118
1092	89
1093	100
1094	100
1095	105
1096	93
1097	90
1098	88
1099	91
1100	76
1101	110
1102	10
1103	5.4

Compound	cAMP production activity (%)
1104	16
1106	58
1107	24
1108	99
1109	29
1110	92
1111	79
1112	76
1113	99
1114	95
1115	140
1116	106
1117	88
1118	79
1119	136
1120	124
1121	118
1122	150
1123	122
1124	119
1125	93
1126	106
1127	91
1128	119
1129	102
1130	100
1131	96
1132	80
1133	113
1134	50
1135	84
1136	112
1137	82
1138	77
1139	86
1140	55
1141	83
1142	68
1143	78
1144	124
1145	102
1146	107
1147	112
1148	100
1149	98
1150	107
1151	105
1152	0.8
1153	106

Compound	cAMP production activity (%)
1154	115
1155	83
1156	77
1157	44
1158	103
1159	87
1160	84
1161	84
1162	112
1163	101
1164	4.7
1165	4.3
1166	144
1167	115
1168	27
1169	50
1170	24
1171	28
1172	73
1173	85
1174	91
1175	81
1176	82
1178	84
1179	65
1180	73
1181	98
1182	109
1183	90
1184	108
1185	102
1186	110
1187	75
1188	99
1189	104
1190	108
1191	66
1192	100
1193	86
1194	62
1195	82
1196	76
1197	74
1198	88
1199	77
1200	73
1201	77
1202	91
1203	90

Compound	cAMP production activity (%)
1204	83
1205	83
1206	88
1207	112
1208	65
1209	94
1210	86
1211	99
1212	96
1213	80
1214	79
1215	74
1216	61
1217	68
1218	90
1219	67
1220	80
1221	75
1222	77
1223	54
1224	88
1225	90
1226	51
1227	77
1228	68
1229	56
1230	64
1231	88
1232	106
1233	78
1234	114
1235	98
1236	99
1237	96
1238	73
1239	91
1240	95
1241	101
1242	106
1243	77
1244	96
1245	115
1246	85
1247	70
1248	81
1249	62
1250	67
1251	56
1252	72

Compound	cAMP production activity (%)
1253	81
1254	87
1255	66
1256	72
1257	98
1258	116
1259	101
1260	81
1261	99
1262	90
1263	73
1264	77
1265	89
1266	96
1267	74
1268	33
1269	92
1270	61
1271	92
1272	71
1273	81
1274	81
1275	89
1276	140
1277	95
1278	95
1279	113
1280	74
1281	95
1282	63
1283	18
1284	2.5
1285	67
1286	35
1287	64
1288	54
1289	17
1290	6.3
1291	48
1292	14
1293	84
1294	76
1295	73
1296	64
1297	98
1298	117
1299	87
1300	81
1301	49

Compound	cAMP production activity (%)
1302	95
1303	102
1304	107
1305	138
1306	159
1307	116
1308	102
1309	109
1310	104
1311	79
1312	105
1313	87
1314	78
1315	76
1316	2.9
1317	3.4
1318	19
1319	5.8
1320	10
1321	63
1322	80
1323	78
1324	1.0
1325	113
1326	84
1327	92
1328	93
1329	85
1330	9.2
1331	96
1332	119
1333	109
1334	116
1335	97
1336	133
1337	44
1338	84
1339	86
1340	84
1341	83
1342	114
1343	98
1344	94
1345	107
1346	113
1347	87
1348	95
1349	98
1350	22

Compound	cAMP production activity (%)
1351	100
1352	78
1353	111
1354	128
1355	118
1356	14
1357	13
1358	153
1359	165
1360	121
1361	104
1362	48
1363	80
1364	84
1365	108
1366	103
1367	58
1368	83
1369	30
1370	64
1371	84
1372	36
1373	44
1374	33
1375	23
1376	42
1377	35
1378	116
1379	86
1380	109
1381	102
1382	93
1383	96
1384	78
1385	92
1386	92
1387	68
1388	59
1389	67
1390	69
1391	116
1392	89
1393	84
1394	82
1395	68
1396	133
1397	24
1398	77
1399	20

Compound	cAMP production activity (%)
1400	49
1401	73
1402	69
1403	66
1404	64
1405	51
1406	7.3
1407	6.6
1408	5.1
1409	13
1410	51
1411	38
1412	7.2
1413	57
1414	47
1415	49
1416	48
1417	49
1418	36
1419	70
1420	20
1421	49
1422	15
1423	17
1424	29
1425	32
1426	5.4
1427	72
1428	67
1429	8.9
1430	13
1431	15
1432	18
1433	38
1434	54
1435	19
1436	54
1437	57
1438	40
1439	18
1440	10
1441	5.0
1442	9.3
1443	61
1444	12
1445	62
1446	88
1447	15
1448	9.4

Compound	cAMP production activity (%)
1449	18



Industrial Applicability

[2391] The present invention provides a compound having a PTH-like effect. The present invention also provides a medicine for use in the prevention and/or treatment of osteoporosis, fracture, osteomalacia, arthritis, thrombocytopenia,

hypoparathyroidism, hyperphosphatemia, tumoral calcinosis or the like.

Claims

1. A compound represented by the following general formula (1) or (2):



or a pharmacologically acceptable salt of a compound represented by formula (1) or (2);
[wherein in general formula (1)

W is selected from:

- 1) a single bond,
- 2) C1-C10 alkylene optionally containing a carbonyl group, wherein the alkylene is optionally substituted with a halogen atom(s) and/or a hydroxyl group(s),
- 3) C2-C10 alkenylene optionally substituted with a halogen atom(s),
- 4) C2-C10 alkynylene,
- 5) arylene optionally substituted with a halogen atom(s),
- 6) heteroarylene optionally substituted with a halogen atom(s),
- 7) C1-C10 heteroalkylene optionally substituted with a halogen atom(s),
- 8) -NH-, -NHCH₂- or -NHCH₂CH₂-,
- 9) cycloalkylene and
- 10) -(cycloalkylene)-CH₂-;

X is selected from the following bond or groups:

- 1) a single bond,


- 2) C1-C10 alkylene optionally substituted with a halogen atom(s) or cycloalkyl,
- 3) C2-C10 alkenylene optionally substituted with a halogen atom(s),
- 4) C2-C10 alkynylene optionally substituted with a halogen atom(s),
- 5) C1-10 oxyalkylene optionally substituted with a halogen atom(s) and
- 6) -NR₄₇-

wherein R₄₇ is selected from:

- i) a hydrogen atom and
- ii) C1-C10 alkyl optionally substituted with a halogen atom(s);

Y is selected from:

- 1) an oxygen atom,
- 2) a sulfur atom and
- 3) =NR₃₇,
- or
- 4) Y is -NR₃₈R₃₉ shown in the following formula (A):



which can be tautomeric;

R₃₇ is selected from:

- 1) hydrogen,
- 2) hydroxy and
- 3) C1-C10 alkoxy;

R₃₈ and R₃₉ are independently selected from hydrogen or C1-C10 alkyl optionally substituted with cycloalkyl, or R₃₈ and R₃₉ may be bonded to each other to form a ring selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholinyl, and the ring is optionally substituted with C1-C10 alkyl;
 m represents an integer of 0 to 2;
 n represents an integer of 0 to 2;

R₁ is selected from:

- 1) hydrogen,
- 2) cycloalkyl optionally substituted with a group(s) selected from R₄, R₅ and R₆,
- 3) a heterocycle optionally substituted with a group(s) selected from R₂₅, R₄, R₅ and R₆,
- 4) aryl optionally substituted with a group(s) selected from R₃, R₄, R₅ and R₆ and
- 5) heteroaryl optionally substituted with a group(s) selected from R₂₅, R₄, R₅ and R₆;

R₃ is selected from:

- 1) -CONR₇R₈,
- 2) -OR₉,
- 3) -NR₉R₁₀,
- 4) -N(R₉)COR₁₁,
- 5) -N(R₉)SO₂R₁₂,
- 6) -SO₂R₁₅,

- 7) C1-10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, -COR₁₆ and -NR₁₃R₁₄,
 8) heteroaryl optionally having C1-10 alkyl and/or C1-10 alkoxy as a substituent(s) and
 9) -N(R₉)CSR₁₁;

R₄ is selected from:

- 1) a halogen atom,
- 2) cyano,
- 3) nitro,
- 4) amino,
- 5) -NHCOR₂₆,
- 6) C1-C10 alkyl optionally substituted with a group(s) independently selected from hydroxycarbonyl, C1-C10 alkoxycarbonyl and aminocarbonyl,
- 7) C1-C10 haloalkyl,
- 8) C1-C10 alkoxy,
- 9) C1-C10 haloalkylcarbonyl,
- 10) -COR₁₆,
- 11) C1-C10 hydroxyalkyl and
- 12) C1-C10 heteroalkyl;

R₅ is selected from a halogen atom, C1-C10 alkyl, C1-C10 haloalkyl and C1-C10 alkoxy;

R₆ is selected from a halogen atom, C1-C10 alkyl and C1-C10 haloalkyl;

R₇ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from amino and C1-C10 alkylamino,
- 3) C1-C10 hydroxyalkyl,
- 4) C1-C10 haloalkyl,
- 5) C1-C10 heteroalkyl,
- 6) C1-C10 heteroalkyl optionally substituted with a group(s) selected from a hydroxyl group, C1-C10 alkylamino and C2-C10 alkenyl,
- 7) aryl,
- 8) heteroaryl,
- 9) aryl C1-C10 alkyl,
- 10) a heterocycle optionally substituted with C1-C10 alkyl,
- 11) -(CH₂)_LCOR₁₆ (wherein L represents an integer of 1 to 4),
- 12) C1-C10 alkoxy,
- 13) C2-C10 alkenyl and
- 14) -NR₄₀R₄₁;

R₄₀ and R₄₁ are independently selected from hydrogen, C1-C10 alkyl and C1-C10 alkylcarbonyl, or R₄₀ and R₄₁ may be bonded to each other to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholinyl, and the heterocycle is optionally substituted with C1-C10 alkyl;

R₈ is selected from hydrogen and C1-C10 alkyl optionally substituted with a halogen atom(s) and/or a hydroxyl group(s);

R₇ and R₈ may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from O, N, S, SO and SO₂ and the heterocycle optionally contains carbonyl, and the heterocycle is optionally substituted with a substituent(s) independently selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally having C1-C10 alkylamino as a substituent(s),
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,

- 7) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
 8) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
 9) a heterocycle optionally substituted with C1-C10 alkyl,
 10) heteroaryl optionally substituted with C1-C10 alkyl,
 11) heterocyclyl C1-C10 alkyl,
 12) -COR₁₆,
 13) -NR₁₉R₂₀,
 14) -SO₂R₂₁,
 15) C1-C10 alkoxy-C1-C10 alkyl optionally having a hydroxyl group(s) as a substituent(s) and
 16) C1-C10 hydroxyalkoxy, wherein the hydrogen atom of the hydroxyl group is optionally replaced by C1-C10 hydroxyalkyl, and

the heterocycle may further form a spiro ring together with a 4- to 6-membered heterocycle, and the bonded 4- to 6-membered heterocycle optionally contains O and N as ring-forming elements in addition to carbon atoms, and the carbon atom(s) may be oxidized to form carbonyl, and the 4- to 6-membered heterocycle is optionally further substituted with C1-C10 alkyl;

R₁₆ is selected from:

- 1) a hydroxyl group,
- 2) C1-C10 alkoxy,
- 3) NR₁₇R₁₈ and
- 4) C1-C10 alkyl optionally substituted with a substituent(s) selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, C1-C10 alkoxy-carbonyl or C1-C10 alkylamino;

R₁₇ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) selected from aryl, amino, C1-C10 alkylamino, C1-C10 alkylcarbonylamino and a hydroxyl group,
- 3) heteroaryl and
- 4) C1-C10 alkoxy;

R₁₈ is selected from hydrogen, C1-C10 alkyl and C1-C10 hydroxyalkyl;

R₁₇ and R₁₈ may be bonded to each other to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl, a halogen atom and C1-C10 alkoxy-carbonyl;

R₁₉ is selected from hydrogen, C1-C10 alkyl, C1-C10 haloalkyl, C1-C10 alkylcarbonyl, C1-C10 hydroxyalkyl, C1-C10 aminoalkyl, C1-C10 alkoxy-carbonyl and C1-C10 heteroalkyl;

R₂₀ is selected from hydrogen and C1-C10 alkyl;

R₁₉ and R₂₀ may be bonded to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom;

R₂₁ is selected from:

- 1) C1-C10 alkyl optionally substituted with aryl,
- 2) amino,
- 3) C1-C10 alkylamino and
- 4) aryl optionally substituted with C1-C10 alkyl;

R₉ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from R₂₃,
- 3) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 4) cycloalkyl optionally substituted with a halogen atom(s) or a hydroxyl group(s),

- 5) a heterocycle optionally substituted with a group(s) independently selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy, C1-C10 alkoxy carbonyl, amino and a halogen atom,
 6) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom and a hydroxyl group,
 7) heteroaryl optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy carbonyl and a halogen atom and
 8) cycloalkenyl optionally substituted with a group(s) selected from C1-C10 alkoxy, C1-C10 alkylamino, amino, a hydroxyl group and a halogen atom, wherein the cycloalkenyl optionally contains a carbonyl group;

R_{23} is independently selected from:

- 1) a halogen atom,
- 2) a hydroxyl group,
- 3) a C1-C10 alkylcarbonyloxy group,
- 4) $-COR_{16}$,
- 5) amino,
- 6) C1-C10 alkylamino,
- 7) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy carbonyl and a halogen atom and
- 8) cyano;

R_{10} is selected from:

- 1) hydrogen and
- 2) C1-C10 alkyl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group and aryl; R_9 and R_{10} may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from N, O, S, SO, SO_2 , carbonyl and thiocarbonyl, and the heterocycle is optionally substituted with a substituent(s) independently selected from R_{24} ;

R_{24} is independently selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from C1-C10 alkylamino and C1-C10 alkylcarbonylamino,
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 8) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 9) a heterocycle optionally substituted with C1-C10 alkyl,
- 10) heteroaryl,
- 11) heterocyclyl C1-C10 alkyl,
- 12) $-COR_{16}$,
- 13) $-NR_{19}R_{20}$ and
- 14) $-SO_2R_{21}$;

R_{11} is selected from:

- 1) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a hydroxyl group,
- ii) $-NR_{17}R_{18}$,
- iii) a C1-C10 alkoxy group,
- iv) a halogen atom,

v) C1-C10 alkoxy carbonyl,

vi) aminocarbonyl and

vii) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, amino, C1-C10 alkylamino and -COR₂₂,

5

2) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, C1-C10 alkoxy, amino, C1-C10 alkylamino and -COR₂₂,

3) cycloalkyl optionally substituted with a halogen atom(s),

10

4) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy carbonyl and a halogen atom,

5) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino and a hydroxyl group,

6) amino,

15

7) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino, hydroxycarbonyl and a hydroxyl group and

8) C2-C10 alkenyl;

R₂₂ is selected from C1-C10 alkoxy, a hydroxyl group, amino and C1-C10 alkylamino;

20

R₁₂ is selected from:

1) C1-C10 alkyl,

2) amino and

3) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from amino, C1-C10 alkylamino and a hydroxy group;

25

R₁₃ is selected from:

1) hydrogen,

30

2) C1-C10 alkyl,

3) C1-C10 alkylcarbonyl, wherein the alkyl is optionally substituted with a hydroxyl group(s),

4) C1-C10 alkoxy carbonyl,

5) aminocarbonyl,

6) C1-C10 alkylaminocarbonyl and

35

7) heterocyclic carbonyl optionally substituted with C1-C10 alkyl;

R₁₄ is selected from:

1) hydrogen and

40

2) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino;

R₁₃ and R₁₄ may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from O, N, S, SO and SO₂, and the heterocycle optionally contains carbonyl, and the heterocycle is optionally substituted with C1-C10 alkyl;

45

R₁₅ is selected from:

1) C1-C10 alkyl and

2) -NR₃₅R₃₆;

50

R₃₅ is selected from:

1) hydrogen,

2) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

55

i) a halogen atom,

ii) a hydroxyl group,

iii) C1-C10 alkylcarbonylamino,

- iv) $-\text{COR}_{16}$,
- v) amino,
- vi) C1-C10 alkylamino,
- vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),
- viii) heteroaryl optionally substituted with a C1-C10 alkyl group(s) and
- ix) a heterocycle,

- 3) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 4) cycloalkyl optionally substituted with a group(s) independently selected from a halogen atom and a hydroxyl group,
- 5) a heterocycle optionally substituted with a group(s) independently selected from C1-C10 alkyl, a halogen atom and aryl C1-C10 alkyl,
- 6) heteroaryl optionally substituted with C1-C10 alkyl and
- 7) C1-C10 alkylcarbonyl;

R_{36} is selected from:

- 1) hydrogen and
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group and aryl;

R_{35} and R_{36} may be bonded to each other to form a ring selected from azetidiny, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom;

R_{25} is selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) aryl,
- iii) heteroaryl,
- iv) a heterocycle optionally substituted with a C1-C10 alkyl group(s),
- v) $-\text{COR}_{16}$,
- vi) $-\text{NR}_{13}\text{R}_{14}$ and
- vii) $-\text{SO}_2\text{R}_{21}$,

- 3) C1-C10 heteroalkyl optionally substituted with a hydroxyl group(s),
- 4) C1-C10 hydroxyalkyl, wherein each hydroxyl group may be independently substituted with a group(s) selected from C1-C10 alkyl, aryl C1-C10 alkyl and C1-C10 alkylcarbonyl,
- 5) $-\text{COR}_{16}$,
- 6) $-\text{SO}_2\text{R}_{21}$,
- 7) aryl and
- 8) cyano;

R_2 is selected from:

- 1) C1-C10 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a substituent(s) independently selected from R_{42} ,
- 2) C2-C10 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a substituent(s) independently selected from R_{42} ,
- 3) C2-C10 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a substituent(s) independently selected from R_{42} ,
- 4) cycloalkyl optionally substituted with a group(s) independently selected from:
 - i) a halogen atom,
 - ii) C2-C10 alkenyl or C1-C10 alkyl,

- iii) aryl optionally substituted with 1 to 3 substituents independently selected from C1-C10 alkyl, a halogen atom, C1-C10 alkoxy, C1-C10 alkylamino and C1-C10 alkylcarbonyl,
- iv) cycloalkyl,
- v) C2-C10 alkenyl optionally substituted with halogen,
- vi) C1-C10 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with a halogen atom(s),
- vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),
- viii) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom or C1-C10 alkoxy optionally substituted with a halogen atom(s),
- ix) C2-C10 alkynyl and
- x) $-\text{Si}(\text{R}_{43})_3$,

5) a heterocycle, wherein the heterocycle is optionally substituted with a group(s) independently selected from:

- i) a C1-C10 alkyl group,
- ii) C1-C10 alkylcarbonyl, wherein the alkyl group is optionally substituted with R_{27} ,
- iii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- iv) heteroarylcarbonyl,
- v) C1-C10 alkoxy carbonyl, wherein the alkyl group is optionally substituted with a group(s) independently selected from a halogen atom, aryl and C1-C10 alkoxy,
- vi) aryloxy carbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) and/or C1-C10 alkyl,
- vii) $-\text{CONR}_{28}\text{R}_{29}$,
- viii) $-\text{SO}_2\text{R}_{21}$,
- ix) a halogen atom,
- x) cycloalkylcarbonyl optionally fused with an aryl group and
- xi) C2-C10 alkenylcarbonyl, wherein the alkenyl group is optionally substituted with aryl, wherein the aryl is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl or C1-C10 alkoxy,

6) aryl optionally substituted with a group(s) independently selected from R_{44} ,

7) heteroaryl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) C1-C10 alkyl and
- iii) C1-C10 alkoxy;

8) C1-C10 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a substituent(s) independently selected from R_{42} ,

9) $-\text{S}(\text{O})_q\text{R}_{43}$ (wherein q is an integer of 0 to 2) and

10) cycloalkenyl optionally substituted with C1-C10 alkyl;

R_{44} is selected from:

- 1) a halogen atom,
- 2) cyano,
- 3) C1-C10 alkyl optionally substituted with a group(s) independently selected from:

- i) a hydroxyl group,
- ii) $-\text{OR}_{26}$,
- iii) cyano,
- iv) aryloxy optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl optionally substituted with a halogen atom(s) or C1-C10 alkoxy optionally substituted with a halogen atom(s) and
- v) a halogen atom,

- 4) cycloalkyl optionally substituted with a group(s) independently selected from a halogen atom or C1-C10 alkyl optionally substituted with a halogen atom(s),
- 5) C1-C10 alkoxy optionally substituted with a halogen atom(s) or a C2-C6 alkenyl group(s),
- 6) -COR₃₀,
- 7) C1-C10 alkylcarbonylamino,
- 8) C1-C10 alkoxycarbonylamino, wherein the alkoxy group is optionally substituted with aryl,
- 9) C1-C10 heteroalkyl optionally substituted with a halogen atom(s),
- 10) aryl optionally substituted with a substituent(s) independently selected from:

- i) a halogen atom,
- ii) C1-C10 alkyl,
- iii) C1-C10 alkoxy and
- iv) aryl optionally substituted with aryl optionally substituted with C1-C10 alkyl,

- 11) heteroaryl optionally substituted with a C1-C10 alkyl group(s),
- 12) -SO₂R₄₃,
- 13) -SOR₄₃,
- 14) C1-C10 alkylthio optionally substituted with a halogen atom(s),
- 15) -Si(R₄₃)₃ and
- 16) -SF₅;

R₄₂ is selected from:

- 1) hydrogen,
- 2) aryl optionally substituted with a group(s) independently selected from C1-C10 alkyl optionally substituted with halogen, a halogen atom and C1-C10 alkoxy,
- 3) hydroxycarbonyl,
- 4) C1-C10 alkoxycarbonyl,
- 5) aminocarbonyl,
- 6) C1-C10 alkylaminocarbonyl,
- 7) C1-C10 alkoxycarbonylamino,
- 8) amino,
- 9) a hydroxyl group and
- 10) oxetane, tetrahydrofuran or tetrahydropyran optionally substituted with C1-C10 alkyl;

R₄₃ represents a C1-C10 alkyl group;

R₂₆ is aryl, or C1-C10 alkyl optionally substituted with a halogen atom(s);

R₂₇ is selected from:

- 1) aryl optionally substituted with a group independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- 2) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with aryl,
- 3) a hydroxyl group,
- 4) amino,
- 5) C1-C10 alkylamino,
- 6) hydroxycarbonyl,
- 7) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl and/or aryl, and
- 8) heteroaryloxy;

R₂₈ is selected from hydrogen or C1-C10 alkyl optionally substituted with aryl;

R₂₉ is selected from hydrogen or C1-C10 alkyl optionally substituted with aryl;

R₂₈ and R₂₉ may be bonded to form a ring selected from azetidyl, pyrrolidinyl, piperidinyl, piperazinyl and morpholinyl, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl and a halogen atom;

R₃₀ is selected from a hydroxyl group, C1-C10 alkoxy and -NR₃₁R₃₂;

R₃₁ and R₃₂ are independently selected from:

- 1) hydrogen,

- 2) C1-C10 alkyl optionally substituted with aryl and
3) aryl;


R_{31} and R_{32} may be bonded to form a ring selected from azetidiny, pyrrolidiny, piperidiny, piperaziny and morpholiny, and the ring is optionally substituted with a group(s) selected independently of each other from C1-C10 alkyl, a halogen atom and C1-C10 alkoxy carbonyl; and

R_{33} and R_{34} are independently selected from:

- 1) hydrogen and
2) C1-C10 alkyl]; and

[wherein in general formula (2) W, Y, m, n, R_4 , R_5 , R_6 , R_7 , R_{11} , R_{16} , R_{19} , R_{20} , R_{21} , R_{33} , R_{34} and R_{44} are as defined for the formula (1) ;


U represents a bond, C1-C10 alkylene or any group selected from groups represented by the following formula:



A is selected from O, NH and CH_2 ;

R_{46} is selected from hydrogen or R_{44} ;

T is selected from aryl and heteroaryl; V is selected from:




E is a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) selected from O, N, S, SO and SO_2 , and the heterocycle is optionally substituted with a group(s) selected from:

- 1) hydrogen,
- 2) a halogen atom,
- 3) C1-C10 alkyl optionally having a hydroxyl group(s) independently selected from C1-C10 alkylamino, a halogen atom and a hydroxyl group,
- 4) a hydroxyl group,
- 5) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 6) aryl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) C1-C10 heteroalkyl optionally substituted with a group(s) independently selected from a halogen atom,

- a hydroxyl group, amino and C1-C10 alkylamino,
 8) a heterocycle optionally substituted with C1-C10 alkyl,
 9) heteroaryl optionally substituted with C1-C10 alkyl,
 10) heterocyclyl C1-C10 alkyl,
 11) $-\text{COR}_{16}$,
 12) $-\text{NR}_{19}\text{R}_{20}$ and
 13) $-\text{SO}_2\text{R}_{21}$;

Z is a divalent group selected from:

- 1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s) and/or a hydroxyl group(s), wherein the carbon atom(s) may be oxidized to form carbonyl;
 2) C2-C10 alkenylene or C2-C10 heteroalkenylene optionally substituted with a halogen atom(s) and/or a hydroxyl group(s), wherein the carbon atom(s) may be oxidized to form carbonyl; and
 3) a group selected from:



G is a divalent group selected from:

- 1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s); and
 2) C2-C10 alkenylene or C2-C10 heteroalkenylene optionally substituted with a halogen atom(s);

J is a divalent group selected from:

- 1) C1-C10 alkylene or C1-C10 heteroalkylene optionally substituted with a halogen atom(s); and
 2) C2-C10 alkenylene or C2-C10 heteroalkenylene - optionally substituted with a halogen atom(s);

B is selected from a heterocycle or heteroaryl; and R_{45} is selected from hydrogen or C1-C10 alkyl]; and wherein in the compound represented by general formula (1) or (2) the term "heteroalkyl" means a group containing 1 to 5 heteroatoms in the alkyl backbone, the term "heteroalkylene" means a group containing 1 to 5 heteroatoms in the alkylene backbone, and the term "heteroalkenylene" means a group containing 1 to 5 heteroatoms in the alkenylene backbone.

2. The compound or a pharmacologically acceptable salt thereof according to claim 1, wherein

W is selected from:

- 1) a single bond,
 2) C1-C10 alkylene optionally containing a carbonyl group, wherein the alkylene is optionally substituted with a halogen atom(s) or hydroxy,
 3) C2-C10 alkenylene optionally substituted with a halogen atom(s),
 4) C2-C10 alkynylene,
 5) arylene,
 6) heteroarylene,
 7) $-\text{NH}-$, $-\text{NHCH}_2-$ or $-\text{NHCH}_2\text{CH}_2-$,
 8) cycloalkylene and
 9) $-(\text{cycloalkylene})-\text{CH}_2-$;

X is selected from the following bond or groups:

- 1) a single bond,
- 2) C1-C10 alkylene optionally substituted with cycloalkyl,
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) C1-C10 oxyalkylene;

R₁ is selected from:

- 1) hydrogen,
- 2) cycloalkyl optionally substituted with a group selected from R₄,
- 3) a heterocycle optionally substituted with a group(s) selected from R₂₅ and R₄,
- 4) aryl optionally substituted with a group(s) selected from R₃, R₄, R₅ and R₆ and
- 5) heteroaryl optionally substituted with a group(s) selected from R₂₅, R₄ and R₅;

R₉ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from R₂₃,
- 3) cycloalkyl optionally substituted with a halogen atom(s) or a hydroxyl group(s),
- 4) a heterocycle optionally substituted with a group(s) selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxy, C1-C10 alkoxycarbonyl, amino and a halogen atom,
- 5) C1-C10 heteroalkyl optionally substituted with a group(s) selected from a halogen atom and a hydroxyl group,
- 6) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl, C1-C10 alkylcarbonyl, C1-C10 alkoxycarbonyl and a halogen atom and
- 7) cycloalkenyl optionally substituted with a group(s) selected from C1-C10 alkoxy, C1-C10 alkylamino, amino, 1 to 3 hydroxyl groups and 1 to 4 halogen atoms, wherein the cycloalkenyl optionally contains a carbonyl group;

R₁₀ is selected from: -

- 1) hydrogen and
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group and aryl;

R₉ and R₁₀ may be bonded to form a 4- to 7-membered heterocycle optionally containing an additional element(s) or group(s) independently selected from N, O, S, SO, SO₂, carbonyl and thiocarbonyl, and the heterocycle is optionally substituted with a substituent(s) independently selected from R₂₄;

R₂₄ is selected from:

- 1) a halogen atom,
- 2) C1-C10 alkyl optionally substituted with a group(s) independently selected from C1-C10 alkylamino and C1-C10 alkylcarbonylamino,
- 3) C1-C10 haloalkyl,
- 4) a hydroxyl group,
- 5) C1-C10 hydroxyalkyl,
- 6) C1-C10 alkoxy optionally substituted with a group(s) independently selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 7) aryl optionally substituted with a group(s) selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 8) C1-C10 heteroalkyl optionally substituted with 1 to 2 groups selected from a halogen atom, a hydroxyl group, amino and C1-C10 alkylamino,
- 9) -COR₁₆ and
- 10) -NR₁₉R₂₀;

R₁₁ is selected from:

- 1) C1-C10 alkyl optionally substituted with 1 to 3 substituents independently selected from:

- i) a hydroxyl group,
- ii) $-NR_{17}R_{18}$,
- iii) a C1-C10 alkoxy group,
- iv) a halogen atom,
- v) C1-C10 alkoxycarbonyl and
- vi) aminocarbonyl,

- 2) aryl,
- 3) aryl C1-C10 alkyl,
- 4) cycloalkyl optionally substituted with a halogen atom(s),
- 5) a heterocycle optionally substituted with C1-C10 alkyl,
- 6) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino and a hydroxyl group,
- 7) amino,
- 8) C1-C10 alkylamino, wherein the alkyl group is optionally substituted with a group(s) independently selected from C1-C10 alkylcarbonylamino, amino, C1-C10 alkylamino, hydroxycarbonyl and a hydroxyl group and
- 9) C2-C10 alkenyl; and

R_{33} and R_{34} are hydrogen,
wherein R_3 , R_4 , R_5 , R_6 , R_{16} , R_{17} , R_{18} , R_{19} , R_{20} , R_{23} and R_{25} are as defined in claim 1, respectively.

3. The compound or a pharmacologically acceptable salt thereof according to claim 1 or 2, wherein

W is selected from:

- 1) a single bond,
- 2) C1-C10 alkylene optionally substituted with a halogen atom(s),
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) heteroarylene,

X is selected from the following bond or groups:

- 1) a single bond,
- 2) C1-C10 alkylene,
- 3) C2-C10 alkenylene,
- 4) C2-C10 alkynylene and
- 5) C1-C10 oxyalkylene, wherein the oxyalkylene is bonded to a 1,3,8-triaza-spiro[4.5]dec-1-en-4-one ring or a 1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione ring through a carbon atom in the oxyalkylene;

R_1 is selected from:

- 1) aryl optionally substituted with a group(s) selected from R_3 , R_4 and R_5 and
- 2) heteroaryl optionally substituted with a group(s) selected from R_{25} and R_4 ;

R_3 is selected from:

- 1) $-CONR_7R_8$,
- 2) $-OR_9$,
- 3) $-NR_9R_{10}$,
- 4) $-N(R_9)COR_{11}$,
- 5) $-N(R_9)SO_2R_{12}$,
- 6) $-SO_2R_{15}$,
- 7) C1-C10 alkyl optionally substituted with a group(s) selected from $-COR_{16}$ and $-NR_{13}R_{14}$ and
- 8) $-N(R_9)CSNH_2$;

R_4 is selected from:

- 1) halogen,
- 2) cyano,
- 3) amino,
- 4) C1-C10 alkyl,
- 5) C1-C10 haloalkyl,
- 6) C1-C10 alkoxy,
- 7) C1-C10 haloalkylcarbonyl,
- 8) -COR₁₆ and
- 9) C1-C10 heteroalkyl;

R₂ is selected from:

- 1) C1-C10 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a group selected from R₄₂,
- 2) C2-C10 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a group selected from R₄₂,
- 3) C2-C10 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a group selected from R₄₂,
- 4) cycloalkyl optionally substituted with a group(s) independently selected from:

- i) a halogen atom,
- ii) C2-C10 alkenyl or C1-C10 alkyl,
- iii) aryl optionally substituted with a group(s) independently selected from C1-C10 alkyl, a halogen atom and C1-C10 alkoxy,
- iv) cycloalkyl,
- v) C2-C10 haloalkenyl or C1-C10 haloalkyl,
- vi) C1-C10 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with a halogen atom(s),
- vii) C1-C10 alkoxy optionally substituted with a halogen atom(s),
- viii) C1-C10 alkyl substituted with C1-C10 alkoxy, wherein the alkyl and/or the alkyl in the alkoxy is optionally substituted with a halogen atom(s),
- ix) C2-C10 alkynyl and
- x) -Si(R₄₃)₃,

- 5) a heterocycle, wherein the heterocycle is optionally substituted with a group(s) selected from:

- i) a C1-C10 alkyl group,
- ii) C1-C10 alkylcarbonyl, wherein the alkyl group is optionally substituted with R₂₇,
- iii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- iv) heteroarylcarbonyl,
- v) C1-C10 alkoxycarbonyl, wherein the alkyl group is optionally substituted with a group(s) independently selected from a halogen atom, aryl and C1-C10 alkoxy,
- vi) aryloxycarbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) and/or C1-C10 alkyl,
- vii) -CONR₂₈R₂₉ and
- viii) -SO₂R₂₁,

- 6) aryl optionally substituted with a group(s) independently selected from R₄₄,

- 7) heteroaryl optionally substituted with any of the following groups:

- i) C1-C10 alkyl,

- 8) C1-C10 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a group selected from R₄₂,

- 9) -S(O)_qR₄₃ (wherein q is an integer of 0 to 2) and

- 10) cycloalkenyl optionally substituted with C1-C10 alkyl,

R₄₄ is selected from:

- 1) a halogen atom,
- 2) cyano,
- 3) C1-C10 alkyl optionally substituted with any of the following groups:
 - i) a hydroxyl group,
 - ii) -OR₂₆,
 - iii) cyano and
 - iv) aryloxy optionally substituted with a group(s) selected from a halogen atom, C1-C10 alkyl, C1-C10 haloalkyl or C1-C10 haloalkoxy,
- 4) C1-C10 haloalkyl,
- 5) cycloalkyl optionally substituted with a group(s) selected from a halogen atom and C1-C10 haloalkyl,
- 6) C1-C10 alkoxy optionally substituted with a halogen atom(s) or a C2-C6 alkenyl group(s),
- 7) -COR₃₀,
- 8) C1-C10 heteroalkyl optionally substituted with a halogen atom(s),
- 9) aryl optionally substituted with a group(s) independently selected from:
 - i) C1-C10 alkyl and
 - ii) aryl,
- 10) heteroaryl optionally substituted with a C1-C10 alkyl group(s),
- 11) -SO₂R₄₃,
- 12) C1-C10 alkylthio optionally substituted with a halogen atom(s),
- 13) -Si(R₄₃) and
- 14) -SF₅; and

R₂₇ is selected from:

- 1) aryl optionally substituted with a group(s) independently selected from a halogen atom, C1-C10 alkyl and C1-C10 alkoxy,
- 2) C1-C10 alkoxy, wherein the alkyl group is optionally substituted with aryl,
- 3) heteroaryl optionally substituted with a group(s) independently selected from C1-C10 alkyl and aryl and
- 4) heteroaryloxy,

wherein R₅, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₂₁, R₂₅, R₂₆, R₂₈, R₂₉, R₃₀, R₄₂ and R₄₃ are as defined in claim 1 or 2 from which claim 3 depends, respectively.

4. The compound or a pharmacologically acceptable salt thereof according to any of claims 1 to 3, wherein

W is selected from:

- 1) C1-C6 alkylene optionally substituted with a fluorine atom(s),
- 2) C1-C6 alkenylene and
- 3) thiophene,

X is selected from the following bond or groups:

- 1) a single bond,
- 2) C1-C6 alkylene and
- 3) C1-C6 oxyalkylene optionally substituted with a halogen atom(s), wherein the oxyalkylene is bonded to a 1,3,8-triaza-spiro[4.5]dec-1-en-4-one ring or a 1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione ring through a carbon atom in the oxyalkylene;

Y represents an oxygen atom;

m represents 1; and

n represents 1.

5. The compound or a pharmacologically acceptable salt thereof according to any of claims 1 to 4, wherein

W is selected from:

- 1) ethylene,
- 2) vinylene and
- 3) thiophene,

X represents a single bond;


R₃ is selected from:

- 1) -CONR₇R₈,
- 2) -OR₉,
- 3) -NR₉R₁₀,
- 4) -N(R₉)COR₁₁,
- 5) -N(R₉)SO₂R₁₂,
- 6) -SO₂R₁₅ and
- 7) C1-C6 alkyl optionally substituted with a group(s) selected from -COR₁₆ and -NR₁₃R₁₄;

R₂ is selected from:

- 1) C1-C10 alkyl optionally substituted with a halogen atom(s), wherein the alkyl group is optionally further substituted with a group selected from R₄₂,
- 2) C2-C10 alkenyl optionally substituted with a halogen atom(s), wherein the alkenyl group is optionally further substituted with a group selected from R₄₂,
- 3) C2-C10 alkynyl optionally substituted with a halogen atom(s), wherein the alkynyl group is optionally further substituted with a group selected from R₄₂,
- 4) cycloalkyl optionally substituted with a group(s) independently selected from:
 - i) a halogen atom,
 - ii) C2-C6 alkenyl or C1-C6 alkyl,
 - iii) aryl optionally substituted with a group(s) independently selected from C1-C6 alkyl, a halogen atom, C1-C6 alkoxy, C1-C6 alkylamino and C1-C6 alkylcarbonyl,
 - iv) cycloalkyl,
 - v) C2-C6 haloalkenyl or C1-C6 haloalkyl,
 - vi) C1-C6 alkylidene, wherein the alkylidene is bonded to the cycloalkyl by a double bond and the alkylidene is optionally substituted with a halogen atom(s),
 - vii) C1-C6 alkoxy optionally substituted with a halogen atom(s),
 - viii) C1-C6 alkyl substituted with C1-C6 alkoxy,
 - ix) C2-C6 alkynyl and
 - x) -Si(R₄₃)₃,

5) a group represented by the following general formula (B):



(wherein Ra represents a group selected from:

- i) C1-C6 alkylcarbonyl, wherein the alkyl group is optionally substituted with R₂₇,
- ii) arylcarbonyl, wherein the aryl group is optionally substituted with a group(s) independently selected from a halogen atom, C1-C6 alkyl and C1-C6 alkoxy,

- iii) C1-C6 alkoxy carbonyl, wherein the alkyl group is optionally substituted with a group(s) selected from a halogen atom, aryl and C1-C6 alkoxy,
- iv) aryloxy carbonyl, wherein the aryl group is optionally substituted with a halogen atom(s) or C1-C6 alkyl,
- v) $-\text{CONR}_{28}\text{R}_{29}$ and
- vi) $-\text{SO}_2\text{R}_{21}$,

- 6) aryl optionally substituted with a group(s) independently selected from R_{44} ,
- 7) heteroaryl optionally substituted with any of the following groups:

- i) a halogen atom,
- ii) C1-C6 alkyl and
- iii) C1-C6 alkoxy;

- 8) C1-C6 alkoxy optionally substituted with a halogen atom(s), wherein the alkoxy group is optionally further substituted with a group selected from R_{42} ,
- 9) $-\text{S}(\text{O})_q\text{R}_{43}$ (wherein q is an integer of 0 to 2) and
- 10) cycloalkenyl optionally substituted with C1-C6 alkyl; and

R_{44} is selected from:

- 1) a halogen atom,
- 2) cyano,
- 3) C1-C6 alkyl optionally substituted with any of the following groups:


- i) a hydroxyl group,
- ii) $-\text{OR}_{26}$,
- iii) cyano and
- iv) aryloxy optionally substituted with a group(s) selected from a halogen atom, C1-C6 alkyl, C1-C6 haloalkyl or C1-C6 haloalkoxy,

- 4) C1-C6 haloalkyl,
- 5) cycloalkyl optionally substituted with a group(s) selected from a halogen atom and C1-C6 haloalkyl,
- 6) C1-C6 alkoxy optionally substituted with a halogen atom(s),
- 7) $-\text{COR}_{30}$,
- 8) C1-C6 heteroalkyl optionally substituted with a halogen atom(s),
- 9) aryl optionally substituted with a group(s) independently selected from:

- i) C1-C6 alkyl and
- ii) aryl,


- 10) heteroaryl optionally substituted with a C1-C6 alkyl group(s),
- 11) $-\text{SO}_2\text{R}_{43}$,
- 12) C1-C6 alkylthio optionally substituted with a halogen atom(s),
- 13) $-\text{Si}(\text{R}_{43})_3$ and
- 14) $-\text{SF}_5$, wherein R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{21} , R_{26} , R_{27} , R_{28} , R_{29} , R_{30} , R_{42} and R_{43} are as defined in claims 1 to 4 from which claim 5 depends, respectively.

6. The compound or a pharmacologically acceptable salt thereof according to claim 3, wherein R_1 is a group represented by the following general formula (3):




wherein R_3 , R_4 and R_5 are as defined for R_3 , R_4 and R_5 in claim 3.

7. The compound or a pharmacologically acceptable salt thereof according to claim 3, wherein R_1 is a group represented by the following general formula (4):




wherein R_4 and R_{25} are as defined for R_4 and R_{25} in claim 3.

8. The compound or a pharmacologically acceptable salt thereof according to claim 5, wherein R_1 is a group represented by the following general formula (3):



wherein R_3 , R_4 and R_5 are as defined for R_3 , R_4 and R_5 in claim 5.

9. The compound or a pharmacologically acceptable salt thereof according to claim 5, wherein R_1 is a group represented by the following general formula (4):



(4)

wherein R_4 and R_{25} are as defined for R_4 and R_{25} in claim 5.

10. A compound according to claim 1, wherein the compound is:

- (1) 8-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (2) 8-(3-chloro-benzenesulfonyl)-2-pyridin-4-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (3) 8-(3-chloro-benzenesulfonyl)-2-propyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (4) 8-(3-chloro-benzenesulfonyl)-2-isopropyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (5) 8-(3-chloro-benzenesulfonyl)-2-(3-methoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (6) 8-(3-chloro-benzenesulfonyl)-2-(3-chloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (7) 2-benzyl-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (8) 8-(3-chloro-benzenesulfonyl)-2-methyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (10) 2-biphenyl-2-yl-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one
- (11) 8-(3-chloro-benzenesulfonyl)-2-o-tolyl-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- (12) 8-(3-chloro-benzenesulfonyl)-2-(4-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (13) 8-(3-chloro-benzenesulfonyl)-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (14) 8-(3-chloro-benzenesulfonyl)-2-(1-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (15) 2-(1-acetyl-piperidin-4-yl)-8-(3-chlorobenzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (16) 2-tert-butyl-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (17) 8-(3-chloro-benzenesulfonyl)-2-(4-trifluoromethylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (18) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butyl ester;
- (19) 8-(3-chloro-benzenesulfonyl)-2-(3-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (20) 8-(3-chloro-benzenesulfonyl)-2-(2-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (21) 8-(3-chloro-benzenesulfonyl)-2-(4-propylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (22) 2-(1R,2S,4S)-bicyclo[2.2.1]hept-2-yl-8-(3-chlorobenzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (23) 2-(1R,2R,4S)-bicyclo[2.2.1]hept-2-yl-8-(3-chlorobenzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (24) 8-(3-chloro-benzenesulfonyl)-2-(4-methoxycyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (25) 8-(3-chloro-benzenesulfonyl)-2-(4-methoxycyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (26) 2-(4-tert-butyl-cyclohexyl)-8-(3-chlorobenzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (27) 8-(3-chloro-benzenesulfonyl)-2-(4-fluoromethylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (28) 8-(3-chloro-benzenesulfonyl)-2-cyclohexylmethyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (29) 8-(3-chloro-benzenesulfonyl)-2-phenethyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (30) 8-(3-chloro-benzenesulfonyl)-2-(2-cyclohexyl-ethyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (31) 8-(3-chloro-benzenesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (32) 8-(3-chloro-benzenesulfonyl)-2-(3-methanesulfonyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (33) 8-(3-chloro-benzenesulfonyl)-2-(1-phenylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (34) 8-(3-chloro-benzenesulfonyl)-2-(2-naphthalen-1-yl-ethyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (35) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (36) 2-tert-butyl-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (37) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (38) 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butyl ester;

- (39) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(2-naphthalen-1-yl-ethanesulfonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (40) 8-(3-chloro-benzenesulfonyl)-2-[1-(propane-1-sulfonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (41) 2-(1-benzenesulfonyl-piperidin-3-yl)-8-(3-chloro-benzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (42) 8-(3-chloro-benzenesulfonyl)-2-(1-phenylmethanesulfonyl-piperidin-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (43) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-(1-phenylmethanesulfonyl-piperidin-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (44) 8-(4-chloro-benzenesulfonyl)-2-(2,4-dichlorophenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (45) 2-(2,4-dichloro-phenyl)-8-(2-trifluoromethylbenzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (46) 8-(butane-1-sulfonyl)-2-(2,4-dichloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (47) 2-(2,4-dichloro-phenyl)-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (48) 2-(2,4-dichloro-phenyl)-8-(quinoline-8-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (49) 8-(3-chloro-4-fluoro-benzenesulfonyl)-2-(2,4-dichloro-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (50) 8-(3-chloro-benzenesulfonyl)-2-(2,4-dichlorophenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (51) 2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-benzoic acid methyl ester;
- (52) 2-cyclohexyl-8-(5-methyl-3-phenyl-isoxazole-4-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (53) 8-(benzo[b]thiophene-3-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (54) 8-(benzo[b]thiophene-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (55) 8-(5-chloro-thiophene-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (56) 2-cyclohexyl-8-(thiophene-2-sulfonyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one;
- (57) 2-cyclohexyl-8-(naphthalene-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (58) 2-cyclohexyl-8-(2,4-dimethyl-thiazole-5-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (59) 2-cyclohexyl-8-(3,5-dimethyl-isoxazole-4-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (60) 2-cyclohexyl-8-(2,3-dihydro-benzo[1,4]dioxine-6-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (61) 2-cyclohexyl-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (62) 3-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-thiophene-2-carboxylic acid methyl ester;
- (63) 5-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-4-methyl-thiophene-2-carboxylic acid methyl ester;
- (64) 2-cyclohexyl-8-(2,5-dimethyl-thiophene-3-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (65) 8-(5-bromo-thiophene-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (66) 8-(5-chloro-thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (67) 8-(naphthalene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (68) 8-(benzo[b]thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (69) 8-(5-bromo-thiophene-2-sulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (70) 8-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.6]undec-1-en-4-one;
- (71) 7-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,7-triaza-spiro[4.5]dec-1-en-4-one;
- (72) 7-(3-chloro-benzenesulfonyl)-2-cyclohexyl-1,3,7-triaza-spiro[4.4]non-1-en-4-one;
- (73) 4-{2-[2-(2,4-dichloro-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
- (74) 4-[2-(2-tert-butyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;
- (75) 3,N,N-trimethyl-4-{2-[4-oxo-2-(4-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (76) 3,N,N-trimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (77) 4-[2-(2-cyclopentyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethylbenzamide;
- (78) 4-{2-[2-(2,6-difluoro-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethylbenzamide;
- (79) 4-{2-[2-(2,6-dimethoxy-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethylbenzamide;
- (80) 4-{2-[2-(3-methoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethylbenzamide;
- (81) 4-{2-[2-(3-chloro-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethylbenzamide;
- (82) 4-{2-[2-(3,5-bis-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethylbenzamide;

- (83) 4-[2-(2-benzyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;
 (84) 3,N,N-trimethyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzamide;
 (85) 4-[2-[2-(2-chloro-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,N,N-trimethyl-benzamide;
 5 (86) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 (87) 4-{2-[2-(4-chloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (88) 4-{2-[2-(2,3-dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethylbenzamide;
 10 (89) 4-{2-[2-(3-chloro-4-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (90) 4-{2-[2-(2-chloro-4-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 15 (91) 4-{2-[2-(3-bromo-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (92) 4-{2-[2-(2,2-difluoro-benzo[1,3]dioxol-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (93) 4-{2-[2-(3-chloro-2-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 20 (94) 3,N,N-trimethyl-4-(2-{4-oxo-2-[4-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;
 (95) 4-{2-[2-(4-chloro-3-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 25 (96) 4-{2-[2-(3-fluoro-4-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (97) 3,N,N-trimethyl-4-(2-[2-(6-methyl-pyridin-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzamide;
 (98) 4-{2-[2-(3,4-dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 30 (99) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 (100) 4-{2-[2-(2,4-bis-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 35 (101) 3,N,N-trimethyl-4-[2-(4-oxo-2-phenethyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzamide;
 (102) 4-{2-[2-(2,4-dimethyl-thiazol-5-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (103) 4-[2-(2-cyclohexylmethyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;
 40 (104) 3,N,N-trimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 (105) 3,N,N-trimethyl-4-{2-[4-oxo-2-(2-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 (106) 3,N,N-trimethyl-4-{2-[4-oxo-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
 45 (107) 4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (108) 4-{2-[2-(2-cyclohexyl-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 50 (109) 4-{2-[2-(3-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 (110) 4-{2-[2-(3-methanesulfonyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;
 55 (111) 3,N,N-trimethyl-4-[2-[2-(2-methyl-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzamide;
 (112) 3,N,N-trimethyl-4-[2-[2-(2-methyl-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzamide;
 (113) 4-{2-[2-(2,3-dimethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-

benzamide;

(114) 4-{2-[2-(3-fluoro-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(115) 4-{2-[2-(3-fluoro-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(116) 4-{2-[2-(2-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(117) 4-{2-[2-(4-fluoro-3-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(118) 4-{2-[2-(4-difluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(119) 4-{2-[2-(2-methoxy-pyridin-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(120) 3,N,N-trimethyl-4-{2-[2-(5-methyl-pyrazin-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(121) 3,N,N-trimethyl-4-{2-[4-oxo-2-thiazol-4-yl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(122) 3,N,N-trimethyl-4-{2-[2-(1-methyl-1H-imidazol-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(123) 3,N,N-trimethyl-4-{2-[4-oxo-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(124) 4-{2-[2-(4-chloro-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(125) 4-{2-[2-(4-fluoro-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(126) 4-{2-[2-(4-chloro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(127) 4-{2-[2-(3-chloro-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(128) 4-{2-[2-(4-chloro-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(129) 4-{2-[2-(4-chloro-2-fluoro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(130) 4-{2-[2-(3-isopropoxymethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(131) 4-{2-[2-(2,4-dichloro-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(133) 4-{2-[2-(3-chloro-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(134) 4-{2-[2-(2,4-dichloro-benzyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(135) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-benzyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(136) 3,N,N-trimethyl-4-{2-[4-oxo-2-(1-trifluoromethyl-cyclopropyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(137) 3,N,N-trimethyl-4-{2-[4-oxo-2-(1-trifluoromethyl-cyclobutyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(138) 3,N,N-trimethyl-4-{2-[4-oxo-2-(1-trifluoromethyl-cyclopentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(139) 4-{2-[2-(2-fluoro-4-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(140) 3,N,N-trimethyl-4-{2-[4-oxo-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(141) 3,N,N-trimethyl-4-{2-[4-oxo-2-((E)-propenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(142) 3,N,N-trimethyl-4-{2-[4-oxo-2-((E)-styryl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(143) 4-{2-[2-(benzothiazol-6-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(144) 4-{2-[2-(4-methoxy-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-

3,N,N-trimethyl-benzamide;

(145) N-{3-methyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-acetamide;

(146) N-(4-{2-[2-(2,3-dimethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methylphenyl)-acetamide;

(147) N-(4-{2-[2-(2,3-dichloro-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methylphenyl)-acetamide;

(148) N-(3-methyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(149) N-(2-hydroxy-ethyl)-N-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(150) acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester;

(151) acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-[2-[4-oxo-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester;

(152) acetic acid (S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-[2-[4-oxo-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-amino]-ethyl ester;

(153) acetic acid (S)-1-acetoxymethyl-2-[acetyl-(4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-amino]-ethyl ester;

(154) 8-[2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl]-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(155) 8-[2-[4-((S)-2,3-dihydroxy-propylamino)-2-methylphenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(156) 8-[2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl]-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(157) 2-(3-chloro-phenyl)-8-[2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(158) 2-(4-chloro-phenyl)-8-[2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(159) 8-[2-[4-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(160) 8-[2-[4-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(161) 8-[2-[4-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl]-2-(4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(162) {4-[2-(2-tert-butyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-carbamic acid tert-butyl ester;

(163) (4-{2-[2-(4,4-difluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzyl)-carbamic acid tert-butyl ester;

(164) (4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-benzyl)-carbamic acid tert-butyl ester;

(165) (3-methyl-4-[2-[4-oxo-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzyl)-carbamic acid tert-butyl ester;

(166) 8-[2-(3-amino-phenyl)-ethanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(167) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-acetamide;

(168) 3,N,N-trimethyl-4-(2-{4-oxo-2-[3-(2,2,2-trifluoroethoxymethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;

(169) 4-(2-{2-[3-(2,2-difluoro-ethoxymethyl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide;

(170) 3,N,N-trimethyl-4-(2-{4-oxo-2-[3-(2,2,3,3-tetrafluoro-propoxymethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;

(171) 4-(2-{2-[3-(3,5-dimethyl-isoxazol-4-yl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,N,N-trimethyl-benzamide;

(172) 4-[2-(2-biphenyl-3-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamide;

(173) 3,N,N-trimethyl-4-[2-[4-oxo-2-(3-pyridin-3-yl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzamide;

(174) 3-{8-[2-(4-dimethylcarbamoyl-2-methyl-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl}-benzoic acid methyl ester;

- (175) 4-(2-[1-(2,4-dichloro-phenoxy)-1-methyl-ethyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-3,N,N-trimethyl-benzamide;
- (176) 3,N,N-trimethyl-4-{2-[4-oxo-2-(2,3,4-trifluorophenoxy)methyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-benzamide;
- (177) 2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic acid (3-ethyl-phenyl)-amide;
- (178) 2-cyclohexyl-8-[(E)-2-(1H-indol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (179) 2-cyclohexyl-8-[(E)-2-(2-trifluoromethyl-phenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (180) 2-cyclohexyl-8-[(E)-2-(3-methoxy-phenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (181) 2-cyclohexyl-8-[(E)-2-(2-methoxy-phenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (182) 2-cyclohexyl-8-[(E)-2-(1H-indol-4-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (183) 2-cyclohexyl-8-[(E)-2-(2-fluoro-6-trifluoromethylphenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (184) 2-cyclohexyl-8-[(E)-2-(2,3-difluoro-phenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (185) 2-cyclohexyl-8-[(E)-2-(3-fluoro-2-trifluoromethylphenyl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (186) 2-cyclohexyl-8-[(E)-2-o-tolyl-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (187) 2-cyclohexyl-8-[(E)-2-[4-(2-hydroxy-ethyl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (188) 2-cyclohexyl-8-[(E)-2-(2-oxo-2,3-dihydrobenzoxazol-7-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (189) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indole-1-carboxylic acid dimethylamide;
- (190) N-{2-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-phenyl}-acetamide;
- (191) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,3-dihydroxypropyl)-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (192) 2-cyclohexyl-8-[(E)-2-[1-(2-morpholin-4-yl-ethyl)-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (193) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl]-N,N-dimethyl-acetamide;
- (194) 3-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl]-1,1-dimethyl-urea;
- (195) cyclopropanecarboxylic acid {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl}-amide;
- (196) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl}-2-hydroxy-acetamide;
- (197) {3-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-4-methyl-phenyl}-carbamic acid methyl ester;
- (198) 1-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl]-3-(2-hydroxy-ethyl)-urea;
- (199) 2-cyclohexyl-8-[(E)-2-[5-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (200) 2-cyclohexyl-8-[(E)-2-[2-(2-hydroxy-ethylamino)-6-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (201) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenylamino]-N,N-dimethyl-acetamide;
- (202) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-3-fluoro-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (203) 2-cyclohexyl-8-[(E)-2-[2-methyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (204) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethylphenyl}-acetamide;
- (205) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-benzyl}-carbamic acid tert-butyl ester;
- (206) 2-cyclohexyl-8-[(E)-2-[4-((R)-2,3-dihydroxypropoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (207) 2-cyclohexyl-8-[(E)-2-[2-methyl-4-[(2-oxo-oxazolidin-5-ylmethyl)-amino]-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (208) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-N-(2-dimethylamino-ethyl)-acetamide;
- (209) 3-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzonitrile;
- (210) 8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (211) 8-[(E)-2-[4-(2-hydroxy-ethylamino)-2-trifluoromethyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (212) 8-[(E)-2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (213) 3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (214) N-(2-hydroxy-ethyl)-3,N-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (215) N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (216) 8-[(E)-2-[2-methyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethenesulfonyl]-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (217) 3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-N-(2,2,2-trifluoro-ethyl)-benzenesulfonamide;
- (218) 3-methyl-N-(2-morpholin-4-yl-ethyl)-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (219) 3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-N-pyridin-3-yl-benzenesulfonamide;
- (220) 8-[(E)-2-[4-[(R)-4-hydroxy-2-oxo-pyrrolidin-1-yl]-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (221) 8-[(E)-2-[4-[(R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl]-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (222) 8-[(E)-2-[4-[(R)-5-hydroxymethyl-2-oxo-oxazolidin-3-yl]-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (223) 8-[(E)-2-[4-[3-(3-dimethylamino-propoxy)-azetidine-1-carbonyl]-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (224) N-(4-hydroxy-butyl)-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (225) N-methyl-N-(4-methyl-3-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (226) 8-[(E)-2-(3-hydroxy-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (227) 8-[(E)-2-(5-hydroxy-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (228) N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-(1-methyl-piperidin-4-yl)-acetamide;
- (229) N-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-methanesulfonamide;
- (230) 8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (231) 3,N,N-trimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (232) N-(2-hydroxy-ethyl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (233) 3-fluoro-N,N-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (234) 8-[(E)-2-[2,5-dichloro-4-(3,4-dihydroxy-butoxy)-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (235) 8-[(E)-2-[4-(3-dimethylamino-propoxy)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (236) N-methyl-N-(2-methyl-3-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (237) N-methyl-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-

sulfonyl]-vinyl]-phenyl)-acetamide;

(238) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;

(239) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-methyl-acetamide;

(240) 2-cycloheptyl-8-((E)-2-[1-((S)-2,3-dihydroxypropyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(241) 2-(4,4-difluoro-cyclohexyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(242) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(243) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(2-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(244) 2-(2-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(245) 8-((E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(246) 8-((E)-2-[1-(2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(tetrahydropyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(247) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(2-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(248) 2-(2-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(249) 2-(2,4-dichloro-phenyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(250) 2-(4-chloro-phenyl)-8-((E)-2-[1-((S)-2,3-dihydroxypropyl)-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(251) 2-cyclohexyl-8-((E)-4-(1H-indol-4-yl)-but-3-ene-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(252) 2-cyclohexyl-8-((E)-5-(1H-indol-4-yl)-pent-4-ene-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(253) 2-cyclohexyl-8-((E)-3-[1-(2,3-dihydroxy-propyl)-1H-indol-4-yl]-prop-2-ene-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(254) 8-((E)-2-[3-(3,4-dihydroxy-butoxy)-2-methylphenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(255) 8-((E)-2-[5-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(256) 8-((E)-2-[3-(2-hydroxy-ethoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(257) N-(2-hydroxy-ethyl)-N-(4-methyl-3-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;

(258) 8-((E)-2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(259) N-(4-((E)-2-[2-(2-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(260) N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-piperidin-4-yl-acetamide;

(261) 2-cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(262) 2-cyclohexyl-8-[5-(1H-indol-4-yl)-pentane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(263) 2-cyclohexyl-8-[4-(1H-indol-4-yl)-butane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(264) N-methyl-N-(2-methyl-3-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(265) N-methyl-N-(4-methyl-3-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(266) 8-[2-[3-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(267) 8-[2-[5-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(268) 8-[2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluorome-

thyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(269) 2-cyclohexyl-8-[2-(1H-indol-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(270) 2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenylamino]-N,N-dimethyl-acetamide;

(271) cyclopropanecarboxylic acid {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl}-amide;

(272) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-benzyl}-carbamic acid tert-butyl ester;

(273) 3-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl}-1,1-dimethyl-urea;

(274) {3-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-4-methyl-phenyl}-carbamic acid methyl ester;

(275) 2-cyclohexyl-8-{2-[5-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(276) N-{2-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamide;

(277) 2-cyclohexyl-8-(2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(278) 1-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl}-3-(2-hydroxy-ethyl)-urea;

(279) 2-cyclohexyl-8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(280) 2-cyclohexyl-8-{2-[4-(2-hydroxy-ethyl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(281) 2-cyclohexyl-8-(2-{2-methyl-4-[(2-oxo-oxazolidin-5-ylmethyl)-amino]-phenyl}-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(282) 2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-N-(2-dimethyl-amino-ethyl)-acetamide;

(283) 2-cyclohexyl-8-{2-[2-methyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(284) 8-{2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(285) 8-{2-[4-(4-hydroxy-piperidin-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(286) 8-{2-[4-(2-hydroxy-ethylamino)-2-trifluoromethylphenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(287) N-(4-hydroxy-butyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;

(288) 8-{2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(289) 8-{2-[2-methyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(290) 3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-(2,2,2-trifluoro-ethyl)-benzenesulfonamide;

(291) 3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;

(292) 3-methyl-N-(2-morpholin-4-yl-ethyl)-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;

(293) 3-fluoro-N,N-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(294) 3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-pyridin-3-yl-benzenesulfonamide;

(295) 8-{2-[4-(3-dimethylamino-propoxy)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(296) 8-{2-[4-((R)-5-hydroxymethyl-2-oxo-oxazolidin-3-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(297) 8-{2-[4-((R)-4-hydroxy-2-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(298) N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

- (299) N-methyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (300) 2-hydroxy-N-methyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- 5 (301) 1-methyl-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-urea;
- (302) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethylphenyl}-acetamide;
- (303) 2-cyclohexyl-8-[2-(1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 10 (304) 2-(3-trifluoromethoxy-phenyl)-8-[2-(2-trifluoromethyl-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (305) 8-[2-(1H-indol-4-yl)-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (306) 8-[2-(1H-indol-4-yl)-ethanesulfonyl]-2-(4-methylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (307) 8-[2-(2,2-difluoro-benzo[1,3]dioxol-4-yl)-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 15 (308) 8-[2-(1-methyl-1H-indol-7-yl)-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (309) 2-cyclohexyl-8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 20 (310) 4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-indole-1-carboxylic acid dimethylamide;
- (311) 8-[3-(1H-indol-4-yl)-propane-1-sulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (312) 2-cyclohexyl-8-[2-(1-methanesulfonyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 25 (313) 2-cyclohexyl-8-[2-[1-(2-dimethylamino-ethyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (314) 8-[2-(4-amino-2-trifluoromethyl-phenyl)-ethanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (315) 2-cyclohexyl-8-[3-(3-trifluoromethyl-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 30 (316) 2-cyclohexyl-8-(3-m-tolyl-propane-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (317) 2-cyclohexyl-8-[3-(3-hydroxy-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (318) 2-cyclohexyl-8-[3-(2-hydroxy-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (319) 8-(2-benzo[b]thiophen-3-yl-ethanesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (320) 2-cyclohexyl-8-(2-isoquinolin-5-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 35 (321) 2-cyclohexyl-8-[2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (322) 8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl]-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (323) 2-cyclohexyl-8-[2-(2-trifluoromethyl-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 40 (324) 4-[3-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-propyl]-N,N-dimethyl-benzamide;
- (325) 8-[2-[4-((R)-2,3-dihydroxy-propoxy)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (326) N-(2-hydroxy-ethyl)-N-(4-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- 45 (327) N-(2-hydroxy-ethyl)-N-(2-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (328) 8-[3-(3-amino-phenyl)-propane-1-sulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (329) N,N-dimethyl-4-{5-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-benzamide;
- 50 (330) N,N-dimethyl-3-{5-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-benzamide;
- (331) 3,N,N-trimethyl-4-{5-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-benzamide;
- (332) N-{4-[5-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-thiophen-2-yl]-3-methyl-phenyl}-acetamide;
- 55 (333) 4-{5-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-3,N,N-trimethyl-benzamide;
- (334) 3-{5-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-

yl)-4,N,N-trimethyl-benzamide;

(335) 2-methyl-3'-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-biphenyl-4-carboxylic acid dimethylamide;

(336) 2-methyl-4'-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-biphenyl-4-carboxylic acid dimethylamide;

(337) 3,N,N-trimethyl-4-{6-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-pyridin-2-yl}-benzamide;

(338) 2-cyclohexyl-8-{2-[2-(3,5-dimethyl-isoxazol-4-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(339) 2-cyclohexyl-8-{2-[2-(6-methoxy-pyridin-3-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(340) 2-cyclohexyl-8-{2-[2-(1-methyl-1H-pyrazol-4-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(341) 8-{5-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-thiophene-2-sulfonyl}-2-(4-fluoro-3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(342) N-(4-{5-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-thiophen-2-yl}-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(343) 2-cyclohexyl-8-((E)-2-thiazol-2-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(344) 2-cyclohexyl-8-((E)-2-cyclopentyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(345) 2-cyclohexyl-8-(2-thiazol-2-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(346) 2-cyclohexyl-8-[3-(4-methoxy-phenyl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(347) N-benzyl-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-benzamide;

(348) 8-(3-chloro-benzenesulfonyl)-2-[3-(morpholine-4-carbonyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(349) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-N-methyl-N-phenyl-benzamide;

(350) N-benzyl-3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-benzamide;

(351) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-N,N-dimethyl-benzamide;

(352) 8-[2-[4-(4-methanesulfonyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(353) N-[(R)-1-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzoyl)-pyrrolidin-3-yl]-acetamide;

(354) 8-[2-[2-methyl-4-((S)-2-trifluoromethyl-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(355) 8-[2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(356) 8-[2-[4-((S)-2-hydroxymethyl-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(357) 8-(2-[4-[4-(2-hydroxy-ethyl)-piperidine-1-carbonyl]-2-methyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(358) 8-(2-[2-methyl-4-[4-(3-methyl-[1,2,4]oxadiazol-5-yl)-piperidine-1-carbonyl]-phenyl]-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(359) 8-[2-[2-methyl-4-(4-pyrimidin-2-yl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(360) 4-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzoyl)-piperazine-1-sulfonic acid dimethylamide;

(361) 8-[2-[2-methyl-4-(4-pyridin-2-yl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(362) 8-[2-[4-[4-[2-(2-hydroxy-ethoxy)-ethyl]-piperazine-1-carbonyl]-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(363) 8-[2-[2-methyl-4-[4-(2-morpholin-4-yl-ethyl)-piperazine-1-carbonyl]-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(364) 8-[2-[2-methyl-4-(4-thiazol-2-yl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(365) 8-[2-[4-(4,4-difluoro-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(366) 8-(2-[4-[4-(3-hydroxy-propyl)-piperazine-1-carbonyl]-2-methyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (367) (S)-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoyl)-pyrrolidine-2-carboxylic acid amide;
- (368) 8-{2-[2-methyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (369) (R)-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoyl)-pyrrolidine-2-carboxylic acid amide;
- (370) 8-{2-[4-((S)-3-hydroxy-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (371) 8-{2-[4-((R)-2-hydroxymethyl-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (372) 8-{2-[4-((S)-3-dimethylamino-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (373) 8-{2-[4-(4-tert-butyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (374) 8-(2-{4-[4-(3-dimethylamino-propyl)-piperazine-1-carbonyl]-2-methyl-phenyl}-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (375) 8-(2-{4-[4-(4-fluoro-phenyl)-piperazine-1-carbonyl]-2-methyl-phenyl}-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (376) 8-{2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (377) 8-{2-[4-(3-hydroxy-azetidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (378) 8-{2-[4-(3-fluoro-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (379) 8-{2-[4-(3-fluoro-azetidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (380) 8-{2-[2-methyl-4-(piperidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (381) 8-{2-[4-(azetidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (382) 4-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzoyl)-piperazine-1-carboxylic acid dimethylamide;
- (383) 3,5,N,N-tetramethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (384) 8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (385) 8-[(E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (386) 8-[(E)-2-[4-(4-hydroxy-4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (387) 8-[(E)-2-[4-((3R,4R)-3-dimethylamino-4-hydroxy-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (388) 8-[(E)-2-[2,6-dimethyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (389) N-(3-hydroxy-propyl)-3,5,N-trimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (390) N-(2-dimethylamino-ethyl)-3,5,N-trimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (391) N-(3-dimethylamino-propyl)-3,5,N-trimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (392) N-carbamoylmethyl-3,5,N-trimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (393) 3,5,N-trimethyl-N-(1-methyl-piperidin-4-yl)-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;
- (394) 8-[(E)-2-[4-(4-acetyl-piperazine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (395) 4-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzoyl)-piperazine-1-carboxylic acid dimethylamide;

- (396) 4-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoyl)-piperazine-1-carboxylic acid amide;
- (397) 8-((E)-2-[4-[4-(3-dimethylamino-propyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (398) 8-((E)-2-[2,6-dimethyl-4-((R)-3-methylamino-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (399) 8-((E)-2-[4-((R)-3-dimethylamino-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (400) 8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (401) 8-((E)-2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (402) 8-((E)-2-[4-((S)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (403) 8-((E)-2-[4-(4-hydroxy-4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (404) 8-((E)-2-[4-((R)-3-amino-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (405) 8-((E)-2-[4-(4-dimethylamino-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (406) 8-((E)-2-[4-((3S,4S)-3-hydroxy-4-isopropylamino-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (407) 8-((E)-2-[4-[4-(2-dimethylamino-ethoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (408) 8-((E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (409) 2,N,N-trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (410) 2-methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (411) 8-{2-[2-methyl-3-(morpholine-4-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (412) 4,N,N-trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (413) N-(2-hydroxy-ethyl)-4,N-dimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (414) 8-{2-[2-methyl-5-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (415) 4,N-dimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (416) 8-{2-[4-[4-(2-hydroxy-ethyl)-piperazine-1-carbonyl]-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (417) 8-{2-[2-methyl-4-(thiazolidine-3-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (418) 2-fluoro-5,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (419) 8-{2-[5-fluoro-4-(4-hydroxy-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (420) N-benzyl-2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-1H-indol-3-yl]-acetamide;
- (421) 2-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-N-methyl-benzamide;
- (422) 3-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-N-pyridin-4-yl-benzamide;
- (423) 8-(3-chloro-benzenesulfonyl)-2-[1-(3,3-dimethyl-butyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (424) 8-(3-chloro-benzenesulfonyl)-2-[1-(2-hydroxy-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (425) 8-(3-chloro-benzenesulfonyl)-2-[1-(4-chlorobenzoyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (426) 8-(3-chloro-benzenesulfonyl)-2-[1-(3-methoxy-benzoyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (428) 2-[1-(1H-indole-5-carbonyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (429) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(3-phenyl-propionyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (430) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(1,2,3,4-tetrahydro-naphthalene-2-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (431) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(quinoline-6-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (432) 2-[1-(2-3H-imidazol-4-yl-acetyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (433) 2-[1-[2-(2,5-dimethyl-thiazol-4-yl)-acetyl]-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (434) 2-[1-[2-(5-methyl-2-phenyl-oxazol-4-yl)-acetyl]-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (435) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(2-pyridin-2-yl-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (436) 2-[1-(4-benzyloxy-butyryl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (437) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(4-phenyl-butyryl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (438) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-(2-pyridin-4-yl-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (439) 2-[1-(2-tert-butoxy-acetyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (441) 8-(2-naphthalen-1-yl-ethanesulfonyl)-2-[1-[(E)-(3-phenyl-acryloyl)]-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (442) 2-[1-(2-amino-acetyl)-piperidin-3-yl]-8-(3-chlorobenzenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (444) 8-[2-[2-methyl-4-(3-methylamino-pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (445) 8-[2-[4-(3-amino-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (446) 8-[(E)-2-[2,6-dimethyl-4-(piperazine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (447) 8-[(E)-2-[4-[4-(2-hydroxy-acetyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (448) 4-[4-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzoyl)-piperazin-1-yl]-4-oxo-butyric acid methyl ester;
- (449) 8-[(E)-2-[4-[4-(4-dimethylamino-butyryl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (450) 2-methoxy-N-methyl-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (451) 2-hydroxy-N-methyl-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (452) [(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzoyl)-methyl-amino]-acetic acid;
- (453) 8-(3-chloro-benzenesulfonyl)-2-[1-(3,3-dimethylbutyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (454) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid tert-butylamide;
- (455) 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid phenethyl-amide;
- (456) 8-(3-chloro-benzenesulfonyl)-2-[1-(piperidine-1-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (457) 3-(2-dimethylamino-ethyl)-1-methyl-1-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-urea;
- (458) 1-methyl-1-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-

sulfonyl]-vinyl)-phenyl)-urea;

(459) 1-{3-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-2-methyl-phenyl}-3-methyl-urea;

(460) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid 2-methoxy-ethyl ester;

(461) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid benzyl ester;

(462) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid isobutyl ester;

(463) 3-[8-(3-chloro-benzenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid methyl ester;

(464) 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid benzyl ester;

(465) 3-[8-(2-naphthalen-1-yl-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidine-1-carboxylic acid 2,2-dimethyl-propyl ester;

(466) methyl-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-carbamic acid 2-dimethylamino-ethyl ester;

(467) methyl-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-carbamic acid methyl ester;

(468) methyl-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-carbamic acid methyl ester;

(469) 2-cyclohexyl-8-(2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-4-thione;

(470) 1-(3,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(471) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;

(472) 1-(3,5-dimethyl-4-[2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(473) 1-(3,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-thiourea;

(475) 8-[2-[2,6-dimethyl-4-(methyl-thiazol-2-yl-amino)-phenyl]-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(476) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-acetic acid methyl ester;

(477) 2-cyclohexyl-8-[(E)-2-[1-(morpholine-4-carbonyl)-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(478) 2-cyclohexyl-8-[(E)-2-[1-(4-hydroxy-butyl)-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(479) 2-cyclohexyl-8-[(E)-2-(1H-indol-7-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(480) 2-cyclohexyl-8-[(E)-2-[1-(3,4-dihydroxy-butyl)-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(481) N-(2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-ethyl)-acetamide;

(482) N-(1-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-ethyl)-acetamide;

(483) 2-cyclohexyl-8-[(E)-2-(1-thiazol-2-ylmethyl-1H-indol-4-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(484) 3-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-propane-1-sulfonic acid amide;

(485) 4-methyl-piperazine-1-carboxylic acid {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenyl}-amide;

(486) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-benzyl}-carbamic acid isobutyl ester;

(487) 2-cyclohexyl-8-[(E)-2-[1-[2-(2-hydroxy-ethoxy)-ethyl]-1H-indol-5-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(488) 2-cyclohexyl-8-[(E)-2-(6-methyl-1H-indol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(489) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-benzyl}-2-hydroxy-acetamide;

- (490) 3-(2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-ethyl)-1,1-dimethyl-urea;
- (491) 8-[(E)-2-(4-amino-2,6-bis-trifluoromethyl-phenyl)-ethenesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (492) 2-cyclohexyl-8-[(E)-2-(6-trifluoromethyl-1H-indol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (493) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenylamino}-acetic acid methyl ester;
- (494) 2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenylamino}-N-(2-hydroxy-ethyl)-acetamide;
- (495) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-N-(2-hydroxyethyl)-N-methyl-3-trifluoromethyl-benzenesulfonamide;
- (496) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-N,N-bis-(2-hydroxyethyl)-3-trifluoromethyl-benzenesulfonamide;
- (497) 2-cyclohexyl-8-[(E)-2-[4-(2-dimethylamino-ethylamino)-2-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (498) N-(2-acetyl-amino-ethyl)-2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethyl-phenylamino}-acetamide;
- (499) 2-cyclohexyl-8-[(E)-2-(6-trifluoromethyl-1H-benzimidazol-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (500) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-5-fluoro-1H-indol-4-yl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (501) 2-cyclohexyl-8-[(E)-2-[4-[(2-hydroxy-ethyl)-methyl-amino]-2-trifluoromethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (502) 2-cyclohexyl-8-[(E)-2-[4-(1,1-dioxo-1 λ ⁶-thiomorpholine-4-carbonyl)-2-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (503) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-N-(2-dimethylamino-ethyl)-3-methyl-benzenesulfonamide;
- (504) 3-(3-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (505) 1-methyl-3-(3-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (506) N-(3-isopropoxy-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (507) 3-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (508) 3-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1,5,5-trimethyl-imidazolidine-2,4-dione;
- (509) N-(3-ethoxy-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (510) N-(3-ethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (511) 5,5-dimethyl-3-(3-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (512) N-(3-methoxymethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (513) 2-(4-methyl-cyclohexyl)-8-[(E)-2-[2-methyl-4-(2-oxo-oxazolidin-3-yl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (514) N-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-(2-methoxy-ethyl)-acetamide;
- (515) 8-[(E)-2-[2,6-dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (516) 8-[(E)-2-[2,6-dimethyl-4-(2-oxo-piperidin-1-yl)-phenyl]-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (517) N-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzyl)-N-methyl-acetamide;
- (518) 3-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzyl)-imidazolidine-2,4-dione;

- (519) 3-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (520) 8-((E)-2-[2,6-dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethenesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (521) 1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl)-pyrrolidine-2,5-dione;
- (522) N-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl)-acetamide;
- (523) N-(3,5-difluoro-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide;
- (524) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (525) [(S)-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-2,5-dioxo-pyrrolidin-3-yl]-carbamic acid tert-butyl ester;
- (526) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (527) 8-((E)-2-[2-methyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (528) 8-((E)-2-[2-methyl-4-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethylamino]-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (529) 2-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenylamino)-acetamide;
- (530) 3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-N-pyridin-4-yl-benzenesulfonamide;
- (531) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (532) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-2,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3-bis-(2-hydroxy-ethyl)-urea;
- (533) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3-bis-(2-hydroxy-ethyl)-3-methyl-urea;
- (534) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-(tetrahydro-pyran-4-yl)-methanesulfonamide;
- (535) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (536) 3-[3-methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-imidazolidine-2,4-dione;
- (537) 3-[4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-trifluoromethylphenyl]-imidazolidine-2,4-dione;
- (538) 4-[3-methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-morpholine-3,5-dione;
- (539) 3-[3,5-dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-imidazolidine-2,4-dione;
- (540) 5,5-dimethyl-3-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-imidazolidine-2,4-dione;
- (541) 8-((E)-2-[1-((R)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (542) N-(2-methoxy-ethyl)-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-acetamide;
- (543) N-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-N-(2-methoxy-ethyl)-acetamide;
- (544) N-[2-(2-hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-acetamide;
- (545) N-[2-(2-fluoro-ethoxy)-ethyl]-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-acetamide;
- (546) N-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-N-methyl-acetamide;
- (547) 3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid;

- (548) 3-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-imidazolidine-2,4-dione;
- (549) 8-((E)-2-{4-[4-(2-fluoro-ethyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl}-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (550) 8-((E)-2-[2,6-dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (551) N-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-N-[2-(2-fluoro-ethoxy)-ethyl]-acetamide;
- (552) 3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzoic acid N'-acetyl-hydrazide;
- (553) 8-[(E)-2-(4-hydroxymethyl-2,6-dimethyl-phenyl)-ethenesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (554) 3-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-imidazolidine-2,4-dione;
- (555) 3-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-5,5-dimethyl-imidazolidine-2,4-dione
- (556) 8-((E)-2-[2,6-dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (557) 1-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-pyrrolidine-2,5-dione;
- (558) N-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-acetamide;
- (559) N-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-benzyl]-N-methyl-acetamide;
- (560) 3-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidine-2,4-dione;
- (561) N-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-[2-(2-methoxy-ethoxy)-ethyl]-acetamide;
- (562) N-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-[2-(2-fluoro-ethoxy)-ethyl]-acetamide;
- (563) 3-(4-((E)-2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidine-2,4-dione;
- (564) 3-(4-((E)-2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;
- (565) 3-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-benzyl]-1,1-dimethyl-urea;
- (566) (2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-ethyl)-carbamic acid methyl ester;
- (567) 4-methyl-piperazine-1-carboxylic acid 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-benzylamide;
- (568) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-phenyl}-N',N'-dimethyl-sulfamide;
- (569) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-benzyl}-(3-dimethylamino-propyl)-carbamic acid tert-butyl ester;
- (570) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-N-(2,2-dimethyl-propyl)-acetamide;
- (571) 2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl}-N-(2-diisopropylamino-ethyl)-acetamide;
- (572) 2-cyclohexyl-8-((E)-2-[2-methyl-4-(2-oxo-azetidin-1-yl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (573) 2-cyclohexyl-8-((E)-2-[2,6-dimethyl-4-(thiazol-2-ylamino)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (574) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethylphenyl}-N-(3-methyl-oxetan-3-ylmethyl)-acetamide;
- (575) 3-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione;
- (576) N-(3-ethoxymethyl-4-((E)-2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;

- (577) 2-(4-methyl-cyclohexyl)-8-[(E)-2-[2-methyl-4-(4-oxo-oxazolidin-3-yl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (578) 4-(3-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-3-oxo-piperazine-1-carboxylic acid tert-butyl ester;
- (579) 2-(4-methyl-cyclohexyl)-8-[(E)-2-[2-methyl-4-(4-methyl-2-oxo-oxazolidin-3-yl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (580) 2-(4-methyl-cyclohexyl)-8-[(E)-2-[2-methyl-4-(3-oxo-isoxazolidin-2-yl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (581) 3-(2,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (582) 3-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-5-methyl-imidazolidine-2,4-dione;
- (583) 3-(2,6-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (584) N-(2,6-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-methyl-acetamide;
- (585) 8-[(E)-2-(3-methoxy-2-methyl-phenyl)-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (586) 8-[(E)-2-[4-[(1H-imidazol-2-yl)-methyl-amino]-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (587) 1-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-dihydro-pyrimidine-2,4-dione;
- (588) 2-(4-methyl-cyclohexyl)-8-[(E)-2-[2-methyl-4-(5-oxo-pyrazolidin-1-yl)-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (589) 5-tert-butoxymethyl-3-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-imidazolidine-2,4-dione;
- (590) N,N-dimethyl-2-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonylamino)-acetamide;
- (591) 3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-N-pyridin-3-ylmethyl-benzenesulfonamide;
- (592) N-(4-hydroxy-cyclohexyl)-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (593) 8-[(E)-2-(4-methanesulfonyl-2-methyl-phenyl)-ethenesulfonyl]-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (594) N-acetyl-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (595) N-(2-hydroxy-1,1-bis-hydroxymethyl-ethyl)-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (596) N-(1-benzyl-piperidin-4-yl)-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (597) 8-[(E)-2-[1-[(S)-2,3-dihydroxy-propyl]-1H-indol-5-yl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (598) 8-[(E)-2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (599) N-[(R)-2,3-dihydroxy-propyl]-3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzenesulfonamide;
- (600) 8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-3,5-difluoro-phenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (601) 8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-3,5-dimethylphenyl]-ethenesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (602) 4-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester;
- (603) N-isopropyl-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (604) 4-[acetyl-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester;
- (605) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-(2-hydroxy-ethyl)-urea;

- (606) 8-((E)-2-[4-((R)-2-(isopropylamino-methyl)-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (607) N-(2-dimethylamino-ethyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (608) 2-dimethylamino-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide;
- (609) N-(2-dimethylamino-ethyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methanesulfonamide;
- (610) N-((R)-1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-5-oxo-pyrrolidin-2-ylmethyl)-acetamide;
- (611) 8-((E)-2-[4-((R)-2-dimethylaminomethyl-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (612) N-cyanomethyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (613) N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-(2,2,2-trifluoro-ethyl)-acetamide;
- (614) 3-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,5,5-trimethyl-imidazolidine-2,4-dione;
- (615) 3-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (616) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-(tetrahydro-pyran-4-yl)-acetamide;
- (617) 8-((E)-2-[2-methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (618) 3-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione;
- (619) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (620) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (621) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-((E)-2-[4-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (622) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (623) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl)-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (624) 1-(4-((E)-2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidine-2,4-dione;
- (625) 1-(4-((E)-2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-5-methyl-imidazolidine-2,4-dione;
- (626) 1-(4-((E)-2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3-methyl-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (627) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (628) 3,5-dimethyl-4-((E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-benzoic acid;
- (629) 3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid trimethylhydrazide;
- (630) 8-((E)-2-[1-((R)-2,3-dihydroxy-propyl)-4,6-dimethyl-1H-indol-5-yl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (631) 3-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;
- (632) N-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-(3-methyl-oxetan-3-ylmethyl)-acetamide;
- (633) 2-cyclohexyl-8-((E)-2-quinolin-8-yl-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (634) 2-cyclohexyl-8-((E)-2-(2-oxo-2,3-dihydro-1H-indol-4-yl)-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (635) 3-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-imidazolidine-2,4-dione;

- (636) 3-(4-((E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;
- (637) 2-cyclohexyl-8-[(E)-2-(2-methyl-1H-indol-4-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (638) 2-cyclohexyl-8-[(E)-2-(1-methyl-1,2,3,4-tetrahydro-quinolin-5-yl)-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (639) 8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (640) 4-[(E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl-[1,1,1-²H₃]methyl-carbamic acid tert-butyl ester;
- (641) 8-[(E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (642) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (643) 8-[(E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethenesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (644) 8-[(E)-2-[2,6-dimethyl-4-(2-oxa-7-aza-spiro[3.5]nonane-7-carbonyl)-phenyl]-ethenesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (645) 2-cyclohexyl-8-[2-(1,2,3,4-tetrahydro-quinolin-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (646) 4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-1H-indole-3-carbonitrile;
- (647) 7-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-indole-1-carboxylic acid dimethylamide;
- (648) 2-cyclohexyl-8-(2-quinolin-5-yl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (649) 2-cyclohexyl-8-[2-[3-(2,2,2-trifluoro-acetyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (650) 2-cyclohexyl-8-[2-(1-isopropyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (651) 2-cyclohexyl-8-[2-[1-(4-hydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (652) N-[3-cyano-4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-acetamide;
- (653) 8-(2-isoquinolin-5-yl-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (654) 8-(2-quinolin-5-yl-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (655) N-(3-methoxy-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;
- (656) 8-[2-[4-(3,3-dimethyl-1,1,4-trioxo-1 λ ⁶-[1,2,5]thiadiazolidin-2-yl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (657) 8-[2-[2-methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (658) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzyl}-carbamic acid isobutyl ester;
- (659) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzyl}-2-hydroxy-acetamide;
- (660) 2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenylamino]-N-(4-hydroxy-butyl)-acetamide;
- (661) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl}-carbamic acid isobutyl ester;
- (662) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl}-benzamide;
- (663) 2-cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzothiazol-6-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (664) N-(2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzenesulfonylamino]-ethyl)-acetamide;
- (665) N-(2-dimethylamino-ethyl)-2-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenylamino)-acetamide;
- (666) 8-[2-[4-((2R,6S)-2,6-dimethyl-morpholine-4-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (667) N-(2,2,3,3,4,4,4-heptafluoro-butyl)-3-methyl-4-[2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-

spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzenesulfonamide;

(668) 8-[2-(3-methoxy-2-methyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(669) (2-methoxy-3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-carbamic acid tert-butyl ester;

(670) 8-[2-[4-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(671) 8-[2-(5,7-dimethyl-2-oxo-2,3-dihydro-benzoxazol-6-yl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(672) (2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-ethyl)-carbamic acid methyl ester;

(673) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-methyl-carbamic acid tert-butyl ester;

(674) 8-(2-[4-[3-(2-hydroxy-ethyl)-2-oxo-imidazolidin-1-yl]-2-methyl-phenyl]-ethanesulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(675) 2-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-benzenesulfonylamino)-acetamide;

(676) 8-{2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethylphenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(677) 3-(4-[2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethylphenyl)-imidazolidine-2,4-dione;

(678) N-cyclopentyl-N-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(679) 1-(2,3-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(680) 1-(2,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(681) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester;

(682) 1-(3-methoxy-5-methyl-4-[2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(683) 1-(3-chloro-5-methyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(684) 1-(2-methoxy-3,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(685) 5,7-dimethyl-6-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-1H-quinazoline-2,4-dione;

(686) 8-[(E)-2-[1-(2-amino-ethyl)-1H-indol-4-yl]-ethenesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(687) 8-[(E)-2-(4-aminomethyl-phenyl)-ethenesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(688) (S)-2-amino-3-hydroxy-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-propionamide;

(689) N-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-N-piperidin-4-yl-methanesulfonamide;

(690) 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(691) 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethenesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(692) 1-[3,5-dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-1-methyl-urea;

(693) 1-(2-methoxy-3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;

(694) 1-[3,5-dimethyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl]-1-(2-fluoro-ethyl)-urea;

(695) 1-(3-chloro-5-methyl-4-[(E)-2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;

(696) 1-[3,5-dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

- (697) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (698) 8-((E)-2-[4-[2-((R)-2,3-dihydroxy-propoxy)-ethoxy]-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (699) N-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide;
- (700) N-[2-(2-hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-((E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-phenyl]-acetamide
- (701) N-[3,5-dimethyl-4-((E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-phenyl]-N-(2-hydroxy-ethyl)-acetamide;
- (702) 2-cyclohexyl-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (704) 8-((E)-2-[1-[2-((S)-2,3-dihydroxy-propoxy)-ethyl]-1H-indol-4-yl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (705) 8-((E)-2-[2,6-dimethyl-4-((2S,3S)-2,3,4-trihydroxy-butoxy)-phenyl]-ethenesulfonyl)-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (706) 2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-8-((E)-2-[1-[2-((2S,3S)-2,3,4-trihydroxy-butoxy)-ethyl]-1H-indol-4-yl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (707) N-((S)-2,3-dihydroxy-propyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (708) {4-((E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbamic acid tert-butyl ester;
- (709) N-[4-((E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl)-3,5-dimethyl-phenyl]-N-(4-hydroxy-cyclohexyl)-acetamide;
- (710) 2-cyclohexyl-8-((E)-2-[4-((R)-2-(2-hydroxy-ethoxymethyl)-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (711) N-(2-hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-isobutylamide;
- (712) 8-((E)-2-[4-((R)-5-hydroxymethyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (713) 8-((E)-2-[4-((R)-5-hydroxymethyl-3-methyl-2-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (714) 3-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-(2-hydroxy-ethyl)-5,5-dimethyl-imidazolidine-2,4-dione;
- (715) N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-(4-hydroxy-cyclohexyl)-methanesulfonamide;
- (716) N-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide;
- (717) N-(2-fluoro-5-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-(2-hydroxy-ethyl)-acetamide;
- (718) N-(4-((E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;
- (719) N-(2-hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methanesulfonamide;
- (720) N-(2-hydroxy-ethyl)-N-(3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl)-methanesulfonamide;
- (721) N-(4-[2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;
- (722) 2-hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (723) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
- (724) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-[2-(2-hydroxy-ethoxy)-ethyl]-urea;
- (725) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-(2-methoxy-ethyl)-urea;
- (726) 1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-isopropyl-urea;

- (727) 1-(3-ethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- (728) 1-(3-methoxy-5-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- (729) 1-cyanomethyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea;
- (730) 1-cyclopentyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea;
- (731) 1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-(2,2,2-trifluoro-ethyl)-urea;
- (732) 1-(4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethylphenyl)-1-methyl-urea;
- (733) 1-((S)-2,3-dihydroxy-propyl)-1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-urea;
- (734) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3-methyl-phenyl}-N-cyclopentyl-acetamide;
- (735) (S)-2-amino-N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-3-methyl-butylamide;
- (736) 2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-N-pyridin-4-yl-acetamide;
- (737) 2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (738) 2-cyclohexyl-8-[(E)-2-[4-(4,5-dihydro-thiazol-2-ylamino)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (739) 2-cyclohexyl-8-[(E)-2-[2-methyl-4-(3-methyl-oxetan-3-ylmethoxy)-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (740) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropyl}-phenyl)-1-methyl-urea;
- (741) 1-(3,5-dimethyl-4-((E)-1-methyl-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- (742) N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-sulfamide;
- (743) N-(3-hydroxy-propyl)-N'-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-sulfamide;
- (744) N-methyl-N-(3-methyl-4-((E)-3-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-propenyl)-phenyl)-acetamide;
- (745) 2-cyclohexyl-8-[2-(2-methyl-1H-indol-4-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (746) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (747) 2-cyclohexyl-8-[2-(1-methyl-1,2,3,4-tetrahydro-quinolin-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (748) 2-cyclohexyl-8-[2-[1-(3,4-dihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (749) 3-(2,6-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;
- (750) 2-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (751) N-(3-ethoxy-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (752) N-(3-ethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (753) N-(3-ethoxymethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (754) N-(3-methoxymethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (755) N-(2-methoxy-ethyl)-N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-acetamide;
- (756) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-

sulfonyl)-ethyl)-phenyl]-N-(2-methoxy-ethyl)-acetamide;

(757) N-[2-(2-fluoro-ethoxy)-ethyl]-N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-acetamide;

(758) N-[2-(2-hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-acetamide;

(759) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-N-methyl-acetamide;

(760) 3-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-1-methyl-imidazolidine-2,4-dione;

(761) 8-(2-{4-[4-(2-fluoro-ethyl)-piperazine-1-carbonyl]-2,6-dimethyl-phenyl}-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(764) 8-{2-[2,6-dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(765) 8-{2-[2,6-dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(766) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-phenyl]-N-[2-(2-fluoro-ethoxy)-ethyl]-acetamide;

(767) 3-{3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl}-5-methyl-imidazolidine-2,4-dione;

(768) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-benzoic acid;

(769) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-benzoic acid N'-acetyl-hydrazide;

(770) 8-[2-(4-hydroxymethyl-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(771) 8-{2-[2,6-dimethyl-4-(2-oxo-piperidin-1-yl)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(772) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-benzyl]-acetamide;

(773) N-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl)-acetamide;

(774) 3-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-benzyl]-imidazolidine-2,4-dione;

(775) 3-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-benzyl]-5,5-dimethyl-imidazolidine-2,4-dione;

(776) 8-{2-[2,6-dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(777) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-benzyl]-pyrrolidine-2,5-dione;

(778) 3-(2,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-imidazolidine-2,4-dione

(779) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl)-imidazolidine-2,4-dione;

(780) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl)-5,5-dimethyl-imidazolidine-2,4-dione;

(781) 8-{2-[2,6-dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(782) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl)-pyrrolidine-2,5-dione;

(783) N-(2,6-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-N-methyl-acetamide;

(784) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl)-benzyl]-N-methyl-acetamide;

(785) N-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl)-N-methyl-acetamide;

(786) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-dihydro-pyrimidine-2,4-dione;

(787) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sul-

fonyl)-ethyl)-benzoic acid trimethylhydrazide;

(788) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(789) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(790) 3-{3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-imidazolidine-2,4-dione;

(791) 8-{2-[1-((R)-2,3-dihydroxy-propyl)-4,6-dimethyl-1H-indol-5-yl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(792) 3-(3-methyl-4-[2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-imidazolidine-2,4-dione;

(793) 8-{2-[2-methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(794) N-(4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(795) 3-{4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-trifluoromethyl-phenyl}-imidazolidine-2,4-dione;

(796) 3,5-dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzoic acid;

(797) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(798) 8-{2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethylphenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(799) 3-(3,5-dimethyl-4-[2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-imidazolidine-2,4-dione;

(800) 3-(4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;

(801) 8-{2-[4-(3,4-dihydroxy-butoxy)-2,6-dimethylphenyl]-ethanesulfonyl}-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(802) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(803) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(804) 8-{2-[2,6-dimethyl-4-(2-oxa-7-aza-spiro[3.5]nonane-7-carbonyl)-phenyl]-ethanesulfonyl}-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(812) N-(3-isopropoxy-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(817) 1-{3,5-dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;

(818) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-(2-fluoro-ethyl)-urea;

(820) N-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide;

(821) 8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethylphenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(822) 8-(2-{1-[2-((S)-2,3-dihydroxy-propoxy)-ethyl]-1H-indol-4-yl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(823) 2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-8-(2-{1-[2-((2S,3S)-2,3,4-trihydroxy-butoxy)-ethyl]-1H-indol-4-yl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(824) N-((S)-2,3-dihydroxy-propyl)-N-(3,5-dimethyl-4-[2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-acetamide;

(825) N-[2-(2-hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-acetamide;

(826) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-[2-(2-hydroxy-ethoxy)-ethyl]-urea;

(827) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-(2-methoxy-ethyl)-urea;

(828) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-isopropyl-urea;

- (829) 1-(3-ethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (830) 1-((S)-2,3-dihydroxy-propyl)-1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-urea;
- (831) 8-{2-[4-(4-hydroxy-4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (832) 3-{3,5-dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-imidazolidine-2,4-dione;
- (833) 5-tert-butoxymethyl-3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;
- (834) 2-(4-methyl-cyclohexyl)-8-{2-[2-methyl-4-(5-oxo-pyrazolidin-1-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (835) 2-cyclohexyl-8-[2-(1H-indol-7-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (836) 2-cyclohexyl-8-[2-(6-trifluoromethyl-1H-indol-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (837) 2-cyclohexyl-8-[2-(6-methyl-1H-indol-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (838) 2-cyclohexyl-8-{2-[1-[2-(2-hydroxy-ethoxy)-ethyl]-1H-indol-5-yl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (839) 2-cyclohexyl-8-[2-(6-trifluoromethyl-1H-benzimidazol-5-yl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (840) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (841) 2-cyclohexyl-8-{2-[4-(1,1-dioxo-1 λ ⁶-thiomorpholine-4-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (842) 2-cyclohexyl-8-{2-[4-(2-dimethylamino-ethylamino)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (843) 8-{2-[2-methyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (844) 4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-N-(2-dimethylamino-ethyl)-3-methyl-benzenesulfonamide;
- (845) 8-(2-{2-methyl-4-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethylamino]-phenyl}-ethanesulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (846) 3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-pyridin-4-yl-benzenesulfonamide;
- (847) 2-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenylamino)-acetamide;
- (848) N-(1-benzyl-piperidin-4-yl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;
- (849) 8-{2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (850) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (851) N-((R)-2,3-dihydroxy-propyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;
- (852) 3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-N-pyridin-3-ylmethyl-benzenesulfonamide;
- (853) N-(4-hydroxy-cyclohexyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;
- (854) N-(2-hydroxy-1,1-bis-hydroxymethyl-ethyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzenesulfonamide;
- (855) 8-{2-[4-(3,4-dihydroxy-butoxy)-3,5-dimethyl-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (856) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (857) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (858) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-{2-[4-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (859) 4-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfo-

nyl]-ethyl)-phenoxy)-piperidine-1-carboxylic acid tert-butyl ester;

(860) 8-{2-[4-(3,4-dihydroxy-butoxy)-3,5-difluoro-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(861) 2-cyclohexyl-8-{2-[2-methyl-4-(2-oxo-azetidin-1-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(862) 1-methyl-3-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(863) 5,5-dimethyl-3-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(864) 5,5-dimethyl-3-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(865) 2-(4-methyl-cyclohexyl)-8-{2-[2-methyl-4-(2-oxo-oxazolidin-3-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(866) 2-(4-methyl-cyclohexyl)-8-{2-[2-methyl-4-(4-oxo-oxazolidin-3-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(867) 8-{2-[1-((R)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(868) 8-(2-[4-[(1H-imidazol-2-yl)-methyl-amino]-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(869) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;

(870) 2-(4-methyl-cyclohexyl)-8-{2-[2-methyl-4-(2-oxo-piperazin-1-yl)-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(872) 2-cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(873) N-(2-fluoro-5-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-N-(2-hydroxy-ethyl)-acetamide;

(874) N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-sulfamide;

(875) N-(3-hydroxy-propyl)-N'-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-sulfamide;

(876) 1-cyanomethyl-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-urea;

(877) 1-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-(2,2,2-trifluoro-ethyl)-urea;

(878) 1-(4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione;

(879) 1-(4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-5-methyl-imidazolidine-2,4-dione;

(880) 3-(4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione;

(881) 3-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(882) 3-(4-{2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidine-2,4-dione;

(883) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-imidazolidine-2,4-dione;

(884) 3-(4-{2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-imidazolidine-2,4-dione;

(885) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1,5,5-trimethyl-imidazolidine-2,4-dione;

(886) (S)-2-amino-3-hydroxy-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-propionamide;

(887) 1-(4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-5,5-dimethyl-imidazolidine-2,4-dione;

(888) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzoic acid N,N'-dimethyl-hydrazide;

(889) 8-{2-[2-methyl-4-(piperidin-4-yloxy)-phenyl]-ethanesulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-

spiro[4.5]dec-1-en-4-one hydrochloride;

(890) 4-{2-[2-(4,4-difluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(891) 4-{2-[2-(4-methanesulfonyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(892) 3,N,N-trimethyl-4-(2-{4-oxo-2-[(E)-2-(3-trifluoromethyl-phenyl)-vinyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;

(893) 3,N,N-trimethyl-4-[2-(4-oxo-2-phenylethynyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-benzamide;

(894) 4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(895) 4-{2-[2-(4-tert-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(896) 4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(897) 3,N,N-trimethyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;

(898) 4-{2-[4-(cyclohexylmethyl-amino)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-diene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(899) 4-{2-[4-dimethylamino-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-diene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(900) 4-{2-[4-[(Z)-hydroxyimino]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(901) 4-{2-[4-[(Z)-methoxyimino]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(902) 4-(2-[2-[4-(2-methoxy-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3,N,N-trimethyl-benzamide;

(903) N-{4-[2-(2-cyclopentyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamide;

(904) N-(3-methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(905) N-(4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(906) N-(4-{2-[2-(4-cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(907) N-(4-{2-[2-(3-cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(908) N-(4-{2-[2-(2-cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(909) N-(4-{2-[2-(4-tert-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(910) N-(4-{2-[2-(4-ethoxymethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(911) N-(3-methyl-4-{2-[4-oxo-2-(4-propoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(912) N-(4-{2-[2-(4-butoxy-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(913) N-(4-{2-[2-(4-isopropoxymethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(914) N-[4-(2-[2-[4-(3-fluoro-propoxy)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;

(915) N-[4-(2-[2-[4-(3-fluoro-propoxy)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;

(916) N-[4-(2-{2-[4-((E)-3,3-difluoro-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;

(917) N-[4-[2-(2-cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamide;

(918) N-[4-[2-(2-adamantan-1-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamide;

nyl)-acetamide;

(919) N-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethoxymethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;

(920) N-[4-(2-{2-[4-(4-chloro-phenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;

(921) N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propoxy)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;

(922) N-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoroethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;

(923) N-(3-methyl-4-{2-[4-oxo-2-((1S,3R)-3-propylcyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl)-acetamide;

(924) N-(3-methyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl)-acetamide;

(925) N-(4-{2-[2-(4,4-dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;

(926) N-(3-methyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-acetamide;

(927) N-[4-(2-{2-[4-(2-methoxy-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;

(928) [3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-carbamic acid tert-butyl ester;

(929) N-(2-hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-isobutylamide;

(930) 2-hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-acetamide;

(931) N-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-2,2,2-trifluoro-N-methyl-acetamide;

(932) N-(3,5-dimethyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-2,2,2-trifluoro-N-methyl-acetamide;

(933) N-(3,5-dimethyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-2,2,2-trifluoro-N-methyl-acetamide;

(934) 1-14-[2-(2-cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl]-1-methyl-urea;

(935) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(936) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoroethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea;

(937) 1-[2-chloro-3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea;

(938) 1-(4-{2-[2-(4,4-dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(939) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propylidene)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea;

(940) 1-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3,5-dimethyl-phenyl]-1-methyl-urea;

(941) 1-(3,5-dimethyl-4-{2-[8-oxo-6-(3-trifluoromethoxyphenyl)-2,5,7-triaza-spiro[3.4]oct-5-ene-2-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(942) N-(4-{(E)-1-fluoro-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3-methyl-phenyl)-N-methyl-acetamide;

(943) 3-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-imidazolidine-2,4-dione;

(944) 3-[3-methyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-imidazolidine-2,4-dione;

(945) 3-[4-(2-{2-[4-((E)-3,3-difluoro-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidine-2,4-dione;

(946) 3-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidine-2,4-dione;

(947) 3-(3-methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfo-

nyl]-ethyl)-phenyl)-imidazolidine-2,4-dione;

(948) 3-[3-methyl-4-(2-[4-oxo-2-[4-(2,2,2-trifluoro-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-imidazolidine-2,4-dione;

(949) 3-(3-methyl-4-[2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-imidazolidine-2,4-dione;

(950) 3-(3-methyl-4-[2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-imidazolidine-2,4-dione;

(951) 3-(4-[2-[2-(4,4-dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3-methyl-phenyl)-imidazolidine-2,4-dione;

(952) 3-(3-methyl-4-[2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-imidazolidine-2,4-dione;

(953) N-(2-chloro-4-[2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-5-methyl-phenyl)-acetamide;

(954) N-(2-chloro-4-[2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3-methyl-phenyl)-acetamide;

(955) N-[4-(2-[2-[4-(3,3-difluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;

(956) 3-[4-(2-[2-[4-(3,3-difluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidine-2,4-dione;

(957) 2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-N-phenyl-benzamide;

(958) 8-(2-[2-methyl-4-[4-(1-methyl-piperidin-4-yl)-piperazine-1-carbonyl]-phenyl]-ethanesulfonyl-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(959) 8-[(E)-2-[4-((R)-3-fluoro-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(960) 8-[(E)-2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(961) 3,5,N,N-tetramethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;

(962) 3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzamide;

(963) 8-[(E)-2-[4-(3-fluoro-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(964) 8-[2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(965) 8-[2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(966) 8-[2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(967) 8-[(E)-2-[2,6-dimethyl-4-(2-oxa-6-aza-spiro[3.3]heptane-6-carbonyl)-phenyl]-ethanesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(968) 8-[(E)-2-[2,6-dimethyl-4-(3-oxo-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(969) 8-[(E)-2-[4-((3R,5S)-3,5-dimethyl-piperazine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(970) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-[(E)-2-[4-(3-hydroxy-3-methyl-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(971) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(972) 8-[(E)-2-[2,6-dimethyl-4-(4-oxo-piperidine-1-carbonyl)-phenyl]-ethanesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(973) 2-[4-(3,3-difluoro-allyl)-cyclohexyl]-8-[2-[4-(4-fluoro-4-hydroxymethyl-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(974) 2-[4-(3,3-difluoro-allyl)-cyclohexyl]-8-[2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(975) 2-(4-ethyl-cyclohexyl)-8-(2-[4-[4-(2-fluoro-ethyl)-piperazine-1-carbonyl]-2-methyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(976) 2-(4-ethyl-cyclohexyl)-8-[2-[2-methyl-4-(4-oxetan-3-yl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (977) 8-{2-[2-methyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (978) 8-{2-[4-(azetidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (979) 8-{2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2-methyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (980) 8-{2-[2,6-dimethyl-4-(pyrrolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (981) 8-{2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (982) 8-{2-[4-(azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (983) 8-{2-[2,6-dimethyl-4-(piperidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (984) 8-{2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (985) 8-{2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (986) 8-{2-[2,6-dimethyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (987) N,N-dimethyl-2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (988) N-methoxy-2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-acetamide;
- (989) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (990) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (991) 8-((E)-2-[2,6-dimethyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (992) 8-((E)-2-[4-(isoxazolidine-2-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (993) 8-((E)-2-[4-[(R)-2,3-dihydroxy-propoxy]-piperidine-1-carbonyl]-2,6-dimethyl-phenyl)-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (994) 3,5,N,N-tetramethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;
- (995) 8-{2-[2,6-dimethyl-4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (996) 3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzoic acid hydrazide;
- (997) 8-((E)-2-[2,6-dimethyl-4-(pyrazolidine-1-carbonyl)-phenyl]-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (998) N-methoxy-3,5,N-trimethyl-4-((E)-2-[2-(4-methylcyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;
- (999) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1000) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1001) 8-((E)-2-[4-(isoxazolidine-2-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1002) 8-((E)-2-[4-(1,1-dioxo-1 λ ⁶-thiomorpholine-4-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1003) 3,5,N,N-tetramethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzamide;
- (1006) 2-cyclohexyl-8-((E)-2-[4-(4-hydroxy-4-trifluoromethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1007) 2-cyclohexyl-8-((E)-2-[2,6-dimethyl-4-(2-oxo-oxazolidine-3-carbonyl)-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1008) 2-(4-butyl-cyclohexyl)-8-((E)-2-[4-[2-(2-hydroxy-ethoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1009) 2-(4-butyl-cyclohexyl)-8-[(E)-2-[4-[2-(2-hydroxy-ethoxy)-ethoxy]-piperidine-1-carbonyl]-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1010) 8-((E)-2-[4-[2-(2,3-dihydroxy-propoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1011) 2-amino-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamide;
- (1012) (3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-methyl-carbamic acid 2-hydroxy-ethyl ester;
- (1013) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3,3-trimethyl-urea;
- (1014) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1,3-dimethyl-urea;
- (1015) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- (1016) 8-((E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl)-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1017) 1-(4-((E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1018) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
- (1019) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[3-(4,4,4-trifluoro-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
- (1020) 1-[3,5-dimethyl-4-((E)-2-[4-oxo-2-[4-(4,4,4-trifluoro-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
- (1021) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(4-pentyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- (1022) 1-(3,5-dimethyl-4-((E)-2-[2-(7-methylsulfanyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;
- (1023) 1-[3,5-dimethyl-4-((E)-2-[2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
- (1024) 1-[3,5-dimethyl-4-[(E)-2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-1-methyl-urea;
- (1025) 1-[3,5-dimethyl-4-[(E)-2-(2-non-3-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-1-methyl-urea;
- (1026) 1-[3,5-dimethyl-4-((E)-2-[2-[10-(3-methyl-oxetan-3-yl)-decyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl]-1-methyl-urea;
- (1027) 1-[3,5-dimethyl-4-[(E)-2-(2-non-1-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-phenyl]-1-methyl-urea;
- (1028) 1-(4-[2-[2-(4-fluoro-3-trifluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1029) 1-(4-[2-[2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1030) 1-(4-[2-[2-(4-tert-butyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1031) 1-[3,5-dimethyl-4-(2-[2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1032) 1-[3,5-dimethyl-4-(2-[2-[10-(3-methyl-oxetan-3-yl)-decyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1033) 2-[8-(2-[4-(tert-butoxycarbonyl-methyl-amino)-2,6-dimethyl-phenyl]-ethanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-pyrrolidine-1-carboxylic acid tert-butyl ester;
- (1034) (4-[2-[2-(1,1-difluoro-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-methyl-carbamic acid tert-butyl ester;
- (1035) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(8,8,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1036) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1037) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(4-isopropylidene-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1038) 1-[3,5-dimethyl-4-(2-[4-oxo-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1039) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(5,6,7,8-tetrahydro-naphthalen-2-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- (1040) 1-(4-{2-[2-((1S,3S,5R)-3,5-dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1041) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(1-propyl-butyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- (1042) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1043) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(6,6,7,7,7-pentafluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- (1044) N-(3,5-dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-N-methyl-acetamide;
- (1045) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- (1046) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- (1047) N-(3,5-dimethyl-4-{2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-N-methyl-acetamide;
- (1048) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[1-(4,4,4-trifluoro-butyl)-cyclopropyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1050) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- (1051) 1-(4-{2-[2-(cis-3,4-dimethyl-cyclopentyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1052) 1-{4-[2-(2-dicyclopropylmethyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;
- (1053) 1-{4-[2-(2-bicyclo[3.3.1]non-9-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;
- (1054) 1-{4-[2-((1R,5S)-2-bicyclo[3.2.1]oct-3-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;
- (1055) 1-(3,5-dimethyl-4-{2-[2-(3-methyl-cyclopentyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;
- (1056) 1-{4-[2-(2-bicyclo[2.2.1]hept-7-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;
- (1057) 1-(4-{2-[2-((1S,2S)-2-hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1058) 1-{3,5-dimethyl-4-[2-(4-oxo-2-spiro[2.5]oct-6-yl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- (1059) 1-(4-{2-[2-(4-difluoromethylene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1060) N-[4-(2-{2-[4-(2,2-difluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;
- (1061) N-[4-(2-{2-[4-(2-fluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;
- (1062) N-[4-(2-{2-[4-(4-butyl-4-fluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide];
- (1063) N-[4-(2-{2-[4-((E)-3-fluoro-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide];
- (1064) N-[4-(2-{2-[4-(3-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide];
- (1065) N-[4-(2-{2-[4-(3-chloro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide];
- (1066) N-[4-(2-{2-[4-(3-fluoro-propylidene)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide];

- (1067) N-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1068) N-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- 5 (1069) N-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;
- (1070) N-[4-(2-{2-[4-((E)-3-fluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1071) N-[4-(2-{2-[4-(2,2-difluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- 10 (1072) N-[4-(2-{2-[4-(2-fluoro-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1073) N-[4-(2-{2-[4-fluoro-4-(3-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- 15 (1074) N-(4-{2-[2-(4-ethynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;
- (1075) N-(4-{2-[2-(4-difluoromethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;
- (1076) N-[4-(2-{2-[4-(3,3-difluoro-propylidene)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- 20 (1077) N-[4-(2-{2-[4-(2-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- (1078) N-[4-(2-{2-[4-(1-fluoro-1-methyl-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-3-methyl-phenyl]-acetamide;
- 25 (1079) N-(4-{2-[2-(4-butylidene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;
- (1080) N-[3-methyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-acetamide;
- (1081) 1-{3,5-dimethyl-4-[2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- 30 (1082) 1-{3,5-dimethyl-4-[2-(2-non-3-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- (1083) 1-{3,5-dimethyl-4-[2-(2-non-1-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- 35 (1084) 1-(3,5-dimethyl-4-{1-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-cyclopropylmethyl}-phenyl)-1-methyl-urea;
- (1085) 1-(4-{2,2-difluoro-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1086) 1-{4-[2,2-difluoro-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-urea;
- 40 (1087) 1-{3,5-dimethyl-4-[2-(4-oxo-2-[1,1';2',1"]terphenyl-3-yl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl}-1-methyl-urea;
- (1088) 1-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea;
- 45 (1089) 1-[3,5-dimethyl-4-((E)-2-{4-oxo-2-[3-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-vinyl)-phenyl]-1-methyl-urea;
- (1090) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-propyl}-phenyl)-1-methyl-urea;
- (1091) 2-[4-fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 50 (1092) 2-[4-fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1093) 2-(4-chloro-3-trifluoromethoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- 55 (1094) 2-(3-fluoro-4-trifluoromethoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1095) 2-(3,4-bis-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1096) 8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1097) 8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(2,2,2-trifluoro-ethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1098) 2-(4-fluoro-3-methyl-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1099) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1100) 8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1101) 8-[1,1-difluoro-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1102) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-[(E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1103) 8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1104) 6-(4-methyl-cyclohexyl)-2-(2-naphthalen-1-yl-ethanesulfonyl)-2,5,7-triaza-spiro[3.4]oct-5-en-8-one;
- (1106) (11-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-undecyl)-carbamic acid tert-butyl ester;
- (1107) 8-[(E)-2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-(9-hydroxy-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1108) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1109) 14-(8-[(E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecanoic acid;
- (1110) 1-(3,5-dimethyl-4-[(E)-2-[2-(4-[11,1-²H₃]methyl-(4-²H₁)cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1111) 1-(4-[(E)-2-[2-(4-fluoro-3-trifluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1112) 1-(3,5-dimethyl-4-[(E)-2-[2-(3-nonafluorobutyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1113) 1-(4-[(E)-2-[2-(4-chloro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1114) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1115) 1-(4-[(E)-2-[2-(6-ethoxy-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1116) 1-[3,5-dimethyl-4-[(E)-2-[4-oxo-2-[3-(6,6,6-trifluoro-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl]-1-methyl-urea;
- (1117) 1-[3,5-dimethyl-4-[(E)-2-[4-oxo-2-[4-(6,6,6-trifluoro-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl]-1-methyl-urea;
- (1118) 1-(4-[(E)-2-[2-(11-fluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1119) 1-(4-[(E)-2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1120) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-((E)-6-phenyl-hex-5-enyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1121) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-((E)-9-phenyl-non-8-enyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1122) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(5-propylsulfanyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1123) 1-(4-[(E)-2-[2-(7-methoxy-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1124) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(5-propoxy-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;
- (1125) 1-(4-[(E)-2-[2-(11,11-difluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1126) 1-(3,5-dimethyl-4-[(E)-2-[4-oxo-2-(3-propyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;

nyl)-phenyl)-1-methyl-urea;

(1127) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(2-propyl-benzofuran-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;

(1128) 1-(4-((E)-2-[2-(3-methoxy-4-pentyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1129) 1-(4-((E)-2-[2-(4-[1,1,2,2,2-²H₅]ethyl-cyclohex-3-enyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1130) 1-(4-((E)-2-[2-(4-[1,1,2,2,2-²H₅]ethyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-urea;

(1131) 12-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid;

(1132) 12-(8-((E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid ethyl ester;

(1133) 1-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;

(1134) 1-(3,5-dimethyl-4-((Z)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-phenyl)-1-methyl-urea;

(1135) 1-[3,5-dimethyl-4-(2-[4-oxo-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

(1136) 1-(3,5-dimethyl-4-[2-[4-oxo-2-(3'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(1137) 1-(3,5-dimethyl-4-[2-[4-oxo-2-(4-trimethylsilanyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(1138) 1-(4-[2-[2-((1R,3R,5S)-3,5-bis-trifluoromethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1139) 1-(3,5-dimethyl-4-[2-[4-oxo-2-(4-trimethylsilanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(1140) 1-[3,5-dimethyl-4-[2-(4-oxo-2-tridecyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-1-methyl-urea;

(1141) 1-[3,5-dimethyl-4-[2-(4-oxo-2-undecyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-1-methyl-urea;

(1142) 1-[3,5-dimethyl-4-[2-(2-octyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-phenyl]-1-methyl-urea;

(1143) 1-(4-[2-[2-((1S,3R)-3-hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1144) 1-[3,5-dimethyl-4-[2-[2-[3-(3-methyl-butyl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl]-1-methyl-urea;

(1145) 1-(4-[2-[2-((1S,3R)-3-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1146) 1-[3,5-dimethyl-4-(2-[2-((1S,3S)-3-(3-methyl-butyl)-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

(1147) 1-(3,5-dimethyl-4-12-[4-oxo-2-(3,3,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1148) 1-(4-[2-[2-(4-[1,1,2,2,2-²H₅]ethyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1149) 1-(4-[2-[2-(4-[1,1,2,2,2-²H₅]ethyl-cyclohex-3-enyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1150) 1-(4-[2-[2-(4-chloro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1151) 1-(3,5-dimethyl-4-12-[4-oxo-2-(1,9,9,9-tetrafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl)-1-methyl-urea;

(1152) 1-(3,5-dimethyl-4-[2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-phenyl)-1-methyl-urea;

(1153) 1-(4-[2-[2-(11-fluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1154) 1-(4-[2-[2-(6-ethylsulfanyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-urea;

(1155) 1-[3,5-dimethyl-4-[2-(4-oxo-2-[1,1';3',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfo-

nyl)-ethyl]-phenyl)-1-methyl-urea;

(1156) 1-(4-{2-[2-(11,11-difluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;

(1157) 1-{3,5-dimethyl-4-[(E)-2-(4-oxo-2-[1,1',3',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-1-methyl-urea;

(1158) 8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1159) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1160) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1161) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-[(E)-2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1162) 2-(4-fluoro-3-trifluoromethyl-phenyl)-8-[(E)-2-[4-(4-hydroxymethyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1163) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1164) 8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-2-(4-fluoro-3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1165) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1166) 8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl]-2-(4-fluoro-3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1167) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1168) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl)-vinyl]-indol-1-yl}-acetic acid;

(1169) [3-(3-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-ureido]-acetic acid;

(1170) 12-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid;

(1171) 10-(8-{(E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid;

(1172) 10-(8-{(E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid amide;

(1173) 12-(8-{(E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid amide;

(1174) 12-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid amide;

(1175) 14-(8-{(E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethenesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecanoic acid amide;

(1176) 14-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecanoic acid amide;

(1178) 8-[2-(2-amino-5,7-dimethyl-benzoxazol-6-yl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1179) 2-cyclohexyl-8-[2-[2,6-dimethyl-4-(2-oxo-oxazolidine-3-carbonyl)-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1180) N-methoxy-3,5,N-trimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(1181) 8-{2-[4-(isoxazolidine-2-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one ;

(1182) 8-{2-[4-(isoxazolidine-2-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1183) 8-{2-[4-(1,1-dioxo-1λ⁶-thiomorpholine-4-carbonyl) - 2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1184) 8-(2-[4-[4-((R)-2,3-dihydroxy-propoxy)-piperidine-1-carbonyl]-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

(1185) 2-(3,4-dichloro-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1186) 2-(3-chloro-4-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1187) [3-(8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-phenyl]-acetonitrile;
- (1188) 2-(3-chloro-4-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1189) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-pentafluorosulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1190) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-pentafluorosulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1191) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(3-trifluoromethyl-phenoxy-methyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1192) 2-(4-fluoro-2,3-dimethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1193) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1194) 2-benzo[1,3]dioxol-5-yl-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1195) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-pentafluoroethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1196) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-pentafluoroethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1197) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1198) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1199) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,3,3-tetrafluoro-propoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1200) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,2-trifluoro-ethoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1201) 2-[3-chloro-4-(2,2,3,3-tetrafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1202) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(4,4,5,5-pentafluoro-pentyloxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1203) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,3,3-pentafluoro-propoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1204) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(4,4,4-trifluoro-butoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1205) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(3-trifluoromethyl-phenoxy-methyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1206) 2-[3-(1,1-difluoro-ethyl)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1207) 2-[3-fluoro-4-(4,4,5,5-pentafluoro-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1208) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(1H-indol-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1209) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1210) 2-[4-chloro-3-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1211) 2-[3-(1,1,2,2,3,3,4,4,4-²H₉)butoxy-4-fluoro-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1212) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(2,2,3,3-tetrafluoro-2,3-dihydro-benzo[1,4]dioxin-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1213) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(1,1,2,2-tetrafluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1214) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-methyl-4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1215) 2-[4-fluoro-3-(3-fluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1216) 2-[3-chloro-4-(2,2,2-trifluoro-ethoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1217) 2-(3-difluoromethyl-4-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1218) 2-[4-fluoro-3-(4,4,4-trifluoro-butoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1219) 2-(4-difluoromethyl-3-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1220) 2-(4-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1221) 2-(4-[1,1,2,2,2-²H₅]ethoxy-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1222) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-[1,2,2,2,2,2-²H₇]isopropoxy-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1223) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1224) 2-(3-[1,1,2,2-²H₅]ethoxy-4-fluoro-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1225) 2-(4-fluoro-3-[1,2,2,2,2-²H₇]isopropoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1226) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[3-(2,2,3,3-tetrafluoro-propoxy)-5-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1227) 2-[4-chloro-3-(4,4,4-trifluoro-butoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1228) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[3-(2,2,3,3,3-pentafluoro-propoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1229) 2-[4-chloro-3-(2,2,3,3-tetrafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1230) 2-[4-chloro-3-(2,2,3,3,3-pentafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1231) 2-(3-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-4-chlorophenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1232) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-[3-(4,4,4-trifluoro-butoxy)-4-trifluoromethoxy-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1233) 2-(3-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-5-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1234) 8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1235) 8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(3-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1236) 2-(3-chloro-4-trifluoromethyl-phenyl)-8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1237) 2-(3,4-dimethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1238) 8-((E)-2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(4-methoxy-3-methyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1239) 2-(4-fluoro-2,5-dimethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1240) 2-(4-fluoro-2,3-dimethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1241) 2-(3-chloro-4-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1242) 2-(3-chloro-4-fluoro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1243) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl)-2-(3-pentafluoroethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1244) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-pentafluoroethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1245) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-pentafluorosulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1246) 2-[3-chloro-4-(2,2,2-trifluoro-ethoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1247) 2-[3-chloro-4-(2,2,3,3-tetrafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1248) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,2-trifluoro-ethoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1249) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,3,3,3-pentafluoro-propoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1250) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(4,4,4-trifluoro-butoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1251) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,3,3-tetrafluoro-propoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1252) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(4,4,5,5-pentafluoro-pentyloxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1253) 2-(4-[1,1,2,2,2-²H₅]ethoxy-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1254) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1255) 2-(4-difluoromethyl-3-fluoro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1256) 2-(3-difluoromethyl-4-fluoro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1257) 2-(3,4-dichloro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1258) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(3-methyl-4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1259) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(2-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1260) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-fluoro-4-(2,2,2-trifluoro-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1261) 2-(4-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-3-trifluoromethyl-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1262) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-[1,2,2,2,2,2-²H₆]isopropoxy-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1263) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1264) 2-[4-fluoro-3-(3-fluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1265) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[4-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-3-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1266) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-[3-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1267) 2-[3-(1,1-difluoro-ethyl)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1268) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(1H-indol-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1269) 8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1270) 2-[4-chloro-3-(2,2,3,3,3-pentafluoro-propoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1271) 2-[4-chloro-3-(2,2,2-trifluoro-1,1-dimethyl-ethoxy)-phenyl]-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1272) 2-(3-[1,1,2,2,2-²H₅]ethoxy-4-fluoro-phenyl)-8-((E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethenesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1273) 8-[(E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(4,4,4-trifluoro-butoxy)-4-trifluoromethoxy-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1274) 8-[(E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1275) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1276) 8-[2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1277) 8-[2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(3-methyl-4-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1278) 2-(3-chloro-4-trifluoromethyl-phenyl)-8-[2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1279) 8-[2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluoromethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1280) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1281) 8-[2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-2-(2,2,3,3-tetrafluoro-2,3-dihydro-benzo[1,4]dioxin-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1282) [3-(8-[(E)-2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-phenyl]-acetonitrile;
- (1283) 8-[(E)-2-[2-methyl-4-(piperidin-4-yloxy)-phenyl]-ethanesulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1284) 8-[2-(4-aminomethyl-2-methyl-phenyl)-ethanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1285) 2-(4-methyl-cyclohexyl)-8-[(E)-2-[2-methyl-4-(2-oxo-piperazin-1-yl)-phenyl]-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1286) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-2-pyrrolidin-2-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (1287) 2-(1,1-difluoro-ethyl)-8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethanesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (1288) 8-[(E)-2-(2,6-dimethyl-4-[1,1,1-²H₃]-methylamino-phenyl)-ethanesulfonyl]-2-(4-fluoro-3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one hydrochloride;
- (1289) 8-[(E)-2-(2,6-dimethyl-4-methylaminomethyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1290) 8-[(E)-2-(4-aminomethyl-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1291) 8-[(E)-2-(4-aminomethyl-2,6-dimethyl-phenyl)-ethanesulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1292) 8-[(E)-2-(2,6-dimethyl-4-methylaminomethyl-phenyl)-ethanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1293) 8-[2-(4-amino-3-chloro-2-methyl-phenyl)-ethanesulfonyl]-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1294) N-(1-acetyl-piperidin-4-yl)-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-phenyl)-acetamide;
- (1295) 8-[2-[4-(4,5-dihydro-thiazol-2-ylamino)-2-methyl-phenyl]-ethanesulfonyl]-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1296) N-[4-(2-[2-[4-(4-fluoro-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamide;
- (1297) 1-(3,5-dimethyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1298) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1299) 3-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-5-hydroxymethyl-imidazolidine-2,4-dione;
- (1300) 1-[3,5-dimethyl-4-((E)-2-[14-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl)-benzyl]-1-methyl-urea;
- (1301) 1-(3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl]-benzyl)-1-methyl-urea;

- (1302) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1303) 1-(3,5-dimethyl-4-{(E)-2-[4-oxo-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea;
- (1304) 1-(3,5-dimethyl-4-{(E)-2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea;
- (1305) 1-(3,5-dimethyl-4-{(E)-2-[4-oxo-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea;
- (1306) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-trifluoromethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1307) 1-(4-{(E)-2-[2-(11-hydroxy-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1308) 11-(8-{(E)-2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-undecyl)-carbamic acid tert-butyl ester;
- (1309) 1-(4-{(E)-2-[2-(9-hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1310) 1-(4-{2-[2-(4-isopropylidene-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1311) 1-(4-{2-[2-(1,1-difluoro-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1312) 1-(4-{(E)-2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-[1,1,1-²H₃]methyl-urea;
- (1313) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl]-urea;
- (1314) 1-[3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl]-urea;
- (1315) 1-(4-{(E)-2-[2-(11-amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1316) 1-[3,5-dimethyl-4-{(E)-2-[4-oxo-2-[5-(propane-1-sulfinyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea;
- (1317) 1-(4-{2-[2-(6-ethanesulfinyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1318) 1-[3,5-dimethyl-4-{(E)-2-[4-oxo-2-[5-(propane-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea;
- (1319) 1-(4-{(E)-2-[2-(7-methanesulfonyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1320) 1-(4-{2-[2-(6-ethanesulfonyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1321) 1-(4-{(E)-2-[2-(9,9-difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1322) 1-(4-{2-[2-(9-hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1323) 1-(4-{2-[2-(9,9-difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1324) 1-(4-{2-[2-(9-amino-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1325) 1-(4-{(E)-2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1326) 1-(4-{(Z)-2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1327) 1-(4-{2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1328) 8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[3-(4,4,5,5,5-pentafluoropentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1329) 8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-fluoro-3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1330) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;

- (1331) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1332) 8-{2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethanesulfonyl}-2-(4-fluoro-3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1333) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1334) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-{2-[4-(4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1335) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1336) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1337) 2-(4-fluoro-3-trifluoromethoxy-phenyl)-8-{2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1338) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[3-(4,4,4-trifluoro-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea;
- (1339) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-phenyl]-1-methyl-urea;
- (1340) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1341) 1-(3,5-dimethyl-4-{2-[2-(3-nonafluorobutyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1342) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(4-pentyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1343) 1-(3,5-dimethyl-4-{2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1344) 1-(4-{2-[2-(6-ethoxy-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1345) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[3-(6,6,6-trifluoro-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1346) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(6,6,6-trifluoro-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1347) 12-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecanoic acid ethyl ester;
- (1348) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(9-phenyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1349) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(6-phenyl-hexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1350) 14-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecanoic acid;
- (1351) 1-(4-{2-[2-(7-methoxy-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1352) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(5-propoxy-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1353) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(3-propyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1354) 1-(4-{2-[2-(3-methoxy-4-pentyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1355) 1-(3,5-dimethyl-4-{2-[4-oxo-2-(2-propyl-benzofuran-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1356) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[5-(propane-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1357) 1-(4-{2-[2-(7-methanesulfonyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1358) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;
- (1359) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[3-(4,4,4-trifluoro-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-phenyl]-1-methyl-urea;

- (1360) [11-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-undecyl]-carbamic acid tert-butyl ester;
- (1361) 1-(4-{2-[2-(11-hydroxy-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1362) 10-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid;
- (1363) 10-(8-{2-[2,6-dimethyl-4-(1-methyl-ureido)-phenyl]-ethanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decanoic acid amide;
- (1364) 1-(3,5-dimethyl-4-{2-[2-(4-[1,1,1-²H₃]-methyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-1-methyl-urea;
- (1365) 1-(3,5-dimethyl-4-{(E)-2-[4-oxo-2-(9-phenyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-vinyl}-phenyl)-1-methyl-urea;
- (1366) 1-(4-{2-[2-(11-amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1367) 3-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-4-ethoxy-cyclobut-3-ene-1,2-dione;
- (1368) 3-amino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dione;
- (1369) 3-dimethylamino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dione;
- (1370) N-(4-{2-[2-(4-ethynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamide;
- (1371) 1-(4-{2-[4-{(Z)-hydroxyimino}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-urea;
- (1372) 2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic acid 2-methyl-benzylamide;
- (1373) 2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonic acid (2-o-tolyl-ethyl)-amide;
- (1374) 2-cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-2-hydroxy-ethanesulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1375) 2-cyclohexyl-8-(2-oxo-2-o-tolyl-ethanesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1376) 2-cyclohexyl-8-(2-o-tolyl-ethynesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1377) 2-cyclohexyl-8-[2-(1H-indol-4-yl)-ethynesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1378) 3,5,N,N-tetramethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzamide;
- (1379) 3,5,N,N-tetramethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;
- (1380) 8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1381) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1382) 8-{2-[4-(4-hydroxy-4-methyl-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1383) 8-{2-[4-(4-fluoromethyl-4-hydroxy-piperidine-1-carbonyl)-2,6-dimethyl-phenyl]-ethanesulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1384) 8-(2-[4-{(R)-2,3-dihydroxy-propoxy}-piperidine-1-carbonyl]-2,6-dimethyl-phenyl)-ethanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1385) 3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzoic acid hydrazide;
- (1386) 8-{2-[2,6-dimethyl-4-(pyrazolidine-1-carbonyl)-phenyl]-ethanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-one;
- (1387) N,N-dimethyl-2-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (1388) N-methoxy-2-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;
- (1389) 1-[3,5-dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl)-benzyl]-1-methyl-urea;
- (1390) 1-(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzyl)-1-methyl-urea;
- (1391) 1-(4-{2-[2-(4-fluoro-3-trifluoromethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-

3,5-dimethyl-phenyl)-1-[1,1,1-²H₃]methyl-urea;

(1392) 2-cyclohexyl-8-{2-[1-((2S,3S)-2,3,4-trihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl}-1,3,8-triazaspiro[4.5]dec-1-en-4-one;

(1393) 8-{2-[1-((2S,3S)-4-benzyloxy-2,3-dihydroxy-butyl)-1H-indol-4-yl]-ethanesulfonyl}-2-cyclohexyl-1,3,8-triazaspiro[4.5]dec-1-en-4-one;

(1394) N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzyl]-acetamide;

(1395) 3,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenylamino)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(1396) 3,N,N-trimethyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenylamino)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(1397) 3,N,N-trimethyl-4-{2-[2-(methyl-phenyl-amino)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-benzamide;

(1398) 4-{2-[2-(benzyl-methyl-amino)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(1399) 4-{2-[2-(cyclohexylmethyl-amino)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(1400) 3,N,N-trimethyl-4-(2-{2-[methyl-(4-trifluoromethyl-phenyl)-amino]-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl}-ethyl)-benzamide;

(1401) N-(3-methyl-4-{2-[4-oxo-2-(4-trifluoromethyl-phenylamino)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(1402) N-(3-methyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenylamino)-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-phenyl)-acetamide;

(1403) 4-{2-[2-(4-butyl-piperidin-1-yl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(1404) 4-{2-[2-(4-butyl-cyclohexylamino)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamide;

(1405) 4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-N-pent-4-enyl-benzamide;


(1419) N-(2-allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluoromethyl-phenyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamide;

(1443) dec-9-enoic acid {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-amide; or

(1446) hept-6-enoic acid {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triazaspiro[4.5]dec-1-ene-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-amide,

or a pharmacologically acceptable salt thereof.


11. The compound or a pharmacologically acceptable salt thereof according to any of claims 1 to 5, wherein U represents C1-C6 alkylene or any group selected from groups represented by the following formula:



A is O;

R_{46} is selected from hydrogen, C1-C10 alkyl, C1-C10 haloalkyl and C1-C10 hydroxyalkyl;

V is selected from:




E is pyrrolidine or piperidine optionally substituted with a hydroxyl group; and
 R₇ is selected from:

- 1) hydrogen,
- 2) C1-C10 alkyl and
- 3) C1-C10 hydroxyalkyl.


12. A pharmaceutical composition comprising the compound or a pharmacologically acceptable salt thereof according to any of claims 1 to 11 as an active ingredient.
13. A prophylactic or therapeutic agent for use in the treatment or prevention of osteoporosis, fracture, osteomalacia, arthritis, thrombocytopenia, hypoparathyroidism, hyperphosphatemia or tumoral calcinosis, comprising the compound or a pharmacologically acceptable salt thereof according to any of claims 1 to 11 as an active ingredient.
14. Use of the compound or a pharmacologically acceptable salt thereof according to any of claims 1 to 11 for the manufacture of a prophylactic or therapeutic agent for use in the treatment or prevention of osteoporosis, fracture, osteomalacia, arthritis, thrombocytopenia, hypoparathyroidism, hyperphosphatemia or tumoral calcinosis.
15. The compound or a pharmacologically acceptable salt thereof according to any of claims 1 to 11 for use in the treatment or prevention of osteoporosis, fracture, osteomalacia, arthritis, thrombocytopenia, hypoparathyroidism, hyperphosphatemia or tumoral calcinosis.

Patentansprüche

1. Eine Verbindung, dargestellt durch die folgende allgemeine Formel (1) oder (2):



(1)



oder ein pharmakologisch verträgliches Salz einer Verbindung, dargestellt durch Formel (1) oder (2):

[wobei in der allgemeinen Formel (1)
W ausgewählt ist aus:

- 1) einer Einfachbindung;
- 2) C1-C10-Alkylen, das gegebenenfalls einen Carbonylrest enthält, wobei das Alkylen gegebenenfalls mit einem Halogenatom(en) und/oder einem Hydroxylrest(en) substituiert ist,
- 3) C2-C10-Alkenylen, gegebenenfalls substituiert mit einem Halogenatom(en),
- 4) C2-C10-Alkinylen,
- 5) Arylen, gegebenenfalls substituiert mit einem Halogenatom(en),
- 6) Heteroarylen, gegebenenfalls substituiert mit einem Halogenatom(en),
- 7) C1-C10-Heteroalkylen, gegebenenfalls substituiert mit einem Halogenatom(en),
- 8) -NH-, -NHCH₂- oder -NHCH₂CH₂-,
- 9) Cycloalkylen und
- 10) -(Cycloalkylen)-CH₂-;

X ausgewählt ist aus der folgenden Bindung oder den folgenden Resten:


- 1) einer Einfachbindung,
 - 2) C1-C10-Alkylen, gegebenenfalls substituiert mit einem Halogenatom(en) oder Cycloalkyl,
 - 3) C2-C10-Alkenylen, gegebenenfalls substituiert mit einem Halogenatom(en),
 - 4) C2-C10-Alkinylen, gegebenenfalls substituiert mit einem Halogenatom(en),
 - 5) C1-C10-Oxyalkylen, gegebenenfalls substituiert mit einem Halogenatom(en), und
 - 6) -NR₄₇-
- wobei R₄₇ ausgewählt ist aus

- i) einem Wasserstoffatom und
- ii) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Halogenatom(en);

Y ausgewählt ist aus:

- 1) einem Sauerstoffatom,
- 2) einem Schwefelatom und
- 3) =NR₃₇

oder 4) Y das in der folgenden Formel (A) gezeigte -NR₃₈R₃₉ ist:



10 das tautomer sein kann;
R₃₇ ausgewählt ist aus:

- 15
- 1) Wasserstoff,
 - 2) Hydroxy und
 - 3) C1-C10-Alkoxy;

20 R₃₈ und R₃₉ unabhängig ausgewählt sind aus Wasserstoff oder C1-C10-Alkyl, gegebenenfalls substituiert mit Cycloalkyl, oder R₃₈ und R₃₉ aneinander gebunden sein können, um einen Ring, ausgewählt aus der Gruppe bestehend aus Azetidiny, Pyrrolidiny, Piperidiny, Piperaziny und Morpholiny, zu bilden, und der Ring gegebenenfalls mit C1-C10-Alkyl substituiert ist;

m eine ganze Zahl von 0 bis 2 darstellt;

n eine ganze Zahl von 0 bis 2 darstellt;

R₁ ausgewählt ist aus:

- 25
- 1) Wasserstoff,
 - 2) Cycloalkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R₄, R₅ und R₆,
 - 3) einem Heterocyclus, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R₂₅, R₄, R₅ und R₆,
 - 4) Aryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R₃, R₄, R₅ und R₆, und
 - 5) Heteroaryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R₂₅, R₄, R₅ und R₆;

30 R₃ ausgewählt ist aus:

- 35
- 1) -CONR₇R₈,
 - 2) -OR₉,
 - 3) -NR₉R₁₀,
 - 4) -N(R₉)COR₁₁,
 - 5) -N(R₉)SO₂R₁₂,
 - 6) -SO₂R₁₅,
 - 7) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, -COR₁₆ und NR₁₃R₁₄,
 - 8) Heteroaryl, das gegebenenfalls C1-C10-Alkyl und/oder C1-C10-Alkoxy als einen Substituent(en) aufweist, und
 - 9) -N(R₉)CSR₁₁;

45 R₄ ausgewählt ist aus:

- 50
- 1) einem Halogenatom,
 - 2) Cyano,
 - 3) Nitro,
 - 4) Amino,
 - 5) -NHCOR₂₆,
 - 6) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus Hydroxycarbonyl, C1-C10-Alkoxycarbonyl und Aminocarbonyl,
 - 7) C1-C10-Halogenalkyl,
 - 8) C1-C10-Alkoxy,
 - 9) C1-C10-Halogenalkylcarbonyl,
 - 10) -COR₁₆,
 - 11) C1-C10-Hydroxyalkyl und
- 55

12) C1-C10-Heteroalkyl;

R₅ ausgewählt ist aus einem Halogenatom, C1-C10-Alkyl, C1-C10-Halogenalkyl und C1-C10-Alkoxy;

R₆ ausgewählt ist aus einem Halogenatom, C1-C10-Alkyl und C1-C10-Halogenalkyl;

R₇ ausgewählt ist aus:

- 1) Wasserstoff,
- 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus Amino und C1-C10-Alkylamino,
- 3) C1-C10-Hydroxyalkyl,
- 4) C1-C10-Halogenalkyl,
- 5) C1-C10-Heteroalkyl,
- 6) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Hydroxylrest, C1-C10-Alkylamino und C2-C10-Alkenyl,
- 7) Aryl,
- 8) Heteroaryl,
- 9) Aryl-C1-C10-alkyl,
- 10) einem Heterocyclus, gegebenenfalls substituiert mit C1-C10-Alkyl,
- 11) $-(CH_2)_L COR_{16}$ (wobei L eine ganze Zahl von 1 bis 4 bedeutet),
- 12) C1-C10-Alkoxy,
- 13) C2-C10-Alkenyl und
- 14) $-NR_{40}R_{41}$;

R₄₀ und R₄₁ unabhängig ausgewählt sind aus Wasserstoff, C1-C10-Alkyl und C1-C10-Alkylcarbonyl, oder R₄₀ und R₄₁ aneinander gebunden sein können, um einen Ring, ausgewählt aus Azetidiny, Pyrrolidiny, Piperidiny, Piperaziny und Morpholiny, zu bilden, und der Heterocyclus gegebenenfalls mit C1-C10-Alkyl substituiert ist; R₈ ausgewählt ist aus Wasserstoff und C1-C10-Alkyl, gegebenenfalls substituiert mit einem Halogenatom(en) und/oder einem Hydroxylrest(en);

R₇ und R₈ gebunden sein können, um einen 4- bis 7-gliedrigen Heterocyclus zu bilden, der gegebenenfalls ein zusätzliches Element(e) oder einen zusätzlichen Rest(e), unabhängig ausgewählt aus O, N, S, SO und SO₂, enthält, und der Heterocyclus gegebenenfalls Carbonyl enthält und der Heterocyclus gegebenenfalls mit einem Substituent(en) substituiert ist, unabhängig ausgewählt aus:

- 1) einem Halogenatom,
- 2) C1-C10-Alkyl, das gegebenenfalls C1-C10-Alkylamino als einen Substituent(en) aufweist,
- 3) C1-C10-Halogenalkyl,
- 4) einem Hydroxylrest,
- 5) C1-C10-Hydroxyalkyl,
- 6) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 7) Aryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 8) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 9) einem Heterocyclus, gegebenenfalls substituiert mit C1-C10-Alkyl,
- 10) Heteroaryl, gegebenenfalls substituiert mit C1-C10-Alkyl,
- 11) Heterocyclyl-C1-C10-alkyl,
- 12) $-COR_{16}$,
- 13) $-NR_{19}R_{20}$,
- 14) $-SO_2R_{21}$,
- 15) C1-C10-Alkoxy-C1-C10-alkyl, das gegebenenfalls einen Hydroxylrest(e) als einen Substituent(en) aufweist, und
- 16) C1-C10-Hydroxyalkyloxy, wobei das Wasserstoffatom des Hydroxylrests gegebenenfalls durch C1-C10-Hydroxyalkyl ersetzt ist, und

der Heterocyclus weiter einen Spiroring zusammen mit einem 4- bis 6-gliedrigen Heterocyclus bilden kann und der gebundene 4- bis 6-gliedrige Heterocyclus gegebenenfalls O und N als ringbildende Elemente zusätzlich zu Kohlenstoffatomen enthält und das Kohlenstoffatom/die Kohlenstoffatome oxidiert sein können, um Carbonyl

zu bilden, und der 4- bis 6-gliedrige Heterocyclus gegebenenfalls weiter mit C1-C10-Alkyl substituiert ist;
 R_{16} ausgewählt ist aus:

- 1) einem Hydroxylrest,
- 2) C1-C10-Alkoxy,
- 3) $NR_{17}R_{18}$ and
- 4) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Substituent(en), ausgewählt aus einem Halogenatom, einem Hydroxylrest, C1-C10-Alkoxycarbonyl oder C1-C10-Alkylamino;

R_{17} ausgewählt ist aus:

- 1) Wasserstoff,
- 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus Aryl, Amino, C1-C10-Alkylamino, C1-C10-Alkylcarbonylamino und einem Hydroxylrest,
- 3) Heteroaryl und
- 4) C1-C10-Alkoxy;

R_{18} ausgewählt ist aus Wasserstoff, C1-C10-Alkyl und C1-C10-Hydroxyalkyl;

R_{17} und R_{18} aneinander gebunden sein können, um einen Ring, ausgewählt aus Azetidiny, Pyrrolidiny, Piperidiny, Piperaziny und Morpholiny, zu bilden, und der Ring gegebenenfalls mit einem Rest(en), unabhängig voneinander ausgewählt aus C1-C10-Alkyl, einem Halogenatom und C1-C10-Alkoxycarbonyl, substituiert ist;
 R_{19} ausgewählt ist aus Wasserstoff, C1-C10-Alkyl, C1-C10-Halogenalkyl, C1-C10-Alkylcarbonyl, C1-C10-Hydroxyalkyl, C1-C10-Aminoalkyl, C1-C10-Alkoxycarbonyl und C1-C10-Heteroalkyl;

R_{20} ausgewählt ist aus Wasserstoff und C1-C10-Alkyl;

R_{19} und R_{20} gebunden sein können, um einen Ring, ausgewählt aus Azetidiny, Pyrrolidiny, Piperidiny, Piperaziny und Morpholiny, zu bilden, und der Ring gegebenenfalls mit einem Rest(en), unabhängig voneinander ausgewählt aus C1-C10-Alkyl und einem Halogenatom, substituiert ist;

R_{21} ausgewählt ist aus:

- 1) C1-C10-Alkyl, gegebenenfalls substituiert mit Aryl,
- 2) Amino,
- 3) C1-C10-Alkylamino und
- 4) Aryl, gegebenenfalls substituiert mit C1-C10-Alkyl;

R_9 ausgewählt ist aus:

- 1) Wasserstoff,
- 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus R_{23} ,
- 3) Aryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 4) Cycloalkyl, gegebenenfalls substituiert mit einem Halogenatom(en) oder einem Hydroxylrest(en),
- 5) einem Heterocyclus, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkyl, C1-C10-Alkylcarbonyl, C1-C10-Alkoxy, C1-C10-Alkoxycarbonyl, Amino und einem Halogenatom,
- 6) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom und einem Hydroxylrest,
- 7) Heteroaryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus C1-C10-Alkyl, C1-C10-Alkylcarbonyl, C1-C10-Alkoxycarbonyl und einem Halogenatom, und
- 8) Cycloalkenyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus C1-C10-Alkoxy, C1-C10-Alkylamino, Amino, einem Hydroxylrest und einem Halogenatom, wobei das Cycloalkenyl gegebenenfalls einen Carbonylrest enthält;

R_{23} unabhängig ausgewählt ist aus:

- 1) einem Halogenatom,
- 2) einem Hydroxylrest,
- 3) einem C1-C10-Alkylcarbonyloxyrest,
- 4) $-COR_{16}$,

- 5) Amino,
 6) C1-C10-Alkylamino,
 7) einem Heterocyclus, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus C1-C10-Alkyl, C1-C10-Alkylcarbonyl, C1-C10-Alkoxycarbonyl und einem Halogenatom, und
 8) Cyano;

R₁₀ ausgewählt ist aus:

- 1) Wasserstoff und
 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom, einem Hydroxylrest und Aryl;

R₉ und R₁₀ gebunden sein können, um einen 4- bis 7-gliedrigen Heterocyclus zu bilden, der gegebenenfalls ein zusätzliches Element/zusätzliche Elemente oder einen zusätzlichen Rest/zusätzliche Reste, unabhängig ausgewählt aus N, O, S, SO, SO₂, Carbonyl und Thiocarbonyl, enthält, und der Heterocyclus gegebenenfalls mit einem Substituent(en), unabhängig ausgewählt aus R₂₄, substituiert ist;
 R₂₄ unabhängig ausgewählt ist aus:

- 1) einem Halogenatom,
 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkylamino und C1-C10-Alkylcarbonylamino,
 3) C1-C10-Halogenalkyl,
 4) einem Hydroxylrest,
 5) C1-C10-Hydroxyalkyl,
 6) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
 7) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
 8) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
 9) einem Heterocyclus, gegebenenfalls substituiert mit C1-C10-Alkyl,
 10) Heteroaryl,
 11) Heterocyclyl-C1-C 10-alkyl,
 12) -COR₁₆,
 13) -NR₁₉R₂₀ und
 14) -SO₂R₂₁;

R₁₁ ausgewählt ist aus:

- 1) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:
 i) einem Hydroxylrest,
 ii) -NR₁₇R₁₈,
 iii) einem C1-C10-Alkoxyrest,
 iv) einem Halogenatom,
 v) C1-C10-Alkoxycarbonyl,
 vi) Aminocarbonyl und
 vii) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, C1-C10-Alkoxy, Amino, C1-C10-Alkylamino und -COR₂₂,
 2) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, C1-C10-Alkoxy, Amino, C1-C10-Alkylamino und -COR₂₂,
 3) Cycloalkyl, gegebenenfalls substituiert mit einem Halogenatom(en),
 4) einem Heterocyclus, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus C1-C10-Alkyl, C1-C10-Alkylcarbonyl, C1-C10-Alkoxycarbonyl und einem Halogenatom,
 5) C1-C10-Alkoxy, wobei der Alkylrest gegebenenfalls mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkylcarbonylamino, Amino, C1-C10-Alkylamino und einem Hydroxylrest, substituiert ist,
 6) Amino,

- 7) C1-C10-Alkylamino, wobei der Alkylrest gegebenenfalls mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkylcarbonylamino, Amino, C1-C10-Alkylamino, Hydroxycarbonyl und einem Hydroxylrest, substituiert ist, und
8) C2-C10-Alkenyl,

R_{22} aus C1-C10-Alkoxy, einem Hydroxylrest, Amino und C1-C10-Alkylamino ausgewählt ist;
 R_{12} ausgewählt ist aus:

- 1) C1-C10-Alkyl,
2) Amino und
3) C1-C10-Alkylamino, wobei der Alkylrest gegebenenfalls mit einem Rest(en), unabhängig ausgewählt aus Amino, C1-C10-Alkylamino und einem Hydroxylrest, substituiert ist;

R_{13} ausgewählt ist aus:

- 1) Wasserstoff,
2) C1-C10-Alkyl,
3) C1-C10-Alkylcarbonyl, wobei das Alkyl gegebenenfalls mit einem Hydroxylrest(en) substituiert ist,
4) C1-C10-Alkoxycarbonyl,
5) Aminocarbonyl,
6) C1-C10-Alkylaminocarbonyl und
7) heterocyclischem Carbonyl, gegebenenfalls substituiert mit C1-C10-Alkyl;

R_{14} ausgewählt ist aus:

- 1) Wasserstoff und
2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino;

R_{13} und R_{14} gebunden sein können, um einen 4- bis 7-gliedrigen Heterocyclus zu bilden, der gegebenenfalls ein zusätzliches Element/zusätzliche Elemente oder einen zusätzlichen Rest/zusätzliche Reste, unabhängig ausgewählt aus O, N, S, SO und SO_2 , enthält, und der Heterocyclus gegebenenfalls Carbonyl enthält und der Heterocyclus gegebenenfalls mit C1-C10-Alkyl substituiert ist;

R_{15} ausgewählt ist aus:

- 1) C1-C10-Alkyl und
2) $-NR_{35}R_{36}$;

R_{35} ausgewählt ist aus:

- 1) Wasserstoff,
2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:
i) einem Halogenatom,
ii) einem Hydroxylrest,
iii) C1-C10-Alkylcarbonylamino,
iv) $-COR_{16}$,
v) Amino,
vi) C1-C10-Alkylamino,
vii) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en),
viii) Heteroaryl, gegebenenfalls substituiert mit einem C1-C10-Alkylrest(en), und
ix) einem Heterocyclus,

- 3) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
4) Cycloalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom und einem Hydroxylrest,
5) einem Heterocyclus, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-

- C10-Alkyl, einem Halogenatom und Aryl-C1-C10-alkyl,
 6) Heteroaryl, gegebenenfalls substituiert mit C1-C10-Alkyl, und
 7) C1-C10-Alkylcarbonyl;

R₃₆ ausgewählt ist aus:

- 1) Wasserstoff und
 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest und Aryl;

R₃₅ und R₃₆ aneinander gebunden sein können, um einen Ring, ausgewählt aus Azetidiny, Pyrrolidiny, Piperidiny, Piperaziny und Morpholiny, zu bilden, und der Ring gegebenenfalls mit einem Rest(en), unabhängig voneinander ausgewählt aus C1-C10-Alkyl und einem Halogenatom, substituiert ist;

R₂₅ ausgewählt ist aus:

- 1) einem Halogenatom,
 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:

- i) einem Halogenatom,
 ii) Aryl,
 iii) Heteroaryl,
 iv) einem Heterocyclus, gegebenenfalls substituiert mit einem C1-C10-Alkylrest(en),
 v) -COR₁₆,
 vi) -NR₁₃R₁₄ und
 vii) -SO₂R₂₁,

- 3) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Hydroxylrest(en),
 4) C1-C10-Hydroxyalkyl, wobei jeder Hydroxylrest unabhängig mit einem Rest(en), ausgewählt aus C1-C10-Alkyl, Aryl-C1-C10-alkyl und C1-C10-Alkylcarbonyl, substituiert sein kann,
 5) -COR₁₆,
 6) -SO₂R₂₁,
 7) Aryl und
 8) Cyano;

R₂ ausgewählt ist aus:

- 1) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkylrest gegebenenfalls weiter mit einem Substituent(en), unabhängig ausgewählt aus R₄₂, substituiert ist,
 2) C2-C10-Alkenyl, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkenylrest gegebenenfalls weiter mit einem Substituent(en), unabhängig ausgewählt aus R₄₂, substituiert ist,
 3) C2-C10-Alkiny, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkinyrest gegebenenfalls weiter mit einem Substituent(en), unabhängig ausgewählt aus R₄₂, substituiert ist,
 4) Cycloalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:

- i) einem Halogenatom,
 ii) C2-C10-Alkenyl oder C1-C10-Alkyl,
 iii) Aryl, gegebenenfalls substituiert mit 1 bis 3 Substituenten, unabhängig ausgewählt aus C1-C10-Alkyl, einem Halogenatom, C1-C10-Alkoxy, C1-C10-Alkylamino und C1-C10-Alkylcarbonyl,
 iv) Cycloalkyl,
 v) C2-C10-Alkenyl, gegebenenfalls substituiert mit Halogen,
 vi) C1-C10-Alkyliden, wobei das Alkyliden an das Cycloalkyl über eine Doppelbindung gebunden ist und das Alkyliden gegebenenfalls mit einem Halogenatom(en) substituiert ist,
 vii) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en),
 viii) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom oder C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en),
 ix) C2-C10-Alkiny und
 x) -Si(R₄₃)₃,

5) einem Heterocyclus, wobei der Heterocyclus gegebenenfalls mit einem Rest(en) substituiert ist, unabhängig ausgewählt aus:

- i) einem C1-C10-Alkylrest,
- ii) C1-C10-Alkylcarbonyl, wobei der Alkylrest gegebenenfalls mit R_{27} substituiert ist,
- iii) Arylcarbonyl, wobei der Arylrest gegebenenfalls mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, C1-C10-Alkyl und C1-C10-Alkoxy, substituiert ist,
- iv) Heteroarylcarbonyl,
- v) C1-C10-Alkoxycarbonyl, wobei der Alkylrest gegebenenfalls mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, Aryl und C1-C10-Alkoxy, substituiert ist,
- vi) Aryloxycarbonyl, wobei der Arylrest gegebenenfalls mit einem Halogenatom(en) und/oder C1-C10-Alkyl substituiert ist,
- vii) $-CONR_{28}R_{29}$,
- viii) $-SO_2R_{21}$,
- ix) einem Halogenatom,
- x) Cycloalkylcarbonyl, gegebenenfalls kondensiert mit einem Arylrest, und
- xi) C2-C10-Alkenylcarbonyl, wobei der Alkenylrest gegebenenfalls mit Aryl substituiert ist, wobei das Aryl gegebenenfalls mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, C1-C10-Alkyl oder C1-C10-Alkoxy, substituiert ist,

6) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus R_{44} ,

7) Heteroaryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:

- i) einem Halogenatom,
- ii) C1-C10-Alkyl und
- iii) C1-C10-Alkoxy;

8) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkoxyrest gegebenenfalls weiter mit einem Substituent(en), unabhängig ausgewählt aus R_{42} , substituiert ist,

9) $-S(O)_qR_{43}$ (wobei q eine ganze Zahl von 0 bis 2 ist) und

10) Cycloalkenyl, gegebenenfalls substituiert mit C1-C10-Alkyl;

R_{44} ausgewählt ist aus:

- 1) einem Halogenatom,
- 2) Cyano,
- 3) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:

- i) einem Hydroxylrest,
- ii) $-OR_{26}$,
- iii) Cyano,
- iv) Aryloxy, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, C1-C10-Alkyl, gegebenenfalls substituiert mit einem Halogenatom(en), oder C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en), und
- v) einem Halogenatom,

4) Cycloalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom oder C1-C10-Alkyl, gegebenenfalls substituiert mit einem Halogenatom(en),

5) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en) oder einem C2-C6-Alkenylrest(en),

6) $-COR_{30}$,

7) C1-C10-Alkylcarbonylamino,

8) C1-C10-Alkoxycarbonylamino, wobei der Alkoxyrest gegebenenfalls mit Aryl substituiert ist,

9) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Halogenatom(en),

10) Aryl, gegebenenfalls substituiert mit einem Substituent(en), unabhängig ausgewählt aus:

- i) einem Halogenatom,
- ii) C1-C10-Alkyl,

iii) C1-C10-Alkoxy und

iv) Aryl, gegebenenfalls substituiert mit Aryl, gegebenenfalls substituiert mit C1-C10-Alkyl,

11) Heteroaryl, gegebenenfalls substituiert mit einem C1-C10-Alkylrest(en),

12) $-\text{SO}_2\text{R}_{43}$,

13) $-\text{SOR}_{43}$,

14) C1-C10-Alkylthio, gegebenenfalls substituiert mit einem Halogenatom(en),

15) $-\text{Si}(\text{R}_{43})_3$ und

16) $-\text{SF}_5$;

R_{42} ausgewählt ist aus:

1) Wasserstoff,

2) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkyl, gegebenenfalls substituiert mit Halogen, einem Halogenatom und C1-C10-Alkoxy,

3) Hydroxycarbonyl,

4) C1-C10-Alkoxycarbonyl,

5) Aminocarbonyl,

6) C1-C10-Alkylaminocarbonyl,

7) C1-C10-Alkoxycarbonylamino,

8) Amino,

9) einem Hydroxylrest und

10) Oxetan, Tetrahydrofuran oder Tetrahydropyran, gegebenenfalls substituiert mit C1-C10-Alkyl;

R_{43} einen C1-C10-Alkylrest darstellt;

R_{26} Aryl oder C1-C10-Alkyl, gegebenenfalls substituiert mit einem Halogenatom(en), ist;

R_{27} ausgewählt ist aus:

1) Aryl, gegebenenfalls substituiert mit einem Rest, unabhängig ausgewählt aus einem Halogenatom, C1-C10-Alkyl und C1-C10-Alkoxy,

2) C1-C10-Alkoxy, wobei der Alkylrest gegebenenfalls mit Aryl substituiert ist,

3) einem Hydroxylrest,

4) Amino,

5) C1-C10-Alkylamino,

6) Hydroxycarbonyl,

7) Heteroaryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkyl und/oder Aryl, und

8) Heteroaryloxy;

R_{28} ausgewählt ist aus Wasserstoff oder C1-C10-Alkyl, gegebenenfalls substituiert mit Aryl;

R_{29} ausgewählt ist aus Wasserstoff oder C1-C10-Alkyl, gegebenenfalls substituiert mit Aryl;

R_{28} und R_{29} gebunden sein können, um einen Ring, ausgewählt aus Azetidiny, Pyrrolidiny, Piperidiny, Piperaziny und Morpholiny, zu bilden, und der Ring gegebenenfalls mit einem Rest(en), unabhängig voneinander ausgewählt aus C1-C10-Alkyl und einem Halogenatom, substituiert ist;

R_{30} ausgewählt ist aus einem Hydroxylrest, C1-C10-Alkoxy und $-\text{NR}_{31}\text{R}_{32}$;

R_{31} und R_{32} unabhängig ausgewählt sind aus:

1) Wasserstoff,

2) C1-C10-Alkyl, gegebenenfalls substituiert mit Aryl, und

3) Aryl,;


R_{31} und R_{32} gebunden sein können, um einen Ring, ausgewählt aus Azetidiny, Pyrrolidiny, Piperidiny, Piperaziny und Morpholiny, zu bilden, und der Ring gegebenenfalls mit einem Rest(en), unabhängig voneinander ausgewählt aus C1-C10-Alkyl, einem Halogenatom und C1-C10-Alkoxycarbonyl, substituiert ist; und

R_{33} und R_{34} unabhängig ausgewählt sind aus:


1) Wasserstoff und

2) C1-C10-Alkyl]; und

[wobei in der allgemeinen Formel (2) W, Y, m, n, R₄, R₅₃, R₆, R₇, R₁₁, R₁₆, R₁₉, R₂₀, R₂₁, R₃₃, R₃₄ und R₄₄ wie für die Formel (1) definiert sind;
U eine Bindung, C1-C10-Alkylen oder einen Rest, ausgewählt aus Resten, dargestellt durch die folgende Formel, bedeutet:



A ausgewählt ist aus O, NH und CH₂;
R₄₆ ausgewählt ist aus Wasserstoff oder R₄₄;
T ausgewählt ist aus Aryl und Heteroaryl;
V ausgewählt ist aus:




E ein 4- bis 7-gliedriger Heterocyclus ist, der gegebenenfalls ein zusätzliches Element/zusätzliche Elemente oder einen zusätzlichen Rest/zusätzliche Reste, ausgewählt aus O, N, S, SO und SO₂, enthält, und der Heterocyclus gegebenenfalls mit einem Rest(en) substituiert ist, ausgewählt aus:

- 1) Wasserstoff,
- 2) einem Halogenatom,
- 3) C1-C10-Alkyl, das gegebenenfalls einen Hydroxylrest(e) enthält, unabhängig ausgewählt aus C1-C10-Alkylamino, einem Halogenatom und einem Hydroxylrest,
- 4) einem Hydroxylrest,
- 5) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 6) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 7) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 8) einem Heterocyclus, gegebenenfalls substituiert mit C1-C10-Alkyl,
- 9) Heteroaryl, gegebenenfalls substituiert mit C1-C10-Alkyl,
- 10) Heterocycl-C1-C10-alkyl,
- 11) -COR₁₆,
- 12) -NR₁₉R₂₀ und
- 13) -SO₂R₂₁;

Z ein zweiwertiger Rest ist, ausgewählt aus:

- 1) C1-C10-Alkylen oder C1-C10-Heteroalkylen, gegebenenfalls substituiert mit einem Halogenatom(en) und/oder einem Hydroxylrest(en), wobei das Kohlenstoffatom/die Kohlenstoffatome oxidiert sein können, um Carbonyl zu bilden;

- 2) C2-C10-Alkenylen oder C2-C10-Heteroalkenylen, gegebenenfalls substituiert mit einem Halogenatom(en) und/oder einem Hydroxylrest(en), wobei das Kohlenstoffatom/die Kohlenstoffatome oxidiert sein können, um Carbonyl zu bilden;
 3) einem Rest, ausgewählt aus:



G ein zweiwertiger Rest ist, ausgewählt aus:

- 1) C1-C10-Alkylen oder C1-C10-Heteroalkylen, gegebenenfalls substituiert mit einem Halogenatom(en); und
 2) C2-C10-Alkenylen oder C2-C10-Heteroalkenylen, gegebenenfalls substituiert mit einem Halogenatom(en);

J ein zweiwertiger Rest ist, ausgewählt aus:

- 1) C1-C10-Alkylen oder C1-C10-Heteroalkylen, gegebenenfalls substituiert mit einem Halogenatom(en); und
 2) C2-C10-Alkenylen oder C2-C10-Heteroalkenylen, gegebenenfalls substituiert mit einem Halogenatom(en);

B ausgewählt ist aus einem Heterocyclus oder Heteroaryl; und

R₄₅ ausgewählt ist aus Wasserstoff oder C1-C10-Alkyl;

und wobei in der durch allgemeine Formel (1) oder (2) dargestellten Verbindung der Ausdruck "Heteroaryl" einen Rest bedeutet, der 1 bis 5 Heteroatome in dem Alkylgerüst enthält, der Ausdruck "Heteroalkylen" einen Rest bedeutet, der 1 bis 5 Heteroatome in dem Alkylengerüst enthält, und der Ausdruck "Heteroalkenylen" einen Rest bedeutet, der 1 bis 5 Heteroatome in dem Alkenylengerüst enthält.

2. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach Anspruch 1, wobei

W ausgewählt ist aus:

- 1) einer Einfachbindung,
 2) C1-C10-Alkylen, das gegebenenfalls einen Carbonylrest enthält, wobei das Alkylen gegebenenfalls mit einem Halogenatom(en) oder Hydroxy substituiert ist,
 3) C2-C10-Alkenylen, gegebenenfalls substituiert mit einem Halogenatom(en),
 4) C2-C10-Alkenylen,
 5) Arylen,
 6) Heteroarylen,
 7) -NH-, -NHCH₂- oder -NHCH₂CH₂-,
 8) Cycloalkylen und
 9) -(Cycloalkylen)-CH₂-;

X ausgewählt ist aus der folgenden Bindung oder den folgenden Resten:

- 1) einer Einfachbindung,
 2) C1-C10-Alkylen, gegebenenfalls substituiert mit Cycloalkyl,
 3) C2-C10-Alkenylen,
 4) C2-C10-Alkinylen und
 5) C1-C10-Oxyalkylen;

R₁ ausgewählt ist aus:

- 1) Wasserstoff,
- 2) Cycloalkyl, gegebenenfalls substituiert mit einem Rest, ausgewählt aus R_4 ,
- 3) einem Heterocyclus, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R_{25} und R_4 ,
- 4) Aryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R_3 , R_4 , R_5 und R_6 , und
- 5) Heteroaryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R_{25} , R_4 und R_5 ;

R_9 ausgewählt ist aus:

- 1) Wasserstoff,
- 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus R_{23} ,
- 3) Cycloalkyl, gegebenenfalls substituiert mit einem Halogenatom(en) oder einem Hydroxylrest(en),
- 4) einem Heterocyclus, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus C1-C10-Alkyl, C1-C10-Alkylcarbonyl, C1-C10-Alkoxy, C1-C10-Alkoxycarbonyl, Amino und einem Halogenatom,
- 5) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom und einem Hydroxylrest,
- 6) Heteroaryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkyl, C1-C10-Alkylcarbonyl, C1-C10-Alkoxycarbonyl und einem Halogenatom, und
- 7) Cycloalkenyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus C1-C10-Alkoxy, C1-C10-Alkylamino, Amino, 1 bis 3 Hydroxylresten und 1 bis 4 Halogenatomen, wobei das Cycloalkenyl gegebenenfalls einen Carbonylrest enthält;

R_{10} ausgewählt ist aus:

- 1) Wasserstoff und
- 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest und Aryl;

R_9 und R_{10} gebunden sein können, um einen 4- bis 7-gliedrigen Heterocyclus zu bilden, der gegebenenfalls ein zusätzliches Element/zusätzliche Elemente oder einen zusätzlichen Rest/zusätzliche Reste, unabhängig ausgewählt aus N, O, S, SO, SO_2 , Carbonyl und Thiocarbonyl, enthält, und der Heterocyclus gegebenenfalls mit einem Substituent(en), unabhängig ausgewählt aus R_{24} , substituiert ist;

R_{24} ausgewählt ist aus:

- 1) einem Halogenatom,
- 2) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkylamino und C1-C10-Alkylcarbonylamino,
- 3) C1-C10-Halogenalkyl,
- 4) einem Hydroxylrest,
- 5) C1-C10-Hydroxyalkyl,
- 6) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 7) Aryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 8) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit 1 bis 2 Resten, ausgewählt aus einem Halogenatom, einem Hydroxylrest, Amino und C1-C10-Alkylamino,
- 9) $-COR_{16}$ und
- 10) $NR_{19}R_{20}$;

R_{11} ausgewählt ist aus:

- 1) C1-C10-Alkyl, gegebenenfalls substituiert mit 1 bis 3 Substituenten, unabhängig ausgewählt aus:
 - i) einem Hydroxylrest,
 - ii) $-NR_{17}R_{18}$,
 - iii) einem C1-C10-Alkoxyrest,
 - iv) einem Halogenatom,
 - v) C1-C10-Alkoxycarbonyl und
 - vi) Aminocarbonyl,

- 2) Aryl,
- 3) Aryl-C1-C10-alkyl,
- 4) Cycloalkyl, gegebenenfalls substituiert mit einem Halogenatom(en),
- 5) einem Heterocyclus, gegebenenfalls substituiert mit C1-C10-Alkyl,
- 6) C1-C10-Alkoxy, wobei der Alkylrest gegebenenfalls mit einem Rest(en) substituiert ist, unabhängig ausgewählt aus C1-C10-Alkylcarbonylamino, Amino, C1-C10-Alkylamino und einem Hydroxylrest,
- 7) Amino,
- 8) C1-C10-Alkylamino, wobei der Alkylrest gegebenenfalls mit einem Rest(en) substituiert ist, unabhängig ausgewählt aus C1-C10-Alkylcarbonylamino, Amino, C1-C10-Alkylamino, Hydroxycarbonyl und einem Hydroxylrest, und
- 9) C2-C10-Alkenyl; und

R₃₃ und R₃₄ Wasserstoff sind,
wobei R₃, R₄, R₅, R₆, R₁₆, R₁₇, R₁₈, R₁₉, R₂₀, R₂₃ und R₂₅ jeweils wie in Anspruch 1 definiert sind.

3. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach Anspruch 1 oder 2, wobei

W ausgewählt ist aus:

- 1) einer Einfachbindung;
- 2) C1-C10-Alkylen, gegebenenfalls substituiert mit einem Halogenatom(en),
- 3) C2-C10-Alkenylen,
- 4) C2-C10-Alkinylen und
- 5) Heteroarylen,

X ausgewählt ist aus der folgenden Bindung oder den folgenden Resten:

- 1) einer Einfachbindung,
- 2) C1-C10-Alkylen,
- 3) C2-C10-Alkenylen,
- 4) C2-C10-Alkinylen und
- 5) C1-C10-Oxyalkylen, wobei das Oxyalkylen an einen 1,3,8-Triaza-spiro[4.5]dec-1-en-4-on-Ring oder einen 1,3,8-Triaza-spiro[4.5]dec-1-en-4-thion-Ring über ein Kohlenstoffatom in dem Oxyalkylen gebunden ist;

R₁ ausgewählt ist aus:

- 1) Aryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R₃, R₄ und R₅, und
- 2) Heteroaryl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus R₂₅ und R₄;

R₃ ausgewählt ist aus:

- 1) -CONR₇R₈,
- 2) -OR₉,
- 3) -NR₉R₁₀,
- 4) -N(R₉)COR₁₁,
- 5) -N(R₉)SO₂R₁₂,
- 6) -SO₂R₁₅,
- 7) C₁-C₁₀-Alkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus -COR₁₆ und -NR₁₃R₁₄, und
- 8) -N(R₉)CSNH₂;

R₄ ausgewählt ist aus:

- 1) Halogen,
- 2) Cyano,
- 3) Amino,
- 4) C1-C10-Alkyl,
- 5) C1-C10-Halogenalkyl,
- 6) C1-C10-Alkoxy,

- 7) C1-C10-Halogenalkylcarbonyl,
- 8) -COR₁₆ und
- 9) C1-C10-Heteroalkyl;

R₂ ausgewählt ist aus:

- 1) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkylrest gegebenenfalls weiter mit einem Rest, ausgewählt aus R₄₂, substituiert ist,
- 2) C2-C10-Alkenyl, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkenylrest gegebenenfalls weiter mit einem Rest, ausgewählt aus R₄₂, substituiert ist,
- 3) C2-C10-Alkinyl, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkinylrest gegebenenfalls weiter mit einem Rest, ausgewählt aus R₄₂, substituiert ist,
- 4) Cycloalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:

- i) einem Halogenatom,
- ii) C2-C10-Alkenyl oder C1-C10-Alkyl,
- iii) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkyl, einem Halogenatom und C1-C10-Alkoxy,
- iv) Cycloalkyl,
- v) C2-C10-Halogenalkenyl oder C1-C10-Halogenalkyl,
- vi) C1-C10-Alkyliden, wobei das Alkyliden an das Cycloalkyl über eine Doppelbindung gebunden ist und das Alkyliden gegebenenfalls mit einem Halogenatom(en) substituiert ist,
- vii) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en),
- viii) C1-C10-Alkyl, substituiert mit C1-C10-Alkoxy, wobei das Alkyl und/oder das Alkyl in dem Alkoxy gegebenenfalls mit einem Halogenatom(en) substituiert ist,
- ix) C2-C10-Alkinyl und
- x) -Si(R₄₃)₃,

5) einem Heterocyclus, wobei der Heterocyclus gegebenenfalls mit einem Rest(en) substituiert ist, ausgewählt aus:

- i) einem C1-C10-alkylrest,
- ii) C1-C10-Alkylcarbonyl, wobei der Alkylrest gegebenenfalls mit R₂₇ substituiert ist,
- iii) Arylcarbonyl, wobei der Arylrest gegebenenfalls mit einem Rest(en) substituiert ist, unabhängig ausgewählt aus einem Halogenatom, C1-C10-Alkyl und C1-C10-Alkoxy,
- iv) Heteroarylcarbonyl,
- v) C1-C10-Alkoxycarbonyl, wobei der Alkylrest gegebenenfalls mit einem Rest(en) substituiert ist, unabhängig ausgewählt aus einem Halogenatom, Aryl und C1-C10-Alkoxy,
- vi) Aryloxycarbonyl, wobei der Arylrest gegebenenfalls mit einem Halogenatom(en) und/oder C1-C10-Alkyl substituiert ist,
- vii) -CONR₂₈R₂₉ und
- viii) -SO₂R₂₁,

6) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus R₄₄,

7) Heteroaryl, gegebenenfalls substituiert mit einem der folgenden Reste:

- i) C1-C10-Alkyl,

8) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkoxyrest gegebenenfalls weiter mit einem Rest, ausgewählt aus R₄₂, substituiert ist,

9) -S(O)_qR₄₃ (wobei q eine ganze Zahl von 0 bis 2 ist) und

10) Cycloalkenyl, gegebenenfalls substituiert mit C1-C10-Alkyl,

R₄₄ ausgewählt ist aus:

- 1) einem Halogenatom,
- 2) Cyano,
- 3) C1-C10-Alkyl, gegebenenfalls substituiert mit einem der folgenden Reste:

- i) einem Hydroxylrest,
- ii) $-OR_{26}$,
- iii) Cyano und
- iv) Aryloxy, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom, C1-C10-Alkyl, C1-C10-Halogenalkyl oder C1-C10-Halogenalkoxy,

- 4) C1-C10-Halogenalkyl,
- 5) Cycloalkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom und C1-C10-Halogenalkyl,
- 6) C1-C10-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en) oder einem C2-C6-Alkenylrest(en),
- 7) $-COR_{30}$,
- 8) C1-C10-Heteroalkyl, gegebenenfalls substituiert mit einem Halogenatom(en),
- 9) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:

- i) C1-C10-Alkyl und
- ii) Aryl,

- 10) Heteroaryl, gegebenenfalls substituiert mit einem C1-C10-Alkylrest(en),
- 11) $-SO_2R_{43}$,
- 12) C1-C10-Alkylthio, gegebenenfalls substituiert mit einem Halogenatom(en),
- 13) $-Si(R_{43})_3$ und
- 14) $-SF_5$; und

R_{27} ausgewählt ist aus:

- 1) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, C1-C10-Alkyl und C1-C10-Alkoxy,
- 2) C1-C10-Alkoxy, wobei der Alkylrest gegebenenfalls mit Aryl substituiert ist,
- 3) Heteroaryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C10-Alkyl und Aryl, und
- 4) Heteroaryloxy,

wobei R_5 , R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{21} , R_{25} , R_{26} , R_{28} , R_{29} , R_{30} , R_{42} und R_{43} jeweils wie in Anspruch 1 oder 2, von dem Anspruch 3 abhängt, definiert sind.

4. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach einem der Ansprüche 1 bis 3, wobei

W ausgewählt ist aus:

- 1) C1-C6-Alkylen, gegebenenfalls substituiert mit einem Fluoratom(en),
- 2) C1-C6-Alkenylen und
- 3) Thiophen,

X ausgewählt ist aus der folgenden Bindung oder den folgenden Resten:

- 1) einer Einfachbindung,
- 2) C1-C6-Alkylen und
- 3) C1-C6-Oxyalkylen, gegebenenfalls substituiert mit einem Halogenatom(en), wobei das Oxyalkylen an einen 1,3,8-Triaza-spiro[4.5]dec-1-en-4-on-Ring oder einen 1,3,8-Triaza-spiro[4.5]dec-1-en-4-thion-Ring über ein Kohlenstoffatom in dem Oxyalkylen gebunden ist;

Y ein Sauerstoffatom bedeutet;
m 1 darstellt; und
n 1 darstellt.

5. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach einem der Ansprüche 1 bis 4, wobei W ausgewählt ist aus:

- 1) Ethylen,
- 2) Vinylen und
- 3) Thiophen,

X eine Einfachbindung bedeutet;
 R_3 ausgewählt ist aus:


- 1) $-\text{CONR}_7\text{R}_8$,
- 2) $-\text{OR}_9$,
- 3) $-\text{NR}_9\text{R}_{10}$,
- 4) $-\text{N}(\text{R}_9)\text{COR}_{11}$,
- 5) $-\text{N}(\text{R}_9)\text{SO}_2\text{R}_{12}$,
- 6) $-\text{SO}_2\text{R}_{15}$ und
- 7) C1-C6-Alkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus $-\text{COR}_{16}$ und $-\text{NR}_{13}\text{R}_{14}$;

R_2 ausgewählt ist aus:

- 1) C1-C10-Alkyl, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkylrest gegebenenfalls weiter mit einem Rest, ausgewählt aus R_{42} , substituiert ist,
- 2) C2-C10-Alkenyl, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkenylrest gegebenenfalls weiter mit einem Rest, ausgewählt aus R_{42} , substituiert ist,
- 3) C2-C10-Alkynyl, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkynylrest gegebenenfalls weiter mit einem Rest, ausgewählt aus R_{42} , substituiert ist,
- 4) Cycloalkyl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:

- i) einem Halogenatom,
- ii) C2-C6-Alkenyl oder C1-C6-Alkyl,

- 7) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus C1-C6-Alkyl, einem Halogenatom, C1-C6-Alkoxy, C1-C6-Alkylamino und C1-C6-Alkylcarbonyl,
- 8) Cycloalkyl,
- 9) C2-C6-Halogenalkenyl oder C1-C6-Halogenalkyl,
- 10) C1-C6-Alkyliden, wobei das Alkyliden an das Cycloalkyl über eine Doppelbindung gebunden ist und das Alkyliden gegebenenfalls mit einem Halogenatom(en) substituiert ist,
- 11) C1-C6-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en),
- 12) C1-C6-Alkyl, substituiert mit C1-C6-Alkoxy, wobei das Alkyl und/oder das Alkyl in dem Alkoxy gegebenenfalls mit Halogen substituiert ist,
- 13) C2-C6-Alkynyl und
- 14) $-\text{Si}(\text{R}_{43})_3$,
- 5) einem Rest, dargestellt durch die folgende allgemeine Formel (B):



(wobei R_a einen Rest bedeutet, ausgewählt aus:

- i) C1-C6-Alkylcarbonyl, wobei der Alkylrest gegebenenfalls mit R_{27} substituiert ist,
- ii) Arylcarbonyl, wobei der Arylrest gegebenenfalls mit einem Rest(en), unabhängig ausgewählt aus einem Halogenatom, C1-C6-Alkyl und C1-C6-Alkoxy, substituiert ist,
- iii) C1-C6-Alkoxy, wobei der Alkylrest gegebenenfalls mit einem Rest(en), ausgewählt aus einem Halogenatom, Aryl und C1-C6-Alkoxy, substituiert ist,
- iv) Aryloxy, wobei der Arylrest gegebenenfalls mit einem Halogenatom(en) oder C1-C6-Alkyl substituiert ist,

v) $-\text{CONR}_{28}\text{R}_{29}$ und

vi) $-\text{SO}_2\text{R}_{21}$),

6) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus R_{44} ,

7) Heteroaryl, gegebenenfalls substituiert mit einem der folgenden Reste:

i) einem Halogenatom,

ii) C1-C6-Alkyl und

iii) C1-C6-Alkoxy;

15) C1-C6-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en), wobei der Alkoxyrest gegebenenfalls weiter mit einem Rest, ausgewählt aus R_{42} , substituiert ist,

16) $-\text{S}(\text{O})_q\text{R}_{43}$ (wobei q eine ganze Zahl von 0 bis 2 ist) und

17) Cycloalkenyl, gegebenenfalls substituiert mit C1-C6-Alkyl;

und

R_{44} ausgewählt ist aus:

1) einem Halogenatom;

2) Cyano,

3) C1-C6-Alkyl, gegebenenfalls substituiert mit einem der folgenden Reste:

i) einem Hydroxylrest,

ii) $-\text{OR}_{26}$,

iii) Cyano und

iv) Aryloxy, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom, C1-C6-Alkyl, C1-C6-Halogenalkyl oder C1-C6-Halogenalkoxy,

4) C1-C6-Halogenalkyl,

5) Cycloalkyl, gegebenenfalls substituiert mit einem Rest(en), ausgewählt aus einem Halogenatom und C1-C6-Halogenalkyl,

6) C1-C6-Alkoxy, gegebenenfalls substituiert mit einem Halogenatom(en),

7) $-\text{COR}_{30}$,

8) C1-C6-Heteroalkyl, gegebenenfalls substituiert mit einem Halogenatom(en),

9) Aryl, gegebenenfalls substituiert mit einem Rest(en), unabhängig ausgewählt aus:

i) C1-C6-Alkyl und

ii) Aryl,

10) Heteroaryl, gegebenenfalls substituiert mit einem C1-C6-Alkylrest(en),

11) $-\text{SO}_2\text{R}_{43}$,


12) C1-C6-Alkylthio, gegebenenfalls substituiert mit einem Halogenatom(en),

13) $-\text{Si}(\text{R}_{43})_3$ und

14) SF_5 ,

wobei R_7 , R_8 , R_9 , R_{10} , R_{11} , R_{12} , R_{13} , R_{14} , R_{15} , R_{16} , R_{21} , R_{26} , R_{27} , R_{28} , R_{29} , R_{30} , R_{42} und R_{43} jeweils wie in den Ansprüchen 1 bis 4, von dem Anspruch 5 abhängt, definiert sind.


6. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach Anspruch 3, wobei R_1 ein durch die folgende allgemeine Formel (3) dargestellter Rest ist:



(3)

wobei R_3 , R_4 und R_5 wie für R_3 , R_4 und R_5 in Anspruch 3 definiert sind.


7. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach Anspruch 3, wobei R_1 ein durch die folgende allgemeine Formel (4) dargestellter Rest ist:



(4)

wobei R_4 und R_{25} wie für R_4 und R_{25} in Anspruch 3 definiert sind.


8. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach Anspruch 5, wobei R_1 ein durch die folgende allgemeine Formel (3) dargestellter Rest ist:



(3)

wobei R_3 , R_4 und R_5 wie für R_3 , R_4 und R_5 in Anspruch 5 definiert sind.

9. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach Anspruch 5, wobei R_1 ein durch die folgende allgemeine Formel (4) dargestellter Rest ist:



(4)

wobei R_4 und R_{25} wie für R_4 und R_{25} in Anspruch 5 definiert sind.

10. Eine Verbindung nach Anspruch 1, nämlich:

- (1) 8-(3-Chlor-benzolsulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (2) 8-(3-Chlor-benzolsulfonyl)-2-pyridin-4-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (3) 8-(3-Chlor-benzolsulfonyl)-2-propyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (4) 8-(3-Chlor-benzolsulfonyl)-2-isopropyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (5) 8-(3-Chlor-benzolsulfonyl)-2-(3-methoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (6) 8-(3-Chlor-benzolsulfonyl)-2-(3-chlor-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (7) 2-Benzyl-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (8) 8-(3-Chlor-benzolsulfonyl)-2-methyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (9) 2-Biphenyl-2-yl-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (10) 8-(3-Chlor-benzolsulfonyl)-2-o-tolyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (11) 8-(3-Chlor-benzolsulfonyl)-2-(4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (12) 8-(3-Chlor-benzolsulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (13) 8-(3-Chlor-benzolsulfonyl)-2-(1-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (14) 2-(1-Acetyl-piperidin-4-yl)-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (15) 2-tert-Butyl-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (16) 8-(3-Chlor-benzolsulfonyl)-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (17) 3-[8-(3-Chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäure-tert-butylester;
- (18) 8-(3-Chlor-benzolsulfonyl)-2-(3-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (19) 8-(3-Chlor-benzolsulfonyl)-2-(2-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (20) 8-(3-Chlor-benzolsulfonyl)-2-(4-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (21) 2-(1R,2S,4S)-Bicyclo[2.2.1]hept-2-yl-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (22) 2-(1R,2R,4S)-Bicyclo[2.2.1]hept-2-yl-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (23) 8-(3-Chlor-benzolsulfonyl)-2-(4-methoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (24) 8-(3-Chlor-benzolsulfonyl)-2-(4-methoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (25) 2-(4-tert-Butyl-cyclohexyl)-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (26) 8-(3-Chlor-benzolsulfonyl)-2-(4-fluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (27) 8-(3-Chlor-benzolsulfonyl)-2-cyclohexylmethyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (28) 8-(3-Chlor-benzolsulfonyl)-2-phenethyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (29) 8-(3-Chlor-benzolsulfonyl)-2-(2-cyclohexyl-ethyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (30) 8-(3-Chlor-benzolsulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (31) 8-(3-Chlor-benzolsulfonyl)-2-(3-methansulfonyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (32) 8-(3-Chlor-benzolsulfonyl)-2-(1-phenyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (33) 8-(3-Chlor-benzolsulfonyl)-2-(2-naphthalen-1-yl-ethyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (34) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (35) 2-tert-Butyl-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (36) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (37) 3-[8-(2-Naphthalen-1-yl-ethansulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäure-tert-butylester;
- (38) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-[1-(2-naphthalen-1-yl-ethansulfonyl)-piperidin-3-yl]-1,3,8-triaza-

spiro[4.5]dec-1-en-4-on;

(40) 8-(3-Chlor-benzolsulfonyl)-2-[1-(propan-1-sulfonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(41) 2-(1-Benzolsulfonyl-piperidin-3-yl)-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(42) 8-(3-Chlor-benzolsulfonyl)-2-(1-phenylmethansulfonyl-piperidin-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(43) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-(1-phenylmethansulfonyl-piperidin-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(44) 8-(4-Chlor-benzolsulfonyl)-2-(2,4-dichlor-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(45) 2-(2,4-Dichlor-phenyl)-8-(2-trifluormethyl-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(46) 8-(Butan-1-sulfonyl)-2-(2,4-dichlor-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(47) 2-(2,4-Dichlor-phenyl)-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(48) 2-(2,4-Dichlor-phenyl)-8-(chinolin-8-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(49) 8-(3-Chlor-4-fluor-benzolsulfonyl)-2-(2,4-dichlor-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(50) 8-(3-Chlor-benzolsulfonyl)-2-(2,4-dichlor-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(51) 2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-benzoesäuremethylester;

(52) 2-Cyclohexyl-8-(5-methyl-3-phenyl-isoxazol-4-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(53) 8-(Benzo[b]thiophen-3-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(54) 8-(Benzo[b]thiophen-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(55) 8-(5-Chlor-thiophen-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(56) 2-Cyclohexyl-8-(thiophen-2-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(57) 2-Cyclohexyl-8-(naphthalen-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(58) 2-Cyclohexyl-8-(2,4-dimethyl-thiazol-5-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(59) 2-Cyclohexyl-8-(3,5-dimethyl-isoxazol-4-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(60) 2-Cyclohexyl-8-(2,3-dihydro-benzo[1,4]dioxin-6-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(61) 2-Cyclohexyl-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(62) 3-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-thiophen-2-carbonsäuremethylester;

(63) 5-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-4-methyl-thiophen-2-carbonsäuremethylester;

(64) 2-Cyclohexyl-8-(2,5-dimethyl-thiophen-3-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(65) 8-(5-Brom-thiophen-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(66) 8-(5-Chlor-thiophen-2-sulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(67) 8-(Naphthalen-2-sulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(68) 8-(Benzo[b]thiophen-2-sulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(69) 8-(5-Brom-thiophen-2-sulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(70) 8-(3-Chlor-benzolsulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.6]undec-1-en-4-on;

(71) 7-(3-Chlor-benzolsulfonyl)-2-cyclohexyl-1,3,7-triaza-spiro[4.5]dec-1-en-4-on;

(72) 7-(3-Chlor-benzolsulfonyl)-2-cyclohexyl-1,3,7-triaza-spiro[4.4]non-1-en-4-on;

(73) 4-{2-[2-(2,4-Dichlor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(74) 4-[2-(2-tert-Butyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamid;

(75) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(76) 3,N,N-Trimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(77) 4-[2-(2-Cyclopentyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamid;

(78) 4-{2-[2-(2,6-Difluor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(79) 4-{2-[2-(2,6-Dimethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(80) 4-{2-[2-(3-Methoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(81) 4-{2-[2-(3-Chlor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(82) 4-{2-[2-(3,5-Bis-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(83) 4-[2-(2-Benzyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamid;

(84) 3,N,N-Trimethyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-benzamid;

(85) 4-{2-[2-(2-Chlor-5-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

- (86) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (87) 4-{2-[2-(4-Chlor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- 5 (88) 4-{2-[2-(2,3-Dichlor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (89) 4-{2-[2-(3-Chlor-4-fluor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (90) 4-{2-[2-(2-Chlor-4-fluor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- 10 (91) 4-{2-[2-(3-Brom-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (92) 4-{2-[2-(2,2-Difluor-benzo[1,3]dioxol-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- 15 (93) 4-{2-[2-(3-Chlor-2-fluor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (94) 3,N,N-Trimethyl-4-{2-[4-oxo-2-[4-(1,1,2,2-tetrafluor-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (95) 4-{2-[2-(4-Chlor-3-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- 20 (96) 4-{2-[2-(3-Fluor-4-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (97) 3,N,N-Trimethyl-4-{2-[2-(6-methyl-pyridin-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- 25 (98) 4-{2-[2-(3,4-Dichlor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (99) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (100) 4-{2-[2-(2,4-Bis-trifluorinethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- 30 (101) 3,N,N-Trimethyl-4-{2-[4-oxo-2-phenethyl-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (102) 4-{2-[2-(2,4-Dimethyl-thiazol-5-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (103) 4-{2-[2-(2-Cyclohexylmethyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamid};
- 35 (104) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (105) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(2-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- 40 (106) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(4-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (107) 4-{2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (108) 4-{2-[2-(2-Cyclohexyl-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- 45 (109) 4-{2-[2-(3-Fluor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (110) 4-{2-[2-(3-Methansulfonyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- 50 (111) 3,N,N-Trimethyl-4-{2-[2-(2-methyl-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (112) 3,N,N-Trimethyl-4-{2-[2-(2-methyl-5-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (113) 4-{2-[2-(2,3-Dimethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- 55 (114) 4-{2-[2-(3-Fluor-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (115) 4-{2-[2-(3-Fluor-5-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-

trimethyl-benzamid;

(116) 4-{2-[2-(2-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(117) 4-{2-[2-(4-Fluor-3-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(118) 4-{2-[2-(4-Difluoromethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(119) 4-{2-[2-(2-Methoxy-pyridin-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(120) 3,N,N-Trimethyl-4-{2-[2-(5-methyl-pyrazin-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(121) 3,N,N-Trimethyl-4-{2-(4-oxo-2-thiazol-4-yl-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl}-benzamid;

(122) 3,N,N-Trimethyl-4-{2-[2-(1-methyl-1H-imidazol-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(123) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(124) 4-{2-[2-(4-Chlor-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(125) 4-{2-[2-(4-Fluor-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(126) 4-{2-[2-(4-Chlor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(127) 4-{2-[2-(3-Chlor-5-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(128) 4-{2-[2-(4-Chlor-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(129) 4-{2-[2-(4-Chlor-2-fluor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(130) 4-{2-[2-(3-Isopropoxymethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(131) 4-{2-[2-(2,4-Dichlor-phenoxy-methyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(133) 4-{2-[2-(3-Chlor-2-methyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(134) 4-{2-[2-(2,4-Dichlor-benzyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(135) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(3-trifluormethyl-benzyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(136) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(1-trifluormethyl-cyclopropyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(137) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(1-trifluormethyl-cyclobutyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(138) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(1-trifluormethyl-cyclopentyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(139) 4-{2-[2-(2-Fluor-4-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(140) 3,N,N-Trimethyl-4-{2-[4-oxo-2-[3-(1,1,2,2-tetrafluor-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(141) 3,N,N-Trimethyl-4-{2-[4-oxo-2-((E)-propenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(142) 3,N,N-Trimethyl-4-{2-[4-oxo-2-((E)-styryl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;

(143) 4-[2-(2-Benzothiazol-6-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamid;

(144) 4-{2-[2-(4-Methoxy-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(145) N-{3-Methyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-acetamid;

(146) N-(4-{2-[2-(2,3-Dimethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

- (147) N-(4-{2-[2-(2,3-Dichlor-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;
- (148) N-(3-Methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (149) N-(2-Hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (150) Essigsäure-(S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-amino]-ethylester;
- (151) Essigsäure-(S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-{2-[4-oxo-2-(4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-amino]-ethylester;
- (152) Essigsäure-(S)-1-acetoxymethyl-2-[acetyl-(3-methyl-4-{2-[4-oxo-2-(4-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-amino]-ethylester;
- (153) Essigsäure-(S)-1-acetoxymethyl-2-[acetyl-(4-{2-[2-(4-fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-amino]-ethylester;
- (154) 8-{2-[4-((S)-2,3-Dihydroxy-propylamino)-2-methyl-phenyl]-ethansulfonyl}-2-(4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (155) 8-{2-[4-((S)-2,3-Dihydroxy-propylamino)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (156) 8-{2-[4-((S)-2,3-Dihydroxy-propylamino)-2-methyl-phenyl]-ethansulfonyl}-2-(4-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (157) 2-(3-Chlor-phenyl)-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (158) 2-(4-Chlor-phenyl)-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (159) 8-{2-[4-(2-Hydroxy-ethylamino)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (160) 8-{2-[4-(2-Hydroxy-ethylamino)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (161) 8-{2-[4-(2-Hydroxy-ethylamino)-2-methyl-phenyl]-ethansulfonyl}-2-(4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (162) {4-[2-(2-tert-Butyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-benzyl}-carbaminsäure-tert-butylester;
- (163) (4-{2-[2-(4,4-Difluor-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-benzyl)-carbaminsäure-tert-butylester;
- (164) (4-{2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-benzyl)-carbaminsäure-tert-butylester;
- (165) (3-Methyl-4-{2-[4-oxo-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzyl)-carbaminsäure-tert-butylester;
- (166) 8-[2-(3-Amino-phenyl)-ethansulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (167) N-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-acetamid;
- (168) 3,N,N-Trimethyl-4-(2-{4-oxo-2-[3-(2,2,2-trifluor-ethoxymethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzamid;
- (169) 4-(2-[2-[3-(2,2-Difluor-ethoxymethyl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3,N,N-trimethyl-benzamid;
- (170) 3,N,N-Trimethyl-4-(2-[4-oxo-2-[3-(2,2,3,3-tetrafluor-propoxymethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzamid;
- (171) 4-(2-[2-[3-(3,5-Dimethyl-isoxazol-4-yl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3,N,N-trimethyl-benzamid;
- (172) 4-[2-(2-Biphenyl-3-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,N,N-trimethyl-benzamid;
- (173) 3,N,N-Trimethyl-4-[2-[4-oxo-2-(3-pyridin-3-yl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzamid;
- (174) 3-[8-[2-(4-Dimethylcarbamoyl-2-methyl-phenyl)-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-benzoesäuremethylester;
- (175) 4-(2-[2-[1-(2,4-Dichlor-phenoxy)-1-methyl-ethyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3,N,N-trimethyl-benzamid;
- (176) 3,N,N-Trimethyl-4-[2-[4-oxo-2-(2,3,4-trifluor-phenoxy-methyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzamid;
- (177) 2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonsäure-(3-ethyl-phenyl)-amid;

- (178) 2-Cyclohexyl-8-[(E)-2-(1H-indol-5-yl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (179) 2-Cyclohexyl-8-[(E)-2-(2-trifluornethyl-phenyl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (180) 2-Cyclohexyl-8-[(E)-2-(3-methoxy-phenyl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (181) 2-Cyclohexyl-8-[(E)-2-(2-methoxy-phenyl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (182) 2-Cyclohexyl-8-[(E)-2-(1H-indol-4-yl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (183) 2-Cyclohexyl-8-[(E)-2-(2-fluor-6-trifluormethyl-phenyl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (184) 2-Cyclohexyl-8-[(E)-2-(2,3-difluor-phenyl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (185) 2-Cyclohexyl-8-[(E)-2-(3-fluor-2-trifluormethyl-phenyl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (186) 2-Cyclohexyl-8-[(E)-2-o-tolyl-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (187) 2-Cyclohexyl-8-[(E)-2-[4-(2-hydroxy-ethyl)-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (188) 2-Cyclohexyl-8-[(E)-2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (189) 4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-indol-1-carbonsäuredimethylamid;
 (190) N-{2-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-methyl-phenyl}-acetamid;
 (191) 2-Cyclohexyl-8-[(E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (192) 2-Cyclohexyl-8-[(E)-2-[1-(2-morpholin-4-yl-ethyl)-1H-indol-4-yl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (193) 2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-indol-1-yl}-N,N-dimethyl-acetamid;
 (194) 3-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-phenyl}-1,1-dimethyl-harnstoff;
 (195) Cyclopropan-carbonsäure-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-phenyl}-amid;
 (196) N-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-phenyl}-2-hydroxy-acetamid;
 (197) {3-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-4-methyl-phenyl}-carbaminsäuremethylester;
 (198) 1-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-phenyl}-3-(2-hydroxy-ethyl)-harnstoff;
 (199) 2-Cyclohexyl-8-[(E)-2-[5-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (200) 2-Cyclohexyl-8-[(E)-2-[2-(2-hydroxy-ethylamino)-6-methyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (201) 2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-phenyl}-amino-N,N-dimethyl-acetamid;
 (202) 2-Cyclohexyl-8-[(E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-3-fluor-1H-indol-4-yl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (203) 2-Cyclohexyl-8-[(E)-2-[2-methyl-4-(3,4,4,4-tetrafluor-pyrrolidin-1-carbonyl)-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (204) N-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-acetamid;
 (205) {4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-benzyl}-carbaminsäure-tert-butylester;
 (206) 2-Cyclohexyl-8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (207) 2-Cyclohexyl-8-[(E)-2-{2-methyl-4-[(2-oxo-oxazolidin-5-ylmethyl)-amino]-phenyl}-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (208) 2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl}-N-(2-dimethyl-amino-ethyl)-acetamid;
 (209) 3-[(E)-2-[2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-benzonitril;
 (210) 8-[(E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethensulfonyl]-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro [4.5]dec-1-en-4-on;
 (211) 8-[(E)-2-[4-(2-Hydroxy-ethylamino)-2-trifluormethyl-phenyl]-ethensulfonyl]-2-(3-trifluormethyl-phenyl)-

1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(212) 8-((E)-2-[2-Methyl-4-(pyrrolidin-1-carbonyl)-phenyl]-ethensulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(213) 3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonamid;

(214) N-(2-Hydroxy-ethyl)-3,N-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;

(215) N-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;

(216) 8-((E)-2-[2-Methyl-4-(4-methyl-piperazin-1-carbonyl)-phenyl]-ethensulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(217) 3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-N-(2,2,2-trifluor-ethyl)-benzolsulfonamid;

(218) 3-Methyl-N-(2-morpholin-4-yl-ethyl)-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonamid;

(219) 3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-N-pyridin-3-yl-benzolsulfonamid;

(220) 8-((E)-2-[4-((R)-4-Hydroxy-2-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(221) 8-((E)-2-[4-((R)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(222) 8-((E)-2-[4-((R)-5-Hydroxymethyl-2-oxo-oxazolidin-3-yl)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(223) 8-((E)-2-[4-(3-(3-Dimethylamino-propoxy)-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(224) N-(4-Hydroxy-butyl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonamid;

(225) N-Methyl-N-(4-methyl-3-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;

(226) 8-((E)-2-(3-Hydroxy-2-methyl-phenyl)-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(227) 8-((E)-2-(5-Hydroxy-2-methyl-phenyl)-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(228) N-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-(1-methyl-piperidin-4-yl)-acetamid;

(229) N-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-methansulfonamid;

(230) 8-((E)-2-[4-(4-Hydroxy-piperidin-1-carbonyl)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(231) 3,N,N-Trimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;

(232) N-(2-Hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;

(233) 3-Fluor-N,N-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;

(234) 8-((E)-2-[2,5-Dichlor-4-(3,4-dihydroxy-butoxy)-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(235) 8-((E)-2-[4-(3-Dimethylamino-propoxy)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(236) N-Methyl-N-(2-methyl-3-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;

(237) N-Methyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;

(238) N-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;

(239) N-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamid;

(240) 2-Cycloheptyl-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

ro[4.5]dec-1-en-4-on;

(241) 2-(4,4-Difluor-cyclohexyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(242) 8-((E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(243) 8-((E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(2-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(244) 2-(2-Fluor-3-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(245) 8-((E)-2-[1-((S)-2,2-Dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-4-yl]-ethensulfonyl)-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(246) 8-((E)-2-[1-(2,2-Dimethyl-[1,3]dioxolan-4-ylmethyl)-1H-indol-4-yl]-ethensulfonyl)-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(247) 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethensulfonyl)-2-(2-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(248) 2-(2-Fluor-3-trifluormethyl-phenyl)-8-((E)-2-[2-methyl-4-(pyrrolidin-1-carbonyl)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(249) 2-(2,4-Dichlor-phenyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(250) 2-(4-Chlor-phenyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(251) 2-Cyclohexyl-8-((E)-4-(1H-indol-4-yl)-but-3-en-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(252) 2-Cyclohexyl-8-((E)-5-(1H-indol-4-yl)-pent-4-en-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(253) 2-Cyclohexyl-8-((E)-3-[1-(2,3-dihydroxy-propyl)-1H-indol-4-yl]-prop-2-en-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(254) 8-((E)-2-[3-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(255) 8-((E)-2-[5-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(256) 8-((E)-2-[3-(2-Hydroxy-ethoxy)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(257) N-(2-Hydroxy-ethyl)-N-(4-methyl-3-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;

(258) 8-((E)-2-[4-((R)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(259) N-(4-((E)-2-[2-(2-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamid;

(260) N-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-piperidin-4-yl-acetamid;

(261) 2-Cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(262) 2-Cyclohexyl-8-[5-(1H-indol-4-yl)-pentan-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(263) 2-Cyclohexyl-8-[4-(1H-indol-4-yl)-butan-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(264) N-Methyl-N-(2-methyl-3-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(265) N-Methyl-N-(4-methyl-3-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(266) 8-{2-[3-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(267) 8-{2-[5-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(268) 8-{2-[4-((R)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(269) 2-Cyclohexyl-8-[2-(1H-indol-5-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(270) 2-[4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-phenyl]anilino-N,N-dimethyl-acetamid;

(271) Cyclopropancarbonsäure-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-phenyl}-amid;

- (272) {4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-benzyl}-carbaminsäure-tert-butylester;
- (273) 3-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-phenyl}-1,1-dimethyl-harnstoff;
- (274) {3-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-4-methyl-phenyl}-carbaminsäuremethylester;
- (275) 2-Cyclohexyl-8-{2-[5-(2-hydroxy-ethylamino)-2-methyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (276) N-{2-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamid;
- (277) 2-Cyclohexyl-8-(2-o-tolyl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (278) 1-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-phenyl}-3-(2-hydroxy-ethyl)-harnstoff;
- (279) 2-Cyclohexyl-8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (280) 2-Cyclohexyl-8-{2-[4-(2-hydroxy-ethyl)-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (281) 2-Cyclohexyl-8-(2-{2-methyl-4-[(2-oxo-oxazolidin-5-ylmethyl)-amino]-phenyl}-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (282) 2-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-N-(2-dimethylamino-ethyl)-acetamid;
- (283) 2-Cyclohexyl-8-{2-[2-methyl-4-(3,3,4,4-tetrafluor-pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (284) 8-{2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (285) 8-{2-[4-(4-Hydroxy-piperidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (286) 8-{2-[4-(2-Hydroxy-ethylamino)-2-trifluormethyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (287) N-(4-Hydroxy-butyl)-3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzolsulfonamid;
- (288) 8-{2-[2-Methyl-4-(pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (289) 8-{2-[2-Methyl-4-(4-methyl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (290) 3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-N-(2,2,2-trifluor-ethyl)-benzolsulfonamid;
- (291) 3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzolsulfonamid;
- (292) 3-Methyl-N-(2-morpholin-4-yl-ethyl)-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzolsulfonamid;
- (293) 3-Fluor-N,N-dimethyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (294) 3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-N-pyridin-3-yl-benzolsulfonamid;
- (295) 8-{2-[4-(3-Dimethylamino-propoxy)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (296) 8-{2-[4-((R)-5-Hydroxymethyl-2-oxo-oxazolidin-3-yl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (297) 8-{2-[4-((R)-4-Hydroxy-2-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (298) N-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (299) N-Methyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (300) 2-Hydroxy-N-methyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (301) 1-Methyl-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-harnstoff;

- (302) N-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-acetamid;
- (303) 2-Cyclohexyl-8-[2-(1H-indol-4-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (304) 2-(3-Trifluormethoxy-phenyl)-8-[2-(2-trifluormethyl-phenyl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (305) 8-[2-(1H-Indol-4-yl)-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (306) 8-[2-(1H-Indol-4-yl)-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (307) 8-[2-(2,2-Difluor-benzo[1,3]dioxol-4-yl)-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (308) 8-[2-(1-Methyl-1H-indol-7-yl)-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro [4.5] dec-1-en-4-on;
- (309) 2-Cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (310) 4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-indol-1-carbonsäuredimethylamid;
- (311) 8-[3-(1H-Indol-4-yl)-propan-1-sulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (312) 2-Cyclohexyl-8-[2-(1-methansulfonyl-1H-indol-4-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (313) 2-Cyclohexyl-8-{2-[1-(2-dimethylamino-ethyl)-1H-indol-4-yl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (314) 8-[2-(4-Amino-2-trifluormethyl-phenyl)-ethansulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (315) 2-Cyclohexyl-8-[3-(3-trifluormethyl-phenyl)-propan-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (316) 2-Cyclohexyl-8-(3-m-tolyl-propan-1-sulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (317) 2-Cyclohexyl-8-[3-(3-hydroxy-phenyl)-propan-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (318) 2-Cyclohexyl-8-[3-(2-hydroxy-phenyl)-propan-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (319) 8-(2-Benzo[b]thiophen-3-yl-ethansulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (320) 2-Cyclohexyl-8-(2-isochinolin-5-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (321) 2-Cyclohexyl-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-methyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (322) 8-[2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethansulfonyl]-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (323) 2-Cyclohexyl-8-[2-(2-trifluormethyl-phenyl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (324) 4-[3-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-propyl]-N,N-dimethyl-benzamid;
- (325) 8-{2-[4-((R)-2,3-Dihydroxy-propoxy)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (326) N-(2-Hydroxy-ethyl)-N-(4-methyl-3-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (327) N-(2-Hydroxy-ethyl)-N-(2-methyl-3-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl -phenyl)-acetamid;
- (328) 8-[3-(3-Amino-phenyl)-propan-1-sulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (329) N,N-Dimethyl-4-{5-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-thiophen-2-yl}-benzamid;
- (330) N,N-Dimethyl-3-{5-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-thiophen-2-yl}-benzamid;
- (331) 3,N,N-Trimethyl-4-{5-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-thiophen-2-yl}-benzamid;
- (332) N-{4-[5-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-thiophen-2-yl]-3-methyl-phenyl}-acetamid;
- (333) 4-{5-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-thiophen-2-yl}-3,N,N-trimethyl-benzamid;
- (334) 3-{5-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-thiophen-2-yl}-4,N,N-trimethyl-benzamid;
- (335) 2-Methyl-3'-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-biphenyl-4-carbonsäuredimethylamid;
- (336) 2-Methyl-4'-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-biphenyl-4-carbonsäuredimethylamid;
- (337) 3,N,N-Trimethyl-4-{6-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-pyridin-2-yl}-benzamid;
- (338) 2-Cyclohexyl-8-{2-[2-(3,5-dimethyl-isoxazol-4-yl)-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-

4-on;

(339) 2-Cyclohexyl-8-{2-[2-(6-methoxy-pyridin-3-yl)-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(340) 2-Cyclohexyl-8-{2-[2-(1-methyl-1H-pyrazol-4-yl)-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(341) 8-{5-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-thiophen-2-sulfonyl}-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(342) N-(4-{5-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-thiophen-2-yl}-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamid;

(343) 2-Cyclohexyl-8-((E)-2-thiazol-2-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(344) 2-Cyclohexyl-8-((E)-2-cyclopentyl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(345) 2-Cyclohexyl-8-(2-thiazol-2-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(346) 2-Cyclohexyl-8-[3-(4-methoxy-phenyl)-propan-1-sulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(347) N-Benzyl-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-benzamid;

(348) 8-(3-Chlor-benzolsulfonyl)-2-[3-(morpholin-4-carbonyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(349) 3-[8-(3-Chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-N-methyl-N-phenyl-benzamid;

(350) N-Benzyl-3-[8-(3-chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-benzamid;

(351) 3-[8-(3-Chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-N,N-dimethyl-benzamid;

(352) 8-[2-[4-(4-Methansulfonyl-piperazin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl]-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(353) N-[(R)-1-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzoyl)-pyrrolidin-3-yl]-acetamid;

(354) 8-{2-[2-Methyl-4-((S)-2-trifluormethyl-pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(355) 8-{2-[4-((R)-3-Hydroxy-pyrrolidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(356) 8-{2-[4-((S)-2-Hydroxymethyl-pyrrolidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(357) 8-(2-[4-[4-(2-Hydroxy-ethyl)-piperidin-1-carbonyl]-2-methyl-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(358) 8-(2-[2-Methyl-4-[4-(3-methyl-[1,2,4]oxadiazol-5-yl)-piperidin-1-carbonyl]-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(359) 8-{2-[2-Methyl-4-(4-pyrimidin-2-yl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(360) 4-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzoyl)-piperazin-1-sulfonsäure-dimethylamid;

(361) 8-{2-[2-Methyl-4-(4-pyridin-2-yl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(362) 8-[2-(4-[4-[2-(2-Hydroxy-ethoxy)-ethyl]-piperazin-1-carbonyl]-2-methyl-phenyl)-ethansulfonyl]-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(363) 8-(2-[2-Methyl-4-[4-(2-morpholin-4-yl-ethyl)-piperazin-1-carbonyl]-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(364) 8-{2-[2-Methyl-4-(4-thiazol-2-yl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(365) 8-{2-[4-(4,4-Difluor-piperidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(366) 8-(2-[4-[4-(3-Hydroxy-propyl)-piperazin-1-carbonyl]-2-methyl-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(367) (S)-1-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzoyl)-pyrrolidin-2-carbonsäureamid;

(368) 8-{2-[2-Methyl-4-(3,3,4,4-tetrafluor-pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(369) (R)-1-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzoyl)-pyrrolidin-2-carbonsäureamid;

(370) 8-{2-[4-((S)-3-Hydroxy-pyrrolidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(371) 8-{2-[4-((R)-2-Hydroxymethyl-pyrrolidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (372) 8-{2-[4-((S)-3-Dimethylamino-pyrrolidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (373) 8-{2-[4-(4-tert-Butyl-piperazin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (374) 8-(2-{4-[4-(3-Dimethylamino-propyl)-piperazin-1-carbonyl]-2-methyl-phenyl}-ethansulfonyl)-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (375) 8-(2-{4-[4-(4-Fluor-phenyl)-piperazin-1-carbonyl]-2-methyl-phenyl}-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (376) 8-{2-[4-(4-Isopropyl-piperazin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (377) 8-{2-[4-(3-Hydroxy-azetidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (378) 8-{2-[4-(3-Fluor-pyrrolidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (379) 8-{2-[4-(3-Fluor-azetidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (380) 8-{2-[2-Methyl-4-(piperidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (381) 8-{2-[4-(Azetidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (382) 4-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzoyl)-piperazin-1-carbonsäuredimethylamid;
- (383) 3,5,N,N-Tetramethyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (384) 8-((E)-2-[4-(4-Hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (385) 8-((E)-2-[4-(3-Hydroxy-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (386) 8-((E)-2-[4-(4-Hydroxy-4-hydroxymethyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (387) 8-((E)-2-[4-((3R,4R)-3-Dimethylamino-4-hydroxy-pyrrolidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (388) 8-((E)-2-[2,6-Dimethyl-4-(pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (389) N-(3-Hydroxy-propyl)-3,5,N-trimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (390) N-(2-Dimethylamino-ethyl)-3,5,N-trimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (391) N-(3-Dimethylamino-propyl)-3,5,N-trimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (392) N-Carbamoylmethyl-3,5,N-trimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (393) 3,5,N-Trimethyl-N-(1-methyl-piperidin-4-yl)-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (394) 8-((E)-2-[4-(4-Acetyl-piperazin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (395) 4-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzoyl)-piperazin-1-carbonsäuredimethylamid;
- (396) 4-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzoyl)-piperazin-1-carbonsäureamid;
- (397) 8-((E)-2-[4-[4-(3-Dimethylamino-propyl)-piperazin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (398) 8-((E)-2-[2,6-Dimethyl-4-((R)-3-methylamino-pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (399) 8-((E)-2-[4-((R)-3-Dimethylamino-pyrrolidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (400) 8-((E)-2-[4-(4-Hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (401) 8-[(E)-2-[4-[(R)-3-Hydroxy-pyrrolidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (402) 8-[(E)-2-[4-[(S)-3-Hydroxy-pyrrolidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (403) 8-[(E)-2-[4-(4-Hydroxy-4-hydroxymethyl-piperidin-1-carbonyl)-2,6-dimethylphenyl]-ethansulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (404) 8-[(E)-2-[4-[(R)-3-Amino-pyrrolidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (405) 8-[(E)-2-[4-(4-Dimethylamino-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (406) 8-[(E)-2-[4-[(3S,4S)-3-Hydroxy-4-isopropylamino-pyrrolidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (407) 8-[(E)-2-[4-(2-Dimethylamino-ethoxy)-piperidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (408) 8-[(E)-2-[4-(3-Hydroxy-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (409) 2,N,N-Trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (410) 2-Methyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (411) 8-{2-[2-Methyl-3-(morpholin-4-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (412) 4,N,N-Trimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (413) N-(2-Hydroxy-ethyl)-4,N-dimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (414) 8-{2-[2-Methyl-5-(4-methyl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (415) 4,N-Dimethyl-3-{2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (416) 8-{2-[4-[4-(2-Hydroxy-ethyl)-piperazin-1-carbonyl]-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (417) 8-{2-[2-Methyl-4-(thiazolidin-3-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (418) 2-Fluor-5,N,N-trimethyl-4-{2-[4-oxo-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamide;
- (419) 8-{2-[5-Fluor-4-(4-hydroxy-piperidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluoromethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (420) N-Benzyl-2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-1H-indol-3-yl}-acetamid;
- (421) 2-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-N-methyl-benzamid;
- (422) 3-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-N-pyridin-4-yl-benzamid;
- (423) 8-(3-Chlor-benzolsulfonyl)-2-[1-(3,3-dimethyl-butyryl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (424) 8-(3-Chlor-benzolsulfonyl)-2-[1-(2-hydroxy-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (425) 8-(3-Chlor-benzolsulfonyl)-2-[1-(4-chlor-benzoyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (426) 8-(3-Chlor-benzolsulfonyl)-2-[1-(3-methoxy-benzoyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (428) 2-[1-(1H-Indol-5-carbonyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (429) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-[1-(3-phenyl-propionyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (430) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-[1-(1,2,3,4-tetrahydro-naphthalen-2-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (431) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-[1-(chinolin-6-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (432) 2-[1-(2-3H-Imidazol-4-yl-acetyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (433) 2-{1-[2-(2,5-Dimethyl-thiazol-4-yl)-acetyl]-piperidin-3-yl}-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-tria-

za-spiro[4.5]dec-1-en-4-on;

(434) 2-{1-[2-(5-Methyl-2-phenyl-oxazol-4-yl)-acetyl]-piperidin-3-yl}-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(435) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-[1-(2-pyridin-2-yl-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(436) 2-[1-(4-Benzoyloxy-butyryl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(437) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-[1-(4-phenyl-butyryl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(438) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-[1-(2-pyridin-4-yl-acetyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(439) 2-[1-(2-tert-Butoxy-acetyl)-piperidin-3-yl]-8-(2-naphthalen-1-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(441) 8-(2-Naphthalen-1-yl-ethansulfonyl)-2-{1-[(E)-(3-phenyl-acryloyl)]-piperidin-3-yl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(442) 2-[1-(2-Amino-acetyl)-piperidin-3-yl]-8-(3-chlor-benzolsulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(444) 8-{2-[2-Methyl-4-(3-methylamino-pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(445) 8-{2-[4-(3-Amino-pyrrolidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-onhydrochlorid;

(446) 8-[(E)-2-[2,6-Dimethyl-4-(piperazin-1-carbonyl)-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-onhydrochlorid;

(447) 8-[(E)-2-[4-[4-(2-Hydroxy-acetyl)-piperazin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(448) 4-[4-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-benzoyl)-piperazin-1-yl]-4-oxo-buttersäuremethylester;

(449) 8-[(E)-2-[4-[4-(4-Dimethylamino-butyryl)-piperazin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(450) 2-Methoxy-N-methyl-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-acetamid;

(451) 2-Hydroxy-N-methyl-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-acetamid;

(452) [(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-benzoyl)-methyl-amino]-essigsäure;

(453) 8-(3-Chlor-benzolsulfonyl)-2-[1-(3,3-dimethyl-butyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(454) 3-[8-(3-Chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäure-tert-butylamid;

(455) 3-[8-(2-Naphthalen-1-yl-ethansulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäurephenethyl-amid;

(456) 8-(3-Chlor-benzolsulfonyl)-2-[1-(piperidin-1-carbonyl)-piperidin-3-yl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(457) 3-(2-Dimethylamino-ethyl)-1-methyl-1-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-harnstoff;

(458) 1-Methyl-1-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-harnstoff;

(459) 1-[3-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-2-methyl-phenyl]-3-methyl-harnstoff;

(460) 3-[8-(3-Chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäure-2-methoxyethylester;

(461) 3-[8-(3-Chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäurebenzylester;

(462) 3-[8-(3-Chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäureisobutylester;

(463) 3-[8-(3-Chlor-benzolsulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäuremethylester;

(464) 3-[8-(2-Naphthalen-1-yl-ethansulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbonsäurebenzylester;

(465) 3-[8-(2-Naphthalen-1-yl-ethansulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl]-piperidin-1-carbon-

säure-2,2-dimethyl-propylester;

(466) Methyl-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-carbaminsäure-2-dimethylamino-ethylester;

(467) Methyl-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-carbaminsäuremethylester;

(468) Methyl-(3-methyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-carbaminsäuremethylester;

(469) 2-Cyclohexyl-8-(2-o-tolyl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-thion;

(470) 1-(3,5-Dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-1-methyl-harnstoff;

(471) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;

(472) 1-(3,5-Dimethyl-4-[2-[4-oxo-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-1-methyl-harnstoff;

(474) 1-(3,5-Dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-1-methyl-thioharnstoff;

(475) 8-{2-[2,6-Dimethyl-4-(methyl-thiazol-2-yl-amino)-phenyl]-ethansulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(476) {4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-indol-1-yl}-essigsäuremethylester;

(477) 2-Cyclohexyl-8-[(E)-2-[1-(morpholin-4-carbonyl)-1H-indol-4-yl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(478) 2-Cyclohexyl-8-[(E)-2-[1-(4-hydroxy-butyl)-1H-indol-4-yl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(479) 2-Cyclohexyl-8-[(E)-2-(1H-indol-7-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(480) 2-Cyclohexyl-8-[(E)-2-[1-(3,4-dihydroxy-butyl)-1H-indol-4-yl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(481) N-(2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl}-ethyl)-acetamid;

(482) N-(1-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl}-ethyl)-acetamid;

(483) 2-Cyclohexyl-8-[(E)-2-(1-thiazol-2-ylmethyl-1H-indol-4-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(484) 3-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-indol-1-yl}-propan-1-sulfonsäureamid;

(485) 4-Methyl-piperazin-1-carbonsäure-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-phenyl}-amid;

(486) {4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-benzyl}-carbaminsäureisobutylester;

(487) 2-Cyclohexyl-8-[(E)-2-[1-[2-(2-hydroxy-ethoxy)-ethyl]-1H-indol-5-yl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(488) 2-Cyclohexyl-8-[(E)-2-(6-methyl-1H-indol-5-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(489) N-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-benzyl}-2-hydroxyacetamid;

(490) 3-(2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl}-ethyl)-1,1-dimethyl-harnstoff;

(491) 8-[(E)-2-(4-Amino-2,6-bis-trifluormethyl-phenyl)-ethansulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(492) 2-Cyclohexyl-8-[(E)-2-(6-trifluormethyl-1H-indol-5-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(493) {4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-phenylamino}-essigsäuremethylester;

(494) 2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-trifluormethyl-phenylamino}-N-(2-hydroxy-ethyl)-acetamid;

(495) 4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-N-(2-hydroxy-ethyl)-N-methyl-3-trifluormethyl-benzolsulfonamid;

(496) 4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-N,N-bis-(2-hydroxy-ethyl)-3-trifluormethyl-benzolsulfonamid;

- (497) 2-Cyclohexyl-8-((E)-2-[4-(2-dimethylamino-ethylamino)-2-methyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (498) N-(2-Acetylamino-ethyl)-2-[4-((E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-3-trifluormethyl-phenylamino]-acetamid;
- (499) 2-Cyclohexyl-8-((E)-2-(6-trifluormethyl-1H-benzimidazol-5-yl)-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (500) 2-Cyclohexyl-8-((E)-2-[1-((S)-2,2-dimethyl-[1,3]dioxolan-4-ylmethyl)-5-fluor-1 H-indol-4-yl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (501) 2-Cyclohexyl-8-((E)-2-[4-[(2-hydroxy-ethyl)-methyl-amino]-2-trifluormethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (502) 2-Cyclohexyl-8-((E)-2-[4-(1,1-dioxo-1 λ ⁶-thiomorpholine-4-carbonyl)-2-methylphenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (503) 4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-N-(2-dimethylamino-ethyl)-3-methyl-benzolsulfonamid;
- (504) 3-(3-Methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (505) 1-Methyl-3-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (506) N-(3-Isopropoxy-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (507) 3-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-5,5-dimethyl-imidazolidin-2,4-dion;
- (508) 3-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1,5,5-trimethyl-imidazolidin-2,4-dion;
- (509) N-(3-Ethoxy-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (510) N-(3-Ethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (511) 5,5-Dimethyl-3-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (512) N-(3-Methoxymethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (513) 2-(4-Methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(2-oxo-oxazolidin-3-yl)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (514) N-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-(2-methoxy-ethyl)-acetamid;
- (515) 8-((E)-2-[2,6-Dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethensulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (516) 8-((E)-2-[2,6-Dimethyl-4-(2-oxo-piperidin-1-yl)-phenyl]-ethensulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (517) N-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzyl)-N-methyl-acetamid;
- (518) 3-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzyl)-imidazolidin-2,4-dion;
- (519) 3-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzyl)-5,5-dimethyl-imidazolidin-2,4-dion;
- (520) 8-((E)-2-[2,6-Dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethensulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (521) 1-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzyl)-pyrrolidin-2,5-dion;
- (522) N-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzyl)-acetamid;
- (523) N-(3,5-Difluor-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamid;
- (524) 8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (525) [(S)-1-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-2,5-dioxo-pyrrolidin-3-yl]-carbaminsäure-tert-butylester;

- (526) 8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (527) 8-((E)-2-[2-Methyl-4-(4-methyl-piperazin-1-carbonyl)-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (528) 8-((E)-2-[2-Methyl-4-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethylamino]-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (529) 2-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenylamino)-acetamid;
- (530) 3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-N-pyridin-4-yl-benzolsulfonamid;
- (531) 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (532) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1,3-bis-(2-hydroxy-ethyl)-harnstoff;
- (533) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1,3-bis-(2-hydroxy-ethyl)-3-methyl-harnstoff;
- (534) N-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-(tetrahydro-pyran-4-yl)-methansulfonamid;
- (535) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (536) 3-(3-Methyl-4-((E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (537) 3-[4-((E)-2-(2-Nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-3-trifluormethyl-phenyl]-imidazolidin-2,4-dion;
- (538) 4-[3-Methyl-4-((E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-phenyl]-moipholin-3,5-dion;
- (539) 3-[3,5-Dimethyl-4-((E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-phenyl]-imidazolidin-2,4-dion;
- (540) 5,5-Dimethyl-3-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-imidazolidin-2,4-dion;
- (541) 8-((E)-2-[1-((R)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (542) N-(2-Methoxy-ethyl)-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-acetamid;
- (543) N-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-N-(2-methoxy-ethyl)-acetamid;
- (544) N-[2-(2-Hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-acetamid;
- (545) N-[2-(2-Fluor-ethoxy)-ethyl]-N-[3-methyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-acetamid;
- (546) N-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-N-methyl-acetamid;
- (547) 3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzoesäure;
- (548) 3-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-imidazolidin-2,4-dion;
- (549) 8-((E)-2-[4-[4-(2-Fluor-ethyl)-piperazin-1-carbonyl]-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (550) 8-((E)-2-[2,6-Dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethensulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (551) N-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-N-[2-(2-fluor-ethoxy)-ethyl]-acetamid;
- (552) 3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzoesäure-N'-acetyl-hydrazid;
- (553) 8-((E)-2-(4-Hydroxymethyl-2,6-dimethyl-phenyl)-ethensulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (554) 3-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzyl]-imidazolidin-2,4-dion;

- (555) 3-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl}-vinyl)-benzyl]-5,5-dimethyl-imidazolidin-2,4-dion;
- (556) 8-((E)-2-[2,6-Dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethensulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (557) 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl}-vinyl)-benzyl]-pyrrolidin-2,5-dion;
- (558) N-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl}-vinyl)-benzyl]-acetamid;
- (559) N-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl}-vinyl)-benzyl]-N-methyl-acetamid;
- (560) 3-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidin-2,4-dion;
- (561) N-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-[2-(2-methoxy-ethoxy)-ethyl]-acetamid;
- (562) N-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-[2-(2-fluor-ethoxy)-ethyl]-acetamid;
- (563) 3-(4-((E)-2-[2-(4-Isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidin-2,4-dion;
- (564) 3-(4-((E)-2-[2-(4-Isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidin-2,4-dion;
- (565) 3-(4-((E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-benzyl)-1,1-dimethyl-harnstoff;
- (566) 2-(4-((E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-phenyl)-ethyl)-carbaminsäuremethylester;
- (567) 4-Methyl-piperazin-1-carbonsäure-4-((E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-benzylamid;
- (568) N-(4-((E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-3-methyl-phenyl)-N',N'-dimethyl-sulfamid;
- (569) {4-((E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-3-methyl-benzyl}-(3-dimethylamino-propyl)-carbaminsäure-tert-butylester;
- (570) 2-(4-((E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-phenyl)-N-(2,2-dimethyl-propyl)-acetamid;
- (571) 2-(4-((E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-phenyl)-N-(2-diisopropylamino-ethyl)-acetamid;
- (572) 2-Cyclohexyl-8-((E)-2-[2-methyl-4-(2-oxo-azetidin-1-yl)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (573) 2-Cyclohexyl-8-((E)-2-[2,6-dimethyl-4-(thiazol-2-ylamino)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (574) N-(4-((E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl)-3,5-dimethyl-phenyl)-N-(3-methyl-oxetan-3-ylmethyl)-acetamid;
- (575) 3-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (576) N-(3-Ethoxymethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (577) 2-(4-Methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(4-oxo-oxazolidin-3-yl)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (578) 4-(3-Methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-3-oxo-piperazin-1-carbonsäure-tert-butylester;
- (579) 2-(4-Methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(4-methyl-2-oxo-oxazolidin-3-yl)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (580) 2-(4-Methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(3-oxo-isoxazolidin-2-yl)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (581) 3-(2,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (582) 3-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-5-methyl-imidazolidin-2,4-dion;
- (583) 3-(2,6-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;

- (584) N-(2,6-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamid;
- (585) 8-[(E)-2-(3-Methoxy-2-methyl-phenyl)-ethensulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (586) 8-((E)-2-[4-((1H-Imidazol-2-yl)-methyl-amino)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (587) 1-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-dihydro-pyrimidin-2,4-dion;
- (588) 2-(4-Methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(5-oxo-pyrazolidin-1-yl)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (589) 5-tert-Butoxymethyl-3-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (590) N,N-Dimethyl-2-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonylamino)-acetamid;
- (591) 3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-N-pyridin-3-ylmethyl-benzolsulfonamid;
- (592) N-(4-Hydroxy-cyclohexyl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonamid;
- (593) 8-[(E)-2-(4-Methansulfonyl-2-methyl-phenyl)-ethensulfonyl]-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (594) N-Acetyl-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonamid;
- (595) N-(2-Hydroxy-1,1-bis-hydroxymethyl-ethyl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonamid;
- (596) N-(1-Benzyl-piperidin-4-yl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonamid;
- (597) 8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-5-yl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (598) 8-((E)-2-[4-(4-Isopropyl-piperazin-1-carbonyl)-2-methyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (599) N-((R)-2,3-Dihydroxy-propyl)-3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzolsulfonamid;
- (600) 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-3,5-difluor-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (601) 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-3,5-dimethyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (602) 4-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenoxy)-piperidin-1-carbonsäure-tert-butylester;
- (603) N-Isopropyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (604) 4-[Acetyl-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-amino]-piperidin-1-carbonsäure-tert-butylester;
- (605) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-(2-hydroxy-ethyl)-harnstoff;
- (606) 8-((E)-2-[4-((R)-2-(Isopropylamino-methyl)-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (607) N-(2-Dimethylamino-ethyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (608) 2-Dimethylamino-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamid;
- (609) N-(2-Dimethylamino-ethyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-methansulfonamid;
- (610) N-[(R)-1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-5-oxo-pyrrolidin-2-ylmethyl]-acetamid;
- (611) 8-((E)-2-[4-((R)-2-Dimethylaminomethyl-5-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (612) N-Cyanomethyl-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;

- (613) N-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-(2,2,2-trifluor-ethyl)-acetamid;
- (614) 3-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1,5,5-trimethyl-imidazolidine-2,4-dion;
- (615) 3-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-5,5-dimethyl-imidazolidine-2,4-dion;
- (616) N-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-(tetrahydro-pyran-4-yl)-acetamid;
- (617) 8-((E)-2-[2-Methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (618) 3-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (619) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-((E)-2-[2-methyl-4-(pyrrolidin-1-carbonyl)-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (620) 2-(4-fluor-3-trifluormethyl-phenyl)-8-(E)-2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (621) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-((E)-2-[4-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (622) 8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (623) 8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-2-(8,8,9,9,9-pentafluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (624) 1-(4-((E)-2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3-methyl-phenyl)-imidazolidin-2,4-dion;
- (625) 1-(4-((E)-2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3-methyl-phenyl)-5-methyl-imidazolidin-2,4-dion;
- (626) 1-(4-((E)-2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3-methyl-phenyl)-5,5-dimethyl-imidazolidin-2,4-dion;
- (627) 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (628) 3,5-Dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-benzoesäure;
- (629) 3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzoesäuretrimethylhydrazid;
- (630) 8-((E)-2-[1-((R)-2,3-Dihydroxy-propyl)-4,6-dimethyl-1H-indol-5-yl]-ethensulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (631) 3-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidin-2,4-dion;
- (632) N-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-(3-methyl-oxetan-3-ylmethyl)-acetamid;
- (633) 2-Cyclohexyl-8-((E)-2-chinolin-8-yl-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (634) 2-Cyclohexyl-8-[(E)-2-(2-oxo-2,3-dihydro-1H-indol-4-yl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (635) 3-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-imidazolidin-2,4-dion;
- (636) 3-(4-((E)-2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-imidazolidin-2,4-dion;
- (637) 2-Cyclohexyl-8-[(E)-2-(2-methyl-1H-indol-4-yl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (638) 2-Cyclohexyl-8-[(E)-2-(1-methyl-1,2,3,4-tetrahydro-chinolin-5-yl)-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (639) 8-((E)-2-[4-(3,4-Dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (640) (4-((E)-2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-[1,1,1,1-²H₃]methyl-carbaminsäure-tert-butylester;
- (641) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (642) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (643) 8-((E)-2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethensulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-

1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(644) 8-[(E)-2-[2,6-Dimethyl-4-(2-oxa-7-aza-spiro[3.5]nonan-7-carbonyl)-phenyl]-ethansulfonyl]-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(645) 2-Cyclohexyl-8-[2-(1,2,3,4-tetrahydro-chinolin-5-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(646) 4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-1H-indol-3-carbonitril;

(647) 7-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-indol-1-carbonsäuredimethylamid;

(648) 2-Cyclohexyl-8-(2-chinolin-5-yl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(649) 2-Cyclohexyl-8-[2-[3-(2,2,2-trifluor-acetyl)-1H-indol-4-yl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(650) 2-Cyclohexyl-8-[2-(1-isopropyl-1H-indol-4-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(651) 2-Cyclohexyl-8-[2-[1-(4-hydroxy-butyl)-1H-indol-4-yl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(652) N-{3-Cyano-4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-acetamid;

(653) 8-(2-Isochinolin-5-yl-ethansulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(654) 8-(2-Chinolin-5-yl-ethansulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(655) N-(3-Methoxy-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-acetamid;

(656) 8-[2-[4-(3,3-Dimethyl-1,1,4-trioxo-1 λ ⁶-[1,2,5]thiadiazolidin-2-yl)-2,6-dimethylphenyl]-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(657) 8-[2-[2-Methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethansulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(658) {4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-benzyl}-carbaminsäureisobutylester;

(659) N-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-benzyl}-2-hydroxy-acetamid;

(660) 2-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-phenylamino}-N-(4-hydroxy-butyl)-acetamid;

(661) {4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-phenyl}-carbaminsäureisobutylester;

(662) N-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-phenyl}-benzamid;

(663) 2-Cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzothiazol-6-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(664) N-(2-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-benzolsulfonylamino}-ethyl)-acetamid;

(665) N-(2-Dimethylamino-ethyl)-2-(3-methyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenylamino)-acetamid;

(666) 8-[2-[4-((2R,6S)-2,6-Dimethyl-morpholin-4-carbonyl)-2-methyl-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(667) N-(2,2,3,3,4,4,4-Heptafluor-butyl)-3-methyl-4-[2-[4-oxo-2-(3-trifluormethylphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzolsulfonamid;

(668) 8-[2-(3-Methoxy-2-methyl-phenyl)-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(669) (2-Methoxy-3,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-methyl-carbaminsäure-tert-butylester;

(670) 8-[2-[4-((S)-2-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethansulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(671) 8-[2-(5,7-Dimethyl-2-oxo-2,3-dihydro-benzoxazol-6-yl)-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(672) (2-{4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-ethyl)-carbaminsäuremethylester;

(673) {4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-benzyl}-methyl-carbaminsäure-tert-butylester;

(674) 8-(2-{4-[3-(2-Hydroxy-ethyl)-2-oxo-imidazolidin-1-yl]-2-methyl-phenyl}-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(675) 2-(3-Methyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfo-

nyl]-ethyl)-benzolsulfonylamino)-acetamid;

(676) 8-{2-[4-(3,4-Dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(677) 3-(4-{2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-imidazolidin-2,4-dion;

(678) N-Cyclopentyl-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(679) 1-(2,3-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;

(680) 1-(2,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;

(681) {4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbaminsäure-tert-butylester;

(682) 1-(3-Methoxy-5-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;

(683) 1-(3-Chlor-5-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;

(684) 1-(2-Methoxy-3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;

(685) 5,7-Dimethyl-6-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-1H-chinazolin-2,4-dion;

(686) 8-[(E)-2-[1-(2-Amino-ethyl)-1H-indol-4-yl]-ethensulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(687) 8-[(E)-2-(4-Aminomethyl-phenyl)-ethensulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(688) (S)-2-Amino-3-hydroxy-N-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-propionamid;

(689) N-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-N-piperidin-4-yl-methansulfonamid;

(690) 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethensulfonyl]-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(691) 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethensulfonyl]-2-(8,8,9,9,9-pentafluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(692) 1-{3,5-Dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl}-1-methyl-harnstoff;

(693) 1-(2-Methoxy-3,5-dimethyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;

(694) 1-[3,5-Dimethyl-4-[(E)-2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl}-vinyl]-phenyl]-1-(2-fluor-ethyl)-harnstoff;

(695) 1-(3-Chlor-5-methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;

(696) 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;

(697) 8-[(E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethensulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(698) 8-[(E)-2-[4-2-((R)-2,3-Dihydroxy-propoxy)-ethoxy]-2,6-dimethyl-phenyl]-ethensulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(699) N-[3,5-Dimethyl-4-[(E)-2-4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl]-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamid;

(700) N-[2-(2-Hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl]-acetamid;

(701) N-[3,5-Dimethyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl]-N-(2-hydroxy-ethyl)-acetamid;

(702) 2-Cyclohexyl-8-[(E)-3-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(704) 8-[(E)-2-[1-2-((S)-2,3-Dihydroxy-propoxy)-ethyl]-1H-indol-4-yl]-ethensulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(705) 8-[(E)-2-[2,6-Dimethyl-4-((2S,3S)-2,3,4-trihydroxy-butoxy)-phenyl]-ethensulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (706) 2-[4-(3,3,3-Trifluor-propyl)-cyclohexyl]-8-((E)-2-{1-[2-((2S,3S)-2,3,4-trihydroxy-butoxy)-ethyl]-1H-indol-4-yl}-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (707) N-((S)-2,3-Dihydroxy-propyl)-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (708) {4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-methyl-benzyl}-(2-hydroxy-ethyl)-carbaminsäure-tert-butylester;
- (709) N-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-N-(4-hydroxy-cyclohexyl)-acetamid;
- (710) 2-Cyclohexyl-8-((E)-2-{4-[(R)-2-(2-hydroxy-ethoxymethyl)-5-oxo-pyrrolidin-1-yl]-2,6-dimethyl-phenyl}-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (711) N-(2-Hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl-phenyl)-isobutylamid;
- (712) 8-((E)-2-[4-((R)-5-Hydroxymethyl-3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (713) 8-((E)-2-[4-((R)-5-Hydroxymethyl-3-methyl-2-oxo-pyrrolidin-1-yl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (714) 3-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-(2-hydroxy-ethyl)-5,5-dimethyl-imidazolidin-2,4-dion;
- (715) N-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-(4-hydroxy-cyclohexyl)-methansulfonamid;
- (716) N-(4-((E)-2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamid;
- (717) N-(2-Fluor-5-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-(2-hydroxy-ethyl)-acetamid;
- (718) N-(4-((E)-2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamid;
- (719) N-(2-Hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-methansulfonamid;
- (720) N-(2-Hydroxy-ethyl)-N-(3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl)-methansulfonamid;
- (721) N-(4-{2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-N-(2-hydroxy-ethyl)-acetamid;
- (722) 2-Hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (723) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (724) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-[2-(2-hydroxy-ethoxy)-ethyl]-harnstoff;
- (725) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-(2-methoxy-ethyl)-harnstoff;
- (726) 1-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-isopropyl-harnstoff;
- (727) 1-(3-Ethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-methyl-harnstoff;
- (728) 1-(3-Methoxy-5-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-methyl-harnstoff;
- (729) 1-Cyanomethyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-harnstoff;
- (730) 1-Cyclopentyl-1-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-harnstoff;
- (731) 1-(3-Methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-(2,2,2-trifluor-ethyl)-harnstoff;
- (732) 1-(4-[2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (733) 1-((S)-2,3-Dihydroxy-propyl)-1-(3,5-dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-harnstoff;
- (734) N-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3-methyl-phenyl}-N-cyclopentyl-acetamid;

- (735) (S)-2-Amino-N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl}-3-methyl-butylamid;
- (736) 2-{4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-indol-1-yl}-N-pyridin-4-yl-acetamid;
- (737) 2-(3-Methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-acetamid;
- (738) 2-Cyclohexyl-8-[(E)-2-[4-(4,5-dihydro-thiazol-2-ylamino)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (739) 2-Cyclohexyl-8-[(E)-2-[2-methyl-4-(3-methyl-oxetan-3-ylmethoxy)-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (740) 1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-cyclopropyl}-phenyl)-1-methyl-harnstoff;
- (741) 1-(3,5-Dimethyl-4-[(E)-1-methyl-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (742) N-(3-Methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triazaspiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-sulfamid;
- (743) N-(3-Hydroxy-propyl)-N'-(3-methyl-4-[(E)-2-[4-oxo-2-(3-trifluormethoxyphenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-sulfamid;
- (744) N-Methyl-N-(3-methyl-4-[(E)-3-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro [4.5]dec-1-en-8-sulfonyl]-propenyl]-phenyl)-acetamid;
- (745) 2-Cyclohexyl-8-[2-(2-methyl-1H-indol-4-yl)-ethansulfonyl]-1,3,8-triazaspiro[4.5]dec-1-en-4-on;
- (746) 8-{2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (747) 2-Cyclohexyl-8-[2-(1-methyl-1,2,3,4-tetrahydro-chinolin-5-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (748) 2-Cyclohexyl-8-[2-[1-(3,4-dihydroxy-butyl)-1H-indol-4-yl]-ethansulfonyl]-1,3,8-triaza-spiro [4.5]dec-1-en-4-on;
- (749) 3-(2,6-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-imidazolidin-2,4-dion;
- (750) 2-(3-Methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (751) N-(3-Ethoxy-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (752) N-(3-Ethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (753) N-(3-Ethoxymethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (754) N-(3-Methoxymethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (755) N-(2-Methoxy-ethyl)-N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (756) N-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-N-(2-methoxy-ethyl)-acetamid;
- (757) N-[2-(2-Fluor-ethoxy)-ethyl]-N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluorpropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-acetamid;
- (758) N-[2-(2-Hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluorpropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-acetamid;
- (759) N-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-N-methyl-acetamid;
- (760) 3-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-imidazolidin-2,4-dion;
- (761) 8-{2-[4-[4-(2-Fluor-ethyl)-piperazin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl}-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (764) 8-{2-[2,6-Dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethansulfonyl}-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (765) 8-{2-[2,6-Dimethyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethansulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (766) N-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sul-

fonyl)-ethyl)-phenyl]-N-[2-(2-fluor-ethoxy)-ethyl]-acetamid;
 (767) 3-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-5-methyl-imidazolidin-2,4-dion;
 (768) 3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl)-benzoesäure;
 (769) 3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl)-benzoesäure N'-acetyl-hydrazid;
 (770) 8-[2-(4-Hydroxymethyl-2,6-dimethyl-phenyl)-ethansulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5] dec-1-en-4-on;
 (771) 8-[2-[2,6-Dimethyl-4-(2-oxo-piperidin-1-yl)-phenyl]-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (772) N-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl)-ethyl)-benzyl]-acetamid;
 (773) N-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl)-acetamid;
 (774) 3-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl)-benzyl]-imidazolidin-2,4-dion;
 (775) 3-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl)-benzyl]-5,5-dimethyl-imidazolidin-2,4-dion;
 (776) 8-[2-[2,6-Dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethansulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (777) 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl)-benzyl]-pyrrolidin-2,5-dion;
 (778) 3-(2,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-imidazolidin-2,4-dion;
 (779) 3-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl)-imidazolidin-2,4-dion;
 (780) 3-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl)-5,5-dimethyl-imidazolidin-2,4-dion;
 (781) 8-[2-[2,6-Dimethyl-4-(2-oxo-pyrrolidin-1-ylmethyl)-phenyl]-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (782) 1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl)-pyrrolidin-2,5-dion;
 (783) N-(2,6-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-N-methyl-acetamid;
 (784) N-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl)-benzyl]-N-methyl-acetamid;
 (785) N-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl)-N-methyl-acetamid;
 (786) 1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-dihydro-pyrimidin-2,4-dion;
 (787) 3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl)-ethyl)-benzoesäuretrimethylhydrazid;
 (788) 8-[2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethansulfonyl]-2-(8,8,9,9,9-pentafluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (789) 8-[2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethansulfonyl]-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (790) 3-{3-Methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-imidazolidin-2,4-dion;
 (791) 8-[2-[1-((R)-2,3-Dihydroxy-propyl)-4,6-dimethyl-1H-indol-5-yl]-ethansulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (792) 3-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-imidazolidin-2,4-dion;
 (793) 8-[2-[2-Methyl-4-(3-oxo-morpholin-4-yl)-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
 (794) N-(4-[2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-N-(2-hydroxy-ethyl)-acetamid;
 (795) 3-{4-[2-(2-Nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-trifluormethyl-phenyl}-imida-

zolidin-2,4-dion;

(796) 3,5-Dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-benzoesäure;

(797) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-[2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(798) 8-[2-[4-(3,4-Dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(799) 3-(3,5-Dimethyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-imidazolidin-2,4-dion;

(800) 3-(4-[2-[2-(4-Fluor-3-trifluornethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-3,5-dimethyl-phenyl)-imidazolidin-2,4-dion;

(801) 8-[2-[4-(3,4-Dihydroxy-butoxy)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(802) 8-[2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(803) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-[2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(804) 8-[2-[2,6-Dimethyl-4-(2-oxa-7-aza-spiro[3.5]nonan-7-carbonyl)-phenyl]-ethansulfonyl]-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(812) N-(3-Isopropoxy-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-acetamid;

(817) 1-[3,5-Dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl]-1-methyl-harnstoff;

(818) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-(2-fluor-ethyl)-harnstoff;

(820) N-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamid;

(821) 8-[2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(822) 8-(2-[1-[2-((S)-2,3-Dihydroxy-propoxy)-ethyl]-1H-indol-4-yl]-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(823) 2-[4-(3,3,3-Trifluor-propyl)-cyclohexyl]-8-(2-[1-[2-((2S,3S)-2,3,4-trihydroxy-butoxy)-ethyl]-1H-indol-4-yl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(824) N-((S)-2,3-Dihydroxy-propyl)-N-(3,5-dimethyl-4-[2-[4-oxo-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-acetamid;

(825) N-[2-(2-Hydroxy-ethoxy)-ethyl]-N-[3-methyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl]-acetamid;

(826) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-[2-(2-hydroxy-ethoxy)-ethyl]-harnstoff;

(827) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl]-1-(2-methoxy-ethyl)-harnstoff;

(828) 1-(3,5-Dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-1-isopropyl-harnstoff;

(829) 1-(3-Ethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-1-methyl-harnstoff;

(830) 1-((S)-2,3-Dihydroxy-propyl)-1-(3,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-harnstoff;

(831) 8-[2-[4-(4-Hydroxy-4-hydroxymethyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(832) 3-[3,5-Dimethyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl]-imidazolidin-2,4-dion;

(833) 5-tert-Butoxymethyl-3-(3,5-dimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-imidazolidin-2,4-dion;

(834) 2-(4-Methyl-cyclohexyl)-8-[2-[2-methyl-4-(5-oxo-pyrazolidin-1-yl)-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(835) 2-Cyclohexyl-8-[2-(1H-indol-7-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(836) 2-Cyclohexyl-8-[2-(6-trifluormethyl-1H-indol-5-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(837) 2-Cyclohexyl-8-[2-(6-methyl-1H-indol-5-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(838) 2-cyclohexyl-8-(2-[1-[2-(2-hydroxy-ethoxy)-ethyl]-1H-indol-5-yl]-ethansulfonyl)-1,3,8-triaza-spi-

ro[4.5]dec-1-en-4-on;

(839) 2-Cyclohexyl-8-[2-(6-trifluormethyl-1H-benzimidazol-5-yl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(840) 8-[2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethansulfonyl]-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(841) 2-Cyclohexyl-8-[2-[4-(1,1-dioxo-1λ⁶-thiomorpholin-4-carbonyl)-2-methylphenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(842) 2-Cyclohexyl-8-[2-[4-(2-dimethylamino-ethylamino)-2-methyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(843) 8-[2-[2-Methyl-4-(4-methyl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(844) 4-[2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-N-(2-dimethylamino-ethyl)-3-methyl-benzolsulfonamid;

(845) 8-[2-[2-Methyl-4-[2-(4-methyl-piperazin-1-yl)-2-oxo-ethylamino]-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(846) 3-Methyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-N-pyridin-4-yl-benzolsulfonamid;

(847) 2-(3-Methyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenylamino)-acetamid;

(848) N-(1-Benzyl-piperidin-4-yl)-3-methyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzolsulfonamid;

(849) 8-[2-[4-(4-Isopropyl-piperazin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(850) 8-[2-[1-((S)-2,3-Dihydroxy-propyl)-1H-indol-5-yl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(851) N-((R)-2,3-Dihydroxy-propyl)-3-methyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzolsulfonamid;

(852) 3-Methyl-4-[2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-N-pyridin-3-ylmethyl-benzolsulfonamid;

(853) N-(4-Hydroxy-cyclohexyl)-3-methyl-4-[2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzolsulfonamid;

(854) N-(2-Hydroxy-1,1-bis-hydroxymethyl-ethyl)-3-methyl-4-[2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzolsulfonamid;

(855) 8-[2-[4-(3,4-Dihydroxy-butoxy)-3,5-dimethyl-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(856) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-[2-[2-methyl-4-(pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(857) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-[2-[4-((R)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(858) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-[2-[4-((S)-2-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-2-methyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(859) 4-(3-Methyl-4-[2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenoxy)-piperidin-1-carbonsäure-tert-butylester;

(860) 8-[2-[4-(3,4-Dihydroxy-butoxy)-3,5-difluor-phenyl]-ethansulfonyl]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(861) 2-Cyclohexyl-8-[2-[2-methyl-4-(2-oxo-azetidin-1-yl)-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(862) 1-Methyl-3-(3-methyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-imidazolidin-2,4-dion;

(863) 5,5-Dimethyl-3-(3-methyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-phenyl)-imidazolidin-2,4-dion;

(864) 5,5-Dimethyl-3-[3-methyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-imidazolidin-2,4-dion;

(865) 2-(4-Methyl-cyclohexyl)-8-[2-[2-methyl-4-(2-oxo-oxazolidin-3-yl)-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(866) 2-(4-Methyl-cyclohexyl)-8-[2-[2-methyl-4-(4-oxo-oxazolidin-3-yl)-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(867) 8-[2-[1-((R)-2,3-Dihydroxy-propyl)-1H-indol-4-yl]-ethansulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-

1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(868) 8-(2-{4-[(1H-Imidazol-2-yl)-methyl-amino]-2,6-dimethyl-phenyl}-ethansulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(869) 3-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-5,5-dimethyl-imidazolidin-2,4-dion;

(870) 2-(4-Methyl-cyclohexyl)-8-{2-[2-methyl-4-(2-oxo-piperazin-1-yl)-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(872) 2-Cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(873) N-(2-Fluor-5-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-N-(2-hydroxy-ethyl)-acetamid;

(874) N-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-sulfamid;

(875) N-(3-Hydroxy-propyl)-N'-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-sulfamid;

(876) 1-Cyanomethyl-1-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-harnstoff;

(877) 1-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-(2,2,2-trifluor-ethyl)-harnstoff;

(878) 1-(4-{2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidin-2,4-dion;

(879) 1-(4-{2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-5-methyl-imidazolidin-2,4-dion;

(880) 3-(4-{2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidin-2,4-dion;

(881) 3-(3-Methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-imidazolidin-2,4-dion;

(882) 3-(4-{2-[2-(4-Isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidin-2,4-dion;

(883) 3-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-imidazolidin-2,4-dion;

(884) 3-(4-{2-[2-(4-Isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-imidazolidin-2,4-dion;

(885) 3-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1,5,5-trimethyl-imidazolidin-2,4-dion;

(886) (S)-2-Amino-3-hydroxy-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-propionamid;

(887) 1-(4-{2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-5,5-dimethyl-imidazolidin-2,4-dion;

(888) 3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzoesäure-N,N'-dimethyl-hydrazid;

(889) 8-{2-[2-Methyl-4-(piperidin-4-yloxy)-phenyl]-ethansulfonyl}-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on-hydrochlorid;

(890) 4-{2-[2-(4,4-Difluor-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(891) 4-{2-[2-(4-Methansulfonyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(892) 3,N,N-Trimethyl-4-(2-{4-oxo-2-[(E)-2-(3-trifluormethyl-phenyl)-vinyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzamid;

(893) 3,N,N-Trimethyl-4-[2-(4-oxo-2-phenylethynyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzamid;

(894) 4-{2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(895) 4-{2-[2-(4-tert-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(896) 4-{2-[2-(4-Ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(897) 3,N,N-Trimethyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluor-butyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfo-

nyl)-ethyl)-benzamid;

(898) 4-{2-[4-(Cyclohexylmethyl-amino)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-dien-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(899) 4-{2-[4-Dimethylamino-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]deca-1,3-diene-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(900) 4-{2-[4-[(Z)-Hydroxyimino]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(901) 4-{2-[4-[(Z)-Methoxyimino]-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(902) 4-{2-[2-[4-(2-Methoxy-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(903) N-{4-[2-(2-Cyclopentyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamid;

(904) N-(3-Methyl-4-{2-[4-oxo-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(905) N-(4-{2-[2-(4-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(906) N-(4-{2-[2-(4-Cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(907) N-(4-{2-[2-(3-Cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(908) N-(4-{2-[2-(2-Cyano-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(909) N-(4-{2-[2-(4-tert-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(910) N-(4-{2-[2-(4-Ethoxymethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(911) N-(3-Methyl-4-{2-[4-oxo-2-(4-propoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(912) N-(4-{2-[2-(4-Butoxy-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(913) N-(4-{2-[2-(4-Isopropoxymethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(914) N-[4-(2-[2-[4-(3-Fluor-propoxy)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamid;

(915) N-[4-(2-[2-[4-(3-Fluor-propoxy)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamid;

(916) N-[4-(2-[2-[4-((E)-3,3-Difluor-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamid;

(917) N-{4-[2-(2-Cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl}-acetamid;

(918) N-(4-[2-(2-Adamantan-1-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl)-acetamid;

(919) N-[3-Methyl-4-(2-[4-oxo-2-[4-(2,2,2-trifluor-ethoxymethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-acetamid;

(920) N-[4-(2-[2-[4-(4-Chlor-phenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamid;

(921) N-[3-Methyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propoxy)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-acetamid;

(922) N-[3-Methyl-4-(2-[4-oxo-2-[4-(2,2,2-trifluor-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-acetamid;

(923) N-(3-Methyl-4-{2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(924) N-(3-Methyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(925) N-(4-{2-[2-(4,4-Dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(926) N-(3-Methyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfo-

nyl]-ethyl)-phenyl)-acetamid;

(927) N-[4-(2-[2-[4-(2-Methoxy-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;

(928) 3-Methyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl]-carbaminsäure-tert-butylester;

(929) N-(2-Hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-isobutylamid;

(930) 2-Hydroxy-N-(2-hydroxy-ethyl)-N-(3-methyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(931) N-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-2,2,2-trifluor-N-methyl-acetamid;

(932) N-(3,5-Dimethyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-2,2,2-trifluor-N-methyl-acetamid;

(933) N-(3,5-Dimethyl-4-{2-[4-oxo-2-(3,3,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-2,2,2-trifluor-N-methyl-acetamid;

(934) 1-{4-[2-(2-Cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-harnstoff;

(935) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;

(936) 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluor-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;

(937) 1-[2-Chlor-3,5-dimethyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluor-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;

(938) 1-(4-{2-[2-(4,4-Dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(939) 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyliden)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;

(940) 1-[4-(2-{2-[4-(3,3-Difluor-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3,5-dimethyl-phenyl]-1-methyl-harnstoff;

(941) 1-(3,5-Dimethyl-4-{2-[8-oxo-6-(3-trifluormethoxy-phenyl)-2,5,7-triaza-spiro[3.4]oct-5-en-2-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;

(942) N-(4-((E)-1-Fluor-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3-methyl-phenyl)-N-methyl-acetamid;

(943) 3-[3-Methyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-imidazolidin-2,4-dion;

(944) 3-[3-Methyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluor-butyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-imidazolidin-2,4-dion;

(945) 3-[4-(2-{2-[4-((E)-3,3-Difluor-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidin-2,4-dion;

(946) 3-[4-(2-{2-[4-(3,3-Difluor-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidin-2,4-dion;

(947) 3-(3-Methyl-4-{2-[4-oxo-2-(4-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-imidazolidin-2,4-dion;

(948) 3-[3-Methyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluor-ethyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-imidazolidin-2,4-dion;

(949) 3-(3-Methyl-4-{2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-imidazolidin-2,4-dion;

(950) 3-(3-Methyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-imidazolidin-2,4-dion;

(951) 3-(4-{2-[2-(4,4-Dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-imidazolidin-2,4-dion;

(952) 3-(3-Methyl-4-{2-[4-oxo-2-(3,3,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-imidazolidin-2,4-dion;

(953) N-(2-Chlor-4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-5-methyl-phenyl)-acetamid;

(954) N-(2-Chlor-4-{2-[2-(4-ethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;

(955) N-[4-(2-[2-[4-(3,3-Difluor-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-

methyl-phenyl]-acetamid;

(956) 3-[4-(2-[2-[4-(3,3-Difluor-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3-methyl-phenyl]-imidazolidin-2,4-dion;

(957) 2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-N-phenyl-benzamid;

(958) 8-(2-[2-Methyl-4-[4-(1-methyl-piperidin-4-yl)-piperazin-1-carbonyl]-phenyl]-ethansulfonyl)-2-(3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(959) 8-((E)-2-[4-((R)-3-Fluor-pyrrolidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(960) 8-((E)-2-[4-(4-Hydroxymethyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(961) 3,5,N,N-Tetramethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;

(962) 3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;

(963) 8-((E)-2-[4-(3-Fluor-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(964) 8-{2-[4-(4-Hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(965) 8-{2-[4-(3-Hydroxy-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(966) 8-{2-[4-(4-Hydroxymethyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-(3-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(967) 8-((E)-2-[2,6-Dimethyl-4-(2-oxa-6-aza-spiro[3.3]heptane-6-carbonyl)-phenyl]-ethansulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(968) 8-((E)-2-[2,6-Dimethyl-4-(3-oxo-piperazin-1-carbonyl)-phenyl]-ethansulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(969) 8-((E)-2-[4-((3R,5S)-3,5-Dimethyl-piperazin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(970) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-((E)-2-[4-(3-hydroxy-3-methyl-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(971) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-hydroxymethyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(972) 8-((E)-2-[2,6-Dimethyl-4-(4-oxo-piperidin-1-carbonyl)-phenyl]-ethansulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(973) 2-[4-(3,3-Difluor-allyl)-cyclohexyl]-8-{2-[4-(4-fluor-4-hydroxymethyl-piperidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(974) 2-[4-(3,3-Difluor-allyl)-cyclohexyl]-8-{2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(975) 2-(4-Ethyl-cyclohexyl)-8-(2-[4-[4-(2-fluor-ethyl)-piperazin-1-carbonyl]-2-methyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(976) 2-(4-Ethyl-cyclohexyl)-8-{2-[2-methyl-4-(4-oxetan-3-yl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(977) 8-{2-[2-Methyl-4-(pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(978) 8-{2-[4-(Azetidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(979) 8-{2-[4-((R)-3-Hydroxy-pyrrolidin-1-carbonyl)-2-methyl-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(980) 8-{2-[2,6-Dimethyl-4-(pyrrolidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(981) 8-{2-[4-((R)-3-Hydroxy-pyrrolidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(982) 8-{2-[4-(Azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(983) 8-{2-[2,6-Dimethyl-4-(piperidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(984) 8-{2-[4-(4-Hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (985) 8-{2-[4-(3-Hydroxy-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (986) 8-{2-[2,6-Dimethyl-4-(4-methyl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (987) N,N-Dimethyl-2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (988) N-Methoxy-2-(3-methyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (989) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (990) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethylphenyl]-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (991) 8-((E)-2-[2,6-Dimethyl-4-(4-methyl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (992) 8-((E)-2-[4-(Isoxazolidin-2-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (993) 8-((E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-piperidin-1-carbonyl]-2,6-dimethyl-phenyl)-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (994) 3,5,N,N-Tetramethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (995) 8-{2-[2,6-Dimethyl-4-(4-methyl-piperazin-1-carbonyl)-phenyl]-ethansulfonyl}-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (996) 3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzoesäurehydrazid;
- (997) 8-((E)-2-[2,6-Dimethyl-4-(pyrazolidin-1-carbonyl)-phenyl]-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (998) N-Methoxy-3,5,N-trimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (999) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1000) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1001) 8-((E)-2-[4-(Isoxazolidin-2-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1002) 8-((E)-2-[4-(1,1-Dioxo-1 λ ⁶-thiomorpholin-4-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1003) 3,5,N,N-Tetramethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl)-benzamid;
- (1006) 2-Cyclohexyl-8-((E)-2-[4-(4-hydroxy-4-trifluormethyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1007) 2-Cyclohexyl-8-((E)-2-[2,6-dimethyl-4-(2-oxo-oxazolidin-3-carbonyl)-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1008) 2-(4-Butyl-cyclohexyl)-8-((E)-2-[4-[4-(2-hydroxy-ethoxy)-piperidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1009) 2-(4-Butyl-cyclohexyl)-8-((E)-2-[4-[4-[2-(2-hydroxy-ethoxy)-ethoxy]-piperidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1010) 8-((E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-piperidin-1-carbonyl]-2,6-dimethyl-phenyl)-ethansulfonyl)-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1011) 2-Amino-N-(3,5-dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-N-methyl-acetamid;
- (1012) (3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-methyl-carbaminsäure 2-hydroxy-ethylester;
- (1013) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1,3,3-trimethyl-harnstoff;
- (1014) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1,3-dimethyl-harnstoff;
- (1015) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-methyl-harnstoff;

- (1016) 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethansulfonyl]-2-(3-trifluormethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1017) 1-(4-[(E)-2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1018) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[3-(1,1,2,2-tetrafluor-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1019) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[3-(4,4,4-trifluor-butoxy)-phenyl]-1,3,8-triaza-spiro [4.5] dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1020) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(4,4,4-trifluor-butoxy)-phenyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1021) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(4-pentyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1022) 1-(3,5-Dimethyl-4-[(E)-2-[2-(7-methylsulfanyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1023) 1-[3,5-Dimethyl-4-((E)-2-[2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1024) 1-(3,5-Dimethyl-4-[(E)-2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl)-1-methyl-harnstoff;
- (1025) 1-(3,5-Dimethyl-4-[(E)-2-(2-non-3-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl)-1-methyl-harnstoff;
- (1026) 1-[3,5-Dimethyl-4-((E)-2-[2-[10-(3-methyl-oxetan-3-yl)-decyl]-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1027) 1-(3,5-Dimethyl-4-[(E)-2-(2-non-1-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-phenyl)-1-methyl-harnstoff;
- (1028) 1-(4-{2-[2-(4-Fluor-3-trifluormethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1029) 1-(4-{2-[2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1030) 1-(4-{2-[2-(4-tert-Butyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1031) 1-[3,5-Dimethyl-4-(2-[2-[8-(3-methyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- (1032) 1-[3,5-Dimethyl-4-(2-[2-[10-(3-methyl-oxetan-3-yl)-decyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- (1033) 2-[8-(2-[4-(tert-Butoxycarbonyl-methyl-amino)-2,6-dimethyl-phenyl]-ethansulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-pyrrolidin-1-carbonsäure-tert-butylester;
- (1034) 4-{2-[2-(1,1-Difluor-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-methyl-carbaminsäure-tert-butylester;
- (1035) 8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethansulfonyl]-2-(8,8,9,9,9-pentafluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1036) 8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethansulfonyl]-2-(3-trifluormethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1037) 8-[2-(2,6-Dimethyl-4-methylamino-phenyl)-ethansulfonyl]-2-(4-isopropyliden-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1038) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[3-(1,1,2,2-tetrafluor-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- (1039) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(5,6,7,8-tetrahydro-naphthalen-2-yl)-1,3,8-triaza-spiro 4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-1-methyl-harnstoff;
- (1040) 1-(4-{2-[2-((1S,3S,5R)-3,5-Dimethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1041) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(1-propyl-butyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1042) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(2,2,3,3,3-pentafluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- (1043) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(6,6,7,7,7-pentafluor-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1044) N-(3,5-Dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluor-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-N-methyl-acetamid;

- (1045) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(7,7,7-trifluor-heptyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1046) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- 5 (1047) N-(3,5-Dimethyl-4-{2-[4-oxo-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-N-methyl-acetamid;
- (1048) 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[1-(4,4,4-trifluor-butyl)-cyclopropyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1050) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3-trifluormethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- 10 (1051) 1-(4-{2-[2-(cis-3,4-Dimethyl-cyclopentyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1052) 1-[4-[2-(2-Dicyclopropylmethyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl]-1-methyl-harnstoff;
- 15 (1053) 1-[4-[2-(2-Bicyclo[3.3.1]non-9-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl]-1-methyl-harnstoff;
- (1054) 1-[4-[2-((1R,SS)-2-Bicyclo[3.2.1]oct-3-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl]-1-methyl-harnstoff;
- (1055) 1-(3,5-Dimethyl-4-{2-[2-(3-methyl-cyclopentyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- 20 (1056) 1-[4-[2-(2-Bicyclo[2.2.1]hept-7-yl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl]-1-methyl-harnstoff;
- (1057) 1-(4-{2-[2-((1S,2S)-2-Hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- 25 (1058) 1-(3,5-Dimethyl-4-[2-(4-oxo-2-spiro[2.5]oct-6-yl-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl)-1-methyl-harnstoff;
- (1059) 1-(4-{2-[2-(4-Difluormethylen-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1060) N-[4-(2-[2-[4-(2,2-Difluor-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- 30 (1061) N-[4-(2-[2-[4-(2-Fluor-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- (1062) N-(4-[2-[2-(4-Butyl-4-fluor-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-3-methyl-phenyl)-acetamid;
- 35 (1063) N-[4-(2-[2-[4-((E)-3-Fluor-propenyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- (1064) N-[4-(2-[2-[4-(3-Fluor-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- (1065) N-[4-(2-[2-[4-(3-Chlor-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- 40 (1066) N-[4-(2-[2-[4-(3-Fluor-propylidene)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- (1067) N-[4-(2-[2-[4-(3,3-Difluor-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- 45 (1068) N-[4-(2-[2-[4-(3,3-Difluor-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- (1069) N-[3-Methyl-4-(2-[4-oxo-2-[4-(2,2,3,3,3-pentafluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl]-acetamid;
- (1070) N-[4-(2-[2-[4-((E)-3-Fluor-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- 50 (1071) N-[4-(2-[2-[4-(2,2-Difluor-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- (1072) N-[4-(2-[2-[4-(2-Fluor-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- 55 (1073) N-[4-(2-[2-[4-Fluor-4-(3-fluor-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3-methyl-phenyl]-acetamid;
- (1074) N-(4-[2-[2-(4-Ethynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-3-methyl-phenyl)-acetamid;

- (1075) N-(4-{2-[2-(4-Difluormethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;
- (1076) N-[4-(2-{2-[4-(3,3-Difluor-propyliden)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;
- (1077) N-[4-(2-{2-[4-(2-Fluor-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;
- (1078) N-[4-(2-{2-[4-(1-Fluor-1-methyl-ethyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;
- (1079) N-(4-{2-[2-(4-Butyliden-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;
- (1080) N-[3-Methyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-acetamid;
- (1081) 1-{3,5-Dimethyl-4-[2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-1-methyl-harnstoff;
- (1082) 1-{3,5-Dimethyl-4-[2-(2-non-3-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-1-methyl-harnstoff;
- (1083) 1-{3,5-Dimethyl-4-[2-(2-non-1-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-1-methyl-harnstoff;
- (1084) 1-(3,5-Dimethyl-4-{1-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-cyclopropylmethyl}-phenyl)-1-methyl-harnstoff;
- (1085) 1-(4-{2,2-Difluor-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1086) 1-{4-[2,2-Difluor-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-1-methyl-harnstoff;
- (1087) 1-{3,5-Dimethyl-4-[2-(4-oxo-2-[1,1';2',1'']terphenyl-3-yl-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl}-1-methyl-harnstoff;
- (1088) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(4,4,4-trifluor-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1089) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[3-(4,4,4-trifluor-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1090) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-propyl}-phenyl)-1-methyl-harnstoff;
- (1091) 2-[4-Fluor-3-(4,4,5,5,5-pentafluor-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1092) 2-[4-Fluor-3-(4,4,5,5,5-pentafluor-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1093) 2-(4-Chlor-3-trifluormethoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1094) 2-(3-Fluor-4-trifluormethoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1095) 2-(3,4-Bis-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1096) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[3-(4,4,5,5,5-pentafluor-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1097) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[3-(2,2,2-trifluor-ethyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1098) 2-(4-Fluor-3-methyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1099) 2-(4-Fluor-3-trifluormethoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1100) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[3-(1,1,2,2-tetrafluor-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1101) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1102) 2-(4-Fluor-3-trifluormethoxy-phenyl)-8-((E)-2-[4-(3-hydroxy-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1103) 8-((E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[3-(4,4,5,5,5-pentafluor-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (1104) 6-(4-Methyl-cyclohexyl)-2-(2-naphthalen-1-yl-ethansulfonyl)-2,5,7-triaza-spiro[3.4]oct-5-en-8-on;
- (1106) 11-{8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethensulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl}-undecyl)-carbaminsäure-tert-butylester;
- (1107) 8-[(E)-2-(2,6-Dimethyl-4-methylamino-phenyl)-ethensulfonyl]-2-(9-hydroxy-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1108) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1109) 14-(8-[(E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethensulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecansäure;
- (1110) 1-(3,5-Dimethyl-4-[(E)-2-[2-(4-[1,1,1-²H₃]methyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1111) 1-(4-[(E)-2-[2-(4-Fluor-3-trifluormethoxy-phenyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1112) 1-(3,5-Dimethyl-4-[(E)-2-[2-(3-nonafluorbutyl-phenyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1113) 1-(4-[(E)-2-[2-(4-Chlor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1114) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1115) 1-(4-[(E)-2-[2-(6-Ethoxy-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1116) 1-[3,5-Dimethyl-4-[(E)-2-[4-oxo-2-[3-(6,6,6-trifluor-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl]-1-methyl-harnstoff;
- (1117) 1-[3,5-Dimethyl-4-[(E)-2-[4-oxo-2-[4-(6,6,6-trifluor-hexyl)-phenyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl]-1-methyl-harnstoff;
- (1118) 1-(4-[(E)-2-[2-(11-Fluor-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1119) 1-(4-[(E)-2-(2-Hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1120) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-((E)-6-phenyl-hex-5-enyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1121) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-((E)-9-phenyl-non-8-enyl)-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1122) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(5-propylsulfanyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1123) 1-(4-[(E)-2-[2-(7-Methoxy-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1124) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(5-propoxy-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1125) 1-(4-[(E)-2-[2-(11,11-Difluor-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1126) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(3-propyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1127) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(2-propyl-benzofuran-6-yl)-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1128) 1-(4-[(E)-2-[2-(3-Methoxy-4-pentyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1129) 1-(4-[(E)-2-[2-(4-[1,1,2,2,2-²H₅]Ethyl-cyclohex-3-enyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1130) 1-(4-[(E)-2-[2-(4-[1,1,2,2,2-²H₅]Ethyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1131) 12-(8-[(E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethensulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecansäure;
- (1132) 12-(8-[(E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethensulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecansäureethylester;
- (1133) 1-(3,5-Dimethyl-4-[(E)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-1-methyl-harnstoff;
- (1134) 1-(3,5-Dimethyl-4-[(Z)-2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vi-

nyl)-phenyl)-1-methyl-harnstoff;

(1135) 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[3-(4,4,5,5,5-pentafluor-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl}-ethyl)-phenyl]-1-methyl-harnstoff;

(1136) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-1-methyl-harnstoff;

(1137) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4-trimethylsilanyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-1-methyl-harnstoff;

(1138) 1-(4-{2-[2-((1R,3R,5S)-3,5-Bis-trifluormethyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1139) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4-trimethylsilanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-1-methyl-harnstoff;

(1140) 1-(3,5-Dimethyl-4-[2-(4-oxo-2-tridecyl-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl)-1-methyl-harnstoff;

(1141) 1-(3,5-Dimethyl-4-[2-(4-oxo-2-undecyl-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl)-1-methyl-harnstoff;

(1142) 1-(3,5-Dimethyl-4-[2-(2-octyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-phenyl)-1-methyl-harnstoff;

(1143) 1-(4-{2-[2-((1S,3R)-3-Hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1144) 1-[3,5-Dimethyl-4-(2-{2-[3-(3-methyl-butyl)-phenyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl}-ethyl)-phenyl]-1-methyl-harnstoff;

(1145) 1-(4-{2-[2-((1S,3R)-3-Butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1146) 1-[3,5-Dimethyl-4-(2-{2-[1,3,3,3-tetrafluor-3-(3-methyl-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl}-ethyl)-phenyl]-1-methyl-harnstoff;

(1147) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3,3,9,9,9-pentafluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-1-methyl-harnstoff;

(1148) 1-(4-{2-[2-(4-[1,1,2,2,2-²H₅]Ethyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1149) 1-(4-{2-[2-(4-[1,1,2,2,2-²H₅]Ethyl-cyclohex-3-enyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1150) 1-(4-{2-[2-(4-Chlor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1151) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(1,9,9,9-tetrafluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-1-methyl-harnstoff;

(1152) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4'-propyl-biphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-1-methyl-harnstoff;

(1153) 1-(4-{2-[2-(11-Fluor-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1154) 1-(4-{2-[2-(6-Ethylsulfanyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1155) 1-(3,5-Dimethyl-4-[2-(4-oxo-2-[1,1';3',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl)-1-methyl-harnstoff;

(1156) 1-(4-{2-[2-(11,11-Difluor-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

(1157) 1-(3,5-Dimethyl-4-[(E)-2-(4-oxo-2-[1,1';3',1'']terphenyl-3-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-methyl-harnstoff;

(1158) 8-[(E)-2-[4-(4-Hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl]-2-[3-(4,4,5,5,5-pentafluor-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1159) 2-(4-Fluor-3-trifluormethoxy-phenyl)-8-[(E)-2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1160) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1161) 2-(4-Fluor-3-trifluormethoxy-phenyl)-8-[(E)-2-[4-(4-hydroxymethyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1162) 2-(4-Fluor-3-trifluormethyl-phenyl)-8-[(E)-2-[4-(4-hydroxymethyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1163) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dime-

thyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1164) 8-[(E)-2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethensulfonyl]-2-(4-fluor-3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1165) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1166) 8-[(E)-2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethensulfonyl]-2-(4-fluor-3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1167) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-2-methyl-phenyl]-ethensulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1168) 4-[(E)-2-(2-Cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-vinyl]-indol-1-yl]-essigsäure;

(1169) [3-(3-Methyl-4-[(E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-vinyl]-phenyl)-ureido] -essigsäure;

(1170) 12-(8-[2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-2-yl)-dodecansäure;

(1171) 10-(8-[(E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decansäure;

(1172) 10-(8-[(E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decansäureamid;

(1173) 12-(8-[(E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecansäureamid;

(1174) 12-(8-[2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecansäureamid;

(1175) 14-(8-[(E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecansäureamid;

(1176) 14-(8-[2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecansäureamid;

(1178) 8-[2-(2-Amino-5,7-dimethyl-benzoxazol-6-yl)-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1179) 2-Cyclohexyl-8-[2-[2,6-dimethyl-4-(2-oxo-oxazolidin-3-carbonyl)-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1180) N-Methoxy-3,5,N-trimethyl-4-[2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl]-benzamid;

(1181) 8-[2-[4-(Isoxazolidin-2-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1182) 8-[2-[4-(Isoxazolidin-2-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1183) 8-[2-[4-(1,1-Dioxo-1λ⁶-thiomorpholine-4-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1184) 8-[2-[4-[(R)-2,3-Dihydroxy-propoxy]-piperidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1185) 2-(3,4-Dichlor-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1186) 2-(3-Chlor-4-fluor-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1187) [3-(8-[(E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-phenyl]-acetonitril;

(1188) 2-(3-Chlor-4-trifluormethyl-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1189) 8-[(E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(4-pentafluorsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1190) 8-[(E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-pentafluorsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1191) 8-[(E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-[4-(3-trifluormethyl-phenoxy-methyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1192) 2-(4-Fluor-2,3-dimethyl-phenyl)-8-[(E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

(1193) 8-[(E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-methyl-4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (1194) 2-Benzo[1,3]dioxol-5-yl-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1195) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(4-pentafluorethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1196) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(3-pentafluorethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1197) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[4-(2,2,2-trifluor-1,1-dimethyl-ethoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1198) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1199) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[4-(2,2,3,3-tetrafluor-propoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1200) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[4-(2,2,2-trifluor-ethoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1201) 2-[3-Chlor-4-(2,2,3,3-tetrafluor-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1202) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[4-(4,4,5,5-pentafluor-pentyloxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1203) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[4-(2,2,3,3-pentafluor-propoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1204) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[4-(4,4,4-trifluor-butoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1205) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[3-(3-trifluormethyl-phenoxy-methyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1206) 2-[3-(1,1-Difluor-ethyl)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1207) 2-[3-Fluor-4-(4,4,5,5-pentafluor-pentyloxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1208) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(1H-indol-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1209) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[3-(2,2,2-trifluor-1,1-dimethyl-ethoxy)-4-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1210) 2-[4-Chlor-3-(2,2,2-trifluor-1,1-dimethyl-ethoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1211) 2-(3-[1,1,2,2,3,3,4,4,4-²H₉]Butoxy-4-fluor-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1212) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(2,2,3,3-tetrafluor-2,3-dihydro-benzo[1,4]dioxin-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1213) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-[4-(1,1,2,2-tetrafluor-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1214) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(3-methyl-4-trifluoromethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1215) 2-[4-Fluor-3-(3-fluor-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1216) 2-[3-Chlor-4-(2,2,2-trifluor-ethoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1217) 2-(3-Difluormethyl-4-fluor-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1218) 2-[4-Fluor-3-(4,4,4-trifluor-butoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1219) 2-(4-Difluormethyl-3-fluor-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1220) 2-(4-[1,1,2,2,3,3,4,4,4-²H₉]Butoxy-3-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1221) 2-(4-[1,1,2,2,2-²H₅]Ethoxy-3-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1222) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethensulfonyl)-2-(4-[1,2,2,2,2,2-²H₇]isopropoxy-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (1223) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[3-(2,2,3,3-tetrafluor-propoxy)-4-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1224) 2-(3-[1,1,2,2,2,2- H_5]Etoxy-4-fluor-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 5 (1225) 2-(4-fluor-3-[1,2,2,2,2,2- H_7]Isopropoxy-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1226) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[3-(2,2,3,3-tetrafluor-propoxy)-5-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1227) 2-[4-Chlor-3-(4,4,4-trifluor-butoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 10 (1228) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[3-(2,2,3,3,3-pentafluor-propoxy)-4-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1229) 2-[4-Chlor-3-(2,2,3,3-tetrafluor-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 15 (1230) 2-[4-Chlor-3-(2,2,3,3,3-pentafluor-propoxy)-phenyl]-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1231) 2-(3-[1,1,2,2,3,3,4,4,4- H_9]Butoxy-4-chlor-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1232) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[3-(4,4,4-trifluor-butoxy)-4-trifluormethoxy-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 20 (1233) 2-(3-[1,1,2,2,3,3,4,4,4- H_9]Butoxy-5-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1234) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 25 (1235) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-methyl-4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1236) 2-(3-Chlor-4-trifluormethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1237) 2-(3,4-Dimethyl-phenyl)-8-((E)-2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 30 (1238) 8-((E)-2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-methoxy-3-methyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1239) 2-(4-Fluor-2,5-dimethyl-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 35 (1240) 2-(4-Fluor-2,3-dimethyl-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1241) 2-(3-Chlor-4-trifluormethyl-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxypiperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1242) 2-(3-Chlor-4-fluor-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 40 (1243) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethylphenyl]-ethansulfonyl)-2-(3-pentafluorethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1244) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-pentafluorethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 45 (1245) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-pentafluorsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1246) 2-[3-Chlor-4-(2,2,2-trifluor-ethoxy)-phenyl]-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1247) 2-[3-Chlor-4-(2,2,3,3-tetrafluor-propoxy)-phenyl]-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 50 (1248) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-(2,2,2-trifluor-ethoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1249) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethylphenyl]-ethansulfonyl)-2-[4-(2,2,3,3,3-pentafluor-propoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- 55 (1250) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-(4,4,4-trifluor-butoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1251) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-(2,2,3,3-tetrafluor-propoxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (1252) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-(4,4,5,5,5-pentafluor-pentyloxy)-3-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1253) 2-(4-[1,1,2,2,2-²H₅]Ethoxy-3-trifluormethyl-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1254) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-methyl-4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1255) 2-(4-Difluormethyl-3-fluor-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxypiperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1256) 2-(3-Difluormethyl-4-fluor-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1257) 2-(3,4-Dichlor-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1258) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(3-methyl-4-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1259) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(2-methyl-4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1260) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[3-fluor-4-(2,2,2-trifluor-ethoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1261) 2-(4-[1,1,2,2,3,3,4,4,4-²H₉]Butoxy-3-trifluormethyl-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1262) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(4-[1,2,2,2,2,2-²H₇]isopropoxy-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1263) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-fluor-3-(4,4,5,5,5-pentafluor-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1264) 2-[4-Fluor-3-(3-fluor-propoxy)-phenyl]-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1265) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-(2,2,2-trifluor-1,1-dimethyl-ethoxy)-3-trifluormethylphenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1266) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[3-(2,2,2-trifluor-1,1-dimethyl-ethoxy)-4-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1267) 2-[3-(1,1-Difluor-ethyl)-phenyl]-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1268) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-(1H-indol-6-yl)-1,3,8-triaza-spiro [4.5]dec-1-en-4-on;
- (1269) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethylphenyl]-ethansulfonyl)-2-(4-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1270) 2-[4-Chlor-3-(2,2,3,3,3-pentafluor-propoxy)-phenyl]-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1271) 2-[4-Chlor-3-(2,2,2-trifluor-1,1-dimethyl-ethoxy)-phenyl]-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1272) 2-(3-[1,1,2,2,2-²H₅]Ethoxy-4-fluor-phenyl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1273) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[3-(4,4,4-trifluor-butoxy)-4-trifluormethoxy-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1274) 8-((E)-2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[3-(2,2,3,3-tetrafluor-propoxy)-4-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1275) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-[2-[4-(4-fluormethyl-4-hydroxypiperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1276) 8-[2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(4-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1277) 8-[2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(3-methyl-4-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1278) 2-(3-Chlor-4-trifluormethyl-phenyl)-8-[2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1279) 8-[2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-[3-(2,2,3,3-tetrafluor-propoxy)-4-trifluormethyl-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1280) 2-(2,2-difluor-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(4-fluormethyl-4-hydroxypiperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;

- (1281) 8-[2-[4-(4-fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-2-(2,2,3,3-tetrafluor-2,3-dihydro-benzo[1,4]dioxin-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1282) [3-(8-((E)-2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-phenyl]-acetonitrile;
- (1283) 8-((E)-2-[2-methyl-4-(piperidin-4-yloxy)-phenyl]-ethansulfonyl)-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1284) 8-[2-(4-aminomethyl-2-methyl-phenyl)-ethansulfonyl]-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1285) 2-(4-methyl-cyclohexyl)-8-((E)-2-[2-methyl-4-(2-oxo-piperazin-1-yl)-phenyl]-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1286) 8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethansulfonyl]-2-pyrrolidin-2-yl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on hydrochloride;
- (1287) 2-(1,1-difluor-ethyl)-8-[2-(2,6-dimethyl-4-methylamino-phenyl)-ethansulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on hydrochloride;
- (1288) 8-((E)-2-(2,6-dimethyl-4-[1,1,1-²H₃]-methylamino-phenyl)-ethansulfonyl)-2-(4-fluor-3-trifluormethyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on hydrochloride;
- (1289) 8-((E)-2-(2,6-dimethyl-4-methylaminomethyl-phenyl)-ethansulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1290) 8-((E)-2-(4-aminomethyl-2,6-dimethyl-phenyl)-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1291) 8-((E)-2-(4-aminomethyl-2,6-dimethyl-phenyl)-ethansulfonyl)-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1292) 8-((E)-2-(2,6-dimethyl-4-methylaminomethyl-phenyl)-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1293) 8-[2-(4-amino-3-chlor-2-methyl-phenyl)-ethansulfonyl]-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1294) N-(1-Acetyl-piperidin-4-yl)-N-(3-methyl-4-((E)-2-[4-oxo-2-(3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-acetamid;
- (1295) 8-[2-[4-(4,5-Dihydro-thiazol-2-ylamino)-2-methyl-phenyl]-ethansulfonyl]-2-(4-ethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1296) N-[4-(2-[2-[4-(4-Fluor-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-3-methyl-phenyl]-acetamid;
- (1297) 1-(3,5-Dimethyl-4-{2-[2-((1S,3R)-3-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1298) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3,3,5,5-tetramethyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1299) 3-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-5-hydroxymethyl-imidazolidine-2,4-dion;
- (1300) 1-[3,5-Dimethyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzyl]-1-methyl-harnstoff;
- (1301) 1-(3,5-Dimethyl-4-((E)-2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-benzyl)-1-methyl-harnstoff;
- (1302) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(8,8,9,9,9-pentafluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1303) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(8,8,9,9,9-pentafluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-methyl-harnstoff;
- (1304) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(9,9,9-trifluor-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-methyl-harnstoff;
- (1305) 1-(3,5-Dimethyl-4-((E)-2-[4-oxo-2-(3-trifluormethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl)-1-methyl-harnstoff;
- (1306) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3-trifluormethylsulfanyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1307) 1-(4-((E)-2-[2-(11-Hydroxy-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1308) [11-(8-((E)-2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-undecyl]-carbaminsäure-tert-butylester;
- (1309) 1-(4-((E)-2-[2-(9-Hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-3,5-dimethyl-phenyl)-1-methyl-harnstoff;

- (1310) 1-(4-{2-[2-(4-Isopropyliden-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1311) 1-(4-{2-[2-(1,1-Difluor-ethyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1312) 1-(4-{(E)-2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-[1,1,1-²H₃]methyl-harnstoff;
- (1313) [3,5-Dimethyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl]-harnstoff;
- (1314) (3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzyl)-harnstoff;
- (1315) 1-(4-{(E)-2-[2-(11-Amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1316) 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[5-(propan-1-sulfinyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1317) 1-(4-{2-[2-(6-Ethansulfinyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1318) 1-[3,5-Dimethyl-4-((E)-2-{4-oxo-2-[5-(propan-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl)-phenyl]-1-methyl-harnstoff;
- (1319) 1-(4-{(E)-2-[2-(7-Methansulfonyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1320) 1-(4-{2-[2-(6-Ethansulfonyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1321) 1-(4-{(E)-2-[2-(9,9-Difluor-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1322) 1-(4-{2-[2-(9-Hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1323) 1-(4-{2-[2-(9,9-Difluor-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1324) 1-(4-{2-[2-(9-Amino-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1325) 1-(4-{(E)-2-[2-(9-Fluor-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1326) 1-(4-{(Z)-2-[2-(9-Fluor-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1327) 1-(4-{2-[2-(9-Fluor-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1328) 8-{2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-[3-(4,4,5,5,5-pentafluor-pentyloxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1329) 8-{2-[4-((R)-2,3-Dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-(4-fluor-3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1330) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-dimethyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1331) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-{2-[4-(3,4-dihydroxy-butoxy)-2-methylphenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1332) 8-{2-[4-(3,4-Dihydroxy-butoxy)-2-methyl-phenyl]-ethansulfonyl}-2-(4-fluor-3-trifluormethoxy-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1333) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1334) 2-(4-Fluor-3-trifluormethoxy-phenyl)-8-{2-[4-(4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1335) 2-(4-Fluor-3-trifluormethoxy-phenyl)-8-{2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1336) 2-(2,2-Difluor-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1337) 2-(4-Fluor-3-trifluormethoxy-phenyl)-8-{2-[4-(3-hydroxy-azetidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1338) 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[3-(4,4,4-trifluor-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;

- (1339) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(4,4,4-trifluor-butoxy)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- (1340) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(5-phenyl-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- 5 (1341) 1-(3,5-Dimethyl-4-{2-[2-(3-nonafluorbutyl-phenyl)-4-oxo-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1342) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(4-pentyl-phenyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- 10 (1343) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1344) 1-(4-{2-[2-(6-Ethoxy-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1345) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[3-(6,6,6-trifluor-hexyl)-phenyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- 15 (1346) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(6,6,6-trifluor-hexyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- (1347) 12-(8-{2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-dodecansäureethylester;
- (1348) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(9-phenyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- 20 (1349) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(6-phenyl-hexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1350) 14-(8-{2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-tetradecansäure;
- 25 (1351) 1-(4-{2-[2-(7-Methoxy-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1352) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(5-propoxy-pentyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1353) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(3-propyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- 30 (1354) 1-(4-{2-[2-(3-Methoxy-4-pentyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1355) 1-(3,5-Dimethyl-4-{2-[4-oxo-2-(2-propyl-benzofuran-6-yl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- 35 (1356) 1-[3,5-Dimethyl-4-(2-{4-oxo-2-[5-(propan-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- (1357) 1-(4-{2-[2-(7-Methansulfonyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl} -3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1358) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(4,4,4-trifluor-butyl)-phenyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- 40 (1359) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[3-(4,4,4-trifluor-butyl)-phenyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-phenyl]-1-methyl-harnstoff;
- (1360) [11-(8-{2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-undecyl]-carbaminsäure-tert-butylester;
- 45 (1361) 1-(4-{2-[2-(11-Hydroxy-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1362) 10-(8-{2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decansäure;
- (1363) 10-(8-{2-[2,6-Dimethyl-4-(1-methyl-ureido)-phenyl]-ethansulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-2-yl)-decansäureamid;
- 50 (1364) 1-(3,5-Dimethyl-4-{2-[2-(4-[1,1,1-²H₃]-methyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-1-methyl-harnstoff;
- (1365) 1-(3,5-Dimethyl-4-{(E)-2-[4-oxo-2-(9-phenyl-nonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-vinyl}-phenyl)-1-methyl-harnstoff;
- 55 (1366) 1-(4-{2-[2-(11-Amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl} -3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1367) 3-[(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-4-ethoxy-cyclobut-3-en-1,2-dion;

- (1368) 3-Amino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-ene-1,2-dion;
- (1369) 3-Dimethylamino-4-[(3,5-dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-methyl-amino]-cyclobut-3-en-1,2-dion;
- (1370) N-(4-{2-[2-(4-Ethynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3-methyl-phenyl)-acetamid;
- (1371) 1-(4-{2-[4-[(Z)-Hydroxyimino]-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-methyl-harnstoff;
- (1372) 2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonsäure-2-methyl-benzylamid;
- (1373) 2-(4-Methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonsäure-(2-o-tolyl-ethyl)-amid;
- (1374) 2-Cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-2-hydroxy-ethansulfonyl}-1,3,8-triaza-spiro [4.5]dec-1-en-4-on;
- (1375) 2-Cyclohexyl-8-(2-oxo-2-o-tolyl-ethansulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1376) 2-Cyclohexyl-8-(2-o-tolyl-ethinesulfonyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1377) 2-Cyclohexyl-8-[2-(1H-indol-4-yl)-ethinesulfonyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1378) 3,5,N,N-Tetramethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzamid;
- (1379) 3,5,N,N-Tetramethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (1380) 8-{2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1381) 8-{2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1382) 8-{2-[4-(4-Hydroxy-4-methyl-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1383) 8-{2-[4-(4-Fluormethyl-4-hydroxy-piperidin-1-carbonyl)-2,6-dimethyl-phenyl]-ethansulfonyl}-2-(4-methyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1384) 8-(2-[4-[4-((R)-2,3-Dihydroxy-propoxy)-piperidin-1-carbonyl]-2,6-dimethyl-phenyl]-ethansulfonyl)-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1385) 3,5-Dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]dec-1-en-8-sulfonyl]-ethyl)-benzoesäurehydrazid;
- (1386) 8-{2-[2,6-Dimethyl-4-(pyrazolidin-1-carbonyl)-phenyl]-ethansulfonyl}-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1387) N,N-Dimethyl-2-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (1388) N-Methoxy-2-(3-methyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;
- (1389) 1-[3,5-Dimethyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl]-1-methyl-harnstoff;
- (1390) 1-(3,5-Dimethyl-4-{2-[2-(4-methyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzyl)-1-methyl-harnstoff;
- (1391) 1-(4-{2-[2-(4-Fluor-3-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,5-dimethyl-phenyl)-1-[1,1,1-²H₃]methyl-harnstoff;
- (1392) 2-Cyclohexyl-8-{2-[1-((2S,3S)-2,3,4-trihydroxy-butyl)-1H-indol-4-yl]-ethansulfonyl}-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1393) 8-{2-[1-((2S,3S)-4-Benzyloxy-2,3-dihydroxy-butyl)-1H-indol-4-yl]-ethansulfonyl}-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-on;
- (1394) N-[3-Methyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluor-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl)-benzyl]-acetamid;
- (1395) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (1396) 3,N,N-Trimethyl-4-{2-[4-oxo-2-(4-trifluormethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (1397) 3,N,N-Trimethyl-4-{2-[2-(methyl-phenyl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-benzamid;
- (1398) 4-{2-[2-(Benzyl-methyl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;
- (1399) 4-{2-[2-(Cyclohexylmethyl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trime-

thyl-benzamid;

(1400) 3,N,N-Trimethyl-4-(2-{2-[methyl-(4-trifluormethyl-phenyl)-amino]-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl}-ethyl)-benzamid;

(1401) N-(3-Methyl-4-{2-[4-oxo-2-(4-trifluormethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(1402) N-(3-Methyl-4-{2-[4-oxo-2-(3-trifluormethyl-phenylamino)-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-phenyl)-acetamid;

(1403) 4-{2-[2-(4-Butyl-piperidin-1-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(1404) 4-{2-[2-(4-Butyl-cyclohexylamino)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N,N-trimethyl-benzamid;

(1405) 4-{2-[2-(3-But-3-enyloxy-5-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N-dimethyl-N-pent-4-enyl-benzamid;


(1419) N-(2-Allyloxy-ethyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluormethyl-phenyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl]-ethyl}-3,N-dimethyl-benzamid;

(1443) Dec-9-ensäure-{4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-amid; oder

(1446) Hept-6-ensäure-{4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-en-8-sulfonyl)-ethyl]-3,5-dimethyl-phenyl}-methyl-amid,

oder ein pharmakologisch verträgliches Salz davon.


11. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach einem der Ansprüche 1 bis 5, wobei U C1-C6-Alkylen oder einen Rest, ausgewählt aus durch die folgende Formel dargestellten Resten, bedeutet:



A gleich O ist;

R₄₆ ausgewählt ist aus Wasserstoff, C1-C10-Alkyl, C1-C10-Halogenalkyl und C1-C10-Hydroxyalkyl;

V ausgewählt ist aus:



E Pyrrolidin oder Piperidin, gegebenenfalls substituiert mit einem Hydroxylrest, ist; und R₇ ausgewählt ist aus:


- 1) Wasserstoff,
- 2) C1-C10-Alkyl und
- 3) C1-C10-Hydroxyalkyl.

12. Ein Arzneimittel, umfassend die Verbindung oder ein pharmakologisch verträgliches Salz davon nach einem der Ansprüche 1 bis 11 als einen Wirkstoff.


13. Ein prophylaktisches oder therapeutisches Mittel zur Verwendung bei der Behandlung oder Prävention von Osteoporose, Fraktur, Osteomalazie, Arthritis, Thrombozytopenie, Hypoparathyreoidismus, Hyperphosphatämie oder tumoraler Calcinose, umfassend die Verbindung oder ein pharmakologisch verträgliches Salz davon nach einem der Ansprüche 1 bis 11 als einen Wirkstoff.
14. Verwendung der Verbindung oder eines pharmakologisch verträglichen Salzes davon nach einem der Ansprüche 1 bis 11 zur Herstellung eines prophylaktischen oder therapeutischen Mittels zur Verwendung bei der Behandlung oder Prävention von Osteoporose, Fraktur, Osteomalazie, Arthritis, Thrombozytopenie, Hypoparathyreoidismus, Hyperphosphatämie oder tumoraler Calcinose.
15. Die Verbindung oder ein pharmakologisch verträgliches Salz davon nach einem der Ansprüche 1 bis 11 zur Verwendung bei der Behandlung oder Prävention von Osteoporose, Fraktur, Osteomalazie, Arthritis, Thrombozytopenie, Hypoparathyreoidismus, Hyperphosphatämie oder tumoraler Calcinose.

Revendications

1. Composé représenté par la formule générale (1) ou (2) suivante :



(1)



(2),

ou un sel pharmacologiquement acceptable d'un composé représenté par la formule (1) ou (2) ;
[où dans la formule générale (1)

W est choisi parmi :

- 1) une liaison simple,
- 2) un groupe alkylène en C₁ à C₁₀ contenant éventuellement un groupe carbonyle, où le groupe alkylène est éventuellement substitué par un ou plusieurs atomes d'halogène et/ou un ou plusieurs groupes hydroxyle,
- 3) un groupe alcénylène en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène,

- 4) un groupe alcynylène en C_2 à C_{10} ,
- 5) un groupe arylène éventuellement substitué par un ou plusieurs atomes d'halogène,
- 6) un groupe hétéroarylène éventuellement substitué par un ou plusieurs atomes d'halogène,
- 7) un groupe hétéroalkylène en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène,
- 8) un groupe $-NH-$, $-NHCH_2-$ ou $-NHCH_2CH_2-$,
- 9) un groupe cycloalkylène et
- 10) un groupe $-(cycloalkylène)-CH_2-$;

X est choisi parmi la liaison ou les groupes suivants :

- 1) une liaison simple,
- 2) un groupe alkylène en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène ou un groupe cycloalkyle,
- 3) un groupe alcénylène en C_2 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène,
- 4) un groupe alcynylène en C_2 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène,
- 5) un groupe oxyalkylène en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène, et
- 6) un groupe $-NR_{47}-$


où R_{47} est choisi parmi :

- i) un atome d'hydrogène et
- ii) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène ;

Y est choisi parmi :

- 1) un atome d'oxygène,
- 2) un atome de soufre et
- 3) un groupe $=NR_{37}$,

ou 4) Y représente un groupe $-NR_{38}R_{39}$ représenté par la formule (A) suivante :



qui peut être tautomère ;

R_{37} est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe hydroxy et
- 3) un groupe alcoxy en C_1 à C_{10} ;

R_{38} et R_{39} sont choisis indépendamment parmi un atome d'hydrogène ou un groupe alkyle en C_1 à C_{10} éventuellement substitué par un groupe cycloalkyle, ou

R_{38} et R_{39} peuvent être liés l'un à l'autre pour former un cycle choisi dans le groupe constitué des cycles azétidinyle, pyrrolidinyle, pipéridinyle, pipérazinyle et morpholinyle, et le cycle est éventuellement substitué par un groupe alkyle en C_1 à C_{10} ; m représente un nombre entier de 0 à 2 ;

n représente un nombre entier de 0 à 2 ;

R_1 est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe cycloalkyle éventuellement substitué par un ou plusieurs groupes choisis parmi R_4 , R_5 et R_6 ,
- 3) un hétérocycle éventuellement substitué par un ou plusieurs groupes choisis parmi R_{25} , R_4 , R_5 et R_6 ,

- 4) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis parmi R_3 , R_4 , R_5 et R_6 , et
 5) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes choisis parmi R_{25} , R_4 , R_5 et R_6 ;

5 R_3 est choisi parmi :

- 1) un groupe $-\text{CONR}_7\text{R}_8$,
 2) un groupe $-\text{OR}_9$,
 3) un groupe $-\text{NR}_9\text{R}_{10}$,
 10 4) un groupe $-\text{N}(\text{R}_9)\text{COR}_{11}$,
 5) un groupe $-\text{N}(\text{R}_9)\text{SO}_2\text{R}_{12}$,
 6) un groupe $-\text{SO}_2\text{R}_{15}$,
 7) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, un groupe hydroxyle, $-\text{COR}_{16}$ et $-\text{NR}_{13}\text{R}_{14}$,
 15 8) un groupe hétéroaryle ayant éventuellement un groupe alkyle en C_1 à C_{10} et/ou un groupe alcoxy en C_1 à C_{10} en tant que substituant(s) et
 9) un groupe $-\text{N}(\text{R}_9)\text{CSR}_{11}$;

20 R_4 est choisi parmi :

- 1) un atome d'halogène,
 2) un groupe cyano,
 3) un groupe nitro,
 4) un groupe amino,
 25 5) un groupe NHCOR_{26} ,
 6) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes hydroxycarbonyle, alcoxy en C_1 à C_{10} -carbonyle et aminocarbonyle,
 7) un groupe halogénoalkyle en C_1 à C_{10} ,
 8) un groupe alcoxy en C_1 à C_{10} ,
 30 9) un groupe halogénoalkyl en C_1 à C_{10} -carbonyle,
 10) un groupe $-\text{COR}_{16}$,
 11) un groupe hydroxyalkyle en C_1 à C_{10} et
 12) un groupe hétéroalkyle en C_1 à C_{10} ;

35 R_5 est choisi parmi un atome d'halogène, les groupes alkyle en C_1 à C_{10} , halogénoalkyle en C_1 à C_{10} et alcoxy en C_1 à C_{10} ;

R_6 est choisi parmi un atome d'halogène, les groupes alkyle en C_1 à C_{10} et halogénoalkyle en C_1 à C_{10} ;

R_7 est choisi parmi :

- 40 1) un atome d'hydrogène,
 2) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes amino et alkyl en C_1 à C_{10} -amino,
 3) un groupe hydroxyalkyle en C_1 à C_{10} ,
 4) un groupe halogénoalkyle en C_1 à C_{10} ,
 45 5) un groupe hétéroalkyle en C_1 à C_{10} ,
 6) un groupe hétéroalkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes hydroxyle, alkyl en C_1 à C_{10} -amino et alcényle en C_2 à C_{10} ,
 7) un groupe aryle,
 8) un groupe hétéroaryle,
 50 9) un groupe arylalkyle en C_1 à C_{10} ,
 10) un hétérocycle éventuellement substitué par un groupe alkyle en C_1 à C_{10} ;
 11) un groupe $-(\text{CH}_2)_L\text{COR}_{16}$ (où L représente un nombre entier de 1 à 4),
 12) un groupe alcoxy en C_1 à C_{10} ,
 13) un groupe alcényle en C_2 à C_{10} et
 55 14) un groupe $\text{NR}_{40}\text{R}_{41}$;

R_{40} et R_{41} sont choisis indépendamment parmi un atome d'hydrogène, les groupes alkyle en C_1 à C_{10} et alkyl en C_1 à C_{10} -carbonyle, ou R_{40} et R_{41} peuvent être liés l'un à l'autre pour former un cycle choisi parmi les cycles

azétidinyle, pyrrolidinyle, pipéridinyle, pipérazinyle et morpholinyle, et l'hétérocycle est éventuellement substitué par un groupe alkyle en C₁ à C₁₀ ;

R₈ est choisi parmi un atome d'hydrogène et un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène et/ou un ou plusieurs groupes hydroxyle ;

R₇ et R₈ peuvent être liés pour former un hétérocycle de 4 à 7 chaînons contenant éventuellement un ou plusieurs éléments ou groupes supplémentaires choisis indépendamment parmi les atomes O, N, S, les groupes SO et SO₂, et l'hétérocycle contient éventuellement un groupe carbonyle, et l'hétérocycle est éventuellement substitué par un ou plusieurs substituants choisis indépendamment parmi :

- 1) un atome d'halogène,
- 2) un groupe alkyle en C₁ à C₁₀ ayant éventuellement un ou plusieurs groupes alkyl en C₁ à C₁₀-amino en tant que substituant(s),
- 3) un groupe halogénoalkyle en C₁ à C₁₀,
- 4) un groupe hydroxyle,
- 5) un groupe hydroxyalkyle en C₁ à C₁₀,
- 6) un groupe alcoxy en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino,
- 7) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino,
- 8) un groupe hétéroalkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino,
- 9) un hétérocycle éventuellement substitué par un groupe alkyle en C₁ à C₁₀,
- 10) un groupe hétéroaryle éventuellement substitué par un groupe alkyle en C₁ à C₁₀,
- 11) un groupe hétérocyclyalkyle en C₁ à C₁₀,
- 12) un groupe -COR₁₆,
- 13) un groupe -NR₁₉R₂₀,
- 14) un groupe -SO₂R₂₁,
- 15) un groupe alcoxy en C₁ à C₁₀-alkyle en C₁ à C₁₀ ayant éventuellement un ou plusieurs groupes hydroxyle en tant que substituant(s) et
- 16) un groupe hydroxyalkyl en C₁ à C₁₀-oxy, dans lequel l'atome d'hydrogène du groupe hydroxyle est éventuellement remplacé par un groupe hydroxyalkyle en C₁ à C₁₀, et

l'hétérocycle peut former en outre un cycle spiro conjointement avec un hétérocycle de 4 à 6 chaînons, et l'hétérocycle de 4 à 6 chaînons lié contient éventuellement des atomes O et N en tant qu'élément formant le cycle en plus des atomes de carbone, et les un ou plusieurs atomes de carbone peuvent être oxydés pour former un groupe carbonyle, et l'hétérocycle de 4 à 6 chaînons est éventuellement en outre substitué par un groupe alkyle en C₁ à C₁₀ ;

R₁₆ est choisi parmi :

- 1) un groupe hydroxyle,
- 2) un groupe alcoxy en C₁ à C₁₀,
- 3) un groupe NR₁₇R₁₈ et
- 4) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs substituants choisis parmi un atome d'halogène, les groupes hydroxyle, alcoxy en C₁ à C₁₀-carbonyle ou alkyl en C₁ à C₁₀-amino ;

R₁₇ est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes aryle, amino, alkyl en C₁ à C₁₀-amino, alkyl en C₁ à C₁₀-carbonylamino et hydroxyle,
- 3) un groupe hétéroaryle et
- 4) un groupe alcoxy en C₁ à C₁₀ ;

R₁₈ est choisi parmi un atome d'hydrogène, les groupes alkyle en C₁ à C₁₀ et hydroxyalkyle en C₁ à C₁₀ ;

R₁₇ et R₁₈ peuvent être liés l'un à l'autre pour former un cycle choisi parmi les cycles azétidinyle, pyrrolidinyle, pipéridinyle, pipérazinyle et morpholinyle, et le cycle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment les uns des autres parmi un groupe alkyle en C₁ à C₁₀, un atome d'halogène et un groupe alcoxy en C₁ à C₁₀-carbonyle ;

R₁₉ est choisi parmi un atome d'hydrogène, les groupes alkyle en C₁ à C₁₀, halogénoalkyle en C₁ à C₁₀, alkyl en C₁ à C₁₀-carbonyle, hydroxyalkyle en C₁ à C₁₀, aminoalkyle en C₁ à C₁₀, alcoxy en C₁ à C₁₀-carbonyle et hétéroalkyle en C₁ à C₁₀ ;

R₂₀ est choisi parmi un atome d'hydrogène et un groupe alkyle en C₁ à C₁₀ ;

R₁₉ et R₂₀ peuvent être liés pour former un cycle choisi parmi les cycles azétidinyle, pyrrolidinyle, pipéridinyle, pipérazinyle et morpholinyle, et le cycle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment les uns des autres parmi un groupe alkyle en C₁ à C₁₀ et un atome d'halogène ;

R₂₁ est choisi parmi :

- 1) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un groupe aryle,
- 2) un groupe amino,
- 3) un groupe alkyl en C₁ à C₁₀-amino et
- 4) un groupe aryle éventuellement substitué par un groupe alkyle en C₁ à C₁₀ ;

R₉ est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes R₂₃,
- 3) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino,
- 4) un groupe cycloalkyle éventuellement substitué par un ou plusieurs atomes d'halogène ou un ou plusieurs groupes hydroxyle,
- 5) un hétérocycle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyle en C₁ à C₁₀, alkyl en C₁ à C₁₀-carbonyle, alcoxy en C₁ à C₁₀, alcoxy en C₁ à C₁₀-carbonyle, amino et un atome d'halogène,
- 6) un groupe hétéroalkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène et un groupe hydroxyle,
- 7) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes alkyle en C₁ à C₁₀, alkyl en C₁ à C₁₀-carbonyle, alcoxy en C₁ à C₁₀-carbonyle et un atome d'halogène et
- 8) un groupe cycloalcényle éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes alcoxy en C₁ à C₁₀, alkyl en C₁ à C₁₀-amino, amino, hydroxyle et un atome d'halogène, où le groupe cycloalcényle contient éventuellement un groupe carbonyle ;

R₂₃ est choisi indépendamment parmi :

- 1) un atome d'halogène,
- 2) un groupe hydroxyle,
- 3) un groupe alkyl en C₁ à C₁₀-carbonyloxy,
- 4) un groupe -COR₁₆,
- 5) un groupe amino,
- 6) un groupe alkyl en C₁ à C₁₀-amino,
- 7) un hétérocycle éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes alkyle en C₁ à C₁₀, alkyl en C₁ à C₁₀-carbonyle, alcoxy en C₁ à C₁₀-carbonyle et un atome d'halogène et
- 8) un groupe cyano ;

R₁₀ est choisi parmi :

- 1) un atome d'hydrogène et
- 2) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis parmi un atome d'halogène, les groupes hydroxyle et aryle ;

R₉ et R₁₀ peuvent être liés pour former un hétérocycle de 4 à 7 chaînons contenant éventuellement un ou plusieurs éléments ou groupes supplémentaires choisis indépendamment parmi les atomes N, O, S, les groupes SO, SO₂, carbonyle et thiocarbonyle, et l'hétérocycle est éventuellement substitué par un ou plusieurs substituants choisis indépendamment parmi les groupes R₂₄ ;

R₂₄ est choisi indépendamment parmi :

- 1) un atome d'halogène,
- 2) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyl en C_1 à C_{10} -amino et alkyl en C_1 à C_{10} -carbonylamino,
- 3) un groupe halogénoalkyle en C_1 à C_{10} ,
- 4) un groupe hydroxyle,
- 5) un groupe hydroxyalkyle en C_1 à C_{10} ,
- 6) un groupe alcoxy en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C_1 à C_{10} -amino,
- 7) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C_1 à C_{10} -amino,
- 8) un groupe hétéroalkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C_1 à C_{10} -amino,
- 9) un hétérocycle éventuellement substitué par un groupe alkyle en C_1 à C_{10} ,
- 10) un groupe hétéroaryle,
- 11) un groupe hétérocyclylalkyle en C_1 à C_{10} ,
- 12) un groupe $-COR_{16}$,
- 13) un groupe $-NR_{19}R_{20}$ et
- 14) un groupe $-SO_2R_{21}$;

R_{11} est choisi parmi :

- 1) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

- i) un groupe hydroxyle,
- ii) un groupe $-NR_{17}R_{18}$,
- iii) un groupe alcoxy en C_1 à C_{10} ,
- iv) un atome d'halogène,
- v) un groupe alcoxy en C_1 à C_{10} -carbonyle,
- vi) un groupe aminocarbonyle et
- vii) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, alcoxy en C_1 à C_{10} , amino, alkyl en C_1 à C_{10} -amino et $-COR_{22}$,

- 2) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, alcoxy en C_1 à C_{10} , amino, alkyl en C_1 à C_{10} -amino et $-COR_{22}$,
- 3) un groupe cycloalkyle éventuellement substitué par un ou plusieurs atomes d'halogène,
- 4) un hétérocycle éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes alkyle en C_1 à C_{10} , alkyl en C_1 à C_{10} -carbonyle, alcoxy en C_1 à C_{10} -carbonyle et un atome d'halogène,
- 5) un groupe alcoxy en C_1 à C_{10} , dans lequel le groupe alkyle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyl en C_1 à C_{10} -carbonylamino, amino, alkyl en C_1 à C_{10} -amino et hydroxyle,
- 6) un groupe amino,
- 7) un groupe alkyl en C_1 à C_{10} -amino, dans lequel le groupe alkyle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyl en C_1 à C_{10} -carbonylamino, amino, alkyl en C_1 à C_{10} -amino, hydroxycarbonyle et hydroxyle et
- 8) un groupe alcényle en C_2 à C_{10} ;

R_{22} est choisi parmi les groupes alcoxy en C_1 à C_{10} , hydroxyle, amino et alkyl en C_1 à C_{10} -amino ;

R_{12} est choisi parmi :

- 1) un groupe alkyle en C_1 à C_{10} ,
- 2) un groupe amino et
- 3) un groupe alkyl en C_1 à C_{10} -amino, dans lequel le groupe alkyle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes amino, alkyl en C_1 à C_{10} -amino et hydroxyle ;

R_{13} est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe alkyle en C₁ à C₁₀,
- 3) un groupe alkyl en C₁ à C₁₀-carbonyle, dans lequel le groupe alkyle est éventuellement substitué par un ou plusieurs groupes hydroxyle,
- 4) un groupe alcoxy en C₁ à C₁₀-carbonyle,
- 5) un groupe aminocarbonyle,
- 6) un groupe alkyl en C₁ à C₁₀-aminocarbonyle et
- 7) un groupe carbonyle hétérocyclique éventuellement substitué par un groupe alkyle en C₁ à C₁₀;

R₁₄ est choisi parmi :

- 1) un atome d'hydrogène et
- 2) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino ;

R₁₃ et R₁₄ peuvent être liés pour former un hétérocycle de 4 à 7 chaînons contenant éventuellement un ou plusieurs éléments ou groupes supplémentaires choisis indépendamment parmi les atomes O, N, S, les groupes SO et SO₂, et l'hétérocycle contient éventuellement un groupe carbonyle, et l'hétérocycle est éventuellement substitué par un groupe alkyle en C₁ à C₁₀ ;

R₁₅ est choisi parmi :

- 1) un groupe alkyle en C₁ à C₁₀ et
- 2) un groupe -NR₃₅R₃₆ ;

R₃₅ est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

- i) un atome d'halogène,
- ii) un groupe hydroxyle,
- iii) un groupe alkyl en C₁ à C₁₀-carbonylamino,
- iv) un groupe -COR₁₆,
- v) un groupe amino,
- vi) un groupe alkyl en C₁ à C₁₀-amino,
- vii) un groupe alcoxy en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène,
- viii) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes alkyle en C₁ à C₁₀ et
- ix) un hétérocycle,

- 3) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino,
- 4) un groupe cycloalkyle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène et un groupe hydroxyle,
- 5) un hétérocycle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un groupe alkyle en C₁ à C₁₀, un atome d'halogène et un groupe arylalkyle en C₁ à C₁₀,
- 6) un groupe hétéroaryle éventuellement substitué par un groupe alkyle en C₁ à C₁₀ et
- 7) un groupe alkyl en C₁ à C₁₀-carbonyle ;

R₃₆ est choisi parmi :

- 1) un atome d'hydrogène et
- 2) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle et aryle ; R₃₅ et R₃₆ peuvent être liés l'un à l'autre pour former un cycle choisi parmi les cycles azétidinyle, pyrrolidinyle, pipéridinyle, pipérazinyle et morpholinyle, et le cycle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment les uns des autres parmi un groupe alkyle en C₁ à C₁₀ et un atome d'halogène ;

R₂₅ est choisi parmi :

- 1) un atome d'halogène,
- 2) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

- i) un atome d'halogène,
- ii) un groupe aryle,
- iii) un groupe hétéroaryle,
- iv) un hétérocycle éventuellement substitué par un ou plusieurs groupes alkyle en C₁ à C₁₀,
- v) un groupe -COR₁₆,
- vi) un groupe -NR₁₃R₁₄ et
- vii) un groupe -SO₂R₂₁,

- 3) un groupe hétéroalkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes hydroxyle,
- 4) un groupe hydroxyalkyle en C₁ à C₁₀, dans lequel chaque groupe hydroxyle peut être indépendamment substitué par un ou plusieurs groupes choisis parmi les groupes alkyle en C₁ à C₁₀, arylalkyle en C₁ à C₁₀ et alkyl en C₁ à C₁₀-carbonyle,
- 5) un groupe -COR₁₆,
- 6) un groupe -SO₂R₂₁,
- 7) un groupe aryle et
- 8) un groupe cyano ;

R₂ est choisi parmi :

- 1) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alkyle est éventuellement en outre substitué par un ou plusieurs substituants choisis indépendamment parmi les groupes R₄₂,
- 2) un groupe alcényle en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alcényle est éventuellement en outre substitué par un ou plusieurs substituants choisis indépendamment parmi les groupes R₄₂,
- 3) un groupe alcynyle en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alcynyle est éventuellement en outre substitué par un ou plusieurs substituants choisis indépendamment parmi les groupes R₄₂,
- 4) un groupe cycloalkyle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

- i) un atome d'halogène,
- ii) un groupe alcényle en C₂ à C₁₀ ou alkyle en C₁ à C₁₀,
- iii) un groupe aryle éventuellement substitué par 1 à 3 substituants choisis indépendamment parmi un groupe alkyle en C₁ à C₁₀, un atome d'halogène, les groupes alcoxy en C₁ à C₁₀, alkyl en C₁ à C₁₀-amino et alkyl en C₁ à C₁₀-carbonyle,
- iv) un groupe cycloalkyle,
- v) un groupe alcényle en C₂ à C₁₀ éventuellement substitué par un atome d'halogène,
- vi) un groupe alkylidène en C₁ à C₁₀, où le groupe alkylidène est lié au groupe cycloalkyle par une double liaison et le groupe alkylidène est éventuellement substitué par un ou plusieurs atomes d'halogène,
- vii) un groupe alcoxy en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène,
- viii) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène ou un groupe alcoxy en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène,
- ix) un groupe alcynyle en C₂ à C₁₀ et
- x) un groupe -Si(R₄₃)₃,

- 5) un hétérocycle, où l'hétérocycle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

- i) un groupe alkyle en C₁ à C₁₀,

ii) un groupe alkyl en C_1 à C_{10} -carbonyle, dans lequel le groupe alkyle est éventuellement substitué par R_{27} ,

iii) un groupe arylcarbonyle, dans lequel le groupe aryle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes alkyle en C_1 à C_{10} et alcoxy en C_1 à C_{10} .

iv) un groupe hétéroarylcabonyle,

v) un groupe alcoxy en C_1 à C_{10} -carbonyle, dans lequel le groupe alkyle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes aryle et alcoxy en C_1 à C_{10} .

vi) un groupe aryloxy carbonyle, dans lequel le groupe aryle est éventuellement substitué par un ou plusieurs atomes d'halogène et/ou un groupe alkyle en C_1 à C_{10} ,

vii) un groupe $-\text{CONR}_{28}\text{R}_{29}$,

viii) un groupe $-\text{SO}_2\text{R}_{21}$,

ix) un atome d'halogène,

x) un groupe cycloalkylcarbonyle éventuellement fusionné avec un groupe aryle et

xi) un groupe alcényle en C_2 à C_{10} -carbonyle, dans lequel le groupe alcényle est éventuellement substitué par un groupe aryle, où le groupe aryle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes alkyle en C_1 à C_{10} ou alcoxy en C_1 à C_{10} ,

6) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes R_{44} ,

7) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

i) un atome d'halogène,

ii) un groupe alkyle en C_1 à C_{10} et

iii) un groupe alcoxy en C_1 à C_{10} ,

8) un groupe alcoxy en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alcoxy est éventuellement en outre substitué par un ou plusieurs substituants choisis indépendamment parmi les groupes R_{42} ,

9) un groupe $-\text{S}(\text{O})_q\text{R}_{43}$ (où q est un nombre entier de 0 à 2) et

10) un groupe cycloalcényle éventuellement substitué par un groupe alkyle en C_1 à C_{10} ;

R_{44} est choisi parmi :

1) un atome d'halogène,

2) un groupe cyano,

3) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

i) un groupe hydroxyle,

ii) un groupe $-\text{OR}_{26}$,

iii) un groupe cyano,

iv) un groupe aryloxy éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène ou un groupe alcoxy en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène et

v) un atome d'halogène,

4) un groupe cycloalkyle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène ou un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène,

5) un groupe alcoxy en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène ou un ou plusieurs groupes alcényle en C_2 à C_6 ,

6) un groupe $-\text{COR}_{30}$,

7) un groupe alkyl en C_1 à C_{10} -carbonylamino,

8) un groupe alcoxy en C_1 à C_{10} -carbonylamino, dans lequel le groupe alcoxy est éventuellement substitué par un groupe aryle,
 9) un groupe hétéroalkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène,
 10) un groupe aryle éventuellement substitué par un ou plusieurs substituants choisis indépendamment
 5 parmi :

- i) un atome d'halogène,
- ii) un groupe alkyle en C_1 à C_{10} ,
- iii) un groupe alcoxy en C_1 à C_{10} et
- iv) un groupe aryle éventuellement substitué par un groupe aryle éventuellement substitué par un
 10 groupe alkyle en C_1 à C_{10} ,

- 11) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes alkyle en C_1 à C_{10} ,
- 12) un groupe $-SO_2R_{43}$,
- 13) un groupe $-SOR_{43}$,
- 14) un groupe alkyl en C_1 à C_{10} -thio éventuellement substitué par un ou plusieurs atomes d'halogène,
- 15) un groupe $-Si(R_{43})_3$ et
- 16) un groupe $-SF_5$;

20 R_{42} est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi
 un groupe alkyle en C_1 à C_{10} éventuellement substitué par un atome d'halogène, un atome d'halogène et
 25 un groupe alcoxy en C_1 à C_{10} ,
- 3) un groupe hydroxycarbonyle,
- 4) un groupe alcoxy en C_1 à C_{10} -carbonyle,
- 5) un groupe aminocarbonyle,
- 6) un groupe alkyl en C_1 à C_{10} -aminocarbonyle,
- 7) un groupe alcoxy en C_1 à C_{10} -carbonylamino,
- 8) un groupe amino,
- 9) un groupe hydroxyle et
- 10) un groupe oxétane, tétrahydrofurane ou tétrahydropyrane éventuellement substitué par un groupe
 30 alkyle en C_1 à C_{10} ;

35 R_{43} représente un groupe alkyle en C_1 à C_{10} ;

R_{26} représente un groupe aryle, ou un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène ;

R_{27} est choisi parmi :

- 1) un groupe aryle éventuellement substitué par un groupe choisi indépendamment parmi un atome
 d'halogène, les groupes alkyle en C_1 à C_{10} et alcoxy en C_1 à C_{10} ,
- 2) un groupe alcoxy en C_1 à C_{10} , dans lequel le groupe alkyle est éventuellement substitué par un groupe
 45 aryle,
- 3) un groupe hydroxyle,
- 4) un groupe amino,
- 5) un groupe alkyl en C_1 à C_{10} -amino,
- 6) un groupe hydroxycarbonyle,
- 7) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment
 50 parmi les groupes alkyle en C_1 à C_{10} et/ou aryle, et
- 8) un groupe hétéroaryloxy ;

R_{28} est choisi parmi un atome d'hydrogène ou un groupe alkyle en C_1 à C_{10} éventuellement substitué par un
 groupe aryle ;

55 R_{29} est choisi parmi un atome d'hydrogène ou un groupe alkyle en C_1 à C_{10} éventuellement substitué par un
 groupe aryle ;

R_{28} et R_{29} peuvent être liés pour former un cycle choisi parmi les cycles azétidinyle, pyrrolidinyle, pipéridinyle,
 pipérazinyle et morpholinyle, et le cycle est éventuellement substitué par un ou plusieurs groupes choisis

indépendamment les uns des autres parmi un groupe alkyle en C₁ à C₁₀ et un atome d'halogène ;

R₃₀ est choisi parmi les groupes-hydroxyle, alcoxy en C₁ à C₁₀ et -NR₃₁R₃₂ ;

R₃₁ et R₃₂ sont choisis indépendamment parmi :

- 1) un atome d'hydrogène,
- 2) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un groupe aryle et
- 3) un groupe aryle ;

R₃₁ et R₃₂ peuvent être liés pour former un cycle choisi parmi les cycles azétidinyle, pyrrolidinyle, pipéridinyle, pipérazinyle et morpholinyle, et le cycle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment les uns des autres parmi un groupe alkyle en C₁ à C₁₀, un atome d'halogène et un groupe alcoxy en C₁ à C₁₀-carbonyle ; et


R₃₃ et R₃₄ sont choisis indépendamment parmi :

- 1) un atome d'hydrogène et
- 2) un groupe alkyle en C₁ à C₁₀ ; et

[où dans la formule générale (2)

W, Y, m, n, R₄, R₅, R₆, R₇, R₁₁, R₁₆, R₁₉, R₂₀, R₂₁, R₃₃, R₃₄ et R₄₄ sont tels que définis pour la formule (1) ;

U représente une liaison, un groupe alkylène en C₁ à C₁₀ ou tout groupe choisi parmi des groupes représentés par la formule suivante :




A est choisi parmi un atome O, les groupes NH et CH₂ ;

R₄₆ est choisi parmi un atome d'hydrogène ou R₄₄ ;

T est choisi parmi les groupes aryle et hétéroaryle ;

V est choisi parmi :




E représente un hétérocycle de 4 à 7 chaînons contenant éventuellement un ou plusieurs éléments ou groupes supplémentaires choisis parmi les atomes O, N, S, les groupes SO et SO₂, et l'hétérocycle est éventuellement substitué par un ou plusieurs groupes choisis parmi :

- 1) un atome d'hydrogène,
- 2) un atome d'halogène,
- 3) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un groupe alkyl en C₁ à C₁₀-amino, un atome d'halogène et un groupe hydroxyle,
- 4) un groupe hydroxyle,
- 5) un groupe alcoxy en C₁ à C₁₀ éventuellement, substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino,

- 6) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino,
 7) un groupe hétéroalkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C₁ à C₁₀-amino,
 8) un hétérocycle éventuellement substitué par un groupe alkyle en C₁ à C₁₀,
 9) un groupe hétéroaryle éventuellement substitué par un groupe alkyle en C₁ à C₁₀,
 10) un groupe hétérocyclyalkyle en C₁ à C₁₀,
 11) un groupe -COR₁₆,
 12) un groupe -NR₁₉R₂₀ et
 13) un groupe -SO₂R₂₁ ;

Z représente un groupe bivalent choisi parmi :

- 1) un groupe alkylène en C₁ à C₁₀ ou hétéroalkylène en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène et/ou un ou plusieurs groupes hydroxyle, où les un ou plusieurs atomes de carbone peuvent être oxydés pour former un groupe carbonyle ;
 2) un groupe alcénylène en C₂ à C₁₀ ou hétéroalcénylène en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène et/ou un ou plusieurs groupes hydroxyle, où les un ou plusieurs atomes de carbone peuvent être oxydés pour former un groupe carbonyle ; et
 3) un groupe choisi parmi :



G représente un groupe bivalent choisi parmi :

- 1) un groupe alkylène en C₁ à C₁₀ ou hétéroalkylène en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène ; et
 2) un groupe alcénylène en C₂ à C₁₀ ou hétéroalcénylène en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène ;

J représente un groupe bivalent choisi parmi :

- 1) un groupe alkylène en C₁ à C₁₀ ou hétéroalkylène en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène ; et
 2) un groupe alcénylène en C₂ à C₁₀ ou hétéroalcénylène en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène ;

B est choisi parmi un hétérocycle ou un groupe hétéroaryle ; et

R₄₅ est choisi parmi un atome d'hydrogène ou un groupe alkyle en C₁ à C₁₀ ;

et où dans le composé représenté par les formules générales (1) ou (2) le terme « hétéroalkyle » signifie un groupe contenant 1 à 5 hétéroatomes dans le squelette alkyle, le terme « hétéroalkylène » signifie un groupe contenant 1 à 5 hétéroatomes dans le squelette alkylène, et le terme « hétéroalcénylène » signifie un groupe contenant 1 à 5 hétéroatomes dans le squelette alcénylène.

2. Composé ou sel pharmacologiquement acceptable de celui-ci selon la revendication 1, dans lequel

W est choisi parmi :

- 1) une liaison simple,
 2) un groupe alkylène en C₁ à C₁₀ contenant éventuellement un groupe carbonyle, où le groupe alkylène est éventuellement substitué par un ou plusieurs atomes d'halogène ou un groupe hydroxy,
 3) un groupe alcénylène en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène,

- 4) un groupe alcynylène en C_2 à C_{10} ,
- 5) un groupe arylène,
- 6) un groupe hétéroarylène,
- 7) un groupe $-NH-$, $-NHCH_2-$ ou $-NHCH_2CH_2-$,
- 8) un groupe cycloalkylène et
- 9) un groupe $-(cycloalkylène)-CH_2-$;

X est choisi parmi la liaison ou les groupes suivants :

- 1) une liaison simple,
- 2) un groupe alkylène en C_1 à C_{10} éventuellement substitué par un groupe cycloalkyle,
- 3) un groupe alcénylène en C_2 à C_{10} ,
- 4) un groupe alcynylène en C_2 à C_{10} et
- 5) un groupe oxyalkylène en C_1 à C_{10} ;

R_1 est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe cycloalkyle éventuellement substitué par un groupe choisi parmi les groupes R_4 ,
- 3) un hétérocycle éventuellement substitué par un ou plusieurs groupes choisis parmi R_{25} et R_4 ,
- 4) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis parmi R_3 , R_4 , R_5 et R_6 , et
- 5) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes choisis parmi R_{25} , R_4 et R_5 ;

R_9 est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes R_{23} ,
- 3) un groupe cycloalkyle éventuellement substitué par un ou plusieurs atomes d'halogène ou un ou plusieurs groupes hydroxyle,
- 4) un hétérocycle éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes alkyle en C_1 à C_{10} , alkyl en C_1 à C_{10} -carbonyle, alcoxy en C_1 à C_{10} , alcoxy en C_1 à C_{10} -carbonyle, amino et un atome d'halogène,
- 5) un groupe hétéroalkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène et un groupe hydroxyle,
- 6) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyle en C_1 à C_{10} , alkyl en C_1 à C_{10} -carbonyle, alcoxy en C_1 à C_{10} -carbonyle et un atome d'halogène et
- 7) un groupe cycloalcényle éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes alcoxy en C_1 à C_{10} , alkyl en C_1 à C_{10} -amino, amino, 1 à 3 groupes hydroxyle et 1 à 4 atomes d'halogène, où le groupe cycloalcényle contient éventuellement un groupe carbonyle ;

R_{10} est choisi parmi :

- 1) un atome d'hydrogène et
- 2) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle et aryle ; R_9 et R_{10} peuvent être liés pour former un hétérocycle de 4 à 7 chaînons contenant éventuellement un ou plusieurs éléments ou groupes supplémentaires choisis indépendamment parmi les atomes N, O, S, les groupes SO, SO_2 , carbonyle et thiocarbonyle, et l'hétérocycle est éventuellement substitué par un ou plusieurs substituants choisis indépendamment parmi les groupes R_{24} ;

R_{24} est choisi parmi :

- 1) un atome d'halogène,
- 2) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyl en C_1 à C_{10} -amino et alkyl en C_1 à C_{10} -carbonylamino,
- 3) un groupe halogénoalkyle en C_1 à C_{10} ,

- 4) un groupe hydroxyle,
 5) un groupe hydroxyalkyle en C_1 à C_{10} ,
 6) un groupe alcoxy en C_1 à C_{10} éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C_1 à C_{10} -amino,
 7) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C_1 à C_{10} -amino,
 8) un groupe hétéroalkyle en C_1 à C_{10} éventuellement substitué par un ou deux groupes choisis parmi un atome d'halogène, les groupes hydroxyle, amino et alkyl en C_1 à C_{10} -amino,
 9) un groupe $-COR_{16}$ et
 10) un groupe $-NR_{19}R_{20}$;

R_{11} est choisi parmi :

- 1) un groupe alkyle en C_1 à C_{10} éventuellement substitué par 1 à 3 substituants choisis indépendamment parmi :
- i) un groupe hydroxyle,
 - ii) un groupe $-NR_{17}R_{18}$,
 - iii) un groupe alcoxy en C_1 à C_{10} ,
 - iv) un atome d'halogène,
 - v) un groupe alcoxy en C_1 à C_{10} -carbonyle et
 - vi) un groupe aminocarbonyle,
- 2) un groupe aryle,
 3) un groupe arylalkyle en C_1 à C_{10} ,
 4) un groupe cycloalkyle éventuellement substitué par un ou plusieurs atomes d'halogène,
 5) un hétérocycle éventuellement substitué par un groupe alkyle en C_1 à C_{10} ,
 6) un groupe alcoxy en C_1 à C_{10} , dans lequel le groupe alkyle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyl en C_1 à C_{10} -carbonylamino, amino, alkyl en C_1 à C_{10} -amino et hydroxyle,
 7) un groupe amino,
 8) un groupe alkyl en C_1 à C_{10} -amino, dans lequel le groupe alkyle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyl en C_1 à C_{10} -carbonylamino, amino, alkyl en C_1 à C_{10} -amino, hydroxycarbonyle et hydroxyle et
 9) un groupe alcényle en C_2 à C_{10} ; et

R_{33} et R_{34} représentent un atome d'hydrogène,
 où $R_3, R_4, R_5, R_6, R_{16}, R_{17}, R_{18}, R_{19}, R_{20}, R_{23}$ et R_{25} sont tels que définis dans la revendication 1, respectivement.

3. Composé ou sel pharmacologiquement acceptable de celui-ci selon la revendication 1 ou 2, où

W est choisi parmi :

- 1) une liaison simple,
- 2) un groupe alkylène en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène,
- 3) un groupe alcénylène en C_2 à C_{10} ,
- 4) un groupe alcynylène en C_2 à C_{10} et
- 5) un groupe hétéroarylène,

X est choisi parmi la liaison ou les groupes suivants :

- 1) une liaison simple,
- 2) un groupe alkylène en C_1 à C_{10} ,
- 3) un groupe alcénylène en C_2 à C_{10} ,
- 4) un groupe alcynylène en C_2 à C_{10} et
- 5) un groupe oxyalkylène en C_1 à C_{10} , où le groupe oxyalkylène est lié à un cycle 1,3,8-triaza-spiro[4.5]déc-1-én-4-one ou à un cycle 1,3,8-triaza-spiro[4.5]déc-1-ène-4-thione par l'intermédiaire d'un atome de carbone dans le groupe oxyalkylène ;

R₁ est choisi parmi :

- 1) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis parmi R₃, R₄ et R₅ et
- 2) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes choisis parmi R₂₅ et R₄ ;

R₃ est choisi parmi :

- 1) un groupe -CONR₇R₈,
- 2) un groupe -OR₉,
- 3) un groupe -NR₉R₁₀,
- 4) un groupe -N(R₉)COR₁₁,
- 5) un groupe N(R₉)SO₂R₁₂,
- 6) un groupe -SO₂R₁₅,
- 7) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes -COR₁₆ et -NR₁₃R₁₄ et
- 8) un groupe -N(R₉)CSNH₂ ;

R₄ est choisi parmi :

- 1) un atome d'halogène,
- 2) un groupe cyano,
- 3) un groupe amino,
- 4) un groupe alkyle en C₁ à C₁₀,
- 5) un groupe halogénoalkyle en C₁ à C₁₀,
- 6) un groupe alcoxy en C₁ à C₁₀,
- 7) un groupe halogénoalkyl en C₁ à C₁₀-carbonyle,
- 8) un groupe -COR₁₆ et
- 9) un groupe hétéroalkyle en C₁ à C₁₀ ;

R₂ est choisi parmi :

- 1) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alkyle est éventuellement en outre substitué par un groupe choisi parmi les groupes R₄₂,
- 2) un groupe alcényle en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alcényle est éventuellement en outre substitué par un groupe choisi parmi les groupes R₄₂,
- 3) un groupe alcynyle en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alcynyle est éventuellement en outre substitué par un groupe choisi parmi les groupes R₄₂,
- 4) un groupe cycloalkyle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

- i) un atome d'halogène,
- ii) un groupe alcényle en C₂ à C₁₀ ou alkyle en C₁ à C₁₀,
- iii) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un groupe alkyle en C₁ à C₁₀, un atome d'halogène, et un groupe alcoxy en C₁ à C₁₀,
- iv) un groupe cycloalkyle,
- v) un groupe halogénoalcényle en C₂ à C₁₀ ou halogénoalkyle en C₁ à C₁₀,
- vi) un groupe alkylidène en C₁ à C₁₀, où le groupe alkylidène est lié au groupe cycloalkyle par une double liaison et le groupe alkylidène est éventuellement substitué par un ou plusieurs atomes d'halogène,
- vii) un groupe alcoxy en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène,
- viii) un groupe alkyle en C₁ à C₁₀ substitué par un groupe alcoxy en C₁ à C₁₀, où le groupe alkyle et/ou le groupe alkyle dans le groupe alcoxy est éventuellement substitué par un ou plusieurs atomes d'halogène,
- ix) un groupe alcynyle en C₂ à C₁₀ et
- x) un groupe -Si(R₄₃)₃,

- 5) un hétérocycle, où l'hétérocycle est éventuellement substitué par un ou plusieurs groupes choisis parmi :

- i) un groupe alkyle en C_1 à C_{10} ,
- ii) un groupe alkyl en C_1 à C_{10} -carbonyle, dans lequel le groupe alkyle est éventuellement substitué par R_{27} ,
- iii) un groupe arylcarbonyle, dans lequel le groupe aryle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes alkyle en C_1 à C_{10} et alcoxy en C_1 à C_{10} ,
- iv) un groupe hétéroarylcabonyle,
- v) un groupe alcoxy en C_1 à C_{10} -carbonyle, dans lequel le groupe alkyle est éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes aryle et alcoxy en C_1 à C_{10} ,
- vi) un groupe aryloxy carbonyle, dans lequel le groupe aryle est éventuellement substitué par un ou plusieurs atomes d'halogène et/ou un groupe alkyle en C_1 à C_{10} ,
- vii) un groupe $-\text{CONR}_{28}\text{R}_{29}$ et
- viii) un groupe $-\text{SO}_2\text{R}_{21}$,

6) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes R_{44} ,

7) un groupe hétéroaryle éventuellement substitué par un des groupes suivants :

- i) un groupe alkyle en C_1 à C_{10} ,

8) un groupe alcoxy en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alcoxy est éventuellement en outre substitué par un groupe choisi parmi les groupes R_{42} ,

9) un groupe $-\text{S}(\text{O})_q\text{R}_{43}$ (où q est un nombre entier de 0 à 2) et

10) un groupe cycloalcényle éventuellement substitué par un groupe alkyle en C_1 à C_{10} ,

R_{44} est choisi parmi :

1) un atome d'halogène,

2) un groupe cyano,

3) un groupe alkyle en C_1 à C_{10} éventuellement substitué par un des groupes suivants :

- i) un groupe hydroxyle,

- ii) un groupe $-\text{OR}_{26}$,

- iii) un groupe cyano et

- iv) un groupe aryloxy éventuellement substitué par un ou plusieurs groupes choisis parmi un atome d'halogène, un groupe alkyle en C_1 à C_{10} , halogénoalkyle en C_1 à C_{10} ou halogénoalcoxy en C_1 à C_{10} ,

4) un groupe halogénoalkyle en C_1 à C_{10} ,

5) un groupe cycloalkyle éventuellement substitué par un ou plusieurs groupes choisis parmi un atome d'halogène et un groupe halogénoalkyle en C_1 à C_{10} ,

6) un groupe alcoxy en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène ou un ou plusieurs groupes alcényle en C_2 à C_6 ,

7) un groupe $-\text{COR}_{30}$,

8) un groupe hétéroalkyle en C_1 à C_{10} éventuellement substitué par un ou plusieurs atomes d'halogène,

9) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

- i) un groupe alkyle en C_1 à C_{10} et

- iii) un groupe aryle,

10) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes alkyle en C_1 à C_{10} ,

11) un groupe $-\text{SO}_2\text{R}_{43}$,

12) un groupe alkyl en C_1 à C_{10} -thio éventuellement substitué par un ou plusieurs atomes d'halogène,

13) un groupe $-\text{Si}(\text{R}_{43})_3$ et

14) un groupe $-\text{SF}_5$ et

R_{27} est choisi parmi :

- 1) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi un atome d'halogène, les groupes alkyle en C₁ à C₁₀ et alcoxy en C₁ à C₁₀,
 2) un groupe alcoxy en C₁ à C₁₀, dans lequel le groupe alkyle est éventuellement substitué par un groupe aryle,
 3) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi les groupes alkyle en C₁ à C₁₀ et aryle et
 4) un groupe hétéroaryloxy,

où R₅, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₂₁, R₂₅, R₂₆, R₂₈, R₂₉, R₃₀, R₄₂ et R₄₃ sont tels que définis dans les revendications 1 ou 2 dont la revendication 3 dépend, respectivement.

4. Composé ou sel pharmacologiquement acceptable de celui-ci selon l'une des revendications 1 à 3, où

W est choisi parmi :

- 1) un groupe alkylène en C₁ à C₆ éventuellement substitué par un ou plusieurs atomes de fluor,
 2) un groupe alcénylène en C₁ à C₆ et
 3) un groupe thiophène,

X est choisi parmi la liaison ou les groupes suivants :

- 1) une liaison simple,
 2) un groupe alkylène en C₁ à C₆ et
 3) un groupe oxyalkylène en C₁ à C₆ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe oxyalkylène est lié à un cycle 1,3,8-triaza-spiro[4.5]déc-1-én-4-one ou à un cycle 1,3,8-triaza-spiro[4.5]déc-1-ène-4-thione par l'intermédiaire d'un atome de carbone dans le groupe oxyalkylène ;

Y représente un atome d'oxygène ;

m représente 1 ; et

n représente 1.

5. Composé ou sel pharmacologiquement acceptable de celui-ci selon l'une des revendications 1 à 4, où

W est choisi parmi :

- 1) un groupe éthylène,
 2) un groupe vinylène et
 3) un groupe thiophène,

X représente une liaison simple ;

R₃ est choisi parmi :

- 1) un groupe -CONR₇R₈,
 2) un groupe -OR₉,
 3) un groupe -NR₉R₁₀,
 4) un groupe -N(R₉)COR₁₁,
 5) un groupe -N(R₉)SO₂R₁₂,
 6) un groupe -SO₂R₁₅ et
 7) un groupe alkyle en C₁ à C₆ éventuellement substitué par un ou plusieurs groupes choisis parmi les groupes -COR₁₆ et -NR₁₃R₁₄ ;


R₂ est choisi parmi :

- 1) un groupe alkyle en C₁ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alkyle est éventuellement en outre substitué par un groupe choisi parmi les groupes R₄₂,
 2) un groupe alcényle en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le groupe alcényle est éventuellement en outre substitué par un groupe choisi parmi les groupes R₄₂,
 3) un groupe alcynyle en C₂ à C₁₀ éventuellement substitué par un ou plusieurs atomes d'halogène, où le

groupe alcynyle est éventuellement en outre substitué par un groupe choisi parmi les groupes R_{42} ,
4) un groupe cycloalkyle éventuellement substitué par un ou plusieurs groupes choisis indépendamment
parmi :

- i) un atome d'halogène,
- ii) un groupe alcényle en C_2 à C_6 ou alkyle en C_1 à C_6 ,
- iii) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment
parmi un groupe alkyle en C_1 à C_6 , un atome d'halogène, les groupes alcoxy en C_1 à C_6 , alkyl en C_1
à C_6 -amino et alkyl en C_1 à C_6 -carbonyle,
- iv) un groupe cycloalkyle,
- v) un groupe halogénoalcényle en C_2 à C_6 ou halogénoalkyle en C_1 à C_6 ,
- vi) un groupe alkylidène en C_1 à C_6 , où le groupe alkylidène est lié au groupe cycloalkyle par une
double liaison et le groupe alkylidène est éventuellement substitué par un ou plusieurs atomes d'halogène,
- vii) un groupe alcoxy en C_1 à C_6 éventuellement substitué par un ou plusieurs atomes d'halogène,
- viii) un groupe alkyle en C_1 à C_6 substitué par un groupe alcoxy en C_1 à C_6 , où le groupe alkyle et/ou
le groupe alkyle dans le groupe alcoxy est éventuellement substitué par un atome d'halogène,
- ix) un groupe alcynyle en C_2 à C_6 et
- x) un groupe $-Si(R_{43})_3$,

5) un groupe représenté par la formule générale (B) suivante :



(où Ra représente un groupe choisi parmi :

- i) un groupe alkyl en C_1 à C_6 -carbonyle, dans lequel le groupe alkyle est éventuellement substitué par
 R_{27} ,
- ii) un groupe arylcarbonyle, dans lequel le groupe aryle est éventuellement substitué par un ou plusieurs
groupes choisis indépendamment parmi un atome d'halogène, les groupes alkyle en C_1 à C_6 et alcoxy
en C_1 à C_6 ,
- iii) un groupe alcoxy en C_1 à C_6 -carbonyle, dans lequel le groupe alkyle est éventuellement substitué
par un ou plusieurs groupes choisis parmi un atome d'halogène, les groupes aryle et alcoxy en C_1 à C_6 ,
- iv) un groupe aryloxy carbonyle, dans lequel le groupe aryle est éventuellement substitué par un ou
plusieurs atomes d'halogène ou un groupe alkyle en C_1 à C_6 ,
- v) un groupe $-CONR_{28}R_{29}$ et
- vi) un groupe $-SO_2R_{21}$,

6) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi
les groupes R_{44} ,

7) un groupe hétéroaryle éventuellement substitué par l'un quelconque des groupes suivants :

- i) un atome d'halogène,
- ii) un groupe alkyle en C_1 à C_6 et
- iii) un groupe alcoxy en C_1 à C_6 ;

8) un groupe alcoxy en C_1 à C_6 éventuellement substitué par un ou plusieurs atomes d'halogène, où le
groupe alcoxy est éventuellement en outre substitué par un groupe choisi parmi les groupes R_{42} ,

9) un groupe $-S(O)_qR_{43}$ (où q est un nombre entier de 0 à 2) et

10) un groupe cycloalcényle éventuellement substitué par un groupe alkyle en C_1 à C_6 ; et

R_{44} est choisi parmi :

- 1) un atome d'halogène,
- 2) un groupe cyano,
- 3) un groupe alkyle en C₁ à C₆ éventuellement substitué par l'un quelconque des groupes suivants :

- i) un groupe hydroxyle,
- ii) un groupe -OR₂₆,
- iii) un groupe cyano et
- iv) un groupe aryloxy éventuellement substitué par un ou plusieurs groupes choisis parmi un atome d'halogène, les groupes alkyle en C₁ à C₆; halogénoalkyle en C₁ à C₆ ou halogénoalcoxy en C₁ à C₆,


- 4) un groupe halogénoalkyle en C₁ à C₆,
- 5) un groupe cycloalkyle éventuellement substitué par un ou plusieurs groupes choisis parmi un atome d'halogène et un groupe halogénoalkyle en C₁ à C₆,
- 6) un groupe alcoxy en C₁ à C₆ éventuellement substitué par un ou plusieurs atomes d'halogène,
- 7) un groupe -COR₃₀,
- 8) un groupe hétéroalkyle en C₁ à C₆ éventuellement substitué par un ou plusieurs atomes d'halogène,
- 9) un groupe aryle éventuellement substitué par un ou plusieurs groupes choisis indépendamment parmi :

- i) un groupe alkyle en C₁ à C₆ et
- ii) un groupe aryle,

- 10) un groupe hétéroaryle éventuellement substitué par un ou plusieurs groupes alkyle en C₁ à C₆,
- 11) un groupe -SO₂R₄₃,
- 12) un groupe alkyl en C₁ à C₆-thio éventuellement substitué par un ou plusieurs atomes d'halogène,
- 13) un groupe -Si(R₄₃)₃ et
- 14) un groupe -SF₅,

où R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₂₁, R₂₆, R₂₇, R₂₈, R₂₉, R₃₀, R₄₂ et R₄₃ sont tels que définis dans les revendications 1 à 4 dont la revendication 5 dépend, respectivement.


6. Composé ou sel pharmacologiquement acceptable de celui-ci selon la revendication 3, dans lequel R₁ est un groupe représenté par la formule générale (3) suivante :



(3)

dans laquelle R₃, R₄ et R₅ sont tels que définis pour R₃, R₄ et R₅ dans la revendication 3.


7. Composé ou sel pharmacologiquement acceptable de celui-ci selon la revendication 3, dans lequel R₁ est un groupe représenté par la formule générale (4) suivante :



(4)

dans laquelle R_4 et R_{25} sont tels que définis pour R_4 et R_{25} dans la revendication 3.


8. Composé ou sel pharmacologiquement acceptable de celui-ci selon la revendication 5, dans lequel R_1 est un groupe représenté par la formule générale (3) suivante :



(3)

dans laquelle R_3 , R_4 et R_5 sont tels que définis pour R_3 , R_4 et R_5 dans la revendication 5.

9. Composé ou sel pharmacologiquement acceptable de celui-ci selon la revendication 5, dans lequel R_1 est un groupe représenté par la formule générale (4) suivante :



(4)

dans laquelle R_4 et R_{25} sont tels que définis pour R_4 et R_{25} dans la revendication 5.

10. Composé selon la revendication 1, dans laquelle le composé est:

- (1) 8-(3-chloro-benzènesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (2) 8-(3-chloro-benzènesulfonyl)-2-pyridin-4-yl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (3) 8-(3-chloro-benzènesulfonyl)-2-propyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (4) 8-(3-chloro-benzènesulfonyl)-2-isopropyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (5) 8-(3-chloro-benzènesulfonyl)-2-(3-méthoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (6) 8-(3-chloro-benzènesulfonyl)-2-(3-chloro-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (7) 2-benzyl-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 5 (8) 8-(3-chloro-benzènesulfonyl)-2-méthyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (10) 2-biphényl-2-yl-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (11) 8-(3-chloro-benzènesulfonyl)-2-o-tolyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (12) 8-(3-chloro-benzènesulfonyl)-2-(4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (13) 8-(3-chloro-benzènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 10 (14) 8-(3-chloro-benzènesulfonyl)-2-(1-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (15) 2-(1-acetyl-pipéridin-4-yl)-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (16) 2-tert-butyl-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (17) 8-(3-chloro-benzènesulfonyl)-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro [4.5]déc-1-én-4-one;
- (18) ester tert-butyle de l'acide 3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pi-
- 15 péridine-1-carboxylique;
- (19) 8-(3-chloro-benzènesulfonyl)-2-(3-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (20) 8-(3-chloro-benzènesulfonyl)-2-(2-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (21) 8-(3-chloro-benzènesulfonyl)-2-(4-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (22) 2-(1R,2S,4S)-bicyclo[2.2.1]hept-2-yl-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 20 (23) 2-(1R,2R,4S)-bicyclo[2.2.1]hept-2-yl-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (24) 8-(3-chloro-benzènesulfonyl)-2-(4-méthoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (25) 8-(3-chloro-benzènesulfonyl)-2-(4-méthoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (26) 2-(4-tert-butyl-cyclohexyl)-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (27) 8-(3-chloro-benzènesulfonyl)-2-(4-fluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 25 (28) 8-(3-chloro-benzènesulfonyl)-2-cyclohexylméthyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (29) 8-(3-chloro-benzènesulfonyl)-2-phénéthyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (30) 8-(3-chloro-benzènesulfonyl)-2-(2-cyclohexyl-éthyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (31) 8-(3-chloro-benzènesulfonyl)-2-(3-trifluorométhyl-phenyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (32) 8-(3-chloro-benzènesulfonyl)-2-(3-méthanesulfonyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 30 (33) 8-(3-chloro-benzènesulfonyl)-2-(1-phényl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (34) 8-(3-chloro-benzènesulfonyl)-2-(2-naphthalèn-1-yl-éthyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (35) 8-(2-naphthalèn-1-yl-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (36) 2-tert-butyl-8-(2-naphthalèn-1-yl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (37) 8-(2-naphthalèn-1-yl-éthanesulfonyl)-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-
- 35 one;
- (38) ester tert-butyle de l'acide 3-[8-(2-naphthalèn-1-yl-éthanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- (39) 8-(2-naphthalèn-1-yl-éthanesulfonyl)-2-[1-(2-naphthalèn-1-yl-éthanesulfonyl)-pipéridin-3-yl]-1,3,8-triaza-
- 40 spiro[4.5]déc-1-én-4-one;
- (40) 8-(3-chloro-benzènesulfonyl)-2-[1-(propane-1-sulfonyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-
- one;
- (41) 2-(1-benzènesulfonyl-pipéridin-3-yl)-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (42) 8-(3-chloro-benzènesulfonyl)-2-(1-phénylméthanesulfonyl-pipéridin-3-yl)-1,3,8-triaza-spiro[4.5]déc-1-én-
- 4-one;
- 45 (43) 8-(2-naphthalèn-1-yl-éthanesulfonyl)-2-(1-phénylméthanesulfonyl-pipéridin-3-yl)-1,3,8-triaza-
- spiro[4.5]déc-1-én-4-one;
- (44) 8-(4-chloro-benzènesulfonyl)-2-(2,4-dichloro-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (45) 2-(2,4-dichloro-phényl)-8-(2-trifluorométhyl-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (46) 8-(butane-1-sulfonyl)-2-(2,4-dichloro-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 50 (47) 2-(2,4-dichloro-phényl)-8-(2-naphthalèn-1-yl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (48) 2-(2,4-dichloro-phényl)-8-(quinoline-8-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (49) 8-(3-chloro-4-fluoro-benzènesulfonyl)-2-(2,4-dichloro-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (50) 8-(3-chloro-benzènesulfonyl)-2-(2,4-dichloro-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (51) ester méthyle de l'acide 2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-benzoïque;
- 55 (52) 2-cyclohexyl-8-(5-méthyl-3-phényl-isoxazole-4-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (53) 8-(benzo[b]thiophène-3-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (54) 8-(benzo[b]thiophène-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (55) 8-(5-chloro-thiophène-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (56) 2-cyclohexyl-8-(thiophène-2-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (57) 2-cyclohexyl-8-(naphthalène-1-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (58) 2-cyclohexyl-8-(2,4-diméthyl-thiazole-5-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (59) 2-cyclohexyl-8-(3,5-diméthyl-isoxazole-4-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (60) 2-cyclohexyl-8-(2,3-dihydro-benzo[1,4]dioxine-6-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (61) 2-cyclohexyl-8-(2-naphthalèn-1-yl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (62) ester méthyle de l'acide 3-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-thiophène-2-carboxylique;
- (63) ester méthyle de l'acide 5-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-4-méthyl-thiophène-2-carboxylique;
- (64) 2-cyclohexyl-8-(2,5-diméthyl-thiophène-3-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (65) 8-(5-bromo-thiophène-2-sulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (66) 8-(5-chloro-thiophène-2-sulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (67) 8-(naphthalène-2-sulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (68) 8-(benzo[b]thiophène-2-sulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (69) 8-(5-bromo-thiophène-2-sulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (70) 8-(3-chloro-benzènesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.6]undéc-1-én-4-one;
- (71) 7-(3-chloro-benzènesulfonyl)-2-cyclohexyl-1,3,7-triaza-spiro[4.5]déc-1-én-4-one;
- (72) 7-(3-chloro-benzènesulfonyl)-2-cyclohexyl-1,3,7-triaza-spiro[4.4]non-1-én-4-one;
- (73) 4-{2-[2-(2,4-dichloro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (74) 4-[2-(2-tert-butyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,N,N-triméthylbenzamide;
- (75) 3,N,N-triméthyl-4-{2-[4-oxo-2-(4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (76) 3,N,N-triméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (77) 4-[2-(2-cyclopentyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl)-éthyl]-3,N,N-triméthylbenzamide;
- (78) 4-{2-[2-(2,6-difluoro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (79) 4-{2-[2-(2,6-diméthoxy-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (80) 4-{2-[2-(3-méthoxy-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (81) 4-{2-[2-(3-chloro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (82) 4-{2-[2-(3,5-bis-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (83) 4-[2-(2-benzyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,N,N-triméthylbenzamide;
- (84) 3,N,N-triméthyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-benzamide;
- (85) 4-{2-[2-(2-chloro-5-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (86) 3,N,N-triméthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (87) 4-{2-[2-(4-chloro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (88) 4-{2-[2-(2,3-dichloro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (89) 4-[2-[2-(3-chloro-4-fluoro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl]-éthyl]-3,N,N-triméthylbenzamide;
- (90) 4-[2-[2-(2-chloro-4-fluoro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3,N,N-triméthylbenzamide;
- (91) 4-{2-[2-(3-bromo-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (92) 4-{2-[2-(2,2-difluoro-benzo[1,3]dioxol-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthylbenzamide;
- (93) 4-[2-[2-(3-chloro-2-fluoro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3,N,N-triméthylbenzamide;
- (94) 3,N,N-triméthyl-4-(2-{4-oxo-2-[4-(1,1,2,2-tetrafluoro-éthoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-

sulfonyl)-éthyl)-benzamide;

(95) 4-{2-[2-(4-chloro-3-méthyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(96) 4-{2-[2-(3-fluoro-4-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(97) 3,N,N-triméthyl-4-{2-[2-(6-méthyl-pyridin-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(98) 4-{2-[2-(3,4-dichloro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(99) 3,N,N-triméthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(100) 4-{2-[2-(2,4-bis-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(101) 3,N,N-triméthyl-4-{2-(4-oxo-2-phénéthyl-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl)-éthyl}-benzamide;

(102) 4-{2-[2-(2,4-diméthyl-thiazol-5-yl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(103) 4-{2-(2-cyclohexylméthyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl}-3,N,N-triméthyl-benzamide;

(104) 3,N,N-triméthyl-4-{2-[4-oxo-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(105) 3,N,N-triméthyl-4-{2-[4-oxo-2-(2-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(106) 3,N,N-triméthyl-4-{2-[4-oxo-2-(4-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(107) 4-{2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(108) 4-{2-[2-(2-cyclohexyl-éthyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(109) 4-{2-[2-(3-fluoro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(110) 4-{2-[2-(3-methanesulfonyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(111) 3,N,N-triméthyl-4-{2-[2-(2-méthyl-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(112) 3,N,N-triméthyl-4-{2-[2-(2-méthyl-5-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(113) 4-{2-[2-(2,3-diméthyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(114) 4-{2-[2-(3-fluoro-2-méthyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(115) 4-{2-[2-(3-fluoro-5-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(116) 4-{2-[2-(2-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(117) 4-{2-[2-(4-fluoro-3-méthyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(118) 4-{2-[2-(4-difluorométhoxy-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(119) 4-{2-[2-(2-méthoxy-pyridin-4-yl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(120) 3,N,N-triméthyl-4-{2-[2-(5-méthyl-pyrazin-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(121) 3,N,N-triméthyl-4-{2-(4-oxo-2-thiazol-4-yl-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl}-benzamide;

(122) 3,N,N-triméthyl-4-{2-[2-(1-méthyl-1H-imidazol-2-yl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(123) 3,N,N-triméthyl-4-{2-[4-oxo-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(124) 4-{2-[2-(4-chloro-phénoxy-méthyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

thyl-benzamide;

(125) 4-{2-[2-(4-fluoro-phénoxy-méthyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(126) 4-{2-[2-(4-chloro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(127) 4-{2-[2-(3-chloro-5-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(128) 4-{2-[2-(4-chloro-2-méthyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(129) 4-{2-[2-(4-chloro-2-fluoro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(130) 4-{2-[2-(3-isopropoxyméthyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(131) 4-{2-[2-(2,4-dichloro-phénoxy-méthyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(133) 4-{2-[2-(3-chloro-2-méthyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(134) 4-{2-[2-(2,4-dichloro-benzyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(135) 3,N,N-triméthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-benzyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(136) 3,N,N-triméthyl-4-{2-[4-oxo-2-(1-trifluorométhyl-cyclopropyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(137) 3,N,N-triméthyl-4-{2-[4-oxo-2-(1-trifluorométhyl-cyclobutyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(138) 3,N,N-triméthyl-4-{2-[4-oxo-2-(1-trifluorométhyl-cyclopentyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(139) 4-{2-[2-(2-fluoro-4-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(140) 3,N,N-triméthyl-4-{2-[4-oxo-2-[3-(1,1,2,2-tetrafluoro-éthoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(141) 3,N,N-triméthyl-4-{2-[4-oxo-2-((E)-propényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(142) 3,N,N-triméthyl-4-{2-[4-oxo-2-((E)-styryl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(143) 4-{2-[2-(benzothiazol-6-yl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(144) 4-{2-[2-(4-méthoxy-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(145) N-{3-méthyl-4-[2-(4-oxo-2-m-tolyl-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-acétamide;

(146) N-(4-{2-[2-(2,3-diméthyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-acétamide;

(147) N-(4-{2-[2-(2,3-dichloro-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-acétamide;

(148) N-(3-méthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;

(149) N-(2-hydroxy-éthyl)-N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;

(150) ester (S)-1-acetoxyméthyl-2-[acetyl-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-amino]-éthyle de l'acide acétique;

(151) ester (S)-1-acetoxyméthyl-2-[acetyl-(3-méthyl-4-{2-[4-oxo-2-(4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-amino]-éthyle de l'acide acétique;

(152) ester (S)-1-acetoxyméthyl-2-[acetyl-(3-méthyl-4-{2-[4-oxo-2-(4-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-amino]-éthyle de l'acide acétique;

(153) ester (S)-1-acetoxyméthyl-2-[acetyl-(4-{2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-amino]-éthyle de l'acide acétique;

(154) 8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-méthyl-phényl]-éthanesulfonyl}-2-(4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(155) 8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-

- 1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (156) 8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-méthyl-phényl]-éthanesulfonyl}-2-(4-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (157) 2-(3-chloro-phényl)-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-méthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (158) 2-(4-chloro-phényl)-8-{2-[4-((S)-2,3-dihydroxy-propylamino)-2-méthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (159) 8-{2-[4-(2-hydroxy-éthylamino)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (160) 8-{2-[4-(2-hydroxy-éthylamino)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (161) 8-{2-[4-(2-hydroxy-éthylamino)-2-méthyl-phényl]-éthanesulfonyl}-2-(4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (162) ester tert-butyle de l'acide {4-[2-(2-tert-butyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-méthyl-benzyl-carbamique};
- (163) ester tert-butyle de l'acide (4-{2-[2-(4,4-difluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-benzyl)-carbamique;
- (164) ester tert-butyle de l'acide (4-{2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-benzyl)-carbamique;
- (165) ester tert-butyle de l'acide (3-méthyl-4-{2-[4-oxo-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl]-éthyl}-benzyl)-carbamique;
- (166) 8-[2-(3-amino-phényl)-éthanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]dec-1-én-4-one;
- (167) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl)-éthyl]-phényl}-acétamide;
- (168) 3,N,N-triméthyl-4-(2-{4-oxo-2-[3-(2,2,2-trifluoro-éthoxyméthyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-benzamide;
- (169) 4-(2-{2-[3-(2,2-difluoro-éthoxyméthyl)-phényl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-3,N,N-triméthyl-benzamide;
- (170) 3,N,N-triméthyl-4-(2-{4-oxo-2-[3-(2,2,3,3-tetrafluoro-propoxyméthyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-benzamide;
- (171) 4-(2-{2-[3-(3,5-diméthyl-isoxazol-4-yl)-phényl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-3,N,N-triméthyl-benzamide;
- (172) 4-[2-(2-biphényl-3-yl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,N,N-triméthyl-benzamide;
- (173) 3,N,N-triméthyl-4-{2-[4-oxo-2-(3-pyridin-3-yl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (174) ester méthyle de l'acide 3-{8-[2-(4-diméthylcarbamoyl-2-méthyl-phényl)-éthanesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl}-benzoïque;
- (175) 4-(2-{2-[1-(2,4-dichloro-phénoxy)-1-méthyl-éthyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-3,N,N-triméthyl-benzamide;
- (176) 3,N,N-triméthyl-4-{2-[4-oxo-2-(2,3,4-trifluoro-phénoxyméthyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (177) (3-éthyl-phényl)-amide de l'acide 2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonique;
- (178) 2-cyclohexyl-8-[(E)-2-(1H-indol-5-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (179) 2-cyclohexyl-8-[(E)-2-(2-trifluorométhyl-phényl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (180) 2-cyclohexyl-8-[(E)-2-(3-méthoxy-phényl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (181) 2-cyclohexyl-8-[(E)-2-(2-méthoxy-phényl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (182) 2-cyclohexyl-8-[(E)-2-(1H-indol-4-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (183) 2-cyclohexyl-8-[(E)-2-(2-fluoro-6-trifluorométhyl-phényl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (184) 2-cyclohexyl-8-[(E)-2-(2,3-difluoro-phényl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (185) 2-cyclohexyl-8-[(E)-2-(3-fluoro-2-trifluorométhyl-phényl)-éthènesulfomyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (186) 2-cyclohexyl-8-[(E)-2-o-tolyl-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (187) 2-cyclohexyl-8-[(E)-2-[4-(2-hydroxy-éthyl)-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (188) 2-cyclohexyl-8-[(E)-2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (189) diméthylamide de l'acide 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vi-

nyl]-indole-1-carboxylique;

(190) N-{2-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-méthyl-phényl}-acétamide;

(191) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(192) 2-cyclohexyl-8-[(E)-2-[1-(2-morpholin-4-yl-éthyl)-1H-indol-4-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(193) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-indol-1-yl]-N,N-diméthyl-acétamide;

(194) 3-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phényl]-1,1-diméthyl-urée;

(195) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phényl}-amide de l'acide cyclopropanecarboxylique;

(196) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phényl}-2-hydroxy-acétamide;

(197) ester méthyle de l'acide {3-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-4-méthyl-phényl}-carbamique;

(198) 1-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phényl}-3-(2-hydroxy-éthyl)-urée;

(199) 2-cyclohexyl-8-[(E)-2-[5-(2-hydroxy-éthylamino)-2-méthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(200) 2-cyclohexyl-8-[(E)-2-[2-(2-hydroxy-éthylamino)-6-méthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(201) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phénylamino]-N,N-diméthyl-acétamide;

(202) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,2-diméthyl-[1,3]dioxolan-4-ylméthyl)-3-fluoro-1H-indol-4-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(203) 2-cyclohexyl-8-[(E)-2-[2-méthyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(204) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3,5-diméthyl-phényl}-acétamide;

(205) ester tert-butyle de l'acide {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro [4.5] déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-benzyl}-carbamique;

(206) 2-cyclohexyl-8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(207) 2-cyclohexyl-8-[(E)-2-[2-méthyl-4-[(2-oxo-oxazolidin-5-ylméthyl)-amino]-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(208) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl]-N-(2-diméthylamino-éthyl)-acétamide;

(209) 3-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzonitrile;

(210) 8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(211) 8-[(E)-2-[4-(2-hydroxy-éthylamino)-2-trifluorométhyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(212) 8-[(E)-2-[2-méthyl-4-(pyrrolidine-1-carbonyl)-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(213) 3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzènesulfonamide;

(214) N-(2-hydroxy-éthyl)-3,N-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;

(215) N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-acétamide;

(216) 8-[(E)-2-[2-méthyl-4-(4-méthyl-piperazine-1-carbonyl)-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(217) 3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-N-(2,2,2-trifluoro-éthyl)-benzènesulfonamide;

(218) 3-méthyl-N-(2-morpholin-4-yl-éthyl)-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzènesulfonamide;

- (219) 3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-N-pyridin-3-yl-benzènesulfonamide;
- (220) 8-((E)-2-[4-((R)-4-hydroxy-2-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (221) 8-((E)-2-[4-((R)-2-hydroxyméthyl-5-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (222) 8-((E)-2-[4-((R)-5-hydroxyméthyl-2-oxo-oxazolidin-3-yl)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (223) 8-((E)-2-[4-[3-(3-diméthylamino-propoxy)-azetidine-1-carbonyl]-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (224) N-(4-hydroxy-butyl)-3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzènesulfonamide;
- (225) N-méthyl-N-(4-méthyl-3-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (226) 8-((E)-2-(3-hydroxy-2-méthyl-phényl)-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (227) 8-((E)-2-(5-hydroxy-2-méthyl-phényl)-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (228) N-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-(4-méthyl-pipéridin-4-yl)-acétamide;
- (229) N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-méthanesulfonamide;
- (230) 8-((E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (231) 3,N,N-triméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzamide;
- (232) N-(2-hydroxy-éthyl)-N-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (233) 3-fluoro-N,N-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzamide;
- (234) 8-((E)-2-[2,5-dichloro-4-(3,4-dihydroxy-butoxy)-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (235) 8-((E)-2-[4-(3-diméthylamino-propoxy)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (236) N-méthyl-N-(2-méthyl-3-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (237) N-méthyl-N-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (238) N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (239) N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-méthyl-acétamide;
- (240) 2-cycloheptyl-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (241) 2-(4,4-difluoro-cyclohexyl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (242) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (243) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(2-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (244) 2-(2-fluoro-3-trifluorométhyl-phényl)-8-((E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (245) 8-((E)-2-[1-((S),2,2-diméthyl-[1,3]dioxolan-4-yl)méthyl)-1H-indol-4-yl]-éthènesulfonyl)-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (246) 8-((E)-2-[1-(2,2-diméthyl-[1,3]dioxolan-4-yl)méthyl)-1H-indol-4-yl]-éthènesulfonyl)-2-(tetrahydro-pyran-4-yl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (247) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthènesulfonyl)-2-(2-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (248) 2-(2-fluoro-3-trifluorométhyl-phényl)-8-((E)-2-[2-méthyl-4-(pyrrolidine-1-carbonyl)-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (249) 2-(2,4-dichloro-phényl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 5 (250) 2-(4-chloro-phényl)-8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (251) 2-cyclohexyl-8-((E)-4-(1H-indol-4-yl)-but-3-ène-1-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (252) 2-cyclohexyl-8-((E)-5-(1H-indol-4-yl)-pent-4-ène-1-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (253) 2-cyclohexyl-8-((E)-3-[1-(2,3-dihydroxy-propyl)-1H-indol-4-yl]-prop-2-ène-1-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 10 (254) 8-((E)-2-[3-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (255) 8-((E)-2-[5-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 15 (256) 8-((E)-2-[3-(2-hydroxy-éthoxy)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (257) N-(2-hydroxy-éthyl)-N-(4-méthyl-3-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (258) 8-((E)-2-[4-((R)-2-hydroxyméthyl-5-oxo-pyrrolidin-1-yl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 20 (259) N-(4-((E)-2-[2-(2-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3-méthyl-phényl)-N-(2-hydroxy-éthyl)-acétamide;
- (260) N-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-pipéridin-4-yl-acétamide;
- 25 (261) 2-cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzoxazol-7-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (262) 2-cyclohexyl-8-[5-(1H-indol-4-yl)-pentane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (263) 2-cyclohexyl-8-[4-(1H-indol-4-yl)-butane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (264) N-méthyl-N-(2-méthyl-3-[2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- 30 (265) N-méthyl-N-(4-méthyl-3-[2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- (266) 8-{2-[3-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 35 (267) 8-{2-[5-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (268) 8-{2-[4-((R)-2-hydroxyméthyl-5-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (269) 2-cyclohexyl-8-[2-(1H-indol-5-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 40 (270) 2-[4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-phényl-amino]-N,N-diméthyl-acétamide;
- (271) {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-phényl}-amide de l'acide cyclopropanecarboxylique;
- (272) ester tert-butyle de l'acide{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-benzyl}-carbamique;
- 45 (273) 3-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-phényl}-1,1-diméthyl-urée;
- (274) ester méthyle de l'acide {3-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-4-méthyl-phényl}-carbamique;
- 50 (275) 2-cyclohexyl-8-{2-[5-(2-hydroxy-éthylamino)-2-méthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (276) N-{2-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-méthyl-phényl}-acétamide;
- (277) 2-cyclohexyl-8-(2-o-tolyl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 55 (278) 1-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-phényl}-3-(2-hydroxy-éthyl)-urée;
- (279) 2-cyclohexyl-8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (280) 2-cyclohexyl-8-[2-[4-(2-hydroxy-éthyl)-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (281) 2-cyclohexyl-8-(2-{2-méthyl-4-[(2-oxo-oxazolidin-5-ylméthyl)-amino]-phényl}-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (282) 2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-N-(2-diméthylamino-éthyl)-acétamide;
- (283) 2-cyclohexyl-8-{2-[2-méthyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (284) 8-[2-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthanesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (285) 8-{2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (286) 8-[2-[4-(2-hydroxy-éthylamino)-2-trifluorométhyl-phényl]-éthanesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (287) N-(4-hydroxy-butyl)-3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonamide;
- (288) 8-{2-[2-méthyl-4-(pyrrolidine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (289) 8-{2-[2-méthyl-4-(4-méthyl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (290) 3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-N-(2,2,2-trifluoro-éthyl)-benzènesulfonamide;
- (291) 3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonamide;
- (292) 3-méthyl-N-(2-morpholin-4-yl-éthyl)-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonamide;
- (293) 3-fluoro-N,N-diméthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (294) 3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-N-pyridin-3-yl-benzènesulfonamide;
- (295) 8-{2-[4-(3-diméthylamino-propoxy)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (296) 8-{2-[4-((R)-5-hydroxyméthyl-2-oxo-oxazolidin-3-yl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (297) 8-[2-[4-((R)-4-hydroxy-2-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthanesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (298) N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;
- (299) N-méthyl-N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;
- (300) 2-hydroxy-N-méthyl-N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;
- (301) 1-méthyl-1-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-urée;
- (302) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl}-acétamide;
- (303) 2-cyclohexyl-8-[2-(1H-indol-4-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (304) 2-(3-trifluorométhoxy-phényl)-8-[2-(2-trifluorométhyl-phényl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (305) 8-[2-(1H-indol-4-yl)-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (306) 8-[2-(1H-indol-4-yl)-éthanesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (307) 8-[2-(2,2-difluoro-benzo[1,3]dioxol-4-yl)-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (308) 8-[2-(1-méthyl-1H-indol-7-yl)-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (309) 2-cyclohexyl-8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (310) diméthylamide de l'acide 4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-indole-1-carboxylique;

- (311) 8-[3-(1H-indol-4-yl)-propane-1-sulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (312) 2-cyclohexyl-8-[2-(1-méthanesulfonyl-1H-indol-4-yl)-éthanesulfonyl]-1,3,8-triaza-spiro [4.5]déc-1-én-4-one;
- (313) 2-cyclohexyl-8-[2-[1-(2-diméthylamino-éthyl)-1H-indol-4-yl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (314) 8-[2-(4-amino-2-trifluorométhyl-phényl)-éthanesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (315) 2-cyclohexyl-8-[3-(3-trifluorométhyl-phényl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (316) 2-cyclohexyl-8-(3-m-tolyl-propane-1-sulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (317) 2-cyclohexyl-8-[3-(3-hydroxy-phényl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (318) 2-cyclohexyl-8-[3-(2-hydroxy-phényl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (319) 8-(2-benzo[b]thiophén-3-yl-éthanesulfonyl)-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (320) 2-cyclohexyl-8-(2-isoquinolin-5-yl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (321) 2-cyclohexyl-8-[2-[4-((S)-2,3-dihydroxy-propylamino)-2-méthyl-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (322) 8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthanesulfonyl]-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (323) 2-cyclohexyl-8-[2-(2-trifluorométhyl-phényl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (324) 4-[3-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-propyl]-N,N-diméthyl-benzamide;
- (325) 8-[2-[4-((R)-2,3-dihydroxy-propoxy)-2-méthyl-phényl]-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (326) N-(2-hydroxy-éthyl)-N-(4-méthyl-3-[2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- (327) N-(2-hydroxy-éthyl)-N-(2-méthyl-3-[2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- (328) 8-[3-(3-amino-phényl)-propane-1-sulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (329) N,N-diméthyl-4-{5-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-thiophén-2-yl}-benzamide;
- (330) N,N-diméthyl-3-{5-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-thiophén-2-yl}-benzamide;
- (331) 3,N,N-triméthyl-4-{5-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-thiophén-2-yl}-benzamide;
- (332) N-{4-[5-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-thiophén-2-yl]-3-méthyl-phényl}-acétamide;
- (333) 4-{5-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-thiophén-2-yl}-3,N,N-triméthyl-benzamide;
- (334) 3-{5-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-thiophén-2-yl}-4,N,N-triméthyl-benzamide;
- (335) diméthylamide de l'acide 2-méthyl-3'-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-biphényl-4-carboxylique;
- (336) diméthylamide de l'acide 2-méthyl-4'-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-biphényl-4-carboxylique;
- (337) 3,N,N-triméthyl-4-{6-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-pyridin-2-yl}-benzamide;
- (338) 2-cyclohexyl-8-[2-[2-(3,5-diméthyl-isoxazol-4-yl)-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (339) 2-cyclohexyl-8-[2-[2-(6-méthoxy-pyridin-3-yl)-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (340) 2-cyclohexyl-8-[2-[2-(1-méthyl-1H-pyrazol-4-yl)-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (341) 8-[5-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-thiophène-2-sulfonyl]-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (342) N-(4-[5-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-thiophén-2-yl]-3-méthyl-phényl)-N-(2-hydroxy-éthyl)-acétamide;
- (343) 2-cyclohexyl-8-((E)-2-thiazol-2-yl-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (344) 2-cyclohexyl-8-((E)-2-cyclopentyl-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (345) 2-cyclohexyl-8-(2-thiazol-2-yl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (346) 2-cyclohexyl-8-[3-(4-méthoxy-phényl)-propane-1-sulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (347) N-benzyl-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-benzamide;
- (348) 8-(3-chloro-benzènesulfonyl)-2-[3-(morpholine-4-carbonyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (349) 3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-N-méthyl-N-phényl-benzamide;
- (350) N-benzyl-3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-benzamide;
- (351) 3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-N,N-diméthyl-benzamide;
- (352) 8-{2-[4-(4-méthanesulfonyl-piperazine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (353) N-[(R)-1-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzoyl)-pyrrolidin-3-yl]-acétamide; (354) 8-{2-[2-méthyl-4-((S)-2-trifluorométhyl-pyrrolidine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (355) 8-{2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (356) 8-{2-[4-((S)-2-hydroxyméthyl-pyrrolidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (357) 8-(2-[4-[4-(2-hydroxy-éthyl)-pipéridine-1-carbonyl]-2-méthyl-phényl]-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (358) 8-(2-[2-méthyl-4-[4-(3-méthyl-[1,2,4]oxadiazol-5-yl)-pipéridine-1-carbonyl]-phényl]-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (359) 8-{2-[2-méthyl-4-(4-pyrimidin-2-yl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (360) diméthylamide de l'acide 4-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzoyl)-piperazine-1-sulfonique;
- (361) 8-{2-[2-méthyl-4-(4-pyridin-2-yl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (362) 8-[2-(4-[4-[2-(2-hydroxy-éthoxy)-éthyl]-piperazine-1-carbonyl]-2-méthyl-phényl)-éthanesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (363) 8-(2-[2-méthyl-4-[4-(2-morpholin-4-yl-éthyl)-piperazine-1-carbonyl]-phényl]-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (364) 8-{2-[2-méthyl-4-(4-thiazol-2-yl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (365) 8-{2-[4-(4,4-difluoro-pipéridine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (366) 8-(2-[4-[4-(3-hydroxy-propyl)-piperazine-1-carbonyl]-2-méthyl-phényl]-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (367) amide de l'acide (S)-1-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzoyl)-pyrrolidine-2-carboxylique;
- (368) 8-{2-[2-méthyl-4-(3,3,4,4-tetrafluoro-pyrrolidine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (369) amide de l'acide (R)-1-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzoyl)-pyrrolidine-2-carboxylique;
- (370) 8-{2-[4-((S)-3-hydroxy-pyrrolidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (371) 8-{2-[4-((R)-2-hydroxyméthyl-pyrrolidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (372) 8-{2-[4-((S)-3-diméthylamino-pyrrolidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (373) 8-{2-[4-(4-tert-butyl-piperazine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (374) 8-(2-[4-[4-(3-diméthylamino-propyl)-piperazine-1-carbonyl]-2-méthyl-phényl]-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (375) 8-(2-[4-[4-(4-fluoro-phényl)-piperazine-1-carbonyl]-2-méthyl-phényl]-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (376) 8-{2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (377) 8-{2-[4-(3-hydroxy-azetidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (378) 8-{2-[4-(3-fluoro-pyrrolidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (379) 8-{2-[4-(3-fluoro-azetidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (380) 8-{2-[2-méthyl-4-(pipéridine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (381) 8-{2-[4-(azetidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (382) diméthylamide de l'acide 4-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzoyl)-piperazine-1-carboxylique;
- (383) 3,5,N,N-tetraméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;
- (384) 8-[(E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (385) 8-[(E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (386) 8-[(E)-2-[4-(4-hydroxy-4-hydroxyméthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (387) 8-[(E)-2-[4-((3R,4R)-3-diméthylamino-4-hydroxy-pyrrolidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (388) 8-[(E)-2-[2,6-diméthyl-4-(pyrrolidine-1-carbonyl)-phényl]-éthènesulfonyl]-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (389) N-(3-hydroxy-propyl)-3,5,N-triméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;
- (390) N-(2-diméthylamino-éthyl)-3,5,N-triméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;
- (391) N-(3-diméthylamino-propyl)-3,5,N-triméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;
- (392) N-carbamoylméthyl-3,5,N-triméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;
- (393) 3,S,N-triméthyl-N-(1-méthyl-pipéridin-4-yl)-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;
- (394) 8-[(E)-2-[4-(4-acetyl-piperazine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (395) diméthylamide de l'acide 4-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzoyl)-piperazine-1-carboxylique;
- (396) amide de l'acide 4-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzoyl)-piperazine-1-carboxylique;
- (397) 8-[(E)-2-[4-(3-diméthylamino-propyl)-piperazine-1-carbonyl]-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (398) 8-[(E)-2-[2,6-diméthyl-4-((R)-3-méthylamino-pyrrolidine-1-carbonyl)-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (399) 8-[(E)-2-[4-((R)-3-diméthylamino-pyrrolidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (400) 8-[(E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (401) 8-[(E)-2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (402) 8-[(E)-2-[4-((S)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (403) 8-[(E)-2-[4-(4-hydroxy-4-hydroxyméthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (404) 8-[(E)-2-[4-((R)-3-amino-pyrrolidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (405) 8-[(E)-2-[4-(4-diméthylamino-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (406) 8-[(E)-2-[4-((3S,4S)-3-hydroxy-4-isopropylamino-pyrrolidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (407) 8-((E)-2-{4-[4-(2-diméthylamino-éthoxy)-pipéridine-1-carbonyl]-2,6-diméthyl-phényl}-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (408) 8-((E)-2-{4-(3-hydroxy-azetidine-1-carbonyl)-2,6-diméthyl-phényl}-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (409) 2,N,N-triméthyl-3-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (410) 2-méthyl-3-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (411) 8-{2-[2-méthyl-3-(morpholine-4-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (412) 4,N,N-triméthyl-3-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (413) N-(2-hydroxy-éthyl)-4,N-diméthyl-3-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (414) 8-{2-[2-méthyl-5-(4-méthyl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (415) 4,N-diméthyl-3-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (416) 8-(2-{4-[4-(2-hydroxy-éthyl)-piperazine-1-carbonyl]-2-méthyl-phényl}-éthanesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (417) 8-(2-[2-méthyl-4-(thiazolidine-3-carbonyl)-phényl]-éthanesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (418) 2-fluoro-5,N,N-triméthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;
- (419) 8-{2-[5-fluoro-4-(4-hydroxy-pipéridine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (420) N-benzyl-2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-1H-indol-3-yl}-acétamide;
- (421) 2-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-N-méthyl-benzamide;
- (422) 3-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-N-pyridin-4-yl-benzamide;
- (423) 8-(3-chloro-benzènesulfonyl)-2-[1-(3,3-diméthyl-butyryl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (424) 8-(3-chloro-benzènesulfonyl)-2-[1-(2-hydroxy-acetyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (425) 8-(3-chloro-benzènesulfonyl)-2-[1-(4-chloro-benzoyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (426) 8-(3-chloro-benzènesulfonyl)-2-[1-(3-méthoxy-benzoyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (428) 2-[1-(1H-indole-5-carbonyl)-pipéridin-3-yl]-8-(2-naphthalèn-1-yl)-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (429) 8-(2-naphthalèn-1-yl)-éthanesulfonyl)-2-[1-(3-phényl-propionyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (430) 8-(2-naphthalèn-1-yl)-éthanesulfonyl)-2-[1-(1,2,3,4-tetrahydro-naphthalène-2-carbonyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (431) 8-(2-naphthalèn-1-yl)-éthanesulfonyl)-2-[1-(quinoline-6-carbonyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (432) 2-[1-(2-3H-imidazol-4-yl-acetyl)-pipéridin-3-yl]-8-(2-naphthalèn-1-yl)-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (433) 2-[1-[2-(2,5-diméthyl-thiazol-4-yl)-acetyl]-pipéridin-3-yl]-8-(2-naphthalèn-1-yl)-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (434) 2-[1-[2-(5-méthyl-2-phényl-oxazol-4-yl)-acetyl]-pipéridin-3-yl]-8-(2-naphthalèn-1-yl)-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]dc-1-én-4-one;
- (435) 8-(2-naphthalèn-1-yl)-éthanesulfonyl)-2-[1-(2-pyridin-2-yl-acetyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (436) 2-[1-(4-benzyloxy-butyryl)-pipéridin-3-yl]-8-(2-naphthalèn-1-yl)-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (437) 8-(2-naphthalèn-1-yl)-éthanesulfonyl)-2-[1-(4-phényl-butyryl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (438) 8-(2-naphthalèn-1-yl-éthanesulfonyl)-2-[1-(2-pyridin-4-yl-acetyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (439) 2-[1-(2-tert-butoxy-acetyl)-pipéridin-3-yl]-8-(2-naphthalèn-1-yl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 5 (441) 8-(2-naphthalèn-1-yl-éthanesulfonyl)-2-{1-[(E)-(3-phényl-acryloyl)]-pipéridin-3-yl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (442) 2-[1-(2-amino-acetyl)-pipéridin-3-yl]-8-(3-chloro-benzènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (444) 8-{2-[2-méthyl-4-(3-méthylamino-pyrrolidine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 10 (445) 8-{2-[4-(3-amino-pyrrolidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one hydrochloride;
- (446) 8-[(E)-2-[2,6-diméthyl-4-(piperazine-1-carbonyl)-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one hydrochloride;
- (447) 8-[(E)-2-[4-(2-hydroxy-acetyl)-piperazine-1-carbonyl]-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 15 (448) ester méthyle de l'acide 4-[4-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzoyl]-piperazin-1-yl]-4-oxo-butyrique
- (449) 8-[(E)-2-[4-(4-diméthylamino-butyryl)-piperazine-1-carbonyl]-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 20 (450) 2-méthoxy-N-méthyl-N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-acétamide;
- (451) 2-hydroxy-N-méthyl-N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-acétamide;
- (452) acide [(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzoyl)-méthyl-amino]-acétique ;
- 25 (453) 8-(3-chloro-benzènesulfonyl)-2-[1-(3,3-diméthyl-butyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (454) tert-butylamide de l'acide 3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- 30 (455) phénéthyl-amide de l'acide 3-[8-(2-naphthalèn-1-yl-éthanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- (456) 8-(3-chloro-benzènesulfonyl)-2-[1-(pipéridine-1-carbonyl)-pipéridin-3-yl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (457) 3-(2-diméthylamino-éthyl)-1-méthyl-1-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phénol)-urée;
- 35 (458) 1-méthyl-1-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-urée;
- (459) 1-{3-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-2-méthyl-phényl}-3-méthyl-urée;
- 40 (460) ester 2-méthoxy-éthyle de l'acide 3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- (461) ester benzyle de l'acide 3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- (462) ester isobutyle de l'acide 3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- 45 (463) ester méthyle de l'acide 3-[8-(3-chloro-benzènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- (464) ester benzyle de l'acide 3-[8-(2-naphthalèn-1-yl-éthanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- 50 (465) ester 2,2-diméthyl-propyle de l'acide 3-[8-(2-naphthalèn-1-yl-éthanesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl]-pipéridine-1-carboxylique;
- (466) ester 2-diméthylamino-éthyle de l'acide méthyl-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-carbamique;
- (467) ester méthyle de l'acide méthyl-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-carbamique;
- 55 (468) ester méthyle de l'acide méthyl-(3-méthyl-4-[2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-carbamique ;
- (469) 2-cyclohexyl-8-(2-o-tolyl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-4-thione;

- (470) 1-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;
- (471) 1-(3,5-diméthyl-4-{(E)-2-[4-oxo-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl}-phényl)-1-méthyl-urée;
- (472) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;
- (474) 1-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-thiourée;
- (475) 8-{2-[2,6-diméthyl-4-(méthyl-thiazol-2-yl-amino)-phényl]-éthanesulfonyl}-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (476) ester méthyle de l'acide {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-indol-1-yl}-acétique
- (477) 2-cyclohexyl-8-[(E)-2-[1-(morpholine-4-carbonyl)-1H-indol-4-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (478) 2-cyclohexyl-8-[(E)-2-[1-(4-hydroxy-butyl)-1H-indol-4-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (479) 2-cyclohexyl-8-[(E)-2-(1H-indol-7-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (480) 2-cyclohexyl-8-[(E)-2-[1-(3,4-dihydroxy-butyl)-1H-indol-4-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (481) N-(2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl}-éthyl)-acétamide;
- (482) N-(1-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl}-éthyl)-acétamide;
- (483) 2-cyclohexyl-8-[(E)-2-(1-thiazol-2-ylméthyl-1H-indol-4-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (484) amide de l'acide 3-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-indol-1-yl}-propane-1-sulfonique;
- (485) {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phényl}-amide de l'acide 4-méthyl-piperazine-1-carboxylique;
- (486) ester isobutyle de l'acide {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-benzyl}-carbamique;
- (487) 2-cyclohexyl-8-[(E)-2-{1-[2-(2-hydroxy-éthoxy)-éthyl]-1H-indol-5-yl}-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (488) 2-cyclohexyl-8-[(E)-2-(6-méthyl-1H-indol-5-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (489) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-benzyl}-2-hydroxy-acétamide;
- (490) 3-(2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl}-éthyl)-1,1-diméthyl-urée;
- (491) 8-[(E)-2-(4-amino-2,6-bis-trifluorométhyl-phényl)-éthènesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (492) 2-cyclohexyl-8-[(E)-2-(6-trifluorométhyl-1H-indol-5-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (493) ester méthyle de l'acide {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phénylamino}-acétique;
- (494) 2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phénylamino}-N-(2-hydroxy-éthyl)-acétamide;
- (495) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-N-(2-hydroxy-éthyl)-N-méthyl-3-trifluorométhyl-benzènesulfonamide;
- (496) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-N,N-bis-(2-hydroxy-éthyl)-3-trifluorométhyl-benzènesulfonamide;
- (497) 2-cyclohexyl-8-[(E)-2-[4-(2-diméthylamino-éthylamino)-2-méthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (498) N-(2-acétylamino-éthyl)-2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phénylamino}-acétamide;
- (499) 2-cyclohexyl-8-[(E)-2-(6-trifluorométhyl-1H-benzimidazol-5-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (500) 2-cyclohexyl-8-[(E)-2-[1-((S)-2,2-diméthyl-[1,3]dioxolan-4-ylméthyl)-5-fluoro-1H-indol-4-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (501) 2-cyclohexyl-8-((E)-2-[4-[(2-hydroxy-éthyl)-méthyl-amino]-2-trifluorométhyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (502) 2-cyclohexyl-8-((E)-2-[4-(1,1-dioxo-1λ⁶-thiomorpholine-4-carbonyl)-2-méthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (503) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-N-(2-diméthylamino-éthyl)-3-méthyl-benzènesulfonamide;
- (504) 3-(3-méthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-imidazolidine-2,4-dione;
- (505) 1-méthyl-3-(3-méthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-imidazolidine-2,4-dione;
- (506) N-(3-isopropoxy-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (507) 3-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-5,5-diméthyl-imidazolidine-2,4-dione;
- (508) 3-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1,5,5-triméthyl-imidazolidine-2,4-dione;
- (509) N-(3-éthoxy-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (510) N-(3-éthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (511) 5,5-diméthyl-3-(3-méthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-imidazolidine-2,4-dione;
- (512) N-(3-méthoxyméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (513) 2-(4-méthyl-cyclohexyl)-8-((E)-2-[2-méthyl-4-(2-oxo-oxazolidin-3-yl)-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (514) N-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-(2-méthoxy-éthyl)-acétamide;
- (515) 8-((E)-2-[2,6-diméthyl-4-(3-oxo-morpholin-4-yl)-phényl]-éthènesulfonyl)-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (516) 8-((E)-2-[2,6-diméthyl-4-(2-oxo-pipéridin-1-yl)-phényl]-éthènesulfonyl)-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (517) N-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]dec-1-ène-8-sulfonyl]-vinyl)-benzyl)-N-méthyl-acétamide;
- (518) 3-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl)-imidazolidine-2,4-dione;
- (519) 3-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl)-5,5-diméthyl-imidazolidine-2,4-dione;
- (520) 8-((E)-2-[2,6-diméthyl-4-(2-oxo-pyrrolidin-1-ylméthyl)-phényl]-éthènesulfonyl)-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (521) 1-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl)-pyrrolidine-2,5-dione;
- (522) N-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl)-acétamide;
- (523) N-(3,5-difluoro-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-méthyl-acétamide;
- (524) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (525) ester tert-butyle de l'acide [(S)-1-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-2,5-dioxo-pyrrolidin-3-yl]-carbamique;
- (526) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (527) 8-((E)-2-[2-méthyl-4-(4-méthyl-piperazine-1-carbonyl)-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (528) 8-((E)-2-[2-méthyl-4-[2-(4-méthyl-piperazin-1-yl)-2-oxo-éthylamino]-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (529) 2-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phénylamino)-acétamide;

- (530) 3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-N-pyridin-4-yl-benzènesulfonamide;
- (531) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (532) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1,3-bis-(2-hydroxy-éthyl)-urée;
- (533) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1,3-bis-(2-hydroxy-éthyl)-3-méthyl-urée;
- (534) N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-(tetrahydro-pyran-4-yl)-méthanesulfonamide;
- (535) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-((E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (536) 3-{3-méthyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl}-imidazolidine-2,4-dione;
- (537) 3-{4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-trifluorométhyl-phényl}-imidazolidine-2,4-dione;
- (538) 4-{3-méthyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl}-morpholine-3,5-dione;
- (539) 3-{3,5-diméthyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl}-imidazolidine-2,4-dione;
- (540) 5,5-diméthyl-3-[3-méthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-imidazolidine-2,4-dione;
- (541) 8-((E)-2-[1-(R)-2,3-dihydroxy-propyl]-1H-indol-4-yl)-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (542) N-(2-méthoxy-éthyl)-N-[3-méthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-acétamide;
- (543) N-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-N-(2-méthoxy-éthyl)-acétamide;
- (544) N-[2-(2-hydroxy-éthoxy)-éthyl]-N-[3-méthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-acétamide;
- (545) N-[2-(2-fluoro-éthoxy)-éthyl]-N-[3-méthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-acétamide;
- (546) N-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-N-méthyl-acétamide;
- (547) acide 3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzoïque;
- (548) 3-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-imidazolidine-2,4-dione;
- (549) 8-((E)-2-[4-[4-(2-fluoro-éthyl)-piperazine-1-carbonyl]-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (550) 8-((E)-2-[2,6-diméthyl-4-(3-oxo-morpholin-4-yl)-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (551) N-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-N-[2-(2-fluoro-éthoxy)-éthyl]-acétamide;
- (552) N'-acetyl-hydrazide de l'acide 3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzoïque;
- (553) 8-[(E)-2-(4-hydroxyméthyl-2,6-diméthyl-phényl)-éthènesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (554) 3-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl]-imidazolidine-2,4-dione;
- (555) 3-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl]-5,5-diméthyl-imidazolidine-2,4-dione;
- (556) 8-((E)-2-[2,6-diméthyl-4-(2-oxo-pyrrolidin-1-ylméthyl)-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (557) 1-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl]-pyrrolidine-2,5-dione;
- (558) N-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl]-acétamide;

- (559) N-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzyl]-N-méthyl-acétamide;
- (560) 3-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3-méthyl-phényl)-imidazolidine-2,4-dione;
- (561) N-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-N-[2-(2-méthoxy-éthoxy)-éthyl]-acétamide;
- (562) N-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-N-[2-(2-fluoro-éthoxy)-éthyl]-acétamide;
- (563) 3-(4-((E)-2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3-méthyl-phényl)-imidazolidine-2,4-dione-;
- (564) 3-(4-((E)-2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-imidazolidine-2,4-dione;
- (565) 3-(4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-benzyl)-1,1-diméthyl-urée;
- (566) ester méthyle de l'acide (2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl]-éthyl)-carbamique;
- (567) 4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-benzylamide de l'acide 4-méthyl-piperazine-1-carboxylique;
- (568) N-(4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-méthyl-phényl)-N',N'-diméthyl-sulfamide;
- (569) ester tert-butyle de l'acide {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-méthyl-benzyl}-(3-diméthylamino-propyl)-carbamique;
- (570) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl]-N-(2,2-diméthyl-propyl)-acétamide;
- (571) 2-[4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl]-N-(2-diisopropylamino-éthyl)-acétamide;
- (572) 2-cyclohexyl-8-[(E)-2-[2-méthyl-4-(2-oxo-azetidin-1-yl)-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (573) 2-cyclohexyl-8-[(E)-2-[2,6-diméthyl-4-(thiazol-2-ylamino)-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (574) N-(4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3,5-diméthyl-phényl)-N-(3-méthyl-oxetan-3-ylméthyl)-acétamide;
- (575) 3-(3,5-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-imidazolidine-2,4-dione;
- (576) N-(3-éthoxyméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-acétamide;
- (577) 2-(4-méthyl-cyclohexyl)-8-[(E)-2-[2-méthyl-4-(4-oxo-oxazolidin-3-yl)-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (578) ester tert-butyle de l'acide 4-(3-méthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-3-oxo-piperazine-1-carboxylique;
- (579) 2-(4-méthyl-cyclohexyl)-8-[(E)-2-[2-méthyl-4-(4-méthyl-2-oxo-oxazolidin-3-yl)-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (580) 2-(4-méthyl-cyclohexyl)-8-[(E)-2-[2-méthyl-4-(3-oxo-isoxazolidin-2-yl)-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (581) 3-(2,5-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-imidazolidine-2,4-dione;
- (582) 3-(3,5-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-5-méthyl-imidazolidine-2,4-dione;
- (583) 3-(2,6-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-imidazolidine-2,4-dione;
- (584) N-(2,6-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-N-méthyl-acétamide;
- (585) 8-[(E)-2-(3-méthoxy-2-méthyl-phényl)-éthènesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (586) 8-[(E)-2-[4-[(1H-imidazol-2-yl)-méthyl-amino]-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (587) 1-(3,5-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-dihydro-pyrimidine-2,4-dione;

- (588) 2-(4-méthyl-cyclohexyl)-8-((E)-2-[2-méthyl-4-(5-oxo-pyrazolidin-1-yl)-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (589) 5-tert-butoxyméthyl-3-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-imidazolidine-2,4-dione;
- (590) N,N-diméthyl-2-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzènesulfonylamino)-acétamide;
- (591) 3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-N-pyridin-3-ylméthyl-benzènesulfonamide;
- (592) N-(4-hydroxy-cyclohexyl)-3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzènesulfonamide;
- (593) 8-((E)-2-(4-méthanesulfonyl-2-méthyl-phényl)-éthènesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (594) N-acetyl-3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzènesulfonamide;
- (595) N-(2-hydroxy-1,1-bis-hydroxyméthyl-éthyl)-3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzènesulfonamide;
- (596) N-(1-benzyl-pipéridin-4-yl)-3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzènesulfonamide;
- (597) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (598) 8-((E)-2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-méthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (599) N-((R)-2,3-dihydroxy-propyl)-3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzènesulfonamide;
- (600) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-3,5-difluoro-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (601) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-3,5-diméthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (602) ester tert-butyle de l'acide 4-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phénoxy)-pipéridine-1 carboxylique;
- (603) N-isopropyl-N-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (604) ester tert-butyle de l'acide 4-[acetyl-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-amino]-pipéridine-1-carboxylique;
- (605) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-(2-hydroxy-éthyl)-urée;
- (606) 8-((E)-2-[4-((R)-2-(isopropylamino-méthyl)-5-oxo-pyrrolidin-1-yl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (607) N-(2-diméthylamino-éthyl)-N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phénol)-acétamide;
- (608) 2-diméthylamino-N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-méthyl-acétamide;
- (609) N-(2-diméthylamino-éthyl)-N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-méthanesulfonamide;
- (610) N-((R)-1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-5-oxo-pyrrolidin-2-ylméthyl)-acétamide;
- (611) 8-((E)-2-[4-((R)-2-diméthylaminométhyl-5-oxo-pyrrolidin-1-yl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (612) N-cyanométhyl-N-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (613) N-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-(2,2,2-trifluoro-éthyl)-acétamide;
- (614) 3-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1,5,5-triméthyl-imidazolidine-2,4-dione;
- (615) 3-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-5,5-diméthyl-imidazolidine-2,4-dione;
- (616) N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-(tetrahydro-pyran-4-yl)-acétamide;

- (617) 8-((E)-2-[2-méthyl-4-(3-oxo-morpholin-4-yl)-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (618) 3-(3-méthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-imidazolidine-2,4-dione;
- (619) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-((E)-2-[2-méthyl-4-(pyrrolidine-1-carbonyl)-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (620) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-((E)-2-[4-((R)-2-hydroxyméthyl-5-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (621) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-((E)-2-[4-((S)-2-hydroxyméthyl-5-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (622) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (623) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (624) 1-(4-((E)-2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3-méthyl-phényl)-imidazolidine-2,4-dione;
- (625) 1-(4-((E)-2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3-méthyl-phényl)-5-méthyl-imidazolidine-2,4-dione;
- (626) 1-(4-((E)-2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3-méthyl-phényl)-5,5-diméthyl-imidazolidine-2,4-dione;
- (627) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (628) acide 3,5-diméthyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-benzoïque;
- (629) triméthylhydrazide de l'acide 3,5-diméthyl-4-[(E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-vinyl]-benzoïque;
- (630) 8-((E)-2-[1-((R)-2,3-dihydroxy-propyl)-4,6-diméthyl-1H-indol-5-yl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (631) 3-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-imidazolidine-2,4-dione;
- (632) N-(4-((E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-N-(3-méthyl-oxetan-3-ylméthyl)-acétamide;
- (633) 2-cyclohexyl-8-((E)-2-quinolin-8-yl-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (634) 2-cyclohexyl-8-[(E)-2-(2-oxo-2,3-dihydro-1H-indol-4-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (635) 3-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-imidazolidine-2,4-dione;
- (636) 3-(4-((E)-2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-imidazolidine-2,4-dione;
- (637) 2-cyclohexyl-8-[(E)-2-(2-méthyl-1H-indol-4-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (638) 2-cyclohexyl-8-[(E)-2-(1-méthyl-1,2,3,4-tetrahydro-quinolin-5-yl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (639) 8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (640) ester tert-butyle de l'acide (4-((E)-2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-[1,1,1-²H₃]méthyl-carbamique;
- (641) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (642) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (643) 8-((E)-2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthènesulfonyl)-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (644) 8-((E)-2-[2,6-diméthyl-4-(2-oxa-7-aza-spiro[3.5]nonane-7-carbonyl)-phényl]-éthènesulfonyl)-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (645) 2-cyclohexyl-8-[2-(1,2,3,4-tetrahydro-quinolin-5-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (646) 4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-1H-indole-3-carbonitrile;
- (647) diméthylamide de l'acide 7-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-indole-1-carboxylique;

- (648) 2-cyclohexyl-8-(2-quinolin-5-yl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (649) 2-cyclohexyl-8-{2-[3-(2,2,2-trifluoro-acetyl)-1H-indol-4-yl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (650) 2-cyclohexyl-8-[2-(1-isopropyl-1H-indol-4-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (651) 2-cyclohexyl-8-{2-[1-(4-hydroxy-butyl)-1H-indol-4-yl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (652) N-{3-cyano-4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-acétamide;
- (653) 8-(2-isoquinolin-5-yl-éthanesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (654) 8-(2-quinolin-5-yl-éthanesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (655) N-(3-méthoxy-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;
- (656) 8-{2-[4-(3,3-diméthyl-1,1,4-trioxo-1 λ ⁶-[1,2,5]thiadiazolidin-2-yl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (657) 8-{2-[2-méthyl-4-(3-oxo-morpholin-4-yl)-phényl]-éthanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (658) ester isobutyle de l'acide {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-benzyl}-carbamique;
- (659) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-benzyl}-2-hydroxy-acétamide;
- (660) 2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-phényl-amino}-N-(4-hydroxy-butyl)-acétamide;
- (661) ester isobutyle de l'acide {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-phényl}-carbamique;
- (662) N-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-phényl}-benzamide;
- (663) 2-cyclohexyl-8-[2-(2-oxo-2,3-dihydro-benzothiazol-6-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (664) N-(2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-méthyl-benzènesulfonylamino}-éthyl)-acétamide;
- (665) N-(2-diméthylamino-éthyl)-2-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phénylamino)-acétamide;
- (666) 8-[2-[4-((2R,6S)-2,6-diméthyl-morpholine-4-carbonyl)-2-méthyl-phényl]-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (667) N-(2,2,3,3,4,4,4-heptafluoro-butyl)-3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonamide;
- (668) 8-[2-(3-méthoxy-2-méthyl-phényl)-éthanesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (669) ester tert-butyle de l'acide (2-méthoxy-3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-méthyl-carbamique;
- (670) 8-{2-[4-((S)-2-hydroxyméthyl-5-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthanesulfonyl}-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (671) 8-[2-(5,7-diméthyl-2-oxo-2,3-dihydro-benzoxazol-6-yl)-éthanesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (672) ester méthyle de l'acide (2-{4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-éthyl)-carbamique;
- (673) ester tert-butyle de l'acide {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-méthyl-benzyl}-méthyl-carbamique;
- (674) 8-(2-{4-[3-(2-hydroxy-éthyl)-2-oxo-imidazolidin-1-yl]-2-méthyl-phényl}-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (675) 2-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonylamino)-acétamide;
- (676) 8-{2-[4-(3,4-dihydroxy-butoxy)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (677) 3-(4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-imidazolidine-2,4-dione;
- (678) N-cyclopentyl-N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-

sulfonyl]-éthyl)-phényl)-acétamide;

(679) 1-(2,3-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(680) 1-(2,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(681) ester tert-butyle de l'acide {4-[2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-méthyl-benzyl}-(2-hydroxy-éthyl)-carbamique;

(682) 1-(3-méthoxy-5-méthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(683) 1-(3-chloro-5-méthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(684) 1-(2-méthoxy-3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(685) 5,7-diméthyl-6-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-1H-quinazoline-2,4-dione;

(686) 8-[(E)-2-[1-(2-amino-éthyl)-1H-indol-4-yl]-éthènesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(687) 8-[(E)-2-(4-aminométhyl-phényl)-éthènesulfonyl]-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(688) (S)-2-amino-3-hydroxy-N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-propionamide;

(689) N-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-N-pipéridin-4-yl-méthanésulfonamide;

(690) 8-[(E)-2-(2,6-diméthyl-4-méthylamino-phényl)-éthènesulfonyl]-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(691) 8-[(E)-2-(2,6-diméthyl-4-méthylamino-phényl)-éthènesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(692) 1-{3,5-diméthyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl}-1-méthyl-urée;

(693) 1-(2-méthoxy-3,5-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-1-méthyl-urée;

(694) 1-{3,5-diméthyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl}-1-(2-fluoro-éthyl)-urée;

(695) 1-(3-chloro-5-méthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-1-méthyl-urée;

(696) 1-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(697) 8-[(E)-2-[4-[(R)-2,3-dihydroxy-propoxy]-2,6-diméthyl-phényl]-éthènesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(698) 8-[(E)-2-[4-[2-[(R)-2,3-dihydroxy-propoxy]-éthoxy]-2,6-diméthyl-phényl]-éthènesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(699) N-[3,5-diméthyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl]-N-[2-(2-hydroxy-éthoxy)-éthyl]-acétamide;

(700) N-[2-(2-hydroxy-éthoxy)-éthyl]-N-(3-méthyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl)-acétamide;

(701) N-[3,5-diméthyl-4-[(E)-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl]-N-(2-hydroxy-éthyl)-acétamide;

(702) 2-cyclohexyl-8-[(E)-2-[1-[(S)-2,3-dihydroxy-propyl]-1H-indol-5-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(704) 8-[(E)-2-[1-[2-[(S)-2,3-dihydroxy-propoxy]-éthyl]-1H-indol-4-yl]-éthènesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(705) 8-[(E)-2-[2,6-diméthyl-4-[(2S,3S)-2,3,4-trihydroxy-butoxy]-phényl]-éthènesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(706) 2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-8-[(E)-2-[1-[2-[(2S,3S)-2,3,4-trihydroxy-butoxy]-éthyl]-1H-indol-4-yl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(707) N-[(S)-2,3-dihydroxy-propyl]-N-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-acétamide ;

(708) ester tert-butyle de l'acide {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-méthyl-benzyl}-(2-hydroxy-éthyl)-carbamique;

- (709) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3,5-diméthyl-phényl}-N-(4-hydroxy-cyclohexyl)-acétamide;
- (710) 2-cyclohexyl-8-((E)-2-{4-[(R)-2-(2-hydroxy-éthoxyméthyl)-5-oxo-pyrrolidin-1-yl]-2,6-diméthyl-phényl}-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (711) N-(2-hydroxy-éthyl)-N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-isobutylamide;
- (712) 8-[(E)-2-[4-((R)-5-hydroxyméthyl-3,3-diméthyl-2-oxo-pyrrolidin-1-yl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (713) 8-[(E)-2-[4-((R)-5-hydroxyméthyl-3-méthyl-2-oxo-pyrrolidin-1-yl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (714) 3-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-1-(2-hydroxy-éthyl)-5,5-diméthyl-imidazolidine-2,4-dione;
- (715) N-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro 4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-(4-hydroxy-cyclohexyl)-méthanesulfonamide;
- (716) N-(4-[(E)-2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-3,5-diméthyl-phényl)-N-[2-(2-hydroxy-éthoxy)-éthyl]-acétamide;
- (717) N-(2-fluoro-5-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-N-(2-hydroxy-éthyl)-acétamide;
- (718) N-(4-[(E)-2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl]-3,5-diméthyl-phényl)-N-(2-hydroxy-éthyl)-acétamide;
- (719) N-(2-hydroxy-éthyl)-N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-méthanesulfonamide;
- (720) N-(2-hydroxy-éthyl)-N-{3-méthyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-méthanesulfonamide;
- (721) N-(4-{2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-N-(2-hydroxy-éthyl)-acétamide;
- (722) 2-hydroxy-N-(2-hydroxy-éthyl)-N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-acétamide;
- (723) 1-[3,5-diméthyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl]-1-méthyl-urée;
- (724) 1-[3,5-diméthyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl]-1-[2-(2-hydroxy-éthoxy)-éthyl]-urée;
- (725) 1-[3,5-diméthyl-4-[(E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl]-1-(2-méthoxy-éthyl)-urée;
- (726) 1-(3,5-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-1-isopropyl-urée;
- (727) 1-(3-éthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-1-méthyl-urée;
- (728) 1-(3-méthoxy-5-méthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-1-méthyl-urée;
- (729) 1-cyanométhyl-1-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-urée;
- (730) 1-cyclopentyl-1-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-urée ;
- (731) 1-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-1-(2,2,2-trifluoro-éthyl)-urée;
- (732) 1-(4-{2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- (733) 1-((S)-2,3-dihydroxy-propyl)-1-(3,5-diméthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-urée;
- (734) N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3-méthyl-phényl}-N-cyclopentyl-acétamide;
- (735) (S)-2-amino-N-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-3,5-diméthyl-phényl}-3-méthyl-butylamide;
- (736) 2-{4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-indol-1-yl}-N-pyridin-4-yl-acétamide;
- (737) 2-(3-méthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-acétamide;

- (738) 2-cyclohexyl-8-[(E)-2-[4-(4,5-dihydro-thiazol-2-ylamino)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (739) 2-cyclohexyl-8-[(E)-2-[2-méthyl-4-(3-méthyl-oxetan-3-ylméthoxy)-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (740) 1-(3,5-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-cyclopropyl]-phényl)-1-méthyl-urée;
- (741) 1-(3,5-diméthyl-4-[(E)-1-méthyl-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-1-méthyl-urée;
- (742) N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-sulfamide;
- (743) N-(3-hydroxy-propyl)-N'-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-sulfamide;
- (744) N-méthyl-N-(3-méthyl-4-[(E)-3-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-propényl]-phényl)-acétamide;
- (745) 2-cyclohexyl-8-[2-(2-méthyl-1H-indol-4-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (746) 8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (747) 2-cyclohexyl-8-[2-(1-méthyl-1,2,3,4-tetrahydro-quinolin-5-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (748) 2-cyclohexyl-8-[2-[1-(3,4-dihydroxy-butyl)-1H-indol-4-yl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (749) 3-(2,6-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-imidazolidine-2,4-dione;
- (750) 2-(3-méthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- (751) N-(3-éthoxy-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- (752) N-(3-éthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- (753) N-(3-éthoxyméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- (754) N-(3-méthoxyméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;
- (755) N-(2-méthoxy-éthyl)-N-[3-méthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-acétamide;
- (756) N-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-N-(2-méthoxy-éthyl)-acétamide;
- (757) N-[2-(2-fluoro-éthoxy)-éthyl]-N-[3-méthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-acétamide;
- (758) N-[2-(2-hydroxy-éthoxy)-éthyl]-N-[3-méthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-acétamide;
- (759) N-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-N-méthyl-acétamide;
- (760) 3-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-imidazolidine-2,4-dione;
- (761) 8-(2-[4-[4-(2-fluoro-éthyl)-piperazine-1-carbonyl]-2,6-diméthyl-phényl]-éthanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (764) 8-[2-[2,6-diméthyl-4-(3-oxo-morpholin-4-yl)-phényl]-éthanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (765) 8-[2-[2,6-diméthyl-4-(3-oxo-morpholin-4-yl)-phényl]-éthanesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (766) N-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-N-[2-(2-fluoro-éthoxy)-éthyl]-acétamide;
- (767) 3-(3,5-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-5-méthyl-imidazolidine-2,4-dione;
- (768) acide 3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzoïque;
- (769) N'-acetyl-hydrazide de l'acide 3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-tria-

za-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-benzoïque;

(770) 8-[2-(4-hydroxyméthyl-2,6-diméthyl-phényl)-éthanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(771) 8-[2-[2,6-diméthyl-4-(2-oxo-pipéridin-1-yl)-phényl]-éthanesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(772) N-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-benzyl]-acétamide;

(773) N-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzyl)-acétamide;

(774) 3-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-benzyl]-imidazolidine-2,4-dione;

(775) 3-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-benzyl]-5,5-diméthyl-imidazolidine-2,4-dione;

(776) 8-[2-[2,6-diméthyl-4-(2-oxo-pyrrolidin-1-ylméthyl)-phényl]-éthanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(777) 1-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-benzyl]-pyrrolidine-2,5-dione;

(778) 3-(2,5-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-imidazolidine-2,4-dione;

(779) 3-(3,5-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzyl)-imidazolidine-2,4-dione;

(780) 3-(3,5-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzyl)-5,5-diméthyl-imidazolidine-2,4-dione;

(781) 8-[2-[2,6-diméthyl-4-(2-oxo-pyrrolidin-1-ylméthyl)-phényl]-éthanesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(782) 1-(3,5-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzyl)-pyrrolidine-2,5-dione;

(783) N-(2,6-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-N-méthyl-acétamide;

(784) N-[3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-benzyl]-N-méthyl-acétamide;

(785) N-(3,5-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzyl)-N-méthyl-acétamide;

(786) 1-(3,5-diméthyl-4-[2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-dihydro-pyrimidine-2,4-dione;

(787) triméthylhydrazide de l'acide 3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoropropyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-benzoïque;

(788) 8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthanesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(789) 8-[2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthanesulfonyl]-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(790) 3-{3-méthyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-imidazolidine-2,4-dione;

(791) 8-[2-[1-((R)-2,3-dihydroxy-propyl)-4,6-diméthyl-1H-indol-5-yl]-éthanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(792) 3-(3-méthyl-4-[2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-imidazolidine-2,4-dione;

(793) 8-[2-[2-méthyl-4-(3-oxo-morpholin-4-yl)-phényl]-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(794) N-(4-[2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3,5-diméthyl-phényl)-N-(2-hydroxy-éthyl)-acétamide;

(795) 3-{4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-trifluorométhyl-phényl}-imidazolidine-2,4-dione;

(796) acide 3,5-diméthyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-benzoïque;

(797) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-[2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(798) 8-[2-[4-(3,4-dihydroxy-butoxy)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (799) 3-(3,5-diméthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl)-imidazolidine-2,4-dione;
- (800) 3-(4-{2-[2-(4-fluoro-3-trifluorométhyl-phényl)]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-3,5-diméthyl-phényl)-imidazolidine-2,4-dione;
- 5 (801) 8-{2-[4-(3,4-dihydroxy-butoxy)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (802) 8-{2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (803) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-{2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 10 (804) 8-{2-[2,6-diméthyl-4-(2-oxa-7-aza-spiro[3.5]nonane-7-carbonyl)-phényl]-éthanesulfonyl}-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (812) N-(3-isopropoxy-4-{2-[2-(4-méthyl-cyclohexyl)]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl)-acétamide;
- 15 (817) 1-{3,5-diméthyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-1-méthyl-urée;
- (818) 1-[3,5-diméthyl-4-(2-(4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-phényl]-1-(2-fluoro-éthyl)-urée;
- (820) N-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl]-N-[2-(2-hydroxy-éthoxy)-éthyl]-acétamide;
- 20 (821) 8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (822) 8-(2-{1-[2-((S)-2,3-dihydroxy-propoxy)-éthyl]-1H-indol-4-yl]-éthanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 25 (823) 2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-8-(2-{1-[2-((2S,3S)-2,3,4-trihydroxy-butoxy)-éthyl]-1H-indol-4-yl]-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (824) N-((S)-2,3-dihydroxy-propyl)-N-(3,5-diméthyl-4-{2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl-phényl)-acétamide;
- (825) N-[2-(2-hydroxy-éthoxy)-éthyl]-N-{3-méthyl-4-[2-(2-nonyl-4-oxo-1,3,8-triazaspiro [4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-acétamide ;
- 30 (826) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl}-éthyl-phényl)-1-[2-(2-hydroxy-éthoxy)-éthyl]-urée;
- (827) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl]-1-(2-méthoxy-éthyl)-urée;
- 35 (828) 1-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl)-1-isopropyl-urée;
- (829) 1-(3-éthyl-4-{2-[2-(4-méthyl-cyclohexyl)]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl)-1-méthyl-urée;
- (830) 1-((S)-2,3-dihydroxy-propyl)-1-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl)-urée;
- 40 (831) 8-(2-[4-(4-hydroxy-4-hydroxyméthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (832) 3-(3,5-diméthyl-4-[2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl)-imidazolidine-2,4-dione;
- 45 (833) 5-tert-butoxyméthyl-3-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl)-imidazolidine-2,4-dione;
- (834) 2-(4-méthyl-cyclohexyl)-8-{2-[2-méthyl-4-(5-oxo-pyrazolidin-1-yl)-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (835) 2-cyclohexyl-8-[2-(1H-indol-7-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 50 (836) 2-cyclohexyl-8-[2-(6-trifluorométhyl-1H-indol-5-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (837) 2-cyclohexyl-8-[2-(6-méthyl-1H-indol-5-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (838) 2-cyclohexyl-8-(2-{1-[2-(2-hydroxy-éthoxy)-éthyl]-1H-indol-5-yl]-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (839) 2-cyclohexyl-8-[2-(6-trifluorométhyl-1H-benzimidazol-5-yl)-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 55 (840) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthanesulfonyl}-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (841) 2-cyclohexyl-8-{2-[4-(1,1-dioxo-1λ⁶-thiomorpholine-4-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-1,3,8-

triazaspiro[4.5]déc-1-én-4-one;

(842) 2-cyclohexyl-8-{2-[4-(2-diméthylamino-éthylamino)-2-méthyl-phényl]-éthanesulfonyl}-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(843) 8-{2-[2-méthyl-4-(4-méthyl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(844) 4-{2-(2-cyclohexyl-4-oxo-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl)-éthyl}-N-(2-diméthylamino-éthyl)-3-méthyl-benzènesulfonamide;

(845) 8-(2-{2-méthyl-4-[2-(4-méthyl-piperazin-1-yl)-2-oxo-éthylamino]-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl))-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(846) 3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-N-pyridin-4-yl-benzènesulfonamide;

(847) 2-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phénylamino)-acétamide;

(848) N-(1-benzyl-pipéridin-4-yl)-3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxyphényl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonamide;

(849) 8-{2-[4-(4-isopropyl-piperazine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(850) 8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(851) N-((R)-2,3-dihydroxy-propyl)-3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxyphényl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonamide;

(852) 3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-N-pyridin-3-ylméthyl-benzènesulfonamide;

(853) N-(4-hydroxy-cyclohexyl)-3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonamide;

(854) N-(2-hydroxy-1,1-bis-hydroxyméthyl-éthyl)-3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phényl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzènesulfonamide;

(855) 8-{2-[4-(3,4-dihydroxy-butoxy)-3,5-diméthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(856) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-{2-[2-méthyl-4-(pyrrolidine-1-carbonyl)-phényl]-éthanesulfonyl}-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(857) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-{2-[4-((R)-2-hydroxyméthyl-5-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthanesulfonyl}-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(858) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-{2-[4-((S)-2-hydroxyméthyl-5-oxo-pyrrolidin-1-yl)-2-méthyl-phényl]-éthanesulfonyl}-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(859) ester tert-butyle de l'acide 4-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phénoxy)-pipéridine-1-carboxylique;

(860) 8-{2-[4-(3,4-dihydroxy-butoxy)-3,5-difluoro-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(861) 2-cyclohexyl-8-{2-[2-méthyl-4-(2-oxo-azetidin-1-yl)-phényl]-éthanesulfonyl}-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(862) 1-méthyl-3-(3-méthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-imidazolidine-2,4-dione;

(863) 5,5-diméthyl-3-(3-méthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-imidazolidine-2,4-dione;

(864) 5,5-diméthyl-3-[3-méthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-imidazolidine-2,4-dione;

(865) 2-(4-méthyl-cyclohexyl)-8-{2-[2-méthyl-4-(2-oxo-oxazolidin-3-yl)-phényl]-éthanesulfonyl}-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(866) 2-(4-méthyl-cyclohexyl)-8-{2-[2-méthyl-4-(4-oxo-oxazolidin-3-yl)-phényl]-éthanesulfonyl}-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(867) 8-{2-[1-((R)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-éthanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(868) 8-(2-{4-[1H-imidazol-2-yl]-méthyl-amino}-2,6-diméthyl-phényl)-éthanesulfonyl)-2-(4-méthyl-cyclohexyl)-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

(869) 3-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-5,5-diméthyl-imidazolidine-2,4-dione;

(870) 2-(4-méthyl-cyclohexyl)-8-{2-[2-méthyl-4-(2-oxo-piperazin-1-yl)-phényl]-éthanesulfonyl}-1,3,8-triazaspiro[4.5]déc-1-én-4-one;

spiro[4.5]déc-1-én-4-one;

(872) 2-cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-5-yl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(873) N-(2-fluoro-5-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-N-(2-hydroxy-éthyl)-acétamide;

(874) N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-sulfamide;

(875) N-(3-hydroxy-propyl)-N'-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-sulfamide;

(876) 1-cyanométhyl-1-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-urée;

(877) 1-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-(2,2,2-trifluoro-éthyl)-urée;

(878) 1-(4-{2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-imidazolidine-2,4-dione;

(879) 1-(4-{2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-5-méthyl-imidazolidine-2,4-dione;

(880) 3-(4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-imidazolidine-2,4-dione;

(881) 3-(3-méthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-imidazolidine-2,4-dione;

(882) 3-(4-{2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-imidazolidine-2,4-dione;

(883) 3-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-imidazolidine-2,4-dione;

(884) 3-(4-{2-[2-(4-isopropyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-imidazolidine-2,4-dione;

(885) 3-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1,5,5-triméthyl-imidazolidine-2,4-dione;

(886) (S)-2-amino-3-hydroxy-N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-propionamide;

(887) 1-(4-{2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-5,5-diméthyl-imidazolidine-2,4-dione;

(888) N,N'-diméthyl-hydrazide de l'acide 3,5-diméthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzoïque;

(889) 8-{2-[2-méthyl-4-(pipéridin-4-yloxy)-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one hydrochloride ;

(890) 4-{2-[2-(4,4-difluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(891) 4-{2-[2-(4-méthanesulfonyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(892) 3,N,N-triméthyl-4-(2-[4-oxo-2-[(E)-2-(3-trifluorométhyl-phényl)-vinyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzamide;

(893) 3,N,N-triméthyl-4-[2-(4-oxo-2-phényléthynyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzamide;

(894) 4-{2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(895) 4-{2-[2-(4-tert-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(896) 4-{2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(897) 3,N,N-triméthyl-4-(2-[4-oxo-2-[4-(4,4,4-trifluoro-butyl)-cyclohexyl]-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzamide;

(898) 4-{2-[4-(cyclohexylméthyl-amino)-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]deca-1,3-diène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(899) 4-{2-[4-diméthylamino-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]deca-1,3-diène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(900) 4-{2-[4-[(Z)-hydroxyimino]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-

éthyl)-3,N,N-triméthyl-benzamide;

(901) 4-{2-[4-[(Z)-méthoxyimino]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(902) 4-(2-[2-[4-(2-méthoxy-éthyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,N,N-triméthyl-benzamide;

(903) N-{4-[2-(2-cyclopentyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-méthyl-phényl}-acétamide;

(904) N-(3-méthyl-4-[2-[4-oxo-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;

(905) N-(4-[2-[2-(4-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(906) N-(4-[2-[2-(4-cyano-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(907) N-(4-[2-[2-(3-cyano-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(908) N-(4-[2-[2-(2-cyano-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(909) N-(4-[2-[2-(4-tert-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(910) N-(4-[2-[2-(4-éthoxyméthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(911) N-(3-méthyl-4-[2-[4-oxo-2-(4-propoxy-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;

(912) N-(4-[2-[2-(4-butoxy-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(913) N-(4-[2-[2-(4-isopropoxyméthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(914) N-(4-[2-[2-[4-(3-fluoro-propoxy)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(915) N-(4-[2-[2-[4-(3-fluoro-propoxy)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(916) N-(4-[2-[2-[4-((E)-3,3-difluoro-propényl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(917) N-(4-[2-(2-cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-méthyl-phényl)-acétamide;

(918) N-(4-[2-(2-adamantan-1-yl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3-méthyl-phényl)-acétamide;

(919) N-(3-méthyl-4-(2-[4-oxo-2-[4-(2,2,2-trifluoro-éthoxyméthyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-acétamide;

(920) N-(4-[2-[2-[4-(4-chloro-phényl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(921) N-(3-méthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propoxy)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-acétamide;

(922) N-(3-méthyl-4-(2-[4-oxo-2-[4-(2,2,2-trifluoro-éthyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-acétamide;

(923) N-(3-méthyl-4-[2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;

(924) N-(3-méthyl-4-[2-[2-((1S,3R)-3-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;

(925) N-(4-[2-[2-(4,4-diméthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(926) N-(3-méthyl-4-[2-[4-oxo-2-(3,3,5,5-tetraméthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-phényl)-acétamide;

(927) N-(4-[2-[2-[4-(2-méthoxy-éthyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;

(928) ester tert-butyle de l'acide [3-méthyl-4-(2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzyl]-carbamique;

(929) N-(2-hydroxy-éthyl)-N-(3-méthyl-4-[2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-

ène-8-sulfonyl]-éthyl)-phényl)-isobutylamide;

(930) 2-hydroxy-N-(2-hydroxy-éthyl)-N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-acétamide;

(931) N-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-2,2,2-trifluoro-N-méthyl-acétamide;

(932) N-(3,5-diméthyl-4-{2-[2-((1S,3R)-3-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-2,2,2-trifluoro-N-méthyl-acétamide;

(933) N-(3,5-diméthyl-4-{2-[4-oxo-2-(3,3,5,5-tetraméthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-2,2,2-trifluoro-N-méthyl-acétamide;

(934) 1-{4-[2-(2-cycloheptyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl}-1-méthyl-urée;

(935) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-éthyl-phényl)-1-méthyl-urée;

(936) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-éthyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(937) 1-[2-chloro-3,5-diméthyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-éthyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(938) 1-(4-{2-[2-(4,4-diméthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(939) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propylidène)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(940) 1-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl]-1-méthyl-urée;

(941) 1-(3,5-diméthyl-4-{2-[8-oxo-6-(3-trifluorométhoxy-phényl)-2,5,7-triaza-spiro[3.4] oct-5-ène-2-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(942) N-(4-{(E)-1-fluoro-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl}-3-méthyl-phényl)-N-méthyl-acétamide;

(943) 3-[3-méthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-imidazolidine-2,4-dione;

(944) 3-[3-méthyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-imidazolidine-2,4-dione;

(945) 3-[4-(2-{2-[4-((E)-3,3-difluoro-propényl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-imidazolidine-2,4-dione;

(946) 3-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-imidazolidine-2,4-dione;

(947) 3-(3-méthyl-4-{2-[4-oxo-2-(4-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-imidazolidine-2,4-dione;

(948) 3-[3-méthyl-4-(2-{4-oxo-2-[4-(2,2,2-trifluoro-éthyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-imidazolidine-2,4-dione;

(949) 3-(3-méthyl-4-{2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-imidazolidine-2,4-dione;

(950) 3-(3-méthyl-4-{2-[2-((1S,3R)-3-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-imidazolidine-2,4-dione;

(951) 3-(4-{2-[2-(4,4-diméthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-imidazolidine-2,4-dione;

(952) 3-(3-méthyl-4-{2-[4-oxo-2-(3,3,5,5-tetraméthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-imidazolidine-2,4-dione;

(953) N-(2-chloro-4-{2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-5-méthyl-phényl)-acétamide;

(954) N-(2-chloro-4-{2-[2-(4-éthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-acétamide;

(955) N-[4-(2-{2-[4-(3,3-difluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;

(956) 3-[4-(2-{2-[4-(3,3-difluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-imidazolidine-2,4-dione;

(957) 2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-N-phényl-benzamide;

(958) 8-(2-{2-méthyl-4-[4-(1-méthyl-pipéridin-4-yl)-piperazine-1-carbonyl]-phényl}-éthanesulfonyl)-2-(3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

- (959) 8-[(E)-2-[4-((R)-3-fluoro-pyrrolidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (960) 8-[(E)-2-[4-(4-hydroxyméthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (961) 3,5,N,N-tetraméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;
- (962) 3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl]-benzamide;
- (963) 8-[(E)-2-[4-(3-fluoro-azetidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (964) 8-[2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (965) 8-[2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (966) 8-[2-[4-(4-hydroxyméthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (967) 8-[(E)-2-[2,6-diméthyl-4-(2-oxa-6-aza-spiro[3.3]heptane-6-carbonyl)-phényl]-éthènesulfonyl]-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (968) 8-[(E)-2-[2,6-diméthyl-4-(3-oxo-piperazine-1-carbonyl)-phényl]-éthènesulfonyl]-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (969) 8-[(E)-2-[4-((3R,5S)-3,5-diméthyl-piperazine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (970) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-[(E)-2-[4-(3-hydroxy-3-méthyl-azetidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (971) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-[(E)-2-[4-(4-hydroxy-4-hydroxyméthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (972) 8-[(E)-2-[2,6-diméthyl-4-(4-oxo-pipéridine-1-carbonyl)-phényl]-éthènesulfonyl]-2-(4-fluoro-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (973) 2-[4-(3,3-difluoro-allyl)-cyclohexyl]-8-[2-[4-(4-fluoro-4-hydroxyméthyl-pipéridine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (974) 2-[4-(3,3-difluoro-allyl)-cyclohexyl]-8-[2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (975) 2-(4-éthyl-cyclohexyl)-8-[2-[4-(2-fluoro-éthyl)-piperazine-1-carbonyl]-2-méthyl-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (976) 2-(4-éthyl-cyclohexyl)-8-[2-[2-méthyl-4-(4-oxetan-3-yl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (977) 8-[2-[2-méthyl-4-(pyrrolidine-1-carbonyl)-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (978) 8-[2-[4-(azetidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro [4.5]déc-1-én-4-one;
- (979) 8-[2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2-méthyl-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (980) 8-[2-[2,6-diméthyl-4-(pyrrolidine-1-carbonyl)-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (981) 8-[2-[4-((R)-3-hydroxy-pyrrolidine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (982) 8-[2-[4-(azetidine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (983) 8-[2-[2,6-diméthyl-4-(pipéridine-1-carbonyl)-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (984) 8-[2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (985) 8-[2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (986) 8-[2-[2,6-diméthyl-4-(4-méthyl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl]-2-nonyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (987) N,N-diméthyl-2-(3-méthyl-4-[(E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-phényl)-acétamide;

- (988) N-méthoxy-2-(3-méthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
- (989) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (990) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (991) 8-((E)-2-[2,6-diméthyl-4-(4-méthyl-piperazine-1-carbonyl)-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (992) 8-((E)-2-[4-(isoxazolidine-2-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (993) 8-((E)-2-[4-(4-((R)-2,3-dihydroxy-propoxy)-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (994) 3,5,N,N-tetraméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzamide;
- (995) 8-[2-[2,6-diméthyl-4-(4-méthyl-piperazine-1-carbonyl)-phényl]-éthanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (996) hydrazide de l'acide 3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzoïque;
- (997) 8-((E)-2-[2,6-diméthyl-4-(pyrazolidine-1-carbonyl)-phényl]-éthènesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (998) N-méthoxy-3,5,N-triméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzamide ;
- (999) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1000) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1001) 8-((E)-2-[4-(isoxazolidine-2-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1002) 8-((E)-2-[4-(1,1-dioxo-1 λ^6 -thiomorpholine-4-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1003) 3,5,N,N-tetraméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-benzamide;
- (1006) 2-cyclohexyl-8-((E)-2-[4-(4-hydroxy-4-trifluorométhyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1007) 2-cyclohexyl-8-((E)-2-[2,6-diméthyl-4-(2-oxo-oxazolidine-3-carbonyl)-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1008) 2-(4-butyl-cyclohexyl)-8-((E)-2-[4-[4-(2-hydroxy-éthoxy)-pipéridine-1-carbonyl]-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1009) 2-(4-butyl-cyclohexyl)-8-((E)-2-[4-[4-(2-hydroxy-éthoxy)-éthoxy]-pipéridine-1-carbonyl]-2,6-diméthyl-phényl)-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1010) 8-((E)-2-[4-[4-((R)-2,3-dihydroxy-propoxy)-pipéridine-1-carbonyl]-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1011) 2-amino-N-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-N-méthyl-acétamide;
- (1012) ester 2-hydroxy-éthyle de l'acide (3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-méthyl-carbamique;
- (1013) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1,3,3-triméthyl-urée;
- (1014) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1,3-diméthyl-urée;
- (1015) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;
- (1016) 8-((E)-2-(2,6-diméthyl-4-méthylamino-phényl)-éthènesulfonyl)-2-(3-trifluorométhylsulfanyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1017) 1-(4-((E)-2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;
- (1018) 1-[3,5-diméthyl-4-((E)-2-(4-oxo-2-[3-(1,1,2,2-tetrafluoro-éthoxy)-phényl]-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;

- (1019) 1-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[3-(4,4,4-trifluoro-butoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;
- (1020) 1-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(4,4,4-trifluoro-butoxy)-phényl]-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;
- 5 (1021) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(4-pentyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;
- (1022) 1-(3,5-diméthyl-4-((E)-2-[2-(7-méthylsulfanyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;
- (1023) 1-[3,5-diméthyl-4-((E)-2-[2-[8-(3-méthyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;
- 10 (1024) 1-[3,5-diméthyl-4-((E)-2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl)-phényl]-1-méthyl-urée;
- (1025) 1-[3,5-diméthyl-4-((E)-2-(2-non-3-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl)-phényl]-1-méthyl-urée;
- 15 (1026) 1-[3,5-diméthyl-4-((E)-2-[2-[10-(3-méthyl-oxetan-3-yl)-decyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;
- (1027) 1-[3,5-diméthyl-4-((E)-2-(2-non-1-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl)-phényl]-1-méthyl-urée;
- (1028) 1-(4-{2-[2-(4-fluoro-3-trifluorométhoxy-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- 20 (1029) 1-(4-{2-[2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- (1030) 1-(4-{2-[2-(4-tert-butyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- 25 (1031) 1-[3,5-diméthyl-4-(2-{2-[8-(3-méthyl-oxetan-3-yl)-octyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;
- (1032) 1-[3,5-diméthyl-4-(2-{2-[10-(3-méthyl-oxetan-3-yl)-decyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;
- (1033) ester tert-butyle de l'acide 2-(8-{2-[4-(tert-butoxycarbonyl-méthyl-amino)-2,6-diméthyl-phényl]-éthane-sulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-pyrrolidine-1-carboxylique;
- 30 (1034) ester tert-butyle de l'acide (4-{2-[2-(1,1-difluoro-éthyl)-4-oxo-1,3,8-triazaspiro [4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-méthyl-carbamique;
- (1035) 8-[2-(2,6-diméthyl-4-méthylamino-phényl)-éthanesulfonyl]-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- 35 (1036) 8-[2-(2,6-diméthyl-4-méthylamino-phényl)-éthanesulfonyl]-2-(3-trifluorométhylsulfanyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1037) 8-[2-(2,6-diméthyl-4-méthylamino-phényl)-éthanesulfonyl]-2-(4-isopropylidène-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1038) 1-[3,5-diméthyl-4-(2-[4-oxo-2-[3-(1,1,2,2-tetrafluoro-éthoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;
- 40 (1039) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(5,6,7,8-tetrahydro-naphthalèn-2-yl)-1,3,8-triazaspiro [4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;
- (1040) 1-(4-{2-[2-((1S,3S,SR)-3,5-diméthyl-cyclohexyl)-4-oxo-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- 45 (1041) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(1-propyl-butyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;
- (1042) 1-[3,5-diméthyl-4-(2-(4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl)-phényl]-1-méthyl-urée;
- (1043) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(6,6,7,7,7-pentafluoro-heptyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;
- 50 (1044) N-(3,5-diméthyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-N-méthyl-acétamide;
- (1045) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(7,7,7-trifluoro-heptyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;
- 55 (1046) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;
- (1047) N-(3,5-diméthyl-4-{2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-N-méthyl-acétamide;

- (1048) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[1-(4,4,4-trifluoro-butyl)-cyclopropyl]-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl}-éthyl)-phényl]-1-méthyl-urée;
- (1050) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;
- (1051) 1-(4-{2-[2-(cis-3,4-diméthyl-cyclopentyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- (1052) 1-{4-[2-(2-dicyclopropylméthyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl}-1-méthyl-urée;
- (1053) 1-{4-[2-(2-bicyclo[3.3.1]non-9-yl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl}-1-méthyl-urée;
- (1054) 1-{4-[2-((1R,5S)-2-bicyclo[3.2.1]oct-3-yl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl}-1-méthyl-urée;
- (1055) 1-(3,5-diméthyl-4-{2-[2-(3-méthyl-cyclopentyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;
- (1056) 1-{4-[2-(2-bicyclo[2.2.1]hept-7-yl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl}-1-méthyl-urée;
- (1057) 1-(4-{2-[2-((1S,2S)-2-hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- (1058) 1-{3,5-diméthyl-4-[2-(4-oxo-2-spiro[2.5]oct-6-yl-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-1-méthyl-urée;
- (1059) 1-(4-{2-[2-(4-difluorométhylène-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- (1060) N-[4-(2-{2-[4-(2,2-difluoro-éthyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1061) N-[4-(2-{2-[4-(2-fluoro-éthyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1062) N-(4-{2-[2-(4-butyl-4-fluoro-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-acétamide;
- (1063) N-[4-(2-{2-[4-((E)-3-fluoro-propényl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1064) N-[4-(2-{2-[4-(3-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1065) N-[4-(2-{2-[4-(3-chloro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1066) N-[4-(2-{2-[4-(3-fluoro-propylidène)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1067) N-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1068) N-[4-(2-{2-[4-(3,3-difluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1069) N-[3-méthyl-4-(2-{4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-acétamide;
- (1070) N-[4-(2-{2-[4-((E)-3-fluoro-allyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1071) N-[4-(2-{2-[4-(2,2-difluoro-éthyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1072) N-[4-(2-{2-[4-(2-fluoro-éthyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1073) N-[4-(2-{2-[4-fluoro-4-(3-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1074) N-(4-{2-[2-(4-éthynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-acétamide;
- (1075) N-(4-{2-[2-(4-difluorométhyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-phényl)-acétamide;
- (1076) N-[4-(2-{2-[4-(3,3-difluoro-propylidène)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1077) N-[4-(2-{2-[4-(2-fluoro-propyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;

- (1078) N-[4-(2-[2-[4-(1-fluoro-1-méthyl-éthyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-méthyl-phényl]-acétamide;
- (1079) N-(4-[2-[2-(4-butyldène-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3-méthyl-phényl)-acétamide;
- (1080) N-[3-méthyl-4-(2-[4-oxo-2-[4-(2,2,3,3,3-pentafluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-acétamide;
- (1081) 1-{3,5-diméthyl-4-[2-(2-non-4-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-1-méthyl-urée;
- (1082) 1-{3,5-diméthyl-4-[2-(2-non-3-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-1-méthyl-urée;
- (1083) 1-{3,5-diméthyl-4-[2-(2-non-1-ynyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-1-méthyl-urée;
- (1084) 1-(3,5-diméthyl-4-{1-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-cyclopropylméthyl}-phényl)-1-méthyl-urée;
- (1085) 1-(4-{2,2-difluoro-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;
- (1086) 1-[4-[2,2-difluoro-2-(2-nonyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl]-1-méthyl-urée;
- (1087) 1-{3,5-diméthyl-4-[2-(4-oxo-2-[1,1';2',1'']terphényl-3-yl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl}-1-méthyl-urée;
- (1088) 1-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(4,4,4-trifluoro-butyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;
- (1089) 1-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[3-(4,4,4-trifluoro-butyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;
- (1090) 1-(3,5-diméthyl-4-[2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-propyl]-phényl)-1-méthyl-urée;
- (1091) 2-[4-fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phényl]-8-((E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1092) 2-[4-fluoro-3-(4,4,5,5,5-pentafluoro-pentyloxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1093) 2-(4-chloro-3-trifluorométhoxy-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1094) 2-(3-fluoro-4-trifluorométhoxy-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1095) 2-(3,4-bis-trifluorométhyl-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1096) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1097) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(2,2,2-trifluoro-éthyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1098) 2-(4-fluoro-3-méthyl-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1099) 2-(4-fluoro-3-trifluorométhoxy-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1100) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(1,1,2,2-tetrafluoro-éthoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1101) 8-{1,1-difluoro-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1102) 2-(4-fluoro-3-trifluorométhoxy-phényl)-8-((E)-2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1103) 8-((E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1104) 6-(4-méthyl-cyclohexyl)-2-(2-naphthalèn-1-yl)-éthanesulfonyl)-2,5,7-triaza-spiro[3.4]oct-5-én-8-one;
- (1106) ester tert-butyle de l'acide (11-{8-[(E)-2-(2,6-diméthyl-4-méthylamino-phényl)-éthènesulfonyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl}-undecyl)-carbamique;
- (1107) 8-[(E)-2-(2,6-diméthyl-4-méthylamino-phényl)-éthènesulfonyl]-2-(9-hydroxy-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1108) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sul-

fonyl]-vinyl)-phényl)-1-méthyl-urée;

(1109) acide 14-(8-((E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-tétradécanoïque;

(1110) 1-(3,5-diméthyl-4-((E)-2-[2-(4-[1,1,1,2-H₃]méthyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1111) 1-(4-((E)-2-[2-(4-fluoro-3-trifluorométhoxy-phényl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1112) 1-(3,5-diméthyl-4-((E)-2-[2-(3-nonafluorobutyl-phényl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1113) 1-(4-((E)-2-[2-(4-chloro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1114) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(4'-propyl-biphényl-3-yl)-1,3,8-triaza-spiro 4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1115) 1-(4-((E)-2-[2-(6-éthoxy-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1116) 1-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[3-(6,6,6-trifluoro-hexyl)-phényl]-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1117) 1-[3,5-diméthyl-4-((E)-2-[4-oxo-2-[4-(6,6,6-trifluoro-hexyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1118) 1-(4-((E)-2-[2-(11-fluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1119) 1-[4-((E)-2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1120) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-((E)-6-phényl-hex-5-enyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1121) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-((E)-9-phényl-non-8-enyl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1122) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(5-propylsulfanyl-pentyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1123) 1-(4-((E)-2-[2-(7-méthoxy-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1124) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(5-propoxy-pentyl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1125) 1-(4-((E)-2-[2-(11,11-difluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1126) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(3-propyl-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1127) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(2-propyl-benzofuran-6-yl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1128) 1-(4-((E)-2-[2-(3-méthoxy-4-pentyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1129) 1-(4-((E)-2-[2-(4-[1,1,2,2,2-H₅]éthyl-cyclohex-3-enyl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1130) 1-(4-((E)-2-[2-(4-[1,1,2,2,2-H₅]éthyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-vinyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1131) acide 12-(8-((E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-dodécanoïque;

(1132) ester éthyle de l'acide 12-(8-((E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-dodécanoïque;

(1133) 1-(3,5-diméthyl-4-((E)-2-[4-oxo-2-(5-phényl-pentyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1134) 1-(3,5-diméthyl-4-((Z)-2-[4-oxo-2-(5-phényl-pentyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-1-méthyl-urée;

(1135) 1-[3,5-diméthyl-4-(2-[4-oxo-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1136) 1-(3,5-diméthyl-4-[2-[4-oxo-2-(3'-propyl-biphényl-3-yl)-1,3,8-triazaspiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1137) 1-(3,5-diméthyl-4-[2-[4-oxo-2-(4-triméthylsilanyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfo-

nyl]-éthyl)-phényl)-1-méthyl-urée;

(1138) 1-(4-{2-[2-((1R,3R,3,5)-3,5-bis-trifluorométhyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1139) 1-(3,5-diméthyl-4-[2-[4-oxo-2-(4-triméthylsilanyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1140) 1-{3,5-diméthyl-4-[2-(4-oxo-2-tridecyl-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl)-1-méthyl-urée;

(1141) 1-{3,5-diméthyl-4-[2-(4-oxo-2-undecyl-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl)-1-méthyl-urée;

(1142) 1-{3,5-diméthyl-4-[2-(2-octyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl)-1-méthyl-urée;

(1143) 1-(4-{2-[2-((1S,3R)-3-hexyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1144) 1-{3,5-diméthyl-4-(2-{2-[3-(3-méthyl-butyl)-phényl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1145) 1-(4-{2-[2-((1S,3R)-3-butyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1146) 1-{3,5-diméthyl-4-(2-{2-[(1S,3S)-3-(3-méthyl-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1147) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(3,3,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1148) 1-(4-{2-[2-(4-[1,1,2,2,2-²H₅]éthyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1149) 1-(4-{2-[2-(4-[1,1,2,2,2-²H₅]éthyl-cyclohex-3-enyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1150) 1-(4-{2-[2-(4-chloro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1151) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(1,9,9,9-tetrafluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1152) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(4'-propyl-biphényl-3-yl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1153) 1-(4-{2-[2-(11-fluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1154) 1-(4-{2-[2-(6-éthylsulfanyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée ;

(1155) 1-{3,5-diméthyl-4-[2-(4-oxo-2-[1,1',3',1'']terphényl-3-yl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-phényl)-1-méthyl-urée;

(1156) 1-(4-{2-[2-(11,11-difluoro-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3,5-diméthyl-phényl)-1-méthyl-urée;

(1157) 1-{3,5-diméthyl-4-[(E)-2-(4-oxo-2-[1,1',3',1'']terphényl-3-yl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-phényl)-1-méthyl-urée;

(1158) 8-[(E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-[3-(4,4,5,5,5-pentafluoro-pentyloxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1159) 2-(4-fluoro-3-trifluorométhoxy-phényl)-8-[(E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1160) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1161) 2-(4-fluoro-3-trifluorométhoxy-phényl)-8-[(E)-2-[4-(4-hydroxyméthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro 4.5]déc-1-én-4-one;

(1162) 2-(4-fluoro-3-trifluorométhyl-phényl)-8-[(E)-2-[4-(4-hydroxyméthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1163) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1164) 8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthènesulfonyl]-2-(4-fluoro-3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1165) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[(E)-2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthènesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1166) 8-[(E)-2-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthènesulfonyl]-2-(4-fluoro-3-trifluorométhoxy-

phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1167) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1168) acide {4-[(E)-2-(2-cyclohexyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-vinyl]-indol-1-yl}-acétique ;

(1169) acide [3-(3-méthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-ureido]-acétique;

(1170) acide 12-(8-{(E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-dodécanoïque;

(1171) acide 10-(8-{(E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthènesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-décanoïque;

(1172) amide de l'acide 10-(8-{(E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthènesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-décanoïque;

(1173) amide de l'acide 12-(8-{(E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthènesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-dodécanoïque;

(1174) amide de l'acide 12-(8-{(E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-dodécanoïque;

(1175) amide de l'acide 14-(8-{(E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthènesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-tetradécanoïque;

(1176) amide de l'acide 14-(8-{(E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthanesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-tetradécanoïque;

(1178) 8-[2-(2-amino-5,7-diméthyl-benzoxazol-6-yl)-éthanesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1179) 2-cyclohexyl-8-[2-[2,6-diméthyl-4-(2-oxo-oxazolidine-3-carbonyl)-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1180) N-méthoxy-3,5,N-triméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(1181) 8-{2-[4-(isoxazolidine-2-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1182) 8-{2-[4-(isoxazolidine-2-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1183) 8-{2-[4-(1,1-dioxo-1λ⁶-thiomorpholine-4-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1184) 8-(2-[4-[4-((R)-2,3-dihydroxy-propoxy)-pipéridine-1-carbonyl]-2,6-diméthyl-phényl]-éthanesulfonyl)-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1185) 2-(3,4-dichloro-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1186) 2-(3-chloro-4-fluoro-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1187) [3-(8-{(E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-phényl]-acetonitrile;

(1188) 2-(3-chloro-4-trifluorométhyl-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro [4.5]déc-1-én-4-one;

(1189) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-pentafluorosulfanyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1190) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-pentafluorosulfanyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1191) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(3-trifluorométhyl-phénoxy-méthyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1192) 2-(4-fluoro-2,3-diméthyl-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1193) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-méthyl-4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1194) 2-benzo[1,3]dioxol-5-yl-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1195) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-pentafluoroéthyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one ;

(1196) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-pen-

tafluoroéthyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1197) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(2,2,2-trifluoro-1,1-diméthyl-éthoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1198) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1199) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(2,2,3,3-tetrafluoro-propoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1200) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(2,2,2-trifluoro-éthoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1201) 2-[3-chloro-4-(2,2,3,3-tetrafluoro-propoxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1202) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(4,4,5,5-pentafluoro-pentyloxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1203) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(2,2,3,3,3-pentafluoro-propoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1204) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(4,4,4-trifluoro-butoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1205) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(3-trifluorométhyl-phénoxy)méthyl]-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1206) 2-[3-(1,1-difluoro-éthyl)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1207) 2-[3-fluoro-4-(4,4,5,5-pentafluoro-pentyloxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1208) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(1H-indol-6-yl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1209) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(2,2,2-trifluoro-1,1-diméthyl-éthoxy)-4-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1210) 2-[4-chloro-3-(2,2,2-trifluoro-1,1-diméthyl-éthoxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1211) 2-[3-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-4-fluoro-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1212) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(2,2,3,3-tetrafluoro-2,3-dihydro-benzo[1,4]dioxin-6-yl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1213) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(1,1,2,2-tetrafluoro-éthoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1214) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-méthyl-4-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1215) 2-[4-fluoro-3-(3-fluoro-propoxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1216) 2-[3-chloro-4-(2,2,2-trifluoro-éthoxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1217) 2-(3-difluorométhyl-4-fluoro-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1218) 2-[4-fluoro-3-(4,4,4-trifluoro-butoxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1219) 2-(4-difluorométhyl-3-fluoro-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1220) 2-[4-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-3-trifluorométhyl-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1221) 2-[4-[1,1,2,2,2,2-²H₆]éthoxy-3-trifluorométhyl-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1222) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-[1,2,2,2,2,2-²H₇]isopropoxy-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1223) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1224) 2-[3-[1,1,2,2,2-²H₅]éthoxy-4-fluoro-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1225) 2-(4-fluoro-3-[1,2,2,2,2,2-²H₇]isopropoxy-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbo-

nyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1226) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-[3-(2,2,3,3-tetrafluoro-propoxy)-5-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1227) 2-[4-chloro-3-(4,4,4-trifluoro-butoxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-
 2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1228) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-[3-(2,2,3,3,3-pentafluoro-propoxy)-4-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1229) 2-[4-chloro-3-(2,2,3,3-tetrafluoro-propoxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbo-
 nyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1230) 2-[4-chloro-3-(2,2,3,3,3-pentafluoro-propoxy)-phényl]-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-car-
 bonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1231) 2-(3-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-4-chloro-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbo-
 nyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1232) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-[3-(4,4,4-trifluoro-butoxy)-4-trifluorométhoxy-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1233) 2-(3-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-5-trifluorométhyl-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-
 1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1234) 8-{2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-trifluoro-
 méthoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1235) 8-{2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(3-méthyl-
 4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1236) 2-(3-chloro-4-trifluorométhyl-phényl)-8-{2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-
 phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1237) 2-(3,4-diméthyl-phényl)-8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-
 éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1238) 8-((E)-2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-mé-
 thoxy-3-méthyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1239) 2-(4-fluoro-2,5-diméthyl-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-di-
 méthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1240) 2-(4-fluoro-2,3-diméthyl-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-di-
 méthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1241) 2-(3-chloro-4-trifluorométhyl-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-
 diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1242) 2-(3-chloro-4-fluoro-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-
 phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1243) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-(3-pentafluoroéthyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1244) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-(4-pentafluoroéthyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1245) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-(4-pentafluorosulfanyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1246) 2-[3-chloro-4-(2,2,2-trifluoro-éthoxy)-phényl]-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbo-
 nyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1247) 2-[3-chloro-4-(2,2,3,3-tetrafluoro-propoxy)-phényl]-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-
 carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1248) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-[4-(2,2,2-trifluoro-éthoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1249) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-[4-(2,2,3,3,3-pentafluoro-propoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1250) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-[4-(4,4,4-trifluoro-butoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1251) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-[4-(2,2,3,3-tetrafluoro-propoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1252) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-
 2-[4-(4,4,5,5,5-pentafluoro-pentyloxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1253) 2-(4-[1,1,2,2,2-²H₅]éthoxy-3-trifluorométhyl-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-
 carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1254) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-

- 2-(3-méthyl-4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1255) 2-(4-difluorométhyl-3-fluoro-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1256) 2-(3-difluorométhyl-4-fluoro-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1257) 2-(3,4-dichloro-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1258) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(3-méthyl-4-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1259) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(2-méthyl-4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1260) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-fluoro-4-(2,2,2-trifluoro-éthoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1261) 2-(4-[1,1,2,2,3,3,4,4,4-²H₉]butoxy-3-trifluorométhyl-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1262) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-[1,2,2,2,2,2-²H₇]isopropoxy-3-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1263) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-fluoro-3-(4,4,5,5-pentafluoro-pentyloxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1264) 2-[4-fluoro-3-(3-fluoro-propoxy)-phényl]-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1265) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[4-(2,2,2-trifluoro-1,1-diméthyl-éthoxy)-3-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1266) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(2,2,2-trifluoro-1,1-diméthyl-éthoxy)-4-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1267) 2-[3-(1,1-difluoro-éthyl)-phényl]-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1268) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(1H-indol-6-yl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1269) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-(4-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1270) 2-[4-chloro-3-(2,2,3,3,3-pentafluoro-propoxy)-phényl]-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1271) 2-[4-chloro-3-(2,2,2-trifluoro-1,1-diméthyl-éthoxy)-phényl]-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1272) 2-(3-[1,1,2,2,2-²H₅]éthoxy-4-fluoro-phényl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1273) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(4,4,4-trifluoro-butoxy)-4-trifluorométhoxy-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1274) 8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1275) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-[2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1276) 8-[2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-(4-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-en-4-one;
- (1277) 8-[2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-(3-méthyl-4-trifluorométhyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1278) 2-(3-chloro-4-trifluorométhyl-phényl)-8-[2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1279) 8-[2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-[3-(2,2,3,3-tetrafluoro-propoxy)-4-trifluorométhyl-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1280) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-((E)-2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1281) 8-[2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl]-2-(2,2,3,3-tetrafluoro-2,3-dihydro-benzo[1,4]dioxin-6-yl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
- (1282) [3-(8-((E)-2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthènesulfonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-phényl]-acetonitrile,
- (1283) 8-((E)-2-[2-méthyl-4-(pipéridin-4-yloxy)-phényl]-éthènesulfonyl)-2-(3-trifluorométhoxy-phényl)-1,3,8-

triaza-spiro[4.5]déc-1-én-4-one;
 (1284) 8-[2-(4-aminométhyl-2-méthyl-phényl)-éthanesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-
 triaza-spiro[4.5]déc-1-én-4-one;
 (1285) 2-(4-méthyl-cyclohexyl)-8-[(E)-2-[2-méthyl-4-(2-oxo-piperazin-1-yl)-phényl]-éthènesulfonyl]-1,3,8-tria-
 5 za-spiro[4.5]déc-1-én-4-one;
 (1286) 8-[2-(2,6-diméthyl-4-méthylamino-phényl)-éthanesulfonyl]-2-pyrrolidin-2-yl-1,3,8-triaza-spiro[4.5]déc-
 1-én-4-one hydrochloride;
 (1287) 2-(1,1-difluoro-éthyl)-8-[2-(2,6-diméthyl-4-méthylamino-phényl)-éthanesulfonyl]-1,3,8-triaza-
 spiro[4.5]déc-1-én-4-one hydrochloride;
 10 (1288) 8-[(E)-2-(2,6-diméthyl-4-[1,1,1-²H₃]-méthylamino-phényl)-éthènesulfonyl]-2-(4-fluoro-3-trifluorométhyl-
 phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one hydrochloride;
 (1289) 8-[(E)-2-(2,6-diméthyl-4-méthylaminométhyl-phényl)-éthènesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-
 triaza-spiro[4.5]déc-1-én-4-one;
 (1290) 8-[(E)-2-(4-aminométhyl-2,6-diméthyl-phényl)-éthènesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-
 15 1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 (1291) 8-[(E)-2-(4-aminométhyl-2,6-diméthyl-phényl)-éthènesulfonyl]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-
 spiro[4.5]déc-1-én-4-one;
 (1292) 8-[(E)-2-(2,6-diméthyl-4-méthylaminométhyl-phényl)-éthènesulfonyl]-2-[4-(3,3,3-trifluoro-propyl)-cyclo-
 hexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;
 20 (1293) 8-[2-(4-amino-3-chloro-2-méthyl-phényl)-éthanesulfonyl]-2-(4-éthyl-cyclohexyl)-1,3,8-triaza-
 spiro[4.5]déc-1-én-4-one;
 (1294) N-(1-acetyl-pipéridin-4-yl)-N-(3-méthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhoxy-phényl)-1,3,8-triaza-
 spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl)-acétamide;
 (1295) 8-[2-[4-(4,5-dihydro-thiazol-2-ylamino)-2-méthyl-phényl]-éthanesulfonyl]-2-(4-éthyl-cyclohexyl)-1,3,8-
 25 triaza-spiro[4.5]déc-1-én-4-one;
 (1296) N-[4-(2-[2-[4-(4-fluoro-butyl)-cyclohexyl]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-3-
 méthyl-phényl]-acétamide;
 (1297) 1-(3,5-diméthyl-4-{2-[2-((1S,3R)-3-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfo-
 nyl]-éthyl)-phényl)-1-méthyl-urée;
 30 (1298) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(3,3,5,5-tetraméthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfo-
 nyl]-éthyl)-phényl)-1-méthyl-urée;
 (1299) 3-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-
 éthyl)-phényl)-5-hydroxyméthyl-imidazolidine-2,4-dione;
 (1300) 1-[3,5-diméthyl-4-((E)-2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-
 35 ène-8-sulfonyl]-vinyl)-benzyl]-1-méthyl-urée;
 (1301) 1-(3,5-diméthyl-4-((E)-2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vi-
 nyl)-benzyl)-1-méthyl-urée;
 (1302) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfo-
 nyl]-éthyl)-phényl)-1-méthyl-urée;
 40 (1303) 1-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(8,8,9,9,9-pentafluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sul-
 fonyl]-vinyl)-phényl)-1-méthyl-urée;
 (1304) 1-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(9,9,9-trifluoro-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vi-
 nyl)-phényl)-1-méthyl-urée;
 (1305) 1-(3,5-diméthyl-4-[(E)-2-[4-oxo-2-(3-trifluorométhylsulfanyl-phényl)-1,3,8-triaza-spiro [4.5]déc-1-ène-8-
 45 sulfonyl]-vinyl)-phényl)-1-méthyl-urée;
 (1306) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(3-trifluorométhylsulfanyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sul-
 fonyl]-éthyl)-phényl)-1-méthyl-urée;
 (1307) 1-(4-[(E)-2-[2-(11-hydroxy-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-3,5-dimé-
 thyl-phényl)-1-méthyl-urée;
 50 (1308) ester tert-butyle de l'acide [11-(8-[(E)-2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthènesulfonyl]-4-
 oxo-1,3,8-triaza-spiro[4.5]déc-1-én-2-yl)-undecyl]-carbamique;
 (1309) 1-(4-[(E)-2-[2-(9-hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl]-3,5-diméthyl-
 phényl)-1-méthyl-urée;
 (1310) 1-(4-[2-[2-(4-isopropylidène-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3,5-
 55 diméthyl-phényl)-1-méthyl-urée;
 (1311) 1-(4-[2-[2-(1,1-difluoro-éthyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl]-3,5-diméthyl-
 phényl)-1-méthyl-urée;
 (1312) 1-(4-[(E)-2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vi-

nyl)-3,5-diméthyl-phényl)-1-[1,1,1-²H₃]méthyl-urée;

(1313) [3,5-diméthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-benzyl]-urée;

(1314) (3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzyl)-urée;

(1315) 1-(4-{(E)-2-[2-(11-amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1316) 1-[3,5-diméthyl-4-((E)-2-{4-oxo-2-[5-(propane-1-sulfinyl)-pentyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;

(1317) 1-(4-{2-[2-(6-éthanesulfinyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1318) 1-[3,5-diméthyl-4-((E)-2-{4-oxo-2-[5-(propane-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl)-phényl]-1-méthyl-urée;

(1319) 1-(4-{(E)-2-[2-(7-méthanesulfonyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1320) 1-(4-{2-[2-(6-éthanesulfonyl-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1321) 1-(4-{(E)-2-[2-(9,9-difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl}-3,5-diméthyl-phényl)-1-méthyl-urée ;

(1322) 1-(4-{2-[2-(9-hydroxy-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1323) 1-(4-{2-[2-(9,9-difluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1324) 1-(4-{2-[2-(9-amino-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1325) 1-(4-{(E)-2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1326) 1-(4-{(Z)-2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1327) 1-(4-{2-[2-(9-fluoro-nonyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1328) 8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-[3-(4,4,5,5,5-pentafluoropentyloxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1329) 8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-fluoro-3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1330) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-((R)-2,3-dihydroxy-propoxy)-2,6-diméthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1331) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1332) 8-{2-[4-(3,4-dihydroxy-butoxy)-2-méthyl-phényl]-éthanesulfonyl}-2-(4-fluoro-3-trifluorométhoxy-phényl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1333) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1334) 2-(4-fluoro-3-trifluorométhoxy-phényl)-8-{2-[4-(4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1335) 2-(4-fluoro-3-trifluorométhoxy-phényl)-8-{2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1336) 2-(2,2-difluoro-benzo[1,3]dioxol-5-yl)-8-{2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1337) 2-(4-fluoro-3-trifluorométhoxy-phényl)-8-{2-[4-(3-hydroxy-azetidine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1338) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[3-(4,4,4-trifluoro-butoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(1339) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butoxy)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(1340) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(5-phényl-pentyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl)-1-méthyl-urée;

(1341) 1-(3,5-diméthyl-4-{2-[2-(3-nonafluorobutyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-

éthyl)-phényl)-1-méthyl-urée;

(1342) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(4-pentyl-phényl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;

(1343) 1-(3,5-diméthyl-4-{2-[4-oxo-2-((1S,3R)-3-propyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;

(1344) 1-(4-{2-[2-(6-éthoxy-hexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1345) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[3-(6,6,6-trifluoro-hexyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(1346) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(6,6,6-trifluoro-hexyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(1347) ester éthyle de l'acide 12-(8-{2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthanésulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-2-yl)-dodécanoïque;

(1348) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(9-phényl-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;

(1349) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(6-phényl-hexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;

(1350) acide 14-(8-{2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthanésulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-2-yl)-tétradécanoïque;

(1351) 1-(4-{2-[2-(7-méthoxy-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1352) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(5-propoxy-pentyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;

(1353) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(3-propyl-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;

(1354) 1-(4-{2-[2-(3-méthoxy-4-pentyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1355) 1-(3,5-diméthyl-4-{2-[4-oxo-2-(2-propyl-benzofuran-6-yl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;

(1356) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[5-(propane-1-sulfonyl)-pentyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(1357) 1-(4-{2-[2-(7-méthanésulfonyl-heptyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1358) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(4,4,4-trifluoro-butyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(1359) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[3-(4,4,4-trifluoro-butyl)-phényl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-phényl]-1-méthyl-urée;

(1360) ester tert-butyle de l'acide [11-(8-{2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthanésulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-2-yl)-undecyl]-carbamique;

(1361) 1-(4-{2-[2-(11-hydroxy-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1362) acide 10-(8-{2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthanésulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-2-yl)-décanoïque;

(1363) amide de l'acide 10-(8-{2-[2,6-diméthyl-4-(1-méthyl-ureido)-phényl]-éthanésulfonyl}-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-2-yl)-décanoïque;

(1364) 1-(3,5-diméthyl-4-{2-[2-(4-[1,1,1-²H₃]-méthyl-[4-²H₁]cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-1-méthyl-urée;

(1365) 1-(3,5-diméthyl-4-{(E)-2-[4-oxo-2-(9-phényl-nonyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-vinyl}-phényl)-1-méthyl-urée;

(1366) 1-(4-{2-[2-(11-amino-undecyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1367) 3-[(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-méthyl-amino]-4-éthoxy-cyclobut-3-ène-1,2-dione;

(1368) 3-amino-4-[(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-méthyl-amino]-cyclobut-3-ène-1,2-dione;

(1369) 3-diméthylamino-4-[(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-méthyl-amino]-cyclobut-3-ène-1,2-dione;

(1370) N-(4-{2-[2-(4-éthynyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3-méthyl-

phényl)-acétamide;

(1371) 1-(4-{2-[4-[(Z)-hydroxyimino]-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-méthyl-urée;

(1372) 2-méthyl-benzylamide de l'acide 2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonique;

(1373) (2-o-tolyl-éthyl)-amide de l'acide 2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonique;

(1374) 2-cyclohexyl-8-{2-[1-((S)-2,3-dihydroxy-propyl)-1H-indol-4-yl]-2-hydroxy-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1375) 2-cyclohexyl-8-(2-oxo-2-o-tolyl-éthanesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1376) 2-cyclohexyl-8-(2-o-tolyl-éthynesulfonyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1377) 2-cyclohexyl-8-[2-(1H-indol-4-yl)-éthynesulfonyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1378) 3,5,N,N-tetraméthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-benzamide;

(1379) 3,5,N,N-tetraméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(1380) 8-{2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1381) 8-{2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1382) 8-{2-[4-(4-hydroxy-4-méthyl-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1383) 8-{2-[4-(4-fluorométhyl-4-hydroxy-pipéridine-1-carbonyl)-2,6-diméthyl-phényl]-éthanesulfonyl}-2-(4-méthyl-cyclohexyl)-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1384) 8-(2-{4-[4-((R)-2,3-dihydroxy-propoxy)-pipéridine-1-carbonyl]-2,6-diméthyl-phényl}-éthanesulfonyl)-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1385) hydrazide de l'acide 3,5-diméthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-benzoïque;

(1386) 8-{2-[2,6-diméthyl-4-(pyrazolidine-1-carbonyl)-phényl]-éthanesulfonyl}-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1387) N,N-diméthyl-2-(3-méthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;

(1388) N-méthoxy-2-(3-méthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro 4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;

(1389) 1-[3,5-diméthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro [4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzyl]-1-méthyl-urée;

(1390) 1-(3,5-diméthyl-4-{2-[2-(4-méthyl-cyclohexyl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzyl)-1-méthyl-urée;

(1391) 1-(4-{2-[2-(4-fluoro-3-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,5-diméthyl-phényl)-1-[1,1,1-²H₃]méthyl-urée;

(1392) 2-cyclohexyl-8-{2-[1-((2S,3S)-2,3,4-trihydroxy-butyl)-1H-indol-4-yl]-éthanesulfonyl}-1,3,8-triaza-spiro[4.5]déc-1-én-4-one ;

(1393) 8-{2-[1-((2S,3S)-4-benzyloxy-2,3-dihydroxy-butyl)-1H-indol-4-yl]-éthanesulfonyl}-2-cyclohexyl-1,3,8-triaza-spiro[4.5]déc-1-én-4-one;

(1394) N-[3-méthyl-4-(2-{4-oxo-2-[4-(3,3,3-trifluoro-propyl)-cyclohexyl]-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl)-benzyl]-acétamide;

(1395) 3,N,N-triméthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phénylamino)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(1396) 3,N,N-triméthyl-4-{2-[4-oxo-2-(4-trifluorométhyl-phénylamino)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(1397) 3,N,N-triméthyl-4-{2-[2-(méthyl-phényl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-benzamide;

(1398) 4-{2-[2-(benzyl-méthyl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N',triméthyl-benzamide;


(1399) 4-{2-[2-(cyclohexylméthyl-amino)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;

(1400) 3,N,N-triméthyl-4-(2-{2-[méthyl-(4-trifluorométhyl-phényl)-amino]-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl}-éthyl)-benzamide;

- (1401) N-(3-méthyl-4-{2-[4-oxo-2-(4-trifluorométhyl-phénylamino)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;
 (1402) N-(3-méthyl-4-{2-[4-oxo-2-(3-trifluorométhyl-phénylamino)-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-phényl)-acétamide;
 (1403) 4-{2-[2-(4-butyl-pipéridin-1-yl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;
 (1404) 4-{2-[2-(4-butyl-cyclohexylamino)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N,N-triméthyl-benzamide;
 (1405) 4-{2-[2-(3-but-3-enyloxy-5-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N-diméthyl-N-pent-4-enyl-benzamide;
 (1419) N-(2-allyloxy-éthyl)-4-{2-[2-(3-but-3-enyloxy-5-trifluorométhyl-phényl)-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl]-éthyl}-3,N-diméthyl-benzamide;
 (1443) {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl}-méthyl-amide de l'acide déc-9-énoïque; ou
 (1446) {4-[2-(2-hex-5-enyl-4-oxo-1,3,8-triaza-spiro[4.5]déc-1-ène-8-sulfonyl)-éthyl]-3,5-diméthyl-phényl}-méthyl-amide de l'acide hept-6-énoïque,

ou un sel pharmacologiquement acceptable de ceux-ci.


11. Composé ou sel pharmacologiquement acceptable de celui-ci selon l'une quelconque des revendications 1 à 5, dans lequel U représente un groupe alkylène en C₁ à C₆ ou tout groupe choisi parmi les groupes représentés par la formule suivante :



A représente O ;

R₄₆ est choisi parmi un atome d'hydrogène, les groupes alkyle en C₁ à C₁₀, halogénoalkyle en C₁ à C₁₀ et hydroxyalkyle en C₁ à C₁₀ ;

V est choisi parmi :



E représente un groupe pyrrolidine ou pipéridine éventuellement substitué par un groupe hydroxyle ; et R₇ est choisi parmi :

- 1) un atome d'hydrogène,
- 2) un groupe alkyle en C₁ à C₁₀ et
- 3) un groupe hydroxyalkyle en C₁ à C₁₀.

12. Composition pharmaceutique comprenant le composé ou un sel pharmacologiquement acceptable de celui-ci selon l'une quelconque des revendications 1 à 11 en tant que principe actif.

13. Agent prophylactique ou thérapeutique pour une utilisation dans le traitement ou la prévention de l'ostéoporose, d'une fracture, de l'ostéomalacie, de l'arthrite, d'une thrombocytopénie, d'une hypoparathyroïdie, d'une hyperphosphatémie ou d'une calcinose tumorale, comprenant le composé ou un sel pharmacologiquement acceptable de celui-ci selon l'une quelconque des revendications 1 à 11 en tant que principe actif.

5

14. Utilisation du composé ou d'un sel pharmacologiquement acceptable de celui-ci selon l'une quelconque des revendications 1 à 11 pour la fabrication d'un agent prophylactique ou thérapeutique pour une utilisation dans le traitement ou la prévention de l'ostéoporose, d'une fracture, de l'ostéomalacie, de l'arthrite, d'une thrombocytopénie, d'une hypoparathyroïdie, d'une hyperphosphatémie ou d'une calcinose tumorale.

10

15. Composé ou sel pharmacologiquement acceptable de celui-ci selon l'une quelconque des revendications 1 à 11 pour une utilisation dans le traitement ou la prévention de l'ostéoporose, d'une fracture, de l'ostéomalacie, de l'arthrite, d'une thrombocytopénie, d'une hypoparathyroïdie, d'une hyperphosphatémie ou d'une calcinose tumorale.

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- DE 4234330 [2273]

Non-patent literature cited in the description

- **KARAPLIS, A. C. et al.** *Genes and Dev.*, 1994, vol. 8, 277-289 [0014]
- **LANSKE, B. et al.** *Science*, 1996, vol. 273, 663-666 [0014]
- Parathyroid hormone-related protein: Structure, processing and physiological actions. **BROADUS, A. E. ; STEWART, A. F. et al.** Basic and Clinical Concepts. Raven Press, 1994, 259-294 [0014]
- Parathyroid hormone: Biosynthesis, secretion, chemistry and action. **KRONENBERG, H. M. et al.** Handbook of Experimental Pharmacology. Springer-Verlag, 1993, 185-201 [0014]
- **URENA, P. et al.** *Endocrinology*, 1994, vol. 134, 451-456 [0014]
- **ISHIHARA, T. et al.** *EMBO J.*, 1991, vol. 10, 1635-1641 [0014]
- **LIN, H. Y. et al.** *Science*, 1991, vol. 254, 1022-1024 [0014]
- **JELINEK, L. et al.** *Science*, 1993, vol. 259, 1614-1616 [0014]
- **KOLAKOWSKI, L. F.** *Receptors and Channels*, 1994, vol. 2, 1-7 [0014]
- **USDIN, T. et al.** *J. Biol. Chem.*, 1995, vol. 270, 15455-15458 [0014]
- **BERGWITZ, C. et al.** *J. Biol. Chem.*, 1996, vol. 271, 26469-26472 [0014]
- **ABOU-SAMRA, A. B et al.** *Pro. Natl. Acad. Sci. USA*, 1992, vol. 89, 2732-2736 [0014]
- **BRINGHURST F. R. et al.** *Endocrinology*, 1993, vol. 132, 2090-2098 [0014]
- **NEER RM et al.** *N. Eng. J. Med*, 2003, vol. 344, 1434-1441 [0014]
- **GREENSPAN SL et al.** *Ann of Intern Med*, 2007, vol. 146, 326-339 [0014]
- **TASHJIAN ; GAGEL.** *J. Bone Miner. Res*, 2006, vol. 21, 354-365 [0014]
- **HOARE SRJ. et al.** *Drug Discov. Today*, 2005, vol. 10, 417-427 [0014]
- *Chemistry-A European Journal*, 2002, vol. 8 (2), 439-450 [0219]
- **MARCH.** Advanced Organic Chemistry. John Wiley and Sons, 523, , 524 [0222]
- *J. Org. Chem.*, 1985, vol. 50 (12), 2066-2073 [0225]
- *J. Org. Chem.*, 1984, vol. 49 (26), 5124-5131 [0225]
- *Tetrahedron Lett.*, 1994, vol. 35, 1837-1840 [0226]
- *Chemistry Lett.*, 1992, 1483-1486 [0226]
- *Tetrahedron*, 2001, vol. 57 (37), 7899-7907 [0235] [0236]
- *Synlett*, 2005, vol. 5, 834-838 [0236]
- **MARCH.** Advanced Organic Chemistry [0239] [0249]
- **MARCH.** Advanced Organic Chemistry. John Wiley and Sons [0277] [0278]
- *J. Med. Chem.*, 2005, vol. 48, 6066-6083 [0277] [0278]
- *Organic Syntheses*, 1951, vol. 31, 8-11 [0277] [0278]
- *Bioorg. Med. Chem. Lett.*, 2003, vol. 13, 837-840 [0277] [0278]
- *Chem. Rev.*, 2002, vol. 102, 1359 [0277] [0278]
- *J. Organomet. Chem.*, 1999, vol. 576, 147 [0277] [0278]
- *Chem. Rev.*, 1995, vol. 95, 2457 [0277] [0278]
- Protective Groups in Organic Synthesis. Wiley-Interscience [0277] [0278]
- *Angewandte Chemie, International Edition*, 2004, vol. 43 (41), 5542-5546 [2224]
- *Journal of Organic Chemistry*, 2006, vol. 71 (21), 8183-8189 [2259]
- **TAKASU et al.** *J. Bone. Miner. Res.*, 1999, vol. 14, 11-20 [2388]