



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
16.01.2013 Bulletin 2013/03

(51) Int Cl.:
A61K 31/35 (2006.01) **A01N 43/16** (2006.01)
A61K 31/352 (2006.01) **A61K 31/517** (2006.01)

(21) Application number: **12171867.0**

(22) Date of filing: **28.02.2007**

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR
Designated Extension States:
AL BA HR MK RS

(72) Inventor: **Sherris, David, I.**
Jamaica Plain, MA Massachusetts 02130 (US)

(74) Representative: **Duffy, Assumpta Dympha FRKelly**
27 Clyde Road
Ballsbridge
Dublin 4 (IE)

(30) Priority: **28.02.2006 US 777318 P**
27.04.2006 US 412618

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
07757634.6 / 1 996 021

(71) Applicant: **Paloma Pharmaceuticals, Inc.**
Jamaica Plain, MA 02130 (US)

Remarks:

- This application was filed on 13-06-2012 as a divisional application to the application mentioned under INID code 62.
- Claims filed after the date of receipt of the divisional application (Rule 68(4) EPC).

(54) **Compositions and methods to treat diseases characterized by cellular proliferation and angiogenesis**

(57) Described herein are compositions and methods for preventing and/or treating diseases involving ab-

errant angiogenesis employing one or more benzo[c]chromen-6-one derivatives.




FIG. 2

Description

FIELD OF THE INVENTION

[0001] The present invention relates to compositions and methods for preventing and/or treating diseases associated with cellular proliferation and/or angiogenesis. The current invention is directed in part to a series of chemical compositions that demonstrate therapeutic benefit in diseases involving abnormal cellular proliferation, abnormal angiogenesis or a combination thereof.

BACKGROUND OF THE INVENTION

[0002] Blood vessels that make up the cardiovascular system may be broadly divided into arteries, veins and capillaries. Arteries carry blood away from the heart at relatively high pressure; veins carry blood back to the heart at low pressure, while capillaries provide the link between the arterial and venous blood supply. During embryonic development, vessels are first formed through vasculogenesis, utilizing pluripotent endothelial cell precursors. Later, through arteriogenesis, larger blood vessels are formed possessing a more complex structure of endothelial cells, smooth muscle cells and pericytes (tunica media). Although arteriogenesis is not considered to occur in the adult, blood vessels may be formed in the adult through vasculogenesis and notably a process known as angiogenesis. Under normal conditions, angiogenic neovascularization occurs during such conditions as wound repair, ischemic restoration and the female reproductive cycle (generating endometrium forming the corpus luteum and during pregnancy to create the placenta). The capillaries, relatively simple vessels formed by angiogenesis, lack a developed tunica as they are predominantly composed of endothelial cells and to a lesser extent perivascular cells and basement membrane. Cancer is a disease state characterized by the uncontrolled proliferation of altered tissue cells. Tumors less than a few millimeters in size utilize nearby normal vessels to provide nutrients and oxygen. However, above this critical size, cancer cells utilize angiogenesis to create additional vascular support. Normally, angiogenesis is kept in check by the body naturally creating angiogenic inhibitors to counteract angiogenic factors. However, the cancer cell changes this balance by producing angiogenic growth factors in excess of the angiogenic inhibitors, thus favoring blood vessel growth. Cancer initiated angiogenesis is not unlike angiogenesis observed during normal vessel growth. Angiogenic factors pass from the tumor cell to the normal endothelium, binding the endothelial cell, activating it and inducing endothelial signaling events leading to endothelial cell proliferation. Endothelial tubes begin to form, homing in toward the tumor with the formation of capillary loops. Capillaries then undergo a maturation process to stabilize loop structure.

[0003] There have been several chemotherapeutic approaches targeted against tumor cell proliferation including alkylating agents, antimetabolites, antimitotics, antimetabolites and antibiotics. These act preferentially on the rapidly proliferating tumor cells. Hormonal therapy with antiestrogens or anti-androgens is another approach to attacking cancer cells that work by inhibiting the proliferative action of the required hormone. Although anti-cancer agents fall into specific classifications, it is not uncommon for agents to act by multiple modes of action. For example, the anti-estrogen tamoxifen has been shown to have anti-proliferative activity on cancer cells and endothelial cells (anti-angiogenic) by an estrogen independent mechanism. Taxol, an antimitotic agent acting on microtubules has also demonstrated anti-angiogenic properties, possibly by inducing apoptosis of endothelial cells through Bcl-2 phosphorylation.

[0004] Cancer is but one disease associated with a pathological neovasculature. A wide variety of diseases involving aberrant angiogenesis exist in nature. These diseases utilize the same steps involved in normal capillary growth but do so in an aberrant manner creating capillaries which lack a high degree of stability and function. Agents capable of inhibiting angiogenesis would be expected to exert activity on a variety of pathological neovascular diseases. Angiogenesis may be considered a key component in the pathogenesis of a number of diseases. If through therapeutic intervention angiogenesis could be slowed down or eliminated, anti-angiogenic agents would then be expected to abolish or lessen a variety of neovasculature associated diseases. Anti-angiogenic therapy will likely be very effective at suppressing tumor growth by denying the tumors a blood supply. However, anti-angiogenic therapy may prove more effective in combination with other therapies aimed directly at the tumor cells. Chemical agents that demonstrate both anti-angiogenic and tumor directed properties would be advantageous in this regard.

[0005] Thus, there remains a need to develop agents that demonstrate anti-proliferative effects against human endothelial cells for the treatment of a variety of diseases, including, but not limited to, cancer, in addition to an inhibitory effect directly on cancer cells for the treatment of tumors, or other cells acting as an initiator of angiogenesis for diseases outside of cancer. And at least in some cases, the ability to have a sustained half-life through the creation of analogs with modifications that inhibit metabolism and hence clearance or loss of activity, and show little or no toxicity. The present invention seeks to meet these and other needs.

BRIEF SUMMARY OF THE INVENTION

[0006] The present invention relates to compositions and methods for treating diseases associated with cellular proliferation and/or angiogenesis. The current invention is directed in part to a series of chemical compositions that demonstrate a therapeutic benefit in diseases involving abnormal cellular proliferation, abnormal angiogenesis or a combination thereof.

[0007] One embodiment of the present invention is directed to compositions used to prevent and/or treat abnormal cellular proliferation. In one aspect, the invention is directed to a series of benzo[c]chromen-6-one derivatives that demonstrate enhanced anti-proliferative effects against human endothelial cells for the treatment of a variety of diseases, including, but not limited to, cancer. These agents have a dual anti-angiogenic, tumor cell anti-proliferative activity.

[0008] In another aspect of the present embodiment, the invention is directed to a series of benzo[c]chromen-6-one derivatives that demonstrate enhanced anti-proliferative effects against human endothelial cells for the treatment of a variety of diseases including, but not limited to, cancer with minimal or no anti-proliferative effects directly on tumor cells. In other words, these agents show predominantly anti-angiogenic activity.

[0009] Another embodiment of the present invention is directed to a series of benzo[c]chromen-6-one derivatives that demonstrate enhanced anti-proliferative effects against human endothelial cells for the treatment of a variety of diseases including, but not limited to, cancer. In one aspect, the enhanced anti-proliferative effect against human endothelial cells is complimented by an anti-proliferative inhibitory effect directly on tumor cells for the treatment of cancer. In another aspect, the enhanced anti-proliferative effect against endothelial cells is complimented with an anti-proliferative inhibitory effect on pathologically relevant cells specific to the disease, outside of cancer, for example keratinocytes for skin diseases.

[0010] In yet another embodiment, the present invention is directed toward methods of administering a therapeutically effective amount of one or more compositions described herein to a subject in need thereof. In one aspect, the targeted subject has been diagnosed with or is predisposed toward one or more diseases associated with abnormal cellular proliferation and/or angiogenesis, including for example, a cancer.

[0011] Other features and advantages of the invention will be apparent from the following detailed description of embodiments thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0012]

Figure 1A is a bar graph showing inhibition of bFGF-stimulated endothelial cell proliferation by Palomid 529;
 Figure 1B is a bar graph showing inhibition of VEGF-stimulated endothelial cell proliferation by Palomid 529;
 Figure 2 is a bar graph presenting data on the apoptotic inducing ability of compounds of the present invention;
 Figure 3 is a bar graph presenting data on the apoptotic inducing activity in keratinocytes;
 Figure 4 is a graph showing inhibition of keratinocyte growth;
 Figure 5 is a bar graph showing induction of apoptotic activity at varying concentrations of compound; and
 Figure 6 is a bar graph showing metabolic stability of compounds of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

[0013] The present invention relates to compositions and methods for preventing and/or treating diseases associated with unwanted cellular proliferation and/or angiogenesis. The current invention is directed in part to a series of chemical compositions that demonstrate therapeutic benefit in diseases involving abnormal cellular proliferation, abnormal angiogenesis or a combination thereof. In a particular aspect, the instant invention relates to benzo[c]chromen-6-one derivatives that demonstrate their effect on diseases characterized by abnormal proliferation, abnormal angiogenesis or a combination thereof.


[0014] The term "derivative" is understood by those skilled in the art. For example, a derivative can be understood as a chemical compound that is produced from another compound of similar structure in one or more steps, such as illustrated in Table I (infra) for benzo[c]chromen-6-one.

[0015] Disease therapeutic agents currently under development are based on a variety of targeting strategies. One strategy is the use of natural inhibitors of angiogenesis such as thrombospondin, angiostatin and endostatin. Another strategy is the use of agents that block receptors required to stimulate angiogenesis, such as antagonists of the VEGF receptor. A third strategy is the inhibition of enzymes which allow new blood vessels to invade surrounding tissues, for example, inhibitors of matrix metalloproteinases. Another strategy for inhibiting angiogenesis is through the use of integrin antagonists such as $\alpha v \beta 1$ antibodies or small molecule drugs through the inhibition of endothelial cell adhesion effecting capillary tube formation.

[0016] Angiogenesis is an attractive therapeutic target for cancer treatment due to its selectivity of action. Blood vessels in growing tumors are in a microenvironment conducive to cellular activation and rapid proliferation whereas blood vessels in most normal tissues are quiescent. This microenvironment inducing cellular activation and rapid proliferation are believed to be the physiological differences that allow the selective targeting of blood vessels in the tumor by anti-angiogenic agents.

[0017] The present invention relates to a therapeutic formulation comprising one or more compositions useful in the treatment of unwanted cellular proliferation and/or angiogenesis and/or keratinocyte proliferation. The present invention also relates to a therapeutic formulation comprising one or more compositions useful in the treatment of cancer as well as other diseases characterized by the undesired excessive, abnormal stimulation or proliferation of, for example endothelial cells or other cells resulting in such diseases including, but not limited to, ocular diseases of corneal, retinal or anterior chamber neovasculation, cancer (including, but not limited to, fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, hemangioma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinomas, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, osteosarcoma, meningioma, melanoma, neuroblastoma, retinoblastoma, acoustic neuroma, neurofibromas, trachoma and pyogenic granulomas, acute lymphocytic leukemia and acute myelocytic leukemia, chronic leukemia, polycythemia vera, lymphoma, multiple myeloma, Waldenstrom's macroglobulinemia, and heavy chain disease), hereditary hemorrhagic telangiectasia, solid or blood born tumors, acquired immune deficiency syndrome, post-menopausal symptoms, osteoporosis, cardiovascular disease, Alzheimer's disease, to reduce the incidence of strokes, as an alternative to prior estrogen replacement therapies, vascular malformations, abnormal wound healing, inflammatory and immune disorders, gout, and other ocular diseases (retinal/choroidal, corneal, neoplastic and anterior chamber neovascular, including but not limited to macular degeneration, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, multifocal choroiditis, Best's disease, Stargardt's disease, cblC type of cobalamin deficiency, hyperviscosity syndrome, Sorsby's fundus dystrophy, pseudoxanthoma elasticum, rubeosis iridis, Osler Weber syndrome (Osler-Weber-Rendu disease), keratoconjunctivitis, Vitamin A deficiency, phlyctenulosis, contact lens over wear, infection (including but not limited to bacterial, viral, parasitic, or fungal), atopic and superior limbic dermatitis, chronic uveitis, chronic vitritis, Eales' disease, radial keratotomy, uncharacteristic proliferation of fibrovascular or fibrous tissue including all forms of proliferative vitreoretinopathy whether or not associated with diabetes, chronic retinal detachment, trauma (including but not limited to abrasion, previous surgery with complications such as corneal allograft rejection, alkaline burns, acid burns or hydrocarbon burns, mechanical or thermal damage to Bruch's membrane), pterygium, neoplasia (including but not limited to retinoblastoma and melanoma)), arthritis (rheumatoid and osteoarthritis) and skin diseases (psoriasis, atopic dermatitis, photodamage). Other diseases associated with angiogenesis include Sjögren's syndrome, systemic lupus, polyarteritis, pemphigoid, sickle cell anemia, Paget's disease, vein or artery occlusion, carotid obstructive disease, Lyme disease, Behcet's disease, bartonellosis, arteriosclerosis, induction of amenorrhea to block ovulation or to prevent implantation by the blastula, surgical adhesions and chronic inflammation (including but not limited to ulcerative colitis and Crohn's disease).

[0018] In accordance with the present invention, there is provided pharmaceutical compositions comprising Formula I:



where,

R1 = H or alkyl;

R2 = H, OH, O-alkyl, amino, O-heterocyc, O-aryl, O-substituted alkyl, where substitution is e.g. halo, aryl, or heteroaryl, O-Ac, O-PO3, O-SO3, or OSO2NH2;

R3 = H, OH, O-alkyl, O-CH2Aryl, O-CH2heteroaryl, O-alkylaryl, O-acyl, or nitro;

R4 = H, Alkyl, CH2Aryl, substituted alkyl, OH, O-alkyl, C-aryl, OCH2Aryl, OCH2Heteroaryl, O-Acyl, OPO3, OSO3, or OSO2NH2;

R5 = H, Oxo, aryl, hydroxyl, alkyl, or O-alkyl;


R6=H;

R7 = H, Acyl, substituted alkyl, where substitution is e.g. hydroxyl or sulfamoyl, alkyl, O- -alkyl; or O-substituted alkyl where substitution is O-PO3 or OSO3;

R8 = H; and

X = O, N, or S.

[0019] In accordance with the present invention, there is provided pharmaceutical compositions comprising Formula II:



where,

R1 = H or alkyl;

R2=H, O-alkyl, OH, amino, O-heterocyc, O-aryl, O-substituted alkyl where substitution is e.g. halo, aryl, or heteroaryl, O-Ac, O-PO3, O-SO3, or OSO2NH2;


R3 = H, O-alkyl, O-substituted alkyl where substitution is aryl or heteroaryl, OH, O-acyl, or nitro;

R4 = H, Alkyl, CH2Aryl, substituted alkyl, O), O-alkyl, O-aryl, OCH2Aryl, OCH2Heteroaryl, O-Acyl, OPO3, OSO3, or OSO2NH2;

R5 = H, Aryl, heteroaryl or substituted alkyl; and

R6 = H, Alkyl, or Aryl.

[0020] In accordance with the present invention, there is provided pharmaceutical compositions comprising Formula III:



where,


R1 = alkyl or H;

R2 = alkyl or H;

R3 = Acetyl; and

R4 = H or Alkyl.

[0021] In accordance with the present invention, there is provided pharmaceutical compositions comprising Formula IV:



where,

R1 = H or F;

R2 = H or nitro;

























R3 = H;

































R4 = H; and


R5 = alkyl, substituted alkyl or aryl.


[0022] In accordance with the present invention, there is provided a pharmaceutical composition comprising a therapeutically effective amount of one or more benzo[c]chromen-6-one derivatives having the following structure depicted in Table I:

Table I: Structural formula of benzo[c]chromen-6-one derivatives

5	 SG00526	 SG00527	 SG00528	 SG00529
10	 SG00530	 SG00531	 SG00532	 SG00533
15	 SG00535	 SG00536	 SG00537	 SG00538
20	 SG00539	 SG00540	 SG00272	 SG00541
25	 SG00542	 SG00543	 SG00544	 SG00545
30	 SG00546	 SG00547	 SG00548	 SG00549
35				
40				
45				
50				
55				

5	 SG00550	 SG00551	 SG00552	 SG00553
10	 SG00554	 SG00555	 SG00556	 SG00557
15	 SG00558	 SG00559	 SG00560	 SG00561
20	 SG00562	 SG00563	 SG00564	 SG00565
25	 SG00566	 SG00567	 SG00568	 SG00569
30	 SG00570	 SG00571	 SG00572	 SG00573
35	 SG00574	 SG00575	 SG00576	 SG00577
40	 SG00578	 SG00579	 SG00580	 SG00581
45	 SG00582	 SG00583	 SG00584	 SG00585
50	 SG00586	 SG00587	 SG00588	 SG00589
55	 SG00590	 SG00591	 SG00592	 SG00593





[0023] The individual benzo[c]-chromen-6-one derivatives of Table I are identified by the designation "SG" followed by a number. They are alternatively referred to herein by the designation "Palomid" followed by the same number, i.e. the terms "SG" and "Palomid" are used interchangeably throughout this application.

[0024] The compounds of Table I exhibit anti-angiogenic and/or anti-keratinocyte activities. Those skilled in the art will appreciate that the invention includes other benzo[c]-chromen-6-one derivatives having anti-angiogenic, and/or anti-tumor activities. These characteristics can be determined for each derivative using the assays detailed below and elsewhere in the literature. The process or processes by which benzo[c]chromen-6-one derivatives affect cell growth remains to be fully researched, however, benzo[c]chromen-6-one derivatives may induce changes in the levels and activities of various proteins involved in the progression of the cell cycle. These include cofactors of DNA replication and repair, e.g., proliferating cell nuclear antigen and cell division cycle kinases (and regulators). Benzo[c]chromen-6-one may also up-regulate Death Receptor 5 and caspase 8, inhibit HIF-1 α (a global transcriptional regulator of angiogenesis genes), or inhibit the Akt/mTor signal transduction pathway (a key regulator pathway of cell growth and proliferation with its deregulation associated with human diseases, including but not limited to cancer).

[0025] Assays relevant to these mechanisms of action and inhibition of cell proliferation are well-known in the art. For example, anti-mitotic activity mediated by effects on tubulin polymerization activity can be evaluated by testing the ability of a benzo[c]chromen-6-one derivative to inhibit tubulin polymerization and microtubule assembly in vitro. Other such assays include counting of cells in tissue culture plates or assessment of cell number through metabolic assays or incorporation into DNA of labeled (radio-chemically, e.g., 3H-thymidine or fluorescently labeled) or immuno-reactive (BrdU) nucleotides. In addition, measuring HIF- α activity for example through luciferase reporter groups or Akt/mTor signalling through for example activated intermediates as in the phosphorylation of Akt. Furthermore, anti-angiogenic activity may be evaluated through endothelial cell migration, endothelial cell tubule formation or vessel outgrowth in ex-vivo models of rat aortic rings.

[0026] The present invention also relates to implants or other devices comprised of one or more compositions described herein or prodrugs thereof wherein the composition or prodrug is formulated in a biodegradable or non-biodegradable format for sustained release. Non-biodegradable formats release the drug in a controlled manner through physical or mechanical processes without the format being itself degraded. Bio-degradable formats are designed to gradually be hydrolyzed or solubilized by natural processes in the body, allowing gradual release of the admixed drug or prodrug. Both bio-degradable and non-biodegradable formats and the process by which drugs are incorporated into the formats for controlled release are well known to those skilled in the art. These implants or devices can be implanted in the vicinity where delivery is desired, for example, at the site of a aberrant skin or in the vicinity of aberrant vasculature.

[0027] The present invention also relates to coated vascular stents to prevent restenosis, a renarrowing or blockage of an artery at the same site where treatment, such as angioplasty or stent procedure, has already been done. According to the present invention, the stent or other surgically implantable device is coated with one or more compositions described herein. The coating of such a device is well known to those skilled in the art.

[0028] The present invention also relates to conjugated prodrugs and uses thereof. More particularly, the invention relates to conjugates of benzo[c]chromen-6-one derivatives and the use of such conjugates in the prophylaxis or treatment

of conditions associated with uncharacteristic cell proliferation and/or uncharacteristic angiogenesis. Such diseases include, but are not limited to, excessive, abnormal stimulation or proliferation of cancer cells, endothelial cells or other pathologically involved cells.

[0029] The present invention also provides a conjugated prodrug of a benzo[c]chromen-6-one derivative conjugated to a biological activity modifying agent, e.g., a peptide, an antibody or fragment thereof, or in vivo hydrolysable esters, such as methyl esters, phosphate or sulfate groups, and amides or carbamates. Modifications can include modifying a hydroxyl group with a phosphate group. This derivative would not be expected to have activity due to the modification causing a significant change to the derivative thereby losing biological activity. However the modification imparts better solubility characteristics, i.e., more water soluble, which could facilitate transport through the blood or give it better oral availability to allow it to reach its site of activity. Once it gets into the microenvironment where it is needed to have activity, the modification is cleaved through natural processes, i.e., endogenous enzymes which are present at the site of needed activity. Alternatively it may just keep the derivative in a state to give it better systemic concentration which would then be cleaved again in the systemic circulation and thereby enhance its activity. The incorporation of benzo[c]chromen-6-one derivatives into a disease-dependently activated prodrug enables significant improvement of potency and selectivity in treating one or more disease conditions referred to hereinabove.

[0030] In addition to the compounds of the present invention, the pharmaceutical composition of this invention may also contain or be co-administered (simultaneously or sequentially) with one or more pharmacological agents of value in treating one or more disease conditions referred to hereinabove. Such agents include, but are not limited to, pharmaceutical agents well known to those skilled in the art for their oncolytic or anti-cancer activity. Other agents include those that suppress the side-effects of oncolytic or anti-cancer agents such as those directed toward counter-acting nausea and emesis.

[0031] Furthermore, the benzo[c]chromen-6-one derivatives or prodrugs thereof may be incorporated into bio-degradable or non-degradable formats allowing for sustained release. For example, the formulation being implanted in the proximity of where the delivery is desired, at the site of a tumor or in the vicinity of aberrant vasculature. Alternatively, the pharmaceutical formulation can be packaged into a delivery vehicle that has a chemical moiety that provides for specificity. For example, the moiety can be an antibody or some other such molecule that directs and facilitates delivery of the active agent to the desirable site (or cell/tumor).

[0032] The present invention also relates to use of the benzo[c]chromen-6-one derivatives or prodrugs thereof for the preparation of a medicant for the prophylaxis or treatment of conditions associated with any disease characterized by uncharacteristic cell proliferation and/or uncharacteristic angiogenesis and/or inflammation.

[0033] The present invention also relates to the provision of a pharmaceutical composition comprising benzo[c]chromen-6-one derivatives or prodrugs thereof according to the present invention together with a pharmaceutical acceptable carrier, diluent or excipient.

The pharmaceutical composition may also be used for the prophylaxis or treatment of conditions associated with any disease characterized by uncharacteristic cell proliferation and/or uncharacteristic angiogenesis and/or inflammation.

[0034] The present invention also pertains to methods of prophylaxis or treatment of a condition associated with any disease characterized by uncharacteristic cell proliferation and/or uncharacteristic angiogenesis and/or inflammation, said method including administering to a subject in need of such prophylaxis or treatment an effective amount of benzo[c]chromen-6-one derivatives or prodrugs thereof according to the present invention as described hereinabove. (It should be understood that prophylaxis or treatment of said condition includes amelioration of said condition.)

[0035] By "effective amount" it is meant a therapeutically effective amount that relieves symptoms, partially or completely, associated with a particular disease or syndrome. Such amounts can be readily determined by an appropriately skilled practitioner, taking into account the condition to be treated, the route of administration, and other relevant factors - well known to those skilled in the art. Such a person will be readily able to determine a suitable dose, mode and frequency of administration.

[0036] Pharmaceutically acceptable salts of the benzo[c]chromen-6-one derivatives or prodrugs thereof may be prepared in any conventional manner. In vivo hydrolysable esters, for example, methyl esters, phosphate or sulfate groups, and amides or carbamates may be prepared in any conventional manner.

[0037] The benzo[c]chromen-6-one derivatives or prodrugs thereof can be provided as physiologically acceptable formulations using known techniques and these formulations can be administered by standard routes. The compositions may be administered through means including, but not limited to, topical, oral, rectal or parenteral, for example, intravenous, subcutaneous or intramuscular, route. In addition, the compositions may be incorporated into format allowing for sustained release, the formats being implanted in the proximity of where the delivery is desired, for example, at the site of the skin disease or aging skin or in the vicinity of aberrant vasculature. The dosage of the composition will depend on the condition being treated, the particular derivative used, and other clinical factors such as weight and condition of the subject and the route of administration of the compound - all of which is appreciated by those skilled in the art. For example, a person skilled in the art will be able by reference to standard texts, such as Remington's Pharmaceuticals Sciences 17th edition (the entire teaching of which is incorporated herein by reference), determine how the formulations

are to be made and how these may be administered.

[0038] The formulations including, but not limited to, those suitable for oral, rectal, nasal, inhalation, topical (including, but not limited to, dermal, transdermal, buccal and sublingual), vaginal or parenteral (including, but not limited to, subcutaneous, intramuscular, intravenous, intradermal, intraocular (including, but not limited to, intra-vitreous, sub-conjunctival, sub-Tenon's, trans-scleral), intra-tracheal and epidural) and inhalation administration. The formulations may be conveniently presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association the active ingredient and a pharmaceutical carrier(s) or excipient(s). The formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

[0039] Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil emulsion, etc.

[0040] A tablet may be made by compression or molding, optimally with one or more accessory ingredient. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface-active or dispersing agent. Molded tablets may be made by molding, in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide a slow or controlled release of the active ingredient therein.

[0041] Formulations suitable for administration via the mouth include lozenges comprising the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the ingredient to be administered in a suitable liquid carrier.

[0042] Formulations suitable for topical administration to the skin may be presented as ointments, creams, gels and pastes comprising the ingredient to be administered in a pharmaceutical, cosmeceutical or cosmetic acceptable carrier. A viable delivery system is a transdermal patch containing the ingredient to be administered.

[0043] Formulations for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or a salicylate.

[0044] Formulations suitable for nasal administration, wherein the carrier is a solid, include a coarse powder having a particle size, for example, in the range of 20 to 500 microns which is administered in the manner in which snuff is taken, for example, by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable formulations include wherein the carrier is a liquid for administration, as for example a nasal spray or as nasal drop, including aqueous or oily solutions of the active ingredient.

[0045] Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing, in addition to the active ingredient, ingredients such as carriers as are known in the art to be appropriate.

[0046] Formulation suitable for inhalation may be presented as mists, dusts, powders or spray formulations containing, in addition to the active ingredient, ingredients such as carriers as are known in the art to be appropriate.

[0047] Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostatic agents and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried, lyophilized, conditions requiring only the addition of the sterile liquid, for example, water for injections, immediately prior to use. Extemporaneous injection solution and suspensions may be prepared from sterile powders, granules and tablets of the kinds previously described.

[0048] Acceptable unit dosage formulations are those containing a daily dose or unit, daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the administered ingredient. In addition to the ingredients mentioned above, the formulations of the present invention may include other agents conventional in the art having regard to the type of formulation in question, for example, those suitable for oral administration may include flavoring agents. The present invention includes compositions of about 100% to about 90% pure isomers. In another aspect, the invention pertains to compositions of about 90% to about 80% pure isomer. In yet another aspect, the invention pertains to compositions of about 80% to about 70% pure isomer. In still another aspect, the invention pertains to a composition of about 70% to about 60% pure isomer. In yet a further aspect, the invention pertains to a composition of about 60% to about 50% pure isomer. However, a stereochemical isomer labeled as alpha or beta may be a mixture of both in any ratio, where it is chemically possible by one skilled in the art. Additionally, included by this invention are both classical and non-classical bio-isosteric atom and substituent replacements and are well known by one skilled in the art. Such bio-isosteric replacements include, for example, substitution of =S or =NH for =O.

[0049] Known compounds that are used in accordance with the invention and precursors to novel compounds according

to the invention can be purchased from commercial sources, for example, Sigma-Aldrich. Other compounds according to the invention can be synthesized according to known methods well known to those skilled in the art.

The synthetic route for benzo[c]chromen-6-one derivatives SG00292 and SG00392 are summarized in Scheme 1, *infra*. This synthetic route presents one potential way to prepare this series of derivatives, and other synthetic routes (including modifying the order of synthetic steps or reagents) are possible to someone skilled in the art. In specific cases, the nature of protecting groups or the order of reactions may have to be altered in order to reach the desired products. These changes to the general synthetic schemes are well understood to one skilled in the art.


EXAMPLES

[0050] Benzo[c]chromen-6-one derivatives according to the present invention may be prepared using the following reaction schemes, Scheme 1 and synthetic methods Scheme 2.

The present invention also includes benzo[c]chromen-6-one derivatives prepared from the starting point of Scheme 1. The synthesis of these analogs are described in the synthetic methods shown in Scheme 3 and represents examples from the benzo[c]chromen-6-one derivatives as depicted in Table I.

Scheme 1

[0051]



Scheme 2

Synthesis 5-Benzyloxy-2-bromo-4-methoxybenzaldehyde (2)

[0052] 2-Bromo-5-hydroxy-4-methoxybenzaldehyde (25 g, 0.108 mol) and K_2CO_3 (30 g, 0.216 mol) were added to acetonitrile (250 mL) and flushed with Ar. Benzyl bromide (20 g, 0.12 mol) was added and the mixture was heated under Ar for 20 h at 50°C. After cooling, the mixture was poured into water (200 mL) and extracted with CH_2Cl_2 (300 mL). The CH_2Cl_2 was washed with water (3 x 100 mL), dried and concentrated. Recrystallization with isopropanol: water (3:1) gave 28.8 g (83%) of 2 as a light brown solid. 1H -NMR (400 MHz, $CDCl_3$) δ 3.96 (3H, s, OCH_3), 5.16 (2H, s, CH_2Ph), 7.07-7.48 (7H, m, ArH + CH_2Ph), 10.16 (1H, s, CHO).

5-Benzoyloxy-2-bromo-4-methoxyphenol (3)

[0053] 5-Benzoyloxy-2-bromo-4-methoxy-benzaldehyde **2** (5 g, 16.0 mmol) was added to CH₂Cl₂ (40 mL), flushed with Ar and cooled in an ice bath. A solution of mCPBA (5.2 g) in CH₂Cl₂ (50 mL) was added dropwise. Once the addition was complete the reaction mixture was refluxed under Ar for 14 h. After cooling the mixture was washed with sat. NaHCO₃ (3 x 50 mL), brine, dried and concentrated. The residue was recrystallized from ethyl acetate/hexanes to 4.1 g (85%) of **3** as large tan needles. ¹H-NMR (400 MHz, CDCl₃) δH 3.88 (3H, s, OCH₃), 5.10 (2H, s, CH₂Ph), 6.74 (1H, s, ArH), 7.08 (1H, s, ArH), 7.34 -7.40 (5H, m, CH₂Ph), 8.25 (1H, s, OH).

1-Benzoyloxy-4-bromo-2,5-dimethoxybenzene (4)

[0054] 5-Benzoyloxy-2-bromo-4-methoxy-phenol **3** (2.76 g, 89.0 mmol) and NaH (0.89 g, 13.0 mmol, 60% dispersion in oil) were added to a flask and flushed with Ar. Dry THF (50 mL) was added and the suspension was stirred in an ice bath for 20 min. CH₃I (1.7 mL, 27.0 mmol, filtered through basic alumina) was added and the mixture stirred at room temperature under Ar for 18 h. After cooling the reaction mixture in an ice bath, water was added slowly. The mixture was extracted with ethyl acetate, dried and concentrated to give yellow oil that solidified under vacuum. The oil was purified by silica gel chromatography using silica gel with (10% ethyl acetate/hexanes) to give 2.5 g (88%) of **4** as a white solid. ¹H-NMR (400 MHz, CDCl₃) δH 3.75 (3H, s, OCH₃), 3.84 (3H, s, OCH₃), 5.15 (2H, s, CH₂Ph), 6.57 (1H, s, ArH), 7.07 (1H, s, ArH), 7.32 -7.42 (5H, m, CH₂Ph).

4-Benzoyloxy-2,5-dimethoxyphenylboronic acid (5b)

[0055] 1-Benzoyloxy-4-bromo-2,5-dimethoxybenzene **4** (7.48g, 23.0mmol) was placed in a dry flask and flushed with Ar. Dry THF (75 mL) was added and the solution was cooled to -78°C in a dry ice/acetone bath. nBuLi (11 mL, 2.5M in hexanes) was added and the mixture was stirred for 20 min at -78°C. Triisopropyl borate (10.7 mL, 0.463 mol) was added and the reaction stirred for 2h at -78°C then allowed to come to room temperature at which time a white precipitate began to form. After stirring for an additional 20 h the reaction was quenched with saturated NH₄Cl (25 mL). After separating the organic layer the aqueous layer was extracted with ethyl acetate (2 x 50 mL). The organic layers were combined, dried and concentrated. The residue was triturated with hexanes and filtered to give 4.1 g (62%) of **5b** as a light off-white creamy solid. ¹H-NMR (400 MHz, DMSO-d₆) δH 3.70 (3H, s, OCH₃), 3.79 (3H, s, OCH₃), 5.16 (2H, s, CH₂Ph), 6.77 (1H, s, ArH), 7.18 (1H, s, ArH), 7.33 -7.52 (5H, m, CH₂Ph)

5-Acetyl-2-trifluoromethanesulfonyloxybenzoic acid methyl ester (6)

[0056] Methyl 5-acetylsalicylate (25.0 g, 0.129 mol) was dissolved in CH₂Cl₂ (250 mL) and pyridine (60 mL) under Ar at 0°C. Trifluoromethanesulfonic anhydride (37.9 g, 0.133 mol) was then added over 20 min. The reaction mixture was stirred for an additional 30 min and then quenched with water (500 mL). The organic layer was separated and washed three times with 5% HCl (80 mL). After removing the solvent the solid obtained was dried under vacuum to yield 40.3 g (96%) of **6**. ¹H-NMR (400 MHz, CDCl₃) δH 2.56 (3H, s, COCH₃), 3.89 (3H, s, OCH₃), 7.32 (1H, d, ArH), 8.12 (1H, d, ArH), 8.52 (1H, s, ArH),

4-Acetyl-4'-benzyloxy-2'-methoxybiphenyl-2-carboxylic acid methyl ester (7b)

[0057] 4-Benzoyloxy-2,5-dimethoxyphenylboronic acid **5b** (4.15 g, 14.4 mmol), 5-Acetyl-2-trifluoromethanesulfonyloxybenzoic acid methyl ester **6** (4.69g, 14.4mmol) and K₂CO₃ (3.98 g, 28.8 mmol) were added and the flask was flushed with Ar. Absolute ethanol (83 mL) and DME (94 mL) were added followed by Pd (PPh₃)₄ (0.87 g, 0.785 mmol) and the reaction mixture refluxed for 4 h. After cooling, water (100 mL), ethyl acetate (100 mL) and brine (50 mL) were added. The organic layer was washed with brine (2 x 50 mL) and the combined aqueous fraction was back extracted with ethyl acetate. The combined organic fraction was dried, concentrated and recrystallized from ethyl acetate/hexanes to give 6.5 g (99%) of **7b** as a yellow solid. ¹H-NMR (400 MHz, CDCl₃) δH 2.65 (3H, s, COCH₃), 3.58 (3H, s, OCH₃), 3.68 (3H, s, OCH₃), 3.88 (3H, s, OCH₃), 5.21 (2H, s, CH₂Ph), 6.55 (1H, s, ArH), 6.86 (1H, s, ArH), 7.30-7.48 (6H, m, ArH + CH₂Ph), 8.10 (1H, d, ArH), 8.37 (1H, s, ArH),

4-Acetyl-4'-benzyloxy-2'-methoxybiphenyl-2-carboxylic acid (8b)

[0058] To 4-Acetyl-4'-benzyloxy-2'-methoxybiphenyl-2-carboxylic acid methyl ester **7b** (4.06 g, 9.7 mmol) and NaOH (0.773 g, 19.4 mmol) was added methanol (60 mL) and water (60 mL). The reaction was refluxed under Ar for 7 h then cooled to room temperature. After placing in an ice bath, 1 M HCl was added to give a yellow precipitate that was filtered,

washed with water and recrystallized from THF/hexanes to give 2.7 g (69%) of **8b** as yellow crystals. ¹H-NMR (400 MHz, CDCl₃) δ 2.68 (3H, s, COCH₃), 3.62 (3H, s, OCH₃), 5.16 (2H, s, CH₂Ph), 6.77 (1H, s, ArH), 7.18 (1H, s, ArH), 7.33-7.52 (5H, m, CH₂Ph), 3.90 (3H, s, OCH₃), 5.22 (2H, s, CH₂Ph), 6.58 (1H, s, ArH), 6.90 (1H, s, ArH), 7.34-7.50 (6H, m, ArH + CH₂Ph), 8.17 (1H, d, ArH), 8.50 (1H, s, ArH),

8-Acetyl-3-benzyloxy-2-methoxybenzo[c]chromen-6-one (**9b**)

[0059] 4-Acetyl-4'-benzyloxy-2'-methoxybiphenyl-2-carboxylic acid **8b** (1.0 g, 2.5 mmol) was suspended in 1,2-dichloroethane (30 mL). SOCl₂ (200 mL, 2.7 mmol) was added and the reaction mixture refluxed for 2 h under Ar. After cooling to room temperature (some precipitate formed) AlCl₃ (0.262 g, 0.002 mol) was added turning the mixture red. The reaction was stirred at room temperature for 17 h then quenched with water (30 mL) and diluted with CH₂Cl₂ (100 mL). After washing the organic layer with brine (2 x 50 mL) it was dried and concentrated. The residue was dissolved in hot CHCl₃ and then cooled. Hexanes were added to help precipitate the product. A second recrystallization gave 0.3 g (32%) of **9b** as a yellow solid. ¹H-NMR (400 MHz, CDCl₃) δ 2.73 (3H, s, COCH₃), 4.05 (3H, s, OCH₃), 5.28 (2H, s, CH₂Ph), 6.94 (1H, s, ArH), 7.35-7.50 (6H, s, ArH + CH₂Ph), 8.07 (1H, d, ArH), 8.42 (1H, d, ArH), 8.92 (1H, s, ArH)

8-Acetyl-3-hydroxy-2-methoxybenzo[c]chromen-6-one (**10**)

[0060] Sodium formate (2.18 g, 32 mmol) and formic acid (4.2 mL, 106.8 mmole) were added to a suspension of **9b** (10.0 g, 26.71 mmol) in a 1:1 mixture of dry THF and absolute methanol (1.5 L) in a 3 liter 3-necked flask equipped with an overhead stirrer and a heating mantle. To this mixture was added 100 mg of 10% palladium on carbon and the reaction refluxed under argon for 7 hours. At this time, all of the starting **9b** had gone into solution. The solution was filtered hot to remove the catalyst and the solvent removed by rotary evaporation. (If the solution is allowed to cool down, the product will precipitate and can be separated from the catalyst by extracting the solid with 6 liters of refluxing methanol).

The resulting solid (7.8 g) was purified by silica gel chromatography as described below.

[0061] In a typical run, 3.12 g of crude **10** was mixed with 20 g of silica gel, suspended in 200 mL of methanol and the solvent removed by rotary evaporation. This material was placed on top of a silica gel column (6 cm x 36 cm, 400 g of silica gel), and eluted with a stepwise gradient of 1% acetone / dichloromethane, 10% acetone / dichloromethane and 100% acetone. All pure fractions were combined and evaporated to give 2.4 g (80% yield) of the desired intermediate **10**. ¹H (300 MHz) (DMSO-d₆) δ 2.07 (3H, s), 2.66 (3H, s), 3.92 (3H, s), 6.81 (1H, s), 7.78 (1H, s), 8.33 (1H, d, J = 8.7 Hz), 8.46 (1H, d, J = 8.7 Hz) and 8.66 (1H, d, J = 1.8 Hz).

Scheme 3

[0062] SG00393. SG00392 (1.0 g, 2.67 mmol) and NaBH₄ (0.1 g, 2.67 mmol) were added to a 2:1 mixture of THF (20 mL) and absolute ethanol (10 mL) and left to stir for 1.5 h. The reaction mixture was cooled in an ice bath and 0.5 N HCl added until the color changed from yellow to clear. Water (20 mL) was added and the mixture extracted with CH₂Cl₂, dried, concentrated and the residue purified by silica gel flash column chromatography using CH₂Cl₂:acetone (8/1) to give 0.71 g of SG00393.

[0063] SG00394. SG0093 (0.1 g, 0.27 mmol) was added to anhydrous CH₂Cl₂ (6 mL) and cooled to -78°C giving a heterogeneous mixture. DIBAL (1M in hexanes, 0.66 mL, 0.66 mmol) was added dropwise over 2 h. An additional amount of DIBAL was added (0.2 mL) and after a total time of 2.5 h the reaction was quenched by the addition of methanol (0.8 mL). The reaction mixture was allowed to come to room temperature, CH₂Cl₂ (100 mL), ice and a small amount of acetone were added and the mixture stirred for 15 min. The CH₂Cl₂ layer was washed with sat NaHCO₃, brine, dried and concentrated. The residue was re-dissolved in acetone (40 mL) and pre-adsorbed onto silica gel (1 g). After evaporation of the acetone the residue was purified by silica gel flash column chromatography using CH₂Cl₂:acetone (6/1) to give 69 mg of SG0094.

[0064] SG00395. Crude SG00394 (1.18 g, 3.12 mmol), triethylamine (1.73 mL, 12.5 mmol), acetic anhydride (1.18 mL, 12.5 mmol) and anhydrous CH₂Cl₂ (50 mL) were stirred at room temperature under N₂. Once crystal of DMAP was added, the reaction mixture stirred for 15 min, then extracted with CH₂Cl₂. The CH₂Cl₂ layer was washed with sat NaHCO₃, brine, dried, concentrated and purified by silica gel flash column chromatography using CH₂Cl₂:acetone (10/1) to give 0.89 g of SG00395.

[0065] SG00396. SG00395 (0.83 g, 0.197 mmol) was added to anhydrous CH₂Cl₂ (25 mL) and cooled in a methanol/dry ice bath under N₂. Et₃SiH (0.631 mL, 3.95 mmol) was added followed by BF₃ Et₂O (0.375 mL, 2.96 mmol) dropwise and stirred vigorously for 0.5 h. The reaction mixture was removed from the cooling bath and after 45 minutes quenched with sat NaHCO₃ (3 mL). The reaction mixture was extracted with CH₂Cl₂, washed with sat. NaHCO₃, brine, dried, concentrated and purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/2) to give 0.71 g of SG00396.

[0066] SG00397. SG00396 (0.135 g, 0.334 mmol), formic acid (0.525 mL, 1.34 mmol), sodium formate (27 mg, 0.4 mmol), 10% Pd/C (0.3 mol %), anhydrous THF (4 mL) and absolute ethanol (4 mL) were heated to reflux under N₂ for 1.5 h. The reaction was cooled and approximately half of the reaction mixture evaporated. The silica gel residue was purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/2) to give 50 mg of SG00397.

[0067] SG00398. To the remaining half of the reaction mixture in the preparation of SG00397 was added additional 10% Pd/C and the reaction refluxed for 0.5 h. The Pd/C was filtered off, washed with methanol and silica gel added to the filtrate. After concentrating, the silica gel residue was purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/2) to give 32 mg of SG00398.

[0068] SG00399. SG00395 (0.44 g, 1.09 mmol) and Amberlyst-15 resin (12-15 beads) were stirred in methanol (10 mL) under N₂ for 2 h. The Amberlyst was filtered, washed with methanol and the filtrate concentrated. The residue was purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/2) to give 0.4g of SG00399.

[0069] SG00400. SG00397 (95 mg, 0.304 mmol) was added to methanol (2 mL). To this mixture K₂CO₃ (0.126 g, 0.912 mmol) and water (0.1 mL) were added and the reaction stirred under N₂ for 3 h. The reaction was stopped by the addition of 1% HCl (0.1 mL) and methanol (10 mL). Silica gel was added, the solvent evaporated and the residue was purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/1) to give 72 mg of SG00400. SG00477. SG00292 (0.18 g, 0.63 mmol) was added to anhydrous CH₂Cl₂ (7 mL) with stirring under N₂. Et₃N (0.35 mL, 2.53 mmol), acetic anhydride (0.24 mL, 2.53 mmol) and one crystal of DMAP were added. After stirring for 15 min. CH₂Cl₂ was added and the mixture washed with sat NaHCO₃, brine, dried, concentrated and pre-adsorbed onto silica gel. The silica gel flash column chromatography using ethyl acetate:hexanes (2/1) to give 80 mg of SG00477.

[0070] SG00490. SG00396 (122 mg, 0.3 mmol), K₂CO₃ (125 mg, 0.9 mmol) and water (0.13 mL) were added to methanol (3.3 mL) and stirred under N₂ for 1.5 h then quenched with 1% H₂SO₄. The reaction was extracted with CH₂Cl₂ and divided into two equal portions. One portion was concentrated and purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/1) to give 48 mg of SG00490. The remaining portion was converted to SG00491.

[0071] SG00491. The remaining portion of crude SG00490 was oxidized using the Dess-Martin reagent (37.3 mg, 0.9 mmol) over 1 h. The reaction was extracted with CH₂Cl₂, washed with sat NaHCO₃ brine, dried, concentrated and purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/1) to give 40 mg of SG00491.

[0072] SG00492. Prepared following the method for SG00392 starting with SG00491. Yield 44 mg.

[0073] SG00493. SG492 (116 mg, 0.41 mmol), K₂CO₃ (112 mg, 0.82 mmol) and CH₃I (1 mL) were added to acetone (10 mL) and refluxed for 2 days. Silica gel was added to the reaction mixture, concentrated and purified by silica gel column chromatography using silica gel flash column chromatography using CH₂Cl₂:acetone (9/1) to give 100 mg of SG00493. SGO0494. Prepared following method for SG00493 using 1-(2-chloroethyl)piperidine hydrochloride. Yield 52 mg.

[0074] SG00495. Prepared following method for SG00493 using ethyl bromide. Yield 20 mg.

[0075] SG00496. SG00393 (116 mg, 0.308 mmol) was added to anhydrous THF (10 mL) in an ice bath. NaH (60% dispersion in oil, 22 mg, 0.92 mmol) was added and the mixture stirred for 20 min. CH₃I was added dropwise and the reaction stirred for 0.5 h. The ice bath was removed and the reaction was stirred overnight. Additional CH₃I was added and the reaction mixture refluxed for 5 h. The reaction was quenched with water and distilled to remove the excess CH₃I. CH₂Cl₂ and water were added, and after separating, the CH₂Cl₂ layer was dried, concentrated and purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/1) to give SG00496.

[0076] SG00510. Prepared following the method for SG00393 using SG00493. Yield 48 mg.

[0077] SG00511. Prepared following the method for SG00493 using 2-(bromomethyl) hydrobromide. Yield 170 mg.

[0078] SG00512. Prepared following the method for SG00493 using ethyl bromide. Yield 63 mg.

[0079] SG00513. Prepared following the method for SG00493 using isopropyl bromide. Yield 220 mg.

[0080] SG00514. Prepared following the method of S000493 using 7-hydroxycaurmarin and benzyl bromide. Yield 1.3 g.

[0081] SG00519. Prepared following the method for SG00493 using scopoletin and benzyl bromide. Yield 17 mg.

[0082] SG00520. Prepared following the method for SG00393 using SG00494. Yield 75 mg.

[0083] SG00521. Prepared following the method for SG00393 using SG00512. Yield 118 mg.

[0084] SG00526. SG00511 (50 mg, 0.133 mmol) and NaBH₄ (5.0 mg, 0.133 mmol) were added to a 1:1 mixture of ethanol and THF (10 mL total) and left to stir for 48 h, then refluxed for 2 h. After cooling the reaction mixture was acidified to pH 2 with 1N HCl then taken to pH 8 with sat NaHCO₃ and extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with water, brine, dried and concentrated. The residue was purified by silica gel chromatography using a gradient of hexanes:CHCl₃ (1/1) following by CHCl₃ following by 3% CH₃OH/CHCl₃ to give 40 mg of SG00526.

[0085] SG00527. SG292 (100 mg, 0.35 mmol) was added to a mixture of NaH (15.4 mg, 0.4 mmol) in DMF (10 mL) and the reaction mixture refluxed for 2 h. After cooling down to room temperature 4-methoxybenzyl bromide (0.57 mL, 0.42 mmol) dissolved in DMF was added and the reaction mixture heated to 70°C for 9 h. Water (10 mL) was added

and the reaction mixture extracted with CHCl_3 (3 x 20 mL), the combined organic layers were washed with water, brine, dried and concentrated. The residue was purified by hexaacs: CHCl_3 (1/2) followed by CHCl_3 to give 85 mg of SG00527.

[0086] SG00528. Prepared following method for SG00526 using SG00530. Yield 20 mg.

[0087] SG00529. Prepared following method for SG00526 using SG00527. Yield 40 mg.

[0088] SG00530. Prepared following method for SG00527 using 3-methoxybenzyl bromide. Yield 110mg.

[0089] SG00531. Prepared following method for SG00393 using SG00495. Yield 36 mg.

[0090] SG00532. Prepared following method for SG00393 using SG00513. Yield 71 mg.

[0091] SG40533. Prepared following method for SG00393 using SG00273. Yield 8 mg.

[0092] SG00541. Prepared following methods for SG00527 using 2-methoxybenzyl chloride. Yield 80 mg.

[0093] SG00542. Prepared following method for SG00527 using 2-(chloromethyl)phenyl acetate. Yield 80 mg.

[0094] SG00543. To SG00392 (0.19 g, 0.51 mmol) in anhydrous CH_2Cl_2 (8 mL) was added CH_3MgI (0.2 mL, 1.6 mM) dropwise with stirring at room temperature under N_2 . After 25 min additional CH_3MgI (0.2 mL, 1.6 mM) was added. After 1 h still additional CH_3MgI (0.2 mL, 1.6 mM) was added. The CH_2Cl_2 was separated, washed with slightly acidic water, silica gel added and the CH_2Cl_2 evaporated to pre-adsorb the crude reaction. The dimethyl alcohol was purified by silica gel flash column chromatography using ethyl acetate:hexanes (2/1) and used in the next step. The dimethyl alcohol (58 mg, 0.15 mmol) was debenzylated following the method for SG00292 to give SG00543. Yield 32 mg.

[0095] SG00544. Prepared following method for SG00526 using 2-chloromethylphenyl acetate. Yield 15 mg.

[0096] SG00545. Prepared following method for SG00526 starting with SG00541. Yield 40 mg.

[0097] SG00546. Prepared following method for SG00527 using 3,5-dimethoxybenzyl chloride. Yield 80 mg.

[0098] SG00547. Prepared following method for SG00526 starting with SG00546. Yield 30 mg.

[0099] SG00548. The dimethyl alcohol (56 mg, 0.14 mmol) produced in the preparation of SG00543 was added to anhydrous CH_2Cl_2 (3 mL) containing a catalytic amount of Amberlyst-15 and MgSO_4 and stirred for 6 h and then placed in the freezer overnight. After filtering, the crude dehydration product was purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/2) and used in the next step. The purified dehydration product was dissolved in absolute ethanol (3 mL) and a suspension of 10% Pd-C (30 mg) in absolute ethanol (1.5 mL) was added and a balloon filled with H_2 attached. After stirring for 7 h the catalyst was filtered off, the crude reaction pre-adsorbed onto silica gel and purified by silica gel flash column chromatography using ethyl acetate:hexanes (1/2) to give SG00548.

[0100] SG00549. To a suspension of NaH (0.02 g, 0.55 mmol) in anhydrous DMF (5 mL) was added SG00391 (0.1 g, 0.37 mmol). The resulting yellow opaque mixture was refluxed for 1 h. Benzyl bromide (0.05 mL, 0.41 mmol) was added and the mixture became an orange/yellow clear solution. The reaction mixture was cooled added to water (15 mL) and extracted with ethyl acetate (3 x 12 mL). The organic layer was washed with brine, dried, concentrated and purified by flash silica gel chromatography using 15% ethyl acetate in hexanes to give SG00549 in a quantitative yield.

[0101] SG00550. Prepared following method for SG00549 using 4-methoxybenzyl bromide. Yield 100 mg.

[0102] SG00551. Prepared following method for SG00549 using 2-methoxybenzyl bromide. Yield 62 mg.

[0103] SG00552. Prepared following method for SG00549 using 3-methoxybenzyl bromide. Yield 70 mg.

[0104] SG00553. Prepared following method for SG00526 starting with SG00555. Yield 20 mg.

[0105] SG00554. Prepared following method for SG00526 starting with SG00556. Yield 24 mg.

[0106] SG00555. Prepared following method for SG00527 using 3-chloromethylpyridine hydrochloride. Yield 53 mg.

[0107] SG00556. Prepared following method for SG00527 using 4-chloromethylpyridine hydrochloride. Yield 45 mg.

[0108] SG00557. Prepared following method for SG00527 using 4-(chloromethyl)phenyl acetate. Yield 5 mg.

[0109] SG00558. Prepared following method for SG00527 using 4-(chloromethyl)phenyl. Yield 45 mg.

[0110] SG00559. Prepared following method for SG00527 using 4-methylbenzyl bromide. Yield 58 mg.

[0111] SG00560. Prepared following method for SG00549 using 4-bromobenzyl bromide. Yield 60 mg.

[0112] SG00561. Prepared following method for SG00549 using 3-bromobenzyl bromide. Yield 100 mg.

[0113] SG00562. Prepared following method for SG00549 using 3-chlorobenzyl bromide. Yield 80 mg.

[0114] SG00568. Prepared following method for SG00527 using 2-bromoethyl benzene. Yield 18 mg.

[0115] SG00569. Prepared following method for SG00543 using PhMgBr . Yield 19 mg.

[0116] SG00570. Prepared following the preparation of the dehydration product in the synthesis of

[0117] SG00548 using the diol side product generated in the preparation of SG00549. Yield 21 mg.

[0118] SG00571. Prepared following method for SG00549 using 4-chlorobenzyl bromide. Yield 90 mg.

[0119] SG00572. Prepared following method for SG00549 using 4-fluorobenzyl bromide. Yield 110 mg.

[0120] SG00573. Prepared following method for SG00549 using methyl 4-(bromomethyl)benzoate. Yield 40 mg.

[0121] SG00574. Prepared following method for SG00549 using 4-bromomethyl benzophenone. Yield 30 mg.

[0122] SG00575. Prepared following method for SG00526 starting with SG00559. Yield 25 mg.

[0123] SG00576. Prepared following method for SG00527 using 3-methylbenzyl bromide. Yield 30 mg.

[0124] SG00577. Prepared following method for SG00527 using 3,4,S-trimethoxybenzyl bromide. Yield 45 mg.

[0125] SG00592. Prepared following method for SG00527 using 4-methoxybenzyl chloride and 3-(4-bromophenyl)-7-hydroxycoumarin. Yield 62 mg.

- [0126] SG00593. Prepared following method for SG00592 using 3,5-dimethoxybenzyl bromide. Yield 74 mg.
- [0127] SG00594. Prepared following method for SG00527 using 4-trifluoromethylbenzyl chloride. Yield 22 mg.
- [0128] SG00595. Prepared following method for SG00527 using 4-fluorobenzyl chloride. Yield 43 mg.
- [0129] SG00596. Prepared following method for SG00549 using 3,5-dimethoxybenzyl bromide. Yield 30 mg.
- [0130] SG00597. Prepared following method for SG00527 using ethyl bromoethyl acetate. Yield 15 mg.
- [0131] SG00598. Prepared following method for SG00527 using SG00293 (the ketone of SG00292 reduced to the alcohol). Yield 16 mg.
- [0132] SG00599. From the reaction to prepare SG00569, SG00599 was also isolated. Yield 3.3 mg.
- [0133] SG00609. Prepared following method for SG00526 starting with SG00577. Yield 32 mg.
- [0134] SG00612. SG00292 (0.1g, 0.35 mmol) was added to anhydrous CH₂Cl₂ (10 mL) with stirring. Pyridine (0.05 mL) and benzoyl chloride (0.1 mL) were added and the reaction stirred for 1 h. The reaction was poured into 5% HCl, extracted with CH₂Cl₂, washed with sat NaHCO₃, dried, concentrated and purified by flash silica gel chromatography using ethyl acetate:hexanes (1/1) to give 25 mg of SG00612.
- [0135] SG00613. Prepared following method for SG00612 using 4-methoxybenzyl chloride. Yield 2.3 mg.
- [0136] SG00614. SG00547 (50 mg, 0.114 mmol) was dissolved in a 1:1 mixture of anhydrous diethyl ether and CH₂Cl₂ (6 mL). PBr₃ (124 mg, 0.46 mmol) was added and the reaction stirred over the weekend at room temperature. Sat NaHCO₃ was added and the reaction extracted with CH₂Cl₂, washed with brine, dried, concentrated and purified by flash silica gel chromatography using hexanes then CHCl₃ then 1% methanol in CHCl₃ to give 20 mg of SG00614.
- [0137] SG00615. Prepared following the method for SG00543 starting with SG00546 and EtMgBr. Yield 55 mg.
- [0138] SG00616. Prepared following the method for SG00543 starting with SG00546 and CH₃MgI. Yield 74 mg.
- [0139] SG00617. Prepared following method for SG00527 using SG00293 (the ketone of SG00292 reduced to the alcohol) and 4-bromomethyl benzophenone. Yield 13 mg.
- [0140] SG00618. 4-benzyloxybenzoic acid (1 g, 4.4 mmol) was added to anhydrous CH₂Cl₂ (11 mL). A catalytic amount of DMF (5 drops) was added along with oxalyl chloride in CH₂Cl₂ (2M, 5.75 mL) and the reaction stirred for 2 h. The solvents were evaporated and the crude 4-benzyloxybenzoyl chloride was used directly. SG00618 was prepared following the method for SG00612 using 4-benzylbenzoyl chloride. Yield 49 mg.
- [0141] SG00619. Prepared following the method of SG00527 but using SG00293 (the methyl ketone of SG00292 reduced to the alcohol) and 4-formylbenzyl bromide (prepared by DIBAL reduction of 4-cyanobenzyl bromide, Yield 45 mg).
- [0142] SG00620. Prepared following the method for SG00527 using 4-nitrobenzyl bromide. Yield 20 mg.

Experimental Date

[0143] The following examples refer to representative illustrations from the compounds depicted in Table 1. The aforementioned compounds are found to have anti-proliferative, anti-angiogenic properties and/or other meaningful activities to be described below.

[0144] Example 1: Anti-tumor (anti-proliferative for cancer cells), anti-angiogenic activity (anti-proliferative for endothelial cells) measured in vitro as inhibition of proliferation binding to estrogen receptor alpha and beta HUVEC Proliferation. Inhibition of the proliferation of human umbilical vein endothelial cells, HUVECs, is shown as one measure of anti-angiogenic activity. HUVECS and the required media complements were purchased from Cascade Biologies (Portland, OR) and the growth and maintenance of the cultures was as described by the manufacturer. The proliferation assay was carried out by seeding the HUVECs in 96-well plates at a density of 1,000 cells/well in complete medium. Following a 24 h plating period, the cells were starved for 24 h in 0.5% serum before being treated with SG ("Signal Gene" now "Palomid") angiogenic inhibitors in the presence of 10 ng/ml b-FGF or dosing ranging presence of either b-FGF or VEGF in complete medium. After 48 h, cell number was determined using a calorimetric method as described by the supplier (Promega Corp., Madison, WI). The results were expressed as the percentage of the maximal b-FGF or VEGF response in the absence of angiogenic inhibitors. Non-proliferating endothelial cells were assayed by growing HUVECs to quiescence in 96-well plates and treating with angiogenic inhibitors for 48 h. Initially, 5,000 cells/well were seeded and confluence was achieved the next day. The plates were incubated another 24 h to ensure growth arrest before treatment with angiogenic inhibitors. Cell number was determined as outlined above.

[0145] Cancer Cell Lines. Measurement of the inhibition of tumor cell growth is one measure of anti-cancer activity. Two human cancer cell lines were used to assess the effects of SG angiogenic inhibitors on the proliferation of these cells. The cell lines were MCF-7 breast cancer cells and the colon carcinoma cell line, HCT-116. All cell lines were obtained from American Type Tissue Culture (Manassas, VA) and maintained in their respective media as described by the supplier. The proliferation studies were carried out essentially as described for the proliferating endothelial cells.

[0146] ER Binding Assay. Derivatives which bind and transduce a signal through estrogen receptors would not be considered a positive activity as such an activity could enhance cancer growth as well as induce angiogenesis. Derivatives which either have little or no binding to estrogen receptors ("ER") would be one desired activity. Alternatively, derivatives which bound to estrogen receptors but did not transduce a signal could also be considered a positive activity. Human

cDNAs encoding ERa and ERb were used as templates to express receptor proteins in vitro. The proteins were produced with rabbit reticulocyte lysates as supplied by Promega (TNT kit) that couples transcription and translation in a single reaction. The amount of template used in each reaction was determined empirically and expression was monitored in parallel reactions where [35S]methionine was incorporated into the receptor followed by gel electrophoresis and exposure to film. Binding reactions were carried out in 100 mL final volumes in TEG buffer (10 mM tris, pH 7.5, 1.5 mM EDTA, 10% glycerol). Five (5) mL of in vitro transcribed-translated receptor was used in each binding reaction in the presence of 0.5 nM [3H]estradiol (E2). All compounds were routinely tested from 10⁻¹¹ M to 10⁻⁶ M and were diluted in ethanol. The reactions were incubated at 4°C overnight and bound E2 was quantified by adding 200 mL dextran-coated charcoal. After a 15 min rotation at 4 °C, the tubes were centrifuged for 10 min and 150 mL of the supernatant was added to 5 mL scintillation cocktail for determination of cpm by liquid scintillation counting. Controls for background were included in each experiment using 5 mL unprogrammed rabbit reticulocyte lysate. This value, typically 10 - 15% of the maximal counts was subtracted from all values. The maximum binding was determined by competing bound E2 with only the ethanol vehicle. This value was set to 100% (maximal E2 binding). Values for percent inhibition were calculated based on the maximal E2 binding. The data were plotted and Ki values calculated using the Prism Software. Experiments were conducted at least three times in duplicate.

[0147] The results are shown in Table II and Figure 1. Activity of derivatives show anti-angiogenic activity through inhibition of the proliferation of angiogenic cytokine stimulated endothelial cells. The majority of the derivatives lack the ability to bind to estrogen receptors alpha and beta hence would not be expected to signal through these receptors, a possible stimulator of angiogenesis.

[0148] Activity of derivatives falls into two groups, those which have dual anti-tumor and anti-angiogenic activity and those which have primarily anti-angiogenic activity. See Table II below.

Table II

Palomid	HUVECP % inhibition 3mM	HUVECP % inhibition 0.3mM	HUVECq % inhibition 3mM	Colon % inhibition 3mM	Breast % inhibition 3mM	hERa % binding	hERb % binding
529	113	65	31	20	32	na	na
547	106	42	25	17	35	na	na
575	104	41	33	10	31	na	na
545	100	32	22	21	17	na	na
528	80	<10	25	11	16	41	31
550	77	nd	14	<10	28	na	na
574	74	13	29	24	na	na	na
393	71	nd	21	26	11	na	na
551	62	nd	na	na	25	na	na
573	145	nd	<10	<10	na	na	na
546	100	18	23	na	13	na	na
559	96	72	35	na	17	na	na
568	78	nd	14	na	<10	na	37
560	53	nd	na	na	na	na	na
na, no activity; HUVECP, HUVEC proliferating; HUVECq, HUVEC quiescent; hERa, human estrogen receptor alpha; hERb, human estrogen receptor beta							

Example 2: Apoptotic activity of derivatives

[0149] Apoptosis Assay. The apoptosis assay was conducted to determine if the derivatives inhibited cellular proliferation by inducing programmed cell death. Representative apoptotic activity is shown for endothelial cells with activity implied for other proliferating cells such as keratinocytes. Apoptosis of endothelial cells is yet another means to show anti-angiogenic activity. Cell death is monitored by quantifying the amount of cytoplasmic histone-associated DNA fragments that accumulate in the cell. Apoptosis assay kit was supplied by Roche (cat # 1 544 675) with ELISA detection

and a monoclonal anti-histone antibody. Briefly, HUVECs or keratinocytes were trypsinized, diluted, and aliquoted into microfuge tubes at a concentration of 50,000 cells/tube. Treatment with a compound was for six hours at 37°C followed by cell lysis and analysis using the detection kit according to the manufacturer. Apoptosis was quantified calorimetrically at an absorbance of 405 nm. Controls consisted of a negative vehicle (ctl) control (1% ethanol) and a positive camptothecin (CAM) control at 4 mg/mL in ethanol. See Figure 2. For keratinocyte assay, Palomid 529, a leading clinical candidate available from Paloma Pharmaceuticals, was added at 100 μ M. Results are shown in Figure 3. (Palomid 529 is "SG00529", see Table I).

Example 3: Anti-keratinocyte activity measured in vitro as inhibition of proliferation of keratinocytes and induction of apoptosis

[0150] Skin diseases at least in part are due to abnormal presence and proliferation of keratinocytes. Means to either inhibit said keratinocyte proliferation and/or the ability to induce apoptosis of keratinocytes in said diseases would be expected to aid in the amelioration of abnormal skin pathologies. The following represents illustrative data to support this supposition.

[0151] Keratinocyte proliferation. A benzo[c]chromen-6-one derivative, Palomid 529, was examined with the following protocol. Low passage human keratinocytes (NHEK, neonatal-pooled, p3-5, Cambrex, CC2507) were seeded in black 96-well plates at 1,000 cells per well in complete media (KBM, Cambrex) and incubated overnight. Cell culture media was then removed and replaced with fresh growth media plus either Palomid 529 or vehicle (1% DMSO). Palomid 529 was examined at nine concentrations (1:2 dilutions starting at 100 μ M). Each experimental and control group was examined in replicates of six. Cultures were maintained for 3 days and cellular proliferation was then analyzed by determining metabolically active cells with a fluorescence-based assay (Alamar Blue, Invitrogen, diluted 1:20 in growth media, read following a five hour incubation, 530ex/580em). See Figure 4.

[0152] Keratinocyte apoptosis. Palomid 529 was examined with the following protocol. Low passage human keratinocytes (NHEK, neonatal-pooled, p3-5, Cambrex, CC2507) were seeded in black 96-well plates at 3,000 cells per well in complete media (KBM, Cambrex) and incubated overnight (same plate as cell viability assay, adjacent wells). Cell culture media was then removed and replaced with fresh growth media plus either Palomid 529 or vehicle (1% DMSO). Palomid 529 was examined at seven concentrations (100, 30, 10, 1, 0.3, 0.1, & 0.03 μ M). Each experimental and control group was examined in replicates of three. Following an overnight incubation, culture media was removed and the cells lysed with lysis reagent supplied with the Roche Cell Death ELISA plus kit. Lysed cells were then transferred to clean eppendorf tubes and centrifuged at 200 x g to remove nuclei and cell debris.


[0153] Supernatants, containing the cytoplasmic fraction (including cytoplasmic nucleosomes), were then transferred to streptavidin coated plates and incubated with anti-Histone (biotin conjugated) and anti-DNA (POD conjugated) antibodies for two hours. Wells were then carefully washed. ABTS was then added to the wells and following the development of color (approximately 30 minutes), the reaction stopped by the addition of ABTS Stop solution. Apoptosis was then measured by reading the OD at 405nm with a reference background wavelength of 490nm. See Figure 5.

Example 4: Metabolic stability assay using human primary hepatocytes

[0154] The cell based assay serves to determine the stability or half-life of compounds in cells. These specialized hepatocytes contain all of the necessary phase I and phase II enzymes that can act upon drugs. Compounds that are not or hardly metabolized in these cells are thought to be metabolically stable and would be expected to have a longer half-life in vivo than those that are metabolized by the hepatocytes. Results are shown in Figure 6. Derivatives were incubated with human primary hepatocytes for 120 or 240 minutes. % remaining compound is shown at left. Control derivative capable of metabolism by phase I and/or phase II enzymes is shown as derivative 292.

[0155] Features:

1. A composition comprising a compound according to Formula I:



wherein,

R1 is H or alkyl;

R2 is H, OH, O-alkyl, amino, O-heterocyc, O-aryl, O-Ac, O-PO₃, O-SO₃, OSO₂NH₂ or O-substituted alkyl wherein said substitution is halo, aryl, or heteroaryl;

R3 is H, OH, O-alkyl, O-CH₂Aryl, O-CH₂heteroaryl, O-alkylaryl, O-acyl, or nitro;

R4 is H, Alkyl, CH₂Aryl, substituted alkyl, OH, O-alkyl, O-aryl, OCH₂Aryl, OCH₂Heteroaryl, O-Acyl, OPO₃, OSO₃, or OSO₂NH₂;

R5 is H, Oxo, aryl, hydroxyl, alkyl, or O-alkyl;


R6 is H;

R7 is H, Acyl, alkyl, O-alkyl, substituted alkyl wherein said substitution is hydroxyl or sulfamoyl, or O-substituted alkyl wherein said substitution is O-PO₃ or OSO₃;

R8 is H; and

X is O, N, or S.

2. A composition comprising a compound according to Formula II:



wherein,

R1 is H or alkyl;

R2 is H, O-alkyl, OH, amino, O-heterocyc, O-aryl, O-Ac, O-PO₃, O-SO₃, OSO₂NH₂ or O-substituted alkyl wherein said substitution is halo, aryl, or heteroaryl;


R3 is H, O-alkyl, OH, O-acyl, nitro or O-substituted alkyl wherein substitution is aryl or heteroaryl;

R4 is H, Alkyl, CH₂Aryl, substituted alkyl, OH, O-alkyl, O-aryl, OCH₂Aryl, OCH₂Heteroaryl, O-Acyl, OPO₃, OSO₃, or OSO₂NH₂;

R5 is H, Aryl, heteroaryl or substituted alkyl; and

R6 is H, Alkyl, or Aryl.

3. A composition comprising a compound according to Formula III:



wherein,


R1 is alkyl or H;

R2 is alkyl or H;

R3 is Acetyl; and

R4 is H or Alkyl.

4. A composition comprising a compound according to Formula IV:



wherein,

R1 is H or F;

R2 is H or nitro;

R3 is H;

R4 is H; and

R5 is alkyl, substituted alkyl or aryl.

5. The composition of any one of features 1-4, wherein said composition is one or more compositions selected from the group consisting of Table I.

6. The composition of feature 5, wherein said Table I comprises benzo(c)chromen-6-one derivatives SG00272, SG00273, SG00373, SG00477, SG00519, SG00526, SG00527, SG00528, SG00529, SG00530, SG00531, SG00532, SG00533, SG00535, SG00536, SG00537, SG00538, SG00539, SG00540, SG00541, SG00542, SG00543, SG00544, SG00545, SG00546, SG00547, SG00548, SG00549, SG00550, SG00551, SG00552, SG00553, SG00554, SG00555, SG00556, SG00557, SG00558, SG00559, SG00560, SG00561, SG00562, SG00563, SG00564, SG00565, SG00566, SG00567, SG00568, SG00569, SG00570, SG00571, SG00572, SG00573, SG00574, SG00575, SG00576, SG00577, SG00579, SG00580, SG00581, SG00582, SG00583, SG00584, SG00585, SG00586, SG00587, SG00588, SG00589, SG00590, SG00591, SG00592, SG00593, SG00594, SG00595, SG00596, SG00597, SG00598, SG00599, SG00600, SG00601, SG00602, SG00603, SG00604, SG00605, SG00606, SG00607, SG00608, SG00609, SG00610, SG00611, SG00612, SG00613, SG00614, SG00615, SG00616, SG00617, SG00618, SG00619, SG00620 and SG00629.

7. A composition for preventing or treating a disease characterized by unwanted angiogenesis comprising a compound selected from the group consisting of Formula I, Formula II, Formula III and Formula IV.

8. The composition of feature 7, wherein said disease is further characterized by unwanted cellular proliferation.

9. The composition of feature 8, wherein said composition exhibits anti-angiogenic activity or anti-proliferation activity.
10. The composition of feature 8, wherein said composition exhibits anti-angiogenic activity and anti-proliferation activity.
11. The composition of feature 8, wherein said composition exhibits apoptotic activity.
12. The composition of any one of features 1-11, wherein said composition is formulated in a biodegradable or non-biodegradable format for sustained release.
13. The composition of any one of features 1-11, wherein said composition is conjugated to a prodrug.
14. The composition of feature 13, wherein said prodrug is selected from the group consisting of a peptide, antibody, antibody fragment, hydrolysable ester amides, and carbamates.
15. The composition of feature 13, wherein said prodrug is selected from the group consisting of glycolic acid, lactic acid, 2-hydroxyoctanoic acid, hydroxylauric glycolic acid, and combinations thereof.
16. A method of preventing or treating a disease characterized by unwanted angiogenesis, comprising administering to a subject a therapeutic amount of one or more compositions selected from the group consisting of Formula I, Formula II, Formula III and Formula IV.
17. The method of feature 16, wherein said composition further comprises an acceptable delivery vehicle.
18. The method of feature 16, wherein said subject has or is predisposed toward a cancer.
19. The method of feature 18, wherein said cancer is selected from the group consisting of fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, hemangioma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinomas, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, osteosarcoma, meningioma, melanoma, neuroblastoma, retinoblastoma, acousticneuroma, neurofibromas, trachoma and pyogenic granulomas, acute lymphocytic leukemia and acute myelocytic leukemia, chronic leukemia, polycythemia vera, lymphoma, multiple myeloma, Waldenstrom's macroglobulinemia, and heavy chain disease.
20. The method of feature 16, wherein said subject has or is predisposed toward a disease selected from the group consisting of hereditary hemorrhagic telangiectasia, solid or blood born tumors, acquired immune deficiency syndrome, post-menopausal symptoms, osteoporosis, cardiovascular disease, Alzheimer's disease, strokes, vascular malformations, abnormal wound healing, inflammatory and immune disorders, gout, macular degeneration, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, multifocal choroiditis, Best's disease, Stargardt's disease, cblc-type cobalamin deficiency, hyperviscosity syndrome, Sorsby's fundus dystrophy, pseudoxanthoma elasticum, rubeosis iridis, Osler Weber syndrome, Osler-Weber-Rendu disease, keratoconjunctivitis, Vitamin A deficiency, phlyctenulosis, bacterial infection, viral infection, parasitic infection, fungal infection, atopic and superior limbic dermatitis, chronic uveitis, chronic vitritis, Eales' disease, radial keratotomy, uncharacteristic proliferation of fibrovascular or fibrous tissue, proliferative vitreoretinopathy, chronic retinal detachment, trauma (including but not limited to abrasion, previous surgery with complications such as corneal allograft rejection, alkaline burns, acid burns or hydrocarbon burns, mechanical or thermal damage to Bruch's membrane), pterygium, arthritis (rheumatoid and osteoarthritis), psoriasis, atopic dermatitis, dermal photodamage), Sjögren's syndrome, systemic lupus, polyarteritis, pemphigoid, sickle cell anemia, Paget's disease, vein or artery occlusion, carotid obstructive disease, Lyme disease, Behcet's disease, bartonellosis, arteriosclerosis, induction of amenorrhea to block ovulation or to prevent implantation by the blastula, surgical adhesions, chronic inflammation, ulcerative colitis and Crohn's disease.
21. The method of feature 16 further comprising the co-administration to said subject said one or more compositions

selected from the group consisting of Formula I, Formula II, Formula III and Formula IV and a therapeutic agent directed toward the treatment or prevention of said disease.

22. The method of feature 21, wherein said therapeutic agent is selected from the group consisting of an oncolytic agent, an anti-cancer agent, an anti-nausea agent and an anti-emesis agent.

23. The method of feature 16, wherein said administration includes topical, oral, nasal, rectal, and parenteral administration of said one or more compositions.


24. The method of feature 16, wherein said one or more compositions is associated with an implant.

25. The method of feature 16, wherein said one or more compositions is associated with a device.

26. The method of feature 25, wherein said device is a vascular stent.

Claims

1. A composition comprising a compound according to Formula IV:



wherein,

R1 is H or F;





























R2 is H or nitro;

































R3 is H;


R4 is H; and

R5 is alkyl, substituted alkyl or aryl.

2. A composition, wherein said composition comprises one or more compounds selected from the group consisting of Table I, wherein said Table I comprises benzo(c)chromen-6-one derivatives SG00393, SG00477, SG00526, SG00527, SG00528, SG00529, SG00530, SG00541, SG00542, SG00543, SG00544, SG00545, SG00546, SG00547, SG00548, SG00549, SG00550, SG00551, SG00552, SG00553, SG00554, SG00555, SG00556, SG00557, SG00558, SG00559, SG00560, SG00561, SG00562, SG00568, SG00569, SG00570, SG00571, SG00572, SG00573, SG00574, SG00575, SG00576, SG00577, SG00579, SG00582, SG00583, SG00584, SG00587, SG00594, SG00595, SG00596, SG00597, SG00598, SG00599, SG00600, SG00602, SG00603, SG00604, SG00605, SG00606, SG00607, SG00609, SG00612, SG00613, SG00614, SG00615, SG00616, SG00617, SG00618, SG00619, SG00620 and SG00629 having the structural formulae given in the Table below:

5	 SG00393	 SG00477	 SG00526	 SG00527
10	 SG00528	 SG00529	 SG00530	 SG00541
15	 SG00542	 SG00543	 SG00544	 SG00545
20	 SG00546	 SG00547	 SG00548	 SG00549
25	 SG00550	 SG00551	 SG00552	 SG00553
30	 SG00554	 SG00555	 SG00556	 SG00557
35	 SG00558	 SG00559	 SG00560	 SG00561
40				
45				
50				
55				

5	 SG00562	 SG00568	 SG00569	 SG00570
10	 SG00571	 SG00572	 SG00573	 SG00574
15	 SG00575	 SG00576	 SG00577	 SG00579
20	 SG00582	 SG00583	 SG00584	 SG00587
25	 SG00594	 SG00595	 SG00596	 SG00597
30	 SG00598	 SG00599	 SG00600	 SG00602
35	 SG00603	 SG00604	 SG00605	 SG00606
40	 SG00607	 SG00609	 SG00612	 SG00613
45				
50				



- 15
3. The composition of claim 1 or 2 for use in treating a disease or condition, wherein said disease or condition is a cancer, wherein, optionally, said cancer is selected from the group consisting of fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, hemangioma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinomas, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, osteosarcoma, meningioma, melanoma, neuroblastoma, retinoblastoma, acoustic neuroma, neurofibromas, trachoma and pyogenic granulomas, acute lymphocytic leukemia and acute myelocytic leukemia, chronic leukemia, polycythemia vera, lymphoma, multiple myeloma, Waldenstrom's macroglobulinemia, and heavy chain disease; or
- 20
- 25
- 30
- wherein said disease or condition is selected from the group consisting of hereditary hemorrhagic telangiectasia, solid or blood born tumors, acquired immune deficiency syndrome, post-menopausal symptoms, osteoporosis, cardiovascular disease, Alzheimer's disease, strokes, vascular malformations, abnormal wound healing, inflammatory and immune disorders, gout, macular degeneration, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, multifocal choroiditis, Best's disease, Star-gardt's disease, cblC-type cobalamin deficiency, hyperviscosity syndrome, Sorsby's fundus dystrophy, pseudoxanthoma elasticum, rubeosis iridis, Osler Weber syndrome, Osler-Weber-Rendu disease, keratoconjunctivitis, Vitamin A deficiency, phlyctenulosis, bacterial infection, viral infection, parasitic infection, fungal infection, atopic and superior limbic dermatitis, chronic uveitis, chronic vitritis, Eales' disease, radial keratotomy, uncharacteristic proliferation of fibrovascular or fibrous tissue, proliferative vitreoretinopathy, chronic retinal detachment, trauma (including but not limited to abrasion, previous surgery with complications such as corneal allograft rejection, alkaline burns, acid burns or hydrocarbon burns, mechanical or thermal damage to Bruch's membrane), pterygium, arthritis (rheumatoid and osteoarthritis), psoriasis, atopic dermatitis, dermal photodamage), Sjögren's syndrome, systemic lupus, polyarthritis, pemphigoid, sickle cell anemia, Paget's disease, vein or artery occlusion, carotid obstructive disease, Lyme disease, Behcet's disease, bartonellosis, arteriosclerosis, induction of amenorrhea to block ovulation or to prevent implantation by the blastula, surgical adhesions, chronic inflammation, ulcerative colitis and Crohn's disease.
- 35
- 40
- 45
4. The composition for use of claim 3, wherein the composition further comprises a therapeutic agent directed toward the treatment of said disease or condition for co-administration to a subject with one or more compounds as defined in claim 1 or 2.
- 50
5. The composition for use of claim 4, wherein said therapeutic agent is selected from the group consisting of an oncolytic agent, an anti-cancer agent, an anti-nausea agent and an anti-emesis agent.
- 55
6. The composition for use of any one of claims 3 to 5, wherein said composition is formulated for any one of topical, oral, nasal, rectal, and parenteral administration of said one or more compounds.
7. The composition for use of any one of claims 3 to 6, wherein said one or more compounds is associated with an

implant; or wherein said one or more compounds is associated with a device, preferably a vascular stent.

- 5 8. The composition for use of any one of claims 3 to 7, wherein said condition is selected from the group consisting of macular degeneration, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, multifocal choroiditis, Best's disease, Stargardt's disease, Sorsby's fundus dystrophy, pseudoxanthoma elasticum, rubeosis iridis, kera-
toconjunctivitis, phlyectenulosis, chronic uveitis, chronic vitritis, Eales' disease, radial keratotomy, proliferative vit-
reoretinopathy, and chronic retinal detachment.

- 10 9. The composition for use of any one of claims 3 to 8, wherein said condition is macular degeneration.

15

20

25

30

35

40

45

50

55




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6



EUROPEAN SEARCH REPORT

Application Number
EP 12 17 1867

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	SCHMIDT J M ET AL: "Synthesis and evaluation of a novel nonsteroidal-specific endothelial cell proliferation inhibitor", JOURNAL OF MEDICINAL CHEMISTRY 20030410 US, vol. 46, no. 8, 10 April 2003 (2003-04-10), pages 1289-1292, XP002563534, ISSN: 0022-2623 * compounds 9a, 10; scheme 2; last ? of p. 1289 to second ? of p. 1290; penultimate ? of section "conclusion" on p. 1291 *	1,3-9	INV. A61K31/35 A01N43/16 A61K31/352 A61K31/517
X	US 2002/115711 A1 (SCHMIDT JOHNATHAN MARTIN [CA] ET AL SCHMIDT JONATHAN MARTIN [CA] ET AL) 22 August 2002 (2002-08-22) * ?0001; ?0007; ?0010; claim 1, claim 4; example 3; ?0039; ?0046 *	1,3-9	TECHNICAL FIELDS SEARCHED (IPC) A61K C07D
X	WO 02/086078 A2 (UNIV VIRGINIA [US]; BROWN MILTON L [US]) 31 October 2002 (2002-10-31) * p. 7, l. 11-15; p. 1, l. 10-12; p. 16, l. 21-23; p. 29, table 1, compound 49 *	1,3-9	
X	WO 2005/117876 A1 (UNIV VIRGINIA PATENT FOUNDATION [US]; BROWN MILTON L [US] UNIV VIRGINIA) 15 December 2005 (2005-12-15) * claims 2, 14-15; p. 6, first *	1,3-9	
-/--			
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 5 December 2012	Examiner Dahse, Thomas
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

 2
EPO FORM 1503 03.02 (P04C01)



EUROPEAN SEARCH REPORT

Application Number
EP 12 17 1867

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	ES 484 472 A1 (ICI LTD) 16 May 1980 (1980-05-16) * p. 25-26, example 12 *	1	
X	US 6 313 113 B1 (LOHRAY VIDYA BHUSHAN [IN] ET AL) 6 November 2001 (2001-11-06) * col. 33, example 16; table in col. 36; col. 36, l. 30-36; claims 16, 18 *	1,3-9	
X	ISHIGURO I ET AL: "[Isolation and identification of the 2-methyl-1,2,3,4-tetrahydro-4-quinazolone as a new metabolite in the urine of anthranilic acid injected rats].", YAKUGAKU ZASSHI : JOURNAL OF THE PHARMACEUTICAL SOCIETY OF JAPAN OCT 1974, vol. 94, no. 10, October 1974 (1974-10), pages 1232-1239, XP009165390, ISSN: 0031-6903 * title *	1	
			TECHNICAL FIELDS SEARCHED (IPC)
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 5 December 2012	Examiner Dahse, Thomas
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

2
EPO FORM 1503 03.82 (P04C01)



Application Number

EP 12 17 1867

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing claims for which payment was due.

- ☐ Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for those claims for which no payment was due and for those claims for which claims fees have been paid, namely claim(s):
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for those claims for which no payment was due.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet B

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☒ As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:
- ☐ The present supplementary European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims (Rule 164 (1) EPC).



LACK OF UNITY OF INVENTION
SHEET B

Application Number

EP 12 17 1867

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. claims: 1, 3-9(all partially)

directed to compositions of formula IV, wherein R5 is alkyl or substituted alkyl, and their use in the prevention/treatment of a disease characterized by unwanted angiogenesis and/or cellular proliferation.

2. claims: 1, 3-9(all partially)

directed to compositions of formula IV, wherein R5 is (substituted?) aryl, and their use in the prevention/treatment of a disease characterized by unwanted angiogenesis and/or cellular proliferation.

3. claims: 2(completely); 3-9(partially)

directed to compositions comprising SG00393 to SG00629 of claim 2, falling under general formula type I, and their use in the prevention/treatment of a disease characterized by unwanted angiogenesis and/or cellular proliferation.

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 12 17 1867

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-12-2012

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 2002115711	A1	22-08-2002	AT 396989 T	15-06-2008
			AU 2002234466 B2	16-02-2006
			CA 2438713 A1	29-08-2002
			DK 1373260 T3	29-09-2008
			EP 1373260 A1	02-01-2004
			ES 2307721 T3	01-12-2008
			PT 1373260 E	05-09-2008
			US 2002115711 A1	22-08-2002
			WO 02066467 A1	29-08-2002

WO 02086078	A2	31-10-2002	AU 2002257195 A1	05-11-2002
			CA 2444704 A1	31-10-2002
			CN 1518447 A	04-08-2004
			EP 1389103 A2	18-02-2004
			JP 2004532848 A	28-10-2004
			JP 2008308501 A	25-12-2008
			US 2004122030 A1	24-06-2004
			WO 02086078 A2	31-10-2002

WO 2005117876	A1	15-12-2005	AU 2005249527 A1	15-12-2005
			BR PI0511748 A	02-01-2008
			CA 2568622 A1	15-12-2005
			CN 1988907 A	27-06-2007
			EP 1750706 A1	14-02-2007
			JP 4991527 B2	01-08-2012
			JP 2008501695 A	24-01-2008
			KR 20090024309 A	09-03-2009
			NZ 551775 A	24-12-2010
			US 2007134161 A1	14-06-2007
			US 2007244098 A1	18-10-2007
			US 2012202800 A1	09-08-2012
			WO 2005117876 A1	15-12-2005

ES 484472	A1	16-05-1980	NONE	

US 6313113	B1	06-11-2001	US 6011036 A	04-01-2000
			US 6030973 A	29-02-2000
			US 6313113 B1	06-11-2001

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- Remington's Pharmaceuticals Sciences [0037]