

(19)



(11)

EP 2 559 688 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
20.02.2013 Bulletin 2013/08

(51) Int Cl.:
C07D 249/08 (2006.01) **A01N 43/653** (2006.01)

(21) Application number: **11177556.5**

(22) Date of filing: **15.08.2011**

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

Designated Extension States:
BA ME

(71) Applicant: **BASF SE
67056 Ludwigshafen (DE)**

(72) Inventors:

- **Dietz, Jochen, Dr.
76227 Karlsruhe (DE)**
- **Riggs, Richard, Dr.
68165 Mannheim (DE)**

- **Boudet, Nadege, Dr.
69502 Hemsbach (DE)**
- **Lohmann, Jan Klaus, Dr.
67245 Lambsheim (DE)**
- **Craig, Ian Robert, Dr.
67063 Ludwigshafen (DE)**
- **Haden Egon, Dr.
67346 Speyer (DE)**
- **Lauterwasser, Erica May Wilson, Dr.
68199 Mannheim (DE)**
- **Müller, Bernd Dr.
67227 Frankenthal (DE)**
- **Grammenos, Wassilios, Dr.
67071 Ludwigshafen (DE)**
- **Grote, Thomas, Dr.
67157 Wachenheim (DE)**

(54) **Fungicidal substituted 1-[2-[2-halo-4-(4-halogen-phenoxy)-phenyl]-2-butoxy-ethyl]-1H-[1,2,4]triazole compounds**

(57) The present invention relates to substituted 1-[2-[2-halo-4-(4-halogen-phenoxy)-phenyl]-2-butoxy-ethyl]-1H-[1,2,4]triazole compounds of formula I as defined in the description, and the N-oxides, and salts thereof, processes and intermediates for preparing these compounds and also to compositions comprising at least one


such compound.. The invention also relates to the use of such compounds and compositions for combating harmful fungi and seed coated with at least one such compound.

EP 2 559 688 A1

Description

[0001] The present invention relates to fungicidal 1-{2-[2-halo-4-(4-halogen-phenoxy)-phenyl]-2-butoxy-ethyl}-1H-[1,2,4]triazole compounds and the N-oxides and the salts thereof for combating phytopathogenic fungi, and to the use and methods for combating phytopathogenic fungi and to seeds coated with at least one such compound. The invention also relates to processes for preparing these compounds and to compositions comprising at least one such compound.

[0002] Certain 1-{2-[2-chloro-4-(4-chloro-phenoxy)-phenyl]-2-alkoxy-ethyl}-1H-[1,2,4]triazole compounds of formula




wherein R² is methyl, propyl or iso-butyl, and their use for controlling phytopathogenic fungi are known from EP 0 126 430 A2 and US 4,940,720.

[0003] The compounds according to the present invention differ from those described in the abovementioned publications by the specific substituent for R² instead of iso-butyl.

[0004] In many cases, in particular at low application rates, the fungicidal activity of the known fungicidal compounds is unsatisfactory. Based on this, it was an object of the present invention to provide compounds having improved activity and/or a broader activity spectrum against phytopathogenic harmful fungi.

[0005] This object is achieved by substituted 1-{2-[2-halo-4-(4-halogen-phenoxy)-phenyl]-2-butoxy-ethyl}-1H-[1,2,4]triazole compounds having good fungicidal activity against phytopathogenic harmful fungi.

[0006] Accordingly, the present invention relates to the compounds of formula I:



wherein:

40 X¹,X² independently of each other are selected from halogen;

45 R¹ is hydrogen, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₃-C₈-cycloalkyl, C₃-C₈-cycloalkyl-C₁-C₄-alkyl, phenyl, phenyl-C₁-C₄-alkyl, phenyl-C₂-C₄-alkenyl or phenyl-C₂-C₄-alkynyl;

R² is n-butyl, sec.-butyl or tert.-butyl;

50 wherein the aliphatic moieties of R¹ and/or R² may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^a which independently of one another are selected from:

R^a halogen, CN, nitro, C₁-C₄-alkoxy and C₁-C₄-halogenalkoxy;

55 wherein the cycloalkyl and/or phenyl moieties of R¹ may carry 1, 2, 3, 4, 5 or up to the maximum number of identical or different groups R^b which independently of one another are selected from:

R^b halogen, CN, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-halogenalkyl and C₁-C₄-halogenalkoxy;

and the N-oxides and the agriculturally acceptable salts thereof.


[0007] The present invention furthermore relates to the use of these compounds for combating harmful fungi and seed coated with at least one such compound and also to compositions comprising at least one such compound of formula I.

[0008] The present invention furthermore relates to processes for preparing compounds of formula I and to intermediates such as compounds of formula XII.

[0009] The term "compounds I" refers to compounds of formula I. Likewise, this terminology applies to all sub-formulae, e. g. "compounds I.A" refers to compounds of formula I.A or "compounds XII" refers to compounds of formula XII, etc..


[0010] The compounds I can be obtained by various routes in analogy to prior art processes known (cf. J.Agric. Food Chem. (2009) 57, 4854-4860; EP 0 275 955 A1; DE 40 03 180 A1; EP 0 113 640 A2; EP 0 126 430 A2) and by the synthesis routes shown in the following schemes and in the experimental part of this application.

[0011] In a first process, for example, halo-phenoles II wherein X^1 and X^2 as defined herein, are reacted, in a first step, with bromo derivatives III wherein Y is F or Cl, preferably in the presence of a base. Thereafter, the resulting compounds IV are then transformed into Grignard reagents by the reaction with transmetallation reagents such as isopropylmagnesium halides and subsequently reacted with acetyl chloride preferably under anhydrous conditions and preferably in the presence of a catalyst such as CuCl, AlCl₃, LiCl and mixtures thereof, to obtain acetophenones V. These compounds V can be halogenated e.g. with bromine preferably in an organic solvent such as diethyl ether, methyl tert.-butyl ether (MTBE), methanol or acetic acid. The resulting compounds VI can subsequently reacted with 1H-1,2,4-triazole preferably in the presence of a solvent such as tetrahydrofuran (THF), dimethylformamide (DMF), toluene and in the presence of a base such as potassium carbonate, sodium hydroxide or sodium hydride to obtain compounds VII. These triazole compounds VII are reacted with a Grignard reagent R¹-M wherein R¹ is as defined above and M is MgBr, MgCl, Li or Na (e.g. phenylalkyl-MgBr or an organolithium reagent phenylalkyl-Li), preferably under anhydrous conditions to obtain compounds VIII. These compounds VIII are reacted with R²-LG, wherein wherein R¹ is as defined above and LG represents a nucleophilically replaceable leaving group such as halogen, alkylsulfonyl, alkylsulfonyloxy and arylsulfonyloxy, preferably chloro, bromo or iodo, particularly preferably bromo, preferably in the presence of a base to form compounds I. The preparation of compounds I can be illustrated by the following scheme:




[0012] In a second process to obtain compounds I, bromo derivatives III, in a first step, are reacted with e.g. isopropylmagnesium bromide followed by an acyl chloride agent IX wherein R¹ is as defined above (e.g. acetyl chloride) preferably under anhydrous conditions and preferably in the presence of a catalyst such as CuCl, AlCl₃, LiCl and mixtures thereof, to obtain compounds X. Alternatively, 1,3-dichlorobenzene of formula IIIb can be reacted with an acyl chloride agent IX wherein R¹ is as defined above (e.g. acetyl chloride) preferably in the presence of a catalyst such as AlCl₃. Then, ketones X are reacted with phenoles II preferably in the presence of a base to obtain compounds Va.

[0013] Thereafter, intermediates Va are reacted with trimethylsulf(ox)onium halides preferably iodide preferably in the presence of a base such as sodium hydroxide. Thereafter, the epoxides XI are reacted with 1H-1,2,4-triazole preferably in the presence of a base such as potassium carbonate and preferably in the presence of an organic solvent such as DMF to obtain compounds VIII. These compounds VIII are reacted with R²-LG, wherein wherein R¹ is as defined above and LG represents a nucleophilically replaceable leaving group such as halogen, alkylsulfonyl, alkylsulfonyloxy and arylsulfonyloxy, preferably chloro, bromo or iodo, particularly preferably bromo, preferably in the presence of a base to form compounds I., which can subsequently be alkylated as described above. The preparation of compounds I can be illustrated by the following scheme:



[0014] In a third process, the epoxide ring of intermediates XI which may be obtained according to the second process described herein is cleaved by reaction with alcohols R^2OH preferably under acidic conditions. Thereafter, the resulting compounds XII are reacted with halogenating agents or sulfonating agents such as PBr_3 , PCl_3 , mesyl chloride, tosyl chloride or thionyl chloride to obtain compounds XIII wherein LG is a nucleophilically replaceable leaving group such as halogen, alkylsulfonyl, alkylsulfonyloxy and arylsulfonyloxy, preferably chloro, bromo or iodo, particularly preferably bromo or alkylsulfonyl. Then compounds XII are reacted with 1H-1,2,4-triazole to obtain compounds I. The preparation of compounds I can be illustrated by the following scheme:



[0015] If individual compounds I cannot be obtained by the routes described above, they can be prepared by derivatization of other compounds I.

[0016] The N-oxides may be prepared from the compounds I according to conventional oxidation methods, e. g. by treating compounds I with an organic peracid such as metachloroperbenzoic acid (cf. WO 03/64572 or J. Med. Chem. 38(11), 1892-903, 1995); or with inorganic oxidizing agents such as hydrogen peroxide (cf. J. Heterocycl. Chem. 18(7), 1305-8, 1981) or oxone (cf. J. Am. Chem. Soc. 123(25), 5962-5973, 2001). The oxidation may lead to pure mono-N-oxides or to a mixture of different N-oxides, which can be separated by conventional methods such as chromatography.

[0017] If the synthesis yields mixtures of isomers, a separation is generally not necessarily required since in some cases the individual isomers can be interconverted during work-up for use or during application (e. g. under the action of light, acids or bases). Such conversions may also take place after use, e. g. in the treatment of plants in the treated plant, or in the harmful fungus to be controlled.

[0018] In the definitions of the variables given above, collective terms are used which are generally representative for the substituents in question. The term " C_1-C_6 -alkyl" indicates the number of carbon atoms possible in each case in the substituent or substituent moiety in question.

[0019] The term "halogen" refers to fluorine, chlorine, bromine and iodine.

[0020] The term " C_1-C_6 -alkyl" refers to a straight-chained or branched saturated hydrocarbon group having 1 to 6 carbon atoms, e.g. methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl,

1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl. Likewise, the term "C₂-C₄-alkyl" refers to a straight-chained or branched alkyl group having 2 to 4 carbon atoms, such as ethyl, propyl (n-propyl), 1-methylethyl (iso-propyl), butyl, 1-methylpropyl (sec.-butyl), 2-methylpropyl (iso-butyl), 1,1-dimethylethyl (tert.-butyl).

[0021] The term "C₂-C₄-alkenyl" refers to a straight-chain or branched unsaturated hydrocarbon radical having 2 to 4 carbon atoms and a double bond in any position, e.g. ethenyl, 1-propenyl, 2-propenyl (allyl), 1-methylethenyl, 1-but enyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl. Likewise, the term "C₂-C₆-alkenyl" refers to a straight-chain or branched unsaturated hydrocarbon radical having 2 to 6 carbon atoms and a double bond in any position.

[0022] The term "C₂-C₄-alkynyl" refers to a straight-chain or branched unsaturated hydrocarbon radical having 2 to 4 carbon atoms and containing at least one triple bond, such as ethynyl, prop-1-ynyl, prop-2-ynyl (propargyl), but-1-ynyl, but-2-ynyl, but-3-ynyl, 1-methyl-prop-2-ynyl. Likewise, the term "C₂-C₆-alkynyl" refers to a straight-chain or branched unsaturated hydrocarbon radical having 2 to 6 carbon atoms and at least one triple bond.

[0023] The term "C₁-C₄-halogenalkyl" refers to a straight-chained or branched alkyl group having 1 to 4 carbon atoms, wherein some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as mentioned above, for example chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl and pentafluoroethyl, 2-fluoropropyl, 3-fluoropropyl, 2,2-difluoropropyl, 2,3-difluoropropyl, 2-chloropropyl, 3-chloropropyl, 2,3-dichloropropyl, 2-bromopropyl, 3-bromopropyl, 3,3,3-trifluoropropyl, 3,3,3-trichloropropyl, CH₂-C₂F₅, CF₂-C₂F₅, CF(CF₃)₂, 1-fluoromethyl-2-fluoroethyl, 1-(chloromethyl)-2-chloroethyl, 1-bromomethyl-2-bromoethyl, 4-fluorobutyl, 4-chlorobutyl, 4-bromobutyl or nonafluorobutyl, and the like.

[0024] The term "C₃-C₈-cycloalkyl" refers to monocyclic saturated hydrocarbon radicals having 3 to 8 carbon ring members, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl.

[0025] The term "C₁-C₄-alkoxy" refers to a straight-chain or branched alkyl group having 1 to 4 carbon atoms which is bonded via an oxygen, at any position in the alkyl group, e.g. methoxy, ethoxy, n-propoxy, 1-methylethoxy, butoxy, 1-methyprooxy, 2-methyl-propoxy or 1,1-dimethylethoxy.

[0026] The term "C₁-C₄-halogenalkoxy" refers to a C₁-C₄-alkoxy radical as defined above, wherein some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as mentioned above, e.g., OCH₂F, OCHF₂, OCF₃, OCH₂Cl, OCHCl₂, OCl₃, chlorofluoromethoxy, dichlorofluoromethoxy, chlorodifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2-bromoethoxy, 2-iodoethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 2-chloro-2-fluoroethoxy, 2-chloro-2,2-difluoroethoxy, 2,2-dichloro-2-fluoroethoxy, 2,2,2-trichloroethoxy, OC₂F₅, 2-fluoropropoxy, 3-fluoropropoxy, 2,2-difluoropropoxy, 2,3-difluoropropoxy, 2-chloropropoxy, 3-chloropropoxy, 2,3-dichloropropoxy, 2-bromopropoxy, 3-bromopropoxy, 3,3,3-trifluoropropoxy, 3,3,3-trichloropropoxy, OCH₂-C₂F₅, OCF₂-C₂F₅, 1-fluoromethyl-2-fluoroethoxy, 1-chloromethyl-2-chloroethoxy, 1-bromomethyl-2-bromoethoxy, 4-fluorobutoxy, 4-chlorobutoxy, 4-bromobutoxy or nonafluorobutoxy.

[0027] Agriculturally acceptable salts of compounds I encompass especially the salts of those cations or the acid addition salts of those acids whose cations and anions, respectively, have no adverse effect on the fungicidal action of the compounds I. Suitable cations are thus in particular the ions of the alkali metals, preferably sodium and potassium, of the alkaline earth metals, preferably calcium, magnesium and barium, of the transition metals, preferably manganese, copper, zinc and iron, and also the ammonium ion which, if desired, may carry one to four C₁-C₄-alkyl substituents and/or one phenyl or benzyl substituent, preferably diisopropylammonium, tetramethylammonium, tetrabutylammonium, trimethylbenzylammonium, furthermore phosphonium ions, sulfonium ions, preferably tri(C₁-C₄-alkyl)sulfonium, and sulfoxonium ions, preferably tri(C₁-C₄-alkyl)sulfoxonium. Anions of useful acid addition salts are primarily chloride, bromide, fluoride, hydrogensulfate, sulfate, dihydrogenphosphate, hydrogenphosphate, phosphate, nitrate, bicarbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate, and the anions of C₁-C₄-alkanoic acids, preferably formate, acetate, propionate and butyrate. They can be formed by reacting a compound of formula I with an acid of the corresponding anion, preferably of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid or nitric acid.

[0028] The compounds of formula I can be present in atropisomers arising from restricted rotation about a single bond of asymmetric groups. They also form part of the subject matter of the present invention.

[0029] Depending on the substitution pattern, the compounds of formula I and their N-oxides may have one or more centers of chirality, in which case they are present as pure enantiomers or pure diastereomers or as enantiomer or diastereomer mixtures. Both, the pure enantiomers or diastereomers and their mixtures are subject matter of the present invention.

[0030] In respect of the variables, the embodiments of the intermediates correspond to the embodiments of the compounds I.

[0031] Preference is given to those compounds I and where applicable also to compounds of all sub-formulae such I.A provided herein and to the intermediates such as compounds XII and XIII, wherein the substituents (such as X¹, X², R¹, R², R^a and R^b) have independently of each other or more preferably in combination the following meanings:

[0032] One embodiment relates to compounds I, wherein X¹ is F or Cl, in particular Cl.

[0033] Another embodiment relates to compounds I, wherein X² is F or Cl, in particular Cl.

[0034] A further embodiment relates to compounds I, wherein R¹ is hydrogen, C₁-C₄-alkyl, allyl, C₂-C₄-alkynyl, cyclopropyl, cyclopropylmethyl, phenyl, benzyl, phenylethenyl or phenylethynyl, more preferably hydrogen or C₁-C₄-alkyl, in particular methyl.

[0035] A further embodiment relates to compounds I, wherein R¹ is hydrogen.

[0036] A further embodiment relates to compounds I, wherein R¹ is C₁-C₄-alkyl, alkyl, C₂-C₄-alkynyl, cyclopropyl, cyclopropylmethyl, phenyl, benzyl, phenylethenyl or phenylethynyl, wherein the aforementioned groups may be substituted by R^a and/or R^b as defined above, more preferably they carry 1, 2 or 3 halogen substituents, even more preferably R¹ is C₁-C₂-haloalkyl, in particular R¹ is CF₃.

[0037] A further embodiment relates to compounds I, wherein R¹ is C₃-C₈-cycloalkyl or C₃-C₈-cycloalkyl-C₁-C₄-alkyl, more preferably selected from cyclopropyl and cyclopropylmethyl, wherein the aforementioned groups may be substituted by R^a and/or R^b as defined herein.

[0038] A further embodiment relates to compounds I, wherein R² is n-butyl, wherein the aforementioned groups may be substituted by R^a as defined above.


[0039] A further embodiment relates to compounds I, wherein R² is sec.-butyl, wherein the aforementioned groups may be substituted by R^a as defined above.

[0040] A further embodiment relates to compounds I, wherein R² is tert.-butyl, wherein said tert.-butyl may be substituted by R^a as defined above.

[0041] A further embodiment relates to compounds I, wherein R² is unsubstituted.


[0042] A further embodiment relates to compounds I, wherein R² carries 1, 2, 3, 4 or 5 groups R^a selected from halogen, CN and nitro, more preferably selected from F and Cl.

[0043] A further embodiment relates to compounds, wherein X¹ and X² are Cl and R² is unsubstituted n-butyl, which compounds are of formula I.A:



40

[0044] A further embodiment relates to compounds, wherein X¹ and X² are Cl and R² is unsubstituted tert.-butyl, which compounds are of formula I.B:



55

[0045] Particularly preferred embodiments of the invention relate to compounds I, wherein the combination of X¹, X² and R² (including R^a) is as defined in Table P below.

Table P:

line	X ¹	X ²	R ²
P-1	Cl	Cl	-CH ₂ CH ₂ CH ₂ CH ₃
P-2	Cl	F	-CH ₂ CH ₂ CH ₂ CH ₃
P-3	F	Cl	-CH ₂ CH ₂ CH ₂ CH ₃
P-4	F	F	-CH ₂ CH ₂ CH ₂ CH ₃
P-5	Cl	Cl	-CH(CH ₃)CH ₂ CH ₃
P-6	Cl	F	-CH(CH ₃)CH ₂ CH ₃
P-7	F	Cl	-CH(CH ₃)CH ₂ CH ₃
P-8	F	F	-CH(CH ₃)CH ₂ CH ₃
P-9	Cl	Cl	-C(CH ₃) ₃
P-10	Cl	F	-C(CH ₃) ₃
P-11	F	Cl	-C(CH ₃) ₃
P-12	F	F	-C(CH ₃) ₃
P-13	Cl	Cl	-CH ₂ CH ₂ CH ₂ CH ₂ Cl
P-14	Cl	F	-CH ₂ CH ₂ CH ₂ CH ₂ Cl
P-15	F	Cl	-CH ₂ CH ₂ CH ₂ CH ₂ Cl
P-16	F	F	-CH ₂ CH ₂ CH ₂ CH ₂ Cl
P-17	Cl	Cl	-CH ₂ CH ₂ CH ₂ CH ₂ F
P-18	Cl	F	-CH ₂ CH ₂ CH ₂ CH ₂ F
P-19	F	Cl	-CH ₂ CH ₂ CH ₂ CH ₂ F
P-20	F	F	-CH ₂ CH ₂ CH ₂ CH ₂ F
P-21	Cl	Cl	-CH ₂ CH ₂ CH ₂ CH ₂ CN
P-22	Cl	F	-CH ₂ CH ₂ CH ₂ CH ₂ CN
P-23	F	Cl	-CH ₂ CH ₂ CH ₂ CH ₂ CN
P-24	F	F	-CH ₂ CH ₂ CH ₂ CH ₂ CN
P-25	Cl	Cl	-CH(CH ₃)CH ₂ CH ₂ Cl

line	X ¹	X ²	R ²
P-26	Cl	F	-CH(CH ₃)CH ₂ CH ₂ Cl
P-27	F	Cl	-CH(CH ₃)CH ₂ CH ₂ Cl
P-28	F	F	-CH(CH ₃)CH ₂ CH ₂ Cl
P-29	Cl	Cl	-CH(CH ₃)CH ₂ CH ₂ F
P-30	Cl	F	-CH(CH ₃)CH ₂ CH ₂ F
P-31	F	Cl	-CH(CH ₃)CH ₂ CH ₂ F
P-32	F	F	-CH(CH ₃)CH ₂ CH ₂ F
P-33	Cl	Cl	-C(CH ₂ Cl) ₃
P-34	Cl	F	-C(CH ₂ Cl) ₃
P-35	F	Cl	-C(CH ₂ Cl) ₃
P-36	F	F	-C(CH ₂ Cl) ₃
P-37	Cl	Cl	-C(CH ₂ F) ₃
P-38	Cl	F	-C(CH ₂ F) ₃
P-39	F	Cl	-C(CH ₂ F) ₃
P-40	F	F	-C(CH ₂ F) ₃
P-41	Cl	Cl	-C(CCl ₃) ₃
P-42	Cl	F	-C(CCl ₃) ₃
P-43	F	Cl	-C(CCl ₃) ₃
P-44	F	F	-C(CCl ₃) ₃
P-45	Cl	Cl	-C(CF ₃) ₃
P-46	Cl	F	-C(CF ₃) ₃
P-47	F	Cl	-C(CF ₃) ₃
P-48	F	F	-C(CF ₃) ₃

5

[0046] A skilled person will readily understand that the preferences given in connection with compounds I apply for formula XII as defined above.

[0047] With respect to their use, particular preference is given to the 1440 compounds of formula I compiled in Table 1 to 48 below. The groups mentioned in the Tables for a substituent are furthermore, independently of the combination wherein they are mentioned, a particularly preferred embodiment of the substituent in question.

Table 1: Compounds 1 to 30 of formula I, wherein X¹, X² and R² are defined as in line P-1 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 2: Compounds 31 to 60 of formula I, wherein X¹, X² and R² are defined as in line P-2 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 3: Compounds 61 to 90 of formula I, wherein X¹, X² and R² are defined as in line P-3 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 4: Compounds 91 to 120 of formula I, wherein X¹, X² and R² are defined as in line P-4 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 5: Compounds 121 to 150 of formula I, wherein X¹, X² and R² are defined as in line P-5 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 6: Compounds 151 to 180 of formula I, wherein X¹, X² and R² are defined as in line P-6 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 36: Compounds 1051 to 1080 of formula I, wherein X¹, X² and R² are defined as in line P-36 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 37: Compounds 1081 to 1110 of formula I, wherein X¹, X² and R² are defined as in line P-37 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

5 Table 38: Compounds 1111 to 1140 of formula I, wherein X¹, X² and R² are defined as in line P-38 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 39: Compounds 1141 to 1170 of formula I, wherein X¹, X² and R² are defined as in line P-39 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

10 Table 40: Compounds 1171 to 1200 of formula I, wherein X¹, X² and R² are defined as in line P-40 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 41: Compounds 1201 to 1230 of formula I, wherein X¹, X² and R² are defined as in line P-41 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

15 Table 42: Compounds 1231 to 1260 of formula I, wherein X¹, X² and R² are defined as in line P-42 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 43: Compounds 1261 to 1290 of formula I, wherein X¹, X² and R² are defined as in line P-43 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

20 Table 44: Compounds 1291 to 1320 of formula I, wherein X¹, X² and R² are defined as in line P-44 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 45: Compounds 1321 to 1350 of formula I, wherein X¹, X² and R² are defined as in line P-45 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

25 Table 46: Compounds 1351 to 1380 of formula I, wherein X¹, X² and R² are defined as in line P-46 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table 47: Compounds 1381 to 1410 of formula I, wherein X¹, X² and R² are defined as in line P-47 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

30 Table 48: Compounds 1411 to 1440 of formula I, wherein X¹, X² and R² are defined as in line P-48 of table P and the meaning of R¹ for each individual compound corresponds in each case to one line of table A.

Table A:

line	R ¹
A-1	H
A-2	CH ₃
A-3	CH ₂ CH ₃
A-4	CH ₂ CH ₂ CH ₃
A-5	CH(CH ₃) ₂
A-6	CH ₂ CH ₂ CH ₂ CH ₃
A-7	C ₃ H ₅
A-8	C ₅ H ₉
A-9	C ₆ H ₁₁
A-10	C ₆ H ₅

line	R ¹
A-11	CH ₂ -C ₆ H ₅
A-12	CH ₂ -C ₃ H ₅
A-13	CF ₃
A-14	CHF ₂
A-15	CH ₂ -CN
A-16	CH ₂ CH ₂ -CN
A-17	C≡CH
A-18	C≡CCH ₃
A-19	CH ₂ C≡CH
A-20	4-F-C ₆ H ₄

line	R ¹
A-21	4-Cl-C ₆ H ₄
A-22	2,4-Cl ₂ -C ₆ H ₃
A-23	2,4,6-Cl ₃ -C ₆ H ₂
A-24	2,4,6-F ₃ -C ₆ H ₂
A-25	CH ₂ -C ₆ H ₅
A-26	CH ₂ -(4-F-C ₆ H ₄)
A-27	CH ₂ -(4-Cl-C ₆ H ₄)
A-28	CH=CH-C ₆ H ₅
A-29	CH=CH-(4-F-C ₆ H ₄)
A-30	CH=CH-(4-Cl-C ₆ H ₄)

50 [0048] The compounds I and the compositions according to the invention, respectively, are suitable as fungicides. They are distinguished by an outstanding effectiveness against a broad spectrum of phytopathogenic fungi, including soil-borne fungi, which derive especially from the classes of the Plasmodiophoromycetes, Peronosporomycetes (syn. Oomycetes), Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes (syn. Fungi imperfecti). Some are systemically effective and they can be used in crop protection as foliar fungicides, fungicides for seed dressing and soil fungicides. Moreover, they are suitable for controlling harmful fungi, which inter alia occur in wood or roots of plants.

55 [0049] The compounds I and the compositions according to the invention are particularly important in the control of a multitude of phytopathogenic fungi on various cultivated plants, such as cereals, e. g. wheat, rye, barley, triticale, oats or rice; beet, e. g. sugar beet or fodder beet; fruits, such as pomes, stone fruits or soft fruits, e. g. apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries, blackberries or gooseberries; leguminous plants, such as lentils, peas, alfalfa or soybeans; oil plants, such as rape, mustard, olives, sunflowers, coconut, cocoa beans, castor oil plants,

oil palms, ground nuts or soybeans; cucurbits, such as squashes, cucumber or melons; fiber plants, such as cotton, flax, hemp or jute; citrus fruit, such as oranges, lemons, grape-fruits or mandarins; vegetables, such as spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, cucurbits or paprika; lauraceous plants, such as avocados, cinnamon or camphor; energy and raw material plants, such as corn, soybean, rape, sugar cane or oil palm; corn; 5 tobacco; nuts; coffee; tea; bananas; vines (table grapes and grape juice grape vines); hop; turf; sweet leaf (also called Stevia); natural rubber plants or ornamental and forestry plants, such as flowers, shrubs, broad-leaved trees or evergreens, e. g. conifers; and on the plant propagation material, such as seeds, and the crop material of these plants.

[0050] Preferably, compounds I and compositions thereof, respectively are used for controlling a multitude of fungi on field crops, such as potatoes sugar beets, tobacco, wheat, rye, barley, oats, rice, corn, cotton, soybeans, rape, legumes, 10 sunflowers, coffee or sugar cane; fruits; vines; ornamentals; or vegetables, such as cucumbers, tomatoes, beans or squashes.

[0051] The term "plant propagation material" is to be understood to denote all the generative parts of the plant such as seeds and vegetative plant material such as cuttings and tubers (e. g. potatoes), which can be used for the multiplication of the plant. This includes seeds, roots, fruits, tubers, bulbs, rhizomes, shoots, sprouts and other parts of plants, including 15 seedlings and young plants, which are to be transplanted after germination or after emergence from soil. These young plants may also be protected before transplantation by a total or partial treatment by immersion or pouring.

[0052] Preferably, treatment of plant propagation materials with compounds I and compositions thereof, respectively, is used for controlling a multitude of fungi on cereals, such as wheat, rye, barley and oats; rice, corn, cotton and soybeans.

[0053] The term "cultivated plants" is to be understood as including plants which have been modified by breeding, 20 mutagenesis or genetic engineering including but not limiting to agricultural biotech products on the market or in development (cf. http://www.bio.org/speeches/pubs/er/agri_products.asp). Genetically modified plants are plants, which genetic material has been so modified by the use of recombinant DNA techniques that under natural circumstances cannot readily be obtained by cross breeding, mutations or natural recombination. Typically, one or more genes have been integrated into the genetic material of a genetically modified plant in order to improve certain properties of the plant. 25 Such genetic modifications also include but are not limited to targeted post-translational modification of protein(s), oligo- or polypeptides e. g. by glycosylation or polymer additions such as prenylated, acetylated or farnesylated moieties or PEG moieties.

[0054] Plants that have been modified by breeding, mutagenesis or genetic engineering, e. g. have been rendered 30 tolerant to applications of specific classes of herbicides, such as auxin herbicides such as dicamba or 2,4-D; bleacher herbicides such as hydroxylphenylpyruvate dioxygenase (HPPD) inhibitors or phytoene desaturase (PDS) inhibitors; acetolactate synthase (ALS) inhibitors such as sulfonyl ureas or imidazolinones; enolpyruvylshikimate-3-phosphate synthase (EPSPS) inhibitors, such as glyphosate; glutamine synthetase (GS) inhibitors such as glufosinate; protoporphyrinogen-IX oxidase inhibitors; lipid biosynthesis inhibitors such as acetyl CoA carboxylase (ACCase) inhibitors; or 35 oxynil (i. e. bromoxynil or ioxynil) herbicides as a result of conventional methods of breeding or genetic engineering. Furthermore, plants have been made resistant to multiple classes of herbicides through multiple genetic modifications, such as resistance to both glyphosate and glufosinate or to both glyphosate and a herbicide from another class such as ALS inhibitors, HPPD inhibitors, auxin herbicides, or ACCase inhibitors. These herbicide resistance technologies are 40 e. g. described in Pest Managem. Sci. 61, 2005, 246; 61, 2005, 258; 61, 2005, 277; 61, 2005, 269; 61, 2005, 286; 64, 2008, 326; 64, 2008, 332; Weed Sci. 57, 2009, 108; Austral. J. Agricult. Res. 58, 2007, 708; Science 316, 2007, 1185; and references quoted therein. Several cultivated plants have been rendered tolerant to herbicides by conventional 45 methods of breeding (mutagenesis), e. g. Clearfield® summer rape (Canola, BASF SE, Germany) being tolerant to imidazolinones, e. g. imazamox, or ExpressSun® sunflowers (DuPont, USA) being tolerant to sulfonyl ureas, e. g. tribenuron. Genetic engineering methods have been used to render cultivated plants such as soybean, cotton, corn, beets and rape, tolerant to herbicides such as glyphosate and glufosinate, some of which are commercially available under the trade names RoundupReady® (glyphosate-tolerant, Monsanto, U.S.A.), Cultivance® (imidazolinone tolerant, 50 BASF SE, Germany) and LibertyLink® (glufosinate-tolerant, Bayer CropScience, Germany).

[0055] Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more insecticidal proteins, especially those known from the bacterial genus *Bacillus*, particularly from *Bacillus thuringiensis*, such as 6-endotoxins, e. g. CryIA(b), CryIA(c), CryIF, CryIF(a2), CryIIA(b), CryIIIA, CryIIIB(b1) or Cry9c; 55 vegetative insecticidal proteins (VIP), e. g. VIP1, VIP2, VIP3 or VIP3A; insecticidal proteins of bacteria colonizing nematodes, e. g. *Photobacterium* spp. or *Xenorhabdus* spp.; toxins produced by animals, such as scorpion toxins, arachnid toxins, wasp toxins, or other insect-specific neurotoxins; toxins produced by fungi, such as Streptomyces toxins, plant lectins, such as pea or barley lectins; agglutinins; proteinase inhibitors, such as trypsin inhibitors, serine protease inhibitors, patatin, cystatin or papain inhibitors; ribosome-inactivating proteins (RIP), such as ricin, maize-RIP, abrin, luffin, saporin or bryodin; steroid metabolism enzymes, such as 3-hydroxysteroid oxidase, ecdysteroid-IDP-glycosyl-transferase, cholesterol oxidases, ecdysone inhibitors or HMG-CoA-reductase; ion channel blockers, such as blockers of sodium or calcium channels; juvenile hormone esterase; diuretic hormone receptors (helicokinin receptors); stilben synthase, bibenzyl synthase, chitinases or glucanases. In the context of the present invention these insecticidal proteins

or toxins are to be understood expressly also as pre-toxins, hybrid proteins, truncated or otherwise modified proteins. Hybrid proteins are characterized by a new combination of protein domains, (see, e. g. WO 02/015701). Further examples of such toxins or genetically modified plants capable of synthesizing such toxins are disclosed, e. g., in EP-A 374 753, WO 93/007278, WO 95/34656, EP-A 427 529, EP-A 451 878, WO 03/18810 und WO 03/52073. The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e. g. in the publications mentioned above. These insecticidal proteins contained in the genetically modified plants impart to the plants producing these proteins tolerance to harmful pests from all taxonomic groups of arthropods, especially to beetles (Coleoptera), two-winged insects (Diptera), and moths (Lepidoptera) and to nematodes (Nematoda). Genetically modified plants capable of synthesizing one or more insecticidal proteins are, e. g., described in the publications mentioned above, and some of which are commercially available such as YieldGard® (corn cultivars producing the Cry1Ab toxin), YieldGard® Plus (corn cultivars producing Cry1Ab and Cry3Bb1 toxins), Starlink® (corn cultivars producing the Cry9c toxin), Herculex® RW (corn cultivars producing Cry34Ab1, Cry35Ab1 and the enzyme Phosphinothrinic-N-Acetyltransferase [PAT]); NuCOTN® 33B (cotton cultivars producing the Cry1Ac toxin), Bollgard® I (cotton cultivars producing the Cry1Ac toxin), Bollgard® II (cotton cultivars producing Cry1Ac and Cry2Ab2 toxins); VIPCOT® (cotton cultivars producing a VIP-toxin); NewLeaf® (potato cultivars producing the Cry3A toxin); Bt-Xtra®, NatureGard®, KnockOut®, BiteGard®, Protecta®, Bt11 (e. g. Agrisure® CB) and Bt176 from Syngenta Seeds SAS, France, (corn cultivars producing the Cry1Ab toxin and PAT enzyme), MIR604 from Syngenta Seeds SAS, France (corn cultivars producing a modified version of the Cry3A toxin, c.f. WO 03/018810), MON 863 from Monsanto Europe S.A., Belgium (corn cultivars producing the Cry3Bb1 toxin), IPC 531 from Monsanto Europe S.A., Belgium (cotton cultivars producing a modified version of the Cry1Ac toxin) and 1507 from Pioneer Overseas Corporation, Belgium (corn cultivars producing the Cry1F toxin and PAT enzyme).

[0056] Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the resistance or tolerance of those plants to bacterial, viral or fungal pathogens. Examples of such proteins are the so-called "pathogenesis-related proteins" (PR proteins, see, e. g. EP-A 392 225), plant disease resistance genes (e. g. potato cultivars, which express resistance genes acting against *Phytophthora infestans* derived from the mexican wild potato *Solanum bulbocastanum*) or T4-lysozym (e. g. potato cultivars capable of synthesizing these proteins with increased resistance against bacteria such as *Erwinia amylovora*). The methods for producing such genetically modified plants are generally known to the person skilled in the art and are described, e. g. in the publications mentioned above.

[0057] Furthermore, plants are also covered that are by the use of recombinant DNA techniques capable to synthesize one or more proteins to increase the productivity (e. g. bio mass production, grain yield, starch content, oil content or protein content), tolerance to drought, salinity or other growth-limiting environmental factors or tolerance to pests and fungal, bacterial or viral pathogens of those plants.

[0058] Furthermore, plants are also covered that contain by the use of recombinant DNA techniques a modified amount of substances of content or new substances of content, specifically to improve human or animal nutrition, e. g. oil crops that produce health-promoting long-chain omega-3 fatty acids or unsaturated omega-9 fatty acids (e. g. Nexera® rape, DOW Agro Sciences, Canada).

[0059] Furthermore, plants are also covered that contain by the use of recombinant DNA techniques a modified amount of substances of content or new substances of content, specifically to improve raw material production, e. g. potatoes that produce increased amounts of amylopectin (e. g. Amflora® potato, BASF SE, Germany).

[0060] The compounds I and compositions thereof, respectively, are particularly suitable for controlling the following plant diseases:

Albugo spp. (white rust) on ornamentals, vegetables (e. g. *A. candida*) and sunflowers (e. g. *A. tragopogonis*); *Alternaria* spp. (Alternaria leaf spot) on vegetables, rape (*A. brassicola* or *brassicae*), sugar beets (*A. tenuis*), fruits, rice, soybeans, potatoes (e. g. *A. solani* or *A. alternata*), tomatoes (e. g. *A. solani* or *A. alternata*) and wheat; *Aphanomyces* spp. on sugar beets and vegetables; *Ascochyta* spp. on cereals and vegetables, e. g. *A. tritici* (anthracnose) on wheat and *A. hordei* on barley; *Bipolaris* and *Drechslera* spp. (teleomorph: *Cochliobolus* spp.), e. g. Southern leaf blight (*D. maydis*) or Northern leaf blight (*B. zeicola*) on corn, e. g. spot blotch (*B. sorokiniana*) on cereals and e.g. *B. oryzae* on rice and turfs; *Blumeria* (formerly *Erysiphe*) *graminis* (powdery mildew) on cereals (e. g. on wheat or barley); *Botrytis* *cinerea* (teleomorph: *Botryotinia fuckeliana*: grey mold) on fruits and berries (e. g. strawberries), vegetables (e. g. lettuce, carrots, celery and cabbages), rape, flowers, vines, forestry plants and wheat; *Bremia lactucae* (downy mildew) on lettuce; *Ceratocystis* (syn. *Ophiostoma*) spp. (rot or wilt) on broad-leaved trees and evergreens, e. g. *C. ulmi* (Dutch elm disease) on elms; *Cercospora* spp. (Cercospora leaf spots) on corn (e.g. Gray leaf spot: *C. zeae-maydis*), rice, sugar beets (e. g. *C. beticola*), sugar cane, vegetables, coffee, soybeans (e. g. *C. sojina* or *C. kikuchii*) and rice; *Cladosporium* spp. on tomatoes (e. g. *C. fulvum*: leaf mold) and cereals, e. g. *C. herbarum* (black ear) on wheat; *Claviceps purpurea* (ergot) on cereals; *Cochliobolus* (anamorph: *Helminthosporium* of *Bipolaris*) spp. (leaf spots) on corn (*C. carbonum*), cereals (e. g. *C. sativus*, anamorph: *B. sorokiniana*) and rice (e. g. *C. miyabeanus*, anamorph: *H. oryzae*); *Colletotrichum* (teleomorph: *Glomerella*) spp. (anthracnose)

on cotton (e. g. *C. gossypii*), corn (e. g. *C. graminicola*: Anthracnose stalk rot), soft fruits, potatoes (e. g. *C. coccodes*: black dot), beans (e. g. *C. lindemuthianum*) and soybeans (e. g. *C. truncatum* or *C. gloeosporioides*); *Corticium* spp., e. g. *C. sasakii* (sheath blight) on rice; *Corynespora cassiicola* (leaf spots) on soybeans and ornamentals; *Cycloconium* spp., e. g. *C. oleaginum* on olive trees; *Cylindrocarpon* spp. (e. g. fruit tree canker or young vine decline, teleomorph: *Nectria* or *Neonectria* spp.) on fruit trees, vines (e. g. *C. liriodendri*, teleomorph: *Neonectria liriodendri*: Black Foot Disease) and ornamentals; *Dematophora* (teleomorph: *Rosellinia*) necatrix (root and stem rot) on soybeans; *Diaporthe* spp., e. g. *D. phaseolorum* (damping off) on soybeans; *Drechslera* (syn. *Helminthosporium*, teleomorph: *Pyrenophora*) spp. on corn, cereals, such as barley (e. g. *D. teres*, net blotch) and wheat (e. g. *D. tritici-repentis*: tan spot), rice and turf; *Esca* (dieback, apoplexy) on vines, caused by *Formitiporia* (syn. *Phellinus*) *punctata*, *F. mediterranea*, *Phaeomoniella chlamydospora* (earlier *Phaeoacremonium chlamydosporum*), *Phaeoacremonium aleophilum* and/or *Botryosphaeria obtusa*; *Elsinoe* spp. on pome fruits (*E. pyri*), soft fruits (*E. veneta*: anthracnose) and vines (*E. ampelina*: anthracnose); *Entyloma oryzae* (leaf smut) on rice; *Epicoccum* spp. (black mold) on wheat; *Erysiphe* spp. (powdery mildew) on sugar beets (*E. betae*), vegetables (e. g. *E. pisii*), such as cucurbits (e. g. *E. cichoracearum*), cabbages, rape (e. g. *E. cruciferarum*); *Eutypa lata* (Eutypa canker or dieback, anamorph: *Cytosporina lata*, syn. *Libertella blepharis*) on fruit trees, vines and ornamental woods; *Exserohilum* (syn. *Helminthosporium*) spp. on corn (e. g. *E. turicum*); *Fusarium* (teleomorph: *Gibberella*) spp. (wilt, root or stem rot) on various plants, such as *F. graminearum* or *F. culmorum* (root rot, scab or head blight) on cereals (e. g. wheat or barley), *F. oxysporum* on tomatoes, *F. solani* on soybeans and *F. verticillioides* on corn; *Gaeumannomyces graminis* (take-all) on cereals (e. g. wheat or barley) and corn; *Gibberella* spp. on cereals (e. g. *G. zeae*) and rice (e. g. *G. fujikuroi*: Bakanae disease); *Glomerella cingulata* on vines, pome fruits and other plants and *G. gossypii* on cotton; Grain-staining complex on rice; *Guignardia bidwellii* (black rot) on vines; *Gymnosporangium* spp. on rosaceous plants and junipers, e. g. *G. sabinae* (rust) on pears; *Helminthosporium* spp. (syn. *Drechslera*, teleomorph: *Cochliobolus*) on corn, cereals and rice; *Hemileia* spp., e. g. *H. vastatrix* (coffee leaf rust) on coffee; *Isariopsis clavigrata* (syn. *Cladosporium vitis*) on vines; *Macrophomina phaseolina* (syn. *phaseoli*) (root and stem rot) on soybeans and cotton; *Microdochium* (syn. *Fusarium*) *nivale* (pink snow mold) on cereals (e. g. wheat or barley); *Microsphaera diffusa* (powdery mildew) on soybeans; *Monilinia* spp., e. g. *M. laxa*, *M. fructicola* and *M. fructigena* (bloom and twig blight, brown rot) on stone fruits and other rosaceous plants; *Mycosphaerella* spp. on cereals, bananas, soft fruits and ground nuts, such as e. g. *M. graminicola* (anamorph: *Septoria tritici*, Septoria blotch) on wheat or *M. fijiensis* (black Sigatoka disease) on bananas; *Peronospora* spp. (downy mildew) on cabbage (e. g. *P. brassicae*), rape (e. g. *P. parasitica*), onions (e. g. *P. destructor*), tobacco (*P. tabacina*) and soybeans (e. g. *P. manshurica*); *Phakopsora pachyrhizi* and *P. meibomiae* (soybean rust) on soybeans; *Phialophora* spp. e. g. on vines (e. g. *P. tracheiphila* and *P. tetraspora*) and soybeans (e. g. *P. gregata*: stem rot); *Phoma lingam* (root and stem rot) on rape and cabbage and *P. betae* (root rot, leaf spot and damping-off) on sugar beets; *Phomopsis* spp. on sunflowers, vines (e. g. *P. viticola*: can and leaf spot) and soybeans (e. g. stem rot: *P. phaseoli*, teleomorph: *Diaporthe phaseolorum*); *Physoderma maydis* (brown spots) on corn; *Phytophthora* spp. (wilt, root, leaf, fruit and stem root) on various plants, such as paprika and cucurbits (e. g. *P. capsici*), soybeans (e. g. *P. megasperma*, syn. *P. sojae*), potatoes and tomatoes (e. g. *P. infestans*: late blight) and broad-leaved trees (e. g. *P. ramorum*: sudden oak death); *Plasmiodiophora brassicae* (club root) on cabbage, rape, radish and other plants; *Plasmopara* spp., e. g. *P. viticola* (grapevine downy mildew) on vines and *P. halstedii* on sunflowers; *Podosphaera* spp. (powdery mildew) on rosaceous plants, hop, pome and soft fruits, e. g. *P. leucotricha* on apples; *Polymyxa* spp., e. g. on cereals, such as barley and wheat (*P. graminis*) and sugar beets (*P. betae*) and thereby transmitted viral diseases; *Pseudocercospora herpotrichoides* (eyespot, teleomorph: *Tapesia yallundae*) on cereals, e. g. wheat or barley; *Pseudoperonospora* (downy mildew) on various plants, e. g. *P. cubensis* on cucurbits or *P. humili* on hop; *Pseudopezicula tracheiphila* (red fire disease or 'rotbrenner', anamorph: *Phialophora*) on vines; *Puccinia* spp. (rusts) on various plants, e. g. *P. triticina* (brown or leaf rust), *P. striiformis* (stripe or yellow rust), *P. hordei* (dwarf rust), *P. graminis* (stem or black rust) or *P. recondita* (brown or leaf rust) on cereals, such as e. g. wheat, barley or rye, *P. kuehnii* (orange rust) on sugar cane and *P. asparagi* on asparagus; *Pyrenophora* (anamorph: *Drechslera*) *tritici-repentis* (tan spot) on wheat or *P. teres* (net blotch) on barley; *Pyricularia* spp., e. g. *P. oryzae* (teleomorph: *Magnaporthe grisea*, rice blast) on rice and *P. grisea* on turf and cereals; *Pythium* spp. (damping-off on turf, rice, corn, wheat, cotton, rape, sunflowers, soybeans, sugar beets, vegetables and various other plants (e. g. *P. ultimum* or *P. aphanidermatum*); *Ramularia* spp., e. g. *R. collo-cygni* (Ramularia leaf spots, Physiological leaf spots) on barley and *R. beticola* on sugar beets; *Rhizoctonia* spp. on cotton, rice, potatoes, turf, corn, rape, potatoes, sugar beets, vegetables and various other plants, e. g. *R. solani* (root and stem rot) on soybeans, *R. solani* (sheath blight) on rice or *R. cerealis* (Rhizoctonia spring blight) on wheat or barley; *Rhizopus stolonifer* (black mold, soft rot) on strawberries, carrots, cabbage, vines and tomatoes; *Rhynchosporium secalis* (scald) on barley, rye and triticale; *Sclerotinia* spp. (stem rot or white mold) on vegetables and field crops, such as rape, sunflowers (e. g. *S. sclerotiorum*) and soybeans (e. g. *S. rolfsii* or *S. sclerotiorum*); *Septoria* spp. on various plants, e. g. *S. glycines* (brown spot) on soybeans, *S. tritici* (Septoria blotch) on wheat and *S. (syn. Stagonospora) nodorum* (Stagonospora blotch) on cereals;

5 *Uncinula* (syn. *Erysiphe*) *necator* (powdery mildew, anamorph: *Oidium tucken*) on vines; *Setosphaeria* spp. (leaf blight) on corn (e. g. *S. turicum*, syn. *Helminthosporium turicum*) and turf; *Sphacelotheca* spp. (smut) on corn, (e. g. *S. reiliana*: head smut), sorghum und sugar cane; *Sphaerotheca fuliginea* (powdery mildew) on cucurbits; *Spongospora subterranea* (powdery scab) on potatoes and thereby transmitted viral diseases; *Stagonospora* spp. on cereals, e. g. *S. nodorum* (Stagonospora blotch, teleomorph: *Leptosphaeria* [syn. *Phaeosphaeria*] *nodorum*) on wheat; *Synchytrium endobioticum* on potatoes (potato wart disease); *Taphrina* spp., e. g. *T. deformans* (leaf curl disease) on peaches and *T. pruni* (plum pocket) on plums; *Thielaviopsis* spp. (black root rot) on tobacco, pome fruits, vegetables, soybeans and cotton, e. g. *T. basicola* (syn. *Chalara elegans*); *Tilletia* spp. (common bunt or stinking smut) on cereals, such as e. g. *T. tritici* (syn. *T. caries*, wheat bunt) and *T. controversa* (dwarf bunt) on wheat; *Typhula incarnata* (grey snow mold) on barley or wheat; *Urocystis* spp., e. g. *U. occulta* (stem smut) on rye; *Uromyces* spp. (rust) on vegetables, such as beans (e. g. *U. appendiculatus*, syn. *U. phaseoli*) and sugar beets (e. g. *U. betae*); *Ustilago* spp. (loose smut) on cereals (e. g. *U. nuda* and *U. avaenae*), corn (e. g. *U. maydis*: corn smut) and sugar cane; *Venturia* spp. (scab) on apples (e. g. *V. inaequalis*) and pears; and *Verticillium* spp. (wilt) on various plants, such as fruits and ornamentals, vines, soft fruits, vegetables and field crops, e. g. *V. dahliae* on strawberries, rape, potatoes and tomatoes.

10 [0061] The compounds I and compositions thereof, respectively, are also suitable for controlling harmful fungi in the protection of stored products or harvest and in the protection of materials. The term "protection of materials" is to be understood to denote the protection of technical and non-living materials, such as adhesives, glues, wood, paper and paperboard, textiles, leather, paint dispersions, plastics, colling lubricants, fiber or fabrics, against the infestation and destruction by harmful microorganisms, such as fungi and bacteria. As to the protection of wood and other materials, the particular attention is paid to the following harmful fungi: Ascomycetes such as *Ophiostoma* spp., *Ceratocystis* spp., *Aureobasidium pullulans*, *Sclerophoma* spp., *Chaetomium* spp., *Humicola* spp., *Petriella* spp., *Trichurus* spp.; Basidiomycetes such as *Coniophora* spp., *Coriolus* spp., *Gloeophyllum* spp., *Lentinus* spp., *Pleurotus* spp., *Poria* spp., *Serpula* spp. and *Tyromyces* spp., Deuteromycetes such as *Aspergillus* spp., *Cladosporium* spp., *Penicillium* spp., *Trichorma* spp., *Alternaria* spp., *Paecilomyces* spp. and Zygomycetes such as *Mucor* spp., and in addition in the protection of stored products and harvest the following yeast fungi are worthy of note: *Candida* spp. and *Saccharomyces cerevisiae*.

15 [0062] The compounds I and compositions thereof, respectively, may be used for improving the health of a plant. The invention also relates to a method for improving plant health by treating a plant, its propagation material and/or the locus where the plant is growing or is to grow with an effective amount of compounds I and compositions thereof, respectively.

20 [0063] The term "plant health" is to be understood to denote a condition of the plant and/or its products which is determined by several indicators alone or in combination with each other such as yield (e. g. increased biomass and/or increased content of valuable ingredients), plant vigor (e. g. improved plant growth and/or greener leaves ("greening effect")), quality (e. g. improved content or composition of certain ingredients) and tolerance to abiotic and/or biotic stress. The above identified indicators for the health condition of a plant may be interdependent or may result from each other.

25 [0064] The compounds of formula I can be present in different crystal modifications whose biological activity may differ. They are likewise subject matter of the present invention.

30 [0065] The compounds I are employed as such or in form of compositions by treating the fungi or the plants, plant propagation materials, such as seeds, soil, surfaces, materials or rooms to be protected from fungal attack with a fungicidally effective amount of the active substances. The application can be carried out both before and after the infection of the plants, plant propagation materials, such as seeds, soil, surfaces, materials or rooms by the fungi.

35 [0066] Plant propagation materials may be treated with compounds I as such or a composition comprising at least one compound I prophylactically either at or before planting or transplanting.

40 [0067] The invention also relates to agrochemical compositions comprising an auxiliary and at least one compound I according to the invention.

45 [0068] An agrochemical composition comprises a fungicidally effective amount of a compound I. The term "effective amount" denotes an amount of the composition or of the compounds I, which is sufficient for controlling harmful fungi on cultivated plants or in the protection of materials and which does not result in a substantial damage to the treated plants. Such an amount can vary in a broad range and is dependent on various factors, such as the fungal species to be controlled, the treated cultivated plant or material, the climatic conditions and the specific compound I used.

50 [0069] The compounds I, their N-oxides and salts can be converted into customary types of agrochemical compositions, e. g. solutions, emulsions, suspensions, dusts, powders, pastes, granules, pressings, capsules, and mixtures thereof. Examples for composition types are suspensions (e.g. SC, OD, FS), emulsifiable concentrates (e.g. EC), emulsions (e.g. EW, EO, ES, ME), capsules (e.g. CS, ZC), pastes, pastilles, wettable powders or dusts (e.g. WP, SP, WS, DP, DS), pressings (e.g. BR, TB, DT), granules (e.g. WG, SG, GR, FG, GG, MG), insecticidal articles (e.g. LN), as well as gel formulations for the treatment of plant propagation materials such as seeds (e.g. GF). These and further compositions

types are defined in the "Catalogue of pesticide formulation types and international coding system", Technical Monograph No. 2, 6th Ed. May 2008, Crop-Life International.

[0070] The compositions are prepared in a known manner, such as described by Mollet and Grubemann, Formulation technology, Wiley VCH, Weinheim, 2001; or Knowles, New developments in crop protection product formulation, Agrow Reports DS243, T&F Informa, London, 2005.

[0071] Suitable auxiliaries are solvents, liquid carriers, solid carriers or fillers, surfactants, dispersants, emulsifiers, wetters, adjuvants, solubilizers, penetration enhancers, protective colloids, adhesion agents, thickeners, humectants, repellents, attractants, feeding stimulants, compatibilizers, bactericides, anti-freezing agents, anti-foaming agents, colorants, tackifiers and binders.

[0072] Suitable solvents and liquid carriers are water and organic solvents, such as mineral oil fractions of medium to high boiling point, e.g. kerosene, diesel oil; oils of vegetable or animal origin; aliphatic, cyclic and aromatic hydrocarbons, e.g. toluene, paraffin, tetrahydronaphthalene, alkylated naphthalenes; alcohols, e.g. ethanol, propanol, butanol, benzylalcohol, cyclohexanol; glycols; DMSO; ketones, e.g. cyclohexanone; esters, e.g. lactates, carbonates, fatty acid esters, gamma-butyrolactone; fatty acids; phosphonates; amines; amides, e.g. N-methylpyrrolidone, fatty acid dimethylamides; and mixtures thereof.

[0073] Suitable solid carriers or fillers are mineral earths, e.g. silicates, silica gels, talc, kaolins, limestone, lime, chalk, clays, dolomite, diatomaceous earth, bentonite, calcium sulfate, magnesium sulfate, magnesium oxide; polysaccharides, e.g. cellulose, starch; fertilizers, e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas; products of vegetable origin, e.g. cereal meal, tree bark meal, wood meal, nutshell meal, and mixtures thereof.

[0074] Suitable surfactants are surface-active compounds, such as anionic, cationic, nonionic and amphoteric surfactants, block polymers, polyelectrolytes, and mixtures thereof. Such surfactants can be used as emulsifier, dispersant, solubilizer, wetter, penetration enhancer, protective colloid, or adjuvant. Examples of surfactants are listed in McCutcheon's, Vol.1: Emulsifiers & Detergents, McCutcheon's Directories, Glen Rock, USA, 2008 (International Ed. or North American Ed.).

[0075] Suitable anionic surfactants are alkali, alkaline earth or ammonium salts of sulfonates, sulfates, phosphates, carboxylates, and mixtures thereof. Examples of sulfonates are alkylarylsulfonates, diphenylsulfonates, alpha-olefin sulfonates, lignine sulfonates, sulfonates of fatty acids and oils, sulfonates of ethoxylated alkylphenols, sulfonates of alkoxylated arylphenols, sulfonates of condensed naphthalenes, sulfonates of dodecyl- and tridecylbenzenes, sulfonates of naphthalenes and alkylnaphthalenes, sulfosuccinates or sulfosuccinamates. Examples of sulfates are sulfates of fatty acids and oils, of ethoxylated alkylphenols, of alcohols, of ethoxylated alcohols, or of fatty acid esters. Examples of phosphates are phosphate esters. Examples of carboxylates are alkyl carboxylates, and carboxylated alcohol or alkylphenol ethoxylates.

[0076] Suitable nonionic surfactants are alkoxylates, N-substituted fatty acid amides, amine oxides, esters, sugar-based surfactants, polymeric surfactants, and mixtures thereof. Examples of alkoxylates are compounds such as alcohols, alkylphenols, amines, amides, arylphenols, fatty acids or fatty acid esters which have been alkoxylated with 1 to 50 equivalents. Ethylene oxide and/or propylene oxide may be employed for the alkoxylation, preferably ethylene oxide. Examples of N-substituted fatty acid amides are fatty acid glucamides or fatty acid alkanolamides. Examples of esters are fatty acid esters, glycerol esters or monoglycerides. Examples of sugar-based surfactants are sorbitans, ethoxylated sorbitans, sucrose and glucose esters or alkylpolyglucosides. Examples of polymeric surfactants are home- or copolymers of vinylpyrrolidone, vinylalcohols, or vinylacetate.

[0077] Suitable cationic surfactants are quaternary surfactants, for example quaternary ammonium compounds with one or two hydrophobic groups, or salts of long-chain primary amines. Suitable amphoteric surfactants are alkylbетaines and imidazolines. Suitable block polymers are block polymers of the A-B or A-B-A type comprising blocks of polyethylene oxide and polypropylene oxide, or of the A-B-C type comprising alkanol, polyethylene oxide and polypropylene oxide.

[0078] Suitable polyelectrolytes are polyacids or polybases. Examples of polyacids are alkali salts of polyacrylic acid or polyacid comb polymers. Examples of polybases are polyvinylamines or polyethyleneamines.

[0079] Suitable adjuvants are compounds, which have a neglectable or even no pesticidal activity themselves, and which improve the biological performance of the compound I on the target. Examples are surfactants, mineral or vegetable oils, and other auxiliaries. Further examples are listed by Knowles, Adjuvants and additives, Agrow Reports DS256, T&F Informa UK, 2006, chapter 5.

[0080] Suitable thickeners are polysaccharides (e.g. xanthan gum, carboxymethylcellulose), anorganic clays (organically modified or unmodified), polycarboxylates, and silicates.

[0081] Suitable bactericides are bronopol and isothiazolinone derivatives such as alkylisothiazolinones and benzisothiazolinones.

[0082] Suitable anti-freezing agents are ethylene glycol, propylene glycol, urea and glycerin.

[0083] Suitable anti-foaming agents are silicones, long chain alcohols, and salts of fatty acids.

[0084] Suitable colorants (e.g. in red, blue, or green) are pigments of low water solubility and water-soluble dyes. Examples are inorganic colorants (e.g. iron oxide, titan oxide, iron hexacyanoferrate) and organic colorants (e.g. alizarin-,

azo- and phthalocyanine colorants).

[0084] Suitable tackifiers or binders are polyvinylpyrrolidons, polyvinylacetates, polyvinyl alcohols, polyacrylates, biological or synthetic waxes, and cellulose ethers.

[0085] Examples for composition types and their preparation are:

5 i) Water-soluble concentrates (SL, LS)
10-60 wt% of a compound I and 5-15 wt% wetting agent (e.g. alcohol alkoxylates) are dissolved in water and/or in a water-soluble solvent (e.g. alcohols) ad 100 wt%. The active substance dissolves upon dilution with water.

10 ii) Dispersible concentrates (DC)
5-25 wt% of a compound I and 1-10 wt% dispersant (e.g. polyvinylpyrrolidone) are dissolved in organic solvent (e.g. cyclohexanone) ad 100 wt%. Dilution with water gives a dispersion.

15 iii) Emulsifiable concentrates (EC)
15-70 wt% of a compound I and 5-10 wt% emulsifiers (e.g. calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in water-insoluble organic solvent (e.g. aromatic hydrocarbon) ad 100 wt%. Dilution with water gives an emulsion.

20 iv) Emulsions (EW, EO, ES)
5-40 wt% of a compound I and 1-10 wt% emulsifiers (e.g. calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in 20-40 wt% water-insoluble organic solvent (e.g. aromatic hydrocarbon). This mixture is introduced into water ad 100 wt% by means of an emulsifying machine and made into a homogeneous emulsion. Dilution with water gives an emulsion.

25 v) Suspensions (SC, OD, FS)
In an agitated ball mill, 20-60 wt% of a compound I are comminuted with addition of 2-10 wt% dispersants and wetting agents (e.g. sodium lignosulfonate and alcohol ethoxylate), 0.1-2 wt% thickener (e.g. xanthan gum) and water ad 100 wt% to give a fine active substance suspension. Dilution with water gives a stable suspension of the active substance. For FS type composition up to 40 wt% binder (e.g. polyvinylalcohol) is added.

30 vi) Water-dispersible granules and water-soluble granules (WG, SG)
50-80 wt% of a compound I are ground finely with addition of dispersants and wetting agents (e.g. sodium lignosulfonate and alcohol ethoxylate) ad 100 wt% and prepared as water-dispersible or water-soluble granules by means of technical appliances (e.g. extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active substance.

35 vii) Water-dispersible powders and water-soluble powders (WP, SP, WS)
50-80 wt% of a compound I are ground in a rotor-stator mill with addition of 1-5 wt% dispersants (e.g. sodium lignosulfonate), 1-3 wt% wetting agents (e.g. alcohol ethoxylate) and solid carrier (e.g. silica gel) ad 100 wt%. Dilution with water gives a stable dispersion or solution of the active substance.

40 viii) Gel (GW, GF)
In an agitated ball mill, 5-25 wt% of a compound I are comminuted with addition of 3-10 wt% dispersants (e.g. sodium lignosulfonate), 1-5 wt% thickener (e.g. carboxymethylcellulose) and water ad 100 wt% to give a fine suspension of the active substance. Dilution with water gives a stable suspension of the active substance.

45 ix) Microemulsion (ME)
5-20 wt% of a compound I are added to 5-30 wt% organic solvent blend (e.g. fatty acid dimethylamide and cyclohexanone), 10-25 wt% surfactant blend (e.g. alcohol ethoxylate and arylphenol ethoxylate), and water ad 100 %. This mixture is stirred for 1 h to produce spontaneously a thermodynamically stable microemulsion.

50 x) Microcapsules (CS)
An oil phase comprising 5-50 wt% of a compound I, 0-40 wt% water insoluble organic solvent (e.g. aromatic hydrocarbon), 2-15 wt% acrylic monomers (e.g. methylmethacrylate, methacrylic acid and a di- or triacrylate) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol). Radical polymerization initiated by a radical initiator results in the formation of poly(meth)acrylate microcapsules. Alternatively, an oil phase comprising 5-50 wt% of a compound I according to the invention, 0-40 wt% water insoluble organic solvent (e.g. aromatic hydrocarbon), and an isocyanate monomer (e.g. diphenylmethene-4,4'-diisocyanate) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol). The addition of a polyamine (e.g. hexamethylenediamine) results in the formation of polyurea microcapsules. The monomers amount to 1-10 wt%. The wt% relate to the total CS composition.

55 xi) Dustable powders (DP, DS)
1-10 wt% of a compound I are ground finely and mixed intimately with solid carrier (e.g. finely divided kaolin) ad 100 wt%.

x) Granules (GR, FG)
0.5-30 wt% of a compound I is ground finely and associated with solid carrier (e.g. silicate) ad 100 wt%. Granulation is achieved by extrusion, spray-drying or fluidized bed.

xi) Ultra-low volume liquids (UL)

1-50 wt% of a compound I are dissolved in organic solvent (e.g. aromatic hydrocarbon) ad 100 wt%.

The compositions types i) to xi) may optionally comprise further auxiliaries, such as 0.1-1 wt% bactericides, 5-15 wt% anti-freezing agents, 0.1-1 wt% anti-foaming agents, and 0.1-1 wt% colorants.

5

[0086] The agrochemical compositions generally comprise between 0.01 and 95%, preferably between 0.1 and 90%, and in particular between 0.5 and 75%, by weight of active substance. The active substances are employed in a purity of from 90% to 100%, preferably from 95% to 100% (according to NMR spectrum).

10

[0087] Solutions for seed treatment (LS), Suspoemulsions (SE), flowable concentrates (FS), powders for dry treatment (DS), water-dispersible powders for slurry treatment (WS), water-soluble powders (SS), emulsions (ES), emulsifiable concentrates (EC) and gels (GF) are usually employed for the purposes of treatment of plant propagation materials, particularly seeds. The compositions in question give, after two-to-tenfold dilution, active substance concentrations of from 0.01 to 60% by weight, preferably from 0.1 to 40%, in the ready-to-use preparations. Application can be carried out before or during sowing. Methods for applying compound I and compositions thereof, respectively, on to plant propagation material, especially seeds include dressing, coating, pelleting, dusting, soaking and in-furrow application methods of the propagation material. Preferably, compound I or the compositions thereof, respectively, are applied on to the plant propagation material by a method such that germination is not induced, e. g. by seed dressing, pelleting, coating and dusting.

15

[0088] When employed in plant protection, the amounts of active substances applied are, depending on the kind of effect desired, from 0.001 to 2 kg per ha, preferably from 0.005 to 2 kg per ha, more preferably from 0.05 to 0.9 kg per ha, and in particular from 0.1 to 0.75 kg per ha.

20

[0089] In treatment of plant propagation materials such as seeds, e. g. by dusting, coating or drenching seed, amounts of active substance of from 0.1 to 1000 g, preferably from 1 to 1000 g, more preferably from 1 to 100 g and most preferably from 5 to 100 g, per 100 kilogram of plant propagation material (preferably seeds) are generally required.

25

[0090] When used in the protection of materials or stored products, the amount of active substance applied depends on the kind of application area and on the desired effect. Amounts customarily applied in the protection of materials are 0.001 g to 2 kg, preferably 0.005 g to 1 kg, of active substance per cubic meter of treated material.

30

[0091] Various types of oils, wetters, adjuvants, fertilizer, or micronutrients, and further pesticides (e.g. herbicides, insecticides, fungicides, growth regulators, safeners) may be added to the active substances or the compositions comprising them as premix or, if appropriate not until immediately prior to use (tank mix). These agents can be admixed with the compositions according to the invention in a weight ratio of 1:100 to 100:1, preferably 1:10 to 10:1.

35

[0092] The user applies the composition according to the invention usually from a predosage device, a knapsack sprayer, a spray tank, a spray plane, or an irrigation system. Usually, the agrochemical composition is made up with water, buffer, and/or further auxiliaries to the desired application concentration and the ready-to-use spray liquor or the agrochemical composition according to the invention is thus obtained. Usually, 20 to 2000 liters, preferably 50 to 400 liters, of the ready-to-use spray liquor are applied per hectare of agricultural useful area.

40

[0093] According to one embodiment, individual components of the composition according to the invention such as parts of a kit or parts of a binary or ternary mixture may be mixed by the user himself in a spray tank and further auxiliaries may be added, if appropriate.

45

[0094] Mixing the compounds I or the compositions comprising them in the use form as fungicides with other fungicides results in many cases in an expansion of the fungicidal spectrum of activity being obtained or in a prevention of fungicide resistance development. Furthermore, in many cases, synergistic effects are obtained.

[0095] The following list of active substances, in conjunction with which the compounds I can be used, is intended to illustrate the possible combinations but does not limit them:

50

A) Respiration inhibitors

55

- Inhibitors of complex III at Q_o site (e.g. strobilurins): azoxystrobin, coumethoxystrobin, coumoxystrobin, dimoxystrobin, enestroburin, fenaminstrobin, fenoxyystrobin/flufenoxystrobin, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, picoxyystrobin, pyraclostrobin, pyrametostrobin, pyraoxystrobin, trifloxystrobin, 2-[2-(2,5-dimethyl-phenoxy)methyl]-3-methoxy-acrylic acid methyl ester and 2-(2-(3-(2,6-dichlorophenyl)-1-methyl-allylideneaminoxy)methyl)-phenyl]-2-methoxyimino-N-methyl-acetamide, pyribencarb, triclopyricarb/chlorodincarb, famoxadone, fenamidone;
- inhibitors of complex III at Q_i site: cyazofamid, amisulbrom, [(3S,6S,7R,8R)-8-benzyl-3-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxinan-7-yl] 2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[(3-acetoxymethoxy)-4-methoxy-pyridine-2-carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxinan-7-yl] 2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[(3-isobutoxycarbonyloxy-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxinan-7-yl] 2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[(3-(1,3-benzod-

oxol-5-ylmethoxy)-4-methoxypyridine-2-carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate;

- inhibitors of complex II (e. g. carboxamides): benodanil, bixafen, boscalid, carboxin, fenfuram, fluopyram, flutolanil, fluxapyroxad, furametpyr, isopyrazam, mepronil, oxycarboxin, penflufen, penthiopyrad, sedaxane, tecloftalam, thifluzamide, N-(4'-trifluoromethylthiobiphenyl-2-yl)-3-difluoromethyl-1-methyl-1 H-pyrazole-4-carboxamide, N-(2-(1,3,3-trimethyl-butyl)-phenyl)-1,3-dimethyl-5-fluoro-1 H-pyrazole-4-carboxamide, N-[9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide;
- other respiration inhibitors (e.g. complex I, uncouplers): diflumetorim, (5,8-difluoroquinazolin-4-yl)-{2-[2-fluoro-4-(4-trifluoromethylpyridin-2-yloxy)-phenyl]-ethyl}-amine; nitrophenyl derivates: binapacryl, dinobuton, dinocap, fluazinam; ferimzone; organometal compounds: fentin salts, such as fentin-acetate, fentin chloride or fentin hydroxide; ametoctradin; and silthiofam;

B) Sterol biosynthesis inhibitors (SBI fungicides)

- C14 demethylase inhibitors (DMI fungicides): triazoles: azaconazole, bitertanol, bromuconazole, cyproconazole, difenoconazole, diniconazole, diniconazole-M, epoxiconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imibenconazole, ipconazole, metconazole, myclobutanil, oxoconazole, paclobutrazole, penconazole, propiconazole, prothioconazole, simeconazole, tebuconazole, tetaconazole, triadimefon, triadimenol, triticonazole, uniconazole; imidazoles: imazalil, pefurazoate, prochloraz, triflumizol; pyrimidines, pyridines and piperazines: fenarimol, nuarimol, pyrifenox, triforine;
- Delta14-reductase inhibitors: aldimorph, dodemorph, dodemorph-acetate, fenpropimorph, tridemorph, fenpropidin, piperalin, spiroxamine;
- Inhibitors of 3-keto reductase: fenhexamid;

C) Nucleic acid synthesis inhibitors

- phenylamides or acyl amino acid fungicides: benalaxyl, benalaxyl-M, kiralaxyl, metalaxyl, metalaxyl-M (mefenoxam), ofurace, oxadixyl;
- others: hymexazole, octhilinone, oxolinic acid, bupirimate, 5-fluorocytosine, 5-fluoro-2-(p-tolylmethoxy)pyrimidin-4-amine, 5-fluoro-2-(4-fluorophenylmethoxy)pyrimidin-4-amine;

D) Inhibitors of cell division and cytoskeleton

- tubulin inhibitors, such as benzimidazoles, thiophanates: benomyl, carbendazim, fuberidazole, thiabendazole, thiophanate-methyl; triazolopyrimidines: 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine
- other cell division inhibitors: diethofencarb, ethaboxam, pencycuron, fluopicolide, zoxamide, metrafenone, pyriofenone;

E) Inhibitors of amino acid and protein synthesis

- methionine synthesis inhibitors (anilino-pyrimidines): cyprodinil, mepanipyrim, pyrimethanil;
- protein synthesis inhibitors: blasticidin-S, kasugamycin, kasugamycin hydrochloride-hydrate, mildiomycin, streptomycin, oxytetracyclin, polyoxine, validamycin A;

F) Signal transduction inhibitors

- MAP / histidine kinase inhibitors: fluoroimid, iprodione, procymidone, vinclozolin, fenpiclonil, fludioxonil;
- G protein inhibitors: quinoxyfen;

G) Lipid and membrane synthesis inhibitors

- Phospholipid biosynthesis inhibitors: edifenphos, iprobenfos, pyrazophos, isoprothiolane;
- lipid peroxidation: dicloran, quintozene, tecnazene, tolclofos-methyl, biphenyl, chloroneb, etridiazole;
- phospholipid biosynthesis and cell wall deposition: dimethomorph, flumorph, mandipropamid, pyrimorph, ben-thiavalicarb, iprovalicarb, valifenalate and N-(1-(4-cyano-phenyl)ethanesulfonyl)-but-2-yl) carbamic acid-(4-fluorophenyl) ester;

- compounds affecting cell membrane permeability and fatty acids: propamocarb, propamocarb-hydrochlorid

H) Inhibitors with Multi Site Action

- inorganic active substances: Bordeaux mixture, copper acetate, copper hydroxide, copper oxychloride, basic copper sulfate, sulfur;
- thio- and dithiocarbamates: ferbam, mancozeb, maneb, metam, metiram, propineb, thiram, zineb, ziram;
- organochlorine compounds (e.g. phthalimides, sulfamides, chloronitriles): anilazine, chlorothalonil, captafol, captan, folpet, dichlofuanid, dichlorophen, flusulfamide, hexachlorobenzene, pentachlorphenole and its salts, phthalide, tolylfuanid, N-(4-chloro-2-nitro-phenyl)-N-ethyl-4-methyl-benzenesulfonamide;
- guanidines and others: guanidine, dodine, dodine free base, guazatine, guazatine-acetate, iminoctadine, iminoctadine-triacetate, iminoctadine-tris(albesilate), dithianon;

I) Cell wall synthesis inhibitors

- inhibitors of glucan synthesis: validamycin, polyoxin B; melanin synthesis inhibitors: pyroquilon, tricyclazole, carpropamid, dicyclomet, fenoxyanil;

J) Plant defence inducers

- acibenzolar-S-methyl, probenazole, isotianil, tiadinil, prohexadione-calcium; phosphonates: fosetyl, fosetyl-alumin, phosphorous acid and its salts;

K) Unknown mode of action

- bronopol, chinomethionat, cyflufenamid, cymoxanil, dazomet, debacarb, diclomezine, difenzoquat, difenzoquat-methylsulfate, diphenylamin, fenpyrazamine, flumetover, flusulfamide, flutianil, methasulfocarb, nitrapyrin, nitrothial-isopropyl, oxin-copper, proquinazid, tebuflouquin, tecloftalam, triazoxide, 2-butoxy-6-iodo-3-propylchromen-4-one, N-(cyclopropylmethoxyimino-(6-difluoro-methoxy-2,3-di-fluoro-phenyl)-methyl)-2-phenyl acetamide, N'-(4-(4-chloro-3-trifluoromethyl-phenoxy)-2,5-dimethyl-phenyl)-N-ethyl-N-methyl formamidine, N'-(4-(4-fluoro-3-trifluoromethyl-phenoxy)-2,5-dimethyl-phenyl)-N-ethyl-N-methyl formamidine, N'-(2-methyl-5-trifluoromethyl-4-(3-trimethylsilanyl-propoxy)-phenyl)-N-ethyl-N-methyl formamidine, N'-(5-difluoromethyl-2-methyl-4-(3-trimethylsilanyl-propoxy)-phenyl)-N-ethyl-N-methyl formamidine, 2-{1-[2-(5-methyl-3-trifluoromethyl-pyrazole-1-yl)-acetyl]-piperidin-4-yl}-thiazole-4-carboxylic acid methyl-(1,2,3,4-tetrahydro-naphthalen-1-yl)-amide, 2-{1-[2-(5-methyl-3-trifluoromethyl-pyrazole-1-yl)-acetyl]-piperidin-4-yl}-thiazole-4-carboxylic acid methyl-(R)-1,2,3,4-tetrahydro-naphthalen-1-yl-amide, 1-[4-[4-(2,6-difluorophenyl)-4,5-dihydro-3-isoxazolyl]-2-thiazolyl]-1-piperidinyl]-2-[5-methyl-3-(trifluoromethyl)-1 H-pyrazol-1-yl]ethanone, methoxy-acetic acid 6-tert-butyl-8-fluoro-2,3-dimethyl-quinolin-4-yl ester, N-Methyl-2-{1-[(5-methyl-3-trifluoromethyl-1H-pyrazol-1-yl)-acetyl]-piperidin-4-yl}-N-[(1 R)-1,2,3,4-tetrahydronaphthalen-1-yl]-4-thiazolecarboxamide, 3-[5-(4-methylphenyl)-2,3-dimethylisoxazolidin-3-yl]-pyridine, 3-[5-(4-chloro-phenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine (pyrisoxazole), N-(6-methoxy-pyridin-3-yl) cyclopropanecarboxylic acid amide, 5-chloro-1-(4,6-dimethoxy-pyrimidin-2-yl)-2-methyl-1H-benzoimidazole, 2-(4-chloro-phenyl)-N-[4-(3,4-dimethoxy-phenyl)-isoxazol-5-yl]-2-prop-2-ynylxyacetamide;

L) Antifungal biocontrol agents, plant bioactivators: *Ampelomyces quisqualis* (e.g. AQ 10[®] from Intrachem Bio GmbH & Co. KG, Germany), *Aspergillus flavus* (e.g. AFLAGUARD[®] from Syngenta, CH), *Aureobasidium pullulans* (e.g. BOTECTOR[®] from bio-ferm GmbH, Germany), *Bacillus pumilus* (e.g. NRRL Accession No. B-30087 in SONATA[®] and BALLAD[®] Plus from AgraQuest Inc., USA), *Bacillus subtilis* (e.g. isolate NRRL-Nr. B-21661 in RHAPSODY[®], SERENADE[®] MAX and SERENADE[®] ASO from AgraQuest Inc., USA), *Bacillus subtilis* var. *amylolique-faciens* FZB24 (e.g. TAEGRO[®] from Novozyme Biologicals, Inc., USA), *Candida oleophila* I-82 (e.g. ASPIRE[®] from Ecogen Inc., USA), *Candida saitoana* (e.g. BIOCURE[®] (in mixture with lysozyme) and BIOCOAT[®] from Micro Flo Company, USA (BASF SE) and Arysta), Chitosan (e.g. ARMOUR-ZEN from BotriZen Ltd., NZ), *Clonostachys rosea* f. *catenulata*, also named *Gliocladium catenulatum* (e.g. isolate J1446: PRESTOP[®] from Verdera, Finland), *Coniothyrium minitans* (e.g. CONTANS[®] from Prophita, Germany), *Cryphonectria parasitica* (e.g. *Endothia parasitica* from CNICM, France), *Cryptococcus albidus* (e.g. YIELD PLUS[®] from Anchor Bio-Technologies, South Africa), *Fusarium oxysporum* (e.g. BIOFOX[®] from S.I.A.P.A., Italy, FUSACLEAN[®] from Natural Plant Protection, France), *Metschnikowia fructicola* (e.g. SHEMER[®] from Agrogreen, Israel), *Microdochium dimerum* (e.g. ANTIBOT[®] from Agrauxine, France), *Phlebiopsis gigantea* (e.g. ROTSOP[®] from Verdera, Finland), *Pseudozyma flocculosa* (e.g. SPORODEX[®] from

Plant Products Co. Ltd., Canada), *Pythium oligandrum* DV74 (e.g. POLYVERSUM® from Remeslo SSRO, Biopreparaty, Czech Rep.), *Reynoutria sachlinensis* (e.g. REGALIA® from Marrone BioInnovations, USA), *Talaromyces flavus* V117b (e.g. PROTUS® from Prophita, Germany), *Trichoderma asperellum* SKT-1 (e.g. ECO-HOPE® from Kumiai Chemical Industry Co., Ltd., Japan), *T. atroviride* LC52 (e.g. SENTINEL® from Agrimm Technologies Ltd, NZ), *T. harzianum*T-22 (e.g. PLANTSHIELD® der Firma BioWorks Inc., USA), *T. harzianum* TH 35 (e.g. ROOT PRO® from Mycontrol Ltd., Israel), *T. harzianum* T-39 (e.g. TRICHODEX® and TRICHODERMA 2000® from Mycontrol Ltd., Israel and Makhteshim Ltd., Israel), *T. harzianum* and *T. viride* (e.g. TRICHOPEL from Agrimm Technologies Ltd, NZ), *T. harzianum* ICC012 and *T. viride* ICC080 (e.g. REMEDIER® WP from Isagro Ricerca, Italy), *T. polysporum* and *T. harzianum* (e.g. BINAB® from BINAB Bio-Innovation AB, Sweden), *T. stromaticum* (e.g. TRICOVAB® from C.E.P.L.A.C., Brazil), *T. virens* GL-21 (e.g. SOILGARD® from Certis LLC, USA), *T. viride* (e.g. TRIECO® from Ecosense Labs. (India) Pvt. Ltd., Indien, BIO-CURE® F from T. Stanes & Co. Ltd., Indien), *T. viride* TV1 (e.g. *T. viride* TV1 from Agribiotec srl, Italy), *Ulocladium oudemansii* HRU3 (e.g. BOTRY-ZEN® from Botry-Zen Ltd, NZ);

M) Growth regulators

abscisic acid, amidochlor, ancytidol, 6-benzylaminopurine, brassinolide, butralin, chlormequat (chlormequat chloride), choline chloride, cyclanilide, daminozide, dike-gulac, dimethipin, 2,6-dimethylpuridine, ethephon, flumetralin, flurprimidol, fluthiacet, forchlorfenuron, gibberellic acid, inabenfide, indole-3-acetic acid, maleic hydrazide, mefluclidide, mepiquat (mepiquat chloride), naphthaleneacetic acid, N-6-benzyladenine, paclobutrazol, prohexadione (prohexadione-calcium), prohydrojasmon, thidiazuron, triapenthenol, tributyl phosphorothioate, 2,3,5-tri-iodobenzoic acid, trinexapac-ethyl and uniconazole;

N) Herbicides

- acetamides: acetochlor, alachlor, butachlor, dimethachlor, dimethenamid, flufenacet, mefenacet, metolachlor, metazachlor, napropamide, naproanilide, pethoxamid, pretilachlor, propachlor, thenylchlor;
- amino acid derivatives: bilanafos, glyphosate, glufosinate, sulfosate;
- aryloxyphenoxypropionates: clodinafop, cyhalofop-butyl, fenoxaprop, fluazifop, haloxyfop, metamifop, pro-paquizafop, quizalofop, quizalofop-P-tefuryl;
- Bipyridyls: diquat, paraquat;
- (thio)carbamates: asulam, butylate, carbetamide, desmedipham, dimepiperate, eptam (EPTC), esprocarb, molinate, orbencarb, phenmedipham, prosulfocarb, pyributicarb, thiobencarb, triallate;
- cyclohexanediones: butoxydim, clethodim, cycloxydim, profoxydim, sethoxydim, tepraloxydim, tralkoxydim;
- dinitroanilines: benfluralin, ethalfluralin, oryzalin, pendimethalin, prodiamine, trifluralin;
- diphenyl ethers: acifluorfen, aclonifen, bifenox, diclofop, ethoxyfen, fomesafen, lactofen, oxyfluorfen;
- hydroxybenzonitriles: bomoxynil, dichlobenil, ioxynil;
- imidazolinones: imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr;
- phenoxy acetic acids: clomeprop, 2,4-dichlorophenoxyacetic acid (2,4-D), 2,4-DB, dichlorprop, MCPA, MCPA-thioethyl, MCPB, Mecoprop;
- pyrazines: chlорidazon, flufenpyr-ethyl, fluthiacet, norflurazon, pyridate;
- pyridines: aminopyralid, clopyralid, diflufenican, dithiopyr, fluridone, fluoxypyrr, picloram, picolinafen, thiazopyr;
- sulfonyl ureas: amidosulfuron, azimsulfuron, bensulfuron, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethoxysulfuron, flazasulfuron, flucetosulfuron, flupyrifos, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metazosulfuron, metsulfuron-methyl, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxsulfuron, triflusulfuron, tritosulfuron, 1-((2-chloro-6-propyl-imidazo[1,2-b]pyridazin-3-yl)sulfonyl)-3-(4,6-dimethoxy-pyrimidin-2-yl)urea;
- triazines: ametryn, atrazine, cyanazine, dimethametryn, ethiozin, hexazinone, metamitron, metribuzin, prometryn, simazine, terbutylazine, terbutryn, triaziflam;
- ureas: chlorotoluron, daimuron, diuron, fluometuron, isoproturon, linuron, methabenzthiazuron, tebuthiuron;
- other acetolactate synthase inhibitors: bispyribac-sodium, cloransulam-methyl, diclosulam, florasulam, flucarbazone, flumetsulam, metosulam, ortho-sulfamuron, penoxsulam, propoxycarbazone, pyribambenz-propyl, pyribenzoxim, pyrifalid, pyriminobac-methyl, pyrimisulfan, pyriproxyfen, pyroxasulfone, pyroxsulam;
- others: amicarbazone, aminotriazole, anilofos, beflubutamid, benazolin, bencarbazone, benfluresate, benzofenap, bentazone, benzobicyclon, bicyclopyrone, bromacil, bromobutide, butafenacil, butamifos, cafenstrole, carfentrazone, cinidon-ethyl, chlorthal, cinmethylin, clomazone, cumyluron, cyprosulfamide, dicamba, difenzoquat, diflufenzopyr, *Drechslera monoceras*, endothal, ethofumesate, etobenzanid, fenoxasulfone, fentrazamide, flumiclorac-pentyl, flumioxazin, flupoxam, flurochloridone, flurtamone, indanofan, isoxaben, isoxaflutole, lenacil, propanil, propyzamide, quinclorac, quinmerac, mesotrione, methyl arsonic acid, naptalam, oxadiargyl, oxadiazon, oxaziclofene, pentoxazone, pinoxaden, pyraclonil, pyraflufen-ethyl, pyrasulfotole, pyrazoxyfen, pyra-

5 zolynate, quinoclamine, saflufenacil, sulcotrione, sulfentrazone, terbacil, tefuryltrione, tembotrione, thiencarbazone, topramezone, (3-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)-phenoxy]-pyridin-2-yloxy)-acetic acid ethyl ester, 6-amino-5-chloro-2-cyclopropyl-pyrimidine-4-carboxylic acid methyl ester, 6-chloro-3-(2-cyclopropyl-6-methyl-phenoxy)-pyridazin-4-ol, 4-amino-3-chloro-6-(4-chlorophenyl)-5-fluoropyridine-2-carboxylic acid, 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-pyridine-2-carboxylic acid methyl ester, and 4-amino-3-chloro-6-(4-chloro-3-dimethylamino-2-fluoro-phenyl)-pyridine-2-carboxylic acid methyl ester.

10 O) Insecticides

- 10 - organo(thio)phosphates: acephate, azamethiphos, azinphos-methyl, chlorpyrifos, chlorpyrifos-methyl, chlорfenvinphos, diazinon, dichlorvos, dicrotophos, dimethoate, disulfoton, ethion, fenitrothion, fenthion, isoxathion, malathion, methamidophos, methidathion, methyl-parathion, mevinphos, monocrotophos, oxydemeton-methyl, paraoxon, parathion, phentoate, phosalone, phosmet, phosphamidon, phorate, phoxim, pirimiphos-methyl, profenofos, prothiofos, sulprophos, tetrachlorvinphos, terbufos, triazophos, trichlorfon;
- 15 - carbamates: alanycarb, aldicarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, fenoxy carb, furathiocarb, methiocarb, methomyl, oxamyl, pirimicarb, propoxur, thiodicarb, triazamate;
- 20 - pyrethroids: allethrin, bifenthrin, cyfluthrin, cyhalothrin, cyphenothrin, cypermethrin, alpha-cypermethrin, beta-cypermethrin, zeta-cypermethrin, deltamethrin, esfenvalerate, etofenprox, fenpropothrin, fenvalerate, imiprothrin, lambda-cyhalothrin, permethrin, prallethrin, pyrethrin I and II, resmethrin, silafluofen, tau-fluvalinate, te-fluthrin, tetramethrin, tralomethrin, transfluthrin, profluthrin, dimefluthrin;
- 25 - insect growth regulators: a) chitin synthesis inhibitors: benzoylureas: chlorfluazuron, cyramazin, diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, teflubenzuron, triflumuron; buprofezin, diafenolan, hexythiazox, etoxazole, clofentazine; b) ecdysone antagonists: halofenozide, methoxyfenozide, tebufenozide, azadirachtin; c) juvenoids: pyriproxyfen, methoprene, fenoxy carb; d) lipid biosynthesis inhibitors: spirodiclofen, spiromesifen, spirotetramat;
- 30 - nicotinic receptor agonists/antagonists compounds: clothianidin, dinotefuran, flupyradifurone, imidacloprid, thiamethoxam, nitenpyram, acetamiprid, thiacloprid, 1-(2-chloro-thiazol-5-ylmethyl)-2-nitrimino-3,5-dimethyl-[1,3,5]triazinane;
- 35 - GABA antagonist compounds: endosulfan, ethiprole, fipronil, vaniliprole, pyrafluprole, pyriproxyfen, 5-amino-1-(2,6-dichloro-4-methyl-phenyl)-4-sulfinamoyl-1H-pyrazole-3-carbothioic acid amide;
- macrocyclic lactone insecticides: abamectin, emamectin, milbemectin, lepimectin, spinosad, spinetoram;
- mitochondrial electron transport inhibitor (METI) I acaricides: fenazaquin, pyridaben, tebufenpyrad, tolfenpyrad, flufennerim;
- 40 - METI II and III compounds: acequinocyl, fluacyprin, hydramethylnon;
- Uncouplers: chlорfenapyr;
- oxidative phosphorylation inhibitors: cyhexatin, diafenthiuron, fenbutatin oxide, propargite;
- moulting disruptor compounds: cryomazine;
- mixed function oxidase inhibitors: piperonyl butoxide;
- 45 - sodium channel blockers: indoxacarb, metaflumizone;
- others: benclothiaz, bifenazate, cartap, flonicamid, pyridalyl, pymetrozine, sulfur, thiocyclam, flubendiamide, chlorantraniliprole, cyazypyr (HGW86), cyenopyrafen, flupyrazofos, cyflumetofen, amidoflumet, imicyafos, bis-trifluron, and pyrifluquinazon.

50 **[0096]** The present invention furthermore relates to agrochemical compositions comprising a mixture of at least one compound I (component 1) and at least one further active substance useful for plant protection, e. g. selected from the groups A) to O) (component 2), in particular one further fungicide, e. g. one or more fungicide from the groups A) to L), as described above, and if desired one suitable solvent or solid carrier. Those mixtures are of particular interest, since many of them at the same application rate show higher efficiencies against harmful fungi. Furthermore, combating harmful fungi with a mixture of compounds I and at least one fungicide from groups A) to L), as described above, is more efficient than combating those fungi with individual compounds I or individual fungicides from groups A) to L). By applying compounds I together with at least one active substance from groups A) to O) a synergistic effect can be obtained, i.e. more than simple addition of the individual effects is obtained (synergistic mixtures).

55 **[0097]** This can be obtained by applying the compounds I and at least one further active substance simultaneously, either jointly (e. g. as tank-mix) or separately, or in succession, wherein the time interval between the individual applications is selected to ensure that the active substance applied first still occurs at the site of action in a sufficient amount at the time of application of the further active substance(s). The order of application is not essential for working of the present invention.

[0098] In binary mixtures, i.e. compositions according to the invention comprising one compound I (component 1) and one further active substance (component 2), e.g. one active substance from groups A) to O), the weight ratio of component 1 and component 2 generally depends from the properties of the active substances used, usually it is in the range of from 1:100 to 100:1, regularly in the range of from 1:50 to 50:1, preferably in the range of from 1:20 to 20:1, more preferably in the range of from 1:10 to 10:1 and in particular in the range of from 1:3 to 3:1.

[0099] In ternary mixtures, i.e. compositions according to the invention comprising one compound I (component 1) and a first further active substance (component 2) and a second further active substance (component 3), e.g. two active substances from groups A) to O), the weight ratio of component 1 and component 2 depends from the properties of the active substances used, preferably it is in the range of from 1:50 to 50:1 and particularly in the range of from 1:10 to 10:1, and the weight ratio of component 1 and component 3 preferably is in the range of from 1:50 to 50:1 and particularly in the range of from 1:10 to 10:1.

[0100] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group A) (component 2) and particularly selected from azoxystrobin, dimoxystrobin, fluoxastrobin, kresoxim-methyl, orysastrobin, picoxystrobin, pyraclostrobin, trifloxystrobin; famoxadone, fenamidone; bixafen, boscalid, fluopyram, fluxapyroxad, isopyrazam, penflufen, penthiopyrad, sedaxane; ametoctradin, cyazofamid, fluazinam, fentin salts, such as fentin acetate.

[0101] Preference is given to mixtures comprising a compound of formula I (component 1) and at least one active substance selected from group B) (component 2) and particularly selected from cyproconazole, difenoconazole, epoxiconazole, fluquinconazole, flusilazole, flutriafol, metconazole, myclobutanil, penconazole, propiconazole, prothioconazole, triadimefon, triadimenol, tebuconazole, tetriconazole, triticonazole, prochloraz, fenarimol, triforine; dodemorph, fenpropimorph, tridemorph, fenpropidin, spiroxamine; fenhexamid.

[0102] Preference is given to mixtures comprising a compound of formula I (component 1) and at least one active substance selected from group C) (component 2) and particularly selected from metalaxyl, (metalaxyl-M) mefenoxam, ofurace.

[0103] Preference is given to mixtures comprising a compound of formula I (component 1) and at least one active substance selected from group D) (component 2) and particularly selected from benomyl, carbendazim, thiophanate-methyl, ethaboxam, fluopicolide, zoxamide, metrafenone, pyriofenone.

[0104] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group E) (component 2) and particularly selected from cyprodinil, mepanipyrim, pyrimethanil.

[0105] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group F) (component 2) and particularly selected from iprodione, fludioxonil, vinclozolin, quinoxyfen.

[0106] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group G) (component 2) and particularly selected from dimethomorph, flumorph, iprovalicarb, benthiazalicarb, mandipropamid, propamocarb.

[0107] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group H) (component 2) and particularly selected from copper acetate, copper hydroxide, copper oxychloride, copper sulfate, sulfur, mancozeb, metiram, propineb, thiram, captafol, folpet, chlorothalonil, dichlofluanid, dithianon.

[0108] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group I) (component 2) and particularly selected from carpropamid and fenoxanil.

[0109] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group J) (component 2) and particularly selected from acibenzolar-S-methyl, probenazole, tiadinil, fosetyl, fosetyl-aluminium, H_3PO_3 and salts thereof.

[0110] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group K) (component 2) and particularly selected from cymoxanil, proquinazid and *N*-methyl-2-{1-[(5-methyl-3-trifluoromethyl-1H-pyrazol-1-yl)-acetyl]-piperidin-4-yl}-*N*[(1R)-1,2,3,4-tetrahydronaphthalen-1-yl]-4-thiazole-carboxamide.

[0111] Preference is also given to mixtures comprising a compound I (component 1) and at least one active substance selected from group L) (component 2) and particularly selected from *Bacillus subtilis* strain N RRL No. B-21661, *Bacillus pumilus* strain N RRL No. B-30087 and *Ulocladium oude mansii*.

[0112] Accordingly, the present invention furthermore relates to compositions comprising one compound I (component 1) and one further active substance (component 2), which further active substance is selected from the column "Component 2" of the lines B-1 to B-361 of Table B.

[0113] A further embodiment relates to the compositions B-1 to B-361 listed in Table B, where a row of Table B corresponds in each case to a fungicidal composition comprising one of the in the present specification individualized compounds of formula I (component 1) and the respective further active substance from groups A) to O) (component 2) stated in the row in question. Preferably, the compositions described comprise the active substances in synergistically effective amounts.

Table B: Composition comprising one individualized compound I and one further active substance from groups A) to O)

Mixture	Component 1	Component 2
B-1	one individualized compound I	Azoxystrobin
B-2	one individualized compound I	Coumethoxystrobin
B-3	one individualized compound I	Coumoxystrobin
B-4	one individualized compound I	Dimoxystrobin
B-5	one individualized compound I	Enestroburin
B-6	one individualized compound I	Fenaminstrobin
B-7	one individualized compound I	Fenoxystrobin/Flufenoxystrobin
B-8	one individualized compound I	Fluoxastrobin
B-9	one individualized compound I	Kresoxim-methyl
B-10	one individualized compound I	Metominostrobin
B-11	one individualized compound I	Orysastrobin
B-12	one individualized compound I	Picoxystrobin
B-13	one individualized compound I	Pyraclostrobin
B-14	one individualized compound I	Pyrametostrobin
B-15	one individualized compound I	Pyraoxystrobin
B-16	one individualized compound I	Pyribencarb
B-17	one individualized compound I	Trifloxystrobin
B-18	one individualized compound I	Triclopyricarb/Chlorodincarb
B-19	one individualized compound I	2-[2-(2,5-dimethyl-phenoxy)methyl]-phenyl]-3-methoxy-acrylic acid methyl ester
B-20	one individualized compound I	2-(2-(3-(2,6-dichlorophenyl)-1-methyl-allylideneaminoxy)methyl)-phenyl)-2-methoxyimino-N-methyl-acetamide
B-21	one individualized compound I	Benalaxyll
B-22	one individualized compound I	Benalaxyll-M
B-23	one individualized compound I	Benodanil
B-24	one individualized compound I	Bixafen
B-25	one individualized compound I	Boscalid
B-26	one individualized compound I	Carboxin
B-27	one individualized compound I	Fenfuram
B-28	one individualized compound I	Fenhexamid
B-29	one individualized compound I	Flutolanil
B-30	one individualized compound I	Fluxapyroxad
B-31	one individualized compound I	Furametylpyr
B-32	one individualized compound I	Isopyrazam
B-33	one individualized compound I	Isotianil
B-34	one individualized compound I	Kiralaxyll
B-35	one individualized compound I	Mepronil

(continued)

Mixture	Component 1	Component 2
B-36	one individualized compound I	Metalaxyl
B-37	one individualized compound I	Metalaxyl-M
B-38	one individualized compound I	Ofurace
B-39	one individualized compound I	Oxadixyl
B-40	one individualized compound I	Oxycarboxin
B-41	one individualized compound I	Penflufen
B-42	one individualized compound I	Penthiopyrad
B-43	one individualized compound I	Sedaxane
B-44	one individualized compound I	Tecloftalam
B-45	one individualized compound I	Thifluzamide
B-46	one individualized compound I	Tiadnil
B-47	one individualized compound I	2-Amino-4-methyl-thiazole-5-carboxylic acid anilide
B-48	one individualized compound I	N-(4'-trifluoromethylthiobiphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide
B-49	one individualized compound I	N-(2-(1,3,3-trimethyl-butyl)-phenyl)-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide
B-50	one individualized compound I	N-[9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide
B-51	one individualized compound I	Dimethomorph
B-52	one individualized compound I	Flumorph
B-53	one individualized compound I	Pyrimorph
B-54	one individualized compound I	Flumetover
B-55	one individualized compound I	Fluopicolide
B-56	one individualized compound I	Fluopyram
B-57	one individualized compound I	Zoxamide
B-58	one individualized compound I	Carpropamid
B-59	one individualized compound I	Diclocymet
B-60	one individualized compound I	Mandipropamid
B-61	one individualized compound I	Oxytetracyclin
B-62	one individualized compound I	Silthiofam
B-63	one individualized compound I	N-(6-methoxy-pyridin-3-yl) cyclopro-panecarboxylic acid amide
B-64	one individualized compound I	Azaconazole
B-65	one individualized compound I	Bitertanol
B-66	one individualized compound I	Bromuconazole
B-67	one individualized compound I	Cyproconazole
B-68	one individualized compound I	Difenconazole
B-69	one individualized compound I	Diniconazole

(continued)

Mixture	Component 1	Component 2
B-70	one individualized compound I	Diniconazole-M
B-71	one individualized compound I	Epoxiconazole
B-72	one individualized compound I	Fenbuconazole
B-73	one individualized compound I	Fluquinconazole
B-74	one individualized compound I	Flusilazole
B-75	one individualized compound I	Flutriafol
B-76	one individualized compound I	Hexaconazol
B-77	one individualized compound I	Imibenconazole
B-78	one individualized compound I	Ipconazole
B-79	one individualized compound I	Metconazole
B-80	one individualized compound I	Myclobutanil
B-81	one individualized compound I	Oxoconazol
B-82	one individualized compound I	Paclobutrazol
B-83	one individualized compound I	Penconazole
B-84	one individualized compound I	Propiconazole
B-85	one individualized compound I	Prothioconazole
B-86	one individualized compound I	Simeconazole
B-87	one individualized compound I	Tebuconazole
B-88	one individualized compound I	Tetraconazole
B-89	one individualized compound I	Triadimefon
B-90	one individualized compound I	Triadimenol
B-91	one individualized compound I	Triticonazole
B-92	one individualized compound I	Uronicazole
B-93	one individualized compound I	Cyazofamid
B-94	one individualized compound I	Imazalil
B-95	one individualized compound I	Imazalil-sulfate
B-96	one individualized compound I	Pefurazoate
B-97	one individualized compound I	Prochloraz
B-98	one individualized compound I	Triflumizole
B-99	one individualized compound I	Benomyl
B-100	one individualized compound I	Carbendazim
B-101	one individualized compound I	Fuberidazole
B-102	one individualized compound I	Thiabendazole
B-103	one individualized compound I	Ethaboxam
B-104	one individualized compound I	Etridiazole
B-105	one individualized compound I	Hymexazole
B-106	one individualized compound I	2-(4-Chloro-phenyl)-N-[4-(3,4-dimethoxy-phenyl)-isoxazol-5-yl]-2-prop-2-yn-yloxy-acetamide

(continued)

Mixture	Component 1	Component 2
5 B-107	one individualized compound I	Fluazinam
5 B-108	one individualized compound I	Pyrifenoxy
10 B-109	one individualized compound I	3-[5-(4-Chloro-phenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine (Pyrisoxazole)
10 B-110	one individualized compound I	3-[5-(4-Methyl-phenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine
15 B-111	one individualized compound I	Bupirimate
15 B-112	one individualized compound I	Cyprodinil
15 B-113	one individualized compound I	5-Fluorocytosine
20 B-114	one individualized compound I	5-Fluoro-2-(p-tolylmethoxy)pyrimidin-4-amine
20 B-115	one individualized compound I	5-Fluoro-2-(4-fluorophenylmethoxy)pyrimidin-4-amine
20 B-116	one individualized compound I	Diflumetorim
25 B-117	one individualized compound I	(5,8-Difluoroquinazolin-4-yl)-{2-[2-fluoro-4-(4-trifluoromethylpyridin-2-yloxy)-phenyl]-ethyl}-amine
25 B-118	one individualized compound I	Fenarimol
25 B-119	one individualized compound I	Ferimzone
30 B-120	one individualized compound I	Mepanipyrim
30 B-121	one individualized compound I	Nitrapyrin
30 B-122	one individualized compound I	Nuarimol
30 B-123	one individualized compound I	Pyrimethanil
35 B-124	one individualized compound I	Triforine
35 B-125	one individualized compound I	Fenpiclonil
35 B-126	one individualized compound I	Fludioxonil
40 B-127	one individualized compound I	Aldimorph
40 B-128	one individualized compound I	Dodemorph
40 B-129	one individualized compound I	Dodemorph-acetate
45 B-130	one individualized compound I	Fenpropimorph
45 B-131	one individualized compound I	Tridemorph
45 B-132	one individualized compound I	Fenpropidin
50 B-133	one individualized compound I	Fluoroimid
50 B-134	one individualized compound I	Iprodione
50 B-135	one individualized compound I	Procymidone
55 B-136	one individualized compound I	Vinclozolin
55 B-137	one individualized compound I	Famoxadone
55 B-138	one individualized compound I	Fenamidone
55 B-139	one individualized compound I	Flutianil
55 B-140	one individualized compound I	Ocithilinone
55 B-141	one individualized compound I	Probenazole

(continued)

Mixture	Component 1	Component 2
B-142	one individualized compound I	Fenpyrazamine
B-143	one individualized compound I	Acibenzolar-S-methyl
B-144	one individualized compound I	Ametoctradin
B-145	one individualized compound I	Amisulbrom
B-146	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[(3-isobutyryloxymethoxy-4-methoxypyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-[1,5]dioxinan-7-yl] 2-methylpropanoate
B-147	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[(3-acetoxy-4-methoxypyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxinan-7-yl] 2-methylpropanoate
B-148	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[(3-(acetoxymethoxy)-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxinan-7-yl] 2-methylpropanoate
B-149	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[(3-isobutoxycarbonyloxy-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxinan-7-yl] 2-methylpropanoate
B-150	one individualized compound I	[(3S,6S,7R,8R)-8-benzyl-3-[[3-(1,3-benzodioxol-5-ylmethoxy)-4-methoxy-pyridine-2-carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxinan-7-yl] 2-methylpropanoate
B-151	one individualized compound I	Anilazin
B-152	one individualized compound I	Blasticidin-S
B-153	one individualized compound I	Captafol
B-154	one individualized compound I	Captan
B-155	one individualized compound I	Chinomethionat
B-156	one individualized compound I	Dazomet
B-157	one individualized compound I	Debacarb
B-158	one individualized compound I	Diclomezine
B-159	one individualized compound I	Difenoquat,
B-160	one individualized compound I	Difenoquat-methylsulfate
B-161	one individualized compound I	Fenoxanil
B-162	one individualized compound I	Folpet
B-163	one individualized compound I	Oxolinsäure
B-164	one individualized compound I	Piperalin
B-165	one individualized compound I	Proquinazid
B-166	one individualized compound I	Pyroquilon
B-167	one individualized compound I	Quinoxifen
B-168	one individualized compound I	Triazoxid
B-169	one individualized compound I	Tricyclazole
B-170	one individualized compound I	2-Butoxy-6-ido-3-propyl-chromen-4-one

(continued)

Mixture	Component 1	Component 2
5	B-171	one individualized compound I 5-Chloro-1-(4,6-dimethoxy-pyrimidin-2-yl)-2-methyl-1H-benzoimidazole
10	B-172	one individualized compound I 5-Chloro-7-(4-methyl-piperidin-1-yl)-6-(2,4,6-trifluorophenyl)-[1,2,4]tri-azolo[1,5-a]pyrimidine
15	B-173	one individualized compound I Ferbam
20	B-174	one individualized compound I Mancozeb
25	B-175	one individualized compound I Maneb
30	B-176	one individualized compound I Metam
35	B-177	one individualized compound I Methasulphocarb
40	B-178	one individualized compound I Metiram
45	B-179	one individualized compound I Propineb
50	B-180	one individualized compound I Thiram
55	B-181	one individualized compound I Zineb
	B-182	one individualized compound I Ziram
	B-183	one individualized compound I Diethofencarb
	B-184	one individualized compound I Benthiavalicarb
	B-185	one individualized compound I Iprovalicarb
	B-186	one individualized compound I Propamocarb
	B-187	one individualized compound I Propamocarb hydrochlorid
	B-188	one individualized compound I Valifenalate
	B-189	one individualized compound I N-(1-(1-(4-cyanophenyl)ethanesulfonyl)-but-2-yl) carbamic acid-(4-fluorophenyl) ester
	B-190	one individualized compound I Dodine
	B-191	one individualized compound I Dodine free base
	B-192	one individualized compound I Guazatine
	B-193	one individualized compound I Guazatine-acetate
	B-194	one individualized compound I Iminoctadine
	B-195	one individualized compound I Iminoctadine-triacetate
	B-196	one individualized compound I Iminoctadine-tris(albesilate)
	B-197	one individualized compound I Kasugamycin
	B-198	one individualized compound I Kasugamycin-hydrochloride-hydrate
	B-199	one individualized compound I Polyoxine
	B-200	one individualized compound I Streptomycin
	B-201	one individualized compound I Validamycin A
	B-202	one individualized compound I Binapacryl
	B-203	one individualized compound I Dicloran
	B-204	one individualized compound I Dinobuton
	B-205	one individualized compound I Dinocap
	B-206	one individualized compound I Nitrothal-isopropyl

(continued)

Mixture	Component 1	Component 2
B-207	one individualized compound I	Tecnazen
B-208	one individualized compound I	Fentin salts
B-209	one individualized compound I	Dithianon
B-210	one individualized compound I	Isoprothiolane
B-211	one individualized compound I	Edifenphos
B-212	one individualized compound I	Fosetyl, Fosetyl-aluminium
B-213	one individualized compound I	Iprobenfos
B-214	one individualized compound I	Phosphorous acid (H_3PO_3) and derivatives
B-215	one individualized compound I	Pyrazophos
B-216	one individualized compound I	Tolclofos-methyl
B-217	one individualized compound I	Chlorothalonil
B-218	one individualized compound I	Dichlofuanid
B-219	one individualized compound I	Dichlorophen
B-220	one individualized compound I	Flusulfamide
B-221	one individualized compound I	Hexachlorbenzene
B-222	one individualized compound I	Pencycuron
B-223	one individualized compound I	Pentachlorophenol and salts
B-224	one individualized compound I	Phthalide
B-225	one individualized compound I	Quintozone
B-226	one individualized compound I	Thiophanate Methyl
B-227	one individualized compound I	Tolylfluanid
B-228	one individualized compound I	N-(4-chloro-2-nitro-phenyl)-N-ethyl-4-methyl-benzenesulfonamide
B-229	one individualized compound I	Bordeaux mixture
B-230	one individualized compound I	Copper acetate
B-231	one individualized compound I	Copper hydroxide
B-232	one individualized compound I	Copper oxychloride
B-233	one individualized compound I	basic Copper sulfate
B-234	one individualized compound I	Sulfur
B-235	one individualized compound I	Biphenyl
B-236	one individualized compound I	Bronopol
B-237	one individualized compound I	Cyflufenamid
B-238	one individualized compound I	Cymoxanil
B-239	one individualized compound I	Diphenylamin
B-240	one individualized compound I	Metrafenone
B-241	one individualized compound I	Pyriofenone
B-242	one individualized compound I	Mildiomycin
B-243	one individualized compound I	Oxin-copper

(continued)

Mixture	Component 1	Component 2
5 B-244	one individualized compound I	Prohexadione calcium
B-245	one individualized compound I	Spiroxamine
B-246	one individualized compound I	Tebuflouquin
B-247	one individualized compound I	Tolylfluanid
10 B-248	one individualized compound I	N-(Cyclopropylmethoxyimino-(6-difluoromethoxy-2,3-difluoro-phenyl)-methyl)-2-phenyl acetamide
B-249	one individualized compound I	N'-(4-(4-chloro-3-trifluoromethyl-phenoxy)-2,5-dimethyl-phenyl)-N-ethyl-N-methyl formamidine
15 B-250	one individualized compound I	N'-(4-(4-fluoro-3-trifluoromethyl-phenoxy)-2,5-dimethyl-phenyl)-N-ethyl-N-methyl formamidine
B-251	one individualized compound I	N'-(2-methyl-5-trifluoromethyl-4-(3-trimethylsilylpropoxy)-phenyl)-N-ethyl-N-methyl formamidine
20 B-252	one individualized compound I	N'-(5-difluoromethyl-2-methyl-4-(3-trimethylsilylpropoxy)-phenyl)-N-ethyl-N-methyl formamidine
B-253	one individualized compound I	2-[1-[2-(5-Methyl-3-trifluoromethyl-pyrazole-1-yl)-acetyl]-piperidin-4-yl]-thiazole-4-carboxylic acid methyl-(1,2,3,4-tetrahydro-naphthalen-1-yl)-amide
25 B-254	one individualized compound I	2-[1-[2-(5-Methyl-3-trifluoromethyl-pyrazole-1-yl)-acetyl]-piperidin-4-yl]-thiazole-4-carboxylic acid methyl-(R)-1,2,3,4-tetrahydro-naphthalen-1-yl-amide
30 B-255	one individualized compound I	1-[4-[4-[5-(2,6-difluorophenyl)-4,5-di-hydro-3-isoxazolyl]-2-thiazolyl]-1-piperidinyl]-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone
B-256	one individualized compound I	Methoxy-acetic acid 6-tert-butyl-8-fluoro-2,3-dimethyl-quinolin-4-yl ester
35 B-257	one individualized compound I	N-Methyl-2-[1-[(5-methyl-3-trifluoro-methyl-1H-pyrazol-1-yl)-acetyl]-piperidin-4-yl]-N-[(1R)-1,2,3,4-tetrahydro-naphthalen-1-yl]-4-thiazolecarboxamide
B-258	one individualized compound I	<i>Bacillus subtilis</i> NRRL No. B-21661
40 B-259	one individualized compound I	<i>Bacillus pumilus</i> N RRL No. B-30087
B-260	one individualized compound I	<i>Ulocladium oudemansii</i>
B-261	one individualized compound I	Carbaryl
45 B-262	one individualized compound I	Carbofuran
B-263	one individualized compound I	Carbosulfan
B-264	one individualized compound I	Methomylthiodicarb
B-265	one individualized compound I	Bifenthrin
50 B-266	one individualized compound I	Cyfluthrin
B-267	one individualized compound I	Cypermethrin
B-268	one individualized compound I	alpha-Cypermethrin
B-269	one individualized compound I	zeta-Cypermethrin
B-270	one individualized compound I	Deltamethrin
55 B-271	one individualized compound I	Esfenvalerate

(continued)

Mixture	Component 1	Component 2
B-272	one individualized compound I	Lambda-cyhalothrin
B-273	one individualized compound I	Permethrin
B-274	one individualized compound I	Tefluthrin
B-275	one individualized compound I	Diflubenzuron
B-276	one individualized compound I	Flufenoxuron
B-277	one individualized compound I	Lufenuron
B-278	one individualized compound I	Teflubenzuron
B-279	one individualized compound I	Spirotetramate
B-280	one individualized compound I	Clothianidin
B-281	one individualized compound I	Dinotefuran
B-282	one individualized compound I	Imidacloprid
B-283	one individualized compound I	Thiamethoxam
B-284	one individualized compound I	Flupyradifurone
B-285	one individualized compound I	Acetamiprid
B-286	one individualized compound I	Thiacloprid
B-287	one individualized compound I	Endosulfan
B-288	one individualized compound I	Fipronil
B-289	one individualized compound I	Abamectin
B-290	one individualized compound I	Emamectin
B-291	one individualized compound I	Spinosad
B-292	one individualized compound I	Spinetoram
B-293	one individualized compound I	Hydramethylnon
B-294	one individualized compound I	Chlorfenapyr
B-295	one individualized compound I	Fenbutatin oxide
B-296	one individualized compound I	Indoxacarb
B-297	one individualized compound I	Metaflumizone
B-298	one individualized compound I	Flonicamid
B-299	one individualized compound I	Lubendiamide
B-300	one individualized compound I	Chlorantraniliprole
B-301	one individualized compound I	Cyazypyrr (HGW86)
B-302	one individualized compound I	Cyflumetofen
B-303	one individualized compound I	Acetochlor
B-304	one individualized compound I	Dimethenamid
B-305	one individualized compound I	metolachlor
B-306	one individualized compound I	Metazachlor
B-307	one individualized compound I	Glyphosate
B-308	one individualized compound I	Glufosinate
B-309	one individualized compound I	Sulfosate

(continued)

Mixture	Component 1	Component 2
B-310	one individualized compound I	Clodinafop
B-311	one individualized compound I	Fenoxyprop
B-312	one individualized compound I	Fluazifop
B-313	one individualized compound I	Haloxifop
B-314	one individualized compound I	Paraquat
B-315	one individualized compound I	Phenmedipham
B-316	one individualized compound I	Clethodim
B-317	one individualized compound I	Cycloxydim
B-318	one individualized compound I	Profoxydim
B-319	one individualized compound I	Sethoxydim
B-320	one individualized compound I	Tepraloxydim
B-321	one individualized compound I	Pendimethalin
B-322	one individualized compound I	Prodiamine
B-323	one individualized compound I	Trifluralin
B-324	one individualized compound I	Acifluorfen
B-325	one individualized compound I	Bromoxynil
B-326	one individualized compound I	Imazamethabenz
B-327	one individualized compound I	Imazamox
B-328	one individualized compound I	Imazapic
B-329	one individualized compound I	Imazapyr
B-330	one individualized compound I	Imazaquin
B-331	one individualized compound I	Imazethapyr
B-332	one individualized compound I	2,4-Dichlorophenoxyacetic acid (2,4-D)
B-333	one individualized compound I	Chloridazon
B-334	one individualized compound I	Clopyralid
B-335	one individualized compound I	Fluroxypyr
B-336	one individualized compound I	Picloram
B-337	one individualized compound I	Picolinafen
B-338	one individualized compound I	Bensulfuron
B-339	one individualized compound I	Chlorimuron-ethyl
B-340	one individualized compound I	Cyclosulfamuron
B-341	one individualized compound I	Iodosulfuron
B-342	one individualized compound I	Mesosulfuron
B-343	one individualized compound I	Metsulfuron-methyl
B-344	one individualized compound I	Nicosulfuron
B-345	one individualized compound I	Rimsulfuron
B-346	one individualized compound I	Triflusulfuron
B-347	one individualized compound I	Atrazine

(continued)

Mixture	Component 1	Component 2
B-348	one individualized compound I	Hexazinone
B-349	one individualized compound I	Diuron
B-350	one individualized compound I	Florasulam
B-351	one individualized compound I	Pyroxasulfone
B-352	one individualized compound I	Bentazone
B-353	one individualized compound I	Cinidon-ethyl
B-354	one individualized compound I	Cinmethylin
B-355	one individualized compound I	Dicamba
B-356	one individualized compound I	Diflufenzopyr
B-357	one individualized compound I	Quinclorac
B-358	one individualized compound I	Quinmerac
B-359	one individualized compound I	Mesotrione
B-360	one individualized compound I	Saflufenacil
B-361	one individualized compound I	Topramezone

[0114] The active substances referred to as component 2, their preparation and their activity against harmful fungi is known (cf.: <http://www.alanwood.net/pesticides/>); these substances are commercially available. The compounds described by IUPAC nomenclature, their preparation and their fungicidal activity are also known (cf. Can. J. Plant Sci. 48 (6), 587-94, 1968; EP-A 141 317; EP-A 152 031; EP-A 226 917; EP-A 243 970; E P-A 256 503; E P-A 428 941; E P-A 532 022; EP-A 1 028 125; EP-A 1 035 122; EP-A 1 201 648; EP-A 1 122 244, JP 2002316902; DE 19650197; DE 10021412; DE 102005009458; US 3,296,272; US 3,325,503; WO 98/46608; WO 99/14187; WO 99/24413; WO 99/27783; WO 00/29404; WO 00/46148; WO 00/65913; WO 01/54501; WO 01/56358; WO 02/22583; WO 02/40431; WO 03/10149; WO 03/11853; WO 03/14103; WO 03/16286; WO 03/53145; WO 03/61388; WO 03/66609; WO 03/74491; WO 04/49804; WO 04/83193; WO 05/120234; WO 05/123689; WO 05/123690; WO 05/63721; WO 05/87772; WO 05/87773; WO 06/15866; WO 06/87325; WO 06/87343; WO 07/82098; WO 07/90624, WO 11/028657).


[0115] The mixtures of active substances can be prepared as compositions comprising besides the active ingredients at least one inert ingredient by usual means, e. g. by the means given for the compositions of compounds I.

[0116] Concerning usual ingredients of such compositions reference is made to the explanations given for the compositions containing compounds I.

[0117] The mixtures of active substances according to the present invention are suitable as fungicides, as are the compounds of formula I. They are distinguished by an outstanding effectiveness against a broad spectrum of phytopathogenic fungi, especially from the classes of the Ascomycetes, Basidiomycetes, Deuteromycetes and Peronosporomycetes (syn. Oomycetes). In addition, it is referred to the explanations regarding the fungicidal activity of the compounds and the compositions containing compounds I, respectively.

Claims

1. Compounds of formula I



wherein:

5 X^1, X^2 independently of each other are selected from halogen;

R^1 is hydrogen, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkyl- C_1 - C_4 -alkyl, phenyl, phenyl- C_1 - C_4 -alkyl, phenyl- C_2 - C_4 -alkenyl or phenyl- C_2 - C_4 -alkynyl;

R^2 is n-butyl, sec.-butyl or tert.-butyl;

wherein the aliphatic moieties of R^1 and/or R^2 may carry 1, 2, 3 or up to the maximum possible number of identical or different groups R^a which independently of one another are selected from:

10 R^a halogen, CN, nitro, C_1 - C_4 -alkoxy and C_1 - C_4 -halogenalkoxy;

wherein the cycloalkyl and/or phenyl moieties of R^1 may carry 1, 2, 3, 4, 5 or up to the maximum number of identical or different groups R^b which independently of one another are selected from:

15 R^b halogen, CN, nitro, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -halogenalkyl and C_1 - C_4 -halogenalkoxy;

and the N-oxides and the agriculturally acceptable salts thereof.

2. The compounds according to claim 1, wherein X^1 is Cl.

20 3. The compounds according to any of the claims 1 and 2, wherein X^2 is Cl.



4. The compounds according to any of the claims 1 to 3, wherein R^1 is hydrogen or C_1 - C_4 -alkyl.

25 5. The compounds according to any of the claims 1 to 4, wherein R^2 is tert.-butyl.


6. The compounds according to any of the claims 1 to 4, wherein R^2 is n-butyl.

7. The compounds according to any of the claims 1 to 6, wherein R^2 is unsubstituted.

30 8. A process for preparing compounds of formula I as defined in any of claims 1 to 7, which comprises reacting a compound of formula III




50 wherein X^1 is defined as in any of the claims 1 to 2, under basic conditions; and reacting the resulting compound of formula IV




wherein X^1 and X^2 are defined as in any of the claims 1 to 3,

in the presence of a catalyst with isopropylmagnesium bromide followed by a reaction with acetyl chloride; and halogenating the resulting compound of formula V




10 ,

wherein X¹ and X² are defined as in any of the claims 1 to 3, with bromine; and reacting the resulting compound of formula VI




20 ,

wherein X¹ and X² are defined as in any of the claims 1 to 3, under basic conditions with 1H-1,2,4-triazole; and reacting the resulting compound of formula VII



30 ,

wherein X¹ and X² are defined as in any of the claims 1 to 3, with R¹-M, wherein R¹ is as defined in claims 1 or 4 and M is MgBr, MgCl, Li or Na, and reacting the resulting compound of VIII



45 ,

wherein X¹, X² and R¹ are defined as in any of the claims 1 to 4, under basic conditions with R²-LG, wherein R² is defined as in any of the claims 1, 5, 6 or 7 and LG is a nucleophilically replaceable leaving group, to obtain compounds of formula I.


50

9. A process for preparing compounds of formula I as defined in any of claims 1 to 7, which comprises reacting a compound of formula III




,


wherein X^2 is defined as in any of the claims 1 or 3 and Y is F or Cl,
 in presence of a catalyst with isopropylmagnesium halide followed by a reaction with a compound of formula IX
 $R^1\text{-COCl}$,
 wherein R^1 is as defined in claims 1 or 4;
 and converting the resulting compound of formula X




15 wherein X^2 is defined as in any of the claims 1 or 3, Y is F or Cl and R^1 is as defined in claims 1 or 4,
 under basic conditions with a halo-phenole of formula II



25 wherein X^1 is defined as in any of the claims 1 to 2;
 and reacting the resulting compound of formula Va



35 wherein X^1 , X^2 and R^1 are defined as in any of the claims 1 to 4, with trimethylsulf(ox)onium halide;
 and reacting the resulting compound of formula XI




45 wherein X^1 , X^2 and R^1 are defined as in any of the claims 1 to 4, under basic conditions with 1H-1,2,4-triazole;
 and reacting the resulting compound of formula VIII



55 wherein X^1 , X^2 and R^1 are defined as in any of the claims 1 to 4,
 under basic conditions with $R^2\text{-LG}$, wherein R^2 is defined as in any of the claims 1, 5, 6 or 7 and LG is a nucleophilically


replaceable leaving group,
to obtain compounds of formula I.

5 10. A process for preparing compounds of formula I as defined in any of claims 1 to 7, which comprises reacting a compound of formula XI




,

15 wherein X¹, X² and R¹ are defined as in any of the claims 1 to 4,
under acidic conditions with R²-OH, wherein R² is as defined in any of the claims 1, 5, 6 or 7;
and reacting the resulting compound of formula XII



,


25 wherein X¹, X², R¹ and R² are defined as in any of the claims 1 to 7,
with a halogenating agent or sulfonating agent as defined herein;
and reacting the resulting compound of formula XIII



,

35 40 wherein X¹, X², R¹ and R² are defined as in any of the claims 1 to 7 and LG is a nucleophilically replaceable leaving group with 1H-1,2,4-triazole,
to obtain compounds I.

45 11. Compounds of formula XII



,

55 wherein X¹, X², R¹ and R² are defined as in any of the claims 1 to 7.

12. Agrochemical compositions wherein said composition comprise an auxiliary and at least one compound of formula

I, as defined in any of the claims 1 to 7, an N-oxide or an agriculturally acceptable salt thereof.

13. The compositions according to claim 12, comprising additionally a further active substance.

5 14. Use of compounds of formula I, the N-oxides and the agriculturally acceptable salts thereof, as defined in any of the claims 1 to 7, or of the compositions as defined in any of the claims 12 to 13, for combating phytopathogenic fungi.

15. Seed coated with at least one compound of formula I as defined in any of the claims 1 to 7 or with a composition as defined in any of the claims 12 to 13, in an amount of from 0.1 g to 10 kg per 100 kg of seed.

10

15

20

25

30

35

40

45

50

55



EUROPEAN SEARCH REPORT

Application Number
EP 11 17 7556

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (IPC)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	DE 38 01 233 A1 (CIBA GEIGY AG [CH]) 4 August 1988 (1988-08-04)	1-15	INV. C07D249/08
Y	* claim 1; table 1; compounds 23,25,47,48,50,57 *	1-15	A01N43/653
X,D	----- EP 0 126 430 A2 (CIBA GEIGY AG [CH]) 28 November 1984 (1984-11-28) * page 50; table 1; compound 118 *	1-15	-----
Y	EP 0 354 183 A2 (CIBA GEIGY AG [CH]) 7 February 1990 (1990-02-07) * compound Formula (II) *	1-15	-----
			TECHNICAL FIELDS SEARCHED (IPC)
			C07D A01N
The present search report has been drawn up for all claims			
1	Place of search	Date of completion of the search	Examiner
	Munich	28 October 2011	Timmermans, Michel
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 11 17 7556

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

28-10-2011

Patent document cited in search report		Publication date		Patent family member(s)		Publication date	
DE 3801233	A1	04-08-1988		NONE			
EP 0126430	A2	28-11-1984		AU 607993 B2 AU 1550488 A AU 572145 B2 AU 2835884 A BR 8402388 A CA 1223264 A1 CS 247179 B2 DD 226882 A5 DD 248274 A5 DE 3484968 D1 DK 248184 A ES 8600635 A1 GB 2143815 A GB 2179659 A GR 79954 A1 HU 196693 B IL 71862 A JP 1705328 C JP 3061666 B JP 59222434 A JP 1843250 C JP 2096544 A JP 5052819 B JP 1735883 C JP 2085265 A JP 4020912 B MX 5769 A NZ 208209 A PH 20959 A PH 21667 A PH 22876 A PT 78602 A TR 22379 A US 4945100 A ZA 8403774 A		21-03-1991 21-07-1988 05-05-1988 22-11-1984 02-04-1985 23-06-1987 18-12-1986 04-09-1985 05-08-1987 02-10-1991 20-11-1984 16-01-1986 20-02-1985 11-03-1987 31-10-1984 30-01-1989 31-08-1988 27-10-1992 20-09-1991 14-12-1984 12-05-1994 09-04-1990 06-08-1993 26-02-1993 26-03-1990 07-04-1992 01-10-1993 08-01-1988 10-06-1987 13-01-1988 19-01-1989 01-06-1984 11-03-1987 31-07-1990 30-01-1985	
EP 0354183	A2	07-02-1990		AU 611315 B2 AU 3928489 A BR 8903917 A DE 58903329 D1 DK 381489 A JP 2083304 A NZ 230176 A ZA 8905927 A		06-06-1991 08-02-1990 27-03-1990 04-03-1993 05-02-1990 23-03-1990 29-01-1992 25-07-1990	

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 0126430 A2 [0002] [0010]
- US 4940720 A [0002]
- EP 0275955 A1 [0010]
- DE 4003180 A1 [0010]
- EP 0113640 A2 [0010]
- WO 0364572 A [0016]
- WO 02015701 A [0055]
- EP 374753 A [0055]
- WO 93007278 A [0055]
- WO 9534656 A [0055]
- EP 427529 A [0055]
- EP 451878 A [0055]
- WO 0318810 A [0055]
- WO 0352073 A [0055]
- WO 03018810 A [0055]
- EP 392225 A [0056]
- EP 141317 A [0114]
- EP 152031 A [0114]
- EP 226917 A [0114]
- EP 243970 A [0114]
- EP 256503 A [0114]
- EP 428941 A [0114]
- EP 532022 A [0114]
- EP 1028125 A [0114]
- EP 1035122 A [0114]
- EP 1201648 A [0114]
- EP 1122244 A [0114]
- JP 2002316902 B [0114]
- DE 19650197 [0114]
- DE 10021412 [0114]
- DE 102005009458 [0114]
- US 3296272 A [0114]
- US 3325503 A [0114]
- WO 9846608 A [0114]
- WO 9914187 A [0114]
- WO 9924413 A [0114]
- WO 9927783 A [0114]
- WO 0029404 A [0114]
- WO 0046148 A [0114]
- WO 0065913 A [0114]
- WO 0154501 A [0114]
- WO 0156358 A [0114]
- WO 0222583 A [0114]
- WO 0240431 A [0114]
- WO 0310149 A [0114]
- WO 0311853 A [0114]
- WO 0314103 A [0114]
- WO 0316286 A [0114]
- WO 0353145 A [0114]
- WO 0361388 A [0114]
- WO 0366609 A [0114]
- WO 0374491 A [0114]
- WO 0449804 A [0114]
- WO 0483193 A [0114]
- WO 05120234 A [0114]
- WO 05123689 A [0114]
- WO 05123690 A [0114]
- WO 0563721 A [0114]
- WO 0587772 A [0114]
- WO 0587773 A [0114]
- WO 0615866 A [0114]
- WO 0687325 A [0114]
- WO 0687343 A [0114]
- WO 0782098 A [0114]
- WO 0790624 A [0114]
- WO 11028657 A [0114]

Non-patent literature cited in the description

- *J.Agric. Food Chem.*, 2009, vol. 57, 4854-4860 [0010]
- *J. Med. Chem.*, 1995, vol. 38 (11), 1892-903 [0016]
- *J. Heterocyc. Chem.*, 1981, vol. 18(7), 1305-8 [0016]
- *J. Am. Chem. Soc.*, 2001, vol. 123 (25), 5962-5973 [0016]
- *Pest Managem. Sci.*, 2005, vol. 61, 246 [0054]
- *PEST. MANAGEM. SCI.*, 2005, vol. 61, 258 [0054]
- *PEST. MANAGEM. SCI.*, 2005, vol. 61, 277 [0054]
- *PEST. MANAGEM. SCI.*, 2005, vol. 61, 269 [0054]
- *PEST. MANAGEM. SCI.*, 2005, vol. 61, 286 [0054]
- *PEST. MANAGEM. SCI.*, 2008, vol. 64, 326 [0054]
- *PEST. MANAGEM. SCI.*, 2008, vol. 64, 332 [0054]
- *Weed Sci.*, 2009, vol. 57, 108 [0054]
- *Austral. J. Agricult. Res.*, 2007, vol. 58, 708 [0054]
- *Science*, 2007, vol. 316, 1185 [0054]
- Catalogue of pesticide formulation types and international coding system. Technical Monograph No. 2. Crop-Life International, May 2008 [0069]
- **MOLLET ; GRUBEMANN**. Formulation technology. Wiley VCH, 2001 [0070]
- New developments in crop protection product formulation. **KNOWLES**. Agrow Reports DS243. T&F Informa, 2005 [0070]

EP 2 559 688 A1

- McCutcheon's, Vol.1: Emulsifiers & Detergents. McCutcheon's Directories, 2008, vol. 1 **[0074]**
- **KNOWLES.** Adjuvants and additives, Agrow Reports DS256. T&F Informa UK, 2006 **[0078]**
- *Can. J. Plant Sci.*, 1968, vol. 48 (6), 587-94 **[0114]**