



(11)

EP 2 578 683 B9

(12)

CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:

Corrected version no 1 (W1 B1)**Corrections, see****Description Paragraph(s) 122, 133, 282**

(51) Int Cl.:

C12N 15/115 (2010.01)**C12Q 1/68 (2006.01)****A61K 31/7088 (2006.01)**

(48) Corrigendum issued on:

15.04.2015 Bulletin 2015/16(45) Date of publication and mention
of the grant of the patent:**26.11.2014 Bulletin 2014/48**(21) Application number: **13150168.6**(22) Date of filing: **14.02.2005**(54) **Aptamer therapeutics useful in the treatment of complement-related disorders**

Aptamer-Therapeutika zur Behandlung von mit dem Komplementsystem in Verbindung stehenden Erkrankungen

Agents thérapeutiques à base d'aptamères utiles dans le traitement de troubles liés au complément

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR**(30) Priority: **12.02.2004 US 544542 P****25.02.2004 US 547747 P****21.06.2004 US 581685 P****07.09.2004 US 608048 P**

(43) Date of publication of application:

10.04.2013 Bulletin 2013/15

(60) Divisional application:

14194459.5 / 2 860 251(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:**05713496.7 / 1 727 567**(73) Proprietor: **Archemix LLC****San Francisco, CA 94111 (US)**

(72) Inventors:

- Benedict, Claude**
Cambridge, MA 02140 (US)

- Diener, John**
Cambridge, MA 02139 (US)
- Epstein, David**
Huntington, NY 11743 (US)
- Grate, Dilara**
Waltham, MA 02453 (US)
- Keene, Sara Chesworth**
Westford, MA 01886 (US)
- Kurz, Jeffrey**
Arlington, MA 02474 (US)
- Kurz, Markus**
Newton, MA 02465 (US)
- McCauley, Thomas Green**
Cambridge, MA 02141 (US)
- Rottman, James**
Sudbury, MA 01776 (US)
- Thompson, Kristin**
Arlington, MA 02474 (US)
- Wilson, Charles**
Concord, MA 01742 (US)
- Zoltoski, Anna J.**
325 South Boston, MA 02127 (US)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

(74) Representative: **Clegg, Richard Ian**
Mewburn Ellis LLP
33 Gutter Lane
London
EC2V 8AS (GB)

(56) References cited:
WO-A1-99/41271

- **WATSON S R ET AL:** "Anti-L-selectin aptamers: binding characteristics, pharmacokinetic parameters, and activity against an intravascular target *in vivo*", **ANTISENSE & NUCLEIC ACID DRUG DEVELOPMENT**, MARY ANN LIEBERT, INC., NEW YORK, US, vol. 10, no. 2, 1 April 2000 (2000-04-01), pages 63-75, XP002533820, ISSN: 1087-2906
- **BELL C ET AL:** "Oligonucleotide NX1838 inhibits VEGF165-mediated cellular responses *in vitro*", **IN VITRO CELLULAR & DEVELOPMENTAL BIOLOGY -ANIMAL**, SPRINGER, GERMANY, vol. 35, no. 9, 1 October 1999 (1999-10-01), pages 533-542, XP002979398, ISSN: 1071-2690
- **VOROBEJV P E ET AL:** "Nuclease resistance and RNase H sensitivity of oligonucleotides bridged by oligomethylenediol and oligoethylene glycol linkers", **ANTISENSE & NUCLEIC ACID DRUG DEVELOPMENT**, MARY ANN LIEBERT, INC., NEW YORK, US LNKD- DOI: 10.1089/108729001750171290, vol. 11, 1 April 2001 (2001-04-01), pages 77-85, XP002226341, ISSN: 1087-2906
- **BIESECKER G ET AL:** "Derivation of RNA aptamer inhibitors of human complement C5", **IMMUNOPHARMACOLOGY**, ELSEVIER SCIENCE PUBLISHERS BV, XX LNKD- DOI: 10.1016/S0162-3109(99)00020-X, vol. 42, no. 1-3, 1 May 1999 (1999-05-01), pages 219-230, XP002377259, ISSN: 0162-3109

- **BURCOVICH B ET AL:** "Branched polyethylene glycol (bPEG) conjugated antisense oligonucleotides", **NUCLEOSIDES & NUCLEOTIDES**, MARCEL DEKKER, INC, US, vol. 17, no. 9-11, 1 September 1998 (1998-09-01), pages 1567-1570, XP008082550, ISSN: 0732-8311
- **BRODY E N ET AL:** "APTAMERS AS THERAPEUTIC AND DIAGNOSTIC AGENTS", **JOURNAL OF BIOTECHNOLOGY**, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 74, no. 1, 1 March 2000 (2000-03-01), pages 5-13, XP009031105, ISSN: 0168-1656
- **RIMMELE M:** "NUCLEIC ACID APTAMERS AS TOOLS AND DRUGS: RECENT DEVELOPMENTS", **CHEMBIOCHEM - A EUROPEAN JOURNAL OF CHEMICAL BIOLOGY**, WILEY VCH, WEINHEIM, DE LNKD- DOI: 10.1002/CBIC.200300648, vol. 4, no. 10, 1 January 2003 (2003-01-01), pages 963-971, XP001176680, ISSN: 1439-4227
- **OSTENDORF T ET AL:** "VEGF165 mediates glomerular endothelial repair", **JOURNAL OF CLINICAL INVESTIGATION**, AMERICAN SOCIETY FOR CLINICAL INVESTIGATION, US, vol. 104, no. 7, 1 October 1999 (1999-10-01), pages 913-923, XP002238360, ISSN: 0021-9738, DOI: 10.1172/JCI6740
- **BONORA GIAN MARIA:** "POLYMER-CONJUGATED BIOACTIVE OLIGONUCLEOTIDES", **JOURNAL OF BIOACTIVE AND COMPATIBLE POLYMERS**, LANCASTER, PA, US, vol. 17, no. 5, 1 September 2002 (2002-09-01), pages 375-389, XP008078479, DOI: 10.1177/0883911502017005555

Description**FIELD OF THE INVENTION**

5 [0001] The invention relates generally to the field of nucleic acids and more particularly to aptamers capable of binding to the C5 protein of the complement system, useful as therapeutics in and diagnostics in complement-related cardiac, inflammatory, and auto-immune disorders, ischemic reperfusion injury and/or other diseases or disorders in which C5 mediated complement activation has been implicated. The invention further relates to materials and methods for the administration of aptamers capable of binding to the C5 complement system protein.

10

BACKGROUND OF THE INVENTION

15 [0002] Aptamers are nucleic acid molecules having specific binding affinity to molecules through interactions other than classic Watson-Crick base pairing.

15

[0003] Aptamers, like peptides generated by phage display or monoclonal antibodies ("MAbs"), are capable of specifically binding to selected targets and modulating the target's activity, e.g., through binding aptamers may block their target's ability to function. Created by an *in vitro* selection process from pools of random sequence oligonucleotides, aptamers have been generated for over 100 proteins including growth factors, transcription factors, enzymes, immunoglobulins, and receptors. A typical aptamer is 10-15 kDa in size (30-45 nucleotides), binds its target with sub-nanomolar affinity, and discriminates against closely related targets (e.g., aptamers will typically not bind other proteins from the same gene family). A series of structural studies have shown that aptamers are capable of using the same types of binding interactions (e.g., hydrogen bonding, electrostatic complementarity, hydrophobic contacts, steric exclusion) that drive affinity and specificity in antibody-antigen complexes.

20

20 [0004] Aptamers have a number of desirable characteristics for use as therapeutics and diagnostics including high specificity and affinity, biological efficacy, and excellent pharmacokinetic properties. In addition, they offer specific competitive advantages over antibodies and other protein biologics, for example:

30

1) Speed and control. Aptamers are produced by an entirely *in vitro* process, allowing for the rapid generation of initial leads, including therapeutic leads. *In vitro* selection allows the specificity and affinity of the aptamer to be tightly controlled and allows the generation of leads, including leads against both toxic and non-immunogenic targets.

35

2) Toxicity and Immunogenicity. Aptamers as a class have demonstrated little or no toxicity or immunogenicity. In chronic dosing of rats or woodchucks with high levels of aptamer (10 mg/kg daily for 90 days), no toxicity is observed by any clinical, cellular, or biochemical measure. Whereas the efficacy of many monoclonal antibodies can be severely limited by immune response to antibodies themselves, it is extremely difficult to elicit antibodies to aptamers most likely because aptamers cannot be presented by T-cells via the MHC and the immune response is generally trained not to recognize nucleic acid fragments.

40

3) Administration. Whereas most currently approved antibody therapeutics are administered by intravenous infusion (typically over 2-4 hours), aptamers can be administered by subcutaneous injection (aptamer bioavailability via subcutaneous administration is >80% in monkey studies (Tucker et al., J. Chromatography B. 732: 203-212, 1999)). This difference is primarily due to the comparatively low solubility and thus large volumes necessary for most therapeutic MAbs. With good solubility (>150 mg/mL) and comparatively low molecular weight (aptamer: 10-50 kDa; antibody: 150 kDa), a weekly dose of aptamer may be delivered by injection in a volume of less than 0.5 mL. In addition, the small size of aptamers allows them to penetrate into areas of conformational constrictions that do not allow for antibodies or antibody fragments to penetrate, presenting yet another advantage of aptamer-based therapeutics or prophylaxis.

45

4) Scalability and cost. Therapeutic aptamers are chemically synthesized and consequently can be readily scaled as needed to meet production demand. Whereas difficulties in scaling production are currently limiting the availability of some biologics and the capital cost of a large-scale protein production plant is enormous, a single large-scale oligonucleotide synthesizer can produce upwards of 100 kg/year and requires a relatively modest initial investment. The current cost of goods for aptamer synthesis at the kilogram scale is estimated at \$500/g, comparable to that for highly optimized antibodies. Continuing improvements in process development are expected to lower the cost of goods to < \$100/g in five years.

55

5) Stability. Therapeutic aptamers are chemically robust. They are intrinsically adapted to regain activity following exposure to factors such as heat and denaturants and can be stored for extended periods (>1 yr) at room temperature

as lyophilized powders.

The Complement System

5 [0005] The complement system comprises a set of at least 20 plasma and membrane proteins that act together in a regulated cascade system to attack extracellular forms of pathogens (e.g., bacterium). The complement system includes two distinct enzymatic activation cascades, the classical and alternative pathways (Figure 1), and a non-enzymatic pathway known as the membrane attack pathway.

10 [0006] The first enzymatically activated cascade, known as the classical pathway, comprises several components, C1, C4, C2, C3 and C5 (listed by order in the pathway). Initiation of the classical pathway of the complement system occurs following binding and activation of the first complement component (C1) by both immune and non-immune activators. C1 comprises a calcium-dependent complex of components C1q, C1r and C1s, and is activated through binding of the C1q component. C1q contains six identical subunits and each subunit comprises three chains (the A, B and C chains). Each chain has a globular head region that is connected to a collagen-like tail. Binding and activation of C1q by antigen-antibody complexes occurs through the C1q head group region. Numerous non-antibody C1q activators, including proteins, lipids and nucleic acids, bind and activate C1q through a distinct site on the collagen-like stalk region. The C1qrs complex then catalyzes the activation of complement components C4 and C2, forming the C4bC2a complex which functions as a C3 convertase.

15 [0007] The second enzymatically activated cascade, known as the alternative pathway, is a rapid, antibody-independent route for complement system activation and amplification. The alternative pathway comprises several components, C3, Factor B, and Factor D (listed by order in the pathway). Activation of the alternative pathway occurs when C3b, a proteolytic cleavage form of C3, is bound to an activating surface agent such as a bacterium. Factor B is then bound to C3b, and cleaved by Factor D to yield the active enzyme, Ba. The enzyme Ba then cleaves more C3 to generate more C3b, producing extensive deposition of C3b-Ba complexes on the activating surface.

20 [0008] Thus, both the classical and alternate complement pathways produce C3 convertases that split factor C3 into C3a and C3b. At this point, both C3 convertases further assemble into C5 convertases (C4b2a3b and C3b3bBb). These complexes subsequently cleave complement component C5 into two components: the C5a polypeptide (9 kDa) and the C5b polypeptide (170 kDa). The C5a polypeptide binds to a 7 transmembrane G-protein coupled receptor, which was originally associated with leukocytes and is now known to be expressed on a variety of tissues including hepatocytes and neurons. The C5a molecule is the primary chemotactic component of the human complement system and can trigger a variety of biological responses including leukocyte chemotaxis, smooth muscle contraction, activation of intracellular signal transduction pathways, neutrophil-endothelial adhesion, cytokine and lipid mediator release and oxidant formation.

25 [0009] The larger C5b fragment binds sequentially to later components of the complement cascade, C6, C7, C8 and C9 to form the C5b-9 membrane attack complex ("MAC"). The C5b-9 MAC can directly lyse erythrocytes, and in greater quantities, it is lytic for leukocytes and damaging to tissues such as muscle, epithelial and endothelial cells. In sublytic amounts, the MAC can stimulate upregulation of adhesion molecules, intracellular calcium increase and cytokine release. In addition, the C5b-9 MAC can stimulate cells such as endothelial cells and platelets without causing cell lysis. The non-lytic effects of C5a and the C5b-9 MAC are sometimes quite similar.

30 [0010] Although the complement system has an important role in the maintenance of health, it has the potential to cause or contribute to disease. For example, the complement system has been implicated in side effects relating to coronary artery bypass graft ("CABG") surgery, numerous renal, rheumatological, neurological, dermatological, hematological, vascular/pulmonary, allergy, infectious, and biocompatibility/shock diseases and/or conditions, and diabetic retinopathy. The complement system is not necessarily the only cause of a disease state, but it may be one of several factors that contribute to pathogenesis.

35 [0011] In Fitch et al., Circ. 100:2499-506 (1999), the effects of the anti-C5 single-chain antibody fragment Pexelizumab on patients undergoing coronary artery bypass graft surgery with cardiopulmonary bypass ("CPB") was tested. Individual patients were administered Pexelizumab in a 10 minute, single-bolus dose just prior to CPB at 0.5 mg/kg, 1.0 mg/kg and 2.0 mg/kg. Blood samples were removed and tested for complement activity at pre -dose, 5 min post-dose, after 5 min at 28°C, after initiation of rewarming, after 5 min at 37°C, and up to 7 days after CPB. Pharmacodynamic analysis demonstrated significant dose-dependent inhibition of complement hemolytic activity for up to 14 hours at a dosage of 2 mg/kg, and the generation of proinflammatory complement byproducts (sC5b-9) was effectively inhibited in a dose-dependent fashion. As previously mentioned, however, antibody therapeutics have certain limitations.

40 [0012] Accordingly, it would be beneficial to have novel inhibitors of the complement system for use as therapeutics and diagnostics in the treatment of complement-related disorders.

45 [0013] WO 99/41271 A1 describes methods for identifying and preparing high-affinity Nucleic Acid Ligands to Complement System Proteins C1q, C3 and C5, including RNA ligands identified by the SELEX method.

50 [0014] Watson et al., Antisense & Nucleic Acid Drug Development 10:63-75 (2000) describes *in vitro* binding characteristics, pharmacokinetic parameters and inhibition of lymphocyte trafficking in a SCID mouse/human lymphocyte model

for 2'F pyrimidine 2'OH purine RNA and ssDNA anti-human L-selectin aptamers.

[0015] Bell et al., *In Vitro Cellular & Developmental Biology - Animal* 35: 533-542 (1999) describes characterisation of the nuclease-resistant VEGF-specific aptamer NX1838 (2'-fluoropyrimidine RNA-based oligonucleotide/40-kDa-PEG).

[0016] Vorobjeb et al., *Antisense & Nucleic Acid Drug Development* 10:63-75 (2000) describes properties of chimeric oligodeoxynucleotides made of short sequences (tetramers, pentamers, octamers and decamers) bridged by hexamethylene diol and hexaethylene glyco linkers.

BRIEF DESCRIPTION OF THE DRAWINGS

10 [0017]

Figure 1 is an illustration depicting the classical and alternative pathways of the complement system.

15 Figure 2 is a schematic representation of the *in vitro* aptamer selection (SELEX™) process from pools of random sequence oligonucleotides.

20 Figure 3A is an illustration depicting the nucleotide sequence and secondary structure of an anti-C5 aptamer (SEQ ID NO: 1), in which the underlined residues are either 2'-H pyrimidine residues or 2'-fluoro pyrimidine residues, the boxed residues are either 2'-fluoro pyrimidine residues or 2'-OMe pyrimidine residues, and the residues indicated by an arrow (→) represent residues that must contain a 2'-fluoro modification.

25 Figure 3B is an illustration depicting the nucleotide sequence and secondary structure of the ARC330 anti-C5 aptamer (SEQ ID NO: 2), in which the circled residues are 2'-H residues, the pyrimidine residues are 2'-fluoro substituted, and the majority of purine residues are 2'-OMe substituted, except for the three 2'-OH purine residues shown in outline.

30 Figure 3C is an illustration depicting the nucleotide sequence and secondary structure of the ARC 186 anti-C5 aptamer (SEQ ID NO: 4) in which all 21 pyrimidine residues have 2'-fluoro modifications and the majority of purines (14 residues) have 2'-OMe modifications, except for the three 2'-OH purine residues shown in outline.

35 Figure 4 is an illustration of a 40 kD branched PEG (1,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide)).

40 Figure 5 is an illustration of a 40 kD branched PEG (1,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide)) attached to the 5'end of an aptamer.

45 Figure 6 is an illustration depicting various strategies for synthesis of high molecular weight PEG-nucleic acid conjugates.

50 Figure 7A is a graph comparing dose dependent inhibition of hemolysis by PEGylated anti-C5 aptamers (ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62), and ARC 187 (SEQ ID NO: 5)), to a non-PEGylated anti-C5 aptamer (ARC 186 (SEQ ID NO: 4)); Figure 7B is a table of the IC₅₀ values of the aptamers used in the hemolysis assay depicted in Figure 7A; Figure 7C is a graph comparing dose dependent inhibition of hemolysis by PEGylated anti-C5 aptamers ARC187 (SEQ ID NO: 5), ARC 1537 (SEQ ID NO: 65), ARC1730 (SEQ ID NO (66), and ARC1905 (SEQ ID NO: 67); Figure 7D is a table of the IC₅₀ values of the aptamers used in the hemolysis assay depicted in Figure 7C.

55 Figure 8 is a graph of percent inhibition of hemolysis by the anti- C5 aptamer, ARC658 (SEQ ID NO: 62), of cynomolgus serum complement versus human serum complement.

Figure 9 is a graph depicting the binding of ARC186 (SEQ ID NO: 4) to purified C5 protein at both 37°C and room temperature (23 °C) following a 15 minute incubation.

Figure 10 is another graph depicting the binding of ARC186 (SEQ ID NO: 4) to purified C5 protein at both 37 °C and room temperature (23°C) following a 4 hour incubation.

55 Figure 11 is graph showing a time course of dissociation of C5•ARC186 complex at 23 °C.

Figure 12 is a graph showing a time course of equilibration in the formation of C5•ARC186 complex at 23 °C.

Figure 13 is a graph depicting ARC186 (SEQ ID NO: 4) binding to C5 protein versus protein components upstream and downstream in the complement cascade.

5 Figure 14 is a graph depicting the percentage of radiolabeled ARC186 (SEQ ID NO: 4) that bound C5 in the presence of unlabeled competitor ARC186 (SEQ ID NO: 4), ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62) or ARC187 (SEQ ID NO: 5).

10 Figure 15 is a graph depicting the amount of C5b complement protein produced in blood samples incubated for 5 hours at 25 °C and 37 °C in the presence of varying concentrations of the ARC186 (SEQ ID NO: 4) aptamer.

15 Figure 16 is a graph depicting percent complement inhibition by ARC187 (SEQ ID NO: 5) in the presence of zymosan in undiluted human serum, citrated human whole blood or cynomolgus serum.

15 Figure 17 is a graph showing ARC658 (SEQ ID NO: 62) fully inhibits complement activation (C5a) in the tubing loop model described in Example 1D.

20 Figure 18 is a graph depicting the dissociation constants for Round 10 of the C5 selection pools. Dissociation constants (K_d s) were estimated by fitting the data to the equation: fraction RNA bound = amplitude* K_d /(K_d + [C5]). "ARC520" (SEQ ID NO: 70) refers to the naive unselected dRmY pool and the "+" indicates the presence of competitor (0.1mg/ml tRNA, 0.1mg/ml salmon sperm DNA).

25 Figure 19 is a graph depicting C5 clone dissociation constant curves. Dissociation constants (K_d s) were estimated by fitting the data to the equation: fraction RNA bound = amplitude* K_d /(K_d + [C5]).

Figure 20 is a graph depicting an IC_{50} curve that illustrates the inhibitory effect on hemolysis activity of varying concentrations of anti-C5 aptamer clone ARC913 (SEQ ID NO: 75) as compared to ARC186 (SEQ ID NO: 4).

30 Figure 21 is an illustration depicting the structure of ARC187 (SEQ ID NO: 5).

35 Figure 22 is an illustration depicting the structure of ARC 1905 (SEQ ID NO: 67).

Figure 23 is a table outlining the experimental design of the first isolated perfused heart study.

35 Figure 24 is a graph comparing the pressure tracings for the intraventricular pressure in the left ventricle (LV) of an isolated heart exposed to human plasma (A) with the LVP pressure tracings of an isolated heart exposed to the control aptamer solution (B).

40 Figure 25 is a graph comparing the pressure tracings for the intraventricular pressure in the left ventricle (LV) of the isolated hearts exposed to the molar equivalent, 10X and 50X aptamer/C5 solutions (where a concentration of approximately 500 nM is assumed for C5 in normal, undiluted human plasma).

45 Figure 26 is a graph comparing the heart rate changes in beats per minute (bpm) in isolated mouse hearts after exposure to human plasma and various plasma/aptamer solutions.

Figure 27 is a graph comparing the changes in the heart weight in isolated mouse hearts before and after exposure to human plasma containing 0 -1X molar ratio ARC186 (SEQ ID NO: 4) (failed hearts), or 10-50X molar ratio (hearts protected with C5 aptamer).

50 Figure 28 is a graph comparing the relative C5a production in human plasma, containing varying aptamer concentrations, following perfusion through isolated mouse hearts. Relative C5a concentrations are plotted as absorbance units (Abs), where higher readings reflect the presence of higher C5a levels.

55 Figure 29 is a graph comparing the relative soluble C5b-9 production in human plasma containing varying aptamer concentrations, following perfusion through isolated mouse hearts.

Figure 30 is a graph showing the effect of ARC186 (SEQ ID NO: 4) on C3 cleavage in mouse heart effluent.

Figure 31 is a table showing the immunohistochemistry staining results for the isolated perfused mouse heart study.

Figure 32 is a table showing the molar ratio of ARC658 (SEQ ID NO: 62) necessary, in human or primate serum, to protect the heart from C5b-mediated damage.

5 Figure 33 is a graph showing a log-linear plot of remaining percent of full-length ARC186 as a function of incubation time in both rat and cynomolgus macaque plasma.

Figure 34 is a table showing the experimental design of the pharmacokinetic study conducted Sprague-Dawley rats as described in Example 5.

10 Figure 35 is a table showing mean plasma concentration of ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62) or ARC187 (SEQ ID NO: 5) versus time in Sprague-Dawley rats.

15 Figure 36 is a graph depicting mean plasma concentration of ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5) over time following intravenous administration of aptamer in rats.

Figure 37 is a table showing the noncompartmental analysis of the concentration versus time data depicted in Figures 35 and 36.

20 Figure 38A is a table showing the design for the pharmacokinetic study of ARC 187 (SEQ ID NO: 5) and ARC1905 (SEQ ID NO: 67) in mice; Figure 38B is a graph depicting the pharmacokinetic profile of ARC187 (SEQ ID NO: 5) and ARC1905 (SEQ ID NO: 67) in CD-1 mice after a single IV bolus administration; Figure 38C is a table showing the noncompartmental analysis of the concentration versus time data depicted in Figure 38B.

25 Figure 39 is a table showing detection of the listed aptamers in mouse heart tissue following intravenous administration.

Figure 40 is a table showing the experimental design of animal Study 1, described in Example 5E.

30 Figure 41 is a table showing aptamer plasma concentration versus time following intravenous bolus administration of aptamer to cynomolgus macaques.

Figure 42 is a table listing the pharmacokinetic parameters for ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5) administered intravenously to cynomolgus macaque in Study 1.

35 Figures 43(a) and 43(c) are graphs depicting plasma concentrations of sC5b-9 and C5a over time following intravenous administration of the anti-C5 aptamers ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62), or ARC187 (SEQ ID NO: 5) to cynomolgus macaques; Figures 43(b) and 43(d) are graphs depicting plasma concentrations of sC5b-9 and C5a versus concentration of anti-C5 aptamers, ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62), or ARC187 (SEQ ID NO: 5).

40 Figure 44 is a table showing the experimental design of Study 2, described in Example 5F.

Figure 45 is a graph showing the mean aptamer plasma concentration at various time points following intravenous administration of ARC658 (SEQ ID NO: 62), or ARC187 (SEQ ID NO: 5) to cynomolgus macaques.

45 Figure 46 is a table showing the two compartmental analysis of the concentration versus time data following intravenous bolus aptamer administration to cynomolgus macaque.

50 Figure 47 is a graph depicting C5b-9 concentration versus ARC 187 (SEQ ID NO: 5) or ARC658 (SEQ ID NO: 62) concentration in the presence of zymosan in cynomolgus plasma.

Figure 48 is a graph depicting C5a concentration versus ARC 187 (SEQ ID NO: 5) or ARC658 (SEQ ID NO: 62) concentration in the presence of zymosan in cynomolgus plasma.

55 Figure 49 is a table summarizing the PK-PD study of ARC 187 (SEQ ID NO: 5) during and after IV bolus plus infusion administration to cynomolgus macaques.

Figure 50 is a table summarizing the pharmacokinetic parameters for ARC 187 (SEQ ID NO: 5) in cynomolgus

macaques after IV bolus administration.

Figure 51 is a graph depicting the calculated and actual measured pharmacokinetic profiles of ARC187 (SEQ ID NO: 5) during and after IV bolus plus infusion administration to cynomolgus macaques.

Figure 52 is a graph showing the plasma levels of active ARC 187 (SEQ ID NO: 5) remain constant during and after IV bolus plus infusion administration to cynomolgus macaques.

Figure 53 is a table showing the predicted human dosing requirements for anti-C5 aptamers in CABG surgery.

Figure 54 is a graph depicting ARC 187 (SEQ ID NO: 5) has relatively no *in vitro* effect on coagulation as measured by the prothrombin time (PT) and activated partial thromboplastin time (APTT).

Figure 55 is a table summarizing the *in vitro* effects of ARC 187 (SEQ ID NO: 5) on anti-coagulation activity of heparin, and procoagulation activity of protamine.


Figure 56 is a graph showing ARC187 (SEQ ID NO: 5) does not effect the reversal of heparin anticoagulation *in vivo*.

Figure 57 is graph showing heparin and protamine both have no effect on ARC 187 (SEQ ID NO: 5) anti-complement function, measured by inhibition of complement activation of zymosan.

SUMMARY OF THE INVENTION

[0018] The present invention provides materials and methods for the treatment, prevention and amelioration of complement related disease. In one embodiment, an aptamer comprising a nucleotide sequence according to ARC 186 (SEQ ID NO: 4) having a PEG conjugated to its 5' end is provided. In particular embodiments, this ARC186 aptamer/PEG conjugate comprises substantially the same binding affinity for C5 complement protein as an aptamer comprising the sequence according to SEQ ID NO: 4. Substantially the same binding affinity as used herein means no more than about a 2 to ten fold difference, preferably no more than a 2 to five fold difference in dissociation constants as measured by dot blot analysis. In some embodiments the dissociation constants are measured by competition dot blot analysis as described in Example 1A below. In some embodiments the aptamer/PEG conjugate comprises a half life, preferably the terminal half life in a two compartment model, of at least 15 hours, preferably at least 24 hours, more preferably at least 48 hours in primate. In some embodiments the aptamer/PEG conjugate comprises a half life, preferably the terminal half life in a two compartment model, of at least 10, preferably at least 15 hours in rat. The PEG conjugated to the 5' end of ARC186 (SEQ ID NO: 4) is a 40 kDa PEG. The 40 kDa PEG may be a branched PEG. The branched 40 kDa PEG may be 1,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide) or 2,3-bis(mPEG-[20 kDa])-propyl-1-carbamoyl.

[0019] Where the branched 40 kDa PEG is 1,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide), an aptamer having the structure set forth below is provided:




where,

“” indicates a linker

Aptamer = fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-3T (SEQ ID NO: 4),

wherein fC and fU = 2'-fluoro nucleotides, and mG and mA = 2'-OMe nucleotides and all other nucleotides are 2'-OH.

[0020] Where the branched 40 kDa PEG is 2,3-bis(mPEG-[20 kDa])-propyl-1-carbamoyl, an aptamer having the structure set forth below is provided:



where,


“” indicates a linker

Aptamer = fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfC fCfUmGfCmG-3T (SEQ ID NO: 4),

wherein fC and fU = 2'-fluoro nucleotides, and mG and mA = 2'-OMe nucleotides and all other nucleotides are 2'-OH.

[0021] In some embodiments the linker is an alkyl linker. In particular embodiments, the alkyl linker comprises 2 to 18 consecutive CH₂ groups. In preferred embodiments, the alkyl linker comprises 2 to 12 consecutive CH₂ groups. In particularly preferred embodiments the alkyl linker comprises 3 to 6 consecutive CH₂ groups.

[0022] Disclosed is an aptamer, ARC187 (SEQ ID NO: 5), having the structure set forth below:




where **Aptamer** =

25 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfC fCfUmGfCmG-3T (SEQ ID NO: 4)

wherein fC and fU = 2'-fluoro nucleotides, and mG and mA = 2'-OMe nucleotides and all other nucleotides are 2'-OH.

[0023] Also provided is an aptamer, ARC1905 (SEQ ID NO: 67), having the structure set forth below:



- where **Aptamer** =

fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfC fCfUmGfCmG-3T (SEQ ID NO: 4)

wherein fC and fU = 2'-fluoro nucleotides, and mG and mA = 2'-OMe nucleotides and all other nucleotides are 2'-OH.

[0024] In another aspect, the invention provides pharmaceutical compositions. Described is a pharmaceutical composition comprising a therapeutically effective amount of ARC187 (SEQ ID NO: 5) or ARC1905 (SEQ ID NO: 67) or a salt thereof. The pharmaceutical composition may comprise a pharmaceutically acceptable carrier or diluent. In this aspect an ARC187 (SEQ ID NO: 5) or ARC1905 (SEQ ID NO: 67) pharmaceutical composition for use in the treatment, prevention or amelioration of disease *in vivo* is provided.

[0025] In another aspect of the present disclosure, methods of treatment are provided. In one embodiment, the method comprises treating, preventing or ameliorating a disease mediated by C5 complement protein, and/or its derivatives C5a and C5b-9, the method including administering a pharmaceutical composition comprising ARC187 (SEQ ID NO: 5) or ARC1905 (SEQ ID NO: 67) or a salt thereof to a vertebrate. In some embodiments, the method comprises administering the pharmaceutical composition of the invention to a mammal. In some embodiments, the mammal is a human.

[0026] In some embodiments, the C5 complement protein, C5a and/or C5b-9-mediated disease to be treated is acute ischemic diseases (myocardial infarction, stroke, ischemic/reperfusion injury); acute inflammatory diseases (infectious disease, septicemia, shock, acute/hyperacute transplant rejection); chronic inflammatory and/or immune-mediated diseases (allergy, asthma, rheumatoid arthritis, and other rheumatological diseases, multiple sclerosis and other neurological diseases, psoriasis and other dermatological diseases, myasthenia gravis, systemic lupus erythematosus (SLE), subacute/chronic transplant rejection, glomerulonephritis and other renal diseases). In some embodiments, the C5 complement protein, C5a and/or C5b-9 mediated diseases to be treated include complement activation associated with dialysis or circumstances in which blood is passed over and/or through synthetic tubing and/or foreign material. In some embodiments, the C5 complement protein, C5a and/or C5b-9 mediated disease to be treated is selected from the group

consisting of myocardial injury relating to CABG surgery, myocardial injury relating to balloon angioplasty and myocardial injury relating to restenosis. In some embodiments the C5 complement protein C5a and/or C5b-9- mediated disease to be treated is complications relating to CABG surgery. In a particular embodiment, the disease to be treated is myocardial injury relating to CABG surgery.

5 [0027] In some embodiments, the method of the present disclosure includes administering the pharmaceutical composition comprising ARC187 (SEQ ID NO: 5) or ARC 1905 (SEQ ID NO: 67) to achieve an aptamer plasma concentration that is about .5 to about 10 times that of the endogenous C5 complement protein. In some embodiments, the pharmaceutical ARC187 (SEQ ID NO: 5) or ARC1905 (SEQ ID NO: 67) aptamer compositions are administered to achieve an aptamer plasma concentration that is about .75 to about 5 times, .75 to about 3 times, and 1.5 to about 2 times that of the endogenous C5 complement protein while in other embodiments the aptamer composition is administered to achieve a concentration equivalent to that of the endogenous complement protein. In some embodiments, the pharmaceutical composition comprising ARC 187 (SEQ ID NO: 5) or ARC1905 (SEQ ID NO: 67) is administered to achieve an aptamer plasma concentration of about 5 μ M, about 4 μ M, about 3 μ M, about 2 μ M, about 1.5 μ M, about 1 μ M or of about 500 nM.

10 [0028] Any combination of route, duration, and rate of administration may be used that is sufficient to achieve the 15 aptamer plasma concentrations. In some embodiments the pharmaceutical composition is administered intravenously. In some embodiments, the pharmaceutical composition is administered as a bolus and/or via continuous infusion.

20 [0029] In particular embodiments of treating, preventing and/or ameliorating complications related to CABG surgery, particularly myocardial injury related to CABG surgery, the method of the present disclosure comprises administering the pharmaceutical composition prior to surgery and continuing administration at least 24 hours, in some embodiments 25 about 48 hours or in some embodiments about 72 hours. In a particular embodiment, a plasma aptamer concentration of about two times the endogenous complement protein concentration is achieved by administration of an intravenous bolus of about .75 to 1.25, preferably of about 1 mg of aptamer per kg of the patient to be treated in advance of, simultaneously with or after intravenous infusion of a lower dose of aptamer wherein mg does not include the weight of the conjugated PEG. In some embodiments the lower dose will be infused at a rate selected from the range of 0.001 to 30 0.005 mg/kg/min wherein mg does not include the weight of the conjugated PEG. In a particular embodiment, the lower dose will be infused at a rate of about 0.0013 mg/kg/min. In still other embodiments, where the aptamer/conjugate comprises a sufficiently long half life, the aptamer pharmaceutical composition may be administered once or twice daily 35 as an intravenous bolus dose.

[0030] In another aspect of the invention, diagnostic methods are provided. Described is a diagnostic method that 40 comprises contacting the ARC 187 (SEQ ID NO 5) or ARC1905 (SEQ ID NO 67) with a composition suspected of comprising C5 complement protein or a variant thereof, and detecting the presence or absence of C5 complement protein or a variant thereof. In some embodiments the complement protein or variant are vertebrate, particularly mammalian, and more particularly human. Disclosed is an ARC187 (SEQ ID NO: 5) or ARC1905 (SEQ ID NO: 67) composition for use as an *in vitro* or *in vivo* diagnostic.

45 [0031] The present disclosure provides, an aptamer comprising a nucleotide sequence selected from the group consisting of: ARC 330 (SEQ ID NO: 2) and ARC188-189, ARC250, ARC296-297, ARC331-334, ARC411-440, ARC457-459, ARC473, ARC522-525, ARC532, ARC543-544, ARC550-554, ARC657-658, ARC672, ARC706, ARC1537, ARC1730, (SEQ ID NOS: 6 to SEQ NO: 66). In a particular embodiment, an aptamer comprising a nucleotide sequence comprises a sequence according to SEQ ID NO 1. In a particular embodiment, an aptamer comprising a nucleotide sequence 50 selected from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66 is provided. In some embodiments, where the aptamer comprises a nucleotide sequence selected from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66, the aptamer comprises substantially the same binding affinity for C5 complement protein as an aptamer comprising the sequence according to SEQ ID NO: 4.

[0032] In some embodiments wherein the aptamer comprises a nucleotide sequence selected from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66, the aptamer comprises a half life, preferably the terminal half life in a two compartment model, of at least 15, preferably at least 30 hours in primate. In some embodiments wherein the aptamer comprises a nucleotide sequence selected from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66, the aptamer comprises a half life, preferably the terminal half life in a two compartment model, of at least 1 and a half, preferably at least seven hours in rat.

55 [0033] In some embodiments of this aspect of the present disclosure, wherein the aptamer comprises a nucleotide sequence selected from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66, the aptamer is synthesized with a 5' linker as follows:

$H_2N\text{~~~~~} 5' \text{ Aptamer } 3'$,

wherein "~~~" denotes the linker. In some embodiments the linker is an alkyl linker as follows: $H_2N-(CH_2)_n-5' \text{ Aptamer } 3'$

wherein n=2 to 18, preferably n= 2-12, more preferably n= 3 to 6, more preferably n=6, and wherein **Aptamer** = fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfC fCfUmGfCmG-3T (SEQ ID NO: 4)

wherein fC and fU = 2'-fluoro nucleotides, and mG and mA = 2'-OMe nucleotides and all other nucleotides are 2'-OH.

5 The resulting amine-modified aptamer may be conjugated to a PEG moiety selected from the group consisting of a 10 kDa PEG, 20 kDa PEG, 30 kDa PEG and 40kDa linear PEG. In some embodiments, a pharmaceutical composition comprising a therapeutically effective amount of an aptamer comprising a nucleotide sequence selected from the group consisting of: SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 6 to SEQ NO: 66, particularly from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66 or a salt thereof is provided. The pharmaceutical 10 composition of the present disclosure may comprise a pharmaceutically acceptable carrier or diluent. In this aspect a pharmaceutical composition for use in the treatment, prevention or amelioration of disease *in vivo*, comprising an aptamer which comprises a nucleotide sequence selected from the group consisting of: SEQ ID NO: 2 and SEQ ID NO: 6 to SEQ NO: 66, particularly from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66 is provided.

15 [0034] In another embodiment, a method of treating, preventing or ameliorating a disease mediated by C5 complement protein is provided, comprising administering a pharmaceutical composition comprising an aptamer or a salt thereof, where the aptamer comprises a nucleotide sequence selected from the group consisting of: SEQ ID NO: 2 and SEQ ID NO: 6 to SEQ NO: 66, particularly from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66 to a vertebrate. In some embodiments of this aspect of the present disclosure, the method comprises 20 administering the pharmaceutical composition of the present disclosure to a mammal, preferably a human.

[0035] In some embodiments, the C5 complement protein, C5a and/or C5b-9-mediated disease to be treated is acute ischemic diseases (myocardial infarction, stroke, ischemic/reperfusion injury); acute inflammatory diseases (infectious disease, septicemia, shock, acute/hyperacute transplant rejection); chronic inflammatory and/or immune-mediated diseases (allergy, asthma, rheumatoid arthritis, and other rheumatological diseases, multiple sclerosis and other neurological diseases, psoriasis and other dermatological diseases, myasthenia gravis, systemic lupus erythematosus (SLE), subacute/chronic transplant rejection, glomerulonephritis and other renal diseases). In some embodiments, the C5 complement protein, C5a and/or C5b-9 mediated diseases to be treated include complement activation associated with dialysis or circumstances in which blood is passed over and/or through synthetic tubing and/or foreign material. In some 25 embodiments, the C5 complement protein C5a and/or C5b-9- mediated disease to be treated is selected from the group consisting of myocardial injury relating to CABG surgery, myocardial injury relating to balloon angioplasty and myocardial injury relating to restenosis. In some embodiments the C5 complement protein C5a and/or C5b-9- mediated disease to be treated is complications relating to CABG surgery. In a particular embodiment, the disease to be treated is myocardial 30 injury relating to CABG surgery.

[0036] In some embodiments, the method of the present disclosure includes administering the pharmaceutical composition comprising an aptamer having a nucleotide sequence selected from the group consisting of: SEQ ID NO: 2 and SEQ ID NO: 6 to SEQ NO: 66, particularly from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66, to a patient to achieve an aptamer plasma concentration that is about .5 to about 10 times that of the endogenous C5 complement protein. In some embodiments, the pharmaceutical aptamer compositions are administered to achieve an aptamer plasma concentration that is about .75 to about 5 times, .75 to about 3 times, and 40 1.5 to about 2 times that of the endogenous C5 complement protein while in other embodiments the aptamer composition is administered to achieve a concentration equivalent to that of the endogenous complement protein. In some embodiments, the pharmaceutical composition of the present disclosure administered to achieve an aptamer plasma concentration of about 5 μ M, about 4 μ M, about 3 μ M, about 2 μ M, about 1.5 μ M, about 1 μ M or of about 500 nM.

[0037] Any combination of route, duration, and rate of administration may be used that is sufficient to achieve the aptamer plasma concentrations of the present disclosure. In some embodiments the pharmaceutical composition is administered intravenously. In some embodiments, the pharmaceutical composition is administered as a bolus and/or via continuous infusion.

[0038] In particular embodiments of treating, preventing and/or ameliorating complications related to CABG surgery, particularly myocardial injury related to CABG surgery, the method of the present disclosure comprises administering the pharmaceutical composition prior to surgery and continuing administration at least 24 hours, in some embodiments about 48 hours or in some embodiments about 72 hours. In a particular embodiment of this aspect of the present disclosure, the desired aptamer plasma concentration, e.g., two times the endogenous complement protein concentration in some embodiments, is achieved by administration of an intravenous bolus to the patient to be treated in advance of, simultaneously with, or after intravenous infusion of a lower dose of aptamer. In still other embodiments of this aspect of the present disclosure, where the aptamer/conjugate comprises a sufficiently long half life, the aptamer pharmaceutical composition may be administered once or twice daily as an intravenous bolus dose.

[0039] In another aspect of the present disclosure diagnostic methods are provided. In one embodiment, the diagnostic method comprises contacting a composition suspected of comprising C5 complement protein or a variant thereof with

an aptamer comprising a nucleotide sequence selected from the group consisting of: SEQ ID NO 2 and SEQ ID NO 6 to SEQ NO 66, particularly from the group consisting of SEQ ID NO: 61, SEQ ID NO: 62, and SEQ ID NO: 64 to SEQ ID NO: 66, and detecting the presence or absence of C5 complement protein or a variant thereof. In some embodiments the complement protein or variant is vertebrate, particularly mammalian, and more particularly human. The present disclosure provides an aptamer composition having an aptamer comprising a nucleotide sequence selected from the group consisting of: SEQ ID NO 2 and SEQ ID NO 6 to SEQ NO 66 for use as an *in vitro* or *in vivo* diagnostic.

[0040] In another aspect of the present disclosure, an aptamer comprising a nucleotide sequence that is 80% identical to any one of the sequences selected from the group consisting of ARC915, ARC 874-878, ARC954, SEQ ID NOS: 75 to 81, SEQ ID NO: 83, and SEQ ID NOS: 88 to 98 is provided. In some embodiments, an aptamer comprising a nucleotide sequence that is 80% identical to the unique region of any one of the sequences selected from the group consisting of SEQ ID NOS: 75 to 81 and SEQ ID NOS: 88 to 98 is provided. In another embodiment an aptamer comprising a nucleotide sequence that is 90% identical to any one of the sequences selected from the group consisting of SEQ ID NOS: 75 to 81, SEQ ID NO: 83, and SEQ ID NOS: 88 to 98 is provided. In a particular embodiment, an aptamer comprising a nucleotide sequence that is 90% identical to the unique region of any one of the sequences selected from the group consisting of SEQ ID NOS: 75 to 81 and SEQ ID NOS: 88 to 98 is provided. In yet another embodiment, an aptamer comprising a nucleotide sequence of 40 contiguous nucleotides identical to 40 contiguous nucleotides included in any one of the sequences selected from the group consisting of SEQ ID NOS: 75 to 81 and SEQ ID NOS: 88 to 98 is provided. In another embodiment, an aptamer comprising a nucleotide sequence of 30 contiguous nucleotides identical to 30 contiguous nucleotides included in any one of the sequences selected from the group consisting of SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98 is provided. In yet another embodiment, an aptamer that binds specifically to C5 complement protein comprising a nucleotide sequence of 10 contiguous nucleotides identical to 10 contiguous nucleotides included in any one of the sequences selected from the group consisting of SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98 is provided. In a preferred embodiment an aptamer comprising a nucleotide sequence according to any one of the nucleotide sequences selected from the group consisting of: SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98, is provided.

[0041] In some embodiments, the aptamers of this aspect of the present disclosure described immediately above may further comprise a chemical modification selected from the group consisting: of a chemical substitution at a sugar position; a chemical substitution at a phosphate position; and a chemical substitution at a base position of the nucleic acid sequence. In some embodiments the modification is selected from the group consisting of: incorporation of a modified nucleotide; 3' capping, conjugation to a high molecular weight, non-immunogenic compound; conjugation to a lipophilic compound; and modification of the phosphate back bone.

[0042] In preferred embodiments of this aspect of the present disclosure, the aptamer modulates a function of a C5 complement protein or a variant thereof. In particularly preferred embodiments, the aptamer inhibits a function of C5 complement protein or a variant thereof, preferably *in vivo*, more preferably *in vivo* in humans. In one embodiment of this aspect of the present disclosure, the function modulated, preferably inhibited, by the aptamer is C5 complement protein cleavage.

[0043] In some embodiments of another aspect, the invention provides a pharmaceutical composition comprising a therapeutically effective amount of an aptamer that blocks C5 complement protein cleavage *in vivo* or a salt thereof and a pharmaceutically acceptable carrier or diluent.

[0044] A pharmaceutical composition comprising a therapeutically effective amount of an aptamer comprising a nucleotide sequence 80% identical to, preferably 90% identical to a nucleotide sequence selected from the group consisting of SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98 or a salt thereof is described. A pharmaceutical composition comprising a therapeutically effective amount of an aptamer comprising a nucleotide sequence 80% identical to, preferably 90% identical to the unique region of a nucleotide sequence selected from the group consisting of SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98 or a salt thereof is described. A pharmaceutical composition comprising a therapeutically effective amount of an aptamer having 40, 30 or 10 contiguous nucleotides identical to 40, 30 or 10 nucleotides, respectively, to a nucleotide sequence selected from the group consisting of SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98 is described. The pharmaceutical composition may comprise a pharmaceutically acceptable carrier or diluent. A pharmaceutical composition is described for use in the treatment, prevention or amelioration of disease *in vivo*, where the pharmaceutical composition comprises an aptamer having a nucleotide sequence selected from the group consisting of: SEQ ID NOS: 3 to 4, SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98 or a salt thereof.

[0045] In some embodiments, the C5 complement protein, C5a and/or C5b-9-mediated disease to be treated is acute ischemic diseases (myocardial infarction, stroke, ischemic/reperfusion injury); acute inflammatory diseases (infectious disease, septicemia, shock, acute/hyperacute transplant rejection); chronic inflammatory and/or immune-mediated diseases (allergy, asthma, rheumatoid arthritis, and other rheumatological diseases, multiple sclerosis and other neurological diseases, psoriasis and other dermatological diseases, myasthenia gravis, systemic lupus erythematosus (SLE), subacute/chronic transplant rejection, glomerulonephritis and other renal diseases). In some embodiments, the C5

complement protein, C5a and/or C5b-9 mediated diseases to be treated include complement activation associated with dialysis or circumstances in which blood is passed over and/or through synthetic tubing and/or foreign material. In some embodiments, the C5 complement protein C5a and/or C5b-9- mediated disease to be treated is selected from the group consisting of myocardial injury relating to CABG surgery, myocardial injury relating to balloon angioplasty and myocardial injury relating to restenosis. In some embodiments the C5 complement protein C5a and/or C5b-9- mediated disease to be treated is complications relating to CABG surgery. In a particular embodiment, the disease to be treated is myocardial injury relating to CABG surgery.

[0046] In some embodiments, the method of the present disclosure includes administering the pharmaceutical composition comprising an aptamer having a nucleotide sequence selected from the group consisting of: SEQ ID NOS: 3 to 4, SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98, to a patient to achieve an aptamer plasma concentration that is about .5 to about 10 times that of the endogenous C5 complement protein. In some embodiments, the pharmaceutical aptamer compositions are administered to achieve an aptamer plasma concentration that is about .75 to about 5 times, .75 to about 3 times, and 1.5 to about 2 times that of the endogenous C5 complement protein while in other embodiments the aptamer composition is administered to achieve a concentration equivalent to that of the endogenous complement protein. In some embodiments, the pharmaceutical composition of the present disclosure administered to achieve an aptamer plasma concentration of about 5 μ M, about 4 μ M, about 3 μ M, about 2 μ M, about 1.5 μ M, about 1 μ M or of about 500 nM.

[0047] Any combination of route, duration, and rate of administration may be used that is sufficient to achieve the aptamer plasma concentrations of the present disclosure. In some embodiments the pharmaceutical composition is administered intravenously. In some embodiments, the pharmaceutical composition is administered as a bolus and/or via continuous infusion.

[0048] In particular embodiments of treating, preventing and/or ameliorating complications related to CABG surgery, particularly myocardial injury related to CABG surgery, the method of the present disclosure comprises administering the pharmaceutical composition prior to surgery and continuing administration at least 24 hours, in some embodiments about 48 hours or in some embodiments about 72 hours. In a particular embodiment of this aspect of the present disclosure, the desired aptamer plasma concentration, e.g., two times the endogenous complement protein concentration in some embodiments, is achieved by administration of an intravenous bolus to the patient to be treated in advance of, simultaneously with or after intravenous infusion of a lower dose of aptamer. In still other embodiments of this aspect of the present disclosure, where the aptamer/conjugate comprises a sufficiently long half life, the aptamer pharmaceutical composition may be administered once or twice daily as an intravenous bolus dose.

[0049] In another embodiment, a diagnostic method is provided, the method comprising contacting a composition suspected of comprising C5 complement protein or a variant thereof with an aptamer comprising a nucleotide sequence selected from the group consisting of: SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98 and detecting the presence or absence of C5 complement protein or a variant thereof. In some embodiments the complement protein or variant is vertebrate, particularly mammalian, and more particularly human. The present disclosure provides an aptamer composition having an aptamer comprising a nucleotide sequence selected from the group consisting of: SEQ ID NOS: 75 to 81, SEQ ID NO: 83 and SEQ ID NOS: 88 to 98 for use as an *in vitro* or *in vivo* diagnostic.

[0050] In some embodiments, an aptamer comprising a nucleotide sequence consisting essentially of a nucleotide sequence selected from the group consisting of SEQ ID NO 68 and 69 is provided. In some embodiments, an aptamer comprising a nucleotide sequence consisting of a nucleotide sequence selected from the group consisting of SEQ ID NO 68 and 69 is provided. In some embodiments of this aspect of the present disclosure, the aptamers may be used in a diagnostic method.

DETAILED DESCRIPTION OF THE INVENTION

[0051] The details of one or more embodiments of the invention are set forth in the accompanying description below. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. Other features, objects, and advantages of the invention will be apparent from the description. In the specification, the singular forms also include the plural unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In the case of conflict, the present Specification will control.

The SELEX™ Method

[0052] A suitable method for generating an aptamer is with the process entitled "Systematic Evolution of Ligands by Exponential Enrichment" ("SELEX™") generally depicted in Figure 2. The SELEX™ process is a method for the *in vitro* evolution of nucleic acid molecules with highly specific binding to target molecules and is described in, e.g., U.S. Pat.

No. 5,475,096 entitled "Nucleic Acid Ligands", and U.S. Pat. No. 5,270,163 (see also WO 91/19813) entitled "Nucleic Acid Ligands". Each SELEX™-identified nucleic acid ligand, *i.e.*, each aptamer, is a specific ligand of a given target compound or molecule. The SELEX™ process is based on the unique insight that nucleic acids have sufficient capacity for forming a variety of two- and three-dimensional structures and sufficient chemical versatility available within their monomers to act as ligands (*i.e.*, form specific binding pairs) with virtually any chemical compound, whether monomeric or polymeric. Molecules of any size or composition can serve as targets.

[0053] SELEX™ relies as a starting point upon a large library or pool of single stranded oligonucleotides comprising randomized sequences. The oligonucleotides can be modified or unmodified DNA, RNA, or DNA/RNA hybrids. In some examples, the pool comprises 100% random or partially random oligonucleotides. In other examples, the pool comprises random or partially random oligonucleotides containing at least one fixed and/or conserved sequence incorporated within randomized sequence. In other examples, the pool comprises random or partially random oligonucleotides containing at least one fixed and/or conserved sequence at its 5' and/or 3' end which may comprise a sequence shared by all the molecules of the oligonucleotide pool. Fixed sequences are sequences such as hybridization sites for PCR primers, promoter sequences for RNA polymerases (*e.g.*, T3, T4, T7, and SP6), restriction sites, or homopolymeric sequences, such as poly A or poly T tracts, catalytic cores, sites for selective binding to affinity columns, and other sequences to facilitate cloning and/or sequencing of an oligonucleotide of interest. Conserved sequences are sequences, other than the previously described fixed sequences, shared by a number of aptamers that bind to the same target.

[0054] The oligonucleotides of the pool preferably include a randomized sequence portion as well as fixed sequences necessary for efficient amplification. Typically the oligonucleotides of the starting pool contain fixed 5' and 3' terminal sequences which flank an internal region of 30-50 random nucleotides. The randomized nucleotides can be produced in a number of ways including chemical synthesis and size selection from randomly cleaved cellular nucleic acids. Sequence variation in test nucleic acids can also be introduced or increased by mutagenesis before or during the selection/amplification iterations.

[0055] The random sequence portion of the oligonucleotide can be of any length and can comprise ribonucleotides and/or deoxyribonucleotides and can include modified or non-natural nucleotides or nucleotide analogs. See, *e.g.*, U.S. Patent No. 5,958,691; U.S. Patent No. 5,660,985; U.S. Patent No. 5,958,691; U.S. Patent No. 5,698,687; U.S. Patent No. 5,817,635; U.S. Patent No. 5,672,695, and PCT Publication WO 92/07065. Random oligonucleotides can be synthesized from phosphodiester-linked nucleotides using solid phase oligonucleotide synthesis techniques well known in the art. See, *e.g.*, Froehler et al., *Nucl. Acid Res.* 14:5399-5467 (1986) and Froehler et al., *Tet. Lett.* 27:5575-5578 (1986). Random oligonucleotides can also be synthesized using solution phase methods such as triester synthesis methods. See, *e.g.*, Sood et al., *Nucl. Acid Res.* 4:2557 (1977) and Hirose et al., *Tet. Lett.*, 28:2449 (1978). Typical syntheses carried out on automated DNA synthesis equipment yield 10¹⁴-10¹⁶ individual molecules, a number sufficient for most SELEX™ experiments. Sufficiently large regions of random sequence in the sequence design increases the likelihood that each synthesized molecule is likely to represent a unique sequence.

[0056] The starting library of oligonucleotides may be generated by automated chemical synthesis on a DNA synthesizer. To synthesize randomized sequences, mixtures of all four nucleotides are added at each nucleotide addition step during the synthesis process, allowing for random incorporation of nucleotides. As stated above, in one embodiment, random oligonucleotides comprise entirely random sequences; however, in other embodiments, random oligonucleotides can comprise stretches of nonrandom or partially random sequences. Partially random sequences can be created by adding the four nucleotides in different molar ratios at each addition step.

[0057] The starting library of oligonucleotides may be either RNA or DNA. In those instances where an RNA library is to be used as the starting library it is typically generated by transcribing a DNA library *in vitro* using T7 RNA polymerase or modified T7 RNA polymerases and purified. The RNA or DNA library is then mixed with the target under conditions favorable for binding and subjected to step-wise iterations of binding, partitioning and amplification, using the same general selection scheme, to achieve virtually any desired criterion of binding affinity and selectivity. More specifically, starting with a mixture containing the starting pool of nucleic acids, the SELEX™ method includes steps of: (a) contacting the mixture with the target under conditions favorable for binding; (b) partitioning unbound nucleic acids from those nucleic acids which have bound specifically to target molecules; (c) dissociating the nucleic acid-target complexes; (d) amplifying the nucleic acids dissociated from the nucleic acid-target complexes to yield a ligand-enriched mixture of nucleic acids; and (e) reiterating the steps of binding, partitioning, dissociating and amplifying through as many cycles as desired to yield highly specific, high affinity nucleic acid ligands to the target molecule. In those instances where RNA aptamers are being selected, the SELEX™ method further comprises the steps of: (i) reverse transcribing the nucleic acids dissociated from the nucleic acid-target complexes before amplification in step (d); and (ii) transcribing the amplified nucleic acids from step (d) before restarting the process.

[0058] Within a nucleic acid mixture containing a large number of possible sequences and structures, there is a wide range of binding affinities for a given target. A nucleic acid mixture comprising, for example, a 20 nucleotide randomized segment can have 4²⁰ candidate possibilities. Those which have the higher affinity constants for the target are most likely to bind to the target. After partitioning, dissociation and amplification, a second nucleic acid mixture is generated,

enriched for the higher binding affinity candidates. Additional rounds of selection progressively favor the best ligands until the resulting nucleic acid mixture is predominantly composed of only one or a few sequences. These can then be cloned, sequenced and individually tested for binding affinity as pure ligands or aptamers.

[0059] Cycles of selection and amplification are repeated until a desired goal is achieved. In the most general case, selection/amplification is continued until no significant improvement in binding strength is achieved on repetition of the cycle. The method is typically used to sample approximately 10^{14} different nucleic acid species but may be used to sample as many as about 10^{18} different nucleic acid species. Generally, nucleic acid aptamer molecules are selected in a 5 to 20 cycle procedure. In one embodiment, heterogeneity is introduced only in the initial selection stages and does not occur throughout the replicating process.

[0060] In one embodiment of SELEX™, the selection process is so efficient at isolating those nucleic acid ligands that bind most strongly to the selected target, that only one cycle of selection and amplification is required. Such an efficient selection may occur, for example, in a chromatographic-type process wherein the ability of nucleic acids to associate with targets bound on a column operates in such a manner that the column is sufficiently able to allow separation and isolation of the highest affinity nucleic acid ligands.

[0061] In many cases, it is not necessarily desirable to perform the iterative steps of SELEX™ until a single nucleic acid ligand is identified. The target-specific nucleic acid ligand solution may include a family of nucleic acid structures or motifs that have a number of conserved sequences and a number of sequences which can be substituted or added without significantly affecting the affinity of the nucleic acid ligands to the target. By terminating the SELEX™ process prior to completion, it is possible to determine the sequence of a number of members of the nucleic acid ligand solution family.

[0062] A variety of nucleic acid primary, secondary and tertiary structures are known to exist. The structures or motifs that have been shown most commonly to be involved in non-Watson-Crick type interactions are referred to as hairpin loops, symmetric and asymmetric bulges, pseudoknots and myriad combinations of the same. Almost all known cases of such motifs suggest that they can be formed in a nucleic acid sequence of no more than 30 nucleotides. For this reason, it is often preferred that SELEX™ procedures with contiguous randomized segments be initiated with nucleic acid sequences containing a randomized segment of between about 20 to about 50 nucleotides and in some embodiments of about 30 to about 40 nucleotides. In one example, the 5'-fixed:random:3'-fixed sequence comprises a random sequence of about 30 to about 50 nucleotides.

[0063] The core SELEX™ method has been modified to achieve a number of specific objectives. For example, U.S. Patent No. 5,707,796 describes the use of SELEX™ in conjunction with gel electrophoresis to select nucleic acid molecules with specific structural characteristics, such as bent DNA. U.S. Patent No. 5,763,177 describes SELEX™ based methods for selecting nucleic acid ligands containing photoreactive groups capable of binding and/or photocrosslinking to and/or photoactivating a target molecule. U.S. Patent No. 5,567,588 and U.S. Patent No. 5,861,254 describe SELEX™ based methods which achieve highly efficient partitioning between oligonucleotides having high and low affinity for a target molecule. U.S. Patent No. 5,496,938 describes methods for obtaining improved nucleic acid ligands after the SELEX™ process has been performed. U.S. Patent No. 5,705,337 describes methods for covalently linking a ligand to its target.

[0064] SELEX™ can also be used to obtain nucleic acid ligands that bind to more than one site on the target molecule, and to obtain nucleic acid ligands that include non-nucleic acid species that bind to specific sites on the target. SELEX™ provides means for isolating and identifying nucleic acid ligands which bind to any envisionable target, including large and small biomolecules such as nucleic acid-binding proteins and proteins not known to bind nucleic acids as part of their biological function as well as cofactors and other small molecules. For example, U.S. Patent No. 5,580,737 discloses nucleic acid sequences identified through SELEX™ which are capable of binding with high affinity to caffeine and the closely related analog, theophylline.

[0065] Counter-SELEX™ is a method for improving the specificity of nucleic acid ligands to a target molecule by eliminating nucleic acid ligand sequences with cross-reactivity to one or more non-target molecules. Counter- SELEX™ is comprised of the steps of: (a) preparing a candidate mixture of nucleic acids; (b) contacting the candidate mixture with the target, wherein nucleic acids having an increased affinity to the target relative to the candidate mixture may be partitioned from the remainder of the candidate mixture; (c) partitioning the increased affinity nucleic acids from the remainder of the candidate mixture; (d) dissociating the increased affinity nucleic acids from the target; e) contacting the increased affinity nucleic acids with one or more non-target molecules such that nucleic acid ligands with specific affinity for the non-target molecule(s) are removed; and f) amplifying the nucleic acids with specific affinity only to the target molecule to yield a mixture of nucleic acids enriched for nucleic acid sequences with a relatively higher affinity and specificity for binding to the target molecule. As described above for SELEX™, cycles of selection and amplification are repeated as necessary until a desired goal is achieved.

[0066] One potential problem encountered in the use of nucleic acids as therapeutics and vaccines is that oligonucleotides in their phosphodiester form may be quickly degraded in body fluids by intracellular and extracellular enzymes such as endonucleases and exonucleases before the desired effect is manifest. The SELEX™ method thus encompasses

the identification of high-affinity nucleic acid ligands containing modified nucleotides conferring improved characteristics on the ligand, such as improved *in vivo* stability or improved delivery characteristics. Examples of such modifications include chemical substitutions at the ribose and/or phosphate and/or base positions. SELEX™-identified nucleic acid ligands containing modified nucleotides are described, e.g., in U.S. Patent No. 5,660,985, which describes oligonucleotides containing nucleotide derivatives chemically modified at the 2' position of ribose, 5 position of pyrimidines, and 8 position of purines, U.S. Patent No. 5,756,703 which describes oligonucleotides containing various 2'-modified pyrimidines, and U.S. Patent No. 5,580,737 which describes highly specific nucleic acid ligands containing one or more nucleotides modified with 2'-amino (2'-NH₂), 2'-fluoro (2'-F), and/or 2'-OMe (2'-OMe) substituents.

[0067] Modifications of the nucleic acid ligands include, but are not limited to, those which provide other chemical groups that incorporate additional charge, polarizability, hydrophobicity, hydrogen bonding, electrostatic interaction, and fluxionality to the nucleic acid ligand bases or to the nucleic acid ligand as a whole. Modifications to generate oligonucleotide populations which are resistant to nucleases can also include one or more substitute internucleotide linkages, altered sugars, altered bases, or combinations thereof. Such modifications include, but are not limited to, 2'-position sugar modifications, 5-position pyrimidine modifications, 8-position purine modifications, modifications at exocyclic amines, substitution of 4-thiouridine, substitution of 5-bromo or 5-iodo-uracil; backbone modifications, phosphorothioate or alkyl phosphate modifications, methylations, and unusual base-pairing combinations such as the isobases isocytidine and isoguanidine. Modifications can also include 3' and 5' modifications such as capping.

[0068] In one embodiment, oligonucleotides are provided in which the P(O)O group is replaced by P(O)S ("thioate"), P(S)S ("dithioate"), P(O)NR₂ ("amide"), P(O)R, P(O)OR', CO or CH₂ ("formacetal") or 3'-amine (-NH-CH₂-CH₂-), where in each R or R' is independently H or substituted or unsubstituted alkyl. Linkage groups can be attached to adjacent nucleotides through an -O-, -N-, or -S- linkage. Not all linkages in the oligonucleotide are required to be identical. As used herein, the term phosphorothioate encompasses one or more non-bridging oxygen atoms in a phosphodiester bond replaced by one or more sulfur atoms.

[0069] In further embodiments, the oligonucleotides comprise modified sugar groups, for example, one or more of the hydroxyl groups is replaced with halogen, aliphatic groups, or functionalized as ethers or amines. In one embodiment, the 2'-position of the furanose residue is substituted by any of an OMe, O-alkyl, O-allyl, S-alkyl, S-allyl, or halo group. Methods of synthesis of 2'-modified sugars are described, e.g., in Sproat, et al., Nucl. Acid Res. 19:733-738 (1991); Cotten, et al., Nucl. Acid Res. 19:2629-2635 (1991); and Hobbs, et al., Biochemistry 12:5138-5145 (1973). Other modifications are known to one of ordinary skill in the art. Such modifications may be pre-SELEX™ process modifications or post-SELEX™ process modifications (modification of previously identified unmodified ligands) or may be made by incorporation into the SELEX process.

[0070] Pre-SELEX process modifications or those made by incorporation into the SELEX process yield nucleic acid ligands with both specificity for their SELEX™ target and improved stability, e.g., *in vivo* stability. Post-SELEX™ process modifications made to nucleic acid ligands may result in improved stability, e.g., *in vivo* stability without adversely affecting the binding capacity of the nucleic acid ligand.

[0071] The SELEX™ method encompasses combining selected oligonucleotides with other selected oligonucleotides and non-oligonucleotide functional units as described in U.S. Patent No. 5,637,459 and U.S. Patent No. 5,683,867. The SELEX™ method further encompasses combining selected nucleic acid ligands with lipophilic or non-immunogenic high molecular weight compounds in a diagnostic or therapeutic complex, as described, e.g., in U.S. Patent No. 6,011,020, U.S. Patent No. 6,051,698, and PCT Publication No. WO 98/18480. These patents and applications teach the combination of a broad array of shapes and other properties, with the efficient amplification and replication properties of oligonucleotides, and with the desirable properties of other molecules.

[0072] The identification of nucleic acid ligands to small, flexible peptides via the SELEX™ method has also been explored. Small peptides have flexible structures and usually exist in solution in an equilibrium of multiple conformers, and thus it was initially thought that binding affinities may be limited by the conformational entropy lost upon binding a flexible peptide. However, the feasibility of identifying nucleic acid ligands to small peptides in solution was demonstrated in U.S. Patent No. 5,648,214. In this patent, high affinity RNA nucleic acid ligands to substance P, an 11 amino acid peptide, were identified.

[0073] The aptamers with specificity and binding affinity to the target(s) of the present disclosure are typically selected by the SELEX™ process as described herein. As part of the SELEX™ process, the sequences selected to bind to the target are then optionally minimized to determine the minimal sequence having the desired binding affinity. The selected sequences and/or the minimized sequences are optionally optimized by performing random or directed mutagenesis of the sequence to increase binding affinity or alternatively to determine which positions in the sequence are essential for binding activity. Additionally, selections can be performed with sequences incorporating modified nucleotides to stabilize the aptamer molecules against degradation *in vivo*.

2'Modified SELEX™

[0074] In order for an aptamer to be suitable for use as a therapeutic, it is preferably inexpensive to synthesize, safe and stable *in vivo*. Wild-type RNA and DNA aptamers are typically not stable *in vivo* because of their susceptibility to degradation by nucleases. Resistance to nuclease degradation can be greatly increased by the incorporation of modifying groups at the 2'-position.

[0075] Fluoro and amino groups have been successfully incorporated into oligonucleotide libraries from which aptamers have been subsequently selected. However, these modifications greatly increase the cost of synthesis of the resultant aptamer, and may introduce safety concerns in some cases because of the possibility that the modified nucleotides could be recycled into host DNA by degradation of the modified oligonucleotides and subsequent use of the nucleotides as substrates for DNA synthesis.

[0076] Aptamers that contain 2'-OMe ("2'-OMe") nucleotides, as provided in some embodiments herein, overcome many of these drawbacks. Oligonucleotides containing 2'-OMe nucleotides are nuclease-resistant and inexpensive to synthesize. Although 2'-OMe nucleotides are ubiquitous in biological systems, natural polymerases do not accept 2'-O-methyl NTPs as substrates under physiological conditions, thus there are no safety concerns over the recycling of 2'-OMe nucleotides into host DNA. The SELEX™ method used to generate 2'-modified aptamers is described, e.g., in U.S. Patent Application No. 2004/0197804 and U.S. Patent Application No. 2005/0037394, entitled "Method for *in vitro* Selection of 2'-OMe Substituted Nucleic Acids".

[0077] The present invention includes aptamers that bind to and modulate the function of complement protein C5 which contain modified nucleotides (e.g., nucleotides which have a modification at the 2' position) to make the oligonucleotide more stable than the unmodified oligonucleotide to enzymatic and chemical degradation as well as thermal and physical degradation. Although there are several examples of 2'-OMe containing aptamers in the literature (see, e.g., Green et al., Current Biology 2, 683-695, 1995) these were generated by the *in vitro* selection of libraries of modified transcripts in which the C and U residues were 2'-fluoro (2'-F) substituted and the A and G residues were 2'-OH. Once functional sequences were identified then each A and G residue was tested for tolerance to 2'-OMe substitution, and the aptamer was re-synthesized having all A and G residues which tolerated 2'-OMe substitution as 2'-OMe residues. Most of the A and G residues of aptamers generated in this two-step fashion tolerate substitution with 2'-OMe residues, although, on average, approximately 20% do not. Consequently, aptamers generated using this method tend to contain from two to four 2'-OH residues, and stability and cost of synthesis are compromised as a result. By incorporating modified nucleotides into the transcription reaction which generate stabilized oligonucleotides used in oligonucleotide pools from which aptamers are selected and enriched by SELEX™ (and/or any of its variations and improvements, including those described herein), the methods of the present disclosure eliminate the need for stabilizing the selected aptamer oligonucleotides (e.g., by resynthesizing the aptamer oligonucleotides with modified nucleotides).

[0078] The present disclosure provides aptamers comprising combinations of 2'-OH, 2'-F, 2'-deoxy, and 2'-OMe modifications of the ATP, GTP, CTP, TTP, and UTP nucleotides. Also provided are aptamers comprising combinations of 2'-OH, 2'-F, 2'-deoxy, 2'-OMe, 2'-NH₂, and 2'-methoxyethyl modifications of the ATP, GTP, CTP, TTP, and UTP nucleotides. Also provided are aptamers comprising 5⁶ combinations of 2'-OH, 2'-F, 2'-deoxy, 2'-OMe, 2'-NH₂, and 2'-methoxyethyl modifications of the ATP, GTP, CTP, TTP, and UTP nucleotides.

[0079] 2' modified aptamers of the present disclosure are created using modified polymerases, e.g., A a modified T7 polymerase, having a rate of incorporation of modified nucleotides having bulky substituents at the furanose 2' position that is higher than that of wild-type polymerases. For example, a single mutant T7 polymerase (Y639F) in which the tyrosine residue at position 639 has been changed to phenylalanine readily utilizes 2'deoxy, 2'aminoo-, and 2'fluorouridine triphosphates (NTPs) as substrates and has been widely used to synthesize modified RNAs for a variety of applications. However, this mutant T7 polymerase reportedly can not readily utilize (*i.e.*, incorporate) NTPs with bulky 2'-substituents such as 2'-OMe or 2'-azido (2'-N₃) substituents. For incorporation of bulky 2' substituents, a double T7 polymerase mutant (Y639F/H784A) having the histidine at position 784 changed to an alanine residue in addition to the Y639F mutation has been described and has been used in limited circumstances to incorporate modified pyrimidine NTPs. See Padilla, R. and Sousa, R., Nucleic Acids Res., 2002, 30(24): 138. A single mutant T7 polymerase (H784A) having the histidine at position 784 changed to an alanine residue has also been described. Padilla et al., Nucleic Acids Research, 2002, 30: 138. In both the Y639F/H784A double mutant and H784A single mutant T7 polymerases, the change to a smaller amino acid residue such as alanine allows for the incorporation of bulkier nucleotide substrates, e.g., 2'-O methyl substituted nucleotides.

[0080] Generally, it has been found that under the conditions disclosed herein, the Y693F single mutant can be used for the incorporation of all 2'-OMe substituted NTPs except GTP and the Y639F/H784A double mutant can be used for the incorporation of all 2'-OMe substituted NTPs including GTP. It is expected that the H784A single mutant possesses properties similar to the Y639F and the Y639F/H784A mutants when used under the conditions disclosed herein.

[0081] 2'-modified oligonucleotides may be synthesized entirely of modified nucleotides, or with a subset of modified nucleotides. The modifications can be the same or different. All nucleotides may be modified, and all may contain the

same modification. All nucleotides may be modified, but contain different modifications, e.g., all nucleotides containing the same base may have one type of modification, while nucleotides containing other bases may have different types of modification. All purine nucleotides may have one type of modification (or are unmodified), while all pyrimidine nucleotides have another, different type of modification (or are unmodified). In this way, transcripts, or pools of transcripts are generated using any combination of modifications, including for example, ribonucleotides (2'-OH), deoxyribonucleotides (2'-deoxy), 2'-F, and 2'-OMe nucleotides. A transcription mixture containing 2'-OMe C and U and 2'-OH A and G is referred to as a "rRmY" mixture and aptamers selected therefrom are referred to as "rRmY" aptamers. A transcription mixture containing deoxy A and G and 2'-OMe U and C is referred to as a "dRmY" mixture and aptamers selected therefrom are referred to as "dRmY" aptamers. A transcription mixture containing 2'-OMe A, C, and U, and 2'-OH G is referred to as a "rGmH" mixture and aptamers selected therefrom are referred to as "rGmH" aptamers. A transcription mixture alternately containing 2'-OMe A, C, U and G and 2'-OMe A, U and C and 2'-F G is referred to as a "alternating mixture" and aptamers selected therefrom are referred to as "alternating mixture" aptamers. A transcription mixture containing 2'-OMe A, U, C, and G, where up to 10% of the G's are ribonucleotides is referred to as a "r/mGmH" mixture and aptamers selected therefrom are referred to as "r/mGmH" aptamers. A transcription mixture containing 2'-OMe A, U, and C, and 2'-F G is referred to as a "fGmH" mixture and aptamers selected therefrom are referred to as "fGmH" aptamers. A transcription mixture containing 2'-OMe A, U, and C, and deoxy G is referred to as a "dGmH" mixture and aptamers selected therefrom are referred to as "dGmH" aptamers. A transcription mixture containing deoxy A, and 2'-OMe C, G and U is referred to as a "dAmB" mixture and aptamers selected therefrom are referred to as "dAmB" aptamers, and a transcription mixture containing all 2'-OH nucleotides is referred to as a "rN" mixture and aptamers selected therefrom are referred to as "rN" or "rRrY" aptamers. A "mRmY" aptamer is one containing all 2'-OMe nucleotides and is usually derived from a r/mGmH oligonucleotide by post-SELEX replacement, when possible, of any 2'-OH Gs with 2'-OMe Gs.

[0082] Described examples include any combination of 2'-OH, 2'-deoxy and 2'-OMe nucleotides any combination of 2'-deoxy and 2'-OMe nucleotides. Described examples include any combination of 2'-deoxy and 2'-OMe nucleotides in which the pyrimidines are 2'-OMe (such as dRmY, mRmY or dGmH).

[0083] Incorporation of modified nucleotides into the aptamers of the invention is accomplished before (pre-) the selection process (e.g., a pre-SELEX™ process modification). Optionally, aptamers of the invention in which modified nucleotides have been incorporated by pre-SELEX™ process modification can be further modified by post-SELEX™ process modification (i.e., a post-SELEX™ process modification after a pre-SELEX™ modification). Pre-SELEX™ process modifications yield modified nucleic acid ligands with specificity for the SELEX™ target and also improved *in vivo* stability. Post-SELEX™ process modifications, i.e., modification (e.g., truncation, deletion, substitution or additional nucleotide modifications of previously identified ligands having nucleotides incorporated by pre-SELEX™ process modification) can result in a further improvement of *in vivo* stability without adversely affecting the binding capacity of the nucleic acid ligand having nucleotides incorporated by pre-SELEX™ process modification.

[0084] To generate pools of 2'-modified (e.g., 2'-OMe) RNA transcripts in conditions under which a polymerase accepts 2'-modified NTPs the preferred polymerase is the Y693F/H784A double mutant or the Y693F single mutant. Other polymerases, particularly those that exhibit a high tolerance for bulky 2'-substituents, may also be used. Such polymerases can be screened for this capability by assaying their ability to incorporate modified nucleotides under the transcription conditions disclosed herein.

[0085] A number of factors have been determined to be important for the transcription conditions useful in the methods disclosed herein. For example, increases in the yields of modified transcript are observed when a leader sequence is incorporated into the 5' end of a fixed sequence at the 5' end of the DNA transcription template, such that at least about the first 6 residues of the resultant transcript are all purines.

[0086] Another important factor in obtaining transcripts incorporating modified nucleotides is the presence or concentration of 2'-OH GTP. Transcription can be divided into two phases: the first phase is initiation, during which an NTP is added to the 3'-hydroxyl end of GTP (or another substituted guanosine) to yield a dinucleotide which is then extended by about 10-12 nucleotides; the second phase is elongation, during which transcription proceeds beyond the addition of the first about 10-12 nucleotides. It has been found that small amounts of 2'-OH GTP added to a transcription mixture containing an excess of 2'-OMe GTP are sufficient to enable the polymerase to initiate transcription using 2'-OH GTP, but once transcription enters the elongation phase the reduced discrimination between 2'-OMe and 2'-OH GTP, and the excess of 2'-OMe GTP over 2'-OH GTP allows the incorporation of principally the 2'-OMe GTP.

[0087] Another important factor in the incorporation of 2'-OMe substituted nucleotides into transcripts is the use of both divalent magnesium and manganese in the transcription mixture. Different combinations of concentrations of magnesium chloride and manganese chloride have been found to affect yields of 2'-OMe transcripts, the optimum concentration of the magnesium and manganese chloride being dependent on the concentration in the transcription reaction mixture of NTPs which complex divalent metal ions. To obtain the greatest yields of maximally 2' substituted OMe transcripts (i.e., all A, C, and U and about 90% of G nucleotides), concentrations of approximately 5 mM magnesium chloride and 1.5 mM manganese chloride are preferred when each NTP is present at a concentration of 0.5 mM. When

the concentration of each NTP is 1.0 mM, concentrations of approximately 6.5 mM magnesium chloride and 2.0 mM manganese chloride are preferred. When the concentration of each NTP is 2.0 mM, concentrations of approximately 9.6 mM magnesium chloride and 2.9 mM manganese chloride are preferred. In any case, departures from these concentrations of up to two-fold still give significant amounts of modified transcripts.

5 [0088] Priming transcription with GMP or guanosine is also important. This effect results from the specificity of the polymerase for the initiating nucleotide. As a result, the 5'-terminal nucleotide of any transcript generated in this fashion is likely to be 2'-OH G. The preferred concentration of GMP (or guanosine) is 0.5 mM and even more preferably 1 mM. It has also been found that including PEG, preferably PEG-8000, in the transcription reaction is useful to maximize incorporation of modified nucleotides.

[0089] For maximum incorporation of 2'-OMe ATP (100%), UTP (100%), CTP(100%) and GTP (~90%) ("r/mGmH") into transcripts the following conditions are preferred: HEPES buffer 200 mM, DTT 40 mM, spermidine 2 mM, PEG-8000 10% (w/v), Triton X-100 0.01% (w/v), MgCl₂ 5 mM (6.5 mM where the concentration of each 2'-OMe NTP is 1.0 mM), MnCl₂ 1.5 mM (2.0 mM where the concentration of each 2'-OMe NTP is 1.0 mM), 2'-OMe NTP (each) 500 μ M (more preferably, 1.0 mM), 2'-OH GTP 30 μ M, 2'-OH GMP 500 μ M, pH 7.5, Y639F/H784A T7 RNA Polymerase 15 units/ml, inorganic pyrophosphatase 5 units/ml, and an all-purine leader sequence of at least 8 nucleotides long. As used herein, one unit of the Y639F/H784A mutant T7 RNA polymerase (or any other mutant T7 RNA polymerase specified herein) is defined as the amount of enzyme required to incorporate 1 nmole of 2'-OMe NTPs into transcripts under the r/mGmH conditions. As used herein, one unit of inorganic pyrophosphatase is defined as the amount of enzyme that will liberate 1.0 mole of inorganic orthophosphate per minute at pH 7.2 and 25 °C.

20 [0090] For maximum incorporation (100%) of 2'-OMe ATP, UTP and CTP ("rGmH") into transcripts the following conditions are preferred: HEPES buffer 200 mM, DTT 40 mM, spermidine 2 mM, PEG-8000 10% (w/v), Triton X-100 0.01% (w/v), MgCl₂ 5 mM (9.6 mM where the concentration of each 2'-OMe NTP is 2.0 mM), MnCl₂ 1.5 mM (2.9 mM where the concentration of each 2'-OMe NTP is 2.0 mM), 2'-OMe NTP (each) 500 μ M (more preferably, 2.0 mM), pH 7.5, Y639F T7 RNA Polymerase 15 units/ml, inorganic pyrophosphatase 5 units/ml, and an all-purine leader sequence of at least 8 nucleotides long.

25

[0091] For maximum incorporation (100%) of 2'-OMe UTP and CTP ("rRmY") into transcripts the following conditions are preferred: HEPES buffer 200 mM, DTT 40 mM, spermidine 2 mM, PEG-8000 10% (w/v), Triton X-100 0.01% (w/v), MgCl₂ 5 mM (9.6 mM where the concentration of each 2'-OMe NTP is 2.0 mM), MnCl₂ 1.5 mM (2.9 mM where the concentration of each 2'-OMe NTP is 2.0 mM), 2'-OMe NTP (each) 500 μ M (more preferably, 2.0 mM), pH 7.5, Y639F/H784A T7 RNA Polymerase 15 units/ml, inorganic pyrophosphatase 5 units/ml, and an all-purine leader sequence of at least 8 nucleotides long.

[0092] For maximum incorporation (100%) of deoxy ATP and GTP and 2'-OMe UTP and CTP ("dRmY") into transcripts the following conditions are preferred: HEPES buffer 200 mM, DTT 40 mM, spermine 2 mM, spermidine 2 mM, PEG-8000 10% (w/v), Triton X-100 0.01 % (w/v), MgCl₂ 9.6 mM, MnCl₂ 2.9 mM, 2'-OMe NTP (each) 2.0 mM, pH 7.5, Y639F 35 T7 RNA Polymerase 15 units/ml, inorganic pyrophosphatase 5 units/ml, and an all-purine leader sequence of at least 8 nucleotides long.

[0093] For maximum incorporation (100%) of 2'-OMe ATP, UTP and CTP and 2'-F GTP ("fGmH") into transcripts the following conditions are preferred: HEPES buffer 200 mM, DTT 40 mM, spermidine 2 mM, PEG-8000 10% (w/v), Triton X-100 0.01% (w/v), MgCl₂ 9.6 mM, MnCl₂ 2.9 mM, 2'-OMe NTP (each) 2.0 mM, pH 7.5, Y639F T7 RNA Polymerase 15 units/ml, inorganic pyrophosphatase 5 units/ml, and an all-purine leader sequence of at least 8 nucleotides long.

[0094] For maximum incorporation (100%) of deoxy ATP and 2'-OMe UTP, GTP and CTP ("dAmB") into transcripts the following conditions are preferred: HEPES buffer 200 mM, DTT 40 mM, spermidine 2 mM, PEG-8000 10% (w/v), Triton X-100 0.01 % (w/v), MgCl₂ 9.6 mM, MnCl₂ 2.9 mM, 2'-OMe NTP (each) 2.0 mM, pH 7.5, Y639F T7 RNA Polymerase 15 units/ml, inorganic pyrophosphatase 5 units/ml, and an all-purine leader sequence of at least 8 nucleotides long.

45 [0095] For each of the above (a) transcription is preferably performed at a temperature of from about 20 °C to about
50 °C, preferably from about 30 °C to 45 °C, and more preferably at about 37 °C for a period of at least two hours and
(b) 50-300 nM of a double stranded DNA transcription template is used (200 nM template is used in round 1 to increase
diversity (300 nM template is used in dRmY transcriptions)), and for subsequent rounds approximately 50 nM, a 1/10
dilution of an optimized PCR reaction, using conditions described herein, is used). The preferred DNA transcription
templates are described below (where ARC254 and ARC256 transcribe under all 2'-OMe conditions and ARC255 trans-
scribes under rRmY conditions).

50

ARC254 (SEQ ID NO: 99):

ARC255 (SEQ ID NO: 100):

5 ARC256 (SEQ ID NO: 101):

10 [0096] Under rN transcription conditions of the present disclosure, the transcription reaction mixture comprises 2'-OH adenosine triphosphates (ATP), 2'-OH guanosine triphosphates (GTP), 2'-OH cytidine triphosphates (CTP), and 2'-OH uridine triphosphates (UTP). The modified oligonucleotides produced using the rN transcription mixtures of the present disclosure comprise substantially all 2'-OH adenosine, 2'-OH guanosine, 2'-OH cytidine, and 2'-OH uridine. In a preferred embodiment of rN transcription, the resulting modified oligonucleotides comprise a sequence where at least 80% of all adenosine nucleotides are 2'-OH adenosine, at least 80% of all guanosine nucleotides are 2'-OH guanosine, at least 80% of all cytidine nucleotides are 2'-OH cytidine, and at least 80% of all uridine nucleotides are 2'-OH uridine. In a more preferred embodiment of rN transcription, the resulting modified oligonucleotides comprise a sequence where at least 90% of all adenosine nucleotides are 2'-OH adenosine, at least 90% of all guanosine nucleotides are 2'-OH guanosine, at least 90% of all cytidine nucleotides are 2'-OH cytidine, and at least 90% of all uridine nucleotides are 2'-OH uridine. In a most preferred embodiment of rN transcription, the modified oligonucleotides comprise a sequence where 100% of all adenosine nucleotides are 2'-OH adenosine, 100% of all guanosine nucleotides are 2'-OH guanosine, 100% of all cytidine nucleotides are 2'-OH cytidine, and 100% of all uridine nucleotides are 2'-OH uridine.

[0097] Under rRmY transcription conditions of the present disclosure, the transcription reaction mixture comprises 2'-OH adenosine triphosphates, 2'-OH guanosine triphosphates, 2'-OMe cytidine triphosphates, and 2'-OMe uridine triphosphates. The modified oligonucleotides produced using the rRmY transcription mixtures of the present disclosure comprise substantially all 2'-OH adenosine, 2'-OH guanosine, 2'-OMe cytidine and 2'-OMe uridine. In a preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 80% of all adenosine nucleotides are 2'-OH adenosine, at least 80% of all guanosine nucleotides are 2'-OH guanosine, at least 80% of all cytidine nucleotides are 2'-OMe cytidine and at least 80% of all uridine nucleotides are 2'-OMe uridine. In a more preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 90% of all adenosine nucleotides are 2'-OH adenosine, at least 90% of all guanosine nucleotides are 2'-OH guanosine, at least 90% of all cytidine nucleotides are 2'-OMe cytidine and at least 90% of all uridine nucleotides are 2'-OMe uridine. In a most preferred embodiment, the resulting modified oligonucleotides comprise a sequence where 100% of all adenosine nucleotides are 2'-OH adenosine, 100% of all guanosine nucleotides are 2'-OH guanosine, 100% of all cytidine nucleotides are 2'-OMe cytidine and 100% of all uridine nucleotides are 2'-OMe uridine.

[0098] Under dRmY transcription conditions of the present disclosure, the transcription reaction mixture comprises 2'-deoxy adenosine triphosphates, 2'-deoxy guanosine triphosphates, 2'-O-methyl cytidine triphosphates, and 2'-O-methyl uridine triphosphates. The modified oligonucleotides produced using the dRmY transcription conditions of the present disclosure comprise substantially all 2'-deoxy adenosine, 2'-deoxy guanosine, 2'-O-methyl cytidine, and 2'-O-methyl uridine. In a preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 80% of all adenosine nucleotides are 2'-deoxy adenosine, at least 80% of all guanosine nucleotides are 2'-deoxy guanosine, at least 80% of all cytidine nucleotides are 2'-O-methyl cytidine, and at least 80% of all uridine nucleotides are 2'-O-methyl uridine. In a more preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 90% of all adenosine nucleotides are 2'-deoxy adenosine, at least 90 % of all guanosine nucleotides are 2'-deoxy guanosine, at least 90% of all cytidine nucleotides are 2'-O-methyl cytidine, and at least 90% of all uridine nucleotides are 2'-O-methyl uridine. In a most preferred embodiment, the resulting modified oligonucleotides comprise a sequence where 100% of all adenosine nucleotides are 2'-deoxy adenosine, 100% of all guanosine nucleotides are 2'-deoxy guanosine, 100% of all cytidine nucleotides are 2'-O-methyl cytidine, and 100% of all uridine nucleotides are 2'-O-methyl uridine.

[0099] Under rGmH transcription conditions of the present disclosure, the transcription reaction mixture comprises 2'-OH guanosine triphosphates, 2'-OMe cytidine triphosphates, 2'-OMe uridine triphosphates, and 2'-OMe adenosine triphosphates. The modified oligonucleotides produced using the rGmH transcription mixtures of the present disclosure comprise substantially all 2'-OH guanosine, 2'-OMe cytidine, 2'-OMe uridine, and 2'-OMe adenosine. In a preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 80% of all guanosine nucleotides are 2'-OH guanosine, at least 80% of all cytidine nucleotides are 2'-OMe cytidine, at least 80% of all uridine nucleotides are 2'-OMe uridine, and at least 80% of all adenosine nucleotides are 2'-OMe adenosine. In a more preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 90% of all guanosine nucleotides are 2'-OH guanosine, at least 90% of all cytidine nucleotides are 2'-OMe cytidine, at least 90% of all uridine nucleotides are 2'-OMe uridine, and at least 90% of all adenosine nucleotides are 2'-OMe adenosine.

otides are 2'-OH guanosine, at least 90% of all cytidine nucleotides are 2'-OMe cytidine, at least 90% of all uridine nucleotides are 2'-OMe uridine, and at least 90% of all adenosine nucleotides are 2'-OMe adenosine. In a most preferred embodiment, the resulting modified oligonucleotides comprise a sequence where 100% of all guanosine nucleotides are 2'-OH guanosine, 100% of all cytidine nucleotides are 2'-OMe cytidine, 100% of all uridine nucleotides are 2'-OMe uridine, and 100% of all adenosine nucleotides are 2'-OMe adenosine.

[0100] Under r/mGmH transcription conditions of the present disclosure, the transcription reaction mixture comprises 2'-OMe adenosine triphosphate, 2'-OMe cytidine triphosphate, 2'-OMe guanosine triphosphate, 2'-OMe uridine triphosphate and 2'-OH guanosine triphosphate. The resulting modified oligonucleotides produced using the r/mGmH transcription mixtures of the present disclosure comprise substantially all 2'-OMe adenosine, 2'-OMe cytidine, 2'-OMe guanosine, and 2'-OMe uridine, wherein the population of guanosine nucleotides has a maximum of about 10% 2'-OH guanosine. In a preferred embodiment, the resulting r/mGmH modified oligonucleotides comprise a sequence where at least 80% of all adenosine nucleotides are 2'-OMe adenosine, at least 80% of all cytidine nucleotides are 2'-OMe cytidine, at least 80% of all guanosine nucleotides are 2'-OMe guanosine, at least 80% of all uridine nucleotides are 2'-OMe uridine, and no more than about 10% of all guanosine nucleotides are 2'-OH guanosine. In a more preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 90% of all adenosine nucleotides are 2'-OMe adenosine, at least 90% of all cytidine nucleotides are 2'-OMe cytidine, at least 90% of all guanosine nucleotides are 2'-OMe guanosine, at least 90% of all uridine nucleotides are 2'-OMe uridine, and no more than about 10% of all guanosine nucleotides are 2'-OH guanosine. In a most preferred embodiment, the resulting modified oligonucleotides comprise a sequence where 100% of all adenosine nucleotides are 2'-OMe adenosine, 100% of all cytidine nucleotides are 2'-OMe cytidine, 90% of all guanosine nucleotides are 2'-OMe guanosine, and 100% of all uridine nucleotides are 2'-OMe uridine, and no more than about 10% of all guanosine nucleotides are 2'-OH guanosine.

[0101] Under fGmH transcription conditions of the present disclosure, the transcription reaction mixture comprises 2'-OMe adenosine triphosphates, 2'-OMe uridine triphosphates, 2'-OMe cytidine triphosphates, and 2'-F guanosine triphosphates. The modified oligonucleotides produced using the fGmH transcription conditions of the present disclosure comprise substantially all 2'-OMe adenosine, 2'-OMe uridine, 2'-OMe cytidine, and 2'-F guanosine. In a preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 80% of all adenosine nucleotides are 2'-OMe adenosine, at least 80% of all uridine nucleotides are 2'-OMe uridine, at least 80% of all cytidine nucleotides are 2'-OMe cytidine, and at least 80% of all guanosine nucleotides are 2'-F guanosine. In a more preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 90% of all adenosine nucleotides are 2'-OMe adenosine, at least 90% of all uridine nucleotides are 2'-OMe uridine, at least 90% of all cytidine nucleotides are 2'-OMe cytidine, and at least 90% of all guanosine nucleotides are 2'-F guanosine. In a most preferred embodiment, the resulting modified oligonucleotides comprise a sequence where 100% of all adenosine nucleotides are 2'-OMe adenosine, 100% of all uridine nucleotides are 2'-OMe uridine, 100% of all cytidine nucleotides are 2'-OMe cytidine, and 100% of all guanosine nucleotides are 2'-F guanosine.

[0102] Under dAmB transcription conditions of the present disclosure, the transcription reaction mixture comprises 2'-deoxy adenosine triphosphates, 2'-OMe cytidine triphosphates, 2'-OMe guanosine triphosphates, and 2'-OMe uridine triphosphates. The modified oligonucleotides produced using the dAmB transcription mixtures of the present disclosure comprise substantially all 2'-deoxy adenosine, 2'-OMe cytidine, 2'-OMe guanosine, and 2'-OMe uridine. In a preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 80% of all adenosine nucleotides are 2'-deoxy adenosine, at least 80% of all cytidine nucleotides are 2'-OMe cytidine, at least 80% of all guanosine nucleotides are 2'-OMe guanosine, and at least 80% of all uridine nucleotides are 2'-OMe uridine. In a more preferred embodiment, the resulting modified oligonucleotides comprise a sequence where at least 90% of all adenosine nucleotides are 2'-deoxy adenosine, at least 90% of all cytidine nucleotides are 2'-OMe cytidine, at least 90% of all guanosine nucleotides are 2'-OMe guanosine, and at least 90% of all uridine nucleotides are 2'-OMe uridine. In a most preferred embodiment, the resulting modified oligonucleotides comprise a sequence where 100% of all adenosine nucleotides are 2'-deoxy adenosine, 100% of all cytidine nucleotides are 2'-OMe cytidine, 100% of all guanosine nucleotides are 2'-OMe guanosine, and 100% of all uridine nucleotides are 2'-OMe uridine.

[0103] In each case, the transcription products can then be used as the library in the SELEX™ process to identify aptamers and/or to determine a conserved motif of sequences that have binding specificity to a given target. The resulting sequences are already stabilized, eliminating this step from the process to arrive at a stabilized aptamer sequence and giving a more highly stabilized aptamer as a result. Another advantage of the 2'-OMe SELEX™ process is that the resulting sequences are likely to have fewer 2'-OH nucleotides required in the sequence, possibly none. To the extent 2'OH nucleotides remain they can be removed by performing post-SELEX modifications.

[0104] As described below, lower but still useful yields of transcripts fully incorporating 2' substituted nucleotides can be obtained under conditions other than the optimized conditions described above. For example, variations to the above transcription conditions include:

[0105] The HEPES buffer concentration can range from 0 to 1 M. The present disclosure also contemplates the use of other buffering agents having a pKa between 5 and 10 including, for example, Tris(hydroxymethyl)aminomethane.

[0106] The DTT concentration can range from 0 to 400 mM. The methods of the present disclosure also provide for the use of other reducing agents including, for example, mercaptoethanol.

[0107] The spermidine and/or spermine concentration can range from 0 to 20 mM.

[0108] The PEG-8000 concentration can range from 0 to 50 % (w/v). The methods of the present disclosure also provide for the use of other hydrophilic polymer including, for example, other molecular weight PEG or other polyalkylene glycols.

[0109] The Triton X-100 concentration can range from 0 to 0.1% (w/v). The methods of the present disclosure also provide for the use of other non-ionic detergents including, for example, other detergents, including other Triton-X detergents.

[0110] The $MgCl_2$ concentration can range from 0.5 mM to 50 mM. The $MnCl_2$ concentration can range from 0.15 mM to 15 mM. Both $MgCl_2$ and $MnCl_2$ must be present within the ranges described and in a preferred embodiment are present in about a 10 to about 3 ratio of $MgCl_2$: $MnCl_2$, preferably, the ratio is about 3-5:1, more preferably, the ratio is about 3-4:1.

[0111] The 2'-OMe NTP concentration (each NTP) can range from 5 μ M to 5 mM.

[0112] The 2'-OH GTP concentration can range from 0 μ M to 300 μ M.

[0113] The 2'-OH GMP concentration can range from 0 to 5 mM.

[0114] The pH can range from pH 6 to pH 9. The methods of the present disclosure can be practiced within the pH range of activity of most polymerases that incorporate modified nucleotides. In addition, the methods of the present disclosure provide for the optional use of chelating agents in the transcription reaction condition including, for example, EDTA, EGTA, and DTT.

[0115] The selected aptamers having the highest affinity and specific binding as demonstrated by biological assays as described in the examples below are suitable therapeutics for treating conditions in which the C5 complement protein is involved in pathogenesis.

25 Aptamers with Binding Affinity to Complement System Protein C5

[0116] Although the complement system has an important role in the maintenance of health, it has the potential to cause or contribute to disease. Thus, it is desirable to develop inhibitors of the complement system for therapeutic use. It is particularly desirable to develop inhibitors of complement protein C5 because it is a component of both the classical and alternative pathways of complement activation cascades (Matis and Rollins (1995) *Nature Medicine* 1(8):839-842). Accordingly, inhibition of C5 can prevent complement-mediated damage caused by either pathway. Some complement system proteins, such as C1q and C3, are important in the normal defense mechanisms against microorganisms and in the clearance of immune components and damaged tissue; however, C5 is relatively unimportant for these functions. Thus, C5 function can be inhibited for short or long periods of time without compromising the protective role of the complement system.

[0117] A therapeutic C5 inhibitor is also desirable because inhibiting cleavage of C5 prevents the generation of two potentially damaging complement activities. First, inhibiting the generation of C5a from the cleavage of C5 eliminates the major complement chemotactic and vasoactive activity. Second, inhibiting the generation of C5b from the cleavage of C5 blocks assembly of the cytolytic C5b-9 membrane attack complex ("MAC"). Inhibition of C5 cleavage blocks both the C5a and the C5b effects on leukocytes and on tissue such as endothelial cells (Ward (1996) *Am. J. Pathol.* 149:1079).

[0118] Both C5a and the MAC have been implicated in acute and chronic inflammation associated with human disease, and their role in disease states has been confirmed in animal models. C5a is required for complement and neutrophil dependent lung vascular injury (Ward (1997) *J. Lab. Clin. Med.* 129:400; Mulligan et al., (1998) *J. Clin. Invest.* 98:503), and is associated with neutrophil and platelet activation in shock and in burn injury (Schmid et al., (1997) *Shock* 8:119).

[0119] The MAC mediates muscle injury in acute autoimmune myasthenia gravis (Biesecker and Gomez (1989) *J. Immunol.* 142:2654), organ rejection in transplantation (Baldwin et al., (1995) *Transplantation* 59:797; Brauer et al., (1995) *Transplantation* 59:288; Takahashi et al., (1997) *Immunol. Res.* 16:273), and renal injury in autoimmune glomerulonephritis (Biesecker (1981) *J. Exp. Med.* 39:1779; Nangaku (1997) *Kidney Int.* 52:1570). Both C5a and the MAC are implicated in acute myocardial ischemia (Homeister and Lucchesi (1994) *Annu. Rev. Pharmacol. Toxicol.* 34:17), acute (Bednar et al., (1997) *J. Neurosurg.* 86:139) and chronic CNS injury (Morgan (1997) *Exp. Clin. Immunogenet.* 14:19), leukocyte activation during extracorporeal circulation (Sun et al., (1995) *Nucleic Acids Res.* 23:2909; Spycher and Nydegger (1995) *Infusionsther. Transfusionsmed.* 22:36) and in tissue injury associated with autoimmune diseases including arthritis and lupus (Wang et al., (1996) *Immunology* 93:8563).

[0120] Complement activation has also been implicated in diabetic retinopathy, and can compound or initiate retinal vascular damage (Zhang et al., (2002) *Diabetes* 51:3499). Low level constitutive complement activation normally occurs in the non-diabetic eye, evidenced by the presence of MAC and complement regulatory proteins in the eyes of non-diabetic rats, indicating that complement dysregulation occurs in diabetic patients (Sohn et al., (2000) *IOVS* 41:3492). In addition, C5b-9 deposition has been detected in retinal vessels from diabetic human donors where absent from non-

diabetic human donors (Zhang *et al.*), reduced expression of CD55 and CD59 is shown in diabetic retinas (Zhang *et al.*), and glycated CD59 is present in urine from diabetic patients, but not non-diabetic patients (Acosta *et al.*, (2002) PNAS 97, 5450-5455). Additionally, the complement and vascular system is known to be activated in type I diabetes. See, *e.g.* Hansen, T.K. *et al.*, Diabetes, 53: 1570-1576 (2004). C5a activates endothelial cells via interaction with the immune and complement systems. See, *e.g.*, Albrecht, E.A. *et al.*, Am J Pathology, 164: 849-859 (2004). The vascular system is activated in ocular diseases including diabetic retinopathy. See, *e.g.* Gert, V.B. *et al.*, Invest Ophthalmol Vis Sci, 43: 1104-1108 (2002). The complement system is also activated in diabetic retinopathy. See, See, *e.g.* Gert, V.B. *et al.*, Invest Ophthalmol Vis Sci, 43: 1104-1108 (2002) and Badouin, C *et al.*, Am J Ophthalmol, 105:383-388 (1988).

5 [0120] In some embodiments, the materials of the present disclosure comprise a series of nucleic acid aptamers of about 15 to about 60 nucleotides in length which bind specifically to complement protein C5 and which functionally modulate, *e.g.*, block, the activity of complement protein C5 in *in vivo* and/or cell-based assays.

10 [0121] Aptamers that are capable of specifically binding and modulating complement protein C5 are set forth herein. These aptamers provide a low-toxicity, safe, and effective modality of treating, ameliorating and/or preventing a variety of complement-related diseases or disorders including, for example, complement-related heart disorders (*e.g.*, myocardial injury; C5 mediated complement complications relating to coronary artery bypass graft (CABG) surgery such as post-operative bleeding, systemic neutrophil and leukocyte activation, increased risk of myocardial infarction, and increased cognitive dysfunction; restenosis; and C5 mediated complement complications relating to percutaneous coronary intervention), ischemia-reperfusion injury (*e.g.*, myocardial infarction, stroke, frostbite), complement-related inflammatory disorders (*e.g.*, asthma, arthritis, sepsis, and rejection after organ transplantation), and complement-related autoimmune disorders (*e.g.*, myasthenia gravis, systemic lupus erythematosus (SLE)). Other indications for which C5 inhibition is desirable include, for example, lung inflammation (Mulligan *et al.* (1998) J. Clin. Invest. 98:503), extracorporeal complement activation (Rinder *et al.* (1995) J. Clin. Invest. 96:1564), antibody-mediated complement activation (Biesecker *et al.* (1989) J. Immunol. 142:2654), glomerulonephritis and other renal diseases, ocular indications such as C5 mediated ocular tissue damage, *e.g.* diabetic retinopathy, and paroxysmal nocturnal hemoglobinuria. These aptamers may also be used in diagnostics.

15 [0122] These aptamers may include modifications as described herein including, *e.g.*, conjugation to lipophilic or high molecular weight compounds (*e.g.*, PEG), incorporation of a capping moiety, incorporation of modified nucleotides, and modifications to the phosphate back bone.

20 [0123] Described herein is an isolated, non-naturally occurring aptamer that binds to the C5 complement protein is provided. In some embodiments, the isolated, non-naturally occurring aptamer has a dissociation constant ("K_d") for C5 complement protein of less than 100 μ M, less than 1 μ M, less than 500 nM, less than 100 nM, less than 50 nM, less than 1 nM, less than 500 pM, less than 100 pM, less than 50 pM. In some embodiments of the invention, the dissociation constant is determined by dot blot titration as described in Example 1 below.

25 [0124] In another embodiment, the aptamer modulates a function of the C5 complement protein. In another embodiment, the aptamer inhibits a C5 function while in another embodiment the aptamer stimulates a function of C5. A C5 complement protein variant as used herein encompasses variants that perform essentially the same function as a C5 complement protein function. A C5 complement protein variant preferably comprises substantially the same structure and in some embodiments comprises 80% sequence identity, more preferably 90% sequence identity, and more preferably 95% sequence identity to the amino acid sequence of the C5 complement protein comprising the amino acid sequence below (SEQ ID NO: 102) (cited in Haviland *et al.*, J Immunol. 1991 Jan 1;146(1):362-8).:

45

50

55

```

1 mgllgilcfl iflgktwgqe qtyvisapki frvgaseniv iqvygyteaf datisiksy
61 dkkfsyssgh vhlsenkfq nsailtiqpk qlpggqnpvs yvylevvskh fskskrmpit
121 ydngflfih dkpvypdqs vkvrvyslnd dlkpakretv ltfidpege vdmveeidhi
181 giisfpdfki psnprygmwt ikakykedfs ttgtayfevk eyvlphfsvs iepeynfigy
241 knfknfeiti karyfynkvv teadvyitfg iredlkddqk emmqtamqnt mlingiaqvt
301 fdsetavkel syssledlnn kylyiavtvi estggfseea eipgikyvls pyklnlvatp
361 lflkpgipyp ikvqvkdsld qlvggvpvtl naqtidvnqe tsdldpsksv trvddgvasf
421 vlnlpsgvtv lefnvktdap dlpeenqare gyraiayssl sqsylyidwt dnhkallvge
481 hlniivtpks pyidkithyn ylilskgkii hfgtrekfsd asyqsinipv tqnmvpssrl
541 lvyyivtgeq taelvsdsvw lnieekcgnq lqvhlspdad ayspgqtvsl nmatgmtdswv
601 alaavdsavy gvqrgakkpl ervfqfleks 'dlgcgagggl nnanvhlag ltfltnanad
661 dsqendepck eilrprrtlq kkieeiaky khsvvkkccy dgacvnndet ceqraarisl
721 gprciakafte ccvvasqlra nishkdmqlg rlmktllpv skpeirsyfp eswlwehlv
781 prrkqlqfal pdslltweiq gvgisntgic vadtvkakvf kdvflemnip ysvvrgeqiq
841 lkgtvynyrt sgmqfcvkm s a vegictses pvidhqgtks skcvrpkveg ssshlvftv
901 lpleighni nfsletwfgk eilvktlrvv pegvkresys gtldprgiy gtisrrkefp
961 yripldlvpk teikrilsvk gllvgeilsa vlsqeginil thlpkgsaea elmsvvpvfy
1021 vfhyletgnh wnifhsdpli ekqklkkklk egmlsimssyr nadysysvwk ggsastwlta
1081 falrvlgqvn kyveqgnqnsi cnsllwven yqldngsfke nsqyqpkklq gtlpvareen
1141 slyltaftvi girkafdicp lvkldtalik adnflleentl paqstftlai sayalslgdk
1201 thpqfrsivs alkrealvkg nppiyrfwd nlqhkdsstv ntgtarmvet tayalltsln
1261 lkdinyvnpv ikwlseeqry gggfystqdt inaiegltex sllvkqlrls mdidvsykhk
1321 galhnykmtd knflgrpvev llnddlivst gfgsglatvh vttvvhktst seevcsfylk
1381 idtqdieash yrgygnssdyk rivacasykp sreessssgss havmdislpt gisaneedlk
1441 alvegvdqlf tdyqikdghv ilqlnsipss dflcvrfrif elfevgflsp atftvyeyhr
1501 pdkgctmfys tsnikiqkvc egaackcvea dcggqmeeeld ltisaetrkq tackpeiaya
1561 ykvtsitsitv envfvkykat lldiyktgea vaekdseitf ikkvctctnae lvkgrqyli
1621 gkealqikyn fsfryiypld sltwieywpr dttcscsqaf lanldefaed iflnc

```

30 [0125] The sequence identity of target variants may be determined using BLAST as described below. The terms "sequence identity" in the context of two or more nucleic acid or protein sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection. For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, Adv. Appl. Math. 2: 482 (1981), by the homology alignment algorithm of Needleman & Wunsch, J Mol. Biol. 48: 443 (1970), by the search for similarity method of Pearson & Lipman, Proc. Nat'l. Acad. Sci. USA 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by visual inspection (see generally, Ausubel et al., *infra*).

35 [0126] One example of an algorithm that is suitable for determining percent sequence identity is the algorithm used in the basic local alignment search tool (hereinafter "BLAST"), see, e.g. Altschul et al., J Mol. Biol. 215: 403-410 (1990) and Altschul et al., Nucleic Acids Res., 15: 3389-3402 (1997). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (hereinafter "NCBI"). The default parameters used in determining sequence identity using the software available from NCBI, e.g., BLASTN (for nucleotide sequences) and BLASTP (for amino acid sequences) are described in McGinnis et al., Nucleic Acids Res., 32: W20-W25 (2004).

40 [0127] In another embodiment of the present disclosure, the aptamer has substantially the same ability to bind C5 complement protein as that of an aptamer comprising any one of: SEQ ID NOS 1-2, 5-67, 75-81, 83 or 88-98. In another embodiment, the aptamer has substantially the same structure and ability to bind C5 complement protein as that of an aptamer comprising any one of: SEQ ID NOS 1-2, 5-67, 75-81, 83 or 88-98. In another embodiment, the aptamers have a sequence, including any chemical modifications, according to any one of SEQ ID NOS: 2, 5-67, 75-81, 83 or 88-98. In another embodiment, the aptamers are used as an active ingredient in pharmaceutical compositions. In another embodiment, the aptamers or compositions comprising the aptamers of the present disclosure are used to treat a variety of complement-related diseases or disorders including any one selected from the group consisting of: complement-

related heart disorders (e.g., myocardial injury; C5 mediated complement complications relating to coronary artery bypass graft (CABG) such as post-operative bleeding, systemic neutrophil and leukocyte activation, increased risk of myocardial infarction and increased cognitive dysfunction; restenosis; and C5 mediated complement complications relating to percutaneous coronary intervention), ischemia-reperfusion injury (e.g., myocardial infarction, stroke, frostbite), complement-related inflammatory disorders (e.g., asthma, arthritis, sepsis, and rejection after organ transplantation), and complement-related autoimmune disorders (e.g., myasthenia gravis, systemic lupus erythematosus (SLE), lung inflammation, extra-corporeal complement activation, antibody-mediated complement activation and ocular indications such complement mediated ocular tissue damage such as diabetic retinopathy).

[0128] In one embodiment, the anti-C5 aptamers include a mixture of 2'-fluoro modified nucleotides, 2'-OMe modified nucleotides ("2'-OMe") and 2'-OH purine residues. A descriptive generic sequence (SEQ ID NO: 1) for a modified anti-C5 aptamer is shown below in Table 1, and the structure is shown in Figure 3A. The vast majority of purines (A and G) have been modified to 2'-OMe, excluding only two G residues which remain 2'-OH (residues shown in outline). The circled residues represent a subset of pyrimidines that can be simultaneously modified to 2'-H without substantially altering the anti-C5 activity of the aptamer (see ARC330 in Table 1 below (SEQ ID NO: 2, Figure 3B)). The underlined residues shown in Figure 3A represent pyrimidine residues that can contain either a 2'-fluoro or a 2'-H modification (but not 2'-OMe), while the boxed residues represent pyrimidine residues that can contain either a 2'-fluoro or a 2'-OMe modification (but not 2'-H). The residues indicated with an arrow (→) must contain a 2'-fluoro modification. Without a 2'-fluoro modification at the residues indicated by an arrow (→), resulting hemolytic activity of the resulting aptamer is substantially decreased. In a preferred embodiment, an anti-C5 aptamer of the present disclosure comprises a nucleotide sequence according to SEQ ID NO 1.

[0129] An example of an anti-C5 aptamer according to the present disclosure is ARC186 (SEQ ID NO: 4) which is shown in Figure 3C and described in U.S. Pat. Ser. No. 6,395,888. All 21 pyrimidine residues of ARC186 have 2'-fluoro modifications. The majority of purines (14 residues) have 2'-OMe modifications, except for three 2'-OH purine residues (shown in outline in Figure 3C). The anti-C5 aptamers of the present disclosure can also include different mixtures of 2'-fluoro and 2'-H modifications. For example, another anti-C5 aptamer of the present disclosure is the ARC330 (SEQ ID NO 2) shown in Figure 3B. ARC330 SEQ ID NO 2) contains seven 2'-H modifications (circled residues in Figure 3B), 14 pyrimidine residues with 2'-fluoro modifications, 14 purine residues with 2'-OMe modifications, and three 2'-OH purine residues (shown in outline in Figure 3B).

[0130] Other combinations of aptamers containing a mixture of 2'-fluoro modifications, 2'-OMe modifications, 2'-OH purine residues, and conjugation to non-immunogenic, high molecular weight compounds (e.g., PEG) of varying size, each of which were derived from ARC186 (SEQ ID NO: 4), are described in Table 1 below. The present disclosure provides aptamers as described in Table 1 below as well as aptamers as described below lacking or additionally comprising a 3' cap (e.g., inverted dT).

[0131] Unless indicated otherwise, the nucleotide sequences in Table 1 below are listed in the 5' to 3' direction. For each of the individual sequences in Table 1, all 2'-OMe purine or pyrimidine modifications are indicated by an "m" preceding the corresponding nucleotide; all 2'-fluoro pyrimidine modifications are indicated by an "f" preceding the corresponding nucleotide; all purine or pyrimidine deoxy modifications are indicated by a "d" preceding the corresponding nucleotide; and any purine or pyrimidine appearing without an "m", "f", or "d" preceding the nucleotide indicates a 2'-OH residue.

Table 1:

SEQ ID NO: 1

X₁X₂fCfCrGfCX₃X₄fUX₅X₆X₇X₈X₉X₁₀X₁₁rGX₁₂X₁₃X₁₄X₁₅X₁₆X₁₇X₁₈X₁₉X₂₀X₂₁X₂₂X₂₃fUfUX₂₄X₂₅X₂₆X₂₇X₂₈fCX₂₉

where:

X₁=fC or mC

X₂=rG or mG

X₃=rG or mG

X₄=rG or mG

X₅=fC or dC

X₆=fU or dT

X₇=fC or dC

X₈=rA or mA

X₉=rG or mG

X₁₀=rG or mG

X₁₁= fC or dC

X₁₂=fC or mC

(continued)

5 X₁₃=fU or mU
 X₁₄=rG or mG
 X₁₅=rA or mA
 X₁₆=rG or mG
 X₁₇=fU or dT
 X₁₈=fC or dC
 X₁₉=fU or dT
 10 X₂₀=rG or mG
 X₂₁=rA or mA
 X₂₂=rG or mG
 X₂₃=fU or dT
 15 X₂₄=rA or mA
 X₂₅=fC or dC
 X₂₆=fC or dC
 X₂₇=fU or dT
 20 X₂₈=rG or mG
 X₂₉=rG or mG

ARC330 (SEQ ID NO: 2)

fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

25 ARC185 (SEQ ID NO: 3)
 GAfCGAfUGfCGGfUfCfUfCAfUGfCGfUfCGAGfUGfUGAGfUfUfUAfCfCfUfUfCGfUfC

ARC186 (SEQ ID NO: 4)

30 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-
 3T

ARC187 (SEQ ID NO: 5)

35 40kDa PEG-- NH-
 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-
 3T

Where the branched 40 kDa PEG is ,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide)

40 ARC188 (SEQ ID NO: 6)
 AGGAfCGAfUGfCGGfUfCfUfCAfUGfCGfUfCGAGfUGfUGAGfUfUfUAfCfCfUfUfCGfUfC

ARC189 (SEQ ID NO: 7)

45 AGfCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-
 G

ARC250 (SEQ ID NO: 8)

GGfCGfCfCGfCGGfUfCfUfCAGGfCGfCfUGAGfUfCfUGAGfUfUfUAfCfCfUGfCG

ARC296 (SEQ ID NO: 9)

fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCdCfUmGfCmG-3T

55 ARC297 (SEQ ID NO: 10)
 mCmGmCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCdCfUmGmCmG-
 3T

(continued)

5 ARC331 (SEQ ID NO: 11)
dCmGdCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGdCmG

10 ARC332 (SEQ ID NO: 12)
dCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

15 ARC333 (SEQ ID NO: 13)
fCmGdCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

20 ARC334 (SEQ ID NO: 14)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGdCmG

25 ARC411 (SEQ ID NO: 15)
fCmGmCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

30 ARC412 (SEQ ID NO: 16)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGmCmG

35 ARC413 (SEQ ID NO: 17)
mCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

40 ARC414 (SEQ ID NO: 18)
mCmGmCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGmCmG

45 ARC415 (SEQ ID NO: 19)
fCmGfCdCGfCmGmGfUdCTdCniAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

50 ARC416 (SEQ ID NO: 20)
fCmGfCfCGdCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

55 ARC417 (SEQ ID NO: 21)
fCmGfCdCGdCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

60 ARC418 (SEQ ID NO: 22)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGdCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

65 ARC419 (SEQ ID NO: 23)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCTmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

70 ARC420 (SEQ ID NO: 24)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGdCTmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

75 ARC421 (SEQ ID NO: 25)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

80 ARC422 (SEQ ID NO: 26)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

85 ARC423 (SEQ ID NO: 27)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

(continued)

ARC424 (SEQ ID NO: 28)
5 fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGTTTAfCfCfUmGfCmG

ARC425 (SEQ ID NO: 29)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCTmGfCmG

10 ARC426 (SEQ ID NO: 30)
fCmGfCfCGfCmGmGmUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCdCfUmGfCmG

15 ARC427 (SEQ ID NO: 31)
fCmGfCmCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

ARC428 (SEQ ID NO: 32)
fCmGfCfCGmCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

20 ARC429 (SEQ ID NO: 33)
fCmGfCmCGmCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

25 ARC430 (SEQ ID NO: 34)
fCmGfCfCGfCmGmGfUdCfUdCmAmGmGdCGmCfUmGmAmGfUdCfUmGmAmGfUfUfUAfCfCfUmGfCmG

ARC431 (SEQ ID NO: 35)
fCmGfCfCGfCmGmGfUdCfUdCmAmGmGdCGfCmUmGmAmGfUdCfUmGmAmGfUfUfUAfCfCfUmGfCm

30 ARC432 (SEQ ID NO: 36)
fCmGfCfCGfCmGmGfUdCfUdCmAmGmGdCGmCmUmGmAmGfUdCfUmGmAmGfUfUfUAfCfCfUmGfCmG

35 ARC433 (SEQ ID NO: 37)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGmUfUfUAfCfCfUmGfCmG

ARC434 (SEQ ID NO: 38)
40 fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

ARC435 (SEQ ID NO: 39)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

45 ARC436 (SEQ ID NO: 40)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGmUmUmUAfCfCfUmGfCmG

ARC437 (SEQ ID NO: 41)
50 fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCmUmGfCmG

ARC438 (SEQ ID NO: 42)
fCmGfCfCdGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

55 ARC439 (SEQ ID NO: 43)
fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCdGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG

(continued)

ARC440 (SEQ ID NO: 44)

fCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUdAfCfCfUmGfCmG

5

ARC457 (SEQ ID NO: 45)

mGfCmGfUfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmAfCmG
mC

10

ARC458 (SEQ ID NO: 46)

mGmGmGfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmCmCmC

15

ARC459 (SEQ ID NO: 47)

mGfCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfCmG
mC

20

ARC473 (SEQ ID NO: 48)

mGmGmAfCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUm
GfCmGfUfCfU-3T

25

ARC522 (SEQ ID NO:

49)

mGmGfCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGmCmUmGmAmGTdCTmGmAmGTfUfUAdCdCTmGfC
mGmCmC

30

ARC523 (SEQ ID NO: 50)

mGmGmCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGmCmUmGmAmGTdCTmGmAmGTTAdCdCTmGdC
mGmCmC

35

ARC524 (SEQ ID NO: 51)

mGmGmCmGdCdCGdCmGmGTdCTdCmAmGmGdCGmCmUmGmAmGTdCTmGmAmGTTmAdCdCTmGd
CmGmCmC

40

ARC525 (SEQ ID NO: 52)

mGmGmCmGdCdCGdCmGmGTdCmUmCmAmGmGdCGmCmUmGmAmGmUmCmUmGmAmGTTmAdCd
CTmGdCmGmCmC

45

ARC532 (SEQ ID NO:

53)

Biotin-
AGfCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCm
G

50

ARC543 (SEQ ID NO: 54)

mGmGfCmGfCfCGfCmGmGfUdCTdCmAmGmGdCGfCfUmGmAmGTdCTmGmAmGfUfUfUAfCfCfUmGfC
mGmCmC

55

(continued)

ARC544 (SEQ ID NO:

55)

5 mGmGfCmGfCfCGfCmGmGfUmCmUmCmAmGmGmCGfCfUmGmAmGmUmCmUmGmAmGfUfUfUAfCfCf
UmGfCmGmCmC

ARC550 (SEQ ID NO: 56)

10 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUmAfCfCfUmGfCmG-
3T

ARC551 (SEQ ID NO: 57)

15 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGmCmUmGmAmGfUfCfUmGmAmGfUfUfUfUAfCfCfUmGfCmG-
3T

ARC552 (SEQ ID NO: 58)

20 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-3T

ARC553 (SEQ ID NO: 59)

25 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGmCmUmGmAmGfUfCfUmGmAmGfUfUfUfUmAfCfCfUmGfCmG-
G-3T

25 ARC554 (SEQ ID NO:

60)

30 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGmCmUmGmAmGfUfCfUmGmAmGfUfUfUmAfCfCfUmGfCmG-
-3T

30 ARC 657 (SEQ ID NO: 61)

20 kDa PEG-NH-

35 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUfUAfCfCfUmGfCmG-
3T

ARC 658 (SEQ ID NO: 62)

30 kDa PEG-NH-

40 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUfUAfCfCfUmGfCmG-
3T

ARC 672 (SEQ ID NO: 63)

45 NH2-

45 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-
3T

ARC706 (SEQ ID NO: 64)

50 10 kDa PEG-NH-

50 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUfUAfCfCfUmGfCmG-
3T

55 ARC1537 (SEQ ID NO: 65)

(continued)

40kDa PEG-NH-

fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-
3T

5 ARC1730 (SEQ ID NO: 66)

PEG20K-NH-

fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-
NH-PEG20K

10 ARC1905 (SEQ ID NO: 67)

40K PEG-NH--

fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUAfCfCfUmGfCmG-
3T

15 Where the branched 40 kDa PEG is 2,3-bis(mPEG-[20 kDa])-propyl-1-carbamoyl

20 ARC243 (SEQ ID NO: 68)

GGfCGAfUfUAfCfUGGGAfCGGAfCfUfCGfCGAfUGfUGAGfCfCfCAGAfCGAfCfUfCGfCfC

25 ARC244 (SEQ ID NO: 69)

GGfCfUfUfCfUGAAGAfUfUAfUfUfCGfCGAfUGfUGAAfCfUfCfCAGAfCfCfCfC

30 [0132] Other aptamers that bind complement protein C5 are described below in Example 3.

[0133] In some embodiments aptamer therapeutics of the present disclosure have great affinity and specificity to their targets while reducing the deleterious side effects from non-naturally occurring nucleotide substitutions if the aptamer therapeutics break down in the body of patients or subjects. In some embodiments, the therapeutic compositions containing the aptamer therapeutics of the present disclosure are free of or have a reduced amount of fluorinated nucleotides.

35 [0134] The aptamers of the present disclosure can be synthesized using any oligonucleotide synthesis techniques known in the art including solid phase oligonucleotide synthesis techniques well known in the art (see, e.g., Froehler et al., Nucl. Acid Res. 14:5399-5467 (1986) and Froehler et al., Tet. Lett. 27:5575-5578 (1986)) and solution phase methods such as triester synthesis methods (see, e.g., Sood et al., Nucl. Acid Res. 4:2557 (1977) and Hirose et al., Tet. Lett., 28:2449 (1978)).

Pharmaceutical Compositions

40 [0135] The invention also includes pharmaceutical compositions containing aptamer molecules that bind to complement protein C5. In some embodiments, the compositions are suitable for internal use and include an effective amount of a pharmacologically active compound of the invention, alone or in combination, with one or more pharmaceutically acceptable carriers. The compounds are especially useful in that they have very low, if any toxicity.

45 [0136] Compositions of the invention can be used to treat or prevent a pathology, such as a disease or disorder, or alleviate the symptoms of such disease or disorder in a patient. For example, compositions of the present invention can be used to treat or prevent a pathology associated with complement-related heart disorders (e.g., myocardial injury; C5 mediated complement complications relating to coronary artery bypass graft (CABG) surgery such as post-operative bleeding, systemic neutrophil and leukocyte activation, increased risk of myocardial infarction and increased cognitive dysfunction; restenosis; and C5 mediated complications relating to percutaneous coronary intervention); ischemia-reperfusion injury (e.g., myocardial infarction, stroke, frostbite); complement-related inflammatory disorders (e.g., asthma, arthritis, sepsis, and rejection after organ transplantation); and complement-related autoimmune disorders (e.g., myasthenia gravis, systemic lupus erythematosus (SLE, or lupus); lung inflammation; extracorporeal complement activation; antibody-mediated complement activation; and ocular indications such as diabetic retinopathy. Compositions of the invention are useful for administration to a subject suffering from, or predisposed to, a disease or disorder which is related to or derived from complement protein C5 to which the aptamers of the invention specifically bind.

55 [0137] Compositions of the invention can be used in a method for treating a patient or subject having a pathology. The methods of the present disclosure involve administering to the patient or subject an aptamer or a composition comprising aptamers that bind to complement protein C5, so that binding of the aptamer to complement protein C5 alters

its biological function, thereby treating the pathology.

[0138] The patient or subject having a pathology, *i.e.*, the patient or subject treated by the methods of the present disclosure can be a vertebrate, more particularly a mammal, or more particularly, a human.

[0139] In practice, the aptamers or their pharmaceutically acceptable salts, are administered in amounts which will be sufficient to exert their desired biological activity, *e.g.*, inhibiting the binding of the aptamer target to its receptor, preventing cleavage of a target protein.

[0140] One aspect of the invention comprises an aptamer composition of the invention in combination with other treatments for C5 mediated complement disorders. The aptamer composition of the invention may contain, for example, more than one aptamer. In some examples, an aptamer composition of the invention, containing one or more compounds of the invention, is administered in combination with another useful composition such as an antiinflammatory agent, an immunosuppressant, an antiviral agent, or the like. Furthermore, the compounds of the invention may be administered in combination with a cytotoxic, cytostatic, or chemotherapeutic agent such as an alkylating agent, anti-metabolite, mitotic inhibitor or cytotoxic antibiotic, as described above. In general, the currently available dosage forms of the known therapeutic agents for use in such combinations will be suitable.

[0141] "Combination therapy" (or "co-therapy") includes the administration of an aptamer composition of the invention and at least a second agent as part of a specific treatment regimen intended to provide the beneficial effect from the co-action of these therapeutic agents. The beneficial effect of the combination includes, but is not limited to, pharmacokinetic or pharmacodynamic co-action resulting from the combination of therapeutic agents. Administration of these therapeutic agents in combination typically is carried out over a defined time period (usually minutes, hours, days or weeks depending upon the combination selected).

[0142] "Combination therapy" may, but generally is not, intended to encompass the administration of two or more of these therapeutic agents as part of separate monotherapy regimens that incidentally and arbitrarily result in the combinations of the present disclosure. "Combination therapy" is intended to embrace administration of these therapeutic agents in a sequential manner, that is, wherein each therapeutic agent is administered at a different time, as well as administration of these therapeutic agents, or at least two of the therapeutic agents, in a substantially simultaneous manner. Substantially simultaneous administration can be accomplished, for example, by administering to the subject a single capsule having a fixed ratio of each therapeutic agent or in multiple, single capsules for each of the therapeutic agents.

[0143] Sequential or substantially simultaneous administration of each therapeutic agent can be effected by any appropriate route including, but not limited to, topical routes, oral routes, intravenous routes, intramuscular routes, and direct absorption through mucous membrane tissues. The therapeutic agents can be administered by the same route or by different routes. For example, a first therapeutic agent of the combination selected may be administered by injection while the other therapeutic agents of the combination may be administered topically.

[0144] Alternatively, for example, all therapeutic agents may be administered topically or all therapeutic agents may be administered by injection. The sequence in which the therapeutic agents are administered is not narrowly critical unless noted otherwise. "Combination therapy" also can embrace the administration of the therapeutic agents as described above in further combination with other biologically active ingredients. Where the combination therapy further comprises a non-drug treatment, the non-drug treatment may be conducted at any suitable time so long as a beneficial effect from the co-action of the combination of the therapeutic agents and non-drug treatment is achieved. For example, in appropriate cases, the beneficial effect is still achieved when the non-drug treatment is temporally removed from the administration of the therapeutic agents, perhaps by days or even weeks.

[0145] Therapeutic or pharmacological compositions of the present invention will generally comprise an effective amount of the active component(s) of the therapy, dissolved or dispersed in a pharmaceutically acceptable medium. Pharmaceutically acceptable media or carriers include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Supplementary active ingredients can also be incorporated into the therapeutic compositions of the present invention.

[0146] The preparation of pharmaceutical or pharmacological compositions will be known to those of skill in the art in light of the present disclosure. Typically, such compositions may be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection; as tablets or other solids for oral administration; as time release capsules; or in any other form currently used, including eye drops, creams, lotions, salves, inhalants and the like. The use of sterile formulations, such as saline-based washes, by surgeons, physicians or health care workers to treat a particular area in the operating field may also be particularly useful. Compositions may also be delivered via microdevice, microparticle or sponge.

[0147] Upon formulation, therapeutics will be administered in a manner compatible with the dosage formulation, and in such amount as is pharmacologically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above, but drug release capsules and the like can also be employed.

[0148] In this context, the quantity of active ingredient and volume of composition to be administered depends on the

host animal to be treated. Precise amounts of active compound required for administration depend on the judgment of the practitioner and are peculiar to each individual.

[0149] A minimal volume of a composition required to disperse the active compounds is typically utilized. Suitable regimes for administration are also variable, but would be typified by initially administering the compound and monitoring the results and then giving further controlled doses at further intervals.

[0150] For instance, for oral administration in the form of a tablet or capsule (e.g., a gelatin capsule), the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, magnesium aluminum silicate, starch paste, gelatin, methylcellulose, sodium carboxymethylcellulose and/or polyvinylpyrrolidone, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, silica, talcum, stearic acid, its magnesium or calcium salt and/or polyethyleneglycol and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum starches, agar, alginic acid or its sodium salt, or effervescent mixtures, and the like. Diluents, include, e.g., lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine.

[0151] Injectable compositions are preferably aqueous isotonic solutions or suspensions, and suppositories are advantageously prepared from fatty emulsions or suspensions. The compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. The compositions are prepared according to conventional mixing, granulating or coating methods, respectively, and typically contain about 0.1 to 75%, preferably about 1 to 50%, of the active ingredient.

[0152] The compounds of the invention can also be administered in such oral dosage forms as timed release and sustained release tablets or capsules, pills, powders, granules, elixirs, tinctures, suspensions, syrups and emulsions.

[0153] Liquid, particularly injectable compositions can, for example, be prepared by dissolving, dispersing, etc. The active compound is dissolved in or mixed with a pharmaceutically pure solvent such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, to thereby form the injectable solution or suspension. Additionally, solid forms suitable for dissolving in liquid prior to injection can be formulated.

[0154] The compounds of the present invention can be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions.

[0155] Parenteral injectable administration is generally used for subcutaneous, intramuscular or intravenous injections and infusions. Additionally, one approach for parenteral administration employs the implantation of a slow-release or sustained-released systems, which assures that a constant level of dosage is maintained, according to U.S. Pat. No. 3,710,795.

[0156] Furthermore, preferred compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, inhalants, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Other preferred topical preparations include creams, ointments, lotions, aerosol sprays and gels, wherein the concentration of active ingredient would typically range from 0.01% to 15%, w/w or w/v.

[0157] For solid compositions, excipients include pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like may be used. The active compound defined above, may be also formulated as suppositories using for example, polyalkylene glycols, for example, propylene glycol, as the carrier. In some embodiments, suppositories are advantageously prepared from fatty emulsions or suspensions.

[0158] The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, containing cholesterol, stearylamine or phosphatidylcholines. In some embodiments, a film of lipid components is hydrated with an aqueous solution of drug to a form lipid layer encapsulating the drug, as described in U.S. Pat. No. 5,262,564. For example, the aptamer molecules described herein can be provided as a complex with a lipophilic compound or non-immunogenic, high molecular weight compound constructed using methods known in the art. An example of nucleic-acid associated complexes is provided in U.S. Patent No. 6,011,020.

[0159] The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropyl-methacrylamide-phenol, polyhydroxyethylaspanamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters,

polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

[0160] If desired, the pharmaceutical composition to be administered may also contain minor amounts of non-toxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and other substances such as for example, sodium acetate, and triethanolamine oleate.

5 [0161] The dosage regimen utilizing the aptamers is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular aptamer or salt thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

10 [0162] Oral dosages of the present disclosure, when used for the indicated effects, will range between about 0.05 to 7500 mg/day orally. The compositions are preferably provided in the form of scored tablets containing 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100.0, 250.0, 500.0 and 1000.0 mg of active ingredient. Compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily.

15 [0163] Infused dosages, intranasal dosages and transdermal dosages will range between 0.05 to 7500 mg/day. Subcutaneous, intravenous and intraperitoneal dosages will range between 0.05 to 3800 mg/day.

[0164] Effective plasma levels of the compounds of the present invention range from 0.002 mg/mL to 50 mg/mL.

Modulation of pharmacokinetics and biodistribution of aptamer therapeutics

20 [0165] It is important that the pharmacokinetic properties for all oligonucleotide-based therapeutics, including aptamers, be tailored to match the desired pharmaceutical application. While aptamers directed against extracellular targets do not suffer from difficulties associated with intracellular delivery (as is the case with antisense and RNAi-based therapeutics), such aptamers must still be able to be distributed to target organs and tissues, and remain in the body (unmodified) for a period of time consistent with the desired dosing regimen.

25 [0166] Thus, the present disclosure provides materials and methods to affect the pharmacokinetics of aptamer compositions, and, in particular, the ability to tune aptamer pharmacokinetics. The tunability of (i.e., the ability to modulate) aptamer pharmacokinetics is achieved through conjugation of modifying moieties (e.g., PEG polymers) to the aptamer and/or the incorporation of modified nucleotides (e.g., 2'-fluoro or 2'-OMe) to alter the chemical composition of the nucleic acid. The ability to tune aptamer pharmacokinetics is used in the improvement of existing therapeutic applications, or alternatively, in the development of new therapeutic applications. For example, in some therapeutic applications, e.g., in antineoplastic or acute care settings where rapid drug clearance or turn-off may be desired, it is desirable to decrease the residence times of aptamers in the circulation. Alternatively, in other therapeutic applications, e.g., maintenance therapies where systemic circulation of a therapeutic is desired, it may be desirable to increase the residence times of aptamers in circulation.

30 [0167] In addition, the tunability of aptamer pharmacokinetics is used to modify the biodistribution of an aptamer therapeutic in a subject. For example, in some therapeutic applications, it may be desirable to alter the biodistribution of an aptamer therapeutic in an effort to target a particular type of tissue or a specific organ (or set of organs). In these applications, the aptamer therapeutic preferentially accumulates in a specific tissue or organ(s). In other therapeutic applications, it may be desirable to target tissues displaying a cellular marker or a symptom associated with a given disease, cellular injury or other abnormal pathology, such that the aptamer therapeutic preferentially accumulates in the affected tissue. For example, as described in copending provisional application United States Serial No. 60/550790, filed on March 5, 2004, and entitled "Controlled Modulation of the Pharmacokinetics and Biodistribution of Aptamer Therapeutics", PEGylation of an aptamer therapeutic (e.g., PEGylation with a 20 kDa PEG polymer) is used to target inflamed tissues, such that the PEGylated aptamer therapeutic preferentially accumulates in inflamed tissue.

35 [0168] To determine the pharmacokinetic and biodistribution profiles of aptamer therapeutics (e.g., aptamer conjugates or aptamers having altered chemistries, such as modified nucleotides) a variety of parameters are monitored. Such parameters include, for example, the half-life ($t_{1/2}$), the plasma clearance (C1), the volume of distribution (Vss), the area under the concentration-time curve (AUC), maximum observed serum or plasma concentration (C_{max}), and the mean residence time (MRT) of an aptamer composition. As used herein, the term "AUC" refers to the area under the plot of the plasma concentration of an aptamer therapeutic versus the time after aptamer administration. The AUC value is used to estimate the bioavailability (i.e., the percentage of administered aptamer therapeutic in the circulation after aptamer administration) and/or total clearance (C 1) (i.e., the rate at which the aptamer therapeutic is removed from circulation) of a given aptamer therapeutic. The volume of distribution relates the plasma concentration of an aptamer therapeutic to the amount of aptamer present in the body. The larger the Vss, the more an aptamer is found outside of the plasma (i.e., the more extravasation).

40 [0169] The present disclosure provides materials and methods to modulate, in a controlled manner, the pharmacokinetics and biodistribution of stabilized aptamer compositions *in vivo* by conjugating an aptamer to a modulating moiety

such as a small molecule, peptide, or polymer terminal group, or by incorporating modified nucleotides into an aptamer. As described herein, conjugation of a modifying moiety and/or altering nucleotide(s) chemical composition alters fundamental aspects of aptamer residence time in circulation and distribution to tissues.

[0170] In addition to clearance by nucleases, oligonucleotide therapeutics are subject to elimination *via* renal filtration. As such, a nuclease-resistant oligonucleotide administered intravenously typically exhibits an *in vivo* half-life of <10 min, unless filtration can be blocked. This can be accomplished by either facilitating rapid distribution out of the blood stream into tissues or by increasing the apparent molecular weight of the oligonucleotide above the effective size cut-off for the glomerulus. Conjugation of small therapeutics to a PEG polymer (PEGylation), described below, can dramatically lengthen residence times of aptamers in circulation, thereby decreasing dosing frequency and enhancing effectiveness against vascular targets.

[0171] Aptamers can be conjugated to a variety of modifying moieties, such as high molecular weight polymers, e.g., PEG; peptides, e.g., Tat (a 13-amino acid fragment of the HIV Tat protein (Vives, et al. (1997), *J. Biol. Chem.* 272(25): 16010-7)), Ant (a 16-amino acid sequence derived from the third helix of the *Drosophila* *antennapedia* homeotic protein (Pietersz, et al. (2001), *Vaccine* 19(11-12): 1397-405)) and Arg₇ (a short, positively charged cell-permeating peptides composed of polyarginine (Arg₇) (Rothbard, et al. (2000), *Nat. Med.* 6(11): 1253-7; Rothbard, J et al. (2002), *J. Med. Chem.* 45(17): 3612-8); and small molecules, e.g., lipophilic compounds such as cholesterol. Among the various conjugates described herein, *in vivo* properties of aptamers are altered most profoundly by complexation with PEG groups. For example, complexation of a mixed 2'F and 2'-OMe modified aptamer therapeutic with a 20 kDa PEG polymer hinders renal filtration and promotes aptamer distribution to both healthy and inflamed tissues. Furthermore, the 20 kDa PEG polymer-aptamer conjugate proves nearly as effective as a 40 kDa PEG polymer in preventing renal filtration of aptamers. While one effect of PEGylation is on aptamer clearance, the prolonged systemic exposure afforded by presence of the 20 kDa moiety also facilitates distribution of aptamer to tissues, particularly those of highly perfused organs and those at the site of inflammation. The aptamer-20 kDa PEG polymer conjugate directs aptamer distribution to the site of inflammation, such that the PEGylated aptamer preferentially accumulates in inflamed tissue. In some instances, the 20 kDa PEGylated aptamer conjugate is able to access the interior of cells, such as, for example, kidney cells.

[0172] Overall, effects on aptamer pharmacokinetics and tissue distribution produced by low molecular weight modifying moieties, including cholesterol and cell-permeating peptides are less pronounced than those produced as a result of PEGylation or modification of nucleotides (e.g., an altered chemical composition). While not intending to be bound by theory, it is suggested that cholesterol-mediated associations with plasma lipoproteins, postulated to occur in the case of the antisense conjugate, are precluded in the particular context of the aptamer-cholesterol conjugate folded structure, and/or relate to aspect of the lipophilic nature of the cholesterol group. Like cholesterol, the presence of a Tat peptide tag promotes clearance of aptamer from the blood stream, with comparatively high levels of conjugate appearing in the kidneys at 48 hrs. Other peptides (e.g., Ant, Arg₇) that have been reported in the art to mediate passage of macromolecules across cellular membranes *in vitro*, do not appear to promote aptamer clearance from circulation. However, like Tat, the Ant conjugate significantly accumulates in the kidneys relative to other aptamers. While not intending to be bound by theory, it is possible that unfavorable presentation of the Ant and Arg₇ peptide modifying moieties in the context of three dimensionally folded aptamers *in vivo* impairs the ability of these peptides to influence aptamer transport properties.

[0173] Modified nucleotides can also be used to modulate the plasma clearance of aptamers. For example, an unconjugated aptamer which incorporates both 2'-F and 2'-OMe stabilizing chemistries, which is typical of current generation aptamers as it exhibits a high degree of nuclease stability *in vitro* and *in vivo*, displays rapid loss from plasma (i.e., rapid plasma clearance) and a rapid distribution into tissues, primarily into the kidney, when compared to unmodified aptamer.

PEG-Derivatized Nucleic Acids

[0174] As described above, derivatization of nucleic acids with high molecular weight non-immunogenic polymers has the potential to alter the pharmacokinetic and pharmacodynamic properties of nucleic acids making them more effective therapeutic agents. Favorable changes in activity can include increased resistance to degradation by nucleases, decreased filtration through the kidneys, decreased exposure to the immune system, and altered distribution of the therapeutic through the body.

[0175] The aptamer compositions of the invention may be derivatized with polyalkylene glycol ("PAG") moieties. Examples of PAG-derivatized nucleic acids are found in United States Patent Application Ser. No. 10/718,833, filed on November 21, 2003. Typical polymers used in the present disclosure include poly(ethylene glycol) ("PEG"), also known as poly(ethylene oxide) ("PEO") and polypropylene glycol (including poly isopropylene glycol). Additionally, random or block copolymers of different alkylene oxides (e.g., ethylene oxide and propylene oxide) can be used in many applications. In its most common form, a polyalkylene glycol, such as PEG, is a linear polymer terminated at each end with hydroxyl groups: HO-CH₂CH₂O-(CH₂CH₂O) n-CH₂CH₂-OH. This polymer, alpha-, omega-dihydroxylpoly(ethylene glycol), can also be represented as HO-PEG-OH, where it is understood that the -PEG- symbol represents the following structural

unit: $-\text{CH}_2\text{CH}_2\text{O}-(\text{CH}_2\text{CH}_2\text{O})_n-\text{CH}_2\text{CH}_2-$ where n typically ranges from about 4 to about 10,000.

[0176] As shown, the PEG molecule is di-functional and is sometimes referred to as "PEG diol." The terminal portions of the PEG molecule are relatively non-reactive hydroxyl moieties, the -OH groups, that can be activated, or converted to functional moieties, for attachment of the PEG to other compounds at reactive sites on the compound. Such activated PEG diols are referred to herein as bi-activated PEGs. For example, the terminal moieties of PEG diol have been functionalized as active carbonate ester for selective reaction with amino moieties by substitution of the relatively non-reactive hydroxyl moieties, -OH, with succinimidyl active ester moieties from N-hydroxy succinimide.

[0177] In many applications, it is desirable to cap the PEG molecule on one end with an essentially non-reactive moiety so that the PEG molecule is mono-functional (or mono-activated). In the case of protein therapeutics which generally display multiple reaction sites for activated PEGs, bi-functional activated PEGs lead to extensive cross-linking, yielding poorly functional aggregates. To generate mono-activated PEGs, one hydroxyl moiety on the terminus of the PEG diol molecule typically is substituted with non-reactive methoxy end moiety, $-\text{OCH}_3$. The other, un-capped terminus of the PEG molecule typically is converted to a reactive end moiety that can be activated for attachment at a reactive site on a surface or a molecule such as a protein.

[0178] PAGs are polymers which typically have the properties of solubility in water and in many organic solvents, lack of toxicity, and lack of immunogenicity. One use of PAGs is to covalently attach the polymer to insoluble molecules to make the resulting PAG-molecule "conjugate" soluble. For example, it has been shown that the water-insoluble drug paclitaxel, when coupled to PEG, becomes water-soluble. Greenwald, et al., *J. Org. Chem.*, 60:331-336 (1995). PAG conjugates are often used not only to enhance solubility and stability but also to prolong the blood circulation half-life of molecules.

[0179] Polyalkylated compounds of the present disclosure are typically between 5 and 80 kD in size however any size can be used, the choice dependent on the aptamer and application. Other PAG compounds of the present disclosure are between 10 and 80 kD in size. Still other PAG compounds of the present disclosure are between 10 and 60 kD in size. For example, a PAG polymer may be at least 10, 20, 30, 40, 50, 60, or 80 kD in size. Such polymers can be linear or branched.

[0180] In contrast to biologically-expressed protein therapeutics, nucleic acid therapeutics are typically chemically synthesized from activated monomer nucleotides. PEG-nucleic acid conjugates may be prepared by incorporating the PEG using the same iterative monomer synthesis. For example, PEGs activated by conversion to a phosphoramidite form can be incorporated into solid-phase oligonucleotide synthesis. Alternatively, oligonucleotide synthesis can be completed with site-specific incorporation of a reactive PEG attachment site. Most commonly this has been accomplished by addition of a free primary amine at the 5'-terminus (incorporated using a modified phosphoramidite in the last coupling step of solid phase synthesis). Using this approach, a reactive PEG (e.g., one which is activated so that it will react and form a bond with an amine) is combined with the purified oligonucleotide and the coupling reaction is carried out in solution. In some embodiment the polymers are branched PEG molecules. In still other embodiments the polymers are 40 kDa branched PEG, see, e.g. (1,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide) depicted in Figure 4. In some embodiments the 40 kD branched PEG (1,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide) is attached to the 5' end of the aptamer as depicted in Figure 5.

[0181] The ability of PEG conjugation to alter the biodistribution of a therapeutic is related to a number of factors including the apparent size (e.g., as measured in terms of hydrodynamic radius) of the conjugate. Larger conjugates (>10kDa) are known to more effectively block filtration via the kidney and to consequently increase the serum half-life of small macromolecules (e.g., peptides, antisense oligonucleotides). The ability of PEG conjugates to block filtration has been shown to increase with PEG size up to approximately 50 kDa (further increases have minimal beneficial effect as half life becomes defined by macrophage-mediated metabolism rather than elimination via the kidneys).

[0182] Production of high molecular weight PEGs (>10 kDa) can be difficult, inefficient, and expensive. As a route towards the synthesis of high molecular weight PEG-nucleic acid conjugates, previous work has been focused towards the generation of higher molecular weight activated PEGs. One method for generating such molecules involves the formation of a branched activated PEG in which two or more PEGs are attached to a central core carrying the activated group. The terminal portions of these higher molecular weight PEG molecules, i.e., the relatively non-reactive hydroxyl (-OH) moieties, can be activated, or converted to functional moieties, for attachment of one or more of the PEGs to other compounds at reactive sites on the compound. Branched activated PEGs will have more than two termini, and in cases where two or more termini have been activated, such activated higher molecular weight PEG molecules are referred to herein as, multi-activated PEGs. In some cases, not all termini in a branch PEG molecule are activated. In cases where any two termini of a branch PEG molecule are activated, such PEG molecules are referred to as bi-activated PEGs. In some cases where only one terminus in a branch PEG molecule is activated, such PEG molecules are referred to as mono-activated. As an example of this approach, activated PEG prepared by the attachment of two monomethoxy PEGs to a lysine core which is subsequently activated for reaction has been described (Harris et al., *Nature*, vol.2: 214-221, 2003).

[0183] The present disclosure provides another cost effective route to the synthesis of high molecular weight PEG-

nucleic acid (preferably, aptamer) conjugates including multiply PEGylated nucleic acids. The present disclosure also encompasses PEG-linked multimeric oligonucleotides, e.g., dimerized aptamers. The present disclosure also relates to high molecular weight compositions where a PEG stabilizing moiety is a linker which separates different portions of an aptamer, e.g., the PEG is conjugated within a single aptamer sequence, such that the linear arrangement of the high molecular weight aptamer composition is, e.g., nucleic acid - PEG - nucleic acid (- PEG - nucleic acid)_n where n is greater than or equal to 1.

[0184] High molecular weight compositions of the present disclosure include those having a molecular weight of at least 10 kD. Compositions typically have a molecular weight between 10 and 80 kD in size. High molecular weight compositions of the are at least 10, 20, 30, 40, 50, 60, or 80 kD in size.

[0185] A stabilizing moiety is a molecule, or portion of a molecule, which improves pharmacokinetic and pharmacodynamic properties of the high molecular weight aptamer compositions of the present disclosure. In some cases, a stabilizing moiety is a molecule or portion of a molecule which brings two or more aptamers, or aptamer domains, into proximity, or provides decreased overall rotational freedom of the high molecular weight aptamer compositions of the present disclosure. A stabilizing moiety can be a polyalkylene glycol, such a polyethylene glycol, which can be linear or branched, a homopolymer or a heteropolymer. Other stabilizing moieties include polymers such as peptide nucleic acids (PNA). Oligonucleotides can also be stabilizing moieties; such oligonucleotides can include modified nucleotides, and/or modified linkages, such as phosphorothioates. A stabilizing moiety can be an integral part of an aptamer composition, i.e., it is covalently bonded to the aptamer.

[0186] Compositions of the present disclosure include high molecular weight aptamer compositions in which two or more nucleic acid moieties are covalently conjugated to at least one polyalkylene glycol moiety. The polyalkylene glycol moieties serve as stabilizing moieties. In compositions where a polyalkylene glycol moiety is covalently bound at either end to an aptamer, such that the polyalkylene glycol joins the nucleic acid moieties together in one molecule, the polyalkylene glycol is said to be a linking moiety. In such compositions, the primary structure of the covalent molecule includes the linear arrangement nucleic acid-PAG-nucleic acid. One example is a composition having the primary structure nucleic acid-PEG-nucleic acid. Another example is a linear arrangement of: nucleic acid - PEG - nucleic acid-PEG - nucleic acid.

[0187] To produce the nucleic acid-PEG nucleic acid conjugate, the nucleic acid is originally synthesized such that it bears a single reactive site (e.g., it is mono-activated). In a preferred embodiment, this reactive site is an amino group introduced at the 5'-terminus by addition of a modified phosphoramidite as the last step in solid phase synthesis of the oligonucleotide. Following deprotection and purification of the modified oligonucleotide, it is reconstituted at high concentration in a solution that minimizes spontaneous hydrolysis of the activated PEG. In a preferred embodiment, the concentration of oligonucleotide is 1 mM and the reconstituted solution contains 200 mM NaHCO₃-buffer, pH 8.3. Synthesis of the conjugate is initiated by slow, step-wise addition of highly purified bi-functional PEG. In a preferred embodiment, the PEG diol is activated at both ends (bi-activated) by derivatization with succinimidyl propionate. Following reaction, the PEG-nucleic acid conjugate is purified by gel electrophoresis or liquid chromatography to separate fully-, partially-, and un-conjugated species. Multiple PAG molecules concatenated (e.g., as random or block copolymers) or smaller PAG chains can be linked to achieve various lengths (or molecular weights). Non-PAG linkers can be used between PAG chains of varying lengths.

[0188] The 2'-OMe, 2'-fluoro and other modified nucleotide modifications stabilize the aptamer against nucleases and increase its half life *in vivo*. The 3'-3'-dT cap also increases exonuclease resistance. See, e.g., U.S. Patents 5,674,685; 5,668,264; 6,207,816; and 6,229,002.

PAG-derivatization of a reactive nucleic acid

[0189] High molecular weight PAG-nucleic acid-PAG conjugates can be prepared by reaction of a mono-functional activated PEG with a nucleic acid containing more than one reactive site. In one embodiment, the nucleic acid is bi-reactive, or bi-activated, and contains two reactive sites: a 5'-amino group and a 3'-amino group introduced into the oligonucleotide through conventional phosphoramidite synthesis, for example: 3'-5'-di-PEGylation as illustrated in Figure 6. In alternative embodiments, reactive sites can be introduced at internal positions, using for example, the 5-position of pyrimidines, the 8-position of purines, or the 2'-position of ribose as sites for attachment of primary amines. In such embodiments, the nucleic acid can have several activated or reactive sites and is said to be multiply activated. Following synthesis and purification, the modified oligonucleotide is combined with the mono-activated PEG under conditions that promote selective reaction with the oligonucleotide reactive sites while minimizing spontaneous hydrolysis. In the preferred embodiment, monomethoxy-PEG is activated with succinimidyl propionate and the coupled reaction is carried out at pH 8.3. To drive synthesis of the bi-substituted PEG, stoichiometric excess PEG is provided relative to the oligonucleotide. Following reaction, the PEG-nucleic acid conjugate is purified by gel electrophoresis or liquid chromatography to separate fully-, partially-, and un-conjugated species.

[0190] The linking domains can also have one or more polyalkylene glycol moieties attached thereto. Such PAGs can

be of varying lengths and may be used in appropriate combinations to achieve the desired molecular weight of the composition.

[0191] The effect of a particular linker can be influenced by both its chemical composition and length. A linker that is too long, too short, or forms unfavorable steric and/or ionic interactions with the target will preclude the formation of complex between aptamer and target. A linker, which is longer than necessary to span the distance between nucleic acids, may reduce binding stability by diminishing the effective concentration of the ligand. Thus, it is often necessary to optimize linker compositions and lengths in order to maximize the affinity of an aptamer to a target.

EXAMPLE 1

Anti-C5 Aptamer Activity in the Classical and Alternative Complement Pathways

Example 1A: Hemolysis Assay.

[0192] The CH50 test measures the ability of the complement system in a serum test sample to lyse 50% of cells in a standardized suspension of antibody-coated sheep erythrocytes. A solution of 0.2% human serum was mixed with antibody-coated sheep erythrocytes (Diamedix EZ Complement CH50 Kit, Diamedix Corp., Miami, FL) in the presence or absence of various anti-C5 aptamers. The assay was run according to the kit protocol in veronal-buffered saline containing calcium, magnesium and 1% gelatin (GVB⁺⁺ complement buffer) and incubated for 30 minutes at 37 °C. After incubation, the samples were centrifuged to pellet intact erythrocytes. The optical density at 412 nm (OD₄₁₂) of the supernatant was read to quantify the release of soluble hemoglobin, which is proportional to the extent of hemolysis (Green et al., (1995) *Chem. Biol.* 2:683-95). To verify that the aptamers blocked C5 activation, some hemolysis supernatants were analyzed for the presence of C5a and C5b-9 by ELISA (C5b-9 ELISA kit, Quidel, San Diego, CA; C5a ELISA kit, BD Biosciences, San Diego, CA).

[0193] The addition of a non-PEGylated anti-C5 aptamer (ARC 186) (SEQ ID NO: 4) to the reaction mixture inhibited hemolysis in a dose-dependent manner, as shown in the graph of Figure 7A, with an IC₅₀ of 0.5 ± 0.1 nM, (see Figure 7B), a value that is consistent with the K_D determined by nitrocellulose filtration. At very high aptamer concentrations (>10 nM), the extent of hemolysis was essentially indistinguishable from background (no serum added), indicating that ARC186 (SEQ ID NO: 4) was able to completely block complement activity. Conjugation of the ARC186 (SEQ ID NO: 4) aptamer with 20 kDa (ARC657; SEQ ID NO: 61), 30 kDa (ARC658; SEQ ID NO: 62), branched 40 kDa (1,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide) (ARC187; SEQ ID NO: 5), branched 40 kDa (2,3-bis(mPEG-[20 kDa])-propyl-1-carbamoyl) (ARC1905; SEQ ID NO: 67), linear 40 kDa (ARC1537; SEQ ID NO: 65), and linear 2x20 kDa (ARC1730; SEQ ID NO: 66) PEG groups had little effect on the aptamer inhibitory activity (Figure 7A-Figure 7D). Finally, ELISA analysis of hemolysis supernatants indicated that this functional inhibition correlated with blockade of C5a release. Thus, the hemolysis data show that ARC186 (SEQ ID NO: 4), and its PEGylated conjugates, are highly potent complement inhibitors that function by blocking the convertase-catalyzed activation of C5.

[0194] Hemolysis assays with non-PEGylated material indicated that the anti-C5 aptamer does not cross-react with C5 from a number of non-primate species, including rat, guinea pig, dog and pig. However, significant inhibitory activity was observed in screens of primate serum, including serum from cynomolgus macaque, rhesus macaque and chimpanzee. The *in vitro* efficacy of the anti-C5 aptamer was further investigated in cynomolgus serum using ARC658 (SEQ ID NO: 62), the 30 kDa-PEG analogue of ARC186 (SEQ ID NO: 4). In a side-by-side comparison (n = 3), ARC658 inhibited human complement activity with an IC₅₀ of 0.21 ± 0.0 nM and cynomolgus complement activity with an IC₅₀ of 1.7 ± 0.4 nM (Figure 8). Thus ARC658 (SEQ ID NO: 62) is 8 ± 3 fold less potent in cynomolgus serum compared to human by this measure.

[0195] **Nitrocellulose Filter Binding Assays.** Individual aptamers were ³²P-labeled at the 5' end by incubation with γ -³²P-ATP and polynucleotide kinase (New England Biolabs, Beverly, MA). Radiolabeled aptamer was purified away from free ATP by gel-filtration followed by polyacrylamide gel electrophoresis. To measure anti-C5 aptamer affinity, radiolabeled aptamer (\leq 10 pM) was incubated with increasing concentrations (0.05 - 100 nM) of purified C5 protein (Quidel, San Diego, CA) in phosphate-buffered saline containing 1 mM MgCl₂ at room temperature (23°C) and 37°C, for 15 min and 4 hr time intervals. The binding reactions were analyzed by nitrocellulose filtration using a Minifold I dot-blot, 96-well vacuum filtration manifold (Schleicher & Schuell, Keene, NH). A three-layer filtration medium was used, consisting (from top to bottom) of Protran nitrocellulose (Schleicher & Schuell), Hybond-P nylon (Amersham Biosciences, Piscataway, NJ) and GB002 gel blot paper (Schleicher & Schuell). The nitrocellulose layer, which selectively binds protein over nucleic acid, preferentially retained the anti-C5 aptamer in complex with a protein ligand, while non-complexed anti-C5 aptamer passed through the nitrocellulose and adhered to the nylon. The gel blot paper was included simply as a supporting medium for the other filters. Following filtration, the filter layers were separated, dried and exposed on a phosphor screen (Amersham Biosciences) and quantified using a Storm 860 Phosphorimager® blot imaging system (Amersham Biosciences).

[0196] As shown in Figure 9 and Figure 10, increasing C5 concentrations enhance the proportion of ARC 186 captured on the nitrocellulose membrane. The dependence of bound ARC 186 on increasing C5 concentrations is well-described by a single-site binding model ($C5 + ARC186 \leftrightarrow C5 \cdot ARC186$; % bound = $C_{max} / (1 + K_D / [C5])$; C_{max} is the maximum % bound at saturating [C5]; K_D is the dissociation constant). ARC186 binding curves at two temperatures following either a 15 min or a 4 hr incubation are shown in Figures 9 and 10, respectively. Following a 15 min incubation, the ARC186 binding curves at 23 and 37°C are essentially indistinguishable within error, fitting with K_D values of 0.5 - 0.6 nM (Figure 9). Differences between binding curves at 23 and 37°C become more pronounced when the incubation time is extended. Following a 4 hr incubation (Figure 10), the K_D observed at 23°C. decreases to 0.08 ± 0.01 nM, while the K_D observed at 37°C remains unchanged (0.6 ± 0.1 nM).

[0197] To demonstrate the basis for the long incubation requirement at room temperature, the affinity at this temperature was further explored using kinetic methods. The rate of the reverse reaction describing the dissociation of $C5 \cdot ARC186$ is $v_{rev} = k_{-1}[C5 \cdot ARC186]$, where v_{rev} is the rate (units of $M \text{ min}^{-1}$) and k_{-1} is the first order dissociation rate constant (units of min^{-1}). The rate of the forward reaction describing the formation of the $C5 \cdot ARC186$ complex is $v_{for} = k_1[C5][ARC186]$, where v_{for} is the rate (units of $M \text{ min}^{-1}$) and k_1 is the second order association rate constant (units of $M^{-1}\text{min}^{-1}$). The data were analyzed using the pseudo-first order assumption, where the concentration of one reactant (C5 in this case) is held in vast excess over the other ($[C5] \gg [ARC186]$, and thus remains essentially unchanged over the course of the reaction. Under these conditions, the forward reaction is described by the rate equation for a first order process, $v_{for} = k_1'[ARC186]$, where $k_1' = k_1[C5]$.

[0198] To analyze dissociation of $C5 \cdot ARC186$, radiolabeled ARC186 (≤ 10 pM) was pre-equilibrated with 5 nM C5 protein in phosphate-buffered saline containing 1 mM $MgCl_2$ at room temperature (23 °C). The dissociation reaction was initiated by the addition of non-labeled ARC186 (1 μ M), which acts as a trap for free C5, and stopped by nitrocellulose filtration partitioning of bound and free radiolabeled ARC 186. A timecourse of ARC 186 dissociation was obtained by varying the duration between initiation of the dissociation reaction and filtration. The timecourse of dissociation, observed as a decrease in the percentage of radiolabeled ARC186 captured on the nitrocellulose filter (equal to the percent bound to C5), is well-described by a single-exponential decay where $\% \text{ ARC186 bound} = 100 \times e^{-k_{-1}t}$ (see Figure 11). The value of the dissociation rate constant, k_{-1} , determined by this method is $0.013 \pm 0.02 \text{ min}^{-1}$, corresponding to a half-life ($t_{1/2} = \ln 2 / k_{-1}$) of 53 ± 8 min.

[0199] To analyze the association reaction, the equilibration rate constant (k_{eq}) for the formation of $C5 \cdot ARC186$ was measured in the presence of varying concentrations of C5 protein (1- 5 nM). Complex formation was initiated by mixing together C5 protein and radiolabeled ARC186 in PBS containing 1 mM $MgCl_2$ at room temperature (23°C), and stopped by nitrocellulose filtration partitioning. As described for the dissociation reactions, a timecourse of complex formation was obtained by varying the duration between the initiation of the reaction and filtration. The timecourse of equilibration, observed as an increase in the percentage of radiolabeled ARC186 captured on the nitrocellulose filter, is well described by a single-exponential decay where $\% \text{ ARC186 bound} = 100 \times (1 - e^{-k_{eq}t})$. The timecourses of equilibration for 1, 2 and 4 nM C5 are displayed in Figure 12. As expected, the value of k_{eq} increases linearly with [C5] (k_{eq} (1 nM) = $0.19 \pm 0.02 \text{ min}^{-1}$; k_{eq} (2 nM) = $0.39 \pm 0.03 \text{ min}^{-1}$; k_{eq} (3 nM) = $0.59 \pm 0.05 \text{ min}^{-1}$; k_{eq} (4 nM) = $0.77 \pm 0.06 \text{ min}^{-1}$; k_{eq} (5 nM) = $0.88 \pm 0.06 \text{ min}^{-1}$). Under the conditions of the experiment, the relationship between k_{eq} , k_1 and k_{-1} is $k_{eq} = k_1[C5] + k_{-1}$. Thus, an estimate of k_1 is derived from the slope of a plot of k_{eq} versus [C5] (see Figure 12 inset), in this case $0.18 \pm 0.01 \text{ nM}^{-1}\text{min}^{-1}$.

[0200] These data indicate that, under conditions of low C5 concentration (e.g., 0.1 nM), an extended incubation is required in order for the mixture of C5 and radiolabeled ARC186 to reach equilibrium. Under these conditions, $k_{eq} = (0.18 \pm 0.01 \text{ nM}^{-1}\text{min}^{-1}) (0.1 \text{ nM}) + 0.013 \text{ min}^{-1} = 0.03 \text{ min}^{-1}$, corresponding to a half-life of 22 min. Thus, nearly 2 hours of room temperature incubation (- 5 half-lives) are required for complete (> 95%) equilibration. A short incubation time (e.g., 15 min) will significantly underestimate the actual affinity of the complex, as shown above by the difference in affinities observed for a 15 min ($K_D = 0.5$ nM) versus a 4 hour ($K_D = 0.08$ nM) incubation. An alternative estimate of the room temperature K_D can be calculated from the kinetic data according to the relationship $K_D = k_{-1} / k_1$. In this case, the calculated K_D is 0.07 ± 0.01 nM, which is completely consistent with the K_D determined above by thermodynamic methods.

[0201] The specificity of ARC 186 (SEQ ID NO: 4) for C5 was also assessed in nitrocellulose filtration assays by comparison with complement components both upstream and downstream from C5 in the complement cascade. The results are depicted in Figure 13. The aptamer essentially does not recognize C5a ($K_D \gg 3 \mu\text{M}$), although it does display weak affinity for soluble C5b-9 ($K_D > 0.2 \mu\text{M}$), presumably due to interactions with the C5b component. Other complement components display moderate to weak affinity for the aptamer. Non-activated C3 essentially does not bind to the aptamer; however, factor H ($K_D \sim 100$ nM) and, to a much lesser extent, C1q ($K_D > 0.3 \mu\text{M}$) do bind. Taken together, the data indicate that ARC186 (SEQ ID NO: 4) binds with high affinity to human C5, mainly via recognition of the C5b domain.

[0202] PEG-modified aptamers are not readily evaluated by nitrocellulose filtration assays due to the tendency of these aptamers to adhere to nitrocellulose even in the absence of target protein. However, the relative affinities of these aptamers can be assessed from their comparative ability to compete with radiolabeled, non-PEGylated aptamer (≤ 10 pM) for binding to target. As the concentration of cold (i.e., non-radiolabeled) competitor increases, the percent of

radiolabeled aptamer bound to target protein decreases. As shown in Figure 14, increasing concentrations of cold ARC186 (SEQ ID NO: 4) or PEGylated aptamer (ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62), and ARC187 (SEQ ID NO: 5)) (0.05 -1000 nM) readily compete with radiolabeled ARC186 (SEQ ID NO: 4) for binding in the presence of 2 nM C5 protein. Additionally, the titration curves for all four aptamers nearly overlap, indicating that PEG-conjugation has little or no effect on the affinity of the aptamer for C5.

Example 1B: Whole Blood Assay.

[0203] The effect of the anti-C5 aptamer on the alternative pathway of the complement system was analyzed using the following whole blood assay. In the absence of an anticoagulant, blood was drawn from normal human volunteers. Aliquots of blood (containing no anti-coagulant) were incubated with increasing concentrations of ARC186 (SEQ ID NO: 4) for 5 hours at room temperature or 37 °C. Samples were centrifuged to isolate serum and the presence of C5b in the serum was detected by sC5b-9 ELISA (C5b-9 ELISA kit, Quidel, San Diego, CA). As shown in Figure 15, the anti-complement activity, as reflected in production of C5b-9, between samples incubated at different temperatures diverged at 3 μ M. The room temperature data indicated that the concentration of aptamer required for quantitative inhibition is in the range of 3-6 μ M, whereas the reported concentration of C5 is approximately 400 nM. These results suggest that greater than 10-fold molar excess of anti-C5 aptamer (ARC186; SEQ ID NO: 4) may be required for complete inhibition of C5 activity.

Example 1C: Complement activation by zymosan.

[0204] Zymosan is a polysaccharide component of the yeast cell wall, and a potent activator of the alternative complement cascade. Addition of zymosan to ex vivo samples of blood, plasma or serum results in the accumulation of complement activation products, including C5a and the soluble version of C5b-9 (sC5b-9). Samples of undiluted human serum (Center for Blood Research, Boston, MA), citrated human whole blood (Center for Blood Research, Boston, MA) or cynomolgus serum (Charles River Labs, Wilmington, MA) were spiked with increasing concentrations of ARC658 (SEQ ID NO: 62), the 30K PEG analog of ARC186 (SEQ ID NO: 4). To activate complement, zymosan (Sigma, St. Louis, MO) in a 10X suspension was added to samples to a final concentration of 5 mg/mL. Following a 15 minute incubation at 37°C, zymosan particles were removed by centrifugation and the extent of complement activation was determined by C5a and/or sC5b-9 ELISA (C5b-9 ELISA kit, Quidel, San Diego, CA; C5a ELISA kit, BD Biosciences, San Diego, CA).

[0205] In the absence of aptamer, zymosan treatment activates ~50% of serum or whole blood C5, compared to ~1% activation in untreated sample. Addition of anti-C5 aptamer up to 50 nM (~10% of C5 concentration in blood) had little effect on sC5b-9 formation. However, further titration of C5 with increasing concentrations of ARC658 (SEQ ID NO: 62) inhibited C5 activation in a dose-dependent manner as seen in Figure 16. In human serum or whole blood, quantitative (~99%) inhibition was observed at 0.8 - 1 μ M ARC658 (SEQ ID NO: 62), corresponding to -2 molar equivalents of aptamer to C5. Higher concentrations of aptamer were required to achieve comparable inhibition in cynomolgus serum. In this case, 99% inhibition was achieved only in the presence of 10 μ M aptamer, or -20 molar equivalents of aptamer to C5.

Example 1D: Tubing loop model of complement activation

[0206] To test the ability of anti-C5 aptamer to block complement activation induced by exposure to foreign materials, as found in a cardiopulmonary bypass circuit, we used the tubing loop model described by Nilsson and colleagues (Gong et al, (1996) Journal of Clinical Immunology 16, 222-9; Nilsson et al, (1998) Blood 92, 1661-7). Tygon S-50-HL medical/surgical tubing (1/4" inner diameter) was cut into lengths of approximately 300 mm (approximately 9 mL volume) and filled with 5 mL human donor blood containing 0.4 units/mL heparin (Celsus) and varying concentrations of ARC658 (SEQ ID NO: 62) (0-5 μ M). Each length of Tygon tubing was closed into a loop with a piece of silicone tubing, as described in Gong et al. Tubing loops were rotated for 1 hour at approximately 30 rpm in a 37°C water bath. The loop contents were then poured into polypropylene conical tubes containing EDTA (10 mM final concentration) to quench complement activation. Platelet-poor plasma was isolated by centrifugation and analyzed for C5a and C3a by ELISA (C3a ELISA kit, Quidel, San Diego, CA; C5a ELISA kit, BD Biosciences, San Diego, CA).

[0207] The total complement activation in the absence of aptamer was small compared to the zymosan assay. Typically, C5a levels increased by approximately 6 ng/mL following the 1 hour incubation, corresponding to activation of <1% of the available C5. Nevertheless, this increase was reproducible and inhibited by titration with ARC658 (SEQ ID NO: 62). As shown in Figure 17, 300 - 400 nM ARC658 (SEQ ID NO: 62) was sufficient to achieve 99% inhibition of C5 activation, a level that is approximately equivalent or slightly less than the molar concentration of C5 in blood. While not wishing to be bound by any theory, although less aptamer is required to obtain 99% inhibition of C5 activation in this model than in the zymosan model, this observation could reflect the substantial differences in the complement-activating stimulus used in the two assays. C3a generation was also monitored as a control to verify that ARC658 (SEQ ID NO: 62) did not

block activation steps earlier than C5 in the complement cascade. C3a levels increased by approximately 4000 ng/mL following the 1 hour incubation, corresponding to activation of around 10% of the available C3. In contrast to C5a generation, little dose dependent inhibition of C3a generation was observed upon titration with ARC658 (SEQ ID NO: 62) demonstrating that ARC658 (SEQ ID NO: 62) specifically blocks C5 cleavage.

5

EXAMPLE 2

De Novo Selections and Sequences

10 C5 Selection with dRmY pool

[0208] Two selections were performed to identify dRmY aptamers to human full length C5 protein. The C5 protein (Quidel Corporation, San Diego, CA or Advanced Research Technologies, San Diego, CA) was used in full length ("FL") and partially trypsinized ("TP") form and both selections were direct selections against the protein targets which had been immobilized on a hydrophobic plate. Both selections yielded pools significantly enriched for full length C5 binding versus naïve, unselected pool. All sequences shown in this example are shown 5' to 3'.

[0209] Pool Preparation: A DNA template with the sequence TAATACGACTCACTATAGGGAGAGGAGAACGTTCTACN₍₃₀₎GGTCGAT CGATCGATCATCGATG (ARC520; SEQ ID NO: 70) was synthesized using an ABI EXPEDITE™ DNA synthesizer, and deprotected by standard methods. The templates were amplified with 5' primer TAATACGACT-CACTATAGGGAGAGGAGAACGTTCTAC (SEQ ID NO: 71) and 3' primer CATCGATGATCGATCGATCGAC (SEQ ID NO: 72) and then used as a template for *in vitro* transcription with Y639F single mutant T7 RNA polymerase. Transcriptions were done using 200 mM HEPES, 40 mM DTT, 2 mM spermidine, 0.01 % TritonX-100, 10% PEG-8000, 9.5 mM MgCl₂, 2.9 mM MnCl₂, 2 mM NTPs, 2 mM GMP, 2 mM spermine, 0.01 units/μL inorganic pyrophosphatase, and Y639F single mutant T7 polymerase.

[0210] Selection: In round 1, a positive selection step was conducted on nitrocellulose filter binding columns. Briefly, 1 X 10¹⁵ molecules (0.5 nmoles) of pool RNA were incubated in 100 μL binding buffer (1X DPBS) with 3 uM full length C5 or 2.6 uM partially trypsinized C5 for 1 hour at room temperature. RNA:protein complexes and free RNA molecules were separated using 0.45 um nitrocellulose spin columns from Schleicher & Schuell (Keene, NH). The columns were pre-washed with 1 mL 1X DPBS, and then the RNA:protein containing solutions were added to the columns and spun in a centrifuge at 1500 g for 2 min. Three buffer washes of 1 mL were performed to remove nonspecific binders from the filters, then the RNA:protein complexes attached to the filters were eluted twice with 200 μL washes of elution buffer (7 M urea, 100 mM sodium acetate, 3 mM EDTA, pre-heated to 95 °C). The eluted RNA was precipitated (2 μL glycogen, 1 volume isopropanol, ½ volume ethanol). The RNA was reverse transcribed with the ThermoScript RT-PCR™ system (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions, using the 3' primer described above SEQ ID NO: 72, followed by PCR amplification (20 mM Tris pH 8.4, 50 mM KCl, 2 mM MgCl₂, 0.5 uM primers SEQ ID NO: 71 and SEQ ID NO: 72, 0.5 mM each dNTP, 0.05 units/μL Taq polymerase (New England Biolabs, Beverly, MA)). The PCR templates were purified using Centriprep columns (Princeton Separations, Princeton, NJ) and used to transcribe the next round pool.

[0211] In subsequent rounds of selection, separation of bound and free RNA was done on Nunc Maxisorp hydrophobic plates (Nunc, Rochester, NY). The round was initiated by immobilizing 20 pmoles of both the full length C5 and partially trypsinized C5 to the surface of the plate for 1 hour at room temperature in 100 μL of 1X DPBS. The supernatant was then removed and the wells were washed 4 times with 120 μL wash buffer (1X DPBS). The protein wells were then blocked with a 1X DPBS buffer containing 0.1 mg/mL yeast tRNA and 0.1 mg/mL salmon sperm DNA as competitors. The pool concentration used was always at least in five fold excess of the protein concentration. The pool RNA was also incubated for 1 hour at room temperature in empty wells to remove any plastic binding sequences, and then incubated in a blocked well with no protein to remove any competitor binding sequences from the pool before the positive selection step. The pool RNA was then incubated for 1 hour at room temperature and the RNA bound to the immobilized C5 was reverse transcribed directly in the selection plate by the addition of RT mix (3' primer, SEQ ID NO:72 and Thermoscript RT, Invitrogen) followed by incubation at 65 °C for 1 hour. The resulting cDNA was used as a template for PCR (Taq polymerase, New England Biolabs). Amplified pool template DNA was desalting with a Centriprep column (Princeton Separations) according to the manufacturer's recommended conditions and used to program transcription of the pool RNA for the next round of selection. The transcribed pool was gel purified on a 10 % polyacrylamide gel every round.

[0212] The selection progress was monitored using a sandwich filter binding (dot blot) assay. The 5'-³²P-labeled pool RNA (trace concentration) was incubated with C5, 1X DPBS plus 0.1 mg/mL tRNA and 0.1 mg/mL salmon sperm DNA, for 30 minutes at room temperature, and then applied to a nitrocellulose and nylon filter sandwich in a dot blot apparatus (Schleicher and Schuell). The percentage of pool RNA bound to the nitrocellulose was calculated and monitored approximately every 3 rounds with a single point screen (+/-300 nM C5). Pool K_d measurements were measured using a titration of protein and the dot blot apparatus as described above.

[0213] Selection data: Both selections were enriched after 10 rounds over the naïve pool. See Figure 18. At round 10, the pool K_d was approximately 115 nM for the full length and 150 nM for the trypsinized selection, but the extent of binding was only about 10% at the plateau in both. The R10 pools were cloned using TOPO TA cloning kit (Invitrogen) and sequenced.

[0214] **Sequence Information:** 45 clones from each pool were sequenced. R10 full length pool was dominated by one single clone ARC913 (SEQ ID NO: 75) which made up 24% of the pool, 2 sets of duplicates and single sequences made up the remainder. The R10 trypsinized pool contained 8 copies of the same sequence ARC913 (SEQ ID NO: 75), but the pool was dominated by another sequence (AMX221.A7; 46%). The clone ARC913 (SEQ ID NO: 75) had a K_d about 140 nM and the extent of binding went to 20 %. See Figure 19.

[0215] The individual sequence listed in Table 5 is listed in the 5' to 3' direction, and represents the ribonucleotide sequence of the aptamer that was selected under the dRmY SELEX™ conditions provided. In the embodiments of the present disclosure derived from this selection (and as reflected in the sequence listing) the purines (A and G) are deoxy and the pyrimidines (U and C) are 2'-OMe. The sequence listed in Table 5 may or may not contain capping (e.g., a 3'-inverted dT). The unique sequence of the aptamer below begins at nucleotide 23, immediately following the fixed sequence GGGAGAGGAGAGAACGUUCUAC (SEQ ID NO 73), and runs until it meets the 3'-fixed nucleic acid sequence GGUCGAUCGAUCGAUCGAUG (SEQ ID NO 74)

[0216]

Table 5: Nucleotide sequence of the C5 dRmY aptamer

20 ARC913 (SEQ ID NO: 75)

GGGAGAGGAGAGAACGUUCUACCUUGGUUUGGCACAGGCAUACAUACGCAGGGGUCGAUCGAUC
GAUCAUCGAUG

[0217] **Hemolysis Assay:** The effect of ARC913 (SEQ ID NO: 75) on the classical pathway of the complement system was analyzed using a hemolysis assay previously described, compared to both ARC186 (SEQ ID NO: 4) (Anti-C5 aptamer, positive control) and unselected dRmY pool (negative control). In the assay of hemolytic inhibition, a solution of 0.2% whole human serum was mixed with antibody-coated sheep erythrocytes (Diamedix EZ Complement CH50 Test, Diamedix Corporation, Miami, FL) in the presence of titrated ARC913 (SEQ ID NO: 75). The assay was run in veronal-buffered saline containing calcium, magnesium and 1% gelatin (GVB⁺⁺ complement buffer) and incubated for 1hr at 25 °C. After incubation the samples were centrifuged. The optical density at 415 nm (OD_{415}) of the supernatant was read. The inhibition of hemolysis activity is expressed as % hemolysis activity compared to control. See Figure 20. The IC_{50} of the aptamer was calculated to be about 30 nM.

EXAMPLE 3

Composition and Sequence Optimization

40 Example 3A: Minimization of ABC913:

[0218] Six constructs based on ARC913 (SEQ ID NO: 75) were transcribed, gel purified, and tested in dot blots for binding to C5. ARC954 was similar to the parent clone with a K_d of 130 nM and extent of binding at 20%, while ARC874 (SEQ ID NO: 76) was the only other clone that bound to C5 with a K_d of 1 μ M.

[0219] The individual sequences listed in Table 6 are listed in the 5' to 3' direction and were derived from aptamers that were selected under the dRmY SELEX conditions provided. In the embodiments of the present disclosure derived from this selection (and as reflected in the sequence listing) the purines (A and G) are deoxy and the pyrimidines (U and C) are 2'-OMe. Each of the sequences listed in Table 6 may or may not contain capping (e.g., a 3'-inverted dT).

Table 6. Nucleotide sequences of ABC913 minimized clones

ARC874 (SFQ ID NO: 76)

CCUUGGCUUUGGCACAGGCAUACAUACGGCAGGG

ARC875 (SEQ ID NO: 77)

ARE56 (SEQ ID NO: 11)

ABC876 (SEQ ID NO: 78)

Argonaute (SEQ ID No. 75)

(continued)

ARC877 (SEQ ID NO: 79)

GGGUUUGGCACAGGCAUACAAACCC

ARC878 (SEQ ID NO: 80)

GGCGGCACAGGCAUACAUACGCAGGGGUCGCC

ARC954 (SEQ ID NO: 81)

CGUUCUACCUUGGUUUGGCACAGGCAUACAUACGCAGGGGUCGAUCG

10 Example 3B: Optimization of ARC913: Doped Reselection

15 [0220] In order to both optimize clone ARC913 (SEQ ID NO: 75) for C5 binding affinity and to determine the key binding elements, a doped reselection was conducted. Doped reselections are used to explore the sequence requirements within an active clone or minimer. Selections are carried out with a synthetic, degenerate pool that has been designed based on a single sequence. The level of degeneracy usually varies from 70% to 85% wild type nucleotide. In general, neutral mutations are observed but in some cases sequence changes can result in improvements in affinity. The composite sequence information can then be used to identify the minimal binding motif and aid in optimization efforts.

20 [0221] **Pool preparation:** The template sequence taatcgaactcaataGGGAGAGGAGAGAACGTTCTACN₍₃₀₎ GTTAC-GACTAGCATCGATG (SEQ ID NO: 82) was based on ARC913 (SEQ ID NO: 75) and was synthesized with each residue originating from the random region doped at a 15% level, i.e. at each random ("N") position, the residue has a 85% chance of being the nucleotide found in the wild type sequence CTTGGTTTGGCACAGGCATACATACGCAGGGGTC-GATCG (SEQ ID NO: 83) and a 15% chance of being one of the other three nucleotides.

25 [0222] The template and RNA pool for the doped reselection were prepared essentially as described above. The templates were amplified with the primers taatcgaactcaataGGGAGAGGAGAGAACGTTCTAC (SEQ ID NO: 84) and CATCGATGCTAGTCGTAAC (SEQ ID NO: 85). Two selections were done with full length C5, one selection using a higher concentration of salt in the wash step. The selection protocol was carried out as described above, with two exceptions: 1) Round 1 was done on hydrophobic plates (as well as all subsequent rounds) with only a positive step; and 2) no competitor was used at all during the selection. The C5 concentration and RNA pool concentration were kept constant at 200 nM and 1uM respectively.

30 [0223] **Doped reselection data.** Both the normal and high salt selections were enriched after 5 rounds over the naïve pool. At round 5 the pool K_d was approximately 165 nM for the high salt selection and 175 nM for the normal salt selection. The extent of binding was about 20% at the plateau in both. The R4 pools were cloned using TOPO TA cloning kit (Invitrogen, Carlsbad, CA), and 48 clones from each pool were sequenced. 12 clones from each pool were transcribed and assayed in a single point dot blot assay at 500 nM C5. Dissociation constants (K_ds) were again measured using the dot blot assay previously described. K_ds were estimated for the 11 best clones identified in the single point screen, by fitting the data to the equation: fraction RNA bound = amplitude*K_d/(K_d + [C5]). The clones with the three best K_ds were SEQ ID NO: 91 (73 nM), SEQ ID NO: 96 (84 nM) and SEQ ID NO: 95 (92 nM). The sequences for these 11 clones are listed below in Table 7.

35 [0224] The sequences listed in Table 7 are listed in the 5' to 3' direction and represent the nucleotide sequences of the aptamers that were selected under the dRmY SELEX conditions provided. In the embodiments of the present disclosure derived from this selection (and as reflected in the sequence listing), the corresponding sequences comprising the dRmY combinations of residues, as indicated in the text, wherein the purines (A and G) are deoxy and the pyrimidines (U and C) are 2'-OMe. Each of the sequences listed in Table 7 may or may not contain capping (e.g., a 3'-inverted dT). The unique sequences of each of aptamer below begins at nucleotide 23, immediately following the 5' fixed sequence GGGAGAGGAGAGAACGUUCUAC (SEQ ID NO: 86), and runs until it meets the 3'fixed nucleic acid sequence GU-UACGACUAGCAUCGAUG (SEQ ID NO: 87).

[0225]

Table 7. Nucleotide sequences of clones from doped reselection

(SEQ ID NO: 88)
 GGGAGGGAGGAAACGUUCUACCUUGGUUUGGCACAGGCAUACAUACGCAGGGGUCGAUCGGUUUACGACUAGCAUCGAUG
(SEQ ID NO: 89)
 GGGAGGGAGGAAACGUUCUACCUUGGUUUGGCACAGGCAUACAUACGCAGGGGUCGAUCGGUUUACGACUAGCAUCGAUG
(SEQ ID NO: 90)
 GGGAGGGAGGAAACGUUCUACCUUGGUUUGGCACAGGCAUAAAUAACGCAGGGCUCGAAUCGGUUUACGACUAGCAUCGAUG
(SEQ ID NO: 91)
 GGGAGGGAGGAGAACGUUCUACCUUGGUUUGGCCAGGCAUAAAUAACGCAGGGAUUUGAACUCCGUUACGACUAGCAUCGAUG
(SEQ ID NO: 92)
 GGGAGGGAGGAGAACGUUCUACCUUGGUUUGGCCAGGCAUACAUACGCAGGGGUUACGACUAGCAUCGGAUUACGACUAGCAUCGAUG
(SEQ ID NO: 93)
 GGGAGGGAGGAGAACGUUCUACCUUGGUUUGGCCACAGCCAACCCUACGCACGGCAUCGGCCGGUUUACGACUAGCAUCGAUG
(SEQ ID NO: 94)
 GGGAGGGAGGAGAACGUUCUACCUUGGUUUGGCCACAGGCAUACAUACGCAGGGGUUACGACUAGCAUCGAUA
(SEQ ID NO: 95)
 GGGAGGGAGGAGAACGUUCUACCUUGGUUUCGACUGUCAUACAUACACACGGCAAUCGGUUUACGACUAGCAUCGAUG
(SEQ ID NO: 96)
 GGGAGGGAGGAGAACGUUCUACCUUGGUUUGGCNCAGGCAUANAUACGCACGGGUUCGAUCGGUUUACGACUAGCAU
(SEQ ID NO: 97)
 GGGAGGGAGGAGAACGUUCUACCUUCUCUGCCACAAGCAUACCUUUCGCGGGGUUCUAUUGGUUACGACUAGCAUCGAUG
(SEQ ID NO: 98)
 GGGAGGGAGGAGAACGUUCUACCUUGGUUUGGCACAGGCAUAAAUCGCAGGGGUUCGAUCCGUUACGACUAGCAUCGAUG

5
10
15
20
25
30
35
40
45
50
55

Example 3C: 40 kDa Branched PEG Modification of ARC 186

[0226] The oligonucleotide 5' NH₂-fCmGfCfCGfCmGmGfUfCiUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfUfUfAfCfCfUm GfCmG-3T -3' (SEQ ID NO: 63) was synthesized on an Expedite DNA synthesizer (ABI, Foster City, CA) according to the recommended manufacturer's procedures using standard commercially available 2'-OMe RNA and 2'-F RNA and TBDMS-protected RNA phosphoramidites (Glen Research, Sterling, VA) and a inverted deoxythymidine CPG support. Terminal amine function was attached with a 5'-amino-modifier C6-TFA (Glen Research, Sterling, VA). After deprotection, the oligonucleotides were purified by ion exchange chromatography on Super Q 5PW (30) resin (Tosoh Biosciences) and ethanol precipitated.

[0227] The amine-modified aptamer was conjugated to different PEG moieties post-synthetically. The aptamer was dissolved in a water/DMSO (1:1) solution to a concentration between 1.5 and 3 mM. Sodium carbonate buffer, pH 8.5, was added to a final concentration of 100 mM, and the oligo was reacted overnight with a 1.7 molar excess of the desired PEG reagent (e.g. ARC1905 40 kDa Sunbright GL2-400NP p-nitrophenyl carbonate ester [NOF Corp, Japan], or ARC187 40 kDa mPEG2-NHS ester [Nektar, Huntsville AL]) dissolved in an equal volume of acetonitrile. The resulting products were purified by ion exchange chromatography on Super Q 5PW (30) resin (Tosoh Biosciences), and desalting using reverse phase chromatography performed on Amberchrom CG300-S resin (Rohm and Haas), and lyophilized. The structure of ARC187 (SEQ ID NO: 5) is shown in Figure 21 while the structure of ARC 1905 (SEQ ID NO: 67) is shown in Figure 22.

EXAMPLE 4**Isolated Perfused Heart Model**Example 4A: Proof of Principle with ARC 186

[0228] The average concentration of complement component C5 in human plasma is approximately 500 nM. Upon exposure of isolated mouse hearts perfused with Krebs Heinseleit buffer to 6% human plasma, the human complement cascade is activated, leading to cleavage of C5 into C5a and C5b. Component C5b subsequently forms a complex with complement components C6, C7, C8 and C9 also known as the "membrane attack complex" ("MAC" or C5b-9) which damages heart blood vessels and cardiac myocytes, thus leading to myocardial dysfunction (increased end diastolic pressure, arrhythmias) and asystole (Evans et. al., Molecular Immunology, 32, 1183-1195 (1995)). Previously, monoclonal and single chain antibodies that block human C5 cleavage (Pexelizumab or a single-chain scFv version of Pexelizumab) were tested in this model and shown to inhibit myocardial damage and dysfunction (Evans et al, 1995).

[0229] This model was used to establish that the C5-blocking aptamer ARC 186 (SEQ ID NO: 4), like Pexeluzimab, inhibited human C5-mediated complement damage to isolated perfused mouse hearts. C57 Bl/6 mice were purchased from Charles River Laboratories, (Wilmington, MA). Briefly, following induction of deep anesthesia, each mouse heart was removed and mounted on a blunt needle inserted into the aorta, through which the heart was continuously perfused with Krebs Heinseleit buffer. A pressure transducer (Mouse Specifics, Boston, MA) was inserted into the left ventricle allowing continuous measurement of the heart rate and intraventricular pressure. After a 15-minute period of equilibration during which baseline measurements were taken, hearts were subsequently perfused with buffer and 6% human plasma +/- aptamer at various concentrations (See Figure 23). During these studies and as described in Evans et al., we demonstrated that hearts which were perfused with Krebs Heinseleit buffer + 6% human plasma experienced failure within 5 minutes of adding the plasma to the perfusate, whereas hearts that were continuously perfused with buffer alone continued to beat in excess of two hours. Hence, the length of each experiment was arbitrarily defined as 15 minutes.

The outline of this study with ARC 186 is presented in Figure 23.

[0230] Intraventricular pressure was monitored and recorded continuously resulting in a pressure wave tracing (Figures 24 and 25). The lowest deflection point represents the end diastolic pressure ("EDP") and the highest deflection point represents the systolic pressure ("SP"). Baseline pressure waves appear to the left of the vertical black line marked "0" shown on each tracing. As previously published (Evans et al, 1995), hearts perfused with 6% human plasma experienced a rapid increase in left ventricular end diastolic pressure, ultimately culminating in asystole (the heart stops) within 5 minutes (Figure 24). When irrelevant aptamer was added to the human plasma at 50-fold molar excess, increased EDP and asystole were also observed (Figure 24).

[0231] When ARC 186 was added to the system at molar equivalence, there was also a precipitous increase in EDP, culminating in asystole (Figure 25). In all three groups of hearts that experienced complement-mediated damage, increased EDP and asystole, the heart was visibly edematous and turgid by the end of the experiment. When ARC186 was added to plasma in 10-fold or 50-fold (Figure 25) molar excess, ventricular pressure waves remained normal and asystole was not observed. In addition, the previously described edema and turgidity were not apparent in these groups.

[0232] During each experiment, the heart rate was recorded at 5-minute intervals, and the average heart rate for the

group during each interval was graphed. As shown in Figure 26 hearts perfused without aptamer or with irrelevant aptamer developed asystole quickly, usually within 5 minutes. ARC186 added to the system at molar equivalence slightly delayed the onset of asystole. Hearts in this group ultimately failed, however. ARC186 added to the plasma at 10-fold or 50-fold molar excess preserved the heart rate for the duration of each experiment.

5 [0233] The percent increase in heart weight over baseline was calculated for a representative sample of failed hearts (no aptamer or 50-fold molar excess of irrelevant aptamer) and compared to ARC186-protected hearts (10-fold and 50-fold molar excess of ARC186). As shown in Figure 27, ARC186 protected hearts gained significantly less weight than the failed hearts in the control groups.

10 [0234] Because ARC186 inhibits C5 but not C3 cleavage, C3 cleavage products (C3a) but not C5 cleavage products (C5a or C5b) should be found in the effluent flowing from the isolated hearts protected by ARC186. To directly show that ARC186 inhibited cleavage of human plasma C5, the relative levels of human complement proteins C5a and C5b (C5 cleavage products) and C3a (a C3 cleavage product) were measured in the buffer effluent from the various groups by commercially available ELISA kits (C5b-9 ELISA kit, Quidel, San Diego, CA; C5a and C3a ELISA kit, BD Biosciences, San Diego, CA). ARC186 inhibited human plasma C5 cleavage and the production of C5a (Figure 28) and C5b-9 (Figure 15 29) in a dose-dependent manner. In contrast, ARC186 had no effect on cleavage of human C3 into C3a and C3b (Figure 30) further demonstrating the C5 specificity of the molecule.

20 [0235] Once generated, complement C3b and C5b fragments are deposited locally on tissues in the vicinity of the site of complement activation. Following completion of the experiments, mouse hearts were frozen in OCT media (Sakura Finetek, Torrance, CA), sectioned and then stained using standard immunohistochemistry for the presence of human C3b (clone H11, Chemicon, Temecula, CA), human C5b-9 (clone aE11, DAKO, Carpinteria, CA) or control mouse IgG (Vector Laboratories, Burlingame, CA). Results of the study are presented in Figure 31.

25 [0236] As described in this study, the C5-blocking aptamer ARC186 was tested in an ex vivo model of complement component C5-mediated tissue damage which uses isolated mouse hearts perfused with Krebs Heinseleit buffer and 6% heparinized human plasma, based on a model described in a previously published study that tested the effects of the anti-C5 antibody, Pexeluzimab on the complement system (Evans, Molecular Immunol 32:1183, (1995). Using this model, it was demonstrated that the C5 - blocking aptamer (a) inhibited cleavage of human plasma C5 (but not C3), (b) inhibited deposition of human C5b (but not C3b) on mouse heart tissue and (c) inhibited human C5b-9 mediated myocardial dysfunction at clinically relevant concentrations (5 μ M, a 10-fold molar excess of aptamer vs. C5). These data show that when the human complement cascade is activated in a physiologically relevant manner, C5-blocking aptamers 30 are able to inhibit cleavage of plasma C5 and prevent myocardial damage and dysfunction.

Example 4B: Efficacy of PEGylated Aptamer

35 [0237] The material and methods used in this study were exactly the same as described in Example 4A above. The experimental design and results are presented in Figure 32. The first half of the experiment used human heparinized plasma (Center for Blood Research, Harvard Medical School, Boston, MA) as a source of complement and the second half used heparinized cynomolgus macaque plasma (Charles River Laboratories, Wilmington, MA) as a source of complement. A PEGylated aptamer (ARC658; SEQ ID NO:62) was added to the system at increasing molar ratios. Although 40 all of the relevant ventricular pressure tracings were collected, the table lists the presence or absence of an increase in end diastolic pressure (EDP), whether or not asystole occurred and the time until heart failure (defined as the presence of an elevated EDP and asystole).

45 [0238] During experiments with human plasma, the optimal dose of AR658 (SEQ ID NO: 62) was determined to be molar equivalence (500 nM) whereas during experiments with non-human primate plasma, a 50-fold molar excess (25 μ M) was necessary to protect the heart from C5b-mediated damage (see Figure 32).

50 [0239] These data are consistent with the difference in the affinity of the anti-C5 aptamer for human v. non-human primate C5 indicated by the *in vitro* data. While not wishing to be bound by any theory, during our subsequent cynomolgus macaque PK/PD studies described in Example 5, we additionally demonstrated that a 30-fold molar excess of aptamer was necessary to inhibit zymosan-mediated plasma C5 cleavage, further supporting the notion that the aptamer binds primate C5 with lower affinity than human C5.

55 [0240] Collectively, these studies indicate that both C5-blocking aptamers ARC186 (SEQ ID NO: 4) and to a greater extent ARC658 (SEQ ID NO: 62) are efficacious in the mouse isolated, perfused heart model. This model also demonstrated that significantly more ARC658 (SEQ ID NO: 62) had to be used to inhibit cynomolgus macaque plasma C5-mediated heart damage (30+ molar excess), compared with human C5-mediated heart damage (molar equivalence), further supporting *in vitro* data which indicated that the aptamer had lower affinity for primate C5. Finally, these data indicated that cynomolgus macaques would need to be dosed beyond a 30-fold molar excess in order to demonstrate *in vivo* C5 blockade during PK/PD studies.

EXAMPLE 5**Drug Metabolism & Pharmacokinetics of Anti-C5 Aptamers in Animals**

5 [0241] In Examples 5A-5G, all mass based concentration data refers only to the molecular weight of the oligonucleotide portion of the aptamer, irrespective of the mass conferred by PEG conjugation.

Example 5A: Metabolic stability of the C5 inhibitor ARC186 in primate and rat plasma

10 [0242] The non-PEGylated oligonucleotide precursor of the aptamers (*i.e.*, ARC 186; SEQ ID NO: 4) was tested in rat and cynomolgus macaque plasma (Charles River Labs, Wilmington, MA) in order to assess its stability, rate kinetics, and pathways of degradation. Testing was performed using 5' end-radiolabeled (^{32}P) aptamer incubated at 37°C in 95% pooled plasma (citrated) over the course of 50 hrs. At selected time points, aliquots of aptamer-containing plasma were withdrawn, immediately flash frozen in liquid nitrogen, and stored at -80° C. Detection and analysis of the aptamer and 15 its metabolites in plasma was accomplished using liquid-liquid (phenol-chloroform) extraction followed by gel electrophoresis (on a 10% denaturing polyacrylamide sequencing gel) and high-resolution autoradiography.

15 [0243] Figure 33 shows a log-linear plot of remaining percent of full-length aptamer as a function of incubation time in both rat and cynomolgus macaque plasma. The degradation profile in both species appears to be essentially monophasic, with a rate constant of approximately $k = 0.002 \text{ hr}^{-1}$.

Example sub: Pharmacokinetics of ARC657, ARC658 and ARC187 in the rat following intravenous administration

20 [0244] To assess the pharmacokinetic profile of ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5), and to estimate the required dosing level and frequency in primates and humans, a pharmacokinetic study was conducted in catheterized Sprague-Dawley rats (Charles River Labs, Wilmington, MA). Aptamers were 25 formulated for injection at 10 mg/mL (oligo weight) in standard saline and sterile-filtered (0.2 μm) into a pre-sterilized dosing vial under aseptic conditions. The route of administration used for the rat study was an intravenous bolus via the tail vein at a dose of 10 mg/kg. Study arms consisted of 3 animals per group, from which serial bleeds were taken pre-dose and at specified time points over the course of 48 hours. The study design is outlined in Figure 34. Blood samples were 30 obtained from the surgically implanted jugular vein catheters, transferred directly to EDTA-coated tubes, mixed by inversion, and placed on ice until processing for plasma.

35 [0245] Plasma was harvested by centrifugation of blood-EDTA tubes at 5000 rpm for 5 minutes and supernatant (plasma) was transferred to a fresh pre-labeled tube. Plasma samples were stored at -80° C until the time of analysis. Analysis of plasma samples for ARC 187 was accomplished using a homogeneous assay format utilizing the direct addition of plasma aliquots to assay wells containing the commercially available fluorescent nucleic acid detection reagent Oligreen™ (Molecular Probes, Eugene, OR). After a brief incubation period (5 min) at room temperature, protected from light, the assay plates were read by a fluorescence plate reader (SpectraMax Gemini XS, Molecular Devices, Sunnyvale, CA). The fluorescence signal from each well was proportional to the concentration of aptamer in the well, and sample concentrations were calculated by interpolation of fluorescence values from a fluorescence-concentration standard curve (mean values from duplicate or triplicate curves). Mean plasma concentrations were obtained at each time point from the three animals in each group. Plasma concentration versus time data was subjected to noncompartmental analysis (NCA) using the industry standard pharmacokinetic modeling software WinNonLin™ v.4.0 (Pharsight Corp., Mountain View, CA). Estimates were obtained for the following primary pharmacokinetic parameters: maximum plasma concentration, C_{\max} ; area under the concentration-time curve, AUC; terminal half-life, $t_{1/2}$; terminal clearance, Cl; and volume of distribution at steady state, V_{ss} .

40 [0246] Mean plasma concentration versus time data are shown in Figure 35 and are plotted in Figure 36. The concentration versus time data was subjected to noncompartmental analysis (NCA) using WinNonLin™ v.4.0. This analysis yielded the values presented in Figure 37.

45 [0247] As anticipated, the 40 kDa aptamer ARC 187 (SEQ ID NO: 5) had the longest half-life and the 20 kDa aptamer, ARC657 (SEQ ID NO: 61), the shortest. The observed V_{ss} relative to the known plasma volume (~40 mL/kg) suggested a moderate degree of binding/sequestration of ARC 187 (SEQ ID NO: 5) to proteins and/or tissue matrix in the extravascular space. Assuming a need to maintain a 5-fold molar excess of aptamer, the results of this study suggested that ARC187 (SEQ ID NO: 5) provides a significant advantage in terms of the dosing frequency and total amount of aptamer needed to maintain the desired plasma levels.

55 [0248] Previous studies (data not shown) in rodents and primates with aptamers of similar composition have shown dose proportionality/linearity at doses up to 30 mg/kg, so it is not anticipated that this dosing level will result in nonlinear pharmacokinetic behavior.

Example 5C: Pharmacokinetics of ARC187 and ARC1905 in the mouse following intravenous administration

[0249] To assess the pharmacokinetic profile of the ARC 186 (SEQ ID NO: 4) oligonucleotide backbone conjugated to a different 40 kDa branched PEG than that of ARC187 (SEQ ID NO: 5), a pharmacokinetic study was conducted in 5 female CD-1 mice (obtained from Charles River Labs, Wilmington, MA). Aptamers were formulated for injection at 2.5 mg/mL (oligo weight) in standard saline and sterile-filtered (0.2 μ m) into a pre-sterilized dosing vial under aseptic conditions. The route of administration used for the mouse study was an intravenous bolus via the tail vein at a dose of 10 mg/kg. Study arms consisted of 3 animals per group, from which terminal bleeds were taken pre-dose (i.e., the non-dosed control group) and at specified time points over the course of 72 hours. The study design is outlined in Figure 38A.

[0250] Blood samples were obtained by terminal cardiac puncture, transferred directly to EDTA-coated tubes, mixed by inversion, and placed on ice until processing for plasma. Plasma was harvested by centrifugation of blood-EDTA tubes at 5000 rpm for 5 minutes and supernatant (plasma) was transferred to a fresh pre-labeled tube. Plasma samples were stored at -80° C until the time of analysis. Analysis of plasma samples for ARC 187 and 1905 was accomplished using a homogeneous assay format utilizing the direct addition of plasma aliquots to assay wells containing the 10 commercially available fluorescent nucleic acid detection reagent Oligreen™ (Molecular Probes, Eugene, OR). After a brief 15 incubation period (5 min) at room temperature, protected from light, the assay plates were read by a fluorescence plate reader (SpectraMax Gemini XS, Molecular Devices, Sunnyvale, CA). The fluorescence signal from each well was proportional to the concentration of aptamer in the well, and sample concentrations were calculated by interpolation of fluorescence values from a fluorescence-concentration standard curve (mean values from duplicate or triplicate curves).

20 Mean plasma concentrations were obtained at each time point from the three animals in each group. Plasma concentration versus time data was subjected to noncompartmental analysis (NCA) using the industry standard pharmacokinetic 25 modeling software WinNonLin™ v.4.0 (Pharsight Corp., Mountain View, CA). Estimates were obtained for the following primary pharmacokinetic parameters: maximum plasma concentration, C_{max} ; area under the concentration-time curve, AUC; terminal half-life, $t_{1/2}$; terminal clearance, Cl; and volume of distribution at steady state, V_{ss} . Mean plasma concentration versus time data are plotted in Figure 38B.

[0251] The concentration versus time data was subjected to noncompartmental analysis (NCA) using WinNonLin™ 30 v.4.0. This analysis yielded the values presented in Figure 38C. As anticipated, the 40 kDa PEGs from both vendors showed pharmacokinetic equivalence in mice.

Example 5D: Tissue uptake study of the C5 inhibitors ARC657, ARC658 and ARC187 in the mouse following intravenous bolus administration

[0252] Female CD-1 mice were obtained from Charles River Labs (Wilmington, MA). Formulation of ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5) for injection was in saline at 5 mg/ml. Dosing 35 formulations were sterile-filtered (0.2 μ m) into pre-sterilized dosing vials under aseptic conditions and animals were given an intravenous bolus via the tail vein at a dose of 25 mg/kg. The study consisted of groups of 3 animals for each of four time-points, t=pre-dose, 3, 6, 12 hrs. Following exsanguination, the vasculature of each animal was flushed extensively (V-30 mL) with saline to remove any blood left in the vasculature. Tissues (heart, liver, kidney) were harvested, weighed, then homogenized at 50% w/v in standard saline, and stored at -80° C until the time of analysis.

[0253] Analysis of tissue for ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62), and ARC187 (SEQ ID NO: 5) was 40 accomplished using a hybridization-based ELISA-type assay. In this assay, a biotinylated capture probe was pre-immobilized in the wells of a 96-well microplate at a binding solution concentration of 125 nM for 3 hrs. The plate wells were washed 5 times with 1X PBS. The plates were then blocked with 150 μ l/well of a 1X SuperBlock in TBS (Pierce Chemical, Rockford, IL). Plates were washed again, covered, and stored at 4° C until use. In separate tubes, the samples(s) were 45 annealed in a buffer containing a FAM-labeled (5'-Fluorescein) sample-detection probe at 200 nM at 90°C for 10 min, then quenched on ice. Concentration standards and control samples of plasma/tissue were also pre-annealed with sample-detection probe solutions and then pipetted into assay plate wells containing immobilized biotin capture probe, and annealed at 45° C for 2.5 hrs. Plates were then washed again, and filled with 100 μ l/well of a solution containing 50 1X PBS containing 1 μ g/mL of anti-fluorescein monoclonal antibody conjugated to horse radish peroxidase (anti-FITC MAb-HRP, Molecular Probes, Eugene, OR) in 1X PBS, and incubated for approximately 1 hr. Plates were washed again as above. Assay plate wells were then filled with 100 μ l of a solution containing a fluorogenic HRP substrate (QuantaBlu, Pierce Chemical, Rockford, IL), and incubated for 20-30 min protected from light. After 45 minute incubation, 100 μ l/well of a stop solution was added to quench the fluorescent precipitate-producing reaction. Plates were read immediately on a fluorescence microplate reader (SpectraMax Gemini XS, Molecular Devices, Sunnyvale, CA) with fluorescence excitation at 325 nm and emission detected at 420 nm. Each well was read 10 times. All three aptamers 55 were detectable in the heart tissue at the three timepoints (Figure 39).

Example 5E: Pharmacokinetics and pharmacodynamics of the C5 inhibitors ARC657, ARC658 and ARC187 in the cynomolgus macaque following intravenous administration study 1

[0254] Formulation of ARC657 (SEQ ID NO: 61), ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5) for injection was in standard saline at 10 mg/mL and dosing formulations were sterile-filtered (0.2 μ m) into pre-sterilized dosing vials under aseptic conditions. The route of administration used for the macaque study was an intravenous bolus via a surgically implanted femoral vein catheter at a dose of 30 mg/kg (approximately 50-fold molar excess). The study design is outlined in Figure 40. Blood samples were obtained from the femoral vein catheters, transferred directly to sodium citrate-coated tubes, mixed by inversion, and placed on ice until they were centrifuged to separate plasma (3000 rpm for 5 minutes). Plasma was then divided into 250 μ L aliquots which were stored at -80° C and one aliquot of each sample was evaluated for aptamer concentration using the fluorescence-based Oligreen™ assay previously described in the rat PK section above.

[0255] The primary plasma concentration versus time data is presented in tabular form in Figure 41. As anticipated, the 40 kDa PEG aptamer ARC187 (SEQ ID NO: 5) persisted in plasma for the longest period of time whereas the 20 kDa PEG aptamer ARC657 (SEQ ID NO: 61) persisted for the shortest amount of time. Inspection of the data shown in Figure 41 suggested that the data would best be fit by a two-compartment model. Thus, the pharmacokinetic parameter estimates reported in Figure 42 were derived from the two-compartment model using the industry standard pharmacokinetic modeling software WinNonLin™ v.4.0 (Pharsight Corp., Mountain View, CA).

[0256] As shown in Figure 42, all of the aptamers had a similar Cmax value, between 23 and 30 μ M, indicating that the aptamer dose (30 mg/kg) was sufficient to achieve a 50-fold molar excess of plasma aptamer vs C5 concentration (50 fold molar excess, about 25 μ M). Although they differ by 10,000 molecular weight, ARC657 (20 kDa PEG) (SEQ ID NO: 61) and ARC658 (30 kDa PEG) (SEQ ID NO: 62) had similar exposure (AUC), $t_{1/2}(\alpha)$ and $t_{1/2}(\beta)$ values. In contrast, ARC 187 (SEQ ID NO: 5) had significantly higher exposure (AUC) values, a prolonged $t_{1/2}(\alpha)$ and a slightly longer $t_{1/2}(\beta)$ than the other molecules.

[0257] Additional aliquots of the plasma samples collected during the pharmacokinetics study were subsequently analyzed *in vitro* to determine the efficacy of primate C5 blockade. The zymosan activation assay was run as described above to determine the amount of primate C5b-9 and C5a, generated, respectively. The data were plotted in several different formats including C5b-9 concentration versus sample time (Figure 43a), C5b-9 concentration versus aptamer concentration (Figure 43b), C5a concentration versus sample time (Figure 43c), and C5a concentration versus aptamer concentration (Figure 43d).

[0258] The 40 kDa PEG aptamer ARC187 (SEQ ID NO: 5) inhibited primate C5 cleavage (C5b-9 and C5a concentration) for the longest period of time (Figures 35a and 35c). When the C5b-9 and C5a data were plotted versus aptamer concentration, it indicated that the concentration of C5 blocking aptamer had to exceed 30-fold molar excess, regardless of the size of the PEG molecules, in order for C5 cleavage to be completely inhibited (Figures 35b and 35d).

[0259] In summary, the data from the cynomolgus macaque PK/PD study demonstrate that (a) as anticipated, at least a 30-fold molar excess of aptamer (about 15 μ M plasma aptamer concentration) was necessary to inhibit C5 cleavage *in vivo* in the cynomolgus macaque, regardless of the size of the PEG group, (b) C5-blocking aptamers did not cause overt toxicity in this species, and (c) when animals were dosed at a relatively high levels (50-fold molar excess), plasma aptamer levels were well within the appropriate assay range during the period of sampling to allow calculation of pharmacokinetic parameters

Example 5F: Pharmacokinetics and pharmacodynamics of the C5 inhibitors ARC658 and ARC187 in the cynomolgus macaque following intravenous administration- study 2

[0260] Study 2 was similar in design to study 1 described above, with the following exceptions a) only two compounds were evaluated (ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5); b) the number of animals was increased to four per group; and c) the 1-minute plasma sample was deleted and replaced with a 144 hour sample to ensure that the terminal half-life calculation was based upon more data points. The formulation and dosing of these two aptamers, blood sampling and plasma isolation techniques was identical to the methods described above in study 1. The design for study 2 is summarized in Figure 44.

[0261] Following completion of study 2, plasma aliquots were analyzed as described in study 1 to determine the a) the concentration of aptamer in plasma at various timepoints following intravenous administration, and b) the efficacy of C5 blockade.

[0262] Plasma aptamer concentration was plotted as a function of time (Figure 45) and the primary data for ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5) are presented in tabular form in Figures 39 and 40, respectively. The 40 kDa PEG aptamer ARC187 (SEQ ID NO: 5) persisted in plasma for the longest period of time. Inspection of Figure 45 indicated that the data would be best fit by a two-compartment model. Thus, the pharmacokinetic parameter estimates reported in Figure 46 were derived from the two-compartment model using WinNonLin™ v.4.0 (Pharsight Corp., Mountain

View, CA).

[0263] Comparing the pharmacokinetic parameters generated during the PK/PD study 1 and study 2 above, the data for ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5) were similar with the exception of the $t_{1/2}(\alpha)$ measurement for ARC187. While not wishing to be bound by any theory, the discrepancy in the $t_{1/2}(\alpha)$ measurements for ARC187 between the two studies is likely due to the small sample size in the pilot study.

[0264] As demonstrated in Figure 46, the C_{max} values were similar for ARC658 (SEQ ID NO: 62) and ARC187 (SEQ ID NO: 5). In contrast, drug exposure (AUC) was significantly greater in animals treated with ARC187 (SEQ ID NO: 5). Also, ARC187 had prolonged $t_{1/2}(\alpha)$ and $t_{1/2}(\beta)$ values as compared to ARC658 (SEQ ID NO: 62). These data, along with the data generated during the PK/PD study 1 indicate that of the C5-blocking aptamers ARC187 may provide the most effective *in vivo* C5 blockade for a given dose.

[0265] Additional aliquots of the plasma samples collected during the pharmacokinetics study were subsequently analyzed *in vitro* to determine the efficacy of primate C5 blockade. As before, the zymosan activation assay was run to determine the amount of primate C5b-9 and C5a, respectively, generated. The data were plotted as C5b-9 concentration versus aptamer concentration (Figure 47) and C5a concentration versus aptamer concentration (Figure 48). As previously demonstrated during PK/PD study 1, the concentration of C5 blocking aptamer must exceed a 30-fold molar excess (aptamer to plasma C5 concentration), or approximately 15 μ M, regardless of the size of the PEG molecule, in order for primate C5 cleavage to be completely inhibited (Figures 41 and 42).

[0266] By inspecting the data in the tables of Figures 39 and 40, it is apparent that after a 30-mg/kg I.V. bolus, ARC658 (SEQ ID NO: 62) remains above 15 μ M for approximately 4 hours whereas ARC187 remains above 15 μ M for approximately 8 hours. Thus, given a similar dose of drug, the 40 K aptamer ARC 187 provides clinical efficacy for approximately twice as long as the 30K aptamer ARC658 (SEQ ID NO: 62).

[0267] In summary, cynomolgus macaques must be treated with at least a 30-fold molar excess of aptamer vs plasma C5 in order to block C5 conversion *in vivo*. These data are consistent with previous *in vitro* (hemolysis) and ex-vivo (isolated perfused mouse heart) studies which suggested that the C5-binding aptamers had a lower affinity for primate C5 versus human C5. It has been shown that C5-blocking aptamers can safely be delivered as an intravenous bolus at a dose of up to 30 mg/kg, which equates to approximately, a 50-fold molar excess of aptamer vs C5 concentration.

Example 5G: Pharmacokinetics and pharmacodynamics of the C5 inhibitor ARC187 in the cynomolgus macaque following bolus IV administration and infusion

[0268] The pharmacokinetic (PK) and pharmacodynamic (PD) profiles of ARC187 (SEQ ID NO: 5) were also evaluated in cynomolgus macaques after an intravenous loading bolus followed immediately by the initiation of an intravenous infusion. This study design is shown in Figure 49.

[0269] The loading bolus dose and infusion rate necessary to achieve the target steady state plasma concentration of 1 μ M were calculated using the pharmacokinetic parameters derived from the IV bolus-only study listed in Figure 50.

[0270] A total of three cynomolgus macaques were administered an IV bolus of ARC187 at 1 mg/kg, followed immediately by the initiation of an IV infusion at a rate of 0.0013 mg/kg/min for a period of 48 hrs. Samples of whole blood were collected from 0 to 192 hours post-treatment, stored on wet ice, processed for plasma, and then stored frozen at -80 C. The concentration of ARC 187 in plasma samples was determined using both a fluorescent nucleic acid stain assay (described in Example 5B) and a GLP-validated performance liquid chromatography (HPLC) assay. The HPLC assay method for the determination of ARC187 in monkey plasma was validated by ClinTrials Bio-Research (Montreal, Canada). The validation study complied with the United States Food and Drug Administration (FDA) Good Laboratory Practice (GLP) regulations (21 CFR §58). The HPLC assay method was validated with respect to: selectivity, linearity, lower limit of quantitation (LLOQ), carry-over, intra-assay precision and accuracy, inter-assay precision and accuracy, stock solution stability, injection medium stability, short-term matrix stability, freeze-thaw stability, long-term matrix stability and dilution integrity. The usable linear dynamic concentration range of assay was determined to be 0.080 to 50.0 μ M.

[0271] The measured PK profile of ARC 187 under these conditions conformed well to the calculated profile generated using only the IV bolus PK parameters (see Figure 51). The target plasma concentration of 1 μ M was established in < 5 min post-dose and maintained for the entire duration of infusion. After cessation of the infusion, the aptamer showed a terminal clearance half-life, $t_{1/2}(\beta) \sim 40$ -60 hr.

[0272] The pharmacodynamic activity of ARC187 (SEQ ID NO: 5) in the cynomolgus macaque was evaluated ex-vivo by using plasma samples collected during PK study in the zymosan activation assay previously described with the modification that cynomolgous sample plasma was diluted 10-fold into 10% human plasma and then treated with 5 mg/mL zymosan. C5 activation, as reflected by the appearance of the C5a cleavage product, was measured by ELISA specific to human C5a (C5a ELISA kit, BD Biosciences, San Diego, CA). The concentration of active ARC187 in each sample was then quantified by comparison with a standard curve derived from zymosan assays using samples prepared with known ARC 187 levels (see Figure 52). This study indicates that ARC187 maintains its anti-complement activity throughout the duration of and following infusion, at levels substantially consistent with the pharmacokinetic profile

described above.

Example 5H: Prediction of Human Dosing Requirement

5 [0273] Human dosing requirements for prevention, amelioration, or treatment of complications related to CABG surgery are based on the following assumptions: first, CABG patients will be administered a single intravenous bolus dose of the anti-C5 aptamer prior to initiating surgery, followed by continuous infusion to establish and maintain a steady-state plasma concentration of 1.5 μ M for 24-48 hours post CABG surgery. The bolus dose and infusion rate estimates are based upon calculations using the pharmacokinetic parameters derived from the previously described IV bolus-only and
10 bolus plus infusion studies in cynomolgus macaques. The estimated bolus dose of ARC 187 is 1 mg/kg, and the associated infusion rate is 0.0013 mg/kg/min. For this bolus plus 48 hr infusion regimen, the anticipated total drug requirement is 0.4 g for ARC187, where mass refers to oligonucleotide weight only (see column 7 in the table of Figure 53). Column 2 of the table shown in Figure 53 refers to the weight of the PEG group conjugated to oligonucleotide portion of ARC 187, column three refers to the molecular weight of the oligonucleotide portion of ARC 187 (and will be the same for all
15 aptamers herein that comprise ARC 186 (SEQ ID NO: 4) as its oligonucleotide sequence), column 4 refers to the molecular weight of 40 kDa PEG conjugated to ARC186 (SEQ ID NO: 4) via amine reactive chemistry as described in Example 3C above, column 5 refers to ARC187's α phase half life in a two compartment model, and column six refers to ARC187's β phase half life in a two compartment model.

20 **EXAMPLE 6**

Anti-C5 Aptamers and Heparin/Protamine Interaction

25 [0274] One anticipated application of the anti-C5 aptamer is as a prophylactic for the prevention or mitigation of inflammatory side effects associated with coronary artery bypass graft (CABG) surgery. High concentrations of the anticoagulant heparin (3 - 5 units/mL or 1 - 2 μ M) are typically administered during CABG to prevent thrombosis and maintain patency within components of the bypass pump; reversal of heparin's effect after the procedure, and restoration of normal hemostasis, is achieved by the administration of similarly high concentrations of protamine (- 5 μ M). Given the potential dangers to patients of any interference in the effectiveness of either of these drugs, it was necessary to demonstrate that anti-C5 aptamers (1) do not alter the activities of either drug and (2) do not display inherent effects on hemostasis that could complicate patient anticoagulation treatment.

30 [0275] Heparin is a sulfated polysaccharide with a net negative charge and a mean molecular mass of approximately 15 kDa that exerts an inhibitory effect on a number of proteases in the coagulation cascade by promoting interactions with antithrombin. Protamine, a highly positively charged polypeptide, is able to block heparin activity via a poorly characterized interaction that is at least partially electrostatic in nature. The functional core of ARC187 (SEQ ID NO: 5), like heparin, is highly anionic. Thus, it is conceivable that ARC187 could nonspecifically bind to heparin-binding sites or protamine and interfere with the activities of these molecules. The following studies investigated the inherent (i.e., heparin-like) anticoagulant properties of ARC 187, the effects of ARC 187 on heparin function, the effects of ARC 187 on heparin-neutralization by protamine, and the effects of protamine on the complement inhibiting properties of ARC 187.

40 Example 6A: In vitro effects of ARC187 on coagulation

45 [0276] The inherent effects of ARC187 (SEQ ID NO: 5) on plasma coagulability were investigated using standard clinical tests of the extrinsic and intrinsic arms of the coagulation cascade, the prothrombin time (PT) and activated partial thromboplastin time (aPTT), respectively. As shown in Figure 54, titration of citrated human plasma with concentrations well in excess of projected doses (up to 20 μ M) resulted in no change in the PT, and only a slight elevation in the aPTT.

50 [0277] To assess the *in vitro* effects of ARC 187 on heparin and protamine functions, blood from 3 individuals was drawn into 4-5 units/mL heparin, doses associated with heparin levels used in CABG surgery. The coagulability of these samples was assessed using the activated clot time (ACT), a whole blood coagulation test routinely used to monitor heparin activity during surgery. At these concentrations of heparin, in the absence of other additives, the ACT was significantly prolonged from a baseline value of ~150 seconds to ~500 seconds in the presence of 4 U/mL heparin or ~800 seconds in the presence of 5 U/mL heparin. Addition of 10 μ M ARC 187 to these samples had little effect on clot time, demonstrating that ARC 187 does not interfere with the anticoagulant activity of heparin.

55 [0278] The heparin anticoagulant effect was readily neutralized by titration with protamine up to 6-8 μ M (4 U/mL heparin) or 12 μ M (5 U/mL heparin). ACT values in the presence of heparin and neutralizing concentrations of protamine were essentially indistinguishable from baseline. Since the nucleic acid core of ARC187 (12 kDa) is of larger molecular weight than protamine (5 kDa), one might expect that equimolar concentrations of ARC 187 added to protamine would

be sufficient to *completely* reverse the neutralizing activity of protamine. However, preincubation of protamine with approximately equivalent concentrations of ARC 187 had little effect on the ACT. Blood samples containing neutralizing concentrations of protamine displayed similar ACT values in the presence or absence of 10 μ M ARC 187, indicating that ARC 187 has only a slight if any effect on the procoagulant activity of protamine. These results are summarized in Figure 55.

Example 6B: *In vivo* effects of ARC187 on coagulation

[0279] The interactions between the function of heparin and protamine during concurrent administration of anti-C5 aptamer ARC 187 (SEQ ID NO: 5), at clinical doses of heparin and clinical/subclinical/superclinical doses of protamine were investigated to determine whether the presence of subclinical/superclinical plasma concentrations of ARC 187 would interfere with the reversal of heparin anticoagulation by protamine. The results of the study are summarized in Figure 56. Briefly, the baseline ACT values were unaffected by 10 μ M (i.e., 10-fold molar excess of the clinical dose) of ARC187 at all heparin doses tested. Similarly, the extent of anticoagulation by heparin was unaffected by 10 μ M ARC 187. In the absence of ARC 187, the minimum efficacious dose of protamine was - 30% (clinical dose=100%). Furthermore, the reversal of heparin anticoagulation by 30% protamine was unaffected by 10-fold molar excess of the clinical dose (i.e., 10 μ M) of ARC187. Thus, the use of ARC187 for complement inhibition in a clinical setting (e.g., CABG) should be unaffected by concurrent use of heparin and protamine at typical doses.

Example 6C: Effect of heparin and protamine on ARC187 anti-complement function

[0280] The effects of heparin and protamine on the anti-complement activity of ARC187 (SEQ ID NO: 5) were examined in citrated whole blood samples activated with zymosan, as described in Example 1. Just prior to zymosan activation, ARC 187 was titrated into samples of citrated blood treated under four conditions: 1) no treatment (no heparin or protamine); 2) 4 U/mL heparin; 3) 6 μ M protamine; 4) 4 U/mL heparin + 6 μ M protamine. Following activation with zymosan, C5 activation was quantified by ELISA measurement of sC5b-9 in plasma (C5b-9 ELISA kit, Quidel, San Diego, CA). For each condition, the results, expressed as percent inhibition of C5 activation versus ARC 187 concentration, were indistinguishable within error (see Figure 57). In all cases complete inhibition was achieved with 1-2 μ M ARC187. Thus, heparin and protamine, separately or combined at concentrations relevant to their use in CABG surgery, do not appear to affect the anti-complement activity of ARC187.

[0281] The invention having now been described by way of written description and example, those of skill in the art will recognize that the invention can be practiced in a variety of embodiments and that the description and examples above are for purposes of illustration and not limitation of the following claims.

SEQUENCE LISTING

[0282]

<110> Archemix Corp., et al.
 <120> Aptamer Therapeutics Useful in the Treatment of Complement Related Disorders
 <130> 23239-576-061
 <140> US2005/004606
 <141> 2005-02-14
 <150> 60/608,048
 <151> 2004-09-07
 <150> 60/581,685
 <151> 2004-06-21
 <150> 60/547,747
 <151> 2004-02-25
 <150> 60/544,542
 <151> 2004-02-12

<160> 102

<170> PatentIn version 3.3

5 <210> 1

 <211> 38

 <212> DNA

 <213> artificial sequence

10 <220>

 <223> synthetic aptamer

 <220>

 <221> misc_feature

15 <223> cytosine at positions 3, 4, 6 and 37 are 2'-fluoro

 <220>

 <221> misc_feature

 <223> uridine at positions 9, 30 and 31 are 2'-fluoro

20 <220>

 <221> misc_feature

 <222> (1)..(1)

 <223> n at position 1 is 2'-fluoro cytidine or 2'-O-methyl cytidine

25 <220>

 <221> misc_feature

 <222> (2)..(2)

 <223> n at position 2 is 2'-OH guanosine or 2'-O-methyl guanosine

30 <220>

 <221> misc_feature

 <222> (7)..(7)

 <223> n at position 7 is 2'-OH guanosine or 2'-O-methyl guanosine

35 <220>

 <221> misc_feature

 <222> (8)..(8)

 <223> n at position 8 is 2'-OH guanosine or 2'-O-methyl guanosine

40 <220>

 <221> misc_feature

 <222> (10)..(10)

 <223> n at position 10 is 2'-fluoro cytosine or deoxy cytidine

45 <220>

 <221> misc_feature

 <222> (11)..(11)

 <223> n at position 11 is 2'-fluoro uridine or deoxy thymidine

50 <220>

 <221> misc_feature

 <222> (12)..(12)

 <223> n at position 12 is 2'-fluoro cytosine or deoxy cytidine

55 <220>

 <221> misc_feature

 <222> (13)..(13)

<223> n at position 13 is 2'-OH adenosine or 2'-O-methyl adenosine

<220>

<221> misc_feature

5 <222> (14)..(14)

<223> n at position 14 is 2'-OH guanosine or 2'-O-methyl guanosine

<220>

<221> misc_feature

10 <222> (15)..(15)

<223> n at position 15 is 2'-OH guanosine or 2'-O-methyl guanosine

<220>

<221> misc_feature

15 <222> (16)..(16)

<223> n at position 16 is 2'-fluoro cytosine or deoxy cytidine

<220>

<221> misc_feature

20 <222> (18)..(18)

<223> n at position 18 is 2'-fluoro cytosine or 2'-O-methyl cytosine

<220>

<221> misc_feature

25 <222> (19)..(19)

<223> n at position 19 is 2'-fluoro uridine or 2'-O-methyl uridine

<220>

<221> misc_feature

30 <222> (20)..(20)

<223> n at position 20 is 2'-OH guanosine or 2'-O-methyl guanosine

<220>

<221> misc_feature

35 <222> (21)..(21)

<223> n at position 21 is 2'-OH adenosine or 2'-O-methyl adenosine

<220>

<221> misc_feature

40 <222> (22)..(22)

<223> n at position 22 is 2'-OH guanosine or 2'-O-methyl guanosine

<220>

<221> misc_feature

45 <222> (23)..(23)

<223> n at position 23 is 2'-fluoro uridine or deoxy thymidine

<220>

<221> misc_feature

50 <222> (24)..(24)

<223> n at position 24 is 2'-fluoro cytosine or deoxy cytidine

<220>

<221> misc_feature

55 <222> (25)..(25)

<223> n at position 25 is 2'-fluoro uridine or deoxy thymidine

<220>

5 <221> misc_feature
 <222> (26)..(26)
 <223> n at position 26 is 2'-OH guanosine or 2'-O-methyl guanosine

 10 <220>
 <221> misc_feature
 <222> (27)..(27)
 <223> n at position 27 is 2'-OH adenosine or 2'-O-methyl adenosine

 15 <220>
 <221> misc_feature
 <222> (28)..(28)
 <223> n at position 28 is 2'-OH guanosine or 2'-O-methyl guanosine

 20 <220>
 <221> misc_feature
 <222> (29)..(29)
 <223> n at position 29 is 2'-fluoro uridine or deoxy thymidine

 25 <220>
 <221> misc_feature
 <222> (32)..(32)
 <223> n at position 32 is 2'-OH adenosine or 2'-O-methyl adenosine

 30 <220>
 <221> misc_feature
 <222> (34)..(34)
 <223> n at position 34 is 2'-fluoro cytosine or deoxy cytidine

 35 <220>
 <221> misc_feature
 <222> (35)..(35)
 <223> n at position 35 is 2'-fluoro uridine or deoxy thymidine

 40 <220>
 <221> misc_feature
 <222> (36)..(36)
 <223> n at position 36 is 2'-OH guanosine or 2'-O-methyl guanosine

 45 <220>
 <221> misc_feature
 <222> (38)..(38)
 <223> n at position 38 is 2'-OH guanosine or 2'-O-methyl guanosine

 50 <400> 1
 nnccgcnnun nnnnnngnnn nnnnnnnnnu unnnnnncn 38

 55 <210> 2
 <211> 38
 <212> DNA
 <213> artificial sequence

 <220>

<223> synthetic aptamer

<220>

<221> misc_feature

5 <222> (1)..(38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy, and at positions 11, 23 and 25, which are deoxy thymidine

<220>

10 <221> misc_feature

<222> (1)..(38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

15 <400> 2

cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38

<210> 3

<211> 42

20 <212> RNA

<213> artificial sequence

<220>

25 <223> synthetic aptamer

<220>

<221> misc_feature

<222> (1)..(42)

<223> all pyrimidines are 2'-fluoro

30 <400> 3

gacgaugcgg ucucaugcgu cgagugugag uuuaccuucg uc 42

<210> 4

35 <211> 39

<212> DNA

<213> artificial sequence

<220>

40 <223> synthetic aptamer

<220>

<221> misc_feature

<222> (1)..(38)

45 <223> all pyrimidines are 2'-fluoro

<220>

<221> misc_feature

<222> (1)..(38)

50 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

<220>

<221> misc_feature

55 <222> (39)..(39)

<223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

<400> 4

cgccgcgguc ucaggcgcug agucugaguu uaccugcgt 39

<210> 5

<211> 39

5 <212> DNA

<213> artificial sequence

<220>

<223> synthetic aptamer

10

<220>

<221> misc_feature

<222> (1)..(1)

15 <223> cytosine at position 1 is a modified by a 40 kDa branched (1,3-bis(mPEG-[20 kDa])-propyl-2-(4'-butamide)) PEG attached to the nucleotide via an amine linker

<220>

<221> misc_feature

<222> (1)..(38)

20 <223> all pyrimidines are 2'-fluoro

<220>

<221> misc_feature

<222> (1)..(38)

25 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

<220>

<221> misc_feature

<222> (39)..(39)

30 <223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

<400> 5

cgccgcgguc ucaggcgcug agucugaguu uaccugcgt 39

35

<210> 6

<211> 44

<212> RNA

40 <213> artificial sequence

<220>

<223> synthetic aptamer

45

<220>

<221> misc_feature

<222> (1)..(44)

<223> all pyrimidines are 2'-fluoro

50

<400> 6

aggacgaugc ggucucaugc gucgagugug aguuuuaccuu cguc 44

<210> 7

<211> 40

<212> RNA

55 <213> artificial sequence

<220>

<223> synthetic aptamer

5 <220>
 <221> misc_feature
 <222> (1)..(40)
 <223> all pyrimidines are 2'-fluoro
 10 <220>
 <221> misc_feature
 <222> (1)..(40)
 <223> all purines are 2'-O-methyl; except at positions 2, 7 and 19, wherein guanosine is 2'-OH, and positions 1 and
 34, wherein adenosine is 2'-OH
 15 <400> 7
 agcgccgcgg ucucaggcgc ugagucugag uuuaccugcg 40
 20 <210> 8
 <211> 40
 <212> RNA
 <213> artificial sequence
 25 <220>
 <221> misc_feature
 <222> (1)..(40)
 <223> all pyrimidines are 2'-fluoro
 30 <400> 8
 ggcgcgcgg ucucaggcgc ugagucugag uuuaccugcg 40
 35 <210> 9
 <211> 39
 <212> DNA
 <213> artificial sequence
 40 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32,
 wherein adenosine is 2'-OH
 45 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16, 24, 33 and 34, wherein cytidine is deoxy, and at
 50 positions 11, 23, and 25, which are deoxy thymidine
 55 <220>
 <221> misc_feature
 <222> (39)..(39)
 <223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)
 <400> 9
 cgccgcgguc tcaggcgcug agtctgaguu uaccugcgt 39

5 <210> 10
 <211> 39
 <212> DNA
 <213> artificial sequence
 10 <220>
 <223> synthetic aptamer
 15 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl, except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH
 20 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro, except at positions 10, 12, 16, 24, 33 and 34, wherein cytidine is deoxy; at positions 1, 3, and 37, wherein cytosine is 2'-O-methyl; and at positions 11, 23, and 25, which are deoxy thymidine
 25 <220>
 <221> misc_feature
 <222> (39)..(39)
 <223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)
 30 <400> 10
 cgccgcgguc tcaggcgcug agtctgagu uaccugcgt 39
 35 <210> 11
 <211> 38
 <212> DNA
 <213> artificial sequence
 40 <220>
 <223> synthetic aptamer
 45 <220>
 <221> misc feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl, except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH
 50 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 1, 3, 10, 12, 16, 24 and 37, wherein cytidine is deoxy; and at positions 11, 23 and 25, which are deoxy thymidine
 55 <400> 11
 cgccgcgguc tcaggcgcug agtctgagu uaccugcg 38
 60 <210> 12
 <211> 38
 <212> DNA
 <213> artificial sequence
 65 <220>
 <223> synthetic aptamer

5 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32,
 wherein adenosine is 2'-OH

10 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 1, 10, 12, 16 and 24, wherein cytidine is deoxy, and at positions
 11, 23 and 25, which are deoxy thymidine

15 <400> 12
 cgccgcgguc tcaggcgug agtctgagu uaccugcg 38

20 <210> 13
 <211> 38
 <212> DNA
 <213> artificial sequence

25 <220>
 <223> synthetic aptamer

30 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32,
 wherein adenosine is 2'-OH

35 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 3, 10, 12, 16 and 24, wherein cytidine is deoxy; and positions
 11, 23, and 25, which are deoxy thymidine

40 <400> 13
 cgccgcgguc tcaggcgug agtctgagu uaccugcg 38

45 <210> 14
 <211> 38
 <212> DNA
 <213> artificial sequence

50 <220>
 <223> synthetic aptamer

55 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16, 24 and 37, wherein cytidine is deoxy; and at
 positions 11, 23, and 25, which are deoxy thymidine

5 <400> 14
 cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38
 <210> 15
 <211> 38
 <212> DNA
 <213> artificial sequence
 10 <220>
 <223> synthetic aptamer
 <220>
 <221> misc_feature
 <222> (1)..(38)
 15 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH
 <220>
 <221> misc_feature
 20 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16, and 24, wherein cytidine is deoxy; at position 3, wherein cytosine is 2'-O-methyl; and at positions 11, 23, and 25, which are deoxy thymidine
 25 <400> 15
 cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38
 <210> 16
 <211> 38
 <212> DNA
 30 <213> artificial sequence
 <220>
 <223> synthetic aptamer
 35 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH
 40 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 37, wherein cytosine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine
 45 <400> 16
 cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38
 <210> 17
 <211> 38
 <212> DNA
 <213> artificial sequence
 55 <220>
 <223> synthetic aptamer
 <220>

<221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

5

<220>
 <221> misc feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 1, wherein cytosine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine

10

<400> 17
 cgccgcgguc tcaggcgug agtctgaguu uaccugcg 38

15

<210> 18
 <211> 38
 <212> DNA
 <213> artificial sequence

20

<220>
 <223> synthetic aptamer

25

<220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

30

<220>
 <221> misc feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at positions 1, 3 and 37, wherein cytosine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine

35

<400> 18
 cgccgcgguc tcaggcgug agtctgaguu uaccugcg 38

40

<210> 19
 <211> 38
 <212> DNA
 <213> artificial sequence

45

<220>
 <223> synthetic aptamer

50

<220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

55

<220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all pyrimidines are 2'-fluoro; except at positions 4, 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23, and 25, which are deoxy thymidine

<400> 19

cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38

<210> 20

<211> 38

5 <212> DNA

<213> artificial sequence

<220>

<223> synthetic aptamer

10

<220>

<221> misc_feature

<222> (1)..(38)

15 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

<220>

<221> misc_feature

<222> (1)..(38)

20 <223> all pyrimidines are 2'-fluoro; except at positions 6, 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23 and 25, which are deoxy thymidine

<400> 20

25 cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38

25

<210> 21

<211> 38

<212> DNA

<213> artificial sequence

30

<220>

<223> synthetic aptamer

30

<220>

<221> misc_feature

<222> (1)..(38)

35 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

40

<220>

<221> misc_feature

<222> (1)..(38)

45 <223> all pyrimidines are 2'-fluoro; except at positions 4, 6, 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23 and 25, which are deoxy thymidine

45

<400> 21

50 cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38

50

<210> 22

<211> 38

<212> DNA

<213> artificial sequence

55

<220>

<223> synthetic aptamer

<220>

<221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

5 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16, 18 and 24, wherein cytidine is deoxy; and at positions 11, 23 and 25, which are deoxy thymidine

10

<400> 22

cgccgcgguc tcaggcgug agtctgaguu uaccugcg 38

15 <210> 23

<211> 38

<212> DNA

<213> artificial sequence

20 <220>

20 <223> synthetic aptamer

<220>

<221> misc_feature

<222> (1) .. (38)

25 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

30 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 19, 23 and 25, which are deoxy thymidine

35

<400> 23

35 cgccgcgguc tcaggcgctg agtctgaguu uaccugcg 38

40 <210> 24

<211> 38

<212> DNA

40 <213> artificial sequence

45 <220>

<223> synthetic aptamer

45

<220>

<221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

50

<220>

<221> misc_feature

<222> (1) .. (38)

55 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16, 18 and 24, wherein cytidine is deoxy; and at positions 11, 19, 23 and 25, which are deoxy thymidine

<400> 24

55 cgccgcgguc tcaggcgctg agtctgaguu uaccugcg 38

5 <210> 25
 <211> 38
 <212> DNA
 <213> artificial sequence
 10 <220>
 <223> synthetic aptamer
 15 <220>
 <221> misc_feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH
 20 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23, 25 and 29, which are deoxy thymidine
 25 <210> 26
 <211> 38
 <212> DNA
 <213> artificial sequence
 30 <220>
 <223> synthetic aptamer
 35 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH
 40 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23, 25 and 30, which are deoxy thymidine
 45 <400> 26
 cgccgcgguc tcaggcgug agtctgagtu uaccugcg 38
 50 <210> 27
 <211> 38
 <212> DNA
 <213> artificial sequence
 55 <220>
 <223> synthetic aptamer
 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32,

wherein adenosine is 2'-OH

<220>

<221> misc feature

5 <222> (1)..(38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23, 25 and 31, which are deoxy thymidine

<400> 27

10 cgccgcgguc tcaggcgcug agtctgaguu taccugcg 38

<210> 28

<211> 38

15 <212> DNA

<213> artificial sequence

<220>

<223> synthetic aptamer

20 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

25 <220>

<221> misc_feature

<222> (1) .. (38)

30 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23, 25, 29, 30 and 31, which are deoxy thymidine

<400> 28

cgccgcgguc tcaggcgcug agtctgagtt taccugcg 38

35 <210> 29

<211> 38

<212> DNA

<213> artificial sequence

40 <220>

<223> synthetic aptamer

<220>

<221> misc_feature

45 <222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

50 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-flouro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23, 25 and 35, which are deoxy thymidine

55 <400> 29

cgccgcgguc tcaggcgcug agtctgaguu uaccctgcg 38

<210> 30

5 <211> 38
 <212> DNA
 <213> artificial sequence

 10 5 <220>
 <223> synthetic aptamer

 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

 15 10 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16, 24, 33 and 34, wherein cytidine is deoxy; at position 9, wherein uridine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine

 20 20 <400> 30
 cgccgcgguc tcaggcgcug agtctgagu uaccugcg 38

 <210> 31
 <211> 38
 25 25 <212> DNA
 <213> artificial sequence

 <220>
 <223> synthetic aptamer
 30 30 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

 <220>
 <221> misc feature
 <222> (1) .. (38)
 40 40 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 4, wherein cytosine is 2'-O-methyl; and at positions 11, 23, and 25, which are deoxy thymidine

 <400> 31
 cgccgcgguc tcaggcgcug agtctgagu uaccugcg 38
 45 45 <210> 32
 <211> 38
 <212> DNA
 <213> artificial sequence

 50 50 <220>
 <223> synthetic aptamer

 <220>
 <221> misc_feature
 <222> (1) .. (38)
 55 55 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

5 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 6,
 wherein cytosine is 2'-O-methyl; and at positions 11, 23, and 25, which are deoxy thymidine

 10 <400> 32
 cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38

 15 <210> 33
 <211> 38
 <212> DNA
 <213> artificial sequence

 20 <220>
 <223> synthetic aptamer

 25 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32,
 wherein adenosine is 2'-OH

 30 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 4
 and 6, wherein cytosine is 2'-O-methyl; and at positions 11, 23, and 25, which are deoxy thymidine

 35 <400> 33
 cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38

 40 <210> 34
 <211> 38
 <212> DNA
 <213> artificial sequence

 45 <220>
 <223> synthetic aptamer

 50 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32,
 wherein adenosine is 2'-OH

 55 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at position
 18, wherein cytosine is 2'-O-methyl

 <400> 34
 cgccgcgguc ucaggcgcug agucugaguu uaccugcg 38

 <210> 35
 <211> 38
 <212> DNA

5 <213> artificial sequence

10 <220>

15 <223> synthetic aptamer

20 <220>

25 <221> misc_feature

30 <222> (1) .. (38)

35 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

40 <220>

45 <221> misc_feature

50 <222> (1) .. (38)

55 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at position 19, wherein uridine is 2'-O-methyl

60 <400> 35

65 cggccggcuc ucaggcgcug agucugaguu uaccugcg 38

70 <210> 36

75 <211> 38

80 <212> DNA

85 <213> artificial sequence

90 <220>

95 <223> synthetic aptamer

100 <220>

105 <221> misc_feature

110 <222> (1) .. (38)

115 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

120 <220>

125 <221> misc_feature

130 <222> (1) .. (38)

135 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 18, wherein cytosine is 2'-O-methyl; and at position 19, wherein uridine is 2'-O-methyl

140 <400> 36

145 cggccggcuc ucaggcgcug agucugaguu uaccugcg 38

150 <210> 37

155 <211> 38

160 <212> DNA

165 <213> artificial sequence

170 <220>

175 <223> synthetic aptamer

180 <220>

185 <221> misc_feature

190 <222> (1) .. (38)

195 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

200 <220>

5 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 29,
 wherein uridine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine
 10 <400> 37
 cgccgcgguc tcaggcgug agtctgaguu uaccugcg 38
 <210> 38
 <211> 38
 <212> DNA
 <213> artificial sequence
 15 <220>
 <223> synthetic aptamer
 <220>
 <221> misc_feature
 <222> (1) .. (38)
 20 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32,
 wherein adenosine is 2'-OH
 <220>
 25 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 30,
 wherein uridine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine
 <400> 38
 30 cgccgcgguc tcaggcgug agtctgaguu uaccugcg 38
 <210> 39
 <211> 38
 <212> DNA
 35 <213> artificial sequence
 <220>
 <223> synthetic aptamer
 40 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32,
 wherein adenosine is 2'-OH
 45 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 31,
 50 wherein uridine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine
 <400> 39
 cgccgcgguc tcaggcgug agtctgaguu uaccugcg 38
 55 <210> 40
 <211> 38
 <212> DNA
 <213> artificial sequence

5 <220>

<223> synthetic aptamer

10 <220>

<221> misc feature

<222> (1)..(38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

15 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at positions 29, 30 and 31, wherein uridine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine

20 <400> 40

cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38

25 <210> 41

<211> 38

<212> DNA

<213> artificial sequence

30 <220>

<223> synthetic aptamer

<220>

<221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and position 32, wherein adenosine is 2'-OH

35 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at position 35, wherein uridine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine

40 <400> 41

cgccgcgguc tcaggcgcug agtctgaguu uaccugcg 38

<210> 42

<211> 38

<212> DNA

<213> artificial sequence

45 <220>

<223> synthetic aptamer

50 <220>

<221> misc_feature

<222> (1)..(38)

<223> all purines are 2'-O-methyl; except at position 5, wherein guanosine is deoxy; at position 17, wherein guanosine is 2'-OH; and position 32, wherein adenosine is 2'-OH

55 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23 and 25, which are deoxy thymidine

5 <400> 42

cggccgguc tcaggcgug agtctgaguu uaccugcg 38

<210> 43

<211> 38

<212> DNA

10 <213> artificial sequence

<220>

<223> synthetic aptamer

15 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at position 5, wherein guanosine is 2'-OH; at position 17, wherein guanosine is deoxy; and position 32, wherein adenosine is 2'-OH

20 <220>

<221> misc_feature

<222> (1) .. (38)

25 <223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23 and 25, which are deoxy thymidine

<400> 43

cggccgguc tcaggcgug agtctgaguu uaccugcg 38

30 <210> 44

<211> 38

<212> DNA

<213> artificial sequence

35 <220>

<223> synthetic aptamer

<220>

<221> misc_feature

40 <222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH; and position 32, wherein adenosine is deoxy

<220>

45 <221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; and at positions 11, 23 and 25, which are deoxy thymidine

50 <400> 44

cggccgguc tcaggcgug agtctgaguu uaccugcg 38

<210> 45

<211> 40

55 <212> DNA

<213> artificial sequence

<220>

5 <223> synthetic aptamer

10 <220>

<221> misc_feature

<222> (1)..(40)

<223> all purines are 2'-O-methyl; except at positions 6 and 18, wherein guanosine is 2'-OH; and position 33, wherein adenosine is 2'-OH

15 <220>

<221> misc_feature

<222> (1)..(40)

<223> all pyrimidines are 2'-fluoro; except at positions 11, 13, 17 and 25, wherein cytidine is deoxy; at position 40, wherein cytosine is 2'-O-methyl; and at positions 12, 24 and 26, which are deoxy thymidine

20 <400> 45

gcuucgggu ctcaggcgcu gagtctgagu uuaccuacgc 40

25 <210> 46

<211> 38

<212> DNA

<213> artificial sequence

30 <220>

<223> synthetic aptamer

35 <220>

<221> misc_feature

<222> (1)..(38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH; and position 32, wherein adenosine is deoxy

40 <220>

<221> misc_feature

<222> (1)..(38)

<223> all pyrimidines are 2'-fluoro; except at positions 10, 12, 16 and 24, wherein cytidine is deoxy; at positions 36, 37 and 38 wherein cytosine is 2'-O-methyl; and at positions 11, 23 and 25, which are deoxy thymidine

45 <400> 46

gggcgcgguc tcaggcgcug agtctgagu uaccuccc 38

50 <210> 47

<211> 40

<212> DNA

<213> artificial sequence

55 <220>

<223> synthetic aptamer

60 <220>

<221> misc feature

<222> (1)..(40)

<223> all purines are 2'-O-methyl; except at positions 6 and 18, wherein guanosine is 2'-OH; and position 33, wherein adenosine is 2'-OH

65 <220>

<221> misc_feature

<222> (1)..(40)

<223> all pyrimidines are 2'-fluoro; except at positions 11, 13, 17 and 25, wherein cytidine is deoxy; at position 40,

wherein cytosine is 2'-O-methyl; and at positions 12, 24 and 26, which are deoxy thymidine

<400> 47
gccccgggu ctcaggcgcu gagtctgagu uuaccugcgc 40

5
<210> 48
<211> 45
<212> DNA
<213> artificial sequence

10 <220>
<223> synthetic aptamer

<220>
<221> misc_feature
<222> (1) .. (44)
<223> all purines are 2'-O-methyl; except at positions 8 and 20, wherein guanosine is 2'-OH; and position 35 wherein adenosine is 2'-OH

20 <220>
<221> misc_feature
<222> (1) .. (44)
<223> all pyrimidines are 2'-fluoro

25 <220>
<221> misc_feature
<222> (45) .. (45)
<223> thymidine at position 45 is a 3' inverted deoxy thymidine (3'-3' linked)

30 <400> 48
ggacgcccgcg gucucaggcg cugagucuga guuuaccugc gucut 45

<210> 49
<211> 42

35 <212> DNA
<213> artificial sequence

<220>
<223> synthetic aptamer

<220>
<221> misc_feature
<222> (1) .. (42)
<223> all purines are 2'-O-methyl; except at positions 7 and 19, wherein guanosine is 2'-OH; and at position 34,
wherein adenosine is 2'-OH

<220>
<221> misc_feature
<222> (1) .. (42)
<223> all cytosines are 2'-fluoro; except at positions 12, 14, 18, 26, 35 and 36, which are deoxy cytidine; and at positions 20, 41 and 42, wherein cytosine is 2'-O-methyl

55 <220>
<221> misc_feature
<222> (1) .. (42)
<223> all uridines are 2'-fluoro; except at position 21, wherein uridine is 2'-O-methyl; and at positions 13, 25, 27,
31 and 37, which are deoxy thymidine

5 <400> 49
ggcgccgcgg uctcaggcgc ugagtctgag tuuacctgcg cc 42

10 <210> 50
<211> 42
<212> DNA
<213> artificial sequence

15 <220>
<223> synthetic aptamer

20 <220>
<221> misc_feature
<222> (1) .. (42)
<223> all purines are 2'-O-methyl; except at positions 7 and 19, wherein guanosine is 2'-OH; and at position 34, wherein adenosine is 2'-OH

25 <220>
<221> misc_feature
<222> (1) .. (42)
<223> all cytosines are 2'-fluoro; except at positions 12, 14, 18, 26, 35, 36 and 39, which are deoxy cytidine; and at positions 3, 20, 41 and 42, wherein cytosine is 2'-O-methyl

30 <220>
<221> misc_feature
<222> (1)..(42)
<223> uridine at position 11 is 2'-fluoro; uridine at position 21 is 2'-O-methyl; positions 13, 25, 27, 31, 32, 33 and 37 are deoxy thymidine

35 <400> 50
ggcgccgcgg uctcaggcgc ugagtctgag tttacctgcg cc 42

40 <210> 51
<211> 42
<212> DNA
<213> artificial sequence

45 <220>
<223> synthetic aptamer

50 <220>
<221> misc_feature
<222> (1)..(42)
<223> all purines are 2'-O-methyl; except at positions 7 and 19, wherein guanosine is 2'-OH

55 <220>
<221> misc_feature
<222> (1) .. (42)
<223> cytosine at positions 5, 6, 8, 12, 14, 18, 26, 35, 36 and 39 are deoxy cytidine; and cystosine at positions 3, 20, 41 and 42 are 2'-O-methyl

60 <220>
<221> misc_feature
<222> (1) .. (42)
<223> uridine at position 21 is 2'-O-methyl; positions 11, 13, 25, 27, 31, 32, 33 and 37 are deoxy thymidine

65 <400> 51
ggcgccgcgg ttcaggcgc ugagtctgag tttacctgcg cc 42

5 <210> 52
 <211> 42
 <212> DNA
 <213> artificial sequence
 10 <220>
 <223> synthetic aptamer
 <220>
 15 <221> misc_feature
 <222> (1) .. (42)
 <223> all purines are 2'-O-methyl; except at positions 7 and 19, wherein guanosine is 2'-OH
 <220>
 20 <221> misc_feature
 <222> (1) .. (42)
 <223> uridine at positions 13, 21, 25 and 27 are 2'-O-methyl; positions 11, 31, 32, 33 and 37 are deoxy thymidine
 <220>
 25 <221> misc_feature
 <222> (1) .. (42)
 <223> cytosine at positions 5, 6, 8, 12, 18, 35, 36 and 39 are deoxy cytidine; and cytosine at positions 3, 14, 20, 26, 41 and 42 are 2'-O-methyl
 <400> 52
 ggcgcgcgg tcucaggcgc ugagucugag tttacctgcg cc 42
 <210> 53
 <211> 40
 30 <212> RNA
 <213> artificial sequence
 <220>
 35 <223> synthetic aptamer
 <220>
 <221> misc feature
 <222> (1)..(1)
 <223> adenosine at position 1 has biotin conjugated to the 5' end
 <220>
 40 <221> misc_feature
 <222> (1) .. (40)
 <223> all purines are 2'-O-methyl; except at positions 3, 8 and 20, wherein guanosine is 2'-OH; and at position 2,
 45 wherein adenosine is 2'-OH
 <220>
 <221> misc_feature
 <222> (1) .. (40)
 <223> all pyrimidines are 2'-fluoro
 <400> 53
 agcgccgcgg ucucaggcgc ugagucugag uuuaccugcg 40
 <210> 54
 <211> 42
 <212> DNA
 <213> artificial sequence

5 <220>
 <223> synthetic aptamer

 10 <220>
 <221> misc_feature
 <222> (1) .. (42)
 <223> all purines are 2'-O-methyl; except at positions 7 and 19, wherein guanosine is 2'-OH; and at position 34, wherein adenosine is 2'-OH

 15 <220>
 <221> misc feature
 <222> (1)..(42)
 <223> all cytosines are 2'-fluoro; except at positions 12, 14, 18 and 26, which are deoxy cytidine; and at positions 41 and 42, wherein cytosine is 2'-O-methyl

 20 <220>
 <221> misc_feature
 <222> (1) .. (42)
 <223> all uridines are 2'-fluoro; positions 13, 25, and 27 are deoxy thymidine

 25 <210> 54
 <211> 42
 <212> RNA
 <213> artificial sequence

 30 <220>
 <223> synthetic aptamer

 35 <220>
 <221> misc_feature
 <222> (1) .. (42)
 <223> all purines are 2'-O-methyl; except at positions 7 and 19, wherein guanosine is 2'-OH; and at position 34, wherein adenosine is 2'-OH

 40 <220>
 <221> misc_feature
 <222> (1) .. (42)
 <223> all cytosines are 2'-fluoro; except at positions 12, 14, 18, 26, 41 and 42, wherein cytosine is 2'-O-methyl

 45 <220>
 <221> misc_feature
 <222> (1) .. (42)
 <223> all uridines are 2'-fluoro; except at positions 13, 25, and 27, wherein uridine is 2'-O-methyl

 50 <400> 55
 ggcgcgcgg uctcaggcgc ugagtctgag uuuaccugcg cc 42

 55 <210> 56
 <211> 39
 <212> DNA
 <213> artificial sequence

 <220>
 <223> synthetic aptamer

5 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro

 10 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH

 15 <220>
 <221> misc_feature
 <222> (39) .. (39)
 <223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

 20 <400> 56
 cgccgcgguc ucaggcgcug agucugaguu uaccugcgt 39

 25 <210> 57
 <211> 39
 <212> DNA
 <213> artificial sequence

 30 <220>
 <223> synthetic aptamer

 35 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at position 18, wherein cytosine is 2'-O-methyl; and at position 19 wherein uridine is 2'-O-methyl

 40 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH; and at position 32, wherein adenosine is 2'-OH

 45 <220>
 <221> misc_feature
 <222> (39) .. (39)
 <223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

 50 <400> 57
 cgccgcgguc ucaggcgcug agucugaguu uaccugcgt 39

 55 <210> 58
 <211> 39
 <212> DNA
 <213> artificial sequence

 60 <220>
 <223> synthetic aptamer

 65 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at position 29, which is deoxy thymidine

5 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH; and at position 32,
 wherein adenosine is 2'-OH

10 <220>
 <221> misc_feature
 <222> (39) .. (39)
 <223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

15 <400> 58
 cgccgcgguc ucaggcgcug agucugagt uaccugcgt 39

20 <210> 59
 <211> 39
 <212> DNA
 <213> artificial sequence

25 <220>
 <221> misc feature
 <222> (1)..(38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH

30 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at position 18, wherein cytosine is 2'-O-methyl; and position 19, wherein
 uridine is 2'-O-methyl

35 <220>
 <221> misc_feature
 <222> (39) .. (39)
 <223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

40 <400> 59
 cgccgcgguc ucaggcgcug agucugagu uaccugcgt 39

45 <210> 60
 <211> 39
 <212> DNA
 <213> artificial sequence

50 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH

55 <220>
 <221> misc_feature
 <222> (1) .. (38)
 <223> all pyrimidines are 2'-fluoro; except at position 18, wherein cytosine is 2'-O-methyl; at position 19, wherein

uridine is 2'-O-methyl; and at position 29, which is deoxy thymidine

5 <220>

<221> misc feature

<222> (39)..(39)

<223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

10 <400> 60

cgccgcgguc ucaggcgcug agucugagt uaccugcgt 39

15 <210> 61

<211> 39

<212> DNA

<213> artificial sequence

20 <220>

<221> misc_feature

<222> (1) .. (1)

<223> cytosine at position 1 is modified by a 20 kDa PEG attached to the nucleotide via an amine linker

25 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro

30 <220>

<221> misc_feature

<222> (1)..(38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

35 <220>

<221> misc_feature

<222> (39) .. (39)

<223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

40 <400> 61

cgccgcgguc ucaggcgcug agucugagu uaccugcgt 39

45 <210> 62

<211> 39

<212> DNA

<213> artificial sequence

50 <220>

<223> synthetic aptamer

55 <220>

<221> misc_feature

<222> (1) .. (1)

<223> cytosine at position 1 is modified by a 30 kDa PEG attached to the nucleotide via an amine linker

55 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro

<220>

<221> misc_feature

5 <222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

<220>

10 <221> misc_feature

<222> (39) .. (39)

<223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

<400> 62

15 cgccgcgguc ucaggcgcug agucugaguu uaccugcgt 39

<210> 63

<211> 39

<212> DNA

20 <213> artificial sequence

<220>

<223> synthetic aptamer

25 <220>

<221> misc_feature

<222> (1) .. (1)

<223> cytosine at position 1 is modified by a 5' amine linker

30 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro

35 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

40 <220>

<221> misc_feature

<222> (39) .. (39)

<223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

45 <400> 63

cgccgcgguc ucaggcgcug agucugaguu uaccugcgt 39

<210> 64

50 <211> 39

<212> DNA

<213> artificial sequence

<220>

55 <223> synthetic aptamer

<220>

<221> misc_feature

<222> (1)..(1)

<223> cytosine at position 1 is modified by a 10 kDa PEG attached to the nucleotide via an amine linker

5 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro

10 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

15 <220>

<221> misc_feature

<222> (39) .. (39)

<223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

20 <400> 64

cgccgcgguc ucaggcgcug agucugaguu uaccugcgt 39

<210> 65

<211> 39

25 <212> DNA

<213> artificial sequence

30 <220>

<223> synthetic aptamer

35 <220>

<221> misc_feature

<222> (1) .. (1)

<223> cytosine at position 1 is modified by a linear 40 kDa PEG attached to the nucleotide via an amine linker

40 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro

45 <220>

<221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

50 <220>

<221> misc_feature

<222> (39) .. (39)

<223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

<400> 65

cgccgcgguc ucaggcgcug agucugaguu uaccugcgt 39

55 <210> 66

<211> 38

<212> RNA

<213> artificial sequence

<220>

<223> synthetic aptamer

<220>

5 <221> misc_feature

<222> (1) .. (1)

<223> cytosine at position 1 is modified by a 20 kDa PEG attached to the nucleotide via an amine linker

<220>

10 <221> misc_feature

<222> (1) .. (38)

<223> all pyrimidines are 2'-fluoro

<220>

15 <221> misc_feature

<222> (1) .. (38)

<223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

<220>

20 <221> misc_feature

<222> (38) .. (38)

<223> guanosine at position 38 is modified by a 20 kDa PEG attached to the nucleotide via an amine linker

25 <400> 66

cgccgcgguc ucaggcgcug agucugaguu uaccugcg 38

<210> 67

<211> 39

30 <212> DNA

<213> artificial sequence

<220>

35 <223> synthetic aptamer

<220>

<221> misc_feature

<222> (1) .. (1)

40 <223> cytosine at position 1 is modified by a 40 kDa branched (2,3-bis(mPEG-[20 kDa])-propyl-1-carbamoyl) PEG attached to the nucleotide via an amine linker

<220>

<221> misc feature

<222> (1)..(38)

45 <223> all pyrimidines are 2'-fluoro

<220>

<221> misc_feature

<222> (1) .. (38)

50 <223> all purines are 2'-O-methyl; except at positions 5 and 17, wherein guanosine is 2'-OH, and at position 32, wherein adenosine is 2'-OH

<220>

<221> misc_feature

<222> (39) .. (39)

55 <223> thymidine at position 39 is a 3' inverted deoxy thymidine (3'-3' linked)

<400> 67

cggcgccguc ucaggcgcug agucugaguu uaccugcgt 39
 <210> 68
 <211> 46
 5 <212> RNA
 <213> artificial sequence

 <220>
 <223> synthetic aptamer
 10 <220>
 <221> misc feature
 <222> (1)..(46)
 <223> all pyrimidines are 2'-fluoro
 15 <400> 68
 ggcgauuacu gggacggacu cgcauguga gcccagacga cucgcc 46

 <210> 69
 20 <211> 40
 <212> RNA
 <213> artificial sequence

 <220>
 25 <223> synthetic aptamer

 <220>
 <221> misc_feature
 <222> (1) .. (40)
 30 <223> all pyrimidines are 2'-fluoro

 <400> 69
 ggcuucugaa gauuauuucg cgaugugaac uccagacccc 40

 35 <210> 70
 <211> 92
 <212> DNA
 <213> artificial sequence

 40 <220>
 <223> synthetic template

 <220>
 <221> misc_feature
 45 <222> (40) .. (69)
 <223> n may be any nucleotide (a, c, g, or t)

 <400> 70

 50 taatacgact cactataggg agaggagaga acgttctacn nnnnnnnnnn nnnnnnnnnn 60
 nnnnnnnnnng gtcgatcgat cgatcatcga tg 92

 55 <210> 71
 <211> 39
 <212> DNA
 <213> artificial sequence

5 <220> 71
 taatacact cactatagg agaggagaga acgttctac 39
 <210> 72
 <211> 23
 <212> DNA
 <213> artificial sequence
 10 <220>
 <223> synthetic primer
 15 <400> 72
 catcgatcat cgatcgatcg acc 23
 <210> 73
 <211> 22
 <212> RNA
 <213> artificial sequence
 20 <220>
 <223> synthetic fixed region
 25 <400> 73
 gggagaggag agaacguucu ac 22
 <210> 74
 <211> 23
 <212> RNA
 <213> artificial sequence
 30 <220>
 <223> synthetic fixed region
 35 <400> 74
 ggucgaucga ucgaucaucg aug 23
 <210> 75
 <211> 75
 <212> DNA
 <213> artificial sequence
 40 <220>
 <223> synthetic aptamer
 45 <220>
 <221> misc_feature
 <222> (1) .. (75)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl
 50 <400> 75
 55 gggagaggag agaacguucu accuugguuu ggcacaggca uacauacgca gggucgauc
 gaucgaucau cgaug 60
 65 75

5 <210> 76
 <211> 32
 <212> DNA
 <213> artificial sequence
 10 <220>
 <223> synthetic aptamer
 <220>
 15 <221> misc_feature
 <222> (1) .. (32)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl
 <400> 76
 20 ccuugguuug gcacaggcau acauacgcag gg 32
 <210> 77
 <211> 32
 <212> DNA
 25 <213> artificial sequence
 <220>
 <223> synthetic aptamer
 <220>
 30 <221> misc feature
 <222> (1)..(32)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl
 <400> 77
 35 ccuugguuug gcacaggcau acaaacgcag gg 32
 <210> 78
 <211> 25
 <212> DNA
 40 <213> artificial sequence
 <220>
 <223> synthetic aptamer
 <220>
 45 <221> misc_feature
 <222> (1) .. (25)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl
 <400> 78
 50 ggguuuggca caggcauaca uaccc 25
 <210> 79
 <211> 25
 <212> DNA
 <213> artificial sequence
 <220>
 55 <223> synthetic aptamer
 <220>
 <221> misc_feature

<222> (1) .. (25)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

5 <400> 79
 ggguuuggca caggcauaca aaccc 25

<210> 80
 <211> 32
 <212> DNA
 10 <213> artificial sequence

<220>
 <223> synthetic aptamer

15 <220>
 <221> misc_feature
 <222> (1) .. (32)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

20 <400> 80
 ggcggcacag gcauacauac gcagggguucg cc 32

<210> 81
 <211> 47
 25 <212> DNA
 <213> artificial sequence

<220>
 <223> synthetic aptamer

30 <220>
 <221> misc_feature
 <222> (1) .. (47)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

35 <400> 81
 cguucuaccc ugguuuggca caggcauaca uacgcagggg ucgaucg 47

<210> 82
 <211> 88
 <212> DNA
 <213> artificial sequence

40 <220>
 <223> synthetic template

<220>
 <221> misc_feature
 <222> (40) .. (69)
 50 <223> n may be any nucleotide (a, t, c, or g)

<400> 82

55 taatacgact cactataggg agaggagaga acgttctacn nnnnnnnnnn nnnnnnnnnn
 nnnnnnnnnng ttacgactag catcgatg 60
 88

5	<210> 83 <211> 39 <212> DNA <213> artificial sequence	
10	<220> <223> synthetic template	
15	<400> 83 cttggtttgg cacaggcata catacgagg ggtcgatcg	39
20	<210> 84 <211> 39 <212> DNA <213> artificial sequence	
25	<220> <223> synthetic primer	
30	<400> 84 taatacgact cactataggg agaggagaga acgttctac	39
35	<210> 85 <211> 19 <212> DNA <213> artificial sequence	
40	<220> <223> synthetic primer	
45	<400> 85 catcgatgct agtcgtaac	19
50	<210> 86 <211> 22 <212> RNA <213> artificial sequence	
55	<220> <223> synthetic fixed region	
60	<400> 86 gggagaggag agaacguucu ac	22
65	<210> 87 <211> 19 <212> RNA <213> artificial sequence	
70	<220> <223> synthetic fixed region	
75	<400> 87 guuacgacua gcaucgaug	19
80	<210> 88 <211> 80 <212> DNA	

<213> artificial sequence

<220>

<223> synthetic aptamer

5

<220>

<221> misc_feature

<222> (1)..(80)

<223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

10

<400> 88

gggagaggag agaacguucu accuugguuu ggcacaggca uacauacgca gggucgauc . 60

15

gguuacgacu agcaucgau g 80

<210> 89

<211> 80

20

<212> DNA

<213> artificial sequence

25

<220>
<223> synthetic aptamer

<220>

<221> misc_feature

<222> (1) .. (80)

<223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

30

<400> 89

gggagaggag agaacguucu accuugguuu ggcacaggca uacauacgca gggucgauc 60

35

uguuacgacu agcaucgau g 80

<210> 90

<211> 80

40

<212> DNA

<213> artificial sequence

45

<220>
<223> synthetic aptamer

<220>

<221> misc_feature

<222> (1)..(80)

<223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

50

<400> 90

gggagaggag agaacguucu accuugguuu ggcacaggca uaaauacgca gggucgauc 60

55

gguuacgacu agcaucgau g 80

<210> 91

5 <211> 80
 <212> DNA
 <213> artificial sequence

 10 <220>
 <223> synthetic aptamer

 15 <220>
 <221> misc_feature
 <222> (1) .. (80)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

 20 <400> 91

 25 <210> 92
 <211> 78
 <212> DNA
 <213> artificial sequence

 30 <220>
 <223> synthetic aptamer

 35 <220>
 <221> misc_feature
 <222> (1) .. (78)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

 40 <400> 92

 45 <210> 93
 <211> 80
 <212> DNA
 <213> artificial sequence

 50 <220>
 <223> synthetic aptamer

 55 <220>
 <221> misc_feature
 <222> (1) .. (80)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

 60 <400> 93

 65 <210> 94
 <211> 80
 <212> DNA
 <213> artificial sequence

 70 <220>
 <223> synthetic aptamer

 75 <220>
 <221> misc_feature
 <222> (1) .. (80)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

 80 <400> 94

 85 <210> 95
 <211> 80
 <212> DNA
 <213> artificial sequence

 90 <220>
 <223> synthetic aptamer

5 <210> 94
<211> 69
<212> DNA
<213> artificial sequence

10 <220>
<223> synthetic aptamer

15 <220>
<221> misc_feature
<222> (1) .. (69)
<223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

20 <400> 94

25 <210> 95
<211> 79
<212> DNA
<213> artificial sequence

30 <220>
<223> synthetic aptamer

35 <220>
<221> misc_feature
<222> (1) .. (79)
<223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

40 <400> 95

45 <210> 96
<211> 75
<212> DNA
<213> artificial sequence

50 <220>
<223> synthetic aptamer

55 <220>
<221> misc_feature
<222> (1) .. (75)
<223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

60 gggagaggag agaacguucu accuugguuu ggcacaggca uacauacgca ggucgaucgg
uuacgacua 69

65 gggagaggag agaacguucu accuuaggua cgcacugua uacauacaca cggcaaucg
guuacgacua gcaucgaug 79

70 <210> 96
<211> 75
<212> DNA
<213> artificial sequence

75 <220>
<223> synthetic aptamer

80 <220>
<221> misc_feature
<222> (34) .. (34)
<223> n may be any nucleotide (a, u, c, or g)

5 <220>
 <221> misc_feature
 <222> (43) .. (43)
 <223> n may be any nucleotide (a, u, c, or g)

6 <400> 96

10 gggagaggag agaacguucu accuugguuu ggcncaggca uanauacgca cgggucgauc 60
 gguuacgacu agcau 75

15 <210> 97
 <211> 80
 <212> DNA
 <213> artificial sequence

20 <220>
 <223> synthetic aptamer

25 <220>
 <221> misc_feature
 <222> (1) .. (80)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

26 <400> 97

30 gggagaggag agaacguucu accuuucucu gccacaagca uaccuucgcg ggguucuauu 60
 gguuacgacu agcaucgaug 80

35 <210> 98
 <211> 79
 <212> DNA
 <213> artificial sequence

40 <220>
 <223> synthetic aptamer

45 <220>
 <221> misc_feature
 <222> (1) .. (79)
 <223> wherein all purines are deoxy, and all pyrimidines are 2'-O-methyl

46 <400> 98

50 gggagaggag agaacguucu accuugguuu ggcacaggca uauauacgca gggucgaucc 60
 guuacgacua gcaucgaug 79

55 <210> 99
 <211> 93
 <212> DNA
 <213> artificial sequence

<220>
<223> synthetic template

5 <220>
<221> misc_feature
<222> (25) .. (54)
<223> n may be any nucleotide (a, t, c, or g)

10 <400> 99

15 catcgatgct agtcgtaacg atccnnnnnn nnnnnnnnnn nnnnnnnnnn nnnncgagaa 60
 cgttctctcc tctccctata gtgagtcgta tta 93

20 <210> 100
<211> 92
<212> DNA
<213> artificial

25 <220>
<223> synthetic template

30 <220>
<221> misc_feature
<222> (24) .. (53)
<223> n may be any nucleotide (a, t, c, or g)

35 <400> 100

40 catgcatcgc gactgactag ccgnnnnnnn nnnnnnnnnn nnnnnnnnnn nnngtagaac 60
 gttctctcct ctccctatacg tgagtcgtat ta 92

45 <210> 101
<211> 92
<212> DNA
<213> artificial

50 <220>
<223> synthetic template

55 <220>
<221> misc_feature
<222> (24) .. (53)
<223> n may be any nucleotide (a, t, c, or g)

60 <400> 101

65 catcgatcga tcgatcgaca gcgnnnnnnn nnnnnnnnnn nnnnnnnnnn nnngtagaac 60
 gttctctcct ctccctatacg tgagtcgtat ta 92

70 <210> 102
<211> 1676

<212> PRT
 <213> artificial

5 <220>
 <223> synthetic C5

<400> 102

10	Met	Gly	Leu	Leu	Gly	Ile	Leu	Cys	Phe	Leu	Ile	Phe	Leu	Gly	Lys	Thr
	1				5					10				15		
15	Trp	Gly	Gln	Glu	Gln	Thr	Tyr	Val	Ile	Ser	Ala	Pro	Lys	Ile	Phe	Arg
						20			25					30		
20	Val	Gly	Ala	Ser	Glu	Asn	Ile	Val	Ile	Gln	Val	Tyr	Gly	Tyr	Thr	Glu
						35			40				45			
25	Ala	Phe	Asp	Ala	Thr	Ile	Ser	Ile	Lys	Ser	Tyr	Pro	Asp	Lys	Lys	Phe
						50			55				60			
30	Ser	Tyr	Ser	Ser	Gly	His	Val	His	Leu	Ser	Ser	Glu	Asn	Lys	Phe	Gln
						65			70			75			80	
35	Asn	Ser	Ala	Ile	Leu	Thr	Ile	Gln	Pro	Lys	Gln	Leu	Pro	Gly	Gly	Gln
						85			90				95			
40	Asn	Pro	Val	Ser	Tyr	Val	Tyr	Leu	Glu	Val	Val	Ser	Lys	His	Phe	Ser
						100			105				110			
45	Lys	Ser	Lys	Arg	Met	Pro	Ile	Thr	Tyr	Asp	Asn	Gly	Phe	Leu	Phe	Ile
						115			120				125			
50	His	Thr	Asp	Lys	Pro	Val	Tyr	Thr	Pro	Asp	Gln	Ser	Val	Lys	Val	Arg
						130			135				140			
55	Val	Tyr	Ser	Leu	Asn	Asp	Asp	Leu	Lys	Pro	Ala	Lys	Arg	Glu	Thr	Val
						145			150			155			160	
60	Leu	Thr	Phe	Ile	Asp	Pro	Glu	Gly	Ser	Glu	Val	Asp	Met	Val	Glu	Glu
						165			170				175			
65	Ile	Asp	His	Ile	Gly	Ile	Ile	Ser	Phe	Pro	Asp	Phe	Lys	Ile	Pro	Ser
						180			185				190			

EP 2 578 683 B9

Asn Pro Arg Tyr Gly Met Trp Thr Ile Lys Ala Lys Tyr Lys Glu Asp
195 200 205

5 Phe Ser Thr Thr Gly Thr Ala Tyr Phe Glu Val Lys Glu Tyr Val Leu
210 215 220

10 Pro His Phe Ser Val Ser Ile Glu Pro Glu Tyr Asn Phe Ile Gly Tyr
225 230 235 240

15 Lys Asn Phe Lys Asn Phe Glu Ile Thr Ile Lys Ala Arg Tyr Phe Tyr
245 250 255

20 Asn Lys Val Val Thr Glu Ala Asp Val Tyr Ile Thr Phe Gly Ile Arg
260 265 270

25 Glu Asp Leu Lys Asp Asp Gln Lys Glu Met Met Gln Thr Ala Met Gln
275 280 285

30 Asn Thr Met Leu Ile Asn Gly Ile Ala Gln Val Thr Phe Asp Ser Glu
290 295 300

35 Thr Ala Val Lys Glu Leu Ser Tyr Tyr Ser Leu Glu Asp Leu Asn Asn
305 310 315 320

40 Lys Tyr Leu Tyr Ile Ala Val Thr Val Ile Glu Ser Thr Gly Gly Phe
325 330 335

45 Ser Glu Glu Ala Glu Ile Pro Gly Ile Lys Tyr Val Leu Ser Pro Tyr
340 345 350

50 Lys Leu Asn Leu Val Ala Thr Pro Leu Phe Leu Lys Pro Gly Ile Pro
355 360 365

55 Tyr Pro Ile Lys Val Gln Val Lys Asp Ser Leu Asp Gln Leu Val Gly
370 375 380

Gly Val Pro Val Thr Leu Asn Ala Gln Thr Ile Asp Val Asn Gln Glu
385 390 395 400

Thr Ser Asp Leu Asp Pro Ser Lys Ser Val Thr Arg Val Asp Asp Gly
405 410 415

Val Ala Ser Phe Val Leu Asn Leu Pro Ser Gly Val Thr Val Leu Glu
420 425 430

Phe Asn Val Lys Thr Asp Ala Pro Asp Leu Pro Glu Glu Asn Gln Ala

435

440

445

5 Arg Glu Gly Tyr Arg Ala Ile Ala Tyr Ser Ser Leu Ser Gln Ser Tyr
 450 455 460

10 Leu Tyr Ile Asp Trp Thr Asp Asn His Lys Ala Leu Leu Val Gly Glu
 465 470 475 480

15 His Leu Asn Ile Ile Val Thr Pro Lys Ser Pro Tyr Ile Asp Lys Ile
 485 490 495

20 Thr His Tyr Asn Tyr Leu Ile Leu Ser Lys Gly Lys Ile Ile His Phe
 500 505 510

25 Gly Thr Arg Glu Lys Phe Ser Asp Ala Ser Tyr Gln Ser Ile Asn Ile
 515 520 525

30 Pro Val Thr Gln Asn Met Val Pro Ser Ser Arg Leu Leu Val Tyr Tyr
 530 535 540

35 Ile Val Thr Gly Glu Gln Thr Ala Glu Leu Val Ser Asp Ser Val Trp
 545 550 555 560

40 Leu Asn Ile Glu Glu Lys Cys Gly Asn Gln Leu Gln Val His Leu Ser
 565 570 575

45 Pro Asp Ala Asp Ala Tyr Ser Pro Gly Gln Thr Val Ser Leu Asn Met
 580 585 590

50 Ala Thr Gly Met Asp Ser Trp Val Ala Leu Ala Ala Val Asp Ser Ala
 595 600 605

55 Val Tyr Gly Val Gln Arg Gly Ala Lys Lys Pro Leu Glu Arg Val Phe
 610 615 620

60 Gln Phe Leu Glu Lys Ser Asp Leu Gly Cys Gly Ala Gly Gly Leu
 625 630 635 640

65 Asn Asn Ala Asn Val Phe His Leu Ala Gly Leu Thr Phe Leu Thr Asn
 645 650 655

70 Ala Asn Ala Asp Asp Ser Gln Glu Asn Asp Glu Pro Cys Lys Glu Ile
 660 665 670

75 Leu Arg Pro Arg Arg Thr Leu Gln Lys Lys Ile Glu Glu Ile Ala Ala
 675 680 685

EP 2 578 683 B9

Lys Tyr Lys His Ser Val Val Lys Lys Cys Cys Tyr Asp Gly Ala Cys
690 695 700

5 Val Asn Asn Asp Glu Thr Cys Glu Gln Arg Ala Ala Arg Ile Ser Leu
705 710 715 720

10 Gly Pro Arg Cys Ile Lys Ala Phe Thr Glu Cys Cys Val Val Ala Ser
725 730 735

15 Gln Leu Arg Ala Asn Ile Ser His Lys Asp Met Gln Leu Gly Arg Leu
740 745 750

20 His Met Lys Thr Leu Leu Pro Val Ser Lys Pro Glu Ile Arg Ser Tyr
755 760 765

25 Phe Pro Glu Ser Trp Leu Trp Glu Val His Leu Val Pro Arg Arg Lys
770 775 780

30 Gln Leu Gln Phe Ala Leu Pro Asp Ser Leu Thr Thr Trp Glu Ile Gln
785 790 795 800

35 Gly Val Gly Ile Ser Asn Thr Gly Ile Cys Val Ala Asp Thr Val Lys
805 810 815

40 Ala Lys Val Phe Lys Asp Val Phe Leu Glu Met Asn Ile Pro Tyr Ser
820 825 830

45 Val Val Arg Gly Glu Gln Ile Gln Leu Lys Gly Thr Val Tyr Asn Tyr
835 840 845

50 Arg Thr Ser Gly Met Gln Phe Cys Val Lys Met Ser Ala Val Glu Gly
850 855 860

55 Ile Cys Thr Ser Glu Ser Pro Val Ile Asp His Gln Gly Thr Lys Ser
865 870 875 880

Ser Lys Cys Val Arg Gln Lys Val Glu Gly Ser Ser Ser His Leu Val
885 890 895

Thr Phe Thr Val Leu Pro Leu Glu Ile Gly Leu His Asn Ile Asn Phe
900 905 910

55 Ser Leu Glu Thr Trp Phe Gly Lys Glu Ile Leu Val Lys Thr Leu Arg
915 920 925

Val Val Pro Glu Gly Val Lys Arg Glu Ser Tyr Ser Gly Val Thr Leu
 930 935 940

5 Asp Pro Arg Gly Ile Tyr Gly Thr Ile Ser Arg Arg Lys Glu Phe Pro
 945 950 955 960

10 Tyr Arg Ile Pro Leu Asp Leu Val Pro Lys Thr Glu Ile Lys Arg Ile
 965 970 975

15 Leu Ser Val Lys Gly Leu Leu Val Gly Glu Ile Leu Ser Ala Val Leu
 980 985 990

Ser Gln Glu Gly Ile Asn Ile Leu Thr His Leu Pro Lys Gly Ser Ala
 995 1000 1005

20 Glu Ala Glu Leu Met Ser Val Val Pro Val Phe Tyr Val Phe His
 1010 1015 1020

25 Tyr Leu Glu Thr Gly Asn His Trp Asn Ile Phe His Ser Asp Pro
 1025 1030 1035

30 Leu Ile Glu Lys Gln Lys Leu Lys Lys Lys Leu Lys Glu Gly Met
 1040 1045 1050

Leu Ser Ile Met Ser Tyr Arg Asn Ala Asp Tyr Ser Tyr Ser Val
 1055 1060 1065

35 Trp Lys Gly Gly Ser Ala Ser Thr Trp Leu Thr Ala Phe Ala Leu
 1070 1075 1080

40 Arg Val Leu Gly Gln Val Asn Lys Tyr Val Glu Gln Asn Gln Asn
 1085 1090 1095

45 Ser Ile Cys Asn Ser Leu Leu Trp Leu Val Glu Asn Tyr Gln Leu
 1100 1105 1110

Asp Asn Gly Ser Phe Lys Glu Asn Ser Gln Tyr Gln Pro Ile Lys
 1115 1120 1125

50 Leu Gln Gly Thr Leu Pro Val Glu Ala Arg Glu Asn Ser Leu Tyr
 1130 1135 1140

55 Leu Thr Ala Phe Thr Val Ile Gly Ile Arg Lys Ala Phe Asp Ile
 1145 1150 1155

EP 2 578 683 B9

Cys Pro Leu Val Lys Ile Asp Thr Ala Leu Ile Lys Ala Asp Asn
 1160 1165 1170

5 Phe Leu Leu Glu Asn Thr Leu Pro Ala Gln Ser Thr Phe Thr Leu
 1175 1180 1185

10 Ala Ile Ser Ala Tyr Ala Leu Ser Leu Gly Asp Lys Thr His Pro
 1190 1195 1200

15 Gln Phe Arg Ser Ile Val Ser Ala Leu Lys Arg Glu Ala Leu Val
 1205 1210 1215

20 Lys Gly Asn Pro Pro Ile Tyr Arg Phe Trp Lys Asp Asn Leu Gln
 1220 1225 1230

25 His Lys Asp Ser Ser Val Pro Asn Thr Gly Thr Ala Arg Met Val
 1235 1240 1245

30 Glu Thr Thr Ala Tyr Ala Leu Leu Thr Ser Leu Asn Leu Lys Asp
 1250 1255 1260

35 Ile Asn Tyr Val Asn Pro Val Ile Lys Trp Leu Ser Glu Glu Gln
 1265 1270 1275

40 Arg Tyr Gly Gly Gly Phe Tyr Ser Thr Gln Asp Thr Ile Asn Ala
 1280 1285 1290

45 Ile Glu Gly Leu Thr Glu Tyr Ser Leu Leu Val Lys Gln Leu Arg
 1295 1300 1305

50 Leu Ser Met Asp Ile Asp Val Ser Tyr Lys His Lys Gly Ala Leu
 1310 1315 1320

His Asn Tyr Lys Met Thr Asp Lys Asn Phe Leu Gly Arg Pro Val
 1325 1330 1335

Glu Val Leu Leu Asn Asp Asp Leu Ile Val Ser Thr Gly Phe Gly
 1340 1345 1350

Ser Gly Leu Ala Thr Val His Val Thr Thr Val Val His Lys Thr
 1355 1360 1365

Ser Thr Ser Glu Glu Val Cys Ser Phe Tyr Leu Lys Ile Asp Thr
 1370 1375 1380

55 Gln Asp Ile Glu Ala Ser His Tyr Arg Gly Tyr Gly Asn Ser Asp

EP 2 578 683 B9

1385

1390

1395

5 Tyr Lys Arg Ile Val Ala Cys Ala Ser Tyr Lys Pro Ser Arg Glu
 1400 1405 1410

Glu Ser Ser Ser Gly Ser Ser His Ala Val Met Asp Ile Ser Leu
1415 1420 1425

10

Pro Thr Gly Ile Ser Ala Asn Glu Glu Asp Leu Lys Ala Leu Val
 1430 1435 1440

15 Glu Gly Val Asp Gln Leu Phe Thr Asp Tyr Gln Ile Lys Asp Gly
1445 1450 1455

His	Val	Ile	Leu	Gln	Leu	Asn	Ser	Ile	Pro	Ser	Ser	Asp	Phe	Leu
	1460				1465							1470		

Cys Val Arg Phe Arg Ile Phe Glu Leu Phe Glu Val Gly Phe Leu
 1475 1480 1485

25

Ser Pro Ala Thr Phe Thr Val Tyr Glu Tyr His Arg Pro Asp Lys
 1490 1495 1500

30 Gln Cys Thr Met Phe Tyr Ser Thr Ser Asn Ile Lys Ile Gln Lys
 1505 1510 1515

Val Cys Glu Gly Ala Ala Cys Lys Cys Val Glu Ala Asp Cys Gly
 1520 . . . 1525 . . . 1530 . . .

35

Gln Met Gln Glu Glu Leu Asp Leu Thr Ile Ser Ala Glu Thr Arg
 1535 1540 1545

40

Lys Gln Thr Ala Cys Lys Pro Glu Ile Ala Tyr Ala Tyr Lys Val
 1550 1555 1560

Ser Ile Thr Ser Ile Thr Val Glu Asn Val Phe Val Lys Tyr Lys
 1565 1570 1575

50

Ala Thr Leu Leu Asp Ile Tyr Lys Thr Gly Glu Ala Val Ala Glu
1580 1585 1590

50

Lys Asp Ser Glu Ile Thr Phe Ile Lys Lys Val Thr Cys Thr Asn
 1595 1600 1605

66

Ala Glu Leu Val Lys Gly Arg Gln Tyr Leu Ile Met Gly Lys Glu

Ala Leu Gln Ile Lys Tyr Asn Phe Ser Phe Arg Tyr Ile Tyr Pro
1625 1630 1635

5 Leu Asp Ser Leu Thr Trp Ile Glu Tyr Trp Pro Arg Asp Thr Thr
1640 1645 1650

10 Cys Ser Ser Cys Gln Ala Phe Leu Ala Asn Leu Asp Glu Phe Ala
1655 1660 1665

15 Glu Asp Ile Phe Leu Asn Gly Cys
1670 1675

Claims

1. An aptamer/PEG conjugate or a pharmaceutically acceptable salt thereof having an aptamer sequence of:

20 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfU
fUAfCfCfUmGfCmG-3T (SEQ ID NO: 4),

25 wherein fC and fU = 2'-fluoro nucleotides, and mG and mA= 2'-OMe nucleotides and all other nucleotides are 2'-OH and 3T indicates an inverted deoxy thymidine,
wherein the 5' end of the aptamer sequence is conjugated to a 40 kDa linear polyethylene glycol (PEG) moiety via a linker.

30 2. The aptamer/PEG conjugate or salt of claim 1, wherein the 5' end of the aptamer sequence has a 5' linker as follows: H₂N~~~~~5'Aptamer3', wherein ~~~~~ denotes the linker, and wherein the amine-modified aptamer is conjugated to the 40 kDa linear polyethylene glycol (PEG) moiety via the linker.

35 3. The aptamer/PEG conjugate or salt of claim 1 or 2, wherein the linker is an alkyl linker that comprises 6 consecutive CH₂ groups, an alkyl linker comprising between 2 to 18 consecutive CH₂ groups, an alkyl linker comprising between 2 to 12 consecutive CH₂ groups, or an alkyl linker comprising between 3 to 6 consecutive CH₂ groups.

40 4. A composition comprising a therapeutically effective amount of the aptamer/PEG conjugate of any one of claims 1 to 3 or a salt thereof and a pharmaceutically acceptable carrier or diluent.

45 5. Use of the aptamer/PEG conjugate or salt of any one of claims 1 to 3 in the manufacture of a medicament for treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal.

50 6. The aptamer/PEG conjugate or salt of any one of claims 1 to 3 for use in a method of treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal.

7. The use of claim 5 or the aptamer/PEG conjugate or salt for use in a method of treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal of claim 6, wherein the mammal is a human.

55 8. The use of claim 5 or the aptamer/PEG conjugate or salt for use in a method of treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal of claim 6, wherein the disease is an acute ischemic disease, an acute inflammatory disease, or a chronic inflammatory or immune-mediated disease.

9. The use of claim 5 or the aptamer/PEG conjugate or salt for use in a method of treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal of claim 6, wherein the disease is myocardial infarction, stroke, ischemia/reperfusion injury, infectious disease, septicemia, shock, acute transplant

rejection, hyperacute transplant rejection, allergy, asthma, rheumatoid arthritis or another rheumatological disease, multiple sclerosis or another neurologic disease, psoriasis or another dermatological disease, myasthenia gravis, systemic lupus erythematosus (SLE), subacute transplant rejection, chronic transplant rejection or glomerulonephritis or another renal disease.

5 **10.** The use of claim 5 or the aptamer/PEG conjugate or salt for use in a method of treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal of claim 6, wherein the disease is myocardial injury relating to coronary artery bypass surgery (CABG) surgery, myocardial injury relating to balloon angioplasty or myocardial injury relating to restenosis.

10 **11.** The use of claim 5 or the aptamer/PEG conjugate or salt for use in a method of treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal of claim 6, wherein the disease to be treated is a C5, C5a and/or C5b-9 complement protein mediated complication relating to CABG surgery.

15 **12.** The use of claim 5 or the aptamer/PEG conjugate or salt for use in a method of treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal of claim 6, wherein the aptamer/PEG conjugate is administered peri-operatively.

20 **13.** The use of claim 5 or the aptamer/PEG conjugate or salt for use in a method of treating, preventing or ameliorating a disease mediated by C5, C5a and/or C5b-9 complement protein in a mammal of claim 6, wherein the aptamer/PEG conjugate is administered intravenously.

25 **14.** An *in vitro* diagnostic method comprising contacting the aptamer/PEG conjugate or salt of any one of claims 1 to 3 with a composition suspected of comprising C5 complement protein or a variant thereof and detecting the presence or absence of C5 complement protein or a variant thereof.

30 **15.** The diagnostic method of claim 14, wherein the C5 complement protein or variant thereof is a human C5 complement protein or variant thereof.

35 **16.** An aptamer/PEG conjugate or salt according to any one of claims 1 to 3, for use as an *in vitro* diagnostic.

40 **17.** An aptamer/PEG conjugate or salt according to any one of claims 1 to 3, for use as an *in vivo* diagnostic.

45 **18.** An aptamer/PEG conjugate or salt according to any one of claims 1 to 3, for use in the treatment, prevention or amelioration of disease *in vivo*.

Patentansprüche

40 **1.** Aptamer/PEG-Konjugat oder ein pharmazeutisch annehmbares Salz davon mit der folgenden Aptamersequenz:

45 fCmGfCfCGfCmGmGfUfCfUfCmAmGmGfCGfCfUmGmAmGfUfCfUmGmAmGfUfU
 fUAfCfCfUmGfCmG-3T (Seq.-ID Nr. 4),

50 worin fC und fU = 2'-Fluor-Nucleotide sind und mG und mA = 2'-OMe-Nucleotide sind und alle anderen Nucleotide 2'-OH sind und 3T ein invertiertes Desoxythymidin bezeichnet,
 worin das 5'-Ende der Aptamersequenz über einen Linker an eine unverzweigte 40-kDa-Polyethylenglykol- (PEG) Gruppierung konjugiert ist.

55 **2.** Aptamer/PEG-Konjugat oder Salz nach Anspruch 1, worin das 5'-Ende der Aptamersequenz einen 5'-Linker wie folgt aufweist: H₂N~~~~~5'Aptamer3', worin ---- den Linker bezeichnet und worin das Amin-modifizierte Aptamer über den Linker an die unverzweigte 40-kDa-Polyethylenglykol- (PEG) Gruppierung konjugiert ist.

60 **3.** Aptamer/PEG-Konjugat oder Salz nach Anspruch 1 oder 2, worin der Linker ein Alkyl-Linker, der 6 aufeinanderfolgende CH₂-Gruppen umfasst, ein Alkyl-Linker, der zwischen 2 bis 18 aufeinanderfolgende CH₂-Gruppen umfasst,

ein Alkyl-Linker, der zwischen 2 bis 12 aufeinanderfolgende CH₂-Gruppen umfasst, oder ein Alkyl-Linker ist, der zwischen 3 bis 6 aufeinanderfolgende CH₂-Gruppen umfasst.

4. Zusammensetzung, die eine therapeutisch wirksame Menge des Aptamer/PEG-Konjugats nach einem der Ansprüche 1 bis 3 oder ein Salz davon und einen pharmazeutisch annehmbaren Träger oder Verdünner umfasst.
5. Verwendung des Aptamer/PEG-Konjugats oder eines Salzes nach einem der Ansprüche 1 bis 3 zur Herstellung eines Medikaments zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird.
10. 6. Aptamer/PEG-Konjugat oder Salz nach einem der Ansprüche 1 bis 3 zur Verwendung in einem Verfahren zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird.
15. 7. Verwendung nach Anspruch 5 oder Aptamer/PEG-Konjugat oder Salz zur Verwendung in einem Verfahren zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird, nach Anspruch 6, worin das Säugetier ein Mensch ist.
20. 8. Verwendung nach Anspruch 5 oder Aptamer/PEG-Konjugat oder Salz zur Verwendung in einem Verfahren zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird, nach Anspruch 6, worin die Erkrankung eine akute ischämische Erkrankung, eine akute Entzündungskrankheit oder eine chronische Entzündungs- oder Immunkrankheit ist.
25. 9. Verwendung nach Anspruch 5 oder Aptamer/PEG-Konjugat oder Salz zur Verwendung in einem Verfahren zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird, nach Anspruch 6, worin die Erkrankung ein Myokardinfarkt, ein Schlaganfall, eine ischämische/Reperfusions-Verletzung, eine Infektionskrankheit, Sepsis, Schock, eine akute Transplantatabstoßung, eine hyperakute Transplantatabstoßung, eine Allergie, Asthma, rheumatoide Arthritis oder eine andere rheumatologische Erkrankung, multiple Sklerose oder eine andere neurologische Erkrankung, Psoriasis oder eine andere dermatologische Erkrankung, Myasthenia gravis, systemischer Lupus erythematoses (SLE), eine subakute Transplantatabstoßung, chronische Transplantatabstoßung oder Glomerulonephritis oder eine andere Nierenerkrankung ist.
30. 10. Verwendung nach Anspruch 5 oder Aptamer/PEG-Konjugat oder Salz zur Verwendung in einem Verfahren zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird, nach Anspruch 6, worin die Erkrankung eine Myokardverletzung in Verbindung mit einer koronaren Arterienbypassoperation (CABG), eine Myokardverletzung in Verbindung mit einer Ballon-Angioplastie oder eine Myokardverletzung in Verbindung mit Restenose ist.
35. 11. Verwendung nach Anspruch 5 oder Aptamer/PEG-Konjugat oder Salz zur Verwendung in einem Verfahren zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird, nach Anspruch 6, worin die zu behandelnde Erkrankung eine durch C5-, C5a- und/oder C5b-9-Komplementprotein vermittelte Komplikation in Verbindung mit einer CABG-Operation ist.
40. 12. Verwendung nach Anspruch 5 oder Aptamer/PEG-Konjugat oder Salz zur Verwendung in einem Verfahren zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird, nach Anspruch 6, worin das Aptamer/PEG-Konjugat perioperativ verabreicht wird.
45. 13. Verwendung nach Anspruch 5 oder Aptamer/PEG-Konjugat oder Salz zur Verwendung in einem Verfahren zur Behandlung, Prävention oder Linderung einer Erkrankung, die durch C5-, C5a- und/oder C5b-9-Komplementprotein in einem Säugetier vermittelt wird, nach Anspruch 6, worin das Aptamer/PEG-Konjugat intravenös verabreicht wird.
50. 14. In-vitro-Diagnoseverfahren, das das Kontaktieren des Aptamer/PEG-Konjugats oder Salzes nach einem der Ansprüche 1 bis 3 mit einer Zusammensetzung, von der vermutet wird, dass sie C5-Komplementprotein oder eine Variante davon enthält, und das Detektieren der Gegenwart oder Abwesenheit von C5-Komplementprotein oder einer Variante davon umfasst.
55. 15. Diagnoseverfahren nach Anspruch 14, worin das C5-Komplementprotein oder die Variante davon ein menschliches

C5-Komplementprotein oder eine Variante davon ist.

16. Aptamer/PEG-Konjugat oder Salz nach einem der Ansprüche 1 bis 3 zur Verwendung als ein In-vitro-Diagnostikum.
- 5 17. Aptamer/PEG-Konjugat oder Salz nach einem der Ansprüche 1 bis 3 zur Verwendung als ein In-vivo-Diagnostikum.
18. Aptamer/PEG-Konjugat oder Salz nach einem der Ansprüche 1 bis 3 zur Verwendung bei der Behandlung, Prävention oder Linderung einer Erkrankung in vivo.

10

Revendications

1. Conjugué d'aptamère/PEG ou un sel acceptable du point de vue pharmaceutique de celui-ci ayant une séquence d'aptamères de :

15

fCmGfCfCGfCmGmGfUFfCfUFfCmAmGmGfCGfCfUmGmAmGfUFfCfUmGmAmGfUFU
fUAfCfCfUmGfCmG-3T (SEQ ID NO 4),

20

où fC et fU = 2'-fluoro nucleotides, et mG et mA = 2'-OMe nucléotides et tous les autres nucléotides sont 2'-OH, et 3T indique un déoxy thymidine inversé, où la fin 5' de la séquence d'aptamères est conjuguée à un fragment de polyéthylène glycol linéaire 40 kDa (PEG) via un lieu.

25

2. Conjugué d'aptamère/PEG ou un sel selon la revendication 1, où la fin 5' de la séquence d'aptamères présente un lieu 5' comme suit : H₂N~~~~~5'aptamère3', où ----- indique le lieu, et où l'aptamère modifié par une amine est conjugué au fragment de polyéthylène glycol linéaire 40 kDa (PEG) par le lieu.
- 30 3. Conjugué d'aptamère/PEG ou un sel selon la revendication 1 ou 2, où le lieu est un alkyle lieu qui comprend 6 groupes CH₂ consécutifs, un alkyle lieu comprenant entre 2 à 18 groupes CH₂ consécutifs, un alkyle lieu comprenant entre 2 à 12 groupes CH₂ consécutifs, ou bien un alkyle lieu comprenant entre 3 à 6 groupes CH₂ consécutifs.
4. Composition comprenant une quantité efficace du point de vue thérapeutique d'un conjugué d'aptamère/PEG selon l'une quelconque des revendications 1 à 3 ou un sel de celui-ci et un support ou diluent acceptable du point de vue pharmaceutique.
- 35 5. Utilisation du conjugué d'aptamère/PEG ou sel selon l'une quelconque des revendications 1 à 3 dans la fabrication d'un médicament pour le traitement, la prévention ou l'amélioration d'un trouble régulé par une protéine de complément C5, C5a et/ou C5b-9 chez un mammifère.
- 40 6. Conjugué d'aptamère/PEG ou un sel selon l'une quelconque des revendications 1 à 3 pour utilisation dans une méthode de traitement, de prévention ou d'amélioration d'un trouble régulé par une protéine de complément C5, C5a et/ou C5b-9 chez un mammifère.
- 45 7. Utilisation selon la revendication 5 ou le conjugué d'aptamère/PEG ou sel pour utilisation dans une méthode de traitement, de prévention ou d'amélioration d'un trouble régulé par une protéine de complément C5, C5a et/ou C5b-9 chez un mammifère selon la revendication 6, où le mammifère est un être humain.
- 50 8. Utilisation selon la revendication 5 ou le conjugué d'aptamère/PEG ou sel pour utilisation dans une méthode de traitement, de prévention ou d'amélioration d'un trouble régulé par une protéine de complément C5, C5a et/ou C5b-9 chez un mammifère de la revendication 6, où le trouble est un trouble ischémique aigu, un trouble inflammatoire aigu ou un trouble chronique inflammatoire ou d'origine immunitaire.
- 55 9. Utilisation selon la revendication 5 ou le conjugué d'aptamère/PEG ou sel pour l'utilisation dans une méthode de traitement, de prévention ou d'amélioration d'un trouble régulé par une protéine de complément C5, C5a et/ou C5b-9 chez un mammifère de la revendication 6, où le trouble est l'infarctus du myocarde, un accident cérébrovasculaire, une lésion d'ischémie/reperfusion, une maladie infectieuse, une septicémie, un choc, un rejet de transplantation

aigu, un rejet de transplantation hyper-aigu, une allergie, l'asthme, l'arthrite rhumatoïde ou un autre trouble rhumatologique, la sclérose multiple ou un autre trouble neurologique, le psoriasis ou une autre maladie dermatologique, la myasthénie grave, la maladie systémique lupus érythémateux (SLE), le rejet de transplantation subaigu, le rejet de transplantation chronique ou la gloméronéphrite ou une autre maladie rénale.

5

10. Utilisation selon la revendication 5 ou le conjugué d'aptamère/PEG ou sel pour utilisation dans une méthode de traitement, de prévention ou d'amélioration d'un trouble régulé par la protéine de complément C5, C5a et/ou C5b-9 chez un mammifère de la revendication 6, où le trouble est une lésion du myocarde liée à une chirurgie de pontage aorto-coronarien (CABG), la lésion du myocarde liée à l'angioplastie à ballonnet ou bien la lésion du myocarde liée à une resténose.

15

11. Utilisation selon la revendication 5 ou le conjugué d'aptamère/PEG ou sel pour utilisation dans une méthode de traitement, de prévention ou d'amélioration d'un trouble régulé par la protéine de complément C5, C5a et/ou C5b-9 chez un mammifère de la revendication 6, où le trouble à traiter est une complication ayant pour origine la protéine de complément C5, C5a et/ou C5b-9 liée à une chirurgie CABG.

20

12. Utilisation selon la revendication 5 ou le conjugué d'aptamère/PEG ou sel pour utilisation dans une méthode de traitement, de prévention ou d'amélioration d'un trouble régulé par la protéine de complément C5, C5a et/ou C5b-9 chez un mammifère de la revendication 6, où le conjugué d'aptamère/PEG est administré de manière péri-opératoire.

25

13. Utilisation selon la revendication 5 ou le conjugué d'aptamère/PEG ou sel pour utilisation dans une méthode de traitement, de prévention ou d'amélioration d'un trouble régulé par la protéine de complément C5, C5a et/ou C5b-9 chez un mammifère de la revendication 6, où le conjugué d'aptamère/PEG est administré de manière intraveineuse.

30

14. Méthode de diagnostic *in vitro* comprenant la mise en contact d'un conjugué d'aptamère/PEG ou sel selon l'une quelconque des revendications 1 à 3 avec une composition dont on suspecte qu'elle contient une protéine de complément C5 ou une variante de celle-ci, et la détection de la présence ou de l'absence d'une protéine de complément C5 ou d'une variante de celle-ci.

35

15. Méthode de diagnostic selon la revendication 14, dans laquelle la protéine de complément C5 ou la variante de celle-ci est une protéine de complément C5 humaine ou une variante de celle-ci.

16. Conjugué d'aptamère/PEG ou sel selon l'une quelconque des revendications 1 à 3, pour utilisation comme un diagnostic *in vitro*.

40

17. Conjugué d'aptamère/PEG ou sel selon l'une quelconque des revendications 1 à 3, pour utilisation comme un diagnostic *in vivo*.

18. Conjugué d'aptamère/PEG ou sel selon l'une quelconque des revendications 1 à 3, pour utilisation dans le traitement, la prévention ou l'amélioration d'un trouble *in vivo*.

45

50

55




Fig. 1




Fig. 2




Fig. 3A




Fig. 3B




Fig. 3C




Fig. 4





Fig. 5






Fig. 7A

ARC#	PEG MODIFICATION	IC50 (nM)
186	NO PEG	0.5 ± 0.1
657	ARC186+20 kDa	0.6 ± 0.1
658	ARC186+30 kDa	0.5 ± 0.1
187	40 kDa BRANCHED 1,3-BIS (mPEG-[20 kDa])-PROPYL-2-(4'-BUTAMIDE)	0.6 ± 0.2

Fig. 7B




Fig. 7C

ARC#	PEG MODIFICATION	IC50 (nM)
187	40K BRANCHED 1,3-BIS(mPEG-[20 kDa])-PROPYL-2-(4'-BUTAMIDE)	0.38 ± 0.03
1537	40K LINEAR	0.39 ± 0.06
1730	2 x 20K LINEAR	0.48 ± 0.04
1905	40K BRANCHED 2,3-BIS(mPEG-[20 kDa])-PROPYL-1-CARBAMOYL	0.40 ± 0.03

Fig. 7D




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21




Fig. 22

GROUP	# MICE	TEST ARTICLE	DOSE		MEASUREMENTS
			CONC (nM)	PERFUSION RATE mL/MIN	
1	5	NO APTAMER	0	3	HEART RATE, RHYTHM, LEFT VENTRICULAR PRESSURE
3	3	IRRELEVANT APTAMER AT 50X MOLAR RATIO	1500	3	HEART RATE, RHYTHM, LEFT VENTRICULAR PRESSURE
4	3	ARC186 AT MOLAR EQUIVALENCE	30	3	HEART RATE, RHYTHM, LEFT VENTRICULAR PRESSURE
5	5	ARC186 AT 10X MOLAR RATIO	300	3	HEART RATE, RHYTHM, LEFT VENTRICULAR PRESSURE
6	4	ARC186 AT 50X MOLAR RATIO	1500	3	HEART RATE, RHYTHM, LEFT VENTRICULAR PRESSURE

Fig. 23




Fig. 24A




Fig. 24B




Fig. 25




Fig. 26




Fig. 27




Fig. 28




Fig. 29




Fig. 30

ANIMAL	TREATMENT	IRRELEVANT	C3b	C5b-9
1-001 (1/26)	NONE	NEG	POS	POS
1-002 (1-26)	NONE	NEG	POS	POS
1-001	NONE	NEG	POS	POS
1-002	NONE	NEG	POS	POS
1-003	NONE	NEG	POS	POS
3-001	IRRELEVANT APTAMER 50X MOLAR	NEG	POS	POS
3-002	IRRELEVANT APTAMER 50X MOLAR	NEG	POS	POS
3-003	IRRELEVANT APTAMER 50X MOLAR	NEG	POS	POS
4-001	C5 APTAMER MOLAR EQUIVALENCE	NEG	POS	POS
4-002	C5 APTAMER MOLAR EQUIVALENCE	NEG	POS	POS
4-003	C5 APTAMER MOLAR EQUIVALENCE	NEG	POS	POS
5-001	C5 APTAMER 10X MOLAR EQUIVALENCE	NEG	POS	NEG
5-002	C5 APTAMER 10X MOLAR EQUIVALENCE	NEG	POS	NEG
5-003	C5 APTAMER 10X MOLAR EQUIVALENCE	NEG	POS	NEG
5-004	C5 APTAMER 10X MOLAR EQUIVALENCE	NEG	POS	NEG
5-005	C5 APTAMER 10X MOLAR EQUIVALENCE	NEG	POS	NEG
6-001	C5 APTAMER 50X MOLAR EQUIVALENCE	NEG	POS	NEG
6-002	C5 APTAMER 50X MOLAR EQUIVALENCE	NEG	POS	NEG
6-003	C5 APTAMER 50X MOLAR EQUIVALENCE	NEG	POS	NEG
6-004	C5 APTAMER 50X MOLAR EQUIVALENCE	NEG	POS	NEG

Fig. 31

ANIMAL	COMPOUND	SERUM	MOLAR RATIO	INCREASED EDP	FAILURE (Y OR NO)	TIME TO FAILURE
1-001	ARC658	HUMAN	0	Y	Y	4 MIN 30 SEC
1-002	ARC658	HUMAN	0	Y	Y	4 MIN 38 SEC
1-003	ARC658	HUMAN	0	Y	Y	3 MIN 20 SEC
1-004	ARC658	HUMAN	0	Y	Y	12 MIN 24 SEC
1-005	ARC658	HUMAN	0	Y	N	NA
1-006	ARC658	HUMAN	0	Y	Y	8 MIN 30 SEC
1a-001	ARC658	HUMAN	0.5	Y	Y	7 MIN 22 SEC
1a-002	ARC658	HUMAN	0.5	Y	Y	7 MIN 30 SEC
1a-003	ARC658	HUMAN	0.5	Y	Y	8 MIN 13 SEC
2-001	ARC658	HUMAN	1	N	N	NA
2-002	ARC658	HUMAN	1	N	N	NA
2-003	ARC658	HUMAN	1	N	N	NA
2-004	ARC658	HUMAN	1	N	N	NA
3-001	ARC658	HUMAN	3	N	N	NA
3-002	ARC658	HUMAN	3	N	N	NA
3-003	ARC658	HUMAN	3	N	N	NA
6-001	ARC658	PRIMATE	0	Y	Y	2 MIN 30 SEC
6-002	ARC658	PRIMATE	0	Y	Y	2 MIN 47 SEC
6-003	ARC658	PRIMATE	0	Y	Y	3 MIN 30 SEC
6-004	ARC658	PRIMATE	0	Y	Y	1 MIN 28 SEC
6a-001	ARC658	PRIMATE	3	Y	Y	5 MIN 2 SEC
6a-002	ARC658	PRIMATE	3	Y	Y	4 MIN 30 SEC
6a-003	ARC658	PRIMATE	3	Y	Y	3 MIN 43 SEC
6a-004	ARC658	PRIMATE	3	Y	Y	8 MIN 0 SEC
7-001	ARC658	PRIMATE	10	Y	N	NA
7-002	ARC658	PRIMATE	10	Y	N	NA
7-003	ARC658	PRIMATE	10	Y	Y	9 MIN 15 SEC
7-004	ARC658	PRIMATE	10	Y	N	NA
9-001	ARC658	PRIMATE	20	Y	Y	12 MIN 50 SEC
11-001	ARC658	PRIMATE	30	Y	N	NA
11-002	ARC658	PRIMATE	30	Y	Y	10 MIN 0 SEC
11-003	ARC658	PRIMATE	30	Y	N	NA
13-001	ARC658	PRIMATE	50	N	N	NA
13-002	ARC658	PRIMATE	50	N	N	NA
13-003	ARC658	PRIMATE	50	N	N	NA

Fig. 32





Fig. 33

GROUP NUMBER	NUMBER OF ANIMALS	TEST ARTICLE	DOSAGE LEVEL (mg/kg)	DOSAGE VOLUME (mL/kg)	DOSING REGIMEN	SAMPLE COLLECTION
1	3	ARC657 20 kDa PEG	10	1	INTRAVENOUS BOLUS ON DAY 1	t=PRE-DOSE, 0.5, 1, 2, 4, 8, 16, 24, 32, 48 HRS
2	3	ARC658 30 kDa PEG	10	1	INTRAVENOUS BOLUS ON DAY 1	t=PRE-DOSE, 0.5, 1, 2, 4, 8, 16, 24, 32, 48 HRS
3	3	ARC187 40 kDa PEG	10	1	INTRAVENOUS BOLUS ON DAY 1	t=PRE-DOSE, 0.5, 1, 2, 4, 8, 16, 24, 32, 48 HRS

Fig. 34

PK OF ANTI-C5 APTAMER-PEG CONJUGATES IN SPRAGUE-DAWLEY RAT						
APTAMER	ARC657		ARC658		ARC187	
PEG GROUP	20 kDa		30 kDa		40 kDa	
TIME (HRS)	MEAN	STDEV	MEAN	STDEV	MEAN	STDEV
	Cp (µM)					
0	0.00	0.00	0.00	0.00	0.00	0.00
0.5	12.77	2.67	2.97	0.62	6.52	1.77
1	13.38	2.08	4.09	1.44	7.94	1.08
2	6.63	1.87	3.40	0.66	6.57	1.56
4	2.38	0.87	2.22	0.39	4.56	0.54
8	0.72	0.18	2.87	1.84	1.53	0.57
12	0.31	0.03	0.60	0.09	1.45	0.17
24	<LLOQ	<LLOQ	0.47	0.17	0.73	0.03
32	<LLOQ	<LLOQ	<LLOQ	<LLOQ	0.49	0.01
48	<LLOQ	<LLOQ	<LLOQ	<LLOQ	0.27	0.02

Fig. 35

**Fig. 36**

SPECIES	DOSE (mg/kg)	ADMIN ROUTE	PEG GROUP	C_{max} ($\mu\text{g}/\text{mL}$)	$AUC_{0-\infty}$ ($\mu\text{g}\cdot\text{hr}/\text{mL}$)	$t_{1/2}(\alpha)$ (HR)	$t_{1/2}(\beta)$ (HR)	Cl (mL/min/kg)	V_{ss} (mL/kg)
RAT	10	IV	20 kD	169.97	458	-	2.05	0.36	65
RAT	10	IV	30 kD	52.00	497	-	7.40	0.34	220
RAT	10	IV	40 kD	100.85	886	-	15.00	0.19	188

Fig. 37

GROUP NO.	NO. ANIMALS	TEST ARTICLE	DOSE		DOSE ROUTE	SAMPLE COLLECTION
			(mg/kg)	(mL/kg)		
1	33F	ARC187	10	4	IV	t=PRE-DOSE, 0.5, 1, 2, 4, 8, 16, 24, 32, 48, 72 HRS
2	33F	ARC1905	10	4	IV	t=PRE-DOSE, 0.5, 1, 2, 4, 8, 16, 24, 32, 48, 72 HRS

Fig. 38A




Fig. 38B

NONCOMPARTMENTAL PK ANALYSIS (NCA)			
ATTRIBUTE	UNIT	ARC1905	ARC187
C _{max}	μM	12	10
AUC _{0-∞}	μM-HR	88	89
MRT _{0-∞}	HR	11	11
CL	mL/min. kg	0.15	0.15
V _{ss}	mL/kg	95	96
K ₁₀	1/HR	0.151	0.121
t _{1/2} (K ₁₀)	HR	5	6

Fig. 38C

APTAMER	PRE	1 HR	3 HR	6 HR
ARC657	NO	YES	YES	YES
ARC658	NO	YES	YES	YES
ARC187	NO	YES	YES	YES

Fig. 39

GROUP NUMBER	NUMBER OF ANIMALS	TEST ARTICLE	DOSAGE LEVEL (mg/kg)	DOSAGE VOLUME (mL/kg)	DOSING REGIMEN	SAMPLE COLLECTION
1	1	ARC657 20 kDa PEG	30	3	INTRAVENOUS BOLUS ON DAY 1	t=PRE-DOSE, 1, 5, 10, 15, AND 30 MINUTES AND AT 1, 2, 4, 6, 8, 12, 24, 48, 72, 96, AND 192 HRS
2	1	ARC658 30 kDa PEG	30	3	INTRAVENOUS BOLUS ON DAY 1	t=PRE-DOSE, 1, 5, 10, 15, AND 30 MINUTES AND AT 1, 2, 4, 6, 8, 12, 24, 48, 72, 96, AND 192 HRS
3	1	ARC187 40 kDa PEG	30	3	INTRAVENOUS BOLUS ON DAY 1	t=PRE-DOSE, 1, 5, 10, 15, AND 30 MINUTES AND AT 1, 2, 4, 6, 8, 12, 24, 48, 72, 96, AND 192 HRS

Fig. 40

PK OF ANTI-C5 APTAMER -PEG CONJUGATES IN CYNOMOLGUS MACAQUE			
APTAMER	ARC657	ARC658	ARC187
PEG GROUP	20 kDa	30 kDa	40 kDa
	MEAN	MEAN	MEAN
TIME (HRS)	C _p (μM)	C _p (μM)	C _p (μM)
0	0.00	0.00	0.00
0.08	16.41	21.99	18.00
0.25	16.64	22.67	17.26
0.5	19.38	26.16	24.17
1	19.36	27.08	21.79
4	7.73	14.16	21.69
8	2.03	6.02	16.68
12	1.10	3.04	13.11
24	0.36	0.95	7.24
48	0.15	0.42	2.30
72	0.07	0.25	0.96
96	0.03	0.23	0.54
192	0.01	0.04	0.14

Fig. 41

SPECIES	DOSE (mg/kg)	ADMIN ROUTE	PEG GROUP	C _{max} (μg/ml)	AUC _{0-∞} (μg·hr/mL)	t _{1/2} (α) (HR)	t _{1/2} (β) (HR)	Cl (mL/min/kg)	V _{ss} (mL/kg)
PRIMATE	30	IV	20 kD	293	1191	2.35	31.80	0.41	273
PRIMATE	30	IV	30 kD	385	2484	3.33	40.08	0.20	233
PRIMATE	30	IV	40 kD	316	6752	12.38	60.11	0.07	141

Fig. 42




Fig. 43A




Fig. 43B




Fig. 43C




Fig. 43D

GROUP NUMBER	NUMBER OF ANIMALS	TEST ARTICLE	DOSAGE LEVEL (mg/kg)	DOSAGE VOLUME (mL/kg)	DOSING REGIMEN	SAMPLE COLLECTION
1	4	ARC658 30 kDa PEG	30	3	INTRAVENOUS BOLUS ON DAY 1	t=PRE-DOSE, 5, 10, 15, AND 30 MINUTES AND AT 1, 2, 4, 6, 8, 12, 24, 48, 72, 96, 144 AND 192 HRS
2	4	ARC187 40 kDa PEG	30	3	INTRAVENOUS BOLUS ON DAY 1	t=PRE-DOSE, 5, 10, 15, AND 30 MINUTES AND AT 1, 2, 4, 6, 8, 12, 24, 48, 72, 96, 144 AND 192 HRS

Fig. 44




Fig. 45

SPECIES	DOSE (mg/kg)	ADMIN ROUTE	PEG GROUP	C_{max} (μ M)	$AUC_{0-\infty}$ (μ g_hr/mL)	$t_{1/2}(\alpha)$ (HR)	$t_{1/2}(\beta)$ (HR)	Cl (mL/min_kg)	V_{ss} (mL/kg)
PRIMATE	30	IV	30 kD	519	3059	2.35	30.45	0.16	213
PRIMATE	30	IV	40 kD	609	8061	4.48	53.13	0.06	168

Fig. 46




Fig. 47




Fig. 48

GROUP NUMBER	NUMBER OF ANIMALS	TEST ARTICLE	DOSAGE LEVEL (mg/kg)	DOSE CONC. (mg/mL)	DOSE VOLUME (mL/kg)	DOSING REGIMEN	LAST DAY ON TEST
1	3	ARC187	1 mg/kg IV BOLUS AND 0.0013 mg/kg/min IV INFUSION	10	0.1 mL/kg IV BOLUS AND 0.1872 mL/kg/day, IV INFUSION	IV BOLUS AND 48 HOUR INFUSION ON DAY 1	DAY 8

Fig. 49

SPECIES	DOSE (mg/kg)	ADMIN ROUTE	PEG GROUP	C _{max} (μg/ml)	AUC _{0-∞} (μg·hr/mL)	t _{1/2} (α) (HR)	t _{1/2} (β) (HR)	Cl (mL/min·kg)	V _{ss} (mL/kg)
PRIMATE	30	IV	40 kD	610	8061	4.48	53.13	0.06	168

Fig. 50




Fig. 51




Fig. 52

APTAMER	PEG GROUP	MOL. WT. (OLIGO)	MOL. WT. (TOTAL)	$t_{1/2}(\alpha)$ (HR)	$t_{1/2}(\beta)$ (HR)	TOTAL (OLIGO) DOSE FOR 1.5 μ M (g)
ARC187	40 kDa	12,703	52,703	4.48	53.13	0.40

Fig. 53




Fig. 54

TREATMENT	ACT (SECONDS)		
	DONOR 1 ^a	DONOR 2 ^b	DONOR 3 ^b
BASELINE	133 \pm 30	145 \pm 12	138 \pm 1
+ HEPARIN	842 \pm 95	519 \pm 16	450 \pm 15
+ HEPARIN & ARC187	nd	618 \pm 55	492 \pm 28
+ HEPARIN & PROTAMINE	160 \pm 11	155 \pm 6	159 \pm 9
+ HEPARIN, PROTAMINE & ARC187	200 \pm 5	172 \pm 1	169 \pm 8

^a5 U/mL HEPARIN; ^b4 U/mL HEPARIN; nd, NOT DETERMINED

Fig. 55




Fig. 56




Fig. 57

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9941271 A1 [0013]
- US 5475096 A [0052]
- US 5270163 A [0052]
- WO 9119813 A [0052]
- US 5958691 A [0055]
- US 5660985 A [0055] [0066]
- US 5698687 A [0055]
- US 5817635 A [0055]
- US 5672695 A [0055]
- WO 9207065 A [0055]
- US 5707796 A [0063]
- US 5763177 A [0063]
- US 5567588 A [0063]
- US 5861254 A [0063]
- US 5496938 A [0063]
- US 5705337 A [0063]
- US 5580737 A [0064] [0066]
- US 5756703 A [0066]
- US 5637459 A [0071]
- US 5683867 A [0071]
- US 6011020 A [0071] [0158]
- US 6051698 A [0071]
- WO 9818480 A [0071]
- US 5648214 A [0072]
- US 20040197804 A [0076]
- US 20050037394 A [0076]
- US 6395888 B [0129]
- US 3710795 A [0155]
- US 5262564 A [0158]
- US 60550790 B [0167]
- US 10718833 B [0175]
- US 5674685 A [0188]
- US 5668264 A [0188]
- US 6207816 A [0188]
- US 6229002 A [0188]
- US 2005004606 A [0282]
- US 60608048 B [0282]
- US 60581685 B [0282]
- US 60547747 B [0282]
- US 20040225 B [0282]
- US 60544542 B [0282]

Non-patent literature cited in the description

- **TUCKER et al.** *J. Chromatography B*, 1999, vol. 732, 203-212 [0004]
- **FITCH et al.** *Circ.*, 1999, vol. 100, 2499-506 [0011]
- **WATSON et al.** *Antisense & Nucleic Acid Drug Development*, 2000, vol. 10, 63-75 [0014]
- **BELL et al.** *In Vitro Cellular & Developmental Biology - Animal*. *In Vitro Cellular & Developmental Biology - Animal*, 1999, vol. 35, 533-542 [0015]
- **VOROBJEB et al.** *Antisense & Nucleic Acid Drug Development*, 2000, vol. 10, 63-75 [0016]
- **FROEHLER et al.** *Nucl. Acid Res.*, 1986, vol. 14, 5399-5467 [0055] [0134]
- **FROEHLER et al.** *Tet. Lett.*, 1986, vol. 27, 5575-5578 [0055] [0134]
- **SOOD et al.** *Nucl. Acid Res.*, 1977, vol. 4, 2557 [0055] [0134]
- **HIROSE et al.** *Tet. Lett.*, 1978, vol. 28, 2449 [0055] [0134]
- **SPROAT et al.** *Nucl. Acid Res.*, 1991, vol. 19, 733-738 [0069]
- **COTTEN et al.** *Nucl. Acid Res.*, 1991, vol. 19, 2629-2635 [0069]
- **HOBBS et al.** *Biochemistry*, 1973, vol. 12, 5138-5145 [0069]
- **GREEN et al.** *Current Biology*, 1995, vol. 2, 683-695 [0077]
- **PADILLA, R. ; SOUSA, R.** *Nucleic Acids Res.*, 2002, vol. 30 (24), 138 [0079]
- **PADILLA et al.** *Nucleic Acids Research*, 2002, vol. 30, 138 [0079]
- **MATIS ; ROLLINS.** *Nature Medicine*, 1995, vol. 1 (8), 839-842 [0116]
- **WARD.** *Am. J. Pathol.*, 1996, vol. 149, 1079 [0117]
- **WARD.** *J. Lab. Clin. Med.*, 1997, vol. 129, 400 [0118]
- **MULLIGAN et al.** *J. Clin. Invest.*, 1998, vol. 98, 503 [0118] [0121]
- **SCHMID et al.** *Shock*, 1997, vol. 8, 119 [0118]
- **BIESECKER ; GOMEZ.** *J. Immunol.*, 1989, vol. 142, 2654 [0118]
- **BALDWIN et al.** *Transplantation*, 1995, vol. 59, 797 [0118]
- **BRAUER et al.** *Transplantation*, 1995, vol. 59, 288 [0118]
- **TAKAHASHI et al.** *Immunol. Res.*, 1997, vol. 16, 273 [0118]
- **BIESECKER.** *J. Exp. Med.*, 1981, vol. 39, 1779 [0118]
- **NANGAKU.** *Kidney Int.*, 1997, vol. 52, 1570 [0118]

- **LUCCHESI.** *Annu. Rev. Pharmacol. Toxicol.*, 1994, vol. 34, 17 [0118]
- **BEDNAR et al.** *J. Neurosurg.*, 1997, vol. 86, 139 [0118]
- **SUN et al.** *Nucleic Acids Res.*, 1995, vol. 23, 2909 [0118]
- **SPYCHER ; NYDEGGER.** *Infusionsther. Transfusionsmed.*, 1995, vol. 22, 36 [0118]
- **WANG et al.** *Immunology*, 1996, vol. 93, 8563 [0118]
- **ZHANG et al.** *Diabetes*, 2002, vol. 51, 3499 [0119]
- **SOHN et al.** *IOVS*, 2000, vol. 41, 3492 [0119]
- **ACOSTA et al.** *PNAS*, 2002, vol. 97, 5450-5455 [0119]
- **HANSEN, T.K. et al.** *Diabetes*, 2004, vol. 53, 1570-1576 [0119]
- **ALBRECHT, E.A. et al.** *Am J Pathology*, 2004, vol. 164, 849-859 [0119]
- **GERT, V.B. et al.** *Invest Ophthalmol Vis Sci*, 2002, vol. 43, 1104-1108 [0119]
- **BADOUIN, C et al.** *Am J Ophthalmol*, 1988, vol. 105, 383-388 [0119]
- **RINDER et al.** *J. Clin. Invest.*, 1995, vol. 96, 1564 [0121]
- **BIESECKER et al.** *J. Immunol.*, 1989, vol. 142, 2654 [0121]
- **HAVILAND et al.** *J Immunol.*, 01 January 1991, vol. 146 (1), 362-8 [0124]
- **SMITH ; WATERMAN.** *Adv. Appl. Math.*, 1981, vol. 2, 482 [0125]
- **NEEDLEMAN ; WUNSCH.** *J Mol. Biol.*, 1970, vol. 48, 443 [0125]
- **PEARSON ; LIPMAN.** *Proc. Nat'l. Acad. Sci. USA*, 1988, vol. 85, 2444 [0125]
- **ALTSCHUL et al.** *J Mol. Biol.*, 1990, vol. 215, 403-410 [0126]
- **ALTSCHUL et al.** *Nucleic Acids Res.*, 1997, vol. 15, 3389-3402 [0126]
- **MCGINNIS et al.** *Nucleic Acids Res.*, 2004, vol. 32, W20-W25 [0126]
- **VIVES et al.** *J. Biol. Chem.*, 1997, vol. 272 (25), 16010-7 [0171]
- **PIETERSZ et al.** *Vaccine*, 2001, vol. 19 (11-12), 1397-405 [0171]
- **ROTHBARD et al.** *Nat. Med.*, 2000, vol. 6 (11), 1253-7 [0171]
- **ROTHBARD, J et al.** *J. Med. Chem.*, 2002, vol. 45 (17), 3612-8 [0171]
- **GREENWALD et al.** *J. Org. Chem.*, 1995, vol. 60, 331-336 [0178]
- **HARRIS et al.** *Nature*, 2003, vol. 2, 214-221 [0182]
- **GREEN et al.** *Chem. Biol.*, 1995, vol. 2, 683-95 [0192]
- **GONG et al.** *Journal of Clinical Immunology*, 1996, vol. 16, 222-9 [0206]
- **NILSSON et al.** *Blood*, 1998, vol. 92, 1661-7 [0206]
- **EVANS.** *Molecular Immunology*, 1995, vol. 32, 1183-1195 [0228]
- **EVANS.** *Molecular Immunol*, 1995, vol. 32, 1183 [0236]