



(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Claims EN 7

(51) Int Cl.:
C07D 213/79^(2006.01) C07D 213/803^(2006.01)

(86) International application number:
PCT/US2012/022291

(48) Corrigendum issued on:
21.06.2017 Bulletin 2017/25

(87) International publication number:
WO 2012/103047 (02.08.2012 Gazette 2012/31)

(45) Date of publication and mention
of the grant of the patent:
01.03.2017 Bulletin 2017/09

(21) Application number: **12739953.3**

(22) Date of filing: **24.01.2012**

(54) **PROCESS FOR THE PREPARATION OF
4-AMINO-5-FLUORO-3-HALO-6-(SUBSTITUTED)PICOLINATES**

VERFAHREN ZUR HERSTELLUNG VON
4-AMINO-5-FLUOR-3-HALO-6-(SUBSTITUT-)PICOLINATEN

PROCÉDÉ DE PRÉPARATION DE
4-AMINO-5-FLUORO-3-HALOGÉNO-6-(SUBSTITUÉ)PICOLINATES

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **25.01.2011 US 201161435958 P**

(43) Date of publication of application:
04.12.2013 Bulletin 2013/49

(73) Proprietor: **Dow AgroSciences LLC**
Indianapolis, IN 46268 (US)

(72) Inventors:
• **WHITEKER, Gregory, T.**
Carmel
IN 46032 (US)
• **ARNDT, Kim, E.**
Carmel
IN 46032 (US)
• **RENGA, James, M.**
Indianapolis
IN 46220 (US)

- **ZHU, Yuanming**
Carmel
IN 46033 (US)
- **LOWE, Christian, T.**
Westfield
IN 46074 (US)
- **SIDDALL, Thomas, L.**
Zionsville
IN 46077 (US)
- **PODHOREZ, David, E.**
Midland
MI 48640 (US)
- **ROTH, Gary, Alan**
Midland
MI 48640 (US)
- **WEST, Scott, P.**
Midland
MI 48640 (US)

(74) Representative: **f & e patent**
Fleischer, Engels & Partner mbB, Patentanwälte
Braunsberger Feld 29
51429 Bergisch Gladbach (DE)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

(56) References cited:

WO-A1-2006/062979	WO-A1-2011/144891
US-A- 3 285 925	US-A- 3 325 272
US-A- 3 629 424	US-A- 3 803 159
US-A1- 2009 088 322	US-A1- 2010 041 556
US-B1- 6 297 197	US-B2- 6 784 137

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification


Description

[0001] The present invention concerns a process for the preparation of 4-amino-5-fluoro-3-halo-6-(substituted)picol-
 inates. More particularly, the present invention concerns a process for the preparation of 4-amino-5-fluoro-3-halo-6-(sub-
 stituted)picolinate in which the 5-fluoro substituent is introduced by a halogen exchange early in the process scheme.

[0002] U.S. Patent 6,297,197 B1 describes *inter alia* certain 6-(alkoxy or aryloxy)-4-amino-3-chloro-5-fluoropicolinate
 compounds and their use as herbicides. U.S. Patents 6,784,137 B2 and 7,314,849 B2 describe *inter alia* certain 6-(aryl)-
 4-amino-3-chloro-5-fluoropicolinate compounds and their use as herbicides. U.S. Patent 7,432,227 B2 describes *inter*
alia certain 6-(alkyl)-4-amino-3-chloro-5-fluoropicolinate compounds and their use as herbicides. Each of these patents
 describes the manufacture of 4-amino-3-chloro-5-fluoropicolinate starting materials by fluorination of the corresponding
 5-unsubstituted pyridines with 1-(chloromethyl)-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate). It would
 be advantageous to produce 4-amino-5-fluoro-3-halo-6-(substituted)picolinate without having to rely on direct fluorina-
 tion of the 5-position of the pyridine ring with an expensive fluorinating agent like 1-(chloromethyl)-4-fluoro-1,4-diazoni-
 abicyclo[2.2.2]octane bis(tetrafluoroborate).

WO 2011/144891 A1 relates to certain substituted picolinic acid derivatives and their use in controlling plant growth and
 describes *inter alia* 4-amino-3-bromo-6-chloro-5-fluoropyridine-2-carboxylic acid methyl ester as synthetic intermediate.
 US 2009/0088322 A1 relates to herbicidal 6-amino-4-pyrimidinecarboxylic acids having alkyl, alkenyl or alkynyl substit-
 uents in the 5-position and 4-aminopicolinic acids having alkyl, alkenyl or alkynyl substituents in the 3-position and *inter*
alia describes their preparation using amination, coupling in the presence of a transition metal and halogen exchange
 steps. WO 2006/062979 A1 relates to substituted 6-cyclopropyl-4-aminopicolinic acid derivatives useful as herbicides
 and describes *inter alia* certain 3,5-dihalo-substituted derivatives thereof.

[0003] The present invention concerns a process for the preparation of 4-amino-5-fluoro-3-halo-6-(substituted)picol-
 inates from 4,5,6-trichloropicolinate. More particularly, the present invention concerns a process for the preparation of
 a 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of the Formula I



wherein


W represents Cl, Br or I;

R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl or phenyl substituted with from 1 to 4 substituents independently
 selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy; and

R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl; which substituted C₇-C₁₁ arylalkyl
 has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano,
 C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl,
 or where two adjacent substituents are taken together as -O(CH₂)_nO-, wherein n=1 or 2;


which comprises the following steps:

a) fluorinating a 4,5,6-trichloropicolinate of Formula A




wherein R¹ is as previously defined;

with a source of fluoride ion to produce a 4,5,6-trifluoropicolinate of Formula B




wherein R¹ is as previously defined;

b) aminating the 4,5,6-trifluoropicolinate of Formula B with ammonia to produce a 4-amino-5,6-difluoropicolinate of Formula C




wherein R¹ is as previously defined;

c) exchanging the fluoro substituent in the 6-position of the 4-amino-5,6-difluoro-picolinate of Formula C with an iodo, bromo or chloro substituent by treating with an iodide, bromide or chloride source to produce a 4-amino-5-fluoro-6-halopicolinate of Formula D



wherein X represents Cl, Br or I;
and R¹ is as previously defined;

d) halogenating the 4-amino-5-fluoro-6-halopicolinate of Formula D with a halogen source to produce a 4-amino-3,6-dihalo-5-fluoropicolinate of Formula E



wherein W and X independently represent Cl, Br or I;
and R¹ is as previously defined; and

e) coupling the 4-amino-3,6-dihalo-5-fluoropicolinate of Formula E with an aryl, alkyl or alkenyl metal compound of the Formula F


R-Met

F

wherein R is as previously defined and Met represents Zn-halide, Zn-R, tri-(C₁-C₄ alkyl)tin, copper, or B(OR²)(OR³), where R² and R³ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group in the presence of a transition metal catalyst to produce the 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of Formula I.


[0004] The steps a) through e) may be performed in the order listed, as depicted in Scheme I.

Scheme I




[0005] Alternatively, the order in which the steps are performed can be rearranged as illustrated, for example, in Schemes II and III.

Scheme II



[0006] In accordance with Scheme II, the present invention concerns a process for the preparation of a 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of the Formula I



10 wherein


W represents Cl, Br or I;

R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl, or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy; and

15 R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO-, wherein n=1 or 2;


20 which comprises the following steps:

a) fluorinating a 4,5,6-trichloropicolinate of Formula A




30 wherein R¹ is as previously defined;

with a source of fluoride ion to produce a 4,5,6-trifluoropicolinate of Formula B




40 wherein R¹ is as previously defined;

45 b) aminating the 4,5,6-trifluoropicolinate of Formula B with ammonia to produce a 4-amino-5,6-difluoropicolinate of Formula C



55 wherein R¹ is as previously defined;

c) exchanging the fluoro substituent in the 6-position of the 4-amino-5,6-difluoro picolinate of Formula C with an iodo, bromo or chloro substituent by treating with an iodide, bromide or chloride source to produce a 4-amino-5-fluoro-6-halopicolinate of Formula D




wherein X represents Cl, Br or I; and
R¹ is as previously defined;

d) coupling the 4-amino-5-fluoro-6-halopicolinate of Formula D with an aryl, alkyl or alkenyl metal compound of the Formula F




wherein R is as previously defined and Met represents Zn-halide, Zn-R, tri-(C₁-C₄ alkyl)tin, copper, or B(OR²)(OR³), where R² and R³ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group in the presence of a transition metal catalyst to produce the 4-amino-5-fluoro-6-(substituted)-picolinate of Formula G.




wherein R and R¹ are as previously defined; and

e) halogenating the 4-amino-5-fluoro-6-(substituted)picolinate of Formula G with a halogen source to produce a 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of Formula I.

Scheme III



[0007] In Scheme III, the iodine, bromine or chlorine exchange step c) is not necessary. Thus, the present invention also concerns a process for the preparation of a 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of the Formula I



wherein


W represents Cl, Br or I;

R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy; and

R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO-, wherein n=1 or 2;

which comprises the following steps:

a) coupling a 4,5,6-trichloropicolinate of Formula A




wherein R¹ is as previously defined;

with an aryl, alkyl or alkenyl metal compound of the Formula F




wherein R is as previously defined and Met represents Zn-halide, Zn-R, tri-(C₁-C₄ alkyl)tin, copper, or B(OR²)(OR³), where R² and R³ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group in the presence of a transition metal catalyst to produce a 4,5-dichloro-6-(substituted)picolinate of Formula H



wherein R and R¹ are as previously defined;


b) fluorinating the 4,5-dichloro-6-(substituted)picolinate of Formula H with a fluoride ion source to produce a 4,5-difluoro-6-(substituted)picolinate of Formula J



J

wherein R¹ is as previously defined;

c) aminating the 4,5-difluoro-6-(substituted)picolinate of Formula J with ammonia to produce a 4-amino-5-fluoro-6-(substituted)picolinate of Formula K



K


wherein R and R¹ are as previously defined; and

d) halogenating the 4-amino-5-fluoro-6-(substituted)picolinate of Formula K with a halogen source to produce the 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of Formula I.

[0008] At any step in Schemes I-III, the ester substituent, R¹, may optionally be exchanged with a different R¹ substituent. These esters, including unsubstituted or substituted C₇-C₁₁ arylalkyl esters, can be prepared by direct esterification or transesterification reactions using techniques which are well known in the art.


[0009] Another aspect of the present invention is the novel intermediates produced during the present process, viz., compounds selected from the group consisting of:

a)




wherein R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy and R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl, which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO-, wherein n=1 or 2;


b)



wherein X represents I, Br, Cl or F, Y¹ represents H, Cl, Br, or I with the proviso that when X is Cl, Y¹ is H, Br or I, and R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl, which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO-, wherein n=1 or 2; and with the proviso that the compound does not have the following structure




c)y



wherein Y² represents H, and R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy, and R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl, which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO-, wherein n=1 or 2; and

d)



wherein R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl, or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy, and R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl, which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO-, wherein n=1 or 2.

[0010] The terms "alkyl," "alkenyl" and "alkynyl," as well as derivative terms such as "alkoxy," "acyl," "alkylthio" and "alkylsulfonyl," as used herein, include within their scope straight chain and branched chain moieties. Unless specifically stated otherwise, each may be unsubstituted or substituted with one or more substituents selected from but not limited to halogen, hydroxy, alkoxy, alkylthio, C₁-C₆ acyl, formyl, cyano, or aryloxy, provided that the substituents are sterically compatible and the rules of chemical bonding and strain energy are satisfied. The terms "alkenyl" and "alkynyl" are intended to include one or more unsaturated bonds.

[0011] The term "arylalkyl," as used herein, refers to a phenyl substituted alkyl group having a total of 7 to 11 carbon atoms, such as benzyl (-CH₂C₆H₅), 2-methylnaphthyl (-CH₂C₁₀H₇) and 1- or 2-phenethyl (-CH₂CH₂C₆H₅ or -CH(CH₃)C₆H₅). The phenyl group may itself be unsubstituted or substituted with one or more substituents independently

selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO- wherein n=1 or 2, provided that the substituents are sterically compatible and the rules of chemical bonding and strain energy are satisfied.


[0012] Unless specifically limited otherwise, the term "halogen," as well as derivative terms such as "halo," refers to fluorine, chlorine, bromine and iodine.

[0013] The 6-phenyl groups substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy may be of any orientation, but 4-substituted phenyl, 2,4-disubstituted phenyl, 2,3,4-trisubstituted phenyl, 2,4,5-trisubstituted phenyl, and 2,3,4,6-tetrasubstituted phenyl isomers are preferred.

[0014] The 4-amino-5-fluoro-3-halo-6-(substituted)picolinates are prepared from 4,5,6-trichloropicolinates by a series of steps involving fluorine exchange, amination, halogen exchange, halogenation and transition metal assisted coupling. The individual steps may be performed in different sequences.

[0015] The 4,5,6-trichloropicolinate starting materials are known compounds; see, for example, Example 3 in U.S. Patent 6,784,137 B2. Higher esters, including unsubstituted or substituted C₇-C₁₁ arylalkyl esters, can be prepared by direct esterification or transesterification reactions using techniques which are well known in the art.

[0016] In the fluorine exchange reaction, the fluorinated picolinate is prepared by reacting the corresponding chlorinated picolinate with at least one equivalent of fluoride ion source for each ring chlorine substituent to be exchanged.




Typical fluoride ion sources are alkali metal fluorides which include sodium fluoride (NaF), potassium fluoride (KF) and cesium fluoride (CsF), with KF and CsF being preferred. Fluoride salts such as tetrabutylammonium fluoride (*n*-Bu₄NF) may also be used. Preferably, the reaction is carried out in a polar aprotic solvent or reaction medium such as, dimethylsulfoxide (DMSO), *N*-methylpyrrolidinone (NMP), *N,N*-dimethylformamide (DMF), hexamethylphosphoramide (HMPA) or sulfolane. Additives such as crown ethers or phase transfer agents which are known to increase the rate of fluoride exchange may also be used. The temperature at which the reaction is conducted is not critical but usually is from 70 °C to 180 °C and preferably from 80 °C to 120 °C. Depending upon which solvent is employed in a particular reaction, the optimum temperature will vary. Generally speaking the lower the temperature the slower the reaction will proceed. The present reaction is typically conducted in the presence of vigorous agitation sufficient to maintain an essentially uniformly dispersed mixture of the reactants.

[0017] In conducting the fluorination reaction, neither the rate, nor the order, of addition of the reactants is critical. Usually, the solvent and alkali metal fluoride are mixed before the chlorinated picolinate is added to the reaction mixture. A typical reaction generally requires from 2 to 100 hours and is usually conducted at ambient atmospheric pressure.

[0018] While the exact amount of reactants is not critical, it is preferred to employ an amount of alkali metal fluoride which will supply at least an equimolar amount of fluorine atoms based on the number of chlorine atoms to be exchanged in the starting material, *i.e.*, at least an equimolar amount of alkali metal fluoride. After the reaction is completed the desired product is recovered by employing standard separation and purification techniques.

[0019] In the amination, a 4-fluoropicolinate is allowed to react with ammonia to replace the fluorine atom with an amino group.




While only a stoichiometric amount of ammonia is required, it is often convenient to use a large excess of ammonia. The reaction is carried out in an inert solvent, preferably, a polar aprotic solvent or reaction medium such as DMSO, NMP, DMF, HMPA or sulfolane. Alternatively, aqueous ammonium hydroxide can be used, with or without use of an

organic solvent. The temperature at which the reaction is conducted is not critical but usually is from 0 °C to 45 °C and preferably from 10 °C to 30 °C.

[0020] In conducting the amination reaction, the 4-fluoropicolinate is dissolved in the solvent, and the ammonia is added to the reaction mixture with cooling. Excess ammonia gas is typically bubbled into the reaction mixture. A typical reaction generally requires from 0.5 to 5 hours and is usually conducted at ambient atmospheric pressure.


[0021] The amine-containing products or intermediates obtained by any of these processes can be recovered by conventional means, such as evaporation or extraction, and can be purified by standard procedures, such as recrystallization or chromatography. Purification of the amine-containing products or intermediates can also be affected by protonation with an acid to form a salt which is isolated in higher purity by crystallization, precipitation or extraction. A variety of acids, such as hydrochloric acid, hydrobromic acid, nitric acid, acetic acid or sulfuric acid, can be used. Anhydrous hydrochloric acid is a preferred acid. The purified salt is then neutralized with a base to form the neutral amine-containing product or intermediate. Inorganic bases, such as sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate or sodium bicarbonate may be used. Organic bases such as triethylamine are preferred. Purification of the amine-containing product or intermediate may be performed in this manner immediately after the amination step, or after subsequent reactions, e.g., halogenation, coupling, have been preformed.

[0022] In the halogen (iodine, bromine or chlorine) exchange reaction, the 6-iodinated, 6-brominated or 6-chlorinated picolinate is prepared by reacting the corresponding 6-fluorinated picolinate with at least one equivalent of iodide, bromide or chloride.



Typically, the halogen exchange reaction is carried out in the presence of a large excess of anhydrous hydrogen iodide (HI), hydrogen bromide (HBr) or hydrogen chloride (HCl). The reaction is typically performed in the absence of water to minimize the formation of byproducts. The halogen exchange generally requires from 5 to 50 equivalents of HI, HBr or HCl, preferably from 10 to 20 equivalents. The reaction is carried out in an inert solvent, preferably, a polar solvent such as dioxane or acetic acid. The temperature at which the reaction is conducted is not critical but usually is from 75 °C to 150 °C and preferably from 100 °C to 125 °C. The reaction is typically performed in a sealed pressure reactor which is capable of containing HI, HBr or HCl gas. A typical reaction generally requires from 0.5 to 5 hours.

[0023] In the halogenation reaction, a chlorine, bromine or iodine atom is introduced into the 3-position of the picolinate by reacting the 3-unsubstituted picolinate with a halogen source in an inert solvent.



[0024] When the halogen atom at the 3-position is Cl, the chlorine source can be chlorine (Cl₂) itself or reagents such as sulfonyl chloride, *N*-chlorosuccinimide or 1,3-dichloro-5,5-dimethylhydantoin. When chlorine or sulfonyl chloride are used, a large excess of chlorinating agent is used. When chlorine gas is used, the reaction is performed in an inert solvent, preferably, a solvent such as dichloromethane, dichloromethane-water or acetic acid. When sulfonyl chloride is used, the reaction can be performed in an inert solvent, such as dichloromethane or in neat sulfonyl chloride. The temperature at which the reaction is conducted is not critical but usually is from 0 °C to 45 °C and preferably from 10 °C to 30 °C. A typical reaction generally requires from 0.5 to 5 hours. The chlorination reaction is usually conducted at ambient atmospheric pressure.

[0025] When the chlorinating agent used is *N*-chlorosuccinimide or 1,3-dichloro-5,5-dimethylhydantoin, the reaction is carried out using a stoichiometric amount of chlorinating reagent. For chlorinations using 1,3-dichloro-5,5-dimethylhydantoin as the chlorinating agent, both chlorines in the hydantoin are found to react. The reaction is performed in an inert polar solvent, such as DMF or acetonitrile. The temperature at which the reaction is conducted is not critical but usually is from 20 °C to 85 °C and preferably from 50 °C to 80 °C. When acetonitrile is used as solvent, it is convenient


to carry out the reaction at the reflux temperature. A typical reaction generally requires from 0.5 to 5 hours. The chlorination reaction is usually conducted at ambient atmospheric pressure.

[0026] When the halogen atom at the 3-position is Br, the bromine source can be bromine (Br₂) itself or reagents such as sulfonyl bromide, *N*-bromosuccinimide or 1,3-dibromo-5,5-dimethylhydantoin. When Br₂ is used as the brominating agent, a large excess can be employed, and the reaction is performed in an inert solvent, preferably, a solvent such as dichloromethane, dichloromethane-water or acetic acid. The temperature at which the reaction is conducted is not critical but usually is from 0 °C to 45 °C and preferably from 10 °C to 30 °C. A typical reaction generally requires from 0.5 to 5 hours. The bromination reaction is usually conducted at ambient atmospheric pressure.

[0027] When the brominating agent used is *N*-bromosuccinimide or 1,3-dibromo-5,5-dimethylhydantoin, the reaction is carried out using a stoichiometric amount of brominating reagent. The reaction is performed in an inert polar solvent, such as DMF or acetonitrile. The temperature at which the reaction is conducted is not critical but usually is from 20 °C to 85 °C and preferably from 50 °C to 80 °C. When acetonitrile is used as solvent, it is convenient to carry out the reaction at the reflux temperature. A typical reaction generally requires from 0.5 to 5 hours. The bromination reaction is usually conducted at ambient atmospheric pressure.

[0028] When the halogen atom at the 3-position is I, the iodine source can be iodine (I₂) itself or reagents such as iodine monochloride or *N*-iodosuccinimide. Periodic acid may be used in conjunction with I₂. When I₂ is used as the iodinating agent, a large excess of I₂ can be employed, and the reaction is performed in an inert solvent, preferably, a solvent such as dichloromethane, dichloromethane-water, methyl alcohol or acetic acid. The temperature at which the reaction is conducted is not critical but usually is from 0 °C to 45 °C and preferably from 10 °C to 30 °C. A typical reaction generally requires from 0.5 to 5 hours. The iodination reaction is usually conducted at ambient atmospheric pressure.

[0029] In the coupling reaction, a 6-iodo, bromo or choropicolinate is reacted with an aryl, alkyl or alkenyl metal compound where the metal is a Zn-halide, Zn-R, tri-(C₁-C₄ alkyl)tin, copper, or B(OR²)(OR³), where R² and R³ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group, in the presence of a transition metal catalyst.



"Catalyst" is a transition metal catalyst, in particular a palladium catalyst such as palladium diacetate or bis(triphenylphosphine)palladium(II) dichloride, or a nickel catalyst such as nickel(II) acetylacetonate or bis(triphenylphosphine)nickel(II) dichloride. In addition, catalysts can be prepared *in situ* from metal salts and ligands, such as palladium acetate and triphenylphosphine or nickel(II) chloride and triphenylphosphine. These *in situ* catalysts can be prepared by prior reaction of metal salt and ligand, followed by addition to the reaction mixture, or by separate addition of the metal salt and ligand directly to the reaction mixture.

[0030] Typically, coupling reactions are carried out in the absence of oxygen using an inert gas, such as nitrogen or argon. Techniques used to exclude oxygen from coupling reaction mixtures, such as sparging with inert gas, are well known to those skilled in the art. Examples of such techniques are described in *The Manipulation of Air-Sensitive Compounds*, 2nd ed.; Shriver, D. F., Drezzdon, M. A., Eds.; Wiley-Interscience, 1986. Sub-stoichiometric amounts of a catalyst are used, typically from 0.0001 equivalents to 0.1 equivalents. Additional amounts of ligand may optionally be added to increase catalyst stability and activity. In addition, additives such as Na_2CO_3 , K_2CO_3 , KF, CsF and NaF are typically added to the coupling reaction. The coupling reaction generally requires from 1 to 5 equivalents of such additive, preferably from 1 to 2 equivalents. Water may optionally be added to the coupling reaction to increase the solubility of these additives. The coupling reaction generally requires from 1 to 3 equivalents of an aryl, alkyl or alkenyl metal compound, preferably from 1 to 1.5 equivalents. The reaction is carried out in an inert solvent, such as toluene, tetrahydrofuran (THF), dioxane or acetonitrile. The temperature at which the reaction is conducted is not critical but usually is from 25 °C to 150 °C and preferably from 50 °C to 125 °C. A typical reaction generally requires from 0.5 to 24 hours. No particular order of addition of reactants is typically required. It is often operationally simpler to combine all reactants except the catalyst and then deoxygenate the reaction solution. Following deoxygenation, the catalyst can be added to commence the coupling reaction.

[0031] When the Met portion of the aryl, alkyl or alkenyl metal compound is a Zn-halide, Zn-R or copper, protection of reactive functional groups may be necessary. For example, if an amino substituent (-NHR or -NH₂) is present, protection of these reactive groups may be required. A variety of groups are known in the art for protection of amino groups from reaction with organometallic reagents. Examples of such protecting groups are described in Protective Groups in Organic

Synthesis, 3rd ed.; Greene, T. W.; Wuts, P. G. M., Eds.; Wiley-Interscience, 1999. The choice of which metal to use in R-Met is influenced by a number of factors, such as cost, stability, reactivity and the need to protect reactive functional groups.

[0032] The products obtained by any of these processes can be recovered by conventional means, such as evaporation or extraction, and can be purified by standard procedures, such as recrystallization or chromatography.


[0033] The following examples are presented to illustrate the invention.

Examples

Preparation of Starting Material

Example A. Propan-2-yl 4,5,6-trichloropicolinate


[0034]



[0035] Methyl 4,5,6-trichloropicolinate (14.19 grams (g), 59.0 millimoles (mmol)) was slurried in 2-propanol (150 milliliters (mL)) in a 250 mL round bottom flask equipped with a Dean-Stark trap and a reflux condenser. Sulfuric acid (98% H_2SO_4 ; 8.07 g, 82 mmol) was added, and the reaction mixture was heated to reflux. After 20 hours (h) at reflux, the majority of the 2-propanol (100 mL) was distilled overhead. The pot solidified upon cooling to room temperature. The resulting solid was stirred with ethyl acetate (EtOAc ; 500 mL) and saturated (satd) aqueous (aq) sodium bicarbonate solution (NaHCO_3 ; 500 mL). The organic layer was separated, washed with brine and then filtered through Celite. The organic extract was concentrated to 150 mL by rotary evaporation. Hexane (100 mL) was added, and the solution was stored at -20°C overnight. Crystals were collected, washed with hexane and dried in air (7.58 g, mp $104.6\text{--}105.7^\circ\text{C}$). A second crop was obtained by concentration of the filtrate to give a total of 10.36 g (65%) ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 8.23 (s, 1H, pyridine H), 5.16 (septet, $J = 6.3$ Hz, 1H, CHMe_2), 1.34 (d, $J = 6.3$ Hz, 6H, CHMe_2); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 161.9 (CO_2R), 150.6, 145.9, 145.0, 133.1, 125.4 (C3), 70.7 (CHMe_2), 21.7 (Me). Anal. Calcd for $\text{C}_9\text{H}_8\text{Cl}_3\text{NO}_2$: C, 40.26; H, 3.00; N, 5.22. Found: C, 40.25; H, 3.02; N, 5.22.

Example B. Benzyl 4,5,6-trichloropicolinate


[0036]



[0037] A mixture of methyl 4,5,6-trichloropicolinate (25 g, 0.10 moles (mol)) and benzyl alcohol (100 g, 0.2 mol) in a 250 mL three-neck round bottom flask was heated under nitrogen at 100°C . Titanium isopropoxide (0.6 g, 0.02 mol) was added. After 4 h at 100°C , the nearly colorless solution was cooled and transferred to a 250 mL round bottom single neck flask. Excess benzyl alcohol was removed under vacuum to give a nearly white solid (31 g, 94%): mp $125\text{--}126.5^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 8.08 (s, 1H, pyridine H), 7.42 (m, 2H, phenyl), 7.31 (m, 3H, phenyl), 5.40 (s, 2H, CH_2Ph); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 162.0 (CO_2R), 150.4, 145.0, 144.9, 134.7, 133.1, 128.3 (phenyl CH), 125.4 (pyridine CH), 67.88 (CH_2Ph).

Example C. Benzyl 4,5,6-trichloropicolinate

[0038]




[0039] A 22 L round bottom flask was fitted with a thermocouple, mechanical stirrer and a Dean-Stark trap which was connected to a nitrogen bubbler. The vessel was purged with nitrogen and then 4,5,6-trichloropicolinate (2547 g, 10.07 mol), pyridinium-p-toluenesulfonate (PPTS; 130 g, 0.52 mol), benzyl alcohol (2249 g, 20.8 mol) and xylenes (10278 g) were added. Stirring was started, and the pot was heated to 140 to 145 °C. The xylenes/water azeotrope was collected in the Dean-Stark trap over 5 h. The total amount of distillate collected was 4750 g (415 g was water). After the water stopped distilling overhead, a reactor sample was taken and analyzed by high-performance liquid chromatography (HPLC) to ensure less than 1.5 area% of starting carboxylic acid remained. The reaction was allowed to cool to room temperature and stirred overnight. Xylenes (4000 g) were removed by vacuum distillation. The solution was cooled to 85-100 °C and then vacuum transferred to a 30 L jacketed crystallization vessel that had been fitted with a mechanical stirrer and thermocouple. The vacuum was released with nitrogen and a nitrogen bubbler was placed on the crystallization vessel. To the xylene solution was added isopropyl alcohol (IPA; 6200 g) over 15 minutes (min). The resulting slurry was allowed to cool slowly to room temperature and then cooled further to 5 °C. The solid was collected by filtration, and the cake was washed with cold (5-10 °C) IPA (3731 g). The solid was dried in air to a constant weight providing white crystals (2765 g, gas chromatography (GC) internal standard purity 96.5 %, 84.3%).

Fluorine exchange

Example 1a. Propan-2-yl 4,5,6-trifluoropicolinate

[0040]




[0041] A 250 mL three-neck flask was equipped with a mechanical stirrer, a Dean-Stark trap with nitrogen inlet and a thermocouple. The flask was purged with nitrogen and CsF (23.38 g, 154 mmol) was added. Anhydrous DMSO (124 mL) was added, and the suspension was evacuated/backfilled (5x) with nitrogen. The suspension was heated at 80 °C for 30 min. DMSO (20 mL) was distilled off under vacuum at 75 °C to remove any residual water. Propan-2-yl 4,5,6-trichloropicolinate (13.45 g, 50.1 mmol) was added against a nitrogen purge. The reaction mixture was evacuated/backfilled (3x) and heated at 100 °C for 1 h with vigorous stirring.

[0042] A second 250 mL three-neck flask was equipped with a mechanical stirrer, a Dean-Stark trap with nitrogen inlet and a thermocouple. The flask was purged with nitrogen and CsF (24.41 g, 0.160 mmol) was added. Anhydrous DMSO (30 mL) was added, and the suspension was evacuated/backfilled (5x) with nitrogen. The suspension was heated to 80 °C for 30 min. DMSO (22 mL) was distilled off under vacuum at 75 °C to remove residual water. The cooled reaction mixture in the first flask was cannula filtered into the second flask under nitrogen. The reaction mixture was evacuated/backfilled (5x) and then heated to 100 °C for 1 h and then for an additional 90 min at 110 °C. Analysis of an aliquot by GC showed 96% propan-2-yl 4,5,6-trifluoropicolinate with only 1.4% propan-2-yl 5-chloro-4,6-difluoropicolinate present. The crude product solution was used directly in the amination step without further purification. Alternatively, the product can be isolated by aqueous workup, extraction with EtOAc and drying to give a light tan oil: ¹H NMR (400 MHz, CDCl₃) δ 7.94 (dd, *J*_{F-H} = 4.5, 8.7 Hz, 1H, H3), 5.30 (septet, *J*_{H-H} = 6.3 Hz, 1H, CHMe₂), 1.44 (d, *J*_{H-H} = 6.3 Hz, 6H, CHMe₂); ¹³C {¹H} NMR (101 MHz, CDCl₃) δ 161.2 (s, CO₂iPr), 157.3 (ddd, *J*_{F-C} = 266, 8, 6 Hz, C4/C6), 152.2 (ddd, *J*_{F-C}

= 241, 12, 5 Hz, C4/C6), 141.1 (dt, J_{F-C} = 14, 7 Hz, C2), 137.0 (ddd, J_{F-C} = 270, 31, 13 Hz, C5), 113.8 (dd, J_{F-C} = 17, 4 Hz, C3), 70.4 (s, CHMe₂), 21.33 (s, Me); ¹⁹F NMR (376 MHz, CDCl₃) δ -74.29 (dd, J_{F-F} = 24, 22 Hz, F6), -112.67 (ddd, J_{F-F} = 22, 19, J_{F-H} = 8.3 Hz, F4), -151.58 (ddd, J_{F-F} = 24, 19, J_{F-H} = 4.7 Hz, F5).

Example 1b. Propan-2-yl 4,5-difluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-picolinate

[0043]




[0044] A 250 mL three-neck flask was equipped with a distillation head, a nitrogen inlet, a mechanical stirrer and a thermocouple. The flask was charged with CsF (14.2 g, 93.0 mmol). Anhydrous DMSO (80 mL) was added, and the suspension was evacuated/backfilled (5x) with nitrogen. The suspension was heated at 80 °C for 30 min. DMSO (20 mL) was distilled off under vacuum to remove any residual water. Solid propan-2-yl 4,5-dichloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (10.44 g, 26.6 mmol) was added, and the solution was evacuated/backfilled with nitrogen (5x). The reaction mixture was heated to 105 °C under nitrogen. After 4 h at 105 °C, analysis of an aliquot by GC showed a 91:6 ratio of difluoro to monofluoro products. The reaction mixture was allowed to cool to room temperature.

[0045] A second 250 mL three-neck flask was equipped with a mechanical stirrer, a distillation head with a nitrogen inlet and a thermocouple. The flask was purged with nitrogen and CsF (7.5 g, 49.4 mmol) was added. Anhydrous DMSO (20 mL) was added, and the suspension was evacuated/backfilled (5x) with nitrogen. The suspension was heated at 80 °C for 30 min. DMSO (15 mL) was distilled off under vacuum to remove residual water. The cooled reaction mixture in the first flask was cannula filtered into the second flask under nitrogen. The reaction mixture was evacuated/backfilled (5x) and then heated to 100 °C for 2 h. Analysis of an aliquot by GC showed a 93:2 ratio of desired product to monofluoro intermediate. The reaction mixture was poured into ice-water (550 g) and was extracted with EtOAc (3 x 200 mL). The combined organic extracts were washed with water (5 x 100 mL) and brine, dried over magnesium sulfate (MgSO₄) and concentrated under reduced pressure to give a brown oil (8.57 g) which crystallized upon standing. The solid was purified by silica gel chromatography (330 g silica column; 0-50% EtOAc-hexane gradient) to give a white solid (4.98 g, 52%): mp 98.4-100.0 °C; ¹H NMR (400 MHz, acetone-*d*₆) δ 8.16 (dd, J_{F-H} = 10.0, 5.6 Hz, 1H, pyridine H), 7.43 (m, 2H, phenyl), 5.24 (hept, J_{H-H} = 6.3 Hz, 1H, CHMe₂), 4.01 (d, J_{F-H} = 1.1 Hz, 3H, OMe), 1.37 (d, J_{H-H} = 6.3 Hz, 6H, CHMe₂); ¹³C{¹H} NMR (101 MHz, acetone-*d*₆) δ 163.1 (CO₂R), 157.1 (dd, J_{F-C} = 264, 12 Hz, C4/C5), 154.8 (d, J_{F-C} = 254 Hz, C2' phenyl), 148.6 (dd, J_{F-C} = 267, 11 Hz, C4/C5), 147.4 (t, J_{F-C} = 6 Hz), 145.5 (d, J_{F-C} = 13 Hz), 144.6 (d, J_{F-C} = 13 Hz), 131.0, 126.8, 126.6 (d, J_{F-C} = 3.7 Hz), 123.2, 115.8 (d, J_{F-C} = 16 Hz), 70.6 (CHMe₂), 62.1 (d, J_{F-C} = 4 Hz, OMe), 21.9 (CHMe₂); ¹⁹F NMR (376 MHz, CDCl₃) δ -124.82 (dd, J_{F-F} = 21 Hz, J_{F-H} = 9.9 Hz, F4), -129.45 (dd, J_{F-F} = 27.8 Hz, J_{F-H} = 6.9 Hz, phenyl F), -141.81 (m, F5). Anal. Calcd for C₁₆H₁₃ClF₃NO₃: C, 53.42; H, 3.64; N, 3.89. Found: C, 53.77; H, 3.70; N, 3.95. Analysis of an aliquot by GC showed that the product was 95.5% pure with 1.7% monofluoro impurity.

Example 1c. Benzyl 4,5-difluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate


[0046]



[0047] A 250 mL three-neck flask was equipped with a distillation head, a nitrogen inlet, a mechanical stirrer and a thermocouple. The flask was charged with CsF (21.07 g, 139.0 mmol). Anhydrous DMSO (100 mL) was added, and the suspension was evacuated/backfilled (5x) with nitrogen. The suspension was heated at 80 °C for 30 min. DMSO (30 mL) was distilled off under vacuum to remove any residual water. Solid benzyl 4,5-dichloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (15.34 g, 34.8 mmol) was added, and the solution was evacuated/backfilled with nitrogen (5x). The reaction mixture was heated to 105 °C under nitrogen. After 6 h at 105 °C, analysis of an aliquot by GC showed no peak for the monofluoro intermediate. The reaction mixture was allowed to cool to room temperature. The reaction mixture was poured into ice-water (400 g) and was extracted with EtOAc (3 x 200 mL). The combined organic extracts were washed with satd NaHCO₃ solution, water (5 x 100 mL) and brine. The extracts were dried (MgSO₄) and concentrated under reduced pressure to give a tan solid (12.97 g). The solid was purified by flash chromatography (330 g silica column; 0-20% EtOAc-gradient) to give a white solid (9.95 g; 70%); mp 114-116 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.01 (dd, J_{F-H} = 9.4, 5.5 Hz, 1H, pyridine H), 7.53 - 7.20 (m, 7H, phenyl), 5.44 (s, 2H, CH₂Ph), 3.99 (d, J_{F-H} = 1.2 Hz, 3H, OMe); ¹³C NMR (101 MHz, CDCl₃) δ 162.8 (d, J_{F-C} = 3 Hz, CO₂Bn), 156.2 (dd, J_{F-C} = 267, 12 Hz), 153.9 (d, J_{F-C} = 255 Hz), 148.0 (dd, J_{F-C} = 269, 11 Hz), 145.4 (t, J_{F-C} = 7 Hz), 144.7 (d, J_{F-C} = 13 Hz), 144.6 (dd, J_{F-C} = 13, 2 Hz), 135.2 (s), 130.6 (d, J_{F-C} = 3 Hz), 125.6 (d, J_{F-C} = 4 Hz), 125.4 (d, J_{F-C} = 2 Hz), 122.0 (d, J_{F-C} = 14 Hz), 115.0 (d, J_{F-C} = 16 Hz), 67.9 (s, CH₂Ph), 61.6 (d, J_{F-C} = 5 Hz, OMe); ¹⁹F{¹H} NMR (376 MHz, CDCl₃) δ -123.90 (d, J_{F-F} = 19.7 Hz, F4), -128.37 (d, J_{F-F} = 33.5 Hz, F2'), -139.64 (dd, J_{F-F} = 33.5, 19.7 Hz, F5). Anal. Calcd for C₂₀H₁₃ClF₃NO₃: C, 58.91; H, 3.21; N, 3.43. Found: C, 59.03; H, 3.20; N, 3.39.

Example 1d. Benzyl 4,5-difluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate

[0048]




[0049] A 22 L straight wall jacketed reactor was fitted with an overhead stirrer, condenser, nitrogen inlet and outlet, and a stoppered solids loading port. The reactor was purged with nitrogen for 2 days. The loading port was opened, and benzyl 4,5-dichloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (2032 g, 4.12 mol, 89.3% purity) was quickly added to the reactor. CsF (2500 g, 16.46 mol) was quickly poured into the reactor. The reactor was next loaded with anhydrous (<100 ppm water) DMSO (8869 g). The mixture was heated to 110 °C for 2 h. The mixture was cooled to 35 °C and then filtered. The filtered salts were washed with DMSO (2 x 1108 g). The combined filtrate was cooled to 15-20 °C, and water (3023 g) was added with stirring over the course of 1 h. The mixture was cooled to 10-12 °C and then filtered. The collected solid was washed with 3:1 DMSO/water (1814 g) and then with water (2000 g). The resulting tan solid was dried to give the title compound (1626 g, 85.7 wt% HPLC purity (hexanophenone internal standard), 83%).

Amination

Example 2a. Propan-2-yl 4-amino-5,6-trifluoropicolinate

[0050]




[0051] The reaction mixture from Example 1a was filtered to remove Cs salts, and the salts were washed with DMSO

(50 mL). The DMSO washing solution was added to the DMSO solution (150 mL) which had been saturated with ammonia for 15 min. The flask was kept in a cold bath which kept the temperature near 16 °C. Ammonia was bubbled through the reaction mixture for 30 min, during which time a white precipitate formed. After 90 min, analysis of an aliquot by GC showed a single major peak for the 4-amino product. The reaction mixture was quenched by addition of satd aq ammonium chloride (NH₄Cl) solution (100 mL) followed by water (400 mL). The aqueous solution was extracted into ether (Et₂O 3 x 150 mL) and then EtOAc (3 x 150 mL). The combined organic extracts were washed with water (5 x 150 mL) and then brine. The extracts were dried (MgSO₄) and evaporated to a tan solid which was washed with 1:1 hexane-ether to give a light tan powder (5.57 g, 51.4% overall): mp 168-170 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.42 (d, *J*_{F-H} = 5.5 Hz, 1H, pyridine H), 5.22 (septet, *J* = 6.2 Hz, 1H, CHMe₂), 4.75 (s, 2H, NH₂), 1.35 (d, *J* = 6.2 Hz, 6H, CHMe₂); ¹³C{¹H} NMR (101 MHz, DMSO-*d*₆) δ 162.8 (CO₂R), 151.2 (dd, *J*_{F-C} = 228, 12 Hz, C6), 146.5 (dd, *J*_{F-C} = 9, 6 Hz, C2/C4), 139.3 (dd, *J*_{F-C} = 16, 5 Hz, C2/C4), 133.8 (dd, *J*_{F-C} = 252, 31 Hz, C5), 112.3 (C3), 68.8 (CHMe₂), 21.5 (Me); ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -91.9 (d, *J*_{F-F} = 26.6 Hz, F6), -163.9 (dd, *J*_{F-F} = 26.6, *J*_{H-F} = 5.6 Hz, F5). Anal. Calcd for C₉H₁₀F₂N₂O₂: C, 50.00; H, 4.66; N, 12.96. Found: C, 49.96; H, 4.65; N, 12.91.

Example 2b. Propan-2-yl 4-amino-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-picolinate


[0052]



[0053] Propan-2-yl 4,5-difluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (4.89 g, 13.9 mmol) was dissolved in DMSO (100 mL). Ammonia was bubbled through the solution for a total of 100 min over the course of 48 h. The reaction mixture was poured into ice-water (500 mL). The product was extracted into EtOAc (3 x 250 mL). The combined organic extracts were washed with water (5 x 100 mL) and then brine, dried (MgSO₄) and concentrated under reduced pressure to give a white solid (4.36 g, 88%): mp 180.2-181.9 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.54 (d, *J*_{F-H} = 6.5 Hz, 1H, pyridine H), 7.27 (m, 2H, phenyl), 5.27 (heptet, *J*_{H-H} = 6.3 Hz, 1H, CHMe₂), 4.69 (s, 2H, NH₂), 3.96 (d, *J*_{F-H} = 0.9 Hz, 3H, OMe), 1.38 (d, *J*_{H-H} = 6.3 Hz, 6H, CHMe₂); ¹³C{¹H} NMR (101 MHz, DMSO-*d*₆) δ 163.7 (CO₂R), 153.2 (d, *J*_{F-C} = 252 Hz), 146.8 (d, *J*_{F-C} = 254 Hz), 144.2 (d, *J*_{F-C} = 4 Hz), 143.9, 143.7, 139.0 (d, *J*_{F-C} = 14 Hz), 128.2 (d, *J*_{F-C} = 3 Hz), 126.0 (d, *J* = 3 Hz), 125.4 (d, *J*_{F-C} = 3 Hz), 123.9 (dd, *J*_{F-C} = 14, 3 Hz), 112.5 (d, *J*_{F-C} = 5 Hz), 68.5 (CHMe₂), 61.5 (d, *J*_{F-C} = 4 Hz, OMe), 21.56 (CHMe₂); ¹⁹F NMR (376 MHz, CDCl₃) δ -128.43 (dd, *J*_{F-F} = 32.0, *J*_{F-H} = 6.6 Hz), -142.27 (dd, dd, *J*_{F-F} = 32.0, *J*_{F-H} = 6.3 Hz). Anal. Calcd for C₁₆H₁₅ClF₂N₂O₃: C, 53.87; H, 4.24; N, 7.85. Found: C, 53.65; H, 4.28; N, 7.75.

Example 2c. Benzyl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate

[0054]




[0055] Benzyl 4,5-difluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (4.99 g, 12.2 mmol) was slurried in DMSO (100 mL). Ammonia was bubbled through the solution for 30 min. After stirring overnight, the reaction mixture was poured into ice-water (500 mL). The product was extracted into EtOAc (3 x 150 mL). The combined organic extracts were washed with water (5 x 100 mL) and brine, dried (MgSO₄) and concentrated under reduced pressure to give a white

solid (4.99 g, 101 %); ^1H NMR (400 MHz, CDCl_3) δ 7.52 (d, $J_{\text{F-H}} = 6.5$ Hz, 1H, pyridine H3), 7.45 - 7.38 (m, 2H), 7.37 - 7.17 (m, 5H), 5.38 (s, 2H, CH_2Ph), 4.67 (br s, 2H, NH_2), 3.94 (d, $J_{\text{F-H}} = 1.1$ Hz, 3H, OMe); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, CDCl_3) δ 164.4 (CO_2R), 153.9 (d, $J_{\text{F-C}} = 254$ Hz), 147.6 (d, $J_{\text{F-C}} = 256$ Hz), 144.4 (d, $J_{\text{F-C}} = 14$ Hz), 144.0 (d, $J_{\text{F-C}} = 5$ Hz), 142.2 (d, $J_{\text{F-C}} = 12$ Hz), 140.4 (d, $J_{\text{F-C}} = 15$ Hz), 135.6 (s), 129.5 (d, $J_{\text{F-C}} = 3$ Hz), 128.5 (CH), 128.3 (CH), 128.3 (CH), 125.6 (d, $J_{\text{F-C}} = 3$ Hz, CH), 125.2 (d, $J_{\text{F-C}} = 4$ Hz, CH), 123.3 (dd, $J_{\text{F-C}} = 14$, 4 Hz), 113.1 (d, $J_{\text{F-C}} = 4$ Hz, C3), 67.3 (s, CH_2Ph), 61.5 (d, $J_{\text{F-C}} = 4$ Hz, OMe); $^{19}\text{F}\{^1\text{H}\}$ NMR (376 MHz, CDCl_3) δ -128.54 (dd, $J = 30.7$, 5.2 Hz, F_2'), -141.84 (dd, $J = 30.8$, 6.5 Hz, F_5). HRMS-ESI (m/z): $[\text{M}]^+$ calcd for $\text{C}_{20}\text{H}_{15}\text{ClF}_2\text{N}_2\text{O}_3$, 404.0739; found, 404.0757.

Example 2d. Benzyl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate

[0056]




[0057] A solution of benzyl 4,5-difluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (65.0 g, 0.16 mol, 87% purity) in DMSO (400 mL) was prepared in a 1 L 4-neck flask fitted with a mechanical stirrer, thermometer, ammonia input and gas outlet. Ammonia gas (8.1 g, 0.48 mol, 3 equiv) was added by bubbling through a Teflon tube below the surface of the DMSO solution for 20 min. During the ammonia addition, the reaction changed to a pink/light red color and the internal temperature rose to 30 °C. After 5 h of stirring, additional ammonia gas (6.1 g, 0.36 mol, 2.25 equiv) was added over 20 min. After stirring for an additional 2.5 h, HPLC analysis indicated complete consumption of starting material. Nitrogen was bubbled into the reaction mixture, and the reaction was allowed to stir overnight. The reaction mixture was filtered to remove salts formed during the reaction, and the solids were washed with DMSO (50 mL). To the DMSO solution was added water (225 mL) dropwise over 1 h. The resulting precipitate was filtered and then washed with DMSO/water (2:1, 2 x 40 mL) followed by water (2 x 50 mL). The solid was dried to give the title compound (55.35 g, 87%, 87% purity by HPLC (hexanophenone internal standard)).

Halogen exchange

Example 3a. Propan-2-yl 4-amino-6-chloro-5-fluoropicolinate

[0058]




Propan-2-yl 4-amino-5,6-difluoropicolinate (4.25 g, 19.7 mmol) was dissolved in hydrochloric acid (HCl, 4 M in dioxane; 65 mL) in a 100 mL Hastalloy stirred Parr reactor. The reactor was heated to 100 °C for 2 h. Upon standing at room temperature overnight, a yellow crystalline solid formed. This solid was not soluble in EtOAc, but did dissolve upon shaking with satd aq NaHCO_3 solution (500 mL) and EtOAc (300 mL). The aqueous layer was extracted with EtOAc (2 x 250 mL). The combined organic extracts were washed with water (5 x 50 mL) and then brine. The extracts were dried (MgSO_4) and concentrated under vacuum to provide an off-white solid. The crude product was purified by column chromatography (120 g silica column; 0-100% hexane-EtOAc gradient) to give a white solid (2.11 g, 46%): mp 190.7-192.4 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.543 (d, $J_{\text{F-H}} = 5.7$ Hz, 1H), 6.91 (br s, 2H, NH_2), 5.09 (septet, $J = 6$ Hz, 1H, CHMe_2), 1.29 (d, $J = 6$ Hz, 6H, CHMe_2); $^{13}\text{C}\{^1\text{H}\}$ NMR (101 MHz, $\text{DMSO}-d_6$) δ 162.8 (CO_2R), 144.8 (d, $J_{\text{F-C}} = 12$ Hz, C2/C4), 143.4 (d, $J_{\text{F-C}} = 254$ Hz, C5), 142.7 (d, $J_{\text{F-C}} = 4.8$ Hz, C2/C4), 136.5 (d, $J_{\text{F-C}} = 17$ Hz, C6), 112.8 (d, $J_{\text{F-C}} = 5$

Hz, C3), 68.9 (CHMe₂), 21.6 (Me); ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ -141.0 (d, *J*_{F-H} = 6 Hz). Anal. Calcd for C₉H₁₀ClF₂N₂O₂: C, 46.47; H, 4.33; N, 12.04. Found: C, 46.50; H, 4.33; N, 11.96.

Halogenation

Example 4a. Propan-2-yl 4-amino-3,6-dichloro-5-fluoropicolinate


[0059]



[0060] Propan-2-yl 4-amino-6-chloro-5-fluoropicolinate (1.191 g, 5.12 mmol) was almost completely dissolved in CH₂Cl₂ (40 mL). Water (40 mL) was added. Chlorine was bubbled through the solution for 5 min. After 30 min, an aliquot of the reaction mixture was analyzed by GC, showing desired product and only 1.7% starting material. The aqueous layer was separated and extracted with CH₂Cl₂ (50 mL). The combined organic extracts were washed with satd aq NaHCO₃ solution and then brine. The extracts were dried (MgSO₄) and concentrated under reduced pressure to yield an orange oil. Flash chromatography (120 g silica column; 0-50% EtOAc-hexane gradient) afforded a bright yellow crystalline solid (394 mg, 28%): ¹H NMR (400 MHz, CDCl₃) δ 5.29 (septet, *J* = 6.3 Hz, 1H, CHMe₂), 5.19 (br s, 2H, NH₂), 1.40 (d, *J* = 6.3 Hz, 6H, CHMe₂); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 163.2 (CO₂R), 143.3 (d, *J*_{F-C} = 5 Hz, C2), 142.8 (d, *J*_{F-C} = 270 Hz, C5), 141.0 (d, *J*_{F-C} = 26 Hz, C4), 135.3 (d, *J*_{F-C} = 17 Hz, C6), 114.9 (s, C3), 70.6 (CHMe₂), 21.6 (s, Me); ¹⁹F NMR (376 MHz, CDCl₃) δ -136.5.

Example 4b. Propan-2-yl 4-amino-3,6-dichloro-5-fluoropicolinate


[0061]



[0062] Propan-2-yl 4-amino-6-chloro-5-fluoropicolinate (634 milligrams (mg), 2.73 mmol) was slurried in acetonitrile (11 mL). 1,3-Dichloro-5,5-dimethyl-hydantoin (303 mg, 1.54 mmol) was added as a solid, and the reaction mixture was stirred at reflux for 2.5 h. Additional 1,3-dichloro-5,5-dimethylhydantoin (50 mg, 0.25 mmol) was added, and the reaction mixture was stirred at reflux for an additional hour. Water (20 mL) was added. Acetonitrile was then removed by rotary evaporation to give an oily, yellow solid which was extracted with EtOAc (2 x 20 mL). The combined organic extracts were washed with 10% sodium bisulfite (NaHSO₃) solution, satd aq NaHCO₃ solution and brine, dried (MgSO₄) and concentrated under reduced pressure to provide a pale orange solid (671 mg, 92%): ¹H NMR (400 MHz, CDCl₃) δ 5.29 (septet, *J* = 6.3 Hz, 1H, CHMe₂), 5.19 (br s, 2H, NH₂), 1.40 (d, *J* = 6.3 Hz, 6H, CHMe₂); ¹⁹F NMR (376 MHz, CDCl₃) δ -136.5.

Example 4c. Methyl 4-amino-3-bromo-6-chloro-5-fluoropicolinate


[0063]



[0064] Methyl 4-amino-6-chloro-5-fluoropicolinate (1.0 g, 4.9 mmol) was combined with 1,3-dibromo-5,5-dimethylhydantoin (1.7 g, 5.9 mmol) in 1,2-dichloroethane (15 mL) and heated at reflux (83 °C) for 4 h. The cooled mixture was stirred with 10% NaHSO₃ solution and EtOAc (30 mL). The organic phase was separated, washed with water (2 x 20 mL), brine (10 mL), dried (Na₂SO₄) and concentrated. The residue was purified by silica gel chromatography (5-50% EtOAc-hexane) to give an orange solid (840 mg, 61%): mp 138-139 °C; EIMS *m/z* 282, 284; ¹H NMR (400 MHz, CDCl₃) δ 5.09 (s, 2H, NH₂), 3.97 (s, 3H, Me); ¹⁹F NMR (376 MHz, CDCl₃) δ -135.55 (s).

Example 4d. Methyl 4-amino-6-chloro-5-fluoro-3-iodopicolinate


[0065]



[0066] Methyl 4-amino-6-chloro-5-fluoropicolinate (2.2 g, 10.8 mmol) was dissolved in methyl alcohol (CH₃OH; 20 mL). The solution was treated with periodic acid (880 mg, 3.9 mmol) and iodine (2.2 g, 8.6 mmol) and then heated to reflux for 20 h. The mixture was cooled, and the volatiles were removed under vacuum. The residue was dissolved in EtOAc (50 mL) and then stirred with 10% NaHSO₃ solution (20 mL) for 10 min. The organic phase was separated and washed with brine (10 mL), dried (Na₂SO₄) and evaporated. The residue was purified by silica gel chromatography (5-50% EtOAc-hexane gradient) to give the title compound as a light orange solid (2.5 g, 70%): mp 149-151 °C; ESIMS *m/z* 330 ([M]⁺); ¹H NMR (400 MHz, CDCl₃) δ 5.17 (s, 2H, NH₂), 3.97 (s, 3H, OMe); ¹⁹F NMR (376 MHz, CDCl₃) δ -135.79 (s).

Example 4e. Propan-2-yl 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate


[0067]



[0068] Propan-2-yl 4-amino-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (3.065 g, 8.59 mmol) was dissolved in sulfolane (150 mL). The solution was stirred under nitrogen at room temperature for 8 h. During this time, a white precipitate formed. Hexane (100 mL) was added, and the mixture was stored overnight at -20 °C. The product was filtered, washed with hexane and then slurried in EtOAc (100 mL). The organic suspension was neutralized with satd aq NaHCO₃ solution, which caused all solids to dissolve. The organic layer was separated and washed with 10% aq NaHSO₃ solution and brine, dried (MgSO₄), and concentrated under reduced pressure to give a white solid (1.962 g, 58%): mp 109-111 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.25 (m, 2H), 5.32 (septet, *J* = 6.3 Hz, 1H, CHMe₂), 5.07 (br s, 2H), 3.97 (d, *J*_{F-H} = 1.0 Hz, 3H, OMe), 1.40 (d, *J* = 6.3 Hz, 6H, CHMe₂); ¹⁹F NMR (376 MHz, CDCl₃) δ -128.16 (dd, *J*_{F-F} = 33.3 Hz, *J*_{F-H} = 2.5 Hz, phenyl F), -138.35 (d, *J*_{F-F} = 33.4 Hz, pyridine F5).

Example 4f. Benzyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate


[0069]



[0070] Benzyl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate (2.07 g, 5.12 mmol) was slurried in acetonitrile (20 mL) in a scintillation vial. 1,3-Dichloro-5,5-dimethylhydantoin (554 mg, 2.181 mmol) was added as a solid, and the reaction mixture was stirred at reflux for 1 h. After cooling to room temperature, water (40 mL) was added to precipitate the product. The solid was collected on a Buchner funnel and washed with water. Drying under vacuum at 55 °C gave a white solid (2.187 g, 97%): ¹H NMR (400 MHz, CDCl₃) δ 7.50 - 7.41 (m, 2H, aromatic), 7.41 - 7.20 (m, 5H, aromatic), 5.42 (s, 2H, CH₂Ph), 4.92 (br s, 2H, NH₂), 3.97 (d, *J* = 1.2 Hz, 3H, OMe); ¹⁹F{¹H} NMR (CDCl₃) δ -128.19 (d, *J* = 33.9 Hz, F2'), -137.79 (d, *J* = 33.8 Hz, F5).

Example 4g. Benzyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate


[0071]



[0072] A solution of benzyl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate (53.0 g, 0.131 mol, 95% purity) in acetonitrile (450 mL) was prepared in a 1 L 3-neck round bottom flask equipped with mechanical stirrer and thermometer. 1,3-Dichloro-5,5-dimethylhydantoin 14.2 g, 0.072 mol, 0.55 equiv) was added. The reaction mixture was heated at 80 °C for 3 h. The reaction mixture was allowed to cool to room temperature and was then added dropwise to a dilute sodium bisulfite (NaHSO₃) solution (990 mL, 7.5 g of NaHSO₃) over 1 h. The resulting precipitate was isolated by filtration, washed with acetonitrile-water (1:1 v/v, 2 x 50 mL) and then water (2 x 50 mL). The solid was dried to give a pale yellow powder (53.44 g, 94%, 96.1% HPLC purity (octanophenone internal standard)).

Example 4h. Propan-2-yl 4-amino-3-iodo-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate


[0073]



[0074] Propan-2-yl 4-amino-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (600 mg, 1.682 mmol) was dissolved in acetic acid (5.6 mL). Sodium acetate (1.5 g, 18.50 mmol) was added followed by iodine monochloride (2.2 g, 13.45 mmol). An exotherm of approximately 10 °C was observed during the addition. The reaction mixture was heated at 80 °C for 20 h. The reaction mixture was then diluted with water and extracted with EtOAc. The organic layer was washed with water, satd aq NaHCO₃ solution, dried (MgSO₄) and then concentrated to dryness. The residue was applied to a silica gel column (80 g) then eluted (0-70% acetone-hexanes gradient) to give an orange solid (343 mg, 42%): mp 134-135 °C; ¹H NMR (400 MHz, acetone-*d*₆) δ 7.41 (dd, *J* = 8.5, 1.6 Hz, 1H), 7.33 (dd, *J* = 8.5, 6.8 Hz, 1H), 6.29 (s, 2H), 5.29 - 5.14 (septet, 1H), 3.98 (d, *J* = 1.1 Hz, 3H), 1.37 (d, *J* = 6.3 Hz, 6H); EIMS *m/z* 396.

Example 4i. Propan-2-yl 4-amino-3-bromo-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate

[0075]




[0076] Propan-2-yl 4-amino-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (500 mg, 1.402 mmol) was dissolved in dichloromethane (3.2 mL). *N*-Bromo-succinimide (299 mg, 1.682 mmol) was added, and the solution was stirred at ambient temperature for 20 h. The reaction mixture was then concentrated to dryness. The residue was purified by chromatography (40 g silica gel column; 0-70% EtOAc-hexanes) to give a tan solid (504 mg, 83%): ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.47 (dd, *J* = 8.5, 1.6 Hz, 1H), 7.29 (dd, *J* = 8.5, 7.1 Hz, 1H), 7.01 (s, 2H), 5.22 - 5.10 (m, 1H), 3.93 (d, *J* = 0.9 Hz, 3H), 1.32 (d, *J* = 6.3 Hz, 6H); EIMS *m/z* 350.

Coupling

Example 5a. Propan-2-yl 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate

[0077]




[0078] A 50 mL Schlenk flask was charged with propan-2-yl 4-amino-3,6-dichloro-5-fluoropicolinate (2.162 g, 8.09 mmol), 2-(4-chloro-2-fluoro-3-methoxyphenyl)-1,3,2-dioxaborinane (2.775 g, 11.35 mmol) and CsF (2.601 g, 17.12 mmol). Acetonitrile (20 mL) and water (7 mL) were added. The solution was evacuated/backfilled with nitrogen (5x). Solid bis(triphenylphosphine)palladium(II) dichloride (Pd(PPh₃)₂Cl₂; 281 mg, 4.9 mol%) was added. The solution was evacuated/backfilled with nitrogen (5x) and then heated at 70 °C under nitrogen for 3 h. After 3 h, the reaction mixture was allowed to cool to room temperature. The aqueous layer was separated. Water (20 mL) was added to the organic layer. The resulting dark brown precipitate was filtered and washed with water. The solid was dissolved in EtOAc (60 mL) and filtered to remove a small amount of black solid. The EtOAc solution was treated with activated carbon (175 mg) and filtered to give a wine-colored solution. Evaporation under reduced pressure gave a dark red solid. Purification (120 g silica column; 0-50% EtOAc-hexane gradient) gave a white solid (2.59 g, 82%): mp 110.6-112.1 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.25 (m, 2H), 5.32 (septet, *J* = 6.3 Hz, 1H, CHMe₂), 5.07 (br s, 2H), 3.97 (d, *J*_{F-H} = 1.0 Hz, 3H, OMe), 1.40 (d, *J* = 6.3 Hz, 6H, CHMe₂); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 164.2 (CO₂R), 153.8 (d, *J*_{F-C} = 254 Hz,

C5/C2'), 145.5 (d, $J_{F-C} = 258$ Hz, C5/C2'), 145.0 (d, $J_{F-C} = 5$ Hz), 144.4 (d, $J_{F-C} = 14$ Hz), 140.0 (d, $J_{F-C} = 13$ Hz), 137.5 (d, $J_{F-C} = 14$ Hz), 129.7 (d, $J_{F-C} = 3$ Hz), 125.4 (d, $J_{F-C} = 2$ Hz, CS'/C6'), 125.2 (d, $J_{F-C} = 3$ Hz, C5'C6'), 122.7 (dd, $J_{F-C} = 14, 4$ Hz, Cl'), 114.6 (C3), 70.2 (CHMe₂), 61.5 (d, $J_{F-C} = 4$ Hz, OMe), 21.6 (CHMe₂); ¹⁹F NMR (376 MHz, CDCl₃) δ -128.16 (dd, $J_{F-F} = 33.3$ Hz, $J_{F-H} = 2.5$ Hz, phenyl F), -138.35 (d, $J_{F-F} = 33.4$ Hz, pyridine F5). Anal. Calcd for C₁₆H₁₄Cl₂F₂N₂O₃: C, 49.12; H, 3.61; N, 7.16. Found: C, 49.30; H, 3.69; N, 7.08. The product was found to be 97.5% pure by HPLC.

Example 5b. Propan-2-yl 4,5-dichloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-picolinate


[0079]



[0080] A 100 mL Schlenk flask was charged with propan-2-yl 4,5,6-trichloropicolinate (10.46 g, 39.0 mmol), 2-(4-chloro-2-fluoro-3-methoxyphenyl)-1,3,2-dioxaborinane (13.27 g, 54.3 mmol) and CsF (11.76 g, 77.0 mmol). Acetonitrile (75 mL) and water (25 mL) were added. The reaction mixture was evacuated/backfilled with N₂ (5x). Solid Pd(PPh₃)₂Cl₂ (1.331 g, 1.896 mmol) was added. The solution was evacuated/backfilled with N₂ (5x) and then stirred at reflux for 2 h. A white solid precipitated upon cooling to room temperature. The solid was filtered, washed with water and dried in air (10.56 g, 69%): mp 123.8-127.7 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.20 (s, 1H, pyridine), 7.28 (dd, $J_{H-H} = 8.5$ Hz, $J_{F-H} = 1.6$ Hz, 1H), 7.13 (dd, $J_{H-H} = 8.5$ Hz, $J_{F-H} = 6.8$ Hz, 1H), 5.32 (septet, $J = 6.3$ Hz, 1H, CHMe₂), 3.99 (d, $J = 1.2$ Hz, 3H, OMe), 1.41 (d, $J = 6.3$ Hz, 6H, CHMe₂); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 162.7 (CO₂R), 153.8, 153.6 (d, $J_{F-C} = 253$ Hz, C2'), 146.7, 144.5 (d, $J_{F-C} = 13$ Hz), 144.0, 134.0, 130.0 (d, $J_{F-C} = 3.4$ Hz), 125.9, 125.3 (d, $J_{F-C} = 3$ Hz), 125.1 (d, $J_{F-C} = 3$ Hz), 70.4 (CHMe₂), 61.6 (d, $J_{F-C} = 4$ Hz, OMe), 21.7 (CHMe₂). Anal. Calcd for C₁₆H₁₃Cl₃NO₃: C, 48.94; H, 3.34; N, 3.57. Found: C, 48.91; H, 3.50; N, 3.51.

Example 5c. Methyl 4,5-dichloro-6-ethylpicolinate


[0081]



[0082] A 100 mL three-neck flask equipped with a reflux condenser and nitrogen inlet was charged with methyl 4,5,6-trichloropicolinate (2.40 g, 9.98 mmol). Anhydrous THF (50 mL) was added followed by *N,N*-dimethylethanolamine (0.20 g). The reaction mixture was sparged with nitrogen for 15 min. Solid Pd(PPh₃)₂Cl₂ (140 mg, 0.2 mmol) was added. The reaction mixture was stirred under nitrogen for 20 min. Diethylzinc (1 M solution in hexanes; 10 mL, 10 mmol) was added in 2 mL portions. When starting material was no longer observed through analysis by GC, the reaction mixture was quenched with water and extracted into EtOAc. The combined organic extracts were washed with brine, dried (MgSO₄) and concentrated under reduced pressure to a white solid (2.34 g). Analysis by GC-MS showed the solid contained 11% starting methyl 4,5,6-trichloropicolinate. ¹H NMR (400 MHz, CDCl₃) δ 8.06 (s, 1H, pyridine H), 4.01 (s, 3H, CO₂Me) 3.10 (q, $J = 8$ Hz, 2H, CH₂), 1.33 (t, $J = 8$ Hz, 3H, CH₂CH₃); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 164.4, 162.8, 145.4, 143.3, 132.9, 124.5 (C3), 53.1 (CO₂Me), 30.0 (CH₂), 12.3 (CH₂CH₃).

Example 5d. Benzyl 4,5-dichloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate


[0083]



[0084] A 250 mL three-neck flask equipped with a reflux condenser and nitrogen inlet was charged with benzyl 4,5,6-trichloropicolinate (17.77 g, 56.10 mmol), 2-(4-chloro-2-fluoro-3-methoxyphenyl)-1,3,2-dioxaborinane (19.20 g, 79.0 mmol) and CsF (17.04 g, 112.0 mmol). Acetonitrile (100 mL) and water (30 mL) were added. The reaction mixture was evacuated/backfilled with nitrogen (5x). Solid Pd(PPh₃)₂Cl₂ (1.724 g, 2.456 mmol) was added. The solution was evacuated/backfilled with nitrogen (5x) and then stirred at reflux for 90 min. A white solid precipitated upon cooling to room temperature. The solid was filtered, washed with water and dried in air (18.66 g, 75%): ¹H NMR (400 MHz, CDCl₃) δ 8.23 (s, 1H, pyridine H), 7.52 - 7.32 (m, 5H, phenyl), 7.27 (dd, J_{H-H} = 8.4 Hz, J_{F-H} = 1.7 Hz, 1H, aromatic), 7.10 (dd, J_{H-H} = 8.4 Hz, J_{F-H} = 6.8 Hz, 1H, aromatic), 5.44 (s, 2H, CH₂Ph), 3.98 (d, J_{F-H} = 1.3 Hz, 3H, OMe); ¹³C{¹H} NMR (101 MHz, CDCl₃) δ 163.0, 153.7, 153.5 (d, J_{F-C} = 253 Hz, C2'), 146.0, 144.5 (d, J_{F-C} = 13 Hz), 144.1, 135.0, 134.2, 129.9 (d, J_{F-C} = 3 Hz), 128.5, 126.1, 125.8 (d, J_{F-C} = 14 Hz), 125.3 (d, J_{F-C} = 3 Hz), 124.9 (d, J_{F-C} = 2 Hz), 67.9 (CH₂), 61.5 (d, J_{F-C} = 4 Hz, OMe). Anal. Calcd for C₂₀H₁₃Cl₃FO₃: C, 54.51; H, 2.97; N, 3.18. Found: C, 54.60; H, 3.08; N, 3.16.

Example 5e. Benzyl 4,5-dichloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate

[0085]




[0086] A 22-L straight wall reactor equipped with a mechanical stirrer was charged with tap water (3403 g). Dipotassium phosphate (K₂HPO₄; 2596 g, 14.9 mol) was then added, and the mixture was stirred until all solids dissolved while being purged with a nitrogen stream. After all the solids had dissolved, acetonitrile (8173 g) was loaded into the reactor. To a separate 30-L straight-wall jacketed reactor equipped with a bottom drain and a mechanical overhead stirrer was loaded 2-(4-chloro-2-fluoro-3-methoxyphenyl)-1,3,2-dioxaborinane (1724 g, 7.01 mol) and benzyl 4,5,6-trichloropicolinate (1630 g, 4.97 mol). The reactor was evacuated and filled with nitrogen (3x). The acetonitrile/H₂O mixture containing K₂HPO₄ was then transferred into the 30-L reactor, and the lines were rinsed with acetonitrile (1434 g). The slurry was sparged with nitrogen for 30 min, and then triphenylphosphine (114.8 g, 0.44 mol) was added. The slurry was then sparged with nitrogen for 15 min followed by addition of bis(benzonitrile)palladium(II) chloride (83.8 g, 0.22 mol). The bright yellow slurry was sparged with nitrogen for 15 min, and then the mixture was heated to 74-75 °C. After stirring at 74 °C for 3.3 h the reaction was deemed complete by HPLC analysis. At this stage the reactor cool-down temperature was set to 5 °C and immediately, cold water (4448 g, approximately 3 °C) was added to the reactor. The resulting precipitate was filtered to give a cream-white colored cake. The filter cake was washed with cold acetonitrile/H₂O (3345 g, 1.4:1, 8-10 °C) to afford an off-white wet cake. The cake was dried under a stream of nitrogen to a constant weight of 2044 g. HPLC analysis using an internal standard (tetraphenylethylene) showed the product was 90.0% pure and contained 1840 g (84.0%) of the product.

Purification of Ammonium Salts

Example 6a. Propan-2-yl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate


[0087]



[0088] Propan-2-yl 4-amino-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (1.50 g, 71% purity by LC area) was added to tetrahydrofuran (THF; 10 mL) and heated to 40 °C to provide a clear yellow solution. The solution was allowed to cool to room temperature and HCl (4 M in dioxane; 1.3 mL, 5.2 mmol) was added. After addition of HCl, solids precipitated from the solution and the reaction mixture was cooled to 0 °C. The solids were isolated by vacuum filtration and washed with cold THF (5 mL). The salt wet cake was added to THF (10 mL) and water (5 mL). Triethylamine (Et₃N; 0.8 mL, 5.7 mmol) was added to the mixture and the reaction mixture formed a clear biphasic solution. The reaction mixture was transferred to a separatory funnel, and the organic layer was separated. Hexanes (20 mL) was added to the organic layer and solids precipitated from solution. The reaction mixture was cooled to 0 °C and allowed to stir for 30 min. The solids were isolated by vacuum filtration, washed with hexanes (10 mL), and dried in a vacuum oven at 40 °C to provide propan-2-yl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate as a white solid (0.90 g, 89% purity by LC area): mp 174-176 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.54 (d, *J* = 6.4 Hz, 1H), 7.34 - 7.22 (m, 2H), 5.27 (septet, *J* = 6.3 Hz, 1H), 4.56 (br s, 2H), 3.97 (d, *J* = 1.0 Hz, 3H), 1.39 (d, *J* = 6.3 Hz, 6H); ¹³C NMR (101 MHz, CDCl₃) δ 164.03, 154.07 (d, *J* = 253.7 Hz), 147.70 (d, *J* = 256.1 Hz), 144.92 (d, *J* = 5.0 Hz), 144.48 (d, *J* = 13.9 Hz), 142.01 (d, *J* = 12.2 Hz), 140.58 (d, *J* = 14.4 Hz), 129.59 (d, *J* = 3.4 Hz), 125.85 (d, *J* = 3.7 Hz), 125.29 (d, *J* = 3.8 Hz), 123.57 (dd, *J* = 14.1, 3.7 Hz), 112.85 (d, *J* = 3.7 Hz), 69.58 (s), 61.63 (d, *J* = 4.5 Hz), 21.86 (s); ¹⁹F NMR (376 MHz, CDCl₃) δ -128.44 (d, *J* = 32.7 Hz), -142.30 (d, *J* = 31.3 Hz).

Example 6b. Propan-2-yl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl) 5-fluoropicolinate

[0089]




[0090] Propan-2-yl 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (1.75 g, 95% purity by LC area) was added to dichloromethane (15 mL), and the mixture was heated to 40 °C to provide a clear yellow solution. The solution was allowed to cool to room temperature and HCl (4 M in dioxane; 1.25 mL, 5 mmol) was added. After addition of HCl, solids precipitated from the solution, and the reaction mixture was cooled to 0 °C. The solid was isolated by vacuum filtration and washed with cold dichloromethane (5 mL). The salt wet cake was added to dichloromethane (10 mL) and water (5 mL). Et₃N (0.6 mL, 4.3 mmol) was added to the mixture, and the reaction mixture formed a clear biphasic solution. The reaction mixture was transferred to a separatory funnel, and the organic layer was separated. Hexanes (20 mL) was added to the organic layer and solids precipitated from solution. The reaction mixture was cooled to 0 °C, and the solids were isolated by vacuum filtration and washed with hexanes (10 mL). The solid was dried in a vacuum oven at 40 °C to provide propan-2-yl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate as a white solid. (1.02 g, 99% purity by LC area): mp 115-117 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.31 - 7.22 (m, 2H), 5.32

(septet, $J = 6.3$ Hz, 1H), 4.93 (br s, 2H), 3.98 (d, $J = 1.1$ Hz, 3H), 1.41 (d, $J = 6.3$ Hz, 6H); ^{13}C NMR (101 MHz, CDCl_3) δ 164.28, 153.93 (d, $J = 254.4$ Hz), 145.79 (d, $J = 230.6$ Hz), 145.23 (d, $J = 4.9$ Hz), 144.44 (d, $J = 13.8$ Hz), 139.97 (d, $J = 13.5$ Hz), 137.72 (d, $J = 13.8$ Hz), 129.90 (d, $J = 3.4$ Hz), 125.59 (d, $J = 3.2$ Hz), 125.41 (d, $J = 3.7$ Hz), 122.82 (dd, $J = 14.0, 4.4$ Hz), 114.82, 70.36, 61.66 (d, $J = 4.7$ Hz), 21.76; ^{19}F NMR (376 MHz, CDCl_3) δ -128.15 (d, $J = 34.1$ Hz), -138.44 (d, $J = 34.1$ Hz).

Example 6c. Methyl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate

[0091]




[0092] Benzyl 4-amino-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (3.00 g, 7.41 mmol) was added to CH_3OH (35 mL). Sodium methoxide (25 wt% in CH_3OH ; 2.0 mL, 8.9 mmol) was added to the reaction mixture and allowed to stir for 24 h. Water (50 mL) was added to the reaction mixture and the mixture was concentrated under reduced pressure to remove most of the CH_3OH . The mixture was extracted with EtOAc (2 x 40 mL), and the combined organic layers were washed with water (40 mL) and satd sodium chloride (40 mL) and concentrated under reduced pressure to provide methyl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate as a pale yellow solid (1.86 g; 67% purity by LC area).

[0093] Methyl 4-amino-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)picolinate (1.50 g, 67% purity by LC area) was added to THF (10 mL) and dichloromethane (5 mL), and the mixture was heated to 40°C to provide a clear yellow solution. The solution was allowed to cool to room temperature and HCl (4 M in dioxane; 1.4 mL, 5.6 mmol) was added. After addition of HCl, solids precipitated from the solution, and the reaction mixture was cooled to 0°C . The solid was isolated by vacuum filtration and washed with cold THF (5 mL). The salt wet cake was added to THF (25 mL) and water (10 mL). Et_3N (0.8 mL, 5.7 mmol) was added to the mixture, and the reaction mixture formed a clear biphasic solution. The reaction mixture was transferred to a separatory funnel, and the organic layer was separated and concentrated to a solution of ~10 mL. Hexanes (20 mL) was added to the organic layer, and solids precipitated from solution. The reaction mixture was cooled to 0°C , and the solids were isolated by vacuum filtration and washed with hexanes (10 mL). The solid was dried in a vacuum oven at 40°C to provide methyl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate as a white solid (0.53 g, 89% purity by LC area): mp $203\text{--}204^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ 7.60 (d, $J = 6.4$ Hz, 1H), 7.28 - 7.25 (m, 2H), 4.58 (br s, 2H), 3.97 (d, $J = 1.1$ Hz, 3H), 3.96 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ 165.27, 154.05 (d, $J = 254.1$ Hz), 147.83 (d, $J = 256.3$ Hz), 144.56 (d, $J = 13.7$ Hz), 144.31 (d, $J = 5.1$ Hz), 142.16 (d, $J = 12.4$ Hz), 140.63 (d, $J = 14.5$ Hz), 129.69 (d, $J = 3.5$ Hz), 125.63 (d, $J = 3.1$ Hz), 125.41 (d, $J = 3.7$ Hz), 123.43 (dd, $J = 14.1, 3.6$ Hz), 113.05 (d, $J = 3.7$ Hz), 61.64 (d, $J = 4.5$ Hz), 52.98; ^{19}F NMR (376 MHz, CDCl_3) δ -128.71 (d, $J = 28.6$ Hz), -141.94 (d, $J = 28.6$ Hz).

Example 6d. Methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl) 5-fluoropicolinate

[0094]




[0095] Benzyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate (3.00 g, 6.83mmol) was added to CH₃OH (35 mL). Sodium methoxide (25 wt% in CH₃OH; 1.9 mL, 8.2mmol) was added to the reaction mixture, and the reaction mixture was allowed to stir for 24 h. Water (50 mL) was added to the reaction mixture, and the mixture was concentrated under reduced pressure to remove most of the CH₃OH. The mixture was extracted with EtOAc (2 x 40 mL), and the combined organic layers were washed with water (40 mL) and satd sodium chloride (40 mL). Concentration of the volatiles under reduced pressure provided methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate as a pale yellow solid (2.04 g, 88% purity by LC area).

[0096] Methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate (1.25 g, 88% purity by LC area) was added to dichloromethane (15 mL) and THF (3 mL), and the mixture was heated to 40 °C to provide a clear yellow solution. The solution was allowed to cool to room temperature, and HCl (4 M in dioxane; 0.95 mL, 3.8mmol) was added. After addition of HCl, solids precipitated from the solution, and the reaction mixture was cooled to 0 °C. The solid was isolated by vacuum filtration and washed with cold dichloromethane (5 mL). The salt wet cake was added to dichloromethane (20 mL) and water (10 mL). Et₃N (0.6 mL, 4.3mmol) was added to the mixture, and the reaction mixture formed a clear biphasic solution. The reaction mixture was transferred to a separatory funnel, and the organic layer was separated and concentrated under reduced pressure to a solution of ~10 mL. Hexanes (20 mL) was added to the organic layer and solids precipitated from solution. The reaction mixture was cooled to 0 °C, and the solids were isolated by vacuum filtration and washed with hexanes (10 mL). The solid was dried in a vacuum oven at 40 °C to provide methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate as a white solid (0.51 g, 96% purity by LC area): mp 170-171 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.30 - 7.20 (m, 2H), 4.98 (br s, 2H), 3.98 (m, 6H); ¹³C NMR (101 MHz, CDCl₃) δ 164.67, 153.95 (d, J = 254.7 Hz), 145.93 (d, J = 245.8 Hz), 144.57 (s), 143.65 (d, J = 4.6 Hz), 140.21 (d, J = 13.3 Hz), 137.71 (d, J = 13.9 Hz), 130.02 (d, J = 3.5 Hz), 125.49 (d, J = 7.1 Hz), 125.49, 122.70 (dd, J = 14.1, 4.3 Hz), 115.89 (d, J = 1.5 Hz), 61.67 (d, J = 4.5 Hz), 53.06; ¹⁹F NMR (376 MHz, CDCl₃) δ -128.34 (d, J = 31.3 Hz), -137.60 (d, J = 32.7 Hz).

Example 6e. Benzyl 4-amino-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate

[0097]




[0098] A slurry of benzyl 4-amino-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-2-pyridinecarboxylate (98 g, 0.209 mol, 86.5% purity) and THF (300 mL) was gently heated to 28 °C to give a clear amber solution. The solution was allowed to cool to 20 °C, and HCl (4 M in 1,4-dioxane; 55 mL, 0.219 mol, 1.05 equiv) was added via syringe over 1 min. The solution rapidly became cloudy with solid formation, and the temperature of the mixture reached 28 °C. The mixture was cooled to below 10 °C. The precipitate was filtered and rinsed with cold THF (2 x 40 mL) to give a white solid (120.4 g). This solid was stirred with THF (300 mL) and water (100 mL). To this slurry was added Et₃N (30.5 mL, 0.219 mol) via syringe over 1 min, and the solids dissolved to give a cloudy mixture. The organic layer was separated. Hexanes (450 mL) was added with stirring, and the solution was cooled to below 10 °C. The resulting precipitate was filtered and rinsed

with hexanes (2 x 40 mL) to give a white solid (76.1 g, 95.1 wt% HPLC purity (hexanophenone internal standard)). An additional 7.56 g of 93.9% pure product was obtained from concentration of the filtrate. The ^1H and ^{19}F NMR spectra of the isolated product were identical to that observed in example 2c.

Example 6f. Benzyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate


[0099]



[0100] To a solution of benzyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropicolinate (5.00 g, 87% purity by LC area, 88 wt% by LC assay) in dichloromethane (25 mL) was added HCl (4 M in 1,4-dioxane; 3.1 mL, 12.4 mmol) via syringe. The solution became cloudy with solids formation after 1 min of stirring. The mixture was cooled in an ice bath to below 10 °C, filtered and rinsed with cold dichloromethane (5 mL) to give a white solid. This solid was slurried with dichloromethane (30 mL) and water (15 mL), and Et_3N (1.98 mL, 14.2 mmol) was added. The solids dissolved to give a two phase mixture. After stirring for 15 min, the mixture was transferred to a separatory funnel and the phases were allowed to separate over 15 min. The organic layer was separated, hexanes (60 mL) was added, and the mixture was cooled to below 10 °C. The solution quickly became cloudy and solids precipitated from the mixture. Vacuum filtration of the mixture provided a white solid, which was dried in a vacuum oven at 40 °C to afford benzyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoro-2-pyridinecarboxylate as a white solid. (3.11 g, 70%, 97% purity by LC area, 97 wt% by LC assay). The ^1H and ^{19}F NMR spectra of the isolated product were identical to that observed in example 4f.

Claims

1. A process for the preparation of a 4-amino-5-fluoro-3-halo-6-(substituted)-picolinate of the Formula I



wherein


W represents Cl, Br or I;

R represents $\text{C}_1\text{-C}_4$ alkyl, cyclopropyl, $\text{C}_2\text{-C}_4$ alkenyl or phenyl substituted with from 1 to 4 substituents independently selected from halogen, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_1\text{-C}_4$ alkoxy or $\text{C}_1\text{-C}_4$ haloalkoxy; and

R^1 represents $\text{C}_1\text{-C}_{12}$ alkyl or an unsubstituted or substituted $\text{C}_7\text{-C}_{11}$ arylalkyl which substituted $\text{C}_7\text{-C}_{11}$ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, halogenated $\text{C}_1\text{-C}_6$ alkyl, halogenated $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ alkylthio, $\text{C}(\text{O})\text{OC}_1\text{-C}_6$ alkyl, or where two adjacent substituents are taken together as $-\text{O}(\text{CH}_2)_n\text{O}-$ wherein $n=1$ or 2;


which comprises the following steps:

a) fluorinating the 4,5,6-trichloropicolinate of Formula A




A

wherein R¹ is as previously defined;
with a source of fluoride ion to produce a 4,5,6-trifluoropicolinate of Formula B




B

wherein R¹ is as previously defined;
b) aminating the 4,5,6-trifluoropicolinate of Formula B with ammonia to produce a 4-amino-5,6-difluoropicolinate of Formula C




C

wherein R¹ is as previously defined;
c) exchanging the fluoro substituent in the 6-position of the 4-amino-5,6-difluoropicolinate of Formula C with an iodo, bromo or chloro substituent by treating with an iodide, bromide or chloride source to produce a 4-amino-5-fluoro-6-halopicolinate of Formula D



D

wherein X represents Cl, Br or I,
and R¹ is as previously defined;
d) halogenating the 4-amino-5-fluoro-6-halopicolinate of Formula D with a halogen source to produce a 4-amino-3,6-dihalo-5-fluoropicolinate of Formula E



E

wherein W and X independently represent Cl, Br or I,

and R¹ is as previously defined; and

e) coupling the 4-amino-3,6-dihalo-5-fluoropicolinate of Formula E with an aryl, alkyl or alkenyl metal compound of the Formula F


R-Met

F

wherein R is as previously defined and Met represents Zn-halide, Zn-R, tri-(C₁-C₄ alkyl)tin, copper, or B(OR²)(OR³), where R² and R³ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group in the presence of a transition metal catalyst to produce the 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of Formula I.

2. The process of Claim 1 in which an amine-containing product or intermediate is purified by: a) protonating with an acid to form a salt, b) isolating the salt in higher purity by crystallization, precipitation or extraction, and c) neutralizing the purified salt with a base to form the purified neutral amine-containing product or intermediate.

3. A process for the preparation of a 4-amino-5-fluoro-3-halo-6-(substituted)-picolinate of the Formula I



wherein


W represents Cl, Br or I;

R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl, or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy; and

R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl, which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO- wherein n=1 or 2;


which comprises the following steps:

a) fluorinating a 4,5,6-trichloropicolinate of Formula A




wherein R¹ is as previously defined;

with a source of fluoride ion to produce a 4,5,6-trifluoropicolinate of Formula B




wherein R¹ is as previously defined;

b) aminating the 4,5,6-trifluoropicolinate of Formula B with ammonia to produce a 4-amino-5,6-difluoropicolinate of Formula C



wherein R¹ is as previously defined;

c) exchanging the fluoro substituent in the 6-position of the 4-amino-5,6-difluoropicolinate of Formula C with an iodo, bromo or chloro substituent by treating with an iodide, bromide or chloride source to produce a 4-amino-5-fluoro-6-halopicolinate of Formula D




wherein X represents Cl, Br or I; and

R¹ is as previously defined;

d) coupling the 4-amino-5-fluoro-6-halopicolinate of Formula D with an aryl, alkyl or alkenyl metal compound of the Formula F




wherein R is as previously defined and Met represents Zn-halide, Zn-R, tri-(C₁-C₄ alkyl)tin, copper, or B(OR²)(OR³), where R² and R³ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group in the presence of a transition metal catalyst to produce the 4-amino-5-fluoro-6-(substituted)-picolinate of Formula G.



wherein R, R¹ are as previously defined; and

e) halogenating the 4-amino-5-fluoro-6-(substituted)picolinate of Formula G with a halogen source to produce a 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of Formula I.

- 50
4. The process of Claim 3 in which an amine-containing product or intermediate is purified by: a) protonating with an acid to form a salt, b) isolating the salt in higher purity by crystallization, precipitation or extraction, and c) neutralizing the purified salt with a base to form the purified neutral amine-containing product or intermediate.
5. A process for the preparation of a 4-amino-5-fluoro-3-halo-6-(substituted)-picolinate of the Formula I
- 55



wherein


W represents Cl, Br or I;

R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy; and

R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO- wherein n=1 or 2;

which comprises the following steps:

a) coupling a 4,5,6-trichloropicolinate of Formula A




wherein R¹ is as previously defined;

with an aryl, alkyl or alkenyl metal compound of the Formula F




wherein R is as previously defined and Met represents Zn-halide, Zn-R, tri-(C₁-C₄ alkyl)tin, copper, or B(OR²)(OR³), where R² and R³ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group in the presence of a transition metal catalyst to produce a 4,5-dichloro-6-(substituted)picolinate of Formula H



wherein R and R¹ are as previously defined;


b) fluorinating the 4,5-dichloro-6-(substituted)picolinate of Formula H with a fluoride ion source to produce a 4,5-difluoro-6-(substituted)picolinate of Formula J



J

wherein R¹ is as previously defined;

c) aminating the 4,5-difluoro-6-(substituted)picolinate of Formula J with ammonia to produce a 4-amino-5-fluoro-6-(substituted)picolinate of Formula K



K


wherein R and R¹ are as previously defined; and

d) halogenating the 4-amino-5-fluoro-6-(substituted)picolinate of Formula K with a halogen source to produce the 4-amino-5-fluoro-3-halo-6-(substituted)picolinate of Formula I.

6. The process of Claim 5 in which an amine-containing product or intermediate is purified by: a) protonating with an acid to form a salt, b) isolating the salt in higher purity by crystallization, precipitation or extraction, and c) neutralizing the purified salt with a base to form the purified neutral amine-containing product or intermediate.


7. A compound selected from the group consisting of :

a)




wherein R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy and R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)ₙO- wherein n=1 or 2;

b)




wherein X represents I, Br, Cl or F, Y¹ represents H, Cl, Br, or I with the proviso that when X is Cl, Y¹ is H, Br or I, and R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl which substituted C₇-C₁₁

arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO- wherein n=1 or 2 and with the proviso that the compound does not have the following structure




c)



wherein Y² represents H, and R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy and R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO- wherein n=1 or 2; and


d)



wherein R represents C₁-C₄ alkyl, cyclopropyl, C₂-C₄ alkenyl, or phenyl substituted with from 1 to 4 substituents independently selected from halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy and R¹ represents C₁-C₁₂ alkyl or an unsubstituted or substituted C₇-C₁₁ arylalkyl which substituted C₇-C₁₁ arylalkyl has a phenyl group substituted with one or more substituents independently selected from halogen, nitro, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogenated C₁-C₆ alkyl, halogenated C₁-C₆ alkoxy, C₁-C₆ alkylthio, C(O)OC₁-C₆ alkyl, or where two adjacent substituents are taken together as -O(CH₂)_nO- wherein n=1 or 2.

Patentansprüche

1. Ein Verfahren für die Herstellung von einem 6-substituierten 4-Amino-5-fluor-3-halogenpicolinsäureester der Formel I,



wobei


W Cl, Br oder I darstellt;

R C₁-C₄-Alkyl, Cyclopropyl, C₂-C₄-Alkenyl oder Phenyl, substituiert mit 1 bis 4 Substituenten, die unabhängig voneinander aus Halogen, C₁-C₄-Alkyl, C₁-C₄-Halogenalkyl, C₁-C₄-Alkoxy oder C₁-C₄-Halogenalkoxy ausgewählt sind, darstellt; und

R¹ C₁-C₁₂-Alkyl oder ein unsubstituiertes oder substituiertes C₇-C₁₁-Arylalkyl darstellt, wobei das substituierte C₇-C₁₁-Arylalkyl eine Phenylgruppe, substituiert mit einem oder mehreren Substituenten, die unabhängig voneinander aus Halogen, Nitro, Cyano, C₁-C₆-Alkyl, C₁-C₆-Alkoxy, halogeniertem C₁-C₆-Alkyl, halogeniertem C₁-C₆-Alkoxy, C₁-C₆-Alkylthio, C(O)OC₁-C₆-Alkyl ausgewählt sind, aufweist oder wobei zwei benachbarte Substituenten als -O(CH₂)_nO- zusammengefasst werden, wobei n=1 oder 2 ist;


welches die folgenden Schritte umfasst:

a) Fluorieren des 4,5,6-Trichlorpicolinsäureesters der Formel A,




wobei R¹ wie vorstehend definiert ist;

mit einer Quelle für Fluoridionen, um einen 4,5,6-Trifluorpicolinsäureester der Formel B herzustellen,



wobei R¹ wie vorstehend definiert ist;

b) Aminieren des 4,5,6-Trifluorpicolinsäureesters der Formel B mit Ammoniak, um einen 4-Amino-5,6-difluorpicolinsäureester der Formel C herzustellen,




wobei R¹ wie vorstehend definiert ist;

c) Austauschen des Fluorsubstituenten in der 6-Position des 4-Amino-5,6-difluorpicolinsäureesters der Formel


EP 2 667 714 B9

C mit einem Iod-, Brom- oder Chlorsubstituenten durch Behandlung mit einer Iodid-, Bromid- oder Chloridquelle, um einen 4-Amino-5-fluor-6-halogenpicolinsäureester der Formel D herzustellen,



wobei X Cl, Br oder I darstellt
und R¹ wie vorstehend definiert ist;

d) Halogenieren des 4-Amino-5-fluor-6-halogenpicolinsäureesters der Formel D mit einer Halogenquelle, um einen 4-Amino-3,6-dihalogen-5-fluorpicolinsäureester der Formel E herzustellen,



wobei W und X unabhängig voneinander Cl, Br oder I darstellen und R¹ wie vorstehend definiert ist; und


e) Kuppeln des 4-Amino-3,6-dihalogen-5-fluorpicolinsäureesters der Formel E mit einer Aryl-, Alkyl- oder Alkenylmetallverbindung der Formel F,



wobei R wie vorstehend definiert ist und Met Zn-Halogenid, Zn-R, Tri-(C₁-C₄-Alkyl)zinn, Kupfer oder B(OR²)(OR³) darstellt, wobei R² und R³ unabhängig voneinander Wasserstoff oder C₁-C₄-Alkyl darstellen oder, wenn sie zusammengefasst werden, eine Ethylen- oder Propylengruppe bilden, in der Gegenwart eines Übergangsmetallkatalysators, um den 6-substituierten 4-Amino-5-fluor-3-halogenpicolinsäureester der Formel I herzustellen.

2. Das Verfahren gemäß Anspruch 1, in welchem ein aminhaltiges Produkt oder Zwischenprodukt gereinigt wird durch:
a) Protonieren mit einer Säure, um ein Salz zu bilden, b) Isolieren des Salzes in höherer Reinheit durch Kristallisation, Ausfällung oder Extraktion und c) Neutralisieren des gereinigten Salzes mit einer Base, um das gereinigte, neutrale aminhaltige Produkt oder Zwischenprodukt zu bilden.

3. Ein Verfahren für die Herstellung eines 6-substituierten 4-Amino-5-fluor-3-halogenpicolinsäureesters der Formel I,



wobei

W Cl, Br oder I darstellt;


R C₁-C₄-Alkyl, Cyclopropyl, C₂-C₄-Alkenyl oder Phenyl, substituiert mit 1 bis 4 Substituenten, die unabhängig voneinander aus Halogen, C₁-C₄-Alkyl, C₁-C₄-Halogenalkyl, C₁-C₄-Alkoxy oder C₁-C₄-Halogenalkoxy ausgewählt sind, darstellt; und

R¹ C₁-C₁₂-Alkyl oder ein unsubstituiertes oder substituiertes C₇-C₁₁-Arylalkyl darstellt, wobei das substituierte

C₇-C₁₁-Arylalkyl eine Phenylgruppe, substituiert mit einem oder mehreren Substituenten, die unabhängig voneinander aus Halogen, Nitro, Cyano, C₁-C₆-Alkyl, C₁-C₆-Alkoxy, halogeniertem C₁-C₆-Alkyl, halogeniertem C₁-C₆-Alkoxy, C₁-C₆-Alkylthio, C(O)OC₁-C₆-Alkyl ausgewählt sind, aufweist oder wobei zwei benachbarte Substituenten als -O(CH₂)_nO- zusammengekommen werden, wobei n=1 oder 2 ist;


welches die folgenden Schritte umfasst:

a) Fluorieren eines 4,5,6-Trichlorpicolinsäureesters der Formel A,




wobei R¹ wie vorstehend definiert ist;

mit einer Quelle für Fluoridionen, um einen 4,5,6-Trifluorpicolinsäureester der Formel B herzustellen,




wobei R¹ wie vorstehend definiert ist;

b) Aminieren des 4,5,6-Trifluorpicolinsäureesters der Formel B mit Ammoniak, um einen 4-Amino-5,6-difluorpicolinsäureester der Formel C herzustellen,



wobei R¹ wie vorstehend definiert ist;

c) Austauschen des Fluorsubstituenten in der 6-Position des 4-Amino-5,6-difluorpicolinsäureesters der Formel C mit einem Iod-, Brom- oder Chlorsubstituenten durch Behandlung mit einer Iodid- Bromid- oder Chloridquelle, um einen 4-Amino-5-fluor-6-halogenpicolinsäureester der Formel D herzustellen,



wobei X Cl, Br oder I darstellt


und R¹ wie vorstehend definiert ist;

d) Kuppeln des 4-Amino-5-fluor-6-halogenpicolinsäureesters der Formel D mit einer Aryl-, Alkyl- oder Alkenylmetallverbindung der Formel F,

R-Met

F


wobei R wie vorstehend definiert ist und Met Zn-Halogenid, Zn-R, Tri-(C₁-C₄-Alkyl)zinn, Kupfer oder B(OR²)(OR³) darstellt, wobei R² und R³ unabhängig voneinander Wasserstoff oder C₁-C₄-Alkyl darstellen oder, wenn sie zusammengekommen werden, eine Ethylen- oder Propylengruppe bilden, in der Gegenwart eines Übergangsmetallkatalysators, um den 6-substituierten 4-Amino-5-fluorpicolinsäureester der Formel G herzustellen,



wobei R, R¹ wie vorstehend definiert sind; und

e) Halogenieren des 6-substituierten 4-Amino-5-fluorpicolinsäureesters der Formel G mit einer Halogenquelle, um einen 6-substituierten 4-Amino-5-fluor-3-halogenpicolinsäureester der Formel I herzustellen.

4. Das Verfahren gemäß Anspruch 3, in welchem ein aminhaltiges Produkt oder Zwischenprodukt gereinigt wird durch:
 - a) Protonieren mit einer Säure, um ein Salz zu bilden, b) Isolieren des Salzes in höherer Reinheit durch Kristallisation, Ausfällung oder Extraktion und c) Neutralisieren des gereinigten Salzes mit einer Base, um das gereinigte, neutrale aminhaltige Produkt oder Zwischenprodukt zu bilden.
5. Ein Verfahren für die Herstellung von einem 6-substituierten 4-Amino-5-fluor-3-halogenpicolinsäureester der Formel I,



wobei


W Cl, Br oder I darstellt;

R C₁-C₄-Alkyl, Cyclopropyl, C₂-C₄-Alkenyl oder Phenyl, substituiert mit 1 bis 4 Substituenten, die unabhängig voneinander aus Halogen, C₁-C₄-Alkyl, C₁-C₄-Halogenalkyl, C₁-C₄-Alkoxy oder C₁-C₄-Halogenalkoxy ausgewählt sind, darstellt; und

R¹ C₁-C₁₂-Alkyl oder ein unsubstituiertes oder substituiertes C₇-C₁₁-Arylalkyl darstellt, wobei das substituierte C₇-C₁₁-Arylalkyl eine Phenylgruppe, substituiert mit einem oder mehreren Substituenten, die unabhängig voneinander aus Halogen, Nitro, Cyano, C₁-C₆-Alkyl, C₁-C₆-Alkoxy, halogeniertem C₁-C₆-Alkyl, halogeniertem C₁-C₆-Alkoxy, C₁-C₆-Alkylthio, C(O)OC₁-C₆-Alkyl ausgewählt sind, aufweist oder wobei zwei benachbarte Substituenten als -O(CH₂)_nO- zusammengekommen werden, wobei n=1 oder 2 ist;

welches die folgenden Schritte umfasst:

- a) Kuppeln eines 4,5-6-Trichlorpicolinsäureesters der Formel A,




A

wobei R¹ wie vorstehend definiert ist;
mit einer Aryl-, Alkyl- oder Alkenylmetallverbindung der Formel F,




wobei R wie vorstehend definiert ist und Met Zn-Halogenid, Zn-R, Tri-(C₁-C₄-Alkyl)zinn, Kupfer oder B(OR²)(OR³) darstellt, wobei R² und R³ unabhängig voneinander Wasserstoff oder C₁-C₄-Alkyl darstellen oder, wenn sie zusammengekommen werden, eine Ethylen- oder Propylengruppe bilden, in der Gegenwart eines Übergangsmetallkatalysators, um einen 6-substituierten 4,5-Dichlorpicolinsäureester der Formel H herzustellen,




H

wobei R und R¹ wie vorstehend definiert sind;
b) Fluorieren des 6-substituierten 4,5-Dichlorpicolinsäureesters der Formel H mit einer Quelle für Fluoridionen, um einen 6-substituierten 4,5-Difluorpicolinsäureester der Formel J herzustellen,



J

wobei R¹ wie vorstehend definiert ist;
c) Aminieren des 6-substituierten 4,5-Difluorpicolinsäureesters der Formel J mit Ammoniak, um einen 6-substituierten 4-Amino-5-fluorpicolinsäureester der Formel K herzustellen,



K


wobei R und R¹ wie vorstehend definiert sind; und
d) Halogenieren des 6-substituierten 4-Amino-5-fluorpicolinsäureesters der Formel K mit einer Halogenquelle, um den 6-substituierten 4-Amino-5-fluor-3-halogenpicolinsäureester der Formel I herzustellen.

6. Das Verfahren gemäß Anspruch 5, in welchem ein aminhaltiges Produkt oder Zwischenprodukt gereinigt wird durch:
a) Protonieren mit einer Säure, um ein Salz zu bilden, b) Isolieren des Salzes in höherer Reinheit durch Kristallisation, Ausfällung oder Extraktion und c) Neutralisieren des gereinigten Salzes mit einer Base, um das gereinigte, neutrale


aminhaltige Produkt oder Zwischenprodukt zu bilden.

7. Eine Verbindung ausgewählt aus der Gruppe bestehend aus:


a)




wobei R C₁-C₄-Alkyl, Cyclopropyl, C₂-C₄-Alkenyl oder Phenyl, substituiert mit 1 bis 4 Substituenten, die unabhängig voneinander aus Halogen, C₁-C₄-Alkyl, C₁-C₄-Halogenalkyl, C₁-C₄-Alkoxy oder C₁-C₄-Halogenalkoxy ausgewählt sind, darstellt und R¹ C₁-C₁₂-Alkyl oder ein unsubstituiertes oder substituiertes C₇-C₁₁-Arylalkyl darstellt, wobei das substituierte C₇-C₁₁-Arylalkyl eine Phenylgruppe, substituiert mit einem oder mehreren Substituenten, die unabhängig voneinander aus Halogen, Nitro, Cyano, C₁-C₆-Alkyl, C₁-C₆-Alkoxy, halogeniertem C₁-C₆-Alkyl, halogeniertem C₁-C₆-Alkoxy, C₁-C₆-Alkylthio, C(O)OC₁-C₆-Alkyl ausgewählt sind, aufweist oder wobei zwei benachbarte Substituenten als -O(CH₂)_nO- zusammengefasst werden, wobei n=1 oder 2 ist;



wobei X I, Br, Cl oder F darstellt, Y¹ H, Cl, Br oder I darstellt, unter der Voraussetzung, dass, wenn X Cl darstellt, Y¹ H, Br oder I darstellt und R¹ C₁-C₁₂-Alkyl oder ein unsubstituiertes oder substituiertes C₇-C₁₁-Arylalkyl darstellt, wobei das substituierte C₇-C₁₁-Arylalkyl eine Phenylgruppe, substituiert mit einem oder mehreren Substituenten, die unabhängig voneinander aus Halogen, Nitro, Cyano, C₁-C₆-Alkyl, C₁-C₆-Alkoxy, halogeniertem C₁-C₆-Alkyl, halogeniertem C₁-C₆-Alkoxy, C₁-C₆-Alkylthio, C(O)OC₁-C₆-Alkyl ausgewählt sind, aufweist oder wobei zwei benachbarte Substituenten als -O(CH₂)_nO- zusammengefasst werden, wobei n=1 oder 2 ist und unter der Voraussetzung, dass die Verbindung nicht die folgende Struktur aufweist




c)



wobei Y² H darstellt und R C₁-C₄-Alkyl, Cyclopropyl, C₂-C₄-Alkenyl oder Phenyl, substituiert mit 1 bis 4 Sub-

stituenten, die unabhängig voneinander aus Halogen, C₁-C₄-Alkyl, C₁-C₄-Halogenalkyl, C₁-C₄-Alkoxy oder C₁-C₄-Halogenalkoxy ausgewählt sind, darstellt und R¹ C₁-C₁₂-Alkyl oder ein unsubstituiertes oder substituiertes C₇-C₁₁-Arylalkyl darstellt, wobei das substituierte C₇-C₁₁-Arylalkyl eine Phenylgruppe, substituiert mit einem oder mehreren Substituenten, die unabhängig voneinander aus Halogen, Nitro, Cyano, C₁-C₆-Alkyl, C₁-C₆-Alkoxy, halogeniertem C₁-C₆-Alkyl, halogeniertem C₁-C₆-Alkoxy, C₁-C₆-Alkylthio, C(O)OC₁-C₆-Alkyl ausgewählt sind, aufweist oder wobei zwei benachbarte Substituenten als -O(CH₂)_nO-zusammengenommen werden, wobei n=1 oder 2 ist; und


d)



wobei R C₁-C₄-Alkyl, Cyclopropyl, C₂-C₄-Alkenyl oder Phenyl, substituiert mit 1 bis 4 Substituenten, die unabhängig voneinander aus Halogen, C₁-C₄-Alkyl, C₁-C₄-Halogenalkyl, C₁-C₄-Alkoxy oder C₁-C₄-Halogenalkoxy ausgewählt sind, darstellt und R¹ C₁-C₁₂-Alkyl oder ein unsubstituiertes oder substituiertes C₇-C₁₁-Arylalkyl darstellt, wobei das substituierte C₇-C₁₁-Arylalkyl eine Phenylgruppe, substituiert mit einem oder mehreren Substituenten, die unabhängig voneinander aus Halogen, Nitro, Cyano, C₁-C₆-Alkyl, C₁-C₆-Alkoxy, halogeniertem C₁-C₆-Alkyl, halogeniertem C₁-C₆-Alkoxy, C₁-C₆-Alkylthio, C(O)OC₁-C₆-Alkyl ausgewählt sind, aufweist oder wobei zwei benachbarte Substituenten als -O(CH₂)_nO-zusammengenommen werden, wobei n=1 oder 2 ist.

Revendications

1. Procédé de préparation d'un 4-amino-5-fluoro-3-halogéno-6-substituant-picolinate, de formule I :




dans laquelle

- W représente un atome de chlore, de brome ou d'iode,
- R représente un groupe alkyle en C₁-C₄, cyclopropyle, alcényle en C₂-C₄, ou phényle porteur de 1 à 4 substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes alkyle en C₁-C₄, halogéno-alkyle en C₁-C₄, alkoxy en C₁-C₄ et halogéno-alkoxy en C₁-C₄,
- et R¹ représente un groupe alkyle en C₁-C₁₂, ou un groupe aryl-alkyle en C₇-C₁₁ avec ou sans substituant(s), lequel groupe aryl-alkyle en C₇-C₁₁, s'il est porteur de substituant(s), comprend un groupe phényle qui porte un ou plusieurs substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes nitro, cyano, alkyle en C₁-C₆, alkoxy en C₁-C₆, halogéno-alkyle en C₁-C₆, halogéno-alkoxy en C₁-C₆, alkyl-thio en C₁-C₆ et C(O)O(alkyle en C₁-C₆), ou dont deux substituants adjacents forment ensemble un chaînon -O(CH₂)_nO- où l'indice n vaut 1 ou 2 ;


lequel procédé comporte les étapes suivantes :

- a) effectuer la fluoration d'un 4,5,6-trichloro-picolinate, de formule A :




A

dans laquelle R¹ a la signification indiquée plus haut,
au moyen d'une source d'ions fluorure, de manière à produire un 4,5,6-trifluoro-picolinate, de formule B :




B

dans laquelle R¹ a la signification indiquée plus haut ;
b) effectuer une amination de ce 4,5,6-trifluoro-picolinate de formule B, avec de l'ammoniac, de manière à
produire un 4-amino-5,6-difluoro-picolinate, de formule C :




C

dans laquelle R¹ a la signification indiquée plus haut ;
c) effectuer l'échange du substituant fluoro en position 6 de ce 4-amino-5,6-difluoro-picolinate de formule C,
contre un substituant iodo, bromo ou chloro, par traitement avec une source d'iodure, de bromure ou de chlorure,
de manière à produire un 4-amino-5-fluoro-6-halogéno-picolinate, de formule D :



D

dans laquelle X représente un atome de chlore, de brome ou d'iode, et R¹ a la signification indiquée plus haut ;
d) effectuer une halogénéation de ce 4-amino-5-fluoro-6-halogéno-picolinate de formule D, avec une source
d'halogène, de manière à produire un 4-amino-3,6-dihalogéno-5-fluoro-picolinate, de formule E :



E

dans laquelle W et X représentent chacun, indépendamment, un atome de chlore, de brome ou d'iode, et R¹ a la signification indiquée ci-dessus ;

e) et effectuer un couplage de ce 4-amino-3,6-dihalogéno-5-fluoro-picolinate de formule E, avec un composé de type aryl-métal, alkyl-métal ou alcényl-métal, de formule F :

R-Mét


F

dans laquelle R a la signification indiquée plus haut et le symbole Mét représente un fragment zinc-halogénure ou de formule Zn-R, un fragment tri(alkyle en C₁-C₄)-étain, un atome de cuivre ou un groupe de formule B(OR²)(OR³) où R² et R³ représentent chacun, indépendamment l'un de l'autre, un atome d'hydrogène ou un groupe alkyle en C₁-C₄, ou représentent des entités qui forment conjointement un groupe éthanediyle ou propandiyle,

en présence d'un catalyseur à base d'un métal de transition, de manière à produire un 4-amino-5-fluoro-3-halogéno-6-substituant-picolinate de formule I.

2. Procédé selon la revendication 1, dans lequel on purifie un produit ou un intermédiaire porteur de groupe(s) amino en réalisant les étapes suivantes : (a) effectuer une protonation avec un acide, de manière à former un sel ; (b) isoler ce sel à un degré de pureté plus élevé, par cristallisation, précipitation ou extraction, (c) et neutraliser le sel ainsi purifié avec une base, de manière à obtenir le produit ou l'intermédiaire porteur de groupe(s) amino purifié à l'état neutre.

3. Procédé de préparation d'un 4-amino-5-fluoro-3-halogéno-6-substituant-picolinate, de formule I :




I

dans laquelle

- W représente un atome de chlore, de brome ou d'iode,
- R représente un groupe alkyle en C₁-C₄, cyclopropyle, alcényle en C₂-C₄, ou phényle porteur de 1 à 4 substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes alkyle en C₁-C₄, halogéno-alkyle en C₁-C₄, alcoxy en C₁-C₄ et halogéno-alcoxy en C₁-C₄, - et R¹ représente un groupe alkyle en C₁-C₁₂, ou un groupe aryl-alkyle en C₇-C₁₁ avec ou sans substituant(s), lequel groupe aryl-alkyle en C₇-C₁₁, s'il est porteur de substituant(s), comprend un groupe phényle qui porte un ou plusieurs substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes nitro, cyano, alkyle en C₁-C₆, alcoxy en C₁-C₆, halogéno-alkyle en C₁-C₆, halogéno-alcoxy en C₁-C₆, alkyl-thio en C₁-C₆ et C(O)O(alkyle en C₁-C₆), ou dont deux substituants adjacents forment ensemble un chaînon -O(CH₂)_nO- où l'indice n vaut 1 ou 2 ;


lequel procédé comporte les étapes suivantes :

- a) effectuer la fluoration d'un 4,5,6-trichloro-picolinate, de formule A :




A

dans laquelle R¹ a la signification indiquée plus haut,
au moyen d'une source d'ions fluorure, de manière à produire un 4,5,6-trifluoro-picolinate, de formule B :




B

dans laquelle R¹ a la signification indiquée plus haut ;
b) effectuer une amination de ce 4,5,6-trifluoro-picolinate de formule B, avec de l'ammoniac, de manière à produire un 4-amino-5,6-difluoro-picolinate, de formule C :



C

dans laquelle R¹ a la signification indiquée plus haut ;
c) effectuer l'échange du substituant fluoro en position 6 de ce 4-amino-5,6-difluoro-picolinate de formule C, contre un substituant iodo, bromo ou chloro, par traitement avec une source d'iodure, de bromure ou de chlorure, de manière à produire un 4-amino-5-fluoro-6-halogéno-picolinate, de formule D :



D

dans laquelle X représente un atome de chlore, de brome ou d'iode, et R¹ a la signification indiquée plus haut ;
d) effectuer un couplage de ce 4-amino-5-fluoro-6-halogéno-picolinate de formule D, avec un composé de type aryl-métal, alkyl-métal ou alcényl-métal, de formule F :


R-Mét

F

dans laquelle R a la signification indiquée plus haut et le symbole Mét représente un fragment zinc-halogénure

ou de formule $Zn-R$, un fragment tri(alkyle en C_1-C_4)-étain, un atome de cuivre ou un groupe de formule $B(OR^2)(OR^3)$ où R^2 et R^3 représentent chacun, indépendamment l'un de l'autre, un atome d'hydrogène ou un groupe alkyle en C_1-C_4 , ou représentent des entités qui forment conjointement un groupe éthanediyle ou propa-

nediyle,
en présence d'un catalyseur à base d'un métal de transition, de manière à produire un 4-amino-5-fluoro-6-substituant-picolinate, de formule G :




G

dans laquelle R et R^1 ont les significations indiquées plus haut ;

e) et effectuer une halogénéation de ce 4-amino-5-fluoro-6-substituant-picolinate de formule G, avec une source d'halogène, de manière à produire un 4-amino-5-fluoro-3-halogéno-6-substituant-picolinate de formule I.

4. Procédé selon la revendication 3, dans lequel on purifie un produit ou un intermédiaire porteur de groupe(s) amino en réalisant les étapes suivantes : (a) effectuer une protonation avec un acide, de manière à former un sel ; (b) isoler ce sel à un degré de pureté plus élevé, par cristallisation, précipitation ou extraction, (c) et neutraliser le sel ainsi purifié avec une base, de manière à obtenir le produit ou l'intermédiaire porteur de groupe(s) amino purifié à l'état neutre.

5. Procédé de préparation d'un 4-amino-5-fluoro-3-halogéno-6-substituant-picolinate, de formule I :




I

dans laquelle

- W représente un atome de chlore, de brome ou d'iode,
- R représente un groupe alkyle en C_1-C_4 , cyclopropyle, alcényle en C_2-C_4 , ou phényle porteur de 1 à 4 substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes alkyle en C_1-C_4 , halogéno-alkyle en C_1-C_4 , alcoxy en C_1-C_4 et halogéno-alcoxy en C_1-C_4 ,
- et R^1 représente un groupe alkyle en C_1-C_{12} , ou un groupe aryl-alkyle en C_7-C_{11} avec ou sans substituant(s), lequel groupe aryl-alkyle en C_7-C_{11} , s'il est porteur de substituant(s), comprend un groupe phényle qui porte un ou plusieurs substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes nitro, cyano, alkyle en C_1-C_6 , alcoxy en C_1-C_6 , halogéno-alkyle en C_1-C_6 , halogéno-alcoxy en C_1-C_6 , alkyl-thio en C_1-C_6 et $C(O)O$ (alkyle en C_1-C_6), ou dont deux substituants adjacents forment ensemble un chaînon $-O(CH_2)_nO-$ où l'indice n vaut 1 ou 2 ;

lequel procédé comporte les étapes suivantes :

- a) effectuer le couplage d'un 4,5,6-trichloro-picolinate, de formule A :




A

dans laquelle R¹ a la signification indiquée plus haut,
avec un composé de type aryl-métal, alkyl-métal ou alcényl-métal, de formule F :




dans laquelle R a la signification indiquée plus haut et le symbole Mét représente un fragment zinc-halogénure ou de formule Zn-R, un fragment tri(alkyle en C₁-C₄)-étain, un atome de cuivre ou un groupe de formule B(OR²)(OR³) où R² et R³ représentent chacun, indépendamment l'un de l'autre, un atome d'hydrogène ou un groupe alkyle en C₁-C₄, ou représentent des entités qui forment conjointement un groupe éthanediyle ou propandiyyle,
en présence d'un catalyseur à base d'un métal de transition, de manière à produire un 4,5-dichloro-6-substituant-picolinate, de formule H :




H

dans laquelle R et R¹ ont les significations indiquées plus haut ;
b) effectuer une fluoration de ce 4,5-dichloro-6-substituant-picolinate de formule H, au moyen d'une source d'ions fluorure, de manière à produire un 4,5-difluoro-6-substituant-picolinate, de formule J :



J

dans laquelle R¹ a la signification indiquée plus haut ;
c) effectuer une amination de ce 4,5-difluoro-6-substituant-picolinate de formule J, avec de l'ammoniac, de manière à produire un 4-amino-5-fluoro-6-substituant-picolinate, de formule K :



K


dans laquelle R et R¹ ont les significations indiquées plus haut ;

d) et effectuer une halogénéation de ce 4-amino-5-fluoro-6-substituant-picolinate de formule K, avec une source d'halogène, de manière à produire un 4-amino-5-fluoro-3-halogéno-6-substituant-picolinate de formule I.

6. Procédé selon la revendication 5, dans lequel on purifie un produit ou un intermédiaire porteur de groupe(s) amino en réalisant les étapes suivantes : (a) effectuer une protonation avec un acide, de manière à former un sel ; (b) isoler ce sel à un degré de pureté plus élevé, par cristallisation, précipitation ou extraction, (c) et neutraliser le sel ainsi purifié avec une base, de manière à obtenir le produit ou l'intermédiaire porteur de groupe(s) amino purifié à l'état neutre.


7. Composé choisi dans l'ensemble formé par ceux répondant aux formules suivantes :

a)




dans laquelle R représente un groupe alkyle en C₁-C₄, cyclopropyle, alcényle en C₂-C₄, ou phényle porteur de 1 à 4 substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes alkyle en C₁-C₄, halogéno-alkyle en C₁-C₄, alcoxy en C₁-C₄ et halogéno-alcoxy en C₁-C₄, et R¹ représente un groupe alkyle en C₁-C₁₂, ou un groupe aryl-alkyle en C₇-C₁₁ avec ou sans substituant(s), lequel groupe aryl-alkyle en C₇-C₁₁, s'il est porteur de substituant(s), comprend un groupe phényle qui porte un ou plusieurs substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes nitro, cyano, alkyle en C₁-C₆, alcoxy en C₁-C₆, halogéno-alkyle en C₁-C₆, halogéno-alcoxy en C₁-C₆, alkyl-thio en C₁-C₆ et C(O)O(alkyle en C₁-C₆), ou dont deux substituants adjacents forment ensemble un chaînon -O(CH₂)_nO- où l'indice n vaut 1 ou 2 ;

b)




dans laquelle X représente un atome de fluor, de chlore, de brome ou d'iode, Y¹ représente un atome d'hydrogène, de chlore, de brome ou d'iode, étant entendu que si X représente un atome de chlore, Y¹ représente un atome d'hydrogène, de brome ou d'iode, et R¹ représente un groupe alkyle en C₁-C₁₂, ou un groupe aryl-alkyle en C₇-C₁₁ avec ou sans substituant(s), lequel groupe aryl-alkyle en C₇-C₁₁, s'il est porteur de substituant(s), comprend un groupe phényle qui porte un ou plusieurs substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes nitro, cyano, alkyle en C₁-C₆, alcoxy en C₁-C₆, halogéno-alkyle en C₁-C₆,

halogéno-alcoxy en C₁-C₆, alkyl-thio en C₁-C₆ et C(O)O(alkyle en C₁-C₆), ou dont deux substituants adjacents forment ensemble un chaînon -O(CH₂)_nO- où l'indice n vaut 1 ou 2, et étant entendu que le composé ne présente pas la structure suivante :




c)



dans laquelle Y² représente un atome d'hydrogène, R représente un groupe alkyle en C₁-C₄, cyclopropyle, alcényle en C₂-C₄, ou phényle porteur de 1 à 4 substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes alkyle en C₁-C₄, halogéno-alkyle en C₁-C₄, alcoxy en C₁-C₄ et halogéno-alcoxy en C₁-C₄, et R¹ représente un groupe alkyle en C₁-C₁₂, ou un groupe aryl-alkyle en C₇-C₁₁ avec ou sans substituant(s), lequel groupe aryl-alkyle en C₇-C₁₁, s'il est porteur de substituant(s), comprend un groupe phényle qui porte un ou plusieurs substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes nitro, cyano, alkyle en C₁-C₆, alcoxy en C₁-C₆, halogéno-alkyle en C₁-C₆, halogéno-alcoxy en C₁-C₆, alkylthio en C₁-C₆ et C(O)O(alkyle en C₁-C₆), ou dont deux substituants adjacents forment ensemble un chaînon -O(CH₂)_nO- où l'indice n vaut 1 ou 2 ;

d) et



dans laquelle R représente un groupe alkyle en C₁-C₄, cyclopropyle, alcényle en C₂-C₄, ou phényle porteur de 1 à 4 substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes alkyle en C₁-C₄, halogéno-alkyle en C₁-C₄, alcoxy en C₁-C₄ et halogéno-alcoxy en C₁-C₄, et R¹ représente un groupe alkyle en C₁-C₁₂, ou un groupe aryl-alkyle en C₇-C₁₁ avec ou sans substituant(s), lequel groupe aryl-alkyle en C₇-C₁₁, s'il est porteur de substituant(s), comprend un groupe phényle qui porte un ou plusieurs substituant(s) choisi(s) indépendamment parmi les atomes d'halogène et les groupes nitro, cyano, alkyle en C₁-C₆, alcoxy en C₁-C₆, halogéno-alkyle en C₁-C₆, halogéno-alcoxy en C₁-C₆, alkyl-thio en C₁-C₆ et C(O)O(alkyle en C₁-C₆), ou dont deux substituants adjacents forment ensemble un chaînon -O(CH₂)_nO- où l'indice n vaut 1 ou 2.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 6297197 B1 [0002]
- US 6784137 B2 [0002] [0015]
- US 7314849 B2 [0002]
- US 7432227 B2 [0002]
- WO 2011144891 A1 [0002]
- US 20090088322 A1 [0002]
- WO 2006062979 A1 [0002]

Non-patent literature cited in the description

- The Manipulation of Air-Sensitive Compounds. Wiley-Interscience, 1986 [0030]
- Protective Groups in Organic Synthesis. Wiley-Interscience, 1999 [0031]