



EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
21.06.2017 Bulletin 2017/25

(51) Int Cl.:
G01N 33/564 ^(2006.01) **C07K 16/18** ^(2006.01)
C12N 15/09 ^(2006.01) **G01N 33/53** ^(2006.01)

(21) Application number: **12853028.4**

(86) International application number:
PCT/JP2012/079776

(22) Date of filing: **16.11.2012**

(87) International publication number:
WO 2013/080811 (06.06.2013 Gazette 2013/23)

(54) **BIOMARKER FOR LYMPHOCYTIC INFUNDIBULONEUROHYPOPHYSITIS, AND USE
APPLICATIONS THEREOF**

BIOMARKER FÜR LYMPHOZYTISCHE INFUNDIBULONEUROHYPOPHYSITIS UND SEINE
VERWENDUNG

BIOMARQUEUR POUR UNE INFUNDIBULO-NEURO HYPOPHYSITE LYMPHOCYTAIRE ET
APPLICATIONS D'UTILISATION ASSOCIÉES

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **28.11.2011 JP 2011258387**

(43) Date of publication of application:
08.10.2014 Bulletin 2014/41

(73) Proprietor: **National University Corporation
Nagoya University
Nagoya-shi, Aichi 464-8601 (JP)**

(72) Inventors:
• **SUGIMURA, Yoshihisa
Nagoya-shi
Aichi 464-8601 (JP)**
• **OISO, Yutaka
Nagoya-shi
Aichi 464-8601 (JP)**
• **IWAMA, Shintaro
Nagoya-shi
Aichi 464-8601 (JP)**
• **ENOMOTO, Atsushi
Nagoya-shi
Aichi 464-8601 (JP)**
• **KATO, Takuya
Nagoya-shi
Aichi 464-8601 (JP)**

(74) Representative: **Steinecke, Peter
Müller Fottner Steinecke
Rechtsanwalts- und Patentanwaltspartnerschaft
mbB
Römerstrasse 16 b
52428 Jülich (DE)**

(56) References cited:
• **JINN-SHIUN CHEN ET AL: "Detection of
autoantibodies against Rabphilin-3A-like protein
as a potential biomarker in patient's sera of
colorectal cancer", CLINICA CHIMICA ACTA,
ELSEVIER BV, AMSTERDAM, NL, vol. 412, no. 15,
12 April 2011 (2011-04-12), pages 1417-1422,
XP028218491, ISSN: 0009-8981, DOI:
10.1016/J.CCA.2011.04.020 [retrieved on
2011-04-19]**
• **HIRAI H ET AL: "Selective screening of secretory
vesicle-associated proteins for autoantigens in
type 1 diabetes: VAMP2 and NPY are new minor
autoantigens", CLINICAL IMMUNOLOGY,
ACADEMIC PRESS, US, vol. 127, no. 3, 1 June
2008 (2008-06-01), pages 366-374, XP022673823,
ISSN: 1521-6616, DOI:
10.1016/J.CLIM.2008.01.018 [retrieved on
2008-03-24]**
• **"Clinical features and diagnosis
of "lymphocytic" infundibulo-hypophysitis",
FOLIA ENDOCRINOLOGICA JAPONICA, vol. 88,
no. Supplement-2, 1 August 2012 (2012-08-01),
pages 68-78, XP055194001, ISSN: 0029-0661,
DOI: 10.1507/endocrine.88.Supplement-2_68**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- | | |
|---|--|
| <ul style="list-style-type: none"> • SHINTARO IWAMA ET AL.: 'Proteomics ni yoru Lymphocyte-sei Roto Kasuitai Koyoen Shinki Jiko Kogen no Dotei to sono Shikkan Tokusei ni Kansuru Kento, O3-8-17' FOLIA ENDOCRINOLOGICA JAPONICA vol. 87, no. 1, 01 April 2011, page 295, XP008173898 • YOSHIHISA SUGIMURA: 'Kasuitai Koyo Hormone no Saizensen II. Nyohosho Oyobi Kanren Shikkan Lymphocytic infundibuloneurohypophysitis' HORMONE TO RINSHO vol. 59, no. 5, 01 May 2011, pages 471 - 474, XP008174040 | <ul style="list-style-type: none"> • YUTAKA OISO ET AL.: 'Kanno Kasuitai Kino Shogai ni Kansuru Chosa Kenkyu Lymphocyte-sei Roto Kasuitai Koyoen no Shindan Marker to shite no Ko 76kD Tanpaku Kotai no Yuyosei no Kensho to, ES-AVP Saibo Baiyokei o Mochiita 76kD Tanpaku no AVP Bunpi eno Kan'yo no Kaiseki' KANNO KASUITAI KINO SHOGAI NI KANSURU CHOSA KENKYU HEISEI 23 NENDO SOKATSU ? BUNTAN KENKYU HOKOKUSHO March 2012, pages 141 - 146, XP008174034 |
|---|--|

Description

TECHNICAL FIELD

5 **[0001]** The present invention relates to a biomarker for lymphocytic infundibuloneurohypophysitis, and more specifically to a biomarker for lymphocytic infundibuloneurohypophysitis and an assay using the biomarker. This application is based upon and claims the benefit of priority from prior Japanese Patent Application No. 2011-258387, filed November 28, 2011.

BACKGROUND ART

10 **[0002]** Lymphocytic hypophysitis is an autoimmune chronic inflammatory disease caused by infiltration of lymphocytes and plasmacytes into the hypothalamic infundibulum and pituitary. Lymphocytic infundibuloneurohypophysitis (hereinafter abbreviated as "LINH") is a disease accompanied by inflammation limited to infundibulum and posterior lobe, and shows central diabetes insipidus. The diagnosis of LINH is sometimes difficult, and its definite diagnosis requires biopsy
 15 of the pituitary. However, such diagnosis is invasive and thus is scarcely carried out, and not a few subjects are erroneously diagnosed and subjected to needless surgery. The etiology of the disease is unknown, and there is no established therapy. **[0003]** Some diagnostic markers (biomarkers) for lymphocytic hypophysitis are reported (for example, see Non-Patent Literatures 1 to 5), but none of them is clinically applied because none of them combines high sensitivity and high
 20 specificity. In addition, there is no report on the diagnosis marker specific to LINH.

PRIOR ART DOCUMENT

NON-PATENT DOCUMENT

25 **[0004]**

Non-patent Literature 1: J Clin Endocrinol Metab. 87 (2): 752-7, 2002

Non-patent Literature 2: Horm Res. 55 (6): 288-92, 2001

Non-patent Literature 3: Eur J Endocrinol. 147 (6): 767-75, 2002

30 Non-patent Literature 4: J Endocrinol Invest, 30 (2): 153-62, 2007

Non-patent Literature 5: Clin Endocrinol. 69 (2): 269-78, 2008

SUMMARY OF THE INVENTION

35 PROBLEM TO BE SOLVED BY THE INVENTION

[0005] In consideration of the above-described background, the present invention is intended to provide an LINH-specific biomarker and applications thereof, more specifically the technique useful for the examination of LINH, thereby allowing the discrimination of LINH and highly confident diagnosis. Specifically, the purpose of the present invention is
 40 to provide a biomarker useful for the examination of LINH, an assay using the biomarker, and a test reagent used in the assay.

MEANS FOR SOLVING PROBLEM

45 **[0006]** The inventors made investigations for finding an LINH-specific biomarker. Specifically, firstly, serums were collected from LINH patients, disease control subjects showing central diabetes insipidus because of, for example, cerebral tumor, and healthy subjects, and IgG was purified. On the other hand, an extract of posterior pituitary protein was prepared from rat pituitary. The IgG purified from the serum and the extract of posterior pituitary protein was mixed, and subjected to immunoprecipitation using protein G beads, and then the antigen was eluted from the immune precipitate.
 50 The eluted antigen was subjected to analysis (shotgun proteomics) using an LC-MS/MS system, Pradigm MS4-PAL-LTQ Orbitrap XL (Thermo Fisher Scientific Inc.). The MS data was analyzed using a public database and an analysis software, and the potential autoantigen protein was identified. Secondly, recombinant proteins were synthesized for six proteins as autoantigen protein candidates, and then the reactivity between these recombinant proteins and the serums of the patients was studied by the Western blot method. As a result of this, an autoantibody against rabphilin 3a was
 55 found in eight subjects of the nine LINH patients (8/9). On the other hand, the cases wherein the autoantibody was found in the control were 1/10 for the healthy subjects, 0/3 for the subjects with lymphocytic adenohypophysitis, 1/3 for the subjects with IgG4-related hypophysitis accompanied by diabetes insipidus, and 1/10 for the subjects with diabetes insipidus accompanying tumor and others. Accordingly, rabphilin 3a was proved to be an autoantigen having high

specificity to LINH. In other words, it has been proved that the autoantibody (anti-rabphilin 3a antibody) against rabphilin 3a is markedly useful as a biomarker for LINH, and that the use of the antibody as the indicator allows the examination of LINH (for example, examination for discrimination) with markedly high sensitivity and high specificity.

[0007] As a result of further study, it was found that rabphilin 3a is expressed also in the rat hypothalamic paraventricular nuclei, supraoptic nuclei, and posterior pituitary. This finding suggests that rabphilin 3a is directly involved in the pathogenesis of LINH, and proved that the autoantibody against the molecule, more specifically the anti-rabphilin 3a antibody is useful for grasping the disease state of LINH.

[0008] As a result of study using more specimens, markedly high specificity was found in the discrimination between diabetes insipidus (DI) accompanying to tumor lesion and LINH, and the discrimination between idiopathic or secondary central diabetes insipidus and LINH, in addition, high specificity was found also in the discrimination of other autoimmune disease and LINH. Therefore, it was further corroborated that the anti-rabphilin 3a antibody is useful as a biomarker for LINH. The following aspects of the present invention are mainly based on the above-described facts.

[1] A lymphocytic infundibuloneurohypophysitis marker including an anti-rabphilin 3a antibody.

[2] An assay for lymphocytic infundibuloneurohypophysitis, using the level of an anti-rabphilin 3a antibody in a specimen as an indicator.

[3] The assay according to [2], which includes the following steps (1) to (3):

(1) a step of providing a specimen derived from a subject;

(2) a step of detecting an anti-rabphilin 3a antibody in the specimen; and

(3) a step of determining whether the developed disease is lymphocytic infundibuloneurohypophysitis or not, or the possibility of developing lymphocytic infundibuloneurohypophysitis, based on the detection result.

[4] The assay according to [3], wherein the high level of the anti-rabphilin 3a antibody is used as the indicator showing that the developed disease is lymphocytic infundibuloneurohypophysitis, and as the indicator showing that the possibility of developing lymphocytic infundibuloneurohypophysitis is high.

[5] The assay according to [3] or [4], wherein the determination in the step (3) is carried out based on the comparison between the detected value obtained in the step (2) and the detected value obtained in the control specimen.

[6] The assay according to [3] or [4], wherein the determination in the step (3) is carried out based on the comparison between the detected value obtained in the step (2) and the detected value in the specimen taken from the same subject in the past.

[7] The assay according to any of [2] to [6], wherein the specimen is blood, plasma, serum, cerebrospinal fluid, or urine.

[8] A test reagent for lymphocytic infundibuloneurohypophysitis, including rabphilin 3a or its antibody-binding fragment.

[9] A kit for testing lymphocytic infundibuloneurohypophysitis, including the lymphocytic infundibuloneurohypophysitis test reagent according to [8].

BRIEF DESCRIPTION OF DRAWING

[0009]

Fig. 1 shows the result of a Western blot test using the blood serums of the patients and rabphilin 3a. Reaction to rabphilin 3a was found in the LINH patients (subjects 1 to 3). On the other hand, similar reaction was not found in secondary diabetes insipidus (DI) or healthy subjects.

Fig. 2 shows the summary of the autoantibody positive rate for rabphilin 3a. The proportion of the positive subjects was shown for each disease. For example, 8/9 indicates that eight of nine subjects were autoantibody-positive.

Fig. 3 shows the reactivity between the HEK293FT cells which had been transfected with rabphilin 3a and the blood serums of LINH patients (result of immunohistochemistry). The reactivity to an anti-V5 antibody (left), reactivity to the blood serums (center), and the merge of two images (right) are shown.

Fig. 4 shows the expression of an LINH-specific antigen. The immunohistochemistry of rat posterior pituitary (upper row), and hypothalamic supraoptic nucleus (SON) (lower row). The localization of AVP (left) and rabphilin 3a (center) was studied. The right is the merge of two images.

Fig. 5 shows the summary of the autoantibody positive rate for rabphilin 3a. Using more specimens, the usefulness of rabphilin 3a was studied. The autoantibody positive rate was studied also for the specimens of autoimmune disease patients.

Fig. 6 shows the summary of the sensitivity and specificity of discrimination.

DESCRIPTION OF EMBODIMENT

(First aspect of the present invention: biomarker for lymphocytic infundibuloneurohypophysitis (LINH))

[0010] A first aspect of the present invention relates to a biomarker for LINH (hereinafter may be referred to as "the biomarker of the present invention"). The biomarker of the present invention is a useful indicator for the discrimination of LINH, or for evaluating the possibility of developing LINH. The term "discrimination" herein means the determination whether the disease developed in the subject is LINH or other disease. Especially, the biomarker of the present invention is useful for ascertaining whether the patient with diabetes insipidus has LINH or not. In particular, the biomarker of the present invention is important as an indicator for discriminating central diabetes insipidus accompanying LINH and diabetes insipidus accompanying tumor lesion.

[0011] "Possibility of developing LINH" includes current and future possibilities of developing LINH. The "current possibility of developing LINH" represents whether LINH is developed or not, or the probability of developing LINH at the time of assay. On the other hand, "future possibility of developing LINH" represents the future possibility (risk) of developing LINH.

[0012] The biomarker of the present invention is composed of an antibody (autoantibody) against an LINH-specific antigen molecule rabphilin 3a, more specifically an anti-rabphilin 3a antibody. Rabphilin 3a is a protein coded by an RPH3A gene. Rabphilin 3a is an effector of RAB3A, and known to participate in the exocytosis of neurotransmitters and hormones. The amino acid sequence of rabphilin 3a (DEFINITION: Homo sapiens rabphilin 3A homolog (mouse), mRNA (cDNA clone MGC: 29559 IMAGE: 3510158), complete cds. ACCESSION: BC017259) and the nucleotide sequence coding the amino acid sequence are set forth in SEQ ID NO. 1 and SEQ ID NO. 2 in the sequence list, respectively. As a general rule, the biomarker of the present invention is not the anti-rabphilin 3a antibody existing in a living body, but the anti-rabphilin 3a antibody in a specimen taken from (or separated from) a living body.

(Second aspect of the present invention: LINH assay)

[0013] A second aspect of the present invention relates to a use of the biomarker of the present invention, and provides a method for testing whether the developed disease is LINH or not, or the possibility of developing LINH (hereinafter may be referred to as "the assay of the present invention"). The assay of the present invention allows the discrimination of LINH (when determining whether the developed disease is LINH or not). The assay of the present invention is useful as a means for determining whether LINH is developed not at present, or a means for evaluating the possibility of developing LINH in future (when the possibility of developing LINH is determined). Thus, the assay of the present invention provides useful information for the diagnosis of LINH. The assay of the present invention is, for example, useful for simple and objective definite diagnosis of LINH.

[0014] In the assay of the present invention, the level of the biomarker of the present invention in the specimen derived from the subject is used as the indicator. The "level" herein typically means "amount" or "concentration". However, according to convention and technical common knowledge, the term "level" is used also when showing whether the biomarker of the present invention can be detected or not (more specifically the presence or absence of apparent presence).

[0015] The assay of the present invention includes the following steps:

- (1) a step of providing a specimen derived from a subject;
- (2) a step of detecting an anti-rabphilin 3a antibody in the specimen;
- (3) a step of determining whether the developed disease is lymphocytic infundibuloneurohypophysitis or not, or the possibility of developing lymphocytic infundibuloneurohypophysitis, based on the detection result.

[0016] In the step (1), a specimen derived from the subject is provided. The specimen may be, for example, blood, plasma, serum, cerebrospinal fluid, or urine of the subject. The subject is not particularly limited. More specifically, the present invention is widely applicable to the subject requiring the discrimination of LINH, and the subject requiring the determination of the possibility of developing LINH at present or in future (more specifically, whether LINH is developed or not, the degree of possibility of having LINH, and the degree of possibility of developing LINH in future). For example, the subjects having the symptom of central diabetes insipidus correspond to "the subject requiring the discrimination of LINH". In addition, when the present invention is applied to the patient who has been diagnosed to have LINH based on health questionnaire or other test by physician, the propriety of the diagnosis can be determined based on an objective indicator, or the level of the biomarker. More specifically, according to the assay of the present invention, information aiding or supporting the prior art diagnosis can be obtained. This information is useful for the determination of more appropriate course of treatment, and promotes the improvement of therapeutic effect and the patient's QOL (quality of life). On the other hand, the present invention may be used for monitoring the disease state, thereby preventing the

increase of the difficulty of treatment, seriousness, and recurrence.

[0017] The subjects in whom hyperplasia of the pituitary stalk or pituitary enlargement is found by image examination such as MRI examination are presumed to have high risk of developing LINH (high risk subjects), and are suitable subjects. The application of the present invention to these subjects before the development of the symptom of LINH allows prevention or delay of the development, or therapeutic intervention in the early stage. The present invention is useful for specifying the subjects having high risk of developing LINH. Such specification allows the decrease of possibility of developing LINH (the possibility of having LINH) by, for example, precautions or improvement of lifestyle habits. Those having no subjective symptom, who can't be or is hard to be determined whether having LINH or not by prior art diagnosis, are also preferred subjects of the present invention. The present invention may be carried out as an item of health examination.

[0018] In the step (2), the biomarker in the specimen is detected. It is not essential to strictly determine the biomarker level. More specifically, the biomarker level is detected so as to allow the determination in the subsequent step (3). For example, the detection may be carried out so as to allow the determination whether the biomarker level in the specimen exceeds the specified value.

[0019] The method for detecting the biomarker is not particularly limited, but is preferably an immunological method. Immunological methods allow prompt and highly sensitive detection. In addition, the operation is simple. The measurement by an immunological method uses a substance which specifically binds to the biomarker. As the substance, in typical cases, rabphilin 3a or its antibody-binding fragment is used. The "antibody-binding fragment" is a fragment of rabphilin 3a containing the epitope binding to the anti-rabphilin 3a antibody, and binds to the anti-rabphilin 3a antibody in the same manner as full length rabphilin 3a.

[0020] Examples of the measurement method include latex agglutination assay, fluoroimmunoassay (FIA method), enzyme immunoassay (EIA method), radioimmunoassay method (RIA method), and Western blot method. Examples of preferred measurement methods include the FIA method and EIA method (including ELISA method). These methods allow highly sensitive, prompt, and simple detection. The FIA method uses a fluorescent-labeled antibody, and detects an antigen-antibody complex (immune complex) using fluorescence as the signal. On the other hand, the EIA method uses an enzyme-labeled antibody, and detects an immune complex using the color development or luminescence based on enzyme reaction as the signal.

[0021] The ELISA method has many advantages such as high detection sensitivity, high specificity, marked quantitative performance, simple operation, and suitability for simultaneous treatment of many specimens. An example of the specific operation method for using the ELISA method is shown below. Firstly, the anti-rabphilin 3a antibody prepared by immunizing a rabbit or goat is immobilized on an insoluble support. Specifically, for example, the surface of a microplate is immunized (coated) by an anti-rabphilin 3a antibody. The antibody thus solid-phased is brought into contact with the specimen and enzyme-labeled rabphilin 3a. As a result of this operation, the solid-phased antibody binds to the rabphilin 3a to form an immune complex. However, if the anti-rabphilin 3a antibody (autoantibody) is present in the specimen, it competes with the solid-phased antibody, whereby the bonding amount between the solid-phased antibody and rabphilin 3a decreases. After removing the non-specific binding component by washing operation, color is developed by the reaction with a substrate of the enzyme. Subsequently, the immune complex is detected using the amount of color development as the indicator. Details about the ELISA method are described in many publications and literatures, and they may be referred when the experimental method and experimental condition are established. The method may be competitive or non-competitive. Alternatively, a method including contacting the specimen with the solid-phased rabphilin 3a (or its antibody-binding fragment) to form an immune complex, and then detecting the immune complex using a secondary antibody or the like, or a sandwich method may be used.

[0022] Alternatively, a means such as a protein array or protein chip which can detect many specimens simultaneously may be used.

[0023] In the step (3), whether the developed disease is LINH or not is determined based on the detection result. In other embodiment, the possibility of developing LINH is determined based on the detection result. In order to allow highly accurate determination, it is preferred that the determination be carried out after comparing the detected value obtained in the step (2) with the detected value of the control specimen (control). The control specimen may be, for example, a specimen from a healthy subject, a specimen from a patient with diabetes insipidus accompanying tumor lesion, a specimen from a patient with idiopathic or secondary central diabetes insipidus, or a specimen from a patient with other autoimmune disease. The determination may be qualitative or quantitative. As is apparent from the criteria, the determination herein may be carried out automatically/mechanically without depending on the judgment by a specialist such as a physician or laboratory technician.

(Example of qualitative determination 1: discrimination)

[0024] The determination that "the developed disease is LINH" is made when the detected value (for example, labeled amount or fluorescence amount) is higher than the standard value, and the determination that "the developed disease

is not LINH" is made when the detected value is lower than the standard value. When discrimination from a specific disease is made, in typical cases, the determination that "the developed disease is LINH" is made when the detected value is higher than the standard value, and the determination that "the developed disease is the specific disease" is made when the detected value is lower than the standard value. The "specific disease" herein means, for example, lymphocytic adenohypophysitis, IgG4-related hypophysitis accompanied by diabetes insipidus, and tumor lesion accompanied by diabetes insipidus (the same applies to the following examples).

(Example of qualitative determination 2: discrimination)

[0025] The determination that "the developed disease is LINH" is made when reactivity is found (positive), and the determination that "the developed disease is not LINH" is made when no reactivity is found (negative). When discrimination from a specific disease is made, in typical cases, the determination that "the developed disease is LINH" is made when reactivity is found (positive), and the determination that "the developed disease is the specific disease" is made when no reactivity is found (negative).

(Example of qualitative determination 3: possibility of developing LINH)

[0026] The determination that "the possibility of having LINH is high" is made when the detected value (for example, labeled amount or fluorescence amount) is higher than the standard value, and the determination that "LINH is not developed" or "the possibility of not having LINH is high" is made when the detected value is lower than the standard value. When determining the possibility of developing the disease in future, the determination that "LINH will be developed" or "the possibility of developing LINH is high" is made when the detected value is higher than the standard value, and the determination that "LINH will not be developed" or "the possibility of not developing LINH is high" is made when the detected value is lower than the standard value.

(Example of qualitative determination 4: possibility of developing LINH)

[0027] The determination that "LINH is developed" or "the possibility of having LINH is high" is made when reactivity is found (positive), and the determination that "LINH is not developed" or "the possibility of not having LINH is high" is made when no reactivity is found (negative). When determining the possibility of developing the disease in future, the determination that "LINH will be developed" or "the possibility of developing LINH is high" is made when reactivity is found (positive), and the determination that "LINH will not be developed" or "the possibility of not developing LINH is high" is made when no reactivity is found (negative).

(Example of qualitative determination 1: discrimination)

[0028] As shown below, "the possibility that the developed disease is LINH (%)" is established for each range of the detected value, and "the possibility that the developed disease is LINH (%)" is determined from the detected value.

Detected values a to b: the possibility of LINH is 10% or less
 Detected values b to c: the possibility of LINH is from 10% to 30%
 Detected values c to d: the possibility of LINH is from 30% to 50%
 Detected values d to e: the possibility of LINH is from 50% to 70%
 Detected values e to f: the possibility of LINH is from 70% to 90%

(Example of qualitative determination 2: possibility of developing LINH)

[0029] As shown below, "the possibility of developing LINH (%)" is established for each range of the measured value, and "the possibility of developing LINH (%)" is determined from the measured value.

Measured values a to b: the possibility of developing LINH is 10% or less
 Measured values b to c: the possibility of developing LINH is from 10% to 30%
 Measured values c to d: the possibility of developing LINH is from 30% to 50%
 Measured values d to e: the possibility of developing LINH is from 50% to 70%
 Measured values e to f: the possibility of developing LINH is from 70% to 90%

[0030] The number of determination categories, and the biomarker level and determination result associated with each determination category will not be limited to the above examples, and may be freely established based on, for example,

preliminary experiments. For example, the "threshold" when making the determination (discrimination) whether LINH or not using a predetermined threshold as the boundary or determining the level of the possibility of developing LINH, and the "biomarker level range" associated with the category according to the determination (discrimination) whether LINH or not, or the level of the possibility of developing LINH may be established based on statistical analysis using many specimens.

[0031] According to one embodiment of the present invention, the biomarker level measured at one point is compared with the biomarker level measured in the past in the same subject, thereby determining the presence or absence of the fluctuations and/or the degree of fluctuations in the biomarker level. The data thus obtained regarding the change in the biomarker level is useful information for monitoring the presence or absence of the development of LINH or the possibility of developing LINH, grasping the therapeutic effect, or estimating prognosis. Specifically, for example, on the basis of the fluctuation in the biomarker level, the increase, decrease, or no change in the possibility of developing LINH from the last test to the present test can be determined. If this evaluation is carried out in parallel with the treatment of LINH, the therapeutic effect can be confirmed, and the sign of recurrence of LINH can be grasped in advance. This allows the determination of more suitable course of treatment. In this manner, the present invention will markedly contribute to the maximization of the therapeutic effect and the improvement of patient's QOL (quality of life).

(Third aspect of the present invention: reagent and kit for testing the possibility of developing LINH)

[0032] The present invention further provides a reagent and kit used for the assay of LINH. The reagent of the present invention includes rabphilin 3a or its antibody-binding fragment (hereinafter referred to as "active ingredient"), which is a substance showing specific binding properties to the biomarker of the present invention. The active ingredient of the reagent of the present invention can be prepared in the form of, for example, a recombinant protein. For example, it can be prepared by transforming an appropriate host cell with a DNA which codes the target active ingredient (rabphilin 3a or its antibody-binding fragment), and collecting the protein expressed in the transformant. The collected protein is purified as appropriate according to the intended use. In this manner, various modifications can be made by obtaining a binding molecule as a recombinant protein. For example, the active ingredient composed of a recombinant protein linked to any desired peptide can be obtained by inserting a DNA coding a binding molecule and other appropriate DNA into the same vector, and producing a recombinant protein using the vector. Alternatively, modification which causes the addition of a sugar chain and/or lipid, or processing of N-terminal or C-terminal may be made. These modifications allow extraction of a recombinant protein, simplification of purification, or addition of biological function. The method for preparing the active ingredient is not limited to genetic engineering method. For example, the active ingredient of the reagent of the present invention may be prepared from a natural material using a standard method (for example, fracturing, extraction, or purification).

[0033] The active ingredient may be prepared using a cell-free synthesis system. The cell-free synthesis system (cell-free transcription system, cell-free transcription/translation system) synthesizes an mRNA or protein in vitro from the nucleic acid (DNA or mRNA) template coding for it, using not living cells but the ribosome or transcription/translation factor derived from living cells (or obtained by a genetic engineering method). The cell-free synthesis system commonly uses a cell extract obtained by purifying a cell lysate as necessary. The cell extract commonly contains ribosome, various factors such as an initiation factor, and various enzymes such as tRNA necessary for protein synthesis. When protein synthesis is carried out, other substances necessary for protein synthesis, such as energy sources including various amino acids, ATP, and GTP, and creatine phosphate are added to the cell extract. As a matter of course, during the protein synthesis, ribosome prepared separately, various factors, and/or various enzymes may be added as necessary.

[0034] Development of a transcription/translation system constructed by reconstituting various molecules (factors) necessary for protein synthesis was reported (Shimizu, Y. et al.: Nature Biotech., 19, 751-755, 2001). In this synthesis system, the genes of 31 factors composing the protein synthesis system of a bacterium, which includes three initiation factors, three elongation factors, four factors involved with termination, 20 aminoacyl-tRNA synthetases which make an amino acid bind to tRNA, and methionyl tRNA formyl transferase are amplified from Escherichia coli, and a protein synthesis system is reconstituted using them in vitro. The present invention may use such a reconstituted synthesis system.

[0035] The cell-free protein synthesis system has the following advantages. Firstly, the system does not require the maintenance of living cells, so that provides good operability and a high degree of freedom. Accordingly, synthesis systems variously corrected and modified according to the properties of the intended protein can be designed. Secondly, a cell-based synthesis system basically cannot synthesize a protein toxic to the cells contained in the system, but a cell-free system can produce such a toxic protein. Furthermore, many proteins can be synthesized simultaneously and quickly, which facilitates high throughput. This system has an advantage that it allows easy separation and purification of produced proteins, which is advantageous for high throughput. The system also has an advantage that it can synthesize nonnatural proteins by the uptake of nonnatural amino acids.

[0036] Examples of the cell-free protein synthesis system widely used at present include the followings. More specif-

ically, an Escherichia coli S30 extract system (procaryotic cell system), a wheat germ extract system (eucaryotic cell system), and a solubilized rabbit reticulocyte system (eucaryotic cell system). These systems are commercially available in the form of kits, and can be easily used.

[0037] The wheat germ extract system has an advantage that it can efficiently synthesizes high quality eucaryotic proteins, and is often used for synthesizing eukaryotic proteins which are difficult to be synthesized by the Escherichia coli S30 extract system. Recently, it was reported that a highly efficient and stable synthesis system is constructed by preparing an extract from the germ from which the seed albumen component has been removed by washing, and receives widespread attention (Madin, K. et al.: Proc. Natl. Acad. Sci. USA, 97: 559-564, 2000). Thereafter, techniques such as an untranslated mRNA sequence having high translation promotion capacity, a protein synthesis method using PCR for multi-item function analysis, and construction of special high expression vectors are developed (Sawasaki, T. et al.: Proc. Natl. Acad. Sci. USA, 99: 14652-14657, 2002), and are expected to be applied to various fields.

[0038] The wheat germ extract can be obtained by centrifuging ground wheat germ, and then separating the supernatant liquid by gel filtration. The translation reaction may refer to the method of Anderson et al. (Anderson, C. W. et al.: Methods Enzymol., 101, 638-644 (1983)). Improved methods are also reported, and for example, the method of Kawarasaki et al. (Kawarasaki, Y. et al.: Biotechnol. Prog., 16, 517-521 (2000)) and the method of Madin et al. (Madin, K. et al.: Proc. Natl. Acad. Sci. USA, 97: 559-564, 2000) may be referred to. For the wheat germ extract system, refer to WO 00/68412 A1, WO 01/27260 A1, WO 2002/024939 A1, WO 2005/063979 A1, Japanese Unexamined Patent Application Publication No. 6-7134, Japanese Unexamined Patent Application Publication No. 2002-529531, Japanese Unexamined Patent Application Publication No. 2005-355513, Japanese Unexamined Patent Application Publication No. 2006-042601, Japanese Unexamined Patent Application Publication No. 2007-097438, and Japanese Unexamined Patent Application Publication No. 2008-029203.

[0039] The cell-free synthesis system used for carrying out the present invention is not limited to the above-described one, and may be a system constructed based on, for example, an extract of a bacterium other than Escherichia coli, a plant other than wheat, an extract derived from an insect, an extract from animal cells, or genome information.

[0040] By labeling the active ingredient of the reagent of the present invention, the amount of the bound antibody can be directly detected using the labeled amount as the indicator. This allows the construction of a more simple assay. On the other hand, the use of an indirect detection method, such as a method using a secondary antibody bound to a labeling agent, or a method using a polymer bound to a secondary antibody and a labeling agent is also preferred from the viewpoint of detection sensitivity.

[0041] Examples of the labeling agent include enzymes such as peroxidase, microperoxidase, horseradish peroxidase (HRP), alkaline phosphatase, β -D-galactosidase, glucose oxidase, and glucose-6-phosphate dehydrogenase, fluorescent substances such as fluorescein isothiocyanate (FITC), tetramethylrhodamineisothiocyanate (TRITC), and europium, chemiluminescence substances such as luminol, isoluminol, and acridinium derivatives, coenzymes such as NAD, biotin, and radioactive materials such as ^{131}I and ^{125}I .

[0042] According to one embodiment, the reagent of the present invention is solid-phased according to the use application. The insoluble support used for making the solid phase is not particularly limited. For example, the insoluble support may be composed of a water-insoluble substance, such as a resin including a polystyrene resin, a polycarbonate resin, a silicon resin, a nylon resin, and glass. Loading on the insoluble support may be achieved by physical adsorption or chemical adsorption.

[0043] The kit of the present invention includes the reagent of the present invention as a main component. The kit may further include other reagent used for carrying out the assay (for example, a buffer solution, a blocking reagent, an enzyme substrate, and a color-producing reagent) and/or an apparatus or instrument (for example, a container, a reaction apparatus, and a fluorescence reader). In addition, the kit preferably includes the active ingredient of the reagent of the present invention (rabphilin 3a or its antibody-binding fragment) as the standard sample. In addition, usually, an instruction manual is attached to the kit of the present invention.

EXAMPLE

1. Search of LINH-specific antigen

[0044] The blood serums of LINH patients, the blood serums of disease control subjects with diabetes insipidus accompanying cerebral tumor and others, and the blood serums of healthy subjects were collected, and IgG was purified by ordinary procedure. On the other hand, an extract of posterior pituitary protein was purified from rat pituitary. The purified IgG was mixed with the extract of posterior pituitary protein, the mixture was subjected to immunoprecipitation using protein G beads, and then an antigen was eluted from the immune precipitate. The eluted antigen was reduced, alkylated, and then subjected to trypsin digestion (in-solution digestion). The sample thus prepared was subjected to analysis (shotgun proteomics) using an LC-MS/MS system, Pradigm MS4-PAL-LTQ Orbitrap XL (Thermo Fisher Scientific Inc.). The MS data thus obtained was analyzed based on NCBI RefSeq and Swiss Prot database, using Mascot (Matrix

Science Ltd.) software. As a result of the analysis, six autoantigen protein candidates were identified.

2. Reactivity between LINH-specific antigen and blood serums of patients

(1) Western blot

[0045] Subsequently, six proteins as autoantigen protein candidates were subcloned into an expression vector (pcDNA3.1D/V5-His-TOPO (registered trademark)). Thereafter, the expression vector was transfected into a mammal cell line (HEK293FT) using Lipofectamine 2000 (registered trademark), thereby synthesizing recombinant proteins. Six recombinant proteins of autoantigen candidates synthesized using the HEK293FT cells were separated by SDS-polyacrylamide electrophoresis (PAGE), and the reactivity between the blood serums of the patients and recombinant proteins was studied by the Western blot using the blood serums of the patients as the primary antibodies. The blood serums used in the study are as follows:

- (a) LINH 9 subjects
- (b) Lymphocytic adenohypophysitis (LAH) 3 subjects
- (c) IgG4-related hypophysitis 3 subjects
- (d) Disease control (10 subjects in total)
- Germ cell tumor + diabetes insipidus (DI) 3 subjects
- After surgery of suprasellar pineal tumor + DI 1 subject
- After surgery of hypothalamus tumor + DI 1 subject
- Rathke's cleft cyst + DI 1 subject
- Craniopharyngioma + DI 2 subjects
- After surgery of glioma + DI 1 subject
- Sarcoidosis + DI 1 subject
- (e) Healthy control (healthy subject) 10 subjects

[0046] The results of the Western blot (representative examples) are shown in Fig. 1. In eight of the nine patients with LINH (8/9), an autoantibody against rabphilin 3a, which is a 76 kDa protein, was found. On the other hand, the subject in whom an autoantibody was found in the control was 1/10 for the healthy subjects, 0/3 for the lymphocytic adenohypophysitis subjects, 1/3 for the subjects with IgG4-related hypophysitis accompanied by diabetes insipidus, and 1/10 for the subjects with diabetes insipidus accompanying tumor (Fig. 2). For all the studied subjects, sensitivity of the autoantibody was 88.9% (8/9), specificity was 87.0% (20/23) (88.5% (23/26) when the subjects with IgG4-related hypophysitis were added), and specificity of 90% (9/10) was shown in the discrimination from the subjects with DI accompanying tumor lesion, which is difficult to be clinically discriminated from LINH (Fig. 2).

(2) Immunohistochemistry

[0047] The reactivity between the HEK293FT cells transfected with rabphilin 3a and the blood serum of an LINH patient was studied by immunohistochemistry. The results are shown in Fig. 3. The LINH blood serum (the highest row) showed reactivity in agreement with the cells recognized by the anti-V5 antibody (co localization). When the blood serum of other LINH patient was used (third row from the top), the same staining property was found. On the other hand, when an empty vector was transfected (the second row from the top), no reaction was found neither of the anti-V5 antibody nor blood serum. When the blood serum of a patient with diabetes insipidus after tumor surgery was used (the lowest row), no colocalization was found, and it was suggested that the antibody against rabphilin 3a is specific to the patients with LINH.

3. Expression of LINH-specific antigen

[0048] The expression of rabphilin 3a in the vasopressin (AVP) neurons in the rat posterior pituitary and hypothalamic supraoptic nucleus (SON) was studied by immunohistochemistry (double staining by an anti-AVP antibody and anti-rabphilin antibody). The result of immunohistochemistry is shown in Fig. 4. The expression of rabphilin 3a was found in the AVP neurons in the posterior pituitary and hypothalamus SON. The fact that the expression of rabphilin 3a was found in the lesion region of LINH strongly suggests that the possibility of involvement of the protein in the disease state of LINH.

4. Summary

[0049] The LINH-specific antigen protein rabphilin 3a was successfully found. The anti-rabphilin 3a antibody is used

as a biomarker for LINH. The use of the anti-rabphilin 3a antibody as the indicator allows the test of LINH with markedly high sensitivity and specificity (for example, discrimination). The anti-rabphilin 3a antibody is also useful for grasping the disease state of LINH.

5 5. Study using more specimens (including specimens of other autoimmune diseases)

10 **[0050]** The type of the specimen and subject was increased, and usefulness of the anti-rabphilin 3a antibody as a biomarker was further studied. The results are shown in FIGs. 5 and 6. The specificity was as high as 92.8% and 95.2% for the discrimination between the diabetes insipidus accompanying tumor lesion and LINH, and the discrimination between idiopathic or secondary central diabetes insipidus and LINH, respectively, and a high specificity (89.9%) was exhibited for the discrimination between various autoimmune diseases and LINH. Accordingly, more specifically, it was supported that the anti-rabphilin 3a antibody is markedly useful as a biomarker for LINH.

INDUSTRIAL APPLICABILITY

15 **[0051]** In the assay of the present invention, discrimination of LINH and others are carried out using the anti-rabphilin 3a antibody as the indicator. According to the assay of the present invention, LINH can be discriminated with markedly high sensitivity and specificity. Accordingly, the assay of the present invention can be regarded as markedly useful for the prevention of erroneous diagnosis, and determination of appropriate course of treatment.

SEQUENCE LISTING

[0052]

25 <110> NATIONAL UNIVERSITY CORPORATION NAGOYA UNIVERSITY
SUGIMURA, Yoshihisa
OISO, Yutaka
IWAMA, Shintaro
ENOMOTO, Atsushi
30 KATO, Takuya

<120> BIOMARKER OF LYMPHOCYTIC INFUNDIBULO-NEUROHYPOPHYSITIS AND USE THEREOF

<130> NU11009P

35 <150> JP P2011-258387
<151> 2011-11-28

<160> 2

40 <170> PatentIn version 3.5

<210> 1

<211> 690

45 <212> PRT

<213> Homo sapiens

<400> 1

50

55

EP 2 787 346 B1

	Met	Thr	Asp	Thr	Val	Phe	Ser	Asn	Ser	Ser	Asn	Arg	Trp	Met	Tyr	Pro	
	1				5					10					15		
5	Ser	Asp	Arg	Pro	Leu	Gln	Ser	Lys	Leu	Gln	Ala	Gly	Trp	Ser	Val	His	
				20					25					30			
10	Pro	Gly	Gly	Gln	Pro	Asp	Arg	Gln	Arg	Lys	Gln	Glu	Glu	Leu	Thr	Asp	
			35					40					45				
15	Glu	Glu	Lys	Glu	Ile	Ile	Asn	Arg	Val	Ile	Ala	Arg	Ala	Glu	Lys	Met	
		50					55					60					
20	Glu	Glu	Met	Glu	Gln	Glu	Arg	Ile	Gly	Arg	Leu	Val	Asp	Arg	Leu	Glu	
	65					70					75					80	
25	Asn	Met	Arg	Lys	Asn	Val	Ala	Gly	Asp	Gly	Val	Asn	Arg	Cys	Ile	Leu	
					85					90					95		
30	Cys	Gly	Glu	Gln	Leu	Gly	Met	Leu	Gly	Ser	Ala	Cys	Val	Val	Cys	Glu	
				100					105					110			
35	Asp	Cys	Lys	Lys	Asn	Val	Cys	Thr	Lys	Cys	Gly	Val	Glu	Thr	Asn	Asn	
			115					120					125				
40	Arg	Leu	His	Ser	Val	Trp	Leu	Cys	Lys	Ile	Cys	Ile	Glu	Gln	Arg	Glu	
	130						135					140					

EP 2 787 346 B1

	Val	Trp	Lys	Arg	Ser	Gly	Ala	Trp	Phe	Phe	Lys	Gly	Phe	Pro	Lys	Gln	
	145					150					155					160	
5	Val	Leu	Pro	Gln	Pro	Met	Pro	Ile	Lys	Lys	Thr	Lys	Pro	Gln	Gln	Pro	
					165					170					175		
	Val	Ser	Glu	Pro	Ala	Ala	Pro	Glu	Gln	Pro	Ala	Pro	Glu	Pro	Lys	His	
10				180					185					190			
	Pro	Ala	Arg	Ala	Pro	Ala	Arg	Gly	Asp	Ser	Glu	Asp	Arg	Arg	Gly	Pro	
			195					200					205				
15	Gly	Gln	Lys	Thr	Gly	Pro	Asp	Pro	Ala	Ser	Ala	Pro	Gly	Arg	Gly	Asn	
	210						215						220				
	Tyr	Gly	Pro	Pro	Val	Arg	Arg	Ala	Ser	Glu	Ala	Arg	Met	Ser	Ser	Ser	
20	225					230					235					240	
	Ser	Arg	Asp	Ser	Glu	Ser	Trp	Asp	His	Ser	Gly	Gly	Ala	Gly	Asp	Ser	
25					245					250					255		
	Ser	Arg	Ser	Pro	Ala	Gly	Leu	Arg	Arg	Ala	Asn	Ser	Val	Gln	Ala	Ser	
				260					265					270			
30	Arg	Pro	Ala	Pro	Gly	Ser	Val	Gln	Ser	Pro	Ala	Pro	Pro	Gln	Pro	Gly	
			275					280						285			
	Gln	Pro	Gly	Thr	Pro	Gly	Gly	Ser	Arg	Pro	Gly	Pro	Gly	Pro	Ala	Gly	
35	290						295					300					
	Arg	Phe	Pro	Asp	Gln	Lys	Pro	Glu	Val	Ala	Pro	Ser	Asp	Pro	Gly	Thr	
	305					310					315					320	
40	Thr	Ala	Pro	Pro	Arg	Glu	Glu	Arg	Thr	Gly	Gly	Val	Gly	Gly	Tyr	Pro	
					325					330					335		
	Ala	Val	Gly	Ala	Arg	Glu	Asp	Arg	Met	Ser	His	Pro	Ser	Gly	Pro	Tyr	
45				340					345					350			
	Ser	Gln	Ala	Ser	Ala	Ala	Ala	Pro	Gln	Pro	Ala	Ala	Ala	Arg	Gln	Pro	
50			355					360					365				
	Pro	Pro	Pro	Glu	Glu	Glu	Glu	Glu	Glu	Ala	Asn	Ser	Tyr	Asp	Ser	Asp	
		370					375					380					
55	Glu	Ala	Thr	Thr	Leu	Gly	Ala	Leu	Glu	Phe	Ser	Leu	Leu	Tyr	Asp	Gln	
	385					390					395					400	

EP 2 787 346 B1

	Asp	Asn	Ser	Ser	Leu	Gln	Cys	Thr	Ile	Ile	Lys	Ala	Lys	Gly	Leu	Lys	
					405					410					415		
5	Pro	Met	Asp	Ser	Asn	Gly	Leu	Ala	Asp	Pro	Tyr	Val	Lys	Leu	His	Leu	
				420					425					430			
10	Leu	Pro	Gly	Ala	Ser	Lys	Ser	Asn	Lys	Leu	Arg	Thr	Lys	Thr	Leu	Arg	
			435					440					445				
15	Asn	Thr	Arg	Asn	Pro	Ile	Trp	Asn	Glu	Thr	Leu	Val	Tyr	His	Gly	Ile	
	450						455					460					
20	Thr	Asp	Glu	Asp	Met	Gln	Arg	Lys	Thr	Leu	Arg	Ile	Ser	Val	Cys	Asp	
	465					470					475					480	
25	Glu	Asp	Lys	Phe	Gly	His	Asn	Glu	Phe	Ile	Gly	Glu	Thr	Arg	Phe	Ser	
				485						490					495		
30	Leu	Lys	Lys	Leu	Lys	Pro	Asn	Gln	Arg	Lys	Asn	Phe	Asn	Ile	Cys	Leu	
				500					505					510			
35	Glu	Arg	Val	Ile	Pro	Met	Lys	Arg	Ala	Gly	Thr	Thr	Gly	Ser	Ala	Arg	
			515					520					525				
40	Gly	Met	Ala	Leu	Tyr	Glu	Glu	Glu	Gln	Val	Glu	Arg	Val	Gly	Asp	Ile	
	530						535					540					
45	Glu	Glu	Arg	Gly	Lys	Ile	Leu	Val	Ser	Leu	Met	Tyr	Ser	Thr	Gln	Gln	
	545					550					555					560	
50	Gly	Gly	Leu	Ile	Val	Gly	Ile	Ile	Arg	Cys	Val	His	Leu	Ala	Ala	Met	
				565					570						575		
55	Asp	Ala	Asn	Gly	Tyr	Ser	Asp	Pro	Phe	Val	Lys	Leu	Trp	Leu	Lys	Pro	
				580					585					590			
60	Asp	Met	Gly	Lys	Lys	Ala	Lys	His	Lys	Thr	Gln	Ile	Lys	Lys	Lys	Thr	
			595					600					605				
65	Leu	Asn	Pro	Glu	Phe	Asn	Glu	Glu	Phe	Phe	Tyr	Asp	Ile	Lys	His	Ser	
	610						615					620					
70	Asp	Leu	Ala	Lys	Lys	Ser	Leu	Asp	Ile	Ser	Val	Trp	Asp	Tyr	Asp	Ile	
	625					630					635					640	
75	Gly	Lys	Ser	Asn	Asp	Tyr	Ile	Gly	Gly	Cys	Gln	Leu	Gly	Ile	Ser	Ala	

645

650

655

5 Lys Gly Glu Arg Leu Lys His Trp Tyr Glu Cys Leu Lys Asn Lys Asp
660 665 670

10 Lys Lys Ile Glu Arg Trp His Gln Leu Gln Asn Glu Asn His Val Ser
675 680 685

15 Ser Asp
690

<210> 2
<211> 865
<212> DNA
<213> Homo sapiens

20 <400> 2

atgactgaca ccgtgttcag caacagttct aaccgttgga tgtaccccag tgaccggccc 60
cttcaatcaa agctccaggc aggctggtcc gtccaccccg gtggtcagcc tgacaggcag 120
25 aggaagcagg aagagctgac tgatgaggag aaagaaatca tcaacagggt gattgctcga 180
gctgagaaaa tggaagagat ggagcaggag cgaatcggac gcctggtgga ccgcctagaa 240
aacatgagga agaacgtggc tggagatggg gtgaaccgct gcatactgtg tggagaacag 300
30 ctggggatgc tgggctctgc ctgtgtagta tgtgaggact gtaagaagaa cgtctgcacc 360
aagtgcggag tggagaccaa caaccgcctg cattctgtgt ggctctgcaa aatctgcatt 420
35 gagcagaggg aggtgtggaa gcgttctgga gcgtggttct tcaaaggctt ccccaaacag 480
gtcctccac agcctatgcc tataaagaag accaagcccc agcagcctgt cagtgagcct 540
gctgcccctg aacagcctgc tcctgagccc aagcaccctg cccgggctcc agctcgagg 600
40 gacagtgaag ataggagggg cccgggtcag aagacaggcc ctgaccagc ctctgctccc 660
gggcgaggaa actatgggcc tcccgtgcgc agggcctccg aggcacgaat gagctcatct 720
agccgagatt cagagagctg ggaccacagt gggggtgctg gagactccag ccggagccca 780
45 gcaggtttga gacgggcca ctcagtccag gcctccagac ctgccccagg ctcggtgcag 840
agcccagcgc cacctcagcc tgggc 865

Claims

1. Use of an anti-rabphilin 3a antibody for the *in vitro* detection of lymphocytic infundibuloneurohypophysitis.
2. An assay for lymphocytic infundibuloneurohypophysitis, using the level of an anti-rabphilin 3a antibody in a specimen as an indicator.
3. The assay according to claim 2, which includes the following steps (1) to (3):

- (1) a step of providing a specimen derived from a subject;
- (2) a step of detecting an anti-rabphilin 3a antibody in the specimen; and
- (3) a step of determining whether the developed disease is lymphocytic infundibuloneurohypophysitis or not, or the possibility of developing lymphocytic infundibuloneurohypophysitis, based on the detection result.

- 4. The assay according to claim 3, wherein the high level of the anti-rabphilin 3a antibody is used as the indicator showing that the developed disease is lymphocytic infundibuloneurohypophysitis, and as the indicator showing that the possibility of developing lymphocytic infundibuloneurohypophysitis is high.
- 5. The assay according to claim 3 or 4, wherein the determination in the step (3) is carried out based on the comparison between the detected value obtained in the step (2) and the detected value obtained in the control specimen.
- 6. The assay according to claim 3 or 4, wherein the determination in the step (3) is carried out based on the comparison between the detected value obtained in the step (2) and the detected value in the specimen taken from the same subject in the past.
- 7. The assay of according to any of claims 2 to 6, wherein the specimen is blood, plasma, serum, cerebrospinal fluid, or urine.
- 8. Use of a test reagent comprising rabphilin 3a or its antibody-binding fragment for the assay of lymphocytic infundibuloneurohypophysitis *in vitro*.
- 9. Use of a kit, comprising the test reagent as defined in claim 8 for testing lymphocytic infundibuloneurohypophysitis *in vitro*.

Patentansprüche

- 1. Verwendung von einem Anti-Rabphilin 3a Antikörper zur *in vitro* Detektion der lymphozytischen Infundibuloneurohypophysitis.
- 2. Analyse von lymphozytischer Infundibuloneurohypophysitis unter Verwendung eines Anti-Rabphilin 3a Antikörper Niveaus in einer Probe als ein Indikator.
- 3. Analyse gemäß Anspruch 2, welche die folgenden Schritte (1) bis (3) enthält:
 - (1) ein Schritt zur Verfügungstellung einer Probe erlangt aus einem Probanden;
 - (2) ein Schritt zur Detektion eines Anti-Rabphilin 3a Antikörpers in der Probe; und
 - (3) ein Schritt zur Bestimmung ob die entwickelte Krankheit lymphozytische Infundibuloneurohypophysitis ist oder nicht, oder die Möglichkeit der Entwicklung einer lymphozytischen Infundibuloneurohypophysitis, basierend auf dem Detektionsergebnis.
- 4. Analyse gemäß Anspruch 3, wobei das hohe Niveau des Anti-Rabphilin 3a Antikörpers als Indikator verwendet wird um zu zeigen, dass die entwickelte Krankheit lymphozytische Infundibuloneurohypophysitis ist, und als Indikator um zu zeigen, dass die Möglichkeit der Entwicklung der lymphozytischen Infundibuloneurohypophysitis hoch ist.
- 5. Analyse gemäß Anspruch 3 oder 4, wobei die Bestimmung in dem Schritt (3) basierend auf dem Vergleich zwischen dem detektierten Niveau erhalten in dem Schritt (2) und dem detektierten Niveau erhalten in der Kontrollprobe ausgeführt wird.
- 6. Analyse gemäß Anspruch 3 oder 4, wobei die Bestimmung in dem Schritt (3) basierend auf dem Vergleich zwischen dem detektierten Niveau erhalten in dem Schritt (2) und dem detektierten Niveau in einer Probe entnommen von dem gleichen Probanden in der Vergangenheit ausgeführt wird.
- 7. Analyse gemäß einem der Ansprüche 2 bis 6, wobei die Probe Blut, Plasma, Serum, cerebrospinale Flüssigkeit oder Urin ist.
- 8. Verwendung von einer Testreagenz umfassend Raphilin 3a oder dessen Antikörper bindendes Fragment zur Analyse


der lymphozytischen Infundibuloneurohypophysitis *in vitro*.

9. Verwendung eines Kits umfassend die Testreagenz wie definiert in Anspruch 8 zur Untersuchung von lymphozytischer Infundibuloneurohypophysitis *in vitro*.

Revendications

1. Utilisation d'un anticorps anti-rabphiline 3a pour la détection *in vitro* de la neurohypophysite infundibulaire lymphocytaire.
2. Test pour la neurohypophysite infundibulaire lymphocytaire, utilisant le niveau d'un anticorps anti-rabphiline 3a dans un échantillon comme indicateur.
3. Test selon la revendication 2, qui inclut les étapes (1) à (3) suivantes:
 - (1) une étape de fourniture d'un échantillon dérivé d'un sujet;
 - (2) une étape de détection d'un anticorps anti-rabphiline 3a dans l'échantillon; et
 - (3) une étape de détermination si la maladie développée est la neurohypophysite infundibulaire lymphocytaire ou pas, ou de la possibilité de développer une neurohypophysite infundibulaire lymphocytaire, sur la base du résultat de la détection.
4. Test selon la revendication 3, où le haut niveau de l'anticorps anti-rabphiline 3a est utilisé comme indicateur montrant que la maladie développée est la neurohypophysite infundibulaire lymphocytaire, et comme indicateur montrant que la possibilité de développer une neurohypophysite infundibulaire lymphocytaire est élevée.
5. Test selon la revendication 3 ou 4, où la détermination dans l'étape (3) est accomplie sur la base de la comparaison entre la valeur détectée obtenue dans l'étape (2) et la valeur détectée obtenue dans l'échantillon témoin.
6. Test selon la revendication 3 ou 4, où la détermination dans l'étape (3) est accomplie sur la base de la comparaison entre la valeur détectée obtenue dans l'étape (2) et la valeur détectée dans l'échantillon prélevé auprès du même sujet dans le passé.
7. Test selon l'une quelconque des revendications 2 à 6, où l'échantillon est du sang, du plasma, du sérum, du liquide céphalo-rachidien ou de l'urine.
8. Utilisation d'un réactif de test comprenant de la rabphiline 3a ou son fragment de liaison d'anticorps pour le test de la neurohypophysite infundibulaire lymphocytaire *in vitro*.
9. Utilisation d'un kit, comprenant le réactif de test tel que défini dans la revendication 8 pour tester la neurohypophysite infundibulaire lymphocytaire *in vitro*.

[Fig. 1]



[Fig. 2]

LINH..... 9 cases **8/9**

LAH..... 3 cases **0/3**

IgG4-related hypophysitis 3 cases **1/3**

Disease control 10 cases **1/10**

Germ cell tumor + DI 3 cases **1/3**

After surgery of suprasellar of
pineal tumor + DI 1 case **0/1**

After surgery of hypothalamus
tumor + DI 1 case **0/1**

Rathke's cleft cyst + DI 1 case **0/1**

Craniopharyngioma + DI 2 cases **0/2**

After surgery of Glioma + DI 1 case **0/1**

Sarcoidosis + DI 1 case **0/1**

Healthy control ... 10 cases **1/10**

All Cases

Sensitivity 88.9% (8/9)

Specificity 87.0% (20/23)


88.5% (23/26)

↑ Value including IgG4

Discrimination from tumor lesion + DI

Specificity 90% (9/10)


[Fig. 3]




[Fig. 4]

Posterior pituitary


AVP



Rabphilin 3a




merge




SON


AVP



Rabphilin 3a



merge



[Fig. 5]

Lymphocytic infundibuloneurohypophysitis	11/12	Healthy subject	3/12
Lymphocytic adenohypophysitis	0/5		
Lymphocytic hypophysitis	2/6		
IgG4-related hypophysitis	2/5		
Idiopathic central diabetes insipidus	0/6	Autoimmune disease	7/69
		Mixed connective-tissue disease	0/2
After surgery of cerebral tumor DI	1/14	Multiple myositis/dermatomyositis	0/8
Germ cell tumor	1/4	Chorionitis	1/14
After surgery of suprasellar of pineal tumor	0/1	Chronic Arthritis rheumatoid	4/20
After surgery of hypothalamus tumor	0/1	Sjogren syndrome	1/10
Rathke's cleft cyst	0/2	SLE	1/15
Craniopharyngioma	0/2	Healthy subject (foreigner)	5/30
After surgery of Glioma	0/1		
After surgery of teratoma	0/2		
Sarcoidosis	0/1		
Empty sella DI	0/1		
Isolated ACTH deficiency	2/5		

[FIG. 6]

<u>All cases</u>	<u>Discrimination between other autoimmune disease and LINH</u>
Sensitivity 91.7% (11/12)	
Specificity 85.6% (131/153)	Specificity 89.9% (62/69)
<u>Discrimination from tumor lesion + DI</u>	<u>Discrimination between healthy subject and LINH</u>
Specificity 92.8% (13/14)	Specificity 80.9% (34/42)
<u>Discrimination between other autoimmune disease and LINH</u>	
Specificity 95.2% (20/21)	

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- JP 2011258387 A [0001]
- WO 0068412 A1 [0038]
- WO 0127260 A1 [0038]
- WO 2002024939 A1 [0038]
- WO 2005063979 A1 [0038]
- JP 6007134 A [0038]
- JP 2002529531 A [0038]
- JP 2005355513 A [0038]
- JP 2006042601 A [0038]
- JP 2007097438 A [0038]
- JP 2008029203 A [0038]
- JP 2011258387 P [0052]

Non-patent literature cited in the description

- *J Clin Endocrinol Metab*, 2002, vol. 87 (2), 752-7 [0004]
- *Horm Res.*, 2001, vol. 55 (6), 288-92 [0004]
- *Eur J Endocrinol.*, 2002, vol. 147 (6), 767-75 [0004]
- *J Endocrinol Invest*, 2007, vol. 30 (2), 153-62 [0004]
- *Clin Endocrinol.*, 2008, vol. 69 (2), 269-78 [0004]
- SHIMIZU, Y. et al. *Nature Biotech.*, 2001, vol. 19, 751-755 [0034]
- MADIN, K. et al. *Proc. Natl. Acad. Sci. USA*, 2000, vol. 97, 559-564 [0037] [0038]
- SAWASAKI, T. et al. *Proc. Natl. Acad. Sci. USA*, 2002, vol. 99, 14652-14657 [0037]
- ANDERSON, C. W. et al. *Methods Enzymol.*, 1983, vol. 101, 638-644 [0038]
- KAWARASAKI, Y. et al. *Biotechnol. Prog.*, 2000, vol. 16, 517-521 [0038]