



(11)

EP 2 834 263 B9

(12)

## CORRECTED EUROPEAN PATENT SPECIFICATION

(15) Correction information:

**Corrected version no 1 (W1 B1)****Corrections, see****Description Paragraph(s) 82, 83, 89, 94**

(51) Int Cl.:

**C07K 14/135 (2006.01)****C12N 5/10 (2006.01)****C12N 7/04 (2006.01)****C12N 15/40 (2006.01)****A61K 39/12 (2006.01)****G01N 33/564 (2006.01)****G01N 33/569 (2006.01)**

(48) Corrigendum issued on:

**27.02.2019 Bulletin 2019/09**

(86) International application number:

**PCT/US2013/035408**(45) Date of publication and mention  
of the grant of the patent:**10.10.2018 Bulletin 2018/41**

(87) International publication number:

**WO 2013/152274 (10.10.2013 Gazette 2013/41)**(21) Application number: **13719249.8**(22) Date of filing: **05.04.2013****(54) EPITOPE-SCAFFOLD IMMUNOGENS AGAINST RESPIRATORY SYNCYTIAL VIRUS (RSV)****EPITOPE-SCAFFOLD IMMUNOGENISCHE PEPTIDE GEGEN RSV****PEPTIDES IMMUNOGENIQUES EPITOPE-SCAFFOLD CONTRE LE RSV**

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB  
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO  
PL PT RO RS SE SI SK SM TR**(30) Priority: **05.04.2012 US 201261620797 P  
05.04.2012 US 201261620804 P**(43) Date of publication of application:  
**11.02.2015 Bulletin 2015/07**(73) Proprietor: **University of Washington through its  
Center for  
Commercialization  
Seattle, WA 98105-4608 (US)**

(72) Inventors:

- SCHIEF, William, R.  
Encinitas, CA 92024 (US)**
- CORREIA, Bruno, E.  
San Diego, CA 92109 (US)**

(74) Representative: **Zwicker, Jörk  
ZSP Patentanwälte PartG mbB  
Hansastraße 32  
80686 München (DE)**

(56) References cited:

**WO-A2-2012/048115**

- JASON S. MCLELLAN ET AL: "Design and Characterization of Epitope Scaffold Immunogens That Present the Motavizumab Epitope from Respiratory Syncytial Virus", JOURNAL OF MOLECULAR BIOLOGY, vol. 409, no. 5, 1 April 2011 (2011-04-01) , pages 853-66, XP055000825, ISSN: 0022-2836, DOI: 10.1016/j.jmb.2011.04.044**
- J. S. MCLELLAN ET AL: "Structure of Respiratory Syncytial Virus Fusion Glycoprotein in the Postfusion Conformation Reveals Preservation of Neutralizing Epitopes", JOURNAL OF VIROLOGY, vol. 85, no. 15, 1 August 2011 (2011-08-01), pages 7788-7796, XP002710047, ISSN: 0022-538X, DOI: 10.1128/JVI.00555-11**
- J. S. MCLELLAN ET AL: "Structure of a Major Antigenic Site on the Respiratory Syncytial Virus Fusion Glycoprotein in Complex with Neutralizing Antibody 101F", JOURNAL OF VIROLOGY, vol. 84, no. 23, 1 December 2010 (2010-12-01), pages 12236-12244, XP055000827, ISSN: 0022-538X, DOI: 10.1128/JVI.01579-10**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- ARBIZA J ET AL: "CHARACTERIZATION OF TWO ANTIGENIC SITES RECOGNIZED BY NEUTRALIZING MONOCLOnal ANTIBODIES DIRECTED AGAINST THE FUSION GLYCOPROTEIN OF HUMAN RESPIRATORY SYNCYTIAL VIRUS", JOURNAL OF GENERAL VIROLOGY, SOCIETY FOR GENERAL MICROBIOLOGY, SPENCERS WOOD, GB, vol. 73, no. 9, 1 January 1992 (1992-01-01), pages 2225-2234, XP002548166, ISSN: 0022-1317, DOI: 10.1099/0022-1317-73-9-2225
- Q. ZHU ET AL: "Analysis of Respiratory Syncytial Virus Preclinical and Clinical Variants Resistant to Neutralization by Monoclonal Antibodies Palivizumab and/or Motavizumab", JOURNAL OF INFECTIOUS DISEASES. JID, vol. 203, no. 5, 5 January 2011 (2011-01-05), pages 674-682, XP055285525, CHICAGO, IL. ISSN: 0022-1899, DOI: 10.1093/infdis/jiq100

**Description****Background**

5 [0001] Respiratory Syncytial Virus (RSV) is the leading cause of viral death in infants worldwide and also causes disease in the elderly and immune-compromised. The current method for preventing RSV infection is passive immunization with Palivizumab (Pali), an FDA-licensed humanized monoclonal antibody that binds the F protein on the RSV surface. Though effective at preventing RSV infection, Pali treatment is not economically or logically feasible on a global scale. McLellan et al. (J Mol Biol. 2011 June 24; 409(5): 853-866) disclose the design and characterization of 10 epitope-scaffold immunogens that present the Motavizumab epitope from RSV and elicit F-binding RSV antibodies in mice.

**Summary of the Invention**

15 [0002] In a first aspect, the present invention provides isolated polypeptides comprising an amino acid sequence according to any of SEQ ID NOS: 1, 3 and 18-29.  
 [0003] In another aspect, the present invention provides virus-like particles comprising the polypeptide of the invention.  
 [0004] In further aspects, the present invention provides isolated nucleic acids encoding the polypeptides of the invention; recombinant expression vectors comprising the isolated nucleic acids of the invention operatively linked to a 20 promoter; and recombinant host cells comprising the recombinant expression vectors of the invention.  
 [0005] In a still further aspect, the present invention provides pharmaceutical compositions, comprising the polypeptide or virus-like particles of the invention, and a pharmaceutically acceptable carrier.

**Brief Description of the Drawings**

25 [0006]

Figure 1 is an overview of the computational procedure Fold From Loops. The procedure takes a functional site (such as the helical hairpin shown) that will be used as the folding nucleus and remain in fixed backbone conformation 30 throughout the procedure. A target topology is supplied and distance constraints are (optionally) derived from the target topology structure to guide the folding trajectory. The polypeptide chain is extended from the folding nucleus and the chain is then folded. If the models produced are more than a cutoff root mean square deviation (rmsd) (e.g. 5 Å) away from the target topology, they are discarded. Otherwise, they enter cycles of design and full-atom optimization. The figure depicts 3 cycles of iterative design and optimization as a reasonable choice, but the number 35 of cycles is to be chosen at the discretion of the user.

Figure 2 shows Motavizumab (Mota) in complex with its peptide epitope from the RSVF protein. A) Side-view of the complex. B) Back-view of the complex. C) Side chains on the interface of the complex are shown in sticks.

Figure 3 is a sequence alignment of the different scaffolds and the sequence of the protein used as target topology (T93). The similarities in many of the positions were imposed by the surface that in the scaffolds was intentionally 40 maintained the same as the target topology.

Figure 4 shows characterization of the oligomeric state of the scaffolds by size exclusion chromatography and static light scattering. All the molecules showed a single monodisperse species and had a molecular weight close to the expected for a monomeric species, approximately 15 kDa. The UV signal from size exclusion is the upper trace in all the graphs, and the light scattering signal is the lower trace.

45 Figure 5 shows circular dichroism analysis of secondary structure and thermal stability of FFL designs. Wavelength scans (left row) for the designs show the double minima typical for helical proteins. Thermal denaturation curves (right row) indicate cooperative unfolding for most designs, and show that FFL\_005 does not melt up to 95 °C. The high stability of FFL\_005 is exemplified by the wavelength scan at 95 °C (left row). Melting temperatures are given in Table 1.

50 Figure 6 shows binding of the scaffolds to Motavizumab assessed by SPR. The scaffolds were coupled to the biacore chip and Motavizumab was used as analyte. Both data and kinetic fits are shown. Kinetic fit parameters are given in Table 1.

55 Figure 7 shows Mota binding specificity of FFL\_001 assessed by SPR. Mota IgG was the ligand, captured by anti-human IgG on the sensor chip, and FFL\_001 and an epitope point mutant of FFL\_001 (FFL\_001\_K82E) were analytes at a concentration of 22 µM. The interaction between FFL\_001 and Mota was eliminated by the point mutation.

Figure 8. Negative stain TEM of wild-type (left) and lysine-functionalized (right) HepBcAg particles. Scale bar is 50 nM.

## Detailed Description of the Invention

**[0007]** Within this application, unless otherwise stated, the techniques utilized may be found in any of several well-known references such as: Molecular Cloning: A Laboratory Manual (Sambrook, et al., 1989, Cold Spring Harbor Laboratory Press), Gene Expression Technology (Methods in Enzymology, Vol. 185, edited by D. Goeddel, 1991. Academic Press, San Diego, CA), "Guide to Protein Purification" in Methods in Enzymology (M.P. Deutshcer, ed., (1990) Academic Press, Inc.); PCR Protocols: A Guide to Methods and Applications (Innis, et al. 1990. Academic Press, San Diego, CA), Culture of Animal Cells: A Manual of Basic Technique, 2nd Ed. (R.I. Freshney. 1987. Liss, Inc. New York, NY), Gene Transfer and Expression Protocols, pp. 109-128, ed. E.J. Murray, The Humana Press Inc., Clifton, N.J.), and the Ambion 1998 Catalog (Ambion, Austin, TX).

**[0008]** As used herein, the amino acid residues are abbreviated as follows: alanine (Ala; A), asparagine (Asn; N), aspartic acid (Asp; D), arginine (Arg; R), cysteine (Cys; C), glutamic acid (Glu; E), glutamine (Gln; Q), glycine (Gly; G), histidine (His; H), isoleucine (Ile; I), leucine (Leu; L), lysine (Lys; K), methionine (Met; M), phenylalanine (Phe; F), proline (Pro; P), serine (Ser; S), threonine (Thr; T), tryptophan (Trp; W), tyrosine (Tyr; Y), and valine (Val; V).

**[0009]** As used herein, the singular forms "a", "an" and "the" include plural referents unless the context clearly dictates otherwise. "And" as used herein is interchangeably used with "or" unless expressly stated otherwise.

**[0010]** All embodiments of any aspect of the invention can be used in combination, unless the context clearly dictates otherwise.

**[0011]** In a first aspect, the present invention provides isolated polypeptides, comprising or consisting of an amino acid sequence according to the following:

(-/G)(-/S)(M/L/R/F)SD(R/A/I/M/V/L)(R/M)KD(L/A/V)E(E/R/K/D)R(L/F/I/A)DK(L/F/A)  
 (L/V/F/M)EA(A/V/F/I/L)KNK(E/M/L/F/W/V)DK(F/M/E/I/V)KAA(M/L/F/I)RK(R/E/G/D/  
 25 Q)(G/D/P/W/E/Q)(Q/I/P/K/F)(R/Q/K/S/G/H)EER(M/K/R/A)KD(W/L/M/K)(A/M/K/F)K(I/F  
 /L/E/K/D)(A/V/M/F/L)R(D/K/E/Y)E(F/V/A/R/M/L)EQ(F/L/A/V/M)R(K/R)A(V/M/I)RN(F/  
 30 R/V/I/A/Y)(L/E/A)(S)(E)(A/L)(L)(S)K(I)(N/Y)D(Y/M/L)(P)I(T)(N/I)(D)(D/Q/K)(K)(K/E/T  
 /M/Q)(L)(T/I/M/V/A)(S)(N)(D/K)(A/T/L/V/I)(K/L/I)K(F/Y/K/E/R/L)(D/A/V/M)(A)(E/I/R)(  
 V/A/M/L)(A/E/K/F/M/W)KK(L/I/V)E(A/L)(F/M/L/E/I)(K/A/V/M/I/L)AD(A/V/I)E(E/R/D/  
 35 K/I/A)(A/M/K/L/W)(A/F/V/K)TQ(-/G)(-/S)(-/W) **(SEQ ID NO:1).**

**[0012]** The inventors have designed the polypeptides of the invention to elicit neutralizing antibodies with similar specificity as Palivizumab or Motavizumab. Palivizumab is a FDA-licensed therapeutic antibody that potently neutralizes Respiratory Syncytial Virus (RSV) by binding antigenic site A (also called "site II") on the RSV F surface glycoprotein.

**[0013]** Motavizumab is an affinity-matured variant of Palivizumab. Thus, vaccine that elicits RSV-neutralizing antibodies similar to Palivizumab (Pali) or Motavizumab (Mota) is desired to protect against RSV infection. Pali and Mota bind to a conformational epitope on the RSV F protein. As disclosed herein, the inventors have developed a computational method to design de novo protein scaffolds for epitope conformational stabilization and presentation to the immune system. This method was applied to the Mota epitope to design the polypeptides of the invention, which are shown to be monomeric, highly thermostable, and extremely high binding affinities for Mota, indicating that the polypeptides have successfully stabilized the desired epitope conformation, as confirmed by crystal structure analysis. The inventors have also demonstrated that polypeptides falling within the scope of this genus can elicit neutralizing antibodies against RSV.

**[0014]** Parentheses represent variable positions in the polypeptide, with the recited amino acid residues as alternatives in these positions.

**[0015]** Disclosed are polypeptide according to SEQ ID NO:2.  
 (-/G)(-/S)(M/L/R/F)SD(R/A/I/M/V)RKD(L/A/V)E(E/R/K/D)R(L/F/I/A)DK(L/F/A)(L/V/F)EA(A/V/F/L)KNK(M/L/F/V)DK(F/M/  
 E/I)KAA(M/L/F/I)RK(R/E/G/D)(G/D/P/W/Q)(Q/I/P/F)(R/Q/K/S/H)EER(M/K/R/A)KD(W/L/M/K)(A/M/K/F)K(I/F/L/E/K/D)(A/  
 50 V/M/F/L)R(D/K/Y/E)E(F/V/A/R/M)EQ(F/L/A/V/M)R(K/R)A(V/M/I)RN(F/R/V/I/Y)(L/E/A)SE(A/L)LSKND(Y/M/L)PITND(D/  
 /Q/K)KKL(T/I/M/V/A)SND(T/L/V/I)(K/L/I)K(F/Y/K/E/L)(D/A/V/M)A(E/I/R)(V/A/M/L)(E/K/F/W)KK(L/I/V)E(A/L)(F/M/L/E/I)(  
 K/A/V/M/L)AD(A/V/I)E(E/R/D/K/I/A)(A/M/K/L/W)(A/F/V)TQ(-/G)(-/S)(-/W) **(SEQ ID NO:2).**

In one preferred embodiment of the invention, the polypeptides comprise or consist of an amino acid sequence according to SEQ ID NO:29.

(-/G)(-/S)(M/L/R/F)SD(R/A/I/M/V)RKD(L/A/V)E(E/R/K/D)R(L/F/I/A)DK(L/F/A)(L/V/F)EA(A/V/F/L)KNK(M/L/F/V)DK(F/M/

E/I)KAA(M/L/F/I)RK(R/E/G/D)(G/D/P/W/Q)(Q/I/P/F)(R/Q/K/S/H)EER(M/K/R/A)KD(W/L/M/K)(A/M/K/F)K(I/F/L/E/K/D)(A/V/M/F/L)R(D/K/Y/E)E(F/V/A/R/M)EQ(F/L/A/V/M)R(K/R)A(V/M/I)RN(F/R/V/I/Y)(L/E/A)SE(A/L)LSKI(N/Y)D(Y/M/L)PIT(N/I)D(D/Q/K)K(K/E/T/M/Q)L(T/I/M/V/A)SND(T/L/V/I)(K/L/I)K(F/Y/K/E/L)(D/A/V/M)A(E/I/R)(V/A/M/L)(E/K/F/W)KK(L/I/V)E(A/L)(F/M/L/E/I)(K/A/V/M/L)AD(A/V/I)E(E/R/D/K/I/A)(A/M/K/L/W)(A/F/V)TQ(-/G)(-/S)(-/W) **SEQ ID NO:29**

5 [0015] Polypeptides according to this genus are those that are present in those polypeptides demonstrating the best range of activities, as demonstrated in the examples that follow. In a further preferred embodiment, the polypeptides comprise or consist of an amino acid sequence according SEQ ID NO:3.

(-/G)(-/S)(M/L/R/F)SD(I/M)RKD(L/A)E(E/R/D)R(F/A)DK(L/F/A)(V/F)EA(A/V/L)KNK(L/F/W)DK(F/M/I)KAA(L/F/I)RK(E/G/D)(G/D/W/Q)(Q/I/P/F)(Q/K/S/H)EER(M/R/A)KD(W/L/M)(M/K/F)K(F/L/K/D)(A/M/L)R(Y/K)E(V/A/M)EQ(L/A/M)R(K/R)A(V/M/I)RN(F/R/I/Y)(L/E/A)SE(A/L)LSKI(N/Y)D(M/L)PIT(N/I)D(D/Q)K(K/E/T/M/Q)L(I/M/A)SND(L/V/I)(K/L/I)K(F/Y/E/L)(D/A/V/M)A(E/I/R)(V/A/L)(E/F/W)KK(L/I)EA(M/L/I)(K/A/M/L)AD(A/V/I)E(R/D/I/A)(M/K/L/W)(A/F/V)TQ(-/G)(-/S)(-/W) **SEQ ID NO:3**

In one preferred embodiment of the invention, the polypeptides comprise or consist of an amino acid sequence according to SEQ ID NO:30.

15 (-/G)(-/S)(M/L/R/F)SD(I/M)RKD(L/A)E(E/R/D)R(F/A)DK(L/F/A)(V/F)EA(A/V/L)KNK(L/F/W)DK(F/M/I)KAA(L/F/I)RK(E/G/D)(G/D/W/Q)(Q/I/P/F)(Q/K/S/H)EER(M/R/A)KD(W/L/M)(M/K/F)K(F/L/K/D)(A/M/L)R(Y/K)E(V/A/M)EQ(L/A/M)R(K/R)A(V/M/I)RN(F/R/I/Y)(L/E/A)SE(A/L)LSKIND(M/L)PITND(D/Q)KKL(I/M/A)SND(L/V/I)(K/L/I)K(F/Y/E/L)(D/A/V/M)A(E/I/R)(V/A/L)(E/F/W)KK(L/I)EA(M/L/I)(K/A/M/L)AD(A/V/I)E(R/D/I/A)(M/K/L/W)(A/F/V)TQ(-/G)(-/S)(-/W) **SEQ ID NO:30**

20 [0016] Polypeptides according to this genus are those that have been exemplified by the inventors as eliciting neutralizing antibodies against RSV. Further disclosed are polypeptides comprising or consisting of an amino acid sequence selected from the group consisting of

>FFL\_001

25 GSRSDMRKDAERRFDKFVEAAKNFKDKFKAALRKGDIKEERRKDMKKLARKEAEQARRAVRNRLSELLSKINDMPITNDQKKLMSNDVLKFAAEAEKKIEALAADAEDKFTQGSW **(SEQ ID NO:4)**;

30 >FFL\_002

GSLSDVRKDVEKRIDKALEAFKNKMDKEKA AfrKDPPSEERRKDKKEFREEREQV  
35 RKAIRNVLSEALSKINDLPITNDKKLVSDVIKKVAEMKKVELEVADVEKKVTQGSW **(SEQ ID NO:5)**;

>FFL\_004

40 GSMSDARKDLEERLDKLLEAAKNKMDKFKAAMRKRGQREERKKDWAKIVRDEFEQFRKAVRNFLSEALSKINDYPITNDDKKLTSNDTKKFAAEVEKKLEAFKADVEEAATQ  
45 **(SEQ ID NO:6)** ;

>FFL\_005

50 GSMSDIRKDLEERFDKLVEALKNKVDKMKA AfrKDQFHEERMKDWFKDLRKEVEQMRRAVRNYASEALSKINDLPITNDKKLASNDVLKLVAEVWKKLEAILADVEAWFTQ **(SEQ ID NO:7)**;

55 >FFL\_006

GSFSDIRKDAEDRADKAFEAAKNKFDKIKAAIRKDW PSEERA KDL MKKARY YEMEQA  
RRAIRNIESEALSKINDLPITNDQKKLASNDI KEMARLF KKLEALMADIEILVTQ (SEQ

5 ID NO:8); and

>FFL\_007

10 GSLSDIRKDAERRFDKLVEAVKNKLDKMKAA L RKEGQQEER MKDL MKFMR KEVEQ  
LRKAMRNFLSEALSKINDMPITNDKKLISNDLKKYDAIAEKKLEAMKADVERMAT  
QGSW (SEQ ID NO:9).

15 [0017] Each of these polypeptides is demonstrated in the examples that follow to be monomeric, highly thermostable, and have extremely high binding affinities for Mota, indicating that the polypeptides have successfully stabilized the desired epitope conformation, and a number of these polypeptides have been shown to elicit neutralizing antibodies against RSV.

20 [0018] In another embodiment, the polypeptides comprise resistance mutants for Motavizumab and/or Palivizumab. These polypeptides can be used, for example, in a vaccine to protect against RSV strains that are resistant to Mota-like or Pali-like neutralizing antibodies, or to prevent the emergence of such resistant RSV strains. In such vaccines, it may be desirable to include epitope-scaffolds bearing resistance mutations within the RSV epitope. In this way, vaccination with "resistance mutant epitope-scaffolds" might induce antibodies that neutralize resistance mutant viruses and hence prevent the emergence of those resistance viruses. Similarly, the "resistance mutant epitope-scaffolds" could be used 25 as reagents to isolate antibodies that neutralize resistance mutant viruses.

25 [0019] For Motavizumab, one established resistance mutation is K272E (Zhu et al. JID, 2011). For Palivizumab, several resistance mutations have been identified, including K272N, K272M, K272T, K272Q (Zhu et al. JID, 2011). Arbiza et al. J. Gen Virol 1992 identified resistance mutations K272E, K272T, N262Y and N268I and showed that these reduce binding by antibodies directed to this region on RSVF.

30 [0020] We have shown that the K272E and N262Y mutations introduced onto our epitope-scaffold RSV\_1isea\_FFL\_001 significantly reduce binding by Mota or Pali. By structural similarity among the epitope-scaffolds, we expect that either of these mutations will have the same effect on binding to other scaffolds such as 005 and 007.

35 [0021] Exemplary such polypeptides include the following (note that the residue numbering for the mutations refers to the position in the RSV F protein sequence not in the epitope-scaffold itself).

RSV\_1isea\_FFL\_001\_resistance\_mutants\_at\_pos272

40 GSRSDMRKDAERRFDKFVEAKNKFDKFKAA L RKGDIKEERRKDMKKLARKEAEQ  
ARRAVRNRLSELLSKINDMPITNDQK[E,T,M,Q]LMSNDVLKFAAEAEKKIEALAADA  
EDKFTQGSW (SEQ ID NO:10)

45 RSV\_1isea\_FFL\_005\_resistance\_mutants\_at\_pos272

GSMSDIRKDLEERFDKLVEALKNKVDKMKAAFRKDQFHEERMKDWFKDLRKEVEQ  
MRRAVRNYASEALSKINDLPITNDDK[E,T,M,Q]LASNDVLKVAEVWKKLEAILADV  
50 EAWFTQ (SEQ ID NO:11)

55 RSV\_1isea\_FFL\_007\_resistance\_mutants\_at\_pos272

5 GSLSDIRKDAERRFDKLVEAVKNKLDKMKAAALRKEGQQEERMKDLMKFMRKVEEQ  
LRKAMRNFLSEALSKINDMPITNDDK[E,T,M,Q]LISNDLKKYDAIAEKKLEAMKADV  
ERMATQGSW (SEQ ID NO:12)

RSV\_1isea\_FFL\_001\_resistance\_mutant\_at\_262

10 GSRSDMRKDAERRFDKFVEAAKNFKDKFKAALRKGDIKEERRKDMKKLARKEAEQ  
ARRAVRNRLSELLSKIYDMPITNDQKKLMSNDVLKFAAEAEKKIEALAADAEDKFTQ  
15 GSW (SEQ ID NO:13)

RSV\_1isea\_FFL\_005\_resistance\_mutant\_at\_262

20 GSMSDIRKDLEERFDKLVEALKNVDKMKAAFRKDQFHEERMKDWFKDLRKEVEQ  
MRRAVRNYASEALSKIYDLPITNDDKKLASNDVLKLVAEVWKKLEAILADVEAWFT  
Q (SEQ ID NO:14)

25 RSV\_1isea\_FFL\_007\_resistance\_mutant\_at\_262

30 GSLSDIRKDAERRFDKLVEAVKNKLDKMKAAALRKEGQQEERMKDLMKFMRKVEEQ  
LRKAMRNFLSEALSKIYDMPITNDDKKLISNDLKKYDAIAEKKLEAMKADVERMAT  
QGSW (SEQ ID NO:15)

35 RSV\_1isea\_FFL\_001\_resistance\_mutant\_at\_268

40 GSRSDMRKDAERRFDKFVEAAKNFKDKFKAALRKGDIKEERRKDMKKLARKEAEQ  
ARRAVRNRLSELLSKINDMPITIDQKKLMSNDVLKFAAEAEKKIEALAADAEDKFTQ  
GSW (SEQ ID NO:16)

45 RSV\_1isea\_FFL\_005\_resistance\_mutant\_at\_268

50 GSMSDIRKDLEERFDKLVEALKNVDKMKAAFRKDQFHEERMKDWFKDLRKEVEQ  
MRRAVRNYASEALSKINDLPITIDDKKLASNDVLKLVAEVWKKLEAILADVEAWFT  
Q (SEQ ID NO:17)

RSV\_1isea\_FFL\_007\_resistance\_mutant\_at\_268

55

5 GSLSDIRKDAERRFDKLVEAVKNKLDKMKAAALRKEGQQEERMKDLMKFMRKEVEQ  
 LRKAMRNFLSEALSKINDMPITIDDKKLISNDLKKYDAIAEKKLEAMKADVERMATQ  
 GSW (SEQ ID NO:18).

10 [0022] The polypeptides may also comprise or consist of smaller epitope-scaffolds with similar properties to those disclosed above. Smaller scaffolds are potentially advantageous because they may contain fewer off-target epitopes, because they may be more amenable to particulate presentation by chemical conjugation, genetic fusion, or other means, and because smaller scaffolds may be more economical to produce commercially.

15 [0023] The smaller scaffolds comprise or consist of an isolated peptide comprising an amino acid of the formula X1-X2, wherein X1 is GS(M/L/C)SD(I/C)(R/C)KD(L/A/C)E(E/R)/FDK(L/G)(SEQ ID NO: 33); and wherein X2 is selected from the group consisting of:

20 VEA(L/V)K(K/N)(L/G)(Q/G)(G/N)(R/G)(Q/E)KEVEQ(M/L)R(R/K)A(V/M)RN  
 (Y/F)(A/L)SEALSKI(N/Y)D(L/M)PIT(N/I)DDK(K/E/T/M/Q)L(A/I)SND(V/L)(K/L)K(L/Y)  
 (V/D)A(E/I)(V/A)(W/E)KKLEA(I/M)(L/K)A(-/G/D)(-/S/V)(-/W/E)(-/R/A)(-/W/M)(-/F/A)(-  
 /T)(-/Q)(-/G)(-/S)(-/W) (SEQ ID NO: 34)

25 and

30 GS(C/L)SD(I/C)(R/C)KD(C/A)ERRFDKGDGGRKA(M/W)RNFLSE(C/F)LS(C/K)I  
 NDMPITNDDKKL(C/I)SND(L/C)KKY(D/L)AIAEK(-/G)(-/S)(-/W) (SEQ ID NO: 35).

35 [0024] X1 may be GS(M/L/C)SDIRKD(L/A)E(E/R)/FDKL(SEQ ID NO: 36); or X1 may be GS(L/C)SD(I/C)(R/C)KD(A/C)ERRFDKG (SEQ ID NO: 37). In various embodiments of the invention, the smaller scaffolds comprise or consist of:

40 RSV\_1isea\_FFL\_005\_min\_A

45 GSMSDIRKDLEERFDKLVEALKKGQGRQKEVEQMRR AVRNYASEALSKINDLPITND  
 DKKLASNDV LVAEVWKKLEAILA (SEQ ID NO:19);

or

45 RSV\_1isea\_FFL\_007\_min\_A

50 GSLSDIRKDAERRFDKLVEAVNLGNGEKEVEQLRKAMRNFLSEALSKINDMPITND  
 DKKLISNDLKKYDAIAEKKLEAMKAGSW (SEQ ID NO:20)

55 [0025] RSV\_1isea\_FFL\_005\_min\_A was designed based on the crystal structure of RSV\_1isea\_FFL\_005, and RSV\_1isea\_FFL\_007\_min\_A was designed based on the design model for RSV\_1isea\_FFL\_007. In both cases, we eliminated some regions of the parent scaffolds (RSV\_1isea\_FFL\_005 and RSV\_1isea\_FFL\_007) that were not directly contacting the RSV epitope. The total epitope-scaffold length was reduced from 112 to 82 for 005 and 115 to 85 for 007 (excluding the purification tag "LEHHHHHH" (SEQ ID NO: 32) at the Cterminus of each construct). To achieve this we employed a minimization protocol composed of two stages: I) deletion of the coordinates in the PDB file of the residues to be eliminated; II) Backbone rebuilding and sequence design of a new connecting loop between helix 1 and helix 2 in

the scaffolds. Residue ranges 25-48 and 104-112 in RSV\_1isea\_FFL\_005 and RSV\_1isea\_FFL\_007 were deleted, and new connecting loops of length 2 or 3 residues were built de novo to create RSV\_1isea\_FFL\_005\_min\_A and RSV\_1isea\_FFL\_007\_min\_A.

[0026] RSV\_1isea\_FFL\_005\_min\_A and RSV\_1isea\_FFL\_007\_min\_A were assessed for their binding to Mota and found to have dissociation constants of 51 and 48 pM, respectively. Thermal stability was also assessed; the melting temperature of RSV\_1isea\_FFL\_007\_min\_A was 60 °C while the melting temperature for RSV\_1isea\_FFL\_005\_min\_A was >75 °C but it did not completely melt at the highest temperature tested (98 °C) so the Tm could not be determined with certainty.

[0027] In further embodiments, the peptides comprise or consist of:

RSV\_1isea\_FFL\_005\_min\_A  
GSMSDIRKDEERFDKLVEALKKGQGRQKEVEQMRRAVRNYASEALSKI(N/Y)DLPIT(N/I)DDK(K/E/T/M/Q)LASN  
DVLKLVAEVWKKLEAILA(D/-)(V/-)(E/-)(A/-)(W/-)(F/-)(T/-)(Q/-)\_(SEQ ID NO:21) or

RSV\_1isea\_FFL\_007\_min\_A  
GSLSDIRKDAERRFDKLVEAVKNLGNGEKEVEQLRKAMRNFLSEALSKI(N/Y)DMPIT(N/I)DDK(K/E/T/M/Q)LISND  
LKKYDAIAEKKLEAMKA(D/-)(V/-)(E/-)(R/-)(M/-)(A/-)(T/-)(Q/-)GSW (SEQ ID NO:22)

[0028] Additional peptides according to the invention are selected from the group consisting of

RSV\_1ISEA\_T\_007\_23L\_D5\_C -> combination of 2 disulfides (D1+D3)

GSCSDIRKDCERRFDKGDGGRKAMRNFLSECLSINKDMPITNDDKKLCN

DLKKYDAIAEKKGWSW (SEQ ID NO:23)

RSV\_1ISEA\_T\_007\_23L\_D6\_C -> combination of 2 disulfides (D2+D3)

GSLSDCRKDCERRFDKGDGGRKAMRNFLSECLSCINDMPITNDDKKLISN

DLKKYDAIAEKKGWSW (SEQ ID NO:24)

RSV\_1ISEA\_T2\_007\_2\_D1\_C -> extra packed bundle + D1 disulfide

GSCSDIRKDAERRFDKGDGGRKAWRNFLSEFLSKINDMPITNDDKKLCN

DLKKYLAIAEKK (SEQ ID NO:25)

RSV\_1ISEA\_T2\_007\_2\_D2\_C

GSLSDCRKDAERRFDKGDGGRKAWRNFLSEFLSCINDMPITNDDKKLISN

DLKKYLAIAEKK (SEQ ID NO:26)

RSV\_1ISEA\_T2\_007\_2\_D3\_C

GSLSDIRKDCERRFDKGDGGRKAWRNFLSECLSINKDMPITNDDKKLCN

DLKKYLAIAEKK (SEQ ID NO:27)

RSV\_1ISEA\_T2\_007\_2\_D4\_C

GSLSDICKDAERRFDKGDRKAWRNFLSEFLSKINDMPITNDDKKLISN

DCKKYLAIAEKK (SEQ ID NO:28)

5 [0029] These peptides may incorporate one or more of the escape mutants disclosed above, and may incorporate the C-terminal deletion removed from the minimized versions disclosed above.

10 [0030] Also disclosed are polypeptides comprising or consisting of a sequence selected from the group consisting of SEQ ID NO:4 (FFL\_001), SEQ ID NO:5 (FFL\_002), SEQ ID NO:7 (FFL\_005), SEQ ID NO:8 (FFL\_006), SEQ ID NO:9 (FFL\_007), SEQ ID NO: 10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28. In a preferred embodiment of the invention, the polypeptide comprises or consists of a sequence selected from the group consisting of SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28. In a further preferred embodiment the polypeptide comprises or consists of a sequence selected from the group consisting of SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO:22. In a further preferred embodiment the polypeptide comprises or consists of a sequence selected from the group consisting of SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28.

15 [0031] The polypeptides may be resurfaced, referring to resurfacing as described in Correia et al J. Mol Biol 2011 or any related application of the concept of resurfacing.

20 [0032] In a further embodiment, the polypeptide includes any variant of the listed sequences obtained by adding one or more disulfide bonds.

25 [0033] As disclosed herein, the inventors have developed a computational method to design protein scaffolds for epitope conformational stabilization and presentation to the immune system. This method was applied to the Mota epitope to design the polypeptides of the invention, which are shown to be monomeric, highly thermostable, and extremely high binding affinities for Mota.

30 [0034] As used throughout the present application, the term "polypeptide" is used in its broadest sense to refer to a sequence of subunit amino acids. The polypeptides of the invention may comprise L-amino acids, D-amino acids (which are resistant to L-amino acid-specific proteases *in vivo*), or a combination of D- and L-amino acids. The polypeptides described herein may be chemically synthesized or recombinantly expressed. The polypeptides may contain any suitable linker, etc. for use in any desired application, such as a peptide tag to facilitate polypeptide purification, or a T-help epitope to enhance the desired immune response. For example, two of the exemplified polypeptides discussed below include a C-terminal "GSW" to facilitate determining protein concentration, as those polypeptides did not include any other 'W' residues.

35 [0035] The polypeptides may be linked to other compounds to promote an increased half-life *in vivo*, such as by PEGylation, HESylation, PASylation, glycosylation, or may be produced as an Fc-fusion or in deimmunized variants. Such linkage can be covalent or noncovalent as is understood by those of skill in the art.

40 [0036] The polypeptides described herein may further comprise a tag, such as a detectable moiety or therapeutic agent. The tag(s) can be linked to the polypeptide through covalent bonding, including, but not limited to, disulfide bonding, hydrogen bonding, electrostatic bonding, recombinant fusion and conformational bonding. Alternatively, the tag(s) can be linked to the polypeptide by means of one or more linking compounds. Techniques for conjugating tags to polypeptides are well known to the skilled artisan. Polypeptides comprising a detectable tag can be used, for example, as probes to isolate B cells that are specific for the epitope present in the polypeptide. However, they may also be used for other detection and/or analytical purposes. Any suitable detection tag can be used, including but not limited to enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals, and nonradioactive paramagnetic metal ions. The tag used will depend on the specific detection/analysis techniques and/or methods used such as flow cytometric detection, scanning laser cytometric detection, fluorescent immunoassays, enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), bioassays (e.g., neutralization assays), Western blotting applications, etc. When the polypeptides of the invention are used for flow cytometric detections, scanning laser cytometric detections, or fluorescent immunoassays, the tag may comprise, for example, a fluorophore. A wide variety of fluorophores useful for fluorescently labeling the polypeptides of the invention are known to the skilled artisan. When the polypeptides are used for *in vivo* diagnostic use, the tag can comprise, for example, magnetic resonance imaging (MRI) contrast agents, such as gadolinium diethylenetriaminepentaacetic acid, to ultrasound contrast agents or to X-ray contrast agents, or by radioisotopic labeling.

45 [0037] The polypeptides can also comprise a tag, such as a linker (including but not limited to an amino acid linker such as cysteine or lysine), for binding to a particle, such as a virus-like particle. As another example, the polypeptides

can usefully be attached to the surface of a microtiter plate for ELISA. The polypeptides can be fused to marker sequences to facilitate purification, as described in the examples that follow. Examples include, but are not limited to, the hexa-histidine tag, the myc tag or the flag tag.

**[0038]** A plurality of the polypeptides may be complexed to a dendrimer. Dendrimers are three dimensional, highly ordered oligomeric and/or polymeric compounds typically formed on a core molecule or designated initiator by reiterative reaction sequences adding the oligomers and/or polymers and providing an outer surface. Suitable dendrimers include, but are not limited to, "starburst" dendrimers and various dendrimer polycations. Methods for the preparation and use of dendrimers are well known to those of skill in the art.

**[0039]** The polypeptides may be fused (via recombinant or chemical means) via their N-terminus, C-terminus, or both N- and C-termini, to an oligomerization domain. Any suitable oligomerization domain can be used. For instance, the polypeptides are fused to GCN4 variants that form trimers (hence trimers or hexamers of the fused polypeptide could be displayed). The polypeptides may also be fused to a fibritin foldon domain that forms trimers. The oligomerization domain could be any protein that assembles into particles, including but not limited to particles made from a (non-viral) lumazine synthase protein and particles made from (non-viral) ferritin or ferritin-like proteins.

**[0040]** The polypeptides may be chemically conjugated to liposomes. For instance, the liposomes may contain a fraction of PEGylated lipid in which the PEG groups are functionalized to carry a reactive group, and the polypeptide is chemically linked to the reactive group on the PEG. Additional immune-stimulating compounds are included within the liposomes, either within the lipid layers or within the interior. In another non-limiting embodiment, specific cell-targeting molecules may be included on the surface of the liposome, including but not limited to molecules that bind to proteins on the surface of dendritic cells.

**[0041]** A plurality (ie: 2 or more; preferably at least 5, 10, 15, 20, 25, 50, 75, 90, or more copies) of the polypeptides may be present in a virus-like particle (VLP), to further enhance presentation of the polypeptide to the immune system. As used herein, a "virus-like particle" refers to a structure that in at least one attribute resembles a virus but which has not been demonstrated to be infectious. Virus-like particles in accordance with the invention do not carry genetic information encoding for the proteins of the virus-like particles. In general, virus-like particles lack a viral genome and, therefore, are noninfectious. In addition, virus-like particles can often be produced in large quantities by heterologous expression and can be easily purified. Preferably, the VLP comprises viral proteins that may undergo spontaneous self-assembly, including but not limited to recombinant proteins of adeno associated viruses, rotavirus, recombinant proteins of norwalkvirus, recombinant proteins of alphavirus, recombinant proteins of foot and mouth disease virus, recombinant proteins of retrovirus, recombinant proteins of hepatitis B virus, recombinant proteins of tobacco mosaic virus, recombinant proteins of flock house virus, and recombinant proteins of human papillomavirus, and Qbeta bacteriophage particles. For instance, the viral proteins could comprise hepatitis B core antigen particles. In other cases, the VLPs could be from lipid-enveloped viruses and include lipid as well as any suitable viral protein, including but not limited to proteins from chikungunya virus, or hepatitis B surface antigen proteins. Methods for producing and characterizing recombinantly produced VLPs have been described for VLPs from several viruses, as reviewed in US 20110236408; see also US 7,229,624. As described in the examples that follow, immunization in the context of a VLP with approximately 75 copies of the FFL\_001 polypeptide (SEQ ID NO:4) conjugated onto Hepatitis B (HepB) core antigen particles results in an increased immune response to the polypeptide.

The VLPs of the invention can be used as vaccines or antigenic formulations for treating or limiting RSV infection, as discussed herein. The VLPs may further comprise other scaffolds presenting other epitopes from RSVF or RSVG, or scaffolds presenting epitopes from additional RSV proteins, such as M, N, G, and/or SH.

**[0042]** The polypeptides may be present on a non-natural core particle, such as a synthetic polymer, a lipid micelle or a metal. Such core particles can be used for organizing a plurality of polypeptides of the invention for delivery to a subject, resulting in an enhanced immune response. By way of example, synthetic polymer or metal core particles are described in U.S. Pat. No. 5,770,380, which discloses the use of a calixarene organic scaffold to which is attached a plurality of peptide loops in the creation of an 'antibody mimic', and U.S. Pat. No. 5,334,394 describes nanocrystalline particles used as a viral decoy that are composed of a wide variety of inorganic materials, including metals or ceramics. Preferred metals in this embodiment include chromium, rubidium, iron, zinc, selenium, nickel, gold, silver, platinum. Preferred ceramic materials in this embodiment include silicon dioxide, titanium dioxide, aluminum oxide, ruthenium oxide and tin oxide. The core particles in this case may be made from organic materials including carbon (diamond). Preferred polymers include polystyrene, nylon and nitrocellulose. For this type of nanocrystalline particle, particles made from tin oxide, titanium dioxide or carbon (diamond) are particularly preferred. A lipid micelle may be prepared by any means known in the art. See US 7,229,624 and references disclosed therein.

**[0043]** In a second aspect, the present invention provides isolated nucleic acids encoding a polypeptide of the present invention. The isolated nucleic acid sequence may comprise RNA or DNA. As used herein, "isolated nucleic acids" are those that have been removed from their normal surrounding nucleic acid sequences in the genome or in cDNA sequences. Such isolated nucleic acid sequences may comprise additional sequences useful for promoting expression and/or purification of the encoded protein, including but not limited to polyA sequences, modified Kozak sequences, and

sequences encoding epitope tags, export signals, and secretory signals, nuclear localization signals, and plasma membrane localization signals. It will be apparent to those of skill in the art, based on the teachings herein, what nucleic acid sequences will encode the polypeptides of the invention.

**[0044]** In a third aspect, the present invention provides recombinant expression vectors comprising the isolated nucleic acid of any aspect of the invention operatively linked to a suitable control sequence. "Recombinant expression vector" includes vectors that operatively link a nucleic acid coding region or gene to any control sequences capable of effecting expression of the gene product. "Control sequences" operably linked to the nucleic acid sequences of the invention are nucleic acid sequences capable of effecting the expression of the nucleic acid molecules. The control sequences need not be contiguous with the nucleic acid sequences, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the nucleic acid sequences and the promoter sequence can still be considered "operably linked" to the coding sequence. Other such control sequences include, but are not limited to, polyadenylation signals, termination signals, and ribosome binding sites. Such expression vectors can be of any type known in the art, including but not limited plasmid and viral-based expression vectors. The control sequence used to drive expression of the disclosed nucleic acid sequences in a mammalian system may be constitutive (driven by any of a variety of promoters, including but not limited to, CMV, SV40, RSV, actin, EF) or inducible (driven by any of a number of inducible promoters including, but not limited to, tetracycline, ecdysone, steroid-responsive). The construction of expression vectors for use in transfecting prokaryotic cells is also well known in the art, and thus can be accomplished via standard techniques. (See, for example, Sambrook, Fritsch, and Maniatis, in: Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory Press, 1989; Gene Transfer and Expression Protocols, pp. 109-128, ed. E.J. Murray, The Humana Press Inc., Clifton, N.J.), and the Ambion 1998 Catalog (Ambion, Austin, TX). The expression vector must be replicable in the host organisms either as an episome or by integration into host chromosomal DNA. The expression vector may be a plasmid. However, the invention is intended to include other expression vectors that serve equivalent functions, such as viral vectors.

**[0045]** In a fourth aspect, the present invention provides host cells that have been transfected with the recombinant expression vectors disclosed herein, wherein the host cells can be either prokaryotic or eukaryotic. The cells can be transiently or stably transfected. Such transfection of expression vectors into prokaryotic and eukaryotic cells can be accomplished via any technique known in the art, including but not limited to standard bacterial transformations, calcium phosphate co-precipitation, electroporation, or liposome mediated-, DEAE dextran mediated-, polycationic mediated-, or viral mediated transfection. (See, for example, Molecular Cloning: A Laboratory Manual (Sambrook, et al., 1989, Cold Spring Harbor Laboratory Press; Culture of Animal Cells: A Manual of Basic Technique, 2nd Ed. (R.I. Freshney. 1987. Liss, Inc. New York, NY). A method of producing a polypeptide according to the invention is also disclosed. The method comprises the steps of (a) culturing a host cell of the invention under conditions conducive to the expression of the polypeptide, and (b) optionally, recovering the expressed polypeptide. The expressed polypeptide can be recovered from the cell free extract, but preferably they are recovered from the culture medium. Methods to recover polypeptide from cell free extracts or culture medium are well known to the man skilled in the art.

**[0046]** In a fifth aspect, the present invention provides pharmaceutical compositions comprising the polypeptide or the VLP of the invention, and a pharmaceutically acceptable carrier. Also disclosed are pharmaceutical compositions comprising one or more polypeptides, VLPs, nucleic acids, recombinant expression vectors, or host cells of the invention and a pharmaceutically acceptable carrier. Such pharmaceutical compositions can be used, for example, in the methods described below. The pharmaceutical composition may comprise in addition to the polypeptide of the invention (a) a lyoprotectant; (b) a surfactant; (c) a bulking agent; (d) a tonicity adjusting agent; (e) a stabilizer; (f) a preservative and/or (g) a buffer.

**[0047]** The buffer in the pharmaceutical composition could be chosen from a Tris buffer, a histidine buffer, a phosphate buffer, a citrate buffer or an acetate buffer. The pharmaceutical composition may also include a lyoprotectant, e.g. sucrose, sorbitol or trehalose. The pharmaceutical composition may include a preservative e.g. benzalkonium chloride, benzethonium, chlorohexidine, phenol, m-cresol, benzyl alcohol, methylparaben, propylparaben, chlorobutanol, o-cresol, p-cresol, chlorocresol, phenylmercuric nitrate, thimerosal, benzoic acid, and various mixtures thereof. The pharmaceutical composition may also include a bulking agent, like glycine. In yet other instances, the pharmaceutical composition may include a surfactant e.g., polysorbate-20, polysorbate-40, polysorbate- 60, polysorbate-65, polysorbate-80 polysorbate-85, poloxamer-188, sorbitan monolaurate, sorbitan monopalmitate, sorbitan monostearate, sorbitan monooleate, sorbitan trilaurate, sorbitan tristearate, sorbitan trioleaste, or a combination thereof. The pharmaceutical composition may also include a tonicity adjusting agent, e.g., a compound that renders the formulation substantially isotonic or isoosmotic with human blood. Exemplary tonicity adjusting agents include sucrose, sorbitol, glycine, methionine, mannitol, dextrose, inositol, sodium chloride, arginine and arginine hydrochloride. The pharmaceutical composition additionally may include a stabilizer, e.g., a molecule which, when combined with a protein of interest substantially prevents or reduces chemical and/or physical instability of the protein of interest in lyophilized or liquid form. Exemplary stabilizers include sucrose, sorbitol, glycine, inositol, sodium chloride, methionine, arginine, and arginine hydrochloride.

**[0048]** The polypeptides may be the sole active agent in the pharmaceutical composition, or the composition may

further comprise one or more other agents suitable for an intended use, including but not limited to adjuvants to stimulate the immune system generally and improve immune responses overall. Any suitable adjuvant can be used. The term "adjuvant" refers to a compound or mixture that enhances the immune response to an antigen. Exemplary adjuvants include, but are not limited to, Adju-Phos, Adjumer™, albumin-heparin microparticles, Algal Glucan, Algammulin, Alum, Antigen Formulation, AS-2 adjuvant, autologous dendritic cells, autologous PBMC, Avridine™, B7-2, BAK, BAY R1005, Bupivacaine, Bupivacaine-HCl, BWZL, Calcitriol, Calcium Phosphate Gel, CCR5 peptides, CFA, Cholera holotoxin (CT) and Cholera toxin B subunit (CTB), Cholera toxin A1-subunit-Protein A D-fragment fusion protein, CpG, CRL1005, Cytokine-containing Liposomes, D-Murapalmidine, DDA, DHEA, Diphtheria toxoid, DL-PGL, DMPC, DMPG, DOC/Alum Complex, Fowlpox, Freund's Complete Adjuvant, Gamma Inulin, Gerbu Adjuvant, GM-CSF, GMDP, hGM-CSF, hIL-12 (N222L), hTNF-alpha, IFA, IFN-gamma in pcDNA3, IL-12 DNA, IL-12 plasmid, IL-12/GMCSF plasmid (Sykes), IL-2 in pcDNA3, IL-2/Ig plasmid, IL-2/Ig protein, IL-4, IL-4 in pcDNA3, Imiquimod, ImmTher™, Immunoliposomes Containing Antibodies to Costimulatory Molecules, Interferon-gamma, Interleukin-1 beta, Interleukin-12, Interleukin-2, Interleukin-7, ISCOM(s)™, Iscoprep 7.0.3™, Keyhole Limpet Hemocyanin, Lipid-based Adjuvant, Liposomes, Loxoribine, LT(R192G), LT-OA or LT Oral Adjuvant, LT-R192G, LTK63, LTK72, MF59, MONTANIDE ISA 51, MONTANIDE ISA 15 720, MPL.TM., MPL-SE, MTP-PE, MTP-PE Liposomes, Murametide, Murapalmidine, NAGO, nCT native Cholera Toxin, Non-Ionic Surfactant Vesicles, non-toxic mutant E112K of Cholera Toxin mCT-E112K, p-Hydroxybenzoique acid methyl ester, pCIL-10, pCIL12, pCMVmCAT1, pCMVN, Peptomer-NP, Pleuran, PLG, PLGA, PGA, and PLA, Pluronic L121, PMMA, PODDS™, Poly rA: Poly rU, Polysorbate 80, Protein Cochleates, QS-21, Quadri A saponin, Quil-A, Rehydragel HPA, Rehydragel LV, RIBI, Ribilike adjuvant system (MPL, TMD, CWS), S-28463, SAF-1, Sclavo peptide, Sendai 20 Proteoliposomes, Sendai-containing Lipid Matrices, Span 85, Specol, Squalane 1, Squalene 2, Stearyl Tyrosine, Tetanus toxoid (TT), Theramide™, Threonyl muramyl dipeptide (TMDP), Ty Particles, and Walter Reed Liposomes. Selection of an adjuvant depends on the subject to be vaccinated. Preferably, a pharmaceutically acceptable adjuvant is used.

**[0049]** Compositions comprising the polypeptides can be stored in any standard form, including, e.g., an aqueous solution or a lyophilized cake. Such compositions are typically sterile when administered to cells or subjects. Sterilization of an aqueous solution is readily accomplished by filtration through a sterile filtration membrane. If the composition is stored in lyophilized form, the composition can be filtered before or after lyophilization and reconstitution.

**[0050]** Disclosed are methods for treating and/or limiting an RSV infection, comprising administering to a subject in need thereof a therapeutically effective amount of one or more polypeptides of the invention, salts thereof, conjugates thereof, VLPs thereof, or pharmaceutical compositions thereof, to treat and/or limit the RSV infection. The method may comprise eliciting an immune response in an individual having or at risk of an RSV infection, comprising administering to a subject in need thereof a therapeutically effective amount of one or more polypeptides of the invention, salts thereof, conjugates thereof, VLPs thereof, or pharmaceutical compositions thereof, to generate an immune response.

**[0051]** "Respiratory Syncytial Virus" and "RSV" refer to a negative-sense, single-stranded RNA virus of the family Paramyxoviridae that causes a respiratory disease, especially in children.

**[0052]** When the method comprises treating an RSV infection, the one or more polypeptides, VLPs, or compositions are administered to a subject that has already been infected with the RSV, and/or who is suffering from symptoms (including but not limited to lower respiratory tract infections, upper respiratory tract infections, bronchiolitis, pneumonia, fever, listlessness, diminished appetite, recurrent wheezing, and asthma) indicating that the subject is likely to have been infected with the RSV. As used herein, "treat" or "treating" means accomplishing one or more of the following: (a) reducing RSV titer in the subject; (b) limiting any increase of RSV titer in the subject; (c) reducing the severity of RSV symptoms; (d) limiting or preventing development of RSV symptoms after infection; (e) inhibiting worsening of RSV symptoms; (f) limiting or preventing recurrence of RSV symptoms in subjects that were previously symptomatic for RSV infection. In one embodiment method, polypeptides, VLPs, or compositions are used as "therapeutic vaccines" to ameliorate the existing infection and/or provide prophylaxis against infection with additional RSV virus.

**[0053]** When the method comprises limiting an RSV infection, the one or more polypeptides, VLPs, or compositions are administered prophylactically to a subject that is not known to be infected, but may be at risk of exposure to the RSV. As used herein, "limiting" means to limit RSV infection in subjects at risk of RSV infection. Groups at particularly high risk include children under age 18 (particularly infants 3 years or younger), adults over the age of 65, and individuals suffering from any type of immunodeficiency. In this method, the polypeptides, VLPs, or compositions are used as vaccines.

**[0054]** As used herein, a "therapeutically effective amount" refers to an amount of the polypeptide that is effective for treating and/or limiting RSV infection. The polypeptides are typically formulated as a pharmaceutical composition, such as those disclosed above, and can be administered via any suitable route, including orally, parentally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional pharmaceutically acceptable carriers, adjuvants, and vehicles. The term parenteral as used herein includes, subcutaneous, intravenous, intra-arterial, intra-muscular, intrasternal, intratendinous, intraspinal, intracranial, intrathoracic, infusion techniques or intraperitoneally. Polypeptide compositions may also be administered via microspheres, liposomes, immune-stimulating complexes (IS-COMs), or other microparticulate delivery systems or sustained release formulations introduced into suitable tissues

(such as blood). Dosage regimens can be adjusted to provide the optimum desired response (e.g., a therapeutic or prophylactic response). A suitable dosage range may, for instance, be 0.1 ug/kg-100 mg/kg body weight; alternatively, it may be 0.5 ug/kg to 50 mg/kg; 1 ug/kg to 25 mg/kg, or 5 ug/kg to 10 mg/kg body weight. The polypeptides can be delivered in a single bolus, or may be administered more than once (e.g., 2, 3, 4, 5, or more times) as determined by an attending physician.

**[0055]** As demonstrated in the examples that follow, RSV infectivity may be neutralized using the polypeptides of the invention. The polypeptides of the invention may prevent RSV from infecting host cells by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to infection of host cells by RSV in the absence of the polypeptides. Neutralization can be measured using standard techniques in the art.

**[0056]** Further disclosed is a pharmaceutical composition, comprising

(a) isolated nucleic acids, recombinant expression vectors, and/or recombinant host cells of the invention ; and  
 (b) a pharmaceutically acceptable carrier. In this aspect, the nucleic acids, expression vectors, and host cells of the invention can be used as polynucleotide-based immunogenic compositions, to express an encoded polypeptide in vivo, in a subject, thereby eliciting an immune response against the encoded polypeptide. Various methods are available for administering polynucleotides into animals. The selection of a suitable method for introducing a particular polynucleotide into an animal is within the level of skill in the art. Polynucleotides of the invention can also be introduced into a subject by other methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), or a DNA vector transporter (see, e.g., Wu et al. (1992) J. Biol. Chem. 267:963-967).

**[0057]** The immune response against the polypeptides, VLPs, or compositions of the invention can be generated by one or more inoculations of a subject with an immunogenic composition of the invention. A first inoculation is termed a "primary inoculation" and subsequent immunizations are termed "booster inoculations". Booster inoculations generally enhance the immune response, and immunization regimens including at least one booster inoculation are preferred. Any polypeptide, VLP, or composition of the invention may be used for a primary or booster immunization. The adequacy of the vaccination parameters chosen, e.g., formulation, dose, regimen and the like, can be determined by taking aliquots of serum from the subject and assaying antibody titers during the course of the immunization program. Alternatively, the T cell populations can be monitored by conventional methods. In addition, the clinical condition of the subject can be monitored for the desired effect, e.g., limiting RSV infection, improvement in disease state (e.g., reduction in viral load), etc. If such monitoring indicates that vaccination is sub-optimal, the subject can be boosted with an additional dose of composition, and the vaccination parameters can be modified in a fashion expected to potentiate the immune response. Thus, for example, the dose of the polypeptide, VLP, or composition, and/or adjuvant, can be increased or the route of administration can be changed.

**[0058]** Further disclosed are methods for monitoring an RSV-induced disease in a subject and/or monitoring response of the subject to immunization by an RSV vaccine, comprising contacting the polypeptides, the VLPs, or the pharmaceutical compositions of the invention with a bodily fluid from the subject and detecting RSV-binding antibodies in the bodily fluid of the subject. By "RSV-induced disease" is intended any disease caused, directly or indirectly, by RSV. The method comprises contacting a polypeptide, VLP, or composition of the invention with an amount of bodily fluid (such as serum, whole blood, etc.) from the subject; and detecting RSV-binding antibodies in the bodily fluid of the subject. The detection of the RSV binding antibodies allows the RSV disease in the subject to be monitored. In addition, the detection of RSV binding antibody also allows the response of the subject to immunization by an RSV vaccine to be monitored. In still other methods, the titer of the RSV binding antibodies is determined. Any suitable detection assay can be used, including but not limited to homogeneous and heterogeneous binding immunoassays, such as radioimmunoassays (RIA), ELISA, immunofluorescence, immunohistochemistry, FACS, BIACORE and Western blot analyses. The methods may be carried in solution, or the polypeptide(s) of the invention may be bound or attached to a carrier or substrate, e.g., microtiter plates (ex: for ELISA), membranes and beads, etc. Carriers or substrates may be made of glass, plastic (e.g., polystyrene), polysaccharides, nylon, nitrocellulose, or teflon, etc. The surface of such supports may be solid or porous and of any convenient shape. The polypeptides of the invention for use in this aspect may comprise a conjugate as disclosed above, to provide a tag useful for any detection technique suitable for a given assay.

**[0059]** Further disclosed are methods for detecting RSV binding antibodies, comprising

(a) contacting the polypeptides, the VLPs, or the compositions of the invention with a composition comprising a candidate RSV binding antibody under conditions suitable for binding of RSV antibodies to the polypeptide, VLP, or composition; and  
 (b) detecting RSV antibody complexes with the polypeptide, VLP, or composition. In this aspect, the methods are performed to determine if a candidate RSV binding antibody recognizes the RSV F epitope present in the polypeptides

of the invention. Any suitable composition may be used, including but not limited to bodily fluid samples (such as serum, whole blood, etc.) from a suitable subject (such as one who has been infected with RSV), naive libraries, modified libraries, and libraries produced directly from human donors exhibiting an RSV-specific immune response. The assays are performed under conditions suitable for promoting binding of antibodies against the polypeptides; such conditions can be determined by those of skill in the art based on the teachings herein. Any suitable detection assay can be used, including but not limited to homogeneous and heterogeneous binding immunoassays, such as radioimmunoassays (RIA), ELISA, immunofluorescence, immunohistochemistry, FACS, BIACORE and Western blot analyses. The methods may be carried in solution, or the polypeptide(s) of the invention may be bound or attached to a carrier or substrate, e.g., microtiter plates (ex: for ELISA), membranes and beads, etc. Carriers or substrates may be made of glass, plastic (e.g., polystyrene), polysaccharides, nylon, nitrocellulose, or teflon, etc. The surface of such supports may be solid or porous and of any convenient shape. The polypeptides of the invention for use in this aspect may comprise a conjugate as disclosed above, to provide a tag useful for any detection technique suitable for a given assay. The RSV F-binding antibodies may be isolated using standard procedures. The methods may comprise isolation of polypeptide-specific memory B cells by fluorescence activated cell sorting (FACS) using standard techniques in the art (see, for example, Science DOI: 10.1126/science. 1187659)

**[0060]** Further disclosed are methods for producing RSV antibodies, comprising

- (a) administering to a subject an amount effective to generate an antibody response of the polypeptides, the VLPs, and/or the compositions of the invention; and
- (b) isolating antibodies produced by the subject.

The polypeptides of the invention can also be used to generate antibodies that recognize the polypeptides of the invention. The method comprises administering to a subject a polypeptide, VLP, or composition of the invention. Such antibodies can be used, for example, in RSV research. A subject employed in this is one typically employed for antibody production, including but not limited to mammals, such as, rodents, rabbits, goats, sheep, etc. The antibodies generated can be either polyclonal or monoclonal antibodies. Polyclonal antibodies are raised by injecting (e.g. subcutaneous or intramuscular injection) antigenic polypeptides into a suitable animal (e.g., a mouse or a rabbit). The antibodies are then obtained from blood samples taken from the animal. The techniques used to produce polyclonal antibodies are extensively described in the literature. Polyclonal antibodies produced by the subjects can be further purified, for example, by binding to and elution from a matrix that is bound with the polypeptide against which the antibodies were raised. Those of skill in the art will know of various standard techniques for purification and/or concentration of polyclonal, as well as monoclonal, antibodies. Monoclonal antibodies can also be generated using techniques known in the art.

### 35 **Example 1**

#### *Fold From Loops*

**[0061]** The FFL protocol was implemented in the ROSETTA™ software package. The FFL protocol requires as starting inputs a structural motif and a pdb file of the target topology to be folded (**Fig 1**). Here, the input structural motif was the RSV peptide-epitope, which has been crystallized in complex with the Fab portion of Motavizumab (PDBid 3IXT). The 3-helix bundle target topology was selected based on the following criteria: the existence of a helix-turn-helix motif in the helical bundle; the high thermal stability of the protein used as the template target topology. The FFL protocol can be partitioned in two-main stages: I) a low-resolution stage with large conformational sampling; II) a full-atom stage with iterations of sequence design and confined conformational sampling.

#### *Low-resolution conformational sampling*

**[0062]** Extended polypeptide chains were appended to the termini of the input motif such that the total number of residues matches that of the template topology. The residues of the topology onto which the input motif were inserted were defined through a loop file; the location of the input motif determined the length of the polypeptide chains to append. At this low-resolution stage the representation of the polypeptide chain was coarse-grained; only the atoms of the main chain were explicitly represented and the side chains were represented as spheres.

**[0063]** The conformational sampling carried out at this stage was performed with a Rosetta™ *ab initio* type protocol. The *ab initio* protocol relies on the insertion of fragments extracted from a large set of known protein structures. The used fragments were collected with the program NNMAKE™, which builds fragment libraries according to a provided sequence and its secondary structure prediction. For the design work described here the fragments were derived from the sequence of the template.

[0064] Optionally, to bias the folding trajectory towards similar structures to that of the target topology, C alpha restraints are extracted [Rohl et al., Methods in Enzymology 383 (2004)] from the target topology and incorporated as a scoring term in the overall scoring function. The Rosetta™ folding protocol which incorporates distance restraints has been previously described by Rohl et al.. The distances between residues were collected if sequence separation was larger than 6 and if both residues were outside of the range of the functional site including five residues upstream and downstream. For the designs described here, a standard deviation of 3 Å for each C $\alpha$ -C $\alpha$  distance extracted from the template protein was allowed.

[0065] In general, throughout the sampling stages the backbone dihedral angles of the input motif were untouched; nevertheless the algorithm implementation allows setting the termini of the motif as moveable. The rationale to allow for these degrees of freedom within the input motif is to favor smooth structural transitions between the input motif and the remaining protein.

#### *Full-atom sequence design and structural optimization*

[0066] After the low-resolution stage during which a large conformational space was explored, the generated models were filtered according to their RMSD relative to the coordinates of native topology. Here, an RMSD threshold of 5 Å was used such that models in the structural vicinity of the target topology would be carried to the sequence design stage, in case a model was above the defined threshold it was automatically discarded. The original side-chain conformations, from the input motif, were recovered and kept fixed throughout the full-atom stage.

[0067] By default, in the sequence design stage the 19 amino-acids (cysteine excluded) were allowed in every residue of the models excluding the input functional motif. The exclusion of cysteine is not required. Generally, cysteines are not used for immunogen design unless disulfide bonds are being designed, since immunogens are intended for the extracellular environment which is oxidizing and hence unpaired cysteines in the extracellular environment will tend to form disulfides by paring with other cysteines. For protein design related to intracellular applications, unpaired cysteines are perfectly acceptable. Also at this stage, additional options were implemented for a finer control of the amino-acid identities allowed in particular residue positions. In the FFL designs described here, some positions of the input motif were not part of the antibody-contacting surface and therefore constituted part of the protein core; in some simulations the non contacting side-chains were allowed to change.

[0068] After each step of sequence design, a step of full-atom refinement (relax) was performed ensuring that the local conformational space was explored. The relax protocol is composed of several rounds of: small backbone perturbations; side-chain repacking; and energy minimization.

[0069] The cycles of iterative sequence design and structural minimization were repeated 3 times. The number of cycles is adjustable by the user.

#### 35 *Design Selection*

##### *Quality filters*

[0070] Typically 10,000 designs were generated by each FFL run, the first filter applied was based on Rosetta™ full-atom energy, and the best 50 designs by Rosetta Energy™ were further considered. Next, a composite filter was applied to select designs with the best structural features. The structural features considered were: Ramachandran score as implemented in Rosetta™; counts of buried polar atoms not involved in hydrogen bonds; and core packing assessment according to the RosettaHoles™ algorithm. Designed models within the top-25 according to the three features were taken to the next stage.

[0071] Some of the designs were selected according the geometrical properties of the models, in particular the bend angle of the helices. Statistics of the bend angles of each helix that composed the helical bundles were collected with the software Helanal™. Bundles with the lowest average bend angle were selected for next stage.

#### *Post-FFL design*

[0072] A long identified culprit of the Rosetta™ energy function is the notorious inability to accurately design solvent exposed residues; one of the possible causes is related with the absence of appropriate electrostatics parameterization. To circumvent this known culprit, the first step of post-FFL design was to impose identical residues in the solvent exposed positions as those of the protein used as the template topology.

[0073] Next, a step of manual design and all-atom refinement was employed to correct core-packing defects and remove polar amino acids from the core. The manual design was performed to: correct the occurrence of polar residues (in particular histidines) in the designs' cores; and designs straight from the automated stage having alanines overrepresented and valines underrepresented when compared to the template topology. The high frequency of alanines is

likely related to the energy term which represents the probability of finding a certain amino-acid given the dihedral angles of the backbone (Ramachandran term), given that the FFL proteins were mostly helical the Ramachandran term will favor the occurrence of alanines and disfavor valines or amino-acids with beta branched side-chains; another potential influencing factor is the proximal arrangement of the helices which in some positions might not allow to fit amino acids larger than alanine.

[0074] Given the nature of the iterative design procedure the generated models were highly intolerable to mutations in the core. Consequently, in the manual design stage, steric clashes were introduced by the mutations and a refinement step was necessary for accurate evaluation of the full-atom energies. The refinement served also as a filtering step to the performed manual mutations, as a given mutation or set of mutations would only be accepted if the full atom energy would recover significantly without causing major distortions in the helical local structure. One of the designs (FFL\_001) selected for experimental characterization was a straight FFL design with no manual changes performed in the core. This polypeptide was thermodynamically stable and bound the antibody of interest with high affinity.

[0075] The designs selected for experimental characterization differed by 6 to 48 mutations when compared to each other. When compared to the sequence of the original template the designs showed between 51 and 59 mutations. A sequence alignment of the FFL designs and the sequence of the original template is shown in **Figure 3**.

[0076] The backbone rmsds within the designed models were of 0.53 to 3.06 Å and between 1.83 and 2.91 when the designs are compared to the coordinate of the initial template.

#### *Design minimization*

[0077] To further reduce the size of the FFL designs, protein segments that didn't contact the mota epitope were eliminated.

### **Experimental Methods**

#### *Expression and Purification*

##### *Non-labeled protein*

[0078] DNA segments encoding scaffold constructs were synthesized with optimized codon usage and RNA structure (Codon Devices, Genscript Corp.), subcloned into pET29 (EMD Biosciences) and transformed into Arctic Express™ *E. coli* (Invitrogen). Single colonies were grown overnight at 37 °C in 10 mL Luria Broth (LB) plus Kanamycin (100 mg/ml). The starter cultures were expanded into 1 L of LB plus Kanamycin and incubated at 37 °C; when cells reached log phase, 250 µM of IPTG was added to the cultures to induce protein expression and the cells were then incubated overnight at 12 °C. Cultures were then pelleted and resuspended in start buffer (160 mM Imidazole, 4 M Sodium Chloride, 160 mM Sodium Phosphate), a tablet of protease inhibitor (Novagen) was added and the cell suspension was frozen at -20 °C.

[0079] The cell suspension was thawed and 10 ml of 10X Bugbuster™ (Novagen), 50 µL of Benzonase Nucleases and 1.7 µL of rLysozyme (Novagen) were added to lyse the cells; the cell suspension was then gently tumbled in an orbital shaker for 20 minutes. Lysed cells were pelleted and the supernatant was filtered through a 0.22 µm filter (Millipore). Supernatants were tumbled with 5 mL of Ni++ Sepharose 6 Fast Flow (GE Healthcare) for 1 hour at 4 °C. The resin was washed 3 times with 30 mL wash buffer (50 mM imidazole, 500 mM Sodium Chloride and 160 mM Sodium Phosphate) and eluted with 20 mL of Elution Buffer (250 mM Imidazole, 500 mM Sodium Chloride and 20 mM Sodium Phosphate). Fractions containing the construct of interest were combined and further purified by preparative size exclusion chromatography (SEC) on Superdex 75 16/60 (GE Healthcare) at room temperature in HBS. Collected fractions were analyzed on a 4-12% SDS denaturing gel (Invitrogen) and positive fractions were combined and concentrated by ultrafiltration (Vivaspin, Bioexpress). Protein concentration was determined by measuring UV absorption signal at 280 nm (Nano-drop™) and calculated from the theoretical extinction coefficient. To facilitate a rapid and accurate protein quantification the sequence GSW was added to all the designs without tryptophans in their primary sequence.

##### *Low-Endotoxin protein*

[0080] In order to prepare low-endotoxin protein for immunization studies, bacterial pellets were alternately resuspended in detergent buffer (50 mM NaH2PO4, 500 mM NaCl, 10 mM imidazole, 0.5 mg/ml lysozyme, 0.01 mg/ml DNase, 0.1% Triton X114) and Ni++ resin was alternately initially washed in 10 mM imidazole, 50 mM NaH2PO4, 500 mM NaCl, 0.1% Triton X114.

*N15-labeled protein*

[0081] Isotopically labeled samples of FFL\_001, FFL\_005, FFL\_006 and FFL\_007 were grown in minimal MOPS medium supplemented with 1g/L of <sup>15</sup>N ammonium chloride. The starter cultures were expanded to 1L of MOPS and were incubated overnight at 37 °C; 3ml of 40% <sup>15</sup>N glucose was added to continue growth, upon reaching an OD<sub>λ=600</sub> 250 µM of IPTG was added to the cultures to induce protein expression and the cells were then incubated overnight at 16 °C.

## Light Scattering

[0082] The monodispersity and molecular weight of purified proteins were further assessed by HPLC (Agilent, 1200 series) coupled in-line to a static light scattering device (miniDAWN TREOS, Wyatt). 100 µl of 1-2 mg/mL protein sample was used and the collected data was analyzed with the ASTRA™ software (Wyatt).

## Circular Dichroism

[0083] Solution thermostabilities (T<sub>m</sub>) were determined by circular dichroism (CD) on an Aviv 62A DS spectrometer. Far-UV wavelength scans (190-260 nm) of 15 to 25 µM protein were collected in a 1 mm path length cuvette. Temperature-induced protein denaturation was followed by change in ellipticity at 210 nm. Experiments were carried over a temperature range from 1-99 °C, with 2 °C increments every 3 minutes, and the resulting data was converted to mean residue ellipticity and fitted to a two-state model.


## NMR

[0084] NMR samples were prepared in 25mM sodium phosphate, 150mM NaCl, pH 7.0, and 90% H<sub>2</sub>O/10% D<sub>2</sub>O at a concentration of 500µM. HSQC spectra for FFL\_001, FFL\_005, FFL\_006 and FFL\_007 were recorded on a Bruker Avance™ 600 MHz NMR spectrometer equipped with an actively shielded z-gradient triple resonance cryo-probe. All spectra were recorded at 25°C. Spectra were processed using NMRPipe™ and NMRView™ (1,2).

## Surface plasmon resonance

[0085] All experiments were carried out on a Biacore 2000 (GE Healthcare) at 25 °C with HBSEP (0.01 M HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA and 0.005% (v/v) Surfactant P20) (GE Healthcare) as running buffer. For binding analysis, 200-500 response units (RUs) of Motavizumab IgG were captured on a CM5 sensor chip containing 8000-9000 RUs of amine-linked mouse anti-human IgG (Human Antibody Capture kit, GE Healthcare). Samples of different protein concentrations were injected in duplicates over this surface at a flow rate of 50-100 µl/min. If necessary, surface regeneration was performed with two 60 seconds injections of 3 M MgCl<sub>2</sub> at a flow rate of 10 µl/min. One flow cell contained anti-human IgG only and its interaction with the analyte was used as reference. Alternatively, another format was used where the epitope where amine-coupled to the sensor chip and Mota Fab was injected with identical flow rates as mentioned above.

Data preparation and analysis were performed using Scrubber 2.0 (BioLogic Software). For kinetic analysis, biosensor data were globally fit to a mass transport limited simple bimolecular binding model:



where A<sub>0</sub> represents injected analyte.

## Results for Example 1

[0086] The structure of the RSV F site A peptide bound to Motavizumab was used as the target binding site for scaffolding. (Figure 2) This peptide structure (chain P in PDBID: 3IXT) is a helix-turn-helix motif and that led us to choose a three-helix bundle as the target topology for these scaffolds. The structure of PDBID: 3LHP, chain S was selected as the particular three-helix bundle. Procedural details of the FFL designs are shown in Table 1. The different parameters and filtering criteria used on the FFL simulations are summarized. The manual intervention stage is also summarized relative to: number of core mutations performed, the initial Rosetta energy of the designs and the Rosetta energy after

the mutations and the full-atom refinement step.

Table 1

Design	Computational Algorithm		Filtering			Manual Intervention	
	SD (Å) <sup>a</sup>	BS design <sup>b</sup>	Energy + Composite Filters	Helix Bend <sup>c</sup>	Rosetta energy	Mutations	Rosetta energy (post-relax) <sup>d</sup>
FFL_001	1.5	✗	✓	✗	-	-	-
FFL_002	3.0	✓	✓	✗	-289	10	-276
FFL_003	3.0	✓	✓	✗	-286	4	-293
FFL_004	3.0	✓	✓	✗	-285	7	-291
FFL_005	1.5	✓	✗	✓	-292	11	-287
FFL_006	3.0	✓	✓	✗	-291	3	-290
FFL_007	1.5	✓	✗	✓	-293	11	-285
FFL_008	1.5	✓	✗	✓	-293	8	-286

<sup>a</sup> SD - standard deviation allowed to the constraints derived from target topology

<sup>b</sup> BS - Binding site design of the positions that are not in direct contact with the antibody

<sup>c</sup> Filtering criteria based on the helix bend angle as implemented in Helanal

<sup>d</sup> Rosetta energy after the mutations have been performed and a step of full atom optimization

**[0087]** Seven scaffolds were designed using the Fold From Loops method (**Table 2**), and their sequences are shown in **Figure 3** along with the sequence of "T93", the template three-helix bundle from 3LHP.

Table 2: Protein scaffolds for the RSV F protein site A epitope

Protein Scaffold	Reference No.
RSVF_siteA_001	FFL 001 (SEQ ID NO:4)
RSVF_siteA_002	FFL 002 (SEQ ID NO:5)
RSVF_siteA_003	FFL 003 (SEQ ID NO: 10)
RSVF_siteA_004	FFL 004 (SEQ ID NO:6)
RSVF_siteA_005	FFL 005 (SEQ ID NO:7)
RSVF_siteA_006	FFL 006 (SEQ ID NO:8)
RSVF_siteA_007	FFL 007 (SEQ ID NO:9)
RSVF_siteA_008	FFL 008 (SEQ ID NO:12)

**[0088]** The designed sequences differ from each other between 8 and 42 mutations. The structural diversity of the computational models varies from each other between 0.53 Å and 3.06 Å.

**[0089]** To assess expression and solubility, the recombinant proteins were expressed in E. Coli, these 7 designed variants were soluble and purifiable through steps of metal affinity chromatography (Ni++) and size exclusion chromatography (SEC) and the yields of expression ranged from 3 to 5 mg L<sup>-1</sup>. To assess the oligomerization state in solution, the seven soluble designs were analyzed by SEC and static light scatter (Figure 4). Six designs were monodisperse and exhibit an apparent molecular weight corresponding to the monomeric protein ( $\approx$ 15 kDa). To evaluate the folding and the thermal stability of the designed molecules we performed circular dichroism spectroscopy (CD) (Figure 5). The six monomeric designs showed typical CD spectra of properly folded helical proteins. Temperature induced denaturation was followed by CD showing that the stability of the designs ranged from 48 to more than 100 °C (**Table 3**). To test whether the functional site (mota epitope) was recreated with fidelity, binding affinities were assessed by Surface Plasmon Resonance (SPR) experiments (Figure 6). The binding constants ( $K_D$ ) were within 30 and 652 pM. (See Table 3) The scaffolds were coupled to the biacore chip and Motavizumab was used as analyte. The binding interaction was readily blocked by a point mutation in the epitope region (K28E), previously reported to have the same effect on the RSV context, therefore showing that the binding specificity was directed to the epitope (Figure 7).

**[0090]** The affinities shown by the best FFL designs represent an improvement by a factor of approximately 7000 over

a previously published  $K_D$  for the peptide-epitope (200 nM). Recently, a side chain grafting strategy was utilized to transplant the RSV epitope to other heterologous scaffolds (McLellan et al., J. Mol. Biol. (2011) 409, 853-866). In that work, the highest affinity design showed a  $K_D$  of 60 nM to the mota antibody. Therefore the FFL designs had KDs improved by a factor of approximately 2000 over the results of McLellan et al.

[0091] To obtain an orthogonal characterization of the solution behavior and structural properties of the designed molecules,  $^{15}\text{N}-^1\text{H}$  hetero-nuclear single-quantum coherence (HSQC) spectra were collected. These spectra showed good peak dispersion typical of protein with well-defined globular folds (not shown). To further evaluate the accuracy of our computational design, an x-ray structure of FFL\_005 was solved. The computational model and the crystal structure of FFL\_005 (not shown) were in close agreement (1.7 Å rmsd over the backbone atoms), demonstrating the validity of the computational methods for designing polypeptides with a desired structural motif and three-dimensional structure. Furthermore, the conformation of the Mota epitope within the crystal structure of FFL\_005 matched the conformation of the Mota-bound peptide from PDB: 3ixt with a rmsd of 0.5 Å, supporting the claim that the FFL method can stabilize the conformation of a structural motif employed as a folding nucleus.

[0092] These studies demonstrate successful creation of novel functional proteins by coupling the *in silico* folding process and sequence design to simultaneously optimize the functional moiety of the molecule and the thermodynamic stability. The described computational strategy is general and flexible such that the target topology is not required to be a naturally occurring protein and back of the envelope topologies can also be used for the design of functionalized proteins. Regarding the structural complexity of the functional sites, the FFL algorithm is suited to deal with discontinuous motifs composed by multiple backbone segments, which are typically required in functional sites of naturally occurring proteins. These results have broad implications for the computational design of functional proteins and the usage of existing protein structures as potential templates.

**Table 3. Mota binding affinities and thermal stabilities of the FFL designs. The binding affinities were assessed by SPR and the thermal stabilities by CD spectroscopy**

Molecule	$T_m$ (°C)	SPR		
		$k_{on}$ (M $^{-1}$ s $^{-1}$ )	$k_{off}$ (s $^{-1}$ )	$k_{off}/k_{on}$ (pM)
FFL_001	75	$3.99 \times 10^6$	$1.19 \times 10^{-4}$	29.98
FFL_002	48	$1.56 \times 10^6$	$7.34 \times 10^{-4}$	469.9
FFL_004	>85	$1.05 \times 10^6$	$8.32 \times 10^{-4}$	795
FFL_005	>100	$2.97 \times 10^6$	$2.09 \times 10^{-4}$	70.3
FFL_006	>85	$3.57 \times 10^6$	$2.32 \times 10^{-4}$	651.9
FFL_007	>85	$1.45 \times 10^6$	$1.36 \times 10^{-4}$	94.1

## Example 2

[0093] FFL\_001 scaffolds were conjugated to the surface of HepBcAg particles to improve immune responses to the epitope. The scaffolds were conjugated via hetero-bifunctional cross-linkers between an engineered cysteine in the scaffold at the opposite end from the epitope, and an engineered lysine on the tip of the major immunodominant region of HepBcAg. This oriented the scaffolds in such a way that the epitope was exposed at the radial exterior of the conjugated particle.

[0094] Particles from HBcAg residues 1-149, a construct that leads to higher expression in bacteria and a predominance of the larger  $T = 4$  particle with 240 HepBcAg monomers (Zlotnick et al., 1996; Wynne et al., 1999), were expressed in *E. coli* and purified via standard sucrose gradients. For chemical coupling of monomeric FFL immunogens, pure lysine-functionalized HBcAg(1-149) particles were expressed and purified using standard techniques, in which a lysine residue was engineered into the tip of the immunodominant spike of every subunit. HBcAg(1-149) WT and lysine-functionalized particles were both full size (30 nm). (Figure 8)

[0095] Conjugation of FFL\_001 scaffolds and HepBcAG were carried out under standard conditions using a 10% Sucrose and 1% CHAPS, resulting in approximately 75 FFL\_001 scaffolds attached to each HepB particle, according to densitometry analysis of SDS-PAGE gels run on purified fractions from sucrose gradient ultracentrifugation. Binding Mota to the FFL\_001-conjugated particles was evaluated by SPR by capturing Mota IgG on the sensor chip and then binding FFL\_001-particles to the Mota IgG-coated surface; subsequently Mota Fab was used as analyte and the kinetics of Mota Fab binding to Mota-IgG-captured-FFL\_001-particles were evaluated; in this manner it was confirmed that the FFL\_001-conjugated-particles bound to Mota with similar high affinity as FFL\_001 monomers (data not shown).

[0096] Table 4 summarizes results from a macaque immunization experiment with FFL scaffold monomers and FFL\_001-conjugated-HepBcAg-particles. Immunogens were scaffold monomers labeled "001", "005", and "007", and

FFL\_001-conjugated-HepBcAg-particles labeled "001-particle". Rhesus macaques (4 animals per immunogen) were immunized by the intramuscular route at 0, 1 and 2 months. Animals were injected with 1 mL total volume of antigen mixed with Adjuplex™ adjuvant, with 0.5 mL injected into each arm. The first immunization included a total of 200 ug of scaffold; subsequent immunizations included a total of 100 ug scaffold. "Naïve" sera was taken from each animal on day 0 before the first immunization. "Imm3" sera was taken from each animal 2 weeks after the 3<sup>rd</sup> immunization. Both the "Naïve" and the "Imm3" sera were evaluated for neutralization in a standard plaque reduction assay at a serum dilution of 1:20. Each sample was run in duplicate (counts for the two individual runs are shown as "Naïve1", "Naïve2", "Imm3\_1", and "Imm3\_2". The average plaque counts "Naïve\_ave" and "Imm3\_ave" were computed from the two runs. The % plaque reduction was calculated as (Naïve\_ave - Imm3\_ave)/Naïve\_ave. The sera were also tested for ELISA reactivity to recombinant RSVF protein. The endpoint titers are given for each animal. The % plaque reduction numbers show a modest linear correlation with the ELISA titers, with a Pearson coefficient of 0.58.

[0097] These data demonstrate that macaque immunization with FFL scaffold monomers or FFL scaffolds presented on HepBcAg particles can result in the production of RSVF-binding antibodies and RSV neutralizing antibodies. The % neutralization (% plaque reduction) was as high as 88% for particle-displayed scaffolds, and as high as 72% for monomeric scaffolds. The average % plaque reduction for VLP-presented scaffolds was 51±25%, which was higher than the average for any of the monomer samples, the highest % plaque reduction for a monomer sample being 33±11% for FFL\_001 monomers. The difference in the average % plaque reduction for particle-001 compared to monomer-001 was not statistically significant. The average RSVF ELISA titer was also higher for the particle-001 sample (94000 ± 20000) compared to the highest titer monomer sample (79000 ± 66000 for FFL\_001), but again the difference was not statistically significant.

Table 4

Immunogen	NHP id	% plaque reduction	RSVF ELISA titer	Naïve_ave	Naïve 1	Naïve 2	Imm3_ave	Imm3_1	Imm3_2
001-particle	D039	0.8757	120000	84.5	83	86	10.5	13	8
007	D030	0.717	110000	79.5	80	79	22.5	21	24
005	D180	0.5924	22000	78.5	72	85	32	35	29
001	C012	0.4536	170000	91.5	89	94	50	46	54
001-particle	C004	0.4491	93000	83.5	74	93	46	50	42
001-particle	D130	0.4207	91000	72.5	68	77	42	43	41
007	D227	0.3716	23000	91.5	84	99	57.5	57	58
007	C010	0.3709	36000	75.5	70	81	47.5	44	51
001	D027	0.3642	12000	75.5	81	70	48	46	50
001	D052	0.3243	65000	92.5	89	96	62.5	63	62
001-particle	D184	0.3038	72000	79	72	86	55	52	58
005	D104	0.2	8500	67.5	64	71	54	54	54
001	D172	0.1954	69000	87	88	86	70	65	75
005	D190	0.1931	4800	72.5	68	77	58.5	50	67
005	D032	0.1243	9500	88.5	85	92	77.5	78	77
007	D087	0.047	15000	74.5	73	76	71	70	72

## SEQUENCE LISTING

[0098]

<110> University of Washington  
Schieff, William  
Correia, Bruno

<120> Polypeptides And Their Use In Treating And Limiting Respiratory Syncytial Virus Infection

<130> 11-1225-PCT3

5 <150> 61/620804  
<151> 2012-04-05

<160> 37

10 <170> PatentIn version 3.5

<210> 1  
<211> 115

<212> PRT

15 <213> Artificial Sequence

<220>  
<223> Synthetic

20 <220>  
<221> MISC\_FEATURE  
<222> (1).. (2)  
<223> optionally absent

25 <220>  
<221> MISC\_FEATURE  
<222> (3)..(3)  
<223> X is M, L, R, or F

30 <220>  
<221> MISC\_FEATURE  
<222> (6).. (6)  
<223> X is R, A, I, M, V or L

35 <220>  
<221> MISC\_FEATURE  
<222> (7)..(7)  
<223> X is R or M

40 <220>  
<221> MISC\_FEATURE  
<222> (10) ..(10)  
<223> X is L, A, or V

45 <220>  
<221> MISC\_FEATURE  
<222> (12)..(12)  
<223> X is E, R, K or D

50 <220>  
<221> MISC\_FEATURE  
<222> (14).. (14)  
<223> X is L, F, I or A

55 <220>  
<221> MISC\_FEATURE  
<222> (17)..(17)  
<223> X is L, F, or A

5           <220>  
 <221> MISC\_FEATURE  
 <222> (18)..(18)  
 <223> X is L, V, F, or M  
  
 10          <220>  
 <221> MISC\_FEATURE  
 <222> (21)..(21)  
 <223> X is A, V, F, I or L  
  
 15          <220>  
 <221> MISC\_FEATURE  
 <222> (25)..(25)  
 <223> X is E, M, L, F, W, or V  
  
 20          <220>  
 <221> MISC\_FEATURE  
 <222> (28)..(28)  
 <223> X is F, M, E, I, or V  
  
 25          <220>  
 <221> MISC\_FEATURE  
 <222> (32)..(32)  
 <223> X is M, L, F, or I  
  
 30          <220>  
 <221> MISC\_FEATURE  
 <222> (35)..(35)  
 <223> X is R, E, G, D, or Q  
  
 35          <220>  
 <221> MISC\_FEATURE  
 <222> (36)..(36)  
 <223> X is G, D, P, W, E, or Q  
  
 40          <220>  
 <221> MISC\_FEATURE  
 <222> (37)..(37)  
 <223> X is Q, I, P, K or F  
  
 45          <220>  
 <221> MISC\_FEATURE  
 <222> (38)..(38)  
 <223> X is R, Q, K, S, G or H  
  
 50          <220>  
 <221> MISC\_FEATURE  
 <222> (42)..(42)  
 <223> X is M, K, R or A  
  
 55          <220>  
 <221> MISC\_FEATURE  
 <222> (45)..(45)  
 <223> X is W, L, K or M  
  
 60          <220>  
 <221> MISC\_FEATURE  
 <222> (46)..(46)

<223> X is A, M, K or F  
  
5       <220>  
      <221> MISC\_FEATURE  
      <222> (48)..(48)  
      <223> X is I, F, L, E, K or D  
  
10      <220>  
      <221> MISC\_FEATURE  
      <222> (49)..(49)  
      <223> X is A, V, M, F, or L  
  
15      <220>  
      <221> MISC\_FEATURE  
      <222> (51)..(51)  
      <223> X is D, K, E, or Y  
  
20      <220>  
      <221> MISC\_FEATURE  
      <222> (53)..(53)  
      <223> X is F, V, A, R, M or L  
  
25      <220>  
      <221> MISC\_FEATURE  
      <222> (56)..(56)  
      <223> X is F, L, A, V or M  
  
30      <220>  
      <221> MISC\_FEATURE  
      <222> (58)..(58)  
      <223> X is K or R  
  
35      <220>  
      <221> MISC\_FEATURE  
      <222> (60)..(60)  
      <223> X is V, M or I  
  
40      <220>  
      <221> MISC\_FEATURE  
      <222> (63)..(63)  
      <223> X is F, R, V, I, A, or Y  
  
45      <220>  
      <221> MISC\_FEATURE  
      <222> (64)..(64)  
      <223> X is L, E, or A  
  
50      <220>  
      <221> MISC\_FEATURE  
      <222> (67)..(67)  
      <223> X is A or L  
  
55      <220>  
      <221> MISC\_FEATURE  
      <222> (72)..(72)  
      <223> X is Nor Y  
  
      <220>

<221> MISC\_FEATURE  
<222> (74)..(74)  
<223> X is Y, M or L

5 <220>  
<221> MISC\_FEATURE  
<222> (78)..(78)  
<223> X is Nor I

10 <220>  
<221> MISC\_FEATURE  
<222> (80)..(80)  
<223> X is D, Q, or K

15 <220>  
<221> MISC\_FEATURE  
<222> (82)..(82)  
<223> X is K, E, T, M or Q

20 <220>  
<221> MISC\_FEATURE  
<222> (84)..(84)  
<223> X is T, I, M, V or A

25 <220>  
<221> MISC\_FEATURE  
<222> (87)..(87)  
<223> X is D or K

30 <220>  
<221> MISC\_FEATURE  
<222> (88)..(88)  
<223> X is A, T, L, V or I

35 <220>  
<221> MISC\_FEATURE  
<222> (89)..(89)  
<223> X is K, L or I

40 <220>  
<221> MISC\_FEATURE  
<222> (91)..(91)  
<223> X is F, Y, K, E, R, or L

45 <220>  
<221> MISC\_FEATURE  
<222> (92)..(92)  
<223> X is D, A, V, or M

50 <220>  
<221> MISC\_FEATURE  
<222> (94)..(94)  
<223> X is E, I or R

55 <220>  
<221> MISC\_FEATURE  
<222> (95)..(95)  
<223> X is V, A, M or L

<220>  
<221> MISC\_FEATURE  
<222> (96)..(96)  
<223> X is A, E, K, F or M

5

<220>  
<221> MISC\_FEATURE  
<222> (99)..(99)  
<223> X is L, I or V

10

<220>  
<221> MISC\_FEATURE  
<222> (101)..(101)  
<223> X is A or L

15

<220>  
<221> MISC\_FEATURE  
<222> (102)..(102)  
<223> X is F, M, L, E or I

20

<220>  
<221> MISC\_FEATURE  
<222> (103)..(103)  
<223> X is K, A, V, M, I or L

25

<220>  
<221> MISC\_FEATURE  
<222> (106)..(106)  
<223> X is A, V, or I

30

<220>  
<221> MISC\_FEATURE  
<222> (108)..(108)  
<223> X is E, R, D, K, I or A

35

<220>  
<221> MISC\_FEATURE  
<222> (109)..(109)  
<223> X is A, M, K, L or W

40

<220>  
<221> MISC\_FEATURE  
<222> (110)..(110)  
<223> X is A, F, V, or K

45

<220>  
<221> MISC\_FEATURE  
<222> (113)..(115)  
<223> optionally absent

50

<400> 1

55

Gly Ser Xaa Ser Asp Xaa Xaa Lys Asp Xaa Glu Xaa Arg Xaa Asp Lys  
 1 5 10 15

5 Xaa Xaa Glu Ala Xaa Lys Asn Lys Xaa Asp Lys Xaa Lys Ala Ala Xaa  
 20 25 30

10 Arg Lys Xaa Xaa Xaa Xaa Glu Glu Arg Xaa Lys Asp Xaa Xaa Lys Xaa  
 35 40 45

15 Xaa Arg Xaa Glu Xaa Glu Gln Xaa Arg Xaa Ala Xaa Arg Asn Xaa Xaa  
 50 55 60

20 Ser Glu Xaa Leu Ser Lys Ile Xaa Asp Xaa Pro Ile Thr Xaa Asp Xaa  
 65 70 75 80

25 Lys Xaa Leu Xaa Ser Asn Xaa Xaa Xaa Lys Xaa Xaa Ala Xaa Xaa Xaa  
 85 90 95

30 Lys Lys Xaa Glu Xaa Xaa Xaa Ala Asp Xaa Glu Xaa Xaa Xaa Thr Gln  
 100 105 110

25 Gly Ser Trp  
 115

30 <210> 2  
 <211> 115  
 <212> PRT  
 <213> Artificial Sequence

35 <220>  
 <223> Synthetic

40 <220>  
 <221> MISC\_FEATURE  
 <222> (1)..(2)  
 <223> optionally absent

45 <220>  
 <221> MISC\_FEATURE  
 <222> (3)..(3)  
 <223> X is M, L, R or F

50 <220>  
 <221> MISC\_FEATURE  
 <222> (6)..(6)  
 <223> X is R, A, I, M or V

55 <220>  
 <221> MISC\_FEATURE  
 <222> (10)..(10)  
 <223> X is L, A, of V

<220>

<221> MISC\_FEATURE  
<222> (12)..(12)  
<223> X is E, R, K or D

5        <220>  
      <221> MISC\_FEATURE  
      <222> (14)..(14)  
      <223> X is L, F, I or A

10      <220>  
      <221> MISC\_FEATURE  
      <222> (17)..(17)  
      <223> X is L, F, or A

15      <220>  
      <221> MISC\_FEATURE  
      <222> (18)..(18)  
      <223> X is L, V, or F

20      <220>  
      <221> MISC\_FEATURE  
      <222> (21)..(21)  
      <223> X is A, V, F, or L

25      <220>  
      <221> MISC\_FEATURE  
      <222> (25)..(25)  
      <223> X is M, L, F or V

30      <220>  
      <221> MISC\_FEATURE  
      <222> (28)..(28)  
      <223> X is F, M, E or I

35      <220>  
      <221> MISC\_FEATURE  
      <222> (32)..(32)  
      <223> X is M, L, F or I

40      <220>  
      <221> MISC\_FEATURE  
      <222> (35)..(35)  
      <223> X is R, E, G or D

45      <220>  
      <221> MISC\_FEATURE  
      <222> (36)..(36)  
      <223> X is G, D, P, W, or Q

50      <220>  
      <221> MISC\_FEATURE  
      <222> (37)..(37)  
      <223> X is Q, I, P, or F

55      <220>  
      <221> MISC\_FEATURE  
      <222> (38)..(38)  
      <223> X is R, Q, K, S or H

5           <220>  
          <221> MISC\_FEATURE  
          <222> (42)..(42)  
          <223> X is M, K, R or A

10           <220>  
          <221> MISC\_FEATURE  
          <222> (45)..(45)  
          <223> X is W, L, M, or K

15           <220>  
          <221> MISC\_FEATURE  
          <222> (46)..(46)  
          <223> X is A, M, K or F

20           <220>  
          <221> MISC\_FEATURE  
          <222> (48)..(48)  
          <223> X is I, F, L, E, K or D

25           <220>  
          <221> MISC\_FEATURE  
          <222> (49)..(49)  
          <223> X is A, V, M, F or L

30           <220>  
          <221> MISC\_FEATURE  
          <222> (51)..(51)  
          <223> X is D, K, Y or E

35           <220>  
          <221> MISC\_FEATURE  
          <222> (53)..(53)  
          <223> X is F, V, A, R, or M

40           <220>  
          <221> MISC\_FEATURE  
          <222> (56)..(56)  
          <223> X is F, L, V, A, or M

45           <220>  
          <221> MISC\_FEATURE  
          <222> (58)..(58)  
          <223> X is K or R

50           <220>  
          <221> MISC\_FEATURE  
          <222> (60)..(60)  
          <223> X is V, M or I

55           <220>  
          <221> MISC\_FEATURE  
          <222> (63)..(63)  
          <223> X is F, R, V, I or Y

          <220>  
          <221> MISC\_FEATURE  
          <222> (64)..(64)

<223> X is L, E or A

<220>

<221> MISC\_FEATURE

5 <222> (67)..(67)  
<223> X is A or L

<220>

<221> MISC\_FEATURE

10 <222> (74)..(74)  
<223> X is Y, M, or L

<220>

<221> MISC\_FEATURE

15 <222> (80)..(80)  
<223> X is D, Q or K

<220>

<221> MISC\_FEATURE

20 <222> (84)..(84)  
<223> X is T, I, M, V, or A

<220>

<221> MISC\_FEATURE

25 <222> (88)..(88)  
<223> X is T, L, V or I

<220>

<221> MISC\_FEATURE

30 <222> (89)..(89)  
<223> X is K, L, or I

<220>

<221> MISC\_FEATURE

35 <222> (91)..(91)  
<223> X is F, Y, K, E or L

<220>

<221> MISC\_FEATURE

40 <222> (92)..(92)  
<223> X is D, A, V or M

<220>

<221> MISC\_FEATURE

45 <222> (94)..(94)  
<223> X is E, I or R

<220>

<221> MISC\_FEATURE

50 <222> (95)..(95)  
<223> X is V, A, M or L

<220>

<221> MISC\_FEATURE

55 <222> (96)..(96)  
<223> X is E, K, F or W

<220>

<221> MISC\_FEATURE  
<222> (99)..(99)  
<223> X is L, I or V

5 <220>  
<221> MISC\_FEATURE  
<222> (101)..(101)  
<223> X is A or L

10 <220>  
<221> MISC\_FEATURE  
<222> (102)..(102)  
<223> X is F, M, L, E or

15 <220>  
<221> MISC\_FEATURE  
<222> (103)..(103)  
<223> X is K A V M or I

20 <220>  
<221> MISC\_FEATURE  
<222> (106)..(106)  
<223> X is A, V or I

25 <220>  
<221> MISC\_FEATURE  
<222> (108)..(108)  
<223> X is E, R, D, K, I or A

30 <220>  
<221> MISC\_FEATURE  
<222> (109)..(109)  
<223> X is A, M, K, L or W

35 <220>  
<221> MISC\_FEATURE  
<222> (110)..(110)  
<223> X is A, E or V

40 <220>  
<221> MISC\_FEATURE  
<222> (113)..(115)  
<222> optionally absent

15 <100> 3

Gly Ser Xaa Ser Asp Xaa Arg Lys Asp Xaa Glu Xaa Arg Xaa Asp Lys  
1 5 10 15

50 Xaa Xaa Glu Ala Xaa Lys Asn Lys Xaa Asp Lys Xaa Lys Ala Ala Xaa  
20 25 30

Arg Lys Xaa Xaa Xaa Xaa Glu Glu Arg Xaa Lys Asp Xaa Xaa Lys Xaa  
 35 40 45

5 Xaa Arg Xaa Glu Xaa Glu Gln Xaa Arg Xaa Ala Xaa Arg Asn Xaa Xaa  
 50 55 60

10 Ser Glu Xaa Leu Ser Lys Ile Asn Asp Xaa Pro Ile Thr Asn Asp Xaa  
 65 70 75 80

15 Lys Lys Leu Xaa Ser Asn Asp Xaa Xaa Lys Xaa Xaa Ala Xaa Xaa Xaa  
 85 90 95

15 Lys Lys Xaa Glu Xaa Xaa Xaa Ala Asp Xaa Glu Xaa Xaa Xaa Thr Gln  
 100 105 110

20 Gly Ser Trp  
 115

<210> 3

<211> 115

25 <212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic

30

<220>

<221> MISC\_FEATURE

<222> (1)..(2)

<223> optionally absent

35

<220>

<221> MISC\_FEATURE

<222> (3)..(3)

<223> X is M, L, R or F

40

<220>

<221> MISC\_FEATURE

<222> (6)..(6)

<223> X is I or M

45

<220>

<221> MISC\_FEATURE

<222> (10)..(10)

<223> X is L or A

50

<220>

<221> MISC\_FEATURE

<222> (12)..(12)

<223> X is E, R or D

55

<220>

<221> MISC\_FEATURE

<222> (14)..(14)

<223> X is F or A

<220>

<221> MISC\_FEATURE

5 <222> (17)..(17)  
<223> X is L, F or A

<220>

<221> MISC\_FEATURE

10 <222> (18)..(18)  
<223> X is V or F

<220>

<221> MISC\_FEATURE

15 <222> (21)..(21)  
<223> X is A, V or L

<220>

<221> MISC\_FEATURE

20 <222> (25)..(25)  
<223> X is L, F, or W

<220>

<221> MISC\_FEATURE

25 <222> (28)..(28)  
<223> X is M, F, or I

<220>

<221> MISC\_FEATURE

30 <222> (32)..(32)  
<223> X is L, F, or I

<220>

<221> MISC\_FEATURE

35 <222> (35)..(35)  
<223> X is E, G, or D

<220>

<221> MISC\_FEATURE

40 <222> (36)..(36)  
<223> X is W, Q, G, or D

<220>

<221> MISC\_FEATURE

45 <222> (37)..(37)  
<223> X is Q, I, P, or F

<220>

<221> MISC\_FEATURE

50 <222> (38)..(38)  
<223> X is Q, K, S, or H

<220>

<221> MISC\_FEATURE

55 <222> (42)..(42)  
<223> X is M, R, or A

<220>

<221> MISC\_FEATURE  
<222> (45)..(45)  
<223> X is W, L, or M

5        <220>  
      <221> MISC\_FEATURE  
      <222> (46)..(46)  
      <223> X is M, K or F

10      <220>  
      <221> MISC\_FEATURE  
      <222> (48)..(48)  
      <223> X is D, L, K or F

15      <220>  
      <221> MISC\_FEATURE <222> (49)..(49)  
      <223> X is A, M, or L

20      <220>  
      <221> MISC\_FEATURE  
      <222> (51)..(51)  
      <223> X is Y or K

25      <220>  
      <221> MISC\_FEATURE  
      <222> (53)..(53)  
      <223> X is V, A or M

30      <220>  
      <221> MISC\_FEATURE  
      <222> (56)..(56)  
      <223> X is L, A or M

35      <220>  
      <221> MISC\_FEATURE  
      <222> (58)..(58)  
      <223> X is K or R

40      <220>  
      <221> MISC\_FEATURE  
      <222> (60)..(60)  
      <223> X is V, M or I

45      <220>  
      <221> MISC\_FEATURE  
      <222> (63)..(63)  
      <223> X is F, R, I or Y

50      <220>  
      <221> MISC\_FEATURE  
      <222> (64)..(64)  
      <223> X is L, E, or A

55      <220>  
      <221> MISC\_FEATURE  
      <222> (67)..(67)  
      <223> X is L or A

<220>  
<221> MISC\_FEATURE <222> (72)..(72)  
<223> X is N or Y

5 <220>  
<221> MISC\_FEATURE  
<222> (74)..(74)  
<223> X is M or L

10 <220>  
<221> MISC\_FEATURE <222> (78)..(78)  
<223> X is N or I

15 <220>  
<221> MISC\_FEATURE  
<222> (80)..(80)  
<223> X is D or Q

20 <220>  
<221> MISC\_FEATURE  
<222> (82)..(82)  
<223> X is K, E, T, M, or Q

25 <220>  
<221> MISC\_FEATURE  
<222> (84)..(84)  
<223> X is I, M, or A

30 <220>  
<221> MISC\_FEATURE  
<222> (88)..(88)  
<223> X is I, L, or V

35 <220>  
<221> MISC\_FEATURE  
<222> (89)..(89)  
<223> X is I, L, or K

40 <220>  
<221> MISC\_FEATURE  
<222> (91)..(91)  
<223> X is F, Y, E or L

45 <220>  
<221> MISC\_FEATURE  
<222> (92)..(92)  
<223> X is D, A, V, or M

50 <220>  
<221> MISC\_FEATURE  
<222> (94)..(94)  
<223> X is E, I or R

55 <220>  
<221> MISC\_FEATURE  
<222> (95)..(95)  
<223> X is V, A, or L

5           <220>  
          <221> MISC\_FEATURE  
          <222> (96)..(96)  
          <223> X is E, F or W

10           <220>  
          <221> MISC\_FEATURE  
          <222> (99)..(99)  
          <223> X is L or I

15           <220>  
          <221> MISC\_FEATURE  
          <222> (102)..(102)  
          <223> X is M, L or I

20           <220>  
          <221> MISC\_FEATURE  
          <222> (103)..(103)  
          <223> X is K, A, M, or L

25           <220>  
          <221> MISC\_FEATURE  
          <222> (106)..(106)  
          <223> X is A, V, or I

30           <220>  
          <221> MISC\_FEATURE  
          <222> (108)..(108)  
          <223> X is R, D, I or A

35           <220>  
          <221> MISC\_FEATURE  
          <222> (109)..(109)  
          <223> X is M, K, L or W

40           <220>  
          <221> MISC\_FEATURE  
          <222> (110)..(110)  
          <223> X is A, F, or V

45           <220>  
          <221> MISC\_FEATURE  
          <222> (113)..(115)  
          <223> optionally absent

46           <400> 3

50

55

Gly Ser Xaa Ser Asp Xaa Arg Lys Asp Xaa Glu Xaa Arg Xaa Asp Lys  
 1 5 10 15

5 Xaa Xaa Glu Ala Xaa Lys Asn Lys Xaa Asp Lys Xaa Lys Ala Ala Xaa  
 20 25 30

10 Arg Lys Xaa Xaa Xaa Glu Glu Arg Xaa Lys Asp Xaa Xaa Lys Xaa  
 35 40 45

15 Xaa Arg Xaa Glu Xaa Glu Gln Xaa Arg Xaa Ala Xaa Arg Asn Xaa Xaa  
 50 55 60

20 Ser Glu Xaa Leu Ser Lys Ile Xaa Asp Xaa Pro Ile Thr Xaa Asp Xaa  
 65 70 75 80

25 Lys Xaa Leu Xaa Ser Asn Asp Xaa Xaa Lys Xaa Xaa Ala Xaa Xaa Xaa  
 85 90 95

30 Lys Lys Xaa Glu Ala Xaa Xaa Ala Asp Xaa Glu Xaa Xaa Xaa Thr Gln  
 100 105 110

25 Gly Ser Trp  
 115

35 <210> 4  
 <211> 115  
 <212> PRT  
 <213> Artificial Sequence

40 <220>  
 <223> Synthetic

<400> 4

45

50

55

Gly Ser Arg Ser Asp Met Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
 1 5 10 15

5 Phe Val Glu Ala Ala Lys Asn Lys Phe Asp Lys Phe Lys Ala Ala Leu  
 20 25 30

10 Arg Lys Gly Asp Ile Lys Glu Glu Arg Arg Lys Asp Met Lys Lys Leu  
 35 40 45

15 Ala Arg Lys Glu Ala Glu Gln Ala Arg Arg Ala Val Arg Asn Arg Leu  
 50 55 60

Ser Glu Leu Leu Ser Lys Ile Asn Asp Met Pro Ile Thr Asn Asp Gln  
 65 70 75 80

20 Lys Lys Leu Met Ser Asn Asp Val Leu Lys Phe Ala Ala Glu Ala Glu  
 85 90 95

25 Lys Lys Ile Glu Ala Leu Ala Ala Asp Ala Glu Asp Lys Phe Thr Gln  
 100 105 110

Gly Ser Trp  
 115

30 <210> 5  
 <211> 115  
 <212> PRT  
 <213> Artificial Sequence

35 <220>  
 <223> Synthetic

40 <400> 5

45

50

55

Gly Ser Leu Ser Asp Val Arg Lys Asp Val Glu Lys Arg Ile Asp Lys  
 1 5 10 15

5 Ala Leu Glu Ala Phe Lys Asn Lys Met Asp Lys Glu Lys Ala Ala Phe  
 20 25 30

10 Arg Lys Asp Pro Pro Ser Glu Glu Arg Arg Lys Asp Lys Lys Lys Glu  
 35 40 45

15 Phe Arg Glu Glu Arg Glu Gln Val Arg Lys Ala Ile Arg Asn Val Leu  
 50 55 60

Ser Glu Ala Leu Ser Lys Ile Asn Asp Leu Pro Ile Thr Asn Asp Lys  
 65 70 75 80

20 Lys Lys Leu Val Ser Asn Asp Val Ile Lys Lys Val Ala Glu Met Lys

85 90 95

25 Lys Lys Val Glu Leu Glu Val Ala Asp Val Glu Lys Lys Val Thr Gln  
 100 105 110

30 Gly Ser Trp  
 115

35 <210> 6  
 <211> 112  
 <212> PRT  
 <213> Artificial Sequence

40 <220>  
 <223> Synthetic

45 <400> 6

50

55

EP 2 834 263 B9

Gly Ser Met Ser Asp Ala Arg Lys Asp Leu Glu Glu Arg Leu Asp Lys  
1 5 10 15

5 Leu Leu Glu Ala Ala Lys Asn Lys Met Asp Lys Phe Lys Ala Ala Met  
20 25 30

10 Arg Lys Arg Gly Gln Arg Glu Glu Arg Lys Lys Asp Trp Ala Lys Ile  
35 40 45

15 Val Arg Asp Glu Phe Glu Gln Phe Arg Lys Ala Val Arg Asn Phe Leu  
50 55 60

Ser Glu Ala Leu Ser Lys Ile Asn Asp Tyr Pro Ile Thr Asn Asp Asp  
65 70 75 80

20 Lys Lys Leu Thr Ser Asn Asp Thr Lys Lys Phe Ala Ala Glu Val Glu  
85 90 95

25 Lys Lys Leu Glu Ala Phe Lys Ala Asp Val Glu Glu Ala Ala Thr Gln  
100 105 110

<210> 7

<211> 112

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Synthetic

35 <400> 7

Gly Ser Met Ser Asp Ile Arg Lys Asp Leu Glu Glu Arg Phe Asp Lys  
1 5 10 15

40 Leu Val Glu Ala Leu Lys Asn Lys Val Asp Lys Met Lys Ala Ala Phe  
20 25 30

45

50

55

Arg Lys Asp Gln Phe His Glu Glu Arg Met Lys Asp Trp Phe Lys Asp  
 35 40 45

5 Leu Arg Lys Glu Val Glu Gln Met Arg Arg Ala Val Arg Asn Tyr Ala  
 50 55 60

10 Ser Glu Ala Leu Ser Lys Ile Asn Asp Leu Pro Ile Thr Asn Asp Asp  
 65 70 75 80

Lys Lys Leu Ala Ser Asn Asp Val Leu Lys Leu Val Ala Glu Val Trp  
 85 90 95

15 Lys Lys Leu Glu Ala Ile Leu Ala Asp Val Glu Ala Trp Phe Thr Gln  
 100 105 110

20 <210> 8  
 <211> 112  
 <212> PRT  
 <213> Artificial Sequence

25 <220>  
 <223> Synthetic

<400> 8

30 Gly Ser Phe Ser Asp Ile Arg Lys Asp Ala Glu Asp Arg Ala Asp Lys  
 1 5 10 15

35 Ala Phe Glu Ala Ala Lys Asn Lys Phe Asp Lys Ile Lys Ala Ala Ile  
 20 25 30

Arg Lys Asp Trp Pro Ser Glu Glu Arg Ala Lys Asp Leu Met Lys Lys  
 35 40 45

40 Ala Arg Tyr Glu Met Glu Gln Ala Arg Arg Ala Ile Arg Asn Ile Glu  
 50 55 60

45 Ser Glu Ala Leu Ser Lys Ile Asn Asp Leu Pro Ile Thr Asn Asp Gln  
 65 70 75 80

50 Lys Lys Leu Ala Ser Asn Asp Ile Ile Lys Glu Met Ala Arg Leu Phe  
 85 90 95

55 Lys Lys Leu Glu Ala Leu Met Ala Asp Ile Glu Ile Leu Val Thr Gln  
 100 105 110

<210> 9  
 <211> 115  
 <212> PRT  
 <213> Artificial Sequence

EP 2 834 263 B9

<220>  
<223> Synthetic

<400> 9

5

Gly Ser Leu Ser Asp Ile Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
1 5 10 15

10 Leu Val Glu Ala Val Lys Asn Lys Leu Asp Lys Met Lys Ala Ala Leu  
20 25 30

15 Arg Lys Glu Gly Gln Gln Glu Glu Arg Met Lys Asp Leu Met Lys Phe  
35 40 45

Met Arg Lys Glu Val Glu Gln Leu Arg Lys Ala Met Arg Asn Phe Leu  
50 55 60

20 Ser Glu Ala Leu Ser Lys Ile Asn Asp Met Pro Ile Thr Asn Asp Asp  
65 70 75 80

25 Lys Lys Leu Ile Ser Asn Asp Leu Lys Lys Tyr Asp Ala Ile Ala Glu  
85 90 95

30 Lys Lys Leu Glu Ala Met Lys Ala Asp Val Glu Arg Met Ala Thr Gln  
100 105 110

Gly Ser Trp  
115

35 <210> 10  
<211> 115  
<212> PRT  
<213> Artificial Sequence

40 <220>  
<223> Synthetic

45 <220>  
<221> MISC\_FEATURE  
<222> (82)..(82)  
<223> X is E, T, M, or Q

<400> 10

50

55

Gly Ser Arg Ser Asp Met Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
 1 5 10 15

5 Phe Val Glu Ala Ala Lys Asn Lys Phe Asp Lys Phe Lys Ala Ala Leu  
 20 25 30

10 Arg Lys Gly Asp Ile Lys Glu Glu Arg Arg Lys Asp Met Lys Lys Leu  
 35 40 45

15 Ala Arg Lys Glu Ala Glu Gln Ala Arg Arg Ala Val Arg Asn Arg Leu  
 50 55 60

20 Ser Glu Leu Leu Ser Lys Ile Asn Asp Met Pro Ile Thr Asn Asp Gln  
 65 70 75 80

25 Lys Xaa Leu Met Ser Asn Asp Val Leu Lys Phe Ala Ala Glu Ala Glu  
 85 90 95

30 Lys Lys Ile Glu Ala Leu Ala Ala Asp Ala Glu Asp Lys Phe Thr Gln  
 100 105 110

Gly Ser Trp  
 115

35 <210> 11  
 <211> 112  
 <212> PRT  
 <213> Artificial Sequence

40 <220>  
 <223> Synthetic

<220>  
 <221> MISC\_FEATURE  
 <222> (82)..(82)  
 <223> X is E, T, M, or Q

45 <400> 11

50

55

Gly Ser Met Ser Asp Ile Arg Lys Asp Leu Glu Glu Arg Phe Asp Lys  
 1 5 10 15

5 Leu Val Glu Ala Leu Lys Asn Lys Val Asp Lys Met Lys Ala Ala Phe  
 20 25 30

10 Arg Lys Asp Gln Phe His Glu Glu Arg Met Lys Asp Trp Phe Lys Asp  
 35 40 45

Leu Arg Lys Glu Val Glu Gln Met Arg Arg Ala Val Arg Asn Tyr Ala  
 50 55 60

15 Ser Glu Ala Leu Ser Lys Ile Asn Asp Leu Pro Ile Thr Asn Asp Asp  
 65 70 75 80

20 Lys Xaa Leu Ala Ser Asn Asp Val Leu Lys Leu Val Ala Glu Val Trp  
 85 90 95

}

25 Lys Lys Leu Glu Ala Ile Leu Ala Asp Val Glu Ala Trp Phe Thr Gln  
 100 105 110

30 <210> 12  
 <211> 115  
 <212> PRT  
 <213> Artificial Sequence

35 <220>  
 <223> Synthetic

40 <220>  
 <221> MISC\_FEATURE  
 <222> (82)..(82)  
 <223> X is E, T, M, or Q

<400> 12

45

50

55

Gly Ser Leu Ser Asp Ile Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
 1 5 10 15

5 Leu Val Glu Ala Val Lys Asn Lys Leu Asp Lys Met Lys Ala Ala Leu  
 20 25 30

10 Arg Lys Glu Gly Gln Gln Glu Glu Arg Met Lys Asp Leu Met Lys Phe  
 35 40 45

15 Met Arg Lys Glu Val Glu Gln Leu Arg Lys Ala Met Arg Asn Phe Leu  
 50 55 60

Ser Glu Ala Leu Ser Lys Ile Asn Asp Met Pro Ile Thr Asn Asp Asp  
 65 70 75 80

20 Lys Xaa Leu Ile Ser Asn Asp Leu Lys Lys Tyr Asp Ala Ile Ala Glu  
 85 90 95

25 Lys Lys Leu Glu Ala Met Lys Ala Asp Val Glu Arg Met Ala Thr Gln  
 100 105 110

Gly Ser Trp  
 115

30 <210> 13  
 <211> 115  
 <212> PRT  
 <213> Artificial Sequence

35 <220>  
 <223> Synthetic  
 <400> 13

40 Gly Ser Arg Ser Asp Met Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
 1 5 10 15

45 Phe Val Glu Ala Ala Lys Asn Lys Phe Asp Lys Phe Lys Ala Ala Leu  
 20 25 30

50

55

Arg Lys Gly Asp Ile Lys Glu Glu Arg Arg Lys Asp Met Lys Lys Leu  
 35 40 45

5 Ala Arg Lys Glu Ala Glu Gln Ala Arg Arg Ala Val Arg Asn Arg Leu  
 50 55 60

10 Ser Glu Leu Leu Ser Lys Ile Tyr Asp Met Pro Ile Thr Asn Asp Gln  
 65 70 75 80

15 Lys Lys Leu Met Ser Asn Asp Val Leu Lys Phe Ala Ala Glu Ala Glu  
 85 90 95

20 Lys Lys Ile Glu Ala Leu Ala Ala Asp Ala Glu Asp Lys Phe Thr Gln  
 100 105 110

25 Gly Ser Trp  
 115

<210> 14

<211> 112

25 <212> PRT

<213> Artificial Sequence

30 <220>

<223> Synthetic

35 <400> 14

Gly Ser Met Ser Asp Ile Arg Lys Asp Leu Glu Glu Arg Phe Asp Lys  
 1 5 10 15

35

Leu Val Glu Ala Leu Lys Asn Lys Val Asp Lys Met Lys Ala Ala Phe  
 20 25 30

40

Arg Lys Asp Gln Phe His Glu Glu Arg Met Lys Asp Trp Phe Lys Asp  
 35 40 45

45

Leu Arg Lys Glu Val Glu Gln Met Arg Arg Ala Val Arg Asn Tyr Ala  
 50 55 60

50

Ser Glu Ala Leu Ser Lys Ile Tyr Asp Leu Pro Ile Thr Asn Asp Asp  
 65 70 75 80

Lys Lys Leu Ala Ser Asn Asp Val Leu Lys Leu Val Ala Glu Val Trp  
 85 90 95

55

Lys Lys Leu Glu Ala Ile Leu Ala Asp Val Glu Ala Trp Phe Thr Gln  
 100 105 110

<210> 15

EP 2 834 263 B9

<211> 115  
<212> PRT  
<213> Artificial Sequence

5 <220>  
<223> Synthetic

<400> 15

10 Gly Ser Leu Ser Asp Ile Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
1 5 10 15

15 Leu Val Glu Ala Val Lys Asn Lys Leu Asp Lys Met Lys Ala Ala Leu  
20 25 30

20 Arg Lys Glu Gly Gln Gln Glu Glu Arg Met Lys Asp Leu Met Lys Phe  
35 40 45

25 Met Arg Lys Glu Val Glu Gln Leu Arg Lys Ala Met Arg Asn Phe Leu  
50 55 60

30 Ser Glu Ala Leu Ser Lys Ile Tyr Asp Met Pro Ile Thr Asn Asp Asp  
65 70 75 80

35 Lys Lys Leu Ile Ser Asn Asp Leu Lys Lys Tyr Asp Ala Ile Ala Glu  
85 90 95

40 Lys Lys Leu Glu Ala Met Lys Ala Asp Val Glu Arg Met Ala Thr Gln  
100 105 110

45 Gly Ser Trp  
115

<210> 16  
<211> 115  
<212> PRT  
<213> Artificial Sequence

45 <220>  
<223> Synthetic

<400> 16

50

55

EP 2 834 263 B9

Gly Ser Arg Ser Asp Met Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
1 5 10 15

5 Phe Val Glu Ala Ala Lys Asn Lys Phe Asp Lys Phe Lys Ala Ala Leu  
20 25 30

10 Arg Lys Gly Asp Ile Lys Glu Glu Arg Arg Lys Asp Met Lys Lys Leu  
35 40 45

15 Ala Arg Lys Glu Ala Glu Gln Ala Arg Arg Ala Val Arg Asn Arg Leu  
50 55 60

20 Ser Glu Leu Leu Ser Lys Ile Asn Asp Met Pro Ile Thr Ile Asp Gln  
65 70 75 80

25 Lys Lys Leu Met Ser Asn Asp Val Leu Lys Phe Ala Ala Glu Ala Glu  
85 90 95

30 Lys Lys Ile Glu Ala Leu Ala Ala Asp Ala Glu Asp Lys Phe Thr Gln  
100 105 110

25 Gly Ser Trp  
115

30 <210> 17  
<211> 112  
<212> PRT  
<213> Artificial Sequence

35 <220>  
<223> Synthetic

<400> 17

40

45

50

55

EP 2 834 263 B9

Gly Ser Met Ser Asp Ile Arg Lys Asp Leu Glu Glu Arg Phe Asp Lys  
1 5 10 15

5 Leu Val Glu Ala Leu Lys Asn Lys Val Asp Lys Met Lys Ala Ala Phe  
20 25 30

10 Arg Lys Asp Gln Phe His Glu Glu Arg Met Lys Asp Trp Phe Lys Asp  
35 40 45

15 Leu Arg Lys Glu Val Glu Gln Met Arg Arg Ala Val Arg Asn Tyr Ala  
50 55 60

20 Ser Glu Ala Leu Ser Lys Ile Asn Asp Leu Pro Ile Thr Ile Asp Asp  
65 70 75 80

25 Lys Lys Leu Ala Ser Asn Asp Val Leu Lys Leu Val Ala Glu Val Trp  
85 90 95

30 Lys Lys Leu Glu Ala Ile Leu Ala Asp Val Glu Ala Trp Phe Thr Gln  
100 105 110

25 <210> 18  
<211> 115  
<212> PRT  
<213> Artificial Sequence

30 <220>  
<223> Synthetic

35 <400> 18

40 Gly Ser Leu Ser Asp Ile Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys

45

50

55



<210> 20  
 <211> 85  
 <212> PRT  
 <213> Artificial Sequence

5

<220>  
 <223> Synthetic

<400> 20

10

Gly Ser Leu Ser Asp Ile Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
 1 5 10 15

15 Leu Val Glu Ala Val Lys Asn Leu Gly Asn Gly Glu Lys Glu Val Glu  
 20 25 30

20 Gln Leu Arg Lys Ala Met Arg Asn Phe Leu Ser Glu Ala Leu Ser Lys  
 35 40 45

25 Ile Asn Asp Met Pro Ile Thr Asn Asp Asp Lys Lys Leu Ile Ser Asn  
 50 55 60

30 Asp Leu Lys Lys Tyr Asp Ala Ile Ala Glu Lys Lys Leu Glu Ala Met  
 65 70 75 80

35 Lys Ala Gly Ser Trp  
 85

<210> 21  
 <211> 90  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Synthetic

40 <220>  
 <221> MISC\_FEATURE  
 <222> (50)..(50)  
 <223> X is N or Y

45 <220>  
 <221> MISC\_FEATURE  
 <222> (56)..(56)  
 <223> X is N or I

50 <220>  
 <221> MISC\_FEATURE  
 <222> (60)..(60)  
 <223> X is K, E, T, M or Q

55 <220>  
 <221> MISC\_FEATURE  
 <222> (83)..(90)  
 <223> optionally absent

&lt;400&gt; 21

5           Gly Ser Met Ser Asp Ile Arg Lys Asp Leu Glu Glu Arg Phe Asp Lys  
       1                   5                   10                   15

10           Leu Val Glu Ala Leu Lys Lys Gly Gln Gly Arg Gln Lys Glu Val Glu  
       20                                   25                           30

15           Gln Met Arg Arg Ala Val Arg Asn Tyr Ala Ser Glu Ala Leu Ser Lys  
       35                           40                           45

20           Ile Xaa Asp Leu Pro Ile Thr Xaa Asp Asp Lys Xaa Leu Ala Ser Asn  
       50                           55                           60

25           Asp Val Leu Lys Leu Val Ala Glu Val Trp Lys Lys Leu Glu Ala Ile  
       65                           70                           75                           80

30           Leu Ala Asp Val Glu Ala Trp Phe Thr Gln  
       85   90

&lt;210&gt; 22

&lt;211&gt; 93

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Synthetic

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (50)..(50)

&lt;223&gt; X is N or Y

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (56)..(56)

&lt;223&gt; X is N or I

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (60)..(60)

&lt;223&gt; X is K, E, T, M, or Q

&lt;220&gt;

&lt;221&gt; MISC\_FEATURE

&lt;222&gt; (83)..(90)

&lt;223&gt; optionally absent

&lt;400&gt; 22

Gly Ser Leu Ser Asp Ile Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
 1 5 10 15

5 Leu Val Glu Ala Val Lys Asn Leu Gly Asn Gly Glu Lys Glu Val Glu  
 20 25 30

10 Gln Leu Arg Lys Ala Met Arg Asn Phe Leu Ser Glu Ala Leu Ser Lys  
 35 40 45

15 Ile Xaa Asp Met Pro Ile Thr Xaa Asp Asp Lys Xaa Leu Ile Ser Asn  
 50 55 60

Asp Leu Lys Lys Tyr Asp Ala Ile Ala Glu Lys Lys Leu Glu Ala Met  
 65 70 75 80

20 Lys Ala Asp Val Glu Arg Met Ala Thr Gln Gly Ser Trp  
 85 90

<210> 23

<211> 65

25 <212> PRT

<213> Artificial Sequence

<220>

30 <223> Synthetic

<400> 23

35 Gly Ser Cys Ser Asp Ile Arg Lys Asp Cys Glu Arg Arg Phe Asp Lys  
 1 5 10 15

Gly Asp Gly Gly Arg Lys Ala Met Arg Asn Phe Leu Ser Glu Cys Leu  
 20 25 30

40 Ser Lys Ile Asn Asp Met Pro Ile Thr Asn Asp Asp Lys Lys Leu Cys  
 35 40 45

45 Ser Asn Asp Leu Lys Lys Tyr Asp Ala Ile Ala Glu Lys Lys Gly Ser  
 50 55 60

50 Trp  
 65

<210> 24

<211> 65

55 <212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic



Gly Ser Leu Ser Asp Cys Arg Lys Asp Ala Glu Arg Arg Phe Asp Lys  
 1 5 10 15

5 Gly Asp Gly Gly Arg Lys Ala Trp Arg Asn Phe Leu Ser Glu Phe Leu  
 20 25 30

10 Ser Cys Ile Asn Asp Met Pro Ile Thr Asn Asp Asp Lys Lys Leu Ile  
 35 40 45

Ser Asn Asp Leu Lys Lys Tyr Leu Ala Ile Ala Glu Lys Lys  
 50 55 60

15 <210> 27  
 <211> 62  
 <212> PRT  
 <213> Artificial Sequence

20 <220>  
 <223> Synthetic

<400> 27

25 Gly Ser Leu Ser Asp Ile Arg Lys Asp Cys Glu Arg Arg Phe Asp Lys  
 1 5 10 15

30 Gly Asp Gly Gly Arg Lys Ala Trp Arg Asn Phe Leu Ser Glu Cys Leu  
 20 25 30

35 Ser Lys Ile Asn Asp Met Pro Ile Thr Asn Asp Asp Lys Lys Leu Ile  
 35 40 45

40 Ser Asn Asp Leu Lys Lys Tyr Leu Ala Ile Ala Glu Lys Lys  
 50 55 60

45 <210> 28  
 <211> 62  
 <212> PRT  
 <213> Artificial Sequence

50 <220>  
 <223> Synthetic

<400> 28

55

Gly Ser Leu Ser Asp Ile Cys Lys Asp Ala Glu Arg Arg Phe Asp Lys  
 1 5 10 15

5 Gly Asp Gly Gly Arg Lys Ala Trp Arg Asn Phe Leu Ser Glu Phe Leu  
 20 25 30

10 Ser Lys Ile Asn Asp Met Pro Ile Thr Asn Asp Asp Lys Lys Leu Ile  
 35 40 45

Ser Asn Asp Cys Lys Lys Tyr Leu Ala Ile Ala Glu Lys Lys  
 50 55 60

15 <210> 29  
 <211> 115  
 <212> PRT  
 <213> Artificial Sequence

20 <220>  
 <223> Synthetic

25 <220>  
 <221> MISC\_FEATURE  
 <222> (1)..(2)  
 <223> optinally absent

30 <220>  
 <221> MISC\_FEATURE  
 <222> (3)..(3)  
 <223> X is M, L, R or F

35 <220>  
 <221> MISC\_FEATURE  
 <222> (6)..(6)  
 <223> X is R, A, I, M or V

40 <220>  
 <221> MISC\_FEATURE  
 <222> (10)..(10)  
 <223> X is L, A or V

45 <220>  
 <221> MISC\_FEATURE  
 <222> (12)..(12)  
 <223> X is E, R, K, or D

50 <220>  
 <221> MISC\_FEATURE  
 <222> (14)..(14)  
 <223> X is L, F, I or A

55 <220>  
 <221> MISC\_FEATURE  
 <222> (17)..(17)  
 <223> X is L, F, or A

<220>

<221> MISC\_FEATURE  
<222> (18).. (18)  
<223> X is L, V, or F

5        <220>  
      <221> MISC\_FEATURE  
      <222> (21)..(21)  
      <223> X is E, V, F, or L

10      <220>  
      <221> MISC\_FEATURE  
      <222> (25)..(25)  
      <223> X is M, L, F, or V

15      <220>  
      <221> MISC\_FEATURE  
      <222> (28)..(28)  
      <223> X is M, E, F, or I

20      <220>  
      <221> MISC\_FEATURE  
      <222> (32)..(32)  
      <223> X is M, L, F, or I

25      <220>  
      <221> MISC\_FEATURE  
      <222> (35)..(35)  
      <223> X is R, E ,G or D

30      <220>  
      <221> MISC\_FEATURE  
      <222> (36)..(36)  
      <223> X is G, D, P, W or Q

35      <220>  
      <221> MISC\_FEATURE  
      <222> (37)..(37)  
      <223> X is Q, I, P or F

40      <220>  
      <221> MISC\_FEATURE  
      <222> (38)..(38)  
      <223> X is R, Q, K, S or H

45      <220>  
      <221> MISC\_FEATURE  
      <222> (42)..(42)  
      <223> X is M, K, R or A

50      <220>  
      <221> MISC\_FEATURE  
      <222> (45)..(45)  
      <223> X is W, L, M or K

55      <220>  
      <221> MISC\_FEATURE  
      <222> (46)..(46)  
      <223> X is A, F, M or K

5           <220>  
          <221> MISC\_FEATURE  
          <222> (48)..(48)  
          <223> X is I, F, L, E, D or K

10           <220>  
          <221> MISC\_FEATURE  
          <222> (49)..(49)  
          <223> X is A, V, M, F, or L

15           <220>  
          <221> MISC\_FEATURE  
          <222> (51)..(51)  
          <223> X is D, K, Y, or E

20           <220>  
          <221> MISC\_FEATURE  
          <222> (53)..(53)  
          <223> X is F, V, A, R or M

25           <220>  
          <221> MISC\_FEATURE  
          <222> (56)..(56)  
          <223> X is F, V, A, L or M

30           <220>  
          <221> MISC\_FEATURE  
          <222> (58)..(58)  
          <223> X is K or R

35           <220>  
          <221> MISC\_FEATURE  
          <222> (60)..(60)  
          <223> X is V, M or I

40           <220>  
          <221> MISC\_FEATURE  
          <222> (63)..(63)  
          <223> X is F, R, V, I or Y

45           <220>  
          <221> MISC\_FEATURE  
          <222> (64)..(64)  
          <223> X is L, E or A

50           <220>  
          <221> MISC\_FEATURE  
          <222> (67)..(67)  
          <223> X is L or A

55           <220>  
          <221> MISC\_FEATURE  
          <222> (72)..(72)  
          <223> X is N or Y

          <220>  
          <221> MISC\_FEATURE  
          <222> (74)..(74)

<223> X is M, L or Y

5       <220>  
      <221> MISC\_FEATURE  
      <222> (78)..(78)  
      <223> X is N or I

10      <220>  
      <221> MISC\_FEATURE  
      <222> (80)..(80)  
      <223> X is D, Q or K

15      <220>  
      <221> MISC\_FEATURE  
      <222> (82)..(82)  
      <223> X is K, E, T, M or Q

20      <220>  
      <221> MISC\_FEATURE  
      <222> (84)..(84)  
      <223> X is I, T, M, V or A

25      <220>  
      <221> MISC\_FEATURE  
      <222> (88)..(88)  
      <223> X is L, T, V or I

30      <220>  
      <221> MISC\_FEATURE  
      <222> (89)..(89)  
      <223> X is L, K or I

35      <220>  
      <221> MISC\_FEATURE  
      <222> (91)..(91)  
      <223> X is F, Y, K, E or L

40      <220>  
      <221> MISC\_FEATURE  
      <222> (92)..(92)  
      <223> X is D, A, V or M

45      <220>  
      <221> MISC\_FEATURE  
      <222> (94)..(94)  
      <223> X is E, I or R

50      <220>  
      <221> MISC\_FEATURE  
      <222> (95)..(95)  
      <223> X is V, A, M or L

55      <220>  
      <221> MISC\_FEATURE  
      <222> (96)..(96)  
      <223> X is E, K, F or W

      <220>

<221> MISC\_FEATURE

<222> (99)..(99)

<223> X is L, I or V

5 <220>

<221> MISC\_FEATURE

<222> (101)..(101)

<223> X is L or A

10 <220>

<221> MISC\_FEATURE

<222> (102)..(102)

<223> X is F, M, L, E or I

15 <220>

<221> MISC\_FEATURE

<222> (103)..(103)

<223> X is L, K, V, M or A

20 <220>

<221> MISC\_FEATURE

<222> (106)..(106)

<223> X is A, V, or I

25 <220>

<221> MISC\_FEATURE

<222> (108)..(108)

<223> X is E, R, D, K, I or A

30 <220>

<221> MISC\_FEATURE

<222> (109)..(109)

<223> X is A, M, K, L or W

35 <220>

<221> MISC\_FEATURE

<222> (110)..(110)

<223> X is A, F or V

40 <220>

<221> MISC\_FEATURE

<222> (113)..(115)

<223> optionally absent

45 <400> 29

Gly Ser Xaa Ser Asp Xaa Arg Lys Asp Xaa Glu Xaa Arg Xaa Asp Lys  
1 5 10 15

50

Xaa Xaa Glu Ala Xaa Lys Asn Lys Xaa Asp Lys Xaa Lys Ala Ala Xaa  
20 25 30

55

Arg Lys Xaa Xaa Xaa Glu Glu Arg Xaa Lys Asp Xaa Xaa Lys Xaa  
35 40 45

Xaa Arg Xaa Glu Xaa Glu Gln Xaa Arg Xaa Ala Xaa Arg Asn Xaa Xaa  
 50 55 60

5 Ser Glu Xaa Leu Ser Lys Ile Xaa Asp Xaa Pro Ile Thr Xaa Asp Xaa  
 65 70 75 80

10 Lys Xaa Leu Xaa Ser Asn Asp Xaa Xaa Lys Xaa Xaa Ala Xaa Xaa Xaa  
 85 90 95

15 Lys Lys Xaa Glu Xaa Xaa Ala Asp Xaa Glu Xaa Xaa Xaa Thr Gln  
 100 105 110

Gly Ser Trp  
 115

20 <210> 30  
 <211> 115  
 <212> PRT  
 <213> Artificial Sequence

25 <220>  
 <223> Synthetic

30 <220>  
 <221> MISC\_FEATURE  
 <222> (1)..(2)  
 <223> optimally absent

35 <220>  
 <221> MISC\_FEATURE  
 <222> (3)..(3)  
 <223> X is M, L, R or F

40 <220>  
 <221> MISC\_FEATURE  
 <222> (6)..(6)  
 <223> X is I or M

45 <220>  
 <221> MISC\_FEATURE  
 <222> (10)..(10)  
 <223> X is L or A

50 <220>  
 <221> MISC\_FEATURE  
 <222> (12)..(12)  
 <223> X is E, R, or D

55 <220>  
 <221> MISC\_FEATURE  
 <222> (14)..(14)  
 <223> X is F or A

<220>

<221> MISC\_FEATURE  
 <222> (17)..(17)  
 <223> X is L, F or A

5 <220>  
 <221> MISC\_FEATURE  
 <222> (18)..(18)  
 <223> X is V or F

10 <220>  
 <221> MISC\_FEATURE  
 <222> (21)..(21)  
 <223> X is A, V or L

15 <220>  
 <221> MISC\_FEATURE  
 <222> (25)..(25)  
 <223> X is F, W or L

20 <220>  
 <221> MISC\_FEATURE  
 <222> (28)..(28)  
 <223> X is F, M or I

25 <220>  
 <221> MISC\_FEATURE  
 <222> (32)..(32)  
 <223> X is F, L or I

30 <220>  
 <221> MISC\_FEATURE  
 <222> (35)..(35)  
 <223> X is E, G or D

35 <220>  
 <221> MISC\_FEATURE  
 <222> (36)..(36)  
 <223> X is E, G, Q, or W

40 <220>  
 <221> MISC\_FEATURE  
 <222> (37)..(37)  
 <223> X is I, P, Q, or F

45 <220>  
 <221> MISC\_FEATURE  
 <222> (38)..(38)  
 <223> X is K, S, Q, or H

50 <220>  
 <221> MISC\_FEATURE  
 <222> (42)..(42)  
 <223> X is M, R, or A

55 <220>  
 <221> MISC\_FEATURE <222> (45)..(45)  
 <223> X is W, L, or M

5           <220>  
 <221> MISC\_FEATURE  
 <222> (46)..(46)  
 <223> X is K, F, or M  
  
 10          <220>  
 <221> MISC\_FEATURE <222> (48)..(48)  
 <223> X is L, F, K or D  
  
 15          <220>  
 <221> MISC\_FEATURE  
 <222> (49)..(49)  
 <223> X is A, M or L  
  
 20          <220>  
 <221> MISC\_FEATURE  
 <222> (51)..(51)  
 <223> X is Y or K  
  
 25          <220>  
 <221> MISC\_FEATURE  
 <222> (56)..(56)  
 <223> X is L, A or M  
  
 30          <220>  
 <221> MISC\_FEATURE  
 <222> (58)..(58)  
 <223> X is K or R  
  
 35          <220>  
 <221> MISC\_FEATURE  
 <222> (60)..(60)  
 <223> X is V, M or I  
  
 40          <220>  
 <221> MISC\_FEATURE  
 <222> (63)..(63)  
 <223> X is F, R, I or Y  
  
 45          <220>  
 <221> MISC\_FEATURE  
 <222> (64)..(64)  
 <223> X is L, E or A  
  
 50          <220>  
 <221> MISC\_FEATURE  
 <222> (67)..(67)  
 <223> X is L or A  
  
 55          <220>  
 <221> MISC\_FEATURE  
 <222> (74)..(74)  
 <223> X is L or M

5           <220>  
 <221> MISC\_FEATURE  
 <222> (80)..(80)  
 <223> X is D or Q  
  
 10          <220>  
 <221> MISC\_FEATURE <222> (84)..(84)  
 <223> X is I, M or A  
  
 15          <220>  
 <221> MISC\_FEATURE  
 <222> (88)..(88)  
 <223> X is I, L or V  
  
 20          <220>  
 <221> MISC\_FEATURE  
 <222> (89)..(89)  
 <223> X is I, L or K  
  
 25          <220>  
 <221> MISC\_FEATURE  
 <222> (91)..(91)  
 <223> X is F, Y, E, or L  
  
 30          <220>  
 <221> MISC\_FEATURE  
 <222> (94)..(94)  
 <223> X is E, I or R  
  
 35          <220>  
 <221> MISC\_FEATURE  
 <222> (95)..(95)  
 <223> X is V, A or L  
  
 40          <220>  
 <221> MISC\_FEATURE  
 <222> (96)..(96)  
 <223> X is E, F or W  
  
 45          <220>  
 <221> MISC\_FEATURE  
 <222> (99)..(99)  
 <223> X is L or I  
  
 50          <220>  
 <221> MISC\_FEATURE  
 <222> (102)..(102)  
 <223> X is M, L or I  
  
 55          <220>  
 <221> MISC\_FEATURE  
 <222> (103)..(103)  
 <223> X is K, A, M or L

<220>  
 <221> MISC\_FEATURE  
 <222> (106)..(106)  
 <223> X is A, V or I

5

<220>  
 <221> MISC\_FEATURE  
 <222> (108)..(108)  
 <223> X is R, D, I or A

10

<220>  
 <221> MISC\_FEATURE  
 <222> (109)..(109)  
 <223> X is M, K, L or W

15

<220>  
 <221> MISC\_FEATURE  
 <222> (110)..(110)  
 <223> X is A, F, or V

20

<220>  
 <221> MISC\_FEATURE  
 <222> (113)..(115)  
 <223> optimally absent

25

<400> 30

Gly Ser Xaa Ser Asp Xaa Arg Lys Asp Xaa Glu Xaa Arg Xaa Asp Lys  
 1 5 10 15

30

Xaa Xaa Glu Ala Xaa Lys Asn Lys Xaa Asp Lys Xaa Lys Ala Ala Xaa  
 20 25 30

35

Arg Lys Xaa Xaa Xaa Xaa Glu Glu Arg Xaa Lys Asp Xaa Xaa Lys Xaa  
 35 40 45

40

Xaa Arg Xaa Glu Xaa Glu Gln Xaa Arg Xaa Ala Xaa Arg Asn Xaa Xaa  
 50 55 60

45

Ser Glu Xaa Leu Ser Lys Ile Asn Asp Xaa Pro Ile Thr Asn Asp Xaa  
 65 70 75 80

Lys Lys Leu Xaa Ser Asn Asp Xaa Xaa Lys Xaa Xaa Ala Xaa Xaa Xaa  
 85 90 95

50

Lys Lys Xaa Glu Ala Xaa Xaa Ala Asp Xaa Glu Xaa Xaa Xaa Thr Gln  
 100 105 110

55

Gly Ser Trp  
 115

<210> 31  
 <211> 116

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

5 &lt;223&gt; Synthetic

&lt;400&gt; 31

10	Gly Ser Ile Ser Asp Ile Arg Lys Asp Ala Glu Val Arg Met Asp Lys	1	5	10	15
15	Ala Val Glu Ala Phe Lys Asn Lys Leu Asp Lys Phe Lys Ala Ala Val	20	25	30	
20	Arg Lys Val Phe Pro Thr Glu Glu Arg Ile Lys Asp Trp Leu Lys Ile	35	40	45	
25	Val Arg Gly Glu Ala Trp Gln Ala Arg Val Ala Val Arg Asn Val Gly	50	55	60	
30	Arg Asp Ala Asn Asp Lys Ala Ala Ala Leu Gly Lys Asp Lys Glu Ile	65	70	75	80
35	Asn Trp Phe Asp Ile Ser Gln Ser Leu Trp Asp Val Gln Lys Leu Thr	85	90	95	
40	Asp Ala Ala Ile Lys Lys Ile Glu Ala Ala Leu Ala Asp Met Glu Ala	100	105	110	
45	Trp Leu Thr Gln	115			

&lt;210&gt; 32

&lt;211&gt; 8

&lt;212&gt; PRT

40 &lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Synthetic

45 &lt;400&gt; 32

Leu Glu His His His His His His	1	5
---------------------------------	---	---

50 &lt;210&gt; 33

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

55 &lt;220&gt;

&lt;223&gt; Synthetic

&lt;220&gt;

<221> MISC\_FEATURE  
 <222> (3)..(3)  
 <223> X is M, L or C

5 <220>  
 <221> MISC\_FEATURE  
 <222> (6)..(6)  
 <223> X is I or C

10 <220>  
 <221> MISC\_FEATURE  
 <222> (7)..(7)  
 <223> X is R or C

15 <220>  
 <221> MISC\_FEATURE  
 <222> (10)..(10)  
 <223> X is L, A, or C

20 <220>  
 <221> MISC\_FEATURE  
 <222> (12)..(12)  
 <223> X is E, R, or absent

25 <220>  
 <221> MISC\_FEATURE  
 <222> (16)..(16)  
 <223> X is L or G

30 <400> 33

Gly	Ser	Xaa	Ser	Asp	Xaa	Xaa	Lys	Asp	Xaa	Glu	Xaa	Phe	Asp	Lys	Xaa
1				5					10				15		

35 <210> 34  
 <211> 76  
 <212> PRT  
 <213> Artificial Sequence

40 <220>  
 <223> Synthetic

45 <220>  
 <221> MISC\_FEATURE  
 <222> (4)..(4)  
 <223> X is L or V

50 <220>  
 <221> MISC\_FEATURE  
 <222> (6)..(6)  
 <223> X is K or N

55 <220>  
 <221> MISC\_FEATURE  
 <222> (7)..(7)  
 <223> X is L or G

5           <220>  
          <221> MISC\_FEATURE  
          <222> (8)..(8)  
          <223> X is Q or G

10           <220>  
          <221> MISC\_FEATURE  
          <222> (9)..(9)  
          <223> X is N or G

15           <220>  
          <221> MISC\_FEATURE  
          <222> (10)..(10)  
          <223> X is R or G

20           <220>  
          <221> MISC\_FEATURE  
          <222> (11)..(11)  
          <223> X is Q or E

25           <220>  
          <221> MISC\_FEATURE  
          <222> (17)..(17)  
          <223> X is L or M

30           <220>  
          <221> MISC\_FEATURE <222> (19)..(19)  
          <223> X is R or K

35           <220>  
          <221> MISC\_FEATURE  
          <222> (21)..(21)  
          <223> X is V or M

40           <220>  
          <221> MISC\_FEATURE  
          <222> (24)..(24)  
          <223> X is Y or F

45           <220>  
          <221> MISC\_FEATURE  
          <222> (25)..(25)  
          <223> X is A or L

50           <220>  
          <221> MISC\_FEATURE  
          <222> (33)..(33)  
          <223> X is N or Y

55           <220>  
          <221> MISC\_FEATURE  
          <222> (35)..(35)  
          <223> X is L or M

55           <220>  
          <221> MISC\_FEATURE  
          <222> (39)..(39)  
          <223> X is N or I

<220>  
<221> MISC\_FEATURE  
<222> (43)..(43)  
<223> X is K, E, T, M, or Q

5

<220>  
<221> MISC\_FEATURE  
<222> (45)..(45)  
<223> X is A or I

10

<220>  
<221> MISC\_FEATURE  
<222> (49)..(49)  
<223> X is V or L

15

<220>  
<221> MISC\_FEATURE  
<222> (50)..(50)  
<223> X is K or L

20

<220>  
<221> MISC\_FEATURE  
<222> (52)..(52)  
<223> X is Y or L

25

<220>  
<221> MISC\_FEATURE  
<222> (53)..(53)  
<223> X is V or D

30

<220>  
<221> MISC\_FEATURE  
<222> (55)..(55)  
<223> X is E or I

35

<220>  
<221> MISC\_FEATURE  
<222> (56)..(56)  
<223> X is V or A

40

<220>  
<221> MISC\_FEATURE  
<222> (57)..(57)  
<223> X is W or E

45

<220>  
<221> MISC\_FEATURE  
<222> (63)..(63)  
<223> X is I or M

50

<220>  
<221> MISC\_FEATURE  
<222> (64)..(64)  
<223> X is L or K

55

<220>  
<221> MISC\_FEATURE <222> (66)..(76)  
<223> optionally absent

<220>  
 <221> MISC\_FEATURE  
 <222> (66)..(66)  
 <223> X is G or D

5

<220>  
 <221> MISC\_FEATURE  
 <222> (67)..(67)  
 <223> X is S or V

10

<220>  
 <221> MISC\_FEATURE  
 <222> (68)..(68)  
 <223> X is W or E

15

<220>  
 <221> MISC\_FEATURE  
 <222> (69)..(69)  
 <223> X is R or A

20

<220>  
 <221> MISC\_FEATURE  
 <222> (70)..(70)  
 <223> X is W or M

25

<220>  
 <221> MISC\_FEATURE  
 <222> (71)..(71)  
 <223> X is F or A

30

<400> 34

Val Glu Ala Xaa Lys Xaa Xaa Xaa Xaa Xaa Xaa Lys Glu Val Glu Gln  
 1 5 10 15

35

Xaa Arg Xaa Ala Xaa Arg Asn Xaa Xaa Ser Glu Ala Leu Ser Lys Ile  
 20 25 30

40

Xaa Asp Xaa Pro Ile Thr Xaa Asp Asp Lys Xaa Leu Xaa Ser Asn Asp

35

40

45

45

Xaa Xaa Lys Xaa Xaa Ala Xaa Xaa Xaa Lys Lys Leu Glu Ala Xaa Xaa  
 50 55 60

50

Ala Xaa Xaa Xaa Xaa Xaa Xaa Thr Gln Gly Ser Trp  
 65 70 75

55

<210> 35  
 <211> 65  
 <212> PRT  
 <213> Artificial Sequence

<220>

<223> Synthetic

<220>  
 5 <221> MISC\_FEATURE  
 <222> (3)..(3)  
 <223> X is C or L

<220>  
 10 <221> MISC\_FEATURE  
 <222> (6)..(6)  
 <223> X is I or C

<220>  
 15 <221> MISC\_FEATURE  
 <222> (7)..(7)  
 <223> X is R or C

<220>  
 20 <221> MISC\_FEATURE  
 <222> (10)..(10)  
 <223> X is A or C

<220>  
 25 <221> MISC\_FEATURE  
 <222> (24)..(24)  
 <223> X is M or W

<220>  
 30 <221> MISC\_FEATURE  
 <222> (31)..(31)  
 <223> X is C or F

<220>  
 35 <221> MISC\_FEATURE  
 <222> (34)..(34)  
 <223> X is C or K

<220>  
 40 <221> MISC\_FEATURE <222> (48)..(48)  
 <223> X is C or I

<220>  
 45 <221> MISC\_FEATURE  
 <222> (52)..(52)  
 <223> X is C or L

<220>  
 50 <221> MISC\_FEATURE  
 <222> (56)..(56)  
 <223> X is D or L

<220>  
 55 <221> MISC\_FEATURE  
 <222> (63)..(65)  
 <223> optionally absent

<400> 35

Gly Ser Xaa Ser Asp Xaa Xaa Lys Asp Xaa Glu Arg Arg Phe Asp Lys  
 1 5 10 15

5 Gly Asp Gly Gly Arg Lys Ala Xaa Arg Asn Phe Leu Ser Glu Xaa Leu  
 20 25 30

10 Ser Xaa Ile Asn Asp Met Pro Ile Thr Asn Asp Asp Lys Lys Leu Xaa  
 35 40 45

Ser Asn Asp Xaa Lys Lys Tyr Xaa Ala Ile Ala Glu Lys Lys Gly Ser  
 50 55 60

15 Trp  
 65

20 <210> 36  
 <211> 16  
 <212> PRT  
 <213> Artificial Sequence

25 <220>  
 <223> Synthetic

30 <220>  
 <221> MISC\_FEATURE  
 <222> (3)..(3)

35 <223> X is M, L or C  
 <220>  
 <221> MISC\_FEATURE

<222> (10)..(10)  
 35 <223> X is L or A

40 <220>  
 <221> MISC\_FEATURE  
 <222> (12)..(12)  
 <223> X is E, R, or absent

<400> 36

45 Gly Ser Xaa Ser Asp Ile Arg Lys Asp Xaa Glu Xaa Phe Asp Lys Leu  
 1 5 10 15

50 <210> 37  
 <211> 17  
 <212> PRT  
 <213> Artificial Sequence

55 <220>  
 <223> Synthetic

<220>  
 <221> MISC\_FEATURE  
 <222> (3)..(3)  
 <223> X is L or C

5           <220>  
           <221> MISC\_FEATURE  
           <222> (6)..(6)  
           <223> X is I or C

10           <220>  
           <221> MISC\_FEATURE  
           <222> (7)..(7)  
           <223> X is R or C

15           <220>  
           <221> MISC\_FEATURE  
           <222> (10)..(10)  
           <223> X is A or C

1           <400> 37

20           Gly Ser Xaa Ser Asp Xaa Xaa Lys Asp Xaa Glu Arg Arg Phe Asp Lys  
           1                   5                   10                   15

25           Gly

## 25       Claims

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, and SEQ ID NO:28.
2. An isolated polypeptide, comprising an amino acid sequence according to SEQ ID NO:1.
3. The isolated polypeptide of claim 2, comprising an amino acid sequence according to SEQ ID NO:29 or SEQ ID NO:3.
- 35       4. The isolated polypeptide of claim 2 or 3, wherein the polypeptide comprises a sequence selected from the group consisting of SEQ ID NO: 10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, and SEQ ID NO:18.
- 40       5. A virus-like particle (VLP) comprising the polypeptide of any one of claims 1-4.
6. An isolated nucleic acid encoding the polypeptide of any one of claims 1-4.
7. A recombinant expression vector comprising the isolated nucleic acid of claim 6 operatively linked to a promoter.
- 45       8. A recombinant host cell comprising the recombinant expression vector of claim 7.
9. A pharmaceutical composition, comprising the polypeptide of any one of claims 1-4, or the VLP of claim 5, and a pharmaceutically acceptable carrier.

## 50       Patentansprüche

1. Ein isoliertes Polypeptid umfassend eine Aminosäuresequenz ausgewählt aus der Gruppe bestehend aus SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27 und SEQ ID NO:28.
2. Ein isoliertes Polypeptid umfassend eine Aminosäuresequenz gemäß SEQ ID NO:1.

3. Das isolierte Polypeptid des Anspruchs 2, umfassend eine Aminosäuresequenz gemäß SEQ ID NO:29 oder SEQ ID NO:3.
5. Das isolierte Polypeptid des Anspruchs 2 oder 3, wobei das Polypeptid eine Sequenz ausgewählt aus der Gruppe bestehend aus SEQ ID NO: 10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17 und SEQ ID NO:18 umfasst.
10. Ein virusähnlicher Partikel (VLP) umfassend das Polypeptid einem der Ansprüche 1-4.
15. Eine isolierte Nukleinsäure kodierend für das Polypeptid einem der Ansprüche 1-4.
20. Ein rekombinanter Expressionsvektor umfassend die isolierte Nukleinsäure des Anspruchs 6 operativ verbunden mit einem Promotor.
25. Eine rekombinante Wirtszelle umfassend den Expressionsvektor des Anspruchs 7.
30. Eine pharmazeutische Zusammensetzung, umfassend das Polypeptid einem der Ansprüche 1-4 oder den VLP des Anspruchs 5, und einen pharmazeutisch annehmbaren Träger.

20 **Revendications**

1. Polypeptide isolé comprenant une séquence d'acides aminés choisie dans l'ensemble constitué par la SEQ ID NO: 19, la SEQ ID NO: 20, la SEQ ID NO: 21, la SEQ ID NO: 22, la SEQ ID NO: 23, la SEQ ID NO:24, la SEQ ID NO: 25, la SEQ ID NO: 26, la SEQ ID NO: 27 et la SEQ ID NO: 28.
2. Polypeptide isolé, comprenant une séquence d'acides aminés selon la SEQ ID NO: 1.
3. Polypeptide isolé selon la revendication 2, comprenant une séquence d'acides aminés selon la SEQ ID NO: 29 ou la SEQ ID NO: 3.
4. Polypeptide isolé selon la revendication 2 ou 3, où le polypeptide isolé comprend une séquence choisie dans l'ensemble constitué par la SEQ ID NO: 10, la SEQ ID NO: 11, la SEQ ID NO: 12, la SEQ ID NO: 13, la SEQ ID NO: 14, la SEQ ID NO: 15, la SEQ ID NO: 16, la SEQ ID NO: 17 et la SEQ ID NO: 18.
35. Particule pseudo-virale (PPV) comprenant le polypeptide selon l'une quelconque des revendications 1 à 4.
6. Acide nucléique isolé codant le polypeptide selon l'une quelconque des revendications 1 à 4.
40. Vecteur d'expression recombinant comprenant l'acide nucléique isolé de la revendication 6 lié de façon opérationnelle à un promoteur.
8. Cellule hôte recombinante comprenant le vecteur d'expression recombinant de la revendication 7.
45. 9. Composition pharmaceutique, comprenant le polypeptide selon l'une quelconque des revendications 1 à 4, ou la PPV selon la revendication 5, et un véhicule pharmaceutiquement acceptable.

50

55




Fig. 1




Figure 2

GSMSDRRKDLEERLDKLLLEAKNKEDDKFKAAMRKGGORETERMKDWAQKIADEFEQFRKAV 60 FFL\_003  
 GSMSDARKDLEERLDKLLLEAKNKMDKPKAAMRKGGOREERKKDWAKIVRDEFQFRKAV 60 FFL\_004  
 GSLSDIRKDAERRFDFKLVEAVKNKLQKMKAAALKKEGQQEERMKDLMKFMRKEVEOLRKAM 60 FFL\_007  
 GSRSDMRKDAERRFDFKVEAAKNKPPDKFKAALRKDIKEERRKDMKLLARKEAEQARRAV 60 FFL\_001  
 GSLSDVRKDVEKRIDKALEAFKNKMDKEKLAFFRKDPFSEERRKDKKKFREEREQVRKAI 60 FFL\_002  
 GSPSDIRKDAEDRADKAFFEAALKNKFDFKIAAALKDWPSEERAALKDMMKKARYMEQARRAI 60 FFL\_006  
 GSLSDLMKDLEKRFDFKMEAALKNWKDVKAALKDQEKGERAKDMFKIFREELEQLRKAI 60 FFL\_008  
 GSMSDIRKDLERFDKLVEALKNWKDVKAALKDQFHEERMKDWFQDLRKEVEQMRRAV 60 FFL\_005  
 GSISDIRKDAEVURMDKAVEAFKNKLQDKFKLAVRKVFPTERIKDWLKVIRGEAEQARVAV 60 T93  
 \* ;  
  
 RNFLSEALSKIN----DYPITNDDKKLTSNDAKKFDAEVAKKLEAFFKADAEEAATO---- 112 FFL\_003  
 RNFLSEALSKIN----DYPITNDDKKLTSNDDEKKFAAEVKKLEAFFKADAEEAATO---- 112 FFL\_004  
 RNFLSEALSKIN----DMPITNDDKKLISNDLKKYDATAEKKLEAMKADVERMATQGSW 115 FFL\_007  
 RNRLSELLSKIN----DMPITNDQKKLMSNDVLIKFKRAAEAKKIEALAAADAEDKFTQGSW 115 FFL\_001  
 RAVLSEALSKIN----DLPIITNDAKKKLUSNDVKKVAEMKKVKELEVADVERKVTQGSW 115 FFL\_002  
 RNIESEALSKIN----DLPIITNDQKKLKLASNDI1KEMARLFFKKLEALMADIEILVTO---- 112 FFL\_006  
 RNALSEALSKIN----DLPIITNDDKKLKLASNKAKKRAARVKKVEAFIADEVAKKTQ---- 112 FFL\_008  
 RNYASEALSKIN----DLPIITNDDKKLASNDVLIKVAEVWKLEA1LADVEAWFTQ---- 112 FFL\_005  
 RNVGGRDANDKAAALGRDKEINWFDISQSLWDYQKLEDAAIKKIEAALADMEAWLTO---- 116 T93  
 \* ;

Figure 3




Fig 4




Figure 5




Fig 6




Figure 7




Figure 8

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- US 20110236408 A [0041]
- US 7229624 B [0041] [0042]
- US 5770380 A [0042]
- US 5334394 A [0042]
- WO 61620804 A [0098]

## Non-patent literature cited in the description

- **MCLELLAN et al.** *J Mol Biol.*, 24 June 2011, vol. 409 (5), 853-866 [0001]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1989 [0007] [0045]
- Methods in Enzymology. Gene Expression Technology. Academic Press, 1991, vol. 185 [0007]
- Methods in Enzymology. Guide to Protein Purification. Academic Press, Inc, 1990 [0007]
- PCR Protocols: A Guide to Methods and Applications. Academic Press, 1990 [0007]
- **LISS, INC.** Culture of Animal Cells: A Manual of Basic Technique. 1987 [0007]
- Gene Transfer and Expression Protocols. The Human Press Inc, 109-128 [0007] [0044]
- **ZHU et al.** *JID*, 2011 [0019]
- **ARBIZA et al.** *J. Gen Virol*, 1992 [0019]
- **CORREIA et al.** *J. Mol Biol.*, 2011 [0031]
- **SAMBROOK ; FRITSCH ; MANIATIS.** Molecular Cloning, A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1989 [0044]
- **R.I. FRESHNEY.** Culture of Animal Cells: A Manual of Basic Technique. Liss, Inc, 1987 [0045]
- **WU et al.** *J. Biol. Chem.*, 1992, vol. 267, 963-967 [0056]
- **Science** [0059]
- **ROHL et al.** *Methods in Enzymology*, 2004, vol. 383 [0064]
- **MCLELLAN et al.** *J. Mol. Biol.*, 2011, vol. 409, 853-866 [0090]