



(12) **CORRECTED EUROPEAN PATENT APPLICATION**

(15) Correction information:
Corrected version no 1 (W1 A1)
Corrections, see
Description Paragraph(s) 56

(51) Int Cl.:
C12N 15/82 (2006.01)

(48) Corrigendum issued on:
05.10.2016 Bulletin 2016/40

(43) Date of publication:
25.02.2015 Bulletin 2015/09

(21) Application number: **13181236.4**

(22) Date of filing: **21.08.2013**

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR

(60) Divisional application:
16160385.7

(71) Applicant: **Industry-Academic Cooperation**
Foundation,
Yonsei University
Seoul 120-749 (KR)


(72) Inventors:
• **Kim, Woo Taek**
411-370 Gyeonggi-do (KR)
• **Kim, Eun Yu**
110-035 Seoul (KR)
• **Seo, Ji Ho**
110-035 Seoul (KR)
• **Park, Ki Youl**
133-070 Seoul (KR)

(74) Representative: **Plougmann Vingtoft a/s**
Rued Langgaards Vej 8
2300 Copenhagen S (DK)

(54) **Gene implicated in abiotic stress tolerance and growth accelerating and use thereof**

(57) The present invention provides a composition for improving the tolerance of a plant to an abiotic stress and a composition for promoting growing of a plant, comprising a nucleotide sequence encoding the AtSRP (*Arabidopsis thaliana* stress related protein) of a plant. The present nucleotide sequence is involved in abiotic stress tolerance such as drought, low-temperature and salt stresses of plants. Therefore, the overexpressing transgenic plants have excellent tolerances to these abiotic stresses, whereby they may be useful as novel functional crops which are affected by climates and environments of the cultivated areas. In addition, where the plants are transformed with the present nucleotide sequence, the growth abilities of the transgenic plants are remarkably enhanced, whereby they may effectively used for cultivating the plants with novel function of rapid growing, and biomass.

Fig. 7b



Description**BACKGROUND OF THE INVENTION****FIELD OF THE INVENTION**

[0001] The present invention relates to a gene implicated in abiotic stress tolerance and growth promotion and a composition for improving abiotic stress tolerance and promoting growing of transformed plants with the same.

DESCRIPTION OF THE RELATED ART

[0002] Due to their sessile nature, higher plants are constantly faced with various adverse environmental factors, including drought, high salt, heavy metals, cold, heat shock, and ozone, during their whole life span. These abiotic stresses are a limiting factor for the growth and development of crop plants. Water deficiency causes dramatic reduction of crop production globally, and the decreasing availability of fresh water may pose a future threat to humans and higher plants. Plants have cellular and genetic defense mechanisms to enhance their tolerance to transient and long-term water shortages by triggering signaling network pathways and inducing stress-responsive genes (Shinozaki and Yamaguchi-Shinozaki, 2007). However, for stress tolerance or sensitivity, our knowledge concerning the biological functions of stress-related genes in higher plants is still rudimentary. Therefore, it is important to study the functions of stress responsive genes to increase the productivity of crop plants.

[0003] Plants have diverse defense strategies to reinforce their tolerance against unfavorable conditions by triggering signaling network pathways and inducing the stress-responsive genes. The present inventors have found that three *AtSRP* (*Arabidopsis thaliana* stress related protein) genes which encode homologs to the small rubber particle protein (SRPP) in rubber trees (*Hevea brasiliensis*) were rapidly induced by dehydration in *Arabidopsis thaliana*. Natural rubber is cis-1,4-polyisoprene produced through mevalonate pathway in cytosolic fractions (latex) of latex vessel tissues (Oh et al. 1999; Sookmark et al. 2002 ; Kim et al. 2004; Chow et al. 2007). Since *Arabidopsis thaliana* does not produce natural rubber, it was unexpected that the *Arabidopsis thaliana* water stress-induced genes shared sequence homology with rubber biosynthetic genes.

[0004] SRPP, originally known as a latex allergin, is a protein tightly bound on a small rubber particle in the latex of rubber trees (*Hevea brasiliensis*).

[0005] Bark tissue of rubber trees is constantly stripped, which is known as a tapping process, to collect rubber latex. Therefore, plugging of latex vessels is essential for rubber trees to prevent the loss of their cytoplasmic components, such as primary metabolites, and to prevent pathogen infection of the latex vessel tissues (Wititsuwannakul et al., 2008b). In the process of latex vessel plugging, hevein or Hevea latex lectin (HLL) interacts with rubber particle (RP) protein to form a rubber latex coagulum (Gidrol et al., 1994; Wititsuwannakul et al., 2008a). Recently, SRPP was purified as an RP glycoprotein that bound HLL and was, therefore, termed HLL-binding protein (HLLBP) (Wititsuwannakul et al., 2008c). Interaction between an N-acetylglucosamine moiety in SRPP and HLL may modulate the degree of latex coagulation in response to tapping and mechanical wounding.

[0006] However, roles of the SRPP associated with abiotic stresses such as drought, high salt and cold have been not known yet.

[0007] Throughout this application, various publications and patents are referred and citations are provided in parentheses. The disclosures of these publications and patents in their entities are hereby incorporated by references into this application in order to fully describe this invention and the state of the art to which this invention pertains.

SUMMARY OF THE INVENTION

[0008] The present inventors have made intensive studies to increase the productivity of crop plants by developing genes for improving a tolerance to abiotic stresses including drought, high salt and cold of the plants. As results, they have discovered that the improved tolerance to abiotic stresses may be obtained when expressions of three *AtSRP* (*Arabidopsis thaliana* stress related protein) genes which encode homologs to the small rubber particle protein (SRPP) in rubber trees (*Hevea brasiliensis*) were increased in plants.

[0009] Accordingly, it is an object of this invention to provide a composition for improving the tolerance of a plant to an abiotic stress.

[0010] It is another object of this invention to provide a composition for promoting growth of a plant.

[0011] It is still another object of this invention to provide a method for improving the tolerance of a plant to an abiotic stress.

[0012] It is further object of this invention to provide a method for promoting growth of a plant.

[0013] Other objects and advantages of the present invention will become apparent from the following detailed de-

scription together with the appended claims and drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014]

Fig. 1 represents that *AtSRPs* are expressed in various organs of Arabidopsis. Fig. 1a represents results of analyzing *AtSRPs* expression by RT-PCR at 7 developmental stages. 18s rRNA was used as a loading control.

WS; whole seedling, RT; root, RL; rosette leaf, ST; stem, F; Flower, SQ; silique, SD; seed.

Fig. 1b represents results of analyzing *AtSRPs* expression by real-time RT-PCR in various organs of Arabidopsis. Results are means $\pm$ S.D. from four repeated experiments.

Fig. 2 represents that *AtSRPs* are up-regulated by ABA and abiotic stresses. Total RNA was prepared from light-grown, 7-day-old Arabidopsis seedlings, which had been treated with 100 mM ABA (1.5-3 hrs), drought (1.5-3 hrs), cold (4°C for 4.5-24 hrs), or high salinity (300 mM NaCl for 3 hrs). Induction patterns of *AtREFPs* were investigated by RT-PCR (Fig. 2a) or real-time qRT-PCR (Fig. 2b). The *RAB18* and *RD29A* genes were used as positive controls for ABA and abiotic stress, respectively. *AtACT8* transcript levels were used as loading controls. Data represent means $\pm$ SD from four independent experiments.

Fig. 3 represents that *AtSRP1*, membrane associated protein, possessed lectin binding property. Fig. 3a represents results of the biochemical analysis using affinity chromatography with concanavalin A. Fig. 3b represents results of the membrane association test. *AtSRPs* were transiently over-expressed using transient assay. Then, total membrane pellets from Arabidopsis seedlings were resuspended in 1% (v/v) Triton X-100, 2 M urea, 1 M NaCl, or 0.1 M Na₂CO₃. Suspensions were recentrifuged at 125,000g to separate solubilized fractions (S) and insoluble fractions (P), which were analyzed by immunoblotting using Flag antibody.

Fig. 4 represents phenotypic characterizations of T3 *35S:AtSRPs* transgenic Arabidopsis plants and knock-out mutants. *AtSRPs*-overexpressors grew more rapidly than wild type plants. In contrast, knock-out mutants showed opposite phenotypes. Fig. 4a represents morphological comparisons of light-grown 3-day-old wild type and *35S:AtSRPs* transgenic seedlings and knock-out mutants. Fig. 4b represents differences in growth patterns of early roots in wild type, *AtSRPs*-overexpressing, and knock-out seedlings. Error bars indicate mean $\pm$ SD (n = 70). Fig. 4c represents determination of the dimensional parameters of second leaves 2 weeks after germination. The second leaves were detached from wild type, transgenic, and knock-out mutant lines, scanned for image analysis using SCIONIMAGE program, and their blade widths and lengths, petiole lengths, and blade areas were calculated. Error bars indicate mean $\pm$ SD (n = 35). Fig. 4d represents growth morphology of wild type, *35S:AtSRPs* transgenic Arabidopsis, and knock-out mutant plants grown under normal growth conditions 28 days after germination (left panel). Right panel represents differences in growth patterns of inflorescence stems of wild type and transgenic Arabidopsis plants 20 to 28 days after germination. Error bars indicate mean $\pm$ SD (n = 80).

Fig. 5 represents faster-growing phenotypes of *35S:AtSRPs* transgenic plants correlated with enhanced cell cycle progression. Fig. 5a represents longitudinal views of root tips from 3-day-old wild type and *AtSRPs*-overexpressing transgenic and knock-out mutant seedlings. Root sections were stained with propidium iodide and analyzed by confocal microscopy. Fig. 5b represents results of confocal microscope demonstrating the epidermal and palisade mesophyll cells from the second leaves of wild type, *AtSRPs*-overexpressors, and knock-out mutants. Scale bars = 20 μ m.

Fig. 6 represents that loss of function of *AtSRPs* showed susceptible phenotype against salt stress. It shows morphologies of 12-day-old wild-type, *AtSRPs*-overexpressing transgenic, and knock-out mutant seedlings in 100 mM NaCl media.

Fig. 7 represents that *AtSRPs* play a role as a positive factor for drought defensive responses. Fig. 7a represents survival rates of wild type and *35S:AtSRPs* transgenic plants after drought stress. Light-grown 25-day-old wild type, 3-week-old transgenic plants, and 26-day-old knock-out mutants were further grown for 8 days without watering. Plants were then re-watered and their survival rates were determined after 3 days. The survival rate was considered as the ability of plants to continue to grow after returned to normal water conditions following eight-day drought stress. The survival rates are following: wild type, 26.97% (24 out of 89 plants); *35S:AtSRP1* transgenic line, 64.84% (59 out of 91 plants); *35S:AtSRP2* transgenic line, 66.29% (59 out of 89 plants); *35S:AtSRP3* transgenic line, 58.02% (47 out of 81 plants); *atsrp1* mutant line, 3.90% (3 out of 77 plants) and *atsrp3* mutant line, 0.00% (0 out of 101 plants). Fig. 7b represents results of measurement of cut rosette water loss (CRWL) rates. The rosette leaves were detached from light-grown 3-week-old wild type, *35S:AtSRPs* transgenic plants, and knock-out mutants, placed on open-lid Petri dishes, and incubated for 0-5 hrs at room temperature. Decreases in fresh weights were determined. Water loss is expressed as the percentage of initial fresh weight of the detached leaves. Error bars indicate mean $\pm$ SD (n=20). Fig. 7c represents amount (mg/g DW) of chlorophyll a + chlorophyll b in leaves of wild, *35S:AtSRPs*-overexpressing plants, and knock-out mutants. Error bars indicate mean $\pm$ SD (n=3).

Fig. 8 represents analysis results of proline content of leaf discs before and after drought treatments. Data were obtained from three independent experiments. Bars represent means $\pm$ SD ($n = 3$). 35S:AtSRPs transgenic leaves retained their leaf water more effectively than did wild-type leaves under drought conditions. In contrast, knockout mutants showed opposite phenotypes.

Fig. 8 schematically represents AtSRP1, AtSRP2 and AtSRP3 cDNA clones.

DETAILED DESCRIPTION OF THE INVENTION

[0015] In one aspect of this invention, there is provided a composition comprising a nucleotide sequence encoding the amino acid sequence selected from the group consisting of SEQ ID NOs:4 to 6 for use in the improvement of the tolerance of a plant to an abiotic stress.

[0016] The present inventors have made intensive studies to increase the productivity of crop plants by developing genes for improving a tolerance to abiotic stresses including drought, high salt and cold of the plants. As results, they have discovered that the improved tolerance to abiotic stresses may be obtained when expressions of three AtSRP (Arabidopsis thaliana stress related protein) genes which encode homologs to the small rubber particle protein (SRPP) in rubber trees (*Hevea brasiliensis*) were increased in plants.

[0017] According to a preferred embodiment, the present nucleotide sequence encoding the amino acid sequence of SEQ ID NOs:4, 5 and 6 comprises the nucleotide sequence as set forth in SEQ ID NO: 1, 2 and 3, respectively.

[0018] According to a preferred embodiment, the amino acid sequence of SEQ ID NOs:4, 5 and 6 is the sequence of SRPP-like proteins present in *Arabidopsis*. These proteins are named as AtSRP (Arabidopsis thaliana stress related protein) 1, AtSRP 2 and AtSRP 3, respectively. As shown in Examples, the present inventors have found that expressions of these proteins or their coding genes were increased by various abiotic stresses, and the improved tolerance to abiotic stresses may be obtained when these genes were over-expressed in plants.

[0019] It would be obvious to the skilled artisan that the nucleotide sequences used in this invention are not limited to those listed in the appended Sequence Listings. The nucleotide sequences described herein are illustrative and their biological equivalents may be also used in this invention for enhancement of tolerance to a drought stress and promotion of flowering or growing in plants. In this regard, the sequence variations should be construed to be covered by the present invention.

[0020] For nucleotides, the variations may be purely genetic, i.e., ones that do not result in changes in the protein product. This includes nucleic acids that contain functionally equivalent codons, or codons that encode the same amino acid, such as six codons for arginine or serine, or codons that encode biologically equivalent amino acids.

[0021] Considering biologically equivalent variations described hereinabove, the nucleic acid molecule of this invention may encompass sequences having substantial identity to them. Sequences having the substantial identity show at least 80%, more preferably at least 90%, most preferably at least 95% similarity to the nucleic acid molecule of this invention, as measured using one of the sequence comparison algorithms. Methods of alignment of sequences for comparison are well-known in the art. Various programs and alignment algorithms are described in: Smith and Waterman, Adv. Appl. Math. 2:482(1981); Needleman and Wunsch, J. Mol. Bio. 48:443(1970); Pearson and Lipman, Methods in Mol. Biol. 24:307-31(1988); Higgins and Sharp, Gene 73:237-44(1988); Higgins and Sharp, CABIOS 5:151-3(1989) Corpet et al., Nuc. Acids Res. 16:10881-90(1988) Huang et al., Comp. Appl. BioSci. 8:155-65(1992) and Pearson et al., Meth. Mol. Biol. 24:307-31(1994). The NCBI Basic Local Alignment Search Tool (BLAST) [Altschul et al., J. Mol. Biol. 215:403-10(1990)] is available from several sources, including the National Center for Biological Information (NCBI, Bethesda, Md.) and on the Internet, for use in connection with the sequence analysis programs blastp, blastm, blastx, tblastn and tblastx. It can be accessed at <http://www.ncbi.nlm.nih.gov/BLAST/>. A description of how to determine sequence identity using this program is available at <http://www.ncbi.nlm.nih.gov/BLAST/blasChelp.html>.

[0022] According to a preferred embodiment, the abiotic stress is selected from the group consisting of a drought stress, a low-temperature stress and a salt stress.

[0023] To introduce a foreign nucleotide sequence into plant cells or plants may be performed by the methods (Methods of Enzymology, Vol. 153, 1987) known to those skilled in the art. The plant may be transformed using the foreign nucleotide inserted into a carrier (e.g., vectors such as plasmid or virus) or *Agrobacterium tumefaciens* as a mediator (Chilton et al., Cell, 11: 263-271 (1977)) and by directly inserting the foreign nucleotide into plant cells (Lorz et al., Mol. Genet., 199: 178-182 (1985); the disclosure is herein incorporated by reference). For example, electroporation, micro-particle bombardment, polyethylene glycol-mediated uptake may be used in the vector containing no T-DNA region.

[0024] Generally, *Agrobacterium-mediated* transformation is the most preferable (U.S. Pat. Nos. 5,004,863, 5,349,124 and 5,416,011), and the skilled artisan can incubate or culture the transformed cells or seeds to mature plants in appropriate conditions.

[0025] The term "plant(s)" as used herein, is understood by a meaning including a plant cell, a plant tissue and a plant seed as well as a mature plant.

[0026] The plants applicable of the present invention include, but not limited to, food crops such as rice plant, wheat,

barley, corn, bean, potato, Indian bean, oat and Indian millet; vegetable crops such as *Arabidopsis* sp., Chinese cabbage, radish, red pepper, strawberry, tomato, watermelon, cucumber, cabbage, melon, pumpkin, welsh onion, onion and carrot; crops for special use such as ginseng, tobacco plant, cotton plant, sesame, sugar cane, sugar beet, *Perilla* sp., peanut and rape; fruit trees such as apple tree, pear tree, jujube tree, peach tree, kiwi fruit tree, grape tree, citrus fruit tree, persimmon tree, plum tree, apricot tree and banana tree; flowering crops such as rose, gladiolus, gerbera, carnation, chrysanthemum, lily and tulip; and fodder crops such as ryegrass, red clover, orchardgrass, alfalfa, tallfescue and perennial ryegrass.

[0027] According to a preferred embodiment, the nucleotide sequence is contained a recombinant plant expression vector; and the recombinant plant expression vector comprises (i) the nucleotide sequence; (ii) a promoter which is operatively linked to the nucleotide sequence of (i) and generates RNA molecules in plant cells; and (iii) a poly A signal sequence inducing polyadenylation at the 3'-end of the RNA molecules.

[0028] The term "operatively linked" as used herein refers to functional linkage between a nucleic acid expression control sequence (such as a promoter, signal sequence, or array of transcription factor binding sites) and a second nucleotide sequence, wherein the expression control sequence affects transcription and/or translation of the nucleic acid corresponding to the second sequence.

[0029] The vector system of this invention may be constructed in accordance with conventional techniques described in Sambrook et al., *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory Press(2001), teachings of which are incorporated herein by reference.

[0030] The suitable promoter in the present invention includes any one commonly used in the art, for example SP6 promoter, T7 promoter, T3 promoter, PM promoter, maize-ubiquitin promoter, Cauliflower mosaic virus (CaMV)-35S promoter, Nopal synthase (nos) promoter, Figwort mosaic virus 35S promoter, Sugarcane bacilliform virus promoter, commelina yellow mottle virus promoter, photo-inducible promoter of small subunit of Ribulose-1,5-bis-phosphate carboxylase (ssRUBISCO), cytosolic triosphosphate isomerase (TPI) promoter in rice, adenine phosphoribosyltransferase (APRT) or octopine synthase promoter in *Arabidopsis*. Preferably, the promoter used in this invention is CaMV 35S.

[0031] According to a preferred embodiment, the 3'-non-translated region causing polyadenylation includes that from the nopaline synthase gene of *Agrobacterium tumefaciens* (NOS 3' end) (Bevan et al., *Nucleic Acids Research*, 11(2):369-385(1983)), that from the octopine synthase gene of *Agrobacterium tumefaciens*, the 3'-end of the protease inhibitor I or II genes from potato or tomato, the CaMV 35S terminator, and OCS (octopine synthase) terminator. Most preferably, the 3'-non-translated region causing polyadenylation in this invention is NOS.

[0032] Optionally, the present vector for plants may further carry a reporter molecule (e.g., genes for luciferase and β -glucuronidase). In addition, the vector may contain antibiotic resistant genes as selective markers (e.g., neomycin phosphotransferase gene (nptII) and hygromycin phosphotransferase gene (hpt)).

[0033] According to a preferred embodiment, the plant expression vector of this invention is *Agrobacterium* binary vectors.

[0034] The term "binary vector" as used herein, refers to a cloning vector containing two separate vector systems harboring one plasmid responsible for migration consisting of left border (LB) and right border (RB), and another plasmid for target gene-transferring. Any *Agrobacterium* suitable for expressing the nucleotide of this invention may be used, and most preferably, the transformation is carried out using *Agrobacterium tumefaciens* GV3101.

[0035] Introduction of the recombinant vector of this invention into *Agrobacterium* can be carried out by a large number of methods known to one skilled in the art. For example, particle bombardment, electroporation, transfection, lithium acetate method and heat shock method may be used. Preferably, the electroporation is used.

[0036] According to the present invention, the present *AtSRPs*-overexpressing transgenic plants grew more rapidly with longer roots, larger leaves, and bolted earlier with prolonged inflorescences. The growth rates of the transgenic plants are enhanced using properties of the *AtSRPs* genes, whereby they may effectively applied for cultivating the plants with novel function of rapid growing, and biomass.

[0037] In another aspect of this invention, there is provided a method for improving the tolerance of a plant to an abiotic stress, comprising:

- (a) introducing a nucleotide sequence encoding the amino acid sequence selected from the group consisting of SEQ ID NOs:4 to 6 to a cell of the plant; and
- (b) obtaining a transgenic plant exhibiting improved tolerance to an abiotic stress from the cell of the plant.

[0038] In still another aspect of this invention, there is provided a composition comprising a nucleotide sequence encoding the amino acid sequence selected from the group consisting of SEQ ID NOs:4 to 6 for use in the improvement of the growth of a plant.

[0039] In further aspect of this invention, there is provided a method for promoting the growth of a plant, comprising:

- (a) introducing a nucleotide sequence encoding the amino acid sequence selected from the group consisting of

SEQ ID NOs:4 to 6 to a cell of the plant; and

(b) obtaining a transgenic plant exhibiting improved tolerance to an abiotic stress from the cell of the plant.

[0040] As the present amino acid sequence, the present nucleotide sequence and the plant expression vector comprising thereof are mentioned above, they are omitted herein to avoid excessive overlaps.

[0041] The features and advantages of the present invention will be summarized as follows:

(a) The present invention provides a method for improving the tolerance of a plant to an abiotic stress and a method for promoting growing of a plant.

(b) The present nucleotide sequence is involved in abiotic stress tolerance such as drought, low-temperature and salt stresses of plants. Therefore, the overexpressing transgenic plants have excellent tolerances to these abiotic stresses, whereby they may be useful as novel functional crops which are affected by climates and environments of the cultivated areas.

(c) In addition, where the plants are transformed with the present nucleotide sequence, the growth abilities of the transgenic plants are remarkably enhanced,

whereby they may effectively used for cultivating the plants with novel function of rapid growing, and biomass.

[0042] The present invention will now be described in further detail by examples. It would be obvious to those skilled in the art that these examples are intended to be more concretely illustrative and the scope of the present invention as set forth in the appended claims is not limited to or by the examples.

EXAMPLES

Cloning and analysis of AtSRPs gene

[0043] Total mRNA was extracted from 5-day-old *Arabidopsis thaliana* seedlings to perform RT-PCT (Seo *et al.*, 2008). cDNA was synthesized by reverse transcription. To amplify specifically *AtSRP1*, *AtSRP2* and *AtSRP3* genes, primer pairs used in PCR were prepared by searching cDNA sequence to each of *AtSRP1*, *AtSRP2* and *AtSRP3* genes from a database of the Arabidopsis Information Resource (TAIR, <http://www.arabidopsis.org>), and subsequently, tagging BamHI at CDS 5' and SacI at CDS 3'. Each of the amplified cDNA of *AtSRP1*, *AtSRP2* and *AtSRP3* gene was ligated to pGEM-T Easy vector (Promega, <http://www.promega.com>) to cloning, and the sequence was analyzed by sequencing reaction (Table 1). The analyzed sequence was exactly corresponded with that of TAIR (<http://www.arabidopsis.org>). In addition, it was found that At1g67360 (*AtSRP1*) included CDS consisting of 723 bp, At2g47780 (*AtSRP2*) included CDS consisting of 708 bp, and At3g05500 (*AtSRP3*) included CDS consisting of 741 bp.

Expression analysis of AtSRPs genes

[0044] To analyze the *AtSRPs* expression in transcription level in various tissues, RT-PCR and real-time RT-PCR were performed (Seo *et al.*, 2008). Primers used in the analysis are shown in Table 1.

[Table 1]

Primers for cloning, preparations of transgenic plants, and protein expressions		
Gene	Forward primer	Reverse primer
Cloning of <i>AtSRPs</i> ORF, RT-PCR and preparation of transgenic plant		
<i>AtSRP1</i>	5'-gcggatccatggagacagagaagaaaaatag-3'	5'-gcgagctctactccgaatcagacgatg-3'
<i>AtSRP2</i>	5'-gcggatccatggctgaagatgaaatagtagtc-3'	5'-gcgagctctcaatcagctcgacactgatc-3'
<i>AtSRP3</i>	5'-gcggatccatggctactcaaacggatctc-3'	5'-gcgagctctcaatcaagtggatggaactc-3'
<i>18s rRNA</i>	5'-gcattgccaaggatgtttt-3'	5'-gtacaaagggcagggacgta-3'
<i>At4CT8</i>	5'-tactgattacctcatgaagatccttac-3'	5'-aaacgatgtctctttagttagaagc-3'
Quantitative real time RT-PCR primer		
<i>AtSRP1</i>	5'-tgattctctcctgtcttctc-3'	5'-ttctctgtcgcctttagat-3'
<i>AtSRP2</i>	5'-tttctgctgatcgtaagg-3'	5'-aatgctcttgtagtgcca-3'

(continued)

Quantitative real time RT-PCR primer		
AtSRP3	5'-ccctgttgaggtccttaaat-3'	5'-gccacaatgggtgctat-3'

Expression and purification of AtSRPs proteins

[0045] AtSRPs were fused with 2xFlag epitope, expressed to BL21-CodonPlus (DE3) RIL (Stratagene), and purified. The proteins were quantified using BSA as a standard protein (Bradford, 1976).

Measurement for lectin binding properties of AtSRPs

[0046] To investigate binding properties to lectin of AtSRPs, two proteins were used as follows: 2xFlag-AtSRPs were expressed in E. coli, and purified to obtain protein; and 35S:2XFlag-AtSRPs were transiently over-expressed in tobacco to obtain protein.

[0047] The two proteins were subject to concanavalin A binding assay according to the manufacturer's method (Con A Sepharose 4B-GE-71707700AG; GEHealthcare Life Sciences). The resultants were subject to immunoblotting using Flag antibodies. As a result, the only band was elucidated in the lane of 35S:2XFlag-AtSRPs using tobacco system, indicating that the only AtSRP1 glycosylated in plant system binds to lectin. Accordingly, it would be understood that the AtSRP1 is a glycoprotein (Fig. 3a).

Measurement for membrane association properties of AtSRPs

[0048] To investigate association properties to membrane of AtSRPs, 35S:2XFlag-AtSRPs were transiently over-expressed in tobacco to obtain protein. The protein was extracted to use. Then, the protein was subject to membrane association test according to Kim and Bassham (Kim and Bassham, 2011). The resultants were subject to immunoblotting using Flag antibodies. As a result, the band of the supernatant in the 0.1 M Na₂CO₃-treated membrane pellet was darker than that of the precipitate (Fig. 3b), addressing that the AtSRP1 protein binding to membrane was eluted into the supernatant by Na₂CO₃ that weakens binding of the AtSRP1 protein to membrane. Accordingly, it would be understood that AtSRPs proteins are membrane associated proteins.

Preparation of AtSRPs-overexpressing transgenic plants and identification of knock-out mutants

[0049] The AtSRPs cDNA were introduced into the corresponding sites of the binary vector pBI121 to overexpress. The used primers are shown in Table 1. The resulting fusion gene was transferred to *Agrobacterium tumefaciens*. *Agrobacterium* cells were transformed into Arabidopsis plants by means of the floral-dip method (Joo et al., 2004). Seeds were germinated on 0.5 X Murashige and Skoog (MS) medium with 30 µg/ml kanamycin to select independent transgenic plants. *atsrp1* and *atsrp3* used T-DNA insertion lines (*atsrp1*: GABI_3095G05, *atsrp3*: WiscDSLOXHS192) which were purchased from ABRC (www.arabidopsis.org). *atsrp2* used knock-down line, which is prepared by RNAi method, to select plants in which expression in transcription level is lower (Lee and Kim, 2010).

Phenotypic analyses of AtSRPs-overexpressing transgenic plants

[0050] Medium was prepared to contain 0.5 x MS (including vitamin B5), 1% sucrose and 0.8% agar (select agar; Life Technology, Rockville, MD, USA) (pH 5.7). Surfaces of seeds of wild type plants and AtSRPs-overexpressing transgenic plants were sterilized, planted and grown under light condition at 22°C. After inhibition of radicle germination, they were daily observed for 5 days. In order to observe the root growth, seeds were observed until 5th day after germination. During the root growth, the root shoot-tip was marked in outside of culture plates, and the root length was measured using ScionImage software (Scion Corp., Frederic, MD, USA). In order to observe the leaf size, the seeds were planted in sterile soil. After 12 days, sizes of the leaves from a seed leaf to a third leaf in wild type plants and AtSRPs-overexpressing transgenic plants were compared. In addition, the second leaf was measured in leaf length, leaf width and petiole length using ScionImage software (Scion Corp., Frederic, MD, USA).

[0051] In order to observe the flower stalk rising time, seeds of wild type plants and overexpressing transgenic plants were planted in sterile soil, and grown under light condition at 22°C. At 19 days after germination, the flower stalk length was measured. As a result, when wild type plants and AtSRPs-overexpressing transgenic plants were compared each other, there was no difference in germination ability. However, AtSRPs-overexpressing transgenic plants grew more rapidly with longer roots, larger leaves, and longer height. Therefore, it could be demonstrated that AtSRPs act as a

positive regulator in plant growth (Figs. 4a-d).

Phenotypic analyses using microscope

[0052] Leaves of 5-day-old wild-type and transgenic *Arabidopsis thaliana* were washed in a solution of chloral hydrate with the reported method (Kwon et al. 2009). Cell layer was observed by bright-field microscope (BX51 fluorescence microscope, Olympus, Japan). For longitudinal views of root tips, roots of 5-day-old wild-type and transgenic *Arabidopsis thaliana* were stained by propidium iodide, and the images were obtained by confocal microscope with the method of Seo et al. (2008) (Fig. 5).

Survival rate determination of wild-type and transgenic Arabidopsis thaliana plants after high salt stress

[0053] 5-day-old wild type, 35S:CaSRP1 transgenic and each of knock-out plants, which had been grown under normal growth conditions, were subjected to high salt stress by growing in high salt-media containing 100mM of NaCl for 7 days. Then, root lengths of each of plants were examined. As a result, each of the knock-out plants has a more sensitive phenotype to high salt stress compared with wild-type plants. Therefore, it could be demonstrated that AtSRPs act as a positive regulator in high salt stress response (Fig. 6).

Survival rate determination of wild-type and transgenic Arabidopsis thaliana plants after drought stress

[0054] 3-week-old wild type, 35S:CaSRP1 transgenic and each of knock-out plants, which had been grown under normal growth conditions, were subjected to drought stress by withholding water for 8 days. The plants were then re-watered and their phenotypes were examined after 3 days. Survival was defined as the ability to resume growth when returned to normal conditions following water stress (Cho et al. 2008). Each of survival rates was as follows (Fig. 7a):

wild type plants, 26.97% (24 out of 89 plants); 35S:AtSRP1 transgenic plants, 64.84% (59 out of 91 plants); 35S:AtSRP2 transgenic plants, 66.29% (59 out of 89 plants); 35S:AtSRP3 transgenic plants, 58.02% (47 out of 81 plants); atsrp1 mutant plants, 3.90% (3 out of 77 plants); atsrp3 mutant plants, 0.00% (0 out of 101 plants). In addition, transgenic plants showed significantly lower decreases of the moisture contents by the lapse of time, as compared to the wild-type and knock-out mutant plants (Fig. 7b).

Chlorophyll and proline content measurements of wild-type and transgenic Arabidopsis thaliana plants after drought stress

[0055] Chlorophyll was extracted in leaves of wild-type and AtSRPs-overexpressing transgenic *Arabidopsis thaliana* plants after drought stress by the method of Bae et al. (2009) using 80% acetone. The remaining leaf extract was dried at 105°C for 16 hours, and weight of the dry extract was measured (Walti et al. 2002). Proline content of the leaves was measured by the method of Claussen (2005). As a result, transgenic plants exhibited significantly higher levels of proline contents, indicating that transgenic plants retained their leaf water more effectively than did wild-type leaves under drought conditions. In contrast, knock-out mutant plants showed opposite phenotypes (Fig. 8).

[0056] Those skilled in the art will appreciate that the conceptions and specific embodiments disclosed in the foregoing description may be readily utilized as a basis for modifying or designing other embodiments for carrying out the same purposes of the present disclosure. Those skilled in the art will also appreciate that such equivalent embodiments do not depart from the spirit and scope of the disclosure as set forth in the appended claims.

<110> Industry-Academic Coopreation Foundation Yonsei University

<120> Gene Implicated in Abiotic Stress Tolerance and Growth Accelerating and Use Thereof

<130> PN120347

<160> 6

<170> KopatentIn 2.0

<210> 1

<211> 723

<212> DNA

<213> Arabidopsis thaliana

<400> 1

atggagacag agaagaaaaa tagcaaggag gtagcgctaa aacacctagc gttagctgaga 60

attgcaacga tccatattct ggcttccgct tcaaattctt acgaatacgc taaacaaaac 120

tccggctctc tcaaattccg cgtggaaaag gtcgaaggag ccgtcaccac cgtcgtcacc 180

cctgtctatc agaaattcaa agatgttctt gattctctcc tgggtctttct agatcacaag 240

gtgggtgaag tttcgtacaa gtttgatgag catgctcctc caatggctaa gaaagtgggtg 300

aatcaagcac atgtattgat ctacaaggcg acagagaaaag ctcaaagctt tgtgaaagag 360

gctcgtaccg gtgggtccaa agctgcgttt aactatgctg caactgagta caagtttttc 420

gttggtgacca actcgggttaa agtctgggct aaacttaacc agtataaacc aatccatgca 480

atgggtgaca aagctttgcc tgtggctgca cacttttcca gtcgggtacaa cgatttggtg 540

actgatatga ctaatatggg ttactctttg gttgggtatc ttccgttggt tcctgttgat 600

gacattgtta aggcttatga aaaggaagat gcaaggagaa agaaaggagg ggatacggct 660

ggaaagaaaag gagagactac tgatgcggct gatggtgata aatcatcgtc tgattcggag 720

tag 723

<210> 2

<211> 708

<212> DNA

<213> Arabidopsis thaliana

<400> 2

atggctgaag atgaaatagt agtcgaagaa gagcaatcac agccacagga gattactcca 60

gttccaccat cttcttcgtc ttgccttcg ttagtggtgg aagacgacga tgagatgaag 120

ctcaagcact tggaattcat tcaagtgcg gcgggttact tcgccgcctg tttctcgacc 180

ctctacgagt tggctaaaga caacgctggt cactcaaac tcggcgctcga gaacattgaa 240

gattgcgttc gaaccgtgct tgctccatta tacgaaaaat tccacgacgt tcctttcaag 300

cttctccttt tcgtcgatcg taaggtggac gatgtgttct tcgatgttga gacatatgtt 360

ccgtcattgg tgaagcaagc ttctagccaa gcgctcacgg tggccacgga ggtgcagcgc 420

EP 2 840 141 A9

	accggcgcttg tggacactac caagagcatt gcaagaagcg tccgggacaa gtacgagcca	480
	gcgccagaggt attacgcagc gacgctgtgg cgtttactga atcaactgcc gttgttccct	540
5	gaagttgctg atttagtgat cccacggcg ttttactggc cagagaagta caacgatgcg	600
	gttcgttacg ttggcgacag agactacttt gggcgagagc atctacccat gattcctatt	660
	gagaagatct ctgatatttt ggaacaagat cagtgtcgag ctgattga	708
10		
	<210> 3	
	<211> 741	
	<212> DNA	
	<213> Arabidopsis thaliana	
15		
	<400> 3	
	atggctactc aaacggatct cgctcagccc aagcttgata tgaccaagga ggagaaagag	60
	aggttgaagt atttgcaatt cgtgcaagct gctgctgtgg aagctctgct tcgctttgct	120
20	cttattttacg ctaaggcaaa ggacaagtct ggctccttga aacctgggtg tgaatctggt	180
	gaaggagctg tcaagactgt cgttggtcct gtctacgaga aataccacga cgtccctggt	240
	gaggctcctta aatacatgga ccagaagggt gatatgtctg tgactgagct tgaccgtcgt	300
25	gtcccaccag tcgtcaagca agtgtctgcc caagccatct ccgctgctca gatagcacc	360
	attgtggcac gtgcgttggc ctctgagggt cgacgtgctg gtgttgttga aaccgcttct	420
	ggaatggcta aatccgtcta ctccaagtac gagcctgctg ctaaggagtt gtatgcaaac	480
30	tatgagccaa aagcagagca gtgtgccgtt tcagcttga agaagcttaa ccagcttct	540
	ctattcccaa ggctggctca agttgctgta ccaacagctg ctttctgctc tgagaagtac	600
	aatgatactg tggtaaggc tgcagagaaa gggtagagag tcacatcgta catgccattg	660
35	gttccaacag agaggatctc aaaaatcttc gctgaggaga aagctgagac cgagcctttg	720
	gagttccatc cacttgattg a	741
40		
	<210> 4	
	<211> 240	
	<212> PRT	
	<213> Arabidopsis thaliana	
45		
	<400> 4	
	Met Glu Thr Glu Lys Lys Asn Ser Lys Glu Val Ala Leu Lys His Leu	
	1 5 10 15	
	Ala Phe Val Arg Ile Ala Thr Ile His Ile Leu Ala Ser Val Ser Asn	
50	20 25 30	
	Leu Tyr Glu Tyr Ala Lys Gln Asn Ser Gly Pro Leu Lys Ser Ala Val	
	35 40 45	
	Glu Lys Val Glu Gly Ala Val Thr Thr Val Val Thr Pro Val Tyr Gln	
55	50 55 60	

EP 2 840 141 A9

	Lys	Phe	Lys	Asp	Val	Pro	Asp	Ser	Leu	Leu	Val	Phe	Leu	Asp	His	Lys	65	70	75	80
5	Val	Gly	Glu	Val	Ser	Tyr	Lys	Phe	Asp	Glu	His	Ala	Pro	Pro	Met	Ala	85	90	95	
	Lys	Lys	Val	Val	Asn	Gln	Ala	His	Val	Leu	Ile	Tyr	Lys	Ala	Thr	Glu	100	105	110	
10	Lys	Ala	Gln	Ser	Phe	Val	Lys	Glu	Ala	Arg	Thr	Gly	Gly	Pro	Lys	Ala	115	120	125	
	Ala	Phe	Asn	Tyr	Ala	Ala	Thr	Glu	Tyr	Lys	Phe	Phe	Val	Val	Thr	Asn	130	135	140	
15	Ser	Val	Lys	Val	Trp	Ala	Lys	Leu	Asn	Gln	Tyr	Lys	Pro	Ile	His	Ala	145	150	155	160
	Met	Gly	Asp	Lys	Ala	Leu	Pro	Val	Ala	Ala	His	Phe	Ser	Ser	Arg	Tyr	165	170	175	
20	Asn	Asp	Leu	Val	Thr	Asp	Met	Thr	Asn	Met	Gly	Tyr	Ser	Leu	Val	Gly	180	185	190	
	Tyr	Leu	Pro	Leu	Val	Pro	Val	Asp	Asp	Ile	Val	Lys	Ala	Tyr	Glu	Lys	195	200	205	
25	Glu	Asp	Ala	Arg	Arg	Lys	Lys	Gly	Gly	Asp	Thr	Ala	Gly	Lys	Lys	Gly	210	215	220	
	Glu	Thr	Thr	Asp	Ala	Ala	Asp	Gly	Asp	Lys	Ser	Ser	Ser	Asp	Ser	Glu	225	230	235	240
30																				
	<210>		5																	
	<211>		235																	
	<212>		PRT																	
	<213>		Arabidopsis thaliana																	
35																				
	<400>		5																	
	Met	Ala	Glu	Asp	Glu	Ile	Val	Val	Glu	Glu	Glu	Gln	Ser	Gln	Pro	Gln	1	5	10	15
40	Glu	Ile	Thr	Pro	Val	Pro	Pro	Ser	Ser	Ser	Ser	Ser	Pro	Ser	Leu	Val	20	25	30	
	Val	Glu	Asp	Asp	Asp	Glu	Met	Lys	Leu	Lys	His	Leu	Glu	Phe	Ile	Gln	35	40	45	
45	Val	Ala	Ala	Val	Tyr	Phe	Ala	Ala	Cys	Phe	Ser	Thr	Leu	Tyr	Glu	Leu	50	55	60	
	Ala	Lys	Asp	Asn	Ala	Gly	Pro	Leu	Lys	Leu	Gly	Val	Glu	Asn	Ile	Glu	65	70	75	80
50	Asp	Cys	Val	Arg	Thr	Val	Leu	Ala	Pro	Leu	Tyr	Glu	Lys	Phe	His	Asp	85	90	95	
	Val	Pro	Phe	Lys	Leu	Leu	Leu	Phe	Val	Asp	Arg	Lys	Val	Asp	Asp	Val	100	105	110	
55	Phe	Phe	Asp	Val	Glu	Thr	Tyr	Val	Pro	Ser	Leu	Val	Lys	Gln	Ala	Ser	115	120	125	

EP 2 840 141 A9

	Ser	Gln	Ala	Leu	Thr	Val	Ala	Thr	Glu	Val	Gln	Arg	Thr	Gly	Val	Val	
	130						135					140					
5	Asp	Thr	Thr	Lys	Ser	Ile	Ala	Arg	Ser	Val	Arg	Asp	Lys	Tyr	Glu	Pro	
	145					150					155					160	
	Ala	Ala	Glu	Tyr	Tyr	Ala	Ala	Thr	Leu	Trp	Arg	Leu	Leu	Asn	Gln	Leu	
					165					170					175		
10	Pro	Leu	Phe	Pro	Glu	Val	Ala	His	Leu	Val	Ile	Pro	Thr	Ala	Phe	Tyr	
				180					185					190			
	Trp	Ser	Glu	Lys	Tyr	Asn	Asp	Ala	Val	Arg	Tyr	Val	Gly	Asp	Arg	Asp	
			195				200						205				
15	Tyr	Phe	Gly	Ala	Glu	Tyr	Leu	Pro	Met	Ile	Pro	Ile	Glu	Lys	Ile	Ser	
	210						215					220					
	Asp	Ile	Leu	Glu	Gln	Asp	Gln	Cys	Arg	Ala	Asp						
	225					230					235						
20																	
	<210>	6															
	<211>	246															
	<212>	PRT															
	<213>	Arabidopsis thaliana															
25																	
	<400>	6															
	Met	Ala	Thr	Gln	Thr	Asp	Leu	Ala	Gln	Pro	Lys	Leu	Asp	Met	Thr	Lys	
	1				5					10					15		
30	Glu	Glu	Lys	Glu	Arg	Leu	Lys	Tyr	Leu	Gln	Phe	Val	Gln	Ala	Ala	Ala	
				20					25					30			
	Val	Glu	Ala	Leu	Leu	Arg	Phe	Ala	Leu	Ile	Tyr	Ala	Lys	Ala	Lys	Asp	
			35				40						45				
35	Lys	Ser	Gly	Pro	Leu	Lys	Pro	Gly	Val	Glu	Ser	Val	Glu	Gly	Ala	Val	
	50						55					60					
	Lys	Thr	Val	Val	Gly	Pro	Val	Tyr	Glu	Lys	Tyr	His	Asp	Val	Pro	Val	
	65					70					75					80	
40	Glu	Val	Leu	Lys	Tyr	Met	Asp	Gln	Lys	Val	Asp	Met	Ser	Val	Thr	Glu	
					85					90					95		
	Leu	Asp	Arg	Arg	Val	Pro	Pro	Val	Val	Lys	Gln	Val	Ser	Ala	Gln	Ala	
			100						105					110			
45	Ile	Ser	Ala	Ala	Gln	Ile	Ala	Pro	Ile	Val	Ala	Arg	Ala	Leu	Ala	Ser	
			115					120					125				
	Glu	Val	Arg	Arg	Ala	Gly	Val	Val	Glu	Thr	Ala	Ser	Gly	Met	Ala	Lys	
			130				135					140					
50	Ser	Val	Tyr	Ser	Lys	Tyr	Glu	Pro	Ala	Ala	Lys	Glu	Leu	Tyr	Ala	Asn	
	145					150					155					160	
	Tyr	Glu	Pro	Lys	Ala	Glu	Gln	Cys	Ala	Val	Ser	Ala	Trp	Lys	Lys	Leu	
				165						170					175		
55	Asn	Gln	Leu	Pro	Leu	Phe	Pro	Arg	Leu	Ala	Gln	Val	Ala	Val	Pro	Thr	

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Claims

1. A composition comprising a nucleotide sequence encoding the amino acid sequence selected from the group consisting of SEQ ID NOs:4 to 6 for use in the improvement of the growth of a plant.
2. The composition according to claim 1, wherein the nucleotide sequence encoding the amino acid sequence of SEQ ID NO:4 comprises the nucleotide sequence as set forth in SEQ ID NO:1.
3. The composition according to claim 1, wherein the nucleotide sequence encoding the amino acid sequence of SEQ ID NO:5 comprises the nucleotide sequence as set forth in SEQ ID NO:2.
4. The composition according to claim 1, wherein the nucleotide sequence encoding the amino acid sequence of SEQ ID NO:6 comprises the nucleotide sequence as set forth in SEQ ID NO:3.
5. The composition according to claim 1, wherein the nucleotide sequence is contained a recombinant plant expression vector; and the recombinant plant expression vector comprises (i) the nucleotide sequence; (ii) a promoter which is operatively linked to the nucleotide sequence of (i) and generates RNA molecules in plant cells; and (iii) a poly A signal sequence inducing polyadenylation at the 3'-end of the RNA molecules.
6. A method for promoting the growth of a plant, comprising:
 - (a) introducing a nucleotide sequence encoding the amino acid sequence selected from the group consisting of SEQ ID NOs:4 to 6 to a cell of the plant; and
 - (b) obtaining a transgenic plant exhibiting improved tolerance to an abiotic stress from the cell of the plant.
7. A composition comprising a nucleotide sequence encoding the amino acid sequence selected from the group consisting of SEQ ID NOs:4 to 6 for use in the improvement of the tolerance of a plant to an abiotic stress.
8. The composition according to claim 7, wherein the nucleotide sequence encoding the amino acid sequence of SEQ ID NO:4 comprises the nucleotide sequence as set forth in SEQ ID NO:1.
9. The composition according to claim 7, wherein the nucleotide sequence encoding the amino acid sequence of SEQ ID NO:5 comprises the nucleotide sequence as set forth in SEQ ID NO:2.
10. The composition according to claim 7, wherein the nucleotide sequence encoding the amino acid sequence of SEQ ID NO:6 comprises the nucleotide sequence as set forth in SEQ ID NO:3.
11. The composition according to claim 7, wherein the abiotic stress is selected from the group consisting of a drought stress, a low-temperature stress and a salt stress.
12. The composition according to claim 7, wherein the nucleotide sequence is contained a recombinant plant expression vector; and the recombinant plant expression vector comprises (i) the nucleotide sequence; (ii) a promoter which

is operatively linked to the nucleotide sequence of (i) and generates RNA molecules in plant cells; and (iii) a poly A signal sequence inducing polyadenylation at the 3'-end of the RNA molecules.

13. A method for improving the tolerance of a plant to an abiotic stress, comprising:

- 5
- (a) introducing a nucleotide sequence encoding the amino acid sequence selected from the group consisting of SEQ ID NOs:4 to 6 to a cell of the plant; and
 - (b) obtaining a transgenic plant exhibiting improved tolerance to an abiotic stress from the cell of the plant.
- 10
- 15
- 20
- 25
- 30
- 35
- 40
- 45
- 50
- 55

Fig. 1a




Fig. 1b




Fig. 2a





Fig. 2b



Fig. 3a**Lectin binding assay****Fig. 3b****Membrane associated protein**




Fig. 4a

Fig. 4b

5-day-old seedling




Fig. 4c





Fig. 4d



WT 35S:AtSRP1 35S:AtSRP2 35S:AtSRP3




Fig. 5a




Fig. 5b





Fig. 6



12-day-old seedling

Fig. 7a





Fig. 7b




Fig. 7c

Fig. 8**Fig. 9**



EUROPEAN SEARCH REPORT

Application Number
EP 13 18 1236

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	WO 2009/013750 A2 (EVOGENE LTD [IL]; RONEN GIL [IL]; KARCHI HAGAI [IL]; DIBER ALEX [IL];) 29 January 2009 (2009-01-29) * sequences 267,1557 * * page 79, line 30 - page 80, line 2 * * tables 14,37,106,107 * -----	1,2,5-8, 11-13	INV. C12N15/82
X	WO 2011/108795 A1 (UNIV YONSEI IACF [KR]; KIM WOO TAEK [KR]; KIM EUN YU [KR]; SEO JI HO []) 9 September 2011 (2011-09-09) *sequence At1g37360.1* -----	1,2,5-8, 11-13	
			TECHNICAL FIELDS SEARCHED (IPC)
			C12N
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 10 January 2014	Examiner Blanco Urgoiti, B
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.02 (P04C01)



Application Number

EP 13 18 1236

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing claims for which payment was due.

☐ Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for those claims for which no payment was due and for those claims for which claims fees have been paid, namely claim(s):

☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for those claims for which no payment was due.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet B

☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.

☐ As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.

☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:

☒ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:

2, 8(completely); 1, 5-7, 11-13(partially)

☐ The present supplementary European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims (Rule 164 (1) EPC).



LACK OF UNITY OF INVENTION
SHEET B

Application Number

EP 13 18 1236

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. claims: 2, 8(completely); 1, 5-7, 11-13(partially)

A composition comprising the amino acid sequence of SEQ ID NO:4 for use in the improvement of plant growth or tolerance to an abiotic stress; encoded by the polynucleotide of SEQ ID NO:1; comprised in a plant expression vector; methods for promoting plant growth and abiotic stress tolerance thereof.

2. claims: 3, 9(completely); 1, 5-7, 11-13(partially)

A composition comprising the amino acid sequence of SEQ ID NO:5 for use in the improvement of plant growth or tolerance to an abiotic stress; encoded by the polynucleotide of SEQ ID NO:2; comprised in a plant expression vector; methods for promoting plant growth and abiotic stress tolerance thereof.

3. claims: 4, 10(completely); 1, 5-7, 11-13(partially)

A composition comprising the amino acid sequence of SEQ ID NO:6 for use in the improvement of plant growth or tolerance to an abiotic stress; encoded by the polynucleotide of SEQ ID NO:3; comprised in a plant expression vector; methods for promoting plant growth and abiotic stress tolerance thereof.

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 13 18 1236

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-01-2014

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2009013750 A2	29-01-2009	AU 2008278654 A1	29-01-2009
		CA 2694481 A1	29-01-2009
		CN 102037127 A	27-04-2011
		EP 2183371 A2	12-05-2010
		US 2010319088 A1	16-12-2010
		US 2014068819 A1	06-03-2014
		WO 2009013750 A2	29-01-2009

WO 2011108795 A1	09-09-2011	KR 20110100417 A	14-09-2011
		WO 2011108795 A1	09-09-2011

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5004863 A [0024]
- US 5349124 A [0024]
- US 5416011 A [0024]

Non-patent literature cited in the description

- **SMITH ; WATERMAN.** *Adv. Appl. Math.*, 1981, vol. 2, 482 [0021]
- **NEEDLEMAN ; WUNSCH.** *J. Mol. Bio.*, 1970, vol. 48, 443 [0021]
- **PEARSON ; LIPMAN.** *Methods in Mol. Biol.*, 1988, vol. 24, 307-31 [0021]
- **HIGGINS ; SHARP.** *Gene*, 1988, vol. 73, 237-44 [0021]
- **HIGGINS ; SHARP.** *CABIOS*, 1989, vol. 5, 151-3 [0021]
- **CORPET et al.** *Nuc. Acids Res.*, 1988, vol. 16, 10881-90 [0021]
- **HUANG et al.** *Comp. Appl. BioSci.*, 1992, vol. 8, 155-65 [0021]
- **PEARSON et al.** *Meth. Mol. Biol.*, 1994, vol. 24, 307-31 [0021]
- **ALTSCHUL et al.** *J. Mol. Biol.*, 1990, vol. 215, 403-10 [0021]
- *Methods of Enzymology*, 1987, vol. 153 [0023]
- **CHILTON et al.** *Cell*, 1977, vol. 11, 263-271 [0023]
- **LORZ et al.** *Mol. Genet.*, 1985, vol. 199, 178-182 [0023]
- **SAMBROOK et al.** *Molecular Cloning, A Laboratory Manual.* Cold Spring Harbor Laboratory Press, 2001 [0029]
- **BEVAN et al.** *Nucleic Acids Research*, 1983, vol. 11 (2), 369-385 [0031]