



(11) **EP 2 877 446 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:

**Corrected version no 1 (W1 B1)**  
**Corrections, see**  
**Description Paragraph(s) 11**  
**Claims EN 1**  
**Claims FR 1**

(51) Int Cl.:

<b>C07C 259/08</b> <sup>(2006.01)</sup>	<b>C07D 221/20</b> <sup>(2006.01)</sup>
<b>A61K 31/16</b> <sup>(2006.01)</sup>	<b>A61K 31/496</b> <sup>(2006.01)</sup>
<b>A61K 31/4465</b> <sup>(2006.01)</sup>	<b>A61K 31/4523</b> <sup>(2006.01)</sup>
<b>A61K 31/55</b> <sup>(2006.01)</sup>	<b>A61P 7/02</b> <sup>(2006.01)</sup>
<b>C07D 401/04</b> <sup>(2006.01)</sup>	<b>C07D 401/06</b> <sup>(2006.01)</sup>
<b>C07D 401/12</b> <sup>(2006.01)</sup>	<b>C07C 317/14</b> <sup>(2006.01)</sup>
<b>C07C 317/48</b> <sup>(2006.01)</sup>	<b>C07D 409/06</b> <sup>(2006.01)</sup>
<b>C07D 471/10</b> <sup>(2006.01)</sup>	<b>C07D 471/20</b> <sup>(2006.01)</sup>
<b>C07D 265/34</b> <sup>(2006.01)</sup>	<b>C07D 491/107</b> <sup>(2006.01)</sup>
<b>C07D 211/34</b> <sup>(2006.01)</sup>	

(48) Corrigendum issued on:

**25.01.2017 Bulletin 2017/04**

(45) Date of publication and mention of the grant of the patent:

**14.09.2016 Bulletin 2016/37**

(86) International application number:

**PCT/EP2013/065071**

(21) Application number: **13737275.1**

(87) International publication number:

**WO 2014/012964 (23.01.2014 Gazette 2014/04)**

(22) Date of filing: **17.07.2013**

(54) **NEW ANTIFIBRINOLYTIC COMPOUNDS**

NEUE ANTIFIBRINOLYTISCHE VERBINDUNGEN

NOUVEAUX COMPOSÉS ANTIFIBRINOLYTIQUES

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**

(74) Representative: **ZBM Patents - Zea, Barlocchi & Markvardsen**

**Plaza Catalunya, 1 2nd floor**  
**08002 Barcelona (ES)**

(30) Priority: **18.07.2012 EP 12382285**

(56) References cited:

**US-A- 3 868 411 US-A- 4 483 867**  
**US-A1- 2002 099 035**

(43) Date of publication of application:

**03.06.2015 Bulletin 2015/23**

(73) Proprietor: **Proyecto de Biomedicina Cima, S.L.**

**31180 Zizur Mayor (ES)**

- **OSAMU KITAGAWA ET AL: "Stereoselective Iodine Atom Transfer [3 + 2] Cycloaddition Reaction with Alkenes Using Unsymmetrical Allylated Active Methine Radicals", THE JOURNAL OF ORGANIC CHEMISTRY, vol. 69, no. 7, 1 April 2004 (2004-04-01), pages 2607-2610, XP055043559, ISSN: 0022-3263, DOI: 10.1021/jo035771a**
- **BOUYSSI D ET AL: "Rearrangement of oxaspiroheptanes to cyclohexanones mediated by lithium iodide", SYNLETT 2000 DE, no. 5, 2000, pages 749-751, XP008157852, ISSN: 0936-5214**

(72) Inventors:

- **ORBE LOPATEGUI, Josune**  
**E-31008 Pamplona (ES)**
- **OYARZABAL SANTAMARINA, Julen**  
**E-31008 Pamplona (ES)**
- **PÁRAMO FERNÁNDEZ, José Antonio**  
**E-31008 Pamplona (ES)**
- **RODRÍGUEZ GARCÍA, José Antonio**  
**E-31008 Pamplona (ES)**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 2 877 446 B9**

**Description**

**[0001]** The present invention relates to spirocyclic compounds of formula (I), as well as to intermediates used in their preparation. It also relates to pharmaceutical or veterinary compositions containing them, and to their use in medicine, in particular as antifibrinolytic and antihemorrhagic agents.

**BACKGROUND ART**

**[0002]** The haemostatic system is responsible for maintaining circulatory fluidity and for preventing haemorrhage in response to vascular injury. Physiological hemostasis is controlled by mechanisms of coagulation and the formation of fibrin and by those favouring the degradation of fibrin (fibrinolysis).

**[0003]** Hyperfibrinolytic states caused by congenital or acquired conditions predispose to important haemorrhagic complications, often requiring transfusions and the need for re-exploration having a detrimental effect on patient outcome. Hemorrhage is responsible for almost 50% of deaths occurring within 24 hours of traumatic injury and for up to 80% of intraoperative trauma mortality. In western countries about one third of in-hospital deaths due to trauma is caused by abnormal blood loss which is an important contributory factor for other causes of death, particularly multi-organ failure, requiring massive blood transfusion. Failure to start appropriate early management in bleeding trauma patients is a leading cause of preventable death from trauma. Post-partum hemorrhage (PPH) is another leading cause of death in the developing world, accounting for 25% of maternal deaths, and rose in the developed world from 1.5% in 1999 to 4.1 % in 2009. The risk of haemorrhage can also be important in cardiovascular patients on anti-coagulant therapy. Pharmacological approaches are an important part of multimodal therapy aiming to reducing bleeding and transfusion in order to reverse specific defects associated with such states; among them, the role of fibrinolysis inhibitors is growing.

**[0004]** It is well known that subjects who bleed excessively in association with surgery or major trauma and need blood transfusions develop more complications than those who do not experience any bleeding. However, moderate bleeding requiring the administration of human blood products may lead to complications associated with the risk of transferring human viruses. Extensive bleeding requiring massive blood transfusions may lead to the development of multiple organ failure including lung or kidney function. Therefore, a major goal in surgery as well as in the treatment of major tissue damage is to avoid or minimise bleeding in order to ensure the formation of stable and solid haemostatic plugs that are not easily dissolved by fibrinolytic enzymes. Furthermore, it is of importance to ensure quick and effective formation of such plugs or clots.

**[0005]** Antifibrinolytic agents are widely used in major surgery to prevent fibrinolysis and reduce blood loss. Currently two synthetic lysine analogs, epsilon-aminocaproic acid (EACA) and tranexamic acid (TXA), are the only antifibrinolytics commercially available to control bleeding. These agents competitively inhibit activation of plasminogen to plasmin, an enzyme that degrades fibrin clots, fibrinogen and other plasma proteins.

**[0006]** However, there are some concerns with these currently available antifibrinolytic agents due to the potential risk of thrombotic complications.

**[0007]** There is still a need for improved treatment of subjects experiencing bleeding episodes, including subjects where the bleeding episodes are due to surgery, trauma, or other forms of tissue damage, as well as in clinical scenarios characterized by excessive fibrinolysis.

**[0008]** US3868411 A describes isobutyl cyclohexene hydroxamic acid derivatives as antifibrinolytic agents.


**[0009]** US4483867 A sets forth tranexamic acid derivatives antifibrinolytically active.

**[0010]** US2002/099035 A1 reports hydroxamic acid derivatives as anti-thrombotic agents

**SUMMARY OF THE INVENTION**

**[0011]** Inventors have conceived and produced found a novel series of spirocyclic compounds that shows good anti-fibrinolytic and antihemorrhagic properties. In particular, the spirocyclic compounds which comprise spirocyclic ring system containing a carbon atom (spiro atom) attached to a hydroxamic acid and a sulfonyl group, show a significant delay in the lysis time in a thromboelastometry assay. In addition, the spirocyclic compounds of the invention also show an important reduction of the bleeding time in vivo animal models as it will be shown in detail in the examples. These characteristics of the compounds of the invention allow a rapid cessation of hemorrhage; favor an effective formation of plugs or clots; have a sustained action (persistence of the clot and prevention of hemorrhage) and aid in minimizing the adverse effects related to other antifibrinolytic/antihemorrhagic treatments having risk of thrombotic complications.

**[0012]** Therefore, a first aspect of the invention relates to a compound of formula (I) (hereinafter also referred as compounds of the invention), or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt



(I)

wherein

A and B form a spirocyclic ring system wherein the spiro atom connecting A and B is a carbon atom and wherein

A is a known 3- to 8-membered carbocyclic or heterocyclic monocyclic ring, saturated or partially unsaturated; or alternatively

A is a known 6- to 18-membered carbocyclic or heterocyclic polycyclic ring system, saturated, partially unsaturated, or partially aromatic; and

B is a known 4- to 7-membered carbocyclic or heterocyclic monocyclic ring, saturated or partially unsaturated;

C is phenyl or a known 5- to 6-membered heteroaromatic ring;

R<sub>1</sub>-R<sub>3</sub> are independently selected from H, halogen, -NO<sub>2</sub>, -CN, R<sup>a</sup>, -OR<sup>a</sup>, -OC(Y)R<sup>a</sup>, -OC(Y)OR<sup>a</sup>, -OC(Y)NR<sup>b</sup>R<sup>a</sup>, -OSO<sub>2</sub>OR<sup>a</sup>, -NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>C(Y)R<sup>a</sup>, -NR<sup>b</sup>C(Y)OR<sup>a</sup>, -NR<sup>b</sup>C(Y)NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>S(O)<sub>2</sub>R<sup>a</sup>, -NR<sup>b</sup>SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -SR<sup>a</sup>, -S(O)R<sup>a</sup>, -S(O)OR<sup>a</sup>, -SO<sub>2</sub>R<sup>a</sup>, -SO<sub>2</sub>(OR<sup>a</sup>), -SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -SC(Y)NR<sup>b</sup>R<sup>a</sup>, -C(Y)R<sup>a</sup>, -C(Y)OR<sup>a</sup>, -C(Y)NR<sup>b</sup>R<sup>a</sup>, -C(Y)NR<sup>b</sup>OR<sup>a</sup>, and -C(O)NR<sup>b</sup>SO<sub>2</sub>R<sup>a</sup>;

R<sub>4</sub>-R<sub>7</sub> are independently selected from halogen, -NO<sub>2</sub>, -CN, R<sup>c</sup>, -OR<sup>c</sup>, -NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>C(Y)R<sup>c</sup>, -NR<sup>d</sup>C(Y)OR<sup>c</sup>, -NR<sup>d</sup>C(Y)NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>S(O)<sub>2</sub>R<sup>c</sup>, -NR<sup>d</sup>SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>, -SR<sup>c</sup>, -S(O)R<sup>c</sup>, -S(O)OR<sup>c</sup>, -SO<sub>2</sub>R<sup>c</sup>, -SO<sub>2</sub>R(OR<sup>c</sup>), -SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>, -SC(Y)NR<sup>d</sup>R<sup>c</sup>, -C(Y)R<sup>c</sup>, -C(Y)OR<sup>c</sup>, -C(Y)NR<sup>d</sup>R<sup>c</sup>, -C(Y)NR<sup>d</sup>OR<sup>c</sup>, and -C(O)NR<sup>d</sup>SO<sub>2</sub>R<sup>c</sup>

R<sup>a</sup> is a saturated or unsaturated (C<sub>1</sub>-C<sub>12</sub>)alkyl optionally substituted with one or more substituents R<sup>e</sup> and/or one Cy<sup>1</sup>; or alternatively R<sup>a</sup> is Cy<sup>2</sup>;

wherein Cy<sup>1</sup> and Cy<sup>2</sup> are independently optionally substituted with: one Cy<sup>3</sup> and/or one or more substituents R<sup>e</sup>, and/or one or more saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl groups optionally substituted with one or more substituents R<sup>e</sup> and/or one Cy<sup>3</sup>; and

wherein any Cy<sup>3</sup> is optionally substituted with one or more substituents independently selected from R<sup>e</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>e</sup>;

each R<sup>a</sup> and R<sup>b</sup> are independently H or R<sup>a</sup>;

R<sup>c</sup> and each R<sup>d</sup> are independently selected from H, Cy<sup>4</sup>, and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>h</sup> and/or one Cy<sup>5</sup>;

wherein Cy<sup>4</sup> is optionally substituted with one or more substituents independently selected from R<sup>h</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>h</sup>; and

wherein Cy<sup>5</sup> is optionally substituted with one or more substituents independently selected from R<sup>h</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>h</sup>;

each R<sup>e</sup> is independently selected from halogen, -NO<sub>2</sub>, -CN, -OR<sup>f</sup>, -OC(Y)R<sup>f</sup>, -OC(Y)OR<sup>f</sup>, -OC(Y)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>C(Y)R<sup>f</sup>, -NR<sup>g</sup>C(Y)OR<sup>f</sup>, -NR<sup>g</sup>C(Y)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>S(O)<sub>2</sub>R<sup>f</sup>, -NR<sup>g</sup>SO<sub>2</sub>NR<sup>g</sup>R<sup>f</sup>, -SR<sup>f</sup>, -S(O)R<sup>f</sup>, -S(O)OR<sup>f</sup>, -SO<sub>2</sub>R<sup>f</sup>,

## EP 2 877 446 B9

$-\text{SO}_2(\text{OR}^f)$ ,  $-\text{SO}_2\text{NR}^g\text{R}^f$ ,  $-\text{SC}(\text{Y})\text{NR}^g\text{R}^f$ ,  $-\text{C}(\text{Y})\text{R}^f$ ,  $-\text{C}(\text{Y})\text{OR}^f$ ,  $-\text{C}(\text{Y})\text{NR}^g\text{R}^f$ ,  $-\text{C}(\text{Y})\text{NR}^g\text{OR}^f$ , and  $-\text{C}(\text{O})\text{NR}^g\text{SO}_2\text{R}^f$ .

$\text{R}^f$  and each  $\text{R}^g$  are independently selected from H,  $\text{Cy}^6$ , and saturated or unsaturated ( $\text{C}_1\text{-C}_6$ )alkyl optionally substituted with one or more substituents  $\text{R}^h$  and/or one  $\text{Cy}^7$ ;

wherein  $\text{Cy}^6$  is optionally substituted with: one  $\text{Cy}^7$ , and/or one or more substituents  $\text{R}^h$ , and/or one or more saturated or unsaturated ( $\text{C}_1\text{-C}_6$ )alkyl groups optionally substituted with one or more substituents  $\text{R}^h$  and/or one  $\text{Cy}^7$ ; and wherein any  $\text{Cy}^7$  is optionally substituted with one or more substituents independently selected from  $\text{R}^h$  and ( $\text{C}_1\text{-C}_4$ )alkyl optionally substituted with one or more substituents  $\text{R}^h$ ;

each  $\text{R}^h$  is independently selected from halogen,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{OR}^i$ ,  $-\text{OC}(\text{O})\text{R}^i$ ,  $-\text{OC}(\text{O})\text{OR}^i$ ,  $-\text{OC}(\text{O})\text{NR}^i\text{R}^i$ ,  $-\text{NR}^i\text{R}^i$ ,  $-\text{NR}^i\text{C}(\text{O})\text{R}^i$ ,  $-\text{NR}^i\text{C}(\text{O})\text{OR}^i$ ,  $-\text{NR}^i\text{C}(\text{O})\text{NR}^i\text{R}^i$ ,  $-\text{NR}^i\text{S}(\text{O})_2\text{R}^i$ ,  $-\text{NR}^i\text{SO}_2\text{NR}^i\text{R}^i$ ,  $-\text{SR}^i$ ,  $-\text{S}(\text{O})\text{R}^i$ ,  $-\text{SO}_2\text{R}^i$ ,  $-\text{SO}_2(\text{OR}^i)$ ,  $-\text{SO}_2\text{NR}^i\text{R}^i$ ,  $-\text{C}(\text{O})\text{R}^i$ ,  $-\text{C}(\text{O})\text{OR}^i$ ,  $-\text{C}(\text{O})\text{NR}^i\text{R}^i$ , and  $-\text{C}(\text{O})\text{NR}^i\text{OR}^i$

each  $\text{R}^i$  is independently H or ( $\text{C}_1\text{-C}_4$ )alkyl optionally substituted with one or more halogen atoms;

Y is O, S, or  $\text{NR}^i$ ;

$\text{Cy}^1$ ,  $\text{Cy}^2$ ,  $\text{Cy}^4$  and  $\text{Cy}^6$  are independently a C or N-attached known ring system selected from 3- to 8-membered carbocyclic or heterocyclic monocyclic ring, saturated or partially unsaturated; phenyl; 5- or 6-membered heteroaromatic ring; and 6- to 18-membered carbocyclic or heterocyclic polycyclic ring system, saturated, partially unsaturated, aromatic or partially aromatic;

$\text{Cy}^3$ ,  $\text{Cy}^5$  and  $\text{Cy}^7$  are independently a C or N-attached known ring system selected from 3- to 8-membered carbocyclic or heterocyclic monocyclic ring, saturated or partially unsaturated; phenyl; and 5- or 6-membered heteroaromatic ring;

wherein in the carbocyclic rings all ring members are carbon atoms; and in the heterocyclic and heteroaromatic rings one or more ring members are selected from N, O, and S; and wherein in all saturated or partially unsaturated rings one or two members of the rings are optionally  $\text{C}(\text{O})$  and/or  $\text{C}(\text{NH})$  and/or  $\text{C}[\text{N}(\text{C}_1\text{-C}_4)\text{alkyl}]$ .

**[0013]** Another aspect of the invention relates to a pharmaceutical or veterinary composition which comprises an effective amount of a compound of formula (I) as defined above, or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt, together with one or more pharmaceutically or veterinary acceptable excipients or carriers.

**[0014]** As previously described, the compounds of the invention are useful as antifibrinolytic and antihemorrhagic agents. Therefore, another aspect of the invention relates to a compound of formula (I) as defined above, or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt, for use as a medicament.


**[0015]** Another aspect of the invention relates to a compound of formula (I) as defined above, or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt, for use as antifibrinolytic and antihemorrhagic agent. Thus, this aspect relates to the use of a compound of formula (I) as defined above, for the manufacture of a medicament for use as antifibrinolytic and antihemorrhagic agent; and may also be formulated as a method for the treatment and/or prevention of hyperfibrinolysis and/or hemorrhages comprising administering an effective amount of the previously defined compound of formula (I), or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt, and one or more pharmaceutically or veterinary acceptable excipients or carriers, in a subject in need thereof, including a human.

**[0016]** Processes for the preparation of compounds of formula (I) are also part of the present disclosure as well as intermediates used in these processes. Accordingly, another aspect of the invention relates to a compound of formula (III)

5

10

15



(III)

20

wherein A, B, C, R<sub>1</sub>-R<sub>7</sub> are as defined above, and R' is H or a carboxy protective group, more particularly a carboxy protective group selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, benzyl, p-methoxyphenyl, trimethylsilyl and [2-(Trimethylsilyl)ethoxy]methyl (SEM), with the proviso that compound (III) is other than 7-methoxycarbonyl-7-phenylsulphonyl-2-oxaspiro[2.4]-heptane and (2*S*\*,4*R*\*)-2-Phenylsulfonyl-4-iodomethyl-6,11-dioxaspiro[4.6]-undecane-2-carboxylic acid methyl ester.


**[0017]** Another aspect of the invention relates to a compound of formula (II)

25

30

35

40



(II)

wherein A, B, C, R<sub>1</sub>-R<sub>7</sub> are as defined above, and R is an hydroxamic acid protective group, more particularly an hydroxamic acid protective group selected from the group consisting of tetrahydro-2H-pyran-2-yloxy (THP), benzyl, 1-naphthylmethyl and dimethyloxybenzyl (DMB).

45

#### DETAILED DESCRIPTION OF THE INVENTION

**[0018]** All terms as used herein in this application, unless otherwise stated, shall be understood in their ordinary meaning as known in the art. Other more specific definitions for certain terms as used in the present application are as set forth below and are intended to apply uniformly through-out the specification and claims unless an otherwise expressly set out definition provides a broader definition.

50

**[0019]** For the purposes of the present invention, in the spirocyclic ring system formed by the ring system A and ring system B, the spiro atom connecting A and B is a carbon atom.

**[0020]** The term "carbocyclic" ring system refers to a known ring system wherein all the ring members are carbon atoms. The term "heterocyclic" ring system refers to a known ring system wherein one or more of the ring members, preferably 1, 2, 3, or 4 ring members, are selected from N, O, and S, where chemically possible. Unless otherwise specified, the "heterocyclic" ring system may be attached to the rest of the molecule through a C or a N atom. Both the carbocyclic and heterocyclic rings can be saturated or partially unsaturated, and may be unsubstituted or substituted as

55

described herein, being the substituents placed on any available position.

**[0021]** According to the present invention, the term "polycyclic" ring refers to a ring system which is formed by two, three or four rings which can be fused, bridged-fused, spiro-fused or can contain different types of fusion. For the purposes of the present invention, in "fused" rings the fusion occurs through one bond which is common to two adjoining rings; in "bridged-fused" rings the fusion occurs through a sequence of atoms (bridgehead) which is common to two rings; and in "spiro-fused" rings, the fusion occurs through only one atom (spiro atom), preferably a carbon atom, which is common to two adjoining rings (including bridged rings)

**[0022]** The polycyclic ring system can be saturated, partially unsaturated, aromatic (except in the case of ring system A) or partially aromatic; and may be unsubstituted or substituted as described herein, being the substituents placed on any available position.

**[0023]** The term "heteroaromatic" ring refers to a known aromatic ring system, wherein one or more of the ring members, preferably 1, 2, 3, or 4 ring members, are selected from N, O, and S where chemically possible. The heteroaromatic ring and phenyl may be unsubstituted or substituted as described herein, being the substituents placed on any available position.

**[0024]** The term "known" ring system as used herein refers to a ring system which is chemically feasible and is known in the art and so intends to exclude those ring systems that are not chemically possible.

**[0025]** For the purposes of the present invention, in all saturated or partially unsaturated rings, one or two members of the rings are optionally C(O) and/or C(NH) and/or C[N(C<sub>1</sub>-C<sub>4</sub>)alkyl].

**[0026]** The term saturated or unsaturated (C<sub>1</sub>-C<sub>n</sub>)alkyl refers to a saturated branched or linear hydrocarbon chain which contains from 1 to n carbon atoms. When the (C<sub>1</sub>-C<sub>n</sub>)alkyl is saturated it contains only single bonds. When the (C<sub>1</sub>-C<sub>n</sub>)alkyl is unsaturated it contains one or two double bonds and/or one or two triple bonds. The saturated or unsaturated (C<sub>1</sub>-C<sub>n</sub>)alkyl may be substituted or unsubstituted as described herein.

**[0027]** A halogen substituent means fluoro, chloro, bromo or iodo.

**[0028]** "Protective group" (PG) refers to a grouping of atoms that when attached to a reactive group in a molecule masks, reduces or prevents that reactivity.

**[0029]** The expression "substituted with one or more" means that a group can be substituted with one or more, preferably with 1, 2, 3 or 4 substituents, provided that this group has enough positions susceptible of being substituted.

**[0030]** For the purposes of the invention, room temperature is 20-25 °C.

**[0031]** As mentioned above, a first aspect of the invention relates to compounds of formula (I) or pharmaceutically or veterinary acceptable salts thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt. There is no limitation on the type of salt that can be used, provided that these are pharmaceutically or veterinary acceptable when they are used for therapeutic purposes. The term "pharmaceutically or veterinary acceptable salts", embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases.

**[0032]** The preparation of pharmaceutically or veterinary acceptable salts of the compounds of formula (I) can be carried out by methods known in the art. For instance, they can be prepared from the parent compound, which contains a basic or acidic moiety, by conventional chemical methods. Generally, such salts are, for example, prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate pharmaceutically or veterinary acceptable base or acid in water or in an organic solvent or in a mixture of them. The compounds of formula (I) and their salts may differ in some physical properties but they are equivalent for the purposes of the present invention.

**[0033]** The compounds of the invention may be in crystalline form either as free solvation compounds or as solvates (e.g. hydrates) and it is intended that both forms are within the scope of the present invention. Methods of solvation are generally known within the art. In general, the solvated forms with pharmaceutically or veterinary acceptable solvents such as water, ethanol and the like are equivalent to the unsolvated form for the purposes of the invention.

**[0034]** Some compounds of formula (I) can have chiral centres that can give rise to various stereoisomers. The present invention relates to each of these stereoisomers and also mixtures thereof. Moreover, some of the compounds of the present invention can show cis/trans isomers. The present invention relates to each of the geometric isomers and mixtures thereof.

**[0035]** Diastereoisomers can be separated by conventional techniques such as chromatography or fractional crystallization. Optical isomers can be resolved by conventional techniques of optical resolution to give optically pure isomers. This resolution can be carried out on any chiral synthetic intermediate or on products of general formula (I). Optically pure isomers can also be individually obtained using enantiospecific synthesis.

**[0036]** In one embodiment, the invention refers to a compound of formula (I) as defined above, wherein A is a known 3- to 8-membered carbocyclic or heterocyclic monocyclic ring or a known 6- to 10-membered carbocyclic or heterocyclic bicyclic ring system.

**[0037]** In another embodiment, the invention refers to a compound of formula (I), wherein A is a monocyclic ring selected from a 3- to 6-membered carbocyclic ring, and a 5- to 6-membered heterocyclic ring.

**[0038]** In another embodiment, the invention refers to a compound of formula (I), wherein A is a carbocyclic monocyclic

ring; or a polycyclic ring system, preferably a bicyclic ring system, wherein the ring containing the spiro atom attached to the hydroxamic acid and the sulfonyl group is a carbocyclic ring.

[0039] In another embodiment, the invention refers to a compound of formula (I), wherein A is selected from cyclopropane, cyclobutane, cyclopentane, cyclohexane, tetrahydrofuran, pyrrolidine, bicyclo[2.2.1]heptane, 2,3-dihydro-1H-indene, hexahydropyrolizin-3-one, and 4-azaspiro[4.4]nonane.

[0040] In another embodiment, the invention refers to a compound of formula (I), wherein A is unsubstituted, i.e.  $R_7$  is H.


[0041] In another embodiment, the invention refers to a compound of formula (I), wherein B is a 6- to 7-membered carbocyclic or heterocyclic monocyclic ring.


[0042] In another embodiment, the invention refers to a compound of formula (I), wherein B is a saturated monocyclic ring, carbocyclic or heterocyclic, wherein at least one of the ring members of the heterocyclic ring is  $NR_4$ .

[0043] In another embodiment, the invention refers to a compound of formula (I), wherein B is selected from cyclohexane, piperidine, morpholine, azepane, piperazine, pyrrolidine, and azetidine.

[0044] In another embodiment, B is piperidine, morpholine, azepane, pyrrolidine, and azetidine, wherein  $R_4$  is placed on the N atom of these rings and  $R_5$ - $R_6$  are H. In another embodiment, B is piperazine, wherein  $R_4$  and  $R_5$  are placed on the N atoms and  $R_6$  is H.

[0045] In another embodiment, the invention refers to a compound of formula (I), wherein A and B form a spirocyclic ring system selected from the group consisting of:





[0046] In another embodiment, the invention refers to a compound of formula (I), wherein C is phenyl. In another embodiment, C is phenyl substituted with R<sup>1</sup> at the ortho, meta or para position, and R<sub>2</sub> and R<sub>3</sub> are independently selected from H, halogen, R<sup>a</sup>, -OR<sup>a</sup>, and -NR<sup>b</sup>R<sup>a</sup>; wherein R<sup>a</sup>, R<sup>a</sup> and R<sup>b</sup> are independently selected from H and -(C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or more halogen atoms. In another embodiment, C is phenyl substituted with R<sub>1</sub> at the para position and R<sub>2</sub> and R<sub>3</sub> are H. In another embodiment, C is phenyl substituted with R<sup>1</sup> at the ortho position and R<sub>2</sub> and R<sub>3</sub> are H. In another embodiment, C is phenyl substituted with R<sup>1</sup> at the meta position and R<sub>2</sub> and R<sub>3</sub> are H. In another embodiment, C is phenyl substituted with R<sup>1</sup> at the meta position, and R<sub>2</sub> at the para position, R<sub>3</sub> being H; or alternatively, C is phenyl substituted with R<sup>1</sup> at the para position, and R<sub>2</sub> at the meta position, R<sub>3</sub> being H; wherein R<sub>2</sub> is selected from H, halogen, R<sup>a</sup>, -OR<sup>a</sup>, and -NR<sup>b</sup>R<sup>a</sup>; and R<sup>a</sup>, R<sup>a</sup> and R<sup>b</sup> are independently selected from H and -(C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or more halogen atoms.

[0047] In another embodiment, the invention refers to a compound of formula (I), wherein in R<sup>b</sup> (relating to R<sub>1</sub>-R<sub>3</sub>), Cy<sup>1</sup> and Cy<sup>2</sup> are independently optionally substituted with one or more substituents selected from R<sup>e</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>e</sup>; and Cy<sup>6</sup> is optionally substituted with one or more substituents independently selected from R<sup>h</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted as previously defined, more particularly with one or more substituents R<sup>h</sup>.

[0048] In a more particular embodiment, the invention refers to a compound of formula (I), wherein in R<sup>b</sup> (relating to R<sub>1</sub>-R<sub>3</sub>), R<sup>b</sup> is H and saturated or unsaturated (C<sub>1</sub>-C<sub>12</sub>)alkyl optionally substituted with one or more substituents R<sup>e</sup>, more particularly wherein in R<sup>e</sup>, R<sup>f</sup> and each R<sup>g</sup> are independently selected from H and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more fluorine atoms.

[0049] In another embodiment, the invention refers to a compound of formula (I), wherein in R<sub>1</sub>-R<sub>3</sub>, R<sup>e</sup> is selected from halogen, -NO<sub>2</sub>, -CN, -OR<sup>f</sup>, -OC(O)R<sup>f</sup>, -OC(O)OR<sup>f</sup>, -OC(O)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>C(O)R<sup>f</sup>, -NR<sup>g</sup>C(O)OR<sup>f</sup>, -NR<sup>g</sup>C(O)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>S(O)<sub>2</sub>R<sup>f</sup>, -SR<sup>f</sup>, -S(O)R<sup>f</sup>, -SO<sub>2</sub>R<sup>f</sup>, -SO<sub>2</sub>NR<sup>g</sup>R<sup>f</sup>, -C(O)R<sup>f</sup>, -C(O)OR<sup>f</sup>, -C(O)NR<sup>g</sup>R<sup>f</sup>, and -C(O)NR<sup>g</sup>OR<sup>f</sup>.

[0050] In another embodiment, the invention refers to a compound of formula (I), wherein in R<sub>1</sub>-R<sub>3</sub>, R<sup>f</sup> and each R<sup>g</sup> are independently selected from H and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more fluorine atoms.

[0051] In another embodiment, the invention refers to a compound of formula (I), wherein in R<sub>1</sub>-R<sub>3</sub>, Cy<sup>1</sup> and Cy<sup>2</sup> are independently optionally substituted with one or more substituents selected from R<sup>e</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted as previously defined; and Cy<sup>6</sup> is optionally substituted with one or more substituents independently selected from R<sup>h</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted as previously defined.

[0052] In another embodiment, the invention refers to a compound of formula (I), wherein R<sub>1</sub>-R<sub>3</sub> are independently selected from H, halogen, -NO<sub>2</sub>, -CN, R<sup>a</sup>, -OR<sup>a</sup>, -OC(O)R<sup>a</sup>, -OC(O)OR<sup>a</sup>, -OC(O)NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>C(O)R<sup>a</sup>, -NR<sup>b</sup>C(O)OR<sup>a</sup>, -NR<sup>b</sup>C(O)NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>S(O)<sub>2</sub>R<sup>a</sup>, -SR<sup>a</sup>, -S(O)R<sup>a</sup>, -SO<sub>2</sub>R<sup>a</sup>, -SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -C(O)R<sup>a</sup>, -C(O)OR<sup>a</sup>, -C(O)NR<sup>b</sup>R<sup>a</sup>, and -C(O)NR<sup>b</sup>OR<sup>a</sup>.

[0053] In another embodiment, the invention refers to a compound of formula (I), wherein in R<sub>4</sub>-R<sub>7</sub>, R<sup>h</sup> is selected from halogen, -NO<sub>2</sub>, -CN, -OR<sup>i</sup>, -OC(O)R<sup>i</sup>, -OC(O)OR<sup>i</sup>, -OC(O)NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>C(O)R<sup>i</sup>, -NR<sup>i</sup>C(O)OR<sup>i</sup>, -NR<sup>i</sup>C(O)NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>S(O)<sub>2</sub>R<sup>i</sup>, -SR<sup>i</sup>, -S(O)R<sup>i</sup>, -SO<sub>2</sub>R<sup>i</sup>, -SO<sub>2</sub>NR<sup>i</sup>R<sup>i</sup>, -C(O)R<sup>i</sup>, -C(O)OR<sup>i</sup>, and -C(O)NR<sup>i</sup>R<sup>i</sup>.

[0054] In another embodiment, the invention refers to a compound of formula (I), wherein R<sub>4</sub>-R<sub>7</sub> are independently selected from halogen, -NO<sub>2</sub>, -CN, R<sup>c</sup>, -OR<sup>c</sup>, -NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>C(O)R<sup>c</sup>, -NR<sup>d</sup>C(O)OR<sup>c</sup>, -NR<sup>d</sup>C(O)NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>S(O)<sub>2</sub>R<sup>c</sup>, -SR<sup>c</sup>, -S(O)R<sup>c</sup>, -SO<sub>2</sub>R<sup>c</sup>, -SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>, -C(O)R<sup>c</sup>, -C(O)OR<sup>c</sup>, and -C(O)NR<sup>d</sup>R<sup>c</sup>.


[0055] In another embodiment, the invention refers to a compound of formula (I), wherein R<sub>2</sub> and R<sub>3</sub> are independently selected from H, halogen, R<sup>a</sup>, -OR<sup>a</sup>, and -NR<sup>b</sup>R<sup>a</sup>; and R<sub>5</sub>-R<sub>7</sub> are independently selected from H, halogen, R<sup>c</sup>, -OR<sup>c</sup>, and -NR<sup>d</sup>R<sup>c</sup>, wherein R<sup>a</sup>, R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup> and R<sup>d</sup> are independently selected from H and -(C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with

one or more fluorine atoms.


**[0056]** In another embodiment, the invention refers to a compound of formula (I), wherein  $R_1$  is selected from H, halogen,  $-NO_2$ ,  $-CN$ ,  $R^a$ ,  $-OR^a$ ,  $-OR^a$ ,  $-OC(O)R^a$ ,  $-OC(O)OR^a$ ,  $-OC(O)NR^bR^a$ ,  $-NR^bR^a$ ,  $-NR^bC(O)R^a$ ,  $-NR^bC(O)OR^a$ ,  $-NR^bC(O)NR^bR^a$ ,  $-NR^bS(O)_2R^a$ ,  $-SR^a$ ,  $-S(O)R^a$ ,  $-SO_2R^a$ ,  $-SO_2NR^bR^a$ ,  $-C(O)R^a$ ,  $-C(O)OR^a$ ,  $-C(O)NR^bR^a$ , and  $-C(O)NR^bOR^a$ ;  $R_4$  is selected from halogen,  $-NO_2$ ,  $-CN$ ,  $R^c$ ,  $-OR^c$ ,  $-NR^dR^c$ ,  $-NR^dC(O)R^c$ ,  $-NR^dC(O)OR^c$ ,  $-NR^dC(O)NR^dR^c$ ,  $-NR^dS(O)_2R^c$ ,  $-SR^c$ ,  $-S(O)R^c$ ,  $-SO_2R^c$ ,  $-SO_2NR^dR^c$ ,  $-C(O)R^c$ ,  $-C(O)OR^c$ , and  $-C(O)NR^dR^c$ ; and  $R_2$ ,  $R_3$ , and  $R_5$ - $R_7$  are independently selected from H, halogen,  $-(C_1-C_4)alkyl$ ,  $-OH$ ,  $-[(C_1-C_4)alkyl]$ ,  $-NH_2$ ,  $-NH[(C_1-C_4)alkyl]$ ,  $-N[(C_1-C_4)alkyl]_2$ , wherein each  $(C_1-C_4)alkyl$  is independently optionally substituted with one or more fluorine atoms.

**[0057]** The present invention also relates to the combination of any of the specific embodiments defined above for any of the variables A, B, C, and  $R_1$ - $R_7$ .


**[0058]** In another embodiment of the invention, the compound of formula (I) is selected from the group consisting of:




1-01




1-02




1-03




1-04




1-05




1-06



1-07




1-08




1-09


5



**1-10**




**1-11**




**1-12**


15



**1-13**




**1-14**




**1-15**


25



**1-16**




**1-17**




**1-18**


40



**1-19**




**1-20**




**1-21**

55


5



**1-22**




**1-23**




**1-24**

10


15



**1-25**




**1-26**




**1-27**

20


25



**1-28**




**1-29**




**1-30**

30


35



**1-31**



**1-32**



**1-33**


40

45


50

55


5



**1-34**




**1-35**




**1-36**


15



**1-37**



**1-38**



**1-39**


20

25


30

35


40



**1-40**




**1-41**




**1-42**

45


50



**1-43**




**1-44**




**1-45**

55


5



**1-46**




**1-47**




**1-48**


15



**1-49**



**1-50**



**1-51**


20

25


30

35


40



**1-52**



**1-53**




**1-54**


45

50


55



**1-55**




**1-56**




**1-57**


5



**1-58**




**1-59**




**1-60**


10



**1-61**



**1-62**




**1-63**


15

20


25



**1-64**



**1-65**




**1-66**


30

35


40



**1-67**




**1-68**




**1-69**

45


50



**1-70**






**1-71**






**1-72**

55




5






10






20



30




40




55

EP 2 877 446 B9


5



**1-88**




**1-89**




**1-90**


10



**1-91**




**1-92**




**1-93**

15


20



**1-94**




**1-95**




**1-96**

25


30



**1-97**




**1-98**




**1-99**

35


40



**1-100**



**1-101**




**1-102**


50

55


5



**1-103**




**1-104**




**1-105**

10


15



**1-106**




**1-107**




**1-108**

20


25



**1-109**




**1-110**




**1-111**

30


35



**1-112**



**1-113**



**1-114**


40

45


50

55


5



**1-115**




**1-116**




**1-117**


10



**1-118**



**1-119**




**1-120**


15

20


25



**1-121**




**1-122**




**1-123**

30


35



**1-124**




**1-125**




**1-126**

40


45



**1-127**




**1-128**




**1-129**

50


55




**1-130**



**1-131**




**1-132**



[0059] \*Regarding these compounds, no attempt has been made to identify the absolute configurations. However, in the examples it is clearly indicated which of the two isomers is concerned in relative terms by differentiating unambiguously between their physical and/or spectroscopic properties.


[0060] Generally, compounds of formula (I) as defined above may be obtained by reacting a compound of formula (III)



with a hydroxylamine of formula RO-NH<sub>2</sub> (IV), wherein A, B, C, R<sub>1</sub>-R<sub>7</sub> are as previously defined; R' is H; and R is an hydroxamic acid protective group, more particularly an hydroxamic acid protective group selected from the group consisting of tetrahydro-2H-pyran-2-yloxy (THP), benzyl, 1-naphthylmethyl and dimethyloxybenzyl (DMB), to give a compound of formula (II)

50

55



and subsequently removing the protective group of the hydroxamic acid to give a compound of formula (I).

**[0061]** The first conversion can be carried out in the presence of an activating agent such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC.HCl) and Hydroxybenzotriazole (HOBT), preferably in the presence of a base, such as N-methylmorpholine (NMM), in a suitable solvent, such as dichloromethane, chloroform or dimethylformamide, at a temperature comprised from room temperature to the temperature of the boiling point of the solvent, preferably at room temperature.

**[0062]** The removal of the protective group of the hydroxamic acid is carried out by standard methods well-known in the art as described for example in T. W. Green and P. G. M. Wuts, *Protective Groups in Organic Chemistry* (Wiley, 3rd ed. 1999, Chapter 2, pp. 17-200). Representative hydroxy protective groups include those where the hydroxy group is either acylated or alkylated such as benzyl, and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers and allyl ethers. For example, the hydroxamic acid protective group is tetrahydro-2H-pyran-2-yloxy (THP), benzyl, 1-naphthylmethyl or dimethoxybenzyl (DMB). When the hydroxamic acid protective group is THP, the deprotection is carried out in acidic medium, for example with HCl, in a suitable solvent such as dioxane.


**[0063]** Intermediate compounds of formula (II), wherein A, B, C, R<sub>1</sub>-R<sub>7</sub> and R are as previously defined, also form part of the invention.

**[0064]** Compounds of formula (III) wherein R' is H can be obtained by removing the protective group of a compound of formula (III) wherein R' is a carboxy protective group by standard methods well-known in the art as described for example in T. W. Green and P. G. M. Wuts, *Protective Groups in Organic Chemistry* (Wiley, 3rd ed. 1999, Chapter 5, pp. 369-451). Representative carboxy protective groups include alkyl, aryl or benzyl esters, silyl esters, amides or hydrazides. For example, the carboxy protective group is (C<sub>1</sub>-C<sub>6</sub>)alkyl, benzyl, p-methoxyphenyl, trimethylsilyl, or [2-(Trimethylsilyl)ethoxy]methyl (SEM). When the carboxy protective group is (C<sub>1</sub>-C<sub>6</sub>)alkyl, the deprotection is carried out in basic medium, for example with LiOH in a suitable solvent such as tetrahydrofuran-methanol.

**[0065]** Intermediate compounds of formula (III) wherein A, B, C, R<sub>1</sub>-R<sub>7</sub> are as previously defined and R' is H or a carboxy protective group, more particularly a carboxy protective group selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, benzyl, p-methoxyphenyl, trimethylsilyl and [2-(Trimethylsilyl)ethoxy]methyl (SEM) form also part of the invention, with the proviso that compound (III) is other than 7-methoxycarbonyl-7-phenylsulphonyl-2-oxaspiro[2.4]heptane and (2*S*\*,4*R*\*)-2-Phenylsulfonyl-4-iodomethyl-6,11-dioxaspiro[4.6]undecane-2-carboxylic acid methyl ester, form also part of the invention. These compounds are cited in D. Bouyssi et al., "Rearrangement of oxaspiroheptanes to cyclohexanones mediated by lithium iodide", *Synlett* 2000, vol. 5, pp. 749-751 and Osamu Kitagawa et al., "Stereoselective Iodine Atom Transfer [3 + 2] Cycloaddition Reaction with Alkenes Using Unsymmetrical Allylated Active Methine Radicals", *The Journal of Organic Chemistry* 2004, vol. 69, pp. 2607-2610, respectively, which describe only a synthetic process without describing any therapeutical application.


Given that the chemical structure of intermediate compounds of formula (II) and formula (III) coincides largely with the chemical structure of the final compounds of formula (I), the particular embodiments defined above for the compounds of formula (I) regarding the variables A, B, C, and R<sub>1</sub>-R<sub>7</sub> are also particular embodiments of the intermediate compounds of formula (II) and formula (III).

**[0066]** Compounds of formula (III) can be generally obtained by coupling a compound of formula (V) with a compound of formula (VI):



15 wherein A, B, C, R<sub>1</sub>-R<sub>7</sub> are as previously defined and R' is carboxy protective group. This conversion is carried out in the presence of a base such as lithium diisopropylamide (LDA), in a suitable solvent such as tetrahydrofuran, and at a suitable temperature, preferably -78 °C.

[0067] Additionally, compounds of formula (I), wherein R<sup>1</sup> is Q-Cy<sup>2</sup>, wherein Q is O, S, SO or SO<sub>2</sub> and Cy<sup>2</sup> is as defined above (i.e. compound of formula (Ia)) may be obtained in general as shown in the following scheme:



Scheme 1


45 [0068] A compound of formula (V), wherein A, B, R<sub>4</sub>-R<sub>7</sub> are as previously defined and R' is carboxy protective group, is reacted as described above with a compound of formula (Via) wherein C, R<sub>2</sub> and R<sub>3</sub> are as previously defined, to give a compound of formula (IIIa), which is deprotected to give the carboxylic acid of formula (IIIb) and subsequently reacted with a compound of formula (IV) as already described to give compound (IIa).

50 [0069] Compound of formula (IIa) is reacted with a compound of formula H-Q'-Cy<sup>2</sup> (VII), wherein Q' is O or S, and Cy<sup>2</sup> is as defined above, to give a compound of formula (IIb). This reaction is carried out in the presence of a base, such as cesium carbonate, optionally in a suitable solvent or without solvent, and preferably heating.

[0070] Further, a compound of formula (IIb) wherein Q is S can be oxidized to a compound of formula (IIb) wherein Q is SO or SO<sub>2</sub> in the presence of an oxidizing agent, such as m-chloroperbenzoic acid, in a suitable solvent, such as dichloromethane, and preferably at room temperature.

[0071] Deprotection of a compound of formula (IIb) yields a compound of formula (Ia) as defined above.


55 [0072] Additionally, compounds of formula (I), wherein R<sup>1</sup> is Q-alk, Q is O or S, and alk is saturated or unsaturated -(C<sub>1</sub>-C<sub>12</sub>)alkyl optionally substituted (i.e. compound of formula (Ib)) may be obtained in general as shown in the following scheme:



Scheme 2

15 **[0073]** Thus, a compound of formula (IIa) can be reacted with a compound of formula H-Q-alk (VIII) in the presence of a base, such as NaH, optionally in a suitable solvent or without solvent, and preferably heating. After removal of the protective group in the resulting compound of formula (IIc), compound of formula (Ib) is obtained.


20 **[0074]** Compounds of formula (I), wherein R<sup>1</sup> is -NR<sup>b</sup>R<sup>a'</sup>, R<sup>b</sup> is saturated or unsaturated -(C<sub>1</sub>-C<sub>12</sub>)alkyl optionally substituted and R<sup>a'</sup> is H or saturated or unsaturated -(C<sub>1</sub>-C<sub>12</sub>)alkyl optionally substituted (i.e. compound of formula (Ic)) may be obtained in general as shown in the following scheme:



Scheme 3

35 **[0075]** Thus, a compound of formula (IIa) can be reacted with a compound of formula H-NR<sup>a</sup>R<sup>b'</sup> (IX), optionally in a suitable solvent or without solvent, and preferably heating. After removal of the protective group in the resulting compound of formula (IIc), compound of formula (Ic) is obtained.


**[0076]** Compounds of formula (I), wherein R<sup>1</sup> is -NCy<sup>2</sup>R<sup>a'</sup> and Cy<sup>2</sup> is an optionally substituted aromatic, heteroaromatic or aliphatic ring (i.e. compound of formula (Id)) may be obtained in general as shown in the following scheme:



Scheme 4

50 **[0077]** Thus, a compound of formula (IIa) can be reacted with a compound of formula H-NCy<sup>2</sup>R<sup>a'</sup> (X), optionally in a suitable solvent or without solvent, and preferably heating. After removal of the protective group in the resulting compound of formula (IIe), compound of formula (Id) is obtained.


55 **[0078]** Compounds of formula (I), wherein R<sup>1</sup> is -NR<sup>b</sup>C(Y)R<sup>a'</sup>, -NR<sup>b</sup>C(Y)OR<sup>a'</sup>, -NR<sup>b</sup>C(Y)NR<sup>b</sup>R<sup>a'</sup>, -NR<sup>b</sup>S(O)<sub>2</sub>R<sup>a'</sup>, -NR<sup>b</sup>SO<sub>2</sub>NR<sup>b</sup>R<sup>a'</sup> (i.e. compound of formula (Ie)) may be obtained from a compound of formula (IIg), which can be prepared as shown in the following scheme:



Scheme 5

[0079] In a first step, compound of formula (IIa) is reacted with benzylamine, preferably heating to give a compound of formula (IIf) which can be hydrogenated in the presence of Pd/C to give a compound of formula (IIg). This compound can be reacted for example with an acyl halide in the presence of a base, such as triethylamine, in a suitable solvent, such as tetrahydrofuran, at a temperature comprised from room temperature to the reflux temperature of the solvent, to give an amide; or with an isocyanate to give a urea.

[0080] Compounds of formula (I), wherein R<sup>1</sup> is -OC(Y)R<sup>a</sup> (i.e. compound of formula (If)) may be obtained from a compound of formula (III), which can be prepared as shown in the following scheme:




Scheme 6

[0081] In a first step, compound of formula (IIa) is reacted with benzyl alcohol in the presence of a base, such as NaH, in a suitable solvent, such as tetrahydrofuran, to give a compound of formula (IIh) which can be hydrogenated in the presence of Pd/C to give a compound of formula (III).

[0082] This compound can be reacted for example with an acyl halide in the presence of a base, such as triethylamine, in a suitable solvent, preferably at room temperature to give an ester.

[0083] Compounds of formula (I), wherein R<sup>1</sup> is R<sup>a</sup> (i.e. compound of formula (Ilg)) may be obtained from a compound of formula (III), which can be prepared as shown in the following scheme:



Scheme 7

[0084] In a first step, compound of formula (IIg) is reacted with potassium iodide and NaNO<sub>2</sub> in a suitable solvent, such as acetonitrile and then an acid, such as concentrated HCl, is added to give a compound of formula (IIIj). This compound can be reacted for example with boronic acid derivatives in the presence of a palladium catalyst (Suzuki coupling) to give a compound of formula (IIIk).

[0085] The same type of reactions shown above for R<sup>1</sup> can be applied to R<sub>2</sub>-R<sub>7</sub>.

[0086] Alternatively, the reactions described above can be carried out in a different order. Thus, the reactions carried

out on compounds of formula (II) (i.e. on the protected hydroxamic acid) can be carried out on compounds of formula (III) (i.e. the carboxylic ester), and can be deprotected to give the corresponding carboxylic acids, which can be converted to the compounds of formula (I) as described above.

5 [0087] The compounds of formulas (V) to (XI) are commercially available or can be obtained by conventional synthetic processes.

[0088] The present invention also relates to a pharmaceutical or veterinary composition comprising an effective amount of a compound of formula (I) as defined above, or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt, together with pharmaceutically or veterinary acceptable excipients or carriers.

10 [0089] The expression "effective amount" as used herein, refers to the amount of a compound that, when administered, is sufficient to prevent development of, or alleviate to some extent, one or more of the symptoms of the disease which is addressed. The specific dose of the compound of the invention to obtain a therapeutic benefit may vary depending on the particular circumstances of the individual patient including, among others, the size, weight, age and sex of the patient, the nature and stage of the disease, the aggressiveness of the disease, and the route of administration. For example, a dose of from about 0.01 to about 300 mg/kg may be used.

15 [0090] The expression "pharmaceutically or veterinary acceptable excipients or carriers" refers to pharmaceutically or veterinary acceptable materials, compositions or vehicles. Each component must be pharmaceutically or veterinary acceptable in the sense of being compatible with the other ingredients of the pharmaceutical or veterinary composition. It must also be suitable for use in contact with the tissue or organ of humans and animals without excessive toxicity, irritation, allergic response, immunogenicity or other problems or complications commensurate with a reasonable benefit/risk ratio.

[0091] The election of the pharmaceutical or veterinary formulation will depend upon the nature of the active compound and its route of administration. Any route of administration may be used, for example oral, parenteral and topical administration.

25 [0092] For example, the pharmaceutical or veterinary composition may be formulated for oral administration and may contain one or more physiologically compatible carriers or excipients, in solid or liquid form. These preparations may contain conventional ingredients such as binding agents, fillers, lubricants, and acceptable wetting agents.

[0093] The pharmaceutical or veterinary composition may be formulated for parenteral administration in combination with conventional injectable liquid carriers, such as water or suitable alcohols. Conventional pharmaceutical or veterinary excipients for injection, such as stabilizing agents, solubilizing agents, and buffers, may be included in such compositions. These pharmaceutical or veterinary compositions may be injected intramuscularly, intraperitoneally, or intravenously.

30 [0094] The pharmaceutical or veterinary composition may be formulated for topical administration. Formulations include creams, lotions, gels, powders, solutions and patches wherein the compound is dispersed or dissolved in suitable excipients. The topical compositions of the invention may be administered by means of a carrier material, which can be a solid support. Thus, it also forms part of the invention a topical composition comprising a carrier material, which can be a solid support. Illustrative, non-limiting examples of solid supports include intelligent textiles, dressings, coatings, sponges, band-aids, sanitary pads, compresses, plasters, etc. The manufacture of such compositions can be obtained by conventional methods, for example, by mixing the combinations of the invention and the material carrier.

35 [0095] The pharmaceutical or veterinary compositions may be in any form, including, among others, tablets, pellets, capsules, aqueous or oily solutions, suspensions, emulsions, or dry powdered forms suitable for reconstitution with water or other suitable liquid medium before use, for immediate or retarded release.

40 [0096] The appropriate excipients and/or carriers, and their amounts, can readily be determined by those skilled in the art according to the type of formulation being prepared.

[0097] The compounds of the present invention are useful as antihemorrhagic and antifibrinolytic agents and can be used in a broad range of therapeutic applications. In surgery, antifibrinolytic agents, in addition to reducing postoperative hemorrhage, can be an alternative to blood transfusion and other hemoderivatives for example in heart, liver and orthopedic surgery, and also in the setting of oncologic surgery in organs rich in fibrinolysis activators (prostate, uterus). In trauma patients antifibrinolytic agents can reduce all-cause mortality and death due to bleeding. Further, the antifibrinolytic agents of the invention can also be used to control bleeding in trombotic therapy, e.g. in cases of acute heart attack and ischemic stroke, and major or intracranial hemorrhages. Moreover, the antifibrinolytic agents of the invention are useful in the treatment of local hemorrhages, e.g. after teeth extraction, in particular in patients with congenital coagulopathies, such as hemophilia, or patients with diabetes; in the treatment of menorrhage in women associated with congenital or acquired coagulopathies, as well as in post-partum haemorrhage, and in the treatment of hemorrhages of gastrointestinal and urologic origin, including prostatectomy.

55 [0098] Throughout the description and claims the word "comprise" and variations thereof, are not intended to exclude other technical features, additives, components, or steps. Furthermore, the word "comprise" encompasses the case of "consisting of". Additional objects, advantages and features of the invention will become apparent to those skilled in the art upon examination of the description or may be learned by practice of the invention. The following examples are

provided by way of illustration, and they are not intended to be limiting of the present invention. Furthermore, the present invention covers all possible combinations of particular and preferred embodiments described herein.

EXAMPLES

5

General Procedure for Prep-HPLC purification method:

10

**[0099]** The HPLC measurement was performed using Gilson 281 from 233 pump (binary), an autosampler, and a UV detector. The fractions were detected by LC-MS. The MS detector was configured with an electrospray ionization source. The source temperature was maintained at 300-350 °C.

Method 1

15

**[0100]** Reverse phase HPLC was carried out on Luna C18 (100x30 mm; 4µm). Solvent A: water with 0.075% trifluoroacetic acid; Solvent B: acetonitrile with 0.075% trifluoroacetic acid. Gradient: At room temperature, 20% of B to 40% of B within 6 min at 25 mL/min; then 40% B at 25 mL/min over 2 min, UV detector.

**[0101]** The following abbreviations have been used in the examples:

20

Boc: tert-butoxycarbonyl

DCM: dichloromethane

DMF: dimethylformamide

Et<sub>3</sub>N: triethylamine

TLC: Thin Layer Chromatography

PE: Petroleum ether

25

AE/EtOAc: ethyl acetate

ACN: acetonitrile

r.t.: room temperature

Rt: retention time

THF: tetrahydrofuran

30

LDA Lithium diisopropylamide

EDC.HCl: 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride

HOBt: Hydroxybenzotriazole

THP-O-NH<sub>2</sub>: N-(tetrahydro-2H-pyran-2-yloxy)amine

MeOH: methanol

35

NMM: N-Methylmorpholine

m-CPBA: m-chloroperbenzoic acid

DCM: dichloromethane

DMSO: dimethylsulfoxide

40


Synthetic route 1a

**[0102]**

45

50

55



Conditions: a. K<sub>2</sub>CO<sub>3</sub> (2 eq), CH<sub>3</sub>I (1.5 eq) in DMF, r.t.; b. LDA (1.25 M) in THF, 1 hour at -78 °C; then, R-02 (2 eq), 1 hour at -78 °C; c. LiOH.H<sub>2</sub>O (10 eq) in THF/MeOH/H<sub>2</sub>O (3/3/2), r.t.; d. EDC.HCl (2 eq), HOBT (2 eq), THP-O-NH<sub>2</sub> (1.5 eq), NMM (3 eq) in DMF, r.t.; e. R-04 (1.5 - 3 eq), Cs<sub>2</sub>CO<sub>3</sub> (2 - 3 eq), 12 hours at 90 °C; e'. m-CPBA (4 eq) in DCM, r.t. (Q=S to Q'=SO<sub>2</sub>); f.- HCl / Dioxane, r.t. for 1 hour.

**[0103]** In the scheme above Q is O or S, Q' is Q or SO<sub>2</sub> and Cy is phenyl or a 5- to 6-membered heteroaryl and can be optionally substituted.

#### Preparation of intermediate I-01 a: 8-tert-butoxycarbonyl-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester

**[0104]** To a solution of 8-tert-butoxycarbonyl-8-azaspiro[4.5]decane-3-carboxylic acid (20 g, 0.071 mol), commercially available from Wuxi Apptec, in DMF (200mL) was added K<sub>2</sub>CO<sub>3</sub> (18.59 g, 2 eq), then compound CH<sub>3</sub>I (14.34 g, 1.5 eq) was added dropwise. The reaction mixture was stirred at room temperature for 2 h. After TLC (PE/AE 5:1) showed the starting material was consumed, the mixture was quenched with water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude compound 8-tert-butoxycarbonyl-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester (20.38 g, 97.09%) as a pale yellow oil which was used for the next step without further purification. ESI-MS (M+1): 298 calc. for C<sub>16</sub>H<sub>27</sub>NO<sub>4</sub>.

#### Preparation of reagent R-02a: 4-Fluorobenzenesulfonyl fluoride

**[0105]** To a solution of compound 4-Fluorobenzenesulfonyl chloride (50 g, 0.256 mol) in ACN (500 mL) was added KF (74.36 g, 5 eq) and 18-crown-6 (2 g) at r.t., then the mixture was stirred at r.t. overnight. The mixture was detected by LC-MS, then saturated aqueous NaHCO<sub>3</sub> was added and the mixture was extracted with EtOAc, the organic layer was washed with saturated aqueous NaHCO<sub>3</sub>, brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude compound 4-Fluorobenzenesulfonyl fluoride (46.70 g) as pale yellow oil which was used for the next step without further purification. ESI-MS (M+1): 179 calc. for C<sub>6</sub>H<sub>4</sub>F<sub>2</sub>O<sub>2</sub>S.

## EP 2 877 446 B9

### Preparation of intermediate I-02a: 8-tert-butoxycarbonyl-3-(4-fluorophenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester

5 [0106] To a solution of compound I-01a (18.38 g, 0.062 mol) in THF (200mL) was added LDA (102 mL, 1.25 M) at -78°C. After stirring at -78 °C for 1 h, the compound 4-Fluorobenzenesulfonyl fluoride (22.23 g, 2 eq) was added to the solution, the reaction was stirred at -78 °C for 1 hour, and then the mixture was stirred at r.t. overnight. After TLC (PE/AE 5:1) showed the starting material was consumed, the mixture was quenched with aqueous NH<sub>4</sub>Cl and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product which was purified by column chromatography (eluting with EA/PE= 100:1 to 10: 1) to give pure compound I-02a (18.94 g, 67.03%) as a pale yellow oil. ESI-MS (M+1): 456 calc. for C<sub>22</sub>H<sub>30</sub>FNO<sub>6</sub>S.

### Preparation of intermediate I-03a: 8-tert-butoxycarbonyl-3-(4-fluorophenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxylic acid

15 [0107] To a solution of compound I-02a (10 g, 0.022 mol) in THF/MeOH/H<sub>2</sub>O (3/3/2, 80mL) was added LiOH·H<sub>2</sub>O (9.23 g, 10 eq). The resulting mixture was stirred at r.t. for 4 h, after TLC (PE/AE 5:1) showed the starting materials were consumed completely, then the mixture was diluted with water and adjusted pH to 2-3 with 1 N HCl and the mixture was extracted with EtOAc, washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product I-03a (9.60 g, 98.97%) as a pale yellow oil which was used in the next step. ESI-MS (M+1): 442 calc. for C<sub>21</sub>H<sub>28</sub>FNO<sub>6</sub>S.

### Preparation of intermediate I-04a: 8-tert-butoxycarbonyl-3-(4-fluorophenylsulfonyl)-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

25 [0108] To a solution of compound I-03a (9.60 g, 0.022 mol) in DMF (100 mL) was added EDC.HCl (8.40 g, 2 eq), HOBT (5.85 g, 2 eq), THP-O-NH<sub>2</sub> (3.86 g, 1.5 eq), NMM (6.67 g, 3 eq) at r.t., then the mixture was stirred at room temperature overnight. The mixture was quenched with aqueous water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product which was purified by column chromatography (eluting with EA/PE = 50:1 to 2:1) to give pure compound I-04a (9.0 g, 75.76%) as a pale yellow solid. ESI-MS (M+1): 541 calc. for C<sub>26</sub>H<sub>37</sub>FN<sub>2</sub>O<sub>7</sub>S.

### Preparation of intermediate I-05a: 8-tert-butoxycarbonyl-3-[4-(4-methoxyphenoxy)phenylsulfonyl]-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

35 [0109] To a solution of compound I-04a (200 mg, 0.37 mmol) in DMF (5 mL) was added p-methoxyphenol (69 mg, 1.5 eq) and Cs<sub>2</sub>CO<sub>3</sub> (241 mg, 2eq), then the reaction mixture was stirred at 90 °C overnight. After TLC (PE/AE 2:1) showed the starting material was consumed, the mixture was quenched with water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude compound which was purified by Prep-TLC (PE/AE 2:1) to give pure compound I-05a (150 mg, 63.03%) as a pale yellow oil. ESI-MS (M+1): 645 calc. for C<sub>33</sub>H<sub>44</sub>N<sub>2</sub>O<sub>9</sub>S.

### Preparation of compound I-03: N-hydroxy-3-[4-(4-methoxyphenoxy)phenylsulfonyl]-8-azaspiro[4.5]decane-3-carboxamide

45 [0110] A solution of intermediate I-05a (150 mg, 0.233 mmol) in HCl/dioxane (4 M/L, 5mL) was stirred at r.t. for 1 h, then concentrated to give the crude product which was purified by prep-HPLC (General procedure, Method 1) to obtain pure compound I-03 (50 mg, 46.73 %) as a red solid. Rt: 2.63, ESI-MS (M+1): 461 calc. for C<sub>23</sub>H<sub>28</sub>N<sub>2</sub>O<sub>6</sub>S.

### Preparation of compound 1-59: 3-(4-fluorophenylsulfonyl)-N-hydroxy-8-azaspiro[4.5]decane-3-carboxamide

50 [0111] Following the same process as described for compound I-03 but starting from of intermediate I-04a, compound 1-59 was obtained. Rt: 2.11, ESI-MS (M+1): 356 calc. for C<sub>16</sub>H<sub>21</sub>FN<sub>2</sub>O<sub>4</sub>S.


[0112] Following the same synthetic route 1 a, and using the same reagents as for compound I-03 unless otherwise indicated in the table below, the following compounds were obtained:

EP 2 877 446 B9

Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
5 1-01	2.51 and 2.71	475.2	1	4-tert-butoxy-carbonyl-4-azaspiro[5.5]-undecane-9-carboxylic acid (1)
1-06	2.74	489.2	1	4-(methoxycarbonyl) phenol (R-04)
1-07	2.29	475.2	1	4-hydroxybenzoic acid (R-04)
10 1-08	3.04	488.2	1	4-hydroxy-N-methylbenzamide (R-04)
1-09	3.30	487.3	1	4-tert-butylphenol (R-04)
1-12	3.23	462.2	1	6-methoxypyridin-3-ol (R-04)
15 1-13	3.06	488.2	1	N-(4-hydroxyphenyl)acetamide (R-04)
1-15	3.77	477.2	1	4-methoxybenzenethiol (R-04)
1-17	3.31	509.2	1	4-methoxybenzenethiol (R-04) via 1-13
1-19	3.53	445.2	1	p-cresol (R-04)
20 1-22	2.58	488.3	1	4-((Dimethylamino)methyl) phenol (R-04)
1-21	2.50	499.2	1	p-trifluoromethylphenol (R-04)
25 1-43	3.38	433.2	1	6-tert-butoxycarbonyl-6-azaspiro[2.5]octane-2-carboxylic acid (1) and p-methoxyphenol (R-04)
1-54	2.31	461.2	1	p-methoxyphenol (R-04)
1-55	2.18	461.2	1	p-methoxyphenol (R-04)
1-62	3.28	528.2	1	4-hydroxyphenyl-pyrrolidin-1-yl-methanone (R-04)
30 1-63	3.06	502.3	1	4-hydroxy-N,N-dimethyl-benzamide (R-04)
1-64	2.82	474.2	1	4-hydroxybenzamide (R-04)
1-65	3.24	516.2	1	4-hydroxy-N-isopropyl-benzamide (R-04)
35 1-68	3.44	461.2	1	m-methoxyphenol (R-04)
1-69	3.31	461.2	1	o-methoxyphenol (R-04)
1-70	3.08	467.2	1	4-methoxycyclohexanol (R-04)
1-71	3.35	432.1	1	Phenol (R-04)
40 1-72	3.81	489.2	1	4-isopropoxyphenol (R-04)
1-73	3.90	515.2	1	4-(trifluoromethoxy)phenol (R-04)
1-74	3.49	489.3	1	Methyl 3-hydroxybenzoate (R-04)
45 1-75	3.13	475.2	1	3-hydroxybenzoic acid (R-04)
1-76	3.72	466.1	1	4-chlorophenol (R-04)
1-85	2.82	488.1	1	3-hydroxy-N-methylbenzamide (R-04)
1-88	1.89	493.2	1	2-fluoro-4-hydroxy benzoic acid (R-04)
50 1-89	1.78	506.2	1	2-fluoro-4-hydroxy-N-methyl benzamide (R-04)
1-91	3.41	488.1	1	2-hydroxy-N-methylbenzamide (R-04)
1-93	2.10	506.2	1	3-fluoro-4-hydroxy-N-methylbenzamide (R-04)
55 1-94	2.58	507.2	1	Methyl 3-fluoro-4-hydroxybenzoate (R-04)
1-95	3.48	493.2	1	3-fluoro-4-hydroxy-benzoic acid (R-04)
1-110	2.69	518.3	1	4-hydroxy-2-methoxy-N-methyl benzamide (R-04)

(continued)

Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-114	2.78	549.2	1	3-chloro-4-(trifluoromethoxy)phenol (R-04)
1-115	2.60	515.2	1	3-(trifluoromethoxy)phenol (R-04)
1-117	2.37	497.1	1	4-(difluoromethoxy)phenol (R-04)
1-118	2.66	533.2	1	3-fluoro-4-(trifluoromethoxy)phenol (R-04)
1-123	3.39	515.1	1	4-(trifluoromethoxy)phenol (R-04)
1-126	1.71	488.1	1	4-hydroxy-N-methyl benzamide (R-04)
1-127	1.85	475.2	1	4-hydroxy benzoic acid (R-04)

Synthetic route 2a**[0113]**

Conditions: c. R-04 (1.5 - 3 eq), Cs<sub>2</sub>CO<sub>3</sub> (2 - 3 eq), 12 hours at 90 °C; d. HCl in dioxane (4M), r.t.; e. K<sub>2</sub>CO<sub>3</sub> (2 eq), R-08 (1.5 eq) in ACN, r.t.; f. LiOH.H<sub>2</sub>O (10 eq) in THF/MeOH/H<sub>2</sub>O (3/3/2), r.t.; g. EDC.HCl (2 eq), HOBT (2 eq), THP-O-NH<sub>2</sub> (1.5 eq), NMM (3 eq) in DMF, r.t.; h.- HCl / Dioxane, r.t. for 1 hour.

**[0114]** In the scheme above Q is O or S; X is halogen; R' is a hydrocarbon chain optionally substituted; and Cy is phenyl or a 5- to 6-membered heteroaryl and can be optionally substituted.

Preparation of intermediate I-06a: 8-tert-butoxycarbonyl-3-[4-(4-methoxyphenoxy)phenylsulfonyl]-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester

**[0115]** To a solution of compound I-02a (900mg, 1.97 mmol) in DMF (120 mL) was added Cs<sub>2</sub>CO<sub>3</sub> (1.92 g, 5.9 mol)

EP 2 877 446 B9

and 4-methoxyphenol (490 mg, 3.95 mmol), the reaction mixture was stirred at 60 °C for 2h. The mixture was diluted with EtOAc and washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the crude product which was purified by column chromatography (EA:PE=1:30-1:5) to give the compound I-06a (600 mg, 54.0%) as a pale yellow oil. ESI-MS (M+1): 560 calc. for C<sub>29</sub>H<sub>37</sub>NO<sub>8</sub>S.

Preparation of intermediate I-07a: 3-[4-(4-methoxyphenoxy)phenylsulfonyl]-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester

**[0116]** A solution of compound I-06a (500 mg, 0.89 mmol) in HCl/dioxane (4M, 6 mL) was stirred at r.t. for 3h. The reaction mixture was concentrated to give the compound I-07a (500mg, crude). ESI-MS (M+1): 461 calc. for C<sub>24</sub>H<sub>29</sub>NO<sub>6</sub>S.

Preparation of intermediate I-08a: 3-[4-(4-methoxyphenoxy)phenylsulfonyl]-8-methyl-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester

**[0117]** To a solution of intermediate I-07a (181 mg, 0.39 mmol) in CH<sub>3</sub>CN (10 mL) was added K<sub>2</sub>CO<sub>3</sub> (107 mg, 0.78 mmol), Mel (28 mg, 0.19 mmol). The reaction mixture was stirred at room temperature for 30 min. After TLC showed that most of starting material was consumed, the mixture was diluted with EtOAc and washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the crude compound I-08a (110 mg, 59.7%) as a pale yellow oil. ESI-MS (M+1): 588 calc. for C<sub>26</sub>H<sub>33</sub>NO<sub>6</sub>S.

Preparation of intermediate I-09a: 3-[4-(4-methoxyphenoxy)phenylsulfonyl]-8-methyl-8-azaspiro[4.5]decane-3-carboxylic acid

**[0118]** To a solution of intermediate I-08 (110 mg, 0.23 mmol) in THF/MeOH/H<sub>2</sub>O (3/3/2, 8 mL) was added LiOH·H<sub>2</sub>O (195 mg, 4.6 mmol). The resulting mixture was refluxed overnight. After TLC showed that most of the starting materials were consumed completely, the mixture was diluted with water and adjusted pH to 2-3. The mixture was extract with EtOAc and washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the crude product I-09a (100 mg, 94.0%) as a pale yellow solid. ESI-MS (M+1): 460 calc. for C<sub>24</sub>H<sub>29</sub>NO<sub>6</sub>S.

Preparation of intermediate I-10a: 3-[4-(4-methoxyphenoxy)phenylsulfonyl]-8-methyl-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

**[0119]** To a solution of intermediate I-09a (100 mg, 0.22 mmol) in DMF (10 mL) was added EDCI (84 mg, 0.44 mmol), HOBt (59 mg, 0.44 mmol), THP-O-NH<sub>2</sub> (51.5 mg, 0.44 mmol), NMM (66 mg, 0.66 mmol). The mixture was stirred at room temperature overnight. The mixture was diluted with EtOAc and washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated to give the crude product which was purified by column chromatography (EA:PE=1:50-1:4) to give the compound I-10a (78 mg, 64.0%) as a pale yellow solid. ESI-MS (M+1): 559 calc. for C<sub>29</sub>H<sub>38</sub>N<sub>2</sub>O<sub>7</sub>S.

Preparation of compound I-04: N-hydroxy-3-[4-(4-methoxyphenoxy)-phenylsulfonyl]-8-methyl-8-azaspiro[4.5]decane-3-carboxamide

**[0120]** A solution of intermediate I-10a (78 mg, 0.14 mmol) in HCl/dioxane (4 M/L, 10 mL) was stirred at r.t. for 3 h, the reaction mixture was concentrated to give the crude product which was purified by prep-HPLC (General procedure, Method 1) to obtain pure compound 1-04 (14.7 mg, 22.0%) as a pale yellow solid. Rt: 2.66, ESI-MS (M+1): 475 calc. for C<sub>24</sub>H<sub>30</sub>N<sub>2</sub>O<sub>6</sub>S.

**[0121]** Following the same synthetic route 2a, and using the same reagents as for compound 1-04 unless otherwise indicated in the table below, the following compounds were obtained:


Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-02	2.59 and 2.74	489.3	1	4-tert-butoxy-carbonyl-4-azaspiro[5.5]undecane-9-carboxylic acid (1)
1-28	2.56	517.2	1	1-chloropropan-2-one (R-08)
1-29	2.52	519.2	1	tert-butyl 2-bromoacetate (R-08)
1-30	2.96	557.2	1	3,3,3-Trifluoropropyl Bromide (R-08)

(continued)

Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-31	2.80	503.2	1	1-Bromopropane (R-08)
1-32	2.79	501.3	1	Cyclopropyl bromide (R-08)
1-33	3.09	518.2	1	Methylcarbamic chloride (R-08)
1-34	2.39	537.3	1	Iodobenzene (R-08)
1-35	2.26	538.3	1	2-Bromopyridine (R-08)
1-36	3.12	551.3	1	Benzyl bromide (R-08)
1-37	2.52	552.3	1	4-(Bromomethyl) pyridine (R-08)
1-39	2.47	541.2	1	4-(Bromomethyl)-1H-imidazole (R-08)
1-58	3.21	503.2	1	Acetyl chloride (R-08)
1-124	3.53	555.2	1	Cyclopropyl bromide (R-08)

## Synthetic route 1b

## [0122]



Conditions: e. R-05 (4 eq), NaH (0 - 4 eq), after 30 min at r.t., 12h at 90 °C; f.- HCl / Dioxane, r.t. for 1 hour

[0123] In the scheme above R is a hydrocarbon chain which can be optionally substituted.

Preparation of intermediate I-11a: 8-tert-butoxycarbonyl-3-(4-[2-(piperidin-1-yl)ethoxy]phenylsulfonyl)-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

[0124] To a solution of 2-(piperidin-1-yl)ethanol (191 mg, 4 eq) in DMF (5mL) was added NaH (36 mg, 4 eq), then the mixture was stirred at r.t. for 0.5h, intermediate I-04a (200 mg, 0.37 mmol) was added, then the reaction mixture was stirred at 90 °C overnight. After TLC (PE/AE 2:1) showed the starting material was consumed, the mixture was quenched with water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude compound which was purified by Prep-TLC (PE/AE = 2:1) to give pure intermediate I-11a (184 mg, 76.67%) as a pale yellow oil. ESI-MS (M+1): 650 calc. for C<sub>33</sub>H<sub>51</sub>N<sub>3</sub>O<sub>8</sub>S.

Preparation of compound 1-11: N-hydroxy-3-(4-[2-(piperidin-1-yl)ethoxy]phenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxamide


[0125] A solution of intermediate I-11a (180 mg, 0.277 mmol) in HCl/dioxane (5mL) was stirred at r.t. for 1 h, then concentrated to give the crude product by prep-HPLC (General procedure, Method 1) to obtain pure compound 1-11 (58.40 mg, 45.27%) as a pale yellow solid. Rt: 2.14. ESI-MS (M+1): 466 calc. for C<sub>23</sub>H<sub>35</sub>N<sub>3</sub>O<sub>5</sub>S.

[0126] Following the same synthetic route 1b, and using the same reagents as for compound 1-11 unless otherwise indicated in the table below, the following compounds were obtained:

Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-05	2.80	425.2	1	3-methylbutan-1-ol (R-05)
1-10	2.50	466.2	1	1-(2-hydroxyethyl)pyrrolidin-2-one (R-05)
1-18	3.67	437.2	1	Cyclohexanol (R-05)
1-20	3.50	445.2	1	Benzyl alcohol (R-05)
1-92	2.65	451.2	1	Cyclohexylmethanol (R-05)
1-99	3.30	529.0	1	4-(trifluoromethoxy)phenyl methanol (R-05)
1-100	2.96	446.1	1	Pyridin-4-yl methanol (R-05)
1-108	1.98	502.3	1	4-(hydroxymethyl)-N-methyl-benzamide (R-05)
1-112	2.57	435.1	1	1 H-Pyrazol-4-ylmethanol (R-05)
1-113	2.56	369.1	1	Sodium methoxyde (R-05)
1-130	2.97	529.2	1	4-(trifluoromethoxy)phenyl methanol (R-05)

### Synthetic route 1c

#### [0127]



Conditions: e. R-07 (10 eq), 12h at 100 °C; f.- HCl / Dioxane, r.t. for 1 hour

[0128] In the scheme above R' is R or H, and R is a hydrocarbon chain which can be optionally substituted.

Preparation of intermediate I-12a: 8-tert-butoxycarbonyl-3-[4-(4-methoxypiperidin-1-yl)phenylsulfonyl]-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

[0129] A solution of intermediate I-04a (100 mg, 0.185 mmol) and 4-methoxypiperidine (213 mg, 10 eq) was stirred at 100 °C overnight. After TLC (PE/AE 2:1) showed the starting material was consumed, the mixture was quenched with water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude compound which was purified by prep-TLC (PE/AE 1:1) to give pure intermediate I-12a (50 mg, 42.74%) as a pale yellow oil. ESI-MS (M+1): 636 calc. for C<sub>32</sub>H<sub>49</sub>N<sub>3</sub>O<sub>8</sub>S.

Preparation of compound 1-16: N-hydroxy-3-[4-(4-methoxypiperidin-1-yl)phenylsulfonyl]-8-azaspiro[4.5]decane-3-carboxamide

[0130] A solution of intermediate I-12a (50 mg, 0.079 mmol) in HCl/dioxane (3 mL) was stirred at r.t. for 1 h, then concentrated to give the crude product which was purified by prep-HPLC (General procedure, Method 1) to obtain pure compound 1-16 (25.0 mg, 69.44%) as a pale yellow solid. ESI-MS (M+1): 452 calc. for C<sub>22</sub>H<sub>33</sub>N<sub>3</sub>O<sub>5</sub>S.

[0131] Following the same synthetic route 1 c, and using the same reagents as for compound 1-16 unless otherwise indicated in the table below, the following compounds were obtained:


Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-14	3.57	436.3	1	Cyclohexylamine (R-07)

(continued)

Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-61	1.99	354.2	1	Benzylamine (R-07)
1-67	2.51	444.3	1	Benzylamine (R-07)
1-77	3.00	466.3	1	4-methoxycyclohexanamine (R-07)
1-78	1.83	436.3	1	Cyclohexylamine (R-07)
1-79	2.15	436.3	1	Cyclohexylamine (R-07)
1-82	2.02	450.1	1	Cyclohexylmethanamine (R-07)
1-83	2.94	493.2	1	4-amino-N-methylcyclohexanecarboxamide (R-07)
1-84	2.40	480.2	1	4-aminocyclohexanecarboxylic acid (R-07)
1-96	1.47	438.1	1	Tetrahydropyran-4-amine (R-07)
1-97	2.08	422.1	1	Cyclopentanamine (R-07)
1-103	2.58	528.2	1	4-(trifluoromethoxy)phenyl-methanamine (R-07)
1-104	1.56	501.2	1	4-(aminomethyl)-N-methyl benzamide (R-07)
1-105	3.30	502.2	1	Methyl 4-(aminomethyl) benzoate (R-07)
1-106	1.75	488.2	1	4-(aminomethyl) benzoic acid (R-07)
1-109	3.65	472.2	1	4-4-difluorocyclohexylamine (R-07)
1-125	2.20	486.2	1	(4-4-difluorocyclohexyl)methanamine (R-07)

## Synthetic route 1d

[0132]



Conditions: e. Benzyl alcohol (4 eq), NaH (4 eq), after 30 min at r.t., 12 h at 90 °C; f.- Pd/C (0.5 g) at H<sub>2</sub> atmosphere, r.t. for 1 hour; g.- R-09 (1.2 eq), Et<sub>3</sub>N (3 eq) in DCM, r.t. overnight; h.- HCl / Dioxane, r.t. for 1 hour

[0133] In the scheme above R is a hydrocarbon chain optionally substituted, a carbocyclic or heteroaliphatic ring optionally substituted, a phenyl or 5- to 6-membered heteroaryl optionally substituted.

EP 2 877 446 B9

Preparation of intermediate I-13a: 3-[4-(benzyloxy)phenylsulfonyl]-8-tert-butoxycarbonyl-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

[0134] To a solution of phenylmethanol (500 mg, 5eq) in THF (5mL) was added NaH (110 mg, 5eq), then the mixture was stirred at r.t for 0.5 h, intermediate I-04a (500 mg, 0.926 mmol) was added, then the reaction mixture was stirred at 80 °C overnight. After TLC (PE/AE 2:1) showed the starting material was consumed, the mixture was quenched with water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude compound which was purified by prep-TLC (PE/AE 2:1) to give pure intermediate I-13a (380 mg, 65.29%) as a pale yellow oil. ESI-MS (M+1): 629 calc. for C<sub>33</sub>H<sub>44</sub>N<sub>2</sub>O<sub>8</sub>S.

Preparation of intermediate I-14a: 8-tert-butoxycarbonyl-3-[4-hydroxyphenylsulfonyl]-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

[0135] To a solution of intermediate I-13a (380 mg, 0.605 mmol) in MeOH (20 mL) was added Pd/C (0.5 g) at room temperature at H<sub>2</sub> atmosphere, then the mixture was stirred at room temperature for 1 h until TLC (PE/AE 2:1) showed the starting material was consumed, then filtered and the filtrate was concentrated to give the crude product I-14a (180 mg, 55.38%) which was used for the next step without further purification. ESI-MS (M+1): 539 calc. for C<sub>26</sub>H<sub>38</sub>N<sub>2</sub>O<sub>8</sub>S.

Preparation of intermediate I-15a: 8-tert-butoxycarbonyl-3-[4-(4-methoxybenzoyloxy)phenylsulfonyl]-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

[0136] To a solution of intermediate I-14a (60 mg, 0.151 mmol) and Et<sub>3</sub>N (46 mg, 3eq) in DCM (5mL) was added 4-methoxybenzoyl chloride (31 mg, 1.2eq) at room temperature, then the mixture was stirred at r.t overnight. After TLC (PE/AE 2:1) showed the starting material was consumed, the mixture was extracted with DCM, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude compound which was purified by prep-TLC (PE/AE 2:1) to give pure compound I-15a (38 mg, 37.62%) as a pale yellow oil. ESI-MS (M+1): 673 calc. for C<sub>34</sub>H<sub>44</sub>N<sub>2</sub>O<sub>10</sub>S.

Preparation of compound 1-26: N-hydroxy-3-[4-(4-methoxybenzoyloxy)phenylsulfonyl]-8-azaspiro[4.5]decane-3-carboxamide

[0137] A solution of intermediate I-15a (38 mg, 0.057 mmol) in HCl/dioxane (5 mL) was stirred at r.t for 1 h, then concentrated to give the crude product by prep-HPLC (General procedure, Method 1) to obtained pure compound I-26 (22 mg, 78.57 %) as a pale yellow solid. Rt: 2.71, ESI-MS (M+1): 489 calc. for C<sub>24</sub>H<sub>28</sub>N<sub>2</sub>O<sub>7</sub>S.

Preparation of compound 1-60: N-hydroxy-3-(4-hydroxyphenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxamide


[0138] Following the same process as described for compound 1-26 but starting from of intermediate I-14a, compound 1-60 was obtained. Rt: 2.11, ESI-MS (M+1): 354 calc. for C<sub>16</sub>H<sub>22</sub>N<sub>2</sub>O<sub>5</sub>S.

[0139] Following the same synthetic route 1d, and using the same reagents as for compound 1-26 unless otherwise indicated in the table below, the following compounds were obtained:

Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-86	3.12	503.1	1	4-chlorocarbonyl benzoic acid (R-09)
1-87	2.79	516.2	1	4-(methylcarbonyl)benzoyl chloride (R-09)
1-98	2.47	465.3	1	Cyclohexanecarbonyl chloride (R-09)
1-101	1.84	534.2	1	3-fluoro-4-(methylcarbonyl)benzoyl chloride (R-09)
1-102	2.75	543.1	1	4-(trifluoromethoxy)benzoyl chloride (R-09)
1-119	2.20	489.0	1	4-methoxybenzoyl chloride (R-09)
1-120	2.61	543.1	1	4-(trifluoromethoxy)benzoyl chloride (R-09)

Synthetic route 1e

[0140]



25  
Conditions: a. LDA (2.0 M) in THF, 1 hour at  $-78\text{ }^{\circ}\text{C}$ ; then, R-10 (1.2 eq), 1 hour at  $-78\text{ }^{\circ}\text{C}$ ; b. Pd(dppf)Cl<sub>2</sub> (0.1 eq) and Na<sub>2</sub>CO<sub>3</sub> (2 eq) in dioxane/ H<sub>2</sub>O, r.t. overnight; c. LiOH.H<sub>2</sub>O (10 eq) in THF/MeOH/H<sub>2</sub>O (3/3/2), r.t.; d. EDC.HCl (2 eq), HOBT (2 eq), THP-O-NH<sub>2</sub> (1.5 eq), NMM (3 eq) in DMF, r.t.; e.- HCl / Dioxane, r.t. for 1 hour.

30  
[0141] In the scheme above B is boronic acids, boronate esters or trifluoroborate salts and R is phenyl or a 5- to 6-membered heteroaryl or a 3- to 7-heterocyclic or carbocyclic aliphatic ring or a hydrocarbon chain which can be optionally substituted.

#### Preparation of reagent R-10a: 4-bromobenzenesulfonyl fluoride

35  
[0142] To a solution of 4-bromobenzenesulfonyl chloride (2.29 g, 9 mmol) in CH<sub>3</sub>CN (45 mL) was added KF (2.1 g, 36 mmol) and 18-crown-6 (0.5 g), then the mixture was stirred at room temperature overnight. The mixture was quenched with aqueous water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product which was purified by column to give the reagent R-10a (1.6 g, 74.4%) as a pale yellow solid. ESI-MS (M+1): 239.2 calc. for C<sub>6</sub>H<sub>4</sub>BrFO<sub>2</sub>S.

#### Preparation of intermediate I-16a: 8-tert-butoxycarbonyl-3-(4-bromophenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester

45  
[0143] To a solution of compound I-01a (230 mg, 0.77 mmol) in THF (20 mL) was added LDA (0.6 mL, 2.0 M, 1.2 mmol) at  $-78\text{ }^{\circ}\text{C}$ . After stirring at  $-78\text{ }^{\circ}\text{C}$  for 1 hour, the reagent R-10a (214 mg, 0.9 mmol) was added to the solution, the reaction was stirred at  $-78\text{ }^{\circ}\text{C}$  for 1 hour, and then the mixture was stirred at r.t overnight. After TLC (PE/AE 5:1) showed the starting material was consumed, the mixture was quenched with aqueous NH<sub>4</sub>Cl and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product which was purified by column chromatography (eluting with EA/PE= 100:1 to 10: 1) to give pure compound I-16a (150 mg, 38%) as a pale yellow oil. ESI-MS (M-55): 459.9 calc. for C<sub>22</sub>H<sub>30</sub>BrNO<sub>6</sub>S.

#### Preparation of intermediate I-17a: 8-tert-butoxycarbonyl-3-(4-(4-methoxyphenyl)phenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester

55  
[0144] To a suspension of compound I-16a (50 mg, 0.1 mmol) and the commercially available 4-methoxyphenyl boronic acid (20 mg, 0.13 mmol, R-11), in dioxane (10 mL)/ H<sub>2</sub>O (2 mL) was added Pd(dppf)Cl<sub>2</sub> (10 mg) and Na<sub>2</sub>CO<sub>3</sub> (22 mg, 0.2 mmol). The mixture was stirred at reflux under N<sub>2</sub> overnight. The resulting mixture was cooled to room temperature and water was added. The organic layer was separated and the aqueous layer was extracted 3 times with EtOAc. The

combined organic layer was washed with brine, dried over sodium sulfate and concentrated under reduced pressure and purified by prep-TLC to afford the desired product I-17a (52 mg, 95%). ESI-MS (M-55): 488.0 calc. for  $C_{29}H_{37}NO_7S$ .

Preparation of intermediate I-18a: 8-tert-butoxycarbonyl-3-(4-(4-methoxyphenyl)phenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxylic acid

[0145] To a solution of compound I-17a (52 mg, 0.096 mmol) in THF/MeOH/H<sub>2</sub>O (3/3/2, 8 mL) was added LiOH.H<sub>2</sub>O (42 mg, 10 eq). The resulting mixture was stirred at r.t. for 4 h, after TLC (PE/AE 5:1) showed the starting materials were consumed completely, then the mixture was diluted with water and adjusted pH to 3-4 with 1 N HCl and the mixture was extracted with EtOAc, washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product I-18a (50 mg, ~100%) as a pale yellow oil which was used directly in the next step. ESI-MS (M-55): 473.9; calc. for  $C_{28}H_{35}NO_7S$

Preparation of intermediate I-19a: 8-tert-butoxycarbonyl-3-(4-(4-methoxyphenyl)phenylsulfonyl)-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

[0146] To a solution of compound I-18a (50 mg, 0.095 mmol) in DMF (15 mL) was added EDC.HCl (40 mg, 0.2 mmol), HOBT (27 mg, 0.2 mmol), THPONH<sub>2</sub> (24 mg, 0.2 mmol), NMM (40 mg, 0.4 mmol), then the mixture was stirred at room temperature overnight. The mixture was quenched with aqueous water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product which was purified by prep-TLC to give the compound I-19a (45 mg, 72%) as a pale yellow solid. ESI MS (M-139): 488.9; calc. for  $C_{33}H_{44}N_2O_8S$ .

Preparation of compound 1-41: N-hydroxy-3-(4-(4-methoxyphenyl)phenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxamide


[0147] A solution of intermediate I-19a (45 mg, 0.072 mmol) in HCl/dioxane (10 mL, 2 N) was stirred at r.t for 2 h, then concentrated to give the crude product which was purified by prep HPLC twice (General procedure, Method 1) to obtain pure compound 1-41 (6.6 mg, 16.8 %) as a yellow solid. ESI-MS (M+1): 445.2 calc. for  $C_{23}H_{28}N_2O_5S$ . Rt is 2.01.

[0148] Following the same synthetic route 1 e, and using the same reagents as for compound 1-41 unless otherwise indicated in the table below, the following compounds were obtained:

Example	R <sub>t</sub>	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-66	2.26	429.2	1	1,3,2-Dioxaborolane, 4,4,5,5-tetramethyl-2-(phenylmethyl)-(R-11)

Synthetic route 1f

[0149]



Conditions: a. R-09 (1.2 eq), Et<sub>3</sub>N (2 eq) in DCM, r.t. overnight; b. HCl / Dioxane, r.t. for 1 hour.

[0150] In the scheme above R and R' are hydrogen, X is a halogen and R'' is phenyl or a 5- to 6- membered heteroaryl or a 3- to 7- heterocyclic or carbocyclic aliphatic ring or a hydrocarbon chain which can be optionally substituted.

## EP 2 877 446 B9

Preparation of intermediate I-20a: 8-tert-butoxycarbonyl-3-(4-(cyclohexanecarbonylamino)phenylsulfonyl)-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

5 **[0151]** To a solution of compound I-12c (80 mg, 0.149 mmol), intermediate obtained to synthesize 1-61 (following synthetic route 1 c), in DCM (3 mL) was added cyclohexanecarbonyl chloride (26.2mg, 1.2eq) and Et<sub>3</sub>N (30.1 mg, 0.298 mmol), then the reaction mixture was stirred at r.t overnight. After TLC (PE/AE 5:1) showed the starting material was consumed, the mixture was quenched with water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude compound which was purified by Prep-TLC (PE/AE 5:1) to give pure intermediate I-20a (65 mg, 67.50%) as a white solid. ESI-MS (M+1): 648.3 calc. for C<sub>33</sub>H<sub>49</sub>N<sub>3</sub>O<sub>8</sub>S.

10 Preparation of compound 1-81: N-hydroxy-3-(4-(cyclohexanecarbonylamino)phenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxamide

15 **[0152]** A solution of compound I-20a (65 mg, 0.1 mmol) in HCl/dioxane (3 mL) was stirred at r.t for 1 h, then concentrated to give the crude product which was purified by prep-HPLC (General procedure, Method 1) to obtained pure compound 1-81 (22.0 mg, 45.74 %) as a yellow solid. ESI-MS (M+1): 464.2 calc. for C<sub>23</sub>H<sub>33</sub>N<sub>3</sub>O<sub>5</sub>S. Rt is 1.89.

Synthetic route 3a

20 **[0153]**

25

30


35

40

45

50

55



Conditions: a. LDA (1.20 M) in THF, 1 hour at  $-78\text{ }^{\circ}\text{C}$ ; then, R-12 (1.2 eq), 1 hour at  $-78\text{ }^{\circ}\text{C}$ ; b. R-13 (3 - 10 eq), NaH (0 - 3 eq) in THF, r.t. - reflux overnight; c. CuI (2 eq), MeOH/MeONa (25%, 20 eq) in DMF, 20 minutes at  $110\text{ }^{\circ}\text{C}$ ; d. LiOH.H<sub>2</sub>O (10 eq) in THF/MeOH/H<sub>2</sub>O (3/3/2), r.t.; e. EDC.HCl (2 eq), HOBT (2 eq), THP-O-NH<sub>2</sub> (1.5 eq), NMM (3 eq) in DMF, r.t.; f. HCl / Dioxane, r.t. for 1 hour.

**[0154]** In the scheme above Q is O or NH, R is a 3- to 7- heterocyclic or carbocyclic aliphatic ring or a hydrocarbon chain which can be optionally substituted and R' is halogen or alcoxy.

#### Preparation of reagent R-12a: 3-chloro-4-fluorobenzenesulfonyl fluoride

**[0155]** To a solution of the commercially available 3-chloro-4-fluorobenzenesulfonyl chloride (7 g, 30 mmol) in CH<sub>3</sub>CN (40 mL) was added KF (7 g, 120 mmol) and 18-crown-6 (0.5 g), then the mixture was stirred at room temperature overnight. The mixture was quenched with aqueous water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product which was purified by column to give the reagent R-12a (4.2 g, 95.5%) as a pale yellow solid. ESI-MS (M+1): 213.2 calc. for C<sub>6</sub>H<sub>3</sub>ClF<sub>2</sub>O<sub>2</sub>S

#### Preparation of intermediate I-21a: 8-tert-butoxycarbonyl-3-(3-chloro-4-fluorophenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxylic acid methyl ester

**[0156]** To a solution of compound I-01a (743 mg, 2.5 mmol) in THF (20 mL) was added LDA (4.2 mL, 1.2 M, 5 mmol) at  $-78\text{ }^{\circ}\text{C}$ . After stirring at  $-78\text{ }^{\circ}\text{C}$  for 1 hour, the reagent R-12a (640 mg, 3.0 mmol) was added to the solution, the reaction was stirred at  $-78\text{ }^{\circ}\text{C}$  for 1 hour, and then the mixture was stirred at r.t. overnight. After TLC (PE/AE 5:1) showed the starting material was consumed, the mixture was quenched with aqueous NH<sub>4</sub>Cl and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product which was purified

## EP 2 877 446 B9

by column chromatography (eluting with EA/PE= 100:1 to 10: 1) to give pure intermediate I-21a (349 mg, 29%) as a pale yellow oil. ESI-MS (M-55): 434.1 calc. for  $C_{22}H_{29}ClFNO_6S$ .

Preparation of intermediate I-22a: 8-tert-butoxycarbonyl-3-(3-chloro-4-(cyclohexylamino)phenylsulfonyl)-8-aza-spiro[4.5]decane-3-carboxylic acid methyl ester

**[0157]** The intermediate I-21a (240 mg, 0.51 mmol) was dissolved in cyclohexylamine (505 mg, 5.1 mmol), R13a. The solution was stirred at 85 °C overnight. The mixture was quenched with aqueous water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous  $Na_2SO_4$ , concentrated to give the crude product which was purified by prep-TLC to give the compound I-22a (110 mg, 38%) as a pale yellow solid. ESI-MS (M-55): 513; calc. for  $C_{28}H_{41}ClN_2O_6S$ .

Preparation of intermediate I-24a: 8-tert-butoxycarbonyl-3-(3-methoxy-4-(cyclohexylamino)phenylsulfonyl)-8-aza-spiro[4.5]decane-3-carboxylic acid

**[0158]** The compound I-22a (179 mg, 0.326 mmol), CuI (124.5 mg, 0.652 mmol) and MeOH/MeONa (25%, 1.41 g, 6.52 mmol) was dissolved in DMF (10 mL). The solution was stirred at 110 °C for 20 min. The mixture was quenched with aqueous water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous  $Na_2SO_4$ , concentrated to give the crude product which was purified by prep-TLC to give the compound I-24a (160 mg, 88%) as a pale yellow solid. The hydrolysis of the methyl ester to the corresponding carboxylic acid occurred simultaneously together with the methoxylation; thus, compound I-23a was not isolated and the intermediate I-24a was directly obtained from I-22a. ESI-MS (M-55): 495.2; calc. for  $C_{28}H_{42}N_2O_7S$ .

Preparation of intermediate I-25a: 8-tert-butoxycarbonyl-3-(3-methoxy-4-(cyclohexylamino)phenylsulfonyl)-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

**[0159]** To a solution of compound I-24a (160 mg, 0.29 mmol) in DMF (15 mL) was added EDC.HCl (112 mg, 0.58 mmol), HOBT (79 mg, 0.58 mmol), THPONH<sub>2</sub> (68 mg, 0.58 mmol) and NMM (88 mg, 0.87 mmol) at r.t, then the mixture was stirred at room temperature overnight. The mixture was quenched with aqueous water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous  $Na_2SO_4$ , concentrated to give the crude product which was purified by prep-TLC to give the intermediate I-25a (100 mg, 53%) as a pale yellow solid. ESI-MS (M-139): 510.1; calc. for  $C_{33}H_{51}N_3O_8S$ .

Preparation of compound 1-90: N-hydroxy-3-(3-methoxy-4-(cyclohexylamino)phenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxamide

**[0160]** A solution of compound I-25a (100 mg, 0.15 mmol) in HCl/dioxane (10 mL) was stirred at r.t for 2 h, then concentrated to give the crude product which was purified by prep HPLC (General procedure, Method 1) to obtained pure compound 1-90 (39.7 mg, 58 %) as a white solid. ESI-MS (M+1): 466.1 calc. for  $C_{23}H_{35}N_3O_5S$ . Rt is 2.48.

Preparation of compound 1-116: N-hydroxy-3-(3-chloro-4-fluorophenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxamide


**[0161]** Following the same process as described for compound 1-90 but skipping the second and third step (b and c) from synthetic route described above (3a), compound 1-116 was obtained. Rt is 2.96, ESI-MS (M+1): 391.2 calc. for  $C_{16}H_{20}ClFN_2O_4S$ .

**[0162]** Following the same synthetic route 3a, and using the same reagents as for compound 1-90 unless otherwise indicated in the table below, the following compounds were obtained:

Example	Rt	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-80	2.26	470.1	1	Cyclohexylamine (R-13)
1-107	2.42	403.1	1	Sodium trimethylsilanoate (R-13)
1-121	2.64	389.1	1	Sodium trimethylsilanoate (R-13)
1-122	2.49	479.2	1	Benzyl alcohol (R-13)

## Synthetic route 3b

[0163]



Conditions: a. R-04 (1.5 - 3 eq), Cs<sub>2</sub>CO<sub>3</sub> (2 - 3 eq), 12 hours at 90 °C; b. LiOH.H<sub>2</sub>O (10 eq) in THF/MeOH/H<sub>2</sub>O (3/3/2), r.t.; c. EDC.HCl (2 eq), HOBT (2 eq), THP-O-NH<sub>2</sub> (1.5 eq), NMM (3 eq) in DMF, r.t.; d. HCl / Dioxane, r.t. for 1 hour.

30 [0164] In the scheme above Q is O or S and Cy is phenyl or a 5- to 6- membered heteroaryl and can be optionally substituted.

35 Preparation of intermediate I-26a: 8-tert-butoxycarbonyl-3-(3-chloro-4-(4-methoxyphenoxy)phenylsulfonyl)-8-aza-spiro[4.5]decane-3-carboxylic acid methyl ester

35 [0165] To a suspension of compound I-21a (420 mg, 0.86 mmol) and p-methoxyphenol (214 mg, 1.72 mmol), in DMF (5 mL) was added Cs<sub>2</sub>CO<sub>3</sub> (700 mg, 2.2 mmol). The mixture was stirred at reflux under N<sub>2</sub> overnight. The resulting mixture was cooled to room temperature and water was added. The organic layer was separated and the aqueous layer was extracted 3 times with EtOAc. The combined organic layers were washed with brine, dried over sodium sulfate and concentrated under reduced pressure and purified by prep TLC to afford the desired product I-26a (380 mg, 75%). ESI-MS (M-55): 538.2 calc. for C<sub>29</sub>H<sub>36</sub>ClNO<sub>8</sub>S.

40

45 Preparation of intermediate I-27a: 8-tert-butoxycarbonyl-3-(3-chloro-4-(4-methoxyphenoxy)phenylsulfonyl)-8-aza-spiro[4.5]decane-3-carboxylic acid

45 [0166] To a solution of compound I-26a (310 mg, 0.52 mmol) in THF/MeOH/H<sub>2</sub>O (3/3/2, 16 mL) was added LiOH.H<sub>2</sub>O (225 mg, 10 eq). The resulting mixture was stirred at r.t. for 4 h, after TLC (PE/AE 5:1) showed the starting materials were consumed completely, then the mixture was diluted with water and adjusted pH to 3~4 with 1 N HCl and the mixture was extracted with EtOAc, washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated and purified by prep-TLC to afford the desired product I-27a (120 mg, 40%). ESI-MS (M-55): 524.2; calc. for C<sub>28</sub>H<sub>34</sub>ClNO<sub>8</sub>S.

50

Preparation of intermediate I-28a: 8-tert-butoxycarbonyl-3-(3-chloro-4-(4-methoxyphenoxy)phenylsulfonyl)-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

55 [0167] To a solution of intermediate I-27a (120 mg, 0.21 mmol) in DMF (10 mL) was added EDC.HCl (81 mg, 0.42 mmol), HOBT (57 mg, 0.42 mmol), THPONH<sub>2</sub> (50 mg, 0.42 mmol), NMM (64 mg, 0.63 mmol), then the mixture was stirred at room temperature overnight. The mixture was quenched with aqueous water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product I-28a (160

mg, ~100%) as a pale yellow oil which was used directly in the next step. ESI-MS (M-139): 539.2; calc. for C<sub>33</sub>H<sub>43</sub>ClN<sub>2</sub>O<sub>9</sub>S.

Preparation of compound 1-56: N-hydroxy-3-(3-chloro-4-(4-methoxyphenoxy)phenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxamide


[0168] A solution of intermediate I-28a (160 mg, ~0.21 mmol) in HCl/dioxane (10 mL, 1 N) was stirred at r.t for 2 h, then concentrated to give the crude product which was purified by prep-HPLC (General procedure, Method 1) to obtained pure compound 1-56 (30.8 mg, 29.7 %) as a yellow solid. ESI-MS (M+1): 495.2 calc. for C<sub>23</sub>H<sub>27</sub>ClN<sub>2</sub>O<sub>6</sub>S. Rt is 2.66.

[0169] Following the same synthetic route 3b, and using the same reagents as for compound 1-56 unless otherwise indicated in the table below, the following compounds were obtained:

Example	Rt	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-111	3.51	549.2	1	4-(trifluoromethoxy)phenol (R-04)

Synthetic route 3c

[0170]



Conditions: a. R-09 (1.2 eq), Et<sub>3</sub>N (4 eq) in DCM, r.t. overnight; b. HCl / Dioxane, r.t. for 1 hour.

[0171] In the scheme above Q is O or NH, R is hydrogen, X is a halogen and R' is phenyl or a 5- to 6- membered heteroaryl or a 3- to 7- heterocyclic or carbocyclic aliphatic ring or a hydrocarbon chain which can be optionally substituted.

Preparation of intermediate I-29a: 8-tert-butoxycarbonyl-3-(3-chloro-4-(4-(trifluoromethoxy)benzoyloxy)phenylsulfonyl)-N-(tetrahydro-2H-pyran-2-yloxy)-8-azaspiro[4.5]decane-3-carboxamide

[0172] To a solution of compound I-25e (115 mg, 0.2 mmol), intermediate obtained to synthesize 1-121 (following synthetic route 3a), and Et<sub>3</sub>N (61 mg, 0.6 mmol) in DCM (10 mL) was added dropwise the commercially available 4-(Trifluoromethoxy)benzoyl chloride, R-09, (0.3 mmol) in DCM (5 mL) with stirring at 0 °C under N<sub>2</sub>. The mixture solution was stirred at r.t overnight. The mixture solution was added water and extracted with EtOAc, the organic layer was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, concentrated to give the crude product which was purified by prep-TLC to give the intermediate I-29a (40 mg, 26%) as a yellow solid. ESI-MS (M-139): 621.1; calc. for C<sub>34</sub>H<sub>40</sub>ClF<sub>3</sub>N<sub>2</sub>O<sub>10</sub>S.

Preparation of compound 1-128: N-hydroxy-3-(3-chloro-4-(4-(trifluoromethoxy)benzoyloxy)phenylsulfonyl)-8-azaspiro[4.5]decane-3-carboxamide

[0173] A solution of intermediate I-29a (40 mg, 0.053 mmol) in HCl/dioxane (15 mL, 1 N) was stirred at r.t for 4 h, then concentrated to give the crude product which was purified by prep-HPLC (General procedure, Method 1) to obtained pure compound 1-128 (11.2 mg, 37 %) as a yellow solid. ESI-MS (M+1): 577.2 calc. for C<sub>24</sub>H<sub>24</sub>ClF<sub>3</sub>N<sub>2</sub>O<sub>7</sub>S. Rt is 2.96.

[0174] Following the same synthetic route 3c, and using the same reagents as for compound 1-128 unless otherwise indicated in the table below, the following compounds were obtained:

Example	Rt	[M+1] <sup>+</sup>	LC-MS Method	Starting materials
1-129	2.60	523.2	1	4-methoxybenzoyl chloride (R-09)

## EP 2 877 446 B9

**[0175]** Synthesized compounds are obtained as racemic mixtures. Corresponding isomers are purified by supercritical fluid chromatography (SFC) to obtain two enantiomers from each racemic compound.

### Procedure for Supercritical Fluid Chromatography (SFC):

5

**[0176]** Details about the preparative SFC separation methods utilized are as follows:

#### Method 1:

10

- Instrument: Thar SFC Pre-80
- Column: ChiralPak AD-H, 250×30 mmI.D.
- Mobile phase: A for CO<sub>2</sub> and B for Ethanol(0.1%NH<sub>3</sub>•H<sub>2</sub>O)
- Gradient: B 50%
- Flow rate: 60 mL /min
- 15 - Back pressure: 100 bar
- Column temperature: 40 °C
- Wavelength: 220 nm
- Cycletime: ~20 min
- Sample preparation: Compound was dissolved in ethanol to ~12 mg/mL
- 20 - Injection: 2.0 mL per injection.

#### Method 2:

25

- Instrument: Mg II preparative SFC
- Column: ChiralPak AD-H, 250×30mmI.D.
- Mobile phase: A for CO<sub>2</sub> and B for Methanol(0.1%NH<sub>3</sub>•H<sub>2</sub>O)
- Gradient: B 50%
- Flow rate: 40mL /min
- Back pressure: 100bar
- 30 - Column temperature: 38 °C
- Wavelength: 220 nm
- Cycletime: ~20min
- Sample preparation: Compound was dissolved in methanol to ~13mg/ml
- Injection: 3.5 mL per injection.

35

#### Method 3:

40

- Instrument: Mg II preparative SFC
- Column: ChiralPak AD-H, 250×30mmI.D.
- Mobile phase: A for CO<sub>2</sub> and B for IPA(0.1 %NH<sub>3</sub>•H<sub>2</sub>O)
- Gradient: B 35%
- Flow rate: 55ml /min
- Back pressure: 100bar
- Column temperature: 38 °C
- 45 - Wavelength: 254 nm
- Cycletime: ~8 min
- Sample preparation: Compound was dissolved in methanol to ~ 5 mg/mL
- Injection: 1.0 mL per injection.

50

**[0177]** After separation, the fractions were dried off via rotary evaporator at bath temperature 40 °C to get the desired isomers. Then, after concentration, enantiomeric excess (e.e.) for both enantiomers were tested under the analytical separation methods described below:

#### Method 1:

55

- Instrument: SHIMADZU-20A UFLC
- Column: ChiralPak AD-3, 100x4.6 mm
- Mobile phase: A for Hexane (0.1%IPAm) and B for Ethanol(0.05%IPAm)

- Gradient: B 40%
- Flow rate: 1.0 mL/min
- Column temperature: 30 °C
- Wavelength: 220 nm

5

## Method 2:

- Instrument: Thar analytical SFC
- Column: ChiralPak AD-H, 250x4.6mm
- Mobile phase: A for CO<sub>2</sub> and B for Methanol (0.05%DEA)
- Gradient: B 50%
- Flow rate: 2.0mL/min
- Back pressure: 100bar
- Column temperature: 35°C
- Wavelength: 220nm

10

15

## Method 3:

- Instrument: Thar analytical SFC
- Column: ChiralPak AD-H, 250x4.6mm
- Mobile phase: A for CO<sub>2</sub> and B for IPA (0.1 %Ethanolamine)
- Gradient: B 30%
- Flow rate: 2.4 mL/min
- Back pressure: 100 bar
- Column temperature: 35 °C
- Wavelength: 220 nm

20

25

**[0178]** After concentration, as previously reported, LC-MS was used to test the purity for both enantiomers. Then, optical rotation was measured at 20 °C using the Autopol V polarimeter.

30

**[0179]** The following isomers were obtained after chiral separation:

Starting material	Example	R <sub>t</sub> (SFC)	SFC Method	e.e. by chiral HPLC (%)	Chiral HPLC Method	Optical Rotation [α] <sub>D</sub>
1-107	1-131	2.42	1	100	1	+ 19.59°
1-107	1-132	3.15	1	95.3	1	- 19.71°
1-08	1-133	2.98	2	100	2	+ 16.66°
1-08	1-134	5.88	2	100	2	- 17.10°
1-55	1-135	4.44	1	100	1	---
1-55	1-136	6.08	1	97.9	1	---
1-79	1-137	5.85	3	96.1	3	---
1-79	1-138	7.63	3	97.7	3	---

35

40

45

#### Antifibrinolytic effect on whole blood clot formation and lysis

**[0180]** Thromboelastometry is a viscoelastometric method for haemostasis testing in whole blood. TEM® measures the interactions of coagulation factors, inhibitors and cellular components during the phases of clotting and subsequent lysis over time. The rheological conditions of this method mimic the sluggish flow of blood in veins.

50

#### Detection method:

**[0181]** Blood samples were obtained between 8-9 a.m. from healthy volunteers and mice in tubes containing citrate solution (0.129 M sodium citrate, Vacutainer BD) and ROTEM tests were performed following the technical details of the ROTEM® analyser (Pentapharm GmbH, Munich, Germany). A modification of in-tem test as described below was

55

## EP 2 877 446 B9

used for the examination of antifibrinolytic effects of tested compounds and its interaction with platelets in citrated blood. Kits: START-TEM assay as a recalcification reagent (ref#503-01) and IN-TEM assay for activation of intrinsic coagulation pathway (ref #503-02).

### 5 Procedure:

10 **[0182]** In a pre-warmed cuvette and holder 1  $\mu\text{L}$  of Tissue plasminogen activator (tPA) (150,000 U/mL, Actylise), 20  $\mu\text{L}$  of start-tem reagent ( $\text{CaCl}_2$ ), 20  $\mu\text{L}$  of in-tem reagent (activators of coagulation system), 3  $\mu\text{L}$  of DMSO (control) or tested compounds (CMs) in DMSO and 300  $\mu\text{L}$  of citrated blood pre-warmed were pipetted. The cup holder containing the sample mixture was placed immediately on the appropriate channel. The measurement was recorded for 60 min to allow clot formation and lysis.

15 **[0183]** Table 3 shows the results in human blood as effective concentration to delay lysis time by 50% ( $\text{EC}_{50\text{LT}}$ ); where,  $\text{EC}_{50\text{LT}} \geq 25 \mu\text{M}$  (+),  $10 \mu\text{M} \leq \text{EC}_{50\text{LT}} < 25 \mu\text{M}$  (++) ,  $1 \mu\text{M} \leq \text{EC}_{50\text{LT}} < 10 \mu\text{M}$  (+++) ,  $\text{EC}_{50\text{LT}} < 1 \mu\text{M}$  (++++) at all the assayed concentrations (1000-0.2  $\mu\text{M}$ ).

Table 3

Example	$\text{EC}_{50\text{LT}}$	Example	$\text{EC}_{50\text{LT}}$
TXA	+++	1-58	++++
1-01	++++	1-59	+
1-02	++	1-60	+
1-03	++++	1-21	++++
1-04	++++	1-41	+
1-05	++	1-43	++++
1-06	+	1-54	+++
1-07	++++	1-55	++++
1-08	++++	1-56	++++
1-09	++++	1-61	+
1-10	++++	1-62	++++
1-11	++++	1-63	+++
1-12	++	1-64	+
1-13	++++	1-65	+++
1-14	++++	1-66	+
1-15	+	1-67	++++
1-16	+++	1-68	+++
1-17	+++	1-69	+++
1-18	+++	1-70	++++
1-19	++++	1-71	+++
1-20	++++	1-72	+++
1-22	+++	1-73	++++
1-26	++++	1-74	+++
1-28	+++	1-75	+++
1-29	+++	1-76	+
1-30	++++	1-77	+++
1-31	++++	1-78	+
1-32	+++	1-79	++++

EP 2 877 446 B9

(continued)

5  
10  
15  
20  
25  
30  
35  
40  
45  
50

Example	EC <sub>50LT</sub>	Example	EC <sub>50LT</sub>
1-33	+	1-80	++++
1-34	+	1-81	++++
1-35	+++	1-82	++++
1-36	+++	1-83	+
1-37	+	1-84	++++
1-39	+	1-85	+++
1-86	++++	1-113	++++
1-87	+++	1-114	++
1-88	++++	1-115	++++
1-89	+	1-116	++++
1-90	+	1-117	++++
1-91	+	1-118	++++
1-92	+++	1-119	++++
1-93	+	1-120	+
1-94	+	1-121	+++
1-95	+	1-122	+
1-96	++++	1-123	++++
1-97	+++	1-124	+++
1-98	+	1-125	++++
1-99	++++	1-126	+
1-100	++++	1-127	++++
1-101	+++	1-128	+
1-102	+	1-129	+++
1-103	++++	1-130	+++
1-104	++++	1-131	++++
1-105	+	1-132	++++
1-106	++++	1-133	++++
1-107	++++	1-134	++++
1-108	++++	1-135	++++
1-109	+	1-136	++++
1-110	++++	1-137	++++
1-111	+	1-138	++++
1-112	++++		

**[0184]** Table 4 shows the results in mice blood as effective concentration to delay lysis time by 50% (EC<sub>50LT</sub>); where, EC<sub>50LT</sub> ≥ 10 μM (+), 1 μM ≤ EC<sub>50LT</sub> < 10 μM (++) , 1 nM ≤ EC<sub>50LT</sub> < 1 μM (+++) and EC<sub>50LT</sub> < 1 nM (++++) for all the assayed concentrations (1000-0.2 μM).

EP 2 877 446 B9

Table 4

Example	EC <sub>50LT</sub>
TXA	+++
1-03	++++
1-07	++++
1-08	+++
1-14	+++
1-62	++++
1-73	++

5 [0185] As observed in the tables above (Tables 3 and 4), compounds of the invention show significant delay in the lysis time, in most cases higher than TXA.

Antifibrinolytic effect *in vivo* (tail bleeding assay)

20 [0186] Bleeding time was evaluated in 2 months old wild-type C57Bl6 (n=10) mice by removing the tail tip. Mice (20-25 g) were anaesthetized with 2.5% isoflurane and maintained at 37 °C on heating pads. The hemostatic efficacy was evaluated in 2 models, traditional bleeding model and hyperfibrinolytic bleeding model.

25 [0187] Traditional bleeding model consisted in intraperitoneal injection of different compounds (4 mg/mouse, 160 mg/Kg) dissolved in 2% carboxymethyl cellulose or vehicle. Thirty minutes after injection, 5 mm of tail tip were removed using a scalpel blade and the tail tip bathed in 1 mL of sterile saline at 37 °C. The time to cessation of bleeding was measured up to 30 min. The time of bleeding was defined as the interval between initial transection and the visual cessation of bleeding. The results are shown in Table 5.

Table 5

	n=10	n=10	n=10
	Vehicle	1-01	1-02
Bleeding time (s)	66.6±19.4	31.0±3.6*	39.2±4.7
*p<0.05 vs vehicle			

30 [0188] Hyperfibrinolytic bleeding model, consisted in injection of 0.5 mg/kg tPA into the ocular plexus to prolong bleeding time due to excessive fibrinolysis. First, the femoral vein was exposed and cannulated with a saline-filled polyurethane catheter (Microcannula 72-9030, Harvard Apparatus) for agents administration. The catheter was connected to a syringe pump (AL-1000, WPI) for the infusion of 200 µL (10% bolus, 90% perfusion during 40 minutes) of tested agents. Then, tPA (0.5 mg/kg) was injected into the ocular plexus and five minutes after tPA administration, saline or the different compounds was infused through the femoral catheter to ensure systemic distribution of all the agents. Reference compounds, TXA and Aprotinin, were administered at 300 and 10 mg/Kg respectively; however, all compounds of the invention were administered at 1 mg/Kg. Five minutes later, 5 mm of tail tip were removed using a scalpel blade and the tail tip bathed in 1 mL of sterile saline at 37 °C.

35 [0189] The time of bleeding was defined as the interval between initial transections and the visual cessation of bleeding, that was measured up to 30 minutes. A value of 30 min was assigned to those animals bleeding longer than the observation period. Table 6 shows the results reporting bleeding time (BT); where BT ≥ 20 minutes (+), 10 minutes ≤ BT < 20 minutes (++) , 5 minutes ≤ BT < 10 minutes (+++) and BT < 5 minutes (++++), Bleeding time was determined in wild type mice (C57/Bl6), where n ≥ 10 per assayed compound; therefore, BT is reported as a mean value - in the case of saline, BT is reported as mean±ESM.

Table 6

Example	BT
Saline	28.9±0.7
TXA	++**

(continued)

Example	BT
Aprotinin	++**
1-03***	+++**†
1-07	++*
1-08	++++**††
1-14	++++**††
1-20	++*
1-26	++*
1-55	++++**††
1-62	+++**†
1-73	++++**††
1-79	++**
1-80	+++**†
1-82	+++**
1-96	+++**
1-99	++++**††
1-102	++*
1-107	+++**
1-108	++++**††
1-112	+++**
1-113	+++**
*p<0.05;**p<0.01 vs saline; *** compound administered at 10 mg/Kg; †p<0.05; ††p<0.01 vs TXA	

**[0190]** As shown table 6, tested compounds of the invention show a very significant reduction of the bleeding time when compared to the control or TXA. In all the cases the dose of tested compounds were lower than TXA or Aprotinin doses.

#### REFERENCES CITED IN THE APPLICATION

#### **[0191]**


Green and P. G. M. Wuts, Protective Groups in Organic Chemistry, Wiley, 3rd ed. 1999, Chapter 2, pp. 17-200 and Chapter 5, pp. 369-451.

D. Bouyssi et al., "Rearrangement of oxaspiroheptanes to cyclohexanones mediated by lithium iodide", Synlett 2000, vol. 5, pp. 749-751.

Osamu Kitagawa et al., "Stereoselective Iodine Atom Transfer [3 + 2] Cycloaddition Reaction with Alkenes Using Unsymmetrical Allylated Active Methine Radicals", The Journal of Organic Chemistry 2004, vol. 69, pp. 2607-2610.

#### **Claims**

1. A compound of formula (I), or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt



wherein

A and B form a spirocyclic ring system wherein the spiro atom connecting A and B is a carbon atom and wherein A is a known 3- to 8-membered carbocyclic or heterocyclic monocyclic ring, saturated or partially unsaturated; or alternatively

A is a known 6- to 18-membered carbocyclic or heterocyclic polycyclic ring system, saturated, partially unsaturated, or partially aromatic; and

B is a known 4- to 7-membered carbocyclic or heterocyclic monocyclic ring, saturated or partially unsaturated; C is phenyl or a known 5- to 6-membered heteroaromatic ring;

R<sub>1</sub>-R<sub>3</sub> are independently selected from H, halogen, -NO<sub>2</sub>, -CN, R<sup>a</sup>, -OR<sup>a</sup>, -OC(Y)R<sup>a</sup>, -OC(Y)OR<sup>a</sup>, -OC(Y)NR<sup>b</sup>R<sup>a</sup>, -OSO<sub>2</sub>OR<sup>a</sup>, -NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>C(Y)R<sup>a</sup>, -NR<sup>b</sup>C(Y)OR<sup>a</sup>, -NR<sup>b</sup>C(Y)NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>S(O)<sub>2</sub>R<sup>a</sup>, -NR<sup>b</sup>SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -SR<sup>a</sup>, -S(O)R<sup>a</sup>, -S(O)OR<sup>a</sup>, -SO<sub>2</sub>R<sup>a</sup>, -SO<sub>2</sub>(OR<sup>a</sup>), -SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -SC(Y)NR<sup>b</sup>R<sup>a</sup>, -C(Y)R<sup>a</sup>, -C(Y)OR<sup>a</sup>, -C(Y)NR<sup>b</sup>R<sup>a</sup>, -C(Y)NR<sup>b</sup>OR<sup>a</sup>, and -C(O)NR<sup>b</sup>SO<sub>2</sub>R<sup>a</sup>;

R<sub>4</sub>-R<sub>7</sub> are independently selected from halogen, -NO<sub>2</sub>, -CN, R<sup>c</sup>, -OR<sup>c</sup>, -NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>C(Y)R<sup>c</sup>, -NR<sup>d</sup>C(Y)OR<sup>c</sup>, -NR<sup>d</sup>C(Y)NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>S(O)<sub>2</sub>R<sup>c</sup>, -NR<sup>d</sup>SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>, -SR<sup>c</sup>, -S(O)R<sup>c</sup>, -S(O)OR<sup>c</sup>, -SO<sub>2</sub>R<sup>c</sup>, -SO<sub>2</sub>R(OR<sup>c</sup>), -SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>, -SC(Y)NR<sup>d</sup>R<sup>c</sup>, -C(Y)R<sup>c</sup>, -C(Y)OR<sup>c</sup>, -C(Y)NR<sup>d</sup>R<sup>c</sup>, -C(Y)NR<sup>d</sup>OR<sup>c</sup>, and -C(O)NR<sup>d</sup>SO<sub>2</sub>R<sup>c</sup>;

R<sup>a</sup> is a saturated or unsaturated (C<sub>1</sub>-C<sub>12</sub>)alkyl optionally substituted with one or more substituents R<sup>e</sup> and/or one Cy<sup>1</sup>; or alternatively R<sup>a</sup> is Cy<sup>2</sup>;

wherein Cy<sup>1</sup> and Cy<sup>2</sup> are independently optionally substituted with: one Cy<sup>3</sup> and/or one or more substituents R<sup>e</sup>, and/or one or more saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl groups optionally substituted with one or more substituents R<sup>e</sup> and/or one Cy<sup>3</sup>; and

wherein any Cy<sup>3</sup> is optionally substituted with one or more substituents independently selected from R<sup>e</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>e</sup>;

each R<sup>a</sup> and R<sup>b</sup> are independently H or R<sup>a</sup>;

R<sup>c</sup> and each R<sup>d</sup> are independently selected from H, Cy<sup>4</sup>, and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>h</sup> and/or one Cy<sup>5</sup>;

wherein Cy<sup>4</sup> is optionally substituted with one or more substituents independently selected from R<sup>h</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>h</sup>; and

wherein Cy<sup>5</sup> is optionally substituted with one or more substituents independently selected from R<sup>h</sup> and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>h</sup>;

each R<sup>e</sup> is independently selected from halogen, -NO<sub>2</sub>, -CN, -OR<sup>f</sup>, -OC(Y)R<sup>f</sup>, -OC(Y)OR<sup>f</sup>, -OC(Y)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>C(Y)R<sup>f</sup>, -NR<sup>g</sup>C(Y)OR<sup>f</sup>, -NR<sup>g</sup>C(Y)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>S(O)<sub>2</sub>R<sup>f</sup>, -NR<sup>g</sup>SO<sub>2</sub>NR<sup>g</sup>R<sup>f</sup>, -SR<sup>f</sup>, -S(O)R<sup>f</sup>, -S(O)OR<sup>f</sup>, -SO<sub>2</sub>R<sup>f</sup>, -SO<sub>2</sub>(OR<sup>f</sup>), -SO<sub>2</sub>NR<sup>g</sup>R<sup>f</sup>, -SC(Y)NR<sup>g</sup>R<sup>f</sup>, -C(Y)R<sup>f</sup>, -C(Y)OR<sup>f</sup>, -C(Y)NR<sup>g</sup>R<sup>f</sup>, -C(Y)NR<sup>g</sup>OR<sup>f</sup> and -C(O)NR<sup>g</sup>SO<sub>2</sub>R<sup>f</sup>;

R<sup>f</sup> and each R<sup>g</sup> are independently selected from H, Cy<sup>6</sup>, and saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one or more substituents R<sup>h</sup> and/or one Cy<sup>7</sup>;

wherein Cy<sup>6</sup> is optionally substituted with: one Cy<sup>7</sup>, and/or one or more substituents R<sup>h</sup>, and/or one or more saturated or unsaturated (C<sub>1</sub>-C<sub>6</sub>)alkyl groups optionally substituted with one or more substituents R<sup>h</sup> and/or one Cy<sup>7</sup>; and

wherein any Cy<sup>7</sup> is optionally substituted with one or more substituents independently selected from R<sup>h</sup> and (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or more substituents R<sup>h</sup>;

each R<sup>h</sup> is independently selected from halogen, -NO<sub>2</sub>, -CN, -OR<sup>i</sup>, -OC(O)R<sup>i</sup>, -OC(O)OR<sup>i</sup>, -OC(O)NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>C(O)R<sup>i</sup>, -NR<sup>i</sup>C(O)OR<sup>i</sup>, -NR<sup>i</sup>C(O)NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>S(O)R<sup>i</sup>, -NR<sup>i</sup>SO<sub>2</sub>NR<sup>i</sup>R<sup>i</sup>, -SR<sup>i</sup>, -S(O)R<sup>i</sup>, -SO<sub>2</sub>R<sup>i</sup>, -SO<sub>2</sub>(OR<sup>i</sup>), -SO<sub>2</sub>NR<sup>i</sup>R<sup>i</sup>, -C(O)R<sup>i</sup>, -C(O)OR<sup>i</sup>, -C(O)NR<sup>i</sup>R<sup>i</sup>, and -C(O)NR<sup>i</sup>OR<sup>i</sup>;

each  $R^i$  is independently H or  $-(C_1-C_4)$ alkyl optionally substituted with one or more halogen atoms;


Y is O, S, or  $NR^9$ ;


$Cy^1$ ,  $Cy^2$ ,  $Cy^4$  and  $Cy^6$  are independently a C or N-attached known ring system selected from 3- to 8-membered carbocyclic or heterocyclic monocyclic ring, saturated or partially unsaturated; phenyl; 5- or 6-membered heteroaromatic ring; and 6- to 18-membered carbocyclic or heterocyclic polycyclic ring system, saturated, partially unsaturated, aromatic or partially aromatic;

$Cy^3$ ,  $Cy^5$  and  $Cy^7$  are independently a C or N-attached known ring system selected from 3- to 8-membered carbocyclic or heterocyclic monocyclic ring, saturated or partially unsaturated; phenyl; and 5- or 6-membered heteroaromatic ring;

wherein in the carbocyclic rings all ring members are carbon atoms; and in the heterocyclic and heteroaromatic rings one or more ring members are selected from N, O, and S; and wherein in all saturated or partially unsaturated rings one or two members of the rings are optionally  $C(O)$  and/or  $C(NH)$  and/or  $C[N(C_1-C_4)alkyl]$ .

2. The compound of formula (I) according to claim 1, wherein A is a known 3-to 8-membered carbocyclic or heterocyclic monocyclic ring or a known 6- to 10-membered carbocyclic or heterocyclic bicyclic ring system.
3. The compound of formula (I) according to claim 2, wherein A is a monocyclic ring selected from a 3- to 6-membered carbocyclic ring and a 5- to 6-membered heterocyclic ring.
4. The compound of formula (I) according to any of the claims 1-3, wherein B is a 6- to 7-membered carbocyclic or heterocyclic monocyclic ring.
5. The compound of formula (I) according to any of the claims 1-4, wherein A and B form a spirocyclic ring system selected from the group consisting of:






6. The compound of formula (I) according to any of the claims 1-5, wherein C is phenyl.
7. The compound of formula (I) according to any of the claims 1-6, wherein in  $R_1$ - $R_3$ ,  $R^f$  and each  $R^g$  are independently selected from H and saturated or unsaturated ( $C_1$ - $C_6$ )alkyl optionally substituted with one or more fluorine atoms.
8. The compound of formula (I) according to any of the claims 1-7, wherein in  $R_1$ - $R_3$ ,  $Cy^1$  and  $Cy^2$  are independently optionally substituted with one or more substituents selected from  $R^e$  and saturated or unsaturated ( $C_1$ - $C_6$ )alkyl optionally substituted; and  $Cy^6$  is optionally substituted with one or more substituents independently selected from  $R^h$  and saturated or unsaturated ( $C_1$ - $C_6$ )alkyl optionally substituted.
9. The compound of formula (I) according to any of the claims 1-8, wherein  $R_1$ - $R_3$  are independently selected from H, halogen,  $-NO_2$ ,  $-CN$ ,  $R^a$ ,  $-OR^{a'}$ ,  $-OC(O)R^{a'}$ ,  $-OC(O)OR^{a'}$ ,  $-OC(O)NR^bR^{a'}$ ,  $-NR^bR^{a'}$ ,  $-NR^bC(O)R^{a'}$ ,  $-NR^bC(O)OR^{a'}$ ,  $-NR^bC(O)NR^bR^{a'}$ ,  $-NR^bS(O)_2R^{a'}$ ,  $-SR^{a'}$ ,  $-S(O)R^{a'}$ ,  $-SO_2R^{a'}$ ,  $-SO_2NR^bR^{a'}$ ,  $-C(O)R^{a'}$ ,  $-C(O)OR^{a'}$ ,  $-C(O)NR^bR^{a'}$ , and  $-C(O)NR^bOR^{a'}$ .
10. The compound of formula (I) according to any of the claims 1-9, wherein  $R_4$ - $R_7$  are independently selected from halogen,  $-NO_2$ ,  $-CN$ ,  $R^c$ ,  $-OR^c$ ,  $-NR^dR^c$ ,  $-NR^dC(O)R^c$ ,  $-NR^dC(O)OR^c$ ,  $-NR^dC(O)NR^dR^c$ ,  $-NR^dS(O)_2R^c$ ,  $-SR^c$ ,  $-S(O)R^c$ ,  $-SO_2R^c$ ,  $-SO_2NR^dR^c$ ,  $-C(O)R^c$ ,  $-C(O)OR^c$ , and  $-C(O)NR^dR^c$ .
11. The compound of formula (I) according to any of the claims 1-10, wherein  $R_2$  and  $R_3$  are independently selected from H, halogen,  $R^a$ ,  $-OR^{a'}$ , and  $-NR^bR^{a'}$ ; and  $R_5$ - $R_7$  are independently selected from H, halogen,  $R^c$ ,  $-OR^c$ , and  $-NR^dR^c$ , wherein  $R^a$ ,  $R^{a'}$ ,  $R^b$ ,  $R^c$  and  $R^d$  are independently selected from H and ( $C_1$ - $C_4$ )alkyl optionally substituted with one or more fluorine atoms.
12. A pharmaceutical or veterinary composition which comprises an effective amount of a compound of formula (I) as defined in any of the claims 1-11, or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt, together with one or more pharmaceutically or veterinary acceptable excipients or carriers.

13. A compound of formula (I) according to any of the claims 1-11, or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt, for use as a medicament.


14. A compound of formula (I) according to any of the claims 1-11, or a pharmaceutically or veterinary acceptable salt thereof, or any stereoisomer either of the compound of formula (I) or of its pharmaceutically or veterinary acceptable salt, for use as antifibrinolytic and antihemorrhagic agent.

15. A compound of formula (III)



(III)

wherein A, B, C, R<sub>1</sub>-R<sub>7</sub> are as defined in any of the claims 1-11, and R' is H or a carboxy protective group selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>)alkyl, benzyl, p-methoxyphenyl, trimethylsilyl and [2-(Trimethylsilyl)ethoxy]methyl (SEM), with the proviso that compound (III) is other than 7-methoxycarbonyl-7-phenylsulphonyl-2-oxaspiro[2.4]heptane and (2S\*,4R\*)-2-Phenylsulfonyl-4-iodomethyl-6,11-dioxaspiro[4.6]undecane-2-carboxylic acid methyl ester; or  
a compound of formula (II)



(II)

wherein A, B, C, R<sub>1</sub>-R<sub>7</sub> are as defined in any of the claims 1-11, and R is an hydroxamic acid protective group selected from the group consisting of tetrahydro-2H-pyran-2-yloxy (THP), benzyl, 1-naphthylmethyl and dimethyl-oxybenzyl (DMB).

### Patentansprüche


1. Eine Verbindung der Formel (I), oder ein pharmazeutisch oder tiermedizinisch akzeptables Salz davon, oder ein

Stereoisomer der Verbindung der Formel (I) oder von seinem pharmazeutisch oder tiermedizinisch akzeptablen Salz

5

10

15



(I)

wobei

20

A und B ein spirozyklisches Ringsystem bilden, wobei das Spiroatom, welches A und B verbindet, ein Kohlenstoffatom ist und wobei

A ein bekannter 3- bis 8-gliedriger carbocyclischer oder heterocyclischer monocyclischer Ring, gesättigt oder teilweise ungesättigt ist; oder, ersatzweise

A ein bekannter 6- bis 18-gliedriges carbocyclisches oder heterocyclisches polycyclisches Ringsystem, gesättigt, teilweise ungesättigt, oder teilweise aromatisch ist; und

25

B ein bekannter 4- bis 7-gliedriger carbocyclischer oder heterocyclischer monocyclischer Ring, gesättigt oder teilweise ungesättigt ist;

C Phenyl oder ein bekannter 5- bis 6-gliedriger heteroaromatischer Ring ist;

30

R<sub>1</sub>-R<sub>3</sub> unabhängig voneinander aus H, Halogen, -NO<sub>2</sub>, -CN, R<sup>a</sup>, -OR<sup>a</sup>, -OC(Y)R<sup>a</sup>, -OC(Y)OR<sup>a</sup>, -OC(Y)NR<sup>b</sup>R<sup>a</sup>, -OSO<sub>2</sub>OR<sup>a</sup>, -NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>C(Y)R<sup>a</sup>, -NR<sup>b</sup>C(Y)OR<sup>a</sup>, -NR<sup>b</sup>C(Y)NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>S(O)<sub>2</sub>R<sup>a</sup>, -NR<sup>b</sup>SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -SR<sup>a</sup>, -S(O)R<sup>a</sup>, -S(O)OR<sup>a</sup>, -SO<sub>2</sub>R<sup>a</sup>, -SO<sub>2</sub>(OR<sup>a</sup>), -SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -SC(Y)NR<sup>b</sup>R<sup>a</sup>, -C(Y)R<sup>a</sup>, -C(Y)OR<sup>a</sup>, -C(Y)NR<sup>b</sup>R<sup>a</sup>, -C(Y)NR<sup>b</sup>OR<sup>a</sup>, und -C(O)NR<sup>b</sup>SO<sub>2</sub>R<sup>a</sup> ausgewählt sind;

35

R<sub>4</sub>-R<sub>7</sub> unabhängig voneinander aus, -NO<sub>2</sub>, -CN, R<sup>c</sup>, -OR<sup>c</sup>, -NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>C(Y)R<sup>c</sup>, -NR<sup>d</sup>C(Y)OR<sup>c</sup>, -NR<sup>d</sup>C(Y)NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>S(O)<sub>2</sub>R<sup>c</sup>, -NR<sup>d</sup>SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>, -SR<sup>c</sup>, -S(O)R<sup>c</sup>, -S(O)OR<sup>c</sup>, -SO<sub>2</sub>R<sup>c</sup>, -SO<sub>2</sub>R(OR<sup>c</sup>), -SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>, -SC(Y)NR<sup>d</sup>R<sup>c</sup>, -C(Y)R<sup>c</sup>, -C(Y)OR<sup>c</sup>, -C(Y)NR<sup>d</sup>R<sup>c</sup>, -C(Y)NR<sup>d</sup>OR<sup>c</sup>, und -C(O)NR<sup>d</sup>SO<sub>2</sub>R<sup>c</sup> ausgewählt sind;

R<sup>a</sup> ein gesättigtes oder ungesättigtes (C<sub>1</sub>-C<sub>12</sub>)Alkyl ist, das wahlweise durch einen oder mehrere R<sup>e</sup> und/oder einen Cy<sup>1</sup> Substituenten substituiert ist; oder ersatzweise R<sup>a</sup> Cy<sup>2</sup> ist;

40

wobei Cy<sup>1</sup> und Cy<sup>2</sup> unabhängig voneinander wahlweise durch: einen Cy<sup>3</sup> und/oder einen oder mehrere R<sup>e</sup> Substituenten substituiert sind, und/oder durch einen oder mehrere gesättigte oder ungesättigte (C<sub>1</sub>-C<sub>6</sub>)Alkyl-Gruppen substituiert sind, die wahlweise durch einen oder mehrere R<sup>e</sup> und/oder einen Cy<sup>3</sup> Substituenten substituiert sind, und

wobei jedes Cy<sup>3</sup> wahlweise durch einen oder mehrere Substituenten substituiert ist, die unabhängig voneinander aus R<sup>e</sup> und gesättigtem oder ungesättigtem (C<sub>1</sub>-C<sub>6</sub>)Alkyl ausgewählt sind, das wahlweise durch einen oder mehrere R<sup>e</sup> Substituenten substituiert ist;

jedes R<sup>a</sup> und R<sup>b</sup> unabhängig voneinander H oder R<sup>a</sup> sind;

45

R<sup>c</sup> und jedes R<sup>d</sup> unabhängig voneinander ausgewählt aus H, Cy<sup>4</sup>, und gesättigtem oder ungesättigtem (C<sub>1</sub>-C<sub>6</sub>)Alkyl sind, das wahlweise durch einen oder mehrere R<sup>h</sup> Substituenten und/oder einen Cy<sup>5</sup> Substituenten substituiert ist;

wobei Cy<sup>4</sup> wahlweise durch einen oder mehrere Substituenten substituiert ist, die unabhängig voneinander aus R<sup>h</sup> und gesättigtem oder ungesättigtem (C<sub>1</sub>-C<sub>6</sub>)Alkyl ausgewählt sind, das wahlweise durch einen oder mehrere R<sup>h</sup> Substituenten substituiert ist; und

50

wobei Cy<sup>5</sup> wahlweise durch einen oder mehrere Substituenten substituiert ist, die unabhängig voneinander aus R<sup>h</sup> und gesättigtem oder ungesättigtem (C<sub>1</sub>-C<sub>6</sub>)Alkyl ausgewählt sind, das wahlweise durch einen oder mehrere R<sup>h</sup> Substituenten substituiert ist;

55

jedes R<sup>e</sup> unabhängig voneinander aus Halogen, -NO<sub>2</sub>, -CN, -OR<sup>f</sup>, -OC(Y)R<sup>f</sup>, -OC(Y)OR<sup>f</sup>, -OC(Y)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>C(Y)R<sup>f</sup>, -NR<sup>g</sup>C(Y)OR<sup>f</sup>, -NR<sup>g</sup>C(Y)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>S(O)<sub>2</sub>R<sup>f</sup>, -NR<sup>g</sup>SO<sub>2</sub>NR<sup>g</sup>R<sup>f</sup>, -SR<sup>f</sup>, -S(O)R<sup>f</sup>, -S(O)OR<sup>f</sup>, -SO<sub>2</sub>R<sup>f</sup>, -SO<sub>2</sub>(OR<sup>f</sup>), -SO<sub>2</sub>NR<sup>g</sup>R<sup>f</sup>, -SC(Y)NR<sup>g</sup>R<sup>f</sup>, -C(Y)R<sup>f</sup>, -C(Y)OR<sup>f</sup>, -C(Y)NR<sup>g</sup>R<sup>f</sup>, -C(Y)NR<sup>g</sup>OR<sup>f</sup> und -C(O)NR<sup>g</sup>SO<sub>2</sub>R<sup>f</sup> ausgewählt ist;

R<sup>f</sup> und jedes R<sup>g</sup> unabhängig voneinander aus H, Cy<sup>6</sup>, und gesättigtem oder ungesättigtem (C<sub>1</sub>-C<sub>6</sub>)Alkyl aus-

gewählt sind, das wahlweise durch einen oder mehrere  $R^h$  Substituenten und/oder einen  $Cy^7$  Substituenten substituiert ist;

wobei  $Cy^6$  wahlweise durch: einen  $Cy^7$  und/oder einen oder mehrere  $R^h$  Substituenten substituiert ist, und/oder durch einen oder mehrere gesättigte oder ungesättigte ( $C_1-C_6$ )Alkyl-Gruppen substituiert ist, die wahlweise durch einen oder mehrere  $R^h$  Substituenten und/oder einen  $Cy^7$  Substituenten substituiert sind, und wobei jedes  $Cy^7$  wahlweise durch einen oder mehrere Substituenten substituiert ist, die unabhängig voneinander aus  $R^h$  und ( $C_1-C_4$ )Alkyl ausgewählt sind, das wahlweise durch einen oder mehrere  $R^h$  Substituenten substituiert ist; jedes  $R^h$  unabhängig voneinander aus Halogen,  $-NO_2$ ,  $-CN$ ,  $-OR^i$ ,  $-OC(O)R^i$ ,  $-OC(O)OR^i$ ,  $-OC(O)NR^iR^i$ ,  $-NR^iR^i$ ,  $-NR^iC(O)R^i$ ,  $-NR^iC(O)OR^i$ ,  $-NR^iC(O)NR^iR^i$ ,  $-NR^iS(O)_2R^i$ ,  $-NR^iSO_2NR^iR^i$ ,  $-SR^i$ ,  $-S(O)R^i$ ,  $-SO_2R^i$ ,  $-SO_2(OR^i)$ ,  $-SO_2NR^iR^i$ ,  $-C(O)R^i$ ,  $-C(O)OR^i$ ,  $-C(O)NR^iR^i$ , und  $-C(O)NR^iOR^i$  ausgewählt ist;

jedes  $R^i$  unabhängig voneinander H oder  $-(C_1-C_4)$ Alkyl ist, das wahlweise durch ein oder mehrere Halogenatome substituiert ist;


Y O, S, oder  $NR^g$  ist;


$Cy^1$ ,  $Cy^2$ ,  $Cy^4$  und  $Cy^6$  unabhängig voneinander ein mit C- oder N-verbundenes bekanntes Ringsystem sind, das aus einem 3- bis 8-gliedrigen carbocyclischen oder heterocyclischen monocyclischen Ring, gesättigt oder teilweise ungesättigt; Phenyl; einem 5- oder 6-gliedrigen heteroaromatischen Ring; und einem 6- bis 18-gliedrigen carbocyclischen oder heterocyclischen polycyclischen Ringsystem, gesättigt, teilweise ungesättigt, aromatisch oder teilweise aromatisch ausgewählt ist;

$Cy^3$ ,  $Cy^5$ , und  $Cy^7$  unabhängig voneinander ein mit C- oder N-verbundenes bekanntes Ringsystem ist, das aus einem 3- bis 8-gliedrigen carbocyclischen oder heterocyclischen monocyclischen Ring, gesättigt oder teilweise ungesättigt; Phenyl; und einem 5- oder 6-gliedrigen heteroaromatischen Ring ausgewählt ist;

wobei in den carbocyclischen Ringen alle Ringglieder Kohlenstoffatome sind; und in den heterocyclischen und heteroaromatischen Ringen ein oder mehrere Ringglieder ausgewählt aus N, O, und S sind; und wobei in allen gesättigten oder teilweise ungesättigten Ringen ein oder zwei Glieder der Ringe wahlweise C(O) und/oder C(NH) und/oder C[N( $C_1-C_4$ )Alkyl] sind.

2. Die Verbindung der Formel (I) nach Anspruch 1, wobei A ein bekannter 3- bis 8-gliedriger carbocyclischer oder heterocyclischer monocyclischer Ring oder ein bekanntes 6- bis 10-gliedriges carbocyclisches oder heterocyclisches bicyclisches Ringsystem ist.
3. Die Verbindung der Formel (I) nach Anspruch 2, wobei A ein monocyclischer Ring ausgewählt aus einem 3- bis 6-gliedrigen carbocyclischen Ring und einem 5-bis 6-gliedrigen heterocyclischen Ring ist.
4. Die Verbindung der Formel (I) nach einem der Ansprüche 1-3, wobei B ein 6- bis 7-gliedriger carbocyclischer oder heterocyclischer monocyclischer Ring ist.
5. Die Verbindung der Formel (I) nach einem der Ansprüche 1-4, wobei A und B ein spirocyclisches Ringsystem bilden, das ausgewählt ist aus der Gruppe bestehend aus:





6. Die Verbindung der Formel (I) nach einem der Ansprüche 1-5, wobei C Phenyl ist.

7. Die Verbindung der Formel (I) nach einem der Ansprüche 1-6, wobei in  $R_1$ - $R_3$ ,  $R^f$  und jedes  $R^g$  unabhängig voneinander aus H und gesättigtem oder ungesättigtem ( $C_1$ - $C_6$ )Alkyl, wahlweise substituiert durch ein oder mehrere Fluoratome, ausgewählt sind.

8. Die Verbindung der Formel (I) nach einem der Ansprüche 1-7, wobei in  $R_1$ - $R_3$ ,  $Cy^1$  und  $Cy^2$  unabhängig voneinander wahlweise durch einen oder mehrere Substituenten substituiert sind, welche aus  $R^e$  und gesättigtem oder ungesättigtem ( $C_1$ - $C_6$ )Alkyl, wahlweise substituiert, ausgewählt sind; und  $Cy^6$  wahlweise durch einen oder mehreren Substituenten substituiert ist, welche unabhängig voneinander aus  $R^h$  und gesättigtem oder ungesättigtem ( $C_1$ - $C_6$ )Alkyl, wahlweise substituiert, ausgewählt sind.

9. Die Verbindung der Formel (I) nach einem der Ansprüche 1-8, wobei  $R_1$ - $R_3$  unabhängig voneinander aus H, Halogen,  $-NO_2$ ,  $-CN$ ,  $R^a$ ,  $-OR^a$ ,  $-OC(O)R^a$ ,  $-OC(O)OR^a$ ,  $-OC(O)NR^bR^a$ ,  $-NR^bR^a$ ,  $-NR^bC(O)R^a$ ,  $-NR^bC(O)OR^a$ ,  $-NR^bC(O)NR^bR^a$ ,  $-NR^bS(O)_2R^a$ ,  $-SR^a$ ,  $-S(O)R^a$ ,  $-SO_2R^a$ ,  $-SO_2NR^bR^a$ ,  $-C(O)R^a$ ,  $-C(O)OR^a$ ,  $-C(O)NR^bR^a$ , und  $-C(O)NR^bOR^a$  ausgewählt sind.

10. Die Verbindung der Formel (I) nach einem der Ansprüche 1-9, wobei  $R_4$ - $R_7$  unabhängig voneinander aus Halogen,  $-NO_2$ ,  $-CN$ ,  $R^c$ ,  $-OR^c$ ,  $-NR^dR^c$ ,  $-NR^dC(O)R^c$ ,  $-NR^dC(O)OR^c$ ,  $-NR^dC(O)NR^dR^c$ ,  $-NR^dS(O)_2R^c$ ,  $-SR^c$ ,  $-S(O)R^c$ ,  $-SO_2R^c$ ,  $-SO_2NR^dR^c$ ,  $-C(O)R^c$ ,  $-C(O)OR^c$ , und  $-C(O)NR^dR^c$  ausgewählt sind.


11. Die Verbindung der Formel (I) nach einem der Ansprüche 1-10, wobei  $R_2$  und  $R_3$  unabhängig voneinander ausgewählt aus H, Halogen,  $R^a$ ,  $-OR^{a'}$ , und  $-NR^bR^{a'}$  sind; und  $R_5$ - $R_7$  unabhängig voneinander ausgewählt aus H, Halogen,  $R^c$ ,  $-OR^c$ , und  $-NR^dR^c$  sind, wobei  $R^a$ ,  $R^{a'}$ ,  $R^b$ ,  $R^c$  und  $R^d$  unabhängig voneinander aus H und  $-(C_1-C_4)$ Alkyl, wahlweise durch ein oder mehrere Fluoratome substituiert, ausgewählt sind.

12. Eine pharmazeutische oder tiermedizinische Zusammensetzung, welche eine wirksame Menge von einer Verbindung der Formel (I) wie in einem der Ansprüche 1-11 definiert umfasst, oder ein pharmazeutisch oder tiermedizinisch akzeptables Salz davon, oder ein Stereoisomer der Verbindung der Formel (I) oder von seinem pharmazeutisch oder tiermedizinisch akzeptablen Salz, zusammen mit einem oder mehreren pharmazeutisch oder tiermedizinisch akzeptablen Hilfsstoffen oder Trägerstoffen.


13. Eine Verbindung der Formel (I) nach einem der Ansprüche 1-11, oder ein pharmazeutisch oder tiermedizinisch akzeptables Salz davon, oder ein Stereoisomer der Verbindung der Formel (I) oder von seinem pharmazeutisch oder tiermedizinisch akzeptablen Salz, zur Verwendung als Arzneimittel.

14. Eine Verbindung der Formel (I) nach einem der Ansprüche 1-11, oder ein pharmazeutisch oder tiermedizinisch akzeptables Salz davon, oder ein Stereoisomer der Verbindung der Formel (I) oder von seinem pharmazeutisch oder tiermedizinisch akzeptablen Salz, zur Verwendung als antifibrinolytisches Mittel und Hämostatikum.

15. Eine Verbindung der Formel (III)



wobei A, B, C,  $R_1$ - $R_7$  wie in einem der Ansprüche 1-11 definiert sind, und  $R'$  H oder eine Carboxy-schützende Gruppe ist, die aus der Gruppe bestehend aus  $(C_1-C_6)$ Alkyl, Benzyl, p-Methoxyphenyl, Trimethylsilyl und [2-(Trimethylsilyl)-ethoxy]methyl (SEM) ausgewählt ist, mit der Voraussetzung, dass die Verbindung (III) nicht 7-Methoxycarbonyl-7-phenylsulphonyl-2-oxaspiro[2.4]heptan oder Methylester aus  $(2S^*, 4R^*)$ -2-Phenylsulfonyl-4-iodomethyl-6,11-dioxaspiro[4.6]undecan-2-carbonsäure ist; oder eine Verbindung der Formel (III)




(II)

wobei A, B, C, R<sub>1</sub>-R<sub>7</sub> wie in einem der Ansprüche 1-11 definiert sind, und R eine Hydroxamsäure-schützende Gruppe ist, die aus der Gruppe bestehend aus Tetrahydro-2H-pyran-2-yloxy (THP), Benzyl, 1-Naphthylmethyl und Dimethylxybenzyl (DMB) ausgewählt ist.

### Revendications

1. Un composé de formule (I), ou un sel acceptable sur le plan pharmaceutique ou vétérinaire de celui-ci, ou un stéréoisomère quelconque du composé de formule (I) ou de son sel acceptable sur le plan pharmaceutique ou vétérinaire



(I)

dans laquelle

A et B forment un système annulaire spirocyclique dans lequel l'atome spiro reliant A et B est un atome de carbone et dans lequel

A est un anneau connu monocyclique carbocyclique ou hétérocyclique de 3 à 8 chaînons, saturé ou partiellement insaturé; ou, de façon alternative,

A est un système annulaire connu polycyclique carbocyclique ou hétérocyclique de 6 à 18 chaînons, saturé, partiellement insaturé, ou partiellement aromatique; et

B est un anneau connu monocyclique carbocyclique ou hétérocyclique de 4 à 7 chaînons, saturé ou partiellement insaturé;

C est phényle ou un anneau connu hétéroaromatique de 5 à 6 chaînons;

R<sub>1</sub>-R<sub>3</sub> sont indépendamment choisis parmi H, halogène, -NO<sub>2</sub>, -CN, R<sup>a</sup>, -OR<sup>a</sup>, -OC(Y)R<sup>a</sup>, -OC(Y)OR<sup>a</sup>, -OC(Y)NR<sup>b</sup>R<sup>a</sup>, -OSO<sub>2</sub>OR<sup>a</sup>, -NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>C(Y)R<sup>a</sup>, -NR<sup>b</sup>C(Y)OR<sup>a</sup>, -NR<sup>b</sup>C(Y)NR<sup>b</sup>R<sup>a</sup>, -NR<sup>b</sup>S(O)<sub>2</sub>R<sup>a</sup>, -NR<sup>b</sup>SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -SR<sup>a</sup>, -S(O)R<sup>a</sup>, -S(O)OR<sup>a</sup>, -SO<sub>2</sub>R<sup>a</sup>, -SO<sub>2</sub>(OR<sup>a</sup>), -SO<sub>2</sub>NR<sup>b</sup>R<sup>a</sup>, -SC(Y)NR<sup>b</sup>R<sup>a</sup>, -C(Y)R<sup>a</sup>, -C(Y)OR<sup>a</sup>, -C(Y)NR<sup>b</sup>R<sup>a</sup>, -C(Y)NR<sup>b</sup>OR<sup>a</sup>, et -C(O)NR<sup>b</sup>SO<sub>2</sub>R<sup>a</sup>;

R<sub>4</sub>-R<sub>7</sub> sont indépendamment choisis parmi halogène, -NO<sub>2</sub>, -CN, R<sup>c</sup>, -OR<sup>c</sup>, -NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>C(Y)R<sup>c</sup>, -NR<sup>d</sup>C(Y)OR<sup>c</sup>,

-NR<sup>d</sup>C(Y)NR<sup>d</sup>R<sup>c</sup>, -NR<sup>d</sup>S(O)<sub>2</sub>R<sup>c</sup>, -NR<sup>d</sup>SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>, -SR<sup>c</sup>, -S(O)R<sup>c</sup>, -S(O)OR<sup>c</sup>, -SO<sub>2</sub>R<sup>c</sup>, -SO<sub>2</sub>R(OR<sup>c</sup>), -SO<sub>2</sub>NR<sup>d</sup>R<sup>c</sup>,  
 -SC(Y)NR<sup>d</sup>R<sup>c</sup>, -C(Y)R<sup>c</sup>, -C(Y)OR<sup>c</sup>, -C(Y)NR<sup>d</sup>R<sup>c</sup>, -C(Y)NR<sup>d</sup>OR<sup>c</sup>,  
 et -C(O)NR<sup>d</sup>SO<sub>2</sub>R<sup>c</sup>;

R<sup>a</sup> est un (C<sub>1</sub>-C<sub>12</sub>)alkyle saturé ou insaturé facultativement substitué par un ou plusieurs substituants R<sup>e</sup> et/ou un Cy<sup>1</sup>; ou alternativement R<sup>a</sup> est Cy<sup>2</sup>;

dans lequel Cy<sup>1</sup> et Cy<sup>2</sup> sont indépendamment facultativement substitués par: un Cy<sup>3</sup> et/ou un ou plusieurs substituants R<sup>e</sup>, et/ou un ou plusieurs groupes (C<sub>1</sub>-C<sub>e</sub>)alkyle saturés ou insaturés facultativement substitués par un ou plusieurs substituants R<sup>e</sup> et/ou un Cy<sup>3</sup>; et

dans lequel un Cy<sup>3</sup> quelconque est facultativement substitué avec un ou plusieurs substituants indépendamment choisis parmi R<sup>e</sup> et (C<sub>1</sub>-C<sub>6</sub>)alkyle saturé ou insaturé facultativement substitué avec un ou plusieurs substituants R<sup>e</sup>;

chaque R<sup>a</sup> et R<sup>b</sup> sont indépendamment H ou R<sup>a</sup>;

R<sup>c</sup> et chaque R<sup>d</sup> sont indépendamment choisis parmi H, Cy<sup>4</sup>, et (C<sub>1</sub>-C<sub>6</sub>)alkyle saturé ou insaturé facultativement substitué par un ou plusieurs substituants R<sup>h</sup> et/ou un Cy<sup>5</sup>;

dans lequel Cy<sup>4</sup> est facultativement substitué par un ou plusieurs substituants indépendamment choisis parmi R<sup>h</sup> et (C<sub>1</sub>-C<sub>6</sub>)alkyle saturé ou insaturé facultativement substitué par un ou plusieurs substituants R<sup>h</sup>; et

dans lequel Cy<sup>5</sup> est facultativement substitué avec un ou plusieurs substituants indépendamment choisis parmi R<sup>h</sup> et (C<sub>1</sub>-C<sub>6</sub>)alkyle saturé ou insaturé facultativement substitué par un ou plusieurs substituants R<sup>h</sup>;

chaque R<sup>e</sup> est indépendamment choisi parmi halogène, -NO<sub>2</sub>, -CN, -OR<sup>f</sup>, -OC(Y)R<sup>f</sup>, -OC(Y)OR<sup>f</sup>, -OC(Y)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>C(Y)R<sup>f</sup>, -NR<sup>g</sup>C(Y)OR<sup>f</sup>, -NR<sup>g</sup>C(Y)NR<sup>g</sup>R<sup>f</sup>, -NR<sup>g</sup>S(O)<sub>2</sub>R<sup>f</sup>, -NR<sup>g</sup>SO<sub>2</sub>NR<sup>g</sup>R<sup>f</sup>, -SR<sup>f</sup>, -S(O)R<sup>f</sup>, -S(O)OR<sup>f</sup>, -SO<sub>2</sub>R<sup>f</sup>, -SO<sub>2</sub>(OR<sup>f</sup>), -SO<sub>2</sub>NR<sup>g</sup>R<sup>f</sup>, -SC(Y)NR<sup>g</sup>R<sup>f</sup>, -C(Y)R<sup>f</sup>, -C(Y)OR<sup>f</sup>, -C(Y)NR<sup>g</sup>R<sup>f</sup>, -C(Y)NR<sup>g</sup>OR<sup>f</sup> et -C(O)NR<sup>g</sup>SO<sub>2</sub>R<sup>f</sup>;

R<sup>f</sup> et chaque R<sup>g</sup> sont indépendamment choisis parmi H, Cy<sup>6</sup>, et (C<sub>1</sub>-C<sub>6</sub>)alkyle saturé ou insaturé facultativement substitué par un ou plusieurs substituants R<sup>h</sup> et/ou un Cy<sup>7</sup>;

dans lequel Cy<sup>6</sup> est facultativement substitué par: un Cy<sup>7</sup> et/ou un ou plusieurs substituants R<sup>h</sup>, et/ou un ou plusieurs groupes (C<sub>1</sub>-C<sub>6</sub>)alkyle saturés ou insaturés facultativement substitués par un ou plusieurs substituants R<sup>h</sup> et/ou un Cy<sup>7</sup>; et dans lequel Cy<sup>7</sup> est facultativement substitué par un ou plusieurs substituants indépendamment choisis parmi R<sup>h</sup> et (C<sub>1</sub>-C<sub>4</sub>)alkyle facultativement substitué par un ou plusieurs substituants R<sup>h</sup>;

chaque R<sup>h</sup> est indépendamment choisi parmi halogène, -NO<sub>2</sub>, -CN, -OR<sup>i</sup>, -OC(O)R<sup>i</sup>, -OC(O)OR<sup>i</sup>, -OC(O)NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>C(O)R<sup>i</sup>, -NR<sup>i</sup>C(O)OR<sup>i</sup>, -NR<sup>i</sup>C(O)NR<sup>i</sup>R<sup>i</sup>, -NR<sup>i</sup>S(O)<sub>2</sub>R<sup>i</sup>, -NR<sup>i</sup>SO<sub>2</sub>NR<sup>i</sup>R<sup>i</sup>, -SR<sup>i</sup>, -S(O)R<sup>i</sup>, -SO<sub>2</sub>R<sup>i</sup>, -SO<sub>2</sub>(OR<sup>i</sup>), -SO<sub>2</sub>NR<sup>i</sup>R<sup>i</sup>, -C(O)R<sup>i</sup>, -C(O)OR<sup>i</sup>, -C(O)NR<sup>i</sup>R<sup>i</sup>, et -C(O)NR<sup>i</sup>OR<sup>i</sup>;


chaque R<sup>i</sup> est indépendamment H ou (C<sub>1</sub>-C<sub>4</sub>)alkyle facultativement substitué par un ou plusieurs atomes d'halogène;

Y est O, S, ou NR<sup>g</sup>;

Cy<sup>1</sup>, Cy<sup>2</sup>, Cy<sup>4</sup> et Cy<sup>6</sup> sont indépendamment un système annulaire connu attaché à C ou à N choisi parmi un anneau monocyclique carbocyclique ou hétérocyclique de 3 à 8 chaînons, saturé ou partiellement insaturé; phényle; un anneau hétéroaromatique de 5 ou 6 chaînons; et un système annulaire polycyclique carbocyclique ou hétérocyclique de 6 à 18 chaînons, saturé, partiellement insaturé, aromatique ou partiellement aromatique; Cy<sup>3</sup>, Cy<sup>5</sup>, et Cy<sup>7</sup> sont indépendamment un système annulaire connu attaché à C ou à N choisi parmi un anneau monocyclique carbocyclique ou hétérocyclique de 3 à 8 chaînons, saturé ou partiellement insaturé; phényle; et un anneau hétéroaromatique de 5 ou 6 chaînons;

dans lequel dans les anneaux carbocycliques tous les chaînons des anneaux sont des atomes de carbone; et dans les anneaux hétérocycliques et hétéroaromatiques un ou plusieurs chaînons des anneaux sont choisis parmi N, O, et S; et dans lequel dans tous les anneaux saturés ou partiellement insaturés un ou deux chaînons des anneaux sont facultativement C(O) et/ou C(NH) et/ou alkyle en C[N(C<sub>1</sub>-C<sub>4</sub>)].

2. Le composé de formule (I) selon la revendication 1, dans lequel A est un anneau connu monocyclique carbocyclique ou hétérocyclique de 3 à 8 chaînons ou un système annulaire connu bicyclique carbocyclique ou hétérocyclique de 6 à 10 chaînons.
3. Le composé de formule (I) selon la revendication 2, dans lequel A est un anneau monocyclique choisi parmi un anneau carbocyclique de 3 à 6 chaînons et un anneau hétérocyclique de 5 à 6 chaînons.
4. Le composé de formule (I) selon l'une quelconque des revendications 1-3, dans lequel B est un anneau monocyclique carbocyclique ou hétérocyclique de 6 à 7 chaînons.
5. Le composé de formule (I) selon l'une quelconque des revendications 1-4, dans lequel A et B forment un système annulaire spirocyclique choisi dans le groupe constitué de :




6. Le composé de formule (I) selon l'une quelconque des revendications 1-5, dans lequel C est phényle.

50 7. Le composé de formule (I) selon l'une quelconque des revendications 1-6, dans lequel dans  $R_1$ - $R_3$ ,  $R^f$  et chaque  $R^g$  sont indépendamment choisis parmi H et ( $C_1$ - $C_6$ )alkyle saturé ou insaturé facultativement substitué par un ou plusieurs atomes de fluor.


55 8. Le composé de formule (I) selon l'une quelconque des revendications 1-7, dans lequel dans  $R_1$ - $R_3$ ,  $Cy^1$  et  $Cy^2$  sont indépendamment facultativement substitués par un ou plusieurs substituants choisis parmi  $R^e$  et ( $C_1$ - $C_6$ )alkyle saturé ou insaturé facultativement substitué; et  $Cy^6$  est facultativement substitué par un ou plusieurs substituants indépendamment choisis parmi  $R^h$  et ( $C_1$ - $C_6$ )alkyle saturé ou insaturé facultativement substitué.

9. Le composé de formule (I) selon l'une quelconque des revendications 1-8, dans lequel  $R_1$ - $R_3$  sont indépendamment choisis parmi H, halogène,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $R^a$ ,  $-\text{OR}^a$ ,  $-\text{OC}(\text{O})\text{R}^a$ ,  $-\text{OC}(\text{O})\text{OR}^a$ ,  $-\text{OC}(\text{O})\text{NR}^b\text{R}^a$ ,  $-\text{NR}^b\text{R}^a$ ,  $-\text{NR}^b\text{C}(\text{O})\text{R}^a$ ,  $-\text{NR}^b\text{C}(\text{O})\text{OR}^a$ ,  $-\text{NR}^b\text{C}(\text{O})\text{NR}^b\text{R}^a$ ,  $-\text{NR}^b\text{S}(\text{O})_2\text{R}^a$ ,  $-\text{SR}^a$ ,  $-\text{S}(\text{O})\text{R}^a$ ,  $-\text{SO}_2\text{R}^a$ ,  $-\text{SO}_2\text{NR}^b\text{R}^a$ ,  $-\text{C}(\text{O})\text{R}^a$ ,  $-\text{C}(\text{O})\text{OR}^a$ ,  $-\text{C}(\text{O})\text{NR}^b\text{R}^a$ , et  $-\text{C}(\text{O})\text{NR}^b\text{OR}^a$ .
10. Le composé de formule (I) selon l'une quelconque des revendications 1-9, dans lequel  $R_4$ - $R_7$  sont indépendamment choisis parmi halogène,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $R^c$ ,  $-\text{OR}^c$ ,  $-\text{NR}^d\text{R}^c$ ,  $-\text{NR}^d\text{C}(\text{O})\text{R}^c$ ,  $-\text{NR}^d\text{C}(\text{O})\text{OR}^c$ ,  $-\text{NR}^d\text{C}(\text{O})\text{NR}^d\text{R}^c$ ,  $-\text{NR}^d\text{S}(\text{O})_2\text{R}^c$ ,  $-\text{SR}^c$ ,  $-\text{S}(\text{O})\text{R}^c$ ,  $-\text{SO}_2\text{R}^c$ ,  $-\text{SO}_2\text{NR}^d\text{R}^c$ ,  $-\text{C}(\text{O})\text{R}^c$ ,  $-\text{C}(\text{O})\text{OR}^c$ , et  $-\text{C}(\text{O})\text{NR}^d\text{R}^c$ .
11. Le composé de formule (I) selon l'une quelconque des revendications 1-10, dans lequel  $R_2$  et  $R_3$  sont indépendamment choisis parmi H, halogène,  $R^a$ ,  $-\text{OR}^a$ , et  $-\text{NR}^b\text{R}^a$ ; et  $R_5$ - $R_7$  sont indépendamment choisis parmi H, halogène,  $R^c$ ,  $-\text{OR}^c$ , et  $-\text{NR}^d\text{R}^c$ , dans lequel  $R^a$ ,  $R^a$ ,  $R^b$ ,  $R^c$  et  $R^d$  sont indépendamment choisis parmi H et  $(\text{C}_1\text{-C}_4)$ alkyle facultativement substitué par un ou plusieurs atomes de fluor.
12. Une composition pharmaceutique ou vétérinaire qui comprend une quantité efficace d'un composé de formule (I) tel que définie dans l'une quelconque des revendications 1-11, ou un sel acceptable sur le plan pharmaceutique ou vétérinaire de celui-ci, ou un stéréoisomère quelconque du composé de formule (I) ou de son sel acceptable sur le plan pharmaceutique ou vétérinaire, conjointement avec un ou plusieurs excipients ou véhicules acceptables sur le plan pharmaceutique ou vétérinaire.
13. Un composé de formule (I) selon l'une quelconque des revendications 1-11, ou un sel acceptable sur le plan pharmaceutique ou vétérinaire de celui-ci, ou un stéréoisomère quelconque du composé de formule (I) ou de son sel acceptable sur le plan pharmaceutique ou vétérinaire, pour l'utilisation comme médicament.
14. Un composé de formule (I) selon l'une quelconque des revendications 1-11, ou un sel acceptable sur le plan pharmaceutique ou vétérinaire de celui-ci, ou un stéréoisomère quelconque du composé de formule (I) ou de son sel acceptable sur le plan pharmaceutique ou vétérinaire, pour l'utilisation comme agent antifibrinolytique et anti-hémorragique.
15. Un composé de formule (III)



(III)

dans lequel A, B, C,  $R_1$ - $R_7$  sont tels que définis dans l'une quelconque des revendications 1-11, et  $R'$  est H ou un groupe protecteur de carboxyle choisi dans le groupe constitué de  $(\text{C}_1\text{-C}_6)$ alkyle, benzyle, p-méthoxyphényle, triméthylsilyle et [2-(triméthylsilyl)éthoxy]méthyle (SEM), avec la condition que le composé (III) ne soit pas le 7-méthoxycarbonyl-7-phénylsulfonyl-2-oxaspiro[2.4]héptane ni le méthylester de l'acide (2*S*\*,4*R*\*)-2-phénylsulfonyl-4-iodométhyl-6,11-dioxaspiro[4.6]undécane-2-carboxylique; ou un composé de formule (II)



(II)

dans laquelle A, B, C, R<sub>1</sub>-R<sub>7</sub> sont tels que définis dans l'une quelconque des revendications 1-11, et R est un groupe protecteur d'acide hydroxamique choisi dans le groupe constitué du tétrahydro-2H-pyran-2-yloxy (THP), du benzyle, du 1-naphthylméthyle et du diméthoxybenzyle (DMB).

**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Patent documents cited in the description**

- US 3868411 A [0008]
- US 4483867 A [0009]
- US 2002099035 A1 [0010]

**Non-patent literature cited in the description**

- **T. W. GREEN ; P. G. M. WUTS.** Protective Groups in Organic Chemistry. Wiley, 1999, 17-200 [0062]
- **T. W. GREEN ; P. G. M. WUTS.** Protective Groups in Organic Chemistry. Wiley, 1999, 369-451 [0064]
- **D. BOUYSSI et al.** Rearrangement of oxaspiroheptanes to cyclohexanones mediated by lithium iodide. *Synlett*, 2000, vol. 5, 749-751 [0065] [0191]
- **OSAMU KITAGAWA et al.** Stereoselective Iodine Atom Transfer [3 + 2] Cycloaddition Reaction with Alkenes Using Unsymmetrical Allylated Active Methine Radicals. *The Journal of Organic Chemistry*, 2004, vol. 69, 2607-2610 [0065] [0191]
- **GREEN ; P. G. M. WUTS.** Protective Groups in Organic Chemistry. Wiley, 1999, 17-200 [0191]
- PROTECTIVE GROUPS IN ORGANIC CHEMISTRY. vol. 5, 369-451 [0191]