



(11) **EP 2 881 146 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
10.06.2015 Bulletin 2015/24

(51) Int Cl.:
A61Q 5/10 (2006.01) **A45D 19/02** (2006.01)
A61K 8/22 (2006.01) **A45D 19/00** (2006.01)

(21) Application number: **14181705.6**

(22) Date of filing: **21.08.2014**

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME

(72) Inventor: **Schofield, Stephen**
Egham, Surrey TW20 9NW (GB)

(74) Representative: **Boubel, Thomas**
Procter & Gamble Service GmbH
IP Department
Frankfurter Strasse 145
61476 Kronberg im Taunus (DE)

(30) Priority: **23.08.2013 EP 13181445**

(71) Applicant: **The Procter & Gamble Company**
Cincinnati, OH 45202 (US)

(54) **Hair colouring method**

(57) Method for colouring hair wherein a first and a second hair colouring composition are applied to the hair, wherein the concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents and/or

the concentration of the oxidizing agents comprised in the second hair colouring composition is lower than in the first hair colouring composition.

EP 2 881 146 A2

Description

FIELD OF THE INVENTION

[0001] The invention relates to a method for colouring hair.

BACKGROUND OF THE INVENTION

[0002] Consumers desiring to colour their hair typically have two options available namely to use a commercially available retail product or kit or use the services of a professional salon. The latter whilst providing a highly desirable colour outcome, is considerably more expensive than the retail option and thus not available to many consumers particularly those who colour regularly.

[0003] For consumers who have previously coloured their hair, the colour and condition of the hair is not homogenous along the entire length. The hair strands will comprise root virgin hair or new growth hair which has not been previously coloured and conversely at the tips hair which has experienced one or multiple hair colouring treatments. The tips of the hair are typically the most damaged portions of the hair and are characterised by an overly deposition of dyes or over-bleaching due to previous oxidative hair colorations. The intermediate hair lengths are typically a medium between these two extreme conditions. However, the problem with current retail hair colour products is that they do not take into account the differences of properties between the different portions of the hair. It is rather challenging even for experienced home colour users to control the dye deposition or the bleaching provided by a retail hair colour product in order to not overly deposit hair dyes on or over-bleaching hair lengths and tips since the instructions provided in hair colour retail kits are typically imprecise, often not followed and the results achieved are not comparable with those from a salon stylist. It is known that different hair colouring compositions may be applied on the hair roots vs. the hair lengths and tips. However, the problem with such compositions is that they do not take into account the differences of properties between the intermediate hair lengths and the tips and therefore the final colour of the hair tips may be darker than the final colour of the intermediate hair lengths which therefore differs from the colour of natural hair where the tips are usually lighter than the rest of the hair.

[0004] Therefore, there is a need to provide a simple method as well as a simple retail kit for colouring or bleaching hair which gives the user the possibility of obtaining a different hair colour or bleaching result on different portions of the hair, for example a different colour result on hair roots vs. hair lengths and tips and a more natural overall colour result.

SUMMARY OF THE INVENTION

[0005] The present invention relates to a method for colouring hair comprising the steps of:

- i) mixing a first dye composition with a first developer composition to obtain a first hair colouring composition, wherein the first dye composition comprises one or more oxidative dye precursors and/or one or more alkalizing agents and the first developer composition comprises one or more oxidizing agents;
- ii) applying a portion or all of the first hair colouring composition to the hair roots and to one or more portions of the hair lengths and tips extending from the hair roots;
- iii) mixing a second dye composition with a second developer composition to obtain a second hair colouring composition, wherein the second dye composition comprises one or more oxidative dye precursors and/or one or more alkalizing agents and the second developer composition comprises one or more oxidizing agents, wherein the concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents and/or the concentration of the oxidizing agents comprised in the second hair colouring composition is lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents and/or the oxidizing agents comprised in the first hair colouring composition or providing a second hair colouring composition comprising one or more direct dyes;
- iv) applying a portion or all of the second hair colouring composition to the hair;
- v) optionally rinsing the hair.

DETAILED DESCRIPTION OF THE INVENTION

[0006] While the specification concludes with claims, which particularly point out and distinctly claim the invention, it is believed the present invention will be better understood from the following description.

[0007] As used herein the term "hair" to be treated may be "living" i.e. on a living body or may be "non-living" i.e. in a wig, hairpiece or other aggregation of non-living keratinous fibres. Mammalian, preferably human hair is preferred. However wool, fur and other keratin containing fibres are suitable substrates for the compositions according to the

present invention. The terms "root", "hair roots", "root hair line" and "virgin hair" all refer to hair which has not been previously treated with a hair colouring or bleaching composition.

[0008] By "hair colouring" composition it is meant a composition suitable for changing the color of hair. The hair colouring composition can comprise oxidative dye precursors, direct dyes or even no, or substantially no, dyes in case of bleaching only compositions where the change of colour is mainly caused by the degradation of the natural melanin contained in the hair shaft by the oxidizing agent. The terms "hair colouring composition" and "method for colouring hair" as used herein also encompasses respectively the terms "hair bleaching composition" and "method for bleaching hair".

[0009] In the preferred embodiment according to the present invention, the hair colouring compositions are applied to hair which has already been previously coloured with hair colouring compositions. In such a case, the terms "root", "hair roots", "root hair line" and "virgin hair" all refer to the portion of hair having grown, since the last hair colouration, said portion of hair being virgin, i.e. naturally-coloured and the terms "hair lengths and tips" refer to the remaining portion of hair having been already previously coloured.

[0010] All percentages are by weight of the total composition unless specifically stated otherwise. All ratios are weight ratios unless specifically stated otherwise.

Method for colouring hair

[0011] The present invention relates to a method for colouring hair as stated hereinbefore.

[0012] The concentration of the oxidative dye precursors comprised in the second hair colouring composition is lower than the concentration of the oxidative dye precursors comprised in the first hair colouring composition and/or the concentration of the alkalizing agents comprised in the second hair colouring composition is lower than the concentration of the alkalizing agents comprised in the first hair colouring composition and/or the concentration of the oxidizing agents comprised in the second hair colouring composition is lower than the concentration of the oxidizing agents comprised in the first hair colouring composition.

[0013] Whilst not wishing to be bound by theory it is believed that by having the concentration of active species such as dye precursors, oxidizing agents such as hydrogen peroxide and alkalizing agents (if present) in the second composition which is lower than in the first composition or wherein the second composition comprises direct dyes, it is possible to obtain a second composition having a lower overall effective lightening and colouring capacity than the first composition which can then be applied to the remaining pre-coloured hair length which requires less lightening and dye deposition. Furthermore, by applying the first composition not only to the hair roots but also to one or more portions of the hair lengths and tips extending from the hair roots, the intermediate hair lengths between the hair roots and the hair tips may be provided with a darker colour than the colour of the hair tips. The overall colour result may therefore be closer to what is observed on natural hair where the hair tips are usually lighter than the rest of the hair.

[0014] In step iv) of the method according to the present invention, the portion or all of the second hair colouring composition may be applied to the rest of the hair lengths and tips which are free of the first hair colouring composition and optionally to one or more portions of the hair to which the first hair colouring composition has been applied in step ii).

[0015] The concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents comprised in the second dye composition may be lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents comprised in the first dye composition. In this case, the oxidative dye precursors and/or the alkalizing agents comprised in the first and second dye compositions may be the same.

[0016] Alternatively, the concentration of oxidizing agents comprised in the second developer composition may be lower than the concentration of oxidizing agents comprised in the first developer composition. In this case, the oxidizing agents comprised in the first and second dye compositions may be the same.

[0017] The first and/or second developer compositions may comprise from 1% to 20% by weight of the oxidizing agents.

[0018] The first and/or second dye compositions comprise from 0.005% to 10% by weight of the oxidative dye precursors and/or from 0.1% to 20% or from 0.1% to 10% by weight of the alkalizing agents.

[0019] The first dye composition may be mixed with the first developer composition at a mixing ratio of from 2:1 to 1:3, or from 1:1 to 1:2. The second dye composition may be mixed with the second developer composition at a mixing ratio of from 2:1 to 1:3, or from 1:1 to 1:2. The mixing ratios of the first dye composition with the first developer composition and of the second dye composition with the second developer composition may be the same. Having the same mixing ratio to obtain the first and the second compositions may be easier for the user to avoid any mistake concerning the mixing ratio when the product is applied on hair.

[0020] The first and second developer compositions may be the same and the concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents comprised in the second dye composition may be lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents comprised in the first dye composition. The oxidative dye precursors and/or the alkalizing agents comprised in the first and second dye compositions may be the same. The concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents comprised in the second dye composition may be at least 1% or at least 5% or from 1% to 12%, or

from 5% to 12% lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents comprised in the first dye composition.

[0021] Alternatively, the first and second dye compositions may be the same and the concentration of oxidizing agents comprised in the second developer composition may be lower than the concentration of oxidizing agents comprised in the first developer composition. The oxidizing agents comprised in the first and second developer compositions may be the same. The concentration of the oxidizing agents comprised in the second developer composition may be at least 15% or at least 25% or from 15% to 85 %, or from 25% to 75% lower than the concentration of the oxidizing agents comprised in the first developer composition.

[0022] In some embodiments, the method may further comprise the step of waiting for a time period of from 10 min to 40 min, or from 15 min to 30 min or from 20 min, which is performed between steps ii) and iv) and the step of waiting for a time period of from 5 min to 20 min or 10 min which is performed between steps iv) and v). These embodiments may be used for colouring hair with non-tonal hair shades such as blonde, brown or black hair shades. In these embodiments, step iii) is typically performed during this time period. It may be performed immediately after step ii) or immediately prior to step iv).

[0023] Alternatively, in some embodiments, step iii) may be performed immediately after step ii) and step iv) may be performed immediately after step iii) and the method further comprises the step of waiting for a time period of from 5 min to 40 min, or from 20 min to 40 min, or of 30 min, which is performed between steps iv) and v). These embodiments may be used for colouring hair with more vibrant, tonal shades such as red, violet, copper.

[0024] In step v) of the method, the hair may be rinsed with water and/or shampoo. After rinsing, they may be further dried and styled as usual. A conditioner composition may be applied to the hair after rinsing, preferably prior to drying and styling.

[0025] The volume of the first and second dye compositions may be from 10 mL to 120 mL, or from 40 mL to 70 mL, or from 55 mL to 65 mL. The volume of the first and second developer compositions may be from 10 mL to 120 mL, or from 40 mL to 70 mL, or from 55 mL to 65 mL.

[0026] The present description is not limited to the embodiments wherein only two different hair colouring compositions having two different concentrations of oxidative dye precursors and/or alkalizing agents and/or oxidizing agents are applied to the hair. It may be envisaged to apply more than two different hair colouring compositions to the hair wherein each time the concentration of oxidative dye precursors and/or alkalizing agents and/or oxidizing agents of the next composition is lower than the one of the previous composition.

Hair colouring compositions

Other ingredients

[0027] The first and/or second hair colouring compositions of the present invention may comprise, in addition to the ingredients indicated above, further ingredients in order to further enhance the properties of the compositions, as long as these are not excluded by the claims.

[0028] Suitable further ingredients include, but not limited to: solvents; oxidizing agents; alkalizing agents; oxidative dye precursors, direct dyes; chelants; radical scavengers; pH modifiers and buffering agents; thickeners and/or rheology modifiers; carbonate ion sources; peroxydicarbonate ion sources; anionic, cationic, nonionic, amphoteric or zwitterionic surfactants, and mixtures thereof; anionic, cationic, nonionic, amphoteric or zwitterionic polymers, and mixtures thereof; fragrances; enzymes; dispersing agents; peroxide stabilizing agents; antioxidants; natural ingredients (such as proteins, protein compounds, and plant extracts); conditioning agents (such as silicones and cationic polymers); ceramides; preserving agents; opacifiers and pearling agents (such as titanium dioxide and mica); and mixtures thereof. Suitable further ingredients referred to above, but not specifically described below, are listed in the International Cosmetics Ingredient Dictionary and Handbook, (8th ed.; The Cosmetics, Toiletry, and Fragrance Association). Particularly, vol. 2, sections 3 (Chemical Classes) and 4 (Functions), which are useful in identifying specific adjuvants to achieve a particular purpose or multipurpose. A few of these ingredients are discussed hereinbelow, whose disclosure is of course non-exhaustive.

Solvents

[0029] The first and/or the second hair colouring compositions according to the present invention may further comprise a solvent. The solvent may be selected from water, or a mixture of water and at least one organic solvent to dissolve the compounds that would not typically be sufficiently soluble in water.

[0030] Suitable organic solvents include, but are not limited to: C1 to C4 lower alkanols (such as ethanol, propanol, isopropanol); aromatic alcohols (such as benzyl alcohol and phenoxyethanol); polyols and polyol ethers (such as carbitols, 2-butoxyethanol, propylene glycol, propylene glycol monomethyl ether, diethylene glycol monoethyl ether, monomethyl

ether, hexylene glycol, glycerol, ethoxy glycol, butoxydiglycol, ethoxydiglycerol, dipropyleneglocol, polyglycerol); propylene carbonate; and mixtures thereof.

[0031] The solvent may be selected from the group consisting of water, ethanol, propanol, isopropanol, glycerol, 1,2-propylene glycol, hexylene glycol, ethoxy diglycol, and mixtures thereof.

[0032] Typically, the first and/or the second hair colouring compositions may comprise water as a main ingredient, particularly in a total amount ranging from at least 50%, alternatively from at least 60%, alternatively from at least 70%, by weight of the total composition. Typically, when present, the first and/or the second hair colouring compositions may comprise a total amount of organic solvents ranging from 1% to 30%, by weight of the total composition.

Oxidizing agents

[0033] The first and/or the second hair colouring compositions according to the present invention may further comprise at least one source of an oxidizing agent. Any oxidizing agent known in the art may be used. Preferred oxidizing agents are water-soluble peroxygen oxidizing agents. As used herein, "water-soluble" means that in standard conditions at least 0.1g, preferably 1g, more preferably 10g of the oxidizing agent can be dissolved in 1 liter of deionized water at 25°C. The oxidizing agents are valuable for the initial solubilisation and decolorisation of the melanin (bleaching) and accelerate the oxidation of the oxidative dye precursors (oxidative dyeing) in the hair shaft.

[0034] Typically, the first and/or the second hair colouring compositions may comprise a total amount of oxidizing agents ranging from 0.1% to 10%, alternatively from 1% to 7%, alternatively from 2% to 5%, by weight of the total composition.

[0035] Suitable water-soluble oxidizing agents include, but are not limited to: inorganic peroxygen materials capable of yielding hydrogen peroxide in an aqueous solution.

[0036] Suitable water-soluble peroxygen oxidizing agents include, but are not limited to: hydrogen peroxide; inorganic alkali metal peroxides (such as sodium periodate and sodium peroxide); organic peroxides (such as urea peroxide and melamine peroxide); inorganic perhydrate salt bleaching compounds (such as the alkali metal salts of perborates, percarbonates, perphosphates, persilicates, persulphates and the like); and mixtures thereof. Inorganic perhydrate salts may be incorporated for example as monohydrates, tetrahydrates. Alkyl/aryl peroxides and/or peroxidases may also be used. Mixtures of two or more such oxidizing agents can be used if desired. The oxidizing agents may be provided in aqueous solution or as a powder which is dissolved prior to use.

[0037] The first and/or the second hair colouring compositions may comprise a water-soluble oxidizing agent selected from the group consisting of hydrogen peroxide, percarbonates (which may be used to provide a source of both oxidizing agent and carbonate ions and or ammonium ions), persulphates, and mixtures thereof.

[0038] When the first and/or the second hair colouring compositions of the present invention are obtained by mixing a developer composition and a tint composition prior to use, the oxidizing agent may be present in the developer composition. The developer composition may be based on any desired formulation chassis, including any commercial product, for example an oil-in-water emulsion. Typical developer compositions comprise 6% or 9% of the H₂O₂ relative to the total weight of the developer composition. A commercial example is the Welloxon® Emulsion with respectively 6% and 9% H₂O₂, marketed by Wella and comprising as INCI ingredients: Water, H₂O₂, Cetearyl Alcohol, Cetareth-25, Salicylic Acid, Phosphoric Acid, Disodium Phosphate, Etidronic Acid.

Alkalizing agents

[0039] The first and/or the second hair colouring compositions according to the present invention may further comprise one or more alkalizing agents. Any alkalizing agent known in the art may be used.

[0040] Typically, the first and/or the second hair colouring compositions may comprise a total amount of alkalizing agents ranging from 0.1% to 10%, alternatively from 0.5% to 6%, alternatively from 1% to 4%, by weight of the total composition.

[0041] Suitable alkalizing agents include, but are not limited to: ammonia; alkanolamines (such as monoethanolamine, diethanolamine, triethanolamine, monopropanolamine, dipropanolamine, tripropanolamine); 2-amino-2-methyl-1,3-propanediol, 2-amino-2-methyl-1-propanol, and 2-amino-2-hydroxymethyl-1,3-propanediol; guanidium salts; alkali metal and ammonium hydroxides (such as sodium hydroxide); alkali metal and ammonium carbonates; and mixtures thereof. Typical alkalizing agents are ammonia and/or monoethanolamine. Preferably, if present, the ammonium ions and carbonate ions are present in the composition at a weight ratio of from 3:1 to 1:10, alternatively from 2:1 to 1:5.

[0042] When the first and/or the second hair colouring compositions of the present invention are obtained by mixing a developer composition and a dye composition prior to use, the alkalizing agent is generally present in the dye composition.

Oxidative dye precursors

[0043] The first and/or the second hair colouring compositions according to the present invention may further comprise oxidative dye precursors, which are usually classified either as primary intermediates (also known as developers) or couplers (also known as secondary intermediates). Various couplers may be used with primary intermediates in order to obtain different shades. Oxidative dye precursors may be free bases or the cosmetically acceptable salts thereof. Typically, The first and/or the second hair colouring compositions may comprise a total amount of oxidative dye precursors ranging up to 12%, alternatively from 0.1% to 10%, alternatively from 0.3% to 8%, alternatively from 0.5% to 6%, by weight of the total composition.

[0044] Suitable primary intermediates include, but are not limited to: toluene-2,5-diamine, p-phenylenediamine, N-phenyl-p-phenylenediamine, N,N-bis(2-hydroxyethyl)-p-phenylenediamine, 2-hydroxyethyl-p-phenylenediamine, hydroxypropyl-bis-(N-hydroxyethyl-p-phenylenediamine), 2-methoxymethyl-p-phenylenediamine, 2-(1,2-dihydroxyethyl)-p-phenylenediamine, 2,2'-(2-(4-aminophenylamino)ethylazanediyl)diethanol, 2-(2,5-diamino-4-methoxyphenyl)propane-1,3-diol, 2-(7-amino-2H-benzo[b][1,4]oxazin-4(3H)-yl)ethanol, 2-chloro-p-phenylenediamine, p-aminophenol, p-(methylamino)phenol, 4-amino-m-cresol, 6-amino-m-cresol, 5-ethyl-o-aminophenol, 2-methoxy-p-phenylenediamine, 2,2'-methylenebis-4-aminophenol, 2,4,5,6-tetraminopyrimidine, 2,5,6-triamino-4-pyrimidinol, 1-hydroxyethyl-4,5-diaminopyrazole sulfate, 4,5-diamino-1-methylpyrazole, 4,5-diamino-1-ethylpyrazole, 4,5-diamino-1-isopropylpyrazole, 4,5-diamino-1-butylpyrazole, 4,5-diamino-1-pentylpyrazole, 4,5-diamino-1-benzylpyrazole, 2,3-diamino-6,7-dihydropyrazolo[1,2-a]pyrazol-1(5H)-one dimethosulfonate, 4,5-diamino-1-hexylpyrazole, 4,5-diamino-1-heptylpyrazole, methoxymethyl-1,4-diaminobenzene, N,N-bis(2-hydroxyethyl)-N-(4-aminophenyl)-1,2-diaminoethane, salts thereof and mixtures thereof.

[0045] Suitable couplers include, but are not limited to: resorcinol, 4-chlororesorcinol, 2-chlororesorcinol, 2-methylresorcinol, 4,6-dichlorobenzene-1,3-diol, 2,4-dimethylbenzene-1,3-diol, m-aminophenol, 4-amino-2-hydroxytoluene, 2-methyl-5-hydroxyethylaminophenol, 3-amino-2,6-dimethylphenol, 3-amino-2,4-dichlorophenol, 5-amino-6-chloro-o-cresol, 5-amino-4-chloro-o-cresol, 6-hydroxybenzomorpholine, 2-amino-5-ethylphenol, 2-amino-5-phenylphenol, 2-amino-5-methylphenol, 2-amino-6-methylphenol, 2-amino-5-ethoxyphenol, 5-methyl-2-(methylamino)phenol, 2,4-diaminophenoxyethanol, 2-amino-4-hydroxyethylaminoanisole, 1,3-bis-(2,4-diaminophenoxy)-propane, 2,2'-(2-methyl-1,3-phenylene)bis(azanediyl)diethanol, benzene-1,3-diamine, 2,2'-(4,6-diamino-1,3-phenylene)bis(oxy)diethanol, 3-(pyrrolidin-1-yl)aniline, 1-(3-(dimethylamino)phenyl)urea, 1-(3-aminophenyl)urea, 1-naphthol, 2-methyl-1-naphthol, 1,5-naphthalenediol, 2,7-naphthalenediol or 1-acetoxy-2-methylnaphthalene, 4-chloro-2-methylnaphthalen-1-ol, 4-methoxy-2-methylnaphthalen-1-ol, 2,6-dihydroxy-3,4-dimethylpyridine, 2,6-dimethoxy-3,5-pyridinediamine, 3-amino-2-methylamino-6-methoxypyridine, 2-amino-3-hydroxypyridine, 2,6-diaminopyridine, pyridine-2,6-diol, 5,6-dihydroxyindole, 6-hydroxyindole, 5,6-dihydroxyindoline, 3-methyl-1-phenyl-1H-pyrazol-5(4H)-one, 1,2,4-trihydroxybenzene, 2-(benzo[d][1,3]dioxol-5-ylamino)ethanol (also known as hydroxyethyl-3,4-methylenedioxyaniline), and mixtures thereof.

[0046] When the first and/or second hair colouring compositions of the invention are obtained by mixing a dye composition and a developer composition, the primary intermediates and couplers are usually incorporated into the dye composition.

Direct Dyes

[0047] The first and/or the second hair colouring compositions according to the present invention may further comprise compatible direct dyes, in an amount sufficient to provide additional coloring, particularly with regard to intensity. Typically, the compositions may comprise a total amount of direct dyes ranging from 0.001% to 5%, by weight of the total composition.

[0048] Suitable direct dyes include but are not limited to: Acid dyes such as Acid Yellow 1, Acid Orange 3, Acid Black 1, Acid Black 52, Acid Orange 7, Acid Red 33, Acid Yellow 23, Acid Blue 9, Acid Violet 43, HC Blue 16, Acid Blue 62, Acid Blue 25, Acid Red 4; Basic Dyes such as Basic Brown 17, Basic Red 118, Basic Orange 69, Basic Red 76, Basic Brown 16, Basic Yellow 57, Basic Violet 14, Basic Blue 7, Basic Blue 26, Basic Red 2, Basic Blue 99, Basic Yellow 29, Basic Red 51, Basic Orange 31, Basic Yellow 87, 4-(3-(4-amino-9,10-dioxo-9,10-dihydroanthracen-1-ylamino)propyl)-4-methylmorpholin-4-ium-methylsulfate, (E)-1-(2-(4-(4,5-dimethylthiazol-2-yl)diazenyl)phenyl)(ethylamino)ethyl-3-methyl-1H-imidazol-3-ium chloride, (E)-4-(2-(4-(dimethylamino)phenyl)diazenyl)-1-methyl-1H-imidazol-3-ium-3-yl)butane-1-sulfonate, (E)-4-(4-(2-methyl-2-phenylhydrazono)methyl)pyridinium-1-yl)butane-1-sulfonate, N,N-dimethyl-3-(4-(methylamino)-9,10-dioxo-4a,9,9a,10-tetrahydroanthracen-1-ylamino)-N-propylpropan-1-aminium bromide; Disperse Dyes such as Disperse Red 17, Disperse Violet 1, Disperse Red 15, Disperse Black 9, Disperse Blue 3, Disperse Blue 23, Disperse Blue 377; Nitro Dyes such as 1-(2-(4-nitrophenylamino)ethyl)urea, 2-(4-methyl-2-nitrophenylamino)ethanol, 4-nitrobenzene-1,2-diamine, 2-nitrobenzene-1,4-diamine, Picramic acid, HC Red No. 13, 2,2'-(2-nitro-1,4-phenylene)bis(azanediyl)diethanol, HC Yellow No. 5, HC Red No. 7, HC Blue No.2, HC Yellow No. 4, HC Yellow No. 2, HC Orange No. 1, HC Red No. 1, 2-(4-amino-2-chloro-5-nitrophenylamino)ethanol, HC Red No. 3, 4-amino-3-nitrophenol,

4-(2-hydroxyethylamino)-3-nitrophenol, 2-amino-3-nitrophenol, 2-(3-(methylamino)-4-nitrophenoxy)ethanol, 3-(3-amino-4-nitrophenyl)propane-1,2-diol, HC Yellow No. 11, HC Violet No. 1, HC Orange No. 2, HC Orange No. 3, HC Yellow No. 9, HC Red No. 10, HC Red No. 11, 2-(2-hydroxyethylamino)-4,6-dinitrophenol, HC Blue No. 12, HC Yellow No. 6, HC Yellow No. 12, HC Blue No. 10, HC Yellow No. 7, HC Yellow No. 10, HC Blue No. 9, 2-chloro-6-(ethylamino)-4-nitrophenol, 6-nitropyridine-2,5-diamine, HC Violet No. 2, 2-amino-6-chloro-4-nitrophenol, 4-(3-hydroxypropylamino)-3-nitrophenol, HC Yellow No. 13, 6-nitro-1,2,3,4-tetrahydroquinoxaline, HC Red No. 14, HC Yellow No. 15, HC Yellow No. 14, N2-methyl-6-nitropyridine-2,5-diamine, N1-allyl-2-nitrobenzene-1,4-diamine, HC Red No. 8, HC Green No.1, HC Blue No. 14; Natural dyes such as Annato, Anthocyanin, Beetroot, Carotene, Capsanthin, Lycopene, Chlorophyll, Henna, Indigo, Cochineal; and mixtures thereof.

[0049] When the first and/or the second hair colouring compositions are obtained by mixing a dye composition and a developer composition, the direct dyes are usually incorporated into the dye composition.

Chelants

[0050] The first and/or the second hair colouring compositions according to the present invention may further comprise chelants (also known as "chelating agent", "sequestering agent", or "sequestrant") in an amount sufficient to reduce the amount of metals available to interact with formulation components, particularly oxidizing agents, more particularly peroxides. Chelants are well known in the art and a non-exhaustive list thereof can be found in AE Martell & RM Smith, Critical Stability Constants, Vol. 1, Plenum Press, New York & London (1974) and AE Martell & RD Hancock, Metal Complexes in Aqueous Solution, Plenum Press, New York & London (1996), both incorporated herein by reference.

[0051] Typically, the first and/or the second compositions may comprise a total amount of chelants ranging from at least 0.01%, alternatively from 0.01% to 5%, alternatively from 0.25% to 3%, alternatively from 0.5% to 1%, by weight of the total composition.

[0052] Suitable chelants include, but are not limited to: carboxylic acids (such as aminocarboxylic acids), phosphonic acids (such as aminophosphonic acids), polyphosphoric acids (such as linear polyphosphoric acids), their salts thereof, and mixtures thereof. By "salts thereof", it is meant in the context of chelants - all salts comprising the same functional structure as the chelant they are referring to and including alkali metal salts, alkaline earth salts, ammonium salts, substituted ammonium salts, and mixtures thereof; alternatively sodium salts, potassium salts, ammonium salts, and mixtures thereof; alternatively monoethanolammonium salts, diethanolammonium salts, triethanolammonium salts, and mixtures thereof.

[0053] Suitable aminocarboxylic acid chelants comprise at least one carboxylic acid moiety (-COOH) and at least one nitrogen atom. Suitable aminocarboxylic acid chelants include, but are not limited to: diethylenetriamine pentaacetic acid (DTPA), ethylenediamine disuccinic acid (EDDS), ethylenediamine diglutamic acid (EDGA), 2-hydroxypropylenediamine disuccinic acid (HPDS), glycylamide-N,N'-disuccinic acid (GADS), ethylenediamine-N,N'-diglutamic acid (EDDG), 2-hydroxypropylenediamine-N,N'-disuccinic acid (HPDDS), ethylenediaminetetraacetic acid (EDTA), ethylenedicyclic acid (EDC), ethylenediamine-N,N'-bis(ortho-hydroxyphenyl acetic acid) (EDDHA), diaminoalkyldi(sulfosuccinic acids) (DDS), N,N'-bis(2-hydroxybenzyl)ethylenediamine-N,N'-diacetic acid (HBED), their salts thereof, and mixtures thereof. Other suitable aminocarboxylic type chelants include, but are not limited to: iminodiacetic acid derivatives such as N-2-hydroxyethyl N,N diacetic acid or glyceryl imino diacetic acid, iminodiacetic acid-N-2-hydroxypropyl sulfonic acid and aspartic acid N-carboxymethyl N-2-hydroxypropyl-3-sulfonic acid, β -alanine-N,N'-diacetic acid, aspartic acid-N,N'-diacetic acid, aspartic acid-N-monoacetic acid and iminodisuccinic acid chelants, ethanoldiglycine acid, their salts thereof, their derivatives thereof, and mixtures thereof. Further suitable aminocarboxylic type chelants include, but are not limited to: dipicolinic acid, 2-phosphonobutane-1,2,4-tricarboxylic acid, their salts thereof, their derivatives thereof, and mixtures thereof.

[0054] Suitable aminophosphonic acid chelants comprise an aminophosphonic acid moiety (-PO₃H₂) or its derivative -PO₃R₂, wherein R₂ is a C₁ to C₆ alkyl or aryl radical and salts thereof. Suitable aminophosphonic acid chelants include, but are not limited to: aminotri-(1-ethylphosphonic acid), ethylene-diaminetetra-(1-ethylphosphonic acid), aminotri-(1-propylphosphonic acid), aminotri-(isopropylphosphonic acid), their salts thereof, and mixtures thereof; alternatively aminotri-(methylenephosphonic acid), ethylene-diamine-tetra-(methylenephosphonic acid) (EDTMP) and diethylene-tri-amine-penta-(methylenephosphonic acid) (DTPMP), their salts thereof, their derivatives thereof, and mixtures thereof.

[0055] Suitable alternative chelants include, but are not limited to: polyethyleneimines, polyphosphoric acid chelants, etidronic acid, methylglycine diacetic acid, N-(2-hydroxyethyl)iminodiacetic acid, minodisuccinic acid, N,N-Dicarboxymethyl-L-glutamic acid, N-lauroyl-N,N',N''-ethylenediamine diacetic acid, their salts thereof, their derivatives thereof, and mixtures thereof. In a specific embodiment, the first and/or the second hair colouring compositions comprise a chelant selected from the group consisting of diethylenetriamine-N,N',N''-polyacids, diethylenetriaminepentaacetic acid (DTPA), diethylenetriaminepenta(methylene phosphonic acid) (DTPMP), diamine-N,N'-dipolyacid, monoamine monoamide-N,N'-dipolyacid, ethylenediaminedisuccinic acid (EDDS), their salts thereof, their derivatives thereof, and mixtures thereof; alternatively ethylenediaminedisuccinic acid (EDDS).

[0056] When the first and/or the second hair colouring compositions of the invention are obtained by mixing a dye composition and a developer composition, the chelants may be incorporated in the dye composition and/or in the developer composition. A chelant is usually present in the developer composition for stability reason.

5 Radical scavengers

[0057] The first and/or the second hair colouring compositions according to the present invention may further comprise a radical scavenger. As used herein the term "radical scavenger" refers to a species that can react with a radical, preferably a carbonate radical to convert the radical species by a series of fast reactions to a less reactive species. The radical scavenger may be different from the alkalizing agent and/or is present in an amount sufficient to reduce the damage to the hair during the colouring /bleaching process.

[0058] Typically, the first and/or the second hair colouring compositions may comprise a total amount of radical scavengers ranging from 0.1% to 10%, alternatively from 1% by weight to 7%, by weight of the total composition.

[0059] Suitable radical scavengers include, but are not limited to: alkanolamines, amino sugars, amino acids, esters of amino acids, and mixtures thereof; alternatively 3-amino-1-propanol, 4-amino-1-butanol, 5-amino-1-pentanol, 1-amino-2-propanol, 1-amino-2-butanol, 1-amino-2-pentanol, 1-amino-3-pentanol, 1-amino-4-pentanol, 3-amino-2-methylpropan-1-ol, 1-amino-2-methylpropan-2-ol, 3-aminopropane-1,2-diol, glucosamine, N-acetylglucosamine, glycine, arginine, lysine, proline, glutamine, histidine, sarcosine, serine, glutamic acid, tryptophan, their salts thereof, and mixtures thereof; alternatively glycine, sarcosine, lysine, serine, 2 methoxyethylamine, glucosamine, glutamic acid, morpholine, piperidine, ethylamine, 3 amino-1-propanol, and mixtures thereof. As used herein, the term "salts thereof" - in the context of radical scavengers-means particularly potassium salts, sodium salts, ammonium salts, and mixtures thereof.

pH modifier and buffering agents

[0060] The first and/or the second hair colouring compositions according to the present invention may further comprise, in addition to the alkalizing agent discussed above, a pH modifier and/or buffering agent in an amount that is sufficiently effective to adjust the pH of the composition to fall within a range from 3 to 13, alternatively from 8 to 12, alternatively from 9 to 11.

[0061] Suitable pH modifiers and/or buffering agents include, but are not limited to: ammonia; alkanolamines (such as monoethanolamine, diethanolamine, triethanolamine, monopropanolamine, dipropanolamine, tripropanolamine, tripropanolamine, 2-amino-2-methyl-1-propanol, 2-amino-2-hydroxymethyl-1,3,-propanediol); guanidium salts; alkali metal and ammonium hydroxides and carbonates; and mixtures thereof.

[0062] Further pH modifier and/or buffering agents include, but are not limited to: sodium hydroxide; ammonium carbonate; acidulents (such as inorganic and inorganic acids including for example phosphoric acid, acetic acid, ascorbic acid, citric acid or tartaric acid, hydrochloric acid); and mixtures thereof.

Thickeners and/or theology modifiers

[0063] The first and/or the second hair colouring compositions according to the invention may further comprise a thickener in an amount sufficient to provide the composition with a viscosity so that it can be readily applied to the hair without unduly dripping off the hair and causing mess. Typically, the first and/or the second hair colouring compositions may comprise a total amount of thickeners ranging from at least 0.1%, alternatively at least 0.5%, alternatively at least 1%, by weight of the total composition.

[0064] Suitable thickeners include, but are not limited to: associative polymers, polysaccharides, non-associative polycarboxylic polymers, and mixtures thereof.

[0065] As used herein, the expression "associative polymers" means amphiphilic polymers comprising both hydrophilic units and hydrophobic units, for example, at least one C8 to C30 fatty chain and at least one hydrophilic unit. Associative polymers are capable of reversibly combining with each other or with other molecules. Suitable associative thickeners include, but are not limited to: nonionic amphiphilic polymers comprising at least one hydrophilic unit and at least one fatty-chain unit; anionic amphiphilic polymers comprising at least one hydrophilic unit and at least one fatty-chain unit; cationic amphiphilic polymers comprising at least one hydrophilic unit and at least one fatty-chain unit; and amphoteric amphiphilic polymers comprising at least one hydrophilic unit and at least one fatty-chain unit, and mixtures thereof.

[0066] Suitable nonionic amphiphilic polymers comprising at least one fatty chain and at least one hydrophilic unit include, but are not limited to: celluloses modified with groups comprising at least one fatty chain (such as hydroxyethyl-celluloses modified with groups comprising at least one fatty chain chosen from alkyl, alkenyl and alkylaryl groups); hydroxypropyl guar modified with groups comprising at least one fatty chain; polyether urethanes comprising at least one fatty chain (such as C8-C30 alkyl or alkenyl groups); copolymers of vinylpyrrolidone and of fatty-chain hydrophobic monomers; copolymers of C1-C6 alkyl acrylates or methacrylates and of amphiphilic monomers comprising at least one

fatty chain; copolymers of hydrophilic acrylates or methacrylates and of hydrophobic monomers comprising at least one fatty chain, and mixtures thereof.

[0067] Suitable nonionic amphiphilic polymers comprising at least one hydrophilic unit and at least one fatty-chain unit include, but are not limited to: those polymers comprising at least one fatty-chain allyl ether unit and at least one hydrophilic unit comprising an ethylenic unsaturated anionic monomeric unit (such as a vinylcarboxylic acid unit, particularly a unit chosen from units derived from acrylic acids, methacrylic acids, and mixtures thereof), wherein the fatty-chain allyl ether unit corresponds to the monomer of formula (I) below



in which R₁ is chosen from H and CH₃, B is an ethyleneoxy radical, n is chosen from zero and integers ranging from 1 to 100, R is chosen from hydrocarbon-based radicals chosen from alkyl, alkenyl, arylalkyl, aryl, alkylaryl and cycloalkyl radicals, comprising from 8 to 30 carbon atoms, and, further, for example, from 10 to 24 carbon atoms and even further, for example, from 12 to 18 carbon atoms.

[0068] Suitable anionic amphiphilic polymers include, but are not limited to: those polymers comprising at least one hydrophilic unit of unsaturated olefinic carboxylic acid type, and at least one hydrophobic unit of the type such as a (C₈-C₃₀) alkyl ester or (C₈-C₃₀) oxyethylenated alkyl ester of an unsaturated carboxylic acid, wherein the hydrophilic unit of unsaturated olefinic carboxylic acid type corresponds to, for example, the monomer of formula (II) below



in which R₁ is chosen from H, CH₃, C₂H₅ and CH₂COOH (i.e. acrylic acid, methacrylic, ethacrylic and itaconic acid units); and wherein the hydrophobic unit of the type such as a (C₈-C₃₀) alkyl ester or (C₈-C₃₀) oxyethylenated alkyl ester of an unsaturated carboxylic acid corresponds to, for example, the monomer of formula (III) below



in which R₁ is chosen from H, CH₃, C₂H₅ and CH₂COOH (i.e. acrylate, methacrylate, ethacrylate and itaconate units), B is an ethyleneoxy radical, n is chosen from zero and integers ranging from 1 to 100, R₂ is chosen from C₈-C₃₀ alkyl radicals, for example, C₁₂-C₂₂ alkyl radical. Anionic amphiphilic polymers may further be cross-linked. The crosslinking agent can be a monomer comprising a group (IV) below



with at least one other polymerizable group whose unsaturated bonds are not conjugated with respect to one another. Mention may be made, for example, of polyallyl ethers such as polyallylsucrose and polyallyl pentaerythritol.

[0069] Suitable cationic amphiphilic polymers include, but are not limited to: quaternized cellulose derivatives and polyacrylates comprising amino side groups. The quaternized cellulose derivatives are, for example, chosen from quaternized celluloses modified with groups comprising at least one fatty chain, such as alkyl, arylalkyl and alkylaryl groups comprising at least 8 carbon atoms, and mixtures thereof, quaternized hydroxyethylcelluloses modified with groups comprising at least one fatty chain, such as alkyl, arylalkyl and alkylaryl groups comprising at least 8 carbon atoms, and mixtures thereof. The alkyl radicals borne by the above quaternized celluloses and hydroxyethylcelluloses, for example, contain from 8 to 30 carbon atoms. The aryl radicals, for example, are chosen from phenyl, benzyl, naphthyl and anthryl groups.

[0070] Suitable amphoteric amphiphilic polymers comprising at least one hydrophilic unit and at least one fatty-chain unit, may be made, for example, of methacrylamidopropyltrimethylammonium chloride/acrylic acid/C₈-C₃₀ alkyl methacrylate copolymers, wherein the alkyl radical is, for example, a stearyl radical.

[0071] Preferred associative polymers comprise at least one hydrophilic unit which is unsaturated carboxylic acid or its derivatives, and at least one hydrophobic unit which is a C₈ to C₃₀ alkyl ester or oxyethylenated C₈-C₃₀ alkyl ester of unsaturated carboxylic acid. The unsaturated carboxylic acid is preferably acrylic acid, methacrylic acid or itaconic acid. Commercially available materials include those sold as Aculy-22 by Rohm & Haas; Permulen TR1, Carbopol 2020, Carbopol Ultrez-21 by Noveon, Structure 2001/3001 by National Starch. Other preferred associative polymers include polyether polyurethane, commercially available as Aculy-44/-46 by Rohm and Haas. Further preferred associative polymers include cellulose modified with groups comprising at least one C₈ - C₃₀ fatty chain, commercially available under the trade name Natrosol Plus Grade 330 CS by Aqualon.

[0072] Suitable non-associative cross-linked polycarboxylic polymers include, but are not limited to: cross-linked acrylic acid homopolymers, copolymers of acrylic or (meth)acrylic acid and of C₁-C₆ alkyl acrylate or (meth)acrylate, and mixtures thereof. Commercially available materials include those sold as Carbopol 980/981/954/2984/5984 by Noveon,

Synthalen M/Synthalen L/Synthalen K by 3V Sigma, Aculyn-33 by Rohm and Haas.

[0073] Suitable polysaccharides include, but are not limited to: glucans, modified and unmodified starches (such as those derived, for example, from cereals, for instance wheat, corn or rice, from vegetables, for instance yellow pea, and tubers, for instance potato or cassava), amylose, amylopectin, glycogen, dextrans, celluloses and derivatives thereof (methylcelluloses, hydroxyalkylcelluloses, ethyl hydroxyethylcelluloses, and carboxymethylcelluloses), mannans, xy-
 5 lans, lignins, arabans, galactans, galacturonans, chitin, chitosans, glucuronoxylans, arabinoxylans, xyloglucans, gluco-
 mannans, pectic acids and pectins, alginic acid and alginates, arabinogalactans, carrageenans, agars, glycosaminoglu-
 cans, gum arabics, gum tragacanth, ghatti gums, karaya gums, carob gums, galactomannans, such as guar gums,
 10 and nonionic derivatives thereof (hydroxypropyl guar) and bio-polysaccharides, such as xanthan gums, gellan gums,
 welan gums, scleroglucans, succinoglycans, and mixtures thereof. Suitable polysaccharides are described in "Encyclo-
 pedia of Chemical Technology", Kirk-Othmer, Third Edition, 1982, volume 3, pp. 896-900, and volume 15, pp. 439-458,
 in "Polymers in Nature" by E. A. MacGregor and C. T. Greenwood, published by John Wiley & Sons, Chapter 6, pp.
 240-328, 1980, and in "Industrial Gums-Polysaccharides and their Derivatives", edited by Roy L. Whistler, Second Edition,
 15 published by Academic Press Inc., all three being incorporated herein by reference. A preferred polysaccharide is a bio-
 polysaccharide, particularly bio-polysaccharides selected from xanthan gum, gellan gum, welan gum, scleroglucan or
 succinoglycan; commercially available as Keltrol® T by Kelco and Rheozan® by Rhodia Chimie. Another preferred
 polysaccharide is hydroxypropyl starch derivative, particularly hydroxypropyl starch phosphate, commercially available
 as Structure XL® by National Starch. Commercially available salt-tolerant thickeners include, but not limited to: xanthan,
 20 guar, hydroxypropyl guar, scleroglucan, methyl cellulose, ethyl cellulose (commercially available as Aquacote), hydrox-
 yethyl cellulose (Natrosol), carboxymethyl cellulose, hydroxypropylmethyl cellulose, microcrystalline cellulose, hydroxy-
 butylmethyl cellulose, hydroxypropyl cellulose (Klucel), hydroxyethyl ethyl cellulose, cetyl hydroxyethyl cellulose (Na-
 trosol Plus 330), N-vinylpyrrolidone (Povidone), Acrylates/Ceteth-20 Itaconate Copolymer (Structure 3001), hydroxypro-
 pyl starch phosphate (Structure ZEA), polyethoxylated urethanes or polycarbamyl polyglycol ester such as PEG-150/De-
 cyl/SMDI copolymer (Aculyn 44), PEG-150/Stearyl/SMDI copolymer (Aculyn 46), trihydroxystearin (Thixcin), acrylates
 25 copolymer (Aculyn 33) or hydrophobically modified acrylate copolymers (such as Acrylates / Steareth-20 Methacrylate
 Copolymer as Aculyn 22), acrylates/steareth-20 methacrylate crosspolymer (Aculyn 88), acrylates/vinyl neodecanoate
 crosspolymer (Aculyn 38), acrylates/beheneth-25 methacrylate copolymer (Aculyn 28), acrylates/C10-30 alkyl acrylate
 crosspolymer (Carbopol ETD 2020), non-ionic amphiphilic polymers comprising at least one fatty chain and at least
 one hydrophilic unit selected from polyether urethanes comprising at least one fatty chain, blends of Ceteth - 10 phosphate,
 30 Di-cetyl phosphate and Cetearyl alcohol (available as Crodafos CES), and mixtures thereof.

[0074] Thickeners for use in the first and/or second developer compositions may include acrylates copolymer, hydro-
 phobically modified acrylate copolymers (e.g. Acrylates / Steareth-20 Methacrylate Copolymer) and mixtures thereof.
 Thickeners for use in the first and/or second dye compositions may include blends of Ceteth - 10 phosphate, Di-cetyl
 phosphate and Cetearyl alcohol (as CRODAFOS CES).

Gel network thickener system

[0075] The first and/or second hair colouring compositions of the present invention may comprise at least one gel
 network thickener system. Said system may comprise at least one low HLB surfactant and/or amphophile having a high
 40 melting point, and at least one additional second surfactant as specified hereinafter. Suitable gel network thickener
 systems are disclosed in PCT application WO2006/060598A1.

[0076] Said low HLB surfactant and/or amphophile may have preferably an HLB of 6 or less and melting point of at
 least 30°C. It may be selected from the group consisting of cetyl, stearyl, cetostearyl or behenyl alcohols, steareth-2,
 glycerol monostearate and mixtures thereof. Said second surfactant may be anionic, non-ionic or cationic. Anionic
 45 surfactants may be selected from the group consisting of alkyl ether phosphates having in average 1-3 ethylene oxide
 units and comprising an alkyl radical comprising from 8 to 30 carbon atoms. Said non-ionic surfactants may be selected
 from the group consisting of those comprising one or more polyethyleneoxide chain including polyoxyethylene alkyl
 ethers having from 100 to 200 ethylene oxide units (e.g. steareth-100, steareth-150). Said cationic surfactant may be
 selected from the group consisting of behentrimonium chloride, behenamidopropyltrimonium methosulfate, stearamido-
 propyltrimonium chloride, arachidtrimonium chloride and mixtures thereof. A preferred gel network thickening system
 50 comprises fatty alcohols having 14 to 30 carbon atoms (cetyl and/or stearyl alcohol) and alkyl ether phosphates (e.g.
 from 1 to 3 ethylene oxide units). The first and/or second hair colouring compositions of the present invention may
 comprise a total amount of gel network thickening system of from 2% to 10% by weight of respectively the first and/or
 second hair colouring composition. The weight ratio of the low HLB surfactants to the second specified surfactants is
 55 preferably from 10:1 to 1:1.

Carbonate ion sources

[0077] The first and/or the second hair colouring compositions according to the present invention may further comprise a source of carbonate ions, carbamate ions, hydrogen carbonate ions, and mixtures thereof in a sufficient amount to reduce damage to the hair during the coloring process. Typically, the first and/or the second hair colouring compositions may comprise a total amount of a carbonate ion source ranging from 0.1% to 15%, alternatively from 0.1% to 10%, alternatively from 1% to 7%, by weight of the total composition.

[0078] Suitable carbonate ion sources include, but are not limited to: sodium carbonate, sodium hydrogen carbonate, potassium carbonate, potassium hydrogen carbonate, guanidine carbonate, guanidine hydrogen carbonate, lithium carbonate, calcium carbonate, magnesium carbonate, barium carbonate, ammonium carbonate, ammonium hydrogen carbonate and mixtures thereof; alternatively sodium hydrogen carbonate, potassium hydrogen carbonate, and mixtures thereof; alternatively ammonium carbonate, ammonium hydrogen carbonate, and mixtures thereof.

Conditioning agents

[0079] The first and/or the second hair colouring compositions according to the present invention may further comprise a conditioning agent, and/or be used in combination with a composition comprising a conditioning agent.

[0080] Typically, the first and/or the second hair colouring compositions may comprise a total amount of conditioning agents ranging from 0.05% to 20%, alternatively from 0.1% to 15%, alternatively from 0.2% to 10%, alternatively from 0.2% to 2%, alternatively from 0.5% to 2%, by weight of the total composition. The conditioning agent may be included in a separate pre- and/or posttreatment composition.

[0081] Suitable conditioning agents include, but are not limited to: silicones, aminosilicones, fatty alcohols, polymeric resins, polyol carboxylic acid esters, cationic polymers, cationic surfactants, insoluble oils and oil derived materials and mixtures thereof. Additional conditioning agents include mineral oils and other oils such as glycerin and sorbitol.

[0082] Particularly useful conditioning materials are cationic polymers. Conditioners of cationic polymer type can be chosen from those comprising units of at least one amine group chosen from primary, secondary, tertiary and quaternary amine groups that may either form part of the main polymer chain, or be borne by a side substituent that is directly attached to the main polymer chain, described hereinafter.

[0083] Suitable silicones include, but are not limited to: polyalkylsiloxane oils, linear polydimethylsiloxane oils containing trimethylsilyl or hydroxydimethylsiloxane endgroups, polymethylphenylsiloxane, polydimethylphenylsiloxane or polydimethyldiphenylsiloxane oils, silicone resins, organofunctional siloxanes having in their general structure one or a number of organofunctional group(s), the same or different, attached directly to the siloxane chain and mixtures thereof. Said organofunctional group(s) may be selected from: polyethyleneoxy and/or polypropyleneoxy groups, (per)fluorinated groups, thiol groups, substituted or unsubstituted amino groups, carboxylate groups, hydroxylated groups, alkoxyated groups, quaternium ammonium groups, amphoteric and betaine groups. The silicone can either be used as a neat fluid or in the form of a pre-formed emulsion. Suitable silicones also include: silicones containing groups that may be ionized into cationic groups, for example aminosilicones containing at least 10 repeating siloxane ($\text{Si}(\text{CH}_3)_2\text{-O}$) units within the polymer chain, with either terminal, graft, or a mixture of terminal and graft amino functional groups. Example functional groups are not limited to aminoethylaminopropyl, aminoethylaminoisobutyl, aminopropyl. In the case of graft polymers, the terminal siloxane units can be $(\text{CH}_3)_3\text{Si-O}$, $\text{R}_{12}(\text{CH}_3)_2\text{Si-O}$, where R_{12} can be either OH or OR_{13} , where R_{13} is a C1-C8 alkyl group, or a mixture of both terminal groups. These silicones are also available as preformed emulsions. Commercially available aminosilicones include those sold as DC-2-8566, DC 7224, DC-2-8220 by Dow Corning; SF1708, SM2125 by GE Silicones; Wacker Belsil ADM 653/ADM 1100/ADM 1600/ADM 652/ADM 6057E/ADM 8020 by Wacker Silicones; DC929, DC939, DC949 by Dow Corning; SM2059 by GE Silicones. Suitable aminosilicones may also contain additional functional groups, particularly additional functional groups including polyoxyalkylene, the reaction product of amines and carbinols, and alky chains. Commercially available materials are known as methoxy PEG/PPG-7/3 Amino-propyl Dimethicone (e.g. Abil Soft AF100, by Degussa), or as Bis(C13-15 Alkoxy)PG Amodimethicone (e.g. DC 8500, by Dow Corning).

[0084] Suitable cationic polymers include, but are not limited to: polymers comprising units of at least one amine group chosen from primary, secondary, tertiary and quaternary amine groups that may either form part of the main polymer chain or be borne by a side substituent that is directly attached to the main polymer chain. Such cationic polymers generally have a number average molecular mass ranging from 500 to 5×10^6 , alternatively from 1000 to 3×10^6 . Preferably the cationic polymers are selected from polymers of the polyamine, polyamino amide and polyquaternary ammonium type.

[0085] Suitable polymers of the polyamine, polyamino amide and polyquaternary ammonium type include, but are not limited to:

- 1) Homopolymers and copolymers derived from acrylic or methacrylic esters or amides. Copolymers of these pol-

ymers may also comprise at least one unit derived from comonomers which may be chosen from the family of acrylamides, methacrylamides, diacetone acylamides, acrylamides and methacrylicamides substituted on the nitrogen with at least one group chosen from lower (C1-C4) alkyls, acrylic and methacrylic acids and esters thereof, vinylactams such as vinylpyrrolidone and vinylcaprolactam, and vinyl esters. Suitable examples include copolymers of acrylamide and of methacryloyloxyethyltrimethylammonium methosulfate, including polymers known as Polyquaternium-5 (e.g. commercially available under the trade name Reten 210/220/230/240/1104/1105/1006 by Hercules; Merquat 5/5 SF by Nalco); copolymers of vinylpyrrolidone and dimethylaminopropyl methacrylamide, including polymers known as Polyquaternium-28 (e.g. Gafquat HS-100 by ISP); copolymers of vinyl pyrrolidone and dialkylaminoalkyl acrylates or methacrylates, including polymers known as Polquaternium-11 (see Gafquat 440/734/755/755N by ISP; Luviquat PQ11 PM by BASF; Polyquat-11 SL by Sino Lion); copolymers of vinylpyrrolidone, dimethylaminopropyl methacrylamide and methacryloylaminopropyl lauryldimonium chloride, including polymers known as polyquaternium-55 (e.g. Styleze W-20 by ISP); copolymers of acrylic acid, acrylamide and methacrylamidopropyltrimonium chloride, including polymers known as Polyquaternium-53 (e.g. Merquat 2003 by Nalco); copolymers of dimethylaminopropylacrylate (DMAPA), acrylic acid and acrylonitrogens and diethyl sulphate, including polymers known as Polyquaternium-31 (e.g. Hypan QT100 by Lipo); copolymers of acrylamide, acrylamidopropyltrimonium chloride, 2-amidopropylacrylamide sulfonate, and dimethylaminopropylacrylate (DMAPA), including polymers known as polyquaternium-43 (e.g. Bozequat 4000 by Clairant); copolymers of acrylic acid, methylacrylate and methacrylamidopropyltrimonium chloride, including polymers known as Polyquaternium-47 (e.g. Merquat 2001/2001N by Nalco); copolymers of methacryloyl ethyl betaine, 2-hydroxyethyl methacrylate and methacryloyl ethyl trimethyl ammonium chloride, including polymers known as Polyquaternium-48 (e.g. Plascize L-450 by Goo Chemical); copolymers of acrylic acid diallyl dimethyl ammonium chloride and acrylamide, including polymers known as polyquaternium-39 (e.g. Merquat 3330/3331 by Nalco). Further suitable examples include copolymers of methacrylamide methacrylamido-propyltrimonium and methacryloylethyltrimethyl ammonium chloride and their derivatives, either homo or copolymerised with other monomers, including polymers known as Polyquaternium-8, Polyquaternium-9, Polyquaternium-12, Polyquaternium-13 Polyquaternium-14, Polyquaternium-15 (e.g. Rohagit KF 720 F by Rohm), Polyquaternium-30 (e.g. Mexomere PX by Chimex), Polyquaternium-33, Polyquaternium-35, Polyquaternium-36 (e.g. Plex 3074 L by Rhon), Polyquaternium 45 (e.g. Plex 3073L by Rohn), Polyquaternium 49 (e.g. Plascize L-440 by Goo Chemicals), Polyquaternium 50 (e.g. Plascize L-441 by Goo Chemicals), Polyquaternium-52.

2) Cationic polysaccharides, such as cationic celluloses and cationic galactomannan gums. Among the cationic polysaccharides that maybe mentioned, for example, are cellulose ether derivatives comprising quaternary ammonium groups and cationic cellulose copolymers or cellulose derivatives grafted with a water-soluble quaternary ammonium monomer and cationic galactomannan gums. Suitable examples include copolymers of hydroxyethyl-celluloses and diallyldimethyl ammonium chlorides, including polymers known as Polyquaternium-4 (e.g. Celquat L 200 and Celquat H 100 by National Starch); copolymers of hydroxyethylcelluloses and a trimethyl ammonium substituted epoxide, including polymers known as Polyquaternium-10 (e.g. AEC Polyquaternium-10 by A&E Connock; Catinal C-100/HC-35/HC-100/HC-200/LC-100/LC-200 by Toho; Celquat SC-240C/SC-230M by National Starch; Dekaquat 400/3000 by Dekker; Leogard GP by Akzo Nobel; RITA Polyquat 400/3000 by RITA; UCARE Polymer JR-125/JR-400/JR-30M/LK/LR 400/LR 30M by Amerchol); copolymers of hydroxyethylcelluloses and lauryl dimethyl ammonium substituted epoxides, including polymers known as Polyquaternium-24 (e.g. Quatrisoft polymer LM-200 by Amerchol); derivatives of hydroxypropyl guar, including polymers as guar hydroxypropyltrimonium chloride (e.g. Catinal CG-100, Catinal CG-200 by Toho; Cosmedia Guar C-261N, Cosmedia Guar C-261N, Cosmedia Guar C-261N by Cognis; DiaGum P 5070 by Freedom Chemical Diamalt; N-Hance Cationic Guar by Hercules/Aqualon; Hi-Care 1000, Jaguar C-17, Jaguar C-2000, Jaguar C-13S, Jaguar C-14S, Jaguar Excel by Rhodia; Kiproguum CW, Kiproguum NGK by Nippon Starch); hydroxypropyl derivatives of guar hydroxypropyltrimonium chloride, including polymers known as hydroxypropyl guar hydroxypropyltrimonium chloride (e.g. Jaguar C-162 by Rhodia).

3) Polyamino amide derivatives resulting from the condensation of polyalkylene polyamines with polycarboxylic acids followed by alkylation with difunctional agents. Among the derivative, mention may be made for example to adipic acid / dimethylaminohydroxypropyl / diethylenetriamine.

4) Polymers obtained by reaction of a polyalkylene polyamine comprising two primary amines groups and at last one secondary amine group with a decarboxylic acid chosen from diglycolic acids and saturated aliphatic dicarboxylic acids comprising from 3 to 8 carbon atoms. Suitable examples include the polymer adipic acid / epoxypopyl / diethylenetriamine.

5) Cyclopolymers of dialkldiallylamine or of dialkyldiallylammonium, including: Dimethyldiallylammonium chloride polymers, including polymers known as Polyquaternium-6 (e.g. Merquat 100 by Nalco; Mirapol 100 by Rhodia; Rheocare CC6 by Cosmetic Rheologies; AEC polyquaternium-6 by A&E Connock; Agequat 400 by CPS; Conditioner P6 by 3V Inc.; Flocare C106 by SNF; Genamin PDAC by Clariant; Mackernium 006 by McIntyre); copolymers of acrylamides and dimethyldiallylammonium chlorides monomers, including polymers known as Polyquaternium-7

(e.g. AEC Polyquaternium-7 by A&E Connock; Agequat-5008/C-505 by CPS; Conditioner P7 by 3V Inc.; Flocare C 107 by SNF; Mackernium 007/007S by McIntyre; ME Polymer 09W by Toho; Merquat 550/2200/S by Nalco; Mirapol 550 by Rhodia; Rheocare CC7/CCP7 by Cosmetic Rheologies; Salcare HSP-7/SC10/ Super 7 by Ciba); copolymers of dimethyldiallylammoniumchlorides and acrylic acids, including polymers known as polyquaternary-22 (e.g. Merquat 280/Merquat 295 by Nalco).

6) Quaternary diammonium polymers comprising repeat units corresponding to $[-N+(R1)(R2) - A1 - N+(R3)(R4) - B1 -] [2X-]$, in which R1, R2, R3 and R4, which may be identical or different, are chosen from aliphatic, alicyclic and arylaliphatic radicals comprising from 1 to 20 carbon atoms and from lower hydroxyalkylaliphatic radicals, or R1, R2, R3 and R4, together or separately, constitute, with the nitrogen atoms to which they are attached, heterocycles optionally comprising a second heteroatom other than nitrogen, or R1, R2, R3 and R4, are chosen from linear or branched C1-C6 alkyl radicals substituted with at least one group chosen from nitrile, ester, acyl and amide groups and groups of $-CO-O-R5-D$ and $-CO-NH-R5-D$ wherein R5 is chosen from alkylene groups and D is chosen from quaternary ammonium groups. A1 and B1, which may be identical or different, are chosen from linear and branched, saturated or unsaturated polymethylene groups comprising 2 to 20 carbon atoms. The polymethylene groups may comprise, linked to or intercalated in the main ring, at least one entity chosen from aromatic rings, oxygen and sulphur atoms and sulphoxide, sulphone, disulphide, amino, alkylamino, hydroxyl, quaternary, ammonium, ureido, amide and ester groups, and X- is an anion derived from inorganic and organic acids. D is chosen from a glycol residue, a bis-secondary diamine residue, a bis-primary diamine residue or a ureylene group. Suitable examples include polymers known as Hexadimethrine chloride, where R1, R2, R3 and R4 are each methyl radicals, A1 is $(CH_2)_3$ and B1 is $(CH_2)_6$ and X = Cl; as polyquaternium-34 where R1 and R2 are ethyl radicals and R3 and R4 are methyl radicals and A1 is $(CH_2)_3$ and B1 is $(CH_2)_3$ and X = Br (e.g. Mexomere PAX by Chimax).

7) Polyquaternary ammonium polymers comprising repeating units of formula $[-N+(R6)(R7)-(CH_2)_r-NH-CO-(CH_2)_q-(CO)_t-NH-(CH_2)_s-N+(R8)(R9)-A-] [2X-]$, in which R6, R7, R8 and R9 which may be identical or different, are chosen from a hydrogen atom and a methyl, ethyl, propyl, hydroxyethyl, hydroxypropyl, and $-CH_2CH_2(OCH_2CH_2)_pOH$ radicals, wherein p is equal to 0 or an integer ranging from 1 to 6, wherein R6, R7, R8 and R9 do not all simultaneously represent a hydrogen atom. R and s which maybe identical or different are each an integer ranging from 1 to 6, q is equal to 0 or an integer ranging from 1 to 34 and X- is anion such as a halide. T is an integer chosen to be equal to 0 or 1. A is chosen from divalent radicals such as $-CH_2-CH_2-O-CH_2-CH_2-$. Suitable examples include: polymers known as polyquaternium-2, where $r=s=3$, $q=0$, $t=0$, R6, R7, R8 and R9 are methyl groups, and A is $-CH_2-CH_2-O-CH_2-CH_2-$ (e.g. Ethpol PQ-2 from Ethox; Mirapol A-15 by Rhodia); as polyquaternium-17 where $r=s=3$, $q=4$, $t=1$ R6, R7, R8 and R9 are methyl groups, and A is $-CH_2-CH_2-O-CH_2-CH_2-$; as Polyquaternium 18, where $r=s=3$, $q=7$, $t=1$ R6, R7, R8 and R9 are methyl groups, and A is $-CH_2-CH_2-O-CH_2-CH_2-$; as the block copolymer formed by the reaction of Polyquaternium-2 with Polyquaternium-17, which are known as Polyquaternium 27 (e.g. Mirapol 175 by Rhodia).

8) Copolymers of vinylpyrrolidones and of vinylimidazoles and optionally vinylcaprolactams, including polymers known as Polyquaternary-16 formed from methylvinylimidazolium chlorides and vinylpyrrolidones (e.g. Luviquat FC370//FC550/FC905/HM-552 by BASF); copolymers of vinylcaprolactams and vinylpyrrolidones with methylvinylimidazolium methosulfates, including polymers known as Polyquaternium-46 (e.g. Luviquat Hold by BASF); copolymers of vinylpyrrolidones and quaternized imidazolines, including polymers known as polyquaternary 44 (e.g. Luviquat Care by BASF).

9) Polyamines such as Polyquart H sold by Cognis under the reference name polyethylene glycol (15) tallow polyamine.

10) Cross linked methacryloyloxy(C1-C4)alkyltri(C1-C4)alkylammonium salt polymers such as the polymers obtained by homopolymerisation of dimethylaminoethyl methacrylates quaternized with methyl chloride, or by copolymerisation of acrylamides with dimethylaminoethyl methacrylates quaternized with methyl chloride, the homo or copolymerisation being followed by crosslinking with a compound comprising olefinic unsaturation, such as methylenebisacrylamides, including polymers known as Polyquaternium-37 (e.g. Synthalen CN/CR/CU sold by 3V sigma; or as a dispersion in another media such as Salcare SC95/SC96 by Ciba; Rheocare CTH(E) by Cosmetic Rheologies) and polymers known as Polyquaternium-32 (e.g. sold as a dispersion in mineral oil such as Salcare SC92 by Ciba).

11) Further examples of cationic polymers include polymers known as Polyquaternium 51 (e.g. Lipidure-PMB by NOF), as Polyquaternium 54 (e.g. Qualty-Hy by Mitsui), as Polyquaternium 56 (e.g. Hairrol UC-4 by Sanyo chemicals), as Polyquaternium 87 (e.g. Luviquat sensation by BASF).

12) Silicone polymers comprising cationic groups and/or groups which may be ionised into cationic groups. Suitable examples include cationic silicones of the general formula $(R10-N+(CH_3)_2)-R11-(Si(CH_3)_2-O)_x-R11-(N+(CH_3)_2)-R10$, where R10 is an alkyl derived from coconut oil, and R11 is $(CH_2CHOCH_2O(CH_2)_3$ and x is a number between 20 and 2000, including polymers known as Quaternium 80 (e.g. Abil Quat 3272/3474 sold by Goldschmidt); silicones containing groups which may be ionised into cationic groups, for example aminosilicones containing at least 10 repeating siloxane $-(Si(CH_3)_2-O)$ units within the polymer chain, with either terminal, graft or

a mixture of terminal and graft aminofunctional groups. Example functional groups are not limited to aminoethyl-aminopropyl, aminoethylaminoisobutyl, aminopropyl. In the case of graft polymers, the terminal siloxane units can either be $(\text{CH}_3)_3\text{Si}-\text{O}$ or $\text{R}_{12}(\text{CH}_3)_2\text{Si}-\text{O}$, where R_{12} can be either OH or OR_{13} , where R_{13} is a C1-C8 alkyl group, or a mixture of both functional terminal groups. These silicones are also available as preformed emulsions. Polymer with terminal siloxane units of $(\text{CH}_3)_3\text{Si}-\text{O}$ examples includes polymers known as trimethylsilylamodimethicone (e.g. DC-2-8566, DC 7224, DC-2-8220 by Dow Corning; SF1708, SM 2125 GE Silicones; Wacker Belsil ADM 653 by Wacker silicones). Further examples include polymers with terminal siloxane units of $(\text{R}_{12}\text{O})(\text{CH}_3)_2\text{Si}-\text{O}$ where R_{12} can be either OH or OR_{13} , where R_{13} is a C1-C8 alkyl group, or a mixture of both functional terminal groups, known as amodimethicone (e.g. Wacker Belsil ADM 1100/ADM 1600/ADM 652/ADM 6057E/ADM 8020 by Wacker Silicones; DC929, DC939, DC949 by Dow Corning; SM2059 by GE silicones). Silicones containing groups which may be ionised into cationic groups - for example silicones containing at least 10 repeating siloxane $-(\text{Si}(\text{CH}_3)_2-\text{O})$ units within the polymer chain, with either terminal, graft or a mixture of terminal and graft aminofunctional groups, together with additional functional groups. Additional functional groups can include polyoxyalkylene, the reaction product of amines and carbinols, alkyl chains. For example products known as methoxy PEG/PPG-7/3 Aminopropyl Dimethicone (e.g. Abil Soft AF100 by Degussa). For example products known as Bis (C13-15 Alkoxy) PG Amodimethicone (e.g. DC 8500 by Dow Corning).

[0086] The cationic polymer may be selected from the group consisting of polyquaternium 37, polyquaternium 7, polyquaternium 22, polyquaternium 87, and mixtures thereof; particularly from the group consisting of polyquaternium 37, polyquaternium 22, and mixtures thereof.

Surfactants

[0087] The first and/or the second hair colouring compositions according to the present invention may further comprise a surfactant. Suitable surfactants generally have a lipophilic chain length of from 8 to 30 carbon atoms and can be selected from anionic surfactants, nonionic surfactants, amphoteric surfactants, cationic surfactants, and mixtures thereof.

[0088] Typically, the first and/or the second hair colouring compositions may comprise a total amount of surfactants ranging from 1% to 60%, alternatively from 2% to 30%, alternatively from 8% to 25%, alternatively from 10% to 20%, by weight of the total composition.

[0089] The first and/or the second hair colouring compositions may comprise a mixture of an anionic surfactant and an amphoteric surfactant with one or more nonionic surfactants. The compositions may comprise a total amount of anionic surfactant ranging from 0.1% to 20%, alternatively from 0.1% to 15%, alternatively from 5% to 15%, by weight of the total composition; and a total amount of amphoteric and/or nonionic components, which may range independently from each other from 0.1% to 15%, alternatively from 0.5% to 10%, alternatively from 1% to 8%, by weight of the total composition.


[0090] Suitable anionic surfactants include, but are not limited to: salts (such as alkaline salts, for example, sodium salts, ammonium salts, amine salts, amino alcohol salts and magnesium salts) of the following compounds: alkyl sulphates, alkyl ether sulphates, alkylamido ether sulphates, alkylaryl polyether sulphates, monoglyceride sulphates; alkyl sulphonates, alkyl phosphates, alkylamide sulphonates, alkylaryl sulphonates, α -olefin sulphonates, paraffin sulphonates; alkyl sulphosuccinates, alkyl ether sulphosuccinates, alkylamide sulphosuccinates; alkyl sulphosuccinates; alkyl sulphoacetates; alkyl ether phosphates; acyl sarcosinates; acyl isethionates; N-acyltaurates; and mixtures thereof. The alkyl or acyl radical of all of these various compounds, for example, comprises from 8 to 24 carbon atoms, and the aryl radical, for example, is chosen from phenyl and benzyl groups. Among the anionic surfactants, which can also be used, mention may also be made of fatty acid salts such as the salts of oleic, ricinoleic, palmitic and stearic acids, coconut oil acid or hydrogenated coconut oil acid; acyl lactylates in which the acyl radical comprises from 8 to 20 carbon atoms. Weakly anionic surfactants can also be used, such as alkyl-D-galactosiduronic acids and their salts, as well as polyoxyalkylenated (C_6 - C_{24}) alkyl ether carboxylic acids, polyoxyalkylenated (C_6 - C_{24}) alkylaryl ether carboxylic acids, polyoxyalkylenated (C_6 - C_{24}) alkylamido ether carboxylic acids and their salts, for example, those comprising from 2 to 50 ethylene oxide groups, and mixtures thereof. Anionic derivatives of polysaccharides, for example carboxyalkyl ether of alkyl polyglucosides, can be also used. Nonionic surfactants are compounds that are well known (see, for example, in this respect "Handbook of Surfactants" by M. R. Porter, published by Blackie & Son (Glasgow and London), 1991, pp. 116-178). Suitable non-ionic surfactants include, but are not limited to: polyethoxylated, polypropoxylated and polyglycerolated fatty acids, alkyl phenols, α -diols and alcohols comprising a fatty chain comprising, for example, from 8 to 18 carbon atoms, it being possible for the number of ethylene oxide or propylene oxide groups to range, for example, from 2 to 200 and for the number of glycerol groups to range, for example, from 2 to 30. Mention may also be made of copolymers of ethylene oxide and of propylene oxide, condensates of ethylene oxide and of propylene oxide with fatty alcohols; polyethoxylated fatty amides preferably having from 2 to 30 mol of ethylene oxide and their momoethanolamine

and diethanolamine derivatives, polyglycerolated fatty amides, for example, comprising on average from 1 to 5, and such as from 1.5 to 4, glycerol groups; polyethoxylated fatty amines such as those containing from 2 to 30 mol of ethylene oxide; oxyethylenated fatty acid esters of sorbitan having from 2 to 30 mol of ethylene oxide; fatty acid esters of sucrose, fatty acid esters of polyethylene glycol, alkylpolyglycosides, N-alkylglucamine derivatives, amine oxides such as (C₁₀-C₁₄)alkylamine oxides or N-acylaminopropylmorpholine oxides.

[0091] Suitable amphoteric surfactants include, but are not limited to: aliphatic secondary and tertiary amine derivatives in which the aliphatic radical is chosen from linear and branched chains comprising from 8 to 22 carbon atoms and comprising at least one water-soluble anionic group (for example carboxylate, sulphonate, sulphate, phosphate or phosphonate); mention may also be made of (C₈-C₂₀)alkylbetaines, sulphobetaines, (C₈-C₂₀)alkylamido(C₁-C₆)alkylbetaines or (C₈-C₂₀)alkylamido(C₁-C₆)alkylsulphobetaines. Among the amine derivatives, mention may be made of the products sold as Miranol, as described, for example, in U.S. Pat. Nos. 2,528,378 and 2,781,354 and having the structures of: R₂-CONHCH₂CH₂-N⁺(R₃)(R₄)(CH₂COO⁻), (VI) in which: R₂ is chosen from alkyl radicals derived from an acid R₂-COOH present in hydrolysed coconut oil, and heptyl, nonyl and undecyl radicals, R₃ is a β-hydroxyethyl group and R₄ is a carboxymethyl group; and of R₅-CONHCH₂CH₂-N(B)(C) (VII) wherein B represents-CH₂CH₂OX', C represents -(CH₂)_z-Y', with z=1 or 2, X' is chosen from the -CH₂CH₂-COOH group and a hydrogen atom, Y' is chosen from -COOH and -CH₂-CHOH-SO₃H radicals, R₅ is chosen from alkyl radicals of an acid R₅-COOH present in coconut oil or in hydrolysed linseed oil, alkyl radicals, such as C₇, C₉, C₁₁ and C₁₃ alkyl radicals, a C₁₇ alkyl radical and its iso form, and unsaturated C₁₇ radical. These compounds are classified in the CTFA dictionary, 5th edition, 1993, under the names disodium cocoamphodiacetate, disodium lauroamphodiacetate, disodium caprylamphodiacetate, disodium capryloamphodiacetate, disodium cocoamphodipropionate, disodium lauroamphodipropionate, disodium caprylamphodipropionate, disodium capryloamphodipropionate, lauroamphodipropionic acid, and cocoamphodipropionic acid. Salts of diethyl aminopropyl cocoaspartamid can be also used.

[0092] Suitable cationic surfactants include, but are not limited to, the quaternary ammonium salts A) to D) as defined hereinafter:

A) Quaternary ammonium salts of general formula (VIII) below:




wherein X⁻ is an anion chosen from halides (chloride, bromide and iodide), (C₂-C₆)alkyl sulphates, such as methyl sulphate, phosphates, alkyl and alkylaryl sulphonates, and anions derived from organic acids, such as acetate and lactate, and wherein R₁ to R₄ are as below in i) or ii).

i) Radicals R₁ to R₃, which may be identical or different, are chosen from linear and branched aliphatic radicals comprising from 1 to 4 carbon atoms, and aromatic radicals such as aryl and alkylaryl. The aliphatic radicals may comprise at least one hetero atom such as oxygen, nitrogen, sulphur and halogens. The aliphatic radicals may be chosen from: alkyl, alkoxy and alkylamide radicals. R₄ is chosen from linear and branched alkyl radicals comprising from 16 to 30 carbon atoms. A suitable cationic surfactant is, for example, a behenyltrimethylammonium salt (for example chloride).


ii) Radicals R₁ and R₂, which may be identical or different, are chosen from linear and branched aliphatic radicals comprising from 1 to 4 carbon atoms, and aromatic radicals such as aryl and alkylaryl. The aliphatic radicals may comprise at least one hetero atom such as oxygen, nitrogen, sulphur and halogens. The aliphatic radicals may be chosen from alkyl, alkoxy, alkylamide and hydroxyalkyl radicals comprising from 1 to 4 carbon atoms. Radicals R₃ and R₄, which may be identical or different, are chosen from linear and branched alkyl radicals comprising from 12 to 30 carbon atoms, the said alkyl radicals comprise at least one function chosen from ester and amide functions. R₃ and R₄ may be chosen from (C₁₂-C₂₂)alkylamido(C₂-C₆)alkyl and (C₁₂-C₂₂) alkylacetate radicals. A suitable cationic surfactant is, for example, a dicetyldimethyl ammonium salt (for example chloride);

B) Quaternary ammonium salts of imidazolinium of formula (IX) below:




in which R_5 is chosen from alkenyl and alkyl radicals comprising from 8 to 30 carbon atoms, for example fatty acid derivatives of tallow, R_6 is chosen from a hydrogen atom, C_1 - C_4 alkyl radicals and alkenyl and alkyl radicals comprising from 8 to 30 carbon atoms, R_7 is chosen from C_1 - C_4 alkyl radicals, R_8 is chosen from a hydrogen atom and C_1 - C_4 alkyl radicals, and X^- is an anion chosen from halides, phosphates, acetates, lactates, alkyl sulphates, alkyl sulphonates and alkylaryl sulphonates. In one embodiment, R_5 and R_6 are, for example, a mixture of radicals chosen from alkenyl and alkyl radicals comprising from 12 to 21 carbon atoms, such as fatty acid derivatives of tallow, R_7 is methyl and R_8 is hydrogen. Such a product is, for example, Quaternium-27 (CTFA 1997) or Quaternium-83 (CTFA 1997), commercially available as "Rewoquat®" W75/W90/W75PG/W75HPG by Witco.

C) Diquaternary ammonium salts of formula (X):



in which R_9 is chosen from aliphatic radicals comprising from 16 to 30 carbon atoms, R_{10} , R_{11} , R_{12} , R_{13} and R_{14} , which may be identical or different, are chosen from hydrogen and alkyl radicals comprising from 1 to 4 carbon atoms, and X^- is an anion chosen from halides, acetates, phosphates, nitrates and methyl sulphates. Such diquaternary ammonium salts, for example, include propanetallowdium dichloride.

D) Quaternary ammonium salts comprising at least one ester function, of formula (XI) below:



in which: R_{15} is chosen from C_1 - C_6 alkyl radicals and C_1 - C_6 hydroxyalkyl and dihydroxyalkyl radicals; R_{16} is chosen from: a radical $\text{R}_{19}\text{C}(\text{O})-$, linear and branched, saturated and unsaturated C_1 - C_{22} hydrocarbon-based radicals R_{20} , and a hydrogen atom, R_{18} is chosen from: a radical $\text{R}_{21}\text{C}(\text{O})-$, linear and branched, saturated and unsaturated C_1 - C_6 hydrocarbon-based radicals R_{22} , and a hydrogen atom, R_{17} , R_{19} and R_{21} , which may be identical or different, are chosen from linear and branched, saturated and unsaturated C_7 - C_{21} hydrocarbon-based radicals; n , p and r , which may be identical or different, are chosen from integers ranging from 2 to 6; y is chosen from integers ranging from 1 to 10; x and z , which may be identical or different, are chosen from integers ranging from 0 to 10; X^- is an anion chosen from simple and complex, organic and inorganic anions; with the proviso that the sum $x+y+z$ is from 1 to 15, that when x is 0, then R_{16} is R_{20} and that when z is 0, then R_{18} is R_{22} . In one embodiment, the ammonium salts of formula (XI) can be used, in which: R_{15} is chosen from methyl and ethyl radicals, x and y are equal to 1; z is equal to 0 or 1; n , p and r are equal to 2; R_{16} is chosen from: a radical $\text{R}_{19}\text{C}(\text{O})-$, methyl, ethyl and C_{14} - C_{22} hydrocarbon-based radicals, and a hydrogen atom; R_{17} , R_{19} and R_{21} , which may be identical or different, are chosen from linear and branched, saturated and unsaturated C_7 - C_{21} , hydrocarbon-based radicals; R_{18} is chosen from: a radical $\text{R}_{21}\text{C}(\text{O})-$ and a hydrogen atom. Such compounds are commercially available as Dehyquart by Cognis, Stepanquat by Stepan, Noxamium by Ceca, and Rewoquat WE 18 by Rewo-Witco.

Ionic strength

[0093] The first and/or the second hair colouring compositions of the present invention may further have an ionic strength as defined herein of less than 1.35 mole/kg, alternatively from 0.10 to 0.75 mole/kg, alternatively from 0.20 to

0.60 mole/kg. Whilst not being bound by theory, it is believed that the ionic strength value may also affect the resultant viscosity and root adhesion properties of the composition. The ionic strength can be affected by salt resources such as the dyes, sodium sulphate, ammonium carbonate anti-oxidants and chelants such as EDDS. The dye tends to have the greatest effect on the ionic strength and thus the amounts added in order to provide any particular shade need to be considered in terms of ionic strength as well as dye outcome in order to prevent viscosity and root adhesion problems.

[0094] The ionic strength of the composition is a function of the concentration of all ions present in that solution and is determined according to the formula:

$$I = \frac{1}{2} \sum_{i=1}^n m_i z_i^2$$

where m_i = molality of ion i ($M = \text{mol}/\text{Kg H}_2\text{O}$), z_i = charge number of that ion, and the sum is taken over all ions in the solution. For example, for a 1:1 electrolyte such as sodium chloride, the ionic strength is equal to the concentration, but for MgSO_4 the ionic strength is four times higher. Generally multivalent ions contribute strongly to the ionic strength.

[0095] For example the ionic strength of a mixed 0.050 M Na_2SO_4 and 0.020 M NaCl solution is: $I = 1/2((2 \times (+1)^2 \times 0.050) + (+1)^2 \times 0.020 + (-2)^2 \times 0.050 + (-1)^2 \times 0.020) = 0.17 \text{ M}$.

Foam

[0096] The first and/or the second hair colouring compositions of the invention may be provided in the form of foam which is applied to the hair. Foam formation is typically achieved by the use of a foaming agent incorporated within the mixed composition (typically present in either the oxidizing composition or the dye composition or both) in combination with a manually operated foaming device. Such manually operated foaming devices are known in the art and include aerosols devices, squeeze foamers and pump foamers.

[0097] Suitable foaming agents includes surfactants such as anionic, nonionic and amphoteric surfactants, nonionic surfactants being preferred; polysaccharides (as described herein); polyvinyl pyrrolidone and copolymers thereof; acrylic polymers such as Acrylates copolymer (Aculyn 33) and Acrylates/Steareth-20 methacrylates (Aculyn 22); C12-C24 fatty acids such as stearates and mixtures thereof.

Viscosity

[0098] The developer compositions and the dye compositions may be, independently from one another, prepared as so called thin liquids or creams.

[0099] Each of the first and second hair colouring compositions may have a viscosity which induces a shear stress of from 20 to 200 Pa at 10 s^{-1} as measured according to the viscosity test method.

[0100] The first hair colouring composition may have a viscosity which induces a shear stress of from 60 to 200 Pa at 10 s^{-1} as measured according to the viscosity test method and the second hair colouring composition may have a viscosity which induces a shear stress of from 20 to 180 Pa at 10 s^{-1} as measured according to the viscosity test method.

[0101] Each of the first and second hair colouring compositions may have a viscosity which induces a shear stress of from 20 to 60 Pa at 10 s^{-1} , when the first and second hair colouring compositions are applied to the hair with a container to which a nozzle or a separate applicator device such as a comb or a brush is attached.

[0102] The first hair colouring composition may have a viscosity which induces a shear stress of from 30 to 200 Pa at 10 s^{-1} , or from 100 to 200 Pa at 10 s^{-1} or from 130 to 180 Pa at 10 s^{-1} when the non-diluted hair colouring composition is applied to the hair with a brush and bowl applicator. The diluted hair colouring composition may have a viscosity which induces a shear stress of from 20 to 180 Pa at 10 s^{-1} , or from 40 to 180 Pa at 10 s^{-1} , or from 70 to 170 Pa at 10 s^{-1} , when the first hair colouring composition is applied to the hair with a brush and bowl applicator or with the hands or fingers of the user.

[0103] Whilst not being bound by theory, it is believed that the provision of the first composition having viscosity values as described hereinabove enables the first composition to be applied directly to the roots without any dripping or running down the hair lengths and also enables the second composition to be easily applied and distributed along the entire remaining hair length with minimal to no dripping from the hair.

Application means

[0104] Both the first and second hair colouring compositions may be applied to the hair with a brush and bowl applicator.

Alternatively, the first hair colouring composition may be applied to the hair with a brush and bowl applicator whereas the second hair colouring composition may be applied to the hair with the hands and fingers of the user.

[0105] Alternatively, both the first and the second hair colouring compositions may be applied to the hair with a container to which a nozzle or a separate applicator device such as a comb is attached.

[0106] The application means may also include means which assist in achieving particular effects such as highlighting such as highlighting combs, brushes and tools, foils and highlighting caps. Additional application means technology can be used to assist in the penetration of the product into the hair. Examples of such technology include heating devices, ultraviolet light devices and ultrasound devices.

Hair colouring kit

[0107] A hair colouring kit may be used for carrying out the method for colouring hair described hereinbefore. The compositions comprised in the kit may comprise any of the ingredients disclosed in the hair colouring compositions section of this application.

[0108] The kit may comprise a first dye composition, a second dye composition, a first developer composition and a second developer composition. Each of the first and second dye compositions may comprise one or more oxidative dye precursors and/or one or more alkalizing agents and each of the first and second developer compositions may comprise one or more oxidizing agents.

[0109] In the embodiments, wherein the first and second developer compositions are the same, the kit may comprise only a first developer composition instead of a first and a second developer composition.

[0110] In the embodiments wherein the first and second dye compositions are the same, the kit may comprise only a first dye composition instead of a first and a second dye composition.

[0111] The first and/or the second developer compositions may comprise from 1% to 20% by weight of the oxidizing agents and the first and/or the second dye compositions may comprise from 0.005% to 10% by weight of the oxidative dye precursors and/or from 0.1% to 20% or 0.1% to 10% by weight of the alkalising agents.

[0112] The kit may further comprise a shampoo for use in rinsing the hair after colouring and/or a conditioner composition.

[0113] The kit may also comprise a color refresher composition. Such colour refresher composition may comprise at least one pre-formed dye and may be applied to the hair immediately after the oxidative colour. This is typically during the next wash cycle(s) from 1 day to 60 days after the original oxidative application. This colour refresher composition can be used to increase the initial colour obtained and/or boost the colour during the wash and style cycle until the next oxidative colouring or bleaching event.

[0114] The present invention may be utilized in a variety of packaging and dispensing devices. These dispensing devices can come in the form of separate devices which may be used independently or in combination with one another. Typically, the hair colouring or bleaching compositions are contained within separate single or multi compartment containers so that the compositions can be stored separately from one another before use. The compositions are then mixed together by a mixing means and then applied to the consumer's hair by an application means.

[0115] The first and/or the second developer compositions, the first and/or the second dye compositions may be provided as separate containers in the kit. The developer composition containers, the dye composition containers may be a bottle, a tube, an aerosol, or a sachet.

[0116] The first and/or the second developer compositions may be provided as a container such as a bottle, a tube, an aerosol, or a sachet and the first and/or second dye compositions may be provided in an additional compartment within the developer container or in a separate container which may be identical such as a dual sachet or aerosol systems for example or different such as a bottle and tube system.

[0117] The consumer may mix the developer compositions and the dye compositions by any means. This may simply involve the use of a mixing bowl into which the compositions are dispensed and then mixed, preferably using a mixing means such as a tool. Alternatively, it may involve the addition of one of the compositions into the container of the other composition (typically the dye composition is added to the developer composition), followed by manual shaking or mixing with a tool. Another system involves the perforation or displacement of a seal located between the separate compartments of the dye composition and developer composition within a single container or sachet followed by manual mixing within the container or in a separate and/or additional container.

[0118] The hair colouring kit may further comprise an applicator. The applicator may be a brush and bowl applicator. Alternatively, the applicator may be a nozzle which may be attached to one of the containers comprised in the kit in case the developer compositions and/or the dye composition are provided as separate containers in the kit or a separate applicator device such as a comb or a brush. Such combs and brushes can be adapted in order to achieve particular effects, whether it is quick and even coverage or root/hairline touch up, or highlights or streaks.

[0119] Alternatively, one of the containers may be provided with a comb attached to or instead of the dispensing nozzle whereby the product is dispensed through hollow tines and dispensing apertures located in the comb tines. The comb

tines may be provided with single or multiple openings along the tines to improve product application and evenness especially root to tip. Product dispensation can be achieved by mechanical pressure applied to the container for example delaminating bottles or any of the mechanisms described hereinabove. The comb may be provided on the container such as to facilitate easy application and may be positioned vertically (so called verticomb) or at an angle to allow the consumer to access all areas.

[0120] The volume of first and/or second developer compositions in the kit may be 10 mL to 120 mL, or from 40 mL to 70 mL, or from 55 mL to 65 mL. The volume of the first and/or second dye composition in the kit may be from 10 mL to 120 mL, or from 40 mL to 70 mL, or from 55 mL to 65 mL.

[0121] The compositions of the kit can be manufactured utilizing any one of the standard approaches, these include a) 'Oil in water' process, b) 'Phase Inversion' process and c) 'One-pot' process. For example, when using "oil in water" process, surfactants of the present invention are added to approximately 50% of total water amount of the compositions at about 90°C, homogenized for 15 to 30 min, then cooled to room temperature to form a premix; this premix is then mixed cold with remaining amounts of water, other optional components and/or oxidizing agents, thus forming the developer compositions and dye compositions of the above described colouring kit.

[0122] The kit may further comprise a set of instructions comprising instructing the user to colour its hair according to the method defined hereinbefore. The set of instruction may comprise:

i) mixing a first dye composition with a first developer composition to obtain a first hair colouring composition, wherein the first dye composition comprises one or more oxidative dye precursors and/or one or more alkalizing agents and the first developer composition comprises one or more oxidizing agents;

ii) applying a portion or all of the first hair colouring composition to the hair roots and to one or more portions of the hair lengths and tips extending from the hair roots;

iii) mixing a second dye composition with a second developer composition to obtain a second hair colouring composition, wherein the second dye composition comprises one or more oxidative dye precursors and/or one or more alkalizing agents and the second developer composition comprises one or more oxidizing agents, wherein the concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents and/or the concentration of the oxidizing agents comprised in the second hair colouring composition is lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents and/or the oxidizing agents comprised in the first hair colouring composition;

iv) applying a portion or all of the second hair colouring composition to the hair;

v) optionally rinsing the hair.

[0123] The set of instructions may comprise any additional step which is disclosed hereinbefore in the method for colouring hair section of the application.

Viscosity test method:

[0124] The viscosity of a composition is measured using a TA Instruments AR 2000 Rheometer or equivalent device equipped with a Peltier plate and a 6cm flat acrylic plate with cross hatchings. The instrument is calibrated according to the manufacturer's instructions and the Peltier plate is set at 25.0°C. The cone is raised to a position approximately 4.5 cm above the plate.

[0125] Immediately after the mixing, approximately 10g of the mixture is transferred gently onto the centre of the Peltier plate using a spatula. The cone is lowered to obtain the specified gap between the tip of the cone and the upper surface of the Peltier plate. The gap setting is specified by the manufacturer of the cone and is typically approximately 1000 microns. The rheometer is programmed to operate in rotational mode with the shear stress ramped from 0.1 to 600 Pa over a period of 4 minutes, termination at 1000 reciprocal seconds. Rotation is initiated immediately after the specified gap is established. Viscosity data collected during the measurement period are shear stress (Pa) plotted as a function of shear rate (s^{-1}).

Examples:

[0126] The following are non-limiting examples of compositions of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention, which would be recognized by one of ordinary skill in the art. All concentrations are listed as weight percent, unless otherwise specified.

Formulations according to the present invention:

[0127] In the following section all the values are expressed in percentages by weight of the total compositions.

Dye composition formulations

[0128]

	Dye Composition 1 7/0	Dye Composition 2 7/0	Dye Composition 3 7/0
Toluene-2,5-diamine Sulfate	0.69	0.69	0.69
4-Aminophenol	0.083	0.083	0.083
Resorcinol	0.337	0.337	0.337
2,4-Diaminophenoxyethanol HCL	0.01	0.01	0.01
m-Aminophenol	0.074	0.074	0.074
2-Methyl-5-hydroxyethylamino- phenol (Paox)	0.02	0.02	0.02
Ethanolamine		4.25	
Ammonium Hydroxide 25%	7.28		3
Hair colouring chassis	91.506	94.536	95.786

Dye composition 4	
Toluene-2,5-diamine Sulfate	0.674
Hydroxyethyl-3,4-methylenedioxyaniline HCL	0.003
m-Aminophenol	0.024
Resorcinol	0.280
2-Methylresorcino 1	0.036
Ammonium Hydroxide 25%	7.770
Hair colouring chassis	91.213

Dye composition 5	
Toluene-2,5-Diamine Sulfate	1.040
m-Aminophenol	0.050
Resorcinol	0.390
2-Methylresorcino 1	0.090
Ethanolamine	3.910
Hair colouring chassis	94.52

Developer composition formulations

[0129]

EP 2 881 146 A2

Developer composition	1	2	3	4	5
Water Purified	Up to 100	Up to 100	Up to 100	Up to 100	Up to 100
Disodium EDTA.	0.04	0.04	0.04	0.04	0.04
Etidronic Acid	0.08	0.08	0.08	0.08	0.08
Aculyn 33	9.00	9.00	9.00	9.00	9.00
Acrylates Steareth-20 methacrylate copolymer	0.10	0.10	0.10	0.10	0.10
Hydrogen peroxide solution, 50%	12.00	18.00	3.8	6	8
Simethicone Emulsion	0.01	0.01	0.01	0.01	0.01

Developer composition	6	7	8	9	10
Water Purified	Up to 100	Up to 100	Up to 100	Up to 100	Up to 100
Etidronic Acid	0.01	0.01	0.01	0.01	0.01
Hydrogen peroxide solution, 50%	12.00	18.00	3.8	6	8
Cetearyl alcohol	3.4	3.4	3.4	3.4	3.4
Ceteareth-25	0.8	0.8	0.8	0.8	0.8
Salicyclic acid	0.1	0.1	0.1	0.1	0.1
Disodium phosphate	0.08	0.08	0.08	0.08	0.08

Example 1:

[0130] Dye composition 1 is mixed with developer composition 1 or 6 at a mixing ratio of 1:1 to obtain a first hair colouring composition. Dye composition 1 is mixed with developer composition 3 or 8 at a mixing ratio of 1:1 to obtain a second hair colouring composition. The first composition is applied to the hair roots and to a portion of the hair lengths and tips extending from the hair roots and left for 20 min on the hair. Then, the second composition is applied to the rest of the hair lengths and tips which are free of the first hair colouring composition. After 10 min, the hair is rinsed to remove the compositions from the hair.

Example 2:

[0131] Same experimental protocol as for example 1, wherein dye composition 1 is mixed with developer composition 4 or 9 instead of developer composition 3 or 8 to obtain a second hair colouring composition.

Example 3:

[0132] Same experimental protocol as for example 1, wherein dye composition 1 is mixed with developer composition 5 or 10 instead of developer composition 3 or 8 to obtain a second hair colouring composition.

Example 4:

[0133] Same experimental protocol as for example 1, 2 or 3, wherein dye composition 2 is used instead of dye composition 1.

Example 5:

[0134] Same experimental protocol as for example 1, 2 or 3, wherein dye composition 3 is used instead of dye composition 1.

Example 6:

[0135] Dye composition 4 is mixed with developer composition 1 or 6 at a mixing ratio of 1:1 to obtain a first hair colouring composition. Dye composition 5 is mixed with developer composition 3 or 8 at a mixing ratio of 1:1 to obtain a second hair colouring composition. The first composition is applied to the hair roots and to a portion of the hair lengths and tips extending from the hair roots and left for 20 min on the hair. Then, the second composition is applied to the rest of the hair lengths and tips which are free of the first hair colouring composition. After 10 min, the hair is rinsed to remove the compositions from the hair.

Example 7:

[0136] Same experimental protocol as for example 6, wherein dye composition 5 is mixed with developer composition 4 or 9 instead of developer composition 3 or 8 to obtain a second hair colouring composition.

Example 8:

[0137] Same experimental protocol as for example 6, wherein dye composition 5 is mixed with developer composition 5 or 10 instead of developer composition 3 or 8 to obtain a second hair colouring composition.

[0138] The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about 40 mm."

Claims

1. A method for colouring hair comprising the steps of:

- i) mixing a first dye composition with a first developer composition to obtain a first hair colouring composition, wherein the first dye composition comprises one or more oxidative dye precursors and/or one or more alkalizing agents and the first developer composition comprises one or more oxidizing agents;
- ii) applying a portion or all of the first hair colouring composition to the hair roots and to one or more portions of the hair lengths and tips extending from the hair roots;
- iii) mixing a second dye composition with a second developer composition to obtain a second hair colouring composition, wherein the second dye composition comprises one or more oxidative dye precursors and/or one or more alkalizing agents and the second developer composition comprises one or more oxidizing agents, wherein the concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents and/or the concentration of the oxidizing agents comprised in the second hair colouring composition is lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents and/or the oxidizing agents comprised in the first hair colouring composition or providing a second hair colouring composition comprising one or more direct dyes;
- iv) applying a portion or all of the second hair colouring composition to the hair;
- v) optionally rinsing the hair.

2. The method according to claim 1, wherein in step iv), the portion or all of the second hair colouring composition is applied to the rest of the hair lengths and tips which are free of the first hair colouring composition and optionally to one or more portions of the hair to which the first hair colouring composition has been applied in step ii).

3. The method according to claim 1 or 2, wherein:

- a. the concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents comprised in the second dye composition is lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents comprised in the first dye composition, preferably wherein the oxidative dye precursors and/or the alkalizing agents comprised in the first and second dye compositions are the same and/or
- b. the concentration of oxidizing agents comprised in the second developer composition is lower than the concentration of oxidizing agents comprised in the first developer composition, preferably wherein the oxidizing agents comprised in the first and second dye compositions are the same.

4. The method according to any of the preceding claims, wherein:

- a. the first and/or second developer compositions comprise from 1 to 20% by weight of the oxidizing agents, and/or
- b. the first and/or second dye compositions comprise from 0.005% to 10% by weight of the oxidative dye precursors and/or from 0.1% to 20% by weight of the alkalizing agents.

5. The method according to any of the preceding claims, wherein the first dye composition is mixed with the first developer composition and/or the second dye composition is mixed with the second developer composition at a mixing ratio of from 2:1 to 1:3, preferably from 1:1 to 1:2, preferably wherein the mixing ratios of the first dye composition with the first developer composition and of the second dye composition with the second developer composition are the same.

6. The method according to any of the preceding claims, wherein the first and second developer compositions are the same and the concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents comprised in the second dye composition is lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents comprised in the first dye composition, preferably wherein the oxidative dye precursors and/or the alkalizing agents comprised in the first and second dye compositions are the same.

7. The method according to claim 6, wherein the concentration of the oxidative dye precursors and/or the concentration of the alkalizing agents comprised in the second dye composition is at least 1%, preferably at least 5% lower than the concentration of respectively the oxidative dye precursors and/or the alkalizing agents comprised in the first dye composition.

8. The method according to any of claims 1 to 5, wherein the first and second dye compositions are the same and the concentration of oxidizing agents comprised in the second developer composition is lower than the concentration of oxidizing agents comprised in the first developer composition, preferably wherein the oxidizing agents comprised in the first and second developer compositions are the same.

9. The method according to claim 8, wherein the concentration of the oxidizing agents comprised in the second developer composition is at least 15%, preferably at least 25% lower than the concentration of the oxidizing agents comprised in the first developer composition.

10. The method according to any of the preceding claims wherein:

- a. the method further comprise the steps of waiting for a time period of from 10 min to 40 min, preferably 15 min to 30 min, more preferably 20 min, which is performed between steps ii) and iv) and waiting for a time period of from 5 min to 20 min, preferably 10 min which is performed between steps iv) and v) or
- b. step iii) is performed immediately after step ii) and step iv) is performed immediately after step iii) and the method further comprises the step of waiting for a time period of from 5 min to 40 min, preferably 20 min to 40 min, more preferably 30 min, which is performed between steps iv) and v).

11. The method according to any of the preceding claims, wherein each of the first and second hair colouring compositions has a viscosity which induces a shear stress of from 20 to 200 Pa at 10 s^{-1} as measured according to the viscosity test method.

12. The method according to any of the preceding claims, wherein the first hair colouring composition has a viscosity which induces a shear stress of from 60 to 200 Pa at 10 s^{-1} as measured according to the viscosity test method and the second hair colouring composition has a viscosity which induces a shear stress of from 20 to 180 Pa at 10 s^{-1} as measured according to the viscosity test method.

13. The method according to any of the preceding claims, wherein the first and second hair colouring compositions are applied to the hair with a brush and bowl applicator or a container to which a nozzle or a separate applicator device such as a comb or a brush is attached.

14. The method according to any of the preceding claims, wherein each of the first and second hair colouring compositions comprises at least one gel network thickener system.

15. The method according to any of the preceding claims, wherein the first developer composition and/or the second

EP 2 881 146 A2

developer composition comprises a thickener selected from acrylates copolymer, hydrophobically modified acrylate copolymers and mixtures thereof.

5

10

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2006060598A1 A [0075]
- US 2528378 A [0091]
- US 2781354 A [0091]

Non-patent literature cited in the description

- International Cosmetics Ingredient Dictionary and Handbook. The Cosmetics, Toiletry, and Fragrance Association, vol. 2 [0028]
- **AE MARTELL ; RM SMITH.** Critical Stability Constants. Plenum Press, 1974, vol. 1 [0050]
- **AE MARTELL ; RD HANCOCK.** Metal Complexes in Aqueous Solution. Plenum Press, 1996 [0050]
- **ROHM ; HAAS.** Permulen TR1, Carbopol 2020, Carbopol Ultrez-21. National Starch, 2001 [0071]
- Encyclopedia of Chemical Technology. 1982, vol. 3, 896-900 [0073]
- Encyclopedia of Chemical Technology. vol. 15, 439-458 [0073]
- **E. A. MACGREGOR ; C. T. GREENWOOD.** Polymers in Nature. John Wiley & Sons, 1980, 240-328 [0073]
- Industrial Gums-Polysaccharides and their Derivatives. Academic Press Inc, [0073]
- **M. R. PORTER.** Handbook of Surfactants. Blackie & Son, 1991, 116-178 [0090]