



(11) **EP 2 884 857 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 91

(51) Int Cl.:
A23L 19/00 ^(2016.01) **A23B 7/16** ^(2006.01)

(86) International application number:
PCT/EP2013/067080

(48) Corrigendum issued on:
05.04.2017 Bulletin 2017/14

(87) International publication number:
WO 2014/027063 (20.02.2014 Gazette 2014/08)

(45) Date of publication and mention
of the grant of the patent:
02.11.2016 Bulletin 2016/44

(21) Application number: **13753134.9**

(22) Date of filing: **15.08.2013**

(54) **METHOD FOR PRODUCING A FOOD PRODUCT**

VERFAHREN ZUR HERSTELLUNG EINES LEBENSMITTELPRODUKTS

PROCÉDÉ DE PRODUCTION D'UN PRODUIT ALIMENTAIRE

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR

(56) References cited:
WO-A1-2004/026042 WO-A1-2007/077546
WO-A1-2008/110513 WO-A1-2010/070010
WO-A2-2008/151807 US-A1- 2007 141 227

(30) Priority: **17.08.2012 EP 12180858**

(43) Date of publication of application:
24.06.2015 Bulletin 2015/26

(73) Proprietor: **Novozymes A/S**
2880 Bagsvaerd (DK)

(72) Inventors:
• **HENDRIKSEN, Hanne Vang**
2840 Holte (DK)
• **MATSUI, Tomoko**
Chiba-shi
Chiba 261-0001 (JP)

- **HENDRIKSEN H V ET AL: "Evaluating the potential for enzymatic acrylamide mitigation in a range of food products using an asparaginase from Aspergillus oryzae", JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY, AMERICAN CHEMICAL SOCIETY, US, vol. 57, no. 10, 27 March 2009 (2009-03-27), pages 4168-4176, XP002575849, ISSN: 0021-8561, DOI: 10.1021/JF900174Q [retrieved on 2009-04-23]**
- **PEDRESCHI ET AL: "The effect of asparaginase on acrylamide formation in French fries", FOOD CHEMISTRY, ELSEVIER LTD, NL, vol. 109, no. 2, 28 December 2007 (2007-12-28), pages 386-392, XP022505550, ISSN: 0308-8146, DOI: 10.1016/J.FOODCHEM.2007.12.057**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 884 857 B9

Description**Reference to sequence listing**

- 5 [0001] This application contains a Sequence Listing in computer readable form. The computer readable form is incorporated herein by reference.

FIELD OF THE INVENTION

- 10 [0002] The present invention relates to a method for producing a heat-treated vegetable-based food product which comprises treatment with asparaginase.

BACKGROUND OF THE INVENTION

- 15 [0003] It is well known that acrylamide formation in heated food products may be reduced by a treatment reducing the amount of asparagine in the food materials, such as by subjecting the food materials to the action of the enzyme asparaginase (see e.g. WO2004/026042).

- [0004] To fit into the production line of an industrial food product, treatment with asparaginase should preferentially take place during an existing step in the production process. It has so far been a challenge to fit asparaginase treatment into the existing industrial production process of, e.g., French fries. One reason being that known asparaginase enzymes may not be stable/active at the typical process temperatures used in an industrial process for production of French fries.

- [0005] WO2004/026042 suggests that in the production of French fries, asparaginase may be added at any suitable stage of the process, such as, e.g., during blanching. However, for an asparaginase to be added during blanching, it should retain a major part of its activity at the high temperature used for blanching. Commercial asparaginase enzymes on the market today are inactivated at such high temperature.

- [0006] WO 2008/110513 discloses a number of thermostable variants of asparaginase from *Aspergillus oryzae*. However, none of these variants are thermostable enough to be added, e.g., during blanching. Treatment of potato sticks for French fries for 5 min at 60°C is disclosed for one of the variants. But 60°C is not an optimal temperature to be used industrially because of potential microbial growth at this temperature.

- 30 [0007] WO2008/151807 discloses a hyper-thermostable asparaginase and its use in food production. Such enzyme could possibly be added, e.g., during blanching. Use of a hyper-thermostable asparaginase in the industrial production of French fries is not optimal though, since such enzyme is not active during subsequent steps in the production process which typically take place at lower temperature.

- [0008] WO 2010/070010 suggests contacting with asparaginase at temperatures as high as 65°C in the presence of, e.g., sodium acid pyrophosphate (SAPP). But even 65°C is not an optimal temperature to be used industrially because of potential microbial growth around this temperature.

- [0009] US 2007/0141227 A1 discloses contacting of, e.g., potato slices with a leaching solution to extract asparagine. In one embodiment, the post-wash extract which contains leached out components is cooled to prevent the enzyme from becoming inactivated and mixed with asparaginase which removes a majority of the asparagine from the extract.

- 40 [0010] WO 2007/077546 discloses a method for reducing the level of asparagine in an asparagine-containing food material which comprises heating of the food material followed by cooling to a temperature of less than about 70°C and adding an asparagine reducing enzyme.

- [0011] It is an object of the present invention to provide an industrially applicable method for producing a heat-treated vegetable-based food product, such as French fries, having a reduced level of acrylamide, where asparaginase treatment of the vegetable-based food material, such as the potato sticks, fits into the existing production process typically applied by the industry.

SUMMARY OF THE INVENTION

- 50 [0012] The present invention provides a method for producing a heat-treated vegetable-based food product comprising:

- (a) contacting of a vegetable-based food material with asparaginase at a temperature of 67-75°C;
 (b) drying the vegetable-based food material at an air temperature of 40 to 90°C; and
 (c) heat-treating the asparaginase treated vegetable-based food material to obtain the heat-treated vegetable-based food product;

wherein the asparaginase has a residual activity after 4 hours' incubation in deionised water with 0.5% SAPP at 70°C, pH 5, of at least 20%, preferably at least 40%, more preferably at least 60%, even more preferably at least 80%, of the

activity without such incubation; and wherein the asparaginase has an activity at 35°C, pH 6, of at least 20%, preferably at least 30%, more preferably at least 35%, of its activity at 50°C, pH 6.

[0013] The invention further relates to use of asparaginase for treatment of a vegetable-based food material at a temperature of 67-75°C; wherein the asparaginase has a residual activity after 4 hours' incubation in deionised water with 0.5% SAPP at 70°C, pH 5, of at least 20%, preferably at least 40%, more preferably at least 60%, even more preferably at least 80%, of the activity without such incubation; and wherein the asparaginase has an activity at 35°C, pH 6, of at least 20%, preferably at least 30%, more preferably at least 35%, of its activity at 50°C, pH 6.

[0014] A number of asparaginase enzymes are provided which are stable at high temperature, i.e. they are still active after prolonged incubation at, e.g., 67-75°C, and also retain their activity at lower temperature, e.g., at 40-60°C. However, any asparaginase having such properties would be useful in the method of the present invention.

[0015] In the method of the present invention, such asparaginase may be added directly, e.g., to the existing chemical dip in the industrial production process of French fries with no process changes required. In the chemical dip often applied by industry, blanched potato strips may be dipped, e.g., in sodium acid pyrophosphate (SAPP) and optionally glucose to control colour. The temperature of the dipping solution is typically kept around 70°C to avoid microbial infection. The chemical dip is typically followed by a drying step. Although the drying step may be performed at high temperature, such as, e.g., about 85°C, the surface temperature of the potato sticks is only about 40-60°C. In the method of the present invention, the asparaginase is active during the drying step. Thus, full effect of the asparaginase treatment is obtained even if the chemical dip is rather short.

[0016] Preferably, the asparaginase is to be active in a pH interval of pH 5-6. pH of a SAPP dipping solution will initially (in pure water) be around 4.8 but gradually increase to 5.5-6 due to buffering from the potatoes.

DETAILED DESCRIPTION OF THE INVENTION

[0017] SEQ ID NO: 1 is the nucleotide sequence of the asparaginase gene from *Aspergillus oryzae*. Based on N-terminal sequencing of the mature asparaginase encoded, nucleotides 1-78 encodes the signal sequence. SEQ ID NO: 2 is the amino acid sequence of asparaginase from *Aspergillus oryzae*. Based on N-terminal sequencing, amino acids 1-26 is the signal sequence. SEQ ID NO: 3 is the nucleotide sequence of a gene encoding a variant of *Aspergillus oryzae* asparaginase having the following substitutions: N70K S307A A323R T327V A349Q S351A V353I. SEQ ID NO: 4 is the amino acid sequence of a variant of *Aspergillus oryzae* asparaginase having the following substitutions: N70K S307A A323R T327V A349Q S351A V353I. SEQ ID NO: 5 is the nucleotide sequence of a gene encoding a variant of *Aspergillus oryzae* asparaginase having the following substitutions: N70K A323R T327V A349Q S351A V353I. SEQ ID NO: 6 is the amino acid sequence of a variant of *Aspergillus oryzae* asparaginase having the following substitutions: N70K A323R T327V A349Q S351A V353I.

[0018] The present invention relates to a method for producing a heat-treated vegetable-based food product comprising:

- (a) contacting of a vegetable-based food material with asparaginase at a temperature of 67-75°C;
- (b) drying the vegetable-based food material under conditions where the asparaginase is active; and
- (c) heat-treating the asparaginase treated vegetable-based food material to obtain the heat-treated vegetable-based food product.

[0019] The vegetable-based food material may be any food material based on vegetables which is to be heat-treated. It may be derived from a vegetable tuber or root such as but not limited to the group consisting of potato, sweet potato, yams, yam bean, parsnip, parsley root, Jerusalem artichoke, carrot, radish, turnip, and cassava.

[0020] The processing of a vegetable tuber or root into a suitable vegetable-based food material may comprise, e.g., rinsing, washing, peeling, cutting, etc., such as to produce tuber or root pieces, e.g., potato pieces, of any size and/or shape, e.g., the form of wedges, sticks or slices, e.g., of a size and shape suitable for further processing into a heat-treated vegetable-based food product such as, e.g., French fries.

[0021] In the context of the present invention, French fries is meant to encompass both the final fries ready for consumption and a par-fried pre-product which is to be finally fried or baked before being consumed.

[0022] In a preferred embodiment, the vegetable-based food material is a potato material such as potato pieces, e.g., potato wedges, potato sticks or potato slices. In a more preferred embodiment, the vegetable-based food material is potato wedges or potato sticks. In an even more preferred embodiment, the vegetable-based food material is potato sticks and the heat-treated vegetable-based food product is French fries.

[0023] Preferably, the vegetable-based food material has been blanched before step (a). Blanching may be performed by any method known in the art, e.g., by wet blanching, steam blanching, microwave blanching or infrared blanching.

[0024] In a typical industrial production of French fries, potatoes are initially washed, sorted, steam peeled and cut. Following cutting, the potato sticks are blanched in 2 to 3 sequential steps typically at 65-90°C for 10-30 min. Blanching is done to inactivate the endogenous enzymes in the potato, to partially cook the potato and to leach out reducing sugars

to prevent excessive browning of the final product. After blanching, the potato strips may quickly be dipped, e.g. for 20-180 seconds, in a warm phosphate salt solution, e.g., a warm solution of sodium acid pyrophosphate (SAPP), to prevent greying of the final product. The dip is optionally combined with a dip in glucose to control the final colour. The potatoes may be dried in a drier with hot circulating air at 45-95°C for 5-20 minutes giving a weight loss of 5-25%. Finally,

the potato sticks are parfried before being quick-frozen and packed. Final frying is done at the restaurant or by consumers. [0025] In the method of the present invention, the contacting with the asparaginase may preferably be combined with dipping in warm phosphate salt solution. I.e., the vegetable based food material, preferably the potato wedges or the potato sticks, more preferably the blanched potato wedges or the blanched potato sticks, is/are dipped or incubated into a solution comprising asparaginase and, e.g., SAPP in one dip bath at a temperature of 67-75°C followed by drying.

Alternatively, the contacting may be performed by spraying the vegetable-based food material, preferably the potato wedges or the potato sticks, more preferably the blanched potato wedges or the blanched potato sticks, with a solution comprising asparaginase and, e.g., SAPP at a temperature of 67-75°C followed by drying. [0026] In a preferred embodiment, step (a) of the method of the invention is dipping vegetable based food material, preferably potato wedges or potato sticks, more preferably blanched potato wedges or blanched potato sticks, into an asparaginase solution having a temperature of 67-75°C, preferably 68-72°C, more preferably 70°C. The asparaginase solution may comprise asparaginase at a concentration of 1,000 to 100,000 ASNU/L, preferably 5,000 to 50,000 ASNU/L, more preferably 10,000 to 30,000 ASNU/L.

[0027] In a more preferred embodiment, the asparaginase solution further comprises SAPP at a concentration of 0.05 to 5%, preferably 0.1 to 1.5%, more preferably 0.5 to 1%. The asparaginase solution may further comprises glucose, e.g., at a concentration of 0.5 to 2%.

[0028] In a preferred embodiment, dipping is performed for a time of 1 second to 10 minutes, preferably 30 seconds to 3 minutes.

[0029] The drying in step (b) is to be performed at an air temperature of 40 to 90°C, preferably 55 to 85°C, more preferably 65 to 75°C, i.e. under conditions where the asparaginase is active. Drying may be performed in an oven where temperature, humidity and/or air flow can be adjusted to the level(s) desired. Drying may be performed, e.g., for between 5 and 30 minutes.

[0030] Preferably, drying is performed so that the surface temperature of the vegetable-based food material, such as the (blanched) potato wedges or potato sticks, during drying in step (b) is 25 to 60°C, preferably 35 to 50°C.

[0031] In a preferred embodiment, the acrylamide content of the heat-treated vegetable-based food product is at least 25% reduced, preferably at least 30%, at least 35%, at least 40%, at least 45% or at least 50% reduced, compared to the acrylamide content of a heat-treated vegetable-based food product produced by a similar method without the addition of asparaginase.

[0032] In a more preferred embodiment, the vegetable-based food material is potato sticks, the heat-treated vegetable-based food product is French fries, and the acrylamide content of the French fries is at least 25% reduced, preferably at least 30%, at least 35%, at least 40%, at least 45% or at least 50% reduced, compared to the acrylamide content of French fries produced by a similar method without the addition of asparaginase.

[0033] The acrylamide content may be determined by any method known in the art, e.g., by the method described in Example 3.

[0034] The invention further relates to a use of asparaginase for treatment of a vegetable-based food material at a temperature of 67-75°C.

[0035] An asparaginase in the context of the present invention means an enzyme having asparaginase activity, i.e., an enzyme that catalyzes the hydrolysis of asparagine to aspartic acid (EC 3.5.1.1).

[0036] Asparaginase activity may be determined according to one of the asparaginase activity assays described in the Examples, e.g., by the ASNU assay. In one embodiment, an asparaginase to be used in the method of the present invention has at least 20%, e.g., at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 100% of the asparaginase activity of the mature polypeptide of SEQ ID NO: 2 when measured at pH 7 and at 37°C.

[0037] The asparaginase may be a microbial asparaginase, e.g., an asparaginase derived from a bacterium, an archaeon or a fungus. It may be a wild type asparaginase, i.e., an asparaginase found in nature, or it may be a variant asparaginase, i.e., an asparaginase comprising an alteration, i.e., a substitution, insertion, and/or deletion, at one or more (e.g., several) positions compared to a parent asparaginase from which it may have been derived. A substitution means replacement of the amino acid occupying a position with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding an amino acid adjacent to and immediately following the amino acid occupying a position.

[0038] An asparaginase to be used according to the invention has a residual activity after 4 hours' incubation in deionised water with 0.5% SAPP at 70°C, pH 5, of at least 20%, preferably at least 40%, more preferably at least 60%, even more preferably at least 80%, of the activity without such incubation. The asparaginase activity may be determined by the rASNU assay as described in the Examples. It may be determined at pH 6 and at 37°C.

[0039] Preferably, an asparaginase to be used according to the invention has a residual activity after 4 hours' incubation in deionised water with 0.5% SAPP at 72°C, pH 5, of at least 5%, preferably at least 10%, more preferably at least 20%, even more preferably at least 40%, such as at least 60% or at least 80%, of the activity without such incubation.

[0040] Preferably, an asparaginase to be used according to the invention has a residual activity after 4 hours' incubation in deionised water with 0.5% SAPP at 74°C, pH 5, of at least 1%, preferably at least 5%, more preferably at least 10%, even more preferably at least 20%, such as at least 40% or at least 60%, of the activity without such incubation.

[0041] An asparaginase to be used according to the invention has an activity at 35°C, pH 6, of at least 20%, preferably at least 30%, more preferably at least 35%, of its activity at 50°C, pH 6.

[0042] Preferably, the asparaginase has an activity at 70°C, pH 6, of at least 20%, preferably preferably at least 25%, more preferably at least 30%, of its activity at 50°C, pH 6.

[0043] More preferably, the asparaginase has (i) an activity at 35°C, pH 6, of at least 20%, preferably at least 30%, more preferably at least 35%, of its activity at 50°C, pH 6, and (ii) an activity at 70°C, pH 6, of at least 20%, preferably at least 25%, more preferably at least 30%, of its activity at 50°C, pH 6.

[0044] Preferably, the asparaginase has an activity at pH 5 of at least 20%, preferably at least 30%, more preferably at least 40%, even more preferably at least 50%, of its activity at pH 6.5. The activity may be determined at 50°C.

[0045] More preferably, the asparaginase has (i) an activity at 35°C, pH 6, of at least 20%, preferably at least 30%, more preferably at least 35%, of its activity at 50°C, pH 6, (ii) an activity at 70°C, pH 6, of at least 20%, preferably at least 25%, more preferably at least 30%, of its activity at 50°C, pH 6, and (iii) an activity at pH 5, 50°C, of at least 20%, preferably at least 30%, more preferably at least 40%, even more preferably at least 50%, of its activity at pH 6.5, 50°C.

[0046] Preferably, the asparaginase has a residual activity after 2 hours' incubation in deionised water with 0.5% SAPP at pH 4 and 70°C of at least 20%, preferably at least 30%, more preferably at least 40%, even more preferably at least 50%, of the activity without such incubation.

[0047] In one embodiment, the asparaginase is obtained from, or is a variant of a parent asparaginase obtained from, a microorganism of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the parent encoded by a polynucleotide is produced by the source or by a strain in which the polynucleotide from the source has been inserted. In one embodiment, the parent is secreted extracellularly.

[0048] The asparaginase or its parent may be a fungal asparaginase. It may be a filamentous fungal asparaginase such as an *Acremonium*, *Agaricus*, *Alternaria*, *Aspergillus*, *Aureobasidium*, *Botryosphaeria*, *Botryotinia*, *Ceriporiopsis*, *Chaetomidium*, *Chrysosporium*, *Claviceps*, *Cochliobolus*, *Coprinopsis*, *Coptotermes*, *Corynascus*, *Cryphonectria*, *Cryptococcus*, *Diplodia*, *Exidia*, *Filibasidium*, *Fusarium*, *Gibberella*, *Holomastigotoides*, *Humicola*, *Irpex*, *Lentinula*, *Leptosphaeria*, *Magnaporthe*, *Melanocarpus*, *Meripilus*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neosartorya*, *Neurospora*, *Paezilomyces*, *Penicillium*, *Phanerochaete*, *Piromyces*, *Poitrasia*, *Pseudoplectania*, *Pseudotrichonympha*, *Rhizomucor*, *Schizophyllum*, *Sclerotinia*, *Scytalidium*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trichoderma*, *Trichophaea*, *Verticillium*, *Volvariella*, or *Xylaria* asparaginase.

[0049] In one embodiment, the asparaginase or its parent is an *Acremonium cellulolyticus*, *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Aspergillus terreus*, *Botryotinia fuckeliana*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola grisea*, *Humicola insolens*, *Humicola lanuginosa*, *Irpex lacteus*, *Mucor miehei*, *Myceliophthora thermophila*, *Neosartorya fischeri*, *Neurospora crassa*, *Penicillium chrysogenum*, *Penicillium citrinum*, *Penicillium funiculosum*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Sclerotinia sclerotiorum*, *Thielavia achromatica*, *Thielavia albolomyces*, *Thielavia albopilosa*, *Thielavia australeinsis*, *Thielavia fimeti*, *Thielavia microspora*, *Thielavia ovispora*, *Thielavia peruviana*, *Thielavia setosa*, *Thielavia spededonium*, *Thielavia subthermophila*, *Thielavia terrestris*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* asparaginase.

[0050] In one preferred embodiment, the asparaginase is a variant of a parent asparaginase obtainable from *Aspergillus*, e.g., from *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae* or *Aspergillus terreus*. Preferably, it is a variant having an increased thermostability as compared to its parent. The thermostability may be determined as described in Example 1.

[0051] In another preferred embodiment, the asparaginase is a variant of a parent asparaginase obtainable from *Aspergillus oryzae*, e.g., the asparaginase of SEQ ID NO: 2 or the mature polypeptide thereof, or a variant of a parent asparaginase obtainable from *Aspergillus niger*, e.g., the asparaginase disclosed in WO2004/030468.

[0052] In another preferred embodiment, the asparaginase has an amino acid sequence which is at least 50% identical to any of SEQ ID NOs: 2 or 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least

90%, at least 95% or at least 98% identical to any of SEQ ID NOs: 2 or 4.

[0053] In another preferred embodiment, the asparaginase is a variant of a parent asparaginase having an amino acid sequence which is at least 50% identical to any of SEQ ID NOs: 2 or 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or 100% identical to any of SEQ ID NOs: 2 or 4.

[0054] In another preferred embodiment, the asparaginase is an asparaginase variant comprising a substitution at one or more positions corresponding to positions 122, 140, 197, 238, 239, 240, 241, 253, 258, 259, 297 or 373 of SEQ ID NO: 2, and having an amino acid sequence which is at least 50% identical to any of SEQ ID NOs: 2 or 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% identical to any of SEQ ID NOs: 2 or 4.

[0055] In another preferred embodiment, the asparaginase is an asparaginase variant comprising (i) one or more of the following substitutions, wherein each position corresponds to a position in SEQ ID NO: 2: T71C, T74A, T74C, K122A, K122R, V139G, T140D, K194L, D197E, I228M, S238C, N239C, K240R, P241E, K253R, I258V, I258Y, R259C, R259V, S297V, S299A, T314A, P333L, S334P, S334W, E337S, S338G, S338W, G356D, K363R or E373H, or (ii) a deletion at one or more positions corresponding to positions 27, 28 or 29 of SEQ ID NO: 2; wherein the asparaginase variant has an amino acid sequence which is at least 50% identical to any of SEQ ID NOs: 2 or 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% identical to any of SEQ ID NOs: 2 or 4.

[0056] In another preferred embodiment, the asparaginase is an asparaginase variant comprising (i) one or more of the following substitutions, wherein each position corresponds to a position in SEQ ID NO: 2: T71C, T74A, T74C, K122A, K122R, V139G, T140D, K194L, D197E, I228M, S238C, N239C, K240R, P241E, K253R, I258V, I258Y, R259C, R259V, S297V, S299A, T314A, P333L, S334P, S334W, E337S, S338G, S338W, G356D, K363R or E373H, or (ii) a deletion at one or more positions corresponding to positions 27, 28 or 29 of SEQ ID NO: 2; wherein the asparaginase variant has an amino acid sequence which is at least 50% identical to SEQ ID NO: 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% identical to SEQ ID NO: 4.

[0057] In another preferred embodiment, the asparaginase is an asparaginase variant comprising one or more of the following substitutions, wherein each position corresponds to a position in SEQ ID NO: 2: T71C, T74A, T74C, K122A, K122R, V139G, T140D, K194L, D197E, I228M, S238C, N239C, K240R, P241E, K253R, I258V, I258Y, R259C, R259V, S297V, S299A, T314A, P333L, S334P, S334W, E337S, S338G, S338W, G356D, K363R or E373H; wherein the asparaginase variant has an amino acid sequence which is at least 50% identical to SEQ ID NO: 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% identical to SEQ ID NO: 4.

[0058] In another preferred embodiment, the asparaginase is an asparaginase variant comprising one or more of the following substitutions, wherein each position corresponds to a position in SEQ ID NO: 2: T74A, K122A, K122R, K194L, K240R, P241E, S299A, S334P, S334W, E337S or S338W; wherein the asparaginase variant has an amino acid sequence which is at least 50% identical to SEQ ID NO: 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% identical to SEQ ID NO: 4.

[0059] In another preferred embodiment, the asparaginase is an asparaginase variant comprising one or more of the following substitutions, wherein each position corresponds to a position in SEQ ID NO: 2: K122A, K240R, P241E, S299A, S334P, E337S or S338W; wherein the asparaginase variant has an amino acid sequence which is at least 50% identical to SEQ ID NO: 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% identical to SEQ ID NO: 4.

[0060] In another preferred embodiment, the asparaginase is an asparaginase variant disclosed in Patent Application No. EP12180861 filed 17 August 2012 which is hereby incorporated by reference.

EXAMPLES

Materials and Methods

Asparaginase activity (ASNU) assay

[0061] The activity of asparaginase may be measured in ASNU. An asparaginase unit (ASNU) is defined as the amount of enzyme needed to generate 1.0 micromole of ammonia in 1 minute at 37°C and pH 7.0, in 0.1 M MOPS buffer with 9.2 mg/ml asparagine.

[0062] Asparaginase hydrolyzes asparagine to aspartic acid and ammonium. The produced ammonium is combined with α -ketoglutarate to form glutamic acid whereby NADH is oxidized to NAD⁺. The reaction is catalysed by a surplus of glutamate dehydrogenase. The consumption of NADH is measured by photometry at 340 nm. NADH has an absorbance at 340 nm, while NAD⁺ has no absorbance. A decrease in color is thus measured, and can be correlated to asparaginase activity.

[0063] Activity is determined relative to an asparaginase standard of known activity. A commercial product having a

declared activity like Acrylaway L may be used as standard.

Asparaginase activity assay - used in Examples 1 and 2

Principle:

[0064] Asparaginase hydrolyses asparagine to aspartic acid and ammonium. Produced ammonium is determined using Nessler's reagent.

[0065] Activity is determined relative to a standard of known activity in ASNU or to a relevant control sample. Activity as determined according to this assay is referred to as nASNU.

Enzyme incubation:

[0066]

<i>Temperature</i>	37°C
<i>pH</i>	6.0
<i>Buffer</i>	20 mM citric acid, pH 6 + 0.001% triton x-100
<i>Asparagine substrate sol.</i>	25 mg/ml in buffer
<i>Stop reagent</i>	1.5 M Trichloroacetic acid (TCA)
<i>Enzyme Standard</i>	0-20 ASNU/ml The enzyme dilutions are made in buffer.
<i>Incubation time</i>	10 min

Procedure:

[0067]

Buffer	750 µL
Asparagine (25 mg/mL)	100 µL
Sample	100 µL
1.5 M TCA (stop reagent)	50 µL
<i>Total volume</i>	1.000 µL

[0068] Mix buffer and asparagine and let it equilibrate at 37°C for 10-15min. Add enzyme standard or sample and incubate for 10 min at 37°C. Add TCA to stop the reaction.

Ammonium determination (Nessler assay):

[0069]

<i>Temperature</i>	Ambient
<i>pH</i>	~ 12, controlled by addition of Nessler's reagent
<i>Nessler's reagent</i>	Mercury(II)chloride, potassium iodide, potassium hydroxide
<i>Incubation time</i>	10 min
<i>A₄₃₆</i>	Endpoint measurement at 440 nm

EP 2 884 857 B9

Procedure:

[0070]

5

	Microtiter plate
MQ Water	140 μ L
Sample	20 μ L
Nessler's reagent	40 μ L
Total volume	200 μ L

10

[0071] MQ water, sample and Nessler's reagent is added to the microtiter plate. Shake for 10 sec and then leave to incubate for 10 min before shaking again and reading at 440 nm.

15

Asparaginase activity assay (rASNU assay) - used in Examples 4, 5, 6 and 7

Reagents:

20

[0072]

- 1 M Potassium phosphate buffer (pH6.0)
1 M KH_2PO_4 (136g/500ml) + 1 M K_2HPO_4 (174g/500ml)
Adjust to pH 6.0
- 100 mM Potassium phosphate buffer (pH 6.0) + 0.1% tritonX-100 (1 L)
100 ml 1 M Potassium phosphate buffer (pH 6.0)
1 g Triton X-100
Adjust to 1000 ml
- 2 M Hydroxylamine (HA) solution (100 ml)
13.9 g hydroxylamine
Adjust to 100 ml with 100 mM potassium phosphate buffer (pH 6)
- Stop solution (500 ml)
23.83 ml acetate
13.88 g $\text{FeCl}_3 \cdot 6 \text{H}_2\text{O}$
84 ml 5 N HCl
Adjust to 500 ml with H_2O
- Substrate solution (100 ml)
10 ml 1 M Potassium phosphate buffer
0.5 g L-asparagine
5 ml 2 M HA soln.
Adjust to 100 ml with H_2O

25

30

35

40

Activity assay:

45

[0073]

- 1 Pipette 20 microL sample into a well.
- 2 Add 100 microL of substrate solution into the well.
- 3 Incubate 20min. at 55°C (or as otherwise indicated).
- 4 Add 100 microL of stop solution into the well.
- 5 Measure A490.

50

[0074] Compare result to that of a control for a relative effect or to a standard of known activity in ASNU. Activity as determined according to this assay is referred to as rASNU.

55

Asparaginase enzymes

[0075] JN065N2 is a variant of wild type *Aspergillus oryzae* asparaginase described in WO2008110513 having the following amino acid substitutions: N70K A323R T327V A349Q S351A V353I as compared to the amino acid sequence of the wild type.

[0076] JN070 is a variant of JN065N2 which further has the amino acid substitution S307A, i.e. it has the following amino acid substitutions as compared to the amino acid sequence of wild type *Aspergillus oryzae* asparaginase: N70K S307A A323R T327V A349Q S351A V353I.

[0077] The following variants (Table 0) of JN070 are used in the present Examples. For the avoidance of any possible doubt: E.g., the variant JN102 has the following amino acid substitutions as compared to the amino acid sequence of wild type *Aspergillus oryzae* asparaginase: N70K K122A S307A A323R T327V A349Q S351 A V353I.

Table 0: Variants of JN070

	Substitutions to JN070
JN102	K122A
JN117	K290V
JN135	S297V S299A
JN140	S334P
JN145	S338W
JN148	K290V S338W
JN152	S299A
JN158	K122A P241E S299A
JN159	K122A K290V S338W
JN161	K122A P241E K290V S299A S338W
JN162	K122A P241E K290V S338W
JN165	K122A S299A S334P
JN166	K122A P241E S334P
JN167	K122A P241E S299A S334P
JN168	K122A K290V S334P S338W
JN178	K122A K240R P241E S299A S334P
JN179	K122A K240R P241E K253R S299A S334P
JN180	K122A K240R P241E S299A S334P E337S
JN181	K122A K240R P241E K253R S299A S334P E337S
JN182	K122A P241E S299A S334P E337S
JN183	K122A P241E S299A S334P S338W
JN184	K122A P241E S299A S334W
JN226	A27* T28* D29* K122A P241E S299A S334P S338W

Example 1*Thermostability in potato blanching water with SAPP*

[0078] 0.5% SAPP was added to potato blanching water (=4 L deionized water which has been used for blanching of 2400 g potatoes at 85°C (16 portions of 150 g for 4 min each)) and pH adjusted to pH 5. This solution is intended to mimic a SAPP solution used for continuous chemical dip treatment of French fries.

[0079] The thermostability of the different asparaginase variants was tested in this solution at 70°C: Each asparaginase variant used at an enzyme dose of 10,000 ASNU/L was incubated in 20 ml of potato blanching water with SAPP at 70°C

EP 2 884 857 B9

for 4 hrs. Samples were taken every ½-1 hr for a total period of 4 hrs and enzyme activity analysed. Results are shown below.

[0080]

Table 1: Residual activity after 4 hrs incubation in "potato blanching water with SAPP" at 70°C compared to the initial activity at time 0.

	%		%		%
Sample	Residual act.	Sample	Residual act.	Sample	Residual act.
JN102	58	JN159	72	JN179	90
JN117	53	JN161	108	JN180	114
JN135	100	JN162	124	JN181	84
JN140	47	JN165	99	JN182	84
JN145	81	JN166	108	JN183	109
JN148	71	JN167	103	JN184	100
JN152	80	JN168	111	JN226	114
JN158	103	JN178	97		

[0081] As seen from the table all variants show increased stability at 70°C compared to the wt *Aspergillus oryzae* asparaginase which has <10% residual activity at similar conditions.

Example 2

Comparing thermostability of the thermostable variants and the wt enzyme in continuous treatment of French fries for acrylamide mitigation

[0082] French fry potatoes (Maris Piper or Bintje) were manually peeled and cut into French fries (size 0.8x0.8x5 cm) using a French fry cutter (Coupe Frites). The potato sticks from the individual potatoes were mixed and held in de-ionized water until use. Portions of 20 g of potato sticks were blanched in two steps; first at 85°C for 4 min (4L deionised water that was reused) and subsequently in 250 ml deionised water at 70°C for 15 minutes (fresh water for each sample). Enzyme treatment was done by holding the blanched potato sticks for 5 min at 70°C in 200 ml enzyme solution (0.5% SAPP in deionised water) dosing 10,000 ASNU/L of the variants or the wt enzyme. Every 5 min a new portion of blanched potato sticks was dipped in the enzyme bath in order to mimic continuous use of the enzyme bath. Samples from the enzyme bath were taken every 30 min for a total period of 4 hours and frozen for later activity analysis.

Results are shown below.

[0083]

Table 2: Residual activity of enzyme solution after 4 hrs continuous use at 70°C of dipping freshly blanched potato strips and holding each portion for 5 min. Residual activity after 4 hrs is calculated as a percentage of the initial measured activity at t=0.

	%		%		%
Sample	Residual act.	Sample	Residual act.	Sample	Residual act.
JN102	81	JN158	112	JN179	111
JN117	107	JN159	86	JN181	100
JN135	83	JN161	80	JN182	100
JN140	73	JN165	122	JN183	108
JN145	87	JN166	108	JN184	128
JN148	85	JN167	73	JN226	121

(continued)

	%		%		%
Sample	Residual act.	Sample	Residual act.	Sample	Residual act.
JN152	77	JN168	80	wt	8

[0084] As seen from the table the variants show a significant increase in stability at 70°C compared to the wt enzyme also under conditions of continuous use in an application set-up.

Example 3

Application performance of the thermostable variants in treatment of French fries for acrylamide mitigation

[0085] French fry potatoes (Maris Piper or Bintje) were manually peeled and cut into French fries (size 0.8x0.8x5 cm) using a French fry cutter (Coupe Frites). The potato sticks from the individual potatoes were mixed and held in de-ionized water until use. Portions of 75 g potato sticks were blanched in two steps; first at 85°C for 4 min (4L deionised water that was reused) and subsequently in 250 ml de-ionized water at 70°C for 15 minutes (fresh water for each sample). Enzyme treatment was done by dipping the blanched potato sticks for 1 min at 70°C in 250 ml enzyme solution (0.5% Sodium Acid Pyrophosphate, pH 5 in deionised water) using a dosage of 10,000 ASNU/L of the variants. For comparison a control sample dipped in 0.5% SAPP without enzyme was included. Testing of the wt enzyme was done at 55°C to ensure activity. Samples were made in triplicate. After enzyme treatment the potato sticks were dried in a ventilated heating cupboard for 10 min at 85°C, 20% RH (relative humidity), and parfried in vegetable oil for 1 min at 175°C. The samples were blast frozen and finally second fried 3 min at 175°C.

[0086] The fries were blended and the acrylamide extracted using acetonitrile and an Automated Solvent Extractor (ASE from Dionex). The extract was treated with Carrez solution I and II (Carrez solution I contains 15 g/100 ml of potassium hexacyanoferrate(II) trihydrate ($K_4[Fe(CN)_6] \times 3H_2O$); Carrez solution II contains 30 g/100 ml of zinc sulfate heptahydrate ($ZnSO_4 \times 7H_2O$)), left overnight in the fridge and filtered using a 0.22 µm filter before HPLC analysis (column: Dionex IonPac ICE-AS1, 9x250 mm, eluent: 5 mM HCl, detection: UV 202 nm). Acrylamide was identified and quantified by comparing with known standards.

Results are given below.

[0087]

Table 3: Calculated reduction in acrylamide formation in final French fries treated with the different variants at a fixed dose of 10,000 ASNU/L and a dip temperature of 70°C. Reduction is calculated by comparing to a control sample dipped in SAPP without enzyme. For the wt enzyme treatment was done at a dip temperature of 55 and 70°C.

Duplicate numbers are from separate repeated trials.

Treatment	Reduction vs. Control sample, %
Control	0
wt (55°C)	45
wt (70°C)	1
JN135	53
JN140	55
JN145	35
JN152	54/50
JN158	52
JN159	49
JN161	41
JN165	37
JN166	36

EP 2 884 857 B9

(continued)

Treatment	Reduction vs. Control sample, %
JN167	46
JN168	42
JN178	41
JN179	51
JN180	52
JN181	51/51
JN182	51/49
JN183	48
JN184	47
JN226	61

[0088] Acrylamide in the final French fry product has been reduced by approximately 50% showing that the variants are active in the application. For the wildtype enzyme the achieved reductions in acrylamide levels were 45% at 55°C and 1% at 70°C.

Example 4

Temperature activity

[0089] Purified enzyme was diluted with 20 mM sodium acetate buffer pH 5.5 to a concentration around 2 rASNU/ml. Ten microliters of enzyme solution was added to 100 microL of substrate solution, which has been pre-incubated at different temperature, of rASNU assay in a 96-well PCR plate and incubated for 20 min. The reaction was stopped by adding 100 microL of stop solution and the absorbance at 490 nm was measured.

Table 4: Temperature activity in percentage of activity at 50°C

Variant No.	Thermoactivity (100% @ 50°C)		
	35°C	60°C	70°C
WT	78%	64%	4%
JN065	67%	68%	24%
JN070	64%	68%	25%
JN117	63%	69%	30%
JN135	79%	72%	29%
JN140	69%	67%	34%
JN145	72%	68%	25%
JN148	67%	75%	29%
JN152	57%	79%	35%
JN158	51%	79%	38%
JN159	67%	68%	27%
JN161	51%	82%	39%
JN162	63%	73%	31%
JN165	55%	77%	38%
JN166	63%	69%	30%

(continued)

Variant No.	Thermoactivity (100% @ 50°C)		
	35°C	60°C	70°C
JN167	53%	83%	42%
JN168	71%	71%	31%
JN178	50%	83%	42%
JN179	45%	83%	41%
JN180	39%	92%	51%
JN181	38%	96%	52%
JN182	41%	92%	50%
JN183	57%	77%	37%
JN184	58%	92%	46%
JN226	59%	84%	41%

Example 5*pH activity*

[0090] Purified enzymes were diluted with 20 mM sodium acetate buffer pH 5.5 to a concentration around 2 rASNU/ml. Ten microliters of enzyme solution was added to 100 microL of preincubated substrate solution with different pH constructed by mixing 2-fold substrate solution of rASNU assay and equal volume of 200 mM Britton-Robinson buffer (pH 2.5 - 11.5). After 20 min incubation at 50°C, the reaction was stopped by adding 100 microL of stop solution and the absorbance at 490 nm was measured.

Table 5: pH activity

Variant No.	pH activity
	pH5/pH6.5 @ 50°C
WT	36%
JN065	41%
JN070	44%
JN102	45%
JN117	47%
JN135	29%
JN140	50%
JN145	43%
JN148	44%
JN152	52%
JN158	53%
JN159	45%
JN161	55%
JN162	45%
JN165	61%
JN166	51%

(continued)

Variant No.	pH activity
	pH5/pH6.5 @ 50°C
JN167	59%
JN168	53%
JN178	59%
JN179	57%
JN180	64%
JN181	62%
JN182	58%
JN183	60%
JN184	55%
JN226	62%

Example 6*pH stability at 70°C*

[0091] Purified enzyme was diluted with 0.5% SAPP pH 4.0 - 7.0 (pH was adjusted using HCl or NaOH) to a concentration around 6 rASNU/ml. Fifty microliters of diluted enzyme solution was transferred to 96-well PCR plate and incubated at 70°C for 2 hours. Immediately after incubation, 10 microL of incubated solution was added to 100 microL of preincubated substrate solution of rASNU assay in a 96-well plate and incubated at 50°C for 20 min. The reaction was stopped by adding 100 microL of stop solution and the absorbance at 490 nm was measured. The residual activity was determined using the activity of a similar sample incubated at 25°C as a control.

Table 6: pH stability

Variant No.	pH stability @70°C, 2h			
	pH4	pH5	pH6	pH7
WT	0%	0%	0%	0%
JN065	10%	12%	0%	0%
JN070	68%	55%	0%	0%
JN102	62%	64%	0%	0%
JN117	77%	78%	0%	0%
JN135	93%	94%	10%	0%
JN140	74%	77%	28%	0%
JN145	93%	99%	46%	0%
JN148	77%	80%	44%	0%
JN152	91%	92%	27%	0%
JN158	93%	98%	73%	0%
JN159	87%	91%	69%	0%
JN161	98%	100%	94%	74%
JN162	98%	102%	93%	1%
JN165	98%	100%	84%	2%

(continued)

Variant No.	pH stability @70°C, 2h			
	pH4	pH5	pH6	pH7
JN166	95%	98%	81%	0%
JN167	105%	103%	92%	40%
JN168	108%	105%	95%	12%
JN178	105%	106%	99%	37%
JN179	109%	104%	94%	4%
JN180	109%	116%	102%	74%
JN181	110%	112%	100%	38%
JN182	109%	112%	101%	74%
JN183	106%	104%	98%	75%
JN184	94%	84%	54%	0%
JN226	99%	99%	92%	80%

Example 7*Enzyme thermostability in 0.5% SAPP*

[0092] Purified enzyme was diluted with 0.5% sodium dihydrogen pyrophosphate (SAPP) pH 5.0 to a concentration around 6 rASNU/ml. Fifty microliters of diluted enzyme solution was transferred to a 96-well PCR plate and incubated at 70 - 80°C for 4 hours by using a thermal cycler. Immediately after incubation, 10 microL of incubated solution was added to 100 microL of pre-incubated substrate solution of rASNU assay in a 96-well plate and incubated at 37 or 50°C for 20 min. The reaction was stopped by adding 100 microL of stop solution and the absorbance at 490 nm was measured. The residual activity was determined using the activity of a similar sample incubated at 25°C as a control.

TABLE 7 Residual activity after 4 hours

Variant No.	residual after 4h in 0.5% SAPP, pH5					
	70 °C	72 °C	74 °C	76 °C	78 °C	80 °C
WT	1%	1%				
JN065	1%	2%				
JN070	35%	1%	2%	1%	1%	
JN102	53%	6%	2%	2%	1%	
JN117	61%	6%	1%	1%	1%	
JN135	88%	72%	21%	1%	1%	
JN140	67%	52%	21%	2%	1%	
JN145	90%	71%	17%	2%	1%	
JN148	74%	62%	26%	2%	1%	
JN152	79%	67%	25%	0%	0%	
JN158	93%	79%	67%	43%	2%	
JN159	86%	75%	47%	3%	0%	
JN161	100%	99%	95%	77%	71%	
JN162	102%	97%	91%	50%	1%	

(continued)

Variant No.	residual after 4h in 0.5% SAPP, pH5					
	70 °C	72 °C	74 °C	76 °C	78 °C	80 °C
JN165	98%	92%	80%	64%	37%	
JN166	100%	95%	85%	61%	9%	
JN167	103%	101%	97%	82%	68%	
JN168	104%	97%	87%	72%	25%	
JN178	103%	103%	99%	92%	78%	33%
JN179	93%	87%	81%	74%	53%	10%
JN180	100%	106%	104%	96%	83%	64%
JN181	100%	100%	96%	87%	72%	35%
JN182	101%	102%	99%	94%	76%	59%
JN183	99%	100%	98%	94%	76%	62%
JN184	94%	87%	64%	48%	22%	0%
JN226	100%	103%	102%	96%	90%	76%

SEQUENCE LISTING

[0093]

<110> Novozymes A/S

<120> Method for producing a food product

<130> 12475-WO-PCT

<160> 6

<170> PatentIn version 3.5

<210> 1

<211> 1137

<212> DNA

<213> *Aspergillus oryzae*

<400> 1

EP 2 884 857 B9

	atgggtgtca atttcaaagt tcttgccctg tcggccttag ctactattag ccatgcttcg	60
	cctctcctat atcctcgagc cacagactcg aacgtcacct atgtgttcac caacccaat	120
5	ggcctgaact ttactcagat gaacaccacc ctgccaaacg tcactatctt cgcgacaggc	180
	ggcacaatcg cgggctccag cggcgacaac accgcaacaa cagggttaca agccggtgca	240
	gtcggcatcc agacactgat cgacgcggtc ccggaaatgc taaacggtgc caacgtcgct	300
10	ggcgtgcaag taaccaatgt cggcagccca gacatcacct ccgacattct cctgcgtctc	360
	tccaaacaga tcaacgaggt ggtctgcaac gacccccacca tggccggtgc agtggtcacc	420
	cacggcaccg acacgctcga agaatccgcc ttcttcctcg acgccacggt caactgtcgc	480
15	aagcccgtgg tcatcgtcgg cgccatgcgc ctttcaaccg ccatctcggc tgacggcccc	540
	ctcaacctcc tgcaatccgt caccgtcgcc gcgagcccca aggcccgaga ccgcggcgcc	600
20	ctgattgtca tgaacgaccg catcgtatcc gccttctacg cctccaagac gaacgccaac	660
	accgtcgata cattcaaggc catcgaaatg ggtaacctgg gcgaggctgt ctccaacaaa	720
	ccctacttct tctaccccc agtcaagcca acaggcaaga cggaagtaga tatccggaac	780
25	atcacctcca tcccagagt cgacatcctc tactcatacg aagacatgca caatgacacc	840
	ctttactccg ccatcgacaa cggcgcaaag ggcacgttta tcgccggctc cggctccggc	900
	tccgtctcca ccccttcag cgccgccatg gaagacatca caaccaaaca caacatcccc	960
30	atcgtagcca gcacgcgcac cggaaacggg gaggtgccgt cctccgccga gtcgagccag	1020
	atcgcaagcg ggtatttgaa ccccgcaaag tcacgcgttt tgcttggtt gttgcttgcc	1080
35	caggggaaga gtattgagga aatgagggcg gtttttgagc ggattggggt tgcttga	1137

<210> 2

<211> 378

<212> PRT

40 <213> *Aspergillus oryzae*

<400> 2

45

50

55

EP 2 884 857 B9

	Met	Gly	Val	Asn	Phe	Lys	Val	Leu	Ala	Leu	Ser	Ala	Leu	Ala	Thr	Ile	
	1				5					10					15		
5	Ser	His	Ala	Ser	Pro	Leu	Leu	Tyr	Pro	Arg	Ala	Thr	Asp	Ser	Asn	Val	
				20					25					30			
	Thr	Tyr	Val	Phe	Thr	Asn	Pro	Asn	Gly	Leu	Asn	Phe	Thr	Gln	Met	Asn	
10			35					40					45				
	Thr	Thr	Leu	Pro	Asn	Val	Thr	Ile	Phe	Ala	Thr	Gly	Gly	Thr	Ile	Ala	
		50					55					60					
15	Gly	Ser	Ser	Ala	Asp	Asn	Thr	Ala	Thr	Thr	Gly	Tyr	Lys	Ala	Gly	Ala	
	65					70					75					80	
	Val	Gly	Ile	Gln	Thr	Leu	Ile	Asp	Ala	Val	Pro	Glu	Met	Leu	Asn	Val	
20					85					90					95		
	Ala	Asn	Val	Ala	Gly	Val	Gln	Val	Thr	Asn	Val	Gly	Ser	Pro	Asp	Ile	
25				100					105					110			
	Thr	Ser	Asp	Ile	Leu	Leu	Arg	Leu	Ser	Lys	Gln	Ile	Asn	Glu	Val	Val	
			115					120					125				
30	Cys	Asn	Asp	Pro	Thr	Met	Ala	Gly	Ala	Val	Val	Thr	His	Gly	Thr	Asp	
		130					135					140					
	Thr	Leu	Glu	Glu	Ser	Ala	Phe	Phe	Leu	Asp	Ala	Thr	Val	Asn	Cys	Arg	
35		145				150					155					160	
	Lys	Pro	Val	Val	Ile	Val	Gly	Ala	Met	Arg	Pro	Ser	Thr	Ala	Ile	Ser	
					165					170					175		
40	Ala	Asp	Gly	Pro	Leu	Asn	Leu	Leu	Gln	Ser	Val	Thr	Val	Ala	Ala	Ser	
				180					185					190			
45	Pro	Lys	Ala	Arg	Asp	Arg	Gly	Ala	Leu	Ile	Val	Met	Asn	Asp	Arg	Ile	
			195				200						205				
	Val	Ser	Ala	Phe	Tyr	Ala	Ser	Lys	Thr	Asn	Ala	Asn	Thr	Val	Asp	Thr	
50		210					215					220					
	Phe	Lys	Ala	Ile	Glu	Met	Gly	Asn	Leu	Gly	Glu	Val	Val	Ser	Asn	Lys	
	225					230					235					240	
55	Pro	Tyr	Phe	Phe	Tyr	Pro	Pro	Val	Lys	Pro	Thr	Gly	Lys	Thr	Glu	Val	
					245					250					255		

EP 2 884 857 B9

	Asp	Ile	Arg	Asn	Ile	Thr	Ser	Ile	Pro	Arg	Val	Asp	Ile	Leu	Tyr	Ser	
				260					265					270			
5	Tyr	Glu	Asp	Met	His	Asn	Asp	Thr	Leu	Tyr	Ser	Ala	Ile	Asp	Asn	Gly	
			275					280					285				
10	Ala	Lys	Gly	Ile	Val	Ile	Ala	Gly	Ser	Gly	Ser	Gly	Ser	Val	Ser	Thr	
		290					295					300					
15	Pro	Phe	Ser	Ala	Ala	Met	Glu	Asp	Ile	Thr	Thr	Lys	His	Asn	Ile	Pro	
	305					310					315					320	
20	Ile	Val	Ala	Ser	Thr	Arg	Thr	Gly	Asn	Gly	Glu	Val	Pro	Ser	Ser	Ala	
					325					330					335		
25	Glu	Ser	Ser	Gln	Ile	Ala	Ser	Gly	Tyr	Leu	Asn	Pro	Ala	Lys	Ser	Arg	
				340					345					350			
30	Val	Leu	Leu	Gly	Leu	Leu	Leu	Ala	Gln	Gly	Lys	Ser	Ile	Glu	Glu	Met	
			355					360					365				
35	Arg	Ala	Val	Phe	Glu	Arg	Ile	Gly	Val	Ala							
							370	375									
40	<210> 3																
	<211> 1137																
	<212> DNA																
	<213> Artificial Sequence																
45	<220>																
	<223> Variant																
50	<400> 3																
55	atgggtgtca	atttcaaagt	tcttgccctg	tcggccttag	ctactattag	ccatgcttcg											60
	cctctcctat	atcctcgagc	cacagactcg	aacgtcacct	atgtgttcac	caaccccaat											120
	ggcctgaact	ttactcagat	gaacaccacc	ctgccaaacg	tcactatctt	cgcgacaggc											180
	ggcacaatcg	cgggctccag	cgccgacaag	accgcaacaa	caggttacaa	agccgggtgca											240
	gtcggcatcc	agacactgat	cgacgcggtc	ccggaaatgc	taaacgttgc	caacgtcgct											300
	ggcgtgcaag	taaccaatgt	cggcagccca	gacatcacct	ccgacattct	cctgcgtctc											360
	tccaaacaga	tcaacgaggt	ggtctgcaac	gacccaccca	tggccggtgc	agtggtcacc											420
	cacggcaccg	acacgctcga	agaatccgcc	ttcttcctcg	acgccacggt	caactgtcgc											480
	aagcccgtgg	tcacgctcgg	cgccatgcgc	ccttcaaccg	ccatctcggc	tgacggcccc											540
	ctcaacctcc	tgcaatccgt	caccgtcgcc	gcgagcccca	aggcccgaga	ccgcggcgcc											600

EP 2 884 857 B9

	ctgattgtca tgaacgaccg catcgtatcc gccttctacg cctccaagac gaacgccaac	660
	accgtcgata cattcaaggc catcgaaatg ggtaacctgg gcgaggtcgt ctccaacaaa	720
5	ccctacttct tctaccccc agtcaagcca acaggcaaga cggaagtaga tatccggaac	780
	atcacctcca tccccagagt cgacatcctc tactcatacg aagacatgca caatgacacc	840
	ctttactccg ccatcgacaa cggcgcaaag ggcacgttta tcgccggctc cggctccggc	900
10	tccgtctcca ccccttcgc ggccgccatg gaagacatca caaccaaaca caacatcccc	960
	atcgtagca gcacccgcgt cggaaacggg gaggtgccgt cctccgccga gtcgagccag	1020
	atcgcaagcg ggtatttgaa cccccagaag gcccgcatct tgcttggtt gttgcttgcc	1080
15	caggggaaga gtattgagga aatgagggcg gtttttgagc ggattggggt tgcttga	1137

<210> 4

<211> 378

20 <212> PRT

<213> Artificial Sequence

<220>

<223> Variant

25

<400> 4

30

35

40

45

50

55

EP 2 884 857 B9

	Met	Gly	Val	Asn	Phe	Lys	Val	Leu	Ala	Leu	Ser	Ala	Leu	Ala	Thr	Ile	
	1				5					10					15		
5	Ser	His	Ala	Ser	Pro	Leu	Leu	Tyr	Pro	Arg	Ala	Thr	Asp	Ser	Asn	Val	
				20					25					30			
10	Thr	Tyr	Val	Phe	Thr	Asn	Pro	Asn	Gly	Leu	Asn	Phe	Thr	Gln	Met	Asn	
			35					40					45				
15	Thr	Thr	Leu	Pro	Asn	Val	Thr	Ile	Phe	Ala	Thr	Gly	Gly	Thr	Ile	Ala	
		50					55					60					
20	Gly	Ser	Ser	Ala	Asp	Lys	Thr	Ala	Thr	Thr	Gly	Tyr	Lys	Ala	Gly	Ala	
	65					70					75					80	
25	Val	Gly	Ile	Gln	Thr	Leu	Ile	Asp	Ala	Val	Pro	Glu	Met	Leu	Asn	Val	
					85					90					95		
30	Ala	Asn	Val	Ala	Gly	Val	Gln	Val	Thr	Asn	Val	Gly	Ser	Pro	Asp	Ile	
				100					105					110			
35	Thr	Ser	Asp	Ile	Leu	Leu	Arg	Leu	Ser	Lys	Gln	Ile	Asn	Glu	Val	Val	
			115					120					125				
40	Cys	Asn	Asp	Pro	Thr	Met	Ala	Gly	Ala	Val	Val	Thr	His	Gly	Thr	Asp	
		130					135					140					

EP 2 884 857 B9

	Thr	Leu	Glu	Glu	Ser	Ala	Phe	Phe	Leu	Asp	Ala	Thr	Val	Asn	Cys	Arg	
	145					150					155					160	
5	Lys	Pro	Val	Val	Ile	Val	Gly	Ala	Met	Arg	Pro	Ser	Thr	Ala	Ile	Ser	
					165					170					175		
10	Ala	Asp	Gly	Pro	Leu	Asn	Leu	Leu	Gln	Ser	Val	Thr	Val	Ala	Ala	Ser	
				180					185					190			
15	Pro	Lys	Ala	Arg	Asp	Arg	Gly	Ala	Leu	Ile	Val	Met	Asn	Asp	Arg	Ile	
			195					200					205				
20	Val	Ser	Ala	Phe	Tyr	Ala	Ser	Lys	Thr	Asn	Ala	Asn	Thr	Val	Asp	Thr	
		210					215					220					
25	Phe	Lys	Ala	Ile	Glu	Met	Gly	Asn	Leu	Gly	Glu	Val	Val	Ser	Asn	Lys	
	225					230					235					240	
30	Pro	Tyr	Phe	Phe	Tyr	Pro	Pro	Val	Lys	Pro	Thr	Gly	Lys	Thr	Glu	Val	
					245					250					255		
35	Asp	Ile	Arg	Asn	Ile	Thr	Ser	Ile	Pro	Arg	Val	Asp	Ile	Leu	Tyr	Ser	
				260					265					270			
40	Tyr	Glu	Asp	Met	His	Asn	Asp	Thr	Leu	Tyr	Ser	Ala	Ile	Asp	Asn	Gly	
			275					280					285				
45	Ala	Lys	Gly	Ile	Val	Ile	Ala	Gly	Ser	Gly	Ser	Gly	Ser	Val	Ser	Thr	
		290					295					300					
50	Pro	Phe	Ala	Ala	Ala	Met	Glu	Asp	Ile	Thr	Thr	Lys	His	Asn	Ile	Pro	
	305					310					315					320	
55	Ile	Val	Arg	Ser	Thr	Arg	Val	Gly	Asn	Gly	Glu	Val	Pro	Ser	Ser	Ala	
					325				330						335		
60	Glu	Ser	Ser	Gln	Ile	Ala	Ser	Gly	Tyr	Leu	Asn	Pro	Gln	Lys	Ala	Arg	
				340					345					350			
65	Ile	Leu	Leu	Gly	Leu	Leu	Leu	Ala	Gln	Gly	Lys	Ser	Ile	Glu	Glu	Met	
		355						360					365				
70	Arg	Ala	Val	Phe	Glu	Arg	Ile	Gly	Val	Ala							
		370					375										
75	<210> 5																
	<211> 1137																
	<212> DNA																
	<213> Artificial Sequence																

EP 2 884 857 B9

<220>

<223> Variant

<400> 5

5
atgggtgtca atttcaaagt tcttgccctg tcggccttag ctactattag ccatgcttcg 60
cctctcctat atcctcgagc cacagactcg aacgtcacct atgtgttcac caacccaat 120
10 ggcctgaact ttactcagat gaacaccacc ctgccaaacg tcactatctt cgcgacaggc 180
ggcacaatcg cgggctccag cgcgcacaag accgcaacaa caggttacaa agccggtgca 240
gtcggcatcc agacactgat cgacgcggtc ccggaaatgc taaacgttgc caacgtcgct 300
15 ggcgtgcaag taaccaatgt cggcagccca gacatcacct ccgacattct cctgcgtctc 360
tccaaacaga tcaacgaggt ggtctgcaac gacccaccca tggccggtgc agtggtcacc 420
cacggcaccg acacgctcga agaatccgcc ttcttcctcg acgccacggt caactgtcgc 480
20 aagcccggtg tcatcgctcg cgccatgcgc cttcaaccg ccattctcggc tgacggcccc 540
ctcaacctcc tgcaatccgt caccgtcgcc gcgagcccca aggcccgaga ccgcggcgcc 600
ctgattgtca tgaacgaccg catcgatatc gccttctacg cctccaagac gaacgccaac 660
25 accgtcgata cattcaaggc catcgaaatg ggtaacctg gcgaggtcgt ctccaacaaa 720
ccctacttct tctaccccc agtcaagcca acaggcaaga cggaagtaga tatccggaac 780
30 atcacctcca tcccagagt cgacatcctc tactcatacg aagacatgca caatgacacc 840
ctttactccg ccatcgacaa cggcgcaaag ggcatcgta tcgccggctc cggctccggc 900
tccgtctcca ccccttcag cgcgcgatg gaagacatca caaccaaaca caacatcccc 960
35 atcgtagca gaccccgct cggaaacggg gaggtgccgt cctccgccga gtcgagccag 1020
atcgcaagcg ggtatttgaa ccccgagaag gcccgcatct tgcttggtt gttgcttgcc 1080
caggggaaga gtattgagga aatgagggcg gtttttgagc ggattggggt tgcttga 1137
40

<210> 6

<211> 378

<212> PRT

<213> Artificial Sequence

45

<220>

<223> Variant

<400> 6

50

Met Gly Val Asn Phe Lys Val Leu Ala Leu Ser Ala Leu Ala Thr Ile
1 5 10 15

55

Ser His Ala Ser Pro Leu Leu Tyr Pro Arg Ala Thr Asp Ser Asn Val
20 25 30

EP 2 884 857 B9

	Thr	Tyr	Val	Phe	Thr	Asn	Pro	Asn	Gly	Leu	Asn	Phe	Thr	Gln	Met	Asn	
			35					40					45				
5	Thr	Thr	Leu	Pro	Asn	Val	Thr	Ile	Phe	Ala	Thr	Gly	Gly	Thr	Ile	Ala	
		50					55					60					
10	Gly	Ser	Ser	Ala	Asp	Lys	Thr	Ala	Thr	Thr	Gly	Tyr	Lys	Ala	Gly	Ala	
	65					70					75					80	
15	Val	Gly	Ile	Gln	Thr	Leu	Ile	Asp	Ala	Val	Pro	Glu	Met	Leu	Asn	Val	
					85					90					95		
20	Ala	Asn	Val	Ala	Gly	Val	Gln	Val	Thr	Asn	Val	Gly	Ser	Pro	Asp	Ile	
				100					105					110			
25	Thr	Ser	Asp	Ile	Leu	Leu	Arg	Leu	Ser	Lys	Gln	Ile	Asn	Glu	Val	Val	
			115					120					125				
30	Cys	Asn	Asp	Pro	Thr	Met	Ala	Gly	Ala	Val	Val	Thr	His	Gly	Thr	Asp	
	130						135					140					
35	Thr	Leu	Glu	Glu	Ser	Ala	Phe	Phe	Leu	Asp	Ala	Thr	Val	Asn	Cys	Arg	
	145					150					155					160	
40	Lys	Pro	Val	Val	Ile	Val	Gly	Ala	Met	Arg	Pro	Ser	Thr	Ala	Ile	Ser	
					165					170					175		
45	Ala	Asp	Gly	Pro	Leu	Asn	Leu	Leu	Gln	Ser	Val	Thr	Val	Ala	Ala	Ser	
				180					185					190			
50	Pro	Lys	Ala	Arg	Asp	Arg	Gly	Ala	Leu	Ile	Val	Met	Asn	Asp	Arg	Ile	
			195				200						205				
55	Val	Ser	Ala	Phe	Tyr	Ala	Ser	Lys	Thr	Asn	Ala	Asn	Thr	Val	Asp	Thr	
	210						215					220					
60	Phe	Lys	Ala	Ile	Glu	Met	Gly	Asn	Leu	Gly	Glu	Val	Val	Ser	Asn	Lys	
	225					230					235					240	
65	Pro	Tyr	Phe	Phe	Tyr	Pro	Pro	Val	Lys	Pro	Thr	Gly	Lys	Thr	Glu	Val	
					245					250					255		
70	Asp	Ile	Arg	Asn	Ile	Thr	Ser	Ile	Pro	Arg	Val	Asp	Ile	Leu	Tyr	Ser	
				260					265					270			
75	Tyr	Glu	Asp	Met	His	Asn	Asp	Thr	Leu	Tyr	Ser	Ala	Ile	Asp	Asn	Gly	

275

280

285

Ala Lys Gly Ile Val Ile Ala Gly Ser Gly Ser Gly Ser Val Ser Thr
 290 295 300

Pro Phe Ser Ala Ala Met Glu Asp Ile Thr Thr Lys His Asn Ile Pro
 305 310 315 320

Ile Val Arg Ser Thr Arg Val Gly Asn Gly Glu Val Pro Ser Ser Ala
 325 330 335

Glu Ser Ser Gln Ile Ala Ser Gly Tyr Leu Asn Pro Gln Lys Ala Arg
 340 345 350

Ile Leu Leu Gly Leu Leu Leu Ala Gln Gly Lys Ser Ile Glu Glu Met
 355 360 365

Arg Ala Val Phe Glu Arg Ile Gly Val Ala
 370 375

Claims

1. A method for producing a heat-treated vegetable-based food product comprising:

- (a) contacting of a vegetable-based food material with asparaginase at a temperature of 67-75°C;
- (b) drying the vegetable-based food material at an air temperature of 40 to 90°C; and
- (c) heat-treating the asparaginase treated vegetable-based food material to obtain the heat-treated vegetable-based food product;

wherein the asparaginase has a residual activity after 4 hours' incubation in deionised water with 0.5% SAPP at 70°C, pH 5, of at least 20%, preferably at least 40%, more preferably at least 60%, even more preferably at least 80%, of the activity without such incubation; and wherein the asparaginase has an activity at 35°C, pH 6, of at least 20%, preferably at least 30%, more preferably at least 35%, of its activity at 50°C, pH 6.

2. The method of claim 1 wherein the vegetable-based food material is potato sticks and the heat-treated vegetable-based food product is French fries.

3. The method of any one of the preceding claims wherein step (a) is dipping vegetable-based food material into an asparaginase solution having a temperature of 67-75°C, preferably 68-72°C, more preferably 70°C.

4. The method of claim 3 wherein the asparaginase solution comprises asparaginase at a concentration of 1,000 to 100,000 ASNU/L, preferably 5,000 to 50,000 ASNU/L, more preferably 10,000 to 30,000 ASNU/L.

5. The method of claim 4 wherein the asparaginase solution further comprises sodium acid pyrophosphate at a concentration of 0.05 to 5%, preferably 0.1 to 1.5%, more preferably 0.5 to 1%.

6. The method of any one of claims 3-5 wherein the dipping is for a time of 1 second to 10 minutes, preferably 30 seconds to 3 minutes.

7. The method of any one of the preceding claims wherein the vegetable-based food material has been blanched prior to step (a).

8. The method of any one of the preceding claims wherein the drying in step (b) is performed at an air temperature of

55 to 85°C, preferably 65 to 75°C.

9. The method of any one of the preceding claims wherein the surface temperature of the vegetable-based food material during drying in step (b) is 25 to 60°C, preferably 35 to 50°C.
10. The method of any one of the preceding claims wherein the drying in step (b) is performed for between 5 and 30 minutes.
11. The method of any one of the preceding claims wherein the acrylamide content of the heat-treated vegetable-based food product is at least 25% reduced, preferably at least 30%, at least 35%, at least 40%, at least 45% or at least 50% reduced, compared to the acrylamide content of a heat-treated vegetable-based food product produced by a similar method without the addition of asparaginase.
12. Use of asparaginase for treatment of a vegetable-based food material at a temperature of 67-75°C; wherein the asparaginase has a residual activity after 4 hours' incubation in deionised water with 0.5% SAPP at 70°C, pH 5, of at least 20%, preferably at least 40%, more preferably at least 60%, even more preferably at least 80%, of the activity without such incubation; and wherein the asparaginase has an activity at 35°C, pH 6, of at least 20%, preferably at least 30%, more preferably at least 35%, of its activity at 50°C, pH 6.
13. The method of any one of claims 1-11 or the use of claim 12 wherein the asparaginase has an activity at 70°C, pH 6, of at least 20%, preferably at least 25%, more preferably at least 30%, of its activity at 50°C, pH 6.
14. The method of any one of claims 1-11 or 13 or the use of any one of claims 12-13 wherein the asparaginase is an asparaginase variant comprising a substitution at one or more positions corresponding to positions 122, 140, 197, 238, 239, 240, 241, 253, 258, 259, 297 or 373 of SEQ ID NO: 2, and having an amino acid sequence which is at least 50% identical to any of SEQ ID NOs: 2 or 4.
15. The method of any one of claims 1-11 or 13 or the use of any one of claims 12-13 wherein the asparaginase is an asparaginase variant comprising (i) one or more of the following substitutions, wherein each position corresponds to a position in SEQ ID NO: 2: T71C, T74A, T74C, K122A, K122R, V139G, T140D, K194L, D197E, I228M, S238C, N239C, K240R, P241E, K253R, I258V, I258Y, R259C, R259V, S297V, S299A, T314A, P333L, S334P, S334W, E337S, S338G, S338W, G356D, K363R or E373H, or (ii) a deletion at one or more positions corresponding to positions 27, 28 or 29 of SEQ ID NO: 2; wherein the asparaginase variant has an amino acid sequence which is at least 50% identical to any of SEQ ID NOs: 2 or 4, preferably at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% identical to any of SEQ ID NOs: 2 or 4.

Patentansprüche

1. Verfahren zur Herstellung eines hitzebehandelten Nahrungsmittelprodukts auf Pflanzenbasis, umfassend:
 - a) Inkontaktbringen eines Nahrungsmittelmaterials auf Pflanzenbasis mit Asparaginase bei einer Temperatur von 67-75°C;
 - b) Trocknen des Nahrungsmittelmaterials auf Pflanzenbasis bei einer Lufttemperatur von 40 bis 90°C; und
 - c) Hitzebehandeln des mit Asparaginase behandelten Nahrungsmittelmaterials auf Pflanzenbasis, wodurch man zu dem hitzebehandelten Nahrungsmittelprodukt auf Pflanzenbasis gelangt;

wobei die Asparaginase nach vierstündiger Inkubation in entionisiertem Wasser mit 0,5% SAPP bei 70°C, pH 5, eine Restaktivität von mindestens 20%, vorzugsweise mindestens 40%, stärker bevorzugt mindestens 60%, noch stärker bevorzugt mindestens 80%, der Aktivität ohne solch eine Inkubation aufweist; und

wobei die Asparaginase eine Aktivität bei 35°C, pH 6, von mindestens 20%, vorzugsweise mindestens 30%, stärker bevorzugt mindestens 35%, ihrer Aktivität bei 50°C, pH 6, aufweist.
2. Verfahren nach Anspruch 1, wobei es sich bei dem Nahrungsmittelmaterial auf Pflanzenbasis um Kartoffelstäbchen und bei dem hitzebehandelten Nahrungsmittelprodukt auf Pflanzenbasis um Pommes Frites handelt.
3. Verfahren nach einem der vorhergehenden Ansprüche, wobei Schritt (a) das Tauchen eines Nahrungsmittelmaterials auf Pflanzenbasis in eine Asparaginaselösung mit einer Temperatur von 67-75°C, vorzugsweise 68-72°C, stärker

bevorzugt 70°C, ist.

4. Verfahren nach Anspruch 3, wobei die Asparaginaselösung Asparaginase in einer Konzentration von 1000 bis 100000 ASNU/L, vorzugsweise 5000 bis 50000 ASNU/L, stärker bevorzugt 10000 bis 30000 ASNU/L, umfasst.
5. Verfahren nach Anspruch 4, wobei die Asparaginaselösung weiterhin Natriumdihydrogenpyrophosphat in einer Konzentration von 0,05 bis 5%, vorzugsweise 0,1 bis 1,5%, stärker bevorzugt 0,5 bis 1%, umfasst.
6. Verfahren nach einem der Ansprüche 3-5, wobei das Tauchen über einen Zeitraum von 1 Sekunde bis 10 Minuten, vorzugsweise 30 Sekunden bis 3 Minuten, erfolgt.
7. Verfahren nach einem der vorhergehenden Ansprüche, wobei das Nahrungsmittelmateriale auf Pflanzenbasis vor Schritt (a) blanchiert worden ist.
8. Verfahren nach einem der vorhergehenden Ansprüche, wobei das Trocknen in Schritt (b) bei einer Lufttemperatur von 55 bis 85°C, vorzugsweise 65 bis 75°C, erfolgt.
9. Verfahren nach einem der vorhergehenden Ansprüche, wobei die Oberflächentemperatur des Nahrungsmittelmateriale auf Pflanzenbasis während des Trocknens in Schritt (b) 25 bis 60°C, vorzugsweise 35 bis 50°C, beträgt.
10. Verfahren nach einem der vorhergehenden Ansprüche, wobei das Trocknen in Schritt (b) über einen Zeitraum von zwischen 5 und 30 Minuten erfolgt.
11. Verfahren nach einem der vorhergehenden Ansprüche, wobei der Acrylamidgehalt des hitzebehandelten Nahrungsmittelprodukts auf Pflanzenbasis im Vergleich zu dem Acrylamidgehalt eines hitzebehandelten Nahrungsmittelprodukts auf Pflanzenbasis, das mit einem ähnlichen Verfahren ohne Zugabe von Asparaginase erzeugt worden ist, um mindestens 25%, vorzugsweise um mindestens 30%, mindestens 35%, mindestens 40%, mindestens 45% oder mindestens 50% vermindert ist.
12. Verwendung von Asparaginase für die Behandlung eines Nahrungsmittelmateriale auf Pflanzenbasis bei einer Temperatur von 67-75°C; wobei die Asparaginase nach vierstündiger Inkubation in entionisiertem Wasser mit 0,5% SAPP bei 70°C, pH 5, eine Restaktivität von mindestens 20%, vorzugsweise mindestens 40%, stärker bevorzugt mindestens 60%, noch stärker bevorzugt mindestens 80%, der Aktivität ohne solch eine Inkubation aufweist; und wobei die Asparaginase eine Aktivität bei 35°C, pH 6, von mindestens 20%, vorzugsweise mindestens 30%, stärker bevorzugt mindestens 35%, ihrer Aktivität bei 50°C, pH 6, aufweist.
13. Verfahren nach einem der Ansprüche 1-11 oder Verwendung nach Anspruch 12, wobei die Asparaginase eine Aktivität bei 70°C, pH 6, von mindestens 20%, vorzugsweise mindestens 25%, stärker bevorzugt mindestens 30%, ihrer Aktivität bei 50°C, pH 6, aufweist.
14. Verfahren nach einem der Ansprüche 1-11 oder 13 oder Verwendung nach einem der Ansprüche 12-13, wobei es sich bei der Asparaginase um eine Asparaginasevariante handelt, die eine Substitution an einer oder mehreren Positionen entsprechend den Positionen 122, 140, 197, 238, 239, 240, 241, 253, 258, 259, 297 oder 373 von SEQ ID NO: 2 umfasst und eine Aminosäuresequenz aufweist, die zu mindestens 50% zu einer der SEQ ID NO: 2 oder 4 identisch ist.
15. Verfahren nach einem der Ansprüche 1-11 oder 13 oder Verwendung nach einem der Ansprüche 12-13, wobei es sich bei der Asparaginase um eine Asparaginasevariante handelt, die (i) eine oder mehrere der folgenden Substitutionen umfasst, wobei jede Position einer Position in SEQ ID NO: 2 entspricht: T71C, T74A, T74C, K122A, K122R, V139G, T140D, K194L, D197E, I228M, S238C, N239C, K240R, P241E, K253R, I258V, I258Y, R259C, R259V, S297V, S299A, T314A, P333L, S334P, S334W, E337S, S338G, S338W, G356D, K363R oder E373H, oder (ii) eine Deletion an einer oder mehreren Positionen entsprechend den Positionen 27, 28 oder 29 von SEQ ID NO: 2 umfasst; wobei die Asparaginasevariante eine Aminosäuresequenz aufweist, die zu mindestens 50% zu einer der SEQ ID NO: 2 oder 4 identisch ist, vorzugsweise zu mindestens 60%, mindestens 70%, mindestens 75%, mindestens 80%, mindestens 85%, mindestens 90%, mindestens 95% oder mindestens 98% zu einer der SEQ ID NO: 2 oder 4 identisch ist.

Revendications

1. Procédé de production d'un produit alimentaire à base de légumes thermo-traité, comprenant les étapes consistant à :

(a) mettre en contact un matériel alimentaire à base de légumes avec une asparaginase à une température de 67 à 75°C ;
 (b) sécher le matériel alimentaire à base de légumes à une température de l'air de 40 à 90°C ; et
 (c) traiter par la chaleur le matériel alimentaire à base de légumes traité avec une asparaginase afin d'obtenir le produit alimentaire à base de légumes thermo-traité ;

dans lequel l'asparaginase a une activité résiduelle, après 4 heures d'incubation dans de l'eau désionisée, avec du SAPP à 0,5% à 70°C, pH 5, d'au moins 20%, de préférence d'au moins 40%, mieux préféré d'au moins 60%, même mieux préféré d'au moins 80% de l'activité sans une telle incubation ; et

dans lequel l'asparaginase a une activité à 35°C, pH 6, d'au moins 20%, de préférence d'au moins 30%, mieux préféré d'au moins 35% de son activité à 50°C, pH 6.

2. Procédé selon la revendication 1, dans laquelle le matériel alimentaire à base de légumes ce sont des bâtonnets de pomme de terre et le produit alimentaire à base de légumes thermo-traité ce sont des frites.

3. Procédé selon l'une quelconque des revendications précédentes, dans laquelle l'étape (a) est une immersion du matériel alimentaire à base de légumes dans une solution d'asparaginase ayant une température de 67 à 75°C, de préférence de 68 à 72°C, mieux préféré de 70°C.

4. Procédé selon la revendication 3, dans laquelle la solution d'asparaginase comprend une asparaginase à une concentration de 1 000 à 100 000 ASNU/L, de préférence de 5 000 à 50 000 ASNU/L, mieux préféré de 10 000 à 30 000 ASNU/L.

5. Procédé selon la revendication 4, dans laquelle la solution d'asparaginase comprend en outre du pyrophosphate acide de sodium à une concentration de 0,05 à 5%, de préférence de 0,1 à 1,5%, mieux préféré de 0,5 à 1%.

6. Procédé selon l'une quelconque des revendications 3 à 5, dans laquelle l'immersion a lieu pendant une durée de 1 seconde à 10 minutes, de préférence de 30 secondes à 3 minutes.

7. Procédé selon l'une quelconque des revendications précédentes, dans laquelle le matériel alimentaire à base de légumes a été blanchi avant l'étape (a).

8. Procédé selon l'une quelconque des revendications précédentes, dans laquelle le séchage dans l'étape (b) est effectué à une température de l'air de 55 à 85°C, de préférence de 65 à 75°C.

9. Procédé selon l'une quelconque des revendications précédentes, dans laquelle la température de surface du matériel alimentaire à base de légumes lors du séchage dans l'étape (b) est de 25 à 60°C, de préférence de 35 à 50°C.

10. Procédé selon l'une quelconque des revendications précédentes, dans laquelle le séchage dans l'étape (b) est effectué pendant un temps compris entre 5 et 30 minutes.

11. Procédé selon l'une quelconque des revendications précédentes, dans laquelle la teneur d'acrylamide du produit alimentaire à base de légumes thermo-traité est réduite d'au moins 25%, de préférence réduite d'au moins 30%, d'au moins 35%, d'au moins 40%, d'au moins 45% ou d'au moins 50%, en comparaison avec la teneur d'acrylamide d'un produit alimentaire à base de légumes thermo-traité produit par un procédé semblable sans l'ajout d'asparaginase.

12. Utilisation d'une asparaginase pour le traitement d'un matériel alimentaire à base de légumes à une température de 67 à 75°C ; dans laquelle l'asparaginase a une activité résiduelle après 4 heures d'incubation dans de l'eau désionisée, avec du SAPP à 0,5% à 70°C, pH 5, d'au moins 20%, de préférence d'au moins 40%, mieux préféré d'au moins 60%, même mieux préféré d'au moins 80% de l'activité sans une telle incubation ; et dans laquelle l'asparaginase a une activité à 35°C, pH 6, d'au moins 20%, de préférence d'au moins 30%, mieux préféré d'au moins 35% de son activité à 50°C, pH 6.

13. Procédé, selon l'une quelconque des revendications 1 à 11, ou utilisation, selon la revendication 12, dans laquelle l'asparaginase a une activité à 70°C, pH 6, d'au moins 20%, de préférence d'au moins 25%, mieux préféré d'au moins 30% de son activité à 50°C, pH 6.

14. Procédé, selon l'une quelconque des revendications 1 à 11 ou 13, ou utilisation, selon l'une quelconque des revendications 12-13, dans laquelle l'asparaginase est un variant d'asparaginase comprenant une substitution en une ou plusieurs position(s) correspondant aux positions 122, 140, 197, 238, 239, 240, 241, 253, 258, 259, 297 ou 373 de la SEQ ID n° : 2, et ayant une séquence d'acides aminés qui est identique, au moins à 50%, à n'importe laquelle des SEQ ID n° : 2 ou 4.

15. Procédé, selon l'une quelconque des revendications 1 à 11 ou 13, ou utilisation, selon l'une quelconque des revendications 12-13, dans laquelle l'asparaginase est un variant d'asparaginase comprenant (i) une ou plusieurs substitution(s) parmi les suivantes, dans lesquelles chaque position correspond à une position dans la SEQ ID n° : 2 : T71C, T74A, T74C, K122A, K122R, V139G, T140D, K194L, D197E, I228M, S238C, N239C, K240R, P241E, K253R, I258V, I258Y, R259C, R259V, S297V, S299A, T314A, P333L, S334P, S334W, E337S, S338G, S338W, G356D, K363R ou E373H, ou (ii) une délétion au niveau d'une ou plusieurs position(s) correspondant aux positions 27, 28 ou 29 de la SEQ ID n° : 2 ; dans laquelle le variant d'asparaginase a une séquence d'acides aminés qui est identique, au moins à 50%, à n'importe laquelle des SEQ ID n° : 2 ou 4, de préférence au moins à 60%, au moins à 70%, au moins à 75%, au moins à 80%, au moins à 85%, au moins à 90%, au moins à 95% ou au moins à 98%, à n'importe laquelle des SEQ ID n° : 2 ou 4.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2004026042 A [0003] [0005]
- WO 2008110513 A [0006] [0075]
- WO 2008151807 A [0007]
- WO 2010070010 A [0008]
- US 20070141227 A1 [0009]
- WO 2007077546 A [0010]
- WO 2004030468 A [0051]
- EP 12180861 A [0060]