



(11)

EP 2 904 099 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:

19.12.2018 Bulletin 2018/51

(51) Int Cl.:

C12N 9/42 (2006.01)

C11D 3/48 (2006.01)

(86) International application number:

PCT/EP2013/070618

(21) Application number: **13771506.6**

(22) Date of filing: **03.10.2013**

(87) International publication number:

WO 2014/053594 (10.04.2014 Gazette 2014/15)

(54) PREVENTING ADHESION OF BACTERIA

VERHINDERN VON BAKTERIENADHÄSION

PRÉVENTION DE L'ADHÉSION DE BACTÉRIES

(84) Designated Contracting States:

AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(74) Representative: **NZ EPO Representatives**

**Krogshøjvej 36
2880 Bagsvaerd (DK)**

(30) Priority: **05.10.2012 EP 12187547**

(56) References cited:

**EP-A1- 2 157 162 WO-A1-2009/148800
WO-A1-2010/069742 WO-A2-2004/053039
WO-A2-2008/007318 WO-A2-2008/007320
US-A- 5 356 803**

(43) Date of publication of application:
12.08.2015 Bulletin 2015/33

(60) Divisional application:

16201125.8 / 3 176 260

- **DATABASE WPI Week 201256 Thomson Scientific, London, GB; AN 2012-J97232 XP002715491, "New Brevundimonas diminuta strain useful in deodorizing agent to remove odor, preferably sulfur bad smell or ammonia bad smell from livestock manure", & KR 2012 0083605 A (LEE E Y) 26 July 2012 (2012-07-26)**

(73) Proprietor: **Novozymes A/S
2880 Bagsvaerd (DK)**

- **LEROY C ET AL: "Effects of commercial enzymes on the adhesion of a marine biofilm-forming bacterium", BIOFOULING (CHUR), HARWOOD ACADEMIC PUBLISHERS, CHUR, CH, vol. 24, no. 1, 1 January 2008 (2008-01-01), pages 11-22, XP008143734, ISSN: 0892-7014, DOI: 10.1080/08927010701784912**

(72) Inventors:

- **GORI, Klaus
DK-2100 Copenhagen OE (DK)**
- **BALTSEN, Lilian Eva Tang
DK-2880 Bagsvaerd (DK)**
- **ALLESEN-HOLM, Marie
DK-3400 Hillerød (DK)**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 904 099 B1

Description**Reference to a Sequence Listing**

- 5 [0001] This application contains a Sequence Listing in computer readable form. The computer readable form is incorporated herein by reference.

FIELD OF THE INVENTION

- 10 [0002] The invention relates to methods for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, using an endo-beta-1,4-glucanase.

BACKGROUND

- 15 [0003] When laundry items like T-shirts or sportswear are used, they are exposed to bacteria from the body of the user and from the rest of the environment in which they are used. These bacteria are a source of bad odor, which develops after use, but which may remain even after wash. The reason for this bad odor is adhesion of bacteria to the textile surface. Because of the adhesion to the textile, the bacteria may remain even after wash, and thereby continue to be a source of bad odor. The references WO2009/148800 relates to a laundering system comprising various bactericides. US5356803 relates to antimicrobial compositions comprising an endoglycosidase and an antimicrobial agent. The endoglucanase is useful in diminishing the adhesion of bacteria. The references WO2008/007318 and WO2008/007320 relates to compositions comprising endo-beta-1,4-glucanase activity and huing agents where the compositions may assist in removal of yellow soil. The references EP2157162, WO2004/053039 and KR20120083605 relates to a bleach composition comprising an enzyme having endo-beta-1,4-glucanase activity.
- 25 [0004] The present invention provides a solution to this problem by reducing the adhesion of bacteria to the textile surface during wash. The selected bacteria are sources of very bad odor, and they were isolated from real-life laundry items.

SUMMARY

- [0005] The invention is defined by the subject-matter of the claims.
- [0006] Described is a method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase.
- 35 [0007] Described is a method for reducing adhesion of bacteria selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp. to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase.
- 40 [0008] Described is a method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase, wherein the aqueous composition comprises at least 1 mg/l of an endo-beta-1,4-glucanase.
- [0009] The present invention provides a method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase, wherein the amino acid sequence of the endo-beta-1,4-glucanase has at least 80% identity, preferably at least 90% identity, more preferably at least 95% identity, and most preferably 100% identity to the amino acid sequence of SEQ ID NO: 1.
- 45 [0010] Described is a method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase, wherein the aqueous composition also comprises a surfactant.
- [0011] Described is a method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase, wherein the surface is a textile surface and the aqueous composition is a laundry detergent composition.
- 55 [0012] Described is a method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase, wherein the bacteria are *Brevundimonas* sp. Described is a method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous com-

position comprising an endo-beta-1,4-glucanase,

wherein the adhesion is reduced by at least 50%, or at least 50% of the bacteria are released from the surface.

[0013] Described is a laundry composition comprising water; textile items; bacteria selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp.; and an endo-beta-1,4-glucanase.

[0014] Described is a laundry composition comprising water; textile items; bacteria selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp.; and an endo-beta-1,4-glucanase, wherein the composition comprises at least 1 mg/l of an endo-beta-1,4-glucanase.

[0015] The present invention describes the use of an endo-beta-1,4-glucanase for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, wherein the amino acid sequence of the endo-beta-1,4-glucanase has at least 80% identity, preferably at least 90% identity, more preferably at least 95% identity, and most preferably 100% identity to the amino acid sequence of SEQ ID NO:1.

[0016] In another aspect, the present invention describes the above use of an endo-beta-1,4-glucanase for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, wherein the bacteria are selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp.

DETAILED DESCRIPTION

Endo-beta-1,4-glucanase

[0017] The endo-beta-1,4-glucanase as specified in the claims is to be used.

endo-beta-1,4-glucanase known in the art is described as follows: Suitable endo-beta-1,4-glucanase includes the polypeptide shown as amino acids 1 to 773 of SEQ ID NO: 1, which is a mature endo-glucanase sequence with a calculated molecular weight of 86 kDa.

[0018] It is believed that positions 1 to about 340 of SEQ ID NO: 1 are the catalytically active domain of the endo-glucanase enzyme. It is also believed that positions from about 340 to about 540 are the cellulose binding domain of the endo-glucanase enzyme.

[0019] The endo-beta-1,4-glucanase enzyme may comprise the amino acid sequence shown as (i) amino acids 1 to 773 of SEQ ID NO: 1,.

[0020] The present invention also provides the use of endo-glucanase polypeptides that are substantially homologous to the polypeptide of (i) above. The term "substantially homologous" is used herein to denote polypeptides being at least 80%, preferably at least 85%, more preferably at least 90%, more preferably at least 95%, even more preferably at least 96% identical, even more preferably at least 97% identical, even more preferably at least 98% identical, and most preferably at least 99% or more identical to the amino acid sequence of SEQ ID NO: 1,.

[0021] For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends Genet. 16: 276-277), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

[0022] ,An endo-beta-1,4-glucanase of SEQ ID NO: 1 may comprise a substitution, deletion, and/or insertion at one or more (e.g., several) positions., the number of amino acid substitutions, deletions and/or insertions introduced into the mature polypeptide of SEQ ID NO: 1 is not more than 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8 or 9. The amino acid changes may be of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of 1-30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

[0023] Examples of conservative substitutions are within the groups of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, In, The Proteins, Academic

Press, New York. Common substitutions are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

[0024] Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

[0025] Essential amino acids in a polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, Science 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for endo-beta-1,4-glucanase activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., 1996, J. Biol. Chem. 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., 1992, Science 255: 306-312; Smith et al., 1992, J. Mol. Biol. 224: 899-904; Wlodaver et al., 1992, FEBS Lett. 309: 59-64. The identity of essential amino acids can also be inferred from an alignment with a related polypeptide.

[0026] Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, Science 241: 53-57; Bowie and Sauer, 1989, Proc. Natl. Acad. Sci. USA 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman et al., 1991, Biochemistry 30: 10832-10837; U.S. Patent No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire et al., 1986, Gene 46: 145; Ner et al., 1988, DNA 7: 127).

[0027] Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness et al., 1999, Nature Biotechnology 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide.

[0028] The polypeptide may be a hybrid polypeptide in which a region of one polypeptide is fused at the N-terminus or the C-terminus of a region of another polypeptide.

[0029] The polypeptide may be a fusion polypeptide or cleavable fusion polypeptide in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide of the present invention. A fusion polypeptide is produced by fusing a polynucleotide encoding another polypeptide to a polynucleotide of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fusion polypeptide is under control of the same promoter(s) and terminator. Fusion polypeptides may also be constructed using intein technology in which fusion polypeptides are created post-translationally (Cooper et al., 1993, EMBO J. 12: 2575-2583; Dawson et al., 1994, Science 266: 776-779).

[0030] A fusion polypeptide can further comprise a cleavage site between the two polypeptides. Upon secretion of the fusion protein, the site is cleaved releasing the two polypeptides. Examples of cleavage sites include, but are not limited to, the sites disclosed in Martin et al., 2003, J. Ind. Microbiol. Biotechnol. 3: 568-576; Svetina et al., 2000, J. Biotechnol. 76: 245-251; Rasmussen-Wilson et al., 1997, Appl. Environ. Microbiol. 63: 3488-3493; Ward et al., 1995, Biotechnology 13: 498-503; and Contreras et al., 1991, Biotechnology 9: 378-381; Eaton et al., 1986, Biochemistry 25: 505-512; Collins-Racie et al., 1995, Biotechnology 13: 982-987; Carter et al., 1989, Proteins: Structure, Function, and Genetics 6: 240-248; and Stevens, 2003, Drug Discovery World 4: 35-48.

[0031] The endo-glucanase enzyme used in the invention may, in addition to the enzyme core comprising the catalytically active domain, i.e. amino acids 1 to about 340 of SEQ ID NO: 1, also comprise a cellulose binding domain (CBD), the cellulose binding domain and the catalytically active domain being operably linked. The cellulose binding domain (CBD) may exist as an integral part of the encoded enzyme as described above and in the appended SEQ ID NO: 1, or be a CBD from another origin, introduced into the endo-glucanase thus creating an enzyme hybrid. In this context, the term "cellulose-binding domain" is intended to be understood as defined by Peter Tomme et al. "Cellulose-Binding Domains: Classification and Properties" in "Enzymatic Degradation of Insoluble Carbohydrates", John N. Saddler and Michael H. Penner (Eds.), ACS Symposium Series, No. 618, 1996. This definition classifies more than 120 cellulose-binding domains into 10 families (I-X), and demonstrates that CBDs are found in various enzymes such as cellulases (endo-glucanases), xylanases, mannanases, arabinofuranosidases, acetyl esterases and chitinases. CBDs have also been found in algae, e.g. the red alga *Porphyra purpurea* as a non-hydrolytic polysaccharide-binding protein, see Tomme et al., *op.cit.* However, most of the CBDs are from cellulases and xylanases, CBDs are found at the N and C termini of proteins or are internal. Enzyme hybrids are known in the art, see e.g. WO 90/00609 and WO 95/16782, and may be prepared by transforming into a host cell a DNA construct comprising at least a fragment of DNA encoding the cellulose-binding domain ligated, with or without a linker, to a DNA sequence encoding the endo-glucanase and growing the host cell to express the fused gene. Enzyme hybrids may be described by the following formula:

CBD-MR-X

wherein CBD is the N-terminal or the C-terminal region of an amino acid sequence corresponding to at least the cellulose-binding domain; MR is the middle region (the linker), and may be a bond, or a short linking group preferably of from about 2 to about 100 carbon atoms, more preferably of from 2 to 40 carbon atoms; or is preferably from about 2 to about 100 amino acids, more preferably of from 2 to 40 amino acids; and X is an N-terminal or C-terminal region of a polypeptide corresponding at least to the catalytically active domain encoded by the DNA sequence of the invention.

[0032] In a similar way, the cellulose binding domain corresponding to amino acids 340 to 540 of SEQ ID NO: 1 can be used to form hybrids with endo-glucanases from sources. Examples of endo-glucanases from other sources replacing the endo-glucanase of amino acids 1 to 340 of SEQ ID NO: 1 include endo-glucanases from: (a) *Bacillus lautus*, for instance *Bacillus lautus* NCIMB 40250 disclosed in WO 91/10732, (b) *Humicola insolens* DSM1800 disclosed in WO 91/17243, (c) *Fusarium oxysporium* DSM2672 disclosed in WO 91/17243, and (d) *Bacillus sp. AC13* NCIMB 40482 disclosed in EP 0651785.

[0033] The concentration of the endo-beta-1,4-glucanase is typically in the range of 0.01-100 ppm enzyme protein, preferably 0.05-50 ppm enzyme protein, more preferably 0.1-50 ppm enzyme protein, more preferably 0.1-30 ppm enzyme protein, more preferably 0.5-20 ppm enzyme protein, and most preferably 0.5-10 ppm enzyme protein.

[0034] In an embodiment, the concentration of the endo-beta-1,4-glucanase is typically in the range of 1-40 ppm enzyme protein, preferably 1-20 ppm enzyme protein, more preferably 1-10 ppm enzyme protein.

Determination of endo-beta-1,4-glucanase activity

ECU method

[0035] In the ECU method, the ability of the enzyme sample to reduce the viscosity of a solution of carboxymethyl-cellulose (CMC) is determined, and the result is given in ECU. The reduction in viscosity is proportional to the endo-cellulase activity. Conditions: CMC type 7LFD from Hercules, pH 7.5 in 0.1M phosphate buffer, CMC concentration 31.1 g per liter reaction at 40°C for 30 minutes. A vibration viscosimeter such as MIVI 3000, Sofraser, France is used to measure the viscosity.

Cellazyme C method

[0036] Cellazyme C is an endo-glucanase assay substrate, supplied in tablet form by Megazyme International Ireland Ltd. Reference is made to Megazyme's pamphlet CZC 7/99 which states: "The substrate is prepared by dyeing and cross-linking HE-cellulose to produce a material which hydrates in water but is water insoluble. Hydrolysis by endo-beta-1,4-glucanase produces water-soluble dyed fragments, and the rate of release of these (increase in absorbance at 590 nm) can be related directly to enzyme activity."

[0037] The enzyme sample is added to 6 ml of a suitable buffer in a test tube, one Cellazyme C tablet is added and dispersed by shaking the tube, then the tube is placed in a water bath at 40°C. The contents are mixed by brief shaking after approximately 15, 30, 45 and 60 minutes. After 60 minutes the solution is filtered through Whatman GF/C filters, 9 cm diameter. The absorbance of the filtered solution is measured at 590nm.

Detergent composition

[0038] In one aspect, the endo-beta-1,4-glucanase is added to and thus becomes a component of a detergent composition.

[0039] The detergent composition of the present invention may be formulated, for example, as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations, or be formulated for hand or machine dishwashing operations.

[0040] The detergent composition may comprise one or more surfactants, which may be anionic and/or cationic and/or non-ionic and/or semi-polar and/or zwitterionic, or a mixture thereof. In a particular embodiment, the detergent composition includes a mixture of one or more nonionic surfactants and one or more anionic surfactants. The surfactant(s) is typically present at a level of from about 0.1% to 60% by weight, such as about 1% to about 40%, or about 3% to about 20%, or about 3% to about 10%. The surfactant(s) is chosen based on the desired cleaning application, and includes any conventional surfactant(s) known in the art.

[0041] When included therein, the detergent will usually contain from about 1% to about 40% by weight, such as from about 5% to about 30%, including from about 5% to about 15%, or from about 20% to about 25% of an anionic surfactant.

Non-limiting examples of anionic surfactants include sulfates and sulfonates, in particular, linear alkylbenzenesulfonates (LAS), isomers of LAS, branched alkylbenzenesulfonates (BABS), phenylalkanesulfonates, alpha-olefinsulfonates (AOS), olefin sulfonates, alkene sulfonates, alkane-2,3-diylbis(sulfates), hydroxyalkanesulfonates and disulfonates, alkyl sulfates (AS) such as sodium dodecyl sulfate (SDS), fatty alcohol sulfates (FAS), primary alcohol sulfates (PAS), alcohol ethersulfates (AES or AEOS or FES, also known as alcohol ethoxysulfates or fatty alcohol ether sulfates), secondary alkanesulfonates (SAS), paraffin sulfonates (PS), ester sulfonates, sulfonated fatty acid glycerol esters, alpha-sulfo fatty acid methyl esters (alpha-SFMe or SES) including methyl ester sulfonate (MES), alkyl- or alkenylsuccinic acid, dodecenylyl/tetradecenyl succinic acid (DTSA), fatty acid derivatives of amino acids, diesters and monoesters of sulfo-succinic acid or soap, and combinations thereof.

[0042] When included therein, the detergent will usually contain from about 0.2% to about 40% by weight of a non-ionic surfactant, for example from about 0.5% to about 30%, in particular from about 1% to about 20%, from about 3% to about 10%, such as from about 3% to about 5%, or from about 8% to about 12%. Non-limiting examples of non-ionic surfactants include alcohol ethoxylates (AE or AEO), alcohol propoxylates, propoxylated fatty alcohols (PFA), alkoxyated fatty acid alkyl esters, such as ethoxylated and/or propoxylated fatty acid alkyl esters, alkylphenol ethoxylates (APE), nonylphenol ethoxylates (NPE), alkylpolyglycosides (APG), alkoxyated amines, fatty acid monoethanolamides (FAM), fatty acid diethanolamides (FADA), ethoxylated fatty acid monoethanolamides (EFAM), propoxylated fatty acid monoethanolamide (PFAM), polyhydroxy alkyl fatty acid amides, or N-acyl N-alkyl derivatives of glucosamine (glucamides, GA, or fatty acid glucamide, FAGA), as well as products available under the trade names SPAN and TWEEN, and combinations thereof.

[0043] The detergent composition may contain about 0-65% by weight of a detergent builder or co-builder, or a mixture thereof. In a dish wash detergent, the level of builder is typically 40-65%, particularly 50-65%. The builder and/or co-builder may particularly be a chelating agent that forms water-soluble complexes with Ca and Mg. Any builder and/or co-builder known in the art for use in laundry detergents may be utilized. Non-limiting examples of builders include zeolites, diphosphates (pyrophosphates), triphosphates such as sodium triphosphate (STP or STPP), carbonates such as sodium carbonate, soluble silicates such as sodium metasilicate, layered silicates (e.g., SKS-6 from Hoechst), ethanolamines such as 2-aminoethan-1-ol (MEA), iminodiethanol (DEA) and 2,2',2"-nitrilotriethanol (TEA), and carboxymethylinulin (CMI), and combinations thereof.

[0044] The detergent composition may contain 0-50% by weight of a bleaching system. Any bleaching system known in the art for use in laundry detergents may be utilized. Suitable bleaching system components include bleaching catalysts, photobleaches, bleach activators, sources of hydrogen peroxide such as sodium percarbonate and sodium perborates, preformed peracids and mixtures thereof. Suitable preformed peracids include, but are not limited to, peroxydicarboxylic acids and salts, percarbonic acids and salts, perimidic acids and salts, peroxymonosulfuric acids and salts, for example, Oxone (R), and mixtures thereof. Non-limiting examples of bleaching systems include peroxide-based bleaching systems, which may comprise, for example, an inorganic salt, including alkali metal salts such as sodium salts of perborate (usually mono- or tetra-hydrate), percarbonate, persulfate, perphosphate, persilicate salts, in combination with a peracid-forming bleach activator. By Bleach activator is meant herein a compound which reacts with peroxygen bleach like hydrogen peroxide to form a Peracid. The peracid thus formed constitutes the activated bleach. Suitable bleach activators to be used herein include those belonging to the class of esters amides, imides or anhydrides. Suitable examples are tetracetyl ethylene diamine (TAED), sodium 3,5,5 trimethyl hexanoyloxybenzene sulphonate, diperoxy dodecanoic acid, 4-(dodecanoyloxy)benzene-sulfonate (LOBS), 4-(decanoyloxy)benzenesulfonate, 4-(decanoyloxy)benzoate (DOBS), 4-(3,5,5-trimethylhexanoyloxy)benzenesulfonate (ISONOBS), tetraacetyl ethylenediamine (TAED) and 4-(nonanoyloxy)benzenesulfonate (NOBS), and/or those disclosed in WO98/17767. A particular family of bleach activators of interest was disclosed in EP624154 and particularly preferred in that family is acetyl triethyl citrate (ATC). ATC or a short chain triglyceride like Triacin has the advantage that it is environmental friendly as it eventually degrades into citric acid and alcohol. Furthermore, acetyl triethyl citrate and triacetin has a good hydrolytical stability in the product upon storage and it is an efficient bleach activator. Finally, ATC provides a good building capacity to the laundry additive. Alternatively, the bleaching system may comprise peroxyacids of, for example, the amide, imide, or sulfone type. The bleaching system may also comprise peracids such as 6-(phthaloylamino)-percapronic acid (PAP). The bleaching system may also include a bleach catalyst.

[0045] Other ingredients of the detergent composition, which are all well-known in art, include hydrotropes, fabric hueing agents, anti-foaming agents, soil release polymers, anti-redeposition agents etc.

[0046] The detergent additive as well as the detergent composition may comprise one or more additional enzymes such as a protease, lipase, cutinase, amylase, carbohydrase, cellulase, pectinase, mannanase, arabinase, galactanase, xylanase, oxidase, e.g., a laccase, and/or peroxidase.

[0047] The endo-beta-1,4-glucanase of the present invention may be added to a detergent composition in an amount corresponding to at least 1 mg of endo-beta-1,4-glucanase protein, such as at least 5 mg of protein, preferably at least 10 mg of protein, more preferably at least 15 mg of protein, even more preferably at least 20 mg of protein, most preferably at least 30 mg of protein, and even most preferably at least 40 mg of protein per liter of wash liquor. Thus, the detergent

composition may comprise at least 0.1% endo-beta-1,4-glucanase protein, preferably at least 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, 0.8%, 1.0%, 1.2%, 1.5%, or 2.0% of endo-beta-1,4-glucanase protein.

[0048] The polypeptide having endo-beta-1,4-glucanase activity may be formulated as a liquid (e.g. aqueous), a solid, a gel, a paste or a dry product formulation. The dry product formulation may subsequently be re-hydrated to form an active liquid or semi-liquid formulation usable in the methods of the invention.

[0049] The compositions of the invention may further comprise auxiliary agents such as wetting agents, thickening agents, buffer(s) for pH control, stabilisers, perfume, colourants, fillers and the like.

[0050] Useful wetting agents are surfactants, i.e. non-ionic, anionic, amphoteric or zwitterionic surfactants. Surfactants are further described above.

Methods and Uses

[0051] Described is a method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase.

[0052] Described is a method for reducing adhesion of Gram-negative and/or Gram-positive bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase.

[0053] Described is a method for reducing adhesion of bacteria selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp. to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase.

[0054] Preferably, the aqueous composition comprises at least 1 mg/l of an endo-beta-1,4-glucanase having at least 80% identity, preferably at least 90% identity, more preferably at least 95% identity, and most preferably 100% identity to the amino acid sequence of SEQ ID NO: 1.

[0055] In an embodiment, the aqueous composition also comprises a surfactant; and optionally also a detergent builder or co-builder. Preferably, the surface is a textile surface and the aqueous composition is a laundry detergent composition. The textile surface may be the surface of any textile item, such as an item made of cotton or a synthetic material, for example a piece of sportswear, a T-shirt, or another piece of clothing which is exposed to sweat when used. The textile surface may also be the surface of bedding, bed linen or towels.

[0056] In an embodiment, the bacteria are *Brevundimonas* sp.

[0057] In an embodiment, the bacterial adhesion is reduced by at least 50%, or at least 50% of the bacteria are released from the surface.

[0058] A detergent composition may comprise at least 3% surfactant, and at least 0.1% of an endo-beta-1,4-glucanase.

[0059] A (laundry) composition may comprise water; textile items; bacteria selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp.; and an endo-beta-1,4-glucanase. Preferably, the composition comprises at least 1 mg/l of an endo-beta-1,4-glucanase. The textile item may be an item made of cotton or a synthetic material, for example a piece of sportswear, a T-shirt, or another piece of clothing which is exposed to sweat when used. The textile item may also be bedding, bed linen or towels.

[0060] The invention also provides for use of the methods and compositions above for reducing adhesion of bacteria selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp. to a surface, or releasing the bacteria from a surface to which they adhere.

[0061] The methods according to the invention may be carried out at a temperature between 0 and 70 degrees Celsius, preferably between 5 and 60 degrees Celsius, more preferably between 5 and 50 degrees Celsius, even more preferably between 5 and 40 degrees Celsius, even more preferably between 5 and 35 degrees Celsius, most preferably between 5 and 30 degrees Celsius, and in particular between 10 and 30 degrees Celsius.

[0062] Methods may employ a treatment time of from 5 minutes to 120 minutes, preferably from 5 minutes to 90 minutes, more preferably from 5 minutes to 60 minutes, more preferably from 5 minutes to 45 minutes, more preferably from 5 minutes to 30 minutes, most preferably from 5 minutes to 20 minutes, and in particular from 5 minutes to 15 minutes.

[0063] Methods may be carried out at pH 3 to pH 11, preferably at pH 5 to pH 10, more preferably at pH 7 to pH 9. Most preferably, the methods are carried out at the pH or temperature optimum of the endo-beta-1,4-glucanase +/- one pH unit.

[0064] The present invention is further described by the following examples which should not be construed as limiting the scope of the invention.

EXAMPLES

[0065] Chemicals used as buffers and substrates were commercial products of at least reagent grade. The endo-beta-

1,4-glucanase (see also WO 02/099091) used in the following Example has an amino acid sequence shown as SEQ ID NO: 1.

EXAMPLE 1

Reducing adhesion of bacteria to a surface using endo-beta-1,4-glucanase

[0066] One of the aims of the present study was to investigate the bacterial diversity in laundry after washing at 15, 40 and 60°C, respectively.

[0067] The study was conducted on laundry collected from Danish households. For each wash, 20 g of laundry items (tea towel, towel, dish cloth, bib, T-shirt armpit, T-shirt collar, socks) in the range 4:3:2:2:1:1:1 was used. Washing was performed in a Laundr-O-Meter (LOM) at 15, 40 and 60°C. For washing at 15 and 40°C, Ariel Sensitive White & Color was used, whereas WFK IEC-A* model detergent was used for washing at 60°C. Ariel Sensitive White & Color was prepared by weighing out 5.1 g and adding tap water up to 1000 ml followed by stirring for 5 minutes. WFK IEC-A* model detergent was prepared by weighing out 5 g and adding tap water up to 1300 ml followed by stirring for 15 min. Washing was performed for 1 hour at 15, 40 and 60°C, respectively, followed by 2 times rinsing for 20 min at 15°C.

[0068] Laundry was sampled immediately after washing at 15, 40 and 60°C, respectively. Twenty grams of laundry was added 0.9% (w/v) NaCl (1.06404; Merck, Darmstadt, Germany) with 0.5% (w/w) Tween 80 to yield a 1:10 dilution in stomacher bag. The mixture was homogenized using a Stomacher for 2 minutes at medium speed. After homogenization, ten-fold dilutions were prepared in 0.9% (w/v) NaCl. Bacteria were enumerated on Tryptone Soya Agar (CM0129, Oxoid, Basingstoke, Hampshire, UK) incubated aerobically at 30°C for 5-7 days. To suppress growth of yeast and moulds, 0.2% sorbic acid (359769, Sigma) and 0.1% cycloheximide (18079; Sigma) were added. Twenty-four bacterial and fungal colonies were selected from countable plates and purified by restreaking twice on Tryptone Soya Agar (TSA). For long time storage, purified isolates were stored at -80°C in TSB containing 20% (w/v) glycerol (49779; Sigma).

[0069] Four strains of *Brevundimonas* sp., one strain of *Sphingomonas* sp., one strain of *Sphingobium* sp., and one strain of *Exiguobacterium* sp. (all isolated as described above) were used in the present study. For long term storage, bacterial strains were maintained at -80°C in Tryptone Soya Broth (TSB) (pH 7.3) (CM0129, Oxoid Ltd, Basingstoke, UK), to which 20% (v/v) glycerol (Merck, Darmstadt, Germany) was added. Bacterial cultures were pre-grown on Tryptone Soya Agar (TSA) (pH 7.3) for 3-5 days at 30°C. From a single colony, a loop-full was transferred to a test tube containing 10 ml TSB and incubated for 1 day at 30°C with shaking (240 rpm).

[0070] After incubation, the bacterial cells were diluted 1000 times in TSB added 0, 0.5, 1, 2, 4, 8, 16, 32, 64, 128 and 256 ppm of endo-beta-1,4-glucanase. One hundred µl was inoculated into a 96-well polystyrene plate (flat bottom) (161093; Nunc, Roskilde, Denmark) and incubated for 3 days at 30°C.

[0071] After incubation, growth was determined by measurement of the optical density, measured at 600 nm using a Spectramax Plus384 reader (Molecular Devices, Sunnyvale, CA, USA).

[0072] Adhesion/biofilm prevention was measured by removing non-adherent cells by washing two times with 0.9% (w/v) NaCl (Merck). To measure adherence, 200 µl of 0.1% (w/v) crystal violet (C0775; Sigma-Aldrich, St. Louis, MO, USA) was added and left for 15 minutes at room temperature. The wells were washed two times with 0.9% (w/v) NaCl, and bound crystal violet was eluted by the addition of 200 µl 96% (w/v) ethanol (201145; Kemetyl, Koge, Denmark) and determined by measurement at 595 nm.

Table 1. Inhibition dosages of the endo-beta-1,4-glucanase, which is the lowest dose at which prevention of adhesion/ biofilm formation was observed.

Bacterial strain	Inhibition dosage (mg/l)
<i>Brevundimonas</i> sp. 1	0.5
<i>Brevundimonas</i> sp. 2	0.5
<i>Brevundimonas</i> sp. 3	0.5
<i>Brevundimonas</i> sp. 4	1
<i>Exiguobacterium</i> sp. 1	1
<i>Sphingobium</i> sp. 1	32
<i>Sphingomonas</i> sp. 1	62

[0073] The study shows that endo-beta-1,4-glucanase decreases the adhesion properties of *Brevundimonas* sp., *Sphingobium* sp., *Sphingomonas* sp. and *Exiguobacterium* sp. found in washed laundry, where they may cause textile

damage or produce malodor when the textiles are used again after being washed.

[0074] Most importantly, inhibition of adhesion properties will prevent transfer of these bacteria between different textile items during the washing process and thus limit the occurrence of these bacteria. Furthermore, inhibition of adhesion properties will minimize the risk of growth of these bacteria inside the washing machine. Growth of bacteria inside the washing machine may cause malodor from the washing machine. Furthermore, detached bacteria may be transferred to textiles during the washing process and later cause malodor from textiles when they are used after the washing process.

SEQUENCE LISTING

[0075]

<110> Novozymes A/S

<120> Preventing adhesion of bacteria

<130> 12523-WO-PCT

<160> 1

<170> PatentIn version 3.5

<210> 1

<211> 773

<212> PRT

<213> Bacillus ap.

<400> 1

EP 2 904 099 B1

	Ala	Glu	Gly	Asn	Thr	Arg	Glu	Asp	Asn	Phe	Lys	His	Leu	Leu	Gly	Asn	
	1				5					10					15		
5	Asp	Asn	Val	Lys	Arg	Pro	Ser	Glu	Ala	Gly	Ala	Leu	Gln	Leu	Gln	Glu	
				20					25					30			
10	Val	Asp	Gly	Gln	Met	Thr	Leu	Val	Asp	Gln	His	Gly	Glu	Lys	Ile	Gln	
			35					40					45				
15	Leu	Arg	Gly	Met	Ser	Thr	His	Gly	Leu	Gln	Trp	Phe	Pro	Asp	Ile	Leu	
		50					55					60					
20	Asn	Asp	Asn	Ala	Tyr	Lys	Ala	Leu	Ala	Asn	Asp	Trp	Glu	Ser	Asn	Met	
	65					70					75					80	
25	Ile	Arg	Leu	Ala	Met	Tyr	Val	Gly	Glu	Asn	Gly	Tyr	Ala	Ser	Asn	Pro	
					85					90					95		
30	Glu	Leu	Ile	Lys	Ser	Arg	Val	Ile	Lys	Gly	Ile	Asp	Leu	Ala	Ile	Glu	
				100					105					110			
35	Asn	Asp	Met	Tyr	Val	Ile	Val	Asp	Trp	His	Val	His	Ala	Pro	Gly	Asp	
			115					120					125				
40	Pro	Arg	Asp	Pro	Val	Tyr	Ala	Gly	Ala	Glu	Asp	Phe	Phe	Arg	Asp	Ile	
		130					135					140					
45	Ala	Ala	Leu	Tyr	Pro	Asn	Asn	Pro	His	Ile	Ile	Tyr	Glu	Leu	Ala	Asn	
	145					150					155					160	
50	Glu	Pro	Ser	Ser	Asn	Asn	Asn	Gly	Gly	Ala	Gly	Ile	Pro	Asn	Asn	Glu	
					165					170					175		
55																	

EP 2 904 099 B1

	Glu Gly Trp Asn Ala Val Lys Glu Tyr Ala Asp Pro Ile Val Glu Met	180	185	190
5	Leu Arg Asp Ser Gly Asn Ala Asp Asp Asn Ile Ile Ile Val Gly Ser	195	200	205
10	Pro Asn Trp Ser Gln Arg Pro Asp Leu Ala Ala Asp Asn Pro Ile Asn	210	215	220
	Asp His His Thr Met Tyr Thr Val His Phe Tyr Thr Gly Ser His Ala	225	230	235
15	Ala Ser Thr Glu Ser Tyr Pro Pro Glu Thr Pro Asn Ser Glu Arg Gly	245	250	255
20	Asn Val Met Ser Asn Thr Arg Tyr Ala Leu Glu Asn Gly Val Ala Val	260	265	270
	Phe Ala Thr Glu Trp Gly Thr Ser Gln Ala Asn Gly Asp Gly Gly Pro	275	280	285
25	Tyr Phe Asp Glu Ala Asp Val Trp Ile Glu Phe Leu Asn Glu Asn Asn	290	295	300
30	Ile Ser Trp Ala Asn Trp Ser Leu Thr Asn Lys Asn Glu Val Ser Gly	305	310	315
	Ala Phe Thr Pro Phe Glu Leu Gly Lys Ser Asn Ala Thr Asn Leu Asp	325	330	335
35	Pro Gly Pro Asp His Val Trp Ala Pro Glu Glu Leu Ser Leu Ser Gly	340	345	350
40	Glu Tyr Val Arg Ala Arg Ile Lys Gly Val Asn Tyr Glu Pro Ile Asp	355	360	365
45	Arg Thr Lys Tyr Thr Lys Val Leu Trp Asp Phe Asn Asp Gly Thr Lys	370	375	380
	Gln Gly Phe Gly Val Asn Ser Asp Ser Pro Asn Lys Glu Leu Ile Ala	385	390	395
50	Val Asp Asn Glu Asn Asn Thr Leu Lys Val Ser Gly Leu Asp Val Ser	405	410	415
55	Asn Asp Val Ser Asp Gly Asn Phe Trp Ala Asn Ala Arg Leu Ser Ala	420	425	430

EP 2 904 099 B1

	Asp	Gly	Trp	Gly	Lys	Ser	Val	Asp	Ile	Leu	Gly	Ala	Glu	Lys	Leu	Thr	
				435					440					445			
5	Met	Asp	Val	Ile	Val	Asp	Glu	Pro	Thr	Thr	Val	Ala	Ile	Ala	Ala	Ile	
				450				455					460				
	Pro	Gln	Ser	Ser	Lys	Ser	Gly	Trp	Ala	Asn	Pro	Glu	Arg	Ala	Val	Arg	
10							470				475					480	
	Val	Asn	Ala	Glu	Asp	Phe	Val	Gln	Gln	Thr	Asp	Gly	Lys	Tyr	Lys	Ala	
					485					490						495	
15	Gly	Leu	Thr	Ile	Thr	Gly	Glu	Asp	Ala	Pro	Asn	Leu	Lys	Asn	Ile	Ala	
				500					505					510			
	Phe	His	Glu	Glu	Asp	Asn	Asn	Met	Asn	Asn	Ile	Ile	Leu	Phe	Val	Gly	
20				515				520					525				
	Thr	Asp	Ala	Ala	Asp	Val	Ile	Tyr	Leu	Asp	Asn	Ile	Lys	Val	Ile	Gly	
25							535					540					
	Thr	Glu	Val	Glu	Ile	Pro	Val	Val	His	Asp	Pro	Lys	Gly	Glu	Ala	Val	
						550				555						560	
30	Leu	Pro	Ser	Val	Phe	Glu	Asp	Gly	Thr	Arg	Gln	Gly	Trp	Asp	Trp	Ala	
					565					570					575		
	Gly	Glu	Ser	Gly	Val	Lys	Thr	Ala	Leu	Thr	Ile	Glu	Glu	Ala	Asn	Gly	
35				580					585					590			
	Ser	Asn	Ala	Leu	Ser	Trp	Glu	Phe	Gly	Tyr	Pro	Glu	Val	Lys	Pro	Ser	
				595				600					605				
40	Asp	Asn	Trp	Ala	Thr	Ala	Pro	Arg	Leu	Asp	Phe	Trp	Lys	Ser	Asp	Leu	
				610			615					620					
45	Val	Arg	Gly	Glu	Asn	Asp	Tyr	Val	Ala	Phe	Asp	Phe	Tyr	Leu	Asp	Pro	
						630					635					640	
	Val	Arg	Ala	Thr	Glu	Gly	Ala	Met	Asn	Ile	Asn	Leu	Val	Phe	Gln	Pro	
50					645					650					655		
	Pro	Thr	Asn	Gly	Tyr	Trp	Val	Gln	Ala	Pro	Lys	Thr	Tyr	Thr	Ile	Asn	
				660					665					670			
55	Phe	Asp	Glu	Leu	Glu	Glu	Ala	Asn	Gln	Val	Asn	Gly	Leu	Tyr	His	Tyr	
				675				680					685				

Glu Val Lys Ile Asn Val Arg Asp Ile Thr Asn Ile Gln Asp Asp Thr
690 695 700

Leu Leu Arg Asn Met Met Ile Ile Phe Ala Asp Val Glu Ser Asp Phe
705 710 715 720

Ala Gly Arg Val Phe Val Asp Asn Val Arg Phe Glu Gly Ala Ala Thr
725 730 735

Thr Glu Pro Val Glu Pro Glu Pro Val Asp Pro Gly Glu Glu Thr Pro
740 745 750

Pro Val Asp Glu Lys Glu Ala Lys Lys Glu Gln Lys Glu Ala Glu Lys
755 760 765

Glu Glu Lys Glu Glu
770

Claims

1. A method for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, comprising contacting the bacteria with an aqueous composition comprising an endo-beta-1,4-glucanase, wherein the amino acid sequence of the endo-beta-1,4-glucanase has at least 80% identity, preferably at least 90% identity, more preferably at least 95% identity, and most preferably 100% identity to the amino acid sequence of SEQ ID NO: 1.
2. The method according to claim 1, wherein the bacteria are selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp.
3. The method of any of claims 1-2, wherein the aqueous composition comprises at least 1 mg/l of an endo-beta-1,4-glucanase.
4. The method of any of claims 1-3, wherein the aqueous composition also comprises a surfactant.
5. The method of any of claims 1-4, wherein the surface is a textile surface and the aqueous composition is a laundry detergent composition.
6. The method of any of claims 1-5, wherein the bacteria are *Brevundimonas* sp.
7. Use of an endo-beta-1,4-glucanase for reducing adhesion of bacteria to a surface, or releasing the bacteria from a surface to which they adhere, wherein the amino acid sequence of the endo-beta-1,4-glucanase has at least 80% identity, preferably at least 90% identity, more preferably at least 95% identity, and most preferably 100% identity to the amino acid sequence of SEQ ID NO: 1.
8. The use according to claim 7, wherein the bacteria are selected from the group consisting of *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., and *Sphingomonas* sp.

Patentansprüche

1. Verfahren zum Verringern von Adhäsion von Bakterien an eine Oberfläche, oder Freisetzen der Bakterien von einer Oberfläche, an die sie adhären, umfassend Inkontaktbringen der Bakterien mit einer wässrigen Zusammensetzung

zung, die eine Endo-beta-1,4-glucanase umfasst, wobei die Aminosäuresequenz der Endo-beta-1,4-glucanase mindestens 80% Identität, bevorzugt mindestens 90% Identität, stärker bevorzugt mindestens 95% Identität und am stärksten bevorzugt 100% Identität zur Aminosäuresequenz von SEQ ID NO: 1 aufweist.

- 5 2. Verfahren nach Anspruch 1, wobei die Bakterien aus der Gruppe bestehend aus *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp. und *Sphingomonas* sp. ausgewählt sind.
3. Verfahren nach einem beliebigen der Ansprüche 1-2, wobei die wässrige Zusammensetzung mindestens 1 mg/l einer Endo-beta-1,4-glucanase umfasst.
- 10 4. Verfahren nach einem beliebigen der Ansprüche 1-3, wobei die wässrige Zusammensetzung auch ein oberflächenaktives Mittel umfasst.
- 15 5. Verfahren nach einem beliebigen der Ansprüche 1-4, wobei die Oberfläche eine Textiloberfläche ist und die wässrige Zusammensetzung eine Wäschedetergensenzusammensetzung ist.
6. Verfahren nach einem beliebigen der Ansprüche 1-5, wobei die Bakterien *Brevundimonas* sp. sind.
- 20 7. Verwendung einer Endo-beta-1,4-glucanase zum Verringern von Adhäsion von Bakterien an eine Oberfläche, oder Freisetzen der Bakterien von einer Oberfläche, an die sie adhären, wobei die Aminosäuresequenz der Endo-beta-1,4-glucanase mindestens 80% Identität, bevorzugt mindestens 90% Identität, stärker bevorzugt mindestens 95% Identität und am stärksten bevorzugt 100% Identität zur Aminosäuresequenz von SEQ ID NO: 1 aufweist.
- 25 8. Verwendung nach Anspruch 7, wobei die Bakterien aus der Gruppe bestehend aus *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp. und *Sphingomonas* sp. ausgewählt sind.

Revendications

- 30 1. Méthode pour réduire l'adhésion de bactéries à une surface, ou libérer les bactéries d'une surface à laquelle elles adhèrent, comprenant la mise en contact des bactéries avec une composition aqueuse comprenant une endo-bêta-1,4-glucanase, dans laquelle la séquence d'acides aminés de l'endo-bêta-1,4-glucanase a une identité d'au moins 80 %, de préférence une identité d'au moins 90 %, mieux encore une identité d'au moins 95 % et tout spécialement une identité de 100 % avec la séquence d'acides aminés de SEQ ID NO : 1.
- 35 2. Méthode selon la revendication 1, dans laquelle les bactéries sont choisies dans le groupe constitué par *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., et *Sphingomonas* sp.
- 40 3. Méthode selon l'une quelconque des revendications 1 et 2, dans laquelle la composition aqueuse comprend au moins 1 mg/l d'une endo-bêta-1,4-glucanase.
4. Méthode selon l'une quelconque des revendications 1 à 3, dans laquelle la composition aqueuse comprend aussi un tensioactif
- 45 5. Méthode selon l'une quelconque des revendications 1 à 4, dans laquelle la surface est une surface textile et la composition aqueuse est une composition détergente pour le linge.
6. Méthode selon l'une quelconque des revendications 1 à 5, dans laquelle les bactéries sont *Brevundimonas* sp.
- 50 7. Utilisation d'une endo-bêta-1,4-glucanase pour réduire l'adhésion de bactéries à une surface, ou libérer les bactéries d'une surface à laquelle elles adhèrent, dans laquelle la séquence d'acides aminés de l'endo-bêta-1,4-glucanase a une identité d'au moins 80 %, de préférence une identité d'au moins 90 %, mieux encore une identité d'au moins 95 % et tout spécialement une identité de 100 % avec la séquence d'acides aminés de la SEQ ID NO : 1.
- 55 8. Utilisation selon la revendication 7, dans laquelle les bactéries sont choisies dans le groupe constitué par *Brevundimonas* sp., *Exiguobacterium* sp., *Sphingobium* sp., et *Sphingomonas* sp.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2009148800 A [0003]
- US 5356803 A [0003]
- WO 2008007318 A [0003]
- WO 2008007320 A [0003]
- EP 2157162 A [0003]
- WO 2004053039 A [0003]
- KR 20120083605 [0003]
- WO 9517413 A [0026]
- WO 9522625 A [0026]
- US 5223409 A [0026]
- WO 9206204 A [0026]
- WO 9000609 A [0031]
- WO 9516782 A [0031]
- WO 9110732 A [0032]
- WO 9117243 A [0032]
- EP 0651785 A [0032]
- WO 9817767 A [0044]
- EP 624154 A [0044]
- WO 02099091 A [0065]

Non-patent literature cited in the description

- NEEDLEMAN ; WUNSCH. *J. Mol. Biol.*, 1970, vol. 48, 443-453 [0021]
- RICE et al. EMBOSS: The European Molecular Biology Open Software Suite. *Trends Genet.*, 2000, vol. 16, 276-277 [0021]
- H. NEURATH ; R.L. HILL. *The Proteins*. Academic Press, 1979 [0023]
- CUNNINGHAM ; WELLS. *Science*, 1989, vol. 244, 1081-1085 [0025]
- HILTON et al. *J. Biol. Chem.*, 1996, vol. 271, 4699-4708 [0025]
- DE VOS et al. *Science*, 1992, vol. 255, 306-312 [0025]
- SMITH et al. *J. Mol. Biol.*, 1992, vol. 224, 899-904 [0025]
- WLODAVER et al. *FEBS Lett.*, 1992, vol. 309, 59-64 [0025]
- REIDHAAR-OLSON ; SAUER. *Science*, 1988, vol. 241, 53-57 [0026]
- BOWIE ; SAUER. *Proc. Natl. Acad. Sci. USA*, 1989, vol. 86, 2152-2156 [0026]
- LOWMAN et al. *Biochemistry*, 1991, vol. 30, 10832-10837 [0026]
- DERBYSHIRE et al. *Gene*, 1986, vol. 46, 145 [0026]
- NER et al. *DNA*, 1988, vol. 7, 127 [0026]
- NESS et al. *Nature Biotechnology*, 1999, vol. 17, 893-896 [0027]
- COOPER et al. *EMBO J.*, 1993, vol. 12, 2575-2583 [0029]
- DAWSON et al. *Science*, 1994, vol. 266, 776-779 [0029]
- MARTIN et al. *J. Ind. Microbiol. Biotechnol.*, 2003, vol. 3, 568-576 [0030]
- SVETINA et al. *J. Biotechnol.*, 2000, vol. 76, 245-251 [0030]
- RASMUSSEN-WILSON et al. *Appl. Environ. Microbiol.*, 1997, vol. 63, 3488-3493 [0030]
- WARD et al. *Biotechnology*, 1995, vol. 13, 498-503 [0030]
- CONTRERAS et al. *Biotechnology*, 1991, vol. 9, 378-381 [0030]
- EATON ; 1986 et al. *Biochemistry*, vol. 25, 505-512 [0030]
- COLLINS-RACIE et al. *Biotechnology*, 1995, vol. 13, 982-987 [0030]
- CARTER et al. *Proteins: Structure, Function, and Genetics*, 1989, vol. 6, 240-248 [0030]
- STEVENS. *Drug Discovery World*, 2003, vol. 4, 35-48 [0030]
- Cellulose-Binding Domains: Classification and Properties" in "Enzymatic Degradation of Insoluble Carbohydrates. PETER TOMME et al. ACS Symposium Series, No. 618. 1996 [0031]