



(11) **EP 3 037 549 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
29.06.2016 Bulletin 2016/26

(51) Int Cl.:
C12Q 1/68 (2006.01)

(21) Application number: **15020211.7**

(22) Date of filing: **02.08.2007**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE
SI SK TR**

(30) Priority: **02.08.2006 US 498976**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
07840663.4 / 2 046 996

(71) Applicant: **Mayo Foundation for Medical
Education and Research
Rochester, MN 55905 (US)**

(72) Inventors:
• **Mrazel, David A.
Rochester, MN 55902 (US)**

• **O'Kane, Dennis J.
Rochester, MN 55902 (US)**
• **Black, John L.
Rochester, MN 55902 (US)**

(74) Representative: **Crump, Julian Richard John
Beresford Crump LLP
16 High Holborn
London WC1V 6BX (GB)**

Remarks:

This application was filed on 29-10-2015 as a
divisional application to the application mentioned
under INID code 62.

(54) **METHODS FOR SELECTING MEDICATIONS**

(57) Methods for selecting a medication for a patient
are described that include determining the patient's geno-
type for a panel of genes and selecting the medication
based on the genotype. Articles of manufacture also are

provided that include nucleic acid molecules for detecting
alleles of genes encoding drug metabolizing enzymes
and genes encoding products involved in neurotransmis-
sion.

EP 3 037 549 A1

Description**TECHNICAL FIELD**

[0001] This invention relates to methods for selecting a medication (e.g., psychotropic medication) for a patient, and more particularly to selecting a patient's medication based on the genotype of genes encoding drug-metabolizing enzymes and genes encoding products involved in, for example, neurotransmission.

SUMMARY

[0002] The invention is based on the identification of a set of genes with polymorphisms that are associated with psychiatric diagnosis and pharmacological response to a medication. As a result, methods of the invention allow the genotype of a patient to be determined and, based on the genotype, a suitable medication to be selected for the patient. Methods of the invention allow the output of multiple genotypic assessments to be integrated, providing important and improved clinical information on which to select and dose medications. Thus, the methods of the invention provide a rational method for the identification of a medication that will result in an optimal response in the patient

[0003] In one aspect, the invention features a method for selecting a psychotropic medication (e.g., an antidepressant, an antipsychotic agent, an anti-anxiety-sedative agent, or a mood-stabilizing agent) for a patient. The method includes providing the patient's genotype for a panel of genes, wherein the panel includes at least three cytochrome P450 genes and a serotonin transporter gene and selecting the psychotropic medication based on the genotype. The at least three cytochrome P450 genes can encode CYP2D6, 1A2, and 2C19 and providing the patient's genotype can include determining if the patient has the CYP1A2*1A or 1A2*3 allele, the CYP2C19*1A, 2C19*1B, or 2C19*2A allele, and the CYP2D6*1A, 2D6*2, 2D6*2N, 2D6*3, 2D6*4, 2D6*5, 2D6*6, 2D6*7, 2D6*8, 2D6*10, 2D6*12, or 2D6*17 allele. The method further can include determining if the patient includes the 2D6*41 allele.

[0004] The panel further can include a serotonin receptor 2A gene. For example, the panel can include cytochrome P450 genes encoding CYP2D6, 2C19, and 1A2, a serotonin transporter gene, and a serotonin receptor 2A gene. Selecting the psychotropic medication can include correlating genotype of the cytochrome P450 genes with capacity of each cytochrome P450 enzyme encoded by each cytochrome P450 gene to metabolize the psychotropic medication and correlating genotype of the serotonin transporter gene and the serotonin receptor gene with ability of the patient to respond to the psychotropic medication.

[0005] The panel can include 4 cytochrome P450 genes (e.g., genes encoding CYP2D6, 2C 19, 3A4, and 1A2). Providing the patient's genotype can include determining if said patient contains the CYP2D6*2, 2D6*3, 2D6*4, 2D6*10, or 2D6*17 allele, the CYP2C19*2A, 2C19*2B, 2C19*3, 2C19*4, 2C19*5A, 2C19*5B, 2C19*6, 2C19*7, or 2C19*8 allele, the CYP3A4*1B, 3A4*2, 3A4*5, 3A4*6, 3A4*12, 3A4*13, 3A4*15A, 3A4*17, 3A4*18A allele, and the CYP1A2*1F allele.

[0006] The panel can include at least 5 cytochrome P450 genes (e.g., CYP1A1, 1A2, 2D6, 2C19, and 3A4). Providing the patient's genotype can include determining if the patient has the CYP1A1*1A, 1A1*2, 1A1*3, or 1A1*4 allele, the CYP1A2*1A or 1A2*3 allele, the CYP2C19*1A, 2C19*1B, or 2C19*2A allele, the CYP2D6*1A, 2D6*2, 2D6*2N, 2D6*3, 2D6*4, 2D6*5, 2D6*6, 2D6*7, 2D6*8, 2D6*10, 2D6*12, 2D6*17, or 2D6*35 allele, and the CYP3A4*1A or 3A4*1B allele.

[0007] The panel further can include a plurality of dopamine receptor genes (e.g., dopamine receptor genes encoding dopamine receptors D1, D2, D3, D4, D5, and D6), a plurality of serotonin receptor genes (e.g., serotonin receptor genes encoding serotonin receptors 1A, 1B, 1D, 2A, or 2C), a tryptophan hydroxylase gene, and/or a catechol-O-methyl transferase gene. Serotonin receptor gene 2A is particularly useful. In some embodiments, the panel of genes contains the CYP2D6, 2C19, 3A4, and 1A2 genes, the serotonin receptor gene, and the serotonin receptor 2A gene.

[0008] In some embodiments, the panel includes at least 10 cytochrome P450 genes (e.g., genes encoding CYP1A1, 1A2, 1B1, 2A6, 2B6, 2C8, 2C9, 2C18, 2C19, 2D6, 2E1, 3A4, and 3A5). Determining the patient's genotype can include determining if the patient has the CYP1A1*1A, 1A1*2, 1A1*3, or 1A1*4 allele; the 1A2*1A or 1A2*3 allele; the CYP1B1*1, 1B1*2, 1B1*3, 1B1*4, 1B1*11, 1B1*14, 1B1*18, 1B1*19, 1B1*20, or 1B1*25 allele; the CYP2A6*1A, 2A6*1B, 2A6*2, or 2A6*5 allele; the CYP2B6*1, 2B6*2, 2B6*3, 2B6*4, 2B6*5, 2B6*6, or 2B6*7 allele; the CYP2C8*1A, 2C8*1B, 2C8*1C, 2C8*2, 2C8*3, or 2C8*4 allele; the CYP2C9*1, 2C9*2, 2C9*3, or 2C9*5 allele; the CYP2C18*m1 or 2C18*m2 allele; the CYP2C19*1A, 2C19*1B, or 2C19*2A allele; the CYP2D6*1A, 2D6*2, 2D6*2N, 2D6*3, 2D6*4, 2D6*5, 2D6*6, 2D6*7, 2D6*8, 2D6*10, 2D6*12, 2D6*17, or 2D6*35 allele; the CYP2E1*1A, 2E1*1C, 2E1*1D, 2E1*2, 2E1*4, 2E1*5, or 2E1*7 allele; the CYP3A4*1A or 3A4*1B allele; and the CYP3A5*1A, 3A5*3, 3A5*5, or 3A5*6 allele.

[0009] In another aspect, the invention features an article of manufacture that includes a substrate. The substrate includes a plurality of discrete regions, wherein each region includes a different population of nucleic acid molecules, wherein the different populations of nucleic acid molecules independently include nucleic acid molecules for detecting the CYP1A2*1A and 1A2*3 alleles; the CYP2C19*1A, 2C19*1B, and 2C19*2A alleles; the CYP2D6*1A, 2D6*2, 2D6*2N, 2D6*3, 2D6*4, 2D6*5, 2D6*6, 2D6*7, 2D6*8, 2D6*10, 2D6*12, 2D6*17, and 2D6*35 alleles; and the 5HTT promoter repeat and exon 2 variable repeat alleles. The population of nucleic acid molecules further can include nucleic acid

molecules for detecting one or more of the following: CYP1A1*1A, 1A1*2, 1A1*3, and 1A1*4 alleles; CYP1B1*1, 1B1*2, 1B1*3, 1B1*4, 1B1*11, 1B1*14, 1B1*18, 1B1*19, 1B1*20, and 1B1*25 alleles, the CYP2A6*1A, 2A6*1B, 2A6*2, and 2A6*5 alleles, the CYP2B6*1, 2B6*2, 2B6*3, 2B6*4, 2B6*5, 2B6*6, and 2B6*7 alleles, the CYP2C8*1A, 2C8*1B, 2C8*1C, 2C8*2, 2C8*3, and 2C8*4 alleles, the CYP2C9*1, 2C9*2, 2C9*3, and 2C9*5 alleles, the CYP2C18*m1 and 2C18*m2 alleles, the CYP2C19*1A, 2C19*1B, and 2C19*2A alleles, the CYP2E1*1A, 2E1*1C, 2E1*1D, 2E1*2, 2E1*4, 2E1*5, and 2E1*7 alleles, the CYP3A4*1A and 3A4*1B alleles; the DAT1 40 bp VNTR and 10 repeat alleles; and the CYP3A5*1A, 3A5*3, 3A5*5, and 3A5*6 alleles. The population further can include nucleic acid molecules for detecting the TPH A218C, A779C, G5806T, A6526G, and (CT)m(CA)n(CT)p alleles.

[0010] The invention also features a method of building a database for use in selecting a medication for a patient. The method includes receiving, in a computer system, a plurality of genotypes for a panel of genes, the panel of genes including a CYP2D6 gene, a CYP2C19 gene, a CYP1A2 gene, a serotonin transporter gene, and a serotonin receptor 2A gene; receiving a plurality of medication profiles specified based on the genotypes; and storing the plurality of genotypes and the medication profiles such that each medication profile is associated with one of the genotypes. The at least one medication profile can identify a medication and the medication can be placed in one of multiple categories included in the medication profile. Such categories can be selected from the group consisting of: medications that are safe to use, medications that should be used with caution, medications that should be closely monitored when used, medications that should be avoided, and combinations thereof. The medication profile can identify a universe of possible medications for the patient's genotype.

[0011] In another aspect, the invention features a computer program product containing executable instructions that when executed cause a processor to perform operations. The operations can include: receive a plurality of genotypes for a panel of genes, wherein the panel of genes includes a CYP2D6 gene, a CYP2C19 gene, a CYP1A2 gene, a serotonin transporter gene, and a serotonin receptor 2A gene; receive a plurality of medication profiles specified based on the genotypes; and store the genotypes and the medication profiles such that each medication profile is associated with one of the genotypes.

[0012] The invention also features a method of selecting a medication for a patient. The method includes receiving, in a computer system, a patient's genotype for a panel of genes, wherein the panel of genes includes a CYP2D6 gene, a CYP2C19 gene, a CYP1A2 gene, a serotonin transporter gene, and a serotonin receptor 2A gene; identifying, in a database comprising a plurality of medication profiles associated with genotypes, a medication profile that is associated with the patient's genotype; and outputting the identified medication profile in response to receiving the patient's genotype. A user can enter the patient's genotype in the computer system or the patient's genotype can be received directly from equipment used in determining the patient's genotype.

[0013] The medication profile can include a ranking of several medications, e.g., based on specific co-factors. The method can include adjusting the ranking before outputting the identified medication profile (e.g., based on receiving a genotypic polymorphism carried by the patient or based on receiving a clinical response relating to the patient). The clinical response can be by a family member of the patient.

[0014] In yet another aspect, the invention features a computer program product containing executable instructions that when executed cause a processor to perform operations that include receiving a patient's genotype for a panel of genes, wherein the panel of genes includes a CYP2D6 gene, a CYP2C19 gene, a CYP1A2 gene, a serotonin transporter gene, and a serotonin receptor 2A gene; identify, in a database including a plurality of medication profiles associated with genotypes, a medication profile that is associated with the patient's genotype; and output the identified medication profile in response to receiving the patient's genotype.

[0015] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used to practice the invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

[0016] Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

DESCRIPTION OF THE DRAWINGS

[0017]

Figure 1 is a block diagram of a computer system 100, according to one embodiment.

Figure 2 is a flow chart of a method 200 for building a database for use in selecting a medication for a patient.

Figure 3 is a flow chart of a method 300 of selecting a medication for a patient.

DETAILED DESCRIPTION

[0018] In general, the invention features a method for selecting a medication (e.g., psychotropic medication) for a patient based on the genotype of genes that are useful for medication selection. Genes to be genotyped typically encode products that influence the metabolism of a medication or that are associated with a better treatment response. An algorithm can be used that initially excludes medications based on problematic genotypes of drug metabolizing enzyme genes (e.g., cytochrome P450 genes). A second stage of such an algorithm can begin with a list of potentially appropriate medications (e.g., antidepressants) and result in further classification of optimal treatments based on the genotype of target genes. For example, for antidepressants, further classification of the potential antidepressants can be based on the genotype of the serotonin transporter (5HTT) and serotonin receptor 2A (HTR2A).

[0019] Psychotropic medications include antipsychotics or neuroleptics, antianxiety-sedative agents, antidepressants or mood elevating agents, and mood-stabilizing drugs such as lithium salts or valproic acid. Non-limiting examples of antipsychotics include the tricyclic phenothiazines (e.g., chlorpromazine, trifluorpromazine, thioridazine, and mesoridazine, fluphenazine, and trifluoperazine), thioxanthenes (e.g., chlorprothixene, clopenthixol, flupenthixol, piflutixol, and thiothixene), and dibenzepines (e.g., loxapine, clozapine, clothiapine, metiapine, zotapine, ICI-204,636, fluperlapine, and olanzapine); butyrophenones such as haloperidol, diphenylbutylpiperidines such as fluspirilene, penfluridol, and pimozide, haloperidol decanoate, and indolones. Non-limiting examples of antianxiety-sedative agents include benzodiazepines such as chlordiazepoxide, diazepam, oxazepam, clorazepate, lorazepam, prazepam, alprazolam, and halazepam). Antidepressants or mood elevating agents include norepinephrine-reuptake inhibitors such as the tertiary amine tricyclics (e.g., amitriptyline, clomipramine, doxepin, and imipramine) and secondary amine tricyclics (e.g., amoxapine, desipramine, maprotiline, protriptyline, and nortriptyline); selective serotonin-reuptake inhibitors (SSRIs) such as fluoxetine, fluvoxamine, paroxetine, sertraline, citalopram, escitalopram, and venlafaxine; atypical antidepressants such as bupropion, nefazodone, and trazodone; noradrenergic and specific serotonergic antidepressants such as mirtazapine; and monoamine oxidase inhibitors such as phenelzine, tranylcypromine, and selegiline.

[0020] In certain ethnic groups as many as 10% of the adolescent population have a 2D6 haplotype that is associated with poor metabolism of many antidepressant medications. See Wong et al. (2001) Ann. Acad. Med. Singapore 29:401-406. Clinical genomic testing of these individuals has clear implications for their treatment and prognosis. In extreme cases, children who were poor metabolizers and who were not identified have had tragic outcomes. These negative case reports have included a reported death of a nine-year-old boy who was not recognized to be a poor 2D6 metabolizer. The treatment of this child with fluoxetine continued despite the development of multiple symptoms because these symptoms were not recognized as being related to his extremely high serum levels of fluoxetine. Sallee et al. (2000) J. Child Adol. Psychiatry 10(1):27-34. While careful clinical surveillance to identify unexpected side effects during the administration of low doses of medication is an alternative clinical strategy to genotyping, genomic testing of a plurality of genes encoding drug metabolizing enzymes (e.g. cytochrome P450 genes) and other target genes (e.g., genes involved in neurotransmission for psychotropic drugs) provides a safe method by which potentially dangerous side effects can be avoided in an affected patient.

Panels of Genes

[0021] The method includes obtaining a biological sample from a patient and obtaining the patient's genotype for a panel of genes. Typically, the panel of genes that are genotyped includes at least three cytochrome P450 genes. The cytochrome P450 genes can be selected from the P450 genes listed in Table 1. For example, the at least three cytochrome P450 genes can encode CYP2D6, 1A2, and 2C 19. In embodiments for selecting antidepressant medications, the genotype of four cytochrome P450 genes (e.g., genes encoding CYP2D6, 2C19, 3A4, and 1A2) can be obtained. In other embodiments, the genotype of at least 5 cytochrome P450 genes (e.g., genes encoding CYP1A1, 1A2, 2D6, 2C 19, and 3A4) can be obtained. In still other embodiments, the genotype of at least 10 cytochrome P450 genes (e.g., genes encoding CYP1A1, 1A2, 1B1, 2A6, 2B6, 2C8, 2C9, 2C18, 2C19, 2D6, 2E1, 3A4, and 3A5) can be obtained. Alleles for each of these cytochrome P450 genes are set forth in Table 1.

[0022] Substrates of CYP2D6 typically are weak bases with the cationic binding site located away from the carbon atom to be oxidized. In particular, substrates of CYP2D6 include amitriptyline, nortriptyline, haloperidol, and desipramine. Venlafaxine is another substrate of CYP2D6 that can be biotransformed to an active metabolite, O-desmethylvenlafaxine, primarily by the CYP2D6 enzyme. Venlafaxine also can be metabolized to an inactive metabolite, N-desmethylvenlafaxine, which requires the CYP2D6 enzyme. Some individuals have altered CYP2D6 gene sequences that result in synthesis of enzymes devoid of catalytic activity or in enzymes with diminished catalytic activity. These individuals metabolize SSRIs (e.g., venlafaxine) and tricyclic antidepressants (TCAs) poorly. Duplication of the functional CYP2D6 gene also has been observed and results in ultrarapid metabolism of SSRIs and other drugs. Individuals without inactivating polymorphisms, deletions, or duplications have the phenotype of an extensive drug metabolizer and are designated as CYP2D6*1. The CYP2D6*2 allele has decreased enzymatic activity resulting from amino acid substitutions. The

CYP2D6*3 and *4 alleles account for nearly 70% of the total deficiencies that result in the poor metabolizer phenotype. The polymorphism responsible for CYP2D6*3 (2549A>del) produces a frame-shift in the mRNA. A polymorphism involved with the CYP2D6*4 allele (1846G>A) disrupts mRNA splicing. These changes produce truncated forms of CYP2D6 devoid of catalytic activity. Other poor metabolizers are CYP2D6*5, *10, and *17. CYP2D6*5 is due to complete gene deletion. The polymorphisms in CYP2D6*10 and *17 produce amino acid substitutions in the CYP2D6 enzyme which have decreased enzyme activity. All of these polymorphisms are autosomal recessive. Consequently, only individuals who are homozygous or who are compound heterozygous for these polymorphisms are poor metabolizers. Individuals who are heterozygous, with one normal gene and one polymorphic gene, will have metabolism intermediate between the extensive (normal) and poor metabolizers.

[0023] CYP1A2 metabolizes many aromatic and heterocyclic amines including clozapine and imipramine. The CYP1A2 *1F allele can result in a product with higher inducibility or increased activity. See Sachse et al. (1999) Br. J. Clin. Pharmacol. 47: 445-449. CYP2C19 also metabolizes many substrates including imipramine, citalopram, and diazepam. The CYP2C19 *2A, *2B, *3, *4, *5A, *5B, *6, *7, and *8 alleles encode products with little or no activity. See Ibeanu et al. (1999) J. Pharmacol. Exp. Ther. 290: 635-640.

[0024] CYP1A1 can be associated with toxic or allergic reactions by extrahepatic generation of reactive metabolites. CYP3A4 metabolizes a variety of substrates including alprazolam. CYP1B1 can be associated with toxic or allergic reactions by extrahepatic generation of reactive metabolites and also metabolizes steroid hormones (e.g., 17 β -estradiol). Substrates for CYP2A6 and CYP2B6 include valproic acid and bupropion, respectively. Substrates for CYP2C9 include Tylenol and antabuse (disulfuram). Substrates for CYP2E1 include phenytoin and carbamazepine. Decreases in activity in one or more of the cytochrome P450 enzymes can impact one or more of the other cytochrome P450 enzymes.

Table 1

Cytochrome P450 Genes		
Cytochrome P450 Gene	Allele	Polymorphism
1A1	*1A *2 *3 *4	None A2455G T3205C C2453A
1A2	*1A *1F *3	None -164C>A G1042A
1B1	*1 *2 *3 *4 *11 *14 *18 *19 *20 *25	None R48G L432V N453S V57C E281X G365W P379L E387K R469W
2A6	*1A *1B *2 *5	None CYP2A7 translocated to 3'-end T479A *1B + G6440T
2B6	*1 *2 *3 *4 *5 *6 *7	R22C S259C K262R R487C Q172H; K262R Q172H; K262R; R487C

EP 3 037 549 A1

(continued)

Cytochrome P450 Genes		
Cytochrome P450 Gene	Allele	Polymorphism
2C8	*1A *1B *1C *2 *3 *4	None -271C>A -370T>G I269F R139K; K399R I264M
2C9	*1 *2 *3 *5	None R144C I359L D360E
2C18	m1 m2	T204A A460T
2C19	*1A *1B *2A *2B *3 *4 *5(A,B) *6 *7 *8	None I331V Splicing defect Splicing defect; E92D New stop codon 636G>A GTG initiation codon, 1A>G 1297C>T, amino acid change (R433W) 395G>A, amino acid change (R132Q) IVS5+2T>A, splicing defect 358T>C, amino acid change (W120R)
2D6	*1A *2 *2N *3 *4 *5 *6 *7 *8 *10 *12 *17 *35	None G1661C, C2850T Gene duplication A2549 deletion G1846A Gene deletion T1707 deletion A2935C G1758T C100T G124A C1023T, C2850T G31A
2E1	*1A *1C, *1D *2 *4 *5 *5 *7 *7 *7	None (6 or 8 bp repeats) G1132A G476A G(-1293)C C(-1053)T T(-333)A G(-71)T A(-353)G
3A4	*1A *1B	None A(-392)G

(continued)

Cytochrome P450 Genes		
Cytochrome P450 Gene	Allele	Polymorphism
	*2	Amino acid change (S222P)
	*5	Amino acid change (P218R)
	*6	Frameshift, 831 ins A
	*12	Amino acid change (L373F)
	*13	Amino acid change (P416L)
	*15A	Amino acid change (R162Q)
	*17	Amino acid change (F189S, decreased)
	*18A	Amino acid change (L293P, increased)
3A5	*1A	None
	*3	A6986G
	*5	T12952C
	*6	G14960A

[0025] Typically, to select a medication, the genotype of other target genes also is obtained in addition to the genotype of the genes encoding drug-metabolizing enzymes. Target genes can encode products that relate to the ability of the patient to respond to a particular class of medication. For example, to select an antidepressant, the target genes can be the serotonin transporter gene and serotonin receptor 2A gene. As such, in one embodiment, the panel of genes can be the CYP2D6 gene, 1A2 gene, and 2C19 gene, the serotonin transporter gene, and serotonin receptor 2A gene. To select an antipsychotic, the target gene can be a dopamine transporter gene.

[0026] Table 2 sets forth the alleles for target genes that can be genotyped in addition to the genes encoding drug-metabolizing enzymes. For example, the panel of genes to be genotyped can include one or more dopamine receptor genes (e.g., genes encoding dopamine receptors D1, D2, D3, D4, D5, and/or D6), a dopamine transporter gene, one or more serotonin receptor genes (e.g., genes encoding serotonin receptors 1A, 1B, 1D, 2A, or 2C), a catechol-O-methyl transferase (COMT) gene, or a tryptophan hydroxylase gene. In one embodiment, a COMT gene and a tryptophan hydroxylase gene are included on the panel with cytochrome P450 genes. In other embodiments, one or more dopamine receptor genes, one or more serotonin receptor genes, a COMT gene, and a tryptophan hydroxylase gene are assessed in combination with the cytochrome P450 genes.

Table 2

Gene	Symbol	Polymorphism
Dopamine Transporter	DAT1,	40 bp VNTR
	SLC6A3	10 repeat allele G710A, Q237R C124T, L42F
Dopamine Receptor D1	DRD1	DRDL1 B2 T244G C179T G127A T11G C81T T595G, S199A G150T, R50S C110G, T37R A109C, T37P
Dopamine Receptor D2	DRD2	TaqI A

EP 3 037 549 A1

(continued)

Gene	Symbol	Polymorphism
		A1051G, T35A C932G, S311C C928, P310S G460A, V154I
Dopamine Receptor D3	DRD3	Ball in exon I MspI DRD3 1 Gly/Ser (allele 2) A25G, S9G
Dopamine Receptor D4	DRD4	48 repeat in exon 3 7 repeat allele 12/13 bp insertion/deletion T581G, V194G C841G, P281A
Dopamine Receptor D5	DRD5	T978C L88F A889C, T297P G1252A, V418I G181A, V61M G185C, C62S T263G, R88L G1354A, W455
Tryptophan Hydroxylase	TPH	A218C A779C G-5806T A-6526G (CT) _m (CA) _n (CT) _p allele 194 in 3' UTR, 5657 bp distant from exon 11
Serotonin Transporter	5-HTTR	Promoter repeat (44 bp insertion (L)/deletion (S) (L = Long form; S = Short form); Intron 2 variable number of repeats (9, 10, 11, or 12); A1815C; G603C; G167C; -3745, T→A (5'FR); -3636, T→C (5'FR); -3631 G→A (5'FR); SNP rs25531, A→G (5'FR); -1090, A→T (5'FR); -1089, A→T (5'FR); -859, A→C (5'FR); -482, T→C (5'FR); -469, C→T (5'FR); -45, C→A (intron 1A); -25, G→A intron 1A; -185, A→C (5' UTR); -149, C→A (5' UTR); G28A (intron 1b); T303C (exon 2); -100, G→A (intron 4); C83T (intron 7); C1149T (exon 8); T204G (intron 8); -131, C→T (intron 11)
Serotonin Receptor 1A	HTR1A	RsaI G815A, G272D G656T, R219L C548T, P551L A82G, I28V G64A, G22S C47T, P16L

(continued)

Gene	Symbol	Polymorphism
Serotonin Receptor 1B	HTR1B	G861C G861C, V287V T371G, F124C T655C, F219L A1099G, I367V G1120A, E374K
Serotonin Receptor 1D	HTR1D	G506T C173T C794T, S265L
Serotonin Receptor 2A	HTR2A	C74A T102C T516C C1340T C1354T
Serotonin Receptor 2C	HTR2C	G796C C10G, L4V G68C, C23S
Catechol-o-methyltransferase	COMT	G158A (Also known as Val/Met) G214T A72S G101C C34S G473A

[0027] For example, to select an antidepressant, the genotype of three (e.g., genes encoding 2D6, 2C19, and 1A2) or four cytochrome P450 genes (e.g., genes encoding 2D6, 2C19, 3A4, and 1A2), the serotonin transporter (5HTTR) gene, and serotonin receptor 2A (HTR2A) gene can be assessed in a patient. In particular, it can be determined if the patient contains the CYP2D6 *2, *3, *4, *10, *17, or *5 del allele, the CYP2C19 *2(A,B), *3, *4, *5 (A,B), *6, *7, *8 alleles, the CYP1A2 *1F allele, the short or long form of the serotonin transporter gene, and the HTR2A T102C polymorphism. In embodiments in which the 3A4 gene is included in the panel, it can be determined if the patient contains the CYP3A4 *1B, *2, *5, *6, *12, *13, *15A, *17, or *18A alleles. Each of these genes influences the metabolism of at least one antidepressant medication or is associated with a better treatment response. As described herein, an algorithm has been created based on a set of six rules relating to the genotype of the six genes (e.g., 2D6 gene, 2C19 gene, 3A4 gene, 1A2 gene, the 5HTTR gene, and HTR2A gene). Similarly, an algorithm can be created based on a set of five rules relating to the five genes (e.g., 2D6 gene, 2C19 gene, 1A2 gene, the 5HTTR gene, and HTR2A gene). Based on these algorithms, medication profiles are provided for a given patient based on the patient's genotype, allowing a clinician to select an acceptable antidepressant without the trial and error of determining if the patient will respond or tolerate a particular antidepressant.

Determining Genotype

[0028] Genomic DNA generally is used to determine genotype, although mRNA also can be used. Genomic DNA is typically extracted from a biological sample such as a peripheral blood sample, but can be extracted from other biological samples, including tissues (e.g., mucosal scrapings of the lining of the mouth or from renal or hepatic tissue). Routine methods can be used to extract genomic DNA from a blood or tissue sample, including, for example, phenol extraction. Alternatively, genomic DNA can be extracted with kits such as the QIAamp® Tissue Kit (Qiagen, Chatsworth, CA),

Wizard® Genomic DNA purification kit (Promega) and the A.S.A.P.™ Genomic DNA isolation kit (Boehringer Mannheim, Indianapolis, IN).

[0029] Typically, an amplification step is performed before proceeding with the genotyping. For example, polymerase chain reaction (PCR) techniques can be used to obtain amplification products from the patient. PCR refers to a procedure or technique in which target nucleic acids are enzymatically amplified. Sequence information from the ends of the region of interest or beyond typically is employed to design oligonucleotide primers that are identical in sequence to opposite strands of the template to be amplified. PCR can be used to amplify specific sequences from DNA as well as RNA, including sequences from total genomic DNA or total cellular RNA. Primers are typically 14 to 40 nucleotides in length, but can range from 10 nucleotides to hundreds of nucleotides in length. General PCR techniques are described, for example in PCR Primer: A Laboratory Manual, Ed. by Dieffenbach, C. and Dveksler, G., Cold Spring Harbor Laboratory Press, 1995. When using RNA as a source of template, reverse transcriptase can be used to synthesize complementary DNA (cDNA) strands. Ligase chain reaction, strand displacement amplification, self-sustained sequence replication or nucleic acid sequence-based amplification also can be used to obtain isolated nucleic acids. See, for example, Lewis (1992) Genetic Engineering News 12(9):1; Guatelli et al. (1990) Proc. Natl. Acad. Sci. USA 87:1874-1878; and Weiss (1991) Science 254:1292-1293.

[0030] Primers typically are single-stranded or double-stranded oligonucleotides that are 10 to 50 nucleotides in length, and when combined with mammalian genomic DNA and subjected to PCR conditions, is capable of being extended to produce a nucleic acid product corresponding to a region of interest within a gene. Typically, PCR products are at least 30 nucleotides in length (e.g., 30, 35, 50, 100, 250, 500, 1000, 1500, or 2000 or more nucleotides in length). Primers such as those listed in Table 7 are particularly useful for producing PCR products for the genes encoding the dopamine transporter, dopamine receptors, tryptophan hydroxylase, serotonin transporter, serotonin receptors, and COMT. Specific regions of mammalian DNA can be amplified (i.e., replicated such that multiple exact copies are produced) when a pair of oligonucleotide primers is used in the same PCR reaction, wherein one primer contains a nucleotide sequence from the coding strand of a nucleic acid and the other primer contains a nucleotide sequence from the non-coding strand of the nucleic acid. The "coding strand" of a nucleic acid is the nontranscribed strand, which has the same nucleotide sequence as the specified RNA transcript (with the exception that the RNA transcript contains uracil in place of thymidine residues), while the "non-coding strand" of a nucleic acid is the strand that serves as the template for transcription.

[0031] A single PCR reaction mixture may contain one pair of oligonucleotide primers. Alternatively, a single reaction mixture may contain a plurality of oligonucleotide primer pairs, in which case multiple PCR products can be generated (e.g., 5, 10, 15, or 20 primer pairs). Each primer pair can amplify, for example, one exon or a portion of one exon. Intron sequences also can be amplified.

[0032] Exons or introns of a gene of interest can be amplified then directly sequenced. Dye primer sequencing can be used to increase the accuracy of detecting heterozygous samples. Alternatively, one or more of the techniques described below can be used to determine genotype.

[0033] For example, allele specific hybridization can be used to detect sequence variants, including complete haplotypes of a mammal. See, Stoneking et al., 1991, Am. J. Hum. Genet. 48:370-382; and Prince et al., 2001, Genome Res., 11(1):152-162. In practice, samples of DNA or RNA from one or more mammals can be amplified using pairs of primers and the resulting amplification products can be immobilized on a substrate (e.g., in discrete regions). Hybridization conditions are selected such that a nucleic acid probe can specifically bind to the sequence of interest, e.g., the variant nucleic acid sequence. Such hybridizations typically are performed under high stringency as some sequence variants include only a single nucleotide difference. High stringency conditions can include the use of low ionic strength solutions and high temperatures for washing. For example, nucleic acid molecules can be hybridized at 42°C in 2X SSC (0.3M NaCl/0.03 M sodium citrate/0.1% sodium dodecyl sulfate (SDS) and washed in 0.1 X SSC (0.015M NaCl/0.0015 M sodium citrate), 0.1% SDS at 65°C. Hybridization conditions can be adjusted to account for unique features of the nucleic acid molecule, including length and sequence composition. Probes can be labeled (e.g., fluorescently) to facilitate detection. In some embodiments, one of the primers used in the amplification reaction is biotinylated (e.g., 5' end of reverse primer) and the resulting biotinylated amplification product is immobilized on an avidin or streptavidin coated substrate (e.g., in discrete regions).

[0034] Allele-specific restriction digests can be performed in the following manner. For nucleotide sequence variants that introduce a restriction site, restriction digest with the particular restriction enzyme can differentiate the alleles. For sequence variants that do not alter a common restriction site, mutagenic primers can be designed that introduce a restriction site when the variant allele is present or when the wild type allele is present. A portion of the nucleic acid of interest can be amplified using the mutagenic primer and a wild type primer, followed by digest with the appropriate restriction endonuclease.

[0035] Certain variants, such as insertions or deletions of one or more nucleotides, change the size of the DNA fragment encompassing the variant. The insertion or deletion of nucleotides can be assessed by amplifying the region encompassing the variant and determining the size of the amplified products in comparison with size standards. For example, a region of a gene of interest can be amplified using a primer set from either side of the variant. One of the primers is

typically labeled, for example, with a fluorescent moiety, to facilitate sizing. The amplified products can be electrophoresed through acrylamide gels with a set of size standards that are labeled with a fluorescent moiety that differs from the primer.

[0036] PCR conditions and primers can be developed that amplify a product only when the variant allele is present or only when the wild type allele is present (MSPCR or allele-specific PCR). For example, patient DNA and a control can be amplified separately using either a wild type primer or a primer specific for the variant allele. Each set of reactions is then examined for the presence of amplification products using standard methods to visualize the DNA. For example, the reactions can be electrophoresed through an agarose gel and the DNA visualized by staining with ethidium bromide or other DNA intercalating dye. In DNA samples from heterozygous patients, reaction products would be detected in each reaction. Patient samples containing solely the wild type allele would have amplification products only in the reaction using the wild type primer. Similarly, patient samples containing solely the variant allele would have amplification products only in the reaction using the variant primer. Allele-specific PCR also can be performed using allele-specific primers that introduce priming sites for two universal energy-transfer-labeled primers (e.g., one primer labeled with a green dye such as fluoroscein and one primer labeled with a red dye such as sulforhodamine). Amplification products can be analyzed for green and red fluorescence in a plate reader. See, Myakishev et al., 2001, *Genome* 11(1):163-169.

[0037] Mismatch cleavage methods also can be used to detect differing sequences by PCR amplification, followed by hybridization with the wild type sequence and cleavage at points of mismatch. Chemical reagents, such as carbodiimide or hydroxylamine and osmium tetroxide can be used to modify mismatched nucleotides to facilitate cleavage.

[0038] Kits also are available commercially to detect many of the cytochrome P450 variants. For example, TAG-IT™ kits are available from Tm Biosciences Corporation (Toronto, Ontario).

Selecting Medications

[0039] After the genotype is determined for each gene on the panel, the medication can be selected. Typically, selecting includes correlating genotype of the cytochrome P450 genes with capacity of each cytochrome P450 enzyme encoded by each cytochrome P450 gene to metabolize the medication. The genotype of other target genes on the panel, e.g., the serotonin transporter and serotonin receptor 2A, can be correlated with the ability of the patient to respond to the medication.

[0040] An algorithm can be used to select the most appropriate medications for an individual patient. The design of the algorithm requires the initial identification of the phenotype, which provides a preliminary identification of the universe of possible medications. At the next step of the algorithm, the results of the target gene analyses can be sequentially entered. The potential list of appropriate medications can be subsequently rank-ordered based on specific co-factors in the algorithmic equation, which assign a positive, negative, or neutral probability score to each of the identified medications in the originally identified set of possible choices. This process adjusts the rank ordering based on the genotypic polymorphism carried by the patient. At the next entry point in the algorithmic equation, the results of the CYP gene analyses can be introduced. This algorithmic analysis is designed to place the medications in three categories: 1) medications that are acceptable for use, i.e., the medication has a high probability of normal metabolism within an individual having a particular genotype, 2) medications that can be used with caution (e.g., medication may require some dosing adjustment based on atypical metabolism); and 3) medications that should be avoided or used with caution and monitoring, e.g., due to potential difficulties in dosing. At this point in the selection process, data related to the medication response of first- and second-degree relatives of the patient can be entered into the algorithmic equation, which pertains to the medication selection of drugs in the first category that has been identified. An adjustment of the rank-ordered, appropriate medications then can be calculated based on clinical responses by family members.

[0041] The selection of an appropriate medication further can be enhanced by including both target data and data related to drug metabolism. This can determine the impact of the CYP products on the clinical response of a particular patient. For example, inclusion of target data and data related to drug metabolism provides the amount of available drug, the ability of the patient to utilize the drug, and information about the quality of the receptor target of the drug, providing a rational approach to selection of medication.

[0042] An example of this process would be the selection of an appropriate antidepressant for a given patient. Once the phenotype of depression is established, the initial universe of antidepressant medications is identified. For example, citalopram, fluvoxamine, bupropion, escitalopram, sertraline, mirtazapine, fluoxetine, venlafaxine, amitriptyline, imipramine, paroxetine, and nortriptyline can be the initial universe of antidepressant medications. It is understood that this initial universe of medications can change as additional antidepressants become available or as additional pharmacogenomic data are available. Subsequently, the data from the target genotyping is entered. If during the first components of the algorithm, it is revealed that the genotype reflects poor metabolism for 2D6 and normal metabolism for 1A2, the rank ordering of medication would be to identify fluvoxamine and bupropion as medications that are likely to be both effective and appropriately managed at recommended doses. At the next stage of the algorithm, if an individual is homozygous for the long form of the serotonin transporter gene, fluvoxamine would be confirmed as being acceptable for use. Output from the algorithm then can be integrated with historical data. For example, if a family member had

responded well to fluvoxamine, this would confirm that the medication is acceptable for use, or, if a first or second degree relative had a problematic response to this medication, an alternative could be chosen.

Computer Systems

[0043] Techniques described herein can be implemented in a computer system having a processor that executes specific instructions in a computer program. The computer system may be arranged to output a medication profile based on receiving a patient's genotype. Particularly, the computer program may include instructions for the system to select the most appropriate medication (e.g., a psychotropic medication) for an individual patient.

[0044] The following are examples of features that may be included in a system. The computer program may be configured such that the computer system can identify the phenotype based on received data and provide a preliminary identification of the universe of possible medications. The system may be able to rank-order the identified medications based on specific co-factors in the algorithmic equation. The system may be able to adjust the rank ordering based on the genotypic polymorphism(s) carried by the patient. The system may be able to adjust the rank ordering based on clinical responses, such as by family members of the patient.

[0045] Figure 1 is a block diagram of a computer system 100 that can be used in the operations described above, according to one embodiment. The system 100 includes a processor 110, a memory 120, a storage device 130 and an input/output device 140. Each of the components 110, 120, 130 and 140 are interconnected using a system bus 150. The system may include analyzing equipment 160 for determining the patient's genotype.

[0046] The processor 110 is capable of processing instructions for execution within the system 100. In one embodiment, the processor 110 is a single-threaded processor. In another embodiment, the processor 110 is a multi-threaded processor. The processor 110 is capable of processing instructions stored in the memory 120 or on the storage device 130, including for receiving or sending information through the input/output device 140.

[0047] The memory 120 stores information within the system 100. In one embodiment, the memory 120 is a computer-readable medium. In one embodiment, the memory 120 is a volatile memory unit. In another embodiment, the memory 120 is a non-volatile memory unit.

[0048] The storage device 130 is capable of providing mass storage for the system 100. In one embodiment, the storage device 130 is a computer-readable medium. In various different embodiments, the storage device 130 may be a floppy disk device, a hard disk device, an optical disk device, or a tape device.

[0049] The input/output device 140 provides input/output operations for the system 100. In one embodiment, the input/output device 140 includes a keyboard and/or pointing device. In one embodiment, the input/output device 140 includes a display unit for displaying graphical user interfaces.

[0050] The system 100 can be used to build a database. Figure 2 shows a flow chart of a method 200 for building a database for use in selecting a medication for a patient. Preferably, the method 200 is performed in the system 100. For example, a computer program product can include instructions that cause the processor 110 to perform the steps of the method 200. The method 200 includes the following steps.

[0051] Receiving, in step 210, a plurality of genotypes 170 for a panel of genes. A computer program in the system 100 may include instructions for presenting a suitable graphical user interface on input/output device 140, and the graphical user interface may prompt the user to enter the genotypes 170 using the input/output device 140, such as a keyboard.

[0052] Receiving, in step 220, a plurality of medication profiles 180. The medication profiles 180 are specified based on the genotypes 170. The user may enter the medication profiles 180 using the input/output device 140, such as a keyboard. For example, the medication profile 180 may include information 190 regarding at least one medication.

[0053] Storing, in step 230, the received genotypes 170 and the medication profiles 180 such that each medication profile 180 is associated with one of the genotypes 170. The system 100 may store the medication profiles 180 and the genotypes 170 in the storage device 130. For example, when the storing is complete, the system 100 can identify a particular one of the medication profiles 180 that is associated with a specific genotype 170. Having identified the medication profile 180, the system 100 can access the information 190 contained within the identified medication profile 180, as will be described in the following example.

[0054] The system 100 may be used for selecting a medication. Figure 3 shows a flow chart of a method 300 of selecting a medication for a patient. Preferably, the method 300 is performed in the system 100. For example, a computer program product can include instructions that cause the processor 110 to perform the steps of the method 300. The method 300 includes the following steps.

[0055] Receiving, in step 310, a patient's genotype for a panel of genes. The genotype may be entered by a user via input/output device 140. For example, the user may obtain the patient's genotype for a panel of genes using the analyzing equipment 160 (which may or may not be connected to the system 100). The user may type the patient's genotype on input/output device 140, such as a keyboard, for receipt by the system 100.

[0056] The genotype may be received directly from the analyzing equipment 160. For example, analyzing equipment

160 may include a processor and suitable software such that it can communicate over a network. The system 100 may be connected to the analyzing equipment 160 through input/output device 140, such as a network adapter, and directly receive the patient's genotype.

[0057] Identifying, in step 320, one of the medication profiles 180 that is associated with the patient's genotype. For example, the system 100 may perform a database search in the storage device 130. Particularly, the system 100 may access the genotype 170 for individual medication profiles 180 until a match is found. Optional step 325 will be described below.

[0058] Outputting, in step 330, the identified medication profile 180 in response to receiving the patient's genotype. The system may output the identified medication profile 180 through input/output device 140. For example, the identified medication profile may be printed or displayed in a suitable graphical user interface on a display device. As another example, the system 100 may transmit the identified medication profile over a network, such as a local area network or the Internet, to which the input/output device 140 is connected.

[0059] The medication profiles 180 can be created such that there is flexibility in how the system 100 outputs them. For example, the information 190 in one or more of the medication profiles 180 may include a ranking of several medications. The program may include instructions for applying rules to the received patient's genotype and adjust the ranking accordingly. In such implementations, the method 300 may include optional step 325 of adjusting the ranking before outputting the identified medication profile. For example, the system 100 may receive a genotypic polymorphism carried by the patient (optionally in the same way the patient's genotype was received) and adjust the ranking accordingly in step 325. As another example, step 325 may involve adjusting the ranking based on a clinical response. The clinical response may be received by the system 100 in the same way as the patient's genotype. For example, the ranking can be adjusted based on a clinical response by a member of the patient's family.

[0060] The medication profiles 180 may be updated as necessary. For example, the introduction of a new medication on the market may prompt a revision of one or more existing medication profiles. A new medication may also be the basis for creating a new medication profile. The adjustment or creation of medication profiles may be done substantially as described above.

[0061] The medication profiles 180 may be used for medication selection in the same system where they were created, or in a different system. That is, the system 100 may first be used for building a database of the medication profiles 180, and the system 100 may thereafter be used to select a medication profile for the genotype of a specific patient. As another example, one or more medication profiles 180 may be transmitted within a computer readable medium such as a global computer network for remote processing according to the invention.

Articles of Manufacture

[0062] In one embodiment, the article of manufacture is a composition containing oligonucleotide primers. Typically, such a composition will contain a first oligonucleotide primer and a second oligonucleotide primer, each 10 to 50 nucleotides in length, which can be combined with genomic DNA from a mammal and subjected to PCR conditions, to produce a nucleic acid product that corresponds to a region of interest within a target gene. A composition also may contain buffers and other reagents necessary for PCR (e.g., DNA polymerase or nucleotides). Furthermore, a composition may contain one or more additional pairs of oligonucleotide primers (e.g., 5, 10, 15, or 20 primer pairs), such that multiple nucleic acid products can be generated.

[0063] In other embodiments, articles of manufacture include populations of nucleic acid molecules immobilized on a substrate. Suitable substrates provide a base for the immobilization of the nucleic acids, and in some embodiments, allow immobilization of nucleic acids into discrete regions. In embodiments in which the substrate includes a plurality of discrete regions, different populations of isolated nucleic acids can be immobilized in each discrete region. The different populations of nucleic acid molecules independently can include nucleic acid molecules for detecting one or more of the alleles set forth in Tables 1 and 2.

[0064] Suitable substrates can be of any shape or form and can be constructed from, for example, glass, silicon, metal, plastic, cellulose or a composite. For example, a suitable substrate can include a multiwell plate or membrane, a glass slide, a chip, or polystyrene or magnetic beads. Nucleic acid molecules or polypeptides can be synthesized *in situ*, immobilized directly on the substrate, or immobilized via a linker, including by covalent, ionic, or physical linkage. Linkers for immobilizing nucleic acids and polypeptides, including reversible or cleavable linkers, are known in the art. See, for example, U.S. Patent No. 5,451,683 and WO98/20019. Immobilized nucleic acid molecules typically are about 20 nucleotides in length, but can vary from about 10 nucleotides to about 1000 or more nucleotides in length.

[0065] The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

EXAMPLES

Example 1

CYP450 Genotype of Patients

[0066] This experiment highlights the results of an assessment of adolescents with atypical responses to selective serotonin reuptake inhibitors. The following CYP2D6 alleles that produce deficiencies were detected: *2 (1661G>C and 2850C>T), *3 (2549A>del), *4 (1661G>C and 1846G>A), *10 (100C>T, 1661G>C and 1846G), *17 (1023C>T and 2850C>T). Deletion of CYP2D6 (*5 del) was detected by electrophoresis following PCR amplification. Tandem duplication of the CYP2D6 gene was detected by amplification of a duplication-specific sequence with detection following electrophoresis.

[0067] Genomic DNA was amplified by the polymerase chain reaction in an Applied Biosystems Thermal Cycler using the primers listed below. The lower case letters "f" and "r" imply the direction of synthesis ("f" indicates a forward/sense synthesis reaction and "r" indicates a reverse/antisense synthesis reaction). The amplified DNA was hybridized with oligonucleotide probes (described below) specific for the normal and polymorphic variants and detected using the Nogen Molecular Biology Workstation. The oligonucleotide probes were fluorescently labeled (e.g., CY3 or CY5).

PreNest Oligonucleotides (Primary Amplification)Amplification Primers:

[0068]

CYP2D6 Prenest B F 5'-CCA GAA GGC TTT GCA GGC TTC A-3' (Bases 1281-1302, SEQ ID NO:1)

CYP2D6 Prenest B R 5'-ACT GAG CCC TGG GAG GTA GGT A-3' (Bases 6357-6378, SEQ ID NO:2)

Nest Oligonucleotides (Secondary Amplification)

[0069]

CYP2D6 (C100T)

Amplification Primers:

[0070]

- CYP2D6 (100)B f 5'GGC CTA CCC TGG GTA AGG GCC TGG AGC AGG A -3' (Bases 1579-1594, SEQ ID NO:3)
- CYP2D6 (100)B r 5'-/5Bio/ CCT GGT CGA AGC AGT AT-3' (Bases 2394-2411, SEQ ID NO:4)

Probes and Stabilizers:

[0071]

CYP2D6 C100T wild type 5'-/5Cy3/GCA CGC TAC C -3' (Bases 1700-1710, SEQ ID NO:5)

CYP2D6 C100T polymorphism 5'-/5Cy5/GCA CGC TAC T -3' (Bases 1700-1710, SEQ ID NO:6)

CYP2D6 C100T stabilizer 5'CAC CAG GCC CCC TGC CAC TG -3' (Bases 1711-1731, SEQ ID NO:7)

CYP2D6 (C1023T)

Amplification Primers:

[0072]

EP 3 037 549 A1

- **CYP2D6 (1023)B f** 5'-CCT GCT CAC TCC TGG TAG CC -3' (Bases 2394-2413, SEQ ID NO:8)
- **CYP2D6(1023) B r** 5'-/5Bio/ CTG TTT CAT GTC CAC GAC C-3' (Bases 2760-2779, SEQ ID NO:9)

Probes and Stabilizers:

[0073]

CYP2D6 (1023) wild type 5'-/5Cy3/TGC CCA TCA C -3' (Bases 2632-2642, SEQ ID NO:10)

CYP2D6 (1023) polymorphism 5'-/5Cy3/TGC CCA TCA T -3' (Bases 2632-2642, SEQ ID NO:11)

CYP2D6 (1023) stabilizer 5'CCA GAT CCT GGG TTT CGG GCC GCG -3' (Bases 2643-2667, SEQ ID NO:12)

CYP2D6 (G1661C and G1846A)

Note: There are 2 polymorphisms on this amplicon.

Amplification Primers:

[0074]

- **CYP2D6 (1661,1846)B f** 5'- CAG AGG CGC TTC TCC G -3' (Bases 3266-3282, SEQ ID NO:13)
- **CYP2D6 (1661,1846)B r** 5'- /5Bio/CTC GGT CTC TCG CTC CGC AC -3' (Bases 3630-3650, SEQ ID NO:14)

Probes and Stabilizers:

[0075]

CYP2D6(1661) wild type 5'-/5Cy3/CTT CTC CGT G -3' (Bases 3266-3276, SEQ ID NO:15)

CYP2D6(1661) polymorphism 5'-/5Cy3/CTT CTC CGT C -3' (Bases 3266-3276, SEQ ID NO:16)

CYP2D6(1661) stabilizer 5'-TCC ACC TTG CGC AAC TTG GGC CTG GG -3' (Bases 3277-3303, SEQ ID NO:17)

CYP2D6(1846) wild type 5'-/5Cy3/CCA CCC CCA G -3' (Bases 3460-3470, SEQ ID NO:18)

CYP2D6(1846) polymorphism 5'-/5Cy3/CCA CCC CCA A -3' (Bases 3460-3470, SEQ ID NO:19)

CYP2D6(1846) stabilizer 5'-GAC GCC CCT TTC GCC CCA ACG -3' (Bases 3471-3492, SEQ ID NO:20)

CYP2D6 (A2549del and C2850T)

Amplification Primers:

[0076]

CYP2D6 (A2549del and C2850T)B f 5'- TGA GAC TTG TCC AGG TGA AC -3' (Bases 4001-4022, SEQ ID NO:21)

▪ **CYP2D6(A2549del and C2850T)B r** 5'-/5Bio/ CCC AGA TGG GCT CAC GCT GC-3' (Bases 4558-4578, SEQ ID NO:22)

Probes and Stabilizers:

[0077]

CYP2D6 (2549) wild type 5'-/5Cy3/ACT GAG CAC A -3' (Bases 4161-4171, SEQ ID NO:23)

CYP2D6 (2549) polymorphism 5'-/5Cy3/ ACT GAG CAC G-3' (Bases 4161-4171, SEQ ID NO:24)

CYP2D6 (2549) stabilizer 5'-GGA TGA CCT GGG ACC CAG CCC AGC C3' (Bases 4172-4199, SEQ ID NO:25)

CYP2D6 (2850) wild type 5'-/5Cy3/GAG AAC CTG C -3' (Bases 4462-4472, SEQ ID NO:26)

CYP2D6 (2850) polymorphism 5'-/5Cy3/ GAG AAC CTG T-3' (Bases 4462-4472, SEQ ID NO:27)

CYP2D6 (2850) stabilizer 5'-GCA TAG TGG TGG CTG ACC TGT TCT CTG -3' (Bases 4473-4500, SEQ ID NO:28)

PCR Amplification.**A. PreNest (Initial Amplification):****[0078]****CYP2D6 PreNest B**

The initial PreNest PCR for the CYP2D6(C1661G), CYP2D6 (G1846A), CYP2D6(A2549Del), CYP2D6(C100T), CYP2D6(C2850T), and CYP2D6 (C1023T) polymorphisms was performed with the following reagents per sample: 0.6 μ L 25 μ M PreNest B F, 0.6 μ L 25 μ M PreNestB R, 2.5 μ L 10 mM dNTP's, 39.55 μ L dH₂O, 5.0 μ L Expand Buffer 3 (Roche), 0.75 μ L Expand Enzyme Mix (Roche), and 1.0 μ L genomic DNA in a total volume of 50 μ L. PCR was performed by holding at 94°C for 2 minutes then 30 cycles of denaturation at 94°C for 30 seconds, annealing at 67°C for 30 seconds, and extension at 68°C for 7 minutes, followed by a final extension at 72°C for 10 minutes. The expected PCR product was $\approx$ 4.8 - 5Kb.

[0079] All subsequent PCR's were performed using the PreNest B product as the template for the amplification.

CYP2D6(100) B

[0080] The (C100T) nested PCR was performed using 1 μ L of the initial PreNest B amplicon as template. The product of this PCR was used to detect the CYP2D6(C100T) mutation and uses the following reagents (25 μ L total volume): 1.0 μ L 25 μ M CYP2D6(100) B f, 1.0 μ L 25 μ M CYP2D6(100) B r, 12.5 μ L Qiagen Hot Start Master Mix, and 9.5 μ L dH₂O. PCR was performed by holding at 95°C for 15 minutes, then 30 cycles of denaturation at 95°C for 30 sec, annealing at 57°C for 30 sec, and extension at 72°C for 1 min, followed by a final extension at 72°C for 7 min.

CYP2D6(1023) B

[0081] A (C1023T) nested PCR was performed using 1 μ L of PreNest B product as template. The product of this PCR was used to detect the CYP2D6 (C1023T) mutation and uses the following reagents (25 μ L total volume): 1.0 μ L 25 μ M CYP2D6 (1023) B f, 1.0 μ L 25 μ M CYP2D6 (1023)B r, 12.5 μ L Amplitaq Gold Master Mix, and 9.5 μ L dH₂O. PCR was performed by holding at 95°C for 10 min, then 30 cycles of denaturation at 95°C for 30 sec, annealing at 59°C for 30 sec, and extension at 72°C for 1 min, followed by a final extension at 72°C for 7 min.

CYP2D6(C1661G and G1846A) B

[0082] The (C1661 G and G1846A) nested PCR was performed using 1 μ L of PreNest B product as template. The product of this PCR was used to detect the CYP2D6 (C1661G) and CYP2D6 (G1846A) mutations and uses the following reagents (25 μ L total volume): 1.0 μ L 25 μ M CYP2D6 (1661,1846)B f, 1.0 μ L 25 μ M CYP2D6 (1661,1846)B r, 12.5 μ L 2X Amplitaq Gold Master Mix, and 9.5 μ L dH₂O. PCR was performed by holding at 95°C for 10 min, then 30 cycles of denaturation at 95°C for 30 sec, annealing at 63°C for 30 sec, and extension at 72°C for 1 min, followed by a final extension at 72°C for 7 min.

CYP2D6 (A2549Del and CYP2D6C2850T) B

[0083] The (A2549Del and C2850T) nested PCR was performed using 1 μ L of PreNest B product as template. The product of this PCR was used to detect the CYP2D6 (A2549Del) and CYP2D6 (C2850T) mutations and uses the following reagents (25 μ L total volume): 1.0 μ L 25 μ M CYP2D6 (2549,2850)B f, 1.0 μ L 25 μ M CYP2D6 (2549,2850)B r, 12.5 μ L 2X Amplitaq Gold Master Mix, and 9.5 μ L dH₂O. PCR was performed by holding at 95°C for 10 min, then 30 cycles of

denaturation at 95°C for 30 sec, annealing at 57°C for 30 sec, and extension at 72°C for 1 min, followed by a final extension at 72°C for 10 min.

[0084] Polymorphisms were detected using the Nanogen Molecular Biology Workstation according to the Nanogens User's manual. Samples were desalted on Millipore desalting plates. In a 1.5 ml centrifuge tube, the following reagents were added: 2.0 µl of wild type probe, 2.0 µl of polymorphism probe, 2.0 µl of stabilizer, and 44 µl high salt buffer. Each assay had probes and stabilizers that were labeled and specific to it. The reagents were vortexed, spun briefly, and dispensed over the Nanochip array for 5 minutes and placed in a dark container. The chips were read with a trending method from 24° C to 50° C, with readings every two degrees. It was confirmed that the fluorescent probe hybridized at 24°C and that all probes had de-hybridized at 50°C. If the relative fluorescence units (RFUs) were above 80 on any chip at 50°C, the chips were stripped in 200 µl of 1.0 M NaOH for 10 minutes and washed three times with dH₂O and once with high salt buffer. A heterozygote control (known heterozygote, one to "normalize" or scale the RFUs and the other to confirm the correcting trending pattern observed in previously genotyped heterozygotes) and a negative control (no DNA) were included for each polymorphism. For background subtraction, L-histidine was used.

[0085] Tables 3-5 summarize the total number of participants for different age groups and the CYP2D6 inferred metabolic phenotype detected using the above procedure. Table 6 provides the particular 2D6 haplotypes that were observed in the patients that had antidepressant medication resistant subtypes of depression.

Table 3

Total Number of Participants in Each Grouping (All Ages; N = 86)

CYP2D6 Inferred Metabolic Phenotype

Response to Psychotropic		Poor	Intermediate	Extensive	Ultra-Rapid
	Non-Response	5	16	6	0
	Side-Effect	5	16	7	0
	Both	8	17	5	1
	Total	18	49	18	1

Table 4

Total Number of Participants in Each Grouping (Ages: <19; N = 12)

CYP2D6 Inferred Metabolic Phenotype

Response to Psychotropic Medication		Poor	Intermediate	Extensive	Ultra-Rapid
	Non-Response	0	3	0	0
	Side-Effect	3	3	0	0
	Both	1	1	0	1
	Total	4	7	0	1

Table 5

Total Number of Participants in Each Grouping (Ages: >18; N = 74)

CYP2D6 Inferred Metabolic Phenotype

Response to Psychotropic Medication		Poor	Intermediate	Extensive	Ultra-Rapid
	Non-Response	5	13	6	0
	Side-Effect	2	13	7	0
	Both	7	16	5	0
	Total	14	42	18	0

Table 6

Antidepressant Medication Resistant Subtypes of Depression¹

Sub#	2D6 Haplo	2D6 Enz Act
100MR	*1/*2P Hz*2	I
101MR	Hz *4	I
102MR	*1/*10 H2*10	I
103MR	Hz *4	I
104MR	*2/*2P Homo *2	I-P
105MR	Hz *11	I
106MR	Hz *4	I
107MR	Homo *3	None
108MR	Hz *4	I
109MR	*2/*2 Homo *2	I-P
110MR	Hz *2	I
111MR	Hz *2	I
112MR	WT	N
113MR	Hz *2	I
114MR	WT	N
115MR	WT	N
116MR	*3/*4 Compd Hz *3/*4	P
117MR	Hz *4	I
118MR	Hz *2	I
119MR	Compound Hz *2/*4	P
120MR	Hz *4	I
121MR	Hz *4	I
122MR	Hz *4	I

EP 3 037 549 A1

(continued)

<i>Antidepressant Medication Resistant Subtypes of Depression¹</i>		
Sub#	2D6 Haplo	2D6 Enz Act
123MR	WT	N
124MR	Hz *12	I
125MR	Hz *4	I
126MR	Hz *4	I
128MR	WT	N
129MR	Hz *4	I
130MR	Hz *1E	N
131MR	Homo *4	P
132MR	Hz *4	I
133MR	Hz *2	I
134MR	Compound Hz *2/*3	P
135MR	Compound Hz *3/*4	P
136MR	Hz *	I
137MR	Hz *2	I
138MR	Compound Hz *2/*4	P
139MR	Compound Hz *2/*5	P
141MR	Homo *5	None
142MR	Compound Hz *4/*15	P
143MR	Hz *4	I
144MR	Hz *17 or *34	I
145MR	Hz *3	I
146MR	Compound Hz *4/*5	P
147MR	WT	N
148MR	Hz *2	I
149MR	WT	N
150MR	Hz *2	I
155MR	WT	N
156MR	Hz *4	I
157MR	Hz *2	I
158MR	Hz *4	I
159MR	Hz *2	I
160MR	WT	N
161MR	Hz *2	I
162MR	WT	N
164MR	Hz *2	I
165MR	Hz *2	I
166MR	WT	N

(continued)

Antidepressant Medication Resistant Subtypes of Depression¹		
Sub#	2D6 Haplo	2D6 Enz Act
167MR	Compound Hz *2/*4	P
168MR	Compound Hz *2/*4	P
169MR	WT	N
170MR	Hz *2	I
171MR	Compound Hz *2/*3	P
172MR	Hz *2 w/ duplication	Inc. act.
173MR	Hz *2	I
174MR	Hz *4 w/ balanced duplication	N
175MR	Hz *2	I
176MR	Hz *4	I
177MR	Hz *2	I
179MR	Homo *2	I-P
180MR	Homo *2	I-P
181MR	WT	N
182MR	Hz *4	I
183MR	WT	N
184MR	Hz *2	I
185MR	WT	N
187MR	Compound Hz *2/*4	P
188MR	Hz *4	I
189MR	Hz *10	I
190MR	WT	N
191MR	Hz *2	I
192MR	Hz *4	I
193MR	Hz *2	I
194MR	Hz *3	I
195MR	Hz *17 or *34	I
196MR	Hz *4	I
197MR	Compound Hz *2/*4	P
198MR	Hz *2	I
199MR	Compound Hz *2/*4	P
¹ P=poor; N=normal; I=intermediate; HI = Higher inducibility; WT = wild type		

[0086] All of these adolescents were found to have atypical 2D6 haplotypes (see Tables 3-5). The initial results of our analysis of the cytochrome P450 gene alleles in an adult sample also demonstrated a high degree of variability in the polymorphisms of the 2D6 gene (see Tables 3-5). Adolescents with polymorphisms associated with poor 2D6 metabolism were noted to have clinical histories marked by either a poor response to the medication or by a high frequency of side effects when they had been treated with the selective serotonin reuptake inhibitors that are metabolized by the 2D6 enzyme.

Example 2

Determining Genotype of the Dopamine Transporter, Dopamine Receptors, Tryptophan Hydroxylase, Serotonin Receptors, and COMT genes

5 **[0087]** The following probes and primers of Table 7 can be generated to detect the genotype of the dopamine transporter (DAT1, SLC6A3), dopamine receptors (DRD1, DRD2, DRD3, DRD4, and DRD5), tryptophan hydroxylase (TPH), serotonin transporter (5-HTT), serotonin receptors (HTR1A, HTR1B, HTR1D, HTR2A, and HTR2C), and COMT genes.

10

15

20

25

30

35

40

45

50

55

Symbol	Polymorphism	Forward Primer	Reverse Primer	Wt. Probe	Variant Probe	Stabilizer	SEQ ID
DAT1, SLC6A3	>G710A, Q237R	CCCCGGCTGAC TCCCTCTG	GACCCCGAGCCTCACCT TCC	AGCTGCCACC	AGCTGCCACT	GCGGAGGCCCCAG GT	29-33
	>C124T, L42F	ACCAAGAGGG AAGAAGCACA G	CCAATGACGGACAGGAG AAA	GAGCTGGTGAG	GAGCTGGTGAA	CTGCACTCCGTTT TGCTCCT	34-38
	40 bp VNTR	GATGGGGTC CTGGTATGCT	CTGGAGTCACTGGCTCAA G				39-40
	>T595G, S199A	GTGTGGCATGG ACCTTGTCTG	GCAATGCCGCCGTATTGT TTC	AAAGCTTATTA CAGA	AGCTTATTACAG C	GGATGAGATGGCA TATGTCTGC	41-45
DRD1	>G150T, R50S	CCTGTTTCCTG TCGCTGCTC	CCAATACCTGTCCACGCT GAT	GGTGTGCGAAC	GGTGTGCGAAA	CTGATAACGGCAG CACAGACC	46-50
	>C110G, T37R	GGGCTGGTGGT GGAGAG	TCTGACACAGCCAAAGGAG ATG	CCAGGAGCG	CCCAGGAGCC	TGGACAGGATGAG CAGCGA	51-55
	>A109C, T37P	GGGCTGGTGGT GGAGAG	TCTGACACAGCCAAAGGAG ATG	CCAGGAGCGT	CCAGGAGCGG	GGACAGGATGAGC AGCGACA	56-60
	>A1051G, T351A	TCTCCACAGCA CTCCCGACAG	CCGAGAACAAATGGCGAG CATC	GGTCCGGGT	GGTCCGGGC	TTTGCCATTGGGC ATGGTCTG	61-65
DRD2	>C932G, S311C	CACCAGCCCA CCCGAGAG	ACCGAGAACAAATGGCGA GCAT	CCATGGTGGG	CCATGGTGGC	ACGGTCCGGGA GAGTC	66-70
	>C928T, P310S	CCGGTACAGC CCCATCCC	GCCGACTCACCGAGAAC AATG	GTGGGACCG	GGTGGGACGA	GTCGGGGAGAGTC AGCTGG	71-75
	>G460A, V154I	GGGGACAGGA GGGAGGGTTG AG	GAAAGGAGTGGGACAGG AGATGGTG	GGAGATCATGA C	TGGAGATCATGA T	GGTGACCCGGGCGC TTG	76-80
	>A25G, S9G	AGCAACCAAG CCCCAAAGAG	GAGCGCGCAGTAGGAGA GG	TTCAGGTGGCT	TCAGGTGGCC	ACTCAGCTGGCTC AGAGATGC	81-85

(continued)

Symbol	Polymorphism	Forward Primer	Reverse Primer	Wt. Probe	Variant Probe	Stabilizer	SEQ ID
DRD4	>T581G, V194G	CGCTACAACC GGCAGGTGG G	GCAGCATGAGCGGGCAG GGTA	GGACGAGTAGA	GGACGAGTAGC	CCACGTAGTCGG GTCCTC	86-90
	>C841G, P281A	CGCCTTCCCC ACGCCACC	GTGGAGCCCGAGGGGT CCG	GGCGCGGG	GGCGCGGC	GGCGGCACAGTCG GG	91-95
DRD5	>A889C, T297P	TGCGGTTTTC ATCAAGAAG	GGAGTTGGCCCGCAGAG A	CCGACAGGGT	CGACAGGGG	CTTGAGAACCTTG GTCTCCTTCTTGAT	96-100
	>G1252A, V418I	CAACGCCGAC TTCGGAA	ACATGCGATCGAAAGGA CTCT	CGGGGGGAAC	CGGGGGGAAT	GGCGTTGGGCATC ATGTGG	101-105
	>G181A, V61M	GCAGTCCAGC CCGAAATG	GCACGATGGCTGCCGA	CTGCGGACAC	CTGCGGACAT	CAGCACGTTGCC AG	106-110
	>G185C, C62S	CAACGTGCTG GTG	CATCTTGGCCTTGTA	GGCTGCGC	TGGCTGCGG	ACAACGACACGTT GCCC	111-115
	>T263G, R88L	CGCCAAGATG ACCAAC	CGGAATGAAGGAGATGA	GCCACGAAGA	GCCACGAAGC	GGTCTGACACAGC CAGAGACA	116-120
	>G1354A, W455-	GCAGCTGCCT ACATCCACAT	CATTCGGGGTGAAAGGT GTTA	GTCCAGCTCC	GTCCAGCTCT	CAGACAGACTCTG CAACAGGGTC	121-125

(continued)

Symbol	Polymorphism	Forward Primer	Reverse Primer	Wt. Probe	Variant Probe	Stabilizer	SEQ ID
HTR1A	>G815A, G272D	GGAGAGTCGG GGAGCAGGAA C	GAGCGGGGGGCACAAGG	GCACAGAGCAC	GCACAGAGCAT	CCCCAGCCTTGCT CTCCA	126-130
	>G656T, R219L	CTACATCCCGC TGCTGCTCAT	CCTGCTCCCCGACTCTCC A	GCGCAGCTC	GCGCAGCTA	GGAAATATGCGCCC ATAGAGAACCA	131-135
	>C548T, P551L	GCTCACTTGGC TTATGGCTTC CTC	CATGAGCAGCAGCGGGA TGT	CATGCGTCGG	CATGCGTCGA	GGTCCGAGCGGTC TTCCG	136-140
	>A82G, I28V	AGGGCAACAA CACCACATCA CCAC	GCAGCAACCAACCGAC ACCAT	ACGTCGGAGAT	CGTCGGAGAC	ACCAGTAGTGTG CCGCCG	141-145
HTR2A	>G64A, G22S	TCAGGGCAAC AACACCACAT	GCACCAACACCGACACC A	GTTGCCGCC	GTRGCCGCT	GGTCTCAAAGGA GCCGT	146-150
	>C47T, P16L	TTTCTTCCTT CCCCCTTCC	CCACCACGCACGCATTG	AAGGGAGCCG	AAAGGGAGCCA	GTGGTGATGTGGT GTTGTTGCC	151-155
HTR2A	T102C	GCTCAACTAC GAACTCCTTA A	TCACTACGGCTGTCAGTA AAG	TTAGCTTCTCCA	TTAGCTTCTCCG	GAGTTAAAGTCAT TACTGTAGAGCCT GGT	156-160
HTR1D	>C794T, S265L	GCTCTGCTCGC TCAAC	GATTTAGTGGCTTTCCT TTC	CCGAGTGCG	GCCGAGTGCA	AGTGCCCTCATG GAGGC	161-165
HTR2C	>C10G, L4V	CTCAATTTTAA ACTTTGGTTGC TTA	TTTGAATAGGAAACACC CATAA	GCATTCTCAG	GCATTCTCAC	GTTACCATGATT GCTTCAGTCTTAA GC	166-170
CYP1A2*1F	>G68C, C23S	ATGATGACAA TGATGCTGATG ATGATAATG	TCCACATCGGAGGTATT GAAAATG	TCACAGAAATA TCAC	CACAGAAATATC AG	ATTGCCAAACCAA TAGGCCAATTAGG	171-175
	-164C>A	GGCTTCAGTTT CCCGATCTG	TAGGCAAGGCAGGCAC TAC	AAGGAGCTGG	GAAGGAGCTGA	GTACATGGGGGCC CCC	176-180

Example 3

[0088] The genotype of the CYP2D6, CYP2C19, and 5HTTTR genes was determined in 97 patients using the methods described above. As indicated in Example 1, CYP2D6 *2, *3, *4, *10, *17, and *5 del alleles, CYP2C19 *2 (A,B), *3, *4, *5 (A,B), *6, *7, and *8 alleles, and the short/long form of the 5HTTTR gene were assessed.

[0089] Each patient that was genotyped exhibited either a poor response to at least one antidepressant medication or an intolerable side effect to at least one antidepressant medication. Based on the genotype information, 18 medication profiles were generated (Table 8). In Table 8, PM is poor metabolizer; IM is intermediate metabolizer; EM is extensive metabolizer; s is the short form of the gene; and l is the long form of the gene. Each medication profile provides antidepressant medications that are acceptable for use, antidepressant medications to be used with caution, and antidepressant medications to be avoided or used with caution and close monitoring.

TABLE 8

Medication Algorithm Microarray Results from Pharmacological Study				
Key Gene Genotype Category	2D6	2C19	5HTTTR	Genotype frequencies in pharmacological study
Med Profile 1	PM	PM	s/s	2
Med Profile 2	PM	PM	s/l	2
Med Profile 3	PM	PM	l/l	1
Med Profile 4	PM	EM	s/s	1
Med Profile 5	PM	EM	s/l	6
Med Profile 6	PM	EM	l/l	1
Med Profile 7	IM	PM	s/s	1
Med Profile 8	IM	PM	s/l	7
Med Profile 9	IM	PM	l/l	3
Med Profile 10	EM	PM	s/s	0
Med Profile 11	EM	PM	s/l	6
Med Profile 12	EM	PM	l/l	3
Med Profile 13	IM	EM	s/s	2
Med Profile 14	IM	EM	s/l	17
Med Profile 15	IM	EM	l/l	16
Med Profile 16	EM	EM	s/s	7
Med Profile 17	EM	EM	s/l	10
Med Profile 18	EM	EM	l/l	12

[0090] Medication profile 1 is associated with 2D6 and 2C19 poor metabolizers and the s/s form of the 5HTTTR gene. In patients with medication profile 1, bupropion is acceptable for use, mirtazapine and fluvoxamine can be used with caution, while citalopram, escitalopram, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, nortriptyline, and sertraline should be avoided or used with caution and close monitoring.

[0091] Medication profile 2 is associated with 2D6 and 2C19 poor metabolizers and the s/l form of the 5HTTTR gene. In patients with medication profile 2, fluvoxamine and bupropion are acceptable for use, sertraline and mirtazapine can be used with caution, and citalopram, escitalopram, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and close monitoring.

[0092] Medication profile 3 is associated with 2D6 and 2C19 poor metabolizers and the l/l form of the 5HTTTR gene. In patients with medication profile 3, fluvoxamine and bupropion are acceptable for use, sertraline and mirtazapine can be used with caution, and citalopram, escitalopram, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and close monitoring.

[0093] Medication profile 4 is associated with 2D6 poor metabolizers, 2C 19 extensive metabolizers, and the s/s form

of the 5HTTTR gene. In patients with medication profile 4, bupropion is acceptable for use, mirtazapine, citalopram, fluvoxamine, and escitalopram can be used with caution, and paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, nortriptyline, and sertraline should be avoided or used with caution and close monitoring.

[0094] Medication profile 5 is associated with 2D6 poor metabolizers, 2C19 extensive metabolizers, and the s/s form of the 5HTTTR gene. In patients with medication profile 5, citalopram, fluvoxamine, bupropion and escitalopram are acceptable for use, sertraline and mirtazapine can be used with caution, and paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and close monitoring.

[0095] Medication profile 6 is associated with 2D6 poor metabolizers, 2C19 extensive metabolizers, and the 1/1 form of the 5HTTTR gene. In patients with medication profile 6, citalopram, fluvoxamine, bupropion and escitalopram are acceptable for use, sertraline and mirtazapine can be used with caution, and paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and close monitoring.

[0096] Medication profile 7 is associated with 2D6 intermediate metabolizers, 2C19 poor metabolizers, and the s/s form of the 5HTTTR gene. In patients with medication profile 7, bupropion is acceptable for use, mirtazapine, venlafaxine, amitriptyline, nortriptyline, fluvoxamine, and sertraline can be used with caution, and citalopram, escitalopram, fluoxetine, imipramine, and paroxetine should be avoided or used with caution and close monitoring.

[0097] Medication profile 8 is associated with 2D6 intermediate metabolizers, 2C19 poor metabolizers, and the s/l form of the 5HTTTR gene. In patients with medication profile 8, fluvoxamine, bupropion, and sertraline are acceptable for use, mirtazapine, paroxetine, venlafaxine, amitriptyline, and nortriptyline can be used with caution, and citalopram, escitalopram, fluoxetine, and imipramine should be avoided or used with caution and close monitoring.

[0098] Medication profile 9 is associated with 2D6 intermediate metabolizers, 2C19 poor metabolizers, and the 1/1 form of the 5HTTTR gene. In patients with medication profile 9, fluvoxamine, bupropion, and sertraline are acceptable for use, mirtazapine, paroxetine, venlafaxine, amitriptyline, and nortriptyline can be used with caution, and citalopram, escitalopram, fluoxetine, and imipramine should be avoided or used with caution and close monitoring.

[0099] Medication profile 10 is associated with 2D6 extensive metabolizers, 2C19 poor metabolizers, and the s/s form of the 5HTTTR gene. In patients with medication profile 10, bupropion and venlafaxine are acceptable for use, mirtazapine, nortriptyline, imipramine, amitriptyline, sertraline, paroxetine, and fluoxetine can be used with caution, and fluvoxamine, citalopram, and escitalopram should be avoided or used with caution and close monitoring.

[0100] Medication profile 11 is associated with 2D6 extensive metabolizers, 2C19 poor metabolizers, and the s/l form of the 5HTTTR gene. In patients with medication profile 11, sertraline, venlafaxine, paroxetine, bupropion, and fluoxetine are acceptable for use, and mirtazapine, fluvoxamine, nortriptyline, citalopram, escitalopram, imipramine, and amitriptyline can be used with caution.

[0101] Medication profile 12 is associated with 2D6 extensive metabolizers, 2C19 poor metabolizers, and the 1/1 form of the 5HTTTR gene. In patients with medication profile 12, sertraline, venlafaxine, paroxetine, bupropion, and fluoxetine are acceptable for use, and mirtazapine, fluvoxamine, nortriptyline, citalopram, escitalopram, imipramine, and amitriptyline can be used with caution.

[0102] Medication profile 13 is associated with 2D6 intermediate metabolizers, 2C19 extensive metabolizers, and the s/s form of the 5HTTTR gene. In patients with medication profile 13, bupropion and mirtazapine are acceptable for use, venlafaxine, amitriptyline, imipramine, nortriptyline, citalopram, fluvoxamine, escitalopram, and sertraline can be used with caution, and paroxetine and fluoxetine should be avoided or used with caution and close monitoring.

[0103] Medication profile 14 is associated with 2D6 intermediate metabolizers, 2C19 extensive metabolizers, and the s/l form of the 5HTTTR gene. In patients with medication profile 14, citalopram, fluvoxamine, bupropion, escitalopram, sertraline, and mirtazapine are acceptable for use, and paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline can be used with caution.

[0104] Medication profile 15 is associated with 2D6 intermediate metabolizers, 2C19 extensive metabolizers, and the 1/1 form of the 5HTTTR gene. In patients with medication profile 15, citalopram, fluvoxamine, bupropion, escitalopram, sertraline, and mirtazapine are acceptable for use, and paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline can be used with caution.

[0105] Medication profile 16 is associated with 2D6 extensive metabolizers, 2C19 extensive metabolizers, and the s/s form of the 5HTTTR gene. In patients with medication profile 16, bupropion, mirtazapine, venlafaxine, amitriptyline, imipramine, and nortriptyline are acceptable for use, and paroxetine, fluoxetine, fluvoxamine, sertraline, escitalopram, and citalopram can be used with caution.

[0106] Medication profile 17 is associated with 2D6 extensive metabolizers, 2C19 extensive metabolizers, and the s/l form of the 5HTTTR gene. Medication profile 18 is associated with 2D6 extensive metabolizers, 2C19 extensive metabolizers, and the 1/1 form of the 5HTTTR gene. In patients with medication profiles 17 or 18, citalopram, fluvoxamine, bupropion, escitalopram, sertraline, mirtazapine, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline are acceptable for use.

Example 4

[0107] The genotype of the CYP2D6, CYP2C19, CYP3A4, CYP1A2, and HTR2A genes was determined in ten patients using the methods described above. The genotype of the 5HTTR gene was determined by amplifying the polymorphic region of the gene using primers flanking the region. Reaction products were sized using electrophoresis (Agilent Technologies). CYP3A4 *1B, *2, *5, *6, *12, *13, *15A, *17, and *18A alleles, CYP1A2 *1 F allele, and the HTR2A 102 polymorphism were detected

[0108] Based on the CYP2D6 genotype, patients were classified as being poor, intermediate, extensive, or ultra rapid metabolizers. For each of the CYP2C19, CYP3A4, and CYP1A2 genotypes, patients were classified as being either extensive or not extensive (i.e., poor) metabolizers. For the 5HTTR genotype, patients were classified as having the short/short (s/s) form of the gene, the short/long (s/l) form of the gene, or the long/long (l/l) form of the gene. For the HR2A genotype, patients were classified as having either the C/C allele or as having the T/T or T/C alleles.

[0109] The following six rules were used to sort the genotypes into 192 treatment recommendations. The first rule relates to the CYP2D6 genotype. For poor metabolizers and ultra rapid metabolizers, CYP2D6 substrates were placed in the "use very cautiously with close monitoring" category. For intermediate metabolizers, exclusive CYP2D6 substrates were placed in the "avoid or use very cautiously with close monitoring" category. For intermediate metabolizers, substrates that are partially metabolized by CYP2D6 were placed in the "use cautiously" category. For extensive metabolizers, CYP2D6 substrates were placed in the "acceptable for use" category.

[0110] After applying the first rule, the second rule relating to the 2C19 genotype was applied. For poor metabolizers based on their CYP2C 19 genotype, exclusive CYP2C 19 substrates were placed in the "avoid or use very cautiously with close monitoring" category. For extensive or intermediate metabolizers based on their CYP2C19 genotype, CYP2C19 substrates were placed in the "acceptable for use" category. Substrates that are primarily metabolized by both CYP2D6 and CYP2C19 were placed in the "avoid or use very cautiously with close monitoring" category if CYP2C19 was poor and CYP2D6 was either intermediate or poor, or placed in the "use cautiously" category if CYP2C 19 was poor and CYP2D6 was extensive.

[0111] A third rule relating to the 3A4 genotype was applied after the first and second rules were applied. For poor metabolizers based on their CYP3A4 genotype, exclusive CYP3A4 substrates were placed in the "avoid or use very cautiously with close monitoring" category. For extensive or intermediate metabolizers based on their CYP3A4 genotype, CYP3A4 substrates were placed in the "acceptable for use" category. Substrates that are primarily metabolized by any two of the following three enzymes: CYP2D6, CYP2C 19, or CYP3A4, were placed in the "avoid or use very cautiously with close monitoring" category if any two of the three were intermediate or poor metabolizers or placed in the "use cautiously" category if one of the two most primary enzymes was a poor metabolizer.

[0112] The fourth rule relating to the 1A2 genotype was applied after applying the first three rules. For poor metabolizers based on CYP1A2 genotype, CYP1A2 substrates were placed in the "avoid or use very cautiously with close monitoring" category (include fluvoxamine as a primary CYP1A2 substrate). For poor metabolizers based on the CYP1A2 genotype, substrates having the CYP1A2 pathway as one of their primary pathways were placed in the "use with caution" category.

[0113] After applying the first four rules, the fifth rule relating to the 5HTTR genotype was applied. For the homozygous short genotype (s/s) of the serotonin transporter gene, SSRI's that were categorized as "acceptable for use" were shifted to the "use with caution" category. For the homozygous short genotype (s/s) of the serotonin transporter gene, SSRI's that were categorized as "use with caution" were shifted to the "avoid or use very cautiously with close monitoring" category. Antidepressants, outside the SSRI class, do not shift category based on 5HTTR genotype.

[0114] Rule six, which relates to the HR2A genotype, was applied after applying the first five rules. For the "mutant form" of the 5-hydroxytryptamine serotonin receptor 2A gene, if paroxetine was in the "acceptable for use" category, it was shifted to the "use with caution" category. If paroxetine was in the "use with caution" category, it was shifted to the "avoid or use very cautiously with close monitoring."

[0115] Based on the genotype of the patients and associated medication profile, medications appropriate for the patient were selected. Table 9 summarizes the data for the 10 patients. Case studies of four of the seven patients are presented below.

[0116] The first case is a 16-year-old white female who was genotyped and found to be a poor metabolizer for CYP2D6 (*3/*9) as well as being a poor metabolizer for CYP2C19 (*2/*2). She was diagnosed at 6 years of age as having ADHD and treated unsuccessfully. She subsequently developed symptoms of depression and mood dysregulation and was placed on many medications that were metabolized by CYP2D6, such as EFFEXOR. Additionally, she was homozygous for a short form of the serotonin transporter gene. Based on her specific genotype of these key genes, the algorithm demonstrates that there are a relatively small number of potential strategies for controlling her depression. The medication profile generated based on her genotype of HR2A TT/TC, s/s form of 5HTTR, extensive metabolizer for 1A2 and 3A4, and poor metabolizer for 2C19 and 2D6 indicated that bupropion was acceptable for use, mirtazapine and fluvoxamine should be used with caution, and citalopram, escitalopram, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, nortriptyline, and sertraline should be avoided or used with caution and monitoring.

[0117] Before genotyping, she had a particularly unusual side effect to citalopram in that she developed hallucinations (incidence less than 1%). This atypical and dangerous side effect could have been avoided if she had been genotyped prior to receiving this medication. The metabolic enzyme that is primarily involved in the metabolism of citalopram is CYP2C19 and a secondary metabolic pathway is CYP2D6. Both of these pathways were nonfunctional in this patient.

Therefore, even on a modest dose of citalopram, this patient developed acute auditory and visual hallucinations, which were subsequently treated with an atypical antipsychotic medication. Additionally, based on an inability to develop an effective treatment plan for this patient, she required three acute hospitalizations and a partial hospitalization experience prior to being referred to a state hospital for two months. She was unsuccessfully tried on more than a dozen psychotropic medications. There is no question that her care could have been greatly improved and perhaps some of her hospitalizations could have been avoided if clinicians had been aware of her inability to use many of the drugs that had been prescribed at their standard doses. Additionally, her serotonin transporter genotype (C/C) would have suggested a poor response to many of the medications that had been prescribed. Subsequent to genotyping, she was placed on a combination of two mood stabilizers that are metabolized by an alternative pathway (LAMICTAL (lamotrigine) and TRILEPTAL (oxcarbazepine)). She is also tolerating treatment with ABILIFY (aripiprazole), which in addition to being metabolized by CYP2D6 is metabolized by CYP3A4, for which she has an internal metabolic pathway. She is now doing reasonably well.

[0118] The second case is a 42-year-old Caucasian female with a persistent depressive illness and pervasive anxiety and a long-standing history of psychotropic medication intolerance. This patient had a comprehensive neurological evaluation, which failed to provide an organic explanation for her symptoms. This patient has been treated with at least nine psychotropic medications in the past. Striking examples of CYP2D6 intolerance include the development of acute hallucinations while taking paroxetine and a very low tolerance to amitriptyline. This patient had poor CYP2D6 metabolism and intermediate CYP2C19 metabolism. This represents a rare genotype that would suggest difficulties in tolerating treatment with citalopram. Indeed, she did have irritability and lethargy when taking citalopram. To compound this problem, she also was homozygous for the short form of the serotonin transporter, which is associated with a poor response to serotonin receptive inhibitors. She had experienced multiple outpatient visits for her psychiatric problems. Awareness of her genotype earlier in her history would clearly have allowed her clinician to avoid a number of medications that she could not tolerate. The medication profile generated based on her genotype of HR2A TT/TC, s/s form of 5HTTR, extensive metabolizer for 1A2 and 3A4, and poor metabolizer for 2C19 and 2D6 indicated that bupropion was acceptable for use, mirtazapine and fluoxetine should be used with caution, and citalopram, escitalopram, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, nortriptyline, and sertraline should be avoided or used with caution and monitoring.

[0119] The third case history is a 67-year-old woman with a diagnosis of major depressive disorder, who had been treated with seven different medications in the past. These had not been effective, and she had required two psychiatric hospitalizations. It was determined that she was a poor metabolizer for both CYP2D6 (*4/*9) and CYP2C19 (*2/*2). Additionally, she was homozygous for the long form of the serotonin transporter gene. Based on her genotype of HR2A C/C, I/I form of 5HTTR, extensive metabolizer for 1A2 and 3A4, and poor metabolizer for 2C 19 and 2D6, the associated medication profile indicated that fluoxetine and bupropion were acceptable for use, sertraline and mirtazapine should be used with caution, and citalopram, escitalopram, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and monitoring. If her genotype had been known, it would have been possible to anticipate that she would have had side effects to both EFFEXOR (Venlafaxine) and LEXAPRO (escitalopram oxalate). This could have been predicted, given that two of the three metabolic pathways involved in the metabolism of LEXAPRO were completely inactive. Subsequent to her genotyping, she was successfully treated with electroconvulsive therapy (ECT) and concurrently successfully treated with REMERON (mirtazapine), trazodone, and KLONOPIN (clonazepam), which have alternative pathways of metabolism to CYP2D6 and CYP2C19.

[0120] The fourth case is a 50-year-old Caucasian woman who had been diagnosed as having dysthymia and chronic anxiety. She had been placed on multiple psychotropic drugs, including two CYP2D6 medications that she could not tolerate. Her genotype indicated that she was a poor CYP2D6 metabolizer (*4/*4) and an intermediate CYP2C19 metabolizer (*1/*2). She was heterozygous for the long and short form of the serotonin transporter gene. Based on her genotype of HR2A C/C, I/s form of 5HTTR, extensive metabolizer for 1A2 and 3A4, and poor metabolizer for 2C19 and 2D6, the associated medication profile indicated that fluoxetine and bupropion were acceptable for use, sertraline and mirtazapine should be used with caution, and citalopram, escitalopram, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and monitoring.

[0121] This patient had multiple hospitalizations at considerable cost. She also found it difficult to tolerate a range of nonpsychiatric medications, including being able to take only one dose of PROZAC (fluoxetine) and ZOLOFT (sertraline) before finding that she was acutely uncomfortable. Of particular interest, she ultimately responded to a subclinical dose of paroxetine, which represents an example of using the algorithm to find a method to administer an effective medication that in normal doses could not have been tolerated.

TABLE 9

Sub. No	Age	Gender	Race	Diagnosis	2D6	2C19	3A4	1A2	5HTTR	HTR2A
1	16	F	Cauc.	ADHD; adjustment disorder NOS; Depression NOS; Psychosis NOS; Bipolar Disorder	3/9 Poor Freq: *3 = 1% *9 = 3%	2/2 Poor Freq: *2 = 14%	*1*1	*1/*1F	s/s	TC
2	42	F	Cauc.	Generalized anxiety disorder; Depressive Disorder NOS; Panic disorder	4/4 Poor Freq: *4=18%	1/2 Poor Freq: *1=86% *2=14%	*1*1	*1/*1F	s/s	TC
3	67	F	Other	Major depressive disorder	4/9 Poor Freq: *9=3%	2/2 Poor Freq: *2=14%	*1*1	*1/*1	1/1	CC
4	50	F	Cauc.	Panic disorder; generalized anxiety disorder; dysthymia, somatogorm disorder NOS	4/4 Poor Freq: *4=18%	1/2 Int. Freq: *1=86% *2=14%	*1*1	*1/*1F	s/l	CC
5	39	F	Cauc.	Generalized anxiety disorder	3/4 Poor Freq: *3=33% *4=18%	1/2 Poor Freq: *1=86% *2=14%	*1*1	*1/*1F	l/l	CC
6	12	F	Hispanic	Bipolar disorder NOS	1/1 ext. freq. *1=37%	1/2 Int. Freq: *1=86% *2=14%	*1/*1B	*1/*1	S/1	CC
7	54	M		Bipolar disorder	2P/2P ext. freq: *2P=unk.	1/1 Ext. Freq: *1=86%	*1*1	*1/*1F	s/l	TC

												HTR2A
Sub. No	Age	Gender	Race	Diagnosis	2D6	2C19	3A4	1A2	5HTTR			T102C
8	47	F	Cauc.	Major depressive disorder, sev w/o psychotic	1/5 Int. Freq: *1=37% *5=4%	1/1 Ext. Freq: *1=37%	*1/*1B	*1/*1	I/I			TC
9	25	F	Cauc.	Panic disorder	2/2P ext. freq: *2=33% *2P=unk.	1/1 Ext. Freq: *1=37%	*1*1	*1/*1	s/I			TC
10	16	F	Asian-American	Major depressive disorder vs. bipolar	4/10 Poor Freq: *4=18% *10=2%	1/1 Ext. Freq: *1=86%	*1*1	*1F/*1F	s/I			TT

[0122] Based on patient no. 5's genotype of HR2A C/C, I/I form of 5HTTR, extensive metabolizer for 1A2 and 3A4, and poor metabolizer for 2C19 and 2D6, the associated medication profile indicated that fluvoxamine and bupropion were acceptable for use, sertraline and mirtazapine should be used with caution, and citalopram, escitalopram, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and monitoring.

[0123] The medication profile associated with patient no. 6's genotype (HTR2A CC; 5HTTR I/s; 1A2, 3A4, and 2D6 extensive; and 2C19 poor) indicates that sertraline, venlafaxine, bupropion, and fluoxetine are acceptable for use, and mirtazapine, fluvoxamine, nortriptyline, citalopram, escitalopram, imipramine, amitriptyline, and paroxetine can be used with caution.

[0124] The medication profile associated with patient no. 7's genotype (HTR2A TT/TC; 5HTTR I/s; 1A2, 3A4, and 2C19 extensive; and 2D6 ultra rapid) indicates that citalopram, fluvoxamine, bupropion, and escitalopram are acceptable for use; sertraline and mirtazapine can be used with caution, and paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and close monitoring.

[0125] The medication profile associated with patient no. 8's genotype (HTR2A TT/TC; 5HTTR 1/1; 1A2, 3A4, and 2C19 extensive; and 2D6 intermediate) indicates that citalopram, fluvoxamine, bupropion, escitalopram, sertraline, and mirtazapine are acceptable for use; and paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and close monitoring.

[0126] The medication profile associated with patient no. 9's genotype (HTR2A TT/TC; 5HTTR I/s; and 1A2, 3A4, 2C19, and 2D6 extensive) indicates that citalopram, fluvoxamine, bupropion, escitalopram, sertraline, mirtazapine, paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline are acceptable for use.

[0127] The medication profile associated with patient no. 10's genotype (HTR2A TT/TC; 5HTTR I/s; 1A2, 3A4, and 2C19 extensive; and 2D6 poor) indicates that citalopram, fluvoxamine, bupropion, and escitalopram are acceptable for use, sertraline and mirtazapine should be used with caution, and paroxetine, fluoxetine, venlafaxine, amitriptyline, imipramine, and nortriptyline should be avoided or used with caution and close monitoring.

Example 5

[0128] The genotype of the CYP2D6, CYP2C19, CYP1A2, and HTR2A genes is determined in patients using the methods described above. Based on the CYP2D6 genotype, patients are classified as being poor, intermediate, extensive, or ultra rapid metabolizers. For the CYP2C 19 genotype, patients are classified as being either extensive, intermediate, or poor metabolizers. For the 1A2 genotype, patients are classified as being poor, intermediate, extensive, or ultra rapid metabolizers. For the 5HTTR genotype, patients are classified as having the short/short (s/s) form of the gene, the short/long (s/l) form of the gene, or the long/long (1/1) form of the gene. For the HR2A genotype, patients are classified as having the C/C allele, the T/T allele, or T/C allele.

[0129] The following five rules can be used to sort the genotypes into treatment recommendations. The first rule relates to the CYP2D6 genotype. For poor metabolizers and ultra rapid metabolizers, CYP2D6 substrates can be placed in the "use very cautiously with close monitoring" category. For intermediate metabolizers, exclusive CYP2D6 substrates can be placed in the "avoid or use very cautiously with close monitoring" category. For intermediate metabolizers, substrates that are partially metabolized by CYP2D6 can be placed in the "use cautiously" category. For extensive metabolizers, CYP2D6 substrates can be placed in the "acceptable for use" category.

[0130] After applying the first rule, the second rule relating to the 2C 19 genotype can be applied. For poor metabolizers based on their CYP2C19 genotype, exclusive CYP2C19 substrates can be placed in the "avoid or use very cautiously with close monitoring" category. For extensive or intermediate metabolizers based on their CYP2C19 genotype, CYP2C 19 substrates can be placed in the "acceptable for use" category. Substrates that are primarily metabolized by both CYP2D6 and CYP2C 19 can be placed in the "avoid or use very cautiously with close monitoring" category if CYP2C19 was poor and CYP2D6 was either intermediate or poor, or can be placed in the "use cautiously" category if CYP2C19 was poor and CYP2D6 was extensive.

[0131] A third rule relating to the 1A2 genotype can be applied after the first and second rules were applied. For poor and ultra-rapid metabolizers based on their CYP1A2 genotype, exclusive CYP1A2 substrates can be placed in the "avoid or use very cautiously with close monitoring" category (include fluvoxamine as a primary CYP1A2 substrate). For poor metabolizers based on the CYP1A2 genotype, substrates having the CYP1A2 pathway as one of their primary pathways can be placed in the "use with caution" category. For extensive or intermediate metabolizers based on their CYP1A2 genotype, CYP1A2 substrates can be placed in the "acceptable for use" category. In other embodiments, intermediate metabolizers can be placed in the "use with caution" category.

[0132] After applying the first three rules, the four rule relating to the 5HTTR genotype can be applied. For the homozygous short genotype (s/s) of the serotonin transporter gene, SSRI's that are categorized as "acceptable for use" can be shifted to the "use with caution" category. For the homozygous short genotype (s/s) of the serotonin transporter gene, SSRI's that are categorized as "use with caution" can be shifted to the "avoid or use very cautiously with close monitoring" category. Antidepressants, outside the SSRI class, do not shift category based on 5HTTR genotype.

[0133] Rule five, which relates to the HR2A genotype, can be applied after applying the first four rules. For the "mutant form" of the 5-hydroxytryptamine serotonin receptor 2A gene, if paroxetine is in the "acceptable for use" category, it can be shifted to the "use with caution" category. If paroxetine is in the "use with caution" category, it can be shifted to the "avoid or use very cautiously with close monitoring."

[0134] Based on the genotype of the patients and associated medication profile, medications appropriate for the patient can be selected.

Example 6

[0135] This examples describes an electronic chart review conducted of patients who had been enrolled in a pharmacogenomic study. Thirty-eight (38) patients had been treated with venlafaxine and genotyped for the 2D6 gene. Participants who were slow (also called "poor") metabolizers, as defined by having either two inactive 2D6 alleles or one inactive and one reduced 2D6 allele, had not been maintained on a dose of greater than 75 mg of venlafaxine. In contrast, 26 of the remaining 33 subjects (79%) who had at least one fully functioning 2D6 allele tolerated a dose of 150 mg or more of venlafaxine. ($p < .002$).

[0136] Clinical guidelines recommend a typical starting dose for venlafaxine to be 75 mg/day. However, an initial dose of 37.5 mg/day is regularly prescribed for geriatric patients. For patients who do not respond to an initial dose of 75 mg/day, increasing the daily dose by increments of 75 mg is recommended up to a maximum dose of 225 mg/day. Patients with no fully active 2D6 gene are predicted to have higher serum levels of both venlafaxine and N-desmethyl-venlafaxine (i.e., an inactive metabolite) for any given dose of venlafaxine when compared to patients with one or two functional copies of the 2D6 gene. Consequently, compromised metabolizers are less able to tolerate higher dosages of 2D6 substrate medications. Some patients with limited 2D6 metabolic capability can be expected to achieve a therapeutic clinical response at a relatively low dose.

Method

[0137] A sample of 199 patients, with a documented history of atypical response to psychotropic medications, were genotyped for the 2D6 gene as part of a clinical research protocol. Atypical response was defined as the development of adverse side effects or a non-response to the medication. Fourteen variant alleles were examined using a TM Bioscience platform. Test-retest reliability for determination of the 2D6 genotype was greater than 99% accurate. An electronic medical record review revealed that 38 of the participants in this protocol had previously been treated with venlafaxine by clinicians who had no knowledge of their 2D6 genotype before or during the course of their treatment. Table 10 provides the 2D6 alleles and the corresponding activity level.

Table 10

Allele	Activity Level
*1	normal
*2A	increased
*2BD	decreased
*3	none
*4	none
*5	none
*6	none
*7	none
*8	none
*9	decreased
*10	decreased
*11	none
*12	decreased
*17	decreased

(continued)

Allele	Activity Level
*41	decreased

[0138] Five "slow metabolizers" in this group of 38 patients were identified. Each of these patients had one completely inactive allele (e.g., *3 or *4). Additionally each patient had a second allele with reduced activity (e.g., *9, *10, *17 or *41). All of the other 33 patients treated with venlafaxine had at least one fully active allele.

[0139] None of the five patients with a slow 2D6 metabolism had been maintained on a dose of venlafaxine that was greater than 75 mg. In contrast, 26 of the remaining 33 subjects (79%) were maintained on dosage levels of 150 mg or 225 mg of venlafaxine. This difference between dosage levels was significant using a Fisher's exact test ($p < .002$). Clinical descriptions of the five subjects who lacked a functional 2D6 gene are shown below.

[0140] **Case 1:** A 44-year-old Caucasian woman was diagnosed with dysthymic disorder and was prescribed 75 mg of venlafaxine. She experienced dry mouth and an increase in appetite, but reported a decrease in depressive symptoms. Her dose was increased to 112.5 mg, at which time, her depression became worse and she complained of fatigue. Her dose was subsequently decreased to 75 mg and she did well for four months on this dose. She had a *3 2D6 allele and a *9 2D6 allele.

[0141] **Case 2:** A 16-year-old Caucasian teenager was diagnosed as having attention deficit hyperactivity disorder at 6 years of age. She subsequently became depressed at age 13, developed psychotic symptoms at age 15 and was diagnosed with bipolar disorder at age 16. She was prescribed 75 mg of venlafaxine for one month after which the medication was discontinued due to excessive somnolence and no improvement in her mood. She had a *3 2D6 allele and a *9 2D6 allele.

[0142] **Case 3:** A 54-year-old Caucasian woman who was diagnosed with major depressive disorder was initially prescribed 37.5 mg of venlafaxine. She discontinued the medication after four days due to intolerable nausea, insomnia, and decreased appetite. She had a *4 2D6 allele and a *17 2D6 allele.

[0143] **Case 4:** A 46-year-old Caucasian man was diagnosed with generalized anxiety disorder. He was prescribed 37.5 mg of venlafaxine a day and became more anxious over the next four days. As his complaints of anxiety increased, he also developed physiological symptoms that included palpitations of his heart. When his venlafaxine was discontinued, his anxiety symptoms improved. He had a *4 2D6 allele and a *41 2D6 allele.

[0144] **Case 5:** A 15-year-old Asian-American teenager was diagnosed as having major depressive disorder and was documented to have a past psychiatric history of three suicide attempts by overdose and five hospitalizations. Although there was concern that she had reported some mood fluctuations, she was prescribed 75 mg of venlafaxine. The venlafaxine was subsequently discontinued after five days because of intolerable side effects. She had previously experienced side effects when she had been placed on low doses of other 2D6 substrate medications (e.g., paroxetine and fluoxetine). She had a *4 2D6 allele and a *10 2D6 allele.

[0145] The analyses of this clinical sample of psychiatric patients indicates that individuals who lacked a full functioning copy of the 2D6 gene did not tolerate a maintenance dose of venlafaxine above 75 mg. In one case, a subject had briefly been able to take 112.5 mg before the treating clinician decreased the dose back to 75 mg (Case 5). This was the only patient with an inactive and partially inactive 2D6 allele that tolerated venlafaxine at any dose. In contrast, 79% of patients who had at least one fully functional copy of the 2D6 gene tolerated doses of venlafaxine of 150 mg or greater.

[0146] In reviewing the five patients who were identified as "slow metabolizers," two patterns were demonstrated. Four patients did not tolerate venlafaxine because of a range of side effects that included nausea, insomnia, anorexia, palpitation and somnolence. One subject who was a "slow metabolizer" had a therapeutic response at 75 mg, but she did not tolerate a dose increase to 112.5 mg.

[0147] One clinical implication of these analyses is that for patients who lack a functional copy of the 2D6 allele, venlafaxine should be initiated at a dose of 37.5 mg and the dosage should be increased cautiously. An alternative strategy would be to select an antidepressant with a primary metabolic pathway other than 2D6, such as fluvoxamine or an antidepressant with alternative pathways of metabolism, such as escitalopram or sertraline.

[0148] A strength of this clinical report is that none of the treating psychiatrists who cared for these patients were aware of their 2D6 genotypes. Consequently, dosing decisions were based on the clinical responses of the patients rather than predictions of phenotype based on genotypic information.

[0149] One limitation of this report is that the clinical data was retrieved from the electronic medical record. While venlafaxine dosage was always documented precisely, the descriptions of the effects and the therapeutic responses of patients were not reported in a standardized manner. The inclusion criteria for participating in the study was demonstrating an atypical response to antidepressant medication, defined as the development of adverse side effects or a non-response to the medication. Consequently, the decision to maintain, raise or lower medication dosage was based on the tolerance and response of the patient to the medication and was independent of diagnosis and comorbidity.

[0150] In summary, patients who did not have at least one fully functional allele of the 2D6 gene were not maintained on doses of venlafaxine greater than 75 mg. This association of compromised genotypes and a clinical history of intolerance to higher doses of venlafaxine, indicates that genetic variations of cytochrome P450 2D6 may contribute to the long observed patient specific variation in response to antidepressant medications.

OTHER EMBODIMENTS

[0151] It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims

[0152] Disclosed are the following items:

1. A method for selecting a psychotropic medication for a patient, said method comprising

(a) providing said patient's genotype for a panel of genes, wherein said panel comprises a CYP2D6 gene, a CYP2C 19 gene, a CYP1A2 gene, a serotonin transporter gene, and a serotonin receptor 2A gene; and
(b) selecting said psychotropic medication based on said genotype.

2. The method of item 1, wherein providing said patient's genotype comprises determining if said patient comprises the CYP1A2*1A or 1A2*3 allele, the CYP2C19*1A, 2C19*1B, or 2C19*2A allele, and the CYP2D6*1A, 2D6*2, 2D6*2N, 2D6*3, 2D6*4, 2D6*5, 2D6*6, 2D6*7, 2D6*8, 2D6*10, 2D6*12, or 2D6* 17 allele.

3. The method of item 1, wherein providing said patient's genotype further comprises determining if said patient comprises the 2D6*41 allele.

4. The method of item 1, wherein said panel comprises 4 cytochrome P450 genes.

5. The method of item 4, wherein said 4 cytochrome P450 genes are CYP2D6, 2C19, 3A4, and 1A2.

6. The method of item 5, wherein providing said patient's genotype comprises determining if said patient comprises the CYP2D6*2, 2D6*3, 2D6*4, 2D6*10, or 2D6*17 allele, the CYP2C19*2A, 2C19*2B, 2C19*3, 2C19*4, 2C19*5A, 2C19*5B, 2C19*6, 2C19*7, or 2C19*8 allele, the CYP3A4*1B, 3A4*2, 3A4*5, 3A4*6, 3A4* 12, 3A4*13, 3A4* 15A, 3A4*17, 3A4* 18A allele, and the CYP1A2*1F allele.

7. The method of item 1, wherein said panel comprises 5 cytochrome P450 genes.

8. The method of item 7, wherein said 5 cytochrome P450 genes are CYP1A1, 1A2, 2D6, 2C19, and 3A4.

9. The method of item 8, wherein providing said patient's genotype comprises determining if said patient comprises the CYP1A1 *1A, 1A1*2, 1A1*3, or 1A1*4 allele, the CYP1A2*1A or 1A2*3 allele, the CYP2C19*1A, 2C19*1B, or 2C19*2A allele, the CYP2D6*1A, 2D6*2, 2D6*2N, 2D6*3, 2D6*4, 2D6*5, 2D6*6, 2D6*7, 2D6*8, 2D6* 10, 2D6* 12, 2D6* 17, or 2D6*35 allele, and the CYP3A4*1A or 3A4* 1B allele.

10. The method of item 1, wherein said panel of genes further comprises a dopamine transporter gene.

11. The method of item 1, wherein said panel further comprises a plurality of dopamine receptor genes.

12. The method of item 11, wherein said dopamine receptor genes encode dopamine receptors D1, D2, D3, D4, D5, and D6.

13. The method of item 1, wherein said panel further comprises a serotonin receptor gene encoding serotonin receptors 1A, 1 B, 1D, or 2C.

14. The method of item 1, wherein said panel further comprises a catechol-O-methyl transferase gene.

15. The method of item 1, wherein said panel comprises at least 10 cytochrome P450 genes.

16. The method of item 15, wherein said at least 10 cytochrome P450 genes are CYP1A1, 1A2, 1B1, 2A6, 2B6,

2C8, 2C9, 2C18, 2C19, 2D6, 2E1, 3A4, and 3A5.

17. The method of item 16, wherein providing said patient's genotype comprises determining if said patient comprises the CYP1A1 *1A, 1A1 *2, 1A1*3, or 1A1*4 allele; the 1A2*1A or 1A2*3 allele; the CYP1B1*1, 1B1*2, 1B1*3, 1B1*4, 1B1*11, 1B1*14, 1B1*18, 1B1*19, 1B1*20, or 1B1 *25 allele; the CYP2A6*1A, 2A6*1B, 2A6*2, or 2A6*5 allele; the CYP2B6*1, 2B6*2, 2B6*3, 2B6*4, 2B6*5, 2B6*6, or 2B6*7 allele; the CYP2C8*1A, 2C8*1B, 2C8*1C, 2C8*2, 2C8*3, or 2C8*4 allele; the CYP2C9*1, 2C9*2, 2C9*3, or 2C9*5 allele; the CYP2C18*m1 or 2C18*m2 allele; the CYP2C19*1A, 2C19* 1B, or 2C19*2A allele; the CYP2D6*1A, 2D6*2, 2D6*2N, 2D6*3, 2D6*4, 2D6*5, 2D6*6, 2D6*7, 2D6*8, 2D6*10, 2D6*12, 2D6* 17, or 2D6*35 allele; the CYP2E1 *1A, 2E1 * 1C, 2E1*1D, 2E1*2, 2E1*4, 2E1*5, or 2E1*7 allele; the CYP3A4*1A or 3A4*1B allele; and the CYP3A5*1A, 3A5*3, 3A5*5, or 3A5*6 allele.

18. The method of item 1, wherein said panel further comprises a tryptophan hydroxylase gene.

19. The method of item 1, wherein said psychotropic medication is an antidepressant.

20. The method of item 1, wherein said psychotropic medication is an antipsychotic agent, an antianxiety-sedative agent, or a mood-stabilizing agent.

21. The method of item 1, wherein selecting said psychotropic medication comprises:

- (a) correlating genotype of said cytochrome P450 genes with capacity of each cytochrome P450 enzyme encoded by each said cytochrome P450 gene to metabolize said psychotropic medication; and
- (b) correlating genotype of said serotonin transporter gene and said serotonin receptor 2A gene with ability of said patient to respond to said psychotropic medication.

22. An article of manufacture comprising a substrate, said substrate comprising a plurality of discrete regions, wherein each said region comprises a different population of nucleic acid molecules, wherein said different populations of nucleic acid molecules independently comprise nucleic acid molecules for detecting the CYP1A2* 1A and 1A2*3 alleles; the CYP2C19*1A, 2C19*1B, and 2C19*2A alleles; the CYP2D6*1A, 2D6*2, 2D6*2N, 2D6*3, 2D6*4, 2D6*5, 2D6*6, 2D6*7, 2D6*8, 2D6*10, 2D6*12, 2D6*17, and 2D6*35 alleles; the serotonin receptor 2A C/T, T/T, C/C alleles; and the 5HTT promoter repeat and exon 2 variable repeat alleles.

EP 3 037 549 A1

SEQUENCE LISTING

<110> Mayo Foundation for Medical Education and Research
 5 <120> METHODS FOR SELECTING MEDICATIONS
 <130> 116-069T1
 <150> EP 07840663.4
 10 <151> 2007-08-02
 <150> PCT/US2007/075092
 <151> 2007-08-02
 <150> 11/498,976
 15 <151> 2006-08-02
 <160> 180
 <170> FastSEQ for Windows Version 4.0
 20 <210> 1
 <211> 22
 <212> DNA
 <213> Artificial Sequence
 <220>
 25 <223> Oligonucleotide
 <400> 1
 ccagaaggct ttgcaggctt ca 22
 <210> 2
 30 <211> 22
 <212> DNA
 <213> Artificial Sequence
 <220>
 35 <223> Oligonucleotide
 <400> 2
 actgagccct gggaggtagg ta 22
 <210> 3
 40 <211> 31
 <212> DNA
 <213> Artificial Sequence
 <220>
 45 <223> Oligonucleotide
 <400> 3
 ggcctaccct gggtaagggc ctggagcagg a 31
 <210> 4
 50 <211> 17
 <212> DNA
 <213> Artificial Sequence
 <220>
 55 <223> Oligonucleotide
 <400> 4
 cctggtcgaa gcagtat 17

	<210> 5	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> Oligonucleotide	
	<400> 5	
	gcacgctacc	10
10	<210> 6	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Oligonucleotide	
	<400> 6	
	gcacgctact	10
20	<210> 7	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Oligonucleotide	
	<400> 7	
	caccaggccc cctgccactg	20
30	<210> 8	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Oligonucleotide	
	<400> 8	
	cctgctcact cctggtagcc	20
40	<210> 9	
	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Oligonucleotide	
	<400> 9	
	ctgtttcatg tccacgacc	19
50	<210> 10	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Oligonucleotide	
	<400> 10	

	tgcccatcac	10
5	<210> 11 <211> 10 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
10	<400> 11 tgcccatcat	10
15	<210> 12 <211> 24 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
20	<400> 12 ccagatcctg ggtttcgggc cgcg	24
25	<210> 13 <211> 16 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
30	<400> 13 cagaggcgct tctcgg	16
35	<210> 14 <211> 20 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
40	<400> 14 ctcgggtctct cgctccgcac	20
45	<210> 15 <211> 10 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
50	<400> 15 cttctccgtg	10
55	<210> 16 <211> 10 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	

	<400> 16	
	cttctccgtc	10
5	<210> 17	
	<211> 26	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
10	<223> Oligonucleotide	
	<400> 17	
	tccaccttgc gcaacttggg cctggg	26
	<210> 18	
15	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
20	<400> 18	
	ccacccccag	10
	<210> 19	
25	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
30	<400> 19	
	ccacccccaa	10
	<210> 20	
35	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
40	<400> 20	
	gacgcccctt tcgcccac g	21
	<210> 21	
	<211> 20	
45	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
50	<400> 21	
	tgagacttgt ccaggtgaac	20
	<210> 22	
	<211> 20	
55	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	

	<400> 22 cccagatggg ctcacgctgc	20
5	<210> 23 <211> 10 <212> DNA <213> Artificial Sequence	
10	<400> 23 actgagcaca	10
	<210> 24 <211> 10 <212> DNA <213> Artificial Sequence	
15	<220> <223> Oligonucleotide	
20	<400> 24 actgagcacg	10
	<210> 25 <211> 25 <212> DNA <213> Artificial Sequence	
25	<220> <223> Oligonucleotide	
30	<400> 25 ggatgacctg ggacccagcc cagcc	25
	<210> 26 <211> 10 <212> DNA <213> Artificial Sequence	
35	<220> <223> Oligonucleotide	
40	<400> 26 gagaacctgc	10
	<210> 27 <211> 10 <212> DNA <213> Artificial Sequence	
45	<220> <223> Oligonucleotide	
50	<400> 27 gagaacctgt	10
	<210> 28 <211> 27 <212> DNA <213> Artificial Sequence	
55	<220> <223> Oligonucleotide	
	<400> 28	

	gcatagtggt ggctgacctg ttctctg	27
	<210> 29	
	<211> 20	
5	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
10	<400> 29	
	ccccgctga ctcccctctg	20
	<210> 30	
	<211> 21	
15	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
20	<400> 30	
	gacccccgag cctcaccttc c	21
	<210> 31	
	<211> 10	
25	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
30	<400> 31	
	agctgccacc	10
	<210> 32	
	<211> 10	
35	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
40	<400> 32	
	agctgccact	10
	<210> 33	
	<211> 15	
45	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
50	<400> 33	
	gcggaggccc caggt	15
	<210> 34	
	<211> 21	
55	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	

	<400> 34	
	accaagaggg aagaagcaca g	21
5	<210> 35	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
10	<223> Oligonucleotide	
	<400> 35	
	ccaatgacgg acaggagaaa	20
	<210> 36	
15	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
20	<400> 36	
	gagctggtga g	11
	<210> 37	
25	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
30	<400> 37	
	gagctggtga a	11
	<210> 38	
35	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
40	<400> 38	
	ctgcactccg ttctgctcct	20
	<210> 39	
45	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
50	<400> 39	
	gatgggggtc ctggtatgtc t	21
	<210> 40	
55	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	

	<400> 40	
	ctggagggtca cggctcaag	19
5	<210> 41	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
10	<223> Oligonucleotide	
	<400> 41	
	gtgtggcatg gaccttgtct g	21
	<210> 42	
15	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
20	<400> 42	
	gcaatgcgcc gtatttgttt c	21
	<210> 43	
25	<211> 15	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
30	<400> 43	
	aaagcttatt acaga	15
	<210> 44	
35	<211> 13	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
40	<400> 44	
	agcttattac agc	13
	<210> 45	
45	<211> 23	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
50	<400> 45	
	ggatgagatg gcatatgtcc tgc	23
	<210> 46	
55	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	

	<223> Oligonucleotide	
	<400> 46	
	cctgtttcct gtcgctgctc	20
5	<210> 47	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Oligonucleotide	
	<400> 47	
	ccaatacctg tccacgctga t	21
15	<210> 48	
	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Oligonucleotide	
	<400> 48	
	ggtgtcggaa c	11
25	<210> 49	
	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> Oligonucleotide	
	<400> 49	
	ggtgtcggaa a	11
35	<210> 50	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
40	<220>	
	<223> Oligonucleotide	
	<400> 50	
	ctgataacgg cagcacagac c	21
45	<210> 51	
	<211> 17	
	<212> DNA	
	<213> Artificial Sequence	
50	<220>	
	<223> Oligonucleotide	
	<400> 51	
	gggctggtgg tggagag	17
55	<210> 52	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	

	<220>		
	<223> Oligonucleotide		
	<400> 52		
5	tctgacacag ccaaggagat g		21
	<210> 53		
	<211> 9		
	<212> DNA		
10	<213> Artificial Sequence		
	<220>		
	<223> Oligonucleotide		
	<400> 53		
15	ccaggagcg		9
	<210> 54		
	<211> 10		
	<212> DNA		
	<213> Artificial Sequence		
20	<220>		
	<223> Oligonucleotide		
	<400> 54		
25	cccaggagcc		10
	<210> 55		
	<211> 19		
	<212> DNA		
	<213> Artificial Sequence		
30	<220>		
	<223> Oligonucleotide		
	<400> 55		
35	tggacaggat gagcagcga		19
	<210> 56		
	<211> 17		
	<212> DNA		
	<213> Artificial Sequence		
40	<220>		
	<223> Oligonucleotide		
	<400> 56		
45	gggctggtgg tggagag		17
	<210> 57		
	<211> 21		
	<212> DNA		
	<213> Artificial Sequence		
50	<220>		
	<223> Oligonucleotide		
	<400> 57		
55	tctgacacag ccaaggagat g		21
	<210> 58		
	<211> 10		
	<212> DNA		
	<213> Artificial Sequence		

	<220>		
	<223> Oligonucleotide		
5	<400> 58		10
	ccaggagcgt		
	<210> 59		
	<211> 10		
	<212> DNA		
10	<213> Artificial Sequence		
	<220>		
	<223> Oligonucleotide		
15	<400> 59		10
	ccaggagcgg		
	<210> 60		
	<211> 20		
	<212> DNA		
20	<213> Artificial Sequence		
	<400> 60		20
	ggacaggatg agcagcgaca		
	<210> 61		
25	<211> 21		
	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Oligonucleotide		
30	<400> 61		21
	tctccacagc actcccgaca g		
	<210> 62		
	<211> 21		
35	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Oligonucleotide		
40	<400> 62		21
	ccgagaacaa tggcgagcat c		
	<210> 63		
	<211> 9		
45	<212> DNA		
	<213> Artificial Sequence		
	<220>		
	<223> Oligonucleotide		
50	<400> 63		9
	ggtccgggt		
	<210> 64		
	<211> 9		
55	<212> DNA		
	<213> Artificial Sequence		
	<220>		

	<223> Oligonucleotide	
	<400> 64	
	ggtccgggc	9
5		
	<210> 65	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
10		
	<220>	
	<223> Oligonucleotide	
	<400> 65	
	tttgccattg ggcattgtct g	21
15		
	<210> 66	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
20		
	<220>	
	<223> Oligonucleotide	
	<400> 66	
	caccagccca cccgagag	18
25		
	<210> 67	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
30		
	<220>	
	<223> Oligonucleotide	
	<400> 67	
	accgagaaca atggcgagca t	21
35		
	<210> 68	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
40		
	<220>	
	<223> Oligonucleotide	
	<400> 68	
	ccatggtggg	10
45		
	<210> 69	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
50		
	<220>	
	<223> Oligonucleotide	
	<400> 69	
	ccatggtggc	10
55		
	<210> 70	
	<211> 17	
	<212> DNA	
	<213> Artificial Sequence	

	<220>		
	<223> Oligonucleotide		
	<400> 70		
5	acgggtcggg gagagtc		17
	<210> 71		
	<211> 18		
	<212> DNA		
10	<213> Artificial Sequence		
	<220>		
	<223> Oligonucleotide		
	<400> 71		
15	ccggtacagc cccatccc		18
	<210> 72		
	<211> 21		
	<212> DNA		
	<213> Artificial Sequence		
20	<220>		
	<223> Oligonucleotide		
	<400> 72		
25	gccgactcac cgagaacaat g		21
	<210> 73		
	<211> 9		
	<212> DNA		
	<213> Artificial Sequence		
30	<220>		
	<223> Oligonucleotide		
	<400> 73		
35	gtgggacgg		9
	<210> 74		
	<211> 10		
	<212> DNA		
	<213> Artificial Sequence		
40	<220>		
	<223> Oligonucleotide		
	<400> 74		
	ggtgggacga		10
45	<210> 75		
	<211> 19		
	<212> DNA		
	<213> Artificial Sequence		
50	<220>		
	<223> Oligonucleotide		
	<400> 75		
	gtcggggaga gtcagctgg		19
55	<210> 76		
	<211> 22		
	<212> DNA		
	<213> Artificial Sequence		

	<220>	
	<223> Oligonucleotide	
	<400> 76	
5	ggggacagga gggaggggtg ag	22
	<210> 77	
	<211> 25	
	<212> DNA	
10	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
	<400> 77	
15	gaagaggagt gggcaggaga tgggtg	25
	<210> 78	
	<211> 12	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
	<400> 78	
25	ggagatcatg ac	12
	<210> 79	
	<211> 13	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
	<400> 79	
35	tggagatcat gat	13
	<210> 80	
	<211> 16	
	<212> DNA	
40	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
	<400> 80	
45	ggtgacccgg cgcttg	16
	<210> 81	
	<211> 20	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
	<400> 81	
55	agcaaccaag ccccaaagag	20
	<210> 82	
	<211> 19	
	<212> DNA	

	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
5	<400> 82	
	gagcgcgcag taggagagg	19
	<210> 83	
	<211> 11	
10	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
15	<400> 83	
	ttcaggtggc t	11
	<210> 84	
	<211> 10	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
25	<400> 84	
	tcaggtggcc	10
	<210> 85	
	<211> 21	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
35	<400> 85	
	actcagctgg ctcagagatg c	21
	<210> 86	
	<211> 21	
	<212> DNA	
40	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
45	<400> 86	
	cgctacaacc ggcaggggtgg g	21
	<210> 87	
	<211> 21	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
55	<400> 87	
	gcagcatgag cgggcagggt a	21
	<210> 88	
	<211> 11	

	<212> DNA	
	<213> Artificial Sequence	
	<220>	
5	<223> Oligonucleotide	
	<400> 88	
	ggacgagtag a	11
	<210> 89	
10	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
15	<223> Oligonucleotide	
	<400> 89	
	ggacgagtag c	11
	<210> 90	
20	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
25	<223> Oligonucleotide	
	<400> 90	
	ccacgtagtc gcggtcctc	19
	<210> 91	
30	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
35	<223> Oligonucleotide	
	<400> 91	
	cgccttcccc cagccacc	19
	<210> 92	
40	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
45	<223> Oligonucleotide	
	<400> 92	
	gttggagccg caggggtccg	20
	<210> 93	
50	<211> 8	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
55	<223> Oligonucleotide	
	<400> 93	
	ggcgcggg	8
	<210> 94	

	<211> 8	
	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> Oligonucleotide	
	<400> 94	
	ggcgcggc	8
10	<210> 95	
	<211> 15	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Oligonucleotide	
	<400> 95	
	gggcgcacag tcggg	15
20	<210> 96	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Oligonucleotide	
	<400> 96	
	tgcggttttc catcaagaag	20
30	<210> 97	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Oligonucleotide	
	<400> 97	
	ggagttggcc cagcagaa	18
40	<210> 98	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Oligonucleotide	
	<400> 98	
	ccgacagggt	10
50	<210> 99	
	<211> 9	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Oligonucleotide	
	<400> 99	
	cgacagggg	9

	<210> 100	
	<211> 27	
	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> Oligonucleotide	
	<400> 100	
	cttgagaacc ttggtctcct tcttgat	27
10	<210> 101	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Oligonucleotide	
	<400> 101	
	caacgccgac ttctggaa	18
20	<210> 102	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Oligonucleotide	
	<400> 102	
	acatgcatc gaaaggactc t	21
30	<210> 103	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Oligonucleotide	
	<400> 103	
	cggggggaac	10
40	<210> 104	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Oligonucleotide	
	<400> 104	
	cggggggaat	10
50	<210> 105	
	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Oligonucleotide	
	<400> 105	
	ggcgttgggc atcatgtgg	19

	<210> 106	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> Oligonucleotide	
	<400> 106	
10	gcagtccagc ccgaaatg	18
	<210> 107	
	<211> 16	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Oligonucleotide	
	<400> 107	
20	gcacgatggc tgcgga	16
	<210> 108	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Oligonucleotide	
	<400> 108	
30	ctgcggacac	10
	<210> 109	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Oligonucleotide	
	<400> 109	
40	ctgcggacat	10
	<210> 110	
	<211> 15	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Oligonucleotide	
	<400> 110	
50	cagcacgttg cccag	15
	<210> 111	
	<211> 13	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Oligonucleotide	
	<400> 111	

	caacgtgctg gtg	13
5	<210> 112 <211> 15 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
10	<400> 112 catcttgccg ttgta	15
15	<210> 113 <211> 8 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
20	<400> 113 ggctgccc	8
25	<210> 114 <211> 9 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
30	<400> 114 tggctgccc	9
35	<210> 115 <211> 17 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
40	<400> 115 acaccagcac gttgccc	17
45	<210> 116 <211> 16 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	
50	<400> 116 cgccaagatg accaac	16
55	<210> 117 <211> 17 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	

	<400> 117 cggaatgaag gagatga	17
5	<210> 118 <211> 10 <212> DNA <213> Artificial Sequence	
10	<220> <223> Oligonucleotide	
	<400> 118 gccacgaaga	10
15	<210> 119 <211> 10 <212> DNA <213> Artificial Sequence	
20	<220> <223> Oligonucleotide	
	<400> 119 gccacgaagc	10
25	<210> 120 <211> 21 <212> DNA <213> Artificial Sequence	
30	<220> <223> Oligonucleotide	
	<400> 120 ggtctgacac agccagagac a	21
35	<210> 121 <211> 20 <212> DNA <213> Artificial Sequence	
40	<220> <223> Oligonucleotide	
	<400> 121 gcagctgcct acatccacat	20
45	<210> 122 <211> 21 <212> DNA <213> Artificial Sequence	
50	<220> <223> Oligonucleotide	
	<400> 122 cattcgggggt gaaaggtggt a	21
55	<210> 123 <211> 10 <212> DNA <213> Artificial Sequence	
	<220> <223> Oligonucleotide	

	<400> 123 gtccagctcc	10
5	<210> 124 <211> 10 <212> DNA <213> Artificial Sequence	
10	<220> <223> Oligonucleotide	
	<400> 124 gtccagctct	10
15	<210> 125 <211> 23 <212> DNA <213> Artificial Sequence	
20	<220> <223> Oligonucleotide	
	<400> 125 cagacagact ctgcaacagg gtc	23
25	<210> 126 <211> 21 <212> DNA <213> Artificial Sequence	
30	<220> <223> Oligonucleotide	
	<400> 126 ggagagtcgg ggagcaggaa c	21
35	<210> 127 <211> 17 <212> DNA <213> Artificial Sequence	
40	<220> <223> Oligonucleotide	
	<400> 127 gaggcggggg cacaagg	17
45	<210> 128 <211> 11 <212> DNA <213> Artificial Sequence	
50	<220> <223> Oligonucleotide	
	<400> 128 gcacagagca c	11
55	<210> 129 <211> 11 <212> DNA <213> Artificial Sequence	
	<220>	

	<223> Oligonucleotide	
	<400> 129	
	gcacagagca t	11
5	<210> 130	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Oligonucleotide	
	<400> 130	
	ccccagcctt gctctcca	18
15	<210> 131	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Oligonucleotide	
	<400> 131	
	ctacatcccg ctgctgctca t	21
25	<210> 132	
	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> Oligonucleotide	
	<400> 132	
	cctgctcccc gactctcca	19
35	<210> 133	
	<211> 9	
	<212> DNA	
	<213> Artificial Sequence	
40	<220>	
	<223> Oligonucleotide	
	<400> 133	
	gcgcagctc	9
45	<210> 134	
	<211> 9	
	<212> DNA	
	<213> Artificial Sequence	
50	<220>	
	<223> Oligonucleotide	
	<400> 134	
	gcgcagcta	9
55	<210> 135	
	<211> 24	
	<212> DNA	
	<213> Artificial Sequence	

	<220>		
	<223> Oligonucleotide		
5	<400> 135		24
	ggaatatgcg cccatagaga acca		
	<210> 136		
	<211> 25		
	<212> DNA		
10	<213> Artificial Sequence		
	<220>		
	<223> Oligonucleotide		
15	<400> 136		25
	gctcacttgg cttattggct tcctc		
	<210> 137		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
20	<220>		
	<223> Oligonucleotide		
25	<400> 137		20
	catgagcagc agcgggatgt		
	<210> 138		
	<211> 10		
	<212> DNA		
	<213> Artificial Sequence		
30	<220>		
	<223> Oligonucleotide		
35	<400> 138		10
	catgcgtcgg		
	<210> 139		
	<211> 10		
	<212> DNA		
	<213> Artificial Sequence		
40	<220>		
	<223> Oligonucleotide		
45	<400> 139		10
	catgcgtcga		
	<210> 140		
	<211> 18		
	<212> DNA		
	<213> Artificial Sequence		
50	<220>		
	<223> Oligonucleotide		
55	<400> 140		18
	ggtccgagcg gtcttccg		
	<210> 141		
	<211> 24		
	<212> DNA		
	<213> Artificial Sequence		

	<220>		
	<223> Oligonucleotide		
	<400> 141		
5	agggcaacaa caccacatca ccac		24
	<210> 142		
	<211> 22		
	<212> DNA		
10	<213> Artificial Sequence		
	<220>		
	<223> Oligonucleotide		
	<400> 142		
15	gcagcaccaa caccgacacc at		22
	<210> 143		
	<211> 11		
	<212> DNA		
	<213> Artificial Sequence		
20	<220>		
	<223> Oligonucleotide		
	<400> 143		
25	acgtcggaga t		11
	<210> 144		
	<211> 10		
	<212> DNA		
	<213> Artificial Sequence		
30	<220>		
	<223> Oligonucleotide		
	<400> 144		
35	cgtcggagac		10
	<210> 145		
	<211> 19		
	<212> DNA		
	<213> Artificial Sequence		
40	<220>		
	<223> Oligonucleotide		
	<400> 145		
	accagtagtg ttgccgccg		19
45	<210> 146		
	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
50	<220>		
	<223> Oligonucleotide		
	<400> 146		
	tcagggaac aacaccacat		20
55	<210> 147		
	<211> 18		
	<212> DNA		

	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
5	<400> 147	
	gcaccaacac cgacacca	18
	<210> 148	
	<211> 9	
10	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
15	<400> 148	
	gttgccgcc	9
	<210> 149	
	<211> 9	
20	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
25	<400> 149	
	gttgccgct	9
	<210> 150	
	<211> 19	
30	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
35	<400> 150	
	ggtctcaaag ggagccggt	19
	<210> 151	
	<211> 20	
40	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
45	<400> 151	
	ttttcttccc tcccccttcc	20
	<210> 152	
	<211> 17	
50	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Oligonucleotide	
55	<400> 152	
	ccaccacgca cgcattg	17
	<210> 153	
	<211> 10	

	<212> DNA	
	<213> Artificial Sequence	
	<220>	
5	<223> Oligonucleotide	
	<400> 153	
	aagggagccg	10
	<210> 154	
10	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
15	<223> Oligonucleotide	
	<400> 154	
	aaagggagcc a	11
	<210> 155	
20	<211> 22	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
25	<223> Oligonucleotide	
	<400> 155	
	gtggtgatgt ggtgttgttg cc	22
	<210> 156	
30	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
35	<223> Oligonucleotide	
	<400> 156	
	gctcaactac gaactcccta a	21
	<210> 157	
40	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
45	<223> Oligonucleotide	
	<400> 157	
	tcactacggc tgtcagtaaa g	21
	<210> 158	
50	<211> 12	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
55	<223> Oligonucleotide	
	<400> 158	
	ttagcttctc ca	12
	<210> 159	

	<211> 12	
	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> Oligonucleotide	
	<400> 159	
	ttagcttctc cg	12
10	<210> 160	
	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Oligonucleotide	
	<400> 160	
	gagttaaagt cattactgta gagcctggt	29
20	<210> 161	
	<211> 16	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Oligonucleotide	
	<400> 161	
	gctctgctcg ctcaac	16
30	<210> 162	
	<211> 21	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Oligonucleotide	
	<400> 162	
	gatttttagtg gctttccttt c	21
40	<210> 163	
	<211> 9	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Oligonucleotide	
	<400> 163	
	ccgagtgcg	9
50	<210> 164	
	<211> 10	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Oligonucleotide	
	<400> 164	
	gccgagtgca	10

	<210> 165	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> Oligonucleotide	
	<400> 165	
	agtgccctc atggaggc	18
10	<210> 166	
	<211> 25	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Oligonucleotide	
	<400> 166	
	ctcaatttta aactttgggtt gctta	25
20	<210> 167	
	<211> 23	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Oligonucleotide	
	<400> 167	
	ttttgaatag gaaacaccca taa	23
30	<210> 168	
	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Oligonucleotide	
	<400> 168	
	gcattcctca g	11
40	<210> 169	
	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Oligonucleotide	
	<400> 169	
	gcattcctca c	11
50	<210> 170	
	<211> 28	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Oligonucleotide	
	<400> 170	
	gttcaccatg attgcttcag tcttaagc	28

	<210> 171	
	<211> 30	
	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> Oligonucleotide	
	<400> 171	
10	atgatgacaa tgatgctgat gatgataatg	30
	<210> 172	
	<211> 25	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Oligonucleotide	
	<400> 172	
20	tccaccatcg gaggtattga aaatg	25
	<210> 173	
	<211> 15	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Oligonucleotide	
	<400> 173	
30	tcacagaaat atcac	15
	<210> 174	
	<211> 14	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Oligonucleotide	
	<400> 174	
40	cacagaaata tcag	14
	<210> 175	
	<211> 26	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Oligonucleotide	
	<400> 175	
50	attgccaaac caataggcca attagg	26
	<210> 176	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Oligonucleotide	
	<400> 176	

ggcttcagtt tcccgatctg 20

<210> 177
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide

<400> 177
 taggcaaggg caggcactac 20

<210> 178
 <211> 10
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide

<400> 178
 aaggagctgg 10

<210> 179
 <211> 11
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide

<400> 179
 gaaggagctg a 11

<210> 180
 <211> 16
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide

<400> 180
 gtacatgggg gcccc 16

Claims

1. A method of determining the initial dose of venlafaxine for a patient, the method comprising:

- (1) determining the patient's genotype for a panel of cytochrome P450 CYP2D6 alleles comprising *1, *2A, *2BD, *3, *4, *5, *6, *7, *8, *9, *10, *11, *12,*17, and *41, wherein *1 and *2A are designated as functional alleles, *2BD, *9, *10, *12,*17, and *41 are designated as decreased activity alleles, and *3, *4, *5, *6, *7, *8, and *11 are designated as inactive alleles; and
- (2) determining if the patient is a poor 2D6 metabolizer, wherein a patient lacking a functional 2D6 allele is characterized as a poor 2D6 metabolizer;

wherein the initial dose of venlafaxine is determined to be a reduced initial dose for the patient characterized as a poor 2D6 metabolizer according to steps (1) and (2).

2. The method of claim 1, wherein the reduced initial dose is less than 75 mg/day.
3. The method of claim 2, wherein the reduced initial dose is 37.5 mg/day.
- 5 4. The method of claim 1, wherein the method further comprises receiving, in a computer system, the patient's genotype for the panel of cytochrome P450 CYP2D6 alleles, said computer system comprising a database, wherein said database comprises a plurality of psychotropic medication profiles.
- 10 5. The method of claim 4, wherein one or more of the steps are performed using said computer system.
6. The method of claim 5, wherein the method further comprises the step of outputting from the computer system the initial dose of a psychotropic medication for administering to the patient.
- 15 7. The method of claim 4, wherein a user enters the patient's genotype in the computer system.
8. The method of claim 4, wherein the patient's genotype is received directly from equipment used in determining the patient's genotype.
- 20 9. The method of claim 1, wherein the psychiatric disease is selected from dysthymic disorder, attention deficit hyperactivity disorder, major depressive disorder or generalized anxiety disorder.
10. The method of claim 9, wherein the psychiatric disease is selected from generalized anxiety disorder.
- 25 11. Venlafaxine for use in the treatment of a psychiatric disease wherein the initial dose of said psychotropic medication is determined by the method of claim 1.
12. A computer readable medium containing executable instructions that when executed cause a processor to perform operations comprising:
 - 30 (a) receiving a patient's genotype for a panel of cytochrome P450 CYP2D6 alleles comprising *1, *2A, *2BD, *3, *4, *5, *6, *7, *8, *9, *10, *11, *12, *17, and *41, wherein *1 and *2A are designated as functional alleles, *2BD, *9, *10, *12, *17, and *41 are designated as decreased activity alleles, and *3, *4, *5, *6, *7, *8, and *11 are designated as inactive alleles,
 - 35 (b) assigning a metabolic phenotype to the patient based on the genotype of the cytochrome P450 genes in (a), wherein the patient is assigned a poor 2D6 metabolizer phenotype if the patient lacks a functional 2D6 allele, and
 - (c) outputting an initial dose of venlafaxine for the patient wherein the initial dose is a reduced initial dose of half the typical starting dose recommended by the clinical guidelines for venlafaxine .
- 40 13. The method of claim 12, wherein the reduced initial dose is less than 75 mg/day.
14. The method of claim 13, wherein the reduced initial dose is 37.5 mg/day.




FIGURE 1




FIGURE 2




FIGURE 3



EUROPEAN SEARCH REPORT

Application Number
EP 15 02 0211

5

10

15

20

25

30

35

40

45

50

55

2

EPO FORM 1503 03.82 (P04C01)

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	WO 2004/074456 A2 (MAYO FOUNDATION [US]; MRAZEK DAVID A [US]; O'KANE DENNIS J [US]; BLACK) 2 September 2004 (2004-09-02) * claims 1,2,15 *	1,12	INV. C12Q1/68
A	WO 2005/038049 A2 (HEINRICH GUENTHER [DE]; ROOTS IVAR [DE]) 28 April 2005 (2005-04-28) * page 34 - page 38 *	1,12	
A	UEDA M ET AL: "The impact of CYP2D6 genotypes on the plasma concentration of paroxetine in Japanese psychiatric patients", PROGRESS IN NEURO-PSYCHOPHARMACOLOGY & BIOLOGICAL PSYCHIATRY, OXFORD, GB, vol. 30, no. 3, 1 May 2006 (2006-05-01), pages 486-491, XP025129418, ISSN: 0278-5846, DOI: 10.1016/J.PNPBP.2005.11.007 [retrieved on 2006-05-01] * the whole document *	1,12	TECHNICAL FIELDS SEARCHED (IPC) C12Q
A	ARRANZ M J ET AL: "Pharmacogenetics for the individualization of psychiatric treatment", AMERICAN JOURNAL OF PHARMACOGENOMICS, WOLTERS KLUWER HEALTH, XX, vol. 1, no. 1, 1 January 2001 (2001-01-01), pages 3-10, XP009092573, ISSN: 1175-2203 * the whole document *	1,12	
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 19 May 2016	Examiner Gabriels, Jan
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



EUROPEAN SEARCH REPORT

Application Number
EP 15 02 0211

5

10

15

20

25

30

35

40

45

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	HUAI-RONG LUO ET AL: "Identification of CYP2D6 impaired functional alleles in Mexican Americans", EUROPEAN JOURNAL OF CLINICAL PHARMACOLOGY, SPRINGER, BERLIN, DE, vol. 61, no. 11, 1 December 2005 (2005-12-01), pages 797-802, XP019329993, ISSN: 1432-1041, DOI: 10.1007/S00228-005-0044-4 * the whole document *	1,12	TECHNICAL FIELDS SEARCHED (IPC)
A	----- DATABASE MEDLINE [Online] US NATIONAL LIBRARY OF MEDICINE (NLM), BETHESDA, MD, US; 2006, RASMUSSEN J O ET AL: "CYP2D6 gene test in psychiatric patients and healthy volunteers.", XP002675622, Database accession no. NLM16537246 * abstract * & SCANDINAVIAN JOURNAL OF CLINICAL AND LABORATORY INVESTIGATION 2006 LNKD-PUBMED:16537246, vol. 66, no. 2, 2006, pages 129-136, ISSN: 0036-5513	1,12	
T	----- DATABASE MEDLINE [Online] US NATIONAL LIBRARY OF MEDICINE (NLM), BETHESDA, MD, US; September 2007 (2007-09), MCALPINE DONALD E ET AL: "Cytochrome P450 2D6 genotype variation and venlafaxine dosage.", XP002675623, Database accession no. NLM17803873 * abstract * -/--		
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 19 May 2016	Examiner Gabriels, Jan
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03/82 (P04C01)

50

55



EUROPEAN SEARCH REPORT

 Application Number
 EP 15 02 0211

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	& MAYO CLINIC PROCEEDINGS. MAYO CLINIC SEP 2007 LNKD- PUBMED:17803873, vol. 82, no. 9, September 2007 (2007-09), pages 1065-1068, ISSN: 0025-6196 ----- BISHOP J R AND ELLINGROD V L: "Neuropsychiatric pharmacogenetics: moving toward a comprehensive understanding of predicting risks and response", PHARMACOGENOMICS,, vol. 5, no. 5, 1 July 2004 (2004-07-01), pages 463-477, XP008102578, ISSN: 1462-2416 -----	1,12	
			TECHNICAL FIELDS SEARCHED (IPC)
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 19 May 2016	Examiner Gabriels, Jan
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

 2
 EPO FORM 1503 03/02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 15 02 0211

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

19-05-2016

10

15

20

25

30

35

40

45

50

55

ORM P0459

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2004074456 A2	02-09-2004	DK 2393028 T3	11-01-2016
		EP 1599576 A2	30-11-2005
		EP 2393028 A2	07-12-2011
		ES 2557885 T3	29-01-2016
		HK 1164503 A1	22-04-2016
		JP 5348840 B2	20-11-2013
		JP 5438524 B2	12-03-2014
		JP 5438610 B2	12-03-2014
		JP 2007525154 A	06-09-2007
		JP 2010176661 A	12-08-2010
		JP 2010273681 A	09-12-2010
		US 2007003931 A1	04-01-2007
		US 2013311202 A1	21-11-2013
		US 2015148336 A1	28-05-2015
		WO 2004074456 A2	02-09-2004

WO 2005038049 A2	28-04-2005	NONE	

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5451683 A [0064]
- WO 9820019 A [0064]

Non-patent literature cited in the description

- **WONG et al.** *Ann. Acad. Med. Singapore*, 2001, vol. 29, 401-406 [0020]
- **SALLEE et al.** *J. Child Adol. Psychiatry*, 2000, vol. 10 (1), 27-34 [0020]
- **SACHSE et al.** *Br. J. Clin. Pharmacol.*, 1999, vol. 47, 445-449 [0023]
- **IBEANU et al.** *J. Pharmacol. Exp. Ther.*, 1999, vol. 290, 635-640 [0023]
- PCR Primer: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1995 [0029]
- **LEWIS.** *Genetic Engineering News*, 1992, vol. 12 (9), 1 [0029]
- **GUATELLI et al.** *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 1874-1878 [0029]
- **WEISS.** *Science*, 1991, vol. 254, 1292-1293 [0029]
- **STONEKING et al.** *Am. J. Hum. Genet.*, 1991, vol. 48, 370-382 [0033]
- **PRINCE et al.** *Genome Res.*, 2001, vol. 11 (1), 152-162 [0033]
- **MYAKISHEV et al.** *Genome*, 2001, vol. 11 (1), 163-169 [0036]