



(11)

EP 3 125 875 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:

07.06.2023 Bulletin 2023/23

(51) International Patent Classification (IPC):

A61K 9/72 (2006.01) **A61K 9/14** (2006.01)
A61K 31/436 (2006.01) **A61P 11/00** (2006.01)

(21) Application number: **15718687.5**

(52) Cooperative Patent Classification (CPC):

A61K 9/0075; A61K 9/145; A61K 31/436;
A61K 45/06; A61P 11/00; A61P 35/00;
A61P 39/06; A61P 43/00

(22) Date of filing: **06.04.2015**

(86) International application number:

PCT/US2015/024551

(87) International publication number:

WO 2015/154084 (08.10.2015 Gazette 2015/40)

(54) **AN INHALABLE RAPAMYCIN FORMULATION FOR TREATING AGE-RELATED CONDITIONS**

INHALIERBARE RAPAMYCIN-FORMULIERUNG ZUR BEHANDLUNG ALTERSBEDINGTER ERKRANKUNGEN

PRÉPARATION À INHALER CONTENANT DE LA RAPAMYCINE POUR TRAITER LES PATHOLOGIES LIÉES À L'ÂGE

(84) Designated Contracting States:

AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR

(74) Representative: **Haseltine Lake Kempner LLP**

Cheapside House
138 Cheapside
London EC2V 6BJ (GB)

(30) Priority: **04.04.2014 US 201461975127 P**

(56) References cited:

WO-A1-2015/054280 WO-A1-2016/130645
WO-A2-2008/063581 WO-A2-2008/137148
US-A1- 2013 102 569

(43) Date of publication of application:
08.02.2017 Bulletin 2017/06

(60) Divisional application:
23162835.5

- **CARVALHO SIMONE R ET AL: "Characterization and pharmacokinetic analysis of crystalline versus amorphous rapamycin dry powder via pulmonary administration in rats", EUROPEAN JOURNAL OF PHARMACEUTICS AND BIOPHARMACEUTICS, ELSEVIER SCIENCE PUBLISHERS B.V., AMSTERDAM, NL, vol. 88, no. 1, 22 May 2014 (2014-05-22), pages 136-147, XP029053289, ISSN: 0939-6411, DOI: 10.1016/J.EJPB.2014.05.008**

(73) Proprietor: **Al Therapeutics, Inc.**
Guilford, CT 06437 (US)

(72) Inventors:

- **ARMER, Thomas**
Guilford, CT 06437 (US)
- **MELVIN, Lawrence, S.**
Guilford, CT 06437 (US)
- **ROTHBERG, Jonathan, M.**
Guilford, CT 06437 (US)
- **LICHENSTEIN, Henri**
Guilford, CT 06437 (US)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 3 125 875 B1

- STEFAN W. RYTER ET AL: "Autophagy in lung disease pathogenesis and therapeutics", REDOX BIOLOGY : AN OFFICIAL JOURNAL OF THE SOCIETY FOR FREE RADICAL BIOLOGY AND MEDICINE, AN OFFICIAL JOURNAL OF THE SOCIETY FOR FREE RADICAL RESEARCH-EUROPE, AN AFFILIATE JOURNAL OF THE INTERNATIONAL SOCIETY FOR FREE RADICAL RESEARCH (SFRRI), vol. 4, 1 April 2015 (2015-04-01), pages 215-225, XP055547308, NL ISSN: 2213-2317, DOI: 10.1016/j.redox.2014.12.010

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Description**FIELD OF THE INVENTION**

[0001] The present invention relates to pharmaceutical compositions for use in methods for pulmonary delivery by inhalation, the compositions comprising rapamycin for anti-aging and for the prophylaxis and treatment of age-related conditions diseases and disorders, wherein the age-related disorder is chronic lung disease.

BACKGROUND OF THE INVENTION

[0002] Rapamycin is a macrocyclic triene antibiotic produced by *Streptomyces hygroscopicus*. See e.g., U.S. Pat. No. 3,929,992. Rapamycin is an inhibitor of mTOR. The immunosuppressive and anti-inflammatory properties of rapamycin initially indicated its use in the transplantation field and in the treatment of autoimmune diseases. For example, it was shown to prevent the formation of humoral (IgE-like) antibodies in response to an albumin allergic challenge, to inhibit murine T-cell activation, and to prolong survival time of organ grafts in histoincompatible rodents. In rodent models of autoimmune disease, it suppresses immune-mediated events associated with systemic lupus erythematosus, collagen-induced arthritis, autoimmune type I diabetes, autoimmune myocarditis, experimental allergic encephalomyelitis, graft-versus-host disease, and autoimmune uveoretinitis.

[0003] Rapamycin is also referred to by its generic drug name, sirolimus (see for example, ANDA #201578, by Dr. Reddys Labs Ltd., approved May 28, 2013). Sirolimus is FDA approved and marketed in the United States for the prophylaxis of organ rejection and renal transplantation under the trade name RAPAMUNE by Wyeth (Pfizer). It is in the form of an oral solution (1 mg/ml) or tablet (multiple strengths). Wyeth (Pfizer) also markets a derivative by the tradename TORISEL (temsirolimus) for the treatment of advanced renal cell carcinoma, which is administered intravenously. Temsirolimus is a water-soluble prodrug of sirolimus. Cordis, a division of Johnson & Johnson, markets a sirolimus-eluting coronary stent under the tradename CYPHER. In this context, the antiproliferative effects of sirolimus prevent restenosis in coronary arteries following balloon angioplasty. US 2010/0305150 to Berg et al. (Novartis) describes rapamycin derivatives for treating and preventing neurocutaneous disorders, such as those mediated by TSC including tuberous sclerosis, as well as those mediated by neurofibromatosis type 1 (NF-1). Rapamycin and its derivatives are further described in Nishimura, T. et al. (2001) Am. J. Respir. Crit. Care Med. 163:498-502 and in U.S. Pat. Nos. 6,384,046 and US 6,258,823.

[0004] Rapamycin use in its clinically approved context has several known adverse effects including lung toxicity (the RAPAMUNE label warns that it is not indicated for lung transplant patients), increased cancer risk, and diabetes-like symptoms. Rapamycin is associated with the occurrence of pulmonary toxicity, usually in the form of interstitial pneumonitis, but pulmonary alveolar proteinosis has also been documented. See for example, Nocera et al., Sirolimus Therapy in Liver Transplant Patients: An Initial Experience at a Single Center, Transplantation Proceedings (2008), 40(6), 1950-1952; Perez et al., Interstitial Pneumonitis Associated With Sirolimus in Liver Transplantation: A Case Report, Transplantation Proceedings (2007), 39(10), 3498-3499; Hashemi-Sadraei et al., Sirolimus-associated diffuse alveolar hemorrhage in a renal transplant recipient on long-term anticoagulation, Clinical Nephrology (2007), 68(4), 238-244; Pedroso et al., Pulmonary alveolar proteinosis - a rare pulmonary toxicity of sirolimus, Transplant International (2007), 20(3), 291-296. The cause of rapamycin-induced pulmonary toxicity is not known.

[0005] Severe respiratory adverse events have also been associated with sirolimus use as an anti-cancer therapy under chronic administration resulting in circulating blood concentrations greater than 1 nanogram/mL range. For example, the lung toxicity of the sirolimus prodrug, temsirolimus, was documented in a 2009 report noting that "interstitial lung disease is a rare side effect of temsirolimus treatment in renal cancer patients". Aparicio et al., Clinical & Translational Oncology (2009), 11(8), 499-510; Vahid et al., Pulmonary complications of novel antineoplastic agents for solid tumors, Chest (2008) 133:528-538. In addition, a 2012 meta-analysis concluded that 10% of cancer patients administered temsirolimus or everolimus may experience mild grade toxicity with a worsening of quality of life and, in some case, interruption of therapy. See Iacovelli et al., Incidence and risk of pulmonary toxicity in patients treated with mTOR inhibitors for malignancy. A meta-analysis of published trials, Acta oncologica (2012), 51(7), 873-879. Furthermore, safety pharmacology studies performed in rats with temsirolimus (see Pharm/Tox section of temsirolimus NDA) showed reductions in respiratory rate as well as alveolar macrophage infiltration and inflammation in the lungs (see Pharmacology Review for temsirolimus NDA 22088 available from the US FDA website). These adverse effects were observed under conditions of relatively high concentrations of the drug in the circulating blood volume as a result of systemic administration.

[0006] A U.S. patent application by Lehrer published in 2013 reflects the view that "[r]apamycin (sirolimus) cannot be safely inhaled because of its well-documented lung toxicity, interstitial pneumonitis". See US 20130004436, citing Chhajed et al. (2006) 73:367-374. The Lehrer patent application is directed to compositions and methods for treating and preventing lung cancer and lymphangioleiomyomatosis. Although some earlier publications, such as US Patent No. 5,080,899 to Sturm et al. (filed February 1991) and US Patent No. 5,635,161 (filed June 1995), contain some generic

description of rapamycin for delivery by inhalation, such generic descriptions were unsupported by any evidence and came before the many reported incidences of rapamycin-induced lung toxicity that appeared following its more wide-spread adoption as an immunosuppressant in the transplantation context and as an inhibitor of cellular proliferation in the anti-cancer context, as evidenced by the reports discussed above.

[0007] WO 2011/163600 describes an aerosol formulation of tacrolimus, which like rapamycin is a macrolide lactone. But tacrolimus is a distinct chemical entity from sirolimus and the molecular target of tacrolimus is calcineurin, not mTOR, and unlike rapamycin, tacrolimus did not show lung toxicity and in fact is indicated for preventing rejection following lung transplantations.

[0008] In view of the wide-spread recognition of the potential for rapamycin-induced lung toxicity, a pharmaceutical composition comprising rapamycin for pulmonary delivery was not considered to be a viable therapeutic option in humans.

[0009] Delivery of drugs to the lung by way of inhalation is an important means of treating a variety of conditions, including such common local conditions as cystic fibrosis, pneumonia, bronchial asthma and chronic obstructive pulmonary disease, some systemic conditions, including hormone replacement, pain management, immune deficiency, erythropoiesis, diabetes, lung cancer, *etc.* See review by Yi et al. J. Aerosol Med. Pulm. Drug Deliv. 23:181-7 (2010). Agents indicated for treatment of lung cancer by inhalation include cisplatin, carboplatin, taxanes, and anthracyclines. See *e.g.*, U.S. Pat. Nos. 6,419,900; 6,419,901; 6,451,784; 6,793,912; and U.S. Patent Application Publication Nos. US 2003/0059375 and US 2004/0039047. In addition, doxorubicin and temozolomide administered by inhalation have been suggested for treating lung metastases. See *e.g.*, U.S. Pat. No. 7,288,243 and U.S. Patent Application Publication No. US 2008/0008662.

[0010] US 20100260733 ("the '733 application") describes the signaling pathways of the TOR (target of rapamycin) kinase as integrating diverse cellular signals from agents such as nutrients, mitogenic growth factors, energy, and stress-related signals to regulate the catabolic and anabolic processes of the cell. The TOR pathway is also described as possibly playing an important role in the life span extension induced by caloric restriction in budding yeast, *Ceanorhabditis elegans* and *Drosophila*. Mitochondrial function is described as playing an important role in maintaining senescence induced by telomere dysfunction and caloric restriction in preventing deterioration of the senescent state through the TOR/AMPK/mitochondrial pathway. This mechanism is reported to be conserved in both yeast and human models of telomere dysfunction. The '733 application provides methods for identifying agents that promote mitochondrial function and thereby prevent or treat the deterioration of senescence. According to the '733 application, since many age-related diseases are linked to mitochondrial dysfunction and/or telomere dysfunction, agents identified by this method can potentially be used to prevent age-related diseases or disorders. This is said to be in contrast to the currently prevalent anti-aging strategies, which are mainly focused on inhibiting senescence by taking senescence as the major contributor to the aging process, rather than by maintaining the senescent state. The '733 application includes a number of *in vitro* based experiments using various cell types showing, for example that low doses of rapamycin inhibit mutagen-induced transformation in human fibroblasts and reduce reactive oxygen species and extend life span in cultured neurons. The '733 application includes only two animal studies, both purporting to demonstrate that low doses of rapamycin reduce infarction size in a rat model of stroke or myocardial infarction.

[0011] U.S. Patent No. 8,492,110 ("the '110 patent") issued from the '733 application and claims methods of preventing cell aging by extending the G0 phase in a cell in a subject in need thereof by administering a low dose (defined as a dose that does not inhibit protein translation and cell growth at the G1 phase of the cell cycle) of a TOR inhibitor.

[0012] U.S. Patent Application No. 20130102569 ("the '569 application") describes methods for treating or preventing an age-related disease, condition, or disorder comprising administering a therapeutically effective amount of an inhibitor of TOR to a patient in need thereof. The '569 application describes a somewhat different role of the TOR inhibitor compared with the '733 application, namely the '569 application provides data purportedly demonstrating that rapamycin inhibits the onset of cellular senescence *in vitro*, rather than acting to maintain the senescent state by preventing its deterioration. The '569 application does not provide any animal data in support of its methods of treating or preventing an age-related disease, condition, or disorder.

[0013] WO 2008/137148 relates to methods for treating, stabilizing, preventing and/or delaying pulmonary hypertension by administering nanoparticles that comprise rapamycin or a derivative thereof and/or nanoparticles that comprise a taxane (*e.g.*, paclitaxel) or a derivative thereof. The described compositions (*e.g.*, unit dosage forms) comprise nanoparticles that comprise a carrier protein and rapamycin or a derivative thereof and/or nanoparticles that comprise a carrier protein and a taxane (*e.g.* paclitaxel) or a derivative thereof.

[0014] WO 2008/063581 relates to a method for treatment of respiratory disorders such as asthma, chronic obstructive pulmonary disease, and chronic sinusitis, including cystic fibrosis, interstitial fibrosis, chronic bronchitis, emphysema, bronchopulmonary dysplasia and neoplasia. The method involves administration, preferably oral, nasal or pulmonary administration, of anti-inflammatory and anti-proliferative drugs (rapamycin or paclitaxel and their analogues). The described particles should be aggregates of a drug, for example rapamycin, and an additive.

[0015] There is a need for pharmaceutical formulations of rapamycin, that can safely be delivered directly to the lungs, preferably by inhalation, in order to provide a more effective dosage form for the treatment and prophylaxis of diseases

and disorders affected by the TOR signaling pathway that reduces or eliminates the toxicities and adverse events associated with oral dosage forms of rapamycin.

BRIEF DESCRIPTION OF THE FIGURES

[0016]

Fig. 1: LC-MS/MS Chromatogram of 10.6 ng/mL Rapamycin (top) and Internal Standard (bottom) in Mouse Blood.

Fig. 2: Representative Chromatograms of 10.6 ng/mL Rapamycin (top) and Internal Standard (bottom) in Mouse Lung Homogenate.

Fig. 3: Calibration Curve for Rapamycin in Mouse Blood.

Fig. 4: Calibration Curve for Rapamycin in Mouse Lung Homogenate.

Fig. 5: Representative Chromatograms of Rapamycin (top) and Internal Standard (bottom) in Blood from Mouse 2-07 Administered Rapamycin by OPA.

Fig. 6: Representative Chromatograms of Rapamycin (top) and Internal Standard (bottom) in Lung Homogenate from Mouse 2-07 Administered Rapamycin by OPA.

Fig. 7: Rapamycin inhibits viability of TSC2 mutant cells (bottom) and inhibits S6 phosphorylation (top).

Fig. 8A and B: S6 Phosphorylation in Mouse Lung Following (A) OPA and oral administration of rapamycin, and (B) administration via inhalation.

Fig. 9: Predicted rapamycin blood concentrations for pulmonary administration repeated once daily.

SUMMARY OF THE INVENTION

[0017] The present invention is based, in part, upon the development of a safe and effective aerosol formulation of a rapamycin composition that is capable of delivering amounts of the rapamycin composition to target tissues effective to exert potent biological activity in those target tissues while minimizing rapamycin associated toxicity. In addition, the invention exploits the discovery of the surprising pharmacokinetics of a rapamycin composition formulated as described herein. As discussed in more detail *infra*, the rapamycin composition delivered directly to the lungs produced markedly higher concentrations of drug in the lung tissue. The amount of drug in lung tissue was unexpectedly higher than what was predictable from previous oral and intravenous studies. And surprisingly, even relatively high amounts of rapamycin delivered directly to the lungs did not result in toxicity to the lung tissue. Moreover, the amount of drug in lung tissue achieved by the methods described here is demonstrated to be effective to exert potent biological activities including inhibition of cell growth and viability as well as inhibition of S6 phosphorylation in the target tissue. These biological activities indicate that the delivered dose of the rapamycin composition according to the claimed methods is sufficient to inhibit mTOR signaling in the target tissues. Thus, these results demonstrate that the aerosol formulations described here are capable of delivering a low, yet therapeutically effective, dose of rapamycin providing for very low systemic exposure to the drug combined with high efficacy. The result is a markedly improved therapeutic index for rapamycin when administered according to the present invention.

[0018] The present invention is defined in the appended claims. In particular the present invention provides a pharmaceutical aerosol composition in the form of a dry powder for pulmonary delivery comprising micronized rapamycin particles having a Mass Median Aerodynamic Diameter (MMAD) of from 0.5 to 5 micrometres (μm) (microns) and particles of a carrier for use in a method of treating a chronic lung disease, the method comprising delivering the dry powder composition to a human subject in need thereof by inhalation, wherein the formulation is effective to deliver a therapeutic amount of rapamycin to the lungs, and wherein the chronic lung disease is not Lymphangioleiomyomatosis (LAM).

[0019] The present invention provides pharmaceutical aerosol formulations of a rapamycin composition for delivery directly to the lungs. In this context, the term "aerosol formulation" refers to a dry powder composition, as described in more detail *infra*. An aerosol formulation of the invention is delivered to a subject by inhalation. As used herein, the term "rapamycin composition" may refer to rapamycin itself, preferably in the amorphous form described as sirolimus. In one embodiment, a rapamycin composition of the invention provides an amount of rapamycin effective to inhibit mTOR signaling in a target tissue with low or no toxicity to the tissue, and concomitant blood levels of rapamycin that are less than about 1 ng/ml.

[0020] In one embodiment, a rapamycin composition of the invention provides an improved safety profile, as evidenced by a higher therapeutic index, especially with respect to its chronic or prolonged use, compared to other dosage forms of rapamycin, for example oral or intravenous dosage forms.

[0021] Compositions are described herein for the treatment and prophylaxis of an age related disease or disorder, wherein the age-related disorder is chronic lung disease, by administering once daily to a human subject in need thereof a pharmaceutical aerosol formulation comprising a rapamycin composition in an amount effective to achieve a maximum

blood level of the rapamycin composition of from about 0.25 to 0.75 ng/ml and a blood trough level of from about 0.075 to 0.25 ng/ml. The maximum blood level may be about 0.5 ng/ml, the blood trough level is about 0.1 ng/ml, and the amount of the rapamycin composition in the formulation is from about 25 to 100 micrograms, or about 50 micrograms, delivered once a day.

[0022] The aerosol formulations of the invention may be formulated with a rapamycin composition alone, or in combination with one or more additional therapeutic agents, in the same dosage form. In addition, the aerosol formulations of the invention may be administered alone, or in combination with one or more additional therapies, each administered either by the same or a different route, e.g., orally, intravenously, etc. In one embodiment, the aerosol formulations of the invention may be administered in combination with one or more additional therapeutic regimens for the treatment of the age related disease or disorder.

[0023] In one embodiment, the present invention provides a pharmaceutical aerosol formulation comprising a rapamycin composition in an amount effective to achieve a therapeutic level of the composition in a target tissue. In the present invention the tissue is lung. The therapeutic level may be determined 12 or 24 hours after delivery, preferably 24 hours after delivery. In one embodiment, the therapeutic level is sustained for at least 24 hours after delivery.

[0024] In one embodiment, the target tissue is lung. The lung to blood concentration ratio of the composition 24 hours after delivery may be at least 100, at least 250, or at least 500. The lung to blood concentration ratio of the composition 24 hours after delivery may be from about 100 to 250, 250 to 500, 500 to 750, or 750 to 1000. The lung to blood concentration ratio of the composition 24 hours after delivery may be at least 5, at least 10, at least 20, at least 30, at least 50, at least 60, at least 70, at least 80, or at least 100.

[0025] In one embodiment, the amount of the rapamycin composition in the aerosol formulation is from 5 to 500 micrograms, from 10 to 250 micrograms, from 15 to 150 micrograms, or from 20 to 100 micrograms. In one embodiment, the amount of the rapamycin composition in the aerosol formulation is 20, 40, 50, 100, 125, or 250 micrograms.

[0026] In one embodiment, the rapamycin composition is sirolimus. In an embodiment not according to the invention, the rapamycin composition is selected from the group consisting of everolimus, temsirolimus, ridaforolimus, umirolimus, and zotarolimus.

[0027] In one embodiment, the rapamycin composition is sirolimus and has an isomeric B:C ratio of greater than 30:1 or greater than 35:1. In one embodiment, the rapamycin composition has an isomeric B:C ratio of 3.5% to 10%.

[0028] In the present invention the age-related disease or disorder is chronic lung disease.

[0029] In one embodiment, the pharmaceutical aerosol composition is for use in a method, the method is an anti-aging therapy effective to prevent aging of the lung. In one embodiment, the subject is a geriatric subject, the method comprises administering the composition over a period of time, and the efficacy of the method is measured as an improvement in grip strength or walking ability following administration of the composition for the period of time. In one embodiment, the composition is administered once or twice daily for a period of time selected from 1 to 3 weeks, less than one month, one to 2 months, 2 to 3 months, or 3 to 4 months.

[0030] In an embodiment not according to the invention, the method comprises the step of administering the composition to the subject produces particles comprising rapamycin having an average mean diameter in the range of 0.1 to 10 μm (microns). In another embodiment not according to the invention, the method comprises the step of administering the composition to the subject produces particles comprising rapamycin having an average mean diameter in the range of 0.5 to 6 μm (microns).

[0031] In one embodiment, the method further comprises one or more additional therapies or therapeutic regimens.

[0032] In one embodiment, the aerosol formulation of the invention is adapted for once daily administration and, according the methods described herein, the aerosol formulation is administered once a day.

[0033] In one embodiment, the aerosol formulation is a dry powder composition suitable for delivery by inhalation. In one embodiment, the dry powder comprises the rapamycin composition in the form of microparticles (*i.e.*, microparticulate rapamycin), particles of a carrier, and one or more optional excipients. In one embodiment, the microparticles consist of particles of drug having mean diameters from about 1 to 5 μm (microns). In one embodiment, the particles have a mean diameter of about 1.5 to 4 μm (microns), about 1.5 to 3.5 μm (microns), or about 2 to 3 μm (microns). The carrier may be selected from the group consisting of arabinose, glucose, fructose, ribose, mannose, sucrose, trehalose, lactose, maltose, starches, dextran, mannitol, lysine, leucine, isoleucine, dipalmitoylphosphatidylcholine, lecithin, polylactic acid, poly (lactic-co-glutamic) acid, and xylitol, and mixtures of any of the foregoing. In one embodiment, the carrier comprises or consists of a blend of two different carriers. The particles of carrier may have diameters ranging from to 200 μm (microns), from 30 to 100 μm (microns), or less than 10 μm (microns). Where the carrier consists of a blend of two different carriers, each carrier consists of particles of a different size range, measured as average particle diameter. In one embodiment, the carrier consists of a blend of two different carriers, a first carrier and a second carrier. The first carrier consists of particles having diameters ranging from about 30-100 μm (microns) and the second carrier consists of particles having diameters of less than 10 μm (microns). The ratio of the two different carriers is in the range of from 3:97 to 97:3. In one embodiment, the ratio of the two different carriers is in the range of from 97:3 or from 95-98:2-5. In one embodiment, the carrier consists of a blend of two different lactose carriers. The drug to carrier ratio in the powder

may be from 0.5 % to 2 % (w/w). In one embodiment, the drug to carrier ratio in the powder is 1 % (w/w).

[0034] The amount of the rapamycin composition in the aerosol formulation is from about 0.1 % to 20 % (w/w) based upon total weight of the composition. In one embodiment, the amount is from about 0.25 % to 2 % (w/w).

[0035] In one embodiment, one or more optional excipients is present in the composition and is selected from a phospholipid and a metal salt of a fatty acid, and mixtures of the foregoing. In one embodiment, the phospholipid is selected from dipalmitylphosphatidylcholine and lecithin. In one embodiment, the metal salt of a fatty acid is magnesium stearate. In one embodiment, the excipient or excipients is coated on the carrier particles in a weight ratio of excipient to large carrier particle ranging from 0.01 to 0.5%

[0036] In one embodiment, the amount of the rapamycin composition in the aerosol formulation is an amount effective to inhibit the biological activity of mTORC1. In one embodiment, the amount is an amount effective to inhibit the phosphorylation of the S6K protein.

[0037] In one embodiment, the amount of the rapamycin composition in the aerosol formulation is an amount effective to achieve a respirable dose of from 5 to 500 micrograms delivered to the lung. In one embodiment, the respirable dose is about 5, about 20, about 50, about 100 or about 250 micrograms. In one embodiment, the respirable dose is about 20 micrograms, about 50 micrograms, or about 100 micrograms. In one embodiment, the amount is an amount effective to produce a concentration of the rapamycin composition in the lung tissue of from 1 ng/g to 1 microgram (ug)/g. In one embodiment, the concentration of the rapamycin composition in the lung tissue is from about 5 to 30 ng/g. In one embodiment, the concentration in the lung tissue is about 5 ng/g, about 10 ng/g, about 15 ng/g, about 20 ng/g, about 25 ng/g, about 30 ng/g, about 50 ng/g, about 60 ng/g, about 100 ng/g, or about 200 ng/g. In accordance with the foregoing embodiments, the concomitant blood trough level of the rapamycin composition is less than 5 ng/ml, less than 2 ng/ml, less than 1 ng/ml, less than 0.5 ng/ml, or less than 0.25 ng/ml. In one embodiment, the blood trough level is less than 1 ng/ml, less than 0.5 ng/ml or less than 0.25 ng/ml.

[0038] In one embodiment, the rapamycin composition persists in lung at therapeutic levels of about 1 ng/g, about 5 ng./g, about 10 ng/g, about 15 ng/g, about 20 ng/g, about 25 ng/g, about 50 ng/g, or about 100 ng/g for a period of time after administration, preferably to a human subject, the period of time selected from about 6 to 10 hours, about 6 to 14 hours, about 6 to 24 hours, and about 6 to 72 hours. In one embodiment, the period of time is selected from about 12 hours, about 14 hours, about 24 hours, and about 72 hours.

[0039] In one embodiment, the rapamycin composition persists in lung at therapeutic levels of about 5 to 100 ng/g or from about 5 to 30 ng/g for a period of time that is about 12 or 24 hours. In one embodiment, the rapamycin composition persists in lung at therapeutic levels of about 5 ng/g, about 10 ng/g, about 20 ng/g, about 30 ng/g, about 50 ng/g, about 60 ng/g, about 70 ng/g, about 80 ng/g, or about 90 ng/g. In one embodiment, the rapamycin composition persists in lung at therapeutic levels of at least 5 ng/g, at least 20 ng/g, or at least 30 ng/g. In one embodiment, the rapamycin composition persists in lung at therapeutic levels of from about 20 ng/g to about 30 ng/g or from about 50 ng/g to about 80 ng/g.

[0040] In one embodiment, the formulation has a fine particle fraction (FPF) greater than 20% with a corresponding fine particle dose (FPD) ranging from 5 micrograms to 2 milligrams, preferably less than 0.5 milligrams, following 1 to 12 months or 1 to 36 months of storage. In one embodiment, the respirable dose, which is the dose delivered to the lung, also referred to as the delivered dose (DD) or emitted dose (ED), ranges from 10 micrograms to 2.5 milligrams, preferably less than 0.5 milligrams. In one embodiment, the delivered dose is from about 20 to 100 micrograms, from about 10 to 25 micrograms or from about 30 to 60 micrograms. In one embodiment, the delivered dose is 20 or 50 micrograms. In one embodiment, the delivered dose is 100 micrograms.

[0041] In one embodiment, the respirable dose of the rapamycin composition is about 20 micrograms, the concentration of drug in the lung tissue is from about 5 to 25 ng/g, the C_{max} in blood is less than 1.0 ng/ml, or from about 0.50 ng/ml to 1.0 ng/ml, or about 0.50 ng/ml to 0.90 ng/ml, the blood trough concentration of drug at 24 hrs post-dosing is less than about 0.20 ng/ml, and the steady-state concentration of drug in the blood at 14 days post-dosing is less than about 0.90 ng/ml, or less than about 0.80 ng/ml.

[0042] In one embodiment, the respirable dose of the rapamycin composition is about 50 micrograms, the concentration of drug in the lung tissue is about 2 to 15 ng/g, the C_{max} in blood is less than 2.0 ng/ml, or from about 0.25 ng/ml to 0.1 ng/ml, or about 0.10 ng/ml to 0.5 ng/ml, the blood trough concentration of drug after a single dose, 24 hrs post-dosing is less than about 0.10 ng/ml, and the trough concentration of drug in the blood after 5 days repeated, once-daily, is less than about 1.0 ng/ml, or less than about 0.50 ng/ml. In one embodiment, the formulation is adapted for once daily administration.

[0043] In one embodiment, the formulation further comprises one or more additional therapeutic agents.

[0044] The invention provides compositions of the invention for use in a method for the treatment and prophylaxis of age-related diseases and disorders, wherein the age related disorder is chronic lung disease, in a human subject in need of such treatment. In one embodiment, the invention provides compositions of the invention for use in a method for treatment and prophylaxis of age-related diseases and disorders in a human subject in need of such treatment or prophylaxis, the method comprising administering to the subject via inhalation a composition or unit dosage form described herein.

[0045] Also provided, but not according to the invention, is a unit dosage form comprising an aerosol formulation comprising a rapamycin composition as described herein, wherein the amount of the rapamycin composition is from about 5 to 2500 micrograms, from 20 to 500 micrograms, or from 50 to 250 micrograms. In one embodiment, the amount of the rapamycin composition is from about 50 to 125 micrograms. In one embodiment, the amount of the rapamycin composition is about 40, about 50, about 100, about 125, or about 250 micrograms. In one embodiment, the amount of the rapamycin composition is about 250 micrograms.

[0046] The unit dosage form is a capsule suitable for use in a dry powder inhaler device. In one embodiment, the capsule contains from 1 mg to 100 mg of the powder (total amount, including the rapamycin composition, carrier, and any optional excipients) or from 10 mg or 40 mg of the powder. The capsule may be a gelatin, plastic, or cellulosic capsule, or in the form of a foil/foil or foil/plastic blister suitable for use in a DPI device.

[0047] Also described herein, but not according to the invention, is a pharmaceutical package or kit comprising a composition or unit dosage form described herein, and instructions for use.

[0048] In one embodiment not according to the invention, the formulation is produced by a wet polishing process comprising the steps of preparing an aqueous suspension of drug, subjecting the drug suspension to microfluidization, and spray-drying the resulting particles to form a dry powder.

[0049] In one embodiment, the rapamycin composition is sirolimus, the carrier consists of a blend of two different lactose carriers, the first carrier consists of particles having average diameters ranging from about 30-100 μm (microns) and the second carrier consists of particles having average diameters of less than 10 μm (microns), the ratio of the two different carriers is about 97:3 to 3:97, and the amount of rapamycin is from 25 to 1400 micrograms.

[0050] Also described herein, but not according to the invention, is a dry powder delivery device comprising a reservoir containing a composition or unit dosage form described herein. The reservoir may be an integral chamber within the device, a capsule, or a blister. In the following embodiments, which are not according to the invention, the device is selected from Plastiap[®] RS01 Model 7, Plastiap[®] RS00 Model 8, XCaps[®], Handihaler[®], Flowcaps[®] TwinCaps[®], and Aerolizer[®]. In one embodiment, the device is selected from Plastiap[®] RS01 Model 7 or Plastiap[®] RS00 Model 8. In one embodiment, the device is Plastiap[®] RS00 Model 8.

DETAILED DESCRIPTION OF THE INVENTION

[0051] The present invention provides compositions for use in a method of treatment and prophylaxis of age-related diseases and disorders in a human subject in need of such treatment, wherein the age-related disorder is chronic lung disease. In one embodiment, the methods comprise administering to the subject via inhalation a composition comprising rapamycin in a suitable carrier, and optionally one or more additives. The term "rapamycin" is used generically throughout this invention disclosure to refer to rapamycin (also referred to as sirolimus) as well as pharmaceutically acceptable salts of rapamycin.

[0052] The compositions described herein are referred to as "aerosol formulations" and are meant to describe aerosolizable compositions suitable for producing respirable particles containing a rapamycin composition, which as described above refers to rapamycin, preferably in the amorphous form described as sirolimus. In one embodiment, the rapamycin composition is sirolimus. The aerosol formulations described herein may comprise the rapamycin composition, a carrier, and optionally one or more additives. The aerosol formulations are in the form of a dry powder.

[0053] The present invention also provides compositions for use in methods for the treatment and prophylaxis of age-related diseases and disorders in a human subject in need of such treatment, wherein the age-related disorder is chronic lung disease, the methods comprising the step of pulmonary administration of an aerosol formulation of the invention to the subject. In one embodiment, the administered dose of the rapamycin composition is sufficient to achieve therapeutic levels of rapamycin in the lung tissue while maintaining a low blood level, or blood trough level, in the subject. For example, the therapeutic levels of the rapamycin composition may be from about 1 ng/g, about 5 ng/g, about 10 ng/g, about 15 ng/g, about 20 ng/g, about 25 ng/g, about 50 ng/g and the blood trough level is from 0.01 to 0.15 ng/ml, from 0.075 to 0.350 ng/ml, from 0.150 to 0.750 ng/ml, from 0.750 to 1.5 ng/ml, or from 1.5 to 5 ng/ml. In one embodiment, the administered dose is sufficient to achieve a therapeutic level of drug in the lung of from about 5 ng/g to 50 ng/g, or from about 5 ng/g to 20 ng/g and a blood trough level of drug of less than 5 ng/ml, less than 2 ng/ml, less than 1 ng/ml, or less than 0.5 ng/ml. In one embodiment, the lung to blood concentration ratio of the rapamycin composition 24 hours after delivery is at least 100, at least 250, or at least 500. In one embodiment, the lung to blood concentration ratio of the rapamycin composition 24 hours after delivery is from about 100 to 250, 250 to 500, 500 to 750, or 750 to 1000.

[0054] Preferably, the aforementioned therapeutic levels are achieved by administering an aerosol formulation described herein once a day. In one embodiment, the total daily dose of the rapamycin composition is in the range of from 5 to 100 micrograms, from 20 to 250 micrograms, from 50 to 500 micrograms (0.05 to 0.5 milligrams), from 250 to 1000 micrograms (0.25 to 1 milligrams) or from 500 to 2000 micrograms (0.5 to 2 milligrams). In one embodiment, the total daily dose is less than 500 micrograms, less than 100 micrograms, less than 50 micrograms, less than 20 micrograms, or less than 10 microgram. In one embodiment, the total daily dose is less than 500 micrograms, less than 250 micrograms,

less than 100 micrograms, less than 50 micrograms, or less than 10 micrograms. In one embodiment, the total daily dose administered to the subject is less than 0.5 mg or less than 0.25 mg per day. Further aspects of pulmonary delivery and dosing, including combination therapies, are described in the section below entitled "Pulmonary Administration and Dosing".

[0055] This disclosure comprises compositions for use in a method, the method comprising administering rapamycin via a pulmonary route in combination with one or more additional agents selected from other anti-agent agents and antioxidants. The one or more additional agents is selected from the group consisting of vitamin C, vitamin E, beta carotene and other carotenoids, selenium, lipoic acid, lycopene, lutein, zeaxanthin, coenzyme Q10, glutathione, N-acetyl cysteine, melatonin, genistein, estradiol, tea extract, and grape seed extract, AICAR, EGCG, grapefruit extract, bilberry extract, selenite, genistein, diallyl trisulfide, benzyl isothiocyanate, phenyl isothiocyanate, phenethyl isothiocyanate, resveratrol, lycopene, and allyl isothiocyanate. The one or more additional agents may be administered by the same or a different route of administration as the rapamycin. For example, the agent may be administered by inhalation, intranasally, orally or intravenously.

[0056] The compositions and compositions for use in methods of the invention are effective as anti-aging therapy and for the treatment and prophylaxis of age-related diseases and disorders in a subject, preferably a human subject, wherein the disorder is chronic lung disease. As used herein, the effective amount of a composition of the invention refers to the amount sufficient to reduce or ameliorate the progression, severity, and/or duration of a disease or disorder, or one or more symptoms of a disease or disorder, to prevent the advancement of a disease or disorder, cause the regression of a disease or disorder, to prevent the development or onset of one or more symptoms associated with a disease or disorder, to enhance or improve the prophylactic or therapeutic effect(s) of another therapy (e.g., a prophylactic or therapeutic agent) with respect to the severity or onset of one or more symptoms of a disease or disorder, or with respect to the development or progression of a disease or disorder. Thus, in the context of the methods of the invention, the terms "treat", "treatment", and "treating" refer to the reduction of the severity, duration, or progression of a disease or disorder or one or more symptoms associated with a disease or disorder.

[0057] In certain embodiments, the methods include pulmonary administration of a composition of the invention as the primary therapy. In other embodiments, the administration of a composition of the invention is an adjuvant therapy. In either case, the methods of the invention contemplate the administration of a composition of the invention in combination with one or more additional therapies for the treatment of a disease or disorder. The terms "therapy" and "therapies" refer to any method, protocol and/or agent that can be used in the prevention, treatment, management or amelioration of a disease or disorder, or one or more symptoms thereof.

[0058] The one or more additional therapies may be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a composition of the invention.

[0059] In some embodiments, an additional therapeutic agent is formulated for co-administration with a composition of the invention in a dosage form for pulmonary administration. In other embodiments, an additional therapeutic agent is administered separately from the dosage form that contains rapamycin, and by the same or different route of administration as the rapamycin. The methods of the invention also contemplate a combination of one or more additional therapeutic agents for administration concomitantly with, before, or after the administration of the dosage form comprising rapamycin.

[0060] In certain embodiments, the methods of the invention are effective to manage a disease or disorder in a subject having the disease or disorder. In this context, the terms "manage", "managing", and "management" refer to the beneficial effects that a subject derives from a therapy which does not result in a cure. In one embodiment, the disease or disorder is managed in the subject if its progression is slowed or stopped during treatment with rapamycin according to the methods of the invention. In another embodiment, the disease or disorder is managed in the subject if one or more symptoms associated with the disease or disorder is ameliorated or stabilized (*i.e.*, the symptom does not worsen during the course of treatment).

[0061] In one embodiment, the methods of the invention are directed to subjects who are "non-responsive" or "refractory" to a currently available therapy for the disease or disorder. In this context, the terms "non-responsive" and "refractory" refer to the subject's response to therapy as not clinically adequate to relieve one or more symptoms associated the disease or disorder. The terms "subject" and "patient" are used interchangeably in this invention disclosure. The terms refer to an animal, preferably a mammal including a non-primate (e.g., a cow, pig, horse, cat, dog, rat, and mouse) and a primate (e.g., a chimpanzee, a monkey such as a cynomolgous monkey and a human), and more preferably a human. In a preferred embodiment, the subject is a human.

[0062] The terms "prevent", "preventing" and "prevention" refer to the prevention of the recurrence, development, progression or onset of one or more symptoms of the disease or disorder resulting from the administration of one or

more compounds identified in accordance the methods of the invention or the administration of a combination of such a compound and a known therapy for a disease or disorder.

[0063] Preferably, the administration of a composition according to the methods described herein in combination with one or more additional therapies provides a synergistic response in the subject having a disease or disorder. In this context, the term "synergistic" refers to the efficacy of the combination being more effective than the additive effects of either single therapy alone. In one embodiment, the synergistic effect of combination rapamycin therapy according to the invention permits the use of lower dosages and/or less frequent administration of at least one therapy in the combination compared to its dose and/or frequency outside of the combination. In another embodiment, the synergistic effect is manifested in the avoidance or reduction of adverse or unwanted side effects associated with the use of either therapy in the combination alone.

[0064] In the context of the pharmaceutical compositions of the invention, a "carrier" refers to, for example, a solid material such as a diluent, stabilizer, adjuvant, excipient, auxiliary agent, or vehicle with which rapamycin is formulated for delivery. Examples of pharmaceutically acceptable carriers for use in the compositions of the invention include, without limitation, dry powder carriers such as lactose, mannose, amino acids, cyclodextrin, dipalmitoylphosphatidylcholine, detergents, suspending agents, carbohydrates (e.g., glucose, lactose, sucrose or dextran), antioxidants (e.g., ascorbic acid or glutathione), chelating agents, low molecular weight proteins, or suitable mixtures thereof. Preferably, in the context of the dry powder aerosol formulations of rapamycin, the carrier, if present, is selected from the group consisting of a saccharide and a sugar alcohol. In one embodiment, the carrier, if present, is lactose.


[0065] The term "pharmaceutically acceptable" indicates approval by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia such as the European Pharmacopeia, for use in animals, and more particularly in humans. One method for solubilizing poorly water soluble or water insoluble drugs is to form a salt of the drug or to prepare a prodrug that is more soluble itself or that can be used to form a water soluble salt of the prodrug. Methods for forming salts and pharmaceutically acceptable salt forms are known in the art and include, without limitation, salts of acidic or basic groups that may be present in the drug or prodrug of interest. Compounds that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that can be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds are those that form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, including but not limited to sulfuric, citric, maleic, acetic, oxalic, hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate (i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. Compounds that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include alkali metal or alkaline earth metal salts and, particularly, calcium, magnesium, sodium lithium, zinc, potassium, and iron salts.

[0066] In one embodiment not according to the invention, the methods and compositions of the invention utilize a water soluble rapamycin, preferably temsirolimus or related compound. In one embodiment, the methods and compositions of the invention utilize rapamycin (sirolimus).

Rapamycin

[0067] Rapamycin is a macrocyclic lactone produced by *Streptomyces hygroscopicus*. Its chemical (IUPAC) name is (3S,6R,7E,9R,10R,12R,14S,15E,17E,19E,21S,23S,26R,27R,34aS)-9,10,12,13, -14,21,22,23,24,25,26,27,32,33,34,34a-hexadecahydro-9,27-dihydroxy-3-[(1R)-2-[(1S,3R,4R)-4-hydroxy-3-methoxycyclohexyl]-1-methylethyl]-10,21-dimethoxy-6,8,12,14,20,26-hexamethyl-23,27-epoxy-3H-pyrido[2,1-c][1,4]oxaazacyclohentacontine-1,5,11,28,29(4H,6H,31H)-pentone.


[0068] Its molecular formula is $C_{51}H_{79}NO_{13}$ and its molecular weight is 914.172 g/mol. Its structure is shown below. Isomers of rapamycin are known, e.g., isomer B and isomer C, having structures as shown in U.S. Patent No. 7,384,953. Typically, rapamycin is a mixture of the B and C isomers. In solution, rapamycin isomers B and C inter convert and an equilibrium is achieved. It is common practice in the literature to depict the structure of rapamycin in the form of the B isomer, which is the form shown below.




[0069] Rapamycin is a white to off-white powder and is considered insoluble in water, having a very low solubility of only 2.6 $\mu\text{g/ml}$. It is freely soluble in benzyl alcohol, chloroform, acetone, and acetonitrile. The water insolubility of rapamycin presents special technical problems to its formulation. In the context of its formulation as an oral dosage form, it has been prepared as an oral solution in the form of a solid dispersion (WO 97/03654) and a tablet containing nanosized (less than 400 nm) particles (US 5,989,591). But these procedures suffer from substantial variation in the dissolution of the active and therefore its bioavailability. Another method of formulation utilizes the crystalline powder. According to art-recognized methods, the transformation of the crystalline form of a low solubility drug to its amorphous form can significantly increase its solubility. While this is also true for rapamycin, the amorphous form is extremely chemically unstable. Pharmaceutical dosage forms comprising amorphous rapamycin (sirolimus) are described in WO 06/039237 and WO 06/094507 (modified release formulation comprising sirolimus and glyceryl monostearate at a concentration of 49.25%). An improved stable oral dosage form of rapamycin is described in US 8,053,444. The dosage form employs a fatty acid ester and a polymer (e.g., polyvinylpyrrolidone (PVP), hydroxypropylcellulose (HPC) or hydroxypropylmethylcellulose (HPMC)) in the composition to increase the stability of sirolimus without adversely affecting its release rate. According to US 8,053,444, a fatty acid ester concentration exceeding 10% w/w suppresses the release rate of sirolimus from the formulation and so should be avoided because it can lead to insufficient absorption from the gastrointestinal tract. The preferred concentration of fatty acid ester (glycerol ester) is 1% to 5% or 5% to 9%. In one embodiment, the aerosol rapamycin compositions of the present invention do not contain a fatty acid ester in combination with a polymer. In one embodiment, the aerosol rapamycin compositions of the invention contain a fatty acid ester at a concentration exceeding 10% or exceeding 12% by weight of the composition.

[0070] Rapamycin suitable for use in the compositions and methods of the invention includes rapamycin (sirolimus) which are inhibitors of the mTOR cellular signaling pathway, and preferably inhibitors of mTOR itself.


[0071] The compound for use in the aerosol formulations and methods described herein is a rapamycin (analogue) selected from the group consisting of analogues everolimus, temsirolimus, ridaforolimus, umirolimus, and zotarolimus. These analogues are not according to the invention. The chemical structures of the rapamycin analogues everolimus, temsirolimus, ridaforolimus, umirolimus, and zotarolimus are shown below.




Everolimus (Affinitor)




Temsirolimus (CCI-779)



Ridaforolimus (MK-8669)



Umirolimus



Zotarolimus (ABT-578)

[0072] Particularly preferred for use in the compositions of the invention is sirolimus. Not according to the invention

are temsirolimus and everolimus. In one embodiment, the compound for use in the aerosol formulations and methods of the invention is selected from the group consisting of sirolimus.

Compositions for Inhalation

[0073] The invention provides pharmaceutical compositions adapted for administration by inhalation comprising rapamycin, in the form of an a dry powder.

[0074] The compositions of the invention are aerosolizable formulations of rapamycin suitable for pulmonary drug delivery in a human subject by inhalation of the aerosol. The term "aerosol" is used in this context to mean a colloidal system in which the dispersed phase is composed of solid particles and in which the dispersal medium is a gas. In one embodiment, the gas is air and the formulation is a solution formulation suitable for administration via a nebulizer or a dry powder formulation suitable for administration via dry powder inhaler device. Generally, respirable particles will have a mean diameter in the range of 0.10 to 10 μm (microns). The size of the particles is selected to maximize targeted delivery either to the lungs themselves (i.e., where the lung is the target tissue) or systemically (where the lungs are utilized as an alternative route for systemic administration, which is described for reference only). The size will preferably be in the range of about 0.5 to 5 μm (microns) where the lung itself is the therapeutic target, or less than 3 μm (microns) for systemic delivery via the lung. Size is measured according to methods known in the art and described, for example, in the U.S. Pharmacopeia at Chapters 905 and 601. For example, it is measured as Mass Median Aerodynamic Diameter (MMAD). The average or mean diameter of the particles comprising the compositions described herein is measured as MMAD.

[0075] In one embodiment, the dispersed phase of the aerosol is composed of solid particles. In this embodiment, the formulation of the invention is a dry powder formulation. Micronized particles of this size can be produced by methods known in the art, for example by mechanical grinding (milling), precipitation from subcritical or supercritical solutions, spray-drying, freeze-drying, or lyophilization.

[0076] Generally, inhaled particles are subject to deposition by one of two mechanisms: impaction, which usually predominates for larger particles, and sedimentation, which is prevalent for smaller particles. Impaction occurs when the momentum of an inhaled particle is large enough that the particle does not follow the air stream and encounters a physiological surface. In contrast, sedimentation occurs primarily in the deep lung when very small particles which have traveled with the inhaled air stream encounter physiological surfaces as a result of random diffusion within the air stream. The aerosol formulations of the invention are preferably adapted to maximize their deposition either by impaction (in the upper airways) or by sedimentation (in the alveoli), in order to achieve the desired therapeutic efficacy.

[0077] The amount of drug delivered to the patient from a delivery device, such as a nebulizer, pMDI or DPI device, is referred to as the delivered dose. It can be estimated in vitro by determining the amount of drug emitted from the delivery device in a simulated inhalation maneuver. This is termed emitted dose (ED) as measured according to methods known in the art, for examples those set out in the U.S. and European Pharmacopeias, e.g., at Chapter 601 and Chapter 905 of the USP. Accordingly, "emitted dose" is considered equivalent to the delivered dose.

[0078] The amount of drug delivered from the delivery device to the lungs of the patient is termed the respirable dose. It can be estimated in vitro by determining the fine particle dose (FPD) as measured using cascade impactors, such as a Next Generation Impactor (NGI) according to methods known in the art, for examples those set out in the U.S. and European Pharmacopeias, e.g., at Chapters 601 and 905 of the USP.

[0079] The amount of drug released in fine, inhalable particles from a delivery device is referred to as the fine particle fraction (FPF) of the formulation. FPF is the fraction of drug in the delivered dose that is potentially respirable. Thus, FPF is the ratio of FPD to ED (emitted, or delivered dose). These characteristics of the formulation are measured according to methods known in the art, for examples those set out in the U.S. and European Pharmacopeias, e.g., at Chapter 601 of the USP and monograph 2.9.18 of the Pharm Europa.

[0080] In one embodiment, the aerosolizable rapamycin formulations of the present invention have an FPF greater than 20% with a corresponding FPD ranging from 10 micrograms to 2 milligrams, preferably less than 0.5 milligrams, even after prolonged storage, e.g., after 1 to 12 months or after 1 to 36 months of storage. In one embodiment the dose delivered to the patient, the delivered dose (DD) or emitted dose (ED), ranges from 25 micrograms to 2.5 milligrams, preferably less than 0.5 milligrams.

[0081] In certain embodiments not according to the invention the rapamycin is encapsulated in a pharmaceutically acceptable compound, material, or matrix. In one embodiment not according to the invention, the rapamycin is encapsulated in a liposomal formulation or non-liposomal formulation.

Dry Powder Compositions

[0082] The aerosolizable composition of the invention is a dry powder comprising micronized particles of rapamycin as the therapeutic agent (also referred to as "drug"), the particles may have a mean diameter of between about 0.5 to

4.5 μm (microns), about 1 to 4 μm (microns), about 1 to 3.5 μm (microns), about 1.5 to 3.5 μm (microns), or about 2 to 3 μm (microns). The dry powder formulation is suitable for use in a dry powder inhaler device (DPI). The amount of rapamycin in the dry powder is from about 0.5 to 20% (w/w) based on total weight of the powder. In one embodiment, the amount of rapamycin is about 1% or 2% (w/w).

[0083] In one embodiment, micronized rapamycin is produced by wet polishing or jet milling as described below to generate diameters in the range of about 0.5 to 4.5 μm (microns), about 1 to 4 μm (microns), or about 2 to 3 μm (microns), and the rapamycin particles are blended onto lactose carrier particles in a drug/carrier ratio ranging from 0.5-2% w/w with a preferred ratio of 1%.

[0084] In one embodiment, the drug particles are lightly compacted into a frangible matrix which is contained within the delivery device (a dry powder inhaler). Upon actuation the delivery device abrades a portion of the drug particles from the matrix, and disperses them in the inspiratory breath delivering the drug particles to the respiratory tract. Alternatively the drug particles may be a free flowing powder contained within a reservoir in the delivery device (a dry powder inhaler). The reservoir can be an integral chamber within the device, or a capsule, blister or similar preformed reservoir that is inserted into the device prior to actuation. Upon actuation the device dispersed a portion of the drug particles from the reservoir and disperses them in the inhalation breath delivering the drug particles to the respiratory tract.

[0085] In one embodiment, the dry powder composition consists of drug particles and a carrier selected from the group consisting of arabinose, glucose, fructose, ribose, mannose, sucrose, trehalose, lactose, maltose, starches, dextran, mannitol, leucine, lysine, isoleucine, dipalmitoylphosphatidylcholine, lecithin, polylactic acid, poly (lactic-co-glutamic) acid, and xylitol, or mixtures of any of the foregoing. In one embodiment, the carrier is lactose, particularly in the form of the monohydrate. In one embodiment, the dry powder composition comprises a blend of two or more carriers.

[0086] In one embodiment the dry powder composition comprises drug and a blend of at least two different carriers. In one embodiment, the drug to carrier ratio is in the range of from about 0.5 to 20% (w/w). In one embodiment, the drug particles have a mean diameter of about 1 to 4, 1 to 3.5, or 1.5 to 3.5, or 2 to 3 μm (microns). The carrier particles may have diameters ranging from 2 to 200 μm (microns).

[0087] In one embodiment, the composition is contained in a blister pack or a reservoir of a DPI device. In one embodiment, the dry powder composition is preloaded into a gelatin, starch, cellulosic, or polymeric capsule, or a foil/foil or foil/plastic blister suitable for use in a DPI device. Each capsule or blister may contain from 1 to 100 milligrams of the dry powder composition. The capsules or blisters may be inserted into a dry powder inhaler (DPI) device such as Aerolizer®, Plastiap® RS01 Model 7, and Plastiap® RS00 Model 8, XCaps®, FlowCaps®, Arcus®, Diskhaler® or Microdose®. Upon actuating the DPI device, the capsules or blisters are ruptured and the powder is dispersed in the inspiratory breath, delivering the drug to the respiratory tract.

[0088] In one embodiment, the dry powder composition is contained in a dry powder inhaler (DPI) device selected from Accuhaler®, Conix™, Rotahaler®, TwinCaps®, XCaps®, FlowCaps®, Turbuhaler®, NextHaler®, CycloHaler®, Revolizer™, Diskhaler®, Diskus®, Spinhaler, Handihaler®, Microdose Inhaler, GyroHaler®, OmniHaler®, Clickhaler®, Duohaler® (Vectura), and ARCUS® inhaler (Civitas Therapeutics). In one embodiment, the invention provides a DPI device containing a dry powder composition described herein. In one embodiment the device is selected from the group consisting of XCaps, FlowCaps, Handihaler, TwinCaps, Aerolizer®, Plastiap® RS01 Model 7, and Plastiap® RS00 Model 8. In one embodiment, the device containing the composition is selected from the group consisting of a GyroHaler®, an OmniHaler®, a Clickhaler®, a Duohaler®, and an ARCUS® inhaler.

[0089] The carrier particles are preferably of larger size (greater than 5 μm (microns)) so as to avoid deposition of the carrier material in the deep lung. In one embodiment, the carrier particles have diameters ranging from 1 to 200 μm (microns), from 30 to 100 μm (microns), or less than 10 μm (microns). In one embodiment the carrier particles are a blend of two carriers, one with particles of about 30-100 μm (microns) and the other with particles less than 10 μm (microns). The ratio of the two different carriers is in the range of from 3:97 to 97:3. In one embodiment, the dry powder composition consists of 0.5 -20% (w/w) drug to carrier ratio, the drug particles having diameters from 0.1 to 10 μm (microns) with a mean diameter less than 3.5 μm (microns). In one embodiment, the carrier material is a crystalline carrier material. Preferably, the crystalline carrier material is one which is at least 90%, preferably greater than 95% crystalline and in which no or substantially no water is absorbed by the carrier under conditions of 80% or lower relative humidity at room temperature. Examples of such crystalline carriers are lactose monohydrate and glucose monohydrate. The amount of carrier is from 1 to 99.0 % or more of the formulation by dry weight of the powder, preferably 5 to 99%, 10 to 99%, 20 to 99%, 30 to 99%, 40 to 99%, or 50 to 99%.

[0090] In one embodiment, the dry powder composition is contained within a reservoir in the delivery device (a dry powder inhaler). The reservoir can be an integral chamber within the device, or a capsule, blister or similar preformed reservoir that is inserted into the device prior to actuation. Upon actuation the device dispersed a portion of the drug particles from the reservoir and disperses them in the inspiratory breath delivering the drug particles to the respiratory tract.

[0091] In one embodiment, drug is present as a fine powder with a pharmaceutically acceptable carrier. In the context, the term "fine" refers to a particle size in the inhalable range, as discussed above. The drug is micronized and the MMAD of the particles of rapamycin in the dry powder composition is from 0.5 to 5 μm (microns), from 1 to 4 μm (microns),

from 1 to 3 μm (microns), or from 2 to 3 μm (microns). The MMAD or Dv50 value is the particle size below which 50% of the volume of the population occurs.

[0092] In one embodiment, the dry powder formulation of rapamycin further comprises one or more additives selected from the additives described below. In one embodiment, the one or more additives comprises or consists of magnesium stearate. In one aspect of this embodiment, the magnesium stearate is present in amounts of 0.001 to 10% by dry weight of the powder, preferably in amounts of from 0.01 to 5% or 0.01 to 2%. In another embodiment, the additive comprises or consists of a phospholipid, such as lecithin (which is a mixture of phosphatidylcholines) in an amount of 0.1% to 1% by dry weight of the powder, preferably 0.2% to 0.6%. In one aspect of this embodiment, the additive is coated onto the carrier material prior to or simultaneously with a step of blending the carrier with the particles of rapamycin. This can be accomplished, for example, by utilizing a high energy mixing step to coat the carrier with the additive, or a long duration of low energy mixing, or a combination of low and high energy mixing to achieve the desired level of coated carrier material. Low energy devices for mixing dry powders to form blends are known in the art and include, for example, V-blenders, double cone blenders, slant cone blenders, cube blenders, bin blenders, horizontal or vertical drum blenders, static continuous blenders, and dynamic continuous blenders. Other, higher energy devices include high shear mixers known to those skilled in the art.

[0093] In certain embodiments, the dry powder is contained in a capsule. In one embodiment the capsule is a gelatin capsule, a plastic capsule, or a cellulosic capsule, or is in the form of a foil/foil or foil/plastic blisters. In each instance, the capsule or blister is suitable for use in a DPI device, preferably in dosage units together with the carrier in amounts to bring the total weight of powder in each capsule to from 1 mg to 100 mg. Alternatively, the dry powder may be contained in a reservoir of a multi-dose DPI device.

[0094] The particle size of the rapamycin can be reduced to the desired microparticulate level by conventional methods, for example by grinding in an air-jet mill, ball mill or vibrator mill, by wet polishing, microprecipitation, spray drying, lyophilization or recrystallization from subcritical or supercritical solutions. Jet milling or grinding in this context refers to micronization of dry drug particles by mechanical means. Micronization techniques do not require making a solution, slurry, or suspension of the drug. Instead, the drug particles are mechanically reduced in size. Due to the relatively high energy that is employed by micronization, in certain embodiments it is desirable to include a carrier material in a micronized mixture with the rapamycin. In this context, the carrier material absorbs some of the energy of micronization which otherwise could adversely affect the structure of the rapamycin. In one embodiment, rapamycin particles in a size range of from 1 to 4 or from 2 to 3 μm (microns) are produced by a jet milling method.

[0095] Wet polishing as described in US2013/0203717 involves using high shear to reduce the particle size of the drug particles in a suspension or slurry. Wet polishing can include just the drug particles or additional particulates termed milling media. In one embodiment, the particle size of the rapamycin can be reduced to the desired level using a wet polishing process, which comprises wet milling, specifically by cavitation at elevated pressure, where rapamycin is suspended in water or other solvent where it is insoluble, and then is followed by spray drying of the suspension to obtain rapamycin as a dry powder. In one embodiment, rapamycin particles in a size range of from 1 to 4 or from 2 to 3 μm (microns) are produced by a wet polishing method that comprises preparing a suspension of rapamycin, subjecting the suspension to microfluidization, and spray-drying the resulting particles to form a dry powder. The rapamycin may be suspended in an anti-solvent selected from the group consisting of propyl or butyl alcohol, water, and ethyl acetate. In one embodiment, the suspension is an aqueous suspension.

[0096] Spray drying generally involves making a solution, slurry, or suspension of the drug, atomizing the solution, slurry, or suspension, to form particles and then evaporating the solution, slurry, or suspension media to form the particles. The solution, slurry or suspension, can be formed under subcritical or supercritical conditions. The evaporation step can be accomplished by elevating the temperature of the atmosphere into which the atomization occurs, or by decreasing the pressure, or a combination of both. In one embodiment, the powder formulation comprising rapamycin is made by spray drying an aqueous dispersion of rapamycin to form a dry powder consisting of aggregated particles of rapamycin having a size suitable for pulmonary delivery, as described above. The aggregate particle size can be adjusted (increased or decreased) to target either the deep lung or upper respiratory sites, such as the upper bronchial region or nasal mucosa. This can be accomplished, for example, by increasing the concentration of rapamycin in the spray-dried dispersion or by increasing the droplet size generated by the spray dryer.

[0097] A dry powder can be made by freeze-drying (lyophilization) the aqueous drug solution, dispersion, or emulsion, or by a combination of spray-drying and freeze-drying.

[0098] In one embodiment not according to the invention, the aqueous dispersion of rapamycin and the one or more optional additives further comprises a dissolved diluent such as lactose or mannitol such that when the dispersion is freeze-dried, respirable diluent particles, each containing at least one embedded drug particle and additive particle, if present, are formed.

[0099] In one embodiment not according to the invention, the dry powder formulation is made by freeze-drying an aqueous dispersion of rapamycin, and one or more optional additives. In one embodiment, the powders contain aggregates of rapamycin and an additive, if present, wherein the aggregates are within a respirable size range as described

above.

[0100] In one embodiment not according to the invention, the dry powder comprises rapamycin loaded liposomes. Drug-loaded liposomes can be produced by methods known in the art, for example using the technique described for tacrolimus in M. Chougale, et al. Int. J. Nanomedicine 2:625-688 (2007). Briefly, rapamycin, hydrogenated phosphatidylcholine (HSPC), and cholesterol are dissolved in a mixture of methanol and chloroform and then subjected to dry thin film formation, e.g., in Rotaevaporator. The liposomes are hydrated and the liposomal dispersion is passed through a high-pressure homogenizer for size reduction. The resultant pellets are characterized for vesicle size and percent drug entrapment and pellets equivalent to the desired amount of rapamycin are then dispersed in a suitable medium and subjected to spray-drying to obtain particles of the desired size for inhalation. The spray dried powder can be filled into capsules, canisters, or blister packs for administration.

[0101] In one embodiment not according to the invention the dry powder particles can be produced by precipitation from a supercritical or subcritical solution.

[0102] The dry powder compositions may be contained in a suitable dry powder inhaler device, or in a capsule or blister for use in such a device. Examples of such devices are provided above and include Accuhaler®, Aerolizer®, the Plastiaple® RS01 Model 7, the Plastiaple® RS00 Model 8, Conix™, Rotahaler®, TwinCaps®, XCaps®, FlowCaps®, Turbuhaler®, NextHaler®, CycloHaler®, Revolizer™, Diskhaler®, Diskus®, Spinhaler, Handihaler®, Microdose Inhaler, GyroHaler®, OmniHaler®, Clickhaler®, or Duohaler® (Vectura), or a breath-actuated ARCUS® inhaler (Civitas Therapeutics). In one embodiment not according to the invention, the invention provides a DPI device containing a dry powder composition described herein. In one embodiment the device is selected from the group consisting of XCaps, FlowCaps, Handihaler, TwinCaps, Aerolizer®, the Plastiaple® RS01 Model 7, and the Plastiaple® RS00 Model 8.

Propellant-Based Formulations not according to the invention

[0103] In another embodiment, the rapamycin is formulated in a propellant-based formulation which may also be referred to generically herein as "a pMDI formulation". A pMDI formulation is suitable for delivery by a device such as a pressurized metered dose inhaler (pMDI). In one embodiment, the composition comprises rapamycin, a propellant, and a vegetable oil or pharmaceutically acceptable derivative of a vegetable oil. The propellant is preferably selected from 1,1,1,2-tetrafluoroethane (HFA134a) and 1,1,1,2,3,3,3-heptafluoropropane (HFA227), or mixtures thereof. In one embodiment, the vegetable oil is selected from olive oil, safflower oil, and soybean oil. The rapamycin may be in solution or in suspension in the propellant. In this context, "in suspension" refers to where the rapamycin is present in particulate form dispersed in the propellant. In one embodiment, the rapamycin is micronized and is present in suspension in the propellant. In one embodiment, the formulation further comprises a wetting agent or co-solvent such as ethanol. In one embodiment, the formulation further comprises a polyhydroxy alcohol such as propylene glycol.

[0104] Suitable propellants are known in the art and include, for example, halogen-substituted hydrocarbons, for example fluorine-substituted methanes, ethanes, propanes, butanes, cyclopropanes or cyclobutanes, particularly 1,1,1,2-tetrafluoroethane (HFA134a) and 1,1,1,2,3,3,3-heptafluoropropane (HFA227), or mixtures thereof.

[0105] In one embodiment, the formulation comprises micronized rapamycin, ethanol, a suitable propellant such as HFA 134a, HFA 227, or a mixture of suitable propellants, and optionally one or more surfactants. In one embodiment, the formulation further comprises a lubricant.

[0106] In one embodiment, the formulation comprises rapamycin, a propellant, and a vegetable oil. In one aspect, the formulation does not comprise an additive or surfactant. For example, the formulation does not comprise ethanol, a polyhydroxy alcohol (e.g., propylene glycol), or a surfactant (e.g., sorbitan trioleate, sorbitan monooleate, or oleic acid).

[0107] In one embodiment, the propellant-based formulation comprises compressed air, carbon dioxide, nitrogen or a liquefied propellant selected from the group consisting of n-propane, n-butane, isobutane or mixtures thereof, or 1,1,1,2-tetrafluoroethane (HFA134a) and 1,1,1,2,3,3,3-heptafluoropropane (HFA227), or mixtures thereof, with or without a polar co-solvent such as an alcohol. The composition can be a solution or a suspension. For suspensions the drug particles have diameters from 0.1 to 10 microns with a mean diameter less than 3.5 microns.

[0108] The propellant-based formulation is prepared by methods known in the art, for example by wet milling the coarse rapamycin, and optional additive, in liquid propellant, either at ambient pressure or under high pressure conditions. In certain embodiments, the additive is a surfactant which serves to prevent aggregation (caking or crystallization), to facilitate uniform dosing, and (or alternatively) to provide a favorable fine particle fraction (FPF). In one aspect, the surfactant is selected from sorbitan trioleate, sorbitan monooleate, or oleic acid. Alternatively, dry powders containing drug particles are prepared by spray-drying or freeze-drying aqueous dispersions of the drug particles as discussed above and the resultant powders dispersed into suitable propellants for use in conventional pressurized metered dose inhalers (pMDIs). In one embodiment, the inhalation device is a RespiMat™.

[0109] In one embodiment, the propellant-based aerosol rapamycin formulations of the invention are stable against particle size growth or change in the crystal morphology of the rapamycin over prolonged periods of time.

Process for Manufacturing Sterile Unit Dose Forms

[0110] In one embodiment, the compositions of the invention are sterile compositions. In one embodiment, the sterile compositions are sterile unit dose forms. In one embodiment, the sterile unit dosage form is a capsule suitable for use in a nebulizer device.

[0111] In one embodiment not according to the invention, the finished composition is sterilized in its container-closure by heat, e.g., autoclaving, or by radiation. In one embodiment not according to the invention, the component parts of the composition are first sterilized by a suitable process including sterile filtration for liquid components and radiation or autoclaving for solids or liquids, the process further comprising maintaining the sterility of the sterile components by packaging in hermetic containers, combining the components in a mixing vessel in the appropriate proportions, and filling the resulting product into a container closure, all performed in an aseptic suite. This process has the disadvantage of being expensive and requiring difficult aseptic handling techniques. Accordingly, it is used primarily to process particulate suspensions or colloidal dispersions, liposomal formulations, or emulsions, which cannot be passed through a sub- μm (submicron) filter for sterilization. Finally, in one embodiment not according to the invention, the finished composition is sterile filtered through a sub- μm (submicron) filter, preferably a 0.2 μm (micron) filter. In one embodiment not according to the invention, the compositions are single-phase aqueous solutions sterilized via a filtration sterilization process. In contrast, emulsions and liposomal formulations are typically not sufficiently stable under the high shear conditions of a filtration sterilization process and so are not preferred for this process. In one embodiment not according to the invention, the compositions of the invention are single-phase aqueous solutions which are filled into a container-closure, e.g., a vial, formed of a polymer, preferably polyethylene, or alternatively a glass vial. Autoclaving and radiation are not suitable where the vial is a polymer vial because of the high likelihood of creating chemical instability in the drug and/or formulation excipients, as well as in the container, and due to the generation of undesirable impurities. In one embodiment not according to the invention, the compositions of the invention are sterilized by a process that does not include heat (autoclaving) or radiation, and instead includes a filtration sterilization process. Preferably, in accordance with this embodiment, the single-phase aqueous solutions of rapamycin are sterilized by filtration through a filter having a pore size less than or equal to 0.2 μm (microns). In one embodiment not according to the invention, the sterile filtrate is collected in a collection vessel located in an aseptic suite. In one embodiment not according to the invention, the sterile filtrate is transferred from the collection vessel into a container closure in an aseptic suite. Preferably the container closure is a polymer vial, preferably a unit dose vial, and most preferably a polyethylene unit dose vial. In one embodiment, the polymer vial is formed by blowmolding immediately before it is filled and then thermally sealed immediately after filling. This technique may be also referred to as "form-fill-seal" or a "blow-fill". This technique is particularly advantageous in the context of the compositions of the invention which are single-phase aqueous solutions of rapamycin because this process does not require heat or radiation, both of which may degrade either the drug itself, the formulation excipients, or the container closure.

Pulmonary Administration and Dosing

[0112] The present invention provides compositions for use in methods for the treatment and prophylaxis of age-related diseases and disorders in a human subject, wherein the age-related disorder is chronic lung disease, by administering rapamycin to the lungs, by inhalation. Pulmonary delivery is preferably accomplished by inhalation of the aerosol through the mouth and throat into the lungs, but may also be accomplished by inhalation of the aerosol through the nose. Thus, in one embodiment the composition is delivered intranasally.

[0113] The compositions and compositions for use in methods of the invention advantageously provide for the targeted delivery of a therapeutically effective amount of rapamycin to the lungs while simultaneously reducing to very low or undetectable levels the amount of rapamycin in the blood and available systemically. In one embodiment, the amount of rapamycin in a single dose of a dry powder composition described herein is from about 5 to 500 micrograms or from about 100 to 300 micrograms, or from about 50 to 250 micrograms. The targeted delivery of low dose rapamycin directly to the lungs while minimizing systemic exposure provides for an improved therapeutic index compared to oral dosage forms.

[0114] In one embodiment, administration of rapamycin by inhalation according to the methods of the invention increases the therapeutic index of rapamycin. In this context, as applied to human subjects, the therapeutic index is a ratio that compares the dose that produces a therapeutic effect (ED_{50}) to the dose that produces a toxicity (TD_{50}) in 50% of the population. The ratio is represented as $\text{TD}_{50}/\text{ED}_{50}$. In one embodiment, administration of rapamycin by inhalation according to the methods of the invention reduces one or more toxicities associated with orally administered rapamycin, thereby increasing the therapeutic index of rapamycin.

[0115] The invention includes aerosolizable formulations in the form of powders. Accordingly, the rapamycin is administered according to the methods of the invention in the form of a dry powder aerosol.

[0116] In one embodiment, the administered dose of rapamycin produces a blood trough level in the subject of from

of from 0.01 to 0.15 ng/ml, from 0.075 to 0.350 ng/ml, from 0.150 to 0.750 ng/ml, from 0.750 to 1.5 ng/ml or from 1.5 to 5 ng/ml. In one embodiment, the administered dose of rapamycin produces a blood trough level in the subject of less than 5 ng/ml, less than 2 ng/ml, less than 1 ng/ml, or less than 0.5 ng/ml.

[0117] In one embodiment, the administered dose of rapamycin is sufficient to produce a concentration of rapamycin in lung tissue in the range of from 1 ng/g to 1 ug/g, preferably from about 5 ng/g to 100 ng/g, from about 5 ng/g to about 20 ng/g, or from about 5 ng/g to about 30 ng/g.

[0118] In one embodiment, the administered dose of rapamycin is from 5 to 100 micrograms, from 20 to 100 micrograms, from 20 to 250 micrograms, from 50 to 500 micrograms (0.05 to 0.5 milligrams), from 250 to 1000 micrograms (0.25 to 1 milligrams) or from 500 to 2000 micrograms (0.5 to 2 milligrams). In one embodiment, the amount of rapamycin administered is less than 500 micrograms, less than 100 micrograms, less than 50 micrograms, less than 20 micrograms, or less than 10 micrograms. Preferably, the amount of rapamycin administered is less than 0.5 milligrams or less than 0.25 milligrams.

[0119] In one embodiment, the rapamycin is administered once daily.

[0120] In one embodiment, the total daily dose of rapamycin is in the range of from 5 to 100 micrograms, from 20 to 250 micrograms, from 50 to 500 micrograms (0.05 to 0.5 milligrams), from 250 to 1000 micrograms (0.5 to 1 milligrams) or from 500 to 2000 micrograms (0.5 to 2 milligrams). In one embodiment, the total daily dose of rapamycin is less than 500 micrograms, less than 100 micrograms, less than 50 micrograms, less than 20 micrograms, or less than 10 micrograms. In one embodiment, the total daily dose of rapamycin administered to the subject is less than 0.5 milligrams or less than 0.25 milligrams per day.

[0121] In one embodiment, a composition of the invention is administered once per day to the subject. In one embodiment, a composition of the invention is administered twice or three times a day. Preferably, the composition is administered once or twice daily, or less than once daily.

[0122] In one embodiment, the methods of the invention comprise administering rapamycin via a pulmonary route in combination with one or more additional therapeutic agents. The one or more additional agents may be administered by the same or a different route of administration as the rapamycin. For example, the agent may be administered by inhalation, intranasally, orally or intravenously.

[0123] In one embodiment, the methods of the invention comprise administering rapamycin via a pulmonary route in combination with one or more additional therapies.

[0124] In certain embodiments, the methods include pulmonary administration of a composition of the invention as the primary therapy. In other embodiments, the administration of a composition of the invention is an adjuvant therapy. In either case, the methods of the invention contemplate the administration of a composition of the invention in combination with one or more additional therapies for the treatment of a disease or disorder. The terms "therapy" and "therapies" refer to any method, protocol and/or agent that can be used in the prevention, treatment, management or amelioration of a disease or disorder, or one or more symptoms thereof.

[0125] Preferably, the administration of a pharmaceutical composition comprising rapamycin according to the methods of the invention in combination with one or more additional therapies provides a synergistic response in the subject. In this context, the term "synergistic" refers to the efficacy of the combination being more effective than the additive effects of either single therapy alone. In one embodiment, the synergistic effect of combination rapamycin therapy according to the invention permits the use of lower dosages and/or less frequent administration of at least one therapy in the combination compared to its dose and/or frequency outside of the combination. In another embodiment, the synergistic effect is manifested in the avoidance or reduction of adverse or unwanted side effects associated with the use of either therapy in the combination alone.

Dry Powder Delivery

[0126] In one embodiment, the dry powder compositions of the invention are delivered by a non-propellant based dry powder inhaler (DPI) device. In one embodiment, the powder is contained in capsules of gelatin or plastic, or in blisters, suitable for use in a DPI device. In one embodiment, the powder is supplied in unit dosage form and in dosage units of from 5 mg to 100 mg of powder per capsule. In another embodiment, the dry powder is contained in a reservoir of a multi-dose dry powder inhalation device. In one embodiment, the inhaler device comprises an aerosol vial provided with a valve adapted to deliver a metered dose, such as 10 to 100 μ l, e.g. 25 to 50 μ l, of the composition, i.e. a device known as a metered dose inhaler.

[0127] In one embodiment, the DPI device is a blister based device such as the GyroHaler® or the OmniHaler® (both from Vectura), a reservoir based device such as the Clickhaler® or Duohaler® (Vectura), and the ARCUS® inhaler (Civitas Therapeutics). In one embodiment, the DPI device is selected from Pulmatrix™, and Hovione Twincaps and XCaps™. In one embodiment the device is selected from the group consisting of XCaps, Plastiap® RS01 Model 7, and Plastiap® RS00 Model 8.

[0128] In one embodiment, the DPI device is selected from the group consisting of Accuhaler®, Aerolizer®, the Plas-

tiap[®] RS01 Model 7, the Plastiap[®] RS00 Model 8, Conix[™], Rotahaler[®], TwinCaps[®], XCaps[®], FlowCaps[®], Turbuhaler[®], NextHaler[®], CycloHaler[®], Revolizer[™], Diskhaler[®], Diskus[®], Spinhaler, Handihaler[®], Microdose Inhaler, GyroHaler[®], OmniHaler[®], Clickhaler[®], or Duohaler[®] (Vectura), or a breath-actuated ARCUS[®] inhaler (Civitas Therapeutics).

[0129] In one embodiment, the DPI device is selected from the group consisting of Arcus[™], Aspi[™], Axahaler[™], Breezhaler[™], Clickhaler[™], Conix Dry[™], Cricket[™], Dreamboat[™], Genuair[™], Gemini[™], Inspiromatic[™], iSPERSE[™], MicroDose[™], Next DPI[™], Prohaler[™], Pulmojet[™], Pulvinal[™], Solis[™], Taifun[™], Taper Dry[™], Trival[™], Novolizer[™], Podhaler[™], Skyehaler[™], Spiromax[™], Twincaps/Flowcaps[™], and Turbuhaler[™]. In one embodiment, the DPI device is adapted to deliver the dry powder from a capsule or blister containing a dosage unit of the dry powder or a multi-dose dry powder inhalation device adapted to deliver, for example, 5-25 mg of dry powder per actuation.

pMDI Delivery not according to the invention

[0130] In another embodiment, the rapamycin is delivered in the form of aerosolized particles from a pressurized container or dispenser that contains a suitable propellant as described above in connection with propellant-based formulations. In one embodiment, the inhaler is a propellant driven inhaler, such as a pMDI device, which releases a metered dose of rapamycin upon each actuation. A typical pMDI device comprises a canister containing drug, a drug metering valve, and a mouthpiece. In one aspect of this embodiment, the rapamycin is formulated as a suspension in the propellant. In the context of this embodiment, the rapamycin is made into a fine powder which is suspended in the liquefied propellant or propellant blend. The suspension is then stored in a sealed canister under sufficient pressure to maintain the propellant in liquid form. In another embodiment, the rapamycin is formulated as a solution. In the context of this embodiment, the rapamycin is solubilized in the liquefied propellant or propellant blend. In one embodiment, the formulation further comprises a stabilizer in an amount suitable to stabilize the formulation against settling, creaming or flocculation for a time sufficient to allow reproducible dosing of the rapamycin after agitation of the formulation. The stabilizer may be present in excess in an amount of about 10 part by weight to about 5000 parts by weight based on one million parts by total weight of the aerosol formulation. In one embodiment, the fluid carrier is 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane or a mixture thereof. In another embodiment, the fluid carrier is a hydrocarbon (e.g., n-butane, propane, isopentane, or a mixture thereof). The composition may further comprise a co-solvent (e.g., ethanol or other suitable co-solvent).

[0131] In one embodiment of the methods of the invention, the aerosol formulation comprising rapamycin further comprises an additional drug.

Additives

[0132] The aerosol compositions of the invention may contain one or more additives in addition to any carrier or diluent (such as lactose or mannitol) that is present in the formulation. In one embodiment, the one or more additives comprises or consists of one or more surfactants. Surfactants typically have one or more long aliphatic chains such as fatty acids which enables them to insert directly into the lipid structures of cells to enhance drug penetration and absorption. An empirical parameter commonly used to characterize the relative hydrophilicity and hydrophobicity of surfactants is the hydrophilic-lipophilic balance ("HLB" value). Surfactants with lower HLB values are more hydrophobic, and have greater solubility in oils, while surfactants with higher HLB values are more hydrophilic, and have greater solubility in aqueous solutions. Thus, hydrophilic surfactants are generally considered to be those compounds having an HLB value greater than about 10, and hydrophobic surfactants are generally those having an HLB value less than about 10. However, these HLB values are merely a guide since for many surfactants, the HLB values can differ by as much as about 8 HLB units, depending upon the empirical method chosen to determine the HLB value.

[0133] Among the surfactants for use in the aerosol compositions of the invention are polyethylene glycol (PEG)-fatty acids and PEG-fatty acid mono and diesters, PEG glycerol esters, alcohol-oil transesterification products, polyglyceryl fatty acids, propylene glycol fatty acid esters, sterol and sterol derivatives, polyethylene glycol sorbitan fatty acid esters, polyethylene glycol alkyl ethers, sugar and its derivatives, polyethylene glycol alkyl phenols, polyoxyethylene-polyoxypropylene (POE-POP) block copolymers, sorbitan fatty acid esters, ionic surfactants, fat-soluble vitamins and their salts, water-soluble vitamins and their amphiphilic derivatives, amino acids and their salts, and organic acids and their esters and anhydrides. Each of these is described in more detail below.

PEG Fatty Acid Esters

[0134] Although polyethylene glycol (PEG) itself does not function as a surfactant, a variety of PEG-fatty acid esters have useful surfactant properties. Among the PEG-fatty acid monoesters, esters of lauric acid, oleic acid, and stearic acid are most useful in embodiments of the present invention. Preferred hydrophilic surfactants include PEG-8 laurate, PEG-8 oleate, PEG-8 stearate, PEG-9 oleate, PEG-10 laurate, PEG-10 oleate, PEG-12 laurate, PEG-12 oleate, PEG-

15 oleate, PEG-20 laurate and PEG-20 oleate. The HLB values are in the range of 4-20.

[0135] Polyethylene glycol fatty acid diesters are also suitable for use as surfactants in the compositions of embodiments of the present invention. Most preferred hydrophilic surfactants include PEG-20 dilaurate, PEG-20 dioleate, PEG-20 distearate, PEG-32 dilaurate and PEG-32 dioleate. The HLB values are in the range of 5-15.

[0136] In general, mixtures of surfactants are also useful in embodiments of the present invention, including mixtures of two or more commercial surfactants as well as mixtures of surfactants with another additive or additives. Several PEG-fatty acid esters are marketed commercially as mixtures or mono- and diesters.

Polyethylene Glycol Glycerol Fatty Acid Esters

[0137] Preferred hydrophilic surfactants are PEG-20 glyceryl laurate, PEG-30 glyceryl laurate, PEG-40 glyceryl laurate, PEG-20 glyceryl oleate, and PEG-30 glyceryl oleate.

Alcohol-Oil Transesterification Products

[0138] A large number of surfactants of different degrees of hydrophobicity or hydrophilicity can be prepared by reaction of alcohols or polyalcohol with a variety of natural and/or hydrogenated oils. Most commonly, the oils used are castor oil or hydrogenated castor oil, or an edible vegetable oil such as corn oil, olive oil, peanut oil, palm kernel oil, apricot kernel oil, or almond oil. Preferred alcohols include glycerol, propylene glycol, ethylene glycol, polyethylene glycol, sorbitol, and pentaerythritol. Among these alcohol-oil transesterified surfactants, preferred hydrophilic surfactants are PEG-3 5 castor oil (Incrocas-35), PEG-40 hydrogenated castor oil (Cremophor RH 40), PEG-25 trioleate (TAGAT.RTM. TO), PEG-60 corn glycerides (Crovol M70), PEG-60 almond oil (Crovol A70), PEG-40 palm kernel oil (Crovol PK70), PEG-50 castor oil (Emalex C-50), PEG-50 hydrogenated castor oil (Emalex HC-50), PEG-8 caprylic/capric glycerides (Labrasol), and PEG-6 caprylic/capric glycerides (Softigen 767). Preferred hydrophobic surfactants in this class include PEG-5 hydrogenated castor oil, PEG-7 hydrogenated castor oil, PEG-9 hydrogenated castor oil, PEG-6 corn oil (Labrafil.RTM. M 2125 CS), PEG-6 almond oil (Labrafil.RTM. M 1966 CS), PEG-6 apricot kernel oil (Labrafil.RTM. M 1944 CS), PEG-6 olive oil (Labrafil.RTM. M 1980 CS), PEG-6 peanut oil (Labrafil.RTM. M 1969 CS), PEG-6 hydrogenated palm kernel oil (Labrafil.RTM. M 2130 BS), PEG-6 palm kernel oil (Labrafil.RTM. M 2130 CS), PEG-6 triolein (Labrafil.RTM. M 2735 CS), PEG-8 corn oil (Labrafil.RTM. WL 2609 BS), PEG-20 corn glycerides (Crovol M40), and PEG-20 almond glycerides (Crovol A40).

Polyglyceryl Fatty Acids

[0139] Polyglycerol esters of fatty acids are also suitable surfactants for use in embodiments of the present invention. Among the polyglyceryl fatty acid esters, preferred hydrophobic surfactants include polyglyceryl oleate (Plurol Oleique), polyglyceryl-2 dioleate (Nikkol DGDO), polyglyceryl-10 trioleate, polyglyceryl stearate, polyglyceryl laurate, polyglyceryl myristate, polyglyceryl palmitate, and polyglyceryl linoleate. Preferred hydrophilic surfactants include polyglyceryl-10 laurate (Nikkol Decaglyn 1-L), polyglyceryl-10 oleate (Nikkol Decaglyn 1-0), and polyglyceryl-10 mono, dioleate (Caprol.RTM. PEG 860), polyglyceryl-10 stearate, polyglyceryl-10 laurate, polyglyceryl-10 myristate, polyglyceryl-10 palmitate, polyglyceryl-10 linoleate, polyglyceryl-6 stearate, polyglyceryl-6 laurate, polyglyceryl-6 myristate, polyglyceryl-6 palmitate, and polyglyceryl-6 linoleate. Polyglyceryl polyricinoleates (Polymuls) are also preferred surfactants.

Propylene Glycol Fatty Acid Esters

[0140] Esters of propylene glycol and fatty acids are suitable surfactants for use in embodiments of the present invention. In this surfactant class, preferred hydrophobic surfactants include propylene glycol monolaurate (Lauroglycol FCC), propylene glycol ricinoleate (Propymuls), propylene glycol monooleate (Myverol P-06), propylene glycol dicaprylate/dicaprate (Captex.RTM. 200), and propylene glycol dioctanoate (Captex.RTM. 800).

Sterol and Sterol Derivatives

[0141] Sterols and derivatives of sterols are suitable surfactants for use in embodiments of the present invention. Preferred derivatives include the polyethylene glycol derivatives. A preferred surfactant in this class is PEG-24 cholesterol ether (Solulan C-24).

Polyethylene Glycol Sorbitan Fatty Acid Esters

[0142] A variety of PEG-sorbitan fatty acid esters are available and are suitable for use as surfactants in embodiments

of the present invention. Among the PEG-sorbitan fatty acid esters, preferred surfactants include PEG-20 sorbitan monolaurate (Tween-20), PEG-20 sorbitan monopalmitate (Tween-40), PEG-20 sorbitan monostearate (Tween-60), and PEG-20 sorbitan monooleate (Tween-80).

5 Polyethylene Glycol Alkyl Ethers

[0143] Ethers of polyethylene glycol and alkyl alcohols are suitable surfactants for use in embodiments of the present invention. Preferred ethers include PEG-3 oleyl ether (Volpo 3) and PEG-4 lauryl ether (Brij 30).

10 Sugar and its Derivatives

[0144] Sugar derivatives are suitable surfactants for use in embodiments of the present invention. Preferred surfactants in this class include sucrose monopalmitate, sucrose monolaurate, decanoyl-N-methylglucamide, n-decyl- β -D-glucopyranoside, n-decyl- β -D-maltopyranoside, n-dodecyl- β -D-glucopyranoside, n-dodecyl- β -D-maltoside, heptanoyl-N-methylglucamide, n-heptyl- β -D-glucopyranoside, n-heptyl- β -D-thioglucoside, n-hexyl- β -D-glucopyranoside, nonanoyl-N-methylglucamide, n-nonyl- β -D-glucopyranoside, octanoyl-N-methylglucamide, n-octyl- β -D-glucopyranoside, and octyl- β -D-thioglucopyranoside.

20 Polyethylene Glycol Alkyl Phenols

[0145] Several PEG-alkyl phenol surfactants are available, such as PEG-10-100 nonyl phenol and PEG-15-100 octyl phenol ether, Tyloxapol, octoxynol, nonoxynol, and are suitable for use in embodiments of the present invention.

25 Polyoxyethylene-Polyoxypropylene (POE-POP) Block Copolymers

[0146] The POE-POP block copolymers are a unique class of polymeric surfactants. The unique structure of the surfactants, with hydrophilic POE and hydrophobic POP moieties in well-defined ratios and positions, provides a wide variety of surfactants suitable for use in embodiments of the present invention. These surfactants are available under various trade names, including Synperonic PE series (ICI); Pluronic.RTM. series (BASF), Emkalix, Lutrol (BASF), Supronic, Monolan, Pluracare, and Plurodac. The generic term for these polymers is "poloxamer" (CAS 9003-11-6). These polymers have the formula: $\text{HO}(\text{C}_2\text{H}_4\text{O})_a(\text{C}_3\text{H}_6\text{O})_b(\text{C}_2\text{H}_4\text{O})_a\text{H}$ where "a" and "b" denote the number of polyoxyethylene and polyoxypropylene units, respectively.

[0147] Preferred hydrophilic surfactants of this class include Poloxamers 108, 188, 217, 238, 288, 338, and 407. Preferred hydrophobic surfactants in this class include Poloxamers 124, 182, 183, 212, 331, and 335.

35 Sorbitan Fatty Acid Esters

[0148] Sorbitan esters of fatty acids are suitable surfactants for use in embodiments of the present invention. Among these esters, preferred hydrophobic surfactants include sorbitan monolaurate (Arlacel 20), sorbitan monopalmitate (Span-40), sorbitan monooleate (Span-80), sorbitan monostearate.

[0149] The sorbitan monopalmitate, an amphiphilic derivative of Vitamin C (which has Vitamin C activity), can serve two important functions in solubilization systems. First, it possesses effective polar groups that can modulate the micro-environment. These polar groups are the same groups that make vitamin C itself (ascorbic acid) one of the most water-soluble organic solid compounds available: ascorbic acid is soluble to about 30 wt/wt % in water (very close to the solubility of sodium chloride, for example). And second, when the pH increases so as to convert a fraction of the ascorbyl palmitate to a more soluble salt, such as sodium ascorbyl palmitate.

Ionic Surfactants

[0150] Ionic surfactants, including cationic, anionic and zwitterionic surfactants, are suitable hydrophilic surfactants for use in embodiments of the present invention. Preferred ionic surfactants include quaternary ammonium salts, fatty acid salts and bile salts. Specifically, preferred ionic surfactants include benzalkonium chloride, benzethonium chloride, cetylpyridinium chloride, docetyl trimethyl ammonium bromide, sodium docecylsulfates, dialkyl methylbenzyl ammonium chloride, edrophonium chloride, domiphen bromide, dialkylesters of sodium sulfonsuccinic acid, sodium dioctyl sulfosuccinate, sodium cholate, and sodium taurocholate. These quaternary ammonium salts are preferred additives. They can be dissolved in both organic solvents (such as ethanol, acetone, and toluene) and water. This is especially useful for medical device coatings because it simplifies the preparation and coating process and has good adhesive properties. Water insoluble drugs are commonly dissolved in organic solvents.

Fat-Soluble Vitamins and Salts Thereof

[0151] Vitamins A, D, E and K in many of their various forms and provitamin forms are considered as fat-soluble vitamins and in addition to these a number of other vitamins and vitamin sources or close relatives are also fat-soluble and have polar groups, and relatively high octanol-water partition coefficients. Clearly, the general class of such compounds has a history of safe use and high benefit to risk ratio, making them useful as additives in embodiments of the present invention.

[0152] The following examples of fat-soluble vitamin derivatives and/or sources are also useful as additives: Alpha-tocopherol, beta-tocopherol, gamma-tocopherol, delta-tocopherol, tocopherol acetate, ergosterol, 1-alpha-hydroxycholecalciferol, vitamin D2, vitamin D3, alpha-carotene, beta-carotene, gamma-carotene, vitamin A, fursultiamine, methylolriboflavin, octotiamine, prosultiamine, riboflavin, vintiamol, dihydrovitamin K1, menadiol diacetate, menadiol dibutyrate, menadiol disulfate, menadiol, vitamin K1, vitamin K1 oxide, vitamins K2, and vitamin K-S(II). Folic acid is also of this type, and although it is water-soluble at physiological pH, it can be formulated in the free acid form. Other derivatives of fat-soluble vitamins useful in embodiments of the present invention may easily be obtained via well known chemical reactions with hydrophilic molecules.

Water-Soluble Vitamins and their Amphiphilic Derivatives

[0153] Vitamins B, C, U, pantothenic acid, folic acid, and some of the menadione-related vitamins/provitamins in many of their various forms are considered water-soluble vitamins. These may also be conjugated or complexed with hydrophobic moieties or multivalent ions into amphiphilic forms having relatively high octanol-water partition coefficients and polar groups. Again, such compounds can be of low toxicity and high benefit to risk ratio, making them useful as additives in embodiments of the present invention. Salts of these can also be useful as additives in the present invention. Examples of water-soluble vitamins and derivatives include, without limitation, acetiamine, benfotiamine, pantothenic acid, cetotiamine, cyclothiamine, dexpantenol, niacinamide, nicotinic acid, pyridoxal 5-phosphate, nicotinamide ascorbate, riboflavin, riboflavin phosphate, thiamine, folic acid, menadiol diphosphate, menadione sodium bisulfite, menadoxime, vitamin B12, vitamin K5, vitamin K6, vitamin K6, and vitamin U. Also, as mentioned above, folic acid is, over a wide pH range including physiological pH, water-soluble, as a salt.

[0154] Compounds in which an amino or other basic group is present can easily be modified by simple acid-base reaction with a hydrophobic group-containing acid such as a fatty acid (especially lauric, oleic, myristic, palmitic, stearic, or 2-ethylhexanoic acid), low-solubility amino acid, benzoic acid, salicylic acid, or an acidic fat-soluble vitamin (such as riboflavin). Other compounds might be obtained by reacting such an acid with another group on the vitamin such as a hydroxyl group to form a linkage such as an ester linkage, etc. Derivatives of a water-soluble vitamin containing an acidic group can be generated in reactions with a hydrophobic group-containing reactant such as stearylamine or riboflavin, for example, to create a compound that is useful in embodiments of the present invention. The linkage of a palmitate chain to vitamin C yields ascorbyl palmitate.

Amino Acids and Their Salts

[0155] Alanine, arginine, asparagines, aspartic acid, cysteine, cystine, glutamic acid, glutamine, glycine, histidine, proline, isoleucine, leucine, lysine, methionine, phenylalanine, serine, threonine, tryptophan, tyrosine, valine, and their derivatives are other useful additives in embodiments of the invention.

[0156] Certain amino acids, in their zwitterionic form and/or in a salt form with a monovalent or multivalent ion, have polar groups, relatively high octanol-water partition coefficients, and are useful in embodiments of the present invention. In the context of the present disclosure we take "low-solubility amino acid" to mean an amino acid which has solubility in unbuffered water of less than about 4% (40 mg/ml). These include cystine, tyrosine, tryptophan, leucine, isoleucine, phenylalanine, asparagine, aspartic acid, glutamic acid, and methionine.

Organic Acids and Their Esters and Anhydrides

[0157] Examples are acetic acid and anhydride, benzoic acid and anhydride, acetylsalicylic acid, diflunisal, 2-hydroxyethyl salicylate, diethylenetriaminepentaacetic acid dianhydride, ethylenediaminetetraacetic dianhydride, maleic acid and anhydride, succinic acid and anhydride, diglycolic anhydride, glutaric anhydride, ascorbic acid, citric acid, tartaric acid, lactic acid, oxalic acid aspartic acid, nicotinic acid, 2-pyrrolidone-5-carboxylic acid, and 2-pyrrolidone.

[0158] These esters and anhydrides are soluble in organic solvents such as ethanol, acetone, methyl ethyl ketone, ethyl acetate. The water insoluble drugs can be dissolved in organic solvent with these esters and anhydrides, then coated easily on to the medical device, then hydrolyzed under high pH conditions. The hydrolyzed anhydrides or esters are acids or alcohols, which are water soluble and can effectively carry the drugs off the device into the vessel walls.

EXAMPLES

[0159] The invention is further described in the following examples, which do not limit the scope of the invention described in the claims.

Example 1 (Comparative example): Aqueous aerosol formulation

[0160] An exemplary aqueous formulation of rapamycin was prepared using the following components.

Component	Amount (g)	Mass Fraction (w/w)
rapamycin	0.1	0.01 %
ethanol	250	25 %
propylene glycol	250	25 %
polysorbate 80	0.02	0.002 %
water	500	50 %
Total	1000	

[0161] Blending Procedure: in a 1000 ml amber volumetric flask, blend 250 propylene glycol with 250 ethanol until uniform. Then sequentially dissolve first 100 mg rapamycin then 20 mg polysorbate 80 in the propylene glycol and ethanol solution. Add water to bring the volumetric to 1000 ml and stir or sonicate until uniform and all the rapamycin is dissolved. Store at controlled temperature away from light.

Example 2: Dry Powder Formulation

[0162] Batch 06RP68.HQ00008 and 06RP68.HQ00009. These two formulations are each a blend of micronized drug (rapamycin) particles dispersed onto the surface of lactose carrier particles. The final composition of each batch comprises 1% (w/w) drug particles having a mean diameter of about 2.60 μm (microns) and 3.00 μm (microns), respectively. Drug particles having a suitable size range are made by wet polishing (06RP68.HQ00008) or jet milling (06RP68.HQ00009), as described below. While this example used 1% (w/w) rapamycin, a range 0.5 to 20% is practicable. The carrier particles consist of a blend of two carriers, Respitose[®] SV003, present at 95.5% (w/w) and having particle sizes of about 30 to 100 μm (microns) (equivalent spherical diameter), and Respitose[®] LH300 (Lactohale 300) present at 5.5% (w/w) and having particle sizes less than 10 μm (microns) (equivalent spherical diameter). After blending, the blends were assayed to confirmed homogeneity and drug content of 1%.

[0163] To reduce drug particle agglomeration and aid in the aerosolization of drug particles several other excipients are optionally included. Optional excipients include phospholipids, such as dipalmitoylphosphatidylcholine (DPPC) and lecithin, and metal fatty acid salts, such as magnesium stearate. These can be coated on the carrier particles in weight ratio of excipient to large carrier particle ranging from 0.01 to 0.5%.

[0164] Capsule Filling: 20 milligrams of the powder blends from Batch 06RP68.HQ00008 and Batch 06RP68.HQ00009 were loaded into size #3 HPMC capsules to produce drug product. For these blends it was feasible to load from 5 to 35 milligrams of drug into #3 size capsules and empty greater than 95% of the loaded blend from the capsule upon actuation in Plastiap[®] RS01 Model 7 or Plastiap[®] RS00 Model 8 devices at flow rates ranging from 60 to 100 liters per minute.

Reference

Example 3: Determination of rapamycin in lung and blood following administration by oropharyngeal aspiration (OPA) and oral gavage to C57BL6 mice

[0165] This study was conducted to evaluate the concentration of rapamycin in male C57BL/6 mice after administration of rapamycin at a very high target dose of 1 mg/kg by gavage and oropharyngeal aspiration (OPA). A method for the analysis of rapamycin in mouse blood and lung homogenate was developed using liquid chromatography with tandem mass spectrometry detection (LC-MS/MS). Calibration curves of rapamycin using triplicate concentrations were analyzed between 1 ng/mL and 2000 ng/mL in mouse blood, and between 2 ng/mL and 20,000 ng/mL in mouse lung homogenate. Accuracy, precision and linearity were within expected ranges.

[0166] In pilot studies, the efficiency of vehicle delivery to the lungs via oropharyngeal aspiration with a volume of 50

μL per mouse was evaluated by administration of Evans Blue dye. The presence of blue dye only in lungs was verified visually, and the absence of blue dye in the stomach demonstrated that delivery to the stomach was avoided in the procedure used.

[0167] Rapamycin was administered to male C57BL/6 mice (N=6) by gavage at a dose of 1.0 mg/kg either orally or via OPA. The oral dose was formulated using pharmaceutical oral liquid formulation Rapamune Oral® (Pfizer). Rapamycin for OPA was prepared by dissolving the test article in an appropriate volume of ethanol, and then addition of an appropriate volume of water to prepare a 10% ethanol solution at a concentration of 1 mg rapamycin/mL. Rapamycin was administered to 2 groups of 6 male C57BL/6 mice by OPA under isoflurane anesthesia. An additional group of 6 mice received vehicle only (10% ethanol in water). At 1 h after administration a group of 6 mice receiving oral and OPA rapamycin were euthanized, and blood was obtained by cardiac puncture, and the lungs removed. The remaining mice in each group administered rapamycin or vehicle by OPA were observed for an additional 3 days. At the 72-h necropsy, blood was obtained by cardiac puncture and the lungs removed. No adverse effects were observed in rapamycin- or vehicle-treated mice in the 72 h period following dosing.

[0168] The concentration of rapamycin was determined in the collected blood and in lung homogenate by LC-MS/MS. At 1 h following OPA of rapamycin, the concentration of rapamycin was ~6 fold higher in lung tissue (3794 ± 1259 ng/g tissue) than in blood (641 ± 220 ng/mL). Following oral administration of a similar dose of rapamycin, the 1-h lung and blood concentrations of rapamycin were 71 ± 43 ng/g and 23 ± 16 ng/mL, respectively. Lung homogenate concentrations following OPA were 53-fold higher than those measured following oral administration of the same high dose (1 mg/kg) of rapamycin. The data suggests that delivery of lower doses of rapamycin to lung (dose levels that do not saturate system) will result in rapamycin levels in the lung that can be achieved by oral dosing but with significantly less rapamycin in the blood than occurs with oral dosing.

Materials and Methods

[0169] Test Substance: Sirolimus (Rapamune, Rapamycin) MW 914.172, C₅₁N₇₉NO₁₂, CAS NUMBER: 53123-88-9. Source (for oral gavage): Rapamune Oral® (Pfizer) for oral administration, Lot No.: MWGT, Expiration: 07/16. Source (for OPA): Rapamycin (Sirolimus) solid, LC Laboratories, Woburn MA, Lot No.: ASW-127, Expiration: 12/2023.

[0170] Animals: Male C57BL/6 mice, approximately 8 weeks of age, from Charles River Laboratories, Inc, Raleigh, NC. Animals were fed Certified Purina Rodent Chow #5002 and were furnished tap water *ad libitum*. The analysis of each feed batch for nutrient levels and possible contaminants was performed by the supplier, examined by the Study Director, and maintained in the study records. The feed was stored at approximately 16-21 °C (60-70 °F) and the period of use did not exceed six months from the milling date. Mice were housed (one per cage) in polycarbonate cages with stainless steel bar lids accommodating a water bottle. Cage sizes are approximately 29.2 cm × 19.0 cm × 12.7 cm (11.5" × 7.5" × 5") high (451 cm² (70 sq. in. floor space)) for mice. Contact bedding was Sani-Chips hardwood chips (P. J. Murphy Forest Products Co., Montville, NJ). Mice were quarantined for a period of 5 days before use on a study. A veterinarian or qualified designee examined the animals prior to their release from quarantine. Temperature and relative humidity in RTI animal rooms were continuously monitored, controlled, and recorded using an automated system (Siebe/Barber-Colman Network 8000 System with Revision 4.4.1 for Signal® software [Siebe Environmental Controls (SEC)/Barber-Colman Company; Loves Park, IL]). The target environmental ranges were 64-79 °F (18 °C - 26 °C) for temperature and 30-70% relative humidity, with a 12-h light cycle per day. At the end of the in-life phase, the mice were euthanized by overexposure to carbon dioxide.

[0171] Test Chemical Preparation: Evans Blue was prepared at 0.5% w/v in sterile distilled water. Rapamune Oral® was administered as supplied for oral dosing. Rapamycin (solid) was dissolved in ethanol and diluted with sterile distilled water to provide a final concentration of 0.5 mg/mL in 10% ethanol.

[0172] Dosing: Each animal was weighed prior to dosing to determine the amount of dose to be administered. A single gavage dose was administered using a 100-μL glass syringe (Hamilton, Reno, NV) fitted with a ball-tipped 20-G stainless steel gavage dosing needle (Popper & Sons Inc., New Hyde Park, NY). The dose administered to each animal was determined from the weight of the full syringe minus that of the empty syringe. The dosing time was recorded. Dosing of animals was spaced apart to allow blood collection at the appropriate times. The dose formulations administered to each group are shown below.

[0173] For oropharyngeal aspiration group animals, a single dose of rapamycin (50 μL) was administered to each mouse under isoflurane anesthesia, using a 100 μL glass syringe (Hamilton, Reno, NV) fitted with a ball-tipped 24-G stainless steel gavage dosing needle (Popper & Sons Inc., New Hyde Park, NY). The mouse was weighed prior to dosing, and the dose of rapamycin administered was recorded by weight. Each mouse was anesthetized with isoflurane, and restrained with the mouth open. The tongue was held to one side of the mouth with forceps, and the dose was slowly injected into the distal part of the oral cavity. The nostrils were covered with a finger for two breaths to ensure aspiration (Rao et al., 2003).

Table 1: Study Design Summary

Dose Group	Route	Compound	No. Animals	Target Dose (mg/ml)	Target Dose (ul)	Target Dose (mg/kg)	Collection Time	Samples Collected
1	OA	Evans Blue	6	-	50	0	1	blood, lung
2	OA	Rapamycin	6	0.5	50	1.0	1	blood, lung
3	Gavage	Rapamune Oral	6	1.0	25	1.0	1	blood, lung
4	OA	Vehicle	6	0	50	1.0	72	blood, lung
5	OA	Rapamycin	6	0.5	50	1.0	72	blood, lung

[0174] Collection of Blood and Lung Samples: At study termination (1 or 72 h after dosing), mice were anesthetized by exposure to CO₂, and blood was collected by cardiac puncture with dipotassium EDTA as anticoagulant. Lung tissue was excised and divided into the right and left lung. The left lung was used for analysis, and the right lung flash frozen in liquid nitrogen and stored at -70 °C for further analysis.

[0175] Analysis of Samples for Rapamycin by LC-MS/MS: An LC-MS/MS method for analysis of rapamycin in lung and blood was prepared based on the published method of Wu et al. (2012). The volumes of blood and lung homogenate were reduced substantially from the published method. Triamcinolone was used as internal standard.

[0176] Lung homogenate was prepared by homogenization of weighed lung samples with 2.8-mm ball bearings in a homogenizer with tissue + deionized water (1:3 w/v) in a SPEX SamplePrep 2010 Geno/Grinder.

[0177] The concentrations of standards were arranged so that each standard came from an alternate stock standard. A six-point calibration curve, each made in triplicate, was employed for analyte quantitation. A simple linear regression model with or without weighting was employed for curve fitting. The concentration range determined was from 1-2000 ng/mL in blood and 2-2000 ng/mL in lung homogenate.

[0178] The following method performance parameters were considered acceptable; the coefficient of determination, r^2 , of ≥ 0.98 for concentration-response relationship; an accuracy of $\leq \pm 15\%$ (for concentrations above LOQ) or $\leq \pm 20\%$ (for concentration at LOQ) of the nominal value. r^2 was greater than 0.999 in all analysis.

[0179] Thirty (30) μL of matrix, 30 μL of spiking solution (methanol for blanks and samples), 10 μL Internal standard solution (in MeOH) and 90 μL of MeOH were pipetted into microcentrifuge tubes, vortexed briefly, then centrifuged for 6 min at 10,000 RPM at $\sim 4^\circ\text{C}$. Aliquots (90 μL) of supernatant were transferred to LC vial inserts, and then analyzed by LC-MS/MS (Table 2).

Table 2: LC-MS/MS Method

Column	Waters Acquity UPLC HSS T3 1.8 μm , 2.1 x 50 mm with VanGuard 2.1 x 5 mm HSS T3 1.8 μm .
Mobile Phase A	10 mM Ammonium Acetate in water, 0.1% acetic acid
Mobile Phase B	MeOH
Injection Vol	2 μL
Flow Rate	0.5 ml/min
Gradient	70% A for 1 min, a linear gradient to 5% A from 1-3 min, held for 1 min, a linear gradient to 70% A from 4-5.1 min, and held at 70% until 6 min
Rapamycin MRM	931.70 $\rightarrow$ 864.70
Triamcinolone (IS MRM)	395.30 $\rightarrow$ 357.20

[0180] Data Collection and Reporting: Study data was collected and reported in the Debra™ system version 5.5.10.72

(Lablogic Systems Ltd., Sheffield, England). This includes data for animal body weights, dose administered, dose time, and sample collection times. Calculations of dose administered and sample collection times were reported with the Debra™ system.

Results

[0181] Rapamycin Analysis: The analysis of rapamycin was set up of sample volumes of 30 µL of blood and lung homogenate. Example chromatograms are shown for rapamycin and internal standard in blood and lung (Figures 1 and 2). Prior to the generation of study samples, triplicate calibration curves were generated for lung and blood, to verify method performance. The calibration range was from 1.0 -2000 ng/mL for blood and 1 - 20,000 ng/mL for lung homogenate. Lung homogenate was prepared with 1 g of lung tissue homogenized in 3 volumes of water, to yield a 1:4 homogenate. Calibration curves are shown in Figures 3 and 4 for blood, lung homogenate, and solvent.

[0182] Oropharyngeal Aspiration: Prior to the administration of rapamycin by oropharyngeal aspiration, administration of Evans Blue was used to verify that the OPA delivered the dose to the lungs. Mice were anaesthetized with isoflurane and administered Evans Blue by OPA, using a syringe equipped with a blunt needle. Immediately following OPA, the mice were euthanized and the lungs and stomach examined visually to ensure that the Evans Blue dye was delivered to the lungs, and was not delivered to the stomach. Four mice were successfully administered Evans Blue with all of the dye appearing to be located in the lungs and none in the stomach.

[0183] Rapamycin Administration: The weight of dose solution administered was determined by weighing the charged syringe with dose solution prior to dosing, and weighing following dosing. The weight of dose solution administered was used to calculate the amount of rapamycin administered. The time of dosing was recorded as 0. Animals in groups 2 and 3 were euthanized at 1 h after dosing. Animals in groups 4 and 5 were observed for 72 h after dosing. No significant clinical signs were observed in any of the groups.

[0184] Rapamycin Analysis in Blood and Lung: Rapamycin was analyzed in mouse blood and left lung homogenate in all of the samples collected (Figures 5 and 6). Samples of the right lung from each animal were saved for potential further analysis. Summary data for the samples are provided in Table 3.

Table 3: Concentration of Rapamycin in Blood and Lung Following Oral and Oropharyngeal (OPA) Administration of Rapamycin to Mice (1 mg/kg)

Animal No.	Route of Admin	Time post-dose (h)	Lung (ng/g tissue)	Blood (ng/ml)
2-07	OPA	1	5040	615.5
2-08	OPA	1	2642	455.5
2-09	OPA	1	4500	622
2-10	OPA	1	1874	364.5
2-11	OPA	1	4006	937
2-12	OPA	1	4700	848.5
Mean			3794	641
SD			1259	220
3-13	Gavage	1	109.8	49.85
3-14	Gavage	1	24.66	11.3
3-15	Gavage	1	122.8	28.7
3-16	Gavage	1	54	28.35
3-17	Gavage	1	<LOQ	2.845
3-18	Gavage	1	43	19.35
Mean			71	23
SD			43	16
5-25	OPA	72	11.7	<ELOQ
5-26	OPA	72	11.4	<ELOQ
5-27	OPA	72	15.7	<ELOQ
5-28	OPA	72	10.8	<ELOQ
5-29	OPA	72	11.9	<ELOQ
5-30	OPA	72	13.6	<ELOQ

(continued)

Animal No.	Route of Admin	Time post-dose (h)	Lung (ng/g tissue)	Blood (ng/ml)
Mean			12.5	
SD			1.9	

[0185] For all sample sets, a triplicate calibration curve was analyzed with the sequence of standard set, sample replicate 1, standard set, sample replicate sample 2, standard set. At 1 h following OPA of rapamycin, the concentration of rapamycin was ~6 fold higher in lung tissue (3794 ± 1259 ng/g tissue) than in blood (641 ± 220 ng/ml). Following oral administration of a similar dose of rapamycin, the 1-h lung and blood concentrations of rapamycin were 71 ± 43 ng/g and 23 ± 16 ng/mL, respectively. Lung homogenate concentrations following OPA were 53-fold higher than those measured following oral administration of same high dose (1 mg/kg) of rapamycin.

Discussion

[0186] This study investigated the concentration of rapamycin in blood and lung tissue following administration of rapamycin by gavage in a commercial oral formulation, and by oropharyngeal administration (OPA) as a suspension prepared in 10% aqueous ethanol. No adverse effects were observed in rapamycin- or vehicle-treated mice up to 72 h following dosing via OPA. Prior to administration of rapamycin, an analytical method was developed, and the administration of a dye into the lung by OPA was verified. The concentrations of rapamycin in lung following OPA were 6-fold higher than in blood. At 72 h after OPA, rapamycin was below the limit of quantitation in blood, but was detectable in lung. This study indicated that rapamycin is available systemically following pulmonary administration, and that lung tissue concentrations greatly exceed that of blood at early and late time points following delivery to the lung.

[0187] These results further demonstrate that rapamycin delivered directly to the lung achieves an unexpectedly high local concentration of drug in lung tissue compared to the blood. This result was entirely unexpected from what is known about the pharmacology of rapamycin, which predicts an approximately equal concentration of the drug in lung tissue and the blood because rapamycin is known to distribute evenly throughout bodily tissues and should be cleared rapidly from the lung due to its high lipophilicity. Accordingly, these results indicate that direct administration of rapamycin to the lungs should be able to achieve a high enough delivered dose for therapeutic efficacy while at the same time achieving almost undetectable systemic availability, thereby eliminating the toxicities associated with oral administration that are due to systemic exposure to the drug. While toxicity to the lung itself is also of concern in view of earlier studies, the results here further unexpectedly indicate that relatively high amounts of rapamycin were not acutely toxic to lung tissue.

Reference

Example 4: Rapamycin inhibits the viability of TSC2 mutant cells and inhibits S6 phosphorylation

[0188] The anti-proliferative activity of rapamycin was tested against the angiomyolipoma (AML) derived TSC2 deficient TRI-AML101 cell line. The TRI-AML101 cell line was derived from a TSC2 deficient primary human AML provided by Dr. Elizabeth Henske (Fox Chase Cancer Center, Philadelphia, PA). The tumor cells were immortalized by a two step process. First the cells were infected with the amphotropic retrovirus LXS16E6E7 that encodes the HPV16 E6 and E7 open reading frames and neomycin resistance cassette. Cells were expanded and neomycin-selected. Individual clones were isolated and frozen down. Next, the human Telomerase gene (hTERT) with hygromycin resistance cassette (pLXS16E6E7-hTERT-hyg plasmid) was transfected in and a stable line was selected by hygromycin selection.

[0189] The activity of rapamycin was tested on the TRI-AML101 cells by performing 10 point dose response analysis of cell viability. Two thousand cells in 50 μ L of growth media (DMEM, 10% FBS, and 1% Penicillin/Streptomycin) were plated per well in a 96 well plate. 24 hours after plating cells another 50 μ L of growth medium containing rapamycin (0.0005-5000 nM, 10-fold dilutions, 0.1% final DMSO concentration) or DMSO only was added to the cells. 72 hours after compound addition, relative cell viability was determined by CellTiter-Glo[®] luminescence assay (Promega) and expressed as a percentage relative to vehicle (DMSO) treated control cells. Rapamycin inhibited viability at concentrations as low as 0.05 nM (Fig. 7, bottom). Inhibition of the mTOR pathway was also demonstrated by measuring the levels of phosphorylated S6 by western blot. AML cells were incubated with 20 nM rapamycin for 24 hours. Western blot analysis was then performed and demonstrated that rapamycin potently inhibits S6 phosphorylation (Fig. 7, top).

Reference

Example 5: S6 Phosphorylation in Mouse Lung Following Oral and OPA Administration of Rapamycin

[0190] As discussed above, our experiments showing the tissue distribution of rapamycin in lung and blood following oral administration and OPA demonstrated that direct administration of rapamycin to the lungs should be able to achieve a high enough delivered dose for therapeutic efficacy while at the same time achieving very low systemic exposure to the drug, thereby simultaneously improving therapeutic efficacy and eliminating many of the toxicities associated with oral administration of rapamycin. To validate this approach, we used the presence of phosphorylated S6 protein in murine lung tissue as a biomarker for mTOR activity. In the mouse strain used (C57bl/6), the mouse airway and alveolar epithelial cells have constitutively active (phosphorylated, "p") S6 protein. The S6 protein is typically phosphorylated by S6K which is downstream of mTORC1 and is activated, for example, downstream of growth factors such as epidermal growth factor (EGF), AKT, ERK, and RSK. mTORC1 promotes cell growth and proliferation by stimulating anabolic processes such as biosynthesis of lipids, proteins, and organelles, and suppressing catabolic processes such as autophagy. The mTORC1 pathway senses and integrates intracellular and extracellular signals, including growth factors, oxygen, amino acids, and energy status, in order to regulate a wide range of processes, such as protein and lipid synthesis and autophagy. mTORC1 is acutely sensitive to rapamycin.

[0191] In the present study, lung tissue was taken from the C57bl/6 mice treated as discussed above, either with vehicle (n=6), or 1 mg/kg rapamycin administered via OPA (n=6) or via oral gavage (n=6) at two time points post dosing, 1 hr and 72 hours. As discussed above, following OPA at 1 hr, rapamycin was detected at 641 ng/ml in the blood and 3794 ng/g tissue in the lung, and at 72 hrs was still detectable in the lung at 12.5 ng/g while being undetectable in the blood at that time point. Conversely, following oral (gavage) administration, at 1 hr, rapamycin was detected at 23 ng/ml in the blood and 71 ng/g tissue in the lung, and at 72 hrs was undetectable in either the lung or blood. As shown by the data in Figure 8A, the level of phosphorylated S6 (pS6) was reduced substantially by both OPA and orally administered rapamycin at 1 hr and remained suppressed at 72 hr for OPA. pS6 was highest in the vehicle control because these mice have constitutively active mTOR signaling. These data show that a delivered dose of rapamycin sufficient to achieve about 70 ng/g drug in the lung substantially abrogates mTOR signaling in the lung tissue as measured by pS6 protein and that mTOR signaling remains suppressed at levels as low as 12.5 ng/g. These results validate our approach to utilize inhaled rapamycin for the treatment of diseases and disorders characterized by aberrantly high mTOR pathway activity by demonstrating that inhaled rapamycin can be delivered at much lower doses than orally administered rapamycin to simultaneously achieve high therapeutic efficacy and very low toxicity.

Example 6: Inhaled rapamycin inhibits S6 phosphorylation in lung tissue

[0192] Normal Sprague-Dawley Rats were dosed by inhalation to achieve target dose of 0.354 mg/kg of rapamycin (LAM-001) (N=36) and subgroups of 6 animals were sacrificed at the following time points (1) Predose, (2) Midway dosing, (3) Immediate post dosing, (4) 2 hours post dosing, (5) 4 hours post dosing and (6) 12 hours post dosing on Study Day 1. Charles River determined the lung concentration of rapamycin for each sacrificed animal at their subgroup time point, the average rapamycin concentration in nanograms rapamycin per gram of tissue (ng/g) for each group is reported in the table below.

Table 4: Rapamycin in lung tissue following administration by inhalation

Target dose 11 ug Day 1	Midway Dosing	Immediate post-dosing	2 hr post-dosing	4 hr post-dosing	12 hr post-dosing
Avg Lung rapa (ng/g):	462	560	250	192	95

[0193] Lungs samples from each animal were collected and snap frozen. The individual frozen lung samples were homogenized (Qiagen TissueLyser LT according to manufacturer's protocols) in 1X RIPA Buffer with protease and phosphatase inhibitors. The lung homogenates were analyzed by western blot analysis of the mTOR downstream target, Phospho-S6 Ribosomal Protein (Ser240/244) (Cell Signaling Technology antibody, clone D68F8) as compared to total S6 protein S6 Ribosomal Protein (Cell Signaling Technology antibody, clone 5G10) levels. The Western blot images were analyzed by NIH imageJ v1.48 to generate the respective antibody reactivity/intensity and create the ratio of S6 phosphorylation (S6-P) to total S6 intensities for each lung sample. The S6-P/ total S6 ratios (y-axis) for sample organized by timepoint group (X-axis) were plotted on a one grouping variable scatter plot vertical graph (GraphPad, version 4.0), all samples in groups are represented by filled in black dots (●) on the graph and average is shown by the horizontal

line between the dots within each respective group (Fig. 8B).

Example 7: Inhaled rapamycin shows unexpected biodistribution to the lung

[0194] It had been reported in the literature that rapamycin collects in the lungs after high oral or IV doses (Yanez, J. et. al., *Pharmacometrics and Delivery of Novel Nanoformulated PEG-b-poly(ϵ -caprolactone) Micelles of Rapamycin, Cancer Chemotherapy and Pharmacology*, 61 (1), 133-144 2007). A study reported that after administering a single dose of 10.0 milligrams/kilogram (mg/kg) to Sprague-Dawley (SD) rats, the amount of rapamycin in the lungs after allowing time for distribution through the tissue compartments (24 hours) was 721 nanograms/gram (ng/g), approximately 19 times the concentration in the blood (Table 5).

Table 5. Biodistribution of Rapamycin in the lung and blood after IV administration per Yanez *et al.*

Yanez 1 Day (24 hours after single dose)				
IV Dose (mg/kg/day)	Rapamycin in Lung (ng/g)	Lung/Blood Ratio	Daily Human Dose Equiv (mg)	Rapamycin in Blood (ng/ml)
10	721	19	600	31

[0195] In an earlier separate study by Napoli (Napoli, K., et. al., *Distribution of Sirolimus in Rat Tissue, Clinical Biochemistry*, 30(2): 135-142, 1997) a range of rapamycin doses to SD rats were administered daily, by oral and intravenous (IV) administration routes. After 14 days of IV administration rapamycin concentrations in lung tissue ranged, dose proportionally, from a 200 to 900 ng/g, approximately 23 to 44 times the higher than the concentrations in the blood. But for oral administration much lower levels of rapamycin accumulated in the lung for the same doses, even though the ratios of lung to blood concentrations of rapamycin were approximately the same (Table 6).

Table 6: Biodistribution of Rapamycin 24 hrs after the 14th once daily IV administration over 14 days per Napoli *et al.*

Napoli 14 Day QD (24 hours after 14th dose)				
IV Dose (mg/kg/day)	Rapamycin in Lung (ng/g)	Lung/Blood Ratio	Daily Human Dose Equivalent (mg)	Rapamycin in Blood (ng/ml)
0.04	219	23	2.4	9.4
0.08	457	30	4.8	15.0
0.16	677	33	9.6	20.6
0.40	868	44	24	19.6
Oral Dose (mg/kg/day)	Rapamycin in Lung (ng/g)	Lung/Blood Ratio	Daily Human Dose Equiv. (mg)	Rapamycin in Blood (ng/ml)
0.4	42	53	3.4	0.8
0.8	59	28	6.7	2.1
1.6	175	24	13.4	7.4

[0196] When we examined the biodistribution of rapamycin following administration via inhalation, we found that rapamycin accumulates much higher in the lungs than would have been predicted based upon Napoli and Yanez, even while the lung to blood ratios were similar.

[0197] In a first study, rapamycin was administered to SD rats by inhalation at two doses of (1) 1.0 mg/kg/day and (2) 0.0360 mg/kg/day, for a single day. After allowing 12 hours for distribution to tissue compartments, the rapamycin trough concentrations in the lungs for the high dose was about 14,800 ng/g and the concentration of rapamycin in the lungs was approximately 23 times higher than in the blood (Table 7). For the low dose, the concentration in the lungs was 24 times higher than the concentration in the blood (Table 7). Table 8 shows the trough lung concentration, the maximum and trough blood concentrations after repeated, once a day dosing for 5 days at the same two doses used in the previous experiment, i.e., 1.0 mg/kg/day and 0.0360 mg/kg/day.

Table 7: Biodistribution of Rapamycin via inhalation 12 hrs after a single dose

Inhaled Dose (mg/kg/day)	Rapamycin in Lung (ng/g)	Lung/Blood Ratio	Daily Human Dose Equivalent (mg)	Rapamycin in Blood (ng/ml)
1.00	14831	23	60	645
0.0360	95	24	2.16	4

Table 8: Biodistribution of Rapamycin via inhalation once a day (measured at trough day 5)

Inhaled Dose (mg/kg/day)	Rapamycin in Lung (ng/g)	Lung/Blood Ratio	Daily Human Dose Equivalent (mg)	Rapamycin in Blood (ng/ml)
1.0000	17163	23	60	746
0.0360	87	22	2.16	4

[0198] The results of this initial study indicate that delivery of rapamycin to the lungs by inhalation produced markedly higher concentrations of drug in the lung tissue than could be achieved by alternative routes of administration, e.g. oral or intravenous, according to the previous work of Yanez and Napoli. Moreover, the high amounts of rapamycin in the lung following delivery via inhalation were unexpectedly higher based upon what would have been predicted from Yanez and Napoli. As both intravenous and inhaled routes of administration have high bioavailability of rapamycin, the inhaled 1 mg/kg dose would have been predicted to achieve lung concentrations about 2.5 times those observed by Napoli's 0.4 mg/kg/day IV dose. Instead, the levels of rapamycin in the lung were approximately 17 times higher when administered via inhalation (compare Table 7, 1 mg/kg/day via inhalation produced 14,831 ng/g drug in lung versus Table 6 (Napoli), 0.40 mg/kg/day IV produced 868 ng/g lung; $14,831/868 = 17$). Similarly the 10 mg/kg intravenous dose administered by Yanez would have been predicted to achieve a lung concentration of rapamycin about 10 times higher than that achieved by the 1 mg/kg inhaled dose. Instead, the intravenous dose achieved lung concentrations approximately 20 times less than the inhaled dose (compare Table 7, 1 mg/kg/day via inhalation produced 14,831 ng/g drug in lung versus Table 5 (Yanez), 10 mg/kg/day IV produced 721 ng/g in lung; $14,831/721 = 21$). This could possibly be due to low metabolic activity in the lungs and slow passive or active transport of rapamycin from the lung tissue compartment into systemic circulation. Regardless of the precise mechanism, these results indicate that delivery of rapamycin to the lungs results in persistently high local concentrations, while circulatory concentrations remain low.

[0199] The results of this initial study were replicated and expanded in additional rat studies and dog studies. These subsequent studies were structured to determine the repeat dose toxicity and toxicokinetics of a dry powder aerosol formulation of 1% (w/w) rapamycin blended with lactose, administered by inhalation to normal Sprague-Dawley (SD) rats and Beagle dogs. In the first study, standard cylindrical flow-through nose-only inhalation chambers were utilized to administer the dry powder formulation to SD rats. For five consecutive days, animals were subjected to test article for 300 minutes each day to achieve a target dose of 0.354 mg/kg of rapamycin. Two sets of animals were used to perform the toxicokinetic measurements for this study. The first set of animals were dosed for 300 minutes on Study Day 1 and blood samples (N=36) and lung samples (N=36) were taken from animals were sacrificed in subgroups of 6 at the following time points: (1) Predose, (2) Midway dosing, (3) Immediate post dosing, (4) 2 hours post dosing, (5) 4 hours post dosing and (6) 12 hours post dosing. The second set of animals were dosed for 300 minutes for 5 consecutive days, and on Study Day 5, blood samples (N=36) and lung samples (N=36) were taken from animals were sacrificed in subgroups of 6 at the following time points (1) Predose, (2) Midway dosing, (3) Immediate post dosing, (4) 2 hours post dosing, (5) 4 hours post dosing and (6) 12 hours post dosing. The maximum concentration of rapamycin in whole blood (ng/ml) and lung tissue samples (ng/g), as well as the concentration of rapamycin 12 hours post-dose.

[0200] A second repeat exposure study was conducted to evaluate the repeat dose toxicity and toxicokinetics of the dry powder formulation administered to SD rats and Beagle dogs for 28 days.

[0201] For the rat study, standard cylindrical flow-through nose-only inhalation chambers were utilized, as above. For 28 consecutive days, animals were exposed to test article for 300 minutes each day to achieve a target doses of 0.167, 4.75 and 9.50 mg/kg of rapamycin respectively. For each of the three dosing groups, one set of animals (N=36) were dosed for 300 minutes each day for 28 consecutive days. On Study Days 1 and 28, blood samples were taken from animals in subgroups of 6 at the following time points: (1) Predose, (2) Midway dosing, (3) Immediate post dosing, (4) 2 hours post dosing, (5) 4 hours post dosing, (6) 12 hours post dosing and (7) 24 hours post dosing.

[0202] For the dog study, a positive flow delivery system (PFDS) consisting of a central plenum and delivery arms was utilized. The central plenum was of modular design with separate ports into which were attached 5 delivery arms

fitted with oronasal exposure masks fitted with inlet and outlet tubes. The mask was fitted over the dog's muzzle in such a way that the nose was inside the mask, allowing entrance and exit of air. During exposure, animals wore a harness and were placed on a restraint platform. The harness was attached to two side poles on the platform in order to restrict lateral movement of the dog. The front part of the harness was loosely attached to a hook on the front of the platform to prevent the animal from turning around. Dogs were exposed to test article for 60 minutes each day to achieve a target doses of 0.020 and 0.053 mg/kg of rapamycin respectively. For each dosing group, one set of animals (N=6) was dosed for 60 minutes each day for 28 consecutive days. On Study Days 1 and 28, blood samples were taken from animals in subgroups of 6 at the following time points: (1) Predose, (2) post dosing (T=0), (3) 1 hour post dosing, (4) 4 hours post dosing, (5) 8 hours post dosing, (6) 12 hours post dosing and (7) 24 hours post dosing. On Day 29, the dogs were sacrificed and a portion of their lung tissue was removed and minced for analysis of rapamycin content.

[0203] The maximum concentration of rapamycin in whole blood (ng/ml) and the trough concentration of rapamycin are presented below along with extrapolations to human dosing. Also included in this table are the trough levels of rapamycin in dog lung (ng/g) after 28-days of repeat dosing.

Table 9: Inhaled rapamycin rat data and extrapolations for human dosing

	Emitted Dose (ug)	Repeat inhaled daily dose (ug)	Max blood levels (ng/ml)	Blood trough levels (ng/ml)	Lung 24 hr trough levels (ng/g)
Rat*	50	5	7.6	2	N/A
	1425	143	70.5	23.3	N/A
	2850	285	77.8	21.0	N/A
Dog**	160	40	7.4	1.9	32
	424	106	27.3	5.9	61
Human	87	35	0.65	0.15	2
	175	70	1.3	0.3	4
*rat 28-day repeat daily dose study 7300225 - blood data are average of D28					
**dog 28-day repeat daily dose study 7300227 - blood and lung data are average of D28					
** blood and lung estimates are extrapolated from the 7300225 and 7300227 study results					

[0204] Notably, based upon the results presented here, a therapeutically effective dose of rapamycin in the lung in the range of about 5 ng/g in humans could be achieved by administering less than about 100 micrograms to the lungs by inhalation. In contrast, achieving a comparable lung concentration by oral delivery according to Yanez would require 4 to 16 milligrams. To achieve a comparable lung concentration by IV delivery according to Napoli, 60 to 600 micrograms would be required.

[0205] In addition, based upon the results presented here, the therapeutic range of about 5 ng/g in the lung could be achieved with a lung to blood partitioning ratio of 13:1 when rapamycin is delivered via inhalation. This means that while rapamycin is within the therapeutic range in the lung tissue, maximum concentrations of only 650 to 1500 picograms/ml rapamycin would circulate in the blood. This low systemic exposure to rapamycin is expected to reduce the toxicities and adverse drug events associated with the much higher systemic exposure to rapamycin resulting from the higher levels of dosing required with oral or IV administration.

[0206] In summary, the results described here demonstrate that administering rapamycin to the lungs via inhalation advantageously provides for a low dose of rapamycin required to achieve a therapeutically effective dose in the lung in the range of about 5 ng/g, combined with low systemic exposure to the drug, resulting in markedly improved therapeutic index for rapamycin.

Example 8: Size Reduction of Rapamycin for Inhalable Compositions

[0207] Particle size of rapamycin was reduced to a target range of $2.0\mu\text{m} < \text{Dv}_{50} < 3.0\mu\text{m}$ using either a wet polishing or jet milling process. For jet milling, a lab scale MCOne unit from Jetpharma was used with the following operating conditions: venturi pressure 2-4 bar, milling pressure 3-5 bar, feed rate 90 g/h. For wet polishing, feed suspensions were prepared using purified water. A microfluidics high pressure homogenizer was used for the size reduction step and the resulting suspension was spray-dried. Details of the wet polishing process are set forth below.

[0208] The high pressure homogenizer used for the size reduction step of the wet polishing process was a pilot-scale Microfluidics High Pressure Homogenizer equipped with an auxiliary processing module (200 micron) and a 100 micron

interaction chamber was used. The unit was operated at -455 bar (-30 bar in the intensifier module hydraulic pressure). After microfluidization the fluid was removed by spray drying to generate a dry powder. A laboratory scale spray dryer, SD45 (BUCHI, model B-290 Advanced) was equipped with a two fluid nozzle (cap and diameter were 1.4 and 0.7 mm, respectively). Two cyclones in series were used (being the first the standard Buchi cyclone and the second the high-performance Buchi cyclone) to collect the dried product. The spray drying unit was operated with nitrogen and in single pass mode, i.e. without recirculation of the drying nitrogen. The aspirator, blowing nitrogen, was set at 100% of its capacity (flow rate at maximum capacity is approximately 40 kg/h). The flow rate of the atomization nitrogen was adjusted to a value in the rotameter of 40 ± 5 mm. Before feeding the product suspension, the spray dryer was stabilized with purified water, during which the flow rate was adjusted to 6 ml/min (20% in the peristaltic pump). The inlet temperature was adjusted to achieve the target outlet temperature (45°C). After stabilization of the temperatures, the feed of the spray dryer was commuted from purified water to the product suspension (keeping the same flow rate used during stabilization) and the inlet temperature once again adjusted in order to achieve the target outlet temperature. At the end of the stock suspension, the feed was once more commuted to purified water in order to rinse the feed line and perform a controlled shut down. The dry product in the collection flasks under both cyclones was weighed and the yield calculated as the mass percentage of the dry product in relation to the total solids in the suspension fed to the high pressure homogenizer.

[0209] Particle size distribution was analyzed by laser diffraction. Solid state characterization (for polymorphic form and purity) was performed by high pressure liquid chromatography (HPLC), X-ray powder diffraction (XRPD), and differential scanning calorimetry (mDSC). Water content was determined by the Karl Fischer method.

[0210] Jet milling produced crystalline rapamycin powder with a monodisperse particle size distribution having Dv10 of 1.5 μm (microns), a Dv50 of 2.7 μm (microns) and a Dv 90 of 4.9 μm (microns), as shown in Table 10 below.

[0211] Wet polishing produced crystalline rapamycin powder with a monodisperse particle size distribution having a Dv10 of 1.0 μm (microns), a Dv50 of 2.4 μm (microns) and a Dv 90 of 5.0 μm (microns) (Table 11).

[0212] Both methods produced particles of rapamycin within the target range and neither process showed an impact on polymorphic form or purity of the rapamycin. The tables below show in-process control data for the jet milling and wet polishing processes. The data indicate that both processes were able to produce API particle sizes within the target range without impacting API purity or polymorphic form.

Table 10: Jet Milling Data

PSD	μm	Dv10	Dv50	Dv90
		1.51	2.74	4.91
XRPD	-	Similar to API (sirolimus) diffractogram		
mDSC	(T _{onset} , °C)	182.2		
KF	% w/w	0.30		
HPLC (% area)	Assay (% w/w)	99.5		
	Sirolimus	99.35		

Table 11: Wet Polishing Data

PSD	μm	Dv10	Dv50	Dv90
		1.05	2.42	4.97
XRPD	-	Similar to API (sirolimus) diffractogram		
mDSC	(T_onset, °C)	185.7		
KF			% w/w	0.30
HPLC (% area)		Assay (% w/w)		99.0
		Sirolimus		99.42

EP 3 125 875 B1

Example 9: Aerosol Performance Testing of Dry Powder Compositions

[0213] The capsules produced in the example above were placed into the device indicated in the tables below and actuated. The aerosol performance delivered from the devices/capsules containing blends from Batch 06RP68.HQ00008 and Batch 06RP68.HQ00009 were characterized using a next generation impactor (NGI) according to the methods described in Chapters 905 and 601 of the USP. The aerosols were tested at flow rates of 60 and 100 liters per minute (LPM). The fine particle dose (FPD) and fine particle fraction (FPF) are shown in the tables below. Mass median aerodynamic diameters (MMAD) and geometric standard deviations (GSD) are also shown.

Table 12: 06RP68.HQ00008 (Wet Polished) + Plasiatpe RS01 Model 7

	60 LPM		100 LPM	
	Mean	%RSD	Mean	%RSD
FPD (μg)	57.31	2.37	67.21	12.46
FPF (%)	39.49	1.85	44.12	8.99
MMAD (μm)	2.81	2.22	2.49	11.97
GSD	2.02	0.99	2.19	8.25

Table 13: 06RP68.HQ00008 (Wet Polished) + Plastiaple RS00 Model 8

	60 LPM		100 LPM	
	Mean	%RSD	Mean	%RSD
FPD (μg)	58.40	0.98	62.39	6.35
FPF (%)	39.68	1.68	41.34	3.70
MMAD (μm)	2.63	7.28	2.58	6.00
GSD	2.05	3.69	2.15	6.32

Table 14: 06RP68.HQ00009 (Jet Milled) + Plastiaple RS01 Model 7

	60 LPM		100 LPM	
	Mean	%RSD	Mean	%RSD
FPD (μg)	52.33	6.72	58.51	15.84
FPF (%)	33.73	3.91	36.69	9.86
MMAD (μm)	3.32	2.27	3.02	4.14
GSD	2.05	1.02	2.24	1.79

Table 15: 06RP68.HQ00009 (Jet Milled)+ Plastiaple RS00 Model 8

	60 LPM		100 LPM	
	Mean	%RSD	Mean	%RSD
FPD (μg)	52.56	2.02	59.11	4.74
FPF (%)	33.97	0.86	36.01	4.20
MMAD (μm)	3.06	1.91	2.93	0.98
GSD	2.04	0.98	2.21	2.73

[0214] Based on these aerosol performance data, the wet polished drug particles are preferred. They resulted in a

higher fine particle dose, higher fine particle fraction, a particle size distribution that would exhibit penetration into both the central and peripheral lung regions, and would have less oral deposition.

Reference Example 10: Pharmacokinetic modeling of rapamycin

[0215] Based on the aerosol performance 06RP68.HQ00008 (Wet Polished) + PlasiTape RS01 Model as shown above, and the results of animal experiments in Example 3, it can be expected that delivery of inhaled rapamycin directly to the lung in humans will similarly result in persistent lung concentrations that are sufficiently high to be therapeutically effective, but with low systemic exposure (low blood concentrations) thereby effectively minimizing side effects due to systemic exposure. A two compartment, pharmacokinetic model was developed to predict the concentrations in the blood and lungs in humans after repeat QD dosing using the formulation and DPI inhaler in Table 11. For the pharmacokinetic model, human PK parameters from the Rapamune® (NDA 21-110, and NDA 21-083) summary basis of approval were used: the volume of distribution was assumed to be 780 liters, clearance was 0.0003/minute, and elimination half-life was 42.3 hours (assuming equivalency to rapamycin IV dosing). Absorption half-life of rapamycin from the lung was estimated to be approximately 0.5 hours, similar to other highly lipophilic compounds, such as fluticasone propionate for which lung absorption data is available. Bioavailability of rapamycin depositing in the lung was assumed to be approximately 100%. Bioavailability of rapamycin absorbed by the GI route through oropharyngeal deposition or removal from the upper airways by mucociliary clearance was assumed to be 14% as reported in the Rapamune® summary basis of approval. For a typical human inspiratory maneuver at a flow rate of 60 liters per minute, as shown in Table 11, the fine particle dose was 57 micrograms, and the fine particle fraction was 40%.

[0216] The model predicts achieving an average steady state concentration after 11 days as shown in Figure 9. From the figure it can be seen that once daily repeat dosing of 57 micrograms delivered to the lungs results in trough blood concentrations of approximately 50 picograms/ml, and maximum concentrations below 200 picograms/ml, substantially below the concentrations of 5-15 ng/ml reported in McCormack et al. (2011), "Efficacy and safety of sirolimus in lymphangioleiomyomatosis", N Engl J Med 364:1595-1606. Assuming a lung tissue mass of 850 grams, no metabolism in the lung and a lung absorption half life of 30 minutes, 57 micrograms rapamycin delivered to the lungs would result in therapeutic levels in the lung tissue, with local lung concentrations of rapamycin as high as approximately 14 ng/gram.

EQUIVALENTS

[0217] Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein.

Claims

1. A pharmaceutical aerosol composition in the form of a dry powder for pulmonary delivery comprising micronized rapamycin particles having a Mass Median Aerodynamic Diameter (MMAD) of from 0.5 to 5 micrometres (μm) (microns) and particles of a carrier for use in a method of treating a chronic lung disease, the method comprising delivering the dry powder composition to a human subject in need thereof by inhalation, wherein the formulation is effective to deliver a therapeutic amount of rapamycin to the lungs, and wherein the chronic lung disease is not Lymphangioleiomyomatosis (LAM).
2. The composition for the use of claim 1, wherein the method comprises administering a total daily dose of from 20 to 250 micrograms of rapamycin.
3. The composition for the use of claim 1 or 2, wherein the amount of rapamycin in the composition is from 0.1 % to 20 % (w/w) or from 0.25 % to 2 % (w/w), based upon total weight of the composition.
4. The composition for the use of claim 1 or 3, wherein the micronized rapamycin particles have an MMAD of from 1 to 5 μm (microns), from 1.5 to 4 μm (microns), or from 1.5 to 3.5 μm (microns).
5. The composition for the use of any of claims 1 to 4, wherein the carrier is selected from the group consisting of arabinose, glucose, fructose, ribose, mannose, sucrose, trehalose, lactose, maltose, starches, dextran, and mannitol.
6. The composition for the use of any of claims 1 to 5, wherein the particles of the carrier have diameters ranging from 1 to 200 μm (microns), from 30 to 100 μm (microns), or less than 10 μm (microns).

7. The composition for the use of any of claims 1 to 6, wherein the carrier comprises or consists of a blend of two different carriers, a first carrier and a second carrier, optionally wherein the carrier consists of a blend of two different lactose carriers.
8. The composition for the use of claim 7, wherein the first carrier consists of particles having diameters ranging from 30-100 μm (microns) and the second carrier consists of particles having diameters of less than 10 μm (microns).
9. The composition for the use of any of claims 1-8, wherein the subject is a geriatric subject.

Patentansprüche

1. Pharmazeutische Aerosolzusammensetzung in der Form eines Trockenpulvers zur pulmonalen Abgabe, umfassend mikronisierte Rapamycin-Partikel mit einem massenmedianen aerodynamischen Durchmesser (MMAD) von 0,5 bis 5 Mikrometer (μm) (Mikron) und Partikel eines Trägers zur Verwendung bei einem Verfahren zum Behandeln einer chronischen Lungenerkrankung, wobei das Verfahren ein Abgeben der Trockenpulverzusammensetzung an ein menschliches Subjekt, das dessen bedarf, durch Inhalation umfasst, wobei die Formulierung wirksam ist, um eine therapeutische Menge Rapamycin an die Lunge abzugeben, und wobei die chronische Lungenerkrankung keine Lymphangioleiomyomatose (LAM) ist.
2. Zusammensetzung zur Verwendung nach Anspruch 1, wobei das Verfahren ein Verabreichen einer täglichen Gesamtdosis von 20 bis 250 Mikrogramm Rapamycin umfasst.
3. Zusammensetzung zur Verwendung nach Anspruch 1 oder 2, wobei die Menge an Rapamycin in der Zusammensetzung von 0,1 Gew.-% bis 20 Gew.-% oder von 0,25 Gew.-% bis 2 Gew.-%, bezogen auf das Gesamtgewicht der Zusammensetzung beträgt.
4. Zusammensetzung zur Verwendung nach Anspruch 1 oder 3, wobei die mikronisierten Rapamycin-Partikel einen MMAD von 1 bis 5 μm (Mikron), von 1,5 bis 4 μm (Mikron) oder von 1,5 bis 3,5 μm (Mikron) aufweisen.
5. Zusammensetzung zur Verwendung nach einem der Ansprüche 1 bis 4, wobei der Träger aus der Gruppe bestehend aus Arabinose, Glucose, Fructose, Ribose, Mannose, Saccharose, Trehalose, Lactose, Maltose, Stärken, Dextran und Mannit ausgewählt ist.
6. Zusammensetzung zur Verwendung nach einem der Ansprüche 1 bis 5, wobei die Partikel des Trägers Durchmesser im Bereich von 1 bis 200 μm (Mikron), von 30 bis 100 μm (Mikron) oder weniger als 10 μm (Mikron) aufweisen.
7. Zusammensetzung zur Verwendung nach einem der Ansprüche 1 bis 6, wobei der Träger eine Mischung aus zwei unterschiedlichen Trägern, einem ersten Träger und einem zweiten Träger umfasst oder daraus besteht, wobei optional der Träger aus einer Mischung aus zwei unterschiedlichen Lactose-Trägern besteht.
8. Zusammensetzung zur Verwendung nach Anspruch 7, wobei der erste Träger aus Partikeln besteht, die Durchmesser im Bereich von 30-100 μm (Mikron) aufweisen, und der zweite Träger aus Partikeln besteht, die Durchmesser von weniger als 10 μm (Mikron) aufweisen.
9. Zusammensetzung zur Verwendung nach einem der Ansprüche 1-8, wobei das Subjekt ein geriatrisches Subjekt ist.

Revendications

1. Composition pharmaceutique en aérosol sous la forme d'une poudre sèche pour administration pulmonaire comprenant des particules de rapamycine micronisées ayant un diamètre aérodynamique médian en masse (MMAD) de 0,5 à 5 micromètres (μm) (microns) et des particules d'un support pour une utilisation dans un procédé de traitement d'une maladie pulmonaire chronique, le procédé comprenant l'administration de la composition de poudre sèche à un sujet humain en ayant besoin par inhalation, dans laquelle la formulation est efficace pour administrer une quantité thérapeutique de rapamycine aux poumons, et dans laquelle la maladie pulmonaire chronique n'est pas la lymphangioleiomyomatose (LAM).

EP 3 125 875 B1

2. Composition pour l'utilisation selon la revendication 1, dans laquelle le procédé comprend l'administration d'une dose quotidienne totale de 20 à 250 microgrammes de rapamycine.
3. Composition pour l'utilisation selon la revendication 1 ou 2, dans laquelle la quantité de rapamycine dans la composition est de 0,1 % à 20 % (p/p) ou de 0,25 % à 2 % (p/p), sur la base du poids total de la composition.
4. Composition pour l'utilisation selon la revendication 1 ou 3, dans laquelle les particules de rapamycine micronisées ont un MMAD de 1 à 5 μm (microns), de 1,5 à 4 μm (microns) ou de 1,5 à 3,5 μm (microns).
5. Composition pour l'utilisation selon l'une quelconque des revendications 1 à 4, dans laquelle le support est choisi dans le groupe constitué de l'arabinose, du glucose, du fructose, du ribose, du mannose, du saccharose, du tréhalose, du lactose, du maltose, des amidons, du dextrane et du mannitol.
6. Composition pour l'utilisation selon l'une quelconque des revendications 1 à 5, dans laquelle les particules du support ont des diamètres allant de 1 à 200 μm (microns), de 30 à 100 μm (microns) ou inférieurs à 10 μm (microns).
7. Composition pour l'utilisation selon l'une quelconque des revendications 1 à 6, dans laquelle le support comprend ou est constitué d'un mélange de deux supports différents, un premier support et un second support, éventuellement dans laquelle le support est constitué d'un mélange de deux supports de lactose différents.
8. Composition pour l'utilisation selon la revendication 7, dans laquelle le premier support est constitué de particules ayant des diamètres allant de 30 à 100 μm (microns) et le second support est constitué de particules ayant des diamètres inférieurs à 10 μm (microns).
9. Composition pour l'utilisation selon l'une quelconque des revendications 1 à 8, dans laquelle le sujet est un sujet gériatrique.

LC-MS/MS Chromatogram of 10.6 ng/mL Rapamycin (top) and Internal Standard
(bottom) in Mouse Blood




FIG. 1

Representative Chromatograms of 10.6 ng/mL Rapamycin (top) and Internal Standard
(bottom) in Mouse Lung Homogenate




FIG. 2

Calibration Curve for Rapamycin in Mouse Blood




FIG. 3

Calibration Curve for Rapamycin in Mouse Lung Homogenate




FIG. 4

Representative Chromatograms of Rapamycin (top) and Internal Standard (bottom) in
Blood from Mouse 2-07 Administered Rapamycin by OPA Administration




FIG. 5

Representative Chromatograms of Rapamycin (top) and Internal Standard (bottom) in Lung Homogenate from Mouse 2-07 Administered Rapamycin by OPA Administration




FIG. 6






FIG. 7

LAM-001 Lung Samples (S6 Phosphorylation)**FIG. 8A**




FIG. 8B




FIG. 9

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 3929992 A [0002]
- US 20100305150 A, Berg [0003]
- US 6384046 B [0003]
- US 6258823 B [0003]
- US 20130004436 A, Chhajed [0006]
- US 5080899 A, Sturm [0006]
- US 5635161 A [0006]
- WO 2011163600 A [0007]
- US 6419900 B [0009]
- US 6419901 B [0009]
- US 6451784 B [0009]
- US 6793912 B [0009]
- US 20030059375 A [0009]
- US 20040039047 A [0009]
- US 7288243 B [0009]
- US 20080008662 A [0009]
- US 20100260733 A [0010]
- US 8492110 B [0011]
- US 20130102569 A [0012]
- WO 2008137148 A [0013]
- WO 2008063581 A [0014]
- US 7384953 B [0068]
- WO 9703654 A [0069]
- US 5989591 A [0069]
- WO 06039237 A [0069]
- WO 06094507 A [0069]
- US 8053444 B [0069]
- US 20130203717 A [0095]

Non-patent literature cited in the description

- **NISHIMURA, T. et al.** *Am. J. Respir. Crit. Care Med.*, 2001, vol. 163, 498-502 [0003]
- **NOCERA et al.** Sirolimus Therapy in Liver Transplant Patients: An Initial Experience at a Single Center. *Transplantation Proceedings*, 2008, vol. 40 (6), 1950-1952 [0004]
- **PEREZ et al.** Interstitial Pneumonitis Associated With Sirolimus in Liver Transplantation: A Case Report. *Transplantation Proceedings*, 2007, vol. 39 (10), 3498-3499 [0004]
- **HASHEMI-SADRAEI et al.** Sirolimus-associated diffuse alveolar hemorrhage in a renal transplant recipient on long-term anticoagulation. *Clinical Nephrology*, 2007, vol. 68 (4), 238-244 [0004]
- **PEDROSO et al.** Pulmonary alveolar proteinosis - a rare pulmonary toxicity of sirolimus. *Transplant International*, 2007, vol. 20 (3), 291-296 [0004]
- **APARICIO et al.** *Clinical & Translational Oncology*, 2009, vol. 11 (8), 499-510 [0005]
- **VAHID et al.** Pulmonary complications of novel anti-neoplastic agents for solid tumors. *Chest*, 2008, vol. 133, 528-538 [0005]
- **IACOVELLI et al.** Incidence and risk of pulmonary toxicity in patients treated with mTOR inhibitors for malignancy. A meta-analysis of published trials. *Acta oncologica*, 2012, vol. 51 (7), 873-879 [0005]
- **YI et al.** *J. Aerosol Med. Pulm. Drug Deliv.*, 2010, vol. 23, 181-7 [0009]
- **M. CHOUGALE et al.** *Int. J. Nanomedicine*, 2007, vol. 2, 625-688 [0100]
- **CHEMICAL ABSTRACTS**, 9003-11-6 [0146]
- **CHEMICAL ABSTRACTS**, 53123-88-9 [0169]
- **P. J. MURPHY.** Contact bedding was Sani-Chips hardwood chips. Forest Products Co, [0170]
- **YANEZ, J.** Pharmacometrics and Delivery of Novel Nanoformulated PEG-b-poly(ϵ -caprolactone) Micelles of Rapamycin. *Cancer Chemotherapy and Pharmacology*, 2007, vol. 61 (1), 133-144 [0194]
- **NAPOLI, K.** Distribution of Sirolimus in Rat Tissue. *Clinical Biochemistry*, 1997, vol. 30 (2), 135-142 [0195]
- **MCCORMACK et al.** Efficacy and safety of sirolimus in lymphangioleiomyomatosis. *N Engl J Med*, 2011, vol. 364, 1595-1606 [0216]