



Europäisches
Patentamt
European
Patent Office
Office européen
des brevets



(11)

EP 3 165 232 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
10.05.2017 Bulletin 2017/19

(51) Int Cl.:
A61K 38/00 (2006.01) **A61K 38/48** (2006.01)
C12N 9/64 (2006.01)

(21) Application number: 16196780.7

(22) Date of filing: 23.12.2013

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: 24.12.2012 US 201261745674 P
15.03.2013 US 201361787026 P

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
13867801.6 / 2 938 351

(71) Applicant: **Bayer Healthcare LLC**
Whippany, NJ 07981-0915 (US)

(72) Inventors:
• **Bauzon, Maxine**
Hercules, CA 94547 (US)
• **Hermiston, Terry**
Mill Valley, CA 94941 (US)

(74) Representative: **BIP Patents**
c/o Bayer Intellectual Property GmbH
Alfred-Nobel-Straße 10
40789 Monheim am Rhein (DE)

Remarks:

This application was filed on 02-11-2016 as a divisional application to the application mentioned under INID code 62.

(54) SHORT-ACTING FACTOR VII POLYPEPTIDES

(57) Short-acting Factor VII peptides are disclosed. A shortened half-life is desirable for treatment of acute bleeding and similar disorders. Modification of the sia-

lylation and/or glycosylation of Factor VII and variants thereof produced peptides useful in treating conditions of acute bleeding.






Figure 3

Description

CROSS REFERENCE TO RELATED APPLICATIONS

5 **[0001]** This application claims priority to U.S App. Ser. No. 61/745,674, filed December 24, 2012, and U.S. App. Ser. No. 61/787,026, filed March 15, 2013, which are hereby incorporated by reference in their entirety.

FIELD OF THE DISCLOSURE

10 **[0002]** Human coagulation Factor VII variants and the polynucleotides encoding such variants, vectors and host cells comprising and expressing such variants, methods of obtaining such variants, methods of using such variants, compositions of the variants, and additional inventive features related thereto are provided herein.

BACKGROUND

15 **[0003]** Blood coagulation is a process consisting of a complex interaction of various blood components (or factors) that eventually gives rise to a fibrin clot. Generally, the blood components, which participate in what has been referred to as the coagulation "cascade," are enzymatically inactive proteins (proenzymes or zymogens) that are converted to proteolytic enzymes by the action of an activator (which itself is an activated clotting factor). Coagulation factors that have undergone such a conversion are generally referred to as "active factors" and are designated by the addition of the letter "a" to the name of the coagulation factor (e.g., Factor VIIa).

20 **[0004]** Initiation of the haemostatic process is mediated by the formation of a complex between tissue factor, which is exposed to the circulating blood following injury to the vessel wall, and Factor VIIa, which is present in the circulation in an amount corresponding to about 1% of the total Factor VII protein mass. This complex is anchored to the tissue factor-bearing cell and converts Factors IX and X to their active forms Factor IXa and Factor Xa on the cell surface. Factor Xa converts prothrombin to thrombin on the tissue factor-bearing cell, which activates Factor VIII, Factor V, Factor XI, and Factor XIII. Furthermore, the limited amount of thrombin formed in this initial step of haemostasis also activates the platelets. Following the action of thrombin on the platelets, the platelets change shape and expose charged phospholipids on their surface. This activated platelet surface forms the template for further Factor X activation and the full thrombin generation. The further Factor X activation on the activated platelet surface occurs via a Factor IXa and Factor VIIa complex formed on the surface of the activated platelet, and Factor Xa then converts prothrombin into thrombin while still on the surface. Thrombin then converts fibrinogen into fibrin, which is insoluble and which stabilizes the initial platelet plug. This process is localized to the site of the tissue factor exposure thereby minimizing the risk of a systemic activation of the coagulation system. In recent years, Factor VII and tissue factor have been found to be the main initiators of blood coagulation.

30 **[0005]** Factor VIIa is produced from its precursor, Factor VII, which is synthesized in the liver and secreted into the blood where it circulates as a single-chain glycoprotein (molecular weight of about 50,000 Da). Wild-type Factor VII as used herein has the amino acid sequence and nucleotide sequence disclosed in FIGs. 1 and 2. The term "Factor VII" is meant to encompass Factor VII polypeptides in their uncleaved form (the zymogen form) as well as those that have been proteolytically or otherwise processed to yield their respective bioactive forms, which may be referred to as Factor VIIa. Wild type Factor VII is cleaved typically between residues 152 and 153 to produce Factor VIIa.

35 **[0006]** Factor VII is converted *in vitro* into the two-chain form Factor VIIa by Factor Xa, Factor XIIa, Factor IXa, or thrombin. Like several other plasma proteins involved in haemostasis, Factor VII is dependent on Vitamin K for its activity, which is required for the gamma-carboxylation of multiple glutamic acid residues that are clustered close to the amino terminus of the protein. These gamma-carboxylated glutamic acids are required for the metal ion-induced interaction of Factor VII with phospholipids. In the presence of tissue factor, phospholipids, and calcium ions, the two-chain Factor VIIa rapidly activates Factor X or Factor IX by limited proteolysis. Factor VIIa is susceptible to proteolytic cleavage, giving rise to a number of degradation products that do not have clotting activity.

40 **[0007]** Factor VII variants having an amino acid sequence derived from wild type Factor VII by substitution, deletion, and/or insertion of one or more amino acids have been published. For example, Dickinson et al. (Proc. Natl. Acad. Sci. USA (1996) 93, 14379-14384) relates to Factor VII variants wherein Lys157, Val158, Glu296, Met298, Asp334, Ser336, or Lys227 have been individually replaced by Ala. Iwanaga et al. (Thromb. Haemost. (supplement August 1999), 466, abstract 1474) relates to Factor VIIa variants wherein residues 316-320 are deleted or residues 311-322 are replaced with the corresponding residues from trypsin. U.S. Pat. App. Pub. 2008/0058255 A1 to Bolt et al. relates to Factor VII variants having a glycosylation-disrupting substitution at either N145 or N322, or at both N145 and N322. Toso et al. reported a series of Factor VII structure-function studies based on naturally occurring mutations. The mutant recombinant Factor VII proteins included T324M, E385K, and two mutant Factor VII proteins lacking glycosylation core sequences in either the Factor VII heavy chain (N322Q) or the Factor VII light chain (N145Q). Toso et al., "Lack of Heavy Chain

Glycosylation in Patient with Factor VII Deficiency Not Responsible for Mutant FVIIa Activity," Blood, vol. 96, no. 11, part 2 (16 Nov. 2000), p. 79b (42nd Annual Meeting of the American Society of Hematology).

[0008] Most naturally occurring peptides and proteins contain carbohydrate moieties attached to the peptide or protein via specific linkages to a select number of amino acids along the length of the primary peptide or protein chain. Thus, many naturally occurring peptides and proteins are termed "glycopeptides" or "glycoproteins," respectively. The variability of the glycosylation pattern on any given peptide or protein can impact the function of that peptide or protein. For example, the structure of the N-linked glycans on a peptide or protein can impact various characteristics of the peptide or protein, including the protease susceptibility, intracellular trafficking, secretion, tissue targeting, biological half-life, and antigenicity of the peptide or protein in a cell or organism. The alteration of one or more of these characteristics can affect the efficacy of a peptide or protein in its natural setting, and can also affect the efficacy of the peptide or protein as a therapeutic agent in situations where the peptide or protein has been generated for that purpose.

[0009] The carbohydrate structure attached to the peptide or protein chain is known as a "glycan" molecule. The specific glycan structure present on a peptide or protein affects the solubility and aggregation characteristics of the peptide or protein, the folding of the primary peptide or protein chain, and, therefore, its functional or enzymatic activity, the resistance of the peptide or protein to proteolytic attack, and the control of proteolysis leading to the conversion of inactive forms of the peptide or protein to active forms. For example, terminal sialic acid residues present on the glycan molecule affect the length of the half-life of the peptide or protein in the mammalian circulatory system. Peptides and proteins whose glycans do not contain terminal sialic acid residues generally are more rapidly removed from the circulation by the liver.

[0010] The glycan structures found in naturally occurring glycopeptides and glycoproteins are typically divided into two classes, N-linked and O-linked glycans. Wild type Factor VIIa contains two N-linked and two O-linked glycosylation sites. N-linked glycosylation is the most common covalent modification in eukaryotes. N-linked glycosylation occurs at the consensus sequence Asn-X-Ser/Thr, where the glycan attaches to the amine group of asparagine and X represents any amino acid except proline. N-linked glycans are based on the common core pentasaccharide, Man₃(GlcNAc)₂, which can be further modified by the addition of monosaccharides such as N-acetyl galactosamine, galactose, neuraminic acid, N-acetylglucosamine, fructose, mannose, and fucose. The Man₃(GlcNAc)₂ core with various monosaccharides including terminal sialic acids may be attached via a N-acetylglucosamine to at the Asn in the Asn-X-Ser/Thr consensus sequence. This chemically complex co-translational modification serves many purposes and affects the biology of the protein in diverse ways including proper folding, functional group orientation, and clearance rates.

[0011] A variety of methods have been proposed in the art to customize the glycosylation pattern of a peptide or protein, including those described in U.S. Patent No. 8,008,252 to DeFrees et al.

[0012] It is often desirable to stimulate or improve the coagulation cascade in a subject. Factor VIIa has been used to control bleeding disorders caused by clotting factor deficiencies (e.g., haemophilia A and B or deficiency of coagulation Factors XI or VII) or clotting factor inhibitors. Recombinant Factor VIIa, manufactured and sold by Novo Nordisk under the trade name NovoSeven®, is approved for the treatment of bleeding episodes in hemophilia A or B patients with inhibitors to Factor VIII or Factor IX and in patients with acquired hemophilia; prevention of bleeding in surgical interventions or invasive procedures in hemophilia A or B patients with inhibitors to Factor VIII or Factor IX and in patients with acquired hemophilia; treatment of bleeding episodes in patients with congenital Factor VII deficiency and prevention of bleeding in surgical interventions or invasive procedures in patients with congenital Factor VII deficiency. U.S. Pat. No. 5,180,583 to Hedner discloses using Factor VIIa to control excessive bleeding in situations not caused by clotting factor defects or clotting factor inhibitors. Hedner discloses treating bleeding disorders caused for example by a defective platelet function, thrombocytopenia, or von Willebrand's disease, and compositions for those uses.

[0013] There is a need to treat bleeding from disorders not caused by congenital or developed clotting factor deficiencies or inhibitors to clotting factors. Several clinical trials have demonstrated the efficacy of recombinant Factor VIIa to control bleeds. However, there are concerns over an increase in undesirable thromboembolic events from use of this molecule. Bleeding is a major problem in many disorders, such as in connection with surgery, complications following surgery, stem and organ transplants, intracranial hemorrhage, aortic aneurysm, and trauma, or overdose of certain anti-coagulants.

50 BRIEF SUMMARY

[0014] It is an object to treat bleeding disorders and episodes with Factor VII polypeptides that are short-acting. One object of the present work is to provide compositions of Factor VII polypeptides (wild-type or variant) that are short-acting, characterized by one or more pharmacokinetic traits such as a shortened half-life. It is an object to provide such a Factor VII molecule with reduced opportunity for thrombotic events outside of the target site and the treatment time-frame. It is an object to provide Factor VII polypeptides (wild-type or variant) with enhanced clearance due to altered glycosylation patterns.

[0015] Described herein is a composition of variant Factor VII polypeptides, in which the variant Factor VII polypeptide

comprises an amino acid sequence having at least two sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least two sequence alterations are (1) a glutamine residue substituted for the proline residue in position 10, and (2) a glutamic acid residue substituted for the lysine residue in position 32; and wherein the ratio of moles of conjugated sialic acid to moles of N-linked glycan in the composition is less than 0.05, less than 0.1, less than 1.0, less than 2.0, less than 3.0, less than 4.0, less than 5.0 or less than 6.0. Also described herein is a composition of variant Factor VII polypeptides in which the variant Factor VII polypeptide comprises an amino acid sequence having at least two sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least two sequence alterations are (1) a glutamine residue substituted for the proline residue in position 10, and (2) a glutamic acid residue substituted for the lysine residue in position 32; and wherein the ratio of moles of conjugated sialic acid per mole of N-linked glycan is within a range selected from the group consisting of (1) from 0 to 5; (2) from 0 to 4; (3) from 0 to 3; (4) from 0 to 2; (5) from 0 to 1 and (6) from 0 to 0.5.

[0016] Also described herein is an isolated variant Factor VII polypeptide comprising an amino acid sequence having at least two sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least two sequence alterations are (1) a glutamine residue substituted for the proline residue in position 10, and (2) a glutamic acid residue substituted for the lysine residue in position 32, wherein the polypeptide has a ratio of moles of conjugated sialic acid to moles of N-linked glycan of less than 0.05, less than 0.1, less than 1.0, less than 2.0, less than 3.0, less than 4.0, less than 5.0 or less than 6.0. Also described herein is a composition of Factor VII polypeptides, wherein the Factor VII polypeptides comprise the amino acid sequence of SEQ ID NO: 16 (wild type Factor VII) and the ratio of moles of conjugated sialic acid to moles of N-linked glycan in the composition is within a range selected from the group consisting of (1) from 1 to 5; (2) from 1 to 4; (3) from 1 to 3; (4) from 1 to 2; and (5) from 0.5 to 1; or conjugated sialic acid is undetectable.

[0017] Also described herein is an isolated variant Factor VII polypeptide selected from the group consisting of:

- (1) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, and (3) a sequence alteration such that N-linked glycosylation at position 145 is disrupted;
- (2) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, and (3) a sequence alteration such that N-linked glycosylation at position 322 is disrupted;
- (3) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, and (2) a glutamic acid residue substituted for the lysine residue in position 32, and (3) sequence alterations such that N-linked glycosylation at positions 145 and 322 is disrupted;
- (4) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, and (2) a glutamic acid residue substituted for the lysine residue in position 32, wherein positions 145 and 322 are asparagine and have attached N-linked glycosylation;
- (5) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the alanine residue in position 34, (4) a glutamic acid residue substituted for the arginine residue in position 36, and (5) a sequence alteration such that N-linked glycosylation at position 145 is disrupted;
- (6) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the alanine residue in position 34, (4) a glutamic acid residue substituted for the arginine residue in position 36, and (5) a sequence alteration such that N-linked glycosylation at position 322 is disrupted;
- (7) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the alanine residue in position 34, (4) a glutamic acid residue substituted for the arginine residue in position 36, and (5) a sequence alterations such that N-linked glycosylation at positions 145 and 322 is disrupted; and
- (8) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino

acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the alanine residue in position 34, and (4) a glutamic acid residue substituted for the arginine residue in position 36, wherein positions 145 and 322 are asparagine and have attached N-linked glycosylation.

[0018] Also described herein are Factor VII polypeptides having reduced conjugation of sialic acid with the Factor VII polypeptide. In certain examples, the Factor VII polypeptide is a variant polypeptide that produces altered glycosylation pattern. In other examples, the Factor VII polypeptide is a wild-type Factor VII polypeptide which has reduced conjugation of sialic acid. In certain embodiments, reduced sialic acid conjugation can be effectuated by treatment of the polypeptide with a sialidase enzyme. In other embodiments, reduced sialic acid conjugation can be effectuated by producing recombinant Factor VII polypeptides in a cell line that is partially or completely deficient in sialylation of peptides. In further embodiments, the reduced sialic acid conjugation can be effectuated by coexpressing the recombinant Factor VII polypeptide and a recombinant or exogenous sialidase enzyme in a cell line.

[0019] Also described is a method for treating a mammal having a disease or a disorder wherein blood clot formation is desirable, comprising administering to a mammal in need thereof an effective amount of a Factor VII polypeptide that has reduced sialic acid conjugation. In certain embodiments, the ratio of moles of conjugated sialic acid to moles of N-linked glycan is less than 0.05. In other embodiments, the Factor VII polypeptide comprises the amino acid sequence of SEQ ID NO: 16. In further embodiments, the Factor VII polypeptide comprises wild-type factor VII. In additional embodiments, the disease or disorder being treated is selected from the group consisting of a hemorrhage, gastrointestinal bleeding, uncontrolled bleeding, bleeding in a mammal undergoing transplantation or resection or surgery, variceal bleeding, thrombocytopenia, hemophilia, intracranial hemorrhage, aortic aneurysm, and over administration of an anti-coagulant.

[0020] Further variants, compositions, methods and related products and processes are disclosed in detail below.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021]

FIGS. 1A-1H show the nucleotide sequences for three Factor VII molecules used in the present application. "V1" is a variant of human Factor VII having four amino acid mutations relative to wild type human amino acid sequence of SEQ ID NO: 16 : (P10Q, K32E, T106N and V253N). "V2" is a variant of human Factor VII having six amino acid mutations relative to wild type human amino acid sequence of SEQ ID NO: 16 : (P10Q, K32E, A34E, R36E, T106N and V253N). FIG. 1 also shows the nucleotide sequences for various constructs used in the examples.

FIG. 2 shows the amino acid sequences for three Factor VII molecules used in the present application. Wild type human Factor VII as used herein has the amino acid sequence of SEQ ID NO: 16. V1 has the amino acid sequence of SEQ ID NO: 17. V2 has the amino acid sequence of SEQ ID NO: 18. In V1 and V2, the changes from wild type Factor VII of SEQ ID NO: 16 are shown in bold.

FIG. 3 is a scheme depicting three Factor VII molecules used in the examples of the present application. Attachment of glycans at N-glycosylation sites is shown. For the depiction of the glycans, a solid box represents N-acetylgalactosamine, a shaded oval represents mannose, an open oval represents galactose, a dark diamond represents sialic acid (also known as N-acetylneurameric acid) and a closed triangle represents fucose. The glycan structure is a depiction using one possible variant of a glycan and does not represent an actual measured glycan.

FIG. 4 is a scheme depicting an N-linked glycan showing attachment at the Asn in the Asn-X-Ser/Thr consensus sequence. The $\text{Man}_3(\text{GlcNAc})_2$ core with various monosaccharides including terminal sialic acids are shown.

FIG. 5 is a scheme depicting two approaches used in the present disclosure to decrease the half-life of Factor VII variants, exemplified with reference to V2.

FIG. 6 is a table of hypoglycosylated Factor VII molecules.

FIG. 7 shows the results of an LC-MS method to identify the sialic acid remaining on the heavy chain of V2 after desialylation according to the conditions of the experiment.

FIG. 8 shows the analysis of desialylated V2 for sialic acid content.

FIG. 9 shows the results of a phospholipid FX activation assay.

FIG. 10 shows the results of a PL-TGA assay on desialylated proteins.

FIG. 11 shows expression of hypoglycosylated Factor VII variants.

FIG. 12 is a table showing determination of "specific activity" of hypoglycosylated FVII variants using transfection supernatants.

FIG. 13 shows the results of a PL-TGA assay on purified hypoglycosylated variant pMB121.

FIG. 14 shows the *in vitro* hepatocyte clearance of desialylated V2 compared to wild type Factor VII.

FIG. 15 shows the results of *in vitro* hepatocyte clearance with Factor VII variants. The hypoglycosylated variants did not display an increase in clearance in this model. This result suggests a different clearance mechanism for these molecules from that utilized by the desialylated V2.

5 FIG. 16 shows pharmacokinetic study results in rat. Half-lives of the desialylated V2 and V1 were significantly shorter than their unmodified parental molecules in Sprague Dawley rats as measured by Factor VII ELISA.

FIG. 17 shows a pharmacokinetic study results in HemA mice.

FIG. 18 shows a desialylated V2 efficacy study in HemA mice.

FIG. 19 shows a desialylated V2 efficacy study in TVT HemA model.

10 FIG. 20 shows the results of thrombin-antithrombin ("TAT") generation in HemA mice with desialylated V2 compared to Factor VII.

FIG. 21 shows a desialylated V2 efficacy study in coagulation-competent mice.

FIG. 22 shows the *in vitro* hepatocyte clearance of desialylated wild-type Factor VII (dWT VIIa) compared to wild-type Factor VII with normal conjugation of sialic acid.

15 FIG. 23 shows tail cut study results in the human tissue factor knock-in (TFKI) mice for dWT VIIa compared to wild-type Factor VII. Desialylated Factor VII was found to be significantly more efficacious than wild-type Factor VII.

FIG. 24 shows results of an ELISA analysis of Thrombin Anti-Thrombin (TAT) complexes after administration of either dWT VIIa or wild-type Factor VII

FIG. 25 shows the results of an analysis of thrombus formation in a FeCl_3 thrombosis model. The given dose of dWT VIIa produced greatly reduced thrombus formation as compared to wild-type Factor VII.

20 FIG. 26 shows the apparent binding affinities of dWT VIIa and wild-type Factor VII for soluble tissue factor as measured by a fluorogenic substrate.

FIG. 27 shows the conversion of Factor X to Factor Xa by a complex of soluble tissue factor and either dWT VIIa or wild-type Factor VII.

25 DETAILED DESCRIPTION

[0022] Methods for modulating the pharmacokinetics of recombinant Factor VII polypeptides (wild-type or variant) to limit thrombotic complications in treatment of acute bleeding are described herein. Also described are Factor VII polypeptides with reduced sialic acid conjugation. Further described are variants of recombinant Factor VII with enhanced clearance from the blood and a decrease in the duration of efficacy. Such variants have a shorter half-life *in vivo* than recombinant wild type Factor VII, due to altered glycosylation patterns. Also described are methods of production and use of such short-acting Factor VII polypeptides.

[0023] To explain Factor VII and glycosylation, FIGs. 3 and 4 are provided. FIG. 3 shows schematically three examples of Factor VII molecules with their domains. Factor VII is a protein consisting of a Gla, EGF, and catalytic domain and containing 2 N-linked Glycans (N145 and N322). V1 is a Factor VII variant with four mutations (P10Q, K32E, T106N, V253N). V2 is a Factor VII variant with six mutations (P10Q, K32E, A343, R36E, T106N, V253N). V1 and V2 both have increased affinity for activated platelets and contain two additional N-glycosylation sites resulting in longer half-lives as compared to wild type Factor VII. The two mutations found solely in V2 (A34E, R36E) are believed to account for its tissue-factor-independence.

[0024] FIG. 4 shows schematically an example of an N-linked glycan showing attachment at the Asn in the Asn-X-Ser/Thr consensus sequence. The $\text{Man}_3(\text{GlcNAc})_2$ core with various monosaccharides including terminal sialic acids are shown.

[0025] Methods of preparing a Factor VII polypeptide having a desired short half-life is provided herein. Two general methods are provided to make a short-acting Factor VII polypeptide, which methods can be used separately or in combination. As shown schematically in FIG. 5 using one example of a Factor VII variant, a glycosylated Factor VII variant can be processed by desialylation or deglycosylation to alter the glycosylation pattern of the variant and thereby to alter, and preferably shorten, its half-life. This method could also be used to desialylate a wild-type Factor VII polypeptide.

[0026] Desialylation may occur by any method known in the art. Examples of suitable methods include enzymatic desialylation by contact with any known enzyme that functions to desialylate including, without limitation, sialidases including neuraminidase-agarose beads (Sigma N5254) and the neuraminidase from *Clostridium perfringens* identified at GI:40479 and in FEBS Lett. 238 (1), 31-34 (1988). Such desialylation may be accomplished by contacting a partially purified recombinant Factor VII polypeptide with a sialidase *in vitro* under suitable conditions, or by co-expression of the sialidase in the host cell expressing the recombinant Factor VII polypeptide. The contacting *in vitro* may be of such duration that only partial desialylation occurs. For example, where a desired half-life can be obtained from a molecule with a ratio of from 0.5 to 1 moles of conjugated sialic acid to moles of N-linked glycan in the composition of Factor VII polypeptides, then contacting with a sialidase for a limited period of time before full desialylation occurs is recommended. Partial desialylation may also be obtained by using a modified sialidase, by contacting the Factor VII polypeptide with

the sialidase under conditions that slow or impair the full functioning of the sialidase, or by other methods apparent to those skilled in the art to produce only partially desialylated polypeptides. Partial desialylation may be measured by comparison to the ratio of conjugated sialic acid to glycan in a reference preparation having been fully desialylated.

[0027] Desialylation may also be accomplished through expression of the Factor VII polypeptide (wild-type or variant) in a cell line that lacks or is deficient in one or more cellular components needed for sialic acid addition. Certain cell lines have been or may be modified to reduce or remove sialylation. For example, Lec2 cells with Chinese hamster ovary ("CHO") origin produce glycoproteins with approximately ten-fold less sialic acid than the wild type cell. It is believed that desialylation of a glycan results in a molecule that can be actively cleared by liver receptors including the Asialoglycoprotein Receptor (ASGPR) and for this reason it shortens half-life.

[0028] The second approach is to deglycosylate a Factor VII variant and thereby obtain a molecule with a shortened half-life. Reduction in glycosylation enhances clearance of Factor VII through renal clearance (50-60Kd cut off, rev. in Caliceti P and Veronese FM, "Pharmacokinetic and biodistribution properties of poly(ethylene glycol)-protein conjugates," *Adv Drug Deliv Rev.* 2003;55(10):1261-77, Weinstein T et al., "Distribution of glycosaminoglycans in rat renal tubular epithelium," *J Am Soc Nephrol.* 1997;8(4):586-95, Choi HS et al., "Renal clearance of quantum dots," *Nat Biotechnol.* 2007;25(10):1165-70), surface charge and isoelectric point (pI) change (which have been related to the increase in glycoprotein circulation, see review in Byrne B. et al., "Sialic acids: carbohydrate moieties that influence the biological and physical properties of biopharmaceutical proteins and living cells," *Drug Discovery Today* 2007;12(7-8):319), and through less glycoprotein-mediated protection from any number of plasma proteases (Ton G., et al., 2005, Nie Y et al., 2006).

[0029] Deglycosylation as used herein includes, without limitation, a genetic modification of a Factor VII polypeptide that results in an altered amino acid sequence as compared to a reference Factor VII polypeptide, which alteration removes an N-linked glycosylation site. For example, a Factor VII variant can be produced with a glycosylation-disrupting alteration at one or more amino acid residues required for the N-linked glycan consensus sequence, i.e., Asn-X-Ser/Thr where X represents any amino acid except proline. As used herein a "glycosylation-disrupting alteration" of a Factor VII amino acid sequence refers to an alteration relative to wild type Factor VII that results in a substitution, addition, or deletion of one or more amino acid residues and that results in a loss of one or more sites for N-linked glycosylation. For example, N-linked glycosylation sites may be removed by replacing N145 and/or N322, both present in wild type Factor VII, with any amino acid (naturally occurring or non-naturally occurring). Glycosylation sites should be identified that have minimal effect on activity when altered to disrupt glycosylation. In another example, deglycosylation may occur by expression of the Factor VII polypeptide (wild-type or variant) in a cell line that lacks the machinery for glycosylation. For example, Factor VII produced in bacterial cells is expected to be completely unglycosylated because bacterial cells lack the cellular machinery for glycosylation. In another embodiment, the Factor VII polypeptide is produced in a cell line that lacks terminal glycosylation enzymes or that has such enzymes but one or more have activity that is less than that found in the wild type cell line. See, e.g., Appa R. et al., 201, Narita M et al., 1998, Seested et al., 2010. In another embodiment, the Factor VII polypeptide is produced in a cell line that harbors a defect in an enzyme involved in the synthesis or attachment of a glycan to Factor VII or a defect in an enzyme involved in the synthesis of CMP-sialic acid transporter. In another embodiment, the Factor VII polypeptide is treated with deglycosylase or chemicals to deglycosylate.

[0030] Treatment by sialidase, deglycosylase, or chemicals to reduce or remove glycans from a Factor VII polypeptide may occur during expression, purification, or post-purification.

[0031] In one embodiment, at least one of the N-linked glycosylation sites in Factor VII variant V1 (N322, N145) or Factor VII variant V2 (N322, N145, N106, N253) was selectively removed with a minimal effect on activity. The N-glycan site was obliterated at the DNA level by disrupting the N-glycan consensus sequence. This was done by removal of the N (Asparagine) codon and replacement with the Q (Glutamine) codon. FIG. 6 is a table showing examples of hypoglycosylated variants. Glycosylation variants were made on wild type Factor VII (referred to herein as "F7"), V1, and V2 backbones. The engineered N-Glycan sites (N106, N253) in V1 and V2 were reverted back to their wild type sequence (T106, V253). Variants pMB113, pMB117, and pMB121 are wild type Factor VII, V1, and V2 constructs respectively containing the two endogenous N-glycosylation sites (N145, N322). All other variants in FIG. 6 have had one or both of their endogenous N-glycan sites removed by introducing N to Q mutations (N145Q, N322Q). This deglycosylation approach results in faster clearance.

[0032] In one aspect of the present disclosure, deglycosylation and desialylation are combined to result in Factor VII polypeptides having desirable shortened half-lives. For example, a Factor VII molecule may be genetically modified to include additional N-linked glycosylation sites beyond the two present in wild type Factor VII. This variant may then be desialylated using one of the methods described herein. The resulting molecule may then retain the glycan structure at each N-linked glycosylation site without the terminal sialic acid. In experiments reported herein, Applicants report such variants that have a faster elimination time than a similar desialylated Factor VII variant that had fewer N-linked glycosylation sites. Similarly, a Factor VII polypeptide having only the two N-linked glycosylation sites found in wild type Factor VII may be deglycosylated at one of these sites and then subjected to desialylation. The resulting Factor VII variant

having one N-linked glycan lacking sialic acid has different pharmacokinetics than the similar Factor VII polypeptide that did not lack a second N-linked glycosylation site based on the experimental evidence reported herein.

Definitions and Embodiments

5

[0033] Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Generally, the nomenclature used herein and the laboratory procedures in cell culture, molecular genetics, organic chemistry, and nucleic acid chemistry and hybridization are those well-known and commonly employed in the art. Standard techniques are used for 10 nucleic acid and polypeptide synthesis. The nomenclature used herein and the laboratory procedures in analytical chemistry and organic synthesis described below are those well-known and commonly employed in the art. Standard techniques, or modifications thereof, are used for chemical syntheses and chemical analyses. Procedures used for genetic engineering are well known and can be found, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, N.Y.

15

[0034] The term "sialic acid" or "sialyl" refers to any member of a family of nine-carbon carboxylated sugars. The most common member of the sialic acid family is N-acetyl-neuraminic acid (2-keto-5-acetamido-3,5-dideoxy-D-glycero-D-galactononulopyranos-1-onic acid (often abbreviated as Neu5Ac, NeuAc, or NANA)).

20

[0035] The terms "polypeptide" and "protein" are used interchangeably herein and refer to a polymer in which the monomers are amino acids and are joined together through amide bonds. Additionally, unnatural amino acids, for example, β -alanine, phenylglycine, and homoarginine, are also included. Amino acids that are not gene-encoded can also be used with the technology disclosed herein. Furthermore, amino acids that have been modified to include reactive groups, glycosylation sites, polymers, therapeutic moieties, biomolecules, and the like can also be used. All of the amino acids used herein can be either the D- or L- isomer. The L-isomer is generally preferred. As used herein, "polypeptide" and "protein" refer to both glycosylated and unglycosylated polypeptides and proteins, respectively.

25

[0036] The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. "Amino acid analogs" refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, i.e., an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g. homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (e.g. norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. "Amino acid mimetics" refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that function in a manner similar to a naturally occurring amino acid.

30

[0037] The term "half-life" or "t_{1/2}," as used herein in the context of administering a polypeptide or protein drug to a patient, is defined as the time required for plasma concentration of a drug in a patient to be reduced by one half.

35

[0038] Half-life can be determined in test animals, for example, by administering a dose of about 25-250 microgram/kg of the preparation; obtaining plasma samples at predetermined times after administration; and determining the content of the Factor VII polypeptide in the samples using one or more of a clotting assay (or any bioassay), an immunoassay, or an equivalent. The data can be displayed graphically and then the bioavailability will be determined as the area under the curve. In certain examples, rat or murine models are used for half-life measurements. Relative bioavailability of a Factor VII polypeptide or composition thereof refers to the ratio of the area under the curve of the short-acting Factor VII polypeptide to that of wild-type Factor VII or another appropriate comparator polypeptide or protein. Any Factor VII variant that has blood coagulation activity of Factor VII is useful for the purposes and methods described herein. Factor VII variants as used herein are polypeptides. The terms "variant Factor VII polypeptides" and "Factor VII variants" are used interchangeably herein. In one embodiment, the Factor VII variants have an amino acid sequence derived from wild type Factor VII (SEQ ID NO: 16) by substitution, deletion, and/or insertion of one or more amino acids. In designating amino acid substitutions, the first letter represents the amino acid present in the wild type human Factor VII at a position. The following number represents the position in human wild type Factor VII. The second letter represents the amino acid replacing the amino acid found in the wild type. For example, "P10Q" represents a substitution of a glutamine (Q) for a proline (P) at amino acid position 10.

40

[0039] In certain examples, the Factor VII variant comprises one or more amino acid substitutions selected from the group consisting of P10Q, K32E, R36E, A34E, T106N, and V253N. In other examples, the Factor VII variant comprises at least 2, 3, 4, 5, or 6 of these substitutions. In further examples, the Factor VII variant comprises an amino acid sequence having at least two sequence alterations relative to the amino acid sequence of SEQ ID NO: 16 (wild type human Factor VII), wherein the at least two sequence alterations are (1) a glutamine residue substituted for the proline residue in position 10, and (2) a glutamic acid residue substituted for the lysine residue in position 32. In another example, the Factor VII variant comprises an amino acid sequence having at least three sequence alterations relative to the amino

acid sequence of SEQ ID NO: 16, wherein the at least three sequence alterations are (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, and (3) a glutamic acid residue substituted for the arginine residue in position 36. In a further example, the Factor VII variant comprises an amino acid sequence having at least four sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least four sequence alterations are (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the arginine residue in position 36, and (4) a glutamic acid residue substituted for the alanine residue in position 34. In one particular example, the Factor VII variant comprises an amino acid sequence having at least six sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least six or six sequence alterations are (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the arginine residue in position 36, (4) a glutamic acid residue substituted for the alanine residue in position 34, (5) an asparagine residue substituted for the threonine residue in position 106 and (6) an asparagine residue substituted for the valine residue in position 253. In another particular example, the Factor VII variant comprises only these six alterations. More details on these variants are found in WO 200158935 to Maxygen, and U.S. Pat. No. 7,371,543 to Pedersen et al., both of which are incorporated by reference herein in their entireties.

[0040] The Factor VII variants described herein can be designed using any functional Factor VII polypeptide as a starting polypeptide. In certain embodiments, the Factor VII polypeptide is a human Factor VII polypeptide. In further embodiments, the Factor VII polypeptide is the human Factor VII polypeptide of SEQ ID NO: 16, or a modified form or allelic variant thereof. Useful starting polypeptides also include modified or variant Factor VII polypeptides comprising an amino acid sequence at least 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, or 66% identical to the sequence of wild type human Factor VII (SEQ ID NO: 16) that also possess Factor VII activity. Further, in certain examples, the variant Factor VII polypeptides of the present disclosure include any polypeptide with at least about 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, or 66% identity to the sequence of SEQ ID NO: 16 that possess Factor VII functionality and that also contain one or more of the amino acid alterations discussed herein relative to SEQ ID NO: 16. In another embodiment, the Factor VII polypeptide comprises an amino acid sequence having more than 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, or 66% homology to SEQ ID NO: 16 and has Factor VII activity, and that also has one or more of the amino acid alterations referenced herein.

[0041] Factor VII variants as used herein also includes glycosylation variants of wild type Factor VII. For example, a partially desialylated wild type Factor VII variant and compositions thereof can be useful because it has a shorter half-life than wild type Factor VII. Also useful herein are pharmaceutical formulations of partially or completely desialylated wild type Factor VII and use of such polypeptides and formulations in the treatment of the diseases recited herein that benefit from a short-acting polypeptide having Factor VII activity. Partial or complete desialylation can be measured by the ratio of moles of conjugated sialic acid to moles of N-linked glycan in a composition of Factor VII polypeptides as described herein.

[0042] Nucleotide sequences encoding the Factor VII variants herein are also useful. In one embodiment, the Factor VII polypeptides are encoded by a nucleotide sequence having at least 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, or 66% identity across the full length to the nucleotide sequence of wild type Factor VII (SEQ ID NO: 1) and that encode a functional Factor VII polypeptide. In certain examples, the nucleotide sequence also encodes a polypeptide containing one or more of the amino acid alterations discussed herein relative to SEQ ID NO: 16. In another embodiment, the Factor VII polypeptide is encoded by a nucleotide sequence having more than 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81%, 80%, 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, or 66% homology to the nucleotide sequence of wild type Factor VII (SEQ ID NO: 1) and that encodes a functional Factor VII polypeptide. In certain examples, the nucleotide sequence also encodes a polypeptide containing one or more of the amino acid alterations discussed herein relative to SEQ ID NO: 16.

[0043] The percent identity values are calculated over the entire amino acid or nucleic acid sequence region. A series of programs based on a variety of algorithms are available to the skilled worker for comparing different sequences. In at least one embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch algorithm (Needleman 1970, J. Mol. Biol. (48):444-453), which has been incorporated into the needle program in the EMBOSS software package (EMBOSS: The European Molecular Biology Open Software Suite, Rice, P., Longden, I., and Bleasby, A, Trends in Genetics 16(6), 276-277, 2000), using either a BLOSUM 45 or PAM250 scoring matrix for distantly related proteins, or either a BLOSUM 62 or PAM160 scoring matrix for closer related proteins, and a gap opening penalty of 16, 14, 12, 10, 8, 6, or 4 and a gap extension penalty of 0.5, 1, 2, 3, 4, 5, or 6. Guides for

local installation of the EMBOSS package as well as links to WEB-Services can be found at emboss.sourceforge.net. A non-limiting example of parameters to be used for aligning two amino acid sequences using the needle program are the default parameters, including the EBLOSUM62 scoring matrix, a gap opening penalty of 10, and a gap extension penalty of 0.5. In yet another embodiment, the percent identity between two nucleotide sequences is determined using the needle program in the EMBOSS software package (EMBOSS: The European Molecular Biology Open Software Suite, Rice, P., Longden, I., and Bleasby, A, Trends in Genetics 16(6), 276-277, 2000) using the EDNAFULL scoring matrix with a gap opening penalty of 16, 14, 12, 10, 8, 6, or 4 and a gap extension penalty of 0.5, 1, 2, 3, 4, 5, or 6. A non-limiting example of parameters to be used for aligning two amino acid sequences using the needle program are the default parameters, including the EDNAFULL scoring matrix, a gap opening penalty of 10, and a gap extension penalty of 0.5. The nucleic acid and protein sequences can further be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the BLAST series of programs (version 2.2) of Altschul et al. (Altschul 1990, J. Mol. Biol. 215:403-10). BLAST using nucleic acid sequences of the present disclosure as query sequence can be performed with the BLASTn, BLASTx, or tBLASTx program using default parameters to obtain either nucleotide sequences (BLASTn, tBLASTx) or amino acid sequences (BLASTx) homologous to sequences encoded by the nucleic acid sequences of the present disclosure. BLAST using protein sequences encoded by the nucleic acid sequences of the present disclosure as query sequence can be performed with the BLASTp or the tBLASTn program using default parameters to obtain either amino acid sequences (BLASTp) or nucleic acid sequences (tBLASTn) homologous to sequences of the present disclosure. To obtain gapped alignments for comparison purposes, Gapped BLAST using default parameters can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25(17):3389-3402.

[0044] The polynucleotides of the present disclosure either essentially consist of the aforementioned nucleotide sequences or comprise the aforementioned nucleotide sequences. Thus, they can contain further nucleotide sequences as well. In certain embodiments, the polynucleotide can comprise, in addition to an open reading frame, further untranslated sequence at the 3' and/or at the 5' terminus of the coding gene region, for example at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, or more nucleotides of the sequence upstream of the 5' terminus of the coding region and/or at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, or more nucleotides of the sequence downstream of the 3' terminus of the coding gene region. Furthermore, the polynucleotides can encode fusion proteins wherein one partner of the fusion protein is a polypeptide being encoded by a nucleotide sequence recited above. Such fusion proteins can comprise so called "tags" which may serve as a detectable marker or as an auxiliary measure for purification purposes. Tags for the different purposes are well known in the art and comprise FLAG-tags, 6-histidine-tags, MYC-tags and the like. In one embodiment, the polynucleotide further comprises an expression control sequence operatively linked to the nucleotide sequence.

[0045] In certain embodiments, a nucleic acid sequence encoding the Factor VII polypeptide is inserted into a suitable vector. Numerous vectors useful for various purposes are well known in the art and persons skilled in the art would be able to readily select an appropriate vector for their desired application. In certain examples, the vector may be a cloning vector or an expression vector. In other examples, the vector may be a plasmid, a viral vector, a cosmid, or an artificial chromosome. In certain examples, the nucleic acid encoding the Factor VII polypeptide may be placed adjacent to and/or under the control of an appropriate promoter. Numerous promoters useful for various purposes are well known in the art and persons skilled in the art would be able to readily select an appropriate promoter for their desired application. In certain examples, the promoter may be a constitutive promoter, an inducible promoter, or a tissue specific promoter.

[0046] In certain embodiments, the Factor VII polypeptides are recombinantly produced in a cell, tissue, or organism. In certain embodiments, such recombinant production is accomplished by transforming or transfecting a host cell with a nucleic acid molecule encoding the variant polypeptide or a vector containing such a nucleic acid. Numerous methods of transformation and transfection are well known in the art and persons skilled in the art would be able to readily select an appropriate method for their desired application.

[0047] Such recombinant production can also be accomplished using any suitable host cell, tissue, or organism. Suitable cells, tissues, and organisms are well known in the art and persons skilled in the art would be able to readily select an appropriate host for their desired application. In some embodiments, the host cell is mammalian. Examples of suitable mammalian cell lines are the COS-1 (ATCC CRL 1650), baby hamster kidney (BHK), HEK293 (ATCC CRL 1573; Graham et al., J. Gen. Virol. 36:59-72, 1977), HEK293T (ATCC CRL 11268; DSM ACC 2494), and HEK293F (Invitrogen R79007) cell lines. A useful BHK cell line is the tk³¹ ts13 BHK cell line (Waechter and Baserga, Proc. Natl. Acad. Sci. USA 79:1106-1110, 1982, incorporated herein by reference), hereinafter referred to as BHK 570 cells. The BHK 570 cell line has been deposited with the American Type Culture Collection, 12301 Parklawn Dr., Rockville, Md. 20852, under ATCC accession number CRL 10314. A tk⁻ ts13 BHK cell line is also available from the ATCC under accession number CRL 1632. In addition, a number of other cell lines can be used within the present disclosure, including Rat Hep I (Rat hepatoma; ATCC CRL 1600), Rat Hep II (Rat hepatoma; ATCC CRL 1548), TCMK (ATCC CCL 139), Human lung (ATCC HB 8065), NCTC 1469 (ATCC CCL 9.1), CHO (ATCC CCL 61), CHO K1 (ATCC CCI61), DUKX cells (Urlaub and Chasin, Proc. Natl. Acad. Sci. USA 77:4216-4220, 1980) and CHO-DG44 cells (Urlaub et al. Cell 33:

405-412, 1983).

[0048] Compositions of Factor VII polypeptides are useful in which the Factor VII polypeptides are defined as herein and the ratio of moles of conjugated sialic acid per mole of N-linked glycan in the composition is less than 0.05, less than 0.1, less than 1.0, less than 2.0, less than 3.0, less than 4.0, less than 5.0 or less than 6.0, or compositions wherein the ratio of moles of conjugated sialic acid per mole of N-linked glycan is within a range selected from the group consisting of (1) from 0 to 8; (2) from 0 to 7; (3) from 0 to 6; (4) from 0 to 5; (5) from 0 to 4; (6) from 0 to 3; (7) from 0 to 2; (8) from 0 to 1 and (9) from 0 to 0.5, or ratios of from 1 to 8, 1 to 7, 1 to 6, 1 to 5, 1 to 4, 1 to 3, 1 to 2, 2 to 8, 2 to 7, 2 to 6, 2 to 5, 2 to 4, 2 to 3, 3 to 8, 3 to 7, 3 to 6, 3 to 5, 3 to 4, 4 to 8, 4 to 7, 4 to 6, 4 to 5, and 0.1 to 1. The ratio is a measurement of the moles of sialic acid bound to a glycoprotein relative to the number of glycans on the glycoprotein. The number of glycans refers to the number of sugar moieties attached to an N-linked glycan in the glycoprotein, where one N-linked glycosylation site can support only one glycan as defined herein for purposes of this ratio. The ratio is determined using a sialic acid fluorescence labeling kit such as that sold by Takara Bio Inc. (cat. #4400). Such a sialic acid fluorescence labeling kit includes a step for the release of sialic acid from the bound glycoprotein, such as by partial acid hydrolysis or by use of sialidase, such as *Arthrobacter ureafaciens* sialidase. The free sialic acids are then labeled with a fluorophore such as 1,2-diamino-4,5-methylenoxybenzene ("DMB"). The labeled sialic acids are then quantitatively measured using HPLC and comparing peak heights to a calibration curve. Thus, the ratio measured is a ratio of moles of sialic acid per mole of glycan released from all the Factor VII polypeptides of the composition.

[0049] In one series of embodiments, the compositions of Factor VII polypeptides or the isolated polypeptides themselves have a half-life as measured in human or mammalian plasma, for example murine or rat plasma, of less than 2 hours, less than 1.5 hours, less than 1 hour, less than .75 hour, less than .5 hour, less than .25 hour, less than 0.1 hour, or so short that it cannot reasonably be measured.

[0050] As used herein, Factor VII activity is a biological activity that may be quantified by measuring the ability of a preparation to promote blood clotting using Factor VII-deficient plasma and thromboplastin, as is well known in the art. In certain examples, a Factor VII polypeptide having Factor VII activity shows at least 25%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% of the activity of wild type Factor VII as measured under the same conditions.

[0051] Pharmaceutical formulations of the Factor VII polypeptides and compositions thereof comprising the Factor VII polypeptide and a pharmaceutically acceptable excipient or carrier are also useful. In certain examples, the pharmaceutical formulations are for parenteral administration, such as by intravenous, subcutaneous or intramuscular administration, and dosing may be as a single bolus dose, intermittent dosing, or as a continuous intravenous infusion. Topical formulations are also useful. One embodiment comprises a pharmaceutical formulation comprising an isolated Factor VII polypeptide as described herein, or comprising a composition of Factor VII polypeptides as described herein, in a lyophilized preparation that is reconstituted at the time of use. Alternatively, the pharmaceutical formulation can be a stable liquid ready-to-use formulation not requiring reconstitution. The pharmaceutical formulation can be a lyophilized powder in single-use vials of 1, 2, 5, or 8 mg of Factor VII polypeptide. After reconstitution with a specified volume of liquid, such as sterile water containing histidine, the final solution can contain any suitable amount of Factor VII polypeptide that produces a therapeutic effect, such as, without limitation, 1 mg/mL (1000 micrograms/mL), 2 mg/mL, 3 mg/mL, 4 mg/mL, 5 mg/mL, 1-2 mg/mL, 1-3 mg/mL, 1-5 mg/mL, 1-10 mg/mL, 0.5-1 mg/mL, or 0.5-2 mg/mL of Factor VII polypeptide. Proper dosage for administration to a patient can be readily determined by persons skilled in the art based upon, for example, the weight of the patient, the type of bleeding disorder or episode being treated, and the activity of the particular Factor VII polypeptide being employed. In certain examples, dosing can be in the range of 70-110 micrograms/kg, 70-90 micrograms/kg, or 80-100 micrograms/kg and can be 90 micrograms/kg. The lyophilized powder may be reconstituted with an aqueous carrier, such as water, buffered water, 0.4% saline, 0.3% glycine, etc. Actual methods for preparing parenterally administrable compositions will be known or apparent to those skilled in the art and are described in more detail in, for example, Remington's Pharmaceutical Sciences, 18th ed., Mack Publishing Company, Easton, Pa. (1990). Topical application, such as can be advisable in the case of trauma, can be carried out by means of a spray, perfusion, catheters, stent, vascular graft or stent, ointment, or other preparation known in the art. In certain examples, topical administration can be by way of a solid or semi-solid matrix, such as a surgical sponge or collagen matrix, which has been treated with, infused with, coated with, or soaked in a composition comprising the Factor VII variant. Methods of preparing such matrices are well known in the art (see, e.g., Thrombosis/Hemostasis 12:445, 2006) and the skilled artisan would be able to readily determine an appropriate dose and method of application of the composition onto the given matrix.

[0052] In one embodiment, the present disclosure relates to kits comprising the Factor VII polypeptide. In certain examples, the kit contains a vial containing ready-to-use liquid containing the Factor VII polypeptide in a suitable pharmaceutical composition. In other examples, the kit contains a vial containing lyophilized Factor VII polypeptide, or a lyophilized formulation comprising the polypeptide, and also a diluent for reconstitution. In other examples, the kit contains a topical formulation of the Factor VII polypeptide, for example, an ointment, spray, or liquid, and a matrix such as a sponge or other medical matrix to which the topical formulation may be applied before administration to the patient.

[0053] Compositions of the Factor VII polypeptides described herein are also useful. Factor VII exists in mixture with its natural degradation products. Accordingly, a composition of Factor VII polypeptides includes polypeptides having one of the full amino acid sequences as recited herein and degradation products having partial amino acid sequences of those described herein. Furthermore, because Factor VII is a glycoprotein, compositions of Factor VII can be expected to contain a heterogeneous mixture of Factor VII polypeptides wherein each glycoprotein in the composition does not have exactly the same glycosylation as the others. Reference to compositions of Factor VII polypeptides or isolated Factor VII polypeptides is meant to encompass mixtures of such polypeptides wherein the individual polypeptides have different glycosylation, and thus the terms "composition" or "isolated Factor VII polypeptide" encompass a heterogeneity of the glycosylation patterns within the polypeptides.

[0054] The Factor VII polypeptides and compositions described herein are useful for the treatment of blood clotting disorders, and those disorders that benefit from blood coagulation, and particularly for coagulation with a drug having a shorter half-life than wild type Factor VII. Accordingly, the Factor VII polypeptides and compositions herein are useful for penetrating traumatic injury; blunt traumatic injury; bleeding in elective surgery; bleeding in cardiac surgery; bleeding in spinal surgery; orthopedic surgery; neurosurgery; oncology surgery; post-partum surgery; menorrhagia; bleeding in stem cell transplantation; bleeding in liver transplantation; gastrointestinal bleeding; active variceal bleeding in cirrhosis; non variceal bleeding in cirrhosis; diffuse alveolar hemorrhage; aortic aneurysm; intracerebral hemorrhage; traumatic brain injury; brain contusion; reversal of warfarin; reversal of heparin; reversal of anticoagulants; reversal of anti-thrombotics; Factor VII deficiency; burns; prophylaxis in hemophilia patients with inhibitors; partial hepatectomy for non-cirrhotic and cirrhotic patients; acquired hemophilia; idiopathic thrombocytopenic purpura; Glanzmann's Thrombasthenia; Glanzmann's Thrombasthenia refractory to platelet transfusion and Bernard-Soulier Syndrome.

[0055] Also disclosed herein is a useful assay for measuring the half-life of coagulation factors such as Factor VII. There is a method of determining the half-life of a coagulation factor comprising incubating viable rat hepatocyte cells with a blood coagulation factor, removing a sample at testing time point 1, separating supernatant from cells in the sample and quantifying the activity or amount of the blood coagulation factor in the supernatant in the sample, wherein the activity or amount of the blood coagulation factor is determined using a double-antibody sandwich ELISA assay. The method may be repeated at different time points to develop a plot of activity or amount of blood clotting factor over time.

EXAMPLES

30 Methods to Obtain Desialylated Factor VII Polypeptides

[0056] Numerous methods were employed to generate desialylated Factor VII polypeptides (both wild-type and variant), including enzymatic desialylation of the polypeptide, production of the Factor VII polypeptide in a sialylation-deficient cell line, and co-expression of Factor VII and a sialidase in a recombinant cell.

35 *Generation of Sialic Acid Deficient Cell Line*

[0057] Endogenous sialic acid is synthesized in mammalian cells involving a complex pathway consisting of 32 enzymes (Wickramasinghe and Medrano 2011). The biosynthesis of sialic acid starts in cytosol converting UDP-N-acetylglucosamine (UDP-GlcNAc) to Neu5Ac involving several enzymes, such as UDP-N-acetylglucosamine-2-epimerase/N-acetylmannosamine kinase (GNE), sialic acid 9-phosphate synthase (NANS), and sialic acid 9-phosphate phosphatase (NAMP). Neu5Ac in cytosol is imported into the nucleus through nuclear pores and converted into CMP-Neu5Ac by an enzyme called CMP-Sia synthase (CMAS). Synthesized CMP-Neu5Ac is again transported back into the cytosol via nuclear pores for further modification and conjugation in the Golgi apparatus. Conversion of Neu5Ac into Neu5Gc in cytosol is catalyzed by the enzyme CMP-NeuAc-hydroxylase (CMAH). Then, CMP-Neu5Ac and CMP-Neu5Gc are transported into the Golgi compartment via a hydrophobic type 3 membrane transporter, CMP-sialic acid transporter (SLC35A1), located in the membrane of the median trans-Golgi. CMP-sialic acid transporter is a key element in the cellular sialylation pathway (Hirschberg, et al. 1998). A homozygous mutation of this gene causes post-natal lethality in the mouse (MGI 4.32, Homologene). In humans mutations in SLC35A1 are associated with the reduction or complete loss of sialyl conjugates. Some insertion and deletion mutations in SLC35A1 are associated with congenital disorders of glycosylation in humans leading to defects in nervous system development, coagulation, and immune deficiency (Martinez-Duncker, et al., 2005). Once CMP-Neu5Ac/CMP-Neu5Gc is transported into the Golgi apparatus they are conjugated with carbohydrates, glycoproteins, and glycolipids by enzymes in the sialyltransferase (ST) family with 20 members.

[0058] CMP-sialic acid transporter (SLC35A1) is the key molecule supporting sialic acid conjugation in the Golgi apparatus, and mutations to this transporter protein lead to synthesis of proteins lacking proper sialylation. To produce desialylated Factor VII, a Factor VII production cell line with CMP-sialic acid transporter gene knockout is produced. Alternatively, desialylation could be achieved by expression of the Factor VII variant in a cell line that produces protein

therapeutics with a very low level or no sialylation on the therapeutic molecules. This technology could be used to produce therapeutic proteins with short T1/2 in patients.

[0059] Lec2 cell with Chinese hamster ovary (CHO) origin was identified with the property of producing approximately 10-fold less sialic acid in glycoproteins and glycolipids than the respective wild-type cells (Stanley and Siminovitch, 1977, 5 Stanley, 1980 and 1983). A later study showed that Lec2 mutants were unable to translocate CMP-sialic acid across membranes of Golgi vesicles in an *in vitro* assay, while translocation of other nucleotide derivatives was comparatively normal in mutant cells (Deutscher, et al., 1984). By using expression cloning, the gene encoding the CMP-sialic acid transporter from Lec2 cells was reported (Eckhardt et al., 1996). Further investigation indicated that the deletion of 10 nucleotides 575 - 751 in the CMP-sialic acid transporter gene was responsible for the Lec2 phenotype (Eckhardt, et al., 1998). Other mutations in the CMP-sialic acid transporter gene, such as in the case of 1E3, 6B2, 8G8, and 9D3 cells, also lead to the Lec2 phenotype (Eckhardt, et al., 1998).

Experiment 1

[0060] This experiment is designed to determine whether the mutation in the gene of CMP-sialic acid transporter, such as in the case of Lec2 cells, results in expressed recombinant protein (e.g., Factor VII) deficient in sialylation in comparison with the same protein expressed from normal CHO cells.

[0061] (1) Testing the expression of recombinant protein, such as Factor VII, from Lec2 cells. Expression vectors containing a Factor VII variant gene (e.g., pMB117 and pMB121) are transfected into Lec2 cells and normal CHO cells under normal transfection conditions. Expression levels of Factor VII from cell culture of these cells is monitored by a Factor VII activity assay. The culture of transfected cells is scaled-up and the culture conditioned media is harvested for the purification of Factor VII.

[0062] (2) Testing the sialic acid content of purified Factor VII expressed from Lec2 cells in comparison to the same protein expressed from normal CHO cells. Purification of Factor VII from these conditioned media is conducted following the normal Factor VII purification methods. The purified Factor VII from either Lec2 cells or normal CHO cells is analyzed for sialic acid content of the purified Factor VII. The biological activity and pharmacokinetic (PK) parameters are analyzed as described herein.

Experiment 2

[0063] In order to produce a manufacturing cell line to express a Factor VII variant with no sialic acid on the expressed recombinant proteins, gene deletion methods are used targeting the CMP-sialic acid transporter gene to modify a cell line that expresses Factor VII (e.g., a CHO cell line). In order to completely inhibit the sialylation in the cell, other targets, such as UDP-N-acetylglucosamine-2-epimerase/N-acetylmannosamine kinase (GNE), sialic acid 9-phosphate synthase (NANS), sialic acid 9-phosphate phosphatase (NANP), and CMP-Sia synthase (CMAS) as listed above in the introduction, 35 optionally can also be deleted to enhance the inhibition of the CMP-Neu5Ac biosynthesis which provides the substrate for the CMP-sialic acid transporter.

[0064] Two gene deletion technologies, TALE nucleases (TALENs) from Life Technologies and ZFP Nucleases (ZFNs) from Sangamo BioSciences/Sigma-Aldrich, can be used for conducting the knockout of CMP-sialic acid transporter 40 gene, or the knockout of multiple genes in the sialic acid synthesis pathway.

[0065] The Factor VII expression cell line with CMP-sialic acid transporter gene knockout is evaluated to confirm the deletion of CMP-sialic acid transporter gene. The confirmed cell line is cultured to produce Factor VII. Factor VII from the CMP-sialic acid transporter gene-deleted cell line is purified and evaluated as described herein and compared to the Factor VII from the parental Factor VII expression cell line for sialic acid content on the molecule.

45 Production of Desialylated Factor VII by Coexpression of Factor VII and a Bacterial Sialidase

[0066] To generate a desialylated form of FVII, we co-expressed FVII along with a bacterial sialidase variant derived from the *Arthrobacter ureafaciens* sialidase (AU sialidase) (N-acetylneuraminate glycohydrolase, EC 3.2.1.18). CHO 50 cells stably express FVII (e.g., SEQ ID NO 16 with P10Q, K32E, A34E and R36E mutations) were further transfected with AU sialidase using a pJ608 expression vector. The expressed protein has a growth hormone signal sequence at the N-terminus to promote secretion of the protein. Stable clones were selected that produced AU sialidase as detected by a chromogenic assay for sialidase in the medium. We also showed that the cells continued to express high levels of 55 FVII protein, which was detected by ELISA assay, SDS PAGE and Western blotting, using FVII specific antibodies as the probe. Using a lectin blotting assay, we were not able to detect sialic acid on FVII protein in the condition media, using purified FVII. In contrast, similar levels of purified FVII derived from cells not transfected with sialidase showed a strong lectin binding signal. Taken together, our data show that the AU sialidase was expressed in the CHO cell medium at levels sufficient to efficiently desialylate FVII that was co-expressed by the cells.

Enzymatic Preparation of Desialylated Factor VII Using a Soluble Sialidase Treatment

[0067] The following starting materials were utilized in this experimental trial:

5 Factor VII: 20mg Wild Type Factor VIIa, concentration about 1 mg/ml
 Sialidase: 20ug, 0.25mg/ml, 50000U/ml, P0720L, purchased from New England BioLabs
 Buffer solution A: 25mM histidine, 50mM NaCl, pH 6.4
 Buffer solution B: 25mM histidine, 1M NaCl, pH 6.4
 FVIIa formulation buffer: 2.3mg/ml sodium chloride, 1.5mg/ml calcium chloride dehydrate, 1.3mg/ml glycylglycine, 10
 10 0.1mg/ml polysorbate 80, 25mg/ml mannitol, 10mg/ml sucrose, 0.5mg/ml methionine, 1.6mg/ml histidine, pH 6.0
 Purification column: 5ml HiTrap Q Sepharose HP column

Using these materials, the following procedure was carried out:

15 1. To 20 mgs of FVIIa (about 1 mg/ml), add 20ug of sialidase (0.25mg/ml, 1:1000 mass ratio) of sialidase
 2. Incubate the reaction at room temperature overnight (about 19hr) before chromatographically purifying de-
 sialylated FVIIa as described below.
 3. Purify desialylated FVIIa on a 5ml HiTrapQ Sepharose HP column as follows:

20 a) Equilibrate the Q-Sepharose column with 5CV of buffer A (25mM histidine, 50mM NaCl, pH 6.4).
 b) Before applying to the column, dilute the FVIIa and sialidase reaction with 200ml of buffer A and adjust
 pH to 6.4.
 c) Load at a flow rate of 2.5ml/min using an AKTA Explorer system while monitoring A280. Collect Flow
 through fraction.
 d) After loading is completed, wash the column with 10CV of Buffer A.
 25 e) Elute the column with 20CV of 0-50% buffer B (25mM histidine, 1M NaCl, pH 6.4) in 40min. Collect peak
 fractions (Desialylated NovoSeven)
 f) Dialyze the peak fractions vs. FVIIa formulation buffer at 4°C overnight
 g) Freeze the sample at -80°C in aliquots.

30 [0068] The product was shown to be highly pure by SDS-PAGE, aSEC, and active in biological assays for FVIIa. Assays for sialic acid content showed no residual sialic acid and LC-MS analysis of the heavy chain showed no significant alteration of glycan structure other than removal of sialic acid.

Enzymatic Preparation of Desialylated Factor VII Using Neuramidase-Agarose Beads

35 [0069] Recombinant wild type Factor VII as used herein is NovoSeven® obtained from Novo Nordisk and referred to herein as "F7." Other starting materials are V1 and V2 as described supra.
 [0070] Frozen starting material was quick-thawed in a 37°C water bath and pooled. The protein was concentrated 2.5-fold by centrifugation; concentrate was gently mixed by pipetting to minimize any super-concentration (aggregation) at the protein-filter interface.
 40 [0071] The V2 was buffer-exchanged from its V2 formulation buffer (containing histidine, CaCl₂, trehalose, methionine, and trace levels of Tween®-20, at a pH 6.4 - 6.6) into MES buffer (containing 10mM MES, 10mM CaCl₂, 50mM NaCl, pH 6.0, sterile-filtered). This was achieved one of three ways. In the first option, V2 was buffer-exchanged with NAP-10 gravity flow columns (GE, 17-0854-01), which were pre-washed 3-5 times with 3 column-volumes each of MES buffer.
 45 The V2 was then loaded onto the column and eluted with 1.5 times the load volume of MES buffer. In the second option, V2 was buffer-exchanged by overnight dialysis in MES buffer. Dialysis cassettes were pre-soaked in MES buffer and the V2 was loaded by syringe into the 3.500 MWCO slide-a-lyzer cassettes (Thermo Scientific, 66130), overnight at 4°C in 10L pitchers with sterile-filtered MES buffer. In the third option, V2 was buffer-exchanged into MES buffer through a Sephadex G-25 (Sigma, G-25-80) gel-filtration column, and equilibrated with MES buffer.
 50 [0072] The buffer-exchanged V2 was desialylated with neuraminidase-agarose (Sigma N5254). The agarose bead product is provided in a 50% slurry mixture, stored in ammonium sulfate buffer; beads were pre-washed 3-5 times in MES buffer; the bead/buffer mixture was separated by centrifugation at 1000rcf for 3min at 4°C, and the supernatant liquid was pipetted off and discarded. To the washed beads, the buffer-exchanged V2 was added and gently mixed by rotation at room temperature for 16 to 22 hours. 2.08 mL of packed beads per mg of protein were used for desialylation; for a larger scale preparation, this was reduced 1:10, to 0.208 mL of beads per mg of protein. Afterward, the desialylated V2 was recovered by centrifugation and pipetted off. The beads were washed once gently for 5 minutes by rotation in 1:1 by volume of fresh MES buffer; the wash mixture was centrifuged as before and the supernatant was pooled with the V2. The beads were finally removed either by sterile-filtration through a 0.2 micron syringe filter or by vacuum filtration

through at 0.45 micron filter.

[0073] Several rounds of endotoxin removal were performed with EndoTrap HD resin (Hyglos). The resin was washed 3-5 times in MES buffer and wash buffer was discarded. In two batches, 1-3mL of washed resin was gently mixed with the desialylated V2 overnight at room temperature. The resin was removed by centrifugation and then filtered through 5 syringe or vacuum filter.

[0074] The desialylated V2 was concentrated 4.75-fold to 2.1mg/mL by centrifugation in Ultracels for 10-minute cycles; the concentrates were gently mixed by pipetting to reduce any aggregation at the protein-filter interface.

[0075] The desialylated V2 was further separated from higher-molecular weight species (and aggregated endotoxin) with HiLoad 26/60 Superdex 200 size exclusion column. The column and the AKTA purifier system was pre-sanitized 10 with 0.1N NaOH + 20% EtOH. The system was pH-neutralized, rinsed with water, and equilibrated with reconstituted and pooled V2 formulation buffer. Several batches of concentrate were manually injected into a 12mL sample loop and loaded onto size exclusion column at a flow rate of 3mL/min; the eluate was recovered and fractionated with a Frac-900 15 into polystyrene tubes (17X100mm, Fisherbrand, 14-956-6D). Early, high-molecular weight peaks were excluded and the desired V2 fractions were pooled and tested for endotoxin levels and concentration using Charles River EndoSafe PTS and NanoDrop ND-1000. V2 buffer was used to elute the desialylated V2.

[0076] Five batches of size-exclusion were performed and the collected eluates were pooled into one batch which 20 was concentrated to 1.0mg/mL in Ultracels. The final preparation was sterile-filtered through 0.2 micron syringe filters and tested for endotoxin and concentration. 1mL aliquots were pipetted into labeled 2mL tubes (Sarstedt, 72.694.006), flash-frozen in an ethanol/dry-ice batch, and stored in a labeled box at -80°C until use.

Characterization - Protein Analytics and *In Vitro* Assay

[0077] The final prep material as well as the non-treated starting material were characterized by protein gel analysis 25 with 4-12% Bis-Tris NuPAGE (Novex NP0335BOX) in MES running buffer and by analytical size exclusion (TSK3000 column; running buffer: 200mM KH2PO4, 150mM KCl, pH 6.8, flow rate: 0.15ml/min, fluorescence detection). Small test samples were analyzed by LC-MS for sialic acid content on the Factor VII heavy chain as well as DMB-labeled sialic acid quantification of total protein using the Takara Bio Inc. kit discussed herein. The activity was tested by phospholipid-dependent Factor X activation and thrombin generation assays.

Sialic Acid Content Analysis

[0078] An LC-MS method was used to identify the sialic acid on the N-glycan of the heavy chain of Factor VII for the non-treated control and desialylated Factor VII. 10 μ g of protein was reduced with 10mM DTT mix at 37°C for 30 min 35 then analyzed on the Agilent 1200 Capillary LC System: Column: PLRP-S 8 μ m 4000A, 0.3x150mm, 75°C. Buffer systems: A: Water with 0.2% Formic Acid+0.01%TFA; B: ACN with 0.2% Formic Acid+0.01%TFA. Gradient: 50 μ L/min, 10% B in 2 min, to 90%B in 25 min, 90%B wash 5 min, 10%B equilibration for 5 min.

[0079] Agilent 6520 Q-TOF system: DualEsi source, gas temp: 350°C, drying gas: 7psi, nebulizer: 10psi, scan range: 500-3000 amu, 1 spectra/s. Reference ions: 1221.990637 and 2421.91399 amu, 50ppm window, Min 1000 counts.

[0080] The results are reported in FIG. 7.

Sialic Acid Quantification using DMB Labeling Kit

[0081] Sialic Acid Fluorescence Labeling Kit (Takara Bio Inc., Cat#4400) is for quantitative and highly sensitive analysis 45 of sialoglycoconjugates. This HPLC-based sialic acid fluorescence labeling technique using 1,2-diamino-4,5-methyle-neoxybenzene (DMB) is a simple and highly sensitive quantitative method. In this method, free sialic acids are analyzed by reverse phase HPLC (GlycosepR, from Glyko, # 1-4727) after labeling by DMB.

Conclusion

[0082] The V2 heavy chain has two N-glycosylation sites. The N-glycans are fucosylated, heavily sialylated bi,tri-and 50 tetra-structures. No terminal sialic acids are found on the de-sialylated sample, which suggests the sample is fully desialylated and that >99.9% of the sialic acid on the Factor VII N-glycan has been removed.

Half-Life Assay Using Rat Hepatocytes

Preparation of Hepatocytes

[0083] Cryopreserved primary rat hepatocytes were obtained from CellzDirect (Invitrogen). Each vial containing ap-

proximately 5 million cells was thawed and the cells were added to 10 ml of Thawing Medium, followed by centrifugation at 60 g for 3 minutes. The cells were resuspended in Incubation Medium + 0.25% BSA (about 4 ml) and the cells were counted using a hemacytometer. Viable cells were counted after staining with Trypan blue to identify dead cells. Cell viability was 80-82%. Cells were used in the clearance assay immediately after counting.

5 [0084] Thawing Medium: Invitrogen CM3000 Thawing/Plating Supplement Pack added to 500 ml Williams E Medium. Incubation Medium: Invitrogen CM4000 Cell Maintenance Supplement Pack added to 500 ml Williams E Medium.

In Vitro Hepatocyte Clearance Assay

10 [0085] Primary rat hepatocytes, 1 million viable cells per ml, were incubated with 25 ng/ml of various Factor VII variants in CellzDirect Incubation Medium + 0.25% BSA, in Eppendorf tubes with gentle end over end mixing at 37°C in a starting volume of 1.2 ml. At each of the indicated time points, 0.25 ml of the mixture was removed and immediately centrifuged to pellet the cells (1000 rpm, 3 minutes in Eppendorf centrifuge). 0.18 ml of the clarified supernatant was removed, quick frozen and stored overnight at -80°C. On the next day, Factor VII in the supernatants was quantified using an ELISA
15 assay in which the corresponding purified mutant protein was used as the standard. No-cell control supernatants in which Factor VII variants were incubated for 2 hours at 37°C in medium alone were used as the zero time point values. Each incubation was done in triplicate. Intrinsic clearance values were calculated based on the method of Lu et al. (Lu
ref.) using the equation $CL_{int} = 0.693 / in vitro T_{1/2}$, normalized for the incubation volume and the number of cells. *In vitro* half-life ($T_{1/2}$) was calculated using the program WinNonLin (Pharsight Corporation, Sunnyvale, CA). Supernatants
20 from hepatocyte incubations were assayed using a double-antibody sandwich ELISA format. 0.1 ml per well of anti-Factor VII monoclonal antibody (1.0 μ g/ml, in PBS) was added to Greiner Microlon 655061 96-well plates. After incubation overnight at 4°C, plates were blocked with 0.2 ml per well of 1% casein blocking buffer (50 mM TrisHCl, 100 mM NaCl, 0.05% Tween 20 pH7.2) for 1.5 hours at 37°C. Plates were washed four times with 0.3 ml per well PBS + 0.05% Tween
25 20 (using a BioTek ELx405 plate washer), and then relevant Factor VII standard and unknown samples were added to the plates. 0.18 ml of each hepatocyte supernatant was diluted two-fold by adding 0.18 ml of Dilution Buffer (50 mM TrisHCl, 100 mM NaCl, 0.1% casein, 0.05% Tween 20 pH7.2). 0.10 ml of each diluted supernatant was added in triplicate to the ELISA plate. Standards were made from the corresponding purified Factor VII variant diluted in Dilution Buffer. Two-fold serial dilutions of the standard were made in Dilution Buffer to yield dilutions in the range of 50 to 0.8 ng/ml
30 final concentration. Factor VII standards and samples (0.1 ml per well) were incubated 2 hours at room temperature (21°C). Plates were washed four times as described above, and then biotinylated detection antibody, 1 μ g/ml in Dilution Buffer (50 mM TrisHCl, 100 mM NaCl, 0.1 % casein, 0.05% Tween 20 pH7.2) was added (0.1 ml per well) followed by incubation for 1.5 hours at room temperature. Plates were washed four times as described above, and then Streptavidin-horseradish peroxidase, diluted 1/1000 in Dilution Buffer was added (0.1 ml per well) followed by incubation for 1 hour at room temperature. Plates were washed again and Ultra-TMB was added, 0.1 ml per well. After incubation for 10 to
35 15 minutes at room temperature, the reaction was stopped with the addition of 0.05 ml per well 2 M H_2SO_4 . Absorbance was read at 450 nm using a Molecular Devices Spectramax M2 plate reader. Data analysis was performed using Softmax Pro 5.4 (Molecular Devices).

40 [0086] Cryopreserved rat hepatocytes, Thawing Medium, and Incubation Medium (CellzDirect), were from Invitrogen/Life Technologies (Grand Island, NY). 1-Step Ultra-TMB (One Step) substrate, catalog no. 34028, was from Thermo Scientific (Rockford, IL) Streptavidin-horseradish peroxidase (SA-HRP), catalog no. DY998, was from R&D Systems, Minneapolis, MN. Phosphate-buffered saline, pH 7.2 was from Invitrogen (Carlsbad, CA). Sprague-Dawley Rat plasma (5% sodium citrate anticoagulant) was from Bioreclamation (Westbury, NY). Greiner Microlon plates (cat. no. 655061) were obtained through Fisher Scientific (Pittsburgh, PA).

45 Methods to Obtain Deglycosylated Variants: Molecular Variants

50 [0087] Wild type Factor VIIa has two N-glycans (N322 and N145), and V1 and V2 each have 4 N-glycans (N106, N145, N253, N322). The additional 2 N-glycans (N106, N253) found in V1 and V2 were originally designed to increase half-life. For this work, these sites are removed by reverting them back to the endogenous amino acid sequence of wild type Factor VII (T106, V253). The remaining 2 endogenous N-glycan sites (N145 and N322) were then removed at the DNA level by engineering in N→Q mutations at these sites. (FIG. 6)

55 [0088] Wild type Factor VII was cloned into pmCMV to make pMB113. Inserts containing a single N to Q mutation at positions aa145 or 322, as well as the double mutant (aa 145 and 322) were synthesized and cloned into pMB113 using the XbaI and PmlI site resulting in clones pMB114-116. Inserts encoding the Gla domains of V1 and V2 were then cloned into pMB113-116 using Ascl and AfeI and resulted in constructs pMB117-120 (V1-based variants) and pMB121-124 (V2-based variants). All constructs were sequence verified (McLab). Mammalian cells (CHO derived cell line) were transiently transfected via electroporation with each construct in a 6 well format. 4 days post transfection, supernatant was collected and assayed by Western Blot for expression, followed by a hFVII ELISA (AssayPro) and FVII activity

assay. A subset of variants was then single cell cloned. The V2 2-Nglycan variant referred to as pMB121 was purified and activated for further analysis.

Method for Purification/Activation of FVIIa from 10L WAVE Expression

5

Summary of Method

[0089] Purification and activation of FVII from diafiltered concentrated conditioned media was accomplished using a multi-step process taking place over several days. The media is first thawed and centrifuged to remove any aggregate that may have formed during freeze/thaw. A psuedoaffinity capture step employing an anion exchange column (Q-Sepharose) eluted with CaCl_2 is used to further concentrate the FVII protein and to change buffer. Next, a hydroxyapatite column is used to further purify the FVII protein. A smaller Q-Sepharose column is then used to further purify FVII before it is activated for 24 hours in solution at pH 7.8 - 8.2. The activation reaction is stopped by lowering the pH to 4.0. The FVIIa is finally dialyzed into Formulation Buffer (pH 6.5) and stored frozen.

10

[0090] The final purified protein is characterized by SDS-PAGE, aSEC, ELISA, glycoanalysis, endotoxin, and FVIIa activity assay.

FVII ELISA

20

[0091] Zymutest FVII enzyme-linked immunoassay (Aniara, West Chester, Ohio). The ELISA is a two-site immunoassay with a rabbit anti-FVII polyclonal antibody bound to the wells of a 96-well microplate. Sample is introduced followed by a rabbit anti-FVII polyclonal antibody coupled to horse radish peroxidase (HRP). The assays were performed following the manufacturer's instructions. Briefly, samples and calibrator were diluted in assay buffer in a 96-well round bottom polypropylene dilution plate. 50 μL aliquots of diluted sample were transferred to the provided rabbit anti-FVII coated plate and incubated for 15 minutes at room temperature. 200 μL HRP coupled rabbit anti-FVII was added and incubated for 1 hour at room temperature. The plate was washed 5 times with 300 μL of the provided wash buffer. TMB was added at 200 μL /well and incubated for approximately 5 minutes at room temperature. The reaction was stopped by introducing 50 μL 0.45 M sulfuric acid. Absorbance was read at 450 nm. Factor VII levels were derived by comparing sample values to a V2 calibration curve generated using a 4-parameter curve fit.

30

Factor VII Chromogenic Assay

35

[0092] Biophen FVII chromogenic assay (Aniara, West Chester, Ohio) was used. The chromogenic assay principle involves formation of an enzyme complex consisting of Factor VII from the sample and rabbit thromboplastin (tissue factor) supplied by the manufacturer. FX, added in excess, is activated to FXa, which in turn cleaves a FXa specific chromogenic substrate (SXa-11) generating pNA. The amount of pNA released is directly proportional to the FXa activity. The assay was performed following the manufacturer's instructions. Briefly, samples and calibrator were diluted in Tris-BSA assay buffer in a 96-well round bottom polypropylene dilution plate (Fisher Scientific). Kit reagents, R1, R2, and R3 and a 96 well flat bottom polystyrene assay plate (Costar) were warmed to 37°C prior to use. 30 μL of sample and calibrator were transferred from dilution plate to assay plate followed by 30 μL of reagent R2 then 60 μL of reagent R1. The assay plate was mixed and incubated for 7 minutes at 37°C in a jitterbug plate shaker (Boekel Scientific). 60 μL of R3 was added and the rate of change in absorbance (change in OD at 405 nm/min) was measured at 37°C using a SpectraMax Plus microplate reader (Molecular Devices). Factor VII levels were derived by comparing sample values to a V2 calibration curve generated using a 4-parameter curve fit.

40

45 Phospholipid-Dependent Thrombin Generation Assay

45

[0093] Compared to wild-type FVIIa, the modification of Gla-domain (P10Q/K32E) increases potency in FX activation, thrombin generation, and whole blood clotting in the presence of anionic phospholipids or activated platelets as a result of additional γ -carboxylation. PL-dependent TGA was designed to measure rFVIIa activity in the presence of anionic phospholipids and performed using thrombin calibrator and substrate reagent, FluCa-kit, from Thrombinoscope, BV. Phospholipid (PL) vesicles composed of 20% phosphatidylserine (PS), 40% phosphatidylethanolamine (PE), and 40% phosphatidylcholine (PC) were from Avanti Polar Lipids and prepared by sonication in 100 mM NaCl, 50 mM Tris-HCl (pH 7.2) for 10 minutes.

55

[0094] Twenty μL of PL-vesicles (500 μM) or thrombin calibrators were dispensed into 96 well plates. Varying concentrations of rFVIIa were diluted into human HemA plasma, added in triplicate to the PL mixture, and equilibrated to 37°C for 10 min. Thrombin generation reactions were initiated by adding FluCa solution and the reactions were continuously monitored for 60 minutes following the Calibrated Automated Thrombography (CAT) method outlined by

Thrombinoscope^{BV}. Data were acquired and analyzed using Thrombinoscope^{BV} (3.4.0) software, which corrected for α_2 -macroglobulin activity using a thrombin calibrator. The analysis parameter 'peak height' represented the maximum level of thrombin generated, 'endogenous thrombin potential' (ETP) corresponded to the total amount of thrombin generated. Thrombin generation parameters were analyzed with 4-parameter non-linear curve fitting method using Prism 4.0 (GraphPad Inc).

5 Phospholipid Dependent FX Activation Assay

10 [0095] The ability of FVIIa to activate FX in the presence of phospholipid vesicles without tissue factor was measured using a PL dependent FX activation assay. Factor VIIa or FVIIa variants are incubated with FX in the presence of phospholipid vesicles. Activation of FX is measured by addition of S-2765, a chromogenic substrate for FXa. Briefly, calibrator and samples are diluted in a polypropylene round bottom plate in Tris-HCl buffer. 30 μ L of 4 μ g/mL FX (Haematologic Technologies Inc.) was added to all wells of a 96-well flat bottom polystyrene plate followed by 30 μ L of phospholipid vesicles consisting of phosphatidylserine, phosphatidylcholine, and phosphatidylethanolamine at a wt. % 15 ratio of 20:40:40. 30 μ L of diluted sample and calibrator were transferred to the FX/phospholipid mixture. The plate was sealed, mixed gently, and incubated for 20 - 23 hours at 37°C. 40 μ L of a 5 mM solution of S-2765 (DiaPharma) was added to all wells. The plate was sealed and incubated for 6 hours at 37°C. Absorbance was read at 405 nm in a microplate reader. Activity of samples was determined by comparing FX activation levels of samples to a F7 calibration curve.

20 Rat PK Study-Animals, Study Protocol (inject of preps, sampling of blood and prep, ELISA, data analysis, sacrifice of animals).

25 [0096] Proteins (F7, V2, V1, dV2 and dV1) were administered intravenously at 0.1mg/kg into Sprague Dawley rats. Plasma samples were taken starting at 1min post administration and analyzed by FVII ELISA.

25 HemA-PK Study

30 [0097] Proteins (F7, dV1) were administered i.v. at 1.0mg/Kg into HemA mice. Plasma samples were taken starting at 5min post administration and analyzed by FVII ELISA and sTF-PT assay.

30 FVII ELISA on Plasma Samples

Materials

35 [0098] Monoclonal antibodies against FVIIa were used. One monoclonal antibody furthermore was biotinylated. Purified FVIIa variants (wild type or de-sialylated) are used as assay calibrators and assay controls. Blocking buffer is 1% (w/v) casein in 30mM Tris pH 7.2, 60mM NaCl, 0.03% Tween-20. Assay dilution buffer (ADB) is 0.1% (w/v) casein, 50mM Tris pH 7.2, 0.1M NaCl, 0.05% Tween-20. Assay wash buffer is PBS + 0.05% Tween-20. Immunoassay plates are Greiner Microlon high binding plates (#655061). Streptavidin-horse radish peroxidase (SA-HRP) is from R&D Systems. HRP substrate Ultra-TMB is from ThermoFisher Pierce. Blank mouse plasma was obtained from CD1 or HemA mice 40 either commercially (Bioreclamation) or through in-house sources. All other materials (casein, Tris, NaCl, Tween-20, PBS, sulfuric acid) are of reagent-grade quality.

Method for FVIIa Sandwich Immunoassay

45 [0099] 96-well assay plates are coated with 0.1ml/well of antibody against FVIIa, 1 μ g/ml in PBS, overnight at 4°C. The plates are aspirated and blocked with 0.2ml/well of blocking buffer for at least 2hr at room temperature with rotation (150rpm). Following blocking, the wells are washed 4 x 0.3ml/well of wash buffer. FVIIa samples or standards are diluted 1:20 to a final concentration of 5% plasma in ADB and incubated 0.1ml/well for at least 1.5hr at room temperature with rotation. All standards, controls, and samples are measured in triplicate wells. After washing the plates as previously 50 described, biotinylated antibody against FVIIa is added, 42ng/ml in ADB, 0.1ml/well, and the plates are incubated for at least 1 hour at room temp with rotation. The plates are washed, followed by incubation with streptavidin-HRP, 1:1000 in ADB, incubating at least 1 hour at room temp with rotation. After a final plate wash, the wells are developed with 0.1ml/well of Ultra-TMB, stopping the reaction with 0.05ml/well of 2M sulfuric acid. The stopped reactions are read at OD-450nm, and the data are analyzed and calibrated. The lower limit of quantitation (LLOQ) for the assay is typically 55 15-30 ng/ml FVIIa in 100% plasma.

Soluble Tissue Factor (sTF)-Based Modified PT Assay to Measure rFVIIa Activity

[0100] A Prothrombin Time (PT) assay was performed to measure the activity of human rFVIIa in HemA mouse *ex vivo* plasma samples.

[0101] Briefly, 50 μ L of sample containing 10% of HemA mouse plasma and 50% of human FVII-deficient plasma (George King Inc) in aPTT buffer (0.15 M NaCl, 0.05 M Tris pH 7.5, 0.1% BSA) were mixed with 50 μ L of sTF-PT reagent and incubated at 37°C for 30 sec. The sTF-PT reagent was composed of 1 volume of 2 μ M recombinant human soluble TF (sTF₁₋₂₂₁) and 1 volume of 8 μ M phospholipid vesicles (PS²⁰:PC⁴⁰:PE⁴⁰). Clotting was initiated by adding 50 μ L of 25 mM of CaCl₂ and the clotting time was recorded on a STA Coagulation Analyzer (Diagnostica Stago Inc). The standards consisted of rFVIIa (wt-rFVII or modified rFVIIa variants) diluted 2-fold serially from 200 to 0.78 ng/mL.

Efficacy of Desialylated V2 in Hemophilia A (HemA) Mice

Acute Tail Cut Efficacy Study

[0102] To determine the blood loss, mice were anesthetized with isoflurane and the tails were placed in 37-38°C warmed 0.9% saline in 15 ml plastic tubes for 10 min. The tail was cut at 4 mm from the tip by scalpel and immediately placed back into a separate pre-warmed 15 ml plastic tube containing 10 ml of saline. The mouse was allowed to bleed freely over 40 min. Desialylated V2 and F7 were dosed intravenously either 5 minutes after or 15- and 30-minutes before tail cut injury. Blood loss was quantified gravimetrically by weighing tubes before and after blood was collected.

Tail Vein Transection Efficacy (TVT) Study

[0103] HemA mice were dosed with desialylated V2 or F7 by tail vein injection at 1 hr before or 5 min after tail vein transection injury. Appropriate anesthesia was used. The tail vein was transected with a #11 scalpel straight blade, and a timer was started. The mouse was then returned to its individual clean cage with white paper bedding (Versi-Dri™) placed on top of a 4X8 inch heating pad. The animal activity status was monitored hourly for the next 9 hrs and at the 24 hr time point. Any mouse that showed signs of reduced activity level was noted on the monitor form and any mouse that showed signs of excessive blood loss was euthanized immediately.

Thrombin-Antithrombin (TAT) Assay in HemA Mouse Plasma

Reagents:

[0104] (1) Capture Antibody: Anti-thrombin polyclonal antibody from Enzyme Research Labs, Cat#TAT-EIA-C.; (2) Detection Antibody: HRP-conjugated anti-AT-III polyclonal antibody, from Enzyme Research Labs, Cat#TAT-EIA-D, (3) Assay Diluent: from Enzyme Research Labs, Cat#TAT-EIA-D, (4) HRP Substrate: Amplex Red, Invitrogen, cat#A12216, (5) Alpha-Thrombin: from Enzyme Research Labs, Cat#HT-1002a, stored at -80°C, (6) AT-III: from Enzyme Research Labs, Cat# HAT, stored at -80°C, (7) BSA: from Sigma, Cat# A-7030; (8) AT-III Deficient Plasma: Purchased from Enzyme Research Labs, Cat: AT-DP, stored at -80°C.

Buffers

[0105] (1) TAT standard buffer: 20 mM Tris-HCl, pH7.4, 0.15M NaCl, 1 mM EDTA, 0.05 U/mL heparin; (2) Coating Buffer: 1 tablet of bi-carbonate + 100ml dH2O, store at 4°C; (3) Blocking buffer: 2%BSA-PBS; (4) Sample dilution buffer: add 0.1M HEPES, pH7.4, 0.15M NaCl, 1% BSA, 0.05% Tween20, filter and aliquot, store at -20°C; (5) Substrate buffer: add 50 μ L of 5 mg/mL Amplex Red, 20 μ L of 3% H2O2 to PBS buffer. Mix, fresh prepared before adding to plates; (6) Preparation of 1 μ M TAT standard stock: add 100 μ L of human AT-III at 1.36 mg/ml and 5.93 μ L of human thrombin at 3.28 mg/mL to 419 μ L of TAT buffer, mix, incubate 10-20 min at 37°C; (7) Preparation of 60 nM TAT standard stock: Add 50 μ L of 1 μ M TAT complex to 783 μ L of AT-III deficient plasma, mix. Aliquot 50 μ L/vials, store at -80 °C.

Assay Procedure

[0106]

1. Dilute anti-thrombin pAb (capture antibody) in bicarbonate buffer (1:100 dilution: for one 96-well plate, add 110 μ L of antibody to 11 mL of bicarbonate buffer).
2. Add 100 μ L diluted coating antibody to each well on a 2HB Immulon 96-well plate. Tap plate gently to ensure all

liquid to cover bottom of plate. Seal the plate and incubate overnight at 4°C.

3. Wash 4 times with 300 μ L Wash buffer in an automated plate washer. Following the last wash, invert the plate and tap it against a clean paper towel.

5. Add 150 μ L of Blocking buffer (2% BSA-PBS) to each well. Seal the plate and incubate at Room temperature for 1.5 hour.

5. Wash 4 times with 300 μ L Wash buffer with an automated plate washer. Following the last wash, invert the plate and tap it against a clean paper towel.

6. Add 100 μ L of Standard, sample and QC to each well in triplicates and incubate plates at room temperature for 2 hr at room temperature.

10. Wash 4 times with 300 μ L Wash buffer with an automated plate washer. Following the last wash, invert the plate and tap it against a clean paper towel.

7. Add 100 μ L HRP-detection antibody (1/100, add 110 μ L of antibody to 11 mL of conjugate diluent), to each well. Seal the plate and incubate for 1 hour at room temperature.

9. Wash 4 times with 300 μ L Wash buffer with an automated plate washer. Following the last wash, invert the plate and tap it against a clean paper towel.

15. Add 70 μ L of Amplex Rd substrate (fresh prepared) to each well.

11. Place the plate in the dark at room temperature and incubate for 15-30 min.

12. Read the plate at OD485nm/595nm.

13. Plot standards with 4-parameter curve fit; Concentration from Controls and each sample were calculated from standard in each ELISA plate.

Efficacy of Desialylated V2 in Coagulation-Competent Mice

[0107] An acute tail cut study was performed to determine the efficacy of dV2 in coagulation-competent mice. Coagulation-competent mice were anesthetized with isoflurane and the tails were placed in 37-38°C warmed 0.9% saline in 15 ml plastic tubes for 10 min. After an iv administration of 5 mg/kg tissue plasminogen activator (tPA), the tail was cut at 50 mm from the tip by scalpel and placed back into a separate pre-warmed 15 ml plastic tube containing 10 ml of saline. Desialylated V2 and F7 were dosed intravenously immediately after tail cut injury. The mouse was allowed to bleed freely over 45 min. Blood loss was quantified gravimetrically by weighing tubes before and after blood was collected.

Results

In vitro Characterization of Desialylated or Deglycosylated Proteins

[0108] The heavy chain of dV2 was analyzed by LC-TOF MS. Analysis showed that the N-Glycans on the heavy chain contained no sialic acid post sialidase treatment. Such analysis on the light chain was complicated by the presence of the Gla domain. To get a global picture of sialic acid content of the treated molecule, Sialic Acid Fluorescence Labeling was carried out. This method showed that greater than 99.9% of sialic acid was removed on V2 during the desialylation process. FIG. 7 shows the analysis of desialylated V2 for sialic acid content. LC-TOF MS analysis and Sialic Acid Fluorescent Labeling were utilized. Sialic acid content analysis was carried out on dV1 as well with similar results (data not shown). The desialylated molecules were tested for activity by both the phospholipid-dependent Xa activation and phospholipid-dependent TGA assay. The PL-Xa and PL-TGA assays demonstrated that activity of the proteins post desialylation was not reduced. (See FIG. 9 and FIG. 10). FIG. 9 shows PL-FXa activation assay on desialylated proteins. Desialylated V1 and V2 (dV1, dV2) were tested for activity using the Phospho-lipid FXa activation assay. Both desialylated proteins had slightly higher activity in this assay as compared to their unmodified parental molecules. FIG. 10 shows PL-TGA assay on desialylated proteins. By PL-TGA, dV2 and dV1 exhibited slightly increased activity over their unmodified parental molecules. Results were normalized to F7.

[0109] The PL-Xa assay consistently showed a measurable increase in activity of dV2 and dV1 over their unmodified parental molecules. The hypoglycosylated wtFVIIa, V2, and V1 molecules were expressed (FIG. 11) and tested as crude expression extracts for both expression and activity. FIG. 11 shows expression of hypoglycosylated FVII variants. Media samples 4 days post electroporation were analyzed for FVII expression. Western blot analysis using an anti-Gla domain antibody shows expression of the variants. Removal of the N-Glycan sites did not appear to affect activity when normalized for expression levels. FIG. 12 shows determination of "specific activity" of hypoglycosylated FVII variants using transfection supernatants. The activity of the crude expression supernatants from two transient transfections of the hypoglycosylated variants were assayed by the Xa activation assay. When normalized for expression as measured by ELISA, no decrease in activity as a result of N-glycan removal was noted. As expected in this assay, V1 and F7 proteins had similar activities while V2 molecules had lower activity, a result of its TF-independence. This was further demonstrated by the PL-TGA activity assay carried out on purified hypoglycosylated V2 with only 2N-Glycans (N322 and N145) referred

to as pMB121. FIG. 13 shows PL-TGA assay on purified hypoglycosylated variant pMB121. By PL-TGA assay, pMB1212 shows enhanced activity over F7 similar to unmodified V2. *In vitro* clearance of these molecules was tested in a hepatocyte clearance model. dV2 demonstrated significant clearance in this model over unmodified V2 (FIG. 14), while marginal or no increase in clearance was seen for the hypoglycosylated variants (FIG. 15).

5

Rat PK and HemA PK

10

[0110] Pharmacokinetic studies in Sprague Dawley rats demonstrated that desialylated and hypoglycosylated proteins cleared significantly faster than their unmodified counterparts as measured by a FVII ELISA. FIG. 16 shows the rat pharmacokinetic results. Half-lives of the desialylated V2 and V1 were significantly shorter than their unmodified parental molecules in Sprague Dawley rats as measured by FVII ELISA. This was true for desialylated V2, desialylated V1, and pMB121 (hypoglycosylated V2). The t_{1/2} for both the desialylated molecules was less than 1 min whereas the t_{1/2} of their parental proteins was approximately 2.5 hrs. The clearance of the hypoglycosylated V2 molecule pMB121 was equivalent to that of F7 with a t_{1/2} of 1.6 hrs. The PK study in HemA mice had a similar result with dV2 and F7 having half-lives of approximately 3 min and 2.6 hrs, respectively (FIG. 17 (A)). The short half-life was confirmed by sTF-PTT clotting assay (FIG. 17 (B)). FIG. 17 shows HemA PK results. The half-life of desialylated V2 was significantly shorter than its unmodified parental molecule in HemA mice as measured by A) FVII ELISA and B) the sTF-PT assay.

15

HemA Efficacy Models

20

[0111] dV2 was tested in HemA mice for efficacy. Using the HemA tail cut model, dV2 was shown to be efficacious at a dose of 1mg/kg (bolus, iv). By comparison, in this model, the efficacious dose for F7 was 2.5mg/kg (bolus, iv). These results demonstrate that dV2 is more efficacious than F7 (FIG. 18 (A)). This model was also utilized to show that the efficacy of dV2 is cleared faster than that of F7 (FIG. 18 (B)). FIG. 18 shows the results of the desialylated V2 efficacy study in HemA mice. Studies with dV2 show that this molecule is A) more efficacious than and B) has faster efficacy clearance than F7 in the HemA tail cut model.

25

[0112] Using the more sensitive TTV model, the faster efficacy clearance of dV2 over F7 was also demonstrated and the efficacious dose confirmed. FIG. 19 shows a dV2 efficacy study in TTV HemA model. TTV studies using an efficacy model (TTV) with higher sensitivity confirmed dV2 has faster efficacy clearance than F7. Thrombin Anti-Thrombin (TAT) measurements as a marker of thrombogenicity performed at 30 and 60min post administration in HemA mice showed significantly lower levels for dV2. FIG. 20 shows TAT measurements. In HemA mice, dV2 administered at its efficacious dose (1mg/kg) generated less Thrombin Anti-Thrombin (TAT) than the efficacious dose of F7 (2.5mg/kg). This data, taken with the efficacy data, would suggest that dV2 has a more favorable therapeutic index than F7.

35

Efficacy in Coagulation Competent Mice

40

[0113] dV2 was tested in tPA-treated, coagulation-competent mice for efficacy. Using the tail cut model, dV2 was shown to be efficacious at doses of 0.3-1mg/kg (bolus, iv). By comparison, in this model, the efficacious dose for F7 was 5mg/kg (bolus, iv). These results demonstrate that dV2 is more efficacious than F7 (FIG. 21). FIG. 21 shows the results of the desialylated V2 efficacy study in tPA-treated, coagulation-competent mice.

Clearance and Efficacy of Desialylated Wild-Type Factor VII (dWT VIIa)

45

[0114] Desialylated wild-type Factor VII (dWT VIIa) was produced as described above using NovoSeven® obtained from Novo Nordisk as the starting Factor VII material and desialylating that starting polypeptide using soluble sialidase enzyme, as described above. The dWT VIIa was found to have a purity of >99%, low endotoxin, and no detectable sialic acid. Additionally, mass spectrometry analysis showed selective removal of sialic acid.

50

[0115] The activity of this dWT VIIa material was analyzed and compared to wild-type Factor VII using the Biophen FVII chromogenic assay and modified PT Assay, as described above. Each of these analyses showed the dWT VIIa to have nearly identical activity to the wild-type Factor VII polypeptide.

[0116] Clearance of dWT VIIa and wild-type Factor VII (1 mg/kg) were also analyzed and compared using the human tissue factor knock-in (TFKI) mice mouse model. As shown in Fig. 22, the half-life of dWT VIIa was significantly shorter than wild-type Factor VII and clearance (ml/h/kg) was more than 40 times faster.

55

[0117] The efficacy of dWT VIIa in comparison to wild-type Factor VII was investigated using TFKI mice and the tail cut method described above. Briefly, 5 mg/kg tPA was injected intravenously into the mice, followed by clipping of the tail 50 mm from the tip. Wild-type Factor VII (NovoSeven®) or dWT VIIa were then injected intravenously with dosages ranging from 1-6mg/kg. Blood was then collected from the tail for 45 minutes, with unstable clots being disrupted every six minutes throughout the collection period. As shown in Fig. 23, dWT VIIa was surprisingly found to be significantly

more efficacious than wild-type Factor VII. More specifically, a 3 mg/kg dose of dWT VIIa caused reduced blood loss in comparison to a 6 mg/kg dose of wild-type Factor VII. Given the results of this analysis, it was determined that 2 mg/kg dWT VIIa is a bioequivalent dose to 6 mg/kg wild-type factor VII.

[0118] The ability of dWT VIIa and wild-type Factor VII (NovoSeven®) to cause systemic coagulation was also investigated by the Thrombin Anti-Thrombin (TAT) method described above. Mice were treated with bioequivalent doses of dWT VIIa (2 mg/kg) and wild-type Factor VII (6 mg/kg) and then the formation of TAT complexes was measured by ELISA. As shown in Fig. 24, the wild-type NovoSeven® Factor VII generated a significantly higher level of TAT than dWT VIIa. Given the fact that dWT VIIa generated only baseline TAT levels, this experiment suggests that this dose of dWT VIIa produces no observable systemic coagulation, despite the fact that the polypeptide is as efficacious as wild-type Factor VII.

[0119] Further, the ability of dWT VIIa and wild-type Factor VII (NovoSeven®) to cause thrombus formation was also investigated in a FeCl_3 thrombosis model. Mice were treated with bioequivalent doses of dWT VIIa (2 mg/kg) and wild-type Factor VII (6 mg/kg) 15 minutes prior to initiation of the thrombosis study. Thrombosis was then initiated by administration of a 3.25 % FeCl_3 solution and then thrombus formation was measured by Doppler for 30 minutes. The resulting blood flow data was plotted on a blood flow versus time graph and then the percentage of the area under the curve for the control sample was calculated to determine the reduction in blood flow caused by thrombus formation for each of the Factor VII treatment groups. As shown in Fig. 25, the wild-type NovoSeven® Factor VII generated a significantly reduced blood flow (average approx.. 40%), while the dWT VIIa showed nearly no reduction in blood flow (average >90%). This experiment demonstrated that the given dose of dWT VIIa produced greatly reduced thrombus formation as compared to wild-type Factor VII.

[0120] The activity and efficacy of dWT VIIa compared to wild-type Factor VII was further investigated by examining the apparent binding affinities of these peptides for soluble tissue factor (sTF) using an SN-17c tripeptide fluorogenic substrate (HTI). As shown in Fig. 26, this analysis demonstrated that dWT VIIa (dF7) and wild-type Factor VII (F7) had equivalent apparent biding affinities for sTF. However, as shown in Fig. 27, in an experimental model examining the ability of these peptides to activate Factor X by titrating the Factor X concentration in the presence of sTF-Factor VII complexes (0.5 nM Factor VII [dWT VIIa or wild-type], 125 nM sTF), the Michaelis Menten kinetics for dWT VIIa and wild-type Factor VII demonstrate that dWT VIIa (dF7) can activate Factor X more effectively (approximately 2-fold) than wild-type Factor VII (F7). This data suggests that dWT VIIa is able to convert more Factor X to Factor Xa per Factor VII active site than its wild-type counterpart.

30 Discussion

[0121] There is an unmet medical need to develop a therapeutic drug that is efficacious for the treatment of acute bleeding but with reduced thrombogenicity. An efficacious Factor VII polypeptide with a short half-life would potentially result in a molecule with a larger therapeutic window suitable for use in acute bleeds.

[0122] V2 and V1 are two Factor VIIa variants (FIGS. 1- 3). These variants contain mutations to their Gla domains that increase their affinity for activated platelets and, in the case of V2, result in tissue factor independence. Both variants also have two additional N-Glycosylation sites, which results in a prolonged half-life as compared to wild type Factor VIIa, a trait that is advantageous for the treatment of hemophilia. However their use as treatments for acute bleeding would benefit from decreases in half-life. This modification would reduce the risk of off-target effects and, as a result, increase their therapeutic index. We have shown here that removal of the sialic acids present on the carbohydrate chains of V2 and V1 result in significantly faster clearance of the molecules in an *in vitro* hepatocyte clearance model. The hypoglycosylated variants did not clear faster in this *in vitro* model, which suggests that the mechanism of clearance between desialylated and hypoglycosyed molecules differ. *In vivo* studies conducted in Sprague Dawley rats demonstrated that the desialylated molecules (dV2 and dV1), as well as the hypoglycosylated variant pMB121, had a significantly decreased half-life. Interestingly, desialylated V2 and V1 both had increased clearance rates as compared to the reported rate for desialylated wild type FVIIa. (Appa et al, Thrombosis and Haemostasis 104.2/2010), a characteristic that may be due to their 2 additional N-glycans. One possible theory for this activity, without limitation on what is claimed herein, is that these extra N-Glycans would, upon desialylation, become additional ligands for the ASGPR or similar receptor and mediate faster clearance. Activity of these molecules was retained or increased in comparison to their parental molecules as measured by *in vitro* activity assays. Faster clearance of dV2 was further verified *in vivo* in a HemA mouse PK study and was shown to be efficacious in HemA tail clip and TVT studies.

[0123] Further, desialylation of wild-type Factor VII produced a Factor VII polypeptide that was cleared much more rapidly than wild-type, while also providing the surprising result of increased efficacy as shown in numerous experimental models.

[0124] Removal of the N-glycans or modification of the monosaccharide composition of the N-glycans of Factor VIIa or Factor VIIa variants results in faster clearing molecules. These molecules retain activity and are efficacious *in vivo*. The development of these fast clearing Factor VIIa molecules would be beneficial for the treatment of acute bleeding

EP 3 165 232 A1

indications, as well as potentially being an antidote for the various anti-coagulants on the market.

5

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

5 <110> Bayer HealthCare LLC
 Bauzon, Maxine
 Hermiston, Terry
 10 <120> Short Acting Factor VII Peptides
 15 <130> BHC115011 PCT
 20 <150> 61/754, 674
 <151> 2012-12-24
 25 <150> 61/787, 026
 <151> 2013-03-15
 30 <160> 18
 35 <170> PatentIn version 3.5
 40 <210> 1
 45 <211> 1218
 <212> DNA
 <213> Homo sapiens
 50 <400> 1
 55 gccaaacgcgt tcctggagga gctgcggccg ggctccctgg agagggagtg caaggaggag 60
 cagtgctct tagaggaggc ccgggagatc ttcaaggacg cggagaggac gaagctgttc 120
 tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc 180
 60 tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggaac 240
 65 tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag 300
 70 tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg 360
 75 ctggcagacg gggtgtcctg cacacccaca gttgaatatc catgtggaaa aatacctatt 420
 80 ctagaaaaaa gaaatgccag caaacccaa ggccgaattt tggggggcaa ggtgtgcccc 480
 85 aaaggggagt gtccatggca ggtcctgttg ttggtaatg gagctcagtt gtgtgggggg 540
 90 accctgatca acaccatctg ggtggtctcc gggcccaact gtttcacaa aatcaagaac 600
 95 tggaggaacc tgatcgcggt gctggcgag cacgacactca gcgagcacga cggggatgag 660
 100 cagagccggc gggtggcgca ggtcatcatc cccagcacgt acgtcccgaa caccaccaac 720
 105 cacgacatcg cgctgcgtccg cctgcaccag cccgtggtcc tcactgacca tgtggtgccc 780
 110 ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcattt 840
 115 gtcagcggct gggccagct gctggaccgt ggccgcacgg ccctggagct catggtcctc 900
 120 aacgtgcccc ggctgtatgac ccaggactgc ctgcagcagt cacggaaagggt gggagactcc 960
 125 ccaaatatca cggagtacat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc 1020
 130 aagggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc 1080
 135 atcgtcagct gggccaggc ctgcgcacc gtggccact ttgggtgtta caccagggtc 1140

tcccagtaca tcgagtggct gcaaaagctc atgcgctcag agccacgccc aggagtcctc	1200
ctgcgagccc catttccc	1218

5
 <210> 2
 <211> 1219
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> Variant human FVII nucleotide sequence

<400> 2	
gccaacgcgt tcctggagga gctgcggcag ggctccctgg agagggagtg caaggaggag	60
cagtgcctc tcgaggagga ccggagatc ttcaagacg cggagaggac gaagctgttc	120
tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc	180
tgcaaggacc agtccagtc statatctgc ttctgcctcc ctgccttcga gggccggAAC	240
tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag	300
tactgcagtg accacaacgg caccaagcgc tcctgtcggt gccacgaggg gtactctctg	360
ctggcagacg gggtgtcctg cacacccaca gttgaatatc catgtggaaa aatacctatt	420
ctagaaaaaa gaaatgccag caaacccaa ggcgaattt tggggggcaa ggtgtgcccc	480
aaaggggagt gtccatggca ggtcctgttg ttggtaatg gagctcagtt gtgtgggggg	540
accctgatca acaccatctg ggtggctcggc gcccggact gttcgacaa aatcaagaac	600
tggaggaacc tgatcgcggt gctggcgag cacgacactca gcgagcacga cggggatgag	660
cagagccggc gggtgtcgca ggtcatcatc cccagcacgt acgtcccggg caccaccaac	720
cacgacatcg cgctgcctcg cctgcaccag cccgtgaacc tcactgacca tgtggtgccc	780
ctctgcctgc ccgaacggac gttcttgag aggacgctgg cttcgtcgcttctcattt	840
gtcagcggct gggccagct gctggaccgt ggcggccacgg ccctggagct catggtcctc	900
aacgtgcccc ggctgtatgac ccaggactgc ctgcagcagt cacggaaagg gggagactcc	960
ccaaatatca cggagtacat gttcttgcc ggctactcgg atggcagcaa ggactcctgc	1020
aagggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc	1080
atcgtcagct gggccagggt ctgcgcaacc gtggccact ttgggtgtt caccagggtc	1140
tcccagtaca tcgagtggct gcaaaagctc atgcgctcag agccacgccc aggagtcctc	1200
ctgcgagccc catttccc	1219

55 <210> 3
 <211> 1218
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Variant human FVII nucleotide sequence

<400> 3

5	gccaacgcgt tcctggagga gctgcggcag ggctccctgg agagggagtg caaggaggag	60
	cagtgcct tcgaggaggc ccgggagatc ttcaaggacg cggagaggac gaagctgttc	120
	tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc	180
10	tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggaac	240
	tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag	300
	tactgcagtg accacaacgg caccaagcgc tcctgtcggt gccacgaggg gtactctctg	360
15	ctggcagacg gggtgtcctg cacacccaca gttgaatatc catgtggaaa aatacctatt	420
	ctagaaaaaa gaaatgccag caaacccaa ggccgaattt tgggggcaa ggtgtgcccc	480
20	aaaggggagt gtccatggca ggtcctgttg ttggtaatg gagctcagtt gtgtgggggg	540
	accctgatca acaccatctg ggtggtctcc gcgcccccact gtttcgacaa aatcaagaac	600
	tggaggaacc tgatcgcggt gctggcgag cacgacactca gcgagcacga cggggatgag	660
25	cagagccggc gggtgtcgca ggtcatcatc cccagcacgt acgtccggg caccaccaac	720
	cacgacatcg cgctgcctcg cctgcaccag cccgtgaacc tcactgacca tgtggtgc	780
	ctctgcctgc ccaaacggac gttctctgag aggacgctgg cttcgtcg cttctcattt	840
30	gtcagcggct ggggccagct gctggaccgt ggccgcacgg ccctggagct catggcctc	900
	aacgtgcccc ggctgatgac ccaggactgc ctgcagcagt cacgaaaggt gggagactcc	960
	ccaaatatca cggagttacat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc	1020
35	aagggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc	1080
	atcgtcagct ggggccaggg ctgcgcaacc gtggccact ttgggtgtta caccagggtc	1140
	tcccagtaca tcgagtggct gcaaaagctc atgcgctcag agccacgccc aggagtcc	1200
40	ctgcgagccc catttccc	1218

<210> 4

<211> 1218

<212> DNA

<213> Artificial Sequence

<220>

<223> Plasmid sequence

<400> 4

50	gccaacgcgt tcctggagga gctgcggccg ggctccctgg agagggagtg caaggaggag	60
	cagtgcct tcgaggaggc ccgggagatc ttcaaggacg cggagaggac gaagctgttc	120
55	tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc	180
	tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggaac	240

5	tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag	300
	tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg	360
	ctggcagacg gggtgtcctg cacaccaca gttgaatatic catgtggaaa aatacctatt	420
	ctagaaaaaa gaaatgccag caaacccaa ggccgaattt tggtggcaaa ggtgtgcccc	480
10	aaaggggagt gtccatggca ggtcctgttg ttgggtaatg gagctcagtt gtgtgggggg	540
	accctgatca acaccatctg ggtggtctcc gcggccact gttcgacaa aatcaagaac	600
	tggaggaacc tggatcgcggt gttggcgag cacgacactca gcgagcacga cggggatgag	660
15	cagagccggc gggtgtcgca ggtcatcatc cccagcacgt acgtcccgcc caccaccaac	720
	cacgacatcg cgctgctccg cctgcaccag ccgtggtcc tcactgacca tgtggtgccc	780
	ctctgcctgc ccgaacggac gttcttgag aggacgctgg cttcgtcg cttcttattt	840
20	gtcagcggt gggccagct gttggaccgt ggccgcacgg ccctggagct catggtcctc	900
	aacgtgcccc ggctgtatgac ccaggactgc ctgcagcagt cacgaaagggt gggagactcc	960
	ccaaatatca cggagtacat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc	1020
25	aagggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc	1080
	atcgtcagct gggccagggt ctgcgcaacc gtggccact ttgggtgtta caccagggtc	1140
	tcccagtaca tcgagtggct gcaaaagctc atgcgctcag agccacgccc aggagtcc	1200
30	ctgcgagccc catttccc	1218

<210> 5
 <211> 1218
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Plasmid sequence

40	<400> 5 gcacacgcgt tcctggagga gctgcggccg ggctccctgg agagggagtg caaggaggag	60
	cagtgccttc tcgaggaggc ccgggagatc ttcaaggacg cggagaggac gaagctgttc	120
45	tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc	180
	tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggAAC	240
	tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag	300
50	tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg	360
	ctggcagacg gggtgtcctg cacaccaca gttgaatatic catgtggaaa aatacctatt	420
	ctagaaaaaa gacaggccag caaacccaa ggccgaattt tggtggcaaa ggtgtgcccc	480
55	aaaggggagt gtccatggca ggtcctgttg ttgggtaatg gagctcagtt gtgtgggggg	540

5	accctgatca acaccatctg ggtggtctcc gcggcccact gtttcacaa aatcaagaac	600
	tggaggaacc tgatcgcggt gctggcgag cacgacctca gcgagcacga cggggatgag	660
	cagagccggc gggtggcgca ggtcatcatc cccagcacgt acgtcccggg caccaccaac	720
	cacgacatcg cgctgctccg cctgcaccag cccgtggtcc tcactgacca tgtggtgcac	780
10	ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcattg	840
	gtcagcggct gggccagct gctggaccgt ggccacgg ccctggagct catggtcctc	900
	aacgtgcccc ggctgatgac ccaggactgc ctgcagcagt cacggaaggt gggagactcc	960
15	ccaaatatca cggagttacat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc	1020
	aaggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc	1080
	atcgtcagct gggccaggg ctgcgcaacc gtggccact ttgggtgta caccagggtc	1140
20	tcccagtaca tcgagtggtc gcaaaagctc atgcgctcag agccacgccc aggagtctc	1200
	ctgcgagccc catttccc	1218
25	<210> 6	
	<211> 1218	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> Plasmid sequence	
	<400> 6	
	gcacacgcgt tcctggagga gctgcggccg ggctccctgg agagggagtg caaggaggag	60
	cagtgcctc tcgaggaggc ccggagatc ttcaaggacg cggagaggac gaagctgttc	120
35	tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggtctcc	180
	tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggAAC	240
	tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag	300
40	tactgcagtgc accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg	360
	ctggcagacg ggggtgtcctg cacacccaca gttgaatatc catgtggaaa aatacctatt	420
	ctagaaaaaa gaaatgccag caaacccaa ggccgaattt tggggggcaa ggtgtgcccc	480
45	aaaggggagt gtccatggca ggtcctgttg ttgggtgaatg gagctcagtt gtgtgggggg	540
	accctgatca acaccatctg ggtggtctcc gcggcccact gtttcacaa aatcaagaac	600
	tggaggaacc tgatcgcggt gctggcgag cacgacctca gcgagcacga cggggatgag	660
50	cagagccggc gggtggcgca ggtcatcatc cccagcacgt acgtcccggg caccaccaac	720
	cacgacatcg cgctgctccg cctgcaccag cccgtggtcc tcactgacca tgtggtgcac	780
55	ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcattg	840
	gtcagcggct gggccagct gctggaccgt ggccacgg ccctggagct catggtcctc	900

5	aacgtcccc ggctgatgac ccaggactgc ctgcagcagt cacggaaggt gggagactcc	960
	ccacagatca cggagtagatcat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc	1020
	aaggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacggc	1080
	atcgtcagct gggccaggg ctgcgcaacc gtggccact ttgggtgt a caccagggtc	1140
10	tccca gtttaca tcgagtggct gcaaaagctc atgcgctca g agccacgccc aggagtcc	1200
	ctgcgagccc catttccc	1218
15	<210> 7	
	<211> 1218	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Plasmid sequence	
25	<400> 7	
	gcacacgcgt tcctggagga gctgcggccg ggctccctgg agagggagtg caaggaggag	60
	cagtgcatact tagaggagga cggggagatc ttcaaggacg cggagaggac gaagctgttc	120
30	tggatttctt acagtgatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc	180
	tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggAAC	240
	tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag	300
35	tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacaggg gtactctcg	360
	ctggcagacg gggtgtcctg cacaccaca gttgaatatc catgtggaaa aatacattt	420
	ctagaaaaaa gacaggccag caaacccaa ggccgaattt tggggggcaa ggtgtcccc	480
40	aaaggggagt gtccatggca ggtcctgttg ttggtaatg gagctcagtt gtgtgggggg	540
	accctgatca acaccatctg ggtggcttcc gcggccact gttcgacaa aatcaagaac	600
45	tggaggaacc tgatcgccgt gctggcgag cacgacactca gcgagcacga cggggatgag	660
	cagagccggc gggtgtcgca ggtcatcatc cccagcacgt acgtcccgagg caccaccaac	720
	cacgacatcg cgctgctccg cctgcaccag cccgtggtcc tcactgacca tgtggtgccc	780
50	ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcatg	840
	gtcagcggct gggccagct gctggaccgt ggccacgg ccctggagct catggcctc	900
	aacgtcccc ggctgatgac ccaggactgc ctgcagcagt cacggaaggt gggagactcc	960
55	ccacagatca cggagtagatcat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc	1020
	aaggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacggc	1080
	atcgtcagct gggccaggg ctgcgcaacc gtggccact ttgggtgt a caccagggtc	1140
	tccca gtttaca tcgagtggct gcaaaagctc atgcgctca g agccacgccc aggagtcc	1200

5	ctgcgagccc catttccc	1218
	<210> 8	
	<211> 1218	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Plasmid sequence	
	<400> 8	
	gccaacgcgt tcctggagga gctgcggcag ggctccctgg agagggagtg caaggaggag	60
	cagtgcctc tcgaggaggc ccgggagatc ttcaagacg cggagaggac gaagctgttc	120
15	tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc	180
	tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggaac	240
20	tgtgagacgc acaaggatga ccagctgatc tttgtgaacg agaacggcgg ctgtgagcag	300
	tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg	360
	ctggcagacg ggggtgcctg cacaccaca gttgaatatc catgtggaaa aatacctatt	420
25	ctagaaaaaa gaaatgccag caaacccaa gcccgaattt tggggggcaa ggtgtgcccc	480
	aaaggggagt gtccatggca ggtcctgttg ttggtaatg gagctcagtt gtgtgggggg	540
	accctgatca acaccatctg ggtggctcc gcccggact gttcgacaa aatcaagaac	600
30	tggaggaacc tgatgcggc gctggcgcag cacgaccta gacgacacg cggggatgag	660
	cagagccggc ggggtggcgca ggtcatcatc cccagcacgt acgtccggg caccaccaac	720
	cacgacatcg cgctgcctcg cctgcaccag cccgtggtcc tcactgacca tttgtgccc	780
35	ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcattt	840
	gtcagccggct gggccagct gctggaccgt ggcgccacgg ccctggagct catggccctc	900
	aacgtgcccc ggctgtatgac ccaggactgc ctgcagcagt cacggaaagg gggagactcc	960
40	ccaaatatca cgaggtacat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc	1020
	aagggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc	1080
45	atcgtagct gggccaggg ctgcgcaacc gtggccact ttgggtgtt caccagggtc	1140
	tcccagtaca tcgagtggct gcaaaagctc atgcgctcag agccacgccc aggagtcc	1200
	ctgcgagccc catttccc	1218
50	<210> 9	
	<211> 1218	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Plasmid sequence	

<400> 9
gccaacgcgt tcctggagga gctgcggcag ggctccctgg agagggagtg caaggaggag 60
cagtgcct tcgaggaggc ccgggagatc ttcaagacg cgagaggac gaagctgttc 120
tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc 180
tgcaaggacc agtccagtc ctatatctgc ttctgcctcc ctgcattcga gggccggAAC 240
tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag 300
tactgcagt accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg 360
ctggcagacg gggtgtcctg cacaccaca gttaatatc catgtggaaa aatacctatt 420
ctagaaaaaa gacaggccag caaacccaa gcccgaattt tggggggcaa ggtgtgcccc 480
aaaggggagt gtccatggca gtcctgttg ttggtaatg gagtcagtt gtgtgggggg 540
accctgatca acaccatctg ggtggctcc gcggccact gttcgacaa aatcaagaac 600
tggaggaacc tgatcgcggt gtcggcgag cacgacactca gcgagcacga cggggatgag 660
cagagccggc gggtgtcgca gtcatcatc cccagcacgt acgtccggg caccaccaac 720
cacgacatcg cgctgcctcg cctgcaccag cccgtggtcc tcactgacca tgtggtgc 780
ctctgcctgc ccgaacggac gttcttgag aggacgctgg cttcgctcg cttctcattt 840
gtcagcggt gggccagct gtcggaccgt ggcgccacgg ccctggagct catggtcctc 900
aacgtgcccc ggctgtatgac ccaggactgc ctgcagcagt cacggaaagggt gggagactcc 960
ccaaatatca cggagtagat gttcttgcc ggctactcgg atggcagcaa ggactcctgc 1020
aagggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc 1080
atcgtcagct gggccaggg ctgcgcaacc gtggccact ttggggtgta caccagggtc 1140
tcccagtaca tcgagtggct gcaaaagctc atgcgctcag agccacgccc aggagtcc 1200
ctgcgacccc catttccc 1218

```
<210> 10
<211> 1218
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Plasmid sequence

<400> 10 gccaacgcgt tcctggagga gctgcggcag ggctccctgg agagggagtg caaggaggag 60
cagtgccttc tcgaggaggc ccggggagatc ttcaagacg cggagaggac gaagctgttc 120
tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc 180
tgcaaggacc agctccagtc ctatatactgc ttctgcctcc ctgccttcga gggccggaaac 240
tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag 300

EP 3 165 232 A1

5	tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg	360
	ctggcagacg gggtgtcctg cacaccaca gttgaatatac catgtggaaa aatacctatt	420
	ctagaaaaaa gaaatgccag caaacccaa ggccgaattt tgggggcaa ggtgtgcccc	480
10	aaaggggagt gtccatggca ggtcctgtt gttgtgaatg gagctcagtt gtgtgggggg	540
	accctgatca acaccatctg ggtggtctcc gggcccact gtttcgacaa aatcaagaac	600
	tggaggaacc tgatcgcggt gctggcgag cacgaccta gcgagcacga cggggatgag	660
	cagagccggc gggtggcgca ggtcatcatc cccagcacgt acgtccccggg caccaccaac	720
15	cacgacatcg cgctgctccg cctgcaccag cccgtggtcc tcactgacca tgggtgccc	780
	ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcattt	840
	gtcagcggct gggccagct gctggaccgt ggccacgg ccctggagct catggtcctc	900
20	aacgtgcccc ggctgtatgac ccaggactgc ctgcagcagt cacggaaggt gggagactcc	960
	ccacagatca cggagtacat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc	1020
	aaggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc	1080
25	atcgtcagct gggccaggg ctgcgcaacc gtggccact ttgggtgtt caccagggtc	1140
	tcccagtaca tcgagtggct gcaaaagctc atgcgctcag agccacgccc aggagtcc	1200
	ctgcgagccc catttccc	1218
30	<210> 11	
	<211> 1218	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Plasmid sequence	
	<400> 11	
	gccaacgcgt tcctggagga gctgcggcag ggctccctgg agagggagtg caaggaggag	60
40	cagtgcctc tcgaggaggc ccggagatc ttcaagacg cggagaggac gaagctgttc	120
	tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgcacaa tgggggtctcc	180
	tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggaac	240
45	tgtgagacgc acaaggatga ccagctgatc tgggtgaacg agaacggcgg ctgtgagcag	300
	tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg	360
	ctggcagacg gggtgtcctg cacaccaca gttgaatatac catgtggaaa aatacctatt	420
50	ctagaaaaaa gacaggccag caaacccaa ggccgaattt tgggggcaa ggtgtgcccc	480
	aaaggggagt gtccatggca ggtcctgtt gttgtgaatg gagctcagtt gtgtgggggg	540
55	accctgatca acaccatctg ggtggtctcc gggcccact gtttcgacaa aatcaagaac	600
	tggaggaacc tgatcgcggt gctggcgag cacgaccta gcgagcacga cggggatgag	660

5	cagagccggc gggtgtcgca ggtcatcatc cccagcacgt acgtcccgaa caccaccaac	720
	cacgacatcg cgctgctccg cctgcaccag cccgtggtcc tcactgacca tgtggtgccc	780
10	ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcatg gtcagcggct gggccagct gctggaccgt ggccacgg ccctggagct catggtcctc aacgtgcccc ggctgatgac ccaggactgc ctgcagcagt cacggaaggt gggagactcc ccacagatca cggagtacat gttctgtgcc ggctactcgg atggcagcaa ggactcctgc aaggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cctgacgggc atcgtcagct gggccaggg ctgcgaacc gtggccact ttgggtgta caccagggtc tcccaagtaca tcgagtggct gcaaaagctc atgcgctcaag agccacgccc aggagtcctc ctgcgagccc catttccc	1218
15		
20	<210> 12 <211> 1218 <212> DNA <213> Artificial Sequence	
25	<220> <223> Plasmid sequence	
30	<400> 12 gccaacgcgt tcctggagga gctgcggcag ggctccctgg agagggagtg caaggaggag cagtgctcct tcgaggagga ccggagatc ttogaagacg aagaggaaac gaagctgtc tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc tgcaaggacc agctccagtc cstatatctgc ttctgcctcc ctgccttcga gggccggAAC tggagacgc acaaggatga ccagctgatc tggtaacg agaacggcgg ctgtgagcag tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg ctggcagacg gggtgtcctg cacaccaca gttgaatatc catgtggaaa aatacctatt ctagaaaaaa gaaatgccag caaacccaa ggccgaattt tggggggcaa ggtgtgcccc aaaggggagt gtccatggca ggtcctgttg ttggtaatg gagctcagtt gtgtgggggg accctgatca acaccatctg ggtggctc gcgcccaact gttcgacaa aatcaagaac tggaggaacc tgatgcggc gctggcgag cacgacactca gcgagcacga cggggatgag cagagccggc gggtgtcgca ggtcatcatc cccagcacgt acgtcccgaa caccaccaac cacgacatcg cgctgctccg cctgcaccag cccgtggtcc tcactgacca tgtggtgccc ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcatg gtcagcggct gggccagct gctggaccgt ggccacgg ccctggagct catggtcctc aacgtgcccc ggctgatgac ccaggactgc ctgcagcagt cacggaaggt gggagactcc	60 120 180 240 300 360 420 480 540 600 660 720 780 840 900 960
35		
40		
45		
50		
55		

ccaaatatca	cgaggatcacat	gttctgtgcc	ggctactcgg	atggcagcaa	ggactcctgc	1020	
aagggggaca	gtggaggccc	acatgccacc	caactaccggg	gcacgtggta	cctgacgggc	1080	
5	atcgtcagct	ggggccaggg	ctgcgaacc	gtgggccact	ttggggtgta	caccagggtc	1140
tcccagtaca	tcgagtggct	gcaaaagctc	atgcgctcaag	agccacgccc	aggagtcctc	1200	
ctgcgagccc	catttccc					1218	
10							
<210>	13						
<211>	1218						
<212>	DNA						
<213>	Artificial Sequence						
15							
<220>							
<223>	Plasmid sequence						
<400>	13						
20	gccaacgcgt	tcctggagga	gctgcggcag	ggctccctgg	agagggagtg	caaggaggag	60
cagtgcct	tcgaggaggo	ccgggagatc	ttcgaagacg	aagaggaaac	gaagctgttc		120
tggatttttt	acagtgtatgg	ggaccagtgt	gcctcaagtc	catgccagaa	tgggggctcc		180
25	tgcaaggacc	agctccagtc	ctatatctgc	ttctgcctcc	ctgccttcga	ggcccggaac	240
tgtgagacgc	acaaggatga	ccagctgatc	tgtgtgaacg	agaacggcgg	ctgtgagcag		300
tactgcagt	accacacggg	caccaagcgc	tcctgtcggt	gccacgaggg	gtactctctg		360
30	ctggcagacg	gggtgtcctg	cacaccaca	gttgaatatc	catgtggaaa	aatacattt	420
ctagaaaaaa	gacaggccag	caaacccaa	ggccgaattt	tggggggcaa	gtgtgcccc		480
35	aaaggggagt	gtccatggca	ggtcctgttg	ttgggtgaatg	gagctcagtt	gtgtgggggg	540
accctgatca	acaccatctg	ggtggcttcc	gcggcccaact	gtttcgacaa	aatcaagaac		600
tggaggaacc	tgatcgcggt	gctggcgag	cacgacactca	gcgagcacga	cggggatgag		660
40	cagagccggc	gggtggcgca	ggtcatcatc	cccagcacgt	acgtcccggg	caccaccaac	720
cacgacatcg	cgctgctccg	cctgcaccag	cccggtggtcc	tcactgacca	tgtggtgccc		780
ctctgcctgc	ccgaacggac	gttctctgag	aggacgctgg	cttcgtgcg	cttctcattt		840
45	gtcagcggt	ggggccagct	gctggaccgt	ggcgccacgg	ccctggagct	catggtcctc	900
aacgtgcccc	ggctgtatgac	ccaggactgc	ctgcagcagt	cacgaaagg	gggagactcc		960
ccaaatatca	cgaggatcacat	gttctgtgcc	ggctactcgg	atggcagcaa	ggactcctgc	1020	
50	aagggggaca	gtggaggccc	acatgccacc	caactaccggg	gcacgtggta	cctgacgggc	1080
atcgtcagct	ggggccaggg	ctgcgaacc	gtgggccact	ttggggtgta	caccagggtc		1140
tcccagtaca	tcgagtggct	gcaaaagctc	atgcgctcaag	agccacgccc	aggagtcctc		1200
55	ctgcgagccc	catttccc					1218

<210> 14
 <211> 1218
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Plasmid sequence

 <400> 14

gccaacgcgt	tcctggagga	gctgcggcag	ggctccctgg	agagggagtg	caaggaggag	60
cagtgcct	tcgaggaggc	ccgggagatc	ttcgaagacg	aagaggaaac	gaagctgttc	120
tggatttctt	acagtgtatgg	ggaccagtgt	gcctcaagtc	catgccagaa	tgggggctcc	180
tgcaaggacc	agctccagtc	ctataatctgc	ttctgcctcc	ctgccttcga	gggcggaaac	240
tgtgagacgc	acaaggatga	ccagctgatc	tgtgtgaacg	agaacggcgg	ctgtgagcag	300
tactgcagtg	accacacggg	caccaagcgc	tcctgtcggt	gccacgaggg	gtactctctg	360
ctggcagacg	gggtgtcctg	cacacccaca	gttgaatatc	catgtggaaa	aatacctatt	420
ctagaaaaaa	gaaatgccag	caaacccaa	ggccgaattt	tggggggcaa	ggtgtgcccc	480
aaaggggagt	gtccatggca	ggtcctgttg	tttgtgaatg	gagctcagtt	gtgtgggggg	540
accctgatca	acaccatctg	ggtggctcc	gccccccact	gttgcacaa	aatcaagaac	600
tggaggaacc	tgatgcgggt	gctggcgag	cacgacactca	gcgagcacga	cggggatgag	660
cagagccggc	gggtggcgca	ggtcatcatc	cccagcacgt	acgtccccgg	caccaccaac	720
cacgacatcg	cgctgctccg	cctgcaccag	ccctgtggcc	tcactgacca	tgtggtgccc	780
ctctgcctgc	ccgaacggac	gttctctgag	aggacgctgg	cattcgtgcg	cttctcattt	840
gtcagcggct	ggggccagct	gctggaccgt	ggccacacgg	ccctggagct	catggtcctc	900
aacgtgcccc	ggctgtatgac	ccaggactgc	ctgcagcagt	cacgaaagg	gggagactcc	960
ccacagatca	cggagtacat	gttctgtgcc	ggctactcgg	atggcagcaa	ggactcctgc	1020
aagggggaca	gtggaggccc	acatgcacc	cactaccggg	gcacgtggta	cctgacgggc	1080
atcgtcagct	ggggccaggg	ctgcgcaacc	gtggggccact	ttgggggtgta	caccagggtc	1140
tcccagtaca	tcgagtggct	gcaaaagctc	atgcgctcag	agccacgccc	aggagtccctc	1200
ctgcgagccc	catttccc					1218

<210> 15
 <211> 1218
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Plasmid sequence

 <400> 15

gccaacgcgt	tcctggagga	gctgcggcag	ggctccctgg	agagggagtg	caaggaggag	60
------------	------------	------------	------------	------------	------------	----

cagtgcctc tcgaggaggc ccgggagatc ttcaagacg aagaggaaac gaagctgttc 120
 tggatttctt acagtgtatgg ggaccagtgt gcctcaagtc catgccagaa tgggggctcc 180
 5 tgcaaggacc agctccagtc ctatatctgc ttctgcctcc ctgccttcga gggccggaac 240
 tgtgagacgc acaaggatga ccagctgatc tgtgtgaacg agaacggcgg ctgtgagcag 300
 tactgcagtg accacacggg caccaagcgc tcctgtcggt gccacgaggg gtactctctg 360
 10 ctggcagacg ggggtgtcctg cacaccaca gttgaatatc catgtggaaa aatacctatt 420
 ctagaaaaaa gacaggccag caaacccaa ggcgaatttgg tggggggcaa ggtgtgcccc 480
 15 aaaggggagt gtccatggca ggtcctgttg ttggtaatg gagctcagtt gtgtgggggg 540
 accctgatca acaccatctg ggtggcttcc gggcccaact gtttcgacaa aatcaagaac 600
 tggaggaacc tgatcgcgtt gctggcgag cacgacctca gcgagcacga cggggatgag 660
 20 cagagccggc ggggtggcgca ggtcatcatc cccagcacgt acgtcccgcc caccaccaac 720
 cacgacatcg cgctgcctcg cctgcaccag cccgtggtcc tcactgacca tgtggtgccc 780
 ctctgcctgc ccgaacggac gttctctgag aggacgctgg cttcgtgcg cttctcattg 840
 25 gtcagcggct gggccagct gctggaccgt ggcgccacgg ccctggagct catggcctc 900
 aacgtgcccc ggctgtatgac ccaggactgc ctgcagcagt cacgaaagggt gggagactcc 960
 ccacagatca cggagtacat gttctgtgcc ggctactcgg atggcagcaa gactcctgc 1020
 30 aaggggaca gtggaggccc acatgccacc cactaccggg gcacgtggta cttgacgggc 1080
 atcgtcagct gggccaggg ctgcgcaacc gtggccact ttgggtgtta caccagggtc 1140
 tcccagtaca tcgagtggt gcaaaagctc atgcgctcaag agccacgccc aggagtcctc 1200
 35 ctgcgagccc catttccc 1218

<210> 16
 <211> 406
 40 <212> PRT
 <213> Homo sapiens

 <400> 16

45 Ala Asn Ala Phe Leu Glu Glu Leu Arg Pro Gly Ser Leu Glu Arg Glu
 1 5 10 15

50 Cys Lys Glu Glu Gln Cys Ser Phe Glu Glu Ala Arg Glu Ile Phe Lys
 20 25 30

Asp Ala Glu Arg Thr Lys Leu Phe Trp Ile Ser Tyr Ser Asp Gly Asp
 35 40 45

55 Gln Cys Ala Ser Ser Pro Cys Gln Asn Gly Gly Ser Cys Lys Asp Gln
 50 55 60

EP 3 165 232 A1

Leu Gln Ser Tyr Ile Cys Phe Cys Leu Pro Ala Phe Glu Gly Arg Asn
65 70 75 80

5 Cys Glu Thr His Lys Asp Asp Gln Leu Ile Cys Val Asn Glu Asn Gly
85 90 95

10 Gly Cys Glu Gln Tyr Cys Ser Asp His Thr Gly Thr Lys Arg Ser Cys
100 105 110

15 Arg Cys His Glu Gly Tyr Ser Leu Leu Ala Asp Gly Val Ser Cys Thr
115 120 125

20 Pro Thr Val Glu Tyr Pro Cys Gly Lys Ile Pro Ile Leu Glu Lys Arg
130 135 140

25 Asn Ala Ser Lys Pro Gln Gly Arg Ile Val Gly Gly Lys Val Cys Pro
145 150 155 160

30 Lys Gly Glu Cys Pro Trp Gln Val Leu Leu Val Asn Gly Ala Gln
165 170 175

35 Leu Cys Gly Gly Thr Leu Ile Asn Thr Ile Trp Val Val Ser Ala Ala
180 185 190

40 His Cys Phe Asp Lys Ile Lys Asn Trp Arg Asn Leu Ile Ala Val Leu
195 200 205

45 Gly Glu His Asp Leu Ser Glu His Asp Gly Asp Glu Gln Ser Arg Arg
210 215 220

50 Val Ala Gln Val Ile Ile Pro Ser Thr Tyr Val Pro Gly Thr Thr Asn
225 230 235 240

55 His Asp Ile Ala Leu Leu Arg Leu His Gln Pro Val Val Leu Thr Asp
245 250 255

60 His Val Val Pro Leu Cys Leu Pro Glu Arg Thr Phe Ser Glu Arg Thr
260 265 270

65 Leu Ala Phe Val Arg Phe Ser Leu Val Ser Gly Trp Gly Gln Leu Leu
275 280 285

70 Asp Arg Gly Ala Thr Ala Leu Glu Leu Met Val Leu Asn Val Pro Arg
290 295 300

75 Leu Met Thr Gln Asp Cys Leu Gln Gln Ser Arg Lys Val Gly Asp Ser
305 310 315 320

EP 3 165 232 A1

Pro Asn Ile Thr Glu Tyr Met Phe Cys Ala Gly Tyr Ser Asp Gly Ser
325 330 335

5 Lys Asp Ser Cys Lys Gly Asp Ser Gly Gly Pro His Ala Thr His Tyr
340 345 350

10 Arg Gly Thr Trp Tyr Leu Thr Gly Ile Val Ser Trp Gly Gln Gly Cys
355 360 365

15 Ala Thr Val Gly His Phe Gly Val Tyr Thr Arg Val Ser Gln Tyr Ile
370 375 380

20 Glu Trp Leu Gln Lys Leu Met Arg Ser Glu Pro Arg Pro Gly Val Leu
385 390 395 400

25 Leu Arg Ala Pro Phe Pro
405

25 <210> 17

<211> 407

<212> PRT

<213> Artificial Sequence

30 <220>

<223> Variant human FVII peptide sequence

35 <220>

<221> misc_feature

<222> (407)..(407)

<223> Xaa can be any naturally occurring amino acid

40 <400> 17

45 Ala Asn Ala Phe Leu Glu Glu Leu Arg Gln Gly Ser Leu Glu Arg Glu
1 5 10 15

40 Cys Lys Glu Glu Gln Cys Ser Phe Glu Glu Ala Arg Glu Ile Phe Glu
20 25 30

45 Asp Ala Glu Arg Thr Lys Leu Phe Trp Ile Ser Tyr Ser Asp Gly Asp
35 40 45

50 Gln Cys Ala Ser Ser Pro Cys Gln Asn Gly Gly Ser Cys Lys Asp Gln
50 55 60

55 Leu Gln Ser Tyr Ile Cys Phe Cys Leu Pro Ala Phe Glu Gly Arg Asn
65 70 75 80

55 Cys Glu Thr His Lys Asp Asp Gln Leu Ile Cys Val Asn Glu Asn Gly

EP 3 165 232 A1

85

90

95

5 Gly Cys Glu Gln Tyr Cys Ser Asp His Asn Gly Thr Lys Arg Ser Cys
 100 105 110

10 Arg Cys His Glu Gly Tyr Ser Leu Leu Ala Asp Gly Val Ser Cys Thr
 115 120 125

15 Pro Thr Val Glu Tyr Pro Cys Gly Lys Ile Pro Ile Leu Glu Lys Arg
 130 135 140

20 Asn Ala Ser Lys Pro Gln Gly Arg Ile Val Gly Gly Lys Val Cys Pro
 145 150 155 160

25 Lys Gly Glu Cys Pro Trp Gln Val Leu Leu Val Asn Gly Ala Gln
 165 170 175

30 Leu Cys Gly Gly Thr Leu Ile Asn Thr Ile Trp Val Val Ser Ala Ala
 180 185 190

35 His Cys Phe Asp Lys Ile Lys Asn Trp Arg Asn Leu Ile Ala Val Leu
 195 200 205

40 Gly Glu His Asp Leu Ser Glu His Asp Gly Asp Glu Gln Ser Arg Arg
 210 215 220

45 Val Ala Gln Val Ile Ile Pro Ser Thr Tyr Val Pro Gly Thr Thr Asn
 225 230 235 240

50 His Asp Ile Ala Leu Leu Arg Leu His Gln Pro Val Asn Leu Thr Asp
 245 250 255

55 His Val Val Pro Leu Cys Leu Pro Glu Arg Thr Phe Ser Glu Arg Thr
 260 265 270

Leu Ala Phe Val Arg Phe Ser Leu Val Ser Gly Trp Gly Gln Leu Leu
 275 280 285

60 Asp Arg Gly Ala Thr Ala Leu Glu Leu Met Val Leu Asn Val Pro Arg
 290 295 300

65 Leu Met Thr Gln Asp Cys Leu Gln Gln Ser Arg Lys Val Gly Asp Ser
 305 310 315 320

70 Pro Asn Ile Thr Glu Tyr Met Phe Cys Ala Gly Tyr Ser Asp Gly Ser
 325 330 335

EP 3 165 232 A1

Lys Asp Ser Cys Lys Gly Asp Ser Gly Gly Pro His Ala Thr His Tyr
340 345 350

5 Arg Gly Thr Trp Tyr Leu Thr Gly Ile Val Ser Trp Gly Gln Gly Cys
355 360 365

10 Ala Thr Val Gly His Phe Gly Val Tyr Thr Arg Val Ser Gln Tyr Ile
370 375 380

Glu Trp Leu Gln Lys Leu Met Arg Ser Glu Pro Arg Pro Gly Val Leu
385 390 395 400

15 Leu Arg Ala Pro Phe Pro Xaa
405

20 <210> 18
<211> 406
<212> PRT
<213> Artificial Sequence

25 <220>
<223> Variant human FVII peptide sequence
<400> 18

30 Ala Asn Ala Phe Leu Glu Glu Leu Arg Gln Gly Ser Leu Glu Arg Glu
1 5 10 15

35 Cys Lys Glu Glu Gln Cys Ser Phe Glu Glu Ala Arg Glu Ile Phe Glu
20 25 30

40 Asp Glu Glu Glu Thr Lys Leu Phe Trp Ile Ser Tyr Ser Asp Gly Asp
35 40 45

45 Gln Cys Ala Ser Ser Pro Cys Gln Asn Gly Gly Ser Cys Lys Asp Gln
50 55 60

55 Leu Gln Ser Tyr Ile Cys Phe Cys Leu Pro Ala Phe Glu Gly Arg Asn
65 70 75 80

60 Cys Glu Thr His Lys Asp Asp Gln Leu Ile Cys Val Asn Glu Asn Gly
85 90 95

65 Gly Cys Glu Gln Tyr Cys Ser Asp His Asn Gly Thr Lys Arg Ser Cys
100 105 110

70 Arg Cys His Glu Gly Tyr Ser Leu Leu Ala Asp Gly Val Ser Cys Thr
115 120 125

75 Pro Thr Val Glu Tyr Pro Cys Gly Lys Ile Pro Ile Leu Glu Lys Arg

EP 3 165 232 A1

130 135 140

5 Asn Ala Ser Lys Pro Gln Gly Arg Ile Val Gly Gly Lys Val Cys Pro
145 150 155 16010 Lys Gly Glu Cys Pro Trp Gln Val Leu Leu Leu Val Asn Gly Ala Gln
165 170 175Leu Cys Gly Gly Thr Leu Ile Asn Thr Ile Trp Val Val Ser Ala Ala
180 185 19015 His Cys Phe Asp Lys Ile Lys Asn Trp Arg Asn Leu Ile Ala Val Leu
195 200 20520 Gly Glu His Asp Leu Ser Glu His Asp Gly Asp Glu Gln Ser Arg Arg
210 215 220Val Ala Gln Val Ile Ile Pro Ser Thr Tyr Val Pro Gly Thr Thr Asn
225 230 235 24025 His Asp Ile Ala Leu Leu Arg Leu His Gln Pro Val Asn Leu Thr Asp
245 250 25530 His Val Val Pro Leu Cys Leu Pro Glu Arg Thr Phe Ser Glu Arg Thr
260 265 27035 Leu Ala Phe Val Arg Phe Ser Leu Val Ser Gly Trp Gly Gln Leu Leu
275 280 285Asp Arg Gly Ala Thr Ala Leu Glu Leu Met Val Leu Asn Val Pro Arg
290 295 30040 Leu Met Thr Gln Asp Cys Leu Gln Gln Ser Arg Lys Val Gly Asp Ser
305 310 315 32045 Pro Asn Ile Thr Glu Tyr Met Phe Cys Ala Gly Tyr Ser Asp Gly Ser
325 330 33550 Lys Asp Ser Cys Lys Gly Asp Ser Gly Gly Pro His Ala Thr His Tyr
340 345 350Arg Gly Thr Trp Tyr Leu Thr Gly Ile Val Ser Trp Gly Gln Gly Cys
355 360 36555 Ala Thr Val Gly His Phe Gly Val Tyr Thr Arg Val Ser Gln Tyr Ile
370 375 380

Glu Trp Leu Gln Lys Leu Met Arg Ser Glu Pro Arg Pro Gly Val Leu
 385 390 395 400

Leu Arg Ala Pro Phe Pro
405

10 **Claims**

1. An isolated variant Factor VII polypeptide comprising an amino acid sequence having at least one sequence alteration relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least one sequence alteration is selected from the group consisting of P10Q, K32E, R36E, A34E, T106N, and V253N, wherein the polypeptide has a ratio of moles of conjugated sialic acid to moles of N-linked glycan between 0 and 5.0.
2. The isolated variant Factor VII polypeptide of claim 1, wherein the variant Factor VII polypeptide comprises two or more sequence alterations selected from the group consisting of P10Q, K32E, R36E, A34E, T106N, and V253N.
3. The isolated variant Factor VII polypeptide of claim 1, wherein the variant Factor VII polypeptide comprises three or more sequence alterations selected from the group consisting of P10Q, K32E, R36E, A34E, T106N, and V253N.
4. The isolated variant Factor VII polypeptide of claim 1, wherein the variant Factor VII polypeptide comprises four or more sequence alterations selected from the group consisting of P10Q, K32E, R36E, A34E, T106N, and V253N.
5. The isolated variant Factor VII polypeptide of claim 1, wherein the variant Factor VII polypeptide comprises the sequence alterations of P10Q, K32E, R36E, A34E, T106N, and V253N.
6. The isolated variant Factor VII polypeptide of claim 1, wherein the variant Factor VII polypeptide comprises the sequence alterations P10Q, K32E, T106N, and V253N.
7. The isolated variant Factor VII polypeptide of claim 1, wherein the amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO: 16 having the sequence alterations P10Q, K32E, R36E, A34E, T106N, and V253N.
8. The isolated variant Factor VII polypeptide of claim 1, wherein the amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO: 16 having the sequence alterations P10Q, K32E, T106N, and V253N.
9. The isolated variant Factor VII polypeptide of any of claims 1-8, wherein the ratio of moles of conjugated sialic acid to moles of N-linked glycan is less than 0.1.
10. A method of preparing the isolated variant Factor VII polypeptide of claim 1, the method comprising
 - (1) obtaining a sialylated Factor VII polypeptide comprising an amino acid sequence having at least one sequence alteration relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least one sequence alteration is selected from the group consisting of P10Q, K32E, R36E, A34E, T106N, and V253N;
 - (2) contacting the sialylated Factor VII polypeptide with sialidase under conditions such that sufficient amounts of covalently attached sialic acid residues are removed from the sialylated Factor VII polypeptide to produce a desialylated Factor VII polypeptide having a ratio of moles of conjugated sialic acid to moles of N-linked glycan between 0 and 5.0 and
 - (3) isolating the variant Factor VII polypeptide thereby produced.
11. A method of preparing the isolated variant Factor VII polypeptide of claim 1, said method comprising
 - (1) producing a Factor VII polypeptide comprising an amino acid sequence having at least one sequence alteration relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least one sequence alteration is selected from the group consisting of P10Q, K32E, R36E, A34E, T106N, and V253N, in a recombinant cell line that is deficient in its ability to sialylate peptides such that it produces a desialylated Factor VII polypeptide

having a ratio of moles of conjugated sialic acid to moles of N-linked glycan between 0 and 5.0; and
 (2) isolating the variant Factor VII polypeptide thereby produced.

12. A method of preparing the isolated variant Factor VII polypeptide of claim 1, said method comprising

5 (1) obtaining a recombinant cell line that coexpresses (a) a recombinant Factor VII polypeptide comprising an amino acid sequence having at least one sequence alteration relative to the amino acid sequence of SEQ ID NO: 16, wherein the at least one sequence alteration is selected from the group consisting of P10Q, K32E, R36E, A34E, T106N, and V253N, and (b) a recombinant sialidase enzyme;

10 (2) culturing said recombinant cell line to allow expression of both the recombinant Factor VII polypeptide and the recombinant sialidase enzyme, wherein said recombinant sialidase enzyme removes sufficient amounts of covalently attached sialic acid residues to produce a desialylated Factor VII polypeptide having a ratio of moles of conjugated sialic acid to moles of N-linked glycan between 0 and 5.0; and

15 (3) isolating the variant Factor VII polypeptide thereby produced.

13. An isolated variant Factor VII polypeptide selected from the group consisting of:

20 (1) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, and (3) a sequence alteration such that N-linked glycosylation at position 145 is disrupted;

25 (2) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, and (3) a sequence alteration such that N-linked glycosylation at position 322 is disrupted;

30 (3) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, and (2) a glutamic acid residue substituted for the lysine residue in position 32, and (3) sequence alterations such that N-linked glycosylation at positions 145 and 322 is disrupted;

35 (4) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, and (2) a glutamic acid residue substituted for the lysine residue in position 32, wherein positions 145 and 322 are asparagine and have attached N-linked glycosylation;

40 (5) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the alanine residue in position 34, (4) a glutamic acid residue substituted for the arginine residue in position 36, and (5) a sequence alteration such that N-linked glycosylation at position 145 is disrupted;

45 (6) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the alanine residue in position 34, (4) a glutamic acid residue substituted for the arginine residue in position 36, and (5) a sequence alteration such that N-linked glycosylation at position 322 is disrupted;

50 (7) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the alanine residue in position 34, (4) a glutamic acid residue substituted for the arginine residue in position 36, and (5) a sequence alterations such that N-linked glycosylation at positions 145 and 322 is disrupted; and

55 (8) a polypeptide comprising a Factor VII amino acid sequence having sequence alterations relative to the amino acid sequence of SEQ ID NO: 16, wherein the sequence alterations consist of (1) a glutamine residue substituted for the proline residue in position 10, (2) a glutamic acid residue substituted for the lysine residue in position 32, (3) a glutamic acid residue substituted for the alanine residue in position 34, and (4) a glutamic acid residue substituted for the arginine residue in position 36, wherein positions 145 and 322 are asparagine and have attached N-linked glycosylation.

14. The isolated variant Factor VII polypeptide of any of claims 1-9 or 13 for use in the treatment of a disease or a disorder wherein blood clot formation is desirable, wherein the disease or disorder is selected from the group consisting of a hemorrhage, gastrointestinal bleeding, uncontrolled bleeding, bleeding in a mammal undergoing transplantation or resection or surgery, variceal bleeding, thrombocytopenia, hemophilia, intracranial hemorrhage, aortic aneurysm, and over administration of an anticoagulant.

5

15. A pharmaceutical composition comprising the isolated variant Factor VII polypeptide of any of claims 1-9 or 13 and a pharmaceutically acceptable excipient.

10

15

20

25

30

35

40

45

50

55

WILD TYPE FACTOR VII

GCCAACCGCTTCCTGGAGGAGCTCGGGCAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTTCAAGGACCGGGAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGCTGTGAG
 CAGTACTGCAGTGACCACACGGCACCAAGCGCTCTGCGGTGCCACGAGGGTACTCT
 CTGCTGGCAGACGGGTGTCTGCACACCCACAGTGAAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAACCCCAAGGCCGAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGAGTGTCCATGGCAGGTCTGTTGTTGGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTATCATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACCGACATCGCGCTGCTCCGCTGCACCAGCCGTGGTCTACTGACCATGTG
 GTGCCCCCTGCTGCCGAACGGACGTTCTGAGAGGGACGCTGGCCTTCGTGCGCTTC
 TCATTGGTCAGCGGCTGGGCCAGCTGCTGGACCGTGGCGCCACGGGCTGGAGCTCATG
 GTCCTCAACGTGCCCCGGCTGATGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGG
 AGACTCCCCAAATATCAGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGACGTGGTACC
 TGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 ACCAGGTCTCCAGTACATCGAGTGGCTGCAAAAGCTATGCGCTCAGAGGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 1)

V1

GCCAACCGCTTCCTGGAGGAGCTCGGGCAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTTCAAGGACCGGGAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGCTGTGAG
 CAGTACTGCAGTGACCACAACGGCACCAAGCGCTCTGCGGTGCCACGAGGGTACTCT
 CTGCTGGCAGACGGGTGTCTGCACACCCACAGTGAAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAACCCCAAGGCCGAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGAGTGTCCATGGCAGGTCTGTTGTTGGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTATCATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACCGACATCGCGCTGCTCCGCTGCACCAGCCGTGAACCTCACTGACCATGTG
 GTGCCCCCTGCTGCCGAACGGACGTTCTGAGAGGGACGCTGGCCTTCGTGCGCTTC
 TCATTGGTCAGCGGCTGGGCCAGCTGCTGGACCGTGGCGCCACGGGCTGGAGCTCATG
 GTCCTCAACGTGCCCCGGCTGATGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGG
 AGACTCCCCAAATATCAGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGACGTGGTACC
 TGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 ACCAGGTCTCCAGTACATCGAGTGGCTGCAAAAGCTATGCGCTCAGAGGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCCCT

Figure 1A

V2

GCCAACCGCTTCTGGAGGAGCTCGGCCAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTCGAAGACGAAGAGGAAACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGTGTGAG
 CAGTACTGCAGTGACCACAACGGCACCAAGCGCTCTGTCGGTGCACAGTTGAATATCCATGTGGAAAAATACCT
 CTGCTGGCAGACGGGTGTCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAACCCCAAGGCCAATTGTGGGGGGCAAGGTGTG
 CCCCAAAGGGAGTGTCCATGGCAGGTCTGTTGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCACTCATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACCGACATCGCGCTGCTCCGCTGCACCCAGGCCGTGAACCTCACTGACCATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCGTGGCCTCGTGCCTTC
 TCATTGGTCAGCGCTGGGGCCAGCTGCTGGACCGTGGCGCCACGGCCCTGGAGGCTCATG
 GTCCTCAACGTGCCCGGCTGATGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGGG
 AGACTCCCCAAATATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTACC
 TGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 ACCAGGGTCTCCAGTACATCGAGTGGCTGCAAAAGCTATGCGCTCAGAGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 3)

pMB113

GCCAACCGCTTCTGGAGGAGCTCGGCCAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTCGAAGACGCAGAGGAGAACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGTGTGAG
 CAGTACTGCAGTGACCACACGGCACCAAGCGCTCTGTCGGTGCACAGGGTACTCT
 CTGCTGGCAGACGGGTGTCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAACCCCAAGGCCAATTGTGGGGGGCAAGGTGTG
 CCCCAAAGGGAGTGTCCATGGCAGGTCTGTTGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCACTCATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACCGACATCGCGCTGCTCCGCTGCACCCAGGCCGTGGTCTCGTGCCTTC
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCGTGGCCTCGTGCCTTC
 TCATTGGTCAGCGCTGGGCCAGCTGCTGGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGGG
 AGACTCCCCAAATATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTACC
 TGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 ACCAGGGTCTCCAGTACATCGAGTGGCTGCAAAAGCTATGCGCTCAGAGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 4)

Figure 1B

pMB114

GCCAACCGCGTCTGGAGGAGCTCGGGCCGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTTCAAGGACCGGGAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGAACGAGAACGGCGGCTGTGAG
 CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCACAGTTGAATATCCATGTGGAAAAATACCT
 CTGCTGGCAGACGGGTGCTCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGACAGGCCAGCAAACCCCAAGGCCGAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCAATCATCCCCAGCACGTACGTCCCCGGCACC
 ACCAACCAACGACATCGCGCTGCTCCGCCTGCACCCAGGCCGTGGTCCACTGACCATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCCTGGCCTTGTGCGCTTC
 TCATTGGTCAGCGGCTGGGCCAGCTGCTGGACCGTGGCCTGGCAGGCCACTGGGCTTGTGCGCTTC
 GTCCTCAACGTGCCCCGGCTGATGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGGG
 AGACTCCCCAAATATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTACC
 TGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 ACCAGGGTCTCCAGTACATCGAGTGGCTGCAAAAGCTATGCGCTCAGAGGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 5)

pMB115

GCCAACCGCGTCTGGAGGAGCTCGGGCCGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTTCAAGGACCGGGAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGAACGAGAACGGCGGCTGTGAG
 CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCACAGGCCACTGTGGAAAAATACCT
 CTGCTGGCAGACGGGTGCTCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAAACCCCAAGGCCGAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCAATCCCCAGCACGTACGTCCCCGGCACC
 ACCAACCAACGACATCGCGCTGCTCCGCCTGCACCCAGGCCGTGGTCCACTGACCATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCCTGGCCTTGTGCGCTTC
 TCATTGGTCAGCGGCTGGGCCAGCTGCTGGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGGG
 AGACTCCCCACAGATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGG
 ACTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTAC
 CTGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 CACCAGGGTCTCCAGTACATCGAGTGGCTGCAAAAGCTATGCGCTCAGAGGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 6)

Figure 1C

pMB116

GCCAACCGCGTCTGGAGGAGCTCGGCCGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCCTCTCGAGGAGGCCGGAGATCTCAAGGACGCCAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGTGTGAG
 CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCACAGTTGAATATCCATGTGGAAAAATACCT
 CTGCTGGCAGACGGGTGCTGACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGACAGGCCAGCAAACCCCAAGGCCAATTGTGGGGGCAAGGTGTG
 CCCCAAAGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCACTCATCCCCAGCACGTACGTCCCCGGCACC
 ACCAACACGACATCGCGCTGCTCCGCTGCACCCAGCCGTGGTCTCACTGACCATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCCTGGCCTTCGTGCGCTTC
 TCATTGGTCAGCGCTGGGGCCAGGGCTGCGAACCGTGGGCCACTTGGGTGTAC
 CTGACGGGCATCGTCAGCTGGGGCCAGGGCTGCGAACCGTGGGCCACTTGGGTGTAC
 CACCAAGGGTCTCCAGTACATCGAGTGGCTGCAAAAGCTCATGCGCTCAGAGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 7)

pMB117

GCCAACCGCGTCTGGAGGAGCTCGGCCAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCCTCTCGAGGAGGCCGGAGATCTCGAAGACGCCAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGTGTGAG
 CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCACAGGGTACTCT
 CTGCTGGCAGACGGGTGCTGACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAAACCCCAAGGCCAATTGTGGGGGCAAGGTGTG
 CCCCAAAGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCACTCATCCCCAGCACGTACGTCCCCGGCACC
 ACCAACACGACATCGCGCTGCTCCGCTGCACCCAGCCGTGGTCTCACTGACCATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCCTGGCCTTCGTGCGCTTC
 TCATTGGTCAGCGCTGGGGCCAGCTGCTGGACCGTGGCGCCACGGGCCCTGGAGCTCATG
 GTCCCTCAACGTGCCCCGGCTGATGACCCAGGACTGCTGCAGCAGTCACGGAAAGGTGG
 AGACTCCCCAAATATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTACC
 TGACGGGCATCGTCAGCTGGGGCCAGGGCTGCGAACCGTGGGCCACTTGGGTGTAC
 ACCAAGGGTCTCCAGTACATCGAGTGGCTGCAAAAGCTCATGCGCTCAGAGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 8)

Figure 1D

pMB118

GCCAACCGCGTCTGGAGGAGCTCGGGAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTCGAAGACGCCAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGCCGG
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGCCGGCTGTGAG
 CAGTACTGCAGTGACCACACGGCACCAAGCGCTCTGTCGGTGCACAGTTGAATATCCATGTGGAAAAATACCT
 CTGCTGGCAGACGGGTGTCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGACAGGCCAGCAAACCCAAAGGCCAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCGTATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCAATCATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACACGACATCGCGCTGCTCCGCTGCACCAAGGCCGTGGTCTCACTGACCATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCGTGGCCTCGTGCCTTC
 TCATTGGTCAGCGCTGGGGCCAGCTGCTGGCGCAGGTCAATCATCCCCAGCACGTCCCGGGC
 GTCCTCAACGTGCCCGGCTGATGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGGG
 AGACTCCCCAATATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTACC
 TGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 ACCAGGGTCTCCAGTACATCGAGTGGCTGCAAAGCTATGCGCTCAGAGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 9)

pMB119

GCCAACCGCGTCTGGAGGAGCTCGGGAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTCGAAGACGCCAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGCCGG
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGCCGGCTGTGAG
 CAGTACTGCAGTGACCACACGGCACCAAGCGCTCTGTCGGTGCACAGGCCAGAGGGTACTCT
 CTGCTGGCAGACGGGTGTCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAAACCCAAAGGCCAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCGTATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCAATCATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACACGACATCGCGCTGCTCCGCTGCACCAAGGCCGTGGTCTCACTGACCATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCGTGGCCTCGTGCCTTC
 TCATTGGTCAGCGCTGGGCCAGCTGCTGGACCGTGGCGCCACGGCCCTGGAGCTCATG
 GTCCTCAACGTGCCCGGCTGATGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGGG
 AGACTCCCCACAGATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGG
 ACTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTAC
 CTGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 CACCAGGGTCTCCAGTACATCGAGTGGCTGCAAAGCTATGCGCTCAGAGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 10)

Figure 1E

pMB120

GCCAACCGCGTCTGGAGGAGCTCGGGAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTCGAAGACCGGGAGAGGACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGTGTGAG
 CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCACAGTTGAATATCCATGTGGAAAAATACCT
 CTGCTGGCAGACGGGTGTCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGACAGGCCAGCAAACCCAAAGGCCAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGAGTGTCCATGGCAGGTCTGTTGTGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCACTCATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACCGACATCGCGCTGCTCCGCTGCACCCAGCCGTGGTCTCAGTACCGACATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCGCTGGCCTTCGTGCGCTTC
 TCATTGGTCAGCGGCTGGGCCAGCTGCTGGCGAGGTCACTCATCCCCAGCACGTACGTCCCGGGCACC
 GTCCTCAACGTGCCCGGCTGATGACCCAGGACTGCCTGCAGCAGTCACGGAAAGGTGGG
 AGACTCCCCACAGATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGG
 ACTCCTGCAAGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGACGTGGTAC
 CTGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGGCCACTTGGGTGTA
 CACCAAGGGCTCCCACTACATCGAGTGGCTGCAAAAGCTCATGCGCTCAGAGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 11)

pMB121

GCCAACCGCGTCTGGAGGAGCTCGGGAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
 GCAGTGCTCCTCGAGGAGGCCGGAGATCTCGAAGACGAAGAGGAAACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGTGTGAG
 CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCACAGGGTACTCT
 CTGCTGGCAGACGGGTGTCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAAACCCAAAGGCCAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGAGTGTCCATGGCAGGTCTGTTGTGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTCACTCATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACCGACATCGCGCTGCTCCGCTGCACCCAGCCGTGGTCTCAGTACCGACATGTG
 GTGCCCTCTGCCTGCCGAACGGACGTTCTGAGAGGAGCGCTGGCCTTCGTGCGCTTC
 TCATTGGTCAGCGGCTGGGCCAGCTGCTGGACCCAGGACTGCCTGCAGCAGTCACGGAAAGGTGGG
 AGACTCCCCAAATATCACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGACGTGGTACCC
 TGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGGCCACTTGGGTGTA
 ACCAAGGGCTCCCACTACATCGAGTGGCTGCAAAAGCTCATGCGCTCAGAGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 12)

Figure 1F

pMB122

GCCAACCGCGTCTGGAGGAGCTCGGGAGGGCTCCCTGGAGAGGGAGTGCAGGAGGA
 GCAGTGCTCCTTCGAGGAGGCCGGAGATCTCGAAGACGAAGAGGAAACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGTGTGAG
 CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCACAGGGTACTCT
 CTGCTGGCAGACGGGTGCTGACACCCACAGTGAAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGACAGGCCAGCAAACCCCAAGGCCGAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGCTCCCGGGCCACTGTTGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTACATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACCAACGACATCGCGCTGCTCCGCTGCACCCAGCCGTGGCTACTGACCATGTG
 GTGCCCCCTGCTGCCGAACGGACGTTCTGAGAGGGACGCTGGCCTTGTGCGCTTC
 TCATTGGTCAGCGGCTGGGCCAGCTGCTGGCGCAGGTACATCCCCAGCACGTACGTCCCGGG
 GTCCTCAACGTGCCCCGGCTGATGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGGG
 AGACTCCCCAAATATCACGGAGTACATGTTCTGCGCGCTACTCGGATGGCAGCAAGGA
 CTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTACC
 TGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 ACCAGGGTCTCCAGTACATCGAGTGGCTGCAAAGCTATGCGCTCAGAGGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 13)

pMB123

GCCAACCGCGTCTGGAGGAGCTCGGGAGGGCTCCCTGGAGAGGGAGTGCAGGAGGA
 GCAGTGCTCCTTCGAGGAGGCCGGAGATCTCGAAGACGAAGAGGAAACGAAGCTGT
 TCTGGATTCTTACAGTGATGGGGACCAGTGTGCCTCAAGTCCATGCCAGAATGGGGCT
 CCTGCAAGGACCAGCTCCAGTCTATATCTGCTTCTGCCTCCCTGCCTCGAGGGCCGGA
 ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGTGTGAG
 CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCACAGGGTACTCT
 CTGCTGGCAGACGGGTGCTGACACCCACAGTGAAATATCCATGTGGAAAAATACCT
 ATTCTAGAAAAAAAGAAATGCCAGCAAACCCCAAGGCCGAATTGTGGGGGCAAGGTGTG
 CCCAAAGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
 GGGGACCCCTGATCAACACCATCTGGGTGCTCCCGGGCCACTGTTGACAAAATCAA
 GAACTGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
 ATGAGCAGAGCCGGGGTGGCGCAGGTACATCCCCAGCACGTACGTCCCGGGCACC
 ACCAACCAACGACATCGCGCTGCTCCGCTGCACCCAGCCGTGGCTACTGACCATGTG
 GTGCCCCCTGCTGCCGAACGGACGTTCTGAGAGGGACGCTGGCCTTGTGCGCTTC
 TCATTGGTCAGCGGCTGGGCCAGCTGCTGGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGGG
 AGACTCCCCACAGATCACGGAGTACATGTTCTGCGCGCTACTCGGATGGCAGCAAGG
 ACTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGGCACGTGGTAC
 CTGACGGGCATCGTCAGCTGGGCCAGGGCTGCGCAACCGTGGCCACTTGGGTGTAC
 CACCAGGGTCTCCAGTACATCGAGTGGCTGCAAAGCTATGCGCTCAGAGGCCACGCC
 AGGAGTCCTCCTGCGAGCCCCATTCCC (SEQ ID NO: 14)

Figure 1G

pMB124

GCCAAACGCGTTCTGGAGGAGCTCGGGAGGGCTCCCTGGAGAGGGAGTGCAAGGAGGA
GCAGTGCTCCTCGAGGAGGCCGGAGATCTTCGAAGACGAAGAGGAAACGAAGCTGT
TCTGGATTTCTTACAGTGATGGGACCAAGTGTGCCCTCAAGTCCATGCCAGAATGGGGCT
CCTGCAAGGACCAGCTCCAGTCCATATCTGCTTCTGCCCTGCCCTCGAGGGCCGGA
ACTGTGAGACGCACAAGGATGACCAGCTGATCTGTGTGAACGAGAACGGCGGCTGTGAG
CAGTACTGCAGTGACCACACGGGACCAAGCGCTCTGTCGGTGCCACGAGGGTACTCT
CTGCTGGCAGACGGGTGTCCTGCACACCCACAGTTGAATATCCATGTGGAAAAATACCT
ATTCTAGAAAAAAAGACAGGCCAGCAAACCCCAAGGCCGAATTGTGGGGGCAAGGTGTG
CCCCAAAGGGGAGTGTCCATGGCAGGTCTGTTGGTGAATGGAGCTCAGTTGTGTGG
GGGGACCCCTGATCAACACCATCTGGGTGGTCTCCCGGGCCACTGTTCGACAAAATCAA
GAACGGAGGAACCTGATCGCGGTGCTGGCGAGCACGACCTCAGCGAGCACGACGGGG
ATGAGCAGAGCCGGGGTGGCGCAGGTCATCATCCCCAGCACGTACGTCCCCGGCACC
ACCAACCACGACATCGCGCTGCTCCGCTGCACCAAGCCGTGGCCTCCTGACCATGTG
GTGCCCTCTGCCCTGCCGAACGGACGTTCTGAGAGGAGCGCTGCCCTCGTGGCTTC
TCATTGGTCAGCGCTGGGCCAGCTGCTGGACCCAGGACTGCCTGCAGCAGTCACGGAAGGTGG
AGACTCCCCACAGATACGGAGTACATGTTCTGTGCCGGCTACTCGGATGGCAGCAAGG
ACTCCTGCAAGGGGGACAGTGGAGGCCACATGCCACCCACTACCGGGCACGTGGTAC
CTGACGGGCATCGTCAGCTGGGCCAGGGCTGCCAACCGTGGGCCACTTGGGTGTA
CACCAAGGGCTCCAGTACATCGAGTGGCTGCAAAAGCTCATGCGCTCAGAGGCCACGCC
AGGAGTCCTCCTGCGAGCCCCATTCCCC (SEQ ID NO: 15)

Figure 1H

Wild Type Factor VII Peptide

ANAFLEELRPGSLERECKEEQCSFEEAREIFKDAERTKLFWISYSDGDQCASSPCQNGGS
CKDQLQSYICFCLPAFEGRNCETHKDDQLICVNENGGCEQYCSDDHTGKRSRRCHEGYSL
LADGVSCPTVEYPCGKIPILEKRNASKPQGRIVGGKVCVKGECPWQVLLVNGAQLCGG
TLINTIWVVAACFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTN
HDIAALLRLHQPVVLTDHVVPLCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVL
NVPRLMQTQDCLQQSRKVGDSPNITEYMFCAGYSDGSKDSCKGDSGGPHATHYRGTWYLTG
IVSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFP (SEQ ID NO: 16)

V1 Peptide

ANAFLEELRQGSLEERECKEEQCSFEEAREIFEDAERTKLFWISYSDGDQCASSPCQNGGS
CKDQLQSYICFCLPAFEGRNCETHKDDQLICVNENGGCEQYCSDDHNGKRSRRCHEGYSL
LADGVSCPTVEYPCGKIPILEKRNASKPQGRIVGGKVCVKGECPWQVLLVNGAQLCGG
TLINTIWVVAACFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTN
HDIAALLRLHQPVNLTDHVVPCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVL
NVPRLMQTQDCLQQSRKVGDSPNITEYMFCAGYSDGSKDSCKGDSGGPHATHYRGTWYLTG
IVSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFP (SEQ ID NO: 17)

V2 Peptide

ANAFLEELRQGSLEERECKEEQCSFEEAREIFEDEEETKLFWISYSDGDQCASSPCQNGGS
CKDQLQSYICFCLPAFEGRNCETHKDDQLICVNENGGCEQYCSDDHNGKRSRRCHEGYSL
LADGVSCPTVEYPCGKIPILEKRNASKPQGRIVGGKVCVKGECPWQVLLVNGAQLCGG
TLINTIWVVAACFDKIKNWRNLIAVLGEHDLSEHDGDEQSRRVAQVIIPSTYVPGTTN
HDIAALLRLHQPVNLTDHVVPCLPERTFSERTLAFVRFSLVSGWGQLLDRGATALELMVL
NVPRLMQTQDCLQQSRKVGDSPNITEYMFCAGYSDGSKDSCKGDSGGPHATHYRGTWYLTG
IVSWGQGCATVGHFGVYTRVSQYIEWLQKLMRSEPRPGVLLRAPFP (SEQ ID NO: 18)

Figure 2





Figure 3






Figure 5

Figure 6

pMB number	Backbone	N-glycosylation sites
pMB113		N145, N322
pMB114	F7wt	N145Q, N322
pMB115		N145, N322Q
pMB116		N145Q, N322Q
pMB117	V1	N145, N322
pMB118		N145Q, N322Q
pMB119		N145Q, N322Q
pMB120		N145, N322
pMB121		N145Q, N322
pMB122	V2	N145, N322Q
pMB123		N145Q, N322Q
pMB124		N145Q, N322Q

Human FVII-B0189, heavy chain

Reduced vs reduced/desialylated




Figure 7




Figure 8





Figure 9

In vitro activity comparisons of rFVIIa variants
(PL-TGA assay in severe hemA patient plasma)




Amount of thrombin generated by 1300 ng/ml of various rFVIIa proteins

	V1 no treatment	V1 treatment	V2 no treatment	V2 treatment	FVII
Peak Thrombin	463 nM	492 nM	480 nM	502 nM	178 nM
Fold Over FVII	2.60	2.76	2.70	2.82	1

Figure 10

Figure 11



	FVII Chromo (ng/ml)	FVII ELISA (ng/ml)	Chromo/ELISA ratio
pMB113	740.4	788.2	0.9
pMB114	619.8	601.0	1.0
pMB115	554.2	426.8	1.3
pMB116	498.9	430.0	1.2
pMB117	1248.3	870.8	1.4
pMB118	1272.1	890.5	1.4
pMB119	601.7	346.8	1.7
pMB120	292	236.8	1.2
pMB121	166.3	808.2	0.2
pMB122	155.6	711.9	0.2
pMB123	181.5	514.5	0.4
pMB124	78.9	277.2	0.3

Figure 12

**In vitro activity comparisons of rFVIIa variants
(PL-TGA assay in severe hemA patient plasma)**





Figure 13

Figure 14






Figure 15






Figure 16

Figure 17

A) ELISA Assay





B) soluble TF PTT clotting assay



	$T_{1/2}$ (h)	MRT (h)	CL (ml/h/kg)	V_{ss} (ml/kg)
rFVII	2.6	1.9	79	152
dV2	~3 min	n.c.	n.c.	n.c.

Figure 18

**B)**




Figure 19




Figure 20




Figure 21




Figure 22

Tail Cut Study

1. 5 mg/kg tPA iv
2. Clip tail at 50 mm from tip
- 3 dWT VIIa or Novo7 by IV
4. Disrupt unstable clots every 6 min through the collection period
5. Collect blood for 45 minutes




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27



EUROPEAN SEARCH REPORT

Application Number

EP 16 19 6780

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10 X	WO 2004/111242 A1 (MAXYGEN HOLDINGS LTD; MAXYGEN APS [DK]; HAANING JESPER MORTENSEN [DK];) 23 December 2004 (2004-12-23) * Variants of Factor VII/VIIa; page 14, line 24 - page 17, line 23; claims 1, 31; sequence SEQ ID NO: 1 * * Coupling to a sugar moiety; page 31, line 4 - page 31, line 15 * * Preparation of variants; page 33, line 16 - page 34, line 9 * * Pharmaceutical compositions and use; page 34, line 11 - page 37, line 25 * -----	1-9,11, 13-15	INV. A61K38/00 A61K38/48 C12N9/64
15 X	WO 2004/083361 A2 (BAYER HEALTHCARE LLC) 30 September 2004 (2004-09-30) * Glycosylation site insertions.; page 20, line 12 - page 22, line 30; sequence SEQ ID NO: 2 * * glycosylation and polypeptide variant production; page 27, line 28 - page 30, line 20 * * Pharmaceutical composition of the invention and use.; page 30, line 22 - page 34, line 3 *	1,2,9, 11,14,15	
20 A	----- page 20, line 12 - page 22, line 30; sequence SEQ ID NO: 2 * * glycosylation and polypeptide variant production; page 27, line 28 - page 30, line 20 * * Pharmaceutical composition of the invention and use.; page 30, line 22 - page 34, line 3 *	3-8,10, 12,13	
25			TECHNICAL FIELDS SEARCHED (IPC)
30 X	----- US 2009/291890 A1 (MADISON EDWIN L [US] ET AL) 26 November 2009 (2009-11-26) * paragraph [[0328]] - paragraph [[370]] * * paragraph [[0387]] - paragraph [[0388]] * * paragraph [[0282]] - paragraph [[0286]] *	1-15	C12N A61K
35	-----		
40	-----	-/-	
45			
50 1	The present search report has been drawn up for all claims		
55	Place of search Munich	Date of completion of the search 18 January 2017	Examiner Bretherick, James
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



EUROPEAN SEARCH REPORT

Application Number

EP 16 19 6780

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10 X	WO 2005/123916 A2 (NOVO NORDISK HEALTHCARE AG [CH]; BOLT GERT [DK]; STEENSTRUP THOMAS DOC) 29 December 2005 (2005-12-29) * Glycosylation disrupted Factor VII variants; page 4, line 1 - page 4, line 16; examples 2-4; sequence SEQ ID NO: 1 * * Glycosylation site insertion; page 6, line 35 - page 6, line 35; sequence SEQ ID NO: 1 * -----	1-15	
15 A	Annette Hrzscovics ' ET AL: "Glycoprotein biosynthesis in yeast", , 1 April 1993 (1993-04-01), XP055336075, Retrieved from the Internet: URL: http://www.fasebj.org/content/7/6/540.full.pdf [retrieved on 2017-01-17] * the whole document *	1-15	
20 A	SEESTED TORBEN ET AL: "The unsialylated subpopulation of recombinant activated factor VII binds to the asialo-glycoprotein receptor (ASGPR) on primary rat hepatocytes", THROMBOSIS AND HAEMOSTASIS, SCHATTAUER GMBH, DE, vol. 104, no. 6, 10 December 2010 (2010-12-10), pages 1166-1173, XP008139984, ISSN: 0340-6245, DOI: 10.1160/TH10-06-0356 * the whole document * * Introduction * -----	1-15	TECHNICAL FIELDS SEARCHED (IPC)
25			
30 A			
35			
40			
45			
50 1	The present search report has been drawn up for all claims		
55	Place of search Munich	Date of completion of the search 18 January 2017	Examiner Bretherick, James
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



EUROPEAN SEARCH REPORT

Application Number

EP 16 19 6780

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10	A APPA RUPA S ET AL: "Investigating clearance mechanisms for recombinant activated factor VII in a perfused liver model", THROMBOSIS AND HAEMOSTASIS, SCHATTAUER GMBH, DE, vol. 104, no. 2, 1 August 2010 (2010-08-01), pages 243-251, XP009156273, ISSN: 0340-6245 * the whole document * * Especially * * page 247; figure 3 * -----	1-15	
15			
20			
25			
30			TECHNICAL FIELDS SEARCHED (IPC)
35			
40			
45			
50	1 The present search report has been drawn up for all claims		
55	Place of search Munich	Date of completion of the search 18 January 2017	Examiner Bretherick, James
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 16 19 6780

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18-01-2017

10	Patent document cited in search report	Publication date	Patent family member(s)	Publication date
15	WO 2004111242 A1 23-12-2004	AT 458057 T AU 2004247799 A1 AU 2010200793 A1 BR PI0411650 A CA 2529828 A1 CN 1839203 A CY 1109984 T1 DK 1644504 T3 EP 1644504 A1 ES 2338425 T3 HK 1095357 A1 HR P20100264 T1 IL 172364 A JP 4915918 B2 JP 5580283 B2 JP 2006527982 A JP 2012105650 A KR 20060022283 A MX PA05013769 A NZ 544728 A NZ 573412 A PT 1644504 E RU 2373282 C2 SI 1644504 T1 US 2005164932 A1 US 2006240526 A1 US 2006241041 A1 US 2006252128 A1 US 2007117756 A1 WO 2004111242 A1	15-03-2010 23-12-2004 25-03-2010 08-08-2006 23-12-2004 27-09-2006 10-09-2014 25-05-2010 12-04-2006 07-05-2010 14-09-2012 31-08-2010 30-01-2014 11-04-2012 27-08-2014 14-12-2006 07-06-2012 09-03-2006 08-03-2006 30-04-2009 30-09-2010 22-03-2010 20-11-2009 30-06-2010 28-07-2005 26-10-2006 26-10-2006 09-11-2006 24-05-2007 23-12-2004	15-05-2009 30-09-2004 15-04-2010 30-09-2004 15-06-2009 06-08-2012 28-12-2005 05-08-2009 23-10-2009 07-08-2012 28-12-2011 21-09-2006 24-05-2012 08-03-2007 30-12-2010
20	WO 2004083361 A2 30-09-2004	AT 431403 T AU 2004221761 A1 AU 2010201201 A1 CA 2519873 A1 DK 1608745 T3 DK 2085470 T3 EP 1608745 A2 EP 2085470 A1 ES 2327044 T3 ES 2386010 T3 JP 4847856 B2 JP 2006521098 A JP 2012095641 A US 2007054366 A1 US 2010330669 A1	15-05-2009 30-09-2004 15-04-2010 30-09-2004 15-06-2009 06-08-2012 28-12-2005 05-08-2009 23-10-2009 07-08-2012 28-12-2011 21-09-2006 24-05-2012 08-03-2007 30-12-2010	
25				
30				
35				
40				
45				
50				
55				

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 16 19 6780

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18-01-2017

10	Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		WO	2004083361 A2	30-09-2004
15	US 2009291890 A1	26-11-2009	AU 2009234390 A1	15-10-2009
			BR PI0911060 A2	13-10-2015
			CA 2721038 A1	15-10-2009
			CN 102083971 A	01-06-2011
			CN 105039291 A	11-11-2015
20			CO 6331368 A2	20-10-2011
			DK 2281037 T3	11-01-2016
			DK 2687596 T3	29-02-2016
			EA 201001628 A1	30-06-2011
			EA 201301350 A1	30-07-2014
25			EP 2281037 A2	09-02-2011
			EP 2679678 A1	01-01-2014
			EP 2687596 A2	22-01-2014
			EP 3064580 A1	07-09-2016
			ES 2556596 T3	19-01-2016
30			ES 2560236 T3	18-02-2016
			ES 2577055 T3	12-07-2016
			HK 1154273 A1	05-08-2016
			HK 1192905 A1	06-01-2017
			HK 1194096 A1	05-08-2016
35			IL 208373 A	31-07-2014
			IL 225740 A	30-04-2015
			JP 5659312 B1	28-01-2015
			JP 5976021 B2	23-08-2016
			JP 2011517942 A	23-06-2011
40			JP 2014113157 A	26-06-2014
			JP 2015017093 A	29-01-2015
			KR 20110008084 A	25-01-2011
			NZ 588322 A	30-03-2012
			PT 2281037 E	27-01-2016
45			SG 190560 A1	28-06-2013
			SG 193779 A1	30-10-2013
			SI 2281037 T1	29-02-2016
			TW 200942258 A	16-10-2009
			TW 201431877 A	16-08-2014
50			US 2009291890 A1	26-11-2009
			US 2014044701 A1	13-02-2014
			US 2016376577 A1	29-12-2016
			WO 2009126307 A2	15-10-2009
55			WO 2005123916 A2	29-12-2005
			EP 1761630 A2	14-03-2007
			JP 2008503541 A	07-02-2008
			US 2008058255 A1	06-03-2008
			WO 2005123916 A2	29-12-2005

EPO FORM P0459
For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 16 19 6780

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18-01-2017

10	Patent document cited in search report	Publication date	Patent family member(s)	Publication date
<hr/>				
15				
20				
25				
30				
35				
40				
45				
50				
55				

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61745674 B [0001]
- US 61787026 B [0001]
- US 20080058255 A1, Bolt [0007]
- US 8008252 B, DeFrees [0011]
- US 5180583 A, Hedner [0012]
- WO 200158935 A, Maxygen [0039]
- US 7371543 B, Pedersen [0039]

Non-patent literature cited in the description

- **DICKINSON et al.** *Proc. Natl. Acad. Sci USA*, 1996, vol. 93, 14379-14384 [0007]
- **TOSO et al.** Lack of Heavy Chain Glycosylation in Patient with Factor VII Deficiency Not Responsible for Mutant FVIIa Activity. *Blood*, 16 November 2000, vol. 96 (11), 79b [0007]
- **FEBS Lett.**, 1988, vol. 238 (1), 31-34 [0026]
- **CALICETI P ; VERONESE FM.** Pharmacokinetic and biodistribution properties of poly(ethylene glycol)-protein conjugates. *Adv Drug Deliv Rev.*, 2003, vol. 55 (10), 1261-77 [0028]
- **WEINSTEIN T et al.** Distribution of glycosaminoglycans in rat renal tubular epithelium. *J Am Soc Nephrol.*, 1997, vol. 8 (4), 586-95 [0028]
- **CHOI HS et al.** Renal clearance of quantum dots. *Nat Biotechnol.*, 2007, vol. 25 (10), 1165-70 [0028]
- **BYRNE B. et al.** Sialic acids: carbohydrate moieties that influence the biological and physical properties of biopharmaceutical proteins and living cells. *Drug Discovery Today*, 2007, vol. 12 (7-8), 319 [0028]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor [0033]
- **NEEDLEMAN.** *J. Mol. Biol.*, 1970, vol. 48, 444-453 [0043]
- **RICE, P. ; LONGDEN, I. ; BLEASBY, A.** EMBOSS: The European Molecular Biology Open Software Suite. *Trends in Genetics*, 2000, vol. 16 (6), 276-277 [0043]
- **ALTSCHUL.** *J. Mol. Biol.*, 1990, vol. 215, 403-10 [0043]
- **ALTSCHUL et al.** *Nucleic Acids Res.*, 1997, vol. 25 (17), 3389-3402 [0043]
- **GRAHAM et al.** *J. Gen. Virol.*, 1977, vol. 36, 59-72 [0047]
- **WAECHTER ; BASERGA.** *Proc. Natl. Acad. Sci. USA*, 1982, vol. 79, 1106-1110 [0047]
- **URLAUB ; CHASIN.** *Proc. Natl. Acad. Sci. USA*, 1980, vol. 77, 4216-4220 [0047]
- **URLAUB et al.** *Cell*, 1983, vol. 33, 405-412 [0047]
- Remington's Pharmaceutical Sciences. Mack Publishing Company, 1990 [0051]