



(11) **EP 3 385 263 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
10.10.2018 Bulletin 2018/41

(51) Int Cl.:
C07D 401/14 ^(2006.01) **A61K 31/435** ^(2006.01)
A61K 31/445 ^(2006.01) **A61P 37/00** ^(2006.01)

(21) Application number: **18171833.9**

(22) Date of filing: **07.06.2013**

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME

(30) Priority: **08.06.2012 US 201261657360 P**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
13800042.7 / 2 858 499

(71) Applicants:
• **Biogen MA Inc.**
Cambridge, MA 02142 (US)
• **Sunesis Pharmaceuticals, Inc.**
South San Francisco, CA 94080 (US)

(72) Inventors:
• **HOPKINS, Brian T.**
Newton, MA Massachusetts 02458 (US)
• **CONLON, Patrick**
Wakefield, MA Massachusetts 01880 (US)

- **CHAN, Timothy R.**
Newton, MA Massachusetts 02458 (US)
- **JENKINS, Tracy J.**
Belmont, MA Massachusetts 02478 (US)
- **CAI, Xiongwei**
Weston, MA Massachusetts 02493 (US)
- **HUMORA, Michael**
Weston, MA Massachusetts 02493 (US)
- **SHI, Xianglin**
Weston, MA Massachusetts 02493 (US)
- **MILLER, Ross A.**
South Plainfield, NJ New Jersey 07080 (US)
- **THOMPSON, Andrew**
South Plainfield, NJ New Jersey 07080 (US)

(74) Representative: **Tostmann, Holger Carl**
Wallinger Ricker Schlotter Tostmann
Patent- und Rechtsanwälte Partnerschaft mbB
Zweibrückenstrasse 5-7
80331 München (DE)

Remarks:

This application was filed on 11-05-2018 as a
divisional application to the application mentioned
under INID code 62.

(54) **PYRIMIDINYL TYROSINE KINASE INHIBITORS**

(57) The present invention provides pyrimidinyl compounds and compositions thereof which are useful as inhibitors of Bruton's tyrosine kinase and which exhibit desirable characteristics for the same.

EP 3 385 263 A1

Description**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] The present application claims priority to U.S. provisional application serial number 61/657,360, filed June 8, 2012, the entire contents of which are hereby incorporated herein by reference.


BACKGROUND OF THE INVENTION

[0002] The Tec kinases are non-receptor tyrosine kinases including: Tec (tyrosine kinase expressed in hepatocellular carcinoma), Btk (Bruton's tyrosine kinase), Itk (interleukin-2 (IL-2)-inducible T-cell kinase; also known as Emt or Tsk), Rlk (resting lymphocyte kinase; also known as Txk), Lck (lymphocyte-specific protein tyrosine kinase), and Bmx (bone-marrow tyrosine kinase gene on chromosome X; also known as Etk)). These kinases are primarily expressed in haematopoietic cells, although expression of Bmx and Tec has been detected in endothelial and liver cells. Tec kinases (Itk, Rlk and Tec) are expressed in T cells and are all activated downstream of the T-cell receptor (TCR). Btk is a downstream mediator of B cell receptor (BCR) signaling which is involved in regulating B cell activation, proliferation, and differentiation. More specifically, Btk contains a PH domain that binds phosphatidylinositol (3,4,5)-trisphosphate (PIP3). PIP3 binding induces Btk to phosphorylate phospholipase C (PLC γ), which in turn hydrolyzes PIP2 to produce two secondary messengers, inositol triphosphate (IP3) and diacylglycerol (DAG), which activate protein kinase PKC, which then induces additional B-cell signaling. Mutations that disable Btk enzymatic activity result in XLA syndrome (X-linked agammaglobulinemia), a primary immunodeficiency. Given the critical roles which Tec kinases play in both B-cell and T-cell signaling, Tec kinases are targets of interest for autoimmune disorders.

[0003] Consequently, there is a great need in the art for effective inhibitors for Tec kinases such as Btk. The present invention fulfills these and other needs.

SUMMARY OF THE INVENTION

[0004] In certain embodiments, the present invention provides a compound of formula I:

**I**

or a pharmaceutically acceptable salt thereof, wherein R¹ and Ring A are as defined and described herein. Such compounds are inhibitors of the Tec kinase family, including Btk. Accordingly, provided compounds can be used in a variety of methods including *in vitro* screening and activity assays as well as *in vivo* pre-clinical, clinical, and therapeutic settings, as described in detail herein.


[0005] In certain embodiments, the present invention provides pharmaceutical formulations comprising provided compounds.

[0006] In certain embodiments, the present invention provides a method of decreasing enzymatic activity of Btk. In some embodiments, such methods include contacting Btk with an effective amount of a Btk inhibitor.

[0007] In certain embodiments, the present invention provides a method of treating a disorder responsive to Btk inhibition in a subject in need thereof. Such disorders and methods are described in detail herein.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

[0008] In certain embodiments, the present invention provides a compound of formula I:



or a pharmaceutically acceptable salt thereof;
wherein:

each R^1 is independently hydrogen, an optionally substituted C_{1-6} aliphatic group, an optionally substituted 3-7 membered monocyclic heterocyclic group, or an optionally substituted heterocyclalkyl group having 3-7 carbon atoms and 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;


or two R^1 groups are taken together with their intervening atoms to form an optionally substituted 3-7 membered saturated or partially unsaturated monocyclic heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

wherein optionally substituted groups may be substituted with halogen, $-NO_2$, $-CN$, $-OR$, $-SR$, $-N(R)_2$, $-C(O)R$, $-CO_2R$, $-N(R)C(O)OR$, $-C(O)N(R)_2$, $-OC(O)R$, $-N(R)C(O)R$, $-S(O)R$, $-S(O)_2R$, or $-S(O)_2N(R)_2$;

each R is independently hydrogen or C_{1-6} aliphatic;

or two R groups attached to the same nitrogen are taken together with their intervening atoms to form an optionally substituted 3-7 membered saturated or partially unsaturated monocyclic heterocyclic ring having 1-2 heteroatoms, in which any second heteroatom is independently selected from nitrogen, oxygen, or sulfur;

Ring A is



R^2 is $-Cl$ or $-F$; and

R^3 is $-CF_3$, $-OCF_3$, or $-F$.

Definitions

[0009] Compounds of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry", 5th Ed., Ed.: Smith, M.B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.

[0010] The abbreviations used herein have their conventional meaning within the chemical and biological arts. The chemical structures and formulae set forth herein are constructed according to the standard rules of chemical valency known in the chemical arts.

[0011] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle," "cycloaliphatic" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups

contain 1-6 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-5 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-4 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-3 aliphatic carbon atoms, and in yet other embodiments, aliphatic groups contain 1-2 aliphatic carbon atoms. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

[0012] As used herein, the term "cycloaliphatic" (or "carbocycle" or "cycloalkyl") refers to a monocyclic C₃-C₆ hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule.

[0013] As used herein, the terms "heterocycle," "heterocyclyl," and "heterocyclic ring" are used interchangeably and refer to a stable 3- to 7-membered monocyclic heterocyclic moiety that is either saturated or partially unsaturated, and having, in addition to carbon atoms, one or more, preferably one to four, heteroatoms, as defined above. When used in reference to a ring atom of a heterocycle, the term "nitrogen" includes a substituted nitrogen. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl), or ⁺NR (as in N-substituted pyrrolidinyl). The term "heterocyclylalkyl" refers to an alkyl group substituted by a heterocyclyl, wherein the alkyl and heterocyclyl portions independently are optionally substituted.

[0014] A heterocyclic ring can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure and any of the ring atoms can be optionally substituted. Examples of such saturated or partially unsaturated heterocyclic radicals include, without limitation, tetrahydrofuranyl, tetrahydrothiophenyl pyrrolidinyl, piperidinyl, pyrrolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, diazepinyl, oxazepinyl, thiazepinyl, morpholinyl, and quinuclidinyl.

[0015] As used herein, the term "partially unsaturated" refers to a ring moiety that includes at least one double or triple bond. The term "partially unsaturated" is intended to encompass rings having multiple sites of unsaturation, but is not intended to include aryl or heteroaryl moieties, as herein defined.

[0016] The term "heteroatom" means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or NR⁺ (as in N-substituted pyrrolidinyl)).


[0017] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in J. Pharmaceutical Sciences, 1977, 66, 1-19, incorporated herein by reference.

[0018] In certain embodiments, the neutral forms of the compounds are regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. In some embodiments, the parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents.

[0019] Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, Z and E double bond isomers, and Z and E conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures including the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ¹³C- or ¹⁴C-enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools, as probes in biological assays, or as therapeutic agents in accordance with the present invention.

Compounds

[0020] As described above, in certain embodiments the present invention provides a compound of formula I:



I

of a pharmaceutically acceptable salt thereof, wherein R^1 and Ring A are as defined and described herein.

[0021] Compounds of formula I have unexpectedly been found to exhibit advantageous properties over known inhibitors of Btk. In certain embodiments, compounds of formula I have increased potency. Without wishing to be bound by any particular theory, it is believed that compounds disclosed herein possess improved potency as Btk inhibitors, an improved off-target profile as measured by hERG inhibition or PXR induction assays, or a combination thereof. Experimental data showing such advantageous properties is provided in the ensuing Examples.

[0022] In some embodiments, both R^1 are hydrogen. In some embodiments, each R^1 is independently C_{1-6} aliphatic. In some embodiments, each R^1 is independently C_{1-5} aliphatic. In some embodiments, each R^1 is independently C_{1-4} aliphatic. In some embodiments, each R^1 is independently C_{1-3} aliphatic. In some embodiments, each R^1 is independently C_{1-2} aliphatic. In some embodiments, both R^1 are methyl.

[0023] In some embodiments, each R^1 is independently hydrogen or C_{1-6} aliphatic. In some embodiments, each R^1 is independently hydrogen or C_{1-5} aliphatic. In some embodiments, each R^1 is independently hydrogen or C_{1-4} aliphatic. In some embodiments, each R^1 is independently hydrogen or C_{1-3} aliphatic. In some embodiments, each R^1 is independently hydrogen or C_{1-2} aliphatic. In some embodiments, each R^1 is independently hydrogen or methyl.


[0024] In some embodiments, one R^1 is hydrogen or and the other R^1 is C_{1-6} aliphatic. In some embodiments, one R^1 is hydrogen and the other R^1 is methyl. In some embodiments, one R^1 is hydrogen and the other R^1 is ethyl. In some embodiments, one R^1 is hydrogen and the other R^1 is C_{1-6} (cycloalkyl)alkyl. In some embodiments, one R^1 is hydrogen and the other R^1 is C_{1-6} (cycloalkyl).

[0025] In some embodiments, one R^1 is hydrogen and the other R^1 is C_{1-6} aliphatic optionally substituted with -OR, wherein R is hydrogen or C_{1-6} aliphatic.

[0026] In some embodiments, one R^1 is hydrogen and the other R^1 is a heterocyclylalkyl group having 3-7 carbon atoms and 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, one R^1 is hydrogen and the other R^1 is an optionally substituted 3-7 membered monocyclic heterocycle.


[0027] In some embodiments, two R^1 groups are taken together with their intervening atoms to form an optionally substituted 3-5 membered saturated or partially unsaturated monocyclic heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, two R^1 groups are taken together with their intervening atoms to form an optionally substituted piperazine ring.

[0028] As described above, Ring A is




[0029] In certain embodiments, R^2 is -Cl. In other embodiments, R^2 is -F. In some embodiments, R^3 is -CF₃. In some embodiments, R^3 is -OCF₃. In some embodiments, R^3 is -F.

[0030] In certain embodiments, Ring A is selected from the group consisting of:



[0031] In some embodiments, in compounds described herein there is a *trans* stereochemical relationship between the piperidine substituent bearing the carboxamide group and the piperidine substituent bearing the lactam group.


[0032] In some embodiments, the present invention provides a compound of formula **II-a**:



II-a

or a pharmaceutically acceptable salt thereof, wherein each of R^1 , R^2 , and R^3 is as defined above and described in classes and subclasses herein.


[0033] In some embodiments, the present invention provides a compound of formula **II-b**:



II-b

or a pharmaceutically acceptable salt thereof, wherein each of R^1 , R^2 , and R^3 is as defined above and described in classes and subclasses herein.


[0034] In some embodiments, the present invention provides a compound of formula **III**:



III

or a pharmaceutically acceptable salt thereof, wherein each of R^2 and R^3 is as defined above and described in classes and subclasses herein.

[0035] In some embodiments, the present invention provides a compound of formula **IV**:




IV

or a pharmaceutically acceptable salt thereof, wherein each of R^1 and R^3 is as defined above and described in classes and subclasses herein. In some embodiments, both R^1 are hydrogen. In some embodiments, one R^1 is hydrogen and the other R^1 is methyl.

[0036] In some embodiments, a provided compound is a compound depicted in Table 1, below, or a pharmaceutically acceptable salt thereof.






Table 1 - Selected compounds of formula I.

[0037]



I-1


I-2

**I-3****I-4****I-5****I-6****I-7**

[0038] Compounds of the invention are synthesized by an appropriate combination of generally well known synthetic methods. Techniques useful in synthesizing the compounds of the invention are both readily apparent and accessible to those of skill in the relevant art. The discussion below is offered to illustrate certain of the diverse methods available for use in assembling the compounds of the invention. However, the discussion is not intended to define the scope of reactions or reaction sequences that are useful in preparing the compounds of the present invention.


[0039] Compounds of formula I may be generally prepared according to Scheme 1.

Scheme 1




[0040] Compounds of formula I may also be generally prepared according to Schemes 2, 3, 3a, 4, 4a, 5, or 5a.


Scheme 2




Scheme 3




Scheme 3a




Scheme 4




Scheme 4a



Scheme 5



Scheme 5a



[0041] The PG, PG₁, PG₂, and PG₃ groups of compounds in Schemes 1 through 5a are each independently a suitable protecting group. Such ester and amine protecting groups are known in the art and are described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, the entirety of which is incorporated herein by reference. In some embodiments, a protecting group is a Boc group.

[0042] In certain embodiments, each of the synthetic steps in Schemes 1 through 5a may be performed sequentially with isolation of each intermediate performed after each step. Alternatively, each of the steps as depicted in Schemes 1, 2, 3, and 4 above, may be performed in a manner whereby no isolation of one or more intermediates is performed.

[0043] In certain embodiments, all the steps of the aforementioned synthesis may be performed to prepare the desired final product. In other embodiments, two, three, four, five, or more sequential steps may be performed to prepare an intermediate or the desired final product.

Uses, Formulation and Administration

[0044] In certain embodiments, compounds of the present invention are for use in medicine. In some embodiments, the present invention provides method of decreasing enzymatic activity of a kinase in the Tec kinase family (e.g., Tec, Btk, Itk, Txk, Lck, and Bmx). In some embodiments, such methods include contacting a kinase of the Tec kinase family with an effective amount of a Tec kinase family inhibitor. Therefore, the present invention further provides methods of inhibiting Tec kinase family enzymatic activity by contacting a Tec kinase family member with a Tec kinase family inhibitor of the present invention. As used herein, the term "Tec kinase family member" refers to any non-receptor tyrosine kinase in the Tec kinase family. In some embodiments, Tec kinase family members are Tec, Btk, Itk, Txk, Lck, and Bmx.

[0045] In some embodiments, the present invention provides methods of decreasing Btk enzymatic activity. In some embodiments, such methods include contacting a Btk with an effective amount of a Btk inhibitor. Therefore, the present invention further provides methods of inhibiting Btk enzymatic activity by contacting a Btk with a Btk inhibitor of the present invention.

[0046] Btk enzymatic activity, as used herein, refers to Btk kinase enzymatic activity. For example, where Btk enzymatic activity is decreased, PIP3 binding and/or phosphorylation of PLC γ is decreased. In some embodiments, the half maximal inhibitory concentration (IC₅₀) of the Btk inhibitor against Btk is less than 1 μ M. In some embodiments, the IC₅₀ of the Btk inhibitor against Btk is less than 500 nM. In some embodiments, the IC₅₀ of the Btk inhibitor against Btk is less than

100 nM. In some embodiments, the IC₅₀ of the Btk inhibitor against Btk is less than 10 nM. In some embodiments, the IC₅₀ of the Btk inhibitor against Btk is less than 1 nM. In some embodiments, the IC₅₀ of the Btk inhibitor against Btk is from 0.1 nM to 10 μM. In some embodiments, the IC₅₀ of the Btk inhibitor against Btk is from 0.1 nM to 1 μM. In some embodiments, the IC₅₀ of the Btk inhibitor against Btk is from 0.1 nM to 100 nM. In some embodiments, the IC₅₀ of the Btk inhibitor against Btk is from 0.1 nM to 10 nM.

[0047] In some embodiments, inhibitors of such Tec kinases are useful for the treatment of diseases and disorders that may be alleviated by inhibiting (i.e., decreasing) enzymatic activity of one or more Tec kinases. The compounds of the invention are effective inhibitors of Tec family kinases and would thus be useful in treating diseases associated with the activity of one or more of the Tec family kinases. The term "diseases" means diseases, syndromes, or disease symptoms. Thus, the present invention provides methods of treating autoimmune disorders, inflammatory disorders, and cancers in a subject in need thereof. Such methods include administering to the subject a therapeutically effective amount of an inhibitor of Tec, Btk, Itk, Txk, Lck, and/or Bmx kinase.

[0048] The term "autoimmune disorders" includes diseases or disorders involving inappropriate immune response against native antigens, such as acute disseminated encephalomyelitis (ADEM), Addison's disease, alopecia areata, antiphospholipid antibody syndrome (APS), hemolytic anemia, autoimmune hepatitis, bullous pemphigoid (BP), Coeliac disease, dermatomyositis, diabetes mellitus type 1, Good Pasture's syndrome, Graves' disease, Guillain-Barré syndrome (GBS), Hashimoto's disease, idiopathic thrombocytopenic purpura, lupus or systemic lupus erythematosus (SLE), mixed connective tissue disease, multiple sclerosis, myasthenia gravis, pemphigus vulgaris, hemophilia with inhibitors, pernicious anaemia, polymyositis, primary biliary cirrhosis, Sjögren's syndrome, temporal arteritis, and Wegener's granulomatosis. The term "inflammatory disorders" includes diseases or disorders involving acute or chronic inflammation such as allergies, asthma (e.g., allergic asthma), atopic dermatitis, prostatitis, glomerulonephritis, pelvic inflammatory disease (PID), inflammatory bowel disease (IBD, e.g., Crohn's disease, ulcerative colitis), reperfusion injury, rheumatoid arthritis, transplant rejection (including transplant patients with a positive cross-match) and vasculitis. In certain embodiments, the present invention provides methods of treating disease, disorders, or conditions that approved for treatment with rituximab (a monoclonal antibody against CD20), including non-Hodgkin's lymphoma (NHL), chronic lymphocytic leukemia (CLL), RA, Wegener's granulomatosis (WG), and microscopic polyangiitis (MPA). In some embodiments, the present invention provides a method of treating rheumatoid arthritis (RA), SLE, or atopic dermatitis using compounds disclosed herein.

[0049] The term "cancer" includes diseases or disorders involving abnormal cell growth and/or proliferation, such as glioma, thyroid carcinoma, breast carcinoma, lung cancer (e.g. small-cell lung carcinoma, non-small-cell lung carcinoma), gastric carcinoma, gastrointestinal stromal tumors, pancreatic carcinoma, bile duct carcinoma, ovarian carcinoma, endometrial carcinoma, prostate carcinoma, renal cell carcinoma, lymphoma (e.g., anaplastic large-cell lymphoma), leukemia (e.g. acute myeloid leukemia, T-cell leukemia, chronic lymphocytic leukemia), multiple myeloma, malignant mesothelioma, malignant melanoma, and colon cancer (e.g. microsatellite instability-high colorectal cancer). In some embodiments, the present invention provides a method of treating leukemia or lymphoma.

[0050] The term "subject," as used herein, refers to a mammal to whom a pharmaceutical composition is administered. Exemplary subjects include humans, as well as veterinary and laboratory animals such as horses, pigs, cattle, dogs, cats, rabbits, rats, mice, and aquatic mammals.

Selected Indications and B Cell Inhibition

[0051] As described above, provided compounds are useful for the treatment of disease, including RA and SLE. As described in more detail below, these diseases are affiliated with B cells. Thus, the present disclosure encompasses the recognition that provided compounds are useful as therapeutics for these and other indications.

[0052] Dysregulation of the immune system is central to the pathogenesis (Panayi GS, et al. Rheum Dis Clin North Am 2001; 27:317-334) of RA. While most of the infiltrating leukocytes in the synovium are T lymphocytes (primarily activated CD4+ T cells) and cells of monocyte/macrophage origin (which release pro-inflammatory cytokines such as IL-1, TNF-alpha and IL-6 and proteolytic enzymes including collagenases and metalloproteinases), B-cells and plasma cells are also found in the synovial fluid (Zhang Z, Bridges SL. Rheum Dis Clin North Am 2001; 27:335-353). A clear role for B cells and their associated effector functions in RA have been demonstrated by the efficacy of rituximab, a selective B cell depleting therapeutic, which is approved for treatment of RA (Cohen SB, et al.; REFLEX Trial Group. Arthritis Rheum. 2006 Sep; 54(9):2793-806).

[0053] Although the etiology of SLE is not fully understood, pathogenic autoantibodies and deposition of immune complexes are felt to be critical to the development of widespread tissue damage (Klippel JH, et al. Primer on the rheumatic diseases. Atlanta: Arthritis Foundation; 2001). Autoantibody and immune-complex mediated activation can be studied by measuring inhibition of macrophage activation by macrophages stimulated through Fc receptors (see exemplification - FcγR activation of primary human macrophages). Loss of tolerance to self-antigens ultimately lead to the stimulation of B cells to produce auto-antibodies often directed against nuclear or cytoplasmic components. Antibodies

against nuclear components (antinuclear antibodies [ANA]) target nuclear antigens including DNA (typically double-stranded DNA [dsDNA]), RNA, histones and small nuclear ribonucleoproteins. These antibodies combine with self-antigens forming immune complexes which deposit in tissues, incite inflammatory reactions and lead to tissue injury. In addition to their roles in pathogenic autoantibody production, B cells also function as antigen-presenting cells (APCs) to T-cells thus playing a role in the initiation of an antigen-specific response. Given the central role of the humoral arm of the immune system in the pathogenesis of SLE, B cells or the B-cell pathway represent desirable therapeutic targets. Belimumab, a monoclonal antibody recently approved for SLE, blocks the binding BAFF to its receptors that are expressed B cells. These receptors serve to activate and potentiate the survival of B cells consistent with a reduction of circulating B cells observed following treatment with belimumab. See also Chan OT, et al. *Immunol Rev.* 1999b;169:107-121; Navarra SV, et al. *Lancet.* 2011 Feb 26;377(9767):721-31; Furie R, et al. *Arthritis Rheum.* 2011 Dec;63(12):3918-30. The role of B cells and myeloid lineage cells in autoimmune diseases such as SLE is further supported by a recent publication which describes efficacy in a preclinical SLE animal model when mice are treated with a small molecule irreversible Btk inhibitor (Honigberg, L.A. *PNAS.* 2010; 107: 13075).

Combinations

[0054] In certain embodiments, a compound of the present invention is administered in combination with another agent. In some embodiments, a compound of the present invention is useful for treating RA and is administered in combination with a disease-modifying antirheumatic drugs (DMARD), including without limitation: methotrexate, abatacept, azathioprine, certolizumab, chloroquine and hydroxychloroquine, cyclosporin, D-penicillamine, adalimumab, etanercept, golimumab, gold salts (including auranofin and sodium aurothiomalate), infliximab, leflunomide, minocycline, rituximab, sulfasalazine, tocilizumab, or combinations thereof. In some embodiments, a compound of the present invention is administered in combination with a NSAID or corticosteroid. In some embodiments, a compound of the present invention is useful for treating SLE and is administered in combination with an agent for the treatment of SLE, including without limitation: corticosteroids, antimalarials, belimumab, mycophenolate mofetil (MMF) or mycophenolate sodium, azathioprine, or combinations thereof. In some embodiments, a compound of the present invention is useful for treating atopic dermatitis and is administered in combination with a topical agent for the treatment of atopic dermatitis, including without limitation: topical steroids, tacrolimus, methotrexate, mometasone furoate (MMF), azathioprine, retinoids, or combinations thereof.

Assays

[0055] To develop useful Tec kinase family inhibitors, candidate inhibitors capable of decreasing Tec kinase family enzymatic activity may be identified *in vitro*. The activity of the inhibitor compounds can be assayed utilizing methods known in the art and/or those methods presented herein.

[0056] Compounds that decrease Tec kinase family members' enzymatic activity may be identified and tested using a biologically active Tec kinase family member, either recombinant or naturally occurring. Tec kinases can be found in native cells, isolated *in vitro*, or co-expressed or expressed in a cell. Measuring the reduction in the Tec kinase family member enzymatic activity in the presence of an inhibitor relative to the activity in the absence of the inhibitor may be performed using a variety of methods known in the art, such as the POLYGAT-LS assays described below in the Examples. Other methods for assaying the activity of Btk and other Tec kinases are known in the art. The selection of appropriate assay methods is well within the capabilities of those of skill in the art.

[0057] Once compounds are identified that are capable of reducing Tec kinase family members' enzymatic activity, the compounds may be further tested for their ability to selectively inhibit a Tec kinase family member relative to other enzymes.

[0058] Compounds may be further tested in cell models or animal models for their ability to cause a detectable changes in phenotype related to a Tec kinase family member activity. In addition to cell cultures, animal models may be used to test Tec kinase family member inhibitors for their ability to treat autoimmune disorders, inflammatory disorders, or cancer in an animal model.

Pharmaceutical Compositions

[0059] In another aspect, the present invention provides pharmaceutical compositions comprising a compound of formula I or a compound of formula I in combination with a pharmaceutically acceptable excipient (e.g., carrier).

[0060] The pharmaceutical compositions include optical isomers, diastereomers, or pharmaceutically acceptable salts of the inhibitors disclosed herein. The compound of formula I included in the pharmaceutical composition may be covalently attached to a carrier moiety, as described above. Alternatively, the compound of formula I included in the pharmaceutical composition is not covalently linked to a carrier moiety.

[0061] A "pharmaceutically acceptable carrier," as used herein refers to pharmaceutical excipients, for example, pharmaceutically, physiologically, acceptable organic or inorganic carrier substances suitable for enteral or parenteral application that do not deleteriously react with the active agent. Suitable pharmaceutically acceptable carriers include water, salt solutions (such as Ringer's solution), alcohols, oils, gelatins, and carbohydrates such as lactose, amylose or starch, fatty acid esters, hydroxymethylcellulose, and polyvinyl pyrrolidone. Such preparations can be sterilized and, if desired, mixed with auxiliary agents such as lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, and/or aromatic substances and the like that do not deleteriously react with the compounds of the invention.

[0062] The compounds of the invention can be administered alone or can be co-administered to the subject. Co-administration is meant to include simultaneous or sequential administration of the compounds individually or in combination (more than one compound). The preparations can also be combined, when desired, with other active substances (e.g. to reduce metabolic degradation).

[0063] Compounds of the present invention can be prepared and administered in a wide variety of oral, parenteral, and topical dosage forms. Thus, the compounds of the present invention can be administered by injection (e.g. intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally). Also, the compounds described herein can be administered by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. It is also envisioned that multiple routes of administration (e.g., intramuscular, oral, transdermal) can be used to administer the compounds of the invention.

[0064] For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substance that may also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

[0065] In powders, the carrier is a finely divided solid in a mixture with the finely divided active component. In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

[0066] The powders and tablets preferably contain from 5% to 70% of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

[0067] For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

[0068] Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water/propylene glycol solutions. For parenteral injection, liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

[0069] When parenteral application is needed or desired, particularly suitable admixtures for the compounds of the invention are injectable, sterile solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. In particular, carriers for parenteral administration include aqueous solutions of dextrose, saline, pure water, ethanol, glycerol, propylene glycol, peanut oil, sesame oil, polyoxyethylene-block polymers, and the like. Ampoules are convenient unit dosages. The compounds of the invention can also be incorporated into liposomes or administered via transdermal pumps or patches. Pharmaceutical admixtures suitable for use in the present invention include those described, for example, in *Pharmaceutical Sciences* (17th Ed., Mack Pub. Co., Easton, PA) and WO 96/05309, the teachings of both of which are hereby incorporated by reference.

[0070] Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizers, and thickening agents as desired. Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

[0071] Also included are solid form preparations that are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

[0072] The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the

appropriate number of any of these in packaged form.

[0073] The quantity of active component in a unit dose preparation may be varied or adjusted from 0.1 mg to 10000 mg, more typically 1.0 mg to 1000 mg, most typically 10 mg to 500 mg, according to the particular application and the potency of the active component. The composition can, if desired, also contain other compatible therapeutic agents.

[0074] Some compounds may have limited solubility in water and therefore may require a surfactant or other appropriate co-solvent in the composition. Such co-solvents include: Polysorbate 20, 60, and 80; Pluronic F-68, F-84, and P-103; cyclodextrin; and polyoxyl 35 castor oil. Such co-solvents are typically employed at a level between about 0.01 % and about 2% by weight.

[0075] Viscosity greater than that of simple aqueous solutions may be desirable to decrease variability in dispensing the formulations, to decrease physical separation of components of a suspension or emulsion of formulation, and/or otherwise to improve the formulation. Such viscosity building agents include, for example, polyvinyl alcohol, polyvinyl pyrrolidone, methyl cellulose, hydroxy propyl methylcellulose, hydroxyethyl cellulose, carboxymethyl cellulose, hydroxy propyl cellulose, chondroitin sulfate and salts thereof, hyaluronic acid and salts thereof, and combinations of the foregoing. Such agents are typically employed at a level between about 0.01% and about 2% by weight.

[0076] The compositions of the present invention may additionally include components to provide sustained release and/or comfort. Such components include high molecular weight, anionic mucomimetic polymers, gelling polysaccharides, and finely-divided drug carrier substrates. These components are discussed in greater detail in U.S. Pat. Nos. 4,911,920; 5,403,841; 5,212,162; and 4,861,760. The entire contents of these patents are incorporated herein by reference in their entirety for all purposes.

Effective Dosages

[0077] Pharmaceutical compositions provided by the present invention include compositions wherein the active ingredient is contained in a therapeutically effective amount, i.e., in an amount effective to achieve its intended purpose. The actual amount effective for a particular application will depend, *inter alia*, on the condition being treated. For example, when administered in methods to treat cancer, such compositions will contain an amount of active ingredient effective to achieve the desired result (e.g. decreasing the number of cancer cells in a subject).

[0078] The dosage and frequency (single or multiple doses) of compound administered can vary depending upon a variety of factors, including route of administration; size, age, sex, health, body weight, body mass index, and diet of the recipient; nature and extent of symptoms of the disease being treated (e.g., the disease responsive to Btk inhibition); presence of other diseases or other health-related problems; kind of concurrent treatment; and complications from any disease or treatment regimen. Other therapeutic regimens or agents can be used in conjunction with the methods and compounds of the invention.

[0079] For any compound described herein, the therapeutically effective amount can be initially determined from cell culture assays. Target concentrations will be those concentrations of active compound(s) that are capable of decreasing kinase enzymatic activity as measured, for example, using the methods described.

[0080] Therapeutically effective amounts for use in humans may be determined from animal models. For example, a dose for humans can be formulated to achieve a concentration that has been found to be effective in animals. The dosage in humans can be adjusted by monitoring kinase inhibition and adjusting the dosage upwards or downwards, as described above. In certain embodiments, the administered dose is in the range of about 10 mg to about 1000 mg per day, either once, twice, or more than twice daily.

[0081] Dosages may be varied depending upon the requirements of the patient and the compound being employed. The dose administered to a patient, in the context of the present invention, should be sufficient to effect a beneficial therapeutic response in the patient over time. The size of the dose also will be determined by the existence, nature, and extent of any adverse side effects. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under circumstances is reached. In some embodiments, the dosage range is 0.001 % to 10% w/v. In some embodiments, the dosage range is 0.1 % to 5% w/v.

[0082] Dosage amounts and intervals can be adjusted individually to provide levels of the administered compound effective for the particular clinical indication being treated. This will provide a therapeutic regimen that is commensurate with the severity of the individual's disease state.

[0083] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

EXEMPLIFICATION


[0084] As depicted in the Examples below, in certain exemplary embodiments, compounds are prepared according

to the following general procedures. It will be appreciated that, although the general methods depict the synthesis of certain compounds of the present invention, the following general methods, and other methods known to one of ordinary skill in the art, can be applied to all compounds and subclasses and species of each of these compounds, as described herein.

Example 1

Synthetic of (3'R,4'S)-1'-tert-butyl 4'-ethyl 2-oxo-[1,3'-bipiperidine]-1',4'-dicarboxylate

[0085]



[0086] Preparation of ethyl 3-oxopiperidine-4-carboxylate intermediate. Ethyl 1-benzyl-3-oxopiperidine-4-carboxylate **1** (15.0 g, 50.5 mmol, 1.0 equiv) was hydrogenated in the presence of 10% Pd/C (1.5 g) catalyst under H₂ at atmospheric pressure in MeOH (250 mL) for 16 h. The catalyst was filtered off and the solvent was *concentrated in vacuo* to give ethyl 3-oxopiperidine-4-carboxylate **2** as a light yellow solid (10.2 g, yield: 98.0%). ESI-MS (M+H)⁺: 172.1. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 4.23 (q, 2H), 3.75 (s, 2H), 3.37 (s, 2H), 3.20-3.16 (m, 2H), 2.44 (t, 1H), 1.25 (t, 3H).

[0087] Preparation of 1-tert-butyl 4-ethyl 3-oxopiperidine-1,4-dicarboxylate. Ethyl 3-oxopiperidine-4-carboxylate **2** (10.2 g, 60.0 mmol, 1.0 equiv) was dissolved in dry MeOH (200 mL), and Et₃N (33.1 mL, 240 mmol, 4.0 equiv) was added. The mixture was stirred for 1 h and Boc₂O (19.5 g, 90.0 mmol, 1.5 equiv) was added and stirred for 16 h. The solvent was *concentrated in vacuo* and the crude was purified by column chromatography (silica, petroleum ether/EtOAc =9:1) to give 1-tert-butyl 4-ethyl 3-oxopiperidine-1,4-dicarboxylate **3** light yellow oil (11.5 g, yield: 86%). ESI-MS (M+H)

56) δ : 216.0. ^1H NMR (400 MHz, CDCl_3) δ : 4.24 (q, 2H), 4.03 (s, 2H), 3.49 (t, 2H), 2.33 (t, 2H), 1.47 (s, 9H), 1.31 (t, 3H).

[0088] Preparation of (S)-1-tert-butyl 4-ethyl 3-((1-phenylethyl)amino)-5,6-dihydro pyridine-1,4-(2H)-dicarboxylate. In a dry flask equipped with a Dean-stark trap and reflux condenser, 1-tert-butyl 4-ethyl 3-oxopiperidine-1,4-dicarboxylate **3** (10.0 g, 37.0 mmol, 1.1 equiv) was dissolved in toluene (100 mL). S-(-)- α -Methylbenzylamine (4.9 g, 40.5 mmol, 1.1 equiv) and *p*-toluenesulfonic acid monohydrate (0.7 g, 3.7 mmol, 0.1 equiv) were added and the mixture was heated to reflux for 16 h. The crude reaction mixture was *concentrated in vacuo* to give (S)-1-tert-butyl 4-ethyl 3-((1-phenylethyl)amino)-5,6-dihydro pyridine-1,4(2H)-dicarboxylate **4** (12.0 g, Y: 88%) as thick orange oil which was used in next step without further purification, ESI-MS (M+H) $^+$: 375.2.

[0089] Preparation of 1-tert-butyl 4-ethyl 3-(((S)-1-phenylethyl)amino)-5,6-dihydro pyridine-1,4(2H) -dicarboxylate. 1-tert-Butyl 4-ethyl 3-(((S)-1-phenylethyl)amino)piperidine-1,4-dicarboxylate **4** (11.2 g, 30.0 mmol, 1.0 equiv) was dissolved in a mixture of CH_3CN (60 mL) and acetic acid (60 mL) and cooled to 0 °C. $\text{NaBH}(\text{OAc})_3$ (19.0 g, 90.0 mmol, 3.0 equiv) was slowly added and the reaction mixture was allowed to stir for 2 h at 0 °C. Saturated NaHCO_3 was slowly added to neutralize the solution to maintain the internal temperature of the flask below 10 °C. The mixture was extracted with EtOAc (50 mL x 3). The combined organic layer was dried (Na_2SO_4), filtered, *concentrated in vacuo*, and then purified by column chromatography (silica, petroleum ether/EtOAc = 9:1) to give 4-ethyl 3-(((S)-1-phenylethyl)amino)-5,6-dihydro pyridine-1,4(2H)-dicarboxylate **5** (8.2 g, Y: 73%) as light yellow oil. ESI-MS (M+H) $^+$: 377.2. ^1H NMR (400 MHz, CD_3OD) δ : 7.31-7.22 (m, 5H), 4.20 (q, 2H), 4.11-3.86 (m, 3H), 3.15 (s, 1H), 3.00-2.90 (m, 2H), 2.64 (d, 2H), 1.87-1.85 (m, 1H), 1.68 (s, 1H), 1.50-1.25 (m, 15H).

[0090] Preparation of trans-1-tert-butyl 4-ethyl 3-(((S)-1-phenylethyl)amino) piperidine-1,4-dicarboxylate. The 1-tert-butyl 4-ethyl 3-(((S)-1-phenylethyl)amino)piperidine-1,4- dicarboxylate **5** (8.0 g, 21.2 mmol, 1.0 equiv) was dissolved in dry EtOH (20 mL) under N_2 . In a separate flame-dried Schlenk flask was placed dry EtOH (150 mL), and sodium (0.450 g, 63.6 mmol, 3.0 equiv) was added portion-wise under N_2 . The mixture was kept under N_2 and vented to remove evolved gases until all of the sodium had dissolved. The clear solution of 1-tert-butyl 4-ethyl 3-(((S)-1-phenylethyl)amino)piperidine-1,4-dicarboxylate was then transferred to the NaOEt solution, and the mixture was stirred at 80 °C under N_2 for 16 h. The solvent was removed under *in vacuo*, and brine (150 mL) was added and the mixture was brought to pH = 10 with 1 N NaOH and extracted with EtOAc (100 mL x 3). The combined organic layers were dried (Na_2SO_4) and *concentrated in vacuo*. The residue was purified by column chromatography (silica, petroleum ether/EtOAc = 5:1) to give (trans)-1-tert-butyl 4-ethyl 3-(((S)-1-phenylethyl) amino)piperidine-1,4-dicarboxylate **6** as a slight yellow solid (3.7 g, yield: 46%). ESI-MS (M+H) $^+$: 377.2.

[0091] Preparation of trans-1-tert-butyl 4-ethyl 3-aminopiperidine-1,4-dicarboxylate. Trans-1-tert-butyl 4-ethyl 3-(((S)-1-phenylethyl)amino)piperidine-1,4-dicarboxylate **6** (3.7 g, 8.3 mmol, 1.0 equiv) was hydrogenated in the presence of 10% Pd/C (0.37 g) catalyst under H_2 at 30 atmospheric pressure in MeOH (100 mL) at 50 °C for 8 h. The catalyst was filtered off and the solvent was removed *in vacuo* to give (trans)-1-tert-butyl 4-ethyl 3-aminopiperidine-1,4-dicarboxylate **7** as light yellow oil (2.5 g, yield: 92%). ESI-MS (M+H) $^+$: 273.1. ^1H NMR (400 MHz, CDCl_3) δ : 4.18 (q, 2H), 3.97-3.94 (m, 2H), 3.37 (s, 1H), 3.07-3.02 (m, 1H), 2.89-2.85 (m, 1H), 2.60-2.55 (m, 1H), 2.01-1.91 (m, 1H), 1.70-1.54 (m, 3H), 1.46 (s, 9H), 1.28 (t, 3H).


[0092] Synthesis of trans-1-tert-butyl 4-ethyl 3-(5-bromopentanamido)piperidine-1,4-dicarboxylate. To a solution of trans-1-tert-butyl 4-ethyl 3-aminopiperidine -1,4-dicarboxylate **7** (2.5 g, 9.2 mmol, 1.0 equiv) in CH_2Cl_2 (50 mL), Et_3N (2.5 mL, 18.4 mmol, 2.0 equiv) was added at rt. After the reaction solution was stirred at rt for 10 min, 5-bromovaleryl chloride (1.9 g, 9.6 mmol, 1.05 eq) was added. The reaction solution was stirred at rt for 2 h. The mixture was quenched with H_2O (20 mL) and extracted with CH_2Cl_2 (50 mL x 3). The organic layer was collected, *concentrated in vacuo*, and the residue was purified by column chromatography (silica, petroleum ether/EtOAc = 1:1) to give (trans)-1-tert-butyl 4-ethyl 3-(5-bromopentanamido)piperidine -1,4-dicarboxylate **8** as yellow oil (3.2 g, yield: 80%). ESI-MS (M+H-56) $^+$: 379.0. ^1H NMR (400 MHz, CDCl_3) δ : 5.99 (d, 1H), 4.39-4.38 (m, 1H), 4.15 (q, 2H), 3.79-3.74 (m, 1H), 3.66-3.60 (m, 1H), 3.41 (t, 2H), 3.30-3.26 (m, 1H), 3.21-3.14 (m, 1H), 2.78-2.74 (m, 1H), 2.19 (t, 2H), 1.99-1.85 (m, 3H), 1.80-1.72 (m, 3H), 1.45 (s, 9H), 1.27 (t, 3H).

[0093] Synthesis of trans-1'-tert-butyl 4'-ethyl 2-oxo-[1,3'-bipiperidine] -1',4'-dicarboxylate. To a solution of trans-1-tert-butyl 4-ethyl 3-(5-bromopentanamido)piperidine-1,4-dicarboxylate **8** (3.0 g, 6.9 mmol, 1.0 equiv) in THF (20 mL), NaH (276 mg, 6.9 mmol, 1.0 equiv) was carefully added in small portions at 0 °C. The reaction solution was stirred at reflux condition for 4 h. The mixture was quenched with H_2O (20 mL), and extracted with EtOAc (30 mL x 3). The organic layer was collected, dried (Na_2SO_4), filtered, and *concentrated in vacuo*. The residue was purified by column chromatography (silica, petroleum ether/EtOAc = 1:2) to give (trans)-1'-tert-butyl 4'-ethyl 2-oxo-[1,3'-bipiperidine] -1',4'-dicarboxylate **9** as a slight yellow oil (2.1 g, yield: 88%). ESI-MS (M+H-56) $^+$: 299.1. ^1H NMR (400 MHz, CDCl_3) δ : 4.10 (q, 4H), 3.38-3.19 (m, 4H), 2.70-2.61 (m, 1H), 2.36-2.31 (m, 2H), 1.95-1.92 (m, 1H), 1.75-1.71 (m, 6H), 1.46 (s, 9H), 1.23 (t, 3H).

Example 2

Preparation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide

[0094]



[0095] Synthesis of *trans*-1'-*tert*-butyl 4'-ethyl-3-iodo-2-oxo-[1,3'-bipiperidine]-1',4'-dicarboxylate. To the solution of *trans*-1'-*tert*-butyl 4'-ethyl-2-oxo-[1,3'-bipiperidine]-1',4'-dicarboxylate **8** (141 mg, 2.58 mmol, 1.0 equiv) in dry toluene (10 mL) at 0 °C, TMEDA (0.89 g, 7.7 mmol, 3.0 equiv) and TMSCl (0.6 mg, 1.0 mmol, 2.0 equiv) were added successively under N₂. After 0.5 h, I₂ (0.98 g, 3.87 mmol, 1.5 equiv) was carefully added in small portions. The reaction solution was stirred at 0 °C to rt for 16 h. The mixture was diluted with EtOAc (100 mL), washed with saturated Na₂S₂O₃ (20 mL × 2) and brine (20 mL), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The crude product **9** (2.2 g, Y: 81%) was used directly in the next step without further purification. ESI-MS (M+H-56)⁺: 424.9. ¹H NMR (400 MHz, CDCl₃) δ: 4.78-4.73 (m, 1H), 4.19-4.04 (m, 4H), 3.55-3.30 (m, 4H), 3.24-3.16 (m, 2H), 2.73-2.60 (m, 1H), 2.22-2.14 (m, 2H), 1.96-1.78 (m, 2H), 1.70-1.60 (m, 1H), 1.44 (s, 9H), 1.25 (t, J = 7.2 Hz, 3H).

[0096] Synthesis of *trans*-1'-*tert*-butyl 4'-ethyl 3-((3-chloro-5-(trifluoromethyl) phenyl)amino)-2-oxo-[1,3'-bipiperidine]-1',4'-dicarboxylate. A 1.0 M solution of lithium bis(trimethyldisilyl)amide in THF (13 mL, 12 mmol, 2.0 equiv) was added through an addition funnel at 10-15 °C to a solution of 3-chloro-5-(trifluoromethyl)aniline (15 g, 78 mmol, 1.2 equiv) in THF (13 mL). The mixture was allowed to stir at room temperature for 20 min and a solution of crude *trans*-1'-*tert*-butyl 4'-ethyl-3-iodo-2-oxo-[1,3'-bipiperidine]-1',4'-dicarboxylate **9** (3.7 g, 65 mmol, 1.0 equiv) in THF (13 mL) was added through an addition funnel at 10-15 °C over 30 min. After addition, the reaction was allowed to stir at the temperature for 30 min. Upon completion, the reaction was cooled to 5 °C and quenched slowly with water (10 mL), keeping the temperature below 20 °C. The quenched reaction was extracted with EtOAc (2 × 30 mL). The combined organic layers were washed with saturated brine (30 mL), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The resulting crude


product was purified over silica gel eluting with a gradient of 10% to 75% of EtOAc in heptanes to give the desired product 10. ESI-MS (M+H-56)⁺: 463.1. ¹H NMR (400 MHz, CDCl₃) δ: 6.92 (s, 1H), 6.71-6.69 (m, 2H), 4.17-4.06 (m, 4H), 3.78-3.68 (m, 2H), 3.46-3.36 (m, 3H), 3.23-3.07 (m, 2H), 2.73-2.65 (m, 1H), 2.44-2.37 (m, 1H), 2.03-1.85 (m, 3H), 1.71-1.61 (m, 2H), 1.46 (s, 9H), 1.27-1.19 (m, 3H).

[0097] Synthesis of *trans*-1'-(*tert*-butoxycarbonyl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid. To a solution of *trans*-1'-(*tert*-butyl 4'-ethyl 3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-1',4'-dicarboxylate 10 (180 mg, 0.33 mmol, 1.0 equiv) in EtOH (5 mL) was added NaOH (40 mg, 0.99 mmol, 3.0 equiv) and solution was stirred at 80 °C for 1 h. The solvent was concentrated *in vacuo* and the residue was suspended in water (10 mL) and adjusted to pH = 6 with HCl (4 N). The precipitate was filtered to afford (*trans*)-1'-(*tert*-butoxycarbonyl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid 11 (150 mg, Y: 82%) as yellow solid which was used next step without further purification. ESI-MS (M+H-85)⁺: 463.1. ¹H NMR (400 MHz, CDCl₃) δ: 6.85 (s, 1H), 6.82 (s, 1H), 6.78 (s, 1H), 4.12-3.96 (m, 4H), 3.53-3.37 (m, 2H), 3.11-3.04 (m, 2H), 2.75-2.67 (m, 1H), 2.24-2.18 (m, 1H), 1.98-1.89 (m, 3H), 1.71-1.58 (m, 2H), 1.44 (s, 9H).

[0098] Synthesis of *trans*-*tert*-butyl 4'-carbamoyl-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-1'-carboxylate. To the solution of *trans* 1'-(*tert*-butoxycarbonyl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid 11 (70 mg, 0.14 mmol, 1.0 equiv) in DMF (2 mL), was added NH₄Cl (22 mg, 0.41 mmol, 3.0 equiv), HBTU (103 mg, 0.270 mmol, 2.0 equiv) and DIPEA (52 mg, 0.41 mmol, 3.0 equiv). The reaction solution was stirred at rt for 16 h, diluted with EtOAc (10 mL) and washed with water (5 mL) and brine (5 mL). The organic phase was separated and concentrated *in vacuo* to afford a crude oil which was purified by pre-HPLC (MeOH/H₂O with 0.05% TFA as mobile phase) to give the compound (*trans*)-*tert*-butyl 4'-carbamoyl-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-1'-carboxylate 12 (60 mg, yield: 86%) as a light solid. ESI-MS (M+H-56)⁺: 463.1. ¹H NMR (400 MHz, CD₃OD) δ: 6.87-6.86 (m, 1H), 6.84-6.83 (m, 1H), 6.80 (s, 1H), 4.11-4.03 (m, 3H), 3.53-3.35 (m, 2H), 3.20-3.08 (m, 2H), 2.77-2.74 (m, 1H), 2.25-2.18 (m, 1H), 1.99-1.88 (m, 3H), 1.70-1.60 (m, 2H), 1.46 (s, 9H).


[0099] Synthesis of *trans*-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. To the solution of *trans*-*tert*-butyl 4'-carbamoyl-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-1'-carboxylate 12 (60 mg, 0.11 mmol) in CH₂Cl₂ (1.0 mL) was added CF₃CO₂H (1.0 mL) at rt. The reaction mixture was stirred at rt for 2 h, concentrated *in vacuo* to give desired product 13 (43 mg, 90%) which was used directly in the next step without further purification. ESI-MS (M+H)⁺: 419.0.

[0100] Synthesis of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. To a solution of *trans*-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide 13 (42 mg, 0.10 mmol, 1.0 equiv) in 1-butanol (2 mL), 6-chloro-5-fluoropyrimidin-4-amine (18 mg, 0.12 mmol, 1.2 equiv) was added DIPEA (26 mg, 0.20 mmol, 2.0 equiv). The reaction solution was stirred at 120 °C for 16 h. The mixture was diluted with EtOAc (20 mL), washed with H₂O (10 mL) and brine (10 mL), dried (Na₂SO₄), filtered, and concentrated *in vacuo*. The crude was purified by pre-HPLC (MeOH/H₂O with 0.05% TFA as mobile phase) to give the compound (*trans*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide 14 (44 mg, yield: 83%) as a yellow solid. ESI-MS (M+H)⁺: 530.0. HPLC: (214 nm: 100%, 254 nm: 100%). ¹H NMR (400 MHz, CD₃OD) δ: 7.97 (s, 1H), 6.84 (s, 1H), 6.81 (s, 1H), 6.76 (s, 1H), 4.58-4.52 (m, 2H), 4.09-4.03 (m, 1H), 3.52-3.35 (m, 3H), 3.29-3.27 (m, 4H), 3.12-3.05 (m, 1H), 2.24-2.17 (m, 1H), 2.02-1.91 (m, 3H), 1.80-1.63 (m, 2H).




[0101] (3*R*,3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers of compound 14 was separated into three peaks by SFC (IA(2 x 15 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min) and the title compound corresponded to peak 3. LCMS (Agilent460, 254 nm): ES (+) MS m/e = 530.1 (M+1) @ 1.20 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.77 (d, J = 2.01 Hz, 1H), 7.38 (br. s., 1H), 6.94 (s, 2H), 6.75 - 6.87 (m, 2H), 6.41 - 6.66 (m, 3H), 4.29 (br. s., 1H), 4.23

(d, J = 13.05 Hz, 1H), 3.96 - 4.18 (m, 2H), 3.44 (td, J = 6.15, 12.30 Hz, 1H), 3.24 - 3.33 (m, 1H), 3.10 (br. s., 1H), 2.88 (br. s., 1H), 2.82 (t, J = 12.30 Hz, 1H), 2.13 (qd, J = 5.94, 12.30 Hz, 1H), 1.74 - 1.93 (m, 3H), 1.58 - 1.74 (m, 1H), 1.41 - 1.58 (m, 1H).




[0102] (3S,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers of compound **14** was separated into three

peaks by SFC (IA(2 x 15 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min) and the title compound corresponded to peak 2. LCMS (Agilent460, 254 nm): ES (+) MS m/e = 530.1 (M+1) @ 1.19 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, J = 1.76 Hz, 1H), 7.39 (br. s., 1H), 6.98 (s, 1H), 6.96 (s, 1H), 6.72 - 6.88 (m, 2H), 6.57 (s, 2H), 6.54 (d, J = 7.78 Hz, 1H), 4.05 - 4.33 (m, 4H), 3.37 (t, J = 6.27 Hz, 2H), 3.11 (br. s., 1H), 2.94 (br. s., 1H), 2.82 (t, J = 12.30 Hz, 1H), 2.02 - 2.16 (m, 1H), 1.75 - 1.92 (m, 3H), 1.57 - 1.74 (m, 1H), 1.36 - 1.54 (m, 1H).



[0103] (3S,3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers of compound **14** was separated into three

peaks by SFC (IA(2 x 15 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min). Peak 1 of 3 was further purified SFC (AD-H (2 x 15cm), 30% *i*PrOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min) to afford the title compound. LCMS (Agilent 460, 254 nm): ES (+) MS m/e = 530.1 (M+1) @ 1.20 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, J = 1.76 Hz, 1H), 7.38 (br. s., 1H), 6.94 (s, 2H), 6.83 (s, 1H), 6.80 (s, 1H), 6.42 - 6.66 (m, 3H), 4.18 - 4.47 (m, 2H), 3.95 - 4.18 (m, 2H), 3.39 - 3.52 (m, 1H), 3.24 - 3.31 (m, 1H), 3.10 (br. s., 1H), 2.88 (br. s., 1H), 2.82 (t, J = 12.30 Hz, 1H), 2.13 (qd, J = 5.91, 12.39 Hz, 1H), 1.73 - 1.92 (m, 3H), 1.58 - 1.73 (m, 1H), 1.42 - 1.58 (m, 1H).



[0104] (3R,3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-


oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers of compound **14** was separated into three peaks by SFC (IA(2 x 15cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min). Peak 1 of 3 was further purified SFC (AD-H (2 x 15cm), 30% *i*PrOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min) to afford the titled compound. LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 530.1 (M+1) @ 1.20 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, *J*= 1.76 Hz, 1H), 7.39 (br. s., 1H), 6.98 (s, 1H), 6.96 (s, 1H), 6.73 - 6.88 (m, 2H), 6.57 (s, 2H), 6.54 (d, *J*= 7.78 Hz, 1H), 4.05 - 4.35 (m, 4H), 3.37 (t, *J*= 6.15 Hz, 2H), 3.12 (br. s., 1H), 2.94 (br. s., 1H), 2.82 (t, *J*= 12.30 Hz, 1H), 2.09 (sxt, *J*= 5.80 Hz, 1H), 1.74 - 1.92 (m, 3H), 1.56 - 1.73 (m, 1H), 1.36 - 1.52 (m, 1H).

Example 3


Alternative synthesis of (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl) phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide.

[0105] In addition to the methods described in Example 2, (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl) phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide (compound **I-1**) was also synthesized according to Scheme 6.


in EtOH (50 L) under N₂ was added (Boc)₂O (4.2 kg, 19.1 mol, 1.0 equiv), Et₃N (1.9 kg, 19.1 mol, 1.0 equiv) and 10% Pd(OH)₂/C (250 g, 10%w/w). After evacuated and refilled with hydrogenation three times, the mixture was stirred under 1 atm of hydrogen at 50 °C for 15 hr. LC-MS indicated completely consumption of **3-1**. After the mixture was cooled to ambient temperature, the catalyst was filtered through a layer of celite and washed with EtOH (2.5 L). The filtrate was concentrated *in vacuo* to afford crude **3-2** (5.2 kg) as an oil, which was used in next step without further purification.




[0107] (S)-1-tert-butyl 4-ethyl 3-((1-phenylethyl)amino)-5,6-dihydropyridine-1,4 (2H)-dicarboxylate (3-3). To a 100 L reactor equipped with Dean-Stark apparatus was charged toluene (20 L), crude compound **3-2** (5.2 kg, 19.1 mol, 1.0 equiv) rinsed with toluene (30 L), pTSA (329 g, 0.2 mol, 0.01 equiv), and *S*-(-)- α -methylbenzylamine .95 kg, 16.2 mol, 0.85 equiv). The mixture was heated to reflux with a nitrogen blanket and the water was removed through Dean-Stark. After 18 hours, LC-MS indicated complete consumption of **3-2**. The mixture was then cooled to the ambient temperature. The insolubles were removed by filtration, and the filtrate was concentrated *in vacuo* to dryness to afford crude **3-3** as a thick oil. This crude product was used in the next step without further purification. 3-10% of amide byproduct formed in this reaction and its structure was tentatively assigned based on LC-MS data.




[0108] (3R)-1-tert-butyl 4-ethyl 3-(((S)-1-phenylethyl)amino)piperidine-1,4-dicarboxylate (3-4). To a 100 L reactor charged with NaBH₄ (1.16 kg, 30.5 mol, 2.0 equiv) and anhydrous THF (60 L) under nitrogen was added TFA (10.5 kg, 92 mol, 6.0 equiv) slowly over 30 min while maintaining temperature at 0~5 °C. The mixture was then cooled to -45 °C. In a separation reactor, crude product **3-3** was dissolved in anhydrous acetonitrile (30 L), which was added slowly to the above solution of NaBH₄/TFA while maintaining the internal temperature between -45~-30 °C. The mixture was stirred at -45 °C for 1 h, after which time, HPLC indicated complete consumption of compound **3-3**. The mixture was slowly diluted with ice water (50 kg) and the mixture was then warmed to 10 °C. The product was extracted with EtOAc (2 x 40 L) and the combined organic layers were washed with saturated NaHCO₃ solution (20 L). pH of the aqueous was ~8. The organic layers were dried (Na₂SO₄) and concentrated *in vacuo* to nearly dryness to afford a residue which was further azeotroped with MeOH (10 L x 3) to remove excess EtOAc. In the end, a 10 L solution of crude **3-4** in MeOH was obtained, which was used directly in the subsequent step without further purification. ESI-MS (M+H-1)⁺: 377.2. ¹H NMR (400 MHz, CD₃OD) δ 7.31-7.22 (m, 5H), 4.20 (q, 2H), 4.11-3.86 (m, 3H), 3.15 (s, 1H), 3.00-2.90 (m, 2H), 2.64 (d, 2H), 1.87-1.85 (m, 1H), 1.68 (s, 1H), 1.50-1.25 (m, 15H).



[0109] (3R)-1-(tert-butoxycarbonyl)-3-(((S)-1-phenylethyl)amino)piperidine-4-carboxylic acid (3-5). To a 100 L reactor were charged a with THF/MeOH (1:1, 80 L), was added a solution of LiOH·H₂O (2.5 kg, 60 mol, 4.0 equiv) in water (10 L) and a solution of crude 3-4 in MeOH (10 L) from the above step. The resulting mixture was stirred at 22 °C for 18 hours, at which time LC/MS indicated complete consumption of starting material 3-4. The solution was diluted with MTBE (40 L) and stirred for 20 min. The aqueous layer was separated, cooled to 0 °C and neutralized with 3N HCl solution to pH between 7-8, while maintaining the internal temperature below 10 °C. The solution was washed with DCM (5 x 30 L) or until the LC/MS indicated no product 3-5 remained in the aqueous layer. The combined organic layers were concentrated *in vacuo* to dryness, suspended in EtOAc and petroleum ether (2:1, 10 L) and stirred for 2 hours, the solids were filtered, washed by petroleum ether (5 L) and dried under vacuum at 50 °C for 18 hours to give product (3.5 Kg, 53% yield) as a solid with 95% purity. Compound 3-5 is a mixture of ~30:70 trans/cis at C-4 and -93:7 R:S at C-3. The average overall yield from 3-1 is 43-55%. ESI-MS (M+H-1)⁺: 349.2. ¹H NMR (400 MHz, CD₃OD) δ: 8.22-8.06 (m, 5H) 4.11 (m, 1H), 3.86-3.82 (m, 1H), 3.59-3.56 (m, 1H), 2.79-2.65 (m, 1H), 3.22- 2.62 (m, 2H), 2.06 - 2.16 (m, 12H).




[0110] (3R)-1-(6-amino-5-fluoropyrimidin-4-yl)-3-(((S)-1-phenylethyl)amino)piperidine-4-carboxylic acid (3-7). To a 50 L reactor was charged with 10 L of 2N HCl and **3-5** (850 g, 2.44 mol, 1.0 equiv). The mixture was warmed to 30 °C and stirred for 2 hours, at which time HPLC indicted complete consumption of starting **3-5**. The solution was diluted with MTBE (4 L) and stirred for 20 min, layers were separated and to the aqueous layer was added solid K₂CO₃ (660 g) over 1 hour to pH ~7. Additional K₂CO₃ (660 g, 4.8 mol, 2.0 equiv) was added following by 6-chloro-5-fluoropyrimidine-4-ylamine (360 g, 2.44 mole 1.0 equiv) and 1,4-dioxane (5 L). The mixture was heated to gentle reflux at 100 °C and stirred at this temperature for 16 hours. HPLC indicated <2% of compound **3-6** remained. The mixture was washed with DCM (2 x 5 L) and the organic wash solutions were discarded. The aqueous layer was treated with active carbon (425 g) by stirring the slurry for 1 hour at 30 °C followed by filtration through diatomite. This active carbon treatment was repeated. The resulting aqueous solution was neutralized to pH -7 with concentrated HCl, and stirred at 22 °C for 3 hours, the resulting slurry was filtered and the wet cake was washed with washed with 1,4-dioxane/water (1:1, 1.2 L), dried under vacuum at 50 °C for 18 hr until KF ~0.5%. of product **3-7** was obtained as a pale white solid (690 g, 81% yield) with purity of 98.6%. The product contains a mixture of 1:9 cis/trans iomers at the C3 and C4 positions.. ESI-MS (M+H-1)⁺: 460.2. ¹H NMR (400 MHz, CD₃OD) δ: 8.49 (d, J= 2.01 Hz, 1H), 8.21-8.14 (m, 5H) 4.94-4.90 (m, 1H), 4.63 (d, J= 11.55 Hz, 1H), 4.42 (m, 1H), 4.03 (m, 2H), 3.59 - 3.72 (m, 3H), 2.84 - 2.93 (m, 1H), 2.20 - 2.31 (m, 1H), 2.15 (d, J= 6.78 Hz, 3H).




[0111] (3R)-3-amino-1-(6-amino-5-fluoropyrimidin-4-yl)piperidine-4-carboxylic acid (3-8). To a 10 L reaction were charged under N_2 , *i*-PrOH (3.5 L), H_2O (3.5 L), 3-7 (1.0 equiv, 0.97 mol, 350 g), potassium fluoride mono hydrate (290 g, 3.0 eq, 3.0 mol) and 35 g of 20% $Pd(OH)_2/C$ (10% v/w). After evacuated/refilled with hydrogen three times, the mixture was warmed to 40-50 °C and vigorously stirred at that temperature under 1 atmosphere of hydrogen. After 18 hours, LC/MS indicated <1% of starting material 3-7 remained. The mixture was purged with N_2 for 20 min, cooled to 22 °C, and filtered. Both the wet cake and filtrate contained the product and processed separately.

[0112] The filtrate was concentrated *in vacuo* at 50 °C to a volume of ~200 mL. After cooled to 20 °C and stirred at this temperature for 2 hours, a slurry was obtained, and the solid was filtered, washed with water (400 mL) and dried under vacuum and at 50 °C to give product 3-8 (65 g). The wet cake from the reaction filtration was stirred in 1N HCl (1 L) for 2 hours to dissolve the product and the remaining catalyst solid was then removed by filtration. The acidic filtrate was neutralized with solid LiOH to pH ~7 to precipitate the product 3-8. The product was washed with water (200 mL), dried under vacuum and at 50 °C to give 120 g of product. A total of 185 g of product was obtained with 98.7% purity and in 75% yield based on H NMR. All the mother liquors were combined and concentrated to a volume of ~400 mL result in a slurry, filtration, wash with water and drying gave additional 64 g solid with ~50% purity. 1H NMR (400 MHz, D_2O): δ 7.77 (s, 1H) 4.12 (d, J = 14.05 Hz, 1H), 4.01 (d, J = 13.05 Hz, 1H), 3.26 (d, J = 13.80 Hz, 1H), 2.99 - 3.10 (m, 1H), 2.64 - 2.73 (m, 1H), 1.98 (dd, J = 3.39, 14.18 Hz, 1H), 1.74 - 1.87 (m, 1H).



[0113] ((3R,3'R)-1'-(tert-butoxycarbonyl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid (3-10). To a solution of 3-8 (440 g, 1.9 mole, 1.0 equiv) in DMSO (10 L) was added 3-8A sequentially (640 g, 1.9 mole, 1.0 equiv) and sodium triacetoxyborohydride (STAB, 402.0 g, 3.8 mole, 2 equiv) and Et_3N (190 g, 1.9 mol, 1.0 equiv). The mixture was heated to 50 °C and stirred for 3 hours to show complete conversion by HPLC to intermediate 3-9.

[0114] The solution was diluted with MeOH (182 g, 5.7 mol, 3.0 eq.) to quench the excess of STAB, and the reaction was heated to 70-80 °C. After 16 hours, HPLC indicated 22% of product 3-10 formed and 61% intermediate 3-9 remained and chiral HPLC indicated ~3% lactam epimer. The mixture was held at 70-80 °C for additional 24 hours to give 50% 3-10, 35% 3-9, and 7% lactam epimer. After another 40 hours stirring, 80% 3-10 formed, 4% 3-9 remained, and the lactam epimer increased to 14%. The mixture was cooled to 22 °C, and quenched with 2N NH_4Cl solution (5 L) to give a slurry mixture. After 30 minute stirring, the mixture was filtered and the wet cake was washed with water (3 L), dried under vacuum and at 55 °C until KF < 0.1. Crude 3-10 was obtained as a brown solid (850 g, 97.7%); chiral HPLC indicated 12.5% lactam epimer. This product was used directly without further purification.



[0115] (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid (3-10-trans).

To a 10 L reactor was charged under N₂ with **3-10** (850 g, 1.9 mol, 1.0 equiv) in DMF (4.25 L, 5 v/w) to give a clear solution, was added 4-dimethylaminopyridine (DMAP 116g, 0.95 mol, 0.5 equiv) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI, 36.5 g, 0.19 mol, 0.1 equiv). After the mixture was stirred at 15 to 22 °C for about 1 hour, additional EDCI (36.5 g, 0.19 mol, 0.1 equiv) was added and stirred for another 1 hour. HPLC indicated a 69:1 trans/cis mixture. The product **3-10-trans** was not isolated and was converted to compound **I-1** in one-pot. ¹H NMR (300 MHz, DMSO *d*₆): δ 1.47-1.55 (m, 1H), 1.63-1.68 (m, 1H), 1.81-1.87 (m, 1H), 1.90-1.97 (m, 1H), 2.93-3.19 (m, 1H), 3.16-3.23 (m, 1H), 3.33-3.45 (m, 2H) 4.07-4.33 (m, 3H), 6.80 (m, 1H), 6.94-6.98 (m, 1H), 7.10-7.16 (m, 2H), 7.91 (s, 1H).

[0116] (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide (I-1).


To the above reaction mixture, was charged at 22 °C, with O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU, 600 g, 1.9 mol, 1.0 equiv), N,N-diisopropylethylamine (DIPEA, 1.0 kg, 9.5 mol, 5.0 equiv) and finally NH₄Cl (260 g, 5.7 mol, 3.0 equiv). The resulting mixture was stirred at 15 °C for 1 hour, HPLC indicated complete consumption of **3-10-trans**, the mixture was poured into brine (25 L) and extracted with EtOAc (2 x 2L). The combined organics were washed with brine (2 x 2L) and concentrated *in vacuo* below 45 °C to dryness to result in a crude I-1, which was purified by chromatograph with EtOAc/petroleum ether/MeOH (1:1:0 to 50:50:10) to give three fractions, which contained 316 g, 98.8% chemical purity and 10.8% epimer, 160 g, 82.3% chemical purity and 17.5% epimer and 180 g, 61% purity and 11.3% epimer, respectively. The above first two fractions were combined and further purified by prep-HPLC to give 200 g product with >99% purity and <1% epimer. ¹H NMR (400 MHz, DMSO *d*₆): δ 1.48-1.53 (m, 1H), 1.66-1.69 (m, 1H), 1.77-1.79 (m, 3H), 2.11-2.16 (m, 1H), 2.80-2.88 (m, 2H), 3.11 (s, 1H), 3.42-3.48 (m, 1H), 4.0-4.25 (m, 4H), 6.58 (s, 3H), 6.80-6.85 (d, *J* = 10.2, 2H), 6.95 (s, 2H), 7.40 (s, 1H), 7.77 (s, 1H).

Example 4


Alternative syntheses of (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide.

[0117] In addition to the methods described in Examples 2 and 3, (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide (compound I-1) was also synthesized according to Scheme 6.

Scheme 7




[0118] (3R)-3-amino-1-(*tert*-butoxycarbonyl)piperidine-4-carboxylic acid (4-L-6). To a 10 L reactor charged under nitrogen with compound 4-5 (100 g, 0.287 mole), MeOH (6 L, 60 v/w) and 10 g 20% Pd(OH)₂/C. The reactor was evacuated/refilled with hydrogen three times and the mixture was warmed to 40-50 °C while stirring under 3 Mpa of hydrogen for 40 hours. LC/MS indicated complete consumption of starting material 4-5. The mixture was cooled to 22 °C and filtered, and the filtrate was concentrated *in vacuo* to dryness to afford a solid product. This crude product was slurried in EtOH (500 mL) at 22 °C for 2 hours, filtered and dried under vacuum at 50 °C to afford a 85% yield of product 4-L-6 (60 g, 0.245 mole) as a white solid.




[0119] (3R)-1-(tert-butoxycarbonyl)-3-(((R)-4-((3-chloro-5-(trifluoromethyl)phenyl)amino)-5-ethoxy-5-oxopentyl)amino)piperidine-4-carboxylic acid (4-L-8). To a solution of 4-L-6 (48.4 g, 0.197 mole) in DMSO (450 mL) was added Et₃N (20.2 g, 0.199 mole, 1 equiv), 3-8A (67.4 g, 0.199 mole, 1 equiv) and sodium triacetoxyborohydride (STAB, 84.8 g, 0.40 mole, 2.0 equiv). The mixture was heated to 50 °C over 30 min and stirred at that temperature for 3 hours. LC/MS indicated consumption of most of starting material 4-L-6 and formation of 4-L-8.


[0120] The reaction was quenched by adding EtOH (35 mL) and stirring at 50 °C for 30 min. The mixture was heated at 75-85 °C for 3 days. The mixture was cooled to 18 °C and transferred slowly into water (6 L) while vigorously stirring to afford a slurry. After 2 hours, the solids were filtered and washed with water (3 x 3 L), dried under vacuum at 60-70 °C for 24 hours to give 4-L-9 (114 g) as a brown solid. The solid was used directly in the subsequent step.



[0121] (3R,3'R,4'S)-1'-(tert-butoxycarbonyl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid (4-L-9-trans). To a solution of crude 4-L-9 (100 g), in DMF (500 mL) was added 4-dimethylaminopyridine (11 g, 0.09 mole, 0.5 equiv) and stirred at 20 °C for 10 min. 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (7.0 g, 0.036 mole, 0.2 equiv) was added and the reaction was stirred 20 °C for 3 hours. HPLC indicated a ratio of 57:43 cis/trans mixture and additional EDAC (3.5 g, 0.018 mole, 0.1 equiv) was added. After 5 hours, HPLC indicated complete conversion to 4-L-9-trans. The mixture was transferred to water (2.25 L) slowly and the mixture was extracted with EtOAc (2 x 500 mL), and the organic layers were washed with brine (500 mL) and water (500 mL), concentrated *in vacuo* to dryness to give crude 4-L-9-trans (100 g) as a brown solid. The crude was dissolved in EtOAc (135 mL) at 60 °C and then cooled to 20 °C over 1 hour followed by adding 50 mL petroleum ether. The mixture was aged for 2 hours. The solids were filtered and washed with 3:1 EtOAc/petroleum ether (50 mL), dried under vacuum at 50 °C for 16 hours to give 4-L-9-trans (23 g, 22% yield with 99% purity). ¹H NMR (400 MHz, DMSO-*d*₆) δ 6.94 (s, 2H), 6.81 (s, 1H), 6.54 - 6.61 (m, 1H), 3.99 - 4.08 (m, 1H), 3.42 - 3.38 (m, 2H), 2.07 - 2.16 (m, 1H), 1.74 - 1.92 (m, 3H), 1.39 (s, 9H).




[0122] (3R,3'R,4'S)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid hydrochloride (4-L-10 trans). To a solution of 0.5N HCl in EtOAc (76 mL) was added 4-L-9 trans (20 g, 38 mmol) and heated at 20 °C for 18 h to give a slurry. The solid was filtered, washed with EtOAc (5 mL) and dried under vacuum at 45 °C for 18 h to afford 4-L-10 as the HCl salt (17 g, 97% yield).



[0123] (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid (**3-10-trans**). A solution of **4-L-10** (2.0 g, 4.38 mole), 6-chloro-5-fluoro-pyrimidin-4-ylamine (711 mg, 4.82 mmole, 1.1 equiv), DIPEA (1.52 mL, 8.77 mole, 2 eq.) in 40 mL nBuOH was heated to 130-140 °C for 72 h. The mixture was cooled to 22 °C and *concentrated in vacuo* to afford a residue which purified by column to give **3-10-trans** (1.1 g, 47%). A relatively minor amount of epimer (3S,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid was also observed. Intermediate **3-10-trans** can be converted to compound **I-1** via procedure described above.

[0124] Compound **I-1** was also synthesized according to Scheme 8.

Scheme 8



[0125] The synthesis of (*R*)-*tert*-butyl 1-(3-chloro-5-(trifluoromethyl)phenyl)-5-oxopyrrolidine-2-carboxylate **4-e** was synthesized using a similar procedure as in Phillips, D. P.; Zhu, X. -F.; Lau, T. L.; Yang, K.; Liu, H. Tetrahedron Letters, 2009, 50, 7293, whereby the (*S*)-methyl 5-oxopyrrolidine-2-carboxylate and 1-chloro-4-iodobenzene were substituted for the (*R*)-*tert*-butyl 5-oxopyrrolidine-2-carboxylate and 1-chloro-3-iodo-5-(trifluoromethyl)benzene.

[0126] **(2R)-tert-butyl 1-(3-chloro-5-(trifluoromethyl)phenyl)-5-hydroxypyrrolidine-2-carboxylate.** An anhydrous solution of **4-e** (11 g, 30 mmol) in Me-THF (100 mL) was cooled to -35 °C under an atmosphere of nitrogen. A solution of DIABL-H (5.9 g, 42 mmol) in toluene (42 mL) was added dropwise while maintaining the temperature at -35 °C. The reaction was monitored by HPLC and upon completion a solution of 1N Rochell salt (100 mL) was added while maintaining the reaction temperature below 0 °C. The organic phase was separated, washed with 1N Rochell salt (50 mL x 3) and separated, diluted with Et₃N (4 mL), dried (Na₂SO₄) and concentrated *in vacuo* to afford **4-f** (8.3 g) as an oil.

[0127] **(3R)-1-(6-amino-5-fluoropyrimidin-4-yl)-3-(((R)-5-(tert-butoxy)-4-((3-chloro-5-(trifluoromethyl)phenyl)amino)-5-oxopentyl)amino)piperidine-4-carboxylic acid.** A solution of **4-f** (37.4 g, 0.146 mmol) in DMF (700 mL) was treated with **3-8** (40.2 g, 0.11 mmol), Et₃N (10.1 g, 0.1 mmol) STAB (42.4 g, 0.2 mmol) and the mixture was heated to 55 °C for 5 h. The reaction was diluted with water (2.5 L), extracted with EtOAc (500 mL x 3), the organic phases were combined and washed with brine, separated, dried (Na₂SO₄) and concentrated *in vacuo* to afford **4-a** (40.2 g) as solid which was used without any additional purification.

[0128] **(3R)-1-(6-amino-5-fluoropyrimidin-4-yl)-3-(((R)-4-carboxy-4-((3-chloro-5-(trifluoromethyl)phenyl)amino)butyl)amino)piperidine-4-carboxylic acid.** To a solution of 5 N HCl (250 mL) was added *t*-butyl ester **4-a** and the suspension was heated to 55 °C for 5 h while the hydrolysis was monitored by HPLC. Upon complete formation of the product the water was removed *in vacuo* resulting in a solid **4-b** which was dried under vacuum and used without any additional purification.


[0129] **(3R,3'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxylic acid.** To a solution of acid **4-b** (55 g, 0.1 mol) in DMF (500 mL) was added a DIEA (64.5 g, 0.5 mol), CDI (32.5 g, 0.2 mol) at 0 °C. The solution was stirred for 1.5 h at 0 °C, diluted with water (3 L), adjusted to a pH 3 with HCl and extracted with EtOAc (2 L x 3). The organic phase were combined, dried (Na₂SO₄) and concentrated *in vacuo* to afford **4-c** (48 g).

[0130] The remaining steps to compound **1-1** are completed via procedures described above.

Example 5


Synthesis of *trans*-tert-butyl 3-((3-chloro-5-(trifluoromethyl)phenyl) amino)-4'-(methylcarbamoyl)-2-oxo-[1,3'-bipiperidine]-1'-carboxylate.

[0131]




[0132] **Synthesis of *trans*-tert-butyl 3-((3-chloro-5-(trifluoromethyl)phenyl) amino)-4'-(methylcarbamoyl)-2-oxo-[1,3'-bipiperidine]-1'-carboxylate.** A similar procedure was used as described for the synthesis of (3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide **14** to afford the crude material which was purified by pre-HPLC (MeOH/H₂O with 0.05% NH₃·H₂O as mobile phase) to give the title compound (360 mg, yield: 67%) as a yellow solid. ESI-MS (M+H)⁺: 544.18. HPLC: (214 nm: 100.0%, 254 nm:


100.0%). ^1H NMR (400 MHz, CD_3OD) (mixture of isomers) δ 7.69-7.68 (m, 1H), 6.78 (s, 1H), 6.75 (s, 1H), 6.71 (s, 1H), 4.39-4.36 (m, 2H), 4.09-4.03 (m, 1H), 3.53-3.31 (m, 3H), 3.20-3.10 (m, 1H), 2.99-2.92 (m, 1H), 2.55 (s, 3H), 2.28-2.19 (m, 1H), 1.96-1.77 (m, 5H), 1.68-1.58 (m, 1H).




[0133] (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N-methyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into two peaks by SFC (AD-H (2 x 15 cm), 50% 1:1 IPA:methanol (0.1% DEA)/ CO_2 , 100 bar, 60 ml/min). Peak 2 was further purified by SFC separation (AD-H (2 x 15 cm), 30% *i*PrOH(0.15% DEA)/ CO_2 , 100 bar, 60 ml/min) to afford the title compound. LCMS (Agilent 460, 254 nm): ES (+) MS m/e = 544.1 (M+1) @ 1.24 min. ^1H NMR (400 MHz, DMSO-*d*₆) δ 7.72 - 7.85 (m, 2H), 6.92 (s, 2H), 6.81 (s, 1H), 6.43 - 6.64 (m, 3H), 4.34 (br. s., 1H), 4.23 (d, J = 13.05 Hz, 1H), 3.93 - 4.19 (m, 2H), 3.37 - 3.49 (m, 1H), 3.22 - 3.30 (m, 1H), 3.13 (br. s., 1H), 2.84 (t, J = 12.05 Hz, 2H), 2.57 (d, J = 4.52 Hz, 3H), 2.13 (qd, J = 6.05, 12.45 Hz, 1H), 1.60 - 1.89 (m, 4H), 1.40 - 1.58 (m, 1H).



[0134] (3S,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N-methyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into two peaks by SFC (AD-H (2 x 15 cm), 50% 1:1 IPA:methanol (0.1% DEA)/ CO_2 , 100 bar, 60 ml/min). Peak 2 was further purified by SFC separation (AD-H (2 x 15 cm), 30% *i*PrOH (0.15% DEA)/ CO_2 , 100 bar, 60 ml/min) to afford the title compound. LCMS (Agilent 460, 254 nm): ES (+) MS m/e = 544.1 (M+1) @ 1.24 min. ^1H NMR (400 MHz, DMSO-*d*₆) δ 7.83 (q, J = 4.60 Hz, 1H), 7.77 (d, J = 1.76 Hz, 1H), 6.97 (d, J = 6.78 Hz, 2H), 6.81 (s, 1H), 6.58 (s, 2H), 6.53 (d, J = 7.78 Hz, 1H), 4.23 (d, J = 13.05 Hz, 2H), 3.90 - 4.19 (m, 2H), 3.14 (br. s., 1H), 2.92 (br. s., 1H), 2.74 - 2.90 (m, 1H), 2.55 (d, J = 4.52 Hz, 3H), 2.00 - 2.18 (m, 1H), 1.74 - 1.89 (m, 3H), 1.56 - 1.74 (m, 1H), 1.34 - 1.50 (m, 1H).




[0135] (3*S*,3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N*-methyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into two peaks by SFC (AD-H (2 x 15 cm), 50% 1:1 IPA:methanol (0.1% DEA)/CO₂, 100 bar, 60 ml/min). Peak 1 was further purified by SFC (AD-H (2 x 15cm), 30% MeOH (0.15% DEA)/CO₂, 100 bar, 60 ml/min) to afford the titled compound. LCMS (Agilent 460, 254nm): ES (+) MS *m/e* = 544.1 (M+1) @ 1.23 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.72 - 7.85 (m, 2H), 6.92 (s, 2H), 6.81 (s, 1H), 6.57 (s, 2H), 6.54 (d, *J* = 7.53 Hz, 1H), 4.23 (d, *J* = 13.30 Hz, 1H), 4.15 (dd, *J* = 3.26, 12.30 Hz, 1H), 4.08 (td, *J* = 7.06, 10.48 Hz, 1H), 3.36 - 3.47 (m, 1H), 3.23 - 3.30 (m, 1H), 3.13 (br. s., 1H), 2.84 (t, *J* = 11.80 Hz, 2H), 2.57 (d, *J* = 4.52 Hz, 3H), 2.13 (qd, *J* = 6.17, 12.61 Hz, 1H), 1.62 - 1.91 (m, 4H), 1.42 - 1.57 (m, 1H).



[0136] *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N*-methyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into two peaks by SFC (AD-H (2 x 15 cm), 50% 1:1 IPA:methanol (0.1% DEA)/CO₂, 100 bar, 60 ml/min). Peak 1 was further purified by SFC (AD-H (2 x 15 cm), 30% MeOH (0.15% DEA)/CO₂, 100 bar, 60 ml/min) to afford the titled cmpd. LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 544.1 (M+1) @ 1.23 min. LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 544.1 (M+1) @ 1.24 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.80 - 7.89 (m, 1H), 7.72 - 7.80 (m, 1H), 6.97 (d, *J* = 6.53 Hz, 2H), 6.81 (s, 1H), 6.58 (s, 2H), 6.53 (d, *J* = 7.78 Hz, 1H), 4.23 (d, *J* = 13.05 Hz, 2H), 3.89 - 4.19 (m, 2H), 3.13 (br. s., 1H), 2.74 - 3.02 (m, *J* = 12.42, 12.42 Hz, 2H), 2.55 (d, *J* = 4.52 Hz, 3H), 2.08 (qd, *J* = 5.97, 12.20 Hz, 1H), 1.81 (td, *J* = 6.24, 12.36 Hz, 3H), 1.56 - 1.74 (m, 1H), 1.33 - 1.51 (m, 1H).


Example 6

[0137]




[0138] Synthesis of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-

no)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. A similar procedure was used as described for the synthesis of (3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide 14 to afford the crude material which was purified by pre-HPLC (MeOH/H₂O with 0.05% TFA as mobile phase) to give the title compound (45 mg, yield: 90%) as a yellow solid. ESI-MS (M+H)⁺: 558.0. HPLC: (214 nm: 98.2%, 254 nm: 100.0%). ¹H NMR (400 MHz, CD₃OD) δ : 7.77 (s, 1H), 6.92-6.89 (m, 1H), 6.83-6.81 (m, 1H), 6.79 (s, 1H), 4.38-4.38 (m, 2H), 4.05-4.00 (m, 2H), 3.55-3.53 (m, 1H), 3.45-3.40 (m, 2H), 3.15-2.89 (m, 7H), 2.21-2.16 (m, 1H), 1.90-1.86 (m, 3H), 1.66-1.56 (m, 2H).




[0139] (3*R*,3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The title compound was obtained from chiral separation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide using a two step chiral SFC separation. Firstly, the mixture was separated into two peaks containing a mixture of two diastereomers ((3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide and (3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide) using a ChiralPak IC(2 x 15 cm, 30% methanol w/0.1 DEA) column, and then the resulting mixture containing a pair of isomers was further separated into the single enantiomers using a ChiralPak IA (2 x 15 cm, 30% methanol w/0.1% DEA 100 bar) column. ESI-MS (M+H)⁺: 558.0 ¹H NMR (400 MHz, CDCl₃) δ : 7.92 (d, *J* = 1.76 Hz, 1H), 6.94 (s, 1H), 6.72 (d, *J* = 7.78 Hz, 2H), 5.21 (d, *J* = 3.51 Hz, 1H), 4.70 (s, 2H), 4.45 (dd, *J* = 2.76, 12.80 Hz, 2H), 4.17 - 4.32 (m, 1H), 3.64 - 3.80 (m, 2H), 3.44 - 3.58 (s, 3H), 3.09 (s, 3H), 3.00 - 3.09 (m, 1H), 2.95 (s, 3H), 2.40 (dd, *J* = 5.52, 13.30 Hz, 1H), 1.63 - 1.99 (m, 3H), 1.26 - 1.43 (m, 1H).




[0140] (3*S*,3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The title compound was obtained from chiral separation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide using a two step chiral SFC separation. Firstly, the mixture was separated into two peaks containing a mixture of two diastereomers ((3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide and (3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide) using a ChiralPak IC(2 x 15 cm, 30% methanol w/0.1 DEA) column, and then the resulting mixture containing a pair of isomers was further separated into the single enantiomers using a ChiralPak IA (2 x 15 cm, 30% methanol w/0.1% DEA 100 bar) column. ESI-MS (M+H)⁺: 558.0 ¹H NMR (400 MHz, CDCl₃) δ : 7.93 (d, *J* = 1.26 Hz, 1H), 6.93 (s, 1H), 6.69 (br. s., 2H), 5.06 (d, *J* = 4.27 Hz, 1H), 4.71 (s, 1H), 4.45 (d, *J* = 12.55 Hz, 2H), 4.16 - 4.26 (m, 1H), 3.66 - 3.81 (m, 2H), 3.54 - 3.64 (m, 1H), 3.40 - 3.54 (m, 2H), 3.01 - 3.10 (m, 4H), 2.95 (s, 3H), 2.37 (dd, *J* = 5.27, 13.05 Hz, 1H), 1.91 - 2.02

(m, 2H), 1.48 - 1.75 (m, 2H).




[0141] (3S,3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The title compound was obtained from chiral separation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide using a two step chiral SFC separation. Firstly, the mixture was separated into two peaks containing a mixture of two diastereomers ((3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide and (3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide) using a ChiralPak IC(2 x 15 cm, 30% methanol w/0.1 DEA) column, and then the resulting mixture containing a pair of isomers was further separated into the single enantiomers using a ChiralPak IA (2 x 15 cm, 30% methanol w/0.1% DEA 100 bar) column. ESI-MS (M+H)⁺: 558.0 ¹H NMR (400 MHz, CDCl₃) δ : 7.92 (d, *J* = 1.76 Hz, 1H), 6.94 (s, 1H), 6.72 (d, *J* = 7.78 Hz, 2H), 5.21 (d, *J* = 3.51 Hz, 1H), 4.70 (s, 2H), 4.45 (dd, *J* = 2.76, 12.80 Hz, 2H), 4.19 - 4.30 (m, 1H), 3.66 - 3.79 (m, 2H), 3.47 - 3.56 (m, 3H), 3.09 (s, 3H), 2.97 - 3.07 (m, 1H), 2.95 (s, 3H), 2.40 (dd, *J* = 5.52, 13.30 Hz, 1H), 1.81 - 1.97 (m, 3H), 1.73 (dd, *J* = 3.76, 12.80 Hz, 1H).




[0142] (3R,3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide The title compound was obtained from chiral separation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl) amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide using a two step chiral SFC separation. Firstly, the mixture was separated into two peaks containing a mixture of two diastereomers ((3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide and (3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-N,N-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide) using a ChiralPak IC(2 x 15 cm, 30% methanol w/0.1 DEA) column, and then each mixture containing a pair of isomers was further separated into the single enantiomers using a ChiralPak IA (2 x 15 cm, 30% methanol w/0.1% DEA 100 bar) column. ESI-MS (M+H)⁺: 558.0 ¹H NMR (400 MHz, CDCl₃) δ : 7.93 (d, *J* = 1.26 Hz, 1H), 6.93 (s, 1H), 6.69 (br. s., 2H), 5.06 (d, *J* = 4.27 Hz, 1H), 4.71 (s, 1H), 4.45 (d, *J* = 12.55 Hz, 2H), 4.16 - 4.26 (m, 1H), 3.66 - 3.81 (m, 2H), 3.54 - 3.64 (m, 1H), 3.40 - 3.54 (m, 2H), 3.01 - 3.10 (m, 4H), 2.95 (s, 3H), 2.37 (dd, *J* = 5.27, 13.05 Hz, 1H), 1.91 - 2.02 (m, 2H), 1.48 - 1.75 (m, 2H).

Example 7


[0143]




[0144] **Synthesis of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-4'-(4-methylpiperazine-1-carbonyl)-[1,3'-bipiperidin]-2-one.** A similar procedure was used as described for the synthesis of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide 14 to afford 23 which was purified by reverse phase HPLC (MeOH/H₂O with 0.05% NH₃·H₂O as mobile phase) to afford the title compound (100 mg, yield: 70%) as a yellow solid. ESI-MS (M+H)⁺: 613.24. ¹H NMR (400 MHz, CDCl₃) δ: 7.94 (s, 1H), 6.94 (s, 1H), 6.73-6.67 (m, 2H), 5.25-5.03 (m, 1H), 4.71 (s, 2H), 4.49-4.40 (m, 2H), 4.35-4.16 (m, 1H), 3.82-3.64 (m, 3H), 3.62-3.41 (m, 6H), 3.08-2.97 (m, 1H), 2.52-2.34 (m, 3H), 2.30-2.25 (m, 2H), 2.20 (s, 3H), 1.85-1.64 (m, 2H), 1.72-1.64 (m, 2H), 1.49-1.31 (m, 1H).




[0145] **(3*R*,3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-4'-(4-methylpiperazine-1-carbonyl)-[1,3'-bipiperidin]-2-one.** The title compound was obtained from chiral separation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide 23 using a two step chiral SFC separation. Firstly, the mixture was separated into two peaks containing a mixture of two diastereomers ((3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide and (3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide) using a ChiralPak IC (2 x 15 cm, 30% methanol w/0.1 DEA) column, and then the resulting mixture containing a pair of isomers was further separated into the single enantiomers using a ChiralPak IA (2 x 15 cm, 30% methanol w/0.1% DEA 100 bar) column. ESI-MS (M+H)⁺: 613.2 ¹H NMR (400 MHz, CD₃OD) δ: 7.95 (s, 1H), 6.83 - 6.97 (m, 3H), 4.50 - 4.68 (m, 2H), 4.28 - 4.40 (m, 1H), 3.66 - 3.91 (m, 8H), 3.30 - 3.52 (m, 4H), 3.14 (t, *J* = 12.42 Hz, 2H), 2.73 (br. s., 2H), 2.27 (dd, *J* = 5.65, 12.93 Hz, 1H), 1.83 - 2.05 (m, 2H), 1.54 - 1.79 (m, 2H).



[0146] (3S,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-4'-(4-methylpiperazine-1-carbonyl)-[1,3'-bipiperidin]-2-one. The title compound was obtained from chiral separation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl) amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide **23** using a two step chiral SFC separation. Firstly, the mixture was separated into two peaks containing a mixture of two diastereomers ((3'*R*,4'5)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide and (3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide) using a ChiralPak IC(2 x 15 cm, 30% methanol w/0.1 DEA) column, and then the resulting mixture containing a pair of isomers was further separated into the single enantiomers using a ChiralPak IA (2 x 15 cm, 30% methanol w/0.1% DEA 100 bar) column. ESI-MS (M+H)⁺: 613.2. ¹H NMR (400 MHz, CD₃OD) δ : 7.91 - 7.99 (m, 1H), 6.84 - 6.97 (m, 3H), 4.57 (dd, *J* = 14.18, 19.70 Hz, 2H), 4.34 (br. s., 1H), 3.42 - 3.53 (m, 1H), 3.37 (d, *J* = 1.51 Hz, 3H), 3.05 - 3.19 (m, 1H), 2.86 (t, *J* = 7.53 Hz, 1H), 2.68 - 2.78 (m, 2H), 2.26 (dd, *J* = 5.65, 12.93 Hz, 1H), 1.82 - 2.03 (m, 3H), 1.55 - 1.79 (m, 2H).




[0147] (3S,3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-4'-(4-methylpiperazine-1-carbonyl)-[1,3'-bipiperidin]-2-one. The title compound was obtained from chiral separation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl) amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide **23** using a two step chiral SFC separation. Firstly, the mixture was separated into two peaks containing a mixture of two diastereomers ((3'*R*,4'*R*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide and (3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide) using a ChiralPak IC(2 x 15 cm, 30% methanol w/0.1 DEA) column, and then the resulting mixture containing a pair of isomers was further separated into the single enantiomers using a ChiralPak IA (2 x 15 cm, 30% methanol w/0.1% DEA 100 bar) column. ESI-MS (M+H)⁺: 613.2. ¹H NMR (400 MHz, CD₃OD) δ : 7.94 (s, 1H), 6.87 - 6.93 (m, 3H), 4.57 (dd, *J* = 14.18, 19.70 Hz, 1H), 4.34 (br. s., 1H), 3.43 - 3.51 (m, 1H), 3.36 - 3.38 (m, 2H), 3.13 (t, *J* = 12.30 Hz, 1H), 2.86 (t, *J* = 7.53 Hz, 1H), 2.73 (br. s., 1H), 2.26 (dd, *J* = 5.65, 12.93 Hz, 1H), 1.96 - 2.03 (m, 1H), 1.83 - 1.94 (m, 1H), 1.51 - 1.75 (m, 1H).




[0148] ((3R,3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-4'-(4-methylpiperazine-1-carbonyl)-[1,3'-bipiperidin]-2-one. The title compound was obtained from chiral separation of *trans*-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl) amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide **23** using a two step chiral SFC separation. Firstly, the mixture was separated into two peaks containing a mixture of two diastereomers ((3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide and (3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-*N,N*-dimethyl-2-oxo-[1,3'-bipiperidine]-4'-carboxamide) using a ChiralPak IC(2 x 15 cm, 30% methanol w/0.1 DEA) column, and then each mixture containing a pair of isomers was further separated into the single enantiomers using a ChiralPak IA (2 x 15 cm, 30% methanol w/0.1% DEA 100 bar) column. ¹H NMR (400 MHz, CD₃OD) δ : 7.96 (br. s., 1H), 6.90 (br. s., 1H), 6.76 (d, *J* = 10.04 Hz, 2H), 4.60 (t, *J* = 14.06 Hz, 2H), 4.19 - 4.32 (m, 1H), 3.67 - 3.78 (m, 1H), 3.43 - 3.54 (m, 3H), 3.35 - 3.38 (m, 3H), 3.16 (t, *J* = 12.42 Hz, 1H), 2.86 (t, *J* = 7.40 Hz, 2H), 2.79 (s, 3H), 2.32 (dd, *J* = 5.02, 12.80 Hz, 1H), 1.89 - 2.07 (m, 4H), 1.62 - 1.76 (m, 5H),

Example 8

[0149]




[0150] The synthesis of 1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-fluoro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. A similar procedure was used as described for the synthesis of (3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide **14** to afford the crude material which was purified by pre-HPLC (MeOH/H₂O with 0.05% NH₃.H₂O as mobile phase) to give the title compound (175 mg, yield: 69%) as a yellow solid. ESI-MS (M+H)⁺: 514.19. HPLC: (214 nm: 96.13%, 254 nm: 96.53%). ¹H NMR (400 MHz, CD₃OD) δ : 7.79-7.78 (m, 1H), 6.76 (s, 1H), 6.63-6.54 (m, 2H), 4.41-4.36 (m, 2H), 4.08-4.06 (m, 1H), 3.55-3.41 (m, 3H), 3.28-3.25 (m, 1H), 2.99-2.93 (m, 1H), 2.28-2.21 (m, 1H), 1.98-1.78 (m, 6H).




[0151] (3R,3R,4S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-fluoro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into three peaks by SFC

(IA (3 x 15 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 70 ml/min) and peak 3 corresponded to the titled compound. LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 514.0 (M+1) @ 1.09 min. ¹H NMR (400 MHz, CDCl₃) δ : 7.91 (br. s., 1H), 6.66 (d, *J* = 8.53 Hz, 1H), 6.63 (s, 1H), 6.46 (d, *J* = 10.79 Hz, 1H), 6.12 (br. s., 1H), 5.47 (br. s., 1H), 5.16 (d, *J* = 3.51 Hz, 1H), 4.91 (br. s., 2H), 4.35 - 4.54 (m, 2H), 3.82 (td, *J* = 5.11, 10.60 Hz, 2H), 3.51 - 3.60 (m, 1H), 3.34 - 3.48 (m, 3H), 2.96 (t, *J* = 12.30 Hz, 1H), 2.35 - 2.47 (m, 1H), 1.91 - 2.06 (m, 3H), 1.84 (dq, *J* = 3.89, 12.76 Hz, 1H), 1.48 - 1.62 (m, 1H).




[0152] (3*S*,3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-fluoro-5-(trifluoromethyl)phenyl)amino)-2-

oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into three peaks by SFC (IA (3 x 15 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 70 ml/min). Peak 2 of 3 corresponded to desired compound. LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 514.0 (M+1) @ 1.10 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, *J* = 1.76 Hz, 1H), 7.39 (s, 1H), 6.85 (s, 1H), 6.81 (s, 1H), 6.74 (d, *J* = 12.30 Hz, 1H), 6.47 - 6.66 (m, 4H), 4.23 (d, *J* = 12.80 Hz, 2H), 3.90 - 4.18 (m, 2H), 3.34 - 3.46 (m, 2H), 3.12 (br. s., 1H), 2.94 (br. s., 1H), 2.82 (t, *J* = 12.42 Hz, 1H), 2.10 (qd, *J* = 5.75, 12.11 Hz, 1H), 1.74 - 1.92 (m, 3H), 1.56 - 1.72 (m, 1H), 1.37 - 1.52 (m, 1H).



[0153] (3*R*,3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-fluoro-5-(trifluoromethyl)phenyl)amino)-2-


oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into three peaks by SFC (IA (3 x 15 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 70 ml/min). Peak 1 of 3 was further purified by SFC (IA (3 x 15 cm), 30% *i*PrOH (0.1% DEA)/CO₂, 100 bar, 70 ml/min) to afford the titled compound. LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 514.0 (M+1) @ 1.10 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, 1.5 Hz, 1H), 7.39 (s., 1H), 6.85 (s, 1H), 6.81 (s., 1H), 6.74 (d, *J* = 12.30 Hz, 1H), 6.47 - 6.66 (m, 4H), 4.16 - 4.46 (m, 2H), 3.95 - 4.16 (m, 2H), 3.34 - 3.48 (m, 2H), 3.12 (br. s., 1H), 2.87 - 3.01 (m, 2H), 2.82 (t, *J* = 12.30 Hz, 1H), 2.10 (qd, *J* = 5.75, 12.11 Hz, 1H), 1.74 - 1.92 (m, 3H), 1.54 - 1.74 (m, 1H), 1.35 - 1.52 (m, 1H).




[0154] (3*S*,3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-fluoro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide.

The mixture of four diastereomers was separated into three peaks by SFC (IA (3 x 15 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 70 ml/min). Peak 1 of 3 was further purified by SFC (IA (3 x 15cm), 30% *i*PrOH (0.1% DEA)/CO₂, 100 bar, 70 ml/min) to afford the titled compound.


LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 514.0 (M+1) @ 1.10 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, *J* = 1.76 Hz, 1H), 7.38 (br. s., 1H), 6.84 (s, 2H), 6.70 (d, *J* = 12.30 Hz, 1H), 6.47 - 6.65 (m, 4H), 4.18 - 4.48 (m, 2H), 3.92 - 4.18 (m, 2H), 3.38 - 3.49 (m, 1H), 3.20 - 3.30 (m, 1H), 3.11 (br. s., 1H), 2.88 - 2.99 (m, 1H), 2.83 (t, *J* = 12.30 Hz, 1H), 2.14 (qd, *J* = 6.03, 12.52 Hz, 1H), 1.74 - 1.92 (m, 3H), 1.59 - 1.74 (m, 1H), 1.41 - 1.58 (m, 1H).

Example 9**[0155]**


[0156] The synthesis of (3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-fluorophenyl) amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. A similar procedure was used as described for the synthesis of (3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide 14 to afford the crude material which was purified by pre-HPLC (MeOH/H₂O with 0.05% NH₃·H₂O as mobile phase) to give the title compound (141 mg, Y: 30%) as a white solid. ESI-MS (M+H)⁺: 479.9. HPLC: (214 nm: 100%, 254 nm: 100%). ¹H NMR (400 MHz, DMSO *d*₆) δ : 7.78-7.77 (m, 1H), 7.40-7.38 (m, 1H), 6.86-6.82 (m, 1H), 6.62-6.55 (m, 3H), 6.45-6.37 (m, 3H), 4.26-3.94 (m, 4H), 3.47-3.39 (m, 1H), 3.20-3.03 (m, 2H), 2.90-2.78 (m, 2H), 2.18-2.04 (m, 1H), 1.86-1.34 (m, 5H).



[0157] (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-fluorophenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into three peaks by SFC (IC (2 x 15 cm), 25% MeOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min) to afford the title compound as peak 3 respectively. LCMS (Agilent 460, 254nm): ES (+) MS m/e = 480.0 (M+1) @ 1.01 min. ¹H NMR (400 MHz, CDCl₃) δ: 7.90 (br. s., 1H), 6.44 (d, J = 8.53 Hz, 1H), 6.40 (s, 1H), 6.30 (br. s., 1H), 6.23 (d, J = 11.04 Hz, 1H), 5.63 (br. s., 1H), 5.09 (br. s., 1H), 4.94 (br. s., 2H), 4.45 (d, J = 12.80 Hz, 2H), 3.68 - 3.95 (m, 2H), 3.49 - 3.56 (m, 2H), 3.35 - 3.46 (m, 2H), 2.89 - 2.99 (m, 1H), 2.31 - 2.44 (m, 1H), 1.90 - 2.04 (m, 3H), 1.76 - 1.89 (m, 1H), 1.49 - 1.61 (m, 1H).




[0158] (3S,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-fluorophenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into three peaks by SFC (IC (2 x 15 cm), 25% MeOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min) to afford the title compound as peak 1 respectively. LCMS (Agilent 460, 254nm): ES (+) MS m/e = 480.0 (M+1) @ 1.01 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.77 (d, J = 1.76 Hz, 1H), 7.39 (s., 1H), 6.81 (s, 1H), 6.57 (s, 3H), 6.46 (d, J = 12.30 Hz, 1H), 6.39 (dd, J = 1.76, 8.78 Hz, 1H), 6.34 (d, J = 7.53 Hz, 1H), 4.28 (br. s., 1H), 4.23 (d, J = 13.05 Hz, 1H), 4.13 (dd, J = 2.76, 12.30 Hz, 1H), 4.03 (td, J = 6.56, 10.98 Hz, 1H), 3.33 - 3.46 (m, 2H), 3.11 (br. s., 1H), 2.94 (br. s., 1H), 2.82 (t, J = 12.30 Hz, 1H), 2.09 (qd, J = 5.75, 12.11 Hz, 1H), 1.72 - 1.92 (m, 3H), 1.56 - 1.72 (m, 1H), 1.29 - 1.50 (m, 1H).



[0159] (3S,3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-fluorophenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into three peaks by SFC (IC (2 x 15 cm), 25% MeOH(0.1% DEA)/CO₂, 100 bar, 60 ml/min). Peak 2 of 3 was further purified by SFC (IA (3 x 15 cm), 30% iPrOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min) to afford the title compound. LCMS (Agilent460, 254 nm): ES (+) MS m/e = 480.0 (M+1) @ 1.01 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.77 (d, J = 2.01 Hz, 1H), 7.37 (br. s., 1H), 6.84 (s, 1H), 6.57 (s,


2H), 6.55 (br. s., 1H), 6.30 - 6.48 (m, 3H), 4.28 (br. s., 1H), 4.23 (d, $J = 13.05$ Hz, 1H), 4.13 (dd, $J = 3.39, 12.67$ Hz, 1H), 3.97 (td, $J = 6.84, 10.42$ Hz, 1H), 3.38 - 3.50 (m, 1H), 3.22 - 3.29 (m, 1H), 3.10 (br. s., 1H), 2.71-2.97 (m, 1H), 2.83 (t, $J = 12.30$ Hz, 1H), 2.13 (qd, $J = 6.13, 12.49$ Hz, 1H), 1.73 - 1.91 (m, 3H), 1.58 - 1.73 (m, 1H), 1.39 - 1.53 (m, 1H).




[0160] (3R,3'S,4'R)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-fluorophenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was separated into three peaks by SFC (IC (2x15cm), 25% MeOH(0.1% DEA)/CO₂, 100 bar, 60 ml/min). Peak 2 of 3 was further purified by SFC (IA (3 x 15 cm), 30% *i*PrOH (0.1% DEA)/CO₂, 100 bar, 60 ml/min) to afford the title compound. LCMS (Agilent460, 254 nm): ES (+) MS $m/e = 480.0$ (M+1) @. 1.01 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.77 (d, $J = 1.76$ Hz, 1H), 7.39 (s, 1H), 6.81 (s, 1H), 6.57 (s, 3H), 6.46 (td, $J = 2.01, 12.30$ Hz, 1H), 6.39 (td, $J = 1.95, 8.66$ Hz, 1H), 6.34 (d, $J = 7.53$ Hz, 1H), 4.18 - 4.48 (m, 2H), 4.09 - 4.18 (m, 1H), 3.79 - 4.09 (m, 1H), 3.33 - 3.45 (m, 2H), 3.11 (br. s., 1H), 2.92 (br. s., 1H), 2.82 (t, $J = 12.42$ Hz, 1H), 2.09 (qd, $J = 5.75, 12.11$ Hz, 1H), 1.74 - 1.93 (m, 3H), 1.51 - 1.74 (m, 1H), 1.32 - 1.51 (m, 1H).

Example 10


[0161]




[0162] The synthesis of (3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethoxy) phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. A similar procedure was used as described for the synthesis of (3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethyl)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide **14** to afford the crude material which was purified by pre-HPLC (MeOH/H₂O with 0.05% NH₃.H₂O as mobile phase) to give the title compound (320 mg, yield: 44%) as a yellow solid. ESI-MS (M+H)⁺: 546.16. HPLC: (214 nm: 98.4%, 254 nm: 98.0%). ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (s, 1H), 7.39 (s, 1H), 6.84-6.81 (m, 1H), 6.75-6.72 (m, 1H), 6.62-6.57 (m, 3H), 6.52-6.47 (m, 2H), 4.24-3.98 (m, 4H), 3.47-3.40 (m, 1H), 3.17-2.99 (m, 2H), 2.86-2.79 (m, 2H), 2.15-1.99 (m, 1H), 1.86-1.60 (m, 4H), 1.48-1.39 (m, 1H).




[0163] (3*R*,3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethoxy)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was purified by SFC (AD-H(2 x 25 cm), 30% EtOH(0.1% DEA)/CO₂, 100 bar, 70 ml/min) to afford the title compound. LCMS (Agilent460, 254 nm): ES (+) MS *m/e* = 546.0 (M+1) @ 1.20 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, *J* = 1.76 Hz, 1H), 7.39 (br. s., 1H), 6.81 (s, 1H), 6.75 (s, 1H), 6.62 (s, 1H), 6.57 (s, 2H), 6.38 - 6.52 (m, 2H), 4.23 (d, *J* = 13.05 Hz, 2H), 3.92 - 4.18 (m, 2H), 3.34 - 3.45 (m, 2H), 3.11 (br. s., 1H), 2.93 (br. s., 1H), 2.82 (t, *J* = 12.30 Hz, 1H), 2.08 (qd, *J* = 5.75, 12.11 Hz, 1H), 1.74 - 1.92 (m, 3H), 1.56 - 1.73 (m, 1H), 1.36 - 1.50 (m, 1H).



[0164] (3*S*,3'*S*,4'*R*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethoxy)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was purified by SFC (AD-H(2 x 25 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 70 ml/min) to afford the title compound. LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 546.0 (M+1) @ 1.23 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, *J* = 2.01 Hz, 1H), 7.37 (br. s., 1H), 6.84 (s, 1H), 6.72 (s, 1H), 6.60 (s, 1H), 6.57 (s, 2H), 6.43 - 6.54 (m, 2H), 4.23 (d, *J* = 13.05 Hz, 1H), 4.22 (br. s., 1H), 4.13 (dd, *J* = 3.26, 12.30 Hz, 1H), 4.01 (td, *J* = 6.81, 10.23 Hz, 1H), 3.44 (td, *J* = 6.18, 12.49 Hz, 1H), 3.20 - 3.29 (m, 1H), 3.11 (br. s., 1H), 2.88 (br. s., 1H), 2.83 (t, *J* = 12.30 Hz, 1H), 2.12 (qd, *J* = 6.05, 12.46 Hz, 1H), 1.73 - 1.91 (m, 3H), 1.59 - 1.73 (m, 1H), 1.48 (td, *J* = 9.41, 19.33 Hz, 1H).



[0165] (3*S*,3'*R*,4'*S*)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethoxy)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was purified by SFC (AD-H (2 x 25 cm), 30% EtOH (0.1% DEA)/CO₂, 100 bar, 70 ml/min) to afford the title compound. LCMS (Agilent 460, 254 nm): ES (+) MS *m/e* = 546.0 (M+1) @ 1.22 min. ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (d, *J* = 2.01 Hz, 1H), 7.38 (br. s., 1H), 6.81 (s, 1H), 6.75 (s, 1H), 6.62 (s, 1H), 6.57 (s, 2H), 6.42 - 6.51 (m, 2H), 4.23 (d, *J* = 12.80 Hz, 2H), 3.97 - 4.18 (m, 2H), 3.34 - 3.44 (m, 2H), 3.10 (br. s., 1H), 2.93 (br. s., 1H), 2.82 (t, *J* = 12.17 Hz, 1H), 2.03 - 2.15 (m, 1H), 1.77 - 1.90 (m, 3H), 1.57 - 1.73 (m, 1H), 1.37 - 1.48 (m, 1H).




[0166] (3R,3'R,4'S)-1'-(6-amino-5-fluoropyrimidin-4-yl)-3-((3-chloro-5-(trifluoromethoxy)phenyl)amino)-2-oxo-[1,3'-bipiperidine]-4'-carboxamide. The mixture of four diastereomers was purified by SFC (AD-H (2 x 25cm), 30% EtOH(0.1% DEA)/CO₂, 100bar, 70ml/min) to afford the titled compound. LCMS(Agilent 460, 254nm): ES (+) MS m/e = 546.0 (M+1) @. 1.23 min. ¹H NMR (400 MHz, CDCl₃) δ : 7.93 (s, 1H), 6.60 (s, 1H), 6.52 (s, 1H), 6.35 (s, 1H), 5.85 (br. s., 1H), 5.32 (br. s., 1H), 4.97 - 5.15 (m, 1H), 4.78 (br. s., 2H), 4.46 (d, *J* = 13.05 Hz, 2H), 3.67 - 3.87 (m, 2H), 3.34 - 3.60 (m, 4H), 2.99 (t, *J* = 12.17 Hz, 1H), 2.34 - 2.49 (m, 1H), 1.91 - 2.08 (m, 3H), 1.83 (dq, *J* = 3.76, 12.72 Hz, 1H), 1.47 - 1.74 (m, 1H).

Example 11

[0167] *In vitro* BTK kinase assay: BTK-POLYGAT-LS ASSAY. The purpose of the BTK *in vitro* assay was to determine compound potency against BTK through the measurement of IC₅₀. Compound inhibition was measured after monitoring the amount of phosphorylation of a fluorescein-labeled polyGAT peptide (Invitrogen PV3611) in the presence of active BTK enzyme (Upstate 14-552), ATP, and inhibitor. The BTK kinase reaction was done in a black 96 well plate (costar 3694). For a typical assay, a 24 μ L aliquot of an ATP/peptide master mix (final concentration; ATP 10 μ M, polyGAT 100 nM) in kinase buffer (10 mM Tris-HCl pH 7.5, 10 mM MgCl₂, 200 μ M Na₃PO₄, 5 mM DTT, 0.01% Triton X-100, and 0.2 mg/ml casein) was added to each well. Next, 1 μ L of a 4-fold, 40X compound titration in 100% DMSO solvent was added, followed by adding 15 μ L of BTK enzyme mix in 1X kinase buffer (with a final concentration of 0.25 nM). The assay was incubated for 30 minutes before being stopped with 28 μ L of a 50 mM EDTA solution. Aliquots (5 μ L) of the kinase reaction were transferred to a low volume white 384 well plate (Corning 3674), and 5 μ L of a 2X detection buffer (Invitrogen PV3574, with 4 nM Tb-PY20 antibody, Invitrogen PV3552) was added. The plate was covered and incubated for 45 minutes at room temperature. Time resolved fluorescence (TRF) on Molecular Devices M5 (332 nm excitation; 488 nm emission; 518 nm fluorescein emission) was measured. IC₅₀ values were calculated using a four parameter fit with 100% enzyme activity determined from the DMSO control and 0% activity from the EDTA control.

[0168] Selected compounds of formula I were tested and found to be active in the polyGAT assay. Compounds **I-1**, **I-2**, **I-3**, **I-4**, **I-5** and **I-7** gave IC₅₀ values of 0.73 nM, 0.68 nM, 2.07 nM, 0.63 nM, 1.6 nM, and 1.2 nM respectively. Compound **I-6** has an IC₅₀ value less than 1 nM. Comparator compound **I^C**, shown below, produced an IC₅₀ value of 2.0 nM.



I^C

Example 12

STUDY PROTOCOL TO DETERMINE ACTIVATION OF THE PXR NUCLEAR RECEPTOR IN HUMAN DPX2 CELLS

[0169]

a) Protocol Summary: PXR has been shown to be a primary nuclear receptor that mediates drug-induced expression of CYP3A4 (Bertilsson G, et al.; Proc Natl Acad Sci U S A. 1998 Oct 13;95(21):12208-13). Based on this pathway of CYP3A4 induction, cell-based PXR reporter gene assay is commonly used to screen new molecular entities (NMEs) in early drug discovery stage, for their potential to induce CYP3A4 (Luo G, et al. Drug Metab Dispos. 2002 Jul;30(7):795-804.) Studies were designed to evaluate the effect of new molecular entities (NMEs) on the activation of human PXR in DPX2 cells. Cell lines stably transfected with the PXR nuclear receptor and corresponding response elements were seeded into 96-well plates. Twenty-four hr after seeding, cells were treated with 6 distinct concentrations of NMEs in triplicate wells (see below), and cells then returned to the incubator for an additional 24 hr. At the end of this incubation period, the number of viable cells/well were determined using Promega's Cell Titer Fluor cytotoxicity assay. Following this assay, Promega's ONE-Glo was added to the same wells and reporter gene activity assessed.

b) Test System: The test system consisted of the stably transformed DPX2 tumor cell line plated on 96-well microtiter plates. An expression vector harboring the PXR nuclear receptor plus the appropriate enhancers and promoters linked to the luciferase reporter gene have been stably integrated into these tumor cell lines. Receptor activation was assessed by monitoring reporter gene activity, and by comparing the results to vehicle-treated cells. Positive controls consist of cells treated with 6 different concentrations (0.1, 0.5, 1, 5, 10, and 20 μ M) of rifampicin. In this manner, compounds activating PXR can be easily and rapidly identified. Since stably-integrated cell lines were used, it is possible to observe from 3- to 70-fold receptor activation.

c) Data Processing and Receptor Activation Kinetics: Data processed using MS-Excel was calculated as the mean ($n = 3$) and % CV of the fold PXR activation relative to vehicle-treated cells at each of the 6 different doses. All activation data was normalized to the number of viable cells/well. Results were also expressed as a percentage of the response given by the appropriate positive control at a 10 μ M dose. EC_{50} and E_{max} values were derived for test compounds that give receptor activation using nonlinear regression of typical log dose-response curves (Prism V5.0c, GraphPad Software, San Diego, CA). Agents exhibiting atypical dose-response curves were not analyzed in this fashion.

d) New Molecular Entities (NMEs): Test compounds were tested at 0.05, 0.1, 0.5, 1, 2.5, and 10 μ M

[0170] Selected compounds of formula I were tested in the PXR assay. Compounds **I-1**, **I-2**, **I-3**, **I-4**, and **I-5** gave PXR % induction (relative to 10 μ M rifampin) of 62%, 42%, 47%, 67%, and 90%, respectively. Comparator compound **I^c**, shown above, produced a PXR % induction of 95%.

Example 13

Protocol for FastPatch hERG inhibition assay:

[0171] The cardiac potassium channel, hERG, is responsible for a rapid delayed rectifier current (I_{Kr}) in human ventricle and inhibition of I_{Kr} is the most common cause of cardiac action potential prolongation by non-cardiac drugs (see, e.g., Weirich and Antoni, Basic Res. Cardiol., 93, Suppl. 1, 125-32, 1998; Yap and Camm, Clin. Exp. Allergy, 29, Suppl. 3, 174-81, 1999). Increased action potential duration has been cited as a factor in causing prolongation of the QT interval that has been associated with a dangerous ventricular arrhythmia, torsade de pointes (Brown and Rampe, Pharmaceutical News, 7, 15-20, 2000).

[0172] The *in vitro* effects of provided compounds was investigated on the hERG (human ether-à-go-go-related gene) potassium channel current (a surrogate for I_{Kr} , the rapidly activating, delayed rectifier cardiac potassium current) expressed in human embryonic kidney (HEK293) cells stably transfected with hERG cDNA. Cells were placed in HEPES-buffered physiological saline solution in a glass-lined 96-well plate and loaded with appropriate amounts of test and control solutions for a duration of a 3-minute exposure at each concentration. Test compound was diluted in 0.3% DMSO. An automated parallel patch clamp system, QPatch HT (Sophion Bioscience A/S, Denmark), was used to evaluate at various concentrations (e.g., 10 μ M). The IC_{50} values was estimated based on the hERG inhibition data. The study was performed at ChanTest (14656 Neo Parkway, Cleveland, OH). The QPatch screen is further described by Janzen and

Bernasconi (eds.), High Throughput Screening, Methods and Protocols, Second Edition, vol. 565, chapter 10, pg. 209-223, 2009.

[0173] Selected compounds of formula I were tested in the hERG assay. Compounds I-1, I-2, I-3, and I-4 gave hERG IC₅₀ of 15.6 uM, 30 uM, 14.6 uM, and 13.7 uM, respectively. Compound I-5 shows no observable activity in the hERG assay at 10 uM (no IC₅₀ available). Compound I-6 shows little observable hERG activity (< 20% inhibition) at 10uM (no IC₅₀ available). Compound I-7 shows hERG activity (65% inhibition) at 10uM (no IC₅₀ available). Comparator compound IC, shown above, produced a hERG IC₅₀ of 1.18 uM or activity (87% inhibition) at 10 uM.

Example 14

GSH Trapping in Human Liver Microsome : Protocol

[0174] Test compound (final concentration 10 uM) is incubated with either human or rat liver microsomes (final concentration 1 mg/mL), along with activating cofactors NADPH (final concentration 1 mM), potassium phosphate (final concentration 100 mM pH 7.4), magnesium chloride (final concentration 3.3 mM) and the trapping agent GSH (final concentration 5 mM). The incubation mixture is incubated for 60 min at 37 °C and terminated with ice cold acetonitrile (equal volume as incubation mixture) and the supernatants isolated. The supernatants are either injected directly for LC/MS/MS analysis or dried under N₂ and reconstituted in water:acetonitrile (80:20) mixture before LC/MS/MS analysis. The corresponding GSH conjugate is evaluated via LC/MS/MS, using a Triple TOF5600/Xevo Qtof MSe.

Example 15

Rat Collagen-Induced Arthritis Model

[0175] The collagen induced arthritis (CIA) model in female Lewis rats requires primary T and B cell immune responses to type II collagen (CII) immunization for the development of a severe inflammatory disease (see Goldschmidt TJ, Holmdahl R. Cell Immunol. 154(1):240-8, 1994; Helfgott, S. M., et al.; Clin. Immunol. Immunopathol. 31:403, 1984; Holmdahl R. et al., J Autoimmun. 7(6):739-52, 1994; and Stuart, J. M., et al., J. Exp. Med. 155:1, 1982). Clinical disease onsets after a secondary CII challenge and the disease progresses over the following eight days.

[0176] Generally, female Lewis rats are immunized with bovine collagen type II in incomplete Freund's adjuvant. Rats (N=10/group) receive daily oral administration of test compound or vehicle BID by oral gavage beginning on day 1 (therapeutic). Clinical severity of arthritis is assessed by caliper measurements of ankles taken every day beginning on Day 0.

[0177] Detailed protocol: Female Lewis rats are immunized subcutaneously with bovine collagen type II (1:1 emulsion of 2 mg/ml bovine CII in 0.01 N acetic acid: Incomplete Freund's Adjuvant) at three sites of back skin. Six days post immunization rats receive a second subcutaneous injection of bovine CII. A compound of formula I suspension or vehicle (0.5% CMC, 0.1% Tween 80) is administered by oral gavage BID beginning on day 0 (prophylactic) (n=10 animals/group). Clinical severity of CIA is assessed by caliper measurements of ankles taken every day beginning on Day 9. Baseline ankle caliper measurements are taken and confirmed as clinically normal (0.260-0.264 in) for prophylactic treatment. Baseline ankle caliper measurements for established disease animals is assessed on day 1 of therapeutic dosing and animals are randomly assigned to treatment groups after confirmation of clinical disease onset (0.2751-0.2755 in). Data are analyzed across all groups using a one-way analysis of variance (1-way ANOVA), along with an appropriate multiple comparison post-test. Significance for all tests is set at p ≤0.05.

Example 16

Analysis of BCR pathway activation via inhibition of phosphorylation of PLCγ2

[0178] Protocol: One day before treatment, Ramos cells are plated at a density of 3x10⁵ cells per well in 200 μL of complete medium in a 96-well tissue culture filter plates (Millipore, Billerica, MA). On the day of treatment, used medium is removed by filtration and the cells resuspended in 200 μL serum free medium containing serial compound dilutions and DMSO to 0.1%, then incubated for 2 hours at 37 °C. Cells are stimulated for 5 minutes with 10 μg/mL goat anti-human IgM at 37 °C. All medium is removed by filtration and the cells are rinsed with ice cold PBS then lysed on ice for 1 hour with lysis buffer containing; 20 mM Tris (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 2 mM Na₃VO₄, 1% Triton X-100, 0.1% SDS, protease inhibitor cocktail, 1 mM phenylmethylsulfonyl fluoride, (PMSF), Phosphatase inhibitor mix 2 (Sigma cat # P5726 from Sigma, St. Louis, Mo.), and Phosphatase inhibitor mix 3 (Sigma cat # P0044 from Sigma, St. Louis, Mo.). Lysates are subsequently transferred to standard MSD plates (Meso Scale Discovery, (MSD), (Gaithersburg, Maryland)), pretreated with capture antibody (anti-total PLCγ2 antibody B10, (SantaCruz Biotechnologies (Santa

Cruz, CA)) and blocked with BSA according to the manufacturer's directions. Lysates are incubated in the prepared MSD plates overnight at 4 °C with gentle agitation. Wells are washed three times with TBST and treated with anti pPLC γ 2 (SantaCruz) in 1% BSA in PBS for 1 hour at room temperature. Wells are again washed three times with TBST and treated with anti-rabbit sulfo-tag antibody (MSD), for 1 hour at room temperature. After washing with TBST, MSD read buffer is added and the luminescence is measured in an MSD SECTOR Imager 6000. Maximum response is determined as the average luminescence in wells containing stimulated cells treated with anti-IgM and DMSO. Minimal response is determined as the average luminescence in wells containing unstimulated cells treated with DMSO alone. The maximal and minimal values are used to normalize luminescence in compound treatment wells. The normalized values are plotted against compound concentration on a log scale then analyzed using Prism software (GraphPad Software, Inc.). A sigmoidal dose-response equation with variable slope is used to fit the data and generate 50% inhibition concentration (IC₅₀).

[0179] Ramos cells are incubated in 96 well plates with a range of concentrations of a compound of formula **I** for 2 hours, stimulated with 10 μ g/mL anti-IgM for 5 minutes, and PLC γ 2 phosphorylation measured using an electrochemical-luminescent immunoassay. The EC₅₀ is calculated using GraphPad Prism software.

Example 17

Inhibition BCR-induced human B cell proliferation

[0180] Human CD19+ B cells are stimulated with an anti-IgM antibody and the activity of a compound of formula **I** is evaluated in terms of altering cellular metabolism after 72 hours. In this context, cellular metabolism directly correlates with cellular activation and proliferation, and can also reflect relative cell survival during proliferation. Anti-IgM antibody is evaluated for effects on B cell proliferation and determined to exhibit a half-maximal concentration for activation of 10 μ g/mL. Using these activation conditions, varying concentrations of test compound are assayed, in triplicate in 0.1% DMSO, for impact on cellular metabolism of CD 19+ B cells isolated from different donors.

[0181] Protocol: Human B cells are isolated from peripheral blood mononuclear cells or unpurified buffy coats using Ficoll-Hypaque gradients (Amersham) and negatively selected by magnetic cell sorting (Human B Cell Isolation Kit II, Miltenyi Biotec). Target cell purity is determined by flow cytometry by staining for markers of B cells, T cells and monocytes (CD19, CD3, CD14, respectively; BD Biosciences). Data are collected on a FACsCaliber flow cytometer and analyzed using FloJo software (BD Biosciences). Purity of human B cell preparations is routinely greater than 95%. Negatively selected human B cells are stimulated with 10 μ g/mL anti-IgM F(ab')₂ (Jackson ImmunoResearch) in 96 well plates. 100,000 B cells in 0.2 mL RPMI + 10% FBS are treated with varying concentrations (titrated from 5000 nM to 0 nM in 0.5% DMSO) of a compound of formula **I** in triplicate wells or vehicle control in 0.5% DMSO final concentration for 30 minutes at 37 °C, 5% CO₂, then cells are stimulated with 10 μ g/mL anti-IgM F(ab')₂. B cells are stimulated for 72 hr at 37 °C, 5% CO₂. Proliferation is measured using the CellTiter-Glo reagent (Promega), as measured on a luminometer. Mean values are plotted against maximum proliferation and IC₅₀ values are determined using GraphPad Prism v5 software.

Example 18

Evaluation of the effect of compounds on myeloid cell activation *in vitro*

[0182] Fc γ R activation of primary human macrophages. Autoantibody and immune-complex mediated activation through Fc γ R can be modeled by activation of macrophages with immobilized IgG. Primary human macrophages derived from GM-CSF treated monocytes up-regulate activation markers such as CD80, CD86, MHC antigens and the Fc γ RIII receptor. Human monocyte derived macrophages can be activated by plate-bound purified human IgG. This stimulation crosslinks the Fc γ RIII receptor and induces the secretion of pro-inflammatory cytokines such as TNF α , IL-6, IL1 β and MCP-1. Compound of formula **I** are evaluated for inhibition of cytokine expression following FcR activation of human macrophages.

[0183] Generally, macrophages are cultured in plates previously incubated with purified IgG then washed. Titrations of test compound (10,000 nM to 0 nM) are added to these cultures. Cell culture supernatants are analyzed by ELISA for the expression of TNF α and IL-6.

[0184] Protocol: Human monocytes are isolated from buffy coats of healthy donors and negatively selected by magnetic cell sorting (Monocyte Isolation Kit II, Miltenyi Biotec). Purified monocytes are cultured in standard media supplemented with low-IgG FBS and 100 ng/mL GM-CSF for 5-7 days to induce macrophage differentiation. Cultured macrophages are stimulated with 100 μ g/mL plate-bound purified IgG $\pm$ a titration of test compound (10 μ M to 0 nM). Supernatants are collected after 4 hrs and 18 hrs and analyzed for TNF α and IL-6, respectively.

Example 19

Efficacy in Mouse Collagen Antibody-Induced Arthritis

[0185] This Example relates not only to arthritis, but also evaluates the activity of autoantibodies and immune complexes *in vivo* and therefore is relevant to other inflammatory disorders such as SLE. In this experiment, the activity of autoantibodies and immune complexes produce a pathological endpoint that is dependent on FcR signalling, and the Fc portion of such antibodies is inhibited by administration of a compound of formula I.

[0186] The collagen antibody-induced arthritis (CAIA) model in female DBA/1 mice does not require cognate T and B cell responses for the induction of inflammation but rather relies on immune effector mechanisms for the development of clinical disease. A cocktail of four anti-collagen II (CII) specific monoclonal antibodies and immune stimulatory lipopolysaccharide (LPS) administered 3 days after CII specific antibody transfer promote antibody-Fc-Receptor engagement (Kagari T. et al.; J Immunol. 170:4318-24 (2003)), immune complex formation, complement activation (Banda NK, et al.; Clin Exp Immunol. 159:100-8 (2010)) and pro-inflammatory cytokine production to induce a severe inflammatory disease over a 10 day period.


[0187] Generally, arthritis is induced by injection of a cocktail of monoclonal anti-collagen antibodies into DBA/1 mice on day 0. Mice (N=10/group) receive daily oral administration of test compound either QD or BID as indicated beginning on day 0. Paw inflammation is evaluated daily.

[0188] Protocol: Female DBA/1 mice 6-8 weeks of age receive 2 mg of an arthritogenic four clone monoclonal antibody cocktail (Chondrex#10100) i.v. on day 0 followed by a 50 ug dose of LPS on 3 days later. Test compound suspension or vehicle (0.5% CMC, 0.1% Tween 80) is administered BID by oral gavage beginning on day 0 (10 animals/group) just prior to i.v. transfer of antibody cocktail. Clinical severity of CIA is assessed by monitoring inflammation on all four paws, applying a scale ranging from 0 to 4. Each paw is graded as follows: 0, normal; 1, mild but definite redness and swelling of the ankle or wrist, or redness and swelling of any severity for 1 or 2 digits; 2, moderate to severe redness and swelling of the ankle or wrist, or more than two digits; 3, redness and swelling (pronounced edema) of the entire paw; and 4, maximally inflamed limb with involvement of multiple joints. The sum of the four individual scores is the arthritis index, with a maximal possible score of 16 for each animal.

Numbered Embodiments

[0189] The present invention also relates to the following numbered embodiments.

1. A compound of formula I:



or a pharmaceutically acceptable salt thereof,
wherein:

each R^1 is independently hydrogen, an optionally substituted C_{1-6} aliphatic group, an optionally substituted 3-7 membered monocyclic heterocyclic group, or an optionally substituted heterocyclylalkyl group having 3-7 carbon atoms and 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;


or two R^1 groups are taken together with their intervening atoms to form an optionally substituted 3-7 membered saturated or partially unsaturated monocyclic heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

wherein optionally substituted groups may be substituted with halogen, $-\text{NO}_2$, $-\text{CN}$, $-\text{OR}$, $-\text{SR}$, $-\text{N}(\text{R})_2$, $-\text{C}(\text{O})\text{R}$, $-\text{CO}_2\text{R}$, $-\text{N}(\text{R})\text{C}(\text{O})\text{OR}$, $-\text{C}(\text{O})\text{N}(\text{R})_2$, $-\text{OC}(\text{O})\text{R}$, $-\text{N}(\text{R})\text{C}(\text{O})\text{R}$, $-\text{S}(\text{O})\text{R}$, $-\text{S}(\text{O})_2\text{R}$, or $-\text{S}(\text{O})_2\text{N}(\text{R})_2$;

each R is independently hydrogen or C_{1-6} aliphatic;

or two R groups attached to the same nitrogen are taken together with their intervening atoms to form an optionally substituted 3-7 membered saturated or partially unsaturated monocyclic heterocyclic ring having 1-2 heteroatoms, in which any second heteroatom is independently selected from nitrogen, oxygen, or sulfur;


Ring A is



R^2 is $-\text{Cl}$ or $-\text{F}$; and


R^3 is $-\text{CF}_3$, $-\text{OCF}_3$, or $-\text{F}$.

2. The compound of embodiment 1, wherein the compound is of formula **II-a**:




or a pharmaceutically acceptable salt thereof.

3. The compound of embodiment 1, wherein the compound is of formula **II-b**:




or a pharmaceutically acceptable salt thereof.

4. The compound of embodiment 1, wherein the compound is of formula III:



III

5. The compound of embodiment 1, wherein the compound is formula IV:



IV

6. The compound of embodiment 5, wherein R³ is -CF₃.

7. The compound of embodiment 5, wherein R³ is -F.

8. The compound of embodiment 5, wherein both R¹ are hydrogen.

9. The compound of embodiment 5, wherein one R¹ is hydrogen and the other R¹ is an optionally substituted C₁₋₆ aliphatic.

10. The compound of embodiment 9, wherein one R¹ is hydrogen and the other R¹ is methyl.

11. The compound of embodiment 5, wherein both R¹ are optionally substituted C₁₋₆ aliphatic groups.

12. The compound of embodiment 1, wherein one R¹ is hydrogen and the other an is optionally substituted C₁₋₆ aliphatic.

13. The compound of embodiment 1, wherein both R¹ are optionally substituted C₁₋₆ aliphatic groups.

14. The compound of embodiment 1, wherein both R¹ are hydrogen.

15. The compound of embodiment 1, wherein R² is -Cl.


16. The compound of embodiment 1, wherein R² is -F.

17. The compound of embodiment 1, wherein R³ is -CF₃.


18. The compound of embodiment 1, wherein R³ is -OCF₃.

19. The compound of embodiment 1, wherein R³ is -F.


20. A compound selected from the group consisting of:




I-1




I-2




I-3




I-4



I-5



I-6

**I-7**

21. A method of decreasing the enzymatic activity of Bruton's tyrosine kinase comprising contacting Bruton's tyrosine kinase with an effective amount of a compound of any one of embodiments 1-20 or a composition thereof.

22. A method of treating a disorder responsive to inhibition of Bruton's tyrosine kinase comprising administering to a subject an effective amount of a compound of any one of embodiments 1-20 or a composition thereof.

23. A method of treating a disorder selected from the group consisting of autoimmune disorders, inflammatory disorders, and cancers comprising administering to a subject an effective amount of a compound of any one of embodiments 1-20 or a composition thereof.

24. The method of embodiment 23, wherein the disorder is rheumatoid arthritis.


25. The method of embodiment 23, wherein the disorder is systemic lupus erythematosus.






26. The method of embodiment 23, wherein the disorder is atopic dermatitis.

27. The method of embodiment 23, wherein the disorder is leukemia or lymphoma.

Claims

1. A pharmaceutically acceptable salt of a compound selected from the group consisting of:

**I-1****I-2**

**I-3****I-4****I-5****I-6****I-7**

2. A pharmaceutically acceptable salt of a compound of claim 1 or a composition thereof for use in decreasing the enzymatic activity of Bruton's tyrosine kinase.
3. A pharmaceutically acceptable salt of a compound of claim 1 or a composition thereof for use in the treatment of a disorder responsive to inhibition of Bruton's tyrosine kinase.
4. A pharmaceutically acceptable salt of a compound of claim 1 or a composition thereof for use in the treatment of a disorder selected from the group consisting of autoimmune disorders, inflammatory disorders, and cancers.
5. The use of claim 4, wherein the disorder is rheumatoid arthritis.
6. The use of claim 4, wherein the disorder is systemic lupus erythematosus.
7. The use of claim 4, wherein the disorder is atopic dermatitis.
8. The use of claim 4, wherein the disorder is leukemia or lymphoma.

EP 3 385 263 A1

9. A pharmaceutically acceptable salt of a compound of claim 1 or a composition thereof for use in treatment of a disorder associated with dysregulated BCR signaling.
- 5 10. A pharmaceutically acceptable salt of a compound of claim 1 or a composition thereof for use in the treatment of a disorder responsive to B-cell depletion.
- 10 11. A pharmaceutically acceptable salt of a compound of claim 1 or a composition thereof for use in the treatment of a disorder selected from the group consisting of autoimmune disorders, inflammatory disorders, and cancers, wherein each of which is associated with B-cell activation, proliferation, or differentiation.
12. The use of claim 11, wherein the disorder is Non-Hodgkin's Lymphoma.
13. The use of claim 11, wherein the disorder is chronic lymphocytic leukemia.

15

20

25

30

35

40

45

50

55



EUROPEAN SEARCH REPORT

Application Number
EP 18 17 1833

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	WO 2011/029046 A1 (BIOGEN IDEC INC [US]; ERLANSON DANIEL A [US]; MARCOTTE DOUG [US]; KUMA) 10 March 2011 (2011-03-10) * paragraph [0134]; claims 68,71 * -----	1-8	INV. C07D401/14 A61K31/435 A61K31/445 A61P37/00
			TECHNICAL FIELDS SEARCHED (IPC)
			C07D A61K A61P
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 1 August 2018	Examiner Johnson, Claire
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03/82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 18 17 1833

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

01-08-2018

10

15

20

25

30

35

40

45

50

55

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2011029046 A1	10-03-2011	AR 078320 A1	02-11-2011
		AU 2010289313 A1	12-04-2012
		AU 2016202601 A1	19-05-2016
		AU 2018201668 A1	05-04-2018
		CA 2771822 A1	10-03-2011
		CN 102711473 A	03-10-2012
		CN 107011330 A	04-08-2017
		EP 2473049 A1	11-07-2012
		JP 5699149 B2	08-04-2015
		JP 6326533 B2	16-05-2018
		JP 2013503905 A	04-02-2013
		JP 2015091898 A	14-05-2015
		JP 2017186373 A	12-10-2017
		JP 2018111726 A	19-07-2018
		NZ 598985 A	26-07-2013
		TW 201120040 A	16-06-2011
		TW 201713640 A	16-04-2017
		US 2012157443 A1	21-06-2012
		US 2014309212 A1	16-10-2014
		US 2016376281 A1	29-12-2016
		US 2018065974 A1	08-03-2018
		WO 2011029046 A1	10-03-2011

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61657360 A [0001]
- WO 9605309 A [0069]
- US 4911920 A [0076]
- US 5403841 A [0076]
- US 5212162 A [0076]
- US 4861760 A [0076]

Non-patent literature cited in the description

- Additionally, general principles of organic chemistry. Handbook of Chemistry and Physics [0009]
- **THOMAS SORRELL**. Organic Chemistry. University Science Books, 1999 [0009]
- March's Advanced Organic Chemistry. John Wiley & Sons, 2001 [0009]
- **S. M. BERGE et al.** pharmaceutically acceptable salts. *J. Pharmaceutical Sciences*, 1977, vol. 66, 1-19 [0017]
- **T. W. GREENE ; P. G. M. WUTS**. Protecting Groups in Organic Synthesis. John Wiley & Sons, 1999 [0041]
- **PANAYI GS et al.** *Rheum Dis Clin North Am*, 2001, vol. 27, 317-334 [0052]
- **ZHANG Z ; BRIDGES SL**. *Rheum Dis Clin North Am*, 2001, vol. 27, 335-353 [0052]
- **COHEN SB et al.** Arthritis Rheum. REFLEX Trial Group, September 2006, vol. 54, 2793-806 [0052]
- **KLIPPEL JH et al.** Primer on the rheumatic diseases. Arthritis Foundation, 2001 [0053]
- **CHAN OT et al.** *Immunol Rev.*, 1999, vol. 169, 107-121 [0053]
- **NAVARRA SV et al.** *Lancet*, 26 February 2011, vol. 377 (9767), 721-31 [0053]
- **FURIE R et al.** *Arthritis Rheum.*, December 2011, vol. 63 (12), 3918-30 [0053]
- **HONIGBERG, L.A.** *PNAS*, 2010, vol. 107, 13075 [0053]
- Pharmaceutical Sciences. Mack Pub. Co, [0069]
- **PHILLIPS, D. P. ; ZHU, X. -F. ; LAU, T. L. ; YANG, K. ; LIU, H.** *Tetrahedron Letters*, 2009, vol. 50, 7293 [0125]
- **BERTILSSON G et al.** *Proc Natl Acad Sci U S A*, 13 October 1998, vol. 95 (21), 12208-13 [0169]
- **LUO G et al.** *Drug Metab Dispos*, July 2002, vol. 30 (7), 795-804 [0169]
- **WEIRICH ; ANTONI.** *Basic Res. Cardiol.*, 1998, vol. 93 (1), 125-32 [0171]
- **YAP ; CAMM.** *Clin. Exp. Allergy*, 1999, vol. 29 (3), 174-81 [0171]
- **BROWN ; RAMPE.** *Pharmaceutical News*, 2000, vol. 7, 15-20 [0171]
- High Throughput Screening, Methods and Protocols. 2009, vol. 565, 209-223 [0172]
- **GOLDSCHMIDT TJ ; HOLMDAHL R.** *Cell Immunol.*, 1994, vol. 154 (1), 240-8 [0175]
- **HELFGOTT, S. M. et al.** *Clin. Immunol. Immunopathol.*, 1984, vol. 31, 403 [0175]
- **HOLMDAHL R. et al.** *J Autoimmun.*, 1994, vol. 7 (6), 739-52 [0175]
- **STUART, J. M. et al.** *J. Exp. Med.*, 1982, vol. 155 (1) [0175]
- **KAGARI T. et al.** *J Immunol.*, 2003, vol. 170, 4318-24 [0186]
- **BANDA NK et al.** *Clin Exp Immunol.*, 2010, vol. 159, 100-8 [0186]