



(11) **EP 3 415 501 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:  
**Corrected version no 1 (W1 B1)**  
**Corrections, see**  
**Claims EN 1**

(48) Corrigendum issued on:  
**02.12.2020 Bulletin 2020/49**

(45) Date of publication and mention  
of the grant of the patent:  
**08.04.2020 Bulletin 2020/15**

(21) Application number: **18185739.2**

(22) Date of filing: **23.07.2014**

(51) Int Cl.:  
**C07D 215/54** <sup>(2006.01)</sup> **C07D 401/12** <sup>(2006.01)</sup>  
**C07D 401/14** <sup>(2006.01)</sup> **A61K 31/4709** <sup>(2006.01)</sup>  
**A61K 31/5377** <sup>(2006.01)</sup> **A61P 9/00** <sup>(2006.01)</sup>  
**A61P 13/12** <sup>(2006.01)</sup> **A61P 25/02** <sup>(2006.01)</sup>  
**A61P 35/00** <sup>(2006.01)</sup> **A61P 35/02** <sup>(2006.01)</sup>  
**A61P 35/04** <sup>(2006.01)</sup> **A61P 37/02** <sup>(2006.01)</sup>  
**A61P 43/00** <sup>(2006.01)</sup>

(54) **QUINOLINE DERIVATIVE**  
CHINOLINDERIVAT  
DÉRIVÉ DE QUINOLÉINE

(84) Designated Contracting States:  
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB**  
**GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO**  
**PL PT RO RS SE SI SK SM TR**

(30) Priority: **24.07.2013 JP 2013153350**

(43) Date of publication of application:  
**19.12.2018 Bulletin 2018/51**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**14828976.2 / 3 026 045**

(73) Proprietor: **ONO Pharmaceutical Co., Ltd.**  
**Osaka-shi, Osaka 541-8526 (JP)**

(72) Inventors:  
• **INUKAI, Takayuki**  
**Mishima-gun, Osaka 618-8585 (JP)**

• **TAKEUCHI, Jun**  
**Mishima-gun, Osaka 618-8585 (JP)**  
• **YASUHIRO, Tomoko**  
**Mishima-gun, Osaka 618-8585 (JP)**

(74) Representative: **Grünecker Patent- und**  
**Rechtsanwälte**  
**PartG mbB**  
**Leopoldstraße 4**  
**80802 München (DE)**

(56) References cited:  
**WO-A1-2009/137429 WO-A1-2012/011548**  
**WO-A1-2012/028332 WO-A1-2013/074633**

Remarks:

The file contains technical information submitted after  
the application was filed and not included in this  
specification


Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 3 415 501 B9**

## Description

[Technical Field]

**[0001]** The present invention relates to a compound represented by general formula (I):




wherein all of the symbols have the same meanings as given below, a salt thereof, a solvate thereof, or an N-oxide thereof (hereinafter, also abbreviated as the compound of the present invention).

[Background Art]


**[0002]** Axl (also known as: UFO, ARK, Tyro7) is a receptor tyrosine kinase belonging to a TAM family (Axl, Mer and Tyro3) cloned from tumor cells. Gas6 (growth-arrest-specific protein 6) cloned as a gene specifically expressed at the time of cell proliferation arrest is known as a ligand for Axl. Axl activated by binding of Gas6 transfers a signal via phosphorylation. Since the signal activates an Erk1/2 pathway or a PI3K/Akt pathway, the activation of Axl is known to be involved in pathologic conditions of cancers, immune system diseases, circulatory system diseases, and the like (see, Non-Patent Literature 1).

**[0003]** In particular, the relation between Axl and various types of cancers is well known. For example, it is known that the expression of Axl is involved in metastasis and prognosis of breast cancer (see, Non-Patent Literature 2), and that Axl is involved in the pathologic conditions of acute myeloid leukemia (AML) (see Non-Patent Literature 3). Therefore, it is considered that compounds which inhibit the activation of Axl are useful for treatment of various type of cancers, immune system diseases, and circulatory system diseases.

**[0004]** By the way, as prior art of the compound of the present invention, a compound represented by general formula (A):




(wherein  $A^A$  represents C- $R^{10A}$  and N;  $B^A$  represents C- $R^{11A}$  and N;  $D^A$  represents heterocycles of the following general formulae, and the like.




(wherein  $R^{1A}$ ,  $R^{4A}$ , and  $R^{88A}$  are independently -H, -F, -Cl, -Br, -I, -OH, -NH<sub>2</sub>, -OCH<sub>3</sub>, -OC<sub>2</sub>H<sub>5</sub>, or the like;  $R^{2A}$  and  $R^{3A}$  are independently -R<sup>88A</sup> or the like;  $R^{5A}$  and  $R^{6A}$  may be the same as each other or different from each other, and represent -H, -F, -Cl, -Br, -I, -CN, -NO<sub>2</sub>, -CH<sub>3</sub>, or the like;  $R^{7A}$ ,  $R^{8A}$ ,  $R^{10A}$ , and  $R^{11A}$  may be the same as each other or different from each other, and represent -H, -F, -Cl, -Br, -I, -CN, -NO<sub>2</sub>, -CH<sub>3</sub>, or the like;  $R^{9A}$  represents -H or the like;  $R^{12A}$  represents -CN, phenyl, or the like;  $R^{13A}$  represents -H, -F, -Cl, -Br, -I, -CN, -NO<sub>2</sub>, -CH<sub>3</sub>, or the like;  $R^{14A}$  represents -H, -F, -Cl, -Br, -I, -NO<sub>2</sub>, -CN, or the like (where the definitions of the groups are excerpted)) is known to be an Axl inhibitor (see, Non-Patent Literature 1).

**[0005]** Furthermore, a compound represented by general formula (B):



(wherein  $E^B$  and  $G^B$  are independently a hydrogen atom, a C1-6 alkyl group optionally substituted with one to six  $R^{19B}$ , a C6-11 aryl group optionally substituted with one to six  $R^{19B}$  or the like;  $X^B$  represents N or C- $R^{4B}$ ;  $Y^B$  represents N or C- $R^{1dB}$ ;  $D^B$  represents -O-, -S-, -NH- or the like;  $W^B$  represents CH or N;  $R^{aB}$ ,  $R^{bB}$ ,  $R^{cB}$ ,  $R^{dB}$ ,  $R^{1aB}$ ,  $R^{1bB}$ ,  $R^{1cB}$ ,  $R^{1dB}$  and  $R^{4B}$  independently represent a hydrogen atom, -OR<sup>110B</sup>, or the like;  $R^{19B}$  represents a halogen atom, -CN, or the like; and  $R^{110B}$  represents a hydrogen atom, a C1-6 alkyl group optionally substituted with one to six  $R^{129B}$  (where the definitions of the groups are excerpted)) is known to be an Axl inhibitor (see Patent Literature 2).

**[0006]** On the other hand, a compound having a quinoline skeleton and represented by the following general formula (C):




is known to have an ASK1 inhibitory activity, and be an agent for preventing and/or treating amyotrophic lateral sclerosis (ALS) (see Patent Literature 3).


**[0007]** Furthermore, a compound represented by general formula (D):



(wherein  $R^D$  represents




or the like;  $T^D$  represents phenyl or the like;  $Z^D$  represents N or  $CR^{7D}$ ;  $W^D$  represents a substituted or unsubstituted phenyl, substituted or unsubstituted 6-membered nitrogen-containing heteroaryl or the like;  $X^D$  represents O, S,  $S(=O)$ , or the like;  $Y^D$  represents  $-NR^{aD}C(=O)-(CR^{3D}R^{4D})_p-$  or the like;  $R^{aD}$  represents, a hydrogen atom, an alkyl group, or the like; and  $R^{1D}$  represents




or the like;  $J^{2D}$  represents O or  $CR^{4aD}R^{4aD}$ ;  $Q^D$  represents 1- to 5-membered saturated or partially unsaturated alkyl chain or the like;  $R^{1D}$  represents optionally substituted phenyl or may be fused to optionally substituted 5- to 6-membered heterocycle;  $R^{3D}$  and  $R^{4D}$  each independently represents a hydrogen atom, an alkyl group, an aryl group, or the like;  $R^{4aD}$  is absent or represents a hydrogen atom, a halogen atom, or the like (where the definitions of the groups are excerpted)) is known to be a c-Met inhibitor (see Patent Literature 4).

**[0008]** Furthermore, a compound represented by general formula (E):



(wherein  $R^{1E}$ ,  $R^{2E}$  and  $R^{4E}$  independently represent H, F, Cl, Br, I, CN,  $OR^{10E}$ , Cl-C12 alkyl, or the like;  $L^E$  represents a C3-C12 carbon ring, C6-C20 aryl, or the like;  $R^{5E}$  represents  $-C(=Y^E)R^{13E}$ ,  $-C(=Y^E)R^{10E}R^{13E}$ ,  $-NR^{10E}C(=Y^E)R^{13E}$ , or the like;  $R^{10E}$  represents H, C1-C12 alkyl, a C3-C12 carbon ring, a C2-C20 heterocycle, or the like;  $R^{13E}$  represents H, C1-C6 alkyl, or the like; and  $Y^E$  represents O or S (where the definitions of the groups are excerpted)) is known to be a c-Met inhibitor (see Patent Literature 5).


**[0009]** However, any of the prior art literatures neither mention nor suggest that a quinoline derivative as a compound of the present invention, having a bicyclic structure in which a saturated carbon ring is fused to a pyridone ring, represented by the following structural formula.



has a significant Axl inhibitory activity.

Patent Literature 6 discloses methods for treating cancer comprising administering to a patient therapeutically effective amounts of:

a) a compound of the formula A:



A

or a pharmaceutically acceptable salt thereof; and

(b) an erbB inhibitor that inhibits erbB-1 or erbB-2 or erb-3 receptor or a combination thereof; wherein,

$R^1$  is  $C_1$ - $C_6$ -alkyl;

$R^2$  is  $C_1$ - $C_6$ -alkyl or  $-(CH_2)_n-N(R^5)_2$ ;

$R^3$  is Cl or F;

$R^4$  is Cl or F;

each  $R^5$  is independently  $C_1$ - $C_6$ -alkyl or, together with the nitrogen atom to which they are attached, form a morpholino, piperidinyl, or pyrazinyl group;

n is 2, 3, or 4;

p is 0 or 1; and

q is 0, 1, or 2.

[Prior art Literatures]

[Patent Literatures]

**[0010]**

[Patent Literature 1] WO2012/028332  
 [Patent Literature 2] WO2013/074633  
 [Patent Literature 3] WO2012/011548  
 [Patent Literature 4] WO2006/116713  
 [Patent Literature 5] WO2007/146824  
 [Patent Literature 6] WO 2009/137429

[Non-Patent Literatures]

**[0011]**

[Non-Patent Literature 1] Clinical Science, Vol. 122, p. 361-368, 2012  
 [Non-Patent Literature 2] Proceedings of the national academy of sciences of the United States of America, Vol. 107, No. 3, p. 1124-1129, 2010  
 [Non-Patent Literature 3] Blood, Vol. 121, p. 2064-2073, 2013


[Summary of Invention]

[Technical Problem]

**[0012]** A problem to be solved by the present invention is to find a compound having an Axl inhibitory activity, which is useful for treatment of cancer such as AML, and to provide the compound as pharmaceuticals whose side effects are reduced.

[Solution to Problem]


**[0013]** In order to solve the above-mentioned problem, the inventors of the present invention have keenly studied to find a compound strongly inhibiting Axl. As a result, surprisingly, the inventors have found that a bicyclic structure, in which a saturated carbon ring is fused to a pyridone ring, represented by the following structural formula:



improves the Axl inhibitory activity, and have completed the present invention.

**[0014]** That is to say, the present invention relates to:

[1] a compound represented by general formula (I)



wherein  $R^1$  represents (1) a C1-8 alkyl group optionally substituted with one to five  $R^{11}$ , (2) a C3-7 carbon ring optionally substituted with one to five  $R^{12}$ , or (3) a 4- to 7-membered heterocycle optionally substituted with one to five  $R^{13}$ , wherein when the C1-8 alkyl group represented by  $R^1$  is a branched alkyl group, the C1-3 alkyl group branched from the same carbon atom, together with the carbon atom bound thereto, optionally forms a saturated C3-7 carbon ring,

$R^2$  represents (1) a C1-4 alkyl group, (2) a halogen atom, (3) a C1-4 haloalkyl group, (4) an oxo group, (5) an  $-OR^{21}$  group, or (6) an  $=NR^{22}$  group,

$R^3$  represents (1) a C1-4 alkyl group, (2) a halogen atom, or (3) a C1-4 haloalkyl group,

$R^4$  represents (1) a C1-4 alkoxy group, (2) a C1-4 haloalkyl group, (3) an  $-OR^{41}$  group, (4) a C1-4 alkyl group, (5) a C2-4 alkenyloxy group, or (6) a C2-4 alkynyloxy group,

$R^5$  represents (1) a hydrogen atom, (2) a C1-4 alkyl group, (3) a halogen atom, (4) a C1-4 haloalkyl group, or (5) an  $-OR^{21}$  group,

$R^{11}$  represents (1) an  $-OR^{101}$  group, (2) an  $SO_2R^{102}$  group, (3) an  $NR^{103}R^{104}$  group, or (4) a C3-7 carbon ring optionally substituted with one to three halogen atoms,

$R^{12}$  represents (1) a C1-8 alkyl group optionally substituted with a hydroxyl group, or (2) a halogen atom,

$R^{13}$  represents (1) a C1-8 alkyl group optionally substituted with a hydroxyl group, or (2) a halogen atom,

$R^{21}$  represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,

$R^{22}$  represents (1) a hydroxyl group, or (2) C1-4 alkoxy group,

$R^{41}$  represents

(1) a hydrogen atom;

(2) a C1-8 alkyl group substituted with one to two substituents selected from the group consisting of (a) 5- to 7-membered cyclic group optionally substituted with one to two substituents selected from the group consisting of (i) a C1-4 alkyl group, (ii) a C1-4 haloalkyl group, and (iii) a halogen atom, (b)  $NR^{401}R^{402}$ , (c) a hydroxyl group, and (d) an  $SO_2R^{403}$  group;

(3) a C2-8 alkenyl group substituted with one to two substituents selected from the group consisting of (a) 5- to 7-membered cyclic group optionally substituted with one to two substituents selected from the group consisting of (i) a C1-4 alkyl group, (ii) a C1-4 haloalkyl group, and (iii) a halogen atom, (b)  $NR^{401}R^{402}$ , (c) a hydroxyl group, and (d) an  $SO_2R^{403}$  group; or

(4) a C2-8 alkynyl group substituted with one to two substituents selected from the group consisting of (a) 5- to 7-membered cyclic group optionally substituted with one to two substituents selected from the group consisting of (i) a C1-4 alkyl group, (ii) a C1-4 haloalkyl group, and (iii) a halogen atom, (b)  $NR^{401}R^{402}$ , (c) a hydroxyl group, and (d) an  $SO_2R^{403}$  group,

$R^{101}$  represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,

$R^{102}$  represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,

$R^{103}$  and  $R^{104}$  each independently represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,

$R^{401}$  and  $R^{402}$  each independently represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,

$R^{403}$  represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,

A represents (1) CH, or (2) a nitrogen atom,

L represents (1)  $-O-$ , (2)  $-NH-$ , (3)  $-C(O)-$ , (4)  $-CR^6R^7-$ , (5)  $-S-$ , (6)  $-S(O)-$ , or (7)  $-S(O)_2-$ ,

$R^6$  and  $R^7$  each independently represents (1) a hydrogen atom, (2) a halogen atom, (3) a C1-4 alkyl group, (4)

a hydroxyl group, or (5)  $\text{NH}_2$ ,  
 ring1 represents benzene or pyridine,  
 represents a single bond or a double bond,  
 m is an integer from 0 to 5,  
 n is an integer from 0 to 5,  
 p is an integer from 0 to 2,  
 q is an integer from 0 to 4,


when m is two or more, a plurality of  $\text{R}^2$  may be the same as or different from each other, and when two of  $\text{R}^2$  represent a C1-3 alkyl group and are on the same carbon atom, the  $\text{R}^2$ , together with a carbon atom bound thereto, may form a saturated C3-7 carbon ring, when n is two or more, a plurality of  $\text{R}^3$  may be the same as or different from each other, and

when q is two or more, a plurality of  $\text{R}^4$  may be the same as or different from each other, a salt thereof, a solvate thereof, or an N-oxide thereof,

[2] the compound according to the above-mentioned [1], wherein m is one or more, and one of two  $\text{R}^2$  is necessarily an oxo group,

[3] the compound according to the above-mentioned [1] or [2], wherein L is (1) -O-, (2) -NH-, or (3) -C(O)-,

[4] the compound according the above-mentioned [1], which is represented by general formula (1-1)



wherein  $\text{R}^{2-1}$  represents (1) a C1-4 alkyl group, (2) a halogen atom, (3) a C1-4 haloalkyl group, (4) an  $-\text{OR}^{21}$  group, or (5) an  $=\text{NR}^{22}$  group,

m-1 is an integer from 0 to 4,

$\text{L}^1$  is (1) -O-, (2) -NH-, or (3) -C(O)-,

ring 1-1 represents benzene or pyridine,

when m-1 is two or more, a plurality of  $\text{R}^{2-1}$  may be the same as or different from each other, and when two of  $\text{R}^{2-1}$  represent a C1-3 alkyl group and are on the same carbon atom, the  $\text{R}^{2-1}$ , together with a carbon atom bound thereto, may form a saturated C3-7 carbon ring,

and other symbols have the same meanings as defined in the above-mentioned [1],

[5] the compound according to any one of the above-mentioned [1] to [4], which is: (2) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-7,7-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (3) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (4) N-{5-[(7-[3-(4-morpholinyl)propoxy]-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (5) N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]-3-fluorophenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (6) N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]phenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (7) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(4-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (8) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (9) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (10) N-{5-[(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (11) N-{5-[(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinyl}-1-(4-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (12) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(2S)-1-hydroxy-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (13) N-{4-[(6,7-



dimethoxy-4-quinolinyl)oxy]-3-fluorophenyl)-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (14) N-[5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl]-6,6-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (15) N-[5-[(6-methoxy-7-[3-(4-morpholinyl)propoxy]-4-quinolinyl)oxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (16) N-(5-[7-(3-hydroxy-3-methylbutoxy)-6-methoxy-4-quinolinyl]oxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, or (17) N-[5-[(6-methoxy-7-[3-(1-pyrrolidinyl)propoxy]-4-quinolinyl)oxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

[6] a pharmaceutical composition containing a compound represented by general formula (I) as defined in the above-mentioned [1], a salt thereof, a solvate thereof, or an N-oxide thereof,

[7] the pharmaceutical composition according to the above-mentioned [6], which is an Axl inhibitor,

[8] the pharmaceutical composition according to the above-mentioned [6], which is an agent for preventing and/or treating an Axl-related disease,

[9] the pharmaceutical composition according to the above-mentioned [8], wherein the Axl-related diseases includes cancer, kidney diseases, immune system diseases, or circulatory system diseases,

[10] the pharmaceutical composition according to the above-mentioned [9], wherein the cancer is acute myeloid leukemia, chronic myeloid leukemia, acute lymphatic leukemia, melanoma, breast cancer, pancreatic cancer, glioma, esophageal adenocarcinoma, large intestine cancer, renal cell carcinoma, thyroid cancer, non-small cell lung cancer, prostate cancer, stomach cancer, liver cancer, uveal malignant melanoma, ovarian cancer, endometrial cancer, lymphoma, head and neck cancer, or sarcoma,

[11] the pharmaceutical composition according to the above-mentioned [6], which is a metastasis suppressing agent for cancer cells,

[12] a compound represented by general formula (I) according to the above-mentioned [1], a salt thereof, a solvate thereof, or an N-oxide thereof for use in preventing and/or treating an Axl-related disease,

[13] the compound for use according to the above-mentioned [12], wherein the Axl-related disease includes a cancer, a kidney disease, an immune system disease, or a circulatory system disease, and

[14] the compound for use according to the above-mentioned [13], wherein the cancer is acute myeloid leukemia, chronic myeloid leukemia, acute lymphatic leukemia, melanoma, breast cancer, pancreatic cancer, glioma, esophageal adenocarcinoma, large intestine cancer, renal cell carcinoma, thyroid cancer, non-small cell lung cancer, prostate cancer, stomach cancer, liver cancer, uveal malignant melanoma, ovarian cancer, endometrial cancer, lymphoma, head and neck cancer, or sarcoma.

#### [Effects of Invention]

**[0015]** A compound of the present invention has a strong Axl inhibitory activity, has an Axl-selective inhibitory activity to a specific kinase, and has reduced CYP inhibitory effect, and therefore is useful as a therapeutic drug for acute myeloid leukemia, has less side effect and has little concern about drug interaction.

#### [Description of Embodiments]

**[0016]** The present invention will be described in detail hereinafter.

**[0017]** In the present invention, a halogen atom denotes a fluorine atom, a chlorine atom, a bromine atom, and an iodine atom.

**[0018]** In the present invention, the C1-8 alkyl group includes a straight or branched C1-8 alkyl group. Examples thereof include methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, isopropyl, isobutyl, sec-butyl, tert-butyl, and an isomer thereof.

**[0019]** In the present invention, the C1-4 alkyl group includes a straight or branched C1-4 alkyl group. Examples thereof include methyl, ethyl, propyl, butyl, isopropyl, isobutyl, sec-butyl, and tert-butyl.

**[0020]** In the present invention, the C1-3 alkyl group includes a straight or branched C1-3 alkyl group. Examples thereof include a methyl group, an ethyl group, a propyl group, and an isopropyl.

**[0021]** In the present invention, the C1-4 haloalkyl group denotes, for example, a fluoromethyl group, a chloromethyl group, a bromomethyl group, an iodomethyl group, a difluoromethyl group, a trifluoromethyl group, a 1-fluoroethyl group, a 2-fluoroethyl group, a 2-chloroethyl group, a pentafluoroethyl group, a 1-fluoropropyl group, a 2-chloropropyl group, a 3-fluoropropyl group, a 3-chloropropyl group, a 4,4,4-trifluorobutyl group, and a 4-bromobutyl group.

**[0022]** In the present invention, the C2-8 alkenyl group denotes, for example, a vinyl group, a propenyl group, a butenyl group, a pentenyl group, a hexenyl group, a heptenyl group, and an octenyl group, and an isomer thereof.

**[0023]** In the present invention, the C2-8 alkynyl group denotes, for example, an ethynyl group, a propynyl group, a butynyl group, a pentynyl group, a hexynyl group, a heptynyl group, an octynyl group, and an isomer thereof.

**[0024]** In the present invention, examples of the C1-4 alkoxy group include a methoxy group, an ethoxy group, a

propoxy group, an isopropoxy group, a butoxy group, an isobutoxy group, a sec-butoxy group, or a tert-butoxy group.

**[0025]** In the present invention, the C2-4 alkenyloxy group denotes, for example, vinyloxy, propenyloxy, butenyloxy, and an isomer thereof.

**[0026]** In the present invention, the C2-4 alkynyloxy group denotes, for example, ethynyloxy, propynyloxy, butynyloxy, and an isomer thereof.

**[0027]** In the present invention, the C3-7 carbon ring denotes a C3-7 monocyclic carbon ring, and a carbon ring which may be partially or completely saturated, and examples thereof include cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, cyclobutene, cyclopentene, cyclohexene, cycloheptene, cyclobutadiene, cyclopentadiene, cyclohexadiene, cycloheptadiene, or benzene ring.

**[0028]** In the present invention, the C5-7 carbon ring denotes a C5-7 monocyclic carbon ring, and a carbon ring which may be partially or completely saturated, and examples thereof include cyclopentane, cyclohexane, cycloheptane, cyclopentene, cyclohexene, cycloheptene, cyclopentadiene, cyclohexadiene, cycloheptadiene, or benzene ring.

**[0029]** In the present invention, examples of the saturated C3-7 carbon ring include cyclopropane, cyclobutane, cyclopentane, cyclohexane, and cycloheptane.


**[0030]** In the present invention, the 4- to 7-membered heterocycle denotes 4- to 7-membered monocyclic heterocycle, which includes one to five heteroatoms selected from an oxygen atom, a nitrogen atom and a sulfur atom, and a part or all of which is saturated. Example thereof include azetidine, pyrroline, pyrrolidine, imidazoline, imidazolidine, triazoline, triazolidine, tetrazoline, tetrazolidine, pyrazoline, pyrazolidine, dihydropyridine, tetrahydropyridine, piperidine, dihydropyrazine, tetrahydropyrazine, piperazine, dihydropyrimidine, tetrahydropyrimidine, perhydropyrimidine, dihydropyridazine, tetrahydropyridazine, perhydropyridazine, dihydroazepine, tetrahydroazepine, perhydroazepine, dihydrodiazepine, tetrahydrodiazepine, perhydrodiazepine, oxetan, dihydrofuran, tetrahydrofuran, dihydropyran, tetrahydropyran, dihydrooxepine, tetrahydrooxepine, perhydrooxepine, thietane, dihydrothiophene, tetrahydrothiophene, dihydrothiopyran, tetrahydrothiopyran, dihydrothiepin, tetrahydrothiepin, perhydrothiepin, dihydrooxazole, tetrahydrooxazole (oxazolidine), dihydroisoxazole, tetrahydroisoxazole (isoxazolidine), dihydrothiazole, tetrahydrothiazole (thiazolidine), dihydroisothiazole, tetrahydroisothiazole (isothiazolidine), dihydrofurazan, tetrahydrofurazan, dihydrooxadiazole, tetrahydrooxadiazole, (oxadiazolidine), dihydrooxazine, tetrahydrooxazine, dihydrooxadiazine, tetrahydrooxadiazine, dihydrooxazepine, tetrahydrooxazepine, perhydrooxazepine, dihydrooxadiazepine, tetrahydrooxadiazepine, perhydrooxadiazepine, dihydrothiadiazole, tetrahydrothiadiazole (thiadiazolidine), dihydrothiazine, tetrahydrothiazine, dihydrothiadiazine, tetrahydrothiadiazine, dihydrothiazepine, tetrahydrothiazepine, perhydrothiazepine, dihydrothiadiazepin, tetrahydrothiadiazepin, perhydrothiadiazepin, morpholine, thiomorpholine, oxathiane, dioxolane, dioxane, dithiolane, dithiane, pyrrole, imidazole, triazole, tetrazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, azepine, diazepine, furan, pyran, oxepin, thiophene, thiopyran, thiepin, oxazole, isoxazole, thiazole, isothiazole, furazan, oxadiazole, oxazine, oxadiazine, oxazepine, oxadiazepine, thiadiazole, thiazine, thiadiazine, thiazepine, or thiadiazepin ring.

**[0031]** In the present invention, the 5- to 7-membered cyclic group denotes C5-7 carbon ring and 5- to 7-membered heterocycle. Herein, the C5-7 carbon ring has the same meaning as defined above, the 5- to 7-membered heterocycle includes 5- to 7-membered unsaturated heterocycle and 5- to 7-membered saturated heterocycle. Examples of 5- to 7-membered heterocycle include pyrroline, pyrrolidine, imidazoline, imidazolidine, triazoline, triazolidine, tetrazoline, tetrazolidine, pyrazoline, pyrazolidine, dihydropyridine, tetrahydropyridine, piperidine, dihydropyrazine, tetrahydropyrazine, piperazine, dihydropyrimidine, tetrahydropyrimidine, perhydropyrimidine, dihydropyridazine, tetrahydropyridazine, perhydropyridazine, dihydroazepine, tetrahydroazepine, perhydroazepine, dihydrodiazepine, tetrahydrodiazepine, perhydrodiazepine, dihydrofuran, tetrahydrofuran, dihydropyran, tetrahydropyran, dihydrooxepine, tetrahydrooxepine, perhydrooxepine, dihydrothiophene, tetrahydrothiophene, dihydrothiopyran, tetrahydrothiopyran, dihydrothiepin, tetrahydrothiepin, perhydrothiepin, dihydrooxazole, tetrahydrooxazole (oxazolidine), dihydroisoxazole, tetrahydroisoxazole (isoxazolidine), dihydrothiazole, tetrahydrothiazole (thiazolidine), dihydroisothiazole, tetrahydroisothiazole (isothiazolidine), dihydrofurazan, tetrahydrofurazan, dihydrooxadiazole, tetrahydrooxadiazole, (oxadiazolidine), dihydrooxazine, tetrahydrooxazine, dihydrooxadiazine, tetrahydrooxadiazine, dihydrooxazepine, tetrahydrooxazepine, perhydrooxazepine, dihydrooxadiazepine, tetrahydrooxadiazepine, perhydrooxadiazepine, dihydrothiadiazole, tetrahydrothiadiazole (thiadiazolidine), dihydrothiazine, tetrahydrothiazine, dihydrothiadiazine, tetrahydrothiadiazine, dihydrothiazepine, tetrahydrothiazepine, perhydrothiazepine, dihydrothiadiazepin, tetrahydrothiadiazepin, perhydrothiadiazepin, morpholine, thiomorpholine, oxathiane, dioxolane, dioxane, dithiolane, dithiane, pyrrole, imidazole, triazole, tetrazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, azepine, diazepine, furan, pyran, oxepin, thiophene, thiopyran, thiepin, oxazole, isoxazole, thiazole, isothiazole, furazan, oxadiazole, oxazine, oxadiazine, oxazepine, oxadiazepine, thiadiazole, thiazine, thiadiazine, thiazepine, or thiadiazepin ring.


**[0032]** In the present invention, the 6-membered cyclic group denotes C6 carbon ring and 6-membered heterocycle. Examples thereof include cyclohexane, cyclohexene, cyclohexadiene, benzene, pyridine, pyrazine, pyrimidine, pyridazine, pyran, thiopyran, oxazine, oxadiazine, thiazine, thiadiazine, dihydropyridine, tetrahydropyridine, piperidine, dihydropyrazine, tetrahydropyrazine, piperazine, dihydropyrimidine, tetrahydropyrimidine, perhydropyrimidine, dihydropyridazine, tetrahydropyridazine, perhydropyridazine, dihydropyran, tetrahydropyran, dihydrothiopyran, tetrahydrothi-

opyran, dihydrooxazine, tetrahydrooxazine, dihydrooxadiazine, tetrahydrooxadiazine, dihydrothiazine, tetrahydrothiazine, dihydrothiadiazine, tetrahydrothiadiazine, morpholine, thiomorpholine, oxathiane, dioxane, and dithiane ring.

**[0033]** In the present invention, "when the C1-8 alkyl group represented by  $R^1$  is a branched alkyl group, the C1-3 alkyl group branched from the same carbon atom optionally forms a saturated C3-7 carbon ring together" denotes that in a partial structure of the following general formula (I):




(wherein all of the symbols have the same meanings as defined above), for example, when  $R^1$  is a branched alkyl chain as represented in the above-mentioned general formula, the alkyl chain branched from the same carbon atom, together with the carbon atom bound thereto, forms a saturated carbon ring, as shown in the following general formula:




(wherein all of the symbols have the same meanings as defined above).

**[0034]** In the present invention, "when two of  $R^2$  represent a C1-3 alkyl group and are on the same carbon atom, the  $R^2$ , together with a carbon atom bound thereto, may form a saturated C3-7 carbon ring" denotes that in a partial structure of the following general formula (I):



(wherein all of the symbols have the same meanings as defined above), for example, when  $R^2$  is an alkyl group as represented in the above-mentioned general formula and are on the same carbon atom, the  $R^2$ , together with a carbon atom bound thereto, forms a saturated carbon ring, as shown in the following general formula:



(wherein all of the symbols have the same meanings as defined above).

**[0035]** In the present invention, "when two of  $R^{2-1}$  represent a C1-3 alkyl group and are on the same carbon atom, the  $R^{2-1}$ , together with a carbon atom bound thereto, may form a saturated C3-7 carbon ring" has the same definition as that for  $R^2$  in the phrase: "when two of  $R^2$  represent a C1-3 alkyl group and are on the same carbon atom, the  $R^2$ , together with a carbon atom bound thereto, may form a saturated C3-7 carbon ring."

**[0036]** In the present invention, it is preferable that  $m$  is one or more, and one of  $R^2$  is necessarily an oxo group.


**[0037]** In the present invention, it is preferable that  $A$  is CH.

**[0038]** In the present invention, it is preferable that  $R^4$  is a C1-4 alkoxy group or an  $-OR^{41}$  group.

**[0039]** In the present invention, it is preferable that  $L$  is  $-O-$ ,  $-NH-$ , or  $-C(O)-$ .

**[0040]** In the present invention, ring 1 is benzene or pyridine.

**[0041]** In the present invention, it is preferable that the compound represented by general formula (I) is a compound represented by general formula (I-1):



(wherein all of the symbols have the same meanings as defined above).

**[0042]** In the present invention, it is preferable that two binding arms in the ring1 and ring1-1 are bound to a para position.

**[0043]** In the present invention, in general formula (I-1),  $A$  is preferably CH, and  $R^4$  is preferably a C1-4 alkoxy group or an  $-OR^{41}$  group.




**[0044]** In the present invention, preferable compounds preferably include the compounds described in Examples, and the following (2) to (17) are more preferable: (2) N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-7,7-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (3) N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (4) N-{5-[(7-[3-(4-morpholinyl)propoxy]-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (5) N-{4-[(6,7-dimethoxy-4-quinolinyloxy)-3-fluorophenyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (6) N-{4-[(6,7-dimethoxy-4-quinolinyloxy)phenyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (7) N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-1-(4-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (8) N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (9) N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-1-(2-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (10) N-{5-[(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (11) N-{5-[(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinyl}-1-(4-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (12) N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-1-[(2S)-1-hydroxy-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (13) N-{4-[(6,7-dimethoxy-4-quinolinyloxy)-3-fluorophenyl]-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (14) N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-6,6-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (15) N-{5-[(6-methoxy-7-[3-(4-morpholinyl)propoxy]-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (16) N-{5-[(7-[3-hydroxy-3-methylbutoxy]-6-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, (17) N-{5-[(6-methoxy-7-[3-(1-pyrrolidinyl)propoxy]-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide.

[Isomer]

**[0045]** In the present invention, unless specifically directed, all of the isomers are included. For example, an alkyl group includes straight chain and branched chain groups. In addition, all of geometrical isomers of double bonds, rings,

and fused rings (E-, Z-, cis-, trans-isomers), optical isomers by the presence of an asymmetric carbon atom (R-, S-isomer,  $\alpha$ -,  $\beta$ -configurations, enantiomers, diastereomers), optical active isomers having optical rotation property (D, L, d, l-isomers), polar isomers according to chromatographic separation (more polar isomer, less polar isomer), equilibrium compound, rotamers, mixtures thereof at any rate, and racemic mixtures are included in the present invention. Furthermore, the present invention also encompasses all isomers by tautomers.

**[0046]** Furthermore, the optical isomer of the present invention is not only limited to an optical isomer having purity of 100%, but also may include other optical isomers having purity of less than 50%.

**[0047]** In the present invention, unless otherwise noted, as apparent to a person skilled in the art, a symbol:  represents binding toward the back side of the plane of the paper (that is to say, the  $\alpha$ -configuration),  represents binding toward the front side of the plane of the paper (that is to say, the  $\beta$ -configuration), and  represents  $\alpha$ -configuration,  $\beta$ -configuration or an arbitrary mixture thereof.

**[0048]** The compound represented by general formula (I) is converted into a corresponding salt by the well-known method. A salt is preferably a water-soluble salt. Examples of a suitable salt include salts of an alkali metal (e.g., potassium, sodium), salts of an alkaline earth metal (e.g., calcium, magnesium), ammonium salts, or salts of a pharmaceutically acceptable organic amine (e.g., tetramethylammonium, triethylamine, methylamine, dimethylamine, cyclopentylamine, benzylamine, phenethylamine, piperidine, monoethanolamine, diethanolamine, tris(hydroxymethyl)aminomethane, lysine, arginine, N-methyl-D-glucamine), acid addition salts (inorganic acid salts (e.g., hydrochloride, hydrobromide, hydroiodide, sulfate, phosphate, nitrate), organic acid salts (e.g., acetate, trifluoroacetate, lactate, tartrate, oxalate, fumarate, maleate, benzoate, citrate, methanesulfonate, ethanesulfonate, benzenesulfonate, toluenesulfonate, isethionate, glucuronate, gluconate)).


**[0049]** The compound represented by general formula (I) and a salt thereof can be also converted into a solvate. It is preferable that the solvate is low-toxic and water-soluble. Examples of a suitable solvate include solvates with water, or an alcoholic solvent (for example, ethanol).

**[0050]** The N-oxide of the compound represented by general formula (I) denotes compounds represented by general formula (I) in which a nitrogen atom is oxidized. Furthermore, the N-oxide of the compound represented by general formula (I) may be salts of alkali (earth) metal salt, ammonium salt, organic amine salt, and acid addition salt mentioned above.

[Process for producing compound of the present invention]


**[0051]** The compound of the present invention can be produced by the well-known methods, for example, the method described in Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2nd Edition (Richard C. Larock, John Wiley & Sons Inc, 1999), or methods described in Examples, with appropriate modification and in combination thereof.

**[0052]** A compound represented by general formula (I) wherein L is an oxygen atom, and R<sup>2</sup> is an oxo group, that is, a compound represented by general formula (I-A):



(wherein all of the symbols have the same meanings as defined above) can be produced by the process represented by the following reaction process schemes 1 and 2:

## Reaction process scheme 1




(wherein each of  $X^1$  and  $X^2$  independently represents a halogen atom,  $X^1$  and  $X^2$  may be the same as or different from each other, and the other symbols have the same meanings as defined above).

**[0053]** In the reaction process scheme 1, the reaction 1 can be carried out by subjecting a compound represented by general formula (a) and a compound represented by general formula (II) to the aromatic nucleophilic substitution reaction. The aromatic nucleophilic substitution reaction is well known, and is carried out, for example, in an organic solvent (e.g., chlorobenzene, N,N-dimethyl sulfoxide, N,N-dimethyl acetamide, N,N-dimethylformamide, chloroform, dichloromethane, diethyl ether, tetrahydrofuran, methyl t-butyl ether), in the presence or absence of a catalyst (e.g., 4-dimethylaminopyridine (DMAP)), and in the presence or absence of a base (e.g., sodium hydride, triethylamine, cesium carbonate), at 0 to 200°C.

**[0054]** In the reaction process scheme 1, the reaction 2 is carried out by reacting a compound represented by general formula (b) in an organic solvent (e.g., tetrahydrofuran), in the presence of a palladium catalyst (e.g., tris(dibenzylideneacetone)dipalladium(0) chloroform complex), in the presence of a base (e.g., lithium bis(trimethylsilyl)amide (LHMDS), potassium bis(trimethylsilyl)amide (KHMDS), sodium bis(trimethylsilyl)amide (NaHMDS)), a phosphine compound (e.g., 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (S-Phos), tri-tert-butylphosphine ( $P(t-Bu)_3$ )) at 0 to 100°C, and then reacting by adding inorganic acid (e.g., hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, nitric acid) at 0 to 150°C. Alternatively, the production method for aryl amine described in Organic Letters, Vol 3, No. 17, p.2729-2732, 2001 can be employed.

## Reaction process scheme 2



(wherein  $R^d$  represents a C1-4 alkyl group, and the other symbols have the same meanings as defined above).

**[0055]** In the reaction process scheme 2, the reaction 3 is carried out by a reaction of a compound represented by general formula (d) and a compound represented by general formula (III). The reaction is well known, and, for example, can be carried out at 0 to 100°C in an organic solvent (e.g., N,N-dimethyl formamide), in the presence of a base (e.g., tert-butoxy potassium).

**[0056]** In the reaction process scheme 2, the reaction 4 can be carried out by subjecting a compound represented by general formula (e) and a compound represented by general formula (IV) to addition reaction. The reaction is well known, and, for example, can be carried out by reacting at 0 to 100°C in an alcohol solvent (e.g., methanol, ethanol).

**[0057]** In the reaction process scheme 2, the reaction 5 can be carried out by using and subjecting the compound

represented by general formula (c) and the compound represented by general formula (f) to an amidation reaction. The amidation reaction is well known, and examples thereof include:

- (1) a method using an acid halide,
- (2) a method using a mixed acid anhydride, and
- (3) a method using a condensing agent.

**[0058]** These methods are specifically described below:


(1) The method using an acid halide is carried out, for example, by reacting a carboxylic acid with an acid halogenating agent (e.g., oxalyl chloride, thionyl chloride) in an organic solvent (e.g., chloroform, dichloromethane, diethyl ether, tetrahydrofuran) at -20°C to reflux temperature, and then reacting the obtained acid halide in the presence of a base (e.g., pyridine, triethylamine, dimethylaniline, dimethylaminopyridine, diisopropylethylamine) in amine and an organic solvent (e.g., chloroform, dichloromethane, diethyl ether, tetrahydrofuran) at 0 to 40°C. Additionally, the method can be also carried out by reacting the obtained acid halide with an amine at 0 to 40°C by using an alkaline aqueous solution (e.g., sodium bicarbonate water or sodium hydroxide solution) in an organic solvent (e.g., dioxane, tetrahydrofuran).

(2) The method using a mixed acid anhydride is carried out, for example, by reacting carboxylic acid with an acid halide (e.g., pivaloyl chloride, tosyl chloride, mesyl chloride) or an acid derivative (e.g., ethyl chloroformate, isobutyl chloroformate) in the presence of a base (e.g., pyridine, triethylamine, dimethylaniline, dimethylaminopyridine, diisopropylethylamine) in an organic solvent (e.g., chloroform, dichloromethane, diethyl ether, tetrahydrofuran) or in the absence of any solvent at 0 to 40°C, and then reacting the obtained mixed acid anhydride with amine in an organic solvent (e.g., chloroform, dichloromethane, diethyl ether, tetrahydrofuran) at 0 to 40°C.

(3) The method using a condensing agent is carried out, for example, by reacting a carboxylic acid with an amine in an organic solvent (e.g., chloroform, dichloromethane, dimethyl formamide, diethyl ether, tetrahydrofuran) or in the absence of any solvent at 0 to 40°C in the presence or absence of a base (e.g., diisopropylethylamine (DIPEA), pyridine, triethylamine, dimethylaniline, dimethylaminopyridine), using a condensing agent (e.g., O-(7-Aza-1-benzotriazolyl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), (1,3-dicyclohexylcarbodiimide (DCC), 1-ethyl-3-[3-(dimethylamino)propyl]carbodiimide (EDC), 1,1'-carbonyldiimidazole (CDI), 2-chloro-1-methylpyridinium iodine, 1-propylphosphonic acid cyclic anhydride (PPA)) and using, or not using, 1-hydroxybenzotriazole (HOBt).

**[0059]** These reactions (1), (2), and (3) are desirably carried out under an inert gas (e.g., argon, nitrogen) atmosphere in anhydrous conditions.

**[0060]** Furthermore, the compound represented by general formula (c) can be also produced by the reaction process scheme 3.



(wherein all of the symbols have the same meanings as defined above).

**[0061]** In the reaction process scheme 3, a compound represented by general formula (b)' can be produced by the same method as in the above-mentioned reaction 1 using a compound represented by general formula (a)' and the compound represented by general formula (V).

**[0062]** In the reaction process scheme 3, the reaction 6 can be carried out by subjecting the compound represented by general formula (b)' to a reduction reaction of the nitro group. The reduction reaction of a nitro group is well known, and can be carried out by, for example, the following method.

(1) The reduction reaction is carried out in, for example, a solvent [ethers (e.g., tetrahydrofuran, dioxane, dimethoxyethane, diethyl ether), alcohols (e.g., methanol, ethanol), benzenes (e.g., benzene, toluene), ketones (e.g., acetone, methyl ethyl ketone), nitriles (e.g., acetonitrile), amides (e.g., dimethylformamide), water, ethyl acetate,

acetic acid or a mixture solvent of two or more thereof] in the presence of a hydrogenation catalyst (e.g., palladium-carbon, palladium black, palladium, palladium hydroxide, platinum dioxide, platinum-carbon, nickel, Raney-nickel, ruthenium chloride), in the presence or absence of acids (e.g., hydrochloric acid, sulfuric acid, hypochlorous acid, boric acid, tetrafluoroboric acid, acetic acid, p-toluenesulfonic acid, oxalic acid, trifluoroacetic acid, formic acid), at normal pressure or reduced pressure under a hydrogen atmosphere, in the presence of formic acid ammonium or in the presence of hydrazine, at 0 to 200°C.

(2) The reaction is carried out, for example, in a water-miscible solvent (e.g., ethanol, methanol, tetrahydrofuran) in the presence or absence of an acid (e.g., hydrochloric acid, hydrobromic acid, ammonium chloride, acetic acid, ammonium formate) using a metal reagent (e.g., zinc, iron, tin, tin chloride, iron chloride, samarium, indium, sodium borohydride-Nickel chloride) at 0 to 150°C.

**[0063]** In the reaction process schemes 1 to 3, when a compound represented by each general formula includes a protective group, a deprotection reaction can be carried out if necessary. The deprotection reaction of the protective group is known, and can be carried out by the methods mentioned below. Examples thereof include: (1) deprotection reactions by alkaline hydrolysis, (2) deprotection reaction in acidic conditions, (3) deprotection reaction by hydrogenolysis, (4) deprotection reaction of a silyl group, (5) deprotection reaction using metal, (6) deprotection reaction using a metal complex.

**[0064]** These methods are specifically described:

(1) The deprotection reaction by alkaline hydrolysis condition is carried out, for example, in an organic solvent (for example, methanol, tetrahydrofuran, dioxane) with hydroxide of alkaline metal (for example, sodium hydroxide, potassium hydroxide, lithium hydroxide), hydroxide of alkaline earth metal (for example, barium hydroxide, calcium hydroxide), or carbonate (for example, sodium carbonate or potassium carbonate), or an aqueous solution thereof or a mixture thereof at 0 to 40°C.

(2) The deprotection reaction in acidic conditions is carried out, for example, in an organic solvent (for example, dichloromethane, chloroform, dioxane, ethyl acetate, methanol, isopropyl alcohol, tetrahydrofuran, anisole), organic acid (for example, acetic acid, trifluoroacetic acid, methanesulfonic acid, p-tosyl acid), or inorganic acid (for example, hydrochloric acid, sulfuric acid), or a mixture thereof (for example, hydrogen bromide/acetic acid) in the presence or absence of 2,2,2-trifluoroethanol at 0 to 100°C.

(3) The deprotection reaction by hydrogenolysis is carried out, for example, in a solvent (for example, ethers (e.g., tetrahydrofuran, dioxane, dimethoxyethane, diethyl ether), alcohols (for example, methanol, ethanol), benzenes (for example, benzene, toluene), ketones (for example, acetone, methyl ethyl ketone), nitriles (for example, acetonitrile), amides (for example, N,N-dimethylformamide), water, ethyl acetate, acetic acid, or a mixture of two or more thereof) in the presence of a catalyst (for example, palladium-carbon, palladium black, palladium hydroxide-carbon, platinum oxide, Raney nickel) under hydrogen atmosphere at normal pressure or elevated pressure, or in the presence of ammonium formate at 0 to 200°C.

(4) The deprotection reaction of a silyl group is carried out, for example, in a water-miscible organic solvent (for example, tetrahydrofuran, acetonitrile), by using tetrabutylammonium fluoride at 0 to 40°C. The reaction is also carried out, for example, in organic acid (for example, acetic acid, trifluoroacetic acid, methanesulfonic acid, p-tosyl acid), or in inorganic acid (for example, hydrochloric acid, sulfuric acid) or a mixture thereof (for example, hydrogen bromide/acetic acid) at -10 to 100°C.

(5) The deprotection reaction using a metal is carried out, for example, in an acidic solvent (for example, acetic acid, a buffer of pH 4.2 to 7.2, a mixed solution of the solution and an organic solvent such as tetrahydrofuran) in the presence of powder zinc, if necessary, with an ultrasonic wave applied at 0 to 40°C.

(6) The deprotection reaction using a metal complex is carried out, for example, in an organic solvent (for example, dichloromethane, N,N-dimethylformamide, tetrahydrofuran, ethyl acetate, acetonitrile, dioxane, ethanol), water or a mixed solvent thereof in the presence of a trap reagent (for example, tributyltin hydride, triethylsilane, dimedone, morpholine, diethylamine, pyrrolidine), an organic acid (for example, acetic acid, formic acid, 2-ethylhexanoic acid) and/or in the presence of an organic acid salt (for example, sodium 2-ethylhexanoate, potassium 2-ethylhexanoate) in the presence or absence of a phosphine reagent (for example, triphenylphosphine) using a metal complex (for example, tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium (II), palladium acetate (II), chlorotris(triphenylphosphine)rhodium (I)) at 0 to 40°C.

**[0065]** In addition to the above-mentioned methods, the deprotection reaction can be carried out by the method described in for example, T. W. Greene, Protective Groups in Organic Synthesis, Wiley, New York, 1999.

**[0066]** Examples of a protective group for a hydroxyl group include a methyl group, a trityl group, a methoxymethyl (MOM) group, a 1-ethoxyethyl (EE) group, a methoxyethoxymethyl (MEM) group, a 2-tetrahydropyranyl (THP) group, a trimethylsilyl (TMS) group, a triethylsilyl (TES) group, a t-butyldimethylsilyl (TBDMS) group, a t-butyldiphenylsilyl (TB-



DPS) group, an acetyl (Ac) group, a pivaloyl group, a benzoyl group, a benzyl (Bn) group, a p-methoxybenzyl group, an allyloxycarbonyl (Alloc) group, and a 2,2,2-trichloroethoxycarbonyl (Troc) group.

**[0067]** Examples of a protective group for an amino group include a benzyloxycarbonyl group, a t-butoxycarbonyl group, an allyloxycarbonyl (Alloc) group, a 1-methyl-1-(4-biphenyl)ethoxycarbonyl (Bpoc) group, a trifluoroacetyl group, a 9-fluororenylmethoxycarbonyl group, a benzyl (Bn) group, a p-methoxybenzyl group, a benzyloxymethyl (BOM) group, and a 2-(trimethylsilyl)ethoxymethyl (SEM) group.

**[0068]** The protective groups for a hydroxyl group and an amino group are not particularly limited to the above-described groups, and groups are included, in addition to the above-mentioned groups, as long as the groups can be detached easily and selectively. For example, those described in Protective Groups in Organic Synthesis (T. W. Greene, John Wiley & Sons Inc., 1999) may be used.

**[0069]** In each reaction in the present specification, compounds used as starting raw material, for example, the compound represented by general formula (a), (a)', (d), (II), (III), (IV) or (V) is well known or can be produced by well-known methods.

**[0070]** In each reaction in the present specification, as apparent to the skilled persons in the art, the reactions involving heating can be carried out using a water bath, an oil bath, a sand bath or a microwave.

**[0071]** In each reaction in the present specification, a solid-supported reagent which is supported on a high molecular polymer (e.g., polystyrene, polyacrylamide, polypropylene, polyethylene glycol) may be appropriately used.

**[0072]** In each reaction in the present specification, the reaction products can be purified by conventional purification methods, for example, by distillation at normal or reduced pressure, by high performance liquid chromatography using silica gel or magnesium silicate, thin layer chromatography, ion-exchange resin, scavenger resin, or column chromatography, washing, or recrystallization. The purification may be done after each reaction or after several reactions.

[Toxicity]

**[0073]** The toxicity of the compound of the present invention is sufficiently low, and the compound can be safely used as pharmaceuticals.

[Application to pharmaceuticals]

**[0074]** Since the compound of the present invention has an Axl inhibitory activity, it can be used as an agent for preventing and/or treating an Axl-related disease in mammals, especially in human.

**[0075]** In the present invention, examples of the Axl-related diseases include cancer, kidney diseases, immune system disease, and circulatory system disease.

**[0076]** In the present invention, the cancer includes acute myeloid leukemia, chronic myeloid leukemia, acute lymphatic leukemia, melanoma, breast cancer, pancreatic cancer, glioma, esophageal adenocarcinoma, large intestine cancer, renal cell carcinoma, thyroid cancer, non-small cell lung cancer, prostate cancer, stomach cancer, liver cancer, uveal malignant melanoma, ovarian cancer, endometrial cancer, lymphoma, head and neck cancer, and sarcoma.

**[0077]** In the present invention, examples of the kidney diseases include glomerular nephritis, chronic nephritis, IgA nephritis, sequential (secondary) nephritis, nephrosis nephritis, acute renal failure, chronic renal failure, diabetic nephropathy, gouty nephropathy, interstitial nephritis, and nephropylitis.

**[0078]** In the present invention, examples of the immune system disease include psoriasis, and rheumatoid arthritis.

**[0079]** In the present invention, examples of the circulatory system disease include atherosclerosis and thrombosis.

**[0080]** Furthermore, since the compound of the present invention has an Axl inhibitory activity, it can be used as a metastasis suppressing agent to cancer cell.

**[0081]** The compound of the present invention may be administered as a combination drug in combination with other drugs in order to accomplish the following purposes:

- 1) to supplement and/or enhance the preventive and/or therapeutic effect of the compound;
- 2) to improve the kinetics, improvement of absorption, and reduction of the dose of the compound; and/or
- 3) to eliminate the side effects of the compound.

**[0082]** A combination drug of the compound of the present invention and other drugs may be administered in the form of a compounding agent including these components mixed into one formulation, or may be administered in separate formulations. Administration as separate formulations includes simultaneous administration and administration at different times. In the administration at different times, the compound of the present invention may be administered before the other drug. Alternatively, the other drug may be administered before the compound of the present invention. The method for the administration of these drugs may be the same as each other or different from each other.

**[0083]** Diseases on which the preventive and/or therapeutic effect of the above-mentioned combination drug works

are not particularly limited but may be those in which the preventive and/or therapeutic effect of the compound of the present invention is supplemented and/or enhanced.

**[0084]** The other drugs for supplementing and/or enhancing the preventive and/or therapeutic effect of the compound of the present invention against cancer include, for example, alkylating agents, antimetabolites, anticancer antibiotics, plant alkaloids, hormones, platinum compounds, anti-CD20 antibodies, anti-CD52 antibodies, anti-PD-1 antibodies, G-CSF formulations, acute promyelocytic leukemia differentiation-inducing agents, kinase inhibitors, topoisomerase inhibitors, aromatase inhibitors, and other anticancer drugs.

**[0085]** The other drug for supplementing and/or enhancing the preventive and/or therapeutic effect of the compound of the present invention against kidney diseases include, for example, steroids, immunosuppressants, angiotensin II antagonistic drugs, angiotensin-converting enzyme inhibitors, antiplatelet drugs, and anticoagulant drugs.

**[0086]** The other drugs for supplementing and/or enhancing the preventive and/or therapeutic effect of the compound of the present invention against immune system diseases include, for example, immunosuppressants, steroid, disease-modifying anti-rheumatic drugs, prostaglandins, prostaglandin synthase inhibitors, phosphodiesterase inhibitors, metalloprotease inhibitors, anti-cytokine protein formulations such as anti-TNF- $\alpha$  formulations, anti-IL-1 formulations, and anti-IL-6 formulation, cytokine inhibitors, and nonsteroidal anti-inflammatory agents.

**[0087]** The other drugs for supplementing and/or enhancing the preventive and/or therapeutic effect of the compound of the present invention against circulatory system diseases include antiplatelet drugs, angiotensin II antagonistic drugs, angiotensin-converting enzyme inhibitors, HMG-CoA reductase inhibitors, and thiazolidine derivatives.

**[0088]** Examples of the alkylating agents include nitrogen mustard N-oxide hydrochloride, cyclophosphamide, ifosfamide, melphalan, thiotepa, carboquone, busulfan, nimustine hydrochloride, dacarbazine, ranimustine, carmustine, chlorambucil, bendamustine, and mechlorethamine.

**[0089]** Examples of the antimetabolites include methotrexate, mercaptopurine, 6-mercaptopurine riboside, fluorouracil, tegafur, tegafur uracil, carmofur, doxifluridine, cytarabine, encitabine, tegafur gimestat otastat potassium, gemcitabine hydrochloride, cytarabine ocfosfate, procarbazine hydrochloride, and hydroxycarbamide.

**[0090]** Examples of the anticancer antibiotics include actinomycin D, mitomycin C, daunorubicin hydrochloride, doxorubicin hydrochloride, aclarubicin hydrochloride, neocarzinostatin, pirarubicin hydrochloride, epirubicin (hydrochloride), idarubicin hydrochloride, chromomycin A3, bleomycin (hydrochloride), peplomycin sulfate, therarubicin, zinostatin stimalamer, and gemtuzumab ozogamicin.

**[0091]** Examples of the plant formulations include vinblastine sulfate, vincristine sulfate, vindesine sulfate, irinotecan hydrochloride, etoposide, flutamide, vinorelbine tartrate, docetaxel hydrate, and paclitaxel.

**[0092]** Examples of the hormones include estramustine phosphate sodium, mepitiostane, epitio stanol, goserelin acetate, fosfestrol (diethylstilbestrol phosphate), tamoxifen citrate, toremifene citrate, fadrozole hydrochloride hydrate, medroxyprogesterone acetate, bicalutamide, leuporelin acetate, anastrozole, aminoglutethimide, androgen bicalutamide, and fulvestrant.

**[0093]** Examples of the platinum compounds include carboplatin, cisplatin, nedaplatin, and oxaliplatin.

**[0094]** Examples of the anti-CD20 antibodies include rituximab, ibritumomab, ibritumomab tiuxetan, and ocrelizumab.

**[0095]** Examples of the anti-CD52 antibodies include alemtuzumab.

**[0096]** Examples of the anti-PD-1 antibodies include nivolumab, and pembrolizumab.

**[0097]** Examples of the G-CSF formulation include pegfilgrastim, filgrastim, lenograstim, and nartograstim.

**[0098]** Examples of the differentiation-inducing agent for acute promyelocytic leukemia include tamibarotene, tretinoin, and arsenic trioxide formulations.

**[0099]** Examples of the kinase inhibitors include EGFR inhibitors including erlotinib hydrochloride, gefitinib, cetuximab, and panitumumab; HER2 inhibitors including lapatinib and trastuzumab; BCR-ABL inhibitors including imatinib, dasatinib, and nilotinib; multikinase inhibitors including sunitinib, vandetanib, crizotinib, and sorafenib.

**[0100]** Examples of the topoisomerase inhibitor include topotecan, teniposide, irinotecan, and sobuzoxane.

**[0101]** Examples of the aromatase inhibitor include exemestane.

**[0102]** Examples of the other anticancer agents include L-asparaginase, octreotide acetate, porfimer sodium, mitoxantrone acetate, aceglatone, ubenimex, eribulin mesilate, cladribine, krestin, bexarotene, denileukin difitox, temozolomide, nelarabine, fludarabine, bevacizumab, pemetrexed, pentostatin, bortezomib, lenalidomide, and calcium folinate.

**[0103]** Examples of the immunosuppressant include azathioprine, ascomycin, everolimus, salazosulapyridine, cyclosporine, cyclophosphamide, sirolimus, tacrolimus, bucillamine, methotrexate, and leflunomide.

**[0104]** Examples of the steroid include amcinonide, hydrocortisone sodium succinate, prednisolone sodium succinate, methylprednisolone sodium succinate, ciclesonide, difluprednate, betamethasone propionate, dexamethasone, deflazacort, triamcinolone, triamcinolone acetate, halcinonide, dexamethasone palmitate, hydrocortisone, flumetasone pivalate, prednisolone butylacetate, budesonide, prasterone sulfate, mometasone furoate, fluocinonide, fluocinolone acetate, fludoxycortide, flunisolide, prednisolone, alclometasone propionate, clobetasol propionate, dexamethasone propionate, deprodone propionate, fluticasone propionate, beclometasone propionate, betamethasone, methylprednisolone, methylprednisolone suleptanate, methylprednisolone sodium succinate, dexamethasone sodium phosphate,

hydrocortisone sodium phosphate, prednisolone sodium phosphate, diflucortolone valerate, dexamethasone valerate, betamethasone valerate, prednisolone valerate acetate, cortisone acetate, diflorasone acetate, dexamethasone acetate, triamcinolone acetate, paramethasone acetate, halopredone acetate, fludrocortisone acetate, prednisolone acetate, methylprednisolone acetate, clobetasone butyrate, hydrocortisone butyrate, hydrocortisone butyrate propionate, and betamethasone butyrate propionate.

**[0105]** Examples of the angiotensin II antagonistic drug include losartan, candesartan, valsartan, irbesartan, olmesartan, and telmisartan.

**[0106]** Examples of the angiotensin-converting enzyme inhibitor include alacepril, imidapril hydrochloride, quinapril hydrochloride, temocapril hydrochloride, delapril hydrochloride, benazepril hydrochloride, captopril, trandolapril, perindopril erbumine, enalapril maleate, and lisinopril.

**[0107]** Examples of the antiplatelet drugs include dipyridamole, and dilazep hydrochloride hydrate.

**[0108]** Examples of the anticoagulant drugs include warfarin and heparin.

**[0109]** Examples of the disease-modifying anti-rheumatic drugs include D-penicillamine, actarit, auranofin, salazosulfapyridine, hydroxychloroquine, bucillamine, methotrexate, leflunomide, lobenzarit sodium, aurothioglucose, and sodium aurothiomalate.

**[0110]** Examples of the prostaglandins (hereinafter, abbreviated as "PG") include PGE1 formulations (examples: alprostadil alfadex, alprostadil), PGI2 formulations (example: beraprost sodium), PG receptor agonists, and PG receptor antagonists. Examples of the PG receptor include PGE receptors (EP1, EP2, EP3, and EP4), PGD receptors (DP, and CRTH2), PGF receptors (FP), PGI2 receptors (IP), and TX receptors (TP).

**[0111]** Examples of the prostaglandin synthase inhibitor include salazosulfapyridine, mesalazine, olsalazine, 4-aminosalicylic acid, JTE-522, auranofin, carprofen, diphenpyramide, flunoxaprofen, flurbiprofen, indometacin, ketoprofen, lornoxicam, loxoprofen, meloxicam, oxaprozin, pearsamide, naproxen, piroxicam, piroxicam cinnamate, zaltoprofen, and pranoprofen.

**[0112]** Examples of the phosphodiesterase inhibitor include rolipram, cilomilast, Bay19-8004, NIK-616, roflumilast (BY-217), cipamfylline (BRL-61063), atizoram (CP-80633), ONO-6126, SCH-351591, YM-976, V-11294A, PD-168787, D-4396, and IC-485.

**[0113]** Examples of the anti-TNF- $\alpha$  formulation include anti-TNF- $\alpha$  antibodies, soluble TNF- $\alpha$  receptor, anti-TNF- $\alpha$  receptor antibodies, and soluble TNF- $\alpha$  binding protein, and particularly infliximab and etanercept.

**[0114]** Examples of the anti-IL-1 formulation include anti-IL-1 antibodies, soluble IL-1 receptor, anti-IL-1Ra antibodies and/or anti-IL-1 receptor antibodies and particularly anakinra.

**[0115]** Examples of the anti-IL-6 formulation include anti-IL-1 antibodies, soluble IL-6 receptor, and anti-IL-6 receptor antibodies, and particularly tocilizumab.

**[0116]** Examples of the cytokine inhibitor include suplastat tosylate, T-614, SR-31747, and sonatimod.

**[0117]** Examples of the HMG-CoA reductase inhibitor include atorvastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin.

**[0118]** Examples of the thiazolidine derivative include pioglitazone, ciglitazone, rosiglitazone, and troglitazone.

**[0119]** Furthermore, the combination drugs to be combined with a compound of the present invention includes not only ones discovered to date, but also ones that may be discovered in the future.

**[0120]** The compound of the present invention is usually administered systemically or locally, by oral or parenteral administration. Examples of oral agents include liquid medicines for internal use (for example, elixirs, syrups, pharmaceutically acceptable water-based agents, suspensions, and emulsions), and solid medicine for internal use (for example, tablets (including sublingual tablets and orally disintegrating tablets), pills, capsules (including hard capsules, soft capsules, gelatin capsules, and microcapsules), powders, granules, and lozenges). Examples of parenteral agents include liquid medicines (for example, injection agents (e.g., subcutaneous injection agents, intravenous injection agents, intramuscular injection agents, intraperitoneal injection agents, and drip agents), eye drops (for example, aqueous eye drops (e.g., aqueous eye drops, aqueous eye drop suspensions, viscous eye drops, and solubilized eye drops), and nonaqueous eye drops (for example, nonaqueous eye drops and nonaqueous eye drop suspensions)), agents for external use (for example, ointments (e.g., ophthalmic ointments)), and ear drops. These formulations may be controlled release agents such as rapid release formulations, and sustained release formulations. These formulations can be produced by well-known methods, for example, by the methods described in The Japanese Pharmacopoeia.

**[0121]** Liquid medicines for internal use as the oral agent can be produced by, for example, dissolving or suspending an active ingredient in a generally used diluent (for example, purified water, ethanol, or mixture liquid thereof). The liquid medicine may include a wetting agent, a suspension agent, a sweetening agent, a flavoring material, an aromatic substance, a preservative, or a buffer agent.

**[0122]** Solid medicines for internal use as the oral agent are formulated by, for example, mixing the active ingredient with, for example, a vehicle (for example, lactose, mannitol, glucose, microcrystalline cellulose, starch), a binder (for example, hydroxypropyl cellulose, polyvinylpyrrolidone, magnesium metasilicate aluminate), a disintegrant (for example, sodium carboxymethylcellulose), a lubricant (for example, magnesium stearate), a stabilizer, a dissolution adjuvant (for

example, glutamic acid, aspartic acid), and formulating according to standard methods. As necessary, coating may be carried out with a coating agent (for example, sugar, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose phthalate), and coating of two or more layers may be employed.

**[0123]** Agents for external use as parenteral agents are produced by well-known methods or generally used prescriptions. For example, an ointment may be produced by incorporation or melting of an active ingredient into base material. The ointment base material is selected from well-known material or generally used material. For example, a single material or a mixture of two or more of materials are selected from higher fatty acids and higher fatty acid esters (for example, adipic acid, myristic acid, palmitic acid, stearic acid, oleic acid, adipate esters, myristate esters, palmitate esters, stearate esters, oleate esters), waxes (for example, beeswax, spermaceti, ceresin), surfactants (for example, polyoxyethylene alkyl ether phosphate esters), higher alcohols (for example, cetanol, stearyl alcohol, etostearyl alcohol), silicone oils (for example, dimethylpolysiloxane), hydrocarbons (for example, hydrophilic petrolatum, white petrolatum, purified lanolin, liquid paraffin), glycols (for example, ethylene glycol, diethylene glycol, propylene glycol, polyethylene glycol, macrogol), plant oils (for example, castor oil, olive oil, sesame oil, turpentine oil), animal oils (for example, mink oil, egg yolk oil, squalane, squalene), water, absorption promoters, and anti-irritants. Furthermore, a humectant, preservative, stabilizer, antioxidant, or fragrance, may be included.

**[0124]** The injection agents as parenteral agents include solutions, suspensions, emulsions and solid injection agents to be dissolved or suspended in a solvent before use. The injection agent is used by, for example, dissolving, suspending or emulsifying an active ingredient in a solvent. Examples of the solvent include distilled water for injection, physiological saline, vegetable oils, alcohols such as propylene glycol, polyethylene glycol, ethanol, and mixtures thereof. Furthermore, the injection agent may contain a stabilizer, a dissolution aid (e.g., glutamic acid, aspartic acid, and Polysorbate 80 (registered trademark)), a suspending agent, an emulsifying agent, a soothing agent, a buffer, or a preservative. Such an injection agent is produced by sterilizing at the final step or employing an aseptic process. Furthermore, it is also possible to employ an aseptic solid product such as a freeze-dried product produced and sterilized or dissolved in aseptic distilled water for injection or other solvent before use.

**[0125]** When the compound of the present invention or combination agents of the compound of the present invention and other agents are used for the above-mentioned purposes, they are usually administered systemically or locally, usually by oral or parenteral administration. The doses to be administered are different depending upon ages, body weights, symptoms, therapeutic effects, administration method, and treatment time. The doses per adult person are generally from 1 ng to 1000 mg per dose, once or several times per day, by oral administration, from 0.1 ng to 100 mg per dose, once or several times per day, by parenteral administration, or continuous administration 1 to 24 hours per day intravenously. Needless to say, as mentioned above, the doses to be used vary dependent upon various conditions. Therefore, doses lower than the ranges specified above may be sufficient in some cases, and doses higher than the ranges specified above are needed in some cases.

[Examples]

**[0126]** Hereinafter, the present invention is described in detail with reference to Examples mentioned below, but the present invention is not limited thereto.

**[0127]** Solvents given in parentheses shown in chromatographic separation and TLC each indicate the eluting solvent or the developing solvent used, and the ratio is expressed in ratio by volume. The description "NH silica" denotes that CHROMATOREX NH TLC PLATE (catalog No.: 3800003) manufactured by FUJI SILYSIA CHEMICAL LTD. is used; and "DNH silica" denotes that CHROMATOREX NH TLC PLATE (catalog No.: 3800403) manufactured by FUJI SILYSIA CHEMICAL LTD. is used;

**[0128]** LC-MS/ELSD was carried out in the following conditions:

{Column: Waters ACQUITY C<sub>18</sub> (particle diameter:  $1.7 \times 10^{-6}$ ; column length:  $30 \times 2.1$  mm I.D.); flow rate: 1.0 mL/min; column temperature: 40°C; mobile phase (A): 0.1 % formic acid aqueous solution; mobile phase (B): 0.1 % formic acid-acetonitrile solution; gradient (rate of mobile phase (A) : mobile phase (B)): [0 min] 95:5; [0.1 min] 95:5; [1.2 min] 5:95; [1.4 min] 5:95; [1.41 min] 95:5; [1.5 min] 95:5; detector: UV (PDA), ELSD, MS}

**[0129]** The description in a parenthesis in the NMR data shows a solvent used for measurement.

**[0130]** Name of the compounds used in this specification are named by using ACD/Name (registered trademark) manufactured by Advanced Chemistry Development Inc., which is a computer program for naming compounds according to the regulation of IUPAC, or named according to the naming method of IUPAC.

Example 1

4-[(6-chloro-3-pyridinyl)oxy]-6,7-dimethoxy quinoline

**[0131]** Under the stream of nitrogen, a solution of 4-chloro-6,7-dimethoxy quinoline (1.00 g) (CAS registration No.:

35654-56-9) in chlorobenzene (9 mL), 6-chloropyridine-3-ol (0.65 g), and triethyl amine (11.3 mL) were placed in a 100-mL four-necked flask, and the mixture was stirred at a bath temperature (140°C) for five days. The resulting solution was left to cool to room temperature, water and ethyl acetate were added thereto, and the solution was separated. The water layer was extracted again with ethyl acetate, and the combined organic layer was washed with a saturated saline solution, and dried over anhydrous sodium sulfate. The solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1:8) to obtain the title compound (1.16 g) having the following physical property values.

TLC: Rf 0.22 (hexane : ethyl acetate = 1:3);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) : δ 8.52, 8.48, 7.87 - 7.85, 7.66, 7.49, 7.43, 6.65, 3.95, 3.93.

Example 2:

5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinamine

**[0132]** Under the stream of nitrogen, a solution of the compound (1.15 g) produced in Example 1 in tetrahydrofuran (THF) (18 mL), 1.0 mol/L lithium bis(trimethylsilyl)amide (LHDMS) (5.45 mL), tris(dibenzylideneacetone)dipalladium(0) chloroform complex (0.19 g), and 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (0.15 g) were placed in a 200-mL four-necked flask, and the mixture was stirred at a bath temperature (80°C) for 16.5 hours. Furthermore, 6 mol/L hydrochloric acid (10 mL) was added thereto, and the mixture was stirred at a bath temperature (80°C) for two hours. The mixture was left to cool to room temperature, then a sodium hydrogen bicarbonate aqueous solution and ethyl acetate were added, and the resulting solution was separated. The water layer was extracted again with ethyl acetate, and the combined organic layer was washed with a saturated saline solution, and then dried over anhydrous sodium sulfate. The solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (ethyl acetate → ethyl acetate : methanol = 9:1) to obtain the title compound (0.80 g) having the following physical property values.

TLC: Rf 0.51 (ethyl acetate : methanol = 4:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) : δ 8.45, 7.89, 7.51, 7.38 - 7.36, 6.56, 6.42, 6.05, 3.94.

Example 3:

ethyl 2,5-dioxo-5,6,7,8-tetrahydro-2H-chromene-3-carboxylate

**[0133]** 1,3-cyclohexanedione (CAS registration No.: 504-02-9) (13.25g) was dissolved in N,N-dimethyl formamide (DMF) (200 mL) at room temperature, and tert-butoxy potassium (13.26 g) and ethyl (E)-2-cyano-3-ethoxy-2-propenoate (CAS registration No.: 94-05-3) (20.00 g) were added thereto. The mixture was stirred for 21 hours. The reaction solution was diluted with ethyl acetate, 2 mol/L hydrochloric acid aqueous solution was added thereto, and the mixture was stirred. Ethyl acetate and water were further added, and the organic layer was extracted. The extract was washed with a saturated saline solution, then dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure to obtain the title compound (23.62 g) having the following physical property values.

TLC: Rf 0.35 (hexane : ethyl acetate = 1:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 1.37, 2.19, 2.61, 2.92, 4.36, 8.63.

Example 4:

2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinoline carboxylic acid

**[0134]** The compound (10.00 g) produced in Example 3 was dissolved in ethanol (200 mL) at room temperature, aniline (3.94 g) was added thereto, and the mixture was stirred for six hours. Solids precipitated from the reaction solution were collected by filtration through Kiriama funnel, and washed with ethanol. The obtained residue was dried under reduced pressure at 60°C. The title compound (4.01 g) having the following physical property values was obtained.


TLC: Rf 0.37 (dichloromethane : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : δ 2.11, 2.60, 7.25, 7.63, 9.21.

Example 5 (Reference):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0135]**



**[0136]** The compound (105 mg) produced in Example 4 and O-(7-aza-1-benzotriazolyl)-N,N,N',N'-tetramethyl uronium hexafluorophosphate (HATU) (192 mg) were dissolved in DMF (2 mL) at room temperature, diisopropylethylamine (DIPEA) (0.17 mL) and the compound (100 mg) produced in Example 2 were added thereto, and the mixture was stirred for 21 hours. The solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel chromatography (hexane : ethyl acetate = 30:70 → 0:100 → ethyl acetate : methanol = 70:30) to obtain the title compound (116 mg) having the following physical property values.

TLC: Rf 0.76 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.13, 2.60, 4.05, 6.44, 7.25, 7.42, 7.53, 7.63, 8.22, 8.48, 8.51, 9.32, 11.93.


Examples 5(1) to 5(54)

**[0137]** The following Example compounds were obtained by the procedure having the same purpose as in Example 5 using the compound produced in Example 2 and corresponding carboxylic acid derivatives in place of the compound produced in Example 4.

Example 5(1):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-7,7-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0138]**



**[0139]** TLC: Rf 0.75 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.06, 2.43, 2.48, 4.05, 6.45, 7.25, 7.43, 7.54, 7.55-7.65, 8.22, 8.48, 8.51, 9.32, 11.92.

Example 5(2):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2',5'-dioxo-1'-phenyl-2',5',6',8'-tetrahydro-1'H-spiro[cyclopropane-1,7'-quinoline]-3'-carboxamide

**[0140]** TLC: Rf 0.69 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.39, 0.54, 2.41, 2.48, 4.05, 6.45, 7.22, 7.43, 7.53, 7.55-7.62, 8.22, 8.49, 8.51, 9.36, 11.92.

(Byproduct) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-8'-hydroxy-2',5'-dioxo-1'-phenyl-2',5',6',8'-tetrahydro-1'H-spiro[cyclopropane-1,7'-quinoline]-3'-carboxamide

TLC: Rf 0.68 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.44, 0.61, 1.89, 3.39, 3.45, 4.10, 4.14, 6.76, 7.19, 7.47, 7.58-7.65, 7.86, 8.25, 8.63, 9.27, 12.05.

Example 5(3):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-ethyl-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0141]** (LC-MS/ELSD): (retention time: 0.70 min);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.41-1.46, 2.25-2.29, 2.62-2.65, 3.07-3.11, 4.06, 4.26-4.30, 6.45-6.47, 7.43, 7.55, 7.55-7.60, 8.29-8.31, 8.50-8.53, 9.21, 12.23.

Example 5(4):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(4-fluorobenzyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0142]** (LC-MS/ELSD): (retention time: 0.80 min);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.14-2.20, 2.58-2.63, 2.96-3.01, 4.06, 5.50, 6.45-6.47, 6.89-7.10, 7.26-7.37, 7.43, 7.55, 7.57-7.61, 8.29-8.30, 8.50-8.54, 9.29, 12.10.

Example 5(5):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-(tetrahydro-2H-pyran-4-yl)-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide


**[0143]** (LC-MS/ELSD): (retention time: 0.67 min);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.20-2.30, 2.60-2.64, 3.00-3.30, 3.45-3.55, 4.06, 4.15-4.20, 4.40-4.60, 6.45-6.47, 7.44, 7.55-7.61, 8.30-8.31, 8.49-8.52, 9.20, 12.17.

Example 5(6):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0144]**




**[0145]** TLC: Rf 0.54 (ethyl acetate : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.99, 2.09, 2.56, 3.19, 3.94, 4.25, 6.54, 7.40, 7.53, 7.86, 8.39, 8.48, 8.89, 12.19.

Example 5(7):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(4-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

[0146]




[0147] TLC: Rf 0.59 (dichloromethane : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.13, 2.60, 4.05, 6.44, 7.24 - 7.35, 7.43, 7.54, 7.57, 8.23, 8.50, 9.32, 11.88.

Example 5(8):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-6,6-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

[0148]



[0149] TLC: Rf 0.51 (dichloromethane : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.13, 1.80-1.90, 2.40-2.60, 3.92, 3.94, 6.53-6.55, 7.40, 7.50-7.53, 7.57-7.66, 7.84-7.88, 8.34-8.36, 8.40-8.43, 8.47-8.49, 8.99, 11.98.

Example 5(9):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-isobutyl-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

TLC: Rf 0.50 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 0.94, 2.05-2.25, 2.49-2.65, 3.15-3.24, 3.93, 4.13, 6.54, 7.04, 7.53, 7.86, 8.38-8.45, 8.48, 8.89, 12.24.



Example 5(10):


N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(2R)-1-hydroxy-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide TLC: Rf 0.20 (ethyl acetate, NH silica);

**[0151]**  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.70, 1.11, 2.13, 2.41, 2.87-2.99, 3.35, 4.02, 4.07, 4.19, 4.88, 6.17, 7.19, 7.53, 7.86, 8.17, 8.37, 8.58, 8.61, 12.18.

Example 5(11):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(2S)-1-hydroxy-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0152]**




**[0153]** TLC: Rf 0.20 (ethyl acetate, NH silica);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.70, 1.11, 2.13, 2.41, 2.87-2.99, 3.35, 4.02, 4.07, 4.19, 4.88, 6.17, 7.19, 7.53, 7.86, 8.17, 8.37, 8.58, 8.61, 12.18.

Example 5(12):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0154]**



**[0155]** TLC: Rf 0.56 (ethyl acetate : methanol = 9:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  2.13, 2.60, 4.05, 6.44, 7.07, 7.32, 7.43, 7.54, 7.59, 8.23, 8.49, 9.32, 11.85.

Example 5(13):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[1-(hydroxymethyl)cyclobutyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0156]** TLC: Rf 0.15 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.81-1.95, 2.16-2.29, 2.40-2.75, 2.83-3.01, 3.45, 4.05, 4.06, 4.18, 4.48, 6.41, 7.39, 7.55, 7.61, 8.28, 8.46, 8.54, 9.08, 11.99.

Example 5(14):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(1R)-1-(4-fluorophenyl)ethyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0157]** TLC: Rf 0.51 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.00-2.20, 2.45-3.07, 4.05, 4.06, 6.45, 7.03-7.12, 7.18-7.23, 7.43, 7.55, 7.58, 8.28, 8.50, 8.51, 9.23, 12.07.

Example 5(15):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(1S)-1-(4-fluorophenyl)ethyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide


**[0158]** TLC: Rf 0.50 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.00-2.20, 2.45-3.07, 4.05, 4.06, 6.45, 7.03-7.12, 7.18-7.23, 7.43, 7.55, 7.58, 8.28, 8.50, 8.51, 9.23, 12.07.

Example 5(16):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0159]**



**[0160]** TLC: Rf 0.52 (ethyl acetate : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.15, 2.63, 4.05, 6.44, 7.26 - 7.61, 8.22, 8.49, 9.33, 11.85.

Example 5(17):

**[0161]** N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2',5'-dioxo-1'-phenyl-2',5',6',8'-tetrahydro-1'H-spiro[cyclobutane-1,7'-quinoline]-3'-carboxamide

**[0162]** TLC: Rf 0.71 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.78-1.93, 2.64, 2.73, 2.93, 3.02, 4.05, 6.45, 7.24, 7.42, 7.53-7.67, 8.22, 8.49, 9.29, 11.92.

Example 5(18):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-[(1S)-1-phenylethyl]-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0163]** TLC: Rf 0.69 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.01, 2.55, 2.95, 4.05, 4.06, 6.44, 7.19, 7.29-7.43, 7.55, 7.57, 8.28, 8.49-8.53, 9.25, 12.13.

Example 5(19):

1-cyclopropyl-N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0164]** (LC-MS/ELSD): (retention time: 0.69 min);

MASS (ESI, Pos.): 527(M+H)<sup>+</sup>.

Example 5(20):

1-(1-cyclopropyl-2-hydroxyethyl)-N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0165]** (LC-MS/ELSD): (retention time: 0.71 min);

MASS (ESI, Pos.): 571(M+H)<sup>+</sup>.

Example 5(21):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[2-(methylsulfonyl)-1-phenylethyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0166]** TLC: Rf 0.67 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.20-2.80, 3.14, 3.92, 3.93, 4.61, 6.29, 6.49, 7.33, 7.39, 7.50, 7.83, 8.34, 8.45, 8.92, 11.80.

Example 5(22):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-(3-pentanyl)-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0167]** (LC-MS/ELSD): (retention time: 0.80 min);

MASS (ESI, Pos.): 557(M+H)<sup>+</sup>.

Example 5(23):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2-hydroxy-2-methylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0168]** (LC-MS/ELSD): (retention time: 0.66 min);

MASS (ESI, Pos.): 559(M+H)<sup>+</sup>.

Example 5(24):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(1-hydroxy-3-methyl-2-butanyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0169]** (LC-MS/ELSD): (retention time: 0.71 min);

MASS (ESI, Pos.): 573(M+H)<sup>+</sup>.

Example 5(25):

1-cyclobutyl-N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0170]** (LC-MS/ELSD): (retention time: 0.74 min);  
 MASS (ESI, Pos.): 541(M+H)<sup>+</sup>.

Example 5(26):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[1-(4-fluorophenyl)ethyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0171]** (LC-MS/ELSD): (retention time: 0.80 min);  
 MASS (ESI, Pos.): 609 (M+H)<sup>+</sup>.

Example 5(27):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-(3-pyridinyl)-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0172]** (LC-MS/ELSD): (retention time: 0.63 min);  
 MASS (ESI, Pos.): 564(M+H)<sup>+</sup>.

Example 5(28):

N-[5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl]-1-[2-(dimethylamino)ethyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0173]** (LC-MS/ELSD): (retention time: 0.51 min);  
 MASS (ESI, Pos.): 558(M+H)<sup>+</sup>.

Example 5(29):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2-methoxy-2-methylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0174]** TLC: R<sub>f</sub> 0.65 (dichloromethane : methanol = 9:1);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.19-1.35, 2.18, 2.62, 2.81, 3.16, 3.83, 4.06, 4.07, 4.99, 6.45, 7.27, 7.44, 7.56, 7.59, 8.31, 8.51, 8.54, 9.23, 12.24.

Example 5(30):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(3-hydroxy-3-methyl-2-butanyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0175]** (LC-MS/ELSD): (retention time: 0.70 min);  
 MASS (ESI, Pos.): 573 (M+H)<sup>+</sup>.

Example 5(31):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(3-oxetanyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0176]** (LC-MS/ELSD): (retention time: 0.68 min);  
 MASS (ESI, Pos.): 543 (M+H)<sup>+</sup>.

Example 5(32):

1-(4,4-difluorocyclohexyl)-N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0177]** (LC-MS/ELSD): (retention time: 0.80 min);  
 MASS (ESI, Pos.): 605(M+H)<sup>+</sup>.

Example 5(33):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(1-methyl-1H-pyrazole-4-yl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0178]** (LC-MS/ELSD): (retention time: 0.63 min);  
 MASS (ESI, Pos.): 567(M+H)<sup>+</sup>.

Example 5(34):

1-(cyclopropylmethyl)-N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0179]** (LC-MS/ELSD): (retention time: 0.75 min);  
 MASS (ESI, Pos.): 541 (M+H)<sup>+</sup>.

Example 5(35):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(3-methyl-2-butanyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0180]** (LC-MS/ELSD): (retention time: 0.82 min);  
 MASS (ESI, Pos.): 557 (M+H)<sup>+</sup>.

Example 5(36):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-hexyl-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0181]** (LC-MS/ELSD): (retention time: 0.88 min);  
 MASS (ESI, Pos.): 571 (M+H)<sup>+</sup>.

Example 5(37):

1-[(1S)-1-cyclohexylethyl]-N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0182]** (LC-MS/ELSD): (retention time: 0.92 min);  
 MASS (ESI, Pos.): 597 (M+H)<sup>+</sup>.

Example 5(38):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-(3-phenylpropyl)-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0183]** (LC-MS/ELSD): (retention time: 0.85 min);  
 MASS (ESI, Pos.): 605 (M+H)<sup>+</sup>.

Example 5(39):

1-[(1S)-1-cyclopropylethyl]-N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0184]** (LC-MS/ELSD): (retention time: 0.79 min);  
 MASS (ESI, Pos.): 555 (M+H)<sup>+</sup>.

Example 5(40):

1-[(1R)-1-cyclopropylethyl]-N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0185]** (LC-MS/ELSD): (retention time: 0.80 min);  
 MASS (ESI, Pos.): 555 (M+H)<sup>+</sup>.

Example 5(41):

N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-1-(4-methylphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0186]** TLC: R<sub>f</sub> 0.66 (ethyl acetate : methanol = 9:1);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.10, 2.48, 2.60, 4.05, 6.44, 7.13, 7.42, 7.53, 7.56, 8.21, 8.49, 8.50, 9.31, 11.94.

Example 5(42):

1-(4-chlorophenyl)-N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0187]** TLC: R<sub>f</sub> 0.60 (ethyl acetate : methanol = 9:1);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.13, 2.60, 4.05, 6.45, 7.22, 7.42, 7.53, 7.57, 7.61, 8.23, 8.48, 8.50, 9.31, 11.85.

Example 5(43):

1-(2,4-difluorophenyl)-N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0188]** TLC: R<sub>f</sub> 0.33 (ethyl acetate);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.17, 2.62, 4.06, 6.44, 7.14, 7.31, 7.43, 7.54, 7.57, 8.24, 8.50, 9.33, 11.79.

Example 5(44):

1-(2-chlorophenyl)-N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0189]** TLC: R<sub>f</sub> 0.29 (ethyl acetate);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.14, 2.54, 2.64, 4.05, 6.44, 7.34, 7.43, 7.55, 7.57, 7.69, 8.23, 8.49, 8.51, 9.35, 11.85.

Example 5(45):

N-{5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-1-(2-methylphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0190]** TLC: R<sub>f</sub> 0.36 (ethyl acetate);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.08-2.15, 2.36, 2.58-2.65, 4.05, 6.44, 7.15, 7.43-7.59, 8.23, 8.48-8.52, 9.35, 11.97.

Example 5(46):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(1R)-2-hydroxy-1-phenylethyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0191]** TLC: Rf 0.15 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.14, 2.48, 3.04, 3.30, 4.04, 4.05, 4.51, 5.11, 6.22, 7.19 - 7.42, 7.53, 7.73, 8.16, 8.40, 8.51, 8.86, 11.89.

Example 5(47):

N-[5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl]-1-[(1S)-2-hydroxy-1-phenylethyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0192]** TLC: Rf 0.51 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.08-2.25, 2.30-2.70, 2.97-3.16, 3.16-3.40, 4.05, 4.06, 4.45-4.60, 4.88-5.19, 5.67-6.15, 6.26, 7.16-7.42, 7.53, 7.72, 8.18, 8.43, 8.52, 8.94, 11.90.

Example 5(48):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-[(1R)-1-phenylethyl]-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0193]** TLC: Rf 0.51 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.99-2.10, 2.40-2.70, 2.91, 4.06, 6.45, 7.21, 7.30-7.43, 7.55, 7.58, 8.29, 8.52, 9.26, 12.14.

Example 5(49):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(2S)-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0194]** TLC: Rf 0.51 (ethyl acetate : methanol = 10:1)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.74, 1.07, 1.70, 2.10-2.40, 2.48-2.73, 2.95-3.15, 3.96, 4.06, 4.07, 6.45, 7.43, 7.52-7.64, 8.31, 8.47-8.56, 9.21, 12.27.

Example 5(50):

1-(3-chlorophenyl)-N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0195]** TLC: Rf 0.72 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.98-2.07, 2.48-2.60, 3.93, 6.54, 7.40, 7.45-7.51, 7.67-7.69, 7.90, 8.36, 8.41, 8.48, 8.97, 11.89.

Example 5(51):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(3-methylphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0196]** TLC: Rf 0.38 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.89-2.04, 2.36-2.62, 3.92-3.94, 6.54, 7.10-7.56, 7.86, 8.35, 8.42, 8.47-8.49, 8.97, 11.97.

Example 5(52):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(2R)-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0197]** TLC: Rf 0.48 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.74, 1.07, 1.70, 2.10-2.38, 2.49-2.75, 2.93-3.15, 3.88-4.02, 4.06, 4.07, 6.45, 7.43, 7.52-7.63, 8.31, 8.46-8.58, 9.21, 12.27.

Example 5(53):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[1-(2-hydroxy-2-methylpropyl)-1H-pyrazole-4-yl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0198]** TLC: Rf 0.15 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.27, 2.14, 2.59, 2.73, 3.20, 4.05, 4.18, 6.44, 7.43, 7.53, 7.54, 7.70, 8.24, 8.49, 8.51, 9.28, 11.88.

Example 5(54):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-2,5,6,7,8,9-hexahydro-1H-cyclohepta[b]piperidine-3-carboxamide


**[0199]** TLC: Rf 0.32 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.80, 1.91, 2.71, 2.77, 4.05, 6.45, 7.25, 7.43, 7.53-7.66, 8.21, 8.48, 8.50, 9.11, 11.99.

Example 6:

1-(4-fluorophenyl)-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0200]**



**[0201]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 1 → Example → Example 3 → Example 4 → Example 5, using 4-chloro-7-methoxy quinoline (CAS registration No.: 68500-37-8) in place of 4-chloro-6,7-dimethoxy quinoline.

TLC: Rf 0.73 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.92-2.09, 2.40-2.70, 3.93, 6.54, 7.29, 7.41-7.60, 7.87, 8.21, 8.36, 8.41, 8.61, 8.97, 11.94.

Example 6(1) to 6(38)

**[0202]** The following Example compounds were obtained by the procedure having the same purpose as in Example 6, using 4-chloro-7-methoxy quinoline or a corresponding quinoline derivative in place of it and the compound produced in Example 4 or a corresponding carboxylic acid derivative in place of it.

Example 6(1):

N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0203]** TLC: Rf 0.30 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.98, 2.45-2.57, 3.92, 6.52, 7.28, 7.40, 7.44, 7.46, 7.57-7.67, 7.85, 8.20, 8.35, 8.40, 8.60, 8.97, 11.95.



Example 6(2):

N-{5-[(6-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

- 5 **[0204]** (LC-MS/ELSD): (retention time: 0.77 min);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.05-2.20, 2.51-2.68, 3.97, 6.52, 7.25-7.26, 7.27-7.29, 7.41, 7.54-7.67, 8.00, 8.22, 8.50, 8.55, 9.33, 11.93.

Example 6(3):

- 10 1-[(2S)-1-hydroxy-3-methyl-2-butanyl]-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide
- [0205]** TLC: R<sub>f</sub> 0.71 (ethyl acetate : methanol = 5:1);  
 15 <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.72, 1.12, 2.18, 2.48, 2.92-3.02, 3.31, 3.95, 4.04, 4.17, 4.77, 5.23, 6.23, 7.21, 7.78, 8.21, 8.24, 8.51, 8.55, 8.77, 12.16.

Example 6(4):

- 20 1-[(2R)-1-hydroxy-3-methyl-2-butanyl]-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide
- [0206]** TLC: R<sub>f</sub> 0.71 (ethyl acetate : methanol = 5:1);  
 25 <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.72, 1.12, 2.18, 2.48, 2.92-3.02, 3.31, 3.95, 4.04, 4.17, 4.77, 5.23, 6.23, 7.21, 7.78, 8.21, 8.24, 8.51, 8.55, 8.77, 12.16.

Example 6(5):

- 30 1-(3-fluorophenyl)-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide
- [0207]** TLC: R<sub>f</sub> 0.43 (ethyl acetate : methanol = 19:1);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.93-2.08, 2.40-2.65, 3.93, 6.54, 7.27-7.55, 7.65-7.76, 7.87, 8.21, 8.34-8.47, 8.62, 8.98, 11.91.

35 Example 6(6):

2,5-dioxo-1-phenyl-N-(5-[(7-(trifluoromethyl)-4-quinolinyl)oxy]-2-pyridinyl)-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

- 40 **[0208]** (LC-MS/ELSD): (retention time: 1.06 min);  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 2.00, 2.52-2.55, 6.88, 7.46-7.48, 7.58-7.66, 7.93-7.96, 8.40-8.44, 8.46, 8.57, 8.85, 8.99, 11.99.

Example 6(7):

- 45 1-cyclobutyl-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide
- [0209]** TLC: R<sub>f</sub> 0.69 (ethyl acetate : methanol = 19:1);  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.70-1.90, 1.95-2.10, 2.35-2.60, 2.73-2.89, 3.04-3.15, 3.94, 4.89-5.03, 6.55, 7.30, 7.42, 7.88, 8.22, 8.40, 8.42, 8.62, 8.83, 12.16.

- 50 Example 6(8):
- 1-(2,2-dimethylpropyl)-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide
- 55 **[0210]** TLC: R<sub>f</sub> 0.65 (ethyl acetate : methanol = 19:1);  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 0.99, 2.00-2.16, 2.48-2.65, 3.15-3.22, 3.94, 4.12-4.38, 6.55, 7.30, 7.42, 7.88, 8.22, 8.39, 8.42, 8.62, 8.90, 12.20.

Example 6(9):

N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-6,6-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0211]** TLC: Rf 0.55 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.13, 1.80-1.90, 2.34-2.56, 3.92, 6.54, 7.29, 7.41, 7.48-7.68, 7.87, 8.20, 8.35, 8.41, 8.61, 8.99, 11.98.

Example 6(10):

1-(4-chlorophenyl)-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0212]** TLC: Rf 0.52 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.91-2.04, 2.37-2.60, 3.93, 6.54, 7.29, 7.41, 7.52, 7.72, 7.88, 8.20, 8.34, 8.41, 8.61, 8.97, 11.91.

Example 6(11):

N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(4-methylphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0213]** TLC: Rf 0.41 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.91-2.03, 2.42, 2.40-2.60, 3.93, 6.54, 7.26-7.46, 7.87, 8.20, 8.36, 8.41, 8.61, 8.96, 11.97.

Example 6(12):

1-(4,4-difluorocyclohexyl)-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0214]** TLC: Rf 0.50 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.81, 2.25, 2.31, 2.62, 3.08, 3.30, 3.98, 4.10, 6.43, 7.25, 7.42, 7.57, 8.24, 8.30, 8.51, 8.61, 9.19, 12.08.

Example 6(13):

1-(cyclopropylmethyl)-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0215]** TLC: Rf 0.55 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.62, 1.15, 2.26, 2.64, 3.15, 3.98, 4.20, 6.43, 7.23, 7.42, 7.57, 8.23, 8.29, 8.51, 8.61, 9.22, 12.23.

Example 6(14):

1-[(1R)-1-cyclopropylethyl]-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0216]** TLC: Rf 0.55 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.06, 0.30, 0.56, 0.77, 1.80, 2.24, 2.60, 2.89, 3.43, 3.98, 6.44, 7.22, 7.43, 7.56, 8.23, 8.30, 8.51, 8.61, 9.20, 12.29.

Example 6(15):

1-[(1S)-1-cyclopropylethyl]-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0217]** TLC: Rf 0.55 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.06, 0.30, 0.56, 0.77, 1.80, 2.24, 2.60, 2.89, 3.43, 3.98, 6.44, 7.22, 7.43, 7.56, 8.23, 8.30, 8.51, 8.61, 9.20, 12.29.

Example 6(16):

1-(4-fluorophenyl)-N-{5-[(6-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0218]** TLC: Rf 0.69 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.13, 2.60, 3.97, 6.52, 7.29-7.35, 7.41, 7.55-7.60, 8.00, 8.23, 8.49, 8.54, 9.32, 11.88.

Example 6(17):

1-isobutyl-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0219]** TLC: Rf 0.34 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 0.94, 2.08-2.20, 2.40-2.60, 3.12-3.26, 3.94, 4.12, 6.55, 7.30, 7.42, 7.88, 8.22, 8.39, 8.41, 8.62, 8.90, 12.24.

Example 6(18):

N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2',5'-dioxo-1'-phenyl-2',5',6',8'-tetrahydro-1'H-spiro[cyclobutane-1,7'-quinoline]-3'-carboxamide

**[0220]** TLC: Rf 0.56 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.65-1.91, 2.63-2.74, 3.93, 7.29, 7.45, 7.46-7.53, 7.58-7.71, 7.87, 8.20, 8.36, 8.41, 8.61, 8.94, 11.94.

Example 6(19):

1-(2,4-difluorophenyl)-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0221]** TLC: Rf 0.51 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.16, 2.62, 3.97, 6.42, 7.10-7.33, 7.42, 7.56, 8.21-8.24, 8.49, 8.60, 9.33, 11.79.

Example 6(20):

1-(2-chlorophenyl)-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0222]** TLC: Rf 0.40 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.14, 2.55, 2.63, 3.97, 6.41, 7.23, 7.35, 7.42, 7.53-7.58, 7.68, 8.21, 8.23, 8.49, 8.60, 9.34, 11.84.

Example 6(21):

N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2-methylphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0223]** TLC: Rf 0.44 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.15-2.20, 2.35, 2.58-2.67, 3.97, 6.41, 7.15, 7.24, 7.41-7.49, 7.56, 8.21, 8.23, 8.49, 8.60, 9.34, 11.96.

Example 6(22):

1-[1-(hydroxymethyl)cyclobutyl]-N-{5-[(6-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0224]** TLC: Rf 0.15 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.87, 2.18, 2.40, 2.62, 2.94, 3.44, 3.98, 4.10, 4.18, 4.50, 6.46, 7.38, 7.58, 7.62, 7.92, 8.28, 8.49, 8.57, 9.02, 12.02.

Example 6(23):

1-[1-(hydroxymethyl)cyclobutyl]-N-{5-[(7-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0225]** TLC: Rf 0.15 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.95, 2.21, 2.38-2.69, 2.95, 3.42, 3.68, 3.97, 4.15, 4.46, 6.40, 7.23, 7.39, 7.61, 8.24, 8.28, 8.53, 8.58, 9.10, 11.99.

Example 6(24):

N-{5-[(6-methoxy-4-quinolinyloxy)-2-pyridinyl]-1-(2-methylphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0226]** TLC: Rf 0.25 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.12, 2.26, 2.60, 3.97, 6.51, 7.15, 7.40-7.49, 7.58, 8.00, 8.23, 8.49, 8.55, 9.34, 11.98.

Example 6(25):

1-(2-chlorophenyl)-N-{5-[(6-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0227]** TLC: Rf 0.25 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.15, 2.57, 2.63, 3.97, 6.51, 7.35-7.61, 7.68, 8.00, 8.23, 8.50, 8.55, 9.34, 11.86.

Example 6(26):

1-(2,4-difluorophenyl)-N-{5-[(6-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0228]** TLC: Rf 0.25 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.18, 2.62, 3.97, 6.51, 7.18, 7.21-7.40, 7.41, 7.57, 8.01, 8.24, 8.50, 8.55, 9.32, 11.79.

Example 6(27):

1-[(1S)-1-cyclopropylethyl]-N-{5-[(6-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0229]** TLC: Rf 0.60 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.03-0.13, 0.20-0.40, 0.49-0.67, 0.69-0.89, 1.81, 2.09-2.42, 2.47-2.72, 2.89, 3.31-3.52, 3.98, 6.53, 7.42, 7.55-7.64, 8.00, 8.31, 8.53, 8.56, 9.21, 12.30.

Example 6(28):

1-[(1S)-2-hydroxy-1-phenylethyl]-N-{5-[(7-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0230]** TLC: Rf 0.54 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.07-2.29, 2.36-2.68, 2.97-3.16, 3.17-3.43, 3.96, 4.45-4.62, 4.95-5.23, 5.68-6.05, 6.22, 7.12-7.43, 7.72, 8.17, 8.23, 8.47-8.56, 8.86, 11.90.

Example 6(29):

1-[(1R)-1-(4-fluorophenyl)ethyl]-N-{5-[(7-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0231]** TLC: Rf 0.58 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.00, 2.05, 2.40-2.80, 2.96, 3.98, 6.43, 7.08, 7.18-7.29, 7.42, 7.58, 8.24, 8.28, 8.51, 8.61, 9.25, 12.07.

Example 6(30):

N-{5-[(7-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-[(1R)-1-phenylethyl]-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0232]** TLC: Rf 0.58 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.99-2.10, 2.40-2.70, 2.90, 3.98, 6.43, 7.22, 7.27-7.43, 7.59, 8.24, 8.28, 8.52, 8.61, 9.26, 12.13.

Example 6(31):

1-(2-fluorophenyl)-N-{5-[(6-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0233]** TLC: Rf 0.60 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.06-2.25, 2.63, 3.97, 6.52, 7.42, 7.51-7.67, 7.99, 8.23, 8.50, 8.55, 9.33, 11.85.

Example 6(32):

1-(3-fluorophenyl)-N-{5-[(6-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0234]** TLC: Rf 0.66 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.04-2.26, 2.53-2.70, 3.97, 6.52, 6.99-7.14, 7.27-7.36, 7.37-7.44, 7.52-7.68, 7.99, 8.23, 8.49, 8.55, 9.32, 11.85.

Example 6(33):

N-{5-[(7-methoxy-4-quinolinyloxy)-2-pyridinyl]-1-[(2S)-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0235]** TLC: Rf 0.57 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.74, 1.07, 1.70, 2.10-2.34, 2.45-2.76, 2.94-3.14, 3.86-3.96, 3.98, 6.42, 7.21-7.25, 7.43, 7.58, 8.24, 8.30, 8.52, 8.61, 9.21, 12.27.

Example 6(34):

1-(2-fluorophenyl)-N-{5-[(7-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0236]** TLC: Rf 0.35 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.96-2.18, 2.30-2.60, 3.93, 6.54, 7.29, 7.41, 7.44-7.72, 7.88, 8.21, 8.36, 8.41, 8.61, 9.00, 11.81.

Example 6(35):

N-{5-[(7-methoxy-4-quinolinyloxy)-2-pyridinyl]-1-(3-methylphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0237]** TLC: Rf 0.50 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.94-2.05, 2.40, 2.45-2.60, 3.93, 6.54, 7.22-7.55, 7.87, 8.21, 8.36, 8.42, 8.62, 8.97, 11.97.

Example 6(36):

1-(3-chlorophenyl)-N-{5-[(7-methoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0238]** TLC: Rf 0.74 (ethyl acetate : methanol = 19:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.98-2.07, 2.45-2.60, 3.93, 6.54, 7.29, 7.41, 7.46-7.51, 7.66-7.69, 7.87, 8.20, 8.36, 8.41, 8.61, 8.97, 11.89.

Example 6(37):

N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(2R)-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0239]** TLC: Rf 0.55 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.74, 1.07, 1.70, 2.12-2.38, 2.47-2.74, 2.95-3.20, 3.90-3.95, 3.98, 6.43, 7.20-7.25, 7.43, 7.58, 8.24, 8.30, 8.52, 8.61, 9.21, 12.26.

Example 6(38):

1-[1-(2-hydroxy-2-methylpropyl)-1H-pyrazole-4-yl]-N-{5-[(7-methoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0240]** TLC: Rf 0.15 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.28, 2.14, 2.62, 2.75, 3.10, 3.97, 4.18, 6.41, 7.22, 7.42, 7.52-7.60, 7.70, 8.21, 8.23, 8.48, 8.60, 9.27, 11.87.

Example 7:

tert-butyl N-[5-[(6,7-dimethoxy-4-quinazolinyl)amino]-2-pyridyl]carbamate

**[0241]** 4-chloro-6,7-dimethoxyquinazoline (CAS registration No.: 13790-39-1) (450 mg) and tert-butyl (5-aminopyridin-2-yl)carbamate (420 mg) were dissolved in N,N-dimethyl acetamide (DMA) (20 mL) at room temperature. 4 mol/L hydrochloric acid - dioxane (0.5 mL) was added thereto at 50°C, and the mixture was stirred. Thereafter, the resulting solution was heated to 80°C and stirred for three hours. The resulting solution was left to cool to room temperature, methyl tert-butyl ether (MTBE) was added to the reaction solution, solids precipitated from the reaction solution were collected by filtration through Kiriya funnel, and washed with MTBE. The resulting residue was dried under reduced pressure at 60°C. The title compound (821 mg) having the following physical property values was obtained.

TLC: Rf 0.45 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.49, 4.00, 7.34, 7.88, 8.05, 8.32, 8.56, 8.84, 10.02, 11.54.

Example 8:

N<sup>5</sup>-(6,7-dimethoxy-4-quinazolinyl)pyridine-2,5-diamine

**[0242]** The compound (800 mg) produced in Example 7 was dissolved in dichloromethane (10 mL) at room temperature, trifluoroacetic acid (0.3 mL) was added thereto, and the mixture was stirred at room temperature for six hours. The reaction solution was diluted with ethyl acetate, and saturated sodium bicarbonate water was added to the reaction solution and stirred. Ethyl acetate and water were further added, and the organic layer was extracted. THF and water were added to the water layer, and the organic layer was extracted. The organic layer was collected, washed with a saturated saline solution, dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. Ethyl acetate and hexane were added to the resulting residue, followed by stirring at room temperature. The solid was washed with ether in a slurry form. The residue collected by filtration through Kiriya funnel was dried under reduced pressure at 60°C to obtain the title compound (598 mg) having the following physical property values.


TLC: Rf 0.16 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 3.91, 5.83, 6.49, 7.12, 7.65, 7.82, 8.09, 8.30, 9.36.

Example 9:

N-{5-[(6,7-dimethoxy-4-quinazolinyl)amino]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0243]**



**[0244]** The compound produced in Example 4 (76 mg) and HATU (154 mg) were dissolved in DMF (1 mL) at room temperature. To the resulting solution, DIPEA (0.14 mL) and the compound produced in Example 8 (80 mg) were added, and the mixture was stirred for 18 hours. The reaction solution was diluted with ethyl acetate, and saturated sodium bicarbonate water was added and stirred. Ethyl acetate and water were further added, and the organic layer was extracted. The organic layer was washed with a saturated saline solution, dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel chromatography (hexane : ethyl acetate = 30:70 → 0:100 → ethyl acetate : methanol = 70:30) to obtain the title compound (57 mg) having the following physical property values.

TLC: Rf 0.59 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.10, 2.58, 4.04, 7.04, 7.13, 7.25, 7.59, 8.20, 8.41, 8.50, 8.64, 9.30, 11.82.

Examples 9(1) to 9(4)

**[0245]** The following Example compounds were obtained by the procedure having the same purpose as in Example 7 → Example 8 → Example 9, using 4-chloro-6,7-dimethoxyquinazoline or a corresponding quinoline derivative in place of it, tert-butyl (5-aminopyridin-2-yl)carbamate or a corresponding amine derivative in place of it, and the compound produced in Example 4.

Example 9(1):

N-{5-[(6,7-dimethoxy-4-quinolinyl)amino]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide trifluoroacetate

**[0246]** TLC: Rf 0.21 (dichloromethane : methanol = 10:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.86-2.14, 2.49-2.62, 3.98, 4.02, 6.75, 7.35-7.40, 7.41-7.52, 7.53-7.73, 7.94-8.07, 8.36-8.56, 8.99, 10.42, 12.03, 14.00.

Example 9(2):

N-{4-[(6,7-dimethoxy-4-quinolinyl)amino]phenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0247]** (LC-MS/ELSD): (retention time: 0.75 min);

<sup>1</sup>H-NMR (CD<sub>3</sub> OD): δ 2.04-2.13, 2.56-2.64, 4.01, 4.02, 6.82, 7.24, 7.35-7.42, 7.59-7.67, 7.69, 7.75-7.79, 8.19, 9.17.

Example 9(3):

N-{5-[(7-methoxy-4-quinolinyl)amino]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0248]** TLC: Rf 0.63 (ethyl acetate : methanol : ammonia water = 9:1:0.5);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.90-2.05, 2.40-2.60, 3.89, 6.70, 7.18, 7.25, 7.42-7.50, 7.54-7.68, 7.86, 8.22-8.40, 8.97, 8.99, 11.85.

Example 9(4):

N-{4-[(6,7-dimethoxy-4-quinazolinyl)amino]phenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

- 5 **[0249]** (LC-MS/ELSD): (retention time: 0.79 min);  
<sup>1</sup>H-NMR (CD<sub>3</sub> OD): δ 1.94-2.07, 2.50-2.58, 3.94, 3.96, 7.18, 7.43-7.51, 7.57-7.68, 7.75, 7.87, 8.50, 8.95, 9.69, 11.48.

Example 10:

- 10 5-[(E)-{3-(benzyloxy)phenyl}imino]methyl]-2,2-dimethyl-1,3-dioxane-4,6-dione

- [0250]** 3-benzyloxyaniline (25 g), meldrum's acid (22 g), ethyl orthoformate (22 g), and ethanol (25 mL) were placed in a 200-mL eggplant flask. The mixture was heated and refluxed for 80 min, and left to cool to room temperature. Then, the precipitated powder was collected by filtration. The powder was washed with ethanol (50 mL) and dried to obtain the title compound (43 g) having the following physical property values.

TLC: Rf 0.48 (hexane : ethyl acetate = 2:1);  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.66, 5.15, 6.88, 7.11, 7.30 - 7.45, 8.60, 11.2.

Example 11:

- 20 7-(benzyloxy)-4(1H)-quinolinone

- [0251]** The compound produced in Example 10 (42 g) and 1,2-dichlorobenzene (420 mL) were placed in a 1 L-eggplant flask. The mixture was heated and refluxed for 5.5 hours, and left to cool to room temperature. Then, the precipitated powder was collected by filtration. The powder was washed with methanol (84 mL) and dried to obtain the title compound (18 g) having the following physical property values.

TLC: Rf 0.60 (ethyl acetate :methanol = 9:1);  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 5.19, 5.92, 6.98, 7.32 - 7.43, 7.77, 7.97.

- 30 Example 12:

7-(benzyloxy)-4-chloroquinolinone

- [0252]** Under argon atmosphere, the compound produced in Example 11 (17 g), toluene (34 mL) and phosphorus oxychloride (10 g) were placed in a 300-mL eggplant flask. The mixture was heated and refluxed for 2.5 hours, left to cool to 70°C, and diluted with ethyl acetate (135 mL). Thereafter, the mixture was left to cool to room temperature, and was neutralized by 2 mol/L sodium hydroxide aqueous solution. The resulting solution was extracted with ethyl acetate, washed with a saturated saline solution and dried over anhydrous sodium sulfate. Thereby, the solvent was distilled off under reduced pressure to obtain the title compound (18 g) having the following physical property values.

- 40 TLC: Rf 0.60 (hexane : ethyl acetate = 2:1);  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 5.22, 7.34 - 7.51, 8.13, 8.86.

Example 13:

- 45 7-(benzyloxy)-4-[(6-chloro-3-pyridinyl)oxy]quinolinone

- [0253]** The compound produced in Example 12 (15 g), 6-chloropyridine-3-ol (8.3 g), 4-dimethylaminopyridine (7.5 g), and toluene (75 mL) were placed in a 300-mL eggplant flask. The mixture was heated at 110°C for 6.5 hours, then left to cool to room temperature. Water and ethyl acetate were added thereto, and the resulting solution was separated. The extract solution was washed with a saturated saline solution, dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. Then, the resulting residue was dissolved in a small amount of ethyl acetate, and crystallized by adding methanol to the solution. The precipitates were then collected by filtration to obtain the title compound (15 g) having the following physical property values.

- 50 TLC: Rf 0.37 (hexane : ethyl acetate = 1:1);  
 55 <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 5.24, 6.46, 7.31 - 7.52, 8.19, 8.34, 8.65.



Example 14:

5-[[7-(benzyloxy)-4-quinolinyl]oxy]-2-pyridinamine

**[0254]** Under argon atmosphere, a solution of the compound produced in Example 13 (5 g) in THF (25 mL), 1.0 mol/L LHMDS (3.5 mL), tris(dibenzylideneacetone) dipalladium(0) chloroform complex (0.63 g), and 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (0.73 g) were placed in a 200-mL 4-diameter eggplant flask, and the mixture was stirred at 70°C. Disappearance of the raw material was confirmed, the resulting product was left to cool to room temperature, then water and ethyl acetate were added, and the solution was separated. After extraction with ethyl acetate, the solution was washed with a saturated saline solution, dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The resulting residue was suspended in acetonitrile (100 mL), and 2.0 mol/L hydrochloric acid (10 mL) was added to the suspension, and the mixture was stirred at room temperature for 2.5 hours. 1.0 mol/L sodium hydroxide aqueous solution, a saturated sodium hydrogen bicarbonate aqueous solution, and ethyl acetate were added, and the solution was separated. The extracted solution was washed with a saturated saline solution, dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1:1→0:1) to obtain the title compound (2.9 g) having the following physical property values.


TLC: Rf 0.33 (dichloromethane : ethyl acetate :methanol = 8:4:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 5.29, 6.06, 6.41, 6.55, 7.31 - 7.52, 7.88, 8.20, 8.56.

Example 15:

N-(5-[[7-(benzyloxy)-4-quinolinyl]oxy]-2-pyridinyl)-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0255]**




**[0256]** The compound produced in Example 14 (800 mg), the compound produced in Example 4 (660 mg), DIPEA (1.2 mL), and DMF (16 mL) were placed in a 30-mL eggplant flask, and finally HATU (1.1 g) was added thereto, followed by stirring at room temperature overnight. After disappearance of the raw material was confirmed, water and ethyl acetate were added, and the resulting solution was separated. The resulting solution was extracted with ethyl acetate, and then washed with water and a saturated saline solution sequentially in this order, and dried over anhydrous sodium sulfate. The solvent was distilled off under reduced pressure. The resulting residue was dissolved in a small amount of ethyl acetate, and crystallized by adding methanol. The precipitate was filtered to obtain the title compound (1.2 g) having the following physical property values. TLC: Rf 0.74 (ethyl acetate :methanol = 9:1); <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.11, 2.59, 5.23, 6.42, 7.25 - 7.63, 8.20, 8.24, 8.49, 8.59, 9.32, 11.92.

Example 16:

N-{5-[[7-(benzyloxy)-4-quinolinyl]oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0257]**



**[0258]** The compound produced in Example 15 (500 mg), 20% palladium hydroxide-carbon (250 mg), ethyl acetate (18 mL), and methanol (2 mL) were placed in a 200-mL eggplant flask. Under hydrogen atmosphere, the mixture was stirred at room temperature for four hours, followed by filtration through Celite. The filtrate was distilled off under reduced pressure to obtain the title compound (360 mg) having the following physical property values.


TLC: R<sub>f</sub> 0.63 (ethyl acetate : methanol = 9:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 2.00, 2.50, 6.44, 7.19, 7.25, 7.46, 7.60, 7.85, 8.15, 8.34, 8.41, 8.53, 8.97, 10.28, 11.95.

Example 17:

N-[5-({7-[3-(dimethylamino)propoxy]-4-quinolinyl}oxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0259]**



**[0260]** The compound produced in Example 16 (50 mg) was dissolved in THF (1 mL) at room temperature, and 3-(dimethylamino)-1-propanol (30 mg), 1,1'-azobis(N,N-dimethyl formamide) (50mg), and tributylphosphine (0.071 mL) were added sequentially. The mixture was stirred at 60°C for 20 hours, and the solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel chromatography (hexane : ethyl acetate = 30:70 → 0:100 → ethyl acetate : methanol = 70:30) to obtain the title compound (22 mg) having the following physical property values.

TLC: R<sub>f</sub> 0.75 (ethyl acetate : methanol = 5:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.11, 2.34, 2.59, 4.19, 6.40, 7.25, 7.40, 7.54, 7.62, 8.20, 8.23, 8.49, 8.59, 9.33, 11.92.

Examples 17(1) to 17(8)

**[0261]** The following Example compounds were obtained by the procedure having the same purpose as in Example 16 → Example 17, using the compound produced in Example 16 and corresponding alcohol derivatives in place of 3-(dimethylamino)-1-propanol.

Example 17(1):

N-(5-([7-(3-hydroxy-3-methylbutoxy)-4-quinolinyl]oxy)-2-pyridinyl)-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0262]** TLC: Rf 0.78 (ethyl acetate : methanol = 5:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.35, 2.09, 2.60, 4.36, 6.41, 7.21, 7.44, 7.55, 7.58-7.65, 8.21, 8.22, 8.49, 8.60, 9.32, 11.92.

Example 17(2):

N-[5-([7-[3-(4-morpholinyl)propoxy]-4-quinolinyl]oxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0263]** TLC: Rf 0.33 (ethyl acetate, DNH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.00-2.20, 2.45-2.52, 2.53-2.65, 3.72-3.75, 4.18-4.22, 6.40-6.42, 7.20-7.30, 7.40-7.41, 7.53-7.70, 8.19-8.22, 8.47-8.50, 8.58-8.60, 9.32, 11.92.

Example 17(3):

N-[5-([7-[2-(4-morpholinyl)ethoxy]-4-quinolinyl]oxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0264]** TLC: Rf 0.80 (ethyl acetate : methanol = 5:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.10, 2.56-2.64, 2.90, 3.76, 4.29, 6.42, 7.22-7.28, 7.40, 7.53-7.65, 8.20, 8.22, 8.49, 8.60, 9.33, 11.92.

Example 17(4):

1-[(2S)-1-hydroxy-3-methyl-2-butanyl]-N-[5-([7-[3-(4-morpholinyl)propoxy]-4-quinolinyl]oxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0265]** TLC: Rf 0.18 (ethyl acetate, DNH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.73, 1.13, 1.98-2.11, 2.12-2.26, 2.38-2.63, 2.82-3.08, 3.18-3.41, 3.70-3.79, 3.99-4.08, 4.09-4.27, 4.56-4.79, 6.26, 7.21, 7.26-7.29, 7.74, 8.18-8.27, 8.48-8.59, 8.86, 12.15.

Example 17(5):

1-(2,2-dimethylpropyl)-N-[5-([7-[3-(4-morpholinyl)propoxy]-4-quinolinyl]oxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0266]** TLC: Rf 0.41 (ethyl acetate, DNH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.06, 2.01-2.13, 2.14-2.30, 2.44-2.52, 2.53-2.69, 3.01-3.16, 3.67-3.79, 4.21, 6.42, 7.23, 7.42, 7.57, 8.23, 8.29, 8.53, 8.60, 9.22, 12.18.

Example 17(6):

1-(1-methyl-1H-pyrazole-4-yl)-N-[5-([7-[3-(4-morpholinyl)propoxy]-4-quinolinyl]oxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0267]** TLC: Rf 0.50 (ethyl acetate : methanol = 10:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.01-2.21, 2.46-2.52, 2.54-2.66, 2.76, 3.69-3.77, 4.03, 4.20, 6.42, 7.22, 7.41, 7.52, 7.53-7.60, 8.17-8.26, 8.48, 8.59, 9.28, 11.90.

Example 17(7):

1-[1-(2-hydroxy-2-methylpropyl)-1H-pyrazole-4-yl]-N-[5-([7-[3-(4-morpholinyl)propoxy]-4-quinolinyl]oxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0268]** TLC: Rf 0.50 (ethyl acetate : methanol = 10:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.28, 2.01-2.22, 2.45-2.53, 2.54-2.66, 2.69-2.80, 3.20, 3.68-3.78, 4.16-4.25, 6.41, 7.19-7.25, 7.41, 7.52-7.60, 7.69, 8.16-8.26, 8.49, 8.59, 9.28, 11.87.

Example 17(8):

1-[(1R)-1-(4-fluorophenyl)ethyl]-N-[5-({7-[3-(4-morpholinyl)propoxy]-4-quinolinyl}oxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0269]** TLC: R<sub>f</sub> 0.62 (ethyl acetate : methanol = 10:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.93-2.17, 2.41-2.64, 2.81-3.13, 3.74, 4.21, 6.42, 7.02-7.13, 7.18-7.26, 7.42, 7.57, 8.23, 8.28, 8.51, 8.60, 9.25, 11.99-12.11.

Example 18:

(4-bromophenyl)(6,7-dimethoxy-4-quinazolinyl)methanone

**[0270]** At room temperature, 4-chloro-6,7-dimethoxyquinazoline (224 mg), 4-bromobenzaldehyde (221 mg), and 1,3-dimethyl imidazolium iodide (74 mg) were placed in a 50-mL eggplant flask, and the mixture was suspended in 1,4-dioxane (3 mL). At room temperature, 60% sodium hydride (52 mg) was added and stirred. Thereafter, the mixture was heated to 100°C and stirred for one hour. The mixture was left to cool to room temperature, the reaction solution was diluted with ethyl acetate (10 mL), and water was added to the solution (10 mL). The precipitated crystals were collected by filtration to obtain the title compound (196 mg) having the following physical property values.

TLC: R<sub>f</sub> 0.31 (hexane : ethyl acetate = 1:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 3.99, 4.10, 7.42, 7.43, 7.64-7.68, 7.84-7.88, 9.23.

Example 19:

2-methyl-2-propanyl {4-[(6,7-dimethoxy-4-quinazolinyl)carbonyl]phenyl}carbamate

**[0271]** Under argon atmosphere, the compound produced in Example 18 (149 mg), tert-butyl carbamate (51 mg), tris(dibenzylideneacetone)dipalladium - chloroform adduct (21 mg), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (35 mg), and cesium carbonate (182 mg) were suspended in 1,4-dioxane (4 mL) in a 50-mL eggplant flask. Thereafter, the suspension was heated to 100°C, stirred for 12 hours, and then left to cool to room temperature. Water was added thereto, followed by extraction with ethyl acetate. The extracted solution was washed with water and a saturated saline solution, and then dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off under reduced pressure, followed by purification by silica gel column chromatography (hexane : ethyl acetate = 1:1) to obtain the title compound (145 mg) having the following physical property values.

TLC: R<sub>f</sub> 0.17 (hexane : ethyl acetate = 1:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.53, 3.96, 4.09, 6.73, 7.33, 7.40, 7.47-7.51, 7.92-7.96, 9.22.

Example 20:

(4-aminophenyl) (6,7-dimethoxy-4-quinazolinyl)methanone hydrochloride

**[0272]** The compound produced in Example 19 (41 mg) was suspended in methanol (0.5 mL) in a 50-mL eggplant flask at room temperature. 4 mol/L hydrogen chloride / ethyl acetate solution (1 mL) was added, and the reaction mixture was stirred at room temperature for one hour, and then concentrated to obtain the title compound (35 mg) having the following physical property values.


TLC: R<sub>f</sub> 0.48 (dichloromethane : methanol = 9:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 3.79, 4.01, 6.57 - 6.60, 7.06, 7.46, 7.54 - 7.57, 9.14.

Example 21:

N-{4-[(6,7-dimethoxy-4-quinazolinyl)carbonyl]phenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0273]**



**[0274]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5 using the compound produced in Example 20 and the compound produced in Example 4.

TLC: Rf 0.55 (dichloromethane : methanol = 9:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  2.06-2.14, 2.54-2.63, 3.95, 4.11, 7.26-7.29, 7.33, 7.41, 7.61-7.70, 7.82-7.86, 7.94-7.97, 9.23, 9.32, 11.70.

Example 22:

4-[(6,7-dimethoxy-4-quinolyl)oxy]aniline

**[0275]** 4-aminophenol (500 mg) was dissolved in dimethyl sulfoxide (DMSO) (5 mL) at room temperature, and 55% sodium hydride (98 mg) was added thereto. Thereafter, 4-chloro-6,7-dimethoxy quinoline (244 mg) was added, and the mixture was stirred at 100°C for three hours. The reaction solution was diluted with ethyl acetate, saturated sodium bicarbonate water was added, and the mixture was stirred. Furthermore, ethyl acetate and water were added, and the organic layer was extracted. The organic layer was washed with a saturated saline solution, and then dried over anhydrous sodium sulfate. The solvent was distilled off under reduced pressure. The resulting residue was washed with methanol to obtain the title compound (442 mg) having the following physical property values.


TLC: Rf 0.57 (ethyl acetate);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  3.71, 4.04, 6.41, 6.76, 6.98, 7.39, 7.57, 8.44.

Example 23:

N-{4-[(6,7-dimethoxy-4-quinolyl)oxy]phenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0276]**



**[0277]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5, using the compound produced in Example 22 and the compound produced in Example 4.

TLC: Rf 0.72 (ethyl acetate : methanol = 5:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  2.11, 2.60, 4.05, 6.49, 7.14, 7.28, 7.43, 7.56, 7.61-7.70, 7.80, 8.47, 9.34, 11.46.

Examples 23(1) to 23(3)

**[0278]** The following Example compounds were obtained by the procedure having the same purpose as in Example

22 → Example 23, using 4-chloro-6,7-dimethoxy quinoline or a corresponding quinoline derivative in place of it, 4-aminophenol or a corresponding phenol derivative in place of it.

Example 23(1):

2,5-dioxo-1-phenyl-N-[4-(4-quinolinylloxy)phenyl]-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0279]** (LC-MS/ELSD): (retention time: 0.79 min);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.05-2.18, 2.53-2.69, 6.56, 7.12-7.20, 7.26-7.34, 7.54-7.85, 8.08, 8.36, 8.66, 9.34, 11.45.

Example 23(2):

N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-methylphenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0280]** TLC: R<sub>f</sub> 0.50 (dichloromethane : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.05-2.15, 2.32, 2.53-2.63, 4.04, 4.05, 6.50, 7.00, 7.06, 7.26-7.30, 7.42, 7.55-7.68, 8.30, 8.48, 9.35, 11.23.

Example 23(3):

N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]-2,6-difluorophenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0281]** TLC: R<sub>f</sub> 0.28 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.06-2.14, 2.55-2.63, 4.03, 4.05, 6.69, 6.78-6.83, 7.26-7.30, 7.41, 7.45, 7.56-7.68, 8.58, 9.32, 10.77.

Example 24:

4-(2-fluoro-4-nitrophenoxy)-6,7-dimethoxy quinoline

**[0282]** 6,7-dimethoxy quinoline-4-ol (5.0 g), cesium carbonate (1.3 g) and 1,2-difluoro-4-nitrobenzene (3.5 mL) were added into DMF (20 mL), and the mixture was stirred at room temperature for six hours. The reaction solution was diluted with ethyl acetate, and then the organic layer was washed with water. The water layer was extracted with ethyl acetate twice, and the combined organic layer was washed with a saturated saline solution, and dried over anhydrous sodium sulfate. The organic layer was concentrated. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 1:1→0:100) to obtain the title compound (2.3 g) having the following physical property values.

TLC: R<sub>f</sub> 0.62 (hexane : ethyl acetate = 1:9);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 4.04, 4.07, 6.55, 7.31-7.38, 7.45, 7.47, 8.11-8.26, 8.19, 8.58.

Example 25:

4-[(6,7-dimethoxyquinoline-4-yl)oxy]-3-fluoroaniline

**[0283]** The compound produced in Example 24 (2.1 g) was dissolved in DMF : water = 3:1 (45 mL). Zinc (3.9 g) and ammonium chloride (1.9 mg) were added to the solution, and the solution was stirred at room temperature for one hour. Then the reaction solution was filtered through Celite. Saturated sodium hydrogen carbonate aqueous solution was added to the filtrate and solid precipitate was removed by filtration through Celite. Ethyl acetate was added thereto, and the water layer was extracted with ethyl acetate. The combined organic layer was washed with a saturated saline solution, and dried over anhydrous sodium sulfate. The organic layer was concentrated. The title compound (1.9 g) having the following physical property values was obtained.


**[0284]** TLC: R<sub>f</sub> 0.35 (hexane : ethyl acetate = 1:9);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 3.93, 5.48, 6.38, 6.42-6.48, 6.54, 7.06, 7.37, 7.49, 8.44.

Example 26:

N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]-3-fluorophenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

[0285]



[0286] The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5 using the compound produced in Example 25 and the compound produced in Example 4.

TLC: R<sub>f</sub> 0.17 (hexane : ethyl acetate = 1:9);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.06-2.20, 2.54-2.68, 4.06, 6.43, 7.15-7.47, 7.57-7.74, 7.94, 8.48, 9.32, 11.55.

Examples 26(1) to 26(3)

[0287] The following Example compounds were obtained by the procedure having the same purpose as in Example 26, using a carboxylic acid derivative was used in place of the compound produced in Example 4.

Example 26(1):

N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]-3-fluorophenyl}-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide


[0288] TLC: R<sub>f</sub> 0.36 (ethyl acetate);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 0.99, 2.00-2.16, 2.54-2.60, 3.11-3.24, 3.94, 4.28, 6.47, 7.40-7.57, 8.05, 8.47, 8.84, 11.81.

Example 26(2):

N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]-3-fluorophenyl}-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

[0289]



[0290] TLC: R<sub>f</sub> 0.32 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.07-2.21, 2.51-2.71, 4.05, 4.06, 6.39-6.45, 7.01-7.13, 7.15-7.23, 7.29-7.40, 7.42, 7.57, 7.60-7.74,

7.85-8.00, 8.48, 9.33, 11.38-11.49.

Example 26(3):

5 N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]-3-fluorophenyl}-2',5'-dioxo-1'-phenyl-2',5',6',8'-tetrahydro-1'H-spiro[cyclopropane-1,7'-quinoline]-3'-carboxamide TLC: Rf 0.33 (ethyl acetate);

10 **[0291]**  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.32-0.44, 0.50-0.61, 2.40, 2.48, 4.05, 4.06, 6.42, 7.14-7.25, 7.34-7.40, 7.42, 7.58, 7.59-7.70, 7.93, 8.49, 9.37, 11.54.

Example 27:

4-[(6-chloro-3-pyridinyl)oxy]-6,7-dimethoxyquinazoline

15 **[0292]** Under argon atmosphere, DMAP (4.4 g) was added to 4-chloro-6,7-dimethoxyquinazoline (8.0 g) and a DMSO suspension (20 mL) of 6-chloropyridine-3-ol (4.6 g), heated and stirred at a bath temperature (80°C) for two hours, and left to cool to room temperature. Thereafter, the reaction solution was diluted with ethyl acetate, and washed with water and saturated sodium hydrogen carbonate aqueous solution. The organic layer was dried over anhydrous sodium sulfate, and concentrated. The resulting residue was washed with hexane - ethyl acetate (3:1) to obtain the title compound (9.1 g) having the following physical property values.

TLC: Rf 0.16 (hexane : ethyl acetate = 1:1);

$^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ):  $\delta$  3.97, 3.99, 7.41, 7.58, 7.69, 7.97, 8.50, 8.57.

Example 28:

5-[(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinamine hydrochloride

25 **[0293]** Under argon atmosphere, 1.0 mol/L LHMDs (4.7 mL), tris(dibenzylideneacetone)dipalladium(0) chloroform complex (140 mg), and 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (170 mg) were added to a THF solution (15 mL) of the compound produced in Example 27 (1.0 g). The mixture was stirred at a bath temperature (70°C) for four hours. After the reaction solution was left to cool to room temperature, and placed into ice water, followed by extraction with ethyl acetate. The organic layer was washed with water, dried over anhydrous sodium sulfate, and concentrated. The resulting residue was suspended in acetonitrile (30 mL), and 2.0 mol/L hydrochloric acid (10 mL) was added thereto and stirred at room temperature for 30 min. Precipitates generated in the reaction solution were collected by filtration to obtain the title compound (591 mg) having the following physical property values.


TLC: Rf 0.16 (ethyl acetate : methanol = 10:1);

$^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ):  $\delta$  3.96, 3.99, 4.24, 7.10, 7.42, 7.53, 8.00-8.20, 8.07, 8.20, 8.61.

Example 29:

40 N-{5-[(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

45 **[0294]**



55 **[0295]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5 using the compound produced in Example 28 and the compound produced in Example 4.



TLC: Rf 0.75 (ethyl acetate : methanol = 5:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.11, 2.59, 4.07, 7.26, 7.33, 7.54, 7.58-7.69, 8.27, 8.52, 8.61, 9.33, 11.91.

Examples 29(1) to 29(6):

**[0296]** The following Example compounds were obtained by the procedure having the same purpose as in Example 27→ Example 28→ Example 29, using corresponding quinoline derivatives in place of 4-chloro-6,7-dimethoxyquinazoline, 6-chloropyridine-3-ol, and the compound produced in Example 4 or corresponding carboxylic acid derivatives in place of it.

Example 29(1):

N-{5-[(7-methoxy-4-quinazolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0297]** (LC-MS/ELSD): (retention time: 0.99 min);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.02-2.21, 2.57, 3.99, 7.24-7.25, 7.27-7.29, 7.30-7.33, 7.55-7.69, 8.23, 8.26, 8.51, 8.67, 9.33, 11.89.

Example 29(2):

1-cyclobutyl-N-{5-[(6,7-dimethoxy-4-quinazolinyloxy)-2-pyridinyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0298]** TLC: Rf 0.47 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.75-1.95, 1.96-2.10, 2.11-2.25, 2.40-2.54, 2.55-2.65, 2.92-3.16, 4.08, 4.09, 4.74-4.93, 7.34, 7.56, 7.68, 8.36, 8.56, 8.62, 9.16, 12.13.

Example 29(3):

N-{5-[(6,7-dimethoxy-4-quinazolinyloxy)-2-pyridinyl]-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0299]** TLC: Rf 0.44 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.06, 2.18-2.27, 2.63, 3.02-3.18, 4.08, 4.09, 7.34, 7.56, 7.69, 8.36, 8.56, 8.61, 9.22, 12.17.

Example 29(4):

N-{5-[(6,7-dimethoxy-4-quinazolinyloxy)-2-pyridinyl]-1-[(2S)-1-hydroxy-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide


**[0300]** TLC: Rf 0.38 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.74, 1.13, 2.09-2.34, 2.41-2.72, 2.77-3.39, 3.90-4.20, 4.07, 4.09, 4.44-4.69, 7.32, 7.55, 7.70, 8.32, 8.53, 8.60, 9.15, 12.07.

Example 29(5):

N-{5-[(6,7-dimethoxy-4-quinazolinyloxy)-2-pyridinyl]-1-(4-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0301]**



**[0302]** TLC: Rf 0.55 (hexane : ethyl acetate = 1:9, NH silica);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  2.12, 2.56-2.63, 4.07, 7.23-7.35, 7.53, 7.66, 8.26, 8.50, 8.60, 9.30, 11.84.

Example 29(6):

N-{5-[(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinyl}-1-[(2R)-1-hydroxy-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0303]** TLC: Rf 0.20 (ethyl acetate, NH silica);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.72, 1.11, 2.22, 2.52, 2.94, 3.34, 4.07, 4.08, 4.16, 4.40, 4.72, 7.27, 7.54, 7.72, 8.24, 8.45, 8.56, 8.86, 91.5, 12.14.

Example 30:

4-(2-fluoro-4-nitrophenyl)-7-methoxyquinazoline

**[0304]** 4-chloro-7-methoxy-quinazoline (50 mg) and 2-fluoro-4-nitrophenol (60 mg) were added to diphenyl ether (10 mL), and the mixture was stirred at 150°C for two hours by using microwave. The reaction solution was cooled, and water was added. Thereafter, the water layer was extracted with ethyl acetate, and the combined organic layer was washed with a saturated saline solution, and then concentrated. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 100:0  $\rightarrow$  0:100) to obtain the title compound (60 mg) having the following physical property values. MASS (ESI, Pos.): 286 (M+H)<sup>+</sup>.

Example 31:

3-fluoro-4-[7-methoxyquinazoline-4-yl]oxy]aniline


**[0305]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 25, using the compound produced in Example 30.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  3.70, 3.98, 6.49-6.57, 7.06, 7.24-7.30, 8.25, 9.47.

Example 32:

N-{3-fluoro-4-[(7-methoxy-4-quinazolinyl)oxy]phenyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0306]**



**[0307]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5, using the compound produced in Example 31 and the compound produced in Example 4. (LC-MS/ELSD): (retention time: 1.02 min);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  2.06-2.20, 2.53-2.68, 3.98, 7.19-7.44, 7.58-7.73, 7.93, 8.25, 8.67, 9.32, 11.51.

Example 33:

tert-butyl [4-(quinazoline-4-yloxy)phenyl]carbamate

**[0308]** 4-chloro-quinazoline (0.95 g), calcium carbonate (3.5 g), and (4-hydroxy-phenyl)-carbamic acid tert-butyl ester (0.90 g) were added to acetonitrile (10 mL), and the mixture was stirred at 100°C for three hours. Water was added to the reaction solution, and the water layer was extracted with ethyl acetate. The combined organic layer was washed with a saturated saline solution, and then concentrated. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 100:0  $\rightarrow$  0:100) to obtain the title compound (1.3 g) having the following physical property values.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.53, 6.63, 7.18-7.20, 7.47, 7.64-7.68, 7.89-7.93, 8.00, 8.37, 8.76.

Example 34:


4-(quinazoline-4-yloxy)aniline hydrochloride

**[0309]** 4 mol/L hydrochloric acid-1,4-dioxane solution (5 mL) was added to a THF solution (5 mL) of the compound produced in Example 33 (100 mg), and the mixture was stirred at room temperature for three hours. The reaction solution was concentrated to obtain the title compound (81 mg) having the following physical property value. MASS (ESI, Pos.): 238 ( $\text{M}+\text{H}$ ) $^+$ .

Example 35:

2,5-dioxo-1-phenyl-N-[4-(4-quinazolinylxy)phenyl]-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0310]**



**[0311]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5, using the compound produced in Example 34 and the compound produced in Example 4.

(LC-MS/ELSD): (retention time: 0.98 min);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  2.08-2.13, 2.54-2.63, 7.21-7.29, 7.60-7.68, 7.82-7.84, 7.91-7.93, 8.00, 8.37, 8.76, 9.34, 11.43.

Example 36:

5-[[3-(3-hydroxy-4-methoxyphenyl)amino]methylene]-2,2-dimethyl-1,3-dioxane-4,6-dione

**[0312]** Under argon atmosphere, a methyl orthoformate (50 mL) solution of meldrum's acid (8.0 g) was heated at 100°C for 5 min, and then 3-hydroxy-4-methoxyaniline (7.0 g) was added to the solution, and the resulting solution was heated and stirred at 105°C for 25 min. The heating was stopped to cool with water, and precipitated powder was collected by filtration, washed with methyl orthoformate and MTBE, and dried under reduced pressure to obtain the title compound (12.3 g) having the following physical property values.

TLC: Rf 0.25 (hexane : ethyl acetate = 1:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.75, 3.92, 5.82, 6.73, 6.84-6.88, 8.53, 11.2.

Example 37:

5-([3-(benzyloxy)-4-methoxyphenyl]amino)methylene)-2,2-dimethyl-1,3-dioxane-4,6-dione

**[0313]** Under argon atmosphere, the compound produced in Example 36 (12.0 g) was dissolved in DMA (80 mL) at 50°C. The temperature was returned to room temperature. Then, potassium carbonate (7.35 g) and benzyl bromide (8.75 g) were added thereto. The resulting solution was heated and stirred at 60°C for two hours. The mixture was left to cool to room temperature, and the solvent was concentrated, followed by adding ethyl acetate and water and shaking thereof. Since deposits were precipitated, the deposits were collected by filtration, washed with water and ethyl acetate, and dried under reduced pressure to obtain the title compound (8.0 g) having the following physical property values.


TLC: Rf 0.49 (hexane : ethyl acetate = 1:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.75, 3.90, 5.17, 6.76-6.85, 6.91, 7.30-7.48, 8.48, 11.2.

Example 38:

N-[5-({6-methoxy-7-[3-(4-morpholinyl)propoxy]-4-quinolinyl}oxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0314]**



**[0315]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 11 → Example 12 → Example 13 → Example 14 → Example 15 → Example 16 → Example 17, using the compound produced in Example 37.

TLC: Rf 0.20 (ethyl acetate, NH silica);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  2.08-2.19, 2.45-2.64, 3.73, 4.02, 4.27, 6.43, 7.25, 7.43, 7.52-7.65, 8.21, 8.47, 8.49, 9.32, 11.92.


Examples 38(1) to 38(7):

**[0316]** The following Example compounds were obtained by the procedure having the same purpose as in Example 38, using a corresponding compound in place of the compound produced in Example 37.

Example 38(1):

N-(5-{[7-(3-hydroxy-3-methylbutoxy)-6-methoxy-4-quinolinyl]oxy}-2-pyridinyl)-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

[0317]



[0318] TLC: Rf 0.20 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.34, 2.07-2.15, 2.56-2.64, 3.28, 4.01, 4.43, 6.45, 7.25, 7.43, 7.53-7.67, 8.21, 8.48, 8.51, 9.33, 11.93.

Example 38(2):

N-[5-({6-methoxy-7-[3-(1-piperidiny)propoxy]-4-quinolinyl}oxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide


[0319] TLC: Rf 0.28 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.42, 1.52-1.64, 2.08-2.15, 2.38-2.45, 2.50-2.63, 4.02, 4.24, 6.42, 7.21-7.28, 7.42, 7.51, 7.55-7.66, 8.21, 8.47, 8.49, 9.32, 11.92.

Example 38(3):

N-[5-({6-methoxy-7-[3-(1-pyrrolidiny)propoxy]-4-quinolinyl}oxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

[0320]



[0321] TLC: Rf 0.25 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.79, 2.15, 2.56, 2.67, 4.02, 4.27, 6.42, 7.25, 7.42, 7.51, 7.53-7.65, 8.21, 8.47, 8.49, 9.32, 11.92.

Example 38(4):

N-[5-({6-methoxy-7-[3-(1-piperazinyl)propoxy]-4-quinolinyloxy}-2-pyridinyl)-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0322]** TLC: Rf 0.20 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.10, 2.55, 2.71, 3.15, 4.01, 4.26, 6.44, 7.26, 7.42, 7.51-7.67, 8.21, 8.49, 9.31, 11.92.

Example 38(5):

N-(5-{[7-(2-buty-1-yloxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0323]** TLC: Rf 0.56 (ethyl acetate : methanol = 10:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.86, 2.00-2.21, 2.48-2.74, 4.04, 4.90, 6.45, 7.22-7.31, 7.50-7.68, 8.22, 8.45-8.54, 9.33, 11.94.

Example 38(6):

N-[5-({6-methoxy-7-[3-(methylsulfonyl)propoxy]-4-quinolinyloxy}-2-pyridinyl)-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0324]** TLC: Rf 0.16 (ethyl acetate : methanol = 10:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.02-2.19, 2.40-2.68, 2.99, 3.29-3.40, 4.02, 4.32, 6.45, 7.22-7.29, 7.40, 7.49-7.69, 8.22, 8.44-8.55, 9.33, 11.94.

Example 38(7):

N-[5-({6-methoxy-7-[3-(4-methyl-1-piperazinyl)propoxy]-4-quinolinyloxy}-2-pyridinyl)-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0325]** TLC: Rf 0.20 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.13, 2.29, 2.48-2.65, 4.02, 4.25, 6.42, 7.21-7.28, 7.42, 7.51, 7.52-7.67, 8.21, 8.47, 8.49, 9.32, 11.92.

Example 39:

N-[5-({7-[(3-methyl-2-buten-1-yl)oxy]-4-quinolinyloxy}-2-pyridinyl)-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0326]** Under argon atmosphere, cesium carbonate (94 mg) and 1-bromo-3-methyl-2-butene (140 mg) were added to a DMF solution (15 mL) of the compound produced in Example 16 (100 mg), and the mixture was stirred at a bath temperature (-10°C) for two hours. An ammonium chloride aqueous solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with a saturated saline solution, dried over anhydrous sodium sulfate, and concentrated, followed by purification by silica gel column chromatography (hexane : ethyl acetate = 1:4, NH silica) to obtain the title compound (55 mg).


TLC: Rf 0.30 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.79, 1.82, 2.08-2.17, 2.46-2.63, 4.67, 5.57, 6.39, 7.23-7.26, 7.46, 7.48-7.65, 8.18, 8.22, 8.45, 8.49, 9.29, 11.90.

Example 40:

N-{5-[(7-[(2R)-2,3-dihydroxy-3-methylbutyl]oxy]-4-quinolinyloxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0327]**



**[0328]** Under argon atmosphere, tert-butyl alcohol (0.5 mL), water (0.5 mL), AD-mix- $\beta$  (50 mg), and methane sulfonamide (7 mg) were added to a dichloromethane solution (0.5 mL) of the compound produced in Example 39 (20 mg), and the mixture was stirred at room temperature for six hours. A sodium thiosulfate aqueous solution was added to the reaction solution, followed by extraction with ethyl acetate. The organic layer was washed with a saturated saline solution, dried over anhydrous sodium sulfate, and concentrated, followed by purification by silica gel column chromatography (ethyl acetate, NH silica) to obtain the title compound (3 mg).

TLC: Rf 0.10 (ethyl acetate, NH silica);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.32, 1.36, 2.11, 2.37, 2.60, 2.87, 3.92, 4.20, 4.32, 6.42, 7.23, 7.43, 7.53-7.66, 8.20, 8.23, 8.48, 8.60, 9.32, 11.93.

Examples 40(1) to 40(6)

**[0329]** The following Example compounds were obtained by the procedure having the same purpose as in Example 39  $\rightarrow$  Example 40, using a corresponding compound in place of the compound produced in Example 16, 1-bromo-3-methyl-2-butene, and AD-mix- $\beta$  or AD-mix- $\alpha$ .

Example 40(1):

N-{5-[(7-[(2S)-2,3-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0330]** TLC: Rf 0.41 (ethyl acetate : methanol = 10:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.32, 1.36, 1.99-2.19, 2.50-2.68, 3.04, 3.15-3.23, 3.74-3.90, 4.01, 4.21-4.33, 4.37-4.48, 6.45, 7.24-7.29, 7.42, 7.54, 7.54-7.69, 8.18-8.23, 8.46-8.52, 9.32, 11.93.

Example 40(2):

N-{5-[(7-[(2R)-2,3-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0331]** TLC: Rf 0.48 (dichloromethane : methanol = 9:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.28-1.32, 2.07-2.15, 2.56-2.64, 2.90-3.40, 3.85, 4.01, 4.27, 4.43, 6.45, 7.25-7.29, 7.42, 7.53-7.66, 8.21, 8.48, 8.50, 9.32, 11.93.

Example 40(3):

N-{5-[(7-[(2S)-2,3-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0332]** TLC: Rf 0.41 (ethyl acetate : methanol = 10:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.06, 1.33, 1.37, 2.11-2.33, 2.55-2.71, 3.04, 3.06-3.15, 3.16-3.24, 3.79-3.91, 4.03, 4.21-4.33, 4.39-4.48, 6.46, 7.43, 7.53-7.61, 8.27-8.33, 8.48-8.56, 9.22, 12.18.

Example 40(4):

N-{5-[(7-[(2R)-2,3-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0333]** TLC: Rf 0.45 (dichloromethane : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.06, 1.33-1.37, 2.18-2.26, 2.63, 3.00-3.20, 3.87, 4.02, 4.28, 4.44, 6.46, 7.43, 7.55-7.60, 8.30, 8.49-8.55, 9.21, 12.18.

Example 40(5):

**[0334]** N-{5-[(7-[(2R)-2,3-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-2',5'-dioxo-1'-phenyl-2',5',6',8'-tetrahydro-1'H-spiro[cyclopropane-1,7'-quinoline]-3'-carboxamide

**[0335]** TLC: Rf 0.49 (dichloromethane : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.37-0.40, 0.53-0.56, 1.32-1.36, 2.41, 2.48, 2.98, 3.19, 3.85, 4.01, 4.27, 4.43, 6.46, 7.21-7.24, 7.42, 7.54-7.65, 8.22, 8.49-8.52, 9.36, 11.94.

Example 40(6):

N-{5-[(7-[(2S)-2,3-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0336]** TLC: Rf 0.56 (ethyl acetate : methanol = 10:1, DNH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.32, 1.36, 2.05-2.22, 2.53-2.71, 2.98-3.09, 3.11-3.26, 3.77-3.92, 4.01, 4.20-4.30, 4.39-4.47, 6.45, 7.00-7.13, 7.28-7.36, 7.42, 7.51-7.69, 8.23, 8.45-8.54, 9.32, 11.86.

Example 41:

bis(2-methyl-2-propanyl)(5-[(7-benzyloxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl)imidodicarbonate

**[0337]** 5-[(7-(benzyloxy)-6-methoxy-4-quinolinyloxy)-2-pyridinyl]amine (1.6 g) was dissolved in a 1,3-dimethyl-3,4,5,6-tetrahydro-2-(1H)-pyrimidinone solution (15 mL) of di-tert-butyl dicarbonate (CAS registration No.:24424-99-5) (3.95 mL). Triethyl amine (9.37 mL) and DMAP (52 mg) were added to the resulting solution, and the solution was stirred at room temperature for five hours. The reaction solution was diluted with hexane : ethyl acetate mixed solution (1:3), and washed with water. Then, the organic layer was concentrated, and purified by silica gel column chromatography (hexane : ethyl acetate = 4:1 → 0:1 → ethyl acetate :methanol = 50:1 → 20:1) to obtain the title compound (2.4 g) having the following physical property values.

TLC: Rf 0.51 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.49, 4.04, 5.33, 6.45, 7.30-7.43, 7.44-7.62, 8.41, 8.50.

Example 42:

bis(2-methyl-2-propanyl){5-[(7-hydroxy-6-methoxy-4-quinolinyloxy)-2-pyridinyl]imidodicarbonate

**[0338]** Under argon atmosphere, ethyl acetate (10 mL) and ethanol (30 mL) were added to and dissolved in the compound produced in Example 41 (1.4 g). Palladium hydroxide (20 wt%, 420 mg) was added thereto, and the mixture was stirred under hydrogen atmosphere at room temperature for five hours. Palladium hydroxide was removed through Celite, and the filtrate was concentrated to obtain the title compound (1.2 g).

TLC: Rf 0.49 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.50, 4.08, 6.44, 7.36, 7.46-7.60, 8.42, 8.53.

Example 43:

bis(2-methyl-2-propanyl)[5-[(6-methoxy-7-[(3-methyl-3-buten-1-yl)oxy]-4-quinolinyloxy)-2-pyridinyl]imidodicarbonate

**[0339]** 3-methyl-3-buten-1-ol (231 mg), N,N,N',N'-tetramethyl azodicarboxamide (TMAD) (462 mg), and tri-n-butylphosphine (544 mg) were added to a THF suspension (20 mL) of the compound produced in Example 42 (500 mg), and the mixture was stirred at room temperature for three hours. The reaction solution was diluted with ethyl acetate, and



concentrated by removing insoluble matters. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 3:2 → 0:1) to obtain the title compound (570 mg) having the following physical property values.

TLC: R<sub>f</sub> 0.52 (hexane : ethyl acetate = 3:7);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.49, 1.86, 2.68, 4.03, 4.33, 4.76-4.99, 6.46, 7.35, 7.45, 7.50, 7.55, 8.42, 8.52.

Example 44:

bis(2-methyl-2-propanyl){5-[[7-[(3S)-3,4-dihydroxy-3-methylbutoxy]-6-methoxy-4-quinolinyl]oxy]-2-pyridinyl}imidodi-carbonate

**[0340]** Dichloromethane : t-butanol : water = 1:1:1 (1.5 mL) was added to and dissolved in the compound produced in Example 43 (170 mg), AD-mix-β (431 mg) and methane sulfonamide (58 mg) were added thereto, and the mixture was stirred at room temperature overnight. The reaction solution was diluted with ethyl acetate, and washed with water. The organic layer was collected and purified by silica gel column chromatography (hexane : ethyl acetate = 4:1 → 0:1 → ethyl acetate : methanol = 9:1) to obtain the title compound (152 mg) having the following physical property values.

TLC: R<sub>f</sub> 0.54 (ethyl acetate : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.29, 1.50, 2.06-2.17, 2.19-2.36, 2.90-3.32, 3.45-3.65, 4.02, 4.28-4.51, 6.48, 7.36, 7.45, 7.51, 7.56, 8.42, 8.52.

Example 45:

4-(4-{4-[(6-amino-3-pyridinyl)oxy]-6-methoxy-7-quinolinyl}oxy)-(2S)-2-methyl-1,2-butanediol

**[0341]** Trifluoroacetic acid (2 mL) was added to a dichloromethane solution (5 mL) of the compound produced in Example 44 (152 mg). The mixture was stirred at room temperature for two hours. The reaction solution was concentrated, and subjected to azeotrope with toluene twice. The resulting residue was purified by silica gel column chromatography (NH silica, ethyl acetate : methanol = 9:1) to obtain the title compound (100 mg) having the following physical property values.


TLC: R<sub>f</sub> 0.35 (dichloromethane : methanol = 9:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.28, 2.01-2.15, 2.17-2.34, 2.74-2.94, 3.08-3.23, 3.44-3.66, 4.03, 4.26-4.46, 4.52, 6.43, 6.61, 7.31, 7.41, 7.56, 8.03, 8.48.

Example 46:

N-{5-[[7-[(3S)-3,4-dihydroxy-3-methylbutyl]oxy]-6-methoxy-4-quinolinyl]oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0342]**



**[0343]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5, using the compound produced in Example 45 and the compound produced in Example 4.

TLC: 0.43 (dichloromethane : methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.28, 2.05-2.30, 2.56-2.64, 2.70-3.30, 3.48-3.60, 4.02, 4.31-4.45, 6.45, 7.25-7.28, 7.33, 7.43-7.67, 8.21, 8.48-8.51, 9.32, 11.93.

Examples 46(1) to 46(3)

**[0344]** The following Example compounds were obtained by the procedure having the same purpose as in Example 46, using the compound produced in Example 45, or a compound produced by subjecting the compound produced in Example 43 to the procedure having the same purpose as in Example 44 → Example 45 using AD-mix- $\alpha$  in place of AD-mix- $\beta$ , and the compound produced in Example 4 or a corresponding carboxylic acid derivative in place of it.

Example 46(1):

N-{5-[(7-[(3R)-3,4-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0345]** TLC: 0.40 (ethyl acetate : methanol = 10:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.28, 1.97-2.17, 2.18-2.35, 2.52-2.69, 2.74-2.89, 3.13, 3.40-3.69, 4.02, 4.23-4.52, 6.45, 7.20-7.30, 7.42, 7.53, 7.54-7.70, 8.21, 8.45-8.52, 9.32, 11.93.

Example 46(2):

N-{5-[(7-[(3S)-3,4-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-2',5'-dioxo-1'-phenyl-2',5',6',8'-tetrahydro-1'H-spiro[cyclopropane-1,7'-quinoline]-3'-carboxamide

**[0346]** TLC: 0.45 (dichloromethane : methanol = 9:1);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.37-0.41, 0.53-0.56, 1.28, 2.05-2.30, 2.42, 2.48, 2.70-3.30, 3.48-3.60, 4.02, 4.31-4.45, 6.45, 7.21-7.24, 7.42, 7.48-7.65, 8.21, 8.48-8.51, 9.36, 11.93.

Example 46(3):

**[0347]** N-{5-[(7-[(3R)-3,4-dihydroxy-3-methylbutyl]oxy)-6-methoxy-4-quinolinyloxy]-2-pyridinyl}-2',5'-dioxo-1'-phenyl-2',5',6',8'-tetrahydro-1'H-spiro[cyclopropane-1,7'-quinoline]-3'-carboxamide

TLC: 0.50 (ethyl acetate : methanol = 10:1, NH silica);

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.33-0.43, 0.50-0.61, 1.28, 1.98-2.14, 2.16-2.33, 2.42, 2.48, 2.73-2.91, 3.08-3.18, 3.42-3.68, 4.02, 4.23-4.51, 6.46, 7.19-7.25, 7.42, 7.54, 7.55-7.66, 8.22, 8.46-8.54, 9.36, 11.94.

Example 47:

(5E)-5-(hydroxyimino)-4-methyl-2-oxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinoline carboxylic acid

**[0348]** A pyridine solution (1.0 mL) of the compound produced in Example 4 (142 mg) and hydroxylamine hydrochloride (208 mg) was heated and refluxed for one hour. The solution was cooled to room temperature. The reaction solution was diluted with ethyl acetate and washed with 1 mol/L hydrochloric acid. The obtained organic layer was concentrated to obtain the title compound (151 mg) having the following physical property values.


TLC: R<sub>f</sub> 0.42 (dichloromethane : methanol = 9:1);

$^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ):  $\delta$  1.63-1.81, 2.26-2.36, 2.60, 7.38-7.46, 7.51-7.68, 8.90, 11.30, 14.03.

Example 48:

N-[(5E)-5-[(6,7-dimethoxy-4-quinolinyloxy)-2-pyridinyl]-5-(hydroxyimino)-2-oxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0349]**



**[0350]** The title compound having the following physical property values were obtained by the procedure having the same purpose as in Example 5, using the compound produced in Example 47 and the compound produced in Example 2. TLC: 0.15 (ethyl acetate, NH silica); <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.77-1.94, 2.38, 2.77, 4.05, 6.46, 7.20-7.25, 7.43, 7.50-7.65, 7.99-8.07, 8.22, 8.48, 8.52, 9.33, 12.23.

Example 48(1):

N-{5-[(6,7-dimethoxy-4-quinolinyloxy]-2-pyridinyl}-(5E)-5-(methoxyimino)-2-oxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0351]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 48, using a corresponding carboxylic acid derivative in place of the compound produced in Example 47 and the compound produced in Example 2.

TLC: 0.18 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.74-1.92, 2.29-2.40, 2.62-2.75, 4.00, 4.05, 6.45, 7.20-7.28, 7.42, 7.50-7.66, 8.20-8.24, 8.46-8.52, 9.36, 12.26.

Example 49:

7-(benzyloxy)-4-(2-fluoro-4-nitrophenoxy)-6-methoxyquinoline

**[0352]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 24, using 7-benzyloxy-6-methoxy-quinoline-4-ol in place of 6,7-dimethoxy quinoline-4-ol. TLC: R<sub>f</sub> 0.76 (hexane : ethyl acetate = 1:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 4.03, 5.34, 6.54, 7.28-7.43, 7.45, 7.47-7.55, 8.10-8.15, 8.19, 8.56.

Example 50:

4-(2-fluoro-4-nitrophenoxy)-6-methoxy-7-quinolinol hydrobromate

**[0353]** 5.1 mol/L hydrobromic acid - acetic acid solution (10 mL) was added to the compound produced in Example 49 (1.6 g). The mixture was stirred at room temperature for five hours. MTBE (50 mL) was added to the reaction solution, and the resulting solution was stirred. Generated precipitates were collected by filtration to obtain the title compound (1.5 g) having the following physical property values.

<sup>1</sup>H-NMR (CD<sub>3</sub> OD): δ 4.14, 7.05, 7.43, 7.76-7.85, 7.86, 8.30-8.38, 8.39-8.46, 8.69.

Example 51:

4-(2-fluoro-4-nitrophenoxy)-6-methoxy-7-[3-(4-morpholinyl)propoxy]quinoline

**[0354]** Cesium carbonate (4.0 g) and 4-(3-chloropropyl)morpholine (517 mg) were added to a DMF solution (9.8 mL) of the compound produced in Example 50 (1.0 g), and the mixture was stirred at 60°C for 16 hours. The mixture solution was diluted with ethyl acetate, and washed with water. The water layer was extracted with ethyl acetate twice. The combined organic layer was dried over anhydrous sodium sulfate and concentrated. The resulting residue was purified

by silica gel column chromatography (NH silica, hexane : ethyl acetate = 1:1 → 0:1) to obtain the title compound (600 mg) having the following physical property values.

TLC: R<sub>f</sub> 0.50 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.90-2.07, 2.32-2.43, 3.25-3.35, 3.48-3.64, 3.92, 4.21, 6.77, 7.45, 7.55-7.67, 8.14-8.25, 8.41-8.50, 8.56.

Example 52:


3-fluoro-4-((6-methoxy-7-(3-morpholinopropoxy)quinoline-4-yl)oxy) aniline

**[0355]** Under argon atmosphere, the compound produced in Example 51 (300 mg) was dissolved in ethyl acetate : ethanol = 1: 1 (30 mL), and palladium hydroxide (20 wt%, 99 mg) was added to the solution, and the argon atmosphere was replaced with a hydrogen atmosphere. After the solution was stirred at room temperature for eight hours, the reaction solution was filtered through Celite, and the filtrate was concentrated to obtain the title compound (240 mg) having the following physical property values. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.03-2.23, 2.43-2.53, 2.57, 3.66-3.77, 3.86, 4.03, 4.26, 6.40, 6.45-6.61, 7.03, 7.42, 7.58, 8.46.

Example 53:

N-[3-fluoro-4-((6-methoxy-7-[3-(4-morpholinyl)propoxyl-4-quinolinyloxy)phenyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0356]**



**[0357]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5, using the compound produced in Example 52 and the compound produced in Example 2.

TLC: R<sub>f</sub> 0.71 (ethyl acetate : methanol = 10:1, NH silica);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 1.88-2.07, 2.34-2.41, 2.41-2.46, 2.51-2.57, 3.31-3.37, 3.52-3.64, 3.94, 4.20, 6.42-6.49, 7.39, 7.40-7.53, 7.54-7.71, 8.04, 8.46, 8.95, 11.64.

Example 54:

methyl 2-oxo-2,5,6,7-tetrahydro-1H-cyclopenta[b]pyridine-3-carboxylate

**[0358]** 2-oxo-1,5,6,7-tetrahydro-cyclopenta[b]pyridine-3-carboxylic acid (CAS registration No.: 115122-63-9) (200 mg) was dissolved in methanol (20 mL) at room temperature, and concentrated sulfuric acid (0.006 mL) was added to the solution. The mixture was stirred at a bath temperature (70°C) for four hours. The solution was left to cool to room temperature, the solvent was distilled off under reduced pressure, and a sodium hydrogen bicarbonate aqueous solution and dichloromethane were added thereto, and the solution was separated. The organic layer was washed with a saturated saline solution, and then was dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure to obtain the title compound (155 mg) having the following physical property values.

TLC: R<sub>f</sub> 0.21 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.13-2.20, 2.80-2.84, 2.97-3.03, 3.91, 8.10.

Example 55:

methyl 2-oxo-1-phenyl-2,5,6,7-tetrahydro-1H-cyclopenta[b]pyridine-3-carboxylate

**[0359]** The compound produced in Example 54 (140 mg) was dissolved in dichloromethane (7 mL) at room temperature. Phenylboronic acid (220 mg), copper acetate (263 mg), and pyridine (0.234 mL) were added to the solution, and the solution was stirred for 20 hours. The solution was filtered through a glass filter, and the solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 8:2 → ethyl acetate) to obtain the title compound (131 mg) having the following physical property values.

TLC: Rf 0.51 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.02-2.12, 2.53-2.58, 2.83-2.88, 3.88, 7.20-7.23, 7.44-7.53, 8.21.

Example 56:

2-oxo-1-phenyl-2,5,6,7-tetrahydro-1H-cyclopenta[b]pyridine-3-carboxylic acid


**[0360]** The compound produced in Example 55 (120 mg) was dissolved in methanol (2 mL) at room temperature. 2 mol/L sodium hydroxide aqueous solution (0.891 mL) was added to the solution, and the solution was stirred for one hour. 2N hydrochloric acid (0.891 mL) and ethyl acetate were added to the reaction solution, and the solution was separated. The organic layer was washed with a saturated saline solution, then was dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure to obtain the title compound (109 mg) having the following physical property values. TLC: Rf 0.64 (ethyl acetate);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.09-2.19, 2.63-2.68, 2.92-2.97, 7.25-7.30, 7.52-7.62, 8.51, 14.24.

Example 57:

N-[5-[(6,7-dimethoxy-4-quinolinyl)oxyl-2-pyridinyl]-2-oxo-1-phenyl-2,5,6,7-tetrahydro-1H-cyclopenta[b]piperidine-3-carboxamide

**[0361]**



**[0362]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5, using the compound produced in Example 56 and the compound produced in Example 2. TLC: Rf 0.65 (ethyl acetate : methanol = 9:1);

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): δ 2.03-2.13, 2.60-2.65, 2.92-2.97, 3.97, 3.99, 6.58, 7.45, 7.47-7.52, 7.57, 7.58-7.68, 7.86-7.90, 8.38, 8.48, 8.53, 8.60, 12.58.

Examples 57(1) to 57(2)

**[0363]** The Example compounds each having the following physical property values were obtained by the procedure having the same purpose as in Example 55 → Example 56 → Example 57, using a corresponding derivative in place of the compound produced in Example 54 and the compound produced in Example 2.

Example 57(1):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2-oxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0364]** TLC: Rf 0.63 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.71-1.83, 2.20-2.29, 2.68-2.76, 4.05, 6.44, 7.19-7.24, 7.42, 7.46-7.62, 8.21, 8.44-8.50, 12.51.

Example 57(2):

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-2,5,6,7-tetrahydro-1H-cyclopenta[b]piperidine-3-carboxamide

**[0365]** TLC: Rf 0.63 (ethyl acetate :methanol = 9:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.68-2.81, 4.05, 6.45, 7.35, 7.43, 7.53, 7.55-7.68, 8.24, 8.49, 8.51, 9.03, 11.91.

Example 58:

N-{5-[(7-[(4S)-2,2-dimethyl-1,3-dioxolane-4-yl]methoxy)-6-methoxy-4-quinolinyl]oxy}-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0366]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 17, using the compound obtained by subjecting the compound produced in Example 37 to procedure having the same purpose as in Example 11 → Example 12 → Example 13 → Example 14 → Example 15 → Example 16, and (S)-(2,2-dimethyl-1,3-dioxolane-4-yl)methanol.


TLC: Rf 0.60 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.42, 1.50, 2.11, 2.60, 4.01, 4.14-4.29, 4.64, 6.44, 7.42, 7.52-7.65, 8.21, 8.47, 8.50, 9.32, 11.92.

Example 59:

N-{5-[(7-[(2R)-2,3-dihydroxypropyl]oxy)-6-methoxy-4-quinolinyl]oxy}-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0367]**



**[0368]** The compound produced in Example 58 (35 mg) was dissolved in THF (1 mL) at room temperature. Methanol (1 mL) and p-toluenesulfonic acid monohydrate (3.0 mg) were added to the solution, and the mixture was stirred room temperature for 16 hours. The reaction solution was diluted with ethyl acetate, and washed with water and a saturated sodium hydrogen carbonate aqueous solution. The organic layer was dried over anhydrous sodium sulfate, and then concentrated. The resulting residue was purified by silica gel chromatography (ethyl acetate : methanol = 100:0 → 70:30) to obtain the title compound (32 mg) having the following physical property values.

TLC: Rf 0.55 (ethyl acetate : methanol = 5:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.11, 2.56-2.64, 3.89, 4.02, 4.21-4.36, 6.45, 7.25-7.28, 7.44, 7.53-7.66, 8.21, 8.48-8.51, 9.32, 11.93.

Examples 59(1) to 59(3)

**[0369]** The following Example compounds were obtained by the procedure having the same purpose as in Example 59, using the compound produced in Example 58 or a corresponding compound in place of it.

Example 59(1):

N-{5-[(7-[(2S)-2,3-dihydroxypropyl]oxy)-6-methoxy-4-quinolinyl]oxy}-2-pyridinyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0370]** TLC: Rf 0.55 (ethyl acetate : methanol = 5:1, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.11, 2.56-2.64, 3.90, 4.02, 4.21-4.36, 6.44, 7.24-7.29, 7.44, 7.52-7.66, 8.22, 8.48-8.51, 9.33, 11.94.

Example 59(2):

N-{5-[(7-[(2S)-2,3-dihydroxypropyl]oxy)-6-methoxy-4-quinolinyl]oxy}-2-pyridinyl-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0371]** TLC: Rf 0.35 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.06, 2.22, 2.63, 3.09, 3.89, 4.02, 4.22, 4.34, 6.44, 7.43, 7.55, 7.58, 8.30, 8.52, 8.54, 9.21, 12.18.

Example 59(3):

N-{5-[(7-[(2R)-2,3-dihydroxypropyl]oxy)-6-methoxy-4-quinolinyl]oxy}-2-pyridinyl-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0372]** TLC: Rf 0.35 (ethyl acetate, NH silica);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.06, 2.22, 2.63, 3.09, 3.89, 4.02, 4.22, 4.34, 6.44, 7.43, 7.55, 7.58, 8.30, 8.52, 8.54, 9.21, 12.18.

Example 60:

5-hydroxy-N-{5-[(7-hydroxy-6-methoxy-4-quinolinyl]oxy}-2-pyridinyl-2-oxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0373]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 11 → Example 12 → Example 13 → Example 14 → Example 15 → Example 16, using the compound produced in Example 37.


TLC: Rf 0.26 (dichloromethane : methanol = 19:1);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.62-1.93, 2.27, 4.08, 4.85, 6.42, 7.23, 7.52-7.62, 8.21, 8.46, 8.51, 8.80, 12.37.

Example 61:

N-{5-[(7-hydroxy-6-methoxy-4-quinolinyl]oxy}-2-pyridinyl-2-oxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0374]**



**[0375]** The compound produced in Example 60 (300 mg) was dissolved in dichloromethane (5 mL). Triethylsilane (127 mg), and trifluoroacetic acid (0.081 mL) were added to the solution, sequentially. The solution was stirred at room temperature for 22 hours. The reaction solution was diluted with ethyl acetate, and washed with water and a saturated sodium hydrogen carbonate aqueous solution. The organic layer was dried over anhydrous sodium sulfate, and then concentrated. The resulting residue was purified by silica gel chromatography (hexane : ethyl acetate = 30:70 → 0:100) to obtain the title compound (183 mg) having the following physical property values.

TLC: R<sub>f</sub> 0.77 (ethyl acetate :methanol = 5:1);

<sup>1</sup> H-NMR (CDCl<sub>3</sub>): δ 1.75, 2.23, 2.72, 4.08, 6.42, 7.22, 7.51-7.60, 8.21, 8.44-8.51, 8.49, 12.51.

Example 62:

4-methyl-7,8-dihydro-2H-chromene-2,5(6H)-dione

**[0376]** Ethyl acetoacetate (CAS registration No.: 141-97-9) (17.40 g), 1,3-cyclohexanedione (CAS registration No.:504-02-9) (10.00 g), DMAP (0.22 g), and pyridine (30 mL) were placed in a 200-mL eggplant flask, and stirred at a bath temperature (140°C) for one day. After the mixture was left to cool to room temperature, the solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 4:1) to obtain the title compound (4.50 g) having the following physical property values.

(LC-MS/ELSD): (retention time: 0.81 min);

<sup>1</sup> H-NMR (CDCl<sub>3</sub>): δ 2.11, 2.48, 2.57, 2.87, 5.99.

Example 63:

3-bromo-4-methyl-7,8-dihydro-2H-chromene-2,5(6H)-dione

**[0377]** N-bromosuccinimide (CAS registration No.: 128-08-5) (2.00 g) was added into a DMF (40 mL) solution of the compound produced in Example 62 in a 100-mL eggplant flask, and the solution was stirred at room temperature for one day. The solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 9:1) to obtain the title compound (1.16 g) having the following physical property values.

(LC-MS/ELSD): (retention time: 0.76 min);

<sup>1</sup> H-NMR (CDCl<sub>3</sub>): δ 2.12, 2.59, 2.69, 2.87.

Example 64:

3-bromo-4-methyl-1-phenyl-7,8-dihydro-2,5(1H,6H)-quinolinedione

**[0378]** The compound produced in Example 63 (1.15 g) and aniline (1.25 g) were placed in a 30-mL eggplant flask. The mixture was stirred at a bath temperature (70°C) for 20 hours. 1 mol/L hydrochloric acid was added to the reaction solution. The reaction solution was extracted with ethyl acetate, washed with water and a saturated saline solution sequentially in this order, and dried over anhydrous sodium sulfate. The solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 7:3) to obtain the title compound (1.31 g) having the following physical property values.

(LC-MS/ELSD): (retention time: 1.12 min);

<sup>1</sup> H-NMR (CDCl<sub>3</sub>): δ 1.96, 2.43, 2.54, 2.79, 7.16-7.20, 7.49-7.58.

Example 65:

methyl 4-methyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinoline carboxylate

**[0379]** The compound produced in Example 64 (1.11 g), potassium acetate (0.66 g), DMF (11 mL), and methanol (11 mL), and, after degassing, [1,1'-bis(diphenylphosphino)ferrocene]palladium(II)dichloride dichloromethane complex (1:1) [PdCl<sub>2</sub> (dppf)<sub>2</sub> CH<sub>2</sub>Cl<sub>2</sub>] (0.27 g) were placed in a 200-mL eggplant flask. After replacement with carbon monoxide was carried out, stirring at a bath temperature (70°C) was carried out for 17 hours. 1 mol/L hydrochloric acid was added to the reaction solution. The reaction solution was extracted with ethyl acetate, washed with water and a saturated saline solution in this order, and dried over anhydrous sodium sulfate. The solvent was distilled off under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane : ethyl acetate = 7:3) to obtain the title compound (0.51 g) having the following physical property values.



(LC-MS/ELSD): (retention time: 0.75 min);

<sup>1</sup> H-NMR (CDCl<sub>3</sub>): δ 1.97, 2.48, 2.53, 2.56, 3.89, 7.17-7.20, 7.48-7.56.

Example 66:

4-methyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydroquinoline-3-carboxylic acid

**[0380]** The compound produced in Example 65 (1.85 g) and 5 mol/L hydrochloric acid (19 mL) were placed in a 50-mL eggplant flask. The mixture was stirred at a bath temperature (50°C) for 26 hours. Precipitate was removed by filtration, and the filtrate solvent was distilled off under reduced pressure. The resulting residue was washed in a slurry form with methanol and ethyl acetate to obtain the title compound (0.89 g) having the following physical property values.


(LC-MS/ELSD): (retention time: 0.66 min);

<sup>1</sup> H-NMR (CD<sub>3</sub> OD): δ 1.97, 2.54, 2.63, 7.29-7.32, 7.54-7.63.

Example 67:

N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-4-methyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0381]**



**[0382]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 5, using the compound produced in Example 66 and the compound produced in Example 2.

TLC: R<sub>f</sub> 0.62 (dichloromethane :methanol = 9:1);

<sup>1</sup> H-NMR (CDCl<sub>3</sub>): δ 2.01, 2.53, 2.58, 2.86, 4.05, 6.42, 7.22, 7.42, 7.49-7.61, 8.20, 8.45, 8.50, 9.73.

Example 67(1):

N-{4-[(6,7-dimethoxy-4-quinolinyl)oxy]phenyl}-4-methyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide

**[0383]** The title compound having the following physical property values was obtained by the procedure having the same purpose as in Example 67, using the compound produced in Example 66 and the compound produced in Example 22.

TLC: R<sub>f</sub> 0.48 (ethyl acetate :methanol = 19:1);

<sup>1</sup> H-NMR (CDCl<sub>3</sub>): δ 2.01, 2.52, 2.59, 2.93, 4.04, 6.45, 7.14, 7.23, 7.41, 7.55-7.68, 7.72, 8.47, 9.79.

[Experiment Example]

**[0384]** Biological Examples are described below. Based on these experiment methods, the effect of the compound of the present invention was verified.

Biological Example 1: Measurement of an Axl inhibitory activity (in vitro test)

**[0385]** An Axl enzyme inhibitory activity was measured by using LanthaScreen (registered trademark) system (Invitrogen) based on the attached instruction. The reagents used are shown below.

### EP 3 415 501 B9

Reaction buffer solution: a solution containing 50 mmol/L HEPES (pH7.5), 0.01% Brij35, 10 mmol/L MgCl<sub>2</sub> and 1 mmol/L EGTA was prepared by using purified water. Test substance solution: a solution containing a test compound of 5-fold concentration with respect to the final concentration was prepared by 20-fold diluting a DMSO solution of test compound of each concentration with the reaction buffer solution.

Enzyme solution: a solution containing 400 ng/mL Axl enzyme was prepared by using the reaction buffer solution. Substrate solution: a solution containing 45  $\mu$ mol/L ATP and 500 nmmol/L Fluorescein-Poly GT (Invitrogen) was prepared by using the reaction buffer solution.

Detection solution: a solution containing 20 mM EDTA and 4 nM PY20 (Invitrogen) was prepared by using Dilution B (Invitrogen).

**[0386]** A 10 mmol/L DMSO solution of the test compound was dispensed into a 96-well plate (Nunc), and, furthermore, a 3-fold dilution series was prepared using DMSO. In each well of the 96-well plate for measurement, 5  $\mu$ L each of the reaction buffer solution containing DMSO was added to a Blank group and a medium group, and 5  $\mu$ L of the test substance solution was added to the test substance group, respectively. Next, 10  $\mu$ L/well of the reaction buffer solution was added to the Blank group, and 10  $\mu$ L/well each of the enzyme solution was added to the medium group and the test compound group, followed by stirring at room temperature for 10 min. After the completion of stirring, 10  $\mu$ L each of the substrate solution was added into each well, followed by stirring at room temperature with light shielded for one hour. After the completion of reaction, 25  $\mu$ L each of the detection solution was added to each well, and stood still at room temperature with light shielded for 30 min. After standing still, fluorescence intensity at 520 nm and 495 nm at the time of irradiation with exciting light of 340 nm was measured by using Analyst GT (Molecular Devices). The phosphorylation of the artificial substrate was quantified by Time-resolved Fluorescence Resonance Energy Transfer (TR-FRET). TR-FRET ratio was calculated by dividing 520 nm fluorescence signal by 495 nm fluorescence signal for each well, and the inhibition rate (%) in the test compound group was calculated based on the following mathematical formula.

$$\text{Inhibition rate (\%)} = [1 - (\text{TR-FRET ratio of test compound group} - A) / (B - A)] \times 100$$


A: average value of TR-FRET ratios of Blank group

B: average value of TR-FRET ratios of medium group


**[0387]** Values of 50% inhibition rate (IC<sub>50</sub> values) of the test compound were calculated from the inhibition curve based on the inhibition rate of the test compounds in each concentration.

**[0388]** As a result, in the compounds of the present invention, IC<sub>50</sub> values of the compound of, for example, Examples 5, 5(1), 5(6), 17(2), and 23(2) were 0.0022  $\mu$ M, 0.0056  $\mu$ M, 0.0043  $\mu$ M, 0.0044  $\mu$ M, and 0.0011  $\mu$ M, respectively.

**[0389]** On the other hand, as comparative compounds, the Axl inhibitory activity of each of the compound of Example 8 described in Patent Literature 1 (Comparative compound A) and the compound 2 of Example 3 described in Patent Literature 3 (Comparative compound B), having the following structures, was measured. In both cases, IC<sub>50</sub> value was higher than 10  $\mu$ M.



**Comparative compound A**



Comparative compound B

Biological Example 2: Measurement of proliferation suppression rate by using mouse pro-B cell line (Ba/F3 Axl) stably expressing Axl

**[0390]** A 0.1 mmol/L DMSO solution of the test compound was dispensed into a 96-well plate, and a 3-fold dilution series was prepared using DMSO. DMSO solutions of test compounds, having various concentrations, were further 500-fold diluted with a RPMI1640 medium (containing 10% HI-FBS, 1% penicillin) and a diluted solution of the test compound having 500-fold concentration with respect to the final concentration was prepared. In each well of the 96-well plate (BD Biosciences) for measurement, 50  $\mu$ L of a RPMI medium was added to the Blank group, 50  $\mu$ L of a RPMI medium containing 0.2% DMSO was added to the medium group, and 50  $\mu$ L of the diluted solution of the test compound was added to the test compound group, respectively. Ba/F3 Axl was diluted with a medium to have a density of  $2 \times 10^5$  cells/mL to prepare a cell suspension. In each well of the 96-well plate for measurement, 50  $\mu$ L each of the RPMI medium was added to the Blank group, and 50  $\mu$ L each of the cell suspension to the medium group and the test compound group, respectively, and the groups were stood still at 37°C at 5%CO<sub>2</sub> for 48 hours. After standing still, Relative Light Unit (RLU) was measured by using CELLTITER-GLO (registered trademark) LUMINESCENT CELL VIABILITY ASSAY (Promega). The measurement was carried out according to the attached instruction. To each well, 100  $\mu$ L each of light-emitting solution was added. The plate was stirred at room temperature for 3 min and then stood still at room temperature with light shielded for 10 min, and RLU was measured by using Microplate Reader (SpectraMax M5e, Molecular Devices). The average values of RLU of the Blank group and the medium group were respectively calculated, and the proliferation suppression rate of the test compound group was calculated.

Proliferation suppression rate (%) =

$$\{1 - (\text{RLU of test compound group} - A) / (B - A)\} \times 100$$

A: average value of RLU of Blank group

B: average value of RLU of medium group


**[0391]** A value of 50% inhibition rate (IC<sub>50</sub> value) of the test compound was calculated from the inhibition curve based on the inhibition rate in each concentration of the test compound.

**[0392]** As a result, in the compounds of the present invention, IC<sub>50</sub> values of the compounds of, for example, Examples 5, 5(1), 5(6), 17(2), and 23(2) were 0.0007  $\mu$ M, 0.0008  $\mu$ M, 0.0078  $\mu$ M, 0.0012  $\mu$ M, and 0.0012  $\mu$ M, respectively.


**[0393]** On the other hand, IC<sub>50</sub> values of the comparative compounds A and B, were 0.62  $\mu$ M and > 10  $\mu$ M, respectively.

Biological Example 3: Evaluation of kinase selectivity (in vitro test)

**[0394]** Similar to Biological Example 1, values of 50% inhibition rate (IC<sub>50</sub> value) with respect to various kinases (KDR, DDR1, FLT4, and ROS) of the test compound were measured. The Axl selective inhibitory activity of the test compound with respect to kinases, for example, KDR, was calculated based on the above-mentioned ratio of the IC<sub>50</sub> values. The calculated values are shown in the following Table 1. As the test compound, for the compound of the present invention, the compounds of Examples 5, 5(1), 17(2), and 23(2) were used, and for the comparative compounds, the compound of Example 5 (Comparative compound C) and the compound of Example 92 (Comparative compound D) described in Patent Literature 5, having the following structure, were used.



Comparative compound C



Comparative compound D

[Table 1]

	KDR[IC50]/Axl[IC50]
Example 5	about 900 times
Comparative compound C	about 0.2 times
Comparative compound D	about 28 times

[0395] Results showed that the compound of the present invention had Axl selective inhibition effect on KDR as compared with the comparative compounds. KDR is kinase also referred to as vascular endothelial growth factor receptor 2 (VEGF Receptor 2). It is known that inhibition of KDR may cause a side effect of increasing blood pressure (Hypertension, vol. 39, p. 1095-1100, 2002). Therefore, it was suggested that the compounds of the present invention were excellent compounds capable of avoiding the side effect, which was a problem in comparative compounds, for example, hypertension. Furthermore, it has been also suggested that the other three types of kinases (DDR1, FLT4, and ROS) might cause side effect to be avoided, from the phenotype of the KO mouse or transgenic mouse. It became apparent that the compound of the present invention has excellent selectivity to such specific kinases and therefore capable of avoiding side effect.

Biological Example 4: Measurement of inhibitory activity of drug-metabolizing enzyme (human CYP2C8 inhibition effect)

[0396] The reaction was carried out in a 384-well plate. As the positive control substance (CYP2C8: quercetin), a solution, which had been adjusted with DMSO to have 300 times higher concentration than the final concentration (CYP2C8: 22.5 and 225  $\mu\text{mol/L}$ ) and been 75-fold diluted with purified water containing 2.7% acetonitrile, was prepared

(CYP2C8: 0.3 and 3  $\mu\text{mol/L}$ ). The test compounds were prepared to have a concentration of 0.3 and 3  $\text{mol/L}$  with DMSO, and then 75-fold diluted with purified water containing 2.7% acetonitrile to be 4 and 40  $\mu\text{mol/L}$ . Then, a reaction mixture solution was prepared by addition of a potassium phosphate buffer (pH 7.4), magnesium chloride (5  $\text{mol/L}$ ), substrate (CYP2C8: Luciferin-ME, 150  $\mu\text{mol/L}$ ), and E. coli-expressed liver microsome CYP2C8 (Cypex, 30  $\text{pmol/L}$ ) (the numerical values are final concentrations). The reaction was started by addition of 8  $\mu\text{L}$  of this reaction mixture, 4  $\mu\text{L}$  each of the test compound and the positive control solution which had been prepared as described above, and 4  $\mu\text{L}$  of NADPH production system solution (5.2 mM NADP, 13.2 mM glucose-6-phosphate, 1.6 U/mL glucose-6-phosphate dehydrogenase) and incubation was carried out at 37°C for 30 min. Thereafter, 16  $\mu\text{L}$  of luciferase solution was added to stop the reaction and to allow luciferin to emit light, and the luminescence intensity of the reaction solution was measured. The inhibition rate is a reduction rate (inhibition rate) of the luminescence intensity when compared with the control in which the reaction was carried out by the addition of DMSO in place of the test compound solution. The inhibition rate was calculated from the following mathematical formula.

$$\text{Inhibition rate (\%)} = 100 - \{(\text{luminescence intensity of test compound} - \text{background luminescence intensity}) / (\text{luminescence intensity of control} - \text{background luminescence intensity}) \times 100\}$$

**[0397]** The IC<sub>50</sub> value was defined to be < 1  $\mu\text{M}$  when the inhibition rate at 1  $\mu\text{mol/L}$  was not less than 50%; and > 10  $\mu\text{M}$  when the inhibition rate at 10  $\mu\text{mol/L}$  was not more than 50%. The range between the above-mentioned range (not more than 50% at 1  $\mu\text{mol/L}$  and not less than 50% at 10  $\mu\text{mol/L}$ ) was calculated using the following mathematical formula:

$$\text{IC}_{50} = (50 - b) / a$$

wherein a and b are the slope and intercept of the linear regression line:  $y = ax + b$  that passes through the two points: the concentration and the inhibition rate at 1  $\mu\text{mol/L}$  and the concentration and the inhibition rate at 10  $\mu\text{mol/L}$ .

**[0398]** The IC<sub>50</sub> values of the comparative compounds and compounds of the present invention were measured using the measurement method described above.

**[0399]** As a result, the IC<sub>50</sub> value of CYP2C8 was 2.6  $\mu\text{M}$  for the comparative compound E (Example 133 described in Patent Literature 4). On the other hand, for the compound of the present invention, the IC<sub>50</sub> values of CYP2C8 were > 10  $\mu\text{M}$  in the compounds of, for example, Examples 5, 5(1), 17(2), and 23(2). Therefore, it was shown that the compound of the present invention had less CYP inhibition effect with respect to the comparative compound.

[Formulation example]

Formulation example 1 (Reference)

**[0400]** The components indicated below were mixed by a standard method, followed by making the mixture into tablets to obtain 10,000 tablets each containing 10 mg of active ingredient.

- N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide ... 100 g
- calcium carboxymethyl cellulose (disintegrant) ... 20 g
- magnesium stearate (lubricant) ... 10 g
- microcrystalline cellulose ... 870 g

Formulation example 2

**[0401]** The components indicated below were mixed by a standard method, filtered through a dust-removing filter, filled into ampoules so that each ampule contains 5 ml, and thermally sterilized in an autoclave to obtain 10,000 ampoules each containing 20 mg active ingredient.

- N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-7,7-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide ... 200g
- mannitol ... 20g


- distilled water ... 50L

[Industrial Applicability]

**[0402]** A compound of the present invention has a strong Axl inhibitory activity, and therefore, is useful for treatment for Axl-related diseases, for example, cancer, kidney diseases, immune system diseases, and circulatory system diseases.

## Claims

1. A compound represented by general formula (1):



wherein  $R^1$  represents (1) a C1-8 alkyl group optionally substituted with one to five  $R^{11}$ , (2) a C3-7 carbon ring optionally substituted with one to five  $R^{12}$ , or (3) a 4- to 7-membered heterocycle optionally substituted with one to five  $R^{13}$ , wherein when the C1-8 alkyl group represented by  $R^1$  is a branched alkyl group, the C1-3 alkyl group branched from the same carbon atom, together with the carbon atom bound thereto, optionally forms a saturated C3-7 carbon ring,

$R^2$  represents (1) a C1-4 alkyl group, (2) a halogen atom, (3) a C1-4 haloalkyl group, (4) an oxo group, (5) an  $-OR^{21}$  group, or (6) an  $=NR^{22}$  group,

$R^3$  represents (1) a C1-4 alkyl group, (2) a halogen atom, or (3) a C1-4 haloalkyl group,

$R^4$  represents (1) a C1-4 alkoxy group, (2) a C1-4 haloalkyl group, (3) an  $-OR^{41}$  group, (4) a C1-4 alkyl group, (5) a C2-4 alkenyloxy group, or (6) a C2-4 alkynyloxy group,

$R^5$  represents (1) a hydrogen atom, (2) a C1-4 alkyl group, (3) a halogen atom, (4) a C1-4 haloalkyl group, or (5) an  $-OR^{21}$  group,

$R^{11}$  represents (1) an  $-OR^{101}$  group, (2) an  $SO_2R^{102}$  group, (3) an  $NR^{103}R^{104}$  group, or (4) a C3-7 carbon ring optionally substituted with one to three halogen atoms,

$R^{12}$  represents (1) a C1-8 alkyl group optionally substituted with a hydroxyl group, or (2) a halogen atom,

$R^{13}$  represents (1) a C1-8 alkyl group optionally substituted with a hydroxyl group, or (2) a halogen atom,

$R^{21}$  represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,

$R^{22}$  represents (1) a hydroxyl group, or (2) C1-4 alkoxy group,


$R^{41}$  represents (1) a hydrogen atom; (2) a C1-8 alkyl group substituted with one to two substituents selected from the group consisting of (a) 5- to 7-membered cyclic group optionally substituted with one to two substituents selected from the group consisting of (i) a C1-4 alkyl group, (ii) a C1-4 haloalkyl group, and (iii) a halogen atom, (b)  $NR^{401}R^{402}$ , (c) a hydroxyl group, and (d) an  $SO_2R^{403}$  group; (3) a C2-8 alkenyl group substituted with one to two substituents selected from the group consisting of (a) 5- to 7-membered cyclic group optionally substituted with one to two substituents selected from the group consisting of (i) a C1-4 alkyl group, (ii) a C1-4 haloalkyl group, and (iii) a halogen atom, (b)  $NR^{401}R^{402}$ , (c) a hydroxyl group, and (d) an  $SO_2R^{403}$  group; or (4) a C2-8 alkynyl group substituted with one to two substituents selected from the group consisting of (a) 5- to 7-membered cyclic group optionally substituted with one to two substituents selected from the group consisting of (i) a C1-4 alkyl group, (ii) a C1-4 haloalkyl group, and (iii) a halogen atom, (b)  $NR^{401}R^{402}$ , (c) a hydroxyl group, and (d) an  $SO_2R^{403}$  group,

$R^{101}$  represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,  
 $R^{102}$  represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,  
 $R^{103}$  and  $R^{104}$  each independently represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,  
 $R^{401}$  and  $R^{402}$  each independently represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,  
 $R^{403}$  represents (1) a hydrogen atom, or (2) a C1-4 alkyl group,  
A represents (1) CH, or (2) a nitrogen atom,  
L represents (1) -O-, (2) -NH-, (3) -C(O)-, (4) -CR<sup>6</sup>R<sup>7</sup>-, (5) -S-, (6) -S(O)-, or (7) -S(O)<sub>2</sub>-,  
 $R^6$  and  $R^7$  each independently represents (1) a hydrogen atom, (2) a halogen atom, (3) a C1-4 alkyl group, (4)  
a hydroxyl group, or (5) NH<sub>2</sub>,  
ring1 represents benzene or pyridine,  
----- represents a single bond or a double bond,  
m is an integer from 0 to 5,  
n is an integer from 0 to 5,  
p is an integer from 0 to 2,  
q is an integer from 0 to 4,  
when m is two or more, a plurality of R<sup>2</sup> may be the same as or different from each other, and when two of R<sup>2</sup>  
represent a C1-3 alkyl group and are on the same carbon atom, the R<sup>2</sup>, together with a carbon atom bound  
thereto, may form a saturated C3-7 carbon ring,  
when n is 2 or more, a plurality of R<sup>3</sup> may be the same as or different from each other, and when q is 2 or more,  
a plurality of R<sup>4</sup> may be the same as or different from each other,  
a salt thereof, a solvate thereof, or an N-oxide thereof.

2. The compound according to claim 1, wherein m is one or more, and one of R<sup>2</sup> is necessarily an oxo group.

3. The compound according to claim 1 or 2, wherein L is (1) -O-, (2) -NH-, or (3) -C(O)-.

4. The compound according to claim 1, which is represented by general formula (I-1):



wherein R<sup>2-1</sup> represents (1) a C1-4 alkyl group, (2) a halogen atom, (3) a C1-4 haloalkyl group, (4) an -OR<sup>21</sup>  
 group, or (5) an =NR<sup>22</sup> group,  
 m-1 is an integer from 0 to 4,  
 L<sup>1</sup> is (1) -O-, (2) -NH-, or (3) -C(O)-,  
 ring1-1 represents benzene or pyridine,  
 when m-1 is 2 or more, a plurality of R<sup>2-1</sup> may be the same as or different from each other, and when two of  
 R<sup>2-1</sup> represent a C1-3 alkyl group and are on the same carbon atom, the R<sup>2-1</sup>, together with a carbon atom  
 bound thereto, may form a saturated C3-7 carbon ring,  
 and the other symbols have the same meanings as defined in claim 1.

5. The compound according to any one of claims 1 to 4, which is:

(2) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-7,7-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-

3-quinolinecarboxamide,

(3) N-{5-[(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(4) N-[5-{7-[3-(4-morpholinyl)propoxy]-4-quinolinyl}oxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(5) N-[4-{(6,7-dimethoxy-4-quinolinyl)oxy}-3-fluorophenyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(6) N-[4-{(6,7-dimethoxy-4-quinolinyl)oxy}phenyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(7) N-[5-{(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(4-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(8) N-[5-{(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(9) N-[5-{(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-(2-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(10) N-[5-{(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(11) N-[5-{(6,7-dimethoxy-4-quinazolinyl)oxy]-2-pyridinyl}-1-(4-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(12) N-[5-{(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-1-[(2S)-1-hydroxy-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(13) N-[4-{(6,7-dimethoxy-4-quinolinyl)oxy]-3-fluorophenyl}-1-(3-fluorophenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(14) N-[5-{(6,7-dimethoxy-4-quinolinyl)oxy]-2-pyridinyl}-6,6-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(15) N-[5-{(6-methoxy-7-[3-(4-morpholinyl)propoxy]-4-quinolinyl)oxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(16) N-[5-{[7-(3-hydroxy-3-methylbutoxy)-6-methoxy-4-quinolinyl]oxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, or

(17) N-[5-{(6-methoxy-7-[3-(1-pyrrolidinyl)propoxy]-4-quinolinyl)oxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide.

6. A pharmaceutical composition containing a compound represented by general formula (I) as defined in claim 1, a salt thereof, a solvate thereof, or an N-oxide thereof.

7. The pharmaceutical composition according to claim 6, which is an Axl inhibitor.

8. The pharmaceutical composition according to claim 6, which is an agent for preventing and/or treating an Axl-related disease.

9. The pharmaceutical composition according to claim 8, wherein the Axl-related disease includes a cancer, a kidney disease, an immune system disease, or a circulatory system disease.

10. The pharmaceutical composition according to claim 9, wherein the cancer is acute myeloid leukemia, chronic myeloid leukemia, acute lymphatic leukemia, melanoma, breast cancer, pancreatic cancer, glioma, esophageal adenocarcinoma, large intestine cancer, renal cell carcinoma, thyroid cancer, non-small cell lung cancer, prostate cancer, stomach cancer, liver cancer, uveal malignant melanoma, ovarian cancer, endometrial cancer, lymphoma, head and neck cancer, or sarcoma.

11. The pharmaceutical composition according to claim 6, which is a metastasis suppressing agent for cancer cells.

12. A compound represented by general formula (I) according to claim 1, a salt thereof, a solvate thereof, or an N-oxide thereof for use in preventing and/or treating an Axl-related disease.

13. The compound for use according to claim 12, wherein the Axl-related disease includes a cancer, a kidney disease, an immune system disease, or a circulatory system disease.


14. The compound for use according to claim 13, wherein the cancer is acute myeloid leukemia, chronic myeloid



leukemia, acute lymphatic leukemia, melanoma, breast cancer, pancreatic cancer, glioma, esophageal adenocarcinoma, large intestine cancer, renal cell carcinoma, thyroid cancer, non-small cell lung cancer, prostate cancer, stomach cancer, liver cancer, uveal malignant melanoma, ovarian cancer, endometrial cancer, lymphoma, head and neck cancer, or sarcoma.

## Patentansprüche

1. Verbindung, dargestellt durch die allgemeine Formel (1):



wobei  $R^1$  (1) eine C1-8-Alkylgruppe, die wahlweise mit einem bis fünf  $R^{11}$  substituiert ist, (2) einen C3-7-Kohlenstoffring, der wahlweise mit einem bis fünf  $R^{12}$  substituiert ist, oder (3) einen 4- bis 7-gliedrigen Heterocyclus, der wahlweise mit einem bis fünf  $R^{13}$  substituiert ist, darstellt, wobei, wenn die durch  $R^1$  dargestellte C1-8-Alkylgruppe eine verzweigte Alkylgruppe ist, die von demselben Kohlenstoffatom verzweigte C1-3-Alkylgruppe zusammen mit dem daran gebundenen Kohlenstoffatom wahlweise einen gesättigten C3-7-Kohlenstoffring bildet,

$R^2$  (1) eine C1-4-Alkylgruppe, (2) ein Halogenatom, (3) eine C1-4-Halogenalkylgruppe, (4) eine Oxogruppe, (5) eine  $-OR^{21}$ -Gruppe oder (6) eine  $=NR^{22}$ -Gruppe darstellt,

$R^3$  (1) eine C1-4-Alkylgruppe, (2) ein Halogenatom oder (3) eine C1-4-Halogenalkylgruppe darstellt,

$R^4$  (1) eine C1-4-Alkoxygruppe, (2) eine C1-4-Haloalkylgruppe, (3) eine  $-OR^{41}$ -Gruppe, (4) eine C1-4-Alkylgruppe, (5) eine C2-4-Alkenyloxygruppe oder (6) eine C2-4-Alkinyloxygruppe darstellt,

$R^5$  (1) ein Wasserstoffatom, (2) eine C1-4-Alkylgruppe, (3) ein Halogenatom, (4) eine C1-4-Halogenalkylgruppe oder (5) eine  $-OR^{21}$ -Gruppe darstellt,

$R^{11}$  (1) eine  $-OR^{101}$ -Gruppe, (2) eine  $SO_2R^{102}$ -Gruppe, (3) eine  $NR^{103}R^{104}$ -Gruppe oder (4) einen C3-7-Kohlenstoffring, der wahlweise mit einem bis drei Halogenatomen substituiert ist, darstellt,

$R^{12}$  (1) eine C1-8-Alkylgruppe, die wahlweise mit einer Hydroxylgruppe substituiert ist, oder (2) ein Halogenatom darstellt,

$R^{13}$  (1) eine C1-8-Alkylgruppe, die wahlweise mit einer Hydroxylgruppe substituiert ist, oder (2) ein Halogenatom darstellt,


$R^{21}$  (1) ein Wasserstoffatom oder (2) eine C1-4-Alkylgruppe darstellt,

$R^{22}$  (1) eine Hydroxylgruppe oder (2) eine C1-4-Alkoxygruppe darstellt,

$R^{41}$  darstellt: (1) ein Wasserstoffatom, (2) eine C1-8-Alkylgruppe, die mit einem bis zwei Substituenten substituiert ist, die aus der Gruppe ausgewählt sind, die aus (a) einer 5- bis 7-gliedrigen cyclischen Gruppe, die wahlweise mit einem bis zwei Substituenten substituiert ist, die aus der Gruppe ausgewählt sind, die aus (i) einer C1-4-Alkylgruppe, (ii) einer C1-4-Halogenalkylgruppe und (iii) einem Halogenatom besteht, (b)  $NR^{401}R^{402}$ , (c) einer Hydroxylgruppe und (d) einer  $SO_2R^{403}$ -Gruppe besteht, (3) eine C2-8-Alkenylgruppe, die mit ein bis zwei Substituenten substituiert ist, die aus der Gruppe ausgewählt sind, die aus (a) einer 5- bis 7-gliedrigen cyclischen Gruppe, die wahlweise mit ein bis zwei Substituenten substituiert ist, die aus der Gruppe ausgewählt sind, die aus (i) einer C1-4-Alkylgruppe, (ii) einer C1-4-Halogenalkylgruppe und (iii) einem Halogenatom besteht, (b)  $NR^{401}R^{402}$ , (c) einer Hydroxylgruppe und (d) einer  $SO_2R^{403}$ -Gruppe besteht, oder (4) eine C2-8-Alkyl-

gruppe, die mit einem bis zwei Substituenten substituiert ist, die aus der Gruppe ausgewählt sind, die aus (a) einer 5- bis 7-gliedrigen cyclischen Gruppe, die wahlweise mit einem bis zwei Substituenten substituiert ist, die aus der Gruppe ausgewählt sind, die aus (i) einer C1-4-Alkylgruppe, (ii) einer C1-4-Halogenalkylgruppe und (iii) einem Halogenatom besteht, (b)  $\text{NR}^{401}\text{R}^{402}$ , (c) einer Hydroxylgruppe und (d) einer  $\text{SO}_2\text{R}^{403}$ -Gruppe besteht,  $\text{R}^{101}$  (1) ein Wasserstoffatom oder (2) eine C1-4-Alkylgruppe darstellt,  $\text{R}^{102}$  (1) ein Wasserstoffatom oder (2) eine C1-4-Alkylgruppe darstellt,  $\text{R}^{103}$  und  $\text{R}^{104}$  jeweils unabhängig voneinander (1) ein Wasserstoffatom oder (2) eine C1-4-Alkylgruppe darstellen,  $\text{R}^{401}$  und  $\text{R}^{402}$  jeweils unabhängig voneinander (1) ein Wasserstoffatom oder (2) eine C1-4-Alkylgruppe darstellen,  $\text{R}^{403}$  (1) ein Wasserstoffatom oder (2) eine C1-4-Alkylgruppe darstellt, A (1) CH oder (2) ein Stickstoffatom darstellt, L (1) -O-, (2) -NH-, (3) -C(O)-, (4) - $\text{CR}^6\text{R}^7$ -, (5) -S-, (6) -S(O)- oder (7) -S(O)<sub>2</sub>- darstellt,  $\text{R}^6$  und  $\text{R}^7$  jeweils unabhängig voneinander (1) ein Wasserstoffatom, (2) ein Halogenatom, (3) eine C1-4-Alkylgruppe, (4) eine Hydroxylgruppe oder (5)  $\text{NH}_2$  darstellen, Ring1 Benzol oder Pyridin darstellt, ----- eine Einfachbindung oder eine Doppelbindung darstellt, m eine ganze Zahl von 0 bis 5 ist, n eine ganze Zahl von 0 bis 5 ist, p eine ganze Zahl von 0 bis 5 ist, q eine ganze Zahl von 0 bis 5 ist, wenn m zwei oder mehr ist, eine Vielzahl von  $\text{R}^2$  gleich oder verschieden voneinander sein kann, und wenn zwei von  $\text{R}^2$  eine C1-3-Alkylgruppe darstellen und sich am selben Kohlenstoffatom befinden, die  $\text{R}^2$  zusammen mit einem daran gebundenen Kohlenstoffatom einen gesättigten C3-7-Kohlenstoffring bilden können, wenn n 2 oder mehr ist, eine Vielzahl von  $\text{R}^3$  gleich oder verschieden voneinander sein kann und wenn q 2 oder mehr ist, eine Vielzahl von  $\text{R}^4$  gleich oder verschieden voneinander sein kann, ein Salz davon, ein Solvat davon oder ein N-Oxid davon.

2. Verbindung gemäß Anspruch 1, wobei m eins oder mehr ist und einer von  $\text{R}^2$  unbedingt eine Oxogruppe ist.
3. Verbindung gemäß Anspruch 1 oder 2, wobei L (1) -O-, (2) -NH- oder (3) -C(O)- ist.
4. Verbindung gemäß Anspruch 1, die durch die allgemeine Formel (I-1) dargestellt wird:



wobei  $\text{R}^{2-1}$  (1) eine C1-4-Alkylgruppe, (2) ein Halogenatom, (3) eine C1-4-Halogenalkylgruppe, (4) eine - $\text{OR}^{21}$ -Gruppe oder (5) eine = $\text{NR}^{22}$ -Gruppe darstellt, m-1 eine ganze Zahl von 0 bis 4 ist,  $\text{L}^1$  (1) -O-, (2) -NH- oder (3) -C(O)- ist, Ring1-1 Benzol oder Pyridin darstellt, wenn m-1 2 oder mehr ist, eine Vielzahl von  $\text{R}^{2-1}$  gleich oder verschieden voneinander sein kann, und wenn zwei von  $\text{R}^{2-1}$  eine C1-3-Alkylgruppe darstellen und sich am selben Kohlenstoffatom befinden, die  $\text{R}^{2-1}$  zusammen

men mit einem daran gebundenen Kohlenstoffatom einen gesättigten C3-7-Kohlenstoffring bilden können, und die anderen Symbole die gleichen Bedeutungen wie in Anspruch 1 definiert haben.

5. Verbindung gemäß einem der Ansprüche 1 bis 4, welche

- (2) N-{5-[(6,7-Dimethoxy-4-chinolinyloxy)-2-pyridinyl]-7,7-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (3) N-{5-[(6,7-Dimethoxy-4-chinolinyloxy)-2-pyridinyl]-1-(2,2-dimethylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (4) N-{5-[(3-(4-Morpholinyloxy)-4-chinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (5) N-{4-[(6,7-Dimethoxy-4-chinolinyloxy)-3-fluorphenyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (6) N-{4-[(6,7-Dimethoxy-4-chinolinyloxy)phenyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (7) N-{5-[(6,7-Dimethoxy-4-chinolinyloxy)-2-pyridinyl]-1-(4-fluorphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (8) N-{5-[(6,7-Dimethoxy-4-chinolinyloxy)-2-pyridinyl]-1-(3-fluorphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (9) N-{5-[(6,7-Dimethoxy-4-chinolinyloxy)-2-pyridinyl]-1-(2-fluorphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (10) N-{5-[(6,7-Dimethoxy-4-chinazolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (11) N-{5-[(6,7-Dimethoxy-4-chinazolinyloxy)-2-pyridinyl]-1-(4-fluorphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (12) N-{5-[(6,7-Dimethoxy-4-chinolinyloxy)-2-pyridinyl]-1-[(2S)-1-hydroxy-3-methyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (13) N-{4-[(6,7-Dimethoxy-4-chinolinyloxy)-3-fluorphenyl]-1-(3-fluorphenyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (14) N-{5-[(6,7-Dimethoxy-4-chinolinyloxy)-2-pyridinyl]-6,6-dimethyl-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (15) N-{5-[(6-Methoxy-7-[3-(4-morpholinyloxy)-4-chinolinyloxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid,
- (16) N-{5-[(7-(3-Hydroxy-3-methylbutoxy)-6-methoxy-4-chinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid oder
- (17) N-{5-[(6-Methoxy-7-[3-(1-pyrrolidinyloxy)-4-chinolinyloxy]-2-pyridinyl]-2,5-dioxo-1-phenyl-1,2,5,6,7,8-hexahydro-3-chinolincarboxamid ist.

6. Pharmazeutische Zusammensetzung, enthaltend eine Verbindung, welche durch die allgemeine Formel (I) wie in Anspruch 1 definiert dargestellt wird, ein Salz davon, ein Solvat davon oder ein N-Oxid davon.

7. Pharmazeutische Zusammensetzung gemäß Anspruch 6, bei der es sich um einen Axl-Inhibitor handelt.

8. Pharmazeutische Zusammensetzung gemäß Anspruch 6, die ein Mittel zum Verhindern und/oder Behandeln einer mit Axl in Beziehung stehenden Krankheit ist.

9. Pharmazeutische Zusammensetzung gemäß Anspruch 8, wobei die mit Axl in Beziehung stehende Krankheit Krebs, eine Nierenerkrankung, eine Erkrankung des Immunsystems oder eine Erkrankung des Kreislaufsystems einschließt.


10. Pharmazeutische Zusammensetzung gemäß Anspruch 9, wobei der Krebs akute myeloische Leukämie, chronische myeloische Leukämie, akute lymphatische Leukämie, Melanom, Brustkrebs, Bauchspeicheldrüsenkrebs, Gliom, Ösophagus-Adenokarzinom, Dickdarmkrebs, Nierenzellkarzinom, Schilddrüsenkrebs, nicht kleinzelliger Lungenkrebs, Prostatakrebs, Magenkrebs, Leberkrebs, malignes Melanom der mittleren Augenhaut, Eierstockkrebs, Endometriumkrebs, Lymphom, Kopf- und Halskrebs oder Sarkom ist.

11. Pharmazeutische Zusammensetzung gemäß Anspruch 6, die ein Metastasen unterdrückendes Mittel für Krebszellen ist.

12. Verbindung, welche durch die allgemeine Formel (I) gemäß Anspruch 1 dargestellt wird, ein Salz davon, ein Solvat davon oder ein N-Oxid davon zur Verwendung beim Verhindern und/oder Behandeln einer mit Axl in Beziehung stehenden Krankheit.
13. Verbindung zur Verwendung gemäß Anspruch 12, wobei die mit Axl in Beziehung stehende Krankheit Krebs, eine Nierenerkrankung, eine Erkrankung des Immunsystems oder eine Erkrankung des Kreislaufsystems einschließt.
14. Verbindung zur Verwendung gemäß Anspruch 13, wobei der Krebs akute myeloische Leukämie, chronische myeloische Leukämie, akute lymphatische Leukämie, Melanom, Brustkrebs, Bauchspeicheldrüsenkrebs, Gliom, Ösophagus-Adenokarzinom, Dickdarmkrebs, Nierenzellkarzinom, Schilddrüsenkrebs, nicht kleinzelliger Lungenkrebs, Prostatakrebs, Magenkrebs, Leberkrebs, malignes Melanom der mittleren Augenhaut, Eierstockkrebs, Endometriumkrebs, Lymphom, Kopf- und Halskrebs oder Sarkom ist.

## Revendications

1. Composé représenté par la formule générale (I) :



dans laquelle  $R^1$  représente (1) un groupe alkyle en C1 à 8 éventuellement substitué par un à cinq  $R^{11}$ , (2) un cycle carboné en C3 à 7 éventuellement substitué par un à cinq  $R^{12}$ , ou (3) un hétérocycle de 4 à 7 chaînons éventuellement substitué par un à cinq  $R^{13}$ , où lorsque le groupe alkyle en C1 à 8 représenté par  $R^1$  est un groupe alkyle ramifié, le groupe alkyle en C1 à 3 ramifié depuis le même atome de carbone forme éventuellement, avec l'atome de carbone auquel il est lié, un cycle carboné saturé en C3 à 7,

$R^2$  représente (1) un groupe alkyle en C1 à 4, (2) un atome d'halogène, (3) un groupe halogénoalkyle en C1 à 4, (4) un groupe oxo, (5) un groupe  $-OR^{21}$ , ou (6) un groupe  $=NR^{22}$ ,

$R^3$  représente (1) un groupe alkyle en C1 à 4, (2) un atome d'halogène, ou (3) un groupe halogénoalkyle en C1 à 4,

$R^4$  représente (1) un groupe alkoxy en C1 à 4, (2) un groupe halogénoalkyle en C1 à 4, (3) un groupe  $-OR^{41}$ , (4) un groupe alkyle en C1 à 4, (5) un groupe alcényloxy en C2 à 4, ou (6) un groupe alcynyloxy en C2 à 4,

$R^5$  représente (1) un atome d'hydrogène, (2) un groupe alkyle en C1 à 4, (3) un atome d'halogène, (4) un groupe halogénoalkyle en C1 à 4, ou (5) un groupe  $-OR^{21}$ ,

$R^{11}$  représente (1) un groupe  $-OR^{101}$ , (2) un groupe  $SO_2R^{102}$ , (3) un groupe  $NR^{103}R^{104}$ , ou (4) un cycle carboné en C3 à 7 éventuellement substitué par un à trois atomes d'halogène,

$R^{12}$  représente (1) un groupe alkyle en C1 à 8 éventuellement substitué par un groupe hydroxyle, ou (2) un atome d'halogène,

$R^{13}$  représente (1) un groupe alkyle en C1 à 8 éventuellement substitué par un groupe hydroxyle, ou (2) un atome d'halogène,


$R^{21}$  représente (1) un atome d'hydrogène, ou (2) un groupe alkyle en C1 à 4,

$R^{22}$  représente (1) un groupe hydroxyle, ou (2) un groupe alkoxy en C1 à 4,

$R^{41}$  représente (1) un atome d'hydrogène ; (2) un groupe alkyle en C1 à 8 substitué par un à deux substituants choisis dans le groupe constitué par (a) un groupe cyclique de 5 à 7 chaînons éventuellement substitué par un

ou deux substituants choisis dans le groupe constitué par (i) un groupe alkyle en C1 à 4, (ii) un groupe halogénoalkyle en C1 à 4 et (iii) un atome d'halogène, (b) NR<sup>401</sup>R<sup>402</sup>, (c) un groupe hydroxyle et (d) un groupe SO<sub>2</sub>R<sup>403</sup> ;  
 (3) un groupe alcényle en C2 à 8 substitué par un à deux substituants choisis dans le groupe constitué par (a) un groupe cyclique de 5 à 7 chaînons éventuellement substitué par un ou deux substituants choisis dans le  
 groupe constitué par (i) un groupe alkyle en C1 à 4, (ii) un groupe halogénoalkyle en C1 à 4 et (iii) un atome d'halogène, (b) NR<sup>401</sup>R<sup>402</sup>, (c) un groupe hydroxyle et (d) un groupe SO<sub>2</sub>R<sup>403</sup> ; ou (4) un groupe alcynyle en  
 C2 à 8 substitué par un à deux substituants choisis dans le groupe constitué par (a) un groupe cyclique de 5 à 7 chaînons éventuellement substitué par un ou deux substituants choisis dans le groupe constitué par (i) un  
 groupe alkyle en C1 à 4, (ii) un groupe halogénoalkyle en C1 à 4 et (iii) un atome d'halogène, (b) NR<sup>401</sup>R<sup>402</sup>,  
 (c) un groupe hydroxyle et (d) un groupe SO<sub>2</sub>R<sup>403</sup> ;  
 R<sup>101</sup> représente (1) un atome d'hydrogène, ou (2) un groupe alkyle en C1 à 4,  
 R<sup>102</sup> représente (1) un atome d'hydrogène, ou (2) un groupe alkyle en C1 à 4,  
 R<sup>103</sup> et R<sup>104</sup> représentent chacun indépendamment (1) un atome d'hydrogène, ou (2) un groupe alkyle en C1 à 4,  
 R<sup>401</sup> et R<sup>402</sup> représentent chacun indépendamment (1) un atome d'hydrogène, ou (2) un groupe alkyle en C1 à 4,  
 R<sup>403</sup> représente (1) un atome d'hydrogène, ou (2) un groupe alkyle en C1 à 4,  
 A représente (1) CH, ou (2) un atome d'azote,  
 L représente (1) -O-, (2), -NH-, (3) -C(O)-, (4) -CR<sup>6</sup>R<sup>7</sup>-, (5) -S-, (6) -S(O)-, ou (7) -S(O)<sub>2</sub>-,  
 R<sup>6</sup> et R<sup>7</sup> représentent chacun indépendamment (1) un atome d'hydrogène, (2) un atome d'halogène, (3) un  
 groupe alkyle en C1 à 4, (4) un groupe hydroxyle, ou (5) NH<sub>2</sub>,  
 ring1 représente un benzène ou une pyridine,  
 ----- représente une liaison simple ou une liaison double,  
 m est un entier de 0 à 5,  
 n est un entier de 0 à 5,  
 p est un entier de 0 à 2,  
 q est un entier de 0 à 4,  
 lorsque m est supérieur ou égal à deux, une pluralité de R<sup>2</sup> peuvent être identiques ou différents les uns des  
 autres, et lorsque deux des R<sup>2</sup> représentent un groupe alkyle en C1 à 3 et sont sur le même atome de carbone,  
 les R<sup>2</sup> peuvent former, avec un atome de carbone qui leur est lié, un cycle carboné saturé en C3 à 7,  
 lorsque n est supérieur ou égal à 2, une pluralité de R<sup>3</sup> peuvent être identiques ou différents les uns des autres, et  
 lorsque q est supérieur ou égal à 2, une pluralité de R<sup>4</sup> peuvent être identiques ou différents les uns des autres,  
 sel de celui-ci, solvate de celui-ci ou N-oxyde de celui-ci.

2. Composé selon la revendication 1, dans lequel m est supérieur ou égal à un, et l'un des R<sup>2</sup> est nécessairement un groupe oxo.
3. Composé selon la revendication 1 ou 2, dans lequel L est (1) -O-, (2) -NH- ou (3) -C(O)-.
4. Composé selon la revendication 1, qui est représenté par la formule générale (I-1) :



dans laquelle R<sup>2-1</sup> représente (1) un groupe alkyle en C1 à 4, (2) un atome d'halogène, (3) un groupe halogé-

noalkyle en C1 à 4, (4) un groupe -OR<sup>21</sup>, ou (5) un groupe =NR<sup>22</sup>,

m-1 est un entier de 0 à 4,

L<sup>1</sup> est (1) -O-, (2) -NH-, ou (3) -C(O)-,

ring1-1 représente un benzène ou une pyridine,

lorsque m-1 est supérieur ou égal à 2, une pluralité de R<sup>2-1</sup> peuvent être identiques ou différents les uns des autres, et lorsque deux des R<sup>2-1</sup> représentent un groupe alkyle en C1 à 3 et sont sur le même atome de carbone, les R<sup>2-1</sup> peuvent former, avec un atome de carbone qui leur est lié, un cycle carboné saturé en C3 à 7, et les autres symboles ont les mêmes significations que défini à la revendication 1.

5. Composé selon l'une quelconque des revendications 1 à 4, qui est :

(2) le N-{5-[(6,7-diméthoxy-4-quinolinyloxy)-2-pyridinyl]-7,7-diméthyl-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(3) le N-{5-[(6,7-diméthoxy-4-quinolinyloxy)-2-pyridinyl]-1-(2,2-diméthylpropyl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(4) le N-{5-[(7-[3-(4-morpholinyloxy)-4-quinolinyloxy]-2-pyridinyl]-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(5) le N-{4-[(6,7-diméthoxy-4-quinolinyloxy)-3-fluorophényl]-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(6) le N-{4-[(6,7-diméthoxy-4-quinolinyloxy)phényl]-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(7) le N-{5-[(6,7-diméthoxy-4-quinolinyloxy)-2-pyridinyl]-1-(4-fluorophényl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(8) le N-{5-[(6,7-diméthoxy-4-quinolinyloxy)-2-pyridinyl]-1-(3-fluorophényl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(9) le N-{5-[(6,7-diméthoxy-4-quinolinyloxy)-2-pyridinyl]-1-(2-fluorophényl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(10) le N-{5-[(6,7-diméthoxy-4-quinazolinoyloxy)-2-pyridinyl]-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(11) le N-{5-[(6,7-diméthoxy-4-quinazolinoyloxy)-2-pyridinyl]-1-(4-fluorophényl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(12) le N-{5-[(6,7-diméthoxy-4-quinolinyloxy)-2-pyridinyl]-1-[(2S)-1-hydroxy-3-méthyl-2-butanyl]-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(13) le N-{4-[(6,7-diméthoxy-4-quinolinyloxy)-3-fluorophényl]-1-(3-fluorophényl)-2,5-dioxo-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(14) le N-{5-[(6,7-diméthoxy-4-quinolinyloxy)-2-pyridinyl]-6,6-diméthyl-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(15) le N-{5-[(6-méthoxy-7-[3-(4-morpholinyloxy)-4-quinolinyloxy]-2-pyridinyl]-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide,

(16) le N-{5-[(7-(3-hydroxy-3-méthylbutoxy)-6-méthoxy-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide, ou

(17) le N-{5-[(6-méthoxy-7-[3-(1-pyrrolidinyl)propoxy]-4-quinolinyloxy)-2-pyridinyl]-2,5-dioxo-1-phényl-1,2,5,6,7,8-hexahydro-3-quinolinecarboxamide.

6. Composition pharmaceutique contenant un composé représenté par la formule générale (I) ainsi que définie dans la revendication 1, un sel de celui-ci, un solvate de celui-ci, ou un N-oxyde de celui-ci.

7. Composition pharmaceutique selon la revendication 6, qui est un inhibiteur d'Axl.

8. Composition pharmaceutique selon la revendication 6, qui est un agent destiné à prévenir et/ou traiter une maladie liée à Axl.

9. Composition pharmaceutique selon la revendication 8, la maladie liée à Axl incluant un cancer, une maladie des reins, une maladie du système immunitaire ou une maladie du système circulatoire.

10. Composition pharmaceutique selon la revendication 9, le cancer étant une leucémie myéloïde aigüe, une leucémie myéloïde chronique, une leucémie lymphatique aigüe, un mélanome, un cancer du sein, un cancer du pancréas, un gliome, un adénocarcinome de l'œsophage, un cancer du gros intestin, un carcinome des cellules rénales, un

cancer de la thyroïde, un cancer du poumon non à petites cellules, un cancer de la prostate, un cancer de l'estomac, un cancer du foie, un mélanome malin uvéal, un cancer de l'ovaire, un cancer de l'endomètre, un lymphome, un cancer de la tête et du cou, ou un sarcome.

- 5      **11.** Composition pharmaceutique selon la revendication 6, qui est un agent anti-métastatique pour les cellules cancéreuses.
12. Composé représenté par la formule générale (I) selon la revendication 1, sel de celui-ci, solvate de celui-ci, ou N-oxyde de celui-ci destiné à être utilisé pour prévenir et/ou traiter une maladie liée à Axl.
- 10      **13.** Composé destiné à être utilisé selon la revendication 12, la maladie liée à Axl incluant un cancer, une maladie des reins, une maladie du système immunitaire ou une maladie du système circulatoire.
- 15      **14.** Composé destiné à être utilisé selon la revendication 13, le cancer étant une leucémie myéloïde aigüe, une leucémie myéloïde chronique, une leucémie lymphatique aigüe, un mélanome, un cancer du sein, un cancer du pancréas, un gliome, un adénocarcinome de l'œsophage, un cancer du gros intestin, un carcinome des cellules rénales, un cancer de la thyroïde, un cancer du poumon non à petites cellules, un cancer de la prostate, un cancer de l'estomac, un cancer du foie, un mélanome malin uvéal, un cancer de l'ovaire, un cancer de l'endomètre, un lymphome, un cancer de la tête et du cou, ou un sarcome.
- 20

25

30

35

40

45

50

55

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- WO 2012028332 A [0010]
- WO 2013074633 A [0010]
- WO 2012011548 A [0010]
- WO 2006116713 A [0010]
- WO 2007146824 A [0010]
- WO 2009137429 A [0010]

## Non-patent literature cited in the description

- *Clinical Science*, 2012, vol. 122, 361-368 [0011]
- *Proceedings of the national academy of sciences of the United States of America*, 2010, vol. 107 (3), 1124-1129 [0011]
- *Blood*, 2013, vol. 121, 2064-2073 [0011]
- **RICHARD C. LAROCK**. *Comprehensive Organic Transformations: A Guide to Functional Group Preparations*. John Wiley & Sons Inc, 1999 [0051]
- *Organic Letters*, 2001, vol. 3 (17), 2729-2732 [0054]
- **T. W. GREENE**. *Protective Groups in Organic Synthesis*. Wiley, 1999 [0065]
- **T. W. GREENE**. *Protective Groups in Organic Synthesis*. John Wiley & Sons Inc, 1999 [0068]
- *Hypertension*, 2002, vol. 39, 1095-1100 [0395]