



(11) **EP 3 473 713 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
24.04.2019 Bulletin 2019/17

(51) Int Cl.:
C12N 9/38 (2006.01)

(21) Application number: **18209685.9**

(22) Date of filing: **07.06.2013**

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **08.06.2012 US 201261657180 P**
08.06.2012 EP 12171335

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
17184821.1 / 3 305 896
13727611.9 / 2 859 098

(71) Applicant: **DuPont Nutrition Biosciences ApS**
1411 Copenhagen K (DK)

(72) Inventors:
• **CRAMER, Jacob Flyvholm**
8270 Højbjerg (DK)
• **LARSEN, Morten Krog**
8471 Sabro (DK)

(74) Representative: **D Young & Co LLP**
120 Holborn
London EC1N 2DY (GB)

Remarks:

This application was filed on 30-11-2018 as a
divisional application to the application mentioned
under INID code 62.

(54) **POLYPEPTIDES HAVING TRANSGALACTOSYLATING ACTIVITY**

(57) The present invention relates to polypeptides, specifically polypeptides having transgalactosylating activity and nucleic acids encoding these, and their uses in e.g. dairy product.

EP 3 473 713 A1

Description**FIELD OF THE INVENTION**

[0001] The present invention relates to polypeptides, specifically polypeptides having transgalactosylating activity and nucleic acids encoding these, and their uses in e.g. dairy product.

BACKGROUND OF THE INVENTION

[0002] Galactooligosaccharides (GOS) are carbohydrates which are nondigestible in humans and animals comprising two or more galactose molecules, typically up to nine, linked by glycosidic bonds. GOS's may also include one or more glucose molecules. One of the beneficial effects of GOS's is their ability of acting as prebiotic compounds by selectively stimulating the proliferation of beneficial colonic microorganisms such as bacteria to give physiological benefits to the consumer. The established health effects have resulted in a growing interest in GOSs as food ingredients for various types of food.

[0003] The enzyme β -galactosidase (EC 3.2.1.23) usually hydrolyses lactose to the monosaccharides D-glucose and D-galactose. In the normal enzyme reaction of β -galactosidases, the enzyme hydrolyses lactose and transiently binds the galactose monosaccharide in a galactose-enzyme complex that transfers galactose to the hydroxyl group of water, resulting in the liberation of D-galactose and D-glucose. However, at high lactose concentrations some β -galactosidases are able to transfer galactose to the hydroxyl groups of D-galactose or D-glucose in a process called transgalactosylation whereby galacto-oligosaccharides are produced. Also at high lactose concentrations some β -galactosidases are able to transfer galactose to the hydroxyl groups of lactose or higher order oligosaccharides.

[0004] The genus *Bifidobacterium* is one of the most commonly used types of bacteria cultures in the dairy industry for fermenting a variety of dairy products. Ingestion of *Bifidobacterium*-containing products furthermore has a health-promoting effect. This effect is not only achieved by a lowered pH of the intestinal contents but also by the ability of *Bifidobacterium* to repopulate the intestinal flora in individuals who have had their intestinal flora disturbed by for example intake of antibiotics. *Bifidobacterium* furthermore has the potential of outcompeting potential harmful intestinal microorganisms.

[0005] Galacto-oligosaccharides are known to enhance the growth of *Bifidobacterium*. This effect is likely achieved through the unique ability of *Bifidobacterium* to exploit galacto-oligosaccharides as a carbon source. Dietary supplement of galacto-oligosaccharides is furthermore thought to have a number of long-term disease protecting effects. For example, galacto-oligosaccharide intake has been shown to be highly protective against development of colorectal cancer in rats. There is therefore a great interest in developing cheap and efficient methods for producing galacto-oligosaccharides for use in the industry for improving dietary supplements and dairy products.

[0006] An extracellular lactase from *Bifidobacterium bifidum* DSM20215 truncated with approximately 580 amino acids (BIF3-d3) has been described as a transgalactosylating enzyme in a solution containing lactose solubilised in water (Jorgensen et al. (2001), Appl. Microbiol. Biotechnol., 57: 647- 652). WO 01/90317 also describes a truncation variant (OLGA347) as being a transgalactosylating enzyme and in WO 2012/010597 OLGA347 was shown to transfer a galactose moiety to D-fucose, N-acetyl-galactosamine and xylose.

[0007] In WO 2009/071539 a differently truncated fragment compared to BIF3-d3 is described as resulting in efficient hydrolysis and very low production of GOS when tested in milk.

[0008] The *Bifidobacterium bifidum* lactase enzymes described above have the drawback of either requiring high lactose concentrations such as above 10% (w/w) in order to produce GOS, or a high surplus of another acceptor molecule to generate heterooligosaccharides. Furthermore, a molecule has been described that predominately having beta-galactosylase (hydrolase) activity.

[0009] There is still a need to develop enzymes that are efficient at producing GOS in applications with low lactose substrate levels such as in milk.

SUMMARY OF THE INVENTION

[0010] It is an object of embodiments of the invention to provide a polypeptide which has a useful ratio of transgalactosylation to hydrolysis activity and thus are efficient producers of GOS when incubated with lactose even at low lactose levels such as in a milk-based product. It is a further object of embodiments of the invention to provide a method for production of galacto-oligosaccharides (GOS) in situ in dairy products. It is a further object of embodiments of the invention to provide a method for developing a cheaper and more efficient method for production of galacto-oligosaccharides (GOS) for use in the industry. It is further object of embodiments of the invention to provide polypeptides which are stable against further truncation such as by proteolytic degradation when produced in a suitable organism such as *Bacillus subtilis* e.g. *Bacillus subtilis* strain BG3594. It is yet a further object of embodiments of the invention to provide

polypeptides which are stable against further truncation during storage after final formulation.

[0011] The present inventors have surprisingly found that the polypeptides disclosed herein are efficient producers of galacto-oligosaccharides for example in situ when incubated in a lactose containing composition such as milk, wherein they have an efficient conversion of lactose into GOS resulting in a lower amount of free lactose. The presence of galacto-oligosaccharides in dairy products or other comestible products has the advantage of enhancing the growth of beneficial microbial strains (probiotics) such as the health-promoting *Bifidobacterium* sp. in the product itself and/or in the human or animal that consumes the product.

[0012] In one aspect, disclosed herein is a polypeptide having transgalactosylating activity, which comprises an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, and wherein said polypeptide, when being an expression product in a *Bacillus subtilis* strain BG3594 of a nucleic acid sequence, which encodes said polypeptide, is the only polypeptide expression product of said nucleic acid sequence that exhibits transgalactosylating activity.

[0013] In one aspect, disclosed herein is a polypeptide having transgalactosylating activity selected from the group consisting of:

a. a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues,

b. a polypeptide comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2, wherein said polypeptide consists of at most 975 amino acid residues,

c. a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues,

d. a polypeptide encoded by a polynucleotide that hybridizes under at least low stringency conditions with i) the nucleic acid sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding the polypeptide of SEQ ID NO: 1, 2, 3, 4, or 5; or ii) the complementary strand of i),

e. a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 70% identity to the nucleotide sequence encoding for the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5 or the nucleotide sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding a mature polypeptide, and

f. a polypeptide comprising a deletion, insertion and/or conservative substitution of one or more amino acid residues of SEQ ID NO: 1, 2, 3, 4 or 5.

[0014] In another aspect disclosed herein is a polypeptide having transgalactosylating activity selected from the group consisting of:

a. a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues,

b. a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues,

c. a polypeptide encoded by a polynucleotide that hybridizes under at least low stringency conditions with i) the nucleic acid sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding the polypeptide of SEQ ID NO: 1, 2, 3, 4, or 5; or ii) the complementary strand of i),

d. a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 70% identity to the nucleotide sequence encoding for the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5 or the nucleotide sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding a mature polypeptide, and

e. a polypeptide comprising a deletion, insertion and/or conservative substitution of one or more amino acid residues of SEQ ID NO: 1, 2, 3, 4 or 5.

[0015] In one aspect, disclosed herein is polypeptide which is a C-terminal truncated fragment of SEQ ID NO:22 having transgalactosylating activity and which are stable against further truncation such as by proteolytic degradation when produced in a suitable organism such as *Bacillus subtilis* e.g. *Bacillus subtilis* strain BG3594 and/or which are stable against further truncation during storage after final formulation.

[0016] In one aspect, disclosed herein is a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues.

[0017] In one aspect, disclosed herein is a polypeptide comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2, wherein said polypeptide consists of at most 975 amino acid residues, is provided.

[0018] In one aspect, disclosed herein is a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues.

In one aspect, disclosed herein is a nucleic acid capable of encoding a polypeptide as described herein.

[0019] In one aspect, disclosed herein is an expression vector and/or a plasmid comprising a nucleic acid as described herein, or capable of expressing a polypeptide as described herein.

[0020] In one aspect, disclosed herein is a cell capable of expressing a polypeptide as described herein.

[0021] In one aspect, disclosed herein is a method of expressing a polypeptide, the method comprising obtaining a cell as described herein and expressing the polypeptide from the cell, and optionally purifying the polypeptide.

[0022] In one aspect, disclosed herein is a composition comprising a polypeptide as described herein, preferably a food composition, more preferably a dairy product.

[0023] In one aspect, disclosed herein is a method for producing a food product such as a dairy product by treating a milk-based substrate comprising lactose with a polypeptide as described herein.

[0024] In one aspect, disclosed herein is a galacto-oligosaccharide or composition thereof obtained by treating a substrate comprising lactose with a polypeptide as described herein.

LEGENDS TO THE FIGURE

[0025]

Figure 1 shows a plasmid map for the BIF_1326 variant for recombinant expression in *Bacillus subtilis*.

Figure 2 shows SDS-PAGE showing truncation variants purified using HyperQ column eluted with a NaCl gradient.

Figure 3 shows the ratio of transgalactosylation activity. Ratio is calculated as ratio between Abs420 with acceptor present divided by Abs420 without acceptor present times 100. Variants at or below index 100 are purely hydrolytic variants, whereas the level above reflects relative transgalactosylating activity.

Figure 4 shows Galacto-oligosaccharides (GOS) generating efficacy of selected variants in a yoghurt matrix at 30°C for 3 hours. In this example GOS is the accumulative amount oligosaccharides at and above DP3.

Figure 5 shows SDS-PAGE gel showing the different variants from table 2 expressed and the degradation fragments detected. Lower panel shows magnification and identification of degradation bands.

SEQUENCE LISTING

[0026]

SEQ ID NO: 1 (also named (BIF_917) herein) is a 887 amino acid truncated fragment of SEQ ID NO: 22.

SEQ ID NO: 2 (also named (BIF_995) herein) is a 965 amino acid truncated fragment of SEQ ID NO: 22.

SEQ ID NO: 3 (also named (BIF_1068) herein) is a 1038 amino acid truncated fragment of SEQ ID NO: 22.

SEQ ID NO: 4 (also named (BIF_1172) herein) is a 1142 amino acid truncated fragment of SEQ ID NO: 22.

SEQ ID NO: 5 (also named (BIF_1241) herein) is a 1211 amino acid truncated fragment of SEQ ID NO: 22.

SEQ ID NO: 6 (also named (BIF_1326) herein) is a 1296 amino acid truncated fragment of SEQ ID NO: 22.

SEQ ID NO: 7 is *Bifidobacterium bifidum* glycoside hydrolase catalytic core

SEQ ID NO: 8 is a nucleotide sequence encoding an extracellular lactase from *Bifidobacterium bifidum* DSM20215

SEQ ID NO: 9 is nucleotide sequence encoding BIF_917

SEQ ID NO: 10 is nucleotide sequence encoding BIF_995

SEQ ID NO: 11 is nucleotide sequence encoding BIF_1068

SEQ ID NO: 12 is nucleotide sequence encoding BIF_1172

SEQ ID NO: 13 is nucleotide sequence encoding BIF_1241

SEQ ID NO: 14 is nucleotide sequence encoding BIF_1326

SEQ ID NO: 15 is forward primer for generation of above BIF variants

SEQ ID NO: 16 is reverse primer for BIF917

SEQ ID NO: 17 is reverse primer for BIF995

SEQ ID NO: 18 is reverse primer for BIF1068

SEQ ID NO: 19 is reverse primer for BIF1241

SEQ ID NO: 20 is reverse primer for BIF1326

SEQ ID NO: 21 is reverse primer for BIF1478

SEQ ID NO: 22 is extracellular lactase from Bifidobacterium bifidum DSM20215.

SEQ ID NO: 23 is signal sequence of extracellular lactase from Bifidobacterium bifidum DSM20215

DETAILED DISCLOSURE OF THE INVENTION

Definitions

[0027] In accordance with this detailed description, the following abbreviations and definitions apply. It should be noted that as used herein, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a polypeptide" includes a plurality of such polypeptides, and reference to "the formulation" includes reference to one or more formulations and equivalents thereof known to those skilled in the art, and so forth.

[0028] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. The following terms are provided below.

[0029] "Transgalactosylase" means an enzyme that, among other things, is able to transfer galactose to the hydroxyl groups of D-galactose or D-glucose whereby galacto-oligosaccharides are produced. In one aspect, a transgalactosylase is identified by reaction of the enzyme on lactose in which the amount of galactose generated is less than the amount of glucose generated at any given time.

[0030] In the present context, the term "transgalactosylating activity" means the transfer of a galactose moiety to a molecule other than water. The activity can be measured as [glucose] - [galactose] generated at any given time during reaction or by direct quantification of the GOS generated at any given time during the reaction. This measurement may be performed in several ways such as by a HPLC method as shown in the examples. When comparing measurements of transgalactosylating activity, they have been performed at a given initial lactose concentration, such as e.g. 3, 4, 5, 6, 7, 8, 9 or 10% (w/w).

[0031] In the present context, the term "β-galactosidase activity" means the ability of an enzyme to hydrolyse β-galactosides such as for example lactose into monosaccharides, glucose and galactose.

[0032] In the context of calculating transgalactosylating activity:β-galactosidase activity, the β-galactosidase activity is measured as [galactose] generated at any given time during reaction. This measurement may be performed in several ways such as by a HPLC method as shown in the examples.

[0033] In the present context, the term "ratio of transgalactosylation activity" using ortho-nitrophenol-β-D-galactopyranoside (ONPG) was calculated as follows: Ratio is calculated as ratio between Abs420 with acceptor present divided by Abs420 without acceptor present times 100. Variant at or below index 100 are purely hydrolytic variants, whereas the level above depicts relative transgalactosylating activity.

[0034] Ratio of transgalactosylation activity = $(\text{Abs420}^{+\text{Cellobiose}} / \text{Abs420}^{-\text{Cellobiose}}) * 100\%$, where Abs420^{+Cellobiose} is the absorbance read at 420nm using the described method 3 below including cellobiose in the reaction and Abs420^{-Cellobiose} is the absorbance read at 420nm using the described method 3 below but without cellobiose in the reaction. The equation above is only valid for dilutions where the absorbance is between 0.5 and 1.0.

[0035] In one aspect, the activity is measured after 15 min. reaction, 30 min. reaction, 60 min. reaction, 90 min. reaction, 120 min. reaction or 180 min. reaction. Thus in one aspect, as an example the relative transgalactosylation activity is measured 15 minutes after addition of enzyme, such as 30 minutes after addition of enzyme, such as 60 minutes after addition of enzyme, such as 90 minutes after addition of enzyme, such as 120 minutes after addition of enzyme or such as 180 minutes after addition of enzyme.

[0036] In the present context, the term "ratio of transgalactosylating activity:β-galactosidase activity" means $([\text{Glucose}] - [\text{Galactose}] / [\text{Galactose}])$.

[0037] In the present context, the term [Glucose] means the glucose concentration in % by weight as measured by HPLC.

[0038] In the present context, the term [Galactose] means the galactose concentration in % by weight as measured by HPLC.

[0039] In the present context, the term "lactose has been transgalactosylated" means that a galactose molecule has been covalently linked to the lactose molecule such as for example covalently linked to any of the free hydroxyl groups in the lactose molecule or as generated by internal transgalatotsylation for example forming allolactose.

[0040] In the present context, the evaluation of performance of polypeptides disclosed herein in galactooligosaccharide (GOS) production were tested in a "milk-based assay" (yogurt application mimic). Batch experiments with a volume of 100 μl were performed in 96 well MTP plates using a yogurt mix, consisting of 98,60%(w/v) fresh pasteurized low-fat

milk (Aria Mini-maelk) and 1.4% (w/v) Nutrilac YQ-5075 whey ingredient (Aria). To completely hydrate Nutrilac YQ-5075 the mixture was left with agitation for 20h and afterwards added 20mM NaPhosphate pH 6.5 to ensure a pH of 6.5. This yogurt-base was either used plain or with various supplements such as additional lactose, fucose, maltose, xylose or salts. 90 μ l of the yogurt was mixed with 10 μ l purified enzyme or crude ferment, sealed with tape and incubated at 43°C for 3 hours. The reaction was stopped by 100 μ l 10% Na₂CO₃. Samples were stored at -20°C. Quantification of galactooligosaccharides (GOS), lactose, glucose and galactose were performed by HPLC. Analysis of samples was carried out on a Dionex ICS 3000. IC parameters were as follows: Mobile phase: 150 mM NaOH, Flow: Isochratic, 0,25ml/min, Column: Carbowax PA1, Column temperature: RT, Injection volume: 10 μ L, Detector: PAD, Integration: Manual, Sample preparation: 100 times dilution in Milli-Q water (0,1 ml sample + 9,9 ml water) and filtration through 0.45 μ m syringe filters, Quantification: Peak areas in percent of peak area of the standard. A GOS syrup (Vivanal GOS, Friesland Campina) was used as standard for GOS quantification. Results of such an evaluation is shown in Figure 4, and further described in example 2.

[0041] In the present context, the term "which polypeptide is freeze-dried" means that the polypeptide has been obtained by freeze-drying a liquid of the polypeptide at an appropriate pressure and for an appropriate period removing the water.

[0042] In the present context, the term "which polypeptide is in solution" relates to a polypeptide which is soluble in a solvent without precipitating out of solution. A solvent for this purpose includes any milieu in which the polypeptide may occur, such as an aqueous buffer or salt solution, a fermentation broth, or the cytoplasm of an expression host.

[0043] In the present context, the term "stabilizer" means any stabilizer for stabilizing the polypeptide e.g., a polyol such as, e.g., glycerol or propylene glycol, a sugar or a sugar alcohol, lactic acid, boric acid, or a boric acid derivative (e.g., an aromatic borate ester). In one aspect, the stabilizer is glycerol.

[0044] In the present context, the term "carbohydrate substrate" means an organic compound with the general formula $C_m(H_2O)_n$, that is, consisting only of carbon, hydrogen and oxygen, the last two in the 2:1 atom ratio such as a disaccharide.

[0045] In the present context, the term "disaccharide" is two monosaccharide units bound together by a covalent bond known as a glycosidic linkage formed via a dehydration reaction, resulting in the loss of a hydrogen atom from one monosaccharide and a hydroxyl group from the other. The formula of unmodified disaccharides is $C_{12}H_{22}O_{11}$. In one aspect, the disaccharide is lactulose, trehalose, rhamnose, maltose, sucrose, lactose, fucose or cellobiose. In a further aspect, the disaccharide is lactose.

[0046] The term "isolated" means that the polypeptide is at least substantially free from at least one other component with which the sequence is naturally associated in nature and as found in nature. In one aspect, "isolated polypeptide" as used herein refers to a polypeptide which is at least 30% pure, at least 40% pure, at least 60% pure, at least 80% pure, at least 90% pure, and at least 95% pure, as determined by SDS-PAGE.

[0047] The term "substantially pure polypeptide" means herein a polypeptide preparation which contains at most 10%, preferably at most 8%, more preferably at most 6%, more preferably at most 5%, more preferably at most 4%, at most 3%, even more preferably at most 2%, most preferably at most 1%, and even most preferably at most 0.5% by weight of other polypeptide material with which it is natively associated. It is, therefore, preferred that the substantially pure polypeptide is at least 92% pure, preferably at least 94% pure, more preferably at least 95% pure, more preferably at least 96% pure, more preferably at least 96% pure, more preferably at least 97% pure, more preferably at least 98% pure, even more preferably at least 99%, most preferably at least 99.5% pure, and even most preferably 100% pure by weight of the total polypeptide material present in the preparation. The polypeptides disclosed herein are preferably in a substantially pure form. In particular, it is preferred that the polypeptides are in "essentially pure form", i.e., that the polypeptide preparation is essentially free of other polypeptide material with which it is natively associated. This can be accomplished, for example, by preparing the polypeptide by means of well-known recombinant methods or by classical purification methods. Herein, the term "substantially pure polypeptide" is synonymous with the terms "isolated polypeptide" and "polypeptide in isolated form."

[0048] The term "purified" or "pure" means that a given component is present at a high level state - e.g. at least about 51% pure, such as at least 51% pure, or at least about 75% pure such as at least 75% pure, or at least about 80% pure such as at least 80% pure, or at least about 90% pure such as at least 90% pure, or at least about 95% pure such as at least 95% pure, or at least about 98% pure such as at least 98% pure. The component is desirably the predominant active component present in a composition.

[0049] The term "microorganism" in relation to the present invention includes any "microorganism" that could comprise a nucleotide sequence according to the present invention or a nucleotide sequence encoding for a polypeptide having the specific properties as defined herein and/or products obtained therefrom. In the present context, "microorganism" may include any bacterium or fungus being able to ferment a milk substrate.

[0050] The term "host cell" - in relation to the present invention includes any cell that comprises either a nucleotide sequence encoding a polypeptide having the specific properties as defined herein or an expression vector as described above and which is used in the production of a polypeptide having the specific properties as defined herein. In one aspect, the production is recombinant production.

[0051] The term "milk", in the context of the present invention, is to be understood as the lacteal secretion obtained from any mammal, such as cows, sheep, goats, buffaloes or camels.

In the present context, the term "milk-based substrate" means any raw and/or processed milk material or a material derived from milk constituents. Useful milk-based substrates include, but are not limited to solutions/suspensions of any milk or milk like products comprising lactose, such as whole or low fat milk, skim milk, buttermilk, reconstituted milk powder, condensed milk, solutions of dried milk, UHT milk, whey, whey permeate, acid whey, or cream. Preferably, the milk-based substrate is milk or an aqueous solution of skim milk powder. The milk-based substrate may be more concentrated than raw milk. In one embodiment, the milk-based substrate has a ratio of protein to lactose of at least 0.2, preferably at least 0.3, at least 0.4, at least 0.5, at least 0.6 or, most preferably, at least 0.7. The milk-based substrate may be homogenized and/or pasteurized according to methods known in the art.

[0052] "Homogenizing" as used herein means intensive mixing to obtain a soluble suspension or emulsion. It may be performed so as to break up the milk fat into smaller sizes so that it no longer separates from the milk. This may be accomplished by forcing the milk at high pressure through small orifices.

[0053] "Pasteurizing" as used herein means reducing or eliminating the presence of live organisms, such as microorganisms, in the milk-based substrate. Preferably, pasteurization is attained by maintaining a specified temperature for a specified period of time. The specified temperature is usually attained by heating. The temperature and duration may be selected in order to kill or inactivate certain bacteria, such as harmful bacteria, and/or to inactivate enzymes in the milk. A rapid cooling step may follow. A "food product" or "food composition" in the context of the present invention may be any comestible food or feed product suitable for consumption by an animal or human.

[0054] A "dairy product" in the context of the present invention may be any food product wherein one of the major constituents is milk-based. Preferable, the major constituent is milk-based. More preferably, the major constituent is a milk-based substrate which has been treated with an enzyme having transgalactosylating activity.

[0055] In the present context, "one of the major constituents" means a constituent having a dry matter which constitutes more than 20%, preferably more than 30% or more than 40% of the total dry matter of the dairy product, whereas "the major constituent" means a constituent having a dry matter which constitutes more than 50%, preferably more than 60% or more than 70% of the total dry matter of the dairy product.

[0056] A "fermented dairy product" in present context is to be understood as any dairy product wherein any type of fermentation forms part of the production process. Examples of fermented dairy products are products like yoghurt, buttermilk, creme fraiche, quark and fromage frais. Another example of a fermented dairy product is cheese. A fermented dairy product may be produced by any method known in the art.

[0057] The term "fermentation" means the conversion of carbohydrates into alcohols or acids through the action of a microorganism such as a starter culture. In one aspect, fermentation comprises conversion of lactose to lactic acid.

[0058] In the present context, "microorganism" may include any bacterium or fungus being able to ferment a milk substrate.

[0059] In the present context the term "Pfam domains" means regions within a protein sequence that are identified as either Pfam-A or Pfam-B based on multiple sequence alignments and the presence of Hidden Markov Motifs ("The Pfam protein families database": R.D. Finn, J. Mistry, J. Tate, P. Coghill, A. Heger, J.E. Pollington, O.L. Gavin, P. Gunasekaran, G. Ceric, K. Forslund, L. Holm, E.L. Sonnhammer, S.R. Eddy, A. Bateman Nucleic Acids Research (2010) Database Issue 38:D211-222.). As examples of Pfam domains mention may be made of Glyco_hydro2N (PF02837), Glyco_hydro (PF00703), Glyco_hydro 2C (PF02836) and Bacterial Ig-like domain (group 4) (PF07532).

[0060] As used herein "a position corresponding to position" means that an alignment as described herein is made between a particular query polypeptide and the reference polypeptide. The position corresponding to a specific position in the reference polypeptide is then identified as the corresponding amino acid in the alignment with the highest sequence identity.

[0061] A "variant" or "variants" refers to either polypeptides or nucleic acids. The term "variant" may be used interchangeably with the term "mutant". Variants include insertions, substitutions, transversions, truncations, and/or inversions at one or more locations in the amino acid or nucleotide sequence, respectively. The phrases "variant polypeptide", "polypeptide variant", "polypeptide", "variant" and "variant enzyme" mean a polypeptide/protein that has an amino acid sequence that either has or comprises a selected amino acid sequence of or is modified compared to the selected amino acid sequence, such as SEQ ID NO: 1, 2, 3, 4 or 5.

[0062] As used herein, "reference enzymes," "reference sequence," "reference polypeptide" mean enzymes and polypeptides from which any of the variant polypeptides are based, e.g., SEQ ID NO: 1, 2, 3, 4 or 5. A "reference nucleic acid" means a nucleic acid sequence encoding the reference polypeptide.

[0063] As used herein, the terms "reference sequence" and "subject sequence" are used interchangeably.

[0064] As used herein, "query sequence" means a foreign sequence, which is aligned with a reference sequence in order to see if it falls within the scope of the present invention. Accordingly, such query sequence can for example be a prior art sequence or a third party sequence.

[0065] As used herein, the term "sequence" can either be referring to a polypeptide sequence or a nucleic acid

sequence, depending of the context.

[0066] As used herein, the terms "polypeptide sequence" and "amino acid sequence" are used interchangeably.

[0067] The signal sequence of a "variant" may be the same or may differ from the signal sequence of the wild-type a Bacillus signal peptide or any signal sequence that will secrete the polypeptide. A variant may be expressed as a fusion protein containing a heterologous polypeptide. For example, the variant can comprise a signal peptide of another protein or a sequence designed to aid identification or purification of the expressed fusion protein, such as a His-Tag sequence.

[0068] To describe the various variants that are contemplated to be encompassed by the present disclosure, the following nomenclature will be adopted for ease of reference. Where the substitution includes a number and a letter, e.g., 592P, then this refers to {position according to the numbering system/substituted amino acid}. Accordingly, for example, the substitution of an amino acid to proline in position 592 is designated as 592P. Where the substitution includes a letter, a number, and a letter, e.g., D592P, then this refers to {original amino acid/position according to the numbering system/substituted amino acid}.

[0069] Accordingly, for example, the substitution of alanine with proline in position 592 is designated as A592P.

[0070] Where two or more substitutions are possible at a particular position, this will be designated by contiguous letters, which may optionally be separated by slash marks "/", e.g., G303ED or G303E/D.

[0071] Position(s) and substitutions are listed with reference to for example either SEQ ID NO: 1, 2, 3, 4 or 5. For example equivalent positions in another sequence may be found by aligning this sequence with either SEQ ID NO: 1, 2, 3, 4 or 5 to find an alignment with the highest percent identity and thereafter determining which amino acid aligns to correspond with an amino acid of a specific position of either SEQ ID NO: 1, 2, 3, 4 or 5. Such alignment and use of one sequence as a first reference is simply a matter of routine for one of ordinary skill in the art.

[0072] As used herein, the term "expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation.

[0073] As used herein, "polypeptide" is used interchangeably with the terms "amino acid sequence", "enzyme", "peptide" and/or "protein". As used herein, "nucleotide sequence" or "nucleic acid sequence" refers to an oligonucleotide sequence or polynucleotide sequence and variants, homologues, fragments and derivatives thereof. The nucleotide sequence may be of genomic, synthetic or recombinant origin and may be double-stranded or single-stranded, whether representing the sense or anti-sense strand. As used herein, the term "nucleotide sequence" includes genomic DNA, cDNA, synthetic DNA, and RNA.

[0074] "Homologue" means an entity having a certain degree of identity or "homology" with the subject amino acid sequences and the subject nucleotide sequences. In one aspect, the subject amino acid sequence is SEQ ID NO: 1, 2, 3, 4 or 5, and the subject nucleotide sequence preferably is SEQ ID NO: 9, 10, 11, 12 or 13.

[0075] A "homologous sequence" includes a polynucleotide or a polypeptide having a certain percent, e.g., 80%, 85%, 90%, 95%, or 99%, of sequence identity with another sequence. Percent identity means that, when aligned, that percentage of bases or amino acid residues are the same when comparing the two sequences. Amino acid sequences are not identical, where an amino acid is substituted, deleted, or added compared to the subject sequence. The percent sequence identity typically is measured with respect to the mature sequence of the subject protein, i.e., following removal of a signal sequence, for example. Typically, homologues will comprise the same active site residues as the subject amino acid sequence. Homologues also retain enzymatic activity, although the homologue may have different enzymatic properties than the wild-type.

[0076] As used herein, "hybridization" includes the process by which a strand of nucleic acid joins with a complementary strand through base pairing, as well as the process of amplification as carried out in polymerase chain reaction (PCR) technologies. The variant nucleic acid may exist as single- or double-stranded DNA or RNA, an RNA/DNA heteroduplex or an RNA/DNA copolymer. As used herein, "copolymer" refers to a single nucleic acid strand that comprises both ribonucleotides and deoxyribonucleotides. The variant nucleic acid may be codon-optimized to further increase expression.

[0077] As used herein, a "synthetic" compound is produced by *in vitro* chemical or enzymatic synthesis. It includes, but is not limited to, variant nucleic acids made with optimal codon usage for host organisms, such as a yeast cell host or other expression hosts of choice.

[0078] As used herein, "transformed cell" includes cells, including both bacterial and fungal cells, which have been transformed by use of recombinant DNA techniques. Transformation typically occurs by insertion of one or more nucleotide sequences into a cell. The inserted nucleotide sequence may be a heterologous nucleotide sequence, i.e., is a sequence that is not natural to the cell that is to be transformed, such as a fusion protein.

[0079] As used herein, "operably linked" means that the described components are in a relationship permitting them to function in their intended manner. For example, a regulatory sequence operably linked to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under condition compatible with the control sequences.

[0080] As used herein, the term "fragment" is defined herein as a polypeptide having one or more (several) amino acids deleted from the amino and/or carboxyl terminus wherein the fragment has activity.

[0081] In one aspect, the term "fragment" is defined herein as a polypeptide having one or more (several) amino acids

deleted from the amino and/or carboxyl terminus of the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5; wherein the fragment has transgalactosylating activity.

[0082] The term "Galactose Binding domain-like" as used herein is abbreviated to and interchangeable with the term "GBD".

Degree of identity

[0083] The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "identity".

[0084] In one embodiment, the degree of sequence identity between a query sequence and a reference sequence is determined by 1) aligning the two sequences by any suitable alignment program using the default scoring matrix and default gap penalty, 2) identifying the number of exact matches, where an exact match is where the alignment program has identified an identical amino acid or nucleotide in the two aligned sequences on a given position in the alignment and 3) dividing the number of exact matches with the length of the reference sequence.

[0085] In one embodiment, the degree of sequence identity between a query sequence and a reference sequence is determined by 1) aligning the two sequences by any suitable alignment program using the default scoring matrix and default gap penalty, 2) identifying the number of exact matches, where an exact match is where the alignment program has identified an identical amino acid or nucleotide in the two aligned sequences on a given position in the alignment and 3) dividing the number of exact matches with the length of the longest of the two sequences.

[0086] In another embodiment, the degree of sequence identity between the query sequence and the reference sequence is determined by 1) aligning the two sequences by any suitable alignment program using the default scoring matrix and default gap penalty, 2) identifying the number of exact matches, where an exact match is where the alignment program has identified an identical amino acid or nucleotide in the two aligned sequences on a given position in the alignment and 3) dividing the number of exact matches with the "alignment length", where the alignment length is the length of the entire alignment including gaps and overhanging parts of the sequences.

[0087] Sequence identity comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs use complex comparison algorithms to align two or more sequences that best reflect the evolutionary events that might have led to the difference(s) between the two or more sequences. Therefore, these algorithms operate with a scoring system rewarding alignment of identical or similar amino acids and penalising the insertion of gaps, gap extensions and alignment of non-similar amino acids. The scoring system of the comparison algorithms include:

- i) assignment of a penalty score each time a gap is inserted (gap penalty score),
- ii) assignment of a penalty score each time an existing gap is extended with an extra position (extension penalty score),
- iii) assignment of high scores upon alignment of identical amino acids, and
- iv) assignment of variable scores upon alignment of non-identical amino acids.

[0088] Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons.

The scores given for alignment of non-identical amino acids are assigned according to a scoring matrix also called a substitution matrix. The scores provided in such substitution matrices are reflecting the fact that the likelihood of one amino acid being substituted with another during evolution varies and depends on the physical/chemical nature of the amino acid to be substituted. For example, the likelihood of a polar amino acid being substituted with another polar amino acid is higher compared to being substituted with a hydrophobic amino acid. Therefore, the scoring matrix will assign the highest score for identical amino acids, lower score for non-identical but similar amino acids and even lower score for non-identical non-similar amino acids. The most frequently used scoring matrices are the PAM matrices (Dayhoff et al. (1978), Jones et al. (1992)), the BLOSUM matrices (Henikoff and Henikoff (1992)) and the Gonnet matrix (Gonnet et al. (1992)).

[0089] Suitable computer programs for carrying out such an alignment include, but are not limited to, Vector NTI (Invitrogen Corp.) and the ClustalV, ClustalW and ClustalW2 programs (Higgins DG & Sharp PM (1988), Higgins et al. (1992), Thompson et al. (1994), Larkin et al. (2007)). A selection of different alignment tools is available from the ExPASy Proteomics server at www.expasy.org. Another example of software that can perform sequence alignment is BLAST (Basic Local Alignment Search Tool), which is available from the webpage of National Center for Biotechnology Information which can currently be found at <http://www.ncbi.nlm.nih.gov/> and which was firstly described in Altschul et al. (1990) J. Mol. Biol. 215; 403-410.

[0090] In a preferred embodiment of the present invention, the alignment program is performing a global alignment program, which optimizes the alignment over the full-length of the sequences. In a further preferred embodiment, the global alignment program is based on the Needleman-Wunsch algorithm (Needleman, Saul B.; and Wunsch, Christian

D. (1970), "A general method applicable to the search for similarities in the amino acid sequence of two proteins", Journal of Molecular Biology 48 (3): 443-53). Examples of current programs performing global alignments using the Needleman-Wunsch algorithm are EMBOSS Needle and EMBOSS Stretcher programs, which are both available at <http://www.ebi.ac.uk/Tools/psa/>.

[0091] EMBOSS Needle performs an optimal global sequence alignment using the Needleman-Wunsch alignment algorithm to find the optimum alignment (including gaps) of two sequences along their entire length.

[0092] EMBOSS Stretcher uses a modification of the Needleman-Wunsch algorithm that allows larger sequences to be globally aligned.

[0093] In one embodiment, the sequences are aligned by a global alignment program and the sequence identity is calculated by identifying the number of exact matches identified by the program divided by the "alignment length", where the alignment length is the length of the entire alignment including gaps and overhanging parts of the sequences.

[0094] In a further embodiment, the global alignment program uses the Needleman-Wunsch algorithm and the sequence identity is calculated by identifying the number of exact matches identified by the program divided by the "alignment length", where the alignment length is the length of the entire alignment including gaps and overhanging parts of the sequences.

[0095] In yet a further embodiment, the global alignment program is selected from the group consisting of EMBOSS Needle and EMBOSS stretcher and the sequence identity is calculated by identifying the number of exact matches identified by the program divided by the "alignment length", where the alignment length is the length of the entire alignment including gaps and overhanging parts of the sequences.

[0096] Once the software has produced an alignment, it is possible to calculate % similarity and % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

[0097] In one embodiment, it is preferred to use the ClustalW software for performing sequence alignments. Preferably, alignment with ClustalW is performed with the following parameters for pairwise alignment:

Substitution matrix:	Gonnet 250
Gap open penalty:	20
Gap extension penalty:	0.2
Gap end penalty:	None

[0098] ClustalW2 is for example made available on the internet by the European Bioinformatics Institute at the EMBL-EBI webpage www.ebi.ac.uk under tools - sequence analysis - ClustalW2. Currently, the exact address of the ClustalW2 tool is www.ebi.ac.uk/Tools/clustalw2.

[0099] In another embodiment, it is preferred to use the program Align X in Vector NTI (Invitrogen) for performing sequence alignments. In one embodiment, Exp10 has been may be used with default settings:

Gap opening penalty: 10
 Gap extension penalty: 0.05
 Gapseparation penalty range: 8

[0100] In a another embodiment, the alignment of one amino acid sequence with, or to, another amino acid sequence is determined by the use of the score matrix: blosum62mt2 and the VectorNTI Pair wise alignment settings

Settings	K-tuple	1
	Number of best diagonals	5
	Window size	5
	Gap Penalty	3
	Gap opening Penalty	10
	Gap extension Penalty	0,1

[0101] In one embodiment, the percentage of identity of one amino acid sequence with, or to, another amino acid sequence is determined by the use of Blast with a word size of 3 and with BLOSUM 62 as the substitution matrix

Polypeptides

[0102] In one aspect, disclosed herein is a polypeptide having a ratio of transgalactosylating activity: β -galactosidase activity of at least 0.5, at least 1, at least 2, at least 2.5, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, or at least 12 at or above a concentration of 3% w/w initial lactose concentration.

[0103] In one aspect, disclosed herein is a polypeptide, wherein the glycoside hydrolase catalytic core has an amino acid sequence of SEQ ID NO:7.

[0104] In one aspect, disclosed herein is a polypeptide containing a Glyco_hydro2N (PF02837), a Glyco_hydro (PF00703) and/or a Glyco_hydro 2C (PF02836) domains.

[0105] In one aspect, disclosed herein is a polypeptide containing the Bacterial Ig-like domain (group 4) (PF07532).

[0106] In one aspect, disclosed herein is a polypeptide having transgalactosylating activity selected from the group consisting of:

a. a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues,

b. a polypeptide comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2, wherein said polypeptide consists of at most 975 amino acid residues,

c. a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues,

d. a polypeptide encoded by a polynucleotide that hybridizes under at least low stringency conditions with i) the nucleic acid sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5; or ii) the complementary strand of i),

e. a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 70% identity to the nucleotide sequence encoding for the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5 or the nucleotide sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding a mature polypeptide, and

f. a polypeptide comprising a deletion, insertion and/or conservative substitution of one or more amino acid residues of SEQ ID NO: 1, 2, 3, 4 or 5.

[0107] In another aspect disclosed herein is a polypeptide having transgalactosylating activity selected from the group consisting of:

a. a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues,

b. a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues,

c. a polypeptide encoded by a polynucleotide that hybridizes under at least low stringency conditions with i) the nucleic acid sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding the polypeptide of SEQ ID NO: 1, 2, 3, 4, or 5; or ii) the complementary strand of i),

d. a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 70% identity to the nucleotide sequence encoding for the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5 or the nucleotide sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding a mature polypeptide, and

e. a polypeptide comprising a deletion, insertion and/or conservative substitution of one or more amino acid residues of SEQ ID NO: 1, 2, 3, 4 or 5.

[0108] In one aspect, disclosed herein is a polypeptide, wherein the amino acid sequence has at least 68%, 70%, 72%, 74%, 76%, 78%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, sequence identity to the mature amino acid sequence of SEQ ID NO: 1, 2, 3, 4 or 5.

[0109] In one aspect, disclosed herein is a polypeptide having 90% sequence identity to the mature amino acid sequence of SEQ ID NO:1.

[0110] In one aspect, disclosed herein is a polypeptide having 90% sequence identity to the mature amino acid sequence of SEQ ID NO:2.

[0111] In one aspect, disclosed herein is a polypeptide having 96.5% sequence identity to the mature amino acid sequence of SEQ ID NO:3.

5 **[0112]** In one aspect, disclosed herein is a polypeptide having 96.5% sequence identity to the mature amino acid sequence of SEQ ID NO:4.

[0113] In one aspect, disclosed herein is a polypeptide having 96.5% sequence identity to the mature amino acid sequence of SEQ ID NO:5.

10 **[0114]** In one aspect, disclosed herein is a polypeptide comprising or consisting of the amino acid sequence of SEQ ID NO:1, 2, 3, 4 or 5.

[0115] In one aspect, disclosed herein is a polypeptide, which is derived from *Bifidobacterium bifidum*.

[0116] In one aspect, disclosed herein is a polypeptide having a pH optimum of 6.5-7.5.

[0117] In one aspect, disclosed herein is a polypeptide having a temperature optimum of 30-60 such as 42-60 degree celcius.

15 **[0118]** Polypeptides having activity on carbohydrates can be classified using either the IUBMB system of classification based on their substrate specificity or on the CaZy assignment into one of the current 125 glycoside hydrolase family. In the CaZy database the assignment is based on both sequence and structural information combined with knowledge of stereochemistry of the substrates and products

20 **[0119]** Disclosed herein are polypeptides which when being an expression product in a *Bacillus subtilis* strain BG3594 of a nucleic acid sequence, which encodes said polypeptide, is the only polypeptide expression product of said nucleic acid sequence that exhibits transgalactosylating activity. This may be evaluated by using the following techniques know to a person skilled in the art. The samples to be evaluated are subjected to SDS-PAGE and visualized using a dye appropriate for protein quantification, such as for example the Bio-Rad Criterion system. The gel is then scanned using appropriate densitometric scanner such as for example the Bio-Rad Criterion system and the resulting picture is ensured to be in the dynamic range. The bands corresponding to any variant/fragment derived from SEQ ID NO: 8 are quantified and the percentage of the polypeptides are calculated as: Percentage of polypeptide in question = polypeptide in ques-
25 tion/(sum of all polypeptides exhibiting transgalactosylating activity) *100.

The total number of polypeptides variants/fragments derived from SEQ ID NO:8 in the composition can be determined by detecting fragment derived from SEQ ID NO:8 by western blotting using a polyclonal antibody by methods know to
30 a person skilled in the art.

[0120] The polypeptide disclosed herein comprises at least two separate functional domains contained within the enzyme. Firstly, the polypeptide should contain a glycoside hydrolase catalytic core as described in the following. The catalytic core should belong to the GH-A clan of related glycoside hydrolase families. The GH-A clan is characterized by cleaving glycosidic bonds via a retaining mechanism and possesses a catalytic domain which is based on a TIM
35 barrel fold (Wierenga, 2001, FEBS Letters, 492(3), p 193-8). The catalytic domain contains two glutamic acid residues which act as proton donor and nucleophile, emanating from strands 4 and 7 of the barrel domain (Jenkins, 1995, FEBS Letters, 362(3), p 281-5). The overall structure of the TIM barrel is a (β/α) 8 fold consisting of 8 beta strands and 8 alpha-helices. In one aspect, the glycoside hydrolase catalytic core disclosed herein belong to either of the glycoside hydrolase families GH-2, and -35 which are all TIM-barrel enzymes belonging to the GH-A clan. In a further aspect, the glycoside
40 hydrolase catalytic core belong to family GH-2 or GH-35. In a further aspect, the glycoside hydrolase catalytic core belong to family GH-2. A common denominator is that these enzymes are so called retaining enzymes, so that the stereochemistry of the substrate is conserved in the product (Henrissat, 1997, Curr Opin Struct Biol, 7(5), 637-44).

[0121] In one aspect, the polypeptides disclosed herein have activity on carbohydrates bonds which has the $\beta(1\rightarrow4)$ conformation. This effectively put the enzymes into the IUBMB EC 3.2.1.23 class of β -galactosidases. This activity may
45 be, but is not confined to, determined by utilizing synthetic substrates such as para-nitrophenol- β -D-galactopyranoside (PNPG), ortho-nitrophenol- β -D-galactopyranoside (ONPG) or β -D-galactopyranoside with chromogenic aglycons (XGal). As an alternative way of determining whether an enzyme belong to the EC 3.2.1.23 class of β -galactosidases is to incubate with a substrate such as lactose and measure the release of glucose by a method such as enzymatic deter-
50 mination, HPLC, TLC or other methods known to persons skilled in the art.

[0122] In order to predict functional entities of polypeptides several available public repositories can be applied such as for example Pfam (Nucl. Acids Res. (2010) 38 (suppl 1): D211-D222. doi: 10.1093/nar/gkp985) and Interpro (Nucl. Acids Res. (2009) 37 (suppl 1): D211-D215. doi: 10.1093/nar/gkn785). It should be specified that when performing such analysis the analysis should be performed on the full length sequence of the polypeptide available from public repository
55 databases.

[0123] In a further aspect, a polypeptide containing one or more Pfam domains selected from: Glyco_hydro2N (PF02837), Glyco_hydro (PF00703), Glyco_hydro 2C (PF02836) and Bacterial Ig-like domain (group 4) (PF07532), is provided. In yet a further aspect, a polypeptide containing the Pfam domains Glyco_hydro2N (PF02837), Glyco_hydro (PF00703), Glyco_hydro 2C (PF02836) and Bacterial Ig-like domain (group 4) (PF07532), is provided. In yet a further

aspect, a polypeptide containing the Glyco_hydro2N (PF02837), Glyco_hydro (PF00703), and Glyco_hydro 2C (PF02836) domains which constitutes the catalytic domain of the polypeptide, is provided.

[0124] In a further aspect, a polypeptide as disclosed herein and having a ratio of transgalactosylating activity: β -galactosidase activity of at least 1, at least 2.5, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, or at least 12 as measured at a concentration of 100ppm in a milk-based assay at 37°C and 5 w/w% lactose after 15, 30 or 180 such as 180 minutes reaction, is provided. In a further aspect, the polypeptide is derived from *Bifidobacterium bifidum*.

[0125] In one aspect, the herein disclosed polypeptide(s) has a transgalactosylating activity such that more than 20%, more than 30%, more than 40%, up to 50% of the initial lactose is transgalactosylated as measured at a concentration of 100ppm in a milk-based assay at 37°C and 5 w/w% lactose after 15, 30 or 180 such as 180 minutes of reaction.

[0126] In a further aspect, the herein disclosed polypeptide(s) has a β -galactosidase activity such that less than 80%, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20% of the lactose has been hydrolysed as measured at a concentration of 100ppm in a milk-based assay at 37°C and 5 w/w% lactose after 15, 30 or 180 such as 180 minutes of reaction.

[0127] In one aspect, the β -galactosidase activity and/or the transgalactosylating activity are measured at a concentration of 100ppm corresponding to 2.13 LAU as specified in method 4.

[0128] In a further aspect, the herein disclosed polypeptide(s) has one or more of the following characteristics:

a) a ratio of transgalactosylating activity: β -galactosidase activity of at least of at least 1, at least 2.5, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, or at least 12 as measured at a concentration of 100ppm in a milk-based assay at 37°C and 5 w/w% lactose after 15, 30 or 180 such as 180 minutes reaction, and/or

b) has a transgalactosylating activity such that more than 20%, more than 30%, more than 40%, and up to 50% of the initial lactose has been transgalactosylated as measured at a concentration of 100ppm in a milk-based assay at 37°C and 5 w/w% lactose after 15, 30 or 180 such as 180 minutes of reaction.

[0129] In one aspect, a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues, is provided. In a further aspect, a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1 such as wherein said sequence identity is at least 95%, such as, e.g. at least 96%, at least 97%, at least 98%, at least 99% or at least 100% sequence identity, and wherein said polypeptide consists of at most 980 amino acid residues, is provided. In a further aspect, a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues, is provided. In yet a further aspect, a polypeptide wherein said polypeptide has at least 90% sequence identity with SEQ ID NO: 1, such as wherein said polypeptide has at least 90%, such as, e.g. at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with SEQ ID NO: 1 is provided. In another aspect, a polypeptide having at least 96.5% sequence identity to SEQ ID NO: 2 such as wherein said polypeptide has at least 97%, such as, e.g. at least 98% or at least 99% sequence identity with SEQ ID NO: 2. In one aspect, the polypeptides disclosed herein consist of at the most 975 amino acid residues, such as, e.g. at most 970 amino acid residues, such as at most 950 amino acid residues, such as at most 940 amino acid residues, at most 930 amino acid residues, at most 920 amino acid residues, at most 910 amino acid residues, at most 900 amino acid residues, at most 895 amino acid residues or at most 890 amino acid residues, is provided. In one aspect, a particular polypeptide consists of 887 or 965 amino acid residues, is provided. In one aspect, a polypeptide comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2 such as wherein said sequence identity is at least 98%, such as, e.g. at least 99% or at least 100% sequence identity, wherein said polypeptide consists of at most 975 amino acid residues, such as, e.g. at most 970 or at least 965 amino acid residues, is provided. In one aspect, a polypeptide comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2, wherein said polypeptide consists of at most 975 amino acid residues, is provided.

[0130] In a further preferred aspect, a polypeptide which comprises SEQ ID NO: 1, 2, 3, 4 or 5, is provided. In yet a preferred aspect, a polypeptide consisting of the amino acid sequence of SEQ ID NO: 1, 2, 3, 4, or 5, especially a polypeptide consisting of the amino acid sequence of SEQ ID NO: 1 or 2, is provided.

[0131] In a further aspect, a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3 such as wherein said sequence identity is at least 97%, such as, e.g. at least 98%, at least 99% or at least 100% sequence identity, wherein said polypeptide consists of at most 1300 amino acid residues, is provided.

[0132] In a further aspect, a polypeptide wherein said polypeptide has at least 98.5%, such as at least 99% or at least 99.5% sequence identity with SEQ ID NO: 5, is provided. In one aspect, such a polypeptide consists of at most 1290 amino acid residues, such as, e.g. at most 1280, at most 1270, at most 1260, at most 1250, at most 1240, at most 1230, at most 1220 or at most 1215 amino acid residues, is provided. In a preferred aspect, a polypeptide which consists of

1211 amino acid residues, is provided.

[0133] In a further aspect, a polypeptide wherein said polypeptide has at least 96% such as at least 97%, such as, e.g., at least 98% or at least 99% sequence identity with SEQ ID NO: 4, is provided. In one aspect, a polypeptide which consists of at most 1210 amino acid residues, such as, e.g. at most 1200, at most 1190, at most 1180, at most

1170, at most 1160, at most 1150 or at most 1145 amino acid residues, such as 1142 amino acid residues, is provided. **[0134]** In a further aspect, a polypeptide wherein said polypeptide has at least 96.5% such as at least 97%, such as, e.g., at least 98% or at least 99% sequence identity with SEQ ID NO: 3, is provided. In one aspect, a polypeptide which consists of at most 1130 amino acid residues, such as, e.g. at the most 1120, at the most 1110, at the most 1100, at the most 1090, at the most 1080, at the most 1070, at the most 1060, at the most 1050, at the most 1055 or at the most 1040 amino acid residues, is provided. In a preferred aspect, a polypeptide which consists of 1038 amino acid residues, is provided.

[0135] In a further aspect, the polypeptides disclosed herein has a ratio of transgalactosylation activity above 100% such as above 150%, 175% or 200%.

[0136] Proteins are generally comprised of one or more functional regions, commonly termed domains. The presence of different domains in varying combinations in different proteins gives rise to the diverse repertoire of proteins found in nature. One way of describing the domains are by the help of the Pfam database which is a large collection of protein domain families as described in "The Pfam protein families database": R.D. Finn, J. Mistry, J. Tate, P. Coghill, A. Heger, J.E. Pollington, O.L. Gavin, P. Gunasekaran, G. Ceric, K. Forslund, L. Holm, E.L. Sonnhammer, S.R. Eddy, A. Bateman Nucleic Acids Research (2010) Database Issue 38:D211-222. Each family is represented by multiple sequence alignments and hidden Markov models (HMMs). In a further aspect, the present inventors have found that the herein provided polypeptide(s) contains one or more of the Pfam domains Glyco_hydro2N (PF02837), Glyco_hydro (PF00703), Glyco_hydro 2C (PF02836) and Bacterial Ig-like domain (group 4) (PF07532). In one aspect, the herein provided polypeptide(s) contains Glyco_hydro2N (PF02837), Glyco_hydro (PF00703), Glyco_hydro 2C (PF02836) and Bacterial Ig-like domain (group 4) (PF07532).

[0137] In one aspect, the polypeptides have useful transgalactosylating activity over a range of pH of 4-9, such as 5-8, such as 5.5-7.5, such as 6.5-7.5.

[0138] The present invention encompasses polypeptides having a certain degree of sequence identity or sequence homology with amino acid sequence(s) defined herein or with a polypeptide having the specific properties defined herein. The present invention encompasses, in particular, peptides having a degree of sequence identity with any one of SEQ ID NO: 1, 2, 3, 4 or 5, defined below, or homologues thereof.

[0139] In one aspect, the homologous amino acid sequence and/or nucleotide sequence should provide and/or encode a polypeptide which retains the functional transgalactosylating activity and/or enhances the transgalactosylating activity compared to a polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5.

[0140] In the present context, a homologous sequence is taken to include an amino acid sequence which may be at least 66%, 70%, 75%, 78%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, identical to the subject sequence. Typically, the homologues will comprise the same active sites etc. as the subject amino acid sequence. Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity.

[0141] Thus, the present invention also encompasses variants, homologues and derivatives of any amino acid sequence of a protein or polypeptide as defined herein, particularly those of SEQ ID NO: 1, 2, 3, 4 or 5 defined below.

[0142] The sequences, particularly those of variants, homologues and derivatives of SEQ ID NO: 1, 2, 3, 4 or 5 defined below, may also have deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent substance. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the secondary binding activity of the substance is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine, glycine, alanine, asparagine, glutamine, serine, threonine, phenylalanine, and tyrosine.

[0143] The present invention also encompasses conservative substitution (substitution and replacement are both used herein to mean the interchange of an existing amino acid residue, with an alternative residue) that may occur i.e. like-for-like substitution such as basic for basic, acidic for acidic, polar for polar etc. Non-conservative substitution may also occur i.e. from one class of residue to another or alternatively involving the inclusion of unnatural amino acids such as ornithine (hereinafter referred to as Z), diaminobutyric acid ornithine

[0144] (hereinafter referred to as B), norleucine ornithine (hereinafter referred to as O), pyriylalanine, thienylalanine, naphthylalanine and phenylglycine.

[0145] Conservative substitutions that may be made are, for example within the groups of basic amino acids (Arginine, Lysine and Histidine), acidic amino acids (glutamic acid and aspartic acid), aliphatic amino acids (Alanine, Valine,

Leucine, Isoleucine), polar amino acids (Glutamine, Asparagine, Serine, Threonine), aromatic amino acids (Phenylalanine, Tryptophan and Tyrosine), hydroxyl amino acids (Serine, Threonine), large amino acids (Phenylalanine and Tryptophan) and small amino acids (Glycine, Alanine).

[0146] In one aspect, the polypeptide sequence used in the present invention is in a purified form. In one aspect, the polypeptide or protein for use in the present invention is in an isolated form.

[0147] In one aspect, the polypeptide of the present invention is recombinantly produced.

[0148] The variant polypeptides include a polypeptide having a certain percent, e.g., at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, of sequence identity with SEQ ID NO: 1 or 2.

[0149] The variant polypeptides include a polypeptide having a certain percent, e.g., at least 96%, 97%, 98%, or 99%, of sequence identity with SEQ ID NO: 3, 4 or 5.

[0150] In one aspect, the polypeptides disclosed herein comprises an amino acid sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to the amino acid sequence of the mature polypeptide encoded by the nucleotide sequence encoding the transgalactosylase contained in *Bifidobacterium bifidum* DSM20215 shown herein as SEQ ID NO: 22. All considerations and limitations relating to sequence identities and functionality discussed in terms of the SEQ ID NO: 1, 2, 3, 4 or 5 apply mutatis mutandis to sequence identities and functionality of these polypeptides and nucleotides.

[0151] In one aspect, the subject amino acid sequence is SEQ ID NO: 1, 2, 3, 4 or 5, and the subject nucleotide sequence preferably is SEQ ID NO: 9, 10, 11, 12 or 13.

[0152] In one aspect, the polypeptide is a fragment having one or more (several) amino acids deleted from the amino and/or carboxyl terminus of the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5; wherein the fragment has transgalactosylating activity.

[0153] In one aspect, a fragment contains at least 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, or 1000 amino acid residues

[0154] In a further aspect, the length of the polypeptide variant is 500 to 1300 amino acid residues. In a further aspect, the length of the polypeptide variant is 600 to 1300 amino acids. In a further aspect, the length of the polypeptide variant is 700 to 1300 amino acids. In a further aspect, the length of the polypeptide variant is 800 to 1300 amino acids. In a further aspect, the length of the polypeptide variant is 800 to 1300 amino acids.

Polypeptide variants of SEQ ID NO: 1, 2, 3, 4 or 5

[0155] In one aspect, a variant of SEQ ID NO: 1, 2, 3, 4 or 5 having a substitution at one or more positions which effects an altered property such as improved transgalactosylation, relative to SEQ ID NO: 1, 2, 3, 4 or 5, is provided. Such variant polypeptides are also referred to in this document for convenience as "variant polypeptide", "polypeptide variant" or "variant". In one aspect, the polypeptides as defined herein have an improved transgalactosylating activity as compared to the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5. In another aspect, the polypeptides as defined herein have an improved reaction velocity as compared to the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5.

[0156] In one aspect, the polypeptides and variants as defined herein exhibit enzyme activity. In one aspect, the polypeptides and the variant polypeptides described herein comprise transgalactosylation activity.

[0157] In one aspect, the ratio of transgalactosylating activity:β-galactosidase activity is at least 0.5, such as at least 1, such as at least 1.5, or such as at least 2 after 30 min. reaction such as above a concentration of 3% w/w initial lactose concentration.

[0158] In one aspect, the ratio of transgalactosylating activity:β-galactosidase activity is at least 2.5, such as at least 3, such as at least 4, such as at least 5, such as at least 6, such as at least 7, such as at least 8, such as at least 9, such as at least 10, such as at least 11, or such as at least 12 after 30 min. reaction such as above a concentration of 3% w/w initial lactose concentration.

[0159] In one aspect, the polypeptides and the variants as defined herein are derivable from microbial sources, in particular from a filamentous fungus or yeast, or from a bacterium. The enzyme may, e.g., be derived from a strain of *Agaricus*, e.g. *A. bisporus*; *Ascovaginospora*; *Aspergillus*, e.g. *A. niger*, *A. awamori*, *A. foetidus*, *A. japonicus*, *A. oryzae*; *Candida*; *Chaetomium*; *Chaetotomastia*; *Dictyostelium*, e.g. *D. discoideum*; *Kluyveromyces*, e.g. *K. fragilis*, *K. lactis*; *Mucor*, e.g. *M. javanicus*, *M. mucedo*, *M. subtilissimus*; *Neurospora*, e.g. *N. crassa*; *Rhizomucor*, e.g. *R. pusillus*; *Rhizopus*, e.g. *R. arrhizus*, *R. japonicus*, *R. stolonifer*; *Sclerotinia*, e.g. *S. libertiana*; *Torula*; *Torulopsis*; *Trichophyton*, e.g. *T. rubrum*; *Whetzelinia*, e.g. *W. sclerotiorum*; *Bacillus*, e.g. *B. coagulans*, *B. circulans*, *B. megaterium*, *B. novalis*, *B. subtilis*, *B. pumilus*, *B. stearothermophilus*, *B. thuringiensis*; *Bifidobacterium*, e.g. *B. longum*, *B. bifidum*, *B. animalis*; *Chryseobacterium*; *Citrobacter*, e.g. *C. freundii*; *Clostridium*, e.g. *C. perfringens*; *Diplodia*, e.g. *D. gossypina*; *Enterobacter*, e.g. *E. aerogenes*, *E. cloacae* *Edwardsiella*, *E. tarda*; *Erwinia*, e.g. *E. herbicola*; *Escherichia*, e.g. *E. coli*; *Klebsiella*, e.g. *K. pneumoniae*; *Miriococcum*; *Myrothesium*; *Mucor*; *Neurospora*, e.g. *N. crassa*; *Proteus*, e.g. *P. vulgaris*; *Providencia*, e.g. *P. stuartii*; *Pycnoporus*, e.g. *Pycnoporus cinnabarinus*, *Pycnoporus sanguineus*; *Ruminococcus*, e.g. *R. torques*; *Salmonella*, e.g. *S. typhimurium*; *Serratia*, e.g. *S. liquefasciens*, *S. marcescens*; *Shigella*, e.g. *S. flexneri*; *Strep-*

tomyces, e.g., *S. antibioticus*, *S. castaneoglobisporus*, *S. violeceoruber*; *Trametes*; *Trichoderma*, e.g., *T. reesei*, *T. viride*; *Yersinia*, e.g. *Y. enterocolitica*.

[0160] An isolated and/or purified polypeptide comprising a polypeptide or a variant polypeptide as defined herein is provided. In one embodiment, the variant polypeptide is a mature form of the polypeptide (SEQ ID NO: 1, 2, 3, 4 or 5).

In one aspect, the variants include a C-terminal domain.

[0161] In one aspect, a variant polypeptide as defined herein includes variants wherein between one and about 25 amino acid residues have been added or deleted with respect to SEQ ID NO: 1, 2, 3, 4 or 5. In one aspect, a variant polypeptide as defined herein includes variants wherein between one and 25 amino acid residues have been substituted, added or deleted with respect to SEQ ID NO: 1, 2, 3, 4 or 5. In one aspect, the variant has the amino acid sequence of SEQ ID NO: 1, 2, 3, 4 or 5, wherein any number between one and about 25 amino acids have been substituted. In a further aspect, the variant has the amino acid sequence of SEQ ID NO: 1, 2, 3, 4 or 5, wherein any number between three and twelve amino acids has been substituted. In a further aspect, the variant has the amino acid sequence of SEQ ID NO: 1, 2, 3, 4 or 5, wherein any number between five and nine amino acids has been substituted.

In one aspect, at least two, in another aspect at least three, and yet in another aspect at least five amino acids of SEQ ID NO: 1, 2, 3, 4 or 5 have been substituted.

In one aspect, the herein disclosed polypeptide(s) has the sequence of 1, 2, 3, 4 or 5.

In one aspect, the herein disclosed polypeptide(s) has the sequence of SEQ ID NO: 1, 2, 3, 4 or 5, wherein the 10, such as 9, such as 8, such as 7, such as 6, such as 5, such as 4, such as 3, such as 2, such as 1 amino acid in the N-terminal end are substituted and/or deleted. Enzymes and enzyme variants thereof can be characterized by their nucleic acid and primary polypeptide sequences, by three dimensional structural modeling, and/or by their specific activity. Additional characteristics of the polypeptide or polypeptide variants as defined herein include stability, pH range, oxidation stability, and thermostability, for example. Levels of expression and enzyme activity can be assessed using standard assays known to the artisan skilled in this field. In another aspect, variants demonstrate improved performance characteristics relative to the polypeptide with SEQ ID NO: 1, 2, 3, 4 or 5, such as improved stability at high temperatures, e.g., 65-85°C.

[0162] A polypeptide variant is provided as defined herein with an amino acid sequence having at least about 66%, 68%, 70%, 72%, 74%, 78%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% identity with the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5.

Nucleotides

[0163] In one aspect, the present invention relates to isolated polypeptides having transgalactosylating activity as stated above which are encoded by polynucleotides which hybridize under very low stringency conditions, preferably low stringency conditions, more preferably medium stringency conditions, more preferably medium-high stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with i) the nucleic acid sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding the mature polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5; ii) the cDNA sequence of i) or iii) the complementary strand of i) or ii), (J. Sambrook, E.F. Fritsch, and T. Maniatis, 1989, Molecular Cloning, A Laboratory Manual, 2d edition, Cold Spring Harbor, New York). A subsequence of SEQ ID NO: 9, 10, 11, 12 or 13 contains at least 100 contiguous nucleotides or preferably at least 200 contiguous nucleotides. Moreover, the subsequence may encode a polypeptide fragment which has lactase activity.

[0164] The nucleotide sequence of SEQ ID NO: 9, 10, 11, 12 or 13 or a subsequence thereof, as well as the amino acid sequence of SEQ ID NO: 1, 2, 3, 4 or 5 or a fragment thereof, may be used to design a nucleic acid probe to identify and clone DNA encoding polypeptides having transgalactosylase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 14, preferably at least 25, more preferably at least 35, and most preferably at least 70 nucleotides in length. It is, however, preferred that the nucleic acid probe is at least 100 nucleotides in length. For example, the nucleic acid probe may be at least 200 nucleotides, preferably at least 300 nucleotides, more preferably at least 400 nucleotides, or most preferably at least 500 nucleotides in length. Even longer probes may be used, e.g., nucleic acid probes which are at least 600 nucleotides, at least preferably at least 700 nucleotides, more preferably at least 800 nucleotides, or most preferably at least 900 nucleotides in length. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ³²P, ³H, ³⁵S, biotin, or avidin). Such probes are encompassed by the present invention.

[0165] A genomic DNA library prepared from such other organisms may, therefore, be screened for DNA which hybridizes with the probes described above and which encodes a polypeptide having lactase activity. Genomic or other DNA from such other organisms may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA which is homologous with SEQ ID NO: 9, 10, 11, 12

or 13 or a subsequence thereof, the carrier material is used in a Southern blot.

[0166] For purposes of the present invention, hybridization indicates that the nucleotide sequence hybridizes to a labelled nucleic acid probe corresponding to the nucleotide sequence shown in SEQ ID NO: 9, 10, 11, 12 or 13, its complementary strand, or a subsequence thereof, under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions can be detected using X-ray film.

[0167] In another preferred aspect, the nucleic acid probe is the mature polypeptide coding region of SEQ ID NO: 9, 10, 11, 12 or 13.

[0168] For long probes of at least 100 nucleotides in length, very low to very high stringency conditions are defined as prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 g/ml sheared and denatured salmon sperm DNA, and either 25% formamide for very low and low stringencies, 35% formamide for medium and medium-high stringencies, or 50% formamide for high and very high stringencies, following standard Southern blotting procedures for 12 to 24 hours optimally.

[0169] For long probes of at least 100 nucleotides in length, the carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS preferably at least at 45°C (very low stringency), more preferably at least at 50°C (low stringency), more preferably at least at 55°C (medium stringency), more preferably at least at 60°C (medium-high stringency), even more preferably at least at 65°C (high stringency), and most preferably at least at 70°C (very high stringency).

[0170] In a particular embodiment, the wash is conducted using 0.2X SSC, 0.2% SDS preferably at least at 45°C (very low stringency), more preferably at least at 50°C (low stringency), more preferably at least at 55°C (medium stringency), more preferably at least at 60°C (medium-high stringency), even more preferably at least at 65°C (high stringency), and most preferably at least at 70°C (very high stringency). In another particular embodiment, the wash is conducted using 0.1X SSC, 0.2% SDS preferably at least at 45°C (very low stringency), more preferably at least at 50°C (low stringency), more preferably at least at 55°C (medium stringency), more preferably at least at 60°C (medium-high stringency), even more preferably at least at 65°C (high stringency), and most preferably at least at 70°C (very high stringency).

[0171] For short probes which are about 15 nucleotides to about 70 nucleotides in length, stringency conditions are defined as prehybridization, hybridization, and washing post-hybridization at about 5°C to about 10°C below the calculated T_m using the calculation according to Bolton and McCarthy (1962, Proceedings of the National Academy of Sciences USA 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1X Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg of yeast RNA per ml following standard Southern blotting procedures.

[0172] For short probes which are about 15 nucleotides to about 70 nucleotides in length, the carrier material is washed once in 6X SSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6X SSC at 5°C to 10°C below the calculated T_m .

[0173] Under salt-containing hybridization conditions, the effective T_m is what controls the degree of identity required between the probe and the filter bound DNA for successful hybridization. The effective T_m may be determined using the formula below to determine the degree of identity required for two DNAs to hybridize under various stringency conditions.

[0174] Effective $T_m = 81.5 + 16.6(\log M[Na^+]) + 0.41(\%G+C) - 0.72(\% \text{ formamide})$ (See www.ndsu.nodak.edu/instruct/mcclean/plsc731/dna/dna6.htm)

The G+C content of SEQ ID NO: 10 is 42% and the G+C content of SEQ ID NO: 11 is 44%. For medium stringency, the formamide is 35% and the Na^+ concentration for 5X SSPE is 0.75 M.

[0175] Another relevant relationship is that a 1% mismatch of two DNAs lowers the T_m by 1.4°C. To determine the degree of identity required for two DNAs to hybridize under medium stringency conditions at 42°C, the following formula is used:

$$\% \text{ Homology} = 100 - [(Effective T_m - Hybridization Temperature)/1.4]$$

(See www.ndsu.nodak.edu/instruct/mcclean/plsc731/dna/dna6.htm)

[0176] The variant nucleic acids include a polynucleotide having a certain percent, e.g., 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%, of sequence identity with the nucleic acid encoding SEQ ID NO: 1, 2, 3, 4 or 5. In one aspect, a nucleic acid capable of encoding a polypeptide as disclosed herein, is provided. In a further aspect, the herein disclosed nucleic acid has a nucleic acid sequence which is at least 60%, such as at least 65%, such as at least 70%, such as at least 75%, such as at least 80%, such as at least 85%, such as at least 90%, such as at least 95%, such as at least 99% identical SEQ ID NO: 9, 10, 11, 12 or 13.

[0177] In one aspect, a plasmid comprising a nucleic acid as described herein, is provided.

In one aspect, an expression vector comprising a nucleic acid as described herein, or capable of expressing a polypeptide as described herein, is provided.

[0178] A nucleic acid complementary to a nucleic acid encoding any of the polypeptide variants as defined herein set

forth herein is provided. Additionally, a nucleic acid capable of hybridizing to the complement is provided. In another embodiment, the sequence for use in the methods and compositions described here is a synthetic sequence. It includes, but is not limited to, sequences made with optimal codon usage for expression in host organisms, such as yeast. The polypeptide variants as provided herein may be produced synthetically or through recombinant expression in a host cell, according to procedures well known in the art. In one aspect, the herein disclosed polypeptide(s) is recombinant polypeptide(s). The expressed polypeptide variant as defined herein optionally is isolated prior to use.

In another embodiment, the polypeptide variant as defined herein is purified following expression. Methods of genetic modification and recombinant production of polypeptide variants are described, for example, in U.S. Patent Nos. 7,371,552, 7,166,453; 6,890,572; and 6,667,065; and U.S. Published Application Nos. 2007/0141693; 2007/0072270; 2007/0020731; 2007/0020727; 2006/0073583; 2006/0019347; 2006/0018997; 2006/0008890; 2006/0008888; and 2005/0137111. The relevant teachings of these disclosures, including polypeptide-encoding polynucleotide sequences, primers, vectors, selection methods, host cells, purification and reconstitution of expressed polypeptide variants, and characterization of polypeptide variants as defined herein, including useful buffers, pH ranges, Ca²⁺ concentrations, substrate concentrations and enzyme concentrations for enzymatic assays, are herein incorporated by reference.

[0179] A nucleic acid sequence is provided encoding the protein of SEQ ID NO: 1, 2, 3, 4 or 5 or a nucleic acid sequence having at least about 66%, 68%, 70%, 72%, 74%, 78%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity with a nucleic acid encoding the protein of SEQ ID NO: 1, 2, 3, 4 or 5. In one embodiment, the nucleic acid sequence has at least about 60%, 66%, 68%, 70%, 72%, 74%, 78%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% sequence identity to the nucleic acid of SEQ ID NO: 9, 10, 11, 12 or 13.

Vectors

[0180] In one aspect, the invention relates to a vector comprising a polynucleotide. In one aspect, a bacterial cell comprises the vector. In some embodiments, a DNA construct comprising a nucleic acid encoding a variant is transferred to a host cell in an expression vector that comprises regulatory sequences operably linked to an encoding sequence. The vector may be any vector that can be integrated into a fungal host cell genome and replicated when introduced into the host cell. The FGSC Catalogue of Strains, University of Missouri, lists suitable vectors. Additional examples of suitable expression and/or integration vectors are provided in Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (2001); Bennett et al., *MORE GENE MANIPULATIONS IN FUNGI*, Academic Press, San Diego (1991), pp. 396-428; and U.S. Patent No. 5,874,276. Exemplary vectors include pFB6, pBR322, PUC18, pUC100 and pENTR/D, pDONTM201, pDONRTM221, pENTRTM, pGEM@3Z and pGEM@4Z. Exemplary for use in bacterial cells include pBR322 and pUC19, which permit replication in *E. coli*, and pE194, for example, which permits replication in *Bacillus*.

[0181] In some embodiments, a nucleic acid encoding a variant is operably linked to a suitable promoter, which allows transcription in the host cell. The promoter may be derived from genes encoding proteins either homologous or heterologous to the host cell. Suitable non-limiting examples of promoters include *cbh1*, *cbh2*, *egl1*, and *egl2* promoters. In one embodiment, the promoter is one that is native to the host cell. For example, when *P. saccharophila* is the host, the promoter is a native *P. saccharophila* promoter. An "inducible promoter" is a promoter that is active under environmental or developmental regulation. In another embodiment, the promoter is one that is heterologous to the host cell.

[0182] In some embodiments, the coding sequence is operably linked to a DNA sequence encoding a signal sequence. In another aspect, a representative signal peptide is SEQ ID NO: 27. A representative signal peptide is SEQ ID NO: 9 which is the native signal sequence of the *Bacillus subtilis* aprE precursor. In other embodiments, the DNA encoding the signal sequence is replaced with a nucleotide sequence encoding a signal sequence from other extra-cellular *Bacillus subtilis* pre-cursors. In one embodiment, the polynucleotide that encodes the signal sequence is immediately upstream and in-frame of the polynucleotide that encodes the polypeptide. The signal sequence may be selected from the same species as the host cell.

In additional embodiments, a signal sequence and a promoter sequence comprising a DNA construct or vector to be introduced into a fungal host cell are derived from the same source. In some embodiments, the expression vector also includes a termination sequence. In one embodiment, the termination sequence and the promoter sequence are derived from the same source. In another embodiment, the termination sequence is homologous to the host cell.

[0183] In some embodiments, an expression vector includes a selectable marker. Examples of suitable selectable markers include those that confer resistance to antimicrobial agents, e.g., hygromycin or phleomycin. Nutritional selective markers also are suitable and include *amdS*, *argB*, and *pyr4*. In one embodiment, the selective marker is the *amdS* gene, which encodes the enzyme acetamidase; it allows transformed cells to grow on acetamide as a nitrogen source. The use of an *A. nidulans amdS* gene as a selective marker is described in Kelley et al., *EMBO J.* 4: 475-479 (1985) and Penttilä et al., *Gene* 61: 155-164 (1987).

A suitable expression vector comprising a DNA construct with a polynucleotide encoding a variant may be any vector that is capable of replicating autonomously in a given host organism or integrating into the DNA of the host. In some

embodiments, the expression vector is a plasmid. In some embodiments, two types of expression vectors for obtaining expression of genes are contemplated. The first expression vector comprises DNA sequences in which the promoter, coding region, and terminator all originate from the gene to be expressed. In some embodiments, gene truncation is obtained by deleting undesired DNA sequences to leave the domain to be expressed under control of its own transcriptional and translational regulatory sequences. The second type of expression vector is preassembled and contains sequences required for high-level transcription and a selectable marker. In some embodiments, the coding region for a gene or part thereof is inserted into this general-purpose expression vector, such that it is under the transcriptional control of the expression construct promoter and terminator sequences. In some embodiments, genes or part thereof are inserted downstream of the strong *cbh1* promoter.

Expression hosts/ host cells

[0184] In a further aspect, a host cell comprising, preferably transformed with, a plasmid as described herein or an expression vector as described herein, is provided.

[0185] In a further aspect, a cell capable of expressing a polypeptide as described herein, is provided.

[0186] In one aspect, the host cell as described herein, or the cell as described herein is a bacterial, fungal or yeast cell.

[0187] In a further aspect, the host cell is selected from the group consisting of *Ruminococcus*, *Bifidobacterium*, *Lactococcus*, *Lactobacillus*, *Streptococcus*, *Leuconostoc*, *Escherichia*, *Bacillus*, *Streptomyces*, *Saccharomyces*, *Kluyveromyces*, *Candida*, *Torula*, *Torulopsis* and *Aspergillus*.

[0188] In a further aspect, the host cell is selected from the group consisting of *Ruminococcus hansenii*, *Bifidobacterium breve*, *Bifidobacterium longum*, *Bifidobacterium infantis*, *Bifidobacterium bifidum* and *Lactococcus lactis*.

[0189] In another embodiment, suitable host cells include a Gram positive bacterium selected from the group consisting of *Bacillus subtilis*, *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. coagulans*, *B. circulans*, *B. lautus*, *B. thuringiensis*, *Streptomyces lividans*, or *S. murinus*; or a Gram negative bacterium, wherein said Gram negative bacterium is *Escherichia coli* or a *Pseudomonas* species. In one aspect, the host cell is a *B. subtilis* or *B. licheniformis*. In one embodiment, the host cell is *B. subtilis*, and the expressed protein is engineered to comprise a *B. subtilis* signal sequence, as set forth in further detail below. In one aspect, the host cell expresses the polynucleotide as set out in the claims.

[0190] In some embodiments, a host cell is genetically engineered to express a polypeptide variant as defined herein with an amino acid sequence having at least about 66%, 68%, 70%, 72%, 74%, 78%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% identity with the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5. In some embodiments, the polynucleotide encoding a polypeptide variant as defined herein will have a nucleic acid sequence encoding the protein of SEQ ID NO: 1, 2, 3, 4 or 5 or a nucleic acid sequence having at least about 66%, 68%, 70%, 72%, 74%, 78%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity with a nucleic acid encoding the protein of SEQ ID NO: 1, 2, 3, 4 or 5. In one embodiment, the nucleic acid sequence has at least about 60%, 66%, 68%, 70%, 72%, 74%, 78%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% sequence identity to the nucleic acid of SEQ ID NO: 9, 10, 11, 12 or 13.

Methods for producing polypeptides

[0191] In a further aspect, a method of expressing a polypeptide as described herein comprises obtaining a host cell or a cell as described herein and expressing the polypeptide from the cell or host cell, and optionally purifying the polypeptide.

[0192] An expression characteristic means an altered level of expression of the variant, when the variant is produced in a particular host cell. Expression generally relates to the amount of active variant that is recoverable from a fermentation broth using standard techniques known in this art over a given amount of time. Expression also can relate to the amount or rate of variant produced within the host cell or secreted by the host cell. Expression also can relate to the rate of translation of the mRNA encoding the variant polypeptide.

Transformation, expression and culture of host cells

[0193] Introduction of a DNA construct or vector into a host cell includes techniques such as transformation; electroporation; nuclear microinjection; transduction; transfection, e.g., lipofection mediated and DEAE-Dextrin mediated transfection; incubation with calcium phosphate DNA precipitate; high velocity bombardment with DNA-coated microprojectiles; and protoplast fusion. General transformation techniques are known in the art. See, e.g., Ausubel *et al.* (1987), *supra*, chapter 9; Sambrook *et al.* (2001), *supra*; and Campbell *et al.*, *Curr. Genet.* 16: 53-56 (1989). The expression of heterologous protein in *Trichoderma* is described, for example, in U.S. Patent No. 6,022,725; U.S. Patent No. 6,268,328; Harkki *et al.*, *Enzyme Microb. Technol.* 13: 227-233 (1991); Harkki *et al.*, *BioTechnol.* 7: 596-603 (1989); EP 244,234;

and EP 215,594. In one embodiment, genetically stable transformants are constructed with vector systems whereby the nucleic acid encoding a variant is stably integrated into a host cell chromosome. Transformants are then purified by known techniques.

[0194] In one non-limiting example, stable transformants including an *amdS* marker are distinguished from unstable transformants by their faster growth rate and the formation of circular colonies with a smooth, rather than ragged outline on solid culture medium containing acetamide. Additionally, in some cases a further test of stability is conducted by growing the transformants on solid non-selective medium, e.g., a medium that lacks acetamide, harvesting spores from this culture medium and determining the percentage of these spores that subsequently germinate and grow on selective medium containing acetamide. Other methods known in the art may be used to select transformants.

Identification of activity

[0195] To evaluate the expression of a variant in a host cell, assays can measure the expressed protein, corresponding mRNA, or β -galactosidase activity. For example, suitable assays include Northern and Southern blotting, RT-PCR (reverse transcriptase polymerase chain reaction), and *in situ* hybridization, using an appropriately labeled hybridizing probe. Suitable assays also include measuring activity in a sample. Suitable assays of the activity of the variant include, but are not limited to, ONPG based assays or determining glucose in reaction mixtures such for example described in the methods and examples herein.

Methods for purifying herein disclosed polypeptides

[0196] In general, a variant produced in cell culture is secreted into the medium and may be purified or isolated, e.g., by removing unwanted components from the cell culture medium. In some cases, a variant may be recovered from a cell lysate. In such cases, the enzyme is purified from the cells in which it was produced using techniques routinely employed by those of skill in the art. Examples include, but are not limited to, affinity chromatography, ion-exchange chromatographic methods, including high resolution ion-exchange, hydrophobic interaction chromatography, two-phase partitioning, ethanol precipitation, reverse phase HPLC, chromatography on silica or on a cation-exchange resin, such as DEAE, chromatofocusing, SDS-PAGE, ammonium sulfate precipitation, and gel filtration using Sephadex G-75, for example. Depending on the intended use the herein disclosed polypeptide(s) may for example be either freeze-dried or prepared in a solution. In one aspect, the herein disclosed polypeptide(s) is freeze-dried form. In another aspect, the herein disclosed polypeptide(s) is in solution.

Compositions, application and use

Methods for immobilising and formulation of the herein disclosed polypeptides

[0197] The polypeptide compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the polypeptide composition may be in the form of a granulate or a microgranulate. The polypeptide to be included in the composition may be stabilized in accordance with methods known in the art. Examples are given below of preferred uses of the polypeptides or polypeptide compositions of the invention.

[0198] In one aspect, disclosed herein is a method for producing a food product by treating a substrate comprising lactose with a polypeptide as described herein.

[0199] In one aspect, disclosed herein is a method for producing a dairy product by treating a milk-based substrate comprising lactose with a polypeptide as described herein.

[0200] In one aspect, the substrate comprising lactose is further treated with a hydrolysing beta-galactosidase.

[0201] The enzyme preparation, such as in the form of a food ingredient prepared according to the present invention, may be in the form of a solution or as a solid - depending on the use and/or the mode of application and/or the mode of administration. The solid form can be either as a dried enzyme powder or as a granulated enzyme.

Examples of dry enzyme formulations include spray dried products, mixer granulation products, layered products such as fluid bed granules, extruded or pelletized granules prilled products, lyophilized products.

[0202] The enzyme preparation, such as in the form of a food ingredient prepared according to the present invention, may be in the form of a solution or as a solid - depending on the use and/or the mode of application and/or the mode of administration. The solid form can be either as a dried enzyme powder or as a granulated enzyme.

[0203] In one aspect, a composition preferably a food composition, more preferably a dairy product comprising a cell or a polypeptide as described herein, is provided.

[0204] Furthermore, disclosed herein is a composition comprising at least 5%, such as e.g. 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50% w/w of one or more polypeptide(s) as disclosed herein based on the total amount of polypep-

tides in the composition having at least 70%, e.g. such as 72%, 74%, 74%, 78%, 80%, 82%, 84%, 86%, 88%, 90% sequence identity with SEQ ID NO: 22. This may be evaluated by using the following techniques known to a person skilled in the art. The samples to be evaluated are subjected to SDS-PAGE and visualized using a dye appropriate for protein quantification, such as for example the Bio-Rad Criterion system. The gel is then scanned using appropriate densitometric scanner such as for example the Bio-Rad Criterion system and the resulting picture is ensured to be in the dynamic range. The bands corresponding to any variant/fragment derived from SEQ ID NO: 8 are quantified and the percentage of the polypeptides are calculated as: Percentage of polypeptide in question = polypeptide in question / (sum of all polypeptides exhibiting transgalactosylating activity) * 100. The total number of polypeptides variants/fragments derived from SEQ ID NO: 8 in the composition can be determined by detecting fragment derived from SEQ ID NO: 8 by western blotting using a polyclonal antibody by methods known to a person skilled in the art.

[0205] In one aspect, the composition according to the present invention comprises one or more polypeptide(s) selected from the group consisting of a polypeptide consisting of SEQ ID NO: 1, 2, 3, 4 and 5. In a further aspect, the composition comprises one or more polypeptide(s) selected from the group consisting of a polypeptide consisting of SEQ ID NO: 1, 2 and 3. In yet a further aspect, the composition comprises one or more polypeptide(s) selected from the group consisting of a polypeptide consisting of SEQ ID NO: 1 and 2.

[0206] In one aspect the invention provides an enzyme complex preparation comprising the enzyme complex according to the invention, an enzyme carrier and optionally a stabilizer and/or a preservative.

[0207] In yet a further aspect of the invention, the enzyme carrier is selected from the group consisting of glycerol or water.

[0208] In a further aspect, the preparation/composition comprises a stabilizer. In one aspect, the stabilizer is selected from the group consisting of inorganic salts, polyols, sugars and combinations thereof. In one aspect, the stabilizer is an inorganic salt such as potassium chloride. In another aspect, the polyol is glycerol, propylene glycol, or sorbitol. In yet another aspect, the sugar is a small-molecule carbohydrate, in particular any of several sweet-tasting ones such as glucose, galactose, fructose and saccharose.

[0209] In yet a further aspect, the preparation comprises a preservative. In one aspect, the preservative is methyl paraben, propyl paraben, benzoate, sorbate or other food approved preservatives or a mixture thereof.

[0210] The method of the invention can be practiced with immobilized enzymes, e.g. an immobilized lactase or other galactooligosaccharide producing enzymes. The enzyme can be immobilized on any organic or inorganic support. Exemplary inorganic supports include alumina, celite, Dowex-1-chloride, glass beads and silica gel. Exemplary organic supports include DEAE-cellulose, alginate hydrogels or alginate beads or equivalents. In various aspects of the invention, immobilization of the lactase can be optimized by physical adsorption on to the inorganic support. Enzymes used to practice the invention can be immobilized in different media, including water, Tris-HCl buffer and phosphate buffered solution. The enzyme can be immobilized to any type of substrate, e.g. filters, fibers, columns, beads, colloids, gels, hydrogels, meshes and the like.

[0211] In one aspect, a method for producing a dairy product by treating a milk-based substrate comprising lactose with a polypeptide as described herein is provided. In a further aspect, a method for producing a dairy product by treating a milk-based substrate comprising lactose with a polypeptide having a relative transgalactosylation activity above 60%, such as above 70%, such as above 75% after 15 min. reaction, is provided. In one aspect, the relative transgalactosylation activity is above 3 after 30 min. reaction. In a further aspect, the relative transgalactosylation activity is above 6 after 30 min. reaction. In yet a further aspect, the relative transgalactosylation activity is above 12 after 30 min. reaction. In one aspect, a method is provided, wherein the treatment with a polypeptide as described herein takes place at an optimal temperature for the activity of the enzyme. In a further aspect, the polypeptide is added to the milk-based substrate at a concentration of 0.01-1000 ppm. In yet a further aspect, the polypeptide is added to the milk-based substrate at a concentration of 0.1-100 ppm. In a further aspect, the polypeptide is added to the milk-based substrate at a concentration of 1-10 ppm. In one aspect, a method further comprising fermenting a substrate such as a dairy product with a microorganism, is provided. In a further aspect, the dairy product is yogurt. In a further aspect, the treatment with the polypeptide and the microorganism is performed essentially at the same time. In one aspect, the polypeptide and the microorganism are added to the milk-based substrate essentially at the same time.

In one aspect, a dairy product comprising a cell or a polypeptide as described herein, is provided. In one aspect, the polypeptide as defined herein is added in a concentration of 0.01-1000 ppm. In one aspect, a dairy product comprising an inactivated polypeptide as defined herein, is provided. In one aspect, a dairy product comprising an inactivated polypeptide as defined herein in a concentration of 0.01-1000 ppm, is provided. In one aspect, a dairy product comprising GOS formed in situ by a polypeptide as defined herein, is provided. In one aspect, a dairy product comprising a cell as defined herein, is provided.

A dairy product as described herein may be, e.g., skim milk, low fat milk, whole milk, cream, UHT milk, milk having an extended shelf life, a fermented milk product, cheese, yoghurt, butter, dairy spread, butter milk, acidified milk drink, sour cream, whey based drink, ice cream, condensed milk, dulce de leche or a flavoured milk drink. A dairy product may be manufactured by any method known in the art.

[0212] A dairy product may additionally comprise non-milk components, e.g. vegetable components such as, e.g., vegetable oil, vegetable protein, and/or vegetable carbohydrates. Dairy products may also comprise further additives such as, e.g., enzymes, flavouring agents, microbial cultures such as probiotic cultures, salts, sweeteners, sugars, acids, fruit, fruit juices, or any other component known in the art as a component of, or additive to, a dairy product.

[0213] In one embodiment of the invention, one or more milk components and/or milk fractions account for at least 50% (weight/weight), such as at least 70%, e.g. at least 80%, preferably at least 90%, of the dairy product.

[0214] In one embodiment of the invention, one or more milk-based substrates having been treated with an enzyme as defined herein having transgalactosylating activity account for at least 50% (weight/weight), such as at least 70%, e.g. at least 80%, preferably at least 90%, of the dairy product.

[0215] In one embodiment of the invention, the dairy product is a dairy product which is not enriched by addition of pre-produced galacto-oligosaccharides.

[0216] In one embodiment of the invention, the polypeptide-treated milk-based substrate is not dried before being used as an ingredient in the dairy product.

[0217] In one embodiment of the invention, the dairy product is ice cream. In the present context, ice cream may be any kind of ice cream such as full fat ice cream, low fat ice cream, or ice cream based on yoghurt or other fermented milk products. Ice cream may be manufactured by any method known in the art.

[0218] In one embodiment of the invention, the dairy product is milk or condensed milk.

In one embodiment of the invention, the dairy product is UHT milk. UHT milk in the context of the present invention is milk which has been subjected to a sterilization procedure which is intended to kill all microorganisms, including the bacterial spores. UHT (ultra high temperature) treatment may be, e.g., heat treatment for 30 seconds at 130°C, or heat treatment for one second at 145°C.

[0219] In one preferred embodiment of the invention, the dairy product is ESL milk. ESL milk in the present context is milk which has an extended shelf life due to microfiltration and/or heat treatment and which is able to stay fresh for at least 15 days, preferably for at least 20 days, on the store shelf at 2-5°C.

[0220] In another preferred embodiment of the invention, the dairy product is a fermented dairy product, e.g., yoghurt.

[0221] The microorganisms used for most fermented milk products are selected from the group of bacteria generally referred to as lactic acid bacteria. As used herein, the term "lactic acid bacterium" designates a gram-positive, micro-aerophilic or anaerobic bacterium, which ferments sugars with the production of acids including lactic acid as the predominantly produced acid, acetic acid and propionic acid. The industrially most useful lactic acid bacteria are found within the order "Lactobacillales" which includes *Lactococcus* spp., *Streptococcus* spp., *Lactobacillus* spp., *Leuconostoc* spp., *Pseudoleuconostoc* spp., *Pediococcus* spp., *Brevibacterium* spp., *Enterococcus* spp. and *Propionibacterium* spp. Additionally, lactic acid producing bacteria belonging to the group of anaerobic bacteria, bifidobacteria, i.e. *Bifidobacterium* spp., which are frequently used as food cultures alone or in combination with lactic acid bacteria, are generally included in the group of lactic acid bacteria. Lactic acid bacteria are normally supplied to the dairy industry either as frozen or freeze-dried cultures for bulk starter propagation or as so-called "Direct Vat Set" (DVS) cultures, intended for direct inoculation into a fermentation vessel or vat for the production of a fermented dairy product. Such cultures are in general referred to as "starter cultures" or "starters".

[0222] Commonly used starter culture strains of lactic acid bacteria are generally divided into mesophilic organisms having optimum growth temperatures at about 30°C and thermophilic organisms having optimum growth temperatures in the range of about 40 to about 45°C. Typical organisms belonging to the mesophilic group include *Lactococcus lactis*, *Lactococcus lactis* subsp. *cremoris*, *Leuconostoc mesenteroides* subsp. *cremoris*, *Pseudoleuconostoc mesenteroides* subsp. *cremoris*, *Pediococcus pentosaceus*, *Lactococcus lactis* subsp. *lactis* biovar. *diacetylactis*, *Lactobacillus casei* subsp. *casei* and *Lactobacillus paracasei* subsp. *paracasei*. Thermophilic lactic acid bacterial species include as examples *Streptococcus thermophilus*, *Enterococcus faecium*, *Lactobacillus delbrueckii* subsp. *lactis*, *Lactobacillus helveticus*, *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Lactobacillus acidophilus*. Also the anaerobic bacteria belonging to the genus *Bifidobacterium* including *Bifidobacterium bifidum*, *Bifidobacterium animalis* and *Bifidobacterium longum* are commonly used as dairy starter cultures and are generally included in the group of lactic acid bacteria. Additionally, species of *Propionibacteria* are used as dairy starter cultures, in particular in the manufacture of cheese. Additionally, organisms belonging to the *Brevibacterium* genus are commonly used as food starter cultures. Another group of microbial starter cultures are fungal cultures, including yeast cultures and cultures of filamentous fungi, which are particularly used in the manufacture of certain types of cheese and beverage. Examples of fungi include *Penicillium roqueforti*, *Penicillium candidum*, *Geotrichum candidum*, *Torula kefir*, *Saccharomyces kefir* and *Saccharomyces cerevisiae*.

[0223] In one embodiment of the present invention, the microorganism used for fermentation of the milk-based substrate is *Lactobacillus casei* or a mixture of *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus*.

[0224] Fermentation processes to be used in a method of the present invention are well known and the person of skill in the art will know how to select suitable process conditions, such as temperature, oxygen, amount and characteristics of microorganism/s, additives such as e.g. carbohydrates, flavours, minerals, enzymes, and process time. Obviously, fermentation conditions are selected so as to support the achievement of the present invention.

As a result of fermentation, pH of the milk-based substrate will be lowered. The pH of a fermented dairy product of the invention may be, e.g., in the range 3.5-6, such as in the range 3.5-5, preferably in the range 3.8-4.8.

[0225] In one aspect, a method of using the polypeptides or using any one or more of the above mentioned cell types for producing oligosaccharides, is provided. The oligosaccharides comprise, but are not limited to fructooligo-saccharides,

[0226] In one embodiment of the invention, the oligosaccharides are produced by incubating the cell expressing the polypeptide in a medium that comprises a disaccharide substrate such as for example lactulose, trehalose, rhamnose, maltose, sucrose, lactose, or cellobiose. The incubation is carried out under conditions where oligosaccharides are produced. The cells may be part of a product selected from the group consisting of yoghurt, cheese, fermented milk products, dietary supplements, and probiotic comestible products. Alternatively, the oligosaccharides can be recovered and subsequently be added to the product of interest before or after its preparation.

[0227] In one aspect, the use of a herein disclosed cell for producing a product selected from the group consisting of yoghurt, cheese, fermented milk product, dietary supplement and probiotic comestible product, is provided.

[0228] In one aspect, the polypeptides described herein may be used to prepare cheese products and in methods for making the cheese products. Cheese products may e.g. be selected from the group consisting of cream cheese, cottage cheese, and process cheese. By adding polypeptides the cheeses may contain significantly increased levels of galacto-oligosaccharides and reduced levels of lactose. In one aspect, the lactose levels in the final cheese product may be reduced by at least about 25 percent, preferably at least about 50 percent, and more preferably at least about 75 percent. The polypeptides may be used to reduce lactose in cheese products to less than about 1 gram per serving, an amount that can be tolerated by most lactose-intolerant individuals.

[0229] The cheese products provided herein are nutritionally-enhanced cheese products having increased soluble fiber content, reduced caloric content, excellent organoleptic properties, improved texture, and flavor. Further, the polypeptides described herein may reduce the glycemic index of the cheese products because GOS are more slowly absorbed than lactose or its hydrolysis products. Finally, the polypeptides may reduce the cost of production of cheese products, particularly cream cheese products, because GOS surprisingly provide improved texture to the cream cheese product, thus permitting reduced use of stabilizers, or by allowing for increased moisture content without syneresis.

[0230] In a further aspect, a composition comprising a polypeptide as described herein and a carbohydrate substrate, is provided. In a further aspect, the carbohydrate substrate is a disaccharide. In a further aspect, the disaccharide is for example lactulose, trehalose, rhamnose, maltose, sucrose, lactose or cellobiose. In yet a further aspect, the carbohydrate substrate is lactose. The composition is prepared such that oligosaccharides are produced. The polypeptide as described herein may be part of a product selected from the group consisting of yoghurt, cheese, fermented milk products, dietary supplements, and probiotic comestible products. In one aspect, a composition comprising a polypeptide as described herein and a stabilizer, is provided. Examples of stabilizers is e.g., a polyol such as, e.g., glycerol or propylene glycol, a sugar or a sugar alcohol, lactic acid, boric acid, or a boric acid derivative (e.g., an aromatic borate ester).

[0231] In one aspect, the use of a transgalactosylating polypeptide as disclosed herein or a cell as disclosed herein, for producing galacto-oligosaccharides, is provided. In one aspect, the use of a transgalactosylating polypeptide as disclosed herein or a cell as disclosed herein, for producing galacto-oligosaccharides to be part of a product selected from the group consisting of yoghurt, cheese, fermented dairy products, dietary supplements and probiotic comestible products, is provided. In one aspect, the product is yoghurt, cheese, or fermented dairy products. In one aspect, the use of a transgalactosylating polypeptide as disclosed herein or a cell as disclosed herein, for producing galacto-oligosaccharides to enhance the growth of Bifidobacterium, is provided. In one aspect, the use of a transgalactosylating polypeptide as disclosed herein or a cell as disclosed herein, for producing galacto-oligosaccharides to enhance the growth of Bifidobacterium in a mixed culture fermentation, is provided.

[0232] In one aspect, a process for producing a transgalactosylating polypeptide as disclosed herein, comprising culturing a cell as disclosed herein in a suitable culture medium under conditions permitting expression of said polypeptide, and recovering the resulting polypeptide from the culture, is provided. A process for producing galacto-oligosaccharides, comprising contacting of an polypeptide of as disclosed herein or a cell as disclosed herein with a milk-based solution comprising lactose, is provided.

[0233] Addition of oligosaccharides may enhance growth of either Bifidobacterium alone or of Bifidobacterium in a mixed culture.

[0234] The treatment of milk products with enzymes that converts lactose into monosaccharides or GOS have several advantages. First the products can be consumed by people with lactose intolerance that would otherwise exhibit symptoms such as flatulence and diarrhea. Secondly, dairy products treated with lactase will have a higher sweetness than similar untreated products due to the higher perceived sweetness of glucose and galactose compared to lactose. This effect is particularly interesting for applications such as yoghurt and ice-cream where high sweetness of the end product is desired and this allows for a net reduction of carbohydrates in the consumed product. Thirdly, in ice-cream production a phenomenon termed sandiness is often seen, where the lactose molecules crystallizes due to the relative low solubility of the lactose. When lactose is converted into monosaccharides or GOS the mouth feeling of the ice-cream is much

improved over the non-treated products. The presence of a sandy feeling due to lactose crystallization can be eliminated and the raw material costs can be decreased by replacement of skimmed milk powder by whey powder. The main effects of the enzymatic treatment were increased sweetness.

[0235] In one aspect, the transgalactosylating polypeptide(s) as disclosed herein may be used together with other enzymes such as proteases such as chymosin or rennin, lipases such as phospholipases, amylases, transferases, and lactases. In one aspect, the transgalactosylating polypeptide(s) as disclosed herein may be used together with lactase. This may especially be useful when there is a desire to reduce residual lactose after treatment with the transgalactosylating polypeptide(s) as disclosed herein especially at low lactose levels. A lactase in the context of the present invention is any glycoside hydrolase having the ability to hydrolyse the disaccharide lactose into constituent galactose and glucose monomers. The group of lactases comprises but is not limited to enzymes assigned to subclass EC 3.2.1.108. Enzymes assigned to other subclasses, such as, e.g., EC 3.2.1.23, may also be lactases in the context of the present invention. A lactase in the context of the invention may have other activities than the lactose hydrolysing activity, such as for example a transgalactosylating activity. In the context of the invention, the lactose hydrolysing activity of the lactase may be referred to as its lactase activity or its beta-galactosidase activity. Enzymes having lactase activity to be used in a method of the present invention may be of animal, of plant or of microbial origin. Preferred enzymes are obtained from microbial sources, in particular from a filamentous fungus or yeast, or from a bacterium. The enzyme may, e.g., be derived from a strain of *Agaricus*, e.g. *A. bisporus*; *Ascovaginospora*; *Aspergillus*, e.g. *A. niger*, *A. awamori*, *A. foetidus*, *A. japonicus*, *A. oryzae*; *Candida*; *Chaetomium*; *Chaetotomastia*; *Dictyostelium*, e.g. *D. discoideum*; *Kluyveromyces*, e.g. *K. fragilis*, *K. lactis*; *Mucor*, e.g. *M. javanicus*, *M. mucedo*, *M. subtilissimus*; *Neurospora*, e.g. *N. crassa*; *Rhizomucor*, e.g. *R. pusillus*; *Rhizopus*, e.g. *R. arrhizus*, *R. japonicus*, *R. stolonifer*; *Sclerotinia*, e.g. *S. libertiana*; *Torula*; *Torulopsis*; *Trichophyton*, e.g. *T. rubrum*; *Wetzelinia*, e.g. *W. sclerotiorum*; *Bacillus*, e.g. *B. coagulans*, *B. circulans*, *B. megaterium*, *B. novalis*, *B. subtilis*, *B. pumilus*, *B. stearothermophilus*, *B. thuringiensis*; *Bifidobacterium*, e.g. *B. longum*, *B. bifidum*, *B. animalis*; *Chryseobacterium*; *Citrobacter*, e.g. *C. freundii*; *Clostridium*, e.g. *C. perfringens*; *Diplodia*, e.g. *D. gossypina*; *Enterobacter*, e.g. *E. aerogenes*, *E. cloacae* *Edwardsiella*, *E. tarda*; *Erwinia*, e.g. *E. herbicola*; *Escherichia*, e.g. *E. coli*; *Klebsiella*, e.g. *K. pneumoniae*; *Miriococcum*; *Myrothesium*; *Mucor*; *Neurospora*, e.g. *N. crassa*; *Proteus*, e.g. *P. vulgaris*; *Providencia*, e.g. *P. stuartii*; *Pycnoporus*, e.g. *Pycnoporus cinnabarinus*, *Pycnoporus sanguineus*; *Ruminococcus*, e.g. *R. torques*; *Salmonella*, e.g. *S. typhimurium*; *Serratia*, e.g. *S. liquefaciens*, *S. marcescens*; *Shigella*, e.g. *S. flexneri*; *Streptomyces*, e.g. *S. antibioticus*, *S. castaneoglobisporus*, *S. violeceoruber*; *Trametes*; *Trichoderma*, e.g. *T. reesei*, *T. viride*; *Yersinia*, e.g. *Y. enterocolitica*. In one embodiment, the lactase is an intracellular component of microorganisms like *Kluyveromyces* and *Bacillus*. *Kluyveromyces*, especially *K. fragilis* and *K. lactis*, and other fungi such as those of the genera *Candida*, *Torula* and *Torulopsis*, are a common source of fungal lactases, whereas *B. coagulans* and *B. circulans* are well known sources for bacterial lactases. Several commercial lactase preparations derived from these organisms are available such as Lactozym.RTM. (available from Novozymes, Denmark), HA-Lactase (available from Chr. Hansen, Denmark) and Maxilact.RTM. (available from DSM, the Netherlands), all from *K. lactis*. All these lactases are so called neutral lactases having a pH optimum between pH 6 and pH 8. When such lactases are used in the production of, e.g., low-lactose yoghurt, the enzyme treatment will either have to be done in a separate step before fermentation or rather high enzyme dosages have to be used, because their activity drop as the pH decreases during fermentation. Also, these lactases are not suitable for hydrolysis of lactose in milk performed at high temperature, which would in some cases be beneficial in order to keep the microbial count low and thus ensure good milk quality. In one embodiment, the enzyme is a lactase from a bacterium, e.g. from the family *Bifidobacteriaceae*, such as from the genus *Bifidobacterium* such as the lactase described in WO 2009/071539.

FURTHER ASPECTS OF THE INVENTION

[0236]

Aspect 1. A polypeptide having transgalactosylating activity, which comprises an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, and wherein said polypeptide, when being an expression product in a *Bacillus subtilis* strain BG3594 of a nucleic acid sequence, which encodes said polypeptide, is the only polypeptide expression product of said nucleic acid sequence that exhibits transgalactosylating activity.

Aspect 2. A polypeptide having transgalactosylating activity selected from the group consisting of:

- a. a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues,
- b. a polypeptide comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2, wherein said polypeptide consists of at most 975 amino acid residues,

c. a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues,

d. a polypeptide encoded by a polynucleotide that hybridizes under at least low stringency conditions with i) the nucleic acid sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding the polypeptide of SEQ ID NO: 1, 2, 3, 4, or 5; or ii) the complementary strand of i),

e. a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 70% identity to the nucleotide sequence encoding for the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5 or the nucleotide sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding a mature polypeptide, and

f. a polypeptide comprising a deletion, insertion and/or conservative substitution of one or more amino acid residues of SEQ ID NO: 1, 2, 3, 4 or 5.

Aspect 3. A polypeptide having transgalactosylating activity selected from the group consisting of:

a. a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues,

b. a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues,

c. a polypeptide encoded by a polynucleotide that hybridizes under at least low stringency conditions with i) the nucleic acid sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding the polypeptide of SEQ ID NO: 1, 2, 3, 4, or 5; or ii) the complementary strand of i),

d. a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 70% identity to the nucleotide sequence encoding for the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5 or the nucleotide sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding a mature polypeptide, and

e. a polypeptide comprising a deletion, insertion and/or conservative substitution of one or more amino acid residues of SEQ ID NO: 1, 2, 3, 4 or 5.

Aspect 4. The polypeptide according to any one of the preceding aspects, wherein said polypeptide, when being an expression product in a *Bacillus subtilis* strain BG3594 of a nucleic acid sequence, which encodes said polypeptide, is the only polypeptide expression product of said nucleic acid sequence that exhibits transgalactosylating activity.

Aspect 5. The polypeptide according to any one of the preceding aspects comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues.

Aspect 6. The polypeptide according to any one of the preceding aspects comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues.

Aspect 7. The polypeptide according to any of aspects 1, 2 and 4 comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2, wherein said polypeptide consists of at most 975 amino acid residues.

Aspect 8. A polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues.

Aspect 9. The polypeptide according to aspect 8, wherein said polypeptide has transgalactosylating activity.

Aspect 10. The polypeptide according to any one of aspects 8-9, wherein said polypeptide, when being an expression product in a *Bacillus subtilis* strain BG3594 of a nucleic acid sequence, which encodes said polypeptide, is the only polypeptide expression product of said nucleic acid sequence that exhibits transgalactosylating activity.

Aspect 11. The polypeptide according to any one of the preceding aspects, wherein said sequence identity is at

least 95%, such as, e.g. at least 96%, at least 97%, at least 98%, at least 99% or at least 100% sequence identity.

Aspect 12. A polypeptide wherein said polypeptide has at least 90% sequence identity with SEQ ID NO: 1.

Aspect 13. The polypeptide according to aspect 12, wherein said polypeptide has transgalactosylating activity.

Aspect 14. The polypeptide according to any one of the preceding aspects, wherein said polypeptide has at least 90% such as, e.g. at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity with SEQ ID NO: 1.

Aspect 15. The polypeptide according to any one of aspects 1-14, wherein the degree of sequence identity between a query sequence and a reference sequence is determined by 1) aligning the two sequences by any suitable alignment program using the default scoring matrix and default gap penalty, 2) identifying the number of exact matches, where an exact match is where the alignment program has identified an identical amino acid or nucleotide in the two aligned sequences on a given position in the alignment and 3) dividing the number of exact matches with the length of the reference sequence.

Aspect 16. The polypeptide according to any one of aspects 1-15, wherein the degree of sequence identity between a query sequence and a reference sequence is determined by 1) aligning the two sequences by any suitable alignment program using the default scoring matrix and default gap penalty, 2) identifying the number of exact matches, where an exact match is where the alignment program has identified an identical amino acid or nucleotide in the two aligned sequences on a given position in the alignment and 3) dividing the number of exact matches with the longest of the two sequences.

Aspect 17. The polypeptide according to any one of aspects 1-16, wherein the degree of sequence identity between a query sequence and a reference sequence is determined by 1) aligning the two sequences by any suitable alignment program using the default scoring matrix and default gap penalty, 2) identifying the number of exact matches, where an exact match is where the alignment program has identified an identical amino acid or nucleotide in the two aligned sequences on a given position in the alignment and 3) dividing the number of exact matches with the "alignment length", where the alignment length is the length of the entire alignment including gaps and overhanging parts of the sequences.

Aspect 18. The polypeptide according to any one of aspects 15-17, wherein the suitable alignment program is a global alignment program.

Aspect 19. The polypeptide according to aspect 18, wherein the global alignment program uses the Needleman-Wunsch algorithm.

Aspect 20. The polypeptide according to any of aspects 18 and 19, wherein the global alignment program is selected from the group consisting of EMBOSS Needle and EMBOSS stretcher.

Aspect 21. The polypeptide according to any one of the preceding aspects, which consists of at most 975 amino acid residues, such as, e.g. at most 970 amino acid residues, such as at most 950 amino acid residues, such as at most 940 amino acid residues, at most 930 amino acid residues, at most 920 amino acid residues, at most 910 amino acid residues, at most 900 amino acid residues, at most 895 amino acid residues or at most 890 amino acid residues.

Aspect 22. The polypeptide according to any one of the preceding aspects, which consists of 887 amino acid residues.

Aspect 23. The polypeptide according to any one of the preceding aspects, which comprises SEQ ID NO:1.

Aspect 24. The polypeptide according to any one of the preceding aspects, which consist of the amino acid sequence of SEQ ID NO:1.

Aspect 25. The polypeptide according to any one of the preceding aspects, wherein said polypeptide consists of 965 amino acid residues.

Aspect 26. The polypeptide according to any of aspects 1-23 and 25, wherein said polypeptide has at least 96,5%

sequence identity to SEQ ID NO: 2.

Aspect 27. The polypeptide according to any of aspects 1-23 and 25-26, wherein said polypeptide has at least 97%, such as, e.g. at least 98% or at least 99% sequence identity with SEQ ID NO: 2.

Aspect 28. The polypeptide according to any one of aspects 1-23 and 25-27, wherein said polypeptide comprises SEQ ID NO: 2.

Aspect 29. The polypeptide according to any one of aspects 1-23 and 25-28, wherein said polypeptide consists of the amino acid sequence of SEQ ID NO: 2.

Aspect 30. A polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues.

Aspect 31. The polypeptide according to aspect 30, wherein said polypeptide has transgalactosylating activity.

Aspect 32. The polypeptide according to any one of aspects 30-31, wherein said polypeptide, when being an expression product in a *Bacillus subtilis* strain BG3594 of a nucleic acid sequence, which encodes said polypeptide, is the only polypeptide expression product of said nucleic acid sequence that exhibits transgalactosylating activity.

Aspect 33. The polypeptide according to any one of aspects 1-5, 15-20 and 30-32, wherein said sequence identity is at least 97%, such as, e.g. at least 98%, at least 99% or at least 100% sequence identity.

Aspect 34. The polypeptide according to any one of aspects 1-5, 15-20 and 30-33, which comprises SEQ ID NO: 3.

Aspect 35. The polypeptide according to any one of aspects 11-5, 15-20 and 30-34, which consists of the amino acid sequence of SEQ ID NO: 3.

Aspect 36. The polypeptide according to any one of aspects 1-5, 15-20 and 30-34, which consists of at most 1290 amino acid residues, such as, e.g. at most 1280, at most 1270, at most 1260, at most 1250, at most 1240, at most 1230, at most 1220 or at most 1215 amino acid residues.

Aspect 37. The polypeptide according to any one of aspects 1-5, 15-20, 30-34 and 36, which consists of 1211 amino acid residues.

Aspect 38. The polypeptide according to any one of aspects 1-5, 15-20, 30-34 and 36-37, wherein said polypeptide has at least 98.5% sequence identity with SEQ ID NO: 5.

Aspect 39. The polypeptide according to any one of aspects 1-5, 15-20, 30-34 and 36-38, wherein said polypeptide has at least 99% or at least 99.5% sequence identity with SEQ ID NO: 5.

Aspect 40. The polypeptide according to any one of aspects 1-5, 15-20, 30-34 and 36-39, which comprises SEQ ID NO: 5.

Aspect 41. The polypeptide according to any one of aspects 1-5, 15-20, 30-34 and 36-40, which consists of the amino acid sequence of SEQ ID NO: 5.

Aspect 42. The polypeptide according to any one of aspects 1-5, 15-20, 30-34 and 36-40, which consists of at most 1210 amino acid residues, such as, e.g. at most 1200, at most 1190, at most 1180, at most 1170, at most 1160, at most 1150 or at most 1145 amino acid residues.

Aspect 43. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40 and 42, which consists of 1142 amino acid residues.

Aspect 44. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40 and 42-43, wherein said polypeptide has at least 96% sequence identity with SEQ ID NO: 4.

Aspect 45. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40 and 42-44, wherein said

EP 3 473 713 A1

polypeptide has at least 97%, such as, e.g., at least 98% or at least 99% sequence identity with SEQ ID NO: 4.

Aspect 46. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40 and 42-45, which comprises SEQ ID NO: 4.

Aspect 47. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40 and 42-46, which consists of the amino acid sequence of SEQ ID NO: 4.

Aspect 48. The polypeptide according to any one aspects 1-5, 15-20, 30-34, 36-40 and 42-46, which consists of at most 1130 amino acid residues, such as, e.g. at the most 1120, at the most 1110, at the most 1100, at the most 1090, at the most 1080, at the most 1070, at the most 1060, at the most 1050, at the most 1055 or at the most 1040 amino acid residues.

Aspect 49. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40, 42-46 and 48, wherein said polypeptide consists of 1038 amino acid residues.

Aspect 50. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40, 42-46 and 48-49, wherein said polypeptide has at least 96.5% sequence identity with SEQ ID NO: 3.

Aspect 51. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40, 42-46 and 48-50, wherein said polypeptide has at least 97%, such as, e.g., at least 98% or at least 99% sequence identity with SEQ ID NO: 3.

Aspect 52. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40, 42-46 and 48-51, which comprises SEQ ID NO:3.

Aspect 53. The polypeptide according to any one of aspects 1-5, 15-20, 30-34, 36-40, 42-46 and 48-52, which consists of the amino acid sequence of SEQ ID NO: 3.

Aspect 54. The polypeptide according to any of the preceding aspects, wherein the ratio of transgalactosylating activity: β -galactosidase activity is at least 0.5, such as at least 1, such as at least 1.5, such as at least 2 after 30 min. reaction such as above a concentration of 3% w/w initial lactose concentration.

Aspect 55. The polypeptide according to any one of the preceding aspects, wherein the ratio of transgalactosylating activity: β -galactosidase activity is at least 2.5, such as at least 3, such as at least 4, such as at least 5, such as at least 6, such as at least 7, such as at least 8, such as at least 9, such as at least 10, such as at least 11, or such as at least 12 after 30 min. reaction such as above a concentration of 3% w/w initial lactose concentration.

Aspect 56. The polypeptide according to any one of aspects 54-55, wherein the initial lactose concentration is 3% w/w.

Aspect 57. The polypeptide according to any one of the preceding aspects, which is isolated and/or purified.

Aspect 58. The polypeptide according to any one of the preceding aspects, which is recombinantly produced.

Aspect 59. The polypeptide according to any one of the preceding aspects, wherein said polypeptide has a ratio of transgalactosylation activity above 100%.

Aspect 60. The polypeptide according to any one of the preceding aspects, wherein said polypeptide has a ratio of transgalactosylation activity above 150%, 175% or 200%.

Aspect 61. The polypeptide according to any one of the preceding aspects, having a glycoside hydrolase catalytic core with an amino acid sequence selected from the group consisting of SEQ ID NO:7.

Aspect 62. The polypeptide according to any one of the preceding aspects containing a Glyco_hydro2N (PF02837), a Glyco_hydro (PF00703) and/or a Glyco_hydro 2C (PF02836) domains.

Aspect 63. The polypeptide according to any one of the preceding aspects containing the Bacterial Ig-like domain (group 4) (PF07532).

EP 3 473 713 A1

Aspect 64. The polypeptide according to any one of the preceding aspects, which is derived from *Bifidobacterium bifidum*.

Aspect 65. The polypeptide according to any one of the preceding aspects having a pH optimum of 6.5-7.5.

Aspect 66. The polypeptide according to any one of the preceding aspects having a temperature optimum of 30-60 such as 42-60 degree celcius.

Aspect 67. The polypeptide according to any one of the preceding aspects, wherein the percentage of identity of one amino acid sequence with, or to, another amino acid sequence is determined by the use of Blast with a word size of 3 and with BLOSUM 62 as the substitution matrix.

Aspect 68. The polypeptide according to any one of the preceding aspects, which polypeptide is a recombinant polypeptide.

Aspect 69. The polypeptide according to any one of the preceding aspects, which polypeptide is freeze-dried.

Aspect 70. The polypeptide according to any one of the preceding aspects, which polypeptide is in solution.

Aspect 71. The polypeptide according to any one of the preceding aspects, which polypeptide is isolated.

Aspect 72. The polypeptide according to any one of the preceding aspects, which polypeptide is purified.

Aspect 73. A nucleic acid capable of encoding a polypeptide according to any one of the preceding aspects.

Aspect 74. An expression vector comprising a nucleic acid according to aspect 73, or capable of expressing a polypeptide according to any one aspects 1-72.

Aspect 75. A cell capable of expressing a polypeptide according to any one of aspects 1-72.

Aspect 76. A method of expressing a polypeptide, the method comprising obtaining a cell according aspect 75 and expressing the polypeptide from the cell, and optionally purifying the polypeptide.

Aspect 77. A composition comprising a polypeptide as defined in any one of aspects 1-72, preferably a food composition, more preferably a dairy product.

Aspect 78. The composition according to aspect 77 comprising at least 5%, such as e.g. 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50% w/w of one or more polypeptide(s) as defined in any one of aspects 1-72 based on the total amount of polypeptides in the composition having at least 70%, e.g. such as 72%, 74%, 74%, 78%, 80%, 82%, 84%, 86%, 88%, 90% sequence identity with SEQ ID NO: 22.

Aspect 79. The composition according to any one of aspects 77-78, wherein the one or more polypeptide(s) is selected from the group consisting of a polypeptide consisting of SEQ ID NO: 1, 2, 3, 4 and 5.

Aspect 80. The composition according to any one of aspects 77-79, wherein the one or more polypeptide(s) is selected from the group consisting of a polypeptide consisting of SEQ ID NO: 1, 2, and 3.

Aspect 81. The composition according to any one of aspects 77-80, wherein the one or more polypeptide(s) is a polypeptide consisting of SEQ ID NO: 1 or 2.

Aspect 82. A method for producing a food product by treating a substrate comprising lactose with a polypeptide as defined in any on of aspects 1-72.

Aspect 83. The method according to aspect 82 for producing a dairy product by treating a milk-based substrate comprising lactose with a polypeptide as defined in any one of aspects 1-72.

Aspect 84. The method according to any one of aspects 82-83 further treating the substrate with a hydrolysing beta-galactosidase.

EP 3 473 713 A1

Aspect 85. A galacto-oligosaccharide or composition thereof obtained by treating a substrate comprising lactose with a polypeptide as defined in any one of aspects 1-72.

Aspect 86. A nucleic acid capable of encoding a polypeptide according to any one of the aspects 1-72.

Aspect 87. The nucleic acid according to aspect 86 having a nucleic acid sequence which is at least 60% identical to SEQ ID NO: 9, 10, 11, 12 or 13.

Aspect 88. A plasmid comprising a nucleic acid according to any one of the aspects 86-87.

Aspect 89. An expression vector comprising a nucleic acid according to any one of the aspects 86-87, or capable of expressing a polypeptide according to any one of the aspects 1-72.

Aspect 90. A host cell comprising, preferably transformed with, a plasmid according to aspect 88 or an expression vector according to aspect 89.

Aspect 91. A cell capable of expressing a polypeptide according to any one of the aspects 1-72.

Aspect 92. The host cell according to aspect 90, or the cell according to aspect 91, which is a bacterial, fungal or yeast cell.

Aspect 93. The cell according to aspect 92, wherein the cell is selected from the group consisting of *Ruminococcus*, *Bifidobacterium*, *Lactococcus*, *Lactobacillus*, *Streptococcus*, *Leuconostoc*, *Escherichia*, *Bacillus*, *Streptomyces*, *Saccharomyces*, *Kluyveromyces*, *Candida*, *Torula*, *Torulopsis* and *Aspergillus*.

Aspect 94. The cell according to aspect 93, wherein the cell is selected from the group consisting of *Ruminococcus hansenii*, *Ruminococcus lactaris*, *Bifidobacterium breve*, *Bifidobacterium longum*, *Bifidobacterium infantis*, *Bifidobacterium bifidum* and *Lactococcus lactis*.

Aspect 95. A method of expressing a polypeptide, the method comprising obtaining a host cell or a cell according to any one of aspects 91-94 and expressing the polypeptide from the cell or host cell, and optionally purifying the polypeptide.

Aspect 96. A composition comprising a polypeptide as defined in any of aspects 1-72 and a stabilizer.

Aspect 97. A composition comprising a polypeptide as defined in any of aspects 1-72 and a carbohydrate substrate.

Aspect 98. The composition according to aspect 97, wherein the carbohydrate substrate is a disaccharide.

Aspect 99. The composition according to aspect 98, wherein the disaccharide is lactose.

Aspect 100. A method for producing a dairy product by treating a milk-based substrate comprising lactose with a polypeptide according to any one of aspects 1-72.

Aspect 101. The method according to aspect 90, wherein the polypeptide has a ratio of transgalactosylation activity as defined above.

Aspect 102. The method according to any one of aspects 100-101, wherein the milk-based substrate is yoghurt, cheese, or fermented dairy products.

Aspect 103. The method according to any one of aspects 100-102, further comprising fermenting said substrate with a microorganism capable of fermenting said substrate.

Aspect 104. The method according to aspect 103, wherein the milk-based substrate is yogurt.

Aspect 105. The method according to any one of aspects 100-104, wherein the treatment with the polypeptide and the microorganism is performed essentially at the same time.

EP 3 473 713 A1

Aspect 106. The method according to any one of aspects 100-105, wherein the polypeptide and the microorganism are added to the milk-based substrate essentially at the same time.

Aspect 107. Use of a cell of any of above aspects for producing a product selected from the group consisting of yoghurt, cheese, fermented milk product, dietary supplement and probiotic comestible product.

Aspect 108. A dairy product comprising a cell of any of above aspects.

Aspect 109. A dairy product comprising a polypeptide as defined in any one of aspects 1-72.

Aspect 110. A dairy product comprising a polypeptide as defined in any one of aspects 1-72 in a concentration of 0.01-1000 ppm.

Aspect 111. A dairy product comprising an inactivated polypeptide as defined in any one of aspects 1-72.

Aspect 112. A dairy product comprising an inactivated polypeptide as defined in any one of aspects 1-72 in a concentration of 0.01-1000 ppm.

Aspect 113. A dairy product comprising GOS formed in situ by a polypeptide as defined in any one of aspects 1-72.

Aspect 114. Use of a transgalactosylating polypeptide of any one of aspects 1-72 or a cell of any one of above aspects, for producing galacto-oligosaccharides.

Aspect 115. Use of a transgalactosylating polypeptide of any one of aspects 1-72 or a cell of any one of above aspects, for producing galacto-oligosaccharides to be part of a product selected from the group consisting of yoghurt, cheese, fermented dairy products, dietary supplements and probiotic comestible products.

Aspect 116. Use of a transgalactosylating polypeptide of any one of aspects 1-72 or a cell of any one of above aspects, for producing galacto-oligosaccharides to enhance the growth of *Bifidobacterium*.

Aspect 117. Use of a transgalactosylating polypeptide of any one of aspects 1-72 or a cell of any one of above aspects, for producing galacto-oligosaccharides to enhance the growth of *Bifidobacterium* in a mixed culture fermentation.

Aspect 118. A process for producing a transgalactosylating polypeptide of any one of aspects 1-72, comprising culturing a cell of any one of above aspects in a suitable culture medium under conditions permitting expression of said polypeptide, and recovering the resulting polypeptide from the culture.

Aspect 119. A process for producing galacto-oligosaccharides, comprising contacting of an polypeptide of any one of aspects 1-72 or a cell of any one of above aspects with a milk-based solution comprising lactose.

Aspect 120. A polypeptide which is a C-terminal truncated fragment of SEQ ID NO:22 having transgalactosylating activity and which are stable against further truncation such as by proteolytic degradation when produced in a suitable organism such as *Bacillus subtilis* e.g. *Bacillus subtilis* strain BG3594 and/or which are stable against further truncation during storage after final formulation.

Aspect 121. The polypeptide according to aspect 120 which is as further defined in any one of aspects 1-72.

MATERIALS AND METHODS

Method 1

Production of polypeptide

[0237] Synthetic genes designed to encode the *Bifidobacterium bifidum* full length (1752 residues) gene with codons optimised for expression in *Bacillus subtilis* were purchased from GeneART (Regensburg, Germany) SEQ ID No. 8

[0238] The *Bifidobacterium bifidum* truncation mutants were constructed using polymerase chain reaction with reverse primers that allowed specific amplification of the selected region of the synthetic gene.

EP 3 473 713 A1

[0239] Forward primer: GGGGTAACTAGTGGAAGATGCAACAAGAAG (Spel underlined) (SEQ ID NO: 15).

Reverse primers:

[0240]

Truncation mutant	Primer sequence
BIF917 (SEQ ID NO: 9)	GCGCTTAATTAATTATGTTTTTCTGTGCTTGTTT SEQ ID NO: 16
BIF995 (SEQ ID NO: 10)	GCGCTTAATTAATTACAGTGCGCCAATTTTCATCAATCA SEQ ID NO: 17
BIF1068 (SEQ ID NO: 11)	GCGCTTAATTAATTATTGAACTCTAATTGTCGCTG SEQ ID NO: 18
BIF1241 (SEQ ID NO: 12)	GCGCTTAATTAATTATGTCGCTGTTTTTCAGTTCAAT SEQ ID NO: 19
BIF1326 (SEQ ID NO: 13)	GCGCTTAATTAATTAAAATTCTTGTTCTGTGCCCA SEQ ID NO: 20
BIF1478 (SEQ ID NO: 14)	GCGCTTAATTAATTATCTCAGTCTAATTTTCGCTTGCGC SEQ ID NO: 21

[0241] The synthetic gene was cloned into the pBNspe *Bacillus subtilis* expression vector using the unique restriction sites Spel and PacI (Figure 1) and the isolated plasmids were transformed into the *Bacillus subtilis* strain BG3594. Transformants were restreaked onto LB plates containing 10 µg/mL Neomycin as selection.

[0242] A preculture was setup in LB media containing 10 µg/mL Neomycin and cultivated for 7 hours at 37°C and 180 rpm shaking. 500 µL of this preculture was used to inoculate 50 mL Grant's modified medium containing 10 µg/mL Neomycin at allowed to grow for 68 hours at 33°C and 180 rpm shaking.

[0243] Cells were lysed by addition directly to the culture media of 1 mg/ml Lysozyme (Sigma-Aldrich) and 10 U/ml Benzonase (Merck) final concentrations and incubated for 1 hr at 33°C at 180 RPM. Lysates were cleared by centrifugation at 10.000 x g for 20 minutes and subsequently sterile filtered.

Grant's modified media was prepared according to the following directions:

[0244]

PART I (Autoclave)

Soytone	10 g
Bring to	500 mL per liter

PART II

1M K ₂ HPO ₄	3 mL
Glucose	75 g
Urea	3.6 g
Grant's 10X MOPS	100 mL
Bring to 400 mL per liter	

PART I (2 w/w % Soytone) was prepared, and autoclaved for 25 minutes at 121°C.

PART II was prepared, and mixed with PART 1 and pH was adjusted to pH to 7.3 with HCl/NaOH.

[0245] The volume was brought to full volume and sterilized through 0.22-µm PES filter.

10 xMOPS Buffer was prepared according to the following directions:

[0246]

83.72 g	Tricine
7.17 g	KOH Pellets
12 g	NaCl
29.22 g	0.276M K ₂ SO ₄
10 mL	0.528M MgCl ₂
10 mL	Grant's Micronutrients 100X

EP 3 473 713 A1

[0247] Bring to app. 900 mL with water and dissolve. Adjust pH to 7.4 with KOH, fill up to 1 L and sterile filter the solution through 0.2 µm PES filter.

100 x Micronutrients was prepared according to the following directions:

[0248]

Sodium Citrate.2H ₂ O	1.47 g
CaCl ₂ .2H ₂ O	1.47 g
FeSO ₄ .7H ₂ O	0.4 g
MnSO ₄ .H ₂ O	0.1 g
ZnSO ₄ .H ₂ O	0.1 g
CuCl ₂ .2H ₂ O	0.05 g
CoCl ₂ .6H ₂ O	0.1 g
Na ₂ MoO ₄ .2H ₂ O	0.1 g

[0249] Dissolve and adjust volume to 1 L with water.

[0250] Sterilization was through 0.2 µm PES filter.

[0251] Storing was at 4°C avoid light.

Method 2

Purification and enzyme preparations

[0252] The filtrated enzyme isolate was concentrated using a VivaSpin ultra filtration device with a 10 kDa MW cut off (Vivaspin 20, Sartorius, Lot#12VS2004) and the concentrate was loaded onto a PD10 desalting column (GE healthcare, Lot# 6284601) and eluted in 20mM Tris-HCl pH 8.6. Chromatography was carried out manually on an Äkta FPLC system (GE Healthcare). 4mL of the desalted sample, containing approximately 20mg protein, was loaded onto a 2 mL HyperQ column (HyperCel™, Q sorbent) equilibrated with 20 mM Tris-HCl pH 8.6 at a flowrate of 1ml/min. The column was thoroughly washed with 30 CV (column volumes) wash buffer and the bound β-galactosidase was eluted with a 100CV long gradient into 20 mM Tris-HCl pH 8.6 250 mM NaCl. Remaining impurities on the column were removed with a one-step elution using 20 mM Tris-HCl pH 8.6 500 mM NaCl. Protein in the flow through and elution was analyzed for β-galactosidase activity and by SDS-page.

[0253] SDS-page gels were run with the Invitrogen NuPage® Novex 4-12% Bis-Tris gel 1.0 mm, 10 well (Cat#NP0321box), See-Blue® Plus2 prestained Standard (Cat# LC5925) and NuPAGE® MES SDS Running Buffer (Cat# NP0002) according to the manufacturer's protocol. Gels were stained with Simply Blue Safestain (Invitrogen, Cat# LC6060) (Figure 2).

Method 3

Measuring β-galactosidase activity

[0254] Enzymatic activity was measured using the commercially available substrate 2-Nitrophenyl-β-D-Galactopyranoside (ONPG) (Sigma N1127).

[0255] ONPG w/o acceptor

100	mM KPO4 pH6.0
12,3	mM ONPG

[0256] ONPG supplemented with acceptor

100	mM KPO4 pH6.0
20	mM Cellobiose
12,3	mM ONPG

[0257] STOP Solution10% Na₂CO₃

[0258] 10 µl dilution series of purified enzyme was added in wells of a microtiter plates containing 90 µl ONPG-buffer with or without acceptor. Samples were mixed and incubated for 10 min at 37°C, subsequently 100 µl STOP Solution were added to each well to terminate reaction. Absorbance measurements were recorded at 420 nm on a Molecular Device SpectraMax platereader controlled by the Softmax software package.

[0259] The ratio of transgalactosylation activity was calculated as follows

Ratio of transgalactosylation activity = (Abs₄₂₀^{+Cellobiose}/Abs₄₂₀^{-Cellobiose})*100, for dilutions where the absorbance was between 0.5 and 1.0 (Figure 3).

Method 4**Determination of LAU activity**Principle:

[0260] The principle of this assay method is that lactase hydrolyzes 2-o-nitrophenyl-β-D-galactopyranoside (ONPG) into 2-o-nitrophenol (ONP) and galactose at 37°C. The reaction is stopped with the sodium carbonate and the liberated ONP is measured in spectrophotometer or colorimeter at 420 nm.

Reagents:

[0261] MES buffer pH 6.4 (100mM MES pH 6.4, 10mM CaCl₂): Dissolve 19,52g MES hydrate (Mw: 195.2 g/mol, Sigma-aldrich #M8250-250G) and 1.470g CaCl₂ di-hydrate (Mw: 147.01g/mol, Sigma-aldrich) in 1000 ml ddH₂O, adjust pH to 6.4 by 10M NaOH. Filter the solution through 0.2µm filter and store at 4°C up to 1 month.

ONPG substrate pH 6.4 (12.28mM ONPG, 100mM MES pH 6.4, 10mM CaCl₂): Dissolve 0.370g 2-o-nitrophenyl-β-D-galactopyranoside (ONPG, Mw: 301.55 g/mol, Sigma-aldrich #N1127) in 100 ml MES buffer pH 6.4 and store dark at 4°C for up to 7 days.

Stop reagent (10% Na₂CO₃): Dissolve 20.0g Na₂CO₃ in 200ml ddH₂O, Filter the solution through 0.2µm filter and store at RT up to 1 month.

Procedure:

[0262] Dilution series of the enzyme sample was made in the MES buffer pH 6.4 and 10µL of each sample dilution were transferred to the wells of a microtiter plate (96 well format) containing 90 µl ONPG substrate pH 6.4. The samples were mixed and incubated for 5 min at 37°C using a Thermomixer (Comfort Thermomixer, Eppendorf) and subsequently 100 µl Stop reagent was added to each well to terminate the reaction. A blank was constructed using MES buffer pH 6.4 instead of the enzyme sample. The increase in absorbance at 420 nm was measured at a ELISA reader (SpectraMax platereader, Molecular Device) against the blank.

Calculation of enzyme activity:

[0263] The molar extinction coefficient of 2-o-nitrophenol (Sigma-aldrich #33444-25G) in MES buffer pH 6.4 was determined ($0.5998 \times 10^{-6} \text{ M}^{-1} \times \text{cm}^{-1}$). One unit (U) of lactase activity (LAU) was defined as that corresponding to the hydrolysis of 1 nmol of ONPG per minute. Using microtitre plates with a total reaction volume of 200µL (light path of 0.52cm) the lactase activity per mL of the enzyme sample may be calculated using the following equation:

$$LAU/ml \left(\frac{nmol}{min \cdot mL} \right) = \frac{Abs_{420} \times 200\mu L \times dilution\ factor}{0.5998 \cdot 10^3 \cdot nM^{-1} \cdot cm^{-1} \times 0.52cm \times 5min \times 0.01mL}$$

Calculation of specific activity for BIF917 shown herein as SEQ ID NO: 1:Determination of BIF917 concentration:

[0264] Quantification of the target enzyme (BIF917) and truncation products were determined using the Criterion Stain free SDS-page system (BioRad). Any kD Stain free precast Gel 4-20% Tris-HCl, 18 well (Comb #345-0418) was used with a Serva Tris - Glycine/SDS buffer (BioRad cat. #42529). Gels were run with the following parameters: 200 V, 120

EP 3 473 713 A1

mA, 25 W, 50 min. BSA (1,43 mg/ml) (Sigma-Aldrich, cat. #500-0007) was used as protein standard and Criterion Stain Free Imager (BioRad) was used with Image Lab software (BioRad) for quantification using band intensity with correlation of the tryptophan content.

5 **[0265]** The specific LAU activity of BIF917 was determined from crude ferment (ultra filtration concentrate) of two independent fermentations (as described in method 1) and using 5 different dilutions (see table 1).

[0266] The specific activity of BIF917 was found to be 21.3 LAU/mg or 0.0213 LAU/ppm.

10

15

20

25

30

35

40

45

50

55

Table 1: Determination of BIF917 specific activity

Sample ID	Enzyme	Fermentation	Dilution factor	Activity LAU/ml	Protein (BIF917) concentration mg/ml	Protein (BIF917) concentration ppm	Specific activity LAU/mg	Specific activity LAU/ppm
1	BIF 917	a	5	26.9	1.23	1232	21.9	0.0219
2	BIF 917	a	10	53.9	2.44	2437	22.1	0.0221
3	BIF 917	a	10	75.4	3.56	3556	21.2	0.0212
4	BIF 917	a	20	163.9	7.78	7778	21.1	0.0211
5	BIF 917	a	30	233.6	11.06	11065	21.1	0.0211
6	BIF 917	b	5	30.26825	1.34	1342	22.6	0.0226
7	BIF 917	b	10	55.91536	2.61	2607	21.4	0.0214
8	BIF 917	b	10	76.96056	3.70	3697	20.8	0.0208
9	BIF 917	b	20	156.986	7.75	7755	20.2	0.0202
10	BIF 917	b	30	236.9734	11.45	11452	20.7	0.0207
				Arg			21.3	0.0213
				Std			0.700976	0.000701

EXAMPLES

Example 1

Determining β -galactosidase activity of BIF truncation variants

[0267] Eight different truncation variants: BIF_917, BIF_995, BIF_1068, BIF_1172, BIF_1241, BIF1326, BIF_1400 and BIF_1478 were constructed as described using method 1 and purified as described in method 2 (see Figure 2). The β -galactosidase activity was determined of all truncation variants in presence and absence of cellobiose using the described method 3 above.

Results

[0268] The ratio of transgalactosylation activity $((\text{Abs}_{420}^{+\text{Cellobiose}}/\text{Abs}_{420}^{-\text{Cellobiose}}) \cdot 100)$ was calculated from the measured β -galactosidase activity for each variant and is shown in Figure 3. Variants having a length of 1241 residues or less shows a ratio of transgalactosylation activity above 100%, indicating that these variants are predominantly transgalactosylating. The variants with a length that is more than 1241 residues shows a ratio of transgalactosylation activity below 100%, indicating that these variants are predominantly hydrolytic. BIF_917 and BIF_995 have the highest ratio of transgalactosylation activity around 250%.

Example 2

GOS generated in a yoghurt matrix

[0269] Evaluation of BIF enzymes in GOS production were tested in an yogurt application mimic. Batch experiments with a volume of 100 μl were performed in 96 well MTP plates using a yogurt mix, consisting of 98,60%(w/v) fresh pasteurized low-fat milk (Mini-mælk, Aria Foods, Denmark) and 1.4% (w/v) Nutrilac YQ-5075 whey ingredient (Aria). To completely hydrate Nutrilac YQ-5075 the mixture was left with agitation for 20h and afterwards added 20mM NaPhosphate pH 6.5 to ensure a pH of 6.5. This milk-base was used plain and the lactose concentration was determined to be 5.5%(w/v), corresponding to 5.3% (w/w) in this solution. The following correlation is valid in the present example: 1%(w/v) lactose = 0.9587 % (w/w) lactose. 90 μl of the milk-base was mixed with 10 μl of the purified enzymes, sealed with tape and incubated at 43°C for 3 hours. The reaction was stopped by 100 μl 10% Na_2CO_3 . Samples stored at -20°C.

HPLC method

[0270] Quantification of galactooligosaccharides (GOS), lactose, glucose and galactose were performed by HPLC. Analysis of samples was carried out on a Dionex ICS 3000. IC parameters were as follows: Mobile phase: 150 mM NaOH, Flow: Isochratic, 0,25ml/min, Column: Carbopac PA1, Column temperature: RT, Injection volume: 10 μL , Detector: PAD, Integration: Manual, Sample preparation: 100 times dilution in Milli-Q water (0,1 ml sample + 9,9 ml water) and filtration through 0.45 μm syringe filters, Quantification: Peak areas in percent of peak area of the standard. A GOS syrup (Vivanal GOS, Friesland Campina) was used as standard for GOS quantification. In this example, the term "GOS" is defined as galactooligosaccharides with a degree of polymerization (DP) of 3 or above.

Results

[0271] The quantified amount of GOS generated in the milk-base by BIF_917, BIF_995 and BIF_1326 is shown in Figure 4. It can be seen that the shorter variants BIF_917 and BIF_995 have a significantly (determined by a student's T-test with 95% confidence) higher GOS production around 1.2% (w/v) compared to BIF_1326 generating below 0.1% (w/v).

Example 3

Degradation pattern of truncation variants

[0272] A library covering the region between BIF1230 and BIF1325 was ordered from GeneART (Regensburg, Germany) (see table 2). The truncation variants were produced as described in method 1. The resulting peptides were subjected to SDS-PAGE analysis and visualized with Simply Blue Safestain (Invitrogen, Cat# LC6060) (Figure 5).

Results

[0273] Surprisingly, most of the variants were proteolytically modified in the final broth with varying amounts of target band appearing at the end of fermentation. The variants generated three distinct bands with varying intensities which was verified using mass spectrometry. The variants have C-terminal truncation with BIF917 corresponding to the termini of SEQ ID NO: 1, BIF995 corresponding to the termini of SEQ ID NO: 2 and BIF1068 corresponding to the termini of SEQ ID NO: 3.

The protein bands cut from the gel (marked with arrows in Figure 5) are digested using three different enzymes, as preparation for mass spectrometry analysis. Trypsin hydrolyzes peptide bonds specifically at the carboxyl side of arginine (R) and lysine (K) residues except when a proline (P) is on the carboxyl side. α -Chymotrypsin hydrolyzes peptide bonds specifically at the carboxyl side of tyrosine (Y), phenylalanine (F), tryptophan (W) and leucine (L) except when a proline (P) is on the carboxyl side. Glu-C preferentially cleaves at the carboxyl side of glutamyl (E) in ammonium bicarbonate buffer pH 8, but also cleaves at the carboxyl side of aspartyl (D) if the hydrolysis is carried out in a phosphate buffer pH 8.

[0274] In order to detect the C-terminal, the protein of interest is prepared for analysis using our basic procedure for protein characterisation (A2963), with one change using 40% ^{18}O -water in the digestion buffer. The theory is that the proteolytic cleavage will incorporate both ^{18}O -water and ^{16}O -water in the resulting peptides, which consequently will appear as doublets. The *protein* C-terminal though will only appear as a single peptide with ^{16}O -water, since it is not cleaved but just the "last peptide" left of the protein. In this way the C-terminal is mapped using MS/MS analysis.

Table 2:

Variants			
Name	Fragment of SEQ ID NO: 6		WELL
BIF1230	1	1201	A01
BIF1231	1	1202	B01
BIF1232	1	1203	C01
BIF1233	1	1204	
BIF1234	1	1205	
BIF1235	1	1206	D01
BIF1236	1	1207	
BIF1237	1	1208	E01
BIF1238	1	1209	F01
BIF1239	1	1210	G01
BIF1240	1	1211	A02
BIF1241	1	1212	B02
BIF1242	1	1213	C02
BIF1243	1	1214	
BIF1244	1	1215	E02
BIF1245	1	1216	F02
BIF1246	1	1217	G02
BIF1247	1	1218	H02
BIF1248	1	1219	A03
BIF1249	1	1220	B03
BIF1250	1	1221	C03
BIF1251	1	1222	D03
BIF1252	1	1223	E03

EP 3 473 713 A1

(continued)

5	Variants		
	Name	Fragment of SEQ ID NO: 6	WELL
10	BIF1253	1 1224	
	BIF1254	1 1225	F03
15	BIF1255	1 1226	G03
	BIF1256	1 1227	H03
20	BIF1257	1 1228	A04
	BIF1258	1 1229	B04
25	BIF1259	1 1230	C04
	BIF1260	1 1231	D04
30	BIF1261	1 1232	
	BIF1262	1 1233	E04
35	BIF1263	1 1234	F04
	BIF1264	1 1235	G04
40	BIF1265	1 1236	H04
	BIF1266	1 1237	A05
45	BIF1267	1 1238	B05
	BIF1268	1 1239	C05
50	BIF1269	1 1240	D05
	BIF1270	1 1241	E05
55	BIF1271	1 1242	F05
	BIF1272	1 1243	G05
	BIF1273	1 1244	H05
	BIF1274	1 1245	A06
	BIF1275	1 1246	B06
	BIF1276	1 1247	C06
	BIF1277	1 1248	D06
	BIF1278	1 1249	E06
	BIF1279	1 1250	
	BIF1280	1 1251	F06
	BIF1281	1 1252	G06
	BIF1282	1 1253	H06
	BIF1283	1 1254	A07
	BIF1284	1 1255	
	BIF1285	1 1256	
	BIF1286	1 1257	
	BIF1287	1 1258	B07
	BIF1288	1 1259	

EP 3 473 713 A1

(continued)

5	Variants		
	Name	Fragment of SEQ ID NO: 6	WELL
10	BIF1289	1 1260	C07
	BIF1290	1 1261	D07
15	BIF1291	1 1262	
	BIF1292	1 1263	E07
20	BIF1293	1 1264	
	BIF1294	1 1265	
25	BIF1295	1 1266	F07
	BIF1296	1 1267	G07
30	BIF1297	1 1268	
	BIF1298	1 1269	H07
35	BIF1299	1 1270	A08
	BIF1300	1 1271	B08
40	BIF1301	1 1272	C08
	BIF1302	1 1273	D08
45	BIF1303	1 1274	E08
	BIF1304	1 1275	F08
50	BIF1305	1 1276	G08
	BIF1306	1 1277	H08
55	BIF1307	1 1278	A09
	BIF1308	1 1279	B09
	BIF1309	1 1280	
	BIF1310	1 1281	
	BIF1311	1 1282	
	BIF1312	1 1283	C09
	BIF1313	1 1284	
	BIF1314	1 1285	
	BIF1315	1 1286	D09
	BIF1316	1 1287	E09
	BIF1317	1 1288	F09
	BIF1318	1 1289	G09
	BIF1319	1 1290	H09
	BIF1320	1 1291	A10
	BIF1321	1 1292	B10
	BIF1322	1 1293	C10
	BIF1323	1 1294	D10
	BIF1324	1 1295	E10

(continued)

Variants		
Name	Fragment of SEQ ID NO: 6	WELL
BIF1325	1 1296	F10

Example 4**GOS generated enzymatically in situ in milkbase and yoghurts**

[0275] In this example, the term "GOS" is defined as galactooligosaccharides with a degree of polymerization (DP) of 3 or above.

[0276] Evaluation of GOS production by BIF917 and BIF995 were tested by in situ application in different set-style yogurts. The β -galactosidase was added to the milk-base simultaneous with addition of the specific yoghurt cultures, resulting in the transgalactosylation reaction running together with the yoghurt fermentation process.

[0277] Initial yoghurt (set-style) batch experiments were made with a 100 mL milkbase (yoghurt mix). The milkbase consisted of 98.60%(w/v) fresh pasteurized conventional (not-organic) low-fat milk (Mini-mælk 0.5% fat, Aria Foods Amba, Denmark) and 1.4% (w/v) Nutrilac YQ-5075 whey ingredient (Aria Foods Ingredients, Denmark), resulting in a lactose concentration of 5.5 % (w/v) corresponding to 5.3% (w/w) (1%(w/v) lactose = 0.9587 % (w/w) lactose in this solution). To completely hydrate Nutrilac YQ-5075 the mixture was left with weak agitation for 20 hr at 4°C. in the initial experiment a freeze-dried YO-MIX 485LYO culture was used consisting of *Lactobacillus delbrueckii subsp bulgaricus* and *Streptococcus thermophilus* (DuPont Nutrition Biosciences, Denmark). An initial dilution of the culture was made, adding 10g of YO-MIX 485LYO to 400 mL UHT conventional (not-organic) milk (Let-mælk 1.5% fat, Aria Foods Amba, Denmark). 1.43 mL of the diluted culture was added per litre of milk-base. 100 mL milkbase were distributed in 250 mL bluecap bottles and enzymes were added in varying concentration (10, 20 and 40 ppm, corresponding to 0.213, 0.426 and 0.853 LAU as described in method 4) constituting 1% (v/v) of the final yogurt-mix. Yoghurt fermentation was performed at 43°C and ended after 10hr by fast cooling on ice. Fermentations were always run in duplicates and yoghurt sugar/oligosaccharide composition was analyzed by HPLC on the day after fermentation (see HPLC method below). Fermented yoghurt samples were always stored at 4°C.

The results of the initial yoghurt experiment are shown in table 3. It can be seen that increased dose of either BIF917 or BIF995 from 10 ppm to 40ppm lead to increased GOS content and decreased amount of DP2 (including lactose) in the final yoghurt. The difference in performance of the two variants is within the variance of the HPLC determination and it may be concluded that they perform similar within the investigated dosages.

Table 3. Content of DP2 saccharides (mainly lactose) and GOS (DP3+) in a fermented yogurt treated with increasing dose of BI F917 and BI D995. All results are calculated as an average of three independent measurements.

		Amount w/v % DP2 incl. Lactose	Std	Amount w/v % GOS (DP3+)	Std
Yogurt	BIF917_10 ppm	2.191	0.092	1.249	0.051
	BIF917_20 ppm	1.296	0.047	1.882	0.056
	BIF917_40 ppm	0.970	0.019	2.346	0.047
	BIF995_10ppm	2.787	0.139	1.158	0.035
	BIF995_20ppm	1.494	0.075	1.649	0.082
	BIF995_40ppm	0.931	0.028	2.392	0.063
	H2O	4.219	0.127	0.000	0.000
Milkbase		5.500	0.156	0.000	0.000

[0278] Set-style yoghurt was made with higher initial lactose concentration of 7.5% w/v (corresponding to 7.1% w/w,

as 1% w/v lactose = 0.9423 % w/w in this solution) to investigate its effect on the GOS concentration achieved in the final yoghurt. The following procedure was applied:

1. All powder ingredients (listed in table 4) are mixed and the dry blend are added to the milk / water under good agitation at 4-5°C, left to hydrate for 20 hours at 4°C.
2. The milkbase is preheated to 65°C (P1)
3. The milkbase is homogenised at 65°C / 200 bar
4. The milkbase is pasteurised 95°C for 6 minutes (P3)
5. The milkbase is cooled to 5°C (K2)

Table 4. SET yogurt ingredients list in % (w/ w)

Ingredients in % (w/w)	
Ingredient Name	
Skimmed milk (Skummet-mælk 0.1% fat, Arla Foods Amba, Denmark)	93.533
Cream 38 % fat (Arla Foods Amba, Denmark)	1.067
Nutrillac YQ5075 (Arla Foods Ingredients, Denmark)	1.400
Lactose (Variolac® 992 BG100, Arla Foods Amba, Denmark)	3.000
Enzyme/H2O	1.000
Total %	100

[0279] Following, the milkbase is heated to 43°C (K1). Dilution of the culture YO-MIX 495 consisting of *Lactobacillus delbrueckii subsp bulgaricus* and *Streptococcus thermophilus* (DuPont Nutrition Biosciences, Denmark) was done at the fermentation temperature. 250 mL milkbase were distributed in 500 mL bluecap bottles and enzymes were added in varying concentration constitution 1% (v/v) of the final yogurt-mix. The starter culture, YO-Mix 495, are added in a dosage of 20 DCU (Danisco Culture Units) per 100 litre, where one DCU is 100 billion cells measured as colony forming units. For each trial three samples were made. Fermentation was carried out to pH 4.6 at 43°C and following stopped by fast cooling to 5°C. Yoghurt sugar/oligosaccharide composition was analyzed by HPLC on the day after fermentation (see HPLC method below). Fermented yoghurt samples were always stored at 4°C.

[0280] The results of the yoghurt experiment are shown in table 5. It can be seen that higher initial lactose (7.5% w/v) increased the total GOS generated in the yoghurt compared to the GOS achieved with 5.5% (w/v) initial lactose, as shown in table 3. A final GOS concentration of 2.954% is achieved with the 50 ppm dose of BIF917 tested, whereas 2.662% GOS was produced with 25 ppm BIF917. In comparison did the conventional hydrolyzing β -galactosidase from *Kluyveromyces lactis* (GODO-YNL2, GODO SHUSEI Co., Ltd., Japan) produced 0,355% GOS at a dose of 25 ppm. A decrease in the lactose concentration of 2.127% was observed in the blank yogurt where the same amount of H2O was added instead of enzyme.

Table 5. Content of DP2 saccharides (mainly lactose) and GOS (DP3+) in a fermented yogurt treated with increasing dose of BI F917 and *K. lactis* β -gal.

		Amount w/v % DP2 incl. Lactose	Std	Amount w/v % DP3+ (GOS)	Std
Yogurt	BIF917_12,5 ppm	3.295	0.224	1.525	0.197
	BIF917_25 ppm	2.090	0.045	2.662	0.003
	BIF917_50 ppm	1.395	0.090	2.954	0.202
	<i>K. lactis</i> β -Gal_25ppm	0.425	0.040	0.355	0.006
	H2O	5.431	0.099	0.000	0.000
Milkbase	H2O	7.558	0.265	0.000	0.000

[0281] To test the influence of acidification in the yogurt fermentation on BIF917 performance studies were made with

EP 3 473 713 A1

three different YO-mix cultures all consisting of *Lactobacillus delbrueckii subsp bulgaricus* and *Streptococcus thermophilus* (DuPont Nutrition Biosciences, Denmark): YO-MIX 495 with a relative slow fermentation time; YO-MIX 495 with a relative fast fermentation time and YO-MIX 601 with prolonged lagphase and a strong acidifying fermentation. All fermentations were performed at 43°C with the same amount of BIF917 (25 ppm).

In addition to test the influence of temperature on BIF917 performance, fermentation with YO-MIX 495 and 25 ppm BIF917 were carried out at 43°C, 45°C and 47°C

The following procedure was applied to produce set-style yogurts with an initial lactose concentration of 7.5% (w/v) (corresponding to 7.1% w/w, as 1% w/v lactose = 0.9423 % w/w in this solution):

1. All powder ingredients (listed in table 6) are mixed and the dry blend are added to the milk / water under good agitation at 4-5°C, left to hydrate for 20 hours at 4°C.
2. The milkbase is preheated to 65°C (P1)
3. The milkbase is homogenised at 65°C / 200 bar
4. The milkbase is pasteurised 95°C for 6 minutes (P3)
5. The milkbase is cooled to 5°C (K2)
6. The milkbase is heated to 43°C (K1). Dilution of the cultures YO-MIX 495, 485 and 601 (DuPont Nutrition Biosciences, Denmark) was done at the fermentation temperature.
7. 100 mL milkbase was distributed in 250 mL bluecap bottles and enzymes were added in varying concentration constitution 1% (v/v) of the final yogurt-mix
8. The starter culture, either YO-Mix 495, 485 or 601 were added in a dosage of 20 DCU (Danisco Culture Units) per 100 litre. For each trial three samples were made.
9. Fermentation was carried out to pH 4.6 at 43°C (for teampurature studies at 43°C, 45°C and 47°C respectively) and following stopped by fast cooling to 5°C
10. Yoghurt sugar/oligosaccharide composition was analyzed by HPLC on the day after fermentation (see HPLC method below). Fermented yoghurt samples were always stored at 4°C.

Table 6. SET yogurt ingredients list in % (w/ w)

Ingredients in % (w/w)	
Ingredient Name	
Skimmed milk (Skummet-mælk 0.1% fat, Arla Foods Amba, Denmark)	93.533
Cream 38 % fat (Arla Foods Amba, Denmark)	1.067
Nutrillac YQ5215 (Arla Foods Ingredients, Denmark)	1.400
Lactose (Variolac® 992 BG100, Arla Foods Amba, Denmark)	3.000
Enzyme/H2O	1.000
Total %	100

[0282] The results of the yoghurt experiment are shown in table 7. It can be seen that the different YO-mix cultures, having different acidification profiles, exert no significant effect on the final GOS yield. On average 3.22 % (w/v) GOS is generated and the highest GOS concentration (3.300%) found in the yoghurt produced with YO-mix 485 is being within the variance of quantification. The change in fermentation temperature from 43°C to 45°C and 47°C do not significantly (using a student T-test with 95% confidence limits) change the amount of GOS produced in any of the yoghurts: 3,258% w/v at 43°C, 3,375% w/v at 45°C and 3,236% w/v at 47°C. Thus, it may be concluded that the action of BIF917 under the conditions investigated are robust for in situ generation of GOS in yogurt using various culture and temperature conditions.

Tabel 7. Content of DP2, DP3, DP4, DP5, DP6, glucose and galactose in a fermented yogurt treated with BIF917.

Sample	Enzyme dose	Fermentation culture	Fermentation temperature	Lactose w/v%	Glucose w/v%	Galactose w/v%	DP3 (GOS) w/v%	DP4 (GOS) w/v%	DP5 (GOS) w/v%	DP6 (GOS) w/v%	DP3+ (GOS) w/v%
Milkbase				7.259	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Yogurt	BIF917_25ppm	YM 495	43°C	5.466	0.000	0.448	0.000	0.000	0.000	0.000	0.000
	BIF917_25ppm	YM 495	43°C	1.770	1.368	0.648	2.067	0.793	0.260	0.074	3.194
	BIF917_25ppm	YM 485	43°C	1.860	1.259	0.609	2.065	0.830	0.302	0.103	3.300
	BIF917_25ppm	YM 601	43°C	1.712	1.418	0.832	2.094	0.780	0.248	0.068	3.189
	BIF917_25ppm	YM 495	43°C	1.761	1.344	0.642	2.086	0.813	0.275	0.084	3.258
	BIF917_25ppm	YM 495	45°C	1.838	1.323	0.625	2.128	0.848	0.301	0.099	3.375
	BIF917_25ppm	YM 495	47°C	1.739	1.406	0.722	2.106	0.799	0.257	0.074	3.236
				STD							
Milkbase				0.252	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Yogurt	BIF917_25ppm	YM 495	43°C	0.225	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	BIF917_25ppm	YM 495	43°C	0.075	0.067	0.039	0.080	0.031	0.010	0.002	0.123
	BIF917_25ppm	YM 485	43°C	0.090	0.063	0.015	0.084	0.033	0.012	0.008	0.136
	BIF917_25ppm	YM 601	43°C	0.013	0.014	0.017	0.018	0.010	0.006	0.002	0.037
	BIF917_25ppm	YM 495	43°C	0.042	0.031	0.014	0.046	0.016	0.005	0.001	0.067
	BIF917_25ppm	YM 495	45°C	0.071	0.050	0.013	0.061	0.028	0.012	0.007	0.108
	BIF917_25ppm	YM 495	47°C	0.009	0.015	0.004	0.003	0.002	0.003	0.003	0.011

HPLC method

[0283] All chemicals used were of analytical grade. D-(+)-Lactose (min 99%, no.17814), D-(+)-glucose (min 99.5%, no G8270-100G, batch# 036K0137), and D-(+)-galactose (min 99%, no G0750-25G, batch# 031M0043V) were obtained from Sigma (St. Louis, Mo, USA). Vivinal GOS Syrup (Prod. No 502675 Batch#649566) was obtained from Friesland Campina Domo (Amersfoort, The Netherlands) containing 57% on dry-matter (DM) galacto-oligosaccharides, 21% on DM anhydrous lactose, 20% on DM anhydrous glucose, and 0.8% on DM anhydrous galactose.

Sample preparation

[0284] All standards: Lactose, Glucose, galactose and GOS were prepared in double distilled water (ddH₂O) and filtered through 0.45 µm syringe filters. A set of each standard was prepared ranging in concentration from 10 to 200000 ppm.

[0285] To evaluate quantification of the above set of sugars in a yogurt/milk matrix, the above standards were spiked into a milk and yogurt sample as internal controls. All milk and yogurt samples containing active β-galactosidase were inactivated by heating the sample to 95°C for 10 min. All milk samples were prepared in 96 well MTP plates (Corning, NY, USA) and diluted minimum 20 times and filtered through 0.20 µm 96 well plate filters before analysis (Corning filter plate, PVDF hydrophile membrane, NY, USA). Samples containing more than 50000 ppm (5% w/v) lactose were heated to 30°C to ensure proper solubilization. All yogurt samples were weighted and diluted 10 times in ddH₂O before homogenization of the sample using an Ultra turrax TP18/10 for a few minutes (Janke & Kunkel Ika-labortechnik, Bie & Berntsen, Denmark). β-galactosidase were inactivated by heat treatment and samples were further diluted in 96 well MTP plates filtered through 0.20 µm 96 well plate filters before analysis (Corning filter plate, PVDF hydrophile membrane, NY, USA). All samples were analyzed in 96 well MTP plates sealed with tape.

Instrumentation

[0286] Quantification of galactooligosaccharides (GOS), Lactose, glucose and galactose were performed by HPLC. Analysis of samples was carried out on a Dionex Ultimate 3000 HPLC system (Thermo Fisher Scientific) equipped with a DGP-3600SD Dual-Gradient analytical pump, WPS-3000TSL thermostated autosampler, TCC-3000SD thermostated column oven, and a RI-101 refractive index detector (Shodex, JM Science). Chromeleon datasystem software (Version 6.80, DU10A Build 2826, 171948) was used for data acquisition and analysis.

Chromatographic conditions

[0287] The samples were analyzed by HPLC using a RSO oligosaccharide column, Ag⁺ 4% crosslinked (Phenomenex, The Netherlands) equipped with an analytical guard column (Carbo-Ag⁺ neutral, AJ0-4491, Phenomenex, The Netherlands) at 70°C. The column was eluted with double distilled water (filtered through a regenerated cellulose membrane of 0.45 µm and purged with helium gas) at a flow rate of 0.3 ml/min. Isocratic flow of 0.3 ml/min was maintained throughout analysis with a total run time of 37 min and injection volume was set to 10 µL. Samples were held at 30°C in the thermostated autosampler compartment to ensure solubilisation of lactose. The eluent was monitored by means of a refractive index detector and quantification was made by the peak area relative to the peak area of the given standard. Peaks with a degree of three or higher (DP3+) in the Vivinal GOS syrup (Friesland Food Domo, The Netherlands) were used as standard for GOS quantification following manufactures declaration on GOS content in the product.

LIST OF SEQUENCES

[0288]

> SEQ ID NO: 1 (BIF_917)

vedatrsdsttqmsstpevvysavdskqrtsdfdanwkwfmlsdsvqaqdpafddsawqqvdlphdysitqkysqsneaesa
 ylpggtgwyrksftidrdlagkriainfdgvymnatvwfngvklgthpygyspfsdltgnakfggentivvkvenrlpssrwysgsg
 5 iyrdrvltitvdgvhvgngngaiktplatqnggdvtmnlttkvandteaaanitlkqtvfpkggktdaaigtvttaskiaagasadv
 stitaasplwsiknplytvrtevlnggkvldtydteygrwtgfdatsgfslngekvklkgvsmhhdqgslgavanrraierqveil
 qkmgvnsirtthnpaakalidvcnekglvveevfdmwnrskngntedygkwfgqaiagdnavlbggkdetwakfdltstinrdr
 10 napsvimwslgnemmegisgsvsgfpatsaklvawtkaadstrpmtgydnkikanwnesntmgdnltanggvvgtnysdga
 nydkirtthpswaiygsetasainsrgiynrttgaqssdkqltsydnavgwgavassawydvvqrdfvagtyvwtgfdylgept
 pwngtgsgavgswpspknsyfgivdtagfpkdttyfyqsqwnddvhtlhlpawnenvvakgsgnnvpvvvytdaakvklyft
 pkgstekrligeksftkktaagytyqvyegsddstahknmyltwnvpwaegtisaeaydennrlipegstegnasvtttgaakl
 15 kadadrktitadgkdsyievdvtdanghivpdaanrvtdvkgagklvgvngsspdhdsyqadnrkafsgkvlaivqstkeage
 itvtakadglqsstvkiaattavpgtstekt

> SEQ ID NO: 2 (BIF_995)

vedatrsdsttqmsstpevvysavdskqrtsdfdanwkwfmlsdsvqaqdpafddsawqqvdlphdysitqkysqsneaesa
 ylpggtgwyrksftidrdlagkriainfdgvymnatvwfngvklgthpygyspfsdltgnakfggentivvkvenrlpssrwysgsg
 25 iyrdrvltitvdgvhvgngngaiktplatqnggdvtmnlttkvandteaaanitlkqtvfpkggktdaaigtvttaskiaagasadv
 stitaasplwsiknplytvrtevlnggkvldtydteygrwtgfdatsgfslngekvklkgvsmhhdqgslgavanrraierqveil
 qkmgvnsirtthnpaakalidvcnekglvveevfdmwnrskngntedygkwfgqaiagdnavlbggkdetwakfdltstinrdr
 napsvimwslgnemmegisgsvsgfpatsaklvawtkaadstrpmtgydnkikanwnesntmgdnltanggvvgtnysdga
 30 nydkirtthpswaiygsetasainsrgiynrttgaqssdkqltsydnavgwgavassawydvvqrdfvagtyvwtgfdylgept
 pwngtgsgavgswpspknsyfgivdtagfpkdttyfyqsqwnddvhtlhlpawnenvvakgsgnnvpvvvytdaakvklyft
 pkgstekrligeksftkktaagytyqvyegsddstahknmyltwnvpwaegtisaeaydennrlipegstegnasvtttgaakl
 35 kadadrktitadgkdsyievdvtdanghivpdaanrvtdvkgagklvgvngsspdhdsyqadnrkafsgkvlaivqstkeage
 itvtakadglqsstvkiaattavpgtstektvrsfyysrnyyvktnkpilpsdvevrysdgtsdrqnvtdavsdqqiakagsfsvagt
 vagqkisvrvtmideigal

> SEQ ID NO: 3 (BIF_1068)

Vedatrsdsttqmsstpevyssavdskqrtsdfdanwkwfmlsdsvqaqdpafddsawqqvdlphdysitqkysqsneaesa
 ylpgggtgwyrksftidrdlagkriainfdgvymnatvwfngvklgthpygyspfsdltgnakfggentivvkvenrlpssrwysgsg
 5 iyrdrvltitvdgvhvgngngaiktplatqnggdvtmnlttkvandteaaanitlkqtvfpkggktdaaigtvttaskiaagasadv
 stitaasplwsiknplytvrtevlnggkvldtydteygfrwtgfdatsgslngekvklkgvsmhhdqgslgavanrraierqveil
 qkmgvnsirtthnpaakalidvcnekglvveevfdmwnrskngntedygkwfgqaiagdnavlbgdkdetwakfdltstinrdr
 10 napsvimwslgnemmegisgsvsgfpatsaklvawtkaadstrpmtgydnkikanwnesntmgdnltanggvvgtnysdga
 nydkirtthpswaiygsetasainsrgiynrttggaqssdkqltsydnsgavwgavassawydvvqrdfvagtyvwtgfdylgept
 pwngtgsgavgswwpspknsyfgivdtagfpkdyfyysqwnddvhtlhlipawnenvvaksgnnpvvyvtdaakvklyft
 pkgstekrligeksftkktaagytyqvyegsddstahknmyltwnvpwaegtisaeaydennrlipegstegnasvtttgaakl
 15 kadadrktitadgkdlisyievdvtdanghivpdaanrvtdvkgagklvgvngsspdhdsyqadnrkafsgkvlavqstkeage
 itvtakadqlsstvkiattavpgtstektrsfyysrnyyvtgknkpilpsdvevrysdgtsdrqnvtdavsdqqiakagsfsvagt
 vagqkisvrvtmideigallnysastpvgtpavlpgrpavlpdgtvtsanfahwtkpadtyvntagtvkvpgtatvfgkefkvtat
 20 irvq

> SEQ ID NO: 4 (BIF_1172)

vedatrsdsttqmsstpevyssavdskqrtsdfdanwkwfmlsdsvqaqdpafddsawqqvdlphdysitqkysqsneaesa
 ylpgggtgwyrksftidrdlagkriainfdgvymnatvwfngvklgthpygyspfsdltgnakfggentivvkvenrlpssrwysgsg
 iyrdrvltitvdgvhvgngngaiktplatqnggdvtmnlttkvandteaaanitlkqtvfpkggktdaaigtvttaskiaagasadv
 25 stitaasplwsiknplytvrtevlnggkvldtydteygfrwtgfdatsgslngekvklkgvsmhhdqgslgavanrraierqveil
 qkmgvnsirtthnpaakalidvcnekglvveevfdmwnrskngntedygkwfgqaiagdnavlbgdkdetwakfdltstinrdr
 30 napsvimwslgnemmegisgsvsgfpatsaklvawtkaadstrpmtgydnkikanwnesntmgdnltanggvvgtnysdga
 nydkirtthpswaiygsetasainsrgiynrttggaqssdkqltsydnsgavwgavassawydvvqrdfvagtyvwtgfdylgept
 pwngtgsgavgswwpspknsyfgivdtagfpkdyfyysqwnddvhtlhlipawnenvvaksgnnpvvyvtdaakvklyft
 35 pkgstekrligeksftkktaagytyqvyegsddstahknmyltwnvpwaegtisaeaydennrlipegstegnasvtttgaakl
 kadadrktitadgkdlisyievdvtdanghivpdaanrvtdvkgagklvgvngsspdhdsyqadnrkafsgkvlavqstkeage
 itvtakadqlsstvkiattavpgtstektrsfyysrnyyvtgknkpilpsdvevrysdgtsdrqnvtdavsdqqiakagsfsvagt
 40 vagqkisvrvtmideigallnysastpvgtpavlpgrpavlpdgtvtsanfahwtkpadtyvntagtvkvpgtatvfgkefkvtat
 irvqrsqvtigssvsgnalrltnipadkqsdtdaikdgsttdantggganpsawtnwayskaghntaeitfeyateqqlgqivm
 yffrdsnavrfpdagtkiqi

> SEQ ID NO: 5 (BIF_1241)

vedatrsdsttqmsstpevyssavdskqrtsdfdanwkwfmlsdsvqaqdpafddsawqqvdlphdysitqkysqsneaesa
 50 ylpgggtgwyrksftidrdlagkriainfdgvymnatvwfngvklgthpygyspfsdltgnakfggentivvkvenrlpssrwysgsg
 iyrdrvltitvdgvhvgngngaiktplatqnggdvtmnlttkvandteaaanitlkqtvfpkggktdaaigtvttaskiaagasadv

stitaasplksiknplytvrtevlnggkvldtydteygrwtgfdatsgfslngekvklkgvsmhhdqgslgavanrraierqveil
 qkmgvnsirtthnpaakalidvcnekglvveevfdmwnrskngntedygkwfgqaiagdnavlggdkdetwakfdltstinrdr
 napsvimwslgnemmegisgsvsgfpatsaklvawtkaadstrpmttygdnkikanwnesntmgdnltanggvvgttnysdga
 nydkirtthpswaiygsetasainsrgiynrttggagssdkqltsydnsgvawgavassawydvvqrdfvagtyvwtgfdylgept
 pwngtgsgavgswwpspknsyfgivdtagfpkdttyfyqsqwnddvhtlhlilpawnnenvvakgsgnnvpvvvytdaakvklyft
 pkgstekrligeksftkkktaagytyqvyegsddstahknmyltwnvpwaegtisaeaydennrlipegstegnasvtttgkaakl
 kadadrktitadgkdsyievdvtdanghivpdaanrvtdvkgagklvgvngsspdhdsyqadnrkafsgkvlaivqstkeage
 itvtakadglqsstvkiaattavpgtstektrsfyysrnyyvtgtnkpilpsdvevrysdgtsdrqnvtdavsdqqiakagsfsvagt
 vagqkisvrvtmideigallnysastpvgtpavlpgrpavlpdgtvtsanfahwtkpadtvynntagtvkvpgtatvfgkefkvtat
 irvqrsqvtigssvsgnalrltqnipadkqsdtldaikdgsttdantggganpsawtnwayskaghntaeitfeyateqqlgqivm
 yffrdsnavrfpdagktkiqisadgknwtdlaaetiaaqessdrvkpytydfapvgatfvkvtvnadtttsgvvcaglteielktat

> SEQ ID NO: 6 (BIF_1326)

vedatrsdsttqmsstpevyssavdskqnrtsdldanwkwfmlsdsvqaqdpafddsawqqvdlphdysitqkysqsneaesa
 ylpgggtgwyksftidrdlagkriainfdgvymnatvwfngvklgthpygyspfsdltgnakfggentivvkenrlpssrwygsg
 iyrdrvltvtdgvhvgngvaiktpslatqnggdvtmnlttkvandteaaaniltkqtvfpkggktdaaigtvttaskiaagasadv
 stitaasplksiknplytvrtevlnggkvldtydteygrwtgfdatsgfslngekvklkgvsmhhdqgslgavanrraierqveil
 qkmgvnsirtthnpaakalidvcnekglvveevfdmwnrskngntedygkwfgqaiagdnavlggdkdetwakfdltstinrdr
 napsvimwslgnemmegisgsvsgfpatsaklvawtkaadstrpmttygdnkikanwnesntmgdnltanggvvgttnysdga
 nydkirtthpswaiygsetasainsrgiynrttggagssdkqltsydnsgvawgavassawydvvqrdfvagtyvwtgfdylgept
 pwngtgsgavgswwpspknsyfgivdtagfpkdttyfyqsqwnddvhtlhlilpawnnenvvakgsgnnvpvvvytdaakvklyft
 pkgstekrligeksftkkktaagytyqvyegsddstahknmyltwnvpwaegtisaeaydennrlipegstegnasvtttgkaakl
 kadadrktitadgkdsyievdvtdanghivpdaanrvtdvkgagklvgvngsspdhdsyqadnrkafsgkvlaivqstkeage
 itvtakadglqsstvkiaattavpgtstektrsfyysrnyyvtgtnkpilpsdvevrysdgtsdrqnvtdavsdqqiakagsfsvagt
 vagqkisvrvtmideigallnysastpvgtpavlpgrpavlpdgtvtsanfahwtkpadtvynntagtvkvpgtatvfgkefkvtat
 irvqrsqvtigssvsgnalrltqnipadkqsdtldaikdgsttdantggganpsawtnwayskaghntaeitfeyateqqlgqivm
 yffrdsnavrfpdagktkiqisadgknwtdlaaetiaaqessdrvkpytydfapvgatfvkvtvnadtttsgvvcaglteielktat
 skfvntsaalssltvngtkvdsvlaagsyntpaiadvkaegegnasvtpahdnvirvitesedhvrktftinlgteqef

>SEQ ID NO: 7 Bifidobacterium bifidum glycoside hydrolase catalytic core

qnrtsdldanwkwfmlsdsvqaqdpafddsawqqvdlphdysitqkysqsneaesaylpgggtgwyksftidrdlagkriainfdgv
 ymnatvwfngvklgthpygyspfsdltgnakfggentivvkenrlpssrwygsggyrdvltvtdgvhvgngvaiktpslatq
 nggdvtmnlttkvandteaaaniltkqtvfpkggktdaaigtvttaskiaagasadvstitaasplksiknplytvrtevlnggk
 vldtydteygrwtgfdatsgfslngekvklkgvsmhhdqgslgavanrraierqveilqkmgvnsirtthnpaakalidvcnekglv
 vveevfdmwnrskngntedygkwfgqaiagdnavlggdkdetwakfdltstinrdrnapsvimwslgnemmegisgsvsgfp
 atsaklvawtkaadstrpmtty

>SEQ ID NO: 8 nucleotide sequence encoding full length

gcagttgaagatgcaacaagaagcgatagcacaacacaaatgtcatcaacaccggaagttgtttatcatcagcggtcgatagcaa
 aaaaaatcgacaaagcgattttgatgcgaactggaaatttatgtgtcagatagcgttcaagcacaagatccggcatttgatgattca
 5 gcatggcaacaagttgatctgccgcatgattatagcatcacacagaaatatagccaaagcaatgaagcagaatcagcatatcttccg
 ggaggcacaggctggtatagaaaaagctttacaattgatagagatctggcaggcaaacgcattgcgattaattttgatggcgtctat
 atgaatgcaacagctggtttaatggcggttaaactgggcacacatccgtatggctattcaccggtttcatttgatctgacaggcaatgca
 aaatttggcggagaaaaacacaattgctgtcaaaagttgaaaatagactgccgtcatcaagatggtattcaggcagcggcatttataga
 10 gatgttacactgacagttacagatggcggttcatgttggaataatggcgctcgcaattaaaacaccgtcactggcaacacaaaatggc
 ggagatgtcacaatgaacctgacaacaaaagtcgcaatgatacagaagcagcagcgaacattacactgaaacagacagttttcc
 gaaaggcggaaaaacggatgcagcaattggcacagttacaacagcatcaaaatcaattgcagcaggcgcatcagcagatgttaca
 15 agcacaattacagcagcaagcccgaactgtggtcaattaaaaacccgaacctgtatacagttagaacagaagttctgaacggagg
 caaagttctggatacatatgatacagaatatggcttctgctggacaggcttgatgcaacatcaggctttcactgaatggcgaaaaa
 gtcaaaactgaaaggcgtagcatgcatcatgatcaaggctcacttggcgagttgcaaatagacgcgcaattgaaagacaagtcga
 aatcttgcaaaaaatggcgctcaatagcattcgacacacataatccggcagcaaaagcactgattgatgtctgcaatgaaaaag
 20 gcgttctggttgcgaagaagtctttgatatgtggaaccgcagcaaaaatggcaacacggaagattatggcaaatggtttggccaag
 caattgcaggcgataatgcagttctgggaggcgataaagatgaaacatgggcgaaatttgatcttacatcaacaattaaccgcgata
 gaaatgcaccgtcagttattatgtggtcactgggcaatgaaatgatggaaggcatttcagggtcagtttcagggtttccggcaacatc
 25 agcaaaactggttgcatggacaaaagcagcagattcaacaagaccgatgacatatggcgataacaaaattaaagcgaactggaa
 cgaatcaaatataatggcgataatctgacagcaaatggcgagttgttggcacaattattcagatggcgcaactatgataaaat
 tcgtacaacacatccgtcatgggcaatttatggctcagaacagcatcagcgattaatagccgtggcatttataatagaacaacagg
 30 cggagcacaatcatcagataaacagctgacaagctatgataattcagcagttggctggggagcagttgcatcatcagcatggtatga
 tgttgttcagagagattttgtcgcaggcacatatgtttggacaggatttgattatctgggcgaaccgacaccgtggaatggcacaggc
 tcaggcgagttggctcatggccgtcaccgaaaaatagctattttggcatcgttgatacagcaggctttccgaaagatacatattttt
 tatcagagccagtggaaatgatgatgttcatacactgcatattcttccggcatggaatgaaatgttgttgcaaaaggctcaggcaata
 35 atgttccggttgcgtttatacagatgcagcgaagtgaaactgtatttacaccgaaaggctcaacagaaaaaagactgatcggcg
 aaaaatcatttacaacaaaaaacaacagcggcagggtatataatcaagtctatgaaggcagcgataaagattcaacagcgcataa
 aaacatgtatctgacatggaatgttccgtgggcagaaggcacaatttcagcgggaagcgatgatgaaaataatcgctgtattccgga
 40 aggcagcacagaaggcaacgcacagttacaacaacaggcaaaagcagcaaaactgaaagcagatgcggatcgcaaaacaatta
 cagcggatggcaagatctgtcatatattgaagtcgatgtcacagatgcaaatggccatattgttccggatgcagcaaatagagtca
 catttgatgttaaaggcgaggcaaaactggttggcgttgataatggctcatcaccggatcatgattcatatcaagcggataaccgcaa
 45 agcattttcaggcaaaagtcctggcaattgttcagtcaacaaaagaagcaggcgaaattacagttacagcaaaagcagatggcctgc
 aatcaagcacagttaaaattgcaacaacagcagttccgggaacaagcacagaaaaaacagtccgcagctttattacagccgcaac
 tattatgcaaaaacaggcaacaacacgattctgcggtcagatgttggaagttcgctattcagatggaacaagcgatagacaaaacggt
 50 acatgggatgcagtttcagatgatcaaattgcaaaagcagggtcattttcagttgcaggcacagttgcaggccaaaaaattagcggtc
 gcgtcacaatgattgatgaaattggcgactgtgaattattcagaagcacaccggttgccacaccggcagttcttccgggatcaag
 accggcagtcctgccggatggcacagtcacatcagcaattttgagtcattggacaaaaccggcagatacagtcataatacagc
 aggcacagtcaaagtaccgggaacagcaacagtttttggcaagaatttaaagtcacagcgacaattagagttcaagaagccaa
 55 gttacaattggctcatcagtttcaggaaatgactgagactgacacaaaatattccggcagataaacaatcagatacactggatgcg
 attaaagatggctcaacaacagttgatgcaaatacaggcggaggcgcaaatccgtcagcatggacaaattgggcatattcaaaagc

aggccataacacagcggaaattacatttgaatatgcgacagaacaacaactgggccagatcgatcatgtatttttcgcatagcaat
 gcagtttagatttccggatgctggcaaaacaaaaattcagatcagcgcagatggcaaaattggacagatctggcagcaacagaaa
 5 caattgcagcgcgaagaatcaagcgatagagtcaaaccgtatacatatgattttgcaccggttgccgcaacatttgttaaagtgcagt
 caaaaacgcagatacaacaacaccgtcaggcgttgttgcgcaggcctgacagaaattgaactgaaaacagcgcagcaagcaattt
 gtcacaaatacatcagcagcactgtcatcacttacagtcaatggcacaaaagtttcagattcagttctggcagcaggctcatataaca
 caccggcaattatcgcatggttaaagcggaaggcgaaggcaatgcaagcgttacagtccttccggcacatgataatgttattcgcg
 10 tcattacagaaagcgaagatcatgtcacacgcaaaacatttacaatcaacctgggcacagaacaagaatttccggctgattcagatg
 aaagagattatccggcagcagatgacagtcacagttggctcagaacaaacatcaggcacagcaacagaaggaccgaaaaaat
 ttgcagtcgatggcaacacatcaacatattggcatagcaattggacaccgacaacagttaatgatctgtggatcgcgtttgaactgca
 15 aaaaccgacaaaactggatgactgagatatcttccgcgtccggcaggctcaaaaaatggcagcgtcacagaatataaagttcagg
 tgcagatgatggaacaaactggacagatgcaggctcaggcacatggacaacggattatggttgaaactggcggaatttaataca
 accggtcacacaaaacatgttagactgaaagcggttcatatcatgcatagcggcaacgataaatttatgagcgcaagcgaaa
 ttagactgagaaaagcggctgatacaacggatatttcaggcgcacagttacagttccggcaaaactgacagttgatagagttgatg
 20 cagatcatccggcaacatttgaacaaaagatgtcacagttacactgggagatgcaacactgagatatggcgttgattatctgctgg
 attatgcaggcaatacagcagttggcaaagcaacagtgacagtttagaggcattgataaatattcaggcacagtcgcgaaaaacattt
 acaattgaactgaaaaatgcaccggcaccggaaccgacactgacatcagtttagcgtcaaaaacaaaaccgagcaaaactgacatatg
 25 ttgtcggagatgcatttgatccggcaggcctggttctgcaacatgatagacaagcagatagacctccgcaaccgctggttgcgcaac
 aagcggatgaacgcggactgacatgcggcacaaagatgcgatagagttgaacaactgcgcaaacatgaaaatagagaagcgcata
 gaacaggcctggatcatctggaatttgttggcgcagcagatggcgcagttggagaacaagcaacatttaaagtcctatgtccatgcag
 30 atcaggggagatggcagacatgatgatgcagatgaacgcgatattgatccgcagttccggctgatcatgcagttggcgaactggca
 agagcagcatgccatcatgttattggcctgagagtcgatacacatagacttaaagcaagcggcttcaaattccggctgatgatatgg
 cagaaatcgatcgcattacaggctttcatcgttttgaacgccatgtc

> SEQ ID NO: 9 nucleotide sequence encoding BIF_917

gttgaagatgcaacaagaagcgatagcacaacacaaatgtcatcaacaccggaagttgtttattcatcagcggctgatagcaaaaca
 aaatcgcaacgagcattttgatgcgaactggaaatttatgtctgcagatagcgttcaagcacaagatccggcatttgatgattcagca
 40 tggcaacaagttgatctgccgatgattatagcatcacacagaaatatagccaaagcaatgaagcagaatcagcatatcttccggga
 ggcacaggctggtatagaaaaagctttacaattgatagagatctggcaggcaaacgcattgcgattaattttgatggcgtctatatga
 atgcaacagtcgtgttaatggcgttaaactgggcacacatccgtatggctattcaccgttttcatttgatctgacaggcaatgcaaaat
 45 ttggcggagaaaaacacaattgtcgtcaaagttgaaaatagactgccgtcatcaagatggatttcaggcagcggcatttatagagatg
 ttacactgacagttacagatggcgttcattgttggaataatggcgtcgcaattaaaacaccgtcactggcaacacaaaatggcggag
 atgtcacaatgaacctgacaacaaaagtcgcgaatgatacagaagcagcagcgaacattacactgaaacagacagttttccgaaa
 50 ggcggaaaaacggatgcagcaattggcacagttacaacagcatcaaaatcaattgcagcaggcgcatcagcagatgttacaagca
 caattacagcagcaagcccgaactgtggtcaattaaaaacccgaacctgtatacagttagaacagaagttctgaacggaggcaaa
 gttctggatacatatgatacagaatatggctttcgtggacaggctttgatgcaacatcaggcctttcactgaatggcgaaaaagtc
 55 actgaaaggcgttagcatgcatcatgatcaaggctcacttggcgcagttgcaaatagacgcgcaattgaaagacaagtcgaaatcct
 gcaaaaaatggcgtcaatagcattcgcacacataatccggcagcaaaagcactgattgatgtctgcaatgaaaaaggcgttc
 tggttgtcgaagaagtcctttgatatgtggaaccgcagcaaaaaatggcaacacggaagattatggcaaatggtttggccaagcaattg

caggcgataatgcagttctgggagggcgataaagatgaaacatgggcgaaatttgatcttacatcaacaattaaccgcgatagaaat
 gcaccgtcagttattatgttggtcactgggcaatgaaatgatggaaggcatttcaggctcagtttcaggcttccggcaacatcagcaa
 aactggttgcatggacaaaagcagcagattcaacaagaccgatgacatatggcgataacaaaattaaagcgaactggaacgaatc
 5 aaatacaatgggcgataatctgacagcaaattggcggagtggttggcacaattattcagatggcgcaaactatgataaaattcgtac
 aacacatccgtcatgggcaatttatggctcagaaacagcatcagcgattaatagccgtggcatttataatagaacaacaggcggagc
 acaatcatcagataaacagctgacaagctatgataattcagcagttggctggggagcagttgcatcatcagcatggatgatgttgtt
 10 cagagagattttgtcgcaggcacatatgtttggacaggatttgattatctggggaacggacaccgtggaatggcacagggtcaggc
 gcagttggctcatggcgtcaccgaaaaatagctattttggcatcgttgatacagcaggcttccgaaagatacatattttttatcag
 agccagtggatgatgatgttcatactgcataattctccggcatggaatgaaaatgttggtgcaaaaggctcaggcaataatgttcc
 ggtgtcgtttatacagatgcagcgaaagtgaactgtattttacaccgaaaggctcaacagaaaaaagactgatcggcgaaaaat
 15 catttcaaaaaaacaacagcggcaggctatacatatcaagtctatgaaggcagcgataaagattcaacagcgcataaaaacat
 gtatctgacatggaatgttccgtgggcagaaggcacaatttcagcggaaagcgtatgatgaaaataatcgctgattccggaaggca
 gcacagaaggcaacgcacagttacaacaacaggcaaagcagcaaaactgaaagcagatgcggatcgcaaaacaattacagcg
 20 gatggcaaagatctgtcatatattgaagtcgatgtcacagatgcaaatggccatattgttccggatgcagcaaatagagtcacatttg
 atgttaaaggcgcaggcaaactggttggcgttgataatggctcatcaccggatcatgattcatatcaagcggataaccgcaaagcat
 tttcaggcaaagtcttgcaattgttcagtcacaaaagaagcaggcgaaattacagttacagcaaaagcagatggcctgcaatca
 25 agcacagttaaaattgcaacaacagcagttccgggaacaagcacagaaaaaaca

> SEQ ID NO: 10 nucleotide sequence encoding BIF_995

gttgaagatgcaacaagaagcgatagcacaacacaaatgtcatcaacaccggaagttgtttattcatcagcggctgatagcaaaca
 30 aaatcgcaacagcagttttgatgcaactggaaatttatgtctgcagatagcgttcaagcacaagatccggcatttgatgattcagca
 tggcaacaagttgatctgccgatgattatagcatcacacagaaatatagccaaagcaatgaagcagaatcagcatatctccggga
 ggcacaggctggtatagaaaaagctttacaattgatagagatctggcaggcaaacgcattgcgattaattttgatggcgtctatatga
 35 atgcaacagctctggttaatggcgttaactgggcacacatccgtatggctattcaccgttttcatttgatctgacaggcaatgcaaat
 ttggcggagaaaaacacaattgtcgtcaaagttgaaaatagactgccgtcatcaagatggtattcaggcagcggcatttatagagatg
 ttactgacagttacagatggcgttcattgttgcaataatggcgtcgcaattaaaacaccgtcactggcaacacaaaatggcggag
 40 atgtcacaatgaacctgacaacaaaagtcgcgaatgatacagaagcagcagcgaacattactgaaacagacagttttccgaaa
 ggcggaaaaacggatgcagcaattggcacagttacaacagcatcaaaatcaattgcagcaggcgcatcagcagatgttacaagca
 caattacagcagcaagcccgaactgtggtcaattaaaaacccgaacctgtatacagttagaacagaagttctgaacggaggcaaa
 gttctggatacatatgatacagaatatggcttctgctggacaggctttgatgcaacatcaggcttttactgaatggcgaaaaagtc
 45 actgaaaggcgttagcatgcatcatgatcaaggctcacttggcgcagttgcaaatagacgcgcaattgaaagacaagtcgaaatcct
 gcaaaaaatggcgctcaatagcattcgcacaacacataatccggcagcaaaagcactgattgatgtctgcaatgaaaaaggcgttc
 tgggtgtcgaagaagtctttgatatgtggaaccgcagcaaaaatggcaacacggaagattatggcaaatggttggccaagcaattg
 50 caggcgataatgcagttctgggagggcgataaagatgaaacatgggcgaaatttgatcttacatcaacaattaaccgcgatagaaat
 gcaccgtcagttattatgttggtcactgggcaatgaaatgatggaaggcatttcaggctcagtttcaggcttccggcaacatcagcaa
 aactggttgcatggacaaaagcagcagattcaacaagaccgatgacatatggcgataacaaaattaaagcgaactggaacgaatc
 55 aaatacaatgggcgataatctgacagcaaattggcggagtggttggcacaattattcagatggcgcaaactatgataaaattcgtac
 aacacatccgtcatgggcaatttatggctcagaaacagcatcagcgattaatagccgtggcatttataatagaacaacaggcggagc

acaatcatcagataaacagctgacaagctatgataattcagcagttggctggggagcagttgcatcatcagcatggtatgatgttggt
 cagagagattttgtcgcaggcacatatgtttggacaggatttgattatctggggaaccgacaccgtggaatggcacagggtcaggc
 5 gcagttggctcatggccgtcaccgaaaaatagctattttggcatcgttgatacagcaggcttccgaaagatacatatttttatcag
 agccagtggaatgatgatgttcatacactgcataattctccggcatggaatgaaaatgttggtgcaaaaggctcaggcaataatgttcc
 ggttgctggtttatacagatgcagcgaaagtgaactgtattttacaccgaaaggctcaacagaaaaagactgatcgggcgaataat
 catttcaaaaaaaacaacagcggcaggctatacatatcaagtctatgaaggcagcgataaagattcaacagcgcataaaaaacat
 10 gtatctgacatggaatgttccgtgggcagaaggcacaatttcagcggaaagcgtatgatgaaaaaatcgctgtattccggaaggca
 gcacagaaggcaacgcacagttacaacaacaggcgaagcagcaaaactgaaagcagatgcggatcgcaaaacaattacagcg
 gatggcaagatctgtcatatattgaagtcgatgtcacagatgcaaatggccatattgttccggatgcagcaaatagagtcacattg
 15 atgttaaaggcgcaggcaaaactggttggcgttgataatggctcatcaccggatcatgattcatatcaagcggataaccgcaagcat
 tttcaggcaaaagtcttgcaattgttcagtcaacaaaagaagcaggcgaattacagttacagcaaaagcagatggcctgcaatca
 agcacagttaaaattgcaacaacagcagttccgggaacaagcacagaaaaaacagtcgcagcttttattacagccgcaactattat
 20 gtcaaaacagggaacaaaccgattctgccgtcagatgttgaagttcgctattcagatggaacaagcagatagacaaaacgttacatg
 ggatgcagtttcagatgatcaaattgcaaaagcagggtcattttcagttgcaggcacagttgcaggccaaaaaattagcgttcgcgtc
 acaatgattgatgaaattggcgactg

> SEQ ID NO: 11 nucleotide sequence encoding BIF_1068

gttgaagatgcaacaagaagcagatagcacaacacaaatgtcatcaacaccggaagttgtttattcatcagcggctgatagcaaaaca
 aaatcgcaacagcagttttgatgcgaactggaaatttatgtctgcagatagcgttcaagcacaagatccggcatttgatgattcagca
 30 tggcaacaagttgatctgccgcatgattatagcatcacacagaaatatagccaaagcaatgaagcagaatcagcatatctccggga
 ggcacaggctggtatagaaaaagctttacaattgatagagatctggcaggcaaacgcattgcgattaattttgatggcgtctatatga
 atgcaacagctctggttaatggcgttaaaactgggcacacatccgtatggctattcaccgttttcatttgatctgacaggcaatgcaaaat
 ttggcggagaaaaacacaattgtcgtcaaagttgaaaatagactgccgtcatcaagatggatttcaggcagcggcatttatagagatg
 35 ttacactgacagttacagatggcgttcattgttgcaataatggcgtcgcaattaaaacaccgtcactggcaacacaaaatggcggag
 atgtcacaatgaacctgacaacaaaagtcgcgaatgatacagaagcagcagcgaacattacactgaaacagacagttttccgaaa
 ggcggaaaaacggatgcagcaattggcacagttacaacagcatcaaaatcaattgcagcaggcgcacagcagatgttacaagca
 40 caattacagcagcaagcccgaactgtggtcaattaaaaacccgaacctgtatacagttagaacagaagttctgaacggaggcaaa
 gttctggatacatatgatacagaatatggctttcgtggacaggctttgatgcaacatcaggcctttcactgaatggcgaaaaagtcac
 actgaaaggcgttagcatgcatcatgatcaaggctcacttggcgcagttgcaaatagacgcgcaattgaaagacaagtcgaaatcct
 45 gcaaaaaatggcgctcaatagcattcgcacacacataatccggcagcaaaagcactgattgatgtctgcaatgaaaaaggcgttc
 tggttgtcgaagaagtctttgatatgtggaaccgcagcaaaaaatggcaacacggaagattatggcaaatggtttggccaagcaattg
 caggcgataatgcagttctgggagggcagataaagatgaaacatggcgaaatttgatcttacatcaacaattaaccgcgatagaat
 gcaccgtcagttattatgttggtcactgggcaatgaaatgatggaaggcatttcaggctcagtttcaggcttccggcaacatcagcaa
 50 aactggttgatggacaaaagcagcagattcaacaagaccgatgacatatggcgataacaaaattaaagcgaactggaacgaatc
 aaatacaatgggcgataatctgacagcaaatggcggagttgttggcacaattattcagatggcgcaaatatgataaaattcgtag
 aacacatccgtcatgggcaatttatggctcagaaacagcatcagcgattaatagccgtggcatttataatagaacaacaggcggagc
 55 acaatcatcagataaacagctgacaagctatgataattcagcagttggctggggagcagttgcatcatcagcatggtatgatgttggt
 cagagagattttgtcgcaggcacatatgtttggacaggatttgattatctggggaaccgacaccgtggaatggcacagggtcaggc

gcagttggctcatggccgtcaccgaaaaatagctattttggcatcgttgatacagcaggctttccgaaagatacatatttttatcag
 agccagtggaatgatgatgttcatacactgcatattcttcggcatggaatgaaaatgttggtgcaaaaggctcaggcaataatgttcc
 5 ggttgctggtttatcacagatgcagcgaaagtgaactgtatttacaccgaaaggctcaacagaaaaaagactgatcggcgaaaaat
 cattacaaaaaaaacaacagcggcaggctatacatatcaagtctatgaaggcagcgataaagattcaacagcgcataaaaacat
 gtatctgacatggaatgttccgtgggcagaaaggcacaatttcagcggaaagcgtatgatgaaaataatcgctgattccggaaggca
 gcacagaaggcaacgcacagttacaacaacaggcaagcagcaaaactgaaagcagatgcggatcgcaaaacaattacagcg
 10 gatggcaagatctgtcatatattgaagtcgatgtcacagatgcaaatggccatattgttccggatgcagcaaatagagtcacatttg
 atgttaaaggcgaggcaaaactggttggcgttgataatggctcatcaccggatcatgattcatatcaagcggataaccgcaaaagcat
 tttcaggcaaaagtcttgcaattgttcagtcaacaaaagaagcaggcgaaattacagttacagcaaaagcagatggcctgcaatca
 15 agcacagttaaaattgcaacaacagcagttccgggaacaagcacagaaaaaacagtcgcgagctttattacagccgcaactattat
 gtcaaaacagggaacaaaccgattctgccgtcagatgttgaagttcgctattcagatggaacaagcgatagacaaaacgttacatg
 ggatgcagtttcagatgatcaaattgcaaaagcaggctcattttcagttgcaggcacagttgcaggccaaaaaattagcgttcgcgtc
 acaatgattgatgaaattggcgactgctgaattattcagcaagcacaccggttggcacaccggcagttcttccgggatcaagaccg
 20 gcagtcctgccggatggcacagtcacatcagcaaatttgcagtccattggacaaaaccggcagatacagtcctataatacagcaggc
 acagtcaaagtaccgggaacagcaacagttttggcaaagaatttaaagtcacagcgacaattagagttcaa

> SEQ ID NO: 12 nucleotide sequence encoding BIF_1172

ggtgaagatgcaacaagaagcgatagcacaacacaaatgtcatcaacaccggaagttgtttattcatcagcggctgatagcaaaaca
 aaatcgcaacagcgattttgatggaactggaatttatgtctgcagatagcgttcaagcacaaagatccggcatttgatgattcagca
 30 tggcaacaagttgatctgccgatgattatagcatcacacagaaatatagccaaagcaatgaagcagaatcagcatatcttcggga
 ggcacaggctggtatagaaaaagctttacaattgatagagatctggcaggcaaacgcattgcgattaattttgatggcgtctatatga
 atgcaacagtcgtgttaatggcgttaaactgggcacacatccgtatggctattcaccgttttcatttgatctgacaggcaatgcaaat
 ttggcggagaaaaacacaattgtcgtcaaagttgaaaatagactgccgtcatcaagatggtattcaggcagcggcatttatagagatg
 35 ttacactgacagttacagatggcgttcattgttgcaataatggcgtcgcaattaaaacaccgtcactggcaacacaaaatggcggag
 atgtcacaatgaacctgacaacaaaagtcgcgaatgatacagaagcagcagcgaacattacactgaaacagacagttttccgaaa
 ggcggaaaaacggatgcagcaattggcacagttacaacagcatcaaaatcaattgcagcaggcgcatcagcagatgttacaagca
 40 caattacagcagcaagcccgaactgtggtcaattaaaaacccgaacctgtatacagttagaacagaagttctgaacggaggcaaa
 gttctggatacatatgatacagaatatggcttctgctggacaggctttgatgcaacatcaggctttcactgaatggcgaaaaagtc
 actgaaaggcgttagcatgcatcatgatcaaggctcacttggcgagttgcaaatagacgcgcaattgaaagacaagtcgaaatcct
 45 gcaaaaaatggcgtcaatagcattcgcacaacacataatccggcagcaaaagcactgattgatgtctgcaatgaaaaaggcgttc
 tggttgtcgaagaagtctttgatatgtggaaccgcagcaaaaatggcaacacggaagattatggcaaatggttggccaagcaattg
 caggcgataatgcagttctgggagggcagataaagatgaaacatgggcgaaatttgatcttacatcaacaattaaccgcgatagaat
 gcaccgtcagttattatgtggtcactgggcaatgaaatgatggaaggcatttcaggctcagtttcaggctttccggcaacatcagcaa
 50 aactggttgatggacaaaagcagcagattcaacaagaccgatgacatatggcgataacaaaattaaagcgaactggaacgaatc
 aaatacaatgggagataatctgacagcaaatggcggagttgttggcacaattattcagatggcgcaactatgataaaattcgtag
 aacacatccgtcatgggcaatttatggctcagaaacagcatcagcgattaatagccgtggcatttataatagaacaacaggcggagc
 55 acaatcatcagataaacagctgacaagctatgataattcagcagttggctggggagcagttgcatcatcagcatggatgatgttgtt
 cagagagattttgtcgcaggcacatatgtttggacaggatttgattatctgggcgaaccgacaccgtggaatggcacaggctcaggc

gcagttggctcatggccgtcaccgaaaaatagctattttggcatcgttgatacagcaggctttccgaaagatacatatttttatcag
 agccagtggaatgatgatgttcatactgcatattcttccggcatggaatgaaaatgttgttgcaaaaggctcaggcaataatgttcc
 5 ggtgtcgtttatacagatgcagcgaaagtgaactgtatttacaccgaaaggctcaacagaaaaaagactgatcggcgaaaaat
 catttcaaaaaaacaacagcggcaggctatacatatcaagtctatgaaggcagcgataaagattcaacagcgcataaaaaacat
 gtatctgacatggaatgttccgtgggcagaaggcacaatttcagcgggaagcgtatgatgaaaataatcgctgattccggaaggca
 gcacagaaggcaacgcacatcagttacaacaacaggcaagcagcaaaactgaaagcagatgcggatcgcaaaacaattacagcg
 10 gatggcaaatctgtcatatattgaagtcgatgtcacagatgcaaatggccatattgttccggatgcagcaaatagatgcacatttg
 atgttaaaggcgcaggcaaaactggttggcgttgataatggctcatcaccggatcatgattcatatcaagcggataaccgcaagcat
 tttcaggcaaaagtcttgcaattgttcagtcaacaaaagaagcaggcgaaattacagttacagcaaaagcagatggcctgcaatca
 agcacagttaaaattgcaacaacagcagttccgggaacaagcacagaaaaaacagtcgcagctttattacagccgcaactattat
 15 gtcaaaacaggcaacaaaccgattctgccgtcagatgttgaagttcgctattcagatggaacaagcगतगacaaaacgttacatg
 ggatgcagtttcagatgatcaaattgcaaaagcagggtcattttcagttgcaggcacagttgcaggccaaaaaattagcgttcgcgtc
 acaatgattgatgaaattggcgactgctgaattattcagcaagcacaccggttggcacaccggcagttcttccgggatcaagaccg
 20 gcagtcctgccggatggcacagtcacatcagcaaattttcagttcattggacaaaaccggcagatacagttctataatacagcaggc
 acagtcaaagtaccgggaacagcaacagttttggcaaagaatttaaagtcacagcgacaattagagttcaaagaagccaagttac
 aattggctcatcagtttcaggaaatgactgagactgacacaaaatattccggcagataaacaatcagatacactggatgcgattaa
 25 agatggctcaacaacagttgatgcaatacaggcggaggcgcaaatccgtcagcatggacaaattgggcataattcaaaagcaggc
 cataacacagcggaaattacatttgaatatgcgacagaacaacaactgggccagatcgtcatgtattttttcgcgatagcaatgcag
 ttagatttccggatgctggcaaaacaaaaattcagatc

> SEQ ID NO: 13 nucleotide sequence encoding BIF_1241

gttgaagatgcaacaagaagcgatagcacaacacaaatgtcatcaacaccggaagttgtttattcatcagcggctgatagcaaaaca
 aaatcgcaacagcagttttgatgcgaactggaatttatgtctgcagatagcgttcaagcacagatccggcatttgatgattcagca
 35 tggcaacaagttgatctgccgatgattatagcatcacacagaaatatagccaaagcaatgaagcagaatcagcatatcttccggga
 ggcacaggctggtatagaaaaagctttacaattgatagagatctggcaggcaaacgcattgcgattaattttgatggcgtctatatga
 atgcaacagtcgtggttaattggcgttaaactgggcacacatccgtatggctattcaccgttttcatttgatctgacaggcaatgcaaaat
 40 ttggcggagaaaaacacaattgtcgtcaaagttgaaaatagactgccgtcatcaagatgggtattcaggcagcggcatttatagagatg
 ttactgacagttacagatggcgttcattgttgcaataatggcgtcgcaattaaaacaccgtcactggcaacacaaaatggcggag
 atgtcacaatgaacctgacaacaaaagtcgcgaatgatacagaagcagcagcgaacattactgaaacagacagttttccgaaa
 45 ggcggaaaaacggatgcagcaattggcacagttacaacagcatcaaaatcaattgcagcaggcgcatcagcagatgttacaagca
 caattacagcagcaagcccgaactgtggtcaattaaaaacccgaacctgtatacagttagaacagaagttctgaacggaggcaaa
 gttctggatacatatgatacagaatatggcttctgctggacaggctttgatgcaacatcaggcttttactgaatggcgaaaaagtc
 actgaaaggcgttagcatgcatcatgatcaaggctcacttggcgcagttgcaaatagacgcgcaattgaaagacaagtcgaaatcct
 50 gcaaaaaatggcgctcaatagcattcgcacacacataatccggcagcaaaagcactgattgatgtctgcaatgaaaaaggcgttc
 tggttgtcgaagaagtctttgatatgtggaaccgcagcaaaaaatggcaacacggaagattatggcaaatggtttggccaagcaattg
 caggcgataatgcagttctgggaggcgataaagatgaaacatgggcgaaatttgatcttacatcaacaattaaccgcgatagaaat
 55 gcaccgtcagttattatgtggtcactgggcaatgaaatgatggaaggcatttcaggctcagtttcaggctttccggcaacatcagcaa
 aactggttgcattggacaaaagcagcagattcaacaagaccgatgacatatggcgataacaaaattaaagcgaactggaacgaatc

aaatacaatgggcgataatctgacagcaaatggcggagttgttggcacaattattcagatggcgcaaactatgataaaattcgtac
 aacacatccgtcatgggcaatttatggctcagaaacagcatcagcgattaatagccgtggcatttataatagaacaacaggcggagc
 5 acaatcatcagataaacagctgacaagctatgataattcagcagttggctggggagcagttgcatcatcagcatggatgatgttgtt
 cagagagattttgtcgcaggcacatatgtttggacaggatttgattatctggggaaccgacaccgtggaatggcacagggtcaggc
 gcagttggctcatggccgtcaccgaaaaatagctattttggcatcgttgatacagcaggctttccgaaagatacatattttttatcag
 10 agccagtggatgatgatgttcatactgcataattcttcggcatggaatgaaaatgttgttgcaaaaggctcaggcaataatgttcc
 gggtgtcgtttatacagatgcagcgaaagtgaactgtattttacaccgaaaggctcaacagaaaaaagactgatcggcgaaaaat
 catttcaaaaaaaacaacagcggcaggctatacatatcaagtctatgaaggcagcgataaagattcaacagcgcataaaaaacat
 gtatctgacatggaatgttccgtgggcagaaggcacaatttcagcgggaagcgtatgatgaaaaaatcgctgtattccggaaggca
 15 gcacagaaggcaacgcacagttacaacaacaggcgaagcagcaaaaactgaaagcagatgcggatcgcaaaacaattacagcg
 gatggcaagatctgtcatatattgaagtcgatgtcacagatgcaaatggccatattgttccggatgcagcaaatagagtcacattg
 atgttaaaggcgaggcaaaactggttggcgttgataatggctcatcaccggatcatgattcatatcaagcggataaccgcaaagcat
 20 tttcaggcaaaagtcttgcaattgttcagtcaaaaaagaagcaggcgaaattacagttacagcaaaagcagatggcctgcaatca
 agcacagttaaaattgcaacaacagcagttccgggaacaagcacagaaaaaacagtcgcgagcttttattacagccgcaactattat
 gtcaaaacagggaacaaaccgattctgccgtcagatgttgaagttcgctattcagatggaacaagcgatagacaaaacgttacatg
 ggatgcagtttcagatgatcaaattgcaaaagcagggtcattttcagttgcaggcacagttgcaggccaaaaaattagcgttcgcgtc
 25 acaatgattgatgaaattggcgactgctgaattattcagcaagcacaccgggtggcacaccggcagttcttcgggatcaagaccg
 gcagtcctgccggatggcacagtcacatcagcaaattttcagtcattggacaaaaccggcagatacagtcataatacagcaggc
 acagtcaaaagtaccgggaacagcaacagtttttgcaagaatttaaaagtcacagcgacaattagagttcaaaagaagccaagttac
 30 aattggctcatcagtttcaggaaatgactgagactgacacaaaatattccggcagataaacaatcagatacactggatgcgattaa
 agatggctcaacaacagttgatgcaatacaggcggaggcgcaaatccgtcagcatggacaaattgggcatattcaaaagcaggc
 cataacacagcggaaattacatttgaatatgacagagaacaacaactgggccagatcgatgtatttttttcgcatagcaatgcag
 ttagatttccggatgctggcaaaacaaaaattcagatcagcgagatggcaaaattggacagatctggcagcaacagaaacaatt
 35 gcagcgcaagaatcaagcgatagagtcaaaccgtatacatatgattttgcaccgggttggcgcaacattgttaaagtgacagtcaca
 aacgcagatacaacaacaccgtcaggcgttgtttgcgcaggcctgacagaaattgaactgaaaacagcgaca

> SEQ ID NO: 14 nucleotide sequence encoding BIF_1326

40 gttgaagatgcaacaagaagcgatagcacaacacaaatgtcatcaaacaccggaagttgtttattcatcagcggctgatagcaaaaca
 aaatcgcaagcagattttgatgcgaactggaaatttatgtgtcagatagcgttcaagcacaagatccggcatttgatgattcagca
 45 tggcaacaagttgatctgccgatgattatagcatcacacagaaatatagccaaagcaatgaagcagaatcagcatatcttcggga
 ggcacaggctggtatagaaaaagctttacaattgatagagatctggcaggcaaacgcattgcgattaattttgatggcgtctatatga
 atgcaacagtcgtgttaattggcgttaactgggcacacatccgtatggctattcaccgttttcatttgatctgacaggcaatgcaaaat
 50 ttggcggagaaaaacacaattgtcgtcaaagttgaaaatagactgccgtcatcaagatgggtattcaggcagcggcatttatagagatg
 ttacttgacagttacagatggcgttcattgttgcaataatggcgtcgcaattaaaacaccgtcactggcaacacaaaatggcggag
 atgtcacaatgaacctgacaacaaaagtcgcgaatgatacagaagcagcagcgaacattactgaaacagacagttttccgaaa
 55 ggcggaaaaacggatgcagcaattggcacagttacaacagcatcaaaatcaattgcagcaggcgcatcagcagatgttacaagca
 caattacagcagcaagcccgaactgtggtcaattaaaaacccgaacctgtatacagttagaacagaagttctgaacggaggcaaa
 gttctggatacatatgatacagaatatggcttctgctggacaggctttgatgcaacatcaggccttttactgaatggcgaaaaagtcaa

actgaaaggcgttagcatgcatcatgatcaaggctcacttggcgcagttgcaaatagacgcgcaattgaaagacaagtcgaaatcct
 gcaaaaaatgggcgtcaatagcattcgcacaacacataatccggcagcaaaagcactgattgatgtctgcaatgaaaaaggcgttc
 5 tggttgtcgaagaagtctttgatatgtggaaccgcagcaaaaatggcaacacggaagattatggcaaatggttggccaagcaattg
 caggcgataatgcagttctgggagggcgataaagatgaaacatgggcgaaatttgatcttacatcaacaattaaccgcgatagaat
 gcaccgtcagttattatgttggtcactgggcaatgaaatgatggaaggcatttcagggtcagtttcagggtttccggcaacatcagcaa
 aactggttgcatggacaaaagcagcagattcaacaagaccgatgacatatggcgataacaaaattaaagcgaactggaacgaatc
 10 aaatacaatgggcgataatctgacagcaaatggcggagttgttggcacaaaattatcagatggcgcaaatatgataaaattcgtag
 aacacatccgtcatgggcaatttatggctcagaaacagcatcagcgattaatagccgtggcattataatagaacaacaggcggagc
 acaatcatcagataaacagctgacaagctatgataattcagcagttggctgggagcagttgcatcatcagcatggtatgatgttgtt
 15 cagagagattttgtcgcaggcacatatgtttggacaggatttgattatctgggcgaaccgacaccgtggaatggcacagggtcaggc
 gcagttggctcatggccgtcaccgaaaaatagctattttggcatcgttgatacagcaggctttccgaaagatacatattttttatcag
 agccagtggaaatgatgatgttcatacactgcatattctccggcatggaatgaaaatgttgttgcaaaaggctcaggcaataatgttcc
 ggttgcgtttatacagatgcagcgaaagtgaactgtattttacaccgaaaggctcaacagaaaaaagactgatcggcgaaaaat
 20 cattacaaaaaaacaacagcggcaggctatacatatcaagtctatgaaggcagcgataaagattcaacagcgcataaaaaacat
 gtatctgacatggaatgttccgtgggcagaaggcacaatttcagcggaaagcgtatgatgaaaataatcgctgattccggaaggca
 gcacagaaggcaacgcacagttacaacaacaggcgaagcagcaaaactgaaagcagatgcggatcgcaaaacaattacagcg
 25 gatggcaagatctgtcatatattgaagtcgatgtcacagatgcaaatggccatattgttccggatgcagcaaatagagtcacatttg
 atgttaaaggcgcaggcaaaactggttggcgttgataatggctcatcaccggatcatgattcatatcaagcggataaccgcaaagcat
 tttcaggcaaaagtcctggcaattgttcagtcacaaaaagaagcaggcgaaattacagttacagcaaaagcagatggcctgcaatca
 30 agcacagttaaaattgcaacaacagcagttccgggaacaagcacagaaaaaacagtcgcgagctttattacagccgcaactattat
 gtcaaaacagggaacaaacagattctgccgtcagatgttgaagttcgctattcagatggaacaagcagatagacaaaacgttacatg
 ggatgcagtttcagatgatcaaattgcaaaagcagggtcattttcagttgcaggcacagttgcaggccaaaaaattagcgttcgcgtc
 acaatgattgatgaaattggcgcactgctgaattattcagcaagcacaccgggttgccacaccggcagttcttccgggatcaagaccg
 35 gcagtcctgccggatggcacagtcacatcagcaaattttgagtcatttgacaaaaccggcagatacagtcataatacagcaggc
 acagtcaaaagtaccgggaacagcaacagttttggcaagaatttaaagtcacagcgacaattagagttcaaaagaagccaagttac
 aattggctcatcagtttcaggaaatgactgagactgacacaaaatattccggcagataaacaatcagatacactggatgcgattaa
 40 agatggctcaacaacagttgatgcaatacaggcggaggcgcaaatccgtcagcatggacaaattgggcatattcaaaagcaggc
 cataacacagcggaaattacatttgaatatgcgacagaacaacaactgggccagatcgatgtattttttcgcgatagcaatgcag
 ttagatttccggatgctggcaaaacaaaaattcagatcagcgcagatggcaaaattggacagatctggcagcaacagaaacaatt
 45 gcagcgcaagaatcaagcagatagagtcacacccgtatacatatgattttgcaccgggttgccgcaacattgttaaagtgacagtcaca
 aacgcagatacaacaacaccgtcaggcgttgtttgcgaggcctgacagaaattgaactgaaaacagcgacaagcaaatgtgtcac
 aaatacatcagcagcactgtcatcacttacagtcfaatggcacaagaatttcagattcagttctggcagcagggtcatataacacaccg
 gcaattatcgagatgttaaagcggaaggcgaaggcaatgcaagcgttacagtccttccggcacatgataatgttattcgcgtcatta
 50 cagaaagcgaagatcatgtcacacgcaaaacatttacaatcaacctgggcacagaacaagaattt

> SEQ ID NO: 15 forward primer for generation of BIF variants

GGGGTAACTAGTGGAAGATGCAACAAGAAG

> SEQ ID NO: 16 reverse primer for BIF917

GCGCTTAATTAATTATGTTTTTCTGTGCTTGTTT

> SEQ ID NO: 17 reverse primer for BIF995

GCGCTTAATTAATTACAGTGCGCCAATTCATCAATCA

>SEQ ID NO: 18 reverse primer for BIF1068

GCGCTTAATTAATTATTGAACTCTAATTGTCGCTG

>SEQ ID NO: 19 reverse primer for BIF1241

GCGCTTAATTAATTATGTCGCTGTTTTAGTTCAAT

> SEQ ID NO: 20 reverse primer for BIF1326

GCGCTTAATTAATTAATAATTCTTGTCTGTGCCCA

> SEQ ID NO: 21 reverse primer for BIF1478

GCGCTTAATTAATTATCTCAGTCTAATTTGCTTGCGC

> SEQ ID NO: 22 Bifidobacterium bifidum BIF1750

vedatrsdsttqmsstpevyssavdskqrntsdndanwkfmlsdsvqaqdpafddsawqqvdlphdysitqkysqsneaesa
ylpggtgwyrksftidrdlagkriainfdgvymnatvwfngvklgthpygyspfsdltgnakfggentivvkenrlpssrwysgsg
iyrdvtltvdgvhvgngvaiktpslatqnggdvtmnltkvandteaaanitlkqtvfpkggktdaaigtvttasksiaagasadv
stitaasplksiknplytvrtevlnggkvldtydteygrwtgfdatsgfsngekvklkgvsmhhdqgslgavanrraierqveil
qkmgvnsirtthnpaakalidvcnekglvveevfdmwnrskngntedygkwfgqaiagdnalvggdkdetwakfdltstinrdr
napsvimwslgnemmegisgsvsgfpatsaklvawtkaadstrpmtgygdnkikanwnesntmgdnltanggvvgtnysdga
nydkirtthpswaiygsetasainsrgiynrttggaqssdkqltsydnsgwgavassawydvvqrdfvagtyvwtgfdylgept
pwngtgsgavgswwpspknsyfgivdtagfpkdttyfyqsqwnddvhtlilpawnnenvvaksgnnpvvytdaakvklyft
pkgstekrligeksftkktaagytyqvyegsddstahknmyltwnvpwaegtisaeaydenrlipegstegnasvtttgaakl
kadadrktitadgkdlisyievdvtdanghivpdaanrvtdvkgagklvgvndngsspdhdsyqadnrkafsgkvlaivqstkeage
itvtakadqlsstvkiattavpgtstektrsvfyysrnyyvtgtnkpilpsdvevrysdgtsdrqnvtdavsdqqiakagsfsvagt
vagqkisvrvtmideigallnysastpvgtpavlpdgtvtsanfahwtkpadtyntagtvkvpgtatvfgkefkvtat
irvqrsqvtigssvsgnalrltqnipadkqsdtdaikdgsttdantggganpsawtnwayskaghntaeitfeyateqqlgqivm
yffrdsnavrfdagtkkiqisadgknwtdlaatetiaaqaessdrvkpytydfapvgatfvkvtvnadtttsgvvcaglteielktat
skfvntsaalssltvngtkvdsdslaagsyntpailadvkaegegnasvtpahdnvirvitesedhvrktftinlgtetqefpadsd
erdypaadmvtvtvgseqtsgtategpkkfavdgntstywhsnwtpptvndlwiafelqkptklalrylprpagskngsvteykvq
vsddgtnwtdagsgtwtdygwklafnqpvttkhvrkavhtyadsgndkfmsaseirrkavdttddisgatvtvpakltvdrvda
dhpatfatkdvtvtlgdatlrygvdyllidyagntavgkatvtvrgidkysgtvaktftielknapeptltsvsvtktpsklityvvgda
fdpaglqlqhdrqadrppqplvgeqadergltcgrcdrveqlrkhenreahrtgldhlefvgaadgavgeqatfkvhvhadqgdgr
hddaderdidphvpvdhavgelaraachhvlgirvdthrlkasgfiqpaddmaeidritgfrferhvg

> SEQ ID NO: 23 The signal sequence of extracellular lactase from Bifidobacterium bifidum DSM20215

Vrskklwisllfaliftmafgtssaq

[0289] Various preferred features and embodiments of the present invention will now be described with reference to the following numbered paragraphs (paras).

1. A polypeptide having transgalactosylating activity selected from the group consisting of:

a. a polypeptide comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues,

b. a polypeptide comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2, wherein said polypeptide consists of at most 975 amino acid residues,

c. a polypeptide comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues,

d. a polypeptide encoded by a polynucleotide that hybridizes under at least low stringency conditions with i) the nucleic acid sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5; or ii) the complementary strand of i),

e. a polypeptide encoded by a polynucleotide comprising a nucleotide sequence having at least 70% identity to the nucleotide sequence encoding for the polypeptide of SEQ ID NO: 1, 2, 3, 4 or 5 or the nucleotide sequence comprised in SEQ ID NO: 9, 10, 11, 12 or 13 encoding a mature polypeptide, and

f. a polypeptide comprising a deletion, insertion and/or conservative substitution of one or more amino acid residues of SEQ ID NO: 1, 2, 3, 4 or 5.

2. The polypeptide according to para 1 comprising an amino acid sequence having at least 90% sequence identity with SEQ ID NO: 1, wherein said polypeptide consists of at most 980 amino acid residues.

3. The polypeptide according to para 1 comprising an amino acid sequence having at least 97% sequence identity with SEQ ID NO: 2, wherein said polypeptide consists of at most 975 amino acid residues.

4. The polypeptide according to para 1 comprising an amino acid sequence having at least 96.5% sequence identity with SEQ ID NO: 3, wherein said polypeptide consists of at most 1300 amino acid residues.

5. The polypeptide according to any one of the preceding paras, which consist of the amino acid sequence of SEQ ID NO: 1, SEQ ID NO:2, SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 4.

6. The polypeptide according to any one of the preceding paras, which consist of the amino acid sequence of SEQ ID NO:1, or SEQ ID NO: 2.

7. The polypeptide according to any one of the preceding paras, wherein said polypeptide has a ratio of transgalactosylation activity above 100%.

8. A nucleic acid capable of encoding a polypeptide according to any one of the preceding paras.

9. An expression vector comprising a nucleic acid according to para 8, or capable of expressing a polypeptide according to any one paras 1-7.

10. A cell capable of expressing a polypeptide according to any one of paras 1-7.

11. A method of expressing a polypeptide, the method comprising obtaining a cell according para 10 and expressing the polypeptide from the cell, and optionally purifying the polypeptide.

12. A composition comprising a polypeptide as defined in any one of paras 1-7, preferably a food composition, more preferably a dairy product.

13. A method for producing a food product by treating a substrate comprising lactose with a polypeptide as defined in any one of paras 1-7.

14. The method according to para 13 for producing a dairy product by treating a milk-based substrate comprising lactose with a polypeptide as defined in any one of paras 1-7.

15. A process for producing galacto-oligosaccharides, comprising contacting of a polypeptide of any one of paras 1-7 or a cell of para 10 with a milk-based solution comprising lactose.

SEQUENCE LISTING

<110> DuPont Nutrition BioSciences ApS

<120> POLYPEPTIDES HAVING TRANSGALACTOSYLATING ACTIVITY

<130> NB40269WOPCT

<160> 23

<170> BiSSAP 1.0

<210> 1

<211> 887

<212> PRT

<213> Bifidobacterium bifidum

<220>

<221> SOURCE

<222> 1..887

<223> /mol_type="protein"

/organism="Bifidobacterium bifidum"

<400> 1

```

Val Glu Asp Ala Thr Arg Ser Asp Ser Thr Thr Gln Met Ser Ser Thr
1      5      10      15
Pro Glu Val Val Tyr Ser Ser Ala Val Asp Ser Lys Gln Asn Arg Thr
20     25     30
Ser Asp Phe Asp Ala Asn Trp Lys Phe Met Leu Ser Asp Ser Val Gln
35     40     45
Ala Gln Asp Pro Ala Phe Asp Asp Ser Ala Trp Gln Gln Val Asp Leu
50     55     60
Pro His Asp Tyr Ser Ile Thr Gln Lys Tyr Ser Gln Ser Asn Glu Ala
65     70     75     80
Glu Ser Ala Tyr Leu Pro Gly Gly Thr Gly Trp Tyr Arg Lys Ser Phe
85     90     95
Thr Ile Asp Arg Asp Leu Ala Gly Lys Arg Ile Ala Ile Asn Phe Asp
100    105    110
Gly Val Tyr Met Asn Ala Thr Val Trp Phe Asn Gly Val Lys Leu Gly
115    120    125
Thr His Pro Tyr Gly Tyr Ser Pro Phe Ser Phe Asp Leu Thr Gly Asn
130    135    140
Ala Lys Phe Gly Gly Glu Asn Thr Ile Val Val Lys Val Glu Asn Arg
145    150    155    160
Leu Pro Ser Ser Arg Trp Tyr Ser Gly Ser Gly Ile Tyr Arg Asp Val
165    170    175
Thr Leu Thr Val Thr Asp Gly Val His Val Gly Asn Asn Gly Val Ala
180    185    190
Ile Lys Thr Pro Ser Leu Ala Thr Gln Asn Gly Gly Asp Val Thr Met
195    200    205
Asn Leu Thr Thr Lys Val Ala Asn Asp Thr Glu Ala Ala Asn Ile
210    215    220
Thr Leu Lys Gln Thr Val Phe Pro Lys Gly Gly Lys Thr Asp Ala Ala
225    230    235    240
Ile Gly Thr Val Thr Thr Ala Ser Lys Ser Ile Ala Ala Gly Ala Ser
245    250    255
Ala Asp Val Thr Ser Thr Ile Thr Ala Ala Ser Pro Lys Leu Trp Ser
260    265    270
Ile Lys Asn Pro Asn Leu Tyr Thr Val Arg Thr Glu Val Leu Asn Gly
275    280    285
Gly Lys Val Leu Asp Thr Tyr Asp Thr Glu Tyr Gly Phe Arg Trp Thr
290    295    300
Gly Phe Asp Ala Thr Ser Gly Phe Ser Leu Asn Gly Glu Lys Val Lys

```

EP 3 473 713 A1

	305				310				315			320
	Leu	Lys	Gly	Val	Ser	Met	His	His	Asp	Gln	Gly	Ser
					325				330			335
	Ala	Asn	Arg	Arg	Ala	Ile	Glu	Arg	Gln	Val	Glu	Ile
5					340				345			350
	Gly	Val	Asn	Ser	Ile	Arg	Thr	Thr	His	Asn	Pro	Ala
					355				360			365
	Ile	Asp	Val	Cys	Asn	Glu	Lys	Gly	Val	Leu	Val	Val
					370				375			380
10	Asp	Met	Trp	Asn	Arg	Ser	Lys	Asn	Gly	Asn	Thr	Glu
	385					390						395
	Trp	Phe	Gly	Gln	Ala	Ile	Ala	Gly	Asp	Asn	Ala	Val
					405				410			415
	Lys	Asp	Glu	Thr	Trp	Ala	Lys	Phe	Asp	Leu	Thr	Ser
					420				425			430
15	Asp	Arg	Asn	Ala	Pro	Ser	Val	Ile	Met	Trp	Ser	Leu
					435				440			445
	Met	Glu	Gly	Ile	Ser	Gly	Ser	Val	Ser	Gly	Phe	Pro
					450				455			460
	Lys	Leu	Val	Ala	Trp	Thr	Lys	Ala	Ala	Asp	Ser	Thr
	465					470						475
20	Tyr	Gly	Asp	Asn	Lys	Ile	Lys	Ala	Asn	Trp	Asn	Glu
					485							490
	Gly	Asp	Asn	Leu	Thr	Ala	Asn	Gly	Gly	Val	Val	Gly
					500				505			510
	Asp	Gly	Ala	Asn	Tyr	Asp	Lys	Ile	Arg	Thr	Thr	His
					515				520			525
25	Ile	Tyr	Gly	Ser	Glu	Thr	Ala	Ser	Ala	Ile	Asn	Ser
					530				535			540
	Asn	Arg	Thr	Thr	Gly	Gly	Ala	Gln	Ser	Ser	Asp	Lys
	545					550						555
	Tyr	Asp	Asn	Ser	Ala	Val	Gly	Trp	Gly	Ala	Val	Ala
					565							570
30	Tyr	Asp	Val	Val	Gln	Arg	Asp	Phe	Val	Ala	Gly	Thr
					580				585			590
	Gly	Phe	Asp	Tyr	Leu	Gly	Glu	Pro	Thr	Pro	Trp	Asn
					595				600			605
	Gly	Ala	Val	Gly	Ser	Trp	Pro	Ser	Pro	Lys	Asn	Ser
					610				615			620
35	Val	Asp	Thr	Ala	Gly	Phe	Pro	Lys	Asp	Thr	Tyr	Tyr
	625					630						635
	Gln	Trp	Asn	Asp	Asp	Val	His	Thr	Leu	His	Ile	Leu
					645				650			655
40	Glu	Asn	Val	Val	Ala	Lys	Gly	Ser	Gly	Asn	Asn	Val
					660				665			670
	Tyr	Thr	Asp	Ala	Ala	Lys	Val	Lys	Leu	Tyr	Phe	Thr
					675				680			685
	Thr	Glu	Lys	Arg	Leu	Ile	Gly	Glu	Lys	Ser	Phe	Thr
					690				695			700
45	Ala	Ala	Gly	Tyr	Thr	Tyr	Gln	Val	Tyr	Glu	Gly	Ser
	705					710						715
	Thr	Ala	His	Lys	Asn	Met	Tyr	Leu	Thr	Trp	Asn	Val
					725							730
	Gly	Thr	Ile	Ser	Ala	Glu	Ala	Tyr	Asp	Glu	Asn	Asn
					740				745			750
50	Glu	Gly	Ser	Thr	Glu	Gly	Asn	Ala	Ser	Val	Thr	Thr
					755				760			765
	Ala	Lys	Leu	Lys	Ala	Asp	Ala	Asp	Arg	Lys	Thr	Ile
					770				775			780
	Lys	Asp	Leu	Ser	Tyr	Ile	Glu	Val	Asp	Val	Thr	Asp
	785					790						795
55	Ile	Val	Pro	Asp	Ala	Ala	Asn	Arg	Val	Thr	Phe	Asp
					805							810
												815

EP 3 473 713 A1

Gly Lys Leu Val Gly Val Asp Asn Gly Ser Ser Pro Asp His Asp Ser
 820 825 830
 Tyr Gln Ala Asp Asn Arg Lys Ala Phe Ser Gly Lys Val Leu Ala Ile
 835 840 845
 Val Gln Ser Thr Lys Glu Ala Gly Glu Ile Thr Val Thr Ala Lys Ala
 850 855 860
 Asp Gly Leu Gln Ser Ser Thr Val Lys Ile Ala Thr Thr Ala Val Pro
 865 870 875 880
 Gly Thr Ser Thr Glu Lys Thr
 885

 <210> 2
 <211> 965
 <212> PRT
 <213> Bifidobacterium bifidum

 <220>
 <221> SOURCE
 <222> 1..965
 <223> /mol_type="protein"
 /organism="Bifidobacterium bifidum"

 <400> 2
 Val Glu Asp Ala Thr Arg Ser Asp Ser Thr Thr Gln Met Ser Ser Thr
 1 5 10 15
 Pro Glu Val Val Tyr Ser Ser Ala Val Asp Ser Lys Gln Asn Arg Thr
 20 25 30
 Ser Asp Phe Asp Ala Asn Trp Lys Phe Met Leu Ser Asp Ser Val Gln
 35 40 45
 Ala Gln Asp Pro Ala Phe Asp Asp Ser Ala Trp Gln Gln Val Asp Leu
 50 55 60
 Pro His Asp Tyr Ser Ile Thr Gln Lys Tyr Ser Gln Ser Asn Glu Ala
 65 70 75 80
 Glu Ser Ala Tyr Leu Pro Gly Gly Thr Gly Trp Tyr Arg Lys Ser Phe
 85 90 95
 Thr Ile Asp Arg Asp Leu Ala Gly Lys Arg Ile Ala Ile Asn Phe Asp
 100 105 110
 Gly Val Tyr Met Asn Ala Thr Val Trp Phe Asn Gly Val Lys Leu Gly
 115 120 125
 Thr His Pro Tyr Gly Tyr Ser Pro Phe Ser Phe Asp Leu Thr Gly Asn
 130 135 140
 Ala Lys Phe Gly Gly Glu Asn Thr Ile Val Val Lys Val Glu Asn Arg
 145 150 155 160
 Leu Pro Ser Ser Arg Trp Tyr Ser Gly Ser Gly Ile Tyr Arg Asp Val
 165 170 175
 Thr Leu Thr Val Thr Asp Gly Val His Val Gly Asn Asn Gly Val Ala
 180 185 190
 Ile Lys Thr Pro Ser Leu Ala Thr Gln Asn Gly Gly Asp Val Thr Met
 195 200 205
 Asn Leu Thr Thr Lys Val Ala Asn Asp Thr Glu Ala Ala Ala Asn Ile
 210 215 220
 Thr Leu Lys Gln Thr Val Phe Pro Lys Gly Gly Lys Thr Asp Ala Ala
 225 230 235 240
 Ile Gly Thr Val Thr Thr Ala Ser Lys Ser Ile Ala Ala Gly Ala Ser
 245 250 255
 Ala Asp Val Thr Ser Thr Ile Thr Ala Ala Ser Pro Lys Leu Trp Ser
 260 265 270
 Ile Lys Asn Pro Asn Leu Tyr Thr Val Arg Thr Glu Val Leu Asn Gly
 275 280 285
 Gly Lys Val Leu Asp Thr Tyr Asp Thr Glu Tyr Gly Phe Arg Trp Thr
 290 295 300
 Gly Phe Asp Ala Thr Ser Gly Phe Ser Leu Asn Gly Glu Lys Val Lys
 305 310 315 320

EP 3 473 713 A1

	Leu	Lys	Gly	Val	Ser	Met	His	His	Asp	Gln	Gly	Ser	Leu	Gly	Ala	Val	
					325					330					335		
	Ala	Asn	Arg	Arg	Ala	Ile	Glu	Arg	Gln	Val	Glu	Ile	Leu	Gln	Lys	Met	
				340					345					350			
5	Gly	Val	Asn	Ser	Ile	Arg	Thr	Thr	His	Asn	Pro	Ala	Ala	Lys	Ala	Leu	
			355					360					365				
	Ile	Asp	Val	Cys	Asn	Glu	Lys	Gly	Val	Leu	Val	Val	Glu	Glu	Val	Phe	
		370					375					380					
	Asp	Met	Trp	Asn	Arg	Ser	Lys	Asn	Gly	Asn	Thr	Glu	Asp	Tyr	Gly	Lys	
	385					390					395					400	
10	Trp	Phe	Gly	Gln	Ala	Ile	Ala	Gly	Asp	Asn	Ala	Val	Leu	Gly	Gly	Asp	
				405					410						415		
	Lys	Asp	Glu	Thr	Trp	Ala	Lys	Phe	Asp	Leu	Thr	Ser	Thr	Ile	Asn	Arg	
				420					425					430			
	Asp	Arg	Asn	Ala	Pro	Ser	Val	Ile	Met	Trp	Ser	Leu	Gly	Asn	Glu	Met	
			435					440					445				
15	Met	Glu	Gly	Ile	Ser	Gly	Ser	Val	Ser	Gly	Phe	Pro	Ala	Thr	Ser	Ala	
		450				455					460						
	Lys	Leu	Val	Ala	Trp	Thr	Lys	Ala	Ala	Asp	Ser	Thr	Arg	Pro	Met	Thr	
		465				470				475						480	
	Tyr	Gly	Asp	Asn	Lys	Ile	Lys	Ala	Asn	Trp	Asn	Glu	Ser	Asn	Thr	Met	
				485					490						495		
20	Gly	Asp	Asn	Leu	Thr	Ala	Asn	Gly	Gly	Val	Val	Gly	Thr	Asn	Tyr	Ser	
				500					505					510			
	Asp	Gly	Ala	Asn	Tyr	Asp	Lys	Ile	Arg	Thr	Thr	His	Pro	Ser	Trp	Ala	
			515					520					525				
25	Ile	Tyr	Gly	Ser	Glu	Thr	Ala	Ser	Ala	Ile	Asn	Ser	Arg	Gly	Ile	Tyr	
		530				535					540						
	Asn	Arg	Thr	Thr	Gly	Gly	Ala	Gln	Ser	Ser	Asp	Lys	Gln	Leu	Thr	Ser	
		545				550					555					560	
	Tyr	Asp	Asn	Ser	Ala	Val	Gly	Trp	Gly	Ala	Val	Ala	Ser	Ser	Ala	Trp	
				565						570					575		
30	Tyr	Asp	Val	Val	Gln	Arg	Asp	Phe	Val	Ala	Gly	Thr	Tyr	Val	Trp	Thr	
			580					585						590			
	Gly	Phe	Asp	Tyr	Leu	Gly	Glu	Pro	Thr	Pro	Trp	Asn	Gly	Thr	Gly	Ser	
			595					600					605				
	Gly	Ala	Val	Gly	Ser	Trp	Pro	Ser	Pro	Lys	Asn	Ser	Tyr	Phe	Gly	Ile	
			610				615					620					
35	Val	Asp	Thr	Ala	Gly	Phe	Pro	Lys	Asp	Thr	Tyr	Tyr	Phe	Tyr	Gln	Ser	
					625		630				635					640	
	Gln	Trp	Asn	Asp	Asp	Val	His	Thr	Leu	His	Ile	Leu	Pro	Ala	Trp	Asn	
				645					650						655		
	Glu	Asn	Val	Val	Ala	Lys	Gly	Ser	Gly	Asn	Asn	Val	Pro	Val	Val	Val	
				660					665					670			
40	Tyr	Thr	Asp	Ala	Ala	Lys	Val	Lys	Leu	Tyr	Phe	Thr	Pro	Lys	Gly	Ser	
			675					680					685				
	Thr	Glu	Lys	Arg	Leu	Ile	Gly	Glu	Lys	Ser	Phe	Thr	Lys	Lys	Thr	Thr	
			690				695					700					
	Ala	Ala	Gly	Tyr	Thr	Tyr	Gln	Val	Tyr	Glu	Gly	Ser	Asp	Lys	Asp	Ser	
					705		710				715					720	
45	Thr	Ala	His	Lys	Asn	Met	Tyr	Leu	Thr	Trp	Asn	Val	Pro	Trp	Ala	Glu	
				725						730					735		
	Gly	Thr	Ile	Ser	Ala	Glu	Ala	Tyr	Asp	Glu	Asn	Asn	Arg	Leu	Ile	Pro	
				740					745					750			
	Glu	Gly	Ser	Thr	Glu	Gly	Asn	Ala	Ser	Val	Thr	Thr	Thr	Gly	Lys	Ala	
				755				760					765				
50	Ala	Lys	Leu	Lys	Ala	Asp	Ala	Asp	Arg	Lys	Thr	Ile	Thr	Ala	Asp	Gly	
					770		775					780					
	Lys	Asp	Leu	Ser	Tyr	Ile	Glu	Val	Asp	Val	Thr	Asp	Ala	Asn	Gly	His	
					785		790				795					800	
	Ile	Val	Pro	Asp	Ala	Ala	Asn	Arg	Val	Thr	Phe	Asp	Val	Lys	Gly	Ala	
					805					810					815		
55	Gly	Lys	Leu	Val	Gly	Val	Asp	Asn	Gly	Ser	Ser	Pro	Asp	His	Asp	Ser	

EP 3 473 713 A1

820 825 830
 Tyr Gln Ala Asp Asn Arg Lys Ala Phe Ser Gly Lys Val Leu Ala Ile
 835 840 845
 Val Gln Ser Thr Lys Glu Ala Gly Glu Ile Thr Val Thr Ala Lys Ala
 850 855 860
 Asp Gly Leu Gln Ser Ser Thr Val Lys Ile Ala Thr Thr Ala Val Pro
 865 870 875 880
 Gly Thr Ser Thr Glu Lys Thr Val Arg Ser Phe Tyr Tyr Ser Arg Asn
 885 890 895
 Tyr Tyr Val Lys Thr Gly Asn Lys Pro Ile Leu Pro Ser Asp Val Glu
 900 905 910
 Val Arg Tyr Ser Asp Gly Thr Ser Asp Arg Gln Asn Val Thr Trp Asp
 915 920 925
 Ala Val Ser Asp Asp Gln Ile Ala Lys Ala Gly Ser Phe Ser Val Ala
 930 935 940
 Gly Thr Val Ala Gly Gln Lys Ile Ser Val Arg Val Thr Met Ile Asp
 945 950 955 960
 Glu Ile Gly Ala Leu
 965

 <210> 3
 <211> 1038
 <212> PRT
 <213> Bifidobacterium bifidum

 <220>
 <221> SOURCE
 <222> 1..1038
 <223> /mol_type="protein"
 /organism="Bifidobacterium bifidum"

 <400> 3
 Val Glu Asp Ala Thr Arg Ser Asp Ser Thr Thr Gln Met Ser Ser Thr
 1 5 10 15
 Pro Glu Val Val Tyr Ser Ser Ala Val Asp Ser Lys Gln Asn Arg Thr
 20 25 30
 Ser Asp Phe Asp Ala Asn Trp Lys Phe Met Leu Ser Asp Ser Val Gln
 35 40 45
 Ala Gln Asp Pro Ala Phe Asp Asp Ser Ala Trp Gln Gln Val Asp Leu
 50 55 60
 Pro His Asp Tyr Ser Ile Thr Gln Lys Tyr Ser Gln Ser Asn Glu Ala
 65 70 75 80
 Glu Ser Ala Tyr Leu Pro Gly Gly Thr Gly Trp Tyr Arg Lys Ser Phe
 85 90 95
 Thr Ile Asp Arg Asp Leu Ala Gly Lys Arg Ile Ala Ile Asn Phe Asp
 100 105 110
 Gly Val Tyr Met Asn Ala Thr Val Trp Phe Asn Gly Val Lys Leu Gly
 115 120 125
 Thr His Pro Tyr Gly Tyr Ser Pro Phe Ser Phe Asp Leu Thr Gly Asn
 130 135 140
 Ala Lys Phe Gly Gly Glu Asn Thr Ile Val Val Lys Val Glu Asn Arg
 145 150 155 160
 Leu Pro Ser Ser Arg Trp Tyr Ser Gly Ser Gly Ile Tyr Arg Asp Val
 165 170 175
 Thr Leu Thr Val Thr Asp Gly Val His Val Gly Asn Asn Gly Val Ala
 180 185 190
 Ile Lys Thr Pro Ser Leu Ala Thr Gln Asn Gly Gly Asp Val Thr Met
 195 200 205
 Asn Leu Thr Thr Lys Val Ala Asn Asp Thr Glu Ala Ala Asn Ile
 210 215 220
 Thr Leu Lys Gln Thr Val Phe Pro Lys Gly Gly Lys Thr Asp Ala Ala
 225 230 235 240
 Ile Gly Thr Val Thr Thr Ala Ser Lys Ser Ile Ala Ala Gly Ala Ser

EP 3 473 713 A1

					245				250				255			
	Ala	Asp	Val	Thr	Ser	Thr	Ile	Thr	Ala	Ala	Ser	Pro	Lys	Leu	Trp	Ser
				260					265					270		
5	Ile	Lys	Asn	Pro	Asn	Leu	Tyr	Thr	Val	Arg	Thr	Glu	Val	Leu	Asn	Gly
			275					280					285			
	Gly	Lys	Val	Leu	Asp	Thr	Tyr	Asp	Thr	Glu	Tyr	Gly	Phe	Arg	Trp	Thr
		290					295					300				
	Gly	Phe	Asp	Ala	Thr	Ser	Gly	Phe	Ser	Leu	Asn	Gly	Glu	Lys	Val	Lys
	305					310					315					320
10	Leu	Lys	Gly	Val	Ser	Met	His	His	Asp	Gln	Gly	Ser	Leu	Gly	Ala	Val
				325						330					335	
	Ala	Asn	Arg	Arg	Ala	Ile	Glu	Arg	Gln	Val	Glu	Ile	Leu	Gln	Lys	Met
			340						345					350		
	Gly	Val	Asn	Ser	Ile	Arg	Thr	Thr	His	Asn	Pro	Ala	Ala	Lys	Ala	Leu
		355						360					365			
15	Ile	Asp	Val	Cys	Asn	Glu	Lys	Gly	Val	Leu	Val	Val	Glu	Glu	Val	Phe
	370						375					380				
	Asp	Met	Trp	Asn	Arg	Ser	Lys	Asn	Gly	Asn	Thr	Glu	Asp	Tyr	Gly	Lys
	385					390					395					400
	Trp	Phe	Gly	Gln	Ala	Ile	Ala	Gly	Asp	Asn	Ala	Val	Leu	Gly	Gly	Asp
				405					410						415	
20	Lys	Asp	Glu	Thr	Trp	Ala	Lys	Phe	Asp	Leu	Thr	Ser	Thr	Ile	Asn	Arg
			420						425						430	
	Asp	Arg	Asn	Ala	Pro	Ser	Val	Ile	Met	Trp	Ser	Leu	Gly	Asn	Glu	Met
		435						440					445			
	Met	Glu	Gly	Ile	Ser	Gly	Ser	Val	Ser	Gly	Phe	Pro	Ala	Thr	Ser	Ala
	450					455						460				
25	Lys	Leu	Val	Ala	Trp	Thr	Lys	Ala	Ala	Asp	Ser	Thr	Arg	Pro	Met	Thr
	465					470				475						480
	Tyr	Gly	Asp	Asn	Lys	Ile	Lys	Ala	Asn	Trp	Asn	Glu	Ser	Asn	Thr	Met
				485					490						495	
30	Gly	Asp	Asn	Leu	Thr	Ala	Asn	Gly	Gly	Val	Val	Gly	Thr	Asn	Tyr	Ser
			500						505					510		
	Asp	Gly	Ala	Asn	Tyr	Asp	Lys	Ile	Arg	Thr	Thr	His	Pro	Ser	Trp	Ala
		515						520					525			
	Ile	Tyr	Gly	Ser	Glu	Thr	Ala	Ser	Ala	Ile	Asn	Ser	Arg	Gly	Ile	Tyr
	530					535						540				
35	Asn	Arg	Thr	Thr	Gly	Gly	Ala	Gln	Ser	Ser	Asp	Lys	Gln	Leu	Thr	Ser
	545					550					555					560
	Tyr	Asp	Asn	Ser	Ala	Val	Gly	Trp	Gly	Ala	Val	Ala	Ser	Ser	Ala	Trp
				565					570						575	
	Tyr	Asp	Val	Val	Gln	Arg	Asp	Phe	Val	Ala	Gly	Thr	Tyr	Val	Trp	Thr
			580					585						590		
40	Gly	Phe	Asp	Tyr	Leu	Gly	Glu	Pro	Thr	Pro	Trp	Asn	Gly	Thr	Gly	Ser
		595						600					605			
	Gly	Ala	Val	Gly	Ser	Trp	Pro	Ser	Pro	Lys	Asn	Ser	Tyr	Phe	Gly	Ile
		610					615					620				
	Val	Asp	Thr	Ala	Gly	Phe	Pro	Lys	Asp	Thr	Tyr	Tyr	Phe	Tyr	Gln	Ser
	625					630					635					640
45	Gln	Trp	Asn	Asp	Asp	Val	His	Thr	Leu	His	Ile	Leu	Pro	Ala	Trp	Asn
				645						650					655	
	Glu	Asn	Val	Val	Ala	Lys	Gly	Ser	Gly	Asn	Asn	Val	Pro	Val	Val	Val
			660						665					670		
	Tyr	Thr	Asp	Ala	Ala	Lys	Val	Lys	Leu	Tyr	Phe	Thr	Pro	Lys	Gly	Ser
		675						680					685			
50	Thr	Glu	Lys	Arg	Leu	Ile	Gly	Glu	Lys	Ser	Phe	Thr	Lys	Lys	Thr	Thr
		690					695					700				
	Ala	Ala	Gly	Tyr	Thr	Tyr	Gln	Val	Tyr	Glu	Gly	Ser	Asp	Lys	Asp	Ser
	705					710					715					720
	Thr	Ala	His	Lys	Asn	Met	Tyr	Leu	Thr	Trp	Asn	Val	Pro	Trp	Ala	Glu
				725						730				735		
55	Gly	Thr	Ile	Ser	Ala	Glu	Ala	Tyr	Asp	Glu	Asn	Asn	Arg	Leu	Ile	Pro
				740					745					750		

EP 3 473 713 A1

Glu Gly Ser Thr Glu Gly Asn Ala Ser Val Thr Thr Thr Gly Lys Ala
 755 760 765
 Ala Lys Leu Lys Ala Asp Ala Asp Arg Lys Thr Ile Thr Ala Asp Gly
 770 775 780
 5 Lys Asp Leu Ser Tyr Ile Glu Val Asp Val Thr Asp Ala Asn Gly His
 785 790 795 800
 Ile Val Pro Asp Ala Ala Asn Arg Val Thr Phe Asp Val Lys Gly Ala
 805 810 815
 Gly Lys Leu Val Gly Val Asp Asn Gly Ser Ser Pro Asp His Asp Ser
 820 825 830
 10 Tyr Gln Ala Asp Asn Arg Lys Ala Phe Ser Gly Lys Val Leu Ala Ile
 835 840 845
 Val Gln Ser Thr Lys Glu Ala Gly Glu Ile Thr Val Thr Ala Lys Ala
 850 855 860
 Asp Gly Leu Gln Ser Ser Thr Val Lys Ile Ala Thr Thr Ala Val Pro
 865 870 875 880
 15 Gly Thr Ser Thr Glu Lys Thr Val Arg Ser Phe Tyr Tyr Ser Arg Asn
 885 890 895
 Tyr Tyr Val Lys Thr Gly Asn Lys Pro Ile Leu Pro Ser Asp Val Glu
 900 905 910
 Val Arg Tyr Ser Asp Gly Thr Ser Asp Arg Gln Asn Val Thr Trp Asp
 915 920 925
 20 Ala Val Ser Asp Asp Gln Ile Ala Lys Ala Gly Ser Phe Ser Val Ala
 930 935 940
 Gly Thr Val Ala Gly Gln Lys Ile Ser Val Arg Val Thr Met Ile Asp
 945 950 955 960
 25 Glu Ile Gly Ala Leu Leu Asn Tyr Ser Ala Ser Thr Pro Val Gly Thr
 965 970 975
 Pro Ala Val Leu Pro Gly Ser Arg Pro Ala Val Leu Pro Asp Gly Thr
 980 985 990
 Val Thr Ser Ala Asn Phe Ala Val His Trp Thr Lys Pro Ala Asp Thr
 995 1000 1005
 30 Val Tyr Asn Thr Ala Gly Thr Val Lys Val Pro Gly Thr Ala Thr Val
 1010 1015 1020
 Phe Gly Lys Glu Phe Lys Val Thr Ala Thr Ile Arg Val Gln
 1025 1030 1035

 <210> 4
 35 <211> 1142
 <212> PRT
 <213> Bifidobacterium bifidum

 <220>
 40 <221> SOURCE
 <222> 1..1142
 <223> /mol_type="protein"
 /organism="Bifidobacterium bifidum"

 <400> 4
 45 Val Glu Asp Ala Thr Arg Ser Asp Ser Thr Thr Gln Met Ser Ser Thr
 1 5 10 15
 Pro Glu Val Val Tyr Ser Ser Ala Val Asp Ser Lys Gln Asn Arg Thr
 20 25 30
 Ser Asp Phe Asp Ala Asn Trp Lys Phe Met Leu Ser Asp Ser Val Gln
 35 40 45
 50 Ala Gln Asp Pro Ala Phe Asp Asp Ser Ala Trp Gln Gln Val Asp Leu
 50 55 60
 Pro His Asp Tyr Ser Ile Thr Gln Lys Tyr Ser Gln Ser Asn Glu Ala
 65 70 75 80
 Glu Ser Ala Tyr Leu Pro Gly Gly Thr Gly Trp Tyr Arg Lys Ser Phe
 85 90 95
 55 Thr Ile Asp Arg Asp Leu Ala Gly Lys Arg Ile Ala Ile Asn Phe Asp
 100 105 110

EP 3 473 713 A1

	Gly	Val	Tyr	Met	Asn	Ala	Thr	Val	Trp	Phe	Asn	Gly	Val	Lys	Leu	Gly	
		115						120					125				
	Thr	His	Pro	Tyr	Gly	Tyr	Ser	Pro	Phe	Ser	Phe	Asp	Leu	Thr	Gly	Asn	
		130					135					140					
5	Ala	Lys	Phe	Gly	Gly	Glu	Asn	Thr	Ile	Val	Val	Lys	Val	Glu	Asn	Arg	
	145					150					155					160	
	Leu	Pro	Ser	Ser	Arg	Trp	Tyr	Ser	Gly	Ser	Gly	Ile	Tyr	Arg	Asp	Val	
					165					170					175		
	Thr	Leu	Thr	Val	Thr	Asp	Gly	Val	His	Val	Gly	Asn	Asn	Gly	Val	Ala	
				180					185					190			
10	Ile	Lys	Thr	Pro	Ser	Leu	Ala	Thr	Gln	Asn	Gly	Gly	Asp	Val	Thr	Met	
		195						200					205				
	Asn	Leu	Thr	Thr	Lys	Val	Ala	Asn	Asp	Thr	Glu	Ala	Ala	Ala	Asn	Ile	
		210					215					220					
	Thr	Leu	Lys	Gln	Thr	Val	Phe	Pro	Lys	Gly	Gly	Lys	Thr	Asp	Ala	Ala	
15	225					230					235					240	
	Ile	Gly	Thr	Val	Thr	Thr	Ala	Ser	Lys	Ser	Ile	Ala	Ala	Gly	Ala	Ser	
					245					250					255		
	Ala	Asp	Val	Thr	Ser	Thr	Ile	Thr	Ala	Ala	Ser	Pro	Lys	Leu	Trp	Ser	
				260					265					270			
	Ile	Lys	Asn	Pro	Asn	Leu	Tyr	Thr	Val	Arg	Thr	Glu	Val	Leu	Asn	Gly	
20			275					280					285				
	Gly	Lys	Val	Leu	Asp	Thr	Tyr	Asp	Thr	Glu	Tyr	Gly	Phe	Arg	Trp	Thr	
		290					295					300					
	Gly	Phe	Asp	Ala	Thr	Ser	Gly	Phe	Ser	Leu	Asn	Gly	Glu	Lys	Val	Lys	
	305					310					315					320	
25	Leu	Lys	Gly	Val	Ser	Met	His	His	Asp	Gln	Gly	Ser	Leu	Gly	Ala	Val	
					325					330					335		
	Ala	Asn	Arg	Arg	Ala	Ile	Glu	Arg	Gln	Val	Glu	Ile	Leu	Gln	Lys	Met	
				340					345					350			
	Gly	Val	Asn	Ser	Ile	Arg	Thr	Thr	His	Asn	Pro	Ala	Ala	Lys	Ala	Leu	
				355				360					365				
30	Ile	Asp	Val	Cys	Asn	Glu	Lys	Gly	Val	Leu	Val	Val	Glu	Glu	Val	Phe	
		370				375						380					
	Asp	Met	Trp	Asn	Arg	Ser	Lys	Asn	Gly	Asn	Thr	Glu	Asp	Tyr	Gly	Lys	
	385					390				395						400	
	Trp	Phe	Gly	Gln	Ala	Ile	Ala	Gly	Asp	Asn	Ala	Val	Leu	Gly	Gly	Asp	
				405					410						415		
35	Lys	Asp	Glu	Thr	Trp	Ala	Lys	Phe	Asp	Leu	Thr	Ser	Thr	Ile	Asn	Arg	
				420					425					430			
	Asp	Arg	Asn	Ala	Pro	Ser	Val	Ile	Met	Trp	Ser	Leu	Gly	Asn	Glu	Met	
			435					440					445				
	Met	Glu	Gly	Ile	Ser	Gly	Ser	Val	Ser	Gly	Phe	Pro	Ala	Thr	Ser	Ala	
		450					455					460					
40	Lys	Leu	Val	Ala	Trp	Thr	Lys	Ala	Ala	Asp	Ser	Thr	Arg	Pro	Met	Thr	
	465					470					475					480	
	Tyr	Gly	Asp	Asn	Lys	Ile	Lys	Ala	Asn	Trp	Asn	Glu	Ser	Asn	Thr	Met	
				485						490					495		
	Gly	Asp	Asn	Leu	Thr	Ala	Asn	Gly	Gly	Val	Val	Gly	Thr	Asn	Tyr	Ser	
45				500					505					510			
	Asp	Gly	Ala	Asn	Tyr	Asp	Lys	Ile	Arg	Thr	Thr	His	Pro	Ser	Trp	Ala	
			515					520					525				
	Ile	Tyr	Gly	Ser	Glu	Thr	Ala	Ser	Ala	Ile	Asn	Ser	Arg	Gly	Ile	Tyr	
		530					535					540					
	Asn	Arg	Thr	Thr	Gly	Gly	Ala	Gln	Ser	Ser	Asp	Lys	Gln	Leu	Thr	Ser	
50	545					550					555					560	
	Tyr	Asp	Asn	Ser	Ala	Val	Gly	Trp	Gly	Ala	Val	Ala	Ser	Ser	Ala	Trp	
				565						570					575		
	Tyr	Asp	Val	Val	Gln	Arg	Asp	Phe	Val	Ala	Gly	Thr	Tyr	Val	Trp	Thr	
				580					585					590			
55	Gly	Phe	Asp	Tyr	Leu	Gly	Glu	Pro	Thr	Pro	Trp	Asn	Gly	Thr	Gly	Ser	
			595					600					605				
	Gly	Ala	Val	Gly	Ser	Trp	Pro	Ser	Pro	Lys	Asn	Ser	Tyr	Phe	Gly	Ile	

EP 3 473 713 A1

[illegible]

EP 3 473 713 A1

Met Tyr Phe Phe Arg Asp Ser Asn Ala Val Arg Phe Pro Asp Ala Gly
 1125 1130 1135
 Lys Thr Lys Ile Gln Ile
 1140

<210> 5
 <211> 1211
 <212> PRT
 <213> Bifidobacterium bifidum

<220>
 <221> SOURCE
 <222> 1..1211
 <223> /mol_type="protein"
 /organism="Bifidobacterium bifidum"

<400> 5
 Val Glu Asp Ala Thr Arg Ser Asp Ser Thr Thr Gln Met Ser Ser Thr
 1 5 10 15
 Pro Glu Val Val Tyr Ser Ser Ala Val Asp Ser Lys Gln Asn Arg Thr
 20 25 30
 Ser Asp Phe Asp Ala Asn Trp Lys Phe Met Leu Ser Asp Ser Val Gln
 35 40 45
 Ala Gln Asp Pro Ala Phe Asp Asp Ser Ala Trp Gln Gln Val Asp Leu
 50 55 60
 Pro His Asp Tyr Ser Ile Thr Gln Lys Tyr Ser Gln Ser Asn Glu Ala
 65 70 75 80
 Glu Ser Ala Tyr Leu Pro Gly Gly Thr Gly Trp Tyr Arg Lys Ser Phe
 85 90 95
 Thr Ile Asp Arg Asp Leu Ala Gly Lys Arg Ile Ala Ile Asn Phe Asp
 100 105 110
 Gly Val Tyr Met Asn Ala Thr Val Trp Phe Asn Gly Val Lys Leu Gly
 115 120 125
 Thr His Pro Tyr Gly Tyr Ser Pro Phe Ser Phe Asp Leu Thr Gly Asn
 130 135 140
 Ala Lys Phe Gly Gly Glu Asn Thr Ile Val Val Lys Val Glu Asn Arg
 145 150 155 160
 Leu Pro Ser Ser Arg Trp Tyr Ser Gly Ser Gly Ile Tyr Arg Asp Val
 165 170 175
 Thr Leu Thr Val Thr Asp Gly Val His Val Gly Asn Asn Gly Val Ala
 180 185 190
 Ile Lys Thr Pro Ser Leu Ala Thr Gln Asn Gly Gly Asp Val Thr Met
 195 200 205
 Asn Leu Thr Thr Lys Val Ala Asn Asp Thr Glu Ala Ala Asn Ile
 210 215 220
 Thr Leu Lys Gln Thr Val Phe Pro Lys Gly Gly Lys Thr Asp Ala Ala
 225 230 235 240
 Ile Gly Thr Val Thr Thr Ala Ser Lys Ser Ile Ala Ala Gly Ala Ser
 245 250 255
 Ala Asp Val Thr Ser Thr Ile Thr Ala Ala Ser Pro Lys Leu Trp Ser
 260 265 270
 Ile Lys Asn Pro Asn Leu Tyr Thr Val Arg Thr Glu Val Leu Asn Gly
 275 280 285
 Gly Lys Val Leu Asp Thr Tyr Asp Thr Glu Tyr Gly Phe Arg Trp Thr
 290 295 300
 Gly Phe Asp Ala Thr Ser Gly Phe Ser Leu Asn Gly Glu Lys Val Lys
 305 310 315 320
 Leu Lys Gly Val Ser Met His His Asp Gln Gly Ser Leu Gly Ala Val
 325 330 335
 Ala Asn Arg Arg Ala Ile Glu Arg Gln Val Glu Ile Leu Gln Lys Met
 340 345 350
 Gly Val Asn Ser Ile Arg Thr Thr His Asn Pro Ala Ala Lys Ala Leu
 355 360 365

EP 3 473 713 A1

	Ile	Asp	Val	Cys	Asn	Glu	Lys	Gly	Val	Leu	Val	Val	Glu	Glu	Val	Phe	
	370						375					380					
	Asp	Met	Trp	Asn	Arg	Ser	Lys	Asn	Gly	Asn	Thr	Glu	Asp	Tyr	Gly	Lys	
	385					390				395						400	
5	Trp	Phe	Gly	Gln	Ala	Ile	Ala	Gly	Asp	Asn	Ala	Val	Leu	Gly	Gly	Asp	
				405						410					415		
	Lys	Asp	Glu	Thr	Trp	Ala	Lys	Phe	Asp	Leu	Thr	Ser	Thr	Ile	Asn	Arg	
				420					425					430			
	Asp	Arg	Asn	Ala	Pro	Ser	Val	Ile	Met	Trp	Ser	Leu	Gly	Asn	Glu	Met	
			435					440					445				
10	Met	Glu	Gly	Ile	Ser	Gly	Ser	Val	Ser	Gly	Phe	Pro	Ala	Thr	Ser	Ala	
		450				455						460					
	Lys	Leu	Val	Ala	Trp	Thr	Lys	Ala	Ala	Asp	Ser	Thr	Arg	Pro	Met	Thr	
	465					470				475						480	
	Tyr	Gly	Asp	Asn	Lys	Ile	Lys	Ala	Asn	Trp	Asn	Glu	Ser	Asn	Thr	Met	
				485						490					495		
15	Gly	Asp	Asn	Leu	Thr	Ala	Asn	Gly	Gly	Val	Val	Gly	Thr	Asn	Tyr	Ser	
				500					505					510			
	Asp	Gly	Ala	Asn	Tyr	Asp	Lys	Ile	Arg	Thr	Thr	His	Pro	Ser	Trp	Ala	
			515					520					525				
20	Ile	Tyr	Gly	Ser	Glu	Thr	Ala	Ser	Ala	Ile	Asn	Ser	Arg	Gly	Ile	Tyr	
		530					535					540					
	Asn	Arg	Thr	Thr	Gly	Gly	Ala	Gln	Ser	Ser	Asp	Lys	Gln	Leu	Thr	Ser	
	545					550					555					560	
	Tyr	Asp	Asn	Ser	Ala	Val	Gly	Trp	Gly	Ala	Val	Ala	Ser	Ser	Ala	Trp	
				565						570					575		
25	Tyr	Asp	Val	Val	Gln	Arg	Asp	Phe	Val	Ala	Gly	Thr	Tyr	Val	Trp	Thr	
				580					585					590			
	Gly	Phe	Asp	Tyr	Leu	Gly	Glu	Pro	Thr	Pro	Trp	Asn	Gly	Thr	Gly	Ser	
			595					600					605				
	Gly	Ala	Val	Gly	Ser	Trp	Pro	Ser	Pro	Lys	Asn	Ser	Tyr	Phe	Gly	Ile	
		610					615					620					
30	Val	Asp	Thr	Ala	Gly	Phe	Pro	Lys	Asp	Thr	Tyr	Tyr	Phe	Tyr	Gln	Ser	
	625					630					635					640	
	Gln	Trp	Asn	Asp	Asp	Val	His	Thr	Leu	His	Ile	Leu	Pro	Ala	Trp	Asn	
				645						650					655		
	Glu	Asn	Val	Val	Ala	Lys	Gly	Ser	Gly	Asn	Asn	Val	Pro	Val	Val	Val	
				660						665				670			
35	Tyr	Thr	Asp	Ala	Ala	Lys	Val	Lys	Leu	Tyr	Phe	Thr	Pro	Lys	Gly	Ser	
			675						680				685				
	Thr	Glu	Lys	Arg	Leu	Ile	Gly	Glu	Lys	Ser	Phe	Thr	Lys	Lys	Thr	Thr	
		690					695					700					
	Ala	Ala	Gly	Tyr	Thr	Tyr	Gln	Val	Tyr	Glu	Gly	Ser	Asp	Lys	Asp	Ser	
	705					710					715					720	
40	Thr	Ala	His	Lys	Asn	Met	Tyr	Leu	Thr	Trp	Asn	Val	Pro	Trp	Ala	Glu	
				725						730					735		
	Gly	Thr	Ile	Ser	Ala	Glu	Ala	Tyr	Asp	Glu	Asn	Asn	Arg	Leu	Ile	Pro	
				740					745					750			
	Glu	Gly	Ser	Thr	Glu	Gly	Asn	Ala	Ser	Val	Thr	Thr	Thr	Gly	Lys	Ala	
			755					760					765				
45	Ala	Lys	Leu	Lys	Ala	Asp	Ala	Asp	Arg	Lys	Thr	Ile	Thr	Ala	Asp	Gly	
		770					775					780					
	Lys	Asp	Leu	Ser	Tyr	Ile	Glu	Val	Asp	Val	Thr	Asp	Ala	Asn	Gly	His	
	785					790					795					800	
50	Ile	Val	Pro	Asp	Ala	Ala	Asn	Arg	Val	Thr	Phe	Asp	Val	Lys	Gly	Ala	
				805						810					815		
	Gly	Lys	Leu	Val	Gly	Val	Asp	Asn	Gly	Ser	Ser	Pro	Asp	His	Asp	Ser	
				820					825					830			
	Tyr	Gln	Ala	Asp	Asn	Arg	Lys	Ala	Phe	Ser	Gly	Lys	Val	Leu	Ala	Ile	
			835					840					845				
55	Val	Gln	Ser	Thr	Lys	Glu	Ala	Gly	Glu	Ile	Thr	Val	Thr	Ala	Lys	Ala	
		850					855					860					
	Asp	Gly	Leu	Gln	Ser	Ser	Thr	Val	Lys	Ile	Ala	Thr	Thr	Ala	Val	Pro	

EP 3 473 713 A1

[illegible]

EP 3 473 713 A1

[illegible]

EP 3 473 713 A1

	Tyr	Asp	Asn	Ser	Ala	Val	Gly	Trp	Gly	Ala	Val	Ala	Ser	Ser	Ala	Trp	
				565						570					575		
	Tyr	Asp	Val	Val	Gln	Arg	Asp	Phe	Val	Ala	Gly	Thr	Tyr	Val	Trp	Thr	
			580						585					590			
5	Gly	Phe	Asp	Tyr	Leu	Gly	Glu	Pro	Thr	Pro	Trp	Asn	Gly	Thr	Gly	Ser	
			595					600					605				
	Gly	Ala	Val	Gly	Ser	Trp	Pro	Ser	Pro	Lys	Asn	Ser	Tyr	Phe	Gly	Ile	
		610					615					620					
	Val	Asp	Thr	Ala	Gly	Phe	Pro	Lys	Asp	Thr	Tyr	Tyr	Phe	Tyr	Gln	Ser	
	625					630					635					640	
10	Gln	Trp	Asn	Asp	Asp	Val	His	Thr	Leu	His	Ile	Leu	Pro	Ala	Trp	Asn	
				645						650					655		
	Glu	Asn	Val	Val	Ala	Lys	Gly	Ser	Gly	Asn	Asn	Val	Pro	Val	Val	Val	
				660						665				670			
	Tyr	Thr	Asp	Ala	Ala	Lys	Val	Lys	Leu	Tyr	Phe	Thr	Pro	Lys	Gly	Ser	
			675						680				685				
15	Thr	Glu	Lys	Arg	Leu	Ile	Gly	Glu	Lys	Ser	Phe	Thr	Lys	Lys	Thr	Thr	
		690					695					700					
	Ala	Ala	Gly	Tyr	Thr	Tyr	Gln	Val	Tyr	Glu	Gly	Ser	Asp	Lys	Asp	Ser	
	705					710					715					720	
	Thr	Ala	His	Lys	Asn	Met	Tyr	Leu	Thr	Trp	Asn	Val	Pro	Trp	Ala	Glu	
20					725						730				735		
	Gly	Thr	Ile	Ser	Ala	Glu	Ala	Tyr	Asp	Glu	Asn	Asn	Arg	Leu	Ile	Pro	
				740						745				750			
	Glu	Gly	Ser	Thr	Glu	Gly	Asn	Ala	Ser	Val	Thr	Thr	Thr	Gly	Lys	Ala	
			755					760					765				
	Ala	Lys	Leu	Lys	Ala	Asp	Ala	Asp	Arg	Lys	Thr	Ile	Thr	Ala	Asp	Gly	
25		770						775				780					
	Lys	Asp	Leu	Ser	Tyr	Ile	Glu	Val	Asp	Val	Thr	Asp	Ala	Asn	Gly	His	
	785					790					795					800	
	Ile	Val	Pro	Asp	Ala	Ala	Asn	Arg	Val	Thr	Phe	Asp	Val	Lys	Gly	Ala	
					805						810				815		
30	Gly	Lys	Leu	Val	Gly	Val	Asp	Asn	Gly	Ser	Ser	Pro	Asp	His	Asp	Ser	
				820					825					830			
	Tyr	Gln	Ala	Asp	Asn	Arg	Lys	Ala	Phe	Ser	Gly	Lys	Val	Leu	Ala	Ile	
			835					840					845				
	Val	Gln	Ser	Thr	Lys	Glu	Ala	Gly	Glu	Ile	Thr	Val	Thr	Ala	Lys	Ala	
							855					860					
35	Asp	Gly	Leu	Gln	Ser	Ser	Thr	Val	Lys	Ile	Ala	Thr	Thr	Ala	Val	Pro	
	865						870				875					880	
	Gly	Thr	Ser	Thr	Glu	Lys	Thr	Val	Arg	Ser	Phe	Tyr	Tyr	Ser	Arg	Asn	
					885						890				895		
	Tyr	Tyr	Val	Lys	Thr	Gly	Asn	Lys	Pro	Ile	Leu	Pro	Ser	Asp	Val	Glu	
				900						905				910			
40	Val	Arg	Tyr	Ser	Asp	Gly	Thr	Ser	Asp	Arg	Gln	Asn	Val	Thr	Trp	Asp	
			915						920					925			
	Ala	Val	Ser	Asp	Asp	Gln	Ile	Ala	Lys	Ala	Gly	Ser	Phe	Ser	Val	Ala	
							935					940					
	Gly	Thr	Val	Ala	Gly	Gln	Lys	Ile	Ser	Val	Arg	Val	Thr	Met	Ile	Asp	
45		945				950					955					960	
	Glu	Ile	Gly	Ala	Leu	Leu	Asn	Tyr	Ser	Ala	Ser	Thr	Pro	Val	Gly	Thr	
					965						970				975		
	Pro	Ala	Val	Leu	Pro	Gly	Ser	Arg	Pro	Ala	Val	Leu	Pro	Asp	Gly	Thr	
				980						985				990			
	Val	Thr	Ser	Ala	Asn	Phe	Ala	Val	His	Trp	Thr	Lys	Pro	Ala	Asp	Thr	
50									1000					1005			
	Val	Tyr	Asn	Thr	Ala	Gly	Thr	Val	Lys	Val	Pro	Gly	Thr	Ala	Thr	Val	
		1010					1015					1020					
	Phe	Gly	Lys	Glu	Phe	Lys	Val	Thr	Ala	Thr	Ile	Arg	Val	Gln	Arg	Ser	
		1025					1030					1035				1040	
	Gln	Val	Thr	Ile	Gly	Ser	Ser	Val	Ser	Gly	Asn	Ala	Leu	Arg	Leu	Thr	
55					1045						1050				1055		
	Gln	Asn	Ile	Pro	Ala	Asp	Lys	Gln	Ser	Asp	Thr	Leu	Asp	Ala	Ile	Lys	

EP 3 473 713 A1

1060 1065 1070
 Asp Gly Ser Thr Thr Val Asp Ala Asn Thr Gly Gly Gly Ala Asn Pro
 1075 1080 1085
 Ser Ala Trp Thr Asn Trp Ala Tyr Ser Lys Ala Gly His Asn Thr Ala
 1090 1095 1100
 5 Glu Ile Thr Phe Glu Tyr Ala Thr Glu Gln Gln Leu Gly Gln Ile Val
 1105 1110 1115 1120
 Met Tyr Phe Phe Arg Asp Ser Asn Ala Val Arg Phe Pro Asp Ala Gly
 1125 1130 1135
 10 Lys Thr Lys Ile Gln Ile Ser Ala Asp Gly Lys Asn Trp Thr Asp Leu
 1140 1145 1150
 Ala Ala Thr Glu Thr Ile Ala Ala Gln Glu Ser Ser Asp Arg Val Lys
 1155 1160 1165
 Pro Tyr Thr Tyr Asp Phe Ala Pro Val Gly Ala Thr Phe Val Lys Val
 1170 1175 1180
 15 Thr Val Thr Asn Ala Asp Thr Thr Thr Pro Ser Gly Val Val Cys Ala
 1185 1190 1195 1200
 Gly Leu Thr Glu Ile Glu Leu Lys Thr Ala Thr Ser Lys Phe Val Thr
 1205 1210 1215
 Asn Thr Ser Ala Ala Leu Ser Ser Leu Thr Val Asn Gly Thr Lys Val
 1220 1225 1230
 20 Ser Asp Ser Val Leu Ala Ala Gly Ser Tyr Asn Thr Pro Ala Ile Ile
 1235 1240 1245
 Ala Asp Val Lys Ala Glu Gly Glu Gly Asn Ala Ser Val Thr Val Leu
 1250 1255 1260
 Pro Ala His Asp Asn Val Ile Arg Val Ile Thr Glu Ser Glu Asp His
 1265 1270 1275 1280
 25 Val Thr Arg Lys Thr Phe Thr Ile Asn Leu Gly Thr Glu Gln Glu Phe
 1285 1290 1295

 <210> 7
 <211> 453
 30 <212> PRT
 <213> Bifidobacterium bifidum

 <220>
 <221> SOURCE
 <222> 1..453
 35 <223> /mol_type="protein"
 /note="Glycoside hydrolase catalytic core"
 /organism="Bifidobacterium bifidum"

 <400> 7
 40 Gln Asn Arg Thr Ser Asp Phe Asp Ala Asn Trp Lys Phe Met Leu Ser
 1 5 10 15
 Asp Ser Val Gln Ala Gln Asp Pro Ala Phe Asp Asp Ser Ala Trp Gln
 20 25 30
 Gln Val Asp Leu Pro His Asp Tyr Ser Ile Thr Gln Lys Tyr Ser Gln
 35 40 45
 45 Ser Asn Glu Ala Glu Ser Ala Tyr Leu Pro Gly Gly Thr Gly Trp Tyr
 50 55 60
 Arg Lys Ser Phe Thr Ile Asp Arg Asp Leu Ala Gly Lys Arg Ile Ala
 65 70 75 80
 Ile Asn Phe Asp Gly Val Tyr Met Asn Ala Thr Val Trp Phe Asn Gly
 85 90 95
 50 Val Lys Leu Gly Thr His Pro Tyr Gly Tyr Ser Pro Phe Ser Phe Asp
 100 105 110
 Leu Thr Gly Asn Ala Lys Phe Gly Gly Glu Asn Thr Ile Val Val Lys
 115 120 125
 Val Glu Asn Arg Leu Pro Ser Ser Arg Trp Tyr Ser Gly Ser Gly Ile
 130 135 140
 55 Tyr Arg Asp Val Thr Leu Thr Val Thr Asp Gly Val His Val Gly Asn
 145 150 155 160

EP 3 473 713 A1

Asn Gly Val Ala Ile Lys Thr Pro Ser Leu Ala Thr Gln Asn Gly Gly
 165 170 175
 Asp Val Thr Met Asn Leu Thr Thr Lys Val Ala Asn Asp Thr Glu Ala
 180 185 190
 5 Ala Ala Asn Ile Thr Leu Lys Gln Thr Val Phe Pro Lys Gly Gly Lys
 195 200 205
 Thr Asp Ala Ala Ile Gly Thr Val Thr Thr Ala Ser Lys Ser Ile Ala
 210 215 220
 Ala Gly Ala Ser Ala Asp Val Thr Ser Thr Ile Thr Ala Ala Ser Pro
 225 230 235 240
 10 Lys Leu Trp Ser Ile Lys Asn Pro Asn Leu Tyr Thr Val Arg Thr Glu
 245 250 255
 Val Leu Asn Gly Lys Val Leu Asp Thr Tyr Asp Thr Glu Tyr Gly
 260 265 270
 Phe Arg Trp Thr Gly Phe Asp Ala Thr Ser Gly Phe Ser Leu Asn Gly
 275 280 285
 15 Glu Lys Val Lys Leu Lys Gly Val Ser Met His His Asp Gln Gly Ser
 290 295 300
 Leu Gly Ala Val Ala Asn Arg Arg Ala Ile Glu Arg Gln Val Glu Ile
 305 310 315 320
 20 Leu Gln Lys Met Gly Val Asn Ser Ile Arg Thr Thr His Asn Pro Ala
 325 330 335
 Ala Lys Ala Leu Ile Asp Val Cys Asn Glu Lys Gly Val Leu Val Val
 340 345 350
 Glu Glu Val Phe Asp Met Trp Asn Arg Ser Lys Asn Gly Asn Thr Glu
 355 360 365
 25 Asp Tyr Gly Lys Trp Phe Gly Gln Ala Ile Ala Gly Asp Asn Ala Val
 370 375 380
 Leu Gly Gly Asp Lys Asp Glu Thr Trp Ala Lys Phe Asp Leu Thr Ser
 385 390 395 400
 Thr Ile Asn Arg Asp Arg Asn Ala Pro Ser Val Ile Met Trp Ser Leu
 405 410 415
 30 Gly Asn Glu Met Met Glu Gly Ile Ser Gly Ser Val Ser Gly Phe Pro
 420 425 430
 Ala Thr Ser Ala Lys Leu Val Ala Trp Thr Lys Ala Ala Asp Ser Thr
 435 440 445
 Arg Pro Met Thr Tyr
 450
 35
 <210> 8
 <211> 5160
 <212> DNA
 <213> Bifidobacterium bifidum
 40
 <220>
 <221> source
 <222> 1..5160
 <223> /mol_type="DNA"
 45 /organism="Bifidobacterium bifidum"
 <400> 8
 gcagttgaag atgcaacaag aagcgatagc acaacacaaa tgtcatcaac accggaagtt 60
 gtttattcat cagcggtcga tagcaaacaa aatcgacaaa gcgattttga tgcgaactgg 120
 50 aaatttatgc tgtcagatag cgttcaagca caagatccgg catttgatga ttcagcatgg 180
 caacaagttg atctgccgca tgattatagc atcacacaga aatatagcca aagcaatgaa 240
 gcagaatcag catatcttcc gggaggcaca ggctgggtata gaaaaagctt tacaattgat 300
 55 agagatctgg caggcaaacg cattgcgatt aattttgatg gcgtctatat gaatgcaaca 360

EP 3 473 713 A1

	gtctggttta atggcgtaa actgggcaca catccgtatg gctattcacc gttttcattt	420
	gatctgacag gcaatgcaaa atttggcga gaaaacacaa ttgtcgtcaa agttgaaaat	480
5	agactgccgt catcaagatg gtattcaggc agcggcattt atagagatgt taaactgaca	540
	gttacagatg gcgttcattg tggcaataat ggcgtcgcaa ttaaaacacc gtactggca	600
	acacaaaatg gcggagatgt cacaatgaac ctgacaacaa aagtcgcgaa tgatacagaa	660
10	gcagcagcga acattacact gaaacagaca gtttttccga aaggcggaaa aacggatgca	720
	gcaattggca cagttacaac agcatcaaaa tcaattgcag caggcgcattc agcagatgtt	780
	acaagcaca ttacagcagc aagcccgaaa ctgtggtcaa ttaaaaaccc gaacctgtat	840
15	acagttagaa cagaagttct gaacggaggc aaagttctgg atacatatga tacagaatat	900
	ggctttcgct ggacaggctt tgatgcaaca tcaggctttt cactgaatgg cgaaaaagtc	960
	aaactgaaag gcgttagcat gcatcatgat caaggctcac ttggcgcagt tgcaaataga	1020
20	cgcgcaattg aaagacaagt cgaaatcctg caaaaaatgg gcgtcaatag cattcgca	1080
	acacataatc cggcagcaaa agcactgatt gatgtctgca atgaaaaagg cgttctgggtt	1140
25	gtcgaagaag tctttgatat gtggaaccgc agcaaaaatg gcaacacgga agattatggc	1200
	aaatggtttg gccaaagcaat tgcaggcgat aatgcagttc tgggaggcga taaagatgaa	1260
	acatgggcga aatttgatct tacatcaaca attaacgcgc atagaaatgc accgtcagtt	1320
30	attatgtggt cactgggcaa tgaaatgatg gaaggcattt caggctcagt ttcaggcttt	1380
	ccggcaacat cagcaaaact gtttgcattg acaaaagcag cagattcaac aagaccgatg	1440
	acatatggcg ataacaaaat taaagcgaac tggaaacgaat caaatacaat gggcgataat	1500
35	ctgacagcaa atggcggagt tgttggcaca aattattcag atggcgcaaa ctatgataaa	1560
	attcgtacaa cacatccgtc atgggcaatt tatggctcag aaacagcattc agcgattaat	1620
	agccgtggca ttataatag aacaacaggc ggagcacaat catcagataa acagctgaca	1680
40	agctatgata attcagcagt tggctgggga gcagttgcat catcagcatg gtatgatgtt	1740
	gttcagagag attttgtcgc aggcacatat gtttggacag gatttgatta tctgggcgaa	1800
	ccgacaccgt ggaatggcac aggctcaggc gcagttggct catggccgtc accgaaaaat	1860
45	agctattttg gcatcggtga tacagcaggc tttccgaaag atacatatta tttttatcag	1920
	agccagtgga atgatgatgt tcatacactg catattcttc cggcatggaa tgaaaatgtt	1980
	gttgcaaaaag gctcaggcaa taatgttccg gttgtcgttt atacagatgc agcgaaagtg	2040
50	aaactgtatt ttacaccgaa aggctcaaca gaaaaaagac tgatcggcga aaaatcattt	2100
	acaaaaaaaa caacagcggc aggctataca tatcaagtct atgaaggcag cgataaagat	2160
55	tcaacagcgc ataaaaacat gtatctgaca tggaatgttc cgtgggcaga aggcacaatt	2220
	tcagcgggaag cgtatgatga aaataatcgc ctgattccgg aaggcagcac agaaggcaac	2280

EP 3 473 713 A1

	gcatcagtta caacaacagg caaagcagca aaactgaaag cagatgcgga tcgcaaaaca	2340
	attacagcgg atggcaaaga tctgtcatat attgaagtgc atgtcacaga tgcaaattggc	2400
5	catattgttc cggatgcagc aaatagagtc acatttgatg ttaaaggcgc aggcaaactg	2460
	gttggcgttg ataatggctc atcaccggat catgattcat atcaagcggga taaccgcaaa	2520
	gcattttcag gcaaagtcct ggcaattgtt cagtcaacaa aagaagcagg cgaaattaca	2580
10	gttacagcaa aagcagatgg cctgcaatca agcacagtta aaattgcaac aacagcagtt	2640
	ccgggaacaa gcacagaaaa aacagtccgc agcttttatt acagccgcaa ctattatgtc	2700
	aaaacaggca acaaaccgat tctgccgtca gatgttgaag ttogctattc agatggaaca	2760
15	agcgatagac aaaacgttac atgggatgca gtttcagatg atcaaattgc aaaagcaggc	2820
	tcattttcag ttgcaggcac agttgcaggc caaaaaatta gcgttcgcgt cacaatgatt	2880
	gatgaaattg gcgcactgct gaattattca gcaagcacac cggttggcac accggcagtt	2940
20	cttccgggat caagaccggc agtcctgccg gatggcacag tcacatcagc aaattttgca	3000
	gtccattgga caaaaccggc agatacagtc tataatacag caggcacagt caaagtaccg	3060
	ggaacagcaa cagtttttgg caaagaattt aaagtcacag cgacaattag agttcaaaga	3120
	agccaagtta caattggctc atcagtttca ggaaatgcac tgagactgac acaaaatatt	3180
	ccggcagata aacaatcaga tacactggat gcgattaaag atggctcaac aacagttgat	3240
30	gcaaatacag gcggaggcgc aaatccgtca gcatggacaa attgggcata ttcaaaagca	3300
	ggccataaca cagcggaaat tacatttgaa tatgcgacag aacaacaact gggccagatc	3360
	gtcatgtatt tttttcgcga tagcaatgca gttagatttc cggatgctgg caaaacaaaa	3420
35	attcagatca gcgcagatgg caaaaattgg acagatctgg cagcaacaga aacaattgca	3480
	gcgcaagaat caagcgatag agtcaaaccg tatacatatg attttgcacc ggttggcgca	3540
	acatttggtta aagtgcagct cacaacgcga gatacaacaa caccgtcagg cgttgtttgc	3600
	gcaggcctga cagaaattga actgaaaaca gcgacaagca aatttgtcac aaatacatca	3660
	gcagcactgt catcacttac agtcaatggc acaaaagttt cagattcagt tctggcagca	3720
45	ggctcatata acacaccggc aattatcgca gatgttaaag cggaaggcga aggcaatgca	3780
	agcgttacag tccttccggc acatgataat gttattcgcg tcattacaga aagcgaagat	3840
	catgtcacac gcaaaacatt tacaatcaac ctgggcacag aacaagaatt tccggctgat	3900
50	tcagatgaaa gagattatcc ggcagcagat atgacagtca cagttggctc agaacaaaca	3960
	tcaggcacag caacagaagg accgaaaaaa tttgcagtcg atggcaacac atcaacatat	4020
	tggcatagca attggacacc gacaacagtt aatgatctgt ggatcgcgtt tgaactgcaa	4080
55	aaaccgacaa aactggatgc actgagatat cttccgcgtc cggcaggctc aaaaaatggc	4140

EP 3 473 713 A1

```

agcgtcacag aatataaagt tcaggtgtca gatgatggaa caaactggac agatgcaggc      4200
tcaggcacat ggacaacgga ttatggctgg aaactggcgg aatttaatca accggtcaca      4260
5 acaaaacatg ttagactgaa agcggttcat acatatgcag atagcggcaa cgataaattt      4320
atgagcgcaa gcgaaattag actgagaaaa gcggtcgata caacggatat ttcaggcgca      4380
acagttacag ttccggcaaa actgacagtt gatagagttg atgcagatca tccggcaaca      4440
10 tttgcaacaa aagatgtcac agttacactg ggagatgcaa cactgagata tggcgttgat      4500
tatctgctgg attatgcagg caatacagca gttggcaaag caacagtgc agttagaggc      4560
attgataaat attcaggcac agtcgcgaaa acatttaca ttgaactgaa aaatgcaccg      4620
15 gcaccggaac cgacactgac atcagtttagc gtcaaaacaa aaccgagcaa actgacatat      4680
gttgctcgag atgcatttga tccggcaggc ctggttctgc aacatgatag acaagcagat      4740
agacctccgc aaccgctggt tggcgaacaa gcggatgaac gcggactgac atgcggcaca      4800
20 agatgcgata gagttgaaca actgcgcaaa catgaaaata gagaagcgca tagaacaggc      4860
ctggatcatc tggaaatttgt tggcgcagca gatggcgcag ttggagaaca agcaacattt      4920
aaagtccatg tccatgcaga tcagggagat ggcagacatg atgatgcaga tgaacgcgat      4980
25 attgatccgc atgttccggt cgatcatgca gttggcgaac tggcaagagc agcatgccat      5040
catgttattg gcctgagagt cgatacacat agacttaaag caagcggctt tcaaattccg      5100
30 gctgatgata tggcagaaat cgatcgcatc acaggctttc atcgttttga acgccatgtc      5160

<210> 9
<211> 2661
<212> DNA
35 <213> Bifidobacterium bifidum

<220>
<221> source
<222> 1..2661
40 <223> /mol_type="DNA"
      /organism="Bifidobacterium bifidum"

<400> 9
gttgaagatg caacaagaag cgatagcaca acacaaatgt catcaacacc ggaagttggt      60
45 tattcatcag cggtcgatag caaacaaaat cgcacaagcg attttgatgc gaactggaaa      120
tttatgctgt cagatagcgt tcaagcacia gatccggcat ttgatgattc agcatggcaa      180
caagttgatc tgccgcatga ttatagcatc acacagaaat atagccaaag caatgaagca      240
50 gaatcagcat atcttccggg aggcacaggc tggatatagaa aaagctttac aattgataga      300
gatctggcag gcaaacgcat tgcgattaat tttgatggcg tctatatgaa tgcaacagtc      360
55 tggtttaatg gcgttaaaact gggcacacat ccgtatggct attcaccggt ttcatattgat      420
ctgacaggca atgcaaaatt tggcggagaa aacacaattg tcgtcaaagt tgaaaataga      480

```

EP 3 473 713 A1

	ctgccgtcat	caagatggta	ttcaggcagc	ggcatttata	gagatgttac	actgacagtt	540
	acagatggcg	ttcatgttgg	caataatggc	gtcgcatta	aaacaccgtc	actggcaaca	600
5	caaaatggcg	gagatgtcac	aatgaacctg	acaacaaaag	tcgcgaatga	tacagaagca	660
	gcagcgaaca	ttacactgaa	acagacagtt	tttccgaaag	gcggaaaaac	ggatgcagca	720
	attggcacag	ttacaacagc	atcaaaatca	attgcagcag	gcgcatcagc	agatgttaca	780
10	agcacaatta	cagcagcaag	cccgaactg	tgggtcaatta	aaaacccgaa	cctgtataca	840
	gttagaacag	aagtcttgaa	cggaggcaaa	gttctggata	catatgatac	agaatatggc	900
	tttcgctgga	caggctttga	tgcaacatca	ggcttttcac	tgaatggcga	aaaagtcaaa	960
15	ctgaaaggcg	ttagcatgca	tcatgatcaa	ggctcacttg	gcgcagttgc	aaatagacgc	1020
	gcaattgaaa	gacaagtcga	aatcctgcaa	aaaatgggcg	tcaatagcat	tcgcacaaca	1080
	cataatccgg	cagcaaaagc	actgattgat	gtctgcaatg	aaaaaggcgt	tctggttgct	1140
20	gaagaagtct	ttgatatgtg	gaaccgcagc	aaaaatggca	acacggaaga	ttatggcaaa	1200
	tggtttgcc	aagcaattgc	aggcgataat	gcagttcttg	gaggcgataa	agatgaaaca	1260
	tgggcgaaat	ttgatcttac	atcaacaatt	aaccgcgata	gaaatgcacc	gtcagttatt	1320
25	atgtggtcac	tgggcaatga	aatgatggaa	ggcatttcag	gctcagtttc	aggctttccg	1380
	gcaacatcag	caaaactggt	tgcattggaca	aaagcagcag	attcaacaag	accgatgaca	1440
	tatggcgata	acaaaattaa	agcgaactgg	aacgaatcaa	atacaatggg	cgataatctg	1500
30	acagcaaatg	gcggagttgt	tggcacaat	tattcagatg	gcgcaaacta	tgataaaatt	1560
	cgtacaacac	atccgtcatg	ggcaatttat	ggctcagaaa	cagcatcagc	gattaatagc	1620
35	cgtggcattt	ataatagaac	aacaggcgga	gcacaatcat	cagataaaca	gctgacaagc	1680
	tatgataaatt	cagcagttgg	ctggggagca	gttgcatcat	cagcatggta	tgatgttggt	1740
	cagagagatt	ttgtcgcagg	cacatatgtt	tggacaggat	ttgattatct	gggcgaaccg	1800
40	acaccgtgga	atggcacagg	ctcaggcgca	gttggctcat	ggccgtcacc	gaaaaatagc	1860
	tatttttgga	tcgttgatac	agcaggcttt	ccgaaagata	catattattt	ttatcagagc	1920
	cagtggaatg	atgatgttca	tacactgcat	attcttccgg	catggaatga	aaatgttggt	1980
45	gcaaaaggct	caggcaataa	tgttccgggt	gtcgtttata	cagatgcagc	gaaagtgaaa	2040
	ctgtatttta	caccgaaagg	ctcaacagaa	aaaagactga	tcggcgaaaa	atcatttaca	2100
	aaaaaaacaa	cagcggcagg	ctatacatat	caagtctatg	aaggcagcga	taaagattca	2160
50	acagcgcata	aaaacatgta	tctgacatgg	aatgttccgt	gggcagaagg	cacaatttca	2220
	gcggaagcgt	atgatgaaaa	taatcgcttg	attccggaag	gcagcacaga	aggcaacgca	2280
55	tcagttacaa	caacaggcaa	agcagcaaaa	ctgaaagcag	atgcggatcg	caaaacaatt	2340
	acagcggatg	gcaaagatct	gtcatatatt	gaagtcgatg	tcacagatgc	aaatggccat	2400

EP 3 473 713 A1

attgttccgg atgcagcaaa tagagtcaca tttgatgtta aaggcgcagg caaactgggt 2460

ggcggttgata atggctcatc accggatcat gattcatatc aagcggataa cgcgaaagca 2520

5 ttttcaggca aagtcctggc aattgttcag tcaacaaaag aagcaggcga aattacagtt 2580

acagcaaaaag cagatggcct gcaatcaagc acagttaaaa ttgcaacaac agcagttccg 2640

ggaacaagca cagaaaaaac a 2661

10

<210> 10
 <211> 2895
 <212> DNA
 <213> Bifidobacterium bifidum

15

<220>
 <221> source
 <222> 1..2895
 <223> /mol_type="DNA"
 /organism="Bifidobacterium bifidum"

20

<400> 10
 gttgaagatg caacaagaag cgatagcaca acacaaatgt catcaacacc ggaagttggt 60

tattcatcag cggtcgatag caaacaaaat cgcacaagcg attttgatgc gaactggaaa 120

25 tttatgctgt cagatagcgt tcaagcacia gatccggcat ttgatgattc agcatggcaa 180

caagttgatc tgccgcatga ttatagcatc acacagaaat atagccaaag caatgaagca 240

gaatcagcat atcttccggg aggcacaggc tggatataga aaagctttac aattgataga 300

30 gatctggcag gcaaacgcat tgcgattaat tttgatggcg tctatatgaa tgcaacagtc 360

tggtttaatg gcgttaaact gggcacacat ccgatgggt attcaccgtt ttcatttgat 420

35 ctgacaggca atgcaaaatt tggcggagaa aacacaattg tcgtcaaagt tgaaaataga 480

ctgccgtcat caagatggta ttcaggcagc ggcatattata gagatgttac actgacagtt 540

acagatggcg ttcatgttgg caataatggc gtcgcaatta aaacaccgtc actggcaaca 600

40 caaaatggcg gagatgtcac aatgaacctg acaacaaaag tcgcgaatga tacagaagca 660

gcagcgaaca ttacactgaa acagacagtt tttccgaaag gcggaaaaac ggatgcagca 720

attggcacag ttacaacagc atcaaaatca attgcagcag gcgcatcagc agatgttaca 780

45 agcacaatta cagcagcaag cccgaaactg tggtaatta aaaacccgaa cctgtatata 840

gtagaacag aagttctgaa cggaggcaaa gttctggata catatgatac agaatatggc 900

tttcgctgga caggctttga tgcaacatca ggcttttcac tgaatggcga aaaagtcaaa 960

50 ctgaaaggcg ttagcatgca tcatgatcaa ggctcacttg gcgcagttgc aaatagacgc 1020

gcaattgaaa gacaagtcga aatcctgcaa aaaatggcg tcaatagcat tcgcacaaca 1080

55 cataatccgg cagcaaaagc actgattgat gtctgcaatg aaaaaggcgt tctggttgct 1140

gaagaagtct ttgatatgtg gaaccgcagc aaaaatggca acacggaaga ttatggcaaa 1200

EP 3 473 713 A1

	tggtttggcc aagcaattgc aggcgataat gcagttctgg gaggcgataa agatgaaaca	1260
	tgggcgaaat ttgatcttac atcaacaatt aaccgcgata gaaatgcacc gtcagttatt	1320
5	atgtggtcac tgggcaatga aatgatggaa ggcatttcag gctcagtttc aggccttccg	1380
	gcaacatcag caaaactggg tgcattggaca aaagcagcag attcaacaag accgatgaca	1440
	tatggcgata acaaaattaa agcgaactgg aacgaatcaa atacaatggg cgataatctg	1500
10	acagcaaattg gcggagttgt tggcacaatt tattcagatg gcgcaaacta tgataaaatt	1560
	cgtacaacac atccgtcatg ggcaatttat ggctcagaaa cagcatcagc gattaatagc	1620
	cgtggcattt ataatagaac aacaggcgga gcacaatcat cagataaaca gctgacaagc	1680
15	tatgataatt cagcagttgg ctggggagca gttgcatcat cagcatggta tgatgttggt	1740
	cagagagatt ttgtcgcagg cacatatggt tggacaggat ttgattatct gggcgaaaccg	1800
	acaccgtgga atggcacagg ctcaggcgca gttggctcat ggccgtcacc gaaaaatagc	1860
20	tattttggca tcgttgatac agcaggcttt ccgaaagata catattattt ttatcagagc	1920
	cagtggaaatg atgatgttca tacactgcat attcttccgg catggaatga aaatgttggt	1980
25	gcaaaaggct caggcaataa tgttccgggt gtcgtttata cagatgcagc gaaagtgaaa	2040
	ctgtatttta caccgaaagg ctcaacagaa aaaagactga tcggcgaaaa atcattttaca	2100
	aaaaaaacaa cagcggcagg ctatacatat caagtctatg aaggcagcga taaagattca	2160
30	acagcgcata aaaacatgta tctgacatgg aatgttccgt gggcagaagg cacaatttca	2220
	gcggaagcgt atgatgaaaa taatcgcctg attccggaag gcagcacaga aggcaacgca	2280
	tcagttacaa caacaggcaa agcagcaaaa ctgaaagcag atgcggatcg caaaacaatt	2340
35	acagcggatg gcaaagatct gtcatatatt gaagtcgatg tcacagatgc aaatggccat	2400
	attgttccgg atgcagcaaa tagagtcaca tttgatgtta aaggcgcagg caaactgggt	2460
40	ggcgttgata atggctcatc accggatcat gattcatatc aagcggataa ccgcaaagca	2520
	ttttcaggca aagtcctggc aattgttcag tcaacaaaag aagcaggcga aattacagtt	2580
	acagcaaaaag cagatggcct gcaatcaagc acagttaaaa ttgcaacaac agcagttccg	2640
45	ggaacaagca cagaaaaaac agtccgcagc ttttattaca gccgcaacta ttatgtcaaa	2700
	acaggcaaca aaccgattct gccgtcagat gttgaagttc gctattcaga tggaacaagc	2760
	gatagacaaa acgttacatg ggatgcagtt tcagatgatc aaattgcaaa agcaggctca	2820
50	ttttcagttg caggcacagt tgcaggccaa aaaattagcg ttcgcgtcac aatgattgat	2880
	gaaattggcg cactg	2895
55	<210> 11 <211> 3114 <212> DNA	

EP 3 473 713 A1

<213> Bifidobacterium bifidum

<220>

<221> source

<222> 1..3114

<223> /mol_type="DNA"
/organism="Bifidobacterium bifidum"

<400> 11

```

gttgaagatg caacaagaag cgatagcaca acacaaatgt catcaacacc ggaagttggt      60
tattcatcag cggtcgatag caaacaaaat cgcacaagcg attttgatgc gaactggaaa      120
tttatgctgt cagatagcgt tcaagcaca gatccggcat ttgatgattc agcatggcaa      180
caagttgatc tgccgcatga ttatagcatc acacagaaat atagccaaag caatgaagca      240
gaatcagcat atcttccggg aggcacaggc tggtagataa aaagctttac aattgataga      300
gatctggcag gcaaacgcat tgcgattaat tttgatggcg tctatatgaa tgcaacagtc      360
tggtttaatg gcgttaaact gggcacacat ccgtatggct attcaccgtt ttcatttgat      420
ctgacaggca atgcaaaatt tggcggagaa aacacaattg tcgtcaaagt tgaaaataga      480
ctgccgtcat caagatggta ttcaggcagc ggcattttata gagatgttac actgacagtt      540
acagatggcg ttcattgttg caataatggc gtcgcaatta aaacaccgtc actggcaaca      600
caaaatggcg gagatgtcac aatgaacctg acaacaaaag tcgcgaatga tacagaagca      660
gcagcgaaca ttacactgaa acagacagtt tttccgaaag gcggaaaaac ggatgcagca      720
attggcacag ttacaacagc atcaaaatca attgcagcag gcgcatcagc agatgttaca      780
agcacaatta cagcagcaag cccgaaactg tggtaatta aaaacccgaa cctgtatata      840
gttagaacag aagttctgaa cggaggcaaa gttctggata catatgatac agaatatggc      900
tttcgctgga caggctttga tgcaacatca ggcttttcac tgaatggcga aaaagtcaaa      960
ctgaaaggcg ttagcatgca tcatgatcaa ggctcacttg gcgcagttgc aaatagacgc     1020
gcaattgaaa gacaagtcga aatcctgcaa aaaatgggcg tcaatagcat tcgcacaaca     1080
cataatccgg cagcaaaagc actgattgat gtctgcaatg aaaaaggcgt tctggttgtc     1140
gaagaagtct ttgatatgtg gaaccgcagc aaaaatggca acacggaaga ttatggcaaa     1200
tggtttggcc aagcaattgc aggcgataat gcagttctgg gaggcgataa agatgaaaca     1260
tgggcgaaat ttgatcttac atcaacaatt aaccgcgata gaaatgcacc gtcagttatt     1320
atgtggtcac tgggcaatga aatgatggaa ggcatttcag gctcagtttc aggctttccg     1380
gcaacatcag caaaactggg tgcattggca aaagcagcag attcaacaag accgatgaca     1440
tatggcgata acaaaattaa agcgaactgg aacgaatcaa atacaatggg cgataatctg     1500
acagcaaatg gcggagttgt tggcacaat tattcagatg gcgcaaaacta tgataaaatt     1560
cgtacaacac atccgtcatg ggcaatttat ggctcagaaa cagcatcagc gattaatagc     1620

```

EP 3 473 713 A1

5 cgtggcattt ataatagaac aacaggcgga gcacaatcat cagataaaca gctgacaagc 1680
 tatgataatt cagcagttgg ctggggagca gttgcatcat cagcatggta tgatgttggt 1740
 10 cagagagatt ttgtcgcagg cacatatgtt tggacaggat ttgattatct gggcgaaccg 1800
 acaccgtgga atggcacagg ctcaggcgca gttggctcat ggccgtcacc gaaaaatagc 1860
 tatttttgga tcgttgatac agcaggcttt ccgaaagata catattatct ttatcagagc 1920
 15 cagtggaatg atgatgttca tacactgcat attcttccgg catggaatga aaatgttggt 1980
 gcaaaaggct caggcaataa tgttccgggt gtcgtttata cagatgcagc gaaagtgaag 2040
 ctgtatttta caccgaaagg ctcaacagaa aaaagactga tcggcgaaaa atcatttaca 2100
 20 aaaaaaacia cagcggcagg ctatacatat caagtctatg aaggcagcga taaagattca 2160
 acagcgcata aaaacatgta tctgacatgg aatgttccgt gggcagaagg cacaatttca 2220
 gcggaagcgt atgatgaaaa taatcgcttg attccggaag gcagcacaga aggcaacgca 2280
 25 tcagttacaa caacaggcaa agcagcaaaa ctgaaagcag atgcggatcg caaaacaatt 2340
 acagcggatg gcaaagatct gtcatatatt gaagtcgatg tcacagatgc aaatggccat 2400
 attgttccgg atgcagcaaa tagagtcaca tttgatgtta aaggcgcagg caaactgggt 2460
 ggcgttgata atggctcatc accggatcat gattcatatc aagcggataa ccgcaaagca 2520
 ttttcaggca aagtcctggc aattgttcag tcaacaaaag aagcaggcga aattacagtt 2580
 30 acagcaaaaag cagatggcct gcaatcaagc acagttaaaa ttgcaacaac agcagttccg 2640
 ggaacaagca cagaaaaaac agtcgcgagc ttttattaca gccgcaacta ttatgtcaaa 2700
 acaggcaaca aaccgattct gccgtcagat gttgaagttc gctattcaga tggaacaagc 2760
 35 gatagacaaa acgttacatg ggatgcagtt tcagatgatc aaattgcaaa agcagggtca 2820
 ttttcagttg caggcacagt tgcaggccaa aaaattagcg ttcgcgtcac aatgattgat 2880
 gaaattggcg cactgctgaa ttattcagca agcacaccgg ttggcacacc ggcagttctt 2940
 40 ccgggatcaa gaccggcagt cctgccggat ggcacagtca catcagcaaa ttttgcagtc 3000
 cattggacaa aaccggcaga tacagtctat aatacagcag gcacagtcaa agtaccggga 3060
 acagcaacag tttttggcaa agaattttaa gtcacagcga caattagagt tcaa 3114
 45
 <210> 12
 <211> 3426
 <212> DNA
 <213> Bifidobacterium bifidum
 50
 <220>
 <221> source
 <222> 1..3426
 <223> /mol_type="DNA"
 55 /organism="Bifidobacterium bifidum"
 <400> 12

EP 3 473 713 A1

	gttgaagatg caacaagaag cgatagcaca acacaaatgt catcaacacc ggaagttggt	60
	tattcatcag cggctgatag caaacaaaat cgcacaagcg attttgatgc gaactggaaa	120
5	tttatgctgt cagatagcgt tcaagcacia gatccggcat ttgatgattc agcatggcaa	180
	caagttgatc tgccgcatga ttatagcatc acacagaaat atagccaaag caatgaagca	240
	gaatcagcat atcttccggg aggcacaggc tggatataga aaagctttac aattgataga	300
10	gatctggcag gcaaacgcat tgcgattaat tttgatggcg tctatatgaa tgcaacagtc	360
	tggtttaatg gcgttaaact gggcacacat ccgtatggct attcaccgtt ttcatttgat	420
	ctgacaggca atgcaaaatt tggcggagaa aacacaattg tcgtcaaagt tgaaaaataga	480
15	ctgccgtcat caagatggta ttcaggcagc ggcattttata gagatgttac actgacagtt	540
	acagatggcg ttcattgttg caataatggc gtcgcaatta aaacaccgtc actggcaaca	600
	caaaatggcg gagatgtcac aatgaacctg acaacaaaag tcgcgaatga tacagaagca	660
20	gcagcgaaca ttacactgaa acagacagtt tttccgaaag gcggaaaaac ggatgcagca	720
	attggcacag ttacaacagc atcaaaatca attgcagcag gcgcatcagc agatgttaca	780
25	agcacaatta cagcagcaag cccgaaactg tggtaatta aaaacccgaa cctgtatata	840
	gttagaacag aagttctgaa cggaggcaaa gttctggata catatgatac agaatatggc	900
	tttcgctgga caggctttga tgcaacatca ggcttttcac tgaatggcga aaaagtcaaa	960
30	ctgaaaggcg ttagcatgca tcatgatcaa ggctcacttg gcgcagttgc aaatagacgc	1020
	gcaattgaaa gacaagtcga aatcctgcaa aaaatggcg tcaatagcat tcgcacaaca	1080
	cataatccgg cagcaaaagc actgattgat gtctgcaatg aaaaaggcgt tctggttgtc	1140
35	gaagaagtct ttgatatgtg gaaccgcagc aaaaatggca acacggaaga ttatggcaaa	1200
	tggtttggcc aagcaattgc aggcgataat gcagttctgg gaggcgataa agatgaaaca	1260
	tgggcgaaat ttgatcttac atcaacaatt aaccgcgata gaaatgcacc gtcagttatt	1320
40	atgtggtcac tgggcaatga aatgatggaa ggcatttcag gctcagtttc aggctttccg	1380
	gcaacatcag caaaaactgg tgcattggaca aaagcagcag attcaacaag accgatgaca	1440
	tatggcgata acaaaattaa agcgaactgg aacgaatcaa atacaatggg cgataatctg	1500
45	acagcaaagc gcggagttgt tggcaciaat tattcagatg gcgcaacta tgataaaatt	1560
	cgtacaacac atccgtcatg ggcaatttat ggctcagaaa cagcatcagc gattaatagc	1620
50	cgtggcattt ataatagaac aacaggcgga gcacaatcat cagataaaca gctgacaagc	1680
	tatgataatt cagcagttgg ctggggagca gttgcatcat cagcatggta tgatgttgtt	1740
	cagagagatt ttgtcgcagg cacatatgtt tggacaggat ttgattatct gggcgaaccg	1800
55	acaccgtgga atggcacagg ctgaggcgca gttggctcat ggccgtcacc gaaaaatagc	1860
	tattttggca tcgttgatac agcaggcttt ccgaaagata catattattt ttatcagagc	1920

EP 3 473 713 A1

	cagtggaatg atgatgttca tacactgcat attcttccgg catggaatga aaatgttggt	1980
	gcaaaaggct caggcaataa tgttccggtt gtcgtttata cagatgcagc gaaagtga	2040
5	ctgtatttta caccgaaagg ctcaacagaa aaaagactga tcggcgaaaa atcatttaca	2100
	aaaaaaacaa cagcggcagg ctatacatat caagtctatg aaggcagcga taaagattca	2160
	acagcgcata aaaacatgta tctgacatgg aatgttccgt gggcagaagg cacaatttca	2220
10	gcggaagcgt atgatgaaaa taatcgccgt attccggaag gcagcacaga aggcaacgca	2280
	tcagttacaa caacaggcaa agcagcaaaa ctgaaagcag atgcggatcg caaaacaatt	2340
	acagcggatg gcaaagatct gtcatatatt gaagtogatg tcacagatgc aaatggccat	2400
15	attgttccgg atgcagcaaa tagagtcaca tttgatgtta aaggcgcagg caaactgggt	2460
	ggcgttgata atggctcatc accggatcat gattcatatc aagcggataa ccgcaaagca	2520
	ttttcaggca aagtcctggc aattgttcag tcaacaaaag aagcaggcga aattacagtt	2580
20	acagcaaaag cagatggcct gcaatcaagc acagttaaaa ttgcaacaac agcagttccg	2640
	ggaacaagca cagaaaaaac agtccgcagc ttttattaca gccgcaacta ttatgtcaaa	2700
25	acaggcaaca aaccgattct gccgtcagat gttgaagttc gctattcaga tggaacaagc	2760
	gatagacaaa acgttacatg ggatgcagtt tcagatgatc aaattgcaaa agcaggctca	2820
	ttttcagttg caggcacagt tgcaggccaa aaaattagcg ttcgcgtcac aatgattgat	2880
30	gaaattggcg cactgctgaa ttattcagca agcacaccgg ttggcacacc ggcagttctt	2940
	ccgggatcaa gaccggcagt cctgccggat ggcacagtca catcagcaaa ttttgcagtc	3000
	cattggacaa aaccggcaga tacagtctat aatacagcag gcacagtcaa agtaccggga	3060
35	acagcaacag tttttggcaa agaatttaaa gtcacagcga caattagagt tcaaagaagc	3120
	caagttacaa ttggctcatc agtttcagga aatgcactga gactgacaca aaatattccg	3180
	gcagataaac aatcagatac actggatgcg attaaagatg gctcaacaac agttgatgca	3240
40	aatacaggcg gaggcgcaaa tccgtcagca tggacaaatt gggcatattc aaaagcaggc	3300
	cataacacag cggaaattac atttgaatat gcgacagAAC aacaactggg ccagatcgtc	3360
45	atgtattttt ttcgcgatag caatgcagtt agatttccgg atgctggcaa aacaaaaatt	3420
	cagatc	3426
50	<210> 13	
	<211> 3633	
	<212> DNA	
	<213> Bifidobacterium bifidum	
55	<220>	
	<221> source	
	<222> 1..3633	
	<223> /mol_type="DNA"	

EP 3 473 713 A1

/organism="Bifidobacterium bifidum"

<400> 13

5	gttgaagatg caacaagaag cgatagcaca acacaaatgt catcaacacc ggaagttgtt	60
	tattcatcag cggtcgatag caaacaaaat cgcacaagcg attttgatgc gaactggaaa	120
	tttatgctgt cagatagcgt tcaagcacaa gatccggcat ttgatgattc agcatggcaa	180
10	caagttgatc tgccgcatga ttatagcatc acacagaaat atagccaaag caatgaagca	240
	gaatcagcat atcttccggg aggcacaggc tggatatagaa aaagctttac aattgataga	300
	gatctggcag gcaaacgcat tgcgattaat tttgatggcg tctatatgaa tgcaacagtc	360
15	tggtttaatg gcgttaaact gggcacacat ccgtatggct attcaccgtt ttcatttgat	420
	ctgacaggca atgcaaaatt tggcggagaa aacacaattg tcgtcaaagt tgaaaataga	480
	ctgccgtcat caagatggta ttcaggcagc ggcatttata gagatgttac actgacagtt	540
20	acagatggcg ttcattgttg caataatggc gtcgcaatta aaacaccgtc actggcaaca	600
	caaatggcg gagatgtcac aatgaacctg acaacaaaag tcgcgaatga tacagaagca	660
	gcagcgaaca ttacttgaa acagacagtt tttccgaaag gcggaaaaac ggatgcagca	720
25	attggcacag ttacaacagc atcaaatca attgcagcag gcgcacagc agatgttaca	780
	agcacaatta cagcagcaag cccgaaactg tggtaatta aaaacccgaa cctgtatata	840
30	gttagaacag aagttctgaa cggaggcaaa gttctggata catatgatac agaatatggc	900
	tttcgctgga caggctttga tgcaacatca ggcttttcac tgaatggcga aaaagtcaaa	960
	ctgaaaggcg ttagcatgca tcatgatcaa ggctcacttg gcgcagttgc aaatagacgc	1020
35	gcaattgaaa gacaagtcga aatcctgcaa aaaatgggcg tcaatagcat tcgcacaaca	1080
	cataatccgg cagcaaaagc actgattgat gtctgcaatg aaaaaggcgt tctggttgtc	1140
	gaagaagtct ttgatatgtg gaaccgcagc aaaaatggca acacggaaga ttatggcaaa	1200
40	tggtttggcc aagcaattgc aggcgataat gcagttctgg gaggcgataa agatgaaaca	1260
	tgggcgaaat ttgatcttac atcaacaatt aaccgcgata gaaatgcacc gtcagttatt	1320
	atgtggtcac tgggcaatga aatgatggaa ggcatttcag gctcagtttc aggctttccg	1380
45	gcaacatcag caaaactggt tgcattggaca aaagcagcag attcaacaag accgatgaca	1440
	tatggcgata acaaaattaa agcgaactgg aacgaatcaa atacaatggg cgataatctg	1500
	acagcaaatg gcggagttgt tggcacaaat tattcagatg gcgcaacta tgataaaatt	1560
50	cgtacaacac atccgtcatg ggcaatttat ggctcagaaa cagcatcagc gattaatagc	1620
	cgtggcattt ataatagaac aacaggcgga gcacaatcat cagataaaca gctgacaagc	1680
55	tatgataatt cagcagttgg ctggggagca gttgcatcat cagcatggta tgatgttgtt	1740
	cagagagatt ttgtcgcagg cacatatgtt tggacaggat ttgattatct gggcgaaccg	1800

EP 3 473 713 A1

	acaccgtgga atggcacagg ctcaggcgca gttggctcat ggccgtcacc gaaaaatagc	1860
	tatTTTTggca tcgttgatac agcagggttt cggaaagata catattattt ttatcagagc	1920
5	cagtggaaatg atgatgttca tacactgcat attcttccgg catggaatga aaatgttggt	1980
	gcaaaaggct caggcaataa tgttccgggt gtctgtttata cagatgcagc gaaagtgaaa	2040
	ctgtatttta caccgaaagg ctcaacagaa aaaagactga tcggcgaaaa atcatttaca	2100
10	aaaaaaacaa cagcggcagg ctatacatat caagtctatg aaggcagcga taaagattca	2160
	acagcgcata aaaacatgta tctgacatgg aatgttccgt gggcagaagg cacaatttca	2220
	gcggaagcgt atgatgaaaa taatcgctg attccggaag gcagcacaga aggcaacgca	2280
15	tcagttacaa caacaggcaa agcagcaaaa ctgaaagcag atgcggatcg caaaacaatt	2340
	acagcggatg gcaaaagatct gtcatatatt gaagtcgatg tcacagatgc aaatggccat	2400
20	attgttccgg atgcagcaaa tagagtcaca tttgatgtta aaggcgcagg caaactgggt	2460
	ggcgttgata atggctcatc accggatcat gattcatatc aagcggataa ccgcaaagca	2520
	ttttcaggca aagtcctggc aattgttcag tcaacaaaag aagcaggcga aattacagtt	2580
25	acagcaaaag cagatggcct gcaatcaagc acagttaaaa ttgcaacaac agcagttccg	2640
	ggaacaagca cagaaaaaac agtccgcagc ttttattaca gccgcaacta ttatgtcaaa	2700
	acaggcaaca aaccgattct gccgtcagat gttgaagttc gctattcaga tggaacaagc	2760
30	gatagacaaa acgttacatg ggatgcagtt tcagatgatc aaattgcaaa agcaggctca	2820
	ttttcagttg caggcacagt tgcaggccaa aaaattagcg ttccgcgtcac aatgattgat	2880
35	gaaattggcg cactgctgaa ttattcagca agcacaccgg ttggcacacc ggcagttctt	2940
	ccgggatcaa gaccggcagt cctgccggat ggcacagtca catcagcaaa ttttgcagtc	3000
	cattggacaa aaccggcaga tacagtctat aatacagcag gcacagtcaa agtaccggga	3060
40	acagcaacag tttttggcaa agaatttaaa gtcacagcga caattagagt tcaaagaagc	3120
	caagttacaa ttggctcatc agtttcagga aatgcactga gactgacaca aaatattccg	3180
	gcagataaac aatcagatac actggatgcg attaaagatg gctcaacaac agttgatgca	3240
45	aatacaggcg gaggcgcaaa tccgtcagca tggacaaatt gggcatattc aaaagcaggc	3300
	cataacacag cggaaattac atttgaatat gcgacagAAC aacaactggg ccagatcgtc	3360
	atgtattttt ttccgcgatag caatgcagtt agatttccgg atgctggcaa aacaaaaatt	3420
50	cagatcagcg cagatggcaa aaattggaca gatctggcag caacagaaac aattgcagcg	3480
	caagaatcaa gcgatagagt caaaccgat acatatgatt ttgcaccggt tggcgcaaca	3540
55	tttgtaaag tgacagtcac aaacgcagat acaacaacac cgtcaggcgt tgtttgcgca	3600
	ggcctgacag aaattgaact gaaaacagcg aca	3633

EP 3 473 713 A1

<210> 14
<211> 3888
<212> DNA
<213> Bifidobacterium bifidum

<220>
<221> source
<222> 1..3888
<223> /mol_type="DNA"
/organism="Bifidobacterium bifidum"

<400> 14
gttgaagatg caacaagaag cgatagcaca acacaaatgt catcaacacc ggaagttggt 60
tattcatcag cggtcgatag caaacaaaat cgcacaagcg attttgatgc gaactggaaa 120
tttatgctgt cagatagcgt tcaagcacia gatccggcat ttgatgattc agcatggcaa 180
caagttgatc tgccgcatga ttatagcatc acacagaaat atagccaaag caatgaagca 240
gaatcagcat atcttccggg aggcacaggc tggatataga aaagctttac aattgataga 300
gatctggcag gcaaacgcat tgcgattaat tttgatggcg tctatatgaa tgcaacagtc 360
tggtttaatg gcgttaaact gggcacacat ccgtatggct attcaccgtt ttcatttgat 420
ctgacaggca atgcaaaatt tggcggagaa aacacaattg tcgtcaaagt tgaaaataga 480
ctgccgtcat caagatggta ttcaggcagc ggcatttata gagatgttac actgacagtt 540
acagatggcg ttcattgttg caataatggc gtgcgaatta aaacaccgtc actggcaaca 600
caaaatggcg gagatgtcac aatgaacctg acaacaaaag tcgcgaatga tacagaagca 660
gcagcgaaca ttacactgaa acagacagtt tttccgaaag gcggaaaaac ggatgcagca 720
attggcacag ttacaacagc atcaaaatca attgcagcag gcgcatcagc agatgttaca 780
agcacaatta cagcagcaag cccgaaactg tggtaatta aaaacccgaa cctgtataga 840
gttagaacag aagttctgaa cggaggcaaa gttctggata catatgatac agaatatggc 900
tttcgctgga caggctttga tgcaacatca ggcttttcac tgaatggcga aaaagtcaaa 960
ctgaaaggcg ttagcatgca tcatgatcaa ggctcacttg gcgcagttgc aaatagacgc 1020
gcaattgaaa gacaagtcga aatcctgcaa aaaatggcg tcaatagcat tcgcacaaca 1080
cataatccgg cagcaaaagc actgattgat gtctgcaatg aaaaaggcgt tctggttgtc 1140
gaagaagtct ttgatatgtg gaaccgcagc aaaaatggca acacggaaga ttatggcaaa 1200
tggtttggcc aagcaattgc aggcgataat gcagttctgg gaggcgataa agatgaaaca 1260
tgggcgaaat ttgatcttac atcaacaatt aaccgcgata gaaatgcacc gtcagttatt 1320
atgtggtcac tgggcaatga aatgatggaa ggcatttcag gctcagtttc aggctttccg 1380
gcaacatcag caaaactggg tgcattggca aaagcagcag attcaacaag accgatgaca 1440
tatggcgata acaaaattaa agcgaactgg aacgaatcaa atacaatggg cgataatctg 1500
acagcaaatg gcggagttgt tggcaciaat tattcagatg gcgcaaacta tgataaaatt 1560

EP 3 473 713 A1

	cgtacaacac atccgtcatg ggcaatttat ggctcagaaa cagcatcagc gattaatagc	1620
	cgtggcattt ataatagaac aacaggcgga gcacaatcat cagataaaca gctgacaagc	1680
5	tatgataatt cagcagttgg ctggggagca gttgcatcat cagcatggta tgatgttggt	1740
	cagagagatt ttgtcgcagg cacatatgtt tggacaggat ttgattatct gggcgaaccg	1800
	acaccgtgga atggcacagg ctcaggcgca gttgggtcat ggccgtcacc gaaaaatagc	1860
10	tatdddggca tcgttgatac agcaggcttt ccgaaagata catattattt ttatcagagc	1920
	cagtggaaatg atgatgttca tacactgcat attcttccgg catggaaatga aaatgttggt	1980
	gcaaaaggct caggcaataa tgttccgggt gtogtttata cagatgcagc gaaagtgaag	2040
15	ctgtatdtda caccgaaagg ctcaacagaa aaaagactga tcggcgaaaa atcatttaca	2100
	aaaaaaacaa cagcggcagg ctatacatat caagtctatg aaggcagcga taaagattca	2160
20	acagcgcata aaaacatgta tctgacatgg aatgttccgt gggcagaagg cacaatttca	2220
	gcggaagcgt atgatgaaaa taatcgctg attccggaag gcagcacaga aggcaacgca	2280
	tcagttacaa caacaggcaa agcagcaaaa ctgaaagcag atgcggatcg caaaacaatt	2340
25	acagcggatg gcaaagatct gtcatatatt gaagtogatg tcacagatgc aaatggccat	2400
	attgttccgg atgcagcaaa tagagtcaca tttgatgtta aaggcgcagg caaactgggt	2460
	ggcgttgata atggctcatc accggatcat gattcatatc aagcggataa ccgcaaagca	2520
30	ttttcaggca aagtcctggc aattgttcag tcaacaaaag aagcaggcga aattacagtt	2580
	acagcaaaag cagatggcct gcaatcaagc acagttaaaa ttgcaacaac agcagttccg	2640
35	ggaacaagca cagaaaaaac agtccgcagc ttttattaca gccgcaacta ttatgtcaaa	2700
	acaggcaaca aaccgattct gccgtcagat gttgaagtgc gctattcaga tggacaagc	2760
	gatagacaaa acgttacatg ggatgcagtt tcagatgatc aaattgcaaa agcaggctca	2820
40	ttttcagttg caggcacagt tgcaggccaa aaaattagcg ttccggtcac aatgattgat	2880
	gaaattggcg cactgctgaa ttattcagca agcacaccgg ttggcacacc ggcagttctt	2940
	ccgggatcaa gaccggcagt cctgccggat ggcacagtca catcagcaaa ttttgcagtc	3000
45	cattggacaa aaccggcaga tacagtctat aatacagcag gcacagtcaa agtaccggga	3060
	acagcaacag tttttggcaa agaattttaa gtcacagcga caattagagt tcaaagaagc	3120
	caagttacaa ttgggtcatc agtttcagga aatgcactga gactgacaca aaatattccg	3180
50	gcagataaac aatcagatac actggatgag attaaagatg gctcaacaac agttgatgca	3240
	aatacaggcg gaggcgcaaa tccgtcagca tggacaaatt gggcatattc aaaagcaggc	3300
55	cataacacag cggaaattac atttgaatat gcgacagaac aacaactggg ccagatcgtc	3360
	atgtatdtdt ttccgcatag caatgcagtt agatttccgg atgctggcaa aacaaaaatt	3420

EP 3 473 713 A1

cagatcagcg cagatggcaa aaattggaca gatctggcag caacagaaac aattgcagcg 3480
 caagaatcaa gcgatagagt caaaccgat acatatgatt ttgcaccggt tggcgcaaca 3540
 5 tttgttaaag tgacagtcac aaacgcagat acaacaacac cgtcaggcgt tgtttgcgca 3600
 ggcctgacag aaattgaact gaaaacagcg acaagcaaat ttgtcacaaa tacatcagca 3660
 gcactgtcat cacttacagt caatggcaca aaagtttcag attcagttct ggcagcaggc 3720
 10 tcatataaca caccggcaat tatcgagat gttaaagcgg aaggcgaagg caatgcaagc 3780
 gttacagtcc ttccggcaca tgataatgtt attcgcgtca ttacagaaag cgaagatcat 3840
 gtcacacgca aaacatttac aatcaacctg ggcacagaac aagaattt 3888
 15
 <210> 15
 <211> 30
 <212> DNA
 <213> artificial sequences
 20
 <220>
 <221> source
 <222> 1..30
 <223> /mol_type="DNA"
 /note="Synthetic DNA primer"
 25 /organism="artificial sequences"
 <400> 15
 ggggtaacta gtggaagatg caacaagaag 30
 30
 <210> 16
 <211> 35
 <212> DNA
 <213> artificial sequences
 35
 <220>
 <221> source
 <222> 1..35
 <223> /mol_type="DNA"
 /note="Synthetic DNA primer"
 40 /organism="artificial sequences"
 <400> 16
 gcgcttaatt aattatgttt tttctgtgct tgttc 35
 45
 <210> 17
 <211> 38
 <212> DNA
 <213> artificial sequences
 50
 <220>
 <221> source
 <222> 1..38
 <223> /mol_type="DNA"
 /note="Synthetic PCR primer"
 /organism="artificial sequences"
 55
 <400> 17
 gcgcttaatt aattacagtg cgccaatttc atcaatca 38

<210> 18
 <211> 35
 <212> DNA
 <213> artificial sequences
 5
 <220>
 <221> source
 <222> 1..35
 <223> /mol_type="DNA"
 10 /note="Synthetic PCR primer"
 /organism="artificial sequences"
 <400> 18
 gcgcttaatt aattattgaa ctctaattgt cgctg 35
 15
 <210> 19
 <211> 36
 <212> DNA
 <213> artificial sequences
 20
 <220>
 <221> source
 <222> 1..36
 <223> /mol_type="DNA"
 25 /note="Synthetic PCR primer"
 /organism="artificial sequences"
 <400> 19
 gcgcttaatt aattatgtcg ctgttttcag ttcaat 36
 30
 <210> 20
 <211> 35
 <212> DNA
 <213> artificial sequences
 35
 <220>
 <221> source
 <222> 1..35
 <223> /mol_type="DNA"
 40 /note="Synthetic PCR primer"
 /organism="artificial sequences"
 <400> 20
 gcgcttaatt aattaaaatt cttgttctgt gccca 35
 45
 <210> 21
 <211> 38
 <212> DNA
 <213> artificial sequences
 50
 <220>
 <221> source
 <222> 1..38
 <223> /mol_type="DNA"
 55 /note="Synthetic PCR primer"
 /organism="artificial sequences"
 <400> 21

gcgcttaatt aattatctca gtctaatttc gcttgccg

38

<210> 22
 <211> 1720
 <212> PRT
 <213> Bifidobacterium bifidum

<220>
 <221> SOURCE
 <222> 1..1720
 <223> /mol_type="protein"
 /organism="Bifidobacterium bifidum"

<400> 22

Val	Glu	Asp	Ala	Thr	Arg	Ser	Asp	Ser	Thr	Thr	Gln	Met	Ser	Ser	Thr
1				5					10					15	
Pro	Glu	Val	Val	Tyr	Ser	Ser	Ala	Val	Asp	Ser	Lys	Gln	Asn	Arg	Thr
		20						25					30		
Ser	Asp	Phe	Asp	Ala	Asn	Trp	Lys	Phe	Met	Leu	Ser	Asp	Ser	Val	Gln
	35					40						45			
Ala	Gln	Asp	Pro	Ala	Phe	Asp	Asp	Ser	Ala	Trp	Gln	Gln	Val	Asp	Leu
	50					55					60				
Pro	His	Asp	Tyr	Ser	Ile	Thr	Gln	Lys	Tyr	Ser	Gln	Ser	Asn	Glu	Ala
65					70					75				80	
Glu	Ser	Ala	Tyr	Leu	Pro	Gly	Gly	Thr	Gly	Trp	Tyr	Arg	Lys	Ser	Phe
				85					90					95	
Thr	Ile	Asp	Arg	Asp	Leu	Ala	Gly	Lys	Arg	Ile	Ala	Ile	Asn	Phe	Asp
			100						105				110		
Gly	Val	Tyr	Met	Asn	Ala	Thr	Val	Trp	Phe	Asn	Gly	Val	Lys	Leu	Gly
		115					120					125			
Thr	His	Pro	Tyr	Gly	Tyr	Ser	Pro	Phe	Ser	Phe	Asp	Leu	Thr	Gly	Asn
	130					135					140				
Ala	Lys	Phe	Gly	Gly	Glu	Asn	Thr	Ile	Val	Val	Lys	Val	Glu	Asn	Arg
145					150					155				160	
Leu	Pro	Ser	Ser	Arg	Trp	Tyr	Ser	Gly	Ser	Gly	Ile	Tyr	Arg	Asp	Val
				165					170					175	
Thr	Leu	Thr	Val	Thr	Asp	Gly	Val	His	Val	Gly	Asn	Asn	Gly	Val	Ala
			180					185					190		
Ile	Lys	Thr	Pro	Ser	Leu	Ala	Thr	Gln	Asn	Gly	Gly	Asp	Val	Thr	Met
		195					200					205			
Asn	Leu	Thr	Thr	Lys	Val	Ala	Asn	Asp	Thr	Glu	Ala	Ala	Ala	Asn	Ile
	210				215					220					
Thr	Leu	Lys	Gln	Thr	Val	Phe	Pro	Lys	Gly	Gly	Lys	Thr	Asp	Ala	Ala
225					230					235				240	
Ile	Gly	Thr	Val	Thr	Thr	Ala	Ser	Lys	Ser	Ile	Ala	Ala	Gly	Ala	Ser
			245						250					255	
Ala	Asp	Val	Thr	Ser	Thr	Ile	Thr	Ala	Ala	Ser	Pro	Lys	Leu	Trp	Ser
			260					265					270		
Ile	Lys	Asn	Pro	Asn	Leu	Tyr	Thr	Val	Arg	Thr	Glu	Val	Leu	Asn	Gly
		275					280					285			
Gly	Lys	Val	Leu	Asp	Thr	Tyr	Asp	Thr	Glu	Tyr	Gly	Phe	Arg	Trp	Thr
	290					295					300				
Gly	Phe	Asp	Ala	Thr	Ser	Gly	Phe	Ser	Leu	Asn	Gly	Glu	Lys	Val	Lys
305					310					315				320	
Leu	Lys	Gly	Val	Ser	Met	His	His	Asp	Gln	Gly	Ser	Leu	Gly	Ala	Val
				325					330					335	
Ala	Asn	Arg	Arg	Ala	Ile	Glu	Arg	Gln	Val	Glu	Ile	Leu	Gln	Lys	Met
			340					345					350		
Gly	Val	Asn	Ser	Ile	Arg	Thr	Thr	His	Asn	Pro	Ala	Ala	Lys	Ala	Leu
	355						360					365			
Ile	Asp	Val	Cys	Asn	Glu	Lys	Gly	Val	Leu	Val	Val	Glu	Glu	Val	Phe
	370					375						380			

EP 3 473 713 A1

	Asp	Met	Trp	Asn	Arg	Ser	Lys	Asn	Gly	Asn	Thr	Glu	Asp	Tyr	Gly	Lys	
	385					390					395					400	
	Trp	Phe	Gly	Gln	Ala	Ile	Ala	Gly	Asp	Asn	Ala	Val	Leu	Gly	Gly	Asp	
				405						410						415	
5	Lys	Asp	Glu	Thr	Trp	Ala	Lys	Phe	Asp	Leu	Thr	Ser	Thr	Ile	Asn	Arg	
				420						425					430		
	Asp	Arg	Asn	Ala	Pro	Ser	Val	Ile	Met	Trp	Ser	Leu	Gly	Asn	Glu	Met	
			435						440					445			
	Met	Glu	Gly	Ile	Ser	Gly	Ser	Val	Ser	Gly	Phe	Pro	Ala	Thr	Ser	Ala	
		450					455					460					
10	Lys	Leu	Val	Ala	Trp	Thr	Lys	Ala	Ala	Asp	Ser	Thr	Arg	Pro	Met	Thr	
	465					470					475					480	
	Tyr	Gly	Asp	Asn	Lys	Ile	Lys	Ala	Asn	Trp	Asn	Glu	Ser	Asn	Thr	Met	
				485							490				495		
	Gly	Asp	Asn	Leu	Thr	Ala	Asn	Gly	Gly	Val	Val	Gly	Thr	Asn	Tyr	Ser	
				500						505					510		
15	Asp	Gly	Ala	Asn	Tyr	Asp	Lys	Ile	Arg	Thr	Thr	His	Pro	Ser	Trp	Ala	
			515						520				525				
	Ile	Tyr	Gly	Ser	Glu	Thr	Ala	Ser	Ala	Ile	Asn	Ser	Arg	Gly	Ile	Tyr	
		530					535					540					
	Asn	Arg	Thr	Thr	Gly	Gly	Ala	Gln	Ser	Ser	Asp	Lys	Gln	Leu	Thr	Ser	
20		545				550					555					560	
	Tyr	Asp	Asn	Ser	Ala	Val	Gly	Trp	Gly	Ala	Val	Ala	Ser	Ser	Ala	Trp	
				565							570					575	
	Tyr	Asp	Val	Val	Gln	Arg	Asp	Phe	Val	Ala	Gly	Thr	Tyr	Val	Trp	Thr	
				580					585						590		
25	Gly	Phe	Asp	Tyr	Leu	Gly	Glu	Pro	Thr	Pro	Trp	Asn	Gly	Thr	Gly	Ser	
		595						600					605				
	Gly	Ala	Val	Gly	Ser	Trp	Pro	Ser	Pro	Lys	Asn	Ser	Tyr	Phe	Gly	Ile	
		610					615					620					
	Val	Asp	Thr	Ala	Gly	Phe	Pro	Lys	Asp	Thr	Tyr	Tyr	Phe	Tyr	Gln	Ser	
		625				630					635					640	
30	Gln	Trp	Asn	Asp	Asp	Val	His	Thr	Leu	His	Ile	Leu	Pro	Ala	Trp	Asn	
				645						650						655	
	Glu	Asn	Val	Val	Ala	Lys	Gly	Ser	Gly	Asn	Asn	Val	Pro	Val	Val	Val	
				660						665					670		
	Tyr	Thr	Asp	Ala	Ala	Lys	Val	Lys	Leu	Tyr	Phe	Thr	Pro	Lys	Gly	Ser	
			675						680					685			
35	Thr	Glu	Lys	Arg	Leu	Ile	Gly	Glu	Lys	Ser	Phe	Thr	Lys	Lys	Thr	Thr	
		690					695					700					
	Ala	Ala	Gly	Tyr	Thr	Tyr	Gln	Val	Tyr	Glu	Gly	Ser	Asp	Lys	Asp	Ser	
		705				710					715					720	
	Thr	Ala	His	Lys	Asn	Met	Tyr	Leu	Thr	Trp	Asn	Val	Pro	Trp	Ala	Glu	
				725							730					735	
40	Gly	Thr	Ile	Ser	Ala	Glu	Ala	Tyr	Asp	Glu	Asn	Asn	Arg	Leu	Ile	Pro	
				740						745					750		
	Glu	Gly	Ser	Thr	Glu	Gly	Asn	Ala	Ser	Val	Thr	Thr	Thr	Gly	Lys	Ala	
			755						760					765			
	Ala	Lys	Leu	Lys	Ala	Asp	Ala	Asp	Arg	Lys	Thr	Ile	Thr	Ala	Asp	Gly	
45		770					775					780					
	Lys	Asp	Leu	Ser	Tyr	Ile	Glu	Val	Asp	Val	Thr	Asp	Ala	Asn	Gly	His	
		785				790					795					800	
	Ile	Val	Pro	Asp	Ala	Ala	Asn	Arg	Val	Thr	Phe	Asp	Val	Lys	Gly	Ala	
				805							810				815		
50	Gly	Lys	Leu	Val	Gly	Val	Asp	Asn	Gly	Ser	Ser	Pro	Asp	His	Asp	Ser	
				820						825					830		
	Tyr	Gln	Ala	Asp	Asn	Arg	Lys	Ala	Phe	Ser	Gly	Lys	Val	Leu	Ala	Ile	
			835					840						845			
	Val	Gln	Ser	Thr	Lys	Glu	Ala	Gly	Glu	Ile	Thr	Val	Thr	Ala	Lys	Ala	
		850					855					860					
55	Asp	Gly	Leu	Gln	Ser	Ser	Thr	Val	Lys	Ile	Ala	Thr	Thr	Ala	Val	Pro	
		865				870					875					880	
	Gly	Thr	Ser	Thr	Glu	Lys	Thr	Val	Arg	Ser	Phe	Tyr	Tyr	Ser	Arg	Asn	

EP 3 473 713 A1

				885					890				895
	Tyr	Tyr	Val	Lys	Thr	Gly	Asn	Lys	Pro	Ile	Leu	Pro	Ser
				900					905				910
	Val	Arg	Tyr	Ser	Asp	Gly	Thr	Ser	Asp	Arg	Gln	Asn	Val
5				915				920				925	Thr
	Ala	Val	Ser	Asp	Asp	Gln	Ile	Ala	Lys	Ala	Gly	Ser	Phe
				930			935					940	Ser
	Gly	Thr	Val	Ala	Gly	Gln	Lys	Ile	Ser	Val	Arg	Val	Thr
	945					950					955		Met
	Glu	Ile	Gly	Ala	Leu	Leu	Asn	Tyr	Ser	Ala	Ser	Thr	Pro
10				965					970				Val
	Pro	Ala	Val	Leu	Pro	Gly	Ser	Arg	Pro	Ala	Val	Leu	Pro
				980					985				Asp
	Val	Thr	Ser	Ala	Asn	Phe	Ala	Val	His	Trp	Thr	Lys	Pro
				995				1000					Ala
	Val	Tyr	Asn	Thr	Ala	Gly	Thr	Val	Lys	Val	Pro	Gly	Thr
15				1010				1015				1020	Ala
	Phe	Gly	Lys	Glu	Phe	Lys	Val	Thr	Ala	Thr	Ile	Arg	Val
	1025					1030					1035		Gln
	Gln	Val	Thr	Ile	Gly	Ser	Ser	Val	Ser	Gly	Asn	Ala	Leu
				1045							1050		Arg
20													Leu
	Gln	Asn	Ile	Pro	Ala	Asp	Lys	Gln	Ser	Asp	Thr	Leu	Asp
				1060					1065				Ala
	Asp	Gly	Ser	Thr	Thr	Val	Asp	Ala	Asn	Thr	Gly	Gly	Gly
				1075				1080					Ala
	Ser	Ala	Trp	Thr	Asn	Trp	Ala	Tyr	Ser	Lys	Ala	Gly	His
	1090					1095						1100	Asn
25	Glu	Ile	Thr	Phe	Glu	Tyr	Ala	Thr	Glu	Gln	Gln	Leu	Gly
	1105					1110					1115		Gln
	Met	Tyr	Phe	Phe	Arg	Asp	Ser	Asn	Ala	Val	Arg	Phe	Pro
				1125							1130		Asp
	Lys	Thr	Lys	Ile	Gln	Ile	Ser	Ala	Asp	Gly	Lys	Asn	Trp
30				1140					1145				Thr
	Ala	Ala	Thr	Glu	Thr	Ile	Ala	Ala	Gln	Glu	Ser	Ser	Asp
				1155				1160					Arg
	Pro	Tyr	Thr	Tyr	Asp	Phe	Ala	Pro	Val	Gly	Ala	Thr	Phe
	1170					1175						1180	Val
	Thr	Val	Thr	Asn	Ala	Asp	Thr	Thr	Thr	Pro	Ser	Gly	Val
35	1185					1190					1195		Val
	Gly	Leu	Thr	Glu	Ile	Glu	Leu	Lys	Thr	Ala	Thr	Ser	Lys
				1205							1210		Phe
	Asn	Thr	Ser	Ala	Ala	Leu	Ser	Ser	Leu	Thr	Val	Asn	Gly
				1220					1225				Thr
	Ser	Asp	Ser	Val	Leu	Ala	Ala	Gly	Ser	Tyr	Asn	Thr	Pro
40				1235				1240					Ala
	Ala	Asp	Val	Lys	Ala	Glu	Gly	Glu	Gly	Asn	Ala	Ser	Val
				1250				1255				1260	Thr
	Pro	Ala	His	Asp	Asn	Val	Ile	Arg	Val	Ile	Thr	Glu	Ser
	1265					1270					1275		Glu
45	Val	Thr	Arg	Lys	Thr	Phe	Thr	Ile	Asn	Leu	Gly	Thr	Glu
				1285						1290			Gln
	Pro	Ala	Asp	Ser	Asp	Glu	Arg	Asp	Tyr	Pro	Ala	Ala	Asp
				1300					1305				Met
	Thr	Val	Gly	Ser	Glu	Gln	Thr	Ser	Gly	Thr	Ala	Thr	Glu
				1315				1320					Gly
50	Lys	Phe	Ala	Val	Asp	Gly	Asn	Thr	Ser	Thr	Tyr	Trp	His
				1330				1335				1340	Ser
	Thr	Pro	Thr	Thr	Val	Asn	Asp	Leu	Trp	Ile	Ala	Phe	Glu
	1345					1350					1355		Leu
	Pro	Thr	Lys	Leu	Asp	Ala	Leu	Arg	Tyr	Leu	Pro	Arg	Pro
				1365					1370				Ala
55	Lys	Asn	Gly	Ser	Val	Thr	Glu	Tyr	Lys	Val	Gln	Val	Ser
				1380					1385				Asp
													Asp
													Gly

EP 3 473 713 A1

```

    Thr Asn Trp Thr Asp Ala Gly Ser Gly Thr Trp Thr Thr Asp Tyr Gly
      1395                      1400                      1405
    Trp Lys Leu Ala Glu Phe Asn Gln Pro Val Thr Thr Lys His Val Arg
      1410                      1415                      1420
5    Leu Lys Ala Val His Thr Tyr Ala Asp Ser Gly Asn Asp Lys Phe Met
    1425                      1430                      1435                      1440
    Ser Ala Ser Glu Ile Arg Leu Arg Lys Ala Val Asp Thr Thr Asp Ile
      1445                      1450                      1455
    Ser Gly Ala Thr Val Thr Val Pro Ala Lys Leu Thr Val Asp Arg Val
      1460                      1465                      1470
10   Asp Ala Asp His Pro Ala Thr Phe Ala Thr Lys Asp Val Thr Val Thr
      1475                      1480                      1485
    Leu Gly Asp Ala Thr Leu Arg Tyr Gly Val Asp Tyr Leu Leu Asp Tyr
      1490                      1495                      1500
15   Ala Gly Asn Thr Ala Val Gly Lys Ala Thr Val Thr Val Arg Gly Ile
    1505                      1510                      1515                      1520
    Asp Lys Tyr Ser Gly Thr Val Ala Lys Thr Phe Thr Ile Glu Leu Lys
      1525                      1530                      1535
    Asn Ala Pro Ala Pro Glu Pro Thr Leu Thr Ser Val Ser Val Lys Thr
      1540                      1545                      1550
20   Lys Pro Ser Lys Leu Thr Tyr Val Val Gly Asp Ala Phe Asp Pro Ala
      1555                      1560                      1565
    Gly Leu Val Leu Gln His Asp Arg Gln Ala Asp Arg Pro Pro Gln Pro
      1570                      1575                      1580
    Leu Val Gly Glu Gln Ala Asp Glu Arg Gly Leu Thr Cys Gly Thr Arg
    1585                      1590                      1595                      1600
25   Cys Asp Arg Val Glu Gln Leu Arg Lys His Glu Asn Arg Glu Ala His
      1605                      1610                      1615
    Arg Thr Gly Leu Asp His Leu Glu Phe Val Gly Ala Ala Asp Gly Ala
      1620                      1625                      1630
    Val Gly Glu Gln Ala Thr Phe Lys Val His Val His Ala Asp Gln Gly
      1635                      1640                      1645
30   Asp Gly Arg His Asp Asp Ala Asp Glu Arg Asp Ile Asp Pro His Val
      1650                      1655                      1660
    Pro Val Asp His Ala Val Gly Glu Leu Ala Arg Ala Ala Cys His His
    1665                      1670                      1675                      1680
35   Val Ile Gly Leu Arg Val Asp Thr His Arg Leu Lys Ala Ser Gly Phe
      1685                      1690                      1695
    Gln Ile Pro Ala Asp Asp Met Ala Glu Ile Asp Arg Ile Thr Gly Phe
      1700                      1705                      1710
    His Arg Phe Glu Arg His Val Gly
      1715                      1720
40
    <210> 23
    <211> 30
    <212> PRT
    <213> Bifidobacterium bifidum
45
    <220>
    <221> SOURCE
    <222> 1..30
    <223> /mol_type="protein"
           /note="Signal sequence of extracellular lactase from Bifidobacter
50           ium bifidum DSM20215"
           /organism="Bifidobacterium bifidum"

    <400> 23
    Val Arg Ser Lys Lys Leu Trp Ile Ser Leu Leu Phe Ala Leu Ala Leu
    1          5          10          15
55   Ile Phe Thr Met Ala Phe Gly Ser Thr Ser Ser Ala Gln Ala
      20          25          30

```

Claims

1. A method for producing a food product by treating a lactose-containing substrate comprising lactose with a polypeptide having transgalactosylating activity, wherein the polypeptide is selected from the group consisting of:
 - d. a polypeptide comprising an amino acid sequence having at least 80% sequence identity with SEQ ID NO: 1,
 - e. a polypeptide comprising an amino acid sequence having at least 80% sequence identity with SEQ ID NO: 2,
 - f. a polypeptide comprising an amino acid sequence having at least 80% sequence identity with SEQ ID NO: 3.
2. The method according to claim 1, wherein the lactose-containing substrate is a milk-based substrate.
3. The method according to claim 2, wherein the milk-based substrate is yoghurt, cheese or a fermented dairy product.
4. The method according to claim 1 or 2, wherein the food product is a dairy product.
5. The method according to claim 4, wherein the dairy product is ice cream, milk, condensed milk, UHT milk, ESL milk, a fermented dairy product or yogurt.
6. The method according to any one of claims 2 to 5, wherein the milk-based substrate that has been treated with the polypeptide account for at least 50%, at least 70%, at least 80% or at least 90%, (weight/weight), of the dairy product.
7. The method according to any one of claims 2 to 6, wherein the polypeptide-treated milk-based substrate is not dried before being used as an ingredient in the dairy product.
8. The method according to any preceding claim, wherein the polypeptide sequence identity is at least 85%, 90%, 95%, 96%, 97%, 98%, or 99%.
9. The method according to any preceding claim, wherein the length of the polypeptide is 500 to 1300 amino acid residues.
10. The method according to claim 9 wherein the polypeptide consists of at most 975, 970, 950, 940, 930, 920, 910, 900, 895 or 890 amino acid residues.
11. The method according to any preceding claim, wherein the polypeptide consists of the amino acid sequence of SEQ ID NO: 1, SEQ ID NO:2 or SEQ ID NO: 3.
12. The method according to any preceding claim, wherein the polypeptide has a ratio of transgalactosylation activity above 100%, preferably above 150%, 175% or 200%, calculated as ratio between Abs420 with acceptor present divided by Abs420 without acceptor present times 100.
13. The method according to any one of claims 4 to 12, wherein the dairy product is a dairy product which is not enriched by addition of pre-produced galacto-oligosaccharides.
14. The method according to any one of claims 2 to 13, wherein the polypeptide-treated milk-based substrate is not dried before being used as an ingredient in the dairy product.
15. Use of a polypeptide having transgalactosylating activity as defined in claim 1 for producing galacto-oligosaccharides to be part of a product selected from the group consisting of yoghurt, cheese, fermented dairy products, dietary supplements and probiotic comestible products.

Figure 1




Figure 2




Figure 3




Figure 4





Figure 5





EUROPEAN SEARCH REPORT

Application Number
EP 18 20 9685

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT				
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)	
X	WO 01/90317 A2 (ARLA FOODS AMBA [DK]; JOERGENSEN FLEMMING [DK]; HANSEN OLEN CAI [DK];) 29 November 2001 (2001-11-29)	1-10, 12-15	INV. C12N9/38	
Y	* whole document esp. seq id no 2; pages 4-5, 7-8; Figure 1; and claims 11,18,26-27 *	1-15		
X,D	WO 2009/071539 A1 (NOVOZYMES AS [DK]; CHR HANSEN AS [DK]; HENRIKSEN HANNE VANG [DK]; ERNS) 11 June 2009 (2009-06-11)	1-10, 12-15		
Y	* whole document esp. seq id no 2, 3; pages 2,16,21; claims 1,13,14 *	1-15		
X	WO 2012/010597 A1 (ARLA FOODS AMBA [DK]; BERTELSEN HANS [DK]; LANGBORG WEJSE PETER [DK]) 26 January 2012 (2012-01-26)	1-10, 12-15		
Y	* whole document esp. seq id no 2; pages 2-3,30 *	1-15	TECHNICAL FIELDS SEARCHED (IPC) C12N	
Y	O. HANSEN ET AL: "High-efficiency synthesis of oligosaccharides with a truncated ?-galactosidase from Bifidobacterium bifidum", APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, vol. 57, no. 5-6, 1 December 2001 (2001-12-01), pages 647-652, XP055542920, DE ISSN: 0175-7598, DOI: 10.1007/s00253-001-0845-z * whole document esp. page 647 *	1-15		
The present search report has been drawn up for all claims				
Place of search Munich		Date of completion of the search 17 January 2019		Examiner Brück, Marianne
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document				

EPO FORM 1503 03.02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 18 20 9685

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

17-01-2019

10

15

20

25

30

35

40

45

50

55

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0190317 A2	29-11-2001	AT 352611 T	15-02-2007
		AU 6206801 A	03-12-2001
		AU 2001262068 B2	06-07-2006
		DE 60126269 T2	21-06-2007
		DK 1283876 T3	19-02-2007
		EP 1283876 A2	19-02-2003
		JP 5042431 B2	03-10-2012
		JP 2003534007 A	18-11-2003
		KR 20030016275 A	26-02-2003
		KR 20080045766 A	23-05-2008
		NZ 522623 A	31-10-2003
		US 2002086358 A1	04-07-2002
		US 2003162280 A1	28-08-2003
		WO 0190317 A2	29-11-2001
WO 2009071539 A1	11-06-2009	CN 101932248 A	29-12-2010
		CN 104322666 A	04-02-2015
		CN 104472683 A	01-04-2015
		DK 2234501 T3	26-09-2016
		EP 2234501 A1	06-10-2010
		EP 3138408 A1	08-03-2017
		ES 2591359 T3	28-11-2016
		PL 2234501 T3	31-08-2017
		US 2010285175 A1	11-11-2010
		US 2017156357 A1	08-06-2017
		US 2018295846 A1	18-10-2018
		WO 2009071539 A1	11-06-2009
WO 2012010597 A1	26-01-2012	AR 082261 A1	21-11-2012
		AU 2011281663 A1	07-03-2013
		BR 112013001109 A2	24-05-2016
		CA 2805997 A1	26-01-2012
		CN 103080329 A	01-05-2013
		DK 2596113 T3	07-01-2019
		EA 201300150 A1	30-08-2013
		EP 2596113 A1	29-05-2013
		JP 2013530719 A	01-08-2013
		KR 20130041953 A	25-04-2013
		MX 348325 B	06-06-2017
		NZ 607149 A	24-12-2014
		US 2013189746 A1	25-07-2013
		US 2016369313 A1	22-12-2016
		WO 2012010597 A1	26-01-2012

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 0190317 A [0006]
- WO 2012010597 A [0006]
- WO 2009071539 A [0007] [0235]
- US 7371552 B [0178]
- US 7166453 B [0178]
- US 6890572 B [0178]
- US 6667065 B [0178]
- US 20070141693 A [0178]
- US 20070072270 A [0178]
- US 20070020731 A [0178]
- US 20070020727 A [0178]
- US 20060073583 A [0178]
- US 20060019347 A [0178]
- US 20060018997 A [0178]
- US 20060008890 A [0178]
- US 20060008888 A [0178]
- US 20050137111 A [0178]
- US 5874276 A [0180]
- US 6022725 A [0193]
- US 6268328 B [0193]
- EP 244234 A [0193]
- EP 215594 A [0193]

Non-patent literature cited in the description

- JORGENSEN et al. *Appl. Microbiol. Biotechnol.*, 2001, vol. 57, 647-652 [0006]
- R.D. FINN ; J. MISTRY ; J. TATE ; P. COGGILL ; A. HEGER ; J.E. POLLINGTON ; O.L. GAVIN ; P. GUNESSEKARAN ; G. CERIC ; K. FORSLUND. The Pfam protein families database. *Nucleic Acids Research*, 2010, vol. 38, D211-222 [0059] [0136]
- ALTSCHUL et al. *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0089]
- NEEDLEMAN, SAUL B. ; WUNSCH, CHRISTIAN D. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *Journal of Molecular Biology*, 1970, vol. 48 (3), 443-53 [0090]
- WIERENGA. *FEBS Letters*, 2001, vol. 492 (3), 193-8 [0120]
- JENKINS. *FEBS Letters*, 1995, vol. 362 (3), 281-5 [0120]
- HENRISSAT. *Curr Opin Struct Biol*, 1997, vol. 7 (5), 637-44 [0120]
- J. SAMBROOK ; E.F. FRITSCH ; T. MANIATIS. *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor, 1989 [0163]
- BOLTON ; MCCARTHY. *Proceedings of the National Academy of Sciences USA*, 1962, vol. 48, 1390 [0171]
- SAMBROOK et al. *MOLECULAR CLONING: A LABORATORY MANUAL*. Cold Spring Harbor Laboratory Press, 2001 [0180]
- BENNETT et al. *MORE GENE MANIPULATIONS IN FUNGI*. Academic Press, 1991, 396-428 [0180]
- KELLEY et al. *EMBO J.*, 1985, vol. 4, 475-479 [0183]
- PENTTILA et al. *Gene*, 1987, vol. 61, 155-164 [0183]
- CAMPBELL et al. *Curr. Genet.*, 1989, vol. 16, 53-56 [0193]
- HARKKI et al. *Enzyme Microb. Technol.*, 1991, vol. 13, 227-233 [0193]
- HARKKI et al. *BioTechnol.*, 1989, vol. 7, 596-603 [0193]