



(11)

EP 3 587 576 A9

(12)

CORRECTED EUROPEAN PATENT APPLICATION

published in accordance with Art. 153(4) EPC

(15) Correction information:

Corrected version no 1 (W1 A1)

Corrections, see

Description Paragraph(s) 140

(51) Int Cl.:

C12N 15/113 ^(2010.01) **A61K 31/7115** ^(2006.01)

A61K 31/712 ^(2006.01) **A61P 43/00** ^(2006.01)

(48) Corrigendum issued on:

29.07.2020 Bulletin 2020/31

(86) International application number:

PCT/JP2018/006061

(43) Date of publication:

01.01.2020 Bulletin 2020/01

(87) International publication number:

WO 2018/155450 (30.08.2018 Gazette 2018/35)

(21) Application number: **18757296.1**

(22) Date of filing: **20.02.2018**

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

• **YOSHIDA, Tokuyuki**

Suita-shi

Osaka 565-0871 (JP)

• **MORIHIRO, Kunihiko**

Suita-shi

Osaka 565-0871 (JP)

• **KASAHARA, Yuya**

Suita-shi

Osaka 565-0871 (JP)

• **MIKAMI, Atsushi**

Suita-shi

Osaka 565-0871 (JP)

(30) Priority: **21.02.2017 JP 2017030489**

(71) Applicant: **Osaka University**

Suita-shi, Osaka 565-0871 (JP)

(74) Representative: **J A Kemp LLP**

14 South Square

Gray's Inn

London WC1R 5JJ (GB)

(72) Inventors:

• **OBIKA, Satoshi**

Suita-shi

Osaka 565-0871 (JP)

• **WAKI, Reiko**

Suita-shi

Osaka 565-0871 (JP)

• **INOUE, Takao**

Suita-shi

Osaka 565-0871 (JP)

(54) **ANTISENSE OLIGONUCLEIC ACID**

(57) The present invention aims to provide an anti-sense oligonucleic acid with reduced hepatotoxicity. The antisense oligonucleic acid according to the present invention is characterized in that it has a base length of not less than 7nt and not more than 30nt, wherein nucleic acid residues of not less than 1nt and not more than 5nt

respectively from the both terminals are 2',4'-bridged nucleic acids, 2',4'-non-bridged nucleic acid residue(s) is(are) present between the above-mentioned both terminals, and one or more bases in the nucleic acid residue(s) of the above-mentioned 2',4'-non-bridged nucleic acid residue(s) is/are modified.

EP 3 587 576 A9

Description

[Technical Field]

5 **[0001]** The present invention relates to an antisense oligonucleic acid with low hepatotoxicity.

[Background Art]

10 **[0002]** Nucleic acid medicines can suppress expression of specific genes in cells unlike conventional pharmaceutical products. They are expected to lead to the creation of innovative pharmaceutical products for diseases that were previously difficult to treat. As a technique relating to nucleic acid medicine, for example, an antisense method has been developed in which an antisense oligonucleic acid complementary to a part of mRNA involved in the disease is introduced from the outside to cause formation of a double strand, which in turn inhibits translation process of the pathogenic mRNA and treats or prevents the disease.

15 **[0003]** However, antisense oligonucleic acid has a defect in that it is easily degraded by nucleases in vivo. In addition, the binding affinity and specificity for target mRNA need to be improved further. Thus, the study group of the present inventors developed a nucleic acid molecule bridged at the 2'- and 4'-positions in an attempt to improve nuclease resistance and binding affinity and specificity to target mRNA by stabilization of the steric structure (patent documents 1 - 5) and, using such bridged nucleic acid, developed an oligonucleotide useful as a therapeutic agent for dyslipidemia which is superior in binding affinity to PCSK9 gene and the like (patent document 6).

20 **[0004]** However, as nucleic acid medicament discovery research in anticipation of human clinical trials progresses, new issues to be solved have become apparent. The most important one is "avoidance of potential toxicity of artificial oligonucleotides".

25 **[0005]** For example, it is known that antisense oligonucleic acid intravenously administered to rats and mice is delivered to the liver in about 12 hours. Some antisense oligonucleic acids exhibit severe hepatotoxicity. While the mechanism is not necessarily clear, suppression of expression of non-target mRNA, activation of natural immunity, promotion of expression of unidentified protein gene and the like are considered. As a means to predict hepatotoxicity of antisense oligonucleic acids, research is being conducted to identify the base sequence common to antisense oligonucleic acids exhibiting hepatotoxicity (non-patent documents 1, 2).

30

[Document List]

[Patent documents]

35 **[0006]**

patent document 1: WO 2011/052436

patent document 2: WO 2014/046212

patent document 3: WO 2014/112463

40 patent document 4: WO 2015/125783

patent document 5: WO 2016/017422

patent document 6: WO 2013/118776

[non-patent document]

45

[0007]

non-patent document 1: Peter H. Hagedorn et al., Nucleic Acid Therapeutics, 23(5), pp.302-310 (2013)

non-patent document 2: Andrew D. Burdick et al., Nucleic Acids Research, 2014, pp.1-10

50

[SUMMARY OF THE INVENTION]

[Problems to be Solved by the Invention]

55 **[0008]** As mentioned above, studies are being conducted to identify the base sequence considered to be involved in the hepatotoxicity of antisense oligonucleic acids. The base sequences considered to be involved in hepatotoxicity in the above-mentioned non-patent documents 1, 2 are only nt2 and nt3, and it is sometimes difficult to design an antisense oligonucleic acid free of such short base sequences. For example, mRNA has a partial self structure for stabilization

such as a loop structure, a double strand and the like, and there is a limitation that antisense oligonucleic acid should be designed to hybridize to a single strand of the target mRNA. Moreover, a base sequence that hybridizes to a single strand of the target mRNA and is free of the above-mentioned base sequence does not necessarily show an excellent activity without hepatotoxicity.

[0009] Therefore, the present invention aims to provide an antisense oligonucleic acid with reduced hepatotoxicity.

[Means of Solving the Problems]

[0010] The present inventors have conducted intensive studies in an attempt to solve the aforementioned problems. As a result, they have found that particularly, hepatotoxicity of an antisense oligonucleic acid can be reduced while maintaining its activity by modifying a base of a nucleic acid residue in a particular region of the antisense oligonucleic acid, which resulted in the completion of the present invention.

[0011] The present invention is shown below.

[1] An antisense oligonucleic acid having a base length of not less than 7nt and not more than 30nt, wherein nucleic acid residues of not less than 1nt and not more than 5nt respectively from the both terminals are 2',4'-bridged nucleic acids,


2', 4' -non-bridged nucleic acid residue(s) is (are) present between the above-mentioned both terminals, and one or more bases in the nucleic acid residue(s) of the above-mentioned 2', 4' -non-bridged nucleic acid residue(s) is/are modified.

[2] The antisense oligonucleic acid of the above-mentioned [1] wherein the above-mentioned 2', 4' -non-bridged nucleic acid residue with the modified base is contained in a sequence of TGC or TCC.

[3] The antisense oligonucleic acid of the above-mentioned [1] wherein the above-mentioned modified base is 5-hydroxycytosine, 4-acetylcytosine, 3-C₁₋₆ alkylcytosine, 5-C₁₋₆ alkylcytosine, 2-thiocytosine, 2-thiopyrimidine, dihydrothymine, pseudo thymine, 2-thiouridine, 5-(carboxyhydroxymethyl)uridine, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluridine, dihydrouridine, pseudo uridine, 5-methylaminomethyluridine, 5-methylaminomethyl-2-thiouridine, 5-methoxycarbonylmethyl-2-thiouridine, 5-methoxycarbonylmethyluridine, 5-methoxyuridine, 8-aminoguanine, 8-halogenoguanine, 7-deaza-7-(2-phenylethynyl)guanine, 7-deaza-7-(2-pyridylethynyl)guanine, 7-deaza-7-[2-(C₁₋₇ alkanoyloxy-C₁₋₆ alkyl)ethynyl]guanine, 7-deaza-7-[2-(hydroxy-C₁₋₆ alkyl) ethynyl] guanine, 1-C₁₋₆ alkylguanine, 2, 2-di (C₁₋₆ alkyl)guanine, 2-C₁₋₆ alkylguanine, 7-deaza-7-C₁₋₆ alkylguanine, 7-deaza-7-C₂₋₆ alkenylguanine, 7-deaza-7-C₂₋₆ alkynylguanine, 7-deaza-7-halogenoguanine, inosine, 1-C₁₋₆ alkylinosine, queuosine, β,D-galactosylqueuosine, β,D-mannosylqueuosine, N⁶-C₂₋₆ alkenyladenine, 1-C₁₋₆ alkyladenine, 2-C₁₋₆ alkyladenine, N⁶-C₁₋₆ alkyladenine or 2-C₁₋₆ alkylthio-N⁶-C₂₋₆ alkenyladenine.

[4] The antisense oligonucleic acid of any of the above-mentioned [1] to [3] wherein the above-mentioned 2',4'-non-bridged nucleic acid residue is DNA.


[5] The antisense oligonucleic acid of any of the above-mentioned [1] to [4] wherein the above-mentioned 2',4'-bridged nucleic acid residue has a structure of any of the following formulas (I) to (III):




wherein

X¹ is O, S, a >N(R³) group, a -C(=O)-O- group or a -C(=O)-N(R⁴)- group (R³ and R⁴ are each independently H or a C₁₋₆ alkyl group),


X² is a guanidino group represented by any of the following formulas (IV) to (VII):




(IV)



(V)



(VI)



(VII)

wherein $R^5 - R^{18}$ are each independently H, a C_{1-6} alkyl group, a C_{3-10} cycloalkyl group, an amino-protecting group or a 2-cyanoethoxycarbonyl group,

X^3 is O, S, a $>N(R^{19})$ group, a $-C(=O)-O-$ group or a $-C(=O)-N(R^{20})-$ group (R^{19} and R^{20} are each independently H or a C_{1-6} alkyl group),

$Y^1 - Y^3$ are each independently O^- or S^- ,

Base is a nucleic acid base group,

R^1 and R^2 are each independently H, a C_{1-6} alkyl group or R^1 and R^2 may be taken together to form a C_{1-4} alkylene group, and

n is an integer of not less than 0 and not more than 2.

[6] The antisense oligonucleic acid of the above-mentioned [5] wherein the above-mentioned 2',4'-bridged nucleic acid residue has a structure of the above-mentioned formula (I), X^1 is O and n is 1.

[7] The antisense oligonucleic acid of any of the above-mentioned [1] to [6] showing reduced hepatotoxicity compared to that before introduction of the above-mentioned modification into the base of the nucleic acid residue.

[0012] In this disclosure, the " C_{1-6} alkyl group" is a linear or branched chain monovalent saturated aliphatic hydrocarbon group with a carbon number of not less than one and not more than 6. For example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, s-butyl, t-butyl, n-pentyl, n-hexyl and the like can be mentioned. Preferred is a C_{1-4} alkyl group, more preferred is a C_{1-2} alkyl group, most preferred is methyl.

[0013] The " C_{2-6} alkenyl group" is a linear or branched chain monovalent unsaturated aliphatic hydrocarbon group with a carbon number of not less than 2 and not more than 6 and having at least one carbon-carbon double bond. Examples include ethenyl(vinyl), 1-propenyl, 2-propenyl(allyl), isopropenyl, 2-butenyl, 3-butenyl, isobutenyl, pentenyl, isopentenyl, hexenyl and the like.

[0014] The " C_{2-6} alkynyl group" is a linear or branched chain monovalent unsaturated aliphatic hydrocarbon group with a carbon number of not less than 2 and not more than 6 and having at least one carbon-carbon triple bond. Examples include ethynyl, 1-propynyl, 2-propynyl(propargyl), 2-butenyl, 3-butenyl, pentynyl, hexynyl and the like. Preferred is a C_{2-4} alkynyl group, more preferred is a C_{2-3} alkynyl group.

[0015] The " C_{6-12} aryl group" is a monovalent aromatic hydrocarbon group with a carbon number of not less than 6 and not more than 12. Examples include phenyl, naphthyl, indenyl, biphenyl and the like.

[0016] Examples of the "halogeno group" include fluoro group, chloro group, bromo group and iodo group, with preference given to chloro group and bromo group.

[0017] The " C_{3-10} cycloalkyl group" is a cyclic monovalent saturated aliphatic hydrocarbon group with a carbon number of not less than 3 and not more than 10. It is, for example, cyclopropyl, methylcyclopropyl, dimethylcyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, adamantyl or the like. Preferred is a C_{3-6} cycloalkyl group.

[0018] Examples of the "amino-protecting group" include alkoxycarbonyl protecting groups such as t-butoxycarbonyl group, methoxycarbonyl group, ethoxycarbonyl group and the like; arylmethoxycarbonyl protecting groups such as benzyloxycarbonyl group, p-methoxybenzyloxycarbonyl group, p-nitrobenzyloxycarbonyl group, o-nitrobenzyloxycarbonyl group and the like; arylmethyl protecting groups such as benzyl group, 4-methoxybenzyl group, triphenylmethyl group and the like; alkanoyl protecting groups such as formyl group, acetyl group and the like; aroyl protecting groups such as benzoyl group and the like; arylsulfonyl protecting groups such as 2,4-di nitrobenzenesulfonyl group, o-nitrobenzenesulfonyl group and the like, and the like.

[0019] The " C_{1-6} alkylene group" is a linear or branched chain divalent saturated aliphatic hydrocarbon group with a carbon number of not less than 1 and not more than 6. It is, for example, methylene, ethylene, methylmethylene, n-propylene, methylethylene, n-butylene, methylpropylene, dimethylethylene, n-pentylene, n-hexylene or the like. It is preferably a C_{1-4} alkylene group, more preferably a C_{1-2} alkylene group, further more preferably methylene.

[0020] The " C_{1-7} alkanoyl group" is an atomic group of aliphatic carboxylic acid with a carbon number of 1 to 7 less OH. It is, for example, formyl, acetyl, ethylcarbonyl, n-propylcarbonyl, isopropylcarbonyl, n-butylcarbonyl, isobutylcarbonyl, t-butylcarbonyl, n-pentylcarbonyl, n-hexylcarbonyl or the like, preferably a C_{1-4} alkanoyl group, more preferably

acetyl.

[Effect of the Invention]

5 **[0021]** The hepatotoxicity of the antisense oligonucleic acid of the present invention is remarkably reduced. Furthermore, the antisense oligonucleic acid of the present invention has high binding affinity to target mRNA, is resistant to nuclease attack in vivo and is stable. Therefore, the antisense oligonucleic acid of the present invention is industrially extremely useful since it may be a highly practical nucleic acid medicament.

10 [Brief Description of the Drawings]

[0022]

15 Fig. 1 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

Fig. 2 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

Fig. 3 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and GR gene expression-suppressing activity evaluated in the Example described later.

20 Fig. 4 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and GR gene expression-suppressing activity evaluated in the Example described later.

Fig. 5 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

25 Fig. 6 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

Fig. 7 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and Rps6kb2 gene expression-suppressing activity evaluated in the Example described later.

Fig. 8 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

30 Fig. 9 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

Fig. 10 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and GR gene expression-suppressing activity evaluated in the Example described later.

35 Fig. 11 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and GR gene expression-suppressing activity evaluated in the Example described later.

Fig. 12 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

Fig. 13 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

40 Fig. 14 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

Fig. 15 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

45 Fig. 16 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

Fig. 17 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and GR gene expression-suppressing activity evaluated in the Example described later.

Fig. 18 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

50 Fig. 19 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

Fig. 20 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and Rps6kb2 gene expression-suppressing activity evaluated in the Example described later.

55 Fig. 21 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

Fig. 22 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and Rps6kb2 gene expression-suppressing activity evaluated in the Example described later.

Fig. 23 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and GR gene expression-

suppressing activity evaluated in the Example described later.

Fig. 24 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

Fig. 25 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

Fig. 26 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

Fig. 27 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and GR gene expression-suppressing activity evaluated in the Example described later.

Fig. 28 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid and PCSK9 gene expression-suppressing activity evaluated in the Example described later.

Fig. 29 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

Fig. 30 is a graph showing the results of hepatotoxicity of antisense oligonucleic acid evaluated in the Example described later.

[Description of Embodiments]

[0023] The antisense oligonucleic acid of the present invention is characterized in that it has a base length of not less than 7nt, not more than 30nt; nucleic acid residues of not less than 1nt and not more than 5nt respectively from the both terminals are 2', 4'-bridged nucleic acids; 2', 4'-non-bridged nucleic acid residue(s) is(are) present between the above-mentioned both terminals; and one or more bases in the nucleic acid residue(s) of the above-mentioned 2', 4'-non-bridged nucleic acid residue(s) is/are modified.

[0024] The base sequence of the antisense oligonucleic acid of the present invention is not particularly limited as long as it can hybridize to the target mRNA with high binding affinity and one or more bases of nucleic acid residue in a particular region is/are modified. The original base sequence before modification of the base can be determined by a conventional method according to the target mRNA.

[0025] For example, when the translation initiation site of target mRNA is selected as a binding target site, an antisense oligonucleic acid with high translation inhibitory activity may be designed. mRNA sometimes forms a hairpin loop structure or a double strand structure in the molecule, and a single strand structure site is kinetically and thermodynamically preferable as a binding target site. Thus, the secondary structure of mRNA is predicted by calculation, and a site predicted to be a single-stranded structure can be selected as a binding target site. It is also possible to identify a sequence most likely to form a double strand with the target mRNA by random screening using RNaseH, which selectively cleaves the site where the double strand is formed.

[0026] The nucleic acid base in the antisense oligonucleic acid of the present invention may be, besides a modified base, general adenine, guanine, uracil, thymine or cytosine. Since methylcytosine can be easily synthesized from thymine, methylcytosine may also be used instead of cytosine.

[0027] The antisense oligonucleic acid of the present invention may also be a thioated oligonucleic acid. In general oligonucleic acids, riboses are linked by a phosphodiester bond. The resistance of oligonucleic acid to exonuclease and endonuclease is improved by binding riboses with a phosphorothioate bond. The oligonucleic acid may be complete thioated oligonucleic acid in which all phosphoric acid groups are thioated or chimeric thioated oligonucleic acid in which a part of the phosphoric acid group is thioated.

[0028] The base length of the antisense oligonucleic acid of the present invention is preferably not less than 7nt and not more than 30nt. When the base length is not less than 7nt, the binding affinity and specificity to the target mRNA can be sufficiently ensured. On the other hand, when the base length is not more than 30nt, the antigenicity is sufficiently suppressed and synthesis is facilitated. The base length is preferably not less than 10nt, more preferably not less than 12nt, and preferably not more than 25nt and preferably not more than 20nt. The number of oligonucleic acids with a base length of 17nt is $4^{17}=1.7 \times 10^{10}$ and exceeds $2 \times 3 \times 10^9$ which is the total number of bases in the human genome. Accordingly, it is also possible to set the base length to not less than 17nt to improve specificity.

[0029] In the antisense oligonucleic acid of the present invention, nucleic acid residues each independently not less than 1nt and not more than 5nt from the both terminals are 2', 4'-bridged nucleic acids. The nucleic acids at at least both terminal sites are 2', 4'-bridged, which renders attacks by various nucleases difficult in vivo, and the antisense oligonucleic acid can be present in the living body for a long time after administration to the living body. In addition, since the structure is stabilized by 2', 4'-bridging, a double strand is easily formed with the target mRNA.

[0030] The 2', 4'-bridged structures in two or more 2', 4'-bridged nucleic acid residues may be identical to or different from one another. The 2', 4'-bridged structure is not particularly limited as long as it stabilizes the structure of the ribose moiety and, for example, the structures of the above-mentioned formulas (I) to (III) can be mentioned.

[0031] The nucleic acid base in the 2', 4'-bridged nucleic acid residue may be a general adenine, guanine, uracil,

thymine or cytosine and may be modified. Since methylcytosine can be easily synthesized from thymine, methylcytosine may also be used instead of cytosine, particularly at the both terminal sites. The number of modified bases at the both terminal sites varies depending on the 2',4'-bridged nucleic acid residues at the both terminal sites and is preferably not more than 5 or not more than 4, more preferably not more than 3, further more preferably 1 or 2. The modified bases in

the both terminal sites may be, for example, those similar to the modified base in the 2',4'-non-bridged nucleic acid residue. **[0032]** The antisense oligonucleic acid according to the present invention has one or more 2',4'-non-bridged nucleic acid residues between both terminals composed of 2',4'-bridged nucleic acid residues. That is, the antisense oligonucleic acid according to the present invention is composed of 2',4'-non-bridged nucleic acid residues except for the both terminal sites thereof. Hereinafter, in the present disclosure, the parts other than the above-mentioned both terminal sites of the antisense oligonucleic acid according to the present invention are sometimes referred to as a "middle part". The 2',4'-non-bridged nucleic acid residue in the middle part may be RNA, DNA, a nucleic acid derivative, or a combination of two or more of these as long as bridging is absent between the 2' position and the 4' position. Examples of the nucleic acid derivative include 2'-C₁₋₆ alkylcarbonyloxynucleic acids such as 2'-halogenonucleic acid, 2'-acetoxynucleic acid and the like, 2'-C₁₋₆ alkyloxynucleic acids such as 2'-methoxynucleic acid and the like, 2'-triC₁₋₆ alkylsilyloxynucleic acids such as 2'-trimethylsilyloxynucleic acid and the like, and the like. It is preferable that all the nucleic acids in the above-mentioned middle part be DNAs. When all the nucleic acids in the middle part are DNAs, an RNA-DNA double strand is formed to the target mRNA, the double strand becomes a substrate for RNaseH, and the target mRNA is cleaved. From the aspect of the substrate of RNaseH, the length of the above-mentioned middle part is preferably not less than 4, more preferably not less than 5, and still more preferably not less than 6. On the other hand, the length of the above-mentioned middle part is preferably not more than 10 nt from the aspect of antigenicity.

[0033] In the antisense oligonucleic acid according to the present invention, the bases of one or more nucleic acid residues in the above-mentioned middle part are modified. Since the original base sequence of the antisense oligonucleic acid of the present invention is complementary to the base sequence of the binding target site of the target mRNA, and has strong binding affinity to the binding target site, the binding affinity is considered to be maintained even if the base of the middle part that forms a double strand with the target mRNA is modified. On the other hand, the hepatotoxicity of the antisense oligonucleic acid is caused by the interaction with compounds other than the target mRNA and the binding affinity to such compounds is highly likely lower than that to the target mRNA. In the present invention, the interaction with such compounds is considered to be reduced to the extent that hepatotoxicity is reduced, by modifying the base of one or more nucleic acid residues in the above-mentioned middle part.

[0034] The number of nucleic acid residues whose bases are modified in the middle part is preferably not more than 5, depending on the number of nucleic acid residues constituting the middle part. It is considered that the lower the number, the more certainly the binding affinity to the target mRNA is maintained. The number is preferably not more than 4 or not more than 3, further more preferably 2 or 1 and particularly preferably 1.

[0035] As modification of the base, for example, substitution of a group, addition of a functional group, removal of a functional group are considered. For modification of a base, two or more modifications may be combined as long as it is within one base. The substitution of a group may be not only substitution of a substituent bound to heterocycle of the base, but also substitution of a group forming the heterocycle of the base. For example, it is also possible to substitute =N-forming heterocycle of the base with =C<. For example, =N- at the 7-position of adenine and guanine may become =N< by the addition of a functional group.

[0036] When cytosine is to be modified, a preferable substituent introduction position is, for example, the 5-position. As the substituent at the 5-position, C₁₋₆ alkyl group, C₂₋₆ alkenyl group, C₂₋₆ alkynyl group, C₆₋₁₂ aryl group, hydroxyl group, amino group, C₁₋₆ alkylamino group, di (C₁₋₆ alkyl) amino group, a carboxy group, sulfonic acid group and the like can be mentioned. In addition, one or two from C₁₋₆ alkyl group, C₂₋₆ alkenyl group and C₂₋₆ alkynyl group may be introduced into the 4-position amino group.


[0037] Preferable modification of cytosine, thymine and uracil includes, for example, substitution of the 2-position carbonyl group with a thiocarbonyl group. In addition, a C₁₋₆ alkyl group may be introduced into the 3-position amino group of thymine or uracil.

[0038] When guanine or adenine is to be modified, a preferable substituent introduction position is, for example, the 8-position. Examples of the substituent at the 8-position include halogeno groups in addition to hydrophilic substituents such as amino group, C₁₋₆ alkylamino group, di (C₁₋₆ alkyl) amino group and the like. Alternatively, the 7-position =N- may be converted to =C< and a phenylethynyl group may be introduced.

[0039] Examples of more specific modified base include cytosine derivatives such as 5-hydroxycytosine, 4-acetylcytosine, 3-C₁₋₆ alkylcytosine, 5-C₁₋₆ alkylcytosine, 2-thiocytosine and the like; thymine derivatives such as 2-thiothymine, dihydrothymine, pseudo thymine and the like; uridine derivatives such as 2-thiouridine, 5-(carboxyhydroxymethyl)uridine, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluridine, dihydrouridine, pseudo uridine, 5-methylaminomethyluridine, 5-methylaminomethyl-2-thiouridine, 5-methoxycarbonylmethyl-2-thiouridine, 5-methoxycarbonylmethyluridine, 5-methoxyuridine and the like; guanine derivatives such as 8-aminoguanine, 8-halogenoguanine, 7-deaza-7-(2-phenylethynyl)guanine, 7-deaza-7-[2-(4-pyridyl)ethynyl]guanine, 7-deaza-7-[2-(2-pyridyl)ethynyl]gua-

nine, 7-deaza-7-[2-(3-pyridyl)ethynyl]guanine, 7-deaza-7-[2-(C₁₋₇ alkanoyloxy-C₁₋₆ alkyl) ethynyl] guanine, 7-deaza-7-[2-(hydroxy-C₁₋₆ alkyl) ethynyl] guanine, 1-C₁₋₆ alkylguanine, 2,2-di(C₁₋₆ alkyl)guanine, 2-C₁₋₆ alkylguanine, 7-deaza-7-C₁₋₆ alkylguanine, 7-deaza-7-C₂₋₆ alkenylguanine, 7-deaza-7-C₂₋₆ alkynylguanine, 7-deaza-7-halogenoguanine, inosine, 1-C₁₋₆ alkylinosine, queuosine, β ,D-galactosylqueuosine, β ,D-mannosylqueuosine and the like; and adenine derivatives such as N⁶-C₂₋₆ alkenyladenine, 1-C₁₋₆ alkyladenine, 2-C₁₋₆ alkyladenine, N⁶-C₁₋₆ alkyladenine, 2-C₁₋₆ alkylthio-N⁶-C₂₋₆ alkenyladenine and the like.

[0040] The present inventors developed the following novel 7-substituted guanosine derivative (VIII) as a compound having a modified base:



wherein

R³¹ is H or a hydroxyl-protecting group selected from a silyl protecting group, a trityl protecting group, a carbamate protecting group and a benzylether protecting group;

R³² and R³³ are each independently a phenyl group optionally substituted by a halogeno group, a nitro group or a C₁₋₆ alkyl group, or a C₁₋₆ alkyl group optionally substituted by a cyano group;

R³⁴ is an amino-protecting group; and

R³⁵ is a halogeno group or R³⁶-C $\equiv$ C-(R³⁶ is a C₆₋₁₂ aryl group, a heterocyclic aromatic ring group, a hydroxy-C₁₋₆ alkyl group or a C₁₋₇ alkanoyloxy-C₁₋₆ alkyl group) .

[0041] Examples of the "silyl protecting group" include trimethylsilyl, triethylsilyl, triisopropylsilyl, dimethylisopropylsilyl, diethylisopropylsilyl, dimethylhexylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, tribenzylsilyl, tri-p-xylylsilyl, triphenylsilyl, diphenylmethylsilyl, t-butylmethoxyphenylsilyl and the like, examples of the "trityl protecting group" include triphenylmethyl, α -naphthyl diphenylmethyl, p-methoxyphenyl diphenylmethyl, di(p-methoxyphenyl)phenylmethyl, tri(p-methoxyphenyl)methyl, 4-(4'-bromophenoxy)phenyl diphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyl oxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 3-(imidazol-1-ylmethyl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl and the like, examples of the "carbamate protecting group" include t-butoxycarbonyl, fluorenylmethoxycarbonyl, benzyloxycarbonyl and the like, examples of the "benzylether protecting group" include benzyl, methoxybenzyl, nitrobenzyl, halobenzyl, phenylbenzyl and the like, and examples of the "alkanoyl protecting group" include formyl, acetyl, halogenated acetyl, pivaloyl, benzoyl and the like. As the "amino-protecting group", those recited above can be mentioned.

[0042] The "heterocyclic aromatic ring group" refers to a 5-membered ring aromatic heterocyclyl group, 6-membered ring aromatic heterocyclyl group or fused ring aromatic heterocyclyl group having at least one hetero atom such as a nitrogen atom, an oxygen atom, a sulfur atom and the like. For example, 5-membered heterocyclic aromatic ring groups such as pyrrolyl, imidazolyl, pyrazolyl, thienyl, furyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazole and the like; 6-membered heterocyclic aromatic ring groups such as pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl and the like; fused ring heterocyclic aromatic ring groups such as indolyl, isoindolyl, quinolyl, isoquinolyl, benzofuranyl, isobenzofuranyl, chromenyl and the like can be mentioned. It is preferably heterocyclic aromatic ring group containing a nitrogen atom, more preferably pyridyl.

[0043] The base to be modified is not particularly limited. For example, since the base length of the antisense oligonucleic acid of the present invention is not less than 7nt and not more than 30nt, the base sequence of the original antisense oligonucleic acid may be determined, the base is modified one by one and a modified base superior in a hepatotoxicity reducing effect may be determined. However, as described in the Examples below, since a favorable hepatotoxicity reducing effect tends to be seen when thymine, guanine or cytosine contained in the sequence of TGC or TCC is modified, modification of a base of the nucleic acid residue contained in these sequences can be one aim.

[0044] In the antisense oligonucleic acid according to the present invention, in particular, the base of one nucleic acid

residue is modified, and thus shows reduced hepatotoxicity compared to the antisense oligonucleic acid before such modification. For example, the serum concentrations after administration of antisense oligonucleic acids of AST (aspartate aminotransferase) and ALT (alanine aminotransferase), which are indices of hepatotoxicity, are preferably not more than 60% or not more than 50%, more preferably not more than 40% or not more than 20%, further more preferably not more than 10%, particularly preferably not more than 5%, of the serum concentration measured under the same conditions except that the antisense oligonucleic acid before the above-mentioned modification is used. In addition, the serum concentration of AST after administration of the antisense oligonucleic acid of the present invention is preferably not more than 100IU/L, more preferably not more than 40IU/L, and the serum concentration of ALT is preferably not more than 100IU/L, more preferably not more than 40IU/L, further more preferably not more than 20IU/L.

[0045] In the present disclosure, the serum concentrations of AST and ALT are determined by, for example, intravenously injecting antisense oligonucleic acid to mice or rats at a predetermined dose of about not less than 10 mg/kg body weight and not more than about 30 mg/kg body weight, collecting blood after lapse of predetermined time of not less than about 70 hours and not more than about 120 hr, and measuring same from a serum obtained by centrifugation.

[0046] The antisense oligonucleic acid according to the present invention can be produced by a conventional method. For example, since the base length of the antisense oligonucleic acid of the present invention is not less than 7nt and not more than 30nt, it can be easily produced using an automatic nucleic acid synthesizer.

[0047] Since the antisense oligonucleic acid according to the present invention has a base sequence complementary to the target mRNA, it can strongly inhibit translation of the target mRNA and can also reduce hepatotoxicity as described above. Therefore, the antisense oligonucleic acid is useful as a medicament for preventing or treating a disease involving a target mRNA.

[0048] The antisense oligonucleic acid of the present invention can be formulated as a parenteral preparation or a liposome preparation by blending with, for example, adjuvants generally used in the technical field of preparation formulation of medicaments such as excipient, binder, preservative, oxidation stabilizer, disintegrant, lubricant, corrigent and the like. In addition, a topical preparation such as liquid, cream, ointment and the like can be formulated by blending with, for example, a pharmaceutical carrier generally used in the pertinent technical field.

[0049] The antisense oligonucleic acids of the present invention can be administered to human and animals other than human. The dose may be appropriately adjusted depending on the patient's age, sex, body weight, condition, the kind of disease, severity, prophylactic or therapeutic use, and the like. For example, a dose per administration and per kg body weight for adult human is not less than 0.01 μ g and not more than 100 g. The dose is preferably not less than 0.1 μ g or not less than 1.0 pg, more preferably not less than 10 μ g or not less than 100 pg, further preferably not less than 1 mg, and preferably not more than 10 g or not more than 1.0 g, more preferably not more than 100 mg or not more than 10 mg, further more preferably not more than 5 mg. The frequency of administration may be appropriately adjusted between once per month and about not more than 3 times per day.

[0050] This application claims the benefit of priority right based on a patent application No. 2017-30489 filed in Japan (filing date: February 21, 2017). The contents of Japanese patent application No. 2017-30489 filed on February 21, 2017 are incorporated in full herein for reference.

[Examples]

[0051] The present invention is more specifically described by way of Examples. However, the present invention is not limited by the following Examples, and appropriate modifications may be made within the range compatible to the above-mentioned and the later-mentioned gist. All of such modifications are encompassed in the technical scope of the present invention.

Example 1: Evaluation of hepatotoxicity of antisense oligonucleic acid


(1) Selection and synthesis of antisense oligonucleic acid

[0052] The mechanism of induction of hepatotoxicity which has been a problem in nucleic acid medicaments is considered to include (i) suppression of expression of non-target gene, (ii) activation of natural immunity, and (iii) signaling pathway via a biomolecule such as a protein and the like related to nucleic acid medicaments. As an antisense oligonucleic acid which avoids the above-mentioned mechanisms (i) and (ii) and induces particularly strong hepatotoxicity by the contribution of the above-mentioned mechanism (iii), #101 was selected. The synthesis of #101 having the sequence shown in Table 1 and derivative #101-C2-7 thereof was committed to Gene Design, Inc. The molecular weight of the obtained antisense oligonucleic acid was measured by mass spectrum. The theoretical value of the molecular weight of #101 was 4589.72, the measured value was 4600.61, and the theoretical value of the molecular weight of #101-C2-7 was 4605.72, the measured value was 4602.22.

[Table 1]

#101	GTTatgccacc _m CTA
#101-C2-7	GTTatg ζ cacc _m CTA

[0053] In the above-mentioned sequence, capital letters indicate 2', 4' -bridged nucleic acid (LNA) having the following structure, small letters indicate DNA, the base in _mC is 5-methylcytosine, and the base in ζ is 5-hydroxycytosine.



[0054] Each antisense oligonucleic acid was dissolved in physiological saline (manufactured by Otsuka Pharmaceutical Co., Ltd.) to give a 2 mg/mL solution, and the solution was cryopreserved at -30°C until used in experiments.

(2) Hepatotoxicity test

[0055] Five-week-old male C57BL/6NCrI mice (Nihon Charles River Co., Ltd.) were quarantined and acclimated for 1 week and used for experiments. The mice were optionally divided into 3 groups (4 mice per group), and physiological saline (manufactured by Otsuka Pharmaceutical Co., Ltd.) or each antisense oligonucleic acid solution was administered once at a dose of 10 mL/kg body weight (20 mg/kg body weight as dose of antisense oligonucleic acid) from the tail vein. On the fourth day (96 hours later), the mice were anesthetized by inhalation of 2.0 to 4.0% isoflurane (manufactured by DS Pharma Animal Health), and blood was collected from the abdominal portion of posterior vena cava as much as possible. The obtained blood was allowed to stand at room temperature for 20 to 60 min and centrifuged at 1700xg for 5 min to give serum. At the time of blood collection, when the blood volume did not reach the amount necessary for analysis, it was diluted with water for injection and used for the analysis. The concentrations of aspartate transaminase (AST) and alanine transaminase (ALT) in the obtained serum were measured using an automatic biochemical analyzer ("JCA-BM6070" manufactured by JEOL Ltd.). The breeding and experiment of the above-mentioned animal experiment were conducted in the animal experiment facility of Safety Laboratory, SHIN NIPPON BIOMEDICAL LABORATORIES, LTD. in accordance with the animal experiment regulations of SHIN NIPPON BIOMEDICAL LABORATORIES, LTD. The results are shown in Fig. 1. In Fig. 1, "***" indicates presence of a significant difference at $p < 0.05$.

[0056] As shown in the results of Fig. 1, AST and ALT are enzymes to be the indices of liver dysfunction, and the serum concentrations of these enzymes due to particularly high hepatotoxicity of #101 could be reduced significantly only by changing cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 5-hydroxycytosine.

Example 2: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0057] In the same manner as in the above-mentioned Example 1 except that #12, #98, #14 or derivatives thereof which induce strong hepatotoxicity by contribution of the above-mentioned hepatotoxicity induction mechanism (iii) were used instead of antisense oligonucleic acid #101, the hepatotoxicity was evaluated. The theoretical value of the molecular weight of #12 was 4614.73 and the measured value was 4614.27, the theoretical value of the molecular weight of #12-C2-10 was 4630.73 and the measured value was 4627.86, the theoretical value of the molecular weight of #98 was 4613.75 and the measured value was 4612.79, the theoretical value of the molecular weight of #98-C2-7 was 4629.75 and the measured value was 4631.10, the theoretical value of the molecular weight of #14 was 4606.71 and the measured value was 4605.66, and the theoretical value of the molecular weight of #14-C2-6 was 4622.71 and the measured value

was 4619.72. The sequence of each antisense oligonucleic acid used in this Example is shown in Table 2, and the measurement results of the hepatotoxicity are shown in Fig. 2. In Fig. 2, "*" indicates presence of a significant difference at $p < 0.05$.

[Table 2]

#12	GT _m CcgcatgccTAA
#12-C2-10	GT _m Ccgcatg ζ cTAA
#98	GATatgcccta _m CTA
#98-C2-7	GATatg ζ ccta _m CTA
#14	GTA _t g _c ctccgTTA
#14-C2-6	GTA _t g ζ ctccgTTA

[0058] As shown in the results of Fig. 2, the serum concentrations of AST and ALT could be reduced significantly only by changing cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 5-hydroxycytosine also in other antisense oligonucleic acids #12, #98 and #14 having high hepatotoxicity. The serum concentrations of AST and ALT were reduced to several percents by the above-mentioned mutation, though not as much as that in #101 having particularly high hepatotoxicity.

Example 3: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0059] GR (Glucocorticoid Receptor) is one kind of steroid receptors and works as a receptor for the steroid hormone hydrocortisone. It also undergoes nuclear translocation in a ligand-dependent manner and also works as a transcription factor. Therefore, Posi12 (Nucleic Acid Ther., 2012, 22, 5, 344-359) was selected as an antisense oligonucleic acid targeting same, and the synthesis of Posi12 having the sequence shown in Table 3 and its derivative Posi12-C2-11 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi12 was 4611.71 and the measured value was 4610.81, and the theoretical value of the molecular weight of Posi12-C2-11 was 4627.71 and the measured value was 4625.65.

[Table 3]

Posi12	GT _m CtctttaccTGG
Posi12-C2-11	GT _m Ctctttac ζ TGG

[0060] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum was measured in the same manner as in the above-mentioned Example 1 (2). The results are shown in Fig. 3 (1).

[0061] In addition, the GR gene expression-suppressing activity of each antisense oligonucleic acid was measured. To be specific, after blood was collected in the above-mentioned hepatotoxicity test, the mouse was euthanized by freezing, and the liver was removed and the weight was measured. About 100 mg was collected as a sample from a site where no abnormality was found macroscopically in the excised liver. The weight of the collected sample was measured, the sample was frozen in liquid nitrogen and stored in an ultra low temperature freezer. The cryopreserved liver sample was homogenized as much as possible using a homogenizer (Shake Master auto, Bio Medical Science Inc.) and TRIzol reagent (Thermo Fisher Scientific) under ice-cooling and total RNA was extracted. The UV absorption spectrum of the extracted total RNA was measured using Nano Vue or Nano Vue Plus (GE Healthcare), and the purity was calculated from the RNA concentration and O.D.260/O.D.280 ratio. Quantitative PCR was performed on total RNA using One Step SYBR PrimeScript RT-PCR Kit (Takara) and Applied Biosystems 7500 (Life Technologies Japan Ltd), the expression ratio of the target gene in the antisense oligonucleic acid administration group relative to the saline administration group was calculated, and the activity was evaluated. The results are shown in Fig. 3 (2).

[0062] As shown in the results of Fig. 3 (1), the hepatotoxicity by Posi12 was reduced to 40% by only changing cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 5-hydroxycytosine. On the other hand, as shown in the results of Fig. 3 (2), GR gene expression-suppressing activity did not change even when cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi12, was changed to 5-hydroxycytosine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 4: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0063] Posi14 (Nucleic Acid Ther., 2012, 22, 5, 344-359) was selected as an antisense oligonucleic acid targeting mouse GR, and the synthesis of Posi14 having the sequence shown in Table 4 and its derivative Posi14-C2-6 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi14 was 4691.78 and the measured value was 4691.10, and the theoretical value of the molecular weight of Posi14-C2-6 was 4707.78 and the measured value was 4705.24.

[Table 4]

Posi14	A G G t g c t t t g g T_m C T
Posi14-C2-6	A G G t g ζ t t t g g T_m C T

[0064] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and GR gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 4.

[0065] As shown in the results of Fig. 4 (1), the hepatotoxicity by Posi14 was reduced to 70% by only changing cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 5-hydroxycytosine, and the degree of reduction became 13% by removing abnormal measurement values. On the other hand, as shown in the results of Fig. 4 (2), GR gene expression-suppressing activity did not change even when cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, was changed to 5-hydroxycytosine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 5: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0066] PCSK9 (Pro-protein Convertase Subtilisin Kexin 9) is one kind of endoproteases. It reduces LDL uptake by binding to and degrading the receptor for LDL, thus resulting in an increase in the blood LDL level. Posi15 was selected as an antisense oligonucleic acid targeting mouse PCSK9 gene, and the synthesis of Posi15 having the sequence shown in Table 5 and its derivatives Posi15-C2-4 and Posi15-C2-11 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi15 was 4586.77 and the measured value was 4586.73, the theoretical value of the molecular weight of Posi15-C2-4 was 4602.77 and the measured value was 4600.61, and the theoretical value of the molecular weight of Posi15-C2-11 was 4602.77 and the measured value was 4602.09.

[Table 5]

Posi15	A_m C A c c a a g t t c T_m C_m C
Posi15-C2-4	A_m C A ζ c a a g t t c T_m C_m C
Posi15-C2-11	A_m C A c c a a g t t ζ T_m C_m C

[0067] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and PCSK9 gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 5.

[0068] As shown in the results of Fig. 5 (1), the hepatotoxicity by Posi15 was reduced to 40% or 60% by only changing cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 5-hydroxycytosine, and the degree of reduction became 30% or 40% by removing abnormal measurement values. On the other hand, as shown in the results of Fig. 5 (2), PCSK9 gene expression-suppressing activity almost did not change even when cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi15, was changed to 5-hydroxycytosine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 6: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0069] Posi17 was selected as an antisense oligonucleic acid targeting mouse PCSK9 gene, and the synthesis of Posi17 having the sequence shown in Table 6 and its derivative Posi17-C2-10 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi17 was 4300.54 and the measured value was 4300.92, and the theoretical value of the molecular weight of Posi17-C2-10 was 4316.54 and the measured value was 4314.46.

[Table 6]

Posi17	mCTgtgatgac_mCT_mC
Posi17-C2-10	mCTgtgatga_ζmCT_mC

[0070] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and PCSK9 gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 6. In Fig. 6, "" indicates presence of a significant difference at $p < 0.05$.

[0071] As shown in the results of Fig. 6 (1), the hepatotoxicity by Posi17 was significantly reduced to 50% by only changing cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 5-hydroxycytosine. On the other hand, as shown in the results of Fig. 6 (2), PCSK9 gene expression-suppressing activity almost did not change even when cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi17, was changed to 5-hydroxycytosine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 7: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0072] Rps6kb2 (Ribosomal protein S6 kinase beta-2) is a serine-threonine kinase identified as an enzyme that phosphorylates ribosome protein S6, activated by stimulation with growth factor, stress and the like, and regulates cell cycle and protein synthesis. No.97 was selected as an antisense oligonucleic acid targeting same, and the synthesis of No.97 having the sequence shown in Table 7 and its derivatives No.97-C2-7, No.97-C2-9 and No.97-C2-10 was committed to Gene Design, Inc. The theoretical value of the molecular weight of No. 97 was 4182.40 and the measured value was 4183.06, the theoretical value of the molecular weight of No.97-C2-7 was 4198.40 and the measured value was 4197.54, the theoretical value of the molecular weight of No.97-C2-9 was 4198.40 and the measured value was 4196.60, and the theoretical value of the molecular weight of No.97-C2-10 was 4198.40 and the measured value was 4199.40.

[Table 7]

No.97	mCGccctcgcc_mCT_mC
No.97-C2-7	mCGccct_ζgcc_mCT_mC
No.97-C2-9	mCGccctcg_ζc_mCT_mC
No.97-C2-10	mCGccctcg_ζc_mCT_mC

[0073] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and Rps6kb2 expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 7. In Fig. 7, "" indicates presence of a significant difference at $p < 0.05$.

[0074] As shown in the results of Fig. 7 (1), the hepatotoxicity by No.97 was reduced to 33%, 20% or 35% by only changing cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 5-hydroxycytosine. On the other hand, as shown in the results of Fig. 7 (2), GR gene expression-suppressing activity almost did not change even when cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of No.97, was changed to 5-hydroxycytosine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.


Example 8: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0075] In the same manner as in the above-mentioned Example 1 except that #101-T6-5 was used as the derivative of antisense oligonucleic acid #101, hepatotoxicity was evaluated. The theoretical value of the molecular weight of #101-T6-5 was 4605.78 and the measured value was 4605.10. The sequence of each antisense oligonucleic acid used in this Example is shown in Table 8, and the measurement results of the hepatotoxicity are shown in Fig. 8. In Fig. 8, "" indicates presence of a significant difference at $p < 0.05$.

[Table 8]

#101	GTTatgccaccc_mCTA
#101-T6-5	GTTa_κgccacc_mCTA

[0076] In the above-mentioned sequences, the base in κ is 2-thiocarbonylthymine and κ has the following structure:



[0077] As shown in the results of Fig. 8, the serum concentrations of AST and ALT could be reduced by converting thymine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of antisense oligonucleic acid #101 having very high hepatotoxicity, to 2-thiocarbonylthymine.

Example 9: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0078] In the same manner as in the above-mentioned Example 1 except that #98, #14 or a derivative thereof was used instead of antisense oligonucleic acid #101, hepatotoxicity was evaluated. The theoretical value of the molecular weight of #98-T6-5 was 4629.81 and the measured value was 4631.05, and the theoretical value of the molecular weight of #14-T6-4 was 4622.77 and the measured value was 4623.11. The sequence of each antisense oligonucleic acid used in this Example is shown in Table 9, and the measurement results of the hepatotoxicity are shown in Fig. 9. In Fig. 9, "***" indicates presence of a significant difference at $p < 0.05$.

[Table 9]

#98	GATatgcccta _m CTA
#98-T6-5	GATa κ gcccta _m CTA
#14	GATatgcctccgTTA
#14-T6-4	GATa κ gcctccgTTA

[0079] The serum concentrations of AST and ALT could be reduced significantly only by changing thymine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 2-thiocarbonylthymine also in other antisense oligonucleic acids #98 and #14 having high hepatotoxicity.

Example 10: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0080] Posi12 (Nucleic Acid Ther., 2012, 22, 5, 344-359) was selected as an antisense oligonucleic acid targeting mouse GR, and the synthesis of Posi12 having the sequence shown in Table 10 and its derivative Posi12-T6-4 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi12-T6-4 was 4627.77 and the measured value was 4629.03.

[Table 10]

Posi12	GT _m CtctttaccTGG
Posi 12-T6-4	GT _m C κ ctttaccTGG

[0081] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and GR gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 10. In Fig. 10, "***" indicates presence of a significant difference at $p < 0.05$.

[0082] As shown in the results of Fig. 10 (1), the hepatotoxicity by Posi12 was reduced to 40% by only changing thymine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 2-thiocarbonylthymine. On

the other hand, as shown in the results of Fig. 10 (2), GR gene expression-suppressing activity did not change even when thymine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi12, was changed to 2-thiocarbonylthymine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 11: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0083] Posi14 (Nucleic Acid Ther., 2012, 22, 5, 344-359) was selected as an antisense oligonucleic acid targeting mouse GR, and the synthesis of Posi14 having the sequence shown in Table 11 and its derivative Posi14-T6-4 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi14-T6-4 was 4707.84 and the measured value was 4708.38.

[Table 11]

Posi14	A G G t g c t t t g g T_m C T
Posi14-T6-4	A G G_κ g c t t t g g T_m C T

[0084] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and GR gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 11. In Fig. 11, "***" indicates presence of a significant difference at p<0.05.

[0085] As shown in the results of Fig. 11 (1), the hepatotoxicity by Posi14 was reduced to 3% by only changing thymine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 2-thiocarbonylthymine. On the other hand, as shown in the results of Fig. 11 (2), GR gene expression-suppressing activity did not change even when thymine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi14, was changed to 2-thiocarbonylthymine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 12: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0086] Posi17 (Nucleic Acid Ther., 2012, 22, 5, 344-359) was selected as an antisense oligonucleic acid targeting mouse PCSK9, and the synthesis of Posi17 having the sequence shown in Table 12 and its derivative Posi17-T6-4 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi17-T6-4 was 4316.60 and the measured value was 4316.08.

[Table 12]

Posi17	m C T g t g a t g a c m C T_m C
Posi17-T6-4	m C T g_κ g a t g a c m C T_m C

[0087] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and PCSK9 gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 12. In Fig. 12, "***" indicates presence of a significant difference at p<0.05.

[0088] As shown in the results of Fig. 12 (1), the hepatotoxicity by Posi17 was reduced to 60% by only changing thymine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 2-thiocarbonylthymine. On the other hand, as shown in the results of Fig. 12 (2), PCSK9 gene expression-suppressing activity almost did not change even when thymine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi17, was changed to 2-thiocarbonylthymine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.


Example 13: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0089] In the same manner as in the above-mentioned Example 1 except that #101-G3-6 was used as the derivative of antisense oligonucleic acid #101, hepatotoxicity was evaluated. The theoretical value of the molecular weight of #101-G3-6 was 4604.73 and the measured value was 4605.45. The sequence of each antisense oligonucleic acid used in this Example is shown in Table 13, and the measurement results of the hepatotoxicity are shown in Fig. 13. In Fig. 13, "***" indicates presence of a significant difference at p<0.05.

[Table 13]

#101	GTTatgccacc _m CTA
#101-G3-6	GTTatλccacc _m CTA

[0090] In the above-mentioned sequences, the base in λ is 8-aminoguanine and λ has the following structure:



[0091] As shown in the results of Fig. 13, the serum concentrations of AST and ALT could be reduced by only converting guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of antisense oligonucleic acid #101 having very high hepatotoxicity, to 8-aminoguanine.

Example 14: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0092] In the same manner as in the above-mentioned Example 1 except that #12, #98, #14 or derivatives thereof were used, hepatotoxicity was evaluated. The theoretical value of the molecular weight of #12-G3-9 was 4629.74 and the measured value was 4628.19, the theoretical value of the molecular weight of #98-G3-6 was 4628.76 and the measured value was 4628.93, and the theoretical value of the molecular weight of #14-G3-5 was 4621.72 and the measured value was 4621.39. The sequence of each antisense oligonucleic acid used in this Example is shown in Table 14, and the measurement results of the hepatotoxicity are shown in Fig. 14. In Fig. 14, "*" indicates presence of a significant difference at $p < 0.05$.

[Table 14]

#12	GT _m CcgcatgccTAA
#12-G3-9	GT _m CcgcatλccTAA
#98	GATatgcccta _m CTA
#98-G3-6	GATatλcccta _m CTA
#14	GTAtgctccgTTA
#14-G3-5	GTAtλcctccgTTA

[0093] As shown in the results of Fig. 14, the serum concentrations of AST and ALT could be reduced significantly only by changing guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 8-aminoguanine also in other antisense oligonucleic acids #12, #98 and #14 having high hepatotoxicity.


Example 15: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0094] In the same manner as in the above-mentioned Example 1 except that #101 or a derivative thereof #101-G4-6 was used, hepatotoxicity was evaluated. The theoretical value of the molecular weight of #101-G4-6 was 4668.61 and the measured value was 4670.31. The sequence of each antisense oligonucleic acid used in this Example is shown in Table 15, and the measurement results of the hepatotoxicity are shown in Fig. 15. In Fig. 15, "*" indicates presence of a significant difference at $p < 0.05$.

[Table 15]

#101	GTTatgccacc _m CTA
#101-G4-6	GTTat μ ccacc _m CTA

[0095] In the above-mentioned sequences, the base in μ is 8-bromoguanine and μ has the following structure:



[0096] As shown in the results of Fig. 15, the serum concentrations of AST and ALT could be reduced by only converting guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of antisense oligonucleic acid #101 having very high hepatotoxicity, to 8-bromoguanine.

Example 16: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0097] In the same manner as in the above-mentioned Example 1 except that #12, #98, #14 or derivatives thereof were used instead of antisense oligonucleic acid #101, hepatotoxicity was evaluated. The theoretical value of the molecular weight of #12-G4-9 was 4693.62 and the measured value was 4692.50, the theoretical value of the molecular weight of #98-G4-6 was 4692.64 and the measured value was 4693.45, the theoretical value of the molecular weight of #14-G4-5 was 4685.60 and the measured value was 4683.86. The sequence of each antisense oligonucleic acid used in this Example is shown in Table 16, and the measurement results of the hepatotoxicity are shown in Fig. 16. In Fig. 16, "*" indicates presence of a significant difference at $p < 0.05$.

[Table 16]

#12	GT _m CcgcattgccTAA
#12-G4-9	GT _m Ccgcatt μ ccTAA
#98	GATatgcccta _m CTA
#98-G4-6	GATat μ cccta _m CTA
#14	GATatgcctccgTTA
#14-G4-5	GATat μ ccctccgTTA

[0098] As shown in the results of Fig. 16, the serum concentrations of AST and ALT could be reduced significantly only by changing guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 8-bromoguanine also in other antisense oligonucleic acids #12, #98 and #14 having high hepatotoxicity.

Example 17: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0099] Posi14 (Nucleic Acid Ther., 2012, 22, 5, 344-359) was selected as an antisense oligonucleic acid targeting mouse GR, and the synthesis of Posi14 having the sequence shown in Table 17 and its derivative Posi14-G4-10 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi14-G4-10 was 4770.67 and the measured value was 4771.46.

[Table 17]

Posi14	A G G t g c t t t g g T_m C T
Posi 14-G4-10	A G G t g c t t t μ g T_m C T

[0100] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and GR gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 17. In Fig. 17, "*" indicates presence of a significant difference at p<0.05.

[0101] As shown in the results of Fig. 17 (1), the hepatotoxicity by Posi14 was reduced to 30% by only changing guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 8-bromoguanine. On the other hand, as shown in the results of Fig. 17 (2), GR gene expression-suppressing activity almost did not change even when guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi14, was changed to 8-bromoguanine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 18: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0102] Posi15 was selected as an antisense oligonucleic acid targeting mouse PCSK9, and the synthesis of Posi15 having the sequence shown in Table 18 and its derivative Posi15-G4-8 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi15-G4-8 was 4665.66 and the measured value was 4666.01.

[Table 18]

Posi15	A_m C A c c a a g t t c T_m C_m C
Posi15-G4-8	A_m C A c c a a μ t t c T_m C_m C

[0103] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and PCSK9 gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 18. In Fig. 18, "*" indicates presence of a significant difference at p<0.05.

[0104] As shown in the results of Fig. 18 (2), PCSK9 gene expression-suppressing activity somewhat decreased when guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi15, was changed to 8-bromoguanine. As shown in the results of Fig. 18 (1), hepatotoxicity due to Posi15 was significantly reduced to 1%. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid too much.

Example 19: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0105] Posi17 was selected as an antisense oligonucleic acid targeting mouse PCSK9, and the synthesis of Posi17 having the sequence shown in Table 19 and its derivatives Posi17-G4-3, Posi17-G4-5 and Posi17-G4-8 was committed to Gene Design, Inc. The theoretical value of the molecular weight of Posi17-G4-3 was 4379.43 and the measured value was 4378.54, the theoretical value of the molecular weight of Posi17-G4-5 was 4379.43 and the measured value was 4377.25, and the theoretical value of the molecular weight of Posi17-G4-8 was 4379.43 and the measured value was 4378.34.

[Table 19]

Posi 17	m C T g t g a t g a c_m C T_m C
Posi17-G4-3	m C T μ t g a t g a c_m C T_m C
Posi17-G4-5	m C T g t μ a t g a c_m C T_m C
Posi 17-G4-8	m C T g t g a t μ a c_m C T_m C

[0106] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and PCSK9 gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 19. In Fig. 19, "*" indicates presence of a significant difference at p<0.05.

[0107] As shown in the results of Fig. 19 (1), the hepatotoxicity by Posi17 was significantly reduced to 3%, 9% or 20%

by only changing guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 8-bromoguanine. The hepatotoxicity reducing effect of Posi17-G4-8 showed reduction to 3% when abnormal measurement values were removed. On the other hand, as shown in the results of Fig. 19 (2), PCSK9 gene expression-suppressing activity almost did not change even when guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi17, was changed to 8-bromoguanine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 20: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0108] No.97 was selected as an antisense oligonucleic acid targeting mouse Rps6kb2, and the synthesis of No.97 having the sequence shown in Table 20 and its derivative No.97-G4-8 was committed to Gene Design, Inc. The theoretical value of the molecular weight of No.97-G4-8 was 4261.29 and the measured value was 4261.11.

[Table 20]

No.97	mCGccctcgccmCTmC
No.97-G4-8	mCGccctcμccmCTmC

[0109] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and Rps6kb2 gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 20. In Fig. 20, "*" indicates presence of a significant difference at $p < 0.05$.

[0110] As shown in the results of Fig. 20 (1), the hepatotoxicity by No.97 was significantly reduced to 9% by only changing guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 8-bromoguanine. On the other hand, as shown in the results of Fig. 20 (2), Rps6kb2 gene expression-suppressing activity almost did not change even when guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of No.97, was changed to 8-bromoguanine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.


Example 21: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0111] In the same manner as in the above-mentioned Example 5 except that Posi15 derivative, Posi15-G6-8, was used, hepatotoxicity and activity were evaluated. The theoretical value of the molecular weight of Posi15-G6-8 was 4685.90 and the measured value was 4683.03. The sequence of each antisense oligonucleic acid used in this Example is shown in Table 21, and the measurement results of the hepatotoxicity are shown in Fig. 21. In Fig. 21, "*" indicates presence of a significant difference at $p < 0.05$.

[Table 21]

Posi 15	A_mCAccaagt_mttcT_mC_mC
Posi15-G6-8	A_mCAccaa_αttcT_mC_mC

[0112] In the above-mentioned sequences, the base in α is 7-phenylethynylguanine and α has the following structure:



[0113] As shown in the results of Fig. 21 (1), the hepatotoxicity by Posi15 was significantly reduced to 5% by only

changing guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 7-phenylethynyl-guanine and removing abnormal measurement values. On the other hand, as shown in the results of Fig. 21 (2), PCSK9 gene expression-suppressing activity almost did not change even when guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi15, was changed to 7-phenylethynylguanine. Thus, according to the

Example 22: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0114] No.97 and Posi14 were respectively selected as antisense oligonucleic acids targeting mouse Rps6kb2 and mouse GR, and the synthesis of No.97, Posi14 having the sequences shown in Table 22 and derivatives in which guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, was changed to 8-aminoguanine was committed to Gene Design, Inc. In addition, Posi15 and Posi17 were respectively selected as antisense oligonucleic acids targeting mouse PCSK9 gene and the synthesis of Posi15, Posi17 having the sequences shown in Table 22 and derivatives in which guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, was changed to 8-aminoguanine was committed to Gene Design, Inc.

[Table 22]

No.97	mCGccctcgcc_mCT_mC
No.97-G3-8	mCGccctcλcc_mCT_mC
Posi14	AGGtgctttggT_mCT
Posi14-G3-5	AGGtλcctttggT_mCT
Posi14-G3-10	AGGtgctttλgT_mCT
Posi15	A_mCAccaagt_mttcT_mC_mC
Posi15-G3-8	A_mCAccaaλttcT_mC_mC
Posi17	mCTgtgatgac_mCT_mC
Posi17-G3-3	mCTλtgatgac_mCT_mC

[0115] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and mouse Rps6kb2 gene, mouse GR gene or mouse PCSK9 gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Figs. 22 - 25.


[0116] As shown in the results of Figs. 22 - 25 (1), the hepatotoxicity by No.97, Posi14, Posi15 and Posi17 was clearly reduced by only changing guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 8-aminoguanine. On the other hand, as shown in the results of Figs. 22 - 25 (2), mouse Rps6kb2 gene expression-, mouse GR gene expression- and mouse PCSK9 gene expression-suppressing activities almost did not change except a part thereof even when guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of No.97, Posi14, Posi15 and Posi17, was changed to 8-aminoguanine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 23: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0117] #12 was selected as an antisense oligonucleic acid that induces strong hepatotoxicity, and the synthesis of #12 having the sequence shown in Table 23 and a derivative in which guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, was changed to 7-substituted guanine having the following structure was committed to Gene Design, Inc. The compound used for introduction of the 7-substituted guanosine derivative is novel and the synthesis method thereof is described later.

[Table 23]

#12	GT_mCcgcatg_mccTAA
#12-G6-9	GT_mCcgcatαccTAA
#12-G7-9	GT_mCcgcatβccTAA
#12-G8-9	GT_mCcgcatγccTAA
#12-G9-9	GTC_mCcgcatδccTAA



[0118] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum was measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 26. As shown in the results of Fig. 26, hepatotoxicity by #12 was effectively decreased by only introducing a substituent into the 7-position of guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part.

Example 24: Evaluation of hepatotoxicity and activity of antisense oligonucleic acid

[0119] Posi14 and Posi15 were respectively selected as an antisense oligonucleic acid targeting mouse GR and mouse PCSK9 gene, and the synthesis of Posi14, Posi15 having the sequences shown in Table 24 and a derivative in which guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, was changed to 7-substituted guanine was committed to Gene Design, Inc.

[Table 24]

Posi14	A G G t g c t t t g g T _m C T
Posi14-G7-5	A G G t _β c t t t g g T _m C T
Posi14-G9-5	A G G t _δ c t t t g g T _m C T
Posi14-G9-10	A G G t g c t t t _δ g T _m C T
Posi15	A _m C A c c a a g t t c T _m C _m C
Posi15-G7-8	A _m C A c c a a _β t t c T _m C _m C
Posi15-G9-8	A _m C A c c a a _δ t t c T _m C _m C

[0120] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum and mouse GR gene or mouse PCSK9 gene expression-suppressing activity were measured in the same manner as in the above-mentioned Example 3. The results are shown in Figs. 27, 28.

[0121] As shown in the results of Fig. 27, 28 (1), hepatotoxicity by Posi14 and Posi15 was clearly reduced by only changing guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 7-substituted guanine. On the other hand, as shown in the results of Figs. 27, 28 (2), mouse GR gene expression- and mouse PCSK9 gene expression-suppressing activities almost did not change even when guanine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part of Posi14 and Posi15, was changed to 7-substituted guanine. Thus, according to the present invention, hepatotoxicity can be reduced without reducing the activity of antisense oligonucleic acid.

Example 25: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0122] #101 was selected as an antisense oligonucleic acid that induces strong hepatotoxicity, and the synthesis of #101 having the sequence shown in Table 25 and a derivative in which cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, was changed to 5-hydroxycytosine was committed to Gene Design, Inc.

[Table 25]

#101	G T T a t g c c a c c _m C T A
#101-C2-7	G T T a t g _ζ c a c c _m C T A
#101-C2-8	G T T a t g c _ζ a c c _m C T A
#101-C2-10	G T T a t g c c a _ζ c _m C T A

(continued)

#101-C2-11	GTTatgccc ζ _m CTA
------------	------------------------------------

[0123] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum was measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 29. As shown in the results of Fig. 29, hepatotoxicity by #101 was effectively decreased by only changing cytosine, which is in one of the 2',4'-non-bridged nucleic acid residues in the middle part, to 5-hydroxycytosine.

Example 26: Evaluation of hepatotoxicity of antisense oligonucleic acid

[0124] #101 was selected as an antisense oligonucleic acid that induces strong hepatotoxicity, and the synthesis of #101 having the sequence shown in Table 26 and a derivative in which 1 to 3 bases in the 2',4'-non-bridged nucleic acid residues in the middle part was(were) changed to 5-hydroxycytosine (ζ), 2-thiocarbonylthymine (κ) or 8-bromoguanine (μ) was committed to Gene Design, Inc.


[Table 26]

#101	GTTatgccc _m CTA
#101-C2-7	GTTatg ζ cacc _m CTA
#101-T6-5	GTTa κ gccacc _m CTA
#101-G4-6	GTTat μ ccacc _m CTA
#101-T6C2	GTTa κ g ζ cacc _m CTA
#101-T6G4	GTTa κ μ ccacc _m CTA
#101-G4C2	GTTat μ ζ cacc _m CTA
#101-T6G4C2	GTTa κ μ ζ cacc _m CTA

[0125] With regard to each of the obtained antisense oligonucleic acids, the ALT concentration in mouse serum was measured in the same manner as in the above-mentioned Example 3. The results are shown in Fig. 30. As shown in the results of Fig. 30, it was demonstrated that hepatotoxicity by #101 was effectively decreased by only changing one base in the 2',4'-non-bridged nucleic acid residues in the middle part to a derivative, and the hepatotoxicity attenuation effect is increased still more by changing two or more bases in the 2',4'-non-bridged nucleic acid residues in the middle part, to the derivative(s).

Example 27: Synthesis of 7-deaza-7-iodoguanosine derivative

[0126]



(1) Synthesis of compound 2

[0127] Compound 1 was synthesized according to F. Seela et al., Synthesis, 2004, 8, 1203-1210 and M. Mackova et al., ChemBioChem, 2015, 16, 2225-2236.

[0128] Tetramethylguanidine (4.3 mL, 35 mmol) and pyridine-2-aldoxime (4.3 g, 35 mmol) were added to a mixed solution of compound 1 (5.1 g, 7.0 mmol) in DMF/1,4-dioxane (170 mL, DMF:1,4-dioxane=1:1), and the mixture was stirred under a nitrogen atmosphere at room temperature overnight. The mixture was diluted with ethyl acetate, washed successively with 1N aqueous hydrochloric acid solution, saturated aqueous sodium hydrogen carbonate and saturated brine, dried over anhydrous sodium sulfate and concentrated to give a residue. The residue was washed with methanol to give compound 2 (4.9 g) as a white solid.

¹H NMR (300 MHz, DMSO-d₆) δH: 1.22(9H,s), 2.38(3H,s), 2.40(3H,s), 2.64(1H,m), 2.91(1H,m), 4.45-4.62(3H,m), 5.63(1H, d, J=5.9Hz), 6.55(1H, dd, J=9.1,5.5Hz), 7.34-7.41(5H,m), 7.88-7.94(4H,m), 10.96(1H,s), 11.95(1H,s)

(2) Synthesis of compound 3

[0129] Under ice-cooling, 1N sodium methoxide (12.4 mL, 12.4 mmol) was added to a mixed solution of compound 2 (4.5 g, 6.2 mmol) in THF/methanol (151.5 mL, THF:methanol=150:1.5), and the mixture was stirred under a nitrogen atmosphere under ice-cooling for 15 min. Under ice-cooling, acetic acid (0.8 mL) was added and the mixture was concentrated to give a residue. The residue was purified by silica gel column chromatography (eluent: chloroform/methanol=9/1) to give compound 3 (3.0 g) as a white solid.

¹H NMR (300 MHz, DMSO-d₆) δH: 1.23(9H,s), 2.09(1H,m), 2.36(1H,m), 3.50(2H,m), 3.76(1H,s), 4.29(1H,s), 4.93(1H, t, J=5.1Hz), 5.23(1H,d,J=3.3Hz), 6.45(1H, dd, J=8.1,5.5Hz), 7.45(1H,s), 10.97(1H,s), 11.90(1H,s)

(3) Synthesis of compound 4

[0130] Dimethoxytrityl chloride (0.8 g, 2.4 mmol) was added to a solution (15 mL) of compound 3 (1.0 g, 2.2 mmol) in pyridine, and the mixture was stirred under a nitrogen atmosphere, overnight. Methanol was added and the mixture was stirred for 1 hr, diluted with chloroform, washed with saturated aqueous sodium hydrogen carbonate and saturated brine, dried over sodium sulfate and concentrated to give a residue. The residue was purified by silica gel column chromatography (eluent: chloroform/methanol/triethylamine=94/5/1) to give compound 4 (1.3 g) as a white solid.


¹H NMR (300 MHz, DMSO-d₆) δH: 1.20(9H,s), 2.15 (1H,m), 2.40(1H,m), 3.04(1H,m), 3.13(1H,m), 3.70(6H,s), 3.86(1H,m), 4.27(1H,m), 5.26(1H,d,J=3.7Hz), 6.43(1H,m), 6.81-6.84(4H,m), 7.15-7.35(10H,m), 10.94(1H,s), 11.90(1H,s)

(4) Synthesis of compound 5

[0131] 2-Cyanoethyl diisopropylchlorophosphoramidite (0.27 mL, 1.2 mmol) was added to a solution (30 mL) of diisopropylethylamine (0.28 mL, 1.7 mmol) and compound 4 (0.5 g, 0.6 mmol) in dichloromethane, and the mixture was stirred under a nitrogen atmosphere for 1 hr, diluted with dichloromethane, washed successively with saturated aqueous sodium hydrogen carbonate and saturated brine, dried over anhydrous sodium sulfate, and concentrated to give a residue. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate/triethylamine=10/9/1) to give the title compound (0.4 g) as a white solid.

³¹P NMR (122 MHz, CDCl₃) δP: 148.53, 148.91

Example 28: Synthesis of 7-deaza-7-(2-phenylethynyl)guanosine derivative

[0132]

(1) Synthesis of compound 6


[0133] To a solution (5 mL) of copper iodide (6 mg, 0.03 mmol), triethylamine (0.09 mL, 0.6 mmol), ethynylbenzene (0.04 mL, 0.4 mmol) and compound 4 (0.2 g, 0.3 mmol) in acetonitrile was added tetrakis(triphenylphosphine)palladium(0) (35 mg, 0.03 mmol), and the mixture was stirred under a nitrogen atmosphere at 80°C for 2 hr. The reaction mixture was filtered through celite, and concentrated to give a residue. The residue was purified by silica gel column chromatography (ethyl acetate:triethylamine=98:2) to give the title compound (0.15 g). ¹H NMR (300 MHz, CDCl₃)δH: 1.30(9H,s), 1.90(1H,d,J=3.7Hz), 2.42(2H,m), 3.28(1H,m), 3.40(1H,m), 3.73(6H,s), 4.03(1H,m), 4.55(1H,m), 6.44(1H,m), 6.81-6.84(4H,m), 7.13(1H,s), 7.17-7.33(10H,m), 7.41-7.43(2H,m), 7.50-7.54(2H,m), 7.91(1H,s), 11.69(1H,s)

(2) Synthesis of compound 7

[0134] 2-Cyanoethyl diisopropylchlorophosphoramidite (0.16 mL, 0.7 mmol) was added to a solution (20 mL) of diisopropylethylamine (0.15 mL, 1.2 mmol) and compound 6 (0.4 g, 0.5 mmol) in dichloromethane, and the mixture was stirred under a nitrogen atmosphere for 1 hr, diluted with dichloromethane, washed successively with saturated aqueous sodium hydrogen carbonate and saturated brine, dried over anhydrous sodium sulfate and concentrated to give a residue. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate/triethylamine=10/9/1) to give the title compound (0.4 g) as a white solid.

³¹P NMR (162 MHz, CDCl₃)δP: 148.70, 148.97

Example 29: Synthesis of 7-deaza-7-[2-(4-pyridyl)ethynyl]guanosine derivative

[0135]

(1) Synthesis of compound 8

[0136] To an acetonitrile solution (2.5 mL) containing copper iodide (2 mg, 0.01 mmol), triethylamine (0.04 mL, 0.3 mmol), p-ethynylpyridine (20 mg, 0.2 mmol) and compound 4 (0.1 g, 0.1 mmol) was added tetrakis(triphenylphosphine)palladium(0) (10 mg, 0.01 mmol), and the mixture was stirred under a nitrogen atmosphere at 80°C for 1 hr. The reaction mixture was filtered through celite and concentrated to give a residue. The residue was purified by silica gel column chromatography (eluent: ethyl acetate/triethylamine=98/2) to give the title compound (0.08 g) as a pale-yellow solid.

¹H NMR (300 MHz, CDCl₃)δH: 1.31(9H,s), 2.21(1H,m), 2.43(2H,m), 3.30(1H,m), 3.40(1H,m), 3.73(6H,s), 4.06(1H,m), 4.57(1H,brs), 6.41(1H,tr,J=6.6Hz), 6.80-6.83(4H,m), 7.28-7.43(10H,m), 7.17-7.22(2H,m), 8.14(1H,s), 8.53-8.55(2H,m), 11.78(1H,s)


(2) Synthesis of compound 9

[0137] 2-Cyanoethyl diisopropylchlorophosphoramidite (0.15 mL, 0.7 mmol) was added to a solution (30 mL) of diisopropylethylamine (0.3 mL, 1.7 mmol) and compound 8 (0.5 g, 0.7 mmol) in dichloromethane, and the mixture was stirred under a nitrogen atmosphere for 1 hr. The mixture was diluted with dichloromethane, washed successively with saturated aqueous sodium hydrogen carbonate and saturated brine, dried over anhydrous sodium sulfate and concentrated to give a residue. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate/triethylamine=10/9/1) to give the title compound (0.5 g) as a white solid.

³¹P NMR (162 MHz, CDCl₃)δP: 148.64, 149.07

Example 30: Synthesis of 7-deaza-7-[2-(acetoxymethyl)ethynyl]guanosine derivative

[0138]



(1) Synthesis of compound 10

[0139] To a solution (2.5 mL) of copper iodide (4 mg, 0.01 mmol), triethylamine (0.09 mL, 0.7 mmol), 1-acetoxy-2-propyne (20 mg, 0.2 mmol) and compound 4 (0.1 g, 0.1 mmol) in acetonitrile was added tetrakis(triphenylphosphine)palladium(0) (22 mg, 0.01 mmol), and the mixture was stirred under a nitrogen atmosphere at 80°C for 1 hr. The reaction mixture was filtered through celite and concentrated to give a residue. The residue was purified by silica gel column chromatography (eluent: chloroform/methanol/triethylamine=98/1/1) and repurified by amino silica gel silica gel column chromatography (eluent: chloroform/methanol/triethylamine=98/1/1) to give the title compound (0.08 g) as a pale-yellow solid.

¹H NMR (300 MHz, CDCl₃)δH: 1.28(9H,s), 2.08(3H,s), 2.15(1H,d,J=3.2Hz), 2.37(2H,m), 3.24(1H,m), 3.37(1H,m), 3.77(6H,s), 4.01(1H,m), 4.50(1H,brs), 4.91(2H,s), 6.38(1H,m), 6.80-6.83(4H,m), 7.07(1H,s), 7.81-7.41(9H,m), 8.03(1H,s), 11.73(1H,s)

(2) Synthesis of compound 11

[0140] 2-Cyanoethyl diisopropylchlorophosphoramidite (0.1 mL, 0.4 mmol) was added to a solution (20 mL) of diisopropylethylamine (0.2 mL, 1.0 mmol) and compound 10 (0.3 g, 0.4 mmol) in dichloromethane, and the mixture was stirred under a nitrogen atmosphere for 1 hr. The mixture was diluted with dichloromethane, washed successively with saturated aqueous sodium hydrogen carbonate and saturated brine, dried over anhydrous sodium sulfate and concentrated to give a residue. The residue was purified by silica gel column chromatography (eluent: hexane/ethyl acetate/triethylamine=10/9/1) to give the title compound (0.3 g) as a white solid.

³¹P NMR (122 MHz, CDCl₃)δP: 148.72, 149.04

SEQUENCE LISTING

<110> Osaka University
 5 <120> Antisense Oligonucleic Acid
 <130> F18-006PCT
 <150> JP2017-30489
 10 <151> 2017-02-21
 <160> 61
 <210> 1
 <211> 14
 15 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> modified_base
 <222> (1)..(1)
 20 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 <222> (2)..(2)
 25 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 <222> (3)..(3)
 30 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 <222> (12)..(12)
 35 <223> n stands for 2',4'-bridged m5c
 <220>
 <221> modified_base
 <222> (13)..(13)
 40 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 <222> (14)..(14)
 45 <223> n stands for 2',4'-bridged adenosine
 <400> 1
 nnnatgccac cnnn

14

<210> 2
 <211> 14
 <212> DNA
 50 <213> Artificial Sequence
 <220>
 <221> modified_base
 <222> (1)..(1)
 55 <223> n stands for 2',4'-bridged guanosine
 <220>

<221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 5 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine
 10 <220>
 <221> modified_base
 <222> (7)..(7)
 <223> n stands for 5-hydroxycytidine
 15 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c
 20 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 25 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 2
 nnnatgncac cnnn
 30 <210> 3
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 35 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 40 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 45 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged m5c
 50 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 55 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged adenosine

14

<220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 5
 <400> 3
 nnnccgcatgc cnnn 14

<210> 4
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 10

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 15

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 20

<220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged m5c
 25

<220>
 <221> modified_base
 <222> (10)..(10)
 <223> n stands for 5-hydroxycytidine
 30

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 35

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged adenosine
 40

<220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 45

<400> 4
 nnnccgcatgn cnnn 14

<210> 5
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 50

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 55

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged adenosine

5

<220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

10

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c

15

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

20

<220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

<400> 5
 nnnatgccct annn

25

14

<210> 6
 <211> 14
 <212> DNA
 <213> Artificial Sequence

30

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

35

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged adenosine

40

<220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

45

<220>
 <221> modified_base
 <222> (7)..(7)
 <223> n stands for 5-hydroxycytidine

50

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c

55

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

5 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 6
 nnnatgncct annn 14

10 <210> 7
 <211> 14
 <212> DNA
 <213> Artificial Sequence

15 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

20 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

25 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine

30 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

35 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

40 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 7
 nnntgcctcc gnnn 14

45 <210> 8
 <211> 14
 <212> DNA
 <213> Artificial Sequence

50 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

55 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

<220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine
 5
 <220>
 <221> modified_base
 <222> (6)..(6)
 <223> n stands for 5-hydroxycytidine
 10
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 15
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 20
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 8
 25 nnntgntcc gnnn
 <210> 9
 <211> 14
 <212> DNA
 30 <213> Artificial Sequence
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 35
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 40
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged m5c
 45
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 50
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged guanosine
 55
 <220>
 <221> modified_base
 <222> (14)..(14)

14

<223> n stands for 2',4'-bridged guanosine

<400> 9
nnntctttac cnnn

14

<210> 10
<211> 14
<212> DNA
<213> Artificial Sequence

<220>
<221> modified_base
<222> (1)..(1)
<223> n stands for 2',4'-bridged guanosine

<220>
<221> modified_base
<222> (2)..(2)
<223> n stands for 2',4'-bridged thymidine

<220>
<221> modified_base
<222> (3)..(3)
<223> n stands for 2',4'-bridged m5c

<220>
<221> modified_base
<222> (11)..(11)
<223> n stands for 5-hydroxycytidine

<220>
<221> modified_base
<222> (12)..(12)
<223> n stands for 2',4'-bridged thymidine

<220>
<221> modified_base
<222> (13)..(13)
<223> n stands for 2',4'-bridged guanosine

<220>
<221> modified_base
<222> (14)..(14)
<223> n stands for 2',4'-bridged guanosine

<400> 10
nnntctttac nnnn

14

<210> 11
<211> 14
<212> DNA
<213> Artificial Sequence

<220>
<221> modified_base
<222> (1)..(1)
<223> n stands for 2',4'-bridged adenosine

<220>
<221> modified_base
<222> (2)..(2)

<223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 5 <222> (3)..(3)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 10 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 15 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 <220>
 <221> modified_base
 20 <222> (14)..(14)
 <223> n stands for 2',4'-bridged thymidine
 <400> 11
 nnntgctttg gnnn
 25 <210> 12
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 30 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine
 <220>
 35 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 40 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 45 <221> modified_base
 <222> (6)..(6)
 <223> n stands for 5-hydroxycytidine
 <220>
 50 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 55 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 <220>
 <221> modified_base

14

<222> (14)..(14)
 <223> n stands for 2',4'-bridged thymidine

 <400> 12
 5 nnntgntttg gnnn 14

 <210> 13
 <211> 14
 <212> DNA
 10 <213> Artificial Sequence

 <220>
 <221> modified_base
 <222> (1)..(1)
 15 <223> n stands for 2',4'-bridged adenosine

 <220>
 <221> modified_base
 <222> (2)..(2)
 20 <223> n stands for 2',4'-bridged m5c

 <220>
 <221> modified_base
 <222> (3)..(3)
 25 <223> n stands for 2',4'-bridged adenosine

 <220>
 <221> modified_base
 <222> (12)..(12)
 30 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 <222> (13)..(13)
 35 <223> n stands for 2',4'-bridged m5c

 <220>
 <221> modified_base
 <222> (14)..(14)
 40 <223> n stands for 2',4'-bridged m5c

 <400> 13
 nnnccaagtt cnnn 14

 <210> 14
 <211> 14
 <212> DNA
 45 <213> Artificial Sequence

 <220>
 <221> modified_base
 <222> (1)..(1)
 50 <223> n stands for 2',4'-bridged adenosine

 <220>
 <221> modified_base
 <222> (2)..(2)
 55 <223> n stands for 2',4'-bridged m5c

 <220>
 <221> modified_base

<222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine

 5 <220>
 <221> modified_base
 <222> (4)..(4)
 <223> n stands for 5-hydroxycytidine

 10 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 15 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

 20 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged m5c

 <400> 14
 nnnncaagtt cnnn

 25 <210> 15
 <211> 14
 <212> DNA
 <213> Artificial Sequence

 30 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine

 35 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged m5c

 40 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine

 45 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 5-hydroxycytidine

 50 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

 55 <220>

14

<221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged m5c
 5 <400> 15
 nnnccaagtt nnnn 14

<210> 16
 <211> 13
 10 <212> DNA
 <213> Artificial Sequence

<220>
 <221> modified_base
 15 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (2)..(2)
 20 <223> n stands for 2',4'-bridged thymidine

<220>
 <221> modified_base
 <222> (11)..(11)
 25 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

30 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

35 <400> 16
 nngtgatgac nnn 13

<210> 17
 <211> 13
 40 <212> DNA
 <213> Artificial Sequence

<220>
 <221> modified_base
 45 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (2)..(2)
 50 <223> n stands for 2',4'-bridged thymidine

<220>
 <221> modified_base
 <222> (10)..(10)
 55 <223> n stands for 5-hydroxycytidine

<220>

<221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 5 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 10 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 15 <400> 17
 nnngtgatgan nnn 13
 20 <210> 18
 <211> 13
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c
 30 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine
 35 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 40 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 45 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 <400> 18
 nnccctcgcc nnn 13
 50 <210> 19
 <211> 13
 <212> DNA
 <213> Artificial Sequence
 55 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c

<221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine
 5 <220>
 <221> modified_base
 <222> (7)..(7)
 <223> n stands for 5-hydroxycytidine
 10 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 15 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 20 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 <400> 19
 nncctngcc nnn
 25
 <210> 20
 <211> 13
 <212> DNA
 <213> Artificial Sequence
 30 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c
 35 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine
 40 <220>
 <221> modified_base
 <222> (9)..(9)
 <223> n stands for 5-hydroxycytidine
 45 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 50 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 55 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

13

<400> 20
nnccctcgnc nnn

13

5 <210> 21
 <211> 13
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c

15 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine

20 <220>
 <221> modified_base
 <222> (10)..(10)
 <223> n stands for 5-hydroxycytidine

25 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c

30 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

35 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

 <400> 21
 nnccctcgcn nnn

13

40 <210> 22
 <211> 14
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

50 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

55 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

```

5
  <220>
  <221> modified_base
  <222> (12)..(12)
  <223> n stands for 2',4'-bridged m5c

  <220>
  <221> modified_base
  <222> (13)..(13)
  <223> n stands for 2',4'-bridged thymidine

10
  <220>
  <221> modified_base
  <222> (14)..(14)
  <223> n stands for 2',4'-bridged adenosine

15
  <400> 22
  nnnatgccac cnnn

20
  <210> 23
  <211> 14
  <212> DNA
  <213> Artificial Sequence

  <220>
  <221> modified_base
  <222> (1)..(1)
  <223> n stands for 2',4'-bridged guanosine

  <220>
  <221> modified_base
  <222> (2)..(2)
  <223> n stands for 2',4'-bridged thymidine

30
  <220>
  <221> modified_base
  <222> (3)..(3)
  <223> n stands for 2',4'-bridged thymidine

35
  <220>
  <221> modified_base
  <222> (5)..(5)
  <223> n stands for thiocarbonylthymidine

40
  <220>
  <221> modified_base
  <222> (12)..(12)
  <223> n stands for 2',4'-bridged m5c

45
  <220>
  <221> modified_base
  <222> (13)..(13)
  <223> n stands for 2',4'-bridged thymidine

50
  <220>
  <221> modified_base
  <222> (14)..(14)
  <223> n stands for 2',4'-bridged adenosine

55
  <400> 23
  nnnangccac cnnn

```

<210> 24
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 10
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged adenosine
 15
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine
 20
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c
 25
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 30
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 24
 nnnatgccct annn
 35
 <210> 25
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 45
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged adenosine
 50
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine
 55
 <220>
 <221> modified_base
 <222> (5)..(5)
 <223> n stands for 2-thiocarbonylthymidine

14

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c
 5
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 10
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 15
 <400> 25
 nnnangccct annn 14
 <210> 26
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 20
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 25
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 30
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine
 35
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 40
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 45
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 50
 <400> 26
 nnntgcctcc gnnn 14
 <210> 27
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 55

<220>
 <221> modified_base
 <222> (1)..(1)
 5 <223> n stands for 2',4'-bridged guanosine

 <220>
 <221> modified_base
 <222> (2)..(2)
 10 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 <222> (3)..(3)
 15 <223> n stands for 2',4'-bridged adenosine

 <220>
 <221> modified_base
 <222> (4)..(4)
 20 <223> n stands for 2-thiocarbonylthymidine

 <220>
 <221> modified_base
 <222> (12)..(12)
 25 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 <222> (13)..(13)
 30 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 <222> (14)..(14)
 35 <223> n stands for 2',4'-bridged adenosine

 <400> 27
 nnnngcctcc gnnn

 <210> 28
 <211> 14
 <212> DNA
 40 <213> Artificial Sequence

 <220>
 <221> modified_base
 <222> (1)..(1)
 45 <223> n stands for 2',4'-bridged guanosine

 <220>
 <221> modified_base
 <222> (2)..(2)
 50 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 <222> (3)..(3)
 55 <223> n stands for 2',4'-bridged m5c

 <220>
 <221> modified_base
 <222> (12)..(12)

14

<223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 5 <222> (13)..(13)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 10 <222> (14)..(14)
 <223> n stands for 2',4'-bridged guanosine
 <400> 28
 nnntctttac cnnn 14
 15 <210> 29
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 20 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 25 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 30 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged m5c
 <220>
 35 <221> modified_base
 <222> (4)..(4)
 <223> n stands for 2-thiocarbonylthymidine
 <220>
 40 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 45 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 50 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged guanosine
 <400> 29
 nnnnctttac cnnn 14
 55 <210> 30
 <211> 14
 <212> DNA

<213> Artificial Sequence
 <220>
 <221> modified_base
 5 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine
 <220>
 <221> modified_base
 10 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 15 <222> (3)..(3)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 20 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 25 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 <220>
 <221> modified_base
 30 <222> (14)..(14)
 <223> n stands for 2',4'-bridged thymidine
 <400> 30
 nnntgctttg gnnn
 14
 <210> 31
 <211> 14
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <221> modified_base
 40 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine
 <220>
 <221> modified_base
 45 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 50 <222> (3)..(3)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 55 <222> (4)..(4)
 <223> n stands for 2-thiocarbonylthymidine
 <220>
 <221> modified_base

<222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 5 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

 <220>
 10 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged thymidine

 <400> 31
 nnnngctttg gnnn 14

 15

 <210> 32
 <211> 13
 <212> DNA
 <213> Artificial Sequence

 20

 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c

 25

 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

 30

 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c

 35

 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 40

 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

 <400> 32
 nngtgatgac nnn 13

 45

 <210> 33
 <211> 13
 <212> DNA
 <213> Artificial Sequence

 50

 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c

 55

 <220>
 <221> modified_base

<222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 5 <221> modified_base
 <222> (4)..(4)
 <223> n stands for 2-thiocarbonylthymidine

 <220>
 10 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c

 <220>
 15 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 20 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

 <400> 33
 nngngatgac nnn

 25 <210> 34
 <211> 14
 <212> DNA
 <213> Artificial Sequence

 30 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

 35 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

 40 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

 45 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c

 50 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

 55 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

 <400> 34

13

nnnatgccac cnnn

14

5 <210> 35
 <211> 14
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

15 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

20 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

25 <220>
 <221> modified_base
 <222> (6)..(6)
 <223> n stands for 8-aminoguanosine

30 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c

35 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

40 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

<400> 35
 nnnatnccac cnnn

14

45 <210> 36
 <211> 14
 <212> DNA
 <213> Artificial Sequence

50 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

55 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

<220>

<221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged m5c
 5 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 10 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged adenosine
 15 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 20 <400> 36
 nnnccgcatgc cnnn 14
 25 <210> 37
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 30 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 35 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 40 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged m5c
 45 <220>
 <221> modified_base
 <222> (9)..(9)
 <223> n stands for 8-aminoguanosine
 50 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 55 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged adenosine
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

<400> 37
nnncgcatnc cnnn

14

5 <210> 38
 <211> 14
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

15 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged adenosine

20 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

25 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c

30 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

35 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

 <400> 38
 nnnatgccct annn

14

40 <210> 39
 <211> 14
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

50 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged adenosine

55 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

<220>
 <221> modified_base
 <222> (6)..(6)
 <223> n stands for 8-aminoguanosine
 5
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c
 10
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 15
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 20
 <400> 39
 nnnatnccct annn 14
 <210> 40
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 30
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 35
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine
 40
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 45
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 50
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 55
 <400> 40
 nnntgcctcc gnnn 14

<210> 41
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 10
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 15
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine
 20
 <220>
 <221> modified_base
 <222> (5)..(5)
 <223> n stands for 8-aminoguanosine
 25
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 30
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 35
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 41
 nnntnctcc gnnn
 40
 <210> 42
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 50
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 55
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

14

5 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c

 10 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

 15 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

 <400> 42
 nnnatgccac cnnn

20 <210> 43
 <211> 14
 <212> DNA
 <213> Artificial Sequence

 25 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

 30 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

 35 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine

 40 <220>
 <221> modified_base
 <222> (6)..(6)
 <223> n stands for 8-bromoguanosine

 45 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c

 50 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

 55 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

 <400> 43
 nnnatnccac cnnn

14

14

5 <210> 44
 <211> 14
 <212> DNA
 <213> Artificial Sequence

 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 10 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

 15 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged m5c

 20 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 25 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged adenosine

 30 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

 <400> 44
 nnncgcatgc cnnn
 35

 <210> 45
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

 45 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

 50 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged m5c

 55 <220>
 <221> modified_base
 <222> (9)..(9)

14

<223> n stands for 8-bromoguanosine
 <220>
 <221> modified_base
 5 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 10 <222> (13)..(13)
 <223> n stands for 2',4'-bridged adenosine
 <220>
 <221> modified_base
 15 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 45
 nnnccgcatnc cnnn 14
 <210> 46
 20 <211> 14
 <212> DNA
 <213> Artificial Sequence
 <220>
 25 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 30 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged adenosine
 <220>
 35 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 40 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c
 <220>
 45 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 50 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 46
 nnnatgccct annn 14
 <210> 47
 55 <211> 14
 <212> DNA

<213> Artificial Sequence
 <220>
 <221> modified_base
 5 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 10 <222> (2)..(2)
 <223> n stands for 2',4'-bridged adenosine
 <220>
 <221> modified_base
 15 <222> (3)..(3)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 20 <222> (6)..(6)
 <223> n stands for 8-bromoguanosine
 <220>
 <221> modified_base
 25 <222> (12)..(12)
 <223> n stands for 2',4'-bridged m5c
 <220>
 <221> modified_base
 30 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 35 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine
 <400> 47
 nnnatnccct annn
 14
 <210> 48
 <211> 14
 <212> DNA
 40 <213> Artificial Sequence
 <220>
 <221> modified_base
 45 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine
 <220>
 <221> modified_base
 50 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 <220>
 <221> modified_base
 55 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine
 <220>
 <221> modified_base

<222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 5 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 10 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged adenosine

 <400> 48
 nnntgcctcc gnnn 14

 15

 <210> 49
 <211> 14
 <212> DNA
 <213> Artificial Sequence

 20

 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged guanosine

 25

 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

 30

 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine

 35

 <220>
 <221> modified_base
 <222> (5)..(5)
 <223> n stands for 8-bromoguanosine

 40

 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 45 <222> (13)..(13)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 <222> (14)..(14)
 50 <223> n stands for 2',4'-bridged adenosine

 <400> 49
 nnntncctcc gnnn 14

 55

 <210> 50
 <211> 14

<212> DNA
 <213> Artificial Sequence

5
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine

10
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine

15
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged guanosine

20
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

25
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

30
 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged thymidine

35
 <400> 50
 nnntgctttg gnnn

40
 <210> 51
 <211> 14
 <212> DNA
 <213> Artificial Sequence

45
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine

50
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine

55
 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged guanosine

<220>
 <221> modified_base
 <222> (10)..(10)
 <223> n stands for 8-bromoguanosine

<220>

14

<221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 5 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 10 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged thymidine
 <400> 51
 15 nnntgctttn gnnn 14
 <210> 52
 <211> 14
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine
 25 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged m5c
 30 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine
 35 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 40 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 45 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged m5c
 <400> 52
 50 nnnccaagtt cnnn 14
 <210> 53
 <211> 14
 <212> DNA
 <213> Artificial Sequence
 55 <220>

<221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine
 5 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged m5c
 10 <220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine
 15 <220>
 <221> modified_base
 <222> (8)..(8)
 <223> n stands for 8-bromoguanosine
 20 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 25 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 30 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged m5c
 <400> 53
 nnnccaantt cnnn
 35 <210> 54
 <211> 13
 <212> DNA
 <213> Artificial Sequence
 40 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c
 45 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 50 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 55 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

14

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 5
 <400> 54
 nnngtgatgac nnn 13

<210> 55
 <211> 13
 <212> DNA
 <213> Artificial Sequence

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

<220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 8-bromoguanosine

<220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

<400> 55
 nnntgatgac nnn 13

<210> 56
 <211> 13
 <212> DNA
 <213> Artificial Sequence

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine

<220>
 <221> modified_base
 <222> (5)..(5)
 <223> n stands for 8-bromoguanosine
 5

<220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 10

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 15

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 20

<400> 56
 nngtnatgac nnn 13

<210> 57
 <211> 13
 <212> DNA
 <213> Artificial Sequence
 25

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c
 30

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged thymidine
 35

<220>
 <221> modified_base
 <222> (8)..(8)
 <223> n stands for 8-bromoguanosine
 40

<220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 45

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 50

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 55

<400> 57
 nngtgatnac nnn 13

<210> 58
 <211> 13
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c
 10
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine
 15
 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 20
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine
 25
 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 <400> 58
 nccctcgcc nnn
 30
 <210> 59
 <211> 13
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged m5c
 40
 <220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged guanosine
 45
 <220>
 <221> modified_base
 <222> (8)..(8)
 <223> n stands for 8-bromoguanosine
 50
 <220>
 <221> modified_base
 <222> (11)..(11)
 <223> n stands for 2',4'-bridged m5c
 55
 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

13

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c
 5
 <400> 59
 nncctcncc nnn 13

<210> 60
 <211> 14
 <212> DNA
 <213> Artificial Sequence

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine

<220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

<220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged m5c

<400> 60
 nnnccaagtt cnnn 14

<210> 61
 <211> 14
 <212> DNA
 <213> Artificial Sequence

<220>
 <221> modified_base
 <222> (1)..(1)
 <223> n stands for 2',4'-bridged adenosine

<220>
 <221> modified_base
 <222> (2)..(2)
 <223> n stands for 2',4'-bridged m5c

<220>
 <221> modified_base
 <222> (3)..(3)
 <223> n stands for 2',4'-bridged adenosine

 <220>
 <221> modified_base
 <222> (8)..(8)
 <223> n stands for 7-phenylethynylguanosine

 <220>
 <221> modified_base
 <222> (12)..(12)
 <223> n stands for 2',4'-bridged thymidine

 <220>
 <221> modified_base
 <222> (13)..(13)
 <223> n stands for 2',4'-bridged m5c


 <220>
 <221> modified_base
 <222> (14)..(14)
 <223> n stands for 2',4'-bridged m5c

 <400> 61
 nnnccaantt cnnn

14

Claims


1. An antisense oligonucleic acid having a base length of not less than 7nt and not more than 30nt, wherein
 nucleic acid residues of not less than 1nt and not more than 5nt respectively from the both terminals are 2',4'-
 bridged nucleic acids,
 2',4'-non-bridged nucleic acid residue(s) is(are) present between the both terminals, and
 one or more bases in the nucleic acid residue(s) of the 2',4'-non-bridged nucleic acid residue(s) is/are modified.
2. The antisense oligonucleic acid according to claim 1 wherein the 2',4'-non-bridged nucleic acid residue with the
 modified base is contained in a sequence of TGC or TCC.
3. The antisense oligonucleic acid according to claim 1 wherein the modified base is 5-hydroxycytosine, 4-acetylcy-
 tosinine, 3-C₁₋₆ alkylcytosine, 5-C₁₋₆ alkylcytosine, 2-thiocytosine, 2-thiothymine, dihydrothymine, pseudo thymine,
 2-thiouridine, 5-(carboxyhydroxymethyl)uridine, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylami-
 nomethyluridine, dihydrouridine, pseudo uridine, 5-methylaminomethyluridine, 5-methylaminomethyl-2-thiouridine,
 5-methoxycarbonylmethyl-2-thiouridine, 5-methoxycarbonylmethyluridine, 5-methoxyuridine, 8-aminoguanine, 8-
 halogenoguanine, 7-deaza-7-(2-phenylethynyl)guanine, 7-deaza-7-(2-pyridylethynyl)guanine, 7-deaza-7-[2-(C₁₋₇
 alkanoyloxy-C₁₋₆ alkyl) ethynyl]guanine, 7-deaza-7-[2-(hydroxy-C₁₋₆ alkyl)ethynyl]guanine, 1-C₁₋₆ alkylguanine, 2,2-
 di(C₁₋₆ alkyl) guanine, 2-C₁₋₆ alkylguanine, 7-deaza-7-C₁₋₆ alkylguanine, 7-deaza-7-C₂₋₆ alkenylguanine, 7-deaza-
 7-C₂₋₆ alkynylguanine, 7-deaza-7-halogenoguanine, inosine, 1-C₁₋₆ alkylinosine, queuosine, β ,D-galactosylqueuo-
 sine, β ,D-mannosylqueuosine, N⁶-C₂₋₆ alkenyladenine, 1-C₁₋₆ alkyladenine, 2-C₁₋₆ alkyladenine, N⁶-C₁₋₆ alkylade-
 nine or 2-C₁₋₆ alkylthio-N⁶-C₂₋₆ alkenyladenine.
4. The antisense oligonucleic acid according to any one of claims 1 to 3 wherein the 2',4'-non-bridged nucleic acid
 residue is DNA.
5. The antisense oligonucleic acid according to any one of claims 1 to 4 wherein the 2',4'-bridged nucleic acid residue
 has a structure of any of the following formulas (I) to (III):



wherein

X¹ is O, S, a >N(R³) group, a -C(=O)-O- group or a -C(=O)-N(R⁴)- group (R³ and R⁴ are each independently H or a C₁₋₆ alkyl group),

X² is a guanidino group represented by any of the following formulas (IV) to (VII):



wherein R⁵ - R¹⁸ are each independently H, a C₁₋₆ alkyl group, a C₃₋₁₀ cycloalkyl group, an amino-protecting group or a 2-cyanoethyloxycarbonyl group,

X³ is O, S, a >N(R¹⁹) group, a -C(=O)-O- group or a -C(=O)-N(R²⁰)- group (R¹⁹ and R²⁰ are each independently H or a C₁₋₆ alkyl group),

Y¹ - Y³ are each independently O⁻ or S⁻,

Base is a nucleic acid base group,

R¹ and R² are each independently H, a C₁₋₆ alkyl group or R¹ and R² may be taken together to form a C₁₋₄ alkylene group, and

n is an integer of not less than 0 and not more than 2.

6. The antisense oligonucleic acid according to claim 5 wherein the 2',4'-bridged nucleic acid residue has a structure of the formula (I), X¹ is O and n is 1.

7. The antisense oligonucleic acid according to any one of claims 1 to 6 showing reduced hepatotoxicity compared to that before introduction of the modification into the base of the nucleic acid residue.

Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5

(2) activity evaluation



(1) hepatotoxicity test (ALT)




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17





Fig. 18

(2) activity evaluation



(1) hepatotoxicity test (ALT)




Fig. 19




Fig. 20





Fig. 21

(2) activity evaluation



(1) hepatotoxicity test (ALT)




Fig. 22




Fig. 23




Fig. 24




Fig. 25




Fig. 26




Fig. 27




Fig. 28




Fig. 29





Fig. 30

hepatotoxicity test (ALT)



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2018/006061

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl. C12N15/113 (2010.01) i, A61K31/7115 (2006.01) n,
A61K31/712 (2006.01) n, A61P43/00 (2006.01) n

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl. C12N15/113, A61K31/7115, A61K31/712, A61P43/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Published examined utility model applications of Japan 1922-1996

Published unexamined utility model applications of Japan 1971-2018

Registered utility model specifications of Japan 1996-2018

Published registered utility model applications of Japan 1994-2018

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

JSTPlus/JMEDPlus/JST7580 (JDreamIII), PubMed, CiNii,

Cplus/MEDLINE/EMBASE/BIOSIS/WPIDS (STN)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MERITS, A., "New antisense inhibitors of hepatitis C virus.", J. Antivir. Antiretrovir., 2011, vol. 3	1-7
Y	issue 4, p. 165, whole document	1-7
X	MUTSO, M. et al., "RNA Interference-Guided Targeting of Hepatitis C Virus Replication with Antisense Locked Nucleic Acid-Based Oligonucleotides Containing 8-oxo-dG Modifications.", PLOS ONE, 2015, 10(6), e0128686, doi:10.1371/journal.pone.0128686, abstract, table 1, fig. 1, 4, 7	1-7
Y		1-7



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 May 2018 (14.05.2018)

Date of mailing of the international search report

22 May 2018 (22.05.2018)

Name and mailing address of the ISA/

Japan Patent Office
3-4-3, Kasumigaseki, Chiyoda-ku,
Tokyo 100-8915, Japan

Authorized officer

Telephone No.

Form PCT/ISA/210 (second sheet) (January 2015)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2018/006061

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP 2016-096826 A (IONIS PHARMACEUTICALS INC.) 20	1-7
Y	May 2016, abstract, claim 1, paragraph [0168], fig. 1 & US 2008/0015162 A1 abstract, claims, table 20, fig. 1 & WO 2007/090071 A2 & EP 2015758 A2	1-7
X	US 2015/0051389 A1 (ISIS PHARMACEUTICALS, INC.) 19	1-7
Y	February 2015, paragraphs [0476]-[0480], [0498], example 83, tables 6, 127, 128 & WO 2013/022984 A1 & EP 2673361 A1	1-7
X	US 2016/0160280 A1 (ISIS PHARMACEUTICALS, INC.) 09	1-7
Y	June 2016, abstract, paragraphs [0304], [0430], table 5 & US 2017/0327893 A1	1-7
A	BURDICK, A. D. et al., "Sequence motifs associated with hepatotoxicity of locked nucleic acid- modified antisense oligonucleotides.", Nucleic Acids Research, 2014, vol. 42, no. 8, pp. 4882- 4891	1-7

Form PCT/ISA/210 (continuation of second sheet) (January 2015)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2011052436 A [0006]
- WO 2014046212 A [0006]
- WO 2014112463 A [0006]
- WO 2015125783 A [0006]
- WO 2016017422 A [0006]
- WO 2013118776 A [0006]
- JP 2017030489 A [0050]

Non-patent literature cited in the description

- **PETER H. HAGEDORN et al.** *Nucleic Acid Therapeutics*, 2013, vol. 23 (5), 302-310 [0007]
- **ANDREW D. BURDICK et al.** *Nucleic Acids Research*, 2014, 1-10 [0007]
- *Nucleic Acid Ther.*, 2012, vol. 22 (5), 344-359 [0059]
[0063] [0080] [0083] [0086] [0099]
- **F. SEELA et al.** *Synthesis*, 2004, vol. 8, 1203-1210 [0127]
- **M. MACKOVA et al.** *ChemBioChem*, 2015, vol. 16, 2225-2236 [0127]