

(19)



(11)

EP 3 647 398 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:

15.05.2024 Bulletin 2024/20

(21) Application number: **18203740.8**

(22) Date of filing: **31.10.2018**

(51) International Patent Classification (IPC):
C11D 3/386 ^(2006.01)

(52) Cooperative Patent Classification (CPC):
**C11D 3/38636; C11D 1/66; C11D 3/1246;
C11D 3/128; C11D 3/227; C11D 3/361;
C11D 3/3742; C11D 3/3769; C11D 3/3776;
C11D 3/378; C11D 3/3905; C11D 3/42; C11D 3/48;
C11D 3/50; C11D 17/042**

(54) **CLEANING COMPOSITIONS CONTAINING DISPERSINS V**

REINIGUNGSZUSAMMENSETZUNGEN MIT DISPERSINEN V

COMPOSITIONS DE NETTOYAGE CONTENANT DES DISPERSINES V

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(43) Date of publication of application:
06.05.2020 Bulletin 2020/19

(73) Proprietor: **Henkel AG & Co. KGaA
40589 Düsseldorf (DE)**

(72) Inventors:

- **WEIDE, Mirko
40223 Düsseldorf (DE)**
- **WIELAND, Susanne
41541 Dormagen/ Zons (DE)**
- **DENGUTH, Ulrike
45739 Oer-Erkenschwick (DE)**
- **VEJBORG, Rebecca
3450 Allerød (DK)**
- **SEGURA, Dorotea Raventos
2880 Bagsvaerd (DK)**
- **SALOMON, Jesper
2840 Holte (DK)**
- **JENSEN, Johanne M.
2880 Bagsvaerd (DK)**
- **MONRAD, Rune Nygaard
3400 Hillerød (DK)**

- **DUE, Anne Vindum
2880 Bagsvaerd (DK)**
- **GUDMAND, Martin
2880 Bagsvaerd (DK)**

(56) References cited:

**EP-A1- 0 425 019 WO-A1-2017/186936
WO-A1-2017/186937 WO-A1-2017/186943**

- **SUGAI M ET AL: "Purification of a 51 kDa endo-beta-N-acetylglucosaminidase from Staphylococcus aureus", FEMS MICROBIOLOGY LETTERS, WILEY-BLACKWELL PUBLISHING LTD, GB, vol. 61, no. 3, 15 October 1989 (1989-10-15), pages 267-272, XP023917264, ISSN: 0378-1097 [retrieved on 1989-10-15]**
- **T. WADSTRÖM ET AL: "Bacteriolytic enzymes from Staphylococcus aureus. Purification of an endo-[beta]-N-acetylglucosaminidase", BIOCHEMICAL JOURNAL, vol. 120, no. 4, 1 December 1970 (1970-12-01), pages 725-734, XP055576278, GB ISSN: 0264-6021, DOI: 10.1042/bj1200725**

Remarks:

The complete document including Reference Table(s) and the Sequence Listing(s) can be downloaded from the EPO website

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 3 647 398 B1

Description**Reference to a Sequence Listing**

[0001] This application contains a Sequence Listing in computer readable form.

Background of the Invention

[0002] The present invention relates to specific compositions, as defined herein, comprising enzymes having hexosaminidase activity comprising the motif D[IV]AR[TK] (SEQ ID NO 12) such as dispersins obtained from *Staphylococcus*. The invention further relates to methods and use of said compositions comprising such enzymes in cleaning processes e.g. for stain removal.

Description of the Related Art

[0003] Enzymes have been used in detergents for decades. Usually a cocktail of various enzymes is added to detergent compositions. The enzyme cocktail often comprises various enzymes, wherein each enzyme targets its specific substrate e.g. amylases are active towards starch stains, proteases on protein stains and so forth. Textiles surface and hard surfaces, such as dishes or the inner space of a laundry machine enduring a number of wash cycles, become soiled with many different types of soiling which may compose of proteins, grease, starch etc. One type of stain may compose of organic matter, such as cell debris, biofilm, EPS, etc. Polypeptides having hexosaminidase activity include Dispersins such as Dispersin B (DspB), which are described as β -N-acetylglucosaminidases belonging to the Glycoside Hydrolase 20 family. WO04061117 A2 (Kane Biotech INC) describe use of compositions comprising DspB for reducing and preventing biofilm caused by poly-N-acetylglucosamine-producing bacteria and Kane et al. describes the use of compositions comprising dispersins for reducing biofilm on medical devices and for wound care. The application WO9850512 (Procter and Gamble) disclose laundry or cleaning products comprising one or more hexosaminidase enzymes. The present invention provides suitable enzymes for use in detergents and for deep cleaning of items such as laundry and cleaning process.

[0004] WO 2017/186943 relates to polypeptides having hexosaminidase activity, and polynucleotides encoding the polypeptides. In particular, it relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

[0005] WO 2017/186936 relates to polypeptides having hexosaminidase activity, and polynucleotides encoding the polypeptides. In particular, it relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

[0006] WO 2017/186937 relates to polypeptides having hexosaminidase activity, and polynucleotides encoding the polypeptides. In particular, it relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

[0007] In the non-patent literature "Purification of a 51 kDa endo-beta-N-acetylglucosaminidase from *Staphylococcus aureus*" by Sugai et al. (FEMS MICROBIOLOGY LETTERS, WILEY-BLACKWELL PUBLISHING LTD, GB, vol. 61, no. 3, 15 October 1989, pages 267-272) a bacteriolytic enzyme was isolated and characterized to find that the purified enzyme acted as an endo- β -N-acetylglucosaminidase.

[0008] In the non-patent literature "Bacteriolytic enzymes from *Staphylococcus aureus*. Purification of an endo-[beta]-N-acetylglucosaminidase" by Wadström et al. (BIOCHEMICAL JOURNAL, vol. 120, no. 4, 1 December 1970, pages 725-734) it was found that on cultivation of *Staphylococcus aureus* in a complex liquid medium bacteriolytic activity is observed extracellularly. The purified enzyme (endo-beta-N-acetylglucosaminidase) was found to give a single band on polyacrylamide-gel and cellulose acetate electrophoresis and was devoid of all 14 staphylococcal enzymes and toxins assayed for.

[0009] EP 0 425 019 relates to an antimicrobial composition comprising endo-beta-N-acetylglucosaminidase and/or endoglycopeptidase, and ruminant stomach lysozyme.

Summary of the Invention

[0010] The present invention and its preferred embodiments are apparent from the appendant set of claims.

[0011] When in the following reference is made to "compositions of the invention" or "compositions as described herein", the compositions (a)-(j) recited in claim 1 are meant. Furthermore, if not indicated otherwise, all references to percentages in relation to the disclosed compositions relate to wt % relative to the total weight of the respective composition. It is understood that when reference is made to compositions that contain a hexosaminidase as defined herein, the respective composition contains at least one of said hexosaminidases but can also comprise two or more of them.

[0012] The hexosaminidase preferably has N-acetylglucosaminidase activity, preferably β -1,6 N-acetylglucosaminidase activity.

[0013] The present invention further relates to a composition, as defined herein, comprising at least 0.01 mg *Staphylococcus* hexosaminidase comprising the motif D[IV]AR[TK] (SEQ ID NO 12) and optionally a further cleaning component, wherein the cleaning component is selected from

- (a) at least one surfactant;
- (b) at least one builder; and
- (c) at least one bleach component.

[0014] The invention further relates to a method of formulating a composition, as defined herein, comprising adding a *Staphylococcus* hexosaminidase comprising the motif D[IV]AR[TK] (SEQ ID NO 12) and at least one cleaning component.

[0015] The invention further relates to a method of treating a fabric comprising;

- (a) contacting the fabric with a composition as defined herein or an aqueous solution thereof;
- (b) and optionally rinsing and drying the textile.

[0016] The invention also relates to a method for cleaning or laundering an item comprising the steps of:

- (a) exposing an item to a composition of the invention or a wash liquor comprising a composition of the invention;
- (b) completing at least one wash cycle; and
- (c) optionally rinsing the item, wherein the item is a fabric.

Figures

[0017]

(Figure 1). The shown polypeptides belong to the *Staphylococcus* clade, which is illustrated as a phylogenetic tree in figure 1. The *Staphylococcus* clade or clade of *Staphylococcus* is a group of enzymes all related to the same ancestor and share common properties. Polypeptides forming a group within the clade (a subclade) of the phylogenetic tree can also share common properties and are more closely related than other polypeptides in the clade. The composition according of the present invention comprises *Staphylococcus* hexosaminidase comprising the motif D[IV]AR[TK] (SEQ ID NO 12)

(Figure 2). Shown is an alignment of the disclosed polypeptides.

Overview of sequences of the *Staphylococcus* clade

[0018]

SEQ ID NO 1 is the DNA encoding the full-length polypeptide from *Staphylococcus cohnii* subsp.

SEQ ID NO 2 is the polypeptide derived from SEQ ID NO 1

SEQ ID NO 3 is the mature polypeptide of SEQ ID NO 2

SEQ ID NO 4 is the DNA encoding the full-length polypeptide from *Staphylococcus fleurettii*

SEQ ID NO 5 is the polypeptide derived from SEQ ID NO 4

SEQ ID NO 6 is the mature polypeptide of SEQ ID NO 5

SEQ ID NO 7 is the *Bacillus clausii* secretion signal

SEQ ID NO 8 is a His-tag sequence

SEQ ID NO 9 is the polypeptide motif GXDE

SEQ ID NO 10 is the polypeptide motif [EQ][NRSHA][YVFL][AGSTC][IVLF][EAQYN][SN]

SEQ ID NO 11 is the polypeptide motif [VLIM][LIV]G[GAV]DE[VI][PSA]

SEQ ID NO 12 is the polypeptide motif D[IV]AR[TK]

Detailed Description of the Invention

[0019] Various enzymes are applied in cleaning processes each targeting specific types of soiling such as protein, starch and grease soiling. Enzymes are standard ingredients in detergents for laundry and dish wash. The effectiveness of these commercial enzymes provides detergents which removes much of the soiling. However, organic stains such

as EPS (extracellular polymeric substance) comprised in much biofilm constitute a challenging type of soiling due to the complex nature of such organic matters. EPS is mostly composed of polysaccharides (exopolysaccharides) e.g. PNAG (poly-N-acetylglucosamine) and proteins, but include other macro-molecules such as eDNA, lipids and other organic substances. Organic stains, like biofilm or components hereof, such as PNAG may be sticky or glueing, which when present on textile, may give rise to redeposition or backstaining of soil resulting in a greying of the textile. Further, when dirty laundry items are washed together with less dirty laundry items the dirt present in the wash liquor tend to stick to organic stains e.g. biofilm or biofilm components as a result, hereof the laundry item is more "soiled" after wash than before wash. This effect may also be termed re-deposition. Another drawback of organic stains is the malodor as various malodor related molecules are often associated with organic stains such as biofilm.

[0020] The present invention relates to compositions comprising hexosaminidases comprising the motif D[IV]AR[TK] (SEQ ID NO 12) obtained from *Staphylococcus*, as defined herein, as well as the use and methods of use thereof. The terms "*Staphylococcus* hexosaminidase" and "hexosaminidase obtained from *Staphylococcus*" are used interchangeably throughout. The hexosaminidases are preferably dispersins and comprises N-acetylglucosaminidase and/or β -1,6-N-acetylglucosaminidase activity.

Polypeptides having hexosaminidase activity

[0021] Dispersin: The term "dispersin" and the abbreviation "Dsp" means a polypeptide having hexosaminidase activity, EC 3.2.1.- that catalyzes the hydrolysis of β -1,6-glycosidic linkages of N-acetyl-glucosamine polymers (poly-N-acetylglucosamine, PNAG) found e.g. in biofilm. Thus, dispersins is an enzyme having beta-1,6 N-acetylglucosaminidase activity.

[0022] Hexosaminidase: The term "hexosaminidases" means a polypeptide having hexosaminidase activity (hexosaminidases), and includes EC 3.2.1. e.g. that catalyzes the hydrolysis of N-acetyl-D-hexosamine or N-acetyl-glucosamine polymers found e.g. in biofilm. The term includes dispersins and includes polypeptides having N-acetylglucosaminidase activity and β -1,6 N-acetylglucosaminidase activity. The term "polypeptide having hexosaminidase activity" may be used interchangeably with the term hexosaminidases and similar the term "polypeptide having beta-1,6-N-acetylglucosaminidase activity" may be used interchangeably with the term beta-1,6-N-acetylglucosaminidases. For the purposes of the present invention, hexosaminidase activity is determined according to the procedure described in Assay 1.

[0023] The polypeptide useful in accordance with the invention is comprised in a specific clade of hexosaminidases. This clade is in the present context termed *Staphylococcus* as the hexosaminidases from the clade are obtained from bacteria within the taxonomic family *Staphylococcaceae*, preferably from the *Staphylococcus* genus. The composition of the present invention comprises *Staphylococcus* hexosaminidase comprising the motif D[IV]AR[TK] (SEQ ID NO 12). For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a polynucleotide is produced by the source or by a strain in which the polynucleotide from the source has been inserted. In one aspect, the polypeptide obtained from a given source is secreted extracellularly.

[0024] The phylogenetic tree of the *Staphylococcus* clade is shown in Figure 1. The polypeptides comprising in the *Staphylococcus* clade, which finds use in cleaning processes and compositions of the invention are listed in the table below. The hexosaminidases of Table 1 have 1,6 N-acetylglucosaminidase activity and are thus dispersins. The dispersins of this group have been found to be particularly useful in cleaning of organic stains e.g. PNAG from textiles. In particular, dispersins of Table 1 may be formulated in cleaning composition, comprising a dispersin obtained from *Staphylococcus* and a detergent adjunct. According to the present invention, the *Staphylococcus* hexosaminidase comprises the motif D[IV]AR[TK] (SEQ ID NO 12). The compositions of the invention are useful in cleaning processes such as laundry.

Table 1 The list of hexosaminidase polypeptides having beta-1,6 N-acetylglucosaminidase activity comprised in the *Staphylococcus* clade

SEQ_ID_NO 3	<i>Staphylococcus cohnii</i> subsp. <i>cohnii</i>
SEQ_ID_NO 6	<i>Staphylococcus fleurettii</i>
UniProtKB/TrEMBL	Additional dispersins
A8DMZ4	<i>Staphylococcus lugdunensis</i>
A8DN10	<i>Staphylococcus lugdunensis</i>
E8SGX9	<i>Staphylococcus pseudintermedius</i>
E6MC86	<i>Staphylococcus lugdunensis</i> M23590
F0P5N7	<i>Staphylococcus pseudintermedius</i>

(continued)

	H3WVE1	<i>Staphylococcus lugdunensis</i> VCU139
5	A0A1Q9II29	<i>Staphylococcus pseudintermedius</i>
	A0A038HZE1	<i>Staphylococcus lugdunensis</i> VCU150
	A0A0M2NYH6	<i>Staphylococcus cohnii</i> subsp. <i>cohnii</i>
	A0A0M2NR77	<i>Staphylococcus pasteurii</i>
10	A0A1F2JPF3	<i>Staphylococcus</i> sp. HMSC13A10
	A0A0X8QI92	<i>Staphylococcus lugdunensis</i>
	A0A178NA26	<i>Staphylococcus cohnii</i>
15	A0A178NED3	<i>Staphylococcus cohnii</i>
	A0A1F1MQ36	<i>Staphylococcus</i> sp. HMSC057A08
	A0A1E8V1N3	<i>Staphylococcus</i> sp. HMSC065E07
	A0A1J4HXT8	<i>Staphylococcus</i> sp. HMSC62D11
20	A0A1J6DK65	<i>Staphylococcus cohnii</i>
	A0A1J6DB75	<i>Staphylococcus cohnii</i>
	A0A1J6CW64	<i>Staphylococcus cohnii</i>
25	A0A1J6CJE7	<i>Staphylococcus cohnii</i>
	A0A1J6CYC9	<i>Staphylococcus cohnii</i>
	A0A1J6CTS7	<i>Staphylococcus cohnii</i>

[0025] The Glyco_hydro_20 domain includes the polypeptides having hexosaminidase, preferably beta-1,6 N-acetylglucosaminidase e.g. PNAG activity, these polypeptides are comprised in three specific clades which are the ENYA, VLG and/or DIARK clades as described below and in example 5 and figure 1.

[0026] The polypeptide sequences containing a Glyco_hydro_20 domain comprises several motifs; one example is GXDE (SEQ ID NO 9), situated in positions 157 to 160 in *Staphylococcus cohnii* subsp. *cohnii* (SEQ ID NO 3). Residues D and E are the key catalytic residues of Glyco_hydro_20 enzymes (position 159 to 160 in SEQ ID NO 3). The composition of the present invention comprises *Staphylococcus* hexosaminidase comprising the motif D[IV]AR[TK] (SEQ ID NO 12).

[0027] The hexosaminidases e.g. the dispersins useful in the compositions of the invention may be divided into clades or domain groups characterized by having various motifs. The composition of the present invention comprises *Staphylococcus* hexosaminidase comprising the motif D[IV]AR[TK] (SEQ ID NO 12). One clade (ENYA), was identified. This clade has not been described previously. The clade is termed IES and polypeptides of this clade comprises Glyco_hydro_20 domain polypeptides of bacterial origin and are in addition to having beta-1,6 N-acetylglucosaminidase and PNAG activity, characterized by comprising certain motifs. The polypeptides of the clade comprise the motif example [EQ][NRSHA][YVFL][AGSTC][IVLF][EAQYN][SN] (SEQ ID NO 10), corresponding to ENYAIES at position 44 to 50 of SEQ ID NO 3.

[0028] Another clade was identified. This clade has not been described previously. The clade is termed VLG and polypeptides of this clade comprise Glyco_hydro_20 domain polypeptides of bacterial origin and are in addition to having beta-1,6 N-acetylglucosaminidase and PNAG activity, characterized by comprising certain motifs. The polypeptides of the clade comprise the motif example [VIMS][LIV]G[GAV]DE[VI][PSA] (SEQ ID NO 11), corresponding to VLGGEVVP (positions 155 to 162 of SEQ ID NO 3), where G and DE (corresponding to positions 157 and 159-160 of SEQ ID NO 3) are fully conserved in VLG clade and part of the active site. Residues D and E are the key catalytic residues of Glyco_hydro_20 enzymes (position 159 to 160 in SEQ ID NO 3).

[0029] Yet another clade termed DIARK comprises the hexosaminidases e.g. dispersins useful in the compositions of the invention. The polypeptides of the clade comprise the motif example D[IV]AR[TK] (SEQ ID NO 12), corresponding to pos 10 to 14 of SEQ ID NO 3, where D and AR are fully conserved in DIARK clade (positions 10 and 12-13 in SEQ ID NO 3). According to the present invention, the *Staphylococcus* hexosaminidases e.g. dispersins, comprise the motif D[IV]AR[TK] (SEQ ID NO 12).

[0030] In one aspect of the invention the hexosaminidase e.g. dispersin comprises one or more of the following motifs GXDE (SEQ ID NO 9), [EQ][NRSHA][YVFL][AGSTC][IVLF][EAQYN][SN] (SEQ ID NO 10), or

[VLIM][LIV]G[GAV]DE[VI][PSA] (SEQ ID NO 11), in addition to D[IV]AR[TK] (SEQ ID NO 12). In one aspect, the hexosaminidase e.g. dispersin comprises all four motifs GXDE (SEQ ID NO 9), [EQ][NRSHA][YVFL][AGSTC][IV-LF][EAQYN][SN] (SEQ ID NO 10), [VLIM][LIV]G[GAV]DE[VI][PSA] (SEQ ID NO 11), and D[IV]AR[TK] (SEQ ID NO 12). In one aspect, the hexosaminidase e.g. dispersin comprises the three motifs GXDE (SEQ ID NO 9), [EQ][NRSHA][YV-FL][AGSTC][IVLF][EAQYN][SN] (SEQ ID NO 10) and D[IV]AR[TK] (SEQ ID NO 12).

[0031] An alignment of the polypeptides useful in the compositions of the invention is shown in Figure 2. A phylogenetic tree of the polypeptides useful in the compositions of the invention is shown in Figure 1.

[0032] A polypeptide useful in the compositions of the present invention preferably has a sequence identity to the mature polypeptide sequence shown in SEQ ID NO: 3 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, wherein the polypeptide has hexosaminidase, preferably 1,6 N-acetylglucosaminidase activity. In one aspect, the polypeptide differs by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the polypeptide shown in SEQ ID NO: 3 and preferably has beta-1,6 N-acetylglucosaminidase activity.

[0033] A polypeptide useful in the compositions of the present invention preferably has a sequence identity to the mature polypeptide sequence shown in SEQ ID NO: 6 of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, wherein the polypeptide has hexosaminidase, preferably 1,6 N-acetylglucosaminidase activity. In one aspect, the polypeptide differs by up to 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10, from the polypeptide shown in SEQ ID NO: 6 and preferably has beta-1,6 N-acetylglucosaminidase activity.

[0034] The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "sequence identity".

[0035] For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends Genet. 16: 276-277), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment}).$$

[0036] Essential amino acids in a polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, Science 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant molecules are tested for hexosaminidase activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., 1996, J. Biol. Chem. 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., 1992, Science 255: 306-312; Smith et al., 1992, J. Mol. Biol. 224: 899-904; Wlodaver et al., 1992, FEBS Lett. 309: 59-64. The identity of essential amino acids can also be inferred from an alignment with a related polypeptide

Compositions

[0037] The invention relates to compositions, as defined herein, comprising *Staphylococcus* hexosaminidases comprising the motif D[IV]AR[TK] (SEQ ID NO 12), preferably dispersins, as well as uses and methods of use thereof

Liquid formulations

[0038] In one aspect the cleaning composition is a liquid composition, as defined herein. The hexosaminidase may be formulated as a liquid enzyme formulation, which is generally a pourable composition, though it may also have a high viscosity. The physical appearance and properties of a liquid enzyme formulation may vary a lot - for example, they may have different viscosities (gel to water-like), be colored, not colored, clear, hazy, and even with solid particles like in slurries and suspensions. The minimum ingredients are the enzyme(s) and a solvent system to make it a liquid.

[0039] The solvent system may comprise water, polyols (such as glycerol, (mono, di, or tri) propylene glycol, sugar alcohol (e.g. sorbitol), polypropylene glycol, and/or polyethylene glycol), ethanol, sugars, and salts. Usually the solvent system also includes a preservation agent and/or other stabilizers.

[0040] A liquid enzyme formulation may be prepared by mixing a solvent system and an enzyme concentrate with a desired degree of purity (or enzyme particles to obtain a slurry/suspension).

[0041] In an embodiment, the liquid enzyme composition comprises

- (a) at least 0.01% w/w active enzyme protein,
- (b) at least 0.5% w/w polyol,
- (c) water, and
- (d) optionally a preservation agent.

[0042] The hexosaminidases e.g. dispersins in the liquid composition of the invention may be stabilized using conventional stabilizing agents, e.g. a polyol such as propylene glycol or glycerol, ethylene glycol, polyethylene glycol, sugar alcohols, sorbitol, mannitol, erythritol, dulcitol, inositol, xylitol and adonitol.

[0043] One embodiment of the invention relates to a composition comprising a *Staphylococcus* hexosaminidase, as defined herein, wherein the composition further comprises;

(a)

- i. one or more polyol(s), preferably selected from glycerol, (mono, di, or tri) propylene glycol, ethylene glycol, polyethylene glycol, sugar alcohols, sorbitol, mannitol, erythritol, dulcitol, inositol, xylitol and adonitol,
- ii. optionally one or more enzyme, preferably selected from proteases, amylases or lipases,
- iii. optionally one or more surfactant, preferably selected from anionic and nonionic surfactants, or
- iv. optionally one or more polymer.

[0044] Another preferred embodiment relates to a composition comprising a *Staphylococcus* hexosaminidase, as defined herein, wherein the composition further comprises;

(a)

- i. one or more polyol(s), preferably selected from glycerol, (mono, di, or tri) propylene glycol, ethylene glycol, polyethylene glycol, sugar alcohols, sorbitol, mannitol, erythritol, dulcitol, inositol, xylitol and adonitol,
- ii. optionally one or more enzyme, preferably selected from proteases, amylases or lipases,
- iii. optionally one or more surfactant, preferably selected from anionic and nonionic surfactants, or
- iv. optionally one or more polymer; wherein the hexosaminidase has N-acetylglucosaminidase activity, preferably β -1,6 N-acetylglucosaminidase activity.

[0045] One preferred aspect relates to a composition comprising a *Staphylococcus* hexosaminidase, e.g. dispersin, as defined herein, wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ ID NO 3, SEQ ID NO 6 or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto and wherein the composition optionally further comprises;

(a)

- i. one or more polyol(s), preferably selected from glycerol, (mono, di, or tri) propylene glycol, ethylene glycol, polyethylene glycol, sugar alcohols, sorbitol, mannitol, erythritol, dulcitol, inositol, xylitol and adonitol,
- ii. optionally one or more enzyme, preferably selected from proteases, amylases or lipases,
- iii. optionally one or more surfactant, preferably selected from anionic and nonionic surfactants, or
- iv. optionally one or more polymer; wherein the hexosaminidase has N-acetylglucosaminidase activity, preferably β -1,6 N-acetylglucosaminidase activity.

[0046] One preferred aspect relates to a composition comprising a *Staphylococcus* hexosaminidase, e.g. dispersin, as defined herein, wherein the *Staphylococcus* hexosaminidase is selected from the group shown in Table 1 comprising the motif D[IV]AR[TK] (SEQ ID NO 12) and wherein the composition optionally further comprises;

(a)

- i. one or more polyol(s), preferably selected from glycerol, (mono, di, or tri) propylene glycol, ethylene glycol, polyethylene glycol, sugar alcohols, sorbitol, mannitol, erythritol, dulcitol, inositol, xylitol and adonitol,

- ii. optionally one or more enzyme, preferably selected from proteases, amylases or lipases,
- iii. optionally one or more surfactant, preferably selected from anionic and nonionic surfactants, or
- iv. optionally one or more polymer; wherein the hexosaminidase has N-acetylglucosaminidase activity, preferably β -1,6 N-acetylglucosaminidase activity.

Granular formulations

[0047] Non-dusting granulates may be produced, e.g. as disclosed in US 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are polyethylene oxide products (polyethyleneglycol, PEG) with mean molar weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591

The *Staphylococcus* hexosaminidase for use in the compositions of the invention may be formulated as a granule for example as a co-granule that combines one or more enzymes or benefit agents such as MnTACN. Each enzyme will then be present in more granules securing a more uniform distribution of enzymes in the detergent. This also reduces the physical segregation of different enzymes due to different particle sizes. Methods for producing multi-enzyme co-granulate for the detergent industry is disclosed in the IP.com disclosure IPCOM000200739D.

[0048] Another example of formulation of enzymes by the use of co-granulates are disclosed in WO 2013/188331, which relates to a detergent composition comprising (a) a multi-enzyme co-granule; (b) less than 10 wt zeolite (anhydrous basis); and (c) less than 10 wt phosphate salt (anhydrous basis), wherein said enzyme co-granule comprises from 10 to 98 wt% moisture sink components and the composition additionally comprises from 20 to 80 wt% detergent moisture sink components. WO 2013/188331 also relates to a method of treating and/or cleaning a surface, preferably a fabric surface comprising the steps of (i) contacting said surface with the detergent composition as claimed and described herein in aqueous wash liquor, (ii) rinsing and/or drying the surface.

[0049] An embodiment of the invention relates to compositions, as defined herein, comprising an enzyme granule/particle comprising a *Staphylococcus* hexosaminidase as described herein. The granule is composed of a core, and optionally one or more coatings (outer layers) surrounding the core. Typically, the granule/particle size, measured as equivalent spherical diameter (volume based average particle size), of the granule is 20-2000 μm , particularly 50-1500 μm , 100-1500 μm or 250-1200 μm . The core may include additional materials such as fillers, fibre materials (cellulose or synthetic fibres), stabilizing agents, solubilising agents, suspension agents, viscosity regulating agents, light spheres, plasticizers, salts, lubricants and fragrances. The core may include binders, such as synthetic polymer, wax, fat, or carbohydrate. The core may comprise a salt of a multivalent cation, a reducing agent, an antioxidant, a peroxide decomposing catalyst and/or an acidic buffer component, typically as a homogenous blend. The core may consist of an inert particle with the enzyme absorbed into it, or applied onto the surface, e.g., by fluid bed coating. The core may have a diameter of 20-2000 μm , particularly 50-1500 μm , 100-1500 μm or 250-1200 μm . The core can be prepared by granulating a blend of the ingredients, e.g., by a method comprising granulation techniques such as crystallization, precipitation, pan-coating, fluid bed coating, fluid bed agglomeration, rotary atomization, extrusion, prilling, spheronization, size reduction methods, drum granulation, and/or high shear granulation. Methods for preparing the core can be found in Handbook of Powder Technology; Particle size enlargement by C. E. Capes; Volume 1; 1980; Elsevier. The core of the enzyme granule/particle may be surrounded by at least one coating, e.g., to improve the storage stability, to reduce dust formation during handling, or for coloring the granule. The optional coating(s) may include a salt coating, or other suitable coating materials, such as polyethylene glycol (PEG), methyl hydroxy-propyl cellulose (MHPC) and polyvinyl alcohol (PVA). Examples of enzyme granules with multiple coatings are shown in WO 93/07263 and WO 97/23606.

[0050] The coating may be applied in an amount of at least 0.1% by weight of the core, e.g., at least 0.5%, 1% or 5%. The amount may be at most 100%, 70%, 50%, 40% or 30%.

[0051] The coating is preferably at least 0.1 μm thick, particularly at least 0.5 μm , at least 1 μm or at least 5 μm . In a one embodiment, the thickness of the coating is below 100 μm . In a more particular embodiment the thickness of the coating is below 60 μm . In an even more particular embodiment the total thickness of the coating is below 40 μm . The coating should encapsulate the core unit by forming a substantially continuous layer. A substantially continuous layer is to be understood as a coating having few or no holes, so that the core unit it is encapsulating/enclosing has few or none uncoated areas. The layer or coating should in preferably be homogeneous in thickness. The coating can further contain other materials as known in the art, e.g., fillers, antisticking agents, pigments, dyes, plasticizers and/or binders, such as titanium dioxide, kaolin, calcium carbonate or talc. A salt coating may comprise at least 60% by weight w/w of a salt, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 99% by weight w/w. The salt may be added from a salt solution where the salt is completely dissolved or from a salt suspension wherein the fine particles is less than 50 μm , such as less than 10 μm or less than 5 μm . The salt coating may comprise a single salt or a mixture of two or more salts. The salt may be water soluble, preferably having a solubility

at least 0.1 grams in 100 g of water at 20°C, preferably at least 0.5 g per 100 g water, e.g., at least 1 g per 100 g water, e.g., at least 5 g per 100 g water.

[0052] The salt may be an inorganic salt, e.g., salts of sulfate, sulfite, phosphate, phosphonate, nitrate, chloride or carbonate or salts of simple organic acids (less than 10 carbon atoms, e.g., 6 or less carbon atoms) such as citrate, malonate or acetate. Examples of cations in these salts are alkali or earth alkali metal ions, the ammonium ion or metal ions of the first transition series, such as sodium, potassium, magnesium, calcium, zinc or aluminium. Examples of anions include chloride, bromide, iodide, sulfate, sulfite, bisulfite, thiosulfate, phosphate, monobasic phosphate, dibasic phosphate, hypophosphite, dihydrogen pyrophosphate, tetraborate, borate, carbonate, bicarbonate, metasilicate, citrate, malate, maleate, malonate, succinate, lactate, formate, acetate, butyrate, propionate, benzoate, tartrate, ascorbate or gluconate. Preferably, alkali- or earth alkali metal salts of sulfate, sulfite, phosphate, phosphonate, nitrate, chloride or carbonate or salts of simple organic acids such as citrate, malonate or acetate may be used.

[0053] The salt in the coating may have a constant humidity at 20°C above 60%, particularly above 70%, above 80% or above 85%, or it may be another hydrate form of such a salt (e.g., anhydrate). The salt coating may be as described in WO 00/01793 or WO 2006/034710.

[0054] Specific examples of suitable salts are NaCl (CH₂₀°C=76%), Na₂CO₃ (CH₂₀°C=92%), NaNO₃ (CH₂₀°C=73%), Na₂HPO₄ (CH₂₀°C=95%), Na₃PO₄ (CH₂₅°C=92%), NH₄Cl (CH₂₀°C = 79.5%), (NH₄)₂HPO₄ (CH₂₀°C = 93,0%), NH₄H₂PO₄ (CH₂₀°C = 93.1%), (NH₄)₂SO₄ (CH₂₀°C=81.1%), KCl (CH₂₀°C=85%), K₂HPO₄ (CH₂₀°C=92%), KH₂PO₄ (CH₂₀°C=96.5%), KNO₃ (CH₂₀°C=93.5%), Na₂SO₄ (CH₂₀°C=93%), K₂SO₄ (CH₂₀°C=98%), KHSO₄ (CH₂₀°C=86%), MgSO₄ (CH₂₀°C=90%), ZnSO₄ (CH₂₀°C=90%) and sodium citrate (CH₂₅°C=86%). Other examples include NaH₂PO₄, (NH₄)H₂PO₄, CuSO₄, Mg(NO₃)₂ and magnesium acetate.

[0055] The salt may be in anhydrous form, or it may be a hydrated salt, i.e. a crystalline salt hydrate with bound water(s) of crystallization, such as described in WO 99/32595. Specific examples include anhydrous sodium sulfate (Na₂SO₄), anhydrous magnesium sulfate (MgSO₄), magnesium sulfate heptahydrate (MgSO₄.7H₂O), zinc sulfate heptahydrate (ZnSO₄.7H₂O), sodium phosphate dibasic heptahydrate (Na₂HPO₄.7H₂O), magnesium nitrate hexahydrate (Mg(NO₃)₂(6H₂O)), sodium citrate dihydrate and magnesium acetate tetrahydrate. Preferably the salt is applied as a solution of the salt, e.g., using a fluid bed.

[0056] In one aspect, the present invention provides a composition, as defined herein, comprising a granule, which comprises:

- (a) a core comprising a *Staphylococcus* hexosaminidase, e.g dispersin according to the invention, and
- (b) optionally a coating consisting of one or more layer(s) surrounding the core.

[0057] One aspect of the invention relates to a composition, as defined herein, comprising a granule, which comprises:

- (a) a core comprising a *Staphylococcus* hexosaminidase e.g. dispersin, wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ ID NO 3, SEQ ID NO 6 or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto, and
- (b) optionally a coating consisting of one or more layer(s) surrounding the core.

[0058] One aspect of the invention relates to a composition, as defined herein, comprising a granule, which comprises:

- (a) a core comprising a *Staphylococcus* hexosaminidase selected from the group shown in Table 1 comprising the motif D[IV]AR[TK] (SEQ ID NO 12), and
- (b) optionally a coating consisting of one or more layer(s) surrounding the core.

[0059] Another aspect relates to a composition, as defined herein, comprising a layered granule comprising

- (a) a (non-enzymatic) core;
- (b) a coating surrounding the core, wherein the coating comprises *Staphylococcus* hexosaminidase, e.g. dispersin, according to the invention; and
- (c) optionally a protective salt coating surrounding the enzyme containing coating.

[0060] Another aspect relates to a composition, as defined herein, comprising a layered granule comprising

- (a) a (non-enzymatic) core;
- (b) a coating surrounding the core, wherein the coating comprises *Staphylococcus* hexosaminidase e.g. dispersin, wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ

ID NO 3, SEQ ID NO 6 or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto; and
(c) optionally a protective salt coating surrounding the enzyme containing coating.

[0061] Another aspect relates to a composition, as defined herein, comprising a layered granule comprising:

- (a) a (non-enzymatic) core;
- (b) a coating surrounding the core, wherein the coating comprises a *Staphylococcus* hexosaminidase selected from the group shown in Table 1 comprising the motif D[IV]AR[TK] (SEQ ID NO 12); and
- (c) optionally a protective salt coating surrounding the enzyme containing coating.

Cleaning compositions

[0062] The compositions of the invention are cleaning compositions comprising a *Staphylococcus* hexosaminidase e.g. dispersin comprising the motif D[IV]AR[TK] (SEQ ID NO 12) in combination with one or more additional cleaning composition components, as defined herein. The choice of additional components is within the skill of the artisan and includes conventional ingredients, including the exemplary non-limiting components set forth below.

[0063] One aspect of the invention relates to a composition, as defined herein, comprising;

- a) at least 0.01 mg/mL of at least one *Staphylococcus* hexosaminidase, e.g. dispersin, according to the invention; and
- b) optionally at least one additional cleaning composition component, preferably selected from surfactants, builders, bleach components, polymers, dispersing agents and additional enzymes.

[0064] One aspect of the invention relates to a composition, as defined herein, comprising;

- a) at least 0.01 mg/mL of at least one *Staphylococcus* hexosaminidase, e.g. dispersin, wherein the *Staphylococcus* hexosaminidase is selected from the group shown in Table 1 comprising the motif D[IV]AR[TK] (SEQ ID NO 12); and
- b) optionally at least one additional cleaning composition component, preferably selected from surfactants, builders, bleach components, polymers, dispersing agents and additional enzymes.

[0065] One aspect of the invention relates to a composition, as defined herein, comprising;

- a) at least 0.01 mg/mL of at least one *Staphylococcus* hexosaminidase, e.g. dispersin, wherein the *Staphylococcus* hexosaminidase is selected from polypeptides having at least 80% sequence identity to the polypeptides shown in SEQ ID NO 3 or 6; and
- b) optionally at least one additional cleaning composition component, preferably selected from surfactants, builders, bleach components, polymers, dispersing agents and additional enzymes.

[0066] The *Staphylococcus* hexosaminidase comprising the motif D[IV]AR[TK] (SEQ ID NO 12) may be included in the compositions e.g. cleaning e.g. detergent composition of the present invention at a level of at least 0.0001 to at least 100, at least 0.001 to at least 100, at least 0.01 to at least 100, at least 0.02 to at least 100, at least 0.01 to at least 100, at least 0.1 to at least 100, at least 0.2 to at least 100, at least 0.5 to at least 100 mg/mL, preferably, the concentration of said *Staphylococcus* hexosaminidase enzyme in the cleaning composition e.g. detergent is in the range 0.01 to 100, 0.1 to 50 or 1 to 10 mg/mL. Thus, the detergent composition may comprise at least 0.00008%, preferably at least 0.002%, 0.003%, 0.004%, 0.005%, 0.006%, 0.008%, 0.01%, 0.02%, 0.03%, 0.05%, 0.1%, 0.2%, 0.3%, 0.4%, 0.6%, 0.7%, 0.8%, 0.9% or 1.0% *Staphylococcus* hexosaminidase comprising the motif D[IV]AR[TK] (SEQ ID NO 12).

[0067] The choice of composition components for liquid and granular compositions and of cleaning components for cleaning composition as described above may include, any of the components mentioned below.


[0068] The cleaning compositions of the invention may comprise one or more surfactants, which may be anionic and/or cationic and/or non-ionic and/or semi-polar and/or zwitterionic, or a mixture thereof. In a particular embodiment, the detergent composition includes a mixture of one or more nonionic surfactants and one or more anionic surfactants. The surfactant(s) is typically present at a level of from about 0.1% to 60% by weight, such as about 1% to about 40%, or about 3% to about 20%, or about 3% to about 10%. The surfactant(s) is chosen based on the desired cleaning application, and may include any conventional surfactant(s) known in the art.

[0069] The compositions as defined herein, if not indicated otherwise, may comprise from 0-65 wt %, from about 2 wt % to about 60 wt %, from about 5 wt % to about 50 wt %, from about 5 wt % to about 40 wt %, from about 5 wt % to about 30 wt %, from about 5 wt % to about 20 wt %, from about 5 wt % to about 10 wt % anionic surfactants, amphoteric and/or non-ionic surfactants. "About", as used herein in relation to a numerical value means said value $\pm 10\%$, preferably

$\pm 5\%$. "About 5 wt %" thus means from 4.5 to 5.5 wt %, preferably from 4.75 to 5.25 wt %.

[0070] If not indicated otherwise, the surfactant may be generally selected among nonionic, anionic and/or amphoteric surfactants. In general, bleach-stable surfactants are preferred. Preferred anionic surfactants are sulphate surfactants and in particular alkyl ether sulphates, especially C9-C15 alcohol ether sulfates, preferably ethoxylates or mixed ethoxylates/propoxylates, such as those with 1 to 30 EO, C12-C15 primary alcohol ethoxylate, , such as those with 1 to 30 EO, C8-C16 ester sulphates and C10-C14 ester sulphates, such as mono dodecyl ester sulphates. Non-limiting examples of anionic surfactants include sulfates and sulfonates, in particular, linear alkylbenzenesulfonates (LAS), in particular C12-C13 alkyl benzene sulfonates, isomers of LAS, branched alkylbenzenesulfonates (BABS), phenylalkanesulfonates, alpha-olefinsulfonates (AOS), olefin sulfonates, alkene sulfonates, alkane-2,3-diylbis(sulfates), hydroxyalkanesulfonates and disulfonates, alkyl sulfates (AS) such as sodium dodecyl sulfate (SDS), fatty alcohol sulfates (FAS), primary alcohol sulfates (PAS), alcohol ether sulfates (AES or AEOS or FES, also known as alcohol ethoxysulfates or fatty alcohol ether sulfates), secondary alkanesulfonates (SAS), paraffin sulfonates (PS), ester sulfonates, sulfonated fatty acid glycerol esters, alpha-sulfo fatty acid methyl esters (alpha-SFMe or SES) including methyl ester sulfonate (MES), alkyl- or alkenylsuccinic acid, dodecenyl/tetradecenyl succinic acid (DTSA), fatty acid derivatives of amino acids, diesters and monoesters of sulfo-succinic acid or salt of fatty acids (soap), and combinations thereof. The anionic surfactants are preferably added to the detergent in the form of salts. Suitable cations in these salts are alkali metal ions, such as sodium, potassium and lithium and ammonium salts, for example (2-hydroxyethyl) ammonium, bis(2-hydroxyethyl) ammonium and tris(2-hydroxyethyl) ammonium salts. Non-limiting examples of nonionic surfactants include alcohol ethoxylates (AE or AEO), alcohol propoxylates, propoxylated fatty alcohols (PFA), alkoxyated fatty acid alkyl esters, such as ethoxylated and/or propoxylated fatty acid alkyl esters, alkylphenol ethoxylates (APE), nonylphenol ethoxylates (NPE), alkylpolyglycosides (APG), alkoxyated amines, fatty acid monoethanolamides (FAM), fatty acid diethanolamides (FADA), ethoxylated fatty acid monoethanolamides (EFAM), propoxylated fatty acid monoethanolamides (PFAM), polyhydroxyalkyl fatty acid amides, or N-acyl N-alkyl derivatives of glucosamine (glucamides, GA, or fatty acid glucamides, FAGA), as well as products available under the trade names SPAN and TWEEN, and combinations thereof. Commercially available nonionic surfactants include Plurafac[™], Lutensol[™] and Pluronic[™] range from BASF, Dehypon[™] series from Cognis and Genapol[™] series from Clariant.


[0071] In various embodiments, said surfactant preferably comprises at least one alkyl ether sulfate. Preferred alkyl ether sulfates are those of formula (I)



[0072] In formula (I) R^1 represents a linear or branched, substituted or unsubstituted alkyl group, preferably a linear, unsubstituted alkyl group, more preferably a fatty alcohol moiety. Preferred R^1 moieties are selected from the group consisting of decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl moieties and mixtures thereof, wherein those groups with an even number of carbon atoms are preferred. Particularly preferred R^1 moieties are derived from C₁₀-C₁₈ fatty alcohols, such as those derived from coconut oil alcohols, tallow fatty alcohols, lauryl, myristyl, cetyl or stearyl alcohol or from C₁₀-C₂₀ oxoalcohols.

[0073] AO represents an ethyleneoxide (EO) or propyleneoxide (PO) group, preferably an ethyleneoxide group. The index n represents an integer from 1 to 50, preferably from 1 to 20 and more preferably from 1 to 10. Particularly preferably, n is 1, 2, 3, 4, 5, 6, 7 or 8. X represents a monovalent cation or the n-th part of an n-valent cation, preferred are alkali metal cations, specifically Na⁺ and K⁺, most preferably Na⁺. Further cations X⁺ may be selected from NH₄⁺, 1/2 Zn²⁺, 1/2 Mg²⁺, 1/2 Ca²⁺, 1/2 Mn²⁺, and combinations thereof.


[0074] In various preferred embodiments, the detergent compositions comprise an alkyl ether sulfate selected from fatty alcohol ether sulfates of formula (II)



wherein k = 9 to 19, and n = 1, 2, 3, 4, 5, 6, 7 or 8. Preferred are C₁₀₋₁₆ fatty alcohol ether sulfates with 1-7 EO (k = 9-15, n = 1-7), such as the C₁₂₋₁₄ fatty alcohol ether sulfates with 1-3, particularly 2 EO (k = 11-13, n = 1-3 or 2), more particularly the sodium salts thereof. One specific embodiment thereof is lauryl ether sulfate sodium salt with 2 EO. The level of ethoxylation is an average value and can, for a specific compound, be an integer or fractional number.


[0075] In various embodiments, the surfactant comprises at least one alkyl benzene sulfonate. Said alkyl benzene sulfonate may be present alternatively to the above alkyl ether sulfate or, preferably, in addition to it.

[0076] Exemplary alkyl benzene sulfonates include, but are not limited to linear and branched alkyl benzene sulfonates, preferably linear alkyl benzene sulfonates. Exemplary compounds are those of formula (III)



wherein R' and R'' are independently H or alkyl and combined comprise 9 to 19, preferably 9 to 15 and more preferably 9 to 13 carbon atoms. Particularly preferred are dodecyl and tridecyl benzene sulfonates, in particular the sodium salts thereof.

[0077] In addition or alternatively, the compositions of the invention may further comprise one or more nonionic surfactants. Preferred nonionic surfactants are those of formula (IV)




wherein R² represents a linear or branched substituted or unsubstituted alkyl moiety, AO represents an ethylene oxide (EO) or propylene oxide (PO) group and m is an integer from 1 to 50.

[0078] In formula (IV) R² preferably represents a linear or branched, substituted or unsubstituted alkyl group, preferably a linear, unsubstituted alkyl group, particularly preferred a fatty alcohol group. Preferred groups are R² are selected from decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, octadecyl, nonadecyl, eicosyl groups and combinations thereof, wherein those groups with an even number of carbon atoms are preferred. Particularly preferred are R² groups derived from C₁₂-C₁₈ fatty alcohols, such as coconut oil alcohol, tallow oil alcohol, lauryl, myristyl, cetyl or stearyl alcohol or from C₁₀-C₂₀ oxoalcohols.

[0079] AO represents an ethyleneoxide (EO) or propyleneoxide (PO) group, preferably an ethyleneoxide group. The index m represents an integer from 1 to 50, preferably from 1 to 20 and more preferably from 1 to 6. Particularly preferably, m is 1, 2, 3, 4 or 5, most preferably 3-5, as higher degrees of ethoxylation may negatively influence viscosity and stability.

[0080] In various preferred embodiments, the detergent compositions comprise an alkyl ether selected from fatty alcohol ethers of formula (V)



wherein k = 11 to 19, m = 1, 2, 3, 4, 5, 6, 7 or 8. Preferred are C₁₂₋₁₈ fatty alcohols with 1-6 EO (k = 11-17, m = 1-5 in formula (V)). More preferred are C₁₂₋₁₄ alcohols having 1-5 EO, most preferred are C₁₂₋₁₄ alkyl ethers with 3-5 EO, in particular lauryl ether with 5 EO.

[0081] The detergent compositions may further include other nonionic surfactants, such as alkyl glucosides of the general formula RO(G)_x, where R is a primary linear or 2-methyl-branched aliphatic radical containing 8 to 22 and preferably 12 to 18 carbon atoms and G stands for a glucose unit. The degree of oligomerization x, which indicates the distribution of monoglucosides and oligoglucosides, is a number of 1 to 10 and preferably a number of 1.2 to 1.4.

[0082] In various embodiments, the composition comprises at least two anionic surfactants, e.g. at least one alkyl ether sulfate and preferably at least one alkyl benzene sulfonate, and optionally an alkyl ether.

[0083] Suitable amphoteric surfactants comprise betaines. Preferred betaines are the alkylbetaines, the alkylamido-betaines, the imidazolinium betaines, the sulfobetaines (INCI Sultaines) and the phosphobetaines. Examples of suitable betaines and sulfobetaines are the following compounds designated as INCI: almondamidopropyl betaines, apricotamidopropyl betaines, avocadamidopropyl betaines, babassuamidopropyl betaines, behenamide idopropyl betaines, behenyl betaines, betaines, canola idopropyl betaines, caprylic / capram idopropyl betaines, carnitines, cetyl betaines, Cocamidoehtyl betaines, cocamidopropyl betaines, cocam idopropyl hydroxysultaines, cocobetaines, coco-hydroxysultaines, coco / oleam idopropyl betaines, coco-sultaines, decyl betaines, dihydroxyethyl oleyl glycinates, dihydroxyethyl soy glycinates, dihydroxyethyl stearyl glycinates, dihydroxyethyl tallow glycinates, dimethicones propyl PG Betaines, erucam idopropyl hydroxysultaines, hydrogenated tallow betaines, isostearam idopropyl betaines, lauram idopropyl betaines, lauryl betaines, lauryl hydroxysultaine, lauryl sultaines, milkamidopropyl betaines, minkam idopropyl betaines, myristamine idopropyl betaines, myristyl betaines, oleam idopropyl betaines, oleam idropy Hydroxysultain, Oleyl Betaine, Olivamidopropyl Betaine, Palmam Idopropyl Betaine, Palm Itam Idopropyl Betaine, Palmitoyl Carnitine, Palm Kernelamidopropyl Betaine, Polytetrafluoroethylene Acetoxypopyl Betaine, Ricinoleam Idopropyl Betaine, Sesamidopropyl Betaine, Soyamidopropyl Betaine, Stearam Idopropyl Betaine, Stearyl Betaine, Tallowam Idopropyl Betaine, Tallowamidopropyl Hydroxysultaine, Tallow Betaine, Tallow Dihydroxyethyl Betaine, Undecylenamidopropyl Betaine

and Wheat Germamidopropyl Betaine. A preferred betaine is, for example, cocamidopropyl betaine (cocoamidopropylbetaine). The betaines are particularly preferred for dishwashing compositions, most preferably hand dishwashing detergent compositions.

[0084] Further suitable surfactants include the amine oxides. The amine oxides suitable in accordance with the invention include alkylamine oxides, in particular alkyl dimethylamine oxides, alkylamidoamine oxides and alkoxyalkylamine oxides. Examples of suitable amine oxides are the following compounds designated as INCI: Almond amidopropylamine oxides, Babassu amidopropylamine oxides, Behenamine oxides, Cocamidopropyl Amine oxides, Cocamidopropylamine oxides, Cocamine oxides, Coco-Morpholine oxides, Decylamine oxides, Decyltetradecylamine oxides, Diaminopyrimidine oxides, Dihydroxyethyl C8-10 alkoxypropylamines oxides, Dihydroxyethyl C9-11 alkoxypropylamines oxides, dihydroxyethyl C12-15 alkoxypropylamines oxides, dihydroxyethyl cocamine oxides, dihydroxyethyl lauramine oxides, dihydroxyethyl stearamines oxides, dihydroxyethyl tallowamine oxides, hydrogenated palm kernel amine oxides, hydrogenated tallowamine oxides, hydroxyethyl hydroxypropyl C12-15 alkoxypropylamines oxides, isostearamidopropylamines Oxides, isostearamidopropyl morpholine oxides, lauramidopropylamine oxides, lauramine oxides, methyl morpholine oxides, milkamidopropyl amine oxides, mincamidopropylamine oxides, myristamine idopropylamine oxides, myristamine oxides, myristyl / cetyl amines Oxides, Oleamidopropylamine oxides, Oleamine oxides, Olivamidopropylamine oxides, Palmitamidopropylamine oxides, Palmitamine oxides, PEG-3 Lauramine oxides, Potassium dihydroxyethyl Cocamine oxides phosphates, Potassium Trisphosphonomethylamine oxides, Sesamidopropylamine oxides, Soyamidopropylamine oxides, Stearamidopropylamine oxides, stearamines Oxides, Tallowamidopropylamine oxides, Tallowamine oxides, Undecylenamidopropylamine oxides and Wheat Germamidopropylamine oxides. A preferred amine oxide is, for example, cocamidopropylamine oxides (cocoamidopropylamine oxide).

[0085] For automatic dishwashing applications, low-foaming nonionic surfactants are preferably used, in particular alkoxyated, especially ethoxylated, low-foaming nonionic surfactants. With particular preference, the automatic dishwashing detergents contain nonionic surfactants from the group of the alkoxyated alcohols. Particular preference is given to nonionic surfactants which have a melting point above room temperature. Nonionic surfactants having a melting point above 20 ° C, preferably above 25 ° C, more preferably between 25 and 60 ° C and especially between 26.6 and 43.3 ° C, are particularly preferred. Preferably used surfactants are those from the groups of alkoxyated nonionic surfactants, in particular the ethoxylated primary alcohols and mixtures of these surfactants with structurally more complex surfactants such as polyoxypropylene / polyoxyethylene / polyoxypropylene ((PO / EO / PO) surfactants). Such (PO / EO / PO) nonionic surfactants are also characterized by good foam control. Particularly preferred nonionic surfactants are those containing alternating ethylene oxide and different alkylene oxide units. Among these, in turn, surfactants with EO-AO-EO-AO blocks are preferred, with one to ten EO or AO groups before one block from the other group follows. Exemplary nonionic surfactants are those having a C9-alkyl group with 1 to 4 ethylene oxide units followed by 1 to 4 propylene oxide units, followed by 1 to 4 ethylene oxide units followed by 1 to 4 propylene oxide units. Preference is given in particular to end-capped, poly (oxyalkylated) nonionic surfactants with the end-cap being a linear or branched, saturated or unsaturated, aliphatic or aromatic hydrocarbon radical R having 1 to 30 carbon atoms. The alkyl groups may also comprise hydroxyl groups. The group of these nonionic surfactants include, for example, the C4-22 fatty alcohol (EO)₁₀₋₅₀-2-hydroxyalkyl ethers, in particular also the C8-12 fatty alcohol (EO)₂₂-2-hydroxydecyl ethers and the C4-22 fatty alcohol (EO)₄₀₋₈₀-2-hydroxyalkyl ethers.

[0086] When included therein the detergent will usually contain from about 1% to about 40% by weight of an anionic surfactant, such as from about 5% to about 30%, including from about 5% to about 15%, or from about 15% to about 20%, or from about 20% to about 25% of an anionic surfactant. Non-limiting examples of anionic surfactants include sulfates and sulfonates, in particular, linear alkylbenzenesulfonates (LAS), isomers of LAS, branched alkylbenzenesulfonates (BABS), phenylalkanesulfonates, alpha-olefinsulfonates (AOS), olefin sulfonates, alkene sulfonates, alkane-2,3-diylbis(sulfates), hydroxyalkanesulfonates and disulfonates, alkyl sulfates (AS) such as sodium dodecyl sulfate (SDS), fatty alcohol sulfates (FAS), primary alcohol sulfates (PAS), alcohol ethersulfates (AES or AEOS or FES, also known as alcohol ethoxysulfates or fatty alcohol ether sulfates), secondary alkanesulfonates (SAS), paraffin sulfonates (PS), ester sulfonates, sulfonated fatty acid glycerol esters, alpha-sulfo fatty acid methyl esters (alpha-SFME or SES) including methyl ester sulfonate (MES), alkyl- or alkenylsuccinic acid, dodecenyldodecylsuccinic acid (DTSA), fatty acid derivatives of amino acids, diesters and monoesters of sulfo-succinic acid or salt of fatty acids (soap), and combinations thereof.

[0087] When included therein the detergent will usually contain from about 1% to about 40% by weight of a cationic surfactant, for example from about 0.5% to about 30%, in particular from about 1% to about 20%, from about 3% to about 10%, such as from about 3% to about 5%, from about 8% to about 12% or from about 10% to about 12%. Non-limiting examples of cationic surfactants include alkyl dimethylethanolamine quat (ADMEAQ), cetyltrimethylammonium bromide (CTAB), dimethyldistearyl ammonium chloride (DSDMAC), and alkylbenzyl dimethyl ammonium, alkyl quaternary ammonium compounds, alkoxyated quaternary ammonium (AQA) compounds, ester quats, and combinations thereof.

[0088] When included therein the detergent will usually contain from about 0.2% to about 40% by weight of a nonionic surfactant, for example from about 0.5% to about 30%, in particular from about 1% to about 20%, from about 3% to

about 10%, such as from about 3% to about 5%, from about 8% to about 12%, or from about 10% to about 12%. Non-limiting examples of nonionic surfactants include alcohol ethoxylates (AE or AEO), alcohol propoxylates, propoxylated fatty alcohols (PFA), alkoxylated fatty acid alkyl esters, such as ethoxylated and/or propoxylated fatty acid alkyl esters, alkylphenol ethoxylates (APE), nonylphenol ethoxylates (NPE), alkylpolyglycosides (APG), alkoxylated amines, fatty acid monoethanolamides (FAM), fatty acid diethanolamides (FADA), ethoxylated fatty acid monoethanolamides (EFAM), propoxylated fatty acid monoethanolamides (PFAM), polyhydroxyalkyl fatty acid amides, or N-acyl N-alkyl derivatives of glucosamine (glucamides, GA, or fatty acid glucamides, FAGA), as well as products available under the trade names SPAN and TWEEN, and combinations thereof.

[0089] When included therein the detergent will usually contain from about 0.01 to about 10 % by weight of a semipolar surfactant. Non-limiting examples of semipolar surfactants include amine oxides (AO) such as alkyldimethylamineoxide, N-(coco alkyl)-N,N-dimethylamine oxide and N-(tallow-alkyl)-N,N-bis(2-hydroxyethyl)amine oxide, and combinations thereof.

[0090] When included therein the detergent will usually contain from about 0.01 % to about 10 % by weight of a zwitterionic surfactant. Non-limiting examples of zwitterionic surfactants include betaines such as alkyldimethylbetaines, sulfobetaines, and combinations thereof.

[0091] The cleaning compositions of the invention may contain about 0-65% by weight, such as about 5% to about 50% of a detergent builder or co-builder, or a mixture thereof. In a dish wash detergent, the level of builder is typically 40-65%, particularly 50-65%. The builder and/or co-builder may particularly be a chelating agent that forms water-soluble complexes with Ca and Mg. Any builder and/or co-builder known in the art for use in cleaning detergents may be utilized. Non-limiting examples of builders include zeolites, diphosphates (pyrophosphates), triphosphates such as sodium triphosphate (STP or STPP), carbonates such as sodium carbonate, soluble silicates such as sodium metasilicate, layered silicates (e.g., SKS-6 from Hoechst), ethanolamines such as 2-aminoethan-1-ol (MEA), diethanolamine (DEA, also known as 2,2'-iminodiethan-1-ol), triethanolamine (TEA, also known as 2,2',2"-nitrilotriethan-1-ol), and (carboxymethyl)inulin (CMI), and combinations thereof. "Co-builder", as used herein, means that the respective component is used in combination with another builder. All compounds disclosed as co-builders herein may also be used as main builders and vice versa, unless indicated otherwise.

[0092] In various embodiments, the composition, as defined herein, may comprise, if not indicated otherwise, from 0-65 wt %, for example about 1 wt% to about 65 wt%, from about 5 wt% to about 50 wt%, preferably from about 40 wt% to 65 wt%, such as 50 - 65 wt %, particularly about 20 wt% to about 65 wt%, particularly from 10 wt% to 50 wt% of at least one builder.

[0093] Generally and if not indicated otherwise, the builder may be preferably selected from citrate, carbonate, silicate, aluminosilicate (zeolite) and combinations thereof. Suitable builders also include phosphonates, polyphosphonates, bicarbonates, borates, and further polycarboxylates. Citrate builders, e.g., citric acid and soluble salts thereof (particularly sodium salt), are particularly suitable water-soluble organic builders. Citrates can be used in combination with zeolite, silicates like the BRITESIL types, and/or layered silicate builders. The builder and/or co-builder may be any chelating agent that forms water-soluble complexes with Ca and Mg. Any builder and/or co-builder known in the art for use in cleaning detergents may be utilized. Non-limiting examples of builders include zeolites, in particular zeolite A or P or X, carbonates such as sodium carbonate, soluble silicates such as sodium metasilicate, layered silicates (e.g., SKS-6 from Hoechst), and (carboxymethyl)inulin (CMI), and combinations thereof. Further non-limiting examples of builders include aminocarboxylates, aminopolycarboxylates, and alkyl- or alkenylsuccinic acid. Additional specific examples include 2,2',2"-nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), iminodisuccinic acid (IDS), ethylenediamine-N,N'-disuccinic acid (EDDS), methylglycine-N,N'-diacetic acid (MGDA), glutamic acid-N,N'-diacetic acid (GLDA), 1-hydroxyethane-1,1-diphosphonic acid, N-(2-hydroxyethyl)iminodiacetic acid (EDG), aspartic acid-N-monoacetic acid (ASMA), aspartic acid-N,N'-diacetic acid (ASDA), aspartic acid-N-monopropionic acid (ASMP), iminodisuccinic acid (IDA), N-(sulfomethyl)aspartic acid (SMAS), N-(2-sulfoethyl)-aspartic acid (SEAS), N-(sulfomethyl)glutamic acid (SMGL), N-(2-sulfoethyl)-glutamic acid (SEGL), N-methyliminodiacetic acid (MIDA), serine-N,N'-diacetic acid (SEDA), isoserine-N,N'-diacetic acid (ISDA), phenylalanine-N,N'-diacetic acid (PHDA), anthranilic acid-N,N'-diacetic acid (ANDA), sulfanilic acid-N,N'-diacetic acid (SLDA), taurine-N,N'-diacetic acid (TUDA) and N'-(2-hydroxyethyl)ethylenediamine-N,N,N'-triacetic acid (HEDTA), diethanolglycine (DEG), and combinations and salts thereof. Phosphonates suitable for use herein include 1-hydroxyethane-1,1-diphosphonic acid (HEDP), ethylenediaminetetrakis (methylenephosphonic acid) (EDTMPA), diethylenetriaminepentakis (methylenephosphonic acid) (DTPMPA or DTPMP), nitrilotris (methylenephosphonic acid) (ATMP or NTMP), 2-phosphonobutane-1,2,4-tricarboxylic acid (PBTC), hexamethylenediaminetetrakis (methylenephosphonic acid) (HDTMP). Particularly preferred are HEDP and DTPMP.

[0094] Suitable silicates are crystalline, layered sodium silicates of the general formula $\text{NaMSi}_x\text{O}_{2x+1} \cdot y\text{H}_2\text{O}$, wherein M is sodium or H, x a number of from 1.9 to 4 and y a number of from 0 to 20 and x is preferably 2, 3 or 4. Such silicates are for example disclosed in EP-A-0 164 514. Preferred are silicates in which M is sodium and is 2 or 3. Particularly preferred are β - and δ -sodium disilicate $\text{Na}_2\text{Si}_2\text{O}_5 \cdot y\text{H}_2\text{O}$.

[0095] Although not preferred, the compositions may also comprise phosphates, diphosphates (pyrophosphates) and/or triphosphates such as sodium triphosphate (STP or STPP). It is however preferred that all compositions disclosed herein are phosphate-free, i.e. do not contain deliberately added phosphate, in particular the phosphate content is below 1 wt %, more preferably less than 0.5 wt %, even more preferably less than 0.1 wt %, relative to the total weight of the composition. In alternative embodiments, the invention also relates to phosphate-free cleaning compositions in general that contain the polypeptides of the invention. In one aspect, the invention thus features a phosphate-free cleaning composition comprising any one or more of the polypeptides having hexosaminidase activity disclosed herein.

[0096] If not indicated otherwise, the composition may also contain 0-50% by weight, such as about 5% to about 30%, of a detergent co-builder. The composition may include a co-builder alone, or in combination with a builder, for example a zeolite builder. Non-limiting examples of co-builders include homopolymers of polyacrylates or copolymers thereof, such as poly (acrylic acid) (PAA) or copoly (acrylic acid/maleic acid) (PAA/PMA) or polyaspartic acid. Further exemplary builders and/or co-builders are described in, e.g., WO 09/102854, US 5977053.

[0097] Preferred as co-builders are acrylate-containing water-soluble polymers, such as alkali metal salts of polyacrylic acid or polymethacrylic acid, for example those having a molecular weight M_w in the range of 600 to 750,000 g / mol, as determined by gel permeation chromatography (GPC) according to DIN 55672-1:2007-08 with THF as an eluent.

[0098] Preferred polymers are polyacrylates with a molecular weight M_w of 1,000 to 15,000 g / mol, more preferred, due to their solubility, are short-chain polyacrylates with a molecular weight M_w of 1,000 to 10,000 g / mol, most preferred from 1,000 to 5,000 g / mol.

[0099] Preferred acrylates for use in the present invention are alkali metal salts of polymers of acrylic acid, preferably the sodium salts, in particular those with molecular weights in the range of 1,000 to 10,000 g / mol or 1,000 to 5,000 g / mol. Suitable acrylates are commercially available, for example under the tradename Acusol® from Dow Chemical. Suitable are also copolymers of acrylates, in particular those of acrylic acid and methacrylic acid, and acrylic acid or methacrylic acid and maleic acid.

[0100] In preferred embodiments, the compositions of the invention comprise a sulfopolymer, preferably a copolymer comprising an ethylenically unsaturated sulfonate/sulfonic acid as a co-monomer. Particularly suitable are monomers of allyl sulfonic acids, such as allyloxybenzene sulfonic acid and methallyl sulfonic acid. Particularly preferred sulfonic acid group-containing monomers are 1-acrylamido propane sulfonic acid-1, 2-acrylamido-2-propanesulfonic acid, 2-acrylamido-2-methyl-1-propanesulfonic acid, 2-methacrylamido-2-methyl-1-propanesulfonic acid, 3-methacrylamido-2-hydroxy-propanesulfonic acid, allylsulfonic acid, methallylsulfonic acid, allyloxybenzenesulfonic acid, methallyloxybenzolsulfonsäure, 2-hydroxy-3-(2-propenyloxy) propanesulfonic acid, 2-methyl-2-propenyl-sulfonic acid, styrenesulfonic acid, vinylsulfonic acid, 3-sulfopropyl, 3-sulfo - propyl, sulfomethacrylamide, sulfomethylmethacrylamide and mixtures of said acids or their water-soluble salts. The sulfopolymers are preferably copolymers of the afore-described monomers with unsaturated carboxylic acids. Especially preferred unsaturated carboxylic acids are acrylic acid, methacrylic acid, ethacrylic acid, chloroacrylic acid, alpha-cyanoacrylic acid, crotonic acid, alpha-phenyl-acrylic acid, maleic acid, maleic anhydride, fumaric acid, itaconic acid, citraconic acid, methylenemalononic acid, sorbic acid, cinnamic acid or mixtures thereof. Usable are of course also the unsaturated dicarboxylic acids. Preferred are copolymers with acrylates, in particular with acrylic acid and methacrylic acid, and acrylic acid or methacrylic acid and maleic acid.

[0101] Such polymers are, for example, commercially available under the trade names Acusol®590 or Acusol® 588 from Dow Chemical.

[0102] In one aspect of the invention, the cleaning compositions of the invention comprise a polypeptide as defined herein and at least one sulfopolymer, as defined above. Such compositions are preferably dishwashing compositions.

[0103] In one preferred embodiment, the builder is a non-phosphorus based builder such as citric acid and/or methylglycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and / or salts thereof.

[0104] Further non-limiting examples of co-builders include citrate, chelators such as aminocarboxylates, aminopolycarboxylates and phosphonates, and alkyl- or alkenylsuccinic acid. Additional specific examples include 2,2',2"-nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), iminodisuccinic acid (IDS), ethylenediamine-N,N'-disuccinic acid (EDDS), methylglycinediacetic acid (MGDA), glutamic acid-N,N-diacetic acid (GLDA), 1-hydroxyethane-1,1-diphosphonic acid (HEDP), ethylenediaminetetra(methylenephosphonic acid) (EDTMPA), diethylenetriaminepentakis(methylenephosphonic acid) (DTMPA or DTPMPA), N-(2-hydroxyethyl)iminodiacetic acid (EDG), aspartic acid-N-monoacetic acid (ASMA), aspartic acid-N,N-diacetic acid (ASDA), aspartic acid-N-monoacetic acid (ASMP), iminodisuccinic acid (IDA), N-(2-sulfomethyl)-aspartic acid (SMAS), N-(2-sulfoethyl)-aspartic acid (SEAS), N-(2-sulfomethyl)-glutamic acid (SMGL), N-(2-sulfoethyl)-glutamic acid (SEGL), N-methyliminodiacetic acid (MIDA), α -alanine-N,N-diacetic acid (α -ALDA), serine-N,N-diacetic acid (SEDA), isoserine-N,N-diacetic acid (ISDA), phenylalanine-N,N-diacetic acid (PHDA), anthranilic acid-N,N-diacetic acid (ANDA), sulfanilic acid-N,N-diacetic acid (SLDA), taurine-N,N-diacetic acid (TUDA) and sulfomethyl-N,N-diacetic acid (SMDA), N-(2-hydroxyethyl)ethylenediamine-N,N',N"-triacetic acid (HEDTA), diethanolglycine (DEG), diethylenetriamine penta(methylenephosphonic acid) (DTPMP), aminotris(methylenephosphonic acid) (ATMP), and combinations and salts thereof. Further exemplary builders and/or co-builders are described in, e.g., WO 09/102854, US 5977053

The composition e.g. cleaning composition may contain 0-50% by weight, such as 1-40%, such as 1-30%, such as about 1% to about 20%, of a bleaching system. Any bleaching system comprising components known in the art for use in cleaning detergents may be utilized. Suitable bleaching system components include sources of hydrogen peroxide; sources of peracids; and bleach catalysts or boosters.

[0105] Suitable sources of hydrogen peroxide are inorganic persalts, including alkali metal salts such as sodium percarbonate and sodium perborates (usually mono- or tetrahydrate), and hydrogen peroxide-urea (1/1).

[0106] Peracids may be (a) incorporated directly as preformed peracids or (b) formed in situ in the wash liquor from hydrogen peroxide and a bleach activator (perhydrolysis) or (c) formed in situ in the wash liquor from hydrogen peroxide and a perhydrolase and a suitable substrate for the latter, e.g., an ester.


a) Suitable preformed peracids include, but are not limited to, peroxycarboxylic acids such as peroxybenzoic acid and its ring-substituted derivatives, peroxy- α -naphthoic acid, peroxyphthalic acid, peroxyauric acid, peroxysebacic acid, ϵ -phthalimidoperoxycaproic acid [phthalimidoperoxyhexanoic acid (PAP)], and o-carboxybenzamidoperoxycaproic acid; aliphatic and aromatic diperoxydicarboxylic acids such as diperoxydodecanedioic acid, diperoxyazelaic acid, diperoxysebacic acid, diperoxybrassylic acid, 2-decyldiperoxybutanedioic acid, and diperoxyphthalic, -isophthalic and -terephthalic acids; perimidic acids; peroxymonosulfuric acid; peroxydisulfuric acid; peroxyphosphoric acid; peroxyisilicic acid; and mixtures of said compounds. It is understood that the peracids mentioned may in some cases be best added as suitable salts, such as alkali metal salts (e.g., Oxone®) or alkaline earth-metal salts.

b) Suitable bleach activators include those belonging to the class of esters, amides, imides, nitriles or anhydrides and, where applicable, salts thereof. Suitable examples are tetraacetylenediamine (TAED), sodium 4-[(3,5,5-trimethylhexanoyl)oxy]benzene-1-sulfonate (ISONOBS), sodium 4-(dodecanoyloxy)benzene-1-sulfonate (LOBS), sodium 4-(decanoyloxy)benzene-1-sulfonate, 4-(decanoyloxy)benzoic acid (DOBA), sodium 4-(nonanoyloxy)benzene-1-sulfonate (NOBS), and/or those disclosed in WO98/17767. A particular family of bleach activators of interest was disclosed in EP624154 and particularly preferred in that family is acetyl triethyl citrate (ATC). ATC or a short chain triglyceride like triacetin has the advantage that they are environmentally friendly. Furthermore, acetyl triethyl citrate and triacetin have good hydrolytical stability in the product upon storage and are efficient bleach activators. Finally, ATC is multifunctional, as the citrate released in the perhydrolysis reaction may function as a builder.

[0107] The bleaching system may also include a bleach catalyst or booster.

[0108] Some non-limiting examples of bleach catalysts that may be used in the compositions of the present invention include manganese oxalate, manganese acetate, manganese-collagen, cobalt-amine catalysts and manganese triazacyclononane (MnTACN) catalysts; particularly preferred are complexes of manganese with 1,4,7-trimethyl-1,4,7-triazacyclononane (Me3-TACN) or 1,2,4,7-tetramethyl-1,4,7-triazacyclononane (Me4-TACN), in particular Me3-TACN, such as the dinuclear manganese complex [(Me3-TACN)Mn(O)3Mn(Me3-TACN)](PF6)2, and [2,2',2''-nitritoltris(ethane-1,2-diylazanylidene- κ N-methanylylidene)triphenolato- κ 3O]manganese(III). The bleach catalysts may also be other metal compounds; such as iron or cobalt complexes.

[0109] In some embodiments, where a source of a peracid is included, an organic bleach catalyst or bleach booster may be used having one of the following formulae:



(iii) and mixtures thereof; wherein each R1 is independently a branched alkyl group containing from 9 to 24 carbons or linear alkyl group containing from 11 to 24 carbons, preferably each R1 is independently a branched alkyl group containing from 9 to 18 carbons or linear alkyl group containing from 11 to 18 carbons, more preferably each R1 is independently selected from the group consisting of 2-propylheptyl, 2-butyloctyl, 2-pentylononyl, 2-hexyldecyl, dodecyl, tetradecyl, hexadecyl, octadecyl, isononyl, isodecyl, isotridecyl and isopentadecyl.

[0110] Other exemplary bleaching systems are described, e.g. in WO2007/087258, WO2007/087244, WO2007/087259, EP1867708 (Vitamin K) and WO2007/087242. Suitable photobleaches may for example be sulfonated zinc or aluminium phthalocyanines.

[0111] The compositions may comprise metal care agents. Metal care agents may prevent or reduce the tarnishing, corrosion or oxidation of metals, including aluminium, stainless steel and non-ferrous metals, such as silver and copper. Suitable examples include one or more of the following:

(a) benzotriazoles, including benzotriazole or bis-benzotriazole and substituted derivatives thereof. Benzotriazole derivatives are those compounds in which the available substitution sites on the aromatic ring are partially or completely substituted. Suitable substituents include linear or branch-chain C₁-C₂₀-alkyl groups (e.g., C₁-C₂₀-alkyl groups) and hydroxyl, thio, phenyl or halogen such as fluorine, chlorine, bromine and iodine.

(b) metal salts and complexes chosen from the group consisting of zinc, manganese, titanium, zirconium, hafnium, vanadium, cobalt, gallium and cerium salts and/or complexes, the metals being in one of the oxidation states II, III, IV, V or VI. In one aspect, suitable metal salts and/or metal complexes may be chosen from the group consisting of Mn(II) sulphate, Mn(II) citrate, Mn(II) stearate, Mn(II) acetylacetonate, K⁺TiF₆ (e.g., K₂TiF₆), K⁺ZrF₆ (e.g., K₂ZrF₆), CoSO₄, Co(NO₃)₂ and Ce(NO₃)₃, zinc salts, for example zinc sulphate, hydrozincite or zinc acetate.;

(c) silicates, including sodium or potassium silicate, sodium disilicate, sodium metasilicate, crystalline phyllosilicate and mixtures thereof.

[0112] Further suitable organic and inorganic redox-active substances that act as silver/copper corrosion inhibitors are disclosed in WO 94/26860 and WO 94/26859. Preferably the composition of the invention comprises from 0.1 to 5% by weight of the composition of a metal care agent, preferably the metal care agent is a zinc salt.

[0113] The composition may comprise e.g. one or more hydrotrope, which is a compound that solubilises hydrophobic compounds in aqueous solutions (or oppositely, polar substances in a non-polar environment). Typically, hydrotropes have both hydrophilic and a hydrophobic character (so-called amphiphilic properties as known from surfactants), however the molecular structure of hydrotropes generally do not favor spontaneous self-aggregation, see e.g. review by Hodgdon and Kaler (2007), Current Opinion in Colloid & Interface Science 12: 121-128. Hydrotropes do not display a critical concentration above which self-aggregation occurs as found for surfactants and lipids forming micellar, lamellar or other well defined meso-phases. Instead, many hydrotropes show a continuous-type aggregation process where the sizes of aggregates grow as concentration increases. However, many hydrotropes alter the phase behavior, stability, and colloidal properties of systems containing substances of polar and non-polar character, including mixtures of water, oil, surfactants, and polymers. Hydrotropes are classically used across industries from pharma, personal care, food, to technical applications. Use of hydrotropes in detergent compositions allow for example more concentrated formulations of surfactants (as in the process of compacting liquid detergents by removing water) without inducing undesired phenomena such as phase separation or high viscosity.

[0114] The cleaning composition may contain 0-10% by weight, for example 0-5% by weight, such as about 0.5 to about 5%, or about 3% to about 5%, of a hydrotrope. Any hydrotrope known in the art for use in detergents may be utilized. Non-limiting examples of hydrotropes include sodium benzenesulfonate, sodium p-toluene sulfonate (STS), sodium xylene sulfonate (SXS), sodium cumene sulfonate (SCS), sodium cymene sulfonate, amine oxides, alcohols and polyglycoethers, sodium hydroxynaphthoate, sodium hydroxynaphthalene sulfonate, sodium ethylhexyl sulfate, and combinations thereof.

[0115] The composition e.g. cleaning composition may contain 0-10% by weight, such as 0.5-5%, 2-5%, 0.5-2% or 0.2-1% of a polymer. Any polymer known in the art for use in detergents may be utilized. The polymer may function as a co-builder as mentioned above, or may provide antiredeposition, fiber protection, soil release, dye transfer inhibition, grease cleaning and/or anti-foaming properties. Some polymers may have more than one of the above-mentioned properties and/or more than one of the below-mentioned motifs. Exemplary polymers include (carboxymethyl)cellulose (CMC), poly(vinyl alcohol) (PVA), poly(vinylpyrrolidone) (PVP), poly(ethyleneglycol) or poly(ethylene oxide) (PEG), ethoxylated poly(ethyleneimine), carboxymethyl inulin (CMI), and polycarboxylates such as PAA, PAA/PMA, poly-aspartic acid, and lauryl methacrylate/acrylic acid copolymers, hydrophobically modified CMC (HM-CMC) and silicones, copolymers of terephthalic acid and oligomeric glycols, copolymers of polyethylene terephthalate and poly(oxyethylene terephthalate) (PET-POET), PVP, poly(vinylimidazole) (PVI), poly(vinylpyridine-N-oxide) (PVPO or PVPNO) and polyvinylpyrrolidone-vinylimidazole (PVPVI). Suitable examples include PVP-K15, PVP-K30, ChromaBond S-400, ChromaBond S-403E and Chromabond S-100 from Ashland Aqualon, and Sokalan[®] HP 165, Sokalan[®] HP 50 (Dispersing agent), Sokalan[®] HP 53 (Dispersing agent), Sokalan[®] HP 59 (Dispersing agent), Sokalan[®] HP 56 (dye transfer inhibitor), Sokalan[®] HP 66 K (dye transfer inhibitor) from BASF. Further exemplary polymers include sulfonated polycarboxylates, polyethylene oxide and polypropylene oxide (PEO-PPO) and diquaternium ethoxy sulfate. Other exemplary polymers are disclosed in, e.g., WO 2006/130575. Salts of the above-mentioned polymers are also contemplated. Particularly preferred polymer is ethoxylated homopolymer Sokalan[®] HP 20 from BASF, which helps to prevent redeposition of soil in the wash liquor.

[0116] The composition e.g. cleaning composition of the present invention may also include fabric hueing agents such as dyes or pigments, which when formulated in detergent compositions can deposit onto a fabric when said fabric is

contacted with a wash liquor comprising said detergent compositions and thus altering the tint of said fabric through absorption/reflection of visible light. Fluorescent whitening agents emit at least some visible light. In contrast, fabric hueing agents alter the tint of a surface as they absorb at least a portion of the visible light spectrum. Suitable fabric hueing agents include dyes and dye-clay conjugates, and may also include pigments. Suitable dyes include small molecule dyes and polymeric dyes. Suitable small molecule dyes include small molecule dyes selected from the group consisting of dyes falling into the Colour Index (C.I.) classifications of Direct Blue, Direct Red, Direct Violet, Acid Blue, Acid Red, Acid Violet, Basic Blue, Basic Violet and Basic Red, or mixtures thereof, for example as described in WO2005/03274, WO2005/03275, WO2005/03276 and EP1876226. The detergent composition preferably comprises from about 0.00003 wt% to about 0.2 wt%, from about 0.00008 wt% to about 0.05 wt%, or even from about 0.0001 wt% to about 0.04 wt% fabric hueing agent. The composition may comprise from 0.0001 wt% to 0.2 wt% fabric hueing agent, this may be especially preferred when the composition is in the form of a unit dose pouch. Suitable hueing agents are also disclosed in, e.g. WO 2007/087257 and WO2007/087243.

[0117] The composition e.g. cleaning composition may comprise one or more additional enzymes such as one or more lipase, cutinase, an amylase, carbohydrase, cellulase, pectinase, mannanase, arabinase, galactanase, xylanase, oxidase, e.g., a laccase, and/or peroxidase.

[0118] In general, the properties of the selected enzyme(s) should be compatible with the selected detergent, (i.e., pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

[0119] Suitable proteases for the compositions of the invention include those of bacterial, fungal, plant, viral or animal origin e.g. vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. It may be an alkaline protease, such as a serine protease or a metalloprotease. A serine protease may for example be of the S1 family, such as trypsin, or the S8 family such as subtilisin. A metalloprotease protease may for example be a thermolysin from e.g. family M4 or other metalloprotease such as those from M5, M7 or M8 families.

[0120] Examples of subtilases are those derived from *Bacillus* such as *Bacillus lentus*, *Bacillus alkalophilus*, *Bacillus subtilis*, *Bacillus amyloliquefaciens*, *Bacillus pumilus* and *Bacillus gibsonii* described in; US7262042 and WO09/021867. *Subtilisin lentus*, *Subtilisin Novo*, *subtilisin Carlsberg*, *Bacillus licheniformis*, *subtilisin BPN'*, *subtilisin 309*, *subtilisin 147* and *subtilisin 168* and e.g. protease PD138 described in (WO93/18140). Other useful proteases may be those described in WO01/016285 and WO02/016547. Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the Fusarium protease described in WO94/25583 and WO05/040372, and the chymotrypsin proteases derived from Cellulomonas described in WO05/052161 and WO05/052146.

[0121] A further preferred protease is the alkaline protease from *Bacillus lentus* DSM 5483, as described for example in WO95/23221, and variants thereof which are described in WO92/21760, WO95/23221, EP1921147 and EP1921148.

[0122] Examples of metalloproteases are the neutral metalloprotease as described in WO07/044993 (Proctor & Gamble/Genencor Int.) such as those derived from *Bacillus amyloliquefaciens*.

[0123] Examples of useful proteases are the variants described in: WO89/06279, WO92/19729, WO96/034946, WO98/20115, WO98/20116, WO99/011768, WO01/44452, WO03/006602, WO04/03186, WO04/041979, WO07/006305, WO11/036263, WO11/036264, especially the variants with substitutions in one or more of the following positions: 3, 4, 9, 15, 24, 27, 42, 55, 59, 60, 66, 74, 85, 96, 97, 98, 99, 100, 101, 102, 104, 116, 118, 121, 126, 127, 128, 154, 156, 157, 158, 161, 164, 176, 179, 182, 185, 188, 189, 193, 198, 199, 200, 203, 206, 211, 212, 216, 218, 226, 229, 230, 239, 246, 255, 256, 268 and 269 wherein the positions correspond to the positions of the *Bacillus lentus* protease shown in SEQ ID NO 1 of WO 2016/001449. More preferred the protease variants may comprise one or more of the mutations selected from the group consisting of: S3T, V4I, S9R, S9E, A15T, S24G, S24R, K27R, N42R, S55P, G59E, G59D, N60D, N60E, V66A, N74D, S85R, A96S, S97G, S97D, S97A, S97SD, S99E, S99D, S99G, S99M, S99N, S99R, S99H, S101A, V102I, V102Y, V102N, S104A, G116V, G116R, H118D, H118N, A120S, S126L, P127Q, S128A, S154D, A156E, G157D, G157P, S158E, Y161A, R164S, Q176E, N179E, S182E, Q185N, A188P, G189E, V193M, N198D, V199I, Y203W, S206G, L211Q, L211D, N212D, N212S, M216S, A226V, K229L, Q230H, Q239R, N246K, N255W, N255D, N255E, L256E, L256D T268A and R269H. The protease variants are preferably variants of the *Bacillus lentus* protease (Savinase®) shown in SEQ ID NO 1 of WO 2016/001449, the *Bacillus amylolichenifaciens* protease (BPN') shown in SEQ ID NO 2 of WO2016/001449. The protease variants preferably have at least 80 % sequence identity to SEQ ID NO 1 or SEQ ID NO 2 of WO 2016/001449.

[0124] A protease variant comprising a substitution at one or more positions corresponding to positions 171, 173, 175, 179, or 180 of SEQ ID NO: 1 of WO2004/067737, wherein said protease variant has a sequence identity of at least 75% but less than 100% to SEQ ID NO: 1 of WO2004/067737.

[0125] Suitable commercially available protease enzymes include those sold under the trade names Alcalase®, Duralase™, Durazym™, Relase®, Relase® Ultra, Savinase®, Savinase® Ultra, Primase®, Polarzyme®, Kannase®, Liquanase®, Liquanase® Ultra, Ovozyme®, Coronase®, Coronase® Ultra, Blaze®, Blaze Evity® 100T, Blaze Evity® 125T, Blaze Evity® 150T, Neutrase®, Everlase® and Esperase® (Novozymes A/S), those sold under the tradename Maxatase®, Maxacal®, Maxapem®, Purafect Ox®, Purafect OxP®, Puramax®, FN2®, FN3®, FN4®, Excellase®, Excellenz P1000™,


Excellenz P1250™, Eraser®, Preferenz P100™, Purafect Prime®, Preferenz P110™, Effectenz P1000™, Purafect®, Effectenz P1050™, Purafect Ox®, Effectenz P2000™, Purafast®, Properase®, Opticlean® and Optimase® (Danisco/DuPont), Axapem™ (Gist-Brocades N.V.), BLAP (sequence shown in Figure 29 of US5352604) and variants hereof (Henkel AG) and KAP (*Bacillus alkalophilus subtilisin*) from Kao.

[0126] The protease, as described above, may be stabilized using conventional stabilizing agents, e.g., a polyol such as glycerol, (mono, di, or tri) propylene glycol, sugar alcohol, polypropylene glycol, and/or polyethylene glycol, preferably polyethylene glycol or polypropylene glycol with a molecular weight in the range of 200-1000; or compounds that act by temporarily reducing the activity of proteases (reversible inhibitors).

[0127] Thus, the composition of the invention may also include a protease inhibitor/stabilizer, which is a reversible inhibitor of protease activity, e.g., serine protease activity. Preferably, the protease inhibitor is a (reversible) subtilisin protease inhibitor. In particular, the protease inhibitor may be a peptide aldehyde, boric acid, or a boronic acid; or a derivative of any of these.

[0128] The protease inhibitor may have an inhibition constant to a serine protease, K_i (mol/L) of from $1E-12$ to $1E-03$; more preferred from $1E-11$ to $1E-04$; even more preferred from $1E-10$ to $1E-05$; even more preferred from $1E-10$ to $1E-06$; and most preferred from $1E-09$ to $1E-07$.

[0129] The protease inhibitor may be a boronic acid or a derivative thereof; preferably, a phenylboronic acid or a derivative thereof. In an embodiment of the invention, the phenyl boronic acid derivative is of the following formula:



wherein R is selected from the group consisting of hydrogen, hydroxy, C1-C6 alkyl, substituted C1-C6 alkyl, C1-C6 alkenyl and substituted C1-C6 alkenyl. Preferably, R is hydrogen, CH₃, CH₃CH₂ or CH₃CH₂CH₂.

[0130] In a preferred embodiment, the protease inhibitor (phenyl boronic acid derivative) is 4-formyl-phenyl boronic acid (4-FPBA).

[0131] In another particular embodiment, the protease inhibitor is selected from the group consisting of thiophene-2 boronic acid, thiophene-3 boronic acid, acetamidophenyl boronic acid, benzofuran-2 boronic acid, naphthalene-1 boronic acid, naphthalene-2 boronic acid, 2-FPBA, 3-FBPA, 4-FPBA, 1-thianthrene boronic acid, 4-dibenzofuran boronic acid, 5-methylthiophene-2 boronic acid, thionaphthrene boronic acid, furan-2 boronic acid, furan-3 boronic acid, 4,4 biphenyl-diboronic acid, 6-hydroxy-2-naphthalene, 4-(methylthio) phenyl boronic acid, 4 (trimethyl-silyl)phenyl boronic acid, 3-bromothiophene boronic acid, 4-methylthiophene boronic acid, 2-naphtyl boronic acid, 5-bromothiophene boronic acid, 5-chlorothiophene boronic acid, dimethylthiophene boronic acid, 2-bromophenyl boronic acid, 3-chlorophenyl boronic acid, 3-methoxy-2-thiophene, p-methyl-phenylethyl boronic acid, 2-thianthrene boronic acid, di-benzothiophene boronic acid, 4-carboxyphenyl boronic acid, 9-anthryl boronic acid, 3,5 dichlorophenyl boronic acid, diphenyl boronic acid anhydride, o-chlorophenyl boronic acid, p-chlorophenyl boronic acid, m-bromophenyl boronic acid, p-bromophenyl boronic acid, p-fluorophenyl boronic acid, p-tolyl boronic acid, o-tolyl boronic acid, octyl boronic acid, 1,3,5 trimethylphenyl boronic acid, 3-chloro-4-fluorophenyl boronic acid, 3-aminophenyl boronic acid, 3,5-bis-(trifluoromethyl) phenyl boronic acid, 2,4 dichlorophenyl boronic acid, and 4-methoxyphenyl boronic acid.

[0132] Further boronic acid derivatives suitable as protease inhibitors in the detergent composition are described in US 4,963,655, US 5,159,060, WO 95/12655, WO 95/29223, WO 92/19707, WO 94/04653, WO 94/04654, US 5442100, US 5488157 and US 5472628.

[0133] The protease stabilizer may have the formula: P-(A)_y-L-(B)_x-B₀-R* wherein:

R* is H (hydrogen), CH₃, CX₃, CHX₂, or CH₂X. Preferably, R*=H so that the stabilizer is a peptide aldehyde with the formula P-(A)_y-L-(B)_x-B₀-H;

X is a halogen atom, particularly F (fluorine);

B₀ is a single amino acid residue with L- or D-configuration of the formula -NH-CH(R)-C(=O)-;

x is 1, 2 or 3;

B_x is independently a single amino acid residue, each connected to the next B or to B₀ via its C-terminal;

L is absent or independently a linker group of the formula -C(=O)-, -C(=O)-C(=O)-, -C(=S)-, -C(=S)-C(=S)- or -C(=S)-C(=O)-;

A is absent if L is absent or is independently a single amino acid residue connected to L via the N-terminal of the amino acid;

P is selected from the group consisting of hydrogen or if L is absent an N-terminal protection group;

y is 0, 1, or 2,

R is independently selected from the group consisting of C₁₋₆ alkyl, C₆₋₁₀ aryl or C₇₋₁₀ arylalkyl, optionally substituted

with one or more, identical or different, substituent's R';

R' is independently selected from the group consisting of halogen, -OH, -OR", -SH, -SR", -NH₂, -NHR", -NR"₂, -CO₂H, -CONH₂, -CONHR", -CONR"₂, -NHC(=N)NH₂; and

R" is a C₁₋₆ alkyl group.

x may be 1, 2 or 3 and therefore B may be 1, 2 or 3 amino acid residues respectively. Thus, B may represent B1, B2-B1 or B3-B2-B1, where B3, B2 and B1 each represent one amino acid residue. y may be 0, 1 or 2 and therefore A may be absent, or 1 or 2 amino acid residues respectively having the formula A1 or A2-A1 wherein A2 and A1 each represent one amino acid residue.

[0134] B0 may be a single amino acid residue with L- or D-configuration, which is connected to H via the C-terminal of the amino acid. B0 has the formula -NH-CH(R)-C(=O)-, wherein R is a C₁₋₆ alkyl, C₆₋₁₀ aryl or C₇₋₁₀ arylalkyl side chain, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, phenyl or benzyl, and wherein R may be optionally substituted with one or more, identical or different, substituents R'. Particular examples of B0 are the D- or L-form of arginine (Arg), 3,4-dihydroxyphenylalanine, isoleucine (Ile), leucine (Leu), methionine (Met), norleucine (Nle), norvaline (Nva), phenylalanine (Phe), m-tyrosine, p-tyrosine (Tyr) and valine (Val). A particular embodiment is when B0 is leucine, methionine, phenylalanine, p-tyrosine and valine.

[0135] B1, which is connected to B0 via the C-terminal of the amino acid, may be an aliphatic, hydrophobic and/or neutral amino acid. Examples of B1 are alanine (Ala), cysteine (Cys), glycine (Gly), isoleucine (Ile), leucine (Leu), norleucine (Nle), norvaline (Nva), proline (Pro), serine (Ser), threonine (Thr) and valine (Val). Particular examples of B1 are alanine, glycine, isoleucine, leucine and valine. A particular embodiment is when B1 is alanine, glycine or valine.

[0136] If present, B2, which is connected to B1 via the C-terminal of the amino acid, may be an aliphatic, hydrophobic, neutral and/or polar amino acid. Examples of B2 are alanine (Ala), arginine (Arg), capreomycin (Cpd), cysteine (Cys), glycine (Gly), isoleucine (Ile), leucine (Leu), norleucine (Nle), norvaline (Nva), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr), and valine (Val). Particular examples of B2 are alanine, arginine, capreomycin, glycine, isoleucine, leucine, phenylalanine and valine. A particular embodiment is when B2 is arginine, glycine, leucine, phenylalanine or valine.

[0137] B3, which if present is connected to B2 via the C-terminal of the amino acid, may be a large, aliphatic, aromatic, hydrophobic and/or neutral amino acid. Examples of B3 are isoleucine (Ile), leucine (Leu), norleucine (Nle), norvaline (Nva), phenylalanine (Phe), phenylglycine, tyrosine (Tyr), tryptophan (Trp) and valine (Val). Particular examples of B3 are leucine, phenylalanine, tyrosine and tryptophan.

[0138] The linker group L may be absent or selected from the group consisting of -C(=O)-, -C(=O)-C(=O)-, -C(=S)-, -C(=S)-C(=S)- or -C(=S)-C(=O)-. Particular embodiments of the invention are when L is absent or L is a carbonyl group -C(=O)-.

[0139] A1, which if present is connected to L via the N-terminal of the amino acid, may be an aliphatic, aromatic, hydrophobic, neutral and/or polar amino acid. Examples of A1 are alanine (Ala), arginine (Arg), capreomycin (Cpd), glycine (Gly), isoleucine (Ile), leucine (Leu), norleucine (Nle), norvaline (Nva), phenylalanine (Phe), threonine (Thr), tyrosine (Tyr), tryptophan (Trp) and valine (Val). Particular examples of A1 are alanine, arginine, glycine, leucine, phenylalanine, tyrosine, tryptophan and valine. A particular embodiment is when B2 is leucine, phenylalanine, tyrosine or tryptophan.

[0140] The A2 residue, which if present is connected to A1 via the N-terminal of the amino acid, may be a large, aliphatic, aromatic, hydrophobic and/or neutral amino acid. Examples of A2 are arginine (Arg), isoleucine (Ile), leucine (Leu), norleucine (Nle), norvaline (Nva), phenylalanine (Phe), phenylglycine, Tyrosine (Tyr), tryptophan (Trp) and valine (Val). Particular examples of A2 are phenylalanine and tyrosine.

[0141] The N-terminal protection group P (if present) may be selected from formyl, acetyl (Ac), benzoyl (Bz), trifluoroacetyl, methoxysuccinyl, aromatic and aliphatic urethane protecting groups such as fluorenylmethyloxycarbonyl (Fmoc), methoxycarbonyl (Moc), (fluoromethoxy)carbonyl, benzyloxycarbonyl (Cbz), t-butyloxycarbonyl (Boc) and adamantyloxycarbonyl; p-methoxybenzyl carbonyl, benzyl (Bn), p-methoxybenzyl (PMB), p-methoxyphenyl (PMP), methoxyacetyl, methylamino carbonyl, methylsulfonyl, ethylsulfonyl, benzylsulfonyl, methylphosphoramidyl (MeOP(OH)(=O)) and benzylphosphoramidyl (PhCH₂OP(OH)(=O)).

[0142] In the case of a tripeptide aldehyde with a protection group (i.e. x=2, L is absent and A is absent), P is preferably acetyl, methoxycarbonyl, benzyloxycarbonyl, methylamino carbonyl, methylsulfonyl, benzylsulfonyl and benzylphosphoramidyl. In the case of a tetrapeptide aldehyde with a protection group (i.e. x=3, L is absent and A is absent), P is preferably acetyl, methoxycarbonyl, methylsulfonyl, ethylsulfonyl and methylphosphoramidyl.

[0143] Suitable peptide aldehydes are described in WO94/04651, WO95/25791, WO98/13458, WO98/13459, WO98/13460, WO98/13461, WO98/13462, WO07/141736, WO07/145963, WO09/118375, WO10/055052 and WO11/036153. More particularly, the peptide aldehyde may be Cbz-Arg-Ala-Tyr-H, Ac-Gly-Ala-Tyr-H, Cbz-Gly-Ala-Tyr-H, Cbz-Gly-Ala-Tyr-CF₃, Cbz-Gly-Ala-Leu-H, Cbz-Val-Ala-Leu-H, Cbz-Val-Ala-Leu-CF₃, Moc-Val-Ala-Leu-CF₃, Cbz-Gly-Ala-Phe-H, Cbz-Gly-Ala-Phe-CF₃, Cbz-Gly-Ala-Val-H, Cbz-Gly-Gly-Tyr-H, Cbz-Gly-Gly-Phe-H, Cbz-Arg-Val-Tyr-H,

Cbz-Leu-Val-Tyr-H, Ac-Leu-Gly-Ala-Tyr-H, Ac-Phe-Gly-Ala-Tyr-H, Ac-Tyr-Gly-Ala-Tyr-H, Ac-Phe-Gly-Ala-Leu-H, Ac-Phe-Gly-Ala-Phe-H, Ac-Phe-Gly-Val-Tyr-H, Ac-Phe-Gly-Ala-Met-H, Ac-Trp-Leu-Val-Tyr-H, MeO-CO-Val-Ala-Leu-H, MeNCO-Val-Ala-Leu-H, MeO-CO-Phe-Gly-Ala-Leu-H, MeO-CO-Phe-Gly-Ala-Phe-H, MeSOz-Phe-Gly-Ala-Leu-H, MeSOz-Val-Ala-Leu-H, PhCH₂O-P(OH)(O)-Val-Ala-Leu-H, EtSOz-Phe-Gly-Ala-Leu-H, PhCH₂SO₂-Val-Ala-Leu-H, PhCH₂O-P(OH)(O)-Leu-Ala-Leu-H, PhCH₂O-P(OH)(O)-Phe-Ala-Leu-H, or MeO-P(OH)(O)-Leu-Gly-Ala-Leu-H. A preferred stabilizer for use in the liquid composition of the invention is Cbz-Gly-Ala-Tyr-H, or a hydrosulfite adduct thereof, wherein Cbz is benzyloxycarbonyl.

[0144] Further examples of such peptide aldehydes include α -MAPI, β -MAPI, Phe-C(=O)-Arg-Val-Tyr-H, Phe-C(=O)-Gly-Gly-Tyr-H, Phe-C(=O)-Gly-Ala-Phe-H, Phe-C(=O)-Gly-Ala-Tyr-H, Phe-C(=O)-Gly-Ala-L-H, Phe-C(=O)-Gly-Ala-Nva-H, Phe-C(=O)-Gly-Ala-Nle-H, Tyr-C(=O)-Arg-Val-Tyr-H, Tyr-C(=O)-Gly-Ala-Tyr-H, Phe-C(=S)-Arg-Val-Phe-H, Phe-C(=S)-Arg-Val-Tyr-H, Phe-C(=S)-Gly-Ala-Tyr-H, Antipain, GE20372A, GE20372B, Chymostatin A, Chymostatin B, and Chymostatin C.

[0145] The protease stabilizer may be a hydrosulfite adduct of the peptide aldehyde described above, e.g. as described in WO 2013/004636. The adduct may have the formula P-(A)y-L-(B)x-N(H)-CHR-CH(OH)-SO₃M, wherein P, A, y, L, B, x and R are defined as above, and M is H or an alkali metal, preferably Na or K.

[0146] An aqueous solution of the hydrosulfite adduct may be prepared by reacting the corresponding peptide aldehyde with an aqueous solution of sodium bisulfite (sodium hydrogen sulfite, NaHSOs); potassium bisulfite (KHSOs) by known methods, e.g., as described in WO 98/47523; US 6,500,802; US 5,436,229; J. Am. Chem. Soc. (1978) 100, 1228; Org. Synth., Coll. vol. 7: 361.

[0147] Particularly preferred peptide aldehyde protease stabilizers have the formula P-B3-B2-B1-B0-H, or a hydrosulfite adduct having the formula P-B3-B2-B1-N(H)-CHR-CHOH-SO₃M, wherein

- i) H is hydrogen;
- ii) B0 is a single amino acid residue with L- or D-configuration of the formula -NH-CH(R)-C(=O)-;
- iii) B1 and B2 are independently single amino acid residues;
- iv) B3 is a single amino acid residue, or is absent;
- v) R is independently selected from the group consisting of C₁₋₆ alkyl, C₆₋₁₀ aryl or C₇₋₁₀ arylalkyl optionally substituted with one or more, identical or different, substituents R';
- vi) R' is independently selected from the group consisting of halogen, -OH, -OR", -SH, -SR", -NH₂, -NHR", -NR"₂, -COzH, -CONH₂, -CONHR", -CONR"₂, -NHC(=N)NH₂;
- vii) R" is a C₁₋₆ alkyl group;
- viii) P is an N-terminal protection group, preferably methoxycarbonyl (Moc) or benzyloxycarbonyl (Cbz); and
- ix) M is H or an alkali metal, preferably Na or K.

[0148] In an even more preferred embodiment, the peptide aldehyde protease stabilizer has the formula P-B2-B1-B0-H or an adduct having the formula P-B2-B1-N(H)-CHR-CHOH-SO₃M, wherein

- i) H is hydrogen;
- ii) B0 is a single amino acid residue with L- or D-configuration of the formula -NH-CH(R)-C(=O)-;
- iii) B1 and B2 are independently single amino acid residues;
- iv) R is independently selected from the group consisting of C₁₋₆ alkyl, C₆₋₁₀ aryl or C₇₋₁₀ arylalkyl optionally substituted with one or more, identical or different, substituents R';
- v) R' is independently selected from the group consisting of halogen, -OH, -OR", -SH, -SR", -NH₂, -NHR", -NR"₂, -COzH, -CONH₂, -CONHR", -CONR"₂, -NHC(=N)NH₂;
- vi) R" is a C₁₋₆ alkyl group;
- vii) P is an N-terminal protection group, preferably methoxycarbonyl (Moc) or benzyloxycarbonyl (Cbz); and
- viii) M is H or an alkali metal, preferably Na or K.

[0149] Preferred embodiments of B0, B1, B2, B3, and P are as described above.

[0150] The molar ratio of the above-mentioned peptide aldehydes (or hydrosulfite adducts) to the protease may be at least 1:1 or 1.5:1, and it may be less than 1000:1, more preferred less than 500:1, even more preferred from 100:1 to 2:1 or from 20:1 to 2:1, or most preferred, the molar ratio is from 10:1 to 2:1.

[0151] Formate salts (e.g., sodium formate) and formic acid have also shown good effects as inhibitor of protease activity. Formate can be used synergistically with the above-mentioned protease inhibitors, as shown in WO 2013/004635. The formate salts may be present in the slurry composition in an amount of at least 0.1% w/w or 0.5% w/w, e.g., at least 1.0%, at least 1.2% or at least 1.5%. The amount is typically below 5% w/w, below 4% or below 3%.

[0152] In an embodiment, the protease is a metalloprotease and the inhibitor is a metalloprotease inhibitor, e.g., a protein hydrolysate based inhibitor (e.g., as described in WO 2008/134343).

[0153] Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g., the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in US 4,435,307, US 5,648,263, US 5,691,178, US 5,776,757 and WO 89/09259.

[0154] Especially suitable cellulases are the alkaline or neutral cellulases having colour care benefits. Examples of such cellulases are cellulases described in EP 0 495 257, EP 0 531 372, WO 96/11262, WO 96/29397, WO 98/08940. Other examples are cellulase variants such as those described in WO 94/07998, EP 0 531 315, US 5,457,046, US 5,686,593, US 5,763,254, WO 95/24471, WO 98/12307 and WO99/001544.

[0155] Other cellulases are endo-beta-1,4-glucanase enzyme having a sequence of at least 97% identity to the amino acid sequence of position 1 to position 773 of SEQ ID NO:2 of WO 2002/099091 or a family 44 xyloglucanase, which a xyloglucanase enzyme having a sequence of at least 60% identity to positions 40-559 of SEQ ID NO: 2 of WO 2001/062903.

[0156] Commercially available cellulases include Celluzyme™, and Carezyme™ (Novozymes A/S) Carezyme Premium™ (Novozymes A/S), Celluclean™ (Novozymes A/S), Celluclean Classic™ (Novozymes A/S), Cellusoft™ (Novozymes A/S), Whitezyme™ (Novozymes A/S), Clazinase™, and Puradax HA™ (Genencor International Inc.), and KAC-500(B)™ (Kao Corporation).

[0157] Suitable mannanases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. The mannanase may be an alkaline mannanase of Family 5 or 26. It may be a wild-type from *Bacillus* or *Humicola*, particularly *B. agaradhaerens*, *B. licheniformis*, *B. halodurans*, *B. clausii*, or *H. insolens*. Suitable mannanases are described in WO1999/064619. A commercially available mannanase is Mannaway (Novozymes A/S).

[0158] Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinus*, e.g., from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257. Commercially available peroxidases include Guardzyme™ (Novozymes A/S).

[0159] Suitable lipases and cutinases include those of bacterial or fungal origin. Chemically modified or protein engineered mutant enzymes are included. Examples include lipase from *Thermomyces*, e.g. from *T. lanuginosus* (previously named *Humicola lanuginosa*) as described in EP258068 and EP305216, cutinase from *Humicola*, e.g. *H. insolens* (WO96/13580), lipase from strains of *Pseudomonas* (some of these now renamed to *Burkholderia*), e.g. *P. alcaligenes* or *P. pseudoalcaligenes* (EP218272), *P. cepacia* (EP331376), *P. sp.* strain SD705 (WO95/06720 & WO96/27002), *P. wisconsinensis* (WO96/12012), GDSL-type *Streptomyces* lipases (WO10/065455), cutinase from *Magnaporthe grisea* (WO10/107560), cutinase from *Pseudomonas mendocina* (US5,389,536), lipase from *Thermobifida fusca* (WO11/084412), *Geobacillus stearothermophilus* lipase (WO11/084417), lipase from *Bacillus subtilis* (WO11/084599), and lipase from *Streptomyces griseus* (WO11/150157) and *S. pristinaespiralis* (WO12/137147).

[0160] Other examples are lipase variants such as those described in EP407225, WO92/05249, WO94/01541, WO94/25578, WO95/14783, WO95/30744, WO95/35381, WO95/22615, WO96/00292, WO97/04079, WO97/07202, WO00/34450, WO00/60063, WO01/92502, WO07/87508 and WO09/109500.

[0161] Preferred commercial lipase products include Lipolase™, Lipex™, Lipolex™ and Lipoclean™ (Novozymes A/S), Lumafast (originally from Genencor) and Lipomax (originally from Gist-Brocades).

[0162] Still other examples are lipases sometimes referred to as acyltransferases or perhydrolases, e.g. acyltransferases with homology to *Candida antarctica* lipase A (WO10/111143), acyltransferase from *Mycobacterium smegmatis* (WO05/56782), perhydrolases from the CE 7 family (WO09/67279), and variants of the *M. smegmatis* perhydrolase in particular the S54V variant used in the commercial product Gentle Power Bleach from Huntsman Textile Effects Pte Ltd (WO10/100028).

[0163] Suitable amylases include alpha-amylases and/or a glucoamylases and may be of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, alpha-amylases obtained from *Bacillus*, e.g., a special strain of *Bacillus licheniformis*, described in more detail in GB 1,296,839.

[0164] Suitable amylases include amylases having SEQ ID NO: 2 in WO 95/10603 or variants having 90% sequence identity to SEQ ID NO: 3 thereof. Preferred variants are described in WO 94/02597, WO 94/18314, WO 97/43424 and SEQ ID NO: 4 of WO 99/019467, such as variants with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 178, 179, 181, 188, 190, 197, 201, 202, 207, 208, 209, 211, 243, 264, 304, 305, 391, 408, and 444.

[0165] Different suitable amylases include amylases having SEQ ID NO: 6 in WO 02/010355 or variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a deletion in positions 181 and 182 and a substitution in position 193.

[0166] Other amylases which are suitable are hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 of WO 2006/066594 or variants having 90% sequence identity thereof. Preferred variants of this hybrid alpha-amylase are those having a substitution, a deletion or an insertion in one of more of the

EP 3 647 398 B1

following positions: G48, T49, G107, H156, A181, N190, M197, I201, A209 and Q264. Most preferred variants of the hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of SEQ ID NO: 4 are those having the substitutions:

M197T;
H156Y+A181T+N190F+A209V+Q264S; or
G48A+T49I+G107A+H156Y+A181T+N190F+I201F+A209V+Q264S.

[0167] Further amylases which are suitable are amylases having SEQ ID NO: 6 in WO 99/019467 or variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a substitution, a deletion or an insertion in one or more of the following positions: R181, G182, H183, G184, N195, I206, E212, E216 and K269. Particularly preferred amylases are those having deletion in positions R181 and G182, or positions H183 and G184.

[0168] Additional amylases which can be used are those having SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 2 or SEQ ID NO: 7 of WO 96/023873 or variants thereof having 90% sequence identity to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7. Preferred variants of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7 are those having a substitution, a deletion or an insertion in one or more of the following positions: 140, 181, 182, 183, 184, 195, 206, 212, 243, 260, 269, 304 and 476, using SEQ ID 2 of WO 96/023873 for numbering. More preferred variants are those having a deletion in two positions selected from 181, 182, 183 and 184, such as 181 and 182, 182 and 183, or positions 183 and 184. Most preferred amylase variants of SEQ ID NO: 1, SEQ ID NO: 2 or SEQ ID NO: 7 are those having a deletion in positions 183 and 184 and a substitution in one or more of positions 140, 195, 206, 243, 260, 304 and 476.

[0169] Other amylases which can be used are amylases having SEQ ID NO: 2 of WO 08/153815, SEQ ID NO: 10 in WO 01/66712 or variants thereof having 90% sequence identity to SEQ ID NO: 2 of WO 08/153815 or 90% sequence identity to SEQ ID NO: 10 in WO 01/66712. Preferred variants of SEQ ID NO: 10 in WO 01/66712 are those having a substitution, a deletion or an insertion in one or more of the following positions: 176, 177, 178, 179, 190, 201, 207, 211 and 264.

[0170] Further suitable amylases are amylases having SEQ ID NO: 2 of WO 09/061380 or variants having 90% sequence identity to SEQ ID NO: 2 thereof. Preferred variants of SEQ ID NO: 2 are those having a truncation of the C-terminus and/or a substitution, a deletion or an insertion in one or more of the following positions: Q87, Q98, S125, N128, T131, T165, K178, R180, S181, T182, G183, M201, F202, N225, S243, N272, N282, Y305, R309, D319, Q320, Q359, K444 and G475. More preferred variants of SEQ ID NO: 2 are those having the substitution in one or more of the following positions: Q87E,R, Q98R, S125A, N128C, T131I, T165I, K178L, T182G, M201L, F202Y, N225E,R, N272E,R, S243Q,A,E,D, Y305R, R309A, Q320R, Q359E, K444E and G475K and/or deletion in position R180 and/or S181 or of T182 and/or G183. Most preferred amylase variants of SEQ ID NO: 2 are those having the substitutions:

N128C+K178L+T182G+Y305R+G475K;
N128C+K178L+T182G+F202Y+Y305R+D319T+G475K;
S125A+N128C+K178L+T182G+Y305R+G475K; or
S125A+N128C+T131I+T165I+K178L+T182G+Y305R+G475K wherein the variants are C-terminally truncated and optionally further comprises a substitution at position 243 and/or a deletion at position 180 and/or position 181.

[0171] Further suitable amylases are amylases having SEQ ID NO: 1 of WO13184577 or variants having 90% sequence identity to SEQ ID NO: 1 thereof. Preferred variants of SEQ ID NO: 1 are those having a substitution, a deletion or an insertion in one or more of the following positions: K176, R178, G179, T180, G181, E187, N192, M199, I203, S241, R458, T459, D460, G476 and G477. More preferred variants of SEQ ID NO: 1 are those having the substitution in one or more of the following positions: K176L, E187P, N192FYH, M199L, I203YF, S241QADN, R458N, T459S, D460T, G476K and G477K and/or deletion in position R178 and/or S179 or of T180 and/or G181. Most preferred amylase variants of SEQ ID NO: 1 are those having the substitutions:

E187P+I203Y+G476K
E187P+I203Y+R458N+T459S+D460T+G476K

wherein the variants optionally further comprise a substitution at position 241 and/or a deletion at position 178 and/or position 179.

[0172] Further suitable amylases are amylases having SEQ ID NO: 1 of WO10104675 or variants having 90% sequence identity to SEQ ID NO: 1 thereof. Preferred variants of SEQ ID NO: 1 are those having a substitution, a deletion or an insertion in one or more of the following positions: N21, D97, V128 K177, R179, S180, I181, G182, M200, L204, E242,

G477 and G478. More preferred variants of SEQ ID NO: 1 are those having the substitution in one of more of the following positions: N21D, D97N, V128I K177L, M200L, L204YF, E242QA, G477K and G478K and/or deletion in position R179 and/or S180 or of I181 and/or G182. Most preferred amylase variants of SEQ ID NO: 1 are those having the substitutions: N21D+D97N+V128I

wherein the variants optionally further comprise a substitution at position 200 and/or a deletion at position 180 and/or position 181.

[0173] Other suitable amylases are the alpha-amylase having SEQ ID NO: 12 in WO01/66712 or a variant having at least 90% sequence identity to SEQ ID NO: 12. Preferred amylase variants are those having a substitution, a deletion or an insertion in one of more of the following positions of SEQ ID NO: 12 in WO01/66712: R28, R118, N174; R181, G182, D183, G184, G186, W189, N195, M202, Y298, N299, K302, S303, N306, R310, N314; R320, H324, E345, Y396, R400, W439, R444, N445, K446, Q449, R458, N471, N484. Particular preferred amylases include variants having a deletion of D183 and G184 and having the substitutions R118K, N195F, R320K and R458K, and a variant additionally having substitutions in one or more position selected from the group: M9, G149, G182, G186, M202, T257, Y295, N299, M323, E345 and A339, most preferred a variant that additionally has substitutions in all these positions.

[0174] Other examples are amylase variants such as those described in WO2011/098531, WO2013/001078 and WO2013/001087.

[0175] Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™, Stainzyme™, Stainzyme Plus™, Natalase™, Liquozyme X and BAN™ (from Novozymes A/S), and Rapidase™, Purastar™/Effectenz™, Powerase, Preferenz S1000, Preferenz S100 and Preferenz S110 (from Genencor International Inc./DuPont).

[0176] A suitable peroxidase may be a peroxidase enzyme comprised by the enzyme classification EC 1.11.1.7, as set out by the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (IUBMB), or any fragment derived therefrom, exhibiting peroxidase activity.

[0177] Suitable peroxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinopsis*, e.g., from *C. cinerea* (EP 179,486), and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257.

[0178] A suitable peroxidase includes a haloperoxidase enzyme, such as chloroperoxidase, bromoperoxidase and compounds exhibiting chloroperoxidase or bromoperoxidase activity. Haloperoxidases are classified according to their specificity for halide ions. Chloroperoxidases (E.C. 1.11.1.10) catalyze formation of hypochlorite from chloride ions. Preferably, the haloperoxidase is a vanadium haloperoxidase, i.e., a vanadate-containing haloperoxidase. Haloperoxidases have been isolated from many different fungi, in particular from the fungus group dematiaceous hyphomycetes, such as *Caldariomyces*, e.g., *C. fumago*, *Alternaria*, *Curvularia*, e.g., *C. verruculosa* and *C. inaequalis*, *Drechslera*, *Ulocladium* and *Botrytis*.

[0179] Haloperoxidases have also been isolated from bacteria such as *Pseudomonas*, e.g., *P. pyrocinia* and *Streptomyces*, e.g., *S. aureofaciens*.

[0180] A suitable oxidase includes in particular, any laccase enzyme comprised by the enzyme classification EC 1.10.3.2, or any fragment derived therefrom exhibiting laccase activity, or a compound exhibiting a similar activity, such as a catechol oxidase (EC 1.10.3.1), an o-aminophenol oxidase (EC 1.10.3.4), or a bilirubin oxidase (EC 1.3.3.5). Preferred laccase enzymes are enzymes of microbial origin. The enzymes may be derived from plants, bacteria or fungi (including filamentous fungi and yeasts). Suitable examples from fungi include a laccase derivable from a strain of *Aspergillus*, *Neurospora*, e.g., *N. crassa*, *Podospora*, *Botrytis*, *Collybia*, *Fomes*, *Lentinus*, *Pleurotus*, *Trametes*, e.g., *T. villosa* and *T. versicolor*, *Rhizoctonia*, e.g., *R. solani*, *Coprinopsis*, e.g., *C. cinerea*, *C. comatus*, *C. friesii*, and *C. plicatilis*, *Psathyrella*, e.g., *P. condelleana*, *Panaeolus*, e.g., *P. papilionaceus*, *Myceliophthora*, e.g., *M. thermophila*, *Schytalidium*, e.g., *S. thermophilum*, *Polyporus*, e.g., *P. pin situs*, *Phlebia*, e.g., *P. radiata* (WO 92/01046), or *Coriolus*, e.g., *C. hirsutus* (JP 2238885). Suitable examples from bacteria include a laccase derivable from a strain of *Bacillus*. A laccase derived from *Coprinopsis* or *Myceliophthora* is preferred; in particular, a laccase derived from *Coprinopsis cinerea*, as disclosed in WO 97/08325; or from *Myceliophthora thermophila*, as disclosed in WO 95/33836.

[0181] The composition e.g. cleaning composition of the present invention can also contain dispersants. In particular, powdered detergents may comprise dispersants. Suitable water-soluble organic materials include the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms. Suitable dispersants are for example described in Powdered Detergents, Surfactant science series volume 71, Marcel Dekker, Inc.

[0182] The composition e.g. cleaning composition of the present invention may also include one or more dye transfer inhibiting agents. Suitable polymeric dye transfer inhibiting agents include, but are not limited to, polyvinylpyrrolidone polymers, polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof. When present in a subject composition, the dye transfer inhibiting agents may be present at levels from about 0.0001 % to about 10%, from about 0.01% to about 5% or even from about 0.1% to about 3% by weight of the composition.

[0183] The composition of the present invention will preferably also contain additional components that may tint articles

being cleaned, such as fluorescent whitening agent or optical brighteners. Where present the brightener is preferably at a level of about 0.01% to about 0.5%. Any fluorescent whitening agent suitable for use in a laundry detergent composition may be used in the composition of the present invention. The most commonly used fluorescent whitening agents are those belonging to the classes of diaminostilbene-sulfonic acid derivatives, diarylpyrazoline derivatives and bisphenyl-distyryl derivatives. Examples of the diaminostilbene-sulfonic acid derivative type of fluorescent whitening agents include the sodium salts of: 4,4'-bis-(2-diethanolamino-4-anilino-s-triazin-6-ylamino) stilbene-2,2'-disulfonate, 4,4'-bis-(2,4-di-anilino-s-triazin-6-ylamino) stilbene-2,2'-disulfonate, 4,4'-bis-(2-anilino-4-(N-methyl-N-2-hydroxy-ethylamino)-s-triazin-6-ylamino) stilbene-2,2'-disulfonate, 4,4'-bis-(4-phenyl-1,2,3-triazol-2-yl)stilbene-2,2'-disulfonate and sodium 5-(2H-naphtho[1,2-d][1,2,3]triazol-2-yl)-2-[(E)-2-phenylvinyl]benzenesulfonate. Preferred fluorescent whitening agents are Tinopal DMS and Tinopal CBS available from Ciba-Geigy AG, Basel, Switzerland. Tinopal DMS is the disodium salt of 4,4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino) stilbene-2,2'-disulfonate. Tinopal CBS is the disodium salt of 2,2'-bis-(phenyl-styryl)-disulfonate. Also preferred are fluorescent whitening agents is the commercially available Parawhite KX, supplied by Paramount Minerals and Chemicals, Mumbai, India. Other fluorescent brighteners suitable for use in the invention include the 1-3-diaryl pyrazolines and the 7-alkylaminocoumarins. Suitable fluorescent brightener levels include lower levels of from about 0.01, from 0.05, from about 0.1 or even from about 0.2 wt % to upper levels of 0.5 or even 0.75 wt%.

[0184] The composition of the present invention may also include one or more soil release polymers which aid the removal of soils from fabrics such as cotton and polyester based fabrics, in particular the removal of hydrophobic soils from polyester based fabrics. The soil release polymers may for example be nonionic or anionic terephthalate based polymers, polyvinyl caprolactam and related copolymers, vinyl graft copolymers, polyester polyamides see for example Chapter 7 in Powdered Detergents, Surfactant science series volume 71, Marcel Dekker, Inc. Another type of soil release polymers is amphiphilic alkoxylated grease cleaning polymers comprising a core structure and a plurality of alkoxylate groups attached to that core structure. The core structure may comprise a polyalkylenimine structure or a polyalkanolamine structure as described in detail in WO 2009/087523. Furthermore, random graft co-polymers are suitable soil release polymers. Suitable graft co-polymers are described in more detail in WO 2007/138054, WO 2006/108856 and WO 2006/113314. Suitable polyethylene glycol polymers include random graft co-polymers comprising: (i) hydrophilic backbone comprising polyethylene glycol; and (ii) side chain(s) selected from the group consisting of: C4-C25 alkyl group, polypropylene, polybutylene, vinyl ester of a saturated C1-C6 mono-carboxylic acid, C1-C 6 alkyl ester of acrylic or methacrylic acid, and mixtures thereof. Suitable polyethylene glycol polymers have a polyethylene glycol backbone with random grafted polyvinyl acetate side chains. The average molecular weight of the polyethylene glycol backbone can be in the range of from 2,000 Da to 20,000 Da, or from 4,000 Da to 8,000 Da. The molecular weight ratio of the polyethylene glycol backbone to the polyvinyl acetate side chains can be in the range of from 1: 1 to 1:5, or from 1: 1.2 to 1:2. The average number of graft sites per ethylene oxide units can be less than 1, or less than 0.8, the average number of graft sites per ethylene oxide units can be in the range of from 0.5 to 0.9, or the average number of graft sites per ethylene oxide units can be in the range of from 0.1 to 0.5, or from 0.2 to 0.4. A suitable polyethylene glycol polymer is Sokalan HP22. Other soil release polymers are substituted polysaccharide structures especially substituted cellulosic structures such as modified cellulose derivatives such as those described in EP 1867808 or WO 2003/040279. Suitable cellulosic polymers include cellulose, cellulose ethers, cellulose esters, cellulose amides and mixtures thereof. Suitable cellulosic polymers include anionically modified cellulose, nonionically modified cellulose, cationically modified cellulose, zwitterionically modified cellulose, and mixtures thereof. Suitable cellulosic polymers include methyl cellulose, carboxy methyl cellulose, ethyl cellulose, hydroxyl ethyl cellulose, hydroxyl propyl methyl cellulose, ester carboxy methyl cellulose, and mixtures thereof.

[0185] The composition of the present invention may also include one or more anti-redeposition agents such as carboxymethylcellulose (CMC), polyvinyl alcohol (PVA), polyvinylpyrrolidone (PVP), polyoxyethylene and/or polyethyleneglycol (PEG), homopolymers of acrylic acid, copolymers of acrylic acid and maleic acid, and ethoxylated polyethylenimine. The cellulose based polymers described under soil release polymers above may also function as anti-redeposition agents.

[0186] The composition of the present invention may also include one or more rheology modifiers, structurants or thickeners, as distinct from viscosity reducing agents. The rheology modifiers are selected from the group consisting of non-polymeric crystalline, hydroxy-functional materials, polymeric rheology modifiers which impart shear thinning characteristics to the aqueous liquid matrix of a liquid detergent composition. The rheology and viscosity of the detergent can be modified and adjusted by methods known in the art, for example as shown in EP 2169040.

[0187] Other suitable cleaning composition components include, but are not limited to, anti-shrink agents, anti-wrinkling agents, bactericides, binders, carriers, dyes, enzyme stabilizers, fabric softeners, fillers, foam regulators, hydrotropes, perfumes, pigments, sud suppressors, solvents, bittering agents, and structurants for liquid detergents and/or structure elasticizing agents.

[0188] In various embodiments, the invention relates to compositions which comprise the polypeptides having hexosaminidase activity, as described herein, and any one or more of an adjunct ingredient selected from bittering agents and organic solvents, such as glycerol and 1,2-propane diol.

Formulation of enzyme products

[0189] The cleaning composition of the invention may be in any convenient form, e.g., a bar, a homogenous tablet, a tablet having two or more layers, a pouch having one or more compartments, such as 2 or more, preferably 2, 3, 4 or 5 compartments, a regular or compact powder, a granule, a paste, a gel, or a regular, compact or concentrated liquid.

[0190] Pouches can be configured as single or multicompartments. It can be of any form, shape and material which is suitable for hold the composition, e.g. without allowing the release of the composition to release of the composition from the pouch prior to water contact. The pouch is made from water soluble film which encloses an inner volume. Said inner volume can be divided into compartments of the pouch. Preferred films are polymeric materials preferably polymers which are formed into a film or sheet. Preferred polymers, copolymers or derivatives thereof are selected polyacrylates, and water soluble acrylate copolymers, methyl cellulose, carboxy methyl cellulose, sodium dextrin, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose, malto dextrin, poly methacrylates, most preferably polyvinyl alcohol copolymers and, hydroxypropyl methyl cellulose (HPMC). Preferably the level of polymer in the film for example PVA is at least about 60%. Preferred average molecular weight will typically be about 20,000 to about 150,000. Films can also be of blended compositions comprising hydrolytically degradable and water soluble polymer blends such as polylactide and polyvinyl alcohol (known under the Trade reference M8630 as sold by MonoSol LLC, Indiana, USA) plus plasticisers like glycerol, ethylene glycerol, propylene glycol, sorbitol and mixtures thereof. The pouches can comprise a solid laundry cleaning composition or part components and/or a liquid cleaning composition or part components separated by the water soluble film. The compartment for liquid components can be different in composition than compartments containing solids: US2009/0011970 A1.

[0191] Detergent ingredients can be separated physically from each other by compartments in water dissolvable pouches or in different layers of tablets. Thereby negative storage interaction between components can be avoided. Different dissolution profiles of each of the compartments can also give rise to delayed dissolution of selected components in the wash solution.

[0192] A liquid or gel detergent, which is not unit dosed, may be aqueous, typically containing at least 20% by weight and up to 95% water, such as up to about 70% water, up to about 65% water, up to about 55% water, up to about 45% water, up to about 35% water. Other types of liquids, including without limitation, alkanols, amines, diols, ethers and polyols may be included in an aqueous liquid or gel. An aqueous liquid or gel detergent may contain from 0-30% organic solvent. A liquid or gel detergent may be non-aqueous.

Uses

[0193] The present invention is also directed to methods for using a composition, as defined herein, comprising a *Staphylococcus* hexosaminidase, e.g. dispersin of the invention and compositions hereof. A *Staphylococcus* hexosaminidase of the invention is useful in cleaning processes typically in laundry/textile/fabric (House hold laundry washing, Industrial laundry washing) or hard surface cleaning (ADW, car wash, Industrial surface).

[0194] One aspect of the invention relates to the use of a composition, as defined herein, comprising a *Staphylococcus* hexosaminidase e.g. dispersin comprising the motif D[IV]AR[TK] (SEQ ID NO 12) for cleaning of an item, wherein the item is a textile or a surface.

[0195] One aspect of the invention relates to the use of a composition, as defined herein, comprising a *Staphylococcus* hexosaminidase e.g. dispersin for cleaning of an item, wherein the item is a textile or a surface, wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ ID NO 3, SEQ ID NO 6, or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto.

[0196] One aspect of the invention relates to the use of a composition, as defined herein, comprising a *Staphylococcus* hexosaminidase of the invention,

- a) for preventing, reducing or removing stickiness of the item;
 - b) for pretreating stains on the item;
 - c) for preventing, reducing or removing redeposition of soil during a wash cycle;
 - d) for preventing, reducing or removing adherence of soil to the item;
 - e) for maintaining or improving whiteness of the item;
 - f) for preventing, reducing or removal malodor from the item,
- wherein the item is a textile.

[0197] One aspect of the invention relates to the use of a composition, as defined herein, comprising a *Staphylococcus* hexosaminidase, wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ ID NO 3, SEQ ID NO 6, or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%,

at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto,

- a) for preventing, reducing or removing stickiness of the item;
 - b) for pretreating stains on the item;
 - c) for preventing, reducing or removing redeposition of soil during a wash cycle;
 - d) for preventing, reducing or removing adherence of soil to the item;
 - e) for maintaining or improving whiteness of the item;
 - f) for preventing, reducing or removal malodor from the item,
- wherein the item is a textile.

[0198] The detergent composition of the present invention may be formulated, for example, as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations, or be formulated for hand or machine dishwashing operations. In a specific aspect, the present invention provides a detergent additive comprising one or more enzymes as described herein.

[0199] One aspect of the invention relates to the use of a composition, as defined herein, preferably a detergent composition comprising a *Staphylococcus* hexosaminidase e.g. dispersin comprising the motif D[IV]AR[TK] (SEQ ID NO 12) for cleaning of an item, wherein the item is a textile or a surface.

[0200] One aspect of the invention relates to the use of a composition, as defined herein, preferably a detergent composition comprising a *Staphylococcus* hexosaminidase e.g. dispersin for cleaning of an item, wherein the item is a textile or a surface, wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ ID NO 3, SEQ ID NO 6, or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto.

[0201] One aspect of the invention relates to the use of a composition, as defined herein, preferably a detergent composition comprising a *Staphylococcus* hexosaminidase e.g. dispersin of the invention,

- a) for preventing, reducing or removing stickiness of the item;
 - b) for pretreating stains on the item;
 - c) for preventing, reducing or removing redeposition of soil during a wash cycle;
 - d) for preventing, reducing or removing adherence of soil to the item;
 - e) for maintaining or improving whiteness of the item;
 - f) for preventing, reducing or removal malodor from the item,
- wherein the item is a textile.

[0202] One aspect of the invention relates to the use of a composition, as defined herein, preferably a detergent composition comprising a *Staphylococcus* hexosaminidase, e.g. dispersin wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ ID NO 3, SEQ ID NO 6, or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto,

- a) for preventing, reducing or removing stickiness of the item;
 - b) for pretreating stains on the item;
 - c) for preventing, reducing or removing redeposition of soil during a wash cycle;
 - d) for preventing, reducing or removing adherence of soil to the item;
 - e) for maintaining or improving whiteness of the item;
 - f) for preventing, reducing or removal malodor from the item,
- wherein the item is a textile.

Methods

[0203] The invention further relates to a method of treating a fabric comprising;

- (a) contacting the fabric with a composition of the invention or an aqueous solution thereof;
- (b) and optionally rinsing and drying the textile.

[0204] One aspect relates to a method of treating a fabric comprising;

- (a) contacting the fabric with a composition of the invention or an aqueous solution thereof, said composition com-

prising a *Staphylococcus* hexosaminidase, e.g. dispersin wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ ID NO 3, SEQ ID NO 6, or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto;

(b) and optionally rinsing and drying the textile.

[0205] The invention further relates to a method for cleaning or laundering an item comprising the steps of:

- a. exposing an item to a composition of the invention or a wash liquor comprising such a composition, said composition comprising a *Staphylococcus* hexosaminidase e.g. dispersin comprising the motif D[IV]AR[TK] (SEQ ID NO 12);
- b. completing at least one wash cycle; and
- c. optionally rinsing the item,

wherein the item is a fabric.

[0206] The invention further relates to a method for cleaning or laundering an item comprising the steps of:

- a. exposing an item to a composition of the invention or a wash liquor comprising a composition of the invention, said composition comprising a *Staphylococcus* hexosaminidase e.g. dispersin, wherein the *Staphylococcus* hexosaminidase is selected from the group consisting of polypeptides shown in SEQ ID NO 3, SEQ ID NO 6, or polypeptides having at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or such as at least 99% sequence identity hereto;
- b. completing at least one wash cycle; and
- c. optionally rinsing the item, wherein the item is a fabric.

[0207] One embodiment relates to a method, wherein the *Staphylococcus* hexosaminidase e.g. dispersin comprises one or more of the following motifs GXDE (SEQ ID NO 9), [EQ][NRSHA][YVFL][AGSTC][IVLF][EAQYN][SN] (SEQ ID NO 10), or [VIMS][LIV]G[GAV]DE[VI][PSA] (SEQ ID NO 11), in addition to D[IV]AR[TK] (SEQ ID NO 12).

[0208] According to the present invention, the method, wherein the *Staphylococcus* hexosaminidase comprises the motif D[IV]AR[TK] (SEQ ID NO 12).

[0209] The pH of the aqueous/liquid solution or wash liquor may be in the range of 1 to 11, such as in the range 5.5 to 11, such as in the range of 7 to 9, in the range of 7 to 8 or in the range of 7 to 8.5.

[0210] The wash liquor may have a temperature in the range of 5°C to 95°C, or in the range of 10°C to 80°C, in the range of 10°C to 70°C, in the range of 10°C to 60°C, in the range of 10°C to 50°C, in the range of 15°C to 40°C or in the range of 20°C to 30°C. In one aspect, the temperature of the wash liquor is 30°C.

[0211] The concentration of the *Staphylococcus* hexosaminidase in the wash liquor is typically in the range of at least 0.00001 ppm to at least 10 ppm, at least 0.00002 ppm to at least 10 ppm, at least 0.0001 ppm to at least 10 ppm, at least 0.0002 ppm to at least 10 ppm, at least 0.001 ppm to at least 10 ppm, at least 0.002 ppm to at least 10 ppm, at least 0.01 ppm to at least 10 ppm, at least 0.02 ppm to at least 10 ppm, at least 0.1 ppm to at least 10 ppm, at least 0.2 ppm to at least 10 ppm, at least 0.5 ppm to at least 5 ppm.

Definitions

[0212] Biofilm may be produced by any group of microorganisms in which cells stick to each other or stick to a surface, such as a textile, dishware or hard surface or another kind of surface. These adherent cells are frequently embedded within a self-produced matrix of extracellular polymeric substance (EPS). Biofilm EPS is a polymeric conglomeration generally composed of extracellular DNA, proteins, and polysaccharides. Biofilms may form on living or non-living surfaces. The microbial cells growing in a biofilm are physiologically distinct from planktonic cells of the same organism, which, by contrast, are single-cells that may float or swim in a liquid medium. Bacteria living in a biofilm usually have significantly different properties from planktonic bacteria of the same species, as the dense and protected environment of the film allows them to cooperate and interact in various ways. One benefit of this environment for the microorganisms is increased resistance to detergents and antibiotics, as the dense extracellular matrix and the outer layer of cells protect the interior of the community. On laundry biofilm producing bacteria can be found among the following species: *Acinetobacter* sp., *Aeromicrobium* sp., *Brevundimonas* sp., *Microbacterium* sp., *Micrococcus luteus*, *Pseudomonas* sp., *Staphylococcus epidermidis*, and *Stenotrophomonas* sp. On hard surfaces biofilm producing bacteria can be found among the following species: *Acinetobacter* sp., *Aeromicrobium* sp., *Brevundimonas* sp., *Microbacterium* sp., *Micrococcus luteus*, *Pseudomonas* sp., *Staphylococcus epidermidis*, *Staphylococcus aureus* and *Stenotrophomonas* sp. In one aspect, the biofilm producing strain is *Brevundimonas* sp. In one aspect, the biofilm producing strain is *Pseudomonas*, e.g. *Pseudomonas alcaliphila* or *Pseudomonas fluorescens*. In one aspect, the biofilm producing strain is *Staphylococcus*

aureus.

[0213] By the term "deep cleaning" is meant disruption or removal of components of organic matter, e.g. biofilm, such as polysaccharides, e.g. PNAG, proteins, DNA, soil or other components present in the organic matter.

[0214] Cleaning component or cleaning adjunct: The Cleaning component or cleaning adjunct is different from the *Staphylococcus* hexosaminidase. The precise nature of these cleaning (adjunct) components, and levels of incorporation thereof, will depend on the physical form of the composition and the nature of the operation for which it is to be used. Suitable cleaning components include, but are not limited to the components described below such as surfactants, builders, flocculating aid, chelating agents, dye transfer inhibitors, enzymes, enzyme stabilizers, enzyme inhibitors, catalytic materials, bleach activators, hydrogen peroxide, sources of hydrogen peroxide, preformed peracids, polymeric agents, clay soil removal/anti-redeposition agents, brighteners, suds suppressors, dyes, perfumes, structure elasticizing agents, fabric softeners, carriers, hydrotropes, builders and co-builders, fabric hueing agents, anti-foaming agents, dispersants, processing aids, and/or pigments.

[0215] Cleaning Composition: The term "cleaning composition" refers to compositions that find use in the removal of undesired compounds from items to be cleaned, such as textiles. The detergent composition may be used to e.g. clean textiles for both household cleaning and industrial cleaning. The terms encompass any materials/compounds selected for the particular type of cleaning composition desired and the form of the product (e.g., liquid, gel, powder, granulate, paste, or spray compositions) and includes, but is not limited to, detergent compositions (e.g., liquid and/or solid laundry detergents and fine fabric detergents; fabric fresheners; fabric softeners; and textile and laundry pre-spotters/pre-treatment). In addition to containing the enzymes described herein, the detergent formulation may contain one or more additional enzymes (such as proteases, amylases, lipases, cutinases, cellulases, endoglucanases, xyloglucanases, pectinases, pectin lyases, xanthanases, peroxidases, haloperoxygenases, catalases and mannanases, or any mixture thereof), and/or detergent adjunct ingredients such as surfactants, builders, chelators or chelating agents, bleach system or bleach components, polymers, fabric conditioners, foam boosters, suds suppressors, dyes, perfume, tannin inhibitors, optical brighteners, bactericides, fungicides, soil suspending agents, anti-corrosion agents, enzyme inhibitors or stabilizers, enzyme activators, transferase(s), hydrolytic enzymes, oxido reductases, bluing agents and fluorescent dyes, antioxidants, and solubilizers.

[0216] The term "hard surface cleaning" is defined herein as cleaning of hard surfaces wherein hard surfaces may include floors, tables, walls, roofs etc. as well as surfaces of hard objects such as cars (car wash) and dishes (dish wash). Dish washing includes but are not limited to cleaning of plates, cups, glasses, bowls, cutlery such as spoons, knives, forks, serving utensils, ceramics, plastics, metals, china, glass and acrylics.

[0217] The term "wash performance" is used as an enzyme's ability to remove stains present on the object to be cleaned during e.g. wash or hard surface cleaning.

[0218] The term "whiteness" is defined herein as the quality or state of a textile of being white. Loss of whiteness may be due to removal of optical brighteners/hueing agents and result in a greying or yellowing of the textiles. Greying and yellowing can be due to soil redeposition, body soils, colouring from e.g. iron and copper ions or dye transfer. Whiteness might include one or several issues from the list below: colourant or dye effects; incomplete stain removal (e.g. body soils, sebum etc.); redeposition (greying, yellowing or other discolourations of the object) (removed soils reassociate with other parts of textile, soiled or unsoiled); chemical changes in textile during application; and clarification or brightening of colours.

[0219] The term "laundering" relates to both household laundering and industrial laundering and means the process of treating textiles with a solution containing a cleaning or detergent composition of the present invention. The laundering process can for example be carried out using e.g. a household or an industrial washing machine or can be carried out by hand.

[0220] By the term "malodor" is meant an odor which is not desired on clean items. The cleaned item should smell fresh and clean without malodors adhered to the item. One example of malodor is compounds with an unpleasant smell, which may be produced by microorganisms. Another example of unpleasant smell can be sweat or body odor adhered to an item which has been in contact with human or animal. Another example of malodor can be the odor from spices, which sticks to items for example curry or other exotic spices which smell strongly, tobacco, cooking smell (fried oil, fish etc.), scents of perfume such as deodorant and eau de cologne.

[0221] The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminus processing, C-terminus truncation, glycosylation, phosphorylation, etc.

[0222] The term "textile" means any textile material including yarns, yarn intermediates, fibers, non-woven materials, natural materials, synthetic materials, and any other textile material, fabrics made of these materials and products made from fabrics (e.g., garments and other articles). The textile or fabric may be in the form of knits, wovens, denims, non-wovens, felts, yarns, and towelling. The textile may be cellulose based such as natural cellulose, including cotton, flax/linen, jute, ramie, sisal or coir or manmade cellulose (e.g. originating from wood pulp) including viscose/rayon, cellulose acetate fibers (tricell), lyocell or blends thereof. The textile or fabric may also be non-cellulose based such as natural polyamides including wool, camel, cashmere, mohair, rabbit and silk or synthetic polymers such as nylon, aramid,

polyester, acrylic, polypropylene and spandex/elastane, or blends thereof as well as blends of cellulose based and non-cellulose based fibers. Examples of blends are blends of cotton and/or rayon/viscose with one or more companion material such as wool, synthetic fiber (e.g. polyamide fiber, acrylic fiber, polyester fiber, polyvinyl chloride fiber, polyurethane fiber, polyurea fiber, aramid fiber), and/or cellulose-containing fiber (e.g. rayon/viscose, ramie, flax/linen, jute, cellulose acetate fiber, lyocell). Fabric may be conventional washable laundry, for example stained household laundry. When the term fabric or garment is used it is intended to include the broader term textiles as well.

[0223] The term "variant" means a polypeptide having the activity of the parent or precursor polypeptide and comprising an alteration, i.e., a substitution, insertion, and/or deletion, at one or more (e.g., several) positions compared to the precursor or parent polypeptide. A substitution means replacement of the amino acid occupying a position with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding an amino acid adjacent to and immediately following the amino acid occupying a position.

[0224] Sequence identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "sequence identity". For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends Genet. 16: 276-277), preferably version 6.6.0 or later. The parameters used are a gap open penalty of 10, a gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment}).$$

[0225] Clade: a group of polypeptides clustered together based on homologous features traced to a common ancestor. Polypeptide clades can be visualized as phylogenetic trees and a clade is a group of polypeptides that consists of a common ancestor and all its lineal descendants

Nomenclature

[0226] For purposes of the present invention, the nomenclature [IV] or [I/V] means that the amino acid at this position may be isoleucine (Ile, I) or valine (Val, V). Likewise, the nomenclature [LVI] and [L/V/I] means that the amino acid at this position may be a leucine (Leu, L), valine (Val, V) or isoleucine (Ile, I), and so forth for other combinations as described herein. Unless otherwise limited further, the amino acid X is defined such that it may be any of the 20 natural amino acids.

[0227] Unless otherwise indicated, or if it is apparent from the context that something else is intended, all percentages are percentage by weight (% w/w).

Examples

Assays

Wash assay

Mini Launder-O-Meter (MiniLOM) Model Wash System

[0228] MiniLOM is a mini wash system in which washes are performed in 50 ml test tubes placed in a Stuart rotator. Each tube simulates one small washing machine and during an experiment, each will contain a solution of a specific detergent/enzyme system to be tested along with the soiled and unsoiled fabrics it is tested on. Mechanical stress is achieved via rotation (typically 20rpm), and the temperature is controlled by placement of the rotator in a heating cabinet/room.

Assay I: testing of hexosaminidase activity

[0229] The hexosaminidase activity of the polypeptides listed in the table below was determined using 4-Methylumbelliferyl N-acetyl-β-D-glucosaminide (Sigma-Aldrich) as substrate. The enzymatic reaction was performed in triplicates in a 96 well flat bottom polystyrene microtiter plate (Thermo Scientific) with the following conditions: 20 mM 3-(N-morpholino)propanesulfonic acid pH 7 buffer, 5 mM 4-Methylumbelliferyl N-acetyl-β-D-glucosaminide, 0.01 vol% Brij 35 (Polyoxyethylene lauryl ether, CAS 9002-92-0) detergent and 50 nM purified enzyme sample in a total reaction volume of 200 μl. Blank samples without polypeptide were run in parallel. The reactions were carried out at room temperature

using a SpectraMax M2e Microplate Reader from Molecular Devices. Excitation wavelength was set to 368 nm and emission wavelength to 448 nm. Fluorescent signal was followed for 15 min in Kinetic Mode. Initial rate of reaction was evaluated in units of RFU/min by calculating the maximum initial increase in fluorescent signal over time as 4-Methylumbelliferyl was released from 4-Methylumbelliferyl N-acetyl- β -D-glucosaminide substrate due to enzymatic reaction.

Composition of model detergent A (liquid)

[0230] Ingredients: 12% LAS, 11% AEO Biosoft N25-7 (NI), 5% AEOS (SLES), 6% MPG (monopropylene glycol), 3% ethanol, 3% TEA, 2.75% coco soap, 2.75% soya soap, 2% glycerol, 2% sodium hydroxide, 2% sodium citrate, 1% sodium formate, 0.2% DTMPA and 0.2% PCA (all percentages are w/w).

[0231] Triple-20 Nonionic Model Detergent was prepared by dissolving 3.33 g/l non-ionic detergent containing NaOH 0.87%, MPG (Monopropylenglycol) 6%, Glycerol 2%, Soap-soy 2.75%, Soap-coco 2.75%, PCA (Sokalon CP-5) 0.2%, AEO Biosoft N25-7(NI) 16%, Sodium formate 1%, Sodium Citrate 2%, DTMPA 0.2%, Ethanol (96%) 3 %, adjustment of pH with NaOH or Citric acid ass water to 100% (all percentages are w/w (weight volume) in water with hardness 15 dH.

Example 1: Strain and DNA

[0232] The gene sequence encoding the hexosaminidase polypeptides (SEQ ID 2 and 5) from the strains *Staphylococcus cohnii* subsp. *cohnii* and *Staphylococcus fleurettii* respectively were found in the public database (Accession number SWISSPROT:A0A0M2NY11 and EMBLWGS:LAKJ01000034 for SEQ ID 1 and SWISSPROT:A0A1T1GHQ2 and EMBLWGS:MWJ01000007 for SEQ ID 4). The codon optimized synthetic DNA encoding the mature peptide sequences of the two hexosaminidases were ordered from the company Genearth. The mature polypeptides are shown in SEQ ID 3 and 6.

Table 1:

SEQ ID	donor	country of origen
SEQ ID 3	<i>Staphylococcus cohnii</i> subsp. <i>cohnii</i>	United Kingdom
SEQ ID 6	<i>Staphylococcus fleurettii</i>	Germany

Example 2: Cloning and expression of glycol_hydro_20 hexosaminidases

[0233] The codon optimized synthetic genes encoding the mature peptide sequences of the hexosaminidase with SEQ ID 3 and 6 were inserted into a *Bacillus* expression vector as described in WO12/025577. Briefly, the DNA encoding the mature peptide of the glycol_hydro_20 hexosaminidase gene was cloned in frame to a *Bacillus clausii* secretion signal (BcSP; with the following amino acid sequence: MKKPLGKIVASTALLISVAFSSSIASA (SEQ ID NO: 7). BcSP replaced the native secretion signal in the gene. Downstream of the BcSP sequence, an affinity tag sequence was introduced to ease the purification process (His-tag; with the following amino acid sequence: HHHHHHPR (SEQ ID NO: 8) The gene that was expressed therefore comprised the BcSP sequence followed by the His-tag sequence followed by the mature wild type glycol_hydro_20 sequence. The final expression plasmid (BcSP-His-tag- glycol_hydro_20) was transformed into a *Bacillus subtilis* expression host. The glycol_hydro_20 BcSP-fusion gene was integrated by homologous recombination into the *Bacillus subtilis* host cell genome upon transformation. The gene construct was expressed under the control of a triple promoter system (as described in WO99/43835). The gene coding for chloramphenicol acetyltransferase was used as maker (as described in (Diderichsen et al., 1993, Plasmid 30: 312-315)). Transformants were selected on LB media agar supplemented with 6 micrograms of chloramphenicol per ml. One recombinant *Bacillus subtilis* clone containing the glycol_hydro_20 expression construct was selected and was cultivated on a rotary shaking table in 500 ml baffled Erlenmeyer flasks each containing 100 ml yeast extract-based media. After 3-5 days' cultivation time at 30 °C to 37°C, the enzyme containing supernatant was harvested by centrifugation and the enzymes was purified by His-tag purification.

Example 3: His tag purification method

[0234] The His-tagged glycol_hydro_20 hexosaminidase enzymes were purified by immobilized metal chromatography (IMAC) using Ni^{2+} as the metal ion on 5 mL HisTrap Excel columns (GE Healthcare Life Sciences). The purification took place at pH 7 and the bound protein was eluted with imidazole. The purity of the purified enzymes was checked by SDS-PAGE and the concentration of the enzyme determined by Absorbance 280 nm after a buffer exchange in 50mM HEPES, 100mM NaCl pH7.0.

SEQ ID NO 7: MKKPLGKIVASTALLISVAFSSSIASA

SEQ ID NO 8: HHHHHHPR

Example 4: Biofilm growth and detachment assay

[0235] *Staphylococcus aureus* 15981 was kindly provided by Iñigo Lasa (Valle et al., Mol Microbiol.2003 May; 48 (4):1075-87). The strain was grown on trypticase soy agar (TSA) at 37°C overnight. Next day, a single colony was transferred to 15 ml trypticase soy broth (TSB) and incubated 5 hours at 37°C under shaking. The culture was diluted 1:100 in TSB+1% glucose and 100 µL of the bacterial suspension was transferred to each well of a 96-well microtiter plates (Thermo Scientific, Nunclon Delta Surface, cat # 167008) and incubated 24 hours at 37°C without shaking. and 100 µL of the bacterial suspension was transferred to each well of a 96-well microtiter plates (Thermo Scientific, Nunclon Delta Surface, cat # 167008) and incubated 24 hours at 37°C without shaking. Supernatant was aspirated and wells were washed with 100 µL of 0.9% sodium chloride and filled with 100 µL of either hard water or 3.3 gr/L non-ionic detergent or 3.3 gr/L model A detergent (composition hard water and non-ionic and model A) containing 0 (control) or 20, 10, 5, 2.5, 1.25, 0.62, 0.31, 0.16, 0.08, 0.04, 0.02 and 0.01 µg/mL of enzyme (SEQ ID 3 and 6). After incubation at 37°C for 1 hour, wells were washed with water and stained for 15 min with 100 µL of 0.095% crystal violet solution (SIGMA V5265). Wells were then rinsed twice with 100 µL water, dried and the plates were scanned. The lowest concentration of each enzyme that could remove the visible formation of biofilm of the *S. aureus* 15981 organism after 1 hour incubation, in the presence and absence of detergent was determined (see Table 2). All enzymes were assayed per duplicate in three independent assays. The average of the minimal concentration of enzyme that removed the visible formation of *S. aureus* 15981 from the three assays is listed in Table 2.

Table 2. Minimal concentration of enzyme that can remove the visible formation of *S. aureus* 15981 after 1 hour incubation in either hard water or model A detergent

SEQ ID	Minimal concentration for biofilm removal in hard water µg/mL	Minimal concentration for biofilm removal in non-ionic detergent µg/mL	Minimal concentration for biofilm removal in model A detergent µg/mL
3	0,64	0,94	6,67
6	0,07	0,21	0,41

Example 5. cleaning properties of hexosaminidase in liquid model detergent

[0236] A crude extract of biofilm extracellular polymeric substances (EPS) was prepared from *Staphylococcus aureus* 15981 (kind gift from Iñigo Lasa (Valle, J., A. Toledo-Arana, C. Berasain, J. M. Ghigo, B. Amorena, J. R. Penades, and I. Lasa. 2003, Mol. Microbiol. 48:1075-1087) as follows: 500 mL of TSB + 1% glucose (24563; Roquette Freres) was inoculated, aliquoted into 50ml conical centrifuge tubes (339652; Thermo Scientific Nunc) and incubated for 24 hours at 37°C under shaking conditions (200 rpm). Following incubation, the cells were pelleted by centrifugation (10min, 6000g, 25°C), pooled and resuspended in 4ml 3M NaCl. The suspension was vortexed vigorously and incubated for 15min at ambient temperature to extract the surface-associated EPS. The cells were then re-pelleted (10min, 5000g, 25°C) and the EPS-containing supernatant was retrieved. Milli-Q water was added (6ml) and the solution was sterile-filtered twice (0.45 µm followed by 0.2 µm). The crude extract was stored at -20°C until further use. For wash performance testing, 50ul aliquots of the crude EPS extract was spotted on sterile textile swatches (WFK20A) and incubated for 15 min at ambient temperature. The swatches (sterile or with EPS) were placed in 50 mL test tubes and 10 mL of wash liquor (15⁰dH water with 0.2 g/L iron(III) oxide nano-powder (544884; Sigma-Aldrich) with 3.33g/L liquid model A detergent or 3.33 g/L nonionic model detergent) and enzyme was added to each tube. Washes without enzyme were included as controls. The test tubes were placed in a Stuart rotator and incubated for 1 hour at 37°C at 20rpm. The wash liquor was then removed, and the swatches were rinsed twice with 15⁰dH water and left to dry on filter paper overnight. The remission (REM^{460nm}) values were measured using a Macbeth Color-Eye 7000 (CE7000), and are displayed in table 3. Delta values (REM^{460nm}_(washed with enzyme) - REM^{460nm}_(washed without enzyme)) are also indicated.

Table 3 cleaning effects of hexosaminidases in liquid model detergent

			Model detergent A		Nonionic model detergent	
Swatch	Enzyme	Enzyme concentration (µg/ml)	REM (460nm)	ΔREM	REM(460nm)	ΔREM
wfk20A	No enzyme	0,0	58,5		59,7	

(continued)

			Model detergent A		Nonionic model detergent	
Swatch	Enzyme	Enzyme concentration (μg/ml)	REM (460nm)	ΔREM	REM(460nm)	ΔREM
EPS	No enzyme	0,0	30,1		30,4	
EPS	SEQ ID NO 3	2,0	50,1	20,0	60,0	29,6
EPS	SEQ ID NO 3	0,2	42,0	11,9	60,9	30,5
EPS	SEQ ID NO 6	2,0	62,5	32,4	59,7	29,2
EPS	SEQ ID NO 6	0,2	62,9	32,8	62,0	31,5

Example 5: Construction of clades and phylogenetic trees

[0237] The Glyco_hydro_20 domain includes the polypeptides having hexosaminidase e.g. PNAG activity and comprises the ENYA, VLG and/or DIARK clades. According to the invention, the *Staphylococcus* hexosaminidase comprises the motif D[IV]AR[TK] (SEQ ID NO 12).

[0238] A phylogenetic tree was constructed, of polypeptide sequences containing a Glyco_hydro_20 domain, as defined in PFAM (PF00728, Pfam version 31.0 Finn (2016). Nucleic Acids Research, Database Issue 44:D279-D285). The phylogenetic tree was constructed from a multiple alignment of mature polypeptide sequences containing at least one Glyco_hydro_20 domain. The sequences were aligned using the MUSCLE algorithm version 3.8.31 (Edgar, 2004. Nucleic Acids Research 32(5): 1792-1797), and the trees were constructed using FastTree version 2.1.8 (Price et al., 2010, PloS one 5(3)) and visualized using iTOL (Letunic & Bork, 2007. Bioinformatics 23(1): 127-128). The polypeptide sequences containing a Glyco_hydro_20 domain comprises several motifs; one example is GXDE (SEQ ID NO 9), situated in positions 157 to 160 in *Staphylococcus cohnii subsp. cohnii* (SEQ ID NO 3). Residues D and E are the key catalytic residues of Glyco_hydro_20 enzymes (position 159 to 160 in SEQ ID NO 3).

[0239] The polypeptides in Glyco_hydro_20 can be separated into multiple distinct sub-clusters, or clades as listed below. The distinct motifs for each clade are described in detail below.

Generation of ENYA clade

[0240] A clade, preferably shared by the polypeptides of the invention, was identified. This clade has not been described previously. The clade is termed IES and polypeptides of this clade comprises Glyco_hydro_20 domain polypeptides of bacterial origin and are in addition to having PNAG activity, characterized by comprising certain motifs. The polypeptides of the clade comprise the motif example [EQ][NRSHA][YVFL][AGSTC][IVLF][EAQYN][SN] (SEQ ID NO 10), corresponding to ENYAIES at position 44 to 50 of SEQ ID NO 3.

Generation of VLG clade

[0241] A clade, preferably shared by the polypeptides of the invention, was identified. This clade has not been described previously. The clade is termed VLG and polypeptides of this clade comprise Glyco_hydro_20 domain polypeptides of bacterial origin and are in addition to having PNAG activity, characterized by comprising certain motifs. The polypeptides of the clade comprise the motif example [VIMS][LIV]G[GAV]DE[VI][PSA] (SEQ ID NO 11), corresponding to VLGGEVP (positions 155 to 162 of SEQ ID NO 3), where G and DE (corresponding to positions 157 and 159-160 of SEQ ID NO 3) are fully conserved in VLG clade and part of the active site. Residues D and E are the key catalytic residues of Glyco_hydro_20 enzymes (position 159 to 160 in SEQ ID NO 3).

Generation of DIARK clade

[0242] The DIARK clade comprises VLG domain polypeptides of bacterial origin, having hexosaminidase e.g. PNAG activity. The polypeptides of the clade comprise the motif example D[IV]AR[TK] (SEQ ID NO 12), corresponding to positions 10 to 14 of SEQ ID NO 3, where D and AR are fully conserved in DIARK clade (positions 10 and 12-13 in SEQ ID NO 3). According to the present invention, the *Staphylococcus* hexosaminidase comprises the motif D[IV]AR[TK] (SEQ ID NO 12).

[0243] An alignment of the polypeptides of the invention is shown in Figure 2. A phylogenetic tree of the polypeptides of the invention is shown in Figure 1.

Example 6 characterization of dispersinsDispersin activity as a function of pH

[0244] *Activity assay:* The activity of the dispersin having SEQ ID NO 6 was measured with 4-Nitrophenyl N-acetyl- β -D-glucosaminide (4-NAG, CAS Number 3459-18-5, CHE00244) as substrate as a function of pH (4-10 in 1-unit increments). The concentrations of substrate and the dispersin having SEQ ID NO 6 were 1 mM and 1.0 μ M, respectively, in all measurements. The dilution buffers comprise: 50 mM MES (CAS Number: 4432-31-9), 50 mM glycine (CAS Number: 56-40-6), 50 mM acetic acid (CAS Number: 64-19-7) adjusted to pH 4-10.

[0245] The substrate solution (10 mM) was prepared by dissolving 34.23 mg 4-NAG in 10.0 mL water. Dissolution required rigorous vortex mixing and gentle heating. The enzyme concentration was determined by UV-Vis (ϵ_{280} = 54760 M⁻¹cm⁻¹).

[0246] The enzyme samples were incubated at the different pH-values in volumes of 200 μ L in a thermomixer (in MTP) for 10 min and 500 rpm at 30 °C. After 10 min, the MTP was incubated at 95 °C and 500 rpm for 10 min in thermomixer to end the reaction. Then the samples were transferred to ice bath and cooled for 2 min. The samples were added 20 μ L 4 M NaOH to deprotonate pNP (induce yellow color). Absorbance at 405 nm was measured for 2 min in 10 sec. intervals. All measurements were produced in triplicates and reference samples were produced for all conditions (buffer instead of enzyme).

[0247] *Results:* The following table display the average absorbance (activity) subtracted the average absorbance of the reference samples measured after 10 min incubation at different pH values:

pH	4	5	6	7	8	9	10
A₄₀₅	0.370009	0.831697	0.911115	0.061912	-0.005915	-0.053345	-0.054276

Dispersin stability as a function of pH and NaCl

[0248] *Stability assay - differential scanning fluorimetry:* The thermal stability of the dispersin having SEQ ID NO 6 was measured as a function of pH (4, 6, 7, 8, 10) and NaCl concentration (100, 200, and 300 mM). The thermal unfolding was monitored using intrinsic fluorescence utilizing a Prometheus NT.48. The concentrations the dispersin having SEQ ID NO 6 was 0.2 mg/mL in all measurements. The enzyme concentration was determined by UV-Vis (ϵ_{280} = 54760 M⁻¹cm⁻¹). The dilution buffers comprise: 50 mM MES (CAS Number: 4432-31-9), 50 mM glycine (CAS Number: 56-40-6), 50 mM acetic acid (CAS Number: 64-19-7) adjusted to pH 4, 6, 7, 8, or 10.

[0249] The enzyme samples were prepared by mixing a 5 M NaCl stock, buffer, water (MQ), and enzyme to obtain the desired concentrations. The total volume of each mixture was 100 μ L. The samples were loaded in the instrument in duplicates and measured from 20 to 95 °C with temperature ramping of 2.0 °C/min.

[0250] *Results:* Melting temperatures (T_m -values) were derived from the thermograms using the PR.ThermControl v.2.0.4 software. The following table display the average T_m -values obtained at the different conditions:

		pH				
		4	6	7	8	10
[NaCl] mM	0	43.5	45.9	40.2	34.8	N/A
	100	40.4	46.6	41.2	35.5	N/A
	200	39.4	47.6	42.3	37.3	N/A
	300	39.1	48.4	43.1	38.4	N/A

Dispersin activity as a function of temperature

[0251] *Activity assay:* The activity of the dispersin having SEQ ID NO 6 was measured with 4-Nitrophenyl N-acetyl- β -D-glucosaminide (4-NAG, CAS Number 3459-18-5, CHE00244) as substrate at pH 7. The concentrations of substrate and the dispersin having SEQ ID NO 6 were 1 mM and 0.5 μ M, respectively, in all measurements. The dilution buffer comprises: 50 mM MES (CAS Number: 4432-31-9), 50 mM glycine (CAS Number: 56-40-6), 50 mM acetic acid (CAS Number: 64-19-7), pH 7.

[0252] The substrate solution (10 mM) was prepared by dissolving 35.9 mg 4-NAG in 10.482 mL water.

[0253] Dissolution required rigorous vortex mixing and gentle heating. The enzyme concentration was determined by UV-Vis (ϵ_{280} = 54760 M⁻¹cm⁻¹). The reaction mixture comprised 15.9 μ L enzyme (6.3 μ M), 20 μ L substrate, and 164.1

μL buffer.

[0254] The enzyme samples were incubated in volumes of 200 μL in a thermomixer for 10 min and 500 rpm at 20, 30, 40, 45, 50, 55, 60, or 70 °C. After 10 min, the samples were transferred to ice bath and cooled for 2 min. The samples were added 10 μL 4 M NaOH to deprotonate pNP (induce yellow color). 180 μL reaction mixture was transferred to a MT plate and absorbance at 405 nm was measured for 1 min in 10 sec. intervals. All measurements were produced in duplicates and reference samples were produced for all conditions (buffer instead of enzyme).

[0255] *Results:* The following table display the average absorbance (activity) subtracted the average absorbance of the reference samples measured after 10 min incubation at different temperatures:

	20 °C	30 °C	40 °C	45 °C	50 °C	55 °C	60 °C	70 °C
A₄₀₅	0.121286	0.2024	0.294707	0.184157	0.054971	0.035657	0.02815	0.020007

Claims

1. A composition comprising a *Staphylococcus* hexosaminidase, wherein the *Staphylococcus* hexosaminidase comprises the motif D[IV]AR[TK] (SEQ ID NO 12) and wherein the composition

(a) is a solid, preferably granular, laundry detergent composition and further comprises

- (a1) at least one zeolite builder, preferably in an amount of 10 to 50 wt.-%, more preferably 20-30 wt.-%;
- (a2) at least one phosphonate builder, preferably in an amount of 0.1 to 5 wt.-%, more preferably 0.4 to 1.5 wt.-%;
- (a3) at least one further enzyme, preferably a cellulase, preferably in an amount of active enzyme of 100 to 5000 ppb, more preferably 1000 to 2000 ppb; and
- (a4) at least one polymer, preferably a polyvinylpyrrolidon polymer, preferably in an amount of 0.01 to 1 wt.-%, more preferably 0.1 to 0.3 wt.-%; or

(b) is a solid laundry detergent composition and further comprises

- (b1) at least one silicate builder, preferably in an amount of 2 to 20 wt.-%, more preferably 5-10 wt.-%;
- (b2) optionally carboxymethylcellulose, preferably in an amount of 0.1 to 10 wt.-%, more preferably 0.1 to 4 wt.-%;
- (b3) at least one further enzyme, preferably a cellulase, preferably in an amount of active enzyme of 0.1 to 100 ppm, more preferably 0.1 to 10 ppm;
- (b4) optionally at least one soil release polymer, preferably a polyvinylpyrrolidon polymer, in an amount of 0.1 to 3 wt.-%, more preferably 0.1 to 1.0 wt.-%; and
- (b5) at least one bleaching system, comprising a bleaching agent, a bleach activator and a bleach catalyst, preferably in an amount of 0.1 to 50 wt.-%, more preferably 0.1 to 30 wt.-%; or

(c) is liquid laundry detergent composition and further comprises

- (c1) at least one surfactant, preferably nonionic surfactant, preferably in an amount of 1 to 20 wt.-%, preferably 3 to 15 wt.-%;
- (c2) optionally at least one phosphonate builder, preferably in an amount of 0.1 to 3 wt.-%, more preferably 0.25 to 1.5 wt.-%
- (c3) optionally at least at least one further enzyme, preferably a cellulase, preferably in an amount of enzyme composition of 0.001 to 1 wt.-%, more preferably 0.001 to 0.6 wt.-%; and
- (c4) optionally at least one organic solvent, preferably glycerol, preferably in an amount of 0.1 to 10 wt.-%, more preferably 0.1 to 5 wt.-%; or

(d) is a liquid laundry detergent in unit dose form, preferably a pouch comprising a water-soluble film, and further comprises

- (d1) water in an amount of up to 20 wt.-%, preferably 5 to 15 wt.-%;
- (d2) optionally at least one bittering agent, preferably Benzyldiethyl(2,6-xylylcarbamoyl)-methylammonium-benzoate, preferably in an amount of 0.00001 to 0.04 wt.-%;

(d3) optionally at least one optical brightener, preferably in an amount of 0.01 to 2 wt.-%, more preferably 0.01 to 1 wt.-%; and
(d4) optionally at least one polymer, preferably in an amount of 0.01 to 7 wt.-%, more preferably 0.1 to 5 wt.-%; or

(e) is a fabric finisher and further comprises

(e1) at least one softening silicone, preferably an amino-functionalized silicone, preferably in an amount of 0.1 to 10 wt.-%, more preferably 0.1 to 2 wt.-%;
(e2) at least one perfume, preferably at least partially encapsulated in microcapsules, more preferably at least partially encapsulated in melamine-formaldehyde microcapsules, preferably in an amount of 0.01 to 3 wt.-%, more preferably 0.1 to 1 wt.-%;
(e3) optionally polyquaternium 10 in an amount of 0.1 to 20 wt.-%, preferably 0.1 to 13 wt.-%;
(e4) optionally polyquaternium 37 in an amount of 0.1 to 20 wt.-%, preferably 0.1 to 13 wt.-%;
(e5) optionally a plant-based esterquat, preferably a canola- or palm-based esterquat, in an amount of 0.1 to 20 wt.-%, preferably 0.1 to 13 wt.-%; and
(e6) optionally adipic acid, in an amount of 0.1 to 20 wt.-%, preferably 0.1 to 13 wt.-%; or

(f) is an acidic cleaning agent, preferably having a pH less than 6, and further comprises

(f1) plant-based or bio-based surfactants, preferably each in an amount of 0.1 to 5, more preferably each in an amount of 0.1 to 2 wt.-%;
(f2) at least one acidic biocide, preferably selected from acids, more preferably HCl and formic acid; and
(f3) at least one soil release, water repellent or water spreading polymer, preferably in an amount of 0.01 to 3 wt.-%, more preferably 0.01 to 0.5 wt.-%; or

(g) is a neutral cleaning agent, preferably having a pH between 6.0 and 7.5, and further comprises

(g1) plant-based or bio-based surfactants, preferably each in an amount of 0.1 to 5, more preferably each in an amount of 0.1 to 2 wt.-%;
(g2) at least one biocide, preferably selected from quaternary ammonium compounds and alcohols; and
(g3) at least one soil release, water repellent or water spreading polymer, preferably in an amount of 0.01 to 3 wt.-%, more preferably 0.01 to 0.5 wt.-%; or

(h) is an alkaline cleaning agent, preferably having a pH of more than 7.5, and further comprises (h1) plant-based or bio-based surfactants, preferably each in an amount of 0.1 to 5, more preferably each in an amount of 0.1 to 2 wt.-%; or

(i) is a hand dishwashing agent, preferably liquid hand dishwashing agent, and further comprises (i1) at least one anionic surfactant, preferably in an amount of 0.1 to 40 wt.-%, more preferably 5 to 30 wt.-%;

(i2) at least one amphoteric surfactant, preferably betain, preferably in an amount of 0.1 to 25 wt.-%, more preferably 1 to 15 wt.-%;
(i3) at least one nonionic surfactant, preferably in an amount of 0.1 to 25 wt.-%, more preferably 2 to 10 wt.-%;
(i4) at least one further enzyme, preferably selected from proteases, amylases and combinations thereof, preferably in an amount of enzyme composition of up to 1 wt.-%, more preferably up to 0.6 wt.-%; or

(j) is an automatic dishwashing composition and further comprises

(j1) at least one builder selected from citrate, aminocarboxylates and combinations thereof, preferably in an amount of 5 to 30 wt.-%, more preferably 10 to 20 wt.-%;
(j2) at least one phosphonate builder, preferably in an amount of 0.1 to 5 wt.-%, more preferably 0.4 to 1.5 wt.-%;
(j3) at least one nonionic surfactant, preferably in an amount of 0.1 to 10 wt.-%, more preferably 1 to 5 wt.-%;
(j4) at least one bleaching system, comprising a bleaching agent, a bleach activator and a bleach catalyst, preferably in an amount of 0.1 to 50 wt.-%, more preferably 0.1 to 30 wt.-%; and
(j5) at least one polymer selected from sulfopolymers, cationic polymers and polyacrylates, preferably in an amount of 0.01 to 15 wt.-%, more preferably 2 to 10 wt.-%.

2. A composition according to claim 1, wherein the composition further comprises;

(a)

- i. one or more polyol(s), preferably selected from glycerol, (mono, di, or tri) propylene glycol, ethylene glycol, polyethylene glycol, sugar alcohols, sorbitol, mannitol, erythritol, dulcitol, inositol, xylitol and adonitol,
- ii. optionally one or more enzyme, preferably selected from proteases, amylases or lipases,
- iii. optionally one or more surfactant, preferably selected from anionic and nonionic surfactants,
- iv. optionally one or more polymer;

or

(b) a granule comprising

- i. a core comprising a *Staphylococcus* hexosaminidase and optionally,
- ii. a coating consisting of one or more layer(s) surrounding the core.

3. A composition according to claim 1 or claim 2, wherein the hexosaminidase has N-acetylglucosaminidase activity, preferably β -1,6 N-acetylglucosaminidase activity.

4. Composition according to any of the preceding claims, wherein the polypeptide having hexosaminidase activity comprises the amino acid sequence of SEQ ID NO 3 or polypeptides having at least 60 % e.g. 80%, 85%, 90%, 95%, 98% or 99% sequence identity hereto.

5. Composition according to any of the preceding claims, wherein the composition further comprises one or more enzymes selected from the group consisting of proteases, lipases, cutinases, amylases, carbohydrases, cellulases, pectinases, mannanases, arabinases, galactanases, xylanases and oxidases

6. Use of a composition according to any one of claims 1 to 5, preferably a detergent composition comprising a *Staphylococcus* hexosaminidase,

- a) for preventing, reducing or removing stickiness of the item;
 - b) for pretreating stains on the item;
 - c) for preventing, reducing or removing redeposition of soil during a wash cycle;
 - d) for preventing, reducing or removing adherence of soil to the item;
 - e) for maintaining or improving whiteness of the item;
 - f) for preventing, reducing or removing malodor from the item,
- wherein the item is a textile.

Patentansprüche

1. Zusammensetzung, umfassend *Staphylococcus*-Hexosaminidase, wobei *Staphylococcus*-Hexosaminidase das Motiv D[IV]AR[TK] (SEQ ID NO 12) umfasst und wobei die Zusammensetzung

(a) eine feste, vorzugsweise körnige Wäschewaschmittelzusammensetzung ist und ferner umfasst

- (a1) mindestens eine Zeolithbuildersubstanz, vorzugsweise in einer Menge von 10 bis 50 Gew.-%, stärker bevorzugt 20-30 Gew.-%;
- (a2) mindestens eine Phosphonatbuildersubstanz, vorzugsweise in einer Menge von 0,1 bis 5 Gew.-%, stärker bevorzugt 0,4 bis 1,5 Gew.-%;
- (a3) mindestens ein weiteres Enzym, vorzugsweise eine Cellulase, vorzugsweise in einer Menge von aktivem Enzym von 100 bis 5000 ppb, stärker bevorzugt 1000 bis 2000 ppb; und
- (a4) mindestens ein Polymer, vorzugsweise ein Polyvinylpyrrolidonpolymer, vorzugsweise in einer Menge von 0,01 bis 1 Gew.-%, stärker bevorzugt 0,1 bis 0,3 Gew.-%; oder

(b) eine feste Wäschewaschmittelzusammensetzung ist und ferner umfasst

- (b1) mindestens eine Silikatbuildersubstanz, vorzugsweise in einer Menge von 2 bis 20 Gew.-%, stärker

bevorzugt 5-10 Gew.-%;

(b2) optional Carboxymethylcellulose, vorzugsweise in einer Menge von 0,1 bis 10 Gew.-%, stärker bevorzugt 0,1 bis 4 Gew.-%;

(b3) mindestens ein weiteres Enzym, vorzugsweise eine Cellulase, vorzugsweise in einer Menge von aktivem Enzym von 0,1 bis 100 ppm, stärker bevorzugt 0,1 bis 10 ppm;

(b4) optional mindestens ein Schmutzfreisetzungspolymer, vorzugsweise ein Polyvinylpyrrolidonpolymer, in einer Menge von 0,1 bis 3 Gew.-%, stärker bevorzugt 0,1 bis 1,0 Gew.-%; und

(b5) mindestens ein Bleichsystem, umfassend ein Bleichmittel, einen Bleichaktivator und einen Bleichkatalysator, vorzugsweise in einer Menge von 0,1 bis 50 Gew.-%, stärker bevorzugt 0,1 bis 30 Gew.-%; oder

(c) eine flüssige Wäschewaschmittelzusammensetzung ist und ferner umfasst

(c1) mindestens ein Tensid, vorzugsweise ein nichtionisches Tensid, vorzugsweise in einer Menge von 1 bis 20 Gew.-%, stärker bevorzugt 3 bis 15 Gew.-%;

(c2) optional mindestens eine Phosphonatbuildersubstanz, vorzugsweise in einer Menge von 0,1 bis 3 Gew.-%, stärker bevorzugt 0,25 bis 1,5 Gew.-%;

(c3) optional mindestens ein weiteres Enzym, vorzugsweise eine Cellulase, vorzugsweise in einer Menge von Enzymzusammensetzung von 0,001 bis 1 Gew.-%, stärker bevorzugt 0,001 bis 0,6 Gew.-%; und

(c4) optional mindestens ein organisches Lösungsmittel, vorzugsweise ein Glycerin, vorzugsweise in einer Menge von 0,1 bis 10 Gew.-%, stärker bevorzugt 0,1 bis 5 Gew.-%; oder

(d) ein flüssiges Wäschewaschmittel in Einheitsdosisform ist, vorzugsweise ein Beutel, umfassend einen waserlöslichen Film und ferner umfasst

(d1) Wasser in einer Menge von bis zu 20 Gew.-%, vorzugsweise 5 bis 15 Gew.-%;

(d2) optional mindestens ein Bitterstoff, vorzugsweise Benzyldiethyl(2,6-xylylcarbamoil)-methylammoniumbenzoat, vorzugsweise in einer Menge von 0,00001 bis 0,04 Gew.-%;

(d3) optional mindestens einen optischen Aufheller, vorzugsweise in einer Menge von 0,01 bis 2 Gew.-%, stärker bevorzugt 0,01 bis 1 Gew.-%; und

(d4) optional mindestens ein Polymer, vorzugsweise in einer Menge von 0,01 bis 7 Gew.-%, stärker bevorzugt 0,1 bis 5 Gew.-%; oder

(e) ein Gewebeveredler ist und ferner umfasst

(e1) mindestens ein Weichmachersilikon, vorzugsweise ein aminofunktionalisiertes Silikon, vorzugsweise in einer Menge von 0,1 bis 10 Gew.-%, stärker bevorzugt 0,1 bis 2 Gew.-%;

(e2) mindestens einen Duftstoff, vorzugsweise mindestens teilweise in Mikrokapseln eingekapselt, stärker bevorzugt mindestens teilweise in MelaminFormaldehyd-Mikrokapseln eingekapselt, vorzugsweise in einer Menge von 0,01 bis 3 Gew.-%, stärker bevorzugt 0,1 bis 1 Gew.-%;

(e3) optional Polyquaternium 10 in einer Menge von 0,1 bis 20 Gew.-%, vorzugsweise 0,1 bis 13 Gew.-%;

(e4) optional Polyquaternium 37 in einer Menge von 0,1 bis 20 Gew.-%, vorzugsweise 0,1 bis 13 Gew.-%;

(e5) optional ein Esterquat auf pflanzlicher Basis, vorzugsweise ein Esterquat auf Raps- oder Palmbasis, in einer Menge von 0,1 bis 20 Gew.-%, vorzugsweise 0,1 bis 13 Gew.-%; und

(e6) optional Adipinsäure in einer Menge von 0,1 bis 20 Gew.-%, vorzugsweise 0,1 bis 13 Gew.-%; oder

(f) ein Säurereinigungsmittel ist, das vorzugsweise einen pH-Wert von weniger als 6 aufweist und ferner umfasst

(f1) Tenside auf pflanzlicher Basis oder auf Bio-Basis, vorzugsweise jeweils in einer Menge von 0,1 bis 5, stärker bevorzugt in einer Menge von 0,1 bis 2 Gew.-%;

(f2) mindestens ein saures Biozid, das vorzugsweise aus Säuren, stärker bevorzugt HCl und Ameisensäure ausgewählt ist; und

(f3) mindestens ein Schmutzfreisetzungs-, Wasserabweisungs- oder Wasserausbreitungspolymer, vorzugsweise in einer Menge von 0,01 bis 3 Gew.-%, stärker bevorzugt 0,01 bis 0,5 Gew.-%; oder

(g) ein neutrales Reinigungsmittel ist, das vorzugsweise einen pH-Wert zwischen 6,0 und 7,5 aufweist, und ferner umfasst (g1) Tenside auf pflanzlicher Basis oder auf Bio-Basis, vorzugsweise jeweils in einer Menge von 0,1 bis 5, stärker bevorzugt in einer Menge von 0,1 bis 2 Gew.-%;

(g2) mindestens ein Biozid, das vorzugsweise aus quaternären Ammoniumverbindungen und Alkoholen ausgewählt ist; und
(g3) mindestens ein Schmutzfreisetzungs-, Wasserabweisungs- oder Wasserausbreitungspolymer, vorzugsweise in einer Menge von 0,01 bis 3 Gew.-%, stärker bevorzugt 0,01 bis 0,5 Gew.-%; oder

(h) ein alkalisches Reinigungsmittel ist, das vorzugsweise einen pH-Wert von mehr als 7,5 aufweist, und ferner umfasst (h1) Tenside auf pflanzlicher Basis oder auf Bio-Basis, vorzugsweise jeweils in einer Menge von 0,1 bis 5, stärker bevorzugt in einer Menge von 0,1 bis 2 Gew.-%; oder

(i) ein Handgeschirrspülmittel ist, vorzugsweise ein flüssiges Handgeschirrspülmittel, und ferner umfasst

(i1) mindestens ein anionisches Tensid, vorzugsweise in einer Menge von 0,1 bis 40 Gew.-%, stärker bevorzugt 5 bis 30 Gew.-%;

(i2) mindestens ein amphoter Tensid, vorzugsweise Betain, vorzugsweise in einer Menge von 0,1 bis 25 Gew.-%, stärker bevorzugt 1 bis 15 Gew.-%;

(i3) mindestens ein nichtionisches Tensid, vorzugsweise in einer Menge von 0,1 bis 25 Gew.-%, stärker bevorzugt 2 bis 10 Gew.-%;

(i4) mindestens ein weiteres Enzym, das vorzugsweise aus Proteasen, Amylasen und Kombinationen davon ausgewählt ist, vorzugsweise in einer Menge von Enzymzusammensetzung von bis zu 1 Gew.-%, stärker bevorzugt bis zu 0,6 Gew.-%; oder

(j) eine automatische Geschirrspülzusammensetzung ist und ferner umfasst

(j1) mindestens eine Buildersubstanz, die aus Citrat, Aminocarboxylaten und Kombinationen davon ausgewählt ist, vorzugsweise in einer Menge von 5 bis 30 Gew.-%, stärker bevorzugt 10 bis 20 Gew.-%;

(j2) mindestens eine Phosphonatbuildersubstanz, vorzugsweise in einer Menge von 0,1 bis 5 Gew.-%, stärker bevorzugt 0,4 bis 1,5 Gew.-%;

(j3) mindestens ein nichtionisches Tensid, vorzugsweise in einer Menge von 0,1 bis 10 Gew.-%, stärker bevorzugt 1 bis 5 Gew.-%;

(j4) mindestens ein Bleichsystem, umfassend ein Bleichmittel, einen Bleichaktivator und einen Bleichkatalysator, vorzugsweise in einer Menge von 0,1 bis 50 Gew.-%, stärker bevorzugt 0,1 bis 30 Gew.-%; und

(j5) mindestens ein Polymer, das aus Sulfopolymeren, kationischen Polymeren und Polyacrylaten ausgewählt ist, vorzugsweise in einer Menge von 0,01 bis 15 Gew.-%, stärker bevorzugt 2 bis 10 Gew.-%.

2. Zusammensetzung nach Anspruch 1, wobei die Zusammensetzung ferner umfasst;

(a)

i. ein oder mehrere Polyol(e), die vorzugsweise aus Glycerin, (Mono-, Di- oder Tri-) Propylenglykol, Ethylenglykol, Polyethylenglykol, Zuckeralkoholen, Sorbit, Mannit, Erythrit, Dulcitol, Inositol, Xylitol und Adonit ausgewählt sind,

ii. optional ein oder mehrere Enzyme, die vorzugsweise aus Proteasen, Amylasen oder Lipasen ausgewählt sind,

iii. optional ein oder mehrere Tenside, die vorzugsweise aus anionischen und nichtionischen Tensiden ausgewählt sind,

iv. optional ein oder mehrere Polymere;
oder

(b) ein Korn, umfassend

i. einen Kern, umfassend: Staphylococcus-Hexosaminidase und optional,

ii. eine Beschichtung, bestehend aus einer oder mehreren Schicht(en), die den Kern umgeben.

3. Zusammensetzung nach Anspruch 1 oder 2, wobei die Hexosaminidase N-Acetylglucosaminidase-Aktivität, vorzugsweise β -1,6-N-Acetylglucosaminidase-Aktivität, aufweist.

4. Zusammensetzung nach einem der vorstehenden Ansprüche, wobei das Polypeptid, das Hexosaminidase-Aktivität aufweist, die Aminosäuresequenz von SEQ ID NO 3 oder Polypeptide umfasst, die mindestens 60 %, z. B. 80 %, 85 %, 90 %, 95 %, 98 % oder 99 % Sequenzidentität hierzu aufweisen.

5. Zusammensetzung nach einem der vorstehenden Ansprüche, wobei die Zusammensetzung ferner ein oder mehrere Enzyme umfasst, die aus der Gruppe ausgewählt sind, bestehend aus Proteasen, Lipasen, Cutinasen, Amylasen, Carbohydrasen, Cellulasen, Pektinasen, Mannanasen, Arabinasen, Galaktanasen, Xylanasen und Oxidasen.

6. Verwendung einer Zusammensetzung nach einem der Ansprüche 1 bis 5, vorzugsweise eine Waschmittelzusammensetzung umfassend eine *Staphylococcus*-Hexosaminidase,

a) zum Verhindern, Reduzieren oder Entfernen von Klebrigkeit des Objekts;

b) zum Vorbehandeln von Flecken auf dem Objekt;

c) zum Verhindern, Reduzieren oder Entfernen von Wiederablagerung von Verunreinigungen während eines Waschzyklus;

d) zum Verhindern, Reduzieren oder Entfernen einer Haftung von Verunreinigungen an dem Objekt;

e) zum Erhalten oder Verbessern einer Weiße des Objekts;

f) zum Verhindern, Reduzieren oder Entfernen von unangenehmem Geruch aus dem Gegenstand, wobei das Objekt ein Textil ist.

Revendications

1. Composition comprenant une hexosaminidase de *Staphylococcus*, dans laquelle l'hexosaminidase de *Staphylococcus* comprend le motif D[IV]AR[TK] (SEQ ID NO 12) et dans laquelle la composition

(a) est une composition solide, de préférence granulaire, de détergent à lessive et comprend en outre

(a1) au moins un adjuvant zéolite, de préférence en une quantité de 10 à 50 % en poids, plus préférentiellement 20 à 30 % en poids ;

(a2) au moins un adjuvant phosphonate, de préférence en une quantité de 0,1 à 5 % en poids, plus préférentiellement 0,4 à 1,5 % en poids ;

(a3) au moins une enzyme supplémentaire, de préférence une cellulase, de préférence en une quantité d'enzyme active de 100 à 5000 ppb, plus préférentiellement 1000 à 2000 ppb ; et

(a4) au moins un polymère, de préférence un polymère de polyvinylpyrrolidone, de préférence en une quantité de 0,01 à 1 % en poids, plus préférentiellement 0,1 à 0,3 % en poids ; ou

(b) est une composition solide de détergent à lessive et comprend en outre

(b1) au moins un adjuvant silicate, de préférence en une quantité de 2 à 20 % en poids, plus préférentiellement 5 à 10 % en poids ;

(b2) facultativement de la carboxyméthylcellulose, de préférence en une quantité de 0,1 à 10 % en poids, plus préférentiellement 0,1 à 4 % en poids ;

(b3) au moins une enzyme supplémentaire, de préférence une cellulase, de préférence en une quantité d'enzyme active de 0,1 à 100 ppm, plus préférentiellement 0,1 à 10 ppm ;

(b4) facultativement au moins un polymère antialissure, de préférence un polymère de polyvinylpyrrolidone, en une quantité de 0,1 à 3 % en poids, plus préférentiellement 0,1 à 1,0 % en poids ; et

(b5) au moins un système de blanchiment, comprenant un agent de blanchiment, un activateur de blanchiment et un catalyseur de blanchiment, de préférence en une quantité de 0,1 à 50 % en poids, plus préférentiellement 0,1 à 30 % en poids ; ou

(c) est une composition liquide de détergent à lessive et comprend en outre

(c1) au moins un agent tensioactif, de préférence un agent tensioactif non ionique, de préférence en une quantité de 1 à 20 % en poids, de préférence 3 à 15 % en poids ;

(c2) facultativement au moins un adjuvant phosphonate, de préférence en une quantité de 0,1 à 3 % en poids, plus préférentiellement 0,25 à 1,5 % en poids

(c3) facultativement au moins une enzyme supplémentaire, de préférence une cellulase, de préférence en une quantité de composition enzymatique de 0,001 à 1 % en poids, plus préférentiellement 0,001 à 0,6 % en poids ; et

(c4) facultativement au moins un solvant organique, de préférence du glycérol, de préférence en une quantité de 0,1 à 10 % en poids, plus préférentiellement 0,1 à 5 % en poids ; ou

EP 3 647 398 B1

(d) est un détergent liquide à lessive sous forme de dose unitaire, de préférence un sachet comprenant un film hydrosoluble, et comprend en outre

- 5 (d1) de l'eau en une quantité allant jusqu'à 20 % en poids, de préférence 5 à 15 % en poids ;
(d2) facultativement au moins un agent amérissant, de préférence le benzoate de benzyldiéthyl(2,6-xylyl-carbamoyl)-méthylammonium, de préférence en une quantité de 0,00001 à 0,04 % en poids ;
(d3) facultativement au moins un azurant optique, de préférence en une quantité de 0,01 à 2 % en poids, plus préférablement 0,01 à 1 % en poids ; et
10 (d4) facultativement au moins un polymère, de préférence en une quantité de 0,01 à 7 % en poids, plus préférablement 0,1 à 5 % en poids ; ou

(e) est un finisseur de tissu et comprend en outre

- 15 (e1) au moins une silicone adoucissante, de préférence une silicone à fonctionnalisation amino, de préférence en une quantité de 0,1 à 10 % en poids, plus préférablement 0,1 à 2 % en poids ;
(e2) au moins un parfum, de préférence au moins partiellement encapsulé dans des microcapsules, plus préférablement au moins partiellement encapsulé dans des microcapsules de mélamine-formaldéhyde, de préférence en une quantité de 0,01 à 3 % en poids, plus préférablement 0,1 à 1 % en poids ;
20 (e3) facultativement du polyquaternium 10 en une quantité de 0,1 à 20 % en poids, de préférence 0,1 à 13 % en poids ;
(e4) facultativement du polyquaternium 37 en une quantité de 0,1 à 20 % en poids, de préférence 0,1 à 13 % en poids ;
(e5) facultativement un esterquat d'origine végétale, de préférence un esterquat à base de canola ou de palme, en une quantité de 0,1 à 20 % en poids, de préférence 0,1 à 13 % en poids ; et
25 (e6) facultativement de l'acide adipique, en une quantité de 0,1 à 20 % en poids, de préférence 0,1 à 13 % en poids ; ou

(f) est un agent de nettoyage acide, ayant de préférence un pH inférieur à 6, et comprend en outre

- 30 (f1) des agents tensioactifs d'origine végétale ou biologique, de préférence chacun en une quantité de 0,1 à 5, plus préférablement chacun en une quantité de 0,1 à 2 % en poids ;
(f2) au moins un biocide acide, choisi de préférence parmi des acides, plus préférablement HCl et acide formique ; et
35 (f3) au moins un polymère antialissure, hydrofuge ou d'étalement d'eau, de préférence en une quantité de 0,01 à 3 % en poids, plus préférablement 0,01 à 0,5 % en poids ; ou

(g) est un agent de nettoyage neutre, ayant de préférence un pH compris entre 6,0 et 7,5, et comprend en outre (g1) des agents tensioactifs d'origine végétale ou biologique, de préférence chacun en une quantité de 0,1 à 5, plus préférablement chacun en une quantité de 0,1 à 2 % en poids ;

- 40 (g2) au moins un biocide, choisi de préférence parmi des composés d'ammonium quaternaire et des alcools ;
et
(g3) au moins un polymère antialissure, hydrofuge ou d'étalement d'eau, de préférence en une quantité de 0,01 à 3 % en poids, plus préférablement 0,01 à 0,5 % en poids ; ou
45

(h) est un agent de nettoyage alcalin, ayant de préférence un pH supérieur à 7,5, et comprend en outre (h1) des agents tensioactifs d'origine végétale ou biologique, de préférence chacun en une quantité de 0,1 à 5, plus préférablement chacun en une quantité de 0,1 à 2 % en poids ; ou

- 50 (i) est un agent de lavage de vaisselle à la main, de préférence un agent liquide de lavage de vaisselle à la main, et comprend en outre

- (i1) au moins un agent tensioactif anionique, de préférence en une quantité de 0,1 à 40 % en poids, plus préférablement 5 à 30 % en poids ;
55 (i2) au moins un agent tensioactif amphotère, de préférence la bétaine, de préférence en une quantité de 0,1 à 25 % en poids, plus préférablement 1 à 15 % en poids ;
(i3) au moins un agent tensioactif non ionique, de préférence en une quantité de 0,1 à 25 % en poids, plus préférablement 2 à 10 % en poids ;
(i4) au moins une enzyme supplémentaire, choisie de préférence parmi protéases, amylases et combinai-

sons de celles-ci, de préférence en une quantité de composition enzymatique allant jusqu'à 1 % en poids, plus préférablement jusqu'à 0,6 % en poids ; ou

(j) est une composition de lavage automatique de la vaisselle et comprend en outre

- (j1) au moins un adjuvant choisi parmi citrate, aminocarboxylates et combinaisons de ceux-ci, de préférence en une quantité de 5 à 30 % en poids, plus préférablement 10 à 20 % en poids ;
- (j2) au moins un adjuvant phosphonate, de préférence en une quantité de 0,1 à 5 % en poids, plus préférablement 0,4 à 1,5 % en poids ;
- (j3) au moins un agent tensioactif non ionique, de préférence en une quantité de 0,1 à 10 % en poids, plus préférablement 1 à 5 % en poids ;
- (j4) au moins un système de blanchiment, comprenant un agent de blanchiment, un activateur de blanchiment et un catalyseur de blanchiment, de préférence en une quantité de 0,1 à 50 % en poids, plus préférablement 0,1 à 30 % en poids ; et
- (j5) au moins un polymère choisi parmi sulfopolymères, polymères cationiques et polyacrylates, de préférence en une quantité de 0,01 à 15 % en poids, plus préférablement 2 à 10 % en poids.

2. Composition selon la revendication 1, dans laquelle la composition comprend en outre ;

(a)

- i. un ou plusieurs polyols, choisis de préférence parmi glycérol, (mono, di, ou tri) propylène glycol, éthylène glycol, polyéthylène glycol, alcools de sucre, sorbitol, mannitol, érythritol, dulcitol, inositol, xylitol et adonitol,
 - ii. facultativement une ou plusieurs enzymes, choisies de préférence parmi protéases, amylases ou lipases,
 - iii. facultativement un ou plusieurs agents tensioactifs, choisis de préférence parmi agents tensioactifs anioniques et non ioniques,
 - iv. facultativement un ou plusieurs polymères ;
- ou

(b) un granulé comprenant

- i. un noyau comprenant une hexosaminidase de *Staphylococcus* et facultativement,
- ii. un revêtement constitué d'une ou plusieurs couches entourant le noyau.

3. Composition selon la revendication 1 ou la revendication 2, dans laquelle l'hexosaminidase a une activité N-acétylglucosaminidase, de préférence une activité β -1,6 N-acétyl-glucosaminidase.

4. Composition selon l'une quelconque des revendications précédentes, dans laquelle le polypeptide ayant une activité hexosaminidase comprend la séquence d'acides aminés de SEQ ID NO 3 ou des polypeptides ayant au moins 60 %, par exemple 80 %, 85 %, 90 %, 95 %, 98 % ou 99 % d'identité de séquence avec celle-ci.

5. Composition selon l'une quelconque des revendications précédentes, dans laquelle la composition comprend en outre une ou plusieurs enzymes choisies dans le groupe constitué de protéases, lipases, cutinases, amylases, carbohydrases, cellulases, pectinases, mannanases, arabinases, galactanases, xylanases et oxydases

6. Utilisation d'une composition selon l'une quelconque des revendications 1 à 5, de préférence une composition détergente comprenant une hexosaminidase de *Staphylococcus*,

- a) pour la prévention, la réduction ou l'élimination du caractère collant de l'article ;
- b) pour le prétraitement de taches sur l'article ;
- c) pour la prévention, la réduction ou l'élimination de redéposition de salissures pendant un cycle de lavage ;
- d) pour la prévention, la réduction ou l'élimination d'adhérence de salissures sur l'article ;
- e) pour le maintien ou l'amélioration de la blancheur de l'article ;
- f) pour la prévention, la réduction ou l'élimination des mauvaises odeurs de l'article, l'article étant un textile.



Phylogenetic tree




Figure 2

Multiple alignment

```

      10      20      30      40      50
SEQ ID NO 3 Staphylococcus cohnii subsp. cohnii QDFQKGINVDIARKDYSLKSLKKIVDTIHENNGQYLQLHFSDNENYAIESQFFKH
SEQ ID NO 6 Staphylococcus fleurettii      ESIQECVSVDIARKEYSLESLSKQIVDTIHENNGQYLQLHFSDDENYAIESDYFSH

      60      70      80      90     100     110
SEQ ID NO 3 Staphylococcus cohnii subsp. cohnii ENTASQNYLSQQELKNLIHYSNKLNIIMVVEFDLPSSKAWLLLLKNNSNTHEN
SEQ ID NO 6 Staphylococcus fleurettii      QGITPNENYLTKEIKSLIAYSNELNMVVPDIDFPSSKALLSLIKNEDKDLYNQ

      120     130     140     150     160
SEQ ID NO 3 Staphylococcus cohnii subsp. cohnii IVSDYSDETIDFFSNQKALEISKRCIKELNLFHQPNFQKEQRIVLGGDEVPCGK
SEQ ID NO 6 Staphylococcus fleurettii      IISDYSDNITDFFSNQKALALSKRHIGETITLFNQPKYNGQQRIVLGGDEVPCGG

      170     180     190     200     210     220
SEQ ID NO 3 Staphylococcus cohnii subsp. cohnii SYGNDFINFMNEIGCYAYQNCYEPQIWNDSITKNGLKLKKNYFSVIPWKQSNNE
SEQ ID NO 6 Staphylococcus fleurettii      AYQSDFISYMMNIGSYAAGQCYEPQMWNDMISHEGIKSLNDTFSILYWKQ..NEN

      230     240     250     260     270
SEQ ID NO 3 Staphylococcus cohnii subsp. cohnii NEPGITVEDFLDYNFKNVYNYNFYSLYFLPSKNYSPTDIEEQTSYISWAYNHNSEFY
SEQ ID NO 6 Staphylococcus fleurettii      SKSDLTVEDFAEYDFKLIYNYNFYSLYFLPSNQFTNADIEEQADYISWAYAYNKKFF

      280     290     300     310     320
SEQ ID NO 3 Staphylococcus cohnii subsp. cohnii YLKNPYEVDLSNIQGSALSFWGEHATGMREEEVLNQELPLIRTYLNK
SEQ ID NO 6 Staphylococcus fleurettii      YTNPEPYQEVDSNVKGSALSFWGEDALNMSQTELINQEPLIKAYFSS

```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 04061117 A2, Kane Biotech INC [0003]
- WO 9850512 A [0003]
- WO 2017186943 A [0004]
- WO 2017186936 A [0005]
- WO 2017186937 A [0006]
- EP 0425019 A [0009]
- US 4106991 A [0047]
- US 4661452 A [0047]
- GB 1483591 A [0047]
- WO 2013188331 A [0048]
- WO 9307263 A [0049]
- WO 9723606 A [0049]
- WO 0001793 A [0053]
- WO 2006034710 A [0053]
- WO 9932595 A [0055]
- EP 0164514 A [0094]
- WO 09102854 A [0096] [0104]
- US 5977053 A [0096] [0104]
- WO 9817767 A [0106]
- EP 624154 A [0106]
- WO 2007087258 A [0110]
- WO 2007087244 A [0110]
- WO 2007087259 A [0110]
- EP 1867708 A [0110]
- WO 2007087242 A [0110]
- WO 9426860 A [0112]
- WO 9426859 A [0112]
- WO 2006130575 A [0115]
- WO 200503274 A [0116]
- WO 200503275 A [0116]
- WO 200503276 A [0116]
- EP 1876226 A [0116]
- WO 2007087257 A [0116]
- WO 2007087243 A [0116]
- US 7262042 B [0120]
- WO 09021867 A [0120]
- WO 9318140 A [0120]
- WO 01016285 A [0120]
- WO 02016547 A [0120]
- WO 9425583 A [0120]
- WO 05040372 A [0120]
- WO 05052161 A [0120]
- WO 05052146 A [0120]
- WO 9523221 A [0121]
- WO 9221760 A [0121]
- EP 1921147 A [0121]
- EP 1921148 A [0121]
- WO 07044993 A [0122]
- WO 8906279 A [0123]
- WO 9219729 A [0123]
- WO 96034946 A [0123]
- WO 9820115 A [0123]
- WO 9820116 A [0123]
- WO 99011768 A [0123]
- WO 0144452 A [0123]
- WO 03006602 A [0123]
- WO 0403186 A [0123]
- WO 04041979 A [0123]
- WO 07006305 A [0123]
- WO 11036263 A [0123]
- WO 11036264 A [0123]
- WO 2016001449 A [0123]
- WO 2004067737 A [0124]
- US 5352604 A [0125]
- US 4963655 A [0132]
- US 5159060 A [0132]
- WO 9512655 A [0132]
- WO 9529223 A [0132]
- WO 9219707 A [0132]
- WO 9404653 A [0132]
- WO 9404654 A [0132]
- US 5442100 A [0132]
- US 5488157 A [0132]
- US 5472628 A [0132]
- WO 9404651 A [0143]
- WO 9525791 A [0143]
- WO 9813458 A [0143]
- WO 9813459 A [0143]
- WO 9813460 A [0143]
- WO 9813461 A [0143]
- WO 9813462 A [0143]
- WO 07141736 A [0143]
- WO 07145963 A [0143]
- WO 09118375 A [0143]
- WO 10055052 A [0143]
- WO 11036153 A [0143]
- WO 2013004636 A [0145]
- WO 9847523 A [0146]
- US 6500802 B [0146]
- US 5436229 A [0146]
- WO 2013004635 A [0151]
- WO 2008134343 A [0152]
- US 4435307 A [0153]
- US 5648263 A [0153]
- US 5691178 A [0153]
- US 5776757 A [0153]
- WO 8909259 A [0153]
- EP 0495257 A [0154]

EP 3 647 398 B1

- EP 0531372 A [0154]
- WO 9611262 A [0154]
- WO 9629397 A [0154]
- WO 9808940 A [0154]
- WO 9407998 A [0154]
- EP 0531315 A [0154]
- US 5457046 A [0154]
- US 5686593 A [0154]
- US 5763254 A [0154]
- WO 9524471 A [0154]
- WO 9812307 A [0154]
- WO 99001544 A [0154]
- WO 2002099091 A [0155]
- WO 2001062903 A [0155]
- WO 1999064619 A [0157]
- WO 9324618 A [0158] [0177]
- WO 9510602 A [0158] [0177]
- WO 9815257 A [0158] [0177]
- EP 258068 A [0159]
- EP 305216 A [0159]
- WO 9613580 A [0159]
- EP 218272 A [0159]
- EP 331376 A [0159]
- WO 9506720 A [0159]
- WO 9627002 A [0159]
- WO 9612012 A [0159]
- WO 10065455 A [0159]
- WO 10107560 A [0159]
- US 5389536 A [0159]
- WO 11084412 A [0159]
- WO 11084417 A [0159]
- WO 11084599 A [0159]
- WO 11150157 A [0159]
- WO 12137147 A [0159]
- EP 407225 A [0160]
- WO 9205249 A [0160]
- WO 9401541 A [0160]
- WO 9425578 A [0160]
- WO 9514783 A [0160]
- WO 9530744 A [0160]
- WO 9535381 A [0160]
- WO 9522615 A [0160]
- WO 9600292 A [0160]
- WO 9704079 A [0160]
- WO 9707202 A [0160]
- WO 0034450 A [0160]
- WO 0060063 A [0160]
- WO 0192502 A [0160]
- WO 0787508 A [0160]
- WO 09109500 A [0160]
- WO 10111143 A [0162]
- WO 0556782 A [0162]
- WO 0967279 A [0162]
- WO 10100028 A [0162]
- GB 1296839 A [0163]
- WO 9510603 A [0164]
- WO 9402597 A [0164]
- WO 9418314 A [0164]
- WO 9743424 A [0164]
- WO 99019467 A [0164] [0167]
- WO 02010355 A [0165]
- WO 2006066594 A [0166]
- WO 96023873 A [0168]
- WO 08153815 A [0169]
- WO 0166712 A [0169] [0173]
- WO 09061380 A [0170]
- WO 13184577 A [0171]
- WO 10104675 A [0172]
- WO 2011098531 A [0174]
- WO 2013001078 A [0174]
- WO 2013001087 A [0174]
- EP 179486 A [0177]
- WO 9201046 A [0180]
- JP 2238885 A [0180]
- WO 9708325 A [0180]
- WO 9533836 A [0180]
- WO 2009087523 A [0184]
- WO 2007138054 A [0184]
- WO 2006108856 A [0184]
- WO 2006113314 A [0184]
- EP 1867808 A [0184]
- WO 2003040279 A [0184]
- EP 2169040 A [0186]
- US 20090011970 A1 [0190]
- WO 12025577 A [0233]
- WO 9943835 A [0233]

Non-patent literature cited in the description

- Purification of a 51 kDa endo-beta-N-acetylglucosaminidase from *Staphylococcus aureus*. **SUGAI et al.** FEMS MICROBIOLOGY LETTERS. WILEY-BLACKWELL PUBLISHING LTD, 15 October 1989, vol. 61, 267-272 [0007]
- **WADSTRÖM et al.** Bacteriolytic enzymes from *Staphylococcus aureus*. Purification of an endo-[beta]-N-acetylglucosaminidase. *BIOCHEMICAL JOURNAL*, 01 December 1970, vol. 120 (4), 725-734 [0008]
- **NEEDLEMAN ; WUNSCH.** *J. Mol. Biol.*, 1970, vol. 48, 443-453 [0035] [0224]
- **RICE et al.** EMBOSS: The European Molecular Biology Open Software Suite. *Trends Genet.*, 2000, vol. 16, 276-277 [0035] [0224]
- **CUNNINGHAM ; WELLS.** *Science*, 1989, vol. 244, 1081-1085 [0036]
- **HILTON et al.** *J. Biol. Chem.*, 1996, vol. 271, 4699-4708 [0036]
- **VOS et al.** *Science*, 1992, vol. 255, 306-312 [0036]

- **SMITH et al.** *J. Mol. Biol.*, 1992, vol. 224, 899-904 [0036]
- **WLODAVER et al.** *FEBS Lett.*, 1992, vol. 309, 59-64 [0036]
- Handbook of Powder Technology. **C. E. CAPES**. Particle size enlargement. Elsevier, 1980, vol. 1 [0049]
- **HODGDON ; KALER.** *Current Opinion in Colloid & Interface Science*, 2007, vol. 12, 121-128 [0113]
- *J. Am. Chem. Soc.*, 1978, vol. 100, 1228 [0146]
- *Org. Synth., Coll.*, vol. 7, 361 [0146]
- Powdered Detergents, Surfactant science series. Marcel Dekker, Inc, vol. 71 [0181] [0184]
- *CHEMICAL ABSTRACTS*, 9002-92-0 [0229]
- **DIDERICHSEN et al.** *Plasmid*, 1993, vol. 30, 312-315 [0233]
- **VALLE et al.** *Mol Microbiol.*, May 2003, vol. 48 (4), 1075-87 [0235]
- **VALLE, J. ; A. TOLEDO-ARANA ; C. BERASAIN ; J. M. GHIGO ; B. AMORENA ; J. R. PENADES ; I. LASA.** *Mol. Microbiol.*, 2003, vol. 48, 1075-1087 [0236]
- **FINN.** *Nucleic Acids Research*, 2016, vol. 44, D279-D285 [0238]
- **EDGAR.** *Nucleic Acids Research*, 2004, vol. 32 (5), 1792-1797 [0238]
- **PRICE et al.** *PloS one*, 2010, vol. 5 (3 [0238]
- **LETUNIC ; BORK.** *Bioinformatics*, 2007, vol. 23 (1), 127-128 [0238]
- *CHEMICAL ABSTRACTS*, 3459-18-5 [0244] [0251]
- *CHEMICAL ABSTRACTS*, 4432-31-9 [0244] [0248] [0251]
- *CHEMICAL ABSTRACTS*, 64-19-7 [0244] [0251]
- *CHEMICAL ABSTRACTS*, 56-40-6 [0251]