



(11) **EP 3 702 373 B9**

(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

- (15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Description Paragraph(s) 10, 14, 30, 40, 41, 50, 78, 80, 84-89, 95-97
Claims EN 10
- (48) Corrigendum issued on:
23.11.2022 Bulletin 2022/47
- (45) Date of publication and mention of the grant of the patent:
24.08.2022 Bulletin 2022/34
- (21) Application number: **20154386.5**
- (22) Date of filing: **13.09.2013**
- (51) International Patent Classification (IPC):
C07K 16/28^(2006.01) A61P 35/00^(2006.01)
A61K 39/395^(2006.01) A61P 25/00^(2006.01)
A61P 31/12^(2006.01)
- (52) Cooperative Patent Classification (CPC):
A61K 39/39591; A61P 25/00; A61P 31/12;
A61P 35/00; C07K 16/2818; A61K 2039/505;
C07K 2317/24; C07K 2317/33; C07K 2317/34;
C07K 2317/52; C07K 2317/55; C07K 2317/565;
C07K 2317/567; C07K 2317/70; C07K 2317/71;
- (Cont.)

(54) **ANTI-PD1 ANTIBODIES AND THEIR USE AS THERAPEUTICS AND DIAGNOSTICS**

ANTI-PD1-ANTIKÖRPER SOWIE VERWENDUNG DAVON IN THERAPEUTIKA UND DIAGNOSTIKA
ANTICORPS ANTI-PD1 ET LEUR UTILISATION EN TANT QU'AGENTS THÉRAPEUTIQUES ET DE DIAGNOSTIC

- (84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
Designated Extension States:
BA ME
- (43) Date of publication of application:
02.09.2020 Bulletin 2020/36
- (60) Divisional application:
22182191.1
- (62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
13893636.4 / 3 044 234
- (73) Proprietor: **BeiGene Switzerland GmbH**
4051 Basel (CH)
- (72) Inventors:
• **SONG, Jing**
Changping, Beijing 102206 (CN)
- **LI, Kang**
Changping, Beijing 102206 (CN)
- **ZHANG, Tong**
Changping, Beijing 102206 (CN)
- **XU, Lanlan**
Changping, Beijing 102206 (CN)
- **LIU, Qi**
Changping, Beijing 102206 (CN)
- **PENG, Hao**
Changping, Beijing 102206 (CN)
- (74) Representative: **Mewburn Ellis LLP**
Aurora Building
Counterslip
Bristol BS1 6BX (GB)
- (56) References cited:
WO-A1-2006/121168 WO-A1-2008/156712
WO-A1-2012/135408 WO-A1-2012/145493
WO-A2-2010/089411

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 3 702 373 B9

- M. J. FULLER ET AL: "Immunotherapy of chronic hepatitis C virus infection with antibodies against programmed cell death-1 (PD-1)", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 110, no. 37, 26 August 2013 (2013-08-26), pages 15001-15006, XP055343159, US ISSN: 0027-8424, DOI: 10.1073/pnas.1312772110
 - WU H ET AL: "Humanization of a murine monoclonal antibody by simultaneous optimization of framework and CDR residues", JOURNAL OF MOLECULAR BIOLOGY, ACADEMIC PRESS, UNITED KINGDOM, vol. 294, no. 1, 19 November 1999 (1999-11-19), pages 151-162, XP004461859, ISSN: 0022-2836, DOI: 10.1006/JMBI.1999.3141
 - JAMES W. STAVE ET AL.: "Antibody and Antigen Contact Residues Define Epitope and Paratope size and Structure", THE JOURNAL OF IMMUNOLOGY, vol. 191, 1 January 2013 (2013-01-01), pages 1428-1435, XP055303910,
 - M. SZNOL ET AL: "Antagonist Antibodies to PD-1 and B7-H1 (PD-L1) in the Treatment of Advanced Human Cancer", CLINICAL CANCER RESEARCH, vol. 19, no. 5, 1 March 2013 (2013-03-01), pages 1021-1034, XP055123957, ISSN: 1078-0432, DOI: 10.1158/1078-0432.CCR-12-2063
 - OMID HAMID ET AL: "Safety and Tumor Responses with Lambrolizumab (Anti-PD-1) in Melanoma", NEW ENGLAND JOURNAL OF MEDICINE, vol. 369, no. 2, 11 July 2013 (2013-07-11) , pages 134-144, XP055182016, ISSN: 0028-4793, DOI: 10.1056/NEJMoa1305133
 - BERGER RAANAN ET AL: "Phase I safety and pharmacokinetic study of CT-011, a humanized antibody interacting with PD-1, in patients with advanced hematologic malignancies", CLINICAL CANCER RESEARCH, THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, US, vol. 14, no. 10, 15 May 2008 (2008-05-15), pages 3044-3051, XP002524925, ISSN: 1078-0432, DOI: 10.1158/1078-0432.CCR-07-4079
 - DAVID M. DORFMAN ET AL: "Programmed Death-1 (PD-1) is a Marker of Germinal Center-associated T Cells and Angioimmunoblastic T-Cell Lymphoma", AMERICAN JOURNAL OF SURGICAL PATHOLOGY., vol. 30, no. 7, 1 July 2006 (2006-07-01), pages 802-810, XP055343182, US ISSN: 0147-5185, DOI: 10.1097/01.pas.0000209855.28282.ce
- (52) Cooperative Patent Classification (CPC): (Cont.)
C07K 2317/73; C07K 2317/76; C07K 2317/92;
C07K 2317/94

Description

INTRODUCTION

5 **[0001]** Programmed Death-1 (PD-1, also termed as CD279) is a 55 KD receptor protein related to CD28/CTLA4 co-stimulatory/inhibitory receptor family (Blank et al., 2005 *Cancer Immunol Immunother* 54:307-314). The genes and cDNAs coding for PD-1 were cloned and characterized in mouse and human (Ishida et al., 1992 *EMBO J* 11:3887-3395; Shinohara et al., 1994 *Genomics* 23:704-706). The full length PD-1 contains 288 amino acid residues (NCBI accession number: NP_005009). Its extracellular domain consists of amino acid residues 1-167, and the cytoplasmic C-terminal

10 tail comprises residues 191-288, which has two hypothetical immune-regulatory motifs, an immunoreceptor tyrosine-based inhibitory motif (ITIM; Vivier et al., 1997 *Immunol Today* 18:286-291) and an immunoreceptor tyrosine switch motif (ITSM; Chemnitz et al., 2004 *J Immunol* 173:945-954).

15 **[0002]** To date, two sequence-related ligands, PD-L1 (B7-H1) and PD-L2 (B7-DC), have been identified to specifically interact with PD-1, inducing intracellular signal transduction that inhibits CD3 and CD28 mediated T-cell activation (Riley, 2009 *Immunol Rev* 229:114-125), which, in turn, attenuates T-cell activities, for example, reduction of cell proliferation, IL-2 and IFN- γ secretion, as well as other growth factor and cytokine secretion.

20 **[0003]** Expression of PD-1 was frequently found in immune cells such as T-cells, B-cells, monocytes and natural killer (NK) cells. It was rarely expressed in other human tissues, such as muscle, epithelium, neuronal tissues, etc. Furthermore, high level of PD-1 expression is often associated with activation of immune cells. For example, when human T-cell line, Jurkat, was activated by phytohaemagglutinin (PHA) or phorbol ester (12-O-tetradecanoylphorbol-13-acetate, or TPA), the expression of PD-1 was up-regulated visible in Western Blot (Vibharka et al., 1997 *Exp Cell Res* 232:25-28). The same phenomenon was observed in stimulated murine T- and B-lymphocytes and in primary human CD4⁺T-cells upon stimulation by anti-CD3 antibody (Agata et al., 1996 *Int Immunol* 8:765-772; Bennett et al., 2003 *J Immunol* 170:711-118). The increase of PD-1 expression following stimulation of T effector cells redirects the activated T-effector cells towards

25 exhaustion and reduced immune activities. Therefore, PD-1 mediated inhibitory signal plays an important role in immune tolerance (Bour-Jordan et al., 2011 *Immunol Rev* 241:180-205).

30 **[0004]** Increase of PD-1 expression in tumor-infiltrating lymphocytes (TILs) and PD-1 ligand expression in tumor cells were reported in varieties of cancers involved in different types of tissues and organs such as lung (Konishi et al., 2004 *Clin Cancer Res* 10:5094-5100), liver (Shi et al., 2008 *Int J Cancer* 128:887-896; Gao et al., 2009 *Clin Cancer Res* 15:971-979), stomach (Wu et al., 2006 *Acta Histochem* 108:19-24), kidney (Thompson et al., 2004 *Proc Natl Acad Sci* 101:17174-17179; Thompson et al., 2007 *Clin Cancer Res* 13:1757-1761), breast (Ghebeh et al., 2006 *Neoplasia* 8:190-198), ovary (Hamanishi et al. 2007 *Proc Natl Acad Sci* 104:3360-3365), pancreas (Nomi et al., 2007 *Clin Cancer Res* 13:2151-2157), melanocytes (Hino et al., 2010 *Cancer* 116:1757-1766) and esophagus (Ohigashi et al., 2005 *Clin Cancer Res* 11:2947-2953). More frequently, the increased expression of PD-1 and PD-L1 in those cancers is associated

35 with poor prognosis of patient survival outcome. Transgenic mice with PD-1 gene knockout inhibiting xenograft cancer cell growth further elucidated the significance of PD-1 signaling in the modulation of immune system for cancer eradication or tolerance (Zhang et al., 2009 *Blood* 114:1545-1552).

40 **[0005]** Not only does up-regulation of PD-1 signaling leads to immune tolerance to cancerous growth, but also to viral infection and expansion in human. The prevalent liver infection viruses, HBV and HCV, induce overexpression of PD-1 ligands in hepatocytes and activate PD-1 signaling in T-effector cells, resulting in T-cell exhaustion and tolerance to the viral infection (Boni et al., 2007 *J Virol* 81:4215-4225; Golden-Mason et al., 2008 *J Immunol* 180:3637-3641). Likewise, HIV infection frequently evades human immune system by similar mechanisms. Therapeutic modulation of PD-1 signaling by antagonist molecules may revert immune cells from tolerance, and reactivated to eradicate cancer and chronic viral infection (Blank et al., 2005 *Cancer Immunol Immunother* 54:307-314; Okazaki et al., 2007 *Int Immunol* 19:813-824).

45 WO2006/121168 and WO2008/156712 disclose high affinity anti-human PD-1 antibodies, relating to Nivolumab and Pembrolizumab, respectively. The invention is defined by the claims and any other aspects or embodiments set forth herein not falling within the scope of the claims are for information only.

SUMMARY OF THE INVENTION

50 **[0006]** The invention provides methods and compositions for immune-inhibition of PD-1. The invention provides an antibody as defined in the claims. The invention also provides an antibody for use in therapy, as defined in the claims. The references in this description to methods of treatment by therapy comprising administration of the antibody of the invention are to be interpreted as references to the antibody of the invention for use in those methods.

55 **[0007]** In one aspect, the invention provides an antibody comprising an antigen binding domain which binds human PD-1, and comprises a heavy chain variable region (Vh) comprising SEQ ID NO: 24, and a light chain variable region (Vk) comprising SEQ ID NO: 26, wherein the antibody is monospecific.

[0008] The antibody of the invention comprises a heavy chain variable region (Vh) and a light chain variable region

(Vk) comprising:

317-4B6 CDR-H1, CDR-H2 and CDR-H3 (SEQ ID NOS:31-33);
CDR-L1, CDR-L2 and CDR-L3 (SEQ ID NOS:34-36).

5

[0009] According to the invention, the domain comprises a heavy chain variable region (Vh) and a light chain variable region (Vk) comprising 317-4B6 (SEQ ID NOS:24 and 26).

[0010] In particular embodiments, the domain specifically binds PD1 residues: (a) K45 and I93 (AA numbering based on 2008 PNAS, 105:10483; equivalent to K58 and I106 in SEQ ID NO 2).

10 **[0011]** In particular embodiments, the domain induces IL-2 release in HuT78/PD-1 cells co-cultured with HEK293/OS8/PD-L1 cells or with HEK293/OS8/PD-L2 cells, and/or inhibits IL-2 secretion in HuT78/P3Z cells co-cultured with HEK293/PD-L1 cells or with HEK293/PD-L2 cells.

[0012] The invention also provides antibodies comprising a subject PD-1 binding domain as defined in the claims and a IgG4 heavy chain effector or constant domain comprising any of SEQ ID NO:83-88, particularly SEQ ID NO 87 or 88.

15 **[0013]** The invention provides an antibody comprising an antibody antigen binding domain which binds human PD-1 and comprises a heavy chain variable region (Vh) comprising SEQ ID NO: 24, and a light chain variable region (Vk) comprising SEQ ID NO: 26, for use in therapy, which encompasses methods of using the subject domains by administering the domain to a person determined to have cancer or a viral infection or to otherwise be in need of PD-1 antagonism. In one embodiment, the use is for treating cancer or viral infection in a human.

20

BRIEF DESCRIPTION OF THE DRAWINGS

[0014]

25 Fig. 1. Schematic presentation of PD-1/Fc (top) and PD-1/His (bottom). ECD: extracellular domain. L: linker. H: His tag. Fc: γ 4Fc fragment from human IgG4. N: N-terminus. C: C-terminus.

Fig. 2. Dose-dependent reaction curves of murine mAbs binding to human PD-1 in ELISA. The murine mAbs were indicated at top-left corner of each figure. MAb 317 and 517 share high degree of homology the variable region of heavy and light chains. The binding signal strength was indicated by direct OD₄₅₀ readings. The antigen, PD-1/His, was coated at increasing concentrations up to 70 nanograms per well in a volume of 50 microliters. The method was described in Example 1.

30

Fig. 3. Dose-dependent reaction curve of murine mAbs binding to human PD-1 expressed on live cells by FACS analyses. Murine antibody codes and EC₅₀ were indicated on each panel. MFI stands for mean fluorescence intensity. HuT78/PD-1 cells were suspended in 96-well plate at 5×10^4 cells per well for FACS. PD-1 mAbs binding to the cell surface target and FACS detection were performed as described in Example 1.

35

Fig. 4. Schematic presentation of the cell co-culture systems used for assaying functional activities of anti-PD-1 mAbs. T-cells (either CD4⁺ or CD8⁺) represent HuT78/PD-1 or primary T-cells in PBMCs. TCR: T-cell receptor. N: nucleus. C: cytoplasm

Fig. 5. Dose-dependent reaction curve of murine mAb-induced IL-2 secretion in HuT78/PD-1 cells co-cultured with HEK293/OS8/PD-L1 cells. Baseline: Average IL-2 release induced by mIgGs at all tested concentrations. Top line: Highest IL-2 release based on regression calculation by Prizm Software.

40

Fig. 6. (A) Histograms showing IFN- γ secretion induced by anti-PD-1 mAbs in PBMCs (Donor-19) co-cultured with cell line HEK293/OS8/PD-L1. (B) Histograms showing IFN- γ secretion induced by anti-PD-1 mAbs in PBMCs (Donor-20) co-cultured with cell line HEK293/OS8/PD-L1.

Fig. 7. (A) and (B) ADCC activities of anti-PD-1 mAbs by co-culture of effector cells (NK92MI/PD-1) and target cells (HuT78/PD-1). Means were calculated from two data points of the representative experiments. The mAbs were added to concentration of 10 μ g/ml. Experiment performed as described in Example 9.

45

Fig. 8. Mapping the binding epitopes of anti-PD-1 mAbs by ELISA (up-panel) and Western Blot (lower panel). Conditioned media containing WT or Mt PD-1 were used to assess binding activity by ELISA and Western Blot. ** indicates the AA residues to which the mAb binding activity reduced to 25-50% of WT PD-1. *** indicates the AA residues to which the mAb binding activity reduced below 25% of WT PD-1.

50

Fig. 9. IFN- γ release induced by humanized anti-PD-1 mAbs in primary human PBMCs from different healthy donors, which were co-cultured with HEK293/OS8/PD-L1 cells.

Fig. 10. Cytotoxicity of NK92MI/PD-1 cells enhanced by humanized anti-PD-1 mAbs, hu317 (A) and hu326 (B). The target lung cancer cells, SK-MES-1/PD-L1, were co-cultured with the effector cells at the (T to E) ratio of 1 to 2, and assayed as described in Example 12.

55

Fig. 11. Individual tumor growth curves in three treatment groups, vehicle (PBS), human IgGs (huIgGs) and anti-PD-1 mAb (hu317-1/IgG4mt2). Each curve represents a tumor growth path, the tumor-bearing mice coded by

numbers indicated on the right of each panel. Hep3B/OS8/PD-L1 cells (established from hepatocellular carcinoma line Hep3B) were seeded at Day 1, PBMCs were implanted at Day 15 and three doses of hu317-1/IgG4mt2 were injected at Day 18, 28 and 38, respectively. Methods described in Example 12.

5 DESCRIPTION OF PARTICULAR EMBODIMENTS OF THE INVENTION

[0015] PD-1 initiates inhibitory signaling in immune cells when engaged by its ligands, PD-L1 or PD-L2. In the cases of cancer outgrowth and viral infection, the activation of PD-1 signaling promotes immune tolerance, leading to the cancers or virus-infected cells escaping from immune surveillance and cancer metastasis or viral load increase. Inhibition of PD-1 mediated cellular signaling by therapeutic agents can activate immune cells including T-cells, B-cells and NK cells, and therefore enhance immune cell functions inhibiting cancer cell growth or viral infection, and restore immune surveillance and immune memory function to treat such human diseases.

[0016] The invention provides antibodies whose functions are antagonistic to the ligand-induced and PD-1-mediated cellular signaling in immune cells. Murine anti-PD-1 antibodies were humanized to a high degree of similarity to human antibodies in the framework regions. The full antibodies made in the modified human IgG4 variant format have a unique set of features in the aspects of effector functions and physicochemical properties. The disclosed anti-PD-1 antibodies are suitable for therapeutic uses in cancer treatment, controlling viral infections and other human diseases that are mechanistically involved in exacerbated immune tolerance.

20 Definitions

[0017] Unless the context indicates otherwise, the term "antibody" is used in the broadest sense and specifically covers antibodies (including full length monoclonal antibodies) and antibody fragments so long as they recognize PD-1. An antibody molecule is usually monospecific, but may also be described as idiospecific, heterospecific, or polyspecific. Antibody molecules bind by means of specific binding sites to specific antigenic determinants or epitopes on antigens. "Antibody fragments" comprise a portion of a full length antibody, generally the antigen binding or variable region thereof. Examples of antibody fragments include Fab, Fab', F(ab').sub.2, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

[0018] Monoclonal antibodies (MAbs) may be obtained by methods known to those skilled in the art. See, for example Kohler et al (1975); U.S. Pat. No. 4,376,110; Ausubel et al (1987-1999); Harlow et al (1988); and Colligan et al (1993). The mAbs of the invention may be of any immunoglobulin class including IgG, IgM, IgE, IgA, and any subclass thereof. A hybridoma producing a mAb may be cultivated in vitro or in vivo. High titers of mAbs can be obtained in in vivo production where cells from the individual hybridomas are injected intraperitoneally into mice, such as pristinely-primed Balb/c mice to produce ascites fluid containing high concentrations of the desired mAbs. MAbs of isotype IgM or IgG may be purified from such ascites fluids, or from culture supernatants, using column chromatography methods well known to those of skill in the art.

[0019] An "isolated polynucleotide" refers to a polynucleotide segment or fragment which has been separated from sequences which flank it in a naturally occurring state, e.g., a DNA fragment which has been removed from the sequences which are normally adjacent to the fragment, e.g., the sequences adjacent to the fragment in a genome in which it naturally occurs. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule (e.g., as a cDNA or a genomic or cDNA fragment produced by PCR or restriction enzyme digestion) independent of other sequences. It also includes a recombinant DNA, which is part of a hybrid gene encoding additional polypeptide sequence.

[0020] A "construct" means any recombinant polynucleotide molecule such as a plasmid, cosmid, virus, autonomously replicating polynucleotide molecule, phage, or linear or circular single-stranded or double-stranded DNA or RNA polynucleotide molecule, derived from any source, capable of genomic integration or autonomous replication, comprising a polynucleotide molecule where one or more polynucleotide molecule has been linked in a functionally operative manner, i.e. operably linked. A recombinant construct will typically comprise the polynucleotides of the invention operably linked to transcriptional initiation regulatory sequences that will direct the transcription of the polynucleotide in the intended host cell. Both heterologous and non-heterologous (i.e., endogenous) promoters can be employed to direct expression of the nucleic acids of the invention.

[0021] A "vector" refers any recombinant polynucleotide construct that may be used for the purpose of transformation, i.e. the introduction of heterologous DNA into a host cell. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome

of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors".

[0022] An "expression vector" as used herein refers to a nucleic acid molecule capable of replication and expressing a gene of interest when transformed, transfected or transduced into a host cell. The expression vectors comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desired, provide amplification within the host. The expression vector further comprises a promoter to drive the expression of the polypeptide within the cells. Suitable expression vectors may be plasmids derived, for example, from pBR322 or various pUC plasmids, which are commercially available. Other expression vectors may be derived from bacteriophage, phagemid, or cosmid expression vectors.

Additional Embodiments

[0023] Mouse monoclonal antibodies were identified from screening murine hybridoma clones.

[0024] The following polynucleotide and protein sequences relate to these antibodies:

- a) The cDNA sequence, SEQ ID NO 3, encoding the heavy chain variable region of murine mAb 317;
- b) The protein sequence of the heavy chain variable region of murine mAb 317 or mu317_Vh (SEQ ID NO 4);
- c) The cDNA sequence, SEQ ID NO 5, encoding the light chain variable region of murine mAb 317;
- d) The protein sequence of the light chain variable region of murine mAb 317 or mu317_Vk (SEQ ID NO 6);
- e) The cDNA sequence, SEQ ID NO 7, encoding the heavy chain variable region of murine mAb 326;
- f) The protein sequence of the heavy chain variable region of murine mAb 326 or mu326_Vh (SEQ ID NO 8);
- g) The cDNA sequence, SEQ ID NO 9, encoding the light chain variable region of murine mAb 326;
- h) The protein sequence of the light chain variable region of murine mAb 326 or mu326_Vk (SEQ ID NO 10).

[0025] The complement determinant region (CDR) sequences of these antibodies, which mediate binding to the target antigens, PD-1, including the CDR sequences of mu317 and m326, are as follows:

- a) The CDR1 of mu317 heavy chain (mu317 H-CDR1) contains amino acid sequence of GFSLTSYGVH (SEQ ID NO 11);
- b) The mu317 H-CDR2 contains amino acid sequence of VIWAGGSTNYNSALMS (SEQ ID NO 12);
- c) The mu317 H-CDR3 contains amino acid sequence of ARAYGNYWYIDV (SEQ ID NO 13);
- d) The CDR1 of mu317 light chain (mu317 L-CDR1) contains amino acid sequence of KASQSVSNDVA (SEQ ID NO 14);
- e) The mu317 L-CDR2 contains amino acid sequence of YAFHRFT (SEQ ID NO 15);
- f) The mu317 L-CDR3 contains amino acid sequence of HQAYSSPYT (SEQ NO 16);
- g) The mu326 H-CDR1 contains amino acid sequence of GYTFTNYGMN (SEQ ID NO 17);
- h) The mu326 H-CDR2 contains amino acid sequence of WINNNNGEPTYAEEFKG (SEQ ID NO 18);
- i) The mu326 H-CDR3 contains amino acid sequence of ARDVMDY (SEQ ID NO 19);
- j) The mu326 L-CDR1 contains amino acid sequence of RASESVDNYGYFSFMH (SEQ ID NO 20);
- k) The mu326 L-CDR2 contains amino acid sequence of RASNLES (SEQ ID NO 21);
- l) The mu326 L-CDR3 contains amino acid sequence of QQSKEYPT (SEQ ID NO 22).

[0026] In one embodiment, the invention provides compositions comprising an antibody comprising the sequences of SEQ ID NO 24 and SEQ ID NO 26, which is a humanization monoclonal antibody emanated from murine mAb mu317:

- a) The humanization mAb hu317-4B6 comprises protein sequence of heavy chain variable region (Vh) as SEQ ID NO 24, which is encoded by
- b) the cDNA of hu317-4B6_Vh (SEQ ID NO 23);
- c) The humanization mAb hu317-4B6 also comprises protein sequence of light chain variable region (Vk) as SEQ ID NO 26, which is encoded by
- d) the cDNA of hu317-4B6 (SEQ ID NO 25).

[0027] Other humanization monoclonal antibodies emanated from murine mAbs mu326 and mu317 comprise the following sequences:

- e) the humanization mAb hu326-4A3 comprises protein sequence of Vh as SEQ ID NO 28, which is encoded by
- f) the cDNA of hu326-4A3-Vh (SEQ ID NO 27);

- g) The humanization mAb hu326-4A3 also comprises protein sequence of Vk as SEQ ID NO 30, which is encoded by h) the cDNA of hu326-4A3_Vk (SEQ ID NO 29);
 i) The protein sequences of hu317-4B2_Vh (SEQ ID NO 43) and hu317-4B2_Vk (SEQ ID NO 44);
 j) The protein sequences of hu317-4B5_Vh (SEQ ID NO 45) and hu317-4B5_Vk (SEQ ID NO 46);
 k) The protein sequence of hu317-1_Vh (SEQ ID NO 48) and the cDNA encoding for hu317-1_Vh (SEQ ID NO 47);
 l) The protein sequence of hu317-1_Vk (SEQ ID NO 50) and the cDNA encoding for hu317-1_Vk (SEQ ID NO 49);
 m) The protein sequences of hu326-3B1_Vh (SEQ ID NO 51) and hu326-3B1_Vk (SEQ ID NO 52);
 n) The protein sequences of hu326-3G1_Vh (SEQ ID NO 53) and hu326-3G1_Vk (SEQ ID NO 54);
 o) The protein sequence of hu326-1_Vh (SEQ ID NO 56) and the cDNA encoding for hu326-1_Vh (SEQ ID NO 55);
 p) The protein sequence of hu326-1_Vk (SEQ ID NO 58) and the cDNA encoding for hu326-1_Vk (SEQ ID NO 57);
 q) The protein sequences of other humanization mAbs emanated from mu317 (SEQ ID NO 63-74);
 r) The protein sequences of other humanization mAbs emanated from mu326 (SEQ ID NO 75-82).

[0028] The antibody of the invention comprises the CDR sequences listed below:

- a) H-CDR1 sequence of GFSLTSYGVH (SEQ ID NO 31), shared throughout humanization mAbs hu317 and mu317 in the heavy chains;
 b) H-CDR3 sequence of ARAYGNYWYIDV (SEQ ID NO 33), shared throughout humanization mAbs hu317 and mu317 in the heavy chains;
 c) L-CDR1 sequence of KSSSEVSNDVA (SEQ ID NO 34), shared throughout humanization mAbs hu317-4B2, hu317-4B5 and hu317-4B6 in the light chains;
 d) L-CDR2 sequence of YAFHRFT (SEQ ID NO 35), shared throughout humanization mAbs hu317 and mu317 in the light chains;
 e) L-CDR3 sequence of HQAYSSPYT (SEQ ID NO 36), shared throughout humanization mAbs hu317 and mu317 in the light chains;
 f) H-CDR2 sequence of VIYADGSTNYNPSLKS (SEQ ID NO 32) in hu317-4B6_Vh.

[0029] The CDR sequences of the other humanized mAbs are as follows:

- g) H-CDR2 sequence of VIYAGGSTNYNPSLKS (SEQ ID NO 60) in hu317-4B2_Vh and hu317-4B5_Vh;
 h) H-CDR2 sequence of VIWAGGSTNYNPSLKS (SEQ ID NO 59) in hu317-1_Vh;
 i) L-CDR1 sequence of KASQSVSNDVA (SEQ ID NO 14) in hu317-1_Vk;
 j) H-CDR1 sequence of GYTFTNYGMN (SEQ ID NO 37), shared throughout humanization mAbs hu326 and mu326 in the heavy chains;
 k) H-CDR3 sequence of ARDVMDY (SEQ ID NO 39), shared throughout humanization mAbs hu326 and mu326 in the heavy chains;
 l) L-CDR1 sequence of RASESVDNYGYSEFMH (SEQ ID NO 40), shared throughout humanization mAbs hu326 and mu326 in the light chains;
 m) L-CDR2 sequence of RASNLES (SEQ ID NO 41), shared throughout humanization mAbs hu326 and mu326 in the light chains;
 n) L-CDR3 sequence of QQSKEYPT (SEQ ID NO 42), shared throughout humanization mAbs hu326 and mu326 in the light chains;
 o) H-CDR2 sequence of WINNNNAEPTYAQDFRG (SEQ ID NO 38) in hu326_4A3_Vh;
 p) H-CDR2 sequence of WINNNNGEPTYAQDFRG (SEQ ID NO 62) in the Vh of hu326_1 and other hu317 mAbs.

[0030] Six critical amino acid (AA) residues in PD-1 required for the ligand binding were mutated individually, and mutant and wild-type PD-1 proteins were used to assess the binding epitopes. The residue whose mutation significantly impaired the antibody binding is recognized as a key or significant binding epitope. Significant binding epitopes of mAbs hu317-4B5 and hu317-4B6 are K45 and I93 (AA numbering based on 2008 PNAS, 105:10483; equivalent to K58 and I106 in SEQ ID NO 2); and significant binding epitopes of mAbs hu326-3B1 and hu317-4A3 are I93, L95 and P97 (AA numbering based on 2008 PNAS, 105:10483; equivalent to I106, L108 and P110 in SEQ ID NO 2).

[0031] In a further aspect, the invention provides compositions comprising antibodies as defined in the claims comprising the constant region sequences of recombinant human IgG4 variants, which may be linked to the variable regions of the subject antibodies, including the humanized anti-PD-1 mAbs, which showed preferred effector functions and physicochemical properties. The sequences are as follows:

[0032] The constant region sequence of IgG4mt10 (SEQ ID NO 88);

- a) A reference sequence of IgG4mt1 (SEQ ID NO 83);

- b) A reference sequence of IgG4mt2 (SEQ ID NO 84);
- c) A reference sequence of IgG4mt6 (SEQ ID NO 85);
- d) A reference sequence of IgG4mt8 (SEQ ID NO 86);
- e) A reference sequence of IgG4mt9 (SEQ ID NO 87).

5

[0033] Anti-PD-1 antibody functions can be assayed using a plasmid expressing the recombinant fusion protein, OS8, to generate stable cell lines, HEK293/OS8/PD-L1 or HEK293/OS8/PD-L2, which co-expresses OS8 (a T cell-activating molecule) and a PD-1 ligand. The cell lines were used to engage T-cells and PBMCs by co-culture to assess the functionality of anti-PD-1 mAbs (see Example 3 and Example 4). Alternatively, another plasmid expressing the recombinant fusion protein, P3Z, was used to generate stable cell line, HuT78/P3Z, in which P3Z functions as molecular sensor and signal transduction mediator. When P3Z is engaged by PD-1 ligand, it will transmit intracellular signal to activate IL-2 release in the HuT78 cells. The systems may be used to assess inhibitory effect of anti-PD-1 mAbs (see Example 3).

10

[0034] In another embodiment, the invention provides the humanized anti-PD-1 antibodies as defined in the claims, including hu317-4B6, having functions to suppress PD-1 mediated signal transduction, and to activate immune cells, which trigger a cascade of immune responses including cytokine secretion and cytotoxicity towards target cells such as cancer cells, and such functional use of the antibodies.

15

[0035] In one aspect, the invention provides humanized anti-PD-1 antibodies as defined in the claims that activate several types of immune cells that express PD-1, including human T-cells, NK-cells and PBMCs, whose functions are to amplify the immune response signals, to mobilize immune system and to act as immune effector cells for clearance of cancer cells and viral infections, and such functional use of the antibodies.

20

[0036] In another aspect, the humanized anti-PD-1 mAbs are used as therapeutic agents to treat human diseases that are involved in suppression of immune cells by PD-1 mediated intracellular signaling, leading to disease progression, particularly cancers and viral infections.

[0037] The compositions of the invention are useful for the treatment of cancer, neurodegenerative and infectious, particularly viral, diseases and other conditions in which inappropriate or detrimental expression of the human PD-1 and/or is a component of the etiology or pathology of the condition. Hence, the invention provides an antibody as defined in the claims for use in therapy, which encompasses methods for treating cancer or inhibiting tumor progression in a subject in need thereof with a subject anti-PD-1 protein.

25

[0038] The disclosure includes all combinations of the recited particular embodiments.

30

EXAMPLES

Example 1. Generation of anti-PD-1 monoclonal antibody

[0039] Anti-PD-1 monoclonal antibodies (mAbs) were generated based on conventional hybridoma fusion technology (Kohler and Milstein 1976 Eur J Immunol 6:511-519; de St Groth and Sheidegger 1980, J Immunol Methods 35:1-21; Mechetner 2007 Methods Mol Biol 378:1-13) with minor modifications. MAbs with high binding activities in enzyme-linked immunosorbent assay (ELISA) and fluorescence-activated cell sorting (FACS) assay were selected for further characterization

35

PD-1 recombinant protein for immunization and binding assays

[0040] Expression plasmid containing full-length human PD-1 cDNA was obtained from Origene (Cat. No. SC117011, NCBI Accession No: NM_005018.1, Beijing, China). The extracellular domain consisting of amino acid (AA) 1-168 of PD-1 (SEQ NO.1, SEQ NO.2) was PCR-amplified, and subcloned in pcDNA3.1-based expression vector (Invitrogen, Carlsbad, CA, USA) with C-terminus fused either to a His₆ tag or to the γ Fc domain of human IgG4 heavy chain, which resulted in two recombinant fusion protein expression plasmids, PD-1-EC/His and PD-1-EC/Fc (abbreviated as PD-1/His and PD-1/Fc). The schematic presentation of immunogen/antigen proteins were shown in Fig 1. For the recombinant fusion protein production, PD-1/His and PD-1/Fc plasmids were transiently transfected into 293-F cells in 1-3 liters of medium (Invitrogen), and cultured for 5-7 days in a CO₂ incubator equipped with rotating shaker. The supernatant containing the recombinant protein was collected and cleared by centrifugation at 15000g for 30 minutes. PD-1/His was purified through immobilized metal affinity chromatography using Ni-Sepharose Fast Flow (Cat. No. 17531801, GE Lifesciences, Shanghai, China), followed by size exclusion chromatography using a HiLoad 16/60 Superdex 200 column (Cat. No. 17106901, GE Lifesciences, Shanghai, China). PD-1/Fc was purified using a Protein G Sepharose Fast Flow column (Cat. No. 17061805, GE Lifesciences). Both PD-1/His and PD-1/Fc proteins were dialyzed against phosphate buffered saline (PBS) and stored in -80°C freezer in small aliquots.

45

50

55

[0041] The cDNA coding for human PD-L1 was chemically synthesized by Genescript (Nanjing, China) based on the published sequence (NCBI Accession No. NM_014143). The PD-L2 expression plasmid was purchased from Origene

EP 3 702 373 B9

(Cat. No. SC108873, NCBI Accession No. NM_025239.2, Beijing, China). Both cDNAs were cloned in pcDNA3.1/Hygro-mycin (Cat. No. V870-20, Invitrogen), and pcDNA3.1/V5-His (Cat. No. V810-20, Invitrogen), respectively.

Stable expression cell line

5
[0042] Stable cell lines expressing human PD-1, PD-L1 or PD-L2 were established by transfection of pcDNA3.1 plasmids containing PD-1, PD-L1 and PD-L2 to HUT78 (ATCC, Manassas, VA, USA) and HEK293 (ATCC), respectively, and followed by selection with medium containing 200 micrograms of hygromycin (Cat. No. 10687-010, Invitrogen) or 1 mg of G418 (Sigma) per milliliter. Single clones were isolated by conventional method, either limited dilution or picking up single colonies from culture-well surface. All clones were screened by Western blot and FACS analysis using anti-PD-1, PD-L1 and PD-L2 antibodies (Cat. No. 12-9969, 17-5983, 12-5888, eBioscience, San Diego, USA), respectively, and the top expression clones were selected for FACS binding assay to screen hybridoma monoclonal antibodies, or used in functional assays.

Immunization, hybridoma fusion and cloning

15
[0043] Eight to twelve week-old Balb/c mice (from BEIJING HFK BIOCSIENCE CO.,LTD, Beijing, China) were immunized subcutaneously with 100ul of adjuvant (Cat. No. KX0210041, KangBiQuan, Beijing, China) containing 5 micrograms of PD-1/Fc. The immunization was conducted by two injections of the above immunogen with three weeks apart. Two weeks after the 2nd immunization, the mice sera were evaluated for PD-1 binding by FACS (following sections). The mice with high anti-PD-1 antibody titers in sera were selected and boosted intraperitoneally with 50 micrograms of PD-1/Fc in the absence of any adjuvant. Three days after boosting, the splenocytes were isolated and fused with the murine myeloma cell line, SP2/0 cells (ATCC), using standard techniques (Geftter, M.L. et al., 1977 Somat Cell Genet, 3:231-236).

Assess PD-1 binding activity of antibodies by ELISA and FACS

25
[0044] The supernatants of hybridoma clones were initially screened by Enzyme-Linked Immuno-Sorbent Assay (ELISA) as described in "Flanagan, M.L. et al. 2007 Methods in Molecular Biology 378:33-52" with some modifications. Briefly, 50-200 nanograms of PD-1/His or PD-1/Fc protein in 50 microliters of phosphate buffered saline (PBS) were coated in 96-well plate (Shenzhen JinCanHua Industry Co., Ltd, Shenzhen, China) on per well base. The HRP-linked anti-mouse IgG antibody (Cat. No. 7076S, Cell Signaling Technology, USA and Shanghai, China) and chemiluminescent reagent (Cat. No. PA107-01, TIANGEN, China) were used to detect and develop the ELISA signal, which were read out by a plate reader (PHREASTAR FS, BMG LABTECH, Germany) at wavelength of 450 nm. The ELISA-positive antibody producer clones were further verified by fluorescence-activated cell sorting (FACS) using a conventional method. PD-1 stable expression cell lines, HuT78/PD-1 (10^5 cells/well), described above, was stained with supernatants from anti-PD-1 hybridomas in V-bottom 96-well plates (Cat. No. 3897, Corning, USA and Shanghai, China). To block human Fc receptors, cells were pre-incubated with human IgG (20 μ g/ml) (Cat. No. H11296, LifeHolder, USA and Shanghai, China). PD-1 antibodies were detected with Dylight™ 649-labelled goat anti-mouse IgG antibody (Cat. No. 405312, Biolegend, San Diego, USA) and cell fluorescence was monitored using a flow cytometer (Guava easyCyte 8HT, Merck-Millipore, USA and Shanghai, China).

35
[0045] The conditioned media of hybridoma cells that showed positive signal in both ELISA and FACS assay were subjected to functional assays to identify antibodies with good functional activity in human immune cell-based assays (herein). The antibodies with positive functional activity were further subcloned and characterized.

Subcloning and adaptation to serum-free or low serum medium

45
[0046] The positive hybridoma clones from primary screening through ELISA, FACS and functional assays were subcloned by the conventional method of limited dilution. Each of the positive clones was plated out in a 96-well plate, cultured in RPMI1640 medium (Cat. No. SH30809.01B, Hyclone, Shanghai, China) with 10% fetal bovine serum (FBS, Cat. No. SH30084.03, Hyclone, Beijing, China) in CO₂ incubator. Three subclones from each limited dilution plate were selected and characterized by FACS and functional assays. The subclones selected through functional assays were defined as monoclonal antibody. The top subclones were adapted for growth in the CDM4MAb medium (Cat. No. SH30801.02, Hyclone) with 1-3% FBS.

Expression and purification of monoclonal antibodies

55
[0047] Either murine monoclonal antibody-producing hybridoma cells or recombinant antibody plasmids-transfected 293-F cells (Cat. No. R79007, Invitrogen) were cultured in CDM4MAb medium (Cat. No. SH30801.02, Hyclone) or

Freestyle293 Expression medium (Cat. No. 12338018, Invitrogen), respectively, in a CO₂ incubator at 37°C for 5 to 7 days. The conditioned medium was collected through centrifugation at 10,000g for 30 minutes to remove all cells and cell debris, and filtrated through a 0.22µm membrane before purification. Murine or recombinant antibodies were applied and bound to a Protein A column (Cat. No. 17127901, GE Life Sciences) following the manufacturer's guidance, washed with PBS, eluted in the buffer containing 20mM citrate, 150mM NaCl, pH3.5. The eluted materials were neutralized with 1M Tris pH8.0, and usually contained antibodies of above 90% purity. The Protein A-affinity purified antibodies were either dialyzed against PBS or further purified using a HiLoad 16/60 Superdex200 column (Cat. No. 17531801, GE Life Sciences) to remove aggregates. Protein concentrations were determined by measuring absorbance at 280nm or by Bradford assay (Cat. No. 1856210, Thermo Scientific, Rockford, IL, USA) using bovine IgG of defined concentration (Cat. No. 23212, Thermo Scientific) as the standards. The purified antibodies were stored in aliquots in -80°C freezer.

Example 2. Comparison of binding activities among anti-PD-1 antibodies

[0048] Through screening thousands of hybridoma clones we identified some top monoclonal antibodies (mAb), which bind to human PD-1 with high specificity and strength. As shown in ELISA assay (Figure 2), three of the top antibodies elicited such binding strength and specificity. FACS analysis results demonstrated the selected monoclonal antibodies bind to the native PD-1 proteins expressed on cell surface. Murine mAb317 (mu317), mu326 and mu150 showed concentration-dependent binding activity, and their binding EC₅₀ (Effective concentration at 50% activity) was significantly lower than that of the control mu55 (Figure 3).

Assess mAb binding affinity by Surface Plasmon Resonance (SPR)

[0049] The mAbs with high binding activities in ELISA and FACS, as well as with potent functional activities in the cell-based assays (herein) were examined for their binding kinetic constant in real time binding reactions. Murine anti-PD-1 mAbs were purified from hybridoma supernatants using protein A Flow column (Cat. No. 17531801, GE Life Sciences) followed by exclusion chromatography using a HiLoad 16/60 Superdex200 column (Cat. No. 17106901, GE Life Sciences). The purified anti-PD-1 antibodies were concentrated to 0.5-1 mg/mL in PBS and stored in aliquots in -80°C freezer.

[0050] For determining binding affinities of PD-1 mAbs, SPR measurements were performed in HBS-N buffer (10 mM HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005% v/v surfactant P20, GE Healthcare) using the BIAcore™ T-200 instrument (GE Life Sciences). Anti-mouse Fc CM5 biosensor chip (GE Healthcare) was generated using a standard primary amine coupling protocol. PD-1 mAbs at 0.3 µg/ml were captured on anti-mouse Fc surface for 1 min at 10 µl/min. PD-1/Fc in a serial dilutions from 3.3nM to 120nM was injected over antibody-bound surface for 3 min at 30 µl/min followed by a 10 min dissociation phase. Association rates (K_a or k_{on}) and dissociation rates (K_d or k_{off}) were calculated using the one-to-one Langmuir binding model (BIA Evaluation Software, GE Life Sciences). The equilibrium dissociation constant (K_D) was calculated as the ratio k_{off}/k_{on} .

[0051] As shown in Table 1, both mu326 and mu517, a cognate sequence family member related to mu317, have a sub-nanomolar K_D equaling to 0.324 nM and 0.289 nM, respectively, which is significantly better than that of mu134. The K_{on} rate was similar among the three mAbs listed in Table 1, yet the K_{off} rate was significantly different, much faster dissociation rate was observed in mu134.

Table 1. Binding constant of certain top antibodies

mAbs	K_{on} (M ⁻¹ , S ⁻¹)	K_{off} (s)	K_D (M)
mu326	2.4×10^5	7.79×10^{-5}	3.24×10^{-10}
mu517	1.96×10^5	5.66×10^{-5}	2.89×10^{-10}
mu134	1.1×10^5	3.69×10^{-4}	3.35×10^{-9}

Affinity determination of anti-PD-1 Fabs by SPR

[0052] Anti-PD-1 mAbs were converted into Fab version by PCR to fuse the variable regions of heavy and light chains to the N-terminus of human IgG2-CH1 and constant region of kappa chain, respectively, and subcloned in pcDNA3.1 vector (Invitrogen). Both expression vectors were co-expressed in 293-F cells using a transient transfection protocol similar to the transient expression of whole antibodies. Briefly, the Fab kappa chain was PCR amplified and subcloned in pcDNA3.1-based expression vector (Invitrogen, Carlsbad, CA, USA). In a separate plasmid, the heavy chain variable region (VH) together with the CH1 coding sequence from human IgG2 was fused with a C-terminal c-Myc-His8 tag by overlapping PCR, and then subcloned in the expression vector. The C232S and C233S (Kabat residue numbering,

Kabat et al. Sequence of proteins of immunologic interest, 5th ed Bethesda, MD, NIH 1991) mutations were introduced in the IgG2 heavy chain to prevent disulfide bond exchange and stabilize human IgG2 in the IgG2-A conformation (Lightle et al. 2010 Protein Sci 19(4): 753-762). Both constructs contained a signal peptide upstream of the Fab mature sequences. Secreted expression of Fab was achieved by co-transfection of above 2 plasmids into 293-F cells and cell culture supernatants were harvested 6-7 days post transfection. His8-tagged Fabs were purified from cell culture supernatants using a Ni-sepharose Fast Flow column (Cat. No. 17531801, GE Life Sciences) followed by size exclusion chromatography using a HiLoad 16/60 Superdex200 column (Cat. No. 17106901, GE Life Sciences). The purified Fabs were concentrated to 0.5-5 mg/mL in PBS and stored in aliquots in -80°C freezer.

[0053] For affinity determinations of anti-PD-1 Fabs, SPR assays were used with the BIAcore™ T-200 instrument (GE Life Sciences). Briefly, human PD-1/His or cynomolgus monkey PD-1/His was coupled to activated CM5 biosensor chips (Cat. No. BR100530, GE Life Sciences) to achieve approximately 100-200 response units (RU), followed by blocking un-reacted groups with 1M ethanolamine. Fab samples of increasing concentration from 0.12nM to 90nM were injected in the SPR running buffer (10 mM HEPES, 150 mM NaCl, 0.05% Tween20, pH7.4) at 30 µL/minute, and binding responses on human PD-1/His or monkey PD-1/His were calculated by subtracting of RU from a blank flow-cell. Association rates (k_{on}) and dissociation rates (k_{off}) were calculated using the one-to-one Langmuir binding model (BIA Evaluation Software, GE Life Sciences). The equilibrium dissociation constant (K_d) was calculated as the ratio k_{off}/k_{on} .

[0054] The SPR-determined binding affinities of anti-PD-1 Fabs were listed in Table 18. Each anti-PD-1 Fab bound with high affinity ($K_d = 0.15-1$ nM) to human PD-1. All Fabs, except 326-3G1, bound with slightly lower but comparable (within 5 fold in K_d) affinities to cynomolgus monkey PD-1.

Example 3. Functional activity of anti-PD-1 antibodies in human T cells.

Generation of stable cell lines

[0055] Retroviral packaging cell line PT67, human T cell lines HuT78 and HEK293 were obtained from the American Type Culture Collection (ATCC, Rockville, MD). A HuT78 subline HuT78/PD-1 that expresses PD-1 was generated by retroviral transduction using pFB-neo vector (Stratagene/Agilent Tech, Santa Clara, CA) containing the PD-1 gene, according to the protocol described previously (Zhang et al. 2005 Blood 106: 1544-1551). The T cell engager, a membrane-anchored chimeric Ab (OS8), was constructed by fusing the single chain variable fragment (scFv) of an anti-human CD3 mAb OKT3 (Kipriyanov et al. 1997, PEDS 10:445-453) to the C-terminal domain (113-220) of mouse CD8 α (NCBI Accession No: NP_001074579.1) which includes hinge, transmembrane and cytoplasmic domains. By doing so, anti-CD3 scFv is anchored to cell surface as a T cell activator. Human PD-L1, PD-L2 and OS8 cDNAs were sub-cloned into pcDNA3.1 vector. Stable cell lines HEK293/OS8/PD-L1, Hep3B/OS8/PD-L1 and HEK293/OS8/PD-L2 that co-express both OS8 and PD-L1 or PD-L2 cDNAs were generated by co-transfection of HEK293 and Hep3B cells (ATCC) with the paired plasmids, followed by hygromycin or G418 selection for 10-14 days. Cell lines were then cloned by limiting dilution as described previously (Fuller SA, et al. Curr Protoc Mol Biol. Chapter 11:Unit11.8., 2001). Chimeric PD-1 receptor, named P3Z, was constructed by fusing the extracellular and transmembrane domains of human PD-1 to the cytoplasmic domain of human CD3 ζ chain (NCBI Accession No. NP_932170.1). P3Z-coding cDNA sequence was cloned into pFB-neo and delivered into HuT78 cells via retroviral transduction to generate HuT78/P3Z cells.

Determination of PD-1 antibody functions by IL-2 release in HuT78/PD-1 cells

[0056] To determine whether anti-PD-1 antibodies can block the interaction of PD-L1-induced PD-1 signaling, HuT78/PD-1 cells (1.5×10^4 cells per well in 96-well plate) were pre-incubated with hybridoma supernatants or PD-1 antibodies for 15 minutes prior to co-culture with HEK293/OS8/PD-L1 or HEK293/OS8/PD-L2 cells (4×10^4 per well) in a flat bottom plate fed with 200 µl of RPMI1640 growth medium per well at 37°C. After 16-18 hours, supernatants of the co-culture were collected. IL-2 was assayed by ELISA using human IL-2 Ready-Set-Go! ELISA kits (Cat. No. 88-7025, eBiosciences, San Diego, CA). In this assay, blockade of PD-1 signaling with anti-PD-1 antibodies resulted in enhanced TCR signaling and IL-2 production (Fig. 4).

[0057] As shown in Fig. 5 and Table 2, murine anti-PD-1 mAb, mu317 and mu326, elicited significantly higher functional activity than mu30, inhibiting PD-L1-induced PD-1 signaling which leads to increased IL-2 secretion. Both had higher IL-2 secretion (top line, Table 2), 675 and 634 pg/ml, respectively, and both had lower EC₅₀ (Effective concentration of mAb at 50% level of IL-2 secretion induction) than mu30 antibody.

Table 2. IL-2 release induced by anti-PD-1 mAbs in HuT78/PD-1 cells co-cultured with HEK293/OS8/PD-L1 cells

Antibody	Baseline (pg/ml)	Top line (pg/ml)	EC ₅₀ (µg/ml)
mu30	95	527	0.229

EP 3 702 373 B9

(continued)

Antibody	Baseline (pg/ml)	Top line (pg/ml)	EC ₅₀ (μg/ml)
mu317	95	675	0.083
mu326	95	634	0.053
mIgGs	95	N/A	N/A

Baseline: Average IL-2 release induced by mIgGs at all tested concentrations, see Fig. 4.
 Top line: Highest IL-2 release based on regression calculation by Prizm Software, Fig. 4.
 N/A: Not applicable

[0058] Not only did the engagement of HuT78/PD-1 cells by anti-PD-1 mAbs block PD-L1 induced T-cell activation, but also blocked PD-L2 induced IL-2 release. Table 3 presented the data showing mu317 and mu326 had much higher potency in activating the T-cells as indicated by the parameters (EC₅₀) of IL-2 secretion than those of mu476 .

Table 3. IL-2 release induced by anti-PD-1 mAbs in HuT78/PD-1 cells co-cultured with HEK293/OS8/PD-L2 cells

Antibody	Baseline (pg/ml)	Top line (pg/ml)	EC ₅₀ (μg/ml)
476	180	599	0.183
317	192	563	0.032
326	218	635	0.038

Baseline: Average IL-2 release induced in the lower tail part of the sigmoid reaction curve.
 Top line: Average IL-2 release induced at the plateau part of the sigmoid reaction curve

Determination of PD-1 antibody functions by reverse signaling of IL-2 release in HuT78/P3Z cells

[0059] In chimeric receptor P3Z, PD-1 signaling domain was replaced with the cytoplasmic domain of CD3ζ. Therefore, P3Z mediates activation upon engagement with PD-L1, rather than inhibition as original PD-1 receptor. In this assay, HuT78/P3Z cells (3×10⁴/well) were pre-incubated with hybridoma supernatants or PD-1 antibodies for 15 minutes prior to co-culture with HEK293/PD-L1 or HEK293/PD-L2 cells (5×10⁴/well) in 96-well flat bottom plates (a total volume of 200 μl/well) at 37°C. After 16-18 hours, supernatants were collected and IL-2 production was assayed by ELISA as described above.

[0060] The functional activity of murine anti-PD-1 mAbs was further confirmed by direct read-out of T-cell activation in reverse signaling assay described above. Consistent to the result described above, mu317 and mu326 had best functional activity among the mAbs we screened. As shown in Table 4 and Table 5, mu317 and mu326 were much more potent than one of the low activity mAbs, mu37, both in terms of IC₅₀ and maximum inhibition.

Table 4. Inhibition of IL-2 secretion by anti-PD-1 mAbs in HuT78/P3Z cells co-cultured with HEK293/PD-L1 cells

Antibody	IC ₅₀ (μg/ml)	Max inhibition, %
37	0.287	86.9
317	0.083	99.3
326	0.039	97.6

Maximum inhibition was calculated as percentage (%) of inhibition with anti-PD-1 mAbs added to the highest level of 10 μg/ml in culture.

Table 5. Inhibition of IL-2 secretion by anti-PD-1 mAbs in HuT78/P3Z cells co-cultured with HEK293/PD-L2 cells

Antibody	IC ₅₀ (μg/ml)	Max inhibition, %
37	0.127	43.3

EP 3 702 373 B9

(continued)

Antibody	IC ₅₀ (μg/ml)	Max inhibition, %
317	0.020	94.3
326	0.018	93.4

Maximum inhibition was calculated as percentage (%) of inhibition with anti-PD-1 mAbs added to the highest level of 10 μg/ml in culture.

Example 4. Activation of IFN-γ secretion by anti-PD-1 mAb in primary human PBMCs co-cultured with HEK293/OS8/PD-L1 cells

[0061] To verify if the selected top mAbs against PD-1 also exert functional effect on primary human immune cells, we assayed the antibody function by using freshly isolated peripheral blood mononuclear cells (PBMCs), which are mainly consisted of T-cells (50-70%), B-cells and NK cells (15-30%), and monocytes (2-10%). Human PBMCs were isolated from healthy donors by density gradient centrifugation using ficoll lymphocyte separation medium (Histopaque-1077; Sigma-Aldrich, MO) according to the manufacturer's instructions. All the human blood collection followed the Internal Procedure of Beigene. PBMCs were then stimulated with anti-CD3 mAb (40 ng/mL) OKT3 (Cat. No. 16-0037, eBioscience, CA) for 3 days prior to assay. FACS analysis (Example 1) showed that PD-1 expression on the activated PBMCs (primarily T cells) was increased to variable degree dependent on individual donors (Table 6). To determine the response of pre-activated T cells to PD-1 ligand-positive tumor cells upon engagement TCR/CD3 complex, PBMCs (1×10⁴) were co-cultured with either HEK293/OS8/PD-L1 or HEK293/OS8/PD-L2 cells (3×10⁴) in 96-well flat-bottom plates for 15-18 hours. Cell-free supernatants were assayed for IFN-γ level by ELISA using Ready-Set-Go! ELISA kits (Cat. No. 88-7316, eBiosciences), which is the most prominent indicator of T-cell activation, as well as of other immune cell activation (Thakur A. et al. 2012 Vaccine, 30:4907-4920).

PBMCs and treatment	Percent gated PD-1 staining positive cells versus total PMBCs stained	
	Donor-3	Donor-4
PBMCs, not stimulated / stained by PD-1 Ab	12.0%	3.2%
PBMCs, stimulated / stained by PD-1 Ab	40.0%	38.1%
PBMCs, not stimulated / stained by control Ab	≤ 0.5%	≤ 0.5%
PBMCs, stimulated / stained by control Ab	≤ 0.5%	≤ 0.5%

Stimulation: freshly isolated PBMCs were cultured for 3 days in presence of anti-CD3 antibody, OKT3, and IL-2.
Without stimulation: fresh PBMCs subjected to antibody staining and FACS analysis.

[0062] Fig. 6 demonstrated that presence of mAbs mu317 and mu326 in the co-culture of pre-activated PBMCs and HEK293/OS8/PD-L1 cells resulted in increasing IFN-γ accumulation in a dose-dependent manner. Although the base level of IFN-γ with control murine IgG treatment varies among different donors, the increase of IFN-γ secretion in PBMCs treated by mu317 or mu326 is statistically significant in the range of 0.1 to 10 μg/ml of antibody treatment. Comparing to the corresponding level of mIgG-treated PBMCs, IFN-γ secretion induced by mu317 and mu326 between the 0.1 to 10 μg/ml concentration levels increased 2.5 to 3.2 fold in PBMCs from Donor-19, and increased 1.4 to 2.3 fold in PBMCs of Donor-20, respectively.

Example 5. Activation of human NK cells by anti-PD1 mAbs

Stable cell lines for functional assay in NK cells

[0063] Primary human NK cells were reported previously to express PD-1 protein in response to IL-2 treatment and inhibiting PD-1-mediated signaling enhanced cytotoxicity of NK cells (2010 Blood, 116: 2286). For quantitative assay of functional effect exerted by anti-PD-1 mAbs in NK cells, human NK cell line NK92MI (ATCC) and lung cancer cell line SK-Mes-1 (ATCC) were engineered to stably express human PD-1 and PD-L1, respectively, by retroviral transduction according to the protocols described previously (Zhang et al. 2005, Blood 106: 1544-1551, Zhang et al. 2006, Cancer Res, 66: 5927). The two stable cell lines were named as NK92MI/PD-1 and SK-Mes-1/PD-L1

Anti-PD-1 Abs promote IFN- γ production and secretion in NK92MI/PD-1 cells

[0064] Functional activity of the anti-PD-1 mAbs on NK cells was assayed by quantitative measurement of IFN- γ production and secretion in NK92MI/PD-1 cells which were co-cultured with lung cancer cell line SK-MES-1/PD-L1 at ratio of 1 to 2 in 96-well flat-bottom plate with total of 6×10^4 cells per well. The anti-PD-1 mAbs were added to NK92MI/PD-1 cells 15 minutes before the co-culture started, then the cells were co-cultured for overnight in CO₂ incubator. Cell-free supernatants were assayed for IFN- γ level by ELISA as described in Example 4.

[0065] All anti-PD-1 mAbs triggered significant increase of IFN- γ production from the baseline with low concentration of antibody treatment to top line with high concentration of antibody treatment. The two top antibodies, mu317 and mu326, had lower EC₅₀, than the comparison antibody 5C, indicating they have more potent activating effect to the NK cells (Table 7).

Table 7. IFN- γ secreted in medium (pg/ml) by NK92MI/PD-1 cells in presence of anti-PD-1 mAb and SK-MES-1/PD-L1 cells

Antibody	Baseline (pg/ml)	Top line (pg/ml)	EC ₅₀ (μ g/ml)
317	28	532	0.40
326	15	509	0.20
5C	20	535	1.17

Baseline: Average IFN- γ release induced in the lower tail part of the sigmoid reaction curve.
Top line: Average IFN- γ release induced at the plateau part of the sigmoid reaction curve

Anti-PD-1 antibody enhances cancer cell killing mediated by NK92MI/PD-1 cells

[0066] Cytotoxicity of NK92MI/PD-1 cells against SK-MES-1/PD-L1 cells was determined by lactate dehydrogenase (LDH) release assay using the CytoTox 96 Non-Radioactive Cytotoxicity Assay kit (Promega, Madison, WI). In brief, NK92MI/PD-1 cells (10^5) were pre-incubated with anti-PD-1 mAbs at final concentrations within the range of 0.004-10 μ g/ml for 15 minutes, and SK-MES-1/PD-L1 cells (2×10^4) were added to the immune cell culture in a 96-well V-bottom plate at an effector to tumor cell (E:T) ratio of 5:1, then co-cultured for 5 hours. The complete tumor cell lysis was set as maximum cell killing, the LDH-release assay readout of each sample was calculated as percentage of maximum cell killing. The cell killings (%) of all samples were normalized cross the plates using 10% of baseline as the common standard.

[0067] In the specific cytotoxicity assay set as above, the selected anti-PD-1 mAbs caused a net tumor cell killing (= top line - baseline) ranging from 19% to 20.2% at high concentration of mAb input. Mu317 and mu326 had lower EC₅₀ than mu336, indicating better potency to trigger NK92MI/PD-1 cell-mediated tumor cell killing (Table 8).

Table 8. Cytotoxicity of NK92MI/PD-1 cells towards tumor cells induced by anti-PD-1 mAb

Antibody	Baseline (%)	Top line (%)	EC ₅₀ (μ g/ml)
317	10	29.06	0.50
326	10	30.19	0.37
336	10	29.72	1.52

Baseline: Percent of tumor cells killed not due to the effect of anti-PD-1 mAbs, normalized to 10% cross plates.
Top line: Average percent of tumor killed in presence of highest concentrations of mAbs, i.e. 3 μ g/ml and 10 μ g/ml

Example 6. Cloning and sequence analyses of PD-1 mAbs

[0068] The murine hybridoma clones secreting a specific mAb were cultured to a density of 3 to 10×10^6 cells in a 100mm-tissue culture dish, and the cells were harvested through centrifugation at 1500 rpm in a swing bucket rotor. Total cellular RNA was isolated using Ultrapure RNA kit (Cat. No. CW0581, CWBIOTECH, Beijing, China) following the manufacturer's protocol. The RNA was resuspended in double-deionized water, concentration measured by NanoDrop (ThermoFisher, Shanghai, China).

[0069] PCR primers used for mAb cDNA cloning were synthesized by Invitrogen (Beijing, China) based on the sequences reported previously (Brocks et al. 2001 Mol Med 7:461-469). The 1st strand cDNA was synthesized using reverse transcriptase (Cat. No. AH301-02, Transgen Biotech, Beijing, China). PCR amplification of specific mAb cDNA

was performed using PCR reagent kit (Cat. No. Ap221-12, TransGen Biotech, Beijing, China) and following manufacturer's protocol. The PCR product was either directly sequenced by service provider (GeneWiz, Beijing, China) or subcloned into a pCR vector (Invitrogen), subsequently sequenced (GeneWiz).

[0070] The protein sequences of murine mAbs were analyzed by sequence homology alignment. MAbs were grouped based on sequence homology and epitope-mapping results (Example 13). Complement determinant regions (CDRs) were identified based on Kabat (Wu, T.T. and Kabat, E.A., 1970 J. Exp. Med. 132: 211-250) and IMGT system (Lefranc M.-P. et al., 1999 Nucleic Acids Research, 27, 209-212) by sequence annotation and by internet-based sequence analysis (http://www.imgt.org/IMGT_vquest/share/textes/index.html and <http://www.ncbi.nlm.nih.gov/igblast/>). As shown in Table 9, the CDRs of mu317 and mu326 are very different in sequence length and identity.

Table 9. CDRs of mu317 and mu326

MAbs	CDR1	SEQ ID NO	CDR2	SEQ ID NO	CDR3	SEQ ID NO
mu317, HC	<u>GFSLSYGVH</u>	11	VIWAGGSTNYNSALMS	12	<u>ARAYGNYWYIDV</u>	13
mu317, LC	KASQSVSNDVA	14	<u>YAFHRFT</u>	15	<u>HQAYSSPYT</u>	16
mu326, HC	<u>GYFTNYGMN</u>	17	WINNNNGEPTYAEFKG	18	<u>ARDVMDY</u>	19
mu326, LC	RASEVDNYGYSFMH	20	<u>RASNLES</u>	21	<u>QQSKEYPT</u>	22

Note: CDRs in bold face are based on Kabat system; CDRs underlined are based IMGT system.

Example 7. Humanization of the murine mAbs

Simulation of antibody 3D structure

[0071] The three dimensional structures were simulated for variable domains of mu317 and mu326 in order to identify framework residues that might be important for supporting CDR loop structures. Potentially important framework residues were kept as the original murine residues in the first round antibody humanization. The previously established structural modeling method for antibodies (Morea et al. Methods 2000 20:267-279) was adopted to simulate 3D structure of anti-PD-1 mAbs based on the known canonical structures of antibodies (Al-Lazikani et al. 1997 Journal of Molecular Biology 273:927-948). Briefly, the sequence of each variable domain (Vk and Vh) of murine antibody was blasted in the PDB database (Protein Data Bank, <http://blast.ncbi.nlm.nih.gov/>) to identify the most homologous antibody sequence with known high resolution structure (resolution less than 2.5 angstrom). Selected structure templates for modeling mu317 and mu326 (listed in Table 10) had the same classes of canonical loop structures in L-CDR1, L-CDR2, L-CDR3, H-CDR1, and H-CDR2 to the target antibodies to be modeled. If the templates for the Vk and the Vh came from different immunoglobulins, they were packed together by a least-squares fit of the main chain atoms to form a hybrid structure of Vk-Vh interface residues, which was used as the templates for structural homology modeling by Swiss-model program (Kiefer et al. 2009 Nucleic Acids Research 37, D387-D392). Certain side chain conformation was adjusted while the main chain conformations were retained. At the sites where the parent structure and the modeled structure had the same residue, the side chain conformation was retained. At sites where the residues were different, side chain conformations were modeled on the basis of template structure, rotamer libraries and packing considerations. After homology modeling, PLOP program (Jacobson et al. 2002 Journal of Physical Chemistry 106:11673-11680) was used to refine the homology models to minimize all-atom energy and optimize Vk and Vh interface. This step was performed to improve the stereochemistry, especially in those regions where segments of structures coming from different antibodies had been joined together.

Table 10. Structure templates used in antibody structure simulations

Antibody chain	PDB code of template structure (PDB template for H-CDR3)	Sequence identity	Sequence similarity
mu317 Vk	3MXV	87%	92%

EP 3 702 373 B9

(continued)

Antibody chain	PDB code of template structure (PDB template for H-CDR3)	Sequence identity	Sequence similarity
mu317 Vh	3VFG	83%	91%
mu326 Vk	1EJO	92%	94%
mu326 Vh	1NCA	88%	90%
317-1 Vk	4HJJ	90%	95%
317-1 Vh	3VFG (1AY1)	75%	87%
326-1 Vk	1EJO	87%	92%
326-1 Vh	3T2N (3CXD)	84%	86%

[0072] The structures were also simulated for CDR-grafted 317-1 and 326-1 in order to guide further rounds of antibody engineering to enhance the extents of humanization and/or enhance antibody stabilities. The selected structure templates are also listed in Table 10. The structure simulations were done in a similar way to above procedure, except that the possible conformations of H-CDR3 were taken from PDB templates 1AY1 for 317-1 and 3CXD for 326-1, respectively, which contained H-CDR3s of similar size and torso region. Energy minimization for grafted H-CDR3 residues was done using PLOP.

Humanization

[0073] For humanization of the anti-PD-1 mAbs, we searched human germline IgG genes homologous to the cDNA sequences of mu317 and mu326 variable regions by blasting the human immunoglobulin gene database in IMGT (<http://www.imgt.org/IMGT/vquest/share/textes/index.html>) and NCBI (<http://www.ncbi.nlm.nih.gov/igblast/>) websites. The human IGVH and IGV κ with high homology to the PD-1 mAbs were selected as the template for humanization.

[0074] Humanization was carried out in principle by CDR-grafting. In the 1st round of humanization, mutations from murine to human amino acid residues in framework sequences of variable regions was guided by the simulated 3D structures, and only the murine amino acid residues whose changes retain the overall antibody and CDR loop structure were mutated to human sequence as described above. The initial versions of humanized mAbs were hu317-1 (SEQ NO 47-50) and hu326-1 (SEQ NO 55-58), which comprise a heavy chain with humanized variable heavy chain (Vh) fused to human IgG2 constant region (NCBI accession No. P01859) and a light chain with humanized variable light chain kappa (V κ) fused to human Ig kappa C-region (NCBI Accession No. P01834). Likewise, we generated chimeric antibodies from mu317 and mu326, which are consisted of a murine VH fused to human IgG2 constant region and a murine V κ fused to human Ig kappa C-region. The full chimeric antibodies were named as ch317 and ch326, respectively. All recombinant mAbs were expressed and purified as described in Example 1.

[0075] FACS and functional assays demonstrated that mAb hu317-1 almost retained the same binding and functional activity as the mu317 and ch317. The EC₅₀ difference in FACS analysis between mu317 versus ch317 and hu317-1 may be interpreted by the fact that two different detection antibodies, a goat anti-mouse IgG and a goat anti-human IgG, were used in FACS. In the two functional assays, all three versions of 317 were treated more equal, and the results also close to each other (Table 11).

[0076] As result of the initial round of humanization for mu326, mAb hu326-1 retained similar functional feature to the parental ch326 and mu326 although functional activity in FACS binding assay and in HuT78/PD-1 cell-based IL-2 release assay may be slightly weaker than ch326 (Table 12).

Table 11. Comparison of mu317, ch317 and hu317-1 by FACS and functional assays

Assay/Parameter		mu317	ch317	hu317-1
FACS	EC ₅₀ (µg/ml)	0.11	0.36	0.46
	Top MFI*	205	217	203
Assay-1	EC ₅₀ (µg/ml)	0.11	0.08	0.09
	Top line (pg/ml)	346	294	386
	Baseline (pg/ml)	98	82	91
Assay-2	IC ₅₀ (µg/ml)	0.11	0.10	0.11
	Max inhibition	99.5%	99.0%	99.8%

*MFI: mean fluorescence intensity from FACS analysis
 Assay-1: IL-2 release induced by the mAbs in HuT78/PD-1 cells co-cultured with HEK293/OS8/PD-L1 cells
 Assay-2: IL-2 release induced by the mAbs in HuT78/P3Z cells co-cultured with HEK293/PD-L1 cells

Table 12. Comparison of mu317, ch317 and hu317-1 by FACS and functional assays

Assay/Parameter		mu326	ch326	hu326-1
FACS	EC ₅₀ (µg/ml)	0.126	0.072	0.117
	Top MFI	195	163	129
Assay-1	EC ₅₀ (µg/ml)	0.038	0.074	0.112
	Top line (pg/ml)	1149	1057	1143
	Baseline (pg/ml)	242	250	283
Assay-2	IC ₅₀ (µg/ml)	0.14	0.12	0.10
	Max inhibition	96.9%	81.0%	84.4%

Assay-1: IL-2 release induced by the mAbs in HuT78/PD-1 cells co-cultured with HEK293/OS8/PD-L1 cells
 Assay-2: IL-2 release induced by the mAbs in HuT78/P3Z cells co-cultured with HEK293/PD-L1 cells

[0077] Based on the 1st round of humanization, we further mutated the other murine amino acid (AA) residues in the framework (FR) of hu317-1_Vh and V_k individually to assess the impact on the antibody function. As shown in Table 13, the seven individual mutations in Vh and one mutation in V_k of hu317-1 all have similar functional activities. Only minor changes were observed in some Vh mutation, such as hu317-2_K71V with slightly weaker inhibitory function among the mutations. However, when all the murine amino acid residues mutated together to human (hu317-3A), the

EP 3 702 373 B9

function is clearly weaker than the rest mutations in FACS and IL-2 release assays.

[0078] In the initial trial described above, hu326-1 reached significant humanization level in the FR except for a few of murine AA residues left. Yet, it has weaker function than the mu326. Therefore, we made more individual mutations either back to murine residues or forward to human residues to explore the contribution of each individual AA to mAb326 function. Table 14 presented all single AA mutations made based on hu326-1_Vh template (SEQ NO 56, SEQ NO 57) and their functional assay results. Majority of the mutations showed better functional activity than those of hu326-1, matching the original mu326 mAb. A couple of mutations (E46K and F95Y) showed slightly less potency in the EC₅₀ or IC₅₀, indicating the role of those residues in the antibody structure and function.

Table 13. Comparison of functional activity of Fabs with humanization mutations in hu317-1 framework

Fab and composition		FACS, EC ₅₀	IL-2 release in HuT78/P3Z	
Vh	V _K		Max inhibition, %	EC ₅₀
hu317-1_Vh	hu317-1 V _K	0.19	98.78	0.30
hu317-2_L48I	hu317-1 V _K	0.14	98.51	0.37
hu317-2_L67V	hu317-1 V _K	0.15	98.57	0.30
hu317-2_K71V	hu317-1 V _K	0.18	96.55	0.48
hu317-2_N73T	hu317-1 V _K	0.15	98.29	0.31
hu317-2_S76N	hu317-1 V _K	0.13	98.56	0.28
hu317-2_V78F	hu317-1 V _K	0.18	98.03	0.38
hu317-2_M82L	hu317-1 V _K	0.13	98.47	0.27
hu317-1_Vh	HU317-2_G100Q	0.21	98.86	0.27
hu317-3A	hu317-1 V _K	0.32	79.66	0.35

Note: Unit for EC₅₀ is μg/ml; mutated amino acid residue numbering is same as in the listed sequences for hu317-1; hu317-3A has all the framework sequence mutated to human.

Table 14. Comparison of functional activity of mAbs with mutations in hu326-1 framework

Antibody	FACS, EC ₅₀ μg/ml	IL-2 release in HuT78/P3Z		IL-2 release in HuT78/PD-1	
		Max inhibition, %	IC ₅₀ , μg/ml	Top line, pg/ml	EC ₅₀ , μg/ml
ch326	0.118	93.05	0.074	993	0.135
hu326-1	0.317	92.38	0.087	987	0.213
hu326-2 S9P ^B	0.145	96.04	0.075	1022	0.136
hu326-2 A16E ^B	0.155	96.33	0.078	1048	0.126
hu326-2 E46K ^B	0.132	95.25	0.079	1244	0.259
hu326-2 G63D ^B	0.139	96.44	0.064	1069	0.120
hu326-2 A76V ^F	0.102	96.65	0.071	1002	0.112
hu326-2 S84N ^B	0.131	96.52	0.060	1015	0.126
hu326-2 S85N ^B	0.110	95.62	0.093	932	0.104
hu326-2 T88N ^B	0.098	95.85	0.102		
hu326-2 F95Y ^F	0.097	95.62	0.166	1028	0.135

^B: Back mutation to murine amino acid; ^F: Forward mutation to human amino acid. All of the mutations were made in hu326-1_Vh (SEQ NO 56), which were paired with hu326-1_Vk (SEQ NO 58).

[0079] To explore the best possible Vh and V_K sequence composition for mAbs 317 and 326 that could be used as

EP 3 702 373 B9

therapeutics in human, we made a variety of combination mutations (including some mutations in the CDR sequences) in considerations of the antibody features, such as humanization level in FR, functional activities, physicochemical properties, antibody-dependent cell-mediated cytotoxicity (ADCC) and complement-dependent cytotoxicity (CDC). Most of the mutations were deemed not passing the qualification standards. Through the engineering process, six of the humanized, recombinant mAbs were selected for their potential therapeutic utility: hu317-4B2 (SEQ ID NO 43-44), hu317-4B5 (SEQ ID NO 45-46), hu317-4B6 (SEQ ID NO 23-26), hu326-3B1 (SEQ ID NO 51-52), hu326-3G1 (SEQ ID NO 53-54) and hu326-4A3 (SEQ ID NO 27-30). The CDRs of the mAb were compared to those of original murine antibodies, shown in Table 15 and Table 16.

[0080] Among the six mAbs, hu317-4B2, hu317-4B5 and hu317-4B6 are closely related to each other in sequences and very similar in their functional activities and strength. On the other hand, hu326-3B1, hu326-3G1 and hu326-4A3 are quite close to each other in sequences and functionalities (Table 17-18). Within each of the two groups of mAbs, they also shared many other features in addition to sequences and function, such as physicochemical properties and binding epitopes (described in Examples 10 and 11) though some minor differences do exist.

Table 15. Comparison of CDRs among different versions of mAbs 317

mAbs	CDR1	SEQ ID NO	CDR2	SEQ ID NO	CDR3	SEQ ID NO
mu317, HC	GFSLSYGVH	11	VIWAGGSTNYNSALMS	12	ARAYGNYWYIDV	13
hu317-1, HC	GFSLSYGVH	11	VIWAGGSTNYN <u>PSLKS</u>	59	ARAYGNYWYIDV	13
hu317-4B2, HC	GFSLSYGVH	11	VIYAGGSTNYN <u>PSLKS</u>	60	ARAYGNYWYIDV	13
hu317-4B5, HC	GFSLSYGVH	11	VIYAGGSTNYN <u>PSLKS</u>	60	ARAYGNYWYIDV	13
hu317-4B6, HC	GFSLSYGVH	11	VIYADGSTNYN <u>PSLKS</u>	32	ARAYGNYWYIDV	13
mu317, LC	KASQSVSNDVA	14	YAFHRFT	15	HQAYSSPYT	16
hu317-1, LC	KASQSVSNDVA	14	YAFHRFT	15	HQAYSSPYT	16
hu317-4B2, LC	<u>K</u> <u>S</u> <u>S</u> <u>E</u> <u>S</u> <u>V</u> <u>S</u> <u>N</u> <u>D</u> <u>V</u> <u>A</u>	61	YAFHRFT	15	HQAYSSPYT	16
hu317-4B5, LC	<u>K</u> <u>S</u> <u>S</u> <u>E</u> <u>S</u> <u>V</u> <u>S</u> <u>N</u> <u>D</u> <u>V</u> <u>A</u>	61	YAFHRFT	15	HQAYSSPYT	16
hu317-4B6, LC	<u>K</u> <u>S</u> <u>S</u> <u>E</u> <u>S</u> <u>V</u> <u>S</u> <u>N</u> <u>D</u> <u>V</u> <u>A</u>	61	YAFHRFT	15	HQAYSSPYT	16

Note: AA residues underlined are changed from murine sequence to human antibody sequences or for improvement of physicochemical properties.

Table 16. Comparison of CDRs among different versions of mAbs 326

mAbs	CDR1	SEQ ID NO	CDR2	SEQ ID NO	CDR3	SEQ ID NO
mu326, HC	GYTFTNYGMN	17	WINNNNGEPTYAEEFKG	18	ARDVMDY	19
hu326-1, HC	GYTFTNYGMN	17	WINNNNGEPTYA <u>QG</u> FRG	62	ARDVMDY	19
hu326-3B1, HC	GYTFTNYGMN	17	WINNNNGEPTYA <u>QD</u> FRG	62	ARDVMDY	19
hu326-3G1, HC	GYTFTNYGMN	17	WINNNNGEPTYA <u>QD</u> FRG	62	ARDVMDY	19
hu326-4A3, HC	GYTFTNYGMN	17	WINNNNA <u>E</u> PTYA <u>QD</u> FRG	38	ARDVMDY	19

EP 3 702 373 B9

(continued)

mAbs	CDR1	SEQ ID NO	CDR2	SEQ ID NO	CDR3	SEQ ID NO
mu326, LC	RASESVDNYGY ^u SFMH	20	RASNLES	21	QQSKEYPT	22
hu326-1, LC	RASESVDNYGY ^u SFMH	20	RASNLES	21	QQSKEYPT	22
hu326-3B1, LC	RASESVDNYGY ^u SFMH	20	RASNLES	21	QQSKEYPT	22
hu326-3G1, LC	RASESVDNYGY ^u SFMH	20	RASNLES	21	QQSKEYPT	22
hu326-4A3, LC	RASESVDNYGY ^u SFMH	20	RASNLES	21	QQSKEYPT	22

Note: AA residues underlined are changed from murine sequence to human antibody sequences or for improvement of physicochemical properties.

Table 17. Binding activities of humanized mAbs assayed by ELISA and FACS

mAbs	ELISA, EC ₅₀ µg/ml	FACS, EC ₅₀ µg/ml
hu317-4B2	0.066	0.129*
hu317-4B5	0.057	0.115*
hu317-4B6	0.061	0.092*
hu326-3B1	0.092	0.165
hu326-3G1	0.088	0.190
hu326-4A3	0.091*	0.142*

* FACS data by using Fab version of antibodies without normalization.
** Data from bridging study and normalized.

Table 18. Binding affinity of Fabs assayed by SPR

Fab	K _{on} (M ⁻¹ , s ⁻¹)	K _{off} (s)	K _D (M)
hu317-4B5	3.89 × 10 ⁵	9.07 × 10 ⁻⁵	2.33 × 10 ⁻¹⁰
hu317-4B6	5.71 × 10 ⁵	8.37 × 10 ⁻⁵	1.47 × 10 ⁻¹⁰
hu326-3B1	2.18 × 10 ⁵	1.90 × 10 ⁻⁴	8.70 × 10 ⁻¹⁰
hu326-3G1	2.00 × 10 ⁵	2.01 × 10 ⁻⁴	1.00 × 10 ⁻⁹

Affinity determination of humanized anti-PD-1 Fabs by SPR

[0081] Anti-PD-1 mAbs were converted into Fab version by PCR to fuse the variable regions of heavy and light chains to the N-terminus of human IgG2-CH1 and constant region of kappa chain, respectively, and subcloned in pcDNA3.1 vector (Invitrogen). Both expression vectors were co-expressed in 293-F cells using a transient transfection protocol similar to the transient expression of whole antibodies. Briefly, the Fab kappa chain was PCR amplified and subcloned in pcDNA3.1-based expression vector (Invitrogen, Carlsbad, CA, USA). In a separate plasmid, the heavy chain variable region (VH) together with the CH1 coding sequence from human IgG2 was fused with a C-terminal c-Myc-His8 tag by overlapping PCR, and then subcloned in the expression vector. The C232S and C233S (Kabat residue numbering, Kabat et al. Sequence of proteins of immunologic interest, 5th ed Bethesda, MD, NIH 1991) mutations were introduced in the IgG2 heavy chain to prevent disulfide bond exchange and stabilize human IgG2 in the IgG2-A conformation (Lightle et al. 2010 Protein Sci 19(4): 753-762). Both constructs contained a signal peptide upstream of the Fab mature sequences. Secreted expression of Fab was achieved by co-transfection of above 2 plasmids into 293-F cells and cell culture

supernatants were harvested 6-7 days post transfection. His8-tagged Fabs were purified from cell culture supernatants using a Ni-sepharose Fast Flow column (Cat. No. 17531801, GE Life Sciences) followed by size exclusion chromatography using a HiLoad 16/60 Superdex200 column (Cat. No. 17106901, GE Life Sciences). The purified Fabs were concentrated to 0.5-5 mg/mL in PBS and stored in aliquots in -80°C freezer.

[0082] For affinity determinations of anti-PD-1 Fabs, SPR assays were used with the BIAcore™ T-200 instrument (GE Life Sciences). Briefly, human PD-1/His or cynomolgus monkey PD-1/His was coupled to activated CM5 biosensor chips (Cat. No. BR100530, GE Life Sciences) to achieve approximately 100-200 response units (RU), followed by blocking un-reacted groups with 1M ethanolamine. Fab samples of increasing concentration from 0.12nM to 90nM were injected in the SPR running buffer (10 mM HEPES, 150 mM NaCl, 0.05% Tween20, pH7.4) at 30µL/minute, and binding responses on human PD-1/His or monkey PD-1/His were calculated by subtracting of RU from a blank flow-cell. Association rates (k_{on}) and dissociation rates (k_{off}) were calculated using the one-to-one Langmuir binding model (BIA Evaluation Software, GE Life Sciences). The equilibrium dissociation constant (K_d) was calculated as the ratio k_{off}/k_{on} .

[0083] The SPR-determined binding affinities of anti-PD-1 Fabs were listed in Table 18. Each anti-PD-1 Fab bound with high affinity ($K_d = 0.15$ -1 nM) to human PD-1. All Fabs, except 326-3G1, bound with slightly lower but comparable (within 5 fold in K_d) affinities to cynomolgus monkey PD-1.

Example 8. Generation and expression of recombinant anti-PD-1 mAbs with modified human IgG4 constant region

[0084] Since PD-1 is primarily expressed in activated T cells, PD-1 blocking antibodies linked to naturally occurring type of IgG- \square Fc moieties are expected to induce \square Fc-mediated effector functions, such as ADCC and CDC, to a variable degree depending on the IgG subclasses, which results in elimination of activated T cells (Natsume A, et al, 2009 Drug Des Devel Ther. 3: 7-16). Human antibody subclass IgG4 was shown in many previous reports that it has modest ADCC and almost no CDC effector function (Moore GL, et al. 2010 MAbs, 2:181-189). On the other hand, natural IgG4 was found less stable in stress conditions such as in acidic buffer or under increasing temperature (Angal, S. 1993 Mol Immunol, 30:105-108; Dall'Acqua, W. et al, 1998 Biochemistry, 37:9266-9273; Aalberse et al. 2002 Immunol, 105:9-19). In order to spare PD-1⁺ T cells from being killed and to improve physicochemical properties of the anti-PD-1 antibodies, the humanized mAbs were linked to IgG4 engineered by combinations of mutations to have reduced or null Fc γ R binding or C1q binding activities, therefore, attenuating or eliminating ADCC and CDC effector functions. Considering physicochemical properties of antibody as a biological drug, one of the less desirable, intrinsic properties of IgG4 is dynamic separation of its two heavy chains in solution to form half antibody, which lead to bi-specific antibodies generated *in vivo* via a process called "Fab arm exchange" (Van der Neut Kolfshoten M, et al. 2007 Science, 317:1554-157). The mutation of serine to proline at position 228 (EU numbering system) appeared inhibitory to the IgG4 heavy chain separation (Angal, S. 1993 Mol Immunol, 30:105-108; Aalberse et al. 2002 Immunol, 105:9-19). Some of the amino acid residues in the hinge and γ Fc region were reported to have impact on antibody interaction with Fc γ receptors (Chappel SM, et al. 1991 Proc. Natl. Acad. Sci. USA, 88:9036-9040; Mukherjee, J. et al., 1995 FASEB J, 9:115-119; Armour, K.L. et al., 1999 Eur J Immunol, 29:2613-2624; Clynes, R.A. et al., 2000 Nature Medicine, 6:443-446; Arnold J.N., 2007 Annu Rev Immunol, 25:21-50). Furthermore, some rarely occurring IgG4 isoforms in human population may also elicit different physicochemical properties (Brusco, A. et al. 1998 Eur J Immunogenet, 25:349-55; Aalberse et al. 2002 Immunol, 105:9-19). However, lumping all the mutations and isoforms previously discovered into a specific antibody does not warrant for an ideal antibody molecule to share all the features for therapeutics such as described above, which may be resulted from contradictory effect of the combined mutations and from impact of variable region to the effector function and physicochemical properties of an antibody (Igawa T. et al., 2010 Prot Eng Design Select, 23:385-392; Perchiacca J.M. and Tessier P.M., 2012 Ann Rev Biomol Eng 3:263-286).

[0085] To generate anti-PD-1 mAbs with least ADCC, CDC and instability, we modified the hinge and γ Fc region of human IgG4 by introduce a number of combinations of mutations, which created IgG4mt1 to IgG4mt12. Some of the modified IgG4 variants were clearly less desirable as indicated by our assay results, several relevant IgG4 variants and modified sequences were listed in Table 19. The assessment of these antibodies is described herein.

Table 19. Sequence modifications of IgG4 variants

IgG4 and variants	Amino acid residues*																
	...	228	229	230	231	232	233	234	235	236	...	265	...	309	...	409	...
IgG4	...	S	C	P	A	P	E	F	L	G	...	D	...	L	...	R	...
IgG4mt1	...	<u>P</u>	C	P	A	P	E	F	L	G	...	D	...	L	...	R	...
IgG4mt2	...	<u>P</u>	C	P	A	P	<u>P</u>	<u>V</u>	<u>A</u>	G	...	D	...	L	...	R	...

(continued)

IgG4 and variants	Amino acid residues*																
	...	228	229	230	231	232	233	234	235	236	...	265	...	309	...	409	...
IgG4mt6	...	<u>P</u>	C	P	A	P	<u>P</u>	<u>V</u>	<u>A</u>	G	...	<u>A</u>	...	L	...	R	...
IgG4mt8	...	<u>P</u>	C	P	A	P	<u>P</u>	<u>V</u>	<u>A</u>	G	...	<u>T</u>	...	L	...	R	...
IgG4mt9	...	<u>P</u>	C	P	A	P	<u>P</u>	<u>V</u>	<u>A</u>	G	...	<u>A</u>	...	L	...	<u>K</u>	...
IgG4mt10	...	<u>P</u>	C	P	A	P	<u>P</u>	<u>V</u>	<u>A</u>	G	...	<u>A</u>	...	<u>V</u>	...	<u>K</u>	...

* Amino acid numbering is based on EU system. Changes are highlighted by underline.

Example 9. IgG4mt10 has no Fc γ R binding, lowest ADCC and CDC effector function

[0086] ADCC is initiated when an antibody binds to cell surface target protein followed by ligation to Fc γ receptors (Fc γ Rs) expressed on effector cells. It was well documented that human IgG1 has significantly higher binding affinity to Fc γ Rs than IgG2 and IgG4, specially, binding to Fc γ R-I and Fc γ R-IIIA, which correlated to the strength of IgG1 to activate ADCC. Reminiscent of ADCC, CDC is activated when an antibody cross-links a cell surface target and C1q protein, which followed by a cascade reaction of complement complex formation and target cell lysis. As proxy of ADCC and CDC, assays for antibody binding to Fc γ Rs and C1q may serve as the fundamental indicator of ADCC and CDC. We therefore systematically assessed the mAbs binding to all the major Fc γ Rs.

Fc γ R binding

[0087] Binding of various IgG4 mutants to Fc γ Rs was determined by flow cytometry. In brief, a series of HEK293 transfectants expressing human Fc γ Rs were established. These transfectants expressed Fc γ RI, Fc γ RIIA, Fc γ RIIB or Fc γ RIIIA. Multi-subunit Fc γ Rs (i.e., Fc γ RI and Fc γ RIIIA) were co-expressed with FcR γ . Polymorphic variants (i.e., Fc γ RIIA H131 and R131, Fc γ RIIIA F158 and V158) were also included. A secondary antibody (goat anti-human IgG F(ab)'2-Alexa Fluor 488, Jackson ImmunoResearch, West Grove, PA, USA) was used to detect the binding of anti-PD-1 mAbs with modified IgG4 variants (Table 19) to Fc γ R⁺ HEK293 cells. As expected, anti-PD-1 mAbs in IgG1 format (hu317-1/IgG1 and hu317-4B6/IgG1) bind strongly to all Fc γ Rs including Fc γ RI, Fc γ RIIA (H131 and R131 alleles), Fc γ RIIB, and Fc γ RIIIA (V158 and F158 alleles) (Table 20). Interestingly, when the two different version of humanization mAbs, hu317-1 and hu317-4B6 (with differences in both V_h and V_k), were generated in the same IgG4 variant format, such as either in IgG4mt1 or in IgG4mt6 format, their binding strength (MFI) vary by an range from a couple fold to close to 100 fold (e.g. 455.2/115.7 = 3.9 fold; 13.6/1.0 = 13.6 fold; 434.6/4.9 = 88.7 fold; and etc., see Table 20). It is consistent to the previous findings by other that the variable regions of antibodies do have significant impact on the binding to FcRs, therefore, exerting the impact on effector function such as ADCC (Igawa T. et al., 2010 Prot Eng Design Select, 23:385-392; Perchiacca J.M. and Tessier P.M., 2012 Ann Rev Biomol Eng 3:263-286).

[0088] As demonstrated in Table 20, when hu317-4B6 and hu326-4A3 were made in IgG4mt10 format, they have the lowest binding activity to Fc γ Rs among the PD-1 mAbs and IgG variant formats listed in the table, as well as many other humanization mAbs and IgG formats we have tested in the study. The uniqueness of hu317-4B6 and hu326-4A3 in IgG4mt10 format in this regard may not be extended to the same family of humanization mAbs with somewhat distant sequence homology, such as hu317-1, as described above.

Table 20. Binding strength (MFI*) of anti-PD-1 mAbs to Fc γ Rs determined by FACS

mAbs	Fc γ RI	Fc γ RIIA (H131)	Fc γ RIIA (R131)	Fc γ RIIB	Fc γ RIIIA (F158)	Fc γ RIIIA (V158)
hu317-1 /IgG1	2152.9	168.7	139.6	442.4	99.7	277.2
hu317-4B6 /IgG1	2771.7	1.7	0.6	1.9	28.0	293.7
hu317-1 /IgG4mt1	455.2	21.3	21.9	434.6	0.6	20.7
hu317-4B6 /IgG4mt1	115.7	0.2	0.0	4.9	0	6.1
hu317-1 /IgG4mt6	13.6	1.0	0.8	1.8	0.9	1.1

(continued)

mAbs	Fc γ RI	Fc γ RIIA (H131)	Fc γ RIIA (R131)	Fc γ RIIB	Fc γ RIIIA (F158)	Fc γ RIIIA (V158)
hu317-4B6 /IgG4mt6	1.0	0	0	0	0	0
hu317-4B6 /IgG4mt10	0.4	0	0	0	0	0
hu326-4A3 /IgG4mt10	0.5	0	0	0	0	0
* MFI: mean fluorescence intensity from FACS analysis						

ADCC

[0089] Classical ADCC involves activation of NK cells by antibodies engaging to Fc γ RIIIA or CD16. To verify whether humanized anti-PD-1 mAbs induce ADCC, NK92MI/CD16V cells, which were generated from NK92MI cells (ATCC) by co-transducing expression plasmids containing *CD16* (V158 allele) and *FcR γ* genes, were used as effector cells, and PD-1-expressing T cell line, HuT78/PD-1, was used as target cells. NK92MI/CD16V cells (4×10^4) were co-cultured with equal number of HuT78/PD-1 cells in 96-well V-bottom plates for 5h. Cytotoxicity was determined by LDH release assay described in previous section. The results confirmed that hu317-4B2/IgG4mt6, hu317-4B6/IgG4mt6, hu317-4B6/IgG4mt10 and hu326-4A3/IgG4mt10 all have base level of ADCC comparing to the positive controls (Fig. 7). The minor difference in ADCC between those 4 mAbs may be attributable to experimental error (see error bars in Fig. 7).

CDC

[0090] Human IgG4 antibodies, in general, do not induce any CDC via classical pathway. Whether anti-PD-1 mAbs in IgG4mt10 format will trigger CDC was evaluated using a PD-1-expressing T cell line, Hut78/PD-1, and fresh human serum from healthy donors. Cell lysis by CDC was determined by Celltiter glo assay kits (Promega, Beijing, China). In brief, HuT78/PD-1 cells (2×10^4) were incubated in serum-free RPMI1640 (Invitrogen) with anti-PD-1 Abs (10 μ g/ml) at 37°C for 15 minutes before adding normal human serum (NHS) to the final concentration of 15% or 50% in 96-well flat-bottom plates in a total volume of 120 μ l. After overnight incubation at 37°C, cells were lysed and assayed for ATP concentration. To test whether humanized anti-PD-1 mAbs in IgG4mt10 can kill PD-1⁺ primary T cells via CDC, PBMCs isolated from healthy donors were pre-activated with anti-CD3 Ab OKT3 (40 ng/ml) for 3 days before co-culture with anti-PD-1 Abs plus NHS. The amount of ATP is directly proportional to the number of cells present in culture. Fluorescence was read using a 96-well fluorometer (PHERA Star FS, BMG LABTECH). The results are expressed in relative fluorescence units (RFU) that are proportional to the number of viable cells. The percent CDC activity was calculated as follows: % CDC activity = [(RFU test - RFU background) / (RFU at total cell lysis - RFU background)] \times 100. In general, we were not able to detect any ADCC mediated by anti-PD-1 mAbs in IgG4mt10 format that bind to activated PBMCs. In hypersensitive experimental conditions, such as using PD-1 highly-expressing cell line, high serum and antibody concentration, we detected very low level of CDC in some occasions, and there is not much differences between different versions and anti-PD-1 mAbs, indicating the anti-PD-1 mAbs in IgG4 variant formats retained the feature of low or no CDC activity as the common form of IgG4.

Example 10. Humanized anti-PD-1 mAbs in IgG4mt10 format have enhanced stability under stress conditions*Stability of anti-PD-1 antibodies in high temperature and acidic conditions*

[0091] Anti-PD-1 antibodies used in stability studies were all purified from protein A column followed by size exclusion chromatography (SEC) as described in previous sections. Following purification, the aggregate contents of purified antibody samples were monitored in analytical size exclusion chromatography-high performance liquid chromatography (SEC-HPLC), which fell within the range of 0%-0.5%.

[0092] For SEC-HPLC analysis, the antibody samples were analyzed using a TSKgel G3000 SWXL column (7.8x300 mm, Cat. No. 08541, Tosoh Bioscience, Shanghai, China) under isocratic elution condition (elution buffer 0.2 M sodium phosphate, pH7.2), and subsequent detection at UV-215 nm. In each run, 10 microliters of antibody sample was loaded onto the column and eluted at a flow rate of 1mL/minute. The dimer or larger aggregate species of antibody were separated from monomeric species and the percentages of dimers and aggregates were determined based on the integrated peak areas from UV traces.

[0093] For speed-enhanced shelf stability study, anti-PD-1 antibodies (10-40mg/mL in PBS) were kept in incubators at 40-50°C for 4-7 days in order to test the stability of antibodies in high temperature condition. The antibody samples were then analyzed for heat-induced formation of dimer and aggregates in SEC-HPLC. For each of the anti-PD-1 antibodies analyzed, less than 2% became higher molecular weight species (dimers and aggregates), indicating the anti-PD-1 antibodies had good stability in high temperature conditions.

[0094] Antibody's stability in acidic condition has been a key challenge in the downstream manufacturing process (Liu et al. 2010 mAbs 2:480-499). Antibody elution from protein A and inactivation of virus usually require incubation of antibody in low pH (2.5-4) conditions. However, such acidic conditions could potentially cause antibody denaturation and aggregation. Human IgG4 has been known to be less stable than IgG1 and IgG2 (2002 Immunology 105:9). Therefore, we assayed the humanized mAbs made with various IgG4 mutant forms. Briefly, Antibody stabilities in low pH conditions were studied by 1:1 volume of each antibody sample (10 mg/mL in PBS) mixed with low pH buffers containing 50 mM sodium citrate, 100mM NaCl at pH3.6, 3.3, 3.0 or 2.7, respectively. After 1 hour incubation at room temperature, the antibody samples in low pH conditions were neutralized by 1:5 dilution into SEC-HPLC elution buffer containing 0.2M sodium phosphate, pH7.2. SEC-HPLC analyses were done as described above and percentages of dimers and aggregates induced by low pH conditions were quantified. The anti-PD-1 mAb 317-4B6 in IgG1 format was most stable in bioprocessing-relevant acidic conditions even when pH value get as low as 2.7. Among the anti-PD-1 mAbs made in several IgG4 variants, hu317-4B6/IgG4mt10 and hu326-4A3/IgG4mt10 were the most stable under the acidic buffer condition (Table 21) as the acid-induced aggregates were significantly reduced to a level that was comparable to that of the IgG1 format of anti-PD-1 mAbs, 317-4B6 and 326-4A3, i.e. the soluble aggregate is less than 2% (Table 21).

Table 21. Dimer and soluble aggregates formed in acidic buffers and assayed by SEC-HPLC

anti-PD-1 mAbs	% of dimer and aggregates				
	pH7.2	pH3.6	pH3.3	pH3.0	pH2.7
317-4B6/IgG1	0.0%	0.0%	0.2%	0.1%	0.2%
317-4B6/IgG4mt1	0.0%	1.0%	11.0%	49.0%	48.0%
317-4B6/IgG4mt3	0.0%	13.0%	31.0%	>50%	>50%
317-4B6/IgG4mt6	0.0%	4.0%	41.0%	>50%	>50%
317-4B6/IgG4mt9	0.0%	0.5%	2.1%	3.3%	2.0%
317-4B6/IgG4mt10	0.0%	0.2%	0.6%	0.6%	1.4%
326-4A3/IgG4mt10	0.0%	0.0%	0.4%	0.5%	1.2%

Example 11. Mapping the binding epitopes of anti-PD-1 mAbs

[0095] Previous reports about the crystal structures of PD-1/PD-L1 and PD-1/PD-L2 complexes had shed light to understanding critical amino acid (AA) residues on PD-1 which are required for the ligand-binding (Zhang et al. 2004 Immunity, 20:337-347; Lin D.Y. et al. 2008 PNAS 105:3011-3016; Lazar-Molnar E. et al. 2008 PNAS, 105:10483-10488). In fact, six of such AA residues were identified on the receptor through point mutation analysis required for PD-L1 binding. Five of the six AA residues were also required for PD-L2 binding (Lin D.Y. et al. 2008 PNAS 105:3011-3016). Based on the information from the structure-guided mutation analysis we hypothesized that most effective way for functional mAbs to block PD-1 mediated signaling is to compete with PD-1 ligands by binding to the six critical AA residues, therefore, occupying the binding epitopes required for the ligand binding. To explore the hypothesis and to understand the mechanism of action by functional PD-1 antibodies, we have made six mutants of PD-1 by replacing each of the six critical AAs to Ala, individually, i.e. K45A, I93A, L95A, P97A, I101A and E103A (AA residue numbering based on Lin D.Y. et al. 2008 PNAS 105:3011-3016). The mutant PD-1/Fc and PD-1/His (Fig. 1) were used as templates for PCR-guided mutagenesis or rolling-circle mutagenesis using Fast Mutagenesis System (Cat. No. FM111, Transgen Biotech, Beijing, China). All mutants were sub-cloned in our pcDNA-based expression vectors, and verified by sequencing. The mutated and wild-type PD-1 proteins were expressed by transient transfection (described in Example 1), and prepared after 4 to 6 days of culture. The conditioned media (CM) were analyzed by Western blot to verify the PD-1 protein expression in terms of quality and quantity. The supernatants (CM), after clearing cell debris, were directly used in ELISA analysis or Western blot for epitope-mapping.

[0096] To study the binding epitopes of humanized anti-PD-1 mAbs, ELISA assays using the wild-type (WT) and mutant (Mt) PD-1 were performed to assess the binding activities of hu317-4B5, hu317-4B6, hu326-3B1 and hu326-4A3. For comparison to check the uniqueness of the antibody binding signature, two reference antibody (Reference Ab-1 and

Reference Ab-2 from US8008449B2 and US8168757B2, respectively) were included in the study. Equal volume of CM containing WT or Mt PD-1 was coated in 96-well plate for all mAbs in the same ELISA assay. All ELISA results were normalized using the mean ELISA readings of WT PD-1 binding signals as the standard. ELISA binding signals to a specific Mt PD-1 were further normalized against the highest antibody binding read-out (set as 100%) to the specific Mt PD-1. For convenience of data analysis, when a mAb's ELISA binding signal for a specific mutant dropped below 50% relative to WT PD-1, it is defined that the amino acid residue is a significant binding epitope because whose mutation significantly abrogated the antibody binding. Likewise, if a mAb's ELISA binding signal for a specific mutant dropped below 25%, it is defined as very significant. As shown in Fig. 8, two of the critical AA residues in PD-1, K45 and I93, are significant or very significant epitopes for mAbs hu317-4B5 and hu317-4B6 binding, and three AA residues, I93, L95 and P97, are either significant or very significant epitopes for hu326-3B1 and hu326-4A3. On the other hand, the two reference antibodies have distinctive binding epitopes, P97 is significant for Reference Ab-1, while L95 and P97 are significant for Reference Ab-2.

[0097] Interestingly, when the PD-1 protein is denatured in Western Blot, mAb hu317-4B5 and -4B6 were still capable of binding to WT PD-1 though the critical binding epitopes (K45 and I93) are not close to each other (non-linear). It indicated that the PD-1 protein became renatured to some degree after denaturation in SDS-PAGE of Western Blot process, which allows the anti-PD-1 mAbs to recognize and bind to it. Taking the advantage of this observation, we performed Western Blot analysis for all six antibodies used in above ELISA study. The overall results from Western Blot corroborated very well to the ELISA results, i.e. the significant or very significant epitopes, whose mutations resulted in low binding signals in ELISA, also gave weakest Western Blot band comparing to the binding to other mutant PD-1 (Fig. 8). Some minor differences between ELISA and Western Blot were also observed, e.g., the ELISA binding signals on I93A and E103A by reference Ab-2 were relatively stronger than those in Western Blot. It may be indicative of that those AA residues may also contribute to the binding because whose mutations impacted the binding though only under stress condition (i.e. denaturation or losing native conformation). As summarized in Table 22, the anti-PD-1 mAbs in this invention have identifiable binding epitopes differing from other anti-PD-1 antibody.

Table 22. Summary* of key epitopes by anti-PD-1 mAbs

	K45A	I93A	L95A	P97A	I101A	E103A
hu317-4B5	***	**				
hu317-4B6	***	**				
hu326-3B1		**	**	**		
hu326-4A3		***	**	**		
Ref. Ab-1				**		
Ref. Ab-2			**	**		
* based on Fig.8						

Example 12. Anti-PD-1 mAbs activate primary human PBMCs and inhibit tumor growth in xenograft mouse models

Humanized anti-PD-1 mAbs activate human PBMCs

[0098] Throughout the humanization processes, the humanized anti-PD-1 mAbs at various stages retained similar functional activities as assessed by ELISA, FACS and immune cell-based cytokine release assays. To confirm the function of final versions of humanized mAbs, we assayed the activating functions of hu317-4B5, hu317-4B6, hu326-3B1 and hu326-4A3 using primary human PBMCs. The results demonstrated that those mAbs throughout the humanization maintained the original murine mAb functions to activate primary PBMCs although the degree of activation differs among the four donors due to the variance of individual's genetic background (Fig. 9).

Humanized anti-PD-1 mAbs enhance NK cell-based cytotoxicity against cancer cells

[0099] Reminiscent of the original murine mAbs, the humanized anti-PD-1 mAbs, hu317-4B5, hu317-4B6, hu326-3B1 and hu326-3G1, enhance NK92MI/PD-1 cell-mediated cytotoxicity against the target lung cancer cells, SK-MES-1/PD-L1, in a dose-dependent manner (Fig. 10, Table 23). It appeared evident that in principle the humanized anti-PD-1 mAbs might function to break immune cell tolerance mediated by PD-1 signaling, enhancing the cancer killing activity by immune cells, e.g. NK cells and cytotoxic T-lymphocytes.

Humanized anti-PD-1 mAb activates human PBMCs and inhibits tumor growth in a mouse xenograft cancer model in vivo

[0100] All above experimental evidence indicated that the anti-PD-1 mAbs might work in mouse cancer models utilizing immune-compromised mice xenografted with human cancer cells, subsequently implanting human PBMCs and applying the mAb treatment to inhibit cancer cell growth *in vivo*. The experiment was designed as follows. Seven-eight week old SCID-male mice (Vital River Laboratories, China) were inoculated subcutaneously at right flank with 3×10^6 Hep3B/OS8-PD-L1 cells in 50% Matrigel (BD Biosciences, New Jersey, USA). Fifteen days after tumor inoculation, the mice bearing tumor size between 100-250 mm³ were randomized and divided into three treatment groups. One hundred microliters of pooled PBMCs (5×10^5) from 2 healthy donors were injected intratumorally. Three days post PBMC-implanting, anti-PD-1 antibodies (Hu317-IgG4mt2) and human IgG were administered via s.c. at a dose of 10 mg/kg, respectively. Antibody treatment was repeated once every 10 days for a total of 3 times. PBS was injected in a parallel group as negative control. Tumors were measured twice a week using a caliper starting on day 7. Tumor volumes were calculated using the following formula: $[D \times (d^2)]/2$, in which D represents the large diameter of the tumor, and d represents the small diameter. All animal studies were performed following Beigene Animal Care and Use Procedure.

[0101] In the *in vivo* study, although 60% of tumors in the control groups were auto-regressed, the rest of *in vivo* experiment is still quite informative, which were presented in Fig. 11. In the control groups, either vehicle-treated or human IgG (hulgG)-treated group, each has 40% tumors (2 of 5 mice) outgrowing larger than the baseline at starting point. The two tumors in PBS-treated group grew much larger (above 2,000 mm³, one tumor-bearing mouse was terminated earlier due to passing tumor size limit by protocol). The two tumors in hulgG-treated group grew to the size of 800 and 1,370 mm³, significantly above the average baseline of 164 mm³ though smaller than the PBS-treated tumors. On the other hand, in the anti-PD-1 mAb (hu317-1/IgG4mt2)-treated group, tumors were either completely regressed or close to baseline size (one tumor = 200 mm³, which grew back slowly after regressed to 50% of baseline at two weeks from PBMC implanting). The results indicated that the anti-PD-1 mAb described above can activate human immune cells inhibiting tumor cells growth in the mouse *in vivo* cancer model, which is consistent to the *in vitro* experimental results described above.

SEQUENCE LISTING

[0102]

<110> BeiGene Li, Kang Zhang, Tong Song, Jing Xu, Lanlan Liu, Qi Peng, Hao

<120> ANTI-PD1 ANTIBODIES AND THEIR USE AS THERAPEUTICS AND DIAGNOSIS

<130> BGB 108

<160> 90

<170> PatentIn version 3.5

<210> 1

<211> 444

<212> DNA

<213> Homo sapiens

<400> 1

EP 3 702 373 B9

```

ccaggatggt tcttagactc cccagacagg ccctggaacc cccccacctt ctccccagcc      60
ctgctcgtgg tgaccgaagg ggacaacgcc accttcacct gcagcttctc caacacatcg      120
5 gagagcttcg tgctaaactg gtaccgcatg agccccagca accagacgga caagctggcc      180
gccttccccg aggaccgcag ccagcccggc caggactgcc gcttccgtgt cacacaactg      240
ccaacggggc gtgacttcca catgagcgtg gtcagggccc ggcgcaatga cagcggcacc      300
10 tacctctgtg gggccatctc cctggccccc aaggcgcaga tcaaagagag cctgcgggca      360
gagctcaggg tgacagagag aagggcagaa gtgcccacag cccaccccag cccctcaccc      420
15 aggccagccg gccagttcca aacc                                             444

```

<210> 2

<211> 148

<212> PRT

20 <213> Homo sapiens

<400> 2

```

25   Pro Gly Trp Phe Leu Asp Ser Pro Asp Arg Pro Trp Asn Pro Pro Thr
     1           5           10           15
30   Phe Ser Pro Ala Leu Leu Val Val Thr Glu Gly Asp Asn Ala Thr Phe
           20           25           30
35   Thr Cys Ser Phe Ser Asn Thr Ser Glu Ser Phe Val Leu Asn Trp Tyr
           35           40           45
40   Arg Met Ser Pro Ser Asn Gln Thr Asp Lys Leu Ala Ala Phe Pro Glu
           50           55           60

```

EP 3 702 373 B9

Asp Arg Ser Gln Pro Gly Gln Asp Cys Arg Phe Arg Val Thr Gln Leu
 65 70 75 80
 5 Pro Asn Gly Arg Asp Phe His Met Ser Val Val Arg Ala Arg Arg Asn
 85 90 95
 10 Asp Ser Gly Thr Tyr Leu Cys Gly Ala Ile Ser Leu Ala Pro Lys Ala
 100 105 110
 15 Gln Ile Lys Glu Ser Leu Arg Ala Glu Leu Arg Val Thr Glu Arg Arg
 115 120 125
 Ala Glu Val Pro Thr Ala His Pro Ser Pro Ser Pro Arg Pro Ala Gly
 130 135 140
 20 Gln Phe Gln Thr
 145

<210> 3
 <211> 354
 <212> DNA
 <213> Mus musculus

<400> 3
 30 caggtgcagc tgaaggagtc aggacctggc ctggtggcgc cctcaaagaa cctgtccatc 60
 acttgcaactg tctctgggtt ttcattaacc agctatgggtg tacactggat tcgccagcct 120
 35 ccaggaaagg gactggaatg gctgggagta atatgggccg gtggaagcac aaattataat 180
 tcggctctca tgtccagact gagcatcagc aaagacaact ccaggagcca agttttctta 240
 agaatgaaca gtctgcaaac tgatgacaca gccatgtact actgtgccag agcctatggt 300
 40 aactactggt acatcgatgt ctggggcgca gggaccacgg tcaccgtctc ctca 354

<210> 4
 <211> 118
 <212> PRT
 <213> Mus musculus

<400> 4
 50 Gln Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Lys
 1 5 10 15
 Asn Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30
 55 Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu
 35 40 45

EP 3 702 373 B9

Gly Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Ser Ala Leu Met
 50 55 60
 5
 Ser Arg Leu Ser Ile Ser Lys Asp Asn Ser Arg Ser Gln Val Phe Leu
 65 70 75 80
 10
 Arg Met Asn Ser Leu Gln Thr Asp Asp Thr Ala Met Tyr Tyr Cys Ala
 85 90 95
 15
 Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Ala Gly Thr
 100 105 110
 20
 Thr Val Thr Val Ser Ser
 115

20 <210> 5
 <211> 321
 <212> DNA
 <213> Mus musculus

25 <400> 5

30 gacattgtga tgaccagac tcccaaattc ctgcttgat cagcaggaga cagggttacc 60
 ataacctgca aggccagtca gagtgtgagt aatgatgtag cttggtacca acagaagcca 120
 gggcagtctc ctaaactgct gataaactat gcatttcacg gcttcactgg agtccctgat 180
 cgtttcactg gcagtggata tgggacggat ttcattttca ccatcagcac tgtgcaggct 240
 35 gaagacctgg cagtttattt ctgtcaccag gcttatagtt ctccgtacac gttcggaggg 300
 gggaccaagc tggaaatgaa a 321

40 <210> 6
 <211> 107
 <212> PRT
 <213> Mus musculus

45 <400> 6

50

55

EP 3 702 373 B9

Asp Ile Val Met Thr Gln Thr Pro Lys Phe Leu Leu Val Ser Ala Gly
 1 5 10 15
 5 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Ser Val Ser Asn Asp
 20 25 30
 10 Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
 35 40 45
 15 Asn Tyr Ala Phe His Arg Phe Thr Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60
 20 Ser Gly Tyr Gly Thr Asp Phe Ile Phe Thr Ile Ser Thr Val Gln Ala
 65 70 75 80
 25 Glu Asp Leu Ala Val Tyr Phe Cys His Gln Ala Tyr Ser Ser Pro Tyr
 85 90 95
 30 Thr Phe Gly Gly Gly Thr Lys Leu Glu Met Lys
 100 105

<210> 7
 <211> 342
 <212> DNA
 <213> Mus musculus

<400> 7

35 cagatccagt tgggtgcagtc tggacctgag ctgaagaagc ctggagagac agtcaagatc 60
 tcctgcaagg cttctgggta taccttcaca aactatggaa tgaactgggt gaagcaggct 120
 40 ccaggaaagg gtttaaagtg gatgggctgg ataaacaata ataatggaga gccaacatat 180
 gctgaagagt tcaagggacg gtttgccttc tctttggaaa cctctgccag cactgcctat 240
 ttgcagatca acaacctcaa aatgaggac acggctacat atttctgtgc aagagatggt 300
 45 atggactatt ggggtcaagg aacctcagtc accgtctcct ca 342

<210> 8
 <211> 114
 <212> PRT
 <213> Mus musculus

<400> 8

55

EP 3 702 373 B9

1 Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
 5 Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 10 Gly Met Asn Trp Val Lys Gln Ala Pro Gly Lys Gly Leu Lys Trp Met
 15 Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Glu Glu Phe
 20 Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
 25 Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys
 30 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Ser Val Thr Val
 35

Ser Ser

30 <210> 9
 <211> 330
 <212> DNA
 <213> Mus musculus

35 <400> 9

gacattgtgc tgaccaatc tccagcttct ttggctgtgt ctctagggca gagggccacc 60
 40 atacctgca gagccagtga aagtgtgat aattatggct atagttttat gcactggtac 120
 cagcagaaac caggacagcc accccaactc ctcatctatc gtgcatccaa cctagaatct 180
 gggatccctg ccaggttcag tggcagtggg tctaggacag gcttcaccct caccattaat 240
 45 cctgtggagg ctgatgatgt tgcaacctat tactgtcagc aaagtaaaga atatccgacg 300
 ttcggtggag gcaccaagct ggaagtcaaa 330

50 <210> 10
 <211> 110
 <212> PRT
 <213> Mus musculus

55 <400> 10

EP 3 702 373 B9

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 5 Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Asn Tyr
 20 25 30
 10 Gly Tyr Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 15 Gln Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala
 50 55 60
 20 Arg Phe Ser Gly Ser Gly Ser Arg Thr Gly Phe Thr Leu Thr Ile Asn
 65 70 75 80
 25 Pro Val Glu Ala Asp Asp Val Ala Thr Tyr Tyr Cys Gln Gln Ser Lys
 85 90 95
 30 Glu Tyr Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Val Lys
 100 105 110

<210> 11
 <211> 10
 <212> PRT
 <213> Mus musculus

<400> 11

Gly Phe Ser Leu Thr Ser Tyr Gly Val His
 1 5 10

<210> 12
 <211> 16
 <212> PRT
 <213> Mus musculus

<400> 12

Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Ser Ala Leu Met Ser
 1 5 10 15

<210> 13
 <211> 12
 <212> PRT
 <213> Mus musculus

<400> 13

Ala Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val
 1 5 10

EP 3 702 373 B9

5
<210> 19
<211> 7
<212> PRT
<213> Mus musculus

<400> 19

10
Ala Arg Asp Val Met Asp Tyr
1 5

15
<210> 20
<211> 15
<212> PRT
<213> Mus musculus

<400> 20

20
Arg Ala Ser Glu Ser Val Asp Asn Tyr Gly Tyr Ser Phe Met His
1 5 10 15

25
<210> 21
<211> 7
<212> PRT
<213> Mus musculus

<400> 21

30
Arg Ala Ser Asn Leu Glu Ser
1 5

35
<210> 22
<211> 8
<212> PRT
<213> Mus musculus

<400> 22

40
Gln Gln Ser Lys Glu Tyr Pro Thr
1 5

45
<210> 23
<211> 354
<212> DNA
<213> Artificial Sequence

50
<220>
<223> 317-4B6 cDNA-Vh

<400> 23

55

EP 3 702 373 B9

caggtgcagc tgcaggagtc gggaccagga ctggtgaagc cttcggagac cctgtccctc 60
 acctgcactg tctctgggtt ttcattaacc agctatggtg tacactggat ccggcagccc 120
 5 ccaggaag gactggagtg gatcggggtc atatacgccg atggaagcac aaattataat 180
 ccctccctca agagtcgagt gaccatatca aaagacacct ccaagaacca ggtttccctg 240
 aagctgagct ctgtgaccgc tgcggacacg gccgtgtatt actgtgcgag agcctatggt 300
 10 aactactggt acatcgatgt ctggggccaa gggaccacgg tcaccgtctc ctca 354

<210> 24
 <211> 118
 15 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 317-4B6 pro-Vh
 20
 <400> 24

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15
 25

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30
 30

Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45
 35

Gly Val Ile Tyr Ala Asp Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60
 40

Ser Arg Val Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val Ser Leu
 65 70 75 80
 45

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95
 50

Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
 100 105 110
 55

Thr Val Thr Val Ser Ser
 115

<210> 25
 <211> 321
 <212> DNA
 55 <213> Artificial Sequence

<220>
 <223> 317-4B6 cDNA-Vk

EP 3 702 373 B9

<400> 25

gacatcgtga tgaccagtc tccagactcc ctggctgtgt ctctgggcga gagggccacc 60
 5 atcaactgca agtccagcga gagtgtgagt aatgatgtag cttggtacca gcagaaacca 120
 ggacagcctc ctaagctgct cattaactat gcatttcacg gcttcactgg ggtccctgac 180
 10 cgattcagtg gcagcgggta tgggacagat ttcactctca ccatcagcag cctgcaggct 240
 gaagatgtgg cagtttatta ctgtcaccag gcttatagtt ctccgtacac gtttggccag 300
 gggaccaagc tggagatcaa a 321

15 <210> 26
 <211> 107
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> 317-4B6 pro-Vk

<400> 26

25 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 30 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Ser Asn Asp
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
 35 35 40 45
 40 Asn Tyr Ala Phe His Arg Phe Thr Gly Val Pro Asp Arg Phe Ser Gly
 50 55 60
 Ser Gly Tyr Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala
 65 70 75 80
 45 Glu Asp Val Ala Val Tyr Tyr Cys His Gln Ala Tyr Ser Ser Pro Tyr
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105

50 <210> 27
 <211> 342
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> 326-4A3 cDNA-Vh

EP 3 702 373 B9

<400> 27

5 caggtgcagc tgggtgcagag cggcagcgag ctgaagaagc ccggcgccag cgtgaaggtg 60
 agctgcaagg ccagcggcta caccttcacc aactacggca tgaactgggt gagacaggcc 120
 cccggccagg gcctgaagtg gatgggctgg atcaacaaca acaacgccga gccacctac 180
 10 gccaggact tcagaggcag attcgtgttc agcctggaca ccagcgccag caccgcctac 240
 ctgcagatca gcagcctgaa gaccgaggac accgccgtgt actactgcbc cagagacgtg 300
 atggactact ggggccaggg caccctggtg accgtgagca gc 342

15 <210> 28
 <211> 114
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> 326-4A3 pro-Vh

<400> 28

25 Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
 1 5 10 15
 30 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 25 30
 35 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
 35 40 45
 40 Gly Trp Ile Asn Asn Asn Asn Ala Glu Pro Thr Tyr Ala Gln Asp Phe
 50 55 60
 45 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 50 Leu Gln Ile Ser Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
 100 105 110

Ser Ser

55 <210> 29
 <211> 330
 <212> DNA
 <213> Artificial Sequence

EP 3 702 373 B9

<220>

<223> 326-4A3 cDNA-Vk

<400> 29

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

```

gacattgtgc tgaccagtc tccagcctcc ttggccgtgt ctccaggaca gagggccacc      60
atcacctgca gagccagtga aagtgttgat aattatggct atagttttat gacttggtat      120
cagcagaaac caggacaacc tcctaaactc ctgatttacc gtgcatccaa cctagaatct      180
ggggtcccag ccaggttcag cggcagtggg tctgggaccg atttcaccct cacaattaat      240
cctgtggaag ctgaggatac tgcaaattat tactgtcagc aaagtaaaga atatccgacg      300
ttcggcggag ggaccaaggt ggagatcaaa      330
  
```

<210> 30

<211> 110

<212> PRT

<213> Artificial Sequence

<220>

<223> 326-4A3 pro-Vk

<400> 30

30
 35
 40
 45
 50
 55

```

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Gly
 1           5           10           15

Gln Arg Ala Thr Ile Thr Cys Arg Ala Ser Glu Ser Val Asp Asn Tyr
          20           25           30

Gly Tyr Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
          35           40           45

Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
          50           55           60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn
65           70           75           80

Pro Val Glu Ala Glu Asp Thr Ala Asn Tyr Tyr Cys Gln Gln Ser Lys
          85           90           95

Glu Tyr Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys

          100           105           110
  
```

<210> 31

<211> 10

<212> PRT

EP 3 702 373 B9

<213> Mus musculus

<400> 31

5

Gly Phe Ser Leu Thr Ser Tyr Gly Val His
1 5 10

<210> 32

10

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

15

<223> 317-4B6 H-CDR2 or CDR-H2

<400> 32

20

Val Ile Tyr Ala Asp Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser
1 5 10 15

<210> 33

25

<211> 12

<212> PRT

<213> Mus musculus

<400> 33

30

Ala Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val
1 5 10

<210> 34

35

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

40

<223> 317-4B6 L-CDR1 or CDR-L1

<400> 34

45

Lys Ser Ser Glu Ser Val Ser Asn Asp Val Ala
1 5 10

<210> 35

50

<211> 7

<212> PRT

<213> Mus musculus

<400> 35

55

Tyr Ala Phe His Arg Phe Thr
1 5

EP 3 702 373 B9

<210> 36
<211> 9
<212> PRT
<213> Mus musculus

5

<400> 36

His Gln Ala Tyr Ser Ser Pro Tyr Thr
1 5

10

<210> 37
<211> 10
<212> PRT
<213> Mus musculus

15

<400> 37

Gly Tyr Thr Phe Thr Asn Tyr Gly Met Asn
1 5 10

20

<210> 38
<211> 17
<212> PRT
<213> Artificial Sequence

25

<220>
<223> 326-4A3 H-CDR2 or CDR-H2

30

<400> 38

Trp Ile Asn Asn Asn Asn Ala Glu Pro Thr Tyr Ala Gln Asp Phe Arg
1 5 10 15

35

Gly

<210> 39
<211> 7
<212> PRT
<213> Mus musculus

40

<400> 39

45

Ala Arg Asp Val Met Asp Tyr
1 5

50

<210> 40
<211> 15
<212> PRT
<213> Mus musculus

55

<400> 40

EP 3 702 373 B9

Arg Ala Ser Glu Ser Val Asp Asn Tyr Gly Tyr Ser Phe Met His
1 5 10 15

5 <210> 41
<211> 7
<212> PRT
<213> Mus musculus

10 <400> 41

Arg Ala Ser Asn Leu Glu Ser
1 5

15 <210> 42
<211> 8
<212> PRT
<213> Mus musculus

20 <400> 42

Gln Gln Ser Lys Glu Tyr Pro Thr
1 5

25 <210> 43
<211> 118
<212> PRT
30 <213> Artificial Sequence

<220>
<223> 317-4B2 pro-Vh

35 <400> 43

40

45

50

55

EP 3 702 373 B9

1 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 5 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 10 Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 15 Gly Val Ile Tyr Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 20 Ser Arg Val Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Phe Ser Leu
 25 Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 30 Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
 35 Thr Val Thr Val Ser Ser
 40
 45
 50
 55

<210> 44

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> 317-4B2 pro-Vk

<400> 44

EP 3 702 373 B9

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 5
 Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Ser Asn Asp
 20 25 30
 10
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
 35 40 45
 15
 Asn Tyr Ala Phe His Arg Phe Thr Gly Val Pro Asp Arg Phe Ser Gly
 50 55 60
 20
 Ser Gly Tyr Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala
 65 70 75 80
 25
 Glu Asp Val Ala Val Tyr Tyr Cys His Gln Ala Tyr Ser Ser Pro Tyr
 85 90 95
 30
 Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 45

<211> 118

<212> PRT

30 <213> Artificial Sequence

<220>

<223> 317-4B5 pro-Vh

35 <400> 45

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15
 40
 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30
 45
 Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45
 50
 Gly Val Ile Tyr Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60
 55

EP 3 702 373 B9

Ser Arg Val Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val Ser Leu
 65 70 75 80

5

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

10

Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
 100 105 110

Thr Val Thr Val Ser Ser
 115

15

<210> 46
 <211> 107
 <212> PRT
 <213> Artificial Sequence

20

<220>
 <223> 317-4B5 pro-Vk

25

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

30

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Glu Ser Val Ser Asn Asp
 20 25 30

35

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
 35 40 45

40

Asn Tyr Ala Phe His Arg Phe Thr Gly Val Pro Asp Arg Phe Ser Gly
 50 55 60

45

Ser Gly Tyr Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala
 65 70 75 80

50

Glu Asp Val Ala Val Tyr Tyr Cys His Gln Ala Tyr Ser Ser Pro Tyr
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105

55

<210> 47
 <211> 354
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> 317-1 cDNA-Vh

EP 3 702 373 B9

<400> 47

```

caggtgcagc tgcaggagtc gggaccagga ctggtgaagc cttcggagac cctgtccctc      60
5  acctgcactg tctctgggtt ttcattaacc agctatggtg tacactggat ccggcagccc      120
ccaggaagg gactggagtg gctgggggtc atatgggccc gtggaagcac aaattataat      180
10 ccctccctca agagtcgact gaccatatca aaagacaact ccaagagcca ggtttccctg      240
aagatgagct ctgtgaccgc tgcggacacg gccgtgtatt actgtgagag agcctatggt      300
aactactggt acatcgatgt ctggggccaa gggaccacgg tcaccgtctc ctca          354

```

15 <210> 48
 <211> 118
 <212> PRT
 <213> Artificial Sequence

20 <220>
 <223> 317-1 pro-Vh

<400> 48

```

25      Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
      1              5              10              15
30      Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
      20              25              30
35      Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu
      35              40              45
40      Gly Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
      50              55              60
45      Ser Arg Leu Thr Ile Ser Lys Asp Asn Ser Lys Ser Gln Val Ser Leu
      65              70              75              80
50      Lys Met Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
      85              90
55      Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
      100              105              110
      Thr Val Thr Val Ser Ser
      115

```

55 <210> 49
 <211> 321
 <212> DNA
 <213> Artificial Sequence

EP 3 702 373 B9

<220>

<223> 317-1 cDNA-Vk

<400> 49

5
 gacatcgtga tgaccagtc tccagactcc ctggctgtgt ctctgggcga gagggccacc 60
 atcaactgca aggccagcca gagtgtgagt aatgatgtag cttggtacca gcagaaacca 120
 10 ggacagcctc ctaagctgct cattaactat gcatttcacg gcttcactgg ggtccctgac 180
 cgattcagtg gcagcgggta tgggacagat ttcactctca ccatcagcag cctgcaggct 240
 gaagatgtgg cagtttatta ctgtcaccag gcttatagtt ctccgtacac gtttggcggg 300
 15 gggaccaagc tggagatcaa a 321

<210> 50

<211> 107

20 <212> PRT

<213> Artificial Sequence

<220>

<223> 317-1 pro-Vk

25 <400> 50

30 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Ser Asn Asp
 20 25 30
 35 Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
 35 40 45
 40 Asn Tyr Ala Phe His Arg Phe Thr Gly Val Pro Asp Arg Phe Ser Gly
 50 55 60
 45 Ser Gly Tyr Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala
 65 70 75 80
 50 Glu Asp Val Ala Val Tyr Tyr Cys His Gln Ala Tyr Ser Ser Pro Tyr
 85 90 95
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 51

<211> 114

55 <212> PRT

<213> Artificial Sequence

EP 3 702 373 B9

<220>

<223> 326-3B1 pro-Vh

<400> 51

5

Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
1 5 10 15

10

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
20 25 30

15

Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
35 40 45

20

Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Asp Phe
50 55 60

Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80

25

Leu Gln Ile Ser Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

30

Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
100 105 110

Ser Ser

35

<210> 52

<211> 110

<212> PRT

<213> Artificial Sequence

40

<220>

<223> 326-3B1 pro-Vk

<400> 52

45

50

55

EP 3 702 373 B9

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Gly
 1 5 10 15
 5
 Gln Arg Ala Thr Ile Thr Cys Arg Ala Ser Glu Ser Val Asp Asn Tyr
 20 25 30
 10
 Gly Tyr Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 15
 Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
 50 55 60
 20
 Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn
 65 70 75 80
 25
 Pro Val Glu Ala Asn Asp Thr Ala Asn Tyr Tyr Cys Gln Gln Ser Lys
 85 90 95
 30
 Glu Tyr Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105 110

<210> 53

<211> 114

<212> PRT

30 <213> Artificial Sequence

<220>

<223> 326-3G1 pro-Vh

35 <400> 53

40

45

50

55

EP 3 702 373 B9

1 Gln Val Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Ala
 5 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 10 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
 15 Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Asp Phe
 20 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 25 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
 30 Ser Ser

30 <210> 54
 <211> 110
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> 326-3G1 pro-Vk

<400> 54

40 Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Gly
 1 5 10 15

45

50

55

EP 3 702 373 B9

Gln Arg Ala Thr Ile Thr Cys Arg Ala Ser Glu Ser Val Asp Asn Tyr
 20 25 30

5 Gly Tyr Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45

10 Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
 50 55 60

15 Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn
 65 70 75 80

20 Pro Val Glu Ala Asn Asp Thr Ala Asn Tyr Tyr Cys Gln Gln Ser Lys
 85 90 95

Glu Tyr Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105 110

25 <210> 55
 <211> 342
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> 326-1 cDNA-Vh
 <400> 55

caggtgcagc tgggtgcagag cggcagcag ctgaagaagc ccggcgccag cgtgaaggtg 60

35 agctgcaagg ccagcggcta caccttcacc aactacggca tgaactgggt gagacaggcc 120

cccggccagg gcctggagtg gatgggctgg atcaacaaca acaacggcga gccacctac 180

40 gcccagggct tcagagggcag attcgtgttc agcctggaca ccagcgccag caccgcctac 240

ctgcagatca gcagcctgaa gaccgaggac accgccgtgt acttctgcgc cagagacgtg 300

atggactact ggggccaggg caccaccgtg accgtgagca gc 342

45 <210> 56
 <211> 114
 <212> PRT
 <213> Artificial Sequence

50 <220>
 <223> 326-1 pro-Vh
 <400> 56

55

EP 3 702 373 B9

1 Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
 5 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 10 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 15 Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Gly Phe
 20 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 25 Leu Gln Ile Ser Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr Phe Cys
 30 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val

Ser Ser

30 <210> 57
 <211> 330
 <212> DNA
 <213> Artificial Sequence

35 <220>
 <223> 326-1 cDNA-Vk

<400> 57

40 gacattgtgc tgaccagtc tccagcctcc ttggccgtgt ctccaggaca gagggccacc 60
 atcacctgca gagccagtga aagtgtgat aattatggct atagttttat gactgggat 120
 cagcagaaac caggacaacc tcctaaactc ctgatttacc gtgcatccaa cctagaatct 180
 45 ggggtcccag ccaggttcag cggcagtggg tctaggaccg atttcaccct cacaattaat 240
 cctgtggaag ctaatgatac tgcaaattat tactgtcagc aaagtaaaga atatccgacg 300
 50 ttcggcggag ggaccaaggt ggagatcaaa 330

50 <210> 58
 <211> 110
 <212> PRT
 <213> Artificial Sequence

55 <220>
 <223> 326-1 pro-Vk

EP 3 702 373 B9

<400> 58

5 Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Gly
 1 5 10 15

10 Gln Arg Ala Thr Ile Thr Cys Arg Ala Ser Glu Ser Val Asp Asn Tyr
 20 25 30

15 Gly Tyr Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45

20 Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
 50 55 60

25 Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn
 65 70 75 80

30 Pro Val Glu Ala Asn Asp Thr Ala Asn Tyr Tyr Cys Gln Gln Ser Lys
 85 90 95

35 Glu Tyr Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105 110

<210> 59
 <211> 16
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 317-1 H-CDR2 or CDR-H2

<400> 59

40 Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser
 1 5 10 15

<210> 60
 <211> 16
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 317-4B2 H-CDR2 or CDR-H2

<400> 60

55 Val Ile Tyr Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser
 1 5 10 15

EP 3 702 373 B9

<210> 61
<211> 11
<212> PRT
<213> Artificial Sequence

5
<220>
<223> 317-4B2 L-CDR1 or CDR-L1

10
<400> 61

Lys Ser Ser Glu Ser Val Ser Asn Asp Val Ala
1 5 10

15
<210> 62
<211> 17
<212> PRT
<213> Artificial Sequence

20
<220>
<223> 326-1 H-CDR2 or CDR-H2

<400> 62

25
Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Gly Phe Arg
1 5 10 15

30
Gly

<210> 63
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> 326-3G1 H-CDR2 or CDR-H2

40
<400> 63

Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Asp Phe Arg
1 5 10 15

45
Gly

<210> 64
<211> 118
<212> PRT
<213> Artificial Sequence

<220>
<223> 317-3A1 pro-Vh

<400> 64

EP 3 702 373 B9

1 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 5 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 10 Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 15 Gly Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 20 Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe Ser Leu
 25 Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 30 Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
 35 Thr Val Thr Val Ser Ser
 40
 45
 50
 55

<210> 65

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> 317-3C1 pro-Vh

<400> 65

EP 3 702 373 B9

Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Val Ser Leu
 65 70 75 80

5

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

10

Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
 100 105 110

Thr Val Thr Val Ser Ser
 115

15

<210> 68
 <211> 118
 <212> PRT
 <213> Artificial Sequence

20

<220>
 <223> 317-3G1 pro-Vh

25

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15

30

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30

35

Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45

40

Gly Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60

45

Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Ser Gln Val Ser Leu
 65 70 75 80

50

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

55

Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
 100 105 110

Thr Val Thr Val Ser Ser
 115

<210> 69
 <211> 118
 <212> PRT

EP 3 702 373 B9

<213> Artificial Sequence

<220>

<223> 317-3H1 pro-Vh

5

<400> 69

10 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
1 5 10 15

15 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
20 25 30

20 Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
35 40 45

25 Gly Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
50 55 60

30 Ser Arg Val Thr Ile Ser Lys Asp Thr Ser Lys Ser Gln Val Ser Leu
65 70 75 80

35 Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
85 90 95

40 Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
100 105 110

45 Thr Val Thr Val Ser Ser
115

<210> 70

<211> 118

40

<212> PRT

<213> Artificial Sequence

<220>

<223> 317-3I1 pro-Vh

45

<400> 70

50

55

EP 3 702 373 B9

1 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 5 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 10 Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 15 Gly Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 20 Ser Arg Val Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Phe Ser Leu
 25 Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 30 Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
 35 Thr Val Thr Val Ser Ser
 40
 45
 50
 55

<210> 72

<211> 118

<212> PRT

<213> Artificial Sequence

<220>

<223> 317-4B3 pro-Vh

<400> 72

EP 3 702 373 B9

1 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 5 5 10 15
 5 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30
 10 Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45
 15 Gly Val Ile Asn Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60
 20 Ser Arg Val Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80
 25 Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95
 30 Arg Ala Tyr Gly Asn Tyr Trp Tyr Ile Asp Val Trp Gly Gln Gly Thr
 100 105 110
 35 Thr Val Thr Val Ser Ser
 115
 40 <210> 73
 <211> 118
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> 317-4B4 pro-Vh
 50 <400> 73
 55 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30
 Gly Val His Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Ile
 35 40 45
 Gly Val Ile Trp Ala Gly Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60

EP 3 702 373 B9

Ser Arg Val Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80

5

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

10

Arg Ala Tyr Gly Asn Tyr Pro Tyr Ile Asp Val Trp Gly Gln Gly Thr
 100 105 110

Thr Val Thr Val Ser Ser
 115

15

<210> 74
 <211> 107
 <212> PRT
 <213> Artificial Sequence

20

<220>
 <223> 317-4A2 pro-Vk

25

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

30

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Ser Asn Asp
 20 25 30

35

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
 35 40 45

40

Asn Tyr Ala Phe His Arg Phe Thr Gly Val Pro Asp Arg Phe Ser Gly
 50 55 60

45

Ser Gly Tyr Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala
 65 70 75 80

50

Glu Asp Val Ala Val Tyr Tyr Cys His Gln Ala Tyr Ser Ser Pro Tyr
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105

55

<210> 75
 <211> 114
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> 326-3A1 pro-Vh

EP 3 702 373 B9

<400> 75

5 Gln Val Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Ala
 1 5 10 15

10 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 30

15 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

20 Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Gly Phe
 50 55 60

25 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80

30 Leu Gln Ile Ser Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

35 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
 100 105 110

40 Ser Ser

<210> 76

<211> 114

<212> PRT

35 <213> Artificial Sequence

<220>

<223> 326-3C1 pro-Vh

40 <400> 76

45

50

55

EP 3 702 373 B9

1 Gln Val Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Ala
 5 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 10 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
 15 Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Asp Phe
 20 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 25 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val
 30 Ser Ser

30 <210> 77
 <211> 114
 <212> PRT
 <213> Artificial Sequence
 35 <220>
 <223> 326-3D1 pro-Vh
 <400> 77

40

45

50

55

EP 3 702 373 B9

1 Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
 5 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 10 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 15 Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Gly Phe
 20 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 25 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
 30 Ser Ser

30 <210> 78
 <211> 114
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> 326-3E1 pro-Vh

<400> 78

40

45

50

55

EP 3 702 373 B9

1 Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
 5 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 10 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
 15 Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Asp Phe
 20 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 25 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val
 Ser Ser
 30 <210> 79
 <211> 114
 <212> PRT
 <213> Artificial Sequence
 35 <220>
 <223> 326-3F1 pro-Vh
 <400> 79
 40 Gln Val Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Ala
 45 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 50 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 55 Gly Trp Ile Asn Asn Asn Asn Gly Glu Pro Thr Tyr Ala Gln Gly Phe
 Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80

EP 3 702 373 B9

Leu Gln Ile Ser Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

5 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val
 100 105 110

Ser Ser

10 <210> 80
 <211> 114
 <212> PRT
 <213> Artificial Sequence

15 <220>
 <223> 326-3B N55D pro-Vh

20 <400> 80

Gln Val Gln Leu Val Gln Ser Gly Ser Glu Leu Lys Lys Pro Gly Ala
 1 5 10 15

25 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 25 30

30 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Lys Trp Met
 35 40 45

35 Gly Trp Ile Asn Asn Asn Asp Gly Glu Pro Thr Tyr Ala Gln Asp Phe
 50 55 60

Arg Gly Arg Phe Val Phe Ser Leu Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80

40 Leu Gln Ile Ser Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

45 Ala Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
 100 105 110

Ser Ser

50 <210> 81
 <211> 110
 <212> PRT
 <213> Artificial Sequence

55 <220>
 <223> 326-4A1 pro-Vk

<400> 81

EP 3 702 373 B9

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Gly
 1 5 10 15
 5 Gln Arg Ala Thr Ile Thr Cys Arg Ala Ser Glu Ser Val Asp Asn Tyr
 20 25 30
 10 Gly Tyr Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 15 Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
 50 55 60
 20 Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn
 65 70 75 80
 25 Pro Val Glu Ala Glu Asp Thr Ala Asn Tyr Tyr Cys Gln Gln Ser Lys
 85 90 95
 30 Glu Tyr Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105 110

<210> 82

<211> 110

<212> PRT

<213> Artificial Sequence

<220>

<223> 326-4A2 pro-Vk

<400> 82

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Gly
 1 5 10 15
 40 Gln Arg Ala Thr Ile Thr Cys Arg Ala Ser Glu Ser Val Asp Asn Tyr
 20 25 30
 45 Gly Tyr Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 50 Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
 50 55 60
 55 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn
 65 70 75 80
 60 Pro Val Glu Ala Asn Asp Thr Ala Asn Tyr Tyr Cys Gln Gln Ser Lys
 85 90 95

EP 3 702 373 B9

Glu Tyr Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105 110

5 <210> 83
 <211> 327
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> hulG4mt1 pro

<400> 83

15 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15

20 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30

25 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60

30 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
 65 70 75 80

35 Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95

40 Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
 100 105 110

Glu Phe Glu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 115 120 125

45 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 130 135 140

50 Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
 145 150 155 160

Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
 165 170 175

55 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
 180 185 190

EP 3 702 373 B9

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
 195 200 205

5 Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 210 215 220

10 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
 225 230 235 240

15 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 245 250 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 260 265 270

20 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 275 280 285

25 Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
 290 295 300

30 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 305 310 315 320

Leu Ser Leu Ser Leu Gly Lys
 325

35 <210> 84
 <211> 327
 <212> PRT
 <213> Artificial Sequence

40 <220>
 <223> hulG4mt2 pro

<400> 84

45

50

55

EP 3 702 373 B9

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15

5 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

10 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

15 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr

20

25

30

35

40

45

50

55

EP 3 702 373 B9

Leu Ser Leu Ser Leu Gly Lys
325

5 <210> 85
<211> 327
<212> PRT
<213> Artificial Sequence

10 <220>
<223> hulG4mt6 pro

<400> 85

15

20

25

30

35

40

45

50

55

EP 3 702 373 B9

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15

5 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

10 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

15 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65 70 75 80

20 Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

25 Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
100 105 110

30 Pro Val Ala Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
115 120 125

Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
130 135 140

35 Ala Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
145 150 155 160

40 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
165 170 175

Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
180 185 190

45 Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu

50

55

EP 3 702 373 B9

		195		200		205										
5	Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg
		210					215					220				
10	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys
	225					230					235					240
15	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp
					245					250					255	
20	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys
				260					265					270		
25	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser
			275					280					285			
30	Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser
		290					295					300				
35	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser
	305					310					315					320
40	Leu	Ser	Leu	Ser	Leu	Gly	Lys									
						325										

<210> 86
 <211> 327
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> hulG4mt8 pro

<400> 86

EP 3 702 373 B9

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15

5 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

10 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

15 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

20 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65 70 75 80

25

30

35

40

45

50

55

EP 3 702 373 B9

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 5 Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
 100 105 110
 10 Pro Val Ala Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 115 120 125
 15 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 130 135 140
 20 Thr Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
 145 150 155 160
 25 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
 165 170 175
 30 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
 180 185 190
 35 Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
 195 200 205
 40 Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 210 215 220
 45 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
 225 230 235 240
 50 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 245 250 255
 55 Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 260 265 270
 60 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 275 280 285
 65 Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
 290 295 300
 70 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 305 310 315 320
 75 Leu Ser Leu Ser Leu Gly Lys

5 <210> 87
<211> 327
<212> PRT
<213> Artificial Sequence

10 <220>
<223> huIgG4mt9 pro

<400> 87

15

20

25

30

35

40

45

50

55

EP 3 702 373 B9

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15

5 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30

10 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45

15 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60

20 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
 65 70 75 80

25 Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95

30 Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
 100 105 110

35 Pro Val Ala Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 115 120 125

40 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 130 135 140

45 Ala Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
 145 150 155 160

50 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
 165 170 175

55 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
 180 185 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
 195 200 205

EP 3 702 373 B9

Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 210 215 220

5
 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
 225 230 235 240

10
 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 245 250 255

15
 Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 260 265 270

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 275 280 285

20
 Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
 290 295 300

25
 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 305 310 315 320

Leu Ser Leu Ser Leu Gly Lys
 325

30
 <210> 88
 <211> 327
 <212> PRT
 <213> Artificial Sequence

35
 <220>
 <223> hulG4mt10 pro

40
 <400> 88

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
 1 5 10 15

45
 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30

50
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45

55
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
 65 70 75 80

EP 3 702 373 B9

Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95

5 Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro
 100 105 110

10 Pro Val Ala Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 115 120 125

15 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 130 135 140

Ala Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp
 145 150 155 160

20 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe
 165 170 175

25 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Val His Gln Asp
 180 185 190

Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu
 195 200 205

30 Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 210 215 220

35 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys
 225 230 235 240

40 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 245 250 255

Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 260 265 270

45 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 275 280 285

50 Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser
 290 295 300

Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 305 310 315 320

55 Leu Ser Leu Ser Leu Gly Lys
 325

EP 3 702 373 B9

<210> 89
 <211> 367
 <212> PRT
 <213> Artificial Sequence

5

<220>
 <223> OS8 pro

10

<400> 89

15

20

25

30

35

40

45

50

55

Met Glu Arg His Trp Ile Phe Leu Leu Leu Leu Ser Val Thr Ala Gly
 1 5 10 15

Val His Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg
 20 25 30

Pro Gly Ala Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 35 40 45

Thr Arg Tyr Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu
 50 55 60

Glu Trp Ile Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn
 65 70 75 80

Gln Lys Phe Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser
 85 90 95

Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val
 100 105 110

Tyr Tyr Cys Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp
 115 120 125

Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Gly Gly Gly Gly Ser Gly
 130 135 140

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gln Ile Val Leu Thr Gln Ser
 145 150 155 160

Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 165 170 175

Ser Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 180 185 190

Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Leu Ala Ser
 195 200 205

EP 3 702 373 B9

Gly Val Pro Ala His Phe Arg Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 210 215 220

5
 Leu Thr Ile Ser Gly Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 225 230 235 240

10
 Gln Gln Trp Ser Ser Asn Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu
 245 250 255

15
 Glu Ile Asn Ser Ser Val Val Pro Val Leu Gln Lys Val Asn Ser Thr
 260 265 270

20
 Thr Thr Lys Pro Val Leu Arg Thr Pro Ser Pro Val His Pro Thr Gly
 275 280 285

25
 Thr Ser Gln Pro Gln Arg Pro Glu Asp Cys Arg Pro Arg Gly Ser Val
 290 300

30
 Lys Gly Thr Gly Leu Asp Phe Ala Cys Asp Ile Tyr Ile Trp Ala Pro
 305 310 315 320

35
 Leu Ala Gly Ile Cys Val Ala Leu Leu Leu Ser Leu Ile Ile Thr Leu
 325 330 335

40
 Ile Cys Tyr His Arg Ser Arg Lys Arg Val Cys Lys Cys Pro Arg Pro
 340 345 350

45
 Leu Val Arg Gln Glu Gly Lys Pro Arg Pro Ser Glu Lys Ile Val
 355 360 365

<210> 90
 <211> 304
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P3Z pro

50
 Met Gln Ile Pro Gln Ala Pro Trp Pro Val Val Trp Ala Val Leu Gln
 1 5 10 15

55
 Leu Gly Trp Arg Pro Gly Trp Phe Leu Asp Ser Pro Asp Arg Pro Trp
 20 25 30

Asn Pro Pro Thr Phe Ser Pro Ala Leu Leu Val Val Thr Glu Gly Asp
 35 40 45

EP 3 702 373 B9

Asn Ala Thr Phe Thr Cys Ser Phe Ser Asn Thr Ser Glu Ser Phe Val
 50 55 60

5 Leu Asn Trp Tyr Arg Met Ser Pro Ser Asn Gln Thr Asp Lys Leu Ala
 65 70 75 80

10 Ala Phe Pro Glu Asp Arg Ser Gln Pro Gly Gln Asp Cys Arg Phe Arg
 85 90 95

Val Thr Gln Leu Pro Asn Gly Arg Asp Phe His Met Ser Val Val Arg
 100 105 110

15 Ala Arg Arg Asn Asp Ser Gly Thr Tyr Leu Cys Gly Ala Ile Ser Leu
 115 120 125

20 Ala Pro Lys Ala Gln Ile Lys Glu Ser Leu Arg Ala Glu Leu Arg Val
 130 135 140

25 Thr Glu Arg Arg Ala Glu Val Pro Thr Ala His Pro Ser Pro Ser Pro
 145 150 155 160

Arg Pro Ala Gly Gln Phe Gln Thr Leu Val Val Gly Val Val Gly Gly
 165 170 175

30 Leu Leu Gly Ser Leu Val Leu Leu Val Trp Val Leu Ala Val Ile Arg
 180 185 190

35 Val Lys Phe Ser Arg Ser Ala Asp Ala Pro Ala Tyr Gln Gln Gly Gln
 195 200 205

Asn Gln Leu Tyr Asn Glu Leu Asn Leu Gly Arg Arg Glu Glu Tyr Asp
 210 215 220

40 Val Leu Asp Lys Arg Arg Gly Arg Asp Pro Glu Met Gly Gly Lys Pro
 225 230 235 240

45 Gln Arg Arg Lys Asn Pro Gln Glu Gly Leu Tyr Asn Glu Leu Gln Lys
 245 250 255

50 Asp Lys Met Ala Glu Ala Tyr Ser Glu Ile Gly Met Lys Gly Glu Arg
 260 265 270

Arg Arg Gly Lys Gly His Asp Gly Leu Tyr Gln Gly Leu Ser Thr Ala
 275 280 285

55 Thr Lys Asp Thr Tyr Asp Ala Leu His Met Gln Ala Leu Pro Pro Arg
 290 295 300

Claims

- 5 1. An antibody comprising an antibody antigen binding domain which binds human PD-1 and comprises a heavy chain variable region (Vh) comprising SEQ ID NO: 24, and a light chain variable region (Vk) comprising SEQ ID NO: 26, wherein the antibody is monospecific.
2. The antibody of claim 1, comprising an IgG4 constant domain comprising any one of SEQ ID NO: 83-88.
- 10 3. The antibody of claim 1, comprising an IgG4 constant domain comprising SEQ ID NO: 87 or SEQ ID NO: 88.
4. The antibody of claim 1, comprising an IgG4 constant domain comprising SEQ ID NO: 88.
- 15 5. An antibody comprising an antibody antigen binding domain which binds human PD-1 and comprises a heavy chain variable region (Vh) comprising SEQ ID NO: 24, and a light chain variable region (Vk) comprising SEQ ID NO: 26, for use in therapy.
6. The antibody for use of claim 5, for use in treating cancer or viral infection in a human.
7. The antibody for use according to claim 6 in treating cancer in a human.
- 20 8. The antibody for use according to any one of claim 5 to 7, comprising an IgG4 constant domain comprising any one of SEQ ID NO: 83-88.
- 25 9. The antibody for use according to any one of claim 5 to 7, comprising an IgG4 constant domain comprising SEQ ID NO: 87 or SEQ ID NO: 88.
- 30 10. The antibody for use according to any one of claim 5 to 7, comprising an IgG4 constant domain comprising SEQ ID NO: 88.

Patentansprüche

- 35 1. Antikörper, der eine Antikörperantigen bindende Domäne umfasst, die menschliches PD-1 bindet und eine variable Schwerekettenregion (Vh), umfassend SEQ ID NO: 24, und eine variable Leichtkettenregion (Vk), umfassend SEQ ID NO: 26, umfasst, wobei der Antikörper monospezifisch ist.
- 40 2. Antikörper nach Anspruch 1, umfassend eine konstante IgG4-Domäne, die eine beliebige der SEQ ID NO: 83-88 umfasst.
3. Antikörper nach Anspruch 1, umfassend eine konstante IgG4-Domäne, die SEQ ID NO: 87 oder SEQ ID NO: 88 umfasst.
4. Antikörper nach Anspruch 1, umfassend eine konstante IgG4-Domäne, die SEQ ID NO: 88 umfasst.
- 45 5. Antikörper, der eine Antikörperantigen bindende Domäne umfasst, die menschliches PD-1 bindet und eine variable Schwerekettenregion (Vh), umfassend SEQ ID NO: 24, und eine variable Leichtkettenregion (Vk), umfassend SEQ ID NO: 26, umfasst, zur Verwendung in der Therapie.
- 50 6. Antikörper zur Verwendung nach Anspruch 5 zur Verwendung bei der Behandlung von Krebs oder einer Virusinfektion bei einem Menschen.
7. Antikörper zur Verwendung nach Anspruch 6 zur Krebsbehandlung bei einem Menschen.
- 55 8. Antikörper zur Verwendung nach einem der Ansprüche 5 bis 7, umfassend eine konstante IgG4-Domäne, die eine beliebige der SEQ ID NO: 83-88 umfasst.
9. Antikörper zur Verwendung nach einem der Ansprüche 5 bis 7, umfassend eine konstante IgG4-Domäne, die SEQ ID NO: 87 oder SEQ ID NO: 88 umfasst.

10. Antikörper zur Verwendung nach einem der Ansprüche 5 bis 7, umfassend eine konstante IgG4-Domäne, die SEQ ID NO: 88 umfasst.

5 **Revendications**

1. Anticorps comprenant un domaine de liaison à un antigène d'anticorps qui se lie au PD-1 humain et comprend une région variable de chaîne lourde (Vh) comprenant SEQ ID NO : 24, et une région variable de chaîne légère (Vk) comprenant SEQ ID NO : 26, dans lequel l'anticorps est monospécifique.
- 10
2. Anticorps selon la revendication 1, comprenant un domaine constant d'IgG4 comprenant l'une quelconque des SEQ ID NO : 83 à 88.
- 15
3. Anticorps selon la revendication 1, comprenant un domaine constant d'IgG4 comprenant SEQ ID NO : 87 ou SEQ ID NO : 88.
4. Anticorps selon la revendication 1, comprenant un domaine constant d'IgG4 comprenant SEQ ID NO : 88.
- 20
5. Anticorps comprenant un domaine de liaison à un antigène d'anticorps qui se lie au PD-1 humain et comprend une région variable de chaîne lourde (Vh) comprenant SEQ ID NO : 24, et une région variable de chaîne légère (Vk) comprenant SEQ ID NO : 26, à utiliser en thérapie.
- 25
6. Anticorps à utiliser selon la revendication 5, à utiliser dans le traitement du cancer ou d'une infection virale chez un être humain.
7. Anticorps à utiliser selon la revendication 6 dans le traitement du cancer chez un être humain.
- 30
8. Anticorps à utiliser selon l'une quelconque des revendication 5 à 7, comprenant un domaine constant d'IgG4 comprenant l'une quelconque de SEQ ID NO : 83 à 88.
9. Anticorps à utiliser selon l'une quelconque des revendications 5 à 7, comprenant un domaine constant d'IgG4 comprenant SEQ ID NO : 87 ou SEQ ID NO : 88.
- 35
10. Anticorps à utiliser selon l'une quelconque des revendications 5 à 7, comprenant un domaine constant d'IgG4 comprenant SEQ ID NO : 88.

40

45

50

55

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6 (A)



(B)




Fig. 7




Fig. 8




Fig. 8 (cont'd)




Fig. 9.




Fig. 10







Fig. 11



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2006121168 A [0005]
- WO 2008156712 A [0005]
- US 4376110 A, Kohler [0018]
- US 8008449 B2 [0096]
- US 8168757 B2 [0096]

Non-patent literature cited in the description

- **BLANK et al.** *Cancer Immunol Immunother*, 2005, vol. 54, 307-314 [0001] [0005]
- **ISHIDA et al.** *EMBO J*, 1992, vol. 11, 3887-3395 [0001]
- **SHINOHARA et al.** *Genomics*, 1994, vol. 23, 704-706 [0001]
- **VIVIER et al.** *Immunol Today*, 1997, vol. 18, 286-291 [0001]
- **CHEMNITZ et al.** *J Immunol*, 2004, vol. 173, 945-954 [0001]
- **RILEY.** *Immunol Rev*, 2009, vol. 229, 114-125 [0002]
- **VIBHARKA et al.** *Exp Cell Res*, 1997, vol. 232, 25-28 [0003]
- **AGATA et al.** *Int Immunol*, 1996, vol. 8, 765-772 [0003]
- **BENNETT et al.** *J Immunol*, 2003, vol. 170, 711-118 [0003]
- **BOUR-JORDAN et al.** *Immunol Rev*, 2011, vol. 241, 180-205 [0003]
- **KONISHI et al.** *Clin Cancer Res*, 2004, vol. 10, 5094-5100 [0004]
- **SHI et al.** *Int J Cancer*, 2008, vol. 128, 887-896 [0004]
- **GAO et al.** *Clin Cancer Res*, 2009, vol. 15, 971-979 [0004]
- **WU et al.** *Acta Histochem*, 2006, vol. 108, 19-24 [0004]
- **THOMPSON et al.** *Proc Natl Acad Sci*, 2004, vol. 101, 17174-17179 [0004]
- **THOMPSON et al.** *Clin Cancer Res*, 2007, vol. 13, 1757-1761 [0004]
- **GHEBEH et al.** *Neoplasia*, 2006, vol. 8, 190-198 [0004]
- **HAMANISHI et al.** *Proc Natl Acad Sci*, 2007, vol. 104, 3360-3365 [0004]
- **NOMI et al.** *Clin Cancer Res*, 2007, vol. 13, 2151-2157 [0004]
- **HINO et al.** *Cancer*, 2010, vol. 116, 1757-1766 [0004]
- **OHIGASHI et al.** *Clin Cancer Res*, 2005, vol. 11, 2947-2953 [0004]
- **ZHANG et al.** *Blood*, 2009, vol. 114, 1545-1552 [0004]
- **BONI et al.** *J Virol*, 2007, vol. 81, 4215-4225 [0005]
- **GOLDEN-MASON et al.** *J Immunol*, 2008, vol. 180, 3637-3641 [0005]
- **OKAZAKI et al.** *Int Immunol*, 2007, vol. 19, 813-824 [0005]
- *PNAS*, 2008, vol. 105, 10483 [0010]
- **KOHLER ; MILSTEIN.** *Eur J Immunol*, 1976, vol. 6, 511-519 [0039]
- **ST GROTH ; SHEIDEGGER.** *J Immunol Methods*, 1980, vol. 35, 1-21 [0039]
- **MECHETNER.** *Methods Mol Biol*, 2007, vol. 378, 1-13 [0039]
- **GEFTER, M.L. et al.** *Somat Cell Genet*, 1977, vol. 3, 231-236 [0043]
- **FLANAGAN, M.L. et al.** *Methods in Molecular Biology*, 2007, vol. 378, 33-52 [0044]
- **KABAT et al.** Sequence of proteins of immunologic interest. NIH, 1991 [0052] [0081]
- **LIGHTLE et al.** *Protein Sci*, 2010, vol. 19 (4), 753-762 [0052] [0081]
- **ZHANG et al.** *Blood*, 2005, vol. 106, 1544-1551 [0055] [0063]
- **KIPRIYANOV et al.** *PEDS*, 1997, vol. 10, 445-453 [0055]
- **FULLER SA et al.** *Curr Protoc Mol Biol.*, 2001 [0055]
- **THAKUR A. et al.** *Vaccine*, 2012, vol. 30, 4907-4920 [0061]
- *Blood*, 2010, vol. 116, 2286 [0063]
- **ZHANG et al.** *Cancer Res*, 2006, vol. 66, 5927 [0063]
- **BROCKS et al.** *Mol Med*, 2001, vol. 7, 461-469 [0069]
- **WU, T.T. ; KABAT, E.A.** *J. Exp. Med.*, 1970, vol. 132, 211-250 [0070]
- **LEFRANC M.-P. et al.** *Nucleic Acids Research*, 1999, vol. 27, 209-212 [0070]
- **MOREA et al.** *Methods*, 2000, vol. 20, 267-279 [0071]
- **AL-LAZIKANI et al.** *Journal of Molecular Biology*, 1997, vol. 273, 927-948 [0071]
- **KIEFER et al.** *Nucleic Acids Research*, 2009, vol. 37, D387-D392 [0071]
- **JACOBSON et al.** *Journal of Physical Chemistry*, 2002, vol. 106, 11673-11680 [0071]

EP 3 702 373 B9

- **NATSUME A et al.** *Drug Des Devel Ther.*, 2009, vol. 3, 7-16 [0084]
- **MOORE GL et al.** *MAbs*, 2010, vol. 2, 181-189 [0084]
- **ANGAL, S.** *Mol Immunol*, 1993, vol. 30, 105-108 [0084]
- **DALL'ACQUA, W. et al.** *Biochemistry*, 1998, vol. 37, 9266-9273 [0084]
- **AALBERSE et al.** *Immunol*, 2002, vol. 105, 9-19 [0084]
- **VAN DER NEUT KOLFSCHOTEN M et al.** *Science*, 2007, vol. 317, 1554-157 [0084]
- **CHAPPEL SM et al.** *Proc. Natl. Acad. Sci. USA*, 1991, vol. 88, 9036-9040 [0084]
- **MUKHERJEE, J. et al.** *FASEB J*, 1995, vol. 9, 115-119 [0084]
- **ARMOUR, K.L. et al.** *Eur J Immunol*, 1999, vol. 29, 2613-2624 [0084]
- **CLYNES, R.A. et al.** *Nature Medicine*, 2000, vol. 6, 443-446 [0084]
- **ARNOLD J.N.** *Annu Rev Immunol*, 2007, vol. 25, 21-50 [0084]
- **BRUSCO, A. et al.** *Eur J Immunogenet*, 1998, vol. 25, 349-55 [0084]
- **IGAWA T. et al.** *Prot Eng Design Select*, 2010, vol. 23, 385-392 [0084] [0087]
- **PERCHIACCA J.M. ; TESSIER P.M.** *Ann Rev Biomol Eng*, 2012, vol. 3, 263-286 [0084] [0087]
- **LIU et al.** *mAbs*, 2010, vol. 2, 480-499 [0094]
- *Immunology*, 2002, vol. 105, 9 [0094]
- **ZHANG et al.** *Immunity*, 2004, vol. 20, 337-347 [0095]
- **LIN D.Y. et al.** *PNAS*, 2008, vol. 105, 3011-3016 [0095]
- **LAZAR-MOLNAR E. et al.** *PNAS*, 2008, vol. 105, 10483-10488 [0095]