



(11) **EP 3 766 348 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
20.01.2021 Bulletin 2021/03

(51) Int Cl.:
A01N 43/12 ^(2006.01) **A01N 25/02** ^(2006.01)
A01P 21/00 ^(2006.01) **A01N 45/00** ^(2006.01)

(21) Application number: **20193002.1**

(22) Date of filing: **07.12.2016**

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **07.12.2015 US 201562263830 P**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
16873712.0 / 3 386 304

(71) Applicant: **Valent BioSciences LLC**
Libertyville, IL 60048 (US)

(72) Inventor: **DEVISETTY, Bala, N.**
Libertyville, IL 60048 (US)

(74) Representative: **Elkington and Fife LLP**
Prospect House
8 Pembroke Road
Sevenoaks, Kent TN13 1XR (GB)

Remarks:

This application was filed on 27-08-2020 as a
divisional application to the application mentioned
under INID code 62.

(54) **CONCENTRATED GIBBERELLIN SOLUTION FORMULATIONS**

(57) The present invention is directed to concentrated agricultural solution formulations comprising: from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of GA3, GA4, GA7 and GA4/7; from about 75 to about 90 % wt/wt of at least one solvent selected from the group consisting of polyethylene glycols with average molecular weights of from

about 190 to about 420 daltons and C8 to C12 fatty acid dimethylamides; from about 0.1 to about 4 % wt/wt of a binder; and from about 0.1 to about 3 % wt/wt of a non-ionic surfactant. The invention is also directed to methods for regulating plant growth comprising treating soil or a plant with an effective amount of the formulations of the present invention.

EP 3 766 348 A1

Description**FIELD OF THE INVENTION**

5 **[0001]** The present invention relates to concentrated gibberellin solution formulations.

BACKGROUND OF THE INVENTION

10 **[0002]** Gibberellins are a class of plant growth regulators which are diterpenoid acids. Gibberellins are commercially produced by fermentation of a natural fungus, *Gibberella fujikuroi*. Gibberellins are marketed under various trade names and are commercially used on a variety of fruit orchards, vegetable crops, row crops, and ornamental crops. The predominantly used gibberellin is gibberellin acid ("GA₃").

15 **[0003]** Gibberellins have been formulated in numerous ways over the years in order to attempt to overcome their low solubility. For example, gibberellins have been formulated for foliar application as solutions, soluble powders, wettable powders, tablets, and water-dispersible granules, and soluble granules. The disadvantages of each category of the prior art formulations are discussed below.

20 **[0004]** Solution formulations of the prior art are disadvantageous in several respects. The formulations are less concentrated due to low solubility of gibberellins, have limited storage stability, and/or contain unacceptable amounts of volatile organic compounds ("VOCs").

25 **[0005]** The low solubility of GA₃, gibberellin₄ ("GA₄"), gibberellin₇ ("GA₇"), or gibberellin_{4/7} ("GA_{4/7}") in some solvents, such as propylene glycol, does not permit preparation of high concentration solution formulations. These low strength solution formulations require larger packaging, more storage space, and higher associated transportation, warehousing, and container disposal costs. Due to very low solubility and undesirable hydrolysis, it has been especially difficult to formulate GA₃ in aqueous systems.

30 **[0006]** In order to overcome solubility issues, some formulations use solvents with amounts of VOCs that are not safe for the environment. For example, isopropyl alcohol and methyl alcohol offer severe disadvantages such as flammability and toxicity, which lead to restrictions in manufacturing, packaging, labeling, transportation, and warehousing of such solutions. Tetrahydrofurfuryl alcohol ("THFA") is considered corrosive to the eye and skin.

35 **[0007]** One way to overcome the solubility issues with GA₃, GA₄, and GA_{4/7} is to prepare soluble powder formulations. These powder formulations dissolve readily when mixed with water and form true solutions. Once the solution is formed, no further mixing or agitation of the tank-mix is required.

40 **[0008]** Another way to overcome the solubility issues is to create a wettable powder. A wettable powder formulation is a dry, finely ground formulation. In this type of formulation, the active ingredient is combined with a finely ground dry carrier, usually a mineral clay, along with other ingredients that enhance the ability of the powder to be suspended in water. Upon mixing the wettable powder with water, a suspension is formed, which is then applied by a spray technique. Often the spray liquid must be continuously mixed to prevent settling of insoluble compositions.

45 **[0009]** However, wettable powders and soluble powder formulations tend to produce dust upon handling, such as when pouring, transferring or measuring them. This dust may pose health hazards. Further, powder formulations tend to wet poorly and also solubilize slowly upon addition to water. Powder formulations thus take longer to wet, disperse and solubilize in a tank-mix. Formation of lumps or partially solubilized spray solutions will lead to uneven distribution of the plant growth regulator in the tank-mix with the potential for reduced field performance. Sometimes, foam in the spray tank caused by spray tank adjuvants can also affect wetting and solubility of wettable and soluble powders. Wettable powder formulations will also leave undesirable insoluble residues both in the tank and on the sprayed foliage and fruit.

50 **[0010]** Another type of agricultural formulation is a tablet. Tablet formulations are premeasured dosage delivery systems. They are useful in small areas, or for ornamental purposes. Tablet formulations may be effervescent, which dissolve in water over a period of two to ten minutes depending upon the type and size of the tablet. However, tablets generally deliver only between 0.1 to 1 gram of active ingredient per tablet. They are not ideal for large-scale field operations. Moreover, effervescent tablets are highly susceptible to humidity and may be slow to dissolve and are expensive.

55 **[0011]** Yet another type of agricultural formulation is a water-dispersible granule. Water-dispersible granules are also known as wettable granules or dry flowables. This type of formulation is similar to a wettable powder, except that the active ingredient is formulated as a dispersible granule. To prepare the water-dispersible granules for spray application, they are dispersed in water and form a suspension upon agitation. Many different water-dispersible granular formulations are known for agricultural chemicals. For example, EP 0 252 897 and U.S. Patent No. 4,936,901 disclose encapsulated plant growth regulators in water dispersible granular formulations; and U.S. Patent No. 5,622,658 discloses an extrudable composition for preparing water-dispersible granules.

60 **[0012]** Water-dispersible granules usually have no greater than eight percent moisture content, and form suspensions when added to aqueous solutions. The resulting suspension must be agitated for a period of time in order to fully disperse

it. Agitation or by-pass recirculation of the tank-mix must also be maintained during application. The quality of water-dispersible granules is highly process- and active-ingredient-dependent; and can result in low yield recoveries, poor attrition resistance leading to dust potential, high manufacturing cost and poor dispersion. Generally, sprays of dissolved water-dispersible granular formulations leave undesirable insoluble residues on the treated foliage and fruit.

[0013] For GA₃, GA₄, GA₇, and GA_{4/7} formulations to be efficacious, the active ingredient must solubilize in tank-mixes prior to application. Otherwise, product efficacy will be severely affected. When water-dispersible granules are used, the grower often may not realize if he has achieved complete solubility of the active ingredient in the spray solutions. In addition, water-dispersible granules can harden over time and thus result in poor dispersibility and solubility of the active ingredient. Dust and caking may be problems with certain water-dispersible granules and powder formulations.

[0014] U.S. Patent No. 6,984,609 B2 discloses a concentrated, water soluble granular plant growth regulatory formulation that is commercially available as ProGibb 40% (available from Valent BioSciences Corporation). The disclosed granules swiftly dissolve in water and provide a true solution at the use rates without any insoluble particulates in the spray mixture. However, some orchard and vegetable growers would prefer solution formulations that are safer to handle and easier to apply. The formulation being highly concentrated is not suitable for direct application to the soil or to rice growing aqueous environments.

[0015] Despite all of these formulation options, currently there are not any commercial gibberellin formulations which are suitable for application to the soil. Soil application is desirable because it places the gibberellin in close proximity to the root system which can absorb the gibberellin.

[0016] Therefore, there is a need for environmentally safe, non-phytotoxic, efficacious, high strength gibberellin solution formulations. The improved formulations should overcome the toxicity, handling, storage, transportation, and solubility issues encountered by prior art formulations. The high strength solution formulations should allow development of granular formulations for direct application to soil. As granules specifically sand based do not have absorption capacity high strength gibberellic acid solutions should allow coating the granular surface uniformly. Granular formulations are ideal for direct application to the soil and plant growing environment such as wetland rice. In addition, high strength solutions are also ideal for reducing costs associated with solvent, storage, and transportation.

SUMMARY OF THE INVENTION

[0017] In one aspect, the present invention is directed to concentrated agricultural solution formulations comprising from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of GA₃, GA₄, GA₇ and GA_{4/7}, from about 75 to about 90 % wt/wt of at least one solvent selected from the group consisting of polyethylene glycols with average molecular weights of from about 190 to about 420 daltons and C8 to C12 fatty acid dimethylamides, from about 0.1 to about 4 % wt/wt of a binder, and from about 0.1 to about 3 % wt/wt of a non-ionic surfactant.

[0018] In a further embodiment, the formulations comprise from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of GA₃, GA₄, GA₇ and GA_{4/7}, from about 75 to about 90 % wt/wt polyethylene glycols with average molecular weights of from about 190 to about 210 daltons, from about 0.1 to about 4 % wt/wt of a binder selected from the group consisting of alkylated vinyl pyrrolidone polymers, a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone, and a copolymer of polyvinylpyrrolidone and 1-eicodene, and from about 0.1 to about 3 % wt/wt of a non-ionic surfactant selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate and polyoxyethylene (20) sorbitan monooleate.

[0019] In another aspect, the invention is directed to methods for regulating plant growth comprising treating soil or a plant with an effective amount of the formulations of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

[0020] Despite gibberellins' known stability issues, Applicant has unexpectedly discovered that formulations with from about 75 to about 90 % wt/wt of at least one solvent selected from the group consisting of polyethylene glycols with average molecular weights of from about 190 to about 420 daltons and C8 to C12 fatty acid dimethylamides, from about 0.1 to about 4 % wt/wt of a binder, and from about 0.1 to about 3 % wt/wt of a non-ionic surfactant allowed for stable, concentrated gibberellin formulations. These concentrated gibberellin formulations are versatile as they can be diluted with water and/or used as pre-mix concentrates to provide formulations for soil application.

[0021] In one embodiment, the present invention is directed to concentrated agricultural solution formulations comprising from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of GA₃, GA₄, GA₇ and GA_{4/7}, from about 75 to about 90 % wt/wt of at least one solvent selected from the group consisting of polyethylene glycols with average molecular weights of from about 190 to about 420 daltons and C8 to C12 fatty acid dimethylamides, from about 0.1 to about 4 % wt/wt of a binder, and from about 0.1 to about 3 % wt/wt of a non-ionic surfactant.

[0022] In a preferred embodiment, the gibberellin is GA₃.

[0023] In another embodiment, the formulation comprises from about 9 to about 12 % wt/wt of at least one gibberellin.

In a preferred embodiment, the formulations comprise from about 9.5 to about 11.5 % wt/wt of at least one gibberellin. In a most preferred embodiment, the formulations comprise about 10.6 % wt/wt at least one gibberellin.

[0024] In a further embodiment, the formulations comprise from about 0.1 to about 4 % wt/wt of a color additive. In a preferred embodiment, the color additive is a dye. In a more preferred embodiment, the dye is a FD&C food grade dye. In an even more preferred embodiment, the dye is selected from the group consisting of FD&C Blue #1, FD&C Blue #2, FD&C Green #3, FD&C Red #3, FD&C Red #40, FD&C Yellow #5, FD&C Yellow #6, and Citrus Red #2. In a most preferred embodiment, the dye is FD&C Blue #1. In another most preferred embodiment, the dye is FD&C Red #40.

[0025] As used herein, "FD&C" refers to the United States' Food, Drug, and Cosmetic Act which is United States Code, Title 21. This code provides the following list (as of October 2015) of food grade color additives which may be added to foods: FD&C Blue #1; FD&C Blue #2; FD&C Green #3; FD&C Red #3; FD&C Red #40; FD&C Yellow #5; FD&C Yellow #6; and Citrus Red #2.

[0026] Polyethylene glycol ("PEG") is a polyether compound with the structure: $\text{H}-(\text{O}-\text{CH}_2-\text{CH}_2)_n-\text{OH}$. PEGs are prepared by polymerization of ethylene oxide and are commercially available over a wide range of molecular weights. The number following "polyethylene glycol," or "PEG," refers to the molecular weight. For example, PEG 200 has a range of molecular weights from 190 to 210, PEG 300 from 285 to 315, and PEG 400 from 380 to 420 daltons. In a preferred embodiment, the solvent is polyethylene glycols with average molecular weights of from about 190 to about 420 daltons. In a more preferred embodiment, the solvent is polyethylene glycols with average molecular weights of from about 190 to about 210 daltons.

[0027] In another embodiment, the solvent is at least one C8 to C12 fatty acid dimethylamide. In a preferred embodiment, the solvent is a mixture of C8 to C10 fatty acid dimethylamides or a mixture of C10 to C12 fatty acid dimethylamides. In a more preferred embodiment, the solvent is a mixture of C8 to C10 fatty acid dimethylamides. A mixture of C8 to C10 fatty acid dimethylamides is commercially available as Agnique® AMD 810 (Agnique is available from BASF and a registered trademark of Cognis Corporation).

[0028] In a further embodiment, the solvent is a mixture of: (1) polyethylene glycols with average molecular weights of from about 190 to about 210 daltons; and (2) a mixture of C8 to C10 fatty acid dimethylamides.

[0029] In an embodiment, the binder is selected from the group consisting of alkylated vinyl pyrrolidone polymers, a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone, and a copolymer of polyvinylpyrrolidone and 1-eicododecene. In a preferred embodiment, the binder is a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone (available from Ashland as Agrimer™ VA-6). Alkylated vinyl pyrrolidone polymers, such as Agrimer™ AL10, and a copolymer of polyvinylpyrrolidone and 1-eicododecene, Agrimer™ VA-71, are also available from Ashland.

[0030] In a further embodiment, the formulations comprise from about 1 to about 3 % wt/wt of the binder. In a preferred embodiment, the formulations comprise from about 1.5 to about 2.5 % wt/wt of the binder. In a more preferred embodiment, the formulations comprise about 2 % wt/wt of the binder.

[0031] In another embodiment, the non-ionic surfactant is selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate and polyoxyethylene (20) sorbitan monooleate. In a preferred embodiment, the non-ionic surfactant is polyoxyethylene (20) sorbitan monolaurate.

[0032] In an embodiment, the formulations comprise from about 0.1 to about 2 % wt/wt of the non-ionic surfactant. In a preferred embodiment, the formulations comprise from about 0.5 to about 1.5 % wt/wt of the non-ionic surfactant. In a more preferred embodiment, the formulations comprise about 1 % wt/wt of the non-ionic surfactant.

[0033] In a further embodiment, the formulations comprise from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of gibberellic acid (GA_3), gibberellin₄ (GA_4), gibberellin₇ (GA_7) and gibberellin_{4/7} ($\text{GA}_{4/7}$), from about 75 to about 90 % wt/wt polyethylene glycols with average molecular weights of from about 190 to about 210 daltons, from about 0.1 to about 4 % wt/wt of a binder selected from the group consisting of alkylated vinyl pyrrolidone polymers, a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone, and a copolymer of polyvinylpyrrolidone and 1-eicododecene, and from about 0.1 to about 3 % wt/wt of a non-ionic surfactant selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate and polyoxyethylene (20) sorbitan monooleate. In a preferred embodiment, the gibberellin is GA_3 . In another preferred embodiment, the binder is a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone. In yet another preferred embodiment, the non-ionic surfactant is polyoxyethylene (20) sorbitan monolaurate.

[0034] In yet another embodiment, the present invention is directed to methods for regulating plant growth comprising treating a plant or soil with an effective amount of a formulation of the present invention.

[0035] The formulations of the present invention are also low VOC formulations. This means that the formulations contain less than or equal to 25 % emission potential, as determined by thermo gravimetric analysis ("TGA"). Gibberellin formulations with greater than 25 % emission potential, as determined by TGA, are considered High-VOC products by CADPR (California Department of Pesticide Regulation). TGA involves heating a sample of the formulation in an environmentally controlled chamber while the rate of sample mass loss is measured. CADPR states that the emission potential of the formulation is determined by taking the mean of three replicate TGA measurements of the formulations and then subtracting the percent water and the exempt compounds from the measurement. The TGA process is well

known by those of skill in the art.

[0036] In a further embodiment, the formulations of the present invention are applied to plants or soil at a rate of from about 0.1 to about 50 grams of gibberellin per acre. In a preferred embodiment, the formulations are applied to the plants or soil at a rate of from about 1 to about 20 grams of gibberellin per acre. In a more preferred embodiment, the formulations

are applied to the plants or soil at a rate of from about 1 to about 10 grams of gibberellin per acre.

[0037] Formulations of the present invention may be used as a pre-mix product. For example, the pre-mix product can be added to a tank mix prior to application to the plants. Alternatively, the formulations of the present invention can be used as a pre-mix to pre-formed granular product for direct application to soil or plant growing environment.

[0038] Formulations of the present invention may be used on any plant in need of gibberellin treatment (or on the soil adjacent to the plant in need of treatment), for example, on: artichokes to accelerate maturity and increase yield; blueberries to improve fruit set and fruit size; bananas to stimulate plant growth and reduce effects of stress, or post-harvest for maintaining fruit quality; carrots to maintain foliage growth during periods of stress; celery to increase plant height and yield; cherries to increase fruit size, firmness and quality or to delay maturity for a more orderly harvest; citrus to increase fruit set and yield, to delay rind aging, reduce physiological disorders, or delay maturity for a more orderly harvest; collard greens to facilitate harvest, increase yield, and improve quality; cotton to promote early season growth and increase seedling vigor; and cucumbers to stimulate fruit set during periods of cool weather; pasture land used for animal grazing; and corn. The formulations can be used post harvest on bananas and citrus, etc. Formulations of the present invention could also be used on grapes, melons, pecans, peppers, pineapples, rice, rhubarb, spinach, stone fruits, sugarcane, wheat, strawberries, watercress and other plants in need of treatment.

[0039] In an embodiment, the formulations of the present invention are applied to a plant selected from the group consisting of rice, cotton, corn, soybeans, sugarcane, wheat and beets. In a preferred embodiment, the formulations of the present invention are applied to rice.

[0040] The disclosed embodiments are simply exemplary embodiments of the inventive concepts disclosed herein and should not be considered as limiting, unless the claims expressly state otherwise.

[0041] The term "effective amount" means the amount of the formulation that will provide the desired effect on the plant that is being treated. The "effective amount" will vary depending on the formulation concentration, the type of plants(s) being treated, and the result desired, among other factors. Thus, it is not always possible to specify an exact "effective amount." However, an appropriate "effective amount" in any individual case may be determined by one of ordinary skill in the art. For example, effective amounts of formulations of the present invention may be from about 0.1 to about 50 grams of gibberellin per acre.

[0042] Other components of the formulation may be included in nominal amounts that do not affect the storage stability characteristics of the present formulations. Additional components include additional surface active agents, crystal growth inhibitors, stickers, spreaders, leaf penetrants, dispersants, systemic acquired resistance inducers, systemic acquired resistance inhibitors, anti-foaming agents, preservatives, pH regulators, cosolvents, humectants, UV protectants, biostimulants, root growth promoters, biopesticides, inoculants, vehicles, sequestrants or other components which facilitate production, storage stability, product handling and application.

[0043] It is also contemplated that the ready-to-mix composition materials of this invention may be used in combination with other active ingredients, such as herbicides, fungicides, insecticides, bactericides, nematocides, biochemical pesticides, biostimulants, biopesticides, plant produced pesticides (botanicals), safeners or plant nutrients.

[0044] As used herein, the term "herbicide" broadly refers to compounds or compositions that are used as herbicides, as well as herbicide safeners and algicides. Herbicides may include, but are not limited to, 1,2,4-triazinones, 1,3,5-triazines, alkanamides (acetamides), anilides, aryloxyalkanoic acids, aryloxyphenoxypropionates, benzamides, benzamides (L), benzenedicarboxylic acids, benzofurans, benzoic acids (auxins), benzonitriles, benzothiadiazinones, benzothiazolones, carbamates (DHP), carbamates, chloroacetamides, cyclohexanedione oximes, dinitroanilines, dinitrophenols, diphenyl ethers, diphenyl ethers (cbi), glycine derivatives, halogenated alkanic acids, hydroxybenzonitriles, imidazolinones, isoxazoles, isoxazolidinones, N-phenylphthalimides, organoarsenics, oxadiazoles, oxazolidinediones, oxyacetamides, phenoxy-carboxylic acids, phenyl carbamate herbicides, phenylpyrazole herbicides, phenylpyridazines, phosphinic acids, phosphorodithioates, phthalamates, pyrazole herbicides, pyridazines, pyridazinones (PDS), pyridazinones (PSII), pyridines, pyridinecarbox-amides, pyridinecarboxylic acids, pyrimidindiones, pyrimidines, pyrimidinyl-oxybenzoics, pyrimidinyl-oxybenzoic analogs, quinolinecarboxylic acids, BI class and IV including thiocarbamate, semicarbazones, sulfonylaminocarbonyl-triazolinones, sulfonylureas, tetrazolinones, thiadiazoles, thiocarbamates, triazoles, triazolinones, triazolopyrimidines, triketones, uracils, ureas, 2,3,6-TBA, 2,4,5-T, 2,4-D, 2,4-D-2-ethylhexyl, 2,4-DB, 2,4-D-dimethylammonium, 2,4-D-isopropyl, 2,4-D-isopropyl, 2,4-D-trolamine (2,4-D-triethanolamine), ACD 10614, ACD 10435, acetochlor, acifluorfen, acifluorfen-sodium, aclonifen, acrolein, AD 67, alachlor, alloxydim-sodium, ametryn, amicarbazone, amidosulfuron, amitrole, ammonium sulfamate, anilofos, asulam, atraton, atrazine, azafenidin, azimsulfuron, aziprotryne, barban, beflubutamid, benazolin, benazolin-ethyl, benfluralin, benfuresate, benoxacor, bensulfuron-methyl, bensulide, bentazone, benzobicyclon, benzofenap, benzoylprop, enzoylprop-ethyl, bifenox, bilanafos-sodium, bispyribac-sodium, borax, bromacil, bromobutide, bromofenoxim, bromoxynil, bromoxynil octanoate, bromoxynil-potassium,

brompyrazon, butachlor, butafenacil, butenachlor, buthidazole, butralin, butroxydim, buturon, cafenstrole, calcium cyanamide, carbetamide, carfentrazone-ethyl, chlormethoxyfen, chloramben, chlorbromuron, chlorbufam, chlorfenac, chlorfenac-sodium, chlorfenprop, chlorfenprop-methyl, chlorfenprop, chlorfenprop-ethyl, chlorflurenol-methyl, chloridazon, chlorimuron-ethyl, chlornitrofen, chloroacetic acid, chloro-toluron, chloroxuron, chlorpropham, chlorsulfuron, chlorthal-dimethyl, chlorthiamid, cinidon-ethyl, cinosulfuron, clethodim, sethoxydim, tepraloxym, tralkoxydim, clodinafop-prop-argyl, clofop, clofop-isobutyl, clomazone, clomeprop, clopyralid, cloquintocet-mexyl, cloransulam-methyl, credazine, cumyluron, cyanamide, cyanazine, cyclosulfamuron, cycloxydim, cycluron, cyhalofop-butyl, cyometrinil, daimuron, dazomet, desmedipham, desmetryn, di-allate, dicamba, dichlobenil, dichlormid, dichlorprop, dichlorprop-isooctyl, dichlorprop-P, diclofop, diclofop-methyl, diclosulam, diethatyl-ethyl; diethatyl, difenoxuron, difenzoquat metilsulfate, diflufenican, diflufenzopyr, dikegulac, dikegulac-sodium, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethipin, dimethylarsinic acid, dinitramine dinoseb, dinoseb acetate, dinoterb, diphenamid, dipropetryn, disul, disul-sodium, dithiopyr, diuron, DNOC, DSMA, eglinazone-ethyl, eglinazone, EL 177, endothal, ethalfluralin, ethametsulfuron-methyl, ethidimuron, ethofumesate, ethoxysulfuron, etobenzanid, fenchlorazole-ethyl, fenclorim, fenoprop, fenoprop-butotyl, fenoxaprop-ethyl, fenoxaprop, fenoxaprop-P, fenoxaprop-P-ethyl, fenthiaprop, fenthiaprop-ethyl, fentrazamide, fenuron, flammoprop-methyl, flammoprop-isopropyl, flammoprop, flammoprop-M-isopropyl, flammoprop-M-methyl, flazasulfuron, florasulam, fluaazifop-butyl, fluaazifop-P, fluaazifop-P-butyl, fluaazolate, flucarbazone-sodium, fluchloralin, flufenacet, flumetsulam, flumiclorac-pentyl, flumioxazin, flumipropyn, fluometuron, fluorodifen, fluoroglycofen-ethyl, fluothuron, flupoxam, flupropanate-sodium, flupyr-sulfuron-methyl-sodium, flurazole, flurenol-butyl, fluridone, flurochloridone, fluroxypyr, fluroxypyr-meptyl, flurtamone, fluthiacet-methyl, fomesafen, foramsulfuron, fosamine-ammonium, furilazole, glufosinate-ammonium, glyphosate, glyphosate-ammonium, glyphosate-isopropylammonium, glyphosate-sodium, glyphosate-trimesium, halosulfuron-methyl, haloxyfop, haloxyfop-etotyl, haloxyfop-P, hexaflurate, hexazinone, imazamethabenzmethyl, imazamox, imazapic, imazapyr, imazapyr-isopropylammonium, imazaquin, imazethapyr, imazosulfuron, indanofan, iodosulfuron-methyl-sodium, ioxynil, ioxynil octanoate, ioxynil-sodium, isocarbamid, isocil, isomethiozin, isonoruron, isoproturon, isouron, isoxaben, isoxaflutole, isoxapyrifop, karbutilate, lactofen, lenacil, linuron, LS830556, maleic hydrazide, MCPA, MCPA-thioethyl, MCPB, MCPB-ethyl, mecoprop, mecoprop-P, medinoterb acetate, medinoterb, mefenacet, mefenpyr-diethyl, mefluidide, mesosulfuron-methyl, mesotrione, metamifop, metamitron, metazachlor, methabenzthiazuron, methazole, methiuron, methoprotetryn, methoxyphenone, methyl isothiocyanate, methylarsonic acid, methylaldimuron, metobenzuron, metobromuron, metolachlor, metosulam, metoxuron, metribuzin, metsulfuron-ethyl, MK-616, monalide, monolinuron, monuron, monuron-TCA, MSMA, naphthalic anhydride, naproanilide, napropamide, naptalam, NC-330, neburon, nicosulfuron, nitratin, nitrofen, norflurazon, orbencarb, oryzalin, oxabetrinil, oxadiargyl, oxadiazon, oxasulfuron, oxaziclomefone, oxyfluorfen, pendimethalin, penoxsulam, pentachlorophenol, pentanochlor, pentoxazone, perfludone, pethoxamid, phenisopham, phenmedipham, phenylmercury acetate, picolinafen, primisulfuron-methyl, pro-diamine, profluralin, proglinazone-ethyl, proglinazone, prometon, prometryn, propachlor, propanil, propaquizafop, propazine, propham, propisochlor, propoxycarbazone-sodium, propyzamide, prosulfuron, pyraflufen-ethyl, pyrazolynate, pyrazosulfuron-ethyl, pyrazoxyfen, pyribenzoxim, pyributicarb, pyridate, pyrifthalid, pyriminobac-methyl, pyri-thiobac-sodium, quinclozac, quinmerac, quincloamine, quizalofop-ethyl, quizalofop-P-ethyl, quizalofop-P-tefuryl, rimsulfuron, se-buthylazine, sebumeton, siduron, simazine, simetryn, S-metolachlor, SMY 1500, sodium chlorate, sulcotrione, sulfentrazone, sulfometuron-methyl, sulfosulfuron, tebuthiuron, tepraloxym, terbacil, terbutometon, terbuthylazine, terbutryn, thenylchlor, thiazafluron, thiazopyr, thidiazimin, thifensulfuron-methyl, thiobencarb, 1-dichloroacetylazepane, tralkoxydim, tri-allat, triasulfuron, tribenuron-methyl, trichloroacetic acid, triclopyr, tridiphane, trietazine, trifloxysulfuron-sodium, trifluralin, and triflurosulfuron-methyl.

[0045] Fungicides may include, but are not limited to, amino acid amide carbamates, anilino-pyrimidines, antibiotics, aromatic hydrocarbons, heteroaromatics, chloro/nitrophenyls, benzamides (F), benzenesulfonamides, benzimidazoles, benzimidazole precursors, benzotriazines, carboxamides, cinnamic acids, cyanoacetamide oximes, dicarboximides, dithiolanes, DMI: imidazoles, piperazines, pyrimidines, and triazoles; enopyranuronic acid antibiotics, heteroaromatic hydroxylanilides, MBI: dehydratases and reductases; morpholine morpholines, morpholine spiroketalamines, multi-site chloronitriles, multi-site dimethyldithiocarbamates, multi-site guanidines, multi-site inorganics, multi-site phthalimides, multi-site quinones, multi-site sulfamides, N-phenyl carbamate fungicides, organotin fungicides, phenylamide: acylalanines, phenylamide: butyrolactones, phenylamide: oxazolidinones, phenylpyrroles, phenylurea fungicides, phosphonates, phosphorothiolates, pyridazinone fungicides, pyrimidinamines, pyrimidinols, Qil, quinolines, SBI class IV: thio-carbamates, strobilurin analog: dihydrodioxazines, strobilurin type: imidazolinones, strobilurin type: methoxyacrylates, strobilurin type: ethoxycarbamates, strobilurin type: oxazolidinediones, strobilurin type: oximinoacetamides, strobilurin type: oximinoacetates, thiazolecarboxamides, thiocarbamate fungicides, and thiophenecarboxamides. Suitable fungicides include 1,2-dichloropropane, 2-methoxyethylmercury chloride, 2-phenylphenol, 8-hydroxy-quinoline sulfate, am-propylfos, anilazine, azaconazole, azoxystrobin, benalaxyl, benodanil, benomyl, benquinox, benthiavalicarb-isopropyl, binapacryl, biphenyl, bis(tributyltin) oxide, bitertanol, blasticidin-S, borax, boscalid, bromuconazole, bupirimate, buthiobate, captafol, captan, carbendazim, carboxin, carpropamid, CGA 80 000, chinomethionat, chlobenthiazole, chloraniformethan, chloroneb, chlorothalonil, chlozolinate, climbazole, copper oxychloride, copper sulfate, copper sulfate (trib-

asic), cuprous oxide, cyazofamid, cyflufenamid, cymoxanil, cyproconazole, cyprodinil, cyprofuram, dazomet, dichloflu-
 anid, dichlone, dichlorophen, diclobutrazol, diclocymet, diclomezine, dicloran, diethofencarb, difenoconazole, difenzo-
 quat metilsulfate, diflumetorim, dimethirimol, dimethomorph, dimoxystrobin, diniconazole, dinobuton, dinocap, diphe-
 nylamine, ditalimfos, dithianon, dodemorph, dodemorph acetate, dodine, drazoxolon, edifenphos, epoxiconazole, eta-
 conazole, etem, ethaboxam, ethirimol, etridiazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fenfuram, fen-
 hexamid, fenitropan, fenoxanil, fencpiclonil, fenpropimorph, fentin acetate, fentin hydroxide, ferimzone, fluazinam, fludi-
 oxonil, flumorph, fluoroimide, fluotrimazole, fluoxastrobin, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol,
 folpet, fosetyl-aluminum, fuberidazole, furalaxyl, furametpyr, furconazole-cis, furnecyclox, glyodin, griseofulvin, halac-
 rinate, hexachlorobenzene, hexaconazole, hymexazol, imazalil, imibenconazole, iminoctadine triacetate, iminoctadine
 tris(albesilate), ipconazole, iprodione, iprovalicarb, isoprothiolane, kasugamycin hydrochloride hydrate, kresoxim-methyl,
 mebenil, mepanipyrim, mepronil, mercuric chloride, metalaxyl, metalaxyl-M, metconazole, methasulfocarb, methfurox-
 am, methyl iodide, methyl isothiocyanate, metominostrobin, metsulfovax, mildiomyacin, myclobutanil, myclozolin, na-
 tamycin, nitrothal-isopropyl, nuarimol, ofurace, oleic acid, fatty acids), oxabetrinil, oxadixyl, oxpoconazole fumarate,
 oxycarboxin, penconazole, pencycuron, pentachlorophenol, phenylmercury acetate, phenylmercury dimethyldithiocar-
 bamate, phenylmercury nitrate, phosphonic acid, phthalide, picoxystrobin, polyoxin B, polyoxorim, potassium bicarbo-
 nate, potassium hydroxyquinoline sulfate, prochloraz, procymidone, propamocarb hydrochloride, propiconazole, pro-
 quinazid, prothiocarb; prothiocarb hydrochloride, prothioconazole, pyracarbolid, pyraclostrobin, pyrazophos, pyributi-
 carb, pyrimethanil, pyroquilon, quinochloramine, quinoxifen, quintozone, silthiofam, simeconazole, sodium bicarbonate,
 spiroxamine, SSF-109, sulfur, tebuconazole, tecnazene, tetraconazole, thiabendazole, thicyofen, thifluzamide, thi-
 ophanate, thiophanate-methyl, thiram, tiadinil, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, triamiphos, triazox-
 ide, trichlamide, tricyclazole, trifloxystrobin, triflumizole, triforine, triticonazole, urbacid, validamycin, vinclozolin, zarilamid,
 ziram, and zoxamide.

[0046] Bactericides may include, but are not limited to, bronopol, dichlorophen, nitrpyrin, nickel dimethyldithiocar-
 bamate, kasugamycin, octhilinone, furancarboxylic acid, oxytetracyclin, probenazole, streptomycin, tecloftalam, copper
 sulfate and other copper preparations.

[0047] Insecticides, acaricides and nematocides may include, but are not limited to, abamectin, ABG-9008, acephate,
 acequinocyl, acetamiprid, acetoprole, acrinathrin, AKD-1022, AKD-3059, AKD-3088, alanycarb, aldicarb, aldoxycarb,
 allethrin, alpha-cypermethrin (alphamethrin), amidoflumet, aminocarb, amitraz, avermectin, AZ-60541, azadirachtin,
 azamethiphos, azinphos-methyl, azinphos-ethyl, azocyclotin, *Bacillus firmus*, *Bacillus popilliae*, *Bacillus sphaericus*,
Bacillus subtilis, *Bacillus thuringiensis*, *Bacillus thuringiensis* strain EG-2348, *Bacillus thuringiensis* strain GC-91, *Bacillus*
thuringiensis strain NCTC-11821, *Bacillus thuringiensis israelensis*, baculoviruses, *Beauveria bassiana*, *Beauveria ten-*
ella, benclothiaz, bendiocarb, benfuracarb, bensultap, benzoximate, beta-cyfluthrin, beta-cypermethrin, bifenazate,
 bifenthrin, binapacryl, bioallethrin, bioallethrin-5-cyclopentyl isomer, bioethanomethrin, biopermethrin, bioresmethrin,
 bistrifluoron, BPMP, brofenprox, bromophos-ethyl, bromopropylate, bromfenvinfos (-methyl), BTG-504, BTG-505, bufen-
 carb, buprofezin, butathiofos, butocarboxim, butoxycarboxim, butylpyridaben, cadusafos, camphechlor, carbaryl, car-
 bofuran, carbophenothion, carbosulfan, cartap, CGA-50439, chinomethionat, chlorantraniliprole, chlordane, chlordime-
 form, chloethocarb, chlorethoxyfos, chlorfenapyr, chlorfenvinphos, chlorfluazuron, chlormephos, chlorobenzilate, chlo-
 ropicrin, chlorproxyfen, chlorpyrifos-methyl, chlorpyrifos (-ethyl), chlovaporthrin, chromafenozide, cis-cypermethrin, cis-
 resmethrin, cis-permethrin, clocythrins, cloethocarb, clofentezine, clothianidin, clothiazoben, codlemone, coumaphos,
 cyanofenphos, cyanophos, cyantraniliprole, cycloprene, cycloprothrin, cyfluthrin, cyflumetofen, cyhalothrin, cyhexatin,
 cypermethrin, cyphenothrin (1R-trans-isomer), cyromazine, DDT, deltamethrin, demeton-5-methyl, demeton-5-methyl-
 sulfone, diafenthiuron, dialifos, diazinon, dichlofenthion, dichlorvos, dicofol, dicrotophos, dicyclanil, diflubenzuron, dime-
 fluthrin, dimethoate, dimethylvinphos, dinobuton, dinocap, dinotefuran, diofenolan, disulfoton, docusat-sodium,
 dofenapyn, DOWCO-439, eflusilanate, emamectin, emamectin-benzoate, empenhrin (1R-isomer), endosulfan, *Ento-*
mophora spp., EPN, esfenvalerate, ethiofencarb, ethion, ethiprole, ethoprophos, etofenprox, etoxazole, etrimfos, fam-
 phur, fenamiphos, fenazaquin, fenbutatin oxide, fenfluthrin, fenitrothion, fenobucarb, fenothiocarb, fenoxacrim, fenoxyc-
 carb, fenpropathrin, fenpyrad, fenpyrithrin, fenpyroximate, fensulfothion, fenthion, fentripanil, fenvalerate, fipronil, floni-
 camid, fluacrypyrim, fluazuron, flubendiamide, flubenzimine, flubrocycythrinate, flucycloxuron, flucythrinate, flufenerim,
 flufenoxuron, flufenprox, flumethrin, flupyrzofos, flutenzin (flufenzine), fluvalinate, fonofos, formetanate, formothion,
 fosmethilan, fosthiazate, fubfenprox (fluproxyfen), furathiocarb, gamma-cyhalothrin, gamma-HCH, gossypure, grand-
 lure, granulos virus, halfenprox, halofenozide, HCH, HCN-801, heptenophos, hexaflumuron, hexythiazox, hydram-
 ethylnone, hydroprene, IKA-2002, imidacloprid, imiprothrin, indoxacarb, iodofenphos, iprobenfos, isazofos, isofenphos,
 isoprocarb, isoxathion, ivermectin, japonilure, kadethrin, nuclear polyhedrosis viruses, kinoprene, lambda-cyhalothrin,
 lindane, lufenuron, malathion, mecarbam, mesulfenfos, metaldehyde, metam-sodium, methacryfos, methamidophos,
 Metharhizium anisopliae, Metharhizium flavoviride, methidathion, methiocarb, methomyl, methoprene, methoxychlor,
 methoxyfenozide, metofluthrin, metolcarb, metoxadiazone, mevinphos, milbemectin, milbemycin, MKI-245, MON-45700,
 monocrotophos, moxidectin, MTI-800, naled, NC-104, NC-170, NC-184, NC-194, NC-196, niclosamide, nicotine, niten-
 pyram, nithiazine, NNI-0001, NNI-0101, NNI-0250, NNI-9768, novaluron, noviflumuron, OK-5101, OK-5201, OK-9601,

OK-9602, OK-9701, OK-9802, omethoate, oxamyl, oxydemeton-methyl, Paecilomyces fumosoroseus, parathion-methyl, parathion (-ethyl), permethrin (cis-, trans-), petroleum, PH-6045, phenothrin (1R-trans isomer), phenthoate, phorate, phosalone, phosmet, phosphamidon, phosphocarb, phoxim, piperonyl butoxide, pirimicarb, pirimiphos-methyl, pirimiphos-ethyl, potassium oleate, prallethrin, profenofos, profluthrin, promecarb, propaphos, propargite, propetamphos, propoxur, prothiofos, prothoate, protrifenbute, pymetrozine, pyraclofos, pyrafluprole, pyresmethrin, pyrethrum, pyridaben, pyridalyl, pyridaphenthion, pyridathion, pyrimidifen, pyriprole, pyriproxyfen, quinalphos, resmethrin, RH-5849, ribavirin, RU-12457, RU-15525, rynaxapyr, S-421, S-1833, salithion, sebufos, SI-0009, silafluofen, spinosad, spiroadiclofen, spiromesifen, sulfluramid, sulfotep, sulprofos, SZI-121, tau-fluvalinate, tebufenozide, tebufenpyrad, tebupirimfos, te-flubenzuron, tefluthrin, temephos, temivinvphos, terbam, terbufos, tetrachlorvinphos, tetradifon, tetramethrin, tetramethrin (1R-isomer), tetrasul, theta-cypermethrin, thiacloprid, thiamethoxam, thiapronil, thiatrithos, thiocyclam hydrogenoxalate, thiodicarb, thiofanox, thiometon, thiosultap-sodium, thuringiensin, tolfenpyrad, tralocyrthrin, tralomethrin, transfluthrin, triarathene, triazamate, triazophos, triazuron, trichlophenidine, trichlorfon, triflumuron, trimethacarb, vamidothion, vaniliprole, verbutin, Verticillium lecanii, WL-108477, WL-40027, yl-5201, yl-5301, yl-5302, XMC, xylylcarb, ZA-3274, zeta-cypermethrin, zolaprofos, ZXI-8901, the compound 3-methylphenyl propylcarbamate (Tsumacide Z), the compound 3-(5-chloro-3-pyridinyl)-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3-.2.1]octane-3-carbonitrile (CAS-Reg. No. 185982-80-3) and the corresponding 3-endo-isomer (CAS-Reg. No. 185984-60-5) (cf. WO-96/37494, WO-98/25923), and also preparations which comprise insecticidally active plant extracts, nematodes, fungi or viruses.

[0048] As used herein, all numerical values relating to amounts, weight percentages and the like are defined as "about" or "approximately" each particular value, namely, plus or minus 10 % ($\pm 10\%$). For example, the phrase "at least 5 % by weight" is to be understood as "at least 4.5 % to 5.5 % by weight." Therefore, amounts within 10 % of the claimed values are encompassed by the scope of the claims.

[0049] As used herein, "% wt/wt" refers to the percent weight of the material in relation to the percent weight of the formulation.

[0050] The following examples are intended to illustrate the present invention and to teach one of ordinary skill in the art how to make the formulations of the invention. They are not intended to be limiting in any way.

EXAMPLES

[0051] The following products were the sources of the materials used in the examples.

[0052] Agrimer™ VA-6 (available from Ashland) was used as the source of the binder and is a mixture of an acetic acid ethenyl ester polymer and 1-ethenyl-2-pyrrolidinone.

[0053] Tween™ 20 (available from Croda) was used as the source of the surfactant and is polyoxyethylene (20) sorbitan monolaurate.

[0054] Polyethylene glycol 200 was used as the solvent and is available from Oxiteno, BASF, and Dow.

[0055] Applicant used GA₃ in the form of Technical Grade Active Ingredient ("TGA") when preparing formulations of the present invention. The percent GA₃ was about 96.6 % w/w. Variations in the amount of GA₃ in the TGA can be accounted for by decreasing or increasing the amount of solvent in order to produce the desired percent of GA₃ in the formulation. This is standard practice within the guidelines of U.S. Environmental Protection Agency per 40 C.F.R. § 158.175(b)(2).

Example 1

Preparation of formulations

[0056] A concentrated GA₃ formulation was prepared using the amounts indicated below in Table 1 and using the following process. The solvent was warmed in a 3000 milliliter glass beaker with a stir/hot plate. Then the GA₃ TGA and surfactant were added and mixed until solubilized. The binder was added and stirred until solubilized. Next, the dye was added and stirred until the formulation was homogenous. The formulation was allowed to cool and then packaged.

[0057] FD&C Red #40 was used as the source of the dye in this formulation.

Table 1

Material	% wt/wt	g/batch	Purpose
GA ₃ TGA (96.6% pure)	10.6	161	Active Ingredient
PEG 200	84.5	1280	Solvent
Polyoxyethylene (20) sorbitan monolaurate	1.0	15	Surfactant

(continued)

Material	% wt/wt	g/batch	Purpose
Copolymer of acetic acid ethenyl ester polymer and 1-ethenyl-2-pyrrolidinone	2.0	30	Binder
Dye	2.0	30	Color additive
Total	100	1515	

[0058] Accordingly, a 10.2% GA₃ formulation was prepared. Given the difficulty of formulating GA₃, it was unexpected that this concentrated formulation was homogeneous and stable.

Example 2

[0059] A concentrated GA₃ formulation was prepared using the amounts indicated below in Table 2 and using the following process. The solvent was warmed in a 3 liter glass beaker with a stir/hot plate. Then the GA₃ TGAI and surfactant were added and mixed until solubilized. The binder was added and stirred until solubilized. Next, the dye was added and stirred until the formulation was homogenous. The formulation was allowed to cool and then packaged.

[0060] FD&C Blue #1 was used as the source of the dye in this formulation.

Table 2

Material	% wt/wt	g/batch	Purpose
GA ₃ TGAI (96.6% pure)	10.6	107	Active Ingredient
PEG 200	84.5	853	Solvent
Polyoxyethylene (20) sorbitan monolaurate	1.0	10	Surfactant
Copolymer of acetic acid ethenyl ester polymer and 1-ethenyl-2-pyrrolidinone	2.0	20	Binder
Dye	2.0	20	Color additive
Total	100	1010	

[0061] Accordingly, a 10.2 % GA₃ formulation was prepared with a batch size of 1010 grams. As indicated above in Example 1, it was unexpected that this concentrated formulation was homogeneous and stable.

1. A concentrated agricultural solution formulation comprising:

from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of gibberellic acid (GA₃), gibberellin₄ (GA₄), gibberellin₇ (GA₇) and gibberellin_{4/7} (GA_{4/7});

from about 75 to about 90 % wt/wt of at least one solvent selected from the group consisting of polyethylene glycols with average molecular weights of from about 190 to about 420 daltons and C8 to C12 fatty acid dimethylamides;

from about 0.1 to about 4 % wt/wt of a binder; and

from about 0.1 to about 3 % wt/wt of a non-ionic surfactant.

2. The formulation of 1 wherein the gibberellin is GA₃.

3. The formulation of 1 comprising from about 9 to about 12 % wt/wt of the at least one gibberellin.

4. The formulation of 1 comprising from about 9.5 to about 11.5 % wt/wt of the at least one gibberellin.

5. The formulation of 1 further comprising from about 0.1 to about 4 % wt/wt of a color additive.

6. The formulation of 5 wherein the color additive is a dye selected from the group consisting of FD&C Blue #1, FD&C Blue #2, FD&C Green #3, FD&C Red #3, FD&C Red #40, FD&C Yellow #5, FD&C Yellow #6, and Citrus Red #2.

7. The formulation of 6 wherein the dye is selected from the group consisting of FD&C Blue #1 and FD&C Red #40.
8. The formulation of 1 wherein the binder is selected from the group consisting of alkylated vinyl pyrrolidone polymers, a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone, and a copolymer of polyvinylpyrrolidone and 1-eicododecene.
9. The formulation of 8 wherein the binder is a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone.
10. The formulation of 1 comprising from about 1 to about 3 % wt/wt of the binder.
11. The formulation of 10 comprising from about 1.5 to about 2.5 % wt/wt of the binder.
12. The formulation of 1 wherein the non-ionic surfactant is selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate and polyoxyethylene (20) sorbitan monooleate.
13. The formulation of 12 wherein the non-ionic surfactant is polyoxyethylene (20) sorbitan monolaurate.
14. The formulation of 1 comprising from about 0.1 to about 2 % wt/wt of the non-ionic surfactant.
15. The formulation of 14 comprising from about 0.5 to about 1.5 % wt/wt of the non-ionic surfactant.
16. A concentrated agricultural solution formulation comprising:
 - from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of gibberellic acid (GA₃), gibberellin₄ (GA₄), gibberellin₇ (GA₇) and gibberellin_{4/7} (GA_{4/7});
 - from about 75 to about 90 % wt/wt polyethylene glycols with average molecular weights of from about 190 to about 210 daltons;
 - from about 0.1 to about 4 % wt/wt of a binder selected from the group consisting of alkylated vinyl pyrrolidone polymers, a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone, and a copolymer of polyvinylpyrrolidone and 1-eicododecene; and
 - from about 0.1 to about 3 % wt/wt of a non-ionic surfactant selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate and polyoxyethylene (20) sorbitan monooleate.
17. The formulation of 16 wherein the gibberellin is GA₃.
18. A method for regulating plant growth comprising treating a plant or soil with an effective amount of the formulation of 1.
19. The method of 18 wherein the effective amount is from about 0.1 to about 50 grams of gibberellin per acre.
20. The method of 18 wherein the plant is rice.

Claims

1. A concentrated agricultural solution formulation comprising:
 - from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of gibberellic acid (GA₃), gibberellin₄ (GA₄), gibberellin₇ (GA₇) and gibberellin_{4/7} (GA_{4/7});
 - from about 75 to about 90 % wt/wt of at least one solvent selected from the group consisting of polyethylene glycols with average molecular weights of from about 190 to about 420 daltons and C8 to C12 fatty acid dimethylamides;
 - from about 0.1 to about 4 % wt/wt of a binder; and
 - from about 0.1 to about 3 % wt/wt of a non-ionic surfactant.
2. The formulation of claim 1 wherein the gibberellin is GA₃.
3. The formulation of claim 1 comprising from about 9 to about 12 % wt/wt of the at least one gibberellin, preferably

EP 3 766 348 A1

from about 9.5 to about 11.5 % wt/wt of the at least one gibberellin.

4. The formulation of claim 1 further comprising from about 0.1 to about 4 % wt/wt of a color additive.

5. The formulation of claim 4 wherein the color additive is a dye selected from the group consisting of FD&C Blue #1, FD&C Blue #2, FD&C Green #3, FD&C Red #3, FD&C Red #40, FD&C Yellow #5, FD&C Yellow #6, and Citrus Red #2, preferably wherein the dye is selected from the group consisting of FD&C Blue #1 and FD&C Red #40.

6. The formulation of claim 1 wherein the binder is selected from the group consisting of alkylated vinyl pyrrolidone polymers, a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone, and a copolymer of polyvinylpyrrolidone and 1-eicododecene.

7. The formulation of claim 6 wherein the binder is a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone.

8. The formulation of claim 1 comprising from about 1 to about 3 % wt/wt of the binder, preferably from about 1.5 to about 2.5 % wt/wt of the binder.

9. The formulation of claim 1 wherein the non-ionic surfactant is selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate and polyoxyethylene (20) sorbitan monooleate, preferably wherein the non-ionic surfactant is polyoxyethylene (20) sorbitan monolaurate.

10. The formulation of claim 1 comprising from about 0.1 to about 2 % wt/wt of the non-ionic surfactant, preferably from about 0.5 to about 1.5 % wt/wt of the non-ionic surfactant.

11. A concentrated agricultural solution formulation comprising:

from about 5 to about 15 % wt/wt of at least one gibberellin selected from the group consisting of gibberellic acid (GA₃), gibberellin₄ (GA₄), gibberellin₇ (GA₇) and gibberellin_{4/7} (GA_{4/7});

from about 75 to about 90 % wt/wt polyethylene glycols with average molecular weights of from about 190 to about 210 daltons;

from about 0.1 to about 4 % wt/wt of a binder selected from the group consisting of alkylated vinyl pyrrolidone polymers, a copolymer of acetic acid ethenyl ester and 1-ethenyl-2-pyrrolidinone, and a copolymer of polyvinylpyrrolidone and 1-eicododecene; and

from about 0.1 to about 3 % wt/wt of a non-ionic surfactant selected from the group consisting of polyoxyethylene (20) sorbitan monolaurate and polyoxyethylene (20) sorbitan monooleate.

12. The formulation of claim 11 wherein the gibberellin is GA₃.

13. A method for regulating plant growth comprising treating a plant or soil with an effective amount of the formulation of claim 1.

14. The method of claim 13 wherein the effective amount is from about 0.1 to about 50 grams of gibberellin per acre.

15. The method of claim 13 wherein the plant is rice.



EUROPEAN SEARCH REPORT

Application Number
EP 20 19 3002

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	WO 2015/095312 A1 (VALENT BIOSCIENCES CORP [US]) 25 June 2015 (2015-06-25)	1-15	INV. A01N43/12 A01N25/02 A01P21/00 A01N45/00
Y	* claims 1-20 * * paragraphs [0037] - [0045], [0057] * * examples 1-12 *	1-15	
Y	----- WO 02/082902 A2 (VALENT BIOSCIENCES CORP [US]; DEVISETTY BALA N [US]; BEACH ROBERT M [U] 24 October 2002 (2002-10-24) * claims 1-26 * * examples 1-3 *	1-15	
Y	----- US 2015/080216 A1 (WIKLEY PHIL [GB] ET AL) 19 March 2015 (2015-03-19) * paragraph [0071] *	1-15	
E	----- WO 2017/011562 A1 (STOLLER ETS [US]) 19 January 2017 (2017-01-19) * table 1 * * claims 1-8,13-21 * -----	1-15	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (IPC)
			A01N
Place of search		Date of completion of the search	Examiner
Munich		6 November 2020	Marie, Gérald
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

 1
EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 20 19 3002

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-11-2020

10

15

20

25

30

35

40

45

50

55

Patent document cited in search report			Publication date		Patent family member(s)		Publication date	
WO 2015095312	A1	25-06-2015	AU	2014364823	A1		16-06-2016	
			CL	2016001569	A1		20-01-2017	
			CR	20160281	A		03-10-2016	
			EP	3082421	A1		26-10-2016	
			MA	39192	A1		30-12-2016	
			MX	358473	B		21-08-2018	
			PE	20161250	A1		12-11-2016	
			US	2015173365	A1		25-06-2015	
			WO	2015095312	A1		25-06-2015	

WO 02082902	A2	24-10-2002	AT	345041	T		15-12-2006	
			AU	2002318111	B2		18-05-2006	
			AU	2006202391	A1		29-06-2006	
			BR	0208863	A		19-10-2004	
			CA	2444280	A1		24-10-2002	
			CL	2003002048	A1		14-01-2005	
			DE	60216066	T2		28-06-2007	
			EP	1377163	A2		07-01-2004	
			ES	2274981	T3		01-06-2007	
			HK	1062990	A1		10-12-2004	
			JP	4408630	B2		03-02-2010	
			JP	2004528322	A		16-09-2004	
			MX	PA03009284	A		16-03-2004	
			PT	1377163	E		31-01-2007	
			US	2003008949	A1		09-01-2003	
			US	2006003898	A1		05-01-2006	
			WO	02082902	A2		24-10-2002	
			ZA	200307904	B		14-09-2004	

US 2015080216	A1	19-03-2015	AU	2014318558	A1		31-03-2016	
			CA	2923886	A1		19-03-2015	
			CN	105960168	A		21-09-2016	
			EP	3043645	A1		20-07-2016	
			GB	2520147	A		13-05-2015	
			US	2015080216	A1		19-03-2015	
			WO	2015038917	A1		19-03-2015	
			ZA	201601699	B		30-05-2018	

WO 2017011562	A1	19-01-2017	AR	105347	A1		27-09-2017	
			AU	2016294422	A1		25-01-2018	
			CA	2991303	A1		19-01-2017	
			CL	2017003376	A1		25-05-2018	
			CN	107847431	A		27-03-2018	
			CO	2018001371	A2		30-04-2018	
			CR	20180028	A		10-04-2018	
			EC	SP18000533	A		31-03-2018	

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 20 19 3002

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-11-2020

10	Patent document cited in search report	Publication date	Patent family member(s)	Publication date
15			EP 3322403 A1	23-05-2018
			KR 20180030559 A	23-03-2018
			PE 20181154 A1	17-07-2018
			PH 12018500115 A1	23-07-2018
			TR 201800349 T1	23-07-2018
			TW 201713210 A	16-04-2017
			US 2017013836 A1	19-01-2017
			US 2019142000 A1	16-05-2019
20			WO 2017011562 A1	19-01-2017
			ZA 201800629 B	19-12-2018
25	-----			
30				
35				
40				
45				
50				

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 0252897 A [0011]
- US 4936901 A [0011]
- US 5622658 A [0011]
- US 6984609 B2 [0014]
- WO 9637494 A [0047]
- WO 9825923 A [0047]

Non-patent literature cited in the description

- CHEMICAL ABSTRACTS, 185982-80-3 [0047]
- CHEMICAL ABSTRACTS, 185984-60-5 [0047]