



(12) **CORRECTED EUROPEAN PATENT SPECIFICATION**

(15) Correction information:
Corrected version no 1 (W1 B1)
Corrections, see
Sequence listing
Remarks
Sequence listing replaced or added

(48) Corrigendum issued on:
09.08.2023 Bulletin 2023/32

(45) Date of publication and mention
of the grant of the patent:
14.06.2023 Bulletin 2023/24

(21) Application number: **19735528.2**

(22) Date of filing: **28.06.2019**

(51) International Patent Classification (IPC):
C07D 401/14 ^(2006.01) **C07D 405/14** ^(2006.01)
C07D 403/10 ^(2006.01) **C07D 405/10** ^(2006.01)
C07D 409/04 ^(2006.01) **C07D 409/14** ^(2006.01)
C07D 413/04 ^(2006.01) **C07D 417/14** ^(2006.01)
C07D 471/04 ^(2006.01) **C07D 487/04** ^(2006.01)
C07D 495/04 ^(2006.01) **C07D 209/18** ^(2006.01)
A61P 1/16 ^(2006.01) **A61K 31/404** ^(2006.01)

(52) Cooperative Patent Classification (CPC):
C07D 409/04; A61P 1/16; C07D 209/18;
C07D 401/14; C07D 403/10; C07D 405/10;
C07D 405/14; C07D 409/14; C07D 413/04;
C07D 417/14; C07D 471/04; C07D 487/04;
C07D 495/04

(86) International application number:
PCT/EP2019/067351

(87) International publication number:
WO 2020/002611 (02.01.2020 Gazette 2020/01)

(54) **NOVEL LXR MODULATORS WITH BICYCLIC CORE MOIETY**

NEUARTIGE LXR-MODULATOREN MIT BICYCLISCHEM KERNTEIL

NOUVEAUX MODULATEURS LXR AVEC UNE FRACTION DE NOYAU BICYCLIQUE

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR

(30) Priority: **28.06.2018 EP 18180450**

(43) Date of publication of application:
05.05.2021 Bulletin 2021/18

(73) Proprietor: **OrsoBio, Inc.**
Palo Alto, CA 94303 (US)

(72) Inventors:
• **GEGE, Christian**
89584 Ehingen (DE)
• **KINZEL, Olaf**
69117 Heidelberg (DE)
• **HAMBRUCH, Eva**
68239 Mannheim (DE)

• **BIRKEL, Manfred**
64342 Seeheim-Jugenheim (DE)
• **KREMOSER, Claus**
69121 Heidelberg (DE)
• **DEUSCHLE, Ulrich**
67346 Speyer (DE)

(74) Representative: **Grünecker Patent- und**
Rechtsanwälte
PartG mbB
Leopoldstraße 4
80802 München (DE)

(56) References cited:
WO-A1-2008/119657

Remarks:

The complete document including Reference
Table(s) and the Sequence Listing(s) can be
downloaded from the EPO website

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention relates to novel compounds which are Liver X Receptor (LXR) modulators and to pharmaceutical compositions containing same. The present invention further relates to the use of said compounds in the prophylaxis and/or treatment of diseases which are associated with the modulation of the Liver X Receptor.

Background:

[0002] The Liver X Receptors, LXR α (NR1H3) and LXR β (NR1H2) are members of the nuclear receptor protein superfamily. Both receptors form heterodimeric complexes with Retinoid X Receptor (RXR α , β or γ) and bind to LXR response elements (e.g. DR4-type elements) located in the promoter regions of LXR responsive genes. Both receptors are transcription factors that are physiologically regulated by binding ligands such as oxysterols or intermediates of the cholesterol biosynthetic pathways such as desmosterol. In the absence of a ligand, the LXR-RXR heterodimer is believed to remain bound to the DR4-type element in complex with co-repressors, such as NCOR1, resulting in repression of the corresponding target genes. Upon binding of an agonist ligand, either an endogenous one such as the oxysterols or steroid intermediates mentioned before or a synthetic, pharmacological ligand, the conformation of the heterodimeric complex is changed, leading to the release of corepressor proteins and to the recruitment of coactivator proteins such as NCOA1 (SRC1), resulting in transcriptional stimulation of the respective target genes. While LXR β is expressed in most tissues, LXR α is expressed more selectively in cells of the liver, the intestine, adipose tissue and macrophages. The relative expression of LXR α and LXR β at the mRNA or the protein level may vary between different tissues in the same species or between different species in a given tissue. The LXR's control reverse cholesterol transport, i.e. the mobilization of tissue-bound peripheral cholesterol into HDL and from there into bile and feces, through the transcriptional control of target genes such as ABCA1 and ABCG1 in macrophages and ABCG5 and ABCG8 in liver and intestine. This explains the anti-atherogenic activity of LXR agonists in dietary LDLR-KO mouse models. The LXRs, however, do also control the transcription of genes involved in lipogenesis (e.g. Srebp1c, Scd1, Fasn) which accounts for the liver steatosis observed following prolonged treatment with LXR agonists.

[0003] The liver steatosis liability is considered a main barrier for the development of non-selective LXR agonists for atherosclerosis treatment.

[0004] Non-alcoholic fatty liver disease (NAFLD) is regarded as a manifestation of metabolic syndrome in the liver and NAFLD has reached epidemic prevalences worldwide (Estes et al., *Hepatology* 2018;67:123; Estes et al., *J. Hepatol.* 2018;69:896). The pathologies of NAFLD range from benign and reversible steatosis to steatohepatitis (nonalcoholic steatohepatitis, NASH) that can develop towards fibrosis, cirrhosis and potentially further towards hepatocellular carcinogenesis. Classically, a two-step model has been employed to describe the progression of NAFLD into NASH, with hepatic steatosis as an initiating first step sensitizing towards secondary signals (exogenous or endogenous) that lead to inflammation and hepatic damage (Day et al., *Gastroenterology* 1998;114:842). Nowadays, the transition from benign NAFLD towards the more aggressive state NASH is regarded as multifactorial with genetic, environmental, lifestyle and nutritional influences playing different roles in different individual setups. Independent from the etiology of the disease there is a very strong unmet medical need to stop progression of NAFLD because of the detrimental sequelae such as liver cirrhosis, hepatocellular carcinoma or other forms of liver related modalities.

[0005] LXR expression levels are firmly associated with the state of NAFLD. Notably, LXR expression was shown to correlate with the degree of fat deposition, as well as with hepatic inflammation and fibrosis in NAFLD patients (Ahn et al., *Dig. Dis. Sci.* 2014;59:2975). Furthermore, serum and liver desmosterol levels are increased in patients with NASH but not in people with simple liver steatosis. Desmosterol has been characterized as a potent endogenous LXR agonist (Yang et al., *J. Biol. Chem.* 2006;281:27816). Given the known involvement of the LXRs as master regulators of hepatic lipidogenesis and lipid metabolism, in general and the aforementioned association of LXR expression levels with the stage of fatty liver disease, NAFLD/NASH patients might therefore benefit from blocking the increased LXR activity in the livers of these patients through small molecule antagonists or inverse agonists that shut off LXRs' activity. While doing so it needs to be taken care that such LXR antagonists or inverse agonists do not interfere with LXRs in peripheral tissues or macrophages to avoid disruption of the anti-atherosclerotic reverse cholesterol transport governed by LXR in these tissues or cells.

[0006] Certain publications (e.g. Feet et al., *Cell* 1998;93:693 and Schultz et al., *Genes Dev.* 2000;14:2831) have highlighted the role of LXR α , in particular, for the stimulation of lipidogenesis and hence establishment of NAFLD in the liver. They indicate that it is mainly LXR α being responsible for the hepatic steatosis, hence an LXR α -specific antagonist or inverse agonist might suffice or be desirable to treat just hepatic steatosis. These data, however, were generated only by comparing LXR α , LXR β or double knockout with wild-type mice with regards to their susceptibility to develop steatosis on a high fat diet. They do not account for a major difference in the relative expression levels of LXR α and LXR β in the human as opposed to the murine liver. Whereas LXR α is the predominant LXR subtype in the rodent liver, LXR β is expressed to about the same if not higher levels in the human liver compared to LXR α (data from Unigene or

other expression databases). This was exemplified by testing an LXR β selective agonist in human phase I clinical studies (Kirchgessner et al., Cell Metab. 2016;24:223) which resulted in the induction of strong hepatic steatosis although it was shown to not activate human LXR α .


[0007] Hence it can be assumed that it should be desirable to have no strong preference of an LXR modulator designed to treat NAFLD or NASH for a particular LXR subtype. A certain degree of LXR-subtype selectivity might be allowed if the pharmacokinetic profile of such a compound clearly ensures sufficient liver exposure and resident time to cover both LXRs in clinical use.

[0008] In summary, the treatment of diseases such as NAFLD or NASH would need LXR modulators that block LXRs in a hepato-selective fashion and this could be achieved through hepatotropic pharmacokinetic and tissue distribution properties that have to be built into such LXR modulators.

[0009] The master control on lipidogenesis is exerted by LXRs in all major cell types studied so far. Cancer cells are also highly dependent on *de novo* lipidogenesis and therefore Flaveny et al. tested the LXR inverse agonist tool compound SR9243 in cancer cells and in animal cancer models (Cancer Cell 2015;28:42). They could show that **SR9243** inhibited lipidogenesis along with the Warburg glycolysis effect, in general, and that this molecular effect led to apoptosis and diminished tumor growth *in vivo*.


Prior Art

[0010] Zuercher et al. describes with the structurally unrelated tertiary sulfonamide GSK2033 the first potent, cell-active LXR antagonists (J. Med. Chem. 2010;53:3412). Later, this compound was reported to display a significant degree of promiscuity, targeting a number of other nuclear receptors (Griffett & Burris, Biochem. Biophys. Res. Commun. 2016;479:424). It is stated, that **GSK2033** showed rapid clearance ($Cl_{int} > 1.0$ mL/min/mg protein) in rat and human liver microsomal assays and that this rapid hepatic metabolism of **GSK2033** precludes its use *in vivo*. As such **GSK2033** is a useful chemical probe for LXR in cellular studies only.



GSK2033

[0011] WO2014/085453 describes the preparation of structurally unrelated small molecule LXR inverse agonists of Formula (A) in addition to structure **GSK2033** above:




[0012] The following compounds from this application, in particular, are further described in some publications, mainly from the same group of inventors/authors: **SR9238** is described as a liver-selective LXR inverse agonist that suppresses hepatic steatosis upon parenteral administration (Griffett et al., ACS Chem. Biol. 2013;8:559). After ester saponification of **SR9238** the LXR inactive acid derivative **SR10389** is formed. This compound then has systemic exposure. In addition, it was described, that **SR9238** suppresses fibrosis in a model of NASH again after parenteral administration (Griffett et

al., Mol. Metab. 2015;4:35). With related **SR9243** the effects on aerobic glycolysis (Warburg effect) and lipogenesis were described (Flaveny et al., Cancer Cell 2015;28:42) and the NASH-suppressing data obtained with **SR9238** was confirmed by Huang et al. (BioMed Res. Int. 2018;8071093) using **SR9243**.

[0013] WO2003/082802 describes structurally unrelated LXR agonists of general Formula (B):

5



10

15


[0014] In all examples the acid containing (hetero)aryl moiety is linked via an oxygen atom to the rest of the molecule. Most interesting examples are **GW3965** (Collins et al. J. Med. Chem. 2002;45:1963) and clinical candidate **RGX-104** from Rgenix.

20

[0015] Yu et al. (J. Org. Chem. 2018;83:323) describes the synthesis of 2,3-disubstituted indoles via the following reaction scheme. The only example with an ortho-substituted aryl in 3-position of the indole is structure **C1**.

25

30




35

[0016] WO2016/207217 discloses bicyclic derivatives of Formula (D), which does not fall within the scope of the present invention, since no $-SO_2-$ linked residue is possible for A, which may represent a bicyclic structure including indole. However intermediate **D1** is disclosed (Example 69, Step E), which is the only example with an ortho-substituted aryl in 3-position of the indole.

40


45



50


[0017] WO2016/106266 discloses azaindoles of Formula (E) as TGF β antagonists

55




wherein R is an optionally substituted heterocyclic or heterobicyclic group, R⁸ is selected from a broad range of substituents including -SO₂R⁹. Residue R⁹ can be selected from a broad range of substituents including C₃-C₈-cycloalkyl and heterocycloalkyl. The only examples wherein both the 2- and 3-position of the azaindole is substituted with a cyclic moiety is structure **E1** and **E2**.

[0018] WO2013/111150 discloses adamantane derivatives of Formula (F) as 17β-hydroxysteroid dehydrogenase type 1 inhibitors




wherein Ar is an optionally substituted C₁-C₁₈-heteroaryl group, A can be -SO₂- and B can be absent. No examples are shown, which fall within the scope of the present invention.

[0019] WO2013/028999 discloses structures of Formula (G) as potential therapeutics for neuropsychiatric disorders




wherein Z may be absent and A represents a ring structure, e.g. 3-substituted indole of Formula (G1). Here R¹⁵ and R¹⁷ can be selected from an optionally substituted aryl and heteroaryl moiety. However for this case, no examples are shown.

[0020] WO2013/012649 discloses azaindoles of Formula (H) for the treatment of HIV




wherein linker element L can be -SO₂-, R⁵ and R⁶ can independently be selected from a broad range of substituents including an optionally substituted cycloalkyl, heterocycloalkyl, aryl and heteroaryl. In most cases, R² is a carboxylic acid or bioisostere thereof. No examples are shown, which fall within the scope of the present invention.

[0021] WO2010/124793 and WO2008/132434 disclose azaindoles of Formula (J) as fungicides




wherein R^4 can be selected from a broad range of substituents including an optionally substituted cyclyl, heterocyclyl, aryl and heteroaryl. R^3 can be selected from a broad range of substituents including $-SO_2R^{12}$, with R^{12} again can be selected from a broad range of substituents including an optionally substituted cyclyl, heterocyclyl, aryl and heteroaryl. The only example wherein both the 2- and 3-position of the azaindole is substituted with a cyclic moiety is structure **J1**.

[0022] WO2010/010186 discloses JAK kinase inhibitors of Formula (**K**)




wherein ring Cy is selected from aryl and heteroaryl. With L^1 equals SO_2 , n equals 0, R^{3a} e.g. unsubstituted cycloalkyl, heterocycloalkyl, aryl or heteroaryl and R^{3b} selected from optionally substituted cycloalkyl, heterocycloalkyl, aryl or heteroaryl derivatives falling in the scope of the present invention can be constructed, however no examples are shown.

[0023] WO2009/032116 discloses indoles of Formula (**L**) for treating viral infections




wherein R^1 can be selected from a broad range of substituents including $-SO_2-$. For R^2 the cyclic moieties (**L1** to **L3**) and for R^3 the cyclic moieties (**L4** and **L5**) are possible. In related application WO2009/032125 and WO2009/064848 even more cyclic moieties for R^3 are possible. R^{10} can be selected from optionally substituted cycloalkyl, cycloalkenyl, heterocycloalkyl, heterocycloalkenyl, aryl and heteroaryl. In WO2009/064852, a cyclic moiety of structure **L6** is possible for R^3 . In all applications, no examples are shown, which fall within the scope of the present invention.

[0024] WO2008/116833 discloses azetidine compounds of Formula (**M**) for treating disorders that respond to modulation of the serotonin 5-hydroxytryptamine-6 (5-HT₆) receptor




wherein X^1 and X^2 are independently N or CR^x . Residue R^x can be selected from a broad range of substituents including an optionally substituted phenyl or C_{3-6} -cycloalkyl. Residue A can be selected from optionally substituted C_{3-6} -cycloalkyl, aryl or heteroaryl. No examples, wherein both the 2- and 3-position of the indole is directly substituted by a cyclic moiety, are disclosed.

[0025] WO2008/003736 discloses azaindoles of Formula (**N**)




wherein R^2 and R^3 can independently comprise a saturated nitrogen-containing heterocyclic moiety (e.g. piperidine) while m can be 0. According to claim 8, Q can represent the protecting group $-\text{SO}_2\text{-Ph}$. No examples, wherein both the 2- and 3-position of the indole is directly substituted by a cyclic moiety, are disclosed.

[0026] WO2007/075555 discloses CB_1 antagonists of Formula (P)




wherein R^1 can be selected from a broad range of substituents including a substituted indole while X can represent a bond. The only example where a cyclic moiety is linked to the 3-position of the indole is structure **P1**. More specifically, indole derivatives of Formula (P) are described in WO2004/000831 as histamine H_3 antagonist, again with structure **P1** as example.

[0027] WO2007/134169 and WO2006/050236 disclose indole derivatives of Formula (Q) as inhibitors of $\text{TNF-}\alpha$ production




wherein X can be SO_2 , Y can be selected from a broad range of substituents including cycloalkyl, heterocycloalkyl, aryl and heterocycle while Z has to be selected from $-\text{B(OR)}_2$, $-\text{CONROR}$ and $-\text{N(OR)COR}$ (with $R = \text{H}$ or alkyl). R^3 and R^8 can be independently selected from a broad range of substituents including cycloalkyl and a 5- or 6-membered organic ring. No examples, wherein both the 2- and 3-position of the indole is substituted by a cyclic moiety, are disclosed.

[0028] WO2005/034941 discloses bicyclic structures of Formula (R) as inhibitors for hepatitis C virus polymerase




wherein Ar^1 and Ar are 5- to 10-membered aromatic rings, A^1 can be a cycloalkyl (optionally substituted with alkoxy) and n can be 0. The closest examples to the present invention are structure **R1** and **R2**.

[0029] WO2005/14000 discloses indoles of Formula (S) for the treatment of 5-HT_6 -receptor-related diseases such as obesity and CNS disorders




wherein R¹ represents a nitrogen-attached saturated or unsaturated heterocyclic ring system, R² can be selected from a broad range of substituents including a saturated or unsaturated cycloalkyl, n is selected from 0 to 4 and residue A and B form a saturated or unsaturated cycloalkyl ring. No examples, wherein both the 2- and 3-position of the indole is directly substituted (i.e. n = 0) by a cyclic moiety, are disclosed.

[0030] WO2002/51837 and WO2002/36562 disclose bicyclic structures of Formulae (T) and (T1), respectively, for the treatment of 5-HT₆-receptor-related diseases




wherein X and Y can independently represent a carbon atom, which is optionally substituted with an aryl or heteroaryl moiety, R⁸ may also represent an optionally substituted aryl or heteroaryl moiety. The cyclic moiety on the left-hand-side of Formula (T1) is usually piperazine. No examples, wherein both the 2- and 3-position of the (aza)indole is substituted by a cyclic moiety (e.g. aryl or heteroaryl), are disclosed. The closest example is structure T1.

[0031] WO2002/32863 discloses indoles of Formula (U) for the treatment of 5-HT₆-receptor-related diseases




wherein Ar can be selected from optionally substituted phenyl, naphthyl or 5- to 10-membered mono- or bicyclic heterocyclic moieties, R² can be an unsubstituted phenyl and R³ is selected from hydrogen or 3-(1-azabicyclo[2.2.2]oct-2-en)yl. However no example with suitable substitution at 2- and 3-position of the indole is shown - the closest examples are structure U1 and U2.

[0032] WO9921851 discloses structures of Formula (V) as agricultural or horticultural fungicides




wherein A can be selected from a very broad range of cyclic systems including optionally substituted indole. However no example with suitable substitution at 2- and 3-position of the indole is shown; the closest example is structure **V1**.

[0033] WO9857931 and WO9822452 disclose bicyclic structures of Formulae (**W**) and (**W1**), respectively, as anti-microbial agents




wherein R^2 can be selected from a very broad range of residues including aryl and heteroaryl; and Y can represent NR, with R selected from a very broad range of residues including a arylsulfonyl moiety. No example with substitution at 2- and 3-position of the indole is shown; the closest example is structure **W1**.

[0034] WO9822457 discloses bicyclic structures of Formula (**X**) as anti-inflammatory agents




wherein R^{10} can be selected from a very broad range of substituents including SO_2R^{30} ; and wherein R^{11} , R^{12} , R^{30} can be selected from optionally substituted aryl and heteroaryl. However no example is shown, where R^{10} has indeed a SO_2 -connected moiety.

[0035] WO2001130343, WO2000/46199, WO2000/46197, WO2000/46195, JP06145150, EP0535926, EP0535925 describe indole derivatives, where in 2-position of the indole moiety a 1H- or 2H-tetrazol-5-yl moiety can be attached as only possible cyclic moiety, which functions as a carboxylic acid bioisostere. The only example with such a directly connected tetrazole moiety is disclosed in JP06145150 (Structure **Y1**).




[0036] WO2008/119657 describes imidazolidinone derivatives of Formula (**Z**) binding to LXR with representative example (**Z1**):



[0037] The following four structures were found to be weak binder on another nuclear receptor target and therefore


were mentioned as initial hits in a confidential collaboration with another pharma company:



Summary of the invention

[0038] The subject matter of the invention is as set out in the appended claims.

[0039] The present invention relates to compounds according to Formula (I)



a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein cycle A, B, C, D and residue L and R¹ are defined as in claim 1.

[0040] The compounds of the present invention have a similar or better LXR inverse agonistic activity compared to the known LXR inverse agonists. Furthermore, the compounds of the present invention exhibit an advantageous liver/blood-ratio after oral administration so that disruption of the anti-atherosclerotic reverse cholesterol transport governed by LXR in peripheral macrophages can be avoided. The incorporation of an acidic moiety (or a bioisoster thereof) can improve additional parameters, e.g. microsomal stability, solubility and lipophilicity.


[0041] Thus, the present invention further relates to a pharmaceutical composition comprising a compound according to Formula (I) and at least one pharmaceutically acceptable carrier or excipient.

[0042] The present invention is further directed to compounds according to Formula (I) for use in the prophylaxis and/or treatment of diseases mediated by LXRs.

[0043] Accordingly, the present invention relates to the prophylaxis and/or treatment of non-alcoholic fatty liver disease, non-alcoholic steatohepatitis, liver inflammation, liver fibrosis, obesity, insulin resistance, type II diabetes, familial hypercholesterolemia, hypercholesterolemia in nephrotic syndrome, metabolic syndrome, cardiac steatosis, cancer, viral myocarditis and hepatitis C virus infection.

Detailed description of the invention

[0044] The desired properties of a LXR modulator in conjunction with hepatoselectivity, can be yielded with compounds that follow the structural pattern represented by Formula (I)



a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein

A

is an annelated 5- to 6-membered cycle forming a 6-membered aryl or a 5- to 6-membered heteroaryl containing 1 to 3 heteroatoms independently selected from N, O and S, wherein this cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, C₁₋₆-alkyl, oxo, C₀₋₆-alkylene-OR¹¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkyleneS(O)_nR¹¹, C₀₋₆-alkylene-NR¹¹S(O)₂R¹¹, C₀₋₆-alkylene-S(O)₂NR¹¹R¹², C₀₋₆-alkylene-NR¹¹S(O)₂NR¹¹R¹², C₀₋₆-alkylene-CO₂R¹¹, O-C₁₋₆-alkylene-CO₂R¹¹, C₀₋₆-alkylene-O-COR¹¹, C₀₋₆-alkylene-CONR¹¹R¹², C₀₋₆-alkylene-NR¹¹-COR¹¹, C₀₋₆-alkylene-NR¹¹-CONR¹¹R¹², C₀₋₆-alkylene-O-CONR¹¹R¹², C₀₋₆-alkylene-NR¹¹-CO₂R¹¹ and C₀₋₆-alkylene-NR¹¹R¹², wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and

wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein the new formed cycle is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

B

is selected from the group consisting of 3- to 10-membered cycloalkyl, 3- to 10-membered heterocycloalkyl containing 1 to 3 heteroatoms independently selected from N, O and S, 6- to 14-membered aryl and 5- to 14-membered heteroaryl containing 1 to 4 heteroatoms independently selected from N, O and S,

wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR²¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR²¹, C₀₋₆-alkylene-NR²¹S(O)₂R²¹, C₀₋₆-alkylene-S(O)₂NR²¹R²², C₀₋₆-alkylene-NR²¹S(O)₂NR²¹R²², C₀₋₆-alkylene-CO₂R²¹, O-C₁₋₆-alkylene-CO₂R²¹, C₀₋₆-alkylene-O-COR²¹, C₀₋₆-alkylene-CONR²¹R²², C₀₋₆-alkylene-NR²¹-COR²¹, C₀₋₆-alkylene-NR²¹-CONR²¹R²², C₀₋₆-alkylene-O-CONR²¹R²², C₀₋₆-alkylene-NR²¹-CO₂R²¹ and C₀₋₆-alkylene-NR²¹R²², wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and wherein optionally two adjacent substituents on the cycloalkyl or heterocycloalkyl moiety form a 5- to 6-membered unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;



5

is selected from the group consisting of 6- or 10-membered aryl and 5- to 10-membered heteroaryl containing 1 to 3 heteroatoms independently selected from N, O and S,

10 wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR³¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-(6-membered aryl), C₀₋₆-alkylene-(5- to 6-membered heteroaryl), C₀₋₆-alkylene-S(O)_nR³¹, C₀₋₆-alkylene-NR³¹S(O)₂R³¹, C₀₋₆-alkylene-S(O)₂NR³¹R³², C₀₋₆-alkylene-NR³¹S(O)₂NR³¹R³², C₀₋₆-alkylene-CO₂R³¹, O-C₁₋₆-alkylene-CO₂R³¹, C₀₋₆-alkylene-O-COR³¹, C₀₋₆-alkylene-CONR³¹R³², C₀₋₆-alkylene-NR³¹-COR³¹, C₀₋₆-alkylene-NR³¹-CONR³¹R³², C₀₋₆-alkylene-O-CONR³¹R³², C₀₋₆-alkylene-NR³¹-CO₂R³¹ and C₀₋₆-alkylene-NR³¹R³²,

15 wherein alkyl, alkylene, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

20 and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5-to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;



30

is selected from the group consisting of 3- to 10-membered cycloalkyl, 3- to 10-membered heterocycloalkyl containing 1 to 3 heteroatoms independently selected from N, O and S, 6- to 14-membered aryl and 5- to 14-membered heteroaryl containing 1 to 4 heteroatoms independently selected from N, O and S,

35

wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR²¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR²¹, C₀₋₆-alkylene-NR²¹S(O)₂R²¹, C₀₋₆-alkylene-S(O)₂NR²¹R²², C₀₋₆-alkylene-NR²¹S(O)₂NR²¹R²², C₀₋₆-alkylene-CR⁴¹(=N-OR⁴¹), C₀₋₆-alkylene-CO₂R²¹, O-C₁₋₆-alkylene-CO₂R²¹, C₀₋₆-alkylene-O-COR²¹, C₀₋₆-alkylene-CONR²¹R²², C₀₋₆-alkylene-NR²¹-COR²¹, C₀₋₆-alkylene-NR²¹-CONR²¹R²², C₀₋₆-alkylene-O-CONR²¹R²², C₀₋₆-alkylene-NR²¹-CO₂R²¹ and C₀₋₆-alkylene-NR²¹R²²,

40 wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, CO-OC₁₋₄-alkyl, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

45 and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5-to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

50 and wherein optionally two adjacent substituents on the cycloalkyl or heterocycloalkyl moiety form a 5- to 6-membered unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

55

wherein



has a substituent from above in 1,2-orientation regarding to the connection towards



or has an annelated additional cycle in 1,2-orientation;

L is selected from the group consisting of a bond, C₁₋₆-alkylene, C₂₋₆-alkenylene, C₂₋₆-alkynylene, 3- to 10-membered cycloalkylene, 3- to 10-membered heterocycloalkylene containing 1 to 4 heteroatoms independently selected from N, O and S, 6- or 10-membered arylene and 5- to 10-membered heteroarylene containing 1 to 4 heteroatoms independently selected from N, O and S,

wherein alkylene, alkenylene, alkynylene, cycloalkylene, heterocycloalkylene, arylene and heteroarylene are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR⁴¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR⁴¹, C₀₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, C₀₋₆-alkyleneS(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-CO₂R⁴¹, O-C₁₋₆-alkylene-CO₂R⁴¹, C₀₋₆-alkylene-O-COR⁴¹, C₀₋₆-alkylene-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-COR⁴¹, C₀₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-alkylene-O-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-CO₂R⁴¹ and C₀₋₆-alkylene-NR⁴¹R⁴², wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the arylene and heteroarylene moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

R¹ is selected from the group consisting of H, halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR⁴¹, Y-C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), Y-C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), Y-C₀₋₆-alkylene-(6-membered aryl), Y-C₀₋₆-alkylene-(5- to 6-membered heteroaryl), C₀₋₆-alkylene-S(=O)(-R⁴¹)=N-R⁷⁵, X-C₁₋₆-alkylene-S(=O)(-R⁴¹)=N-R⁷⁵, C₀₋₆-alkylene-S(O)_nR⁴¹, X-C₁₋₆-alkylene-S(O)_nR⁴¹, C₀₋₆-alkylene-S(=NR⁷¹)R⁴¹, X-C₁₋₆-alkylene-S(=NR⁷¹)R⁴¹, C₀₋₆-alkylene-S(O)(=NR⁷¹)R⁴¹, X-C₁₋₆-alkylene-S(O)(=NR⁷¹)R⁴¹, C₀₋₆-alkylene-S(=NR⁷¹)₂R⁴¹, X-C₁₋₆-alkylene-S(=NR⁷¹)₂R⁴¹, C₀₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, X-C₁₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, Co-s-alkyleneS(O)₂NR⁴¹R⁴², X-C₁₋₆-alkylene-S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-SO₃R⁴¹, X-C₁₋₆-alkylene-SO₃R⁴¹, C₀₋₆-alkylene-CO₂R⁴¹, X-C₁₋₆-alkylene-CO₂R⁴¹, C₀₋₆-alkylene-O-COR⁴¹, X-C₁₋₆-alkylene-O-COR⁴¹, C₀₋₆-alkylene-CONR⁴¹R⁴², X-C₁₋₆-alkylene-CONR⁴¹R⁴², C₀₋₆-alkylene-CONR⁴¹OR⁴¹, X-C₁₋₆-alkylene-CONR⁴¹OR⁴¹, C₀₋₆-alkylene-CONR⁴¹SO₂R⁴¹, X-C₁₋₆-alkylene-CONR⁴¹SO₂R⁴¹, C₀₋₆-alkylene-NR⁴¹-COR⁴¹, X-C₁₋₆-alkylene-NR⁴¹-COR⁴¹, C₀₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-alkylene-O-CONR⁴¹R⁴², X-C₁₋₆-alkylene-O-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-CO₂R⁴¹, X-C₁₋₆-alkylene-NR⁴¹-CO₂R⁴¹, C₀₋₆-alkylene-NR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹R⁴²,

wherein alkyl, alkylene, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the aryl and heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl,

O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

R¹¹, R¹², R²¹, R²², R³¹, R³², R⁴¹, R⁴², R⁵¹ are independently selected from H and C₁₋₄-alkyl, wherein alkyl is unsubstituted or substituted with 1 to 3 substituent independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

or R¹¹ and R¹², R²¹ and R²², R³¹ and R³², R⁴¹ and R⁴², respectively, when taken together with the nitrogen to which they are attached complete a 3- to 6-membered ring containing carbon atoms and optionally containing 1 or 2 heteroatoms independently selected from O, S or N; and

wherein the new formed cycle is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

R⁷¹ is independently selected from H, CN; NO₂, C₁₋₄-alkyl and C(O)-OC₁₋₄-alkyl, wherein alkyl is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

R⁷⁵ is independently selected from C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, 3- to 6-membered heterocycloalkyl, 6-membered aryl and 5- to 6-membered heteroaryl,


wherein alkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, Me, Et, CHF₂, CF₃, OH, oxo, CO₂H, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, OMe, OEt, OCHF₂, and OCF₃;

X is independently selected from O, NR⁵¹, S(O)_n, S(=NR⁷¹), S(O)(=NR⁷¹) and S(=NR⁷¹)₂;

Y is independently selected from a bond, O, NR⁵¹, S(O)_n, S(=NR⁷¹), S(O)(=NR⁷¹) and S(=NR⁷¹)₂;

n is independently selected from 0 to 2;

and with the proviso, that the following structures are excluded:



[0045] In a preferred embodiment in combination with any of the above or below embodiments



is an annelated 5- to 6-membered cycle forming a 6-membered aryl or a 5- to 6-membered heteroaryl containing 1 to 3 heteroatoms independently selected from N, O and S, wherein this cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, C₁₋₆-alkyl, oxo, C₀₋₆-alkylene-OR¹¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkyleneS(O)_nR¹¹, C₀₋₆-alkylene-NR¹¹S(O)₂R¹¹, C₀₋₆-alkylene-S(O)₂NR¹¹R¹², C₀₋₆-alkylene-NR¹¹S(O)₂NR¹¹R¹², C₀₋₆-alkylene-CO₂R¹¹, O-C₁₋₆-alkylene-CO₂R¹¹, C₀₋₆-alkylene-O-COR¹¹, C₀₋₆-alkylene-

CONR¹¹R¹², C₀₋₆-alkylene-NR¹¹-COR¹¹, C₀₋₆-alkylene-NR¹¹-CONR¹¹R¹², C₀₋₆-alkylene-O-CONR¹¹R¹²,
C₀₋₆-alkylene-NR¹¹-CO₂R¹¹ and C₀₋₆-alkylene-NR¹¹R¹²,

wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and

wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein the new formed cycle is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl.


[0046] In a more preferred embodiment in combination with any of the above or below embodiments




is an annelated phenyl, thiophenyl, thiazolyl, pyridyl, pyrimidinyl, pyridazinyl and pyrazinyl, wherein this cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, C₁₋₆-alkyl, oxo, C₀₋₆-alkylene-OR¹¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR¹¹, C₀₋₆-alkylene-NR¹¹S(O)₂R¹¹, C₀₋₆-alkylene-S(O)₂NR¹¹R¹², C₀₋₆-alkylene-NR¹¹S(O)₂NR¹¹R¹², C₀₋₆-alkylene-CO₂R¹¹, O-C₁₋₆-alkylene-CO₂R¹¹, C₀₋₆-alkylene-O-COR¹¹, C₀₋₆-alkylene-CONR¹¹R¹², C₀₋₆-alkylene-NR¹¹-COR¹¹, C₀₋₆-alkylene-NR¹¹-CONR¹¹R¹², C₀₋₆-alkylene-O-CONR¹¹R¹², C₀₋₆-alkylene-NR¹¹-CO₂R¹¹ and C₀₋₆-alkylene-NR¹¹R¹²,

wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl.

[0047] In an even more preferred embodiment in combination with any of the above or below embodiments



is selected from




wherein



is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of F, Cl, Br, CN, OH, oxo, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl, O-halo-C₁₋₄-alkyl, NH₂, NHC₁₋₄-alkyl, N(C₁₋₄-alkyl)₂, SO₂-C₁₋₄-alkyl and SO₂-halo-C₁₋₄-alkyl.

[0048] In a most preferred embodiment in combination with any of the above or below embodiments



wherein



is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of F, Cl, Br, CN, Me, Et, CF₃, CHF₂, OH, OMe, OCF₃ and OCHF₃.

[0049] In a preferred embodiment in combination with any of the above or below embodiments



is selected from the group consisting of 3- to 10-membered cycloalkyl, 3- to 10-membered heterocycloalkyl containing 1 to 3 heteroatoms independently selected from N, O and S, 6- to 14-membered aryl and 5- to 14-membered heteroaryl containing 1 to 4 heteroatoms independently selected from N, O and S,

wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR²¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR²¹, C₀₋₆-alkylene-NR²¹S(O)₂R²¹, C₀₋₆-alkylene-S(O)₂NR²¹R²², C₀₋₆-alkylene-NR²¹S(O)₂NR²¹R²², C₀₋₆-alkylene-CO₂R²¹, O-C₁₋₆-alkylene-CO₂R²¹, C₀₋₆-alkylene-O-COR²¹, C₀₋₆-alkylene-CONR²¹R²², C₀₋₆-alkylene-NR²¹-COR²¹, C₀₋₆-alkylene-NR²¹-CONR²¹R²², C₀₋₆-alkylene-O-CONR²¹R²², C₀₋₆-alkylene-NR²¹-CO₂R²¹ and C₀₋₆-alkylene-NR²¹R²²,

wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the cycloalkyl or heterocycloalkyl moiety form a 5- to 6-membered unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N,

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl.

[0050] In a more preferred embodiment in combination with any of the above or below embodiments



is selected from the group consisting of phenyl, pyridyl and thiophenyl,

wherein phenyl, pyridyl and thiophenyl are substituted with 1 to 4 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR²¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR²¹, C₀₋₆-alkylene-NR²¹S(O)₂R²¹, C₀₋₆-alkylene-S(O)₂NR²¹R²², C₀₋₆-alkylene-NR²¹S(O)₂NR²¹R²², C₀₋₆-alkylene-CO₂R²¹, O-C₁₋₆-alkylene-CO₂R²¹,

C_{0-6} -alkylene-O-COR²¹, C_{0-6} -alkylene-CONR²¹R²², C_{0-6} -alkylene-NR²¹-COR²¹, C_{0-6} -alkylene-NR²¹-CONR²¹R²²,
 C_{0-6} -alkylene-O-CONR²¹R²², C_{0-6} -alkylene-NR²¹-CO₂R²¹ and C_{0-6} -alkylene-NR²¹R²²,
 wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents inde-
 5 pendentlly selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H,
 C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the phenyl and pyridyl moiety form a 5- to 8-membered partially
 unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and
 wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from
 10 halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-
 C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl.

[0051] In a similar more preferred embodiment in combination with any of the above or below embodiments

15 **B**

is selected from the group consisting of phenyl, naphthyl, pyridyl, pyrimidinyl, thiophenyl, thiazolyl, cyclopentyl,
 20 cyclohexyl, bicyclo[1.1.1]pentyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octyl and
 piperidinyl,

wherein the cycle is unsubstituted or substituted with 1 to 3 substituents independently selected from the group
 consisting of F, Cl, Br, CN, OH, oxo, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl, O-halo-C₁₋₄-alkyl, C₁₋₄-alkyl-OH and
 25 halo-C₁₋₄-alkyl-OH; and wherein optionally two adjacent substituents on the phenyl ring form together a -(CH₂)₃-,
 -(CH₂)₄-, -OCF₂O- and -OCH₂O- group.

[0052] In an even more preferred embodiment in combination with any of the above or below embodiments

30 **B**

is selected from the group consisting of phenyl and pyridyl,

wherein phenyl and pyridyl is substituted with 1 to 2 substituents independently selected from the group consisting
 35 of F, Cl, CN, CF₃, CH₂F and CHF₂.

[0053] In a most preferred embodiment in combination with any of the above or below embodiments

40 **B**

is 4-difluoromethylphenyl.

[0054] In a preferred embodiment in combination with any of the above or below embodiments

50 **C**

is selected from the group consisting of 6- or 10-membered aryl and 5- to 10-membered heteroaryl containing 1 to
 3 heteroatoms independently selected from N, O and S,

wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 4 substituents
 55 independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C_{0-6} -alkylene-OR³¹,
 C_{0-6} -alkylene-(3- to 6-membered cycloalkyl), C_{0-6} -alkylene-(3- to 6-membered heterocycloalkyl), C_{0-6} -alkylene-(6-
 membered aryl), C_{0-6} -alkylene-(5- to 6-membered heteroaryl), C_{0-6} -alkylene-S(O)_nR³¹, C_{0-6} -alkylene-
 NR³¹S(O)₂R³¹, C_{0-6} -alkylene-S(O)₂NR³¹R³², C_{0-6} -alkylene-NR³¹S(O)₂NR³¹R³², C_{0-6} -alkylene-CO₂R³¹, O-

C_{1-6} -alkylene- CO_2R^{31} , C_{0-6} -alkylene-O-COR³¹, C_{0-6} -alkylene-CONR³¹R³², C_{0-6} -alkylene-NR³¹-COR³¹,
 C_{0-6} -alkylene-NR³¹-CONR³¹R³², C_{0-6} -alkylene-O-CONR³¹R³², C_{0-6} -alkylene-NR³¹- CO_2R^{31} and C_{0-6} -alkylene-
 NR³¹R³²,

wherein alkyl, alkylene, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO_2H , CO_2-C_{1-4} -alkyl, $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, C_{1-4} -alkyl, halo- C_{1-4} -alkyl, O- C_{1-4} -alkyl and O-halo- C_{1-4} -alkyl; and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO_2H , CO_2-C_{1-4} -alkyl, $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, C_{1-4} -alkyl, halo- C_{1-4} -alkyl, O- C_{1-4} -alkyl and O-halo- C_{1-4} -alkyl.

[0055] In a more preferred embodiment in combination with any of the above or below embodiments



is selected from the group consisting of phenyl, pyridyl and thiophenyl, wherein phenyl, pyridyl and thiophenyl are unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of halogen, CN, SF_5 , NO_2 , oxo, C_{1-4} -alkyl, C_{0-6} -alkylene-OR³¹, C_{0-6} -alkylene-(3- to 6-membered cycloalkyl), C_{0-6} -alkylene-(3- to 6-membered heterocycloalkyl), C_{0-6} -alkylene-(6-membered aryl), C_{0-6} -alkylene-(5- to 6-membered heteroaryl), C_{0-6} -alkylene-S(O)_nR³¹, C_{0-6} -alkylene-NR³¹S(O)₂R³¹, C_{0-6} -alkylene-S(O)₂NR³¹R³², C_{0-6} -alkylene-NR³¹S(O)₂NR³¹R³², C_{0-6} -alkylene- CO_2R^{31} , O- C_{1-6} -alkylene- CO_2R^{31} , C_{0-6} -alkylene-O-COR³¹, C_{0-6} -alkylene-CONR³¹R³², C_{0-6} -alkylene-NR³¹-COR³¹, C_{0-6} -alkylene-NR³¹-CONR³¹R³², C_{0-6} -alkylene-O-CONR³¹R³², C_{0-6} -alkylene-NR³¹- CO_2R^{31} and C_{0-6} -alkylene-NR³¹R³², wherein alkyl, alkylene, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO_2H , CO_2-C_{1-4} -alkyl, $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, C_{1-4} -alkyl, halo- C_{1-4} -alkyl, O- C_{1-4} -alkyl and O-halo- C_{1-4} -alkyl;

and wherein residue -L-R¹ is linked in 1,3-orientation regarding the connection towards



and L is not a bond.

[0056] In an even more preferred embodiment in combination with any of the above or below embodiments



is selected from phenyl, pyridyl and thiophenyl; wherein phenyl, pyridyl and thiophenyl is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of F, Cl, CN, OH, oxo, C_{1-4} -alkyl, halo- C_{1-4} -alkyl, O- C_{1-4} -alkyl and O-halo- C_{1-4} -alkyl; and wherein the residue -L-R¹ is linked in 1,3-orientation regarding the connection towards



and L is not a bond.

[0057] In a most preferred embodiment in combination with any of the above or below embodiments



is phenyl, wherein phenyl is unsubstituted or substituted with F, Cl and Me; and wherein the residue -L-R¹ is linked in 1,3-orientation regarding the connection towards



and L is not a bond.

[0058] In a preferred embodiment in combination with any of the above or below embodiments

L is selected from the group consisting of a bond, C₁₋₆-alkylene, C₂₋₆-alkenylene, C₂₋₆-alkynylene, 3- to 10-membered cycloalkylene, 3- to 10-membered heterocycloalkylene containing 1 to 4 heteroatoms independently selected from N, O and S, 6- or 10-membered arylene and 5- to 10-membered heteroarylene containing 1 to 4 heteroatoms independently selected from N, O and S,

wherein alkylene, alkenylene, alkynylene, cycloalkylene, heterocycloalkylene, arylene and heteroarylene are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR⁴¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR⁴¹, C₀₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, C₀₋₆-alkyleneS(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-CO₂R⁴¹, O-C₁₋₆-alkylene-CO₂R⁴¹, C₀₋₆-alkylene-O-COR⁴¹, C₀₋₆-alkylene-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-COR⁴¹, C₀₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-alkylene-O-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-CO₂R⁴¹ and C₀₋₆-alkylene-NR⁴¹R⁴², wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the arylene and heteroarylene moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl.

[0059] in a more preferred embodiment in combination with any of the above or below embodiments


L is selected from the group consisting of 3- to 10-membered cycloalkylene, 3- to 10-membered heterocycloalkylene containing 1 to 4 heteroatoms independently selected from N, O and S, 6-membered arylene and 5- to 6-membered heteroarylene containing 1 to 2 heteroatoms independently selected from N, O and S,

wherein cycloalkylene, heterocycloalkylene, arylene and heteroarylene are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR⁴¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR⁴¹, C₀₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, C₀₋₆-alkylene-S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-CO₂R⁴¹, O-C₁₋₆-alkylene-CO₂R⁴¹, C₀₋₆-alkylene-O-COR⁴¹, C₀₋₆-alkylene-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-COR⁴¹, C₀₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-alkylene-O-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-CO₂R⁴¹ and C₀₋₆-alkylene-NR⁴¹R⁴², wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the arylene and heteroarylene moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl.

[0060] In an even more preferred embodiment in combination with any of the above or below embodiments


-L-R¹ is selected from



wherein the cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of F, Cl, Br, CN, OH, oxo, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl, O-halo-C₁₋₄-alkyl, C₁₋₄-alkyl-OH, halo-C₁₋₄-alkyl-OH, SO₂-C₁₋₄-alkyl and SO₂halo-C₁₋₄-alkyl; and wherein optionally two adjacent substituents on the phenyl ring form together a -(CH₂)₃-, -(CH₂)₄-, -OCF₂O- and -OCH₂O- group.

[0061] In a most preferred embodiment in combination with any of the above or below embodiments

-L-R¹ is selected from



wherein phenyl is unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of F, Cl, CN, OH, Me and OMe.

[0062] In a preferred embodiment in combination with any of the above or below embodiments

R¹ is selected from the group consisting of H, halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR⁴¹, Y-C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), Y-C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), Y-C₀₋₆-alkylene-(6-membered aryl), Y-C₀₋₆-alkylene-(5- to 6-membered heteroaryl), C₀₋₆-alkylene-S(=O)(-R⁴¹)=N-R⁷⁵, X-C₁₋₆-alkylene-S(=O)(-R⁴¹)=N-R⁷⁵, C₀₋₆-alkyleneS(O)_nR⁴¹, X-C₁₋₆-alkylene-S(O)_nR⁴¹, C₀₋₆-alkylene-S(=NR⁷¹)R⁴¹, X-C₁₋₆-alkylene-S(=NR⁷¹)R⁴¹, C₀₋₆-alkylene-S(O)(=NR⁷¹)R⁴¹, X-C₁₋₆-alkylene-S(O)(=NR⁷¹)R⁴¹, C₀₋₆-alkylene-S(=NR⁷¹)₂R⁴¹, X-C₁₋₆-alkylene-S(=NR⁷¹)₂R⁴¹, C₀₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, X-C₁₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, C₀₋₆-alkyleneS(O)₂NR⁴¹R⁴², X-C₁₋₆-alkylene-S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-SO₃R⁴¹, X-C₁₋₆-alkylene-SO₃R⁴¹, C₀₋₆-alkylene-CO₂R⁴¹, X-C₁₋₆-alkylene-CO₂R⁴¹, C₀₋₆-alkylene-O-COR⁴¹, X-C₁₋₆-alkylene-O-COR⁴¹, C₀₋₆-alkylene-CONR⁴¹R⁴², X-C₁₋₆-alkylene-CONR⁴¹R⁴², C₀₋₆-alkylene-CONR⁴¹OR⁴¹, X-C₁₋₆-alkylene-CONR⁴¹OR⁴¹, C₀₋₆-alkylene-CONR⁴¹SO₂R⁴¹, X-C₁₋₆-alkylene-CONR⁴¹SO₂R⁴¹, C₀₋₆-alkylene-NR⁴¹-COR⁴¹, X-C₁₋₆-C₀₋₆-alkylene-NR⁴¹-COR⁴¹, C₀₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-alkylene-O-CONR⁴¹R⁴², X-C₁₋₆-alkylene-O-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-CO₂R⁴¹, X-C₁₋₆-alkylene-NR⁴¹-CO₂R⁴¹, C₀₋₆-alkylene-NR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹R⁴²,

wherein alkyl, alkylene, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the aryl and heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and





wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-

C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl.


[0063] In a more preferred embodiment in combination with any of the above or below embodiments R¹ is selected from CO₂H, tetrazole, CH₂CO₂H, OCH₂CO₂H, SO₂CH₂CO₂H, CHMeCO₂H, CMe₂CO₂H, C(OH)MeCO₂H, CONHSO₂Me and CONH(OH); and optionally the glycine and tauro conjugate thereof.

[0064] In a most preferred embodiment in combination with any of the above or below embodiments R¹ is selected from CO₂H and C(OH)MeCO₂H; and optionally the glycine and tauro conjugate thereof.

[0065] In a preferred embodiment in combination with any of the above or below embodiments -L-R¹ is selected from




and




and optionally the glycine and tauro conjugate thereof.

[0066] In a more preferred embodiment in combination with any of the above or below embodiments -L-R¹ is selected from



and optionally the glycine and tauro conjugate thereof.

[0067] In a most preferred embodiment in combination with any of the above or below embodiments -L-R¹ is selected from



and optionally the glycine and tauro conjugate thereof.

[0068] In a preferred embodiment in combination with any of the above or below embodiments

(D)

is selected from the group consisting of 3- to 10-membered cycloalkyl, 3- to 10-membered heterocycloalkyl containing 1 to 3 heteroatoms independently selected from N, O and S, 6- to 14-membered aryl and 5- to 14-membered heteroaryl containing 1 to 4 heteroatoms independently selected from N, O and S,

wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR²¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR²¹, C₀₋₆-alkylene-NR²¹S(O)₂R²², C₀₋₆-alkylene-S(O)₂NR²¹R²², C₀₋₆-alkylene-NR²¹S(O)₂NR²¹R²², C₀₋₆-alkylene-CR⁴¹(=N-OR⁴¹), C₀₋₆-alkylene-CO₂R²¹, O-C₁₋₆-alkylene-CO₂R²¹, C₀₋₆-alkylene-O-COR²¹, C₀₋₆-alkylene-CONR²¹R²², C₀₋₆-alkylene-NR²¹-COR²¹, C₀₋₆-alkylene-NR²¹-CONR²¹R²², C₀₋₆-alkylene-O-CONR²¹R²², C₀₋₆-alkylene-NR²¹-CO₂R²¹ and C₀₋₆-alkylene-NR²¹R²², wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, CO-OC₁₄-alkyl, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered

partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the cycloalkyl or heterocycloalkyl moiety form a 5- to 6-membered unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

wherein



has a substituent from above in 1,2-orientation regarding to the connection towards



or has an annelated additional cycle in 1,2-orientation.

[0069] In a more preferred embodiment in combination with any of the above or below embodiments



is selected from the group consisting of 6- or 10-membered aryl and 5- to 10-membered heteroaryl containing 1 to 4 heteroatoms independently selected from N, O and S,

wherein aryl and heteroaryl are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR²¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR²¹, C₀₋₆-alkylene-NR²¹S(O)₂R²¹, C₀₋₆-alkylene-S(O)₂NR²¹R²², C₀₋₆-alkylene-NR²¹S(O)₂NR²¹R²², C₀₋₆-alkylene-CR⁴¹(=N-OR⁴¹), C₀₋₆-alkylene-CO₂R²¹, O-C₁₋₆-alkylene-CO₂R²¹, C₀₋₆-alkylene-O-COR²¹, C₀₋₆-alkylene-CONR²¹R²², C₀₋₆-alkylene-NR²¹-COR²¹, C₀₋₆-alkylene-NR²¹-CONR²¹R²², C₀₋₆-alkylene-O-CONR²¹R²², C₀₋₆-alkylene-NR²¹-CO₂R²¹ and C₀₋₆-alkylene-NR²¹R²²,

wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, CO-OC₁₋₄-alkyl, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

wherein



has a substituent from above in 1,2-orientation regarding to the connection towards




or has an annelated additional cycle in 1,2-orientation.

[0070] In an even more preferred embodiment in combination with any of the above or below embodiments




is selected from the group consisting




wherein

R^2 is selected from Me, F, Cl, CN, Me, CHO, CHF_2 , CF_3 , SO_2Me ,



and



and

wherein




is not further substituted or further substituted with 1 to 2 substituents selected from the group consisting F, Cl, CN, Me, OMe, CHO, CHF_2 and CF_3 .

[0071] In a most preferred embodiment in combination with any of the above or below embodiments



is selected from the group consisting of





[0072] In a preferred embodiment in combination with any of the above or below embodiments


Formula (I) contains a substituent selected from the group consisting of CO_2H , tetrazole, CONHSO_2Me and $\text{CONH}(\text{OH})$; and optionally the glycine and tauro conjugate thereof.

[0073] In a more preferred embodiment in combination with any of the above or below embodiments Formula (I) contains a carboxylic acid moiety and optionally the glycine and tauro conjugate thereof.


[0074] In a most preferred embodiment, the compound is selected from







and



or a glycine conjugate or tauro conjugate thereof; and

an enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

[0075] In an upmost preferred embodiment, the compound is 2-chloro-3'-(3-(2-cyanothiophen-3-yl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid or a glycine conjugate or tauro conjugate thereof and optionally a pharmaceutically acceptable salt thereof. Even more preferred is 2-chloro-3'-(3-(2-cyanothiophen-3-yl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid and optionally a pharmaceutically acceptable salt thereof.

[0076] In a similar upmost preferred embodiment, the compound is 2-chloro-3'-(3-(3-cyanopyrazin-2-yl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid or a glycine conjugate or tauro conjugate thereof and optionally a pharmaceutically acceptable salt thereof. Even more preferred is 2-chloro-3'-(3-(3-cyanopyrazin-2-yl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid and optionally a pharmaceutically acceptable salt thereof.

[0077] In a similar upmost preferred embodiment, the compound is 2-chloro-3'-(3-(2-cyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid or a glycine conjugate or tauro conjugate thereof and optionally a pharmaceutically acceptable salt thereof. Even more preferred is 2-chloro-3'-(3-(2-cyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid and optionally a pharmaceutically acceptable salt thereof.

[0078] In a similar upmost preferred embodiment, the compound is 2-chloro-3'-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid or a glycine conjugate or tauro conjugate thereof and optionally a pharmaceutically acceptable salt thereof. Even more preferred is 2-chloro-3'-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid and optionally a pharmaceutically acceptable salt thereof.

[0079] In a similar upmost preferred embodiment, the compound is 3'-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-2,6-difluoro-[1,1'-biphenyl]-4-carboxylic acid or a glycine conjugate or tauro conjugate thereof and optionally a pharmaceutically acceptable salt thereof. Even more preferred is 3'-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-2,6-difluoro-[1,1'-biphenyl]-4-carboxylic acid and optionally a pharmaceutically acceptable salt thereof.

[0080] In a similar upmost preferred embodiment, the compound is 2-chloro-3'-(3-(2,6-dicyanophenyl)-1-((4-(difluor-

omethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-5-fluoro-[1,1'-biphenyl]-4-carboxylic acid or a glycine conjugate or tauro conjugate thereof and optionally a pharmaceutically acceptable salt thereof. Even more preferred is 2-chloro-3'-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-5-fluoro-[1,1'-biphenyl]-4-carboxylic acid and optionally a pharmaceutically acceptable salt thereof.

[0081] The invention also provides the compound of the invention for use as a medicament.

[0082] Also provided is the compound of the present invention for use in the prophylaxis and/or treatment of diseases amenable for treatment with LXR modulators.

[0083] Also provided is the compound of the invention for use in treating a LXR mediated disease selected from non-alcoholic fatty liver disease, non-alcoholic steatohepatitis, liver inflammation, liver fibrosis, obesity, insulin resistance, type II diabetes, familial hypercholesterolemia, hypercholesterolemia in nephrotic syndrome, metabolic syndrome, cardiac steatosis, cancer, viral myocarditis, hepatitis C virus infection or its complications, and unwanted side-effects of long-term glucocorticoid treatment in diseases such as rheumatoid arthritis, inflammatory bowel disease and asthma.

[0084] In a preferred embodiment, the disease is selected from non-alcoholic fatty liver disease, non-alcoholic steatohepatitis, liver inflammation, liver fibrosis, obesity, insulin resistance, type II diabetes, familial hypercholesterolemia, hypercholesterolemia in nephrotic syndrome, metabolic syndrome or cardiac steatosis.

[0085] In a similar preferred embodiment, the disease is cancer.

[0086] In a similar preferred embodiment, the disease is selected from viral myocarditis, hepatitis C virus infection or its complications.

[0087] The invention further relates to a method for preventing and/or treating diseases mediated by LXRs, the method comprising administering a compound of the present invention in an effective amount to a subject in need thereof.

[0088] More specifically, the invention relates to a method for preventing and treating diseases selected from non-alcoholic fatty liver disease, non-alcoholic steatohepatitis, liver inflammation, liver fibrosis, obesity, insulin resistance, type II diabetes, familial hypercholesterolemia, hypercholesterolemia in nephrotic syndrome, metabolic syndrome, cardiac steatosis, cancer, viral myocarditis, hepatitis C virus infection or its complications, and unwanted side-effects of long-term glucocorticoid treatment in diseases such as rheumatoid arthritis, inflammatory bowel disease and asthma.

[0089] Moreover, the invention also relates to the use of a compound according to the present invention in the preparation of a medicament for the prophylaxis and/or treatment of a LXR mediated disease.

[0090] More specifically, the invention relates to the use of a compound according to the present invention in the preparation of a medicament for the prophylaxis and/or treatment of a LXR mediated disease, wherein the disease is selected from non-alcoholic fatty liver disease, non-alcoholic steatohepatitis, liver inflammation, liver fibrosis, obesity, insulin resistance, type II diabetes, familial hypercholesterolemia, hypercholesterolemia in nephrotic syndrome, metabolic syndrome, cardiac steatosis, cancer, viral myocarditis, hepatitis C virus infection or its complications, and unwanted side-effects of long-term glucocorticoid treatment in diseases such as rheumatoid arthritis, inflammatory bowel disease and asthma.

[0091] Also provided is a pharmaceutical composition comprising the compound of the invention and a pharmaceutically acceptable carrier or excipient.

[0092] In the context of the present invention "C₁₋₆-alkyl" means a saturated alkyl chain having 1 to 6 carbon atoms which may be straight chained or branched. Examples thereof include methyl, ethyl, propyl, isopropyl, *n*-butyl, isobutyl, *tert*-butyl, *n*-pentyl, isopentyl, *n*-hexyl and isohexyl. Similarly, "C₁₋₄-alkyl" means a saturated alkyl chain having 1 to 4 carbon atoms which may be straight chained or branched. Examples thereof include methyl, ethyl, propyl, isopropyl, *n*-butyl, isobutyl, and *tert*-butyl.

[0093] The term "halo-C₁₋₄-alkyl" means that one or more hydrogen atoms in the alkyl chain are replaced by a halogen. A preferred example thereof is CH₂F, CHF₂ and CF₃.


[0094] A "C₀₋₆-alkylene" means that the respective group is divalent and connects the attached residue with the remaining part of the molecule. Moreover, in the context of the present invention, "C₀-alkylene" is meant to represent a bond, whereas C₁-alkylene means a methylene linker, C₂-alkylene means a ethylene linker or a methyl-substituted methylene linker and so on. In the context of the present invention, a C₀₋₆-alkylene preferably represents a bond, a methylene, a ethylene group or a propylene group.

[0095] Similarly, a "C₂₋₆-alkenylene" and a "C₂₋₆-alkynylene" means a divalent alkenyl or alkynyl group which connects two parts of the molecule.

[0096] A 3- to 10-membered cycloalkyl group means a saturated or partially unsaturated mono-, bi-, spiro- or multicyclic ring system comprising 3 to 10 carbon atoms. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, bicyclo[2.2.2]octyl, bicyclo[3.2.1]octanyl, spiro[3.3]heptyl, bicyclo[2.2.1]heptyl, adamantyl and pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octyl. Consequently, a 3- to 6-membered cycloalkyl group means a saturated or partially unsaturated mono- bi-, or spirocyclic ring system comprising 3 to 6 carbon atoms whereas a 5- to 8-membered cycloalkyl group means a saturated or partially unsaturated mono-, bi-, or spirocyclic ring system comprising 5 to 8 carbon atoms.

[0097] A 3- to 10-membered heterocycloalkyl group means a saturated or partially unsaturated 3 to 10 membered carbon mono-, bi-, spiro- or multicyclic ring wherein 1, 2, 3 or 4 carbon atoms are replaced by 1, 2, 3 or 4 heteroatoms,


respectively, wherein the heteroatoms are independently selected from N, O, S, SO and SO₂. Examples thereof include epoxidyl, oxetanyl, pyrrolidinyl, tetrahydrofuranyl, piperidinyl, piperazinyl tetrahydropyranyl, 1,4-dioxanyl, morpholinyl, 4-quinuclidinyl, 1,4-dihydropyridinyl and 6-azabicyclo[3.2.1]octanyl. The heterocycloalkyl group can be connected with the remaining part of the molecule via a carbon, nitrogen (e.g. in morpholine or piperidine) or sulfur atom. An example for a S-linked heterocycloalkyl is the cyclic sulfonimidamide



[0098] A 5- to 14-membered mono-, bi- or tricyclic heteroaromatic ring system (within the application also referred to as heteroaryl) means an aromatic ring system containing up to 6 heteroatoms independently selected from N, O, S, SO and SO₂. Examples of monocyclic heteroaromatic rings include pyrrolyl, imidazolyl, furanyl, thiophenyl (thienyl), pyridinyl, pyrimidinyl, pyrazinyl, pyrazolyl, oxazolyl, isoxazolyl, triazolyl, oxadiazolyl and thiadiazolyl. It further means a bicyclic ring system wherein the heteroatom(s) may be present in one or both rings including the bridgehead atoms. Examples thereof include quinolinyl, isoquinolinyl, quinoxalyl, benzimidazolyl, benzisoxazolyl, benzofuranyl, benzoxazolyl, indolyl, indolizyl, 1,5-naphthyridinyl, 1,7-naphthyridinyl and pyrazolo[1,5-a]pyrimidinyl. Examples of tricyclic heteroaromatic rings include acridinyl, benzo[b][1,5]naphthyridinyl and pyrido[3,2-b][1,5]naphthyridinyl.

[0099] The nitrogen or sulphur atom of the heteroaryl system may also be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide.

[0100] If not stated otherwise, the heteroaryl system can be connected via a carbon or nitrogen atom. Examples for N-linked heterocycles are



[0101] A 6- to 14-membered mono-, bi- or tricyclic aromatic ring system (within the application also referred to as aryl) means an aromatic carbon cycle such as phenyl, naphthyl, anthracenyl or phenanthrenyl.

[0102] The term "N-oxide" denotes compounds, where the nitrogen in the heteroaromatic system (preferably pyridinyl) is oxidized. Such compounds can be obtained in a known manner by reacting a compound of the present invention (such as in a pyridinyl group) with H₂O₂ or a peracid in an inert solvent.

[0103] Halogen is selected from fluorine, chlorine, bromine and iodine, more preferably fluorine or chlorine and most preferably fluorine.

[0104] Any formula or structure given herein, is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds. Isotopically labeled compounds have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the disclosure include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as, but not limited to ²H (deuterium, D), ³H (tritium), ¹¹C, ¹³C, ¹⁴C, ¹⁵N, ¹⁸F, ³¹P, ³²P, ³⁵S, ³⁶Cl and ¹²⁵I. Various isotopically labeled compounds of the present disclosure, for example those into which radioactive isotopes such as ³H, ¹³C and ¹⁴C are incorporated. Such isotopically labelled compounds may be useful in metabolic studies, reaction kinetic studies, detection or imaging techniques, such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT) including drug or substrate tissue distribution assays or in radioactive treatment of patients. Isotopically labeled compounds of this disclosure and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

[0105] The disclosure also includes "deuterated analogs" of compounds of Formula (I) in which from 1 to n hydrogens attached to a carbon atom is/are replaced by deuterium, in which n is the number of hydrogens in the molecule. Such compounds may exhibit increased resistance to metabolism and thus be useful for increasing the half-life of any compound of Formula (I) when administered to a mammal, e.g. a human. See, for example, Foster in Trends Pharmacol. Sci. 1984;5:524. Such compounds are synthesized by means well known in the art, for example by employing starting materials in which one or more hydrogens have been replaced by deuterium.

[0106] Deuterium labelled or substituted therapeutic compounds of the disclosure may have improved DMPK (drug metabolism and pharmacokinetics) properties, relating to distribution, metabolism and excretion (ADME). Substitution with heavier isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic


stability, for example increased *in vivo* half-life, reduced dosage requirements and/or an improvement in therapeutic index. An ^{18}F labeled compound may be useful for PET or SPECT studies.

[0107] The concentration of such a heavier isotope, specifically deuterium, may be defined by an isotopic enrichment factor. In the compounds of this disclosure any atom not specifically designated as a particular isotope is meant to represent any stable isotope of that atom. Unless otherwise stated, when a position is designated specifically as "H" or "hydrogen", the position is understood to have hydrogen at its natural abundance isotopic composition. Accordingly, in the compounds of this disclosure any atom specifically designated as a deuterium (D) is meant to represent deuterium.


[0108] Furthermore, the compounds of the present invention are partly subject to tautomerism. For example, if a heteroaromatic group containing a nitrogen atom in the ring is substituted with a hydroxy group on the carbon atom adjacent to the nitrogen atom, the following tautomerism can appear:




[0109] A cycloalkyl or heterocycloalkyl group can be connected straight or spirocyclic, e.g. when cyclohexane is substituted with the heterocycloalkyl group oxetane, the following structures are possible:




[0110] The term "1,3-orientation" means that on a ring the substituents have at least one possibility, where 3 atoms are between the two substituents attached to the ring system, e.g.



[0111] The term "1,2-orientation" (*ortho*) means that on a ring the substituents have one possibility, where 2 atoms are between the two substituents attached to the ring system, e.g.




alternatively the residue R can be incorporated in an annelated additional cycle, e.g.



[0112] It will be appreciated by the skilled person that when lists of alternative substituents include members which, because of their valency requirements or other reasons, cannot be used to substitute a particular group, the list is

intended to be read with the knowledge of the skilled person to include only those members of the list which are suitable for substituting the particular group.

[0113] The term "glycine conjugate or tauro conjugate thereof" means, that the carboxylic acid moiety in the molecule is connected with glycine or taurine, respectively, to form the conjugate (and potentially a prodrug, solvate or pharmaceutically acceptable salt thereof):



[0114] Where tautomerism, like e.g. keto-enol tautomerism, of compounds of the present invention or their prodrugs may occur, the individual forms, like e.g. the keto and enol form, are each within the scope of the invention as well as their mixtures in any ratio. Same applies for stereoisomers, like e.g. enantiomers, cis/trans-isomers, atropisomers, conformers and the like.

[0115] If desired, isomers can be separated by methods well known in the art, e.g. by liquid chromatography. Same applies for enantiomers by using e.g. chiral stationary phases. Additionally, enantiomers may be isolated by converting them into diastereomers, i.e. coupling with an enantiomerically pure auxiliary compound, subsequent separation of the resulting diastereomers and cleavage of the auxiliary residue. Alternatively, any enantiomer of a compound of the present invention may be obtained from stereoselective synthesis using optically pure starting materials. Another way to obtain pure enantiomers from racemic mixtures would use enantioselective crystallization with chiral counterions.

[0116] The compounds of the present invention can be in the form of a pharmaceutically acceptable salt or a solvate. The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids, including inorganic bases or acids and organic bases or acids. In case the compounds of the present invention contain one or more acidic or basic groups, the invention also comprises their corresponding pharmaceutically or toxicologically acceptable salts, in particular their pharmaceutically utilizable salts. Thus, the compounds of the present invention which contain acidic groups can be present on these groups and can be used according to the invention, for example, as alkali metal salts, alkaline earth metal salts or ammonium salts. More precise examples of such salts include sodium salts, potassium salts, calcium salts, magnesium salts or salts with ammonia or organic amines such as, for example, ethylamine, ethanolamine, triethanolamine or amino acids. The compounds of the present invention which contain one or more basic groups, i.e. groups which can be protonated, can be present and can be used according to the invention in the form of their addition salts with inorganic or organic acids. Examples of suitable acids include hydrogen chloride, hydrogen bromide, phosphoric acid, sulfuric acid, nitric acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acids, oxalic acid, acetic acid, tartaric acid, lactic acid, salicylic acid, benzoic acid, formic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, malic acid, sulfaminic acid, phenylpropionic acid, gluconic acid, ascorbic acid, isonicotinic acid, citric acid, adipic acid, and other acids known to the person skilled in the art. If the compounds of the present invention simultaneously contain acidic and basic groups in the molecule, the invention also includes, in addition to the salt forms mentioned, inner salts or betaines (zwitterions). The respective salts can be obtained by customary methods which are known to the person skilled in the art like, for example, by contacting these with an organic or inorganic acid or base in a solvent or dispersant, or by anion exchange or cation exchange with other salts. The present invention also includes all salts of the compounds of the present invention which, owing to low physiological compatibility, are not directly suitable for use in pharmaceuticals but which can be used, for example, as intermediates for chemical reactions or for the preparation of pharmaceutically acceptable salts.

[0117] Further the compounds of the present invention may be present in the form of solvates, such as those which include as solvate water, or pharmaceutically acceptable solvates, such as alcohols, in particular ethanol.

[0118] Furthermore, the present invention provides pharmaceutical compositions comprising at least one compound of the present invention, or a prodrug compound thereof, or a pharmaceutically acceptable salt or solvate thereof as active ingredient together with a pharmaceutically acceptable carrier.

[0119] "Pharmaceutical composition" means one or more active ingredients, and one or more inert ingredients that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing at least one compound of the present invention and a pharmaceutically acceptable carrier.

[0120] The pharmaceutical composition of the present invention may additionally comprise one or more other compounds as active ingredients like a prodrug compound or other nuclear receptor modulators.

[0121] The compositions are suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and

intravenous), ocular (ophthalmic), pulmonary (nasal or buccal inhalation) or nasal administration, although the most suitable route in any given case will depend on the nature and severity of the conditions being treated and on the nature of the active ingredient. They may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the art of pharmacy.

[0122] The compounds of the present invention act as LXR modulators.

[0123] Ligands to nuclear receptors including LXR ligands can either act as agonists, antagonists or inverse agonists. An agonist in this context means a small molecule ligand that binds to the receptor and stimulates its transcriptional activity as determined by e.g. an increase of mRNAs or proteins that are transcribed under control of an LXR response element. Transcriptional activity can also be determined in biochemical or cellular *in vitro* assays that employ just the ligand binding domain of LXR α or LXR β but use the interaction with a cofactor (i.e. a corepressor or a coactivator), potentially in conjunction with a generic DNA-binding element such as the Gal4 domain, to monitor agonistic, antagonistic or inverse agonistic activity.

[0124] Whereas an agonist by this definition stimulates LXR- or LXR-Gal4- driven transcriptional activity, an antagonist is defined as a small molecule that binds to LXRs and thereby inhibits transcriptional activation that would otherwise occur through an endogenous LXR ligand.

[0125] An inverse agonist differs from an antagonist in that it not only binds to LXRs and inhibits transcriptional activity but in that it actively shuts down transcription directed by LXR, even in the absence of an endogenous agonist. Whereas it is difficult to differentiate between LXR antagonistic and inverse agonistic activity *in vivo*, given that there are always some levels of endogenous LXR agonist present, biochemical or cellular reporter assays can more clearly distinguish between the two activities. At a molecular level an inverse agonist does not allow for the recruitment of a coactivator protein or active parts thereof whereas it should lead to an active recruitment of corepressor proteins are active parts thereof. An LXR antagonist in this context would be defined as an LXR ligand that neither leads to coactivator nor to corepressor recruitment but acts just through displacing LXR agonists. Therefore, the use of assays such as the Gal4-mammalian-two-hybrid assay is mandatory in order to differentiate between coactivator or corepressor-recruiting LXR compounds (Kremoser et al., Drug Discov. Today 2007;12:860; Gronemeyer et al., Nat. Rev. Drug Discov. 2004;3:950).

[0126] Since the boundaries between LXR agonists, LXR antagonists and LXR inverse agonists are not sharp but fluent, the term "LXR modulator" was coined to encompass all compounds which are not clean LXR agonists but show a certain degree of corepressor recruitment in conjunction with a reduced LXR transcriptional activity. LXR modulators therefore encompass LXR antagonists and LXR inverse agonists and it should be noted that even a weak LXR agonist can act as an LXR antagonist if it prevents a full agonist from full transcriptional activation.

[0127] Figure 1 illustrates the differences between LXR agonists, antagonists and inverse agonists exemplified by their different capabilities to recruit coactivators or corepressors.

[0128] The compounds are useful for the prophylaxis and/or treatment of diseases which are mediated by LXRs. Preferred diseases are all disorders associated with steatosis, i.e. tissue fat accumulation. Such diseases encompass the full spectrum of non-alcoholic fatty liver disease including non-alcoholic steatohepatitis, liver inflammation and liver fibrosis, furthermore insulin resistance, metabolic syndrome and cardiac steatosis. An LXR modulator based medicine might also be useful for the treatment of hepatitis C virus infection or its complications and for the prevention of unwanted side-effects of long-term glucocorticoid treatment in diseases such as rheumatoid arthritis, inflammatory bowel disease and asthma.

[0129] A different set of applications for LXR modulators might be in the treatment of cancer. LXR antagonists or inverse agonists might useful to counteract the so-called Warburg effect which is associated with a transition from normal differentiated cells towards cancer cells (see Liberti et al., Trends Biochem. Sci. 2016;41:211; Ward & Thompson, Cancer Cell 2012;21:297-308). Furthermore, LXR is known to modulate various components of the innate and adaptive immune system. Oxysterols, which are known as endogenous LXR agonists were identified as mediators of an LXR-dependent immunosuppressive effect found in the tumor microenvironment (Traversari et al., Eur. J. Immunol. 2014;44:1896). Therefore, it is reasonable to assume that LXR antagonists or inverse agonists might be capable of stimulating the immune system and antigen-presenting cells, in particular, to elicit an anti-tumor immune response. The latter effects of LXR antagonists or inverse agonists might be used for a treatment of late stage cancer, in general, and in particular for those types of cancerous solid tumors that show a poor immune response and highly elevated signs of Warburg metabolism.

[0130] In more detail, anti-cancer activity of the LXR inverse agonist **SR9243** was shown to be mediated by interfering with the Warburg effect and lipogenesis in different tumor cells *in vitro* and SW620 colon tumor cells in athymic mice *in vivo* (see Flaveny et al. Cancer Cell. 2015;28:42; Steffensen, Cancer Cell 2015;28:3).

[0131] Therefore, LXR modulators (preferably LXR inverse agonists) may be useful for the treatment of Warburg-dependent cancers.

[0132] LXR modulators (preferably LXR inverse agonists) may counteract the diabetogenic effects of glucocorticoids without compromising the anti-inflammatory effects of glucocorticoids and could therefore be used to prevent unwanted side-effects of long-term glucocorticoid treatment in diseases such as rheumatoid arthritis, inflammatory bowel disease

and asthma (Patel et al. Endocrinology 2017;158:1034).

[0133] LXR modulators (preferably LXR inverse agonists) may be useful for the treatment of hepatitis C virus mediated liver steatosis (see García-Mediavilla et al. Lab. Invest. 2012;92:1191).

[0134] LXR modulators (preferably LXR inverse agonists) may be useful for the treatment of viral myocarditis (see Papageorgiou et al. Cardiovasc. Res. 2015;107:78).

[0135] LXR modulators (preferably LXR inverse agonists) may be useful for the treatment of insulin resistance (see Zheng et al. PLoS One 2014;9:e101269).


[0136] LXR modulators (preferably LXR inverse agonists) may be useful for the treatment of familial hypercholesterolemia (see Zhou et al. J. Biol. Chem. 2008;283:2129).

[0137] LXR modulators (preferably LXR inverse agonists) may be useful for the treatment of hypercholesterolemia in nephrotic syndrome (see Liu & Vazizi in Nephrol. Dial. Transplant. 2014;29:538).

Experimental Section


[0138] The compounds of the present invention can be prepared by a combination of methods known in the art including the procedures described in Schemes I to V below.

[0139] The synthetic route depicted in Scheme I starts with the preparation of alkynes **I-c** by Sonogashira couplings. Subsequently, the free amino group of **I-c** is reacted with sulfonyl chlorides **I-d** in the presence of an appropriate base and appropriate solvent to afford alkynesulfonamides **I-e**. **I-e** undergoes cyclization and concomitant reaction with aromatic halides **I-f** in the presence of appropriate catalyst (e.g. Pd-catalysts), appropriate solvent and temperature to afford compounds of the present invention (**I**). Further manipulation of functional groups present in R^1 by standard methods, known to persons skilled in the art (e.g. ester hydrolysis, amide bond formation), can give rise to further compounds of the present invention. Alternatively, alkyneamine **I-c** can be transformed into alkynetrifluoroacetamides **I-g** which can also undergo aforementioned cyclization and concomitant reaction with aromatic halides **I-f** to afford intermediates **I-h** with an unsubstituted NH. Reaction with sulfonyl chlorides **I-d** in the presence of an appropriate base and appropriate solvent also affords compounds of Formula (**I**).




Scheme I: Synthesis of compounds of the present invention.

[0140] A variation of the routes shown in Scheme I is shown in Scheme II. Alkynesulfonamide **I-e** is reacted in the presence of NIS to afford iodinated intermediates **II-b** which can be substrates for Suzuki couplings to afford compounds (**I**). Alternatively, cyclization of alkynesulfonamide **I-e** in the presence of appropriate catalyst (e.g. Pd-catalysts), appropriate solvent and temperature but without the presence of halides **I-f** afford 3-unsubstituted intermediates **II-d**. Reactions with NBS afford brominated intermediates **II-e** which are likewise substrates for Suzuki coupling reactions to afford compounds of Formula (**I**).




Scheme II: Synthetic route for compounds of the present invention, with introduction of moiety D via Suzuki coupling.

[0141] A further variation of the synthetic route depicted in Schemes I and II is shown in Scheme III. In the presence of B_2Pin_2 , appropriate catalyst (e.g. Pd-catalysts), appropriate solvent, additives and temperature, intermediates **I-e** can undergo cyclization and concomitant formation of 3-pinacoyl boronic esters **III-b**. These can be substrates for Suzuki coupling reactions to afford compounds of the present invention with Formula (I).




Scheme III: Alternative synthetic route for compounds of the present invention, with introduction of D via Suzuki coupling.

[0142] In Scheme IV is depicted a synthetic route for the late stage introduction of the right hand side moieties -L-R¹ to the compounds of the present invention. Sonogashira coupling of **I-a** with bromo-iodo-aromatics **IV-a** afford bromo-alkyneamines **IV-b** which can be transformed to sulfonamides **IV-c**. These can undergo cyclization and concomitant reaction with aromatic bromides **IV-d** in the presence of appropriate catalysts (e.g. Pd-catalysts), appropriate solvent and temperature to afford advanced intermediates **IV-e** with a bromo substituent on ring C. Finally, intermediates **IV-e** can be used as substrates for Suzuki couplings to afford compounds of Formula (I).



Scheme IV: Synthetic route for compounds of the present invention, with final introduction of –L-R¹ via Suzuki coupling.

25 **[0143]** In Scheme V are summarized the synthetic routes for the preparation of the compounds of the present invention starting from the preformed central pyrolo-annulated bicyclic aromatic. N-protected 2-pinacolyl boronic esters **V-a** can undergo Suzuki coupling with halides **V-b** to afford intermediates **V-c**. After bromination with NBS the 3-bromo intermediates **V-d** are obtained, which, after a second Suzuki coupling, are converted to N-protected advanced intermediates **V-e**. When starting with N-protected 3-pinacolyl boronic esters **V-a**, first Suzuki coupling and then bromination of the 2-position and subsequent second Suzuki coupling affords likewise intermediates **V-e**. After deprotection and reaction of the free NH with sulfonyl chlorides **I-d**, in the presence of an appropriate base and solvent, compounds (I) are obtained.




Scheme V: Synthesis of compounds of the present invention, starting from preformed core aromatic.

Abbreviations**[0144]**

5	Ac	acetyl
	ACN	acetonitrile
	AIBN	azobisisobutyronitrile
	aq.	aqueous
	BINAP	(2,2'-bis(diphenylphosphino)-1,1'-binaphthyl)
10	B ₂ Pin ₂	4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane
	Boc	<i>tert</i> -butoxycarbonyl
	BPO	dibenzoyl peroxide
	<i>m</i> -CPBA	<i>meta</i> -chloroperbenzoic acid
	Cy	cyclohexyl
15	DAST	diethylaminosulfur trifluoride
	dba	dibenzylideneacetone
	DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
	DCM	dichloromethane
	DEA	diethanolamine
20	DEAD	diethyl azodicarboxylate
	DIEA or DIPEA	diisopropylethylamine
	DMAP	4- <i>N,N</i> -dimethylaminopyridine
	DMF	<i>N,N</i> -dimethylformamide
	dppf	1,1'-bis(diphenylphosphino)ferrocene
25	EA	ethyl acetate
	EDCI	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
	FCC	flash column chromatography on silica gel
	h	hour(s)
	HATU	<i>O</i> -(7-azabenzotriazole-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
30	HOBt	hydroxybenzotriazole
	IBX	2-iodoxybenzoic acid
	LDA	lithium diisopropylamide
	LiHMDS	lithium <i>bis</i> (trimethylsilyl)amide
	NBS	<i>N</i> -bromosuccinimide
35	NIS	<i>N</i> -iodosuccinimide
	Pin	pinacolato (OCMe ₂ CMe ₂ O)
	PE	petroleum ether
	prep	preparative
	sat.	saturated (aqueous)
40	Sphos	2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl
	TBAF	tetra- <i>n</i> -butylammonium fluoride
	TEA	triethylamine
	TFA	trifluoroacetic acid
	TFAA	trifluoroacetic acid anhydride
45	THF	tetrahydrofuran
	TLC	thin layer chromatography
	Tr	Trityl
	Xantphos	4,5-bis(diphenylphosphino)-9,9-dimethylxanthene
	XPhos	2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl



Preparative Example P1**[0145]**

55

**P1**

Methyl 2-(4-bromo-3-chlorophenyl)-2-(dimethylamino)acetate (P1)

[0146] To a solution of methyl 2-amino-2-(4-bromo-3-chlorophenyl)acetate (300 mg, 1.08 mmol) in MeOH (6 mL) was added CH₂O (37 wt. % in H₂O; 0.5 mL) and HCOOH (2.0 mL). The mixture was stirred at rt for 30 min, then NaBH(OAc)₃ (572 mg, 2.7 mmol) was added. The mixture was stirred at rt for 2 h, diluted with EA (150 mL) and washed with water (15 mL), sat. NaHCO₃ (15 mL) and brine (10 mL). The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by prep-TLC (EA:PE = 1:4) to afford compound **P1** as a colorless oil.

Preparative Example P2**[0147]**Step 1: 2-((Trimethylsilyl)ethynyl)pyridin-3-amine (P2a)**[0148]**

[0149] Pd(PPh₃)₄ (993 mg, 0.86 mmol), CuI (164 mg, 0.86 mmol) and PPh₃ (225 mg, 0.86 mmol) were combined in a round-bottom flask, then degassed and refilled with N₂ three times. To the mixture was added TEA (43 mL), 2-bromopyridin-3-amine (1.49 g, 8.59 mmol) and ethynyltrimethylsilane (2.43 mL, 18.0 mmol). The mixture was stirred at 60°C for 6 h, cooled to rt, filtered through Celite and washed with EA (40 mL). The filtrate was concentrated to give compound **P2a** as a black solid, which was used in the next step without further purification.

Step 2: 2-Ethynylpyridin-3-amine (P2)













[0150] To a solution of compound **P2a** (2.16 g, 8.59 mmol) in THF (26 mL) was added TBAF (26 mL, 1M in THF, 26 mmol) and the mixture was stirred at rt for 3 h, concentrated and purified by FCC (EA/PE = 1:19 to 1:0) to give compound **P2** as a white solid.



Preparative Example P2/1 to P2/9

[0151] The following Preparative Examples were prepared similar as described for Preparative Example P2 using the appropriate building blocks.

#	building block	structure
P2/1		
P2/2		
P2/3		


(continued)

#	building block	structure
P2/4		
P2/5		
P2/6		
P2/7		
P2/8		
P2/9		

Preparative Example P3**[0152]****Step 1: (4-Bromo-2-mercaptophenyl)methanol (P3a)****[0153]**

[0154] To a solution of 4-bromo-2-mercaptobenzoic acid (1.5 g, 6.5 mmol) in THF (30 mL) was added BH_3 (13 mL, 1M in THF). This mixture was stirred overnight and quenched with water (30 mL) and diluted with EA (20 mL). The organic layer was separated and the aq. layer was washed with EA (3 × 20 mL). The combined organic layer was washed with brine (30 mL), dried over Na_2SO_4 , filtered and concentrated. The yellow solid was used in the next step without purification.

Step 2: Ethyl 2-((5-bromo-2-(hydroxymethyl)phenyl)thio)acetate (P3b)**[0155]**




[0156] To a mixture of compound **P3a** (436 mg, 2.0 mmol) and ethyl 2-bromoacetate (306 mg, 2.0 mmol) in DMF (10 mL) was added Cs_2CO_3 (2.0 g, 6.0 mmol). The mixture was stirred at rt overnight, diluted with water (100 mL) and extracted with EA (3×30 mL). The combined organic layer was washed with brine (30 mL), dried over Na_2SO_4 , filtered, concentrated and purified by FCC (PE:EA = 5:1) to afford compound **P3b** as a white solid.

Step 3: Ethyl 2-((5-bromo-2-(hydroxymethyl)phenyl)sulfonyl)acetate (P3)

[0157] To a stirred solution of compound **P3b** (290 mg, 1.0 mmol) in DCM (5 mL) at 0°C was added *m*-CPBA (610 mg, 3.0 mmol, 85%) and the resulting mixture was stirred at rt for 16 h, diluted with aq. sat. NaHCO_3 solution and extracted with EA (3×20 mL). The combined organic layer was dried over Na_2SO_4 , filtered, concentrated and purified by FCC (PE:EA = 5:1) to afford compound **P3** as a white solid.


Preparative Example P3-1

[0158]



Step 1: 4-Bromo-2-((2-ethoxy-2-oxoethyl)thio)-6-fluorobenzoic acid (P3-1a)


[0159]



[0160] To a mixture of 4-bromo-2,6-difluorobenzoic acid (10.0 g, 42.4 mmol) and ethyl 2-mercaptoacetate (5.10 g, 42.4 mmol) in DMF (100 mL) was added Cs_2CO_3 (41.5 g, 127 mmol) and the mixture was stirred at 80°C overnight, diluted with water (1 L) and adjusted to pH = 3 with 2M HCl and extracted with EA (3×300 mL). The combined organic layer was washed with brine (300 mL), dried over Na_2SO_4 , filtered, concentrated and purified by FCC (PE:EA = 1:1) to give compound **P3-1a** as a yellow oil.

Step 2: Ethyl 2-((5-bromo-3-fluoro-2-(hydroxymethyl)phenyl)thio)acetate (P3-1b)

[0161]




[0162] To the solution of compound **P3-1a** (4.10 g, 12.2 mmol) in THF (40 mL) was added B_2H_6 (24.4 mL, 1M in THF). This mixture was stirred at 70°C overnight, quenched with water (100 mL) and extracted with EA (4×40 mL). The combined organic layer was washed with brine (50 mL), dried over Na_2SO_4 , filtered, concentrated and purified by FCC (PE:EA = 5:1) to give compound **P3-1b** as a white solid.

Step 3: Ethyl 2-((5-bromo-3-fluoro-2-(hydroxymethyl)phenyl)sulfonyl)acetate (P3-1)

[0163] To a stirred solution of compound **P3-1b** (1.00 g, 3.40 mmol) in DCM (30 mL) at 0°C was added *m*-CPBA (1.80 g, 10.2 mmol, 85%) and the mixture was stirred at rt for 16 h, diluted with aq. sat. NaHCO₃ solution and extracted with EA (3 × 20 mL). The combined organic layer was dried over Na₂SO₄, concentrated and purified by FCC (PE:EA = 5:1) to give compound **P3-1** as a white solid.

Preparative Example P4


[0164]

Methyl 2-(3-bromophenyl)-2-methylcyclopropane-1-carboxylate (P4)

[0165] To a solution of compound **P16** (1.00 g, 3.92 mmol) in DMF (15 mL) was added MeI (1.11 g, 7.84 mmol) and K₂CO₃ (1.35 g, 9.80 mmol). The mixture was stirred for 2 h at 50°C, cooled, diluted with EA (100 mL) and washed with water (3 × 20 mL) and brine (20 mL), dried over Na₂SO₄, filtered, concentrated and purified by prep-TLC (EA:PE = 1:6) to give compound **P4** as a yellow oil.

Preparative Example P5

[0166]

Methyl 3'-bromo-2-chloro-[1,1'-biphenyl]-4-carboxylate (P5)





[0167] To a solution of methyl 3-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (47.7 g, 161 mmol) in dioxane (300 mL) was added 1-bromo-3-iodobenzene (50.0 g, 177 mmol), Na₂CO₃ (35.7 g, 337 mmol) and Pd(PPh₃)₄ (11.7 g, 10.1 mmol) under N₂. The mixture was stirred at 90°C overnight under N₂, cooled, filtered, concentrated and purified by FCC (EA:PE = 1:50) to give compound **P5** as a white solid.


Preparative Example P5/1 to P5/3

[0168] The following Preparative Examples were prepared similar as described for Preparative Example P5 using the appropriate building blocks.


#	building block(s)	structure
P511	P8	

(continued)

#	building block(s)	structure
P5/2		
P5/3		

Preparative Example P6**[0169]****4-(Methyl-*d*3)benzenesulfonyl chloride-2,3,5,6-*d*4 (P6)**

[0170] To a solution of toluene-*d*8 (1.00 g, 10.0 mmol) in DCM (10 mL) was added ClSO₃H (5 mL) and the mixture was stirred at rt for 2 h, poured into water (100 mL) and extracted with DCM (100 mL). The organic layer was concentrated to give compound **P6** as a white solid.


Preparative Example P7**[0171]****tert-Butyl 2-(3-bromo-4-cyanophenoxy)acetate (P7)**

[0172] A mixture of 2-(3-bromo-4-cyanophenoxy)acetic acid (200 mg 0.78 mmol), Boc₂O (204 mg 0.94 mmol), DMAP (10 mg, 80 μmol) and pyridine (0.4 mL) in tert-BuOH (10 mL) was stirred at rt overnight, concentrated and purified by FCC (PE:EA = 50:1) to give compound **P7** as a yellow oil.

Preparative Example P7/1


[0173] The following Preparative Example was prepared similar as described for Preparative Example P7 using the appropriate building block.

EP 3 814 331 B9



Preparative Example P8

[0174]




Methyl 5-chloro-2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (**P8**)

[0175] To a solution of methyl 4-bromo-5-chloro-2-fluorobenzoate (2.66 g, 10.0 mmol) in dioxane (30 mL) was added B_2Pin_2 (2.79 g, 11.0 mmol), KOAc (2.45 g, 25.0 mmol) and $Pd(dppf)Cl_2$ (260 mg) under N_2 . The mixture was stirred at 80°C overnight under N_2 , diluted with water (50 mL) and extracted with EA (3×50 mL). The combined organic layer was washed with brine, dried over Na_2SO_4 , filtered, concentrated and purified by FCC (EA:PE = 1:40) to give compound **P8** as a white solid.

Preparative Example P9

[0176]




Methyl 4-(3-bromophenyl)butanoate (**P9**)

[0177] To a solution of 4-(3-bromophenyl)butanoic acid (500 mg, 2.06 mmol) in DMF (50 mL) was added K_2CO_3 (569 mg, 4.11 mmol) and CH_3I (438 mg, 3.09 mmol). The mixture was stirred for 2 h at rt. Insoluble salts were filtered off and washed with EA. The combined organic layer was washed with water (3×50 mL), brine (2×50 mL), dried over Na_2SO_4 and filtered. The solvents were removed under reduced pressure to afford compound **P9** as a yellow solid, which was used in the next step without further purification.

Preparative Example P10

[0178]




Methyl 2-(3-bromophenoxy)acetate (**P10**)

[0179] To a solution of 3-bromophenol (1.72 g, 10.0 mmol) and methyl-bromoacetate (1.01 mL, 11.0 mmol) in ACN (60 mL) was added K_2CO_3 (2.07 g, 15.0 mmol) and the mixture was stirred at 50°C overnight. After insoluble salts are filtered off and washed with ACN, the solvent is removed under reduced pressure and the remainder is taken up in EA and washed subsequently with water and brine. The organic layer is dried over Na_2SO_4 , filtered and concentrated to

give compound **P10** as a colorless semi-solid.


Preparative Examples P10/1

[0180] The following Example was prepared similar as described for Preparative Example **P10** using the appropriate building blocks.



Preparative Example P11

[0181]




Methyl 3-((6-bromopyridin-2-yl)oxy)propanoate (**P11**)

[0182] To a solution of 6-bromopyridin-2(1*H*)-one (800 mg, 4.59 mmol) and PPh_3 (2.39 g, 9.19 mmol) in dry THF (30 mL) under N_2 was added DEAD (1.20 g, 6.89 mmol) and methyl 3-hydroxypropanoate (479 mg, 4.59 mmol). The mixture was stirred at rt overnight, quenched with sat. NH_4Cl (60 mL) and extracted with EA (2×30 mL). The combined organic layer was washed with brine (2×30 mL), dried over Na_2SO_4 , concentrated and purified by prep-TLC (EA:PE = 1:4) to give compound **P11** as a white solid.

Preparative Example P12

[0183]





Methyl 1-(3-bromophenyl)azetidine-3-carboxylate (**P12**)

[0184] To a solution of 1-bromo-3-iodobenzene (500 mg, 1.77 mmol) in dioxane (8 mL) was added methyl azetidine-3-carboxylate hydrochloride (295 mg, 1.94 mol), $\text{Pd}_2(\text{dba})_3$ (35 mg, 40 μmol), XPhos (17 mg, 40 μmol) and Na_2CO_3 (375 mg, 3.53 mmol). The mixture was stirred at 100°C overnight, cooled to rt, filtered, concentrated and purified by prep-TLC (PE:EA = 2:1) to give compound **P12** as a yellow oil.

Preparative Example P12/1


[0185] The following Preparative Example was prepared similar as described for Preparative Example **P12** using the appropriate building block.


**Preparative Example P13****[0186]****Methyl 1-(3-bromophenyl)-3-methylazetidine-3-carboxylate (P13)**

[0187] To a solution of 1-bromo-3-iodobenzene (500 mg, 1.77 mmol) in dioxane (15 mL) was added methyl 3-methylazetidine-3-carboxylate hydrochloride (293 mg, 1.77 mmol), Pd₂(dba)₃ (32 mg, 35 μmol), Xantphos (20 mg, 35 μmol) and Cs₂CO₃ (1.35 g, 3.54 mmol). The mixture was stirred at 100°C overnight under N₂, cooled to rt, diluted with water (150 mL) and extracted with EA (3 × 200 mL). The combined organic layer was washed with brine (2 × 50 mL), dried over Na₂SO₄, concentrated and purified by FCC (EA:PE = 1:5) to afford compound **P13** as a yellow oil.



Preparative Example P13/1 to P13/4

[0188] The following Preparative Examples were prepared similar as described for Preparative Example **P13** using the appropriate building blocks.




Preparative Example P14**[0189]****Methyl 1-(6-bromopyridin-2-yl)azetidine-3-carboxylate (P14)**

[0190] To a solution of 2,6-dibromopyridine (500 mg, 2.11 mmol) in DMF (20 mL) was added methyl azetidine-3-carboxylate hydrochloride (384 mg, 2.53 mmol) and K_2CO_3 (729 mg, 5.28 mmol) and the mixture was stirred overnight at 80°C. After cooling to rt insoluble salts were filtered off and washed with EA. The combined organic solvents were washed with water (3 × 50 mL), brine (2 × 50 mL), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by prep-TLC (PE:EA = 4:1) to afford compound **P14** as a yellow oil.

Preparative Example P15**[0191]****Step 1: 3-(3-Bromophenyl)-3-hydroxycyclobutane-1-carboxylic acid (P15a)****[0192]**

[0193] To a solution of 1-bromo-3-iodobenzene (2.82 g, 10.0 mmol) and 3-oxocyclobutane-1-carboxylic acid (1.14 g, 10.0 mmol) in THF (30 mL) at -78°C was added *n*-BuLi (8 mL, 20 mmol, 2.5 M in THF) and the mixture was stirred at -78°C for 4 h, quenched with NH_4Cl (50 mL), neutralized with 1N aq. HCl and extracted with EA (3 x). The combined organic layer was washed with brine, dried over Na_2SO_4 , filtered, concentrated and purified by FCC (EA:PE = 1:1) to give compound **P15a** as a colorless oil.

Step 2: Methyl 3-(3-bromophenyl)-3-hydroxycyclobutane-1-carboxylate (P15b)**[0194]**


[0195] To a solution of compound **P15a** (1.35 g, 5.00 mmol) in DMF (20 mL) was added K_2CO_3 (1.38 g, 10.0 mmol) and CH_3I (710 mg, 5.00 mmol) and the mixture was stirred at rt for 2 h. Water was added (200 mL) and the mixture was extracted with EA. The combined EA extracts were washed with brine, dried over Na_2SO_4 and filtered. The solvent was removed under reduced pressure and the residue was purified by FCC (EA:PE = 1:10) to give compound **P15b** as a colorless oil.

Step 3: Methyl 3-(3-bromophenyl)cyclobutane-1-carboxylate (**P15**)


[0196] To a solution of compound **P15b** (1.10 g, 3.90 mmol) in TFA (20 mL) at 0°C was added triethylsilane (680 mg, 5.85 mmol) and the mixture was stirred for 2 h. Water was added to the mixture (200 mL) and the mixture extracted with EA. The solvent was removed under reduced pressure and the residue was purified by FCC (EA:PE = 1:10) to give compound **P15** as a colorless oil.

Preparative Example P16

[0197]

Step 1: Methyl (E)-3-(3-bromophenyl)acrylate (**P16a**)

[0198]



[0199] (E)-3-(3-Bromophenyl)acrylic acid (3.00 g, 13.2 mmol) was dissolved in DMF (50 mL), Mel (3.75 g, 26.4 mmol) and K₂CO₃ (2.74 g, 19.8 mmol) were added and the mixture was stirred for 2 h at rt. After insoluble salts were filtered and washed with EA, the solvent was washed with water (3 × 50 mL), brine (2 × 50 mL), dried over Na₂SO₄, filtered and concentrated to give compound **P16a** as a yellow solid which was used in the next step without any purification.

Step 2: *rac*-Methyl (1*R*,2*R*)-2-(3-bromophenyl)cyclopropane-1-carboxylate (**P16**)

[0200] Under argon, NaH (60 %, 680 mg, 17.0 mmol) was initially charged in DMSO (30 mL) and trimethylsulphoxonium iodide (3.74 g, 17.0 mmol) was added in one portion at rt. After the evolution of gas had ceased, compound **P16a** (3.15 g, 13.1 mmol), dissolved in DMSO (10 mL), was slowly added drop-wise. After stirring overnight at 50°C, the mixture was partitioned between EA and water. The aq. layer was extracted with EA. The combined organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC (EA:PE = 1:20) to give compound **P16** as a colorless oil.


Preparative Example P16/1 to P16/2

[0201] The following Preparative Examples were prepared similar as described for Preparative Example **P16** using the appropriate building blocks.

#	building block	structure	chemical name
P16/1			<i>rac</i> -methyl (1 <i>R</i> ,2 <i>R</i>)-2-(5-bromothiophen-2-yl) cyclopropane-1-carboxylate
P16/2			<i>rac</i> -methyl (1 <i>R</i> ,2 <i>R</i>)-2-(3-bromo-5-chlorophenyl) cyclopropane-1-carboxylate

Preparative Example P17

[0202]



Methyl 3'-bromo-[1,1'-biphenyl]-3-carboxylate (**P17**)


[0203] To a solution of (3-bromophenyl)boronic acid (1.50 g, 7.47 mmol) in dioxane (30 mL) was added methyl 3-bromobenzoate (1.93 g, 8.96 mmol), Pd(PPh₃)₄ (173 mg, 0.15 mmol) and Na₂CO₃ (1.58 g, 14.9 mmol). The mixture was stirred at 100°C overnight. After cooling to rt the reaction was filtered, concentrated and purified by FCC to give compound **P17** as a yellow oil.

Preparative Examples P17/1 to P17/4

[0204] The following Examples were prepared similar as described for Preparative Example **P17** using the appropriate building blocks.


Preparative Example P18

[0205]



Step 1: Methyl 3'-((trimethylsilyl)ethynyl)-[1,1'-biphenyl]-4-carboxylate (**P18a**)


[0206]









[0207] Pd(PPh₃)₄ (1.98 g, 1.72 mmol), CuI (327 mg, 1.72 mmol) and PPh₃ (450 mg, 1.72 mmol) were combined in a round-bottom flask and the flask was degassed and refilled with N₂ three times. TEA (86 mL), methyl 3'-bromo-[1,1'-biphenyl]-4-carboxylate (**P5/2**, 5.00 g, 17.2 mmol) and ethynyltrimethylsilane (4.86 mL, 36.1 mmol) were added and the mixture was stirred at 60°C for 6 h. After filtration through kieselgur the filtrate was concentrated under reduced pressure to give compound **P18a** as a black solid, which was used in the next step without further purification.



Step 2: Methyl 3'-ethynyl-[1,1'-biphenyl]-4-carboxylate (**P18**)

[0208] To a solution of compound **P18a** (6.21 g, 17.2 mmol) in THF (25 mL) was added TBAF (25 mL, 1M in THF) and the mixture was stirred at rt for 3 h. After concentration under reduced pressure the residue was purified by FCC (EA:PE = 1:20) to give compound **P18** as a white solid.



(continued)


#	building block	structure
P17/2		
P17/3		
P17/4		

Preparative Example P19**[0209]****Step 1: 3-Bromofuran-2-carboxamide (P19a)****[0210]**

[0211] To a solution of 3-bromofuran-2-carboxylic acid (1.00 g, 5.24 mmol) in DMF (10 mL) was added HATU (2.98 g, 7.85 mmol) and DIPEA (1.69 g, 13.1 mmol) and the mixture was stirred at rt for 1 h. NH_4Cl (333 mg, 6.29 mmol) was added and stirring was continued overnight. Water (30 mL) was added, and the mixture was extracted with EA (3×30 mL). The combined organic layer was dried over Na_2SO_4 , concentrated and purified by FCC to give compound **P19a** as a yellow solid.



Step 2: 3-Bromofuran-2-carbonitrile (P19)

[0212] To a solution of compound **19a** (906 mg, 4.77 mmol) in DCM (10 mL) at 0°C was added TFAA (2.50 g, 11.9 mmol) and the mixture was stirred for 2 h, diluted with water (30 mL) and extracted with DCM (3×30 mL). The combined organic layer was dried over Na_2SO_4 , concentrated and purified by FCC (EA:PE = 1:20) to give compound **P19** as a white solid.


Preparative Example P20**[0213]**

2-Bromo-4-fluoro-6-methoxyaniline (P20)


[0214] NBS (12.4 g, 69.4 mmol) was added to a solution of 4-fluoro-2-methoxyaniline (8.90 g, 63.1 mmol) in dry DCM (217 mL) at -78°C and the mixture was stirred at -78°C for 2 h, then allowed to warm to 0°C and stirred for 2 h. The solvent was removed in vacuum and the resulting residue was purified by FCC (EA:PE = 1:10) to give compound **P20** as a yellow oil.

Preparative Example P21**[0215]**Step 1: 2-(4-Bromophenyl)-2-((trimethylsilyl)oxy)propanenitrile (P21a)**[0216]**

[0217] Trimethylsilyl cyanide (4.96 g, 50.0 mmol) and zinc iodide (50 mg) were added to 1-(4-bromophenyl)ethan-1-one (5.00 g, 50.0 mmol) in DCM (200 mL). This mixture was stirred for 5 h at rt. The mixture was washed with water (2 × 20 mL) and brine (20 mL). The organic layer was dried over Na₂SO₄ and concentrated to afford crude compound **P21a**, which was used in the next step without any purification.

Step 2: 2-(4-Bromophenyl)-2-hydroxypropanoic acid (P21b)**[0218]**

[0219] To the solution of compound **P21a** (12.2 g, 40.9 mmol) in AcOH (50 mL) was added conc. HCl (50 mL). The mixture was stirred overnight at rt and heated at 100°C for 2 h. The solvent was removed under reduced pressure. H₂O was added and the mixture was extracted with EA (3 × 200 mL). The organic layer was dried over Na₂SO₄ and concentrated to give crude compound **P21b** as a yellow oil, which was used in the next step without any purification.

Step 3: Methyl 2-(4-bromophenyl)-2-hydroxypropanoate (P21c)**[0220]**

[0221] To a solution of compound **P21b** (6.50 g, 26.5 mmol) in MeOH (60 mL) was added conc. H₂SO₄ (3 mL). The mixture was stirred overnight at rt. The solvent was removed under reduced pressure, dissolved in EA (300 mL) and washed with H₂O (30 mL) and sat. NaHCO₃ (30 mL). The organic layer was dried over Na₂SO₄, concentrated and


purified by FCC (EA:PE = 1:2) to give compound **P21c** as a colorless oil.

Step 4: Methyl 2-hydroxy-2-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)propanoate (**P21**)

[0222] To a solution of compound **P21c** (200 mg, 0.77 mmol) in dioxane (10 mL) was added B_2Pin_2 (209 mg, 0.93 mmol), KOAc (151 mg, 1.54 mmol) and $Pd(dppf)Cl_2$ (56 mg, 0.08 mmol). The mixture was stirred at 100°C overnight under N_2 . After cooling to rt, the mixture was filtered and the solvent was removed under reduced pressure. The residue was purified by prep-TLC (EA:PE = 1:1) to afford compound **P21** as a white solid.


Preparative Examples P2111

[0223] The following Example was prepared similar as described for Preparative Example **P21** using the appropriate building block.




Preparative Example P22 (mixture of 1- and 2-trityl isomer)

[0224]



Step 1: 5-(4-Bromo-3-chlorophenyl)-1H-tetrazole (**P22a**)

[0225]




[0226] To a solution of 4-bromo-3-chlorobenzonitrile (500 mg, 2.33 mmol) in DMF (10 mL) was added NaN_3 (1.50 g, 23.3 mmol) and NH_4Cl (1.20 g, 23.3 mmol). The mixture was stirred at 100°C under N_2 overnight. Then DCM (100 mL) was added and the mixture was washed with brine (30 mL). The organic layer was dried over Na_2SO_4 , concentrated and purified by FCC (EA:PE = 1:3) to give compound **P22a** as a white solid.

Step 2: 5-(4-Bromo-3-chlorophenyl)-1-trityl-1H-tetrazole (**P22**), (mixture of 1- and 2-trityl isomers)

[0227] To a solution of compound **P22a** (350 mg, 1.36 mmol) in DCM (50 mL) was added triphenylmethyl chloride (556 mg, 2.00 mmol) and TEA (202 mg, 2.00 mmol). The mixture was stirred at rt for 12 h. Then DCM (50 mL) was added and the mixture was washed with brine (30 mL). The organic layer was dried over Na_2SO_4 , concentrated and purified by FCC (EA:PE = 1:7) to afford compound **P22** as a white solid.


Preparative Example P23 (mixture of 1- and 2-trityl isomers)

[0228]



Step 1: N-((3-Bromophenyl)(methyl)(oxo)-λ⁶-sulfaneylidene)-2,2,2-trifluoroacetamide (**P23a**)


[0229]



[0230] To a solution of 1-bromo-3-(methylsulfinyl)benzene (950 mg, 4.38 mmol) in DCM (10 mL) was added MgO (697 mg, 17.4 mmol), 2,2,2-trifluoroacetamide (742 mg, 6.57 mmol), Rh₂(OAc)₄ (100 mg) and (diacetoxy)iodobenzene (2.82 g, 8.76 mmol). The mixture was stirred at 40°C overnight and filtered through a pad of Celite. The solvent was removed under reduced pressure and the crude product was purified by FCC (PE:EA = 1:2) to give compound **P23a** as a white solid.

Step 2: (3-Bromophenyl)(imino)(methyl)-λ⁶-sulfanone (**P23b**)


[0231]



[0232] To a stirred solution of compound **P23a** (680 mg, 2.07 mmol) in MeOH (5 mL) was added K₂CO₃ (713 mg, 5.17 mmol) and stirring was continued at rt for 1 h. Then water was added and the mixture was extracted with EA (3 × 20 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄ and concentrated to give compound **P23b** as a white solid.

Step 3: N-((3-Bromophenyl)(methyl)(oxo)-λ⁶-sulfaneylidene)cyanamide (**P23c**)


[0233]



[0234] To a solution of compound **P23b** (430 mg, 1.86 mmol) in DCM (5 mL) was added cyanic bromide (235 mg, 2.24 mmol) and TEA (376 mg, 3.72 mmol). The mixture was stirred at rt for 3 h, diluted with water and extracted with EA (3 × 20 mL). The combined organic layer was washed with sat. aq. NaHCO₃ (20 mL), dried over Na₂SO₄ and concentrated to give compound **P23c** as a yellow solid.

Step 4: ((1*H*-Tetrazol-5-yl)imino)(3-bromophenyl)(methyl)-λ⁶-sulfanone (**P23d**)

[0235]




[0236] To a stirred solution of compound **P23c** (420 mg, 1.63 mmol) in DMF (5 mL) was added NaN_3 (1.06 g, 16.3 mmol) and NH_4Cl (864 mg, 16.3 mmol). The mixture was stirred and heated to 100°C overnight. After cooling to rt, water was added and the mixture was extracted with EA (3×20 mL). The combined organic layer was dried over Na_2SO_4 , filtered, concentrated and purified by prep-TLC (PE:EA = 1:1) to give compound **P23d** as a white solid.

Step 5: (3-Bromophenyl)(methyl)((1-trityl-1H-tetrazol-5-yl)imino)- λ^6 -sulfanone (**P23**) (mixture of 1- and 2-trityl isomer)

[0237] To a stirred solution of compound **P23d** (350 mg, 1.16 mmol) in DCM (20 mL) was added trityl chloride (388 mg, 1.39 mmol) and TEA (0.3 mL, 2.3 mmol). Stirring was continued at rt overnight. Then water was added and the mixture was extracted with DCM (3×50 mL). The combined organic layer was washed with brine (20 mL), dried over Na_2SO_4 , concentrated and purified by FCC (EA:PE = 1:3) to give compound **P23** as a white solid.

Preparative Example P24

[0238]




rel-Methyl (1R,3r,5S)-8-azabicyclo[3.2.1]octane-3-carboxylate hydrochloride (**P24**)

[0239] rel-(1R,3r,5S)-8-(tert-Butoxycarbonyl)-8-azabicyclo[3.2.1]octane-3-carboxylic acid (500 mg, 1.96 mmol) was dissolved in HCl in MeOH (20 mL). The solution was stirred at rt for 5 h. The solvent was removed under reduced pressure to afford compound **P24** as a white solid.


Preparative Example P25

[0240]



Step 1: (4-Bromo-2-(methylsulfonyl)phenyl)methanol (**P25a**)

[0241]




[0242] To a solution of methyl 4-bromo-2-(methylsulfonyl)benzoate (3.00 g, 10.2 mmol) in MeOH (20 mL) was added

LiBH₄ (4.00 g, 100 mmol) slowly at 0°C. The mixture was stirred at 80°C overnight. Water (40 mL) was added slowly under cooling with an ice bath and the mixture was extracted with EA (3 × 30 mL). The combined organic layer was washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give compound **P25a** as a pale yellow solid, which was directly used in the next step.

Step 2: 2-(4-Bromo-2-(methylsulfonyl)phenyl)acetonitrile (**P25b**)


[0243]



[0244] To a solution of cyanic bromide (712 mg, 6.70 mmol) and PPh₃ (1.76 g, 6.70 mmol) in DCM (30 mL) was added a solution of compound **P25a** (1.50 g, 5.60 mmol) in DCM (50 mL). The mixture was stirred at 15°C for 1 h, then DBU (1.10 g, 6.70 mmol) was added at 0°C. The resulting mixture was stirred at 0~15°C for another 16 h. The solvent was concentrated in vacuum. The residue was purified by FCC (PE:EA = 4:1) to give compound **P25b** as a yellow solid.

Step 3: 2-(4-Bromo-2-(methylsulfonyl)phenyl)-2-methylpropanenitrile (**P25c**)

[0245]




[0246] To a solution of compound **P25b** (200 mg, 1.10 mmol) in THF (20 mL) were added potassium tert-butoxide (502 mg, 4.40 mmol) and iodomethane (624 mg, 4.40 mmol) at -78°C. The mixture was warmed to -20°C and stirred overnight, diluted with aq. NH₄Cl (30 mL) and extracted with EA (3 × 30 mL). The combined organic layer was washed with brine (100 mL), dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE:EA = 100:8) to give compound **P25c** as a yellow solid.

Step 4: 2-(4-Bromo-2-(methylsulfonyl)phenyl)-2-methylpropanoic acid (**P25**)

[0247] To a solution of compound **P25c** (850 mg, 2.80 mmol) in EtOH (5 mL) and H₂O (5 mL) was added KOH (1.20 g, 22.4 mmol). The mixture was stirred at 80°C for 2 d. The pH was adjusted to ca. 5 by addition of 1N aq. HCl and the mixture was extracted with DCM/MeOH (10/1, 3 × 40 mL). The combined organic layer was washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated to give compound **P25** as a yellow solid.

Preparative Example P26

[0248]




4-Bromo-3-(trifluoromethyl)-1-trityl-1H-pyrazole (**P26**)

[0249] To a stirred solution of 4-bromo-5-(trifluoromethyl)-1H-pyrazole (428 mg, 2.00 mmol) in DCM (10 mL) was added TEA (606 mg, 6.00 mmol) and (chloromethanetriyl)tribenzene (1.11 g, 4.00 mmol) and stirring was continued at rt overnight. Then the solvent was removed and H₂O (50 mL) was added and the mixture was extracted with EA (3 × 50 mL). The combined organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by FCC

(PE:EA =5:1) to give compound **P26** as a white solid.


Preparative Example P27

[0250]



Step 1: *tert*-Butyl 3-(2-bromophenyl)-3-hydroxyazetidine-1-carboxylate (**P27a**)

[0251]




[0252] To a solution of 1-bromo-2-iodobenzene (8.43 g, 30.0 mmol) in THF (50 mL) at -78°C was slowly added *i*-PrMgBr in THF (0.90M, 33 mL, 30.0 mmol). After stirring for 2 h, a solution of *tert*-butyl 3-oxoazetidine-1-carboxylate (3.20 g, 19.0 mmol) in THF (20 mL) was added dropwise to the mixture at -78°C. The mixture was stirred at rt for 3 h, diluted with sat. aq. NH₄Cl and extracted with EA. The organic layer was washed with water and brine, dried over Na₂SO₄, concentrated and purified by FCC (PE:DCM = 2:1) to afford compound **P27a** as a white solid.

Step 2: *tert*-Butyl 3-(2-bromophenyl)-3-fluoroazetidine-1-carboxylate (**P27**)

[0253] To a stirred solution of compound **P27a** (4.30 g, 13.1 mmol) in DCM (50 mL) at 0°C was slowly added DAST (4.20 g, 26.2 mmol). After stirring for 4 h, the mixture was poured into water and extracted with EA. The organic layer was washed with brine, dried over Na₂SO₄, concentrated and purified by FCC (PE:DCM = 3:1) to give compound **P27** as a colorless oil.


Preparative Example P28

[0254]



Step 1: 3-(3-Bromothiophen-2-yl)oxetan-3-ol (**P28a**)

[0255]




[0256] To a suspension of 3-bromothiophene (13.0 g, 80.2 mmol) in THF (20 mL) was added LDA (48.0 mL, 2.0M in THF, 96.0 mmol) under N₂ at -60°C. The mixture was stirred at -60°C for 45 min. Then oxetan-3-one (8.70 g, 121 mmol) was added and stirring was continued for 30 min at -60°C. Water was added slowly and the mixture was extracted with EA (3 x). The combined organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE:EA = 10:1 to 5:1) to give compound **P28a** as a brown oil.

Step 2: 3-(3-Bromothiophen-2-yl)oxetane (P28)


[0257] To a mixture of compound **P28a** (17.0 g, 72.6 mmol) in DCM (120 mL) was added $\text{BF}_3 \cdot \text{Et}_2\text{O}$ (18.5 mL, 146 mmol) at 0°C under N_2 . The mixture was stirred at 0°C for 30 min. Then triethylsilane (35.0 mL, 220 mmol) was added and the mixture was stirred at 0°C for 30 min. Further triethylsilane (35.0 mL, 220 mmol) was added and the mixture was stirred at 0°C for 30 min. A third portion of triethylsilane (35.0 mL, 220 mmol) was added and stirring was continued at 0°C 30 min. The mixture was added to a solution aq. NaOH (10%, 200 g) under cooling with an ice bath and extracted with EA (3 x). The combined organic layer was dried over Na_2SO_4 , filtered, concentrated and purified by FCC (PE/DCM = 10:1 to 3:1) to give compound **P28** as a yellow oil. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz) δ : 7.23 (d, J = 5.2 Hz, 1H), 6.94 (d, J = 5.2 Hz, 1H), 5.08-5.04 (m, 2H), 4.80-4.77 (m, 2H), 4.67-4.59 (m, 1H).

Preparative Example P29

[0258]

Step 1: 2-Bromocyclohex-1-ene-1-carboxamide (P29a)

[0259]




[0260] To a solution of 2-bromocyclohex-1-ene-1-carboxylic acid (1.20 g, 5.88 mmol) in DCM (20 mL) was added HATU (3.35 g, 8.82 mmol), DIPEA (2.16 g, 16.7 mmol) and NH_4Cl (3.20 g, 58.9 mmol). The mixture was stirred at rt for 24 h, filtered, concentrated and purified by FCC (PE:EA = 1:1) to give compound **P29a** as a colorless oil.

Step 2: 2-Bromocyclohex-1-ene-1-carbonitrile (P29)


[0261] To a solution of compound **P29a** (510 mg, 2.51 mmol) in DCM (20 mL) was added TFAA (1.05 g, 5.02 mmol) at 0°C. The mixture was stirred at rt for 12 h, poured into water (50 mL) and extracted with DCM (3 × 20 mL). The combined the organic layer was washed with brine (30 mL), dried over Na_2SO_4 , filtered, concentrated and purified by FCC (PE:EA = 2:1) to give compound **P29** as a white solid.

Preparative Example P30

[0262]

Step 1: Methyl 2-chloro-3'-((trimethylsilyl)ethynyl)-[1,1'-biphenyl]-4-carboxylate (P30a)

[0263]



EP 3 814 331 B9


[0264] Pd(PPh₃)₄ (553 mg, 0.48 mmol), CuI (93 mg, 0.48 mmol) and PPh₃ (126 mg, 0.48 mmol) were combined in a round-bottom flask, then degassed and refilled with N₂ three times. To the mixture was added TEA (45 mL), compound **P5** (2.00 g, 6.10 mmol), ethynyltrimethylsilane (786 mg, 10.2 mmol) and then the mixture was stirred at 60°C for 6 h, cooled, filtered through kieselguhr and washed with EA (40 mL). The filtrate was concentrated and purified by FCC (PE:EA = 20:1) to give compound **P30a** as a yellow solid.

Step 2: Methyl 2-chloro-3'-ethynyl-[1,1'-biphenyl]-4-carboxylate (**P30**)

[0265] To a solution of compound **P30a** (2.05 g, 5.89 mmol) in MeOH (5 mL) was added K₂CO₃ (778 mg, 7.07 mmol) and the mixture was stirred at rt for 30 min, poured into ice water (50 mL) and extracted with DCM (2 × 50 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated to give compound **P30** as a yellow solid.

Preparative Example P31

[0266]




1-(2-Chloropyridin-3-yl)azetidin-3-ol (**P31**)

[0267] To a solution of 2-chloro-3-iodopyridine (1.20 g, 5.00 mmol) in toluene (20 mL) was added azetidin-3-ol hydrochloride (1.09 g, 10.0 mmol), Cs₂CO₃ (6.52 g, 20.0 mmol), BINAP (311 mg, 0.50 mmol) and Pd₂(dba)₃ (200 mg) under N₂. The mixture was stirred at 110°C overnight under N₂. After cooling to rt the mixture was filtered and the solvent was removed under reduced pressure. The residue was purified by FCC (EA:PE = 1:3) to give compound **P31** as a yellow solid.

Preparative Example P32

[0268]





Ethyl 2-(4-bromophenyl)-2-hydroxypropanoate (**P32**)

[0269] To a solution of ethyl 2-(4-bromophenyl)-2-oxoacetate (512 mg, 2.00 mmol) in THF (30 mL) was added MeMgBr (2 mL, 1M in THF) at 0°C. The mixture was stirred at 0°C for 1 h, diluted with water (50 mL) and extracted with EA (3 × 50 mL). The combined organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE:EA = 5:1) to give compound **P32** as a white solid.



Preparative Examples P32/1

[0270] The following Preparative Example was prepared similar as described for Preparative Example **P32** using the appropriate building block.



Preparative Example P33**[0271]**4-Bromo-3-chlorobenzenesulfonic acid (P31)


[0272] A solution of 4-bromo-3-chlorobenzenesulfonyl chloride (576 mg, 2.00 mmol) in H₂O (30 mL) was stirred at 100°C for 16 h and concentrated to give compound **P33** as a white solid.

Preparative Example P34**[0273]**Step 1: N-((4-Bromo-3-chlorophenyl)sulfonyl)acetamide (P34a)**[0274]**

[0275] 4-Bromo-3-chlorobenzenesulfonamide (1.5 g, 5.5 mmol) was dissolved in pyridine (5 mL). Then DMAP (22 mg, 0.18 mmol) and Ac₂O (1.1 mL, 12 mmol) were added and the mixture was stirred for 3 h at rt, diluted with EA and washed with aq. NH₄Cl solution (3 x) and water. The organic layer was dried over Na₂SO₄ and concentrated. The resulting oil was triturated with PE and the precipitate was collected by filtration to afford compound **P34a** as a white solid.


Step 2: N-((3-Chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)sulfonyl)acetamide (P34)

[0276] To a solution of compound **P34a** (310 mg, 1.00 mmol) in dioxane (5 mL) was added B₂Pin₂ (381 mg, 1.50 mmol), KOAc (276 mg, 2.00 mmol) and Pd(dppf)Cl₂ (120 mg). The mixture was stirred under N₂ at 90°C for 8 h, cooled, filtrated, concentrated and purified by FCC (PE:EA = 5:1) to give compound **P34** as a yellow solid.

Preparative Example P35**[0277]**

Step 1: 4-Bromopyridine-3,5-dicarboxylic acid (P35a)**[0278]**

5




10 **[0279]** To a solution of 4-bromo-3,5-dimethylpyridine (1.24 g, 6.72 mmol) in water (15 mL) was added KMnO_4 (1.59 g, 10.1 mmol) and the mixture was stirred at 100°C for 1 h. Then an additional amount of KMnO_4 (1.59 g, 10.1 mmol) in water (15 mL) was added and stirring at 100°C was continued for 2 h. Then the mixture was filtered and the solvent concentrated to about 5 mL, adjusted to pH = 2 with conc. HCl and concentrated to give compound **P35a** as a white solid.

15 Step 2: 4-Chloropyridine-3,5-dicarboxamide (P35b)

[0280]

20



25 **[0281]** To a solution of compound **P35a** (1.30 g, 5.30 mmol) in DCM (15 mL) was added SOCl_2 (1.5 mL) and DMF (3 drops). The mixture was stirred at 45°C for 2 h, concentrated and redissolved in dioxane (5 mL). $\text{NH}_3 \cdot \text{H}_2\text{O}$ (20 mL) was added dropwise to the solution at 0°C and then concentrated to give compound **P35b** as a yellow solid.


Step 3: 4-Chloropyridine-3,5-dicarbonitrile (P35)

30 **[0282]** To a solution of compound **P35b** (188 mg, 0.94 mmol) in DMF (5 mL) was added POCl_3 (1 mL) and the mixture was stirred at rt overnight, diluted with water (30 mL) and extracted with EA (3 × 50 mL). The combined organic layer was washed with aq. NaHCO_3 (30 mL), concentrated and purified by FCC (PE:EA = 5:1) to give compound **P35** as a white solid.

35 **[0283]** General remarks: A "C" before the example number means that it is a comparative example while a "P" before the example number means that the example contains a protection group. These examples are not falling within the scope of the claims.

Example 1**[0284]**


45



50

Step 1: 2-((3-Bromophenyl)ethynyl)-4-fluoroaniline (1a)**[0285]**


55



[0286] To a solution of 1-bromo-3-iodobenzene (5.00 g, 17.7 mmol) in Et₃N (50 mL) was added Pd(PPh₃)₄ (1.22 g, 1.06 mmol), CuI (269 mg, 1.41 mmol), PPh₃ (278 mg, 1.06 mmol) and 2-ethynyl-4-fluoroaniline (2.86 g, 21.2 mmol). The mixture was stirred at 60°C under N₂ for 4 h, cooled, filtered, concentrated and purified by FCC (PE:EA = 8:1) to give compound **1a** as a yellow solid.

Step 2: N-(2-((3-Bromophenyl)ethynyl)-4-fluorophenyl)-4-methylbenzenesulfonamide (**1b**)

[0287]



[0288] To a solution of compound **1a** (3.50 g, 12.1 mmol) in DCM (50 mL) was added pyridine (3.5 mL), 4-methylbenzene-1-sulfonyl chloride (4.58 g, 24.1 mmol) and DMAP (350 mg). The mixture was stirred at rt overnight, diluted with CH₂Cl₂ (300 mL) and subsequently washed with 2N HCl (3 × 30 mL) and brine (30 mL). The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE:EA = 5:1) to give compound **1b** as a white solid.

Step 3: 3-(2-((3-Bromophenyl)-5-fluoro-1-tosyl-1*H*-indol-3-yl)thiophene-2-carbonitrile (**1**)


[0289] To a solution of compound **1b** (4.20 g, 9.48 mmol) in CH₃CN (60 mL) was added 3-bromothiophene-2-carbonitrile (3.67 g, 14.2 mmol), K₂CO₃ (2.62 g, 10.0 mmol) and Pd(PPh₃)₄ (1.09 g, 0.95 mmol) under N₂. The mixture was stirred at 100°C for 2 h, cooled, poured into EA (400 mL) and washed with water (50 mL) and brine (50 mL). The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC (EA:PE = 1:3) to give compound **1** as a white solid.

Example 1/1 to 1/149












[0290] The following Examples were prepared similar as described for Example 1 using the appropriate building blocks.

#	building block(s)	structure	analytical data
---	-------------------	-----------	-----------------













1/1













(continued)

#	building block(s)	structure	analytical data
5			
10	1/2  P5/2		
15			
20	1/3 		
25			
30	1/4 		
35			
40	1/5  P4	 first eluting isomer retention time 1.67 min	Separation of both isomers under the following conditions: Instrument: SFC-80 (Thar, Waters) Column: OJ 20×250 mm, 10 μm (Daicel) Column temperature: 35°C Mobile phase: CO ₂ /MeOH (0.2% NH ₄ ⁺ OMe ⁻) = 70/30 Flow rate: 80 g/min Back pressure: 100 bar Detection wavelength: 214 nm Cycle time: 2 min Sample solution: 180 mg dissolved in 30 mL MeOH Injection volume: 1 mL
45	1/6	 second eluting isomer retention time 2.16 min	
50			
55	1/7  P12		













(continued)

#	building block(s)	structure	analytical data
5			
1/8	 P12		
10			
1/9	 P12		
15			
1/10	 P12		
20			
1/11	 P12		
25			
1/12	 P12		
30			
1/13	 P12		
35			
40			
45			
50			
55			













(continued)

#	building block(s)	structure	analytical data
5 10 15 20	 P12		
25 30 35 40	 P12		
45 50 55	 P12		
	 P12		
	 P12		











(continued)

#	building block(s)	structure	analytical data
5 10 15 20 25 30 35	1/19 		
15 20 25 30 35	1/20 		
25 30 35	1/21 		
30 35 40 45	1/22  P16a		¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.25 (d, J = 8.5 Hz, 1H), 8.02 (d, J = 5.5 Hz, 1H), 7.57 (dd, J = 3.0, 5.0 Hz, 1H), 7.49 (dd, J = 4.5, 8.5 Hz, 1H), 7.45-7.43 (m, 3H), 7.36 (d, J = 4.0 Hz, 2H), 7.32 (d, J = 8.5 Hz, 2H), 7.11 (dd, J = 1.3, 4.8 Hz, 1H), 6.98 (d, J = 5.0 Hz, 1H), 2.31 (s, 3H); MS: 482.7 (M+Na) ⁺ .
40 45 50 55	1/23 		
50 55	1/24 		











(continued)

#	building block(s)	structure	analytical data
5 1/25			
15 1/26			$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.21 (d, $J = 8.5$ Hz, 1H), 7.99 (d, $J = 5.0$ Hz, 1H), 7.55-7.51 (m, 2H), 7.46 (d, $J = 7.5$ Hz, 1H), 7.40 (t, $J = 7.5$ Hz, 1H), 6.99 (d, $J = 5.0$ Hz, 1H), 6.94-6.89 (m, 4H), 2.23 (s, 3H), 2.06 (s, 6H); MS: 510.8 ($\text{M}+\text{Na}$) $^+$.
20 1/27			$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.33 (d, $J = 8.0$ Hz, 1H), 8.24 (d, $J = 8.0$ Hz, 1H), 8.08 (d, $J = 8.0$ Hz, 1H), 8.02 (d, $J = 8.0$ Hz, 1H), 7.98 (d, $J = 5.0$ Hz, 1H), 7.70 (d, $J = 7.5$ Hz, 1H), 7.63-7.43 (m, 7H), 7.05-6.95 (m, 3H); MS: 519.3 ($\text{M}+\text{Na}$) $^+$.
30 1128			$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.20 (d, $J = 10.5$ Hz, 1H), 8.02 (d, $J = 6.5$ Hz, 1H), 7.81 (s, 1H), 7.78 (s, 1H), 7.67-7.66 (m, 1H), 7.50-7.30 (m, 4H), 7.09 (dd, $J = 4.5, 6.5$ Hz, 1H), 7.04 (d, $J = 6.0$ Hz, 1H), 3.63 (s, 3H); MS: 450.8 ($\text{M}+1$) $^+$.
40 1/29			
50 1/30			













(continued)

#	building block(s)	structure	analytical data
5 1/31	 P5		
15 1/32	 P5		
25 1/33	 P5		
35 1/34	 P5		
45 1/35	 P5		











(continued)

#	building block(s)	structure	analytical data
1/36	 P5		
1/37	 P5		
1/38	 P5		
1/39			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.16 (d, J = 9.5 Hz, 1H), 8.04 (d, J = 5.0 Hz, 1H), 7.74 (d, J = 4.5 Hz, 1H), 7.60 (d, J = 8.5 Hz, 2H), 7.49 (d, J = 8.5 Hz, 2H), 7.22-7.21 (m, 1H), 7.16-7.12 (m, 2H), 7.03 (d, J = 5.5 Hz, 1H), 6.78 (s, 1H), 3.76 (s, 3H); MS: 510.8 (M+1) ⁺ .
1/40			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.03 (d, J = 5.0 Hz, 1H), 7.76 (d, J = 1.5 Hz, 1H), 7.72 (d, J = 5.0 Hz, 1H), 7.61 (d, J = 8.5 Hz, 2H), 7.55 (d, J = 8.5 Hz, 2H), 7.27 (d, J = 9.0 Hz, 1H), 7.19-7.17 (m, 1H), 7.11-7.09 (m, 1H), 7.05 (dd, J = 2.5, 8.5 Hz, 1H), 7.01 (d, J = 5.5 Hz, 1H), 3.91 (s, 3H); MS: 510.8 (M+1) ⁺ .











(continued)

#	building block(s)	structure	analytical data
5 1/41			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.82 (d, 4.0 Hz, 1H), 8.62 (s, 1H), 8.30 (d, J = 8.0 Hz, 1H), 8.01 (d, J = 4.5 Hz, 1H), 7.87 (d, J = 8.5 Hz, 1H), 7.58-7.53 (m, 2H), 7.46-7.35 (m, 5H), 7.30 (d, J = 7.0 Hz, 2H), 6.97 (d, J = 4.5 Hz, 1H); MS: 441.9 (M+1) ⁺ .
10 1/42			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.19 (d, J = 8.5 Hz, 1H), 8.05-8.02 (m, 2H), 7.57-7.54 (m, 1H), 7.46-7.33 (m, 7H), 6.97 (d, J = 5.0 Hz, 1H), 2.63 (s, 3H); MS: 462.1 (M+1) ⁺ .
15 1/43			
20 1/44			
25 1/45			
30 1/46			













(continued)

#	building block(s)	structure	analytical data
5 1/47			
15 1/48			
25 1/49			
35 1/50			
45 1/51			











(continued)

#	building block(s)	structure	analytical data
5			
10	 P5/1		
15			
20	 P5		
25			
30	 P5		
35			
40	 P5		
45			
50	 P16		
55			













(continued)

#	building block(s)	structure	analytical data
5			
10	1/57  P17		
15	1/58  P17/1		
20			
25	1/59  P17/2		
30			
35	1/60  P16		
40			
45	1/61  P9		
50			
55	1/62  P11		















(continued)

#	building block(s)	structure	analytical data
5			
10	1/63  P14		
15			
20	1/64  P12		
25			
30	1/65  P16/1		
35			
40	1/66  P16/2		
45			
50	1/67  P16		
55			















(continued)

#	building block(s)	structure	analytical data
5			
10	 P12/1		
15	 P16		
20			
25	 P16		
30			
35	 P16		
40			
45	 P16		
50	 P16		
55			

(continued)














#	building block(s)	structure	analytical data
1/74	 P12		
1/75	 P12  P19		
1/76	 P12 		
1/77	 P12 		
C1/78	 P12 		

(continued)













#	building block(s)	structure	analytical data
5			
1/79	 P12		
10			
15	 P12		
20			
25			
1/81	 P12		
30			
35			
1/82	 P12		
40			
45			
1/83	 P12		
50			
55			

55











(continued)

#	building block(s)	structure	analytical data
5			
1/89	 P13/2		
10			
1/90	 P13/3		
15			
1/91	 P12  P2/8		
20			
1/92	 P12  P2/4		
25			
1/93	 P12  P2/5		
30			
35			
40			
45			
50			
55			











(continued)

#	building block(s)	structure	analytical data
1/94	 P12  P2/6		
1/95	 P12  P2/7		
1/96	 P17/3		
1/97	 P5		
1/98	 P17/4		











(continued)

#	building block(s)	structure	analytical data
5			
1/99	 P13/4		
1/ 100	 P10 P16		
1/ 101	 P20 P16		
1/ 102	 P5		
1/ 103	 P5		














(continued)

#	building block(s)	structure	analytical data
1/ 104	 P5		
1/ 105	 P5		
1/ 106	 P5		
1/ 107	 P5		
1/ 108	 P5		











(continued)

#	building block(s)	structure	analytical data
5			
1/ 109	 P5		
15			
1/ 110	 P5		
25			
1/ 111	 P5		
35			
1/ 112	 P13/1		
45			
P11 113	 P26 P5		
55			

(continued)

#	building block(s)	structure	analytical data
5			
1/ 114	 P27  P5		
15			
1/ 115	 P27  P5		
25			
1/ 116	 P28  P5		
35			
1/ 117	 P5		
45			
1/ 118	 P5		
55			

(continued)

#	building block(s)	structure	analytical data
1/ 119	 P5		
1/ 120	 P5		
1/ 121	 P5		
1/ 122	 P5		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.41 (dd, $J = 9.2, 4.4$ Hz, 1H), 8.09 (d, $J = 1.6$ Hz, 1H), 8.00 (dd, $J = 8.0, 1.6$ Hz, 1H), 7.60-7.15 (m, 13H), 6.73 (dd, $J = 2.4, 8.4$ Hz, 1H), 6.71 (t, $J = 55.6$ Hz, 1H), 3.96 (s, 3H), 1.88 (s, 3H).
C1 /123	 P5		

(continued)

#	building block(s)	structure	analytical data
5			
10	1/ 124		
15			
20	1/ 125		
25			
30	1/ 126		
35			
40	1/ 127		
45			
50	1/ 128		
55			

(continued)

#	building block(s)	structure	analytical data
5			
10	1/ 129		
15			
20	1/ 130		
25			
30	1/ 131		
35			
40	1/ 132		
45			
50	1/ 133		
55			

(continued)

#

building block(s)



structure

analytical data

5

**C1/
134**



10



15

1/ 135



20



25

1/ 136



30



35

1/ 137



40



45











1/ 138

50













55

(continued)

#	building block(s)	structure	analytical data
1/ 139			$^1\text{H-NMR}$ (500 MHz, DMSO- d_6) δ : 8.32 (dd, $J = 9.3, 4.3$ Hz, 1H), 8.24 (d, $J = 8.5$ Hz, 2H), 7.80 (t, $J = 8.0$ Hz, 1H), 7.72-7.59 (m, 5H), 7.45 (dt, $J = 2.5, 9.0$ Hz, 1H), 7.35-7.24 (m, 2H), 7.23 (d, $J =$ 2.5 Hz, 1H), 7.25 (d, $J = 8.5$ Hz, 1H), 7.08 (t, $J = 55.0$ Hz, 1H).
1/ 140			
1/ 141			
1/ 142			
1/ 143			




(continued)

#	building block(s)	structure	analytical data
1/ 144			
1/ 145			
1/ 146			
1/ 147			
1/ 148			


(continued)

#	building block(s)	structure	analytical data
---	-------------------	-----------	-----------------

1/ 149

**Example 2****[0291]**

Step 1: 3-(5-Fluoro-2-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1-tosyl-1H-indol-3-yl)thiophene-2-carbonitrile (**2a**)

[0292]











[0293] To a solution of compound **1** (2.61 g, 4.73 mmol) in dioxane (40 mL) was added B₂Pin₂ (1.44 g, 5.68 mmol), KOAc (928 mg, 9.46 mmol) and Pd(dppf)Cl₂ (344 mg, 0.47 mmol). The mixture was stirred at 80°C overnight under N₂, cooled, filtered, concentrated and purified by FCC (EA:PE = 1:3) to give compound **2a** as a white solid.

Step 2: 3-(2-(2'-Chloro-4'-((dimethylamino)methyl)-[1,1'-biphenyl]-3-yl)-5-fluoro-1-tosyl-1H-indol-3-yl)thiophene-2-carbonitrile (**2**)


















[0294] To a solution of compound **2a** (150 mg, 0.25 mmol) in dioxane (8 mL) was added 1-(4-bromo-3-chlorophenyl)-N,N-dimethylmethanamine (65 mg, 0.26 mmol), Cs₂CO₃ (163 mg, 0.50 mmol) and Pd(PPh₃)₄ (30 mg, 25 pmol). The mixture was stirred at 100°C overnight under N₂, cooled, filtered, concentrated and purified by prep-HPLC to give compound **2** as a white solid. ¹H-NMR (500 MHz, DMSO-d₆) δ: 8.29 (dd, J = 9.5, 4.5 Hz, 1H), 8.02 (d, J = 5.0 Hz, 1H), 7.51-7.32 (m, 8H), 7.26-7.17 (m, 4H), 7.02-6.99 (m, 2H), 3.42 (s, 2H), 2.21 (s, 3H), 2.18 (s, 6H); MS: 639.9 (M+1)⁺.

Example 2/1 to 2/34











[0295] The following Examples were prepared similar as described for Example 2 (and optionally for Example 1) using the appropriate building blocks.

#	building block(s)	structure	analytical data
2/1			$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.41 (dd, $J = 9.0, 4.5$ Hz, 1H), 7.83 (d, $J = 5.5$ Hz, 1H), 7.65 (d, $J = 1.5$ Hz, 1H), 7.50-7.43 (m, 4H), 7.35-7.26 (m, 4H), 7.20-7.15 (m, 3H), 7.05 (dd, $J = 8.5, 2.5$ Hz, 1H), 6.96 (d, $J = 5.0$ Hz, 1H), 4.19 (s, 2H), 2.29 (s, 3H); MS: 611.8 ($\text{M}+1$) $^+$.
2/2			$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.41 (dd, $J = 9.0, 4.5$ Hz, 1H), 7.81 (d, $J = 5.0$ Hz, 1H), 7.64 (d, $J = 2.0$ Hz, 1H), 7.52-7.41 (m, 4H), 7.29-7.25 (m, 4H), 7.16-7.14 (m, 2H), 7.06 (dd, $J = 8.5, 3.0$ Hz, 1H), 6.97 (s, 1H), 6.93 (d, $J = 5.0$ Hz, 1H), 2.25 (s, 3H), 1.54 (s, 6H); MS: 639.0 ($\text{M}+1$) $^+$.
2/3			
2/4			
2/5			











(continued)

#	building block(s)	structure	analytical data
5	 P2/2		
10			
15			
20			
25			
30			
35			
40			
45			
50			
55			











(continued)

#	building block(s)	structure	analytical data
2/12	P16 		
2/13			
2/14			
P2/15	 P22		
2/16	 prepared according to ACS Med. Chem. Lett. 2016;7:1207		¹H-NMR (500 MHz, DMSO-d₆) δ: 8.34-8.29 (m, 1H), 8.01 (d, J = 5.0 Hz, 1H), 7.96 (d, J = 1.5 Hz, 1H), 7.86-7.84 (m, 2H), 7.61 (s, 1H), 7.53 (t, J = 7.5 Hz, 1H), 7.41-7.28 (m, 6H), 7.21-7.18 (m, 1H), 7.08 (d, J = 5.5 Hz, 1H), 5.59-5.55 (m, 1H), 4.96-4.94 (m, 2H), 3.45 (s, 3H), 2.27 (s, 3H); MS: 691.8 (M+18)⁺.

(continued)

#	building block(s)	structure	analytical data
P2/17	 P23		
2/18	 P25		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.44-8.41 (m, 1H), 8.16 (s, 1H), 7.85-7.76 (m, 4H), 7.51-7.48 (m, 1H), 7.39-7.26 (m, 5H), 7.19-7.17 (m, 2H), 7.06-7.01 (m, 2H), 3.23 (s, 3H), 2.26 (s, 3H), 1.77 (s, 6H); MS: 730.1 ($\text{M}+18$) $^+$.
2/19			
2/20			
2/21			

(continued)

#	building block(s)	structure	analytical data
2/22			
2/23			
2/24			
2/25			
2/26			



(continued)

#	building block(s)	structure	analytical data
2/27			
2/28			
2/29			
2/30			
2/31			



(continued)

#	building block(s)	structure	analytical data
---	-------------------	-----------	-----------------




2/32



2/33



2/34

**Example 3****[0296]**









2-(2-Chloro-3'-(3-(2-cyanothiophen-3-yl)-5-fluoro-1-tosyl-1H-indol-2-yl)-[1,1'-biphenyl]-4-yl)-2-(dimethylamino)acetic acid (**3**)

[0297] To a solution of compound **2/3** (80 mg, 0.11 mmol) in THF (8 mL), MeOH (3 mL) and H₂O (3 mL) was added LiOH·H₂O (24 mg, 0.57 mmol). The mixture was stirred at rt for 30 min, concentrated, diluted with H₂O (6 mL), adjusted

to pH = 3 with 2N HCl and extracted with EA (2 x 50 mL). The combined organic layer was dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to afford compound **3** as a white solid. ¹H-NMR (500 MHz, DMSO-d₆) δ: 8.41 (dd, J = 9.3, 4.3 Hz, 1H), 7.82 (d, J = 5.0 Hz, 1H), 7.74 (d, J = 2.0 Hz, 1H), 7.57 (dd, J = 7.8, 1.7 Hz, 1H), 7.50-7.41 (m, 4H), 7.30-7.25 (m, 3H), 7.18-7.15 (m, 2H), 7.07-7.05 (m, 2H), 6.95 (d, J = 5.0 Hz, 1H), 4.53 (s, 1H), 2.84 (s, 6H), 2.27 (s, 3H); MS: 683.8 (M+1)⁺.

Example 3/1 to 3/3

[0298] The following Examples were saponified similar as described for Example 3 using the appropriate starting material (ester).

#	starting material	structure	analytical data
3/1	 1/5 first eluting isomer		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.42 (dd, J = 4.5, 9.0 Hz, 1H), 7.80 (d, J = 5.0 Hz, 1H), 7.40 (d, J = 8.0 Hz, 1H), 7.32-7.17 (m, 7H), 7.04 (dd, J = 2.8, 8.8 Hz, 1H), 6.99 (s, 1H), 6.88 (d, J = 5.5 Hz, 1H), 2.36 (s, 3H), 1.83-1.80 (m, 1H), 1.43 (s, 3H), 1.31-1.29 (m, 1H), 1.22-1.19 (m, 1H); MS: 568.8 (M-1) ⁻ .
3/2	 1/6 second eluting isomer		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.43 (dd, J = 4.3, 8.8 Hz, 1H), 7.74 (d, J = 5.0 Hz, 1H), 7.36-7.21 (m, 8H), 7.13-7.04 (m, 2H), 6.75 (d, J = 5.0 Hz, 1H), 2.35 (s, 3H), 1.94-1.92 (m, 1H), 1.63-1.62 (m, 1H), 1.38 (s, 3H), 1.20-1.17 (m, 1H); MS: 568.8 (M-1) ⁻ .
3/3	 1/7		¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.21 (dd, J = 4.5, 9.0 Hz, 1H), 6.68 (d, J = 9.0 Hz, 2H), 7.31-7.26 (m, 3H), 7.17 (t, J = 8.0 Hz, 1H), 6.97 (dd, J = 2.5, 9.0 Hz, 1H), 6.75 (d, J = 6.0 Hz, 1H), 6.64 (d, J = 8.0 Hz, 1H), 6.48 (dd, J = 2.0, 8.0 Hz, 1H), 6.39 (d, J = 6.0 Hz, 1H), 6.26 (s, 1H), 3.93 (t, J = 8.0 Hz, 2H), 3.79 (t, J = 6.5 Hz, 2H), 3.60 (s, 3H), 3.54-3.47 (m, 1H), 2.32 (s, 3H); MS: 577.1 (M+1) ⁺ .
3/4	 1/8		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.36 (dd, J = 4.3, 8.8 Hz, 1H), 7.60 (d, J = 5.5 Hz, 1H), 7.30 (d, J = 8.5 Hz, 2H), 7.25-7.17 (m, 4H), 6.84 (dd, J = 2.5, 8.5 Hz, 1H), 6.75-6.70 (m, 2H), 6.55 (dd, J = 1.8, 8.3 Hz, 1H), 6.19 (t, J = 55.0 Hz, 1H), 6.16 (s, 1H), 3.99-3.82 (m, 4H), 3.58-3.52 (m, 1H), 2.38 (s, 3H); MS: 597.1 (M+1) ⁺ .

(continued)



#

starting material

structure



analytical data

3/5





¹H-NMR (500 MHz, DMSO-d₆) δ:
12.64 (br s, 1H), 9.31 (d, J = 1.0 Hz, 1H), 8.26 (dd, J = 4.0, 9.0 Hz, 1H), 8.10 (dd, J = 1.3, 4.8 Hz, 1H), 7.40-7.32 (m, 5H), 7.14-7.06 (m, 3H), 6.60 (br s, 1H), 6.46 (dd, J = 1.8, 8.3 Hz, 1H), 6.20 (br s, 1H), 3.87 (t, J = 8.0 Hz, 2H), 3.74 (t, J = 6.5 Hz, 2H), 3.54-3.48 (m, 1H), 2.33 (s, 3H); MS: 575.0 (M+1)⁺.

3/6





¹H-NMR (500 MHz, DMSO-d₆) δ:
8.23 (dd, J = 4.5, 9.0 Hz, 1H), 7.46 (d, J = 5.0 Hz, 1H), 7.33-7.27 (m, 5H), 7.14 (t, J = 7.8 Hz, 1H), 6.87 (dd, J = 2.5, 8.5 Hz, 1H), 6.63-6.60 (m, 2H), 6.45 (dd, J = 2.0, 8.0 Hz, 1H), 6.18 (s, 1H), 4.14-4.09 (m, 1H), 3.89-3.73 (m, 5H), 3.54-3.48 (m, 1H), 2.99 (s, 3H), 2.31 (s, 3H); MS: 591.1 (M+1)⁺.

3/7





¹H-NMR (500 MHz, DMSO-d₆) δ:
12.69 (br s, 1H), 11.24 (s, 1H), 8.25 (dd, J = 4.3, 9.3 Hz, 1H), 7.49 (d, J = 5.0 Hz, 1H), 7.38-7.29 (m, 6H), 7.13 (d, J = 7.8 Hz, 1H), 6.83 (dd, J = 2.5, 8.5 Hz, 1H), 6.72 (d, J = 5.5 Hz, 1H), 6.60-6.57 (m, 1H), 6.45 (dd, J = 2.0, 8.0 Hz, 1H), 6.19 (s, 1H), 3.91-3.87 (m, 2H), 3.79-3.75 (m, 2H), 3.54-3.50 (m, 1H), 2.32 (m, 3H); MS: 590.0 (M+1)⁺.

3/8



¹H-NMR (500 MHz, DMSO-d₆) δ:
8.28 (dd, J = 4.8, 9.3 Hz, 1H), 8.24 (d, J = 5.0 Hz, 1H), 7.49 (d, J = 7.5 Hz, 1H), 7.42-7.33 (m, 3H), 7.23 (s, 1H), 7.15-7.11 (m, 2H), 6.94 (d, J = 5.0 Hz, 1H), 6.52 (d, J = 7.5 Hz, 1H), 6.47 (dd, J = 1.3, 8.3 Hz, 1H), 6.17 (s, 1H), 3.86 (t, J = 8.0 Hz, 2H), 3.75 (t, J = 6.5 Hz, 2H), 3.45-3.40 (m, 1H), 2.28 (s, 3H); MS: 569.8 (M-1)⁻.

3/9



¹H-NMR (500 MHz, DMSO-d₆) δ:
8.28 (dd, J = 4.5, 9.5 Hz, 1H), 7.98 (d, J = 4.5 Hz, 1H), 7.71-7.67 (m, 1H), 7.54-7.50 (m, 4H), 7.37-7.33 (m, 1H), 7.14-7.11 (m, 2H), 6.93 (d, J = 5.0 Hz, 1H), 6.53 (d, J = 7.5 Hz, 1H), 6.45 (dd, J = 1.8, 8.3 Hz, 1H), 6.18 (s, 1H), 3.83 (t, J = 7.8 Hz, 2H), 3.74 (t, J = 6.3 Hz, 2H), 3.34-3.30 (m, 1H); MS: 557.9 (M+1)⁺.



(continued)

#



starting material

structure



analytical data

3/
10



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
8.28 (dd, $J = 4.0, 9.0$ Hz, 1H), 7.98 (d, $J = 5.0$ Hz, 1H), 7.44 (d, $J = 8.0$ Hz, 2H), 7.37-7.33 (m, 3H), 7.16-7.11 (m, 2H), 6.94 (d, $J = 5.0$ Hz, 1H), 6.55 (d, $J = 7.5$ Hz, 1H), 6.48 (dd, $J = 1.5, 8.0$ Hz, 1H), 6.21 (s, 1H), 3.89 (t, $J = 7.5$ Hz, 2H), 3.77 (t, $J = 6.3$ Hz, 2H), 3.49-3.44 (m, 1H), 2.63 (q, $J = 7.5$ Hz, 2H), 1.12 (t, $J = 7.5$ Hz, 3H); MS: 586.8 ($\text{M}+1$) $^+$.

3/
11



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
8.28 (dd, $J = 4.3, 9.3$ Hz, 1H), 7.99 (d, $J = 5.0$ Hz, 1H), 7.58 (d, $J = 9.0$ Hz, 2H), 7.52-7.23 (m, 4H), 7.15-7.12 (m, 2H), 6.95 (d, $J = 5.0$ Hz, 1H), 6.53 (d, $J = 7.5$ Hz, 1H), 6.49-6.48 (m, 1H), 6.27 (s, 1H), 3.90 (t, $J = 8.0$ Hz, 2H), 3.79 (t, $J = 6.5$ Hz, 2H), 3.46-3.41 (m, 1H); MS: 623.7 ($\text{M}+1$) $^+$.

3/
12

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
8.28 (dd, $J = 4.5, 9.0$ Hz, 1H), 7.97 (d, $J = 5.0$ Hz, 1H), 7.37-7.30 (m, 3H), 7.27 (s, 1H), 7.15-7.10 (m, 2H), 6.94 (d, $J = 5.5$ Hz, 1H), 6.54 (d, $J = 7.5$ Hz, 1H), 6.46 (dd, $J = 1.8, 8.3$ Hz, 1H), 6.14 (s, 1H), 3.84 (t, $J = 8.0$ Hz, 2H), 3.74 (t, $J = 6.5$ Hz, 2H), 3.41-3.35 (m, 1H), 2.88-2.80 (m, 4H), 2.02-1.97 (m, 2H); MS: 598.2 ($\text{M}+1$) $^+$.













3/
13

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
8.38 (dd, $J = 4.0, 9.5$ Hz, 1H), 8.18 (s, 1H), 8.06-7.96 (m, 4H), 7.76-7.68 (m, 2H), 7.46 (d, $J = 9.0$ Hz, 1H), 7.39-7.36 (m, 1H), 7.15-7.12 (m, 2H), 6.95 (d, $J = 5.0$ Hz, 1H), 6.59 (d, $J = 7.0$ Hz, 1H), 6.45 (d, $J = 7.5$ Hz, 1H), 6.08 (s, 1H), 3.69-3.59 (m, 4H), 3.42-3.32 (m, 1H); MS: 608.2 ($\text{M}+1$) $^+$.

3/
14

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
8.16 (dd, $J = 4.5, 9.0$ Hz, 1H), 7.98 (d, $J = 5.5$ Hz, 1H), 7.36-7.17 (m, 4H), 7.05 (t, $J = 7.5$ Hz, 2H), 6.99 (d, $J = 5.0$ Hz, 1H), 6.45-6.40 (m, 2H), 5.97 (s, 1H), 3.79 (t, $J = 8.0$ Hz, 2H), 3.68 (t, $J = 6.3$ Hz, 2H), 3.45-3.40 (m, 1H), 2.36 (s, 3H); MS: 589.9 ($\text{M}+1$) $^+$.

(continued)

#	starting material	structure	analytical data
3/ 15	 1/19		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.42 (d, $J = 8.5$ Hz, 1H), 8.03-8.01 (m, 1H), 7.83 (d, $J = 5.0$ Hz, 1H), 7.79 (s, 1H), 7.67-7.62 (m, 2H), 7.56-7.53 (m, 1H), 7.42-7.39 (m, 2H), 7.34 (d, $J = 8.0$ Hz, 2H), 7.24 (d, $J = 8.5$ Hz, 2H), 6.97 (d, $J = 5.0$ Hz, 1H), 4.26 (s, 2H), 2.35 (s, 3H); MS: 576.7 ($\text{M}+1$) $^+$.
3/ 16	 1/20		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.07 (br s, 1H), 8.28 (d, $J = 8.0$ Hz, 1H), 8.02 (d, $J = 5.0$ Hz, 1H), 7.53-7.50 (m, 1H), 7.44 (d, $J = 8.0$ Hz, 2H), 7.38 (d, $J = 4.0$ Hz, 2H), 7.31 (d, $J = 8.5$ Hz, 2H), 7.05 (d, $J = 3.5$ Hz, 1H), 6.99 (d, $J = 5.5$ Hz, 1H), 6.83 (d, $J = 4.0$ Hz, 1H), 2.97 (s, 2H), 2.31 (s, 3H), 1.09 (s, 6H); MS: 559.0 ($\text{M}-1$) $^-$.
3/ 17	 1/21		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.26 (d, $J = 11.0$ Hz, 1H), 8.05 (d, $J = 6.0$ Hz, 1H), 7.54-7.32 (m, 8H), 7.04 (d, $J = 6.0$ Hz, 1H), 6.91 (d, $J = 2.0$ Hz, 1H), 2.39-2.34 (m, 1H), 2.32 (s, 3H), 1.75-1.70 (m, 1H), 1.39-1.35 (m, 1H), 1.20-1.15 (m, 1H); MS: 562.1 ($\text{M}+18$) $^+$.
3/ 18	 1/22		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.41 (d, $J = 10.0$ Hz, 1H), 7.80 (d, $J = 6.5$ Hz, 1H), 7.65-7.49 (m, 3H), 7.43-7.21 (m, 9H), 6.96 (d, $J = 6.5$ Hz, 1H), 6.35 (d, $J = 20.0$ Hz, 1H), 2.36 (s, 3H); MS: 524.6 ($\text{M}+1$) $^+$.
3/ 19	 1/24		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.47 (br s, 1H), 8.25 (d, $J = 10.5$ Hz, 1H), 8.07 (d, $J = 6.5$ Hz, 1H), 7.70 (d, $J = 20.0$ Hz, 1H), 7.56-7.51 (m, 4H), 7.42-7.32 (m, 4H), 7.23 (d, $J = 4.5$ Hz, 1H), 7.09 (d, $J = 6.5$ Hz, 1H), 6.18 (d, $J = 20.0$ Hz, 1H), 2.31 (s, 3H); MS: 530.7 ($\text{M}+1$) $^+$.
3/ 20	 1/25		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.39 (br s, 1H), 8.29 (d, $J = 8.5$ Hz, 1H), 7.98 (d, $J = 5.0$ Hz, 1H), 7.51 (t, $J = 7.5$ Hz, 1H), 7.40-7.28 (m, 9H), 7.06 (s, 1H), 6.92 (d, $J = 4.0$ Hz, 1H), 2.30 (s, 3H), 1.36 (s, 6H); MS: 563.1 ($\text{M}+\text{Na}$) $^+$.



(continued)

#



starting material

structure



analytical data

3/
21



$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.26 (dd, $J = 4.5, 9.0$ Hz, 1H), 8.10 (d, $J = 1.0$ Hz, 1H), 7.99 (dd, $J = 1.8, 7.8$ Hz, 1H), 7.86 (d, $J = 5.5$ Hz, 1H), 7.56-7.51 (m, 3H), 7.44 (s, 1H), 7.40 (d, $J = 8.0$ Hz, 1H), 7.25-7.21 (m, 1H), 7.15 (dd, $J = 2.8, 8.8$ Hz, 1H), 6.98 (d, $J = 4.5$ Hz, 1H), 3.48 (d, $J = 12.5$ Hz, 2H), 2.53-2.48 (m, 2H), 1.47 (d, $J = 12.5$ Hz, 2H), 1.33-1.28 (m, 1H), 0.86-0.77 (m, 5H); MS: 633.9 ($\text{M}+1$) $^+$.

3/
22



$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.44 (dd, $J = 4.5, 9.0$ Hz, 1H), 8.09 (d, $J = 2.0$ Hz, 1H), 8.00 (dd, $J = 1.5, 7.5$ Hz, 1H), 7.82 (d, $J = 5.0$ Hz, 1H), 7.52-7.44 (m, 3H), 7.39 (d, $J = 8.0$ Hz, 1H), 7.29-6.95 (m, 4H); MS: 632.0 ($\text{M}-1$) $^-$.

3/
23

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.41 (br s, 1H), 8.31 (dd, $J = 4.5, 9.0$ Hz, 1H), 8.04-7.95 (m, 3H), 7.56-7.48 (m, 3H), 7.42-7.34 (m, 4H), 7.26-7.18 (m, 3H), 7.03 (d, $J = 5.0$ Hz, 1H), 2.08 (s, 3H); MS: 642.9 ($\text{M}-1$) $^-$.

3/
24

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.29-8.26 (m, 1H), 8.01 (d, $J = 5.5$ Hz, 1H), 7.65-7.55 (m, 2H), 7.45-7.35 (m, 3H), 7.26 (d, $J = 5.0$ Hz, 1H), 7.18-7.14 (m, 1H), 7.06-7.03 (m, 1H), 6.97 (d, $J = 5.0$ Hz, 1H), 6.91 (s, 1H), 2.35-2.30 (m, 1H), 1.70 (s, 1H), 1.41-1.37 (m, 1H), 1.17 (s, 1H); MS: 575.0 ($\text{M}-1$) $^-$.

3/
25

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.66 (br s, 1H), 8.31-8.26 (m, 1H), 7.76 (d, $J = 5.5$ Hz, 1H), 7.46-7.33 (m, 6H), 7.20-7.15 (m, 2H), 6.60 (d, $J = 7.5$ Hz, 1H), 6.53-6.49 (m, 1H), 6.23 (s, 1H), 3.92 (t, $J = 8.0$ Hz, 2H), 3.79 (t, $J = 6.5$ Hz, 2H), 3.54-3.50 (m, 1H), 2.34 (s, 3H); MS: 572.1 ($\text{M}+1$) $^+$.



(continued)

#



starting material

structure



analytical data

3/
26



¹H-NMR (500 MHz, CD₃OD) δ: 8.42-8.39 (m, 1H), 7.72 (d, J = 1.5 Hz, 1H), 7.36 (d, J = 8.5 Hz, 2H), 7.28-7.14 (m, 5H), 6.65 (d, J = 8.0 Hz, 1H), 6.66-6.64 (m, 1H), 6.60-6.57 (m, 1H), 6.35 (d, J = 2.0 Hz, 1H), 6.20 (s, 1H), 4.01 (t, J = 7.5 Hz, 2H), 3.91 (t, J = 6.0 Hz, 2H), 3.54-3.50 (m, 1H), 2.38 (s, 3H); MS: 556.2 (M+1)⁺.

3/
27



¹H-NMR (500 MHz, DMSO-d₆) δ: 8.23-8.19 (m, 1H), 7.32-7.26 (m, 6H), 7.14 (t, J = 7.5 Hz, 1H), 6.84-6.81 (m, 1H), 6.66-6.62 (m, 1H), 6.55 (d, J = 5.0 Hz, 1H), 6.45-6.42 (m, 1H), 6.16 (s, 3H), 3.90-3.85 (m, 2H), 3.77-3.73 (m, 2H), 3.52-3.49 (m, 1H), 3.25-3.19 (m, 2H), 2.48-2.33 (m, 2H), 2.31 (s, 1H); MS: 591.0 (M+1)⁺.

3/
28

¹H-NMR (500 MHz, CD₃OD) δ: 8.40-8.37 (m, 1H), 7.54 (d, J = 5.5 Hz, 1H), 7.34 (d, J = 8.0 Hz, 2H), 7.25-7.19 (m, 4H), 7.05-7.03 (m, 1H), 6.75 (d, J = 5.0 Hz, 1H), 6.67-6.59 (m, 2H), 6.26 (s, 1H), 4.05-4.01 (m, 2H), 3.95-3.92 (m, 2H), 3.60-3.57 (m, 1H), 2.65 (s, 3H), 2.38 (s, 3H), 2.33 (s, 3H); MS: 618.0 (M+1)⁺.

C3/
29

¹H-NMR (500 MHz, CD₃OD) δ: 8.41-8.38 (m, 1H), 7.63 (d, J = 1.5 Hz, 1H), 7.40-7.35 (m, 3H), 7.30-7.21 (m, 5H), 6.68-6.61 (m, 2H), 6.18 (d, J = 1.5 Hz, 1H), 4.03-3.99 (m, 2H), 3.93-3.89 (m, 2H), 3.56-3.53 (m, 1H), 2.37 (s, 3H); MS: 572.1 (M+1)⁺.

3/
30

¹H-NMR (500 MHz, CD₃OD) δ: 8.35-8.32 (m, 1H), 7.62 (d, J = 5.0 Hz, 1H), 7.31-7.15 (m, 7H), 6.78-6.75 (m, 1H), 6.66 (br s, 2H), 6.51-6.48 (m, 1H), 3.98-3.79 (m, 4H), 3.55-3.50 (m, 1H), 2.37 (s, 3H); MS: 615.1 (M+1)⁺.



(continued)

#



starting material

structure



analytical data

3/
31



¹H-NMR (500 MHz, DMSO-d₆) δ:
8.27-8.23 (m, 1H), 7.62 (d, J = 7.0 Hz, 1H), 7.41 (t, J = 8.0 Hz, 1H), 7.35-7.28 (m, 5H), 7.20 (t, J = 7.0 Hz, 1H), 7.09 (t, J = 7.0 Hz, 1H), 6.87 (d, J = 7.5 Hz, 1H), 6.63-6.56 (m, 2H), 6.38 (d, J = 6.5 Hz, 1H), 6.05-6.01 (m, 1H), 4.41-4.13 (m, 4H), 3.82-3.79 (m, 2H), 3.70-3.66 (m, 2H), 3.59-3.55 (m, 1H), 3.43-3.39 (m, 1H), 2.33 (s, 3H); MS: 597.2 (M+1)⁺.

3/
32



¹H-NMR (500 MHz, DMSO-d₆) δ:
8.29-8.26 (m, 1H), 7.98 (d, J = 5.1 Hz, 1H), 7.41 (d, J = 8.4 Hz, 2H), 7.37-7.31 (m, 3H), 7.17-7.09 (m, 2H), 6.94 (d, J = 5.1 Hz, 1H), 6.56 (d, J = 7.6 Hz, 1H), 6.48 (dd, J = 8.1, 1.8 Hz, 1H), 6.20 (s, 1H), 3.88 (t, J = 7.8 Hz, 2H), 3.77 (t, J = 6.5 Hz, 2H), 3.47-3.43 (m, 1H), 2.33 (s, 3H); MS: 569.7 (M-1)⁻.

3/
33

¹H-NMR (500 MHz, DMSO-d₆) δ:
8.30-8.26 (m, 1H), 7.98 (d, J = 5.1 Hz, 1H), 7.46 (d, J = 8.9 Hz, 2H), 7.35 (td, J = 9.2, 2.5 Hz, 1H), 7.16-7.10 (m, 2H), 7.02 (d, J = 9.0 Hz, 2H), 6.94 (d, J = 5.1 Hz, 1H), 6.56 (d, J = 7.5 Hz, 1H), 6.48 (d, J = 8.1 Hz, 1H), 6.23 (s, 1H), 3.90 (t, J = 7.9 Hz, 2H), 3.77-3.80 (m, 5H), 3.48-3.44 (m, 1H); MS: 578.8 (M+1)⁺.

3/
34

¹H-NMR (500 MHz, DMSO-d₆) δ:
8.31-8.28 (m, 1H), 7.99 (d, J = 5.1 Hz, 1H), 7.92 (d, J = 8.4 Hz, 2H), 7.70 (d, J = 8.3 Hz, 2H), 7.39 (td, J = 9.1, 2.5 Hz, 1H), 7.15-7.11 (m, 2H), 6.94 (d, J = 5.1 Hz, 1H), 6.54-6.47 (m, 2H), 6.24 (s, 1H), 3.89 (t, J = 7.9 Hz, 2H), 3.78 (t, J = 6.4 Hz, 2H), 3.44-3.47 (m, 1H); MS: 625.8 (M+1)⁺.

3/
35

¹H-NMR (500 MHz, DMSO-d₆) δ:
8.30-8.27 (m, 1H), 7.99 (d, J = 5.1 Hz, 1H), 7.73 (d, J = 8.3 Hz, 2H), 7.65 (d, J = 8.3 Hz, 2H), 7.40-7.35 (m, 1H), 7.15-7.12 (m, 2H), 7.09 (t, J = 55.5 Hz, 1H), 6.96 (d, J = 5.1 Hz, 1H), 6.55 (d, J = 7.5 Hz, 1H), 6.48 (d, J = 8.0 Hz, 1H), 6.24 (s, 1H), 3.88 (t, J = 7.9 Hz, 2H), 3.77 (t, J = 6.5 Hz, 2H), 3.42-3.48 (m, 1H); MS: 607.8 (M+1)⁺.



(continued)

#



starting material

structure



analytical data

3/
36



¹H-NMR (500 MHz, CD₃OD) δ : 8.24-8.21 (m, 1H), 7.79 (d, J = 5.1 Hz, 1H), 7.23-7.15 (m, 2H), 7.11 (dd, J = 8.7, 2.5 Hz, 1H), 6.86 (d, J = 5.1 Hz, 1H), 6.77 (d, J = 7.6 Hz, 1H), 6.56-6.53 (m, 2H), 4.05-4.01 (m, 2H), 3.94-3.90 (m, 2H), 3.50-3.56 (m, 1H), 3.47 (d, J = 12.9 Hz, 2H), 2.48 (t, J = 11.7 Hz, 2H), 1.49 (dd, J = 13.0, 2.1 Hz, 2H), 1.39-1.26 (m, 1H), 0.90-0.76 (m, 5H); MS: 579.0 (M+1)⁺.

3/
37



¹H-NMR (500 MHz, DMSO-d₆) δ : 12.77 (s, 1H), 8.27 (dd, J = 9.2, 4.4 Hz, 1H), 7.99 (d, J = 5.1 Hz, 1H), 7.45-7.28 (m, 5H), 7.19-7.08 (m, 2H), 6.95 (d, J = 5.1 Hz, 1H), 6.59-6.57 (m, 1H), 6.51-6.47 (m, 1H), 6.21 (s, 1H), 3.92 (d, J = 7.2 Hz, 2H), 3.54 (d, J = 7.2 Hz, 2H), 2.33 (s, 3H), 1.52 (s, 3H); MS: 586.2 (M+1)⁺.

3/
38

¹H-NMR (400 MHz, CD₃OD) δ : 8.40 (dd, J = 9.2, 4.4 Hz, 1H), 7.88 (d, J = 5.1 Hz, 1H), 7.41-7.57 (m, 3H), 7.35-7.20 (m, 5H), 7.10-6.98 (m, 3H), 3.73-3.69 (m, 2H), 3.47-3.41 (m, 2H), 2.80-2.67 (m, 1H), 2.36 (s, 3H), 2.26 (d, J = 14.1 Hz, 2H), 2.13-1.98 (m, 2H); MS: 600.2 (M+1)⁺.

3/
39

¹H-NMR (500 MHz, DMSO-d₆) δ : 12.14 (s, 1H), 8.30 (dd, J = 9.2, 4.4 Hz, 1H), 8.00-7.97 (m, 1H), 7.43-7.20 (m, 7H), 7.17-7.08 (m, 2H), 7.06-6.91 (m, 2H), 3.62-3.29 (m, 1H), 3.10-2.92 (m, 1H), 2.50-2.39 (m, 2H), 2.32 (s, 3H), 2.24-2.02 (m, 2H); MS: 568.7 (M-1)⁻.

3/
40

¹H-NMR (500 MHz, DMSO-d₆) δ : 12.68 (s, 1H), 8.28 (dd, J = 9.2, 4.4 Hz, 1H), 7.98 (d, J = 5.1 Hz, 1H), 7.59 (d, J = 8.4 Hz, 2H), 7.44-7.32 (m, 3H), 7.28-7.17 (m, 2H), 6.94 (d, J = 5.1 Hz, 1H), 6.54 (dd, J = 8.7, 2.8 Hz, 1H), 6.40 (d, J = 2.3 Hz, 1H), 4.02-3.93 (m, 2H), 3.90-3.77 (m, 2H), 3.59-3.48 (m, 1H), 2.36 (s, 3H); MS: 606.1 (M+1)⁺.



(continued)

#



starting material

structure



analytical data

3/
41



$^1\text{H-NMR}$ (500 MHz, CDCl_3) δ : 8.30 (dd, $J = 9.2, 4.4$ Hz, 1H), 7.49 (d, $J = 5.1$ Hz, 1H), 7.42 (d, $J = 8.3$ Hz, 2H), 7.20-7.12 (m, 3H), 7.05 (dd, $J = 8.4, 2.4$ Hz, 1H), 6.90 (t, $J = 8.9$ Hz, 1H), 6.83 (d, $J = 5.1$ Hz, 1H), 6.56-6.49 (m, 1H), 6.39-6.35 (m, 1H), 4.12-3.98 (m, 4H), 3.63-3.55 (m, 1H), 2.35 (s, 3H); MS: 590.1 ($\text{M}+1$) $^+$.

3/
42



$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.34-8.29 (m, 1H), 7.78 (d, $J = 5.0$ Hz, 1H), 7.34-7.15 (m, 6H), 6.87 (d, $J = 5.0$ Hz, 1H), 6.62 (d, $J = 7.5$ Hz, 1H), 6.57 (dd, $J = 1.5$ Hz, 1.0 Hz, 1H), 6.18 (s, 1H), 4.00 (t, $J = 8.0$ Hz, 2H), 3.89 (t, $J = 7.5$ Hz, 2H), 3.58-3.54 (m, 1H), 2.38 (s, 3H); MS: 589.7 ($\text{M}+1$) $^+$.

3/
43

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.69 (br s, 1H), 8.40 (d, $J = 8.0$ Hz, 1H), 7.99 (d, $J = 6.5$ Hz, 1H), 7.45-7.32 (m, 5H), 7.14 (t, $J = 9.5$ Hz, 1H), 6.95 (d, $J = 6.0$ Hz, 1H), 6.55-6.48 (m, 2H), 6.15 (s, 1H), 3.89 (t, $J = 9.5$ Hz, 2H), 3.74 (t, $J = 8.5$ Hz, 2H), 3.54-3.49 (m, 1H), 2.34 (s, 3H); MS: 606.1 ($\text{M}+1$) $^+$.

3/
44

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.28 (d, $J = 9.0$ Hz, 1H), 7.99 (d, $J = 5.0$ Hz, 1H), 7.52 (dd, $J = 2.5, 2.0$ Hz, 1H), 7.42 (d, $J = 8.5$ Hz, 2H), 7.35-7.31 (m, 3H), 7.13 (t, $J = 7.5$ Hz, 1H), 6.97 (d, $J = 5.0$ Hz, 1H), 6.55 (d, $J = 7.5$ Hz, 1H), 6.50-6.46 (m, 1H), 6.19 (s, 1H), 3.89-3.86 (m, 2H), 3.78-3.74 (m, 2H), 3.45-3.40 (m, 1H), 2.33 (s, 3H); MS: 587.8 ($\text{M}+1$) $^+$.

3/
45

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.46 (d, $J = 8.5$ Hz, 1H), 8.00 (d, $J = 5.0$ Hz, 1H), 7.95-7.88 (m, 2H), 7.46 (d, $J = 8.5$ Hz, 2H), 7.35 (d, $J = 8.5$ Hz, 2H), 7.16 (t, $J = 8.0$ Hz, 1H), 7.01 (d, $J = 5.0$ Hz, 1H), 6.56 (d, $J = 7.5$ Hz, 1H), 6.50 (dd, $J = 1.5, 2.0$ Hz, 1H), 6.20 (s, 1H), 3.93-3.87 (m, 2H), 3.80-3.75 (m, 2H), 3.54-3.50 (m, 1H), 2.35 (s, 3H); MS: 579.1 ($\text{M}+1$) $^+$.



(continued)

#



starting material

structure



analytical data

3/
46



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
8.12 (d, $J = 9.0$ Hz, 1H), 7.98 (d, $J = 5.0$ Hz, 1H), 7.42-7.28 (m, 5H), 7.13 (t, $J = 7.5$ Hz, 1H), 7.07 (s, 1H), 6.96 (d, $J = 5.0$ Hz, 1H), 6.57 (d, $J = 7.5$ Hz, 1H), 6.46 (dd, $J = 1.5, 2.0$ Hz, 1H), 6.22 (s, 1H), 3.90-3.86 (m, 2H), 3.78-3.74 (m, 2H), 3.45-3.41 (m, 1H), 2.36 (s, 3H), 2.31 (s, 3H); MS: 567.8 ($\text{M}+1$) $^+$.

3/
47



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
8.81 (s, 1H), 8.33-8.29 (m, 1H), 8.10-8.03 (m, 3H), 7.85 (d, $J = 10.0$ Hz, 1H), 7.58-7.52 (m, 2H), 7.42-7.32 (m, 4H), 7.27 (d, $J = 10.0$ Hz, 2H), 7.21 (dd, $J = 3.0, 3.5$ Hz, 1H), 7.08 (d, $J = 6.0$ Hz, 1H), 2.25 (s, 3H); MS: 594.1 ($\text{M}+1$) $^+$.

3/
48

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ :
8.43-8.40 (m, 1H), 8.08 (d, $J = 2.0$ Hz, 1H), 7.99 (dd, $J = 10.0, 2.0$ Hz, 1H), 7.83 (d, $J = 6.5$ Hz, 1H), 7.52-7.44 (m, 3H), 7.38 (d, $J = 10.0$ Hz, 1H), 7.30-7.25 (m, 3H), 7.15 (d, $J = 10.0$ Hz, 2H), 7.08-6.96 (m, 3H), 2.24 (s, 3H); MS: 624.7 ($\text{M}-1$) $^-$.

3/
49

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ :
8.44-8.40 (m, 1H), 7.88-7.83 (m, 2H), 7.72 (dd, $J = 8.5, 1.5$ Hz, 1H), 7.53 (d, $J = 2.0$ Hz, 1H), 7.51-7.45 (m, 2H), 7.34-7.27 (m, 5H), 7.18 (d, $J = 8.5$ Hz, 2H), 7.07 (dd, $J = 8.5, 2.5$ Hz, 1H), 7.00 (d, $J = 5.0$ Hz, 1H), 2.30 (s, 3H); MS: 624.7 ($\text{M}-1$) $^-$.

3/
50

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
12.41 (br s, 1H), 8.29-8.25 (m, 1H), 7.98 (d, $J = 5.0$ Hz, 1H), 7.39-7.29 (m, 5H), 7.20-7.12 (m, 2H), 7.02-6.95 (m, 2H), 6.72-6.65 (m, 2H), 3.33-3.30 (m, 2H), 2.83-2.79 (m, 2H), 2.32 (s, 3H), 2.01-1.99 (m, 2H), 1.48-1.43 (m, 2H), 1.16 (s, 3H); MS: 614.0 ($\text{M}+1$) $^+$.



(continued)

#



starting material

structure



analytical data

3/
51



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.03 (s, 1H), 8.26 (d, $J = 8.5$ Hz, 1H), 8.00 (d, $J = 5.5$ Hz, 1H), 7.61-7.58 (m, 2H), 7.53-7.50 (m, 3H), 7.41-7.34 (m, 2H), 7.28 (t, $J = 8.0$ Hz, 1H), 7.00-6.96 (m, 2H), 6.87 (d, $J = 7.5$ Hz, 1H), 6.82 (s, 1H), 4.63 (s, 2H); MS: 549.0 ($\text{M}+1$) $^+$.

3/
52



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.24 (s, 1H), 8.28 (dd, $J = 9.2, 4.4$ Hz, 1H), 8.19 (s, 1H), 7.98 (d, $J = 5.1$ Hz, 1H), 7.42 (d, $J = 8.4$ Hz, 2H), 7.37-7.29 (m, 3H), 7.17-7.10 (m, 2H), 6.93 (d, $J = 5.1$ Hz, 1H), 6.57-6.55 (m, 1H), 6.48-6.45 (m, 1H), 6.20 (s, 1H), 3.87-3.83 (m, 2H), 3.73-3.70 (m, 2H), 3.52-3.39 (m, 1H), 2.33 (s, 3H), 1.36 (s, 6H); MS: 657.0 ($\text{M}+1$) $^+$.

3/
53

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.00 (s, 1H), 8.08 (d, $J = 5.1$ Hz, 1H), 8.00 (d, $J = 8.3$ Hz, 2H), 7.73 (d, $J = 7.8$ Hz, 1H), 7.61-7.56 (m, 4H), 7.50-7.45 (m, 2H), 7.33-7.28 (m, 3H), 7.08 (d, $J = 5.0$ Hz, 1H), 7.00 (dd, $J = 11.4, 2.1$ Hz, 1H), 6.69 (dd, $J = 8.0, 2.2$ Hz, 1H), 3.80 (s, 1H), 2.33 (s, 3H); MS: 623.0 ($\text{M}+1$) $^+$.

3/
54

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.02 (s, 1H), 10.68 (s, 1H), 8.08 (d, $J = 5.1$ Hz, 1H), 8.00 (d, $J = 8.4$ Hz, 2H), 7.73 (d, $J = 8.0$ Hz, 1H), 7.60-7.38 (m, 6H), 7.30-7.21 (m, 3H), 7.02 (d, $J = 5.1$ Hz, 1H), 6.67 (dd, $J = 10.9, 2.3$ Hz, 1H), 6.51 (dd, $J = 8.1, 2.4$ Hz, 1H), 2.29 (s, 3H); MS: 609.2 ($\text{M}+1$) $^+$.

3/
55

$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.74 (br s, 1H), 8.30 (dd, $J = 9.0, 5.0$ Hz, 1H), 8.01 (d, $J = 5.0$ Hz, 1H), 7.72 (d, $J = 8.0$ Hz, 1H), 7.59 (d, $J = 8.5$ Hz, 2H), 7.49-7.34 (m, 7H), 7.27-7.16 (m, 4H), 7.05 (d, $J = 5.0$ Hz, 1H), 5.83 (br s, 1H), 2.24 (s, 3H), 1.65 (s, 3H); MS: 634.8 ($\text{M}-1$) $^-$.



(continued)

#



starting material

structure



analytical data

3/
56



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
13.39 (s, 1H), 8.27-8.24 (m, 1H), 7.97 (s, 1H), 7.93 (d, $J = 7.0$ Hz, 1H), 7.64-7.25 (m, 12H), 7.11 (s, 1H), 7.01 (br s, 1H), 6.75-6.73 (m, 1H), 4.42-4.37 (m, 2H), 4.17-4.13 (m, 2H), 3.67-3.63 (m, 1H); MS: 670.0 ($\text{M}-1$) $^-$.

3/
57



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ :
8.30-8.27 (m, 1H), 8.04 (d, $J = 5.0$ Hz, 1H), 7.96 (s, 1H), 7.93 (d, $J = 8.0$ Hz, 1H), 7.55-7.51 (m, 4H), 7.45-7.38 (m, 5H), 7.21-7.19 (m, 1H), 7.09 (s, 1H), 7.04 (d, $J = 5.0$ Hz, 1H); MS: 644.9 ($\text{M}-1$) $^-$.

3/
58

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.40 (dd, $J = 9.5, 4.0$ Hz, 1H), 7.71 (d, $J = 5.0$ Hz, 1H), 7.33 (d, $J = 8.5$ Hz, 2H), 7.27-7.16 (m, 4H), 7.04-7.02 (m, 1H), 6.91-6.89 (m, 1H), 6.80 (d, $J = 5.5$ Hz, 1H), 6.61 (d, $J = 8.0$ Hz, 1H), 6.46 (s, 1H), 4.16-4.13 (m, 2H), 2.94-2.90 (m, 1H), 2.35 (s, 3H), 2.11-2.07 (m, 2H), 1.89-1.81 (m, 4H), 1.57-1.54 (m, 2H); MS: 626.2 ($\text{M}+1$) $^+$.

3/
59

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ :
8.41-8.38 (m, 1H), 8.07 (s, 1H), 7.97 (d, $J = 7.5$ Hz, 1H), 7.69 (d, $J = 8.0$ Hz, 1H), 7.60-7.32 (m, 8H), 7.31-7.24 (m, 3H), 7.15-6.97 (m, 2H), 6.77 (t, $J = 55.5$ Hz, 1H), 6.69-6.66 (m, 1H), 4.53-4.43 (m, 2H), 4.35-4.29 (m, 2H), 3.72-3.64 (m, 1H); MS: 686.0 ($\text{M}-1$) $^-$.

3/
60

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.41 (dd, $J = 9.2, 4.4$ Hz, 1H), 7.79 (d, $J = 5.1$ Hz, 1H), 7.60 (d, $J = 8.3$ Hz, 2H), 7.53 (d, $J = 8.4$ Hz, 2H), 7.32-7.15 (m, 2H), 7.08-7.01 (m, 2H), 6.88 (d, $J = 5.0$ Hz, 1H), 6.81 (t, $J = 55.0$ Hz, 1H), 6.73-6.67 (m, 2H), 3.53-3.50 (m, 2H), 2.76-2.70 (m, 2H), 2.49-2.32 (m, 1H), 1.98-1.96 (m, 2H), 1.81-1.73 (m, 2H); MS: 634.0 ($\text{M}-1$) $^-$.



(continued)

#



starting material

structure



analytical data

3/
61



$^1\text{H-NMR}$ (400 MHz, CD_3COCD_3) δ : 8.38-8.34 (m, 1H), 8.07 (s, 1H), 8.00 (d, $J = 7.9$ Hz, 1H), 7.71 (s, 4H), 7.63-7.23 (m, 9H), 7.14-6.72 (m, 3H), 3.61-3.54 (m, 2H), 3.52-2.98 (m, 3H); MS: 705.0 ($\text{M}+1$) $^+$.

3/
62



$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.43 (dd, $J = 9.2, 4.3$ Hz, 1H), 8.09 (s, 1H), 8.00 (br s, 1H), 7.76-7.14 (m, 14H), 6.93-6.63 (m, 2H), 4.61-3.78 (m, 4H), 2.81 (s, 3H); MS: 719.0 ($\text{M}+1$) $^+$.

3/
63

$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.40-8.36 (m, 1H), 8.10 (s, 1H), 8.00 (d, $J = 5.5$ Hz, 1H), 7.58-7.23 (m, 11H), 6.85-6.80 (m, 2H), 6.75 (t, $J = 55.5$ Hz, 1H), 4.76-4.73 (m, 1H), 4.58-4.54 (m, 1H), 4.25-4.12 (m, 1H), 4.10-4.09 (m, 1H), 3.89-3.86 (m, 1H); MS: 691.9 ($\text{M}-1$) $^-$.

3/
64

$^1\text{H-NMR}$ (500 MHz, $\text{DMSO}-d_6$) δ : 11.11 (br s, 1H), 8.30 (dd, $J = 4.5, 9.5$ Hz, 1H), 7.99 (d, $J = 5.5$ Hz, 1H), 7.39-6.94 (m, 11H), 2.13 (s, 3H), 1.79-1.76 (m, 6H), 1.65-1.62 (m, 6H); MS: 622.8 ($\text{M}-1$) $^-$.

3/
65

$^1\text{H-NMR}$ (500 MHz, $\text{DMSO}-d_6$) δ : 8.29 (dd, $J = 4.3, 9.3$ Hz, 1H), 7.99 (d, $J = 5.0$ Hz, 1H), 7.39-6.91 (m, 11H), 3.33-3.26 (m, 1H), 2.95-2.90 (m, 1H), 2.41-1.81 (m, 11H); MS: 608.8 ($\text{M}-1$) $^-$.



(continued)

#



starting material

structure



analytical data

3/
66



$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ :
8.28-8.14 (m, 3H), 7.72-7.37 (m,
11H), 6.79-6.77 (m, 1H), 2.96-2.93
(m, 4H), 1.16-1.10 (m, 4H), 0.75 (s,
6H); MS: 593.7 ($\text{M}-1$) $^-$.

3/
67



$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.43
(dd, $J = 9.3, 4.8$ Hz, 1H), 8.04 (d, $J =$
7.5 Hz, 2H), 7.70 (t, $J = 8.0$ Hz, 1H),
7.61 (d, $J = 9.0$ Hz, 2H), 7.54 (d, $J =$
8.0 Hz, 2H), 7.32 (dt, $J = 2.5, 9.3$ Hz,
1H), 7.12 (t, $J = 8.0$ Hz, 1H), 6.90 (dd,
 $J = 2.5, 8.5$ Hz, 1H), 6.80 (t, $J = 55.8$
Hz, 1H), 6.32 (d, $J = 7.5$ Hz, 1H), 6.53
(dd, $J = 1.5, 8.0$ Hz, 1H), 6.21 (t, $J =$
1.8 Hz, 1H), 3.96 (br s, 2H), 3.85 (br
s, 2H), 3.57-3.51 (rn, 1H); MS: 727.2
($\text{M}+1$) $^+$.

3/
68

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.44
(dd, $J = 9.4, 4.4$ Hz, 1H), 8.06-8.03
(m, 2H), 7.71 (t, $J = 8.3$ Hz, 1H), 7.61
(d, $J = 8.0$ Hz, 2H), 7.49 (d, $J = 8.0$
Hz, 2H), 7.35-7.21 (m, 3H), 7.05-6.84
(m, 4H), 2.42-2.34 (m, 1H), 1.70-1.19
(m, 3H); MS: 610.0 ($\text{M}-1$) $^-$.

3/
69

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.38
(dd, $J = 9.3, 4.3$ Hz, 1H), 8.07 (d, $J =$
1.5 Hz, 1H), 7.97 (dd, $J = 8.0, 1.5$ Hz,
1H), 7.55-7.47 (m, 8H), 7.39 (t, $J =$
8.0 Hz, 1H), 7.33 (d, $J = 8.0$ Hz, 1H),
7.27 (dd, $J = 2.5, 9.2$ Hz, 1H), 7.26
(br s, 1H), 7.10-6.59 (m, 4H); MS:
670.9 ($\text{M}-1$) $^-$.

3/
70

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.57
(s, 1H), 8.09-8.07 (m, 3H), 8.00 (dd,
 $J = 8.3, 1.8$ Hz, 1H), 7.74 (t, $J = 7.8$
Hz, 1H), 7.61-7.43 (m, 7H), 7.36 (d,
 $J = 8.0$ Hz, 1H), 7.02 (s, 1H), 6.70 (t,
 $J = 55.5$ Hz, 1H), 6.14 (d, $J = 1.0$ Hz,
1H); MS: 681.1 ($\text{M}+1$) $^-$.



(continued)

#



starting material

structure



analytical data

3/
71


$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.44 (dd, $J = 4.3, 9.3$ Hz, 1H), 8.07-7.97 (m, 2H), 7.70 (t, $J = 8.0$ Hz, 1H), 7.53-7.44 (m, 7H), 7.34 (dt, $J = 2.5, 9.0$ Hz, 1H), 6.94-6.91 (m, 2H), 6.70-6.65 (m, 2H), 6.68 (t, $J = 55.5$ Hz, 1H), 4.71 (s, 2H); MS: 712.0 (M-H) $^-$.

3/
72

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.45 (dd, $J = 9.0, 4.5$ Hz, 1H), 8.17-7.92 (m, 2H), 7.71 (t, $J = 8.0$ Hz, 1H), 7.57-7.48 (m, 7H), 7.36-7.29 (m, 3H), 6.95-6.92 (m, 2H), 6.65 (t, $J = 55.8$ Hz, 1H), 1.75 (s, 3H); MS: 726.0 (M-H) $^-$.

3/
73

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.45 (dd, $J = 4.3, 9.3$ Hz, 1H), 8.07-8.02 (m, 2H), 7.71 (t, $J = 8.0$ Hz, 1H), 7.56-7.48 (m, 7H), 7.34 (dt, $J = 2.5, 9.5$ Hz, 1H), 7.02 (d, $J = 8.5$ Hz, 2H), 7.01-6.93 (m, 2H), 6.66 (t, $J = 55.5$ Hz, 1H), 3.68 (s, 2H); MS: 652.0 (M-CO $_2\text{H}$) $^-$.

Example 4**[0299]**







trans-2-(3-(3-(2-Cyanothiophen-3-yl)-1-tosyl-1H-pyrrolo[3,2-b]pyridin-2-yl)phenyl)cyclopropane-1-carboxylic acid (**4**)

[0300] A mixture of compound **2/4** (178 mg, 0.32 mmol) and LiOH·H₂O (67 mg, 1.62 mmol) in THF (4.1 mL), MeOH (4.1 mL) and water (0.81 mL) was stirred at rt for 3 h, adjusted to pH = 3 with 1N HCl, concentrated, diluted with EA (50 mL) and washed with water (3 x 5 mL) and brine (5 mL). The organic layer was concentrated under reduced pressure, the residue was dissolved in DMF (2.5 mL), filtrated and the filtrate was purified by prep-HPLC to give compound **4** as a white solid. $^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.90 (dd, $J = 8.5, 1.5$ Hz, 1H), 8.60 (dd, $J = 4.8, 0.8$ Hz, 1H), 7.79 (d, $J =$

5.0 Hz, 1H), 7.62 (dd, $J = 8.5, 4.5$ Hz, 1H), 7.32-7.25 (m, 6H), 7.15-7.13 (m, 1H), 6.95 (d, $J = 5.0$ Hz, 1H), 6.76 (s, 1H), 2.39-2.38 (m, 4H), 1.77-1.73 (m, 1H), 1.55-1.51 (m, 1H), 1.25 (br s, 1H); MS: 540.1 ($M+1$)⁺.


Example 4/1 to 4/3

[0301] The following Examples were prepared similar as described for Example 4 using the appropriate starting material.

#	starting material	structure	analytical data
4/1	 2/5		¹ H-NMR (500 MHz, CD ₃ OD) δ : 9.84 (s, 1H), 8.65 (d, $J = 6.0$ Hz, 1H), 8.03 (d, $J = 6.0$ Hz, 1H), 7.87 (d, $J = 5.0$ Hz, 1H), 7.34-7.28 (m, 6H), 7.12 (d, $J = 6.0$ Hz, 1H), 6.99 (d, $J = 5.0$ Hz, 1H), 6.71 (s, 1H), 2.42-2.36 (m, 4H), 1.76-1.73 (m, 1H), 1.54-1.51 (m, 1H), 1.23 (br s, 1H); MS: 540.1 ($M+1$) ⁺ .
4/2	 2/6		¹ H-NMR (500 MHz, CD ₃ OD) δ : 8.51 (dd, $J = 1.5, 5.0$ Hz, 1H), 7.87 (dd, $J = 1.5, 8.0$ Hz, 1H), 7.79 (d, $J = 8.5$ Hz, 1H), 7.68 (d, $J = 8.5$ Hz, 2H), 7.41 (dd, $J = 5.3, 7.7$ Hz, 1H), 7.34-7.26 (m, 4H), 7.21 (dd, $J = 7.5, 1.5$ Hz, 1H), 7.00 (s, 1H), 6.89 (d, $J = 5.0$ Hz, 1H), 2.47-2.44 (m, 1H), 2.40 (s, 3H), 1.80-1.78 (m, 1H), 1.56-1.52 (m, 1H), 1.30-1.29 (m, 1H); MS: 540.0 ($M+1$) ⁺ .
4/3	 2/7		¹ H-NMR (500 MHz, DMSO- <i>d</i> ₆) δ : 7.91 (d, $J = 5.5$ Hz, 1H), 7.68 (d, $J = 5.5$ Hz, 1H), 7.61-7.60 (m, 1H), 7.38-7.32 (m, 4H), 7.15 (t, $J = 7.8$ Hz, 1H), 6.81 (d, $J = 4.5$ Hz, 1H), 6.53-6.50 (m, 2H), 6.13 (d, $J = 1.5$ Hz, 1H), 3.90 (t, $J = 7.8$ Hz, 2H), 3.76 (t, $J = 6.3$ Hz, 2H), 3.53-3.49 (m, 1H), 2.35 (s, 3H); MS: 560.0 ($M+1$) ⁺ .

Example 5

[0302]













2-((5-(3-(3-(2-Cyanothiophen-3-yl)-1-tosyl-1H-indol-2-yl)phenyl)pyridin-3-yl)sulfonyl)acetic acid (5)

[0303] To a mixture of compound 2/8 (110 mg, 0.17 mmol) in MeOH (2 mL) and THF (1 mL) was added LiOH (2M, 0.3 mL) and the mixture was stirred at rt for 1 h. The mixture was neutralized with 1N HCl and extracted with EA (3 x). The combined organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound 5 as a white solid. ¹H-NMR (500 MHz, CD₃OD) δ : 9.09 (d, $J = 2.0$ Hz, 1H), 9.03 (d, $J = 2.0$ Hz, 1H), 8.44-8.42 (m, 2H), 7.85-7.80 (m, 2H), 7.57-7.39 (m, 6H), 7.33 (d, $J = 8.0$ Hz, 2H), 7.20 (d, $J = 8.0$ Hz, 2H), 7.06 (d, $J = 4.5$ Hz, 1H), 4.54 (s, 2H), 2.29 (s, 3H); MS: 554.1 ($M+1$)⁺.

Example 5/1 to 5/10

[0304] The following Examples were prepared similar as described for Example 5 using the appropriate starting material.

#	starting material	structure	analytical data
5/1	 2/9		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.30 (br s, 1H), 8.30 (d, J = 8.5 Hz, 1H), 8.05-8.00 (m, 2H), 7.92-7.86 (m, 2H), 7.77 (d, J = 8.0 Hz, 1H), 7.55-7.51 (m, 3H), 7.42-7.28 (m, 7H), 7.08 (d, J = 5.5 Hz, 1H), 4.96 (s, 2H), 4.61 (s, 2H), 2.26 (s, 3H); MS: 699.8 ($\text{M}+18$) $^+$.
5/2	 1/1		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.43 (s, 1H), 8.27 (d, J = 8.5 Hz, 1H), 8.01 (d, J = 5.0 Hz, 1H), 7.53-7.49 (m, 1H), 7.38-7.22 (m, 9H), 6.98-6.95 (m, 2H), 2.32 (s, 3H), 2.14 (s, 6H); MS: 562.8 ($\text{M}-1$) $^-$.
5/3	 1/2		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.05 (s, 1H), 8.31 (dd, J = 9.2, 4.4 Hz, 1H), 8.04-8.02 (m, 3H), 7.80 (d, J = 7.7 Hz, 1H), 7.68 (d, J = 8.3 Hz, 2H), 7.56-7.47 (m, 2H), 7.44-7.33 (m, 3H), 7.30-7.26 (m, 3H), 7.19 (dd, J = 8.6, 2.5 Hz, 1H), 7.07 (d, J = 5.1 Hz, 1H), 2.25 (s, 3H); MS: m/z 590.6 ($\text{M}-1$) $^-$.
5/4	 7		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.91 (br s, 1H), 8.27 (dd, J = 9.3, 4.8 Hz, 1H), 8.00 (d, J = 5.0 Hz, 1H), 7.45-7.33 (m, 7H), 7.22-7.17 (m, 3H), 6.97 (d, J = 5.0 Hz, 1H), 2.33 (s, 3H), 1.48 (s, 6H); MS: 583.1 ($\text{M}+1$) $^+$.
5/5	 1/3		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.41 (dd, J = 9.0, 4.5 Hz, 1H), 7.85 (d, J = 5.0 Hz, 1H), 7.64 (d, J = 7.5 Hz, 1H), 7.57-7.50 (m, 2H), 7.39 (d, J = 8.0 Hz, 2H), 7.32-7.23 (m, 6H), 7.07 (dd, J = 8.5, 2.5 Hz, 1H), 7.02 (d, J = 5.0 Hz, 1H), 6.90 (dd, J = 7.3, 1.8 Hz, 1H), 2.33 (s, 3H); MS: 609.9 ($\text{M}+1$) $^+$.

(continued)



#

starting material

structure



analytical data

5/6





$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.73 (d, $J = 5.5$ Hz, 1H), 8.30 (dd, $J = 9.3, 4.3$ Hz, 1H), 8.08-8.04 (m, 5H), 7.92 (s, 1H), 7.48-7.40 (m, 3H), 7.33-7.30 (m, 3H), 7.23 (dd, $J = 8.5, 2.0$ Hz, 1H), 7.12 (d, $J = 5.0$ Hz, 1H), 2.29 (s, 3H); MS: 594.1 ($\text{M}+1$) $^+$.

5/7





$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.28 (dd, $J = 9.0, 4.5$ Hz, 1H), 7.99 (d, $J = 4.5$ Hz, 1H), 7.61 (dd, $J = 6.8, 2.3$ Hz, 2H), 7.50 (dd, $J = 6.8, 1.8$ Hz, 2H), 7.39-7.35 (m, 1H), 7.16-7.12 (m, 2H), 6.95 (d, $J = 5.0$ Hz, 1H), 6.54 (d, $J = 7.5$ Hz, 1H), 6.48 (dd, $J = 8.3, 1.8$ Hz, 1H), 6.23 (s, 1H), 3.89 (t, $J = 7.8$ Hz, 2H), 3.79 (t, $J = 6.3$ Hz, 2H), 3.48-3.43 (m, 1H); MS: 589.6 ($\text{M}-1$) $^-$.

5/8





$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.32-8.27 (m, 2H), 7.99 (d, $J = 5.0$ Hz, 1H), 7.68 (dd, $J = 8.8, 2.8$ Hz, 1H), 7.39-7.35 (m, 1H), 7.17-7.13 (m, 2H), 6.96-6.91 (m, 2H), 6.55 (d, $J = 7.5$ Hz, 1H), 6.48 (dd, $J = 8.0, 2.0$ Hz, 1H), 6.23 (s, 1H), 3.90-3.87 (m, 5H), 3.79 (t, $J = 6.5$ Hz, 2H), 3.46-3.43 (m, 1H); MS: 586.7 ($\text{M}-1$) $^-$.

5/9



$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.41-8.38 (m, 1H), 7.80 (d, $J = 5.0$ Hz, 1H), 7.34-7.21 (m, 7H), 7.09 (d, $J = 7.0$ Hz, 1H), 7.04-7.02 (m, 1H), 6.97 (s, 1H), 6.90 (d, $J = 5.0$ Hz, 1H), 3.11-3.07 (m, 1H), 2.96-2.94 (m, 1H), 2.79-2.74 (m, 1H), 2.50-2.44 (m, 1H), 2.36 (s, 3H), 2.08-1.98 (m, 2H), 1.87-1.82 (m, 1H), 1.59-1.51 (m, 1H), 0.86 (d, $J = 6.0$ Hz, 3H); MS: 614.0 ($\text{M}+1$) $^+$.


5/10



$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.53 (dd, $J = 9.2, 4.3$ Hz, 1H), 8.12 (s, 1H), 8.06-8.04 (m, 1H), 7.99-7.97 (m, 1H), 7.70-7.14 (m, 13H), 6.77 (t, $J = 55.5$ Hz, 1H), 4.37-4.27 (m, 1H), 3.78-3.36 (m, 3H), 3.24-3.20 (m, 1H); MS: 704.1 ($\text{M}+1$) $^+$.

Example 6

[0305]



3'-(3-(2-Cyanothiophen-3-yl)-5-fluoro-1-tosyl-1H-indol-2-yl)-N-hydroxy-[1,1'-biphenyl]-4-carboxamide (6)

[0306] To a mixture of compound **5/3** (120 mg, 0.20 mmol) in DMF (5 mL) was added hydroxylamine hydrochloride (27 mg, 0.40 mmol), HATU (114 mg, 0.30 mmol) and DIPEA (103 mg, 0.80 mmol) and the mixture was stirred at rt overnight, diluted with EA (40 mL) and washed with H₂O (30 mL), 1N HCl (20 mL) and brine (30 mL). The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound **6** as a white solid. ¹H-NMR (500 MHz, DMSO-d₆) δ: 10.69 (br s, 1H), 9.15 (br s, 1H), 8.31 (dd, J = 9.3, 4.3 Hz, 1H), 8.02 (d, J = 5.0 Hz, 1H), 7.86 (d, J = 8.5 Hz, 2H), 7.78 (d, J = 8.0 Hz, 1H), 7.62 (d, J = 8.5 Hz, 2H), 7.50-7.47 (m, 2H), 7.41-7.25 (m, 6H), 7.19 (dd, J = 8.5, 2.5 Hz, 1H), 7.07 (d, J = 5.0 Hz, 1H), 2.25 (s, 3H); MS: 605.8 (M-1)⁻.

Example 6/1 to 6/3



[0307] The following Example was prepared similar as described for Example 6 using the appropriate starting material.

#	starting material	structure	analytical data
6/1	5/9		¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.39 (d, J = 8.5 Hz, 1H), 7.82 (d, J = 5.5 Hz, 1H), 7.53-7.23 (m, 9H), 7.04-7.01 (m, 2H), 6.94 (d, J = 5.0 Hz, 1H), 2.43-2.38 (m, 1H), 1.76-1.72 (m, 1H), 1.54-1.50 (m, 1H), 1.25-1.20 (m, 1H); MS: 574.1 (M+1) ⁺ .
6/2	5/3		¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.29 (dd, J = 9.0, 4.0 Hz, 1H), 8.04-7.96 (m, 4H), 7.78 (d, J = 8.5 Hz, 1H), 7.61 (d, J = 8.5 Hz, 2H), 7.50-7.36 (m, 6H), 7.26 (d, J = 8.0 Hz, 2H), 7.17 (dd, J = 8.0, 2.5 Hz, 1H), 7.06 (d, J = 5.0 Hz, 1H), 2.25 (s, 3H); MS: 589.8 (M-1) ⁻ .


(continued)

#	starting material	structure	analytical data
---	-------------------	-----------	-----------------


6/3



$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.43 (dd, $J = 4.4, 9.2$ Hz, 1H), 8.05 (d, $J = 7.6$ Hz, 2H), 7.96 (d, $J = 1.2$ Hz, 1H), 7.84 (dd, $J = 1.8, 7.8$ Hz, 1H), 7.71 (t, $J = 8.0$ Hz, 1H), 7.54-7.30 (m, 9H), 7.09 (s, 1H), 6.94 (dd, $J = 2.4, 8.4$ Hz, 1H), 6.67 (t, $J = 55.4$ Hz, 1H), 4.03 (s, 2H), 1.51 (s, 9H); MS: 793.1 (M-1)-.

Example 7**[0308]**Methyl 4-(3-(3-(2-cyanothiophen-3-yl)-5-fluoro-1-tosyl-1H-indol-2-yl)phenyl)-2,2-dimethylbut-3-ynoate (**7**)

[0309] To a solution of compound **1** (234 mg, 0.52 mmol) in Et_3N (1.5 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (47 mg), CuI (80 mg), PPh_3 (11 mg) and methyl 2,2-dimethylbut-3-ynoate (78 mg, 0.62 mmol). The mixture was stirred at 60°C under N_2 for 4 h, cooled, filtered, concentrated and purified by FCC (PE:EA = 8:1) to give compound **7** as a yellow solid.









Example 8**[0310]**3-(5-Fluoro-2-(4'-((4-methylpiperazin-1-yl)methyl)-[1,1'-biphenyl]-3-yl)-1-tosyl-1H-indol-3-yl)thiophene-2-carbonitrile (**8**)

[0311] To a solution of compound **1** (250 mg, 0.45 mmol) in dioxane (20 mL) was added (3-((4-methylpiperazin-1-yl)methyl)phenyl)boronic acid (116 mg, 0.50 mmol), Cs_2CO_3 (293 mg, 0.90 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (52 mg, 50 μmol). The mixture was stirred at 100°C overnight under N_2 , cooled, filtered, concentrated and purified by prep-HPLC to give compound **8** as a white solid. $^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.43 (dd, $J = 4.5, 9.5$ Hz, 1H), 7.82 (d, $J = 5.0$ Hz, 1H),

7.67 (d, $J = 8.0$ Hz, 1H), 7.47-7.41 (m, 5H), 7.31-7.15 (m, 7H), 7.06 (dd, $J = 2.5, 8.5$ Hz, 1H), 6.96 (d, $J = 5.0$ Hz, 1H), 3.61 (s, 2H), 2.61-2.48 (m, 8H), 2.31 (s, 3H), 2.27 (s, 3H); MS: 661.0 ($M+1$)⁺.

Example 8/1 to 8/8


[0312] The following Example was prepared similar as described for Example 8 using the appropriate starting materials.

#	starting material(s)	structure	analytical data
8/1			¹ H-NMR (500 MHz, DMSO-d ₆) δ : 8.31 (dd, $J = 4.5, 9.0$ Hz, 1H), 8.02 (d, $J = 5.0$ Hz, 1H), 7.72 (d, $J = 8.0$ Hz, 1H), 7.49-7.35 (m, 9H), 7.27-7.17 (m, 4H), 7.06 (d, $J = 5.0$ Hz, 1H), 3.43 (s, 2H), 2.25 (s, 3H), 2.17 (s, 6H); MS: 606.0 ($M+1$) ⁺ .
8/2	 P21		
8/3	 1/138		¹ H-NMR (500 MHz, CD ₃ OD) δ : 8.60 (d, $J = 5.9$ Hz, 1H), 8.43 (d, $J = 6.0$ Hz, 1H), 8.37 (s, 1H), 8.06 (s, 1H), 7.99-7.97 (m, 1H), 7.76-7.08 (m, 13H), 6.74 (t, $J = 55.5$ Hz, 1H), 4.65-4.56 (m, 1H), 4.53-4.49 (m, 1H), 4.41-4.37 (m, 1H), 4.30-4.26 (m, 1H), 3.90-3.78 (m, 1H); MS: 671.1 ($M+1$) ⁺ .
8/4	 1/139		

(continued)

#	starting material(s)	structure	analytical data
8/5	 1/140		
8/6	 1/141		
8/7	 1/145		¹ H-NMR (500 MHz, CD ₃ OD) δ : 8.81 (dd, J = 8.5, 1.5 Hz, 1H), 8.56 (dd, J = 4.8, 1.3 Hz, 1H), 8.06 (d, J = 1.5 Hz, 1H), 7.98 (d, J = 8.0 Hz, 1H), 7.77 (d, J = 7.5 Hz, 1H), 7.65-7.04 (m, 13H), 6.70 (t, J = 55.5 Hz, 1H); MS: 640.2 (M+1) ⁺ .
8/8	 1/139 Pd(dppf)Cl ₂ K ₂ CO ₃ 95°C, 8 h in dioxane/water (17:1)		¹ H-NMR (400 MHz, CD ₃ OD) δ : 8.42 (dd, J = 9.4, 4.2 Hz, 1H), 8.05-8.03 (m, 3H), 7.96 (dd, J = 8.2, 1.8 Hz, 1H), 7.70 (dd, J = 8.4, 7.6 Hz, 1H), 7.51-7.44 (m, 8H), 7.32 (td, J = 9.2, 2.4 Hz, 1H), 6.97 (s, 1H), 6.94 (dd, J = 8.4, 2.4 Hz, 1H), 6.62 (t, J = 55.4 Hz, 1H), 2.00 (s, 3H); MS: 757.0 (M-1) ⁻ .

Example 9**[0313]**




Methyl 3'-(3-(2,6-bis(difluoromethyl)phenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-2-chloro-[1,1'-biphenyl]-4-carboxylate (**9**)

[0314] To a solution of compound **1/55** (180 mg, 0.26 mmol) in DCM (10.0 mL) was added DAST (209 mg, 1.30 mmol) and the mixture was stirred at rt overnight, poured into EA (200 mL) and washed with H₂O (2 x 20 mL) and brine (20 mL). The organic layer was dried over Na₂SO₄, concentrated and purified by prep-TLC (EA:PE = 1:3) to give compound **9** as a colorless oil.


Example 9/1

[0315] The following Example was prepared similar as described for Example 9 using the appropriate starting material.



Example 10

[0316]











2-(2-Chloro-3'-(3-(2-cyanothiophen-3-yl)-5-fluoro-1-tosyl-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxamido)ethane-1-sulfonic acid (**10**)

[0317] To a solution of compound **3/48** (100 mg, 0.20 mmol) in DMF (10 mL) was added EDCI (100 mg, 0.50 mmol), DMAP (60 mg, 0.50 mmol) and 2-aminoethane-1-sulfonic acid (22 mg, 0.20 mmol). The mixture was stirred at rt for 12 h, diluted with water (100 mL) and extracted with EA (3 x 100 mL). The combined organic layer was washed with brine (50 mL), concentrated and purified by FCC (PE:EA = 1:4) to give compound **10** as a white solid. ¹H-NMR (500 MHz,

DMSO- d_6) δ : 8.68 (t, J = 5.3 Hz, 1H), 8.29 (dd, J = 4.5, 9.0 Hz, 1H), 8.03 (d, J = 5.0 Hz, 1H), 7.92 (d, J = 2.0 Hz, 1H), 7.81 (dd, J = 1.8, 8.3 Hz, 1H), 7.54-7.17 (m, 10H), 7.03-7.01 (m, 2H), 3.56-3.52 (m, 2H), 2.70-2.67 (m, 2H), 2.21 (s, 3H); MS: 731.9 (M-1)⁻.


5 Example 1011 to 10/4

[0318] The following Examples were prepared similar as described for Example 10 using the appropriate starting material.

10	#	starting material	structure	analytical data
15	10/1			¹ H-NMR (500 MHz, CD ₃ OD) δ : 8.45-8.43 (m, 1H), 8.07 (brs, 2H), 7.94 (d, J = 2.0 Hz, 1H), 7.82 (dd, J = 8.0, 1.5 Hz, 1H), 7.78 (t, J = 7.8 Hz, 1H), 7.55-7.43 (m, 7H), 7.36-7.28 (m, 2H), 7.04 (s, 1H), 6.95 (dd, J = 8.5, 2.5 Hz, 1H), 6.68 (t, J = 55.5 Hz, 1H), 3.84 (t, J = 6.5 Hz, 2H), 3.13 (t, J = 6.5 Hz, 2H); MS: 786.9 (M-1) ⁻ .
25	10/2			¹ H-NMR (500 MHz, CD ₃ OD) δ : 8.44 (dd, J = 9.3, 4.3 Hz, 1H), 8.06 (br s, 2H), 7.72 (t, J = 7.8 Hz, 1H), 7.59-7.47 (m, 9H), 7.35-7.32 (m, 2H), 7.02-6.58 (m, 3H), 3.97-3.77 (m, 2H), 3.24-3.10 (m, 5H); MS: 801.0 (M-1) ⁻ .
35	10/3			¹ H-NMR (500 MHz, CD ₃ OD) δ : 8.42 (dd, J = 9.0, 4.5 Hz, 1H), 7.97 (d, J = 1.5 Hz, 1H), 7.85-7.81 (m, 2H), 7.56-7.49 (m, 6H), 7.44 (d, J = 7.0 Hz, 1H), 7.39 (d, J = 8.0 Hz, 1H), 7.31 (td, J = 9.0, 2.5 Hz, 1H), 7.09-7.06 (m, 2H), 6.98 (d, J = 5.0 Hz, 1H), 6.71 (t, J = 55.5 Hz, 1H), 3.84 (t, J = 6.8 Hz, 2H), 3.13 (t, J = 6.8 Hz, 2H); MS: 768.0 (M-1) ⁻ .
45	10/4			¹ H-NMR (400 MHz, CDaOD) δ : 8.37 (dd, J = 9.2, 4.4 Hz, 1H), 7.59 (d, J = 8.0 Hz, 2H), 7.47 (d, J = 8.0 Hz, 2H), 7.40-7.35 (m, 3H), 7.26-7.14 (m, 4H), 6.95-6.64 (m, 3H), 3.69-3.61 (m, 2H), 3.00 (t, J = 6.6 Hz, 2H), 2.33-2.28 (m, 1H), 1.64 (brs, 1H), 1.47-1.42 (m, 1H), 1.06 (br s, 1H); MS: 735.0 and 737.0 (M-1) ⁻ .

55 Example 11

[0319]



2-Chloro-3'-(3-(2-cyanophenyl)-5-fluoro-1-tosyl-1H-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid (**11**)











[0320] To a solution of compound **1/32** (165 mg, 0.26 mmol) in HCl/dioxane (4N, 5 mL) was added H₂O (0.5 mL). The mixture was stirred at 90°C overnight, cooled, concentrated, diluted with water and extracted with EA (3 x). The combined organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound **11** as a white solid. ¹H-NMR (500 MHz, DMSO-d₆) δ: 13.36 (br s, 1H), 8.28 (dd, J = 4.5, 9.5 Hz, 1H), 7.97-7.94 (m, 2H), 7.86 (d, J = 7.5 Hz, 1H), 7.66 (t, J = 7.5 Hz, 1H), 7.55-7.31 (m, 9H), 7.24 (d, J = 8.5 Hz, 2H), 7.02-7.00 (m, 2H), 2.21 (s, 3H); MS: 619.0 (M-1)⁻.

Example 11/1 to 11/69

[0321] The following Examples were prepared similar as described for Example 11 using the appropriate starting material.

#	starting material	structure	analytical data
11/1			¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.41 (dd, J = 9.3, 4.3 Hz, 1H), 8.08 (d, J = 1.0 Hz, 1H), 8.02 (dd, J = 1.8, 8.3 Hz, 1H), 7.69-7.64 (m, 1H), 7.49-7.26 (m, 8H), 7.18-1.15 (m, 3H), 7.02 (s, 1H), 6.97 (dd, J = 2.5, 8.5 Hz, 1H), 2.25 (s, 3H); MS: 637.0 (M-1) ⁻ .
11/2			¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.39 (dd, J = 9.3, 4.3 Hz, 1H), 8.05 (d, J = 1.5 Hz, 1H), 7.98 (dd, J = 8.0, 1.5 Hz, 1H), 7.51-7.43 (m, 4H), 7.39 (d, J = 8.0 Hz, 1H), 7.31 (d, J = 8.0 Hz, 1H), 7.28-6.98 (m, 7H), 6.88 (dd, J = 8.5, 2.5 Hz, 1H), 2.51 (s, 3H), 2.24 (s, 3H); MS: 633.0 (M-1) ⁻ .
11/3			¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.38 (dd, J = 9.0, 4.5 Hz, 1H), 8.07 (d, J = 1.5 Hz, 1H), 8.00 (dd, J = 8.0, 1.5 Hz, 1H), 7.57-7.54 (m, 1H), 7.48-7.44 (m, 3H), 7.37 (d, J = 7.5 Hz, 1H), 7.27-7.14 (m, 6H), 7.03 (s, 1H), 6.88 (dd, J = 8.3, 2.8 Hz, 1H), 6.83 (br s, 1H), 3.95 (s, 3H), 2.24 (s, 3H); MS: 649.0 (M-1) ⁻ .











(continued)

#	starting material	structure	analytical data
11/4	 1/36		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.38 (dd, J = 9.3, 4.3 Hz, 1H), 8.07 (d, J = 1.5 Hz, 1H), 7.99 (dd, J = 8.0, 1.5 Hz, 1H), 7.63-7.54 (m, 1H), 7.45-7.42 (m, 4H), 7.36 (d, J = 8.0 Hz, 1H), 7.30-7.14 (m, 6H), 7.02 (s, 1H), 6.88 (dd, J = 8.5, 2.5 Hz, 1H), 2.39 (s, 3H), 2.24 (s, 3H); MS: 633.0 (M-1) $^-$.
11/5	 1/37		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.39 (dd, J = 9.0, 4.5 Hz, 1H), 8.07 (d, J = 2.0 Hz, 1H), 8.00 (dd, J = 8.0, 1.5 Hz, 1H), 7.62 (d, J = 8.0 Hz, 1H), 7.46-7.44 (m, 3H), 7.36 (d, J = 8.5 Hz, 1H), 7.33-7.14 (m, 7H), 7.03 (s, 1H), 6.89 (dd, J = 8.3, 2.8 Hz, 1H), 2.35 (s, 3H), 2.24 (s, 3H); MS: 633.0 (M-1) $^-$.
11/6	 1/38		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.40 (dd, J = 9.3, 4.3 Hz, 1H), 8.08 (s, 1H), 8.00 (dd, J = 8.0, 1.5 Hz, 1H), 7.58-7.15 (m, 13H), 6.71 (dd, J = 8.0, 2.5 Hz, 1H), 2.25 (s, 3H), 1.90 (s, 3H); MS: 633.0 (M-1) $^-$.
11/7	 1/48		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.43 (dd, J = 9.0, 4.5 Hz, 1H), 8.03 (d, J = 7.0 Hz, 1H), 7.85 (d, J = 5.5 Hz, 1H), 7.57-7.44 (m, 5H), 7.38 (d, J = 8.5 Hz, 2H), 7.30 (dt, J = 2.7, 9.2 Hz, 1H), 7.21 (d, J = 10.5 Hz, 1H), 7.15 (s, 1H), 7.07 (dd, J = 8.5, 2.5 Hz, 1H), 6.99 (d, J = 5.5 Hz, 1H), 5.35 (d, J = 47.0 Hz, 2H); MS: 660.9 (M-1) $^-$.
11/8	 1/49		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.43 (dd, J = 9.3, 4.3 Hz, 1H), 8.03 (d, J = 6.5 Hz, 1H), 7.86 (d, J = 5.0 Hz, 1H), 7.58-7.49 (m, 6H), 7.43 (d, J = 7.5 Hz, 1H), 7.31 (td, J = 9.0, 2.5 Hz, 1H), 7.22 (d, J = 10.5 Hz, 1H), 7.16 (s, 1H), 7.09 (dd, J = 2.5, 8.5 Hz, 1H), 7.01 (d, J = 5.0 Hz, 1H), 6.73 (t, J = 55.5 Hz, 1H); MS: 678.9 (M-1) $^-$.











(continued)

#	starting material	structure	analytical data
11/9	 1/50	 1/50	$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 9.33 (s, 1H), 8.46 (dd, J = 9.3, 4.3 Hz, 1H), 8.04 (d, J = 6.5 Hz, 1H), 7.61-7.50 (m, 6H), 7.41 (d, J = 8.0 Hz, 1H), 7.33 (d, J = 2.5 Hz, 1H), 7.27 (d, J = 11.0 Hz, 1H), 7.21-7.18 (m, 2H), 6.73 (t, J = 55.5 Hz, 1H); MS: 679.9 ($\text{M}-1$) $^-$.
11/10	 1/51	 1/51	$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.72 (d, J = 4.5 Hz, 1H), 8.55 (br s, 1H), 8.44 (dd, J = 4.3, 9.3 Hz, 1H), 8.01 (d, J = 6.5 Hz, 1H), 7.78 (d, J = 5.0 Hz, 1H), 7.60-7.04 (m, 11H), 6.73 (t, J = 55.5 Hz, 1H); MS: 673.9 ($\text{M}-1$) $^-$.
11/11	 1/52	 1/52	$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.90 (d, J = 2.5 Hz, 1H), 8.71 (d, J = 2.5 Hz, 1H), 8.47 (dd, J = 9.5, 4.5 Hz, 1H), 8.01 (d, J = 5.0 Hz, 1H), 7.78-7.18 (m, 11H), 6.73 (t, J = 55.5 Hz, 1H); MS: 674.9 ($\text{M}-1$) $^-$.
11/12	 1/53	 1/53	$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.45 (dd, J = 4.3, 9.3 Hz, 1H), 8.08-8.00 (m, 4H), 7.72 (t, J = 7.8 Hz, 1H), 7.55-7.32 (m, 9H), 7.08 (s, 1H), 6.95 (dd, J = 2.5, 8.5 Hz, 1H), 6.68 (t, J = 55.5 Hz, 1H); MS: 679.9 ($\text{M}-1$) $^-$.
11/13	 1/54	 1/54	$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.39 (dd, J = 4.8, 9.3 Hz, 1H), 8.08 (d, J = 1.5 Hz, 1H), 7.98 (dd, J = 1.5, 8.0 Hz, 1H), 7.56-7.25 (m, 12H), 7.21 (s, 1H), 6.73 (t, J = 55.3 Hz, 1H), 6.68 (dd, J = 2.5, 8.5 Hz, 1H); MS: 697.9 ($\text{M}-1$) $^-$.











(continued)

#	starting material	structure	analytical data
11/14	 29		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.41 (dd, $J = 4.2, 9.0$ Hz, 1H), 8.07 (d, $J = 1.6$ Hz, 1H), 7.94 (dd, $J = 1.6, 8.0$ Hz, 1H), 7.57-7.40 (m, 7H), 7.29-7.15 (m, 4H), 7.04 (d, $J = 8.4$ Hz, 2H), 6.75 (t, $J = 55.6$ Hz, 1H), 6.57 (d, $J = 2.6, 8.2$ Hz, 1H), 1.63 (s, 6H); MS: 658.0 (M-1) $^-$.
11/15	 9		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.44 (dd, $J = 4.4, 9.2$ Hz, 1H), 8.08 (d, $J = 1.6$ Hz, 1H), 7.97 (dd, $J = 1.4, 3.8$ Hz, 1H), 7.83 (d, $J = 7.6$ Hz, 2H), 7.73 (t, $J = 7.8$ Hz, 1H), 7.61-7.55 (m, 4H), 7.47-7.23 (m, 5H), 6.91 (s, 1H), 6.77-6.63 (m, 2H), 6.01 (t, $J = 55.0$ Hz, 2H); MS: 729.9 (M-1) $^-$.
11/16	 35		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.39 (s, 1H), 8.78 (d, $J = 2.5$ Hz, 1H), 8.34 (dd, $J = 9.0, 4.5$ Hz, 1H), 8.12-7.93 (m, 5H), 7.57-7.55 (m, 2H), 7.50-7.41 (m, 3H), 7.23-7.21 (m, 2H), 7.04-7.02 (m, 1H); MS: 679.6 (M-1) $^-$.
11/17	 1/104		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.30 (dd, $J = 9.2, 4.4$ Hz, 1H), 8.04 (d, $J = 5.1$ Hz, 1H), 7.99-7.88 (m, 2H), 7.70-7.49 (m, 6H), 7.47-7.35 (m, 3H), 7.20 (dd, $J = 8.6, 2.6$ Hz, 1H), 7.12-7.10 (m, 1H), 7.05-7.03 (m, 1H), 7.01 (t, $J = 55.0$ Hz, 1H); MS: 660.9 (M-1) $^-$.
11/18	 1/105		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.30 (dd, $J = 9.2, 4.4$ Hz, 1H), 8.04 (d, $J = 5.1$ Hz, 1H), 7.99-7.88 (m, 2H), 7.58-7.45 (m, 4H), 7.45-7.35 (m, 3H), 7.31 (t, $J = 8.8$ Hz, 2H), 7.19 (dd, $J = 8.6, 2.6$ Hz, 1H), 7.14 (s, 1H), 7.03 (d, $J = 4.9$ Hz, 1H); MS: 628.9 (M-1) $^-$.











(continued)

#	starting material	structure	analytical data
11/ 19	 1/106		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.42 (s, 1H), 8.31 (dd, $J = 9.2, 4.5$ Hz, 1H), 8.05-7.93 (m, 3H), 7.55-7.53 (m, 2H), 7.48-7.36 (m, 4H), 7.24-7.11 (m, 4H), 7.05 (d, $J = 4.8$ Hz, 1H), 2.15 (s, 3H); MS: 642.9 (M-1) $^-$.
11/ 20	 1/107		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.46 (s, 1H), 8.31-8.28 (m, 1H), 8.05 (d, $J = 5.1$ Hz, 1H), 8.02-7.92 (m, 4H), 7.59-7.50 (m, 4H), 7.48-7.37 (m, 3H), 7.22-7.20 (m, 1H), 7.12 (s, 1H), 7.04 (d, $J = 5.0$ Hz, 1H); MS: 635.9 (M-1) $^-$.
11/ 21	 1/108		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.36 (dd, $J = 9.2, 4.4$ Hz, 1H), 8.07 (d, $J = 1.6$ Hz, 1H), 7.97 (dd, $J = 8.0, 1.6$ Hz, 1H), 7.64 (d, $J = 5.0$ Hz, 1H), 7.52-7.33 (m, 7H), 7.31-7.22 (m, 2H), 7.12 (s, 1H), 6.91 (dd, $J = 8.5, 2.5$ Hz, 1H), 6.82 (br s, 1H), 6.34 (t, $J = 54.8$ Hz, 1H); MS: 669.8 (M-1) $^-$.
11/ 22	 1/110		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.38 (dd, $J = 9.2, 4.3$ Hz, 1H), 8.11 (d, $J = 1.3$ Hz, 1H), 8.01 (d, $J = 6.9$ Hz, 1H), 7.70 (d, $J = 4.0$ Hz, 1H), 7.57-7.24 (m, 10H), 6.85-6.63 (m, 3H); MS: 703.9 (M-1) $^-$.
11/ 23	 1/111		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.78 (d, $J = 4.6$ Hz, 1H), 8.42 (dd, $J = 9.2, 4.3$ Hz, 1H), 8.21 (d, $J = 7.5$ Hz, 1H), 8.09 (d, $J = 1.5$ Hz, 1H), 8.01 (dd, $J = 8.0, 1.5$ Hz, 1H), 7.70-7.06 (m, 11H), 6.79-6.77 (m, 1H), 6.76 (t, $J = 56.0$, 1H); MS: 699.0 (M-1) $^-$.











(continued)

#	starting material	structure	analytical data
11/ 24	 29/1		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.36 (dd, $J = 9.1, 4.4$ Hz, 1H), 8.07 (d, $J = 1.5$ Hz, 1H), 7.97 (dd, $J = 7.9, 1.6$ Hz, 1H), 7.56-7.43 (m, 7H), 7.33 (d, $J = 8.0$ Hz, 1H), 7.25 (dt, $J = 4.4, 8.8$ Hz, 1H), 7.15 (s, 1H), 6.82-6.79 (m, 1H), 6.73 (t, $J = 55.2$ Hz, 1H), 1.70 (s, 6H); MS: 650.1 (M+H) ⁺ .
11/ 25	 P1/113		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.37 (dd, $J = 9.1, 4.3$ Hz, 1H), 8.09 (d, $J = 1.4$ Hz, 1H), 7.99 (dd, $J = 8.0, 1.5$ Hz, 1H), 7.68-6.89 (m, 11H), 6.88-6.86 (m, 1H), 6.73 (t, $J = 55.0$ Hz, 1H); MS: 688.0 (M-1) ⁻ .
11/ 26	 1/115		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.84 (dd, $J = 5.0, 1.6$ Hz, 1H), 8.45 (dd, $J = 9.2, 4.3$ Hz, 1H), 8.16 (dd, $J = 8.0, 1.5$ Hz, 1H), 8.07 (s, 1H), 8.00 (d, $J = 8.0$ Hz, 1H), 7.82-7.10 (m, 11H), 7.05 (dd, $J = 8.4, 2.5$ Hz, 1H), 6.71 (t, $J = 55.6$ Hz, 1H); MS: 655.9 (M-1) ⁻ .
11/ 27	 2/19		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.28 (dd, $J = 9.0, 4.5$ Hz, 1H), 7.88 (d, $J = 6.0$ Hz, 1H), 7.76-7.36 (m, 12H), 7.12-6.90 (m, 4H), 3.82 (s, 3H); MS: 684.9 (M-1) ⁻ .
11/ 28	 2/20		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.41 (dd, $J = 9.0, 4.0$ Hz, 1H), 7.78 (d, $J = 7.5$ Hz, 2H), 7.61-7.39 (m, 10H), 7.30-7.26 (m, 2H), 6.95 (d, $J = 8.5$ Hz, 1H), 6.90-6.87 (m, 1H), 6.70 (t, $J = 55.5$ Hz, 1H), 6.08 (d, $J = 4.5$ Hz, 2H); MS: 665.0 (M-1) ⁻ .

(continued)

#	starting material	structure	analytical data
11/ 29	 2/21		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.40 (dd, J = 9.2, 4.4 Hz, 1H), 7.79-7.50 (m, 10H), 7.35-7.17 (m, 6H), 6.88 (dd, J = 8.4, 2.4 Hz, 1H), 6.69 (t, J = 55.6 Hz, 1H), 3.83 (s, 3H); MS: 651.0 ($\text{M}-1$) $^-$.
11/ 30	 2/22		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.43-8.40 (m, 2H), 8.29 (dd, J = 8.2, 1.4 Hz, 1H), 7.77-7.75 (m, 1H), 7.67-7.26 (m, 13H), 6.91 (dd, J = 2.8, 8.4 Hz, 1H), 6.74 (t, J = 55.6 Hz, 1H); MS: 646.0 ($\text{M}-1$) $^-$.
11/ 31	 2/23		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.41 (dd, J = 9.2, 4.4 Hz, 1H), 7.87 (dd, J = 1.6, 8.0 Hz, 1H), 7.76 (dd, J = 1.4, 11.0 Hz, 1H), 7.65-7.26 (m, 13H), 6.89 (dd, J = 2.4, 8.4 Hz, 1H), 6.72 (t, J = 55.6 Hz, 1H); MS: 639.0 ($\text{M}-1$) $^-$.
11/ 32	 2/24		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.41 (dd, J = 9.2, 4.4 Hz, 1H), 7.83 (d, J = 8.4 Hz, 1H), 7.78-6.55 (m, 16H); MS: 670.9 ($\text{M}-1$) $^-$.
11/ 33	 1/119		$^1\text{H-NMR}$ (500 MHz, MeOD) δ : 8.40 (dd, J = 9.3, 4.3 Hz, 1H), 8.07 (d, J = 2.0 Hz, 1H), 7.99 (dd, J = 7.8, 1.8 Hz, 1H), 7.77 (d, J = 7.5 Hz, 1H), 7.64 (t, J = 7.5 Hz, 1H), 7.56-7.44 (m, 8H), 7.35-7.26 (m, 3H), 7.09 (br s, 1H), 6.90 (dd, J = 8.5, 2.5 Hz, 1H), 6.70 (t, J = 55.5 Hz, 1H); MS: 670.9 ($\text{M}-1$) $^-$.











(continued)

#	starting material	structure	analytical data
11/ 34	 1/120		$^1\text{H-NMR}$ (400 MHz, MeOD) δ : 8.42 (dd, J = 9.4, 4.2 Hz, 1H), 8.06 (d, J = 1.6 Hz, 1H), 7.98 (dd, J = 7.8, 1.4 Hz, 1H), 7.64-7.41 (m, 10H), 7.34-7.28 (m, 2H), 7.05 (br s, 1H), 6.90 (dd, J = 8.2, 2.6 Hz, 1H), 6.70 (t, J = 55.4 Hz, 1H); MS: 670.9 (M-1) $^-$.
11/ 35	 1/121		$^1\text{H-NMR}$ (500 MHz, MeOD) δ : 8.37 (dd, J = 9.3, 4.3 Hz, 1H), 8.08 (d, J = 1.0 Hz, 1H), 7.99 (dd, J = 7.5, 1.5 Hz, 1H), 7.56-7.23 (m, 12H), 7.10 (br s, 1H), 6.76 (dd, J = 1.8, 8.3 Hz, 1H), 6.72 (t, J = 55.5 Hz, 1H), 3.59 (s, 3H); MS: 670.9 (M-1) $^-$.
11/ 36	 1/122		$^1\text{H-NMR}$ (500 MHz, MeOD) δ : 8.41 (dd, J = 9.3, 4.3 Hz, 1H), 8.08 (d, J = 1.5 Hz, 1H), 8.00 (dd, J = 8.0, 1.5 Hz, 1H), 7.60-7.15 (m, 13H), 6.82-6.60 (m, 2H), 1.88 (s, 3H); MS: 670.9 (M-1) $^-$.
11/ 37	 35/1		$^1\text{H-NMR}$ (400 MHz, MeOD) δ : 8.68 (d, J = 2.0 Hz, 1H), 8.50 (dd, J = 9.2, 4.0 Hz, 1H), 8.09-7.98 (m, 5H), 7.83 (d, J = 8.4 Hz, 1H), 7.74 (t, J = 7.8 Hz, 1H), 7.55-7.38 (m, 5H), 7.20 (s, 1H), 6.97 (dd, J = 8.2, 2.6 Hz, 1H); MS: 698.9 (M-1) $^-$.
11/ 38	 9/1		$^1\text{H-NMR}$ (500 MHz, MeOD) δ : 8.42 (dd, J = 9.0, 4.5 Hz, 1H), 8.09 (d, J = 2.0 Hz, 1H), 7.98 (dd, J = 8.0, 1.5 Hz, 1H), 7.59-7.25 (m, 12H), 7.21 (br s, 1H), 6.77 (t, J = 55.5 Hz, 1H), 6.59 (dd, J = 8.0, 2.5 Hz, 1H), 5.90 (t, J = 55.5 Hz, 1H), 1.70 (s, 3H); MS: 694.0 (M-1) $^-$.











(continued)

#	starting material	structure	analytical data
5 C11/ 39	 C1/123		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.40 (dd, $J = 9.2, 4.4$ Hz, 1H), 8.07 (d, $J = 1.6$ Hz, 1H), 7.97 (dd, $J = 8.0, 1.6$ Hz, 1H), 7.56-7.45 (m, 7H), 7.34-7.21 (m, 5H), 7.14-7.11 (m, 4H), 6.73 (t, $J = 55.6$ Hz, 1H); MS: 630.0 ($\text{M}-1$) $^-$.
15 11/ 40	 1/124		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.41 (dd, $J = 9.0, 4.5$ Hz, 1H), 8.19 (dd, $J = 1.0, 4.5$ Hz, 1H), 8.08 (d, $J = 1.0$ Hz, 1H), 8.01 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.56-7.46 (m, 9H), 7.39 (d, $J = 8.0$ Hz, 1H), 7.29-7.25 (m, 1H), 7.14 (br s, 1H), 7.00 (dd, $J = 9.0, 2.5$ Hz, 1H), 6.74 (t, $J = 55.5$ Hz, 1H), 3.50 (s, 3H); MS: 663.0 ($\text{M}+1$) $^+$.
25 11/ 41	 1/125		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.44-8.40 (m, 2H), 8.07 (d, $J = 1.6$ Hz, 1H), 8.00 (dd, $J = 8.0, 1.2$ Hz, 1H), 7.67 (d, $J = 8.4$ Hz, 1H), 7.57-7.25 (m, 10H), 7.14 (br s, 1H), 6.98 (dd, $J = 8.6, 2.6$ Hz, 1H), 6.73 (t, $J = 55.6$ Hz, 1H), 6.51 (t, $J = 72.6$ Hz, 1H); MS: 699.0 ($\text{M}+1$) $^+$.
35 11/ 42	 1/126		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.32 (dd, $J = 9.0, 4.6$ Hz, 1H), 8.16 (d, $J = 1.6$ Hz, 1H), 8.07 (dd, $J = 7.8, 1.4$ Hz, 1H), 7.62-7.22 (m, 10H), 7.15 (dd, $J = 8.2, 2.6$ Hz, 1H), 6.70 (t, $J = 55.6$ Hz, 1H), 2.43-2.16 (m, 3H), 1.92-1.43 (m, 5H); MS: 659.0 ($\text{M}-1$) $^-$.
45 11/ 43	 1/129		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.67 (br s, 1H), 8.51-8.48 (m, 1H), 8.00-7.93 (m, 2H), 7.75 (d, $J = 9.5$ Hz, 1H), 7.63-7.00 (m, 13H), 6.73 (t, $J = 55.3$ Hz, 1H); MS: 671.0 ($\text{M}-1$) $^-$.











(continued)

#	starting material	structure	analytical data
111/44	 1/130		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.38 (dd, $J = 9.1, 4.3$ Hz, 1H), 8.08 (d, $J = 1.5$ Hz, 1H), 7.99 (dd, $J = 7.8, 1.4$ Hz, 1H), 7.56-7.415 (m, 7H), 7.35 (d, $J = 8.0$ Hz, 1H), 7.27-7.22 (m, 1H), 7.16 (s, 1H), 7.07 (dd, $J = 8.5, 2.5$ Hz, 1H), 6.84-6.60 (m, 4H), 5.65 (s, 2H); MS: 674.0 ($\text{M}-1$) $^-$.
111/45	 1/131		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.45 (dd, $J = 9.3, 4.3$ Hz, 1H), 8.01 (d, $J = 1.5$ Hz, 1H), 7.98 (dd, $J = 8.0, 1.5$ Hz, 1H), 7.85 (d, $J = 9.0$ Hz, 1H), 7.59-7.55 (m, 4H), 7.49-7.28 (m, 7H), 7.07-7.05 (m, 2H), 6.72 (t, $J = 55.5$ Hz, 1H); MS: 671.9 ($\text{M}-1$) $^-$.
111/46	 1/132		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.43-8.40 (m, 2H), 8.22-7.97 (m, 3H), 7.54-6.72 (m, 13H), 3.86 (s, 3H); MS: 725.0 ($\text{M}-1$) $^-$.
111/47	 1/133		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.34 (dd, $J = 9.3, 4.3$ Hz, 1H), 8.08 (s, 1H), 8.01 (d, $J = 8.0$ Hz, 1H), 7.64 (dd, $J = 2.0, 6.5$ Hz, 1H), 7.59-7.42 (m, 8H), 7.23-7.18 (m, 3H), 7.00 (dd, $J = 2.3, 8.8$ Hz, 1H), 6.72 (t, $J = 55.5$ Hz, 1H), 6.27-6.25 (m, 1H), 3.57 (s, 3H); MS: 661.0 ($\text{M}-1$) $^-$.
C111/48	 C1/134		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.42 (dd, $J = 9.0, 4.5$ Hz, 1H), 8.06 (d, $J = 1.5$ Hz, 1H), 7.98 (dd, $J = 8.0, 1.5$ Hz, 1H), 7.64-7.46 (m, 11H), 7.36 (d, $J = 7.5$ Hz, 1H), 7.31-7.26 (m, 1H), 7.18 (dd, $J = 8.8, 2.8$ Hz, 1H), 7.10 (s, 1H), 6.73 (t, $J = 55.8$ Hz, 1H); MS: 654.9 ($\text{M}-1$) $^-$.











(continued)

#	starting material	structure	analytical data
11/ 49	 1/135		$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ : 9.52 (s, 1H), 8.52 (d, $J = 5.2$ Hz, 1H), 7.94-7.88 (m, 3H), 7.71-7.32 (m, 12H), 7.14-6.86 (m, 2H); MS: 638.0 (M-1) $^-$.
11/ 50	 2/25		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.36 (dd, $J = 9.5, 4.0$ Hz, 1H), 8.25 (d, $J = 8.0$ Hz, 2H), 8.03 (d, $J = 8.5$ Hz, 2H), 7.79 (t, $J = 7.8$ Hz, 2H), 7.68 (d, $J = 8.0$ Hz, 2H), 7.60-7.56 (m, 4H), 7.50-7.40 (m, 3H), 7.26 (dd, $J = 2.5, 8.5$ Hz, 1H), 7.19 (d, $J = 7.5$ Hz, 1H), 7.04 (t, $J = 55.3$ Hz, 1H); MS: 646.0 (M-1) $^-$.
11/ 51	 2/26		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.45 (dd, $J = 9.5, 4.0$ Hz, 1H), 8.06 (br s, 2H), 7.89 (dd, $J = 8.0, 1.5$ Hz, 1H), 7.78-7.63 (m, 3H), 7.56-7.29 (m, 9H), 6.95 (dd, $J = 8.5, 2.5$ Hz, 1H), 6.69 (t, $J = 55.8$ Hz, 1H); MS: 664.0 (M-1) $^-$.
11/ 52	 2/27		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.45 (dd, $J = 9.0, 4.5$ Hz, 1H), 8.07 (br s, 2H), 7.77-7.64 (m, 3H), 7.57-7.48 (m, 5H), 7.41-7.29 (m, 3H), 7.21-7.18 (m, 1H), 6.96 (dd, $J = 8.5, 2.5$ Hz, 1H), 6.71 (t, $J = 55.8$ Hz, 1H); MS: 682.0 (M-1) $^-$.
11/ 53	 2/28		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.44 (dd, $J = 9.0, 4.0$ Hz, 1H), 8.07-8.03 (m, 2H), 7.72 (t, $J = 8.3$ Hz, 1H), 7.66-7.50 (m, 9H), 7.33 (td, $J = 9.0, 2.5$ Hz, 1H), 7.00 (s, 1H), 6.95 (dd, $J = 8.0, 2.5$ Hz, 1H), 6.68 (t, $J = 55.5$ Hz, 1H); MS: 682.0 (M-1) $^-$.

(continued)

#	starting material	structure	analytical data
11/ 54	 2/29		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.44 (dd, $J = 9.3, 4.8$ Hz, 1H), 8.12-8.02 (m, 2H), 7.77-7.72 (m, 1H), 7.62-7.51 (m, 7H), 7.35 (td, $J = 9.3, 2.8$ Hz, 1H), 7.06 (s, 1H), 6.97 (dd, $J = 8.0, 2.5$ Hz, 1H), 6.71 (t, $J = 55.5$ Hz, 1H); MS: 737.1 ($\text{M}+18$) $^+$.
11/ 55	 1/136		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.42 (dd, $J = 9.5, 4.0$ Hz, 1H), 8.09-8.04 (m, 3H), 8.00 (dd, $J = 1.5, 8.0$ Hz, 1H), 7.73 (t, $J = 7.8$ Hz, 1H), 7.49-7.44 (m, 3H), 7.38-7.29 (m, 6H), 7.05 (d, $J = 1.0$ Hz, 1H), 6.95 (dd, $J = 8.5, 2.5$ Hz, 1H); MS: 663.9/666.0 ($\text{M}-1$) $^-$.
11/ 56	 1/137		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.43 (d, $J = 8.5$ Hz, 1H), 8.09-8.04 (m, 3H), 7.98 (dd, $J = 7.5, 1.5$ Hz, 1H), 7.72 (t, $J = 8.0$ Hz, 1H), 7.58-7.47 (m, 8H), 7.42 (t, $J = 7.8$ Hz, 1H), 7.34 (d, $J = 8.0$ Hz, 1H), 7.19 (d, $J = 7.5$ Hz, 1H), 7.07 (s, 1H), 6.66 (t, $J = 55.5$ Hz, 1H); MS: 661.9 ($\text{M}-1$) $^-$.
11/ 57	 35/2		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.60 (d, $J = 2.0$ Hz, 1H), 8.44 (dd, $J = 4.4, 9.2$ Hz, 1H), 8.10 (s, 1H), 8.03-7.31 (m, 9H), 7.21 (s, 1H), 7.09 (dd, $J = 2.8, 8.4$ Hz, 1H), 7.00 (d, $J = 4.8$ Hz, 1H), 6.65 (t, $J = 54.6$ Hz, 1H); MS: 663.8 ($\text{M}+1$) $^+$.
11/ 58	 1/138		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.45 (dd, $J = 9.3, 4.3$ Hz, 1H), 8.07-7.98 (m, 2H), 7.69 (t, $J = 8.0$ Hz, 1H), 7.58 (d, $J = 8.5$ Hz, 2H), 7.43 (d, $J = 8.0$ Hz, 2H), 7.37-7.31 (m, 2H), 7.24 (t, $J = 7.8$ Hz, 1H), 7.11 (d, $J = 8.0$ Hz, 1H), 6.97-6.93 (m, 2H), 6.79 (t, $J = 55.8$ Hz, 1H), 1.90-1.87 (m, 6H), 1.70-1.67 (m, 6H); MS: 678.0 ($\text{M}-1$) $^-$.

(continued)

#	starting material	structure	analytical data
11/59	 2/30		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.46 (dd, J = 9.2, 4.3 Hz, 1H), 8.05 (d, J = 7.9 Hz, 2H), 7.75-7.62 (m, 4H), 7.52 (q, J = 8.6 Hz, 4H), 7.47-7.28 (m, 5H), 7.21 (d, J = 7.8 Hz, 1H), 6.96 (dd, J = 8.3, 2.5 Hz, 1H), 6.67 (t, J = 55.6 Hz, 1H), 1.79 (s, 3H); MS: 690.0 (M-1) ⁻ .
11/60	 2/31		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.44 (dd, J = 9.5, 4.3 Hz, 1H), 8.06 (br s, 2H), 7.75-7.70 (m, 2H), 7.62-7.59 (m, 1H), 7.52-7.48 (m, 4H), 7.45-7.43 (m, 2H), 7.35-7.31 (m, 1H), 7.21 (d, J = 7.5 Hz, 1H), 6.98-6.93 (m, 2H), 6.64 (t, J = 55.6 Hz, 1H), 1.77 (s, 3H); MS: 724.0 (M-1) ⁻ .
11/61	 43/1		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.45-8.42 (m, 1H), 8.02 (br s, 2H), 7.68 (t, J = 7.9 Hz, 1H), 7.57 (d, J = 8.3 Hz, 2H), 7.45 (d, J = 8.3 Hz, 2H), 7.37-7.13 (m, 4H), 6.98-6.60 (m, 3H), 2.65 (s, 1H), 2.50-2.45 (m, 1H), 2.17-2.14 (m, 2H), 1.71-1.11 (m, 6H); MS: 652.0 (M-1) ⁻ .
11/62	 43/2		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.44 (dd, J = 9.2, 4.3 Hz, 1H), 8.04-8.00 (m, 2H), 7.69 (t, J = 7.9 Hz, 1H), 7.58 (d, J = 8.3 Hz, 2H), 7.47 (d, J = 8.3 Hz, 2H), 7.32 (td, J = 9.1, 2.5 Hz, 1H), 7.29-7.16 (m, 2H), 7.09-7.07 (m, 1H), 7.01-6.61 (m, 3H), 2.46-2.41 (m, 1H), 2.31-2.25 (m, 1H), 2.08 (d, J = 11.8 Hz, 2H), 1.79 (br s, 2H), 1.55 (qd, J = 13.0, 3.3 Hz, 2H), 1.32 (br s, 2H); MS: 652.0 (M-1) ⁻ .
11/63	 1/142		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.28 (dd, J = 4.3, 9.3 Hz, 1H), 8.12 (d, J = 7.5 Hz, 2H), 9.09 (d, J = 2.0 Hz, 1H), 7.99 (dd, J = 1.5, 8.0 Hz, 1H), 7.77 (t, J = 7.8 Hz, 1H), 7.53-7.33 (m, 6H), 7.04 (dd, J = 2.5, 9.0 Hz, 1H), 3.48-3.43 (m, 1H), 2.04 (br s, 2H), 1.80-1.67 (m, 6H); MS: 672.0 (M-1) ⁻ .



(continued)

#



starting material

structure



analytical data

11/
64



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.29 (d, $J = 7.5$ Hz, 2H), 8.16 (dd, $J = 4.3, 9.3$ Hz, 1H), 7.84-7.81 (m, 2H), 7.71 (dd, $J = 11.3, 1.3$ Hz, 1H), 7.62 (d, $J = 7.0$ Hz, 1H), 7.51-7.43 (m, 4H), 7.38 (d, $J = 8.0$ Hz, 1H), 7.30 (dd, $J = 2.5, 8.5$ Hz, 1H), 3.56-3.51 (m, 1H), 2.30-2.22 (m, 1H), 1.79-1.67 (m, 4H), 1.44 (q, $J = 11.8$ Hz, 2H), 1.22 (q, $J = 11.8$ Hz, 2H); MS: 688.0 (M-1) $^-$.

11/
65



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.44 (dd, $J = 4.3, 9.3$ Hz, 1H), 8.07 (d, $J = 7.0$ Hz, 2H), 8.01 (d, $J = 6.5$ Hz, 1H), 7.73 (t, $J = 8.0$ Hz, 1H), 7.56-7.32 (m, 8H), 7.19 (d, $J = 11.0$ Hz, 1H), 7.11 (s, 1H), 6.96 (dd, $J = 2.5, 8.0$ Hz, 1H), 6.70 (t, $J = 55.8$ Hz, 1H); MS: 697.9 (M-1) $^-$.

11/
66

$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.29 (dd, $J = 9.0, 4.0$ Hz, 1H), 8.13-8.08 (m, 3H), 7.99 (dd, $J = 8.5, 1.5$ Hz, 1H), 7.78-7.74 (m, 1H), 7.57-7.29 (m, 6H), 7.03-6.99 (m, 1H), 3.18 (dd, $J = 8.5, 6.0$ Hz, 1H), 2.38-2.25 (m, 2H), 1.85-1.05 (m, 8H); MS: 648.0 (M-1) $^-$.



11/
67


$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 9.15 (br s, 2H), 8.48 (dd, $J = 9.3, 4.3$ Hz, 1H), 8.08 (d, $J = 1.0$ Hz, 1H), 8.00 (dd, $J = 8.0, 1.5$ Hz, 1H), 7.55-7.48 (m, 6H), 7.44-7.38 (m, 3H), 7.13 (dd, $J = 8.0, 2.5$ Hz, 1H), 7.01 (s, 1H), 6.68 (t, $J = 55.8$ Hz, 1H); MS: 681.0 (M-1) $^-$.

11/
68


$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.37 (dd, $J = 9.2, 4.4$ Hz, 1H), 8.10 (s, 1H), 8.00 (d, $J = 7.6$ Hz, 1H), 7.55-7.41 (m, 8H), 7.27-7.22 (m, 1H), 7.16 (s, 1H), 7.10 (dd, $J = 8.8, 2.4$ Hz, 1H), 6.99 (d, $J = 2.8$ Hz, 1H), 6.70 (t, $J = 55.4$ Hz, 1H), 6.03 (d, $J = 2.8$ Hz, 1H), 3.72 (s, 3H); MS: 658.0 (M-1) $^-$.

(continued)

#	starting material	structure	analytical data
11/ 69			¹ H-NMR (400 MHz, CD ₃ OD) δ: 8.38 (dd, J = 9.2, 4.4 Hz, 1H), 7.58 (d, J = 8.4 Hz, 2H), 7.45 (d, J = 8.4 Hz, 2H), 7.39-7.16 (m, 7H), 6.94-6.65 (m, 3H), 2.38-2.34 (m, 1H), 1.61-1.55 (m, 1H), 1.50-1.45 (m, 1H), 1.10 (br s, 1H); MS: 628.0 and 630.0 (M-1) ⁺ .

Example 12**[0322]**

Step 1: Methyl 2-chloro-3'-(3-(2-cyanothiophen-3-yl)-5-fluoro-1-tosyl-1*H*-indol-2-yl)-5-methyl-[1,1'-biphenyl]-4-carboxylate (**12a**)

[0323]

[0324] To a solution of compound **2a** (200 mg, 0.33 mmol) in dioxane (2 mL) and water (0.4 mL) was added methyl 4-bromo-5-chloro-2-methylbenzoate (105 mg, 0.40 mmol), Cs₂CO₃ (215 mg, 0.66 mmol) and Pd(dppf)Cl₂ (20 mg). The mixture was stirred under N₂ at 100°C for 8 h, cooled to rt, poured into EA (40 mL) and washed with H₂O (40 mL) and brine (40 mL). The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC (EA:PE = 1:8) to give compound **12a** as a white solid.











Step 2: 2-Chloro-3'-(3-(2-cyanothiophen-3-yl)-5-fluoro-1-tosyl-1*H*-indol-2-yl)-5-methyl-[1,1'-biphenyl]-4-carboxylic acid (**12**)

[0325] To a solution of compound **12a** (150 mg, 0.23 mmol) in HCl/dioxane (4N, 5 mL) was added H₂O (0.5 mL) and the mixture was stirred at 90°C overnight, concentrated, diluted with water and extracted with EA (3 x). The combined organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give

compound 12 as a white solid. $^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.29 (dd, $J = 9.0, 4.5$ Hz, 1H), 8.03 (d, $J = 5.0$ Hz, 1H), 7.82 (s, 1H), 7.54-7.46 (m, 3H), 7.40-7.32 (m, 3H), 7.24 (d, $J = 8.0$ Hz, 2H), 7.18 (dd, $J = 8.0, 3.0$ Hz, 2H), 7.04-7.00 (m, 2H), 2.53 (s, 3H), 2.21 (s, 3H); MS: 638.9 (M-1) $^-$.

5 **Example 12/1 to 12/10**

[0326] The following Examples were prepared similar as described for Example 12 using the appropriate building block.

#	building block	structure	analytical data
12/1			$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.29 (dd, $J = 9.3, 4.3$ Hz, 1H), 8.03 (d, $J = 5.0$ Hz, 1H), 7.68 (d, $J = 8.0$ Hz, 1H), 7.53-7.15 (m, 10H), 7.03 (d, $J = 5.0$ Hz, 1H), 6.98 (s, 1H), 2.55 (s, 3H), 2.23 (s, 3H); MS: 639.0 (M-1) $^-$.
12/2			$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.42 (dd, $J = 9.0, 4.5$ Hz, 1H), 7.92 (d, $J = 7.5$ Hz, 1H), 7.84 (d, $J = 5.0$ Hz, 1H), 7.53-7.46 (m, 3H), 7.30-7.26 (m, 3H), 7.19-6.97 (m, 6H), 2.28 (s, 3H); MS: 643.0 (M-1) $^-$.
12/3			$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.42 (dd, $J = 9.3, 4.3$ Hz, 1H), 7.85-7.80 (m, 2H), 7.53-7.47 (m, 3H), 7.30-7.25 (m, 3H), 7.17 (d, $J = 8.0$ Hz, 3H), 7.07 (dd, $J = 7.8, 2.8$ Hz, 1H), 7.04-6.96 (m, 2H), 2.27 (s, 3H); MS: 642.9 (M-1) $^-$.
12/4			$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.29 (dd, $J = 9.3, 4.3$ Hz, 1H), 8.03 (d, $J = 5.0$ Hz, 1H), 7.59-7.50 (m, 3H), 7.43 (d, $J = 7.5$ Hz, 1H), 7.40-7.34 (m, 3H), 7.24 (d, $J = 8.5$ Hz, 2H), 7.17 (dd, $J = 8.5, 2.5$ Hz, 1H), 7.10 (s, 1H), 7.06 (d, $J = 5.5$ Hz, 1H), 6.94 (s, 1H), 3.82 (s, 3H), 2.22 (s, 3H); MS: 655.0 (M-1) $^-$.
12/5			$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.42 (dd, $J = 9.3, 4.3$ Hz, 1H), 7.84 (d, $J = 5.0$ Hz, 1H), 7.69 (d, $J = 7.5$ Hz, 1H), 7.51-7.45 (m, 3H), 7.29-7.24 (m, 3H), 7.16 (d, $J = 8.0$ Hz, 2H), 7.09-7.05 (m, 2H), 6.97-6.95 (m, 2H), 3.95 (s, 3H), 2.27 (s, 3H); MS: 655.0 (M-1) $^-$.



(continued)

building block

structure



analytical data

12/6





$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.30 (dd, $J = 9.2$, 4.4 Hz, 1H), 7.98 (d, $J = 5.1$ Hz, 1H), 7.88 (d, $J = 7.8$ Hz, 1H), 7.61 (s, 1H), 7.48-7.33 (m, 4H), 7.29 (d, $J = 8.3$ Hz, 2H), 7.25-7.13 (m, 2H), 6.99 (d, $J = 5.0$ Hz, 1H), 6.93 (br s, 1H), 2.28 (s, 3H); MS: 581.0 ($\text{M}-1$) $^-$.

12/7





$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.30 (dd, $J = 9.2$, 4.4 Hz, 1H), 7.98 (d, $J = 5.1$ Hz, 1H), 7.61 (br s, 1H), 7.52 (br s, 1H), 7.48-7.28 (m, 7H), 7.20-7.17 (m, 1H), 7.06 (d, $J = 4.5$ Hz, 1H), 6.64 (br s, 1H), 3.76 (br s, 3H), 2.28 (s, 3H); MS: 597.2 ($\text{M}+\text{H}$) $^+$.

12/8




$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.29 (dd, $J = 9.2$, 4.4 Hz, 1H), 8.07 (d, $J = 5.1$ Hz, 1H), 7.59-7.49 (m, 2H), 7.45-7.31 (m, 4H), 7.28 (d, $J = 8.3$ Hz, 2H), 7.20 (dd, $J = 8.6$, 2.6 Hz, 1H), 7.10-7.00 (m, 2H), 6.61 (s, 1H), 4.25-4.22 (m, 1H), 2.26 (s, 3H), 1.30 (d, $J = 6.5$ Hz, 6H); MS: 623.0 ($\text{M}-1$) $^-$.


12/9



$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.31 (dd, $J = 9.2$, 4.4 Hz, 1H), 8.17 (d, $J = 8.2$ Hz, 1H), 8.04 (d, $J = 5.1$ Hz, 1H), 7.80 (d, $J = 7.6$ Hz, 1H), 7.68 (s, 1H), 7.55-7.44 (m, 2H), 7.40-7.36 (m, 4H), 7.28-7.24 (m, 3H), 7.19 (dd, $J = 8.5$, 2.5 Hz, 1H), 7.09 (d, $J = 5.1$ Hz, 1H), 2.22 (s, 3H); MS: 631.0 ($\text{M}-1$) $^-$.


12/
10

P33



$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.44 (dd, $J = 9.5$, 4.5 Hz, 1H), 8.06 (br s, 2H), 7.89 (d, $J = 1.5$ Hz, 1H), 7.80 (dd, $J = 1.3$, 7.8 Hz, 1H), 7.72 (t, $J = 8.0$ Hz, 1H), 7.55-7.48 (m, 7H), 7.36-7.31 (m, 2H), 7.02 (s, 1H), 6.95 (dd, $J = 2.3$, 8.3 Hz, 1H), 6.68 (t, $J = 55.8$ Hz, 1H); MS: 715.9 ($\text{M}-1$) $^-$.

Example 13**[0327]**



3-(1-((4-Chlorophenyl)sulfonyl)-5-hydroxy-2-(thiophen-2-yl)-1H-indol-3-yl)thiophene-2-carbonitrile (**13**)

[0328] To a solution of compound **139** (775 mg, 1.52 mmol) in DCM (10 mL) was added BBr₃ (2 mL, 3N in DCM) at 0°C and the mixture was stirred for 2 h, poured into water (50 mL) and extracted with EA (3 × 20 mL). The combined organic layer was washed with aq. K₂CO₃ (30 mL) and brine (2 × 30 mL), dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound **13** as a yellow solid. ¹H-NMR (500 MHz, DMSO-d₆) δ: 9.63 (s, 1H), 8.05-8.03 (m, 2H), 7.73 (d, J = 5.0 Hz, 1H), 7.59 (d, J = 8.5 Hz, 2H), 7.47 (d, J = 8.5 Hz, 2H), 7.19 (d, J = 3.0 Hz, 1H), 7.12-7.10 (m, 1H), 7.01-6.96 (m, 2H), 6.58 (s, 1H); MS: 496.9 (M+1)⁺.


Example 13/1

[0329] The following Example was prepared similar as described for Example 13 using the appropriate starting material.

#	starting material	structure	analytical data
13/1			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 9.99 (s, 1H), 8.01 (d, J = 5.0 Hz, 1H), 7.70-7.68 (m, 2H), 7.63 (d, J = 9.0 Hz, 2H), 7.50 (d, J = 8.5 Hz, 2H), 7.16-7.15 (m, 2H), 7.09-7.08 (m, 1H), 7.00 (d, J = 5.0 Hz, 1H), 6.87 (dd, J = 2.0, 8.5 Hz, 1H); MS: 496.7 (M+1) ⁺ .

Example 14

[0330]




Methyl 2-((1-((4-chlorophenyl)sulfonyl)-3-(2-cyanothiophen-3-yl)-2-(thiophen-2-yl)-1H-indol-5-yl)oxy)acetate (**14**)

[0331] To a solution of compound **13** (120 mg, 0.25 mmol) in DMF (5 mL) was added K₂CO₃ (69 mg, 0.50 mmol) and methyl 2-bromoacetate (46 mg, 0.30 mmol). The mixture was stirred at rt overnight, diluted with water and extracted with EA (3 × 20 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄, filtered, concentrated and purified by FCC to give compound **14** as a brown solid.

Example 15

[0332]



2-((1-((4-Chlorophenyl)sulfonyl)-3-(2-cyanothiophen-3-yl)-2-(thiophen-2-yl)-1H-indol-5-yl)oxy)acetic acid (**15**)





[0333] To a solution of compound **14** (74 mg, 0.13 mmol) in MeOH (3 mL) was added LiOH (1 mL, 2N) and the mixture was stirred at rt overnight, evaporated, adjusted to pH <2 with 2N HCl and extracted with EA (3 × 20 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound **15** as a white solid. ¹H-NMR (500 MHz, DMSO-d₆) δ: 13.02 (br s, 1H), 8.16 (d, J = 9.0 Hz, 1H), 8.05 (d, J = 5.0 Hz, 1H), 7.74 (d, J = 5.0 Hz, 1H), 7.60 (d, J = 8.5 Hz, 2H), 7.49 (d, J = 8.5 Hz, 2H), 7.21-7.11 (m, 3H), 7.03 (d, J = 5.0 Hz, 1H), 6.77 (d, J = 2.5 Hz, 1H), 4.67 (s, 2H); MS: 554.6 (M+1)⁺.


Example 15/1 to 15/6

[0334] The following Examples were prepared similar as described for Example 14 (optional) and Example 15 using the appropriate building blocks.

#	building block(s)	structure	analytical data
15/1			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 12.12 (s, 1H), 8.14 (d, J = 9.0 Hz, 1H), 8.03 (d, J = 5.0 Hz, 1H), 7.73 (d, J = 4.5 Hz, 1H), 7.59 (d, J = 9.0 Hz, 2H), 7.48 (d, J = 9.0 Hz, 2H), 7.20 (d, J = 2.5 Hz, 1H), 7.15-7.11 (m, 2H), 7.02 (d, J = 5.0 Hz, 1H), 6.77 (d, J = 3.0 Hz, 1H), 3.97 (t, J = 6.5 Hz, 2H), 2.37 (t, J = 7.3 Hz, 2H), 1.93 (t, J = 6.8 Hz, 2H); MS: 582.7 (M+1) ⁺ .
15/2			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 12.03 (brs, 1H), 8.14 (d, J = 9.5 Hz, 1H), 8.04 (d, J = 5.0 Hz, 1H), 7.74 (d, J = 5.0 Hz, 1H), 7.59 (d, J = 8.0 Hz, 2H), 7.48 (d, J = 9.0 Hz, 2H), 7.21-7.20 (m, 1H), 7.15-7.11 (m, 2H), 7.02 (d, J = 4.5 Hz, 1H), 6.77 (d, J = 2.5 Hz, 1H), 3.96 (t, J = 6.0 Hz, 2H), 2.27 (t, J = 7.3 Hz, 2H), 1.75-1.62 (m, 4H); MS: 596.9 (M+1) ⁺ .
15/3			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 13.20 (s, 1H), 8.03 (d, J = 5.0 Hz, 1H), 7.72 (d, J = 5.0 Hz, 2H), 7.60 (d, J = 8.5 Hz, 2H), 7.54 (d, J = 8.5 Hz, 2H), 7.26 (d, J = 9.0 Hz, 1H), 7.19 (d, J = 3.5 Hz, 1H), 7.12-7.10 (m, 1H), 7.05 (dd, J = 2.0, 9.0 Hz, 1H), 7.01 (d, J = 5.0 Hz, 1H), 4.86 (s, 2H); MS: 554.6 (M+1) ⁺ .
15/4			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.02 (d, J = 5.0 Hz, 1H), 7.74 (d, J = 2.0 Hz, 1H), 7.71 (d, J = 5.5 Hz, 1H), 7.61 (d, J = 8.5 Hz, 2H), 7.54 (d, J = 8.0 Hz, 2H), 7.25 (d, J = 9.0 Hz, 1H), 7.18 (d, J = 3.0 Hz, 1H), 7.11-7.00 (m, 3H), 4.13 (t, J = 6.5 Hz, 2H), 2.38 (t, J = 7.3 Hz, 2H), 2.03-1.97 (m, 2H); MS: 583.0 (M+1) ⁺ .


(continued)

#	building block(s)	structure	analytical data
15/5			$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.03 (d, J = 5.0 Hz, 1H), 7.74 (d, J = 1.5 Hz, 1H), 7.71 (d, J = 5.5 Hz, 1H), 7.61 (d, J = 8.5 Hz, 2H), 7.54 (d, J = 8.5 Hz, 2H), 7.25 (d, J = 9.0 Hz, 1H), 7.18 (d, J = 4.0 Hz, 1H), 7.10 (t, J = 4.3 Hz, 1H), 7.05-7.03 (m, 1H), 7.01 (d, J = 5.0 Hz, 1H), 4.12 (t, J = 6.0 Hz, 2H), 2.31 (t, J = 7.3 Hz, 2H) 1.83-1.78 (m, 2H), 1.73-1.68 (m, 2H); MS: 597.0 ($\text{M}+1$) $^+$.
15/6			$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.20 (brs, 1H), 8.26 (d, J = 8.5 Hz, 1H), 7.98 (d, J = 5.0 Hz, 1H), 7.50-7.47 (m, 1H), 7.40-7.32 (m, 4H), 7.22 (d, J = 8.5 Hz, 2H), 7.16 (d, J = 8.5 Hz, 2H), 6.99 (d, J = 9.0 Hz, 2H), 6.93 (d, J = 5.0 Hz, 1H), 3.78 (s, 3H), 2.87 (t, J = 7.5 Hz, 2H), 2.59 (t, J = 7.5 Hz, 2H); MS: 542.9 ($\text{M}+1$) $^+$.

Example 16**[0335]**

tert-Butyl 3-(1-((4-chlorophenyl)sulfonyl)-3-(2-cyanothiophen-3-yl)-5-hydroxy-2-(thiophen-2-yl)-1*H*-indol-6-yl)propanoate (16)

[0336] To a solution of compound 13 (120 mg, 0.25 mmol) in DMF (3 mL) was added K_2CO_3 (69 mg, 0.50 mmol) and methyl 2-bromoacetate (38 mg, 0.30 mmol). The mixture was stirred at rt overnight, diluted with water and extracted with EA (3×20 mL). The combined organic layer was washed with brine (20 mL), dried over Na_2SO_4 , filtered, concentrated and purified by FCC to give compound **16** as a brown solid; MS: 624.7 ($\text{M}+1$) $^+$.

Example 17**[0337]**


3-(1-((4-Chlorophenyl)sulfonyl)-3-(2-cyanothiophen-3-yl)-5-hydroxy-2-(thiophen-2-yl)-1*H*-indol-6-yl)propanoic acid (17)

[0338] To a solution of compound **16** (32 mg, 50 μmol) in MeOH (2 mL) was added NaOH (0.5 mL, 2N) and the mixture

was stirred at rt overnight, concentrated, adjusted to pH <2 with 2N HCl and extracted with EA (3 × 20 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound **17** as a white solid. ¹H-NMR (500 MHz, DMSO-d₆) δ: 12.46 (s, 1H), 10.42 (s, 1H), 7.95 (d, J = 9.0 Hz, 1H), 7.77 (d, J = 5.0 Hz, 1H), 7.70 (d, J = 5.0 Hz, 1H), 7.64 (d, J = 8.5 Hz, 2H), 7.57 (d, J = 9.0 Hz, 2H), 7.18 (d, J = 3.0 Hz, 1H), 7.13-7.11 (m, 1H), 7.06 (d, J = 5.0 Hz, 1H), 6.89 (d, J = 9.0 Hz, 1H), 4.43-4.37 (m, 2H), 2.61 (t, J = 7.8 Hz, 2H); MS: 568.7 (M+1)⁺.


Example 18

[0339]



Step 1: Methyl 3-amino-4-(thiophen-2-ylethynyl)benzoate (**18a**)


[0340]



[0341] Methyl 3-amino-4-iodo-benzoate (1.0 g, 3.6 mmol) was dissolved in degassed THF (10 mL) and the mixture was degassed by bubbling a gentle stream of N₂ through the solution. After ~5 min DIPEA (5.0 mL, 29 mmol) was added and the bubbling was continued for a few minutes before addition of 2-ethynylthiophene (0.41 mL, 4.3 mmol). Then CuI (28 mg, 0.15 mmol) was added directly followed by PdCl₂(PPh₃)₂ (49 mg, 70 μmol). The mixture was stirred at rt for 2.5 h, filtered through Celite and washed with THF. EA (200 mL) was added and the mixture was washed with water (50 mL) and brine (50 mL). The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC to give compound **18a**. ¹H-NMR (400 MHz, CDCl₃): 3.90 (s, 3H), 4.36 (br s, 2H), 7.02-7.04 (m, 1H), 7.30-7.34 (m, 2H), 7.35-7.41 (m, 3H).

Step 2: Methyl 4-(thiophen-2-ylethynyl)-3-(2,2,2-trifluoroacetamido)benzoate (**18b**)


[0342]



[0343] Compound **18a** (450 mg, 1.68 mmol) was dissolved in dry THF (1 mL) and the mixture was cooled to 0°C. Then 2,2,2-trifluoroacetic anhydride (0.47 mL, 3.4 mmol) was added dropwise and the mixture was stirred for 15 min, diluted with EA (100 mL) and washed with NaHCO₃-water (1:1, 50 mL) and brine (50 mL). The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC to give the compound **18b** as a yellow solid. ¹H-NMR (400 MHz, CDCl₃): 3.95 (s, 3H), 7.08-7.10 (m, 1H), 7.37-7.38 (m, 1H), 7.43-7.44 (m, 1H), 7.60-7.63 (m, 1H), 7.90-7.92 (m, 1H), 8.75 (br s, 1H), 8.97 (d, 1H).

Step 3: Methyl 3-(2-cyanothiophen-3-yl)-2-(thiophen-2-yl)-1H-indole-6-carboxylate (**18c**)

[0344]




[0345] A dried microwave-tube was charged with compound **18b** (187 mg, 0.53 mmol), Cs_2CO_3 (259 mg, 0.79 mmol), $\text{Pd}(\text{PPh}_3)_4$ (31 mg, 0.03 mmol) and 3-bromothiophene-2-carbonitrile (149 mg, 0.79 mmol). Degassed CH_3CN (2.5 mL) was added and the tube was purged with N_2 . The mixture was stirred at 100°C for 1 h, cooled, diluted with EA (80 mL) and washed with NaHCO_3 :water (1:1, 40 mL), water (20 mL) and brine (15 mL). The organic layer was dried (Na_2SO_4), filtered and concentrated to give compound **18c** as a yellow solid. $^1\text{H-NMR}$ (400 MHz, DMSO-d_6): 3.88 (s, 3H), 7.17-7.19 (m, 1H), 7.33-7.46 (m, 3H), 7.64-7.71 (m, 2H), 8.10 (s, 1H), 8.19-8.21 (m, 1H), 12.32 (s, 1H).

Step 4: Methyl 1-((4-chlorophenyl)sulfonyl)-3-(2-cyanothiophen-3-yl)-2-(thiophen-2-yl)-1H-indole-6-carboxylate (18**)**

[0346] Compound **18c** (180 mg, 0.49 mmol) was dissolved in THF (15 mL) and NaH (3.9 mmol dispersed in mineral oil) was added followed by 4-chlorobenzenesulfonyl chloride (188 mg, 0.89 mmol). The mixture was stirred at rt for 1 h, quenched by careful addition of water, diluted with EA (250 mL) and washed with water (100 mL) and brine (100 mL). The organic layer was dried over Na_2SO_4 , filtered, concentrated and purified by FCC and then prep-HPLC (55-60% acetonitrile in 15 mM NH_4HCO_3 buffer, pH 10) to give compound **18** as a white solid.

Example 19

[0347]




1-((4-Chlorophenyl)sulfonyl)-3-(2-cyanothiophen-3-yl)-2-(thiophen-2-yl)-1H-indole-6-carboxylic acid (19**)**

[0348] Compound **18** (77 mg, 0.14 mmol) was dissolved in THF (4 mL) and cooled to 0°C . In a separate 4 mL vial LiOH (48 mg, 2.0 mmol) was dissolved in water (4 mL) and cooled to 0°C . The base solution was added dropwise to the solution of compound **18** and the resulting mixture was stirred vigorously overnight (the reaction slowly adapted rt). The reaction was re-cooled to 0°C , quenched with 2N HCl (1.4 mL) and extracted with EA (3×20 mL). The combined organic layer was dried over MgSO_4 , filtered, concentrated and purified by prep-HPLC (Xbridge, 30-60% acetonitrile in 0.1% TFA buffer) to give compound **19** as a white solid. $^1\text{H-NMR}$ (400 MHz, DMSO-d_6): 7.07 (d, 1H), 7.12-7.15 (dd, 1H), 7.24-7.25 (dd, 1H), 7.40-7.50 (m, 3H), 7.55-7.63 (m, 2H), 7.76-7.77 (dd, 1H), 7.97-8.00 (dd, 1H), 8.06 (d, 1H), 8.89 (m, 1H), 13.29 (br s, 1H); MS: 542 ($\text{M}+\text{NH}_3+1$) $^+$.


Example 20

[0349]



Step 1: 2-(Thiophen-2-ylethynyl)aniline (**20a**)


[0350]



[0351] To a mixture of 2-iodoaniline (40.0 g, 183 mmol), CuI (700 mg, 3.70 mmol), Pd(PPh₂)₂Cl₂ (1.30 g, 1.83 mmol) and TEA (120 mL) in ACN (1 L) was added 2-ethynylthiophene (24.0 g, 219 mmol) under N₂ via a syringe. The mixture was stirred at 50°C overnight, cooled, filtered, concentrated and purified by FCC (PE:EA = 50:1) to afford compound **20a** as a yellow solid.

Step 2: 4-Chloro-*N*-(2-(thiophen-2-ylethynyl)phenyl)benzenesulfonamide (**20b**)


[0352]



[0353] To a solution of compound **20a** (10.0 g, 50.3 mmol), 4-chlorobenzenesulfonyl chloride (13.1 g, 62.3 mmol) and pyridine (4.17 g, 52.8 mmol) in DCM (150 mL) was added DMAP (306 mg, 2.5 mmol) at rt. The mixture was heated to reflux overnight, cooled, washed with 2N HCl and extracted with DCM. The organic layer was dried over Na₂SO₄, filtered, concentrated and then the residue was washed with PE to give compound **20b** as a yellow solid.

Step 3: 1-((4-Chlorophenyl)sulfonyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-(thiophen-2-yl)-1*H*-indole (**20c**)

[0354]













[0355] A mixture of compound **20b** (11.0 g, 29.4 mmol), Cs₂CO₃ (19 g, 59 mmol), AsPh₃ (1.36 g, 4.40 mmol), Pd₂(dba)₃ (1.34 g, 1.48 mmol) and B₂Pin₂ (15 g, 59 mmol) in dioxane (175 mL) was stirred under N₂ at 60°C for 2 h, cooled, filtered, concentrated and purified by FCC (PE:EA = 20:1 to 5:1) to afford compound **20c** as a white solid.

Step 4: 1-((4-Chlorophenyl)sulfonyl)-3-(2-chlorothiophen-3-yl)-2-(thiophen-2-yl)-1*H*-indole (**20**)













[0356] A solution of compound **20c** (200 mg, 0.40 mmol), 3-bromo-2-chlorothiophene (79 mg, 0.40 mmol), Pd(PPh₃)₄ (46 mg, 40 μmol) and K₃PO₄ (404 mg, 1.6 mmol) in dioxane/H₂O (10:1, 22 mL) was stirred under N₂ at 100°C overnight, cooled, filtered and washed with DCM. Then the filtrate was concentrated and purified by prep-HPLC to afford compound **20** as a white solid. ¹H-NMR (CDCl₃, 300 MHz) δ: 8.35 (d, J = 6.6 Hz, 1H), 7.40-7.25 (m, 8H), 7.13 (d, J = 0.6 Hz, 1H), 7.05-7.03 (m, 2H), 6.61 (d, J = 4.2 Hz, 1H); MS: 589.0 (M+1)⁺.

Example 20/1 to 20/25













[0357] The following Examples were prepared similar as described for Example 20 using the appropriate building blocks.

#	building block(s)	structure	analytical data
20/1			¹ H-NMR (CDCl ₃ , 300 MHz) δ: 1.02 (s, 9H), 6.67 (s, 1H), 6.98-7.01 (m, 1H), 7.08 (s, 1H), 7.19 (s, 1H), 7.28-7.30 (m, 3H), 7.35-7.43 (m, 3H), 7.47-7.50 (m, 2H), 8.32 (s, 1H).
20/2			¹ H-NMR (CDCl ₃ , 300 MHz) δ: 7.04-7.01 (m, 1H), 7.42-7.19 (m, 8H), 7.48-7.57 (m, 2H), 8.41 (d, J = 8.4 Hz, 1H), 8.55 (s, 1H), 8.70 (d, J = 4.5 Hz, 1H); MS: 476.0 (M+1) ⁺ .
20/3			¹ H-NMR (CDCl ₃ , 300 MHz) δ: 6.29 (t, J = 54.9 Hz, 1H), 6.72-6.73 (m, 1H), 7.03 (dd, J = 3.8, 5.0 Hz, 1H), 7.10 (d, J = 2.7 Hz, 1H), 7.25-7.46 (m, 9H), 8.37 (d, J = 8.4 Hz, 1H); MS: 505.9 (M+1) ⁺ .
20/4			¹ H-NMR (CDCl ₃ , 300 MHz) δ: 7.04 (dd, J = 3.8, 5.3 Hz, 1H), 7.15 (d, J = 3.3 Hz, 1H), 7.20 (dd, J = 0.9, 3.6 Hz, 1H), 7.26-7.47 (m, 8H), 7.94 (d, J = 3.3 Hz, 1H), 8.36 (d, J = 8.4 Hz, 1H); MS: 498.0 (M+18) ⁺ .
20/5			¹ H-NMR (CDCl ₃ , 300 MHz) δ: 2.54 (s, 3H), 6.97 (dd, J = 3.8, 8.6 Hz, 1H), 7.02 (s, 1H), 7.07-7.10 (m, 1H), 7.23-7.57 (m, 10H), 8.14-8.17 (m, 1H), 8.32 (d, J = 8.4 Hz, 1H); MS: 545.0 (M+18) ⁺ .













(continued)

#	building block(s)	structure	analytical data
5 20/6			$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 6.99-7.02 (m, 1H), 7.16-7.42 (m, 11H), 7.47 (t, J = 8.4 Hz, 1H), 8.36 (d, J = 8.4 Hz, 1H); MS: 493.0 ($\text{M}+1$) $^+$.
10 20/7			$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 7.09 (dd, J = 3.9, 5.1 Hz, 1H), 7.26-7.28 (m, 2H), 7.36-7.41 (m, 6H), 7.48-7.52 (m, 1H), 8.43 (d, J = 8.4 Hz, 1H), 8.60 (d, J = 2.4 Hz, 1H), 8.79 (d, J = 2.4 Hz, 1H); MS: 477.0 ($\text{M}+1$) $^+$.
15 20/8			
20 20/9			
25 20/ 10			
30 20/11			$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 6.80 (d, J = 5.1 Hz, 1H), 6.98-7.08 (m, 3H), 7.20-7.22 (m, 1H), 7.33-7.53 (m, 7H), 8.39 (d, J = 8.4 Hz, 1H); MS: 482.0 ($\text{M}+18$) $^+$.






(continued)

#	building block(s)	structure	analytical data
5 20/12			$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 6.74-6.84 (m, 2H), 6.89 (d, J = 5.1 Hz, 1H), 7.00-7.03 (m, 1H), 7.20 (d, J = 2.7 Hz, 1H), 7.32-7.51 (m, 6H), 8.30 (d, J = 9.0 Hz, 1H); MS: 483.0 ($\text{M}+1$) $^+$.
10 20/13			$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 6.81 (d, J = 4.8 Hz, 1H), 7.06 (dd, J = 3.2, 5.0 Hz, 1H), 7.21 (d, J = 2.7 Hz, 1H), 7.29-7.49 (m, 9H), 8.38 (d, J = 8.7 Hz, 1H).
15 20/14			$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 3.79 (s, 3H), 6.77-6.79 (m, 3H), 7.05 (dd, J = 3.8, 5.0 Hz, 1H), 7.19 (dd, J = 0.9, 3.6 Hz, 1H), 7.31-7.49 (m, 7H), 8.41 (d, J = 8.4 Hz, 1H), MS: 477.0 ($\text{M}+1$) $^+$.
20 20/15			$^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ : 2.55-2.60 (m, 2H), 2.93 (t, J = 7.5 Hz, 2H), 3.63 (s, 3H), 6.79 (d, J = 5.1 Hz, 1H), 7.04 (dd, J = 3.9, 5.1 Hz, 1H), 7.15-7.18 (m, 3H), 7.33-7.50 (m, 7H), 8.40 (d, J = 8.4 Hz, 1H); MS: 550.0 ($\text{M}+18$) $^+$.
25 20/16			$^1\text{H-NMR}$ (DMSO-d_6 , 400 MHz) δ : 8.17 (d, J = 8.4 Hz, 1H), 7.48-7.33 (m, 10H), 6.98-6.94 (m, 2H), 3.75 (s, 3H), 2.29-2.08 (m, 3H), 1.77-1.36 (m, 5H); MS: 486.2 ($\text{M}+18$) $^+$.
30 20/17			$^1\text{H-NMR}$ (DMSO-d_6 , 400 MHz) δ : 8.61 (s, 1H), 8.30 (d, J = 8.4 Hz, 1H), 8.12 (d, J = 8.4 Hz, 1H), 8.04 (d, J = 8.0 Hz, 1H), 7.96-7.92 (m, 1H), 7.83-7.79 (m, 1H), 7.55-7.48 (m, 3H), 7.37-7.29 (m, 7H), 7.05-7.03 (m, 2H), 3.79 (s, 3H); MS: 516.1 ($\text{M}+1$) $^+$.

(continued)

#	building block(s)	structure	analytical data
20/18			$^1\text{H-NMR}$ (DMSO- <i>d</i> ₆ , 300 MHz) δ : 8.31 (d, <i>J</i> = 8.4 Hz, 1H), 8.16 (d, <i>J</i> = 8.4 Hz, 1H), 7.61-7.50 (m, 2H), 7.43-7.01 (m, 13H), 3.79 (s, 3H); MS: 538.1 (M+18) ⁺ .
20/19			$^1\text{H-NMR}$ (DMSO- <i>d</i> ₆ , 400 MHz) δ : 8.23 (d, <i>J</i> = 8.4 Hz, 1H), 7.51-7.21 (m, 14H), 7.00 (d, <i>J</i> = 8.8 Hz, 2H), 3.77 (s, 3H), 2.86-2.80 (m, 1H), 2.60-2.49 (m, 2H), 2.15-2.09 (m, 1H); MS: 534.1 (M+18) ⁺ .
20/20			$^1\text{H-NMR}$ (DMSO- <i>d</i> ₆ , 400 MHz) δ : 9.07 (dd, <i>J</i> = 1.2, 4.0 Hz, 1H), 8.44 (d, <i>J</i> = 8.0 Hz, 1H), 8.29 (d, <i>J</i> = 8.4 Hz, 1H), 8.24 (d, <i>J</i> = 8.8 Hz, 1H), 7.79 (dd, <i>J</i> = 4.2, 8.6 Hz, 1H), 7.60 (d, <i>J</i> = 8.8 Hz, 1H), 7.53-7.45 (m, 3H), 7.36-7.27 (m, 7H), 7.03 (d, <i>J</i> = 9.2 Hz, 2H), 3.79 (s, 3H); MS: 516.1 (M+1) ⁺ .
20/21			$^1\text{H-NMR}$ (DMSO- <i>d</i> ₆ , 400 MHz) δ : 8.21 (d, <i>J</i> = 8.4 Hz, 1H), 7.47-7.25 (m, 10H), 7.14 (d, <i>J</i> = 8.0 Hz, 1H), 7.01-8.97 (m, 2H), 6.65 (d, <i>J</i> = 4.8 Hz, 1H), 3.76 (s, 3H), 1.85 (s, 3H); MS: 460.1 (M+1) ⁺ .
20/22			$^1\text{H-NMR}$ (DMSO- <i>d</i> ₆ , 400 MHz) δ : 8.20 (d, <i>J</i> = 8.4 Hz, 1H), 7.44-7.29 (m, 10H), 7.02-6.98 (m, 2H), 8.75 (d, <i>J</i> = 6.0 Hz, 1H), 6.41 (d, <i>J</i> = 5.6 Hz, 1H), 3.77 (s, 3H), 3.54 (s, 3H); MS: 476.1 (M+1) ⁺ .
20/23			$^1\text{H-NMR}$ (DMSO- <i>d</i> ₆ , 400 MHz) δ : 8.26 (d, <i>J</i> = 8.0 Hz, 1H), 7.52-7.28 (m, 10H), 7.01 (d, <i>J</i> = 9.2 Hz, 2H), 6.77 (s, 1H), 3.77 (s, 3H), 2.43 (s, 3H); MS: 502.1 (M+18) ⁺ .

(continued)

#	building block(s)	structure	analytical data
20/ 24			
20/25			
Example 21			
[0358]			
			

3-(1-((4-Chlorophenyl)sulfonyl)-2-(thiophen-2-yl)-1H-indol-3-yl)thiophene-2-carboxylic acid (**21**)



[0359] A solution of compound **20/1** (95 mg, 0.24 mmol) and TFA (0.5 mL) in DCM (2.5 mL) was stirred at rt for 4 h, concentrated and then the residue was triturated with PE including a little amount of EA to give compound **21** as a white solid. ¹H-NMR (CDCl₃, 400 MHz) δ: 6.79 (d, J = 3.9 Hz, 1H), 7.02-7.04 (m, 1H), 7.11 (dd, J = 1.1, 2.6 Hz, 1H), 7.21 (d, J = 5.7 Hz, 1H), 7.26-7.40 (m, 6H), 7.45-7.49 (m, 1H), 7.55 (d, J = 3.9 Hz, 1H), 8.35 (d, J = 6.3 Hz, 1H); MS: 497.9 (M-1)⁻.

Example 21/1 to 21/3

[0360] The following Examples were prepared similar as described for Example 20 using the appropriate starting material.



#	starting material	structure	analytical data
---	-------------------	-----------	-----------------

21/1





¹H-NMR (CDCl₃, 400 MHz) δ: 8.34 (d, J = 6.3 Hz, 1H), 7.58-7.13 (m, 10H), 6.97-6.87 (m, 2H), 6.66 (s, 1H), 4.49 (br s, 2H); MS: 547.0 (M-1)⁻.

21/2



¹H-NMR (DMSO-d₆, 300 MHz) δ: 8.21 (d, J = 8.1 Hz, 1H), 7.65 (d, J = 4.8 Hz, 1H), 7.58-7.48 (m, 6H), 7.37-7.32 (m, 1H), 7.16-7.04 (m, 4H), 6.87 (d, J = 7.8 Hz, 1H), 4.83 (s, 2H); MS: 546.9 (M-1)⁻.


21/3



¹H-NMR (CD₃OD, 400 MHz) δ: 8.43 (dd, J = 9.4, 4.2 Hz, 1H), 8.05 (d, J = 8.0 Hz, 2H), 7.98 (d, J = 2.0 Hz, 1H), 7.86 (dd, J = 8.0, 2.0 Hz, 1H), 7.71 (t, J = 8.0 Hz, 1H), 7.54-7.30 (m, 9H), 7.09 (s, 1H), 6.94 (dd, J = 8.4, 2.8 Hz, 1H), 6.67 (t, J = 55.6 Hz, 1H), 4.12 (s, 2H); MS: 737.0 (M-1)⁻.


Example 22

[0361]



3-(1-((4-Chlorophenyl)sulfonyl)-2-(thiophen-2-yl)-1H-indol-3-yl)thiophene-2-carboxamide (**22**)

[0362] To a solution of compound **21** (90 mg, 0.18 mmol) and HATU (137 mg, 0.36 mmol) in DMF (5 mL) was added DIEA (116 mg, 0.90 mmol) at rt. The solution was stirred for 20 min, then NH₄Cl (19 mg, 0.36 mmol) was added and the mixture was stirred at rt for 2 h, cooled, diluted with water and stirred for 10 min. The mixture was filtered to give compound **22** as a yellow solid. ¹H-NMR (CDCl₃, 300 MHz) δ: 5.16 (br s, 2H), 6.68 (d, J = 5.1 Hz, 1H), 7.03 (dd, J = 3.6, 5.1 Hz, 1H), 7.14 (dd, J = 1.2, 3.6 Hz, 1H), 7.23-7.49 (m, 9H), 8.39 (d, J = 8.4 Hz, 1H); MS: 499.0 (M+1)⁺.

Example 23**[0363]**



1-((4-Chlorophenyl)sulfonyl)-3-(2-cyanothiophen-3-yl)-2-(thiophen-2-yl)-1H-indole-5-carboxylic acid (**23**)

[0364] A solution of compound **20/10** (49 mg, 90 μ mol) and LiOH·H₂O (12 mg, 0.27 mmol) in THF/H₂O (3:1, 8 mL) was stirred at rt overnight, concentrated, adjusted to pH to 5-6 with 1N HCl and purified by prep-HPLC to give compound **23** as a yellow solid. ¹H-NMR (CDCl₃, 300 MHz) δ : 8.48 (d, J = 8.7 Hz, 1H), 8.22-8.18 (m, 2H), 7.53 (d, J = 5.1 Hz, 1H), 7.44-7.26 (m, 6H), 7.10-7.07 (m, 1H), 6.88 (d, J = 4.8 Hz, 1H); MS: 522.8 (M-1)⁻.


Example 23/1 to 23/3

[0365] The following Example was prepared similar as described for Example 23 using the appropriate starting material.


#	starting material	structure	analytical data
23/1	20/15		¹ H-NMR (CDCl ₃ , 400 MHz) δ : 8.37 (d, J = 8.7 Hz, 1H), 7.77 (d, J = 5.1, 1H), 7.52-7.45 (m, 2H), 7.39-7.25 (m, 6H), 7.15 (d, J = 3.6 Hz, 1H), 7.07-7.04 (m, 1H), 6.91 (d, J = 5.4 Hz, 1H), 2.92-2.86 (m, 2H), 2.56 (t, J = 7.4 Hz, 2H); MS: 519.0 (M+1) ⁺ , 536.1 (M+18) ⁺ .
23/2	8/5		
23/3	8/6		

Example 24**[0366]**Step 1: Methyl 3-((2-iodophenyl)amino)propanoate (**24a**)**[0367]**

[0368] A mixture of 2-iodoaniline (50 g, 288 mmol) and methyl acrylate (103 mL, 1.14 mol) in AcOH (60 mL) was stirred at 90°C in a sealed tube for 48 h, cooled and filtered. The filtrate was concentrated, diluted with aq. Na₂CO₃ and extracted with EA (2 × 100 mL). The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated to afford compound **24a** as a yellow oil.

Step 2: Methyl 3-((2-iodophenyl)(methyl)amino)propanoate (**24b**)**[0369]**


[0370] A mixture of compound **24a** (17.7 g, 58.1 mmol), CH₃I (29 mL, 46 mmol) and K₂CO₃ (16.3 g, 118 mmol) in ACN (120 mL) was stirred at 80°C in a sealed tube for 48 h, cooled and filtered. The filtrate was concentrated, diluted with H₂O and extracted with EA (2 × 100 mL). The combined organic layer was washed with brine, dried over Na₂SO₄ and concentrated to afford crude compound **24b** as a yellow oil.

Step 3: Methyl 3-(methyl(2-(thiophen-2-ylethynyl)phenyl)amino)propanoate (**24c**)**[0371]**

[0372] A mixture of compound **24b** (24.8 g, 77.7 mmol), 2-ethynyl-thiophene (15.6 mL, 154 mmol), CuI (296 mg, 1.56 mmol), Pd(PPh₃)₂Cl₂ (546 mg, 0.778 mmol) and TEA (39.3 g, 389 mmol) in ACN (90 mL) was stirred at 80°C in a sealed tube overnight, cooled and filtered. The filtrate was concentrated and purified by FCC (PE:EA = 20:1) to afford compound **24c** as a brown oil.

Step 4: Methyl 3-(3-(2-cyanothiophen-3-yl)-2-(thiophen-2-yl)-1*H*-indol-1-yl)propanoate (**24d**)


[0373]



[0374] A mixture of compound **24c** (23.2 g, 77.6 mmol), 3-bromo-thiophene-2-carbonitrile (16.1 g, 85.6 mmol), *n*-Bu₄NI (2.90 g, 7.76 mmol) and PdCl₂(dppf) (1.70 g, 2.33 mmol) in ACN (150 mL) was stirred at 90°C under N₂ overnight, cooled, quenched with H₂O and extracted with EA (2 × 150 mL). The combined organic layer was washed with brine, dried over Na₂SO₄, concentrated and purified by FCC (PE:EA = 5:1) to afford compound **24d** as a brown oil; MS: 393.0 (M+1)⁺.

Step 5: 3-(2-(Thiophen-2-yl)-1*H*-indol-3-yl)thiophene-2-carbonitrile (**24e**)

[0375]




[0376] A mixture of compound **24d** (12.6 g, 32.1 mmol) and DBU (10.1 mL, 64.2 mmol) in DMF (100 mL) was stirred at 120°C overnight, cooled, diluted with H₂O and extracted with EA (2 × 100 mL). The combined organic layer was washed with H₂O (2 × 100 mL) and brine, dried over Na₂SO₄, concentrated and purified by FCC (PE:EA = 5:1) to afford compound **24e** as a yellow solid; MS: 307.0 (M+1)⁺.


Step 6: 3-(1-(Cyclohexylsulfonyl)-2-(thiophen-2-yl)-1*H*-indol-3-yl)thiophene-2-carbonitrile (**24**)

[0377] To a solution of compound **24e** (200 mg, 0.65 mmol) in THF (20 mL) at -78°C under N₂ was added LiHMDS (1.0 M in THF, 0.8 mL, 0.8 mmol) dropwise. The mixture was stirred at -78°C for 30 min, then cyclohexanesulfonyl chloride (144 mg, 0.80 mmol) was added. The mixture was stirred at -78°C for 2 h, diluted with aq. NH₄Cl and extracted with DCM (3 x). The combined organic layer was dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to afford compound **24** as a yellow solid. ¹H-NMR (CDCl₃, 300 MHz) δ: 1.08-1.12 (m, 3H), 1.46-1.60 (m, 3H), 1.73-1.77 (m, 4H), 3.06-3.16 (m, 1H), 6.86 (d, *J* = 5.4 Hz, 1H), 7.05 (dd, *J* = 3.6, 4.8 Hz, 1H), 7.27-7.51 (m, 6H), 8.19 (d, *J* = 8.4 Hz, 1H); MS: 467.7 (M+Na)⁺.


Example 25

[0378]




Step 1: *tert*-Butyl 2-(thiazol-5-yl)-1*H*-indole-1-carboxylate (**25a**)**[0379]**


[0380] A mixture of *tert*-butyl 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-indole-1-carboxylate (4.20 g, 12.2 mmol), 5-bromo-thiazole (2.00 g, 12.2 mmol), Pd(dppf)Cl₂ (877 mg, 1.20 mmol) and K₂CO₃ (5.10 g, 36.6 mmol) in dioxane/H₂O (50 mL/5 mL) was stirred at 100°C under N₂ overnight, cooled, diluted with EA (300 mL) and washed with brine. The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE:EA = 5:1) to give compound **25a** as a colorless oil.

Step 2: *tert*-Butyl 3-bromo-2-(thiazol-5-yl)-1*H*-indole-1-carboxylate (**25b**)**[0381]**

[0382] A mixture of compound **25a** (2.6 g, 8.7 mmol) and NBS (1.85 g, 10.4 mmol) in DMF (50 mL) was stirred at rt under N₂ overnight, diluted with water and extracted with EA (3 × 50 mL). The combined organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE:EA = 10:1) to give compound **25b** as a yellow solid.

Step 3: *tert*-Butyl 3-(2-cyanothiophen-3-yl)-2-(thiazol-5-yl)-1*H*-indole-1-carboxylate (**25c**)**[0383]**

[0384] A mixture of compound **25b** (620 mg, 1.60 mmol), 3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-thiophene-2-carbonitrile (376 mg, 1.60 mmol), Pd(dppf)Cl₂ (117 mg, 160 μmol) and K₂CO₃ (662 mg, 4.80 mmol) in dioxane/H₂O (20 mL/2 mL) was stirred at 100°C under N₂ overnight, cooled, diluted with EA (200 mL) and washed with brine. The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE:EA = 5:1) to give compound **25c** as a yellow oil.

Step 4: 3-(2-(Thiazol-5-yl)-1*H*-indol-3-yl)thiophene-2-carbonitrile (**25d**)**[0385]**

[0386] A mixture of compound **25c** (320 mg, 0.79 mmol) and TFA (4 mL) in DCM (10 mL) was stirred at rt overnight,


concentrated, neutralized with sat. aq. NaHCO_3 and extracted with EA (3×30 mL). The combined organic layer was washed with brine, dried over Na_2SO_4 , filtered and concentrated to afford compound **25d**, which was used in the next step without further purification.

Step 5: 3-(1-((4-Chlorophenyl)sulfonyl)-2-(thiazol-5-yl)-1H-indol-3-yl)thiophene-2-carbonitrile (25**)**

[0387] To a solution of compound **25d** (200 mg, 0.65 mmol) in THF (10 mL) was added NaH (39 mg, 0.98 mmol) under N_2 at 0°C . The mixture was stirred at 0°C for 30 min, then 4-chloro-benzenesulfonyl chloride (165 mg, 0.78 mmol) was added. The mixture was stirred at 0°C for 30 min, poured into sat. aq. NH_4Cl (50 mL) and extracted with EA (3×50 mL). The combined organic layer was washed with brine, dried over Na_2SO_4 , filtered, concentrated and purified by prep-HPLC ($\text{CH}_3\text{CN}/\text{H}_2\text{O} = 20\%$ to 95% , 5 mmol NH_4HCO_3) to afford compound **25** as a yellow solid. $^1\text{H-NMR}$ (DMSO-d_6 , 400 MHz) δ : 9.28 (d, $J = 0.8$ Hz, 1H), 8.27 (d, $J = 8.4$ Hz, 1H), 8.08 (d, $J = 5.2$ Hz, 1H), 7.91 (d, $J = 0.8$ Hz, 1H), 7.66-7.56 (m, 5H), 7.43-7.39 (m, 2H), 7.11 (d, $J = 5.2$ Hz, 1H); MS: 481.8 ($\text{M}+1$) $^+$.


Example 26

[0388]



Step 1: *tert*-Butyl 3-(2-cyanothiophen-3-yl)-1H-indole-1-carboxylate (26a**)**


[0389]



[0390] To a solution of *tert*-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-1-carboxylate (2.0 g, 5.8 mmol), 3-bromo-thiophene-2-carbonitrile (1.1 g, 5.8 mmol) and K_2CO_3 (2.40 g, 17.4 mmol) in dioxane/ H_2O (20 mL/2 mL) was added $\text{Pd}(\text{dppf})\text{Cl}_2$ (413 mg, 0.58 mmol) under N_2 . The mixture was stirred at 90°C for 4 h, evaporated and purified by FCC (PE:EA = 20:1 to 10:1) to afford compound **26a** as a yellow oil.

Step 2: *tert*-Butyl 2-bromo-3-(2-cyanothiophen-3-yl)-1H-indole-1-carboxylate (26b**)**


[0391]



[0392] To a solution compound **26a** (1.30 g, 4.01 mmol) in CCl_4 (20 mL) were added NBS (1.40 g, 8.02 mmol) and AIBN (65.4 mg, 401 μmol). The mixture was stirred at 100°C for 48 h, evaporated and purified by FCC (PE:EA = 10:1) to afford compound **26b** as a yellow oil.

Step 3: *tert*-Butyl 3-(2-cyanothiophen-3-yl)-2-phenyl-1*H*-indole-1-carboxylate (**26c**)

[0393]




[0394] To a solution of compound **26b** (500 mg, 1.24 mmol), PhB(OH)₂ (302 mg, 2.48 mmol) and K₂CO₃ (513 mg, 3.72 mmol) in dioxane (15 mL) was added Pd(dppf)Cl₂ (88.4 mg, 124 μmol) under N₂.

[0395] The mixture was stirred at 90°C overnight, concentrated and purified by FCC (PE:EA = 10:1) to afford compound **26c** as a yellow oil.

Step 4: 3-(2-Phenyl-1*H*-indol-3-yl)thiophene-2-carbonitrile (**26d**)

[0396]



[0397] To a solution of compound **26c** (616 mg, 1.54 mmol) in DCM (4 mL) was added TFA (2 mL). The mixture was stirred at rt for 2 h, concentrated and purified to afford compound **26d** as a white solid.

Step 5: 3-(1-((4-Chlorophenyl)sulfonyl)-2-phenyl-1*H*-indol-3-yl)thiophene-2-carbonitrile (**26**)

[0398] To a solution of compound **26d** (147 mg, 488 μmol) in DMF (5 mL) was added NaH (78 mg, 2.0 mmol) under N₂ at 0°C. The mixture was stirred at 0°C for 30 min, then 4-chloro-benzenesulfonyl chloride (310 mg, 1.46 mmol) was added. The mixture was stirred at 0°C for 30 min, quenched with sat. aq. NH₄Cl and extracted with DCM (3 × 20 mL). The combined organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE:EA = 20:1) to afford compound **26** as a white solid. ¹H-NMR (DMSO-d₆, 400 MHz) δ: 8.25 (d, J = 8.0 Hz, 1H), 7.99 (d, J = 5.2 Hz, 1H), 7.59 (dd, J = 2.0, 6.8 Hz, 2H), 7.53-7.27 (m, 10H), 6.96 (d, J = 4.8 Hz, 1H); MS: 491.7 (M+18)⁺.


Example 26/1

[0399] The following Example was prepared similar as described for Example 26 using the appropriate starting material.

#	starting material	structure	analytical data
26/1			¹ H-NMR (DMSO-d ₆ , 400 MHz) δ: 8.25 (d, J = 8.4 Hz, 1H), 8.03 (d, J = 5.6 Hz, 1H), 7.61-7.58 (m, 2H), 7.55-7.48 (m, 3H), 7.43-7.32 (m, 4H), 7.25-7.21 (m, 2H), 7.00 (d, J = 4.8 Hz, 1H); MS: 509.6 (M+18) ⁺ .

Example 27

[0400]

**27**

2-((3-(2-Cyanothiophen-3-yl)-2-phenyl-1H-indol-1-yl)sulfonyl)phenoxy)acetic acid (27**)**

[0401] To a solution of compound **1/43** (70 mg, 130 μ mol) in MeOH (5 mL) was added NaOH (2N, 0.5 mL) and the mixture was stirred overnight. Then the MeOH was removed and the solution was adjusted to pH <2 with 2N HCl, extracted with EA (10 mL) and washed with brine (10 mL). The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to afford compound **27** as a white solid. ¹H-NMR (500 MHz, DMSO-d₆) δ : 13.21 (br s, 1H), 8.26 (d, J = 8.5 Hz, 1H), 7.99 (d, J = 5.0 Hz, 1H), 7.52-7.34 (m, 8H), 7.27 (d, J = 7.0 Hz, 2H), 7.01-6.99 (m, 3H), 4.75 (s, 2H); MS: 515.1 (M+1)⁺.


Example 27/1 to 27/3

[0402] The following Examples were prepared similar as described for Example 27 using the appropriate starting material.

#	starting material	structure	analytical data
27/1			¹ H-NMR (500 MHz, DMSO-d ₆) δ : 12.55 (br s, 1H), 8.26 (d, J = 9.0 Hz, 1H), 8.00 (d, J = 5.0 Hz, 1H), 7.55-7.34 (m, 10H), 7.28 (d, J = 7.0 Hz, 2H), 6.98 (d, J = 5.0 Hz, 1H), 3.61 (s, 2H); MS: 499.1 (M+1) ⁺ .
27/2			¹ H-NMR (500 MHz, DMSO-d ₆) δ : 13.18 (br s, 1H), 8.32 (d, J = 8.5 Hz, 1H), 8.17 (s, 1H), 8.00 (d, J = 5.0 Hz, 2H), 7.93 (d, J = 8.0 Hz, 1H), 7.86 (d, J = 8.5 Hz, 2H), 7.63-7.52 (m, 4H), 7.45-7.32 (m, 7H), 6.97 (d, J = 5.0 Hz, 1H); MS: 561.2 (M+1) ⁺ .
27/3			¹ H-NMR (500 MHz, DMSO-d ₆) δ : 12.85 (br s, 1H), 8.21 (d, J = 8.0 Hz, 1H), 8.01 (d, J = 5.5 Hz, 1H), 7.54-7.51 (m, 1H), 7.44-7.32 (m, 8H), 7.00 (d, J = 5.0 Hz, 1H), 6.95 (d, J = 3.5 Hz, 1H), 3.91 (s, 2H); MS: 505.0 (M+1) ⁺ .

Example 28

[0403]




3-(4-(3-(2-Cyanothiophen-3-yl)-1-((4-methoxyphenyl)sulfonyl)-1*H*-indol-2-yl)phenyl)propan-amide (28)

[0404] To a solution of compound **15/6** (200 mg, 0.40 mmol) in DMF (10mL) was added EDCI (100 mg, 0.50 mmol), DMAP (60 mg, 0.50 mmol) and NH_4Cl (70 mg, 0.50 mmol) and the mixture was stirred at rt for 12 h, diluted with water (100 mL) and extracted with EA (3×100 mL). The combined organic layer was washed with brine (50 mL), dried over Na_2SO_4 , filtered, concentrated and purified by prep-HPLC to give compound **28** as a white solid. $^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.26 (d, $J = 8.0$ Hz, 1H), 7.98 (d, $J = 5.5$ Hz, 1H), 7.50-7.47 (m, 1H), 7.40-7.33 (m, 5H), 7.20 (d, $J = 8.0$ Hz, 2H), 7.15 (d, $J = 8.0$ Hz, 2H), 7.00 (d, $J = 9.0$ Hz, 2H), 6.93 (d, $J = 5.0$ Hz, 1H), 6.84 (s, 1H), 3.78 (s, 3H), 2.85 (t, $J = 8.0$ Hz, 2H), 2.41 (t, $J = 8.0$ Hz, 2H); MS: 542.1 ($\text{M}+1$) $^+$.


Example 29

[0405]



Step 1: Methyl 2-chloro-3'-((2-((4-(difluoromethyl)phenyl)sulfonamido)-5-fluorophenyl)ethynyl)-[1,1'-biphenyl]-4-carboxylate (29a)


[0406]



[0407] Compound **29a** was synthesized similar as described in Example 1, Step 1 and 2 using the appropriate building blocks.

Step 2: Methyl 2-chloro-3'-((1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (29b)


[0408]



[0409] To a solution of compound **29a** (400 mg, 0.70 mmol) in MeCN (12.0 mL) was added K_2CO_3 (193 mg, 1.40 mmol) and $Pd(PPh_3)_4$ (81 mg, 70 μ mol) under N_2 . The mixture was stirred at 100°C for 2 h, cooled to rt, poured into EA (200 mL) and washed with H_2O (2×20 mL) and brine (20 mL). The organic layer was dried over Na_2SO_4 , concentrated and purified by FCC (EA:PE = 1:4) to give compound **29b** as a colorless oil.

Step 3: Methyl 3'-(3-bromo-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)-2-chloro-[1,1'-biphenyl]-4-carboxylate (**29c**)

[0410]





[0411] To a solution of compound **29b** (150 mg, 0.26 mmol) in THF (15 mL) was added NBS (56 mg, 0.31 mmol). The mixture was stirred at rt overnight, poured into EA (200 mL) and washed with H_2O (2×20 mL) and brine (20 mL). The organic layer was dried over Na_2SO_4 , concentrated and purified by prep-TLC (EA:PE = 1:4) to give compound **29c** as a white solid.

Step 4: Methyl 2-chloro-3'-(1-((4-(difluoromethyl)phenyl)sulfonyl)-3-(2,6-dimethylphenyl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (**29**)

[0412] To a solution of compound **29c** (150 mg, 0.23 mmol) in dioxane (8 mL) was added 2,6-dimethyl-phenylboronic acid (45 mg, 0.30 mmol), Cs_2CO_3 (176 mg, 0.46 mmol) and $Pd(dppf)Cl_2$ (17 mg, 23 μ mol) under N_2 . The mixture was stirred at 90°C overnight, cooled to rt, poured into EA (200 mL) and washed with H_2O (2×20 mL) and brine (20 mL). The organic layer was dried over Na_2SO_4 , concentrated and purified by prep-TLC (EA:PE = 1:4) to give compound **29** as a colorless oil.

Example 29/1 to 29/3

[0413] The following Examples were prepared similar as described for Example **29** using the appropriate starting material(s).











**Example 30****[0414]**

rac-(1*R*,2*R*)-2-(3-(3-(2-Cyanothiophen-3-yl)-1-tosyl-1*H*-indol-2-yl)phenyl)cyclopropane-1-carboxylic acid (**30**)











[0415] To a solution of compound **1/56** (130 mg, 0.23 mmol) in MeOH (10 mL) was added LiOH·H₂O (49 mg, 1.18 mmol) and the mixture was stirred at rt for 1 h. Then the mixture was concentrated, adjusted to pH <4 with 2N aq. HCl and extracted with EA (3 × 30 mL). The combined organic layer was washed with brine (10 mL), dried over Na₂SO₄, concentrated and purified by prep-HPLC to give compound **30** as a white solid. ¹H-NMR (500 MHz, DMSO-*d*₆) δ: 12.36 (s, 1H), 8.27 (d, *J* = 10.0 Hz, 2H), 8.01 (d, *J* = 5.5 Hz, 1H), 7.52-7.48 (m, 1H), 7.41-7.26 (m, 8H), 7.10-7.08 (m, 1H), 7.02-6.98 (m, 1H), 6.88 (s, 1H), 2.38-2.33 (m, 1H), 2.31 (s, 3H), 1.73-1.68 (m, 1H), 1.44-1.39 (m, 1H), 1.25-1.18 (m, 1H). MS: 521 (M-18+H)⁺.

Example 30/1 to 30/16











[0416] The following Examples were prepared similar as described for Example **30** using the appropriate starting materials.

#	starting material	structure	analytical data
30/1	 1/57		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 13.12 (s, 1H), 8.31 (d, J = 8.5 Hz, 1H), 8.07 (s, 1H), 8.01 (d, J = 5.0 Hz, 1H), 7.96 (d, J = 8.0 Hz, 1H), 7.81 (d, J = 7.5 Hz, 1H), 7.76 (d, J = 8.0 Hz, 1H), 7.63-7.60 (m, 1H), 7.55-7.47 (m, 2H), 7.42-7.37 (m, 4H), 7.31-7.25 (m, 3H), 7.07 (d, J = 5.0 Hz, 1H), 2.24 (s, 3H); MS: 574.8 ($\text{M}+1$) $^+$.
30/2	 1/58		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.43 (d, J = 8.5 Hz, 1H), 7.83 (d, J = 5.0 Hz, 1H), 7.65 (d, J = 8.0 Hz, 1H), 7.51-7.32 (m, 9H), 7.27 (d, J = 8.5 Hz, 2H), 7.22 (s, 1H), 7.12 (d, J = 8.5 Hz, 2H), 6.98 (d, J = 5.0 Hz, 1H), 2.22 (s, 3H), 1.62 (s, 6H); MS: 615.0 ($\text{M}+1$) $^+$.
30/3	 1/59		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.75 (s, 1H), 8.75 (s, 1H), 8.66 (d, J = 2.0 Hz, 1H), 8.28 (d, J = 8.5 Hz, 1H), 8.05-8.02 (m, 2H), 7.84 (d, J = 7.5 Hz, 1H), 7.62 (s, 1H), 7.54-7.25 (m, 9H), 7.11 (d, J = 5.0 Hz, 1H), 2.23 (s, 3H), 1.61 (s, 6H); MS: 618.1 ($\text{M}+1$) $^+$.
30/4	 1/60		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.35 (s, 1H), 8.26 (d, J = 9.0 Hz, 1H), 8.02 (d, J = 5.0 Hz, 1H), 7.61-7.44 (m, 5H), 7.42-7.08 (m, 4H), 7.07-7.06 (m, 1H), 7.00-6.94 (m, 2H), 2.39-2.34 (m, 1H), 1.72 (s, 1H), 1.44-1.40 (m, 1H), 1.25-1.23 (m, 1H); MS: 540.8 ($\text{M}+1$) $^+$.
30/5	 1/61		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.05 (s, 1H), 8.27 (d, J = 8.5 Hz, 1H), 7.98 (d, J = 5.0 Hz, 1H), 7.60-7.46 (m, 5H), 7.40-7.23 (m, 4H), 7.13 (d, J = 7.5 Hz, 1H), 7.03 (s, 1H), 6.96 (d, J = 5.0 Hz, 1H), 2.56 (t, J = 8.0 Hz, 2H), 2.15 (t, J = 7.0 Hz, 2H), 1.74-1.70 (m, 2H); MS: 560.8 ($\text{M}+1$) $^+$.



(continued)


#	starting material	structure	analytical data
30/ 6	 1/63		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.34 (d, $J = 8.5$ Hz, 1H), 7.96 (d, $J = 5.0$ Hz, 1H), 7.83-7.79 (m, 1H), 7.73 (d, $J = 9.0$ Hz, 2H), 7.62-7.58 (m, 1H), 7.53-7.42 (m, 4H), 7.23 (d, $J = 5.0$ Hz, 1H), 6.87 (d, $J = 8.5$ Hz, 1H), 6.77 (d, $J = 7.0$ Hz, 1H), 4.54-4.43 (m, 4H), 3.77-3.73 (m, 1H); MS: 574.7 ($\text{M}+1$) $^+$.
30/ 7	 1/64		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.38 (d, $J = 8.0$ Hz, 1H), 7.77 (d, $J = 5.0$ Hz, 1H), 7.51-7.34 (m, 7H), 7.16 (t, $J = 7.5$ Hz, 1H), 6.88 (d, $J = 5.0$ Hz, 1H), 6.65 (d, $J = 7.5$ Hz, 1H), 6.55-6.53 (m, 1H), 6.23 (s, 1H), 4.01-3.97 (m, 2H), 3.91-3.88 (m, 2H), 3.54-3.50 (m, 1H); MS: 574.1 ($\text{M}+1$) $^+$.
30/ 8	 1/65		$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ : 8.26 (d, $J = 8.8$ Hz, 1H), 8.06 (d, $J = 5.5$ Hz, 1H), 7.64-7.51 (m, 5H), 7.42-7.36 (m, 2H), 7.05 (d, $J = 5.0$ Hz, 1H), 7.02 (d, $J = 5.0$ Hz, 1H), 6.88 (d, $J = 3.5$ Hz, 1H), 2.52-2.45 (m, 1H), 1.78-1.75 (m, 1H), 1.48-1.42 (m, 1H), 1.24-1.18 (m, 1H); MS: 582.1 ($\text{M}+18$) $^+$.
30/ 9	 1/66		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 8.25 (d, $J = 8.5$ Hz, 1H), 8.05 (d, $J = 5.0$ Hz, 1H), 7.60 (d, $J = 8.5$ Hz, 2H), 7.55-7.47 (m, 3H), 7.42-7.36 (m, 3H), 7.10 (s, 1H), 7.05 (d, $J = 5.0$ Hz, 1H), 6.90 (s, 1H), 2.39-2.35 (m, 1H), 1.79-1.72 (m, 1H), 1.43-1.38 (m, 1H), 1.25-1.20 (m, 1H); MS: 610.0 ($\text{M}+18$) $^+$.
30/ 10	 1/67		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.29 (s, 1H), 8.14 (d, $J = 8.5$ Hz, 1H), 8.00 (d, $J = 5.5$ Hz, 1H), 7.89 (s, 1H), 7.82 (s, 1H), 7.46-7.34 (m, 3H), 7.25-7.19 (m, 2H), 7.14-7.10 (m, 2H), 6.99 (d, $J = 5.0$ Hz, 1H), 3.65 (s, 3H), 2.37-2.35 (m, 1H), 1.82-1.75 (m, 1H), 1.45-1.38 (m, 1H), 1.28-1.22 (m, 1H); MS: 529.2 ($\text{M}+1$) $^+$.

(continued)

#	starting material	structure	analytical data
30/ 11	 1/68		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.49 (s, 1H), 8.26 (d, J = 8.5 Hz, 1H), 7.99 (d, J = 5.0 Hz, 1H), 7.58-7.40 (m, 5H), 7.40-7.32 (m, 2H), 7.16-7.13 (m, 1H), 6.98 (d, J = 5.0 Hz, 1H), 6.57 (d, J = 8.0 Hz, 1H), 6.52 (d, J = 7.5 Hz, 1H), 6.27 (s, 1H), 3.33-3.28 (m, 2H), 3.21-3.14 (m, 3H), 2.21-2.13 (m, 2H); MS: 587.8 ($\text{M}+1$) $^+$.
30/ 12	 1/69		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.35 (br s, 1H), 8.13 (d, J = 8.0 Hz, 1H), 8.04 (s, 1H), 7.52-7.39 (m, 3H), 7.27-7.13 (m, 4H), 7.04 (d, J = 5.0 Hz, 1H), 2.88-2.83 (m, 1H), 2.41-2.36 (m, 1H), 1.75-1.72 (m, 1H), 1.43-1.39 (m, 1H), 1.28-1.25 (m, 1H), 0.99-0.94 (m, 4H); MS: 471.0 ($\text{M}-18+\text{H}$) $^+$.
30/ 13	 1/70		$^1\text{H-NMR}$ (500 MHz, CD_3OD) δ : 8.37 (d, J = 8.0 Hz, 1H), 7.46-7.42 (m, 1H), 7.33-7.30 (m, 1H), 7.27-7.17 (m, 9H), 6.74 (s, 1H), 6.60 (d, J = 5.5 Hz, 1H), 2.40-2.36 (m, 1H), 2.35 (s, 3H), 1.76-1.72 (m, 1H), 1.51-1.48 (m, 1H), 1.17-1.13 (m, 1H); MS: 548.0 ($\text{M}+1$) $^+$.
30/ 14	 1/71		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.38 (s, 1H), 8.25 (d, J = 8.0 Hz, 1H), 7.49-7.45 (m, 1H), 7.36-7.13 (m, 8H), 7.15 (d, J = 7.0 Hz, 1H), 6.96-6.92 (m, 2H), 6.48-6.46 (m, 1H), 2.39-2.34 (m, 1H), 2.31 (s, 3H), 1.73-1.69 (m, 1H), 1.45-1.40 (m, 1H), 1.27-1.23 (m, 1H); MS: 530.0 ($\text{M}-1$) $^-$.
30/ 15	 1/72		$^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.35 (br s, 1H), 8.25 (d, J = 8.5 Hz, 1H), 7.88-7.85 (m, 1H), 7.57-7.34 (m, 8H), 7.32-7.18 (m, 3H), 7.06 (s, 1H), 6.84 (s, 1H), 2.32-2.29 (m, 4H), 1.64-1.61 (m, 1H), 1.43-1.39 (m, 1H), 1.19-1.16 (m, 1H); MS: 549.0 ($\text{M}-1$) $^-$.


(continued)

#	starting material	structure	analytical data
30/ 16			$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ : 8.32-8.28 (m, 1H), 7.94-7.88 (m, 3H), 7.65-7.24 (m, 12H), 7.04-6.85 (m, 2H), 5.19 (br s, 1H), 1.23 (s, 6H); MS: 663.0 ($\text{M}-1$) ⁻

Example 31**[0417]**

3-((6-(1-((4-Chlorophenyl)sulfonyl)-3-(2-cyanothiophen-3-yl)-1H-indol-2-yl)pyridin-2-yl)oxy)propanoic acid (**31**)

[0418] A solution of compound **1/62** (110 mg, 0.19 mmol) in 4N HCl in dioxane (30 mL) was stirred at rt overnight. The solvent was removed, EA (20 mL) was added and the mixture was washed with water (10 mL) and brine (10 mL). The organic layer was dried over Na_2SO_4 , concentrated and purified by prep-HPLC to give compound **31** as a white solid. $^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ : 12.38 (s, 1H), 8.13 (d, $J = 8.5$ Hz, 1H), 8.02 (d, $J = 5.0$ Hz, 1H), 7.94-7.91 (m, 2H), 7.77-7.65 (m, 3H), 7.54-7.42 (m, 1H), 7.41-7.37 (m, 2H), 7.13 (d, $J = 6.8$ Hz, 1H), 7.00 (d, $J = 4.8$ Hz, 1H), 6.86 (d, $J = 8.0$ Hz, 1H), 4.32 (t, $J = 6.4$ Hz, 2H), 2.67 (t, $J = 6.4$ Hz, 2H); MS: 563.8 ($\text{M}+1$)⁺.

Example 32**[0419]**











3'-((3-(2-Cyanothiophen-3-yl)-1-tosyl-1H-indol-2-yl)-N-(methylsulfonyl)-[1,1'-biphenyl]-3-carboxamide (**32**)


[0420] A cloudy solution of compound **30/1** (100 mg, 0.17 mmol), methanesulfonamide (17 mg, 0.17 mmol), DMAP (21 mg, 0.17 mmol) and EDCI (50 mg, 0.26 mmol) in DMF (4 mL) was stirred for 14 h at rt. The product was purified from the mixture by prep-HPLC to give compound **32** as a white solid. $^1\text{H-NMR}$ (500 MHz, DMSO-d_6) δ : 12.30 (s, 1H), 8.31-7.92 (m, 4H), 7.83 (t, $J = 7.5$ Hz, 2H), 7.65-7.62 (m, 2H), 7.55-7.49 (m, 2H), 7.42-7.39 (m, 4H), 7.30-7.26 (m, 3H),

7.07 (d, $J = 2.5$ Hz, 1H), 3.42 (s, 3H), 2.25 (s, 3H); MS: 652.1 (M+1)⁺.

Example 32/1 to 32/5


5 **[0421]** The following Examples were prepared similar as described for Example 32 using the appropriate starting materials.

#	starting material	structure	analytical data
10 32/1	 30/4		¹ H-NMR (500 MHz, DMSO-d ₆) δ: 12.09 (s, 1H), 8.25 (d, $J = 8.5$ Hz, 1H), 8.01 (d, $J = 5.0$ Hz, 1H), 7.62-7.59 (m, 2H), 7.54-7.23 (m, 7H), 7.10-6.96 (m, 3H), 3.28 (s, 1H), 2.46-2.42 (m, 1H), 2.09-2.06 (m, 1H), 1.51-1.46 (m, 1H), 1.35-1.32 (m, 1H); MS: 658.0 (M+Na) ⁺ .
15 32/2	 3/32		¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.27 (dd, $J = 9.1, 4.4$ Hz, 1H), 7.98 (d, $J = 5.1$ Hz, 1H), 7.46-7.28 (m, 5H), 7.20-7.07 (m, 2H), 6.94 (d, $J = 5.1$ Hz, 1H), 6.57-6.48 (m, 2H), 6.24 (s, 1H), 3.95 (t, $J = 7.5$ Hz, 2H), 3.89-3.80 (m, 1H), 3.72 (t, $J = 6.2$ Hz, 2H), 3.52 (s, 3H), 2.87 (s, 3H), 2.33 (s, 3H), 1.34 (s, 6H); MS: 685.0 (M+1) ⁺ .
20 32/3	 3/32		
25 32/4	 30/4	 32/4	¹ H-NMR (500 MHz, DMSO-d ₆) δ: 12.32 (s, 1H), 8.31-8.28 (m, 1H), 8.11 (d, $J = 1.5$ Hz, 1H), 8.03 (d, $J = 5.0$ Hz, 1H), 7.99-7.97 (m, 2H), 7.55-7.35 (m, 4H), 7.41-7.33 (m, 3H), 7.26-7.18 (m, 3H), 7.07-7.03 (m, 2H), 3.38 (s, 3H), 2.22 (s, 3H); MS: 701.9 (M-1) ⁻ .
30 32/5	 11/12		¹ H-NMR (500 MHz, CD ₃ OD) δ: 8.44 (dd, $J = 9.0, 4.0$ Hz, 1H), 8.08-8.04 (m, 3H), 7.93 (dd, $J = 8.5, 1.5$ Hz, 1H), 7.73 (t, $J = 8.3$ Hz, 1H), 7.55-7.32 (m, 9H), 7.12 (s, 1H), 6.95 (dd, $J = 8.0, 2.5$ Hz, 1H), 6.69 (t, $J = 55.5$ Hz, 1H), 3.36 (s, 3H); MS: 756.8 (M-1) ⁻ .


Example 33**[0422]**

Methyl 3'-((3-(2-cyanothiophen-3-yl)-5-fluoro-7-hydroxy-1-tosyl-1H-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (**33**)

[0423] To a solution of compound **1/101** (120 mg, 0.19 mmol) in DCM (6 mL) at -78°C was slowly added BBr₃ (10 mL, 1M in DCM). The mixture was stirred at this temperature for 40 min and at rt for 1 h, quenched with H₂O (20 mL) and extracted with EA (2 × 100 mL). The combined organic layer was washed with brine (20 mL), dried over Na₂SO₄ and concentrated. The residue was purified by prep-TLC (EA:PE = 1:1) to afford compound **33** as a yellow oil.

Example 34**[0424]**

Step 1: Ethyl 2-((3'-(3-(2-cyanothiophen-3-yl)-1-tosyl-1H-indol-2-yl)-5-fluoro-4-(hydroxymethyl)-[1,1'-biphenyl]-3-yl)sulfonyl)acetate (**34a**)

[0425]


[0426] To a solution of compound **1** (290 mg, 0.54 mmol) in dioxane (15 mL) was added compound **P3-1** (193 mg, 0.54 mmol), B₂Pin₂ (166 mg, 0.65 mmol), Pd(dppf)Cl₂ (39 mg, 0.05 mmol) and KOAc (107 mg, 1.09 mmol). The mixture was stirred at 100°C overnight. After cooling to rt the mixture was filtered, the filtrate was concentrated under reduced pressure and the residue was purified by prep-TLC (EA:PE = 1:1) to afford compound **34a** as a yellow oil.

Step 2: 2-((3'-(3-(2-Cyanothiophen-3-yl)-1-tosyl-1*H*-indol-2-yl)-5-fluoro-4-(hydroxymethyl)-[1,1'-biphenyl]-3-yl)sulfonyl)acetic acid (**34**)

[0427] To a solution of compound **34a** (90 mg, 0.12 mmol) in EtOH (10 mL) was added LiOH·H₂O (26 mg, 0.62 mmol) and the mixture was stirred at rt for 1.5 h. Then the EtOH was removed, water was added and the pH was adjusted to <4 by addition of 2N HCl. The mixture was extracted with EA (3 × 40 mL) and the combined organic layer was washed with brine (10 mL), dried over Na₂SO₄, concentrated and purified by prep-HPLC to afford compound **34** as a white solid. ¹H-NMR (500 MHz, CD₃OD) δ: 8.42 (d, J = 8.5 Hz, 1H), 7.99 (s, 1H), 7.84 (d, J = 4.5 Hz, 1H), 7.75 (d, J = 8.0 Hz, 1H), 7.64-7.30 (m, 9H), 7.20 (d, J = 8.5 Hz, 2H), 7.03 (d, J = 5.0 Hz, 1H), 5.11 (s, 2H), 4.50 (s, 2H), 2.29 (s, 3H); MS: 718.1 (M+18)⁺.


Example 35

[0428]



Step 1: Methyl 3'-((2-amino-5-fluorophenyl)ethynyl)-2-chloro-[1,1'-biphenyl]-4-carboxylate (**35a**)


[0429]



[0430] To a solution of compound **P5** (5.00 g, 15.4 mmol) in TEA (60 mL) was added Pd(PPh₃)₄ (710 mg, 0.61 mmol), CuI (175 mg, 0.92 mmol), PPh₃ (241 mg, 0.92 mmol), and 2-ethynyl-4-fluoroaniline (2.70 g, 20.0 mmol). The mixture was stirred at 60°C under N₂ overnight. After cooling to rt the mixture was filtered, the filtrate was concentrated and the residue was purified by FCC (PE:EA = 2:1) to give compound **35a** as a light yellow solid.

Step 2: Methyl 2-chloro-3'-((5-fluoro-2-(2,2,2-trifluoroacetamido)phenyl)ethynyl)-[1,1'-biphenyl]-4-carboxylate (**35b**)


[0431]



[0432] To a solution of compound **35a** (300 mg, 0.79 mmol) in DCM (15 mL) was added TFAA (199 mg, 0.95 mmol) and TEA (120 mg, 1.19 mmol). The mixture was stirred at rt for 15 min, then DCM (20 mL) was added and the mixture was washed with H₂O (2 × 10 mL) and brine (20 mL). The organic layer was dried over Na₂SO₄ and concentrated to dryness to afford crude compound **35b** as a yellow solid.

Step 3: Methyl 2-chloro-3'-(3-(2-cyanothiophen-3-yl)-5-fluoro-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (**35c**)

[0433]




[0434] To a solution of compound **35b** (320 mg, 0.67 mmol) in ACN (20 mL) was added 3-bromothiophene-2-carbonitrile (190 mg, 1.01 mmol), K₂CO₃ (185 mg, 1.34 mmol), and Pd(PPh₃)₄ (77 mg, 67 μmol) under N₂ and the mixture was stirred at 100°C for 2 h, cooled to rt, poured into EA (20 mL) and washed with H₂O (2 × 20 mL) and brine (20 mL). The organic layer was dried over Na₂SO₄, concentrated and purified by FCC (EA:PE = 1:3) to give compound **35c** as a yellow solid.


Step 4: Methyl 2-chloro-3'-(3-(2-cyanothiophen-3-yl)-5-fluoro-1-((6-(trifluoromethyl)pyridin-3-yl)sulfonyl)-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (**35**)

[0435] To a solution of compound **35c** (200 mg, 0.41 mmol) in THF (8 mL) at 0°C was added NaH (60% in mineral oil, 50 mg, 1.23 mmol) and 6-(trifluoromethyl)pyridine-3-sulfonyl chloride (201 mg, 0.82 mmol). The mixture was stirred at rt for 1 h and poured into cold sat. aq. NH₄Cl (50 mL). The mixture was extracted with EA (2 × 50 mL) and washed with brine (20 mL). The combined organic layer was dried over Na₂SO₄, concentrated and purified by prep-TLC (EA:PE = 1:3) to give compound **35** as a yellow solid.

Example 35/1 to 35/2

[0436] The following Examples were prepared similar as described for Example **35** using the appropriate starting materials.



Example 36**[0437]**


5-(3-(3-(2-Cyanothiophen-3-yl)-5-fluoro-1-tosyl-1H-indol-2-yl)phenyl)-4-methylpicolinic acid (**36**)

[0438] To a stirred solution of compound **2/13** (150 mg, 0.24 mmol) in THF (10 mL) at rt was added 1N LiOH (1 mL) and stirring was continued at rt for 2 h. The mixture was extracted with EA (100 mL), the organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound **36** as a white solid. ¹H-NMR (500 MHz, DMSO-d₆) δ: 8.39 (s, 1H), 8.28 (dd, J = 10.0, 4.5 Hz, 1H), 8.05 (d, J = 5.0 Hz, 1H), 8.01 (s, 1H), 7.57-7.51 (m, 2H), 7.42-7.36 (m, 4H), 7.27 (d, J = 8.5 Hz, 2H), 7.20-7.18 (m, 1H), 7.13 (s, 1H), 7.06 (d, J = 5.0 Hz, 1H), 2.24 (s, 3H), 2.19 (s, 3H); MS: 606.0 (M-1)⁻.

Example 36/1

[0439] The following Example was prepared similar as described for Example **36** using the appropriate starting material.

#	starting material	structure	analytical data
36/1			¹ H-NMR (500 MHz, DMSO-d ₆) δ: 8.30-8.27 (m, 1H), 8.04 (d, J = 5.0 Hz, 1H), 7.91 (d, J = 7.5 Hz, 1H), 7.65 (d, J = 7.5 Hz, 1H), 7.54-7.48 (m, 2H), 7.40-7.36 (m, 4H), 7.28-7.26 (m, 2H), 7.19-7.17 (m, 1H), 7.12 (s, 1H), 7.05 (d, J = 5.0 Hz, 1H), 2.30 (s, 3H), 2.25 (s, 3H); MS: 607.8 (M+1) ⁺ .



Example 37**[0440]**

3-(2-(2'-Chloro-4'-(1*H*-tetrazol-5-yl)-[1,1'-biphenyl]-3-yl)-5-fluoro-1-tosyl-1*H*-indol-3-yl)thiophene-2-carbonitrile (**37**)

[0441] To a stirred solution of compound **P2/15** (150 mg, 0.17 mmol) in acetone (10 mL) at rt was added 1N HCl (1 mL) and stirring was continued for 2 h. Water was added and the mixture was extracted with EA (100 mL). The organic layer was washed with brine, dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound **37** as a white solid. ¹H-NMR (500 MHz, DMSO-*d*₆) δ: 8.31-8.28 (m, 1H), 8.19 (s, 1H), 8.11-8.09 (m, 1H), 8.03 (d, *J* = 5.0 Hz, 1H), 7.59-7.56 (m, 3H), 7.47-7.20 (m, 7H), 7.11 (s, 1H), 7.04 (d, *J* = 5.0 Hz, 1H), 2.21 (s, 3H); MS: 649.0 (*M*-1)⁻.


Example 37/1

[0442] The following Example was prepared similar as described for Example **37** using the appropriate starting material.

#	starting material	structure	analytical data
37/1			¹ H-NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 8.33-8.30 (m, 1H), 8.16 (s, 1H), 8.02-7.76 (m, 5H), 7.61 (s, 1H), 7.51 (t, <i>J</i> = 7.5 Hz, 1H), 7.41-7.37 (m, 3H), 7.30-7.26 (m, 3H), 7.20 (dd, <i>J</i> = 8.5, 2.5 Hz, 1H), 7.07 (d, <i>J</i> = 5.0 Hz, 1H), 3.71 (s, 3H), 2.24 (s, 3H); MS: 694.0 (<i>M</i> +1) ⁺ .

Example 38

[0443]





rel-Methyl (2*R*,4*R*)-1-(3-(3-(2-cyanothiophen-3-yl)-5-fluoro-1-tosyl-1*H*-indol-2-yl)phenyl)-2-methylpiperidine-4-carboxylate (**38**)

[0444] To a solution of compound **1** (500 mg, 0.91 mmol) in toluene (15 mL) was added *rel*-methyl (2*R*,4*R*)-2-methylpiperidine-4-carboxylate (215 mg, 1.36 mmol), Cs₂CO₃ (869 mg, 2.27 mmol), Pd₂(dba)₃ (83 mg, 90 μmol) and BINAP (113 mg, 0.18 mmol) under N₂. The mixture was stirred at 100°C overnight, cooled to rt, poured into EA (200 mL) and washed with H₂O (30 mL) and brine (30 mL). The organic layer was dried over Na₂SO₄, concentrated and purified by prep-TLC (EA:PE = 1:1) to give compound **38** as a yellow oil.


Example 38/1

[0445] The following Example was prepared similar as described for Example **38** using the appropriate starting materials.

**Example 39****[0446]**

Methyl 2-chloro-3'-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-3-(2-(3-fluoroazetidin-3-yl)phenyl)-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (39)

[0447] To a solution of compound **1/114** (30 mg, 40 μ mol) in DCM (2 mL) was added TFA (0.2 mL) and the mixture was stirred at rt for 4 h. The mixture was poured into water and the pH was adjusted to 8 with sat. aq. NaHCO_3 . Then the mixture was extracted with EA and the organic layer was washed with brine, dried over Na_2SO_4 and concentrated to dryness to give compound **39** as a yellow solid.

Example 40**[0448]**


Methyl 2-chloro-3'-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-3-(2-(3-fluoro-1-methylazetidin-3-yl)phenyl)-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (**40**)

[0449] To a solution of compound **39** (27 mg, 40 μ mol) in MeOH (2 mL) was added formaldehyde (0.2 mL) and the mixture was stirred at rt for 1 h. Then $\text{NaBH}(\text{OAc})_3$ (82 mg, 0.37 mmol) was added and the mixture was stirred at rt for overnight. Water (40 mL) was added and the mixture was extracted with DCM (3×20 mL). The combined organic layer was washed with brine (30 mL), dried over Na_2SO_4 , filtered, concentrated and purified by FCC (PE:EA = 1:1) to afford

compound **40** as a yellow solid.

Example 41/1 and Example 41/2

[0450]




Separated atropisomers of 2-chloro-3'-(3-(2-cyano-6-methylphenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1H-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid (**41/1** and **41/2**)

[0451] Compound **11/36** (300 mg) was separated by chiral-HPLC (instrument: Gilson-281; column: IE 20*250, 10 μ m; mobile phase: n-hexane (0.1% DEA):EtOH (0.1% DEA) = 55:45; run time per injection: 14 min; injection: 0.4 mL; sample solution: 75 mg in 3 mL MeOH) to give as first eluting isomer (retention time: 10.28 min) compound **41/1** and as second eluting isomer (retention time: 14.35 min) compound **41/2**. NMR corresponds with Example **11/36**; MS: 669.0 (M-1).


Example 42

[0452]



Step 1: Methyl 4-(N-(4-fluoro-2-iodophenyl)sulfamoyl)benzoate (**42a**)


[0453]



[0454] To a solution of 4-fluoro-2-iodoaniline (2.00 g, 8.43 mmol) in pyridine (10 mL) was added methyl 4-(chlorosulfonyl)benzoate (2.20 g, 9.40 mmol). The mixture was stirred at rt overnight. Brine (40 mL) was added and the formed solid was filtered off, washed with EA (30 mL) and water (30 mL). The crude product was lyophilized to give compound **42a** as a white solid.

Step 2: *N*-(4-Fluoro-2-iodophenyl)-4-(2-hydroxypropan-2-yl)benzenesulfonamide (**42b**)


[0455]



[0456] To a solution of compound **42a** (3.50 g, 8.04 mmol) in THF (30 mL) was added a solution of MeMgBr (2.0M in THF, 20 mL, 40.0 mmol) at -78°C slowly during 20 min. The mixture was stirred at -78°C for 6 h before the mixture was allowed to warm to rt. Saturated aq. NH₄Cl (50 mL) was added and the resulting mixture was extracted with EA (3 × 50 mL). The combined organic layer was dried over Na₂SO₄ and concentrated in vacuo to afford compound **42b** as a white solid.

Step 3: Methyl 2-chloro-3'-((5-fluoro-2-((4-(2-hydroxypropan-2-yl)phenyl)sulfonamido)phenyl)-ethynyl)-[1,1'-biphenyl]-4-carboxylate (**42c**)


[0457]



[0458] To a solution of compound **42b** (1.30 g, 2.98 mmol) and compound **P30** (740 mg, 2.74 mmol) in dry THF (20 mL) were added CuI (23 mg, 0.12 mmol), Pd(PPh₃)₂Cl₂ (130 mg) and TEA (830 mg, 8.22 mmol). The mixture was stirred at 0°C for 30 min under argon and then stirred at rt overnight, diluted with water (30 mL) and extracted with EA (3 × 40 mL). The combined organic layer was washed by brine (2 × 50 mL), dried over Na₂SO₄, filtered, concentrated and purified by FCC (PE: EA = 5: 1) to give compound **42c** as a pale yellow solid.

Step 4: Methyl 2-chloro-3'--(5-fluoro-1-((4-(2-hydroxypropan-2-yl)phenyl)sulfonyl)-3-iodo-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (**42d**)


[0459]



[0460] To a solution of compound **42c** (500 mg, 0.86 mmol) and K₂CO₃ (368 mg, 2.67 mmol) in ACN (30 mL) was added NIS (608 mg, 2.67 mmol) at -10°C under argon. The mixture was allowed to warm to rt during 30 min and stirred overnight. The mixture was washed with aq. sat. Na₂S₂O₃ (3 × 20 mL) and extracted with DCM (2 × 20 mL). The combined organic layer was dried over Na₂SO₄, filtered, concentrated and purified by prep-HPLC to give compound **42d** as a white solid.

Step 5: Methyl 2-chloro-3'-(3-(2-cyanophenyl)-5-fluoro-1-((4-(2-hydroxypropan-2-yl)phenyl)sulfonyl)-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (**42**)


[0461]



[0462] To a solution of compound **42d** (350 mg, 0.49 mmol), (2-cyanophenyl)boronic acid (217 mg, 1.47 mmol) and K_2CO_3 (210 mg, 1.47 mmol) in a mixture of dioxane and H_2O (15 mL, 10:1) was added $Pd(dppf)Cl_2$ (45 mg) under argon. The mixture was stirred at 60°C for 4 h, cooled, quenched with water (20 mL) and extracted with EA (3 × 20 mL). The combined organic layer was washed with brine (20 mL), dried over Na_2SO_4 , filtered, concentrated and purified by prep-TLC (PE:EA = 8: 5) to give compound **42** as a yellow solid.

Example 43/1 and Example 43/2

[0463]




Separated isomers methyl (1*s*,4*s*)-4-(3-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)phenyl)cyclohexane-1-carboxylate (**43/1**) and methyl (1*r*,4*r*)-4-(3-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-fluoro-1*H*-indol-2-yl)phenyl)cyclohexane-1-carboxylate (**43/2**)

[0464] To a solution of compound **8/4** (665 mg, 1.00 mmol) in MeOH (10 mL) was added Pd/C (100 mg). The mixture was stirred at rt for 16 h under H_2 . The catalyst was filtered off and washed with MeOH (15 mL). The combined filtrates were concentrated. The residue was purified by prep-TLC (EA:PE = 1:3) to give the two separated compounds **43/1** and **43/2** as white solids, respectively.


Example 44

[0465]



Step 1: Methyl 3'-((2-amino-4-fluoro-5-methoxyphenyl)ethynyl)-2-chloro-[1,1'-biphenyl]-4-carboxylate (**44a**)


[0466]



[0467] To a solution of compound **P30** (1.39 g, 5.10 mmol) in TEA (20 mL) was added Pd(PPh₃)₄ (237 mg, 205 μmol), Cul (78 mg, 0.41 mmol), PPh₃ (108 mg, 0.41 mmol) and 2-bromo-5-fluoro-4-methoxyaniline (1.34 g, 6.12 mmol). The mixture was stirred at 60°C under N₂ overnight. The reaction was cooled, filtered, concentrated and purified by FCC (PE:EA = 1:1) to give compound **44a** as a light yellow solid.

Step 2: Methyl 2-chloro-3'-((2-((4-(difluoromethyl)phenyl)sulfonamido)-4-fluoro-5-methoxyphenyl)ethynyl)-[1,1'-biphenyl]-4-carboxylate (**44b**)


[0468]



[0469] To a solution of compound **44a** (818 mg, 2.00 mmol) in DCM (10 mL) was added 4-(difluoromethyl)benzene-1-sulfonyl chloride (542 mg, 2.40 mmol), pyridine (316 mg, 4.00 mmol) and DMAP (89 mg). The mixture was stirred at rt overnight, then DCM (20 mL) was added and the mixture was washed with 2N aq. HCl (2 × 20 mL) and brine (40 mL). The organic layer was dried over Na₂SO₄, concentrated and purified by FCC (PE:DCM = 1:1) to give compound **44b** as a white solid.

Step 3: Methyl 2-chloro-3'-((3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-6-fluoro-5-methoxy-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylate (**44c**)

[0470]




[0471] To a solution of compound **44b** (599 mg, 1.00 mmol) in dioxane (5 mL) was added 2-bromo-isophthalonitrile (310 mg, 1.50 mmol), K_2CO_3 (276 mg, 2.00 mmol) and $Pd(PPh_3)_4$ (47 mg, 40 μ mol) under N_2 . The mixture was stirred at 90°C for 4 h under N_2 . Upon completion, the mixture was cooled to rt, poured into EA (20 mL) and washed with H_2O (2×20 mL) and brine (20 mL). The organic layer was dried over Na_2SO_4 , concentrated and purified by FCC (EA:PE = 1:1) to give compound **44c** as a yellow solid.

Step 4: 2-Chloro-3'-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-6-fluoro-5-hydroxy-1*H*-indol-2-yl)-[1,1'-biphenyl]-4-carboxylic acid (**44**)

[0472] To a solution of compound **44c** (390 mg, 0.53 mmol) in CCl_4 (10 mL) was added iodotrimethylsilane (5 mL) and NaI (159 mg, 1.06 mmol) and the mixture was stirred at 85°C overnight. The solvent was removed and the residue was partitioned between sat. aq. $Na_2S_2O_3$ and EA. The aq. phase was again extracted with EA (3 x 20 mL). The combined organic layer was washed with brine, dried over Na_2SO_4 , filtered, concentrated and purified by prep-HPLC to afford compound **44** as a white solid. 1H -NMR (400 MHz, CD_3OD) δ : 8.12 (d, J = 11.7 Hz, 1H), 8.10-7.93 (m, 4H), 7.70 (t, J = 7.9 Hz, 1H), 7.58-7.29 (m, 8H), 7.00 (s, 1H), 6.83-6.48 (m, 2H). MS: 696.0 (M-1) $^-$.

Example 45

[0473]



2-Chloro-3'-(3-(2,6-dicyanophenyl)-1-((4-(difluoromethyl)phenyl)sulfonyl)-5-hydroxy-1*H*-pyrrolo[2,3-c]pyridin-2-yl)-[1,1'-biphenyl]-4-carboxylic acid (**45**)

[0474] If one were to treat a solution of compound **29/3** in ACN (10 mL) with chlorotrimethylsilane and sodium iodide under reflux one would obtain compound **45**.

[0475] If one were to follow the procedures described above using appropriate building blocks, the following compounds can be prepared:

5

10

15

20

25

30


35


40

45


50

55







5



,




,




,

10


15



,




,




,

20


25



,



,



,

30

35


40

45

50

and

55



Compound stock solutions

[0476] The tested compounds were usually dissolved, tested and stored as 20 mM stock solutions in DMSO. Since sulfonyl acetic acid derivatives tend to decarboxylate under these conditions, these stock solutions were prepared, tested and stored as 20 mM DMSO stock solutions containing 100 mM trifluoroacetic acid (5 equivalents). Sulfonyl acetic acid derivatives are shelf stable as solid at rt for long time as reported by Griesbrecht et al. (Synlett 2010:374) or Faucher et al. (J. Med. Chem. 2004;47:18).

TR-FRET β Activity Assay

[0477] Recombinant GST-LXR β ligand-binding domain (LBD; amino acids 156-461; NP009052; SEQ ID NO:4) was expressed in *E. coli* and purified via glutathione-sepharose affinity chromatography. N-terminally biotinylated NCoA3 coactivator peptide (SEQ ID NO:7) was chemically synthesized (Eurogentec). Assays were done in 384 well format (final assay volume of 25 μ L/well) in a Tris/HCl buffer (pH 6.8) containing KCl, bovine serum albumin, Triton-X-100 and 1 μ M 24(5)-25-epoxycholesterol as LXR-prestimulating agonist. Assay buffer was provided and test articles (potential LXR inverse agonists) were titrated to yield final assay concentrations of 50 μ M, 16.7 μ M, 5.6 μ M, 1.9 μ M, 0.6 μ M, 0.2 μ M, 0.07 μ M, 0.02 μ M, 0.007 μ M, 0.002 μ M with one vehicle control. Finally, a detection mix was added containing anti GST-Tb cryptate (CisBio; 610SAXLB) and Streptavidin-XL665 (CisBio; 610SAXLB) as fluorescent donor and acceptor, respectively, as well as the coactivator peptide and LXR β -LBD protein (SEQ ID NO:4). The reaction was mixed thoroughly, equilibrated for 1 h at 4°C and vicinity of LXR β and coactivator peptide was detected by measurement of fluorescence in a VictorX4 multiplate reader (PerkinElmer Life Science) using 340 nm as excitation and 615 and 665 nm as emission wavelengths. Assays were performed in triplicates.

Final assay concentrations of components:

[0478] 240 mM KCl, 1 μ g/ μ L BSA, 0.002% Triton-X-100, 125 pg/ μ L anti GST-Tb cryptate, 2.5 ng/ μ L Streptavidin-XL665, coactivator peptide (400 nM), LXR β protein (530 μ g/mL, i.e. 76 nM).

LXR Gal4 Reporter Transient Transfection Assays

[0479] LXR α and LXR β activity status was determined via detection of interaction with coactivator and corepressor proteins in mammalian two-hybrid experiments (M2H). For this, via transient transfection the full length (FL) proteins of LXR α (amino acids 1-447; NP005684; SEQ ID NO:1) or LXR β (amino acids 1-461; NP009052; SEQ ID NO:2) or the ligand-binding domains (LBD) of LXR α (amino acids 155-447 SEQ ID NO:3) or LXR β (amino acids 156-461; SEQ ID NO:4) were expressed from pCMV-AD (Stratagene) as fusions to the transcriptional activation domain of NF κ B. As cofactors, domains of either the steroid receptor coactivator 1 (SRC1; amino acids 552-887; SEQ ID NO:5) or of the corepressor NCoR (amino acids 1906-2312; NP006302; SEQ ID NO:6) were expressed as fusions to the DNA binding domain of the yeast transcription factor GAL4 (from pCMV-BD; Stratagene). Interaction was monitored via activation of a coexpressed Firefly Luciferase Reporter gene under control of a promoter containing repetitive GAL4 response elements (vector pFRLuc; Stratagene). Transfection efficiency was controlled via cotransfection of constitutively active pRL-CMV *Renilla reniformis* luciferase reporter (Promega). HEK293 cells were grown in minimum essential medium (MEM) with 2 mM L-glutamine and Earle's balanced salt solution supplemented with 8.3% fetal bovine serum, 0.1 mM non-essential amino acids, 1 mM sodium pyruvate, at 37°C in 5% CO₂. 3.5 $\times$ 10⁴ cells/well were plated in 96-well cell culture plates in growth medium supplemented with 8.3% fetal bovine serum for 16-20 h to ~90% confluency. For transfection, medium was taken off and LXR and cofactor expressing plasmids as well as the reporter plasmids are added in 30 μ L OPTIMEM/well including polyethylene-imine (PEI) as vehicle. Typical amounts of plasmids transfect-

EP 3 814 331 B9

ed/well: pCMV-AD-LXR (5 ng), pCMV-BD-cofactor (5 ng), pFR-Luc (100 ng), pRL-CMV (0.5 ng). Compound stocks were prepared in DMSO, prediluted in MEM to a total volume of 120 μ L, and added 4 h after addition of the transfection mixture (final vehicle concentration not exceeding 0.2%). Cells were incubated for additional 16 h, lysed for 10 min in 1 $\times$ Passive Lysis Buffer (Promega) and Firefly and Renilla luciferase activities were measured sequentially in the same cell extract using buffers containing D-luciferine and coelenterazine, respectively. Measurements of luminescence were done in a BMG-luminometer.

Materials	Company	Cat.No.
HEK293 cells	DSMZ	ACC305
MEM	Sigma-Aldrich	M2279
OPTIMEM	LifeTechnologies	11058-021
FCS	Sigma-Aldrich	F7542
Glutamax	Invitrogen	35050038
Pen/Strep	Sigma Aldrich	P4333
Sodium Pyruvate	Sigma Aldrich	S8636
Non Essential Amino Acids	Sigma Aldrich	M7145
Trypsin	Sigma-Aldrich	T3924
PBS	Sigma Aldrich	D8537
PEI	Sigma Aldrich	40.872-7
Passive Lysis Buffer (5x)	Promega	E1941
D-Luciferine	PJK	260150
Coelenterazine	PJK	26035

Table 1. LXR activity data

Ranges (**EC₅₀**): -: no activity measured; A: >10 μ M, B: 1 μ M to <10 μ M, C: 100 nM to <1 μ M, D: <100 nM; italic numbers indicate that efficacy (compared to GW2033) is below 40%.

Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
1/23		CD	D		
1/26	<i>B</i>	<i>C</i>	<i>D</i>		
1/27	B	C	D		
1/28	B	<i>C</i>	D		
1/39	A	<i>B</i>	-		
1/40	B	C	D		
1/41	B	<i>C</i>	<i>D</i>		
1/42	B	<i>D</i>	D		
1/122	C			D	D
1/139	C			D	D
2	C			D	D
2/1	C			C	C
2/2	C			C	C
2/16	D			D	D
2/18	C			C	D
3	D			D	D
3/1	C			C	C
3/2	B			C	D
Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
3/3	B			C	C
3/4	C			C	D
3/5	C			C	C
3/6	B			C	C
3/7	B			-	D

EP 3 814 331 B9

(continued)

	Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
5	3/8	B	C	D		
	3/9	B	C	D		
	3/10	C			C	D
	3/11	B			C	-
	3/12	C			D	D
10	3/13	B			D	D
	3/14	B			C	C
	3/15	B			-	-
	3/16	B	C	C		
	3/17	B	C	D		
15	3/18	B	C	D		
	3/19	B	C	C		
	3/20	B	C	D		
	3/21	C			D	D
	3/22	D			D	D
20	3/23	C			D	D
	3/24		D	D	D	D
	3/25	B			C	D
	3/26	B			C	D
	3/27	B			C	C
25	3/28	A			-	B
	3/29	B			-	-
	3/30	C			D	D
	3/31	C			C	C
	3/32	C			C	D
30	3/33	B			C	D
	3/34	C			C	D
	3/35	C			C	D
	3/36	B			C	C
	3/37	C			D	D
35	3/38	C			D	D
	3/39	C			C	D
	3/40	B			C	D
	3/41	B			D	D
	3/42	B	C	D	C	D
40	3/43	B	C	C	C	D
	3/44	B	D	D	D	D
	3/45	B			C	C
	3/46	C	D	D	D	D
	3/47	C			D	D
45	3/48	C			D	D
	3/49	C			D	D
	3/50	C			D	D
	3/51	B	C	C		
	3/52	C			C	C
50	3/53	D			D	D
	3/54	C			D	D
	3/55	C			D	D
	3/56	D			D	D
	3/57	C			D	D

EP 3 814 331 B9

(continued)

	Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
5	3/58	B			D	D
	3/59	D			D	D
	3/60	C			D	D
	3/61	C			C	C
	3/62	C			C	D
10	3/63	D			D	D
	3/64	C			D	D
	3/65	C			D	D
	3/66	B			C	C
	3/67	C			C	C
15	3/68	C			D	D
	3/69	C			C	C
	3/70	C			C	C
	3/71	D			D	D
	3/72	D			D	D
20	3/73	D			D	D
	4	B			D	C
	4/1	B			C	D
	4/2	B			C	C
	4/3	B			D	D
25	5	C			B	B
	5/1	C			B	C
	5/2	C			C	C
	5/3	C				D
	5/4	B			C	D
30	5/5	C			C	C
	5/6	C			D	D
	5/7	C			C	D
	5/8	C			C	C
	5/9	C			C	D
35	5/10	B			C	C
	6	C			D	D
	6/1	C			D	D
	6/2	C			D	D
	6/3	D			D	D
40	8	C			C	D
	8/1	C			D	D
	8/3	C			C	D
	8/7	C			C	D
	8/8	D			D	D
45	10	C			D	C
	10/1	D			D	D
	10/2	D			D	D
	10/3	D			D	D
	10/4	D			C	D
50	11	D			D	D
	11/1	D			D	D
	11/2	D			D	D
	11/3	D			D	D
	11/4	C			D	D

EP 3 814 331 B9

(continued)

	Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
5	11/5	D			D	D
	11/6	D			D	D
	11/7	D			D	D
	11/8	D			D	D
	11/9	D			D	D
10	11/10	D			D	D
	11/11	D			D	D
	11/12	D			D	D
	11/13	D			D	D
	11/14	D			D	D
15	11/15	D			D	D
	11/16	C			D	D
	11/17	D			D	D
	11/18	C			D	D
	11/19	C			D	D
20	11/20	C			D	D
	11/21	D			D	D
	11/22	D			D	D
	11/23	D			D	D
	11/24	C			C	C
25	11/25	C			C	D
	11/26	D			D	D
	11/27	D			D	D
	11/28	D			D	D
	11/29	D			D	D
30	11/30	D			D	D
	11/31	D			D	D
	11/32	D			D	D
	11/33	D			D	D
	11/34	D			D	D
35	11/35	D			D	D
	11/36	D			D	D
	11/37	D			D	D
	11/38	D			D	D
	C11/39	C			D	D
40	11/40	C			D	D
	11/41	D			D	D
	11/42	A			C	C
	11/43	C			C	C
	11/44	D			D	D
45	11/45	D			D	D
	11/46	D			D	D
	11/47	C			C	C
	C11/48	D			D	D
	11/49	D			D	D
50	11/50	D			D	D
	11/51	D			D	D
	11/52	D			D	D
	11/53	D			D	D
	11/54	D			D	D

EP 3 814 331 B9

(continued)

	Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
5	11/55	D			D	D
	11/56	D			D	D
	11/57	D			D	D
	11/58	D			D	D
	11/59	D			D	D
10	11/60				D	D
	11/61	C			D	D
	11/62	C			D	D
	11/63	C			D	D
	11/64	C			D	D
15	11/65	D			D	D
	11/66	C			D	D
	11/67	C			D	D
	11/68	D			D	D
	11/69	D			D	D
20	12	C			D	D
	12/1	C			D	D
	12/2	C			D	D
	12/3	C			D	D
	12/4	C			D	D
25	12/5	D			D	D
	12/6	B			C	B
	12/7	C			C	C
	12/8	B			C	C
	12/9	C			C	C
30	12/10	D			D	D
	13	C	D	D		
	13/1	B	D	D		
	15	A	B	A		
	15/1	B	C	C		
35	15/2	A	C	C		
	15/3	A	-	-		
	15/4	B	C	D		
	15/5	B	D	D		
	15/6	B	-	C		
40	17	A	-	B		
	19	A	-	-		
	20	C	D	D		
	20/2	B	C	D		
	20/3	C	D	D		
45	20/4	B	D	D		
	20/5	B	C	C		
	20/6	C	C	D		
	20/7	B	C	D		
	20/11	C	D	D		
50	20/12	B	D	D		
	20/13	C	D	D		
	20/14	C	D	D		
	20/15	C	-	C		
	20/16	B	C	C		

EP 3 814 331 B9

(continued)

	Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
5	20/17	-	-	C		
	20/18	-	C	D		
	20/19	A	C	C		
	20/20	B	B	C		
	20/21	B	C	D		
10	20/22	C	C	C		
	20/23	B	C	C		
	21	A	-	B		
	21/1	B	B	B		
	21/2	B	B	B		
15	21/3	D			D	D
	22	B	C	C		
	23	A	C	C		
	23/1	B	-	D		
	23/2	C			D	D
20	23/3	C			D	D
	24	B	C	D		
	25	B	C	D		
	26	C	D	D		
	26/1	B	C	D		
25	27	B	-	C		
	27/1	B	-	-		
	27/2	B	-	-		
	27/3	A	-	-		
	28	B	B	C		
30	30	C	D	D		
	30/1	C	C	D		
	30/2	B	C	C		
	30/3		C	C		
	30/4	B	C	D		
35	30/5	B	C	D		
	30/6	B	C	C		
	30/7	C	D	D		
	30/8	B	C	C		
	30/9	B	C	D		
40	30/10	A	-	-		
	30/11	C	C	D		
	30/12	B	B	C		
	30/13	C	C	D		
	Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
50	30/14	B	C	C		
	30/15	B	C	C		
	30/16	B			C	C
	31	B	C	C		
	32	C	C	C		
55	32/1	C	C	D		
	32/2	C			C	C
	32/4	D			D	D
	32/5	D			D	D

(continued)

	Ex. #	FRET β	LBD-M2H Gal4 α	LBD-M2H Gal4 β	FL-M2H Gal4 α	FL-M2H Gal4 β
5	36	B			C	C
	36/1	C			D	D
	37	C			D	D
	37/1	C			C	C
	41/1	D			D	D
10	41/2	D			D	D
	44	C			D	D

Pharmacokinetics

15 **[0480]** The pharmacokinetics of the compounds was assessed in mice after single dosing and oral administrations. Blood/plasma and liver exposure was measured via LC-MS.

[0481] The study design was as follows:

Animals: C57/bl6/J (Janvier) males

Diet: standard rodent chow

Dose: 20 mg/kg

Animal handling: animals were withdrawn from food at least 12 h before administration

Design: single dose oral administration, n = 3 animals per group

Sacrifice: at stated time point (4, 12 or 24 h) after administration

Bioanalytics: LC-MS of liver and blood/plasma samples

Table 2. Study results

Ranges:

blood/plasma exposure: **A**: >1 μ M, **B**: 300 nM to $\leq$ 1 μ M, **C**: 100 nM to <300 nM, **D**: <100 nM;

liver exposure: **A**: <300 nM, **B**: 300 nM to $\leq$ 1 μ M, **C**: 1 μ M to $\leq$ 3 μ M, **D**: >3 μ M;

liver/plasma ratio: **A**: <3, **B**: 3 to $\leq$ 10, **C**: 10 to $\leq$ 30, **D**: >30;

	Example #	time point (h)	blood/plasma exposure	liver exposure	liver/blood ratio
40	GSK2033 (comparative example)	4	below LLOQ (14.4 ng/mL)	below LLOQ (9.6 ng/mL)	-
	SR9238 (comparative example)	4	below LLOQ	below LLOQ	-
45	3/24	4	D	C	D
	3/48	12	below LLOQ (1.2 ng/mL)	A	-
	5/3	4	C	C	C
	8	4	B	D	B
	23/2	4	B	D	C
50	30/4	4	C	C	C
	30/7	4	D	B	D

[0482] We confirmed that structurally unrelated LXR inverse agonists **GSK2033** and **SR9238** are not orally bioavailable. We found, that compounds from the present invention are orally bioavailable and the target tissue liver was effectively reached by such compounds and a systemic exposure, which is not desired, could be minimized.

Short term HFD mouse model:

[0483] The *in vivo* transcriptional regulation of several LXR target genes by LXR modulators was assessed in mice.

[0484] For this, C57BL/6J were purchased from Elevage Janvier (Rennes, France) at the age of 8 weeks. After an acclimation period of two weeks, animals were prefed on a high fat diet (HFD) (Ssniff Spezialdiäten GmbH, Germany, Surwit EF D12330 mod, Cat. No. E15771-34), with 60 kcal% from fat plus 1% (w/w) extra cholesterol (Sigma-Aldrich, St. Louis, MO) for 5 days. Animals were maintained on this diet during treatment with LXR modulators. The test compounds were formulated in 0.5% hydroxypropylmethylcellulose (HPMC) and administered in three doses (from 1.5 to 20 mg/kg each) by oral gavage according to the following schedule: on day one, animals received treatment in the morning and the evening (ca. 17:00), on day two animals received the final treatment in the morning after a 4 h fast and were sacrificed 4 h thereafter. Animal work was conducted according to the national guidelines for animal care in Germany.

[0485] Upon termination, liver was collected, dipped in ice cold PBS for 30 seconds and cut into appropriate pieces. Pieces were snap frozen in liquid nitrogen and stored at -80°C. For the clinical chemistry analysis from plasma, alanine aminotransferase (ALT, IU/mL), cholesterol (CHOL, mg/dL) and triglycerides (TG, mg/dL) were determined using a fully-automated bench top analyzer (Respos®910, DiaSys Greiner GmbH, Flacht, Germany) with system kits provided by the manufacturer.

[0486] Analysis of gene expression in liver tissue. To obtain total RNA from frozen liver tissue, samples (25 mg liver tissue) were first homogenized with RLA buffer (4M guanidin thiocyanate, 10 mM Tris, 0.97% w:v β-mercapto-ethanol). RNA was prepared using a SV 96 total RNA Isolation system (Promega, Madison, Wisconsin, USA) following the manufacturer's instructions. cDNAs were synthesized from 0.8-1 µg of total RNA using All-in-One cDNA Supermix reverse transcriptase (Absource Diagnostics, Munich, Germany). Quantitative PCR was performed and analyzed using Prime time Gene expression master mix (integrated DNA Technologies, Coralville, Iowa, USA) and a 384-format ABI 7900HT Sequence Detection System (Applied Biosystems, Foster City, USA). The expression of the following genes was analysed: Stearoyl-CoA desaturase1 (*Scd1*), fatty acid synthase (*Fas*) and sterol regulatory element-binding protein1 (*Srebp1c*). Specific primer and probe sequences (commercially available) are listed in Table 3. qPCR was conducted at 95°C for 3 min, followed by 40 cycles of 95°C for 15 s and 60°C for 30 s. All samples were run in duplicates from the same RT-reaction. Gene expression was expressed in arbitrary units and normalized relative to the mRNA of the house-keeping gene TATA box binding protein (*Tbp*) using the comparative Ct method.

Table 3. Primers used for quantitative PCR

Gene	Forward Primer	Reverse Primer	Sequence Probe
<i>Fasn</i>	CCCCTCTGTTAATTGGC TCC (SEQ ID NO:8)	TTGTGGAAGTGCAGGT TAGG (SEQ ID NO:9)	CAGGCTCAGGGTGTCCC ATGTT (SEQ ID NO:10)
	CTGACCTGAAAGCCGA GAAG (SEQ ID NO:11)	AGAAGGTGCTAACGAA CAGG (SEQ ID NO:12)	TGTTTACAAAAGTCTCGC CCCAGCA (SEQ ID NO:13)
<i>Srebp1c</i>	CCATCGACTACATCCGC TTC (SEQ ID NO:14)	GCCCTCCATAGACACA TCTG (SEQ ID NO:15)	TCTCCTGCTTGAGCTTCT GGTTGC (SEQ ID NO:16)
<i>Tbp</i>	CACCAATGACTCCTATG ACCC (SEQ ID NO:17)	CAAGTTTACAGCCAAG ATTACAG (SEQ ID NO:18)	ACTCCTGCCACACCAGC CTC (SEQ ID NO:19)

Table 4. Study results

Ranges:

plasma **A:** >1 µM, **B:** 300 nM to
exposure: ≤ 1 µM, **C:** 100 nM to <300
 nM, **D:** <100 nM;

EP 3 814 331 B9

(continued)

Ranges:

liver **A:** <300 nM, **B:** 300 nM to

exposure: $\leq 1 \mu\text{M}$, **C:** $1 \mu\text{M}$ to ≤ 10

μM , **D:** $>10 \mu\text{M}$;

liver/plasma **A:** <5, **B:** 5 to ≤ 30 , **C:** 30

ratio: to ≤ 100 , **D:** >100 ;

gene suppression: **A:** >0.9 , **B:** 0.6 to ≤ 0.9 , **C:** 0.3 to ≤ 0.6 , **D:** <0.3 ;

	Example #	dose [mg/kg]	plasma exposure 4 h	liver exposure 4 h	liver/plasma ratio 4 h
	3/48	20	D	C	C
	3/59	20	C	C	B
	3/64	20	C	D	D
15	3/73	10	D	B	C
	5/3	20	D	C	B
	8/8	10	C	B	B
	10/1	10	D	B	C
	10/2	10	D	D	B
20	11/11	20	D	D	D
	11/12	20	D	C	C
	11/13	20	A	C	A
	11/16	20	C	C	B
25	11/17	20	C	D	C
	11/23	20	C	C	B
	11/26	10	C	C	C
	11/27	20	D	C	C
	11/33	20	C	C	C
30	11/37	10	D	C	C
	11/49	20	D	C	D
	11/51	10	D	C	C
	11/53	10	D	C	C
35	11/62	10	C	C	B
	11/63	10	D	C	B
	11/65	10	D	C	C
	21/3	10	D	C	D
	23/2	20	C	A	A
40	32/5	10	D	B	B
	Example #		Fasn suppression compared to vehicle	Srebp1c suppr. compared to vehicle	Scd1 suppression compared to vehicle
45	3/48	20	D	D	D
	3/59	20	C	D	C
	3/64	20	B	B	D
	3/73	10	A	D	D
	5/3	20	D	C	C
50	8/8	10	D	D	D
	10/1	10	D	D	D
	10/2	10	C	C	C
	11/11	20	C	D	D
55	11/12	20	C	D	D
	11/13	20	C	D	D
	11/16	20	A	B	C
	11/17	20	C	D	D

(continued)

Example #		Fasn suppression compared to vehicle	<i>Srebp1c</i> suppr. compared to vehicle	<i>Scd1</i> suppression compared to vehicle
5	11/23	20	C	D
	11/26	10	D	C
	11/27	20	C	D
	11/33	20	B	D
10	11/37	10	D	D
	11/49	20	C	D
	11/51	10	D	D
	11/53	10	D	D
15	11/62	10	D	D
	11/63	10	C	C
	11/65	10	D	D
	21/3	10	C	D
	23/2	20	C	D
20	32/5	10	C	C

[0487] Triple oral dosing over two days (day one morning and evening, day two morning) of compounds from the present invention in mice lead to a high liver exposure with a favourable liver-to-plasma ratio. Hepatic LXR target genes were effectively suppressed. These genes are involved in the transcriptional regulation of hepatic *de-novo* lipogenesis (Wang et al., Nat. Rev. Mol. Cell Biol. 2015;16:678). A suppression of these genes will reduce liver fat (liver triglycerides).

Comparative Examples

[0488] The Comparative Examples illustrate that it can be advantageous, when the cyclic moiety in 3-position of the indole (or analog) has at least one substituent in 1,2-orientation (*ortho*-substituent).

5

10

15

20

25

30


35

40

45

50


55



Comparative


Example C3/29

FRETβ	4228 nM (~102%)
FL-M2H Gal4α	inactive
FL-M2H Gal4β	inactive



Example 3/32


FRETβ	551 nM (~98%)
LBD-M2H Gal4α	106 nM (103%)
LBD-M2H Gal4β	13 nM (81%)



Comparative

Example C1/39


FRETβ	104 nM (~91%)
FL-M2H Gal4α	14 nM (117%)
FL-M2H Gal4β	14 nM (140%)



Comparative

Example C1/48

FRETβ	49 nM (~96%)
FL-M2H Gal4α	62 nM (118%)
FL-M2H Gal4p	32 nM (123%)




Example 11/33

FRETβ	19 nM (~99%)
FL-M2H Gal4α	1.3 nM (164%)
FL-M2H Gal4β	1.7 nM (130%)

Claims

1. A compound represented by Formula (I)



a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein



is an annelated 5- to 6-membered cycle forming a 6-membered aryl or a 5- to 6-membered heteroaryl containing 1 to 3 heteroatoms independently selected from N, O and S, wherein this cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, C₁₋₆-alkyl, oxo, C₀₋₆-alkylene-OR¹¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR¹¹, C₀₋₆-alkylene-NR¹¹S(O)₂R¹¹, C₀₋₆-alkylene-S(O)₂NR¹¹R¹², C₀₋₆-alkylene-NR¹¹S(O)₂NR¹¹R¹², C₀₋₆-alkylene-CO₂R¹¹, O-C₁₋₆-alkylene-CO₂R¹¹, C₀₋₆-alkylene-O-COR¹¹, C₀₋₆-alkylene-CONR¹¹R¹², C₀₋₆-alkylene-NR¹¹-COR¹¹, C₀₋₆-alkylene-NR¹¹-CONR¹¹R¹², C₀₋₆-alkylene-O-CONR¹¹R¹², C₀₋₆-alkylene-NR¹¹-CO₂R¹¹ and C₀₋₆-alkylene-NR¹¹R¹², wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein the new formed cycle is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;



is selected from the group consisting of 3- to 10-membered cycloalkyl, 3- to 10-membered heterocycloalkyl containing 1 to 3 heteroatoms independently selected from N, O and S, 6- to 14-membered aryl and 5- to 14-membered heteroaryl containing 1 to 4 heteroatoms independently selected from N, O and S, wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR²¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR²¹, C₀₋₆-alkylene-NR²¹S(O)₂R²¹, C₀₋₆-alkylene-S(O)₂NR²¹R²², C₀₋₆-alkylene-NR²¹S(O)₂NR²¹R²², C₀₋₆-alkylene-CO₂R²¹, O-C₁₋₆-alkylene-CO₂R²¹, C₀₋₆-alkylene-O-COR²¹, C₀₋₆-alkylene-CONR²¹R²², C₀₋₆-alkylene-NR²¹-COR²¹, C₀₋₆-alkylene-NR²¹-CONR²¹R²², C₀₋₆-alkylene-O-CONR²¹R²², C₀₋₆-alkylene-NR²¹-CO₂R²¹ and C₀₋₆-alkylene-NR²¹R²², wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl, and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl,
 and wherein optionally two adjacent substituents on the cycloalkyl or heterocycloalkyl moiety form a 5- to 6-membered unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N,
 wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;



is selected from the group consisting of 6- or 10-membered aryl and 5- to 10-membered heteroaryl containing 1 to 3 heteroatoms independently selected from N, O and S,

wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR³¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-(6-membered aryl), C₀₋₆-alkylene-(5- to 6-membered heteroaryl), C₀₋₆-alkylene-S(O)_nR³¹, C₀₋₆-alkylene-NR³¹S(O)₂R³¹, C₀₋₆-alkylene-S(O)₂NR³¹R³², C₀₋₆-alkylene-NR³¹S(O)₂NR³¹R³², C₀₋₆-alkylene-CO₂R³¹, O-C₁₋₆-alkylene-CO₂R³¹, C₀₋₆-alkylene-O-COR³¹, C₀₋₆-alkylene-CONR³¹R³², C₀₋₆-alkylene-NR³¹-COR³¹, C₀₋₆-alkylene-NR³¹-CONR³¹R³², C₀₋₆-alkylene-O-CONR³¹R³², C₀₋₆-alkylene-NR³¹-CO₂R³¹ and C₀₋₆-alkylene-NR³¹R³²,
 wherein alkyl, alkylene, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;



is selected from the group consisting of 3- to 10-membered cycloalkyl, 3- to 10-membered heterocycloalkyl containing 1 to 3 heteroatoms independently selected from N, O and S, 6- to 14-membered aryl and 5- to 14-membered heteroaryl containing 1 to 4 heteroatoms independently selected from N, O and S,

wherein cycloalkyl, heterocycloalkyl, aryl and heteroaryl are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR²¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR²¹, C₀₋₆-alkylene-NR²¹S(O)₂R²¹, C₀₋₆-alkylene-S(O)₂NR²¹R²², C₀₋₆-alkylene-NR²¹S(O)₂NR²¹R²², C₀₋₆-alkylene-CR⁴¹(=N-OR⁴¹), C₀₋₆-alkylene-CO₂R²¹, O-C₁₋₆-alkylene-CO₂R²¹, C₀₋₆-alkylene-O-COR²¹, C₀₋₆-alkylene-CONR²¹R²², C₀₋₆-alkylene-NR²¹-COR²¹, C₀₋₆-alkylene-NR²¹-CONR²¹R²², C₀₋₆-alkylene-O-CONR²¹R²², C₀₋₆-alkylene-NR²¹-CO₂R²¹ and C₀₋₆-alkylene-NR²¹R²²,

wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, CO-OC₁₋₄-alkyl, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the aryl or heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

and wherein optionally two adjacent substituents on the cycloalkyl or heterocycloalkyl moiety form a


5- to 6-membered unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N,

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

wherein



has a substituent from above in 1,2-orientation regarding to the connection towards



or has an annelated additional cycle in 1,2-orientation;

L is selected from the group consisting of a bond, C₁₋₆-alkylene, C₂₋₆-alkenylene, C₂₋₆-alkynylene, 3- to 10-membered cycloalkylene, 3- to 10-membered heterocycloalkylene containing 1 to 4 heteroatoms independently selected from N, O and S, 6- or 10-membered arylene and 5- to 10-membered heteroarylene containing 1 to 4 heteroatoms independently selected from N, O and S,

wherein alkylene, alkenylene, alkynylene, cycloalkylene, heterocycloalkylene, arylene and heteroarylene are unsubstituted or substituted with 1 to 6 substituents independently selected from the group consisting of halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR⁴¹, C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), C₀₋₆-alkylene-S(O)_nR⁴¹, C₀₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, C₀₋₆-alkylene-S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-CO₂R⁴¹, O-C₁₋₆-alkylene-CO₂R⁴¹, C₀₋₆-alkylene-O-COR⁴¹, C₀₋₆-alkylene-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-COR⁴¹, C₀₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-alkylene-O-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-CO₂R⁴¹ and C₀₋₆-alkylene-NR⁴¹R⁴²,

wherein alkyl, alkylene, cycloalkyl and heterocycloalkyl is unsubstituted or substituted with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and wherein optionally two adjacent substituents on the arylene and heteroarylene moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

R¹ is selected from the group consisting of H, halogen, CN, SF₅, NO₂, oxo, C₁₋₄-alkyl, C₀₋₆-alkylene-OR⁴¹, Y-C₀₋₆-alkylene-(3- to 6-membered cycloalkyl), Y-C₀₋₆-alkylene-(3- to 6-membered heterocycloalkyl), Y-C₀₋₆-alkylene-(6-membered aryl), Y-C₀₋₆-alkylene-(5- to 6-membered heteroaryl), C₀₋₆-alkylene-S(O)_n(-R⁴¹)=N-R⁷⁵, X-C₁₋₆-alkylene-S(O)(-R⁴¹)=N-R⁷⁵, Co-s-alkyleneS(O)_nR⁴¹, X-C₁₋₆-alkylene-S(O)_nR⁴¹, C₀₋₆-alkylene-S(=NR⁷¹)R⁴¹, X-C₁₋₆-alkylene-S(=NR⁷¹)R⁴¹, C₀₋₆-alkylene-S(O)(=NR⁷¹)R⁴¹, X-C₁₋₆-alkylene-S(O)(=NR⁷¹)R⁴¹, C₀₋₆-alkylene-S(=NR⁷¹)₂R⁴¹, X-C₁₋₆-alkylene-S(=NR⁷¹)₂R⁴¹, C₀₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, X-C₁₋₆-alkylene-NR⁴¹S(O)₂R⁴¹, Co-s-alkylene-S(O)₂NR⁴¹R⁴², X-C₁₋₆-alkylene-S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-alkylene-SO₃R⁴¹, X-C₁₋₆-alkylene-SO₃R⁴¹, C₀₋₆-alkylene-CO₂R⁴¹, X-C₁₋₆-alkylene-CO₂R⁴¹, C₀₋₆-alkylene-O-COR⁴¹, X-C₁₋₆-alkylene-O-COR⁴¹, C₀₋₆-alkylene-CONR⁴¹R⁴², X-C₁₋₆-alkylene-CONR⁴¹R⁴², C₀₋₆-alkylene-CONR⁴¹OR⁴¹, X-C₁₋₆-alkylene-CONR⁴¹OR⁴¹, C₀₋₆-alkylene-CONR⁴¹SO₂R⁴¹, X-C₁₋₆-alkylene-CONR⁴¹SO₂R⁴¹, C₀₋₆-alkylene-NR⁴¹-COR⁴¹, X-C₁₋₆-C₀₋₆-alkylene-NR⁴¹-COR⁴¹, C₀₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-alkylene-O-CONR⁴¹R⁴², X-C₁₋₆-alkylene-O-CONR⁴¹R⁴², C₀₋₆-alkylene-NR⁴¹-CO₂R⁴¹, X-C₁₋₆-alkylene-NR⁴¹-CO₂R⁴¹, C₀₋₆-alkylene-NR⁴¹R⁴², X-C₁₋₆-alkylene-NR⁴¹R⁴²,

wherein alkyl, alkylene, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted

with 1 to 6 substituents independently selected from halogen, CN, oxo, hydroxy, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and wherein optionally two adjacent substituents on the aryl and heteroaryl moiety form a 5- to 8-membered partially unsaturated cycle optionally containing 1 to 3 heteroatoms independently selected from O, S or N, and

wherein this additional cycle is unsubstituted or substituted with 1 to 4 substituents independently selected from halogen, CN, oxo, OH, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

R¹¹, R¹², R²¹, R²², R³¹, R³², R⁴¹, R⁴², R⁵¹ are independently selected from H and C₁₋₄-alkyl, wherein alkyl is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

or R¹¹ and R¹², R²¹ and R²², R³¹ and R³², R⁴¹ and R⁴², respectively, when taken together with the nitrogen to which they are attached complete a 3- to 6-membered ring containing carbon atoms and optionally containing 1 or 2 heteroatoms independently selected from O, S or N; and

wherein the new formed cycle is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

R⁷¹ is independently selected from H, CN; NO₂, C₁₋₄-alkyl and C(O)-OC₁₋₄-alkyl,

wherein alkyl is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, halo-(3- to 6-membered cycloalkyl), 3- to 6-membered heterocycloalkyl, halo-(3- to 6-membered heterocycloalkyl), OH, oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl;

R⁷⁵ is independently selected from C₁₋₄-alkyl, 3- to 6-membered cycloalkyl, 3- to 6-membered heterocycloalkyl, 6-membered aryl and 5- to 6-membered heteroaryl,


wherein alkyl, cycloalkyl, heterocycloalkyl, aryl and heteroaryl is unsubstituted or substituted with 1 to 3 substituents independently selected from halogen, CN, Me, Et, CHF₂, CF₃, OH, oxo, CO₂H, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, OMe, OEt, OCHF₂, and OCF₃;

X is independently selected from O, NR⁵¹, S(O)_n, S(=NR⁷¹), S(O)(=NR⁷¹) and S(=NR⁷¹)₂;


Y is independently selected from a bond, O, NR⁵¹, S(O)_n, S(=NR⁷¹), S(O)(=NR⁷¹) and S(=NR⁷¹)₂;

n is independently selected from 0 to 2;


and with the proviso, that the following structures are excluded:



2. The compound according to claim 1, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, N-oxide, solvate or pharmaceutically acceptable salt thereof, wherein



is selected from



wherein



is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of F, Cl, Br, CN, OH, oxo, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl, O-halo-C₁₋₄-alkyl, NH₂, NHC₁₋₄-alkyl, N(C₁₋₄-alkyl)₂, SO₂-C₁₋₄-alkyl and SO₂-halo-C₁₋₄-alkyl.

3. The compound according to claim 1 or 2, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein




is selected from the group consisting of phenyl, naphthyl, pyridyl, pyrimidinyl, thiophenyl, thiazolyl, cyclopentyl, cyclohexyl, bicyclo[1.1.1]pentyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octyl and piperidinyl,

wherein the cycle is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of F, Cl, Br, CN, OH, oxo, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl, O-halo-C₁₋₄-alkyl, C₁₋₄-alkyl-OH and halo-C₁₋₄-alkyl-OH; and wherein optionally two adjacent substituents on the phenyl ring form together a -(CH₂)₃-, -(CH₂)₄-, -OCF₂O- and -OCH₂O- group.

4. The compound according to any one of claims 1 to 3, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein


⊙ is selected from phenyl, pyridyl and thiophenyl; wherein phenyl, pyridyl and thiophenyl is unsubstituted or substituted with 1 to 3 substituents independently selected from the group consisting of F, Cl, CN, OH, oxo, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl and O-halo-C₁₋₄-alkyl; and wherein residue -L-R¹ is linked in 1,3-orientation regarding the connection towards



and L is not a bond.

5. The compound according to any one of claims 1 to 4, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein

- L-R¹ is selected from






15

20



55




and optionally the glycine and tauro conjugate thereof, or enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

8. The compound according to any one of claims 1 to 7, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein


(D)

is selected from the group consisting of




wherein

R² is selected from Me, F, Cl, CN, Me, CHO, CHF₂, CF₃, SO₂Me,



and



and

wherein


(D)

is optionally further substituted with 1 to 2 substituents selected from the group consisting F, Cl, CN, Me, OMe, CHO, CHF₂ and CF₃.

9. The compound according to any one of claims 1 to 8, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein

(D)

is selected from the group consisting of



25 **10.** The compound according to any one of claims 1 to 9 wherein Formula (I) contains a substituent selected from the group consisting of CO₂H, tetrazole, CONHSO₂Me and CONH(OH); and optionally the glycine and tauro conjugate thereof, or enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

11. The compound according to any one of claims 1 to 10, wherein

-L-R¹ is selected from



40 wherein phenyl is unsubstituted or substituted with 1 to 4 substituents independently selected from the group consisting of F, Cl, CN, OH, Me and OMe; and

R¹ is selected from CO₂H and C(OH)MeCO₂H; and optionally the glycine and tauro conjugate thereof, or enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

45 12. The compound according to any one of claims 1 to 10, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein L-R¹ is



wherein the cycle is unsubstituted or further substituted with 1 to 4 substituents independently selected from the group consisting of F, Cl, Br, CN, OH, oxo, C₁₋₄-alkyl, halo-C₁₋₄-alkyl, O-C₁₋₄-alkyl, O-halo-C₁₋₄-alkyl, C₁₋₄-alkyl-OH, halo-C₁₋₄-alkyl-OH, SO₂-C₁₋₄-alkyl and SO₂-halo-C₁₋₄-alkyl; and wherein optionally two adjacent substituents

on the phenyl ring form together a $-(CH_2)_3-$, $-(CH_2)_4-$, $-OCF_2O-$ and $-OCH_2O-$ group.


13. The compound according to any one of claims 1 to 12, wherein R^1 is C_{0-6} -alkylene- CO_2R^{41} or C_{0-6} -alkylene- $CONR^{41}R^{42}$, or a glycine conjugate or tauro conjugate thereof, or enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

14. The compound according to any one of claims 1 to 13, wherein R^1 is $COOH$, or a glycine conjugate or tauro conjugate thereof, or enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

15. The compound according to any one of claims 1 to 13, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein R^1 is C_{0-6} -alkylene- $CONR^{41}R^{42}$.

16. The compound according to claim 15, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, wherein R^{41} and R^{42} are independently selected from H and C_{1-4} -alkyl, wherein C_{1-4} -alkyl is unsubstituted or substituted with CO_2H .




17. The compound according to any one of claims 1 to 12, wherein $L-R^1$ is





















or a glycine conjugate or tauro conjugate thereof, or enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

18. The compound according to any one of claims 1 to 17, where the compound is a glycine conjugate, or enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.


19. The compound according to any one of claims 1 to 18 selected from








and




or a glycine conjugate or tauro conjugate thereof; and
an enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

20. The compound according to any one of claims 1 to 19, which is



an enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.

21. The compound according to any one of claims 1 to 19, which is



or a glycine conjugate thereof, an enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof.


22. A compound according to any one of claims 1 to 21, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, for use as a medicament.

23. A compound according to any one of claims 1 to 21, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, for use in the prophylaxis and/or treatment of diseases amenable for treatment with LXR modulators, wherein the disease is selected from non-alcoholic fatty liver disease, non-alcoholic steatohepatitis, liver inflammation, liver fibrosis, obesity, insulin resistance, type II diabetes, familial hypercholesterolemia, hypercholesterolemia in nephrotic syndrome, metabolic syndrome, cardiac steatosis, cancer, viral myocarditis, hepatitis C virus infection or its complications, and unwanted side-effects of long-term glucocorticoid treatment in diseases such as rheumatoid arthritis, inflammatory bowel disease and asthma.

24. A pharmaceutical composition comprising a compound according to any one of claims 1 to 21, or a glycine conjugate, tauro conjugate, enantiomer, diastereomer, tautomer, *N*-oxide, solvate or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient.

Patentansprüche

1. Verbindung, dargestellt durch die Formel (I)



ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon, wobei



ein anellierter 5- bis 6-gliedriger Ring ist, der ein 6-gliedriges Aryl oder ein 5- bis 6-gliedriges Heteroaryl bildet, das 1 bis 3 Heteroatome enthält, die unabhängig voneinander aus N, O und S ausgewählt sind, wobei dieser Ring unsubstituiert oder mit 1 bis 4 Substituenten unabhängig ausgewählt aus der Gruppe bestehend aus Halogen, CN, SF₅, NO₂, C₁₋₆-Alkyl, Oxo, C₀₋₆-Alkylen-OR¹¹, C₀₋₆-Alkylen-(3- bis 6-gliedriges Cycloalkyl), C₀₋₆-Alkylene-(3- bis 6-gliedriges Heterocycloalkyl), C₀₋₆-Alkylen-S(O)_nR¹¹, C₀₋₆-Alkylen-NR¹¹S(O)₂R¹¹, C₀₋₆-Alkylen-S(O)₂NR¹¹R¹², C₀₋₆-Alkylen-NR¹¹S(O)₂NR¹¹R¹², C₀₋₆-Alkylen-CO₂R¹¹, O-C₁₋₆-Alkylen-CO₂R¹¹, C₀₋₆-Alkylen-O-COR¹¹, C₀₋₆-Alkylen-CONR¹¹R¹², C₀₋₆-Alkylen-NR¹¹-COR¹¹, C₀₋₆-Alkylen-NR¹¹-CONR¹¹R¹², C₀₋₆-Alkylen-O-CONR¹¹R¹², C₀₋₆-Alkylen-NR¹¹-CO₂R¹¹ und C₀₋₆-Alkylen-NR¹¹R¹² substituiert ist, wobei Alkyl, Alkylen, Cycloalkyl und Heterocycloalkyl unsubstituiert oder mit 1 bis 6 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus Halogen, CN, Oxo, Hydroxy, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl; und

wobei gegebenenfalls zwei benachbarte Substituenten an der Aryl- oder Heteroaryleinheit einen 5- bis 8-gliedrigen teilweise ungesättigten Ring bilden, der gegebenenfalls 1 bis 3 Heteroatome enthält, die unabhängig voneinander aus O, S oder N ausgewählt sind, und

wobei der neu gebildete Ring unsubstituiert oder mit 1 bis 3 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus Halogen, CN, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, 3- bis 6-gliedrigem Cycloalkyl, Halogen-(3- bis 6-gliedriges Cycloalkyl), 3- bis 6-gliedriges Heterocycloalkyl, Halogen-(3- bis 6-gliedriges Heterocycloalkyl), OH, Oxo, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, O-C₁₋₄-Alkyl und O-Halo-C₁₋₄-Alkyl;



ausgewählt ist aus der Gruppe bestehend aus 3- bis 10-gliedrigem Cycloalkyl, 3- bis 10-gliedrigem Heterocycloalkyl, das 1 bis 3 Heteroatome enthält, die unabhängig ausgewählt sind aus N, O und S, 6- bis 14-gliedrigem Aryl und 5- bis 14-gliedrigem Heteroaryl, das 1 bis 4 Heteroatome enthält, die unabhängig ausgewählt sind aus N, O und S,

wobei Cycloalkyl, Heterocycloalkyl, Aryl und Heteroaryl unsubstituiert oder mit 1 bis 6 Substituenten substituiert sind, die unabhängig voneinander ausgewählt sind aus der Gruppe bestehend aus Halogen CN, SF₅, NO₂, Oxo, C₁₋₄-Alkyl, C₀₋₆-Alkylene-OR²¹, C₀₋₆-Alkylene-(3-bis 6-gliedrigem Cycloalkyl), C₀₋₆-Alkylen-(3- bis 6-gliedrigem Heterocycloalkyl), C₀₋₆-Alkylen-S(O)_nR²¹, C₀₋₆-Alkylen-NR²¹S(O)₂R²¹, C₀₋₆-Alkylen-S(O)₂NR²¹R²², C₀₋₆-Alkylen-NR²¹S(O)₂NR²¹R²², C₀₋₆-Alkylen-CO₂R²¹, O-C₁₋₆-Alkylen-CO₂R²¹, C₀₋₆-Alkylen-O-COR²¹, C₀₋₆-Alkylen-CONR²¹R²², C₀₋₆-Alkylen-NR²¹-COR²¹, C₀₋₆-Alkylen-NR²¹-CONR²¹R²², C₀₋₆-Alkylen-O-CONR²¹R²², C₀₋₆-Alkylen-NR²¹-CO₂R²¹ und C₀₋₆-Alkylen-NR²¹R²²,

wobei Alkyl, Alkylen, Cycloalkyl und Heterocycloalkyl unsubstituiert oder mit 1 bis 6 Substituenten substituiert ist, die unabhängig voneinander aus Halogen, CN, Oxo, Hydroxy, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl ausgewählt sind, und wobei gegebenenfalls zwei benachbarte Substituenten an der Aryl- oder Heteroaryleinheit einen 5- bis 8-gliedrigen teilweise ungesättigten Ring bilden, der gegebenenfalls 1 bis 3 Heteroatome enthält, die unabhängig voneinander aus O, S oder N ausgewählt sind, und

wobei dieser zusätzliche Ring unsubstituiert oder mit 1 bis 4 Substituenten substituiert ist, die unabhängig

ausgewählt sind aus Halogen, CN, Oxo, OH, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-alkyl,



ausgewählt ist aus der Gruppe bestehend aus 6- oder 10-gliedrigem Aryl und 5- bis 10-gliedrigem Heteroaryl, das 1 bis 3 Heteroatome enthält, die unabhängig ausgewählt sind aus N, O und S,

wobei Cycloalkyl, Heterocycloalkyl, Aryl und Heteroaryl unsubstituiert oder mit 1 bis 4 Substituenten substituiert sind, die unabhängig voneinander aus der Gruppe ausgewählt sind, die aus Halogen CN, SF₅, NO₂, Oxo, C₁₋₄-Alkyl, C₀₋₆-Alkylen-OR³¹, C₀₋₆-Alkylen-(3- bis 6-gliedrigem Cycloalkyl), C₀₋₆-Alkylen-(3- bis 6-gliedrigem Heterocycloalkyl), C₀₋₆-Alkylen-(6-gliedrigem Aryl), C₀₋₆-Alkylen-(5- bis 6-gliedrigem Heteroaryl), C₀₋₆-Alkylen-S(O)_nR³¹, C₀₋₆-Alkylen-NR³¹S(O)₂R³¹, C₀₋₆-Alkylen-S(O)₂NR³¹R³², C₀₋₆-Alkylen-NR³¹S(O)₂NR³¹R³², C₀₋₆-Alkylen-CO₂R³¹, O-C₁₋₆-Alkylen-CO₂R³¹, C₀₋₆-Alkylen-O-COR³¹, C₀₋₆-Alkylen-CONR³¹R³², C₀₋₆-Alkylen-NR³¹-COR³¹, C₀₋₆-Alkylen-NR³¹-CONR³¹R³², C₀₋₆-Alkylen-O-CONR³¹R³², C₀₋₆-Alkylen-NR³¹-CO₂R³¹ und C₀₋₆-Alkylen-NR³¹R³² besteht,

wobei Alkyl, Alkylen, Cycloalkyl, Heterocycloalkyl, Aryl und Heteroaryl unsubstituiert oder mit 1 bis 6 Substituenten substituiert ist, die unabhängig voneinander aus Halogen, CN, Oxo, Hydroxy, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl ausgewählt sind;

und wobei gegebenenfalls zwei benachbarte Substituenten an der Aryl- oder Heteroaryleinheit einen 5- bis 8-gliedrigen teilweise ungesättigten Ring bilden, der gegebenenfalls 1 bis 3 Heteroatome enthält, die unabhängig voneinander aus O, S oder N ausgewählt sind, und

wobei dieser zusätzliche Ring unsubstituiert oder mit 1 bis 4 Substituenten substituiert ist, die unabhängig voneinander aus Halogen, CN, Oxo, OH, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl ausgewählt sind;



ausgewählt ist aus der Gruppe bestehend aus 3- bis 10-gliedrigem Cycloalkyl, 3- bis 10-gliedrigem Heterocycloalkyl, das 1 bis 3 Heteroatome enthält, die unabhängig voneinander ausgewählt sind aus N, O und S, 6- bis 14-gliedrigem Aryl und 5- bis 14-gliedrigem Heteroaryl, das 1 bis 4 Heteroatome enthält, die unabhängig voneinander ausgewählt sind aus N, O und S,

wobei Cycloalkyl, Heterocycloalkyl, Aryl und Heteroaryl unsubstituiert oder mit 1 bis 6 Substituenten substituiert sind, die unabhängig voneinander aus der Gruppe ausgewählt sind, die aus Halogen, CN, SF₅, NO₂, Oxo, C₁₋₄-Alkyl, C₀₋₆-Alkylen-OR²¹, C₀₋₆-Alkylen-(3- bis 6-gliedrigem Cycloalkyl), C₀₋₆-Alkylen-(3- bis 6-gliedrigem Heterocycloalkyl), C₀₋₆-Alkylen-S(O)_nR²¹, C₀₋₆-Alkylen-NR²¹S(O)₂R²¹, C₀₋₆-Alkylen-S(O)₂NR²¹R²², C₀₋₆-Alkylen-NR²¹S(O)₂NR²¹R²², C₀₋₆-Alkylen-CR⁴¹(=N-OR⁴¹), C₀₋₆-Alkylen-CO₂R²¹, O-C₁₋₆-Alkylen-CO₂R²¹, C₀₋₆-Alkylen-O-COR²¹, C₀₋₆-Alkylen-CONR²¹R²², C₀₋₆-Alkylen-NR²¹-COR²¹, C₀₋₆-Alkylen-NR²¹-CONR²¹R²², C₀₋₆-Alkylen-O-CONR²¹R²², C₀₋₆-Alkylen-NR²¹-CO₂R²¹ und C₀₋₆-Alkylen-NR²¹R²² besteht,

wobei Alkyl, Alkylen, Cycloalkyl und Heterocycloalkyl unsubstituiert oder mit 1 bis 6 Substituenten substituiert ist, die unabhängig voneinander aus Halogen, CN, Oxo, Hydroxy, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, CO-OC₁₋₄-Alkyl, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl ausgewählt sind;

und wobei gegebenenfalls zwei benachbarte Substituenten an der Aryl- oder Heteroaryleinheit einen 5- bis 8-gliedrigen teilweise ungesättigten Ring bilden, der gegebenenfalls 1 bis 3 Heteroatome enthält, die unabhängig voneinander aus O, S oder N ausgewählt sind, und

wobei dieser zusätzliche Zyklus unsubstituiert oder mit 1 bis 4 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus Halogen, CN, Oxo, OH, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl;

und wobei gegebenenfalls zwei benachbarte Substituenten an der Cycloalkyl- oder Heterocycloalkyl-Einheit einen 5- bis 6-gliedrigen ungesättigten Ring bilden, der gegebenenfalls 1 bis 3 Heteroatome enthält, die unabhängig voneinander aus O, S oder N ausgewählt sind,

wobei dieser zusätzliche Ring unsubstituiert oder mit 1 bis 4 Substituenten substituiert ist, die unabhängig voneinander aus Halogen, CN, Oxo, OH, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H,

C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl ausgewählt sind;


wobei

5



einen Substituenten von oben in 1,2-Orientierung bezüglich der Verbindung zu

10



aufweist oder einen weiteren anellierten Ring in 1,2-Orientierung hat;

15

L ausgewählt ist aus der Gruppe bestehend aus einer Bindung, C₁₋₆-Alkylen, C₂₋₆-Alkenylen, C₂₋₆-Alkinylen, 3- bis 10-gliedrigem Cycloalkylen, 3- bis 10-gliedrigem Heterocycloalkylen, das 1 bis 4 Heteroatome enthält, die unabhängig ausgewählt sind aus N, O und S, 6- oder 10-gliedrigem Arylen und 5- bis 10-gliedrigem Heteroarylen, das 1 bis 4 Heteroatome enthält, die unabhängig ausgewählt sind aus N, O und S,

20

wobei Alkylen, Alkenylen, Alkinylen, Cycloalkylen, Heterocycloalkylen, Arylen und Heteroarylen unsubstituiert oder mit 1 bis 6 Substituenten substituiert sind, die unabhängig voneinander aus der Gruppe ausgewählt sind, die aus Halogen, CN, SF₅, NO₂, Oxo, C₁₋₄-Alkyl, C₀₋₆-Alkylen-OR⁴¹, C₀₋₆-Alkylen-(3- bis 6-gliedriges Cycloalkyl), C₀₋₆-Alkylen-(3- bis 6-gliedriges Heterocycloalkyl), C₀₋₆-Alkylen-S(O)_nR⁴¹, C₀₋₆-Alkylen-NR⁴¹S(O)₂R⁴¹, C₀₋₆-Alkylen-S(O)₂NR⁴¹R⁴², C₀₋₆-Alkylen-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-Alkylen-CO₂R⁴¹, O-C₁₋₆-Alkylen-CO₂R⁴¹, C₀₋₆-Alkylen-O-COR⁴¹, C₀₋₆-Alkylen-CONR⁴¹R⁴², C₀₋₆-Alkylen-NR⁴¹-COR⁴¹, C₀₋₆-Alkylen-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-Alkylen-O-CONR⁴¹R⁴², C₀₋₆-Alkylen-NR⁴¹-CO₂R⁴¹ und C₀₋₆-Alkylen-NR⁴¹R⁴²,

25

wobei Alkyl, Alkylen, Cycloalkyl und Heterocycloalkyl unsubstituiert oder mit 1 bis 6 Substituenten substituiert ist, die unabhängig voneinander aus Halogen, CN, Oxo, Hydroxy, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl ausgewählt sind;

30

und wobei gegebenenfalls zwei benachbarte Substituenten an der Arylen- und Heteroaryleneinheit einen 5- bis 8-gliedrigen teilweise ungesättigten Ring bilden, der gegebenenfalls 1 bis 3 Heteroatome enthält, die unabhängig voneinander aus O, S oder N ausgewählt sind, und

35

wobei dieser zusätzliche Ring unsubstituiert oder mit 1 bis 4 Substituenten substituiert ist, die unabhängig voneinander aus Halogen, CN, Oxo, OH, CO₂H, CO₂-C₁₋₄-Alkyl, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl ausgewählt sind;

40

R¹ ausgewählt ist aus der Gruppe bestehend aus H, Halogen, CN, SF₅, NO₂, Oxo, C₁₋₄-Alkyl, C₀₋₆-Alkylen-OR⁴¹, Y-C₀₋₆-Alkylen-(3- bis 6-gliedrigem Cycloalkyl), Y-C₀₋₆-Alkylen-(3- bis 6-gliedrigem Heterocycloalkyl), Y-C₀₋₆-Alkylen-(6-gliedrigem Aryl), Y-C₀₋₆-Alkylen-(5- bis 6-gliedrigem Heteroaryl), C₀₋₆-Alkylen-S(=O)(-R⁴¹)=N-R⁷⁵, X-C₁₋₆-Alkylen-S(=O)(-R⁴¹)=N-R⁷⁵, C₀₋₆-Alkylen-S(O)_nR⁴¹, X-C₁₋₆-alkylene-S(O)_nR⁴¹, C₀₋₆-Alkylen-S(=NR⁷¹)R⁴¹, X-C₁₋₆-Alkylen-S(=NR⁷¹)R⁴¹, C₀₋₆-Alkylen-S(O)(=NR⁷¹)R⁴¹, X-C₁₋₆-Alkylen-S(O)(=NR⁷¹)R⁴¹, C₀₋₆-Alkylen-S(=NR⁷¹)₂R⁴¹, X-C₁₋₆-Alkylen-S(=NR⁷¹)₂R⁴¹, C₀₋₆-Alkylen-NR⁴¹S(O)₂R⁴¹, X-C₁₋₆-Alkylen-NR⁴¹S(O)₂R⁴¹, C₀₋₆-Alkylen-S(O)₂NR⁴¹R⁴², X-C₁₋₆-Alkylen-S(O)₂NR⁴¹R⁴², C₀₋₆-Alkylen-NR⁴¹S(O)₂NR⁴¹R⁴², X-C₁₋₆-Alkylen-NR⁴¹S(O)₂NR⁴¹R⁴², C₀₋₆-Alkylen-SO₃R⁴¹, X-C₁₋₆-Alkylen-SO₃R⁴¹, C₀₋₆-Alkylen-CO₂R⁴¹, X-C₁₋₆-Alkylen-CO₂R⁴¹, C₀₋₆-Alkylen-O-COR⁴¹, X-C₁₋₆-Alkylen-O-COR⁴¹, C₀₋₆-Alkylen-CONR⁴¹R⁴², X-C₁₋₆-Alkylen-CONR⁴¹R⁴², C₀₋₆-Alkylen-CONR⁴¹OR⁴¹, X-C₁₋₆-Alkylen-CONR⁴¹OR⁴¹, C₀₋₆-Alkylen-CONR⁴¹SO₂R⁴¹, X-C₁₋₆-Alkylen-CONR⁴¹SO₂R⁴¹, C₀₋₆-Alkylen-NR⁴¹-COR⁴¹, X-C₁₋₆-C₀₋₆-Alkylen-NR⁴¹-COR⁴¹, C₀₋₆-alkylene-NR⁴¹-CONR⁴¹R⁴², X-C₁₋₆-Alkylen-NR⁴¹-CONR⁴¹R⁴², C₀₋₆-Alkylen-O-CONR⁴¹R⁴², X-C₁₋₆-Alkylen-O-CONR⁴¹R⁴², C₀₋₆-Alkylen-NR⁴¹-CO₂R⁴¹, X-C₁₋₆-Alkylen-NR⁴¹-CO₂R⁴¹, C₀₋₆-Alkylen-NR⁴¹R⁴², X-C₁₋₆-Alkylen-NR⁴¹R⁴²,

45

50


55

wobei Alkyl, Alkylen, Cycloalkyl, Heterocycloalkyl, Aryl und Heteroaryl unsubstituiert oder mit 1 bis 6 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus Halogen, CN, Oxo, Hydroxy, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl;


und wobei gegebenenfalls zwei benachbarte Substituenten an der Aryl- und Heteroaryleneinheit einen 5- bis

8-gliedrigen teilweise ungesättigten Zyklus bilden, der gegebenenfalls 1 bis 3 Heteroatome enthält, die unabhängig voneinander aus O, S oder N ausgewählt sind, und wobei dieser zusätzliche Ring unsubstituiert oder mit 1 bis 4 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus Halogen, CN, Oxo, OH, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl;


R¹¹, R¹², R²¹, R²², R³¹, R³², R⁴¹, R⁴², R⁵¹ unabhängig voneinander aus H und C₁₋₄-Alkyl ausgewählt sind, wobei Alkyl unsubstituiert oder substituiert ist mit 1 bis 3 Substituenten, unabhängig ausgewählt aus Halogen, CN, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, 3- bis 6-gliedrigem Cycloalkyl, Halogen-(3- bis 6-gliedrigem Cycloalkyl), 3- bis 6-gliedrigem Heterocycloalkyl, Halogen-(3- bis 6-gliedrigem Heterocycloalkyl), OH, Oxo, CO₂H, CO₂-C₁₋₄-alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-alkyl; oder R¹¹ und R¹², R²¹ und R²², R³¹ und R³², R⁴¹ und R⁴² jeweils zusammen mit dem Stickstoff, an den sie gebunden sind, einen 3- bis 6-gliedrigen Ring vervollständigen, der Kohlenstoffatome und gegebenenfalls 1 oder 2 Heteroatome enthält, die unabhängig voneinander aus O, S oder N ausgewählt sind; und wobei der neu gebildete Ring unsubstituiert oder mit 1 bis 3 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus Halogen, CN, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, 3- bis 6-gliedrigem Cycloalkyl, Halogen-(3- bis 6-gliedrigem Cycloalkyl), 3- bis 6-gliedrigem Heterocycloalkyl, Halogen-(3- bis 6-gliedrigem Heterocycloalkyl), OH, Oxo, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl; R⁷¹ unabhängig ausgewählt ist aus H, CN; NO₂, C₁₋₄-Alkyl und C(O)-OC₁₋₄-Alkyl, wobei Alkyl unsubstituiert oder substituiert ist mit 1 bis 3 Substituenten, unabhängig ausgewählt aus Halogen, CN, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, 3- bis 6-gliedrigem Cycloalkyl, Halogen-(3- bis 6-gliedrigem Cycloalkyl), 3- bis 6-gliedrigem Heterocycloalkyl, Halogen-(3- bis 6-gliedrigem Heterocycloalkyl), OH, Oxo, CO₂H, CO₂-C₁₋₄-Alkyl, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl; R⁷⁵ unabhängig ausgewählt ist aus C₁₋₄-Alkyl, 3- bis 6-gliedrigem Cycloalkyl, 3- bis 6-gliedrigem Heterocycloalkyl, 6-gliedrigem Aryl und 5- bis 6-gliedrigem Heteroaryl, wobei Alkyl, Cycloalkyl, Heterocycloalkyl, Aryl und Heteroaryl unsubstituiert oder mit 1 bis 3 Substituenten substituiert ist, die unabhängig voneinander aus Halogen, CN, Me, Et, CHF₂, CF₃, OH, Oxo, CO₂H, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, OMe, OEt, OCHF₂ und OCF₃ ausgewählt sind; X unabhängig ausgewählt ist aus O, NR⁵¹, S(O)_n, S(=NR⁷¹), S(O)(=NR⁷¹) und S(=NR⁷¹)₂; Y unabhängig ausgewählt ist aus einer Bindung, O, NR⁵¹, S(O)_n, S(=NR⁷¹), S(O)(=NR⁷¹) und S(=NR⁷¹)₂; n unabhängig ausgewählt ist 0 bis 2; und mit der Maßgabe, dass die folgenden Strukturen ausgeschlossen sind:



2. Verbindung nach Anspruch 1 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, N-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon, wobei



ausgewählt ist aus



wobei

(A)

unsubstituiert oder substituiert ist mit 1 bis 3 Substituenten, unabhängig ausgewählt aus der Gruppe bestehend aus F, Cl, Br, CN, OH, Oxo, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, O-C₁₋₄-Alkyl, O-Halogen-C₁₋₄-alkyl, NH₂, NHC₁₋₄-Alkyl, N(C₁₋₄-Alkyl)₂, SO₂-C₁₋₄-Alkyl und SO₂-Halogen-C₁₋₄-Alkyl.

3. Verbindung nach Anspruch 1 oder 2 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solut oder pharmazeutisch annehmbares Salz davon, wobei

(B)


ausgewählt ist aus der Gruppe bestehend aus Phenyl, Naphthyl, Pyridyl, Pyrimidinyl, Thiophenyl, Thiazolyl, Cyclopentyl, Cyclohexyl, Bicyclo[1.1.1]pentyl, Bicyclo[2.2.2]octyl, Bicyclo[2.2.1]heptyl, Pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octyl und Piperidinyl,

wobei der Ring unsubstituiert oder mit 1 bis 3 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus der Gruppe bestehend aus F, Cl, Br, CN, OH, Oxo, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl, O-Halogen-C₁₋₄-Alkyl, C₁₋₄-Alkyl-OH und Halogen-C₁₋₄-Alkyl-OH; und wobei gegebenenfalls zwei benachbarte Substituenten am Phenylring zusammen eine -(CH₂)₃-, -(CH₂)₄-, -OCF₂O- und -OCH₂O-Gruppe bilden.

4. Verbindung nach einem der Ansprüche 1 bis 3 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solut oder pharmazeutisch annehmbares Salz davon, wobei

(C)


aus Phenyl, Pyridyl und Thiophenyl ausgewählt ist; wobei Phenyl, Pyridyl und Thiophenyl unsubstituiert oder mit 1 bis 3 Substituenten substituiert ist, die unabhängig voneinander ausgewählt sind aus der Gruppe bestehend aus F, Cl, CN, OH, Oxo, C₁₋₄-Alkyl, Halogen-C₁₋₄-Alkyl, O-C₁₋₄-Alkyl und O-Halogen-C₁₋₄-Alkyl; und wobei der Rest -L-R¹ in 1,3-Orientierung in Bezug auf die Verbindung mit




verknüpft ist und L keine Bindung ist.

5. Verbindung nach einem der Ansprüche 1 bis 4 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solut oder pharmazeutisch annehmbares Salz davon, wobei

-L-R¹ ausgewählt ist aus






wobei der Ring unsubstituiert oder weiter substituiert ist mit 1 bis 4 Substituenten, die unabhängig voneinander ausgewählt sind aus der Gruppe bestehend aus F, Cl, Br, CN, OH, Oxo, C₁₋₄-Alkyl, Halogen-C₁₋₄-alkyl, O-C₁₋₄-Alkyl, O-Halogen-C₁₋₄-alkyl, C₁₋₄-Alkyl-OH, Halogen-C₁₋₄-alkyl-OH, SO₂-C₁₋₄-Alkyl und SO₂-Halogen-C₁₋₄-alkyl; und wobei gegebenenfalls zwei benachbarte Substituenten am Phenylring zusammen eine -(CH₂)₃-, -(CH₂)₄-, -OCF₂O- und -OCH₂O-Gruppe bilden.

6. Verbindung nach einem der Ansprüche 1 bis 5, wobei


R¹ ausgewählt ist aus CO₂H, Tetrazol, CH₂CO₂H, OCH₂CO₂H, SO₂CH₂CO₂H, CHMeCO₂H, CMe₂CO₂H, C(OH)MeCO₂H, CONHSO₂Me und CONH(OH); und gegebenenfalls das Glycin- und Tauro-Konjugat davon oder Enantiomer, Diastereomer, Tautomer, N-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon.

7. Verbindung nach einem der Ansprüche 1 bis 6, wobei

-L-R¹ ausgewählt ist aus



und




und gegebenenfalls das Glycin- und Tauro-Konjugat davon oder das Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Sivat oder pharmazeutisch annehmbare Salz davon.

8. Verbindung nach einem der Ansprüche 1 bis 7 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Sivat oder pharmazeutisch annehmbares Salz davon, wobei



(D)

ausgewählt ist aus der Gruppe bestehend aus



wobei

R^2 ausgewählt ist aus Me, F, Cl, CN, Me, CHO, CHF_2 , CF_3 , SO_2Me ,



und

wobei


(D)

gegebenenfalls weiter substituiert ist mit 1 bis 2 Substituenten, ausgewählt aus der Gruppe bestehend aus F, Cl, CN, Me, OMe, CHO, CHF_2 und CF_3 .

9. Verbindung nach einem der Ansprüche 1 bis 8 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Sivat oder pharmazeutisch annehmbares Salz davon, wobei

(D)

ausgewählt ist aus der Gruppe bestehend aus





25

-L-R¹ ausgewählt ist aus



34

4



5


209

14. Verbindung nach einem der Ansprüche 1 bis 13, wobei
 R^1 COOH oder ein Glycin-Konjugat oder Tauro-Konjugat davon oder ein Enantiomer, Diastereomer, Tautomer,
N-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon ist.

15. Verbindung nach einem der Ansprüche 1 bis 13 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastere-
 omer, Tautomer, *N*-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon, wobei
 R^1 C₀₋₆-Alkylen-CONR⁴¹R⁴² ist.

16. Verbindung nach Anspruch 15 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer,
N-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon, wobei
 R^{41} und R^{42} unabhängig voneinander aus H und C₁₋₄-Alkyl ausgewählt sind, wobei C₁₋₄-Alkyl unsubstituiert oder
 mit CO₂H substituiert ist.







17. Verbindung nach einem der Ansprüche 1 bis 12, wobei
 L-R¹


















ist, oder ein Glycin-Konjugat oder Tauro-Konjugat davon, oder ein Enantiomer, Diastereomer, Tautomer, *N*-Oxid,
 Solvat oder pharmazeutisch annehmbares Salz davon.

18. Verbindung nach einem der Ansprüche 1 bis 17, wobei die Verbindung ein Glycin-Konjugat oder ein Enantiomer,
 Diastereomer, Tautomer, *N*-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon ist.


19. Verbindung nach einem der Ansprüche 1 bis 18, ausgewählt aus








und




oder ein Glycin-Konjugat oder Tauro-Konjugat davon; und
ein Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solvat oder ein pharmazeutisch annehmbares Salz davon.

20. Verbindung nach einem der Ansprüche 1 bis 19, die



ein Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon ist.

21. Verbindung nach einem der Ansprüche 1 bis 19, die



ist,

oder ein Glycin-Konjugat davon, ein Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon ist.


22. Verbindung nach einem der Ansprüche 1 bis 21 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon zur Verwendung als Arzneimittel.

23. Verbindung nach einem der Ansprüche 1 bis 21 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon zur Verwendung bei der Prophylaxe und/oder Behandlung von Krankheiten, die für eine Behandlung mit LXR-Modulatoren geeignet sind, wobei die Krankheit ausgewählt ist aus nicht-alkoholischer Fettlebererkrankung, nicht-alkoholischer Steatohepatitis, Leberentzündung, Leberfibrose, Fettleibigkeit, Insulinresistenz, Typ-II-Diabetes, familiäre Hypercholesterinämie, Hypercholesterinämie beim nephrotischen Syndrom, metabolisches Syndrom, Herzsteatose, Krebs, virale Myokarditis, Hepatitis-C-Virusinfektion oder deren Komplikationen und unerwünschte Nebenwirkungen einer Langzeit-Glukokortikoid-Behandlung bei Krankheiten wie rheumatoider Arthritis, entzündlichen Darmerkrankungen und Asthma.

24. Pharmazeutische Zusammensetzung, umfassend eine Verbindung nach einem der Ansprüche 1 bis 21 oder ein Glycin-Konjugat, Tauro-Konjugat, Enantiomer, Diastereomer, Tautomer, *N*-Oxid, Solvat oder pharmazeutisch annehmbares Salz davon und einen pharmazeutisch annehmbaren Träger oder Hilfsstoff.

Revendications

1. Composé représenté par la formule (I)



conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel



est un cycle d'annulation de 5 à 6 chaînons qui forme un aryle à 6 chaînons ou un hétéroaryle de 5 à 6 chaînons contenant 1 à 3 hétéroatomes choisis indépendamment parmi N, O et S, ce cycle étant non substitué ou substitué par 1 à 4 substituants choisis indépendamment dans le groupe constitué par un halogène, CN, SF₅, NO₂, alkyle en C₁₋₆, oxo, alkylène en C₀₋₆-OR¹¹, alkylène en C₀₋₆-(cycloalkyle de 3 à 6 chaînons), alkylène en C₀₋₆-(hété-

rocycloalkyle de 3 à 6 chaînons), alkylène en $C_{0-6}-S(O)_nR^{11}$, alkylène en $C_{0-6}-NR^{11}S(O)_2R^{11}$, alkylène en $C_{0-6}-S(O)_2NR^{11}R^{12}$, alkylène en $C_{0-6}-NR^{11}S(O)_2NR^{11}R^{12}$, alkylène en $C_{0-6}-CO_2R^{11}$, O-alkylène en $C_{1-6}-CO_2R^{11}$, alkylène en $C_{0-6}-O-COR^{11}$, alkylène en $C_{0-6}-CONR^{11}R^{12}$, alkylène en $C_{0-6}-NR^{11}-COR^{11}$, alkylène en $C_{0-6}-NR^{11}-CONR^{11}R^{12}$, alkylène en $C_{0-6}-O-CONR^{11}R^{12}$, alkylène en $C_{0-6}-NR^{11}-CO_2R^{11}$ et alkylène en $C_{0-6}-NR^{11}R^{12}$,

l'alkyle, l'alkylène, le cycloalkyle et l'hétérocycloalkyle étant non substitués ou substitués par 1 à 6 substituants choisis indépendamment parmi un halogène, CN, oxo, hydroxy, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, alkyle en C_{1-4} , haloalkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ; et dans lequel, éventuellement, deux substituants adjacents sur le fragment aryle ou hétéroaryle forment un cycle partiellement insaturé de 5 à 8 chaînons contenant éventuellement de 1 à 3 hétéroatomes choisis indépendamment parmi O, S ou N, et

le nouveau cycle formé étant non substitué ou substitué par 1 à 3 substituants choisis indépendamment parmi un halogène, CN, alkyle en C_{1-4} , haloalkyle en C_{1-4} , cycloalkyle de 3 à 6 chaînons, halo-(cycloalkyle de 3 à 6 chaînons), hétérocycloalkyle de 3 à 6 chaînons, halo-(hétérocycloalkyle de 3 à 6 chaînons), OH, oxo, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ;



est choisi dans le groupe constitué par un cycloalkyle de 3 à 10 chaînons, un hétérocycloalkyle de 3 à 10 chaînons contenant de 1 à 3 hétéroatomes choisis indépendamment parmi N, O et S, un aryle de 6 à 14 chaînons et un hétéroaryle de 5 à 14 chaînons contenant de 1 à 4 hétéroatomes choisis indépendamment parmi N, O et S, le cycloalkyle, l'hétérocycloalkyle, l'aryle et l'hétéroaryle étant non substitués ou substitués par 1 à 6 substituants choisis indépendamment dans le groupe constitué par un halogène, CN, SF_5 , NO_2 , oxo, alkyle en C_{1-4} , alkylène en $C_{0-6}-OR^{21}$, alkylène en C_{0-6} -(cycloalkyle de 3 à 6 chaînons), alkylène en C_{0-6} -(hétérocycloalkyle de 3 à 6 chaînons), alkylène en $C_{0-6}-S(O)_nR^{21}$, alkylène en $C_{0-6}-NR^{21}S(O)_2R^{21}$, alkylène en $C_{0-6}-S(O)_2NR^{21}R^{22}$, alkylène en $C_{0-6}-NR^{21}S(O)_2NR^{21}R^{22}$, alkylène en $C_{0-6}-CO_2R^{21}$, O-alkylène en $C_{1-6}-CO_2R^{21}$, alkylène en $C_{0-6}-O-COR^{21}$, alkylène en $C_{0-6}-CONR^{21}R^{22}$, alkylène en $C_{0-6}-NR^{21}-COR^{21}$, alkylène en $C_{0-6}-NR^{21}-CONR^{21}R^{22}$, alkylène en $C_{0-6}-O-CONR^{21}R^{22}$, alkylène en $C_{0-6}-NR^{21}-CO_2R^{21}$ et alkylène en $C_{0-6}-NR^{21}R^{22}$,

l'alkyle, l'alkylène, le cycloalkyle et l'hétérocycloalkyle étant non substitués ou substitués par 1 à 6 substituants choisis indépendamment parmi un halogène, CN, oxo, hydroxy, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, alkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ,

et dans lequel, éventuellement, deux substituants adjacents sur le fragment aryle ou hétéroaryle forment un cycle partiellement insaturé de 5 à 8 chaînons contenant éventuellement de 1 à 3 hétéroatomes choisis indépendamment parmi O, S ou N, et

ce cycle supplémentaire étant non substitué ou substitué par 1 à 4 substituants choisis indépendamment parmi un halogène, CN, oxo, OH, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, alkyle en C_{1-4} , haloalkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ,

et dans lequel, éventuellement, deux substituants adjacents sur le fragment cycloalkyle ou hétérocycloalkyle forment un cycle insaturé de 5 à 8 chaînons contenant éventuellement de 1 à 3 hétéroatomes choisis indépendamment parmi O, S ou N,

ce cycle supplémentaire étant non substitué ou substitué par 1 à 4 substituants choisis indépendamment parmi un halogène, CN, oxo, OH, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, alkyle en C_{1-4} , haloalkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ;



est choisi dans le groupe constitué par un aryle de 6 ou 10 chaînons et un hétéroaryle de 5 à 10 chaînons contenant de 1 à 3 hétéroatomes choisis indépendamment parmi N, O et S,

le cycloalkyle, l'hétérocycloalkyle, l'aryle et l'hétéroaryle étant non substitués ou substitués par 1 à 4 substituants choisis indépendamment dans le groupe constitué par un halogène, CN, SF_5 , NO_2 , oxo, alkyle en C_{1-4} , alkylène en $C_{0-6}-OR^{31}$, alkylène en C_{0-6} -(cycloalkyle de 3 à 6 chaînons), alkylène en C_{0-6} -(hétérocycloalkyle de 3 à 6 chaînons), alkylène en C_{0-6} -(aryle à 6 chaînons), alkylène en C_{0-6} -(hétéroaryle de 5 à 6 chaînons), alkylène en $C_{0-6}-S(O)_nR^{31}$, alkylène en $C_{0-6}-NR^{31}S(O)_2R^{31}$, alkylène en $C_{0-6}-S(O)_2NR^{31}R^{32}$, alkylène en $C_{0-6}-NR^{31}S(O)_2NR^{31}R^{32}$, alkylène en $C_{0-6}-CO_2R^{31}$, O-alkylène en $C_{1-6}-CO_2R^{31}$, alkylène en $C_{0-6}-O-COR^{31}$, alkylène en $C_{0-6}-CONR^{31}R^{32}$, alkylène en $C_{0-6}-NR^{31}-COR^{31}$, alkylène en

$C_{0-6}-NR^{31}-CONR^{31}R^{32}$, alkylène en $C_{0-6}-O-CONR^{31}R^{32}$, alkylène en $C_{0-6}-NR^{31}-CO_2R^{31}$ et alkylène en $C_{0-6}-NR^{31}R^{32}$,

l'alkyle, l'alkylène, le cycloalkyle, l'hétérocycloalkyle, l'aryle et l'hétéroaryle est non substitués ou substitués par 1 à 6 substituants choisis indépendamment parmi un halogène, CN, oxo, hydroxy, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, alkyle en C_{1-4} , haloalkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ;

et dans lequel, éventuellement, deux substituants adjacents sur le fragment aryle ou hétéroaryle forment un cycle partiellement insaturé de 5 à 8 chaînons contenant éventuellement de 1 à 3 hétéroatomes choisis indépendamment parmi O, S ou N, et

ce cycle supplémentaire étant non substitué ou substitué par 1 à 4 substituants choisis indépendamment parmi un halogène, CN, oxo, OH, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, alkyle en C_{1-4} , haloalkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ;



est choisi dans le groupe constitué par un cycloalkyle de 3 à 10 chaînons, un hétérocycloalkyle de 3 à 10 chaînons contenant de 1 à 3 hétéroatomes choisis indépendamment parmi N, O et S, un aryle de 6 à 14 chaînons et un hétéroaryle de 5 à 14 chaînons contenant de 1 à 4 hétéroatomes choisis indépendamment parmi N, O et S,

le cycloalkyle, l'hétérocycloalkyle, l'aryle et l'hétéroaryle étant non substitués ou substitués par 1 à 6 substituants choisis indépendamment dans le groupe constitué par un halogène, CN, SF_5 , NO_2 , oxo, alkyle en C_{1-4} , alkylène en $C_{0-6}-OR^{21}$, alkylène en C_{0-6} -(cycloalkyle de 3 à 6 chaînons), alkylène en C_{0-6} -(hétérocycloalkyle de 3 à 6 chaînons), alkylène en $C_{0-6}-S(O)_nR^{21}$, alkylène en $C_{0-6}-NR^{21}S(O)_2R^{21}$, alkylène en $C_{0-6}-S(O)_2NR^{21}R^{22}$, alkylène en $C_{0-6}-NR^{11}S(O)_2NR^{21}R^{22}$, alkylène en $C_{0-6}-CR^{41}(=N-OR^{41})$, alkylène en $C_{0-6}-CO_2R^{21}$, O-alkylène en $C_{1-6}-CO_2R^{21}$, alkylène en $C_{0-6}-O-COR^{21}$, alkylène en $C_{0-6}-CONR^{21}R^{22}$, alkylène en $C_{0-6}-NR^{21}-COR^{21}$, alkylène en $C_{0-6}-NR^{21}-CONR^{21}R^{22}$, alkylène en $C_{0-6}-O-CONR^{21}R^{22}$, alkylène en $C_{0-6}-NR^{21}-CO_2R^{21}$ et alkylène en $C_{0-6}-NR^{21}R^{22}$,

l'alkyle, l'alkylène, le cycloalkyle et l'hétérocycloalkyle étant non substitués ou substitués par 1 à 6 substituants choisis indépendamment parmi un halogène, CN, oxo, hydroxy, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, CO-Oalkyle en C_{1-4} , alkyle en C_{1-4} , haloalkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ;

et dans lequel, éventuellement, deux substituants adjacents sur le fragment aryle ou hétéroaryle forment un cycle partiellement insaturé de 5 à 8 chaînons contenant éventuellement de 1 à 3 hétéroatomes choisis indépendamment parmi O, S ou N, et

ce cycle supplémentaire étant non substitué ou substitué par 1 à 4 substituants choisis indépendamment parmi un halogène, CN, oxo, OH, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, alkyle en C_{1-4} , haloalkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ;

et dans lequel, éventuellement, deux substituants adjacents sur le fragment cycloalkyle ou hétérocycloalkyle forment un cycle insaturé de 5 à 6 chaînons contenant éventuellement de 1 à 3 hétéroatomes choisis indépendamment parmi O, S ou N,

ce cycle supplémentaire étant non substitué ou substitué par 1 à 4 substituants choisis indépendamment parmi un halogène, CN, oxo, OH, CO_2H , CO_2 -alkyle en C_{1-4} , $CONHCH_2CO_2H$, $CONH(CH_2)_2SO_3H$, alkyle en C_{1-4} , haloalkyle en C_{1-4} , O-alkyle en C_{1-4} et O-halo-alkyle en C_{1-4} ;



ayant un substituant parmi ceux qui précèdent dans une orientation 1,2 par rapport à la liaison en direction de



ou ayant un cycle d'annulation supplémentaire dans une orientation 1,2 ;

L est choisi dans le groupe constitué par une liaison, un alkylène en C₁₋₆, un alcénylène en C₂₋₆, un alcynylène en C₂₋₆, un cycloalkylène de 3 à 10 chaînons, un hétérocycloalkylène de 3 à 10 chaînons contenant de 1 à 4 hétéroatomes choisis indépendamment parmi N, O et S, un arylène de 6 ou 10 chaînons et un hétéroarylène de 5 à 10 chaînons contenant de 1 à 4 hétéroatomes choisis indépendamment parmi N, O et S,

l'alkylène, l'alcénylène, l'alcynylène, le cycloalkylène, l'hétérocycloalkylène, l'arylène et l'hétéroarylène étant non substitués ou substitués par 1 à 6 substituants choisis indépendamment dans le groupe constitué par un halogène, CN, SF₅, NO₂, oxo, alkyle en C₁₋₄, alkylène en C₀₋₆-OR⁴¹, alkylène en C₀₋₆-(cycloalkyle de 3 à 6 chaînons), alkylène en C₀₋₆-(hétérocycloalkyle de 3 à 6 chaînons), alkylène en C₀₋₆-S(O)_nR⁴¹, alkylène en C₀₋₆-NR⁴¹S(O)₂R⁴¹, alkylène en C₀₋₆-S(O)₂NR⁴¹R⁴², alkylène en C₀₋₆-NR⁴¹S(O)₂NR⁴¹R⁴², alkylène en C₀₋₆-CO₂R⁴¹, O-alkylène en C₁₋₆-CO₂R⁴¹, alkylène en C₀₋₆-O-COR⁴¹, alkylène en C₀₋₆-CONR⁴¹R⁴², alkylène en C₀₋₆-NR⁴¹-COR⁴¹, alkylène en C₀₋₆-NR⁴¹-CONR⁴¹R⁴², alkylène en C₀₋₆-O-CONR⁴¹R⁴², alkylène en C₀₋₆-NR⁴¹-CO₂R⁴¹ et alkylène en C₀₋₆-NR⁴¹R⁴²,

l'alkyle, l'alkylène, le cycloalkyle et l'hétérocycloalkyle étant non substitués ou substitués par 1 à 6 substituants choisis indépendamment parmi un halogène, CN, oxo, hydroxy, CO₂H, CO₂-alkyle en C₁₋₄, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-alkyle en C₁₋₄ et O-halo-alkyle en C₁₋₄ ;

et dans lequel, éventuellement, deux substituants adjacents sur le fragment arylène et hétéroarylène forment un cycle partiellement insaturé de 5 à 8 chaînons contenant éventuellement de 1 à 3 hétéroatomes choisis indépendamment parmi O, S ou N, et

ce cycle supplémentaire étant non substitué ou substitué par 1 à 4 substituants choisis indépendamment parmi un halogène, CN, oxo, OH, CO₂H, CO₂-alkyle en C₁₋₄, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-alkyle en C₁₋₄ et O-halo-alkyle en C₁₋₄ ;

R¹ est choisi dans le groupe constitué par H, un halogène, CN, SF₅, NO₂, oxo, alkyle en C₁₋₄, alkylène en C₀₋₆-OR⁴¹, Y-alkylène en C₀₋₆-(cycloalkyle de 3 à 6 chaînons), Y-alkylène en C₀₋₆-(hétérocycloalkyle de 3 à 6 chaînons), Y-alkylène en C₀₋₆-(aryle à 6 chaînons), Y-alkylène en C₀₋₆-(hétéroaryle de 5 à 6 chaînons), alkylène en C₀₋₆-S(=O)(-R⁴¹)=N-R⁷⁵, X-alkylène en C₁₋₆-S(=O)(-R⁴¹)=N-R⁷⁵, alkylène en C₀₋₆-S(O)_nR⁴¹, X-alkylène en C₁₋₆-S(O)_nR⁴¹, alkylène en C₀₋₆-S(=NR⁷¹)R⁴¹, X-alkylène en C₁₋₆-S(=NR⁷¹)R⁴¹, alkylène en C₀₋₆-S(O)(=NR⁷¹)R⁴¹, X-alkylène en C₁₋₆-S(O)(=NR⁷¹)R⁴¹, alkylène en C₀₋₆-S(=NR⁷¹)₂R⁴¹, X-alkylène en C₁₋₆-S(=NR⁷¹)₂R⁴¹, alkylène en C₀₋₆-NR⁴¹S(O)₂R⁴¹, X-alkylène en C₁₋₆-NR⁴¹S(O)₂R⁴¹, alkylène en C₀₋₆-S(O)₂NR⁴¹R⁴², X-alkylène en C₁₋₆-S(O)₂NR⁴¹R⁴², alkylène en C₀₋₆-NR⁴¹S(O)₂NR⁴¹R⁴², X-alkylène en C₁₋₆-S(O)₂NR⁴¹R⁴², alkylène en C₀₋₆-SO₃R⁴¹, X-alkylène en C₁₋₆-SO₃R⁴¹, alkylène en C₀₋₆-CO₂R⁴¹, X-alkylène en C₁₋₆-CO₂R⁴¹, alkylène en C₀₋₆-O-COR⁴¹, X-alkylène en C₁₋₆-OCOR⁴¹, alkylène en C₀₋₆-CONR⁴¹R⁴², X-alkylène en C₁₋₆-CONR⁴¹R⁴², alkylène en C₀₋₆-CONR⁴¹OR⁴¹, X-alkylène en C₁₋₆-CONR⁴¹OR⁴¹, alkylène en C₀₋₆-CONR⁴¹SO₂R⁴¹, X-alkylène en C₁₋₆-CONR⁴¹SO₂R⁴¹, alkylène en C₀₋₆-NR⁴¹-COR⁴¹, X-alkylène en C₁₋₆-NR⁴¹-COR⁴¹, alkylène en C₀₋₆-NR⁴¹-CONR⁴¹R⁴², X-alkylène en C₁₋₆-NR⁴¹-CONR⁴¹R⁴², alkylène en C₀₋₆-O-CONR⁴¹R⁴², X-alkylène en C₁₋₆-O-CONR⁴¹R⁴², alkylène en C₀₋₆-NR⁴¹-CO₂R⁴¹, X-alkylène en C₁₋₆-NR⁴¹-CO₂R⁴¹, alkylène en C₀₋₆-NR⁴¹R⁴², X-alkylène en C₁₋₆-NR⁴¹R⁴²,

l'alkyle, l'alkylène, le cycloalkyle, l'hétérocycloalkyle, l'aryle et l'hétéroaryle étant non substitués ou substitués par 1 à 6 substituants choisis indépendamment parmi un halogène, CN, oxo, hydroxy, CO₂H, CO₂-alkyle en C₁₋₄, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-alkyle en C₁₋₄ et O-halo-alkyle en C₁₋₄ ;

et dans lequel, éventuellement, deux substituants adjacents sur le fragment aryle et hétéroaryle forment un cycle partiellement insaturé de 5 à 8 chaînons contenant éventuellement de 1 à 3 hétéroatomes choisis indépendamment parmi O, S ou N, et


ce cycle supplémentaire étant non substitué ou substitué par 1 à 4 substituants choisis indépendamment parmi un halogène, CN, oxo, OH, CO₂H, CO₂-alkyle en C₁₋₄, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-alkyle en C₁₋₄ et O-halo-alkyle en C₁₋₄ ;

R¹¹, R¹², R²¹, R²², R³¹, R³², R⁴¹, R⁴², R⁵¹ sont choisis indépendamment parmi H et un alkyle en C₁₋₄,


l'alkyle étant non substitué ou substitué par 1 à 3 substituants choisis indépendamment parmi un halogène, CN, alkyle en C₁₋₄, haloalkyle en C₁₋₄, cycloalkyle de 3 à 6 chaînons, halo-(cycloalkyle de 3 à 6 chaînons), hétérocycloalkyle de 3 à 6 chaînons, halo-(hétérocycloalkyle de 3 à 6 chaînons), OH, oxo, CO₂H, CO₂-alkyle en C₁₋₄, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-alkyle en C₁₋₄ et O-halo-alkyle en C₁₋₄ ;

ou R¹¹ et R¹², R²¹ et R²², R³¹ et R³², R⁴¹ et R⁴², respectivement, lorsqu'ils sont pris avec l'azote auquel ils


sont fixés complètent un cycle de 3 à 6 chaînons contenant des atomes de carbone et contenant éventuellement 1 ou 2 hétéroatomes choisis indépendamment parmi O, S ou N ; et le nouveau cycle formé étant non substitué ou substitué par 1 à 3 substituants choisis indépendamment parmi un halogène, CN, alkyle en C₁₋₄, haloalkyle en C₁₋₄, cycloalkyle de 3 à 6 chaînons, halo-(cycloalkyle de 3 à 6 chaînons), hétérocycloalkyle de 3 à 6 chaînons, halo-(hétérocycloalkyle de 3 à 6 chaînons), OH, oxo, CO₂H, CO₂-alkyle en C₁₋₄, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-alkyle en C₁₋₄ et O-halo-alkyle en C₁₋₄ ; R⁷¹ est choisi indépendamment parmi H, CN ; NO₂, un alkyle en C₁₋₄ et C(O)-O-alkyle en C₁₋₄, l'alkyle étant non substitué ou substitué par 1 à 3 substituants choisis indépendamment parmi un halogène, CN, alkyle en C₁₋₄, haloalkyle en C₁₋₄, cycloalkyle de 3 à 6 chaînons, halo-(cycloalkyle de 3 à 6 chaînons), hétérocycloalkyle de 3 à 6 chaînons, halo-(hétérocycloalkyle de 3 à 6 chaînons), OH, oxo, CO₂H, CO₂-alkyle en C₁₋₄, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, O-alkyle en C₁₋₄ et O-halo-alkyle en C₁₋₄ ; R⁷⁵ est choisi indépendamment parmi un alkyle en C₁₋₄, un cycloalkyle de 3 à 6 chaînons, un hétérocycloalkyle de 3 à 6 chaînons, un aryle à 6 chaînons et un hétéroaryle de 5 à 6 chaînons, l'alkyle, le cycloalkyle, l'hétérocycloalkyle, l'aryle et l'hétéroaryle étant non substitué ou substitué par 1 à 3 substituants choisis indépendamment parmi un halogène, CN, Me, Et, CHF₂, CF₃, OH, oxo, CO₂H, CONHCH₂CO₂H, CONH(CH₂)₂SO₃H, SO₃H, OMe, OEt, OCHF₂ et OCF₃ ; X est choisi indépendamment parmi O, NR⁵¹, S(O)_n, S(=NR⁷¹), S(O)(=NR⁷¹)₂ ; Y est choisi indépendamment parmi une liaison, O, NR⁵¹, S(O)_n, S(=NR⁷¹), S(O)(=NR⁷¹)₂ et S(=NR⁷¹)₂ ; n est choisi indépendamment parmi 0 à 2 ; et à la condition d'exclure les structures suivantes :



2. Composé selon la revendication 1, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel



est choisi parmi



étant non substitué ou substitué par 1 à 3 substituants choisis indépendamment dans le groupe constitué par F, Cl, Br, CN, OH, oxo, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-alkyle en C₁₋₄, O-halo-alkyle en C₁₋₄, NH₂, NH-alkyle en C₁₋₄, N(alkyl en C₁₋₄)₂, SO₂-alkyle en C₁₋₄ et SO₂-halo-alkyle en C₁₋₄.

3. Composé selon la revendication 1 ou 2, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel

B

est choisi dans le groupe constitué par phényle, naphtyle, pyridyle, pyrimidinyle, thiophényle, thiazolyle, cyclopentyle, cyclohexyle, bicyclo[1.1.1]pentyle, bicyclo[2.2.2]octyle, bicyclo[2.2.1]heptyle, pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octyle et pi-péridinyle,

le cycle étant non substitué ou substitué par 1 à 3 substituants choisis indépendamment dans le groupe constitué par F, Cl, Br, CN, OH, oxo, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-haloalkyle en C₁₋₄, alkyle en C₁₋₄-OH et haloalkyle en C₁₋₄-OH ; et où, éventuellement, deux substituants adjacents sur le noyau phényle forment ensemble un groupe -(CH₂)₃-, -(CH₂)₄-, -OCF₂O- et -OCH₂O-.

4. Composé selon l'une quelconque des revendications 1 à 3, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel


C

est choisi parmi un phényle, un pyridyle et un thiophényle ; le phényle, le pyridyle et le thiophényle étant non substitués ou substitués par 1 à 3 substituants choisis indépendamment dans le groupe constitué par F, Cl, CN, OH, oxo, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-alkyle en C₁₋₄ et O-halo-alkyle en C₁₋₄ ; et dans lequel le résidu -L-R¹ est lié dans l'orientation 1,3 par rapport à la liaison en direction de



et L n'est pas une liaison.

5. Composé selon l'une quelconque des revendications 1 à 4, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel
-L-R¹ est choisi parmi







dans lesquels le cycle est non substitué ou substitué en outre par 1 à 4 substituants choisis indépendamment dans le groupe constitué par F, Cl, Br, CN, OH, oxo, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-alkyle en C₁₋₄, O-halo-alkyle en C₁₋₄, alkyle en C₁₋₄-OH, haloalkyle en C₁₋₄-OH, SO₂-alkyle en C₁₋₄ et SO₂-halo-alkyle en C₁₋₄ ; et

dans lequel, éventuellement, deux substituants adjacents sur le noyau phényle forment ensemble un groupe -(CH₂)₃-, -(CH₂)₄-, -OCF₂O- et -OCH₂O-.


6. Composé selon l'une quelconque des revendications 1 à 5 dans lequel

R¹ est choisi parmi CO₂H, tétrazole, CH₂CO₂H, OCH₂CO₂H, SO₂CH₂CO₂H, CHMeCO₂H, CMe₂CO₂H, C(OH)MeCO₂H, CONHSO₂Me et CONH(OH) ; et éventuellement conjugué avec la glycine et avec la taurine de celui-ci, ou énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

7. Composé selon l'une quelconque des revendications 1 à 6 dans lequel
-L-R¹ est choisi parmi



et




et, éventuellement, conjugué avec la glycine et la taurine de celui-ci, ou énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.


8. Composé selon l'une quelconque des revendications 1 à 7, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel

D

est choisi dans le groupe constitué par



dans lesquels R^2 est choisi parmi Me, F, Cl, CN, Me, CHO, CHF_2 , CF_3 , SO_2Me ,



et

20

D


est en outre éventuellement également substitué par 1 à 2 substituants choisis dans le groupe constitué par F, Cl, CN, Me, OMe, CHO, CHF_2 et CF_3 .

- 25
9. Composé selon l'une quelconque des revendications 1 à 8, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel

30

D

est choisi dans le groupe constitué par




10. Composé selon l'une quelconque des revendications 1 à 9, où la formule (I) contient un substituant choisi dans le groupe constitué par CO_2H , tétrazole, CONHSO_2Me et $\text{CONH}(\text{OH})$; et éventuellement conjugué avec la glycine

et la taurine de celui-ci, ou énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

11. Composé selon l'une quelconque des revendications 1 à 10, dans lequel


-L-R¹ est choisi parmi



dans lesquels le phényle est non substitué ou substitué par 1 à 4 substituants choisis indépendamment dans le groupe constitué par F, Cl, CN, OH, Me et OMe ; et

R¹ est choisi parmi CO₂H et C(OH)MeCO₂H ; et éventuellement conjugué avec la glycine et la taurine de celui-ci, ou énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

12. Composé selon l'une quelconque des revendications 1 à 10, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel L-R¹ est



dans lequel le cycle est non substitué ou substitué par 1 à 4 substituants choisis indépendamment dans le groupe constitué par F, Cl, Br, CN, OH, oxo, alkyle en C₁₋₄, haloalkyle en C₁₋₄, O-alkyle en C₁₋₄, O-halo-alkyle en C₁₋₄, alkyle en C₁₋₄-OH, haloalkyle en C₁₋₄-OH, SO₂-alkyle en C₁₋₄ et SO₂-halo-alkyle en C₁₋₄ ; et dans lequel, éventuellement, deux substituants adjacents sur le noyau phényle forment ensemble un groupe -(CH₂)₃-, -(CH₂)₄-, -OCF₂O- et -OCH₂O-.

13. Composé selon l'une quelconque des revendications 1 à 12, dans lequel

R¹ est un alkylène en C₁₋₆-CO₂R⁴¹ ou un alkylène en C₁₋₆-CONR⁴¹R⁴², ou conjugué avec la glycine, ou conjugué avec la taurine, ou énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

14. Composé selon l'une quelconque des revendications 1 à 13, dans lequel


R¹ est COOH, ou conjugué avec la glycine, ou conjugué avec la taurine, ou énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

15. Composé selon l'une quelconque des revendications 1 à 13,

ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel R¹ est un alkylène en C₁₋₆-CONR⁴¹R⁴².

16. Composé selon la revendication 15, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, dans lequel R⁴¹ et R⁴² sont choisis indépendamment parmi H et un alkyle en C₁₋₄, l'alkyle en C₁₋₄ étant non substitué ou substitué par CO₂H.










17. Composé selon l'une quelconque des revendications 1 à 12, dans lequel L-R¹ est












ou conjugué avec la glycine ou conjugué avec la taurine de celui-ci, ou énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.


18. Composé selon l'une quelconque des revendications 1 à 17, le composé étant un conjugué avec la glycine, ou un énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

19. Composé selon l'une quelconque des revendications 1 à 18 choisi parmi






et




ou conjugué avec la glycine ou conjugué avec la taurine de celui-ci ; et énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

20. Composé selon l'une quelconque des revendications 1 à 19, qui est



un énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

21. Composé selon l'une quelconque des revendications 1 à 19, qui est




ou conjugué avec la glycine de celui-ci, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci.

22. Composé selon l'une quelconque des revendications 1 à 21, ou conjugué avec la glycine, conjugué avec la taurine, énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, destiné à être utilisé en tant que médicament.

23. Composé selon l'une quelconque des revendications 1 à 21, ou conjugué avec la glycine, conjugué avec la taurine,

énantiomère, diastéréoisomère, tautomère, *N*-oxyde, solvate ou sel pharmaceutiquement acceptable de celui-ci, destiné à être utilisé dans la prophylaxie et/ou le traitement de maladies se prêtant à être traitées par des modulateurs de LXR, la maladie étant choisie parmi une maladie du foie gras non alcoolique, une stéatose hépatique non alcoolique, une inflammation du foie, une fibrose hépatique, l'obésité, la résistance à l'insuline, le diabète de type II, une hypercholestérolémie familiale, une hypercholestérolémie associée au syndrome néphrotique, un syndrome métabolique, une stéatose cardiaque, un cancer, une myocardite virale, une infection au virus de l'hépatite C ou des complications associées, et des effets secondaires indésirables d'un traitement de longue durée aux glucocorticoïdes dans le cadre de maladies telle la polyarthrite rhumatoïde, une maladie inflammatoire de l'intestin et l'asthme.

24. Composition pharmaceutique comprenant un composé selon l'une quelconque des revendications 1 à 21, ou un conjugué avec la glycine, un conjugué avec la taurine, un énantiomère, un diastéréoisomère, un tautomère, un *N*-oxyde, un solvate ou un sel pharmaceutiquement acceptable de celui-ci, ainsi qu'un véhicule ou un excipient pharmaceutiquement acceptable.



5 Fig. 1: Differences between LXR agonists, antagonists and inverse agonists (NcoR = corepressor, NcoA = coactivator).

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2014085453 A [0011]
- WO 2003082802 A [0013]
- WO 2016207217 A [0016]
- WO 2016106266 A [0017]
- WO 2013111150 A [0018]
- WO 2013028999 A [0019]
- WO 2013012649 A [0020]
- WO 2010124793 A [0021]
- WO 2008132434 A [0021]
- WO 2010010186 A [0022]
- WO 2009032116 A [0023]
- WO 2009032125 A [0023]
- WO 2009064848 A [0023]
- WO 2009064852 A [0023]
- WO 2008116833 A [0024]
- WO 2008003736 A [0025]
- WO 2007075555 A [0026]
- WO 2004000831 A [0026]
- WO 2007134169 A [0027]
- WO 2006050236 A [0027]
- WO 2005034941 A [0028]
- WO 200514000 A [0029]
- WO 200251837 A [0030]
- WO 200236562 A [0030]
- WO 200232863 A [0031]
- WO 9921851 A [0032]
- WO 9857931 A [0033]
- WO 9822452 A [0033]
- WO 9822457 A [0034]
- WO 2001130343 A [0035]
- WO 200046199 A [0035]
- WO 200046197 A [0035]
- WO 200046195 A [0035]
- JP 06145150 B [0035]
- EP 0535926 A [0035]
- EP 0535925 A [0035]
- WO 2008119657 A [0036]

Non-patent literature cited in the description

- ESTES et al. *Hepatology*, 2018, vol. 67, 123 [0004]
- ESTES et al. *J. Hepatol.*, 2018, vol. 69, 896 [0004]
- DAY et al. *Gastroenterology*, 1998, vol. 114, 842 [0004]
- AHN et al. *Dig. Dis. Sci.*, 2014, vol. 59, 2975 [0005]
- YANG et al. *J. Biol. Chem.*, 2006, vol. 281, 27816 [0005]
- FEET et al. *Cell*, 1998, vol. 93, 693 [0006]
- SCHULTZ et al. *Genes Dev.*, 2000, vol. 14, 2831 [0006]
- KIRCHGEISSNER et al. *Cell Metab.*, 2016, vol. 24, 223 [0006]
- FLAVENY et al. tested the LXR inverse agonist tool compound SR9243 in cancer cells and in animal cancer models. *Cancer Cell*, 2015, vol. 28, 42 [0009]
- ZUERCHER et al. *J. Med. Chem.*, 2010, vol. 53, 3412 [0010]
- GRIFFETT ; BURRIS. *Biochem. Biophys. Res. Commun.*, 2016, vol. 479, 424 [0010]
- GRIFFETT et al. *ACS Chem. Biol.*, 2013, vol. 8, 559 [0012]
- GRIFFETT et al. *Mol. Metab.*, 2015, vol. 4, 35 [0012]
- FLAVENY et al. *Cancer Cell*, 2015, vol. 28, 42 [0012]
- HUANG et al. *BioMed Res. Int.*, 2018, 8071093 [0012]
- COLLINS et al. *J. Med. Chem.*, 2002, vol. 45, 1963 [0014]
- YU et al. *J. Org. Chem.*, 2018, vol. 83, 323 [0015]
- FOSTER. *Trends Pharmacol. Sci.*, 1984, vol. 5, 524 [0105]
- KREMOSER et al. *Drug Discov. Today*, 2007, vol. 12, 860 [0125]
- GRONEMEYER et al. *Nat. Rev. Drug Discov.*, 2004, vol. 3, 950 [0125]
- LIBERTI et al. *Trends Biochem. Sci.*, 2016, vol. 41, 211 [0129]
- WARD ; THOMPSON. *Cancer Cell*, 2012, vol. 21, 297-308 [0129]
- TRAVERSARI et al. *Eur. J. Immunol.*, 2014, vol. 44, 1896 [0129]
- FLAVENY et al. *Cancer Cell.*, 2015, vol. 28, 42 [0130]
- STEFFENSEN. *Cancer Cell*, 2015, vol. 28, 3 [0130]
- PATEL et al. *Endocrinology*, 2017, vol. 158, 1034 [0132]
- GARCÍA-MEDIAVILLA et al. *Lab. Invest.*, 2012, vol. 92, 1191 [0133]
- PAPAGEORGIOU et al. *Cardiovasc. Res.*, 2015, vol. 107, 78 [0134]
- ZHENG et al. *PLoS One*, 2014, vol. 9, e101269 [0135]

EP 3 814 331 B9

- **ZHOU et al.** *J. Biol. Chem.*, 2008, vol. 283, 2129 [0136]
- **LIU ; VAZIZI.** *Nephrol. Dial. Transplant.*, 2014, vol. 29, 538 [0137]
- *ACS Med. Chem. Lett.*, 2016, vol. 7, 1207 [0295]
- **GRIESBRECHT et al.** *Synlett*, 2010, 374 [0476]
- **FAUCHER et al.** *J. Med. Chem.*, 2004, vol. 47, 18 [0476]
- **WANG et al.** *Nat. Rev. Mol. Cell Biol.*, 2015, vol. 16, 678 [0487]