



(11) **EP 3 821 982 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 153(4) EPC

(43) Date of publication:
19.05.2021 Bulletin 2021/20

(51) Int Cl.:
B01L 7/00 (2006.01) B01J 19/00 (2006.01)

(21) Application number: **19834598.5**

(86) International application number:
PCT/KR2019/008663

(22) Date of filing: **12.07.2019**

(87) International publication number:
WO 2020/013664 (16.01.2020 Gazette 2020/03)

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
Designated Extension States:
BA ME KH MA MD TN

- **HA, Tai Hwan**
Daejeon 34141 (KR)
- **KIM, Kyung Ho**
Daejeon 34141 (KR)
- **KIM, Jin Yeong**
Daejeon 34141 (KR)
- **PARK, Seon Joo**
Daejeon 34141 (KR)
- **SEO, Seong Eun**
Daejeon 34141 (KR)

(30) Priority: **12.07.2018 KR 20180081088**

(71) Applicant: **Korea Research Institute of Bioscience and Biotechnology**
Daejeon 34141 (KR)


(74) Representative: **Berggren Oy**
P.O. Box 16
Eteläinen Rautatiekatu 10A
00101 Helsinki (FI)

(72) Inventors:
• **KWON, Oh Seok**
Daejeon 34141 (KR)

(54) **SUBSTRATE FOR NUCLEIC ACID AMPLIFICATION, AND METHOD FOR MANUFACTURING SAME**

(57) The present invention relates to a substrate for nucleic acid amplification, and a method for manufacturing same, the substrate for rapid and accurate PCR analysis comprising: a transparent substrate; a metal layer formed on the transparent substrate; an N-heterocyclic carbene compound having one end annealed to the surface of the metal layer; and a primer immobilized on the other end of the N-heterocyclic carbene compound.

FIG. 1



EP 3 821 982 A1

Description**TECHNICAL FIELD**

5 [0001] The present invention relates to a substrate for nucleic acid amplification, and a method for manufacturing same.

BACKGROUND ART

10 [0002] A polymerase chain reaction (PCR) method is technique allowing amplification of a specific region of DNA or RNA to a large amount in vitro. A PCR machine using such technique is used for the amplification of a small amount of DNA included in blood in hospitals, research centers, etc., and the PCR machine is also used for diagnosing various diseases including malaria, tuberculosis, hepatitis, etc.

15 [0003] PCR is conducted through one cycle of three steps including denaturation, annealing and elongation, and the rapid rise and drop of temperature in line with the temperature suitable for each step is very important for reducing PCR reaction time. That is, PCR requires the instant shift of the temperature in line with the temperature set for each step, but if time delay is generated during the process for changing the temperature to the preset temperature, a reaction rate is degraded, total PCR reaction time increases, and there are problems of arising unnecessary reactions to produce by-products. Particularly, the decrease of the PCR time enables prompt diagnosis on the spot of infectious diseases which are recently spread rapidly, and plays an important role of serving early disinfection.

20 [0004] However, the conventionally used PCR machine uses electrical energy as a thermal energy source, and a long time of about 1 hour or more is required for completing nucleic acid amplification reaction, and since accessories including a heating instrument and a heat sink are necessary, there are problems of inconvenience in that the apparatuses occupy much space. Accordingly, the application on the spot requiring the prompt diagnosis of diseases is a little bit inconvenient. In addition, Korean Laid-open Patent Publication No. 2017-0106995 discloses a light energy-based PCR system, but
25 there are problems in that PCR is performed in a liquid phase, multiple diagnoses in one chamber is impossible, and the recovery of a used primer is impossible. Particularly, due to a high price, the enlargement of the apparatus and long detection time (1 hour or more), the conventional PCR machine is inconvenient to carry, and immediate diagnosis on the spot is limited.

30 [0005] Therefore, research on PCR for stable, rapid and precise diagnosis irrespective of high temperature and rapid change of the temperature, is steadily being conducted.

[Prior Art Document]

35 [0006] (Patent Document 1) Korean Laid-open Patent Publication No. 2017-0106995

DISCLOSURE OF THE INVENTION**TECHNICAL PROBLEM**

40 [0007] The present invention has been devised to solve the above-described problems and provides a substrate for nucleic acid amplification serving stable and rapid multiple PCR reactions by immobilizing a primer on the surface of a metal layer using an N-heterocyclic carbene compound as a linker, while amplifying nucleic acid using thermal energy generated by irradiating light energy.

TECHNICAL SOLUTION

45 [0008] The present invention provides a substrate for nucleic acid amplification including a transparent matrix, a metal layer formed on the transparent matrix, an N-heterocyclic carbene compound having one end bonded to the surface of the metal layer and a primer immobilized on the other end of the N-heterocyclic carbene compound.

50 [0009] In addition, the present invention provides a method for manufacturing a substrate for nucleic acid amplification, including forming a metal layer on a transparent matrix, introducing one end of an N-heterocyclic carbene compound onto the surface of the metal layer, and immobilizing a primer on the other end of the N-heterocyclic carbene compound.

ADVANTAGEOUS EFFECTS

55 [0010] In case of using the substrate for nucleic acid amplification of the present invention in polymerase chain reaction (PCR), by immobilizing a primer after annealing an N-heterocyclic carbene compound on the surface (solid phase) of a metal layer, PCR reaction may be more stable at high temperature and completed in a very short time when compared

with a case of using a linker by the bond between a metal and thiol (-SH).

[0011] In addition, according to the PCR reaction on a solid phase (transparent matrix/metal layer), the recovery of a primer after the PCR reaction is easy, and various types of primers could be immobilized on the N-heterocyclic carbene compound at the surface of each metal layer, and thus, the simultaneous amplification of diverse nucleic acids is possible, and there are advantages of performing multiple PCR reactions.

BRIEF DESCRIPTION OF THE INVENTION

[0012]

FIG. 1 is a diagram schematically showing a series of light-based PCR reaction using the substrate for nucleic acid amplification of the present invention.

FIG. 2 is a diagram schematically showing a light-based PCR machine on a solid phase using the substrate for nucleic acid amplification of the present invention.

FIG. 3 is a diagram showing a gold (Au) layer formed on a transparent matrix included in the substrate for nucleic acid amplification of the present invention.

FIG. 4 is a diagram showing ¹H-NMR spectrum of N-heterocyclic carbene compound 1 prepared according to Preparation Example 1 of the present invention.

FIG. 5 is a diagram showing ¹H-NMR spectrum of N-heterocyclic carbene compound 2 prepared according to Preparation Example 2 of the present invention.

FIG. 6 is a diagram showing ¹H-NMR spectrum of N-heterocyclic carbene compound 3 prepared according to Preparation Example 3 of the present invention.

FIG. 7 is a diagram showing ¹H-NMR spectrum of N-heterocyclic carbene compound 4 prepared according to Preparation Example 4 of the present invention.

FIG. 8 is a diagram showing ¹H-NMR spectrum of N-heterocyclic carbene compound 5 prepared according to Preparation Example 5 of the present invention.

FIG. 9 is a diagram showing time required for performing 40 cycles of PCR reaction according to Examples 1 and 2 of the present invention.

FIG. 10 is a diagram showing XPS elemental analysis results of a gold (Au) layer on which N-heterocyclic carbene compound 4 bonded according to Example 1 of the present invention.

FIG. 11 is a diagram schematically showing a series of process for immobilizing a primer on a gold (Au) layer using N-heterocyclic carbene compound 4 according to the present invention.

FIG. 12 is a diagram schematically showing a series of process for immobilizing a primer on a gold (Au) layer using N-heterocyclic carbene compound 4 of the present invention.

FIG. 13 is a diagram showing a transparent matrix and a metal layer formed on the transparent matrix according to an embodiment of the present invention.

FIG. 14 is a diagram showing a transparent matrix and a metal layer formed on the transparent matrix according to an embodiment of the present invention.

FIG. 15 is a diagram showing time required for performing 40 cycles of PCR reaction according to Example 3 of the present invention.

FIG. 16 is a diagram showing detection results using a fluorescence material after performing PCR reaction according to Example 1 of the present invention.

FIG. 17 is a diagram showing detection results using a fluorescence material after performing PCR reaction according to Example 2 of the present invention.

FIG. 18 is a diagram showing detection results using a fluorescence material after performing PCR reaction according to Example 3 of the present invention.

FIG. 19 is a diagram showing detection results using a fluorescence material after performing PCR reaction according to Comparative Example 1 of the present invention.

FIG. 20 is a diagram showing a transparent matrix including a gold mesh according to an embodiment of the present invention.

FIG. 21 is a diagram showing time required for performing 40 cycles of PCR reaction in case of including a metal mesh as in FIG. 20.

FIG. 22 is a diagram showing electrophoretic results confirming the completion of PCR reaction after performing PCR reaction according to Example 1 of the present invention.

[Explanation on symbols]

[0013]

10: transparent matrix
 20: metal layer
 30: thermocouple

5 MODE FOR CARRYING OUT THE INVENTION

[0014] Hereinafter, the present invention will be explained in detail.

[0015] The present invention provides a substrate for nucleic acid amplification including: a transparent matrix; a metal layer formed on the transparent matrix; an N-heterocyclic carbene compound having one end bonded to the surface of the metal layer; and a primer immobilized on the other end of the N-heterocyclic carbene compound.

[0016] The substrate for nucleic acid amplification may be used in polymerase chain reaction (PCR) which is molecular biological technique for duplicating and amplifying a desired segment of DNA or RNA.

[0017] The first step of PCR is a step for denaturing DNA (or RNA). Two strands of DNA may be separated by heating, and each separated DNA plays the role of a template. The denaturation temperature is generally 90°C to 96°C, but is changed according to the amount of bases G+C in DNA and the length of DNA. The second step of PCR is a step of annealing. In this step, two types of primers are annealed to each complementary template DNA. The annealing temperature is an important factor for determining the accuracy of reaction, and if the temperature is too high, the primer may be annealed on DNA too weakly, and the product of amplified DNA may be very small. If the temperature is too low, the primer may be non-specifically annealed, and undesired DNA may be amplified. General annealing temperature is 50°C to 65°C. The third step of PCR is a step of elongation. In this step, DNA polymerase resistant to heat produces new DNA in template DNA. In this case, the elongation temperature is 70°C to 75°C. As described above, there are a series of three steps in the PCR reaction, and if the three steps are set to 1 cycle, the PCR reaction is performed by about 30 to 40 cycles.

[0018] The substrate for nucleic acid amplification may be used in light-based PCR using thermal energy produced by the irradiation of light energy during the above-described PCR.

[0019] The light-based PCR may use plasmonic photothermal conversion by the interaction of photons, electrons and phonons at the surface of a metal. Particularly, if photons reach the surface of a metal layer from an excited energy source, light-absorption occurs, and electrons around the surface are excited to a higher state to form electrons of high temperature. According to the rapid diffusion and uniform distribution of such electrons of high temperature to the whole metal layer, a metal layer of high temperature may heat surrounding solutions. In addition, the electrons of high temperature may be cooled again by energy exchange with lattice phonons.

[0020] The temperature of the metal layer in which plasmons are excited as described above may be elevated to maximum 500°C, and a PCR specimen solution around the metal layer may be heated to 150°C or more in a short time.

[0021] The transparent matrix may be manufactured so that light irradiated from a light source positioned under a counter surface to the metal layer formed is transferred to the metal layer without loss, and adhesiveness with the metal layer is excellent.

[0022] The light source is not limited only if generating ultraviolet rays, visible rays or infrared rays, and may use a halogen lamp, an LED lamp, a fluorescent lamp, an incandescent lamp, an arc source lamp, an infrared lamp, an HMI lamp, a UV lamp, etc., preferably, an LED lamp considering power efficiency and economic feasibility.

[0023] The transparent matrix may be formed using a transparent material so as to transmit light irradiated, and may be formed using a material which is hardly deformed by light or heat. Preferably, a glass matrix, a plastic matrix, a silicon matrix, etc., may be used, but is not limited thereto.

[0024] The transparent matrix has a constant thickness, and the thickness of the transparent matrix may be from 0.1 mm to 10 mm, preferably, from 0.5 mm to 5 mm, more preferably, from 0.5 mm to 1.5 mm. If the thickness of the transparent matrix satisfies the above-described range, thermal energy by light energy irradiated from a light source may be efficiently transferred, and a PCR reaction cycle may be optimized.

[0025] The transparent matrix may include a two-dimensional matrix of a planar type; a three-dimensional matrix selected from a spherical type, a hemispherical type, a polyhedral type, a polyprism type, or a cylindrical type; or a mixture type thereof. For example, as in FIG. 13 and FIG. 14, the transparent matrix may include a matrix of a mixture type of a three-dimensional matrix in which at least one partial surface of the polyhedral type matrix includes a hemispherical type matrix.

[0026] In case where the transparent matrix includes the three-dimensional matrix as described above, PCR reaction may be performed at many faces, and there are advantages in that diverse nucleic acids may be amplified simultaneously, and multiple diagnosis may be possible.

[0027] The substrate for nucleic acid amplification may utilize a metal layer for utilizing thermal energy by the irradiation of light energy. In case of using the metal layer, the temperature may be instantly elevated to maximum 500°C, and the temperature change in the temperature range (from about 50°C to 95°C) of the PCR cycle may be promptly conducted.

[0028] The metal layer may include a metal thin film.

[0029] The metal layer may further include a metal mesh. In case where the metal layer further includes a metal mesh, the metal mesh may be formed first on the transparent matrix, and then the metal layer may be formed on the metal mesh. In case where the metal layer further includes the metal mesh, prompt heating and cooling may be possible when compared with a case of using only a metal layer, the temperature change of a PCR cycle may be promptly conducted, and more efficient PCR reaction may be performed.

[0030] As in FIG. 20, if a metal mesh is provided, the temperature change in the temperature range (from about 50°C to 95°C) of the PCR cycle may be promptly conducted, and as in FIG. 21, it could be confirmed that the time required for performing 40 cycles of the PCR reaction is significantly short and about 7 minutes.

[0031] In case where the metal layer is formed on the transparent matrix, the metal layer may be formed on at least one face of the transparent matrix. In this case, the metal layer may be formed on the whole face of the at least one face of the transparent matrix, or may be partially formed on the at least one face of the transparent matrix. That is, on the transparent matrix, one metal layer may be formed, or multiple metal layers may be formed.

[0032] The metal mesh may have a two-dimensional or three-dimensional structure, and in this case, the mesh may include a polygonal pattern including a triangle, a tetragon, a pentagon and hexagon, or a circular pattern.

[0033] In case where the mesh includes a polygonal pattern, the length of one side of the mesh may be from 50 to 200 μm, and in case where the mesh includes a circular pattern, the diameter of the mesh may be from 50 to 200 μm. In addition, the spacing of the mesh may be from 1 to 5 μm, particularly, 1 to 2 μm. The thickness of the metal mesh may be from 10 nm to 200 nm, particularly, from 100 to 200 nm. In case where the thickness of the metal mesh and the size of the mesh satisfy the above-described ranges, prompt heating and cooling may become possible, and the temperature change of the PCR cycle may be promptly conducted when compared with a case of using only a metal layer.

[0034] The metal layer may include any one selected from the group consisting of copper (Cu), silver (Ag), gold (Au), palladium (Pd), platinum (Pt), rhodium (Rh) and combinations thereof (for example, bimetallic nanoparticles). Preferably, gold (Au) which has excellent bonding properties with an N-heterocyclic carbene compound and stability and rapid light-absorption, may be included.

[0035] The method of coating the metal layer on one side of the transparent matrix may use any coating or deposition technique without limitation, and the transparent matrix may be formed into a uniform thickness by a method of chemical vapor deposition (CVD), physical vapor deposition (PVD), thermal evaporation deposition, sputtering deposition or atomic layer deposition (ALD).

[0036] The thickness of the metal layer may be from 10 nm to 200 nm. If the thickness of the metal layer is greater than the range, there are problems in that the metal layer may be damaged during the step of immobilizing a linker, or a linker may be non-uniformly formed.


[0037] A gold (Au) layer formed on the transparent matrix is shown in FIG. 3.

[0038] As a linker for immobilizing the primer on the surface of the metal layer, an N-heterocyclic carbene compound may be used. The N-heterocyclic carbene compound may be introduced onto the surface of the metal layer through a metal-carbene bond.


[0039] Particularly, in case of a linker having a thiol (-SH) group used in the conventional solid phase PCR, there are problems of degrading the reproducibility of PCR reaction results due to the unstability (cleaving of a metal-sulfur bond) of a metal-sulfur bond at high temperature (about 90°C). In contrast, in case of using the N-heterocyclic carbene compound as a linker for immobilizing the primer onto the metal layer, though the surface temperature of the metal layer temporarily rises to maximum 500°C by a light source, stability at high temperature may be achieved due to a metal-carbene bond, and the application thereof to a PCR machine using light energy is possible.

[0040] The N-heterocyclic carbene compound may be represented by Chemical Formula 1 or 2 below.

[Chemical Formula 1]



[Chemical Formula 2]



[0041] In Chemical Formulae 1 and 2,


R1, R2, R5 and R6 are the same or different, and are each independently hydrogen, an alkyl group of 1 to 20 carbon atoms, a cycloalkyl group of 3 to 20 carbon atoms, an aryl group of 6 to 30 carbon atoms or a heteroaryl group of 2 to 30 carbon atoms,

R3, R4, R7, R8, R9 and R10 are the same or different, and are each independently hydrogen, an alkyl group of 1 to 20 carbon atoms, a cycloalkyl group of 3 to 20 carbon atoms, an aryl group of 6 to 30 carbon atoms, a heteroaryl group of 2 to 30 carbon atoms or a structure represented by Chemical Formula 3 below, where adjacent two or more substituents among R7 to R10 are bonded to form a hydrocarbon ring,

at least one of R3 and R4 is the structure represented by Chemical Formula 3 below, and

at least one of R7 to R10 is substituted with the structure represented by Chemical Formula 3 below, or in case where adjacent two or more substituents among R7 to R10 are bonded to form a hydrocarbon ring, at least one of carbon atoms forming the hydrocarbon ring is substituted with the structure represented by Chemical Formula 3 below,

[Chemical Formula 3]



in Chemical Formula 3,

n is a repeating number of a unit in parenthesis and an integer of 1 to 30, and

A is a nitrogen (N)-containing alkyl group of 1 to 20 carbon atoms or a nitrogen (N)-containing heteroaryl group of 2 to 30 carbon atoms.

[0042] R1, R2, R5 and R6 may be the same or different, and may be each independently hydrogen, an alkyl group of 1 to 20 carbon atoms or an aryl group of 6 to 30 carbon atoms.

[0043] R1, R2, R5 and R6 may be the same or different, and may be each independently hydrogen, an alkyl group of 1 to 10 carbon atoms or an aryl group of 6 to 10 carbon atoms.

[0044] R1, R2, R5 and R6 may be the same or different, and may be each independently hydrogen or an alkyl group of 1 to 10 carbon atoms.

[0045] R1, R2, R5 and R6 may be the same or different, and may be each independently hydrogen, isopropyl or benzyl.

[0046] At least one of R1 and R2 and at least one of R5 and R6 may be the same or different, and may be each independently an alkyl group of 1 to 20 carbon atoms or an aryl group of 6 to 30 carbon atoms.

[0047] At least one of R1 and R2 and at least one of R5 and R6 may be the same or different, and may be each independently isopropyl or benzyl.

[0048] R3, R4, R7, R8, R9 and R10 may be the same or different, and may be each independently hydrogen, an alkyl group of 1 to 20 carbon atoms, a cycloalkyl group of 3 to 20 carbon atoms, an aryl group of 6 to 30 carbon atoms, a heteroaryl group of 2 to 30 carbon atoms or the structure represented by Chemical Formula 3, or two or more adjacent

substituents among R7 to R10 may be bonded to form a hydrocarbon ring.

[0049] R3, R4, R7, R8, R9 and R10 may be the same or different, and may be each independently hydrogen or the structure represented by Chemical Formula 3, or two or more adjacent substituents among R7 to R10 may be bonded to form a hydrocarbon ring.

[0050] In case where two or more adjacent substituents among R7 to R10 are bonded to form a hydrocarbon ring, at least one carbon forming the hydrocarbon ring may be substituted with the structure represented by Chemical Formula 3.

[0051] At least one of R3 and R4 may be the structure represented by Chemical Formula 3. In addition, at least one of R7 to R10 may be the structure represented by Chemical Formula 3.

[0052] The n is a repeating number of the unit in parenthesis and may be an integer of 1 to 30, preferably, 1 to 10. More preferably, 1 to 3.

[0053] The A is a nitrogen (N)-containing alkyl group of 1 to 20 carbon atoms, or a nitrogen (N)-containing heteroaryl group of 2 to 30 carbon atoms. Particularly, A may be azide, phthalimide, or amine.

[0054] In the present invention, an "adjacent" group may mean a substituent directly coupled with an element at which a corresponding substituent is substituted, a sterically most nearly positioned substituent to the corresponding substituent, or another substituent substituted at an element at which the corresponding substituent is substituted. For example, two substituents substituted at ortho position in a benzene ring, and two substituents substituted at the same carbon in an aliphatic ring may be interpreted as the "adjacent" group.

[0055] The alkyl group may be a linear chain or a branched chain, and the carbon number may be 1 to 20, preferably, 1 to 10. More preferably, the carbon number may be 1 to 6. Particular examples of the alkyl group may include methyl, ethyl, propyl, n-propyl, isopropyl, butyl, n-butyl, isobutyl, tert-butyl, sec-butyl, 1-methylbutyl, 1-ethylbutyl, pentyl, n-pentyl, isopentyl, neopentyl, tert-pentyl, hexyl, n-hexyl, 1-methylpentyl, 2-methylpentyl, 4-methyl-2-pentyl, 3,3-dimethyl butyl, 2-ethylbutyl, heptyl, n-heptyl, 1-methylhexyl, cyclopentylmethyl, cyclohexylmethyl, octyl, n-octyl, tert-octyl, 1-methylheptyl, 2-ethylhexyl, 2-propylpentyl, n-nonyl, 2,2-dimethylheptyl, 1-ethylpropyl, 1,1-dimethylpropyl, isohexyl, 4-methylhexyl, 5-methylhexyl, benzyl, etc., but is not limited thereto.

[0056] The cycloalkyl group may have 3 to 20 carbon atoms, preferably, 3 to 10 carbon atoms. Particular examples of the cycloalkyl group may include cyclopropyl, cyclobutyl, cyclopentyl, 3-methylcyclopentyl, 2,3-dimethylcyclopentyl, cyclohexyl, 3-methylcyclohexyl, 4-methylcyclohexyl, 2,3-dimethylcyclohexyl, 3,4,5-trimethylcyclohexyl, 4-tert-butylcyclohexyl, cycloheptyl, cyclooctyl, etc., but is not limited thereto.

[0057] The aryl group may have 6 to 30 carbon atoms, preferably, 6 to 10 carbon atoms. The aryl group may be a monocyclic aryl group or a polycyclic aryl group. Particular examples of the monocyclic aryl group may include a phenyl group, a biphenyl group, a terphenyl group, etc., and particular examples of the polycyclic aryl group may include a naphthyl group, an anthracenyl group, a phenanthryl group, a pyrenyl group, a perylenyl group, a chrysenyl group, a fluorenyl group, a triphenylene group, etc., but is not limited thereto.

[0058] The heteroaryl group is an aromatic ring group including one or more selected from N, O, P, S, Si and Se as heteroatoms, and the carbon number may be 2 to 30, preferably, 2 to 20. Particular examples of the heteroaryl group may include a thiophene group, a furan group, a pyrrole group, an imidazole group, a thiazole group, an oxazole group, an oxadiazole group, a triazole group, a pyridyl group, a pyrimidyl group, a triazine group, a triazole group, an acridyl group, a quinoliny group, a quinazoline group, a quinoxaliny group, a phthalazinyl group, an isoquinoline group, an indole group, a carbazole group, a benzoxazole group, a benzimidazole group, a benzothiazole group, a benzocarbazole group, a benzothiophene group, a dibenzothiophene group, a benzofuranyl group, etc., but is not limited thereto.

[0059] In addition, the alkyl group, the cycloalkyl group, the aryl group, the heteroaryl group or the hydrocarbon ring may be further substituted or unsubstituted with an alkyl group, a cycloalkyl group, an aryl group or a heteroaryl group.

[0060] On the surface of the metal layer, one end of the N-heterocyclic carbene compound may be bonded, and in this case, a metal-carbene bond may be formed by a method of chemical vapor deposition (CVD), physical vapor deposition (PVD), thermal evaporation deposition, sputtering deposition, atomic layer deposition (ALD) or chemical-bath deposition (CBD).

[0061] According to an embodiment of the present invention, a series of mechanism of bonding one end of the N-heterocyclic carbene compound on the surface of the metal layer by a chemical vapor deposition method is shown in FIG. 11, and a series of mechanism of bonding one end of the N-heterocyclic carbene compound on the surface of the metal layer by a chemical-bath deposition method is shown in FIG. 12.

[0062] At the other end of the N-heterocyclic carbene compound, a primer may be bonded and immobilized.

[0063] The other end of the N-heterocyclic carbene compound may be functionalized with azide, phthalimide, or amine.

[0064] If the other end of the N-heterocyclic carbene compound is functionalized with azide, phthalimide, or amine, a primer may be bonded to the azide, phthalimide, or amine using click reaction as shown in FIGS. 11 and 12. Accordingly, in contrast to the general bonding of the primer with the metal layer by an electrostatic attraction, the primer may be immobilized onto the metal layer through a stronger chemical bond, and if used in the PCR reaction of a solid phase of high temperature, effects of excellent stability, and improved storability and ease of storage may be achieved.

[0065] The primer may be a primer of at least one type. The primer of at least one type may mean any one selected

from primers which may make complementary bond with different amplification target template DNA (or RNA).

[0066] The primer may be a primer of two or more different types. In case where the primer is the primer of two or more different types, any one among a pair of primers which may make complementary bond with a first template DNA (or RNA) to be amplified, and any one among a pair of primers which may make complementary bond with a second template DNA (or RNA) to be amplified. Here, the first template DNA (or RNA) and the second template DNA (or RNA) may mean DNA (or RNA) having different base sequences.

[0067] As described above, by immobilizing a primer capable of elongating different template DNA (or RNA) onto the other end of the N-heterocyclic carbene compound which is bonded onto the surface of the metal layer, diverse DNA (or RNA) may be amplified at the same time, and the diagnosis by multiple PCR reactions may be possible in one chamber.

[0068] The present invention also provides a method for manufacturing a substrate for nucleic acid amplification, including: forming a metal layer on a transparent matrix; introducing one end of an N-heterocyclic carbene compound onto the surface of the metal layer; and immobilizing a primer on the other end of the N-heterocyclic carbene compound.

[0069] FIG. 1 is a diagram schematically showing a series of a process of light-based PCR reaction using the substrate for nucleic acid amplification of the present invention.

[0070] The schematic explanation on a process for performing the light energy-based PCR reaction using the substrate for nucleic acid amplification of the present invention is as follows, but the PCR reaction is not limited only to the method described below.

[0071] A PCR specimen including amplification target template DNA, dNTP, Taq polymerase, etc., is prepared. Two types of primers (forward primer, reverse primer) making complementary bond with the template DNA are prepared. A portion of the forward primer is immobilized by bonding with the end of the transparent matrix/metal layer/N-heterocyclic carbene compound linker, which is the substrate for nucleic acid amplification of the present invention, and the remaining forward primer and reverse primer are included in the PCR reaction specimen. Then, light energy is irradiated by a light source (LED, etc.) positioned under the transparent matrix, and PCR reaction is performed.

[0072] In the conventional PCR substrate in which a primer is immobilized on the surface of a metal layer by a metal-sulfur bond using a thiol (-SH) group as a linker, there are problems in that the metal-sulfur bond is liable at high temperature (about 90°C or more), and the linker is easily separated. Accordingly, the application thereof to light energy-based PCR by which the surface temperature by the light energy is elevated to 500°C, is impossible, and the application thereof is possible only to a PCR machine using thermal energy converted from electrical energy.

[0073] However, in case of performing PCR reaction using the substrate for nucleic acid amplification of the present invention, the N-heterocyclic carbene compound is attached onto the surface (solid phase) of the metal layer as a linker, and then the primer is immobilized, and accordingly, stability at high temperature is better in contrast to a linker by the bonding of the metal and thiol (-SH), and thus, the application to a PCR machine using thermal energy converted from light energy is possible, prompt temperature control of the metal layer by the irradiation of light energy is possible, and the PCR reaction may be completed within a very short time with about 12 minutes to 15 minutes for performing 40 cycles for the PCR reaction.

[0074] In addition, as described above, since the amplification of diverse nucleic acids at the same time is possible by using the primer of at least one type (or a primer of different two types), there are advantages in that multiple PCR reactions may be performed and more precise diagnosis may be served. In addition, different from PCR which is performed only in a liquid phase, the substrate for nucleic acid amplification may be easily recovered after the PCR reaction, and the storage of the primer is easy, and the regeneration thereof is possible.


[0075] Hereinafter, the present invention will be explained in more detail referring to preferred embodiments.


[0076] However, these embodiments are for explaining the present invention more particularly, and the scope of the present invention is not limited thereto.

<Preparation Example>

<Preparation Example 1>

[0077]





[0078] According to the reaction above, N-heterocyclic carbene compound 1 was prepared.

[0079] In FIG. 4, the $^1\text{H-NMR}$ spectrum of N-heterocyclic carbene compound 1 is shown.

25

<Preparation Example 2>

[0080]

30


35

40


45

50

55



5




10 [0081] According to the reaction above, N-heterocyclic carbene compound 2 was prepared.

[0082] In FIG. 5, the $^1\text{H-NMR}$ spectrum of N-heterocyclic carbene compound 2 is shown.

<Preparation Example 3>


15 [0083]

20




25

30




35

40



45

50




55 [0084] According to the reaction above, N-heterocyclic carbene compound 3 was prepared.

[0085] In FIG. 6, the $^1\text{H-NMR}$ spectrum of N-heterocyclic carbene compound 3 is shown.

<Preparation Example 4>


[0086]

5




10

15




20




25

30



35



40

[0087] According to the reaction above, N-heterocyclic carbene compound 4 was prepared.


[0088] In FIG. 7, the ¹H-NMR spectrum of N-heterocyclic carbene compound 4 is shown.

45


<Preparation Example 5>

[0089]

50



55



[0090] According to the reaction above, N-heterocyclic carbene compound 5 was prepared.

[0091] In FIG. 8, the ¹H-NMR spectrum of N-heterocyclic carbene compound 5 is shown.

<Example>

<Example 1>

[0092] As template DNA, a lung cancer cell line, cDNA of A549 was used, and a PCR reactant of a liquid phase including dNTP, Taq polymerase, a reverse primer below and a forward primer below was prepared.

[0093] By using N-heterocyclic carbene compound 4 prepared in Preparation Example 4, a forward primer below was immobilized on the surface of a gold (Au) layer deposited by CVD on a planar type glass matrix. The substrate on which the forward primer was immobilized was put in a PCR reactant of a liquid phase, which included a relatively small amount of the forward primer, and by using the machine of FIG. 2, PCR reaction of 40 cycles was performed. The time required for performing 40 cycles of the PCR reaction is shown in FIG. 9, and 5 μl of the PCR reaction resultant was loaded on 1 wt% of an agarose gel, and electrophoresis (Mupid 2Plus of GenomicBase Co.) was performed under conditions of 100 V and 20 minutes. In comparison to the conventional RT PCR reaction resultant, the completion of the PCR reaction using the PCR substrate of the present invention was confirmed through FIG. 22.

[Primer immobilized onto substrate of solid phase]

[0094]

- Forward primer: 5'-Amino(C12)-TTTTTTTTTTGACCCAATCATGAGCACTGCTTT-3'

[Primer in PCR reactant of liquid phase]

[0095]

- 5 - Forward primer: 5'-GACCCAATCATGAGCACTG-3'
- Reverse primer: 5'-TGAAGCGACCCTCTGATG-3'

<Example 2>

10 **[0096]** In Example 1, PCR reaction was performed by the same method in Example 1 except for using a mutated lung cancer cell line, cDNA of NCI-H-1975 instead of the lung cancer cell line, cDNA of A549, as the template DNA.

<Example 3>

15 **[0097]** In Example 1, PCR reaction was performed by the same method in Example 1 except for using a hexahedral glass matrix as in FIG. 13 and FIG. 14 instead of the planar type glass matrix.

<Comparative Example 1>

20 **[0098]** PCR reaction was performed by the same method in Example 1 except for performing the PCR reaction without the template DNA.

[0099] The PCR products of Examples 1 to 3 and Comparative Example 1 were treated with EtBr, fluorescence was measured, and the results are shown in FIG. 16 to FIG. 19.

25 **[0100]** According to FIG. 19, in Comparative Example 1 in which the template DNA was not used, and the PCR reaction was not performed, fluorescence was not generated. In contrast, according to FIG. 9, FIG. 16 and FIG. 17, in Examples 1 and 2 which used the cDNA of A549 cell line and NCI-H-1975 cell line as the templates, sufficient PCR products were produced within a short time of about 15 minutes in both Examples 1 and 2, and the generation of fluorescence was confirmed.

30 **[0101]** According to FIG. 15 and FIG. 18, in Example 3 which used the hexahedral glass matrix, a sufficient PCR product was produced within about 8 to 9 minutes, and the generation of fluorescence was confirmed.

[0102] In the above, only given embodiments of the present invention have been explained in detail, but various variations and modifications are apparent to a person in this art within the technical scope of the present invention, and such variations and modifications are certainly included in the appended claims.

35

Claims

1. A substrate for nucleic acid amplification, comprising:

40

- a transparent matrix;
- a metal layer formed on the transparent matrix;
- an N-heterocyclic carbene compound having one end bonded to the surface of the metal layer; and
- a primer immobilized on the other end of the N-heterocyclic carbene compound.

45

2. The substrate for nucleic acid amplification according to claim 1, wherein the transparent matrix comprises at least one matrix selected from a two-dimensional matrix of a planar type; at least one three-dimensional matrix selected from a spherical type, a hemispherical type, a polyhedral type, a polyprism type, and a cylindrical type; and a mixture type thereof.

50

3. The substrate for nucleic acid amplification according to claim 1, wherein the metal layer further comprises a metal mesh.

4. The substrate for nucleic acid amplification according to claim 1, wherein the metal layer comprises gold (Au) .

55


5. The substrate for nucleic acid amplification according to claim 1, wherein the other end of the N-heterocyclic carbene compound is functionalized with azide, phthalimide, or amine.

6. The substrate for nucleic acid amplification according to claim 1, wherein the N-heterocyclic carbene compound is

represented by the following Chemical Formula 1 or 2:

[Chemical Formula 1]


5



10

[Chemical Formula 2]

15



20

25

in Chemical Formulae 1 and 2,

30

R1, R2, R5 and R6 are the same or different, and are each independently hydrogen, an alkyl group of 1 to 20 carbon atoms, a cycloalkyl group of 3 to 20 carbon atoms, an aryl group of 6 to 30 carbon atoms or a heteroaryl group of 2 to 30 carbon atoms,

35

R3, R4, R7, R8, R9 and R10 are the same or different, and are each independently, hydrogen, an alkyl group of 1 to 20 carbon atoms, a cycloalkyl group of 3 to 20 carbon atoms, an aryl group of 6 to 30 carbon atoms, a heteroaryl group of 2 to 30 carbon atoms or a structure represented by the following Chemical Formula 3, where adjacent two or more substituents among R7 to R10 are bonded to form a hydrocarbon ring,


40

at least one of R3 and R4 is the structure represented by the following Chemical Formula 3, and at least one of R7 to R10 is substituted with the structure represented by the following Chemical Formula 3, or in case where adjacent two or more substituents among R7 to R10 are bonded to form a hydrocarbon ring, at least one of carbon atoms forming the hydrocarbon ring is substituted with the structure represented by the following Chemical Formula 3:

45

[Chemical Formula 3]

50



in Chemical Formula 3,

55

n is a repeating number of a unit in parenthesis and an integer of 1 to 30, and A is a nitrogen (N)-containing alkyl group of 1 to 20 carbon atoms or a nitrogen (N)-containing heteroaryl group of 2 to 30 carbon atoms.

7. The substrate for nucleic acid amplification according to claim 1, wherein the primer is a primer of at least one type.

8. A method for manufacturing a substrate for nucleic acid amplification, the method comprising:

forming a metal layer on a transparent matrix;
 introducing one end of an N-heterocyclic carbene compound onto the surface of the metal layer; and
 immobilizing a primer on the other end of the N-heterocyclic carbene compound.

9. The method for manufacturing a substrate for nucleic acid amplification according to claim 8, wherein the transparent matrix comprises at least one matrix selected from a two-dimensional matrix of a planar type; at least one three-dimensional matrix selected from a spherical type, a hemispherical type, a polyhedral type, a polyprism type, and a cylindrical type; and a mixture type thereof.

10. The method for manufacturing a substrate for nucleic acid amplification according to claim 8, wherein the metal layer further comprises a metal mesh.

11. The method for manufacturing a substrate for nucleic acid amplification according to claim 8, wherein the one end of the N-heterocyclic carbene compound is introduced onto the surface of the metal layer through a metal-carbene bond.


12. The method for manufacturing a substrate for nucleic acid amplification according to claim 8, wherein the other end of the N-heterocyclic carbene compound is functionalized with azide, phthalimide, or amine.

13. The method for manufacturing a substrate for nucleic acid amplification according to claim 8, wherein the primer is immobilized on the other end of the N-heterocyclic carbene compound by click reaction.


14. The method for manufacturing a substrate for nucleic acid amplification according to claim 8, wherein the metal layer comprises gold (Au).

15. The method for manufacturing a substrate for nucleic acid amplification according to claim 8, wherein the N-heterocyclic carbene compound is represented by the following Chemical Formula 1 or 2:

[Chemical Formula 1]



[Chemical Formula 2]



in Chemical Formulae 1 and 2,

EP 3 821 982 A1


R1, R2, R5 and R6 are the same or different, and are each independently hydrogen, an alkyl group of 1 to 20 carbon atoms, a cycloalkyl group of 3 to 20 carbon atoms, an aryl group of 6 to 30 carbon atoms or a heteroaryl group of 2 to 30 carbon atoms,

R3, R4, R7, R8, R9 and R10 are the same or different, and are each independently hydrogen, an alkyl group of 1 to 20 carbon atoms, a cycloalkyl group of 3 to 20 carbon atoms, an aryl group of 6 to 30 carbon atoms, a heteroaryl group of 2 to 30 carbon atoms or a structure represented by the following Chemical Formula 3, where adjacent two or more substituents among R7 to R10 are bonded to form a hydrocarbon ring,

at least one of R3 and R4 is the structure represented by the following Chemical Formula 3, and

at least one of R7 to R10 is substituted with the structure represented by the following Chemical Formula 3, or in case where adjacent two or more substituents among R7 to R10 are bonded to form a hydrocarbon ring, at least one of carbon atoms forming the hydrocarbon ring is substituted with the structure represented by the following Chemical Formula 3:

[Chemical Formula 3]



in Chemical Formula 3,

n is a repeating number of a unit in parenthesis and an integer of 1 to 30, and

A is a nitrogen (N)-containing alkyl group of 1 to 20 carbon atoms or a nitrogen (N)-containing heteroaryl group of 2 to 30 carbon atoms.

16. The method for manufacturing a substrate for nucleic acid amplification according to claim 8, wherein the primer is a primer of at least one type.

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG.9




FIG.10




FIG. 11




FIG. 12




FIG.13




FIG.14




FIG.15




FIG.16




FIG.17




FIG.18






FIG.19



1000 μm

FIG. 20



Gold mesh

FIG.21





FIG.22



INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR2019/008663

5

A. CLASSIFICATION OF SUBJECT MATTER
B01L 7/00(2006.01); B01J 19/00(2006.01);
 According to International Patent Classification (IPC) or to both national classification and IPC

10

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 B01L 7/00; B01L 3/00; C12M 1/34; C12M 1/38; C12N 15/09; C12Q 1/68; B01J 19/00

15

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 Korean utility models and applications for utility models: IPC as above
 Japanese utility models and applications for utility models: IPC as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 eKOMPASS (KIPO internal), STN (Registry, Caplus), Google & Keywords: PCR(Polymerase Chain Reaction), nucleic acid amplification, transparent, metal, gold(au), N-heterocyclic carbene, azide, phthalimide, amine

20

C. DOCUMENTS CONSIDERED TO BE RELEVANT

25

30

35

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2013-105679 A1 (LG ELECTRONICS INC.) 18 July 2013 See paragraph [64]; claims 1-19; and figure 7.	1-16
Y	LI, Z. et al. Self-Assembled N-Heterocyclic Carbene-Based Carboxymethylated Dextran Monolayers on Gold as a Tunable Platform for Designing Affinity-Capture Biosensor Surfaces. ACS applied materials & interfaces. 09 May 2018, vol. 10, no. 21, pages 17560-17570 See abstract.	1-16
Y	WANG, X. et al. Peptide-decorated gold nanoparticles via strain-promoted azide-alkyne cycloaddition and post assembly deprotection. RSC Advance. 02 September 2014, vol. 4, no. 81, pages 43087-43091 See abstract; and page 43090, figure 1.	6,13,15
A	KR 10-2012-0020528 A (SAMSUNG ELECTRONICS CO., LTD.) 08 March 2012 See claims 1, 9; and paragraphs [0021], [0025].	1-16
A	KR 10-2018-0034579 A (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) 04 April 2018 See claims 1-23.	1-16

40

Further documents are listed in the continuation of Box C. See patent family annex.

45

* Special categories of cited documents:
 "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier application or patent but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed
 "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 "&" document member of the same patent family

50

Date of the actual completion of the international search 17 OCTOBER 2019 (17.10.2019)	Date of mailing of the international search report 18 OCTOBER 2019 (18.10.2019)
---	--

55

Name and mailing address of the ISA/KR  Korean Intellectual Property Office Government Complex Daejeon Building 4, 189, Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea Facsimile No. +82-42-481-8578	Authorized officer Telephone No.
---	---

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR2019/008663

5
10
15
20
25
30
35
40
45
50
55

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP 2010-041985 A (HITACHI PLANT TECHNOLOGIES LTD.) 25 February 2010 See claims 1-6.	1-16

Form PCT/ISA/210 (continuation of second sheet) (January 2015)

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/KR2019/008663

5
10
15
20
25
30
35
40
45
50
55

Patent document cited in search report	Publication date	Patent family member	Publication date
WO 2013-105679 A1	18/07/2013	None	
KR 10-2012-0020528 A	08/03/2012	US 2012-0052531 A1 US 8663976 B2	01/03/2012 04/03/2014
KR 10-2018-0034579 A	04/04/2018	CN 107921399 A EP 3328535 A1 EP 3328535 A4 JP 2018-521665 A US 2018-0236451 A1 WO 2017-019768 A1	17/04/2018 06/06/2018 27/02/2019 09/08/2018 23/08/2018 02/02/2017
JP 2010-041985 A	25/02/2010	None	

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- KR 20170106995 [0004] [0006]